# arXiv:2501.11956v1 [gr-qc] 21 Jan 2025

# Modified gravitational wave propagations in linearized gravity with Lorentz and diffeomorphism violations and their gravitational wave constraints

Qiang Wang<sup>*a,b*</sup>, \* Jian-Ming Yan<sup>*a,b*</sup>, † Tao Zhu<sup>*a,b*</sup>, ‡ and Wen Zhao<sup>*c,d*§</sup>

<sup>a</sup> Institute for Theoretical Physics & Cosmology, Zhejiang University of Technology, Hangzhou, 310023, China

<sup>b</sup> United Center for Gravitational Wave Physics (UCGWP),

Zhejiang University of Technology, Hangzhou, 310023, China

<sup>c</sup>CAS Key Laboratory for Research in Galaxies and Cosmology, Department of Astronomy,

University of Science and Technology of China, Hefei 230026, China

<sup>d</sup>School of Astronomy and Space Sciences, University of Science and Technology of China, Hefei, 230026, China

(Dated: January 22, 2025)

The standard model extension (SME) is an effective field theory framework that can be used to study the possible violations of Lorentz symmetry and diffeomorphism invariance in the gravitational interaction. In this paper, we explore both the Lorentz- and diffeomorphism-violating effects on the propagations of gravitational waves in the SME's linearized gravity. It is shown that the violations of Lorentz symmetry and diffeomorphism invariance modify the conventional linear dispersion relation of gravitational waves, leading to anisotropy, birefringence, and dispersion effects in the propagation of gravitational waves. With these modified dispersion relations, we then calculate the dephasing effects due to the Lorentz and diffeomorphism violations in the waveforms of gravitational waves produced by the coalescence of compact binaries. With the distorted waveforms, we perform full Bayesian inference with the help of the open source software BILBY on the gravitational wave events of the compact binary mergers in the LIGO-Virgo-KAGRA catalogs GWTC-3. We consider the effects from the operators with the lowest mass dimension d = 2 and d = 3 due to the Lorentz and diffeomorphism violations in the linearized gravity. No signature of Lorentz and diffeomorphism violations arsing from the SME's linearized gravity are found for most GW events, which allows us to give a 90% confidence interval for each Lorentz- and diffeomorphism-violating coefficient.

### I. INTRODUCTION

General relativity (GR) stands as the preeminent theory of gravity, having undergone rigorous experimental validation across a diverse range of scales with remarkable precision [1-10]. Despite its empirical success, GR encounters significant challenges related to the theoretical singularities and issues of quantization, as well as the unresolved phenomena of dark matter and dark energy. Conversely, several candidate theories of quantum gravity, including string theory [11, 12], loop quantum gravity [13], and brane-world scenarios [14], propose frameworks wherein Lorentz and diffeomorphism invariances in the gravitational sector may be spontaneously violated. This potential breakdown of Lorentz and diffeomorphism invariances presents intriguing avenues for addressing some of the limitations of GR and advancing our understanding of fundamental physics.

A common method to investigate potential violations of Lorentz and diffeomorphism symmetries in gravity is through the approach of effective field theory. The Standard-Model Extension (SME) offers a comprehensive framework for examining deviations from Lorentz and diffeomorphism invariance [15, 16]. Within this approach, any components that might disrupt Lorentz or/and diffeomorphism invariance can be systematically included in the Lagrangian. Over recent decades, the SME has been widely applied to test Lorentz invariance in the matter sector. In the context of gravity, studies leveraging the SME to evaluate the Lorentz violations have encompassed a range of techniques including lunar laser ranging [17, 18], atom interferometry [19], cosmic ray observations [20], precision pulsar timing [21– 26], planetary orbital analyses [27], and superconducting gravimeters [28]. Studies in the gravity domain are primarily concerned with the interaction between gravity and matter [29]. Our focus, however, is on the Lorentz and diffeomorphism-violating effects on gravitational wave (GW) propagation in the pure gravity sector within a linear approximation [30–32].

In linearized Lorentz-violating gravity, the possible Lorentz-violating terms involving quadratic metric variations in the Lagrangian can be classified by their mass dimension, d. These terms are further divided into two categories: diffeomorphism-invariant terms and diffeomorphism-violating terms, depending on whether they remain invariant under the gauge transformation  $h_{\mu\nu} \rightarrow h_{\mu\nu} + \partial_{\mu}\xi_{\nu} + \partial_{\nu}\xi_{\mu}$ . Here,  $h_{\mu\nu}$  represents metric perturbations in Minkowski space-time, and  $\xi_{\mu}$  is an arbitrary small vector field. Diffeomorphism invariance requires that  $d \ge 4$  [32, 33]. The effects of Lorentz violations on gravitational waves (GWs) and their observational constraints have been extensively studied for the diffeomorphism-invariant case; for example, [32–43] and references therein. In contrast, for the diffeomorphismviolating case, the Lorentz-violating terms in the Lagrangian can have mass dimensions d = 2 or/and d = 3

<sup>\*</sup> qiangwang@zjut.edu.cn

<sup>†</sup> yanjm@zjut.edu.cn

<sup>&</sup>lt;sup>‡</sup> Corresponding author: zhut05@zjut.edu.cn

<sup>&</sup>lt;sup>§</sup> wzhao7@ustc.edu.cn

An exhaustive classification of Lorentz-violating Lagrangians with quadratic metric variations, whether diffeomorphism-invariant or -violating, is provided in [32]. These terms modify GW dispersion relations, leading to effects such as anisotropy, birefringence, and dispersion, which alter GW waveforms. Waveform modifications for the diffeomorphism-invariant case are discussed in [43] and can be analyzed using Bayesian inference to compare observed GW signals with theoretical models, constraining the Lorentz-violating coefficients in the SME's linearized gravity sector. A parametrized framework for describing symmetry-breaking effects on GWs is presented in [37], which has been used for calculating the symmetry-breaking effects on the primordial GWs [44]. In this paper, we investigate both Lorentz-violating and diffeomorphism-violating effects on GW propagation and calculate the resulting dephasing in waveforms from compact binary coalescences.

The direct detection of GWs from compact binary coalescences by the LIGO-Virgo-KAGRA (LVK) Collaboration has ushered in a new era of gravitational physics [45–52]. These GWs carry crucial information about the local spacetime properties of compact binaries, enabling tests of fundamental gravitational symmetries. Numerous studies have tested Lorentz and parity symmetries using data from LVK GW events [33, 39–41, 53–69]; see [70] for a recent review. While previous tests in Lorentz-violating linearized gravity for diffeomorphisminvariant cases focused on terms with mass dimension  $d \geq 4$ , this paper examines the effects of diffeomorphism violations induced by Lorentz-violating terms with d = 2 and d = 3, along with their observational constraints from LVK GW events. Using the SME framework for Lorentz-violating linearized gravity, we perform Bayesian inference with modified waveforms incorporating diffeomorphism-violating effects on GW events from the GWTC-3 catalog. We find no significant evidence of diffeomorphism violations in most GW data and provide 90% confidence intervals for each diffeomorphismviolating coefficient.

This paper is organized as follows. In the next section, we provide a brief overview of GW propagation in the SME framework, incorporating both Lorentz and diffeomorphism violations and their modified dispersion relations. Section III examines phase modifications to GW waveforms caused by Lorentz-violating and diffeomorphism-violating coefficients in the SME. In Section IV, we describe the matched-filter analysis within Bayesian inference, and in Section V, we present constraints on each diffeomorphism-violating coefficient using data from GW events in the GWTC-3 catalog. Finally, the conclusions and summary are provided in Section VI.

Throughout this paper, we adopt the metric convention (-, +, +, +), with Greek indices  $(\mu, \nu, ...)$  running over 0, 1, 2, 3 and Latin indices (i, j, k) running over 1, 2, 3. Natural units are used, setting  $\hbar = c = 1$ .

# II. GRAVITATIONAL WAVE PROPAGATIONS IN THE LINEAR GRAVITY OF SME

In this section, we present a brief introduction to the GWs in the linearized gravity sector of the SME and the associated modified dispersion relation of GWs due to the effects of the Lorentz and diffeomorphism violations.

# A. Linearized gravity with Lorentz and diffeomorphism violations

The quadratic Lagrangian density for GWs in the linearized gravity sector of the SME is given by [32]

$$\mathcal{L} = \frac{1}{4} \epsilon^{\mu\rho\alpha\kappa} \epsilon^{\nu\sigma\beta\lambda} \eta_{\kappa\lambda} h_{\mu\nu} \partial_{\alpha} \partial_{\beta} h_{\rho\sigma} + \frac{1}{4} h_{\mu\nu} \sum_{\mathcal{K},d} \hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma} h_{\rho\sigma}, \qquad (2.1)$$

where one expands the metric  $g_{\mu\nu}$  of the spacetime in the form of  $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$  with  $\eta_{\mu\nu}$  being the constant Minkowski metric, and  $\epsilon^{\mu\rho\alpha\kappa}$  is the Levi-Civita tensor. The first term in the above expression represents the quadratic approximation to the Lagrangian density for the Einstein-Hilbert action, while the second term which consists of operators  $\hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma}$  denotes the modifications due to the Lorentz and diffeomorphism violations. The operator  $\hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma}$  is the product of a coefficient  $\mathcal{K}^{(d)\mu\nu\rho\sigma\varepsilon_1\varepsilon_2\cdots\varepsilon_{d-2}}$  with d-2 derivatives  $\partial_{\varepsilon_1}\partial_{\varepsilon_2}\cdots\partial_{\varepsilon_{d-2}}$ , i.e.,

$$\hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma} = \mathcal{K}^{(d)\mu\nu\rho\sigma\varepsilon_1\varepsilon_2\cdots\varepsilon_{d-2}}\partial_{\varepsilon_1}\partial_{\varepsilon_2}\cdots\partial_{\varepsilon_{d-2}}.$$
 (2.2)

The coefficient  $\mathcal{K}^{(d)\mu\nu\rho\sigma\varepsilon_1\varepsilon_2\cdots\varepsilon_{d-2}}$  has mass dimension 4-d and are assumed small and constant over the scales relevant for the gravitational phenomenon considered in this paper. As pointed out in [32], to contribute non-trivially to the equation of motion, the  $\hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma}$  has to satisfy the requirement  $\hat{\mathcal{K}}^{(d)(\mu\nu)(\rho\sigma)} \neq \pm \hat{\mathcal{K}}^{(d)(\rho\sigma)(\mu\nu)}$ , where  $\pm$  corresponds to odd d and even d respectively.

According to the the symmetries of the coefficient  $\mathcal{K}^{(d)\mu\nu\rho\sigma\varepsilon_1\varepsilon_2\cdots\varepsilon_{d-2}}$  under the permutations of its indices,  $\hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma}$  can be divided into three different types,

$$\hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma} = \hat{s}^{(d)\mu\rho\nu\sigma} + \hat{q}^{(d)\mu\rho\nu\sigma} + \hat{k}^{(d)\mu\rho\nu\sigma}.$$
 (2.3)

These three types of operators have different symmetries in their indices. Specifically,  $\hat{s}^{(d)\mu\rho\nu\sigma}$  is anti-symmetric in both " $\mu\rho$ " and " $\nu\sigma$ ",  $\hat{q}^{(d)\mu\rho\nu\sigma}$  is anti-symmetric in " $\mu\rho$ " and symmetric in " $\nu\sigma$ ", and  $\hat{k}^{(d)\mu\rho\nu\sigma}$  is totally symmetric. One can further decompose each type of operator into several irreducible pieces and explore their proerties on the gravitational wave propagations. It is shown in [32] such a decomposition leads to 14 independent classess of operators in total, as preented in Table I, see also Table I of ref. [32]. The s-type operators  $\hat{s}^{(d)\mu\rho\nu\sigma}$  are CPT even, denoting that these operators are invariant under the combined symmetry of change conjugation, parity transformation, and time reversal (CPT). They consist of three irreducible pieces in the decomposion,

$$\hat{s}^{(d)\mu\rho\nu\sigma} = \hat{s}^{(d,0)\mu\rho\nu\sigma} + \hat{s}^{(d,1)\mu\rho\nu\sigma} + \hat{s}^{(d,2)\mu\rho\nu\sigma}, \quad (2.4)$$

with

$$\hat{s}^{(d,0)\mu\rho\nu\sigma} = s^{(d,0)\mu\rho\alpha_{1}\nu\sigma\alpha_{2}\alpha_{3}\dots\alpha_{d-2}}\partial_{\alpha_{1}}\dots\partial_{\alpha_{d-2}},$$
$$\hat{s}^{(d,1)\mu\rho\nu\sigma} = s^{(d,1)\mu\rho\nu\sigma\alpha_{1}\dots\alpha_{d-2}}\partial_{\alpha_{1}}\dots\partial_{\alpha_{d-2}},$$
$$\hat{s}^{(d,2)\mu\rho\nu\sigma} = s^{(d,2)\mu\rho\alpha_{1}\nu\sigma\alpha_{2}\alpha_{3}\dots\alpha_{d-2}}\partial_{\alpha_{1}}\dots\partial_{\alpha_{d-2}}.$$

The q-type operators  $\hat{q}^{(d)\mu\rho\nu\sigma}$  are CPT odd, denoting that they change sign under the symmetry of CPT. They consist of six irreducible pieces in the decomposion,

$$\hat{q}^{(d)\mu\rho\nu\sigma} = \hat{q}^{(d,0)\mu\rho\nu\sigma} + \hat{q}^{(d,1)\mu\rho\nu\sigma} + \hat{q}^{(d,2)\mu\rho\nu\sigma} 
+ \hat{q}^{(d,3)\mu\rho\nu\sigma} + \hat{q}^{(d,4)\mu\rho\nu\sigma} + \hat{q}^{(d,5)\mu\rho\nu\sigma},$$
(2.6)

where

$$\hat{q}^{(d,0)\mu\rho\nu\sigma} = \hat{q}^{(d,0)\mu\rho\alpha_{1}\nu\alpha_{2}\sigma\alpha_{3}\alpha_{4}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}},$$

$$\hat{q}^{(d,1)\mu\rho\nu\sigma} = \hat{q}^{(d,1)\mu\rho\nu\sigma\alpha_{1}\alpha_{2}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}},$$

$$\hat{q}^{(d,2)\mu\rho\nu\sigma} = \hat{q}^{(d,2)\mu\rho\nu\alpha_{1}\sigma\alpha_{2}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}},$$

$$\hat{q}^{(d,3)\mu\rho\nu\sigma} = \hat{q}^{(d,3)\mu\rho\alpha_{1}\nu\sigma\alpha_{2}\alpha_{3}\alpha_{4}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}},$$

$$\hat{q}^{(d,5)\mu\rho\nu\sigma} = \hat{q}^{(d,5)\mu\rho\alpha_{1}\nu\alpha_{2}\sigma\alpha_{3}\alpha_{4}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}}.$$

$$(2.7)$$

And the k-type operators are CPT even and consist of five irreducible pieces,

$$\hat{k}^{(d)\mu\nu\rho\sigma} = \hat{k}^{(d,0)\mu\nu\rho\sigma} + \hat{k}^{(d,1)\mu\nu\rho\sigma} + \hat{k}^{(d,2)\mu\nu\rho\sigma} \\
+ \hat{k}^{(d,3)\mu\nu\rho\sigma} + \hat{k}^{(d,4)\mu\nu\rho\sigma},$$
(2.8)

where

$$\hat{k}^{(d,0)\mu\nu\rho\sigma} = \hat{k}^{(d,0)\mu\alpha_{1}\nu\alpha_{2}\rho\alpha_{3}\sigma\alpha_{4}\alpha_{5}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}},$$

$$\hat{k}^{(d,1)\mu\nu\rho\sigma} = \hat{k}^{(d,1)\mu\nu\rho\sigma\alpha_{1}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}},$$

$$\hat{k}^{(d,2)\mu\nu\rho\sigma} = \hat{k}^{(d,2)\mu\alpha_{1}\nu\rho\sigma\alpha_{1}\alpha_{2}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}},$$

$$\hat{k}^{(d,3)\mu\nu\rho\sigma} = \hat{k}^{(d,3)\mu\alpha_{1}\nu\alpha_{2}\rho\sigma\alpha_{3}\sigma\alpha_{4}\alpha_{5}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}},$$

$$\hat{k}^{(d,4)\mu\nu\rho\sigma} = \hat{k}^{(d,4)\mu\alpha_{1}\nu\alpha_{2}\rho\alpha_{3}\sigma\alpha_{4}\alpha_{5}...\alpha_{d-2}}\partial_{\alpha_{1}}...\partial_{\alpha_{d-2}}.$$
(2.9)

The properties of the above 14 coefficients are summarized in Table. I, see also Table I in [32] for more detailed properties of these coefficients. These 14 classes thus characterize all possible phenomenological effects in linearized gravity, affecting the propagating properties of gravitational waves. If we restrict the theory to be diffeomorphism invariance, the quadratic action  $S \sim \int d^4 x \mathcal{L}$  of the linearized gravity should be invariant under the the gauge transformation  $h_{\mu\nu} \rightarrow h_{\mu\nu} + \partial_{\mu}\xi_{\nu} + \partial_{\nu}\xi_{\mu}$ , which requires the condition  $\hat{\mathcal{K}}^{(d)(\mu\nu)(\rho\sigma)}\partial_{\nu} = \pm \hat{\mathcal{K}}^{(d)(\rho\sigma)(\mu\nu)}\partial_{\nu}$  holds. With this condition, the operator  $\hat{\mathcal{K}}^{(d)(\mu\nu\rho\sigma)}$  can only be decomposed into three independent classes [32], which are represents by  $\hat{s}^{(d,0)\mu\rho\nu\sigma}$ ,  $\hat{q}^{(d,0)\mu\rho\nu\sigma}$ , and  $\hat{k}^{(d,0)\mu\rho\nu\sigma}$ , respectively. It is obvious that the diffeomorphism-invariance case only allows the Lorentz-violating operators with mass dimension  $d \geq 4$ , while the diffeomorphism-violating case allows operators with mass dimension  $d \geq 2$ .

# B. GW propagations with Lorentz- and diffeomorphism-violating effects

The equations of motion for GWs can be derived by varying the quadratic action  $S \sim \int d^4x \mathcal{L}$  with respect to  $h_{\mu\nu}$  with Lagrangian density  $\mathcal{L}$  given by (2.1), which yields

$$\frac{1}{2}\eta_{\rho\sigma}\epsilon^{\mu\rho\alpha\kappa}\epsilon^{\nu\sigma\beta\lambda}\partial_{\alpha}\partial_{\beta}h_{\kappa\lambda} - \delta M^{\mu\nu\rho\sigma}h_{\rho\sigma} = 0, \quad (2.10)$$

where the tensor operators

$$\delta M^{\mu\nu\rho\sigma} = -\frac{1}{4} \left( \hat{s}^{\mu\rho\nu\sigma} + \hat{s}^{\mu\sigma\nu\rho} \right) - \frac{1}{2} \hat{k}^{\mu\nu\rho\sigma} - \frac{1}{8} \left( \hat{q}^{\mu\rho\nu\sigma} + \hat{q}^{\nu\rho\mu\sigma} + \hat{q}^{\mu\sigma\nu\rho} + \hat{q}^{\nu\sigma\mu\rho} \right).$$
(2.11)

Here

$$\hat{s}^{\mu\rho\nu\sigma} = \sum_{d} \hat{s}^{(d)\mu\rho\nu\sigma}, \qquad (2.12)$$

$$\hat{q}^{\mu\rho\nu\sigma} = \sum_{d} \hat{q}^{(d)\mu\rho\nu\sigma}, \qquad (2.13)$$

$$\hat{k}^{\mu\rho\nu\sigma} = \sum_{d} \hat{k}^{(d)\mu\rho\nu\sigma}.$$
(2.14)

In GR, the metric perturbation  $h_{\mu\nu}$  contains only two degenerate, traceless, and transverse tensor modes. However, when Lorentz-violating and diffeomorphismviolating modifications are introduced,  $h_{\mu\nu}$  may acquire additional degrees of freedom, depending on the specific nature of the violations. These include two scalar modes and two vector modes. As shown in [34], in Lorentzviolating linearized gravity, the extra scalar and vector modes of gravitational wave (GW) polarization can be directly induced by the two tensorial modes. In this paper, assuming these additional modes are small, and following a similar treatment to that in [33], we focus exclusively on the effects of Lorentz- and diffeomorphism-violating modifications on the two traceless and transverse tensor modes.

With the above considerations, we restrict our analysis to the modes  $h_{ij}$  that satisfy the conditions

$$\eta^{ij}h_{ij} = 0, \quad \partial^i h_{ij} = 0.$$
 (2.15)

Coefficients $\hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma\alpha_1\alpha_{d-2}}$	Young Tableau	d	number
$s^{(d,0)\mu\rho\alpha_1\nu\sigma\alpha_2\alpha_3\alpha_{d-2}}$	$ \begin{array}{c cccc} \mu & \nu & \alpha_3 \\ \hline \rho & \sigma \\ \hline \alpha_1 & \alpha_2 \end{array} \cdots  \boxed{\alpha_{d-2}} $	even $d \ge 4$	(d-3)(d-2)(d+1)
$s^{(d,1)\mu\rho\nu\sigma\alpha_1\alpha_{d-2}}$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	even $d \ge 2$	(d-1)(d+2)(d+3)
$s^{(d,2)\mu\rho\alpha_1\nu\sigma\alpha_2\alpha_3\alpha_{d-2}}$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	even $d \ge 4$	$\frac{4}{3}(d-2)d(d+2)$
$\hat{q}^{(d,0)\mu\rho\alpha_1\nu\alpha_2\sigma\alpha_3\alpha_4\alpha_{d-2}}$	$ \begin{array}{ c c c c c c c c } \mu & \nu & \sigma & \alpha_4 \\ \hline \rho & \alpha_2 & \alpha_3 \\ \hline \alpha_1 \end{array} & \cdots & \alpha_{d-2} \end{array} $	odd $d \ge 5$	$\frac{5}{2}(d-4)(d-1)(d+1)$
$\hat{q}^{(d,1)\mu\rho\nu\sigma\alpha_1\alpha_2\alpha_{d-2}}$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	odd $d \ge 3$	$\frac{1}{2}(d-3)(d+4)(d+1)$
$\hat{q}^{(d,2)\mu\rho\nu\alpha_1\sigma\alpha_2\alpha_{d-2}}$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	odd $d \ge 3$	(d-1)(d+2)(d+3)
$\hat{q}^{(d,3)\mu\rholpha_1 u\sigmalpha_2lpha_{d-2}}$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	odd $d \ge 3$	$\frac{1}{2}d(d+3)(d+1)$
$\hat{q}^{(d,4)\mu\rho\nu\alpha_1\sigma\alpha_2\alpha_3\alpha_4\alpha_{d-2}}$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	odd $d \ge 5$	$\frac{5}{3}(d-3)(d+2)(d+1)$
$\hat{q}^{(d,5)\mu\rho\alpha_1\nu\alpha_2\sigma\alpha_3\alpha_4\alpha_{d-2}}$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	odd $d \ge 5$	$\frac{4}{3}(d+2)d(d-2)$
$\hat{k}^{(d,0)\mu\alpha_1\nu\alpha_2\rho\alpha_3\sigma\alpha_4\alpha_5\alpha_{d-2}}$	$\begin{array}{ c c c c c c c c } \mu & \nu & \rho & \sigma & \alpha_5 \\ \hline \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \hline \end{array} \qquad \cdots \qquad \boxed{\alpha_{d-2}}$	even $d \ge 6$	$\frac{5}{2}(d-5)d(d+1)$
$\hat{k}^{(d,1)\mu\nu\rho\sigma\alpha_1\alpha_{d-2}}$	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	even $d \ge 2$	$\frac{1}{6}(d+3)(d+4)(d+5)$
$\hat{k}^{(d,2)\mu\alpha_1\nu\rho\sigma\alpha_1\alpha_2\alpha_{d-2}}$	$\begin{array}{ c c c c c c c } \mu & \nu & \rho & \sigma & \alpha_2 & \alpha_3 & \cdots & \alpha_{d-2} \\ \hline \alpha_1 & & & & & \\ \hline \end{array}$	even $d \ge 4$	$\frac{1}{2}(d+1)(d+3)(d+4)$
$\hat{k}^{(d,3)\mu\alpha_1\nu\alpha_2\rho\sigma\alpha_3\alpha_5\alpha_{d-2}}$	$\begin{array}{ c c c c c c } \mu & \nu & \rho & \sigma & \alpha_3 \\ \hline \alpha_1 & \alpha_2 \\ \hline \end{array} & & & & & & & & & & & & & & & & & &$	even $d \ge 4$	(d-1)(d+2)(d+3)
$\hat{k}^{(d,4)\mu\alpha_1\nu\alpha_2\rho\alpha_3\sigma\alpha_4\alpha_5\alpha_{d-2}}$	$\begin{array}{ c c c c c c c c } \mu & \nu & \rho & \sigma & \alpha_4 & \alpha_5 & \cdots & \alpha_{d-2} \\ \hline \alpha_1 & \alpha_2 & \alpha_3 & & & & \\ \hline \end{array}$	even $d \ge 6$	$\frac{5}{3}(d-3)(d+2)(d+1)$

TABLE I. Properties of the coefficients  $\hat{\mathcal{K}}^{(d)\mu\nu\rho\sigma\alpha_1...\alpha_{d-2}}$  in each irreducible class. A similar table can also be found in Table. I of ref. [32].

Under these constraints, the equations of motion for GWs given in Eq. (2.10) reduce to

$$(\partial_t^2 - \nabla^2)h^{ij} + 2\delta M^{ijmn}h_{mn} = 0.$$
 (2.16)

In the linearized gravity sector of SME, it is convenient to decompose the GWs into circular polarization modes. To study the evolution of  $h_{ij}$ , we expand it in terms of spatial Fourier harmonics as

$$h_{ij}(\tau, x^{i}) = \sum_{A=R,L} \int \frac{d^{3}k}{(2\pi)^{3}} h_{A}(\tau, k^{i}) e^{ik_{i}x^{i}} e^{A}_{ij}(k^{i}), \qquad \text{or}$$

$$\begin{pmatrix} \partial_{t}^{2} + k^{2} + 2e^{R}_{ij}\delta M^{ijmn}e^{R}_{mn} & 2e^{R}_{ij}\delta M^{ijmn}e^{R}_{mn} \\ 2e^{L}_{ij}\delta M^{ijmn}e^{R}_{mn} & \partial_{t}^{2} + k^{2} + 2e^{R}_{ij}\delta M^{ijmn}e^{R}_{mn} \\ \end{pmatrix}$$

where  $e_{ij}^{A}$  are the circular polarization tensors, and R and L denote the right-handed and left-handed GW polarizations, respectively. The circular polarization tensors  $e_{ij}^{A}$  satisfy the relation

$$\epsilon^{ijk} n_j e^A_{kl} = i\rho_A e^{iA}_{\ l}, \qquad (2.18)$$

with  $\rho_{\rm R} = 1$  and  $\rho_{\rm L} = -1$ . Using this decomposition, the equations of motion in Eq. (2.16) can be rewritten as

$$\ddot{h}_A + k^2 h_A + 2\epsilon^A_{ij} \delta M^{ijmn} e^B_{mn} h_B = 0, \qquad (2.19)$$

or equivalently, in matrix form:

$$\frac{\partial_t^2 + k^2 + 2e_{ij}^R \delta M^{ijmn} e_{mn}^R}{2e_{ij}^L \delta M^{ijmn} e_{mn}^R} \frac{2e_{ij}^R \delta M^{ijmn} e_{mn}^L}{\partial_t^2 + k^2 + 2e_{ij}^L \delta M^{ijmn} e_{mn}^L} \left( \begin{array}{c} h_R \\ h_L \end{array} \right) = 0.$$

$$(2.20)$$

Then, following methods developed for the study of Lorentz violation in the photon sector of the SME [71] as well as GW propagations with diffeomorphism invariance, the modified dispersion relation of GWs with 4-momentum  $k^{\mu} = (\omega, \mathbf{k})$  can be derived by requiring the determinant of the above 2 × 2 matrix vanishes, which yields (see also in [33])<sup>1</sup>

$$\omega = \left(1 - \zeta^0 \pm |\boldsymbol{\zeta}|\right) |\mathbf{k}|, \qquad (2.21)$$

where

$$\zeta^{0} = -\frac{1}{2|\mathbf{k}|^{2}} \Big( e_{ij}^{R} \delta M^{ijmn} e_{mn}^{R} + e_{ij}^{L} \delta M^{ijmn} e_{mn}^{L} \Big),$$
(2.22)

and

$$|\boldsymbol{\zeta}|^{2} = \frac{1}{4|\mathbf{k}|^{4}} \Big[ (e_{ij}^{R} \delta M^{ijmn} e_{mn}^{R} - e_{ij}^{L} \delta M^{ijmn} e_{mn}^{L})^{2} + 4 (e_{ij}^{R} \delta M^{ijmn} e_{mn}^{L}) (e_{kl}^{L} \delta M^{klpq} e_{pq}^{R}) \Big]. \quad (2.23)$$

And  $|\boldsymbol{\zeta}|^2 \equiv (\zeta^1)^2 + (\zeta^2)^2 + (\zeta^3)^2$  with

$$\zeta^{1} - i\zeta^{2} = \frac{1}{|\mathbf{k}|^{2}} (e_{ij}^{R} \delta M^{ijmn} e_{mn}^{L}), \qquad (2.24)$$

$$\zeta^{1} + i\zeta^{2} = \frac{1}{|\mathbf{k}|^{2}} (e_{ij}^{L} \delta M^{ijmn} e_{mn}^{R}), \qquad (2.25)$$

$$\zeta^{3} = \frac{1}{2|\mathbf{k}|^{2}} (e_{ij}^{R} \delta M^{ijmn} e_{mn}^{R} - e_{ij}^{L} \delta M^{ijmn} e_{mn}^{L}).$$
(2.26)

The modified dispersion relation in the above leads to the phase velocities  $(v = \omega/k)$  of the GWs

$$v_{\pm} = 1 - \zeta^0 \pm |\boldsymbol{\zeta}| \,.$$
 (2.27)

Here " $\pm$ " correspond to two modes propagating at different velocities. Therefore, the two tensorial modes can be decompose into two modes propagate at different velocities, one is the fast mode (denoted by  $h_f$  with velocity  $v_+$ ) while another is the slow mode (denote by  $h_s$  with velocity  $v_-$ ). One can connect  $(h_f, h_s)$  with the circular polarization modes  $(h_L, h_R)$  by normalized matrix as

$$\begin{pmatrix} h_f \\ h_s \end{pmatrix} = \begin{pmatrix} -e^{i\varphi/2}\sin\frac{\vartheta}{2} & e^{-i\varphi/2}\cos\frac{\vartheta}{2} \\ e^{i\varphi/2}\cos\frac{\vartheta}{2} & e^{i\varphi/2}\sin\frac{\vartheta}{2} \end{pmatrix} \begin{pmatrix} h_R \\ h_L \end{pmatrix}.$$
(2.28)

Here the angles  $\varphi$  and  $\vartheta$  are defined as

$$\sin\vartheta = \frac{\sqrt{(\zeta^1)^2 + (\zeta^2)^2}}{|\zeta|}, \qquad (2.29)$$

$$\cos\vartheta = \frac{\zeta^3}{|\zeta|},\tag{2.30}$$

$$e^{\pm i\varphi} = \frac{\zeta^1 \pm \zeta^2}{\sqrt{(\zeta^1)^2 + (\zeta^2)^2}}.$$
 (2.31)

In the modified dispersion relation (2.21), the coefficient  $\zeta^0$  and  $\zeta$  are functions of the frequency  $\omega$  and wave vector **k** [40]. Considering it's also direction-dependent and to describe its effects on the propagation of GWs, it is convenient to expand its coefficients in terms of spinweighted spherical harmonics  ${}_{s}Y_{jm}(\hat{\mathbf{n}})$  as

$$\zeta^{0} = \sum_{d,jm} \omega^{d-4} Y_{jm}(\hat{\mathbf{n}}) k^{(d)}_{(I)jm}, \qquad (2.32)$$

<sup>&</sup>lt;sup>1</sup> It is worth noting here that the Lorentz- and diffeomorphismviolating operators introduced in (2.1) only affect the dispersion and the corresponding phase velocities of GWs. It is shown in [57, 58, 72, 73] that the Lorentz-violating terms with mixed time and spatial derivatives can change the damping rates of GWs.

$$\zeta^{1} \mp i\zeta^{2} = \sum_{d,jm} \omega^{d-4} \pm_{4} Y_{jm}(\hat{\mathbf{n}}) \Big[ k^{(d)}_{(E)jm} \pm k^{(d)}_{(B)jm} \Big],$$

$$\zeta^{3} = \sum_{d,jm} \omega^{d-4} Y_{jm}(\hat{\mathbf{n}}) k_{(V)jm}^{(d)}, \qquad (2.34)$$

where  $\mathbf{n} = -\mathbf{k}$  is the direction of the source,  $Y_{jm}(\hat{\mathbf{n}}) = {}_{0}Y_{jm}(\hat{\mathbf{n}})$  is the scalar spherical harmonics function, and  $|s| \leq j \leq d-2$ . The indice *m* takes  $-j, \dots, j$ . The spherical coefficients for Lorentz and diffeomorphism violations  $k_{(I)jm}^{(d)}$ ,  $k_{(E)jm}^{(d)}$ ,  $k_{(B)jm}^{(d)}$ , and  $k_{(V)jm}^{(d)}$  are linear combinations of the tensor coefficients in Eqs. (2.5, 2.7, 2.9), which obey the relation  $k_{jm}^{(d)*} = (-1)^m k_{j-m}^{(d)}$ . The expansions of the coefficient  $\zeta^0$ ,  $\zeta^1 \pm i\zeta^2$ , and  $\zeta^3$  are also a combination of operators at multiple mass dimensions, but they have different expansion properties. The mass dimension *d* of the expansion coefficients can take even numbers of  $d \geq 2$  for the expansion of  $\zeta^0$ , odd numbers of  $d \geq 3$  for  $\zeta^3$ , and even number of  $d \geq 6$  for  $\zeta^1 \pm i\zeta^2$ .

For convenience, the frequency and direction dependence can be separated, and we introduce several energyindependent coefficients as

$$\zeta_{(d)}^{0}(\mathbf{n}) = \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(I)jm}^{(d)}, \qquad (2.35)$$

$$\zeta_{(d)}^{1}(\mathbf{n}) \mp i\zeta_{(d)}^{2}(\mathbf{n}) = \sum_{jm} {}_{\pm 4}Y_{jm}(\hat{\mathbf{n}}) \Big[ k_{(E)jm}^{(d)} \pm ik_{(B)jm}^{(d)} \Big], (2.36)$$
$$\zeta_{(d)}^{3}(\mathbf{n}) = \sum Y_{im}(\hat{\mathbf{n}})k_{(d)}^{(d)} \quad . \qquad (2.37)$$

$$\zeta_{(d)}^{\circ}(\mathbf{n}) = \sum_{jm} Y_{jm}(\mathbf{\hat{n}}) k_{(V)jm}^{(-)}.$$
(2.37)

Then the phase velocity of the GWs can be rewritten as

$$v_{\pm} = 1 - \omega^{d-4} \Big[ \zeta^{0}_{(d)}(\mathbf{n}) \\ \mp \sqrt{(\zeta^{1}_{(d)}(\mathbf{n})^{2} + (\zeta^{2}_{(d)}(\mathbf{n})^{2} + (\zeta^{3}_{(d)}(\mathbf{n})^{2}) \Big]}, (2.38)$$

for a specific mass dimension d. The new effects arising from the Lorentz and diffeomorphism violations in the linearized gravity of SME are fully characterized by the coefficients,  $k_{(I)jm}^{(d)}$ ,  $k_{(V)jm}^{(d)}$ , and  $k_{(E)jm}^{(E)} \pm i k_{(B)jm}^{(d)}$ . These coefficients determine the speed of the GWs and all lead to frequency-dependent dispersions. Specifically, the coefficients  $k_{(V)jm}^{(d)}$  and  $k_{(I)jm}^{(E)} \pm i k_{(B)jm}^{(d)}$  lead to different velocities of two independent tensorial modes of GWs, a fast mode  $h_f$  and a slow mode  $h_s$ , as we mentioned in Eq. (2.28). Therefore, the arrival times of the fast and slow modes could be different. This is also called the velocity birefringence of the GWs. The coefficients  $k_{(I)jm}^{(d)}$ induce the non-birefringent dispersion of GWs. For this case (except d = 4 case), the two independent tensorial modes have the frequency-dependent velocity in the same form. For d = 4 case, the velocity is independent of the frequency of the GWs. In addition, all the coefficients,  $k_{(I)jm}^{(d)}$ ,  $k_{(V)jm}^{(d)}$ , and  $k_{(E)jm}^{(d)} \pm i k_{(B)jm}^{(d)}$  are also direction-dependent if  $j \neq 0$  and induce the anisotropic phase effects on the propagation of the GWs.

In summary, all the coefficients can provide frequency and direction-dependent phase modifications to the GWs. In the following, we are going to study the phase modifications due to these Lorentz- and diffeomorphismviolating coefficients in detail.

# III. PHASE MODIFICATIONS TO THE WAVEFORM OF GWS

Let us turn to derive the modified waveform of GW with Lorentz- and diffeomorphism-violating effects in the linearized gravity sector of the SME. For this purpose, we closely follow the derivation presented in [43, 54, 74]. Now consider a graviton emitted radially at  $r = r_e$  and received at r = 0, we have

$$\frac{dr}{dt} = -\frac{1}{a} \Big[ 1 - \zeta^0 \pm \sqrt{(\zeta^1)^2 + (\zeta^2)^2 + (\zeta^3)^2} \Big]. \quad (3.1)$$

Integrating this equation from the emission time (when  $r = r_e$ ) to arrival time (when r = 0), one obtains

$$r_{e} = \int_{t_{e}}^{t_{0}} \frac{dt}{a(t)} - \omega^{d-4} \Big[ \zeta_{(d)}^{0}(\mathbf{n}) \\ \mp \sqrt{(\zeta_{(d)}^{1}(\mathbf{n}))^{2} + (\zeta_{(d)}^{2}(\mathbf{n}))^{2} + (\zeta_{(d)}^{3}(\mathbf{n}))^{2}} \Big] \int_{t_{e}}^{t_{0}} \frac{dt}{a^{d-3}}$$
(3.2)

Considering gravitons emitted at two different times  $t_e$ and  $t'_e$ , with wave numbers k and k', and received at corresponding arrival times  $t_0$  and  $t'_0$  ( $r_e$  is the same for both). Assuming  $\Delta \equiv t_e - t'_e \leq a/\dot{a}$ , then, the difference in their arrival times is given by

$$\Delta t_{0} = (1+z)\Delta t_{e} + \left(\omega_{e}^{d-4} - \omega_{e}^{\prime d-4}\right) \left[\zeta_{(d)}^{0}(\mathbf{n}) + \sqrt{\left(\zeta_{(d)}^{1}(\mathbf{n})\right)^{2} + \left(\zeta_{(d)}^{2}(\mathbf{n})\right)^{2} + \left(\zeta_{(d)}^{3}(\mathbf{n})\right)^{2}}\right] \int_{t_{e}}^{t_{0}} \frac{dt}{a^{d-3}},$$
(3.3)

where  $z = 1/a(t_e) - 1$  is the cosmological redshift.

Let us focus on the GW signal generated by nonspinning, quasi-circular inspiral in the post-Newtonian approximation. Relative to the GW in GR, the Lorentzand diffeomorphism-violating effects modify the phase of GWs. In [2, 75], it was proved that the difference of arrival times in (3.3) induces the modification to the phase of GWs  $\Psi$  in the following form,

$$\Psi(f) = \Psi^{\text{GR}}(f) \mp \delta \Psi_1(f, \mathbf{n}) + \delta \Psi_2(f, \mathbf{n}), \qquad (3.4)$$

for  $d \neq 3$ , where  $\mp$  correspond to fast and slow modes, respectively, and

$$\delta \Psi_1(f, \mathbf{n}) = \frac{2^{d-3}}{d-3} \frac{u^{d-3}}{\mathcal{M}^{d-3}}$$

$$\sqrt{\left(\zeta_{(d)}^{1}(\mathbf{n})\right)^{2} + \left(\zeta_{(d)}^{2}(\mathbf{n})\right)^{2} + \left(\zeta_{(d)}^{3}(\mathbf{n})\right)^{2}} \int_{t_{e}}^{t_{0}} \frac{dt}{a^{d-3}},$$
(3.5)

$$\delta\Psi_2(f,\mathbf{n}) = \frac{2^{d-3}}{d-3} \frac{u^{d-3}}{\mathcal{M}^{d-3}} \zeta^0_{(d)}(\mathbf{n}) \int_{t_e}^{t_0} \frac{dt}{a^{d-3}}, \ (3.6)$$

where  $u = \pi \mathcal{M} f$  with  $f = \omega/2\pi$  being the frequency of the GWs,  $\mathcal{M} = (1+z)\mathcal{M}_c$  is the measured chirp mass, and  $\mathcal{M}_c \equiv (m_1m_2)^{3/5}/(m_1+m_2)^{1/5}$  is the chirp mass of the binary system with component masses  $m_1$  and  $m_2$ . When d = 3, the phase corrections  $\delta \Psi_1$  and  $\delta \Psi_2$  are given by

$$\delta \Psi_{1} = \sqrt{\left(\zeta_{(d)}^{1}(\mathbf{n})\right)^{2} + \left(\zeta_{(d)}^{2}(\mathbf{n})\right)^{2} + \left(\zeta_{(d)}^{3}(\mathbf{n})\right)^{2}} \\ \times \ln u \int_{t_{e}}^{t_{0}} dt, \qquad (3.7)$$

$$\delta \Psi_2 = \zeta_{(d)}^0(\mathbf{n}) \ln u \int_{t_e}^{t_0} dt.$$
 (3.8)

Then the fast and slow modes  $(h_f, h_s)$  with phase corrections can expressed as

$$h_f = h_f^{\rm GR} e^{(\delta \Psi_2 + \delta \Psi_1)}, \qquad (3.9)$$

$$h_s = h_f^{\rm GR} e^{(\delta \Psi_2 - \delta \Psi_1)}. \tag{3.10}$$

By using the relation in (2.28), one can transform the above results to the circular polarization modes

$$h_R = e^{i\delta\Psi_2} \Big[ (\cos\delta\Psi_1 - i\cos\vartheta\sin\delta\Psi_1) h_R^{GR} \\ - i\sin\vartheta e^{-i\varphi}\sin\delta\Psi_1 h_L^{GR} \Big], \qquad (3.11)$$

$$h_L = e^{i\delta\Psi_2} \Big[ (\cos\delta\Psi_1 + i\cos\vartheta\sin\delta\Psi_1) h_L^{GR} - i\sin\vartheta e^{i\varphi}\sin\delta\Psi_1 h_R^{GR} \Big].$$
(3.12)

The circular polarization modes  $h_R$  and  $h_L$  relate to the modes  $h_+$  and  $h_{\times}$  via

$$h_{+} = \frac{h_L + h_R}{\sqrt{2}},\tag{3.13}$$

$$h_{\times} = \frac{h_L - h_R}{\sqrt{2i}}.\tag{3.14}$$

The corresponding waveforms for  $h_+$  and  $h_{\times}$  are

$$h_{+} = e^{i\delta\Psi_{2}} \Big[ (\cos\delta\Psi_{1} - i\cos\varphi\sin\vartheta\sin\delta\Psi_{1})h_{+}^{GR} \\ - (\cos\vartheta + i\sin\vartheta\sin\varphi)\sin\delta\Psi_{1}h_{\times}^{GR} \Big], \quad (3.15)$$

$$h_{\times} = e^{i\delta\Psi_{2}} \Big[ (\cos\delta\Psi_{1} + i\cos\varphi\sin\vartheta\sin\delta\Psi_{1})h_{\times}^{GR} \\ + (\cos\vartheta - i\sin\vartheta\sin\varphi)\sin\delta\Psi_{1}h_{+}^{GR} \Big]. \quad (3.16)$$

Here we would like to consider several special limits of the above general waveform. Considering that the operators with the lowest mass dimension are expected to have the dominant Lorentz- and diffeomorphism-violating effects on the propagation of GWs, in the following we only discuss specific cases with relative lower mass dimensions, for example, d = 2, 3, 4, 5, 6. Note that ref. [76] explores isotropic Lorentz-violating effects on GWs, which correspond to operators with mass dimensions d = 7 and d = 8.

# A. Non-birefringent dispersion by the even d coefficients $k_{(I)im}^{(d)}$

The coefficients  $k_{(I)jm}^{(d)}$  for even  $d \ge 2$  induce the nonbirefringent dispersion of GWs,

$$\omega = \left(1 - \omega^{d-4} \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(I)jm}^{(d)}\right) |\mathbf{k}|. \qquad (3.17)$$

This leads to the non-birefringent phase velocity of GWs,

$$v = 1 - \omega^{d-4} \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(I)jm}^{(d)}.$$
 (3.18)

This phase velocity is direction-independent for j = 0 but anisotropic for  $j \neq 0$ . For the effect induced by  $k_{(I)jm}^{(d)}$ , it is obvious to see that the phase modification  $\delta \Psi_1 = 0$ . In the following, we consider the phase corrections for mass dimensions d = 2, d = 4, and d = 6, respectively.

1. 
$$d=2$$

For mass dimension d = 2 case, since  $0 \le j \le d - 2$ , j and m have to take j = 0 = m. For this case,  $k_{(I)jm}^{(2)}$ only has one component  $k_{(I)00}^{(2)}$ , which is obvious direction independent. The phase correction for this case is given by

$$\delta\Psi_2 = -\frac{(\pi f)^{-1}}{4\sqrt{\pi}} k^{(2)}_{(I)00} \int_{t_e}^{t_0} a dt.$$
(3.19)

As we mentioned, the d = 2 case can only be induced by the diffeomorphism violations of the linearized gravity described by the Lagrangian (2.1).

2. d=4

For the d = 4 case, the phase velocity is independent of the frequency of the GWs, so they do not give any observable dephasing effects and modify the speed of GWs in a frequency-independent way. This effect can be constrained by comparison with the arrival time of the photons from the associated electromagnetic counterpart. For the binary neutron star merger GW170817 and its associated electromagnetic counterpart GRB170817A [77], the almost coincident observation of the electromagnetic wave and the GW place an exquisite bound on

$$-7 \times 10^{-17} < \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(I)jm}^{(4)} < 3 \times 10^{-15}.$$
 (3.20)

Since this case does not lead to any dephasing effects, we are not going to include this case in our later analysis with GW signals in GWTC-3. In addition, it is shown in [34] that with the effects of the anisotropic coefficients  $k_{(I)jm}^{(4)}$ , the extra polarizations of GWs can be directly generated by the two tensorial modes under certain conditions.

3. d=6

For the mass dimension d = 6 case, the index j can take 0, 1, 2, 3, 4, and the index m runs from -j to j. Note that each of  $k_{(I)jm}^{(6)}$  are complex function which satisfies  $k_{(I)jm}^{(6)*} = (-1)^m k_{(I)j-m}^{(6)}$ . Thus the number of independent components for coefficients  $k_{(I)jm}^{(6)}$  are  $(d-1)^2 = 25$ . The phase correction in the modified waveform for this case is given by

$$\delta\Psi_2 = \frac{8}{3} (\pi f)^3 \left( \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(I)jm}^{(6)} \right) \int_{t_e}^{t_0} \frac{dt}{a^3}.$$
 (3.21)

The coefficients  $k_{(I)jm}^{(6)}$  can be constrained by comparing the modified waveform with the GW strain data from the GW detectors, see ref. [70] for a review. The analysis with the isotropic effect of Lorentz violation which corresponds to j = 0 in the linearized gravity has been carried out through full Bayesian parameter estimations on the GW events observed by the LIGO/Virgo/KAGRA detectors in a series of papers [37, 57, 67–69], while the anisotropic case has been considered in [38].

# B. Birefringent dispersion by the odd d coefficients $k_{(V)jm}^{(d)}$

The coefficients  $k_{(V)jm}^{(d)}$  for odd  $d \geq 3$  produces frequency-dependent dispersion and birefringence effects in GWs, i.e.,

$$\omega = \left(1 \pm \omega^{d-4} \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(V)jm}^{(d)}\right) |\mathbf{k}|. \quad (3.22)$$

Note that here  $\mp$  corresponds to the fast and slow modes respectively. This leads to the birefringent phase velocity of GWs,

$$v_{\pm} = 1 \pm \omega^{d-4} \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(V)jm}^{(d)}.$$
 (3.23)

Similarly, this velocities are direction-independent for j = 0 but anisotropic for  $j \neq 0$ . In this case, the fast and slow modes are circularly polarized, and we have  $\vartheta = 0, \pi$ . For the effects induced by  $k_{(V)jm}^{(d)}$ , it is obvious that the phase corrections  $\delta \Psi_2 = 0$ .

1. d=3

For mass dimension d = 3 case, the index j can take 0 and 1, and index m runs from -j to j. Considering  $k_{(V)jm}^{(5)*} = (-1)^m k_{(V)j-m}^{(5)}$ , it is known that there are only 4 independent components for the coefficients  $k_{(V)jm}^{(3)}$ , they are  $k_{(V)00}^{(3)}$ ,  $k_{(V)10}^{(3)}$ ,  $\text{Rek}_{(V)11}^{(3)}$ , and  $\text{Im}k_{(V)11}^{(3)}$ , in which  $k_{(V)11}^{(3)}$  is a complex function so it contains two independent components. The phase correction for this case is given by

$$\delta \Psi_1 = \ln u \left( \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(V)jm}^{(3)} \right) \int_{t_e}^{t_0} dt. \quad (3.24)$$

Similarly, one can constrain the coefficients  $k_{(V)jm}^{(3)}$  by comparing the modified waveform with the GW strain data from the GW detectors. The analysis with the isotropic effect for d = 3 case (which corresponds to j = 0case) has been performed in refs. [37, 57, 78].

$$2. \quad d=5$$

For mass dimension d = 5 case, the index j can take 0, 1, 2, 3, and index m runs from -j to j. Here the coefficients  $k_{(V)jm}^{(5)}$  have  $(d-1)^2 = 16$  components in total. The phase corrections for this case are given by

$$\delta \Psi_1 = 2(\pi f)^2 \left( \sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(V)jm}^{(3)} \right) \int_{t_e}^{t_0} a^{-2} dt.$$
(3.25)

The constraints on the isotropic effect for d = 5 case have been performed in a series of papers, see refs. [37, 53, 55, 57, 60], while the the anisotropic effects has been constrained in [39, 40, 42, 79].

# C. Birefringent dispersion by the even d coefficients $k_{(E)jm}^{(d)}$ and $k_{(B)jm}^{(d)}$

The coefficients  $k_{(E)jm}^{(d)}$  and  $k_{(B)jm}^{(d)}$  for even  $d \ge 6$  produces frequency-dependent dispersion and birefringence effects in GWs, i.e.,

$$\omega = \left(1 \pm \frac{1}{2}\omega^{d-4}\sqrt{(\zeta^{1}_{(d)}(\mathbf{n}))^{2} + (\zeta^{2}_{(d)}(\mathbf{n}))^{2}}\right)|\mathbf{k}|. (3.26)$$

For this case, we have  $\vartheta = \pi/2$  and  $\delta \Psi_2 = 0$ .

For mass dimension d = 6, the index j = 4 and index m runs from 0 to 4. In this case, the coefficients  $k_{(E)jm}^{(d)}$  and  $k_{(B)jm}^{(d)}$  have 18 components in total. The phase corrections for this case is given by

$$\delta \Psi_1 = \frac{8}{3} (\pi f)^3 \sqrt{(\zeta_{(6)}^1(\mathbf{n}))^2 + (\zeta_{(6)}^2(\mathbf{n}))^2} \int_{t_e}^{t_0} a^{-3} dt.$$
(3.27)

The constraints on the coefficients  $k_{(E)jm}^{(d)}$  and  $k_{(B)jm}^{(d)}$  for d = 6 case have been performed in [40, 42, 79].

# IV. BAYESIAN INFERENCE AND PARAMETER ESTIMATION

In this section, we present a brief introduction to the Bayesian inference used to constrain the coefficients of Lorentz and diffeomorphism violation in the linearized gravity in the framework of the SME. Bayesian inference plays a pivotal role in modern astronomy, enabling the extraction of physical parameters from observational data. Given GW data  $d_i$ , we compare it with the predicted GW strain incorporating Lorentz- and diffeomorphism-violating effects to infer the posterior distribution of parameters  $\vec{\theta}$  that characterize the waveform model. According to Bayes' theorem, the posterior distribution is expressed as:

$$P(\vec{\theta}|d,H) = \frac{P(d|\vec{\theta},H)P(\vec{\theta}|H)}{P(d|H)},$$
(4.1)

where  $P(\vec{\theta}|d, H)$  represents the posterior probability distributions of the model parameters  $\vec{\theta}$ , and H is the waveform model.  $P(\vec{\theta}|H)$  is the prior distribution for parameters  $\vec{\theta}$ ,  $P(d|\vec{\theta}, H)$  is the likelihood function for a given set of model parameters and P(d|H) is the normalization factor, commonly referred as the "evidence":

$$P(d|H) \equiv \int d\vec{\theta} P(d|\vec{\theta}, H) P(\vec{\theta}|H).$$
(4.2)

In most cases, GW signals are weak and embedded within noise, making matched filtering a crucial method for signal extraction. Assuming Gaussian and stationary noise [80–82], the likelihood function for matched filtering is given by:

$$P(\boldsymbol{d}|\boldsymbol{\theta}, H) \propto \prod_{i=1}^{n} e^{-\frac{1}{2} \langle \boldsymbol{d}_{i} - \boldsymbol{h}(\boldsymbol{\theta}) | \boldsymbol{d}_{i} - \boldsymbol{h}(\boldsymbol{\theta}) \rangle}, \qquad (4.3)$$

where  $h(\theta)$  is the GW strain predicted by the waveform model H, and i indexes the individual GW detectors. The noise-weighted inner product  $\langle A|B \rangle$  is defined as:

$$\langle A|B\rangle = 4 \operatorname{Re}\left[\int_0^\infty \frac{A(f)B(f)^*}{S(f)}df\right],$$
 (4.4)

where \* denotes the complex conjugate, and S(f) is the power spectral density (PSD) of the detector. To ensure stable and reliable parameter estimation, we use the PSD data provided in the LVK posterior sample, which is more robust compared to PSDs derived from strain data using Welch averaging [48, 83, 84].

We consider the cases of Lorentz- and diffeomorphismviolating waveforms in (3.15) and (3.16) with different mass dimension d separately. It is mentioned in the previous section that the cases with mass dimensions d = 5and d = 6 have been explored in refs. [38–40, 42, 79], which will not be considered here. For this reason, we explore the cases with d = 2 and d = 3 cases in this paper. To perform the parameter estimation on the modified waveforms (3.15) and (3.16) with the Lorentz- and diffeomorphism-violating effects, we employ the python package BILBY [85, 86]. We perform Bayesian inference on GW data from the 88 compact binary coalescence events in GWTC-3, which include binary neutron stars like GW170817, neutron star-black hole binaries, and binary black holes. Two events, GW200308\_173609 and GW200322\_091133, are excluded in our analysis due to the possible uncertainties of their inferred source properties [46]. It is also shown in [87] from a new analysis that these two events could be generated by Gaussian noise fluctuations. We use the IMRPhenomXPHM template [88–90] for the GR waveform  $h_{+,\times}^{\text{GR}}(f)$  except for the binary neutron star event GW170817, and use IMRPhenomPv2\_NRTidal for GW170817.

Considering that the spherical expansion coefficient formulae in Eqs. (2.38, 2.36, 2.37) provide a general solution for different events within the same coordinate system, we can combine the posterior distributions of individual events as:

$$P(\vec{\theta}|\{d_i\}, H) \propto \prod_{i=1}^{N} P(\vec{\theta}|d_i, H), \qquad (4.5)$$

where  $d_i$  represents the data from the *i*-th GW event, and N is the total number of selected GW events.

### V. RESULTS

In this section, we present the results of the constraints on the Lorentz- and diffeomorphism-violating coefficients for mass diemsnion d = 2 and d = 3 cases. In the following, we present the results for d = 2 and d = 3 separately.

**A.** d = 2

For the mass dimension d = 2 case, the phase corrections (with  $\delta \Psi_1 = 0$ ) in the modified waveform in (3.15) and (3.16) is expressed

$$\delta \Psi_2 = A_{\bar{\mu}} (\pi f)^{-1}, \tag{5.1}$$

where

$$A_{\bar{\mu}} = -\frac{1}{4\sqrt{\pi}} k_{(I)00}^{(2)} \int_0^z \frac{(1+z')^{-2}}{H_0 \sqrt{\Omega_m (1+z')^3 + \Omega_\Lambda}} dz'.$$
(5.2)

Here we adopt a Planck cosmology with  $\Omega_m = 0.315$ ,  $\Omega_{\Lambda} = 0.685$ , and  $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$  [91]. The parameter  $A_{\bar{\mu}}$  is the parameter we sampled in the Bayesian inference along with other GR parameters. We use the uniform prior for parameter  $A_{\bar{\mu}}$  in our analysis. Then from the marginal posterior distributions of  $A_{\bar{\mu}}$  and the redshift z of the analyzed GW events, one can obtain posterior distributions of  $k_{(I)00}^{(2)}$ . In Fig. 1 we display the marginalized posterior distributions of  $k_{(I)00}^{(2)}$  from selected GW events in the GWTC-3. For most GW events we analyze in each test, we do not find any significant signatures of Lorentz and diffeomorphism violation due to the coefficient  $k_{(I)00}^{(2)}$ . A few events that suggest nonzero values for the non-GR coefficients  $A_{\bar{\mu}}$  are excluded from our analysis due to their contradiction with GR. The posterior posterior distributions for  $k_{(I)00}^{(2)}$  from the excluded GW events are presented in Fig. 2. It is mentioned in [60, 66] that these results may arise from limitations in current waveform approximants, such as systematic errors during the merger phase or unaccounted physical effects like eccentricity. Consequently, we have excluded these events from our analysis. Table II presents the list of excluded events along with the estimated constraints on  $k_{(I)00}^{(2)}$  for each excluded event.

In addition, we consider the coefficient  $k_{(I)00}^{(2)}$  as a universal parameter for all GW events, then we obtain its combined constraint by multiplying the posterior distributions of the individual events together, which gives

$$-0.5 \times 10^{-63} \text{ GeV}^2 < k_{(I)00}^{(2)} < 1.3 \times 10^{-63} \text{ GeV}^2$$
(5.3)

at 90% confidence level. The lower and upper limits of  $k_{(I)00}^{(2)}$  are represented by the vertical dash line in Fig. 1.

**B.** d = 3

For the mass dimension d = 3, the phase correction (with  $\delta \Psi_2 = 0$ ) in the modified waveform in (3.15) and (3.16) takes the form

$$\delta \Psi_1 = A_\mu \ln u, \tag{5.4}$$

with

$$A_{\mu} = \left(\sum_{jm} Y_{jm}(\hat{\mathbf{n}}) k_{(V)jm}^{(3)}\right) \\ \times \int_{0}^{z} \frac{(1+z')^{-1}}{H_{0}\sqrt{\Omega_{m}(1+z')^{3} + \Omega_{\Lambda}}} dz'.$$
(5.5)

TABLE II. The list of the excluded events in the analysis for d = 2 case and their constraints on  $k_{(I)00}^{(2)}$  at 90% confidence interval. It is obvious that these constraints favor non-zero values of the non-GR coefficient  $k_{(I)00}^{(2)}$ .

Coefficient	Events	Constraint $(10^{-62} \text{ Gev}^2)$
	GW190408_181802	(15.2, 20.9)
	GW190503_185404	(3.5, 12.6)
	$GW190519_{153544}$	(1.5, 3.3)
	$GW190630_185205$	(5.9, 8.3)
$k_{(I)00}^{(2)}$	$GW190701_{-203306}$	(11.5, 19.1)
	$GW200112_{-}155838$	(8.2, 12.9)
	GW200216_220804	(2.7, 4.9)
	GW200225_060421	(4.2, 23.5)
	$GW200311\_115853$	(8.1, 11.6)

Here  $A_{\mu}$  is the parameter we sampled in the Bayesian inference along with other GR parameters. We use the uniform prior for parameter  $A_{\mu}$  in our analysis. As we mentioned, for mass dimension d = 3, the coefficients  $k_{(V)jm}^{(3)}$  have four independent components,  $k_{(V)00}^{(3)}$ ,  $k_{(V)10}^{(3)}$ ,  $\operatorname{Rek}_{(V)11}^{(3)}$ , and  $\operatorname{Imk}_{(V)11}^{(3)}$ . These components are entirely tangled together. Here we adopt an approach by using the "maximum-reach" method, with which one can constrain each of these components separately [40, 42, 79]. This implies that when one considers one of these components, the others are set to zero.

Then the posterior samples of each component of  $k_{(V)jm}^{(3)}$  can be calculated from the marginal posterior distributions of  $A_{\mu}$ , right ascension (ra), declination (dec), and redshift z of the analyzed GW events, via Eqs. (5.5). Fig. 3 presents the marginalized posterior distributions of  $k_{(V)00}^{(3)}, k_{(V)10}^{(3)}, \text{Re}k_{(V)11}^{(3)}$ , and  $\text{Im}k_{(V)11}^{(3)}$  from selected GW events in the GWTC-3. The lower and upper limits of each coefficient  $k_{(V)00}^{(3)}$ ,  $k_{(V)10}^{(3)}$ ,  $\operatorname{Re}k_{(V)11}^{(3)}$ , and  $\operatorname{Im}k_{(V)11}^{(3)}$ .  $k_{(I)jm}^{(6)}$  are represented by the vertical dash line in each figure of Fig. 3. For most GW events we analyze in each test, we do not find any significant signatures of Lorentz and diffeomorphism violation due to the coefficient  $k_{(V)ij}^{(3)}$ . In addition, for each coefficient of  $k_{(V)00}^{(3)}$ ,  $k_{(V)10}^{(3)}, \operatorname{Re}k_{(V)11}^{(3)}, \operatorname{and} \operatorname{Im}k_{(V)11}^{(3)},$ we consider each of them as a universal parameter for all GW events, and then we obtain their combined constraints separately by multiplying the posterior distributions of the individual events together. Table. III summarizes the 90% confidence interval of each coefficients  $k_{(V)00}^{(3)}$ ,  $k_{(V)10}^{(3)}$ ,  $\operatorname{Re}k_{(V)11}^{(3)}$ , and  $\operatorname{Im}_{(V)11}^{(3)}$ . From both the Fig. 3 and Table. III, it is obvious that the posterior samples and the 90% confidence interval of each coefficient  $k^{(3)}_{(V)ij}$  are all consistent with zero, which indicates there are no any signatures of the Lorentz and diffeomorphism violations arising in the linearized gravity of SME has been found in the GW signals.



FIG. 1. The posterior distributions for  $k_{(I)00}^{(2)}$  from selected GW events in the LIGO-Virgo-KAGRA catalog GWTC-3. The legend indicates the events that give the six tightest constraints. The vertical dash line denotes the 90% interval for  $k_{(I)00}^{(2)}$  from combined results. Note that we have excluded a few events in our analysis (as shown in the list presented in Table. II).



FIG. 2. The posterior distributions for  $k_{(I)00}^{(2)}$  from 9 excluded GW events (the excluded events are listed in Table. II).

### VI. CONCLUSION

The detection of GW signals by the LIGO-Virgo-KAGRA Collaboration marked the beginning of a new era in testing gravity in the strong-field regime. In this study, we investigate the effects of Lorentz- and diffeomorphism-violating effects in the linearized gravity on GW propagation within the framework of SME. Using an approach similar to that in the photon sector of the SME [71], we derive a modified dispersion relation for GWs with the Lorentz- and -violating effects, which lead to anisotropy, birefringence, and dispersion effects in the propagation of gravitational waves. With these modified dispersion relations, we then calculate the dephasing effects due to the Lorentz and diffeomorphism violations in the waveforms of gravitational waves produced by the

TABLE III. 90% confidence interval of each component of the Lorentz- and diffeomorphism-violating coefficients  $k_{(V)jm}^{(3)}$  from 90 GW events in the GWTC-3 catalog.

j	m	Coefficient	Constraint $(10^{-41} \text{ Gev}^{-1})$
0	0	$k^{(3)}_{(V)00}$	(-0.65, 1.13)
1	0	$k_{(V)10}^{(3)}$	(-0.3, 0.9)
	1	Re $k_{(V)11}^{(3)}$	(-4.2, 3.2)
		Im $k_{(V)11}^{(3)}$	(-1.2, 4.2)

coalescence of compact binaries.

With the distorted waveforms, one can test the Lorentz and diffeomorphism invariance of gravity by comparing the modified waveform with the GW strain data from the GW detectors. Several previous works have been carried out for testing Lorentz symmetry in Lorentz-violating linearized gravity for mass dimension  $d \ge 4$  cases. This study explores the effects of diffeomorphism violations induced by Lorentz-violating terms with mass dimensions d = 2 and d = 3 respectively. Using the SME framework for Lorentz-violating linearized gravity, we perform Bayesian inference with modified waveforms incorporating diffeomorphism-violating effects on GW events from the GWTC-3 catalog. We find no significant evidence of diffeomorphism violations in most GW data and provide 90% confidence intervals for each diffeomorphismviolating coefficients.

Our results, illustrated in Fig. 1 for d = 2 case, and Fig. 3 and Table. III for d = 3 case, show no evidence of Lorentz and diffeomorphism violations. Accordingly, we report constraints on the coefficients  $k_{(I)00}^{(2)}$  describing anisotropic non-birefringent effects for d = 2 and coeffcients  $k_{(V)00}^{(3)}$ ,  $k_{(V)10}^{(3)}$ ,  $\operatorname{Rek}_{(V)11}^{(3)}$ , and  $\operatorname{Imk}_{(V)11}^{(3)}$  describing birefringent anisotropic effects for d = 3. Nevertheless, the medians of all components remain near zero and thus are consistent with the results of GR prediction. Looking to the future, the next generation of GW detectors, with enhanced sensitivity and the ability to observe lighter and more distant binary black hole (BBH) and binary neutron star (BNS) systems, is anticipated to tighten further constraints on Lorentz and diffeomorphism violations induced dispersions in GW propagation.

## ACKNOWLEDGEMENTS

We appreciate the helpful discussions with Bo-Yang Zhang and Yuan-Zhu Wang. This work is supported by the National Key Research and Development Program of China under Grant No. 2020YFC2201503, the National Natural Science Foundation of China under Grants No. 12275238 and No. 11675143, the Zhejiang Provincial Natural Science Foundation of China under Grants No. LR21A050001 and No. LY20A050002, and the Fundamental Research Funds for the Provincial Universities



FIG. 3. The posterior distributions for  $k_{(V)00}^{(3)}$ ,  $k_{(V)10}^{(3)}$ ,  $\operatorname{Rek}_{(V)11}^{(3)}$ , and  $\operatorname{Im}k_{(V)11}^{(3)}$  from selected GW events in the LIGO-Virgo-KAGRA catalog GWTC-3. The legend indicates the events that give the six tightest constraints. The vertical dash line denotes the 90% interval for the coefficients  $k_{(V)00}^{(3)}$ ,  $k_{(V)10}^{(3)}$ ,  $\operatorname{Rek}_{(V)11}^{(3)}$  from combined results.

of Zhejiang in China under Grant No. RF-A2019015. W.Z. is supported by the National Natural Science Foundation of China (Grants No. 12325301 and 12273035) and the National Key Research and Development Program of China (Grant No. 2022YFC2204602 and 2021YFC2203102), the Fundamental Research Funds for the Central Universities.

This research has made use of data or software obtained from the Gravitational Wave Open Science Center (gwosc.org), a service of the LIGO Scientific Collaboration, the Virgo Collaboration, and KAGRA. This material is based upon work supported by NSF's LIGO Laboratory which is a major facility fully funded by the National Science Foundation, as well as the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and construction and operation of the GEO600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. Virgo is funded, through the European Gravitational Observatory (EGO), by the French Centre National de Recherche Scientifique (CNRS), the Italian Istituto Nazionale di Fisica Nucleare (INFN) and the Dutch Nikhef, with contributions by institutions from Belgium, Germany, Greece, Hungary, Ireland, Japan, Monaco, Poland, Portugal, Spain. KAGRA is supported by the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan Society for the Promotion of Science (JSPS) in Japan; National Research Foundation (NRF) and Ministry of Science and ICT (MSIT) in Korea; Academia Sinica (AS) and National Science and Technology Council (NSTC) in Taiwan.

The data analyses and results visualization in this work made use of BILBY [85, 86], dynesty [92], LALSuite [93], Numpy [94, 95], Scipy [96], and matplotlib [97].

 Emanuele Berti *et al.*, "Testing General Relativity with Present and Future Astrophysical Observations," Class. Quant. Grav. **32**, 243001 (2015), arXiv:1501.07274 [gr $\mathbf{qc}].$ 

<sup>[2]</sup> Clifford M. Will, "Bounding the mass of the graviton using gravitational wave observations of inspiralling com-

pact binaries," Phys. Rev. D 57, 2061–2068 (1998), arXiv:gr-qc/9709011.

- [3] C. D. Hoyle, U. Schmidt, Blayne R. Heckel, E. G. Adelberger, J. H. Gundlach, D. J. Kapner, and H. E. Swanson, "Submillimeter tests of the gravitational inverse square law: a search for 'large' extra dimensions," Phys. Rev. Lett. 86, 1418–1421 (2001), arXiv:hep-ph/0011014.
- [4] Bhuvnesh Jain and Justin Khoury, "Cosmological Tests of Gravity," Annals Phys. **325**, 1479–1516 (2010), arXiv:1004.3294 [astro-ph.CO].
- [5] Kazuya Koyama, "Cosmological Tests of Modified Gravity," Rept. Prog. Phys. **79**, 046902 (2016), arXiv:1504.04623 [astro-ph.CO].
- [6] Timothy Clifton, Pedro G. Ferreira, Antonio Padilla, and Constantinos Skordis, "Modified Gravity and Cosmology," Phys. Rept. 513, 1–189 (2012), arXiv:1106.2476 [astro-ph.CO].
- [7] Ingrid H. Stairs, "Testing general relativity with pulsar timing," Living Rev. Rel. 6, 5 (2003), arXiv:astroph/0307536.
- [8] R. N. Manchester, "Pulsars and Gravity," Int. J. Mod. Phys. D 24, 1530018 (2015), arXiv:1502.05474 [gr-qc].
- [9] Norbert Wex, "Testing Relativistic Gravity with Radio Pulsars," (2014), arXiv:1402.5594 [gr-qc].
- [10] Michael Kramer, "Pulsars as probes of gravity and fundamental physics," Int. J. Mod. Phys. D 25, 1630029 (2016), arXiv:1606.03843 [astro-ph.HE].
- [11] V. Alan Kostelecky and Stuart Samuel, "Spontaneous Breaking of Lorentz Symmetry in String Theory," Phys. Rev. D 39, 683 (1989).
- [12] V. Alan Kostelecky and Robertus Potting, "CPT and strings," Nucl. Phys. B 359, 545–570 (1991).
- [13] Rodolfo Gambini and Jorge Pullin, "Nonstandard optics from quantum space-time," Phys. Rev. D 59, 124021 (1999), arXiv:gr-qc/9809038.
- [14] C. P. Burgess, James M. Cline, E. Filotas, J. Matias, and G. D. Moore, "Loop generated bounds on changes to the graviton dispersion relation," JHEP 03, 043 (2002), arXiv:hep-ph/0201082.
- [15] Don Colladay and V. Alan Kostelecky, "CPT violation and the standard model," Phys. Rev. D 55, 6760–6774 (1997), arXiv:hep-ph/9703464.
- [16] V. Alan Kostelecky, "Gravity, Lorentz violation, and the standard model," Phys. Rev. D 69, 105009 (2004), arXiv:hep-th/0312310.
- [17] James B. R. Battat, John F. Chandler, and Christopher W. Stubbs, "Testing for Lorentz Violation: Constraints on Standard-Model Extension Parameters via Lunar Laser Ranging," Phys. Rev. Lett. 99, 241103 (2007), arXiv:0710.0702 [gr-qc].
- [18] Adrien Bourgoin, Christophe Le Poncin-Lafitte, Aurélien Hees, Sébastien Bouquillon, Gérard Francou, and Marie-Christine Angonin, "Lorentz Symmetry Violations from Matter-Gravity Couplings with Lunar Laser Ranging," Phys. Rev. Lett. **119**, 201102 (2017), arXiv:1706.06294 [gr-qc].
- [19] Holger Muller, Sheng-wey Chiow, Sven Herrmann, Steven Chu, and Keng-Yeow Chung, "Atom Interferometry tests of the isotropy of post-Newtonian gravity," Phys. Rev. Lett. **100**, 031101 (2008), arXiv:0710.3768 [gr-qc].
- [20] V. Alan Kostelecký and Jay D. Tasson, "Constraints on Lorentz violation from gravitational Čerenkov radiation," Phys. Lett. B 749, 551–559 (2015),

arXiv:1508.07007 [gr-qc].

- [21] Lijing Shao, "Tests of local Lorentz invariance violation of gravity in the standard model extension with pulsars," Phys. Rev. Lett. **112**, 111103 (2014), arXiv:1402.6452
   [gr-qc].
- [22] Lijing Shao, "New pulsar limit on local Lorentz invariance violation of gravity in the standard-model extension," Phys. Rev. D 90, 122009 (2014), arXiv:1412.2320 [gr-qc].
- [23] Ross J. Jennings, Jay D. Tasson, and Shun Yang, "Matter-Sector Lorentz Violation in Binary Pulsars," Phys. Rev. D 92, 125028 (2015), arXiv:1510.03798 [gr-qc].
- [24] Lijing Shao and Quentin G. Bailey, "Testing velocitydependent CPT-violating gravitational forces with radio pulsars," Phys. Rev. D 98, 084049 (2018), arXiv:1810.06332 [gr-qc].
- [25] Lijing Shao and Quentin G. Bailey, "Testing the Gravitational Weak Equivalence Principle in the Standard-Model Extension with Binary Pulsars," Phys. Rev. D 99, 084017 (2019), arXiv:1903.11760 [gr-qc].
- [26] Lijing Shao, "Lorentz-Violating Matter-Gravity Couplings in Small-Eccentricity Binary Pulsars," Symmetry 11, 1098 (2019), arXiv:1908.10019 [hep-ph].
- [27] Aurélien Hees, Quentin G. Bailey, Christophe Le Poncin-Lafitte, Adrien Bourgoin, Attilio Rivoldini, Brahim Lamine, Frédéric Meynadier, Christine Guerlin, and Peter Wolf, "Testing Lorentz symmetry with planetary orbital dynamics," Phys. Rev. D 92, 064049 (2015), arXiv:1508.03478 [gr-qc].
- [28] Natasha A. Flowers, Casey Goodge, and Jay D. Tasson, "Superconducting-Gravimeter Tests of Local Lorentz Invariance," Phys. Rev. Lett. **119**, 201101 (2017), arXiv:1612.08495 [gr-qc].
- [29] Alan V. Kostelecky and Jay D. Tasson, "Matter-gravity couplings and Lorentz violation," Phys. Rev. D 83, 016013 (2011), arXiv:1006.4106 [gr-qc].
- [30] Quentin G. Bailey and V. Alan Kostelecky, "Signals for Lorentz violation in post-Newtonian gravity," Phys. Rev. D 74, 045001 (2006), arXiv:gr-qc/0603030.
- [31] A. F. Ferrari, M. Gomes, J. R. Nascimento, E. Passos, A. Yu. Petrov, and A. J. da Silva, "Lorentz violation in the linearized gravity," Phys. Lett. B 652, 174–180 (2007), arXiv:hep-th/0609222.
- [32] V. Alan Kostelecký and Matthew Mewes, "Lorentz and Diffeomorphism Violations in Linearized Gravity," Phys. Lett. B 779, 136–142 (2018), arXiv:1712.10268 [gr-qc].
- [33] V. Alan Kostelecký and Matthew Mewes, "Testing local Lorentz invariance with gravitational waves," Phys. Lett. B 757, 510–514 (2016), arXiv:1602.04782 [gr-qc].
- [34] Shaoqi Hou, Xi-Long Fan, Tao Zhu, and Zong-Hong Zhu, "Nontensorial gravitational wave polarizations from the tensorial degrees of freedom: Linearized Lorentzviolating theory of gravity," Phys. Rev. D 109, 084011 (2024), arXiv:2401.03474 [gr-qc].
- [35] Yiming Dong, Ziming Wang, and Lijing Shao, "New limits on the local Lorentz invariance violation of gravity in the standard model extension with pulsars," Phys. Rev. D 109, 084024 (2024), arXiv:2311.11038 [astro-ph.HE].
- [36] Kellie Ault, Tests of General Relativity through Searches for Lorentz and CPT Symmetry Breaking, Ph.D. thesis, Embry-Riddle Aeronautical U. (2023).
- [37] Tao Zhu, Wen Zhao, Jian-Ming Yan, Yuan-Zhu Wang, Cheng Gong, and Anzhong Wang, "Constraints on

parity and Lorentz violations in gravity from GWTC-3 through a parametrization of modified gravitational wave propagations," Phys. Rev. D **110**, 064044 (2024), arXiv:2304.09025 [gr-qc].

- [38] Cheng Gong, Tao Zhu, Rui Niu, Qiang Wu, Jing-Lei Cui, Xin Zhang, Wen Zhao, and Anzhong Wang, "Gravitational wave constraints on nonbirefringent dispersions of gravitational waves due to Lorentz violations with GWTC-3 events," Phys. Rev. D 107, 124015 (2023), arXiv:2302.05077 [gr-qc].
- [39] Leïla Haegel, Kellie O'Neal-Ault, Quentin G. Bailey, Jay D. Tasson, Malachy Bloom, and Lijing Shao, "Search for anisotropic, birefringent spacetime-symmetry breaking in gravitational wave propagation from GWTC-3," Phys. Rev. D 107, 064031 (2023), arXiv:2210.04481 [grqc].
- [40] Rui Niu, Tao Zhu, and Wen Zhao, "Testing Lorentz invariance of gravity in the Standard-Model Extension with GWTC-3," JCAP **12**, 011 (2022), arXiv:2202.05092 [gr-qc].
- [41] Kellie O'Neal-Ault, Quentin G. Bailey, Tyann Dumerchat, Leila Haegel, and Jay Tasson, "Analysis of Birefringence and Dispersion Effects from Spacetime-Symmetry Breaking in Gravitational Waves," Universe 7, 380 (2021), arXiv:2108.06298 [gr-qc].
- [42] Lijing Shao, "Combined search for anisotropic birefringence in the gravitational-wave transient catalog GWTC-1," Phys. Rev. D 101, 104019 (2020), arXiv:2002.01185 [hep-ph].
- [43] Matthew Mewes, "Signals for Lorentz violation in gravitational waves," Phys. Rev. D 99, 104062 (2019), arXiv:1905.00409 [gr-qc].
- [44] Tian-Chen Li, Tao Zhu, Wen Zhao, and Anzhong Wang, "Power spectra and circular polarization of primordial gravitational waves with parity and Lorentz violations," JCAP 07, 005 (2024), arXiv:2403.05841 [gr-qc].
- [45] B. P. Abbott *et al.* (LIGO Scientific, Virgo), "Observation of Gravitational Waves from a Binary Black Hole Merger," Phys. Rev. Lett. **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
- [46] R. Abbott et al. (KAGRA, VIRGO, LIGO Scientific), "GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo during the Second Part of the Third Observing Run," Phys. Rev. X 13, 041039 (2023), arXiv:2111.03606 [gr-qc].
- [47] R. Abbott *et al.* (LIGO Scientific, Virgo), "GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run," Phys. Rev. X **11**, 021053 (2021), arXiv:2010.14527 [gr-qc].
- [48] B. P. Abbott *et al.* (LIGO Scientific, Virgo), "GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs," Phys. Rev. X 9, 031040 (2019), arXiv:1811.12907 [astro-ph.HE].
- [49] B. P. Abbott *et al.* (LIGO Scientific, Virgo), "GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral," Phys. Rev. Lett. **119**, 161101 (2017), arXiv:1710.05832 [gr-qc].
- [50] B. P. Abbott *et al.* (LIGO Scientific, Virgo), "GW150914: The Advanced LIGO Detectors in the Era of First Discoveries," Phys. Rev. Lett. **116**, 131103 (2016), arXiv:1602.03838 [gr-qc].
- [51] B. P. Abbott et al. (LIGO Scientific, Virgo), "Properties

of the Binary Black Hole Merger GW150914," Phys. Rev. Lett. **116**, 241102 (2016), arXiv:1602.03840 [gr-qc].

- [52] B. P. Abbott *et al.* (LIGO Scientific, Virgo), "GW150914: First results from the search for binary black hole coalescence with Advanced LIGO," Phys. Rev. D 93, 122003 (2016), arXiv:1602.03839 [gr-qc].
- [53] Yi-Fan Wang, Rui Niu, Tao Zhu, and Wen Zhao, "Gravitational Wave Implications for the Parity Symmetry of Gravity in the High Energy Region," Astrophys. J. 908, 58 (2021), arXiv:2002.05668 [gr-qc].
- [54] Wen Zhao, Tao Zhu, Jin Qiao, and Anzhong Wang, "Waveform of gravitational waves in the general parityviolating gravities," Phys. Rev. D 101, 024002 (2020), arXiv:1909.10887 [gr-qc].
- [55] Zhi-Chao Zhao, Zhoujian Cao, and Sai Wang, "Search for the Birefringence of Gravitational Waves with the Third Observing Run of Advanced LIGO-Virgo," Astrophys. J. 930, 139 (2022), arXiv:2201.02813 [gr-qc].
- [56] Sai Wang and Zhi-Chao Zhao, "Tests of CPT invariance in gravitational waves with LIGO-Virgo catalog GWTC-1," Eur. Phys. J. C 80, 1032 (2020), arXiv:2002.00396 [gr-qc].
- [57] Tao Zhu, Wen Zhao, and Anzhong Wang, "Gravitational wave constraints on spatial covariant gravities," Phys. Rev. D 107, 044051 (2023), arXiv:2211.04711 [gr-qc].
- [58] Tao Zhu, Wen Zhao, and Anzhong Wang, "Polarized primordial gravitational waves in spatial covariant gravities," Phys. Rev. D 107, 024031 (2023), arXiv:2210.05259 [gr-qc].
- [59] Jin Qiao, Tao Zhu, Wen Zhao, and Anzhong Wang, "Polarized primordial gravitational waves in the ghostfree parity-violating gravity," Phys. Rev. D 101, 043528 (2020), arXiv:1911.01580 [astro-ph.CO].
- [60] Yi-Fan Wang, Stephanie M. Brown, Lijing Shao, and Wen Zhao, "Tests of gravitational-wave birefringence with the open gravitational-wave catalog," Phys. Rev. D 106, 084005 (2022), arXiv:2109.09718 [astro-ph.HE].
- [61] Wen Zhao, Tan Liu, Linqing Wen, Tao Zhu, Anzhong Wang, Qian Hu, and Cong Zhou, "Model-independent test of the parity symmetry of gravity with gravitational waves," Eur. Phys. J. C 80, 630 (2020), arXiv:1909.13007 [gr-qc].
- [62] Zheng Chen, Yang Yu, and Xian Gao, "Polarized gravitational waves in the parity violating scalar-nonmetricity theory," JCAP 06, 001 (2023), arXiv:2212.14362 [gr-qc].
- [63] Zhao Li, Jin Qiao, Tan Liu, Tao Zhu, and Wen Zhao, "Gravitational waveform and polarization from binary black hole inspiral in dynamical Chern-Simons gravity: from generation to propagation," JCAP 04, 006 (2023), arXiv:2211.12188 [gr-qc].
- [64] Jin Qiao, Tao Zhu, Guoliang Li, and Wen Zhao, "Post-Newtonian parameters of ghost-free parity-violating gravities," JCAP 04, 054 (2022), arXiv:2110.09033 [grqc].
- [65] Felipe Sulantay, Macarena Lagos, and Máximo Bañados, "Chiral gravitational waves in Palatini-Chern-Simons gravity," Phys. Rev. D 107, 104025 (2023), arXiv:2211.08925 [gr-qc].
- [66] Sai Wang, "Exploring the CPT violation and birefringence of gravitational waves with ground- and spacebased gravitational-wave interferometers," Eur. Phys. J. C 80, 342 (2020), arXiv:1712.06072 [gr-qc].
- [67] B. P. Abbott *et al.* (LIGO Scientific, Virgo), "Tests of General Relativity with the Binary Black Hole Signals

from the LIGO-Virgo Catalog GWTC-1," Phys. Rev. D **100**, 104036 (2019), arXiv:1903.04467 [gr-qc].

- [68] R. Abbott *et al.* (LIGO Scientific, Virgo), "Tests of general relativity with binary black holes from the second LIGO-Virgo gravitational-wave transient catalog," Phys. Rev. D **103**, 122002 (2021), arXiv:2010.14529 [gr-qc].
- [69] R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA), "Tests of General Relativity with GWTC-3," (2021), arXiv:2112.06861 [gr-qc].
- [70] Jin Qiao, Zhao Li, Tao Zhu, Ran Ji, Guoliang Li, and Wen Zhao, "Testing parity symmetry of gravity with gravitational waves," Front. Astron. Space Sci. 9, 1109086 (2023), arXiv:2211.16825 [gr-qc].
- [71] V. Alan Kostelecky and Matthew Mewes, "Electrodynamics with Lorentz-violating operators of arbitrary dimension," Phys. Rev. D 80, 015020 (2009), arXiv:0905.0031 [hep-ph].
- [72] Bo-Yang Zhang, Tao Zhu, Jing-Fei Zhang, and Xin Zhang, "Forecasts for Constraining Lorentz-violating Damping of Gravitational Waves from Compact Binary Inspirals," (2024), arXiv:2402.08240 [gr-qc].
- [73] Xian Gao and Xun-Yang Hong, "Propagation of gravitational waves in a cosmological background," Phys. Rev. D 101, 064057 (2020), arXiv:1906.07131 [gr-qc].
- [74] Jin Qiao, Tao Zhu, Wen Zhao, and Anzhong Wang, "Waveform of gravitational waves in the ghost-free parity-violating gravities," Phys. Rev. D 100, 124058 (2019), arXiv:1909.03815 [gr-qc].
- [75] Saeed Mirshekari, Nicolas Yunes, and Clifford M. Will, "Constraining Generic Lorentz Violation and the Speed of the Graviton with Gravitational Waves," Phys. Rev. D 85, 024041 (2012), arXiv:1110.2720 [gr-qc].
- [76] Cheng Gong, Tao Zhu, Rui Niu, Qiang Wu, Jing-Lei Cui, Xin Zhang, Wen Zhao, and Anzhong Wang, "Gravitational wave constraints on Lorentz and parity violations in gravity: High-order spatial derivative cases," Phys. Rev. D 105, 044034 (2022), arXiv:2112.06446 [gr-qc].
- [77] B. P. Abbott *et al.* (LIGO Scientific, Virgo, Fermi-GBM, INTEGRAL), "Gravitational Waves and Gamma-rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A," Astrophys. J. Lett. 848, L13 (2017), arXiv:1710.05834 [astro-ph.HE].
- [78] Qiang Wu, Tao Zhu, Rui Niu, Wen Zhao, and Anzhong Wang, "Constraints on the Nieh-Yan modified teleparallel gravity with gravitational waves," Phys. Rev. D 105, 024035 (2022), arXiv:2110.13870 [gr-qc].
- [79] Ziming Wang, Lijing Shao, and Chang Liu, "New Limits on the Lorentz/CPT Symmetry Through 50 Gravitational-wave Events," Astrophys. J. 921, 158 (2021), arXiv:2108.02974 [gr-qc].
- [80] Curt Cutler and Eanna E. Flanagan, "Gravitational waves from merging compact binaries: How accurately can one extract the binary's parameters from the inspiral wave form?" Phys. Rev. D 49, 2658–2697 (1994), arXiv:gr-qc/9402014.
- [81] Joseph D. Romano and Neil J. Cornish, "Detection methods for stochastic gravitational-wave backgrounds: a unified treatment," Living Rev. Rel. 20, 2 (2017), arXiv:1608.06889 [gr-qc].
- [82] Eric Thrane and Colm Talbot, "An introduction to Bayesian inference in gravitational-wave astronomy: parameter estimation, model selection, and hierarchical models," Publ. Astron. Soc. Austral. 36, e010 (2019), [Erratum: Publ.Astron.Soc.Austral. 37, e036 (2020)],

arXiv:1809.02293 [astro-ph.IM].

- [83] Neil J. Cornish and Tyson B. Littenberg, "BayesWave: Bayesian Inference for Gravitational Wave Bursts and Instrument Glitches," Class. Quant. Grav. **32**, 135012 (2015), arXiv:1410.3835 [gr-qc].
- [84] Tyson B. Littenberg and Neil J. Cornish, "Bayesian inference for spectral estimation of gravitational wave detector noise," Phys. Rev. D 91, 084034 (2015), arXiv:1410.3852 [gr-qc].
- [85] I. M. Romero-Shaw *et al.*, "Bayesian inference for compact binary coalescences with bilby: validation and application to the first LIGO–Virgo gravitational-wave transient catalogue," Mon. Not. Roy. Astron. Soc. **499**, 3295– 3319 (2020), arXiv:2006.00714 [astro-ph.IM].
- [86] Gregory Ashton *et al.*, "BILBY: A user-friendly Bayesian inference library for gravitational-wave astronomy," Astrophys. J. Suppl. **241**, 27 (2019), arXiv:1811.02042 [astro-ph.IM].
- [87] Gonzalo Morras, Jose Francisco Nuño Siles, Juan Garcia-Bellido, and Ester Ruiz Morales, "False alarms induced by Gaussian noise in gravitational wave detectors," Phys. Rev. D 107, 023027 (2023), arXiv:2209.05475 [gr-qc].
- [88] Cecilio García-Quirós, Marta Colleoni, Sascha Husa, Héctor Estellés, Geraint Pratten, Antoni Ramos-Buades, Maite Mateu-Lucena, and Rafel Jaume, "Multimode frequency-domain model for the gravitational wave signal from nonprecessing black-hole binaries," Phys. Rev. D 102, 064002 (2020), arXiv:2001.10914 [gr-qc].
- [89] Geraint Pratten et al., "Computationally efficient models for the dominant and subdominant harmonic modes of precessing binary black holes," Phys. Rev. D 103, 104056 (2021), arXiv:2004.06503 [gr-qc].
- [90] Geraint Pratten, Sascha Husa, Cecilio Garcia-Quiros, Marta Colleoni, Antoni Ramos-Buades, Hector Estelles, and Rafel Jaume, "Setting the cornerstone for a family of models for gravitational waves from compact binaries: The dominant harmonic for nonprecessing quasicircular black holes," Phys. Rev. D 102, 064001 (2020), arXiv:2001.11412 [gr-qc].
- [91] N. Aghanim *et al.* (Planck), "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. **641**, A6 (2020), [Erratum: Astron.Astrophys. 652, C4 (2021)], arXiv:1807.06209 [astro-ph.CO].
- [92] Joshua S. Speagle, "dynesty: a dynamic nested sampling package for estimating Bayesian posteriors and evidences," Mon. Not. Roy. Astron. Soc. 493, 3132–3158 (2020), arXiv:1904.02180 [astro-ph.IM].
- [93] LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration, "LVK Algorithm Library - LAL-Suite," Free software (GPL) (2018).
- [94] Charles R. Harris *et al.*, "Array programming with NumPy," Nature 585, 357–362 (2020), arXiv:2006.10256 [cs.MS].
- [95] Stéfan van der Walt, S. Chris Colbert, and Gaël Varoquaux, "The NumPy Array: A Structure for Efficient Numerical Computation," Comput. Sci. Eng. 13, 22–30 (2011), arXiv:1102.1523 [cs.MS].
- [96] Pauli Virtanen et al., "SciPy 1.0–Fundamental Algorithms for Scientific Computing in Python," Nature Meth. 17, 261 (2020), arXiv:1907.10121 [cs.MS].
- [97] John D. Hunter, "Matplotlib: A 2D Graphics Environment," Comput. Sci. Eng. 9, 90–95 (2007).