SOLAR PANEL SELECTION USING EXTENDED WASPAS WITH DISC INTUITIONISTIC FUZZY CHOQUET INTEGRAL OPERATORS: CASPAS METHODOLOGY

MAHMUT CAN BOZYİĞİT¹ AND MEHMET ÜNVER²

ABSTRACT. Renewable energy is crucial for addressing the growing energy demands of modern society while mitigating the adverse effects of climate change. Unlike fossil fuels, renewable energy sources such as solar, wind, hydro, geothermal, and biomass are abundant, sustainable, and environmentally friendly. This study focuses on addressing a critical challenge in renewable energy decision-making by developing a novel framework for optimal solar panel selection, a key component of sustainable energy solutions. Solar panel selection involves evaluating multiple interdependent criteria, such as efficiency, cost, durability, and environmental impact. Traditional multi-criteria decision-making (MCDM) methods often fail to account for the interdependencies among these criteria, leading to suboptimal outcomes. To overcome this limitation, the study introduces the Choquet Aggregated Sum Product Assessment (CASPAS) method, a Choquet integral-based MCDM approach that incorporates fuzzy measures to model interactions among criteria. CASPAS generalizes the Weighted Aggregated Sum Product Assessment (WASPAS) method, thereby enhancing decision-making accuracy and reliability. This study also introduces the concept of disc intuitionistic fuzzy set (D-IFS), a generalization of the concept of circular intuitionistic fuzzy set, which employ a radius function capable of assigning varying values to individual elements instead of relying on a fixed radius. Recognizing that traditional weighted aggregation operators neglect the interaction among criteria, this study proposes disc intuitionistic fuzzy Choquet integral operators by incorporating the concept of fuzzy measures, which are effective in modeling such interactions. The proposed method is applied to a renewable energy problem on selecting optimal solar panels. This application demonstrates how the CASPAS method develops decision-making accuracy by incorporating interdependencies among selection criteria. Comparative analysis with existing MCDM methods, alongside sensitivity and validation analyses, underscore the robustness and effectiveness of the CASPAS method.

1. INTRUDUCTION

Renewable energy sources such as solar, wind, hydroelectric, geothermal, and biomass are essential for achieving a sustainable future. These sources offer significant benefits by reducing greenhouse gas emissions and helping to mitigate climate change [47, 52]. Figure 1 shows all types of renewable energy. Solar energy captures sunlight using photovoltaic cells, while wind energy converts the kinetic energy of wind into electricity through turbines. Hydroelectric power generates electricity by utilizing the potential energy stored in water behind dams. Geothermal energy taps into the heat from the Earth's interior, and biomass energy comes from organic materials. These sources are environmentally friendly, helping to reduce our dependence on fossil fuels, which are major contributors to carbon dioxide emissions and global warming [5]. Although renewable energy technologies often come with high initial costs and can be influenced by geographic and climatic conditions, ongoing advancements in technology and supportive policies are making these sources more accessible and cost-effective [14]. Transitioning to renewable energy not only addresses environmental challenges but also promotes energy security and stimulates economic growth [58].

Solar energy offers several key advantages over other renewable sources. One of its primary benefits is its accessibility and flexibility. Solar panels can be installed on rooftops or open spaces, making it suitable for both urban and rural areas [39]. Unlike wind turbines, which

²⁰²⁰ Mathematics Subject Classification. 94D05, 90B50, 28E10.

Key words and phrases. D-IFSs, Choquet integral operators, CASPAS, MCDM, Solar panels.



FIGURE 1. Types of renewable energy

Region	Year	Hydropower	Wind E.	Solar E.	Bioenergy	Geothermal E.
	2023	1 406 863	$1\ 017\ 390$	$1 \ 418 \ 016$	148 840	15 026
World	2022	$1 \ 394 \ 824$	902 883	$1\ 070\ 851$	$144 \ 286$	$14\ 653$
	2021	$1 \ 362 \ 416$	$824 \ 321$	$870\ 643$	$138 \ 338$	$14 \ 432$
	2023	40 280	8 654	$13 \ 438$	1 899	991
Africa	2022	$39\ 275$	7 745	12 646	1 916	956
	2021	37 529	6 909	$11 \ 582$	$1 \ 917$	870
	2023	630 030	508 599	839 329	62 068	5 011
Asia	2022	$620 \ 391$	$426 \ 482$	$601 \ 390$	$58 \ 729$	4 761
	2021	$594 \ 325$	$384 \ 742$	488 907	54 608	4732
	2023	225 728	$255 \ 615$	288 644	42 825	1 637
Europe	2022	225 804	$240\ 278$	$233 \ 906$	$42 \ 458$	1 637
	2021	$224 \ 212$	$221 \ 517$	$190 \ 928$	41 601	1 633
	2023	199 492	$172 \ 328$	$156\ 000$	14 576	3678
N. America	2022	199 581	$164 \ 389$	129 156	14 684	3647
	2021	198 861	$154 \ 210$	109 579	14 906	3 595
	2023	180 962	39 852	49 392	20 348	83
S. America	2022	$180\ 018$	33 580	34 698	19 552	51
	2021	$178\ 482$	29 737	$21 \ 259$	18 622	40
	2023	15 542	14 045	$33 \ 417$	1 106	1 100
Oceania	2022	14 749	$13 \ 035$	29 566	1 104	1 100
	2021	14 509	$11 \ 522$	24 895	1 104	1 093
	~	() -				

TABLE 1. CAP (MW) by energy source and region between 2021 and 2023 according to IRENA [25]

require specific locations with consistent wind or hydroelectric plants that need large water bodies and can disrupt local ecosystems, solar installations are less intrusive and can be deployed in diverse environments [33, 47]. Moreover, solar energy systems have low maintenance costs and produce no emissions during operation, making them an environmentally friendly choice [39]. The falling costs of solar technology also make it an increasingly affordable option for many consumers [33, 47]. Table 1 presents the global capacity in megawatt (MW) for renewable energy sources from 2021 to 2023 [25]. Table 1 highlights global and regional trends in renewable energy capacities (MW) from 2021 to 2023, showing significant growth in solar and wind energy. Solar energy, the fastest-growing source, nearly doubles globally, with Asia leading the increase, followed by Europe and North America. This rapid growth reflects falling costs of photovoltaic technology, increasing efficiency, and widespread policy support for clean energy transitions. Wind energy also expands rapidly, particularly in Asia and Europe,

while hydropower, the largest contributor, grows more modestly due to its already extensive development. Regions like South America and Africa exhibit slower but steady progress, focusing on solar and hydropower. Bioenergy and geothermal energy maintain stable contributions worldwide. The remarkable rise of solar energy not only emphasizes its critical role in reducing carbon emissions but also highlights its accessibility and scalability, making it a cornerstone of the global shift toward cleaner, more sustainable energy systems. Figure 2 shows the growth of solar photovoltaic production in Gigawatt-hours (GWh) globally and across regions from 2014 to 2022 [25]. Worldwide production increased significantly, from 183,473 GWh in 2014 to 1,281,654 GWh in 2022, reflecting rapid adoption of solar energy. Asia leads with the highest production, growing from 55,392 GWh to 686,755 GWh, followed by Europe and North America, which reached 231,064 GWh and 210,107 GWh, respectively. While Africa, South America, and Oceania started with lower production levels, they also saw notable growth, particularly after 2019, driven by investments in renewable energy and technological advancements. This trend underscores the global shift toward cleaner energy sources.



FIGURE 2. Global solar photovoltaic production data by region between 2014 and 2022

Solar panels, often referred to as photovoltaic panels, are systems that harness sunlight and transform it into electricity [19]. This process, known as the photovoltaic effect, occurs when light energy stimulates electrons within a semiconductor material, producing an electric current [38]. Made up of numerous interconnected solar cells, these panels are utilized extensively across residential, commercial, and industrial sectors, as well as in space missions [53]. This technology, which has an important place among renewable energy sources today, has undergone a long scientific and technological development process. The roots of solar panel technology trace back to 1839, when Becquerel [7] discovered the photovoltaic effect. He found that certain materials could produce an electric current under sunlight [19]. While this discovery was revolutionary, it initially remained a theoretical concept without immediate practical use. Becquerel's observation went unreplicated until 1873, when Smith [51] found that light striking selenium could generate a charge. Building on this finding, Adams and Day [1] conducted experiments to confirm Smith's results and published their work, The Action of Light on Selenium, in 1876. In 1883, Charles Fritts developed the first functional solar cell using selenium [19, 38]. Despite its low efficiency of just 1%, this marked the beginning of the technology capable of directly

converting sunlight into electricity [19]. Einstein [17] made a significant impact on solar panel technology by explaining the photoelectric effect in 1905. He proposed that light consists of tiny energy packets known as photons. When these photons hit a material, they can release electrons from its surface, a process known as the photoelectric effect. This discovery is fundamental to the operation of solar cells [38]. In 1939, Ohl [40] developed the solar cell design that is now commonly used in many modern solar panels. He patented his design in 1941. In 1954, scientists Chapin et al. [11] at Bell Laboratories developed the first modern solar cell using silicon. Their creation had an efficiency of around 6%, marking the start of practical solar panel technology [19]. During this period, solar panels became commercially available, but their high production costs restricted their use primarily to specialized applications like satellites. Solar panels were first used in space on the Vanguard I satellite in 1958. This marked a critical point for solar energy development, as it demonstrated their reliability in extreme conditions [16]. The 1970s oil crisis spurred interest in renewable energy, leading to significant investments in solar technology [48]. During this time, the cost of solar power began to decline as production scaled up. Today, solar panels play a crucial role in global renewable energy strategies, with ongoing innovations continually improving their efficiency and making them more accessible [38]. Figure 3 illustrates the timeline of solar panel development.



FIGURE 3. Timeline of the development of solar panels

Fuzzy sets (FSs), introduced by Zadeh [65], represent a significant extension of classical set theory by providing a means to handle uncertainty and vagueness in data. Unlike crisp sets, where an element either fully belongs to a set or does not belong at all, FSs allow for partial membership. Formally, a FS A in a universe of discourse Δ is defined by a membership function $\mu_A : \Delta \to [0, 1]$, where $\mu_A(\delta)$ quantifies the degree to which an element δ belongs to A. This approach is particularly useful in scenarios where sharp boundaries are not appropriate, such as in the analysis of imprecise or subjective information. By accommodating varying degrees of membership, FSs provide a flexible mathematical framework for modeling real-world phenomena where crisp distinctions are not feasible.

Despite their utility, FSs have limitations in capturing the full spectrum of uncertainty. To address these, Atanassov [2] proposed intuitionistic fuzzy sets (IFSs), which further improve the expressive power of FSs. IFSs expand on the concept of partial membership by associating each element in a set with two parameters: a membership degree and a non-membership degree. These values provide a more nuanced representation of uncertainty by allowing the sum of membership and non-membership degrees to be less than or equal to 1, thereby introducing a third dimension called hesitation. Hesitation reflects the degree of indeterminacy or lack of knowledge about an element's status within the set. This additional flexibility makes IFSs highly effective for applications requiring a deeper and more comprehensive representation of uncertainty, such as decision-making, pattern recognition, and fuzzy control systems. By capturing both membership and non-membership alongside hesitation, IFSs offer a more robust tool for tackling complex problems characterized by ambiguity and incomplete information.

In decision-making problems involving uncertainty and ambiguity, assigning specific intuitionistic fuzzy values (IFVs) to evaluation criteria can be challenging. This difficulty stems from the inherent uncertainty in determining the exact degrees to which an element belongs or does not belong to a set. Traditional IFSs may lack the necessary flexibility to handle situations where multiple decision-makers interpret criteria differently or have diverse perspectives. To address this limitation, Atanassov [3, 4] introduced the concept of circular intuitionistic fuzzy sets (C-IFSs), an extension of the conventional IFS framework. C-IFSs use a circular structure to represent the degrees of membership and non-membership, which inherently captures the vagueness and uncertainty associated with these values. This circular representation provides a more dynamic and sensitive way to reflect changes in both membership and non-membership degrees, making it particularly useful in complex decision-making scenarios. The circular design enhances flexibility by accommodating the imprecision and ambiguity often present in real-world problems. As a result, C-IFSs offer decision-makers a more robust framework for evaluating alternatives, especially when assigning precise IFVs to criteria is impractical. Building on this concept, Bozyiğit et al. [8] introduced circular Pythagorean fuzzy sets (C-PFSs), which expand upon C-IFSs by offering a broader representational range. A C-PFS is visualized as a circle that encapsulates both the degree of membership and the degree of non-membership. The center of this circle is defined by two non-negative real numbers, μ and ν , which must satisfy the condition $0 \le \mu^2 + \nu^2 \le 1$. Since the radius degree remains constant for each C-PFS, disc Pythagorean fuzzy sets (D-PFSs) were proposed by replacing this constant radius with a radius function [30]. This approach introduces flexibility to the radius degree.

Multi-Criteria Decision-Making (MCDM) is a specialized field within decision science that addresses the challenge of identifying the most suitable alternatives when faced with multiple and sometimes conflicting criteria. It offers a structured approach to help decision-makers navigate complex situations by systematically analyzing trade-offs between various objectives. MCDM techniques combine both quantitative and qualitative information, enabling a comprehensive evaluation of options. Through the use of mathematical frameworks and well-defined decision processes, MCDM improves the ability to pinpoint optimal solutions, ensures greater clarity and fairness in decision-making, and supports alignment with overarching strategic objectives. Weighted Aggregated Sum Product Assessment (WASPAS) method, introduced by Zavadskas et al. [66], is a robust and versatile MCDM approach that integrates Weighted Sum Model (WSM) and Weighted Product Model (WPM). By combining these two methodologies, WASPAS aims to develop decision-making accuracy and reliability. It assigns weights to criteria based on their relative importance and evaluates alternatives through a hybrid aggregation mechanism, which includes both additive and multiplicative components. The flexibility of WASPAS allows it to effectively handle a wide range of decision-making scenarios, particularly in cases where decision-makers need to balance between conflicting criteria. Fuzzy WASPAS extends the traditional WASPAS method by incorporating fuzzy set theory to address uncertainties and imprecision in decision-making [59, 67]. In real-world applications, decision-makers often face ambiguity in assigning precise values to criteria weights and alternatives. Fuzzy WAS-PAS employs fuzzy values to capture this vagueness. This enables a more realistic representation of preferences and better models human judgment. The fuzzy extension makes the WASPAS method particularly suitable for complex decision-making environments, such as supply chain management, project selection, and resource allocation, where uncertainty is a significant factor. Using the strengths of both fuzzy logic and the hybrid aggregation mechanism, Fuzzy WASPAS provides a powerful tool for solving intricate MCDM problems.

Choquet [13] introduced the concept of capacity and the Choquet integral, an extension of the Lebesgue integral. When defined using an additive measure, Choquet integral simplifies to a weighted arithmetic mean. Building on this, Sugeno [55] expanded the notion of capacity to fuzzy measure and developed the Sugeno integral. Unlike additive measures, fuzzy measures can better capture the interaction among criteria, making fuzzy integrals particularly effective in such contexts. Fuzzy measures and fuzzy integrals are frequently used tools in decision making and pattern recognition problems ([41, 61, 62]). Among various fuzzy measures, the λ -fuzzy measure, defined by Sugeno [55], is notably significant. This measure is constructed using interaction indices and element weights. Takahagi [56] proposed three algorithms to efficiently generate λ -fuzzy measure, accounting for both interaction indices and weight allocations. Further extending these ideas, Tan and Chen [57] introduced the IF Choquet integral operator, designed to aggregate IFVs for MCDM in place of traditional operators reliant on additive measures. Later, Peng and Yang [44] generalized the Choquet integral operator from IFVs to Pythagorean fuzzy values (PFVs), presenting MABAC method based on PF Choquet integral operator. Bozyiğit et al. [10] proposed C-PF Choquet integral operators, which generalize some weighted arithmetic aggregation operators, to aggregate circular Pythagorean fuzzy values (C-PFVs).

This paper offers four significant contributions:

- Theoretical contributions: To improve the flexibility of the fixed radius degree in C-IFSs, this paper introduces the concept of disc intuitionistic fuzzy sets (D-IFSs), which incorporate a radius function. D-IFSs generalize C-IFSs by providing a more expansive framework. Following this, set operations specific to D-IFSs are introduced. Additionally, algebraic operations on disc intuitionistic fuzzy values (D-IFVs) are presented to extend the radius degree $\sqrt{2}$, which is restricted to 1 in circular intuitionistic fuzzy values (C-IFVs) that overlap with D-IFVs. Weighted aggregation operators are then introduced for aggregating D-IFVs. However, since these aggregation operators do not account for interactions between criteria and such interactions may exist in real-world problems, disc intuitionistic fuzzy Choquet integral operators are proposed in the study. These operators use fuzzy measures to consider the interaction between criteria.
- Methodological Contributions: CASPAS (Choquet Aggregated Sum Product Assessment) method introduces a novel approach to multi-criteria decision-making by incorporating Choquet Sum Model (CSM) and Choquet Product Model (CPM), replacing the traditional WSM and WPM used in WASPAS. These innovations employ the Choquet integral to account for interactions and dependencies among criteria, addressing limitations in additive and multiplicative aggregation methods. By integrating a capacity function, CSM and CPM enable more nuanced evaluations, capturing synergies and conflicts between criteria. Compared to WASPAS, CASPAS offers superior accuracy, adaptability, and applicability, marking a significant methodological contribution to decision-making in complex interdependent systems.
- Applied Contributions: This study examines a MCDM problem related to the selection of solar panels based on the criteria of efficiency, cost, durability, and installation complexity. The proposed CASPAS method is applied to evaluate some solar panels, demonstrating its capability to handle interdependent criteria effectively. Following the application, comparison analysis, sensitivity analysis, and validity analysis are conducted to assess the method's robustness and reliability. These analyses highlight the

practical relevance of CASPAS in solar panel selection and its potential to deliver accurate and consistent decision-making outcomes in real-world applications.

• Interdisciplinary Contributions: Using CASPAS method for solar panel selection, this study brings together multiple disciplines to offer a new approach to decision-making processes. It integrates engineering, energy management, economics, and environmental sciences to effectively evaluate technical, economic, environmental, and installation-related factors in solar panel selection. The use of CASPAS provides an innovative contribution to existing methods in engineering and energy planning by considering the interactions between criteria in MCDM processes. Moreover, this study enables the development of an interdisciplinary approach for sustainable energy solutions and environmental impact assessments. Thus, it allows for a more effective and efficient selection of solar energy systems from both technical engineering and environmental economics perspectives.

The structure of the paper is as follows: Section 2 presents a comprehensive literature review covering C-IFSs, existing extended WASPAS methods, and various types of solar panels. It also introduces the theoretical foundation of the proposed CASPAS method. Section 3 focuses on the development of fundamental set operations for D-IFSs and the definition of score and accuracy functions for ranking D-IFVs. Additionally, algebraic operations for D-IFVs are formulated by extending the radius limit from 1 to $\sqrt{2}$. Using these algebraic operations, weighted arithmetic and geometric aggregation operators for D-IFVs are established. In Section 4, after recalling the fundamental concepts of fuzzy measures and the Choquet integral, disc intuitionistic fuzzy Choquet interval operators are introduced. These operators extend the weighted arithmetic and geometric aggregation operators by incorporating interactions between criteria, addressing challenges commonly found in real-world problems. Section 5 begins with an overview of solar panel types and subsequently proposes CASPAS method for D-IFSs. This method advances the WASPAS approach by integrating CSM and CPM techniques. Section 6 explores the sensitivity of the CASPAS method to various parameters and provides validity analysis. Section 7 evaluates the performance of CASPAS through comparisons with existing methods in the literature. The paper concludes with Section 8, which discusses the findings, draws conclusions, and outlines directions for future research.

2. LITERATURE RESEARCH

This section provides an overview of existing studies and methodologies related to C-IFSs, the types of solar panels and extended WASPAS approaches.

2.1. A comparison of C-IFSs and D-IFSs. C-IFSs are an extension of IFS, designed to better express uncertainty and fuzziness. In C-IFSs, each element is represented as a circle, where the center of the circle indicates the membership and non-membership degrees. The radius of the circle represents the hesitation degree. C-IFSs are particularly useful in MCDM, allowing for more flexible modeling of uncertain information. C-IFSs contain more information compared to IFSs, making them effective in solving more complex problems. Atanassov and Marinov [4] introduced the concept of distance measures for C-IFS and proposed four different distance measures based on the classical metric definition. For decision-making applications, Chen [12] proposed Minkowski distance measures specifically designed for C-IFSs. Bozyiğit and Ünver [9] developed score and accuracy functions to rank C-IFVs. They also introduced parametric distance measures based on Hamming and Euclidean metrics for C-IFVs. Garg et al. [20] proposed algebraic operations for C-IFVs, and based on these operations, they introduced weight aggregation operators based on Archimedean t-norms for C-IFVs. Using these aggregation operators, they developed an extended EDAS method, which is a MCDM approach.

In C-IFSs, the radius degree for each element is treated as a constant, which imposes a limitation. To address this restriction, the concept of D-IFSs is introduced in this study, where the radius is variable rather than fixed. In D-IFSs, the radius is represented by a radius

function, similar to the membership and non-membership functions. This provides flexibility in the radius degree for application areas such as decision-making, pattern recognition, and classification. Figure 4 displays a visual comparison of C-IFSs and D-IFSs. As shown in Figure 4, the radius remains constant in a C-IFS, whereas in a D-IFS, the radius can vary.



FIGURE 4. Comparison of C-IFSs and D-IFSs

2.2. A comparison of WASPAS and CASPAS approaches. WASPAS method is a wellestablished MCDM approach that combines the strengths of WSM and WPM. By integrating these two techniques, WASPAS offers a balance between simplicity and computational efficiency. However, a key limitation of WASPAS lies in its assumption of independence among criteria, which may not hold true in real-world scenarios where criteria often exhibit interdependencies. This limitation can lead to suboptimal or less accurate results in complex decision-making problems. WASPAS method has been extensively studied in the literature due to its simplicity and effectiveness in handling MCDM problems. In Table 2, a summary of key studies on WASPAS and its applications is provided, illustrating the various areas where the method has been applied and the contributions made in the literature.

CASPAS, a novel approach, addresses this limitation by employing Choquet integrals through its CSM and CPM. These models allow CASPAS to account for interdependencies among criteria, enabling it to capture the interactions that influence the decision-making process more accurately. By incorporating these interdependencies, CASPAS enhances the realism and reliability of the results, making it particularly suitable for applications with complex criteria relationships. Furthermore, CASPAS introduces greater flexibility in the aggregation process by providing a mechanism to model the influence of each criterion based on its interaction with others. This feature not only improves the accuracy of the method but also extends its applicability to a broader range of decision-making scenarios. As a result, CASPAS emerges as a superior alternative to WASPAS in contexts where the relationships between criteria play a critical role, such as renewable energy selection, financial analysis, and engineering design problems. Figure 6 displays a table comparing the features of WASPAS and CASPAS. Figure 5 also illustrates the stages and steps involved in the CASPAS method.

2.3. Types of solar panels. Solar panels are advanced systems designed to harness sunlight and convert it into electrical energy, making them a cornerstone of clean and sustainable energy solutions. With rising energy demands and increasing environmental concerns, solar technology has seen significant advancements, resulting in the creation of various panel types tailored to diverse applications [39]. Each type offers distinct benefits in terms of efficiency, costeffectiveness, durability, and suitability for specific uses. The most commonly used types of solar panels are as follows:

Year	Environment	Author(s)	Application area
2012	Crips	Zavadskas et al. [66]	Illustrative example
2014	Interval valued intuitionistic fuzzy sets	Zavadskas et al. [67]	Derelict buildings' redevelop- ment decisions and investment
2015	Fuzzy sets	Turskis et al. [59]	Construction site selection
2016	Interval type-2 fuzzy sets	Keshavarz Ghorabaee et al. [29]	Green supplier selection
2017	Single-valued neutrosophic set	Baušys and Juodagalvienė [6]	Garage location selection
2018	Intuitionistic fuzzy sets	Stanujkić and Karabašević [54]	Website evaluation
2019	Spherical fuzzy sets	Kutlu Gundogdu and Kahraman [32]	Industrial robot selection
2020	q-rung orthopair fuzzy sets	Rani and Mishra [46]	Fuel technology selection
2021	Fermatean fuzzy sets	Mishra and Rani [37]	Healthcare waste disposal loca- tion selection
2022	Picture fuzzy sets	Senapati and Chen [49]	Air-conditioning system selection
2023	Pythagorean fuzzy Sets	Yalcin Kavus et al. [64]	Parcel locker location selection
2024	Complex Fuzzy Sets	Khan et al. [31]	Parameter Selection Impacting Software Reliability

TABLE 2. Literature survey in different fuzzy settings for WASPAS approach



0

Construct decision matrices by experts STAGE 2: Assessment of weights STAGE 3: Calculation of Choquet integral operators

> Constuct a fuzzy measure Calculate Choquet Sum Model (CSM) and Choque Product Model (CPM)



of the significance degrees

Bank alternatives according to the highest score

Find significance degrees of alternatives through CSM and CPM

FIGURE 5. Flowchart of CASPAS method

- Monocrystalline silicon panels: They are widely known for their high efficiency, typically ranging from 18% to 22%. Made from a single, continuous silicon crystal, these panels are ideal for limited spaces as they generate more energy per square meter. They are more expensive to produce due to their manufacturing process but are highly durable and offer long lifespans, making them a popular choice for residential and commercial installations [15, 45].
- Polycrystalline silicon panels: Polycrystalline panels are less efficient than monocrystalline panels, with efficiency levels around 15% to 17%. However, they are more affordable to produce, making them a cost-effective choice for larger installations where space is not a major constraint. These panels are made by melting silicon crystals

MAHMUT CAN BOZYİĞİT¹ AND MEHMET ÜNVER²

Feature	WASPAS	CASPAS
Method Type	Combines Weighted Sum Model (WSM) and Weighted Product Model (WPM).	Utilizes Choquet Sum Model (CSM) and Choquet Product Model (CPM) through Choquet integrals.
Assumption	Assumes criteria are independent.	Considers interdependencies among criteria.
Accuracy	Moderate accuracy, dependent on the independence assumption.	Higher accuracy by incorporating interdependencies.
Flexibility	Limited flexibility in aggregation processes.	Greater flexibility in modeling and aggregation processes.
Applicability	Suitable for simpler decision- making problems with independent criteria.	Ideal for complex scenarios with interacting criteria.

FIGURE 6. Comparison of WASPAS and CASPAS

and forming the panel from the solidified material, which results in a lower efficiency compared to monocrystalline panels [15, 45].

- Thin-film solar Panels: Thin-film solar panels are lightweight and flexible, making them ideal for applications where traditional panels might not be suitable. These panels have lower efficiency (typically 10% to 13%) but are cheaper to manufacture. Thin-film panels can be used on a variety of surfaces, including curved or irregular structures, and are often used for large-scale projects or specific architectural designs. Thin-film solar cells are preferred due to their low material requirements and improving efficiency levels. The primary technologies in this category are amorphous silicon (α -Si), copper indium gallium selenide (CIGS), and cadmium telluride (CdTe) [34].
- **PERC** (Passivated Emitter and Rear Cell) solar panels: PERC panels are an improved version of monocrystalline and polycrystalline panels. They feature an additional layer on the rear side of the cell that reflects unabsorbed light back into the panel, increasing efficiency. This technology boosts energy conversion, particularly in low-light conditions or on cloudy days [21].
- **Bifacial solar panels:** These panels represent an innovative advancement in photovoltaic technology, capable of generating electricity from both sides of the panel. Unlike conventional monofacial panels, which capture sunlight only on the front, bifacial panels feature solar cells on both the front and rear. This dual-sided design enables them to harness light not only directly from the sun but also from light reflected off the ground or nearby surfaces, greatly enhancing their overall energy production [22].

Each of these panel types offers unique benefits, depending on factors such as material, efficiency, cost, and installation requirements.

Decision-making problems associated with solar energy have been extensively explored in the literature. Notably, the selection of solar panel types and determining optimal locations for solar energy production are among the most significant decision-making issues. Table 3 provides an overview of studies focusing on decision-making in solar energy.

Author(s)	Method	Solar energy application			
Bozyiğit et al. [8]	A MCDM method based on aggre- gation operators and cosine similarity measures in C-PFSs	Selecting solar cell			
El-Bayeh et al. [18]	MCDM algorithm based on the concept of Rank-Weight-Rank	Solar panels selection in buildings			
Hosouli and Hassani [23]	Fuzzy Analytic Hierarchy Process method	Solar plant location selection			
Kannan et al. [27]	A hybrid approach based on MCDM methods and Monte Carlo simulation	Sustainable evaluation of potential solar sites			
Kaur et al. [28]	TOPSIS based on entropy technique	Selection of solar panel			
Shayani Mehr et al. [50]	A BWM-MULTIMOOSRAL approach	Solar panel technology selection			
Tüysüz and Kahraman [60]	An integrated picture fuzzy Z-AHP & TOPSIS methodology	Solar panel selection			
TABLE 3. Some MCDM studies from the literature on solar energy					

3. DISC INTUITIONISTIC FUZZY SETS

At an assov [2] introduced the concept of IFS, which is an extension of fuzzy sets FS. Throughout this paper Δ denotes a non-empty finite set.

Definition 1. [2] An IFS I in Δ is defined as:

 $I = \{ \langle \delta_i, \mu_I(\delta_i), \nu_I(\delta_i) \rangle : \delta_i \in \Delta \},\$

where μ_I and ν_I are membership and non-membership functions, respectively with the condition $\mu_I(\delta_i) + \mu_I(\delta_i) \leq 1$ for every $\delta_i \in \Delta$. Additionally, an IFV θ is defined as a pair $\langle \mu_{\theta}, \nu_{\theta} \rangle$, where μ_{θ} and ν_{θ} are both within the range [0, 1] and satisfy the condition $\mu_{\theta} + \nu_{\theta} \leq 1$.

C-IFSs, which was introduced by Atanassov [3], are an extension of IFSs. C-IFSs provide decision makers with a means to express uncertainty by utilizing membership and non-membership degrees represented in the form of a circle. In C-IFSs, the membership and non-membership degrees are visually represented by a circle, allowing decision makers to intuitively understand the degrees of membership and non-membership associated with elements in the set.

Definition 2. [3, 4] Let $r \in [0, \sqrt{2}]$. A C-IFS C_r in Δ is defined as:

 $C_r = \{ \langle \delta_i, (\mu_C(\delta_i), \nu_C(\delta_i)); r \rangle : \delta_i \in \Delta \},\$

where μ_C and ν_C are membership and non-membership functions, respectively with the condition $\mu_C(\delta_i) + \nu_C(\delta_i) \leq 1$ for every $\delta_i \in \Delta$. The value r represents the radius of the circle centered at the point $(\mu_C(\delta_i), \nu_C(\delta_i))$ on the plane. This circle visually represents the membership degree and non-membership degree associated with $\delta_i \in \Delta$.

Definition 3. A D-IFS D_r in Δ is defined as:

$$D_{\hat{r}} = \{ \langle \delta_i, (\mu_D(\delta_i), \nu_D(\delta_i)); \hat{r}_D(\delta_i) \rangle : \delta_i \in \Delta \},\$$

where μ_D and ν_D are membership and non-membership functions, respectively with the condition $\mu_D(\delta_i) + \nu_D(\delta_i) \leq 1$ for every $\delta_i \in \Delta$ and r_D is a radius function with $r_D(\delta_i) \in [0, \sqrt{2}]$ for every $\delta_i \in \Delta$. The value $r_D(\delta_i)$ represents the radius of the circle centered at the point $(\mu_D(\delta_i), \nu_D(\delta_i))$ on the plane. This circle visually represents the membership degree and non-membership degree associated with $\delta_i \in \Delta$.

Definition 4. Consider two values μ_{θ} and ν_{θ} within the range $[0, \sqrt{2}]$, satisfying the condition $\mu_{\theta} + \nu_{\theta} \leq 1$, and a value r_{θ} within the range [0, 1]. The triple $\theta = \langle (\mu_{\theta}, \nu_{\theta}); r_{\theta} \rangle$ is called both C-IFV and D-IFV. That is, C-IFV and D-IFV coincide.

Remark 1. Every C-IFS C_r can be expressed as a special case of a D-IFS by assigning a constant value r to the \hat{r}_D radius function. In other words, we can write any C-IFS C_r as:

 $C_r = D_{\hat{r}=r} = \{ \langle \delta_i, (\mu_D(\delta_i), \nu_D(\delta_i)); r \rangle : \delta_i \in \Delta \}.$

This implies that every C-IFS is also a D-IFS. Each D-IFS may not necessarily be a C-IFS. Hence, D-IFS is considered as an extension of the C-IFS framework.

The following definition outlines the set operations defined for D-IFSs.

Definition 5. Consider two D-IFSs, $D_{\hat{r}}$ and $E_{\hat{r}}$ and they are defined as follows:

 $D_{\hat{r}} = \{ \langle \delta_i, (\mu_D(\delta_i,), \nu_D(\delta_i,)); \hat{r}_D(\delta_i) \rangle : \delta_i \in \Delta \} \text{ and } E_{\hat{r}} = \{ \langle \delta_i, (\mu_E(\delta_i), \nu_E(\delta_i)); \hat{r}_E(\delta_i) \rangle : \delta_i \in \Delta \}.$

Some set operations between two D-IFSs can be defined as follows:

i) $D_{\hat{r}} \subset E_{\hat{r}}$ if and only if $\hat{r}_D(\delta_i) \leq \hat{r}_E(\delta_i)$, $\mu_D(\delta_i) \leq \mu_E(\delta_i)$, and $\nu_D(\delta_i) \geq \nu_E(\delta_i)$ for each $\delta_i \in \Delta$.

ii) $D_{\hat{r}} = E_{\hat{r}}$ *if and only if* $D_{\hat{r}} \subset E_{\hat{r}}$ *and* $E_{\hat{r}} \subset D_{\hat{r}}$.

iii) The complement of $D_{\hat{r}}$ is denoted as $D_{\hat{r}}^c$ and is defined as $D_{\hat{r}}^c = \{ \langle \delta_i, (\nu_D(\delta_i), \mu_D(x_i)); \hat{r}_D(\delta_i) \rangle : \delta_i \in \Delta \}.$

We propose an approach to ranking two DIFVs using both the score function and the accuracy function. These functions are outlined as follows.

Definition 6. Let $\theta = \langle (\mu_{\theta}, \nu_{\theta}); r_{\theta} \rangle$ represent a D-IFV (or C-IFV) and let $\xi \in [0, 1]$. A score function S and the accuracy function \mathcal{H} are defined for D-IFVs as

$$S(\theta) = \xi\left(\frac{\mu_{\theta} - \nu_{\theta} + 1}{2}\right) + (1 - \xi)\frac{r_{\theta}}{\sqrt{2}}$$

and

$$\mathcal{H}(\theta) = \xi(\mu_{\theta} + \nu_{\theta}) + (1 - \xi)\frac{r_{\theta}}{\sqrt{2}},$$

respectively. One can see that $0 \leq S(\theta) \leq 1$ and $0 \leq \mathcal{H}(\theta) \leq 1$. By altering $\xi \in [0, 1]$, a decision maker has the ability to control how much intuitionistic fuzzy information and radius information influence the score function's outcome. When ξ equals 1, the degree of radius is effectively excluded from the decision-making process.

Definition 7. Consider two D-IFVs, $\theta = \langle (\mu_{\theta}, \nu_{\theta}); r_{\theta} \rangle$ and $\kappa = \langle (\mu_{\kappa}, \nu_{\kappa}); r_{\kappa} \rangle$. θ and κ can be ranked by the following procedure:

- (1) If $S(\theta) < S(\kappa)$, then θ is ranked lower than κ , showed by $\theta \leq \kappa$.
- (2) If $\mathcal{S}(\theta) = \mathcal{S}(\kappa)$,
 - (2a) If $\mathcal{H}(\theta) < \mathcal{H}(\kappa)$, then θ is ranked lower than κ .
 - (2b) If $\mathcal{H}(\theta) = \mathcal{H}(\theta)$, then θ is equivalent to κ , showed by $\theta \equiv \kappa$.

Some algebraic operations defined for C-IFVs are also valid for D-IFVs [20]. We extend the existing algebraic operations, which are also valid for C-IFVs, by considering that the radius lies between 0 and $\sqrt{2}$.

Definition 8. Let $\theta = \langle (\mu_{\theta}, \nu_{\theta}); r_{\theta} \rangle$ and $\kappa = \langle (\mu_{\kappa}, \nu_{\kappa}); r_{\kappa} \rangle$ represent two *D*-IFVs (or *C*-IFVs) and $\zeta > 0$. We have some algebraic operations based on radius between *D*-IFVs as follows: a) $\theta \oplus \kappa = /(1 - (1 - \mu_{c})(1 - \mu_{c})) + (r_{\theta}r_{\kappa})$

$$a) \ \theta \oplus_{q} \kappa = \left\langle \left(1 - (1 - \mu_{\theta})(1 - \mu_{\kappa}), \nu_{\theta}\nu_{\kappa}\right); \frac{\delta \kappa}{\sqrt{2}} \right\rangle \\b) \ \theta \oplus_{p} \kappa = \left\langle \left(1 - (1 - \mu_{\theta})(1 - \mu_{\kappa}), \nu_{\phi}\nu_{\kappa}\right); \sqrt{2} - \sqrt{2}\left(1 - \frac{r_{\theta}}{\sqrt{2}}\right)\left(1 - \frac{r_{\kappa}}{\sqrt{2}}\right) \right\rangle \\c) \ \theta \otimes_{q} \kappa = \left\langle \left(\mu_{\theta}\mu_{\kappa}, 1 - (1 - \nu_{\theta})(1 - \nu_{\kappa})\right); \frac{r_{\theta}r_{\kappa}}{\sqrt{2}} \right\rangle$$

$$\begin{aligned} \boldsymbol{d}) \ \theta \otimes_{p} \kappa &= \left\langle \left(\mu_{\theta}\mu_{\kappa}, 1 - (1 - \nu_{\theta})(1 - \nu_{\kappa})\right); \sqrt{2} - \sqrt{2}\left(1 - \frac{r_{\theta}}{\sqrt{2}}\right) \left(1 - \frac{r_{\kappa}}{\sqrt{2}}\right) \right. \\ \boldsymbol{e}) \ \zeta_{q}\theta &= \left\langle (1 - (1 - \mu_{\theta})^{\zeta}, \nu_{\theta}^{\zeta}); \sqrt{2}\left(\frac{r_{\theta}}{\sqrt{2}}\right)^{\zeta} \right\rangle \\ \boldsymbol{f}) \ \zeta_{p}\theta &= \left\langle (1 - (1 - \mu_{\theta})^{\zeta}, \nu_{\theta}^{\zeta}); \sqrt{2} - \sqrt{2}\left(1 - \frac{r_{\theta}}{\sqrt{2}}\right)^{\zeta} \right\rangle \\ \boldsymbol{g}) \ \theta^{\zeta_{q}} &= \left\langle (\mu_{\theta}^{\zeta}, 1 - (1 - \nu_{\theta})^{\zeta}); \sqrt{2}\left(\frac{r_{\theta}}{\sqrt{2}}\right)^{\zeta} \right\rangle \\ \boldsymbol{h}) \ \theta^{\zeta_{p}} &= \left\langle (\mu_{\theta}^{\zeta}, 1 - (1 - \nu_{\theta})^{\zeta}); \sqrt{2} - \sqrt{2}\left(1 - \frac{r_{\theta}}{\sqrt{2}}\right)^{\zeta} \right\rangle \end{aligned}$$

Definition 9. Consider a collection of D-IFVs denoted as $\{\theta_k = \langle (\mu_{\theta_k}, \nu_{\phi_k}); r_{\theta_k} \rangle$: $k = 1, \ldots, m\}$. Weighted arithmetic aggregation operators are mappings defined as

$$D\text{-}IFWAO_q(\theta_1,...,\theta_m) := (q) \bigoplus_{k=1}^m \omega_{k_q} \theta_k.$$

and

$$D\text{-}IFWAO_p(\theta_1,...,\theta_m) := (p) \bigoplus_{k=1}^m \omega_{k_p} \theta_k.$$

respectively. Here, each weight ω_i satisfies $0 \leq \omega_i \leq 1$ for $k = 1, \ldots, m$, and the sum of all weights equals 1.

It can be easily obtained from Definition 9 that

$$D\text{-}IFWAO_q(\theta_1,\ldots,\theta_m) = \left\langle \left(1 - \prod_{k=1}^m (1 - \mu_{\theta_k})^{\omega_k}, \prod_{k=1}^m \nu_{\theta_k}^{\omega_k}\right); \prod_{k=1}^m r_{\theta_k}^{\omega_k} \right\rangle,$$

and

$$D\text{-}IFWAO_p(\theta_1,\ldots,\theta_m) = \left\langle \left(1 - \prod_{k=1}^m (1 - \mu_{\theta_k})^{\omega_k}, \prod_{k=1}^m \nu_{\theta_k}^{\omega_k}\right); \sqrt{2} - \prod_{k=1}^m \left(\sqrt{2} - r_{\theta_k}\right)^{\omega_k} \right\rangle,$$

respectively.

Definition 10. Given a collection of D-IFVs denoted as $\{\theta_k = \langle (\mu_{\theta_k}, \nu_{\theta_k}); r_{\theta_k} \rangle : k = 1, ..., m\}$. Weighted geometric aggregation operators are mappings defined as

$$D$$
-IFWGO_q $(\theta_1, \ldots, \theta_m) := (q) \bigotimes_{k=1}^m \theta_k^{\omega_{k_q}}.$

and

$$D\text{-}IFWGO_p(\theta_1,...,\theta_m) := (p) \bigotimes_{k=1}^m \theta_k^{\omega_{k_p}}.$$

respectively. Here, $0 \le \omega_k \le 1$ for any $k = 1, \ldots, m$, and the weights satisfy the condition $\sum_{k=1}^{m} w_k = 1$.

It can be seen from Definition 10 that

$$D\text{-}IFWGO_q(\theta_1,\ldots,\theta_m) = \left\langle \left(\prod_{k=1}^m \mu_{\theta_k}^{\omega_k}, 1-\prod_{k=1}^m (1-\nu_{\theta_k})^{\omega_k}\right); \prod_{k=1}^m r_{\theta_k}^{\omega_k} \right\rangle,$$

and

$$D\text{-}IFWGO_p(\theta_1,\ldots,\theta_m) = \left\langle \left(\prod_{k=1}^m \mu_{\theta_k}^{\omega_k}, 1-\prod_{i=1}^m (1-\nu_{\theta_k})^{\omega_k}\right); \sqrt{2}-\prod_{k=1}^m \left(\sqrt{2}-r_{\theta_k}\right)^{\omega_k} \right\rangle,$$

respectively.

4. DISC INTUITIONISTIC FUZZY CHOQUET INTEGRAL OPERATOR

This section begins with a review of the fuzzy measure and the Choquet integral, followed by the introduction of Choquet integral operators specifically designed for D-IFVs

Definition 11. [55] Let X be a universal set, and let 2^X represent the power set of X. A fuzzy measure is defined as a set function $\tau : 2^X \to [0,1]$ satisfying the following properties:

1) $\tau(\emptyset) = 0$ and $\tau(X) = 1$.

2) For any two subsets $A, B \subseteq X$, if $A \subseteq B$, then $\tau(A) \leq \tau(B)$.

Sugeno [55] proposed the concept of the λ -fuzzy measure, which incorporates an interaction index and criterion weighting. This measure is particularly useful for capturing synergies between criteria and offers more accurate solutions in complex decision-making contexts.

Definition 12. [55] Let X be a set, and consider a set function $\tau : 2^X \to [0,1]$ with $\lambda \ge -1$. The function τ is called a λ -fuzzy measure if it satisfies the following condition for any disjoint subsets A and B of X:

$$\tau(A \cup B) = \tau(A) + \tau(B) + \lambda \tau(A) \tau(B).$$

Definition 13. Let X be a finite set and let $\tau : 2^X \to [0,1]$ be a fuzzy measure. The discrete Choquet integral a function $f : X \to [0,1]$ with respect to τ is defined by

$$C_{\tau}(f(x_1),\ldots,f(x_m)) := \sum_{k=1}^{m} \left[\tau(E_{(k)}) - \tau(E_{(k+1)})\right] f(x_{(k)})$$

where the sequence $\{x_{(k)}\}_{k=0}^{m}$ indicates the indices permuted such that $0 := f(x_{(0)}) \le f(x_{(1)}) \le \dots \le f(x_{(m)}), E_{(k)} := \{x_{(k)}, x_{(k+1)}, \dots, x_{(m)}\}$ and $E_{(m+1)} = \emptyset$.

We now define some arithmetic aggregation operators.

Definition 14. Consider a collection of D-IFVs denoted as $\{\theta_k = \langle (\mu_{\theta_k}, \nu_{\phi_k}); r_{\theta_k} \rangle : k = 1, \ldots, m\}$. Disc intuitionistic fuzzy Choquet arithmetic integral operators with respect to τ are defined by

$$D\text{-}IFCAIO_q^{\tau}(\theta_1,\ldots,\theta_m) = (q) \bigoplus_{k=1}^m \left[\tau(F_{(k)}) - \tau(F_{(k+1)})\right]_q \theta_{(k)},$$

and

$$D\text{-}IFCAIO_p^{\tau}(\theta_1,\ldots,\theta_m) = (p) \bigoplus_{k=1}^m \left[\tau(F_{(k)}) - \tau(F_{(k+1)})\right]_p \theta_{(k)},$$

where the sequence $\{\theta_{(k)}\}_{k=0}^{m}$ indicates the indices permuted such that $\theta_{(1)} \leq \theta_{(2)} \leq \ldots \leq \theta_{(m)}$, $F_{(k)} := \{(k), (k+1), \ldots, (m)\}$ and $F_{(m+1)} = \emptyset$.

The following theorem formulates the given arithmetic aggregation operators.

Theorem 1. Consider a collection of D-IFVs denoted as $\{\theta_k = \langle (\mu_{\theta_k}, \nu_{\phi_k}); r_{\theta_k} \rangle : k = 1, \dots, m\}$. D-IFCAIO^{τ}_q $(\theta_1, \dots, \theta_m)$ and D-IFCAIO^{τ}_p $(\theta_1, \dots, \theta_m)$ are also a D-IFV and can be expressed by

$$D\text{-}IFCAIO_{q}^{\tau}(\theta_{1},\ldots,\theta_{m}) = \left\langle \left(1 - \prod_{k=1}^{m} (1 - \mu_{\theta_{(k)}})^{\tau(F_{(k)}) - \tau(F_{(k+1)})}, \prod_{k=1}^{m} \nu_{\theta_{(k)}}^{\tau(F_{(k)}) - \tau(F_{(k+1)})}\right); \prod_{k=1}^{m} r_{\theta_{(k)}}^{\tau(F_{(k)}) - \tau(F_{(k+1)})}\right\rangle$$

and

$$D\text{-}IFCAIO_{p}^{\tau}(\theta_{1},\ldots,\theta_{m}) = \left\langle \left(1 - \prod_{k=1}^{m} (1 - \mu_{\theta_{(k)}})^{\tau(F_{(k)}) - \tau(F_{(k+1)})}, \prod_{k=1}^{m} \nu_{\theta_{(k)}}^{\tau(F_{(k)}) - \tau(F_{(k+1)})}\right); \\ \sqrt{2} - \prod_{k=1}^{m} \left(\sqrt{2} - r_{\theta_{(k)}}\right)^{\tau(F_{(k)}) - \tau(F_{(k+1)})} \right\rangle$$

respectively where the sequence $\{\theta_{(k)}\}_{k=0}^{m}$ indicates the indices permuted such that $\theta_{(1)} \leq \theta_{(2)} \leq \ldots \leq \theta_{(m)}, F_{(k)} := \{(k), (k+1), \ldots, (m)\}$ and $F_{(m+1)} = \emptyset$.

Proof. It is clear that D-*IFCAIO*^{τ}_q($\theta_1, \ldots, \theta_m$) constitutes a D-IFV. The validity of the second part becomes apparent through the application of mathematical induction. When k = 2, with the help of Definition 8 we have

$$\left[\tau(F_{(1)}) - \tau(F_{(2)})\right]_{q} \theta_{(1)} = \left\langle \left(1 - (1 - \mu_{\theta_{(1)}})^{\tau(F_{(1)}) - \tau(F_{(2)})}, \nu_{\theta_{(1)}}^{\tau(F_{(1)}) - \tau(F_{(2)})}\right); \sqrt{2} \left(\frac{r_{\theta_{(1)}}}{\sqrt{2}}\right)^{\tau(F_{(1)}) - \tau(F_{(2)})} \right\rangle$$

and

$$\left[\tau(F_{(2)}) - \tau(F_{(3)})\right]_{q}\theta_{(2)} = \left\langle \left(1 - (1 - \mu_{\theta_{(2)}})^{\tau(F_{(2)}) - \tau(F_{(3)})}, \nu_{\theta_{(2)}}^{\tau(F_{(2)}) - \tau(F_{(3)})}\right); \sqrt{2} \left(\frac{r_{\theta_{(2)}}}{\sqrt{2}}\right)^{\tau(F_{(2)}) - \tau(F_{(3)})} \right\rangle.$$

Since $F_{(3)} = \emptyset$, we obtain

$$\begin{aligned} D\text{-}IFCAIO_{q}^{\tau}(\theta_{1},\theta_{2}) &= \left[\tau(F_{(2)}) - \tau(F_{(3)})\right]_{p}\theta_{(2)} \oplus_{p} \left[\tau(F_{(2)}) - \tau(F_{(3)})\right]_{p}\theta_{(2)} \\ &= \left\langle \left(1 - (1 - \mu_{\theta_{(1)}})^{\tau(F_{(1)}) - \tau(F_{(2)})}(1 - \mu_{\theta_{(2)}})^{\tau(F_{(2)}) - \tau(F_{(3)})}, \right. \\ &\left. \nu_{\theta_{(1)}}^{\tau(F_{(1)}) - \tau(F_{(2)})}\nu_{\theta_{(2)}}^{\tau(F_{(2)}) - \tau(F_{(3)})}\right); \sqrt{2} \left(\frac{\tau_{\theta_{(1)}}}{\sqrt{2}}\right)^{\tau(F_{(1)}) - \tau(F_{(2)})} \left(\frac{\tau_{\theta_{(2)}}}{\sqrt{2}}\right)^{\tau(F_{(2)}) - \tau(F_{(3)})} \right\rangle \\ &= \left\langle \left(1 - (1 - \mu_{\theta_{(1)}})^{\tau(F_{(1)}) - \tau(F_{(2)})}(1 - \mu_{\theta_{(2)}})^{\tau(F_{(2)}) - \tau(F_{(3)})}, \right. \\ &\left. \nu_{\theta_{(1)}}^{\tau(F_{(1)}) - \tau(F_{(2)})}\nu_{\theta_{(2)}}^{\tau(F_{(2)}) - \tau(F_{(3)})}\right); r_{\theta_{(1)}}^{\tau(F_{(1)}) - \tau(F_{(2)})}r_{\theta_{(2)}}^{\tau(F_{(2)}) - \tau(F_{(3)})} \right\rangle \end{aligned}$$

Now suppose that the expression

$$D\text{-}IFCAIO_{q}^{\tau}(\theta_{1},\ldots,\theta_{n}) = \left\langle \left(1 - \prod_{k=1}^{n} (1 - \mu_{\theta_{(k)}})^{\tau(F_{(k)}) - \tau(F_{(k+1)})}, \prod_{k=1}^{n} \nu_{\theta_{(k)}}^{\tau(F_{(k)}) - \tau(F_{(k+1)})}\right); \prod_{k=1}^{n} r_{\theta_{(k)}}^{\tau(F_{(k)}) - \tau(F_{(k+1)})}\right\rangle$$

is true for m = n. Then for m = n + 1 it is attain that

$$D\text{-}IFCAIO_{q}^{\tau}(\theta_{1},\ldots,\theta_{n},\theta_{n+1}) = D\text{-}IFCAIO_{p}^{\tau}(\theta_{1},\ldots,\theta_{n}) \oplus_{p} \left[\tau(F_{(n+1)}) - \tau(F_{(n+2)})\right]_{p}\theta_{(n+1)}$$

$$= \left\langle \left(1 - \prod_{k=1}^{n+1} (1 - \mu_{\theta_{k}})^{\tau(F_{(k)}) - \tau(F_{(k+1)})}, \prod_{k=1}^{n+1} \nu_{\theta_{k}}^{\tau(F_{(k)}) - \tau(F_{(k+1)})}\right);$$

$$\prod_{k=1}^{n+1} r_{\theta_{(k)}}^{\tau(F_{(k)}) - \tau(F_{(k+1)})}\right\rangle$$

Therefore, the proof is finished for D-*IFCAIO*^{τ}_q($\theta_1, \ldots, \theta_m$). In similar way, one can easily proof for D-*IFCAIO*^{τ}_p($\theta_1, \ldots, \theta_m$).

In the following we define some geometric aggregation operators.

Definition 15. Consider a collection of D-IFVs denoted as $\{\theta_k = \langle (\mu_{\theta_k}, \nu_{\phi_k}); r_{\theta_k} \rangle : k = 1, \ldots, m\}$. Disc intuitionistic fuzzy Choquet geometric integral operators with respect to τ are defined by

$$D\text{-}IFCGIO_q^{\tau}(\theta_1,\ldots,\theta_m) = (q)\bigotimes_{k=1}^m \theta_{(k)}^{\tau(F_{(k)})-\tau(F_{(k+1)})},$$

and

$$D\text{-}IFCGIO_{p}^{\tau}(\theta_{1},\ldots,\theta_{m}) = (p)\bigotimes_{k=1}^{m} \theta_{(k)}^{\tau(F_{(k)})-\tau(F_{(k+1)})},$$

where the sequence $\{\theta_{(k)}\}_{k=0}^m$ indicates the indices permuted such that $\theta_{(1)} \leq \theta_{(2)} \leq \ldots \leq \theta_{(m)}$, $F_{(k)} := \{(k), (k+1), \ldots, (m)\}$ and $F_{(m+1)} = \emptyset$.

The following theorem formulates the geometric aggregation operators.

Theorem 2. Consider a collection of D-IFVs denoted as $\{\theta_k = \langle (\mu_{\theta_k}, \nu_{\theta_k}); r_{\theta_k} \rangle : k = 1, ..., m\}$. D-IFCGIO^{τ}_q $(\theta_1, ..., \theta_m)$ and D-IFCGIO^{τ}_p $(\theta_1, ..., \theta_m)$ are also D-IFV and can be expressed by

$$D\text{-}IFCGIO_{q}^{\tau}(\theta_{1},\ldots,\theta_{m}) = \left\langle \left(\prod_{k=1}^{m} \mu_{\theta_{(k)}}^{\tau(F_{(k)})-\tau(F_{(k+1)})}, 1 - \prod_{k=1}^{m} (1 - \nu_{\theta_{(k)}})^{\tau(F_{(k)})-\tau(F_{(k+1)})} \right); \\ \prod_{k=1}^{n} r_{\theta_{(k)}}^{\tau(F_{(k)})-\tau(F_{(k+1)})} \right\rangle$$

and

$$D\text{-}IFCGIO_{p}^{\tau}(\theta_{1},\ldots,\theta_{m}) = \left\langle \left(\prod_{k=1}^{m} \mu_{\theta_{(k)}}^{\tau(F_{(k)})-\tau(F_{(k+1)})}, 1-\prod_{k=1}^{m} (1-\nu_{\theta_{(k)}})^{\tau(F_{(k)})-\tau(F_{(k+1)})}\right); \right. \\ \left. \sqrt{2} - \prod_{k=1}^{m} \left(\sqrt{2} - r_{\theta_{(k)}}\right)^{\tau(F_{(k)})-\tau(F_{(k+1)})} \right\rangle$$

respectively where the sequence $\{\theta_{(k)}\}_{k=0}^{m}$ indicates the indices permuted such that $\theta_{(1)} \leq \theta_{(2)} \leq \ldots \leq \theta_{(m)}, F_{(k)} := \{(k), (k+1), \ldots, (m)\}$ and $F_{(m+1)} = \emptyset$.

Proof. It can be proved similar to Theorem 1.

Remark 2. Let $\theta_1 = \langle (\mu_{\theta_1}, \nu_{\theta_1}); r_{\theta_1} \rangle$ and $\theta_2 = \langle (\mu_{\theta_2}, \nu_{\theta_2}); r_{\theta_2} \rangle$ such that $\mu_{\theta_1} \leq \mu_{\theta_2}, \nu_{\theta_1} \geq \nu_{\theta_2}, r_{\theta_1} \leq r_{\theta_2}$. $r_{\theta_1} \leq r_{\theta_2}$ or $\mu_{\theta_1} \geq \mu_{\theta_2}, \nu_{\theta_1} \leq \nu_{\theta_2}, r_{\theta_1} \geq r_{\theta_2}$. Assume that the weighted vector is $\omega = (\omega_1, \omega_2) = (0.3, 0.7)$ and a fuzzy measure τ is constructed by $\tau(\emptyset) = 0, \tau(\{1\}) = 0.22, \tau(\{2\}) = 0.58$, and $\tau(\{1, 2\}) = 1$. Then we have

$$D\text{-}IFWGO_{p}(\theta_{1},\theta_{2}) = \left\langle \left(\mu_{D}\text{-}IFWGO_{p}, \nu_{D}\text{-}IFWGO_{p} \right); r_{D}\text{-}IFWGO_{p} \right\rangle \\ = \left\langle \left(\mu_{\theta_{1}}^{0.3}\mu_{\theta_{2}}^{0.7}, 1 - (1 - \nu_{\theta_{1}})^{0.3}(1 - \nu_{\theta_{2}})^{0.7} \right); \sqrt{2} - (\sqrt{2} - \mu_{\theta_{1}})^{0.3}(\sqrt{2} - \mu_{\theta_{2}})^{0.7} \right\rangle$$

and

$$D\text{-}IFCGIO_p^{\tau}(\theta_1, \theta_2) = \left\langle \left(\mu_{D\text{-}IFCGIO_p^{\tau}}, \nu_{D\text{-}IFCGIO_p^{\tau}} \right); r_{D\text{-}IFCGIO_p^{\tau}} \right\rangle$$

where

$$\begin{split} \mu_{D\text{-}IFCGIO_{p}^{\tau}} &= \begin{cases} \mu_{\theta_{1}}^{0.42} \mu_{\theta_{2}}^{0.58} &, \text{ if } \mu_{\theta_{1}} \leq \mu_{\theta_{2}} \\ \mu_{\theta_{1}}^{0.78} \mu_{\theta_{2}}^{0.22} &, \text{ if } \mu_{\theta_{1}} \geq \mu_{\theta_{2}} \end{cases} \\ \nu_{D\text{-}IFCGIO_{p}^{\tau}} &= \begin{cases} 1 - (1 - \nu_{\theta_{1}})^{0.42} (1 - \nu_{\theta_{2}})^{0.58} &, \text{ if } \nu_{\theta_{1}} \geq \nu_{\theta_{2}} \\ 1 - (1 - \nu_{\theta_{1}})^{0.78} (1 - \nu_{\theta_{2}})^{0.22} &, \text{ if } \nu_{\theta_{1}} \leq \nu_{\theta_{2}} \end{cases} \end{split}$$

$$r_{D\text{-}IFCGIO_{p}^{\tau}} = \begin{cases} \sqrt{2} - (\sqrt{2} - r_{\theta_{1}})^{0.42} (\sqrt{2} - r_{\theta_{2}})^{0.58} &, \text{ if } r_{\theta_{1}} \le r_{\theta_{2}} \\ \sqrt{2} - (\sqrt{2} - r_{\theta_{1}})^{0.78} (\sqrt{2} - r_{\theta_{2}})^{0.22} &, \text{ if } r_{\theta_{1}} \ge r_{\theta_{2}} \end{cases}$$

Figure 7 shows a comparison of operators D-IFWGO_p and D-IFCGIO^{au}_p. It can be seen that these two operators have different characteristics.

Theorem 3. Consider a collection of D-IFVs denoted as $\{\theta_k = \langle (\mu_{\theta_k}, \nu_{\phi_k}); r_{\theta_k} \rangle : k = 1, ..., m\}$ and $\omega = (\omega_1, ..., \omega_m)$ is a weight vector. If τ is an additive measure such that $\tau([m]) = \sum_{k=1}^{m} \omega_k$, then

$$D\text{-}IFCAIO_q^{\tau}(\theta_1, \dots, \theta_m) = D\text{-}IFWAO_q(\theta_1, \dots, \theta_m),$$

$$D\text{-}IFCAIO_p^{\tau}(\theta_1, \dots, \theta_m) = D\text{-}IFWAO_p(\theta_1, \dots, \theta_m),$$

$$D\text{-}IFCGIO_q^{\tau}(\theta_1, \dots, \theta_m) = D\text{-}IFWGO_q(\theta_1, \dots, \theta_m),$$

and

$$D$$
-IFCGIO ^{τ} _p($\theta_1, \ldots, \theta_m$) = D -IFWGO_p($\theta_1, \ldots, \theta_m$).

Proof. The proof can be easily seen.

5. An Application to Solar Panels

This section introduces the CASPAS method following an analysis of the solar panel decision problem. The proposed method is then applied to a specific decision-making scenario.

5.1. An assessment of the types of solar panels. Various kinds of solar panels are available. Different types and technologies have been developed according to various usage needs and conditions. Main types and characteristics of solar panels are summarized Table 4.

The choice of solar panel type depends on factors such as efficiency requirements, available space, budget considerations, and the specific application or installation environment. Ongoing research and technological advancements continue to expand the range of options, making solar energy an increasingly versatile and accessible renewable energy source. In the solar panel selection problem to be examined in this article, the criteria to be used in the evaluation according to the opinion of energy systems experts are listed below:

- Efficiency (T_1) : The efficiency of solar panels refers to their ability to convert sunlight into usable electricity. It is a critical factor in evaluating the performance and economic viability of solar energy systems. Solar panel efficiency is typically expressed as a percentage and represents the ratio of the electrical power output to the solar energy input.
- Cost (T_2) : The cost of solar panels can vary widely depending on several factors, including the type of solar panel, the size of the system, the location of installation, and the specific manufacturer or brand. Prices have been consistently decreasing over the years due to technological advancements and increased market competition.
- **Durability** (T_3) : The durability of solar panels is a measure of their ability to withstand environmental conditions, maintain performance, and continue generating electricity over time without significant degradation. Durability is crucial for the long-term reliability and economic viability of a solar energy system.
- Installation Complexity (T_4) : The installation complexity of solar panels refers to the level of difficulty and intricacy involved in the process of setting up a solar panel system. Several factors contribute to the installation complexity, and they can affect the overall cost, time, and expertise required to successfully install solar panels.

Choosing the right solar panels involves a careful evaluation of these criteria in relation to the project's specific goals, budget, and location. Consulting with solar energy experts and conducting thorough research on available options can help in making an informed decision.



FIGURE 7. A comparison of Choquet geometric integral operator (right side) and weighted geometric aggregation operator (left side)

SHORT TITLE

Solar Panels	Characteristics
Monocrystalline Silicon Panel (P_1)	 Made using single-crystal silicon. Typically high efficiency (around 18-22%). Black cells with a characteristic look, often with cut corners. Generally more expensive than other types. Suitable for smaller spaces where high energy output is required [15, 45].
Polycrystalline Solar Panel (P_2)	 Produced using multi-crystal silicon. Usually slightly lower than monocrystalline panels (around 15-17%). Blue-hued cells with straight edges. More economical, generally lower priced. Preferred for large areas where cost-effectiveness is a priority [15, 45].
Thin-Film Solar Panel (P_3)	 Made from various materials (e.g., amorphous silicon, CdTe, CIGS). Generally lower efficiency (around 10-13%). Flexible structures, often with a darker color. Cheaper to manufacture but may produce less power per square meter. Suitable for large areas; flexible nature advantageous for certain specialized applications [34].
PERC Solar Panel (P_4)	 Additional technologies added to traditional monocrystalline or polycrystalline cells. Higher efficiency compared to standard panels (above 20%). Similar to monocrystalline or polycrystalline panels. The advanced manufacturing process can make PERC panels slightly more expensive than traditional panels, but the cost is often offset by their higher efficiency. Used for achieving high energy efficiency in limited spaces [21].
Bifacial Solar Panel (P_5)	 Cells can absorb light from both front and back surfaces. Bifacial panels are designed with solar cells that are capable of capturing sunlight on both sides. Capable of higher energy production due to dual-sided light collection (often generating 5-30% more energy compared to standard panels under optimal conditions). More effective when mounted over reflective surfaces. Higher upfront costs compared to traditional panels due to their advanced technology and materials. Suitable for a variety of installations, including commercial, residential, and utility-scale solar projects [22].

TABLE 4. Types of solar panels and their main characteristics

5.2. **CASPAS approach.** WASPAS was proposed by Zavadskas et al. [66]. WASPAS integrates the principles of both WSM and WPM using a method that applies weights to each. The method involves assigning weights to different criteria to represent their relative importance, normalizing the decision matrix, and then aggregating the scores for each alternative. WASPAS is known for its flexibility in incorporating both the weighted sum and the weighted product models, allowing decision-makers to adapt the method to different decision contexts.

In this subsection, CASPAS is proposed for disc intuitionistic fuzzy setting. This method enables an accurate modelling to be established in the decision-making process by taking into account the interaction between the criteria. We outline the procedural steps for employing the extended D-IF CASPAS method in the examination of MCGDM problems as follows:

- Step I: Form a MCGDM problem, where there are m alternatives denoted as $P = \{P_1, \ldots, P_m\}$. A group of n experts, identified as $E = \{E_1, \ldots, E_n\}$, assesses these alternatives using k criteria outlined in $T = \{T_1, \ldots, T_k\}$.
- Step II: Every expert assesses the alternatives by employing the linguistic terms provided in Table 5 which is each of the linguistic terms corresponds to a D-IFV. When there is a cost criterion present, the values for this criterion undergo a complement operation. Thus, a disc intuitionistic fuzzy group normalized decision matrix is created.

Linguistic term	D-IFV
Extremely High (EH)	$\langle (0.9, 0.1); 0.9 \rangle$
Very High (VH)	$\langle (0.8, 0.2); 0.8 \rangle$
High (H)	$\langle (0.7, 0.3); 0.7 \rangle$
Moderately High (MH)	$\langle (0.6, 0.4), 0.6 \rangle$
Medium (M)	$\langle (0.5, 0.5); 0.5 \rangle$
Moderately Low (ML)	$\langle (0.4, 0.6); 0.4 \rangle$
Low (L)	$\langle (0.3, 0.7); 0.3 \rangle$
Very Low (VL)	$\langle (0.2, 0.8); 0.2 \rangle$
Extremely Low (EL)	$\langle (0.1, 0.9); 0.1 \rangle$
Extremely Low (EL)	((0.1, 0.9), 0.1)

TABLE 5. Linguistic terms corresponding D-IFV

- Step III: A weight vector of the experts is determined according to the experience of the experts. Then the disc intuitionistic fuzzy group normalized decision matrix is aggregated by utilizing one of operators D-IFWAO_q^A, D-IFWAO_q^A, D-IFWGO_q^A, and D-IFWGO_q^A.
- Step IV: With the help of Table 5, the importance weights of the criteria are evaluated by the experts. The importance weights matrix of criteria is formed.
- Step V: By using weight vector of engineers, the importance weights matrix of criteria is aggregated similarly to Step III.
- Step VI: Compute the score function value S_j of each criterion T_j in aggregated importance weights matrix of criteria. Then normalize score of each criterion T_j using

(5.1)
$$\omega_j = \frac{S_j}{\sum\limits_{j=1}^k S_j}$$

Thus the weight vector of the criteria is obtained.

- Step VII: Construct a fuzzy measure τ to be able compute disc intuitionistic fuzzy Choquet operators after determining an interaction index λ and weight vector of the criteria.
- Step VIII: Calculate results of CSM_i based on D-IFCAIO^{τ}_q and D-IFCAIO^{τ}_p for each alternative P_i using aggregated decision matrix and one of the operators

(5.2)

$$CSM_{i}^{q} = \left\langle \left(1 - \prod_{j=1}^{k} (1 - \mu_{\theta_{T_{(j)}}})^{\tau(F_{(j)}) - \tau(F_{(j+1)})}, \prod_{j=1}^{k} \nu_{\theta_{T_{(j)}}}^{\tau(F_{(j)}) - \tau(F_{(j+1)})} \right); \prod_{j=1}^{k} r_{\theta_{T_{(j)}}}^{\tau(F_{(j)}) - \tau(F_{(j+1)})} \right\rangle$$
or

(5.3)

$$CSM_{i}^{p} = \left\langle \left(1 - \prod_{j=1}^{k} (1 - \mu_{\theta_{T_{(j)}}})^{\tau(F_{(j)}) - \tau(F_{(j+1)})}, \prod_{j=1}^{k} \nu_{\theta_{T_{(j)}}}^{\tau(F_{(j)}) - \tau(F_{(j+1)})} \right); \sqrt{2} - \prod_{j=1}^{k} \left(\sqrt{2} - r_{\theta_{T_{(j)}}} \right)^{\tau(F_{(j)}) - \tau(F_{(j+1)})} \right\rangle$$

where the sequence $\{\theta_{T_{(j)}}\}_{j=0}^k$ indicates the indices permuted such that $\theta_{T_{(1)}} \preceq \theta_{T_{(2)}} \preceq \ldots \preceq \theta_{T_{(k)}}, F_{(j)} := \{T_{(j)}, T_{(j+1)}, \ldots, T_{(k)}\}$ and $F_{(k+1)} = \emptyset$.

20

• Step IX: Calculate results of CPM_i based on D-IFCGIO^{τ} and D-IFCGIO^{τ} for each alternative P_i using aggregated decision matrix obtained in Step III and one of the operators

(5.4)

$$CPM_{i}^{q} = \left\langle \left(\prod_{j=1}^{k} \mu_{\theta_{T_{(j)}}}^{\tau(F_{(j)}) - \tau(F_{(j+1)})}, 1 - \prod_{j=1}^{k} (1 - \nu_{\theta_{T_{(j)}}})^{\tau(F_{(j)}) - \tau(F_{(j+1)})} \right); \prod_{j=1}^{k} r_{\theta_{T_{(j)}}}^{\tau(F_{(j)}) - \tau(F_{(j+1)})} \right\rangle$$

(5.5)

or

$$CPM_{i}^{p} = \left\langle \left(\prod_{j=1}^{k} \mu_{\theta_{T_{(j)}}}^{\tau(F_{(j)}) - \tau(F_{(j+1)})}, 1 - \prod_{j=1}^{k} (1 - \nu_{\theta_{T_{(j)}}})^{\tau(F_{(j)}) - \tau(F_{(j+1)})}\right); \sqrt{2} - \prod_{j=1}^{k} \left(\sqrt{2} - r_{\theta_{T_{(j)}}}\right)^{\tau(F_{(j)}) - \tau(F_{(j+1)})} \right\rangle$$

where the sequence $\{\theta_{T_{(j)}}\}_{j=0}^k$ indicates the indices permuted such that $\theta_{T_{(1)}} \leq \theta_{T_{(2)}} \leq \ldots \leq \theta_{T_{(k)}}, F_{(j)} := \{T_{(j)}, T_{(j+1)}, \ldots, T_{(k)}\}$ and $F_{(k+1)} = \emptyset$.

• Step X: Select the threshold number $\varepsilon \in [0, 1]$ and find significance degree of each P_i by utilizing one of formulas

(5.6)
$$SD_i^q = \varepsilon CSM_i^q \oplus_q (1-\varepsilon)CPM_i^q$$

or

$$(5.7) SD_i^p = \varepsilon CSM_i^p \oplus_p (1-\varepsilon)CPM_i^p$$

for i = 1, ..., m.

- Step XI: Compute the score values of the significance degrees $SD_i (i = 1, ..., m)$.
- Step XII: The alternatives are ranked with the one with the highest score being the best alternative. If two alternatives have equal score values, the values of their accuracy functions may be taken into consideration.

5.3. A numerical example. A university is planning to transition its energy sources to renewable ones by installing solar panels across its campus. This initiative aims to reduce carbon footprint, cut energy costs, and serve as an educational resource for students studying renewable energy technologies. The decision involves multiple criteria and alternatives, making it a complex MCGDM problem.

- Step I: In the decision-making process, the university administration has appointed three engineers: two E_1, E_2 possess PhD degrees, while the third E_3 has a master's degree. The evaluation process of solar panels includes five solar panels: monocrystalline silicon panel (P_1) , polycrystalline solar panel (P_2) , thin-film solar panel (P_3) , PERC solar panel (P_4) , bifacial solar panel (P_5) . There are four criteria: efficiency (T_1) , cost (T_2) , durability (T_3) , installation complexity (T_4) for evaluating these panel types.
- Step II: Three engineers assess the alternatives based on the linguistic terms presented in Table 5. The evaluation results are shown in Table 6. Then, by employing D-IFVs associated with each linguistic term, disc intuitionistic fuzzy group decision matrix can be constructed. Given that T_2 and T_4 represent cost criteria, we take the complement of these values. Consequently, this process results in the creation of the disc intuitionistic fuzzy group normalized decision matrix, as depicted in Table 7.
- Step III: The weight vector of experts according to their experience is $\Omega = (\Omega_1, \Omega_2, \Omega_3) = (0.4, 0.4, 0.2)$. After the disc intuitionistic fuzzy group normalized decision matrix is aggregated by using operator *D*-*IFWAO*^{*A*}_{*q*}, we obtain aggregated disc intuitionistic fuzzy decision matrix shown in Table 8.
- Step IV: The importance weights of the criteria are assessed by engineers with assistance from Table 5, leading to the importance weights matrix of criteria listed in Table 9.

Experts	Solar Panels	T_1	T_2	T_3	T_4
	P_1	EH	MH	\mathbf{EH}	Μ
	P_2	VH	Μ	\mathbf{VH}	Μ
E_1	P_3	MH	\mathbf{L}	ML	ML
	P_4	VH	\mathbf{MH}	Η	Μ
	P_5	H	MH	VH	MH
	P_1	VH	Η	VH	MH
	P_2	MH	Μ	Η	MH
E_2	P_3	M	Μ	Μ	ML
	P_4	VH	\mathbf{MH}	VH	Μ
	P_5	VH	Η	\mathbf{VH}	Μ
	P_1	EH	VH	Н	ML
	P_2	H	ML	MH	Μ
E_3	P_3	ML	VL	ML	VL
	P_4	H	Η	\mathbf{MH}	ML
	P_5	H	\mathbf{VH}	Η	Η

TABLE 6. Group decision matrix with linguistic term

Engineers	Solar Panels	T_1	T_2	T_3	T_4
	P_1	$\langle (0.9, 0.1); 0.9 \rangle$	$\langle (0.4, 0.6); 0.6 \rangle$	$\langle (0.9, 0.1); 0.9 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$
	P_2	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$
E_1	P_3	$\langle (0.6, 0.4); 0.6 \rangle$	$\langle (0.7, 0.3); 0.3 \rangle$	$\langle (0.4, 0.6); 0.4 \rangle$	$\langle (0.6, 0.4); 0.4 \rangle$
	P_4	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.4, 0.6); 0.6 \rangle$	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$
	P_5	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.4, 0.6); 0.6 \rangle$	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.4, 0.6); 0.6 \rangle$
	P_1	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.3, 0.7); 0.7 \rangle$	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.4, 0.6); 0.6 \rangle$
	P_2	$\langle (0.6, 0.4); 0.6 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.4, 0.6); 0.6 \rangle$
E_2	P_3	$\langle (0.5, 0.5); 0.5 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$	$\langle (0.6, 0.4); 0.4 \rangle$
	P_4	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.4, 0.6); 0.6 \rangle$	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$
	P_5	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.3, 0.7); 0.7 \rangle$	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$
	P_1	$\langle (0.9, 0.1); 0.9 \rangle$	$\langle (0.2, 0.8); 0.8 \rangle$	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.6, 0.4); 0.4 \rangle$
	P_2	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.6, 0.4); 0.4 \rangle$	$\langle (0.6, 0.4); 0.6 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$
E_3	P_3	$\langle (0.4, 0.6); 0.4 \rangle$	$\langle (0.8, 0.2); 0.2 \rangle$	$\langle (0.4, 0.6); 0.4 \rangle$	$\langle (0.8, 0.2); 0.2 \rangle$
	P_4	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.3, 0.7); 0.7 \rangle$	$\langle (0.6, 0.4); 0.6 \rangle$	$\langle (0.6, 0.4); 0.4 \rangle$
	P_5	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.2, 0.8); 0.8 \rangle$	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.3, 0.7); 0.7 \rangle$
	$\mathbf{D}_{1} = \mathbf{z} = \mathbf{z} = \mathbf{D}_{1}^{*}$	· · · · · · · · · · · · · · · · · · ·		1. 1 1	

TABLE 7. Disc intuitionistic fuzzy group normalized decision matrix

Solar Panels	T_1	T_2	T_3	T_4
P_1	$\langle (0.87, 0.13); 0.86 \rangle$	$\langle (0.32, 0.68); 0.68 \rangle$	$\langle (0.84, 0.16); 0.82 \rangle$	$\langle (0.49, 0.51); 0.51 \rangle$
P_2	$\langle (0.71, 0.29); 0.69 \rangle$	$\langle (0.52, 0.48); 0.48 \rangle$	$\langle (0.73, 0.27); 0.72 \rangle$	$\langle (0.46, 0.54); 0.54 \rangle$
P_3	$\langle (0.53, 0.47); 0.51 \rangle$	$\langle (0.66, 0.34); 0.34 \rangle$	$\langle (0.44, 0.56); 0.44 \rangle$	$\langle (0.65, 0.35); 0.35 \rangle$
P_4	$\langle (0.78, 0.22); 0.78 \rangle$	$\langle (0.38, 0.62); 0.62 \rangle$	$\langle (0.73, 0.27); 0.72 \rangle$	$\langle (0.52, 0.48); 0.48 \rangle$
P_5	$\langle (0.75, 0.25); 0.74 \rangle$	$\langle (0.32, 0.68); 0.68 \rangle$	$\langle (0.78, 0.22); 0.78 \rangle$	$\langle (0.42, 0.58); 0.58 \rangle$

TABLE 8. Aggregated disc intuitionistic fuzzy decision matrix

• Step V: By utilizing weight vector of experts $\Omega = (\Omega_1, \Omega_2, \Omega_3) = (0.4, 0.4, 0.2)$ and operator $DIFWA_p^A$, the importance weights matrix of criteria given in Table 9 is aggregated. Aggregated importance weights matrix of criteria is also shown in Table 9.

SHORT TITLE

Criteria	E_1	E_2	E_3	Aggregated results
T_1	$\langle (0.8, 0.2); 0.8 \rangle$	$\langle (0.7, 0.3); 0.7 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$	$\langle (0.72, 0.28); 0.69 \rangle$
T_2	$\langle (0.6, 0.4); 0.6 \rangle$	$\langle (0.6, 0.4); 0.6 \rangle$	$\langle (0.4, 0.6); 0.4 \rangle$	$\langle (0.57, 0.43); 0.55 \rangle$
T_3	$\langle (0.6, 0.4); 0.6 \rangle$	$\langle (0.5, 0.5); 0.5 \rangle$	$\langle (0.3, 0.7); 0.3 \rangle$	$\langle (0.51, 0.49); 0.49 \rangle$
T_4	$\langle (0.5, 0.5); 0.5 \rangle$	$\langle (0.4, 0.6); 0.4 \rangle$	$\langle (0.2, 0.8); 0.2 \rangle$	$\langle (0.41, 0.59); 0.38 \rangle$
TABLE 9	. The importance	e weights matrix	of criteria and it	s aggregated results

• Step VI: After computing the score value of each criterion T_j via S introduced in Definition 6 for $\xi = 0.8$, we normalize score of each criterion T_j using Eq. 5.1. Therefore the weight vector of the criteria given in is $\omega = (\omega_1, \omega_2, \omega_3, \omega_4) = (0.326, 0.258, 0.232, 0.184)$.

• Step VII: A fuzzy measure employing the λ -fuzzy methodology can be formulated using Takahagi's algorithm. To specify the construction, let's assign λ a value of 0.5 and consider weights $\omega = (\omega_1, \omega_2, \omega_3, \omega_4) = (0.326, 0.258, 0.232, 0.184)$. This yields the fuzzy measure τ , depicted in Table 10.

$\tau(\varnothing) = 0$	$\tau(\{T_1\}) = 0.282631$	$\tau(\{T_2\}) = 0.220555$
$\tau(\{T_3\}) = 0.197269$	$\tau(\{T_4\}) = 0.154918$	$\tau(\{T_1, T_2\}) = 0.534354$
$\tau(\{T_1, T_3\}) = 0.507777$	$\tau(\{T_1, T_4\}) = 0.459442$	$\tau(\{T_2, T_3\}) = 0.439578$
$\tau(\{T_2, T_4\}) = 0.392557$	$\tau(\{T_3, T_4\}) = 0.367467$	$\tau(\{T_1, T_2, T_3\}) = 0.784329$
$\tau(\{T_1, T_2, T_4\}) = 0.730663$	$\tau(\{T_1, T_3, T_4\}) = 0.702027$	$\tau(\{T_2, T_3, T_4\}) = 0.628546$
$\tau(\{T_1, T_2, T_3, T_4\}) = 1$		
	$\Pi_{\rm LDTD} = 10$ Å f	

TABLE 10. A fuzzy measure

• Step VIII-IX: By computing CSM_i and CPM_i for each P_i , the results of CSM_i and CPM_i are obtained and are summarized in Table 11.

Solar Panels	CSM^q	CSM^p	CPM^q	CPM^p		
P_1	$\langle (0.71, 0.29); 0.72 \rangle$	$\langle (0.71, 0.29); 0.74 \rangle$	$\langle (0.57, 0.43); 0.72 \rangle$	$\langle (0.57, 0.43); 0.74 \rangle$		
P_2	$\langle (0.62, 0.38); 0.59 \rangle$	$\langle (0.62, 0.38); 0.62 \rangle$	$\langle (0.59, 0.40); 0.59 \rangle$	$\langle (0.59, 0.40); 0.62 \rangle$		
P_3	$\langle (0.56, 0.44); 0.42 \rangle$	$\langle (0.56, 0.44); 0.43 \rangle$	$\langle (0.55, 0.45); 0.42 \rangle$	$\langle (0.55, 0.45); 0.43 \rangle$		
P_4	$\langle (0.63, 0.37); 0.65 \rangle$	$\langle (0.63, 0.37); 0.67 \rangle$	$\langle (0.57, 0.43); 0.65 \rangle$	$\langle (0.57, 0.43); 0.67 \rangle$		
P_5	$\langle (0.61, 0.39); 0.69 \rangle$	$\langle (0.61, 0.39); 0.70 \rangle$	$\langle (0.52, 0.48); 0.69 \rangle$	$\langle (0.52, 0.48); 0.70 \rangle$		
L	TABLE 11. The results of CSM_i and CPM_i					

• Step X: If we select $\varepsilon = 0.3$, significance degrees SD_i of each P_i are listed in Table 12.

Solar Panels	SD^q	SD^p
P_1	$\langle (0.619, 0.380); 0.717 \rangle$	$\langle (0.619, 0.380); 0.741 \rangle$
P_2	$\langle (0.601, 0.399); 0.596 \rangle$	$\langle (0.601, 0.399); 0.620 \rangle$
P_3	$\langle (0.553, 0.447); 0.420 \rangle$	$\langle (0.553, 0.447); 0.429 \rangle$
P_4	$\langle (0.593, 0.407); 0.651 \rangle$	$\langle (0.593, 0.407); 0.668 \rangle$
P_5	$\langle (0.553, 0.447); 0.695 \rangle$	$\langle (0.553, 0.447); 0.702 \rangle$

TABLE 12. The results of significance degrees SD_i

- Step XI: Score values of the significance degrees $SD_i(i = 1, ..., m)$ are attained by utilizing S introduced in Definition 6 for $\xi = 0.8$. The results are shown Table 13.
- Step XII: For score values of significance degree SD^q , the order is established as $P_1 > P_4 > P_2 > P_5 > P_3$, whereas for score values associated with significance degree SD^p , the sequence is determined as $P_1 > P_2 > P_4 > P_5 > P_3$. Both SD^q and SD^p

Solar Panels	$\mathcal{S}(SD^q)$	$\mathcal{S}(SD^p)$
P_1	0.596	0.600
P_2	0.565	0.569
P_3	0.501	0.502
P_4	0.567	0.568
P_5	0.540	0.541

TABLE 13. The score values of significance degrees SD^q and SD^p

determine P_1 as the most favorable solar panel. Monocrystalline silicon panel emerges as the best alternative as a result of the decision-making process. The difference in the ranking of P_2 and P_4 solar panels is a result of the interactions between the criteria and the radius change due to the circular structure

6. Assessment of performance

In this section, we conduct sensitivity analysis and validity assessment to observe how variations in different scenarios of the solar panel selection problem may impact the ranking of alternatives.

6.1. Sensitivity analysis. Sensitivity analysis is a technique used to assess how the variation (or uncertainty) in the output of a mathematical model or system can be attributed to variations in the inputs. The main goal of sensitivity analysis is to understand which input parameters have the most significant impact on the output of the model. This analysis helps in identifying critical factors and understanding the robustness of the model. In this subsection, we concentrate on an analysis of the effect of the variation of the parameters (ε and ξ) considered in the proposed D-IF CASPAS method on the ranking of alternatives. The study of these parameters reveals the impact of different aspects of D-IFVs and the CASPAS method on the ranking results of the solar panel selection problem.

• Analysis of parameter ε : In Step X of the proposed CASPAS method given in Subsection 5.2, the CSM^q , CSM^p , CPM^q , and CPM^q values are calculated as D-IFVs by adjusting the threshold number $\varepsilon \in [0,1]$. ε affects the results of the significance degrees SD^q and SD^p obtained from CSM^q , CSM^p , CPM^q , and CPM^q . As ε grows larger, the impact of CSM^q on SD^q becomes more pronounced, whereas the influence of CPM^q diminishes. A similar scenario applies to SD^p as well. In Step X of the proposed CASPAS method, an analysis is applied by varying ε between 0.1 and 0.9 to examine the effect of ε on the results. Figure 8 shows the results of the ranking of the ε parameter. These results are calculated according to the score values $\xi = 0.8$. The ranking results can be respectively expressed as follows. For the significance degree SD^q , two different rankings are obtained. For $\varepsilon = 0.1$ and 0.2 we find the order $P_1 > P_2 > P_4 > P_5 > P_3$, while by varying ε from 0.3 to 0.7 we get the order $P_1 > P_4 > P_2 > P_5 > P_3$. If ε equals 0.8 and 0.9, $P_1 > P_4 > P_5 > P_2 > P_3$ is attained. For the significance degree SD^p , we get also two different rankings. When ε changes from 0.1 and 0.3, the ordering of alternatives remains stable as $P_1 > P_2 > P_4 > P_5 > P_3$. Conversely, the ranking shifts to the ranking $P_1 > P_4 > P_2 > P_5 > P_3$ as ε increases from 0.4 to 0.9. Upon examining the ranking of alternatives, it becomes evident that P_3 consistently emerges as the least favorable alternative and P_1 also appears to be the most favorable alternative across all parameters ε ranging from 0.1 to 0.9 for both SD^q and SD^p . Nonetheless, when the results are analysed, the order of the second, third and fourth ranked alternatives varies according to the epsilon parameter. This is mainly due to the adjustment of the effect of the CSM and CPM in the proposed CASPAS method on the significance SD with the ε parameter. The significant role of the radius in D-IFSs and D-IFVs in altering these rankings is noteworthy. With adjustments in the radius degree, there can be shifts in the alternatives because the score values are affected. Consequently, the use



of D-IFSs and D-IFVs in decision-making leads to more accurate modeling compared to assigning fixed values to criteria, thanks to the adaptable nature of their circular framework.

FIGURE 8. Sensitivity analysis across ε values ranging from 0.1 to 0.9

Analysis of parameter ξ : In Step XI of the CASPAS method proposed in Sub-section 5.2, the score function proposed in Definition 6 is used to calculate the score values of the significance degrees SD^q and SD^p . The most important advantage of this score function is that the effect of the radius degree varies depending on the decision maker's choice of the parameter ξ between 0 and 1. Thus, the effect of the radius in D-IFVs can be adjusted and its effect on the ranking can be analyzed. As ξ approaches 1, the degree of radius becomes essentially irrelevant in the decision-making process. To assess the impact of the parameter ξ as detailed in Step XI of the proposed CASPAS methodology, we perform an analysis by varying ξ from 0.1 to 0.9 for the score values of the significance degrees SD^q and SD^p . The radar chart for the variation of the parameter ξ between 0.1 and 0.9 is shown in Figure 9. The findings obtained from Figure 9 can be summarised as follows. According to significance degree SD^q , there are four different ranking. The consistent ranking of alternatives is $P_1 > P_5 > P_4 > P_2 > P_3$ from $\xi = 0.1$ and 0.4. When ξ equals 0.5, alternatives are ordered as $P_1 > P_4 > P_5 > P_2 > P_3$. When ξ varies from 0.6 and 0.8, the ranking of alternatives is obtained as $P_1 > P_4 > P_2 > P_5 > P_3$. Lastly the ranking $P_1 > P_2 > P_4 > P_5 > P_3$ is obtained for $\xi = 0.9$. According to significance degree SD^p , we reach four different rankings. If ξ varies between 0.1 and 0.3, we attain the ranking of alternatives as $P_1 > P_5 > P_4 > P_2 > P_3$. For $\xi = 0.4$ and 0.5 the ranking is $P_1 > P_4 > P_5 > P_2 > P_3$. When ξ is set to 0.6 and 0.7, the consistent ranking of alternatives is obtained as $P_1 > P_4 > P_2 > P_5 > P_3$. Lastly we get ranking $P_1 > P_2 > P_4 > P_5 > P_3$ for $\xi = 0.8$ and 0.9. When the ranking of the alternatives is analysed, it is seen that P_3 is the worst alternative and P_1 is the best alternative for each parameter ξ from 0.1 and 0.9 according to SD^q and SD^p . However, the ranking of the second, third and fourth alternatives changes. For instance, according to SD^{q} when $\xi = 0.1$, P_5 is the second best alternative, while for $\xi = 0.5$, P_4 becomes the second best alternative. This variation results from the circular structures of D-IFSs and D-IFVs. Since the parameter ξ in the score function is adjustable, the impact of the radius degree on rankings also varies. This causes the order of alternatives to change. Additionally, one of the conclusions that can be drawn from this analysis is the potential for different rankings when the degrees of membership and non-membership are modeled with a radius degree instead of being precisely assigned. This highlights the significance of decision-making processes involving D-IFSs constructed with circular structures.



FIGURE 9. Sensitivity analysis across a range of ξ values spanning from 0.1 to 0.9.

6.2. Validity analysis. Validity analysis in MCDM and MCGDM involves assessing the reliability, relevance, and effectiveness of the decision-making process and the associated outcomes. The efficacy of MCDM method is shaped by various factors such as the interplay between alternatives, the consistency of criteria, and impartial evaluations among decision-makers. Wang and Triantaphyllou [63] introduced three evaluative criteria to validate the presented method in MCDM. These three test conditions are systematically applied to the proposed method as follow:

- **I.Condition:** When replacing a non-optimal alternative with a worse one without altering the significance of each criterion, the best alternative should remain unchanged in a useful MCDM method.
- **II.Condition:** An effective MCDM method should exhibit the transitive property. In other words, if alternative X is favored over alternative Y, and alternative Y is chosen over alternative Z, then it follows that alternative X should be selected over alternative Z.
- **III.Condition:** If a decision-making problem is broken down into multiple sub-problems, the order of the subproblems through the decision-making approach should align with the ranking outcome of the original problem.

To assess the validity of the proposed CASPAS method, the following examinations are carried out.

• **I.Condition:** In order to fulfil this condition, the non-optimal alternative P_3 is substituted with an arbitrary worse option P'_3 , and the evaluation of P'_3 is presented in Table 14 based on the linguistic terms presented in Table 5.

SHORT TITLE

Engineers	T_1	T_2	T_3	T_4
E_1	М	L	L	ML
E_2	ML	Μ	ML	Μ
E_3	L	\mathbf{EL}	L	VL

TABLE 14. The evaluation of P'_3 based on the linguistic terms

We can now explore if there are any shifts in the top-ranked alternative for $\varepsilon = 0.3$ and $\xi = 0.5$. Utilizing the proposed CASPAS method, the hierarchy of alternatives is determined as $P_1 > P_5 > P_4 > P_2 > P'_3$, with P_1 retaining its position as the best alternative. Therefore, I.Conditon satisfies for the proposed method.

• II. and III. Condition: The same solar panel problem is divided into four subproblems, which include $\{P_1, P_2, P_3, P_4\}$, $\{P_1, P_2, P_4, P_5\}$, $\{P_2, P_3, P_4, P_5\}$, and $\{P_1, P_2, P_3, P_5\}$, respectively. By using the proposed WASPAS method to solve these sub-problems, the ranking orders of these sub-problems can be obtained as: $P_1 > P_4 > P_2 > P_3$, $P_1 > P_5 > P_4 > P_2$, $P_5 > P_4 > P_2 > P_3$, and $P_1 > P_5 > P_2 > P_3$. Upon merging the results of these sub-problems, the comprehensive result is derived as: $P_1 > P_5 > P_4 > P_2 > P_3$ which is consistent with with the result of the same problem. Therefore, the proposed CASPAS method is confirmed for I.Condition and II.Condition.

7. Comparative Analysis

The section commences with a comparison between weighted aggregation and Choquet integral operators, then proceeds to conduct a thorough analysis of the proposed CASPAS method in relation to the existing C-IF-TOPSIS and C-IF-VIKOR methods. Ultimately, it assesses the advantages of the newly proposed method over these existing approaches.

7.1. A comparison of weighted aggregation operators with Choquet integral operators. In this subsection, we discuss the comparison between two widely used aggregation operators: weighted sum and Choquet integrals. Our aim is to determine the optimal alternative among a variety of alternatives, each assessed based on distinct criteria. Despite the widespread popularity and simplicity of the weighted sum approach, the practical application of the Choquet integral remains challenging. However, theoretically, the Choquet integral promises superior outcomes aligning more closely with a decision maker's preferences. Meyer and Pirlot [36] have demonstrated that the Choquet integral can consider significantly more preferences compared to the weighted arithmetic mean, with a notable difference especially evident when the number of criteria is high. Furthermore, Lust [35] has shown that as the number of criteria increases, the use of the Choquet integral instead of the weighted arithmetic mean increases the likelihood of achieving a more optimal ranking. When using a weighted arithmetic mean, the interaction between decision maker priorities and criteria cannot be assessed with sufficient precision. In order to make a comparative analysis of the results of weighted operators and Choquet integral operators, in the proposed CASPAS method, in Steps VIII and IX, the ignificance degrees SD are calculated using WSM and WPM defined by weighted operators instead of CSM and CPM defined by Choquet integral operator. WSM and WPM are defined as follows. For each alternative $P_i(i = 1, ..., m)$, we have

(7.1)
$$WSM_{i}^{q} = \left\langle \left(1 - \prod_{j=1}^{k} (1 - \mu_{\theta_{T_{j}}})^{\omega_{j}}, \prod_{j=1}^{k} \nu_{\theta_{T_{j}}}^{\omega_{j}}\right); \prod_{j=1}^{k} r_{\theta_{T_{j}}}^{\omega_{j}}\right\rangle$$

or

(7.2)
$$WSM_{i}^{p} = \left\langle \left(1 - \prod_{j=1}^{k} (1 - \mu_{\theta_{T_{j}}})^{\omega_{j}}, \prod_{j=1}^{k} \nu_{\theta_{T_{j}}}^{\omega_{j}} \right); \sqrt{2} - \prod_{k=1}^{m} \left(\sqrt{2} - r_{\theta_{T_{j}}} \right)^{\omega_{j}} \right\rangle$$

and

(7.3)
$$WPM_{i}^{q} = \left\langle \left(\prod_{j=1}^{k} \mu_{\theta_{T_{j}}}^{\omega_{j}}, 1 - \prod_{j=1}^{k} (1 - \nu_{\theta_{T_{j}}})^{\omega_{j}}\right); \prod_{j=1}^{k} r_{\theta_{T_{j}}}^{\omega_{j}} \right\rangle$$

or

(7.4)
$$WPM_{i}^{p} = \left\langle \left(\prod_{j=1}^{k} \mu_{\theta_{T_{j}}}^{\omega_{j}}, 1 - \prod_{j=1}^{k} (1 - \nu_{\theta_{T_{j}}})^{\omega_{j}}\right); \sqrt{2} - \prod_{k=1}^{m} \left(\sqrt{2} - r_{\theta_{T_{j}}}\right)^{\omega_{j}} \right\rangle$$

The results of the significance degrees SD based on WSM and WPM calculated for $\varepsilon = 0.1$ are listed in Table 15 according to q and p. Table 15 also shows the score values which are obtained by utilizing S introduced in Definition 6 for $\xi = 0.9$.

Solar Panels	SD^q	$\mathcal{S}(SD^q)$	SD^p	$\mathcal{S}(SD^p)$
P_1	$\langle (0.60, 0.39); 0.73 \rangle$	0.596	$\langle (0.60, 0.39); 0.75 \rangle$	0.598
P_2	$\langle (0.62, 0.38); 0.61 \rangle$	0.601	$\langle (0.62, 0.38); 0.62 \rangle$	0.602
P_3	$\langle (0.56, 0.43); 0.42 \rangle$	0.538	$\langle (0.56, 0.43); 0.42 \rangle$	0.538
P_4	$\langle (0.60, 0.40); 0.66 \rangle$	0.587	$\langle (0.60, 0.40); 0.68 \rangle$	0.588
P_5	$\langle (0.56, 0.44); 0.70 \rangle$	0.587	$\langle (0.56, 0.44); 0.71 \rangle$	0.554

TABLE 15. The results of significance degrees SD based on WSM and WPM

When the results in Table 15 are analysed, it is seen that the ranking $P_2 > P_1 > P_4 > P_5 > P_3$ is obtained according to SD based on WSM and WPM. On the other hand, according to the Choquet integral operator based CASPAS method proposed in Subsection 5.2, we get the ranking $P_1 > P_2 > P_4 > P_3 > P_5$ by taking $\varepsilon = 0.1$ and $\xi = 0.9$. As can be seen from the results, according to the weighted aggregation operators, the most favorable solar panel is P_2 and the most unfavorable solar panel is P3, while according to the Choquet integral operator, the most favorable solar panel is P_1 and the most unfavorable solar panel is P_5 . The main reason for this difference is that Choquet integral operators take into account the interaction between criteria. Since weighted aggregation operators do not consider the interaction between criteria, more precise results can be obtained with Choquet integral operators in decision-making applications.

7.2. A comparison with C-IF TOPSIS. In this section, we employ C-IF TOSPIS to use in the solar panel selection, conducting a thorough comparative analysis with our proposed approach. TOPSIS (Technique for Order of Preference by Similarity to Ideal Solution) is a MCDM used to determine the best alternative from a set of options. It aims to identify the alternative that is closest to the ideal solution based on a set of criteria. TOPSIS, pioneered by Hwang and Yoon [24] as a method in MCDM, was crafted with the explicit goal of addressing shortcomings in prevailing decision-making methodologies. Its purpose is to offer a more intuitive and pragmatic framework, mitigating the subjectivity inherent in criterion weighting and enhancing result interpretation. By furnishing decision-makers with a structured and quantitative methodology, TOPSIS empowers them to systematically assess and rank alternatives across various criteria. Chen [12] proposed evolved C-IF Minkowski distance measures for C-IFSs and C-IFVs, which are defined by

(7.5)
$$\mathcal{D}^{\beta}_{(3)}(\theta,\kappa) = \frac{1}{2} \left\{ \frac{1}{\sqrt{2}} |r_{\theta} - r_{\kappa}| + \left[\frac{1}{2} \left(|\mu_{\theta} - \mu_{\kappa}|^{\beta} + |\nu_{\theta} - \nu_{\kappa}|^{\beta} \right) \right]^{\frac{1}{\beta}} \right\},$$

for two C-IFVs $\theta = \langle (\mu_{\theta}, \nu_{\theta}); r_{\theta} \rangle$ and $\kappa = \langle (\mu_{\kappa}, \nu_{\kappa}); r_{\kappa} \rangle$. Here, β is a positive integer and represents the metric parameter. Later, Chen [12] introduced a C-IF TOPSIS methodology

28

based on C-IF Minkowski distance measures and applied it to a site selection issue of largescale epidemic hospitals. To put it briefly, in this C-IF TOPSIS method, after the aggregation process, the displaced ideal and anti-ideal ratings are derived with

(7.6)
$$\theta_{P_*} = \left\langle \max_{1 \le i \le m} \mu_{\theta_{P_i}}, \min_{1 \le i \le m} \nu_{\theta_{P_i}}; \max_{1 \le i \le m} r_{\theta_{P_i}} \right\rangle$$

and

(7.7)
$$\theta_{P_{\neg}} = \left\langle \min_{1 \le i \le m} \mu_{\theta_{P_i}}, \max_{1 \le i \le m} \nu_{\theta_{P_i}}; \max_{1 \le i \le m} r_{\theta_{P_i}} \right\rangle,$$

respectively. Then the relative closeness coefficient is calculated with

(7.8)
$$\mathcal{R}_{(3)}^{\beta*}(P_i) = \frac{\mathcal{D}_{(3)}^{\beta}(\theta_{P_i}, \theta_{P_{\gamma}})}{\mathcal{D}_{(3)}^{\beta}(\theta_{P_i}, \theta_{P_{\gamma}}) + \mathcal{D}_{(3)}^{\beta}(\theta_{P_i}, \theta_{P_{\gamma}})}$$

for i = 1, ..., m. The alternative with the highest relative closeness coefficient is the best alternative.

Let us apply C-IF-TOPSIS to solve the solar panel selection problem discussed in Sub-section 5.3. If the disc intuitionistic decision matrix given in Table 8 is aggregated with operator D-IFWGO_q with the weight vector w = (0.326, 0.258, 0.232, 0.181), the results listed in Table 16 are obtained with the displaced ideal and anti-ideal ratings.

	D - $IFWGO_q$
P_1	$\langle (0.59, 0.41); 0.73 \rangle$
P_2	$\langle (0.61, 0.39); 0.61 \rangle$
P_3	$\langle (0.56, 0.44); 0.42 \rangle$
P_4	$\langle (0.59, 0.41); 0.66 \rangle$
P_5	$\langle (0.55, 0.45); 0.70 \rangle$
P_*	$\langle (0.61, 0.39); 0.73 \rangle$
P_{\neg}	$\langle (0.55, 0.45); 0.73 \rangle$

TABLE 16. Aggregation results with the displaced ideal and anti-ideal ratings

After the calculation for $\beta = 3$, we find the relative closeness coefficients $\mathcal{R}_{(3)}^{\beta*}(P_1) = 0.667, \mathcal{R}_{(3)}^{\beta*}(P_2) = 0.631, \mathcal{R}_{(3)}^{\beta*}(P_3) = 0.459, \mathcal{R}_{(3)}^{\beta*}(P_4) = 0.562$, and $\mathcal{R}_{(3)}^{\beta*}(P_5) = 0.207$. Therefore, the ranking $P_1 > P_2 > P_4 > P_3 > P_5$ is attained. This ranking is consistent with the proposed CASPAS method. In other words, there is consistency between the results of C-IF-TOPSIS and D-IF-CASPAS methods.

7.3. A comparison with C-IF VIKOR. The VIKOR (VIseKriterijumska Optimizacija I Kompromisno Resenje) method is utilized in MCDM and MCGDM, pioneered by Opricovic [42]. Its main goal is to pinpoint a compromise solution that serves as a practical alternative closely resembling the ideal solution, followed by establishing a ranking of these alternatives, considering various evaluation criteria. Subsequently, a variation of the VIKOR method, known as the fuzzy VIKOR method, was devised to handle MCDM challenges in fuzzy environments [43]. This approach emphasizes the proximity of alternatives to the ideal solution, effectively managing decision-making hurdles in fuzzy or uncertain contexts. The VIKOR approach determines a compromise solution based on mutual concessions. Kahraman and Otay [26] proposed an extended VIKOR method for C-IFSs. Then, the waste disposal location selection problem was investigated using this C-IF VIKOR. This method can be briefly summarised as follows. Firstly, the decision matrices of the experts are aggregated and then the aggregated decision matrix is created. C-IF positive and negative ideal solutions are determined as

(7.9)
$$\theta_j^* = \max_{1 \le i \le m} \theta_{i,j} \text{ and } \theta_j^- = \max_{1 \le i \le m} \theta_{i,j}$$

using the relative score function (RSF). Maximum group utility, minimum individual regret and VIKOR index are calculated by using D_p and D_o distances, which are defined in two different ways as pessimistic and optimistic view.

Now, we will implement the C-IF VIKOR method to address the solar panel selection issue outlined in Sub-section 5.3. Firstly, using the aggregated decision matrix given in Table 8, the positive and negative ideal solutions shown in Table 17 are determined according to the RSF.

	T_1	T_2	T_3	T_4
θ^*	$\langle (0.87, 0.13); 0.86 \rangle$	$\langle (0.66, 0.34); 0.34 \rangle$	$\langle (0.84, 0.16); 0.82 \rangle$	$\langle (0.65, 0.35); 0.35 \rangle$
θ^{-}	$\langle (0.53, 0.47); 0.51 \rangle$	$\langle (0.32, 0.68); 0.68 \rangle$	$\langle (0.44, 0.56); 0.44 \rangle$	$\langle (0.42, 0.58); 0.58 \rangle$
TABLE 17. C-IF positive and negative ideal solutions				

By following the necessary steps, according to the VIKOR index, we reach the rankings $P_1 > P_2 > P_4 > P_5 > P_3$ for the optimistic case and $P_1 > P_2 > P_5 > P_4 > P_3$ for the pessimistic case. When compared with the proposed D-IF CASPAS method, it is seen that the results are compatible with C-IF VIKOR.

In this section where the D-IF CASPAS method is compared to other approaches, the similarity in results between the proposed method and C-IF TOPSIS and C-IF VIKOR demonstrates result consistency. Furthermore, contrasting the weighted summation operators with the Choquet integral operator highlights how criteria interactions can influence outcomes. Despite its more intricate calculation process compared to weighted aggregation operators, Choquet integral operators yield more precise results by considering criterion interactions, facilitating more accurate modeling of real-world problems. Figure 10 summarises the comparison of the results of the existing methods and the method proposed in this study.



FIGURE 10. Ranking results of existing methods for the solar panel selection problem

8. Contributions and future studies

This study proposed D-IFSs, a generalization of C-IFSs, and develops a range of operations for their application MCDM. A novel method, CASPAS, is proposed, which builds on the Choquet integral and incorporates interactions between criteria, addressing a significant gap in traditional weighted aggregation methods. D-IFVs are ranked using score and accuracy functions, and algebraic operations are formulated by extending the radius limit from 1 to $\sqrt{2}$, allowing the creation of weighted arithmetic and geometric aggregation operators. Additionally, disc intuitionistic fuzzy Choquet interval operators are introduced, which improve

30

the ability to model complex interdependencies in real-world decision-making scenarios. The CASPAS method is applied to a renewable energy problem, specifically the selection of optimal solar panels, and demonstrates its effectiveness in improving decision-making accuracy by incorporating interdependencies among selection criteria. Comparative analyzes with existing methods, alongside sensitivity and validity analysis, confirm the robustness and applicability of the proposed method. The study's contributions advance the theoretical framework of D-IFSs and provide practical tools for enhancing renewable energy decision-making, supporting the global transition to sustainable energy solutions. Future studies could focus on expanding the CASPAS method to other complex decision-making problems, especially in sectors such as healthcare, transportation, and environmental management. Additionally, further exploration of different Choquet aggregation operators and their real-world applications in MCDM could refine the proposed methodology and contribute to a more comprehensive understanding of decision-making under uncertainty.

Acknowledgments

We gratefully thank Dr. Abdullah Erkam Gündüz (Ankara Yıldırım Beyazıt University, Electrical-Electronics Engineering Department) and Research Assistant Muhammed Selman Erel (Ankara Yıldırım Beyazıt University, Electrical-Electronics Engineering Department) for providing the expert opinions.

References

- Adams, W. G., Day, R. E. (1877). V. The action of light on selenium. Proceedings of the Royal Society of London, 25(171-178), 113-117.
- [2] Atanassov, K. T. (1999). Intuitionistic fuzzy sets. In Intuitionistic fuzzy sets (pp. 1-137). Physica, Heidelberg.
- [3] Atanassov, K. T. (2020). Circular intuitionistic fuzzy sets. Journal of Intelligent & Fuzzy Systems, 39(5), 5981-5986.
- [4] Atanassov, K., Marinov, E. (2021). Four distances for circular intuitionistic fuzzy sets. Mathematics, 9(10), 1121.
- [5] Attanayake, K., Wickramage, I., Samarasinghe, U., Ranmini, Y., Ehalapitiya, S., Jayathilaka, R., Yapa, S. (2024). Renewable energy as a solution to climate change: Insights from a comprehensive study across nations. Plos one, 19(6), e0299807.
- [6] Baušys, R., Juodagalvienė, B. (2017). Garage location selection for residential house by WASPAS-SVNS method. Journal of civil engineering and management, 23(3), 421-429.
- [7] Becquerel, E. (1839). Mémoire sur les effets électriques produits sous l'influence des rayons solaires. Comptes Rendus, 9, 561–567.
- [8] Bozyiğit, M. C., Olgun, M., & Ünver, M. (2023). Circular Pythagorean Fuzzy Sets and Applications to Multi-Criteria Decision Making. Informatica, 1-30.
- [9] Bozyiğit, M. C., Ünver, M. (2024). Parametric circular intuitionistic fuzzy information measures and multicriteria decision making with extended TOPSIS. Granular Computing, 9(2), 43.
- [10] Bozyiğit, M. C., Ünver, M., Olgun, M. (2024, July). Circular Pythagorean Fuzzy Choquet Integral Operators and Applications to Multi-criteria Decision Making. In International Conference on Intelligent and Fuzzy Systems (pp. 635-643). Cham: Springer Nature Switzerland.
- [11] Chapin, D. M., Fuller, C. S., Pearson, G. L. (1954). A new silicon p-n junction photocell for converting solar radiation into electrical power. Journal of applied physics, 25(5), 676.
- [12] Chen, T. Y. (2023). Evolved distance measures for circular intuitionistic fuzzy sets and their exploitation in the technique for order preference by similarity to ideal solutions. Artificial Intelligence Review, 56(7), 7347-7401.
- [13] Choquet, G. (1954). Theory of Capacities. Annels del Institut Fourier, 5, 131–295.
- [14] Dincer, I., Temiz, M. (2024). Renewable Energy Sources. In: Renewable Energy Options for Power Generation and Desalination. Springer, Cham.
- [15] Dobrzański, L. A., Drygała, A., Giedroć, M., Macek, M. (2012). Monocrystalline silicon solar cells applied in photovoltaic system. Journal of achievements in materials and manufacturing engineering, 53(1), 7-13.
- [16] Easton, R. L., Votaw, M. J. (1959). Vanguard I IGY satellite (1958 beta). Review of Scientific Instruments, 30(2), 70-75.
- [17] Einstein, A. (1905, January). Über einen die Erzeugung und Verwandlung des Lichtes betreffenden heuristischen Gesichtspunkt.
- [18] El-Bayeh, C. Z., Alzaareer, K., Brahmi, B., Zellagui, M., Eicker, U. (2021). An original multi-criteria decision-making algorithm for solar panels selection in buildings. Energy, 217, 119396.

- [19] Fraas, L., Partain, L. (2010). Solar cells: a brief history and introduction. Solar cells and their applications, 1-15.
- [20] Garg, H., Ünver, M., Olgun, M., Türkarslan, E. (2023). An extended EDAS method with circular intuitionistic fuzzy value features and its application to multi-criteria decision-making process. Artificial intelligence review, 56(Suppl 3), 3173-3204.
- [21] Green, M. A. (2015). The passivated emitter and rear cell (PERC): From conception to mass production. Solar Energy Materials and Solar Cells, 143, 190-197.
- [22] Guerrero-Lemus, R. V. T. K. A. K. L. S. R., Vega, R., Kim, T., Kimm, A., Shephard, L. E. (2016). Bifacial solar photovoltaics–A technology review. Renewable and sustainable energy reviews, 60, 1533-1549.
- [23] Hosouli, S., Hassani, R. A. (2024). Application of multi-criteria decision making (MCDM) model for solar plant location selection. Results in Engineering, 24, 103162.
- [24] Hwang, C. L., Yoon, K. (1981) Methods for multiple attribute decision making. In Multiple attribute decision making (pp. 58-191). Springer, Berlin, Heidelberg.
- [25] IRENA (2024), Renewable energy statistics 2024, International Renewable Energy Agency, Abu Dhabi.
- [26] Kahraman, C., & Otay, I. (2022). Extension of VIKOR method using circular intuitionistic fuzzy sets. In Intelligent and Fuzzy Techniques for Emerging Conditions and Digital Transformation: Proceedings of the INFUS 2021 Conference, held August 24-26, 2021. Volume 2 (pp. 48-57). Springer International Publishing.
- [27] Kannan, D., Moazzeni, S., mostafayi Darmian, S., Afrasiabi, A. (2021). A hybrid approach based on MCDM methods and Monte Carlo simulation for sustainable evaluation of potential solar sites in east of Iran. Journal of Cleaner Production, 279, 122368.
- [28] Kaur, H., Gupta, S., Dhingra, A. (2023). Selection of solar panel using entropy TOPSIS technique. Materials Today: Proceedings.
- [29] Keshavarz Ghorabaee, M., Zavadskas, E.K., Amiri, M., Esmaeili, A. (2016). Multi-criteria evaluation of green suppliers using an extendedWASPASmethodwith interval type-2 fuzzy sets. Journal of Cleaner Production, 137, 213–229.
- [30] Khan, M. J., Alcantud, J. C. R., Kumam, W., Kumam, P., Alreshidi, N. A. (2023). Expanding Pythagorean fuzzy sets with distinctive radii: Disc Pythagorean fuzzy sets. Complex & Intelligent Systems, 9(6), 7037-7054.
- [31] Khan, F. M., Munir, A., Albaity, M., Mahmood, T. (2024). Parameter Selection Impacting Software Reliability by Utilizing WASPAS Technique Based on Tangent Trigonometric Complex Fuzzy Aggregation Operators. IEEE Access.
- [32] Kutlu Gundogdu, F., Kahraman, C. (2019). Extension of WASPAS with spherical fuzzy sets. Informatica, 30(2), 269-292.
- [33] Lazaroiu, A. C., Gmal Osman, M., Strejoiu, C. V., Lazaroiu, G. (2023). A comprehensive overview of photovoltaic technologies and their efficiency for climate neutrality. Sustainability, 15(23), 16297.
- [34] Lee, T. D., Ebong, A. U. (2017). A review of thin film solar cell technologies and challenges. Renewable and Sustainable Energy Reviews, 70, 1286-1297.
- [35] Lust, T. (2015). Choquet integral versus weighted sum in multicriteria decision contexts. In Algorithmic Decision Theory: 4th International Conference, ADT 2015, Lexington, KY, USA, September 27–30, 2015, Proceedings 4 (pp. 288-304). Springer International Publishing.
- [36] Meyer, P., Pirlot, M. (2012, November). On the expressiveness of the additive value function and the Choquet integral models. In DA2PL 2012: from Multiple Criteria Decision Aid to Preference Learning (pp. 48-56).
- [37] Mishra, A. R., Rani, P. (2021). Multi-criteria healthcare waste disposal location selection based on Fermatean fuzzy WASPAS method. Complex & Intelligent Systems, 7(5), 2469-2484.
- [38] Nag, S. K., Gangopadhyay, T. K., Paserba, J. (2022). Solar photovoltaics: A brief history of technologies [history]. IEEE Power and Energy Magazine, 20(3), 77-85.
- [39] Novas, N., Garcia, R. M., Camacho, J. M., Alcayde, A. (2021). Advances in solar energy towards efficient and sustainable energy. Sustainability, 13(11), 6295.
- [40] Ohl, R. S. (1946). U.S. Patent No. 2,402,662. Washington, DC: U.S. Patent and Trademark Office.
- [41] Olgun, M., Türkarslan, E., Ünver, M., Ye, J. (2021). A cosine similarity measure based on the Choquet integral for intuitionistic fuzzy sets and its applications to pattern recognition. Informatica, 32(4), 849-864.
- [42] Opricovic, S. (1998). Multicriteria optimization of civil engineering systems, Faculty of Civil Engineering, PhD Thesis, Faculty of Civil Engineering, Belgrade
- [43] Opricovic, S. (2011). Fuzzy VIKOR with an application to water resources planning. Expert Systems with Applications, 38(10), 12983-12990.
- [44] Peng, X., Yang, Y. (2016). Pythagorean fuzzy Choquet integral based MABAC method for multiple attribute group decision making. International Journal of Intelligent Systems, 31(10), 989-1020.
- [45] Parida, B., Iniyan, S., Goic, R. (2011). A review of solar photovoltaic technologies. Renewable and sustainable energy reviews, 15(3), 1625-1636.
- [46] Rani, P., Mishra, A. R. (2020). Multi-criteria weighted aggregated sum product assessment framework for fuel technology selection using q-rung orthopair fuzzy sets. Sustainable Production and Consumption, 24, 90-104.

- [47] Sathaye, J., Lucon, O., Rahman, A., Christensen, J. M., Denton, F., Fujino, J., ..., Zhang, Y. (2011). Renewable energy in the context of sustainable development. In IPCC special report on renewable energy sources and climate change mitigation. Cambridge University Press.
- [48] Schumacher, D. (1985). The 1973 oil crisis and its aftermath. In Energy: Crisis or Opportunity? (pp. 21-41). Palgrave, London.
- [49] Senapati, T., Chen, G. (2022). Picture fuzzy WASPAS technique and its application in multi-criteria decision-making. Soft Computing, 26(9), 4413-4421.
- [50] Shayani Mehr, P., Hafezalkotob, A., Fardi, K., Seiti, H., Movahedi Sobhani, F., Hafezalkotob, A. (2022). A comprehensive framework for solar panel technology selection: A BWM-MULTIMOOSRAL approach. Energy Science & Engineering, 10(12), 4595-4625.
- [51] Smith, W. (1873). The action of light on selenium. Journal of the Society of Telegraph Engineers, 2(4), 31-33.
- [52] Sorensen, B. (2004). Renewable energy. Elsevier.
- [53] Spanggaard, H., Krebs, F. C. (2004). A brief history of the development of organic and polymeric photovoltaics. Solar Energy Materials and Solar Cells, 83(2-3), 125-146.
- [54] Stanujkić, D., Karabašević, D. (2018). An extension of the WASPAS method for decision-making problems with intuitionistic fuzzy numbers: a case of website evaluation. Operational Research in Engineering Sciences: Theory and Applications, 1(1), 29-39.
- [55] Sugeno, M. (1974) Theory of Fuzzy Integrals and Its Applications. Ph.D. Thesis, Tokyo Institute of Technology, Tokyo.
- [56] Takahagi, E. (2000). On identification methods of λ -fuzzy measures using weights and λ . Japanese Journal of Fuzzy Sets and Systems, 12, 665–676.
- [57] Tan, C., Chen, X. (2010). Intuitionistic fuzzy Choquet integral operator for multi-criteria decision making. Expert Systems with Applications, 37(1), 149-157.
- [58] Tekbiyik Ersoy, N. (2025). Renewable Energy and the Need for Renewable Energy. In: Energy Efficiency and Renewable Energy Policies. Synthesis Lectures on Renewable Energy Technologies. Springer, Cham.
- [59] Turskis, Z., Zavadskas, E. K., Antuchevičienė, J., Kosareva, N. (2015). A hybrid model based on fuzzy AHP and fuzzy WASPAS for construction site selection.
- [60] Tüysüz, N., Kahraman, C. (2023). An integrated picture fuzzy Z-AHP & TOPSIS methodology: Application to solar panel selection. Applied Soft Computing, 149, 110951.
- [61] Ünver, M., Özçelik, G., Olgun, M. (2018). A fuzzy measure theoretical approach for multi criteria decision making problems containing sub-criteria. Journal of Intelligent & Fuzzy Systems, 35(6), 6461-6468.
- [62] Ünver, M., Ozcelik, G., Olgun, M. (2020). A pre-subadditive fuzzy measure model and its theoretical interpretation. TWMS Journal of Applied and Engineering Mathematics, 10(1), 270-278.
- [63] Wang, X., Triantaphyllou, E. (2008). Ranking irregularities when evaluating alternatives by using some ELECTRE methods. Omega, 36(1), 45-63.
- [64] Yalcin Kavus, B., Ayyildiz, E., Gulum Tas, P., Taskin, A. (2023). A hybrid Bayesian BWM and Pythagorean fuzzy WASPAS-based decision-making framework for parcel locker location selection problem. Environmental Science and Pollution Research, 30(39), 90006-90023.
- [65] Zadeh, L. A., (1965). Fuzzy sets. Information and control. 8(3):338-353.
- [66] Zavadskas, E. K., Turskis, Z., Antucheviciene, J., Zakarevicius, A. (2012). Optimization of weighted aggregated sum product assessment. Elektronika ir elektrotechnika, 122(6), 3-6.
- [67] Zavadskas, E. K., Antucheviciene, J., Hajiagha, S. H. R., Hashemi, S. S. (2014). Extension of weighted aggregated sum product assessment with interval-valued intuitionistic fuzzy numbers (WASPAS-IVIF). Applied soft computing, 24, 1013-1021.

¹ANKARA YILDIRIM BEYAZIT UNIVERSITY, FACULTY OF ENGINEERING AND NATURAL SCIENCES, DEPARTMENT OF MATHEMATICS, 06010 ANKARA TURKEY

Email address: mcbozyigit@aybu.edu.tr

²ANKARA UNIVERSITY, FACULTY OF SCIENCE, DEPARTMENT OF MATHEMATICS, 06100 ANKARA TURKEY *Email address*: munver@ankara.edu.tr