

# Faithful Simulation of Distributed Quantum Measurement with Coding for Computing

Anders Høst-Madsen

Department of Electrical & Computer Engineering

University of Hawaii, Manoa

Honolulu, HI, 96822, Email: ahm@hawaii.edu

**Abstract**—This papers consider a two terminal problem, where Alice and Bob jointly want to perform a measurement on a bipartite quantum system  $\rho^{AB}$ . Alice can transmit the results of her measurements to Bob on a classical channel, and Alice and Bob have common randomness. The question is what is the minimum amount of communications and common randomness needed. The paper derives an achievable rate region.

## I. INTRODUCTION

Faithful simulation of measurements was developed by Winter [1]. The idea is as follows. Alice performs a measurement on a quantum state  $\rho$  and sends some classical bits to Bob, who intends to *faithfully* recover Alice's measurement, preserving correlation with the reference system. The key observation is that if Alice and Bob have common randomness, the number of bits transmitted from Alice to Bob can be decreased below that of classical data compression of Alice's measurement outcomes, while still preserving the correlation with the reference system. We refer the reader to the overview paper [2] and the paper [3] for more details about faithful simulation. One could think of faithful simulation as a method for concretely reducing communication rates, but it also can be used to solve many other problems in quantum information theory [2], [3]. This paper is only concerned with finding communication rates leaving possible applications to later papers.

In [3] the authors considered the follow problem. Alice and Bob have a shared bipartite quantum system  $\rho^{AB}$ . Alice makes some measurements with a POVM  $\{\Lambda_u^A\}$  on  $\rho^A$  and Bob measures with  $\{\Lambda_v^B\}$  on  $\rho^B$ . Alice, Bob, and Charlie shares some common randomness and Alice and Bob transmits some classical information to Charlie. The goal is essentially for Charlie to faithfully simulate a function  $z = g(u, v)$  of the measurements.




Figure 1. System model. The aim is for Bob to faithfully simulate  $\Lambda_z^{AB} = \sum_{u,v:z=g(u,v)} \Lambda_u^A \otimes \Lambda_v^B$  on the bipartite system  $\rho^{AB}$ .

In this paper we consider the seemingly simpler problem in Fig. 1. The first difference from the problem considered in [3]

is that the three terminal function computation problem is not solved in general even for classical systems [4]. On the other hand, Orlitsky and Roche [5] solved the two terminal problem in Fig. 1 in the classical case; that solution will guide us to a quantum solution. The second difference is that Charlie in [3] has no quantum information. On the other hand, in Fig. 1 Bob has access to the quantum system  $\rho^B$  and can use that to decode Alice's transmission in addition to measuring  $\{\Lambda_v^B\}$ , similar to faithful simulation with quantum side-information [2], [6], [7].

The notation and terminology in the paper to a large extent follows the notation in [8]. We will use Theorems and Lemmas from [8] without repeating them here. We will use an implicit notation for probabilities when there is no possible confusion, e.g.,  $p(u^n) = p_u^n(u^n)$ .

## II. PROBLEM STATEMENT

We consider a bipartite composite quantum system  $(A, B)$  represented by a density operator  $\rho^{AB}$  on the Hilbert space  $\mathcal{H}_A \otimes \mathcal{H}_B$ . Alice has access to  $\rho^A$  and Bob has access to  $\rho^B$ . The aim is for Bob to perform the measurement  $\{\Lambda_z^{AB}\}_{z \in \mathcal{Z}}$

$$\Lambda_z^{AB} = \sum_{u \in \mathcal{U}, v \in \mathcal{V}: z=g(u,v)} \Lambda_u^A \otimes \Lambda_v^B$$

where  $\{\Lambda_u^A\}_{u \in \mathcal{U}}$  and  $\{\Lambda_v^B\}_{v \in \mathcal{V}}$  are POVMs on  $\mathcal{H}_A$  respectively  $\mathcal{H}_B$  and  $g$  is a deterministic function. This is called a separable decomposition with deterministic integration in [3]. It is sufficient for Bob to faithfully simulate  $\{\Lambda_z^{AB}\}_{z \in \mathcal{Z}}$  using classical transmission from Alice and common randomness with Alice. To this end, Alice transmits the results of an approximate measurement on  $A^n$  as

$$\tilde{\Lambda}_{u^n}^A = \sum_{m=1}^M p_M(m) \Gamma_{u^n}^{(m)} \quad (1)$$

where  $p_M$  is the common randomness distribution, which in turn gives an approximate measurement of  $z^n$ :

$$\tilde{\Lambda}_{z^n}^{AB} = \sum_{u^n \in \mathcal{U}^n, v^n \in \mathcal{V}^n: z^n=g(u^n, v^n)} \tilde{\Lambda}_{u^n}^A \otimes \Lambda_{v^n}^B$$

The requirement is that  $\tilde{\Lambda}_{z^n}^{AB}$  faithfully simulates  $\Lambda_{z^n}^{AB}$  in the following sense. For all  $\epsilon > 0$  and for sufficiently large  $n$ , [2]

$$\sum_{z^n} \|\sqrt{\omega}(\tilde{\Lambda}_{z^n}^{AB} - \Lambda_{z^n}^{AB})\sqrt{\omega}\|_1 \leq \epsilon \quad (2)$$

Notice that there is no reason for Bob to do approximate measurement as we only consider the communications cost from Alice to Bob; an extension could be to also consider compression of  $z^n$ . Also notice that Alice's approximate measurement does not actually have to be a specific form (1); the only requirement is to have an approximate  $\tilde{\Lambda}_{z^n}^{AB}$  that satisfies (2). The results in [3] can be adapted to this system, and their Theorem 4 gives

$$R \geq I(U; RB) - I(U; V) \quad (3)$$

$$R + S \geq H(U|V) \quad (4)$$

where  $R$  is the rate of communications and  $S$  the rate of common randomness.

In the classical case, the above problem reduces to the one considered in [5]. Alice has access to a random variable  $U \in \mathcal{U}$  and Bob to  $V \in \mathcal{V}$  with a joint distribution  $p(u, v)$  and the goal for Bob is to calculate  $Z = g(U, V)$  (with no distortion). Since Orlitsky and Roche's approach is the basis for our solution, we will need to discuss it in some detail. The key concept is that of independence of points in  $\mathcal{U}$ , which we will simplify as

**Definition 1.**  $u, u' \in \mathcal{U}$  are independent if

$$\forall v \in \mathcal{V} : p(u, v), p(u', v) > 0 \Rightarrow g(u, v) = g(u', v)$$

Let  $\mathcal{G}$  denote the set of independent sets and let  $W$  be a  $\mathcal{G}$ -valued random variable. We get the distribution of  $W$  by choosing a conditional distribution  $p(w|u)$  where  $w$  ranges over all the sets in  $\mathcal{G}$  that contain  $u$ . We can restrict  $\mathcal{G}$  to the maximal independent sets. To clarify this concept, a few examples from [5] are illustrative

**Example 2.** If  $p(u, v) > 0$  for all  $u, v$ , the maximal independent sets are the subsets of  $\mathcal{U}$  where  $g$  is constant, i.e.,  $\forall v : g(u, v) = g(u', v)$  when  $u, u'$  is in the same independent set. The maximal independent sets is a partition of  $\mathcal{U}$  and  $w$  is a deterministic function of  $u$ . Alice can simply transmit  $w$  instead of  $u$ , and with classical Slepian-Wolf coding [9] the rate is  $H(W|V)$  (the results in [5] shows that this is optimum).

**Example 3.** Alice and Bob draw a card in  $\{1, 2, 3\}$  from a bag without replacement. Bob needs to determine who has the largest card. In this case  $\mathcal{G} = \{\{1, 2\}, \{2, 3\}\}$ . It is sufficient for Bob to know  $w \in \mathcal{G}$ . For example, if  $w = \{1, 2\}$ , if Bob has  $v = 1$  he knows Alice has the largest card, but if  $v \in \{2, 3\}$  he knows he has the largest card.

Orlitsky and Roche [5] show that for any  $p(w|u)$  the rate

$$R \geq I(W; U|V) = H(W|V) - H(W|V) \quad (5)$$

is achievable, and further that

$$H_G(U|V) = \min_{p(w|u)} I(W; U|V)$$

is optimum. The idea for the achievable rate is as follows. Alice generates  $s$  iid sequences  $w^n$  according  $p(w)$ . Given a sequence  $u^n$  she finds a  $w^n$  among the  $s$  sequences that is jointly (robust) typical with  $u^n$ ; she then randomly bins the

index into  $t$  bins and transmits the bin index to Bob. If  $s$  and  $t$  are sufficiently large ( $s = 2^{n(I(W;U)+\delta)}$ ), the error probability can be made to approach zero as  $n$  becomes large.

Orlitsky and Roche's proof technique does not mix well with faithful simulation. In short, the issue is that we want to simulate the  $w^n$  – reduce  $s$  below  $2^{nI(W;U)}$  – but the measurement approximation is for  $\Lambda_u$ . There are various ways to overcome this, but in this paper we will approach the problem in the following way. Instead of measuring  $u$  and then finding  $w$  as in the classical case, Alice measures  $\mathcal{G}$  directly. That is, Alice uses the POVMs

$$\Lambda_{w_A}^A = \sum_{u \in w} p_A(w|u) \Lambda_u^A \quad (6)$$

Notice that  $\sum_{w \in \mathcal{G}^A} \sum_{u \in w} p_A(w|u) \Lambda_u^A = I$ .

It should also be noticed that Bob does not necessarily have to measure  $v$  to calculate  $g$ . It might be possible to measure less, which has the advantage of disturbing the  $B$  system less, and that could help Bob decode Alice's transmission. So, we define

**Definition 4.** Let  $\mathcal{G}^A$  denote a set of independent sets spanning  $\mathcal{U}$  for Alice. We define  $v, v' \in \mathcal{V}$  to be independent if

$$\forall w \in \mathcal{G}^A : \forall u, u' \in w, p(u, v), p(u', v') > 0 : g(u, v) = g(u', v') \quad (7)$$

Let  $\mathcal{G}^B$  denote a set of independent sets spanning  $\mathcal{V}$  (not necessarily maximal). Similarly as for Alice, we can define a distribution  $p_B(w_B|v)$  and a measurement  $\Lambda_w^B$ .

We can now define a function  $\tilde{g} : \mathcal{G}^A \times \mathcal{G}^B \rightarrow \mathcal{Z}$  by  $\tilde{g}(w_A, w_B) = g(u, v)$  for any  $u \in w_A, v \in w_B$  with  $p(u, v) > 0$ ; the condition (7) ensures that this is a consistent definition.

We can now state the main result of the paper

**Theorem 5.** Let  $\mathcal{G}^A, \mathcal{G}^B$  be spanning independent sets of  $\mathcal{U}, \mathcal{V}$ . Let  $W_A$  be distributed according to  $p_A(w|u)$  and  $W_B$  according to  $p_B(w|v)$ . There exists a faithful simulation of  $\Lambda_z$  with communication rate  $R$  and common randomness rate  $S$  if

$$R \geq I(W_A; RB) - I(W_A; B|W_B) - I(W_A; W_B) \quad (8)$$

$$R + S \geq H(W_A|W_B) - I(W_A; B|W_B) \quad (9)$$

Here  $I(W_A; RB)$  is evaluated on the state  $\sum_{w_A} |w_A\rangle\langle w_A| \otimes \text{Tr}_A \{(I^R \otimes \Lambda_{w_A}^A \otimes I^B) \phi^{RAB}\}$  and  $I(W_A; B|W_B)$  on  $\sum_{w_A, w_B} |w_A\rangle\langle w_A| \otimes |w_B\rangle\langle w_B| \otimes \text{Tr}_A \{(I^R \otimes \Lambda_{w_A}^A \otimes \Lambda_{w_B}^B) \phi^{RAB}\}$  and  $\Lambda_w$  is given in (6).

**Example 6.** We consider a "quantified" version of Example 3. The ensemble is  $\{\frac{1}{6}, |u\rangle_A \langle u|_B\}_{u, v \in \{1, 2, 3\}, u \neq v}$  and the measurements  $\Lambda_u^A = |u\rangle\langle u|_A$ . The bound (8) gives  $R \geq I(W_A; U) - I(W_A; V)$ , which is the same as classical compression (5), and minimized as 0.541 [5], whereas (3) gives  $R \geq H(U|V) = 1$  (the Slepian-Wolf rate). So, (8) gives the optimum rate, and is strictly better than (3). The bound (9) gives  $R + S \geq H(W_A|V) = 0.874$ , which is less than (4). However, the rate 0.541 is achievable without any common

randomness simply by using classical compression (which also simulates  $\Lambda_z$ ), so (9) cannot be optimum.

The rest of the paper is the proof of Theorem 5.

### III. PROOF OF THEOREM 5

In the following we will prove this theorem. We will let  $m$  uniform on  $[M]$  denote the common randomness.

We first similarly to [2] define

$$\begin{aligned}\hat{\rho}_w^A &= \frac{1}{\text{Tr}\{\Lambda_w^A \rho^A\}} \sqrt{\rho^A} \Lambda_w^A \sqrt{\rho^A} \\ \xi'_{w_A^n} &= \Pi_{A^n}^\delta \Pi_{\hat{\rho}_w^A | w^n}^\delta \hat{\rho}_w^A \Pi_{\hat{\rho}_w^A | w^n}^\delta \Pi_{A^n}^\delta \\ \xi' &= \sum_{w^n} \tilde{p}_A(w^n) \xi'_{w_A^n}\end{aligned}\quad (10)$$

where  $\Pi_{A^n}^\delta$  is the typical projector for  $\rho^A$  and  $\Pi_{\hat{\rho}_w^A | W^n}^\delta$  is the conditional typical projector for the ensemble  $\{p_A(w), \hat{\rho}_w^A\}$

$$\tilde{p}_A(w^n) = \begin{cases} \frac{1}{S} p_A(w^n) & w^n \in T_\delta^{W_A^n} \\ 0 & \text{otherwise} \end{cases}\quad (11)$$

with  $S = \sum_{w^n \in T_\delta^{W_A^n}} p_A(w^n)$ . We let  $\Pi$  be the projector unto the eigenvectors of  $\xi'$  greater than  $\epsilon 2^{-n(H(\rho^A)+\delta)} = \epsilon 2^{-n(H(RB)+\delta)}$  and define

$$\begin{aligned}\Omega &= \Pi \xi' \Pi \\ \xi_{w_A^n} &= \Pi \xi'_{w_A^n} \Pi\end{aligned}$$

For use later in the proof (in (20)), we will prove some technical properties of  $\Omega$ :

$$\text{rank}(\Omega) \leq \text{Tr} \Pi \leq \text{Tr} \Pi_{A^n}^\delta \leq 2^{n(H(RB)+\delta)}$$

The second inequality is due to how  $\xi'_{w_A^n}$  is defined in (10) and the third inequality due to the bound on the dimension of the typical subspace [8]. The implication is that the eigenvalues less than  $\epsilon 2^{-n(H(RB)+\delta)}$  contribute at most  $\epsilon$  to  $\text{Tr} \Omega$ , and therefore

$$\begin{aligned}\text{Tr} \Omega &\geq (1 - \epsilon) \text{Tr} \xi' \geq (1 - \epsilon) E[\text{Tr} \xi'_{w_A^n}] \\ &\geq (1 - \epsilon)(1 - \epsilon - 2\sqrt{\epsilon})\end{aligned}\quad (12)$$

Where the last inequality follows from [2, (23)].

We generate random  $w_A^n(j, m), j \in [s], m \in [M]$  according to  $\tilde{p}_A$ . For  $j \in [s]$  we define the operators

$$\Gamma_j^{(m)} = \frac{S}{(1 + \epsilon)s} \sqrt{\omega^A}^{-1} \left( \xi_{w_A^n(j, m)} \right) \sqrt{\omega^A}^{-1}$$

The  $s$  possible outcomes is binned into  $t$  bins, and the bin index is transmitted to Bob.

In order for this scheme to work we need to prove

- 1) The set  $\Gamma_j^{(m)} = \{\Gamma_j^{(m)}\}_{j=1}^s$  constitutes a sub-POVM (with high probability).
- 2) Upon receiving the bin index  $i$  (and knowing  $m$ ), Bob can decode  $w_A^n(j, m)$  (with high probability).
- 3) The resulting measurement faithfully simulates  $\{\Gamma_z^{AB}\}_z$

1) The set  $\Gamma_j^{(m)} = \{\Gamma_j^{(m)}\}_{j=1}^s$  constitutes a sub-POVM: We will show that the set  $\Gamma_j^{(m)} = \{\Gamma_j^{(m)}\}_{j=1}^s$  is a sub-POVM with high probability. If it is not a sub-POVM we put  $\Gamma_j^{(m)} = \{I\}$ . We calculate

$$\sqrt{\omega^A} \sum_{j=1}^s \Gamma_j^{(m)} \sqrt{\omega^A} = \frac{S}{(1 + \epsilon)} \left( \frac{1}{s} \sum_{j=1}^s \xi_{w_A^n(j, m)} \right)$$

where each  $w^n(j, m)$  is chosen independently according to  $p_{\tilde{W}^n}$ . We notice that

$$\begin{aligned}SE[\xi_{w_A^n(j, m)}] &= S \sum_{w_A^n} \tilde{p}_A(w^n) \xi_{w_A^n} = S \Omega \\ &= \sum_{w^n \in T_\delta^{W_A^n}} p_A(w^n) \Pi \Pi_{A^n}^\delta \Pi_{\hat{\rho}_w^A | w^n}^\delta \hat{\rho}_w^A \Pi_{\hat{\rho}_w^A | w^n}^\delta \Pi_{A^n}^\delta \Pi \\ &\leq \sum_{w^n \in T_\delta^{W_A^n}} p_A(w^n) \Pi \Pi_{A^n}^\delta \hat{\rho}_w^A \Pi_{A^n}^\delta \Pi \\ &\leq \Pi \Pi_{A^n}^\delta \sum_{w^n} p_A(w^n) \hat{\rho}_w^A \Pi_{A^n}^\delta \Pi \\ &= \Pi \Pi_{A^n}^\delta (\rho^A)^{\otimes n} \Pi_{A^n}^\delta \Pi \leq (\rho^A)^{\otimes n} = \omega^A\end{aligned}\quad (13)$$

Let  $E_m$  be the event that

$$\frac{1}{s} \sum_{j=1}^s \beta \xi_{w_A^n(j, m)} \leq \beta \Omega (1 + \epsilon)$$

for some scaling factor  $\beta$ . By (13) this event is equivalent to  $\sum_{j=1}^s \Gamma_j^{(m)} \leq I$ , i.e., that  $\Gamma_j^{(m)}$  is a sub-POVM. We will show that  $E_m$  happens with high probability by using the operator Chernoff bound [8, Lemma 17.3.1]. We notice that by (13) and the definition of  $\Pi$ ,  $E[\beta \xi_{w_A^n(j, m)}] = \beta \Omega \geq \beta \epsilon 2^{-n(H(RB)+\delta)} \Pi$ . Furthermore

$$\begin{aligned}\beta \xi_{w_A^n} &= \beta \Pi \Pi_{\rho^A, \delta}^n \Pi_{\hat{\rho}_w^A, \delta}^n \hat{\rho}_w^A \Pi_{\hat{\rho}_w^A, \delta}^n \Pi_{\rho^A, \delta}^n \Pi \\ &\leq \beta \Pi \Pi_{\rho^A, \delta}^n 2^{-n(H(RB|W_A)-\delta)} \Pi_{\hat{\rho}_w^A, \delta}^n \Pi_{\rho^A, \delta}^n \Pi \\ &\leq \Pi\end{aligned}$$

when  $\beta = 2^{n(H(RB|W_A)-\delta)}$ . The first inequality follows from properties of conditional quantum typicality [8]. Then by the operator Chernoff bound

$$\begin{aligned}P(E_m^c) &= P\left(\frac{1}{s} \sum_{j=1}^s \beta \xi_{w_A^n(j, m)} > \beta \Omega (1 + \epsilon)\right) \\ &\leq 2 \text{rank}(\Pi) \exp\left(-\frac{s \epsilon^2 \beta \epsilon 2^{-n(H(RB)+\delta)}}{4 \ln 2}\right) \\ &\leq 2 \exp\left(-\frac{s \epsilon^3 2^{n(H(RB|W_A)-\delta)} 2^{-n(H(RB)+\delta)}}{4 \ln 2} + n(H(RB) + \delta)\right)\end{aligned}$$

Then with  $s = 2^{n(I(W_A; RB)+3\delta)}$  the error probability goes to zero. The total probability of error then is

$$\begin{aligned}P\left(\bigcup_m E_m^c\right) &\leq \sum_m P(E_m^c) \\ &\leq 2M \exp\left(-\frac{\epsilon^3 2^{n\delta}}{4 \ln 2} + n(H(RB) + \delta) \ln 2\right)\end{aligned}$$

So, as long as  $M \leq O(e^n)$ , the total error probability converges to zero.

2) *Bob can decode  $w_A^n$* : Bob measures  $w_B^n$ , which can be restricted to typical  $w_B^n$ , and also receives the bin index  $i$ . He then looks in bin  $i$  for  $w_A^n$  that are jointly typical with  $w_B^n$ ; call this set  $\mathcal{S}^{(m)}(i, w_B^n)$ . Let the index  $k$  enumerate  $\mathcal{S}^{(m)}(i, w_B^n)$ . The post-measurement states are

$$\tilde{\rho}_{w_B^n, w_A^n}^{B^n} = \frac{1}{p(i, k, m, w_B^n)} \text{Tr}_{A^n} \left\{ ((\Gamma_{i,k}^{(m)})^A \otimes \Lambda_{w_B^n}^B) (\rho^{AB})^{\otimes n} \right\}$$

with probabilities

$$p(i, k, m, w_B^n) = \text{Tr} \left\{ ((\Gamma_{i,k}^{(m)})^A \otimes \Lambda_{w_B^n}^B) (\rho^{AB})^{\otimes n} \right\}$$

We consider the conditional typicality projectors

$$\Pi_{i,k,m} = \Pi_{B^n | w_A^n(i,k,m), w_B^n}^\delta$$

Bob first uses a conditional typical projector  $\Pi_{B^n | w_B^n}^\delta$  followed by sequential decoding with  $\{\Pi_{i,k,m}, \hat{\Pi}_{i,k,m}\}$ , where  $\hat{\Pi}_{i,k,m} = I - \Pi_{i,k,m}$ . The probability of correct decoding of the  $k$ -th message is

$$\begin{aligned} P_c &= \text{Tr} \{ \hat{\Pi}_{i,k,m} \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \hat{\Pi}_{i,k,m} \} \\ \hat{\Pi}_{i,k,m} &= \Pi_{i,k,m} \hat{\Pi}_{i,k-1,m} \cdots \hat{\Pi}_{i,1,m} \Pi_{B^n | w_B^n}^\delta \\ \hat{\Pi}_{i,k,m} &= \Pi_{B^n | w_B^n}^\delta \hat{\Pi}_{i,1,m} \hat{\Pi}_{i,k-1,m} \cdots \Pi_{i,k,m} \end{aligned}$$

And the error probability is

$$P_e = 1 - E \left[ \frac{1}{M} \sum_{w_B^n} \sum_{i,k,m} p(i, k, m, w_B^n) \text{Tr} \{ \hat{\Pi}_{i,k,m} \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \hat{\Pi}_{i,k,m} \} \right] \quad (14)$$

$$= 1 - E \left[ \frac{1}{M} \sum_{w_B^n} \sum_{i,k,m} p(i, k, m, w_B^n) \text{Tr} \{ \Upsilon_{i,k,m} \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \} \right] \quad (15)$$

where the equality is due to the rotation invariance of the trace and where

$$\begin{aligned} \Upsilon_{i,k,m} &= \Pi_{B^n | w_B^n}^\delta \hat{\Pi}_{i,1,m} \hat{\Pi}_{i,k-1,m} \cdots \Pi_{i,k,m} \\ &\quad \times \Pi_{i,k,m} \hat{\Pi}_{i,k-1,m} \cdots \hat{\Pi}_{i,1,m} \Pi_{B^n | w_B^n}^\delta \end{aligned}$$

The outer sum in (14) is explicitly for  $w_B^n \in T_\delta^{W_B^n}$  and the expectation is both over the random choice of  $w_A^n(j, m)$  and the random binning.

The first step in the proof is to show that measuring on the states  $\tilde{\rho}_{w_B^n, w_A^n}^{B^n}$  is almost equivalent to measuring on the tensor product states  $\rho_{w_B^n, w_A^n}^{B^n}$ , which enables using typicality methods. To this end, we first notice that we can have  $w_A^n(j, m) = w_A^n(j', m')$  for  $(j, m) \neq (j', m')$ . That also means that for a given  $w_A^n$  there might be multiple  $\Upsilon_{i,k,m}$  measuring it. For a given  $w_A^n$  we would like to have a single measurement operator, for reasons that will become clear later. We define

$$\Upsilon_{w_A^n, w_B^n} = \arg \min_{\Upsilon_{i,k,m}: w_A^n(i,k,m)=w_A^n} \text{Tr} \{ \Upsilon_{i,k,m} \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \} \quad (16)$$

And  $\Upsilon_{w_A^n, w_B^n} = I$  if there is none. We can then write

$$\begin{aligned} P_e &= \frac{1}{M} \sum_{w_A^n, w_B^n} E \left[ \sum_{i,k,m} p(i, k, m, w_B^n) I_{w_A^n = w_A^n(i,k,m)} \right. \\ &\quad \left. \text{Tr} \{ (I - \Upsilon_{i,k,m}) \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \} \right] \\ &\leq \frac{1}{M} \sum_{w_A^n, w_B^n} E \left[ \sum_{i,k,m} p(i, k, m, w_B^n) I_{w_A^n = w_A^n(i,k,m)} \right. \\ &\quad \left. \text{Tr} \{ (I - \Upsilon_{w_A^n, w_B^n}) \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \} \right] \\ &= \frac{1}{M} \sum_{w_A^n, w_B^n} E \left[ \text{Tr} \left\{ (I - \Upsilon_{w_A^n, w_B^n}) \sum_{i,k,m} p(i, k, m, w_B^n) \right. \right. \\ &\quad \left. \left. I_{w_A^n = w_A^n(i,k,m)} \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \right\} \right] \\ &\leq \sum_{w_A^n, w_B^n} E \left[ \text{Tr} \left\{ (I - \Upsilon_{w_A^n, w_B^n}) p(w_A^n, w_B^n) \rho_{w_A^n, w_B^n}^{B^n} \right\} \right] \\ &\quad + E \left[ \sum_{w_A^n, w_B^n} \left\| p(w_A^n, w_B^n) \rho_{w_A^n, w_B^n}^{B^n} \right. \right. \\ &\quad \left. \left. - \sum_{i,k,m: k \in \mathcal{S}^{(m)}(i, w_B^n)} p(i, k, m, w_B^n) I_{w_A^n = w_A^n(i,k,m)} \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \right\| \right] \quad (17) \end{aligned}$$

with the last inequality due to the trace inequality,  $\text{Tr} \{ \Lambda \rho \} \leq \text{Tr} \{ \Lambda \sigma \} + \|\rho - \sigma\|_1$  and the fact that  $\Upsilon_{w_A^n, w_B^n}$  depends only on  $w_A^n, w_B^n$  so that it could be extracted from the inner sum. The second term in (17) is equivalent to the faithful simulation criterion (2) [2], which will be shown in Section III-3 to be less than  $\epsilon$ . We will bound the first term of (17). We again use rotational invariance of the trace to rewrite it on the form (14) as

$$\begin{aligned} P_e &\leq 1 - E \left[ \sum_{w_B^n} \sum_{i,k,m \in \mathcal{S}'(w_B^n)} p(i, k, m, w_B^n) \right. \\ &\quad \left. \text{Tr} \{ \hat{\Pi}_{i,k,m} \rho_{w_B^n, w_A^n}^{B^n} \hat{\Pi}_{i,k,m} \} \right] \end{aligned}$$

where  $\mathcal{S}'(w_B^n)$  are the indices that achieve minimum in (16). Notice that

$$\begin{aligned} 1 &= \text{Tr} \tilde{\rho}_{w_B^n, w_A^n}^{B^n} = \text{Tr} \{ \Pi_{B^n | w_B^n}^\delta \tilde{\rho}_{w_B^n, w_A^n}^{B^n} \} \\ &\quad + \text{Tr} \{ \hat{\Pi}_{B^n | w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n} \} \\ &\leq \text{Tr} \{ \Pi_{B^n | w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n} \Pi_{B^n | w_B^n}^\delta \} + \epsilon \end{aligned}$$

Then by the non-commutative union bound [8, Section 16.6]

$$P_e \leq 2 \left( \sum_{w_B^n} E \left[ \sum_{i,k,m \in S'(w_B^n)} p(i,k,m,w_B^n) \left( \text{Tr} \{ \hat{\Pi}_{i,k,m} \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(i,k,m) \Pi_{B^n|w_B^n}^\delta \} \right) + \sum_{l=1}^{k-1} \text{Tr} \{ \Pi_{i,l,m} \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(i,k,m) \Pi_{B^n|w_B^n}^\delta \} \right] \right)^{1/2} + \epsilon \quad (18)$$

The first term can be bounded by the trace inequality

$$\begin{aligned} T_1 &= \text{Tr} \{ \hat{\Pi}_{i,k,m} \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(i,k,m) \Pi_{B^n|w_B^n}^\delta \} \\ &\leq \text{Tr} \{ \hat{\Pi}_{i,k,m} \rho_{w_B^n, w_A^n}^{B^n}(i,k,m) \} \\ &\quad + \|\Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(i,k,m) \Pi_{B^n|w_B^n}^\delta - \rho_{w_B^n, w_A^n}^{B^n}(i,k,m)\|_1 \\ &\leq \epsilon + 2\sqrt{\epsilon} \end{aligned}$$

where the  $\epsilon$  follows from conditional typicality, and the second from Gentle Measurement [8, Lemma 9.4.2] as conditional typicality gives  $\text{Tr} \{ \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(i,k,m) \Pi_{B^n|w_B^n}^\delta \} \geq 1 - \epsilon$ .

The second term in (18) can be bounded by

$$\begin{aligned} T_2 &= \sum_{w_B^n} E \left[ \sum_{i,k,m \in S'(w_B^n)} p(i,k,m,w_B^n) \right. \\ &\quad \left. \sum_{l=1}^{k-1} \text{Tr} \{ \Pi_{i,l,m} \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(i,k,m) \Pi_{B^n|w_B^n}^\delta \} \right] \\ &\leq \sum_{w_B^n} E \left[ \sum_{i,k,m \in S'(w_B^n)} p(i,k,m,w_B^n) \right. \\ &\quad \left. \sum_{l \in S^{(m)}(i, w_B^n), l \neq k} \text{Tr} \{ \Pi_{i,l,m} \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(i,k,m) \Pi_{B^n|w_B^n}^\delta \} \right] \end{aligned}$$

The first sum is over a restricted set of the indices where  $w_A^n, w_B^n$  are jointly typical; the sum does not decrease if we instead sum over all indices where  $w_A^n, w_B^n$  are jointly typical. Summing over all  $i$  (bin number) and  $k$  (bin index) is equivalent to summing over all  $j \in [s]$  where  $w_A^n, w_B^n$  are jointly typical; we denote this set  $\mathcal{T}(w_B^n)$ . For the second sum, it is only over those indices where  $w_A^n$  is jointly typical with  $w_B^n$  in bin  $i$ . This is equivalent to summing over all  $w_A^n(j, m)$  jointly typical with  $w_B^n$ . Therefore, with  $B_j$  the bin index of  $w_A^n(j, m)$

$$\begin{aligned} T_2 &\leq \sum_{w_B^n} E \left[ \sum_{j \in \mathcal{T}(w_B^n), m} p(w_A^n(j, m), w_B^n) \sum_{j' \in \mathcal{T}(w_B^n): j' \neq j} I_{B_j=B_{j'}} \right. \\ &\quad \left. \text{Tr} \{ \Pi_{B^n|w_A^n}^\delta(j', m), w_B^n \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(j, m) \Pi_{B^n|w_B^n}^\delta \} \right] \end{aligned}$$

Here we move the expectation over the random binning inside the sum and notice that  $E[I_{B_j=B_{j'}}] = P(B_j = B_{j'}) = \frac{1}{t}$ , so that

$$T_2 \leq \frac{1}{t} \sum_{w_B^n} E \left[ \sum_{j \in \mathcal{T}(w_B^n), m} p(w_A^n(j, m), w_B^n) \sum_{j' \in \mathcal{T}(w_B^n): j' \neq j} \text{Tr} \{ \Pi_{B^n|w_A^n}^\delta(j', m), w_B^n \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(j, m) \Pi_{B^n|w_B^n}^\delta \} \right]$$

By classical typicality,  $\frac{p(w_A^n(j, m), w_B^n)}{2^{-n(H(W_A, W_B) - \delta)}}$ , so with  $T_2 = \frac{1}{t} T_2' 2^{-n(H(W_A, W_B) - \delta)}$   $\leq$

$$\begin{aligned} T_2' &\leq \sum_{w_B^n} E \left[ \sum_{j \in \mathcal{T}(w_B^n), m} \sum_{j' \in \mathcal{T}(w_B^n): j' \neq j} \text{Tr} \{ \Pi_{B^n|w_A^n}^\delta(j', m), w_B^n \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n, w_A^n}^{B^n}(j, m) \Pi_{B^n|w_B^n}^\delta \} \right] \\ &= \sum_{w_B^n} \sum_{j \in \mathcal{T}(w_B^n), m} \sum_{j' \in \mathcal{T}(w_B^n): j' \neq j} \text{Tr} \{ E[\Pi_{B^n|w_A^n}^\delta(j', m), w_B^n] \Pi_{B^n|w_B^n}^\delta E[\rho_{w_B^n, w_A^n}^{B^n}(j, m)] \Pi_{B^n|w_B^n}^\delta \} \\ &\leq \sum_{w_B^n} \sum_{j \in \mathcal{T}(w_B^n), m} \sum_{j' \in \mathcal{T}(w_B^n): j' \neq j} \text{Tr} \{ E[\Pi_{w_A^n}(j', m), w_B^n] \Pi_{B^n|w_B^n}^\delta \rho_{w_B^n}^{B^n} \Pi_{B^n|w_B^n}^\delta \} \end{aligned}$$

For the equality we used that  $w_A^n(j, m)$  and  $w_A^n(j', m)$  are chosen independently. For the last inequality we used  $E[\rho_{w_B^n, w_A^n}^{B^n}(j, m)] \leq \frac{1}{1-\epsilon} \rho_{w_B^n}^{B^n}$  [2] as the expectation is over typical  $w_A^n(j, m)$ . Now, by conditional quantum typicality

$$\begin{aligned} T_2' &\leq \frac{1}{(1-\epsilon)} 2^{-n(H(B|W_B) - \delta)} \sum_{w_B^n} \sum_{j \in \mathcal{T}(w_B^n), m} \sum_{j' \in \mathcal{T}(w_B^n): j' \neq j} \text{Tr} \{ E[\Pi_{w_A^n}(j', m), w_B^n] \Pi_{B^n|w_B^n}^\delta \} \\ &\leq \frac{1}{(1-\epsilon)} 2^{-n(H(B|W_B) - \delta)} 2^{n(H(B|W_A, W_B) + \delta)} \\ &\quad \sum_{w_B^n} \sum_{j \in \mathcal{T}(w_B^n), m} \sum_{j' \in \mathcal{T}(w_B^n): j' \neq j} 1 \end{aligned}$$

The sum can be bounded using classical typicality, and putting it together, we get

$$\begin{aligned} T_2 &\leq \frac{1}{t(1-\epsilon)} 2^{-n(H(W_A, W_B) - \delta)} 2^{-n(H(B|W_B) - \delta)} \\ &\quad 2^{n(H(B|W_A, W_B) + \delta)} 2^{n(H(w_B) + \delta)} \\ &\quad Ms 2^{-n(H(W_A|W_B) - \delta)} s 2^{-n(I(W_A; W_B) - \delta)} \end{aligned}$$

If we insert  $Ms = 2^{n(H(W_A) + 2\delta)}$  and  $s = 2^{n(I(W_A; RB) + 3\delta)}$  we get

$$T_2 \leq \frac{1}{t(1-\epsilon)} 2^{-n(I(W_A; B|W_B) - 2\delta)} 2^{n(I(W; RB) + 3\delta)} 2^{-n(I(W_A; W_B) - \delta)}$$

Thus we can use

$$t = 2^{-n(I(W_A; B|W_B) - 2\delta)} 2^{n(I(W; RB) + 4\delta)} 2^{-n(I(W_A; W_B) - \delta)}$$

3) The measurement faithfully simulates  $\{\Gamma_z^{AB}\}_z$ : We need to simulate

$$\Lambda_{z^n}^{AB} = \sum_{u^n, v^n: g^n(u^n, v^n) = z^n} \Lambda_{u^n}^A \otimes \Lambda_{v^n}^B$$

An alternative is as follows

$$\begin{aligned} \Lambda'_{z^n} &= \sum_{w_A^n, w_B^n: \tilde{g}^n(w_A^n, w_B^n) = z^n} \Lambda_{w_A^n}^A \otimes \Lambda_{w_B^n}^B \\ &= \sum_{w_A^n, w_B^n: \tilde{g}^n(w_A^n, w_B^n) = z^n} \sum_{u^n, v^n: p(w_A^n, w_B^n | u^n, v^n) > 0} p(w_A^n, w_B^n | u^n, v^n) \Lambda_{u^n}^A \otimes \Lambda_{v^n}^B \\ &= \sum_{u^n, v^n: g^n(u^n, v^n) = z^n} \left( \sum_{w_A^n, w_B^n: \tilde{g}^n(w_A^n, w_B^n) = z^n} p(w_A^n, w_B^n | u^n, v^n) \right) \Lambda_{u^n}^A \otimes \Lambda_{v^n}^B \\ &= \Lambda_{z^n}^{AB} \end{aligned}$$

Thus, we can equivalently prove simulation of  $\Lambda'_{z^n}$

From the previous step we know that Bob can decode  $w_A^n(j, m)$  with high probability. Let  $\mathcal{S}_z = \{z^n : \exists j, m, w_B^n : \tilde{g}^n(w_A^n(j, m), w_B^n) = z^n\}$ . For  $z^n \in \mathcal{S}_z$  we consider the following collection of operators

$$\begin{aligned} \tilde{\Lambda}_{z^n}^{AB} &= \frac{1}{M} \sum_m \sum_{w_B^n, j: \tilde{g}^n(w_A^n(j, m), w_B^n) = z^n} \Gamma_j^{(m)} \otimes \Lambda_{w_B^n}^B \\ &= \sum_{w_A^n, w_B^n: \tilde{g}^n(w_A^n, w_B^n) = z^n} \frac{c(w_A^n)}{Ms} \frac{S}{1+\epsilon} \sqrt{\omega^A}^{-1} \xi_{w_A^n} \sqrt{\omega^A}^{-1} \otimes \Lambda_{w_B^n}^B \\ &= \sum_{w_A^n, w_B^n: \tilde{g}^n(w_A^n, w_B^n) = z^n} \tilde{\Lambda}_{w_A^n}^A \otimes \Lambda_{w_B^n}^B \end{aligned}$$

where

$$c(w_A^n) = |\{m, j : w_A^n(m, j) = w_A^n\}|$$

We need to evaluate

$$\begin{aligned} d &= \sum_{z^n} \left\| \sqrt{\omega} (\Lambda'_{z^n} - \tilde{\Lambda}_{z^n}^{AB}) \sqrt{\omega} \right\|_1 \\ &= \sum_{z^n \notin \mathcal{S}_z} \left\| \sqrt{\omega} \Lambda'_{z^n} \sqrt{\omega} \right\|_1 + \sum_{z^n \in \mathcal{S}_z} \left\| \sqrt{\omega} (\Lambda'_{z^n} - \tilde{\Lambda}_{z^n}^{AB}) \sqrt{\omega} \right\|_1 \\ &\leq \epsilon + \sum_{z^n \in \mathcal{S}_z} \left\| \sqrt{\omega} \sum_{w_A^n, w_B^n: \tilde{g}^n(w_A^n, w_B^n) = z^n} (\tilde{\Lambda}_{w_A^n} - \Lambda_{w_A^n}^A) \otimes \Lambda_{w_B^n}^B \sqrt{\omega} \right\|_1 \\ &\leq \epsilon + \sum_{w_A^n \notin T_\delta^{W_A^n}, w_B^n} \left\| \sqrt{\omega} (\Lambda_{w_A^n}^A \otimes \Lambda_{w_B^n}^B) \sqrt{\omega} \right\|_1 \\ &\quad + \sum_{w_A^n \in T_\delta^{W_A^n}, w_B^n} \left\| \sqrt{\omega} (\tilde{\Lambda}_{w_A^n} - \Lambda_{w_A^n}^A) \otimes \Lambda_{w_B^n}^B \sqrt{\omega} \right\|_1 \\ &\leq 2\epsilon + \sum_{w_A^n \in T_\delta^{W_A^n}} \left\| \sqrt{\omega^A} (\tilde{\Lambda}_{w_A^n} - \Lambda_{w_A^n}^A) \sqrt{\omega^A} \right\|_1 \\ &= 2\epsilon + \sum_{w_A^n \in T_\delta^{W_A^n}} \left\| \frac{S}{1+\epsilon} \frac{c(w_A^n)}{Ms} \xi_{w_A^n} - \sqrt{\omega^A} \Lambda_{w_A^n}^A \sqrt{\omega^A} \right\|_1 \\ &= 2\epsilon + \sum_{w_A^n \in T_\delta^{W_A^n}} \left\| \frac{S}{1+\epsilon} \frac{c(w_A^n)}{Ms} \xi_{w_A^n} - p_A^n(w_A^n) \hat{\rho}_{w_A^n} \right\|_1 \end{aligned}$$

The first inequality follows from  $P(\mathcal{S}_z) \geq 1 - \epsilon$ , the second inequality from the triangle inequality and reorganizing the sums, the third inequality from classical typicality and [3, Lemma 3]. We continue to bound the second term

$$\begin{aligned} d_2 &\leq \sum_{w_A^n \in T_\delta^{W_A^n}} p_A(w_A^n) \left\| \xi_{w_A^n} - \hat{\rho}_{w_A^n} \right\|_1 \\ &\quad + \sum_{w_A^n \in T_\delta^{W_A^n}} \left\| \frac{S}{1+\epsilon} \frac{c(w_A^n)}{Ms} \xi_{w_A^n} - p_A(w_A^n) \xi_{w_A^n} \right\|_1 \\ &\leq \sum_{w_A^n \in T_\delta^{W_A^n}} p_A(w_A^n) \left\| \xi_{w_A^n} - \xi'_{w_A^n} \right\|_1 \\ &\quad + \sum_{w_A^n \in T_\delta^{W_A^n}} p_A(w_A^n) \left\| \xi'_{w_A^n} - \hat{\rho}_{w_A^n} \right\|_1 \\ &\quad + \sum_{w_A^n \in T_\delta^{W_A^n}} \left| \frac{1}{1+\epsilon} \frac{c(w_A^n)}{Ms} - \frac{1}{S} p_A(w_A^n) \right| \end{aligned} \quad (19)$$

We bound the first term as follows

$$\begin{aligned} &\sum_{w_A^n \in T_\delta^{W_A^n}} p_A(w_A^n) \left\| \xi_{w_A^n} - \xi'_{w_A^n} \right\|_1 \\ &\leq \sum_{w_A^n \in T_\delta^{W_A^n}} \tilde{p}_A(w_A^n) \left\| \Pi \xi'_{w_A^n} \Pi - \xi'_{w_A^n} \right\| \leq 2\sqrt{\epsilon'} \end{aligned}$$

by Gentle Measurement for Ensembles [8, Lemma 9.4.3], as

$$\sum_{w_A^n \in T_\delta^{W_A^n}} \tilde{p}_A^n(w_A^n) \text{Tr}\{\Pi \xi'_{w_A^n} \Pi\} = \text{Tr} \Omega \geq 1 - \epsilon' \quad (20)$$

with  $\epsilon' = (1 - \epsilon)(1 - \epsilon - 2\sqrt{\epsilon})$  by (12). For the second term, by Gentle Measurement [8, Lemma 9.4.2]

$$\left\| \xi'_{w_A^n} - \hat{\rho}_{w_A^n}^A \right\| = \left\| \Pi_{A^n}^\delta \Pi_{\hat{\rho}^A|w_A^n}^\delta \hat{\rho}_{w_A^n}^A \Pi_{\hat{\rho}^A|w_A^n}^\delta \Pi_{A^n}^\delta - \hat{\rho}_{w_A^n}^A \right\| \leq 2\sqrt{\epsilon'} \quad [7]$$

as

$$\begin{aligned} & \text{Tr}\{\Pi_{A^n}^\delta \Pi_{\hat{\rho}^A|w_A^n}^\delta \hat{\rho}_{w_A^n}^A \Pi_{\hat{\rho}^A|w_A^n}^\delta \Pi_{A^n}^\delta\} \\ &= \text{Tr}\{\Pi_{A^n}^\delta \Pi_{\hat{\rho}^A|w_A^n}^\delta \hat{\rho}_{w_A^n}^A \Pi_{\hat{\rho}^A|w_A^n}^\delta\} \\ &\geq \text{Tr}\{\Pi_{A^n}^\delta \hat{\rho}_{w_A^n}^A\} + \frac{1}{2} \|\Pi_{\hat{\rho}^A|w_A^n}^\delta \hat{\rho}_{w_A^n}^A \Pi_{\hat{\rho}^A|w_A^n}^\delta - \hat{\rho}_{w_A^n}^A\|_1 \\ &\geq 1 - \epsilon - \sqrt{\epsilon} \end{aligned}$$

Where we have used the trace inequality  $\text{Tr}\{\Lambda\rho\} \geq \text{Tr}\{\Lambda\sigma\} + \frac{1}{2}\|\rho - \sigma\|_1$  [8, Corollary 9.1.1]. Thus, we conclude that the first two terms in (19)  $\leq \epsilon$  for some  $\epsilon \rightarrow 0$ .

We now bound the third term in (19). As in [2] we use operator Chernoff bound [8, Lemma 17.3.1]. Let  $P$  be the diagonal matrix with  $\hat{p}_A^n(w_A^n)$  on the diagonal for all  $w_A^n \in T_\delta^{W_A}$ , and let  $C$  be the same for the empirical frequencies  $\frac{c(w_A^n)}{M_S}$ . We have  $E[C] = P$  and  $P \geq 2^{-n(H(W_A)+\delta)}I$ . Let  $E_0$  be the event that

$$(1 - \epsilon)P \leq C \leq (1 + \epsilon)P$$

The operator Chernoff bound then gives

$$P(E_0^c) \leq 2 \cdot 2^{-n(H(W_A)+\delta)} \exp\left(-\frac{M_S \epsilon 2^{-n(H(W_A)+\delta)}}{4 \ln 2}\right)$$

Thus, if  $M_S \geq 2^{-n(H(W_A)+2\delta)}$ , this probability converges to zero. We can then bound the third term in (19) conditioned on  $E_0^c$ :

$$\begin{aligned} \sum_{w_A^n \in T_\delta^{W_A}} \left| \frac{1}{1 + \epsilon} \frac{c(w_A^n)}{M_S} - \frac{1}{S} p_A^n(w_A^n) \right| &= \left\| \frac{1}{1 + \epsilon} C - P \right\|_1 \\ &\leq \frac{1}{1 + \epsilon} (\|\epsilon P\|_1 + \|C - P\|_1) \\ &\leq \frac{2\epsilon}{1 + \epsilon} \end{aligned}$$

These bounds were under the assumption that no error happens. But if an error happens the contribution to  $d$  is at most 1, and the total error probability converges to zero with  $n$ .

## REFERENCES

- [1] A. Winter, ““extrinsic” and “intrinsic” data in quantum measurements: Asymptotic convex decomposition of positive operator valued measures,” *Communications in Mathematical Physics*, vol. 244, no. 1, pp. 157–185, 2004. [Online]. Available: <https://doi.org/10.1007/s00220-003-0989-z>
- [2] M. M. Wilde, P. Hayden, F. Buscemi, and M.-H. Hsieh, “The information-theoretic costs of simulating quantum measurements,” *Journal of Physics A: Mathematical and Theoretical*, vol. 45, no. 45, p. 453001, 2012.
- [3] T. A. Atif, M. Heidari, and S. S. Pradhan, “Faithful simulation of distributed quantum measurements with applications in distributed rate-distortion theory,” *IEEE Transactions on Information Theory*, vol. 68, no. 2, pp. 1085–1118, 2022.
- [4] A. E. Gamal and Y.-H. Kim, *Network Information Theory*. Cambridge University Press, 2011.
- [5] A. Orłitsky and J. Roche, “Coding for computing,” *IEEE Transactions on Information Theory*, vol. 47, no. 3, pp. 903–917, 2001.

- [6] I. Devetak and A. Winter, “Classical data compression with quantum side information,” *Phys. Rev. A*, vol. 68, p. 042301, Oct 2003. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevA.68.042301>
- [7] J. M. Renes and R. Renner, “One-shot classical data compression with quantum side information and the distillation of common randomness or secret keys,” *IEEE Transactions on Information Theory*, vol. 58, no. 3, pp. 1985–1991, 2012.
- [8] M. M. Wilde, *Quantum Information Theory*, 2nd ed. Cambridge University Press, 2017.
- [9] T. Cover and J. Thomas, *Information Theory, 2nd Edition*. John Wiley, 2006.