## GROWTH OF AUTOMORPHISMS OF VIRTUALLY SPECIAL GROUPS

ELIA FIORAVANTI

ABSTRACT. We study the speed of growth of iterates of outer automorphisms of virtually special groups, in the Haglund–Wise sense. We show that each automorphism grows either polynomially or exponentially, and that its stretch factor is an algebraic integer. For coarse-median preserving automorphisms, we show that there are only finitely many growth rates and we construct an analogue of the Nielsen–Thurston decomposition of surface homeomorphisms.

These results are new already for right-angled Artin groups. However, even in this particular case, the proof requires studying automorphisms of arbitrary special groups in an essential way.

As results of independent interest, we show that special groups are accessibile over centralisers, and we construct a canonical JSJ decomposition over centralisers. We also prove that, for any virtually special group G, the outer automorphism group Out(G) is boundary amenable, satisfies the Tits alternative, and has finite virtual cohomological dimension.

#### 1. INTRODUCTION

Given an infinite, finitely generated group G, a fundamental problem is describing the structure of its outer automorphism group Out(G). From a topological perspective, this corresponds to understanding self homotopy equivalences of any classifying space of G. Unfortunately, the properties of G are rarely reflected into those of Out(G) in a straightforward way, and there are virtually no general tools to study the structure of automorphisms.

Some of the most elementary groups — such as free and surface groups — give rise to some of the most interesting and intricate outer automorphism groups — such as  $Out(F_n)$  and mapping class groups — whose study has occupied a central place in low-dimensional topology and geometric group theory in the past decades. At the same time, even harmless-looking groups such as the Baumslag–Solitar group BS(2, 4) can behave wildly: its outer automorphism group is not finitely generated [CL83].

The only general class of groups for which we have a near-complete understanding of automorphisms is arguably that of *Gromov-hyperbolic* groups [Gro87], also known as *negatively curved* groups. Breakthroughs of Rips and Sela have shown that every hyperbolic group G has a canonical JSJ decomposition [RS97], and that Out(G) is finitely generated and completely encoded in this decomposition [RS94, Sel97b], leading to the solution of the isomorphism problem in this case [Sel95, DG11]. Regrettably, our understanding of automorphisms for general families of groups has not evolved qualitatively since Rips and Sela's work in the 90s. Many of their results have been extended to *relatively hyperbolic* groups [DS08, BS08, Gro09, DG08, GL15], though always relying on somewhat similar techniques that require hyperbolicity in fundamental ways.

The main goal of this article is to move past the restrictions of negative curvature and develop a general theory of automorphisms for a broad and natural family of *non-positively curved* groups. We have found that the ideal setting for such a theory is provided by the (compact) special groups of Haglund and Wise [HW08]. This is arguably the most important class of *cubulated* groups — marrying a wealth of examples with a powerful toolkit to study them — so our choice was also motivated by recent questions of Rips on the structure of automorphisms of cubulated groups [Sel23a, p. 826].

The author is supported by Emmy Noether grant 515507199 of the Deutsche Forschungsgemeinschaft (DFG).

New general approaches to study automorphisms of non-(relatively-)hyperbolic groups were recently proposed by Groves and Hull [GH19] and by Sela [Sel23a, Sel23b], both relying on (weakly) acylindrical actions on hyperbolic spaces. What is different in our own approach is that we completely give up on acylindricity, rather embracing the fact that we will have to work with non-small  $\mathbb{R}$ -trees and non-acylindrical graphs of groups, and resolving by other means the issues that this causes. To some extent, this choice is a forced one: there exist special groups G such that Out(G)is infinite and such that G has no small actions on  $\mathbb{R}$ -trees [Fio24, p. 166], so not all automorphisms of G are "seen" by its (strongly) acylindrical actions on hyperbolic spaces.

Proving that Out(G) is finitely generated for all special groups G remains out of reach for the moment. Instead, we focus on the problem of analysing how fast the length of an element  $g \in G$  can grow when we apply iterates of an outer automorphism  $\phi \in Out(G)$ .

Even in the very particular case when G is a right-angled Artin group (RAAG), very little seems to be known on this question, despite the fact that several refined aspects of automorphisms of RAAGs are well-understood [Day09, DW19, BCV23], including finite generation of their automorphism groups. In fact, our approach to analyse growth of automorphisms of RAAGs requires studying automorphisms of *arbitrary* special groups in a key way: it relies on a hierarchical construction inspired by Nielsen–Thurston decompositions and [Sel96], by which we reduce the study of an automorphism of a RAAG to the study of a "simpler" automorphism of a general special group.

We now make precise the notion of "length" of a group element  $g \in G$ , and "growth" of an outer automorphism  $\phi \in \text{Out}(G)$ . For the former, it is natural to consider the *conjugacy length* ||g||, i.e. the minimum word length of an element in the conjugacy class of g, fixing once and for all a finite generating set of G, whose choice will play no role. If G is the fundamental group of a compact Riemannian manifold, then ||g|| is roughly equal to the length of a shortest closed geodesic in the free homotopy class determined by g. We then define the growth rate of g under  $\phi$  as the equivalence class of the sequence  $n \mapsto ||\phi^n(g)||$  up to bi-Lipschitz equivalence. More crudely, one can define the stretch factor of  $\phi$  as

$$\operatorname{str}(\phi) := \sup_{g \in G} \limsup_{n \to +\infty} \|\phi^n(g)\|^{1/n}.$$

The first and most important groups for which automorphism growth came to be fully understood were free abelian groups  $\mathbb{Z}^m$ , surface groups  $\pi_1(S)$ , and free groups  $F_m$ . Growth of elements of  $\operatorname{Out}(\mathbb{Z}^m) = \operatorname{GL}_m(\mathbb{Z})$  can be easily described in terms of the Jordan decomposition, while growth of elements of the mapping class group of a compact surface S is completely encoded in the corresponding Nielsen–Thurston decomposition [Thu88]. Finally, analysing growth of general automorphism of free groups  $F_m$  proved to be a significantly more complex problem, which required the development of refined techniques inspired by train tracks on surfaces [BH92, BFH00, BFH05, BG10] and was finally solved by Levitt [Lev09]. In all of these examples, each outer automorphism admits only finitely many growth rates as the element g varies in G, and each growth rate is bi-Lipschitz equivalent to a sequence  $n \mapsto n^p \lambda^n$  for some  $p \in \mathbb{N}$  and  $\lambda \geq 1$ . Moreover, the number  $\lambda$  happens to be an algebraic integer and a weak Perron number<sup>1</sup>.

Automorphisms of Gromov-hyperbolic groups (and toral relatively hyperbolic groups) have a similar growth behaviour, as was announced by [CHHL]. At the same time, almost nothing is known on growth of automorphisms of non-hyperbolic groups and all classical techniques to approach this problem — mainly train tracks and JSJ decompositions — are known to fail or run into serious issues when one abandons the world of (relatively) hyperbolic groups. Even restricting to the rather tame world of right-angled Artin groups, any growth information seems to be available only under significant restrictions on the type of automorphism under consideration [BQ18].

<sup>&</sup>lt;sup>1</sup>An algebraic integer is a root of a monic polynomial with integer coefficients. It is a weak Perron number if its modulus is not smaller than that of any of its Galois conjugates.

What is more, Coulon recently constructed a menagerie of finitely generated groups whose automorphisms exhibit a variety of "intermediate growth" behaviours [Cou22]: they growth faster than any polynomial and slower than any exponential. It remains unknown if these exotic behaviours can occur for automorphisms of finitely presented groups, but there seems to be no reason to expect the contrary.

Our first result is that growth of automorphisms of special groups is rather well-behaved, at least when it comes to the top growth rate. As a particular case, this applies to all automorphisms of right-angled Artin groups. We refer to Theorem 6.6 for a more precise statement.

**Theorem A.** Let G be a virtually special group. The following hold for every  $\phi \in Out(G)$ .

- (1) The stretch factor  $str(\phi)$  is an algebraic integer and a weak Perron number.
- (2) If  $str(\phi) = 1$ , then  $\phi$  grows at most polynomially.
- (3) If  $str(\phi) > 1$ , then the stretch factor is realised on a subgroup  $H \leq G$  that is either a surface group, a free product, or a group with a free abelian direct factor.

Roughly, part (3) of the theorem says that the only source of exponentially-growing automorphisms of special groups are pseudo-Anosovs on compact surfaces, fully irreducible automorphisms of free products, and skewing automorphisms of direct products (see Theorem 6.6(4)). Even more roughly: automorphisms of surface groups, free groups, and free abelian groups are the building blocks of exponentially-growing automorphisms, though this is true only in a rather weak sense.

Theorem A does *not* describe the growth rates of the elements of G under  $\phi$  up to bi-Lipschitz equivalence. However, if G is 1-ended and  $\operatorname{str}(\phi)$  is not realised on any maximal direct product inside of G, we show that the top growth rate of  $\phi$  is purely exponential. Namely, for every  $g \in G$ , the sequence  $n \mapsto \|\phi^n(g)\|$  is either bi-Lipschitz equivalent to  $n \mapsto \lambda^n$  for  $\lambda = \operatorname{str}(\phi)$ , or exponentially slower (see part (3) of Proposition 6.14).

We obtain a more complete description of growth rates under stronger assumptions on the automorphism. Specifically, we need  $\phi$  to preserve the coarse median structure on G induced by a convex-cocompact embedding into a right-angled Artin group. Despite the technical definition, this simply means that  $\phi$  does not skew the factors of any maximal direct product within G. Automorphisms of right-angled Artin groups are coarse-median preserving precisely when they are *untwisted* [Fio24], a class of automorphisms that is well-studied and important in its own right [CSV17]. In addition, *all* automorphisms of right-angled Coxeter groups are coarse-median preserving, and so are all automorphisms of Gromov-hyperbolic groups. See [FLS24, Theorem D] for further examples.

**Theorem B.** Let G be virtually special, and let  $\phi \in Out(G)$  be coarse-median preserving.

- (1) There exist finitely many weak Perron numbers  $\lambda_1, \ldots, \lambda_m > 1$  and an integer  $P \in \mathbb{N}$  such that the following holds. For every  $g \in G$ , the sequence  $n \mapsto \|\phi^n(g)\|$  is either bi-Lipschitz equivalent to  $n \mapsto n^p \lambda_i^n$  for some integer  $0 \le p \le P$  and some index  $1 \le i \le m$ , or it is eventually bounded by the sequence  $n \mapsto n^P$ .
- (2) For any growth rate  $\mathbf{o} = [n \mapsto n^p \lambda_i^n]$ , let  $\mathcal{K}(\mathbf{o})$  be the family of subgroups of G all of whose elements grow at most at speed  $\mathbf{o}$  under  $\phi$ . Then there are only finitely many G-conjugacy classes of maximal subgroups in  $\mathcal{K}(\mathbf{o})$ , and each of these is quasi-convex<sup>2</sup> in G.

Part (2) can be regarded as an analogue of the Nielsen–Thurston decomposition associated to a surface homeomorphism. Namely,  $\phi \in \text{Out}(G)$  determines finitely many quasi-convex subgroups  $K_1, \ldots, K_s \leq G$  such that all elements  $g \in G$  not conjugate into  $\bigcup_i K_i$  grow at the same speed under  $\phi$ , and all elements of the  $K_i$  grow strictly slower. A power of  $\phi$  preserves each  $K_i$ , and its restriction to each  $K_i$  admits a similar finite collection of quasi-convex subgroups containing elements whose

<sup>&</sup>lt;sup>2</sup>This is meant in the coarse-median sense and implies that the subgroup K is undistorted in G and itself virtually special. Equivalently, a finite-index subgroup of K is convex-cocompact in a finite-index special subgroup of G

growth rate is neither the fastest, nor the second fastest. And so on, down to polynomially-growing elements.

We should clarify that we do not show that sub-polynomial growth rates are *exactly* polynomial in Theorem B and, therefore, we leave open the rather unlikely possibility that there are infinitely many of the latter. We discuss in Remark 7.3 why dealing with sub-polynomial growth rates is so delicate; the main point is that this issue has to be dealt with by techniques that go beyond the scope of this article.

Part of the proofs of Theorem A and Theorem B is based on procedures that relate the group Out(G) to the groups Out(P), where  $P \leq G$  are lower-complexity special groups (Proposition 5.12). One consequence that we deem worthy of mention is the following result (Corollary 5.15). When G is a right-angled Artin group, this was shown in [BGH22, CV09, Hor14] (also see [Ham09, CV86, BFH00, BFH05] for the case when G is a free or surface group).

#### **Theorem C.** For any virtually special group G:

- (1) Out(G) is boundary amenable;
- (2) Out(G) is virtually torsion-free with finite cohomological dimension;
- (3)  $\operatorname{Out}(G)$  satisfies the Tits alternative and contains no Baumslag–Solitar subgroups  $\operatorname{BS}(m,n)$ with  $|m| \neq |n|$ .

These are some of the rare properties known to hold for all outer automorphism groups of virtually special groups; the only other one seems to be residual finiteness [AMS16]. Boundary amenability also implies that Out(G) satisfies the Novikov conjecture on higher signatures [Hig00, BCH94].

On the proof of Theorems A and B. The main difficulty to be overcome is that classical JSJ theory is not well-suited to describing automorphisms of non-hyperbolic groups. Any finitely presented group has a JSJ decomposition over cyclic subgroups [RS97], and more generally over slender subgroups [DS99, DS00, FP06, GL17], though there are two key differences.

- (1) JSJ decompositions of hyperbolic groups are *canonical*: they are graph-of-groups splittings that are preserved by all automorphisms of G [Sel97b, Bow98]. Canonical JSJs are available for more general groups G (the *canonical tree of cylinders* of [GL11]), but always under the assumption that abelian subgroups of G do not interact with each other in complicated ways (the typical example is that of a relatively hyperbolic group). By contrast, many special groups (e.g. freely indecomposable RAAGs) are *thick* in the sense of [BDM09], which is a strong negation of the kind of property needed for canonical trees of cylinders to work.
- (2) Cyclic splittings of hyperbolic groups encode all their automorphisms: if G is a torsion-free, 1-ended hyperbolic group that does not split over  $\mathbb{Z}$  as an amalgamated product or HNN extension, then  $\operatorname{Out}(G)$  is finite [BF95]. This property again badly fails for special groups: there are right-angled Artin groups  $A_{\Gamma}$  that do not split over any abelian subgroups, but nevertheless have infinite  $\operatorname{Out}(A_{\Gamma})$  [Fio24, Figure 1].

One fundamental clue as to how to overcome these two issues comes from our previous work in [Fio23]: if G is special and Out(G) is infinite, then G splits over a *centraliser* or a co-abelian subgroup thereof. Centralisers are not slender or small, and they have a complicated intersection pattern in general, but they also form a rather rigid collection of subgroups of G. They are closed under intersections and each of them is itself a special group of lower complexity than G.

One of the goals of this article — and its most important new tool — is the construction of a canonical JSJ decomposition over centralisers, for any special group G. To describe this, let  $\mathcal{S}(G)$  denote the collection of maximal subgroups of G that virtually split as direct products; we refer to these as the singular subgroups of G (Section 4.1).

**Theorem D.** If G is special and 1-ended, then G has an Aut(G)-invariant graph-of-groups splitting (possibly a single vertex) with the following properties:

- (1) each edge group is either a centraliser, or a cyclic subgroup;
- (2) each subgroup in  $\mathcal{S}(G)$  is conjugate into a vertex group;
- (3) each vertex group is of one of the following two kinds:
  - (a) a quadratically hanging subgroup with trivial fibre;
  - (b) a quasi-convex subgroup of G that is elliptic in all splittings of G over centralisers, relative to  $\mathcal{S}(G)$ .

We refer to Theorem 4.28 for the stronger statement that we will use in practice. Quadratically hanging subgroups are fundamental groups of compact surfaces with boundary; as in the case of hyperbolic groups, they need to be set apart in the above decomposition of G, as they admit too many cyclic splittings and no canonical ones. We instead refer to type (b) vertex groups  $V \leq G$ as the "rigid" ones. Unlike for hyperbolic groups, the restriction map  $Out(G) \rightarrow Out(V)$  can have infinite image even when V is rigid. What is important is that any automorphism of a rigid group V has its top growth rate realised on a lower-complexity special group (usually, a singular subgroup of V), which enables us to study growth by induction on a suitable form of complexity. As a base step, one has to deal with surface groups and free products, which can be done using (relative) train tracks. Modulo technicalities, this outlines the core of the proof of Theorem A.

The enhanced JSJ decomposition described in Theorem D is constructed in two main steps:

- (1) First, one constructs an Out(G)-invariant deformation space of splittings of G over centralisers (in the sense of [For02, GL17]), using an accessibility result over centralisers.
- (2) Then, one finds an Out(G)-invariant splitting in the invariant deformation space, using a new construction of canonical splittings; see Theorem 4.18(1). This construction shares some broad similarities with the canonical trees of cylinders of [GL11], but it can be applied under weaker hypotheses, which importantly allow for a complicated intersection pattern of abelian subgroups.

The accessibility result mentioned in the first step does not follow from classical forms of accessibility [Dun85, BF91, Sel97a], as we are interested in non-acylindrical splittings over non-small subgroups. We instead prove the following, which we believe of independent interest (see Theorem 3.1).

**Theorem E.** Any special group G admits an integer N = N(G) such that the following holds. Any reduced graph-of-groups splitting of G over centralisers has at most N edges.

Theorem B and its complete description of growth rates for coarse-median preserving automorphisms require additional work. The strategy is to consider the maximal subgroups of G whose elements grow at below-top speed under the iterates of  $\phi \in \text{Out}(G)$ . One can hope to show that these subgroups are again special and invariant under a power of  $\phi$ ; one can then restrict  $\phi$  to these subgroups and repeat the procedure. Showing that this terminates in finitely many steps eventually yields a description of all growth rates. A similar strategy was used by Sela to construct analogues of Nielsen–Thurston decompositions for automorphisms of free groups [Sel96].

Unfortunately, the maximal subgroups of G growing at below-top speed under  $\phi$  are not even finitely generated for a general automorphism  $\phi$  (Example A.4). This is where the coarse-median preserving hypothesis comes in to save the day: it guarantees that these maximal slow-growing subgroups are quasi-convex in G, and thus themselves special. This is the core of the proof of Theorem B and it relies on the following two key facts:

(1) If G is special and  $\phi \in \text{Out}(G)$  has infinite order, the Bestvina–Paulin construction [Bes88, Pau91] can be used to produce G-actions on  $\mathbb{R}$ -trees whose arc-stabilisers are co-abelian subgroups of centralisers [Fio23]; these are well-behaved subgroups of G, but they can be infinitely generated in general. Importantly, when  $\phi$  is coarse-median preserving, the arc-stabilisers of these  $\mathbb{R}$ -trees are *actual* centralisers (Theorem 2.27), hence finitely generated.

(2) We can then show that all point-stabilisers of these  $\mathbb{R}$ -trees are finitely generated, and even quasi-convex in G. This uses accessibility over centralisers (Theorem E) and some Rips–Sela theory, as we explain in Appendix B. Results of this type for small  $F_m$ –trees were obtained in [GL95, GJLL98].

Even if one is only interested in growth rates of untwisted automorphisms of right-angled Artin groups, the proof of Theorem B forces us to immediately abandon the world of RAAGs: the subgroups in the first stratum of the "Nielsen–Thurston decomposition" of the automorphism have no reason to be RAAGs in general (see Example 7.5). One then has to understand automorphisms of these more general special groups in order to prove Theorem B in this case. This should further motivate the level of generality pursued in this article: it also serves the very concrete purpose of understanding automorphisms of very concrete groups such as RAAGs.

We conclude by mentioning that some versions of Rips–Sela theory for actions of special groups on higher-dimensional median spaces are available [CRK11, CRK15]. We managed to only rely on actions on  $\mathbb{R}$ -trees for this article, but studying more refined aspects of automorphisms of special groups will likely require the higher-dimensional theory.

Structure of paper. Section 2 introduces important terminology and notation about special groups and growth rates of automorphisms. We also collect various recurring lemmas, recall the basics of the Bestvina–Paulin construction, and review the structural results on  $\mathbb{R}$ –trees arising from automorphisms of special groups that were shown in our previous work [Fio23, Fio24].

Section 3 is concerned with accessibility over centralisers; it contains the proof of Theorem E (see Theorem 3.1). Section 4 is devoted to the construction of the enhanced JSJ decomposition discussed in Theorem D (see Theorem 4.28); its core lies in Section 4.2.3. After this, Section 5 briefly connects the JSJ decomposition to automorphism growth, giving some initial applications and also proving boundary amenability, the Tits alternative and finiteness of virtual cohomological dimension for Out(G) (Theorem C and Corollary 5.15).

Section 6 contains the proof of Theorem A (see Theorem 6.6). The main technical step is contained in Section 6.3 and roughly amounts to showing that, for an exponentially-growing outer automorphism  $\phi \in \text{Out}(G)$  and any representative  $\varphi \in \text{Aut}(G)$ , the top growth rate of *conjugacy length* under  $\phi$  is comparable to the top growth rate of *word length* under  $\varphi$ . This is important in order to relate growth on a graph of groups to growth on the vertex groups (such as those of the enhanced JSJ decomposition). Finally, Section 7 proves Theorem B (see Theorem 7.1).

Appendix A is concerned with groups G that decompose into simpler pieces: direct products, free products, and graphs of groups invariant under some automorphism. In each of these cases, we relate the growth of automorphisms of G to their growth on the pieces of the decomposition. This is all fairly straightforward, but it does not seem to appear in the literature and it can get rather fiddly at times. Finally, Appendix B describes the structure of  $\mathbb{R}$ -trees acted upon by a finitely presented group that is accessible over the collection of arc-stabilisers. The main result (Theorem B.2) is a description of point-stabilisers, which is important in the proofs of both Theorem A and Theorem B.

Acknowledgements. I am particularly grateful to Ric Wade for suggesting that the proof of boundary amenability in Theorem C might show finiteness of vcd as well. I also thank Daniel Groves, Camille Horbez and Ashot Minasyan for useful conversations and references, and the authors of [CHHL] for sharing with me a preliminary version of their work after this article was completed.

#### CONTENTS

1.	Introduction	1
2.	Preliminaries	7
3.	Accessibility over centralisers	20
4.	Enhanced JSJ splittings	27
	0.00 -F0.00	

5.	First consequences of the enhanced JSJ splitting	48
6.	Growth of general automorphisms	58
7.	The coarse-median preserving case	69
App	pendix A. Growth vs decompositions	76
App	pendix B. Accessibility and $\mathbb{R}$ -trees	85
Refe	erences	93

#### 2. Preliminaries

This section collects various important results from the literature, and fixes some of our terminology and notation (though much of the latter will be introduced in the parts of the article where it first becomes relevant). Section 2.1 is concerned with special groups. Most importantly, we discuss the properties of four important classes of subgroups: centralisers, convex-cocompact subgroups, parabolics and semi-parabolics. Section 2.2 introduces important concepts regarding growth rates and automorphism growth. A lot of this is non-standard, but it will provide the most convenient terminology to phrase the material in the rest of the article. Section 2.3 recalls the basics of the Bestvina–Paulin construction [Bes88, Pau91] and its connections to automorphism growth. We also review the results of [Fio23] on the very particular structure of  $\mathbb{R}$ –trees arising from automorphisms of special groups.

2.1. Special groups. We say that a group G is  $special^3$  if it is the fundamental group of a compact special cube complex. Equivalently, G is a convex-cocompact subgroup of a right-angled Artin group  $A_{\Gamma}$ , with respect to the action on the universal cover of the Salvetti complex  $\mathcal{X}_{\Gamma}$  [HW08]; this means that G leaves invariant a convex subcomplex of  $\mathcal{X}_{\Gamma}$  and acts cocompactly on it.

**Remark 2.1.** Though Theorems A and B are stated for *virtually* special groups, we will only work with actual special groups in the main body of the paper. One can always reduce to this case by considering a finite-index, characteristic, special subgroup of any virtually special group.

Throughout the article, we fix a convex-cocompact embedding  $G \hookrightarrow A_{\Gamma}$ . It is important to remember that many of the notions that we will consider will not be intrinsic to G: they will rather depend on such a choice of an embedding. In particular, many of these notions might not be preserved by automorphisms of G (though this will not matter). Our results always hold for *all* choices of the convex-cocompact embedding  $G \hookrightarrow A_{\Gamma}$ .

In some parts of the article (those related to Theorem B), we will refer to a coarse median structure on G. This is an equivalence class of maps  $\mu: G^3 \to G$  pairwise at bounded distance from each other with respect to a word metric (see [Bow13, Fio24] for background). Our coarse median structure will always be the one induced by the median operator on  $\mathcal{X}_{\Gamma}$ , pulled back via the chosen convex-cocompact embedding  $G \hookrightarrow A_{\Gamma}$ . The coarse median structure on G is not unique in general: different convex-cocompact embeddings of G in right-angled Artin groups can induce different coarse median structures on G (see [FLS24] for explicit examples). Our results hold for all such structures. An element of Aut(G) is said to be coarse-median preserving if it preserves the chosen coarse median structure on G; this property is satisfied by all inner automorphisms, so it descends to a well-defined property of elements of Out(G). Coarse-median preserving automorphisms always form a subgroup of both Aut(G) and Out(G).

We will occasionally speak of *contracting* elements of G. These are the elements  $g \in G \setminus \{1\}$  with infinite cyclic centraliser in G, or equivalently those that act as Morse isometries on one/all locally

<sup>&</sup>lt;sup>3</sup>Several authors prefer calling these groups "compact special" and reserve the terminology "special" for fundamental groups of (possibly non-compact) special cube complexes, i.e. arbitrary subgroups of right-angled Artin groups.

finite Cayley graphs of G (in the sense of [Sis16, ACGH17]). For a third equivalent characterisation, explaining our terminology: these are the rank-1 isometries of  $\widetilde{C}$ , where  $\widetilde{C} \subseteq \mathcal{X}_{\Gamma}$  is a G-invariant convex subcomplex on which G acts cocompactly; rank-1 isometries of CAT(0) spaces are precisely those that are contracting in the sense of [BF09, CS15]. (Note however that contracting elements of G will fail to be rank-1 or contracting isometries of the whole  $\mathcal{X}_{\Gamma}$  in general.) See e.g. [Gen21b, Theorem 1.3] and [Sis16] for these equivalences.

The rest of this subsection is devoted to discussing four classes of subgroups of any special group G that will play an important role throughout the paper. They will be related by the implications shown in the following diagram:

 $\stackrel{\text{centraliser}}{\Longrightarrow} \underset{G\text{-parabolic}}{\longrightarrow} G\text{-semi-parabolic} \underset{}{\longrightarrow} \text{convex-cocompact.}$ 

With the exception of centralisers, which are an  $\operatorname{Aut}(G)$ -invariant class, these classes of subgroups depend on the choice of the embedding  $G \hookrightarrow A_{\Gamma}$  and are not preserved by automorphisms of G, in general. It is worth mentioning, however, that the class of convex-cocompact subgroups is preserved by coarse-median preserving automorphisms of G [Fio24, Corollary 3.3].

2.1.1. Centralisers. A subgroup  $H \leq G$  is a centraliser if there exists a subset  $A \subseteq G$  such that  $H = Z_G(A)$ , where  $Z_G(A) := \{g \in G \mid ga = ag, \forall a \in A\}$ . Equivalently, we have  $H = Z_G(Z_G(H))$ . We do not consider G itself to be a centraliser, except when it has nontrivial centre. We write

 $\mathcal{Z}(G) := \{ H \le G \mid H \text{ is a centraliser in } G \}.$ 

Note that, viewing G as a subgroup of some right-angled Artin group  $A_{\Gamma}$ , every element of  $\mathcal{Z}(G)$  is the intersection with G of an element of  $\mathcal{Z}(A_{\Gamma})$ , but the converse does not hold: many such intersections do not lie in  $\mathcal{Z}(G)$ . Importantly, the family  $\mathcal{Z}(G)$  is  $\operatorname{Aut}(G)$ -invariant and closed under taking arbitrary intersections.

Many other properties of centralisers in special groups G follow from the fact that they are convex-cocompact and G-semi-parabolic, two classes of subgroups of G that we discuss below.

2.1.2. *G*-parabolic subgroups. A subgroup of a right-angled Artin group  $A_{\Gamma}$  is parabolic if it is conjugate to the subgroup generated by a proper subset of the standard generators, namely  $A_{\Delta}$  for some  $\Delta \subsetneq \Gamma$ . Consequently, if  $G \leq A_{\Gamma}$  is convex-cocompact, we say that a subgroup  $H \leq G$  is *G*-parabolic if  $H \neq G$  and  $H = G \cap P$  for a parabolic subgroup  $P \leq A_{\Gamma}$ .

Again, we emphasise that the data of which subgroups are G-parabolic depends on the chosen embedding  $G \hookrightarrow A_{\Gamma}$ . Since we will consider such an embedding to be fixed throughout the article, we will write, with an abuse of notation:

$$\mathcal{P}(G) := \{ H \le G \mid H \text{ is } G \text{-parabolic} \}.$$

The family  $\mathcal{P}(G)$  is closed under taking intersections; in particular, the trivial subgroup is G-parabolic. However, unlike the family of centralisers,  $\mathcal{P}(G)$  does not have a purely algebraic characterisation and it fails to be  $\operatorname{Aut}(G)$ -invariant in general.

The main utility of G-parabolic subgroups comes from the following observation; see [Fio23, Corollary 3.21] for a proof.

## **Lemma 2.2.** There are only finitely many *G*-conjugacy classes of *G*-parabolic subgroups.

We say that a subgroup  $H \leq G$  is *root-closed* if, whenever  $g^n \in H$  for some  $g \in G$  and  $n \geq 2$ , we actually have  $g \in H$ . In right-angled Artin groups, parabolic subgroups and centralisers are root-closed. As a consequence, all elements of  $\mathcal{Z}(G) \cup \mathcal{P}(G)$  are root-closed in any special group G.

We also define the following concept for later use.

**Definition 2.3.** Let  $H \leq G$  be a subgroup. The *G*-parabolic closure of *H* (denoted  $\widehat{H}$ ) is the intersection of all *G*-parabolic subgroups containing *H*. If no such subgroup exists, we set  $\widehat{H} := G$ .

When  $\widehat{H} \neq G$ , we have  $\widehat{H} \in \mathcal{P}(G)$  and  $\widehat{H}$  is the smallest *G*-parabolic subgroup containing *H*.

2.1.3. Convex-cocompact subgroups. A subgroup  $H \leq G$  is convex-cocompact, with respect to the chosen embedding  $G \hookrightarrow A_{\Gamma}$ , if the universal cover of the Salvetti complex  $\mathcal{X}_{\Gamma}$  admits an *H*-invariant convex subcomplex on which *H* acts cocompactly.

The simplest examples of convex-cocompact subgroups are provided by centralisers and G-parabolic subgroups (see Lemma 2.5 below). The next lemma collects several preliminary properties of convex-cocompact subgroups of special groups.

**Lemma 2.4.** Let  $G \leq A_{\Gamma}$  and  $H, K \leq G$  all be convex-cocompact.

- (1) The intersection  $H \cap K$  is convex-cocompact.
- (2) The normaliser  $N_G(H)$  is convex-cocompact and virtually splits as  $H \times P$  for some  $P \in \mathcal{P}(G)$ .
- (3) If  $g \in G$  satisfies  $gHg^{-1} \leq H$ , then  $gHg^{-1} = H$ .
- (4) The set  $\{g \in G \mid gHg^{-1} \leq K\}$  equals the product  $K \cdot F \cdot N_G(H)$  for a finite subset  $F \subseteq G$ .
- (5) The set  $\{H \cap qKq^{-1} \mid q \in G\}$  contains only finitely many H-conjugacy classes of subgroups.

*Proof.* Parts (1) and (4) are respectively Lemmas 2.8 and 2.9 in [Fio23], while part (2) follows from [Fio23, Corollary 2.13].

Part (5) was essentially shown in [Fio23, Section 7.1], as we briefly explain. If  $C \subseteq \mathcal{X}_{\Gamma}$  is a convex subcomplex on which H acts cocompactly, there exists an integer N such that, for each  $g \in G$ , the intersection  $H \cap gKg^{-1}$  acts on a convex subcomplex of C with at most N orbits of vertices; this was shown in [Fio23, Lemma 7.5] when H = K, and the general case is identical. Now, for each  $m \geq 1$ , the group H has only finitely many conjugacy classes of convex-cocompact subgroups acting on a convex subcomplex of C with m orbits of vertices (see e.g. the claim in the proof of [Fio23, Lemma 2.9]), and this yields part (5).

Finally, we prove part (3). Let  $C \subseteq \mathcal{X}_{\Gamma}$  be a convex subcomplex on which H acts essentially; this exists by [CS11, Proposition 3.12]. Since H is convex-cocompact, the action  $H \curvearrowright C$  is cocompact; let n be its number of vertex orbits. For  $g \in A_{\Gamma}$ , the group  $gHg^{-1}$  acts essentially on the convex subcomplex  $gC \subseteq \mathcal{X}_{\Gamma}$ , also with n orbits of vertices.

If  $gHg^{-1} \leq H$ , then every  $gHg^{-1}$ -essential hyperplane of  $\mathcal{X}_{\Gamma}$  is H-essential; in particular, every hyperplane crossing gC also crosses C. If C and gC are disjoint, let  $x \in C$  and  $y \in gC$  be a pair of gates; we can write  $y = \gamma x$  for an element  $\gamma \in A_{\Gamma}$ . Every hyperplane of  $\mathcal{X}_{\Gamma}$  separating x and  $\gamma x$  must cross every hyperplane of gC, that is, every  $gHg^{-1}$ -essential hyperplane. It follows that  $\gamma$ commutes with  $gHg^{-1}$ . Thus, up to replacing g with  $\gamma^{-1}g$ , we can assume that  $gC \cap C \neq \emptyset$ , which implies that  $gC \subseteq C$ .

Now, consider the ascending union of convex subcomplexes  $U := \bigcup_{k\geq 0} g^{-k}C$  and the ascending union of subgroups  $V := \bigcup_{k\geq 0} g^{-k}Hg^k$ . Clearly,  $U \subseteq \mathcal{X}_{\Gamma}$  is again a convex subcomplex and it is V-invariant. For each  $k \geq 0$ , the action  $g^{-k}Hg^k \curvearrowright g^{-k}C$  has exactly n orbits of vertices; it follows that the same is true of the action  $V \curvearrowright U$ . This shows that V is convex-cocompact. Observing that g normalises V, part (2) implies that a power of g lies in a product  $V \times P$ , and hence in  $g^{-k}Hg^k \times P$ for some k. Thus a power of g normalises  $g^{-k}Hg^k$ , which is only possible if  $g^{-1}Hg = H$ .  $\Box$ 

**Lemma 2.5.** Let  $G \leq A_{\Gamma}$  be convex-cocompact.

- (1) All subgroups in the collections  $\mathcal{Z}(G)$  and  $\mathcal{P}(G)$  are convex-cocompact.
- (2) For every  $P \in \mathcal{P}(G)$ , the normaliser  $N_G(P)$  lies in  $\mathcal{P}(G)$ , except when  $N_G(P) = G$ .

*Proof.* The centraliser of any subset of a right-angled Artin group is convex-cocompact as a consequence of Servatius' Centraliser Theorem [Ser89] (see the proof of Lemma 2.7 below for details); parabolic subgroups are also evidently convex-cocompact. Intersecting these subgroups with G, it follows that all elements of  $\mathcal{Z}(G)$  and  $\mathcal{P}(G)$  are convex-cocompact as well, since convex-cocompactness is stable under taking finite intersections by Lemma 2.4(1).

Regarding part (2), consider an element  $P \in \mathcal{P}(G)$ . Up to conjugating the whole G by an element of  $A_{\Gamma}$ , there exists  $\Delta \subseteq \Gamma$  such that  $P = G \cap A_{\Delta}$ . Choose  $\Delta$  so that it is minimal with this property (even up to further conjugating G by elements of  $A_{\Gamma}$ ).

Let  $\mathcal{X}_{\Gamma}$  denote the universal cover of the Salvetti complex of  $A_{\Gamma}$ . Let  $C_P \subseteq \mathcal{X}_{\Delta} \subseteq \mathcal{X}_{\Gamma}$  be a convex subcomplex on which P acts essentially (in the sense of [CS11]). Note that every element of  $\Delta$ appears as the label of a hyperplane of  $C_P$ . Otherwise,  $C_P$  would be contained in a subcomplex of the form  $x\mathcal{X}_{\Delta_0}$  for some  $x \in A_{\Gamma}$  and  $\Delta_0 \subsetneq \Delta$ , and hence we would have  $P \leq G \cap xA_{\Delta_0}x^{-1}$  and  $x^{-1}Px \leq x^{-1}Gx \cap A_{\Delta_0}$ , violating minimality of  $\Delta$ .

We conclude the proof by showing that  $N_G(P)$  coincides with the subgroup  $G \cap N_{A_{\Gamma}}(A_{\Delta})$ . The latter is *G*-parabolic (or the entire *G*) because  $N_{A_{\Gamma}}(A_{\Delta}) = A_{\Delta} \times A_{\Delta^{\perp}}$  where  $\Delta^{\perp} := \bigcap_{x \in \Gamma} \operatorname{lk}(x)$ , for instance by [AM15, Proposition 3.13]. The inclusion  $G \cap N_{A_{\Gamma}}(A_{\Delta}) \leq N_G(P)$  is clear.

Conversely, if  $g \in N_G(P)$ , we consider the subcomplex  $gC_P \subseteq \mathcal{X}_{\Gamma}$ , which is acted upon essentially by  $gPg^{-1} = P$ . Thus, the convex subcomplexes  $C_P$  and  $gC_P$  are crossed by the same set  $\mathcal{U}$  of hyperplanes of  $\mathcal{X}_{\Gamma}$ , namely those hyperplanes that are P-essential. As a consequence, the set  $\mathcal{V}$  of hyperplanes separating  $C_P$  from  $gC_P$  is transverse to  $\mathcal{U}$ . Since every element of  $\Delta$  appears as the label of a hyperplane of  $C_P$ , as observed above, it follows that every hyperplane in  $\mathcal{V}$  is labelled by an element of  $\Delta^{\perp}$ . Thus, since  $C_P \subseteq \mathcal{X}_{\Delta}$ , we have  $gC_P \subseteq \mathcal{X}_{\Delta \cup \Delta^{\perp}}$  and hence  $g \in A_{\Delta \cup \Delta^{\perp}}$ . In conclusion,  $g \in G \cap N_{A_{\Gamma}}(A_{\Delta})$  as required.  $\Box$ 

2.1.4. *G*-semi-parabolic subgroups. Let  $G \leq A_{\Gamma}$  be convex-cocompact.

Centralisers are by far the most important class of subgroups of G for this article: they will provide the edge groups for most of the splittings of G that we will be interested in. A particularly useful property of centralisers is that they are close to being G-parabolic; their failure is only due to a (virtual) abelian factor. At the same time, the class  $\mathcal{Z}(G)$  also has a considerable weakness: if  $H \leq G$  is convex-cocompact and  $Z \in \mathcal{Z}(G)$ , the intersection  $H \cap Z$  will not normally lie in  $\mathcal{Z}(H)$ . To circumvent this issue, it is often useful to work with the following larger class of subgroups,

which have a similar structure to centralisers and additionally satisfy the above stability property.

**Definition 2.6.** A subgroup  $Q \leq G$  is G-semi-parabolic<sup>4</sup> if it is convex-cocompact, root-closed, and there exists a G-parabolic subgroup  $R \triangleleft Q$  such that Q/R abelian.

For simplicity, we will say that a subgroup is *co-abelian* if it is normal with abelian quotient.

# **Lemma 2.7.** All elements of $\mathcal{Z}(G)$ are *G*-semi-parabolic.

Proof. Consider  $Z \in \mathcal{Z}(G)$  and pick a subset  $B \subseteq G$  such that  $Z = Z_G(B)$ . Up to conjugating G by an element of  $A_{\Gamma}$ , the centraliser  $Z_{A_{\Gamma}}(B)$  is of the form  $A_{\Delta} \times A$ , for a subgraph  $\Delta \subseteq \Gamma$  and a convexcocompact abelian subgroup  $A \leq A_{\Delta^{\perp}}$  where  $\Delta^{\perp} = \bigcap_{v \in \Delta} \operatorname{lk}(v)$  (see [Ser89, Section III] or [Fio23, Remark 3.7(5) and Corollary 3.25]). The subgroup  $Z_{A_{\Gamma}}(B)$  is convex-cocompact and root-closed in  $A_{\Gamma}$ . Since we have  $Z = G \cap Z_{A_{\Gamma}}(B)$ , it follows that Z is convex-cocompact and root-closed in G. Since  $A_{\Delta}$  is co-abelian in  $Z_{A_{\Gamma}}(B)$ , it follows that the G-parabolic subgroup  $G \cap A_{\Delta}$  is co-abelian in Z, showing that Z is G-semi-parabolic.

The co-abelian G-parabolic subgroup R in Definition 2.6 is not uniquely defined. The next proposition shows that one can consider a more canonical *parabolic part* (Definition 2.9) for any G-semi-parabolic subgroup, which will play an important role throughout the article. What is

<sup>&</sup>lt;sup>4</sup>Using Lemma 2.4, it is not hard to show that this definition is indeed equivalent to [Fio23, Definition 3.29].

particularly useful is that parabolic parts can be defined without any explicit reference to the ambient right-angled Artin group, only requiring algebraic properties and the notion of convexcocompactness.

**Proposition 2.8.** Let  $Q \leq G$  be G-semi-parabolic.

- (1) There exists a unique minimal convex-cocompact, root-closed, co-abelian subgroup  $P \triangleleft Q$ .
- (2) The subgroup P is G-parabolic and it has trivial centre.
- (3) If  $gPg^{-1} \leq Q$  for some  $g \in G$ , then  $gPg^{-1} = P$ .
- (4) Denoting by A the centre of Q, the subgroup  $\langle P, A \rangle \cong P \times A$  has finite index in Q. Moreover, A is root-closed, convex-cocompact and (abstractly) isomorphic to the quotient Q/P.

*Proof.* We begin with the following observation, which is the main point of the proof.

**Claim.** There exists a co-abelian G-parabolic subgroup  $R_0 \triangleleft Q$  with trivial centre.

Proof of claim. Let  $R \triangleleft Q$  be a co-abelian G-parabolic subgroup, as provided by Definition 2.6. Denote by  $C_R$  the centre of R, and note that  $C_R$  is convex-cocompact as it lies in  $\mathcal{Z}(R)$ . Since R is convex-cocompact and normal in Q, Lemma 2.4(2) shows that Q virtually splits as  $R \times B$  for a Q-parabolic group B, which must be abelian as R is co-abelian in Q. Similarly, since  $C_R$  is convex-cocompact and normal in R, we have that R virtually splits as  $C_R \times R_0$  for an R-parabolic subgroup  $R_0$ . Actually, since R is G-parabolic, the fact that  $R_0$  is R-parabolic implies that  $R_0$  is G-parabolic, and  $C_R$  and B are abelian.

Note that  $R_0$  has trivial centre. If g is an element in the centre of  $R_0$ , then g commutes with  $R_0 \times C_R$ . It then follows that g commutes with R, since  $Z_G(g)$  is root-closed and  $R_0 \times C_R$  has finite index in R. Thus g lies the centre of R, and hence  $g \in C_R \cap R_0 = \{1\}$ .

Also note that  $R_0$  is normal in Q. Indeed,  $R_0$  is certainly normal in the product  $R_0 \times C_R \times B$ , which has finite index in Q. At the same time,  $R_0$  is G-parabolic, so  $N_G(R_0)$  is G-parabolic by Lemma 2.5(2), and in particular  $N_G(R_0)$  is root-closed. Thus, it follows that  $Q \leq N_G(R_0)$ , as claimed.

We are left to show that  $R_0$  is co-abelian in Q. For this, we consider the abelian group  $B' := C_R \times B$ , which is central in Q (again because centralisers are root-closed). The centraliser  $Z_{A_{\Gamma}}(B')$  splits as  $\Pi \times B''$ , where  $\Pi \leq A_{\Gamma}$  is parabolic and  $B'' \leq A_{\Gamma}$  is an abelian subgroup with  $B' \leq B''$  (again by the Centraliser Theorem). Now, since  $R_0$  is a convex-cocompact subgroup of the product  $\Pi \times B''$ , it virtually splits as the product  $(R_0 \cap \Pi) \times (R_0 \cap B'')$ , and  $R_0 \cap B''$  is trivial because  $R_0$  has trivial centre. Thus  $R_0 \cap \Pi$  has finite index in  $R_0$ , and since  $\Pi$  is root-closed, it follows that  $R_0 \leq \Pi$ . Conversely, a similar argument shows that  $\Pi \cap Q \leq R_0$ , since  $\Pi$  has trivial intersection with  $C_R \times B$ . In conclusion, Q is contained in the product  $\Pi \times B''$ , where B'' is abelian and  $R_0 = \Pi \cap Q$ . This shows that  $R_0$  is co-abelian in Q, completing the proof of the claim.

Using the claim, we now proceed to prove the four parts of the proposition, beginning with part (1). Let  $H \triangleleft Q$  be convex-cocompact, root-closed and co-abelian in Q. Then the intersection  $H \cap R_0$  is convex-cocompact and co-abelian in  $R_0$ . By Lemma 2.4(2) and the fact that  $R_0$  has trivial centre, the latter implies that  $H \cap R_0$  has finite index in  $R_0$  and, since  $H \cap R_0$  is root-closed, it follows that  $H \cap R_0 = R_0$ . In other words, we have  $R_0 \leq H$ , showing that  $R_0$  is the unique minimal convex-cocompact, root-closed, co-abelian subgroup of Q.

Setting  $P := R_0$ , we have proved parts (1) and (2) of the proposition. Part (4) follows from Lemma 2.4(2), noting that the centre of Q projects injectively to a finite-index subgroup of  $Q/R_0$ , and that the latter is a free abelian group (for instance, because  $Q/R_0$  embeds in the free abelian group B'' considered at the end of the proof of the claim).

We are left to prove part (3). Recall from the end of the proof of the claim that Q is contained in a product  $\Pi \times B''$ , where  $\Pi \leq A_{\Gamma}$  is parabolic, B'' is abelian, and  $R_0 = \Pi \cap Q$ . Up to conjugating the whole G by an element of  $A_{\Gamma}$ , we can assume that  $\Pi = A_{\Delta}$  for some  $\Delta \subseteq \Gamma$  and that  $B'' \leq A_{\Delta^{\perp}}$ , where  $\Delta^{\perp} = \bigcap_{v \in \Delta} \operatorname{lk}(v)$ . Now, thinking of elements of  $R_0$  as elements of the right-angled Artin group  $A_{\Gamma}$ , their cyclically reduced parts only involve standard generators in  $\Delta$ ; the same is true of elements of any conjugate  $gR_0g^{-1}$ . Thus, if we have  $gR_0g^{-1} \leq Q$ , then  $gR_0g^{-1} \leq R_0$ , because  $Q \leq A_{\Delta} \times A_{\Delta^{\perp}}$  and  $Q \cap A_{\Delta} = R_0$ . Finally, the inclusion  $gR_0g^{-1} \leq R_0$  implies that  $gR_0g^{-1} = R_0$ by Lemma 2.4(3), as required.

It is convenient to name the subgroup P described in the previous proposition.

**Definition 2.9.** Let  $Q \leq G$  be *G*-semi-parabolic. The *parabolic part* of *Q* is the unique minimal subgroup  $P \triangleleft Q$  that is convex-cocompact, root-closed and co-abelian.

The following example shows that part (4) of Proposition 2.8 cannot be improved.

**Example 2.10.** If  $Z \in \mathcal{Z}(G)$ , its parabolic part  $P \triangleleft Z$  is not a direct factor of Z in general, if we do not first pass to finite index. For instance, consider  $A_{\Gamma} = \langle a, b \rangle \times \langle c \rangle \cong F_2 \times \mathbb{Z}$  and the index-2 subgroup  $G = \langle a^2, b^2, ab, c^2, ac \rangle$ , which has a further index-2 subgroup splitting as the product  $\langle a^2, b^2, ab \rangle \times \langle c^2 \rangle$ . We have  $Z_G(c^2) = G$  and its parabolic part is  $P = \langle a^2, b^2, ab \rangle$ . Since  $G/P \cong \mathbb{Z} = \langle \overline{ac} \rangle$ , we have  $G = P \rtimes \mathbb{Z}$  and P is not a direct factor of G.

We also record the following observation on chains of semi-parabolic subgroups in special groups.

**Remark 2.11.** Let G be special and let  $Q_1, Q_2 \leq G$  be G-semi-parabolic subgroups with the same parabolic part P. If  $Q_1$  is a proper subgroup of  $Q_2$ , then  $Q_1$  has infinite index in  $Q_2$ ; this is because G-semi-parabolic subgroups are root-closed. It follows that the free abelian group  $Q_1/P$  is an infinite-index subgroup of the free abelian group  $Q_2/P$ , and thus it has strictly lower rank. This shows that ascending chains of G-semi-parabolic subgroups of G with a given parabolic part contain at most d + 1 elements, where d is the largest rank of a free abelian subgroup of G.

More generally, if  $Q_1 \leq Q_2$  are arbitrary *G*-semi-parabolic subgroups, then the parabolic part of  $Q_1$  is contained in that of  $Q_2$ . Observing that chains of *G*-parabolic subgroups of *G* have length at most  $|\Gamma^{(0)}|$ , we conclude that arbitrary chains of *G*-semi-parabolic subgroups of *G* have length at most  $|\Gamma^{(0)}| \cdot (d+1)$ . Of course, this bound is far from tight.

### 2.2. Growth rates.

2.2.1. Abstract growth rates. When we speak of a "growth rate" in this article, we will refer to the following general notion of how fast a sequence can grow.

If  $a, b: \mathbb{N} \to \mathbb{R}_{>0}$  are two sequences, we write  $a \leq b$  if there exists a constant C such that  $a_n \leq Cb_n$  for all  $n \geq 0$ ; equivalently, if we have  $\limsup_n a_n/b_n < +\infty$ . Informally, we will also say that a is *slower* than b, and that b is *faster* than a. We say that a and b are *equivalent*, written  $a \sim b$ , if we have both  $a \leq b$  and  $b \leq a$ . We denote by [1] the equivalence class of constant sequences.

**Definition 2.12.** A growth rate is a  $\sim$ -equivalence class  $[a_n]$  of sequences in  $(\mathbb{R}_{>0})^{\mathbb{N}}$  with  $[a_n] \succeq [1]$ . We denote by  $(\mathfrak{G}, \preceq)$  the set of all growth rates with the poset structure induced by the relation  $\preceq$ .

It will often be convenient to work modulo a non-principal ultrafilter  $\omega \subseteq 2^{\mathbb{N}}$  and the weaker ordering on sequences that it induces. We refer to [DK18, Chapter 10] for generalities on ultrafilters.

Given two sequences  $a, b: \mathbb{N} \to \mathbb{R}_{>0}$ , we write  $a \preceq_{\omega} b$  if we have  $\lim_{\omega} a_n/b_n < +\infty$ . Informally, we say that a is  $\omega$ -slower than b, and that b is  $\omega$ -faster than a. We say that a and b are  $\omega$ -equivalent, written  $a \sim_{\omega} b$ , if we have both  $a \preceq_{\omega} b$  and  $b \preceq_{\omega} a$ .

**Definition 2.13.** Let  $\omega$  be a non-principal ultrafilter on  $\mathbb{N}$ . An  $\omega$ -growth rate is a  $\sim_{\omega}$ -equivalence class  $[a_n]$  of sequences in  $(\mathbb{R}_{>0})^{\mathbb{N}}$  with  $[a_n] \succeq_{\omega} [1]$ . We denote by  $(\mathfrak{G}_{\omega}, \preceq_{\omega})$  the set of all  $\omega$ -growth rates with the total order induced by the relation  $\preceq_{\omega}$ .

Each  $\sim_{\omega}$ -equivalence class of sequences is a union of (uncountably many)  $\sim$ -equivalence classes. As a consequence, there is a natural order-preserving quotient map  $(\mathfrak{G}, \preceq) \to (\mathfrak{G}_{\omega}, \preceq_{\omega})$ . We emphasise that, while  $(\mathfrak{G}, \preceq)$  is only partially ordered, all its quotients  $(\mathfrak{G}_{\omega}, \preceq_{\omega})$  are totally ordered. We also record the following observation.

**Remark 2.14.**  $[a_n] \leq [b_n]$  holds if and only if  $[a_n] \leq_{\omega} [b_n]$  holds for all non-principal ultrafilters  $\omega$ .

We will often simply write  $a_n \leq b_n$ ,  $a_n \sim b_n$ ,  $a_n \leq b_n$ , etc in the rest the paper, omitting the square brackets when this streamlines notation without causing ambiguities.

2.2.2. Lengths on groups. Let G be a group with a finite generating set S.

We denote by  $|\cdot|_S$  and  $||\cdot||_S$  the word length and conjugacy length on G associated to S, as defined in the Introduction. If T is a different finite generating set, then  $|\cdot|_T$  and  $||\cdot||_T$  are bi-Lipschitz equivalent to  $|\cdot|_S$  and  $||\cdot||_S$ , respectively.

If  $G \curvearrowright X$  is an isometric action on a metric space, we also consider the translation length of an element, and the *displacement parameter* of the action:

$$\ell_X(g) := \inf_{x \in X} d(x, gx), \qquad \qquad \tau_X^S := \inf_{x \in X} \max_{s \in S} d(sx, x).$$

Despite our notation, these quantities do not just depend on the space X, but also on the specific action. Again, if T is a different finite generating set, there is a constant C = C(S,T) such that  $\frac{1}{C}\tau_X^S \leq \tau_X^T \leq C\tau_X^S$  for all G-spaces X. The following are straightforward observations.

**Lemma 2.15.** If  $\varphi \in \operatorname{Aut}(G)$ , the map  $\varphi \colon G \to G$  is bi-Lipschitz with respect to  $|\cdot|_S$  and  $||\cdot||_S$ .

**Lemma 2.16.** Consider an isometric action  $G \curvearrowright X$ .

- (1) For every  $g \in G$ , we have  $\ell_X(g) \leq \tau_X^S ||g||_S$ . (2) If the action  $G \curvearrowright X$  is proper and cocompact and X is geodesic, then there exists a constant c = c(S, X) > 0 such that  $\ell_X(g) \ge c ||g||_S$  for all  $g \in G$ .

**Remark 2.17.** Let  $H \leq G$  be generated by a finite subset  $T \subseteq H$ . If H is undistorted, then the word lengths  $|\cdot|_T$  and  $|\cdot|_S$  are bi-Lipschitz equivalent on H. However, undistortion alone does not suffice to conclude that the conjugacy lengths  $\|\cdot\|_T$  and  $\|\cdot\|_S$  are bi-Lipschitz equivalent on H.

If G is special and  $H \leq G$  is convex-cocompact, then  $\|\cdot\|_T$  and  $\|\cdot\|_S$  are indeed bi-Lipschitz equivalent on H. This can be seen by applying Lemma 2.16 twice, once to the action of H on a cocompact cubulation  $G \curvearrowright X$ , and once to its action on an H-cocompact convex subcomplex.

2.2.3. Growth of automorphisms. Consider now a finitely generated group G, an automorphism  $\varphi \in \operatorname{Aut}(G)$  and its outer class  $\phi \in \operatorname{Out}(G)$ . Let  $|\cdot|$  and  $||\cdot||$  be the word and conjugacy lengths on G with respect to some finite generating set, whose choice will play no role.

**Definition 2.18.** The growth rate of an element  $g \in G$  under  $\varphi$  is the  $\sim$ -equivalence class  $[|\varphi^n(g)|]$ in  $\mathfrak{G}$ . Similarly, the growth rate of g under  $\phi$  is<sup>5</sup> the equivalence class  $[\|\phi^n(g)\|] \in \mathfrak{G}$ .

In other words, we consider the equivalence class of the sequences  $n \mapsto |\varphi^n(q)|$  and  $n \mapsto ||\phi^n(q)||$ up to multiplicative constants. A different choice of generating set on G only alters  $|\cdot|$  and  $||\cdot||$ through a bi-Lipschitz equivalence, so it leads to the exact same growth rates within  $\mathfrak{G}$ . Since  $\|\cdot\| \leq |\cdot|$ , we always have

$$\left[\|\phi^n(g)\|\right] \preceq \left[|\varphi^n(g)|\right].$$

Moreover,  $[|\varphi^n(g)|] \sim [1]$  holds if and only if a power of  $\varphi$  fixes g, and  $[||\phi^n(g)||] \sim [1]$  holds if and only if a power of  $\phi$  preserves the conjugacy class of g.

<sup>&</sup>lt;sup>5</sup>Note that  $\phi^n(q)$  is not a well-defined element of G, but it is a well-defined conjugacy class.

We denote by  $\mathcal{G}(\varphi) \subseteq \mathfrak{G}$  and  $\mathfrak{g}(\phi) \subseteq \mathfrak{G}$  (or  $\mathcal{G}(G, \varphi)$  and  $\mathfrak{g}(G, \phi)$  if there is any ambiguity) the sets of all growth rates of the automorphism, as g varies in G. The constant rate [1] always appears in both sets, due to the identity element of G.

Of course, there are also versions of the above concepts modulo any non-principal ultrafilter  $\omega$ . We thus denote by  $\mathcal{G}_{\omega}(\varphi)$  and  $\mathfrak{g}_{\omega}(\phi)$  the sets of all  $\omega$ -growth rates of  $\varphi$  and  $\phi$ , that is, the images of the sets  $\mathcal{G}(\varphi)$  and  $\mathfrak{g}(\phi)$  under the projection  $\mathfrak{G} \to \mathfrak{G}_{\omega}$ .

In keeping with the above conventions, we will generally denote  $(\omega)$ -growth rates of automorphisms by the letter  $\mathcal{O}$ , and  $(\omega)$ -growth rates of outer automorphisms by the letter  $\mathfrak{o}$ .

There are two additional important elements of  $\mathfrak{G}$  that we can associate to  $\varphi$  and  $\phi$ . In some sense, they play the role of a "maximum" for the sets  $\mathcal{G}(\varphi)$  and  $\mathfrak{g}(\phi)$ , but it is important to stress that, a priori, they do **not** lie in  $\mathcal{G}(\varphi)$  or  $\mathfrak{g}(\phi)$ . Fixing any finite generating set  $S \subseteq G$ , we write:

$$\begin{aligned} \overline{\mathcal{O}}_{\mathrm{top}}(\varphi) &:= \big[\sigma^{S}(\varphi^{n})\big], & \text{where} \quad \sigma^{S}(\varphi) &:= \max_{s \in S} |\varphi(s)|, \\ \overline{\mathfrak{o}}_{\mathrm{top}}(\phi) &:= \big[\tau^{S}(\phi^{n})\big], & \text{where} \quad \tau^{S}(\phi) &:= \min_{x \in G} \max_{s \in S} |x\varphi(s)x^{-1}|. \end{aligned}$$

Note that  $\overline{\mathcal{O}}_{top}(\varphi)$  and  $\overline{\mathfrak{o}}_{top}(\varphi)$  are again completely independent of the choice of S. The notation  $\tau^{S}(\phi)$  is consistent with the one introduced in Section 2.2.2; it is exactly the displacement parameter for the action of G on the Cayley graph used to define the word length  $|\cdot|$ , pre-composed with  $\varphi$ .

**Lemma 2.19.** We have 
$$\mathcal{O} \leq \mathcal{O}_{top}(\varphi)$$
 for all  $\mathcal{O} \in \mathcal{G}(\varphi)$ , and  $\mathfrak{o} \leq \overline{\mathfrak{o}}_{top}(\phi)$  for all  $\mathfrak{o} \in \mathfrak{g}(\phi)$ .

*Proof.* For all  $g \in G$  and  $n \in \mathbb{N}$ , we have

$$|\varphi^n(g)| \le |g|_S \cdot \sigma^S(\varphi^n), \qquad \qquad \|\phi^n(g)\| \le \|g\|_S \cdot \tau^S(\phi^n),$$

where the latter inequality is obtained by applying Lemma 2.16(1) to the action of G on one of its Cayley graphs, precomposed with  $\varphi^n$ . The lemma immediately follows from these inequalities.  $\Box$ 

Despite the above lemma, it is not at all clear whether the sets  $\mathcal{G}(\varphi)$  and  $\mathfrak{g}(\phi)$  always have a  $\preceq$ -maximum, even for automorphisms of special groups. I still do not know if this is the case in the setting of Theorem A. This is true under the assumptions of Theorem B, but even in that situation it will only become clear near the end of the article (Theorem 7.1(1)).

What is instead clear, even at this preliminary stage, is that automorphisms of special groups always have a top  $\omega$ -growth rate, for any non-principal ultrafilter  $\omega$ .

**Lemma 2.20.** Let G be a special group. For every  $\varphi \in \operatorname{Aut}(G)$  and  $\phi \in \operatorname{Out}(G)$ , the sets  $\mathcal{G}_{\omega}(\varphi)$ and  $\mathfrak{g}_{\omega}(\phi)$  each have a  $\preceq_{\omega}$ -maximum in  $\mathfrak{G}_{\omega}$ .

*Proof.* We claim that the projections of  $\overline{\mathcal{O}}_{top}(\varphi)$  and  $\overline{\mathfrak{o}}_{top}(\phi)$  to  $\mathfrak{G}_{\omega}$  lie in  $\mathcal{G}_{\omega}(\varphi)$  and  $\mathfrak{g}_{\omega}(\phi)$ , respectively. By Lemma 2.19, these projections are indeed  $\omega$ -faster than any  $\omega$ -growth rate in the respective sets, so it will follow that they are the required  $\preceq_{\omega}$ -maxima.

To prove our claim, we need to find two elements  $g, g' \in G$  with  $[|\varphi^n(g)|] \sim_{\omega} \overline{\mathcal{O}}_{top}(\varphi)$  and  $[\|\phi^n(g')\|] \sim_{\omega} \overline{\mathfrak{o}}_{top}(\phi)$ . Since S is finite, there is an element  $s_0 \in S$  such that the equality  $\sigma^S(\varphi^n) = |\varphi^n(s_0)|$  holds for  $\omega$ -all integers n; we can then take  $g := s_0$ . (This works when G is an arbitrary finitely generated group.) As to the existence of an element g' with  $[\|\phi^n(g')\|] \sim_{\omega} \overline{\mathfrak{o}}_{top}(\phi)$ , it can be quickly shown with the Bestvina–Paulin construction, exploiting some weak properties of asymptotic cones of special groups; we explain this a couple of pages below, in Lemma 2.24.

Consequently, in the setting of the lemma, we will write

$$\mathcal{O}_{\rm top}^{\omega}(\varphi) := \max_{\omega} \mathcal{G}_{\omega}(\varphi), \qquad \qquad \mathfrak{o}_{\rm top}^{\omega}(\phi) := \max_{\omega} \mathfrak{g}_{\omega}(\phi)$$

From the proof of the Lemma 2.20, it follows that  $\mathcal{O}_{top}^{\omega}(\varphi)$  and  $\mathfrak{o}_{top}^{\omega}(\phi)$  are simply the projections to  $\mathfrak{G}_{\omega}$  of  $\overline{\mathcal{O}}_{top}(\varphi)$  and  $\overline{\mathfrak{o}}_{top}(\phi)$ , respectively.

**Remark 2.21.** Let G be special. Given some  $\mathfrak{o} \in \mathfrak{G}$ , we have  $\mathfrak{o} \succeq \mathfrak{o}'$  for all  $\mathfrak{o}' \in \mathfrak{g}(\phi)$  if and only if  $\mathfrak{o} \succeq \overline{\mathfrak{o}}_{top}(\phi)$ . Indeed, the backwards implication follows from Lemma 2.19. As to the forwards one, if  $\mathfrak{o} \succeq \overline{\mathfrak{o}}_{top}(\phi)$ , then there exists a non-principal ultrafilter  $\omega$  such that  $\mathfrak{o} \prec_{\omega} \overline{\mathfrak{o}}_{top}(\phi)$  (Remark 2.14). By the proof of Lemma 2.20, there exists an element  $g \in G$  with  $\|\phi^n(g)\| \sim_{\omega} \overline{\mathfrak{o}}_{top}(\phi)$ , and hence  $\mathfrak{o} \prec_{\omega} \|\phi^n(g)\|$ . Thus,  $\mathfrak{o} \succeq \mathfrak{o}'$  for  $\mathfrak{o}' := [\|\phi^n(g)\|] \in \mathfrak{g}(\phi)$ .

By the same argument, given  $\mathfrak{o} \in \mathfrak{G}$ , we have  $\mathfrak{o} \succeq \mathcal{O}$  for all  $\mathcal{O} \in \mathcal{G}(\varphi)$  if and only if  $\mathfrak{o} \succeq \overline{\mathcal{O}}_{top}(\varphi)$ .

2.2.4. Operations on growth rates. Given two growth rates  $[a_n], [b_n] \in \mathfrak{G}$ , the sum  $[a_n] + [b_n] := [a_n + b_n]$  is a well-defined growth rate. The sum of two  $\omega$ -growth rates is also a well-defined  $\omega$ -growth rate. For  $k \in \mathbb{N}$ , we define  $k * [a_n] := [a_{kn}]$ , that is, the equivalence class of the sequence  $n \mapsto a_{kn}$ . Similarly, we set  $\frac{1}{k} * [a_n] := [a_{\lfloor \frac{n}{k} \rfloor}]$ . The operation \* is only well-defined on  $\mathfrak{G}$ ; it does not descend to an operation on  $\mathfrak{G}_{\omega}$ .

For any finitely generated group G and any  $\psi \in \text{Out}(G)$ , we have the following identities

$$\overline{\mathfrak{o}}_{top}(\psi^k) \sim k * \overline{\mathfrak{o}}_{top}(\psi), \qquad \overline{\mathfrak{o}}_{top}(\psi) \sim \frac{1}{k} * \overline{\mathfrak{o}}_{top}(\psi^k),$$

for all  $k \in \mathbb{N}$ . (The second formula uses Lemma 2.15.) Analogous identities hold for elements of  $\operatorname{Aut}(G)$  and  $\overline{\mathcal{O}}_{top}(\cdot)$ .

2.3. **Degenerations.** Let G be a special group. An infinite sequence in Out(G) and a choice of a non-principal ultrafilter  $\omega$  give rise to an isometric action on a median space  $G \curvearrowright \mathcal{X}_{\omega}$ , which equivariantly embeds in a finite product of  $\mathbb{R}$ -trees. We refer to such actions as *degenerations*.

The purpose of Section 2.3 is to briefly recall how these are constructed (the Bestvina–Paulin construction [Bes88, Pau91]) and to collect many of their properties shown in [Fio23, Fio24]. We begin in a general setting and progressively introduce assumptions, to show where these are needed.

2.3.1. Limits of sequences of homomorphisms. To begin with, let G and A be finitely generated groups. Let X be a locally finite Cayley graph of A.

Let  $\operatorname{Rep}(G, A)$  be the quotient of the set of homomorphisms  $\operatorname{Hom}(G, A)$  by the A-action given by post-composition with inner automorphisms of A. For instance, we have  $\operatorname{Rep}(G, G) = \operatorname{Out}(G)$ .

Consider a sequence  $\phi_n \in \operatorname{Rep}(G, A)$ . If  $\varphi_n$  is any sequence of lifts of  $\phi_n$  to  $\operatorname{Hom}(G, A)$ , we obtain a sequence of actions  $G \curvearrowright_{\varphi_n} X$ , in which each  $g \in G$  acts on X as the element  $\varphi_n(g) \in A$ . If  $S \subseteq G$ is a finite generating set, we can then consider the displacement parameters  $\tau_n^S$  of this sequence of G-actions, as defined in Section 2.2.2. The value of  $\tau_n^S$  only depends on the element  $\phi_n$  and is unaffected by the choice of the specific representative  $\varphi_n$ . Moreover, if  $T \subseteq G$  is another generating set, the abstract growth rates  $[\tau_n^S], [\tau_n^T] \in \mathfrak{G}$  coincide. For this reason, we will simply write " $\tau_n$ " in the coming discussion when the choice of generating set is irrelevant.

The sequence  $\tau_n$  is bounded if and only if the sequence  $\phi_n$  is *finite*, meaning that the  $\phi_n$  all lie in some finite subset of Rep(G, A).

Now, consider an infinite sequence  $\phi_n \in \operatorname{Rep}(G, A)$  and fix a non-principal ultrafilter  $\omega$  on  $\mathbb{N}$ . Pick basepoints  $x_n \in X$  that almost realise the  $\tau_n$ , in the sense that  $\max_{s \in S} d(\varphi_n(s)x_n, x_n) \leq 2\tau_n$ . Then define the isometric G-action

$$(G \curvearrowright \mathcal{X}_{\omega}) := \lim_{\omega} \left( G \curvearrowright_{\varphi_n} \frac{1}{\tau_n} X \right),$$

where  $\frac{1}{\tau_n}X$  denotes the Cayley graph X with its metric rescaled by  $\tau_n$ , and we use the sequence of basepoints  $x_n$  to select the correct component of the ultraproduct. We refer the reader to [DK18, Chapter 10] for basics on ultralimits of spaces and actions.

**Definition 2.22.** We refer to the action  $G \curvearrowright \mathcal{X}_{\omega}$  as the *degeneration* determined by the sequence  $\phi_n \in \operatorname{Rep}(G, A)$ , the ultrafilter  $\omega$ , and the sequence of basepoints  $x_n$ .

Strictly speaking, the choice of generating set S also affects the scaling parameters  $\tau_n^S$  and hence the degeneration  $\mathcal{X}_{\omega}$ , albeit only by a G-equivariant homothety, which we can disregard. The choice of representatives  $\varphi_n$  has no effect, up to G-equivariant isometries.

One of the most important properties of degenerations is that, by construction, no point of  $\mathcal{X}_{\omega}$  is fixed by G. This is precisely what motivates the choice of the  $\tau_n$  as scaling factors.

2.3.2. Homomorphisms into RAAGs. Let now A be a right-angled Artin group  $A_{\Gamma}$ .

We will always endow  $A_{\Gamma}$  with its standard generating set, so that the corresponding Cayley graph is naturally identified with the universal cover  $\mathcal{X}_{\Gamma}$  of the Salvetti complex, equipped with the  $\ell^1$ -metric coming from its cubical structure.

The cube complex  $\mathcal{X}_{\Gamma}$  admits a natural  $A_{\Gamma}$ -equivariant embedding into a finite product of simplicial trees  $T^v$ . This embedding preserves distances if we endow  $\mathcal{X}_{\Gamma}$  and the product with their  $\ell^1$ -metrics. Specifically, the trees  $A_{\Gamma} \curvearrowright T^v$  are the Bass–Serre trees of the HNN splittings

$$A_{\Gamma} = A_{\Gamma \setminus \{v\}} *_{A_{\mathrm{lk}(v)}}$$

where the stable letter of the HNN extension is identified with the standard generator v. Thus, there is one of these trees for each vertex  $v \in \Gamma$ . Equivalently,  $T^v$  can be viewed as the tree dual to the set of hyperplanes of  $\mathcal{X}_{\Gamma}$  labelled by v. Composing the embedding  $\mathcal{X}_{\Gamma} \hookrightarrow \prod_{v \in \Gamma} T^v$  with the factor projections, we obtain  $A_{\Gamma}$ -equivariant cellular maps  $\pi_v \colon \mathcal{X}_{\Gamma} \to T^v$ .

All this additional structure gets transferred to any degeneration. Let  $G \curvearrowright \mathcal{X}_{\omega}$  be the degeneration associated to a finitely generated group G, an infinite sequence  $\phi_n \in \operatorname{Rep}(G, A_{\Gamma})$  and an ultrafilter<sup>6</sup>  $\omega$ , along with a suitable choice of basepoints  $x_n$ . The results of [Bow13, Bow16] imply:

**Proposition 2.23.** The space  $\mathcal{X}_{\omega}$  is a complete, geodesic, finite-rank median space. It embeds G-equivariantly and isometrically in a finite product of  $\mathbb{R}$ -trees  $\prod_{v \in \Gamma} T_{\omega}^v$ , endowed with the  $\ell^1$ -metric. The action  $G \curvearrowright \mathcal{X}_{\omega}$  has unbounded orbits.

More precisely, the action  $G \curvearrowright T^v_{\omega}$  is the ultralimit of the actions  $G \curvearrowright_{\varphi_n} \frac{1}{\tau_n} T^v$ , where we use the sequence  $\pi_v(x_n)$  as basepoints, and where  $\tau_n$  is the sequence of displacement factors for the actions  $G \curvearrowright_{\varphi_n} \mathcal{X}_{\Gamma}$ . It is perfectly possible for G to be elliptic in the tree  $T^v_{\omega}$  for some values of  $v \in \Gamma$ , but not for all.

We have the following straightforward connection between degenerations and growth rates of outer automorphisms. In particular, this completes the proof Lemma 2.20 above.

**Lemma 2.24.** Consider a convex-cocompact embedding  $i: G \hookrightarrow A_{\Gamma}$ , an infinite sequence  $\phi_n \in Out(G)$ , and an ultrafilter  $\omega$ . Consider  $i \circ \phi_n \in Rep(G, A_{\Gamma})$ , with displacement parameters  $\tau_n$ , and let  $G \curvearrowright \mathcal{X}_{\omega}$  be a degeneration determined by it. Then, for any element  $g \in G$ , we have (computing conjugacy lengths in G):

(1) if  $\ell_{\mathcal{X}_{\omega}}(g) > 0$ , the  $\omega$ -growth rates  $[\|\phi_n(g)\|]$  and  $[\tau_n]$  coincide in  $\mathfrak{G}_{\omega}$ ;

(2) if  $\ell_{\mathcal{X}_{\omega}}(g) = 0$ , we have  $[\|\phi_n(g)\|] \prec_{\omega} [\tau_n]$  instead.

Moreover, there do exist elements  $g \in G$  with  $\ell_{\mathcal{X}_{\omega}}(g) > 0$ .

Proof. If  $\|\cdot\|_{A_{\Gamma}}$  denotes conjugacy lengths with respect to the standard generating set of  $A_{\Gamma}$ , we have  $\ell_{\mathcal{X}_{\Gamma}}(h) = \|h\|_{A_{\Gamma}}$  for all  $h \in A_{\Gamma}$ . If  $\|\cdot\|$  denotes conjugacy lengths with respect to any finite generating set of G, there exists a constant C such that  $\frac{1}{C}\|\cdot\| \leq \|\cdot\|_{A_{\Gamma}} \leq C\|\cdot\|$  on G; this follows from Remark 2.17 and it is the only point of the proof where we use convex-cocompactness. Now, we have  $\ell_{\mathcal{X}_{\omega}}(g) = \lim_{\omega} \frac{1}{\tau_n} \ell_{\mathcal{X}_{\Gamma}}(i\phi_n(g))$ , for instance by [Fio24, Lemma 7.9(2)] (the inequality  $\geq$  is trivial, but  $\leq$  needs some properties of right-angled Artin groups so that limits of axes do not disappear in  $\mathcal{X}_{\omega}$ ). Parts (1) and (2) immediately follow, since we have seen that  $[\ell_{\mathcal{X}_{\Gamma}}(i\phi_n(g))] \sim [\|\phi_n(g)\|]$ .

<sup>&</sup>lt;sup>6</sup>We will improperly use the term "ultrafilter" with the meaning of "non-principal ultrafilter" throughout the rest of the article, in order to avoid unnecessarily heavy language. Principal ultrafilters clearly are of no use to us.

The existence of elements  $g \in G$  with  $\ell_{\mathcal{X}_{\omega}}(g) > 0$  is due to the fact that  $G \curvearrowright \mathcal{X}_{\omega}$  does not have a global fixed point and thus has loxodromic elements (for instance, using Serre's lemma [Ser03, p. 64] in the  $\mathbb{R}$ -trees  $T_{\omega}^{v}$ ).

In conclusion, in degenerations determined by sequences in Out(G) for a special group G, loxodromic elements are precisely those whose conjugacy lengths in G grow the fastest under the automorphisms (modulo the ultrafilter), and all strictly slower-growing elements are elliptic.

2.3.3. Degenerations of special groups. Let  $G \leq A_{\Gamma}$  be a convex-cocompact subgroup in the rest of the section. Degenerations  $G \curvearrowright \mathcal{X}_{\omega}$  originating from infinite sequences in  $\operatorname{Out}(G)$  have strong properties, which we now review. Most of this was shown in [Fio23].

We will be mainly interested in analysing the  $\mathbb{R}$ -trees  $G \cap T^v_{\omega}$  with  $v \in \Gamma$ , in whose product  $\mathcal{X}_{\omega}$  is embedded by Proposition 2.23. As mentioned, G can be elliptic in some of these trees and, even when it is not, the action will typically not be minimal (i.e. it will have proper invariant subtrees). What matters is that these trees have extremely well-behaved arc-stabilisers — they are close to being elements of  $\mathcal{Z}(G)$  — which in particular guarantees that the G-action is "stable" in the sense of Bestvina and Feighn [BF95] and we can apply Rips-Sela theory to it.

Given a non-elliptic action on an  $\mathbb{R}$ -tree  $H \curvearrowright T$ , we denote by  $\operatorname{Min}(H, T)$  the H-minimal subtree of T. If  $U \subseteq T$  is a subset, we write  $H_U$  for its H-stabiliser, that is, the subgroup of elements  $h \in H$ with hU = U. By an *arc* in an  $\mathbb{R}$ -tree, we mean a subset homeomorphic to [0, 1] (thus, we do not allow arcs to degenerate into points). If  $\beta$  is an arc, then  $H_\beta$  fixes it pointwise; on the other hand, if  $\alpha$  is a line, then  $H_\alpha$  might translate along it. A subtree  $\tau \subseteq T$  is *stable* if all its arcs have the same G-stabiliser. The action  $H \curvearrowright T$  is *BF*-stable (after [BF95] and [Gui08]) if every arc contains a stable sub-arc.

The following is an important concept in order to understand the trees  $G \curvearrowright T_{\omega}^{v}$ .

**Definition 2.25.** We say that a line  $\alpha \subseteq Min(G, T_{\omega}^{v})$  is *perverse* if all the following hold:

- (1)  $\alpha$  is a stable subtree;
- (2) the action  $G_{\alpha} \curvearrowright \alpha$  is nontrivial, and it does not swap the two ends of  $\alpha$ ;
- (3)  $G_{\alpha} = Z_G(x)$  for some element  $x \in G$ , but the *pointwise* stabiliser of  $\alpha$  does not lie in  $\mathcal{Z}(G)$ .

**Remark 2.26.** We record here the following elementary properties of perverse lines.

- (1) If  $\alpha$  is a perverse line, then all its arcs have the same *G*-stabiliser, which coincides with the pointwise stabiliser of  $\alpha$ . This is the kernel of the nontrivial homomorphism  $\rho: G_{\alpha} \to \mathbb{R}$  given by (signed) translation lengths along  $\alpha$ .
- (2) Distinct perverse lines  $\alpha_1, \alpha_2$  share at most one point. Indeed, if their intersection were to contain an arc  $\beta$ , we would have  $G_{\beta} \leq G_{\alpha_1} \cap G_{\alpha_2}$  since the  $\alpha_i$  are stable. Hence  $G_{\beta} = G_{\alpha_1} \cap G_{\alpha_2}$  and, since the stabilisers  $G_{\alpha_i}$  lie in  $\mathcal{Z}(G)$  by part (3) of the definition, we would have  $G_{\beta} \in \mathcal{Z}(G)$ , contradicting again part (3).
- (3) If  $\alpha$  is a perverse line, we have  $N_G(G_\alpha) = G_\alpha$ . Indeed, the normaliser of  $G_\alpha$  must stabilise the minimal subtree of  $G_\alpha$ , which is  $\alpha$  itself.

The following is the main structural result on the trees  $T^v_{\omega}$  and, consequently, on  $\mathcal{X}_{\omega}$ .

**Theorem 2.27.** Let  $G \leq A_{\Gamma}$  be convex-cocompact,  $\phi_n \in \text{Out}(G)$  an infinite sequence, and  $G \curvearrowright \mathcal{X}_{\omega}$  a resulting degeneration. Consider  $v \in \Gamma$  such that G is non-elliptic in  $T_{\omega}^v$ .

- (1) The action  $G \curvearrowright \operatorname{Min}(G, T_{\omega}^{v})$  is BF-stable.
- (2) For any arc  $\beta \subseteq Min(G, T_{\omega}^{v})$ , either  $G_{\beta} \in \mathcal{Z}(G)$ , or  $\beta$  is contained in a perverse line.
- (3) If the  $\phi_n$  are coarse-median preserving, then there are no perverse lines in  $T^v_{\omega}$ . Moreover, each line  $\alpha \subseteq T^v_{\omega}$  is acted upon discretely by its G-stabiliser.

*Proof.* Almost the entire theorem is already shown in [Fio23]. Part (3) follows from [Fio23, Proposition 5.15(c)]. Part (1) is [Fio23, Corollary 6.13]; in fact, BF–stability is also a consequence of part (2), which implies that arc-stabilisers are co-abelian subgroups of centralisers, a class of subgroups that satisfies the ascending chain condition. Regarding part (2), although the definition of perverse lines does not appear there, everything except stability of  $\alpha$  is explicitly mentioned in [Fio23]: see Proposition 5.12, Proposition 5.15(a) and Remark 5.14.

In order to complete the proof of the theorem, we are only left to show that the lines  $\alpha$  that one gets in part (2) are stable, and thus indeed perverse. This can be quickly deduced from what already appears in the proofs of the above-cited results from [Fio23]; we now briefly recall their setting, but (inevitably) assume that the reader is a little familiar with them.

The action  $G \curvearrowright T_{\omega}^{v}$  is the  $\omega$ -limit of actions  $G \curvearrowright_{\varphi_n} \frac{1}{\tau_n} T^{v}$  for some  $\varphi_n \in \operatorname{Aut}(G)$ . Let  $\beta \subseteq \operatorname{Min}(G, T_{\omega}^{v})$  be an arc such that  $G_{\beta}$  is not a centraliser. Then [Fio23, Proposition 5.12] yields a line  $\alpha \subseteq T_{\omega}^{v}$  such that  $G_{\beta} \leq G_{\alpha}$  and such that  $\alpha$  satisfies parts (2) and (3) of Definition 2.25; in particular, if  $\rho: G_{\alpha} \to \mathbb{R}$  is the homomorphism giving translation lengths along  $\alpha$ , we have  $G_{\beta} = \ker \rho$ . Proposition 5.12 in [Fio23] also tells us that the  $G_{\alpha}$ -minimal subtree for each action  $G \curvearrowright_{\varphi_n} \frac{1}{\tau_n} T^{v}$  is a line  $\alpha_n \subseteq T^{v}$ . These lines  $\alpha_n$  converge to  $\alpha$ . If  $\rho_n: G_{\alpha} \to \mathbb{R}$  gives translation lengths along  $\alpha_n$ , then the  $\rho_n$  converge to  $\rho$ . Since  $\rho$  is nontrivial,  $\omega$ -all  $\rho_n$  are also nontrivial.

Let  $t_n > 0$  be the smallest positive value in the image  $\rho_n(G_\alpha) \subseteq \mathbb{R}$ . Let us first show that  $\lim_{\omega} t_n = 0$ . If this were not the case, there would exist  $\epsilon > 0$  such that, for  $\omega$ -all n, every element of  $G_\alpha$  either fixes  $\alpha_n$  pointwise or translates along it by at least  $\epsilon$ . In this case, each  $g \in G_\beta = \ker \rho$ must fix  $\alpha_n$  for  $\omega$ -all n, so  $G_\beta$  is precisely the  $\omega$ -intersection of the G-fixators of the lines  $\alpha_n$ . Each such fixator is a centraliser in G; this follows from [Fio23, Theorem 4.2], noting that its 2nd case cannot occur as it would imply that  $t_n \leq q/\tau_n \to_\omega 0$ , where q is the uniform constant mentioned in that result. In conclusion,  $G_\beta$  is an  $\omega$ -intersection of centralisers, hence itself a centraliser in G by [Fio23, Proposition 3.38], contradicting our assumptions.

Before continuing, we also remark that the image  $\rho_n(G_\alpha) \subseteq \mathbb{R}$  coincides with the image of the centre of  $G_\alpha$ . This is explained in the proof of [Fio23, Proposition 5.15, part (a)] and also again, more explicitly, in the second half of the proof of part (b) of the same result.

Now that we have shown that  $\lim_{\omega} t_n = 0$ , we can prove that each arc  $\beta' \subseteq \alpha$  has the same G-stabiliser as  $\beta$ , i.e. that  $\alpha$  is stable. Let  $\beta_n, \beta'_n \subseteq \alpha_n$  be sequences of arcs converging to  $\beta$  and  $\beta'$ , respectively. Since  $\lim_{\omega} t_n = 0$ , we can assume without loss of generality that  $\beta'_n = g_n \beta_n$  for some elements  $g_n \in G_{\alpha}$  (where  $g_n$  is acting, as usual, through the action  $G \curvearrowright_{\varphi_n} \frac{1}{\tau_n} T^v$ ). By the previous paragraph, we can actually choose these elements  $g_n$  within the centre of  $G_{\alpha}$ .

Finally, observe that the sequence  $(g_n)$  defines an isometry  $g_{\omega}$  of  $T_{\omega}^v$  mapping each sequence  $(x_n)$  to the sequence  $(g_n x_n)$  (where  $g_n x_n$  is once more meant with respect to the action  $G \curvearrowright_{\varphi_n} \frac{1}{\tau_n} T^v$ ). By construction, we have  $g_{\omega}\beta = \beta'$ . Note that  $g_{\omega}$  will not lie in the image of the homomorphism  $\eta: G \to \text{Isom}(T_{\omega}^v)$  given by the action  $G \curvearrowright T_{\omega}^v$ , but it will still centralise  $\eta(G_{\alpha})$  by construction. It follows that  $G_{\beta'} = g_{\omega}G_{\beta}g_{\omega}^{-1} = G_{\beta}$  as required. This completes the proof of the theorem.

We conclude this section with a comment on the dependence of degenerations  $G \curvearrowright \mathcal{X}_{\omega}$  on the choices of ultrafilter and basepoint sequence.

**Remark 2.28.** Let  $G \leq A_{\Gamma}$  be convex-cocompact and let  $\phi_n \in \text{Out}(G)$  be an infinite sequence.

(1) If G has trivial centre, then the degeneration  $G \curvearrowright \mathcal{X}_{\omega}$  is independent of the choice of basepoints used to define it. This is because G is "uniformly non-elementary" by [Fio24, Theorem I]. Indeed, this property implies that there exists a constant C such that, for every finite generating set  $S \subseteq G$ , if  $x_n, y_n \in \mathcal{X}_{\Gamma}$  are sequences with  $\max_{s \in S} d(\varphi_n(s)x_n, x_n) \leq 2\tau_n$  and  $\max_{s \in S} d(\varphi_n(s)y_n, y_n) \leq 2\tau_n$ , then we have  $d(x_n, y_n) \leq C\tau_n$ . Thus, the sequences  $(x_n)$  and  $(y_n)$  select the same component of the ultraproduct  $\prod_{\omega} \frac{1}{\tau_{\omega}} \mathcal{X}_{\Gamma}$ .

(2) The isometry type of the space  $\mathcal{X}_{\omega}$  is independent of the choice of ultrafilter by [CRHK24]. However, in a degeneration  $G \curvearrowright \mathcal{X}_{\omega}$ , the *G*-action might depend on the ultrafilter, a priori. This will be a source of significant complications later in the paper.

For lack of a better place, we record here the following observation; see e.g. [FK24, Lemma 2.18] for a proof. This is particularly useful when working with groups that are not finitely generated.

**Lemma 2.29.** Let  $G \curvearrowright T$  be an (isometric) group action on an  $\mathbb{R}$ -tree. Suppose that there are no infinite ascending chains of G-stabilisers of rays in T. Then, if  $H \leq G$  is any subgroup, either H fixes a point of T or H contains a loxodromic element.

2.4. Splittings over convex-cocompact subgroups. At many points in the paper, we will consider a graph-of-groups splitting of a special group G whose edge groups are centralisers. We will then seek to understand, and possibly further split, the vertex groups of the graph of groups. A key point is that these vertex groups are again special groups, so that our techniques can again be applied to them. This short subsection is devoted to proving precisely this fact; more generally, convex-cocompactness of edge groups implies convex-cocompactness of vertex groups. This is a typical phenomenon when it comes to "geometric" properties of edge groups (compare e.g. [Bow98, Proposition 1.2] and [HW21]).

We state the result for general cocompact cubulations, as it may be useful elsewhere. A cocompact cubulation  $Q \curvearrowright X$  is a proper cocompact action of a discrete group on a CAT(0) cube complex. As above, we say that a subgroup  $H \leq Q$  is convex-cocompact in the cubulation if H preserves a convex subcomplex of X on which it acts cocompactly.

**Proposition 2.30.** Let  $Q \cap X$  be a cocompact cubulation. If  $Q \cap T$  is a minimal action on a simplicial tree such that Q-stabilisers of edges of T are convex-cocompact in X, then Q-stabilisers of vertices of T are convex-cocompact in X.

Proof. We start by constructing a cocompact cubulation  $Q \curvearrowright Y$  having both actions  $Q \curvearrowright X$  and  $Q \curvearrowright T$  as restriction quotients in the sense of [CS11, p. 860] (also see [Fio24, Section 2.5.1] for details); in other words, X and T can both be obtained from Y by collapsing some Q-orbits of hyperplanes. Such a complex Y can be obtained by applying [Fio23, Proposition 7.9] exactly as in the proof of [FLS24, Theorem 2.17, (3) $\Rightarrow$ (4)]. Omitting some details, one considers the diagonal action  $Q \curvearrowright X \times T$  and finds a Q-invariant, Q-cofinite median subalgebra  $M \subseteq X^{(0)} \times T^{(0)}$  that is mapped onto  $X^{(0)}$  by the factor projection  $X \times T \to X$ . Using Chepoi-Roller duality [Che00, Rol98], the action  $Q \curvearrowright M$  is the action on the 0-skeleton of the required CAT(0) cube complex Y.

Now that we have Y, there is by construction a Q-invariant set  $\mathcal{W}$  of pairwise-disjoint hyperplanes of Y such that its dual tree is Q-equivariantly isomorphic to T. Let  $Y^{\circ}$  be a copy of Y from which we have removed the interiors of the carriers of the hyperplanes in  $\mathcal{W}$  (that is, all open cubes intersecting elements of  $\mathcal{W}$ ). The Q-stabilisers of the vertices of T are precisely the Q-stabilisers of the connected components of  $Y^{\circ}$ . Each of these components is a convex subcomplex of Y and it is acted upon cocompactly by its stabiliser, since  $Q \curvearrowright Y$  is cocompact and  $Y^{\circ} \subseteq Y$  is Q-invariant. This shows that Q-stabilisers of vertices of T are convex-cocompact in Y.

Finally, the equivariant projection  $Y \to X$  takes convex subcomplexes to convex subcomplexes, so any subgroup of Q that is convex-cocompact in Y is also convex-cocompact in X.

We take the chance to make here two loosely related observations, for use in later sections.

**Remark 2.31.** Let G be any group and let  $G \curvearrowright T$  be a minimal action on a tree. If edge-stabilisers are root-closed in G, then vertex-stabilisers are root-closed as well. Indeed, consider  $g \in G$  and  $n \ge 2$  such that  $g^n$  fixes a vertex  $x \in T$ . It follows that g is elliptic, so it fixes a vertex  $y \in T$ . If  $y \ne x$ , let  $e \subseteq T$  be the edge incident to x in the direction of y. Since  $g^n$  fixes both x and y, it must fix e. Now, the fact that  $G_e$  is root-closed implies that  $g \in G_e \le G_x$ , as required.

Recall that  $N_G(H)$  denotes the normaliser in G of a subgroup  $H \leq G$  and, assuming H acts on a tree T, Fix $(H) \subseteq T$  denotes the subtree of H-fixed points.

**Lemma 2.32.** Let  $G \leq A_{\Gamma}$  be convex-cocompact. Let  $G \curvearrowright T$  be a minimal action on a tree with convex-cocompact edge-stabilisers. Then, for any convex-cocompact subgroup  $H \leq G$ , the action  $N_G(H) \curvearrowright \operatorname{Fix}(H)$  is cocompact.

Proof. Since G is finitely generated and acts minimally, the action  $G \curvearrowright T$  is cocompact. Therefore, it suffices to show that, for any edge  $e \subseteq T$ , the normaliser  $N_G(H)$  acts cofinitely on the set of edges of Fix(H) that lie in the G-orbit of e. Thus, consider the set  $\{g \in G \mid ge \subseteq Fix(H)\}$ . If E is the stabiliser of e, this set coincides with  $\{g \in G \mid g^{-1}Hg \leq E\}$ . Since H and E are convex-cocompact, the latter set is a product  $N_G(H) \cdot F \cdot E$  for a finite subset  $F \subseteq G$ , by Lemma 2.4(4). This means that  $N_G(H)$  acts with |F| orbits on the set of H-fixed edges in the G-orbit of e, as we wanted.  $\Box$ 

#### 3. Accessibility over centralisers

In this section we prove the following result. This implies Theorem E from the Introduction, recalling that centralisers are semi-parabolic (Section 2.1.4).

### **Theorem 3.1.** Special groups G are (unconditionally) accessible over G-semi-parabolic subgroups.

We briefly explain our terminology regarding accessibility. An action on a simplicial tree is *minimal* if no proper subtree is left invariant; we always implicitly assume that actions on simplicial trees are without inversions. A *splitting* of a group G over a family of subgroups  $\mathcal{F}$  is a minimal G-action on a simplicial tree T such that T has at least one edge and the G-stabiliser of each edge of T lies in  $\mathcal{F}$ . Importantly (and unusually), we are interested in families  $\mathcal{F}$  that are *not* closed under taking subgroups.

Following [BF91], a splitting  $G \curvearrowright T$  is reduced if there does not exist a vertex  $v \in T$  whose G-stabiliser fixes an incident edge and acts transitively on the remaining incident edges. More weakly, we say that  $G \curvearrowright T$  is *irredundant* if T has no degree-2 vertices, except for those where the vertex-stabiliser swaps the two incident edges; in other words, T was not obtained from a smaller splitting of G simply by subdividing an edge. Reduced splittings are irredundant, but the converse does not hold.

**Definition 3.2.** A group G is accessible over a family of subgroups  $\mathcal{F}$  if there exists a number N(G) such that any reduced splitting of G over  $\mathcal{F}$  has at most N(G) orbits of edges.

We say that G is unconditionally accessible over  $\mathcal{F}$  if the same is true of irredundant splittings.

For instance, finitely presented groups are accessible over small subgroups [BF91, Main Theorem], but already the free group  $F_2$  fails to be unconditionally accessible over cyclic subgroups [BF91, p. 450]. The reason why Theorem 3.1 holds in this form is that semi-parabolics in special groups are more rigid than small subgroups: ascending chains of semi-parabolics have uniformly bounded length (compare Lemma 3.6). In any case, the core of Theorem 3.1 lies in accessibility in the usual sense; the 'unconditional' addendum will just make our life a little easier later in the paper.

Here is a rough idea of the proof of Theorem 3.1. When G is special, every G-semi-parabolic subgroup Q (and in particular, every centraliser) admits a G-parabolic co-abelian subgroup: there exists  $P \in \mathcal{P}(G)$  such that  $P \triangleleft Q$  and Q/P is abelian. One thus hopes to leverage the fact that the quotiented normalisers  $N_G(P)/P$  are accessible over small subgroups, together with the fact that there are only finitely many G-conjugacy classes of elements of  $\mathcal{P}(G)$ . This strategy works provided that, given a reduced splitting  $G \curvearrowright T$ , the induced action on the subtree of P-fixed points  $N_G(P) \curvearrowright \operatorname{Fix}(P)$  is sufficiently close to being minimal and reduced. In order to show that this is indeed the case, we will need to control the normal closure  $\langle \langle P \rangle \rangle_G$  and the pro-p topologies on G will prove rather useful for this (or, alternatively but equivalently, the torsion-free nilpotent quotients of G).

We thus begin by recalling a couple of facts on pro-p topologies in Section 3.1, then prove Theorem 3.1 in Section 3.2 (see Corollary 3.9 and Lemma 3.6). We also prove an alternative accessibility result for special groups, which applies to general edge groups, provided that their parabolic closures are elliptic in the splitting (Corollary 3.10).

3.1. **Pro-p topologies.** Let G be a group and p a prime number. The *pro-p topology* on G has the following as a basis of open sets:

 $\{gN \mid g \in G, N \triangleleft G \text{ and } G/N \text{ is a finite } p\text{-group}\}.$ 

A subgroup  $H \leq G$  is *p*-separable if it is closed in the pro-*p* topology on *G*; equivalently, for every  $g \in G \setminus H$ , there exists a homomorphism  $f: G \to F$  such that *F* is a finite *p*-group and  $f(g) \notin f(H)$ . The group *G* is residually *p*-finite if the trivial subgroup is *p*-separable; equivalently, the identity is the only element of *G* that vanishes in all *p*-group quotients of *G*. The reader can consult [RZ10] for an expanded discussion.

**Remark 3.3.** While every finite-index subgroup of G contains a *normal* finite-index subgroup of G, this statement fails is we replace the two occurrences of the word "finite" with "power-of-p". For instance, the free group  $F_2$  has non-normal subgroups of index 3, which cannot contain any normal subgroups of index  $3^n$ , as maximal proper subgroups of p-groups are always normal. An explicit example is the subgroup  $\langle x, y^3, yxy, yx^{-1}y \rangle$  within  $\langle x, y \rangle \cong F_2$ .

We mention this to emphasise that, unlike the case of the profinite topology, it is important that the above basis for the pro-p topology consists of cosets of *normal* subgroups, rather than cosets of general subgroups of index a power of p.

Our interest in pro-p topologies is due to the following elementary observation.

**Lemma 3.4.** If a proper subgroup H < G is *p*-separable, then *H* is contained in a normal subgroup of *G* of index *p*. In particular, the normal closure of *H* is a proper subgroup of *G*.

*Proof.* The key point is that, if F is a finite p-group and  $F_0$  is a maximal proper subgroup of F, then  $F_0$  is normal in F (and has index p). See for instance [Hal59, Theorem 4.3.2].

Now, given a proper, *p*-separable subgroup H < G, pick an element  $g \in G \setminus H$  and a *p*-group quotient  $\pi: G \twoheadrightarrow F$  with  $\pi(g) \notin \pi(H)$ . In particular,  $\pi(H)$  is a proper subgroup of F and it is contained in an index-*p* subgroup  $F_0 \triangleleft F$ . The preimage  $\pi^{-1}(F_0)$  is the required normal index-*p* subgroup of G containing H.

Let now G be a special group. Haglund and Wise showed that all convex-cocompact subgroups of G are separable in G [HW08, Corollary 7.9]. However, many of these subgroups are not p-separable for any value of p; an example are the subgroups of  $F_2$  described in Remark 3.3. What is important to us is that *retracts*, and in particular G-parabolic subgroups, are p-separable for all p.

# **Lemma 3.5.** If G is a special group, then each $P \in \mathcal{P}(G)$ is p-separable in G for every prime p.

*Proof.* Retracts of residually p-finite groups are p-separable, by an argument analogous to the one described in [HW08, Lemma 9.2]. Moreover, right-angled Artin groups are residually p-finite for all p; for instance, Toinet observed in [Toi13, Theorem 6.1] that this follows from the fact that right-angled Artin groups are residually torsion-free nilpotent [DK92] via [Gru57].

Now, since parabolic subgroups of right-angled Artin groups are clear retracts, they are p-separable. If  $G \leq A_{\Gamma}$  is convex-cocompact, elements of  $\mathcal{P}(G)$  are by definition intersections between G and parabolic subgroups of  $A_{\Gamma}$ , so they are p-separable in G for all p.

3.2. **Proof of accessibility.** The main accessibility argument is explained in Proposition 3.8 below. Before discussing it, we record the following observation showing how to deduce unconditional accessibility from standard accessibility, in the presence of a uniform bound on the length of chains of edge-stabilisers. Recall that chains of G-semi-parabolic subgroups of a special group G indeed have uniformly bounded length (Remark 2.11).

In the coming proofs, we will often speak of *collapses* of G-trees. If  $G \curvearrowright T$  is a splitting and  $\mathcal{E}$  is a G-invariant set of edges, we can form a new tree  $G \curvearrowright T'$  by shrinking each edge in  $\mathcal{E}$  to a point. If  $G \curvearrowright T$  was minimal or reduced, then so is  $G \curvearrowright T'$ . Elliptic subgroups for T are elliptic for T', and edge-stabilisers for T' are edge-stabilisers for T.

**Lemma 3.6.** Let G be a finitely generated group that is accessible over a family  $\mathcal{F}$  of subgroups. If there is a uniform bound on the length of chains of subgroups in  $\mathcal{F}$ , then G is unconditionally accessible over  $\mathcal{F}$ .

Proof. Let  $G \curvearrowright T$  be an irredundant splitting over  $\mathcal{F}$ , and let  $\mathcal{G} := T/G$  be the quotient graph. We denote by  $|\mathcal{G}|$  the number of edges of  $\mathcal{G}$ . Let m be the largest integer such that there exists a chain  $F_1 \leq \cdots \leq F_m$  in  $\mathcal{F}$ . We claim that there exists a collapse  $G \curvearrowright T'$  that is reduced and such that the quotient graph  $\mathcal{G}' := T'/G$  satisfies  $|\mathcal{G}| \leq m|\mathcal{G}'|$ . The lemma immediately follows from this.

To prove the claim, say that a vertex  $v \in \mathcal{G}$  is *bad* if the splitting fails to be reduced at v: this means that v has degree 2 in  $\mathcal{G}$ , and that the vertex group at v coincides with at least one of the two incident edge groups (exactly one, as T is irredundant). There are finitely many edge paths  $\pi_1, \ldots, \pi_k$  in  $\mathcal{G}$  such that the interiors of the  $\pi_i$  are pairwise disjoint, all vertices in the interior of each  $\pi_i$  are bad, and all bad vertices of  $\mathcal{G}$  lie in the interior of some  $\pi_i$ . The endpoints of each  $\pi_i$  are either vertices of degree  $\geq 3$  in  $\mathcal{G}$ , or vertices of degree  $\leq 2$  whose vertex group properly contains all incident edge groups.

Say that an edge of some  $\pi_i$  is a *top edge* if its edge group coincides with both vertex groups placed at its vertices, and a *bottom edge* if its edge group is properly contained in both vertex groups placed at its vertices; we call the remaining edges of  $\pi_i$  transitional. We need the following straightforward observations.

- (1) Let  $\tau \subseteq \pi_i$  be a maximal segment all of whose edges are transitional. As we move along  $\tau$ , we see edge groups strictly increase (or strictly decrease, depending on the direction of movement along  $\tau$ ). Thus,  $\tau$  is preceded by a bottom edge of  $\pi_i$  and followed by a top edge of  $\pi_i$ , unless some of the endpoints of  $\tau$  coincide with endpoints of  $\pi_i$ . Moreover,  $\tau$  contains at most m edges (and so does the union of  $\tau$  with its preceding/following bottom/top edges, when these exist).
- (2) Suppose that some  $\pi_i$  only contains transitional edges. Collapsing all edges of  $\pi_i$  except for one (it does not matter which one), we can remove all bad vertices in the interior of  $\pi_i$ , while the endpoints of  $\pi_i$  remain distinct non-bad vertices of the new graph of groups.
- (3) If we collapse a top or bottom edge of some  $\pi_i$ , then the resulting vertex of the new graph of groups becomes non-bad.

In particular, Observation (3) shows that, up to collapsing all top and bottom edges of the  $\pi_i$ , we can assume that all  $\pi_i$  only contain transitional edges. Observation (2) then shows that we can collapse all edges but one in each of the  $\pi_i$  and thus obtain a collapse of  $\mathcal{G}$  without bad vertices, that is, a reduced graph of groups.

Phrased differently, let  $\mathcal{E}_i$  be a set of edges of  $\pi_i$  obtained by choosing one edge from each of its maximal transitional segments. If  $\mathcal{E}_i$  contains  $n_i$  edges, then  $\pi_i$  has at most  $mn_i$  edges, by Observation (1). We then define a graph  $\mathcal{G}'$  by collapsing all edges of  $\mathcal{G}$  that lie in some  $\pi_i$ , but not in  $\mathcal{E}_i$ ; equivalently, we retain only edges of  $\mathcal{G}$  that either lie in some  $\mathcal{E}_i$  or lie outside  $\pi_1 \cup \cdots \cup \pi_k$ . The previous discussion shows that  $\mathcal{G}'$  corresponds to a reduced graph-of-groups splitting of  $\mathcal{G}$ . Moreover,  $|\mathcal{G}| \leq m|\mathcal{G}|'$  as required. This concludes the proof of the lemma. We denote by  $b_1(\cdot)$  the 1st Betti number of a group or graph, also writing  $b_1(\cdot; \mathbb{K})$  when the field  $\mathbb{K}$  needs specifying. We will use the following estimates, whose proof is standard and omitted.

#### Lemma 3.7.

- (1) Let T be a finite tree with e edges and  $v_i$  vertices of degree i. Then  $e \leq 2v_1 + v_2 3$ .
- (2) Let  $\mathcal{G}$  be a finite graph with no vertices of degree  $\leq 2$ . Then  $\mathcal{G}$  has at most  $3b_1(\mathcal{G}) 3$  edges.

We are now ready to prove the main ingredient of our accessibility results. In order to simplify its statement, we say that a group G is N-accessible over a family of subgroups  $\mathcal{F}$ , for some integer  $N \geq 0$ , if every reduced splitting of G over  $\mathcal{F}$  has at most N orbits of edges.

**Proposition 3.8.** Let G be a finitely generated group with a reduced splitting  $G \curvearrowright T$ . Suppose that there exists a subtree  $U \subseteq T$ , with G-stabiliser denoted  $G_U$ , such that all the following hold:

- (1) the subtrees in the family  $\{gU \mid g \in G\}$  cover T, and distinct ones share at most one point;
- (2)  $G_U$  is p-separable in G for some prime p;
- (3) there exists an integer  $N \ge 0$  such that:
  - (a) either  $G_U$  is elliptic in T and N = 0;
  - (b) or  $G_U$  is N-accessible over the family of stabilisers of edges of U.

Then the quotient T/G has at most  $4b_1(G; \mathbb{F}_p) + N$  edges.

*Proof.* To begin with, note that the actions  $G \curvearrowright T$  and  $G_U \curvearrowright U$  have the same number of edge orbits. Indeed, Condition (1) implies that every edge of T has a G-translate in U, and that two edges of U lie in the same G-orbit if and only if they are in the same  $G_U$ -orbit.

Therefore, our goal in the rest of the proof is to bound the number of edges of the quotient  $U/G_U$ . Unfortunately, we cannot simply appeal to the accessibility of  $G_U$  provided by Condition (3), as the action  $G_U \curvearrowright U$  will badly fail to be minimal or reduced, in general. However, the main point is that this failure must be "located" at vertices where U meets one of its G-translates (since the action  $G \curvearrowright T$  is minimal and reduced), and the number of such vertices can be bounded uniformly in terms of G alone. This is the content of the following claim.

Claim 1. There are at most  $2b_1(G; \mathbb{F}_p)$   $G_U$ -orbits of vertices where U intersects its G-translates. Proof of Claim 1. We will prove the claim by studying an auxiliary splitting of G, which we now describe. Condition (1) says that the family  $\mathcal{Y} := \{gU \mid g \in G\}$  is a transverse covering of T in the sense of [Gui08, Definition 1.4]. As explained there,  $\mathcal{Y}$  gives rise to an action  $G \curvearrowright S$ , where S is the simplicial tree constructed as follows:

- S has a "black" vertex for each element of  $\mathcal{Y}$ ;
- S has a "white" vertex for each point of intersection between distinct subtrees in  $\mathcal{Y}$ ;
- edges of S join the white vertex representing a vertex  $x \in T$  to each black vertex of S representing a subtree in  $\mathcal{Y}$  containing x.

The action  $G \curvearrowright S$  is minimal, since  $G \curvearrowright T$  is.

(3.1)

Since G acts transitively on  $\mathcal{Y}$ , there is a single orbit of black vertices in S. As S is bipartite with respect to the black-white colouring, we can pick a fundamental domain for  $G \curvearrowright S$  that realises G as the fundamental group of the following graph of groups:



This graph has a central black vertex labelled by the group  $G_U$  and a finite number n of adjacent white vertices, labelled by groups  $X_i$ . The first s white vertices are joined to the black vertex by a

unique edge, and the remaining ones by multiple edges. Each vertex group  $X_i$  is the *G*-stabiliser of a vertex of *U* (where *U* intersects one of its *G*-translates) and for  $i \leq s$  each edge group  $Y_i$  is the  $G_U$ -stabiliser of that same vertex of *U*. We do not name the groups labelling the multiple edges.

In particular, note that G-orbits of edges of S are in 1-to-1 correspondence with  $G_U$ -orbits of vertices of T where U intersects one of its G-translates. Thus, the proof of the claim boils down to bounding the size of the quotient graph S/G, which is our next goal.

Since G acts minimally on S, each  $Y_i$  is a proper subgroup of  $X_i$  for  $i \leq s$ . Since  $G_U$  is p-separable in G by Condition (2), each intersection  $Y_i = X_i \cap G_U$  is p-separable in  $X_i$ . Since  $Y_i$  is a proper, p-separable subgroup of  $X_i$ , Lemma 3.4 guarantees the existence of an epimorphism  $\rho_i \colon X_i \to \mathbb{Z}/p\mathbb{Z}$ vanishing on  $Y_i$ .

We can therefore define an epimorphism of G onto a free product F, namely the fundamental group of the following graph of groups with trivial edge groups:



Here the underlying graph is the same as in Splitting 3.1, but we have replaced by the trivial group all edge groups, as well as the black vertex group in the middle, and the left-hand white vertex groups; the right-hand white vertex groups have been replaced by  $\mathbb{Z}/p\mathbb{Z}$ . Concretely, the epimorphism  $G \to F$  vanishes on  $G_U$  and  $X_{s+1}, \ldots, X_n$ , as well as on all edge groups of Splitting 3.1, while it equals  $\rho_i$  on each  $X_i$  with  $i \leq s$ .

If  $\sigma$  is the number of G-orbits of edges in S, and thus also the number of edges in the graph of groups defining F, we have that the rank of the free product F is  $\sigma - (n - s) \ge \frac{1}{2}\sigma$ . Hence  $b_1(G; \mathbb{F}_p) \ge b_1(F; \mathbb{F}_p) \ge \frac{1}{2}\sigma$ , proving Claim 1.

Now, let  $\mathcal{M}_U \subseteq U$  be the  $G_U$ -minimal subtree of U, if  $G_U$  is not elliptic in T; otherwise, define  $\mathcal{M}_U = \{u\}$  for some  $G_U$ -fixed vertex  $u \in U$ . Recall that the action  $G_U \curvearrowright U$  is cocompact, as it has the same number of edge orbits as the action  $G \curvearrowright T$ , which is cocompact because it is a minimal action of a finitely generated group. Thus, U remains within bounded distance of the subtree  $\mathcal{M}_U$ .

Choose finitely many finite subtrees  $\Phi_i \subseteq U$  such that each  $\Phi_i$  intersects  $\mathcal{M}_U$  at a single vertex  $u_i$ , and such that each  $G_U$ -orbit of edges of  $U \setminus \mathcal{M}_U$  intersects the union  $\bigcup \Phi_i$  in a single edge. Let  $\Delta_f$  be the set of vertices that have degree  $\leq 2$  within some  $\Phi_i$  (excluding the base vertex  $u_i$ , if relevant). Let  $\Delta_m \subseteq \mathcal{M}_U$  be a set of representatives for the  $G_U$ -orbits of vertices where the action  $G_U \curvearrowright \mathcal{M}_U$  fails to be reduced: these are vertices whose  $G_U$ -stabiliser fixes one incident edge of  $\mathcal{M}_U$  and acts transitively on the remaining incident edges of  $\mathcal{M}_U$ . If  $G_U$  is elliptic in T, we simply set  $\Delta_m := \emptyset$ .

**Claim 2.** Each vertex in  $\Delta_f \cup \Delta_m$  lies in at least one *G*-translate of *U* other than *U* itself.

Proof of Claim 2. First, note that, by the definition of the  $\Phi_i$ , each degree-1 vertex  $v \in \Phi_i \setminus \{u_i\}$  must also have degree 1 within U. Since T does not have degree-1 vertices, by the minimality of the G-action, it follows that v is incident to an edge of T not contained in U. Hence v lies in a G-translate of U distinct from U, as these translates cover T by assumption.



Second, suppose that some  $w \in \Phi_i \setminus \{u_i\}$  has degree 2 within  $\Phi_i$ . Since  $G_U$  leaves invariant  $\mathcal{M}_U$ , the  $G_U$ -stabiliser of w coincides with the  $G_U$ -stabiliser of the edge of  $\Phi_i$  incident to w in the direction of  $u_i \in \mathcal{M}_U$ , and it acts transitively on the remaining edges of U incident to w, again by the construction of the  $\Phi_i$ . Since the action  $G \curvearrowright T$  is reduced, this again implies that w lies in a G-translate of U distinct from U.

Finally, the same argument applies to the vertices of  $\Delta_m$ , showing that they are also intersection points between distinct translates of U, and concluding the proof of Claim 2.

Now, observing that no two vertices of  $\Delta_f \cup \Delta_m$  are in the same  $G_U$ -orbit, the combination of Claims 1 and 2 shows that  $|\Delta_f \cup \Delta_m| \leq 2b_1(G; \mathbb{F}_p)$ . Let  $\mathcal{E}_{\Phi}$  be the set of edges contained in  $\Phi_i$  for some *i*. Applying Lemma 3.7(1) to the finite tree formed by wedging the  $\Phi_i$  identifying their base vertices  $u_i$ , we see that  $|\mathcal{E}_{\Phi}| \leq 2|\Delta_f|$ , and hence  $|\mathcal{E}_{\Phi} \cup \Delta_m| \leq 4b_1(G; \mathbb{F}_p)$ .

If  $G_U$  is elliptic, this shows that  $U/G_U$  has at most  $4b_1(G; \mathbb{F}_p)$  edges. If  $G_U$  is not elliptic, the above means that, after collapsing at most  $4b_1(G; \mathbb{F}_p)$   $G_U$ -orbits of edges, the action  $G_U \curvearrowright U$  becomes minimal and reduced. This collapse then has at most N edge orbits by Condition (3).

Summing up, the quotient  $U/G_U$  always has at most  $4b_1(G; \mathbb{F}_p) + N$  edges. This is also the number of edges of T/G, finally completing the proof of the proposition.

We now deduce Theorem 3.1 from Proposition 3.8, as well as a further accessibility result that applies to more general edge-stabilisers (Corollary 3.10).

For this we consider a special group G, and we realise G as a convex-cocompact subgroup of a right-angled Artin group  $A_{\Gamma}$ , in order to be able to speak of G-parabolic and G-semi-parabolic subgroups. Recall that every special group G acts by conjugation on the set of G-parabolic subgroups  $\mathcal{P}(G)$ , and the quotient by this action  $\mathcal{P}(G)/G$  is finite by Lemma 2.2. Also recall that, for each  $P \in \mathcal{P}(G)$ , the normaliser  $N_G(P)$  is G-parabolic, and hence the quotient  $N_G(P)/P$  is finitely presented (for instance, because  $N_G(P)$  is finitely presented and P is finitely generated).

By Bestvina and Feighn's accessibility [BF91], each finitely presented group H admits a constant  $\gamma(H)$  such that every reduced splitting of H over abelian subgroups has at most  $\gamma(H)$  edge orbits.

**Corollary 3.9.** Let  $G \leq A_{\Gamma}$  be convex-cocompact. If  $G \curvearrowright T$  is a reduced splitting of G over G-semi-parabolic subgroups, then the number of edges of the quotient T/G is at most:

$$4b_1(G;\mathbb{Q}) \cdot |\mathcal{P}(G)/G| + \sum_{[P] \in \mathcal{P}(G)/G} \gamma(N_G(P)/P).$$

Proof. Recall that each G-semi-parabolic subgroup of G has a parabolic part (Definition 2.9), which lies in  $\mathcal{P}(G)$ . Thus, for each G-conjugacy class of G-parabolic subgroups [P], we can consider the edges of T whose stabiliser has parabolic part in the class [P], and collapse all other edges of T. We obtain a collapse  $G \curvearrowright T_{[P]}$  that is still a reduced splitting, and has gained the property that all its edge-stabilisers are G-semi-parabolic with parabolic part in [P]. Bounding the number of edge orbits in  $T_{[P]}$  for each  $[P] \in \mathcal{P}(G)/G$  gives a bound on the number of edge orbits for the original tree T (namely, the sum of the bounds for the  $T_{[P]}$ ).

Thus, we assume in the rest of the proof that there exists  $P \in \mathcal{P}(G)$  such that all edge-stabilisers of T have parabolic part conjugate to P, and our goal becomes showing that T/G has at most  $4b_1(G;\mathbb{Q}) + \gamma(N_G(P)/P)$  edges. For this, the plan is to invoke Proposition 3.8 for the subtree of P-fixed points  $\operatorname{Fix}(P) \subseteq T$ , so we now proceed to check that its assumptions are satisfied.

Every edge-stabiliser E has a conjugate that has exactly P as its parabolic part (rather than a conjugate of P). In particular, the G-translates of Fix(P) cover P. At the same time, each edge-stabiliser E can only contain a given G-conjugate of P if that is the parabolic part of E, by Proposition 2.8(3). This implies that no two G-translates of Fix(P) share an edge, and also that the G-stabiliser of Fix(P) equals the normaliser  $N_G(P)$ . The latter is G-parabolic in G by Lemma 2.5(2), and hence p-separable in G for all primes p by Lemma 3.5. Finally, consider the action  $N_G(P) \curvearrowright \operatorname{Fix}(P)$  and note that this factors through an action  $N_G(P)/P \curvearrowright \operatorname{Fix}(P)$ , which has free abelian edge-stabilisers. Any reduced splitting of  $N_G(P)/P$  over abelian subgroups has at most  $\gamma(N_G(P)/P)$  orbits of edges by [BF91].

In conclusion, Proposition 3.8 shows that T/G has at most  $4b_1(G; \mathbb{F}_p) + \gamma(N_G(P)/P)$  edges, for all primes p. Choosing p large enough, this bound becomes  $4b_1(G; \mathbb{Q}) + \gamma(N_G(P)/P)$ , since G and its abelianisation are finitely generated. This concludes the proof of the corollary.

Corollary 3.9 implies Theorem 3.1, using Lemma 3.6 to promote accessibility to unconditional accessibility, and recalling that chains of G-semi-parabolic subgroups have uniformly bounded length (Remark 2.11).

With similar techniques, one can prove a more general accessibility result, where edge groups might not be semi-parabolic or even finitely generated; it only requires some restrictions on parabolic closures of edge-stabilisers. Recall that the G-parabolic closure of a subgroup  $H \leq G$  is the smallest G-parabolic subgroup containing H (or the entire G if no such G-parabolic subgroup exists). We will not use this more general result in this article, but we expect it to be useful elsewhere.

**Corollary 3.10.** Let  $G \leq A_{\Gamma}$  be convex-cocompact. Let  $G \curvearrowright T$  be a reduced splitting with arbitrary edge groups. Suppose that, for each edge group E, the G-parabolic closure  $\widehat{E}$  is elliptic in T. Then the number of edges of the quotient T/G is at most:

$$4b_1(G;\mathbb{Q}) \cdot |\mathcal{P}(G)/G| + \sum_{[P] \in \mathcal{P}(G)/G} 3b_1(N_G(P)/P;\mathbb{Q}).$$

*Proof.* Similar to the proof of Corollary 3.9, we can consider, for each G-conjugacy class  $[P] \in \mathcal{P}(G)/G$ , the edges of T whose stabiliser has G-parabolic closure lying in [P], and collapse all other edges of T. We obtain a reduced splitting  $G \curvearrowright T_{[P]}$  in which P is still elliptic, and which has gained the property that all edge-stabilisers have G-parabolic closure in [P]. Again, bounding the number of edge orbits in  $T_{[P]}$  for each  $[P] \in \mathcal{P}(G)/G$  gives a bound on the number of edge orbits for T.

Thus, we assume in the rest of the proof that there exists  $P \in \mathcal{P}(G)$  such that P is elliptic in Tand such that all edge-stabilisers of T have G-parabolic closure conjugate to P. Our goal becomes showing that T/G has at most  $4b_1(G; \mathbb{Q}) + 3b_1(N_G(P)/P; \mathbb{Q})$  edges. Again, the plan is to apply Proposition 3.8 to a suitable subtree.

Let  $U \subseteq T$  be the union of the edges of T whose stabiliser has G-parabolic closure exactly equal to P (rather than to a G-conjugate). We claim that U is connected, and thus a subtree. For this, let  $x \in T$  be a vertex fixed by P. Consider an edge  $e \subseteq U$  and its stabiliser E. Since  $\hat{E} = P$ , the vertex x is fixed by E, and hence the shortest path connecting e to x in T is also entirely fixed by E. This path is thus entirely contained in U; for this, we emphasise that P is not properly contained in any of its G-conjugates, e.g. by Lemma 2.4(3). This shows that U is connected<sup>7</sup> as claimed.

Note that, for any subgroup  $H \leq G$  and any  $g \in G$ , the subgroup  $g\hat{H}g^{-1}$  is the *G*-parabolic closure of  $gHg^{-1}$ . Thus, it is clear from the definition of *U* that the *G*-translates of *U* cover *T*, and that distinct *G*-translates of *U* cannot share an edge. The stabiliser  $G_U$  is precisely the normaliser  $N_G(P)$ , which is *G*-parabolic in *G* by Lemma 2.5(2), and hence *p*-separable in *G* for all primes *p* by Lemma 3.5. If  $N_G(P)$  is not elliptic in *T*, let  $\mathcal{M}$  be its minimal subtree. Since *P* is elliptic in *T*, we have  $\mathcal{M} \subseteq \operatorname{Fix}(P)$ . Since all edge-stabilisers of *T* have *G*-parabolic closure conjugate to *P*, all edges of  $\operatorname{Fix}(P)$  have stabiliser exactly equal to *P*. Thus, the action  $N_G(P) \curvearrowright \mathcal{M}$  factors through an action  $N_G(P)/P \curvearrowright \mathcal{M}$ , which has trivial edge-stabilisers. Note that any reduced free splitting of  $N_G(P)/P$  has at most  $3b_1(N_G(P)/P;\mathbb{F}_p)$  orbits of edges: indeed, if  $\mathcal{G}$  is the quotient graph of the splitting, there is an epimorphism of  $N_G(P)/P \twoheadrightarrow \pi_1(\mathcal{G})$ , hence  $b_1(N_G(P)/P;\mathbb{F}_p) \ge b_1(\mathcal{G})$  and we can invoke Lemma 3.7(2) to bound the number of edges in terms of  $b_1(\mathcal{G})$ .

<sup>&</sup>lt;sup>7</sup>This is the key point where we use the assumptions that G-parabolic closures of edge-stabilisers be elliptic.

Summing up, we can invoke Proposition 3.8 for the subtree  $U \subseteq T$  with  $N = 3b_1(N_G(P)/P; \mathbb{F}_p)$ and conclude that T/G has at most  $4b_1(G; \mathbb{F}_p) + 3b_1(N_G(P)/P; \mathbb{F}_p)$  edges, for all primes p. Choosing p large enough, this bound becomes  $4b_1(G; \mathbb{Q}) + 3b_1(N_G(P)/P; \mathbb{Q})$ . Finally, the bound in the statement of the corollary is obtained by summing this quantity over  $\mathcal{P}(G)/G$ , as discussed in the first paragraph of the proof.

**Remark 3.11.** At first sight, the hypotheses of Corollary 3.10 might seem too weak for the result to be true, as no form of "niceness" is required of the edge-stabilisers (for instance, they are allowed to be infinitely generated). However, one hidden assumption is that the G-parabolic closure of each edge-stabiliser is required to be a *proper* subgroup of G (as it has to be elliptic).

The typical reason why a group is not accessible over arbitrary subgroups is that any epimorphism  $\pi: G \to H$ , where H is an arbitrary group, can be used to pull back splittings of H to splittings of G. However, as we do so, all edge-stabilisers of the resulting splittings of G will contain the normal subgroup ker  $\pi$ . If the special group G does not virtually split as a direct product, then no G-parabolic subgroup can be normal in G, other than the trivial subgroup. Thus, every nontrivial normal subgroup of G has G-parabolic closure that is the entire G, and so does every subgroup of G that contains such a nontrival normal subgroup. For this reason, Corollary 3.10 never applies to such "pulled-back" splittings of G.

### 4. Enhanced JSJ splittings

The goal of this section is to construct, for any 1-ended special group G, a canonical splitting of G over centralisers and cyclic subgroups, relative to maximal subgroups of G that virtually split as direct products (the family  $\mathcal{S}(G)$  of "singular" subgroups). This splitting is a key step in our analysis of automorphisms of G leading to Theorems A and B.

The following is a restatement of Theorem D in the terminology of this section, which we explain right after. This is the main result of this section, together with its strengthening in Theorem 4.28.

**Theorem 4.1.** Let G be a 1-ended special group. There exists an Out(G)-invariant  $(\mathcal{ZZ}(G), \mathcal{S}(G))$ tree  $G \curvearrowright T$  such that the G-stabiliser of each vertex of T is of one of the following kinds:

- (a) a quadratically hanging subgroup<sup>8</sup>, relative to  $\mathcal{S}(G)$ ;
- (b) a convex-cocompact subgroup that is  $(\mathcal{Z}(G), \mathcal{S}(G))$ -rigid in G.

The tree T in Theorem 4.1 is allowed to be a single vertex; when this occurs, it carries the important information that G is either "rigid" or a surface group. The former implies that every element of Out(G) has its top growth rate realised on a singular subgroup of G (Proposition 5.8), which will allow us to analyse growth rates by induction on the complexity of G.

Following [GL17], we say that an action on a simplicial tree  $G \curvearrowright T$  is an  $(\mathcal{A}, \mathcal{H})$ -tree, where  $\mathcal{A}$  and  $\mathcal{H}$  are families of subgroups of G, if the G-stabiliser of each edge of T lies in  $\mathcal{A}$  and if each subgroup in  $\mathcal{H}$  fixes a point of T. We will also say that T is a tree over  $\mathcal{A}$  and relative to  $\mathcal{H}$ , with the same meaning. As above, we speak of a *splitting* when the tree is minimal and not a single point. We again stress that we will **not** consider families  $\mathcal{A}$  that are closed under taking subgroups. This is an important difference to [GL17] and standard JSJ theory, where it is often an implicit standing assumption.

A subgroup  $H \leq G$  is  $(\mathcal{A}, \mathcal{H})$ -rigid in G if it is elliptic in all  $(\mathcal{A}, \mathcal{H})$ -trees of G. A vertex group Q of a G-tree T is quadratically hanging (QH) relative to  $\mathcal{H}$  if we can represent Q as the fundamental group of a compact hyperbolic surface  $\Sigma$  such that all incident edge groups of T, as well as all subgroups of Q contained in elements of  $\mathcal{H}$ , are peripheral (i.e. conjugate into the fundamental

<sup>&</sup>lt;sup>8</sup>We point out that these quadratically hanging vertex-stabilisers are not convex-cocompact in general, as some of the cyclic subgroups that we split along might fail to be convex-cocompact.

groups of the components of  $\partial \Sigma$ , and possibly trivial). In Theorem 4.1, we have  $\partial \Sigma \neq \emptyset$  for all QH vertex groups, except when T is a single point and G is a closed surface group.

A *G*-tree *T* is *invariant* under a subgroup  $\mathcal{O} \leq \text{Out}(G)$  if, for each automorphism  $\varphi \in \text{Aut}(G)$ with outer class in  $\mathcal{O}$ , there exists  $\Phi \in \text{Aut}(T)$  such that  $\Phi(gx) = \varphi(g)\Phi(x)$  for all  $g \in G$  and  $x \in T$ . In Theorem 4.1, Out(G)-invariance simply means that the action  $G \curvearrowright T$  extends to an action  $\text{Aut}(G) \curvearrowright T$  (the group *G* has trivial centre whenever *T* is not a single point).

The family of singular subgroups  $\mathcal{S}(G)$  is studied in Section 4.1 below; it is the family of maximal subgroups of G virtually splitting as direct products of infinite groups. Finally, the family  $\mathcal{ZZ}(G)$  is defined by adding the family Cyc(G) of all infinite cyclic subgroups of G to the family of centralisers:

$$\mathcal{ZZ}(G) := \mathcal{Z}(G) \cup \operatorname{Cyc}(G)$$

**Remark 4.2.** There is a slight mismatch between the properties of edge and vertex groups in Theorem 4.1: We construct a tree  $G \curvearrowright T$  with edge groups in  $\mathcal{ZZ}(G)$  — meaning that we allow cyclic edge groups that are not centralisers — but at the same time only claim that non-QH vertex groups are rigid over  $\mathcal{Z}(G)$ . There is a good reason for this: it is the best one can do if we want rigid vertex groups to be *convex-cocompact*. In turn, convex-cocompactness is essential to carry out many inductive arguments later in the paper.

Here is the plan for Section 4. In Section 4.1, we define and study singular subgroups, namely the elements of the family S(G). Then, in Section 4.2, we develop techniques to construct Out(G)invariant splittings of G over non-small subgroups, particularly over non-cyclic centralisers. Here the main result for special groups is Theorem 4.8, though this material is more broadly applicable. Finally, in Section 4.3, we combine this with classical JSJ theory, in order to canonically split Galso over cyclic subgroups and prove Theorem 4.1 (or rather its strengthening Theorem 4.28).

Afterwards, in Section 5, we will discuss a few immediate consequences of the existence of the enhanced JSJ splitting. In particular, we will connect this splitting to automorphism growth, and we will prove the structural properties of Out(G) mentioned in Theorem C.

4.1. Singular subgroups. Let G be a special group. Let  $\mathcal{VDP}(G)$  be the collection of subgroups of G that virtually split as the direct product of two infinite groups. In this subsection, we are interested in the properties of the maximal elements of  $\mathcal{VDP}(G)$ .

**Definition 4.3.** Let  $\mathcal{S}(G)$  denote the collection of maximal subgroups in  $\mathcal{VDP}(G)$ . We refer to the elements of  $\mathcal{S}(G)$  as the *singular subgroups* of G.

It is clear from the definition that the family  $\mathcal{S}(G)$  is  $\operatorname{Aut}(G)$ -invariant, and we will see below that it consists of finitely many *G*-conjugacy classes of subgroups of *G*. Thus, the family  $\mathcal{S}(G)$  is particularly important because a finite-index subgroup of  $\operatorname{Out}(G)$  preserves the *G*-conjugacy class of each of its elements, which will often allow inductive proofs by induction on complexity. In addition, many (though not all) elements of  $\mathcal{Z}(G)$  are contained in singular subgroups, and this will often motivate us to consider splittings of *G* relative to  $\mathcal{S}(G)$ .

As usual, we fix a realisation of G as a convex-cocompact subgroup  $G \leq A_{\Gamma}$  of a right-angled Artin group. This allows us to speak, for instance, of G-parabolic subgroups. Before discussing singular subgroups in general, we need to focus on the following particular kind.

A subgroup  $H \leq G$  is *isolated* if we have  $Z_G(h) \leq H$  for all  $h \in H \setminus \{1\}$ . We are mainly interested in isolated *abelian* subgroups. Isolated abelian subgroups of rank 1 are plentiful: they are exactly maximal cyclic subgroups generated by a contracting element of G. By contrast, there are only finitely many conjugacy classes of isolated abelian subgroups of rank  $\geq 2$ , as the next result shows. Recall that  $\mathcal{P}(G)$  denotes the collection of G-parabolic subgroups.

**Lemma 4.4.** For every isolated abelian subgroup  $A \leq G$  of rank  $\geq 2$ , we have  $A \in \mathcal{P}(G) \cup \{G\}$ .

*Proof.* Let  $A \leq G$  be an isolated abelian subgroup of rank  $\geq 2$ . Conjugating G by an element of  $A_{\Gamma}$ , we can assume that all elements of A are cyclically reduced in  $A_{\Gamma}$ .

Let  $a, b \in A$  be elements with  $\langle a, b \rangle \cong \mathbb{Z}^2$ . Up to replacing a within  $Z_G(a) = A$  and replacing b within  $Z_G(b) = A$ , we can assume the subgroups  $\langle a \rangle$  and  $\langle b \rangle$  are convex-cocompact. The latter implies that  $Z_{A_{\Gamma}}(a) = \langle \overline{a} \rangle \times A_{\Delta_1}$  and  $Z_{A_{\Gamma}}(b) = \langle \overline{b} \rangle \times A_{\Delta_2}$  for some  $\overline{a}, \overline{b} \in A_{\Gamma}$  with  $a \in \langle \overline{a} \rangle$  and  $b \in \langle \overline{b} \rangle$ ; for all this, see e.g. [Fio23, Remark 3.7(1)–(6)]. Now, choose minimal subgraphs  $\Delta'_i \subseteq \Delta_i$  such that  $Z_G(a) = G \cap (\langle \overline{a} \rangle \times A_{\Delta'_1})$  and  $Z_G(b) = G \cap (\langle \overline{b} \rangle \times A_{\Delta'_2})$ . Thus, each element of  $\Delta'_1 \cup \Delta'_2$  is required in order to write some element of  $A = Z_G(a) = Z_G(b)$ .

Since  $\langle a \rangle$  and  $\langle b \rangle$  are convex-cocompact, incommensurable and commute, we have  $a \in A_{\Delta'_2}$  and  $b \in A_{\Delta'_1}$ . It follows that the union  $\Delta'_1 \cup \Delta'_2$  must split as a join  $\Lambda_1 * \Lambda_2 * (\Delta'_1 \cap \Delta'_2)$ , where  $\Lambda_1$  and  $\Lambda_2$  are the standard generators of  $A_{\Gamma}$  required to write a and b, respectively. Since the subgroups  $A_{\Lambda_1}$ ,  $A_{\Lambda_2}$  and  $A_{\Delta'_1 \cap \Delta'_2}$  all commute with A, so does  $A_{\Delta'_1 \cup \Delta'_2}$ . In conclusion  $A = G \cap A_{\Delta'_1 \cup \Delta'_2}$ , showing that A is G-parabolic as required (except when G = A).

The following proposition collects the main properties of the family  $\mathcal{S}(G)$ . In particular, part (6) should motivate why we call its elements "singular" (viewing, as customary, the contracting directions in the group as "regular").

## **Proposition 4.5.** The following hold.

- (1) Every element of  $\mathcal{VDP}(G)$  is contained in an element of  $\mathcal{S}(G)$ .
- (2) All elements of  $\mathcal{S}(G)$  are G-parabolic, unless  $G \in \mathcal{VDP}(G)$  (in which case  $\mathcal{S}(G) = \{G\}$ ). In particular,  $\mathcal{S}(G)$  contains only finitely many G-conjugacy classes of subgroups.
- (3) We have  $\mathcal{S}(G) = \emptyset$  if and only if G is Gromov-hyperbolic.
- (4) We have  $N_G(S) = S$  for all  $S \in \mathcal{S}(G)$ .
- (5) Each element of  $\mathcal{S}(G)$  is either an isolated abelian subgroup of G of rank  $\geq 2$ , or it is the normaliser in G of the parabolic part of a non-abelian element of  $\mathcal{Z}(G)$ .
- (6) An element  $g \in G \setminus \{1\}$  lies in a subgroup in  $\mathcal{S}(G)$  if and only if  $Z_G(g) \ncong \mathbb{Z}$ . This occurs exactly when g is not contracting in G.

*Proof.* To begin with, consider a subgroup  $H \leq G$  that splits as a nontrivial product  $H_1 \times H_2$ . We claim that H is contained either in an isolated abelian subgroup of G of rank  $\geq 2$ , or in the normaliser in G of the parabolic part of a non-abelian element of  $\mathcal{Z}(G)$ . Note that these two kinds of subgroups of G are either G-parabolic or equal to G, by Lemma 4.4 and Lemma 2.5(2). Moreover, they lie in  $\mathcal{VDP}(G)$  by Lemma 2.4(2) (the parabolic part of a non-abelian centraliser is nontrivial).

To prove our claim, let  $P_1$  be the parabolic part of the centraliser  $Z_G(H_1)$ . The subgroup  $H_2$ normalises  $P_1$  as  $H_2 \leq Z_G(H_1)$ ; at the same time, the subgroup  $H_1$  commutes with  $Z_G(H_1)$  and thus it also normalises  $P_1$ . This shows that  $H \leq N_G(P_1)$ . If  $Z_G(H_1)$  is non-abelian, this proves our claim. If instead  $Z_G(H_1)$  is abelian, then  $H_2$  is abelian and it follows that we have  $H \leq Z_G(H_2) \in \mathcal{Z}(G)$ . In this case, let Z be a maximal element of  $\mathcal{Z}(G)$  containing H. If Z is non-abelian and P is its parabolic part, we have  $H \leq N_G(P)$ , again proving our claim. Finally, suppose that Z is abelian. This implies that  $Z \leq Z_G(g)$  for each  $g \in Z$  and, by maximality of Z, we actually have  $Z = Z_G(g)$ for  $g \neq 1$ . In other words, Z is an isolated abelian subgroup, and it must have rank  $\geq 2$  because it contains the product H. This proves our claim.

In conclusion, each element of  $\mathcal{VDP}(G)$  is virtually contained in another element of  $\mathcal{VDP}(G)$  that has one of the two forms mentioned in part (5) of the proposition, and the latter are G-parabolic (unless  $G \in \mathcal{VDP}(G)$ ). Since G-parabolic subgroups are root-closed, it follows that each element of  $\mathcal{VDP}(G)$  is *entirely* contained in such a G-parabolic element of  $\mathcal{VDP}(G)$ . Since chains of Gparabolic subgroups have bounded length (e.g. by Lemma 2.2 and Lemma 2.4(3)), we immediately obtain parts (1), (2) and (5) of the proposition. The rest of the proposition is routine. We have  $S(G) = \emptyset$  if and only if  $\mathcal{VDP}(G) = \emptyset$ , which occurs if and only if G does not contain  $\mathbb{Z}^2$  as a subgroup. For special groups, this is equivalent to Gromov-hyperbolicity [Gen21a]. If  $S \in S(G)$ , we have  $S \leq N_G(S)$  and  $N_G(S) \in \mathcal{VDP}(G)$  by Lemma 2.4(2); thus, maximality implies that  $N_G(S) = S$ . Finally, regarding part (6), we discussed in Section 2.1 how an element  $g \in G \setminus \{1\}$  is contracting if and only if  $Z_G(g) \cong \mathbb{Z}$ . If g lies in an element of  $\mathcal{VDP}(G)$ , it is clear that these equivalent conditions are not satisfied. Conversely, if g does not lie in any element of  $\mathcal{VDP}(G)$ , then the centraliser  $Z_G(g)$  is also not contained in any element of  $\mathcal{VDP}(G)$ . The parabolic part P of  $Z_G(g)$  must be trivial, otherwise we would have  $P \neq \{1\}$  and  $g \in N_G(P) \in \mathcal{VDP}(G)$ . Thus,  $Z_G(g)$  is abelian and the fact that  $Z_G(g) \notin \mathcal{VDP}(G)$ implies that  $Z_G(g) \cong \mathbb{Z}$ . This completes the proof of the proposition.

We conclude Section 4.1 by discussing two more collections of subgroups. In practice, it will often be useful to remove contracting cyclic subgroups from the collection of centralisers. Thus, we write

$$\mathcal{Z}_c(G) := \{ Z_G(g) \mid g \in G, \ Z_G(g) \cong \mathbb{Z} \},\$$
$$\mathcal{Z}_s(G) := \mathcal{Z}(G) \setminus \mathcal{Z}_c(G).$$

Equivalently,  $\mathcal{Z}_c(G)$  is the collection of maximal cyclic subgroups of G that are contracting. Note that  $\mathcal{Z}_s(G)$  can contain cyclic centralisers, just not any that are centralisers of a single element.

Lemma 4.6. The following hold.

- (1) The collection  $\mathcal{Z}_s(G)$  is  $\operatorname{Aut}(G)$ -invariant and closed under intersections.
- (2) The elements of  $\mathcal{Z}_s(G)$  are precisely the elements of  $\mathcal{Z}(G)$  that are contained in some element of  $\mathcal{S}(G)$  (except for the trivial subgroup, which always lies in  $\mathcal{Z}_s(G)$ ).
- (3) If  $Z \in \mathcal{Z}_s(G)$  and  $Z \neq \{1\}$ , then  $N_G(Z)$  is contained in an element of  $\mathcal{S}(G)$ .

Proof. Part (1) follows from part (2), so we only need to prove the latter. The proof of part (2) is similar to the main argument in the proof of Proposition 4.5, but we recall it here for convenience. Consider a centraliser  $Z \in \mathcal{Z}(G)$ . If Z is non-abelian, then its parabolic part P is nontrivial and we have  $Z \leq N_G(P) \in \mathcal{VDP}(G)$ ; in this case, Z is contained in an element of  $\mathcal{S}(G)$ . If Z is abelian, we have  $Z \leq Z_G(Z)$  and we can consider a maximal element  $Z' \in \mathcal{Z}(G)$  containing Z. If Z' is non-abelian, then Z and Z' are contained in an element of  $\mathcal{S}(G)$  as above. If instead Z' is abelian, then Z' is an isolated abelian subgroup.

This shows that each element  $Z \in \mathcal{Z}(G)$  is contained in an element of  $\mathcal{S}(G)$ , except when it is an isolated abelian subgroup of rank 1. The latter are precisely the centralisers of elements  $g \in G$ with  $Z_G(g) \cong \mathbb{Z}$ , that is, the elements of  $\mathcal{Z}_c(G)$ . This completes the proof of parts (1) and (2).

Part (3) is now immediate from the fact that the normaliser  $N_G(Z)$  has a finite-index subgroup of the form  $Z \times Z'$ , with Z' trivial or infinite, by Lemma 2.4(2).

We will often need to compare centralisers in vertex groups of a splitting of G to centralisers in G. The next result is the main tool for this.

**Lemma 4.7.** Let V be a convex-cocompact<sup>9</sup> vertex group of a tree  $G \curvearrowright T$ .

(1) If 
$$G \curvearrowright T$$
 is a  $(\mathcal{ZZ}(G), \mathcal{S}(G))$ -splitting, then  $\mathcal{Z}_s(V) \subseteq \mathcal{Z}_s(G)$  and  $\mathcal{ZZ}(V) \subseteq \mathcal{ZZ}(G)$ .

(2) If  $G \curvearrowright T$  is a  $(\mathcal{Z}(G), \mathcal{S}(G))$ -splitting, then  $\mathcal{Z}(V) \subseteq \mathcal{Z}(G)$ .

*Proof.* We prove the two parts simultaneously. Let  $G \curvearrowright T$  be a  $(\mathcal{ZZ}(G), \mathcal{S}(G))$ -splitting. Consider a centraliser  $Z \in \mathcal{Z}(V)$  and a subset  $B \subseteq V$  such that  $Z = Z_V(B)$ . We can assume that  $Z \neq \{1\}$ .

Suppose first that the centraliser  $Z_G(B) \in \mathcal{Z}(G)$  does not lie in  $\mathcal{Z}_s(G)$ . Thus,  $Z_G(B)$  is a contracting cyclic subgroup of G, and hence  $Z = V \cap Z_G(B)$  is a contracting cyclic subgroup of V. In particular, we have  $Z \notin \mathcal{Z}_s(V)$  and  $Z \in \text{Cyc}(G) \subseteq \mathcal{ZZ}(G)$ , showing part (1) in this case. Part (2)

<sup>&</sup>lt;sup>9</sup>Convex-cocompactness is automatic for splittings over  $\mathcal{Z}(G)$ , by Proposition 2.30, but not for those over  $\mathcal{ZZ}(G)$ .

follows by observing that, if edge groups of T lie in  $\mathcal{Z}(G)$ , then vertex groups are root-closed by Remark 2.31; hence, the cyclic group  $Z_G(B)$  equals  $Z = V \cap Z_G(B)$ .

Now, suppose instead that  $Z_G(B) \in \mathcal{Z}_s(G)$ . Thus,  $Z_G(B)$  is contained in an element of  $\mathcal{S}(G)$ , and hence it is elliptic in T. If  $Z_G(B)$  fixes a vertex  $v \in T$  of which V is the stabiliser, then  $Z = Z_G(B) \in \mathcal{Z}_s(G)$ . Otherwise  $Z = V \cap Z_G(B)$  fixes an edge of T incident to v, and hence Zis the intersection between  $Z_G(B)$  and an edge-stabiliser of T. In this case,  $Z \in \mathcal{Z}(G)$  if edgestabilisers of T lie in  $\mathcal{Z}(G)$ , and  $Z \in \mathcal{ZZ}(G)$  if edge-stabilisers of T lie in  $\mathcal{ZZ}(G)$ . The latter also shows that  $Z \in \mathcal{Z}_s(G)$  unless  $Z \cong \mathbb{Z}$ .

Thus, we are left to show that  $Z \in \mathcal{Z}_s(G)$  when  $Z \cong \mathbb{Z}$  and  $Z \in \mathcal{Z}_s(V)$ . For this, it suffices to note that  $Z = Z_V(B) = \bigcap_{b \in B} Z_V(b)$ , where each  $Z_V(b)$  again lies in  $\mathcal{Z}_s(V)$ . Now, we have  $Z_V(b) \not\cong \mathbb{Z}$  for each  $b \in B$ , as these are centralisers of single elements (and not contracting). The previous discussion then shows that all centralisers  $Z_V(b)$  lie in  $\mathcal{Z}_s(G)$  and, since the latter collection is closed under taking intersections by Lemma 4.6(1), this finally shows that  $Z \in \mathcal{Z}_s(G)$ .

Note that there is no simultaneous refinement of the two parts of Lemma 4.7: for  $(\mathcal{ZZ}(G), \mathcal{S}(G))$ -splittings, there can be elements of  $\mathcal{Z}_c(V)$  that do not lie in  $\mathcal{Z}(G)$ .

4.2. Constructing canonical splittings. The goal of this section is to develop a general procedure that, starting with a sufficiently nice splitting  $G \curvearrowright T$  of a group, produces an Out(G)-invariant splitting of G with similar edge groups and elliptic subgroups.

Our construction (Section 4.2.3) shares some similarities with Guirardel and Levitt's construction of the canonical tree of cylinders [GL11, GL17], but an important difference is that we allow edgestabilisers to have a complicated intersection pattern, rather than asking that they be separated into "isolated" families with members of different families intersecting trivially. As a consequence, our construction can be applied to groups that are far from being relatively hyperbolic, and to splittings over non-small subgroups, provided that certain conditions are met.

Throughout the section, we work with a general group G and simply require the splittings to satisfy certain axioms. We believe that this level of generality will prove useful in future for further applications. The main consequence for special groups is the following theorem.

A torsion-free group G is said to be 1–ended relative to a family of subgroups  $\mathcal{H}$  if it does not admit any  $(\{\{1\}\}, \mathcal{H})$ -splittings. When  $\mathcal{H} = \emptyset$ , this is the usual notion of 1–endedness [Sta68].

**Theorem 4.8.** Let G be special and let  $\mathcal{O} \leq \operatorname{Out}(G)$  be a subgroup. Suppose that G is 1-ended relative to an  $\mathcal{O}$ -invariant collection of subgroups  $\mathcal{H}$  that contains  $\mathcal{S}(G)$ . Then G admits an  $\mathcal{O}$ invariant  $(\mathcal{Z}_s(G), \mathcal{H})$ -tree whose vertex groups are  $(\mathcal{Z}_s(G), \mathcal{H})$ -rigid in G.

Taking  $\mathcal{O} = \text{Out}(G)$  and  $\mathcal{H} = \mathcal{S}(G)$ , one gets an Out(G)-invariant splitting of G if G is 1-ended and not  $(\mathcal{Z}_s(G), \mathcal{S}(G))$ -rigid. Recall that the collection  $\mathcal{Z}_s(G)$  was defined at the end of Section 4.1.

**Remark 4.9.** If G is 1-ended<sup>10</sup>, then all  $(\mathcal{Z}_s(G), \mathcal{S}(G))$ -trees of G are acylindrical. Indeed, for any such tree  $G \curvearrowright T$ , there are only finitely many G-conjugacy classes of subgroups of the form  $E \cap E'$ , where E, E' are edge-stabilisers; this follows from Lemma 2.4(5). In addition, for any such intersection, the action  $N_G(E \cap E') \curvearrowright \operatorname{Fix}(E \cap E')$  is elliptic by Lemma 4.6 (if  $E \cap E' \neq \{1\}$ ) and cocompact by Lemma 2.32. As a consequence, there is a bound, depending only on T, on the diameter of all sets  $\operatorname{Fix}(E \cap E')$ , proving acylindricity.

However, a priori,  $(\mathcal{Z}_s(G), \mathcal{S}(G))$ -trees of G are not uniformly acylindrical; they might get less and less acylindrical as the number of G-orbits of edges increases. For this reason, we cannot make do with Sela's acylindrical accessibility [Sel97a] in this section, and we will truly have to appeal to our accessibility result over centralisers (Theorem 3.1).

<sup>&</sup>lt;sup>10</sup>Note that G is 1-ended relative to  $\mathcal{S}(G)$  if and only if G is 1-ended in absolute terms. Indeed, all elements of  $\mathcal{S}(G)$  virtually split as direct products of infinite subgroups, and thus they are elliptic in all free splittings of G.

It is convenient to introduce the following notation, which will repeatedly come up in this article.

**Notation 4.10.** Let  $\mathcal{F}$  be a family of subgroups of G. For any subgroup  $H \leq G$ , there are two natural ways of restricting  $\mathcal{F}$  to a family of subgroups of H. We denote them by:

$$\mathcal{F}|^{H} := \{ K \in \mathcal{F} \mid K \le H \}, \qquad \mathcal{F}|_{H} := \{ K \le H \mid \exists F \in \mathcal{F} \text{ such that } K \le F \}.$$

The following is an important observation on the vertex groups of the tree in Theorem 4.8.

**Remark 4.11.** If V is a vertex group of the G-tree T provided by Theorem 4.8, then V is also  $(Z_s(G)|^V, \mathcal{H}|_V)$ -rigid in itself. Indeed, let  $v \in T$  be a vertex of which V is the stabiliser and let  $\mathcal{E}_v$  be the collection of G-stabilisers of edges of T incident to v. Note that  $\mathcal{E}_v \subseteq \mathcal{Z}_s(G)|^V \subseteq \mathcal{S}(G)|_V \subseteq \mathcal{H}|_V$ , using Notation 4.10. Thus, if V were to admit a  $(Z_s(G)|^V, \mathcal{H}|_V)$ -splitting  $V \curvearrowright U$ , then all elements of  $\mathcal{E}_v$  would be elliptic in U, and hence Lemma 4.13(1) would allow us to construct a refinement  $G \curvearrowright T'$  of T in which  $v \in T$  gets blown up to a copy of U. In particular, T' would be a  $(\mathcal{Z}_s(G), \mathcal{H})$ -splitting of G in which V is not elliptic, contradicting the fact that V is  $(\mathcal{Z}_s(G), \mathcal{H})$ -rigid in G.

In particular, this shows that V is  $(Z_s(V), \mathcal{H}|_V)$ -rigid in itself, as we have seen in Lemma 4.7(1) that  $Z_s(V) \subseteq Z_s(G)|_V^V$ .

We now discuss the construction of canonical splittings. After some preliminary material, Section 4.2.2 is concerned with finding an  $\mathcal{O}$ -invariant deformation space using accessibility. Then Section 4.2.3 — which contains the core of the argument — constructs an  $\mathcal{O}$ -invariant splitting within a (suitable)  $\mathcal{O}$ -invariant deformation space. Finally, in Section 4.2.4, we restrict to special groups and prove Theorem 4.8.

4.2.1. Refinements, collapses and deformation spaces. Before we start, we fix terminology and recall some classical results. Let G be a group.

If  $G \curvearrowright T$  and  $G \curvearrowright S$  are trees, a *G*-equivariant map  $\pi: T \to S$  is a *collapse* if it preserves alignment of triples of vertices; equivalently, *S* is obtained from *T* by simply collapsing some *G*orbits of edges. The tree *S* is called a *collapse* of *T*, and *T* is a *refinement* of *S*.

**Remark 4.12.** If a subgroup  $\mathcal{O} \leq \operatorname{Out}(G)$  preserves the *G*-conjugacy class of a subgroup  $H \leq G$ , we can define a *restriction*  $\mathcal{O}|_H \leq \operatorname{Out}(H)$ : one considers the subgroup  $U \leq \operatorname{Aut}(G)$  formed by automorphisms  $\varphi$  with  $\varphi(H) = H$  and outer class in  $\mathcal{O}$ , then one defines  $\mathcal{O}|_H$  as the image of the composition  $U \to \operatorname{Aut}(H) \to \operatorname{Out}(H)$  given by first restricting to H and then projecting to outer automorphisms. The conjugation action  $N_G(H) \curvearrowright H$  also determines a subgroup  $C_H^G \leq \operatorname{Out}(H)$ , and we always have  $C_H^G \triangleleft \mathcal{O}|_H$ . For each  $\phi \in \mathcal{O}$ , the possible restrictions of  $\phi$  to H form a canonical coset  $\phi|_H \cdot C_H^G$  within  $\mathcal{O}|_H$ .

We will often use the following observation allowing us to blow up to trees certain vertices of a G-tree. Part (1) is classical, while part (2) is a little more subtle.

**Lemma 4.13.** Let  $G \curvearrowright T$  be a minimal action on a simplicial tree. Consider a vertex  $v \in T$  and let  $\mathcal{E}_v$  be the collection of G-stabilisers of edges of T incident to v. Suppose that the stabiliser  $G_v$  admits a minimal action on a tree  $G_v \curvearrowright S$ .

- (1) If all subgroups in  $\mathcal{E}_v$  are elliptic in S, then there exists a minimal tree  $G \curvearrowright T'$  with an equivariant collapse map  $\pi: T' \to T$  such that the preimage  $\pi^{-1}(v)$  is  $G_v$ -equivariantly isomorphic to S, while the preimage  $\pi^{-1}(w)$  is a singleton for all vertices  $w \notin G \cdot v$ .
- (2) Let  $\mathcal{O} \leq \operatorname{Out}(G)$  be a subgroup that preserves the G-conjugacy class of  $G_v$  and restricts to a subgroup  $\mathcal{O}_v \leq \operatorname{Out}(G_v)$ . Suppose that  $G \curvearrowright T$  is  $\mathcal{O}$ -invariant, and that  $G_v \curvearrowright S$  is  $\mathcal{O}_v$ -invariant. Suppose moreover that G and  $G_v$  have trivial centre, and that neither T nor S is a line. Finally suppose that, for each  $E \in \mathcal{E}_v$ , the set of fixed points of E in S has finite diameter. Then, the tree  $G \curvearrowright T'$  in part (1) can be constructed so that it is  $\mathcal{O}$ -invariant.

*Proof.* For part (1), see for instance [GL17, Lemma 4.12].

We prove part (2). Let  $\mathcal{O} \leq \operatorname{Aut}(G)$  be the preimage of  $\mathcal{O}$  under the quotient projection  $\operatorname{Aut}(G) \to \operatorname{Out}(G)$ . The subgroup of inner automorphisms of G is contained in  $\tilde{\mathcal{O}}$  and, since the centre of G is trivial, we can view this as a copy  $G \triangleleft \tilde{\mathcal{O}}$ . Indeed, each element  $\varphi \in \tilde{\mathcal{O}}$  acts by conjugation on the subgroup of inner automorphisms precisely as the automorphism  $\varphi \in \operatorname{Aut}(G)$  acts on G. We will denote the group  $\tilde{\mathcal{O}}$  by the more suggestive " $G \rtimes \mathcal{O}$ ", bearing in mind that this is not a semi-direct product, as  $\mathcal{O}$  does not lift to a subgroup of  $G \rtimes \mathcal{O}$  in general. Since  $G_v$  also has trivial centre, the restriction  $\mathcal{O}_v \leq \operatorname{Out}(G_v)$  similarly lifts to a subgroup  $G_v \rtimes \mathcal{O}_v \leq \operatorname{Aut}(G_v)$ .

Let  $\rho: G \to \operatorname{Aut}(T)$  be the homomorphism corresponding to the action  $G \curvearrowright T$ . Since T is G-minimal and  $T \ncong \mathbb{R}$ , the centraliser of  $\rho(G)$  in  $\operatorname{Aut}(T)$  is trivial. Indeed, if  $f \in \operatorname{Aut}(T)$  is an element in this centraliser, then f cannot be loxodromic, otherwise its axis would be G-invariant, violating the fact that T is not a line; thus f is elliptic and, since its fixed set is G-invariant, minimality of T implies that  $f = \operatorname{id}$ . Now, since the centraliser of  $\rho(G)$  is trivial, each automorphism  $\varphi \in G \times \mathcal{O} \leq \operatorname{Aut}(G)$  admits a unique  $\Phi \in \operatorname{Aut}(T)$  such that  $\Phi(gx) = \varphi(g)\Phi(x)$  for all  $g \in G$  and  $x \in T$ . Uniqueness guarantees that we obtain a homomorphism  $\tilde{\rho}: G \times \mathcal{O} \to \operatorname{Aut}(T)$  extending  $\rho$ . Similarly, since  $S \ncong \mathbb{R}$ , the action  $G_v \curvearrowright S$  extends to an action  $G_v \propto \mathcal{O}_v \curvearrowright S$ .

Now, let  $(G \times \mathcal{O})_v$  be the  $G \times \mathcal{O}$ -stabiliser of the vertex  $v \in T$ . Its intersection with G is precisely the stabiliser  $G_v$ , and we have  $G_v \triangleleft (G \times \mathcal{O})_v$ . Thus, all automorphisms of G lying in  $(G \times \mathcal{O})_v$ leave  $G_v$  invariant and, by restriction, we obtain a homomorphism  $(G \times \mathcal{O})_v \rightarrow G_v \times \mathcal{O}_v$ . The action  $G_v \times \mathcal{O}_v \curvearrowright S$  then induces an action  $(G \times \mathcal{O})_v \curvearrowright S$  extending  $G_v \curvearrowright S$ .

We now wish to apply part (1) of the lemma to the actions  $G \times \mathcal{O} \curvearrowright T$  and  $(G \times \mathcal{O})_v \curvearrowright S$ , so as to produce the required blowup  $G \times \mathcal{O} \curvearrowright T'$ , which can be viewed simply as an  $\mathcal{O}$ -invariant G-tree. For this, we need to check that, for every edge  $e \subseteq T$  incident to v, the stabiliser  $(G \times \mathcal{O})_e$  is elliptic in S. As in the case of the stabiliser of v, we have  $G_e \triangleleft (G \times \mathcal{O})_e$ . By hypothesis, the set of fixed points of  $G_e \in \mathcal{E}_v$  in S has finite diameter, and so it has a unique barycentre. The group  $(G \times \mathcal{O})_e$ preserves the fixed set of its normal subgroup  $G_e$ , and hence it fixes its barycentre. This shows that  $(G \times \mathcal{O})_e$  is indeed elliptic in S, concluding the proof.

**Remark 4.14.** We wish to emphasise that the finite-diameter assumption in part (2) of Lemma 4.13 is truly necessary, as we found this to be a subtle point. For instance, consider the free group  $F_3 = \langle a, b, c \rangle$ , along with the quadratically-growing automorphism  $\varphi \in \operatorname{Aut}(F_3)$  given by  $a \mapsto ab$ ,  $b \mapsto bc, c \mapsto c$ , and its outer class  $\phi \in \operatorname{Out}(F_3)$ . Denote by  $\psi$  the restriction of  $\varphi$  to the subgroup  $\langle b, c \rangle$ , and by  $[\psi]$  its outer class. Referring to the Bass–Serre trees of the three splittings pictured below,  $\phi$  preserves the left-hand splitting of G, and  $[\psi]$  preserves the right-hand splitting of  $\langle b, c \rangle$ , However,  $\phi$  does not preserve the free splitting of G pictured in the middle (otherwise  $\phi$  would have to grow linearly).



To better understand the issue, let  $F_3 \curvearrowright L$  and  $\langle b, c \rangle \curvearrowright R$  be the Bass–Serre trees of the HNN extensions on the left and right, respectively. Both actions have trivial edge-stabilisers and uniquely extend to actions  $F_3 \rtimes \langle g \rangle \curvearrowright L$  and  $\langle b, c \rangle \rtimes \langle g \rangle \curvearrowright R$ , where the element g acts as  $\varphi$  on  $F_3$ , and as  $\psi$  on  $\langle b, c \rangle$ . The subgroup  $\langle b, c \rangle \rtimes \langle g \rangle \leq F_3 \rtimes \langle g \rangle$  is the  $F_3 \rtimes \langle g \rangle$ –stabiliser of a vertex  $v \in L$ . Now, if  $e \subseteq L$  is the edge with endpoints v and  $a^{-1}v$ , we see that the  $F_3 \rtimes \langle \varphi \rangle$ –stabiliser of e is the cyclic subgroup  $\langle bg \rangle$ , which acts loxodromically on R. Thus, part (1) of Lemma 4.13 cannot be applied to the action  $F_3 \rtimes \langle g \rangle \curvearrowright L$  in order to blow up v to a copy of R.

Two G-trees are said to lie in the same *deformation space* (in the sense of [For02, GL17]) if they have the same elliptic subgroups. The deformation space of a tree  $G \curvearrowright T_1$  is said to *dominate* the

deformation space of another tree  $G \curvearrowright T_2$  if every subgroup of G that is elliptic in  $T_1$  is also elliptic in  $T_2$  (we also simply say that  $T_1$  dominates  $T_2$ ).

We will need the following slight improvement<sup>11</sup> on [GL17, Proposition 2.2].

**Lemma 4.15.** Let G be finitely generated, and let  $G \curvearrowright T_1$  and  $G \curvearrowright T_2$  be two splittings. Suppose that the G-stabiliser of each edge of  $T_1$  is elliptic in  $T_2$ , and that the G-stabiliser of each edge of  $T_2$  is elliptic in  $T_1$ . Then there exists a splitting  $G \curvearrowright T$  with the following properties:

- (1) T is a refinement of  $T_1$  that dominates  $T_2$ ;
- (2) a subgroup of G is elliptic in T if and only if it is elliptic in both  $T_1$  and  $T_2$ ;
- (3) the G-stabiliser of each edge of T is either the G-stabiliser of an edge of  $T_1$  or  $T_2$ , or the intersection of an edge-stabiliser of  $T_1$  with an edge-stabiliser of  $T_2$ ;
- (4) T does not have any degree-2 vertices whose stabiliser fixes the two incident edges, except for any that  $T_1$  might have already had and have not been blown up.

Proof. Since G is finitely generated and acts minimally on  $T_1$ , there are only finitely many G-orbits of vertices in  $T_1$ . Let  $v_1, \ldots, v_k$  be representatives of these orbits, and let  $G_i$  denote the G-stabiliser of  $v_i$ . If all  $G_i$  are elliptic in  $T_2$ , then  $T_1$  dominates  $T_2$  and we can simply set  $T := T_1$ . Thus, up to discarding some  $v_i$  and re-indexing, we can suppose that each  $G_i$  is non-elliptic in  $T_2$ . Let  $M_i \subseteq T_2$  be the  $G_i$ -minimal subtree; this exists because  $G_i$  is finitely generated relative to finitely many G-stabilisers of edges of  $T_1$  [Gui08, Lemma 1.11], and the latter are elliptic in  $T_2$ .

Now, since G-stabilisers of edges of  $T_1$  are elliptic in  $T_2$ , we can use Lemma 4.13(1) to blow up each  $v_i \in T_1$  (and each of its G-translates) to a copy of  $M_i \subseteq T_2$ . Call  $G \curvearrowright T$  the resulting refinement of  $T_1$ . It is clear that a subgroup of G is elliptic in T if and only if it is elliptic in both  $T_1$  and  $T_2$ , proving properties (1) and (2). Regarding property (4), consider the collapse map  $\pi: T \to T_1$ . Note that every vertex of  $M_i$  has degree  $\geq 2$  within  $M_i$ . Thus, up to collapsing some edge orbits of  $G_i \curvearrowright M_i$  without altering its deformation space, we can assume that, for every vertex  $x \in \pi^{-1}(v_i)$  that has degree 2 within  $\pi^{-1}(v_i)$  and whose G-stabiliser fixes both incident edges, there exists an edge of  $T \setminus \pi^{-1}(v_i)$  that is attached to x. We are only left to show property (3).

Consider an edge  $e \subseteq T$ . If e projects to an edge of  $T_1$ , then its G-stabiliser  $G_e$  is an edgestabiliser of  $T_1$ . Suppose instead that e gets collapsed to a vertex of  $T_1$ , without loss of generality to some  $v_i$ . Thus, there exists an edge  $f \subseteq M_i \subseteq T_2$  such that  $G_e$  equals the  $G_i$ -stabiliser of f; that is,  $G_e = G_i \cap G_f$ . By assumption,  $G_f$  is elliptic in  $T_1$ . If  $G_f$  fixes the vertex  $v_i$ , we have  $G_f \leq G_i$ and hence  $G_e = G_f$  is an edge-stabiliser of  $T_2$ . Otherwise, denoting by  $g \subseteq T_1$  the edge incident to  $v_i$  in the direction of  $\operatorname{Fix}(G_f) \subseteq T_1$ , we have  $G_i \cap G_f = G_g \cap G_f$ . Thus,  $G_e$  is the intersection of an edge-stabiliser of  $T_2$ , as required.

4.2.2. Finding an invariant deformation space. Let G be a finitely generated group. Given a suitable G-tree and provided that G is suitably accessible, our first goal is to construct an  $\mathcal{O}$ -invariant deformation space.

Expanding on Definition 3.2, we say that G is unconditionally  $(\mathcal{F}, \mathcal{K})$ -accessible, for two collections of subgroups  $\mathcal{F}$  and  $\mathcal{K}$ , if there is a uniform bound on the number of edge orbits in irredundant  $(\mathcal{F}, \mathcal{K})$ -splittings of G; the case when  $\mathcal{K} = \emptyset$  corresponds to the above definition of unconditional accessibility over  $\mathcal{F}$ .

**Lemma 4.16.** Let G be finitely generated. Consider two collections of subgroups  $\mathcal{F}$  and  $\mathcal{K}$  such that  $\mathcal{F} \subseteq \mathcal{K}$  and  $\mathcal{F}$  is closed under finite intersections. If G is unconditionally  $(\mathcal{F}, \mathcal{K})$ -accessible, then:

(1) For any collection  $\{G \curvearrowright T_i\}_{i \in I}$  of  $(\mathcal{F}, \mathcal{K})$ -trees, there exists an  $(\mathcal{F}, \mathcal{K})$ -tree  $G \curvearrowright T$  whose elliptic subgroups are precisely those that are elliptic in all  $T_i$ . Moreover, T can be chosen to refine any of the  $T_i$ .

<sup>&</sup>lt;sup>11</sup>The important difference lies in Item (3), where we declare that edge-stabilisers of T are *intersections* of edge-stabilisers of  $T_1$  and  $T_2$ , rather than arbitrary subgroups of these.

# (2) There is a uniform bound to the length of any sequence $\mathfrak{D}_0, \mathfrak{D}_1, \mathfrak{D}_2, \ldots$ of pairwise distinct deformation spaces of $(\mathcal{F}, \mathcal{K})$ -trees such that $\mathfrak{D}_i$ dominates $\mathfrak{D}_j$ for i > j.

Proof. Consider a sequence  $G \curvearrowright T_n$  of  $(\mathcal{F}, \mathcal{K})$ -trees. Since  $\mathcal{F} \subseteq \mathcal{K}$  and  $\mathcal{F}$  is closed under finite intersections, Lemma 4.15 yields a sequence  $G \curvearrowright T'_n$  of  $(\mathcal{F}, \mathcal{K})$ -trees such that  $T'_0 = T_0$ , each  $T'_{n+1}$ is a refinement of  $T'_n$ , and a subgroup of G is elliptic in  $T'_n$  if and only if it is elliptic in all of  $T_0, \ldots, T_n$ . Moreover, the  $T'_n$  all have a bounded number of orbits of degree-2 vertices whose stabiliser fixes the two incident edges. Since G is unconditionally  $(\mathcal{F}, \mathcal{K})$ -accessible, there is a uniform bound on the number of edge orbits of the  $T'_n$ , and hence the sequence of refinements  $T'_n$  eventually stabilises. This proves part (1).

The proof of part (2) is similar. Suppose that  $T_0, \ldots, T_k$  are  $(\mathcal{F}, \mathcal{K})$ -trees such that  $T_i$  dominates  $T_j$  for i > j, and such that their deformation spaces are pairwise distinct. Up to collapsing edge orbits of  $G \curvearrowright T_0$  without altering the deformation space of  $T_0$ , we can assume that  $T_0$  has no degree-2 vertices whose stabiliser fixes the two incident edges (or that  $T_0$  is a line on which G acts vertex-transitively). Using Lemma 4.15 as in the previous paragraph, we obtain  $(\mathcal{F}, \mathcal{K})$ -trees  $T'_i$  that refine  $T_0$  and lie in the deformation space of  $T_i$ . Each  $T'_{i+1}$  is a proper refinement of  $T'_i$ , as the deformation spaces of  $T_i$  and  $T_{i+1}$  are distinct. The tree  $T'_k$  is then an  $(\mathcal{F}, \mathcal{K})$ -tree with at least k + 1 edge orbits and at most one orbit of degree-2 vertices whose stabiliser fixes the two incident edges. Thus, accessibility sets a uniform bound to the index k, proving part (2).

Now, consider a subgroup  $\mathcal{O} \leq \operatorname{Out}(G)$ . If  $\mathcal{F}$  is a collection of subgroups of G that is closed under taking G-conjugates, we say that  $\mathcal{F}$  is  $\mathcal{O}$ -invariant if  $\mathcal{F}$  is preserved by the lift of  $\mathcal{O}$  to a subgroup of  $\operatorname{Aut}(G)$ . We denote by  $\mathcal{F}_{int}$  the collection of finite intersections of elements of  $\mathcal{F}$ .

**Corollary 4.17.** Let G be finitely generated. Consider a splitting  $G \cap T$  and some  $\mathcal{O} \leq \text{Out}(G)$ . Let  $\mathcal{F}$  be an  $\mathcal{O}$ -invariant family of subgroups that contains the G-stabiliser of each edge of T. Let  $\mathcal{K}$  be an  $\mathcal{O}$ -invariant family of subgroups elliptic in T. If the following two conditions are satisfied:

- (1) all elements of  $\mathcal{F}$  are elliptic in T;
- (2) G is unconditionally  $(\mathcal{F}_{int}, \mathcal{K})$ -accessible;

then there exists a refinement  $G \curvearrowright T'$  of  $G \curvearrowright T$  such that:

- (1)' all G-stabilisers of edges of T' belong to  $\mathcal{F}_{int}$ ;
- (2)' the subgroups of G that are elliptic in T' are those whose entire  $\mathcal{O}$ -orbit is elliptic in T.

Proof. Let  $\mathcal{T}$  be the collection of splittings obtained by twisting  $G \curvearrowright T$  by the elements of  $\mathcal{O}$ . All elements of  $\mathcal{T}$  are  $(\mathcal{F}, \mathcal{F} \cup \mathcal{K})$ -splittings of G, by Condition (1), so they are in particular  $(\mathcal{F}_{\text{int}}, \mathcal{F}_{\text{int}} \cup \mathcal{K})$ -splittings. Now, Lemma 4.16(1) implies that there exists a refinement of  $G \curvearrowright T$  that is an  $(\mathcal{F}_{\text{int}}, \mathcal{K})$ -splitting whose elliptic subgroups are precisely those that are elliptic in all splittings in  $\mathcal{T}$ . This proves the corollary.

4.2.3. Finding an invariant splitting. Consider now a group G with trivial centre, a cocompact<sup>12</sup> splitting  $G \curvearrowright T$ , and a subgroup  $\mathcal{O} \leq \operatorname{Out}(G)$ . Let  $\mathscr{E}$  be the collection of subgroups of G that are elliptic in T, and let  $\mathscr{A}$  be the collection of G-stabilisers of edges of T. Throughout Section 4.2.3, we assume that the following two conditions hold:

- (i) the collection  $\mathscr{E}$  is  $\mathcal{O}$ -invariant (i.e. the deformation space of T is  $\mathcal{O}$ -invariant);
- (*ii*) no element of  $\mathscr{A}$  properly contains one of its *G*-conjugates.

In part of the coming discussion, we will assume that T satisfies the following additional condition, which can be viewed as a weak form of acylindricity, plus a ban on edges with trivial stabiliser:

(*iii*) for every subgroup  $A \in \mathscr{A}$ , the fixed set  $Fix(A) \subseteq T$  has finite diameter<sup>13</sup>.

 $<sup>^{12}</sup>$ Cocompactness is immediate if G is finitely generated, but we make no such assumption.

<sup>&</sup>lt;sup>13</sup>Condition (*iii*) implies that normalisers of elements of  $\mathscr{A}$  are elliptic in T; in fact, the latter is an equivalent condition if G is special and the elements of  $\mathscr{A}$  are convex-cocompact, by Lemma 2.32.

Our goal is to prove:

**Theorem 4.18.** Let G be a group with trivial centre,  $G \curvearrowright T$  a cocompact splitting, and  $\mathcal{O} \leq \text{Out}(G)$ .

- (1) Under Conditions (i)–(iii), there is an  $\mathcal{O}$ -invariant splitting  $G \curvearrowright T'$  whose edge-stabilisers are a subset of the edge-stabilisers of T, and whose elliptic subgroups are the same as for T.
- (2) Under just Conditions (i) and (ii), and supposing that no element of  $\mathscr{A}$  is normal in G, there is an  $\mathcal{O}$ -invariant splitting  $G \curvearrowright T''$  such that:
  - (a) for each edge-stabiliser A" of T", there exists some A ∈ A such that A ≤ A" ≤ N<sub>G</sub>(A) and A"/A is a free factor of N<sub>G</sub>(A)/A (possibly equal to {1} or the entire N<sub>G</sub>(A)/A);
    (b) every elliptic subgroup of T is elliptic in T".

The main statement in Theorem 4.18 is part (1). As we will see, part (2) is significantly easier to show, and it is not needed for this article; still, we expect it to be useful elsewhere.

We begin with a few observations and reductions. In particular, there is a fourth and last condition that we will be interested in:

(\*) no vertex-stabiliser of T fixes an edge of T.

As the next result shows, it suffices to prove both parts of Theorem 4.18 in the case when this additional condition is satisfied. Denote by  $\mathscr{E}_{\max} \subseteq \mathscr{E}$  the subset of maximal elliptic subgroups.

**Lemma 4.19.** If  $G \curvearrowright T$  satisfies Condition (ii), then it admits a collapse  $G \curvearrowright T_*$  such that:

• T<sub>\*</sub> satisfies Condition (\*);

• a subgroup of G is elliptic in  $T_*$  if and only if it lies in  $\mathscr{E}$  or normalises a subgroup in  $\mathscr{E}_{\max}$ . Each of Conditions (i), (ii), (iii) is passed on to  $T_*$  if satisfied by T. Moreover, if T satisfies Condition (iii), then T and  $T_*$  have the same elliptic subgroups.

*Proof.* For simplicity, say that (e, v) is a *bad pair* if  $e \subseteq T$  is an edge,  $v \in e$  is a vertex, and the stabiliser of v fixes e. If w is the other vertex of e, the stabilisers of e, v, w then satisfy  $G_v = G_e \leq G_w$ . We say that (e, v) is of the *first kind* if  $G_e = G_w$ , and of the *second kind* if  $G_e \leq G_w$ . Condition (\*) is satisfied precisely when there are no bad pairs, of any kind.

Consider a bad pair (e, v). If  $g \in G$  is an element such that  $v \in ge$  and  $ge \neq e$ , we cannot have gv = v, so we must have v = gw. By Condition (*ii*), the latter can only happen if  $G_v = G_w$ , namely when the pair is of the first kind. In particular, if the pair is of the second kind, then e is the only edge in its G-orbit to contain the vertex v. In this case, the edges in the orbit  $G \cdot e$  come arranged in pairwise-disjoint subtrees of T of diameter  $\leq 2$ : each of these subtrees is a G-translate of the union of the edges in  $G \cdot e$  that contain w. Collapsing these subtrees to points does not affect which subgroups of G are elliptic, that is, the collection  $\mathscr{E}$  is the set of elliptic subgroups also for the new tree. At the same time, such a collapse strictly reduces the number of edges in the quotient graph T/G, which is finite by the assumption that G acts cocompactly.

Thus, after a finite sequence of collapses leaving the collection of elliptic subgroups unchanged, we can assume that any leftover bad pairs are of the first kind.

Now, let (e, v) be a bad pair and consider the fixed subtree  $\operatorname{Fix}(G_e) \subseteq T$ . Since there are no bad pairs of the second kind, all vertices and all edges of  $\operatorname{Fix}(G_e)$  have the same G-stabiliser, namely  $G_e$ . In particular, for any edge  $f \subseteq T$  intersecting  $\operatorname{Fix}(G_e)$  at a single vertex, we have  $G_f \leq G_e$ . This also shows that  $G_e \in \mathscr{E}_{\max}$  and that the G-stabiliser of  $\operatorname{Fix}(G_e)$  is precisely the normaliser  $N_G(G_e)$ . Finally, for any subgroup  $H \in \mathscr{E}_{\max}$ , either  $\operatorname{Fix}(H)$  is a single vertex and  $N_G(H)$  fixes it, or  $\operatorname{Fix}(H)$  contains an edge, in which case all vertices and edges of  $\operatorname{Fix}(H)$  have stabiliser exactly H, and all edges of  $\operatorname{Fix}(H)$  belong to bad pairs. In particular, we have  $\operatorname{Fix}(H) \cap \operatorname{Fix}(H') = \emptyset$  for all distinct subgroups  $H, H' \in \mathscr{E}_{\max}$ .

Summing up, edges belonging to bad pairs are precisely those fixed by elements of  $\mathscr{E}_{\max}$ . Collapsing all these edges ensures Condition (\*), and a subgroup is elliptic in the resulting tree  $T_*$  if and only if it either lies in  $\mathscr{E}$  or normalises an element of  $\mathscr{E}_{\max}$ .
Finally,  $T_*$  satisfies again Condition (ii) because its edge-stabilisers are a subset of those of T. For the same reason,  $T_*$  satisfies Condition (iii) if T does, as collapses do not increase diameters of fixed sets. If the collection  $\mathscr{E}$  is  $\mathcal{O}$ -invariant, it is clear that  $\mathscr{E}_{\max}$  is also  $\mathcal{O}$ -invariant, and so is the collection of normalisers of its elements. Thus, the deformation space of  $T_*$  is  $\mathcal{O}$ -invariant if that of T is (Condition (i)). If T satisfies Condition (iii) and  $E \in \mathscr{E}_{\max}$ , then, after removing all bad pairs of the second kind, either E fixes a unique vertex or it is the G-stabiliser of an edge, by the above discussion. Thus, Condition (iii) implies that  $N_G(E)$  is elliptic in T, and hence the collapse  $T_*$  lies in the same deformation space as T. This concludes the proof of the lemma.

In view of Lemma 4.19, we will assume that the splitting  $G \curvearrowright T$  satisfies Condition (\*) in the rest of Section 4.2.3. To some extent, this allows us to reconstruct part of the splitting T algebraically, as the next remark and lemma show.

**Remark 4.20.** By Condition (\*), vertex-stabilisers of T are the same as elements of  $\mathscr{E}_{\max}$ , and each of these has a unique fixed point in T.

We stratify the collection of edge-stabilisers  $\mathscr{A}$  by inclusion. That is, we first define  $\mathscr{A}_1 \subseteq \mathscr{A}$  as the subset of maximal elements under inclusion, then inductively define  $\mathscr{A}_{i+1}$  as the set of maximal elements of  $\mathscr{A} \setminus \mathscr{A}_i$ . Since G acts cocompactly on T, the collection  $\mathscr{A}$  consists of finitely many Gconjugacy classes of subgroups. Condition (*ii*) then ensures that  $\mathscr{A}_{k+1} = \emptyset$  for some integer  $k \geq 1$ ; we define k as the smallest such integer. Condition (*ii*) also ensures that each  $\mathscr{A}_i$  is conjugacyinvariant, and that  $\mathscr{A} = \mathscr{A}_1 \sqcup \cdots \sqcup \mathscr{A}_k$ .

**Lemma 4.21.** Under Conditions (i), (ii) and (\*), the collection  $\mathscr{A}$  is  $\mathcal{O}$ -invariant.

*Proof.* We have already observed that  $\mathscr{E}_{\max}$  is  $\mathcal{O}$ -invariant, that it is exactly the set of vertex-stabilisers of T, and that each of its elements fixes a unique vertex of T.

Define  $\mathscr{E}^1$  to be the set of intersections  $E \cap E'$  for distinct elements  $E, E' \in \mathscr{E}_{\max}$ . Elements of  $\mathscr{E}^1$  are precisely *G*-stabilisers of non-degenerate arcs of *T*; moreover,  $\mathscr{E}^1$  is again  $\mathcal{O}$ -invariant. The set  $\mathscr{A}_1$  coincides with the set of maximal elements of  $\mathscr{E}^1$ , so  $\mathscr{A}_1$  is  $\mathcal{O}$ -invariant.

Now, inductively for  $i \geq 1$ , let  $\mathscr{E}^{i+1}$  be the set of intersections  $E \cap E'$  for distinct elements  $E, E' \in \mathscr{E}_{\max}$  such that there does not exists a sequence  $A_1, \ldots, A_k \in \mathscr{A}_i$  with  $A_1 \leq E, A_k \leq E'$  and  $\langle A_j, A_{j+1} \rangle \in \mathscr{E}$  for all  $1 \leq j < k$ . In other words, if  $v, v' \in T$  are the unique vertices fixed by E and E' respectively, then the arc  $[v, v'] \subseteq T$  is non-degenerate and it is not covered by the fixed sets of the subgroups in  $\mathscr{A}_i$ , so this arc contains an edge with stabiliser in  $\mathscr{A} \setminus \mathscr{A}_i$ . Since  $\mathscr{A}_i$  is  $\mathcal{O}$ -invariant by the inductive hypothesis, the set  $\mathscr{E}^{i+1}$  is  $\mathcal{O}$ -invariant, because of its algebraic description. It follows that the set of maximal elements of  $\mathscr{E}^{i+1}$  is also invariant, and this set is precisely  $\mathscr{A}_{i+1}$ . In conclusion, this shows that  $\mathscr{A}_i$  is  $\mathcal{O}$ -invariant for every i, and hence  $\mathscr{A}$  is itself  $\mathcal{O}$ -invariant.

From the above data, we define some G-invariant families  $\mathcal{W}_i$  of subtrees of T for  $0 \leq i \leq k$ . Namely, we define  $\mathcal{W}_0 := \{\{v\} \mid v \text{ is a vertex of } T\}$  and, for each  $i \geq 1$ , we set

$$\mathcal{W}_i := \{ \operatorname{Fix}(H) \mid H \in \mathscr{A}_i \}.$$

We are about to use these families to construct a canonical sequence of forests  $\mathcal{G}_1, \ldots, \mathcal{G}_k$ . The construction will only use the following straightforward property of the families  $\mathcal{W}_0, \ldots, \mathcal{W}_k$ :

(★) For  $i \ge 1$ , the intersection of any two distinct elements of  $\mathcal{W}_i$  is covered by the subtrees in the union  $\mathcal{W}_0 \cup \cdots \cup \mathcal{W}_{i-1}$ . The elements of  $\mathcal{W}_0$  are pairwise disjoint.

Also note that the subtrees in the last family  $\mathcal{W}_k$  cover the tree T.

We will make use of additional G-invariant families  $\mathcal{U}_1, \ldots, \mathcal{U}_k$  of subtrees of T, which are obtained from the  $\mathcal{W}_i$  as follows: for each  $1 \leq i \leq k$ , the elements of  $\mathcal{U}_i$  are the connected components of the union within T of the subtrees in the family  $\mathcal{W}_0 \cup \cdots \cup \mathcal{W}_{i-1}$ . In particular,  $\mathcal{U}_1 = \mathcal{W}_0$ . The construction of the forests  $\mathcal{G}_1, \ldots, \mathcal{G}_k$  will be based of an iterated application of the following elementary construction, which is a slightly expanded version of Guirardel's tree dual to a transverse covering [Gui04, Gui08].

**Construction 4.22.** Let  $\mathcal{U}, \mathcal{W}$  be two families of subtrees of a tree  $\mathcal{T}$ . Suppose that the elements of  $\mathcal{U}$  are pairwise disjoint, and that the intersection of any two elements of  $\mathcal{W}$  is contained in an element of  $\mathcal{U}$ . We can form a bipartite graph  $\mathcal{G} = \mathcal{G}(\mathcal{U}, \mathcal{W})$  with vertex set  $\mathcal{U} \sqcup \mathcal{W}$  and edges [U, W] for  $U \in \mathcal{U}$  and  $W \in \mathcal{W}$  with  $U \cap W \neq \emptyset$ . It is straightforward to check that  $\mathcal{G}$  is a forest (see [Gui04, Definition 4.8] and the subsequent discussion, after collapsing the elements of  $\mathcal{U}$  to points).

We will speak of  $\mathcal{U}$ -type and  $\mathcal{W}$ -type vertices of  $\mathcal{G}$ , with the obvious meaning. Given a subgraph  $S \subseteq \mathcal{G}$ , we will also speak of its support supp $(S) \subseteq \mathcal{T}$ , which is the union of the subtrees of  $\mathcal{T}$  corresponding to the vertices of S. If  $S \subseteq \mathcal{G}$  is connected, then supp(S) is a subtree of  $\mathcal{T}$ .

Supports give a 1-to-1 correspondence between the connected components of the forest  $\mathcal{G}$  on the one side, and the connected components of the union within  $\mathcal{T}$  of the subtrees in  $\mathcal{U} \sqcup \mathcal{W}$  on the other. To check this, note that, if two elements of  $\mathcal{U} \sqcup \mathcal{W}$  intersect when viewed as subtrees of  $\mathcal{T}$ , then they are connected by a path of length  $\leq 2$  in  $\mathcal{G}$ , when viewed as vertices of  $\mathcal{G}$ .

Now, using the above families  $\mathcal{W}_0, \ldots, \mathcal{W}_k$  and  $\mathcal{U}_1, \ldots, \mathcal{U}_k$  of subtrees of T, we inductively define forests  $\mathcal{G}_1, \ldots, \mathcal{G}_k$  as follows. It is immediate from their definition that the subtrees in each family  $\mathcal{U}_i$ are pairwise disjoint. In addition,  $(\bigstar)$  implies that the intersection of any two distinct elements of  $\mathcal{W}_i$ is contained in an element of  $\mathcal{U}_i$ . Thus, the pair  $(\mathcal{U}_i, \mathcal{W}_i)$  satisfies the hypotheses of Construction 4.22 for each  $1 \leq i \leq k$ , and we can define:

$$\mathcal{G}_i := \mathcal{G}(\mathcal{U}_i, \mathcal{W}_i)$$

In particular, we have  $\mathcal{G}_1 = \mathcal{G}(\mathcal{W}_0, \mathcal{W}_1)$ . Again, we will speak of  $\mathcal{U}$ -type and  $\mathcal{W}$ -type vertices of  $\mathcal{G}_i$ , and associate to each subgraph of  $\mathcal{G}_i$  a support within T. Using the last paragraph of Construction 4.22, we have the equality

$$\mathcal{U}_i = {\text{supp}(C) \mid C \text{ is a connected component of } \mathcal{G}_{i-1} }.$$

It is clear that each forest  $\mathcal{G}_i$  is equipped with a natural G-action, as the families  $\mathcal{U}_i$  and  $\mathcal{W}_i$  are G-invariant, and their construction only depends on the intersection pattern of the  $\mathcal{W}_i$  within T. In fact, the construction is so canonical that the actions  $G \curvearrowright \mathcal{G}_i$  extend to actions of the group " $G \times \mathcal{O}$ ", as the next result shows. As in the proof of Lemma 4.13, we denote by  $G \times \mathcal{O}$  the preimage of  $\mathcal{O} \leq \operatorname{Out}(G)$  within  $\operatorname{Aut}(G)$ ; this contains the subgroup of inner automorphisms as a normal subgroup, which we identify with G (recalling that G has trivial centre).

**Lemma 4.23.** If  $G \sim T$  satisfies Conditions (i), (ii) and (\*), the following hold.

- (1) The last forest  $\mathcal{G}_k$  is a tree.
- (2) Each action  $G \curvearrowright \mathcal{G}_i$  uniquely extends to an action  $G \times \mathcal{O} \curvearrowright \mathcal{G}_i$  by graph automorphisms.

*Proof.* Since the elements of the last family  $\mathcal{W}_k$  cover T, by construction, each path in T is covered by finitely elements of  $\mathcal{U}_k \cup \mathcal{W}_k$ . This implies that  $\mathcal{G}_k$  is connected, proving part (1).

For part (2), we need to show that each representative  $\varphi \in \operatorname{Aut}(G)$  of an outer automorphism  $\phi \in \mathcal{O}$  is realised by a unique graph automorphism  $\Phi_i \colon \mathcal{G}_i \to \mathcal{G}_i$  satisfying  $\Phi_i(gx) = \varphi(g)\Phi_i(x)$  for all  $g \in G$  and  $x \in \mathcal{G}_i$ . Keeping  $\varphi$  fixed, we prove this by induction on *i*.

In the base step, the forest  $\mathcal{G}_1$  can be equivalently described as the bipartite graph having a black vertex for each subgroup in  $\mathscr{A}_1$  (corresponding to elements of  $\mathcal{W}_1$ ), a white vertex for each subgroup in  $\mathscr{E}_{\max}$  (corresponding to elements of  $\mathcal{U}_1 = \mathcal{W}_0$ ), and edges given by inclusions between these two types of subgroups of G. (This uses Condition (\*) and Remark 4.20.) This description of  $\mathcal{G}_1$  is purely algebraic and thus preserved by the automorphism  $\varphi$ , in view of the  $\varphi$ -invariance of  $\mathscr{E}$  and  $\mathscr{A}$  given by Condition (i) and Lemma 4.21. This shows the existence of the required map  $\Phi_1$ . As to uniqueness, note that  $\Phi_1$  is uniquely determined on the set of white vertices of  $\mathcal{G}_1$ , as their *G*-stabilisers are the elements of  $\mathscr{E}_{\max}$ . Since  $\mathcal{G}_1$  is bipartite with respect to the white-black colouring, and since each black vertex is uniquely determined by its link<sup>14</sup>, it follows that  $\Phi_1$  is uniquely determined on the whole forest  $\mathcal{G}_1$ .

For the inductive step, suppose that the maps  $\Phi_1, \ldots, \Phi_{i-1}$  have been defined for some  $i \ge 2$  and let us define the map  $\Phi_i$ .

Let  $\mathscr{C}$  be the family of subgroups of G that arise as stabilisers of the connected components of the forest  $\mathcal{G}_{i-1}$ . By the existence of the map  $\Phi_{i-1}$ , it follows that  $\mathscr{C}$  is  $\varphi$ -invariant. At the same time, Lemma 4.21 implies that the family  $\mathscr{A}_j$  is  $\varphi$ -invariant for each value of j.

Before continuing, we make two observations. First, distinct elements of  $\mathscr{A}_i$  have different fixed sets within T; this is because elements of  $\mathscr{A}_i$  are G-stabilisers of edges of T, so at least one of their fixed edges has precisely them as their G-stabiliser. Second, suppose that  $C \subseteq \mathcal{G}_{i-1}$  is a component, with G-stabiliser  $G_C$ , and consider a subgroup  $H \in \mathscr{E}_{\max}$ . We claim that the vertex  $v_H \in T$  fixed by H lies in  $\operatorname{supp}(C)$  if and only if  $H \leq G_C$ . The forward direction is clear, as the supports of the components of  $\mathcal{G}_{i-1}$  are pairwise disjoint within T, and each vertex of T lies in one. For the converse, suppose for the sake of contradiction that we have  $H \leq G_C$  and  $v_H \notin \operatorname{supp}(C)$ . Then, since  $H \leq G_C$ , the subgroup H preserves both  $\operatorname{supp}(C)$  and  $v_H$ , and hence it fixes any edge of Tseparating  $v_H$  from  $\operatorname{supp}(C)$ , contradicting the fact that each element of  $\mathscr{E}_{\max}$  has a unique fixed point in T. Note that, as a consequence of the claim, we also have that distinct components of the forest  $\mathcal{G}_{i-1}$  are stabilised by distinct elements of  $\mathscr{C}$ .

Now, the forest  $\mathcal{G}_i$  can be described as having one vertex for each subgroup in the disjoint union  $\mathscr{C} \sqcup \mathscr{A}_i$  and edges [X, Y] for subgroups  $X \in \mathscr{C}$  and  $Y \in \mathscr{A}_i$  such that there exists  $Z \in \mathscr{E}_{\max}$  with  $Y \leq Z \leq X$ ; in other words,  $\operatorname{Fix}(Y) \subseteq T$  intersects the support of the component of  $\mathcal{G}_{i-1}$  stabilised by X at the vertex of T with stabiliser Z. By the  $\varphi$ -invariance of all families of subgroups involved, it follows that there exists a map  $\Phi_i \colon \mathcal{G}_i \to \mathcal{G}_i$  with the required properties.

As to uniqueness, note that  $\Phi_i$  is again uniquely determined on the vertices of  $\mathcal{G}_i$  of  $\mathcal{U}$ -type, by the inductive hypothesis. Observing that each vertex of  $\mathcal{G}_i$  of  $\mathcal{W}$ -type is uniquely determined by its link in  $\mathcal{G}_i$ , it follows that  $\Phi_i$  is uniquely determined on  $\mathcal{G}_i$ , completing the proof of the lemma.  $\Box$ 

Part (2) of Theorem 4.18 quickly follows from the previous lemma, as we now explain.

Proof of Theorem 4.18(2). Recall that we are assuming that  $G \curvearrowright T$  satisfies Conditions (i), (ii), (\*), and now also that no edge-stabiliser of T is normal in G. Our goal is to construct a splitting  $G \times \mathcal{O} \curvearrowright T''$  such that all subgroups elliptic in T are elliptic in T'', and such that G-stabilisers of edges of T'' have the form described in the theorem statement.

By Lemma 4.23, we have an action  $G \times \mathcal{O} \curvearrowright \mathcal{G}_k$ , where  $\mathcal{G}_k$  is a tree. Note that  $\mathcal{G}_k$  has at least two  $\mathcal{W}$ -type vertices: if there were only one, this would correspond to a subgroup  $A \in \mathscr{A}_k$  that is normal in G, violating our assumptions.

If  $S \subseteq \mathcal{G}_k$  is a *G*-invariant subtree, then  $\operatorname{supp}(S) \subseteq T$  is also a *G*-invariant subtree, and hence  $\operatorname{supp}(S) = T$  by the assumption that  $G \curvearrowright T$  is minimal. This implies that *S* must contain all  $\mathcal{W}$ -type vertices of  $\mathcal{G}_k$ , as the support of a  $\mathcal{U}$ -type vertex of  $\mathcal{G}_k$  does not contain any edges of *T* with stabiliser in  $\mathscr{A}_k$ . This shows that the convex hull in  $\mathcal{G}_k$  of the set of  $\mathcal{W}$ -type vertices is the smallest *G*-invariant subtree of  $\mathcal{G}_k$ ; we define  $T'' \subseteq \mathcal{G}_k$  as this convex hull. As we have observed that  $\mathcal{G}_k$  has at least two  $\mathcal{W}$ -type vertices, the subtree T'' is not a single point. Thus,  $G \curvearrowright T''$  is a splitting.

Note that T'' is preserved by the  $G \times \mathcal{O}$ -action on  $\mathcal{G}_k$ . Indeed, if T'' is a proper subtree of  $\mathcal{G}_k$ , then each vertex of  $\mathcal{G}_k \setminus T''$  is a  $\mathcal{U}$ -type vertex of degree 1. At the same time, no vertex of T'' has degree 1, as  $G \curvearrowright T''$  is minimal. The action  $G \times \mathcal{O} \curvearrowright \mathcal{G}_k$  preserves vertex degrees, and so it must leave T'' invariant.

<sup>&</sup>lt;sup>14</sup>Here, note that black vertices always have nonempty link in  $\mathcal{G}_1$ , even though white vertces can have empty link.

If a subgroup of G is elliptic in T, then it is contained in an element of  $\mathscr{E}_{\max}$ , which in turn fixes (at least) a  $\mathcal{U}$ -type vertex of  $\mathcal{G}_k$  (recalling that vertices of T are the elements of the family  $\mathcal{U}_1 = \mathcal{W}_0$ ). Thus all subgroups elliptic in T are elliptic in T''.

We are left to describe G-stabilisers of edges of  $\mathcal{G}_k$ . Each such edge is of the form e = [U, W], where  $W = \operatorname{Fix}(A) \subseteq T$  for some edge-stabiliser  $A \in \mathscr{A}_k$ , and  $U \subseteq T$  is a connected component of the union of the edges in T with stabiliser not in  $\mathscr{A}_k$ , such that  $U \cap W \neq \emptyset$ . The G-stabiliser of Wis precisely the normaliser  $N_G(A)$ , as A is the unique smallest G-stabiliser of an edge of W. Thus, the G-stabiliser  $G_e$  of the edge e = [U, W] coincides with the  $N_G(A)$ -stabiliser of the subtree U, or equivalently that of the intersection  $U \cap W$ .

Consider for a moment the action  $N_G(A) \curvearrowright \operatorname{Fix}(A) \subseteq T$  and note that it factors through an action  $N_G(A)/A \curvearrowright \operatorname{Fix}(A)$ . With respect to the latter action, some edges have trivial stabiliser (the edges of T that have G-stabiliser equal to A), and the remaining edges are precisely those contained in an intersection  $U' \cap \operatorname{Fix}(A)$  for some  $U' \in \mathcal{U}_k$  (necessarily adjacent to  $W = \operatorname{Fix}(A)$  within  $\mathcal{G}_k$ ). Collapsing all edges of  $\operatorname{Fix}(A)$  with stabiliser different from A, we obtain a  $N_G(A)/A$ -action on a tree with trivial edge-stabilisers, and the subgroup  $G_e$  is the preimage in  $N_G(A)$  of a vertexstabiliser of this  $N_G(A)/A$ -action. This means that  $A \triangleleft G_e$  and that  $G_e/A$  is a free factor of the group  $N_G(A)/A$  (possibly the trivial subgroup or the entire  $N_G(A)/A$ ), concluding the proof.  $\Box$ 

Part (1) of Theorem 4.18 will require more work. Roughly, starting with the tree  $\mathcal{G}_k$ , the plan is to repeatedly blow up  $\mathcal{U}$ -type vertices, replacing them with the connected components of the forests  $\mathcal{G}_i$  that they represent. In order to be able to perform this blowing-up procedure equivariantly, we now introduce Condition (*iii*) into our standing assumptions.

As a first consequence of Condition (*iii*), the connected components of the forests  $\mathcal{G}_i$  are usually acted upon minimally by their G-stabilisers. This is explained in the next lemma. Recall that we say that a vertex of one of the forests  $\mathcal{G}_i$  is of  $\mathcal{U}$ -type if it corresponds to a connected component of the forest  $\mathcal{G}_{i-1}$  (or to a vertex of T when i = 1), and we say that a vertex of  $\mathcal{G}_i$  is of  $\mathcal{W}$ -type if it corresponds to the subtree  $\operatorname{Fix}(A) \subseteq T$  for some  $A \in \mathscr{A}_i$ .

**Lemma 4.24.** If  $G \curvearrowright T$  satisfies Conditions (i)–(iii) and (\*), the following holds. For each  $1 \leq i \leq k$  and each connected component  $C \subseteq \mathcal{G}_i$ , each vertex  $v \in C$  lies in the minimal subtree<sup>15</sup> of the G-stabiliser of C and this is non-elliptic, except when

- either v is of  $\mathcal{U}$ -type and  $C = \{v\}$  (i.e. v is an isolated vertex of  $\mathcal{G}_i$ );
- or v is of  $\mathcal{U}$ -type, it has degree 1 in C, and  $\operatorname{supp}(v) \subseteq T$  is a single vertex (in which case, either i = 1, or v corresponds to an isolated  $\mathcal{U}$ -type vertex in each of the forests  $\mathcal{G}_1, \ldots, \mathcal{G}_{i-1}$ ).

*Proof.* The main step in the proof of the lemma is the following claim.

**Claim.** Consider  $1 \leq i \leq k$ , a connected component  $C \subseteq \mathcal{G}_i$  and a  $\mathcal{W}$ -type vertex  $w \in C$ . Then w lies on the axis in C of an element of G that stabilises C and is loxodromic in both T and C.

Proof of claim. The  $\mathcal{W}$ -type vertex  $w \in C$  corresponds to the subtree  $\operatorname{Fix}(A) \subseteq T$  for some  $A \in \mathscr{A}_i$ . Let  $e \subseteq \operatorname{Fix}(A)$  be an edge whose G-stabiliser is exactly equal to A; this exists because  $A \in \mathscr{A}$ , and it shows in particular that the G-stabiliser of  $\operatorname{Fix}(A)$  is precisely the normaliser  $N_G(A)$ .

Let x and y be the vertices of the edge e. Observe that the G-stabilisers of  $x, y \in T$  cannot both be contained in  $N_G(A)$ . Indeed, by Condition (\*) there exist elements  $g_x \in G_x \setminus A$  and  $g_y \in G_y \setminus A$ , and the product  $g_x g_y$  is then loxodromic in T. Since  $N_G(A)$  is elliptic in T by Condition (*iii*), it follows that  $g_x$  and  $g_y$  do not both lie in  $N_G(A)$ .

Thus, without loss of generality, there exists an element  $g_x \in G_x \setminus N_G(A)$ . The sets  $\operatorname{Fix}(A)$  and  $g_x \operatorname{Fix}(A)$  are distinct in T and their intersection contains the vertex x, so there exists a connected component  $D \subseteq \mathcal{G}_{i-1}$  such that  $\operatorname{supp}(D) \supseteq \operatorname{Fix}(A) \cap g_x \operatorname{Fix}(A)$ . Note that  $g_x D = D$ , since  $g_x$  fixes

 $<sup>^{15}</sup>$ In particular, the minimal subtree exists, which is not automatic for actions of infinitely generated groups.

 $x \in \text{supp}(D)$ . Let  $u \in C \subseteq \mathcal{G}_i$  be the  $\mathcal{U}$ -type vertex corresponding to D. We have constructed a length-2 geodesic  $[w, u, g_x w]$  in  $C \subseteq \mathcal{G}_i$  with  $g_x u = u$ .

Now recall that, by Condition (\*), we have an element  $g_y \in G_y \setminus A$ , though we do not know whether this element lies in  $N_G(A)$  or not. The edges  $e, g_y e \subseteq T$  are not contained in the support of any components of  $\mathcal{G}_{i-1}$ , as their G-stabilisers lie in  $\mathscr{A}_i$ . As a consequence, e and  $g_y e$  separate  $\operatorname{supp}(D)$  from  $g_y \operatorname{supp}(D)$  in T. In particular, we have  $g_y u \neq u$  in  $\mathcal{G}_i$ . At this point, we need to distinguish two cases.

If  $g_y \in N_G(A)$ , then  $g_y w = w$  and  $[g_y u, w, u, g_x w]$  is a length-3 geodesic in  $C \subseteq \mathcal{G}_i$ . The open edge (w, u) separates the fixed sets of  $g_y$  and  $g_x$  in C, and hence it is contained in the axis of the loxodromic element  $g_x g_y$ . In particular, w lies on the axis of  $g_x g_y$  in C, and we have already observed above that  $g_x g_y$  is also loxodromic in T.

Suppose instead that  $g_y \notin N_G(A)$ . Then, in the tree T, the intersection  $\operatorname{Fix}(A) \cap g_y \operatorname{Fix}(A) \ni y$ is contained in  $\operatorname{supp}(D')$  for a connected component  $D' \subseteq \mathcal{G}_{i-1}$ . We have  $D' \neq D$ , since the edge e separates the supports of D and D' in T. Thus D' determines a vertex  $u' \neq u$  in  $C \subseteq \mathcal{G}_i$  and, since  $g_y$  fixes  $y \in \operatorname{supp}(D')$ , we have  $g_y u' = u'$ . We obtain a length-4 geodesic  $[g_y w, u', w, u, g_x w]$ in C. The length-2 open arc (u', w, u) separates the fixed sets of  $g_y$  and  $g_x$  in C, so  $g_x g_y$  is again loxodromic in C with axis containing w. As above,  $g_x g_y$  is also loxodromic in T.

Since w was an arbitrary  $\mathcal{W}$ -type vertex of C, this completes the proof of the claim.

Now, consider a connected component  $C \subseteq \mathcal{G}_i$  that is not a single  $\mathcal{U}$ -type vertex, and denote by  $G_C$  its G-stabiliser. By the claim,  $G_C$  is not elliptic in C, and the convex hull of the set of  $\mathcal{W}$ -type vertices in C is the smallest  $G_C$ -invariant subtree of C. In other words, the latter is the  $G_C$ -minimal subtree of C, and we will denote it by  $C_{\min}$ .

If the difference  $C \setminus C_{\min}$  is nonempty, it consists of a single layer of  $\mathcal{U}$ -type vertices hanging off  $C_{\min}$ . Let  $u \in C \setminus C_{\min}$  be one such vertex. The stabiliser  $G_u$  must fix the projection of uto  $C_{\min}$ , which is an adjacent vertex of  $\mathcal{W}$ -type  $w \in C_{\min}$ . The G-stabiliser of w is elliptic in Tby Condition (*iii*), so  $G_u$  is also elliptic in T. If  $i \geq 2$ , the vertex u corresponds to a connected component  $D \subseteq \mathcal{G}_{i-1}$ . If D were to contain a vertex of  $\mathcal{W}$ -type, then the above claim would give an element of  $G_u = G_D$  that is loxodromic in T, contradicting the fact that  $G_u$  is elliptic in T. Thus, the component  $D \subseteq \mathcal{G}_{i-1}$  must consist of a single vertex, corresponding to a connected component  $D' \subseteq \mathcal{G}_{i-2}$ . Repeating this argument up until we obtain a singleton component of  $\mathcal{G}_1$ , we ultimately conclude that  $\operatorname{supp}(D)$  is a single vertex of T, proving the lemma.

We also need the following simple observation.

**Remark 4.25.** If a subgroup  $H \leq G$  is elliptic in T and preserves a connected component  $C \subseteq \mathcal{G}_i$ , for some index i, then H fixes a vertex of C. This can be seen as follows.

Since H leaves C invariant, it preserves the subtree  $\operatorname{supp}(C) \subseteq T$ . The fact that H is elliptic in T thus implies that H fixes a vertex  $x \in \operatorname{supp}(C)$ . If  $x \in U$  for some  $U \in \mathcal{U}_i$  representing a vertex of C, then it is immediate that U is preserved by the G-stabiliser of x, and thus also by H. Recall that the intersection of any two subtrees in  $\mathcal{W}_i$  is contained in a subtree in  $\mathcal{U}_i$ . Thus, if x is not contained in the support of any  $\mathcal{U}$ -type vertices of C, then x must be contained in the support of a unique  $\mathcal{W}$ -type vertex of C. In this case, this vertex of C is again fixed by the G-stabiliser of x, and hence by H.

We can finally reap the fruits of the above construction and prove part (1) of Theorem 4.18.

Proof of Theorem 4.18(1). Recall that we have a splitting  $G \curvearrowright T$  satisfying Conditions (i)-(iii)and (\*), and we wish to construct a splitting  $G \rtimes \mathcal{O} \curvearrowright T'$  with G-stabilisers of edges in  $\mathscr{A}$  (in fact, even in  $\mathscr{A}_1$ ) and exactly  $\mathscr{E}$  as family of elliptic subgroups of G. By Lemma 4.23, we already have a  $G \rtimes \mathcal{O}$ -tree, namely  $\mathcal{G}_k$ , but its edge-stabilisers will not normally lie in  $\mathscr{A}$ . By contrast, we do have a  $G \times \mathcal{O}$ -forest with some edge-stabilisers in  $\mathscr{A}_1$ , namely  $\mathcal{G}_1$  (we show this below); its fault is rather that  $\mathcal{G}_1$  is not connected.

To overcome these issues, we will combine the forests  $\mathcal{G}_1, \ldots, \mathcal{G}_k$  to form a single  $G \times \mathcal{O}$ -tree with the same G-orbits of edges and edge-stabilisers as the forest  $\mathcal{G}_1 \sqcup \cdots \sqcup \mathcal{G}_k$ . Then, at the very end, we will collapse all superfluous edges.

We start with the tree  $T_k := \mathcal{G}_k$  and blow up each  $\mathcal{U}$ -type vertex  $u \in T_k = \mathcal{G}_k$ ; by definition, u corresponds to a connected component  $C_u \subseteq \mathcal{G}_{k-1}$  and we simply replace u with a copy of the tree  $C_u$ . The claim below guarantees that this procedure can be performed so as to obtain a tree  $T_{k-1}$  that is still equipped with a  $G \times \mathcal{O}$ -action. The edge set of  $T_{k-1}$  is naturally identified with the edge set of  $\mathcal{G}_{k-1} \sqcup \mathcal{G}_k$ , and the vertex set of  $T_{k-1}$  is naturally identified with the union of the set of  $\mathcal{W}$ -type vertices of  $\mathcal{G}_{k-1} \sqcup \mathcal{G}_k$  and the set of  $\mathcal{U}$ -type vertices of  $\mathcal{G}_{k-1}$ . In particular, these natural identifications are  $G \times \mathcal{O}$ -equivariant. We then proceed to blow up all  $\mathcal{U}$ -type vertices of  $T_{k-1}$ , forming a  $G \times \mathcal{O}$ -tree  $T_{k-2}$ , and repeat this procedure until we finally obtain a tree  $T_1$  whose  $\mathcal{U}$ -type vertices correspond to vertices of T.

Blowing up  $\mathcal{U}$ -type vertices is not an entirely canonical procedure (we need to decide where to attach incident edges). However, the following claim guarantees that the blow-up can be performed  $G \times \mathcal{O}$ -equivariantly (using only part (1) of Lemma 4.13).

**Claim.** Consider an index  $2 \leq i \leq k$  and a component  $C \subseteq \mathcal{G}_{i-1}$ . Let  $j \geq i$  and let  $e \subseteq \mathcal{G}_j$  be an edge whose  $G \times \mathcal{O}$ -stabiliser leaves C invariant. Then this stabiliser is elliptic in C.

Proof of claim. Let  $G_e$  denote the G-stabiliser of the edge e. We begin by observing that  $G_e$  is elliptic in C. Since  $G_e$  fixes the W-type vertex of e, it normalises an edge-stabiliser of T. Condition (*iii*) thus implies that  $G_e$  is elliptic in T. Since  $G_e$  leaves C invariant, by assumption, Remark 4.25 then implies that  $G_e$  fixes a vertex of C, as required.

We now proceed to show that the  $G \times \mathcal{O}$ -stabiliser of e is also elliptic in C. For this, recall that the edge  $e \subseteq \mathcal{G}_j$  has vertices u and w which correspond, respectively, to a component  $C' \subseteq \mathcal{G}_{j-1}$ and to a subtree  $\operatorname{Fix}_T(A) \subseteq T$  with  $A \in \mathscr{A}_j$  and  $\operatorname{supp}_T(C') \cap \operatorname{Fix}_T(A) \neq \emptyset$  (to avoid confusion later in the proof, we add subscripts specifying in which tree we compute fixed sets and supports). In particular, A must leave C' invariant and it follows that A is contained in the stabiliser  $G_e$ . The latter is elliptic in  $C \subseteq \mathcal{G}_{i-1}$ , as we have seen, so A is also elliptic in C.

We claim that the subset  $\operatorname{Fix}_{C}(A) \subseteq \mathcal{G}_{i-1}$  has finite diameter. Indeed, Condition (*iii*) implies that  $\operatorname{Fix}_{T}(A)$  has finite diameter; let us call it K. If  $\operatorname{Fix}_{T}(A')$  and  $\operatorname{Fix}_{T}(A'')$  represent two  $\mathcal{W}$ -type vertices of  $\operatorname{Fix}_{C}(A)$ , then these sets are A-invariant and must therefore intersect  $\operatorname{Fix}_{T}(A)$ . It follows that  $\operatorname{Fix}_{T}(A')$  and  $\operatorname{Fix}_{T}(A'')$  are at distance at most K in T, and thus the corresponding vertices of C are at distance at most 2K. In conclusion,  $\operatorname{Fix}_{C}(A)$  has diameter at most 2K + 2.

Now, recalling that  $A \leq G_e$ , the set  $\operatorname{Fix}_C(G_e)$  also has finite diameter and thus admits a unique barycentre. Since G is normal in  $G \times \mathcal{O}$ , the subgroup  $G_e$  is normal in the  $G \times \mathcal{O}$ -stabiliser of the edge e. Thus, the latter stabiliser leaves the set  $\operatorname{Fix}_C(G_e)$  invariant and it must fix its barycentre. In conclusion, the  $G \times \mathcal{O}$ -stabiliser of e is elliptic in C, as required.

Applying the inductive procedure described before the claim, we ultimately obtain a  $G \times \mathcal{O}$ -tree  $T_1$  whose vertex set is naturally in bijection with the disjoint union  $\mathcal{W}_0 \sqcup \cdots \sqcup \mathcal{W}_k$  and whose edge set is naturally in bijection with the edge set of the forest  $\mathcal{G}_1 \sqcup \cdots \sqcup \mathcal{G}_k$ . These natural bijections are  $G \times \mathcal{O}$ -equivariant and, in particular, they preserve stabilisers.

Lemma 4.24 implies that the minimal subtree  $T_1^{\min}$  of the action  $G \curvearrowright T_1$  exists and contains all vertices of  $T_1$  corresponding to elements of the union  $\bigsqcup_{i\geq 1} \mathcal{W}_i$ , though it is possible in general that some elements of  $\mathcal{W}_0 \cong T^{(0)}$  determine vertices of  $T_1 \setminus T_1^{\min}$  (with degree 1 in  $T_1$ ). Since G is normal in  $G \rtimes \mathcal{O}$ , the subtree  $T_1^{\min}$  is  $G \rtimes \mathcal{O}$ -invariant.

Note that the G-trees  $T_1^{\min}$  and T have the same elliptic subgroups. In one direction, if a subgroup of G fixes a vertex of T, then it fixes the vertex of  $T_1$  determined by the corresponding element of

 $\mathcal{W}_0$ , and it also fixes its projection to  $T_1^{\min}$ . Conversely, if a subgroup of G fixes a vertex of  $T_1$ , then it preserves an element of  $\bigcup_{i\geq 0} \mathcal{W}_i$ , hence it normalises an edge-stabiliser of T, and it is elliptic in T by Condition (*iii*).

Now, let us show that  $T_1^{\min}$  has an edge with G-stabiliser lying in  $\mathscr{A}_1$ . Recall that, by Lemma 4.24, every  $\mathcal{W}$ -type vertex of the forest  $\mathcal{G}_1$  is contained in the minimal subtree of a (non-elliptic) subgroup of G. Thus,  $T_1^{\min}$  contains a length-2 geodesic of the form [u, w, u'], where w corresponds to the subtree  $\operatorname{Fix}(A) \subseteq T$  for some  $A \in \mathscr{A}_1$ , while u and u' correspond to distinct vertices  $x, x' \in \operatorname{Fix}(A)$ such that there exist  $B, B' \in \mathscr{A}_1$  with  $\operatorname{Fix}(A) \cap \operatorname{Fix}(B) = \{x\}$  and  $\operatorname{Fix}(A) \cap \operatorname{Fix}(B') = \{x'\}$ ; indeed, by maximality of A, B, B' within  $\mathscr{A}$ , these intersections cannot contain any edges.

Condition (*iii*) implies that  $N_G(A)$  fixes a vertex  $z \in Fix(A)$ . By maximality of  $A \in \mathscr{A}$ , no vertex of  $Fix(A) \setminus \{z\}$  can be fixed by an element of  $N_G(A) \setminus A$ . In particular, up to swapping x and x', we can assume that  $G_x \cap N_G(A) = A$ . The *G*-stabiliser of the edge  $[u, w] \subseteq T_1^{\min}$  is precisely this intersection. Thus, [u, w] is the required edge of  $T_1^{\min}$  with stabiliser in  $\mathscr{A}_1$ .

In conclusion,  $G \curvearrowright T_1^{\min}$  is an  $\mathcal{O}$ -invariant splitting, it has  $\mathscr{E}$  as collection of elliptic subgroups, and it has an edge with stabiliser in  $\mathscr{A}_1$ . Recalling that the collection  $\mathscr{A}_1$  is  $\mathcal{O}$ -invariant, we can now collapse all G-orbits of edges in  $T_1^{\min}$  whose G-stabiliser does not lie in  $\mathscr{A}_1$ , thus obtaining the  $\mathcal{O}$ -invariant tree described in the statement of the theorem.  $\Box$ 

Through most of Section 4.2.3, we have assumed that the initial splitting  $G \curvearrowright T$  satisfies Condition (\*), but this was not an hypothesis of Theorem 4.18. It is worth remarking that one gets a little more out of the above construction, if Condition (\*) is given at the start.

**Remark 4.26.** If the splitting  $G \curvearrowright T$  in Theorem 4.18 additionally satisfies Condition (\*), then all edge-stabilisers of the splitting  $G \curvearrowright T'$  constructed in part (1) actually lie in  $\mathscr{A}_1$ , that is, they are *maximal* edge-stabilisers of the initial tree T. This is clear from the proof of Theorem 4.18(1).

We do not get this property in general, as the collapse needed to ensure (\*) (Lemma 4.19) might make all maximal edge-stabilisers disappear.

# 4.2.4. Consequences for special groups. We now use the previous discussion to prove Theorem 4.8.

Proof of Theorem 4.8. Consider the collection  $\mathscr{T}$  of minimal, irredundant,  $\mathcal{O}$ -invariant  $(\mathcal{Z}_s(G), \mathcal{H})$ trees of G. We have  $\mathscr{T} \neq \emptyset$ , as it contains the tree that is a single vertex. We order  $\mathscr{T}$  by refinements: we have  $T_1 \preceq T_2$  if  $T_1$  is a G-equivariant collapse of  $T_2$ . Since G is unconditionally accessible over centralisers (Theorem 3.1), the collection  $\mathscr{T}$  admits a maximal element  $G \curvearrowright T$ . The rest of the proof is devoted to checking that the vertex groups of T are  $(\mathcal{Z}_s(G), \mathcal{H})$ -rigid in G.

For this, let V be the G-stabiliser of a vertex  $v \in T$ . We begin with some preliminary observations on V. First, V is convex-cocompact in G, by Proposition 2.30. Next, let  $\mathcal{E}_v$  be the collection of G-stabilisers of edges of T incident to v. Observing that  $\mathcal{E}_v \subseteq \mathcal{Z}_s(G)|_V \subseteq \mathcal{S}(G)|_V \subseteq \mathcal{H}|_V$  (using Notation 4.10), we see that V is 1-ended relative to  $\mathcal{H}|_V$ : otherwise Lemma 4.13(1) would allow us to refine T into a splitting of G relative to  $\mathcal{H}$  with a trivial edge-stabiliser and, collapsing the other G-orbits of edges, we would obtain a free splitting of G relative to  $\mathcal{H}$ , violating 1-endedness of G. A similar argument shows that V is unconditionally  $(\mathcal{Z}_s(G)|^V, \mathcal{H}|_V)$ -accessible: an irredundant  $(\mathcal{Z}_s(G)|^V, \mathcal{H}|_V)$ -splitting of V can be used to refine T into an irredundant  $(\mathcal{Z}_s(G), \mathcal{H})$ -splitting of G (not  $\mathcal{O}$ -invariant in general), and the size of the latter is bounded by accessibility of G over centralisers (Theorem 3.1).

Now, suppose for the sake of contradiction that V is non-elliptic in some  $(\mathcal{Z}_s(G), \mathcal{H})$ -splitting  $G \cap U$ , and let  $M \subseteq U$  be the V-minimal subtree.

We claim that the action  $V \cap M$  is a  $(\mathcal{Z}_s(G)|^V, \mathcal{H}|_V)$ -splitting. Indeed, the V-stabiliser of an edge of M is an intersection  $V \cap Z$  with  $Z \in \mathcal{Z}_s(G)$ . The centraliser Z is contained in an element of  $\mathcal{S}(G) \subseteq \mathcal{H}$ , so it is elliptic in T. If Z fixes v, then  $V \cap Z = Z$ . If it does not, we have  $Z \cap V = Z \cap G_e$ ,

where  $e \subseteq T$  is an edge incident to v and  $G_e \in \mathcal{Z}_s(G)$ . Either way, we have  $V \cap Z \in \mathcal{Z}_s(G)|^V$ . We stress, however, that  $\mathcal{Z}_s(G)|^V \not\subseteq \mathcal{Z}(V)$  in general.

Let  $\mathcal{O}|_V \leq \operatorname{Out}(V)$  be the restriction of  $\mathcal{O}$ , as defined in Remark 4.12. We now claim that the action  $V \curvearrowright M$  can be promoted to an (irredundant)  $\mathcal{O}|_V$ -invariant  $(\mathcal{Z}_s(G)|^V, \mathcal{H}|_V)$ -splitting of V. Once this is proved, it will complete the proof of the theorem: we can then use this splitting of V and Lemma 4.13(2) to refine T into a larger (irredundant)  $\mathcal{O}$ -invariant  $(\mathcal{Z}_s(G), \mathcal{H})$ -splitting of G, contradicting the definition of T. The assumptions of Lemma 4.13(2) are indeed satisfied: first, since  $G \notin \mathcal{S}(G)$  and  $V \notin \mathcal{S}(V) \subseteq \mathcal{H}|_V$ , both G and V have trivial centre and neither has a line-splitting over centralisers, as the kernel of the latter would be a nontrivial, normal, convex-cocompact subgroup (recall Lemma 2.4(2)). Moreover,  $(\mathcal{Z}_s(\cdot), \mathcal{S}(\cdot))$ -splittings are acylindrical, as shown in Remark 4.9.

Thus, we are left to construct an  $\mathcal{O}|_{V}$ -invariant  $(\mathcal{Z}_{s}(G)|^{V}, \mathcal{H}|_{V})$ -splitting of V. Towards this, recall that the collection  $\mathcal{Z}_{s}(G)$  is  $\mathcal{O}$ -invariant, closed under intersections, and each element of  $\mathcal{Z}_{s}(G)$ is contained in an element of  $\mathcal{S}(G) \subseteq \mathcal{H}$  (Lemma 4.6). Thus, the collection  $\mathcal{Z}_{s}(G)|^{V}$  is  $\mathcal{O}|_{V}$ -invariant and closed under intersections, the collection  $\mathcal{H}|_{V}$  is  $\mathcal{O}|_{V}$ -invariant, and we have  $\mathcal{Z}_{s}(G)|^{V} \subseteq \mathcal{H}|_{V}$ . Moreover, as observed above, V is unconditionally  $(\mathcal{Z}_{s}(G)|^{V}, \mathcal{H}|_{V})$ -accessible. All this means that we can apply Corollary 4.17 to  $V \curvearrowright M$ , with  $\mathcal{F} = \mathscr{F}_{int} = \mathcal{Z}_{s}(G)|^{V}$  and  $\mathcal{K} = \mathcal{H}|_{V}$ . As a result, we obtain a new  $(\mathcal{Z}_{s}(G)|^{V}, \mathcal{H}|_{V})$ -splitting  $V \curvearrowright M'$  with the additional property that its deformation space is  $\mathcal{O}|_{V}$ -invariant.

We then wish to apply Theorem 4.18(1) to  $V \curvearrowright M'$ , for which we need to check the conditions of Section 4.2.3. Condition (i) is satisfied by construction, and V also has trivial centre because V is not contained in any singular subgroup of G (as V splits relative to  $\mathcal{H}|_V$ ). No element of  $\mathcal{Z}_s(G)|^V$  is properly contained in a conjugate, by Lemma 2.4(3), so Condition (ii) is also satisfied. Finally, regarding Condition (iii), note that each edge-stabiliser of M' is an element  $Z \in \mathcal{Z}_s(G)|^V$ with  $Z \neq \{1\}$ , because V is 1-ended relative to  $\mathcal{H}|_V$ , as observed above. Hence the normaliser  $N_G(Z)$  is contained in an element of  $\mathcal{S}(G)$  by Lemma 4.6(3), showing that  $N_V(Z)$  is elliptic in M'. Since  $N_V(Z)$  acts cocompactly on  $\operatorname{Fix}(Z) \subseteq M'$  by Lemma 2.32, it follows that  $\operatorname{Fix}(Z)$  has finite diameter, proving Condition (iii). We can thus indeed appeal to Theorem 4.18(1), which yields the required  $\mathcal{O}|_V$ -invariant  $(\mathcal{Z}_s(G)|^V, \mathcal{H}|_V)$ -splitting  $V \curvearrowright M''$ .

Up to removing any degree-2 vertices, M'' is irredundant and this leads to the required contradiction, as explained earlier in the proof.

4.3. The enhanced JSJ splitting. Let G be special. In this subsection we prove Theorem 4.1, constructing the enhanced JSJ splitting of G. This will be achieved by combining the canonical splitting over centralisers that we constructed in Theorem 4.8 with the classical JSJ decomposition over cyclic subgroups [RS97] and its canonical tree of cylinders [GL11, GL17].

In fact, we will prove a strengthening of Theorem 4.1 that will turn out to be more useful later in the paper (see e.g. Remarks 4.32 and 4.33). In order to state this, we need the following notion.

**Definition 4.27** (Optimal QH subgroup). Let  $G \curvearrowright T$  be a splitting relative to a family  $\mathcal{H}$ . Let  $q \in T$  be a vertex whose stabiliser Q is quadratically hanging relative to  $\mathcal{H}$ , and let  $\Sigma$  be the associated surface with  $Q = \pi_1(\Sigma)$ . We say that Q is *optimal* if all the following hold:

- (1) the surface  $\Sigma$  is not a pair of pants;
- (2) for each edge  $e \subseteq T$  incident to q, the stabiliser  $G_e$  is either trivial or Q-conjugate to the entire fundamental group of a component of  $\partial \Sigma$  (rather than to a subgroup thereof);
- (3) for each component  $B \subseteq \partial \Sigma$ , there is at most one<sup>16</sup> Q-orbit of edges  $e \subseteq T$  incident to q such that  $G_e$  is Q-conjugate to  $\pi_1(B)$ .

When the above are satisfied, we also say that q is an *optimal QH vertex* of the tree T.

<sup>&</sup>lt;sup>16</sup>Exactly one, if G is 1–ended relative to  $\mathcal{H}$ ; see [GL17, Lemma 5.16].

We can now state the main result of this subsection; it is clear that this implies Theorem 4.1.

**Theorem 4.28.** Let  $\mathcal{O} \leq \text{Out}(G)$  be a subgroup, and let G be 1-ended relative to an  $\mathcal{O}$ -invariant collection  $\mathcal{H}$  that contains  $\mathcal{S}(G)$ . Then there exists an  $\mathcal{O}$ -invariant  $(\mathcal{ZZ}(G), \mathcal{H})$ -tree  $G \curvearrowright T$  with the following properties.

- (1) The G-stabiliser of each vertex of T is of one of two kinds:
  - (a) an optimal quadratically hanging subgroup relative to  $\mathcal{H}$ ;
  - (b) a convex-cocompact root-closed subgroup of G that is  $(\mathcal{Z}(G), \mathcal{H})$ -rigid in G.
- (2) Each edge  $e \subseteq T$  with G-stabiliser not in  $\mathcal{Z}(G)$  is incident to a type (a) vertex.
- (3) If the family  $\mathcal{H}$  contains all cyclic subgroups of G whose conjugacy class has finite  $\mathcal{O}$ -orbit, then each type (b) vertex group V is  $(\mathcal{Z}(V), \mathcal{H}|_V)$ -rigid in itself.

We now start working towards the proof of Theorem 4.28. Since the cyclic edge groups of T are not convex-cocompact in general, we will need the following observation (Lemma 4.29) to guarantee that non–QH vertex groups are indeed convex-cocompact. This uses Items (1) and (2) in Definition 4.27.

If  $\Sigma$  is a compact surface, a simple closed curve  $\gamma \subseteq \Sigma$  is *essential* if it is neither nulhomotopic nor homotopic into the boundary of  $\Sigma$ . An *essential multicurve* on  $\Sigma$  is a finite union of pairwise disjoint, pairwise non-homotopic, essential simple closed curves on  $\Sigma$ .

**Lemma 4.29.** Let  $G \curvearrowright T$  be any splitting. Let  $q \in T$  be an optimal, quadratically hanging vertex. Also suppose that every edge of T has a G-translate that is incident to q. Then, for any vertex  $w \in T \setminus (G \cdot q)$ , the stabiliser  $G_w$  is convex-cocompact and root-closed.

*Proof.* Let  $\Sigma$  be the compact surface with  $Q = \pi_1(\Sigma)$ . We can assume that  $\Sigma$  is orientable. Otherwise, let  $\Sigma_0$  be its orientable double cover and note that  $Q_0 := \pi_1(\Sigma_0)$  contains the entire fundamental group of each component of  $\partial \Sigma$ . Thus, G has an index-2 subgroup  $G_0$  that contains the G-stabiliser of each vertex of  $T \setminus (G \cdot q)$ , and has  $G_0 \cap Q = Q_0$ . We can then replace G with  $G_0$ .

Now,  $\Sigma$  is the genus-g surface with  $b \geq 1$  boundary components  $S_{g,b}$ . If g = 0, we have  $b \geq 4$ ; indeed, the pair of pants is ruled out by Item (1) in Definition 4.27, and it also does not arise as the orientable double cover of any surface (the rank-2 free group cannot be a proper finite-index subgroup of another free group, by Nielsen-Schreier).

Let  $B \subseteq \partial \Sigma$  be a component. We claim that there are two essential multicurves  $\gamma_1, \gamma_2 \subseteq \Sigma$ such that, if  $T_1, T_2$  are the corresponding cyclic splittings of  $\pi_1(\Sigma)$ , then  $\pi_1(B)$  is precisely the  $\pi_1(\Sigma)$ -stabiliser of some pair of vertices  $(v_1, v_2) \in T_1 \times T_2$ . If g = 0 and  $b \ge 4$ , we can take  $\gamma_1, \gamma_2$ to be single curves, each bounding a pant with boundary components  $B, \gamma_i, B_i$ , for components  $B_i \subseteq \partial \Sigma$  such that  $B, B_1, B_2$  are pairwise distinct. If  $g \ge 1$  and  $b \ge 1$ , then we can represent  $\Sigma$  as a gluing  $S_{g-1,b+1} \cup P$ , where P is a pair of pants with a component equal to B and the remaining two components glued to two boundary components of  $S_{g-1,b+1}$ . Call  $\gamma_1$  the multicurve in  $\Sigma$  formed by the two boundary components of P other than B (to be precise,  $\gamma_1$  is just one curve for (g, b) = (1, 1)). We then define  $\gamma_2 = \phi \cdot \gamma_1$  for an element  $\phi \in Mod(\Sigma)$  that does not preserve  $\gamma_1$ .

Now, the two splittings  $Q \curvearrowright T_i$  can be used to refine  $G \curvearrowright T$  into two splittings  $G \curvearrowright T'_{B,i}$ (Lemma 4.13(1)). We then collapse all edges of  $T'_{B,i}$  coming from edges of T, obtaining two splittings  $G \curvearrowright T''_{B,i}$  whose edge-stabilisers are conjugate to maximal cyclic subgroups represented by essential simple closed curves on  $\Sigma$ . In particular, these cyclic edge-stabilisers coincide with their centralisers in G, and hence they are root-closed and convex-cocompact. Thus, the G-stabiliser of any vertex of  $T''_{B,i}$  is convex-cocompact by Proposition 2.30, and root-closed by Remark 2.31.

Finally, *G*-stabilisers of vertices of  $T \setminus (G \cdot q)$  are intersections of vertex-stabilisers of the trees  $T''_{B,i}$  as *B* varies among the components of  $\partial \Sigma$  and *i* varies in  $\{1, 2\}$  (this uses Item (2) in Definition 4.27). In conclusion, such stabilisers are root-closed and convex-cocompact, by Lemma 2.4(1).

We also record here the following observation for later use. The proof is straightforward, using the same splittings as in the proof of Lemma 4.29 and recalling that  $\mathcal{Z}_c(G)$  is the collection of contracting maximal cyclic subgroups of G. This is where Item (3) of Definition 4.27 is used.

**Lemma 4.30.** Let  $G \curvearrowright T$  be a splitting relative to a collection  $\mathcal{H}$ . Let  $q \in T$  be an optimal, quadratically hanging vertex relative to  $\mathcal{H}$ . If a subgroup  $H \leq G$  is  $(\mathcal{Z}_c(G), \mathcal{H})$ -rigid in G, then:

- (1) if  $h \in H$  is loxodromic in T, the axis of h does not contain q;
- (2) if H is contained in Q, it is peripheral.

In order to canonically split G over cyclic subgroups, we will use the following result, which can be quickly deduced from [GL17]. The only subtlety is that, since we want non–QH vertex groups to be convex-cocompact, we need to be a little careful over which cyclic subgroups we split. From now on, all QH subgroups will be meant relative to  $\mathcal{H}$  without explicit mention.

**Proposition 4.31.** Let  $\mathcal{O} \leq \text{Out}(G)$  and let G be 1-ended relative to an  $\mathcal{O}$ -invariant collection  $\mathcal{H} \supseteq \mathcal{S}(G)$ . There is an  $\mathcal{O}$ -invariant  $(\text{Cyc}(G), \mathcal{H})$ -tree  $G \curvearrowright T$  such that the following hold.

- (1) Each vertex-stabiliser is of one of two kinds:
  - (a) an optimal quadratically hanging subgroup;
  - (b) a convex-cocompact, root-closed subgroup of G that is  $(\mathcal{Z}_c(G), \mathcal{H})$ -rigid in G.
- (2) For each edge  $e \subseteq T$ , either  $G_e \in \mathcal{Z}_c(G)$  or e is incident to a type (a) vertex.

Proof. We will use the terminology of [GL17]. Normalisers of elements of  $\operatorname{Cyc}(G)$  lie in  $\mathcal{Z}(G)$ , hence they are either cyclic or contained in elements of  $\mathcal{S}(G) \subseteq \mathcal{H}$ ; in particular, they are small in  $(\operatorname{Cyc}(G), \mathcal{H})$ -trees. We can therefore appeal to [GL17, Corollary 9.1], using commensurability as the equivalence relation ~ on  $\operatorname{Cyc}(G)$ . The result is an  $\mathcal{O}$ -invariant  $(\operatorname{Cyc}(G), \mathcal{H})$ -tree  $G \curvearrowright T$ (denoted  $(T_a)_c^*$  in the reference) with the following three properties:

- vertex-stabilisers are either quadratically hanging, or  $(Cyc(G), \mathcal{H})$ -rigid in G;
- T is a JSJ tree for  $(Cyc(G), \mathcal{H})$ : its edge-stabilisers are elliptic in all  $(Cyc(G), \mathcal{H})$ -trees, and its elliptic subgroups are elliptic in all  $(Cyc(G), \mathcal{H})$ -trees with such edge-stabilisers;
- T is compatible with all  $(\operatorname{Cyc}(G), \mathcal{H})$ -trees: if  $G \curvearrowright U$  is a one-edge  $(\operatorname{Cyc}(G), \mathcal{H})$ -splitting, then U is a collapse of T, or a collapse of the refinement of T given by splitting a single QH vertex group over an essential simple closed curve on the associated surface.

(According to [GL17, Corollary 9.1], we would have to worry about a third type of vertex group: subgroups of elements of  $\mathcal{Z}(G)$ . However, recalling Lemma 4.6 and the fact that G is 1-ended relative to  $\mathcal{H}$ , we see that such vertex groups are either contained in elements of  $\mathcal{S}(G) \subseteq \mathcal{H}$ , or commensurable to edge groups of T. Hence these vertex groups are  $(\operatorname{Cyc}(G), \mathcal{H})$ -rigid too.)

Without loss of generality, we can assume that all QH vertex groups of T are optimal. Indeed, QH vertex groups whose associated surface is a pair of pants are  $(\operatorname{Cyc}(G), \mathcal{H})$ -rigid in T (see [GL17, Proposition 5.4]), and we regard them as such. For any other QH vertex  $q \in T$ , with stabiliser Qand associated surface  $\Sigma$ , we can modify the tree T near q as follows. Let  $Q \curvearrowright S_{\Sigma}$  be the Q-action on the diameter-2 tree that has a central vertex fixed by Q, and one Q-orbit of edges for each component of  $B \subseteq \partial \Sigma$ , with Q-stabilisers that are the conjugates of  $\pi_1(B)$  within  $Q = \pi_1(\Sigma)$ . We can replace  $q \in T$  with a copy of  $S_{\Sigma}$ : attach each edge  $e \subseteq T$  incident to q to the unique vertex of  $S_{\Sigma}$  whose Q-stabiliser contains  $G_e$  as a subgroup of finite index (we have  $G_e \neq \{1\}$  because G is 1-ended relative to  $\mathcal{H}$ ). Performing this procedure on all QH vertices of T, we obtain a refinement of T in the same deformation space; we create some new edges and vertices, but their G-stabilisers are commensurable to G-stabilisers of previously existing edges, which are  $(\operatorname{Cyc}(G), \mathcal{H})$ -rigid. Thus, the above three items still hold for the modified tree, and all QH vertices have become optimal. We are left to check that the modified tree is still  $\mathcal{O}$ -invariant. This follows from the fact that  $\mathcal{O}$ preserves the collection of fundamental groups of boundary components of the surfaces associated to the QH vertices q of the original tree T. In turn, this holds because, for each such boundary component B, there exists at least one edge incident to q with stabiliser commensurable to B; indeed, this is due to the fact that G is 1–ended relative to  $\mathcal{H}$ , see [GL17, Lemma 5.16].

Thus, we assume that all QH vertices of T are optimal. Let  $G \curvearrowright T'$  be obtained from T by collapsing certain edges: we retain an edge  $e \subseteq T$  only if e is incident to an (optimal) QH vertex, or if its stabiliser  $G_e$  lies in  $\mathcal{Z}_c(G)$ . Note that  $G \curvearrowright T'$  is still an  $\mathcal{O}$ -invariant ( $\operatorname{Cyc}(G), \mathcal{H}$ )-tree. We claim that T' is the tree that we are looking for. Part (2) is clear, as is the fact that all QH vertex groups of T' are optimal. We only need to prove that the remaining vertex groups are of type (b).

If  $v \in T'$  is a non–QH vertex, then all its incident edges  $e \subseteq T'$  either have stabiliser  $G_e \in \mathcal{Z}_c(G)$ , or their other vertex is QH and optimal. Since all elements of  $\mathcal{Z}_c(G)$  are convex-cocompact and root-closed, the combination of Proposition 2.30, Remark 2.31 and Lemma 4.29 shows that the stabiliser  $G_v$  is convex-cocompact and root-closed.

We are left to show that  $G_v$  is elliptic in all  $(\mathcal{Z}_c(G), \mathcal{H})$ -splittings  $G \curvearrowright U$ . It suffices to show this when there is a single *G*-orbit of edges in *U*. In this case, the fact that *U* is compatible with the (modified) JSJ tree *T* leaves only two options:

- either there is an edge  $e \subseteq T$  with  $G_e \in \mathcal{Z}_c(G)$  such that U is obtained from T by collapsing all edges outside of the orbit  $G \cdot e$ ;
- or there are a QH vertex  $q \in T$  and an essential curve  $\gamma$  on the associated surface such that U is obtained by first refining T, splitting  $G_q$  over  $\langle \gamma \rangle$ , and then collapsing all original edges of T.

Either way, the construction of T' makes it clear that  $G_v$  is elliptic in such a tree U. Indeed, denoting by  $\pi: T \to T'$  the collapse map,  $G_v$  stabilises the fibre  $\pi^{-1}(v) \subseteq T$ , which does not contain any QH vertices of T, nor does it contain any edges with G-stabiliser in  $\mathcal{Z}_c(G)$ .

This completes the proof of the proposition.

The fact that we restricted to optimal QH vertex groups guarantees that we have the following.

**Remark 4.32.** Let  $G \curvearrowright T$  be the splitting provided by Proposition 4.31. Considering a type (b) vertex  $v \in T$  and its stabiliser  $V \leq G$ , we make the following additional observations.

- (1) We have  $\mathcal{Z}_c(V) \subseteq \mathcal{Z}_c(G)$ . Indeed, consider an element  $g \in V$  such that  $Z_V(g) = \langle g \rangle$  and suppose for the sake of contradiction that  $Z_V(g) \leq Z_G(g)$ . Since V is root-closed in G, it follows that  $Z_G(g) \in \mathcal{Z}_s(G)$ ; thus,  $Z_G(g)$  is contained in an element of  $\mathcal{S}(G)$  and it must fix a vertex  $x \in T$ . Since  $Z_V(g) \neq Z_G(g)$ , we have  $x \neq v$ ; let  $e \subseteq T$  be the edge incident to v in the direction of x. Thus, g fixes e and, since  $\langle g \rangle$  is maximal cyclic in V, we have  $G_e = \langle g \rangle$ . Since  $Z_V(g)$  is contained in  $Z_G(g) \in \mathcal{Z}_s(G)$ , we have  $G_e \notin \mathcal{Z}_c(G)$ . In particular, the vertex  $w \in e \setminus \{v\}$  must be an optimal QH vertex. Since  $G_w$  is hyperbolic, we have  $x \neq w$ , and hence there exists an edge  $f \subseteq T$  incident to w in the direction of x. Now, we have  $g \in G_e \cap G_f$ , but  $G_e \cap G_f = \{1\}$  by Definition 4.27(3), yielding the required contradiction.
- (2) If the collection  $\mathcal{H}$  contains all cyclic subgroups of G with finite  $\mathcal{O}$ -orbit, then the group V is  $(\mathcal{Z}_c(V), \mathcal{H}|_V)$ -rigid in itself. Indeed, the assumption on  $\mathcal{H}$  implies that it contains the G-stabiliser of every edge of T incident to v. Thus, a  $(\mathcal{Z}_c(V), \mathcal{H}|_V)$ -splitting of V could be used to refine T and, collapsing the original edges of T, we would obtain a splitting of G over an element of  $\mathcal{Z}_c(V)$  relative to  $\mathcal{H}$ , in which V is not elliptic. Since  $\mathcal{Z}_c(V) \subseteq \mathcal{Z}_c(G)$  by the previous observation, this would violate the fact that V is  $(\mathcal{Z}_c(G), \mathcal{H})$ -rigid in G.
- (3) The splitting  $G \curvearrowright T$  is acylindrical. Indeed, any two distinct edges incident to an optimal QH vertex have G-stabilisers with trivial intersection. Moreover, the fixed set in T of any  $Z \in \mathcal{Z}_c(V)$  has diameter at most the number of edges of the quotient T/G: otherwise, there would exist two distinct edges of T with stabiliser equal to Z in the same G-orbit, which would violate the fact that  $N_G(Z) = Z$ .

We are finally ready to prove Theorem 4.28.

Proof of Theorem 4.28. By Theorem 4.8, there is an  $\mathcal{O}$ -invariant  $(\mathcal{Z}_s(G), \mathcal{H})$ -tree  $G \curvearrowright T'$  whose vertex groups are  $(\mathcal{Z}_s(G), \mathcal{H})$ -rigid in G. By Proposition 2.30 and Remark 2.31, its vertex groups are convex-cocompact and root-closed.

Let V be the G-stabiliser of a vertex of T', and consider the restriction  $\mathcal{O}|_V \leq \operatorname{Out}(V)$  of  $\mathcal{O}$  as discussed in Remark 4.12. As in the proof of Theorem 4.8, the fact that G is 1-ended relative to  $\mathcal{H}$  implies that V is 1-ended relative to  $\mathcal{H}|_V$ . Using Notation 4.10, we also have  $\mathcal{S}(V) \subseteq \mathcal{H}|_V$  and  $Z_s(G)|_V \subseteq \mathcal{H}|_V$ , as a consequence of Lemma 4.6 and the fact that  $\mathcal{S}(G) \subseteq \mathcal{H}$ .

We can thus apply Proposition 4.31 to V. We obtain a  $\mathcal{O}|_V$ -invariant  $(\operatorname{Cyc}(V), \mathcal{H}|_V)$ -tree  $V \curvearrowright T_V$ whose QH vertex groups are optimal, and whose non-QH vertex groups are convex-cocompact, root-closed and  $(\mathcal{Z}_c(V), \mathcal{H}|_V)$ -rigid in V. Moreover, each edge either has an element of  $\mathcal{Z}_c(V)$  as its stabiliser, or it is incident to a QH vertex. Since  $\mathcal{Z}_s(G)|_V \subseteq \mathcal{H}|_V$ , Lemma 4.13(2) allows us to use  $T_V$  to refine T' into an  $\mathcal{O}$ -invariant  $(\mathcal{Z}\mathcal{Z}(G), \mathcal{H})$ -tree of G. To check that the hypotheses of the lemma are satisfied: since  $G \notin \mathcal{S}(G)$  and  $V \notin \mathcal{S}(V) \subseteq \mathcal{H}|_V$ , both G and V have trivial centre, and neither T' nor  $T_V$  is a line. Moreover,  $T_V$  is acylindrical by Remark 4.32(c).

We perform this refinement for all  $\mathcal{O}$ -orbits of vertex groups of T', and call  $G \curvearrowright T$  the result. We claim that T is the G-tree that we are looking for. For this, we are only left to show that the non-QH vertex groups of T satisfy the required rigidity properties. In turn, this simply amounts to showing that each non-QH vertex-stabiliser W of one of the actions  $V \curvearrowright T_V$  is  $(\mathcal{Z}(G), \mathcal{H})$ -rigid in G, and additionally  $(\mathcal{Z}(W), \mathcal{H}|_W)$ -rigid in itself if  $\mathcal{H}$  contains all cyclic subgroups of G whose G-conjugacy class has finite  $\mathcal{O}$ -orbit.

Thus, let W be a vertex-stabiliser of  $V \curvearrowright T_V$ . Clearly, W is  $(\mathcal{Z}_s(G), \mathcal{H})$ -rigid in G, since W is contained in V, which has this property. In addition, W is  $(\mathcal{Z}_c(G), \mathcal{H})$ -rigid in G, because it is  $(\mathcal{Z}_c(V), \mathcal{H}|_V)$ -rigid in V by construction. This shows that W is  $(\mathcal{Z}(G), \mathcal{H})$ -rigid in G.

Finally, suppose that  $\mathcal{H}$  contains all cyclic subgroups of G with finite  $\mathcal{O}$ -orbit. If  $w \in T$  is a vertex of which W is the G-stabiliser, it follows that the collection  $\mathcal{H}|_W$  contains all G-stabilisers of edges of T incident to W: indeed, each of these subgroups has finite  $\mathcal{O}$ -orbit, by the  $\mathcal{O}$ -invariance of T, and all non-cyclic edge-stabilisers lie in  $\mathcal{Z}_s(G)|_W \subseteq \mathcal{S}(G)|_W \subseteq \mathcal{H}|_W$ . Thus, if W were to admit a  $(\mathcal{Z}_s(W), \mathcal{H}|_W)$ -splitting, this could be used to refine  $G \curvearrowright T$  into a  $(\mathcal{Z}(G), \mathcal{H})$ -splitting of G in which W is not elliptic, a contradiction (this uses that  $\mathcal{Z}_s(W) \subseteq \mathcal{Z}(G)$ , see Lemma 4.7). This shows that W is  $(\mathcal{Z}_s(W), \mathcal{H}|_W)$ -rigid in itself, while it is  $(\mathcal{Z}_c(W), \mathcal{H}|_W)$ -rigid by Remark 4.32(2). In conclusion, W is  $(\mathcal{Z}(W), \mathcal{H}|_W)$ -rigid in itself, concluding the proof of the theorem.

Another advantage of having restricted to optimal QH vertex groups is the following observation.

**Remark 4.33.** Let  $G \curvearrowright T$  be the enhanced JSJ splitting provided by Theorem 4.28, relative to a collection  $\mathcal{H}$ . If a subgroup  $H \leq G$  is  $(\mathcal{Z}(G), \mathcal{H})$ -rigid in G, we claim that H is elliptic in T.

Indeed, if H were not elliptic, it would contain an element  $h \in H$  that is loxodromic in T (this holds even when H is not finitely generated, by Lemma 2.29). Let  $\alpha \subseteq T$  be the axis of h. On the one hand, no edge of  $\alpha$  can have G-stabiliser in  $\mathcal{Z}(G)$  because H is  $(\mathcal{Z}(G), \mathcal{H})$ -rigid. On the other, no vertex of  $\alpha$  can be optimal and QH, by Lemma 4.30(1). This is a contradiction.

In conclusion, using Lemma 4.30(2), either H fixes a type (b) vertex of T, or it fixes an edge both of whose vertices are of type (a).

#### 5. First consequences of the enhanced JSJ splitting

This section is devoted to two immediate consequences of the existence of the enhanced JSJ splitting obtained in Section 4. The first (Proposition 5.8) connects the JSJ splitting to automorphism growth, which will be important in the proof of Theorems A and B. The second consequence is the existence of a "complexity-reduction" homomorphism from Out(G) to a finite product of groups  $Out(P_i)$ , where  $P_i \leq G$  are simpler special groups (Proposition 5.12). This will allow us to prove Theorem C from the introduction (Corollary 5.15).

5.1. Beat and controlled subgroups. We begin by introducing some useful notation and terminology in this brief subsection. Let G be a finitely generated group.

Recall that, given two abstract growth rates  $[a_n], [b_n] \in \mathfrak{G}$ , we write  $[a_n] \prec [b_n]$  if we have  $[a_n] \preceq [b_n]$  and  $[a_n] \not\sim [b_n]$ . Equivalently, we have  $\limsup_n a_n/b_n < +\infty$  and  $\liminf_n a_n/b_n = 0$ . We will generally omit square brackets from growth rates to simplify notation.

**Definition 5.1.** Consider some  $\phi \in \text{Out}(G)$  and some abstract growth rate  $\mathfrak{o} \in \mathfrak{G}$ .

- (1) An element  $g \in G$  is  $\mathfrak{o}$ -beat (for  $\phi$ ) if we have  $\|\phi^n(g)\| \prec \mathfrak{o}$ . A subgroup  $H \leq G$  is  $\mathfrak{o}$ -beat if all its elements<sup>17</sup> are  $\mathfrak{o}$ -beat. We denote by  $\mathcal{B}(\mathfrak{o}, \phi)$  (or simply  $\mathcal{B}(\mathfrak{o})$ ) the family of  $\mathfrak{o}$ -beat subgroups of G.
- (2) An element  $g \in G$  is  $\mathfrak{o}$ -controlled if we have  $\|\phi^n(g)\| \leq \mathfrak{o}$ . A subgroup  $H \leq G$  is  $\mathfrak{o}$ -controlled if all its elements are  $\mathfrak{o}$ -controlled. We denote by  $\mathcal{K}(\mathfrak{o}, \phi)$  or  $\mathcal{K}(\mathfrak{o})$  the family of  $\mathfrak{o}$ -controlled subgroups of G.

Given an ultrafilter  $\omega$ , we can also consider the elements  $g \in G$  that are  $\mathfrak{o}$ -beat or  $\mathfrak{o}$ -controlled modulo  $\omega$ , that is, those that satisfy  $\|\phi^n(g)\| \prec_\omega \mathfrak{o}$  and  $\|\phi^n(g)\| \preceq_\omega \mathfrak{o}$ , respectively. We consequently define the families  $\mathcal{B}^{\omega}(\mathfrak{o})$  and  $\mathcal{K}^{\omega}(\mathfrak{o})$  of subgroups that are  $\mathfrak{o}$ -beat or  $\mathfrak{o}$ -controlled modulo  $\omega$ .

In Section 7, we will be particularly interested in the family  $\mathcal{B}_{top}(\phi) := \mathcal{B}(\bar{\mathfrak{o}}_{top}(\phi), \phi)$ , where  $\bar{\mathfrak{o}}_{top}(\phi)$  was introduced in Section 2.2.3, and in the corresponding families  $\mathcal{B}_{top}^{\omega}(\phi)$  modulo ultrafilters.

**Remark 5.2.** For any  $\phi \in \text{Out}(G)$  and  $\mathfrak{o} \in \mathfrak{G}$ , the families  $\mathcal{B}(\mathfrak{o}, \phi)$  and  $\mathcal{K}(\mathfrak{o}, \phi)$  are  $\phi$ -invariant. This is because  $\phi$  is bi-Lipschitz with respect to  $\|\cdot\|$  (Lemma 2.15). The same holds for the families  $\mathcal{B}^{\omega}(\mathfrak{o}, \phi)$  and  $\mathcal{K}^{\omega}(\mathfrak{o}, \phi)$ , for any ultrafilter  $\omega$ .

The following "geometric" characterisation of  $\bar{\mathfrak{o}}_{top}(\phi)$ -beat subgroups will be particularly useful.

**Lemma 5.3.** Let G be a special group and suppose that  $\phi \in \text{Out}(G)$  has infinite order. Fix any (non-principal) ultrafilter  $\omega$ . Then, a subgroup  $B \leq G$  lies in  $\mathcal{B}^{\omega}_{\text{top}}(\phi)$  if and only if B is elliptic in the degeneration  $G \curvearrowright \mathcal{X}_{\omega}$  determined by  $\omega$  and  $\phi$  (as in Section 2.3).

*Proof.* Given a single element  $g \in G$ , we have that g is elliptic in  $\mathcal{X}_{\omega}$  if and only if g is  $\overline{\mathfrak{o}}_{top}(\phi)$ -beat modulo  $\omega$  (Lemma 2.24). Thus, any subgroup of G that is elliptic in  $\mathcal{X}_{\omega}$  lies in  $\mathcal{B}_{top}^{\omega}(\phi)$ .

Conversely, consider a subgroup  $H \in \mathcal{B}^{\omega}_{top}(\phi)$ . Recall from Section 2.3.2 that the degeneration  $\mathcal{X}_{\omega}$ comes with finitely many  $\mathbb{R}$ -trees  $G \curvearrowright T^{v}_{\omega}$  (with  $v \in \Gamma$  for a chosen convex-cocompact embedding  $G \hookrightarrow A_{\Gamma}$ ). The subgroup H is elliptic in  $\mathcal{X}_{\omega}$  if and only if H is elliptic in all  $T^{v}_{\omega}$ . Moreover, every element of H is elliptic in  $\mathcal{X}_{\omega}$ , and hence in all  $T^{v}_{\omega}$ . Now, Serre's lemma implies that every finitely generated subgroup of H is elliptic in all trees  $T^{v}_{\omega}$  [Ser03, p. 64]. Finally, since chains of arc-stabilisers of the G-minimal subtree of  $T^{v}_{\omega}$  have bounded length (see Theorem 2.27), it follows that H is itself elliptic in  $T^{v}_{\omega}$  (using Lemma 2.29). In conclusion, H is elliptic in  $\mathcal{X}_{\omega}$ .

5.2. The singular growth rate. Let G be a special group. We now work to connect the enhanced JSJ decomposition of G (Theorem 4.28) to the speed of growth of its outer automorphisms.

Recall from Section 2.2.3 that we have defined a growth rate  $\overline{\mathfrak{o}}_{top}(\phi) \in \mathfrak{G}$  for each  $\phi \in Out(G)$ . We begin with a simple observation.

**Remark 5.4.** Let  $H \leq G$  be a convex-cocompact subgroup, and let  $\phi \in \text{Out}(G)$  preserve the G-conjugacy class of H. Although the restriction  $\phi|_H \in \text{Out}(H)$  is not uniquely defined in general (Remark 4.12), the growth rate  $\overline{\mathfrak{o}}_{\text{top}}(\phi|_H)$  is well-defined. Indeed, any two possible restrictions  $\phi|_H$ 

<sup>&</sup>lt;sup>17</sup>To avoid any confusion, we stress that we always compute conjugacy lengths with respect to finite generating sets of G, not those of H. In fact, we allow H to be infinitely generated.

differ by the restriction to H of an inner automorphism of G, and so conjugacy lengths grow at the same speed under their powers. (Recall that, since H is convex-cocompact, it does not matter whether we compute conjugacy lengths with respect to a finite generating set of H or G.)

Let  $\operatorname{Out}^0(G) \leq \operatorname{Out}(G)$  be the finite-index subgroup that preserves each *G*-conjugacy class in  $\mathcal{S}(G)$ . This subgroup has indeed finite index because  $\mathcal{S}(G)$  is  $\operatorname{Out}(G)$ -invariant and consists of finitely many conjugacy classes of subgroups.

When studying the top growth rate  $\overline{\mathfrak{o}}_{top}(\phi)$  of an outer automorphism  $\phi \in \text{Out}(G)$ , it is often possible to reduce the problem to understanding the top growth rates of the restrictions of  $\phi$  to the singular subgroups of G. For this reason, we introduce the following auxiliary growth rate.

**Definition 5.5** (Singular growth rate). For  $\phi \in \text{Out}^0(G)$ , the singular growth rate of  $\phi$  is

$$\overline{\mathfrak{o}}_{\mathrm{sing}}(\phi) := \sum_{S} \overline{\mathfrak{o}}_{\mathrm{top}}(\phi|_{S}),$$

where the sum is taken over finitely many representatives  $S \in \mathcal{S}(G)$  of the *G*-conjugacy classes of singular subgroups. For a general element  $\phi \in \text{Out}(G)$ , we set  $\overline{\mathfrak{o}}_{\text{sing}}(\phi) := \frac{1}{k} * \overline{\mathfrak{o}}_{\text{top}}(\phi^k)$  for any integer  $k \in \mathbb{N}$  such that  $\phi^k \in \text{Out}^0(G)$  (using the operations introduced in Section 2.2.4).

When  $\mathcal{S}(G) = \emptyset$  (that is, when G is hyperbolic), we simply set  $\overline{\mathfrak{o}}_{sing}(\phi) := [1]$ . If  $\omega$  is a (non-principal) ultrafilter, we similarly write  $\mathfrak{o}_{sing}^{\omega}(\phi)$  for the projection of  $\overline{\mathfrak{o}}_{sing}(\phi)$  to  $\mathfrak{G}_{\omega}$ .

**Lemma 5.6.** For any  $\phi \in \text{Out}(G)$  and  $g \in G$  with  $Z_G(g) \not\cong \mathbb{Z}$ , we have  $\|\phi^n(g)\| \leq \overline{\mathfrak{o}}_{sing}(\phi)$ .

Proof. Since  $\phi$  is bi-Lipschitz with respect to conjugacy length (Lemma 2.15), it suffices to prove the statement for a power of  $\phi$ . Thus, we can assume that  $\phi \in \text{Out}^0(G)$ . Since  $Z_G(g) \not\cong \mathbb{Z}$ , Proposition 4.5(6) shows that  $g \in S$  for some  $S \in \mathcal{S}(G)$  (unless g = 1 and G is hyperbolic). Thus, recalling Lemma 2.19, we obtain  $\|\phi^n(g)\| \leq \overline{\mathfrak{o}}_{\text{top}}(\phi|_S) \leq \overline{\mathfrak{o}}_{\text{sing}}(\phi)$  as claimed.

We will also need the following lemma. In the special case when G is a  $(\mathcal{Z}(G), \mathcal{S}(G))$ -rigid special group, this implies that  $\overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{sing}(\phi)$  for all  $\phi \in Out(G)$ . As mentioned above, this allows us to reduce the study of the top growth rate to lower-complexity subgroups.

**Lemma 5.7.** Suppose that G is  $(\mathcal{Z}(G), \mathcal{H})$ -rigid, for a collection of subgroups  $\mathcal{H} \supseteq \mathcal{S}(G)$ . Consider an outer automorphism  $\phi \in \text{Out}(G)$  and suppose that there exists an abstract growth rate  $\mathfrak{o} \in \mathfrak{G}$ such that all subgroups in  $\mathcal{H}$  are  $\mathfrak{o}$ -controlled. Then, we have  $\overline{\mathfrak{o}}_{\text{top}}(\phi) \preceq \mathfrak{o}$ .

*Proof.* We can assume that  $G \notin \mathcal{S}(G)$ , otherwise  $G \in \mathcal{H}$  and the lemma is clear using Remark 2.21.

Now, suppose for the sake of contradiction that  $\overline{\mathfrak{o}}_{top}(\phi) \not\preceq \mathfrak{o}$ . This implies that there exists an ultrafilter  $\omega$  such that  $\mathfrak{o}_{top}^{\omega}(\phi) \succ_{\omega} \mathfrak{o}$  (Remark 2.14). In particular, the latter means that  $\mathfrak{o}_{top}^{\omega}(\phi) \succ_{\omega}$  [1], so  $\phi$  has infinite order in Out(G). Realising G as a convex-cocompact subgroup of a right-angled Artin group  $A_{\Gamma}$ , we can thus consider the degeneration  $G \curvearrowright \mathcal{X}_{\omega}$  associated  $\phi \in Out(G)$  and the ultrafilter  $\omega$ , as in Section 2.3.2. Recall that  $\mathcal{X}_{\omega}$  embeds G-equivariantly in a finite product of  $\mathbb{R}$ -trees  $T_{\omega}^{v}$  with  $v \in \Gamma$ . Fix a vertex  $v \in \Gamma$  such that the  $\mathbb{R}$ -tree  $G \curvearrowright T_{\omega}^{v}$  is non-elliptic.

By Lemma 5.3, all subgroups in  $\mathcal{H}$  are elliptic in  $T^{v}_{\omega}$ . Indeed, for any  $h \in \mathcal{H} \in \mathcal{H}$ , we have  $\|\phi^{n}(h)\| \leq \mathfrak{o} \prec_{\omega} \mathfrak{o}^{\omega}_{top}(\phi)$ , by hypothesis, and hence  $\mathcal{H} \subseteq \mathcal{B}^{\omega}_{top}(\phi)$ .

Since  $\mathcal{H} \supseteq \mathcal{S}(G)$ , it follows that all elements of  $\mathcal{S}(G)$  are elliptic in  $T^v_{\omega}$ . This implies that the action  $G \curvearrowright T^v_{\omega}$  has no perverse lines (Definition 2.25). Indeed, the stabiliser of a perverse line  $\alpha$  is always a centraliser  $Z \in \mathcal{Z}(G)$  that is not elliptic in  $T^v_{\omega}$  and such that the kernel of the action  $Z \curvearrowright \alpha$  does not lie in  $\mathcal{Z}(G)$ . Since Z is not elliptic, it cannot be contained in an element of  $\mathcal{S}(G)$ , and hence we must have  $Z \in \mathcal{Z}_c(G)$  and  $Z \cong \mathbb{Z}$ . It follows that the kernel of the action  $Z \curvearrowright \alpha$  is the trivial subgroup  $\{1\}$ , which lies in  $\mathcal{Z}(G)$  (otherwise G would have nontrivial centre, violating the fact that  $G \notin \mathcal{S}(G)$ ).

Now, since there are no perverse lines in  $G \curvearrowright T^v_{\omega}$ , the *G*-stabiliser of every arc of  $\operatorname{Min}(G, T^v_{\omega})$ lies in  $\mathcal{Z}(G)$ , by Theorem 2.27. We can thus appeal to Corollary B.3(3) in the appendix to extract from  $T^v_{\omega}$  a  $(\mathcal{Z}(G), \mathcal{H})$ -splitting of *G*, contradicting the hypothesis that *G* is  $(\mathcal{Z}(G), \mathcal{H})$ -rigid.  $\Box$ 

We are now ready to prove the main result of this section. This is a direct consequence of the enhanced JSJ splitting constructed in Theorem 4.28, and it will prove particularly useful in Sections 6 and 7.

For  $\phi \in \text{Out}(G)$ , we denote by  $\mathcal{K}_{\text{sing}}(\phi)$  the collection of  $\overline{\mathfrak{o}}_{\text{sing}}(\phi)$ -controlled subgroups of G. The collection  $\mathcal{K}_{\text{sing}}(\phi)$  always contains both  $\mathcal{S}(G)$  and the family of cyclic subgroups of G whose G-conjugacy class has finite  $\phi$ -orbit. Moreover,  $\mathcal{K}_{\text{sing}}(\phi)$  is  $\phi$ -invariant (Remark 5.2).

**Proposition 5.8.** Let G be special and 1-ended. For any  $\phi \in \text{Out}(G)$ , there exists a  $\phi$ -invariant  $(\mathcal{ZZ}(G), \mathcal{K}_{\text{sing}}(\phi))$ -tree  $G \curvearrowright T$  such that the G-stabiliser of each vertex of T is:

- (a) either an optimal quadratically hanging subgroup relative to  $\mathcal{K}_{sing}(\phi)$ ;
- (b) or a convex-cocompact root-closed subgroup of G that lies in  $\mathcal{K}_{sing}(\phi)$ .

Moreover, each edge  $e \subseteq T$  with G-stabiliser not in  $\mathcal{Z}(G)$  is incident to a type (a) vertex.

Proof. Theorem 4.28 gives a  $\phi$ -invariant  $(\mathcal{ZZ}(G), \mathcal{K}_{sing}(\phi))$ -tree  $G \curvearrowright T$  such that each vertex group is either of type (a), or it is a convex-cocompact root-closed subgroup  $V \leq G$  that is  $(\mathcal{Z}(V), \mathcal{K}_{sing}(\phi)|_V)$ -rigid in itself. Moreover, every edge of T with G-stabiliser not in  $\mathcal{Z}(G)$  is incident to a type (a) vertex. We wish to show that T is the tree we are looking for, which simply amounts to showing that all vertex groups not of type (a) lie in  $\mathcal{K}_{sing}(\phi)$ .

Thus, consider a convex-cocompact vertex group  $V \leq G$  that is  $(\mathcal{Z}(V), \mathcal{K}_{sing}(\phi)|_V)$ -rigid in itself. Note that, for every subgroup  $H \in \mathcal{K}_{sing}(\phi)|_V$  and every element  $h \in H$ , we have  $\|\phi^n(h)\| \leq \overline{\mathfrak{o}}_{sing}(\phi)$ , by definition (it does not matter whether we compute conjugacy lengths with respect to finite generating sets of V or G, as V is convex-cocompact). Up to raising  $\phi$  to a power, we can assume that it preserves the G-conjugacy class of V, since T is  $\phi$ -invariant. Now, Lemma 5.7 shows that  $\overline{\mathfrak{o}}_{top}(\phi|_V) \leq \overline{\mathfrak{o}}_{sing}(\phi)$ , and hence  $V \in \mathcal{K}_{sing}(\phi)$  as required.  $\Box$ 

We will return to the splitting  $G \curvearrowright T$  provided by Proposition 5.8 in Section 6.4, where we will use it to show that  $\phi$  has only a finite number of growth rates above  $\overline{\mathfrak{o}}_{sing}(\phi)$  and each is realised on a quadratically hanging vertex. For convenience, we refer to T as a JSJ tree adapted to  $\phi$  and we will also denote it by  $G \curvearrowright T_{sing}(\phi)$ .

**Corollary 5.9.** Let G be special and 1-ended. For all  $\phi \in \text{Out}(G)$ , the following hold.

- (1) Every element of  $\mathcal{K}_{sing}(\phi)$  is contained in a maximal element of  $\mathcal{K}_{sing}(\phi)$ . There are only finitely many G-conjugacy classes of maximal elements, and they are all convex-cocompact.
- (2) If a subgroup  $H \leq G$  is  $(\mathcal{Z}(G), \mathcal{S}(G))$ -rigid in G, then  $H \in \mathcal{K}_{sing}(\phi)$ .

*Proof.* Let  $G \curvearrowright T$  be JSJ tree adapted to  $\phi$ , as in Proposition 5.8. We can assume that T is a splitting, otherwise G is either QH or an element of  $\mathcal{K}_{sing}(\phi)$ , in which case the corollary is clear. Note that all peripheral subgroups of the QH vertex groups of T are contained in edge groups of T, because G is 1–ended (see [GL17, Lemma 5.16]).

Now, every subgroup  $H \in \mathcal{K}_{\text{sing}}$  fixes a vertex of T. If all vertices fixed by H are of type (a), then H is peripheral in all of these QH vertex groups. By the previous observation, it follows that either  $H = \{1\}$  or H is contained in the stabiliser of an edge  $e \subseteq T$  both of whose vertices are of type (a). In this case,  $G_e$  is a maximal element of  $\mathcal{K}_{\text{sing}}(\phi)$ , it is infinite cyclic, and  $G_e$  is convex-cocompact by Lemma 4.29 (applied to a suitable collapse of the barycentric subdivision of T).

The other option is that H fixes at least one vertex  $x \in T$  that is of type (b). In this case, we have  $G_x \in \mathcal{K}_{sing}(\phi)$  and, up to replacing x, we can assume that  $G_x$  is maximal among G-stabilisers of vertices of T (recall that no convex-cocompact subgroup properly contains a conjugate of itself, by Lemma 2.4(3)). In conclusion, the maximal elements of  $\mathcal{K}_{sing}(\phi)$  are precisely the G-stabilisers of

the edges  $e \subseteq T$  with both vertices of type (a), as well as the maximal G-stabilisers of the type (b) vertices of T. This proves part (1).

Regarding part (2), Remark 4.33 shows that any  $(\mathcal{Z}(G), \mathcal{S}(G))$ -rigid subgroup  $H \leq G$  must fix either a type (b) vertex or an edge connecting two type (a) vertices. Together with the previous discussion, this completes the proof.

5.3. The complexity-reduction homomorphism. This subsection is devoted to the proof of Theorem C: for any virtually special group U, the outer automorphism group Out(U) is boundary amenable, satisfies the Tits alternative, and has finite virtual cohomological dimension. The main ingredient will be the complexity-reduction homomorphism provided by Proposition 5.12 below.

First of all, we need to introduce the correct notion of "complexity", as the proof of Theorem C will be by induction on this quantity. For this — and through most of the coming discussion — we work with an *actual* special group G.

**Definition 5.10.** The *ambient rank*  $\operatorname{ar}(G)$  is the smallest integer r such that G can be embedded as a convex-cocompact subgroup of a right-angled Artin group  $A_{\Gamma}$  where the graph  $\Gamma$  has r vertices.

If G does not virtually split as a direct product, then  $\operatorname{ar}(S) < \operatorname{ar}(G)$  for all  $S \in \mathcal{S}(G)$ ; indeed, S virtually splits as a product, and so it must be contained in a reducible parabolic subgroup of  $A_{\Gamma}$ .

Before we continue, we need to make the following observation, expanding on Remark 4.12.

**Remark 5.11.** Let G be special, let  $H \leq G$  be convex-cocompact, and let  $\mathcal{O} \leq \operatorname{Out}(G)$  be a subgroup preserving the G-conjugacy class of H. Let  $C_H^G \leq \operatorname{Out}(H)$  be the finite subgroup determined by the conjugation action  $N_G(H) \curvearrowright H$ , as in Remark 4.12. If  $C_H^G = \{1\}$  (for instance, if  $N_G(H) = H$ ), then each  $\phi \in \mathcal{O}$  has a uniquely defined restriction  $\phi|_H \in \operatorname{Out}(H)$ . As a consequence, there is a well-defined restriction homomorphism  $\mathcal{O} \to \operatorname{Out}(H)$  in this case.

In general, we can still always find a finite-index subgroup  $\mathcal{O}' \leq \mathcal{O}$  with a well-defined restriction homomorphism  $\mathcal{O}' \to \operatorname{Out}(H)$ . In order to see this, let  $\tilde{\mathcal{O}} \leq \operatorname{Aut}(G)$  be the group of automorphisms of G that leave H invariant and have outer class in  $\mathcal{O}$ . Each element of  $\mathcal{O}$  is represented by at least one automorphism in  $\tilde{\mathcal{O}}$ . Let  $r \colon \tilde{\mathcal{O}} \to \operatorname{Out}(H)$  denote the composition of the restriction homomorphism  $\tilde{\mathcal{O}} \to \operatorname{Aut}(H)$  with the quotient projection  $\operatorname{Aut}(H) \twoheadrightarrow \operatorname{Out}(H)$ . Since H is special,  $\operatorname{Out}(H)$  is residually finite by [AMS16, Corollary 1.2], and so there exists a finite-index subgroup  $\operatorname{Out}'(H) \leq \operatorname{Out}(H)$  that intersects  $C_H^G$  trivially. Let  $\tilde{\mathcal{O}}' \leq \tilde{\mathcal{O}}$  denote the finite-index subgroup  $r^{-1}(\operatorname{Out}'(H))$ , and let  $\mathcal{O}'$  be its projection to a finite-index subgroup of  $\mathcal{O}$ . By construction, any inner automorphism of G that happens to lie in  $\tilde{\mathcal{O}}'$  must restrict to an inner automorphism of H, and thus the restriction of r to  $\tilde{\mathcal{O}}'$  descends to a well-defined homomorphism  $\bar{r} \colon \mathcal{O}' \to \operatorname{Out}(H)$ .

The following result gives the complexity-reduction homomorphism mentioned above. This is a fairly direct consequence of Theorem 4.28, using the work of Levitt on automorphisms preserving a graph of groups [Lev05].

**Proposition 5.12.** Let G be special, 1-ended, and not virtually a direct product. There exist a finiteindex subgroup  $\text{Out}^1(G) \leq \text{Out}(G)$  and finitely many convex-cocompact subgroups  $H_1, \ldots, H_k \leq G$ such that all the following hold:

- (1) for each i, we have  $N_G(H_i) = H_i$  and the G-conjugacy class of  $H_i$  is  $Out^1(G)$ -invariant;
- (2) the kernel of the resulting restriction homomorphism  $\rho: \operatorname{Out}^1(G) \to \operatorname{Out}(H_1) \times \ldots \times \operatorname{Out}(H_k)$ is isomorphic to a special group;
- (3) each  $H_i$  is either isomorphic to a free or surface group, or it has  $\operatorname{ar}(H_i) < \operatorname{ar}(G)$ .

*Proof.* For simplicity, denote by  $\mathcal{S}(G)^*$  the union of  $\mathcal{S}(G)$  with the collection of cyclic subgroups of G whose conjugacy class has finite  $\operatorname{Out}(G)$ -orbit.

As a warm-up, suppose that G is  $(\mathcal{Z}(G), \mathcal{S}^*(G))$ -rigid. Let  $\operatorname{Out}^0(G) \leq \operatorname{Out}(G)$  be the finiteindex subgroup that preserves each G-conjugacy class of subgroups in  $\mathcal{S}(G)$ . For each  $S \in \mathcal{S}(G)$ , we have a well-defined restriction homomorphism  $\operatorname{Out}^0(G) \to \operatorname{Out}(S)$  because  $N_G(S) = S$  (recall Remark 5.11). Choose representatives  $S_1, \ldots, S_k$  of the finitely many *G*-conjugacy classes in  $\mathcal{S}(G)$ and consider the diagonal homomorphism  $\rho$ :  $\operatorname{Out}^0(G) \to \prod_i \operatorname{Out}(S_i)$ . We have  $\operatorname{ar}(S_i) < \operatorname{ar}(G)$  for all *i*. Now, the fact that *G* is rigid implies that ker  $\rho$  is finite: indeed, any infinite sequence in ker  $\rho$ would give a degeneration  $G \curvearrowright \mathcal{X}_\omega$  in which all elements of  $\mathcal{S}^*(G)$  are elliptic, and we would be able to use Corollary B.3(3) to extract a  $(\mathcal{Z}(G), \mathcal{S}^*(G))$ -splitting of *G*, violating rigidity (see the proof of Lemma 5.7 for more details). Finally, since  $\operatorname{Out}(G)$  is residually finite by [AMS16, Corollary 1.2], a finite-index subgroup  $\operatorname{Out}^1(G) \leq \operatorname{Out}^0(G)$  has trivial intersection with ker  $\rho$ , and so the kernel of the restriction of  $\rho$  to  $\operatorname{Out}^1(G)$  is indeed special (it is trivial). This proves the proposition in the warm-up case.

Now, we consider the general case. Theorem 4.28 provides an  $\operatorname{Out}(G)$ -invariant  $(\mathcal{ZZ}(G), \mathcal{S}^*(G))$ tree  $G \curvearrowright T$  (without loss of generality: a splitting) such that each vertex-stabiliser V is either QH relative to  $\mathcal{S}^*(G)$ , or convex-cocompact and  $(\mathcal{Z}(V), \mathcal{S}^*(G)|_V)$ -rigid in itself. Let  $\operatorname{Out}^1(G) \leq \operatorname{Out}(G)$ be the finite-index subgroup that acts trivially on the quotient graph T/G. Let  $V_1, \ldots, V_s$  be representatives of the G-conjugacy classes of (maximal) vertex-stabilisers of T. By the fact that T is relative to  $\mathcal{S}(G)$ , each normaliser  $N_G(V_i)$  is elliptic in T, and hence  $N_G(V_i) = V_i$  for all i (by maximality of  $V_i$ ). Again, this shows that there is no ambiguity in restricting outer automorphisms, and we obtain a natural homomorphism  $\rho$ :  $\operatorname{Out}^1(G) \to \prod_i \operatorname{Out}(V_i)$ .

Up to passing to a further finite-index subgroup  $\operatorname{Out}^2(G) \leq \operatorname{Out}^1(G)$ , we can assume that we have well-defined restriction homomorphisms  $\operatorname{Out}^2(G) \to \operatorname{Out}(E)$  for each edge group E of T (recall Remark 5.11), and that the image of this homomorphism has trivial intersection with the finite subgroup  $C_E^G \leq \operatorname{Out}(E)$  given by the conjugation action  $N_G(E) \curvearrowright E$ .

The kernel of  $\rho: \operatorname{Out}^2(G) \to \prod_i \operatorname{Out}(V_i)$  can be described using [Lev05, Section 2]. In Levitt's terminology, ker  $\rho$  is generated by *bitwists* around the edges of T/G [Lev05, Proposition 2.2]. In fact, we can say more: for any  $\phi \in \ker \rho$ , the restriction of  $\phi$  to each (maximal) vertex group of T is inner and hence, for each edge group E of T, the restriction  $\phi|_E$  lies in  $C_E^G$ . By our choice of  $\operatorname{Out}^2(G)$ , the latter implies that  $\phi|_E$  is trivial. Thus, any element of ker  $\rho$  restricts to the trivial outer automorphism on each edge group of T. A bitwist that is inner on the corresponding edge group represents the same outer class in G as a product of two *twists* (again in Levitt's terminology), so we conclude that ker  $\rho$  is generated by twists. Now, twists around distinct edges commute in  $\operatorname{Out}(G)$ , while twists around a given edge e generate a subgroup of  $\operatorname{Out}(G)$  isomorphic to a direct product of (virtual) direct factors of the centralisers of  $G_e$  in the two vertex groups of e (this uses Lemma 2.4(2)). From this, one can deduce that ker  $\rho$  is isomorphic to a finite direct product of cyclic edge groups and convex-cocompact subgroups of non-QH vertex groups of T (we leave the details of this to the reader). This implies that ker  $\rho$  is special.

Now, if  $V_i$  is a QH vertex group of T, then  $V_i$  is isomorphic to a free or surface group. However, when  $V_i$  is a non-QH vertex group of T, we might still have  $\operatorname{ar}(V_i) = \operatorname{ar}(G)$ . In order to conclude the proof, we thus consider a vertex  $x \in T$  whose stabiliser V is not QH. Let  $\mathcal{O}$  be the image of the restriction homomorphism  $\operatorname{Out}^2(G) \to \operatorname{Out}(V)$ . We wish to construct a homomorphism with trivial kernel from a finite-index subgroup of  $\mathcal{O}$  to a finite product of outer automorphism groups of subgroups of V with strictly lower ambient rank (this will prove the proposition).

Let  $\mathcal{E}_x$  be the collection of G-stabilisers of edges of T incident to the vertex x, and let  $\operatorname{Cyc}(\mathcal{O})$ be the collection of cyclic subgroups of V whose V-conjugacy class has finite  $\mathcal{O}$ -orbit. Recall that V is  $(\mathcal{Z}(V), \mathcal{S}^*(G)|_V)$ -rigid in itself. Since all elements of  $\mathcal{S}^*(G)$  are elliptic in T, it follows that the maximal elements of  $\mathcal{S}^*(G)|_V$  are either elements of  $\mathcal{E}_x$ , or elements of  $\mathcal{S}^*(G)$  that happen to be contained in V. The latter lie in  $\mathcal{S}(V) \cup \operatorname{Cyc}(\mathcal{O})$ . In conclusion, this shows that the group V is  $(\mathcal{Z}(V), \mathcal{S}(V) \cup \mathcal{E}_x \cup \operatorname{Cyc}(\mathcal{O}))$ -rigid in itself.

We can now argue roughly as in the warm-up case. Let  $\mathcal{O}_0 \leq \mathcal{O}$  be a finite-index subgroup that preserves the V-conjugacy class of each subgroup in  $\mathcal{S}(V) \cup \mathcal{E}_x$ , and let  $H_1, \ldots, H_k$  be representatives

of the finitely many V-conjugacy classes of non-cyclic, maximal subgroups in this collection. Each  $H_i$  is self-normalising (by maximality), and we obtain a homomorphism  $\rho' \colon \mathcal{O}_0 \to \prod_i \operatorname{Out}(H_i)$ . Each  $H_i$  is a subgroup of a singular subgroup of G (using Lemma 4.6), and hence  $\operatorname{ar}(H_i) < \operatorname{ar}(G)$  for all i. Now, the kernel of  $\rho'$  is finite: indeed, any infinite sequence in ker  $\rho'$  would give a degeneration  $V \curvearrowright \mathcal{X}_{\omega}$  in which all elements of  $\mathcal{S}(V) \cup \mathcal{E}_x \cup \operatorname{Cyc}(\mathcal{O})$  are elliptic, and we would be able to use Corollary B.3(3) to extract a  $(\mathcal{Z}(V), \mathcal{S}(V) \cup \mathcal{E}_x \cup \operatorname{Cyc}(\mathcal{O}))$ -splitting of V, violating rigidity (again, see the proof of Lemma 5.7 for details). Finally, we use residual finiteness of  $\operatorname{Out}(V)$  to choose a finite-index subgroup  $\mathcal{O}_1 \leq \mathcal{O}_0$  on which  $\rho'$  is injective.

Summing up, we have constructed a homomorphism  $\rho: \operatorname{Out}^2(G) \to \prod_i \operatorname{Out}(V_i)$  with special kernel and, for each  $V_i$  such that  $\operatorname{ar}(V_i) = \operatorname{ar}(G)$  and  $V_i$  is neither a free or surface group, a further injective homomorphism  $\rho'_i: \mathcal{O}_1^i \to \prod_j \operatorname{Out}(H_j)$  where  $\operatorname{ar}(H_j) < \operatorname{ar}(G)$  and  $\mathcal{O}_1^i$  has finite index in the image of  $\operatorname{Out}^2(G) \to \operatorname{Out}(V_i)$ . Composing  $\rho$  with the product of the  $\rho'_i$  (and some identity homomorphisms), this concludes the proof of the proposition.

It is important to stress that Proposition 5.12 gives no information whatsoever on the *image* of the homomorphism  $\rho: \operatorname{Out}^1(G) \to \prod_i \operatorname{Out}(H_i)$ . Better understanding this image would be a reasonable approach to proving finite generation of  $\operatorname{Out}(G)$ , though this would require further ideas.

Before we continue with the proof of Theorem C, we briefly introduce the three properties under consideration in its statement. Let U be a countable, discrete group.

Boundary amenability is also known as coarse amenability, exactness, or Yu's property A [Yu00, HR00]; we refer to [AD02, Oza06, BGH22] for background. If U is boundary amenable, then it satisfies Novikov's conjecture on higher signatures [Hig00, BCH94]. If in addition U is finitely generated, then U is also known to satisfy the coarse Baum–Connes conjecture and the strong Novikov conjecture. Unfortunately, it remains unknown whether Out(G) is finitely generated for all special groups G (though I expect this to be true).

A group U satisfies the *Tits alternative* if, for each subgroup  $H \leq U$  (possibly not finitely generated), either H is virtually solvable or H contains a subgroup isomorphic to the free group  $F_2$ . We will work with a slightly stronger version that we refer to as the *Tits*<sup>\*</sup> alternative: either H is virtually polycylic or H contains  $F_2$ .

The cohomological dimension cd(U) is the supremum of degrees in which U has nontrivial group cohomology, over all coefficient modules. If U has torsion, then cd(U) is infinite. If U is torsion-free, we have cd(U) = cd(U') for all finite-index subgroups  $U' \leq U$ . Thus, if U is virtually torsion-free, one can define the virtual cohomological dimension (vcd for short) vcd(U) as the cohomological dimension of any finite-index torsion-free subgroup of U. See [Bro94] for further background.

We will need the following observations on these properties.

**Remark 5.13.** Let U be a countable, discrete group. Let  $\mathscr{P}$  mean any of the following four properties: boundary amenability, the Tits<sup>\*</sup> alternative, virtual torsion-freeness, finiteness of vcd. The property  $\mathscr{P}$  is stable under the following basic constructions. (See e.g. [BGH22, Section 2.4] for boundary amenability and [Bro94] for vcd, when we do not give other references.)

- (1) Arbitrary subgroups. If U satisfies  $\mathscr{P}$  and  $V \leq U$  is any subgroup, then V satisfies  $\mathscr{P}$ .
- (2) Finite-index overgroups. If  $U \leq V$  has finite-index and U satisfies  $\mathscr{P}$ , then V satisfies  $\mathscr{P}$ .
- (3) Finite direct products. If  $U_1, \ldots, U_k$  satisfy  $\mathscr{P}$ , then the product  $U_1 \times \ldots \times U_k$  satisfies  $\mathscr{P}$ .
- (4) Extensions. Suppose that we have a short exact sequence  $1 \to N \to U \to Q \to 1$ . If the groups N and Q are boundary amenable, then so is U [KW99, Theorem 5.1]. If N and Q satisfy the Tits<sup>\*</sup> (resp. Tits) alternative, then so does U; indeed, if N and Q are virtually polycyclic (resp. virtually solvable), then so is U (see e.g. [D] and [Din12, Lemma 5.5]). If N and Q have finite vcd and U is virtually torsion-free, then vcd(U)  $\leq$  vcd(N)+vcd(Q)  $< +\infty$ . Finally, if N is torsion-free and Q is virtually torsion-free, then U is virtually torsion-free.

(5) Quotients. Suppose that  $N \triangleleft U$ . If N is amenable and U is boundary amenable, then U/N is boundary amenable. If N is finite and U satisfies the Tits<sup>\*</sup> alternative, then U/N satisfies the Tits<sup>\*</sup> alternative (finite-by-polycyclic groups are polycyclic-by-finite).

The next lemma is needed to extend results on Out(G), with G special, to results on Out(U), with U virtually special. The proof of similar to that of [Krs92, Section 2] and [GL07, Lemma 5.4].

**Lemma 5.14.** Let U be a finitely generated group and let  $L \leq U$  be a finite-index subgroup.

- (1) If Out(L) is boundary amenable, then Out(U) is boundary amenable.
- (2) If  $\operatorname{Out}(L)$  satisfies the Tits alternative, then  $\operatorname{Out}(U)$  satisfies the Tits alternative. If L is normal, has finitely generated centre, and  $\operatorname{Out}(L)$  satisfies the Tits<sup>\*</sup> alternative, then  $\operatorname{Out}(U)$  satisfies the Tits<sup>\*</sup> alternative.
- (3) If U and Out(L) are virtually torsion-free and have finite vcd, and if Out(U) is virtually torsion-free, then Out(U) has finite vcd.
- (4) If L is normal in U, if the centre of L is finitely generated, if Out(L) is virtually torsion-free, and if Out(U) is residually finite, then Out(U) is virtually torsion-free.

*Proof.* For convenience, let  $\mathscr{P}$  denote any of the four properties under consideration: boundary amenability, the Tits<sup>\*</sup> (or Tits) alternative, finiteness of vcd, virtual torsion-freeness.

Let  $K_L \leq \operatorname{Aut}(U)$  be the subgroup of automorphisms that fix L pointwise. To begin with, we need to understand the structure of  $K_L$ . (Note that  $K_L$  can be infinite, for instance when U is the Klein bottle group and L is its index-2 torus subgroup; see e.g. [sTe].)

**Claim.** The group  $K_L$  virtually embeds in a finite direct product  $C \times \ldots \times C$ , where C is the centre of a finite-index subgroup of L. In particular,  $K_L$  is virtually abelian.

Proof of claim. Let  $L_0 \leq L$  be a finite-index subgroup that is normal in U. Let  $K_L^0 \leq K_L$  be a finite-index subgroup acting trivially on the coset space  $U/L_0$ . Write  $U = u_1L_0 \sqcup \cdots \sqcup u_kL_0$ , fixing once and for all some coset representatives  $u_i \in U$ . For any  $\varphi \in K_L^0$  and  $1 \leq i \leq k$ , we have  $\varphi(u_iL_0) = (u_iL_0)$  and so there are elements  $\ell_i(\varphi) \in L_0$  such that  $\varphi(u_i) = u_i\ell_i(\varphi)$ . Note that, for  $\varphi, \psi \in K_L^0$ , we have  $\varphi\psi(u_i) = \varphi(u_i)\ell_i(\psi) = u_i\ell_i(\varphi)\ell_i(\psi)$  and thus each map  $\ell_i \colon K_L^0 \to L_0$  is a homomorphism. It follows that the tuple  $(\ell_1, \ldots, \ell_k)$  gives a group embedding  $K_L^0 \to L_0^k$ . Now, for any  $x \in L_0$ , we have  $u_i x u_i^{-1} \in L_0$  because  $L_0$  is normal in U. Hence, for any  $\varphi \in K_L^0$ , we have

$$u_{i}xu_{i}^{-1} = \varphi(u_{i}xu_{i}^{-1}) = \varphi(u_{i})x\varphi(u_{i})^{-1} = u_{i}\ell_{i}(\varphi)x\ell_{i}(\varphi)^{-1}u_{i}^{-1}$$

which yields  $x = \ell_i(\varphi) x \ell_i(\varphi)^{-1}$  for all  $x \in L_0$ . This shows that each homomorphism  $\ell_i \colon K_L^0 \to L_0$  takes values in the centre of  $L_0$ , proving the claim.

Now, let  $\operatorname{Aut}_L(U) \leq \operatorname{Aut}(U)$  be the subgroup of automorphisms leaving L invariant. Since U is finitely generated, it has only finitely many subgroups of the same index as L, and hence  $\operatorname{Aut}_L(U)$  has finite index in  $\operatorname{Aut}(U)$ . Consider the homomorphisms

$$\alpha : \operatorname{Aut}_L(U) \to \operatorname{Out}(U), \qquad \beta : \operatorname{Aut}_L(U) \to \operatorname{Out}(L),$$

defined as follows:  $\alpha$  is the composition of the inclusion  $\operatorname{Aut}_L(U) \hookrightarrow \operatorname{Aut}(U)$  with the quotient projection  $\operatorname{Aut}(U) \twoheadrightarrow \operatorname{Out}(U)$ , while  $\beta$  is the composition of the restriction  $\operatorname{Aut}_L(U) \to \operatorname{Aut}(L)$ with the quotient projection  $\operatorname{Aut}(L) \twoheadrightarrow \operatorname{Out}(L)$ .

The image  $\operatorname{im}(\alpha)$  has finite index in  $\operatorname{Out}(U)$ , while the image  $\operatorname{im}(\beta)$  is just some subgroup of  $\operatorname{Out}(L)$ . By Remark 5.13(1) we know that  $\operatorname{im}(\beta)$  satisfies property  $\mathscr{P}$ , and by Remark 5.13(2) it suffices to prove that  $\operatorname{im}(\alpha)$  satisfies  $\mathscr{P}$  in order to obtain it for  $\operatorname{Out}(U)$  (in each of the four parts of the lemma). Note that  $\operatorname{ker}(\alpha) \leq \operatorname{Aut}(U)$  is the group of conjugations by elements of the normaliser  $N_U(L)$ . As such, it has a finite-index subgroup  $\operatorname{ker}^0(\alpha)$  formed by conjugations by elements of L, and we have  $\operatorname{ker}^0(\alpha) \triangleleft \operatorname{Aut}_L(U)$ . The group  $\operatorname{ker}(\beta)$  is the group of automorphisms of U restricting to an inner automorphism on L, so it splits as a product  $\operatorname{ker}^0(\alpha) \cdot K_L$ .

Now, we have an epimorphism

$$\eta: \operatorname{Aut}_L(U) / \operatorname{ker}^0(\alpha) \twoheadrightarrow \operatorname{Aut}_L(U) / \operatorname{ker}(\beta) \cong \operatorname{im}(\beta)$$

with kernel  $\ker(\beta)/\ker^0(\alpha) \cong K_L/(K_L \cap \ker^0(\alpha))$ . By the claim,  $K_L$  is virtually abelian, and hence  $\ker(\eta)$  is virtually abelian as well. Moreover,  $\operatorname{im}(\eta)$  satisfies property  $\mathscr{P}$ , and the group  $\operatorname{Aut}_L(U)/\ker^0(\alpha)$  maps onto  $\operatorname{im}(\alpha) \cong \operatorname{Aut}_L(U)/\ker(\alpha)$  with finite kernel  $\ker(\alpha)/\ker^0(\alpha)$ .

Using parts (4) and (5) of Remark 5.13, these observations imply that  $im(\alpha)$  is boundary amenable and satisfies the Tits alternative, provided that the respective property holds for  $im(\beta)$ . As discussed above, this proves part (1) and the first half of part (2) of the lemma.

Let  $\operatorname{Aut}_{L}^{0}(U) \leq \operatorname{Aut}_{L}(U)$  be the finite-index subgroup that acts trivially on the coset space U/L. If L is normal in U, we can take  $L_{0} = L$  in the proof of the claim, which then shows that  $K_{L} \cap \operatorname{Aut}_{L}^{0}(U)$  embeds in a product of finitely many copies of the centre of L. In particular  $K_{L} \cap \operatorname{Aut}_{L}^{0}(U)$  is abelian, and it is finitely generated if the centre of L is finitely generated. Thus, under the assumptions of the second half of part (2),  $K_{L}$  and  $\operatorname{ker}(\eta)$  are virtually polycyclic, and Remark 5.13 implies that  $\operatorname{Out}(U)$  satisfies the Tits<sup>\*</sup> alternative, completing the proof of part (2).

We now prove part (3). Suppose that  $\operatorname{Out}(U)$  is virtually torsion-free, and that U and  $\operatorname{Out}(L)$  have finite vcd. Let  $\mathcal{U} \leq \operatorname{Out}(U)$  and  $\mathcal{L} \leq \operatorname{Out}(L)$  be finite-index torsion-free subgroups, and let  $\mathcal{I}$  be the projection of the intersection  $\alpha^{-1}(\mathcal{U}) \cap \beta^{-1}(\mathcal{L})$  to the quotient  $\operatorname{Aut}_L(U)/\ker^0(\alpha)$ . Thus,  $\mathcal{I}$  is a finite-index subgroup of  $\operatorname{Aut}_L(U)/\ker^0(\alpha)$ , and the image of the restricted homomorphism  $\eta|_{\mathcal{I}}$  is torsion-free with finite cohomological dimension, as it is a subgroup of  $\mathcal{L}$ .

Let  $F \triangleleft \mathcal{I}$  be the finite subgroup obtained by intersecting  $\mathcal{I}$  with  $\ker(\alpha)/\ker^{0}(\alpha)$ . The quotient  $\mathcal{I}/F$  is a finite-index subgroup of  $\operatorname{Aut}_{L}(U)/\ker(\alpha) \cong \operatorname{im}(\alpha)$ , and it is identified with a finite-index subgroup of  $\mathcal{U} \leq \operatorname{Out}(U)$ , so it is torsion-free. Since there is no torsion in the image of  $\eta|_{\mathcal{I}}$ , we have  $F \triangleleft \ker(\eta|_{\mathcal{I}})$  and  $\eta$  yields a short exact sequence

$$1 \to \ker(\eta|_{\mathcal{I}})/F \to \mathcal{I}/F \to \operatorname{im}(\eta|_{\mathcal{I}}) \to 1.$$

The kernel of  $\eta|_{\mathcal{I}}$  is a finite-index subgroup of ker $(\eta) \cong K_L/(K_L \cap \ker^0(\alpha))$ , and so it is virtually abelian with finite vcd by the claim (since U has finite vcd by hypothesis). Any finite-index torsionfree subgroup of ker $(\eta|_{\mathcal{I}})$  projects injectively to a finite-index subgroup of ker $(\eta|_{\mathcal{I}})/F$ , and thus the latter has finite vcd as well. Summing up,  $\mathcal{I}/F$  is torsion-free, and both image and kernel of  $\eta$  restricted to  $\mathcal{I}/F$  have finite vcd. Using Remark 5.13(4), we conclude that  $\mathcal{I}/F$  has finite vcd. Since the latter is a finite-index subgroup of Out(U), this proves part (3).

Finally, we discuss part (4) of the lemma. Let  $\mathcal{U}'$  be the finite-index subgroup of  $\operatorname{Aut}_L(U)/\ker^0(\alpha)$  that is the projection of  $\operatorname{Aut}_L^0(U) \leq \operatorname{Aut}_L(U)$ . We have already observed above that the hypotheses of part (4) imply that  $K_L \cap \operatorname{Aut}_L^0(U)$  is abelian and finitely generated.

Let again  $\mathcal{L} \leq \operatorname{Out}(L)$  be a finite-index torsion-free subgroup, and consider  $\mathcal{L}' := \beta^{-1}(\mathcal{L})/\ker^0(\alpha)$ and  $\mathcal{I}' := \mathcal{U}' \cap \mathcal{L}'$ , which are finite-index subgroups of  $\operatorname{Aut}_L(U)/\ker^0(\alpha)$ . The homomorphism  $\eta$ restricted to  $\mathcal{I}'$  gives a short exact sequence

$$1 \to \ker(\eta|_{\mathcal{I}'}) \to \mathcal{I}' \to \mathcal{L},$$

where as above  $\ker(\eta|_{\mathcal{I}'})$  is a quotient of  $K_L \cap \operatorname{Aut}^0_L(U)$ . In particular, this kernel is abelian and finitely generated. Let  $F' \triangleleft \mathcal{I}'$  be the intersection of  $\mathcal{I}'$  with the finite group  $\ker(\alpha)/\ker^0(\alpha)$ . As above, the fact that  $\mathcal{L}$  is torsion-free implies that F' is contained in the kernel of  $\eta|_{\mathcal{I}'}$  and we obtain:

$$1 \to \ker(\eta|_{\mathcal{I}'})/F' \to \mathcal{I}'/F' \to \mathcal{L}.$$

Now,  $\mathcal{I}'/F'$  is a finite-index subgroup of  $\operatorname{Out}(U)$ , and hence it is residually finite by our hypotheses. Since the group  $\operatorname{ker}(\eta|_{\mathcal{I}'})/F'$  is abelian and finitely generated, it has only finitely many torsion elements. A finite-index subgroup of  $\mathcal{I}'/F'$  then avoids all of the latter, by residual finiteness, and hence this finite-index subgroup is torsion-free by Remark 5.13(4). This is the required finite-index torsion-free subgroup of  $\operatorname{Out}(U)$ , concluding the proof of part (4). We are finally ready to prove Theorem C, which we restate here for convenience.

**Corollary 5.15.** The following hold for every virtually special group G:

- (1) Out(G) is boundary amenable;
- (2) Out(G) satisfies the Tits<sup>\*</sup> alternative: subgroups not containing  $F_2$  are virtually polycyclic;
- (3) Out(G) is virtually torsion-free and it has finite vcd.

*Proof.* To begin with, note that it suffices to prove the corollary for an actual special group G. Indeed, every virtually special group has a finite-index, normal, special subgroup. Special groups have finite cohomological dimension (as they have a finite classifying space) and their abelian subgroups have bounded rank. Moreover, for any virtually special group U, the group Out(U) is residually finite by [AMS16, Corollary 1.2]. Thus, using Lemma 5.14, it suffices to prove each of the three parts of the corollary for a finite-index special subgroup.

In the rest of the proof we therefore assume that G is special. We argue by induction on the ambient rank of G, with the base case being trivial. For the inductive step, suppose that, for every special group H with  $\operatorname{ar}(H) < \operatorname{ar}(G)$ , we know that  $\operatorname{Out}(H)$  satisfies the corollary.

We can assume that G is 1-ended. Indeed, in general G admits a free splitting  $G = G_1 \ast \cdots \ast G_k \ast F_m$ for some  $k, m \ge 0$ , where the  $G_i$  are 1-ended. All the factors  $G_i$  are special (e.g. by Proposition 2.30) and thus they are boundary amenable [CN05, Theorem 13], they satisfy the Tits<sup>\*</sup> alternative [SW05], and they are torsion-free with finite vcd; if  $Z_i$  denotes the centre of  $G_i$ , then the quotient  $G_i/Z_i$  is virtually special by Lemma 2.4(2), and so it has finite vcd as well. Now, if the  $Out(G_i)$  are boundary amenable, then so is Out(G) by [BGH22, Corollary 5.3]; if the  $Out(G_i)$  satisfy the Tits<sup>\*</sup> alternative, then so does Out(G) by [Hor14, Theorem 6.1]; and if the  $Out(G_i)$  are virtually torsion-free and have finite vcd, then so does Out(G) by [GL07, p. 709, Theorem 5.2(i)].

Now, suppose for a moment that G virtually splits as a direct product. Let  $L \leq G$  be a finite-index subgroup of the form  $L_1 \times \ldots \times L_k \times \mathbb{Z}^m$ , where  $k + m \geq 2$  and each  $L_i$  is directly indecomposable and has trivial centre. By Lemma 5.14, it again suffices to prove the corollary for  $\operatorname{Out}(L)$ . Note that  $\operatorname{ar}(L_i) < \operatorname{ar}(G)$  for all *i*. By Lemma A.5, automorphisms of L permute the subgroups  $\langle G_i, \mathbb{Z}^m \rangle$ of L. Let  $\operatorname{Out}^*(L) \leq \operatorname{Out}(L)$  be the group of outer classes of automorphisms of L leaving each  $G_i$ invariant and fixing pointwise the centre  $\mathbb{Z}^m$ . Denoting by  $L_{ab}$  the abelianisation of L, the natural homomorphism  $\eta$ :  $\operatorname{Out}(L) \to \operatorname{Out}(L_{ab})$  has  $\ker(\eta)$  virtually contained in  $\operatorname{Out}^*(L)$ .

The group  $\operatorname{Out}(L_{ab})$  is isomorphic to  $\operatorname{GL}_p(\mathbb{Z})$  for some  $p \in \mathbb{N}$ , which is boundary amenable by [GHW05], satisfies the Tits<sup>\*</sup> alternative by [Tit72, Mal51], is virtually torsion-free by Selberg's lemma, and has finite vcd because it acts properly on the symmetric space of  $\operatorname{SL}_p(\mathbb{R})$ . On the other hand, we have  $\operatorname{Out}^*(L) \cong \prod_i \operatorname{Out}(G_i)$ , which satisfies all parts of the corollary by the inductive hypothesis; by Remark 5.13(1), it follows that ker( $\eta$ ) is boundary amenable, satisfies the Tits<sup>\*</sup> alternative, and has finite vcd. Using Remark 5.13(4), this implies that  $\operatorname{Out}(L)$  is boundary amenable and satisfies the Tits<sup>\*</sup> alternative, as this is true of both the image and the kernel of the homomorphism  $\eta$ . The same argument shows that  $\operatorname{Out}(L)$  has finite vcd, provided that we find a finite-index subgroup of  $\operatorname{Out}(L)$  that has torsion-free intersection with  $\operatorname{Out}^*(L)$ ; the existence of the latter is a straightforward consequence of the description of automorphisms of a product in terms of formal triangular matrices, as in Appendix A.1, using that  $\operatorname{Out}^*(L)$  is virtually torsion-free by the inductive hypothesis. Summing up, this proves the entire corollary when G virtually splits as a direct product.

In conclusion, we can assume that G is 1-ended and that it does not virtually split as a direct product. We can then invoke Proposition 5.12, which yields a short exact sequence

$$1 \longrightarrow K \longrightarrow \operatorname{Out}^{1}(G) \longrightarrow \operatorname{Out}(H_{1}) \times \ldots \times \operatorname{Out}(H_{k}),$$

where  $\operatorname{Out}^1(G) \leq \operatorname{Out}(G)$  has finite index, K is special (in particular, torsion-free), and we have  $\operatorname{ar}(H_i) < \operatorname{ar}(G)$  for each  $H_i$  that is not a free or surface group. Outer automorphism groups

of surface and free groups are boundary amenable [Ham09, BGH22], satisfy the Tits<sup>\*</sup> alternative [McC85, BFH00, BFH05], are virtually torsion-free, and have finite vcd [CV86]. Also using the inductive assumption, it follows that the product  $\prod_i \text{Out}(H_i)$  has all these properties, and so does the group K by specialness. One last application of Remark 5.13 finally implies that Out(G) satisfies these properties as well, concluding the proof.

We can also deduce that Out(G) does not have any non-unimodular Baumslag–Solitar subgroups.

**Corollary 5.16.** If G is virtually special and Out(G) contains a Baumslag–Solitar group BS(m, n), then |m| = |n|.

*Proof.* Without loss of generality, we have  $|m| \leq |n|$ . If  $|m| \neq 1$  and  $|m| \neq |n|$ , then BS(m, n) is not residually finite and so it cannot be a subgroup of Out(G), which is residually finite by [AMS16]. On the other hand, BS(1, n) is solvable and not virtually polycyclic for |n| > 1, which would violate the fact that Out(G) satisfies the Tits<sup>\*</sup> alternative by Corollary 5.15.

### 6. Growth of general automorphisms

In this section we prove Theorem A, describing the top growth rate of a general automorphism of a special group G. We will prove a slightly stronger statement, Theorem 6.6 below, which is the main result of this section.

In order to state the latter, we begin by introducing an important property of automorphisms, *docility*, which will play a fundamental role in the proof. Most of Section 6 is devoted to showing that exponentially-growing automorphisms of special groups are docile, after which Theorem A can be deduced relatively quickly from the existence of the enhanced JSJ splitting (see Section 6.5).

6.1. Sound and docile automorphisms. Recall from Section 2.2 that  $\mathfrak{G}$  is the set of abstract growth rates, i.e. bi-Lipschitz equivalence classes of sequences in  $\mathbb{R}_{>0}$  bounded away from zero.

**Definition 6.1.** An abstract growth rate  $[x_n] \in \mathfrak{G}$  is:

- (1) pure if  $[x_n] \sim [n^p \lambda^n]$  for some  $\lambda > 1$  and  $p \in \mathbb{N}$ ;
- (2)  $(\lambda, p)$ -tame, for some  $\lambda > 1$  and  $p \in \mathbb{N}$ , if we have  $[x_n] \sim [a_n \lambda^n]$  for a weakly increasing sequence  $a_n$  with  $[1] \leq [a_n] \leq [n^p]$ .

Pure growth rates are clearly tame, but the converse does not hold. Note that pure growth rates form a *totally* ordered subset of  $(\mathfrak{G}, \preceq)$ , while tame ones do not. Also note that tame rates are at least exponential, as we ask that  $\lambda > 1$ .

Here in Section 6, we are mostly interested in tame growth rates: we will show that the top growth rate of any automorphism of a special group is either sub-polynomial or tame, which will lead to Theorem A. We do not know if this top growth rate is pure in general, so pure rates will only return in Section 7, where we study coarse-median preserving automorphisms and prove Theorem B.

The following is a technical definition tailored to the needs of this section; we briefly motivate it below. Recall the notation  $\overline{\mathcal{O}}_{top}(\cdot)$  and  $\overline{\mathfrak{o}}_{top}(\cdot)$  introduced in Section 2.2.3. For a finitely generated group G and any  $\varphi \in \operatorname{Aut}(G)$  with outer class  $\phi \in \operatorname{Out}(G)$ , we have  $\overline{\mathcal{O}}_{top}(\varphi) \succeq \overline{\mathfrak{o}}_{top}(\phi)$ .

**Definition 6.2.** Let G be finitely generated. Consider  $\varphi \in \operatorname{Aut}(G)$  with outer class  $\phi \in \operatorname{Out}(G)$ .

- (1) The automorphism  $\varphi$  is *sound* if we have  $\overline{\mathcal{O}}_{top}(\varphi) \sim \overline{\mathfrak{o}}_{top}(\phi)$ . The outer automorphism  $\phi$  is *sound* if all its representatives  $\varphi' \in \operatorname{Aut}(G)$  are sound.
- (2) The automorphism  $\varphi$  is *docile* if, at the same time,  $\varphi$  is sound and  $\overline{\mathcal{O}}_{top}(\varphi)$  is tame. The outer automorphism  $\phi$  is *docile* if  $\varphi$  is docile (this is independent of the choice of representative, by Remark 6.4 below). We will also say that  $\varphi$  and  $\phi$  are  $(\lambda, p)$ -docile if we wish to specify the parameters for which the rates  $\overline{\mathcal{O}}_{top}(\varphi)$  and  $\overline{\mathfrak{o}}_{top}(\phi)$  are  $(\lambda, p)$ -tame.

Thus, automorphisms are sound when, under their iterates, word length does not grow much faster than conjugacy length: applying powers of the automorphism does not yield elements most of whose word length is due to a failure to be "cyclically reduced". Exponentially-growing automorphisms of free and surface groups are all sound, while inner automorphisms clearly are not.

The crux of Section 6 lies in showing that exponentially-growing automorphisms of special groups are sound (and docile). As train-track techniques are not available in this context, settling this harmless-looking point will require some work. In turn, soundness and docility are extremely important, because they are necessary to relate the growth of an automorphism of G to its growth on the vertex groups of the enhanced JSJ decomposition of G (using e.g. Proposition A.8).

The following are two important observations on the above definitions.

**Remark 6.3.** If  $[x_n] \in \mathfrak{G}$  is tame, we have  $\left[\sum_{i \leq n} x_i\right] \sim [x_n]$ . Indeed, since  $[x_n] \sim [a_n \lambda^n]$  for a weakly increasing sequence  $a_n$  by definition, we obtain  $\sum_{i \leq n} a_i \lambda^i \leq a_n \sum_{i \leq n} \lambda^i \leq a_n \lambda^n$ .

The need for Remark 6.3 is precisely what motivated us to require the sequence  $a_n$  in Definition 6.1 to be weakly increasing (and that  $\lambda \neq 1$ ). We will use the previous remark in many forms, the first of which is the following. Given  $\varphi \in \operatorname{Aut}(G)$  and an element  $g \in G$ , we can define the elements  $g_n := g\varphi(g)\varphi^2(g)\ldots\varphi^n(g)$ . If the growth rate  $[|\varphi^n(g)|]$  is tame, then  $|g_n| \leq |\varphi^n(g)|$ . This leads to the next observation:

**Remark 6.4.** If  $\phi \in \text{Out}(G)$  has a docile representative  $\varphi \in \text{Aut}(G)$ , then *all* its representatives are docile. Indeed, suppose that  $\varphi$  is docile and let  $\psi(x) = g\varphi(x)g^{-1}$  be another representative. One the one hand, we immediately have  $\overline{\mathcal{O}}_{\text{top}}(\psi) \succeq \overline{\mathfrak{o}}_{\text{top}}(\phi) \sim \overline{\mathcal{O}}_{\text{top}}(\varphi)$ . On the other, setting again  $g_n := g\varphi(g)\varphi^2(g)\ldots\varphi^n(g)$ , we have

$$|\psi^n(x)| = |g_{n-1}\varphi^n(x)g_{n-1}^{-1}| \le 2|g_{n-1}| + |\varphi^n(x)|,$$

for all  $x \in G$  and  $n \in \mathbb{N}$ . Thus, recalling Remark 6.3 and the definition of  $\overline{\mathcal{O}}_{top}(\cdot)$ , we also get  $\overline{\mathcal{O}}_{top}(\psi) \preceq \overline{\mathcal{O}}_{top}(\varphi)$ . In conclusion  $\overline{\mathcal{O}}_{top}(\psi) \sim \overline{\mathcal{O}}_{top}(\varphi)$ , which implies that  $\psi$  is docile.

Finally, we say that an element  $[x_n] \in \mathfrak{G}$  is *sub-polynomial* if  $[x_n] \leq [n^p]$  for some  $p \in \mathbb{N}$ . An automorphism  $\varphi \in \operatorname{Aut}(G)$  is *sub-polynomial* if  $\overline{\mathcal{O}}_{\operatorname{top}}(\varphi)$  is sub-polynomial. Note that, a priori, the fact that an outer class  $\phi \in \operatorname{Out}(G)$  has sub-polynomial  $\overline{\mathfrak{o}}_{\operatorname{top}}(\phi)$  does not imply that any automorphism representing  $\phi$  is sub-polynomial. However, we have the following analogue of Remark 6.4.

**Remark 6.5.** If  $\phi \in \text{Out}(G)$  has a representative  $\varphi \in \text{Aut}(G)$  with  $\overline{\mathcal{O}}_{\text{top}}(\varphi) \preceq n^p$  for some  $p \in \mathbb{N}$ , then all representatives  $\psi \in \text{Aut}(G)$  of  $\phi$  satisfy the weaker inequality  $\overline{\mathcal{O}}_{\text{top}}(\psi) \preceq n^{p+1}$ . This is shown as in the previous remark: if  $\psi(x) = g\varphi(x)g^{-1}$ , we have  $|\psi^n(x)| \leq 2|g_{n-1}| + |\varphi^n(x)|$  and:

$$|g_n| \le \sum_{i \le n} |\varphi^i(x)| \le \sum_{i \le n} i^p \le n^{p+1}.$$

6.2. Statement of results. The following is the main result of Section 6. Importantly, it says that automorphisms of special groups always grow either sub-polynomially or at least exponentially, and that the latter are sound. It also implies Theorem A from the Introduction.

Recall that we say that a subgroup  $H \leq G$  is  $\phi$ -invariant for some  $\phi \in \text{Out}(G)$ , if the Gconjugacy class of H is preserved by one/all automorphism of G in the outer class  $\phi$ . We say that
a number  $\lambda$  is a *d*-algebraic integer for  $d \in \mathbb{N}$  if  $\lambda$  is the root of a monic polynomial of degree  $\leq d$ with integer coefficients.

**Theorem 6.6.** Let G be a special group. There exist natural numbers  $\pi = \pi(G)$ , d = d(G) and k = k(G) such that the following hold for every  $\varphi \in Aut(G)$  and its outer class  $\phi \in Out(G)$ .

- (1) Either  $\phi$  is docile, or  $\overline{\mathcal{O}}_{top}(\varphi) \preceq [n^{\pi}]$ .
- (2) If  $\phi$  is docile, there exists a d-algebraic integer  $\lambda > 1$  such that  $\overline{\mathfrak{o}}_{top}(\phi)$  is  $(\lambda, \pi)$ -tame.

- (3) If  $\phi$  is  $(\lambda, \pi)$ -docile, there is a  $\phi^k$ -invariant subgroup  $H \leq G$  of one of the following kinds.
  - (i) H is<sup>18</sup> isomorphic to the fundamental group of a compact surface on which  $\phi^k$  is represented by a pseudo-Anosov homeomorphism with stretch factor  $\lambda^k$ . In particular, all non-peripheral elements  $h \in H$  satisfy  $\|\phi^n(h)\| \sim \lambda^n$ .
  - (ii) H is a convex-cocompact, infinitely-ended subgroup with a non-sporadic factor system  $\mathcal{F}$ such that  $\phi^k$  is a fully irreducible element of  $\operatorname{Out}(H, \mathcal{F})$  and  $\lambda^k$  is its Perron–Frobenius eigenvalue. Moreover, every element  $h \in H$  not conjugate into a subgroup in  $\mathcal{F}$  satisfies either  $\|\phi^n(h)\| \sim 1$  or  $\|\phi^n(h)\| \sim \lambda^n$ . (See Appendix A.4 for terminology.)
  - (iii) H is a convex-cocompact subgroup of the form  $H' \times \mathbb{Z}^m$  for some  $m \ge 1$ , and there exist elements  $h \in H$  with  $\|\phi^n(h)\| \sim n^p \lambda^n$  for some  $0 \le p \le \pi$ . Moreover, either H' is a surface group or infinitely ended, or  $\phi^k$  descends to an automorphism of the abelianisation of H having  $\lambda^k$  as the maximum modulus of an eigenvalue.

**Remark 6.7.** Part (3) of Theorem 6.6 morally says that the only source of exponentially-growing automorphisms of special groups are surface groups, free groups, and free abelian groups. In practice, however, free groups need to be replaced by free products, and free abelian groups need to be replaced by direct products (see Appendix A.1 for subtleties related to the latter).

Note that, although part (3) constructs elements with  $\|\phi^n(h)\| \sim \lambda^n$  or  $\|\phi^n(h)\| \sim n^p \lambda^n$ , there is no claim that these growth rates coincide with  $\overline{\mathfrak{o}}_{top}(\phi)$ , or even that  $\overline{\mathfrak{o}}_{top}(\phi)$  be precisely realised on an element. We only know that  $\overline{\mathfrak{o}}_{top}(\phi)$  is comprised between  $[\lambda^n]$  and  $[n^{\pi}\lambda^n]$ . Things will be cleaner for coarse-median preserving automorphisms in Section 7.

**Remark 6.8.** When  $\phi$  is docile, the base of exponential growth  $\lambda$  is not just any algebraic integer: either it is the stretch factor of a pseudo-Anosov on a compact surface, or it is the Perron–Frobenius eigenvalue of a fully irreducible automorphism of a free group, or it is the maximum modulus of an eigenvalue of a matrix in  $\operatorname{GL}_m(\mathbb{Z})$  for some  $m \in \mathbb{Z}$ . This completely characterises which algebraic integers  $\lambda$  can appear in Theorem 6.6.

In the first two of the three listed cases,  $\lambda$  is a *Perron number* [Thu88, BH92]: an algebraic integer whose modulus is strictly larger than that of all its Galois conjugates. As to the third case, a number  $\lambda$  is the maximum modulus of an eigenvalue of an element of  $\operatorname{GL}_m(\mathbb{Z})$  if and only if  $\lambda$ is both an *algebraic unit* (meaning that both  $\lambda$  and  $\lambda^{-1}$  are algebraic integers) and a *weak Perron number* (meaning that the modulus of  $\lambda$  is at least as big as that of all its Galois conjugates). To see this, it suffices to consider the companion matrix of the minimal polynomial of  $\lambda$ .

Note that Theorem 6.6 is a little stronger than Theorem A, and this is not an irrelevant difference. As we will see, docility is essential for the *entire* proof of Theorem 6.6 to work.

We will prove Theorem 6.6 by induction on the ambient rank  $\operatorname{ar}(G)$  (Definition 5.10). The most technical part of the proof consists of showing that automorphisms of special groups are either sound or sub-polynomially-growing, which is the content of Section 6.3 below. Once that is achieved, the rest of Theorem 6.6 will follow fairly easily using the enhanced JSJ splitting, as explained in Sections 6.4 and 6.5.

6.3. The main technical step. In this subsection, we isolate the most complicated part of the proof of Theorem 6.6, which mainly has to do with the proof of soundness. For this, it is convenient to introduce the following auxiliary concept, which will not be used outside of Section 6.3.

**Definition 6.9.** Let H be a special group. If  $\psi \in \operatorname{Aut}(H)$  is an automorphism with outer class  $[\psi] \in \operatorname{Out}(H)$ , we say that  $\psi$  is  $\alpha$ -good, for some  $\alpha \in \mathbb{N}$ , if **both** following conditions hold:

<sup>&</sup>lt;sup>18</sup>This is the only one of the three cases (i)–(iii) in which H might not be convex-cocompact. However, H is still a quadratically hanging vertex group in a cyclic splitting of a  $\phi^k$ -invariant convex-cocompact subgroup of G. As such, all non-peripheral elements of H are convex-cocompact in G.

- (a) either  $\overline{\mathfrak{o}}_{top}([\psi])$  is sub-polynomial or  $\psi$  is docile;
- (b) if  $\overline{\mathfrak{o}}_{top}([\psi]) \preceq [n^p]$  for some  $p \in \mathbb{N}$ , then  $\overline{\mathcal{O}}_{top}(\psi) \preceq [n^{p+\alpha}]$ .

We say that H is  $\alpha$ -good if all automorphisms of H are  $\alpha$ -good. Finally, H is good if there exists  $\alpha$  such that H is  $\alpha$ -good.

Our goal is the following proposition, which implies most of Theorem 6.6(1).

## **Proposition 6.10.** All special groups are good.

In the coming discussion, we fix a convex-cocompact subgroup  $G \leq A_{\Gamma}$  with  $|\Gamma^{(0)}| = \operatorname{ar}(G)$ , and we consider an automorphism  $\varphi \in \operatorname{Aut}(G)$  with outer class  $\phi \in \operatorname{Out}(G)$ .

We will prove Proposition 6.10 by induction on the ambient rank. As a base step, one can take the case when  $\operatorname{ar}(G) = 1$ , which implies that G is cyclic. However, we will use in the inductive step the fact that all free and surface groups are 1–good, which requires train tracks and for which we do not obtain a new argument. In a sense, that is the true base step.

As an outline of the inductive step, the plan is to consider the enhanced JSJ splitting of G (Theorem 4.28) and show that its rigid vertex groups are good, after which Proposition A.8 in the appendix will quickly imply that G is itself good. The core of the inductive step lies in showing that rigid special groups are either good or relatively hyperbolic (Lemma 6.11), and then in proving goodness in the relatively hyperbolic case using the work of Guirardel and Levitt (Lemma 6.12).

Recall that  $\mathcal{P}(G)$  is the family of *G*-parabolic subgroups of *G*.

**Lemma 6.11.** Let G be  $(\mathcal{Z}(G), \mathcal{H})$ -rigid, for a  $\varphi$ -invariant collection  $\mathcal{H} \supseteq \mathcal{S}(G)$  consisting of finitely many G-conjugacy classes of subgroups of G. Suppose that all non-cyclic elements of  $\mathcal{H}$  are convex-cocompact and that  $G \notin \mathcal{S}(G)$ . Also suppose that there exists  $\alpha_0 \in \mathbb{N}$  such that all subgroups in  $\mathcal{H} \cup \mathcal{P}(G)$  are  $\alpha_0$ -good, and that the automorphism  $\varphi$  is not  $(\alpha_0 + 1)$ -good. Then G is hyperbolic relative to  $\mathcal{S}(G)$ , and each non-cyclic subgroup in  $\mathcal{H}$  is contained in an element of  $\mathcal{S}(G)$ .

*Proof.* The proof of this lemma is quite long and intricate, so we subdivide it into five steps.

We begin with some basic observations. First, the projection  $\phi \in \text{Out}(G)$  has infinite order; otherwise it would be 1-good, as inner automorphisms have linear growth. Second, G does not virtually split as a direct product (and so G has trivial centre) because we are assuming that  $G \notin S(G)$ . Third, up to raising  $\varphi$  to a power, which does not affect the lack of goodness, we can assume that  $\varphi$  preserves each G-conjugacy class in  $\mathcal{H}$ .

**Step 1.** The growth rate  $\overline{\mathfrak{o}}_{top}(\phi)$  is either sub-polynomial or tame.

Let  $H_1, \ldots, H_k$  be representatives of the finitely many G-conjugacy classes of subgroups in  $\mathcal{H}$ . We can consider the growth rate given by the sum  $\mathfrak{s} := \sum_{j=1}^k \overline{\mathfrak{o}}_{top}(\phi|_{H_j})$  and note that it is  $\succeq \|\phi^n(h)\|$  for each element h of each subgroup  $H \in \mathcal{H}$ . This is true even when H is a cyclic element of  $\mathcal{H}$ , since the assumption that  $\mathcal{H}$  is  $\phi$ -invariant and consists of finitely many conjugacy classes of subgroups implies that  $\left[\|\phi^n(h)\|\right]$  is bounded for all  $h \in H$  in that case.

Now, since G is  $(\mathcal{Z}(G), \mathcal{H})$ -rigid, Lemma 5.7 shows that  $\overline{\mathfrak{o}}_{top}(\phi) \preceq \mathfrak{s}$ . The opposite inequality is also clear, so we obtain  $\overline{\mathfrak{o}}_{top}(\phi) \sim \mathfrak{s}$ . Finally, each rate  $\overline{\mathfrak{o}}_{top}(\phi|_{H_j})$  is either sub-polynomial or tame, by the hypothesis that the  $H_j$  are good, and a sum of such growth rates is itself either sub-polynomial or tame. In conclusion,  $\overline{\mathfrak{o}}_{top}(\phi)$  is either sub-polynomial or tame.

**Step 2.** Each automorphism  $\psi \in \operatorname{Aut}(G)$  in the outer class  $\phi$  falls into one of the following two cases. Case (i): we have  $\overline{\mathcal{O}}_{top}(\psi) \succ \overline{\mathfrak{o}}_{top}(\phi)$  and  $\overline{\mathfrak{o}}_{top}(\phi)$  is tame. Case (ii): we have  $\overline{\mathfrak{o}}_{top}(\phi) \preceq [n^p]$  for some  $p \in \mathbb{N}$  and  $\overline{\mathcal{O}}_{top}(\psi) \preceq [n^{p+\alpha_0}]$ .

By Step 1,  $\overline{\mathfrak{o}}_{top}(\phi)$  is either tame or sub-polynomial. If it is tame, we cannot have  $\overline{\mathcal{O}}_{top}(\psi) \sim \overline{\mathfrak{o}}_{top}(\phi)$  as this would imply that  $\varphi$  is docile (Remark 6.4); the latter would mean that  $\varphi$  is 0–good, against our hypotheses. If instead  $\overline{\mathfrak{o}}_{top}(\phi) \preceq [n^p]$  for some  $p \in \mathbb{N}$ , we cannot have  $\overline{\mathcal{O}}_{top}(\psi) \preceq [n^{p+\alpha_0}]$  as this

would imply that  $\overline{\mathcal{O}}_{top}(\varphi) \preceq [n^{p+\alpha_0+1}]$  by Remark 6.5, which would again violate the assumption that  $\varphi$  is not  $(\alpha_0 + 1)$ -good.

**Step 3.** For each  $\psi \in \operatorname{Aut}(G)$  in the outer class  $\phi$ , there exists a  $\psi$ -invariant centraliser  $G_0(\psi) \in \mathcal{Z}(G)$  that contains all  $\alpha_0$ -good convex-cocompact subgroups  $K \leq G$  such that  $\psi(K) = K$ .

Choose an ultrafilter  $\omega$  as follows. In Case (i) of Step 2, we have  $\overline{\mathcal{O}}_{top}(\psi) \succ \overline{\mathfrak{o}}_{top}(\phi)$  and we pick  $\omega$  with  $\mathcal{O}^{\omega}_{top}(\psi) \succ_{\omega} \mathfrak{o}^{\omega}_{top}(\phi)$ . In Case (ii), we have  $\overline{\mathcal{O}}_{top}(\psi) \not\preceq [n^{p+\alpha_0}]$  and we choose  $\omega$  so that  $\mathcal{O}^{\omega}_{top}(\psi) \succ_{\omega} [n^{p+\alpha_0}]$ . Then, we define:

$$G_0(\psi) := \{ g \in G \mid |\psi^n(g)| \prec_\omega \mathcal{O}^\omega_{\mathrm{top}}(\psi) \}.$$

Since  $|\psi^n(g_1g_2)| \leq_{\omega} \max_i |\psi^n(g_i)|$  for any  $g_1, g_2 \in G$ , we see that  $G_0(\psi)$  is a subgroup of G. Recalling that  $\psi$  is bi-Lipschitz with respect to  $|\cdot|$  (Lemma 2.15), we also see that  $G_0(\psi)$  is  $\psi$ -invariant.

By our choice of  $\omega$ , the group  $G_0(\psi)$  contains all  $\psi$ -invariant,  $\alpha_0$ -good, convex-cocompact subgroups  $K \leq G$ . Indeed, if  $\overline{\mathfrak{o}}_{top}([\psi|_K])$  is tame, then

$$\mathcal{O}_{\mathrm{top}}(\psi|_K) \sim \overline{\mathfrak{o}}_{\mathrm{top}}([\psi|_K]) \preceq \overline{\mathfrak{o}}_{\mathrm{top}}(\phi) \prec_{\omega} \mathcal{O}_{\mathrm{top}}^{\omega}(\psi).$$

If instead  $\overline{\mathfrak{o}}_{top}([\psi|_K])$  is sub-polynomial, then  $\overline{\mathcal{O}}_{top}(\psi|_K)$  is also sub-polynomial. Thus, if  $\overline{\mathfrak{o}}_{top}(\phi)$  is tame, we have again  $\overline{\mathcal{O}}_{top}(\psi|_K) \prec \overline{\mathfrak{o}}_{top}(\phi) \prec_{\omega} \mathcal{O}_{top}^{\omega}(\psi)$ . Finally, if  $\overline{\mathfrak{o}}_{top}(\phi) \preceq [n^p]$  for some integer p, we obtain  $\overline{\mathfrak{o}}_{top}([\psi|_K]) \preceq [n^p]$  and hence  $\overline{\mathcal{O}}_{top}(\psi|_K) \preceq [n^{p+\alpha_0}] \prec_{\omega} \mathcal{O}_{top}^{\omega}(\psi)$ .

In the rest of Step 3, we show that  $G_0(\psi)$  is a centraliser, which will take more work. Let  $\mathcal{X}_{\Gamma}$  be the universal cover of the Salvetti complex of  $A_{\Gamma}$ , and let  $C \subseteq \mathcal{X}_{\Gamma}$  be a convex subcomplex on which G acts essentially. Let  $G \curvearrowright \mathcal{X}_{\omega}$  be the degeneration determined by the ultrafilter  $\omega$  chosen above and the sequence  $\phi^n$ , as in Section 2.3. We use the sequence of automorphisms  $\psi^n$  as representatives of the  $\phi^n$  (this only affects how we represent points of  $\mathcal{X}_{\omega}$  by sequences of points of  $\mathcal{X}_{\Gamma}$ ). Recall that the scaling factors  $\tau_n$  used to define  $\mathcal{X}_{\omega}$  satisfy  $[\tau_n] \sim_{\omega} \mathfrak{o}_{top}^{\omega}(\phi)$ . Choose a sequence  $x_n \in C$  defining a point of  $\mathcal{X}_{\omega}$ .

Now, let  $\sigma_n$  be a sequence with  $[\sigma_n] \sim_{\omega} \mathcal{O}_{top}^{\omega}(\psi)$  and define  $G \curvearrowright \mathcal{Y}_{\omega}$  as the  $\omega$ -limit of the pointed metric spaces  $\frac{1}{\sigma_n}C$ , together with the  $\psi^n$ -twisted G-actions and again the  $x_n$  as basepoints. Since  $[\sigma_n] \succ_{\omega} [\tau_n]$  by assumption, while we have  $[d(\psi^n(g)x_n, x_n)] \sim_{\omega} [\tau_n]$  for all  $g \in G$ , the point  $(x_n) \in \mathcal{Y}_{\omega}$  is fixed by all elements of G. In fact, all points of  $\mathcal{X}_{\omega}$  get collapsed to a single point of  $\mathcal{Y}_{\omega}$ , which we will denote by  $p_{\mathcal{X}} \in \mathcal{Y}_{\omega}$ .

The action  $G \curvearrowright \mathcal{Y}_{\omega}$  also equivariantly embeds in a product of  $\mathbb{R}$ -trees  $G \curvearrowright \mathscr{T}_{\omega}^{v}$ , which satisfy all conclusions of Theorem 2.27. The proof of this is identical<sup>19</sup> to the case of usual degenerations, which unfortunately requires going through the arguments from [Fio23]. Since G is elliptic in the trees  $\mathscr{T}_{\omega}^{v}$ , there are no perverse lines and all G-stabilisers of arcs of  $\mathscr{T}_{\omega}^{v}$  lie in  $\mathcal{Z}(G)$ .

Finally we claim that, for any point  $x \in C$ , the constant sequence (x) defines a point of  $\mathcal{Y}_{\omega}$ . For this, note that G has trivial centre and therefore it is "uniformly non-elementary" as shown in [Fio24, Theorem I]. Straight from the definition of uniform non-elementarity in [Fio24], it follows that there exists a constant K such that, for any finite generating set  $S \subseteq G$ , we have:

$$d_{\mathcal{Y}_{\omega}}(p_{\mathcal{X}},(x)) = \lim_{\omega} \frac{1}{\sigma_n} d(x_n, x) \le \lim_{\omega} \frac{1}{\sigma_n} \cdot K \max_{s \in S} \left( d(\psi^n(s)x_n, x_n) + d(\psi^n(s)x, x) \right)$$
$$\lesssim \lim_{\omega} \frac{1}{\sigma_n} \cdot K(\tau_n + \sigma_n) < +\infty.$$

Hence  $(x) \in \mathcal{Y}_{\omega}$  as required.

<sup>&</sup>lt;sup>19</sup>In Theorem 2.27, we only claimed properties of the minimal subtrees of the  $T_{\omega}^{v}$ , but this is taken care of by the fact that, here, we took an ultralimit of copies of  $C \subseteq \mathcal{X}_{\Gamma}$  rather than of the whole  $\mathcal{X}_{\Gamma}$ . It is irrelevant that G is now elliptic in the  $\mathscr{T}_{\omega}^{v}$ , the same conclusions hold for the entire  $\mathscr{T}_{\omega}^{v}$ . The key point is [Fio23, Theorem 4.2].

We are now ready to conclude. The subgroup is  $G_0(\psi)$  is precisely the *G*-stabiliser of the point  $(x) \in \mathcal{Y}_{\omega}$ , straight from definitions. Since *G* fixes the point  $p_{\mathcal{X}} \in \mathcal{Y}_{\omega}$ , while  $G_0(\psi)$  is a proper subgroup of *G* by assuption, we have  $(x) \neq p_{\mathcal{X}}$ . Hence there exists  $v \in \Gamma$  such that (x) and  $p_{\mathcal{X}}$  have distinct projections to the  $\mathbb{R}$ -tree  $\mathscr{T}_{\omega}^v$ . In conclusion,  $G_0(\psi)$  is actually the *G*-stabiliser of the nontrivial arc spanned by the projections of (x) and  $p_{\mathcal{X}}$  to this tree  $\mathscr{T}_{\omega}^v$ , and so it is a centraliser.

**Step 4.** Each element of  $\mathcal{S}(G)$  contains all centralisers that it intersects nontrivially.

Consider some  $U \in \mathcal{S}(G)$ . As assumed at the beginning of the proof, the *G*-conjugacy class of *U* is  $\varphi$ -invariant, so there exists a representative  $\psi \in \operatorname{Aut}(G)$  of the outer class  $\phi$  with  $\psi(U) = U$ . Since  $U \in \mathcal{H}$ , our hypotheses imply that *U* is  $\alpha_0$ -good. By Step 3, the subgroup *U* is then contained in the centraliser  $G_0(\psi)$ . This implies that  $U = G_0(\psi)$ , as singular subgroups are maximal elements of  $\mathcal{VDP}(G)$ , and non-cyclic centralisers are contained in elements of  $\mathcal{VDP}(G)$ .

Now, consider the following family of subgroups of  $U = G_0(\psi)$ :

$$\mathscr{Z} := \{ U \cap Z_G(g) \mid g \in G \setminus U \}.$$

The collection  $\mathscr{Z}$  is  $\psi$ -invariant and it satisfies the following property.

**Claim.** Maximal elements of  $\mathscr{Z}$  are either U-parabolic or equal to U.

Proof of claim. Consider a maximal element  $M \in \mathscr{Z}$ , then choose an element  $g \in G \setminus U$  with  $M = U \cap Z_G(g)$ . Up to replacing g, we can assume that  $Z_G(g)$  is a maximal element of the family  $\{Z_G(x) \mid x \in G \setminus U\}$ ; this does not alter the intersection with U because M is maximal in  $\mathscr{Z}$ .

Now, note that  $Z_G(g) = G \cap Z_{A_{\Gamma}}(g)$ . Up to conjugating G by an element of  $A_{\Gamma}$ , we can assume that  $Z_{A_{\Gamma}}(g) = A_{\Delta} \times \langle a_1, \ldots, a_k \rangle$  for some  $k \geq 1$ , where the  $a_i$  are pairwise commuting elements of  $A_{\Gamma}$  such that the cyclic subgroups  $\langle a_i \rangle$  are convex-cocompact and  $g \in \langle a_1, \ldots, a_k \rangle$ . For each index i, there exists an integer  $k_i \geq 1$  such that  $a_i^{k_i} \in G$ , see [Fio24, Remark 3.17]. We have  $Z_{A_{\Gamma}}(g) \leq Z_{A_{\Gamma}}(a_i^{k_i})$  for each i and, since  $g \notin U$ , there exists an index j such that  $a_j^{k_j}$  lies outside U. By the maximality of  $Z_G(g)$ , we then have the equality  $Z_{A_{\Gamma}}(g) = Z_{A_{\Gamma}}(a_j^{k_j})$ . In conclusion, we can assume that k = 1.

Thus  $Z_{A_{\Gamma}}(g) = A_{\Delta} \times \langle \overline{g} \rangle$ , where  $\overline{g}$  is a root of g. The fact that U is convex-cocompact and does not contain g implies that  $M = U \cap A_{\Delta}$ , which is U-parabolic (or the entire M) as required.

We now conclude the proof of Step 4. Suppose for the sake of contradiction that there exists some  $g \in G \setminus U$  such that  $M := U \cap Z_G(g) \neq \{1\}$ . Without loss of generality, we can assume that M is a maximal element of  $\mathscr{Z}$ . By the claim,  $\mathscr{Z}$  contains only finitely many U-conjugacy classes of maximal subgroups (Lemma 2.2). Thus, up to raising  $\psi$  to a power, we can assume that the U-conjugacy class of M is  $\psi$ -invariant. Hence there exists an element  $u \in U$  such that the automorphism  $\psi'(x) := u\psi(x)u^{-1}$  leaves invariant M (as well as U).

The normaliser  $N_G(M)$  is also  $\psi'$ -invariant. It is *G*-parabolic by the claim and Lemma 2.5(2). By our hypotheses, the fact that  $N_G(M) \in \mathcal{P}(G)$  implies that it is  $\alpha_0$ -good. Thus, Step 3 shows that the centraliser  $G_0(\psi')$  contains  $N_G(M)$ , as well as *U*. Since  $U \in \mathcal{S}(G)$ , we again have  $U = G_0(\psi')$ , so this shows that  $N_G(M) \leq U$ . Since  $g \in Z_G(M) \leq N_G(M)$  by construction, this violates the fact that  $g \notin U$ , a contradiction.

In conclusion, we have shown that the collection  $\mathscr{Z}$  only consists of the trivial subgroup. That is, no element of  $U \setminus \{1\}$  commutes with an element of  $G \setminus U$ . Now, if U intersects nontrivially some centraliser  $Z \in \mathcal{Z}(G)$ , then we can pick an element  $g \in G$  such that  $Z \leq Z_G(g)$ . A first application of the previous observation implies that  $g \in U$ , since  $Z_G(g) \cap U \neq \{1\}$ , and a second shows that  $U \geq Z_G(g) \geq Z$ . Summing up, we have shown that U contains all elements of  $\mathcal{Z}(G)$ that it intersects nontrivially and, since U was an arbitrary element of  $\mathcal{S}(G)$ , this completes Step 4.

**Step 5.** The group G is hyperbolic relative to  $\mathcal{S}(G)$ .

We wish to prove relative hyperbolicity by appealing to Genevois' criterion [Gen21a, Theorem 1.3]. Elements of  $\mathcal{S}(G)$  are convex-cocompact (see Proposition 4.5(2)) and contain all non-cyclic abelian subgroups of G. Thus, we only need to show that the elements of  $\mathcal{S}(G)$  form a malnormal collection, i.e. that distinct elements of  $\mathcal{S}(G)$  have trivial intersection.

For this, recall that every singular subgroup  $U \in \mathcal{S}(G)$  virtually splits as a nontrivial direct product. Thus, consider a finite-index subgroup  $U^0 \leq U$  splitting as  $U_1 \times \ldots \times U_k \times A$ , where each  $U_i$  has trivial centre and A is abelian; the  $U_i$  and A can all be chosen to be convex-cocompact. If another element  $V \in \mathcal{S}(G)$  intersects U nontrivially, then it intersects  $U^0$  nontrivially. Since the intersection  $U^0 \cap V$  is convex-cocompact, it contains a nontrivial element u lying either in A or in one of the  $U_i$ . Step 4 then implies that  $Z_G(u) \leq U \cap V$ . If  $u \in A$ , we obtain  $U \leq Z_G(u) \leq V$ . If instead  $u \in U_i$  for some i, this shows that V contains all factors of  $U^0$  other than  $U_i$  and, repeating the same argument with u replaced by an element of V in a different factor of U, we again conclude that  $U \leq V$ . Finally, by maximality of singular subgroups, we obtain U = V and complete Step 5.

Summing up, G is relatively hyperbolic by Step 5. All non-cyclic elements of  $\mathcal{H}$  are contained in non-cyclic centralisers by Step 3 (since  $\mathcal{H}$  consists of finitely many conjugacy classes, each preserved by  $\varphi$ ), and hence they are contained in elements of  $\mathcal{S}(G)$ . This concludes the proof of the lemma.  $\Box$ 

Using Guirardel and Levitt's results on automorphisms of relative hyperbolic groups [GL15], we can now quickly deduce goodness from Lemma 6.11.

**Lemma 6.12.** Let G be 1-ended and hyperbolic relative to S(G). If  $\alpha_0 \ge 1$  is such that all subgroups in S(G) are  $\alpha_0$ -good, then G is  $(\alpha_0 + 2)$ -good.

*Proof.* By [GL15], there exist a finite-index subgroup  $\operatorname{Out}^1(G) \leq \operatorname{Out}(G)$  and an  $\operatorname{Out}^1(G)$ -invariant, minimal tree  $G \curvearrowright T$  with the following properties:

- (1) the subgroup  $\operatorname{Out}^1(G)$  preserves the *G*-conjugacy of each vertex group of *T*;
- (2) edge groups of T are infinite cyclic, or finitely generated subgroups of elements of  $\mathcal{S}(G)$ ;
- (3) each vertex group of T is either quadratically hanging, an element of  $\mathcal{S}(G)$ , or "rigid";
- (4) if V is a "rigid" vertex group, then  $\operatorname{Out}^1(G)$  restricts to a finite subgroup of  $\operatorname{Out}(V)$ .

(See in particular Sections 3.3 and 4.1 in [GL15].)

Now, consider any outer automorphism  $\phi \in \text{Out}(G)$ . Up to raising  $\phi$  to a power, which does not affect goodness, we can suppose that  $\phi$  lies in  $\text{Out}^1(G)$  and restricts to an inner automorphism on all rigid vertex groups of T. The restrictions of  $\phi$  to all QH vertex groups are 1–good, while the restrictions to the elements of  $\mathcal{S}(G)$  are  $\alpha_0$ –good by hypothesis. Finally, non-rigid vertex groups are conjugacy-undistorted (in the terminology of Appendix A.3) because they are either QH or convex-cocompact. In conclusion, we can appeal to Proposition A.8 in the appendix, which shows that  $\phi$  is  $(\alpha_0 + 2)$ –good as required.  $\Box$ 

We are finally ready to prove Proposition 6.10. The *Grushko rank* Gr(G) is the sum k + m in any writing  $G = G_1 * \cdots * G_k * F_m$  where the  $G_i$  are freely indecomposable.

Proof of Proposition 6.10. As mentioned, we proceed by induction on the ambient rank  $\operatorname{ar}(G)$ . The base step  $\operatorname{ar}(G) = 1$  is immediate. For the inductive step, suppose that all special groups H with  $\operatorname{ar}(H) < \operatorname{ar}(G)$  are good.

If  $G \in \mathcal{S}(G)$ , that is, if G virtually splits as a direct product, then G is good. Indeed, there exists  $\alpha \in \mathbb{N}$  such that the virtual direct factors of G are  $\alpha$ -good by the inductive assumption, and Corollary A.6 in the appendix shows that G is itself  $\alpha$ -good. Similarly, if G is freely decomposable and its indecomposable factors are  $\beta$ -good for some  $\beta \in \mathbb{N}$ , then G is  $(\beta + 2\operatorname{Gr}(G))$ -good, where  $\operatorname{Gr}(G)$  denotes the Grushko rank of G. This follows from Corollary A.12 and Proposition A.8 (the increase in the goodness parameter is due to sub-polynomially-growing automorphisms preserving a sporadic free factor). Note that the freely indecomposable factors of G are themselves special groups by Proposition 2.30, and their ambient rank does not exceed that of G.

In conclusion, it suffices to prove the proposition under the assumption that G be 1-ended and  $G \notin \mathcal{S}(G)$ . Denote by  $\mathcal{S}^*(G)$  the union of  $\mathcal{S}(G)$  with the family of cyclic subgroups of G whose conjugacy class has finite  $\operatorname{Out}(G)$ -orbit. Theorem 4.28 yields an  $\operatorname{Out}(G)$ -invariant  $(\mathcal{ZZ}(G), \mathcal{S}^*(G))$ -tree  $G \curvearrowright T$  such that each vertex-stabiliser V is either QH relative to  $\mathcal{S}^*(G)$ , or convex-cocompact and  $(\mathcal{Z}(V), \mathcal{S}^*(G)|_V)$ -rigid in itself. Free and surface groups are 1–good (see e.g. [BH92, Lev09]), so all QH vertex groups of T are 1–good.

Our goal is now to show that rigid vertex groups are also good for all practical purposes.

**Claim.** For every rigid vertex group V there exists  $\alpha_V \in \mathbb{N}$  such that, for every  $\varphi \in \operatorname{Aut}(G)$  with  $\varphi(V) = V$ , the restriction  $\varphi|_V$  is  $\alpha_V$ -good.

Proof of claim. It suffices to prove the claim under the assumption that  $\operatorname{ar}(V) = \operatorname{ar}(G)$ , otherwise it is an immediate consequence of the inductive hypothesis. Similarly, we can assume that  $V \notin \mathcal{S}(V)$ and that V is not cyclic. In particular, V is the G-stabiliser of a unique vertex  $v \in T$ .

We wish to apply Lemma 6.11 to V. Let  $\mathcal{H}_V$  be the union of the family  $\mathcal{S}(V)$  with the family of G-stabilisers of edges of T incident to v. There are only finitely many V-conjugacy classes of subgroups in  $\mathcal{H}_V$ , and all non-cyclic elements of  $\mathcal{H}_V$  are convex-cocompact subgroups with ambient rank strictly smaller than  $\operatorname{ar}(G) = \operatorname{ar}(V)$ . In particular, the inductive hypothesis implies that there exists  $\alpha_V \in \mathbb{N}$  such that all groups in  $\mathcal{H}_V \cup \mathcal{P}(V)$  are  $\alpha_V$ -good. Moreover, every subgroup in  $\mathcal{S}(G)^*|_V$  is contained in an element of  $\mathcal{H}_V$  (see the proof of Proposition 5.12 for details), and hence V is  $(\mathcal{Z}(V), \mathcal{H}_V)$ -rigid in itself. Finally, letting  $\Phi \in \operatorname{Aut}(T)$  be the map representing  $\varphi$ , the fact that  $\varphi(V) = V$  implies that  $\Phi(v) = v$ , and hence the collection  $\mathcal{H}_V$  is  $\varphi|_V$ -invariant.

Now, if the automorphism  $\varphi|_V \in \operatorname{Aut}(V)$  is not  $(\alpha_V + 1)$ -good, Lemma 6.11 shows that V is hyperbolic relative to  $\mathcal{S}(V)$  and that all elements of  $\mathcal{H}_V$  are contained in elements of  $\mathcal{S}(V)$ . Since G is 1-ended, V is 1-ended relative to  $\mathcal{H}_V$ , and the previous fact implies that V is 1-ended relative to  $\mathcal{S}(V)$ ; in fact, since all elements of  $\mathcal{S}(V)$  virtually split as direct products, they are freely indecomposable, so we conclude that V is actually 1-ended in the absolute sense. Finally, Lemma 6.12 shows that V is  $(\alpha_V + 2)$ -good, and hence  $\varphi|_V$  is  $(\alpha_V + 2)$ -good in all cases.

Since T has only finitely many G-orbits of vertices, the claim implies that there exists an integer  $\alpha \in \mathbb{N}$  such that all restrictions of automorphisms of G to the vertex groups of T are  $\alpha$ -good. Recalling that T is Out(G)-invariant and all its vertex groups are conjugacy-undistorted (as they are QH or convex-cocompact), we can finally appeal to Proposition A.8 and conclude that G is  $(\alpha + 2)$ -good, completing the proof.

6.4. **Super-singular growth rates.** In the previous subsection, we have shown that automorphisms of special groups either grow sub-polynomially or are docile (Definition 6.2). We can now use this to characterise all growth rates that are faster than the singular growth rate.

Let G be special and 1-ended. Consider  $\phi \in \text{Out}(G)$  and its singular growth rate  $\overline{\mathfrak{o}}_{\text{sing}}(\phi)$ (Definition 5.5). As in Section 5.2, we denote by  $\mathcal{K}_{\text{sing}}(\phi)$  the collection of subgroups of G all of whose elements satisfy  $\|\phi^n(k)\| \leq \overline{\mathfrak{o}}_{\text{sing}}(\phi)$ .

**Remark 6.13.** By Proposition 6.10, the growth rate  $\overline{\mathfrak{o}}_{top}(\phi)$  is always either sub-polynomial or tame. As sums of tame rates are tame, it follows that  $\overline{\mathfrak{o}}_{sing}(\phi)$  is also sub-polynomial or tame.

Recall that we constructed in Proposition 5.8 a JSJ tree adapted to  $\phi$ , which we denote by  $G \curvearrowright T_{\text{sing}}(\phi)$ . This is a  $\phi$ -invariant, minimal  $(\mathcal{ZZ}(G), \mathcal{K}_{\text{sing}}(\phi))$ -tree (possibly a single vertex) such that the G-stabiliser of each vertex of T is:

- (a) either an optimal quadratically hanging subgroup relative to  $\mathcal{K}_{sing}(\phi)$ ;
- (b) or a convex-cocompact root-closed element of  $\mathcal{K}_{sing}(\phi)$ .

Let  $k \geq 1$  be an integer such that  $\phi^k$  acts trivially on the quotient graph  $T_{\text{sing}}(\phi)/G$ . For every QH vertex group Q, we can consider the restriction  $\phi^k|_Q \in \text{Out}(G)$ . As  $\phi^k|_Q$  preserves the peripheral subgroups of the associated surface  $\Sigma_Q$ , it is represented by an element of the mapping class group of  $\Sigma_Q$ ; we can then consider the Nielsen–Thurston decomposition of this mapping class and its pseudo-Anosov components. We define  $\Lambda_{\text{sing}}(\phi^k) := \{\lambda_1, \ldots, \lambda_\ell\}$ , where the  $\lambda_i$  are the stretch factors of the pseudo-Anosov components of  $\phi^k|_Q$ , as Q varies through the QH vertex groups of T. Finally, we define

$$\Lambda_{\operatorname{sing}}(\phi) := \{\lambda_1^{1/k}, \dots, \lambda_\ell^{1/k}\},\,$$

and note that this set is independent of the choice of the integer k.

**Proposition 6.14.** Let G be special and 1-ended. For any  $\phi \in Out(G)$  and the tree  $G \curvearrowright T_{sing}(\phi)$ , all the following hold.

- (1) If  $\overline{\mathfrak{o}}_{sing}(\phi)$  is not sub-polynomial, then each element  $g \in G$  satisfies either  $\|\phi^n(g)\| \leq \overline{\mathfrak{o}}_{sing}(\phi)$ , or  $\|\phi^n(g)\| \sim \lambda^n$  for some  $\lambda \in \Lambda_{sing}(\phi)$ .
- (2) If  $\overline{\mathfrak{o}}_{sing}(\phi) \leq n^p$  for some  $p \in \mathbb{N}$ , then each element  $g \in G$  satisfies either  $\|\phi^n(g)\| \leq n^{p+2}$  or  $\|\phi^n(g)\| \sim \lambda^n$  for some  $\lambda \in \Lambda_{sing}(\phi)$ .
- (3) Either  $\overline{\mathfrak{o}}_{top}(\phi)$  is sub-polynomial, or  $\overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{sing}(\phi)$ , or  $\overline{\mathfrak{o}}_{top}(\phi) \sim \lambda^n$  for  $\lambda = \max \Lambda_{sing}(\phi)$ .

*Proof.* It suffices to prove parts (1) and (2) of the proposition, as part (3) is an immediate consequence of these (using Remark 2.21).

Throughout, we write  $T := T_{\text{sing}}(\phi)$  for simplicity and we suppose that  $\phi$  acts trivially on the quotient graph T/G, which can be achieved by raising  $\phi$  to a power. We also assume that, for each type (a) vertex group Q of T, the restriction  $\phi|_Q$  is represented by a pseudo-Anosov homeomorphism on the associated surface  $\Sigma_Q$  (fixing  $\partial \Sigma_Q$  pointwise). This can be achieved by refining  $T_{\text{sing}}(\phi)$ , splitting each type (a) vertex group Q according to the Nielsen–Thurston decomposition of  $\phi|_Q$ ; any resulting surface on which  $\phi$  restricts to a finite-order or linearly-growing homeomorphism should be considered of type (b).

Now, let  $\mathcal{E}$  be the set of edges of T both of whose vertices are of type (b). Let  $G \curvearrowright T'$  be the tree obtained from T by collapsing all edges in  $\mathcal{E}$ . Note that T' is still  $\phi$ -invariant and  $\phi$  still acts trivially on T'/G. We can still speak of type (a) and type (b) vertices of T', as each fibre of the collapse map  $\pi: T \to T'$  is either a single type (a) vertex, or a subtree of type (b) vertices.

Consider a type (b) vertex  $w \in T'$  and its stabiliser W. If  $\overline{\mathfrak{o}}_{sing}(\phi) \leq n^p$ , we have  $\overline{\mathfrak{o}}_{top}(\phi|_V) \leq n^p$ for every type (b) vertex group V of T and, applying Proposition A.8 to the tree  $W \curvearrowright \pi^{-1}(w)$ , we see that  $\overline{\mathfrak{o}}_{top}(\phi|_W) \leq n^{p+2}$ . If instead  $\overline{\mathfrak{o}}_{sing}(\phi)$  is not sub-polynomial, then it is tame (Remark 6.13) and applying<sup>20</sup> again Proposition A.8 to the tree  $W \curvearrowright \pi^{-1}(w)$ , we see that  $\overline{\mathfrak{o}}_{top}(\phi|_W) \leq \overline{\mathfrak{o}}_{sing}(\phi)$ . In other words,  $W \in \mathcal{K}_{sing}(\phi)$ , and W was already elliptic in T.

Summing up, if an element  $g \in G$  fixes a type (b) vertex of T', then we have  $\|\phi^n(g)\| \leq n^{p+2}$  if  $\overline{\mathfrak{o}}_{sing}(\phi) \leq n^p$ , and we instead have  $\|\phi^n(g)\| \leq \overline{\mathfrak{o}}_{sing}(\phi)$  if  $\overline{\mathfrak{o}}_{sing}(\phi)$  is not sub-polynomial.

In order to complete the proof of parts (1) and (2), we are left to describe the growth rate  $[\|\phi^n(g)\|]$  when g fixes a type (a) vertex or is loxodromic in T'. In the former case, Nielsen-Thurston theory shows that  $[\|\phi^n(g)\|]$  is either [1], [n], or  $[\lambda^n]$  for some  $\lambda \in \Lambda_{\text{sing}}(\phi)$ . Suppose instead that g is loxodromic and let  $\alpha \subseteq T'$  be its axis. By the construction of T', every edge of T' is incident to at least one type (a) vertex, and thus the axis  $\alpha$  contains a type (a) vertex q and two incident edges e, e'. Since q is an optimal QH vertex, e and e' correspond to distinct full peripheral subgroups of its stabiliser Q, and the pair (e, e') determines a homotopy class  $[\gamma]$  of properly embedded arcs on the associated surface  $\Sigma_q$ . As the the restriction of  $\phi$  to Q is represented

<sup>&</sup>lt;sup>20</sup>We emphasise that tameness is absolutely essential for this application of Proposition A.8, as it uses Remark 6.3 (though only if the quotient graph  $\pi^{-1}(w)/W$  is not simply connected). This seemingly inconsequential point is actually one of the main reasons why we had to go through the whole pain of Section 6.3.

by a pseudo-Anosov homeomorphism  $\psi_q$  of  $\Sigma_q$ , the length of the arcs  $\psi_q^n(\gamma)$  grows like  $\lambda_q^n$  for some  $\lambda_q \in \Lambda_{\text{sing}}(\phi)$ . From this, it is straightforward to deduce that  $\|\phi^n(g)\| \sim \lambda_q^n$  for the maximal  $\lambda_q \in \Lambda_{\text{sing}}(\phi)$  such that q is a type (a) vertex on the axis  $\alpha \subseteq T'$ . This concludes the proof.  $\Box$ 

As always, the tree  $T_{\text{sing}}(\phi)$  can be a single vertex, but this means that  $\overline{\mathfrak{o}}_{\text{top}}(\phi) \sim \overline{\mathfrak{o}}_{\text{sing}}(\phi)$ , thus reducing the study of  $\overline{\mathfrak{o}}_{\text{top}}(\phi)$  to special groups with strictly lower ambient rank than G.

**Corollary 6.15.** If G is special and 1-ended, there exists an integer  $\ell = \ell(G)$  such that the following holds. For each  $\phi \in \text{Out}(G)$ , there are at most  $\ell$  non-sub-polynomial growth rates  $\mathfrak{o} \in \mathfrak{g}(\phi)$  with  $\mathfrak{o} \not\preceq \overline{\mathfrak{o}}_{sing}(\phi)$ . For each such  $\mathfrak{o}$ , we have  $\mathfrak{o} \sim \lambda^n$  with  $\lambda$  the stretch factor of a pseudo-Anosov.

*Proof.* This immediately follows from Proposition 6.14, observing that the cardinality of the set  $\Lambda_{\text{sing}}(\phi)$  is bounded above in terms of the total complexity of the QH vertex groups appearing in  $T_{\text{sing}}(\phi)$ . All QH subgroups of G appear in any JSJ decomposition of G, in the sense of [GL17], and thus their total complexity is bounded independently of the automorphism  $\phi$ .

6.5. Completing the proof. We are now ready to prove Theorem 6.6, which we restate here for convenience. The additional part (4) will be useful in Section 7, where we study coarse-median preserving automorphisms and show that their growth rates are either sub-polynomial or pure.

**Theorem 6.16.** Let G be a special group. There exist natural numbers  $\pi = \pi(G)$ , d = d(G) and k = k(G) such that the following hold for every  $\varphi \in Aut(G)$  and its outer class  $\phi \in Out(G)$ .

- (1) Either  $\phi$  is docile, or  $\overline{\mathcal{O}}_{top}(\varphi) \preceq [n^{\pi}]$ .
- (2) If  $\phi$  is docile, there exists a d-algebraic integer  $\lambda > 1$  such that  $\overline{\mathfrak{o}}_{top}(\phi)$  is  $(\lambda, \pi)$ -tame.
- (3) If  $\phi$  is  $(\lambda, \pi)$ -docile, there is a  $\phi^k$ -invariant subgroup  $H \leq G$  of one of the following kinds.
  - (i) H is isomorphic to the fundamental group of a compact surface on which  $\phi^k$  is represented by a pseudo-Anosov homeomorphism with stretch factor  $\lambda^k$ . In particular, all non-peripheral elements  $h \in H$  satisfy  $\|\phi^n(h)\| \sim \lambda^n$ .
  - (ii) H is a convex-cocompact, infinitely-ended subgroup with a non-sporadic factor system  $\mathcal{F}$ such that  $\phi^k$  is a fully irreducible element of  $\operatorname{Out}(H, \mathcal{F})$  and  $\lambda^k$  is its Perron–Frobenius eigenvalue. Moreover, every element  $h \in H$  not conjugate into a subgroup in  $\mathcal{F}$  satisfies either  $\|\phi^n(h)\| \sim 1$  or  $\|\phi^n(h)\| \sim \lambda^n$ .
  - (iii) H is a convex-cocompact subgroup of the form  $H' \times \mathbb{Z}^m$  for some  $m \ge 1$ , and there exist elements  $h \in H$  with  $\|\phi^n(h)\| \sim n^p \lambda^n$  for some  $0 \le p \le \pi$ . Moreover, either H' is a surface group or infinitely ended, or  $\phi^k$  descends to an automorphism of the abelianisation of H having  $\lambda^k$  as the maximum modulus of an eigenvalue.
- (4) If  $\overline{\mathfrak{o}}_{top}(\phi)$  is pure (Definition 6.1), then there exists a  $\phi^k$ -invariant subgroup  $H \leq G$  containing elements h with  $\|\phi^n(h)\| \sim \overline{\mathfrak{o}}_{top}(\phi)$ , and H is of type (i), (iii), or:
  - (ii)' H is convex-cocompact, infinitely-ended and has a non-sporadic factor system  $\mathcal{F}$  such that  $\phi^k \in \operatorname{Out}(H, \mathcal{F})$  is fully irreducible with Perron–Frobenius eigenvalue  $\lambda^k$ . Each element  $h \in H$  not conjugate into a subgroup in  $\mathcal{F}$  satisfies either  $\|\phi^n(h)\| \sim 1$  or  $\|\phi^n(h)\| \sim \overline{\mathfrak{o}}_{top}(\phi)$ .

*Proof.* Proposition 6.10 shows that either  $\phi$  is docile, or all its representatives in Aut(G) grow subpolynomially. We prove the rest of the theorem by induction on the ambient rank  $\operatorname{ar}(G)$ . This amounts to bounding the integer  $\pi$  in parts (1) and (2) independently of the automorphism, to similarly bounding the degree of the algebraic integer  $\lambda$  in part (2), and to finding a subgroup H as in parts (3) and (4). The base step is trivial, so we assume that the theorem holds for all special groups G' with  $\operatorname{ar}(G') < \operatorname{ar}(G)$ .

Suppose first that G is 1-ended and that  $G \notin \mathcal{S}(G)$ . Then Proposition 6.14 guarantees that there are only three cases to consider:

- (x) either there exists  $p \in \mathbb{N}$  such that  $\overline{\mathfrak{o}}_{sing}(\phi) \preceq n^p$  and  $\overline{\mathfrak{o}}_{top}(\phi) \preceq n^{p+2}$ ;
- (y) or we have  $\overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{sing}(\phi)$  and this growth rate is tame;
- (z) or we have  $\overline{\mathfrak{o}}_{top}(\phi) \sim \lambda^n$  for some  $\lambda > 1$ .

In Case (x), the inductive assumption applied to the finitely many G-conjugacy classes of subgroups in  $\mathcal{S}(G)$  yields  $p \leq \pi'$  for some  $\pi' = \pi'(G)$ . Proposition 6.10 then yields an integer  $\alpha = \alpha(G)$  such that every representative  $\varphi \in \operatorname{Aut}(G)$  of  $\phi$  satisfies  $\overline{\mathcal{O}}_{\operatorname{top}}(\varphi) \preceq n^{p+2+\alpha}$ . Thus, part (1) of the theorem holds and parts (2)-(4) are void.

In Case (y), the theorem similarly holds by the inductive assumption. We only need to notice that, up to raising  $\phi$  to a bounded power, each conjugacy class in  $\mathcal{S}(G)$  is  $\phi$ -invariant, and thus  $\overline{\mathfrak{o}}_{sing}(\phi)$  is a sum of growth rates  $\overline{\mathfrak{o}}_{top}(\phi|_S)$  with S in a finite subset  $\mathcal{S}_0 \subseteq \mathcal{S}(G)$ . Since  $\overline{\mathfrak{o}}_{sing}(\phi)$ is  $(\lambda, p)$ -tame for some  $\lambda > 1$  and  $p \in \mathbb{N}$ , there exists  $S \in \mathcal{S}_0$  such that  $\overline{\mathfrak{o}}_{top}(\phi|_S)$  is  $(\lambda, p)$ -tame. Similarly, if  $\overline{\mathfrak{o}}_{top}(\phi|_S) \sim n^p \lambda^n$ , then there exists  $S \in \mathcal{S}_0$  such that  $\overline{\mathfrak{o}}_{top}(\phi|_S) \sim n^p \lambda^n$ . Applying the inductive hypothesis to this subgroup S, we obtain parts (3) and (4) of the theorem. Part (2) similarly follows from the inductive hypothesis and the equality  $\overline{\mathfrak{o}}_{top}(\phi) \sim \sum_S \overline{\mathfrak{o}}_{top}(\phi|_S)$ .

Finally, in Case (z) the growth rate  $\overline{\mathfrak{o}}_{top}(\phi) \sim \lambda^n$  is realised on a quadratically hanging vertex group Q of a  $\phi$ -invariant splitting of G. A power  $\phi^k$  preserves the conjugacy class of Q, and the proof of Proposition 6.14 shows that we can assume that the restriction  $\phi^k|_Q$  is represented by a pseudo-Anosov homeomorphism of the associated surface. The required integer k is at most q!, where q is the number of QH vertex groups in the Out(G)-invariant JSJ decomposition of G (over cyclic subgroups). As such, k is bounded purely in terms of G. This proves parts (3) and (4) of the theorem in this case. Part (2) is also clear, noting that the degree d of  $\lambda$  is bounded above in terms of the maximum complexity of a QH vertex group of the JSJ decomposition of G.

Summing up, under the inductive hypothesis, we have proved that the theorem holds when G is 1-ended and  $G \notin \mathcal{S}(G)$ .

Suppose now that  $G \in \mathcal{S}(G)$ , i.e. that G virtually splits as a direct product. It is not restrictive to assume that G itself splits as a direct product: given a subgroup  $L \leq G$  of index *i* and supposing that G is generated by r elements, there are at most  $(i!)^r$  subgroups of G of index *i*, and so each automorphism of G has a power of degree at most  $(i!)^r$  preserving L. Thus, we can suppose that  $G = G_1 \times \ldots \times G_a \times \mathbb{Z}^b$ , where the  $G_i$  are directly indecomposable and have trivial centre.

Consider an automorphism  $\varphi \in \operatorname{Aut}(G)$ . Up to raising  $\varphi$  to a power of degree  $\leq a!$ , each subgroup  $\langle G_j, \mathbb{Z}^b \rangle$  is left invariant by  $\varphi$  (Lemma A.5). Thus, we can represent  $\varphi$  as a (formal) triangular matrix as in Appendix A.1, with entries automorphisms  $(\varphi_1, \ldots, \varphi_a) \in \prod_j \operatorname{Aut}(G_j), \psi \in \operatorname{GL}_b(\mathbb{Z})$  and a homomorphism  $\alpha \colon \prod_j G_j \to \mathbb{Z}^b$ . Parts (1) and (2) of the theorem are immediate from Corollary A.6 and the inductive hypothesis. Parts (3) and (4) are similarly immediate if b = 0, in which case we can restrict to the  $G_j$ , which are all  $\varphi$ -invariant.

Thus, suppose that  $b \geq 1$ , so that the centre of G is infinite, and let  $\varphi$  be  $(\lambda, \pi)$ -docile. If none of the  $\varphi_j$  is  $(\lambda, \pi)$ -docile (that is, if they grow exponentially-slower than  $[\lambda^n]$ ), then Corollary A.6 shows that  $\overline{\mathfrak{o}}_{top}(\varphi)$  is realised on the abelianisation of G, and in particular it is pure; in this case, we can simply take H := G in parts (3) and (4) of the theorem. If instead there exists j such that  $\varphi_j$  is  $(\lambda, \pi)$ -docile, then it suffices to consider the subgroup  $H_j \leq G_j$  provided by the inductive hypothesis and set  $H := H_j \times \mathbb{Z}^b$  (when  $\overline{\mathfrak{o}}_{top}(\varphi)$  is pure, one should choose the index j so that  $\overline{\mathfrak{o}}_{top}([\varphi_j]) \sim \overline{\mathfrak{o}}_{top}(\varphi)$ ). This concludes the proof when  $G \in \mathcal{S}(G)$ .

In order to complete the proof of the theorem, we are only left to consider the possibility that G is not 1–ended. In this case, we argue by induction on the Grushko rank Gr(G).

Up to raising  $\phi \in \text{Out}(G)$  to a power (bounded by a function of Gr(G)), we can assume that there exists a factor system  $\mathcal{F}$  such that  $\phi$  is a fully irreducible element of  $\text{Out}(G, \mathcal{F})$  (see Lemma A.10). Each subgroup  $F \in \mathcal{F}$  has strictly lower Grushko rank than G, so the restriction  $\phi|_F$  satisfies the theorem. Moreover, the Perron–Frobenius eigenvalue of  $\phi$  has degree d bounded above in terms

of the Grushko rank Gr(G). Thus, parts (1) and (2) of the theorem immediately follow from Proposition A.11 (if  $\mathcal{F}$  is non-sporadic) and Proposition A.8 (if  $\mathcal{F}$  is sporadic, in which case G has a  $\phi$ -invariant splitting with vertex groups in  $\mathcal{F}$ ).

Regarding part (3), suppose that  $\phi$  is  $(\lambda, \pi)$ -docile. If there exists  $F \in \mathcal{F}$  such that the restriction  $\phi|_F$  is  $(\lambda, \pi)$ -docile, we obtain the required subgroup H by the inductive assumption. Otherwise, all restrictions  $\phi|_F$  are sub-polynomial or  $(\mu, \pi)$ -docile for values  $\mu < \lambda$ . In this case, Proposition A.11 (and its proof) show that all elements  $g \in G$  not conjugate into subgroups in  $\mathcal{F}$  satisfy either  $\|\phi^n(g)\| \sim 1$  or  $\|\phi^n(g)\| \sim \lambda^n$ , and the latter is the growth rate  $\bar{\mathfrak{o}}_{top}(\phi)$ , as required. Part (4) is similar: if  $\bar{\mathfrak{o}}_{top}(\phi)$  is pure and coincides with  $\bar{\mathfrak{o}}_{top}(\phi|_F)$  for some  $F \in \mathcal{F}$ , we conclude by the inductive assumption; if instead  $\bar{\mathfrak{o}}_{top}(\phi)$  is pure and strictly faster than all  $\bar{\mathfrak{o}}_{top}(\phi|_F)$ , then the proof of Proposition A.11 shows that all elements  $g \in G$  not conjugate into a subgroup in  $\mathcal{F}$  satisfy either  $\|\phi^n(g)\| \sim 1$  or  $\|\phi^n(g)\| \sim \bar{\mathfrak{o}}_{top}(\phi)$ , as required.

This finally completes the proof of the theorem.

**Remark 6.17.** It is interesting to pinpoint why we were not able to show that  $\overline{\mathfrak{o}}_{top}(\phi)$  is always pure, for a general automorphism  $\phi \in Out(G)$  growing at least exponentially.

If G is 1-ended, not a virtual direct product, and  $\overline{\mathfrak{o}}_{top}(\phi) \not\sim \overline{\mathfrak{o}}_{sing}(\phi)$ , then we have seen that  $\overline{\mathfrak{o}}_{top}(\phi) \sim \lambda^n$  for some  $\lambda > 1$ , and so  $\overline{\mathfrak{o}}_{top}(\phi)$  is pure in this case.

The problems start if we happen to have  $\overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{sing}(\phi)$ . We might then have  $\overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{top}(\phi|_S)$  for a singular subgroup  $S \in \mathcal{S}(G)$ , and there might be a direct factor S' of S such that  $\overline{\mathfrak{o}}_{top}(\phi|_S) \sim \overline{\mathfrak{o}}_{top}(\phi|_{S'})$  and such S' splits as a nontrivial free product  $S' = S_1 * \cdots * S_a * F_b$ . Now, even if the growth rates  $\overline{\mathfrak{o}}_{top}(\phi|_{S_i})$  are pure for all i, there is no guarantee that the growth rate  $\overline{\mathfrak{o}}_{top}(\phi|_{S'})$  is pure. Indeed, looking at the proof of Proposition A.11, we would need to know that there is at least a *linear* gap between  $\overline{\mathfrak{o}}_{top}(\phi|_{S_i})$  as the next-fastest growth rate on  $S_i$ , for each i.

This seems hard to show in general, but we will prove it in the coarse-median preserving case in Section 7, since we can control *all* growth rates in that case.

### 7. The coarse-median preserving case

Having shown that all outer automorphisms of special groups are either sub-polynomially growing or docile (Theorem 6.6), we now proceed to gather more refined information under the assumption that the outer automorphism be *coarse-median preserving*.

Examples of such automorphisms include all automorphisms of right-angled Coxeter groups, as well as *untwisted* automorphisms of right-angled Artin groups [Fio24, Proposition A]. All automorphisms of Gromov-hyperbolic groups are coarse-median preserving, while only a finite subgroup of  $Out(\mathbb{Z}^n)$  has this property. We refer to [FLS24, Theorem D] and [Fio23, Theorem E] for further examples of coarse-median preserving automorphisms of (virtually) special groups.

Consider a special group G, and fix a realisation of G as a convex-cocompact subgroup of a rightangled Artin group  $A_{\Gamma}$ . We equip G with the induced coarse-median operator  $m_{\Gamma}: G^3 \to G$ , and we denote by  $\operatorname{Aut}(G, m_{\Gamma}) \leq \operatorname{Aut}(G)$  and  $\operatorname{Out}(G, m_{\Gamma}) \leq \operatorname{Out}(G)$  the subgroups of automorphisms coarsely preserving  $m_{\Gamma}$ , as defined at the beginning of Section 2.1 (see [Fio24] for details). These subgroups depend on the choice of convex-cocompact embedding  $G \hookrightarrow A_{\Gamma}$ , in general.

The main result of this section is that coarse-median preserving automorphisms always have finitely many exponential growth rates, and each of them is of the form  $[n^p \lambda^n]$  for some  $p \in \mathbb{N}$  and some algebraic integer  $\lambda > 1$  (Remark 7.2). This proves Theorem B from the Introduction.

**Theorem 7.1.** Let  $G \leq A_{\Gamma}$  be convex-cocompact and let  $\phi \in \text{Out}(G, m_{\Gamma})$ .

- (1) Each growth rate in the set  $\mathfrak{g}(\phi)$  is either pure or sub-polynomial, and  $\overline{\mathfrak{o}}_{top}(\phi) \in \mathfrak{g}(\phi)$ .
- (2) There are only finitely many pure growth rates in  $\mathfrak{g}(\phi)$ . Moreover, there exists  $p \in \mathbb{N}$  such that all sub-polynomial growth rates in  $\mathfrak{g}(\phi)$  are  $\preceq [n^p]$ .

- (3) For every  $\varphi \in \operatorname{Aut}(G)$  in the outer class  $\phi$ , all growth rates in the symmetric difference  $\mathcal{G}(\varphi) \triangle \mathfrak{g}(\phi)$  are sub-polynomial.
- (4) For any pure growth rate  $\mathbf{o} \in \mathbf{g}(\phi)$ , there are only finitely many G-conjugacy classes of maximal subgroups in the family  $\mathcal{B}(\mathbf{o})$  (Definition 5.1). Each of these subgroups is convex-cocompact and its G-conjugacy class is preserved by a power of  $\phi$ .
- (5) For any pure growth rate  $\mathfrak{o} \in \mathfrak{g}(\phi)$ , there are an integer  $k \ge 1$  and a  $\phi^k$ -invariant subgroup  $H \le G$  of one of the following two kinds.
  - (i) H is isomorphic to the fundamental group of a compact surface on which  $\phi^k$  is represented by a pseudo-Anosov homeomorphism with stretch factor  $\lambda^k$ . In particular, all non-peripheral elements  $h \in H$  satisfy  $\|\phi^n(h)\| \sim \mathfrak{o} \sim \lambda^n$ .
  - (ii) *H* is a convex-cocompact, infinitely-ended subgroup with a non-sporadic factor system  $\mathcal{F}$ such that  $\phi^k$  is a fully irreducible element of  $\operatorname{Out}(H, \mathcal{F})$  and  $\lambda^k$  is its Perron–Frobenius eigenvalue. Each element  $h \in H$  not conjugate into a subgroup in  $\mathcal{F}$  satisfies either  $\|\phi^n(h)\| \sim 1$  or  $\|\phi^n(h)\| \sim \mathfrak{o} \sim n^p \lambda^n$  for some  $p \in \mathbb{N}$ .

The main content of Theorem 7.1 lies in its parts (1), (2) and (4). Part (5) is rather similar to the statement we have already proved for general automorphisms, the main difference being that we can now realise *all* pure growth rates on particular subgroups, rather than just the top one. Moreover, we can now realise  $\bar{\sigma}_{top}(\phi)$  exactly on a subgroup H, rather than only up to a polynomial error (as was the case in Theorem 6.6(3)), and we can disregard the situation when H has  $\mathbb{Z}^m$  as a direct factor, since coarse-median preserving automorphisms never skew the factors of a direct product (Remark 7.6).

**Remark 7.2.** Consider  $\phi \in \text{Out}(G, m_{\Gamma})$  and a pure growth rate  $\mathfrak{o} \in \mathfrak{g}(\phi)$ . Writing  $\mathfrak{o} \sim n^p \lambda^n$  for some  $p \in \mathbb{N}$  and  $\lambda > 1$ , the real number  $\lambda$  is always an algebraic integer, and a weak Perron number. This can be quickly deduced from Theorem 6.6 and Theorem 7.1 arguing as follows.

Letting  $\mathfrak{o}_1 \prec \cdots \prec \mathfrak{o}_m$  be the pure elements of  $\mathfrak{g}(\phi)$ , we have  $\mathfrak{o} \sim \mathfrak{o}_i$  for some index *i*. If i = m, then  $\mathfrak{o} \sim \overline{\mathfrak{o}}_{top}(\phi)$  and  $\lambda$  is a weak Perron number by Theorem 6.6(2). Otherwise, i < m and we can consider the family  $\mathcal{B}(\mathfrak{o}_{i+1})$ . By Theorem 7.1(4), each maximal subgroup  $B \in \mathcal{B}(\mathfrak{o}_{i+1})$  is convexcocompact and, up to raising  $\phi$  to a power, we have a restriction  $\phi|_B \in \text{Out}(B, m_{\Gamma})$ . Choosing the subgroup B so that it contains an element realising  $\mathfrak{o}_i$ , we obtain  $\mathfrak{o} \sim \overline{\mathfrak{o}}_{top}(\phi|_B)$ , and hence we can again appeal to Theorem 6.6(2) to conclude that  $\lambda$  is a weak Perron number.

**Remark 7.3.** Theorem 7.1 does not show that the entire set  $\mathfrak{g}(\phi)$  is finite, although this is likely to be true. Since part (2) bounds the exponent of sub-polynomial growth rates, the only missing element would be showing that sub-polynomial growth rates are *exactly* polynomial. In turn, this boils down to showing that, if G is special and  $G \curvearrowright T$  is a  $\phi$ -invariant ( $\mathcal{Z}(G), \mathcal{S}(G)$ )-splitting such that  $\overline{\mathfrak{o}}_{top}(\phi|_V) \preceq n^p$  on each vertex group V, then  $\overline{\mathfrak{o}}_{top}(\phi)$  is either  $\sim n^{p+1}$  or  $\sim n^{p+2}$ , rather than just  $\preceq n^{p+2}$  (confront Proposition 6.14(2)). Even in the very particular case when T is an HNN extension with trivial edge-stabilisers, this is a complicated problem, as it requires understanding the speed of growth of the elements  $a\varphi(a)\varphi^2(a)\ldots\varphi^n(a)$ , for a in a vertex group V and  $\varphi \in \operatorname{Aut}(V)$ representing  $\phi|_V$  (see the proof of Proposition A.8). This is delicate<sup>21</sup> because these products can have large amounts of cancellations.

**Remark 7.4.** We would expect the number of pure growth rates in  $\mathfrak{g}(\phi)$  to be bounded only in terms of the group G, independently of the specific outer automorphism  $\phi$ . However, this seems complicated to show, as some of these growth rates can arise from, say, quadratically hanging subgroups of the maximal subgroups in  $\mathcal{B}_{top}(\phi)$ . In general, these QH subgroups are not QH vertex

 $<sup>^{21}</sup>$ In the non-sporadic fully irreducible case, we overcame a similar issue with the trick in the claim in the proof of part (3) of Proposition A.11. However, this is of no help for an invariant HNN extension.

groups of a splitting of the whole G, and it also seems complicated (or downright impossible) to uniformly bound the "complexity" of the maximal elements of  $\mathcal{B}_{top}(\phi)$  in terms of that of G.

When G is 1-ended, Corollary 6.15 shows that there is only a uniformly bounded number of pure growth rates in  $\mathfrak{g}(\phi)$  that are faster than  $\overline{\mathfrak{o}}_{sing}(\phi)$ . Unfortunately, growth rates  $\mathfrak{o} \in \mathfrak{g}(\phi)$  with  $\mathfrak{o} \preceq \overline{\mathfrak{o}}_{sing}(\phi)$  need not be realised on singular subgroups of G, in general.

**Example 7.5.** Consider a right-angled Artin group  $A_{\Gamma}$  and an untwisted outer automorphism  $\phi \in \text{Out}(A_{\Gamma}, m_{\Gamma})$ . The maximal subgroups in  $\mathcal{B}_{\text{top}}(\phi)$  need not be isomorphic to right-angled Artin groups, and the following is likely the simplest example of this.

Let  $\Gamma$  be the following graph, and let  $\tau_i$  be the transvection mapping  $w_i \mapsto w_i v$  and fixing all other standard generators. Consider the composition  $\varphi := \tau_1 \tau_2 \tau_3$  and let  $\phi$  be its outer class.



The outer automorphism  $\phi$  grows linearly and there is a single conjugacy class of maximal subgroups of  $\mathcal{B}_{top}(\phi)$ , namely that of the fixed subgroup  $Fix(\varphi)$ . Moreover, we have

$$\operatorname{Fix}(\varphi) = \langle a, b, c, v, w_2 w_1^{-1}, w_3 w_1^{-1}, w_1 v w_1^{-1} \rangle.$$

We expect this subgroup to not be isomorphic to any right-angled Artin group, though proving this formally may prove challenging. (It is easy to see that the generating set we have given does not correspond to a right-angled Artin presentation of  $\operatorname{Fix}(\varphi)$ , as a does not commute with  $w_2w_1^{-1}$  or  $w_3w_1^{-1}$ , but it commutes with their product  $(w_2w_1^{-1})(w_3w_1^{-1})^{-1} = w_2w_3^{-1}$ .)

The proof of Theorem 7.1 is spread out over the next two subsections.

7.1. **Preliminaries on coarse-median preserving automorphisms.** Here we collect a few results on coarse-median preserving automorphisms that are needed in the proof of Theorem 7.1. For an expanded discussion, we refer the reader to [Fio24, Fio23, FLS24].

Let  $G \leq A_{\Gamma}$  be convex-cocompact, equipped with the induced coarse median  $m_{\Gamma}$ . If  $H \leq G$  is convex-cocompact and  $\varphi \in \operatorname{Aut}(G, m_{\Gamma})$ , then  $\varphi(H)$  is again convex-cocompact [Fio24, Corollary 3.3]. Because of convex-cocompactness, H inherits a coarse median structure from G and, if  $\varphi(H) = H$ , then the restriction  $\varphi|_{H}$  is again coarse-median preserving.

**Remark 7.6.** Coarse-median preserving automorphisms do not skew the factors of a direct product. More precisely, this means the following: Suppose that  $G = G_1 \times \ldots \times G_k \times A$ , where A is free abelian while the  $G_i$  are directly indecomposable and have trivial centre. By Lemma A.5, each automorphism  $\varphi \in \operatorname{Aut}(G)$  permutes the subgroups  $\langle G_i, A \rangle$ , and  $\varphi$  leaves invariant the centre A. If we have  $\varphi \in \operatorname{Aut}(G, m_{\Gamma})$ , then we can further conclude that  $\varphi$  permutes the subgroups  $G_i$  and that the restriction  $\varphi|_A \in \operatorname{Aut}(A)$  has finite order. (See e.g. [Fio23, Definition 2.23] and note that "orthogonality" is preserved by coarse-median preserving automorphisms.)

The following is the most important property of coarse-median automorphisms for this article. This is the single reason why we were able to obtain the stronger Theorem B in this case (in comparison to Theorem A). This all derives from the fact that degenerations of coarse-median preserving automorphisms do not have any perverse lines (Theorem 2.27), and thus point-stabilisers of the degeneration are convex-cocompact in G (using Corollary B.3 from the appendix).

Recall from Section 5.1 that  $\mathcal{B}_{top}(\phi)$  denotes the collection of subgroups of G none of whose elements realises  $\bar{\mathfrak{o}}_{top}(\phi)$ , and  $\mathcal{B}_{top}^{\omega}(\phi)$  similarly denotes the collection of subgroups on which  $\mathfrak{o}_{top}^{\omega}(\phi)$  is not realised, modulo an ultrafilter  $\omega$ .

**Lemma 7.7.** The following statements hold for every  $\phi \in \text{Out}(G, m_{\Gamma})$  and every ultrafilter  $\omega$ .

- (1) Each subgroup in  $\mathcal{B}^{\omega}_{top}(\phi)$  is contained in a maximal subgroup in  $\mathcal{B}^{\omega}_{top}(\phi)$ .
- (2) There are only finitely many G-conjugacy classes of maximal subgroups in  $\mathcal{B}^{\omega}_{\text{top}}(\phi)$ . Each of them is convex-cocompact, root-closed, and its conjugacy class is preserved by a power of  $\phi$ .
- (3) If B, B' are distinct maximal subgroups in  $\mathcal{B}^{\omega}_{top}(\phi)$ , then the intersection  $B \cap B'$  is contained in an element of  $\mathcal{Z}(G)$  (other than G itself, in case G happens to lie in  $\mathcal{Z}(G)$ ).

*Proof.* Throughout, we suppose that  $\phi \in \text{Out}(G, m_{\Gamma})$  has infinite order, otherwise  $\mathcal{B}^{\omega}_{\text{top}}(\phi) = \emptyset$  and the lemma is vacuously true.

Let  $G \curvearrowright \mathcal{X}_{\omega}$  be the degeneration determined by  $\phi$  and the ultrafilter  $\omega$ . Recall that  $\mathcal{X}_{\omega}$  equivariantly embeds in a finite product of  $\mathbb{R}$ -trees  $\prod_{v \in \Gamma} T_{\omega}^{v}$ , and a subgroup of G is elliptic in  $\mathcal{X}_{\omega}$  if and only if it is elliptic in all  $T_{\omega}^{v}$ . Since  $\phi$  is coarse-median preserving, Theorem 2.27 shows that all arc-stabilisers of the minimal subtrees  $\operatorname{Min}(G, T_{\omega}^{v})$  lie in  $\mathcal{Z}(G)$ . By Lemma 5.3, a subgroup of G lies in  $\mathcal{B}_{\operatorname{top}}^{\omega}(\phi)$  if and only if it fixes a point of  $\mathcal{X}_{\omega}$ .

Let  $\mathcal{I}$  be the collection of subgroups of G of the form  $\bigcap_{v \in \Gamma} P_v$ , where each  $P_v$  is the G-stabiliser of a point of  $\operatorname{Min}(G, T^v_{\omega})$  (excluding any  $v \in \Gamma$  for which G is elliptic in  $T^v_{\omega}$ ). The G-stabiliser of a point of  $\mathcal{X}_{\omega}$  fixes a point in each of the trees  $T^v_{\omega}$ , and each point-stabiliser of  $T^v_{\omega}$  fixes a point in  $\operatorname{Min}(G, T^v_{\omega})$ . Thus, the G-stabiliser of each point of  $\mathcal{X}_{\omega}$  is contained in a subgroup in  $\mathcal{I}$  and, conversely, each element of  $\mathcal{I}$  is elliptic in  $\mathcal{X}_{\omega}$ .

By Corollary B.3(1), stabilisers of points of the trees  $Min(G, T_{\omega}^v)$  are convex-cocompact and root-closed, and hence each subgroup in  $\mathcal{I}$  is convex-cocompact and root-closed. Moreover, Corollary B.3(2) shows that there are only finitely many conjugacy classes of stabilisers of points of  $Min(G, T_{\omega}^v)$ , so an iterated application of Lemma 2.4(5) shows that there are only finitely many G-conjugacy classes of subgroups in  $\mathcal{I}$ . Finally, using Lemma 2.4(3), we conclude that chains of subgroups in  $\mathcal{I}$  have bounded length.

This shows that every point-stabiliser of  $\mathcal{X}_{\omega}$  is contained in a maximal point-stabiliser and that the latter all lie in  $\mathcal{I}$ , and so they are convex-cocompact, root-closed and there are only finitely many conjugacy classes of them. The same then holds for maximal elements of  $\mathcal{B}_{top}^{\omega}(\phi)$ , as they are maximal point-stabilisers of  $\mathcal{X}_{\omega}$ . Finally, recalling that the family  $\mathcal{B}_{top}^{\omega}(\phi)$  is  $\phi$ -invariant by Remark 5.2, we conclude that each conjugacy class of subgroups in  $\mathcal{B}_{top}^{\omega}(\phi)$  is preserved by a power of  $\phi$ . This proves parts (1) and (2) of the lemma.

Regarding part (3), if  $B, B' \in \mathcal{B}_{top}^{\omega}(\phi)$  are distinct maximal elements, there exist distinct points  $p, p' \in \mathcal{X}_{\omega}$  such that  $B = G_p$  and  $B' = G_{p'}$ . Without loss of generality, the projections of p and p' to  $T_{\omega}^{v}$  lie within  $Min(G, T_{\omega}^{v})$  for all  $v \in \Gamma$  for which G is not elliptic in  $T_{\omega}^{v}$ . Since  $p \neq p'$ , there exists  $v \in \Gamma$  such that p and p' project to distinct points of  $Min(G, T_{\omega}^{v})$ , and hence  $B \cap B'$  fixes a nontrivial arc of  $Min(G, T_{\omega}^{v})$ . Thus,  $B \cap B'$  is contained in a centraliser.

Using the previous lemma, we can immediately prove parts (3) and (4) of Theorem 7.1, under the assumption that we have already shown parts (1) and (2):

**Lemma 7.8.** Let  $\phi \in \text{Out}(G, m_{\Gamma})$  be such that  $\mathfrak{g}(\phi)$  contains only finitely many pure growth rates, and such that all other elements of  $\mathfrak{g}(\phi)$  are sub-polynomial. Then we have the following.

- (1) If  $\overline{\mathfrak{o}}_{top}(\phi)$  is not sub-polynomial, then  $\overline{\mathfrak{o}}_{top}(\phi) \in \mathfrak{g}(\phi)$ .
- (2) For any pure growth rate  $\mathbf{o} \in \mathbf{g}(\phi)$ , there are only finitely many G-conjugacy classes of maximal subgroups in the family  $\mathcal{B}(\mathbf{o})$ . Each of these subgroups is convex-cocompact and its G-conjugacy class is preserved by a power of  $\phi$ .
- (3) For any  $\varphi \in \operatorname{Aut}(G, m_{\Gamma})$  in the outer class  $\phi$ , all growth rates in the symmetric difference  $\mathcal{G}(\varphi) \triangle \mathfrak{g}(\phi)$  are sub-polynomial.
*Proof.* If  $\mathfrak{g}(\phi)$  contains at least one pure growth rate, then it has a  $\leq$ -maximum  $\mathfrak{o}_{\max}$  by our hypotheses. Recalling Lemma 2.20, we have  $\overline{\mathfrak{o}}_{top}(\phi) \sim_{\omega} \mathfrak{o}_{\max}$  for every ultrafilter  $\omega$ , and this implies that  $\overline{\mathfrak{o}}_{top}(\phi) \sim \mathfrak{o}_{\max} \in \mathfrak{g}(\phi)$  (Remark 2.14), proving part (1).

We now prove part (2). Since all elements of  $\mathfrak{g}(\phi)$  are either pure or sub-polynomial, we have  $\mathcal{B}(\mathfrak{o}) = \mathcal{B}^{\omega}(\mathfrak{o})$  for any pure growth rate  $\mathfrak{o} \in \mathfrak{G}$  and any ultrafilter  $\omega$ . In particular  $\mathcal{B}_{top}(\phi) = \mathcal{B}^{\omega}_{top}(\phi)$ , so Lemma 7.7(2) implies that there are only finitely many conjugacy classes of maximal subgroups in  $\mathcal{B}_{top}(\mathfrak{o})$ , that all of these maximal subgroups are convex-cocompact and root-closed, and that each has conjugacy class preserved by a power of  $\phi$ . Thus, part (2) of the lemma holds for  $\mathfrak{o} = \overline{\mathfrak{o}}_{top}(\phi)$ . For a general pure growth rate  $\mathfrak{o} \in \mathfrak{g}(\phi)$ , each maximal subgroup in  $\mathcal{B}(\mathfrak{o})$  is contained in a maximal subgroup in  $\mathcal{B}_{top}(\phi)$ , so it suffices to restrict  $\phi$  to the latter (after raising  $\phi$  to a power) and repeat the previous argument. This procedure eventually terminates, by the assumption that  $\mathfrak{g}(\phi)$  contains only finitely many pure growth rates, proving part (2).

Finally, we prove part (3). We can suppose that  $\overline{\mathfrak{o}}_{top}(\phi)$  is not sub-polynomial, otherwise Theorem 6.6 implies that all elements of  $\mathcal{G}(\varphi) \cup \mathfrak{g}(\phi)$  are sub-polynomial. Now, Theorem 6.6 also implies that we have  $\overline{\mathcal{O}}_{top}(\varphi) \sim \overline{\mathfrak{o}}_{top}(\phi)$ , and hence  $|\varphi^n(g)| \leq \overline{\mathfrak{o}}_{top}(\phi)$  for all  $g \in G$ . The following characterises the elements for which this inequality is strict.

**Claim.** If  $g_* \in G$  is an element such that  $|\varphi^n(g_*)| \not\sim \overline{\mathfrak{o}}_{top}(\phi)$ , then there exists a convex-cocompact subgroup  $H \leq G$  such that  $\varphi(H) = H$ ,  $g_* \in H$ , and  $H \in \mathcal{B}_{top}(\phi)$ .

Proof of claim. Since  $|\varphi^n(g_*)| \leq \overline{\mathfrak{o}}_{top}(\phi)$  and  $|\varphi^n(g_*)| \not\sim \overline{\mathfrak{o}}_{top}(\phi)$ , we can fix an ultrafilter  $\omega$  such that  $g_*$  lies in the following subset of G:

$$\Omega := \{ g \in G \mid |\varphi^n(g)| \prec_\omega \mathfrak{o}^\omega_{\mathrm{top}}(\phi) \}.$$

Note that  $\Omega$  is a subgroup, as we have  $|\varphi^n(gh)| \leq |\varphi^n(g)| + |\varphi^n(h)|$  for any  $g, h \in G$ . Moreover, we have  $\varphi(\Omega) = \Omega$ , as  $\varphi$  is bi-Lipschitz with respect to  $|\cdot|$ . Finally, note that, for each  $h \in \Omega$ , we have  $\|\phi^n(h)\| \leq |\varphi^n(h)| \prec_{\omega} \mathfrak{o}_{top}^{\omega}(\phi)$ , and hence  $\Omega \in \mathcal{B}_{top}^{\omega}(\phi) = \mathcal{B}_{top}(\phi)$ .

Let Z be the intersection of all elements of  $\mathcal{Z}(G)$  containing  $\Omega$ , or set Z := G if no such element exists. Note that  $\varphi(Z) = Z$  and that Z is either a centraliser or the whole G, so Z is convexcocompact. If  $\overline{\mathfrak{o}}_{top}(\phi|_Z) \not\sim \overline{\mathfrak{o}}_{top}(\phi)$ , then  $Z \in \mathcal{B}_{top}(\phi)$  and the claim is proved.

Otherwise, we have  $\Omega \in \mathcal{B}_{top}(\phi|_Z)$ . Let  $B \leq Z$  be a maximal element of  $\mathcal{B}_{top}(\phi|_Z)$  containing  $\Omega$ , which exists by Lemma 7.7(1). Since the family  $\mathcal{B}_{top}(\phi|_Z)$  is  $\varphi$ -invariant, the subgroup  $\varphi(B)$  is also a maximal element of  $\mathcal{B}_{top}(\phi|_Z)$ . If we had  $\varphi(B) \neq B$ , then Lemma 7.7(3) would imply that  $\varphi(B) \cap B$  is contained in a centraliser  $Z' \in \mathcal{Z}(Z) \subseteq \mathcal{Z}(G)$  with  $Z' \neq Z$ , and we would have  $\Omega \leq Z \cap Z' \leq Z$ , contradicting our choice of Z. In conclusion, we have  $\varphi(B) = B$ ,  $g_* \in \Omega \leq B$  and  $B \in \mathcal{B}_{top}(\phi|_Z) \subseteq \mathcal{B}_{top}(\phi)$ . Finally, B is convex-cocompact by Lemma 7.7(2), proving the claim.

Now, let  $\mathfrak{o}_1 \prec \cdots \prec \mathfrak{o}_m$  be the list of pure growth rates in  $\mathfrak{g}(\phi)$ , which is finite and contains all non-sub-polynomial elements of  $\mathfrak{g}(\phi)$  by hypothesis. If  $|\varphi^n(g_*)| \not\sim \overline{\mathfrak{o}}_{top}(\phi) = \mathfrak{o}_m$ , the claim (together with Theorem 6.6) implies that  $|\varphi^n(g_*)| \preceq \mathfrak{o}_{m-1}$ . A repeated application of this argument then yields that, for each  $g \in G$ , either the growth rate  $[|\varphi^n(g)|]$  is sub-polynomial or  $|\varphi^n(g)| \sim \mathfrak{o}_i$  for some index  $1 \leq i \leq m$ .

In conclusion, we have show that each element of the difference  $\mathcal{G}(\varphi) \setminus \mathfrak{g}(\phi)$  is sub-polynomial. Conversely, the fact that the elements of  $\mathfrak{g}(\phi) \setminus \mathcal{G}(\varphi)$  are sub-polynomial is straightforward: for each pure growth rate  $\mathfrak{o} \in \mathfrak{g}(\phi)$ , we can pick a maximal element  $B \in \mathcal{B}(\mathfrak{o})$ , which is convex-cocompact and preserved by a power of  $\phi$  by part (2), so we can apply Theorem 6.6 to the restriction to B of a power of  $\phi$  to conclude that  $\mathfrak{o} \in \mathcal{G}(\varphi)$ . This proves part (3), concluding the proof of the lemma.  $\Box$ 

7.2. The core of the proof. This subsection is devoted to the proof of Theorem 7.1, which mainly amounts to the following proposition. We have already seen in Lemma 7.8 how parts (3) and (4) of the theorem can then be deduced from this.

**Proposition 7.9.** Let  $G \leq A_{\Gamma}$  be convex-cocompact. For every  $\phi \in Out(G, m_{\Gamma})$ :

- (1) the set  $\mathfrak{g}(\phi)$  contains only finitely many non-sub-polynomial growth rates;
- (2) each growth rate in  $\mathfrak{g}(\phi)$  is either pure or sub-polynomial.

Proof. We prove the proposition by a double induction. The first induction is on the number of vertices of the finite graph  $\Gamma$ . Thus, suppose that the proposition holds for all coarse-median preserving automorphisms of all convex-cocompact subgroups of (proper) parabolic subgroups of  $A_{\Gamma}$ . We can also suppose that  $G \notin S(G)$ , otherwise the proposition holds for G by a straightforward application of Corollary A.6. Up to raising  $\phi$  to a power, we can further assume that each conjugacy class of subgroups in S(G) is  $\phi$ -invariant. We then define the set of growth rates

$$\mathfrak{g}_{\operatorname{sing}}(\phi) := \bigcup_{S \in \mathcal{S}(G)} \mathfrak{g}_{\operatorname{pure}}(\phi|_S),$$

where  $\mathfrak{g}_{\text{pure}}(\phi|_S) \subseteq \mathfrak{g}(\phi|_S)$  denotes the subset of pure growth rates. Note that  $\mathfrak{g}_{\text{sing}}(\phi) \subseteq \mathfrak{G}$  is finite, since each set  $\mathfrak{g}_{\text{pure}}(\phi|_S)$  is finite by the (first) inductive assumption, and there are only finitely many *G*-conjugacy classes of singular subgroups.

The second level of the induction is on the cardinality  $\#\mathfrak{g}_{sing}(\phi)$ . We have  $\#\mathfrak{g}_{sing}(\phi) = 1$  precisely when  $\phi$  grows sub-polynomially on all singular subgroups of G. We will treat this base step simultaneously with the inductive step, as the argument is the same. Thus, suppose that the proposition also holds for all coarse-median preserving outer automorphisms  $\psi$  of convex-cocompact subgroups of  $A_{\Gamma}$  such that  $\#\mathfrak{g}_{sing}(\psi) < \#\mathfrak{g}_{sing}(\phi)$ .

We now proceed by distinguishing three cases. Throughout, we assume that  $\overline{\mathfrak{o}}_{top}(\phi)$  is not subpolynomial, otherwise all elements of  $\mathfrak{g}(\phi)$  are sub-polynomial and the proposition is trivially true. **Case 1.** We have  $\overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{sing}(\phi)$ .

In the base step, this means that  $\overline{\mathfrak{o}}_{top}(\phi)$  is sub-polynomial, which we have already ruled out. For the inductive step, we argue as follows.

First, we claim that  $\overline{\mathfrak{o}}_{top}(\phi)$  is pure and that  $\overline{\mathfrak{o}}_{top}(\phi) \in \mathfrak{g}_{sing}(\phi)$ . Indeed, each growth rate  $\overline{\mathfrak{o}}_{top}(\phi|_S)$  with  $S \in \mathcal{S}(G)$  is either pure or sub-polynomial by the (first) inductive hypothesis, recalling that  $\overline{\mathfrak{o}}_{top}(\phi|_S) \in \mathfrak{g}(\phi|_S)$  by Lemma 7.8(1). Thus,  $\overline{\mathfrak{o}}_{sing}(\phi)$  is either pure or sub-polynomial, as it is defined as a finite sum of such growth rates (Definition 5.5). Since  $\overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{sing}(\phi)$  by the hypothesis of Case 1, we conclude that this growth rate is pure and that it must thus coincide with  $\overline{\mathfrak{o}}_{top}(\phi|_S)$  for some  $S \in \mathcal{S}(G)$ . Hence  $\overline{\mathfrak{o}}_{top}(\phi) \in \mathfrak{g}_{sing}(\phi)$  as claimed.

Next, we claim that all growth rates in  $\mathfrak{g}(\phi)$  are either pure or sub-polynomial. For this, consider an element  $g \in G$  and recall that  $\|\phi^n(g)\| \leq \overline{\mathfrak{o}}_{top}(\phi)$ . If  $\|\phi^n(g)\| \sim \overline{\mathfrak{o}}_{top}(\phi)$ , we are done. Otherwise, there exists an ultrafilter  $\omega$  such that  $\|\phi^n(g)\| \prec_\omega \mathfrak{o}_{top}^\omega(\phi)$ , and hence we have  $g \in B$  for a maximal element  $B \in \mathcal{B}_{top}^\omega(\phi)$  by Lemma 7.7(1). The subgroup B is convex-cocompact and, up to raising  $\phi$ to a power, the conjugacy class of B is  $\phi$ -invariant by Lemma 7.7(2). Since every singular subgroup of B is contained in a singular subgroup of G, we have  $\mathfrak{g}_{sing}(\phi|_B) \subseteq \mathfrak{g}_{sing}(\phi)$ . Since  $B \in \mathcal{B}_{top}^\omega(\phi)$ , we have  $\overline{\mathfrak{o}}_{top}(\phi) \notin \mathfrak{g}_{sing}(\phi|_B)$ . At the same time, we have seen above  $\overline{\mathfrak{o}}_{top}(\phi) \in \mathfrak{g}_{sing}(\phi)$ , so we conclude that  $\#\mathfrak{g}_{sing}(\phi|_B) < \#\mathfrak{g}_{sing}(\phi)$  and the (second) inductive hypothesis implies that all elements of  $\mathfrak{g}(\phi|_B)$  are either pure or sub-polynomial. Since  $g \in B$ , this proves that the growth rate  $\left[\|\phi^n(g)\|\right]$ is pure or sub-polynomial, as claimed.

Finally, we can show that  $\mathfrak{g}(\phi)$  contains only finitely many pure growth rates. By the previous claim, we have  $\mathcal{B}_{top}^{\omega}(\phi) = \mathcal{B}_{top}(\phi)$  for all ultrafilters  $\omega$ . Thus, Lemma 7.7(2) shows that  $\mathcal{B}_{top}(\phi)$ consists of finitely many *G*-conjugacy classes of convex-cocompact subgroups, each preserved by  $\phi$  (after raising  $\phi$  to some power). Now, each  $\mathfrak{o} \in \mathfrak{g}(\phi)$  satisfies either  $\mathfrak{o} \sim \overline{\mathfrak{o}}_{top}(\phi)$  or  $\mathfrak{o} \in \mathfrak{g}(\phi|_B)$ for some  $B \in \mathcal{B}_{top}(\phi)$ , and each set  $\mathfrak{g}(\phi|_B)$  contains only finitely many pure growth rates by the (second) inductive assumption, as we have seen that  $\#\mathfrak{g}_{sing}(\phi|_B) < \#\mathfrak{g}_{sing}(\phi)$ .

This proves the proposition in Case 1.

**Case 2.** We have  $\overline{\mathfrak{o}}_{sing}(\phi) \prec \overline{\mathfrak{o}}_{top}(\phi)$  and G is 1-ended.

In the base step when  $\overline{\mathfrak{o}}_{sing}(\phi)$  is sub-polynomial, Proposition 6.14(2) shows that  $\mathfrak{g}(\phi)$  contains only finitely many non-sub-polynomial growth rates and that these are all pure.

In general, recall that  $\mathcal{K}_{sing}(\phi)$  is the family of subgroups of G all of whose elements satisfy  $\|\phi^n(k)\| \leq \overline{\mathfrak{o}}_{sing}(\phi)$ . By Corollary 5.9, every subgroup in  $\mathcal{K}_{sing}(\phi)$  is contained in a maximal such subgroup, the latter are all convex-cocompact, and there are only finitely many conjugacy classes of them. Up to raising  $\phi$  to a power, we can assume that each conjugacy class of maximal elements of  $\mathcal{K}_{sing}(\phi)$  is  $\phi$ -invariant. For each maximal element  $K \in \mathcal{K}_{sing}(\phi)$ , we then have

(7.1) 
$$\overline{\mathfrak{o}}_{\operatorname{sing}}(\phi|_K) \preceq \overline{\mathfrak{o}}_{\operatorname{top}}(\phi|_K) \preceq \overline{\mathfrak{o}}_{\operatorname{sing}}(\phi).$$

Recall that  $\overline{\mathfrak{o}}_{sing}(\phi)$  is either sub-polynomial or pure, by the (first) inductive assumption. If  $\overline{\mathfrak{o}}_{sing}(\phi)$  is pure, it is realised on some  $S \in \mathcal{S}(G)$  and so it lies in  $\mathfrak{g}_{sing}(\phi)$ . In conclusion, if  $\overline{\mathfrak{o}}_{sing}(\phi|_K)$  is not sub-polynomial and if we do not have  $\#\mathfrak{g}_{sing}(\phi|_K) < \#\mathfrak{g}_{sing}(\phi)$ , then  $\overline{\mathfrak{o}}_{sing}(\phi) \in \mathfrak{g}_{sing}(\phi|_K)$  and Equation (7.1) shows that  $\overline{\mathfrak{o}}_{sing}(\phi|_K) \sim \overline{\mathfrak{o}}_{top}(\phi|_K)$ . Now, either by the (second) inductive hypothesis or by Case 1, we know that  $\phi|_K$  satisfies the proposition for every maximal element  $K \in \mathcal{K}_{sing}(\phi)$ .

Corollary 6.15 shows that, with the exception of finitely many pure growth rates, each  $\mathfrak{o} \in \mathfrak{g}(\phi)$  that is not sub-polynomial satisfies  $\mathfrak{o} \preceq \overline{\mathfrak{o}}_{sing}(\phi)$ . In turn, if  $\mathfrak{o} \preceq \overline{\mathfrak{o}}_{sing}(\phi)$ , then  $\mathfrak{o}$  is realised on some maximal element  $K \in \mathcal{K}_{sing}(\phi)$  and we know that the proposition holds for  $\phi|_K$ . It follows that the proposition holds for  $\phi$  as well, and this concludes Case 2.

## **Case 3.** The group G is freely decomposable.

Let  $\mathcal{I}$  be the collection of freely indecomposable free factors of G. Up to raising  $\phi$  to a power, we can assume that each conjugacy class in  $\mathcal{I}$  is  $\phi$ -invariant. For each  $I \in \mathcal{I}$ , every singular subgroup of I is contained in a singular subgroup of G, and thus we have  $\mathfrak{g}_{\text{sing}}(\phi|_I) \subseteq \mathfrak{g}_{\text{sing}}(\phi)$ . By Cases 1 and 2 above, it follows that the restriction  $\phi|_I$  satisfies the proposition for each  $I \in \mathcal{I}$ . Letting  $\mathfrak{g}_{\text{ind}}(\phi) \subseteq \mathfrak{g}(\phi)$  be the set of all pure growth rates of the restrictions  $\phi|_I$  as I varies in  $\mathcal{I}$ , it follows that the set  $\mathfrak{g}_{\text{ind}}(\phi)$  is finite.

Let  $\Lambda \subseteq \mathbb{R}_{>1}$  be a finite set and  $P \in \mathbb{N}$  an integer such that every growth rate in  $\mathfrak{g}_{ind}(\phi)$  is of the form  $[n^a \nu^n]$  for some  $\nu \in \Lambda$  and some integer  $0 \leq a \leq P$ . Recall that Gr(G) denotes the Grushko rank of G. We complete Case 3 by proving the following.

**Claim.** Each growth rate in the set  $\mathfrak{g}(\phi)$  is either sub-polynomial or of the form  $[n^a\nu^n]$ , for some integer  $0 \leq a \leq P + \operatorname{Gr}(G)$  and some  $\nu \in \Lambda'$ , for a set  $\Lambda \subseteq \Lambda' \subseteq \mathbb{R}_{>1}$  with  $\#(\Lambda' \setminus \Lambda) \leq \operatorname{Gr}(G)$ .

Proof of claim. We prove the claim via a third inductive procedure, on the value of the pair  $(\#\mathfrak{g}_{ind}(\phi), \operatorname{Gr}(G))$ , ordered lexicographically so that the first entry takes precedence over the Grushko rank  $\operatorname{Gr}(G)$ . The base step is trivial, so we only consider the inductive step.

Up to raising  $\phi$  to a power, Lemma A.10 allows us to find a (possibly sporadic) factor system  $\mathcal{F}$  for G such that that  $\phi$  is a fully irreducible element of  $\operatorname{Out}(G, \mathcal{F})$ . For each  $F \in \mathcal{F}$ , we have  $\mathfrak{g}_{\operatorname{ind}}(\phi|_F) \subseteq \mathfrak{g}_{\operatorname{ind}}(\phi)$  and  $\operatorname{Gr}(F) < \operatorname{Gr}(G)$ , and thus the claim holds for the restriction  $\phi|_F$  by the (third) inductive assumption. Let  $\Lambda_0 \subseteq \mathbb{R}_{>1}$  and  $P_0 \in \mathbb{N}$  be such that each growth rate in the union  $\bigcup_{F \in \mathcal{F}} \mathfrak{g}(\phi|_F)$  is either sub-polynomial or of the form  $[n^a \nu^n]$  with  $\nu \in \Lambda_0$  and  $0 \le a \le P_0$ .

If  $\mathcal{F}$  is non-sporadic, Proposition A.11(3) shows that each growth rate in  $\mathfrak{g}(\phi)$  is either subpolynomial, or of the form  $[n^a\nu^n]$  with  $\nu \in \Lambda_0$  and  $0 \leq a \leq P_0 + 1$ , or of the form  $[\lambda^n]$  for a single new number  $\lambda > 1$ . Note that the hypotheses of Proposition A.11(3) are indeed satisfied: if  $F \in \mathcal{F}$ and  $\varphi \in \operatorname{Aut}(F)$  is a representative of  $\phi|_F$ , then every growth rate in  $\mathcal{G}(\varphi)$  is either sub-polynomial or a pure element of  $\mathfrak{g}(\phi)$ ; this follows from Lemma 7.8(3), using the (third) inductive assumption for  $\phi|_F$ . In conclusion, this proves the claim when  $\mathcal{F}$  is non-sporadic.

Suppose instead that  $\mathcal{F}$  is sporadic, so that there exists a  $\phi$ -invariant free splitting  $G \curvearrowright T$ whose non-trivial vertex groups are the elements of  $\mathcal{F}$ . By Proposition A.8, the growth rate  $\overline{\mathfrak{o}}_{top}(\phi)$ equals the fastest pure growth rate in the union  $\bigcup_{F \in \mathcal{F}} \mathfrak{g}(\phi|_F)$ . Each element of  $\mathfrak{g}(\phi)$  other than  $\overline{\mathfrak{o}}_{top}(\phi)$  is realised on a maximal subgroup  $B \in \mathcal{B}^{\omega}_{top}(\phi)$ , for some ultrafilter  $\omega$ . Moreover, B is convex-cocompact and without loss of generality  $\phi$ -invariant (Lemma 7.7). By Kurosh's theorem, the freely indecomposable factors of B are contained in the freely indecomposable factors of G, so we have  $\mathfrak{g}_{ind}(\phi|_B) \subseteq \mathfrak{g}_{ind}(\phi)$ . At the same time, we have  $\overline{\mathfrak{o}}_{top}(\phi) \in \mathfrak{g}_{ind}(\phi) \setminus \mathfrak{g}_{ind}(\phi|_B)$  by construction, so we can conclude by the (third) inductive assumption

The claim concludes Case 3, completing the proof of Proposition 7.9.

We finally conclude the section by piecing together the proof of Theorem 7.1.

Proof of Theorem 7.1. Recall that the theorem consists of five parts. Parts (1)–(4) mostly follow from the combination of Proposition 7.9 and Lemma 7.8. The only statement not covered by this is the portion of part (2) claiming that all sub-polynomial growth rates in  $\mathfrak{g}(\phi)$  are  $\preceq [n^p]$ , for an integer p depending on  $\phi$ . To prove this missing statement, let  $\mathfrak{o}_{\min}$  be the slowest pure growth rate in  $\mathfrak{g}(\phi)$ . Let  $B_1, \ldots, B_m$  be representatives of the finitely many G-conjugacy classes of maximal subgroups in  $\mathcal{B}(\mathfrak{o}_{\min})$ . The  $B_i$  are convex-cocompact and their conjugacy classes are  $\phi$ -invariant (after raising  $\phi$  to a suitable power). Moreover, each sub-polynomial growth rate in  $\mathfrak{g}(\phi)$  is realised on one of the  $B_i$ , and it is therefore  $\preceq \overline{\mathfrak{o}}_{top}(\phi|_{B_i})$ . To conclude, it now suffices to observe that the rates  $\overline{\mathfrak{o}}_{top}(\phi|_{B_i})$ are all sub-polynomial by Theorem 6.6, as all elements of the  $B_i$  grow sub-polynomially under  $\phi$ . Thus,  $\overline{\mathfrak{o}}_{top}(\phi|_{B_i}) \preceq n^{p_i}$  for some integers  $p_i \in \mathbb{N}$  and we can take  $p := \max_i p_i$ .

We are only left to prove part (5) of the theorem, which amounts to realising each pure growth rate  $\mathfrak{o} \in \mathfrak{g}(\phi)$  on a particular subgroup  $H \leq G$ . Up to replacing G with a maximal  $\mathfrak{o}$ -controlled subgroup, we can suppose that  $\mathfrak{o} \sim \overline{\mathfrak{o}}_{top}(\phi)$ . The required subgroup is then provided by Theorem 6.16(4), except that we need to avoid Case (iii) of that result, where H virtually splits as a direct product  $H' \times \mathbb{Z}^m$  for some  $m \geq 1$ . Say without loss of generality that H' has trivial centre, otherwise it has a virtual  $\mathbb{Z}$ -factor, which can be incorporated into  $\mathbb{Z}^m$ . Up to raising  $\phi$  to a power,  $\phi$  has a representative  $\varphi \in \operatorname{Aut}(G)$  that preserves  $H' \times \mathbb{Z}^m$ . By Remark 7.6, the fact that  $\varphi$  is coarsemedian preserving implies that we have  $\varphi(H') = H'$  and, up to raising  $\varphi$  to a further power, the restriction  $\varphi|_{\mathbb{Z}^m}$  is the identity. By Corollary A.6, we have  $\overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{top}(\phi|_{H'})$ . We can now replace  $\phi$  with  $\phi|_{H'} \in \operatorname{Out}(H')$  and repeat the argument, which eventually terminates because  $\operatorname{ar}(H') \leq \operatorname{ar}(G) - m < \operatorname{ar}(G)$ . This concludes the proof.

## APPENDIX A. GROWTH VS DECOMPOSITIONS

In this appendix, we discuss growth of automorphisms  $\varphi$  of a finitely generated group G, under the assumption that G decomposes into simpler pieces on which the behaviour of  $\varphi$  is known.

This decomposition could be a direct product (Section A.2), a  $\varphi$ -invariant graph of groups (Section A.3), or a free splitting (Section A.4). In each of these cases, it should not be surprising that we can describe the growth of  $\varphi$  on G in terms of that on the pieces, but the details of this are not always straightforward and they will require some work, albeit all based on standard techniques.

The main results used in the article are Corollary A.6, Proposition A.8 and Proposition A.11.

A.1. Abelian factors. Consider a product of the form  $G = H \times A$ , where H is a finitely generated group with trivial centre, and  $A \cong \mathbb{Z}^N$  for some  $N \ge 1$ . The automorphism group  $\operatorname{Aut}(G)$  can be described as follows. Consider the set  $\mathcal{M}(H, A)$  of formal matrices

$$\begin{pmatrix} \varphi & 0 \\ \alpha & \psi \end{pmatrix}$$

where  $\varphi \in \operatorname{Aut}(H)$ ,  $\psi \in \operatorname{Aut}(A) \cong \operatorname{GL}_N(\mathbb{Z})$  and  $\alpha \in \operatorname{H}^1(H, A)$ ; in other words,  $\alpha$  is a homomorphism  $H \to A$ . We can make  $\mathcal{M}(H, A)$  into a group by endowing it with a natural product:

$$\begin{pmatrix} \varphi_1 & 0\\ \alpha_1 & \psi_1 \end{pmatrix} \cdot \begin{pmatrix} \varphi_2 & 0\\ \alpha_2 & \psi_2 \end{pmatrix} = \begin{pmatrix} \varphi_1 \varphi_2 & 0\\ \alpha_1 \varphi_2 + \psi_1 \alpha_2 & \psi_1 \psi_2 \end{pmatrix}.$$

There is an action  $\mathcal{M}(H, A) \curvearrowright G$  given by

$$\begin{pmatrix} \varphi & 0 \\ \alpha & \psi \end{pmatrix} \cdot (h, a) = (\varphi(h), \alpha(h) + \psi(a)),$$

which corresponds to a homomorphism  $\iota: \mathcal{M}(H, A) \to \operatorname{Aut}(G)$ . It is not hard to see that this is actually an isomorphism.

**Lemma A.1.** If  $G = H \times A$  as above, the map  $\iota \colon \mathcal{M}(H, A) \to \operatorname{Aut}(G)$  is a group isomorphism.

Proof. Injectivity is clear, so we only need to show that  $\iota$  is surjective. Since H has trivial centre, A is the centre of G and it is preserved by all elements of  $\operatorname{Aut}(G)$ . Given  $\chi \in \operatorname{Aut}(G)$ , we can set  $\psi := \chi|_A \in \operatorname{Aut}(A)$ . Denoting by  $\pi_H, \pi_A$  the two factor projections of G, we also set  $\alpha := \pi_A \circ \chi|_H \in \operatorname{H}^1(H, A)$  and  $\varphi := \pi_H \circ \chi|_H$ . For the moment  $\varphi$  is just a homomorphism  $H \to H$ , but we can certainly write  $\chi$  as  $\chi(h, a) = (\varphi(h), \alpha(h) + \psi(a))$ . We have  $\chi(G) \subseteq \varphi(H) \times A$  and  $\chi$  is an isomorphism, so  $\varphi : H \to H$  must be surjective. Similarly we have  $\chi(\ker \varphi) \subseteq A$  and  $\chi(A) = A$ , so injectivity of  $\chi$  implies injectivity of  $\varphi$ . This shows that  $\varphi \in \operatorname{Aut}(H)$ , and thus  $\chi$  is in the image of  $\iota : \mathcal{M}(H, A) \to \operatorname{Aut}(G)$ . Since  $\chi \in \operatorname{Aut}(G)$  was arbitrary, this completes the proof of the lemma.  $\Box$ 

To simplify inline notation, from now on we will denote by  $\mathcal{M}(\varphi, \psi, \alpha)$  the element of  $\operatorname{Aut}(G)$  that is the image under  $\iota$  of the matrix  $\begin{pmatrix} \varphi & 0 \\ \alpha & \psi \end{pmatrix} \in \mathcal{M}(H, A)$ . For each  $n \geq 1$ , we have  $\chi^n = \mathcal{M}(\varphi^n, \psi^n, \alpha_n)$ , where  $\alpha_n = \sum_{j=1}^n \psi^{j-1} \alpha \varphi^{n-j}$ . Thus, the growth rates of  $\mathcal{M}(\varphi, \psi, \alpha)$  and its projection to  $\operatorname{Out}(G)$ (as defined in Section 2.2) can be described fairly easily in terms of the growth rates of  $\varphi$  and  $\psi$ .

For a finitely generated group Q, we denote by  $Q_{ab}$  the free part of the abelianisation of Q. Every automorphism  $\chi \in \operatorname{Aut}(Q)$  naturally descends to an automorphism of  $\chi_{ab} \in \operatorname{Aut}(Q_{ab})$ , and each element  $q \in Q$  equivariantly projects to an element  $q_{ab} \in Q_{ab}$ .

**Lemma A.2.** Let  $G = H \times A$  be as above. For each automorphism  $\chi = \mathcal{M}(\varphi, \psi, \alpha) \in \operatorname{Aut}(G)$  and each element  $g = (h, a) \in G$ , we have the following equalities in the set of abstract growth rates  $\mathfrak{G}$ :

$$|\chi^{n}(g)| \sim |\varphi^{n}(h)| + |\chi^{n}_{ab}(g_{ab})|, \qquad \qquad \|\chi^{n}(g)\| \sim \|\varphi^{n}(h)\| + |\chi^{n}_{ab}(g_{ab})|.$$

Moreover, there exist an algebraic integer  $\lambda \geq 1$  and  $p \in \mathbb{N}$  such that  $|\chi_{ab}^n(g_{ab})| \sim n^p \lambda^n$ .

Proof. The element  $\chi^n(g)$  has coordinates  $\varphi^n(h)$  and  $\alpha_n(g) + \psi^n(a)$  in H and A respectively, where  $\alpha_n$  is the homomorphism described above. We have  $G_{ab} = H_{ab} \oplus A$ , and the homomorphisms  $\alpha_n \to H \to A$  factor through  $H_{ab}$ . Thus, the coordinate of  $\chi^n_{ab}(g_{ab})$  along A is identical to that of  $\chi^n(g)$ . We obtain the inequality

$$|\chi^n(g)| \sim |\varphi^n(h)| + |\alpha_n(g) + \psi^n(a)| \leq |\varphi^n(h)| + |\chi^n_{ab}(g_{ab})|$$

as well as the analogous inequality for  $\|\chi^n(g)\|$ . The reverse inequalities are immediate from the fact that the projections  $G \to H$  and  $G \to G_{ab}$  are Lipschitz.

Finally, we have  $|\chi_{ab}(g_{ab})| \sim n^p \lambda^n$  for an algebraic integer  $\lambda \geq 1$  and some  $p \in \mathbb{N}$  because  $G_{ab}$  is free abelian and we can invoke the classical Lemma A.3 below. Algebraic integers are closed under taking complex conjugates, products, square roots and thus also under taking moduli.

**Lemma A.3.** Let  $k \geq 1$ . For each automorphism  $\varphi \in \operatorname{Aut}(\mathbb{Z}^k) = \operatorname{GL}_k(\mathbb{Z})$  and each element  $a \in \mathbb{Z}^k \setminus \{0\}$ , there exist  $p \in \mathbb{N}$  and  $\mu \in \mathbb{C}$  with  $|\mu| \geq 1$  such that

$$|\varphi^n(a)| = \|\varphi^n(a)\| \sim n^p |\mu|^n.$$

Moreover,  $\varphi \in \operatorname{GL}_k(\mathbb{Z}) \leq \operatorname{GL}_k(\mathbb{C})$  has a (p+1)-dimensional Jordan block with eigenvalue  $\mu$ .

**Example A.4.** Consider the product  $G = F_k \times \mathbb{Z}^k$  for  $k \geq 3$ . There exists an automorphism  $\chi \in \operatorname{Aut}(G)$  that has exactly three growth rates — namely [1],  $[\lambda^n]$  and  $[n\lambda^n]$  for some  $\lambda > 1$  — where [1] is realised at the identity,  $[\lambda^n]$  is realised by nontrivial elements of a subgroup  $N \times \mathbb{Z}^k \triangleleft G$ , where

 $N \triangleleft F_n$  is infinitely generated, and  $[n\lambda^n]$  is realised by all remaining elements of G. In particular, this shows that Theorem B(3) fails when the automorphism is not coarse-median preserving.

The construction of such automorphisms  $\chi$  is fairly general. We start with a positive, fully irreducible automorphism<sup>22</sup>  $\varphi \in \operatorname{Aut}(F_k)$ . Let  $\varphi_{ab}$  be the induced automorphism of the abelianisation  $(F_k)_{ab} \cong \mathbb{Z}^k$  and let  $\alpha \colon F_k \to (F_k)_{ab}$  be the quotient projection. Set  $G := F_k \times (F_k)_{ab} \cong F_k \times \mathbb{Z}^k$  and  $\chi := \mathcal{M}(\varphi, \varphi_{ab}, \alpha)$ , in the above notation. We now prove the above statements.

Let  $\lambda$  be the Perron–Frobenius eigenvalue of  $\varphi$ . By the existence of train-track maps, we have  $|\varphi^n(h)| \sim ||\varphi^n(h)|| \sim \lambda^n$  for all nontrivial elements  $h \in F_k$  [BH92]. We also have  $|\varphi^n_{ab}(\alpha(h))| \leq |\varphi^n(h)|$  for all  $h \in F_k$ , in terms of the standard generating sets of  $F_k$  and  $(F_k)_{ab}$ . This inequality is an equality whenever h is a positive element of  $F_k$ , since  $\varphi$  is a positive automorphism.

Note that  $\chi^n = \mathcal{M}(\varphi^n, \varphi_{ab}^n, \alpha_n)$  for each  $n \ge 1$ , where  $\alpha_n = n \cdot (\alpha \circ \varphi^{n-1}) = n \cdot (\varphi_{ab}^{n-1} \circ \alpha)$ . Thus,

$$\chi^n(h,a) = \left(\varphi^n(h), \ \varphi^n_{\rm ab}(a) + n \cdot (\varphi^{n-1}_{\rm ab}\alpha)(h)\right).$$

It follows that  $\lambda^n \leq |\chi^n(g)| \leq n\lambda^n$  for every nontrivial element  $g \in G$ . In addition, we have  $|\chi^n(g)| \sim n\lambda^n$  if  $g = (h, \alpha(h)) \in F_k \times (F_k)_{ab}$  for a positive element  $h \in F_k$ . The same statements hold for conjugacy-length growth.

Now, let  $\beta: F_k \to \mathbb{R}^k$  be the limit for  $n \to +\infty$  of the homomorphisms

$$\frac{1}{n\lambda^n}\alpha_n = \lambda^{-n}(\varphi_{ab}^n \circ \alpha) \colon F_k \to (F_k)_{ab} \cong \mathbb{Z}^k \hookrightarrow \mathbb{R}^k.$$

That this limit exists follows from the fact that a power of  $\varphi_{ab} \in \operatorname{GL}_k(\mathbb{Z})$  is represented by a matrix with positive entries and the Perron–Frobenius eigenvalue of  $\varphi_{ab}$  is  $\lambda$ , so all other eigenvalues of  $\varphi_{ab}$  have modulus  $< \lambda$ . The homomorphism  $\beta$  is nontrivial, as it does not vanish on positive elements of  $F_k$ , by the above discussion.

Any nontrivial element g of ker  $\beta \times \mathbb{Z}^k$  clearly has  $|\chi^n(g)| \sim ||\chi^n(g)|| \sim \lambda^n$ , while all elements of G outside ker  $\beta \times \mathbb{Z}^k$  have  $[n\lambda^n]$  as their growth rate, by construction. This proves all our claims.

A.2. Direct products. Consider now a finitely generated group  $G = G_1 \times \ldots \times G_k \times A$ , where  $k \ge 0$  and A is free abelian. Assume that each  $G_i$  is directly indecomposable and has trivial centre.

**Lemma A.5.** The automorphism group  $\operatorname{Aut}(G)$  permutes the subgroups  $\langle G_i, A \rangle$  for  $1 \leq i \leq k$ .

Proof. Consider some  $\varphi \in \operatorname{Aut}(G)$ . Since A is the centre of G, we have  $\varphi(A) = A$ . Thus, G is generated by the subgroups  $\varphi(G_1), \ldots, \varphi(G_k), A$ . Denoting by  $\pi_1 \colon G \to G_1$  the factor projection, it follows that  $G_1$  is generated by the pairwise-commuting subgroups  $\pi_1\varphi(G_1), \ldots, \pi_1\varphi(G_k)$ . Since  $G_1$  has trivial centre, each subgroup  $\pi_1\varphi(G_i)$  has trivial intersection with the subgroup generated by the other k-1 subgroups  $\pi_1\varphi(G_i)$ . It follows that

$$G_1 = \pi_1 \varphi(G_1) \times \ldots \times \pi_1 \varphi(G_k).$$

Since  $G_1$  is directly indecomposable, we must have  $G_1 = \pi_1 \varphi(G_i)$  for some index i and  $\pi_1 \varphi(G_j) = \{1\}$  for all  $j \neq i$ . Repeating the argument for all factors of G, we obtain a permutation  $\sigma \in \text{Sym}(k)$  such that  $G_i = \pi_i \varphi(G_{\sigma(i)})$  for all  $1 \leq i \leq k$  and  $\pi_i \varphi(G_j) = \{1\}$  for all  $j \neq \sigma(i)$ . This shows that  $\langle \varphi(G_j), A \rangle = \langle G_{\sigma^{-1}(j)}, A \rangle$  for all  $1 \leq j \leq k$ , completing the proof.

Let  $\operatorname{Aut}^0(G) \leq \operatorname{Aut}(G)$  be the finite-index subgroup preserving each subgroup  $\langle G_i, A \rangle$ . When  $A = \{1\}$ , the elements of  $\operatorname{Aut}^0(G)$  are simply products  $\varphi_1 \times \ldots \times \varphi_k$  with  $\varphi_i \in \operatorname{Aut}(G_i)$ . We can now apply Lemma A.2 to immediately deduce the following.

<sup>&</sup>lt;sup>22</sup>A simple such example in  $F_3 = \langle a, b, c \rangle$  is the automorphism  $\varphi$  mapping  $a \mapsto ab, b \mapsto bc, c \mapsto cab$ .

**Corollary A.6.** Let  $G = G_1 \times \ldots \times G_k \times A$  be as above. For each  $\varphi \in \operatorname{Aut}^0(G)$ , there exist automorphisms  $\varphi_i \in \operatorname{Aut}(G_i)$  such that, for each element  $g = (g_1, \ldots, g_k, a) \in G$ , we have:

$$|\varphi^{n}(g)| \sim \sum_{i=1}^{k} |\varphi^{n}_{i}(g_{i})| + |\varphi^{n}_{ab}(g_{ab})|, \qquad \qquad \|\varphi^{n}(g)\| \sim \sum_{i=1}^{k} \|\varphi^{n}_{i}(g_{i})\| + |\varphi^{n}_{ab}(g_{ab})|.$$

Moreover, for each  $\varphi$  and g, there exist an algebraic integer  $\lambda \geq 1$  and  $p \in \mathbb{N}$  with  $|\varphi_{ab}^n(g_{ab})| \sim n^p \lambda^n$ .

A.3. Invariant splittings. We now consider a finitely generated group G and an outer automorphism  $\phi \in \text{Out}(G)$  that preserves a splitting of G as a graph of groups. The next lemma is a straightforward exercise in Bass–Serre theory.

**Lemma A.7.** Consider a group G with a one-edge splitting as

$$G = A *_C B$$
 or  $G = A *_{C,\gamma} = \langle A, t \mid t^{-1}ct = \gamma(c), \forall c \in C \rangle.$ 

If, for some  $\phi \in \text{Out}(G)$ , the Bass–Serre tree  $G \curvearrowright T$  extends to an action  $G \rtimes_{\phi} \mathbb{Z} \curvearrowright T$  without inversions, then  $\phi$  is represented by an automorphism  $\varphi \in \text{Aut}(G)$  of the following form:

- (1) In the amalgamated product case, we have  $\varphi(A) = A$  and  $\varphi(B) = B$ .
- (2) In the HNN case, we have  $\varphi(A) = A$ ,  $\varphi(C) = C$  and  $\varphi(t) = ta$  for some  $a \in A$ .

The next proposition is the main result of this short section. The important takeaway is that, in a  $\phi$ -invariant graph of groups, growth on the whole group cannot be strictly faster than growth on the vertex groups, provided that the latter is sufficiently well-behaved and at least exponential.

Recall that the notation  $\overline{\mathcal{O}}_{top}(\cdot)$  and  $\overline{\mathfrak{o}}_{top}(\cdot)$  was introduced in Section 2.2.3. Docile automorphisms were defined in Section 6.1. Finally, given an outer automorphism  $\phi \in Out(G)$  and a subgroup  $H \leq G$  with  $\phi$ -invariant G-conjugacy class, we can consider restrictions  $\phi|_V \in Out(V)$ , although these are not uniquely defined in general (Remark 4.12).

We say that a finitely generated subgroup  $H \leq G$  is *conjugacy-undistorted* if H is undistorted in G and, in addition, the conjugacy length functions of G and H are bi-Lipschitz equivalent on H. For instance, convex-cocompact subgroups of special groups are conjugacy-undistorted (Remark 2.17).

**Proposition A.8.** Consider a finitely generated group G with a  $\phi$ -invariant splitting  $G \curvearrowright T$  for some  $\phi \in \text{Out}(G)$ . Suppose that  $\phi$  descends to the identity on the finite graph T/G. Additionally, suppose that there exist integers  $p, q \in \mathbb{N}$  and a finite subset  $\Lambda \subseteq \mathbb{R}_{>1}$  such that, for each vertex  $v \in T$ , the G-stabiliser of v (denoted V) satisfies **at least one** of the following conditions:

- (a) for every representative  $\varphi \in \operatorname{Aut}(G)$  of  $\phi$  with  $\varphi(V) = V$  and every  $g \in V$ , we have  $|\varphi^n(g)| \leq n^q$  (computing word lengths with respect to a finite generating set of G);
- (b) V is finitely generated, conjugacy-undistorted in G and, for every representative  $\varphi \in \operatorname{Aut}(G)$ of  $\phi$  with  $\varphi(V) = V$ , the restriction  $\varphi|_V \in \operatorname{Aut}(V)$  is  $(\lambda, p)$ -docile for some  $\lambda \in \Lambda$ .

Then, the following properties hold.

- (1) If there are no type (b) vertex groups, we have  $\overline{\mathcal{O}}_{top}(G,\varphi) \preceq n^{q+2}$  for all representatives  $\varphi \in Aut(G)$  of  $\phi$ .
- (2) If there is at least one type (b) vertex group, then  $\phi \in \text{Out}(G)$  is  $(\mu, p)$ -docile for  $\mu = \max \Lambda$ . Moreover, we can choose finitely many type (b) vertex groups  $V_1, \ldots, V_k \leq G$  and restrictions  $\phi_i \in \text{Out}(V_i)$  of the outer automorphism  $\phi$  such that  $\overline{\mathfrak{o}}_{\text{top}}(\phi) \sim \sum_{i=1}^k \overline{\mathfrak{o}}_{\text{top}}(\phi_i)$ .

*Proof.* To begin with, we prove the proposition when T is a one-edge splitting of G, that is, an amalgamation or an HNN extension. In each of these two cases, we represent  $\phi$  by an automorphism  $\varphi \in \operatorname{Aut}(G)$  of the form in Lemma A.7. At the end of the proof, we will briefly explain how to handle general graphs of groups based on this.

Suppose first that  $G = A *_C B$  with  $\varphi(A) = A$  and  $\varphi(B) = B$ . A general element of G can be written as  $g = a_1b_1 \dots a_kb_k$  with  $a_i \in A$  and  $b_i \in B$ , and we have

$$\varphi^n(g) = \varphi^n(a_1)\varphi^n(b_1)\dots\varphi^n(a_k)\varphi^n(b_k), \quad \forall n \in \mathbb{N}.$$

This yields the simple bound  $|\varphi^n(g)| \leq \sum_{i=1}^k |\varphi^n(a_i)| + \sum_{i=1}^k |\varphi^n(b_i)|$ . In part (1), we immediately obtain  $|\varphi^n(g)| \leq n^q$  for all  $g \in G$ , and hence  $\overline{\mathcal{O}}_{top}(\varphi) \leq n^q$ . Regarding part (2), suppose for simplicity that A is of type (a) and B is of type (b) (the case when both A and B are of type (b) is similar). As  $\varphi|_B$  is docile, every element  $g \in G$  satisfies

$$|\varphi^n(g)| \leq n^q + \overline{\mathcal{O}}_{top}(\varphi|_B) \leq \overline{\mathcal{O}}_{top}(\varphi|_B) \sim \overline{\mathfrak{o}}_{top}([\varphi|_B]).$$

Hence  $\overline{\mathcal{O}}_{top}(\varphi) \preceq \overline{\mathfrak{o}}_{top}([\varphi|_B])$ . At the same time, the fact that B is conjugacy-undistorted implies that  $\overline{\mathfrak{o}}_{top}(\phi) \succeq \overline{\mathfrak{o}}_{top}([\varphi|_B])$ . Thus, we obtain  $\overline{\mathcal{O}}_{top}(\varphi) \sim \overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{top}([\varphi|_B])$ , showing that  $\phi$  is docile, as required. Finally, if both vertex groups A and B are of type (b), the same arguments show that  $\overline{\mathcal{O}}_{top}(\varphi) \sim \overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{top}([\varphi|_A]) + \overline{\mathfrak{o}}_{top}([\varphi|_B])$ .

Suppose now that  $G = A *_C$  with  $\varphi(A) = A$ ,  $\varphi(C) = C$  and  $\varphi(t) = ta$ , where t is the stable letter of the HNN extension and  $a \in A$  is some element. A general element of G can be written as  $g = x_0 t^{\epsilon_1} x_1 \dots t^{\epsilon_k} x_k$  with  $x_i \in A$  and  $\epsilon_i \in \{\pm 1\}$ . We then have

$$\varphi^n(g) = x'_0 t^{\epsilon_1} x'_1 \dots t^{\epsilon_k} x'_k,$$

where, setting  $a_n := a\varphi(a)\varphi^2(a)\ldots\varphi^{n-1}(a)$ , each element  $x'_i$  is one of the following four options (depending on the values of  $\epsilon_i$  and  $\epsilon_{i+1}$ ): either  $\varphi^n(x_i)$ , or  $a_n\varphi^n(x_i)$ , or  $\varphi^n(x_i)a_n^{-1}$ , or  $a_n\varphi^n(x_i)a_n^{-1}$ . This yields the inequality

$$\overline{\mathcal{O}}_{\mathrm{top}}(\varphi) \preceq \overline{\mathcal{O}}_{\mathrm{top}}(\varphi|_A) + |a_n|.$$

In part (1), we have  $|a_n| \leq n^{q+1}$  and we obtain  $\overline{\mathcal{O}}_{top}(\varphi) \leq n^{q+1}$ . In part (2), the fact that  $\varphi|_A$  is docile yields  $|a_n| \leq \overline{\mathcal{O}}_{top}(\varphi|_A)$  (Remark 6.3), and hence  $\overline{\mathcal{O}}_{top}(\varphi) \leq \overline{\mathcal{O}}_{top}(\varphi|_A)$ . At the same time, since A is conjugacy-undistorted, we have  $\overline{\mathfrak{o}}_{top}(\phi) \succeq \overline{\mathfrak{o}}_{top}([\varphi|_A])$  and as above  $\overline{\mathcal{O}}_{top}(\varphi) \sim \overline{\mathfrak{o}}_{top}(\phi) \sim \overline{\mathfrak{o}}_{top}([\varphi|_A])$ , showing that  $\phi$  is docile.

So far, we have only considered a *specific* representative  $\varphi$  of the outer class  $\phi$ . This is irrelevant in part (2), while in part (1) it causes the bound  $\overline{\mathcal{O}}_{top}(\varphi) \preceq n^{q+1}$  to translate itself into the weaker bound  $\overline{\mathcal{O}}_{top}(\varphi') \preceq n^{q+2}$  for a general representative  $\varphi'$  (Remark 6.5).

This proves the proposition when  $G \curvearrowright T$  is a one-edge splitting. In general, we can decompose the  $\phi$ -invariant splitting  $G \curvearrowright T$  as a finite sequence of  $\phi$ -invariant one-edge splittings (collapsing all G-orbits of edges of T but one, and then adding them back one at a time). An iterated application of the one-edge case then immediately proves part (2). Part (1) requires a little more care, as we want the polynomial exponent q to increase by at most 2. For this, it suffices to note that T can be decomposed as a finite sequence of amalgamated products followed by a single multiple HNN extension (a splitting whose quotient graph is a wedge of circles, with a single vertex). Amalgamations do not increase q at all (for a specific representative  $\varphi$ ), while the final multiple HNN splitting increases it by at most 1 (here the argument is identical to the single HNN case, except that we will have finitely many stable letters  $t_i$  and, in general,  $\varphi$  can only be put in the form  $\varphi(t_i) = a_i t a'_i$  with  $a_i, a'_i \in A$ ). Finally, there is a further increase by 1 to handle arbitrary representatives  $\varphi'$  using Remark 6.5. This concludes the proof of the proposition.

**Remark A.9.** If in part (2) of Proposition A.8 the growth rates  $\overline{\mathfrak{o}}_{top}(\phi_i)$  are all pure (Definition 6.1), then  $\overline{\mathfrak{o}}_{top}(\phi) \sim \sum_i \overline{\mathfrak{o}}_{top}(\phi_i)$  simply equals the fastest of the  $\overline{\mathfrak{o}}_{top}(\phi_i)$ .

A.4. Free products. Let G be a finitely generated group. A *(free) factor system* for G is the collection  $\mathcal{F}$  of all G-conjugates of the subgroups  $G_1, \ldots, G_k \leq G$  appearing in a decomposition

$$G = G_1 * \dots * G_k * F_m$$

with  $k, m \ge 0$  and  $k + m \ge 2$ . We require the  $G_i$  to be nontrivial, but not that they be freely indecomposable; we allow  $\mathcal{F}$  to be empty if  $G \cong F_m$  for  $m \ge 2$ . The group G admits a factor system whenever it is neither freely indecomposable nor isomorphic to  $\mathbb{Z}$ .

Following [GH22], we say that the pair  $(G, \mathcal{F})$  is *sporadic* if  $(k, m) \in \{(2, 0), (1, 1)\}$ . A  $(G, \mathcal{F})$ -free factor is a subgroup of G arising as a vertex group in a free splitting of G relative to  $\mathcal{F}$ . A  $(G, \mathcal{F})$ -free factor is *proper* if it is neither the trivial group nor an element of  $\mathcal{F}$ . Let  $Out(G, \mathcal{F}) \leq Out(G)$  be the subgroup of outer automorphisms that leave invariant each G-conjugacy class of subgroups in  $\mathcal{F}$ . An element  $\phi \in Out(G, \mathcal{F})$  is *fully irreducible* if none of its (nontrivial) powers preserves the G-conjugacy class of a proper  $(G, \mathcal{F})$ -free factor.

The following is an easy equivalent characterisation of full irreducibles. If  $\mathcal{F}_1, \mathcal{F}_2$  are factor systems for G, we write  $\mathcal{F}_1 \leq \mathcal{F}_2$  if each subgroup in  $\mathcal{F}_1$  is contained in a subgroup in  $\mathcal{F}_2$ .

**Lemma A.10.** If  $\phi \in \text{Out}(G, \mathcal{F})$  is not fully irreducible, then there exist a factor system  $\mathcal{F}' > \mathcal{F}$ and an integer  $p \geq 1$  such that  $\phi^p \in \text{Out}(G, \mathcal{F}')$ .

As a first step, we can characterise growth of fully irreducible automorphisms in terms of their growth on the elements of the factor system. This is a relatively straightforward (but also rather fiddly) application of train track maps for free products [CT94, FM15, Lym22b].

Recall from Section 2.2.3 that  $\mathcal{G}(\varphi) \subseteq \mathfrak{G}$  and  $\mathfrak{g}(\phi) \subseteq \mathfrak{G}$  denote the sets of growth rates of an automorphism  $\varphi$  and an outer automorphism  $\phi$ , respectively. The rate  $\overline{\mathfrak{o}}_{top}(\phi)$  always bounds all elements of  $\mathfrak{g}(\phi)$  from above, but it might not lie in  $\mathfrak{g}(\phi)$  a priori; the same is true of  $\overline{\mathcal{O}}_{top}(\varphi)$  and  $\mathcal{G}(\varphi)$ . Also recall that we refer to polynomial-times-exponential growth rates as *pure* (Definition 6.1).

**Proposition A.11.** Consider a finitely generated group  $G = G_1 * \cdots * G_k * F_m$ , let  $\mathcal{F}$  be the factor system given by the  $G_i$ , and let  $\phi \in \text{Out}(G, \mathcal{F})$ . Suppose that  $\phi$  is fully irreducible and  $\mathcal{F}$  is nonsporadic. In addition, suppose that each restriction  $\phi_i := \phi|_{G_i} \in \text{Out}(G_i)$  is represented by some  $\varphi_i \in \text{Aut}(G_i)$  such that either  $\overline{\mathcal{O}}_{\text{top}}(\varphi_i)$  is sub-polynomial, or  $\varphi_i$  is  $(\lambda_i, p_i)$ -docile for some  $\lambda_i > 1$ and  $p_i \in \mathbb{N}$ . Then there exists a Perron number  $\lambda > 1$  such that all the following hold.

- (1) The automorphism  $\phi$  is  $(\mu, q)$ -docile for  $\mu = \max\{\lambda_1, \dots, \lambda_k, \lambda\}$  and some  $q \in \mathbb{N}$ . We have  $q \leq p_j + 1$  for the largest integer  $p_j$  such that  $\lambda_j = \mu$ ; if no such index j exists, then q = 0.
- (2) If we have  $[\mu^n] \in \mathfrak{g}(\phi_i)$  whenever  $\lambda_i = \mu$ , then  $[\mu^n] \in \mathfrak{g}(\phi)$ .
- (3) Suppose that there exist a finite set  $\Lambda \subseteq \mathbb{R}_{>1}$  and  $P \in \mathbb{N}$  such that, for all indices i and all representatives  $\varphi'_i \in \operatorname{Aut}(G_i)$  of  $\phi_i \in \operatorname{Out}(G_i)$ , each growth rate in the union  $\mathcal{G}(\varphi'_i) \cup \mathfrak{g}(\phi_i)$  is either sub-polynomial or equal to  $[n^a \nu^n]$ , for some  $\nu \in \Lambda$  and some integer  $0 \leq a \leq P$ . Then, each growth rate in the set  $\mathfrak{g}(\phi)$  is either sub-polynomial, or equal  $[\lambda^n]$ , or equal to  $[n^a \nu^n]$  for some  $\nu \in \Lambda$  and some integer  $0 \leq a \leq P + 1$ .

*Proof.* We begin with a general discussion that only assumes finite generation of G and no additional properties of the restrictions  $\phi|_{G_i}$ ; this discussion will culminate in Equations A.1 and A.3 below, estimating the word-length growth of elements not conjugate into an element of  $\mathcal{F}$ , and the top growth rate of  $\phi$ , respectively. After this, we will add the docility assumption and draw the necessary conclusions. Throughout, we omit some technical details in the interest of overall clarity.

We start by representing G as the fundamental group of a graph of groups  $\mathcal{G}$  with trivial edge groups and precisely the elements of  $\mathcal{F}$  as G-conjugates of the nontrivial vertex groups. We can find  $\mathcal{G}$  so that, in addition,  $\phi$  is realised by a relative train track map  $f: \mathcal{G} \to \mathcal{G}$ ; see e.g. [Lym22a, Section 1]. Let  $\overline{\mathcal{G}}$  be the finite graph underlying the graph of groups  $\mathcal{G}$  and let  $\overline{f}: \overline{\mathcal{G}} \to \overline{\mathcal{G}}$  be the map of graphs underlying f. Choosing a base vertex p and a spanning tree for  $\overline{\mathcal{G}}$ , we identify the vertex groups of  $\mathcal{G}$  with the  $G_i$ , and we fix a representative  $\varphi \in \operatorname{Aut}(G)$  of the outer class  $\phi$ . For notational convenience, choose automorphisms  $\varphi_i \in \operatorname{Aut}(G_i)$  representing the restrictions  $\phi_i \in \operatorname{Out}(G_i)$ , and denote by  $\psi$  the self-bijection of the disjoint union  $\bigsqcup_i G_i$  that equals  $\varphi_i$  on each  $G_i$  (note that  $\psi$  is not the restriction of a single element of  $\operatorname{Aut}(G)$  in general). A generalised path is a string  $\pi = g_0 e_1 g_1 \dots e_s g_s$  such that  $e_1 \dots e_s$  is an (oriented) edge path in  $\overline{\mathcal{G}}$  and each  $g_j$  lies in the vertex group  $G_i$  associated to the terminal vertex of  $e_j$  (which is also the initial vertex of  $e_{j+1}$ ). The edge length of  $\pi$  is  $\ell(\pi) := s$  and the total length is  $|\pi| := s + \sum_j |g_j|$ , where the word lengths  $|g_j|$  are computed with respect to some fixed choice of finite generating sets for the  $G_i$ . The generalised path  $\pi$  is immersed if we do not have  $g_j = 1$  and  $e_{j+1} = e_j^{-1}$  for any index j. Elements of G are in 1-to-1 correspondence with closed, immersed generalised paths based at  $p \in \overline{\mathcal{G}}$ . The word length of an element  $g \in G$  is roughly the same as the total length of the generalised path  $\pi$  representing g, up to a multiplicative constant independent of g.

Since  $\phi$  is fully irreducible, no power of  $\overline{f}$  leaves invariant a proper subgraph of  $\overline{\mathcal{G}}$  with at least one edge (up to collapsing some edges of  $\mathcal{G}$  without altering the vertex groups). In particular, the train track map f has only one stratum containing edges; let  $\lambda \geq 1$  be the Perron–Frobenius eigenvalue of its transition matrix. Up to raising  $\phi$  and f to a power, we can further assume that:

- (i)  $\overline{f}$  fixes every vertex of  $\overline{\mathcal{G}}$ ;
- (ii) for each edge  $e \subseteq \mathcal{G}$ , the path  $\overline{f}(e)$  contains all edges of  $\overline{\mathcal{G}}$ ;
- (iii) for each (oriented) edge  $e \subseteq \mathcal{G}$ , the edge paths  $\overline{f^n}(e) \subseteq \overline{\mathcal{G}}$  all start with the same edge  $e_1$  and end with the same edge  $e_2$ , for  $n \ge 1$ .

Since  $\mathcal{F}$  is non-sporadic, we do not have  $\overline{f}(e) = e$  for any edge e. Thus, the edge lengths  $\ell(f^n(e))$  grow exponentially with n and hence  $\lambda \neq 1$ .

If  $\pi = g_0 e_1 g_1 \dots e_s g_s$  is a generalised path, its image  $f(\pi)$  is  $\psi(g_0) f(e_1) \psi(g_1) \dots f(e_s) \psi(g_s)$ , recalling that we have defined  $\psi = \varphi_i$  on each  $G_i$ . Here, each  $f(e_i)$  is a generalised path having  $\overline{f}(e_i)$  as underlying edge path.

In order to easily estimate the total length of the paths  $f^n(e)$ , we define one last auxiliary concept. If  $\pi = g_0 e_1 g_1 \dots e_s g_s$ , we refer to the slots between consecutive edges as the nodes of  $\pi$ ; thus,  $\pi$  has s+1 nodes and they are occupied by the elements  $g_0, \dots, g_s$ . To each node of  $f^n(\pi)$ , we inductively associate an order, which is an integer between 0 and n. All nodes of  $\pi$  have order 0. If e is an edge of  $f^{n-1}(\pi)$ , then f(e) is a subpath of  $f^n(\pi)$ ; all nodes of  $f^n(\pi)$  that are interior nodes of f(e) for some edge  $e \subseteq f^{n-1}(\pi)$  are also declared to have order 0. Every other node of  $f^n(\pi)$  is the image under f of a node of  $f^{n-1}(\pi)$ ; if the node of  $f^{n-1}(\pi)$  had order m, for some  $m \ge 0$ , then we declare the corresponding node of  $f^n(\pi)$  to have order m + 1. The initial and terminal nodes of  $f^n(\pi)$  thus have order n.

A straightforward computation shows that, for all  $0 \leq i < n$ , the path  $f^n(\pi)$  has exactly  $\ell(f^{n-i}(\pi)) - \ell(f^{n-i-1}(\pi))$  order-*i* nodes, and it has  $\ell(\pi) + 1$  order-*n* nodes. Choose a constant  $C \geq 1$  such that every edge  $e \subseteq \mathcal{G}$  satisfies

$$\frac{1}{C}\lambda^n \le \ell(f^n(e)) \le C\lambda^n, \qquad \forall n \in \mathbb{N}.$$

Since  $\ell(f(e)) \ge 2$  for all edges  $e \subseteq \mathcal{G}$ , we also have  $\ell(f^n(e)) - \ell(f^{n-1}(e)) \ge \ell(f^{n-1}(e)) \ge \frac{1}{\lambda C} \lambda^n$ .

Let  $\Omega$  be the finite set of group elements appearing at the nodes of the paths f(e), as e varies through the edges of  $\mathcal{G}$ ; we add the identity  $1 \in G$  to  $\Omega$  to simplify notation in the coming discussion. For any  $n \geq 0$ , every order-0 node of  $f^n(e)$  is occupied by an element lying in  $\Omega$ . Property (iii) implies that, for  $1 \leq i \leq n$ , every order-*i* node of  $f^n(e)$  is occupied by an element of the form

$$\left(a\psi(a)\psi^2(a)\dots\psi^{i-2}(a)\right)\cdot\psi^{i-1}(b)\psi^i(c)\psi^{i-1}(d)\cdot\left(\psi^{i-2}(e)\dots\psi^2(e)\psi(e)e\right),\quad\text{where }a,b,c,d,e\in\Omega.$$

In particular, fixing *i*, there is only a fixed finite set of elements that can occupy the order-*i* nodes in the paths  $f^n(e)$ , as *n* and *e* are allowed to vary. Moreover, this finite set has at most  $|\Omega|^5$  elements, independently of *i*. Denote by  $N_i$  the average word length of these finitely many elements.

For any edge  $e \subseteq \mathcal{G}$ , we can write

$$|f^{n}(e)| = \ell(f^{n}(e)) + \sum_{i=0}^{n-1} N_{i}^{e,n} \cdot \left(\ell(f^{n-i}(e)) - \ell(f^{n-i-1}(e))\right) + N_{n}^{e,n}(\ell(e)+1),$$

where  $N_i^{e,n}$  is the (weighted) average word length of the elements occupying the order-*i* nodes of  $f^n(e)$ . Recalling property (ii), the  $N_i^{e,n}$  are roughly equal to  $N_i$ , up to a multiplicative constant independent of e, i, n. Hence, up to slightly enlarging the constant C chosen above, we have

$$\frac{1}{C} \cdot \sum_{i=0}^{n} N_i \lambda^{n-i} \le |f^n(e)| \le C \sum_{i=0}^{n} N_i \lambda^{n-i}, \quad \text{for all } n \in \mathbb{N} \text{ and } e \subseteq \mathcal{G}$$

For any immersed path  $\pi = g_0 e_1 g_1 \dots e_s g_s$ , denote by  $|f^n(\pi)|_{\text{pt}}$  the length of the path  $f^n(\pi)$  pulled *tight*, that is, the length of the immersed path homotopic to  $f^n(\pi)$ . If the length  $|f^n(\pi)|_{\rm pt}$  does not stay bounded as n increases, the above inequalities allow us to estimate

(A.1) 
$$|f^{n}(\pi)|_{\text{pt}} \sim \sum_{j=0}^{s} |\psi^{n}(g_{j})| + \ell(\pi) \cdot \sum_{j=0}^{n} N_{j} \lambda^{n-j},$$

where the multiplicative constant implicit in the symbol ~ does not depend on  $\pi$  or the integer n. Here, the inequality  $\gtrsim$  uses bounded backtracking [Coo87] as in [BH92], working in the graph  $\mathcal{G}$ .

Equation (A.1) shows that, for any element  $g \in G$  not conjugate into one of the subgroups  $G_1, \ldots, G_k$ , either the growth rate  $| \| \phi^n(g) \| |$  is bounded, or we have:

(A.2) 
$$\left[ \|\phi^n(g)\| \right] \sim \left[ |\varphi^n(g)| \right] \preceq \sum_{i=1}^k \overline{\mathcal{O}}_{top}(\varphi_i) + \left[ \sum_{j=0}^n N_j \lambda^{n-j} \right].$$

In particular, we claim that this implies the following equality in  $\mathfrak{G}$ :

(A.3) 
$$\overline{\mathcal{O}}_{\rm top}(\varphi) \sim \sum_{i=1}^{k} \overline{\mathcal{O}}_{\rm top}(\varphi_i) + \big[\sum_{j=0}^{n} N_j \lambda^{n-j}\big].$$

The inequality  $\leq$  is immediate from the definition of  $\overline{\mathcal{O}}_{top}(\varphi)$  and Equation (A.2). Conversely, it is clear that  $\overline{\mathcal{O}}_{top}(\varphi) \succeq \overline{\mathcal{O}}_{top}(\varphi_i)$  for all *i*, while it follows from Equation (A.1) that we also have  $\overline{\mathcal{O}}_{top}(\varphi) \succeq \sum_{j=0}^{n} N_j \lambda^{n-j}$ . Thus,  $\overline{\mathcal{O}}_{top}(\varphi)$  is (coarsely) bounded below by the sum of these rates.

The entire discussion up to this point was completely general: we have not made use of any assumptions on the  $G_i$  other than finite generation.

From now on assume that, for each i, either  $\varphi_i \in \operatorname{Aut}(G_i)$  is  $(\lambda_i, p_i)$ -docide for some  $\lambda_i > 1$  and  $p_i \in \mathbb{N}$ , or the growth rate  $\overline{\mathcal{O}}_{top}(\varphi_i)$  is sub-polynomial (in which case, we set  $\lambda_i := 1$  for convenience). We proceed to discuss the claims in the various parts of the proposition.

**Part** (1). Recalling Remark 6.3 and the definition of  $N_n$ , we have

(A.4) 
$$[N_n] \preceq \sum_{i=1}^k \overline{\mathcal{O}}_{top}(\varphi_i) \preceq [n^p \lambda_*^n],$$

where  $\lambda_* := \max\{\lambda_1, \ldots, \lambda_k\}$ , and p is defined as the maximum of the  $p_i$  such that  $\lambda_i = \lambda_*$ . Consequently, Equation (A.3) yields:

- if λ<sub>\*</sub> > λ, then *O*<sub>top</sub>(φ) ~ Σ<sub>i=1</sub><sup>k</sup> *O*<sub>top</sub>(φ<sub>i</sub>) and the latter is (λ<sub>\*</sub>, p)-tame;
  if λ<sub>\*</sub> ≤ λ, then *O*<sub>top</sub>(φ) ~ Σ<sub>i=1</sub><sup>k</sup> *O*<sub>top</sub>(φ<sub>i</sub>) + [Σ<sub>j=0</sub><sup>n</sup> N<sub>j</sub>λ<sup>n-j</sup>], where the latter growth rate satisfies [λ<sup>n</sup>] ≤ [Σ<sub>j=0</sub><sup>n</sup> N<sub>j</sub>λ<sup>n-j</sup>] ≤ [n<sup>p+1</sup>λ<sup>n</sup>]. In particular, rewriting this growth rate as [λ<sup>n</sup> · Σ<sub>j=0</sub><sup>n</sup> N<sub>j</sub>λ<sup>-j</sup>], we see that it is (λ, p + 1)-tame.

We are only left to show that  $\varphi$  is sound, namely that we have  $\mathcal{O}_{top}(\varphi) \preceq \overline{\mathfrak{o}}_{top}(\phi)$ . If  $g \in G$ is not conjugate into any of the  $G_i$ , and its conjugacy class does not have finite  $\phi$ -orbit, then Equation (A.2) shows that  $|\varphi^n(g)| \sim ||\phi^n(g)|| \leq \overline{\mathfrak{o}}_{top}(\phi)$ . If instead g is conjugate into some  $G_i$ , then  $|\varphi^n(g)| \preceq \overline{\mathcal{O}}_{\text{top}}(\varphi_i) \sim \overline{\mathfrak{o}}_{\text{top}}(\phi_i) \preceq \overline{\mathfrak{o}}_{\text{top}}(\phi)$ . In conclusion, we have  $|\varphi^n(g)| \preceq \overline{\mathfrak{o}}_{\text{top}}(\phi)$  for all  $g \in G$ , which yields  $\overline{\mathcal{O}}_{\text{top}}(\varphi) \preceq \overline{\mathfrak{o}}_{\text{top}}(\phi)$  as required.

**Part (2).** If  $\lambda_* < \lambda$ , then Equation (A.4) shows that  $\sum_{j=0}^n N_j \lambda^{n-j} \sim \lambda^n$ . Thus, Equation (A.2) implies that  $\|\phi^n(g)\| \sim \lambda^n$  for all elements  $g \in G$  not conjugate into any  $G_i$ , and whose conjugacy class is not preserved by a power of  $\phi$ . We thus have  $[\lambda^n] \in \mathfrak{g}(\phi)$  in this case. If instead  $\lambda_* \geq \lambda$ , then the assumptions of part (2) imply that  $[\lambda^n_*] \in \mathfrak{g}(\phi_i) \subseteq \mathfrak{g}(\phi)$  for some index *i*. Either way, recalling that  $\mu = \max\{\lambda_*, \lambda\}$ , we have  $[\mu^n] \in \mathfrak{g}(\phi)$ .

**Part (3).** Suppose that there exist a finite set  $\Lambda \subseteq \mathbb{R}_{>1}$  and  $P \in \mathbb{N}$  such that, for all indices *i* and all representatives  $\varphi'_i \in \operatorname{Aut}(G_i)$  of  $\phi_i \in \operatorname{Out}(G_i)$ , each growth rate in the union  $\mathcal{G}(\varphi'_i) \cup \mathfrak{g}(\phi_i)$  is either sub-polynomial or equal to  $[n^a \nu^n]$ , for some  $\nu \in \Lambda$  and some integer  $0 \leq a \leq P$ .

Our goal is to precisely estimate the growth rate  $\lfloor \|\phi^n(g)\| \rfloor$  for all elements  $g \in G$  not conjugate into any  $G_i$  and such that  $\|\phi^n(g)\| \not\sim 1$ . Equation (A.1) shows that  $\lfloor \|\phi^n(g)\| \rfloor$  is a sum of finitely many growth rates in  $\bigcup_i \mathcal{G}(\varphi_i)$  with the growth rate  $\lfloor \sum_{j=0}^n N_j \lambda^{n-j} \rfloor$ . Since a finite sum of pure growth rates equals the fastest among them, it suffices to show that either  $\sum_{j=0}^n N_j \lambda^{n-j} \sim \lambda^n$ , or  $\sum_{j=0}^n N_j \lambda^{n-j} \sim n^a \nu^n$  for some  $\nu \in \Lambda$  and an integer  $0 \leq a \leq P+1$ .

Recall that the integer  $N_{j+1}$  is defined as the average word length of a uniformly bounded number of elements of the form

$$a\psi(a)\ldots\psi^{j-1}(a)\cdot\psi^j(c)\cdot\psi^{j-1}(e)\ldots\psi(e)e,$$

with  $a, c, e \in G_i$  for some index *i*. We will thus need the following observation.

**Claim.** Consider some  $a, c, e \in G_i$  and set  $u_n := a\psi(a) \dots \psi^{n-1}(a) \cdot \psi^n(c) \cdot \psi^{n-1}(e) \dots \psi(e)e$ . Then the growth rate  $[|u_n| + |u_{n+1}|]$  is either sub-polynomial or equal to  $[n^a \nu^n]$  for some  $\nu \in \Lambda$  and an integer  $0 \le a \le P$ .

*Proof of claim.* Setting for simplicity  $e_n := \psi^{n-1}(e) \dots \psi(e)e$ , observe that we have

$$u_n^{-1}u_{n+1} = e_n^{-1}\psi^n (c^{-1}a\psi(c)e)e_n.$$

Recall that  $\psi|_{G_i}$  coincides with the automorphism  $\varphi_i \in \operatorname{Aut}(G_i)$ , and let  $\eta \in \operatorname{Aut}(G_i)$  be the automorphism defined by  $\eta(x) = e^{-1}\varphi_i(x)e$ . For every  $n \ge 1$ , we have  $\eta^n(x) = e_n^{-1}\varphi_i^n(x)e_n$ . Thus, setting  $w := c^{-1}a\psi(c)e$ , we can rewrite the above equality simply as

$$u_n^{-1}u_{n+1} = \eta^n(w).$$

As  $\eta$  is in the same outer class as  $\varphi_i$ , the hypotheses of part (3) guarantee that every growth rate in  $\mathcal{G}(\eta)$  is sub-polynomial or equal to  $[n^a \nu^n]$  for  $\nu \in \Lambda$  and  $0 \leq a \leq P$ . Thus, the same is true of the growth rate  $[|u_n^{-1}u_{n+1}|]$ .

If  $[|u_n^{-1}u_{n+1}|]$  is sub-polynomial, then a telescopic argument shows that  $[|u_n|]$  is sub-polynomial. Suppose instead that  $|u_n^{-1}u_{n+1}| \sim n^a \nu^n$ . Then, we similarly get  $|u_n| \leq \sum_{j=1}^n j^a \nu^j \leq n^a \nu^n$ , while the triangle inequality yields  $|u_n| + |u_{n+1}| \geq |u_n^{-1}u_{n+1}| \sim n^a \nu^n$ . In conclusion, we obtain the equality  $|u_n| + |u_{n+1}| \sim n^a \nu^n$ , as desired.

Now, the claim implies that the growth rate  $[N_n+N_{n+1}]$  is sub-polynomial or equal to  $[n^a\nu^n]$  with  $\nu \in \Lambda$  and  $0 \le a \le P$ . Carrying out the sum two terms at a time, this shows that  $\lambda^n \cdot \sum_{j=0}^n N_j \lambda^{-j}$  is  $\sim \lambda^n$  if  $\nu < \lambda$ , while it is  $\sim n^a \nu^n$  if  $\nu > \lambda$ , and finally  $\sim n^{a+1}\nu^n$  if  $\nu = \lambda$ .

This concludes the proof of part (3) and of the entire proposition.

The restriction to fully irreducible automorphisms in Proposition A.11 was meant to simplify the already technical proof. However it can be easily removed, which we do in the next corollary.

**Corollary A.12.** Let G be finitely generated and infinitely ended, with freely indecomposable free factors  $H_i$ . Consider  $\phi \in \text{Out}(G)$ , let  $\phi_i \in \text{Out}(H_i)$  be the restrictions of  $\phi$ , and let  $\varphi_i \in \text{Aut}(H_i)$ 

be representatives of the  $\phi_i$ . For each *i*, suppose that either  $\overline{\mathcal{O}}_{top}(\varphi_i)$  is sub-polynomial, or  $\varphi_i$  is  $(\lambda_i, p_i)$ -docile for some  $\lambda_i > 1$  and  $p_i \in \mathbb{N}$ . Then all the following hold.

- (1) Either  $\phi$  is represented by some  $\varphi \in \operatorname{Aut}(G)$  with sub-polynomial  $\overline{\mathcal{O}}_{\operatorname{top}}(\varphi)$ , or  $\phi$  is  $(\mu, q)$ docile for some  $\mu > 1$  and  $q \in \mathbb{N}$ , where  $\mu$  is either  $\max_i \lambda_i$  or a larger Perron number.
- (2) If  $[\lambda_i^n] \in \mathfrak{g}(\phi_i)$  whenever  $\lambda_i = \mu$ , then  $[\mu^n] \in \mathfrak{g}(\phi)$ .

Proof. Write  $G = H_1 * \cdots * H_{k'} * F_{m'}$  where the  $H_i$  are freely indecomposable. We can assume that  $k' \geq 1$  since all automorphisms of free groups satisfy the thesis (see e.g. [Lev09]). Defining  $\mathcal{F}'$  as the set of G-conjugates of the  $H_i$ , it is immediate that  $\mathcal{F}'$  is a  $\phi$ -invariant factor system. Up to raising  $\phi$  to a power, we have  $\phi \in \text{Out}(G, \mathcal{F}')$ . Up to further raising  $\phi$  to a power, an iterated application of Lemma A.10 yields a factor system  $\mathcal{F} > \mathcal{F}'$  such that  $\phi \in \text{Out}(G, \mathcal{F})$  and such that  $\phi$  is fully irreducible for  $(G, \mathcal{F})$ .

Arguing by induction on the Grushko rank of G, we can assume that the corollary holds for the restriction of  $\phi$  to each element of  $\mathcal{F}$ . If  $\mathcal{F}$  is non-sporadic, then the corollary was the content of Proposition A.11. If instead  $\mathcal{F}$  is sporadic, then G admits a  $\phi$ -invariant free splitting whose vertex groups are the elements of  $\mathcal{F}$ , and in this case the corollary follows from Proposition A.8.

## Appendix B. Accessibility and $\mathbb{R}$ -trees

This appendix is concerned with actions  $G \curvearrowright T$ , where G is a finitely presented group and T is an  $\mathbb{R}$ -tree. If T is BF-stable and G is accessible over arc-stabilisers, one can deduce strong consequences on the structure T. Above all, G-stabilisers of points of T are finitely generated, provided that intersections of arc-stabilisers are also finitely generated. Showing this latter fact is the main goal of this appendix. We will deduce it — with significant additional technical work — from classical Rips-Sela theory [RS94, GLP94, BF95, Sel97a] and the work of Guirardel [Gui98, Gui08]. We will often rely on the particularly clean treatment in [Gui08] for reference.

Before stating our main result, we need to introduce some terminology and notation. In Section 4, we gave a definition of quadratically hanging subgroups that implicitly restricts to the case with trivial fibre. In this appendix, we need to work with a more general notion that allows for nontrivial fibres (Definition B.1); this is similar but not identical to the classical notion [GL17, Definition 5.13]. For a compact surface  $\Sigma$ , we say that a subgroup  $P \leq \pi_1(\Sigma)$  is:

- peripheral if P is conjugate to a subgroup of the fundamental group of a component of  $\partial \Sigma$ ;
- a full peripheral if either  $P = \{1\}$  or P is conjugate to the (entire) fundamental group of a component of  $\partial \Sigma$ .

We only define optimal QH subgroups (cf. Definition 4.27), as this is all we use in this appendix.

**Definition B.1.** Let G be a group, let  $\mathcal{H}$  be a family of subgroups of G, and let  $F \leq G$  be a subgroup. A subgroup  $Q \leq G$  is an *optimal quadratically hanging subgroup with fibre* F, relative to  $\mathcal{H}$ , if there exists a simplicial G-tree  $G \curvearrowright S$  relative to  $\mathcal{H}$  with the following properties:

- (1) there exists a vertex  $x \in S$  such that Q is the G-stabiliser of x;
- (2) we have  $F \triangleleft Q$  and an identification  $Q/F \cong \pi_1(\Sigma)$  for a compact hyperbolic surface  $\Sigma$  other than the pair of pants;
- (3) for every edge  $e \subseteq S$  incident to x, we have  $F \triangleleft G_e$  and  $G_e/F$  is a full peripheral in  $\pi_1(\Sigma)$ ;
- (4) for each component  $B \subseteq \partial \Sigma$ , there is at most one Q-orbit of edges  $e \subseteq S$  incident to x such that  $G_e/F$  is conjugate to  $\pi_1(B)$  in  $\pi_1(\Sigma)$ ;
- (5) for every  $H \in \mathcal{H}$ , the projection of  $H \cap Q$  to Q/F is peripheral in  $\pi_1(\Sigma)$ .

We also say that x is an optimal QH vertex of S. A subgroup  $P \leq Q$  is a full peripheral subgroup of Q, if we have  $F \triangleleft P$  and P/F is a full peripheral subgroup of  $\pi_1(\Sigma)$ . A subgroup  $E \leq Q$  is essential if it is of the form  $F \rtimes \langle \gamma \rangle$ , where  $\gamma$  is a lift of an essential simple closed curve on the surface  $\Sigma$ .

Given another family  $\mathcal{F}$  of subgroups of G, we denote by  $QH(\mathcal{F}, \mathcal{H})$  the family of (optimal) quadratically hanging subgroups of G with fibre in  $\mathcal{F}$  relative to  $\mathcal{H}$ , and by  $Per(\mathcal{F}, \mathcal{H})$  and  $Ess(\mathcal{F}, \mathcal{H})$  their families of full peripheral and essential subgroups, respectively. Recall that  $\mathcal{F}_{int}$  denotes the family of finite intersections of subgroups in  $\mathcal{F}$ .

The following is the main result of this appendix. We emphasise that all our actions on  $\mathbb{R}$ -trees are implicitly assumed to be by isometries, unless explicitly stated otherwise.

**Theorem B.2.** Let G be finitely presented and torsion-free. Let  $G \curvearrowright T$  be a minimal action on an  $\mathbb{R}$ -tree that is not a single point. Let  $\mathcal{F}$  be the family of G-stabilisers of arcs of T, and let  $\mathscr{E}$  be the family of subgroups of G elliptic in T. Suppose that the following conditions are satisfied.

- (i) The group G is unconditionally accessible over  $\mathcal{F}_{int} \cup Ess(\mathcal{F}, \emptyset)$ .
- (ii) All elements of  $\mathcal{F}_{int}$  are finitely generated and root-closed.
- (iii) Chains of subgroups in  $\mathcal{F}_{int}$  have bounded length.

Then all of the following hold.

- (1) There are only finitely many G-conjugacy classes of G-stabilisers of stable arcs of T.
- (2) For every point  $p \in T$ , both of the following hold.
  - (a) The G-stabiliser  $G_p$  is finitely generated.
    - (b) If  $G_p \notin \mathcal{F} \cup \operatorname{Per}(\mathcal{F}, \mathscr{E})$ , then  $G_p$  coincides with the *G*-stabiliser of a vertex *x* in some  $(\mathcal{F}_{\operatorname{int}} \cup \operatorname{Per}(\mathcal{F}, \mathscr{E}), \mathscr{E})$ -splitting  $G \curvearrowright S$  with the following additional property. For every edge  $e \subseteq S$  incident to *x* such that  $G_e \notin \mathcal{F}_{\operatorname{int}}$ , the vertex of *e* other than *x* is an optimal *QH* vertex of *S* with fibre in  $\mathcal{F}$  relative to  $\mathscr{E}$ .
- (3) There are only finitely many G-orbits of points  $p \in T$  such that  $G_p \notin \mathcal{F}$ .
- (4) If a subgroup  $H \leq G$  is not elliptic in T, then H is non-elliptic in a  $(\mathcal{F}_{int} \cup \operatorname{Ess}(\mathcal{F}, \mathscr{E}), \mathscr{E})$ -splitting of G.

In this article, we only use the theorem through its main consequence for special groups:

**Corollary B.3.** Let G be a special group. Let  $G \curvearrowright T$  be a minimal action on an  $\mathbb{R}$ -tree such that all arc-stabilisers lie in  $\mathcal{Z}(G)$ . Let  $\mathscr{E}$  be the family of subgroups elliptic in T. Then:

- (1) for every  $p \in T$ , the stabiliser  $G_p$  is convex-cocompact;
- (2) there are only finitely many G-conjugacy classes of point-stabilisers of T;
- (3) if  $H \leq G$  is not elliptic in T, then H is non-elliptic in a  $(\mathcal{Z}(G), \mathscr{E})$ -splitting of G;
- (4) for ever  $p \in T$ , either  $G_p$  is trivial, or  $G_p \in \mathcal{Z}_c(G)$ , or  $G_p$  is a subgroup of an element of  $\mathcal{S}(G)$ , or finally  $G_p$  is a vertex group in a  $(\mathcal{Z}(G) \cup \operatorname{Per}(\mathcal{Z}(G), \mathscr{E}), \mathscr{E})$ -splitting of G.

*Proof.* Note that the action  $G \curvearrowright T$  satisfies the hypothesis of Theorem B.2. Indeed, we have  $\operatorname{Ess}(\mathcal{Z}(G), \emptyset) \subseteq \mathcal{Z}(G)$ , as an essential subgroup of the form  $F \rtimes \langle \gamma \rangle$  coincides with the centraliser  $Z_G(\gamma)$ . Moreover, G is unconditionally accessible over  $\mathcal{Z}(G)$  by Theorem 3.1, and chains of centralisers have bounded length by Remark 2.11.

Now, part (3) of the corollary is immediate from Theorem B.2(4). Part (4) of the corollary also directly follows from Theorem B.2(2b), observing that nontrivial elements of  $\mathcal{Z}(G) \cup \text{Per}(\mathcal{Z}(G), \mathscr{E})$ either lie in  $\mathcal{Z}_c(G)$ , or are contained in elements of  $\mathcal{S}(G)$ . Indeed, if  $P \in \text{Per}(\mathcal{Z}(G), \mathscr{E})$  and  $F \in \mathcal{Z}(G)$ is the fibre of the corresponding QH subgroup, then either  $F \neq \{1\}$  (in which case P normalises F, and so either  $P = F \in \mathcal{Z}_c(G)$  or P is contained in an element of  $\mathcal{S}(G)$  by Lemma 4.6(3)), or  $F = \{1\}$  (in which case, P is cyclic, so  $P \leq Z_G(P) \in \mathcal{Z}(G)$ ).

Regarding part (2), Theorem B.2(3) shows that there are only finitely many conjugacy classes of point-stabilisers of T that are not arc-stabilisers, so it suffices to show that arc-stabilisers of Talso fall into only finitely many conjugacy classes. Consider an arc  $[x, y] \subseteq T$ . Since chains in  $\mathcal{Z}(G)$ have bounded length, there exist two points  $x', y' \in (x, y)$  such that the arcs [x, y] and [x', y'] have the same G-stabiliser (with the points x, x', y', y so aligned). Choosing stable arcs  $\beta_x \subseteq [x, x']$  and  $\beta_y \subseteq [y', y]$ , we see that the stabiliser of [x, y] coincides with the intersection of the stabilisers of  $\beta_x$  and  $\beta_y$ . Now, there are only finitely many conjugacy classes of stabilisers of stable arcs, by Theorem B.2(1), and so Lemma 2.4(5) implies that there are only finitely many conjugacy classes of (general) arc-stabilisers, as we wanted.

Finally, part (1) of the corollary follows from Theorem B.2(2b) as we briefly discuss. If  $p \in T$ and  $G_p \notin \mathcal{Z}(G) \cup \operatorname{Per}(\mathcal{Z}(G), \mathscr{E})$ , then  $G_p$  is a vertex group of a particular splitting of G over  $\mathcal{Z}(G) \cup \operatorname{Per}(\mathcal{Z}(G), \mathscr{E})$ . This implies that  $G_p$  is convex-cocompact, appealing to Proposition 2.30, and using Lemma 4.29 and the particular structure of the splitting to work around the fact that the subgroups in  $\operatorname{Per}(\mathcal{Z}(G), \mathscr{E})$  might not be convex-cocompact. Instead, we can use the fact that all elements of  $\operatorname{Ess}(\mathcal{Z}(G), \mathscr{E})$  are convex-cocompact (they are centralisers).

We are left to consider the case when  $G_p \in \mathcal{Z}(G) \cup \operatorname{Per}(\mathcal{Z}(G), \mathscr{E})$ . If  $G_p \in \mathcal{Z}(G)$ , it is automatically convex-cocompact. Not all elements of  $\operatorname{Per}(\mathcal{Z}(G), \mathscr{E})$  are convex-cocompact, but those that can arise as  $G_p$  actually are. This requires looking at the proof of Theorem B.2(2b) at the end of this appendix: we can only have  $G_p \in \operatorname{Per}(\mathcal{Z}(G), \mathscr{E}) \setminus \mathcal{Z}(G)$  if there is a splitting  $G \curvearrowright T$  (over  $\mathcal{Z}(G)$ , relative to  $\mathscr{E}$ ) with a QH vertex group Q such that  $G_p$  is a full peripheral subgroup of Q and  $G_p$ is not commensurable to the G-stabiliser of any incident edge of T; in this case, Lemma 4.29 again shows that  $G_p$  is convex-cocompact.

We now start working towards the proof of Theorem B.2, which will occupy the rest of this appendix.

Since G is finitely presented, there exists a sequence  $G \curvearrowright \mathcal{G}_n$  of geometric  $\mathbb{R}$ -trees converging strongly to T (in the sense of [LP97]). In particular, this means that there are G-equivariant morphisms  $f_n: \mathcal{G}_n \to T$  (continuous maps such that each arc of  $\mathcal{G}_n$  can be subdivided into finitely many arcs on which  $f_n$  is isometric), and also equivariant morphisms  $f_n^m: \mathcal{G}_n \to \mathcal{G}_m$  for m > n such that  $f_n = f_m \circ f_n^m$ . We denote by  $\mathcal{E}_n$  the family of subgroups of G that are elliptic in  $\mathcal{G}_n$ , and observe that  $\mathcal{E}_n \subseteq \mathcal{E}$  and  $\mathcal{E}_n \subseteq \mathcal{E}_m$  for m > n. Strong convergence implies that, in particular, any finitely generated element of  $\mathcal{E}$  eventually lies in  $\mathcal{E}_n$ .

Let  $G \curvearrowright \mathcal{D}_n$  be the simplicial tree dual to the decomposition of  $\mathcal{G}_n$  into indecomposable components and simplicial arcs [Gui08, Proposition 1.25]. The tree  $\mathcal{D}_n$  is bipartite: it has a black vertex for each indecomposable component of  $\mathcal{G}_n$ , as well as for each maximal arc of  $\mathcal{G}_n$  containing no branch points in its interior (a "simplicial arc"); it has a white vertex for every point of intersection between subtrees of  $\mathcal{G}_n$  associated to black vertices; edges correspond to point-subtree inclusions. The tree  $G \curvearrowright \mathcal{D}_n$  is minimal and relative to  $\mathscr{E}_n$ . If  $U \subseteq \mathcal{G}_n$  is an indecomposable component, we denote by  $\mathcal{G}_U$  its G-stabiliser, and by [U] the corresponding black vertex of  $\mathcal{D}_n$ . Note that, if  $e \subseteq \mathcal{D}_n$  is an edge incident to [U], then the G-stabiliser of e is the  $\mathcal{G}_U$ -stabiliser of a point of U.

Observe that the action  $G \curvearrowright T$  is BF-stable because of Condition (*iii*): every arc of T contains a stable sub-arc. If  $U \subseteq \mathcal{G}_n$  is an indecomposable component, it follows that the image  $f_n(U) \subseteq T$ is a stable subtree (see [Gui08, Section 1.6]). If  $K_U$  denotes the G-stabiliser of one/all arcs of  $f_n(U) \subseteq T$ , we have  $G_U \leq N_G(K_U)$  and  $K_U \in \mathcal{F}$ . In general, arc-stabilisers of  $U \subseteq \mathcal{G}_n$  are proper subgroups of  $K_U$ . We will say that U is saturated if  $K_U$  fixes U pointwise, that is, if  $K_U$  is the kernel of the action  $G_U \curvearrowright U$ , as well as the G-stabiliser of every arc of U.

If  $U \subseteq \mathcal{G}_n$  is saturated, we obtain an indecomposable action  $G_U/K_U \curvearrowright U$  with trivial arcstabilisers. The output of the Rips machine now shows that U is of one of three possible types — axial, exotic, or surface; see e.g. Remark 1.29 and Proposition A.6 in [Gui08]. We will use the following facts about these three types:

• If U is axial, then U is isometric to  $\mathbb{R}$  and  $G_U/K_U$  is free abelian of rank  $\geq 2$  (note that  $G_U/K_U$  is torsion-free because  $K_U \in \mathcal{F}$  is root-closed by Condition (*ii*)). Moreover,  $K_U$  is the G-stabiliser of all edges of  $\mathcal{D}_n$  incident to the black vertex [U].

- If U is exotic, then  $G_U/K_U$  admits a (simplicial) free splitting with the same elliptic subgroups as the action  $G_U/K_U \curvearrowright U$  [Gui98, Proposition 7.2]. In particular,  $G_U$  splits over  $K_U$  relative to the G-stabilisers of the edges of  $\mathcal{D}_n$  incident to the black vertex [U]. We can then use this splitting of  $G_U$  to refine the splitting  $G \curvearrowright \mathcal{D}_n$  (Lemma 4.13), if we so desire.
- If U is of surface type, then the action  $G_U/K_U \curvearrowright U$  is dual to an arational measured foliation on a compact hyperbolic surface  $\Sigma_U$  of which  $G_U/K_U$  is the fundamental group. Non-trivial point-stabilisers of  $G_U/K_U \curvearrowright U$  are maximal cyclic subgroups of  $G_U/K_U$ , namely the conjugates of the fundamental groups of the components of  $\partial \Sigma_U$ . The G-stabiliser of each edge of  $\mathcal{D}_n$  incident to [U] is either equal  $K_U$  or to the extension of  $K_U$  by one of these cyclic subgroups. In particular, [U] is an optimal quadratically hanging vertex of  $\mathcal{D}_n$  with fibre  $K_U$  (Definition B.1). Note that  $\Sigma_U$  is indeed not a pair of pants, as it supports an arational measured foliation.

Any essential simple closed curve  $\gamma \subseteq \Sigma_U$  gives a splitting of  $G_U/K_U$  over the maximal cyclic subgroup  $C := \langle \gamma \rangle$ , relative to the fundamental groups of the components of  $\partial \Sigma_U$ . In turn, this gives a splitting of  $G_U$  over the essential subgroup  $K_U \rtimes \langle \gamma \rangle$ , relative to the G-stabilisers of the edges of  $\mathcal{D}_n$  incident to the black vertex [U]. Again, we can use this splitting of  $G_U$  to refine  $G \curvearrowright \mathcal{D}_n$ , if we so desire.

Summing up, any saturated indecomposable component of exotic or surface type  $U \subseteq \mathcal{G}_n$  determines a particular splitting of G over an element of  $\mathcal{F}$  or  $\operatorname{Ess}(\mathcal{F}, \mathcal{E}_n)$ , relative to  $\mathcal{E}_n$ . This splitting is obtained by first refining  $\mathcal{D}_n$  at the vertex [U] and then collapsing (most) edges of  $\mathcal{D}_n$ .

We can now use accessibility (as provided by Condition (i)) to prove part (1) of Theorem B.2.

**Lemma B.4.** There are only finitely many G-conjugacy classes of G-stabilisers of stable arcs of T. Moreover, there exists an integer N such that each tree  $\mathcal{G}_n$  has at most N G-orbits of saturated indecomposable components.

Proof. Let N be the largest number of edge orbits in an irredundant splitting of G over  $\mathcal{F} \cup$ Ess $(\mathcal{F}, \mathscr{E}_n)$ , which exists by Condition (i). Suppose for the sake of contradiction that there exist N + 1 stable arcs  $\overline{\beta}_0, \ldots, \overline{\beta}_N \subseteq T$  whose G-stabilisers  $B_0, \ldots, B_N$  are pairwise not G-conjugate. Since each  $B_i$  is finitely generated, by Condition (ii), there exists  $n \in \mathbb{N}$  such that the stable arcs  $\overline{\beta}_i \subseteq T$  all lift isometrically to (stable) arcs  $\beta_i \subseteq \mathcal{G}_n$  with  $B_i$  as their G-stabiliser (this follows from the definition of strong convergence). Each  $\beta_i$  shares an arc with a subset  $U_i \subseteq \mathcal{G}_n$  that is either an indecomposable component or a maximal simplicial arc.

Now, consider the splitting  $G \curvearrowright \mathcal{D}_n$ . Form a new splitting  $G \curvearrowright \mathcal{D}'_n$  by blowing up, for each indecomposable  $U_i$  of exotic or surface type, the vertex  $[U_i] \in \mathcal{D}_n$  to a one-edge splitting of  $G_{U_i}$ over an element of  $\mathcal{F} \cup \text{Ess}(\mathcal{F}, \mathscr{E}_n)$ , as described above. Then form a third splitting  $G \curvearrowright \mathcal{D}''_n$  by collapsing all G-orbits of edges of  $\mathcal{D}'_n$ , except for those created in the blow-up, and except for the G-orbit of one edge incident to each of the vertices  $[U_i]$  of axial or simplicial type. The result is that  $G \curvearrowright \mathcal{D}''_n$  is a splitting over  $\mathcal{F} \cup \text{Ess}(\mathcal{F}, \mathscr{E}_n)$  with exactly N + 1 orbits of edges. Moreover,  $\mathcal{D}''_n$  is irredundant since the  $B_i$  are pairwise not G-conjugate (unless N = 0 and  $\mathcal{D}''_n \cong \mathbb{R}$ ). This violates accessibility, providing the required contradiction and proving the first half of the lemma.

The second half is similar: Suppose that some  $\mathcal{G}_n$  has 2N + 1 saturated indecomposable components  $U_0, \ldots, U_{2N}$  in pairwise distinct G-orbits. We again consider  $\mathcal{D}_n$  and inflate it to a tree  $\mathcal{D}'_n$  by blowing up each vertex  $[U_i]$  of exotic or surface type to a one-edge splitting of  $G_{U_i}$  over an element of  $\mathcal{F} \cup \mathrm{Ess}(\mathcal{F}, \mathscr{E}_n)$ . Note that the G-stabiliser of each new edge of  $\mathcal{D}'_n$  is properly contained in the G-stabiliser of both incident vertices. Then, we form a splitting  $\mathcal{D}''_n$  by collapsing all edges of  $\mathcal{D}'_n$  coming from edges of  $\mathcal{D}_n$ , except for the G-orbit of one edge incident to each  $[U_i]$  of axial type. Again,  $G \curvearrowright \mathcal{D}''_n$  is a splitting over  $\mathcal{F} \cup \mathrm{Ess}(\mathcal{F}, \mathscr{E}_n)$  with exactly 2N + 1 orbits of edges. Any failure of  $\mathcal{D}''_n$  to be irredundant comes from a degree-2 vertex whose two incident edges e, f have the same G-stabiliser; this can only happen if the other vertices of e and f are, respectively, G-translates of some  $[U_i], [U_j]$  with  $i \neq j$  and both  $U_i, U_j$  axial (with *G*-conjugate kernels). In particular, the edges e and f are in distinct *G*-orbits (as  $\mathcal{D}''_n \not\cong \mathbb{R}$ ) and the problem is solved by collapsing the *G*-orbit of one of them. Ultimately, we obtain an irredundant splitting of G over  $\mathcal{F} \cup \mathrm{Ess}(\mathcal{F}, \mathscr{E}_n)$  with at least N + 1 edge orbits, violating again accessibility.

Using the previous lemma and Guirardel's work in [Gui08, Appendix A], we can now conclude that the indecomposable components of the geometric approximations  $\mathcal{G}_n$  stabilise for large n.

**Lemma B.5.** There exists  $n_0 \in \mathbb{N}$  such that all the following hold for all  $n \geq n_0$ .

- (1) Every indecomposable component  $U \subseteq \mathcal{G}_n$  is saturated.
- (2) For all m > n and each indecomposable component  $U \subseteq \mathcal{G}_n$ , the image  $f_n^m(U) \subseteq \mathcal{G}_m$ is an indecomposable component of  $\mathcal{G}_m$ , and the map  $f_n^m|_U \colon U \to f_n^m(U)$  is an isometry. Moreover, the G-stabilisers of U and  $f_n^m(U)$  coincide.
- (3) For all m > n, each indecomposable component of  $\mathcal{G}_m$  is the image under  $f_n^m$  of a (unique) indecomposable component of  $\mathcal{G}_n$ .

Proof. For all integers m > n and each indecomposable component  $U \subseteq \mathcal{G}_n$ , the image  $f_n^m(U) \subseteq \mathcal{G}_m$ is indecomposable and, therefore, there exists an indecomposable component  $V_m \subseteq \mathcal{G}_m$  such that  $f_n^m(U) \subseteq V_m$  and  $G_U \leq G_{V_m}$  (see again [Gui08, Section 1.6]). We have already seen that the image  $f_n(U) \subseteq T$  is indecomposable and that all its arcs are stable with the same G-stabiliser  $K_U \in \mathcal{F}$ , which is finitely generated by Condition (*ii*). It follows that, for large enough m, an arc of  $f^n(U) \subseteq T$  lifts to an arc of  $f_n^m(U) \subseteq \mathcal{G}_m$  with precisely  $K_U$  as its G-stabiliser. As a consequence, the indecomposable component  $V_m \subseteq \mathcal{G}_m$  is saturated. Summing up, for every n and each indecomposable component  $U \subseteq \mathcal{G}_n$ , the image  $f_n^m(U)$  is contained in a saturated indecomposable component  $V_m \subseteq \mathcal{G}_m$  for all sufficiently large m (depending on n and U).

Now, suppose that  $U \subseteq \mathcal{G}_n$  is saturated to begin with, so that we obtain an indecomposable action with trivial arc-stabilisers  $G_U/K_U \curvearrowright U$ . By [Gui08, Lemma A.7], there exists an integer  $n_U \ge n$  such that, for all  $m \ge k \ge n_U$ , the map  $f_k^m|_{V_k} : V_k \to V_m$  is an isometry (the hypotheses in Guirardel's lemma are slightly different from ours, but the key point is the same, namely [Gui08, Theorem A.11]). In addition, Guirardel's lemma shows that either  $V_k$  is axial or  $G_{V_k} = G_{V_m}$ . In fact, in our situation, we cannot have<sup>23</sup> a proper inclusion  $G_{V_k} \le G_{V_m}$  even when  $V_k$  is axial: indeed,  $V_m$ would also be axial, so the quotient  $G_{V_m}/K_U$  would be abelian, which would imply that  $G_{V_k} \triangleleft G_{V_m}$ ; hence, looking at the action  $G \curvearrowright \mathcal{G}_k$ , it would follow that  $G_{V_m}$  preserves the line  $V_k$ .

Summing up, for each indecomposable component  $U \subseteq \mathcal{G}_n$ , there exists an integer  $n_U$  such that parts (1) and (2) of the lemma hold for  $m \ge n_U$ , for the indecomposable components of the  $\mathcal{G}_m$ containing  $f_n^m(U)$ . Since the number of G-orbits of saturated indecomposable components in the  $\mathcal{G}_n$  is uniformly bounded by Lemma B.4, we get a uniform upper bounded to the required integers  $n_U$ , proving parts (1) and (2) of the lemma. Part (3) then immediately follows from this.

Up to discarding finitely many  $\mathcal{G}_n$ , we assume in the coming discussion that  $n_0 = 0$ . Denoting by  $\mathcal{F}_{stab} \subseteq \mathcal{F}$  the family of *G*-stabilisers of stable arcs of *T*, and recalling that  $\mathcal{F}_{stab}$  consists of finitely many *G*-conjugacy classes of finitely generated subgroups by Lemma B.4 and Condition (*ii*), we can similarly assume that all elements of  $\mathcal{F}_{stab}$  are elliptic in all  $\mathcal{G}_n$ . Since *T* is BF-stable, every element of  $\mathcal{F}$  is contained in an element of  $\mathcal{F}_{stab}$ , and it follows that  $\mathcal{F} \subseteq \mathscr{E}_n$  for all *n*.

Denote by Ind be the family of subgroups of G arising as stabilisers of indecomposable components of the  $\mathcal{G}_n$ ; by Lemma B.4, Ind also consists of finitely many G-conjugacy classes of subgroups. For each  $I \in$  Ind, we denote by  $\partial I$  the family of subgroups of I arising as (entire) point-stabilisers in the action  $I \curvearrowright U$ , where U is the indecomposable component associated to I in one/all  $\mathcal{G}_n$  (by

<sup>&</sup>lt;sup>23</sup>The situation for axial components is more delicate in Guirardel's paper because the geometric approximations are acted upon by different groups there, as they arise from an action on an  $\mathbb{R}$ -tree of a possibly non-finitely-presented group G. In our setting, axial components do not need to be treated as an exception.

Lemma B.5, the value of n plays no role). We denote by  $K_I \triangleleft I$  the kernel of the action  $I \curvearrowright U$  (also denoted  $K_U$  above). Finally, we write  $\partial \text{Ind} := \bigcup_{I \in \text{Ind}} \partial I$ .

It is convenient to give a name to the following type of splittings of G.

**Definition B.6.** A splitting  $G \curvearrowright \Delta$  is *excellent* if it satisfies the following properties:

- (1) for each  $I \in \text{Ind}$ , there exists a vertex  $x_I \in \Delta$  such that the *G*-stabiliser of  $x_I$  is *I*, the *G*-stabiliser of every edge of  $\Delta$  incident to  $x_I$  lies in  $\partial I$ , and at most one edge incident to  $x_I$  has a given element of  $\partial I \setminus \{K_I\}$  as its stabiliser;
- (2) if an edge  $e \subseteq \Delta$  is not incident to  $x_I$  for any  $I \in \text{Ind}$ , then the *G*-stabiliser of *e* lies in  $\mathcal{F}_{\text{int}}$ ;
- (3) all subgroups in  $\mathscr{E}$  are elliptic in  $\Delta$ ;
- (4) if a subgroup of G is elliptic in  $\Delta$ , then it lies in  $\mathscr{E}$  or is contained in an element of Ind.

Our next goal is showing that G indeed admits an excellent splitting. For this, we first need the following observation.

**Lemma B.7.** Let  $G \curvearrowright S$  be a splitting relative to  $\mathcal{F}$  satisfying Items (1) and (2) in Definition B.6. For any vertex  $x \in S$  with  $G_x \notin$  Ind and any subgroup  $F \in \mathcal{F}_{int}$ , we have  $G_x \cap F \in \mathcal{F}_{int}$ .

*Proof.* Set  $\Omega := G_x \cap F$  for simplicity. Since S is a splitting relative to  $\mathcal{F}$ , there exists a vertex  $y \in S$  fixed by F. If  $F \leq G_x$ , then  $\Omega$  coincides with F and lies in  $\mathcal{F}_{int}$ . Suppose instead that  $F \not\leq G_x$  in the rest of the proof. Thus, we have  $y \neq x$  and, letting  $e \subseteq S$  be the edge incident to x in the direction of y, we have  $\Omega = G_e \cap F$ . If  $G_e \in \mathcal{F}_{int}$ , we again have  $\Omega \in \mathcal{F}_{int}$ .

Suppose instead that  $G_e \notin \mathcal{F}_{int}$ . Let z be the vertex of e other than x, and note that its Gstabiliser must be some  $I \in Ind$  (as  $G_x \notin Ind$  by hypothesis). Let  $I \cap U$  be the corresponding
indecomposable component of the  $\mathcal{G}_n$ , and let  $\overline{e} \in U$  be a point with  $G_e$  as its I-stabiliser. If y = z, then  $F \leq I$  and, since we have assumed that  $\mathcal{F} \subseteq \mathscr{E}_n$  for all n, there is a point of U
fixed by F; this point must be distinct from  $\overline{e}$  (otherwise we would have  $F \leq G_e \leq G_x$ ), and this
implies that  $\Omega = G_e \cap F = K_I \cap F \in \mathcal{F}_{int}$ . If instead  $y \neq z$ , let  $f \subseteq S$  be the edge incident to z in the direction of y. Again, there exists a point  $\overline{f} \in U$  with  $G_f$  as its I-stabiliser, and hence  $\Omega = F \cap (G_e \cap G_f) = F \cap K_I \in \mathcal{F}_{int}$ , as we wanted.

We can now prove that excellent splittings indeed exist, provided that the  $\mathcal{G}_n$  are not indecomposable. We will quickly be able to complete the proof of Theorem B.2 once this is shown.

**Proposition B.8.** If  $G \notin \text{Ind}$ , there exists an excellent splitting  $G \curvearrowright \Delta$ .

*Proof.* Let  $\mathscr{S}$  be the family of splittings of G that satisfy Items (1) and (2) in Definition B.6. For each  $k \in \mathbb{N}$ , let  $\mathscr{S}_k \subseteq \mathscr{S}$  be the subset of splittings relative to  $\mathscr{E}_k$ . We have  $\mathscr{S}_k \supseteq \mathscr{S}_m$  for m > k, as  $\mathscr{E}_k \subseteq \mathscr{E}_m$ .

Observe that  $\mathscr{S}_k \neq \emptyset$  for all  $k \in \mathbb{N}$ . For instance, we can consider the tree  $G \curvearrowright \mathcal{D}_k$  defined above, which is not a single vertex because  $G \notin$  Ind. We can then collapse all G-orbits of edges of  $\mathcal{D}_k$  that are not incident to any black vertex of  $\mathcal{D}_k$  representing an indecomposable component of  $\mathcal{G}_k$ . This collapse lies in  $\mathscr{S}_k$ , showing that  $\mathscr{S}_k \neq \emptyset$ .

In order to progress with the proof of the proposition, we will need the following observation.

**Claim 1.** For any two splittings  $S_1, S_2 \in \mathscr{S}_k$ , there exists a third splitting  $S \in \mathscr{S}_k$  that refines  $S_1$  and dominates  $S_2$ .

Proof of Claim 1. This essentially follows from Lemma 4.15, except that we need to check that the refinement of  $S_1$  constructed there truly lies in  $\mathscr{S}$ . For this, consider a vertex  $x \in S_1$  such that  $G_x$  is not elliptic in  $S_2$ , and note that this implies that  $G_x \notin \text{Ind.}$  let  $M \subseteq S_2$  be the  $G_x$ -minimal subtree. We claim that there exists an edge of M whose  $G_x$ -stabiliser lies in  $\mathcal{F}_{\text{int.}}$ 

If there exists an edge  $e \subseteq M$  such that  $G_e \in \mathcal{F}_{int}$ , then the  $G_x$ -stabiliser of e lies in  $\mathcal{F}_{int}$  by Lemma B.7. Thus, we can suppose that no edge of M has G-stabiliser in  $\mathcal{F}_{int}$ . Considering any

edge  $e \subseteq M$ , this implies that there exists a vertex  $y \in e$  whose G-stabiliser is some  $I \in Ind$ . Note that there also exists an edge  $f \subseteq M$  incident to y with  $f \neq e$ . Let  $I \curvearrowright U$  be the indecomposable component associated to I. If y' is the vertex of  $S_1$  fixed by I, we have  $y' \neq x$ , as we have seen above that  $G_x \notin Ind$ . Now, let  $g \subseteq S_1$  be the edge incident to y' in the direction of x. The G-stabilisers of the edges e, f, g equal the I-stabilisers of three points of U, and at least one of the two subgroups  $G_e$  and  $G_f$  must be different from  $G_g$ , unless they all equal  $K_I$  (recall Item (1) in Definition B.6). Thus, assuming without loss of generality that  $G_e \neq G_g$ , we have  $G_e \cap G_g = K_I \in \mathcal{F}$ . It follows that the  $G_x$ -stabiliser of the edge  $e \subseteq M$  equals  $G_x \cap (G_e \cap G_g) \in \mathcal{F}_{int}$  by Lemma B.7.

In conclusion, we have shown that there always exists an edge  $e \subseteq M$  whose  $G_x$ -stabiliser lies in  $\mathcal{F}_{int}$ . Collapsing all other  $G_x$ -orbits of edges of M, we obtain a one-edge splitting of  $G_x \curvearrowright M'$ . We can then blow up the vertex  $x \in S_1$  to a copy of M', thus obtaining another splitting of G in  $\mathscr{S}_k$ . We can only repeat this procedure a finite number of times, as G is unconditionally accessible over  $\mathcal{F}_{int}$  by Condition (i). Thus, we eventually obtain a refinement of  $S_1$  that lies in  $\mathscr{S}_k$  and dominates  $S_2$  (see the proof of Lemma 4.15 for details).

Now observe that, for each  $k \in \mathbb{N}$ , there exists a splitting  $\Delta_k \in \mathscr{S}_k$  that dominates all other splittings in  $\mathscr{S}_k$ . This is proved exactly as in Lemma 4.16(1), using the fact that G is unconditionally accessible over  $\mathcal{F}_{int}$ , and using Claim 1 in place of Lemma 4.15 (we cannot directly use Lemma 4.16, as some edge-stabilisers of the splittings in  $\mathscr{S}$  do not lie in  $\mathcal{F}_{int}$ ).

We now show that, for each  $k \in \mathbb{N}$ , the splitting  $G \curvearrowright \Delta_k$  satisfies Item (4) in Definition B.6.

**Claim 2.** If  $x \in \Delta_k$  is a vertex such that  $G_x \notin \text{Ind}$ , then  $G_x$  is elliptic in the  $\mathbb{R}$ -tree T.

Proof of Claim 2. Observe that the G-stabilisers of the edges of  $\Delta_k$  incident to x lie in  $\partial \operatorname{Ind} \cup \mathcal{F}_{\operatorname{int}}$ ; these subgroups are elliptic in T and in  $\mathcal{G}_n$  for all n. Suppose for the sake of contradiction that  $G_x$  is not elliptic in T.

We will show that, for all sufficiently large integers n, the group  $G_x$  splits over an element of  $\mathcal{F}_{int}$  that it contains, relative to  $\mathscr{E}_n$ . This will imply that  $G \curvearrowright \Delta_k$  can be refined (using Lemma 4.13(1)) into a splitting that still lies in  $\mathscr{S}_k$  and in which  $G_x$  is no longer elliptic, which will contradict the fact that  $\Delta_k$  dominates all splittings in  $\mathscr{S}_k$ .

Now, note that  $G_x$  is finitely generated relative to finitely many elements of  $\partial \operatorname{Ind} \cup \mathcal{F}_{\operatorname{int}}$  by [Gui08, Lemma 1.11]. Thus, the fact that  $G_x$  is not elliptic in T implies that  $G_x$  contains an element  $g \in G_x$ that is loxodromic in T, and hence loxodromic in  $\mathcal{G}_n$  for all n. Let  $\alpha \subseteq T$  and  $\alpha_n \subseteq \mathcal{G}_n$  be the axes of g, and choose a stable arc  $\beta \subseteq \alpha$ . For large n, the arc  $\beta$  lifts to an arc  $\beta_n \subseteq \alpha_n$  with the same G-stabiliser as  $\beta$  (as the elements of  $\mathcal{F}$  are finitely generated). Up to modifying  $\beta_n$  we can assume that either  $\beta_n$  is a maximal simplicial arc of  $\mathcal{G}_n$  or  $\beta_n$  is an arc of some indecomposable component  $U \subseteq \mathcal{G}_n$ . In the former case, we can collapse to a point every arc of  $\mathcal{G}_n$  whose G-translates are all disjoint from the interior of  $\beta_n$ , thus obtaining a one-edge splitting of G over the conjugates of  $G_\beta \in \mathcal{F}$ , in which the elements of  $\mathcal{G}_n$  in this case.

Thus, suppose that the axis  $\alpha_n \subseteq \mathcal{G}_n$  shares an arc with an indecomposable component  $U \subseteq \mathcal{G}_n$ . Note that the intersection  $G_x \cap G_U$  is elliptic in all trees  $\mathcal{G}_m$  (as x is a vertex of a splitting in  $\mathscr{S}$ ). In particular, there is at most one point  $z \in U$  such that the intersection between  $G_x$  and the  $G_U$ -stabiliser of z is not contained in the kernel  $K_U$ . Now, consider the action  $G \curvearrowright \mathcal{D}_n$  and let  $e, f \subseteq \mathcal{D}_n$  be the two edges incident to [U] corresponding to the two points where  $\alpha_n$  exits U. By the previous discussion, up to swapping e and f, we can assume that  $G_x \cap G_e \leq K_U \in \mathcal{F}_{int}$ . It follows that,  $G_x \cap G_e = G_x \cap K_U \in \mathcal{F}_{int}$  by Lemma B.7. In conclusion, after collapsing all edges of  $\mathcal{D}_n$  outside the G-orbit of e, we again obtain a splitting of G over an element of  $\mathcal{F}_{int}$ , in which the elements of  $\mathscr{E}_n$  are elliptic and  $G_x$  is not. This induces the desired splitting of  $G_x$  and, as discussed at the start, it proves the claim. Finally, observe that the splitting  $G \curvearrowright \Delta_k$  dominates the splitting  $G \curvearrowright \Delta_m$  for all m > k, as we have  $\mathscr{S}_k \supseteq \mathscr{S}_m$ . Using again Claim 1 and the fact that G is unconditionally accessible over  $\mathcal{F}_{int}$ , as in the proof of Lemma 4.16(2), we conclude that the deformation spaces of the trees  $\Delta_n$  must stabilise for large n.

In conclusion, setting  $\Delta := \Delta_k$  for a sufficiently large value of k, we obtain a splitting that satisfies Items (1), (2) and (4) of Definition B.6 and the following slightly weaker form of Item (3): all subgroups in the union  $\bigcup_{n \in \mathbb{N}} \mathscr{E}_n$  are elliptic in  $\Delta$ .

In order to conclude that  $\Delta$  is indeed excellent, consider some  $E \in \mathscr{E}$ . Each finitely generated subgroup of E lies in  $\mathscr{E}_n$  for large n, so all finitely generated subgroups of E are elliptic in  $\Delta$ . Since chains in  $\mathcal{F}_{int}$  are bounded by Condition (*iii*), it follows that E is itself elliptic in  $\Delta$  (see Lemma 2.29). This shows that  $G \curvearrowright \Delta$  is excellent, proving the proposition.

Before we continue, it is important to observe that an excellent splitting can be modified (near the vertices fixed by exotic elements of Ind) so as to make its edge groups finitely generated.

**Remark B.9.** Let  $G \curvearrowright \Delta$  be an excellent splitting. Consider an edge  $e \subseteq \Delta$ . If  $G_e \in \mathcal{F}_{int}$ , then  $G_e$  is finitely generated by Condition (*ii*). If  $G_e \notin \mathcal{F}_{int}$ , then exists a vertex  $y \in e$  such that  $G_y$  equals some  $I \in Ind$  and  $G_e \in \partial I$ . If I is of axial type, then  $G_e$  equals the kernel  $K_I \in \mathcal{F}$ , which is finitely generated. Similarly, if I is of surface type, then  $G_e \in Per(\mathcal{F}, \mathscr{E})$ , so  $G_e$  is an extension of  $K_I$  by a cyclic subgroup and it is again finitely generated.

Suppose instead that I is of exotic type. In this case, the G-stabilisers of the incident edge groups can be infinitely generated, in general. However, as mentioned above, I admits a splitting over  $K_I$  whose vertex groups are precisely the elements of  $\partial I$ . We can thus blow up the vertex  $y \in \Delta$  to a copy of this splitting of I, then collapse all edges that were incident to y in  $\Delta$  (and their G-translates). This procedure does not affect the stabilisers of the vertices of  $\Delta \setminus (G \cdot y)$ .

Summing up, there exists a  $(\mathcal{F}_{int} \cup \operatorname{Per}(\mathcal{F}, \mathscr{E}), \mathscr{E})$ -splitting  $G \curvearrowright \Delta'$  whose vertex-stabilisers are the elements of Ind not of exotic type, the kernels of the exotic components, and the vertex-stabilisers of  $\Delta$  outside Ind. All edge-stabilisers of  $\Delta'$  are finitely generated, and hence all vertex-stabilisers of  $\Delta'$  are finitely generated as well. Moreover, edges of  $\Delta'$  with stabiliser in  $\operatorname{Per}(\mathcal{F}, \mathscr{E}) \setminus \mathcal{F}_{int}$  have a vertex whose stabiliser is an element of Ind of surface type, which in particular lies in  $\operatorname{QH}(\mathcal{F}, \mathscr{E})$ .

We can finally complete the proof of Theorem B.2.

Proof of Theorem B.2. We can assume throughout the proof that  $G \notin$  Ind. Otherwise, all  $\mathcal{G}_n$  are indecomposable, and they are equivariantly isometric to each other and to T. In that case, all parts of the theorem follow from the above discussion of indecomposable actions. Furthermore, part (1) of the theorem was shown above in Lemma B.4, so we only discuss parts (2)–(4).

Part (2) can be quickly deduced from Proposition B.8. Let  $G \curvearrowright \Delta'$  be the modification of an excellent splitting, as constructed in Remark B.9. Consider a point  $p \in T$  and its stabiliser  $G_p$ . Since  $G_p \in \mathscr{E}$ , there exists a vertex  $x \in \Delta'$  such that  $G_p \leq G_x$ . If  $G_x$  is some element  $I \in Ind$  and if  $G_p$  fixes no other vertices of  $\Delta'$ , then  $G_p$  equals the *I*-stabiliser of a point of the associated indecomposable component U, which is of axial or surface type. In this case  $G_p \in \mathcal{F} \cup Per(\mathcal{F}, \mathscr{E})$ .

Otherwise, we can assume that  $G_x \notin \text{Ind.}$  Thus, we have  $G_x \in \mathscr{E}$  and there exists a point  $q \in T$ such that  $G_p \leq G_x \leq G_q$ . If  $q \neq p$ , then  $G_p$  coincides with the *G*-stabiliser of the arc  $[p,q] \subseteq T$ , and hence we again have  $G_p \in \mathcal{F}$ . Finally, if q = p, we obtain the equality  $G_p = G_x$  and, as observed in Remark B.9, the stabiliser  $G_p$  is indeed finitely generated and a vertex group of a splitting of Gwith the required properties. This proves part (2) of the theorem.

Regarding part (3), we have just just shown that, for each  $p \in T$ , either  $G_p \in \mathcal{F} \cup \operatorname{Per}(\mathcal{F}, \mathscr{E})$  or  $G_p$  is a vertex-stabiliser in the splitting  $G \curvearrowright \Delta'$ . Since G is finitely generated, the action  $G \curvearrowright \Delta'$  is cocompact. Moreover, there are only finitely many G-conjugacy classes of (relevant) elements of  $\operatorname{Per}(\mathcal{F}, \mathscr{E})$ : by Lemma B.4 there are only finitely many G-conjugacy classes in the family Ind

and, for each  $I \in$  Ind of surface type, there are only finitely many I-conjugacy classes of subgroups in  $\partial I$  (as these correspond to the boundary components of the associated compact surface). In conclusion, there are only finitely many G-conjugacy classes of point-stabilisers of T outside  $\mathcal{F}$ . The latter implies that there are only finitely many G-orbits of points of T whose stabiliser does not fix an arc, proving part (3).

We finally address part (4). If  $H \leq G$  is non-elliptic in T, then H is also non-elliptic in the splitting  $G \curvearrowright \Delta'$ . Note that H contains elements that are loxodromic in  $\Delta'$ , as chains in  $\mathcal{F}_{int}$  have bounded length by Condition (*iii*) (using Lemma 2.29). Thus, H has a well-defined minimal subtree  $M \subseteq \Delta'$ . If the G-stabiliser of at least one edge of M lies in  $\mathcal{F}_{int}$ , then we can collapse all edges of  $\Delta'$  outside the G-orbit of this edge, and thus obtain a  $(\mathcal{F}_{int}, \mathscr{E})$ -splitting of G in which H is not elliptic. Otherwise, there exists a vertex  $x \in M$  such that  $G_x$  is an element of Ind of surface type. In this case, we can refine  $\Delta'$  by blowing up the vertex x to a one-edge splitting of  $G_x$  over an element of  $\mathrm{Ess}(\mathcal{F}, \mathscr{E})$ , relative to  $\partial G_x$ . If we then collapse all edges that have not been created during the blow-up, we obtain a one-edge ( $\mathrm{Ess}(\mathcal{F}, \mathscr{E}), \mathscr{E}$ )-splitting of G in which H is not elliptic, proving part (4) and concluding the proof of Theorem B.2.

## References

- [ACGH17] Goulnara N. Arzhantseva, Christopher H. Cashen, Dominik Gruber, and David Hume. Characterizations of Morse quasi-geodesics via superlinear divergence and sublinear contraction. Doc. Math., 22:1193–1224, 2017.
- [AD02] Claire Anantharaman-Delaroche. Amenability and exactness for dynamical systems and their C\*-algebras. Trans. Amer. Math. Soc., 354(10):4153–4178, 2002.
- [AM15] Yago Antolín and Ashot Minasyan. Tits alternatives for graph products. J. Reine Angew. Math., 704:55– 83, 2015.
- [AMS16] Yago Antolín, Ashot Minasyan, and Alessandro Sisto. Commensurating endomorphisms of acylindrically hyperbolic groups and applications. *Groups Geom. Dyn.*, 10(4):1149–1210, 2016.
- [BCH94] Paul Baum, Alain Connes, and Nigel Higson. Classifying space for proper actions and K-theory of group C\*-algebras. In C\*-algebras: 1943–1993 (San Antonio, TX, 1993), volume 167 of Contemp. Math., pages 240–291. Amer. Math. Soc., Providence, RI, 1994.
- [BCV23] Corey Bregman, Ruth Charney, and Karen Vogtmann. Outer space for RAAGs. Duke Math. J., 172(6):1033–1108, 2023.
- [BDM09] Jason Behrstock, Cornelia Druţu, and Lee Mosher. Thick metric spaces, relative hyperbolicity, and quasiisometric rigidity. Math. Ann., 344(3):543–595, 2009.
- [Bes88] Mladen Bestvina. Degenerations of the hyperbolic space. Duke Math. J., 56(1):143–161, 1988.
- [BF91] Mladen Bestvina and Mark Feighn. Bounding the complexity of simplicial group actions on trees. Invent. Math., 103(3):449–469, 1991.
- [BF95] Mladen Bestvina and Mark Feighn. Stable actions of groups on real trees. Invent. Math., 121(2):287–321, 1995.
- [BF09] Mladen Bestvina and Koji Fujiwara. A characterization of higher rank symmetric spaces via bounded cohomology. Geom. Funct. Anal., 19(1):11–40, 2009.
- [BFH00] Mladen Bestvina, Mark Feighn, and Michael Handel. The Tits alternative for  $Out(F_n)$ . I. Dynamics of exponentially-growing automorphisms. Ann. of Math. (2), 151(2):517–623, 2000.
- [BFH05] Mladen Bestvina, Mark Feighn, and Michael Handel. The Tits alternative for  $Out(F_n)$ . II. A Kolchin type theorem. Ann. of Math. (2), 161(1):1–59, 2005.
- [BG10] Martin R. Bridson and Daniel Groves. The quadratic isoperimetric inequality for mapping tori of free group automorphisms. *Mem. Amer. Math. Soc.*, 203(955):xii+152, 2010.
- [BGH22] Mladen Bestvina, Vincent Guirardel, and Camille Horbez. Boundary amenability of  $Out(F_N)$ . Ann. Sci. Éc. Norm. Supér. (4), 55(5):1379–1431, 2022.
- [BH92] Mladen Bestvina and Michael Handel. Train tracks and automorphisms of free groups. Ann. of Math. (2), 135(1):1–51, 1992.
- [Bow98] Brian H. Bowditch. Cut points and canonical splittings of hyperbolic groups. Acta Math., 180(2):145–186, 1998.
- [Bow13] Brian H. Bowditch. Coarse median spaces and groups. *Pacific J. Math.*, 261(1):53–93, 2013.
- [Bow16] Brian H. Bowditch. Some properties of median metric spaces. *Groups Geom. Dyn.*, 10(1):279–317, 2016.

- [BQ18] Corey Bregman and Yulan Qing. Dilatation of outer automorphisms of right-angled Artin groups. arXiv:1810.06499, 2018.
- [Bro94] Kenneth S. Brown. Cohomology of groups, volume 87 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original.
- [BS08] Igor Belegradek and Andrzej Szczepański. Endomorphisms of relatively hyperbolic groups. Internat. J. Algebra Comput., 18(1):97–110, 2008. With an appendix by Oleg V. Belegradek.
- [Che00] Victor Chepoi. Graphs of some CAT(0) complexes. Adv. in Appl. Math., 24(2):125–179, 2000.
- [CHHL] Rémi Coulon, Arnaud Hilion, Camille Horbez, and Gilbert Levitt. In preparation.
- [CL83] Donald J. Collins and Frank Levin. Automorphisms and Hopficity of certain Baumslag-Solitar groups. Arch. Math. (Basel), 40(5):385–400, 1983.
- [CN05] Sarah Campbell and Graham A. Niblo. Hilbert space compression and exactness of discrete groups. J. Funct. Anal., 222(2):292–305, 2005.
- [Coo87] Daryl Cooper. Automorphisms of free groups have finitely generated fixed point sets. J. Algebra, 111(2):453-456, 1987.
- [Cou22] Rémi Coulon. Examples of groups whose automorphisms have exotic growth. Algebr. Geom. Topol., 22(4):1497–1510, 2022.
- [CRHK24] Montserrat Casals-Ruiz, Mark Hagen, and Ilya Kazachkov. Real cubings and asymptotic cones of hierarchically hyperbolic groups. Preprint, available at www.wescac.net/cones.html, 2024.
- [CRK11] Montserrat Casals-Ruiz and Ilya Kazachkov. On systems of equations over free partially commutative groups. Mem. Amer. Math. Soc., 212(999):viii+153, 2011.
- [CRK15] Montserrat Casals-Ruiz and Ilya Kazachkov. Limit groups over partially commutative groups and group actions on real cubings. *Geom. Topol.*, 19(2):725–852, 2015.
- [CS11] Pierre-Emmanuel Caprace and Michah Sageev. Rank rigidity for CAT(0) cube complexes. Geom. Funct. Anal., 21(4):851–891, 2011.
- [CS15] Ruth Charney and Harold Sultan. Contracting boundaries of CAT(0) spaces. J. Topol., 8(1):93–117, 2015.
- [CSV17] Ruth Charney, Nathaniel Stambaugh, and Karen Vogtmann. Outer space for untwisted automorphisms of right-angled Artin groups. *Geom. Topol.*, 21(2):1131–1178, 2017.
- [CT94] Donald J. Collins and Edward C. Turner. Efficient representatives for automorphisms of free products. Michigan Math. J., 41(3):443–464, 1994.
- [CV86] Marc Culler and Karen Vogtmann. Moduli of graphs and automorphisms of free groups. Invent. Math., 84(1):91–119, 1986.
- [CV09] Ruth Charney and Karen Vogtmann. Finiteness properties of automorphism groups of right-angled Artin groups. Bull. Lond. Math. Soc., 41(1):94–102, 2009.
- [D] Steve D. Extension of a virtually polycyclic group by a virtually polycyclic group. Mathematics Stack Exchange. https://math.stackexchange.com/q/2073210 (version: 2016-12-27).
- [Day09] Matthew B. Day. Peak reduction and finite presentations for automorphism groups of right-angled Artin groups. Geom. Topol., 13(2):817–855, 2009.
- [DG08] François Dahmani and Daniel Groves. The isomorphism problem for toral relatively hyperbolic groups. Publ. Math. Inst. Hautes Études Sci., (107):211–290, 2008.
- [DG11] François Dahmani and Vincent Guirardel. The isomorphism problem for all hyperbolic groups. *Geom. Funct. Anal.*, 21(2):223–300, 2011.
- [Din12] Tien-Cuong Dinh. Tits alternative for automorphism groups of compact Kähler manifolds. Acta Math. Vietnam., 37(4):513–529, 2012.
- [DK92] Gérard Duchamp and Daniel Krob. The lower central series of the free partially commutative group. Semigroup Forum, 45(3):385–394, 1992.
- [DK18] Cornelia Druţu and Michael Kapovich. Geometric group theory, volume 63 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 2018. With an appendix by Bogdan Nica.
- [DS99] Martin J. Dunwoody and Michah E. Sageev. JSJ-splittings for finitely presented groups over slender groups. Invent. Math., 135(1):25–44, 1999.
- [DS00] Martin J. Dunwoody and Eric L. Swenson. The algebraic torus theorem. *Invent. Math.*, 140(3):605–637, 2000.
- [DS08] Cornelia Druţu and Mark V. Sapir. Groups acting on tree-graded spaces and splittings of relatively hyperbolic groups. Adv. Math., 217(3):1313–1367, 2008.
- [Dun85] Martin J. Dunwoody. The accessibility of finitely presented groups. Invent. Math., 81(3):449–457, 1985.
- [DW19] Matthew B. Day and Richard D. Wade. Relative automorphism groups of right-angled Artin groups. J. Topol., 12(3):759–798, 2019.
- [Fio23] Elia Fioravanti. On automorphisms and splittings of special groups. *Compos. Math.*, 159(2):232–305, 2023.

- [Fio24] Elia Fioravanti. Coarse-median preserving automorphisms. *Geom. Topol.*, 28(1):161–266, 2024.
- [FK24] Elia Fioravanti and Alice Kerr. Short positive loxodromics in graph products. arXiv:2410.06751, 2024.
- [FLS24] Elia Fioravanti, Ivan Levcovitz, and Michah Sageev. Coarse cubical rigidity. J. Topol., 17(3):Paper No. e12353, 50, 2024.
- [FM15] Stefano Francaviglia and Armando Martino. Stretching factors, metrics and train tracks for free products. Illinois J. Math., 59(4):859–899, 2015.
- [For02] Max Forester. Deformation and rigidity of simplicial group actions on trees. *Geom. Topol.*, 6:219–267, 2002.
- [FP06] Koji Fujiwara and Panagiotis Papasoglu. JSJ-decompositions of finitely presented groups and complexes of groups. Geom. Funct. Anal., 16(1):70–125, 2006.
- [Gen21a] Anthony Genevois. Algebraic characterisation of relatively hyperbolic special groups. Israel J. Math., 241(1):301–341, 2021.
- [Gen21b] Anthony Genevois. Rank-one isometries of CAT(0) cube complexes and their centralisers. Adv. Geom., 21(3):347–364, 2021.
- [GH19] Daniel Groves and Michael Hull. Homomorphisms to acylindrically hyperbolic groups I: Equationally noetherian groups and families. *Trans. Amer. Math. Soc.*, 372(10):7141–7190, 2019.
- [GH22] Vincent Guirardel and Camille Horbez. Boundaries of relative factor graphs and subgroup classification for automorphisms of free products. *Geom. Topol.*, 26(1):71–126, 2022.
- [GHW05] Erik Guentner, Nigel Higson, and Shmuel Weinberger. The Novikov conjecture for linear groups. Publ. Math. Inst. Hautes Études Sci., (101):243–268, 2005.
- [GJLL98] Damien Gaboriau, Andre Jaeger, Gilbert Levitt, and Martin Lustig. An index for counting fixed points of automorphisms of free groups. *Duke Math. J.*, 93(3):425–452, 1998.
- [GL95] Damien Gaboriau and Gilbert Levitt. The rank of actions on R-trees. Ann. Sci. École Norm. Sup. (4), 28(5):549–570, 1995.
- [GL07] Vincent Guirardel and Gilbert Levitt. The outer space of a free product. Proc. Lond. Math. Soc. (3), 94(3):695-714, 2007.
- [GL11] Vincent Guirardel and Gilbert Levitt. Trees of cylinders and canonical splittings. *Geom. Topol.*, 15(2):977–1012, 2011.
- [GL15] Vincent Guirardel and Gilbert Levitt. Splittings and automorphisms of relatively hyperbolic groups. Groups Geom. Dyn., 9(2):599–663, 2015.
- [GL17] Vincent Guirardel and Gilbert Levitt. JSJ decompositions of groups. Astérisque, (395):vii+165, 2017.
- [GLP94] Damien Gaboriau, Gilbert Levitt, and Frédéric Paulin. Pseudogroups of isometries of R and Rips' theorem on free actions on R-trees. Israel J. Math., 87(1-3):403–428, 1994.
- [Gro87] Misha Gromov. Hyperbolic groups. In Essays in group theory, volume 8 of Math. Sci. Res. Inst. Publ., pages 75–263. Springer, New York, 1987.
- [Gro09] Daniel Groves. Limit groups for relatively hyperbolic groups. I. The basic tools. Algebr. Geom. Topol., 9(3):1423–1466, 2009.
- [Gru57] Karl W. Gruenberg. Residual properties of infinite soluble groups. Proc. London Math. Soc. (3), 7:29–62, 1957.
- [Gui98] Vincent Guirardel. Approximations of stable actions on **R**-trees. Comment. Math. Helv., 73(1):89–121, 1998.
- [Gui04] Vincent Guirardel. Limit groups and groups acting freely on ℝ<sup>n</sup>-trees. Geom. Topol., 8:1427–1470, 2004.
   [Gui08] Vincent Guirardel. Actions of finitely generated groups on ℝ-trees. Ann. Inst. Fourier (Grenoble), 58(1):159–211, 2008.
- [Hal59] Marshall Hall, Jr. The theory of groups. The Macmillan Company, New York, 1959.
- [Ham09] Ursula Hamenstädt. Geometry of the mapping class groups. I. Boundary amenability. Invent. Math., 175(3):545–609, 2009.
- [Hig00] Nigel Higson. Bivariant K-theory and the Novikov conjecture. Geom. Funct. Anal., 10(3):563–581, 2000.
- [Hor14] Camille Horbez. The Tits alternative for the automorphism group of a free product. arXiv:1408.0546, 2014.
- [HR00] Nigel Higson and John Roe. Amenable group actions and the Novikov conjecture. J. Reine Angew. Math., 519:143–153, 2000.
- [HW08] Frédéric Haglund and Daniel T. Wise. Special cube complexes. Geom. Funct. Anal., 17(5):1551–1620, 2008.
- [HW21] Frédéric Haglund and Daniel T. Wise. A note on finiteness properties of graphs of groups. Proc. Amer. Math. Soc. Ser. B, 8:121–128, 2021.
- [Krs92] Sava Krstić. Finitely generated virtually free groups have finitely presented automorphism group. Proc. London Math. Soc. (3), 64(1):49–69, 1992.

- [KW99] Eberhard Kirchberg and Simon Wassermann. Permanence properties of  $C^*$ -exact groups. Doc. Math., 4:513–558, 1999.
- [Lev05] Gilbert Levitt. Automorphisms of hyperbolic groups and graphs of groups. *Geom. Dedicata*, 114:49–70, 2005.
- [Lev09] Gilbert Levitt. Counting growth types of automorphisms of free groups. *Geom. Funct. Anal.*, 19(4):1119–1146, 2009.
- [LP97] Gilbert Levitt and Frédéric Paulin. Geometric group actions on trees. Amer. J. Math., 119(1):83–102, 1997.
- [Lym22a] Rylee A. Lyman. CTs for free products. arXiv:2203.08868, 2022.
- [Lym22b] Rylee A. Lyman. Train track maps on graphs of groups. Groups Geom. Dyn., 16(4):1389–1422, 2022.
- [Mal51] A. I. Mal'cev. On some classes of infinite soluble groups. Mat. Sbornik N.S., 28(70):567–588, 1951.
- [McC85] John McCarthy. A "Tits-alternative" for subgroups of surface mapping class groups. Trans. Amer. Math. Soc., 291(2):583–612, 1985.
- [Oza06] Narutaka Ozawa. Amenable actions and applications. In International Congress of Mathematicians. Vol. II, pages 1563–1580. Eur. Math. Soc., Zürich, 2006.
- [Pau91] Frédéric Paulin. Outer automorphisms of hyperbolic groups and small actions on R-trees. In Arboreal group theory (Berkeley, CA, 1988), volume 19 of Math. Sci. Res. Inst. Publ., pages 331–343. Springer, New York, 1991.
- [Rol98] Martin A. Roller. Poc sets, median algebras and group actions. An extended study of Dunwoody's construction and Sageev's theorem. Preprint, University of Southampton, available at arXiv:1607.07747, 1998.
- [RS94] Eliyahu Rips and Zlil Sela. Structure and rigidity in hyperbolic groups. I. Geom. Funct. Anal., 4(3):337– 371, 1994.
- [RS97] Eliyahu Rips and Zlil Sela. Cyclic splittings of finitely presented groups and the canonical JSJ decomposition. Ann. of Math. (2), 146(1):53–109, 1997.
- [RZ10] Luis Ribes and Pavel Zalesskii. Profinite groups, volume 40 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, second edition, 2010.
- [Sel95] Zlil Sela. The isomorphism problem for hyperbolic groups. I. Ann. of Math. (2), 141(2):217–283, 1995.
- [Sel96] Zlil Sela. The Nielsen-Thurston classification and automorphisms of a free group. I. Duke Math. J., 84(2):379–397, 1996.
- [Sel97a] Zlil Sela. Acylindrical accessibility for groups. Invent. Math., 129(3):527–565, 1997.
- [Sel97b] Zlil Sela. Structure and rigidity in (Gromov) hyperbolic groups and discrete groups in rank 1 Lie groups. II. Geom. Funct. Anal., 7(3):561–593, 1997.
- [Sel23a] Zlil Sela. Automorphisms of groups and a higher rank JSJ decomposition I: RAAGs and a higher rank Makanin-Razborov diagram. Geom. Funct. Anal., 33(3):824–874, 2023.
- [Sel23b] Zlil Sela. Automorphisms of groups and a higher rank JSJ decomposition II: the single ended case. arXiv:2209.12891, 2023.
- [Ser89] Herman Servatius. Automorphisms of graph groups. J. Algebra, 126(1):34–60, 1989.
- [Ser03] Jean-Pierre Serre. *Trees.* Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. Translated from the French original by John Stillwell, Corrected 2nd printing of the 1980 English translation.
- [Sis16] Alessandro Sisto. Quasi-convexity of hyperbolically embedded subgroups. *Math. Z.*, 283(3-4):649–658, 2016.
- [Sta68] John R. Stallings. On torsion-free groups with infinitely many ends. Ann. of Math. (2), 88:312–334, 1968.
   [sTe] sTertooy. Group automorphism fixing finite-index subgroup. Mathematics Stack Exchange. https://math.stackexchange.com/q/4962759 (version: 2024-08-25).
- [SW05] Michah Sageev and Daniel T. Wise. The Tits alternative for CAT(0) cubical complexes. Bull. London Math. Soc., 37(5):706–710, 2005.
- [Thu88] William P. Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. Bull. Amer. Math. Soc. (N.S.), 19(2):417–431, 1988.
- [Tit72] Jacques Tits. Free subgroups in linear groups. J. Algebra, 20:250–270, 1972.
- [Toi13] Emmanuel Toinet. Conjugacy *p*-separability of right-angled Artin groups and applications. *Groups Geom.* Dyn., 7(3):751–790, 2013.
- [Yu00] Guoliang Yu. The coarse Baum-Connes conjecture for spaces which admit a uniform embedding into Hilbert space. *Invent. Math.*, 139(1):201–240, 2000.

Institute of Algebra and Geometry, Karlsruhe Institute of Technology *Email address:* elia.fioravanti@kit.edu