

# SYMMETRIC CARTAN CALCULUS, THE PATTERSON-WALKER METRIC AND SYMMETRIC COHOMOLOGY

FILIP MOUČKA AND ROBERTO RUBIO

ABSTRACT. We develop symmetric Cartan calculus, an analogue of classical Cartan calculus for symmetric differential forms. We first show that the analogue of the exterior derivative, the symmetric derivative, is not unique and its different choices are parametrized by torsion-free affine connections. We use a choice of symmetric derivative to generate the symmetric bracket and the symmetric Lie derivative, and give geometric interpretations of all of them. By proving the structural identities and describing the role of affine morphisms, we reveal an unexpected link of symmetric Cartan calculus with the Patterson-Walker metric, which we recast as a direct analogue of the canonical symplectic form on the cotangent bundle. We show that, in the light of the Patterson-Walker metric, symmetric Cartan calculus becomes a complete analogue of classical Cartan calculus. As an application of this new framework, we introduce symmetric cohomology, which involves Killing tensors, or, in the case of the Levi-Civita connection, Killing vector fields. We characterize the Levi-Civita connections of  $S^1$  in terms of symmetric cohomology and develop further techniques.

## CONTENTS

1. Introduction	2
2. Symmetric Cartan calculus	5
3. Commutation relations of symmetric Cartan calculus	16
4. Affine manifold morphisms	20
5. The Patterson-Walker metric	25
6. Symmetric cohomology	33
7. Further techniques for symmetric cohomology	41
Appendix A. Non-geometric derivations	51
Appendix B. Various lifts to the cotangent bundle	51
Appendix C. Computations for $H_{\nabla}^1(\mathbb{R}^2)$	54
Appendix D. Comparison with classical Cartan calculus	55
References	57

---

This project has been supported by MICIU/AEI/10.13039/501100011033 and the EU FEDER under the grant PID2022-137667NA-I00 (GENTLE). The first author has been partially supported by the Grant Agency of the Czech Technical University in Prague, grant No. OHK4-013/25. The second author has also received support from the MICIU/AEI and the EU FSE under the Ramón y Cajal fellowship RYC2020-030114-I and from the AGAUR under the grant 2021-SGR-01015.

## 1. INTRODUCTION

Classical Cartan calculus, understood as the calculus of vector fields  $\mathfrak{X}(M)$  and differential forms  $\Omega^\bullet(M)$  on a manifold  $M$ , is an ubiquitous language for differential geometry. Its structural identities are the graded commutators of the contraction operator  $\iota$ , the exterior derivative  $d$  and the Lie derivative  $L$ . Namely, for  $X, Y \in \mathfrak{X}(M)$ , we have:

$$(1) \quad \begin{aligned} [\iota_X, \iota_Y]_g &= 0, & [d, d]_g &= 0, & [\iota_X, d]_g &= L_X, \\ [L_X, \iota_Y]_g &= \iota_{[X, Y]}, & [L_X, L_Y]_g &= L_{[X, Y]}, & [d, L_X]_g &= 0. \end{aligned}$$

The exterior derivative plays a geometric generating role. Together with the contraction, which is a linear-algebraic operation (that is, at the level of bundles, not necessarily sections), it recovers the Lie derivative and also the Lie bracket as a derived bracket. Furthermore, the exterior derivative gives rise to the de Rham cohomology.

A very natural question is whether there is an analogous framework for symmetric differential forms and what it might be helpful for. This would first mean finding analogues to the exterior derivative, the Lie derivative and the Lie bracket satisfying similar properties. Surprisingly, the only precedent we are aware of is the definition of a symmetric bracket in [12] (later geometrically interpreted in [21]), which was used, by mimicking the explicit formulas in the usual setting, to define a symmetric differential and a symmetric Lie derivative [16]. We have found no justification that these are the only possible or even reasonable choices, no exhaustive study of the structural identities analogous to (1) and, beyond the symmetric bracket [3, 6], neither a full geometric interpretation nor a significant application.

The aim of the present work is to develop and justify a sound and geometrically meaningful approach to symmetric Cartan calculus, as well as to show its two-way interaction with the Patterson-Walker metric, and introduce a symmetric cohomology theory, which is related to Killing tensors. Key to our strategy is our starting point: we look for all possible symmetric analogues of the exterior derivative that can play an equivalent generating role as a geometric derivation (Definition 2.4).

The first main feature of symmetric Cartan calculus is, precisely, that there is no canonical symmetric derivative for symmetric differential forms, which we denote by  $\Upsilon^\bullet(M)$ . Such an analogue depends on the choice of a connection, which can be, without loss of generality, assumed to be torsion free. In fact, we prove:

**Proposition 2.9.** *The map  $\nabla \mapsto \nabla^s$  gives an isomorphism of  $\Upsilon^2(M, TM)$ -affine spaces:*

$$\left\{ \begin{array}{l} \text{torsion-free} \\ \text{connections on } M \end{array} \right\} \xleftrightarrow{\sim} \left\{ \begin{array}{l} \text{geometric derivations of the algebra} \\ \text{of symmetric forms on } M \end{array} \right\}.$$

An a priori surprising fact is that  $\nabla^s$  does not square to zero. However, this is natural, since the symmetric derivative is a usual derivation and derivations are an algebra for the usual commutator, as opposed to the graded commutator for graded derivations. Thus, squaring to zero is natural in the context of classical Cartan calculus since  $[d, d]_g = 2d \circ d$ , whereas in the symmetric case, we have that the analogous expression of squaring to zero is  $[\nabla^s, \nabla^s] = 0$  for the usual commutator, which is trivially satisfied.

We give a geometric interpretation of  $\nabla^s$  making use of the correspondence between a symmetric form  $\varphi$  and a polynomial in velocities  $\tilde{\varphi}$  (an element of a subspace of  $\mathcal{C}^\infty(TM)$ , as we explain in Section 2.3), together with the geodesic spray  $X_\nabla \in \mathfrak{X}(TM)$ .

**Proposition 2.14.** *Let  $\nabla$  be a connection on  $M$ . For every  $\varphi \in \Upsilon^r(M)$  we have*

$$\widetilde{\nabla^s \varphi} = X_\nabla \tilde{\varphi}.$$

The symmetric Lie derivative with respect to a vector field  $X \in \mathfrak{X}(M)$  is introduced as  $L_X^s = [\iota_X, \nabla^s]$  and we show in Proposition 2.19 (see also Figure 1) that it captures the variation with respect to a combination of the flow of  $X$  and the parallel transport with respect to  $\nabla$ . Finally, the symmetric bracket is derived from the formula  $[L_X^s, \iota_Y] = \iota_{\langle X:Y \rangle_s}$ . Its geometric interpretation in relation to geodesically invariant distributions is recalled in Section 2.7. Overall, we have the following properties for the (ungraded) commutator:

$$\begin{aligned} [\iota_X, \iota_Y] &= 0, & [\nabla^s, \nabla^s] &= 0, & [\iota_X, \nabla^s] &= L_X^s, \\ [L_X^s, \iota_Y] &= \iota_{\langle X:Y \rangle_s}, & [L_X^s, L_Y^s] &\neq L_{\langle X:Y \rangle_s}^s, & [\nabla^s, L_X^s] &\neq 0. \end{aligned}$$

Note that the analogues of the last two formulae of (1) are not true. Instead, we show in Section 3 that

$$\begin{aligned} [L_X^s, L_Y^s] &= L_{[X,Y]}^s + \iota_{2(\nabla_{\nabla X} Y - \nabla_{\nabla Y} X - R_\nabla(X,Y))}, \\ [\nabla^s, L_X^s] &= 2\nabla_{\nabla X}^s + \iota_{2\text{sym} \iota_X R_\nabla + R_\nabla^s X}. \end{aligned}$$

The study of the right-hand side of these expressions leads us to find an unexpected link with the Patterson-Walker metric  $g_\nabla$ , a metric on  $T^*M$  that is determined by the choice of a torsion-free connection  $\nabla$  on  $M$ . On the one hand, by using a lift  $\hat{\nabla}$  of the connection  $\nabla$  to  $T^*M$  (Section 5.2), the symmetric derivative allows us to recast the Patterson-Walker metric as an analogue of the canonical symplectic form on  $T^*M$ .

**Theorem 5.7.** *For any torsion-free connection  $\nabla$  on  $M$ , there holds*

$$\hat{\nabla}^s \alpha_{can} = g_\nabla,$$

where  $\alpha_{can}$  is the canonical 1-form on  $T^*M$ .

On the other hand, the Patterson-Walker metric, together with the notion of the complete lift  $X^c \in \mathfrak{X}(T^*M)$  of a vector field  $X$ , gives a framework in which the structural identities become much simpler.

**Theorem 5.19.** *Let  $\nabla$  be a torsion-free connection on  $M$  and  $X, Y \in \mathfrak{X}(M)$ . All of the six relations*

$$\begin{aligned} [\iota_X, \iota_Y] &= 0, & [\nabla^s, \nabla^s] &= 0, & [\iota_X, \nabla^s] &= L_X^s, \\ [L_X^s, \iota_Y] &= \iota_{\langle X:Y \rangle_s}, & [L_X^s, L_Y^s] &= 0, & [\nabla^s, L_X^s] &= 0 \end{aligned}$$

*are satisfied if and only if  $X^c$  and  $Y^c$  are gradient Killing vector fields for  $g_\nabla$ .*

This seemingly extra condition on  $X$  and  $Y$  implicitly appears in the classical case as well. Indeed, the analogues of gradient Killing vector fields for  $g_\nabla$  are Hamiltonian ones for the canonical 2-form  $\omega_{\text{can}}$ , and complete lifts are always Hamiltonian. This, together with the fact that  $[L_X^s, L_Y^s] = 0 = L_{\langle X:Y \rangle}^s$  if  $X^c$  and  $Y^c$  are gradient Killing, draws a complete analogy between Theorem 5.19 and (1).

The study of the lifts required to prove these last two theorems is recalled in Appendix B.

Finally, just as the exterior derivative plays a generating role in classical Cartan calculus and is the basis of de Rham cohomology, the question of whether there is any cohomology theory associated to the symmetric derivative arises naturally.

Closed symmetric forms for the symmetric derivative  $\nabla^s$  recover the notion of Killing tensors. In the case of the Levi-Civita connection of a (pseudo-)Riemannian metric, Killing tensors are well established. They are related to the separability of Hamilton-Jacobi equation [4], or the symmetries of the Laplacian [13]. Probably the most famous Killing tensor is the *Carter tensor* in the *Kerr-Newman spacetime*, the spacetime of a charged rotating black hole [9, 10, 26].

Defining a cohomology theory with an operator that does not square to zero requires some adjustment, namely, considering the intersection of the image with the kernel. We do this to define symmetric cohomology, and give a geometric interpretation in terms of polynomials in velocities (Corollary 6.14). We focus specially on the first symmetric cohomology group. We prove that its dimension is bounded, and, when dealing with a (pseudo-)Riemannian metric, we can relate it to Killing vector fields modulo gradient Killing vector fields.

Symmetric cohomology depends, as symmetric Cartan calculus does, on the choice of a torsion-free connection. Far from being a hindrance, we take advantage of this fact, to prove our last result, which offers a characterization of Levi-Civita connections on  $S^1$  in terms of symmetric cohomology.

**Proposition 6.26.** *The only connections  $\nabla$  on  $S^1$  such that  $H_\nabla^1(S^1) \neq \{0\}$  (and hence  $\mathbb{R}$ ) are the Levi-Civita connections for any choice of metric on  $S^1$ .*

We present further techniques for symmetric cohomology, namely the Künneth subalgebra, the Patterson-Walker bound and the study of left-invariant symmetric cohomology. These have proved to be useful tools in obtaining information

about the symmetric cohomology groups of product manifolds, cotangent bundles and compact Lie groups. Further applications of symmetric Cartan calculus will come in forthcoming papers about symmetric Poisson structures and generalized geometry.

We finish the paper with a table (Appendix D) that summarizes and compares various aspects of symmetric Cartan calculus with classical Cartan calculus.

**Acknowledgements.** We would like to thank the Weizmann Institute of Science for their hospitality in hosting us as visitors, Andrew Lewis for helpful discussions about the symmetric bracket and Rudolf Šmolka for noticing a mistake in the first version of this text.

**Notation and conventions.** We work on the smooth category for manifolds and maps. We denote an arbitrary manifold (of positive dimension) by  $M$ , its smooth functions by  $\mathcal{C}^\infty(M)$ , its tangent bundle by  $TM$ , its cotangent bundle by  $T^*M$ , its vector fields by  $\mathfrak{X}(M)$  and its differential forms by  $\Omega^\bullet(M)$ .

When we consider a connection on  $M$ , we mean an affine connection (that is, a connection on  $TM$ ), which can determine other connections (for instance, on  $T^*M$  by duality). We will denote these connections with the same symbol.

When we consider a vector space  $V$ , we mean a finite-dimensional vector space over  $\mathbb{R}$  or  $\mathbb{C}$ . We use the Einstein summation convention throughout the text.

Note that our definition of symmetric cohomology will be unrelated to that of symmetric group cohomology introduced in [24].

## 2. SYMMETRIC CARTAN CALCULUS

We first look at the linear-algebraic level before moving to smooth manifolds.

**2.1. The symmetric algebra and its derivations.** Consider a vector space  $V$ , its dual space  $V^*$  and the tensor algebra  $\otimes^\bullet V^*$ . The natural action of a permutation  $\pi \in \mathcal{S}_r$  on a decomposable element (with  $\alpha^j \in V^*$ ),

$$\pi \cdot (\alpha^1 \otimes \dots \otimes \alpha^r) := \alpha^{\pi^1} \otimes \dots \otimes \alpha^{\pi^r},$$

is extended linearly to  $\otimes^r V^*$  and gives rise to the **symmetric projection**  $\text{sym}_r : \otimes^r V^* \rightarrow \otimes^r V^*$ ,

$$\text{sym}_r(\varphi) := \frac{1}{r!} \sum_{\pi \in \mathcal{S}_r} \pi \cdot \varphi,$$

and, more generally,  $\text{sym} := \bigoplus_{r \in \mathbb{N}} \text{sym}_r$ . Consider the  $\mathbb{N}$ -graded vector space

$$\text{Sym}^\bullet V^* := \frac{\otimes^\bullet V^*}{\ker \text{sym}} \cong \text{Im } \text{sym}.$$

The latter isomorphism allows us to realize  $\text{Sym}^\bullet V^*$  as a vector subspace of  $\otimes^\bullet V^*$ . Define the **symmetric product** of  $\varphi \in \text{Sym}^r V^*$  and  $\psi \in \text{Sym}^l V^*$  by

$$\varphi \odot \psi := \frac{(r+l)!}{r!l!} \text{sym}(\varphi \otimes \psi).$$

We thus have the **symmetric algebra**  $(\text{Sym}^\bullet V^*, +, \odot)$ .

At this linear-algebraic level, we can already look at the **contraction operator**. The degree- $(-1)$  map  $\iota_X$ , for any  $X \in V$ , descends from  $\otimes^\bullet V^*$  to  $\text{Sym}^\bullet V^*$ , where it becomes a derivation, that is,

$$(2) \quad \iota_X(\varphi \odot \psi) = \iota_X \varphi \odot \psi + \varphi \odot \iota_X \psi,$$

as we show in the proof of the next lemma. Note that the sign is always  $+$ , that is, it is not graded.

**Lemma 2.1.** *The degree- $(-1)$  derivations of  $\text{Sym}^\bullet V^*$  are in bijective correspondence with  $V$  via the contraction operator  $\iota$ .*

*Proof.* We first prove that the contraction operator is indeed a derivation of  $\text{Sym}^\bullet V^*$ . Given  $\varphi \in \text{Sym}^r V^*$  and  $\psi \in \text{Sym}^l V^*$ , we denote  $n := r + l - 1$ . We shall use the shorthand notation  $\varphi(Y) := \varphi(Y, \dots, Y)$  for a symmetric  $r$ -tensor  $\varphi$  acting on  $r$  copies of a vector  $Y \in V$ . Then, for all  $Y \in V$ ,

$$(\iota_X(\varphi \odot \psi))(Y) = \frac{1}{r!l!} \sum_{\pi \in \mathcal{S}_n} \iota_X(\pi \cdot (\varphi \otimes \psi))(Y).$$

We can write the symmetric group  $\mathcal{S}_n$  as the disjoint union of the two subsets  $\mathcal{S}_n^r := \{\pi \in \mathcal{S}_n \mid 1 \in \pi(\{1, \dots, r\})\}$  and  $\mathcal{S}_n^l := \{\pi \in \mathcal{S}_n \mid 1 \in \pi(\{r+1, \dots, r+l\})\}$ . Therefore,

$$(\iota_X(\varphi \odot \psi))(Y) = \frac{1}{r!l!} \left( \sum_{\pi \in \mathcal{S}_n^r} (\iota_X \varphi)(Y) \psi(Y) + \sum_{\pi \in \mathcal{S}_n^l} \varphi(Y) (\iota_X \psi)(Y) \right).$$

Since the cardinal of the sets  $\mathcal{S}_n^r$  and  $\mathcal{S}_n^l$  is  $n!r$  and  $n!l$ , respectively, we arrive at

$$\begin{aligned} (\iota_X(\varphi \odot \psi))(Y) &= \frac{n!}{(r-1)!l!} (\iota_X \varphi)(Y) \psi(Y) + \frac{n!}{r!(l-1)!} \varphi(Y) (\iota_X \psi)(Y) \\ &= \frac{1}{(r-1)!l!} \sum_{\pi \in \mathcal{S}_n} (\iota_X \varphi)(Y) \psi(Y) + \frac{1}{r!(l-1)!} \sum_{\pi \in \mathcal{S}_n} \varphi(Y) (\iota_X \psi)(Y) \\ &= (\iota_X \varphi \odot \psi + \varphi \odot \iota_X \psi)(Y). \end{aligned}$$

Identity (2) then follows from polarization, so  $\iota_X$  is a derivation. As an algebra,  $\text{Sym}^\bullet V^*$  is generated by  $\text{Sym}^0 V^*$ , that is, the scalars (where degree- $(-1)$  maps act trivially), and  $\text{Sym}^1 V^* = V^*$ , whose linear maps are exactly given by  $(V^*)^* \cong V$  via the contraction  $\iota$ .  $\square$

**2.2. Geometric derivations.** Given a manifold  $M$ , we replace  $V$  in Section 2.1 by  $T_m M$  and consider the  $r$ th-symmetric power bundle

$$\mathrm{Sym}^r T^* M,$$

whose sections we denote by  $\Upsilon^r(M)$ . We define the graded  $\mathcal{C}^\infty(M)$ -module

$$\Upsilon^\bullet(M) := \bigoplus_{r \in \mathbb{N}} \Upsilon^r(M).$$

*Remark 2.2.* Note that  $\Upsilon^0(M) = \mathcal{C}^\infty(M)$  and  $\Upsilon^1(M) = \Omega^1(M)$ , we will use one notation or the other depending on the context. On the other hand, we have avoided talking about  $\mathrm{Sym}^\bullet T^* M$  since it would be a bundle of infinite rank.

*Remark 2.3.* It will be convenient to also introduce the notation

$$\Upsilon^r(M, E) := \Gamma(\mathrm{Sym}^r T^* M \otimes E)$$

for an arbitrary vector bundle  $E$  over  $M$  and, analogously,  $\Upsilon^\bullet(M, E)$ .

The symmetric projection, the symmetric product and the contraction of Section 2.1 are easily extended to tensor fields. We thus obtain a graded algebra

$$(\Upsilon^\bullet(M), +, \odot).$$

The derivations of any graded algebra together with the commutator form a Lie algebra that is moreover graded. In this case, we denote it by  $\mathrm{Der}(\Upsilon^\bullet(M))$ . For derivations of fixed degree, we will use a subindex, e.g.,  $\iota_X \in \mathrm{Der}_{-1}(\Upsilon^\bullet(M))$ .

Since the exterior derivative, together with the contraction, generates the Cartan calculus for  $\Omega^\bullet(M)$ , now we look for analogous derivations of  $\Upsilon^\bullet(M)$ , which we call geometric.

**Definition 2.4.** A **geometric derivation**  $D \in \mathrm{Der}(\Upsilon^\bullet(M))$  is a derivation of degree 1 satisfying, for all  $f \in \mathcal{C}^\infty(M)$  and  $X \in \mathfrak{X}(M)$ ,

$$(Df)(X) = Xf.$$

Every derivation of  $\Upsilon^\bullet(M)$  is a local operator, and since the action on functions is prescribed, any geometric derivation is uniquely characterized by its action on 1-forms. Namely, it is determined by a linear map  $D : \Upsilon^1(M) \rightarrow \Upsilon^2(M)$  such that

$$(3) \quad D(f\alpha) = df \odot \alpha + fD\alpha$$

for all  $f \in \mathcal{C}^\infty(M)$  and  $\alpha \in \Upsilon^1(M) = \Omega^1(M)$ . Formula (3) resembles the defining property of a connection on the cotangent bundle. In fact, given such a connection  $\nabla : \Omega^1(M) \rightarrow \Omega^1(M, T^*M)$ , we define the linear map  $\nabla^s : \Upsilon^1(M) \rightarrow \Upsilon^2(M)$  by

$$(4) \quad \nabla^s := 2 \mathrm{sym} \circ \nabla,$$

which satisfies

$$\nabla^s(f\alpha) = 2 \mathrm{sym}(\nabla(f\alpha)) = 2 \mathrm{sym}(df \otimes \alpha + f\nabla\alpha) = df \odot \alpha + f\nabla^s\alpha.$$

**Definition 2.5.** Let  $\nabla$  be a connection on  $M$ . The **symmetric derivative** corresponding to  $\nabla$  is the geometric derivation  $\nabla^s \in \text{Der}(\Upsilon^\bullet(M))$ , defined by (4).

In fact, these are all the possible geometric derivations.

**Lemma 2.6.** *Every geometric derivation is the symmetric derivative corresponding to some affine connection.*

*Proof.* Consider an arbitrary geometric derivation  $D$  and define

$$\nabla := \frac{1}{2} (D + d)|_{\Omega^1(M)},$$

which is a linear map  $\Omega^1(M) \rightarrow \Omega^1(M, T^*M)$ . Moreover, for all  $f \in \mathcal{C}^\infty(M)$  and  $\alpha \in \Omega^1(M)$ ,

$$\begin{aligned} \nabla(f\alpha) &= \frac{1}{2}(D(f\alpha) + d(f\alpha)) = \frac{1}{2}(df \odot \alpha + fD\alpha + df \wedge \alpha + fd\alpha) \\ &= \frac{1}{2}(df \otimes \alpha + \alpha \otimes df + df \otimes \alpha - \alpha \otimes df) + f\nabla\alpha = df \otimes \alpha + f\nabla\alpha, \end{aligned}$$

thus  $\nabla$  is a connection. As  $d\alpha \in \Omega^2(M)$ ,  $\text{sym}(d\alpha) = 0$ , so  $\nabla^s = 2\text{sym} \circ \nabla = D$ .  $\square$

To any connection  $\nabla$  on  $M$  with torsion  $T_\nabla \in \Omega^2(M, TM)$  we can associate the torsion-free connection  $\nabla^0$  given, for  $X, Y \in \mathfrak{X}(M)$ , by

$$\nabla_X^0 Y := \nabla_X Y - \frac{1}{2} T_\nabla(X, Y).$$

**Lemma 2.7.** *Given two affine connections, their symmetric derivatives are the same if and only if their associated torsion-free connections are the same.*

*Proof.* Note that the symmetric derivative corresponding to a connection  $\nabla$  on  $M$  may be explicitly written as

$$(\nabla^s \alpha)(X, Y) = X\alpha(Y) + Y\alpha(X) - \alpha(\nabla_X Y + \nabla_Y X)$$

for all  $\alpha \in \Upsilon^1(M)$  and  $X, Y \in \mathfrak{X}(M)$ . It follows that two connections  $\nabla$  and  $\nabla'$  on  $M$  induce the same symmetric derivative if and only if

$$(5) \quad \nabla_X Y + \nabla_Y X = \nabla'_X Y + \nabla'_Y X.$$

On the other hand, the associated torsion-free connection to  $\nabla$  is given by

$$(6) \quad \nabla_X^0 Y = \frac{1}{2}(\nabla_X Y + \nabla_Y X + [X, Y]),$$

hence  $\nabla^0 = \nabla'^0$  if and only if (5) is satisfied.  $\square$

*Remark 2.8.* Note that, geometrically, having the same torsion-free connection means that they have the same geodesics. It may be seen easily from that both the geodesic equation and the associated torsion-free connection depend solely on the symmetric part of the original affine connection, see (6).



**Proposition 2.9.** *The map  $\nabla \mapsto \nabla^s$  gives an isomorphism of  $\Upsilon^2(M, TM)$ -affine spaces:*

$$\left\{ \begin{array}{c} \text{torsion-free} \\ \text{connections on } M \end{array} \right\} \xleftrightarrow{\sim} \left\{ \begin{array}{c} \text{geometric derivations of the algebra} \\ \text{of symmetric forms on } M \end{array} \right\}.$$

*Proof.* From Lemmas 2.6 and 2.7, there is a bijective correspondence. The set of torsion-free connections on  $M$  is an affine space modeled on  $\Upsilon^2(M, TM)$ . On the other hand, for geometric derivations  $D'$  and  $D$ ,

$$(D' - D)(f\alpha) = df \odot \alpha + fD'\alpha - df \odot \alpha - fD\alpha = f(D' - D)(\alpha)$$

for  $f \in \mathcal{C}^\infty(M)$  and  $\alpha \in \Upsilon^1(M)$ , so the difference between  $D'|_{\Upsilon^1(M)}$  and  $D|_{\Upsilon^1(M)}$  may be viewed as an element of  $\Upsilon^2(M, TM)$ . Since every geometric derivation is uniquely determined by its restriction to 1-forms, they are also a  $\Upsilon^2(M, TM)$ -affine space. The map  $\nabla \mapsto \nabla^s$  is clearly affine and the result follows.  $\square$

Thus, without loss of generality, we can and will mostly consider (apart from some instances in Sections 4, 5 and Appendix B) connections to be torsion free.

*Remark 2.10.* We have seen in the introduction that it is not natural for the replacement of the exterior derivative to square to zero. Actually, no geometric derivation squares to zero: for any  $f \in \mathcal{C}^\infty(M)$ , if  $D \circ D = 0$ , we have

$$0 = D(Df^2) = D(2fDf) = 2Df \odot Df + 2fD(Df) = 2df \odot df,$$

which is a contradiction (as  $\dim M > 0$ ), hence  $D \circ D \neq 0$ . In Appendix A, we show that the same holds for non-geometric derivations.

The unique extension of the symmetric derivative  $\nabla^s$  defined by (4) to  $\Upsilon^\bullet(M)$  is given, for  $\varphi \in \Upsilon^r(M)$ , by

$$\nabla^s \varphi = (r + 1) \text{sym}(\nabla \varphi).$$

Indeed, the endomorphism on the right-hand side is of degree 1, it coincides with the exterior derivative when  $\varphi$  is a function and it reduces to (4) when  $\varphi$  is a 1-form. Moreover, it is a derivation (as the covariant derivative with respect to any vector field is), so it coincides with  $\nabla^s$ .

Explicitly for  $\varphi \in \Upsilon^r(M)$  and  $X_j \in \mathfrak{X}(M)$  we have

$$(7) \quad (\nabla^s \varphi)(X_1, \dots, X_{r+1}) = \sum_{j=1}^{r+1} (\nabla_{X_j} \varphi)(X_1, \dots, \hat{X}_j, \dots, X_{r+1}).$$

**2.3. Geometric interpretation of the symmetric derivative.** We use the relation between symmetric forms on  $M$  and a certain class of smooth functions on  $TM$  to give a geometric interpretation of  $\nabla^s$ .

**Definition 2.11.** We call a smooth function  $\xi \in \mathcal{C}^\infty(TM)$  a **degree- $r$  polynomial in velocities** on  $M$  if it is a *homogeneous function of degree  $r$* , that is, for every  $\lambda \in \mathbb{R}$  and  $u \in TM$ ,

$$\xi(\lambda u) = \lambda^r \xi(u).$$

The direct sum of the spaces of degree- $r$  polynomials for  $r \in \mathbb{N}$  is a subalgebra of  $\mathcal{C}^\infty(TM)$  that is unital and graded. We call its elements **polynomials in velocities** on  $M$ . The next lemma explains this notation.

**Lemma 2.12.** *Let  $\xi \in \mathcal{C}^\infty(TM)$  be a degree- $r$  polynomial in velocities on  $M$ . If  $r = 0$ , there is a unique  $f \in \mathcal{C}^\infty(M)$  such that  $\xi = \text{pr}^* f$ . Whereas, for  $r > 0$ , given a natural coordinate chart  $(TU, \{x^i\} \cup \{v^j\})$ , there is a unique set of functions  $\{\xi_{j_1 \dots j_r}\} \subseteq \mathcal{C}^\infty(U)$  such that  $\xi|_{TU} = (\text{pr}^* \xi_{j_1 \dots j_r}) v^{j_1} \dots v^{j_r}$ .*

*Proof.* Consider a degree-0 polynomial in velocities  $\xi \in \mathcal{C}^\infty(TM)$ . We have  $\xi(\lambda u) = \xi(u)$ , that is,  $\xi$  is constant along the fibres of  $TM$ . Denote by  $0 \in \mathfrak{X}(M)$  the null vector field and define  $f \in \mathcal{C}^\infty(M)$  by  $f := \xi \circ 0$ . The uniqueness of  $f$  follows from the surjectivity of  $\text{pr}$ .

Assume now that  $r > 0$ . By the choice of a natural coordinate chart on  $TM$ , a degree- $r$  polynomial in velocities  $\xi \in \mathcal{C}^\infty(TM)$  restricted to a fibre of  $TM$  corresponds uniquely to a smooth homogeneous function  $\xi_0 : \mathbb{R}^n \rightarrow \mathbb{R}$  of degree  $r$ , where  $n := \dim M$ . It is now enough to check that  $\xi_0$  is a homogeneous polynomial of degree  $r$ . Acting by the partial derivative  $\frac{\partial}{\partial u^i}$  on the relation  $\xi_0(\lambda u^1, \dots, \lambda u^n) = \lambda^r \xi_0(u^1, \dots, u^n)$ , we obtain  $\lambda \frac{\partial \xi_0}{\partial u^i}(\lambda u^1, \dots, \lambda u^n) = \lambda^r \frac{\partial \xi_0}{\partial u^i}(u^1, \dots, u^n)$ , hence  $\frac{\partial \xi_0}{\partial u^i} : \mathbb{R}^n \rightarrow \mathbb{R}$  is a smooth homogeneous function of degree  $r - 1$ . By *Euler's homogenous function theorem*, every smooth homogenous function  $\eta : \mathbb{R}^n \rightarrow \mathbb{R}$  of degree  $r$  satisfies

$$r \eta(u^1, \dots, u^n) = \sum_{i=1}^n u^i \frac{\partial \eta}{\partial u^i}(u^1, \dots, u^n).$$

Since we have already proven the statement for  $r = 0$ , it follows, by induction on  $r$ , that  $\xi_0$  is a homogeneous polynomial of degree  $r$ . Finally, the smoothness of  $\xi$  gives the existence and smoothness of  $\{\xi_{j_1 \dots j_r}\}$   $\square$

The assignment  $\Upsilon^r(M) \rightarrow \mathcal{C}^\infty(TM) : \varphi \mapsto \tilde{\varphi}$ , where, for  $u \in TM$ ,

$$\begin{aligned} \tilde{\varphi}(u) &:= \frac{1}{r!} \varphi(u, \dots, u) && \text{for } r > 0, \\ \tilde{f} &:= \text{pr}^* f && \text{for } r = 0 \end{aligned}$$

gives, by Lemma 2.12 and polarization, an isomorphism of unital graded algebras

$$(8) \quad \Upsilon^\bullet(M) \xrightarrow{\sim} \{ \text{polynomials in velocities on } M \}.$$

We show that the symmetric derivative is closely related to the **geodesic spray** of  $\nabla$ , which is the vector field  $X_\nabla \in \mathfrak{X}(TM)$  given, for  $u \in TM$ , by

$$(X_\nabla)_u = u^h \in T_u TM.$$

The superscript  $h$  here stands for the *horizontal lift* to  $TM$  (for more details, see Appendix B, where the analogous horizontal lift to  $T^*M$  is recalled).

*Remark 2.13.* In natural local coordinates  $(TU, \{x^i\} \cup \{v^j\})$ , we have

$$(X_\nabla)_u = v^i(u) \partial_{x^i}|_u - v^i(u)v^j(u) \Gamma_{ij}^k(\text{pr}(u)) \partial_{v^k}|_u.$$

**Proposition 2.14.** *Let  $\nabla$  be a connection on  $M$ . For every  $\varphi \in \Upsilon^r(M)$  we have*

$$\widetilde{\nabla^s \varphi} = X_\nabla \tilde{\varphi}.$$

*Proof.* Since  $\nabla^s$  and  $X_\nabla$  are derivations of  $\Upsilon^\bullet(M)$  and  $\mathcal{C}^\infty(TM)$  respectively. It is enough to check that the equality holds for functions and 1-forms on  $M$ . For  $f \in \mathcal{C}^\infty(M)$ , we have

$$(X_\nabla \tilde{f})(u) = (X_\nabla \text{pr}^* f)(u) = uf = (df)(u) = \widetilde{\nabla^s f}(u).$$

On the other hand, for  $\alpha \in \Omega^1(M)$  locally given by  $\alpha|_U = \alpha_i dx^i$ , we get

$$\begin{aligned} (X_\nabla \tilde{\alpha})(u) &= (X_\nabla)_u((\text{pr}^* \alpha_k) v^k) = v^i(u) v^j(u) \left( \frac{\partial \alpha_j}{\partial x^i}(\text{pr}(u)) - \alpha_k(\text{pr}(u)) \Gamma_{ij}^k(\text{pr}(u)) \right) \\ &= (\nabla_u \alpha)(u) = \frac{1}{2} (\nabla^s \alpha)(u, u) = \widetilde{\nabla^s \alpha}(u). \end{aligned}$$

□

**2.4. The symmetric Lie derivative.** Once we have a characterization of the symmetric analogues of the exterior derivative, we can make the following definition, drawing inspiration from Cartan's magic formula and using the fact that the commutator of derivations is again a derivation.

**Definition 2.15.** The **symmetric Lie derivative**  $L_X^s \in \text{Der}_0(\Upsilon^\bullet(M))$  corresponding to  $\nabla$  with respect to  $X \in \mathfrak{X}(M)$  is defined by

$$L_X^s := [\iota_X, \nabla^s] = \iota_X \circ \nabla^s - \nabla^s \circ \iota_X.$$

*Remark 2.16.* Note that  $L_X^s f = Xf$  for any  $f \in \mathcal{C}^\infty(M)$  and, thanks to (7), an explicit formula is given, for  $\varphi \in \Upsilon^r(M)$  and  $X, X_j \in \mathfrak{X}(M)$ , by

$$(9) \quad (L_X^s \varphi)(X_1, \dots, X_r) = (\nabla_X \varphi)(X_1, \dots, X_r) - \sum_{j=1}^r \varphi(\nabla_{X_j} X, X_1, \dots, \hat{X}_j, \dots, X_r).$$

The symmetric Lie derivative is related to the covariant and the usual Lie derivatives as follows.

**Proposition 2.17.** *For any connection  $\nabla$  on  $M$  and  $X \in \mathfrak{X}(M)$ ,*

$$(10) \quad \nabla_X^0 = \frac{1}{2} (L_X^s + L_X)$$

*as elements in  $\text{Der}(\Upsilon^\bullet(M))$ .*

*Proof.* It is straightforward to check that the covariant and Lie derivatives define elements in  $\text{Der}(\Upsilon^\bullet(M))$ . It is then enough to check the equality for a function  $f \in \mathcal{C}^\infty(M)$  (which is trivial, as all the terms are  $Xf$ ), and for a 1-form  $\alpha \in \Upsilon^1(M)$ ,

$$\begin{aligned} \frac{1}{2}(L_X + L_X^s)\alpha &= \frac{1}{2}(\iota_X \circ d + d \circ \iota_X + \iota_X \circ \nabla^s - d \circ \iota_X)\alpha \\ &= \iota_X(\text{skew} \circ \nabla^0 + \text{sym} \circ \nabla^0)\alpha = \iota_X \nabla^0 \alpha = \nabla_X^0 \alpha, \end{aligned}$$

where ‘skew’ denotes the usual skew-symmetric projection analogous to ‘sym’.  $\square$

As a consequence, for a torsion-free connection  $\nabla$  and  $\varphi \in \Upsilon^\bullet(M)$  such that  $\nabla\varphi = 0$ , we have

$$(11) \quad L_X^s \varphi = -L_X \varphi.$$

This applies, in particular, to a (pseudo-)Riemannian metric  $\varphi = g$  and  $\nabla = {}^g\nabla$  (the Levi-Civita connection of  $g$ ), which allows us to determine the symmetric derivative corresponding to  ${}^g\nabla$  in an elegant way.

**Proposition 2.18.** *The symmetric derivative corresponding to the Levi-Civita connection  ${}^g\nabla$  of a (pseudo-)Riemannian metric  $g$  is completely determined by*

$${}^g\nabla^s \alpha = L_{g^{-1}\alpha} g$$

for any  $\alpha \in \Upsilon^1(M)$ .

*Proof.* Using the fact that  ${}^g\nabla g = 0$ , and hence  ${}^g\nabla^s g = 0$ , we find

$${}^g\nabla^s \alpha = {}^g\nabla^s(\iota_{g^{-1}\alpha} g) = -(\iota_{g^{-1}\alpha} {}^g\nabla^s - {}^g\nabla^s \iota_{g^{-1}\alpha})g = -L_{g^{-1}\alpha} g.$$

It follows from (11) that  ${}^g\nabla^s \alpha = L_{g^{-1}\alpha} g$ .  $\square$

**2.5. Geometric interpretation of the symmetric Lie derivative.** Given a vector field  $X \in \mathfrak{X}(M)$ , Lie and covariant derivatives at a point  $m \in M$  have infinitesimal formulas in terms, respectively, of the local flow  $\Psi^X$  and the parallel transport  $P^\gamma$  along an integral curve  $\gamma$  of  $X$  starting at  $m$ . For a tensor field  $\tau$ ,

$$(12) \quad (L_X \tau)_m = \lim_{t \rightarrow 0} \frac{1}{t} ((\Psi_t^X)_m^* \tau_{\Psi_t^X(m)} - \tau_m) = \left. \frac{d}{dt} \right|_{t=0} (\Psi_t^X)_m^* \tau_{\Psi_t^X(m)}$$

$$(13) \quad (\nabla_X \tau)_m = \lim_{t \rightarrow 0} \frac{1}{t} (P_{t,0}^\gamma \tau_{\Psi_t^X(m)} - \tau_m) = \left. \frac{d}{dt} \right|_{t=0} P_{t,0}^\gamma \tau_{\Psi_t^X(m)}.$$

The relation (10) gives us a recipe how to write a similar formula also for a symmetric Lie derivative.

**Proposition 2.19.** *Let  $\nabla$  be a connection on  $M$ . For all  $\varphi \in \Upsilon^\bullet(M)$ ,  $X \in \mathfrak{X}(M)$ , and  $m \in M$ , there holds*

$$(L_X^s \varphi)_m = \lim_{t \rightarrow 0} \frac{1}{t} (P_{2t,0}^\gamma (\Psi_{-t}^X)_{\Psi_{2t}^X(m)}^* \varphi_{\Psi_t^X(m)} - \varphi_m) = \left. \frac{d}{dt} \right|_{t=0} P_{2t,0}^\gamma (\Psi_{-t}^X)_{\Psi_{2t}^X(m)}^* \varphi_{\Psi_t^X(m)},$$

where  $\gamma : I \rightarrow M$  is an integral curve of the vector field  $X$  starting at the point  $m$  and  $P^\gamma$  is the  $\nabla^0$ -parallel transport along  $\gamma$ .

*Proof.* Starting from the right-hand side, we have

$$\begin{aligned} & \lim_{t \rightarrow 0} \frac{1}{t} (P_{2t,0}^\gamma (\Psi_{-t}^X)^*_{\Psi_{2t}^X(m)} \varphi_{\Psi_t^X(m)} - \varphi_m) \\ &= \lim_{t \rightarrow 0} \frac{1}{t} (P_{2t,0}^\gamma (\Psi_{-t}^X)^*_{\Psi_{2t}^X(m)} \varphi_{\Psi_t^X(m)} - P_{2t,0}^\gamma \varphi_{\Psi_{2t}^X(m)} + P_{2t,0}^\gamma \varphi_{\Psi_{2t}^X(m)} - \varphi_m) \\ &= 2 \lim_{t \rightarrow 0} \frac{1}{2t} (P_{2t,0}^\gamma \varphi_{\Psi_{2t}^X(m)} - \varphi_m) - \lim_{t \rightarrow 0} \frac{1}{t} P_{2t,0}^\gamma (\varphi_{\Psi_{2t}^X(m)} - (\Psi_{-t}^X)^*_{\Psi_{2t}^X(m)} \varphi_{\Psi_t^X(m)}). \end{aligned}$$

From (13), the first term is  $2(\nabla_X^0 \varphi)_m$ . We focus on the second term, by additivity of the flow it equals

$$\begin{aligned} & \lim_{t \rightarrow 0} \frac{1}{t} P_{2t,0}^\gamma (\Psi_{-2t}^X)^*_{\Psi_{2t}^X(m)} ((\Psi_{2t}^X)^*_m \varphi_{\Psi_{2t}^X(m)} - (\Psi_t^X)^*_m \varphi_{\Psi_t^X(m)}) \\ &= \lim_{t \rightarrow 0} P_{2t,0}^\gamma (\Psi_{-2t}^X)^*_{\Psi_{2t}^X(m)} \left( 2 \frac{1}{2t} ((\Psi_{2t}^X)^*_m \varphi_{\Psi_{2t}^X(m)} - \varphi_m) - \frac{1}{t} ((\Psi_t^X)^*_m \varphi_{\Psi_t^X(m)} - \varphi_m) \right). \end{aligned}$$

That is, by the continuity and since  $P_{0,0}^\gamma = \text{id} = (\Psi_0^X)^*_{\Psi_0^X(m)}$ ,

$$2 \lim_{t \rightarrow 0} \frac{1}{2t} ((\Psi_{2t}^X)^*_m \varphi_{\Psi_{2t}^X(m)} - \varphi_m) - \lim_{t \rightarrow 0} \frac{1}{t} ((\Psi_t^X)^*_m \varphi_{\Psi_t^X(m)} - \varphi_m).$$

Finally, using (12), we have that the second term is

$$2(L_X \varphi)_m - (L_X \varphi)_m = (L_X \varphi)_m.$$


Adding the first and second terms and using (10) we get

$$\lim_{t \rightarrow 0} \frac{1}{t} (P_{2t,0}^\gamma (\Psi_{-t}^X)^*_{\Psi_{2t}^X(m)} \varphi_{\Psi_t^X(m)} - \varphi_m) = 2(\nabla_X^0 \varphi)_m - (L_X \varphi)_m = (L_X^s \varphi)_m.$$

□

We can regard the map  $P_{2t,0}^\gamma \circ (\Psi_{-t}^X)^*_{\Psi_{2t}^X(m)}$  as a  $\nabla^s$ -**parallel transport**: it moves a tensor along an integral curve of a vector field  $X$  first forwards from  $\Psi_t^X(m)$  to  $\Psi_{2t}^X(m)$  via the flow, and after that it takes it backwards to  $m$  by the  $\nabla^0$ -parallel transport, see Figure 1.

*Remark 2.20.* The interpretation in terms of the  $\nabla^s$ -parallel transport of Proposition 2.19 makes it possible to define the symmetric Lie derivative for arbitrary tensor fields. Formula (10) is then satisfied in general.


 FIGURE 1.  $\nabla^s$ -parallel transport.

**2.6. The symmetric bracket.** By analogy with the formula  $\iota_{[X,Y]} = [L_X, \iota_Y]_g$ , the corresponding bracket  $\langle : \rangle_s$  in the symmetric setting should be determined by

$$(14) \quad \iota_{\langle X:Y \rangle_s} = [L_X^s, \iota_Y] = L_X^s \circ \iota_Y - \iota_Y \circ L_X^s.$$

Since the right-hand side is a derivation of degree  $(-1)$  and, by Lemma 2.1, any such derivation is the interior multiplication by a unique vector field, equation (14) determines uniquely the symmetric bracket  $\langle : \rangle_s$ . We can find an explicit expression as, for  $\alpha \in \Upsilon^1(M)$ ,

$$\begin{aligned} L_X^s \iota_Y \alpha - \iota_Y L_X^s \alpha &= X\alpha(Y) - (\nabla_X \alpha)(Y) + \alpha(\nabla_Y X) \\ &= \alpha(\nabla_X Y + \nabla_Y X) = \iota_{(\nabla_X Y + \nabla_Y X)} \alpha. \end{aligned}$$

**Definition 2.21.** The **symmetric bracket** corresponding to a connection  $\nabla$  on  $M$  is given, for  $X, Y \in \mathfrak{X}(M)$ , by

$$(15) \quad \langle X : Y \rangle_s = \nabla_X Y + \nabla_Y X.$$

**Example 2.22.** Let  $g$  be a (pseudo-)Riemannian metric. We have the following formula for the symmetric bracket corresponding to the Levi-Civita connection  ${}^g\nabla$ :

$$(16) \quad \langle X : Y \rangle_s = g^{-1}(L_X g(Y) + L_Y g(X)) - \text{grad}_g g(X, Y),$$

where  $\text{grad}_g : \mathcal{C}^\infty(M) \rightarrow \mathfrak{X}(M)$  is the natural map given by  $\text{grad}_g := g^{-1} \circ d$ .

*Remark 2.23.* Formula (16) can be reinterpreted, by using  $\vartheta := g^{-1}$ , as a symmetric bracket on the cotangent bundle: for  $\alpha, \beta \in \Omega^1(M)$ ,

$$(17) \quad \langle \alpha : \beta \rangle_\vartheta := L_{\vartheta(\alpha)} \beta + L_{\vartheta(\beta)} \alpha - d\vartheta(\alpha, \beta).$$

The bracket in (17) can be defined for an arbitrary  $\vartheta \in \mathfrak{X}_{\text{sym}}^2(M)$ , possibly degenerate. It turns out to be a symmetric analogue of the Lie algebroid bracket on the

cotangent bundle of a Poisson manifold  $(M, \pi)$  (see, for instance, [11]):

$$[\alpha, \beta]_\pi := L_{\pi(\alpha)}\beta - L_{\pi(\beta)}\alpha - d\pi(\alpha, \beta).$$

Analogues of the usual identities for the Lie bracket, the exterior and Lie derivatives also hold here. On the one hand,

$$(18) \quad L_X^s Y = \langle X : Y \rangle_s = L_Y^s X,$$

which follows by applying  $L_X Y = \nabla_X^0 Y - \nabla_Y^0 X$  and (10). On the other hand, from (7) and (9), we have, for  $\varphi \in \Upsilon^r(M)$  and  $X, X_j \in \mathfrak{X}(M)$ ,

$$\begin{aligned} (\nabla^s \varphi)(X_1, \dots, X_{r+1}) &= \sum_{j=1}^{r+1} X_j \varphi(X_1, \dots, \hat{X}_j, \dots, X_{r+1}) \\ &\quad - \sum_{\substack{i,j=1, \\ i < j}}^{r+1} \varphi(\langle X_i : X_j \rangle_s, X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{r+1}), \\ (L_X^s \varphi)(X_1, \dots, X_r) &= X \varphi(X_1, \dots, X_r) - \sum_{j=1}^r \varphi(\langle X : X_j \rangle_s, X_1, \dots, \hat{X}_j, \dots, X_r). \end{aligned}$$

**2.7. Geometric interpretation of the symmetric bracket.** As a consequence of (18) and Proposition 2.19, the symmetric bracket measures the invariance of a vector field with respect to the  $\nabla^s$ -parallel transport. A different interpretation, in terms of a second derivative of commutator of flows, is given in [3, Sec. 3].

We recall now an additional geometric interpretation of the symmetric bracket [21], which describes the properties of involutive distributions.

For a connection  $\nabla$  on  $M$ , a smooth distribution  $\Delta \subseteq TM$  is  **$\nabla$ -geodesically invariant** if each  $\nabla$ -geodesic  $\gamma : I \rightarrow M$  defined on some open interval  $I \subseteq \mathbb{R}$  possesses the property that existence of  $t_0 \in I$  such that  $\dot{\gamma}(t_0) \in \Delta_{\gamma(t_0)}$  implies that  $\dot{\gamma}(t) \in \Delta_{\gamma(t)}$  for all  $t \in I$ . A submanifold  $S \subseteq M$  is called **totally  $\nabla$ -geodesic** if every  $\nabla$ -geodesic on  $M$  that intersects  $S$  is completely contained in  $S$ .

*Remark 2.24.* Note that if a geodesically invariant distribution is, in addition, integrable, it corresponds to a foliation by totally geodesic submanifolds.

The geometric interpretation for the symmetric bracket is analogous to Frobenius theorem for integrable distributions.

**Theorem 2.25** ([21]). *For a connection  $\nabla$  on  $M$ , a smooth distribution  $\Delta \subseteq TM$  with locally constant rank is  $\nabla$ -geodesically invariant if and only if*

$$\langle \Gamma(\Delta) : \Gamma(\Delta) \rangle_s \subseteq \Gamma(\Delta).$$

In [6], the role of the bracket in theoretical mechanics is discussed.

## 3. COMMUTATION RELATIONS OF SYMMETRIC CARTAN CALCULUS

The classical Cartan calculus on differential forms  $\Omega^\bullet(M)$  is characterized by the graded-commutation relations between  $\iota$ ,  $d$  and  $L$ :

$$\begin{aligned} [\iota_X, \iota_Y]_g &= 0, & [d, d]_g &= 0, & [\iota_X, d]_g &= L_X, \\ [L_X, \iota_Y]_g &= \iota_{[X, Y]}, & [L_X, L_Y]_g &= L_{[X, Y]}, & [d, L_X]_g &= 0. \end{aligned}$$

Just as graded derivations are closed under the graded commutator, derivations are closed under the commutator. Thus, for the symmetric Cartan calculus on  $\Upsilon^\bullet(M)$ , we aim to have similar relations between  $\iota$ ,  $\nabla^s$  and  $L^s$  and the usual commutator.

As  $\Upsilon^\bullet(M)$  is the space of symmetric forms on  $M$  we have

$$[\iota_X, \iota_Y] = 0.$$

The analogue of the second relation

$$[\nabla^s, \nabla^s] = 0$$

is also satisfied trivially because of the skew-symmetry of the commutator. The analogues of the third and fourth formulas were used to define the symmetric Lie derivative (Definition 2.15) and derive the symmetric bracket (14), that is,

$$[\iota_X, \nabla^s] = L_X^s, \quad [L_X^s, \iota_Y] = \iota_{\langle X, Y \rangle_s}.$$

For the remaining two identities, we will find that the analogues are more involved. In the case of  $[L_X, L_Y]_g = L_{[X, Y]}$ , it is clear that  $[L_X^s, L_Y^s] \neq L_{\langle X, Y \rangle_s}^s$  since the left-hand side is skew-symmetric on  $X$  and  $Y$ , whereas the right-hand side is symmetric. Their proof will require the extension of the contraction operator and the symmetric derivative (Sections 3.1 and 3.3).

**3.1. The symmetric contraction operator.** Since this is a linear-algebraic operator, we first work on a vector space  $V$  and then globalize it. Consider  $\sigma \in \text{Sym}^k V^* \otimes V$ . The **symmetric contraction operator**

$$\iota_\sigma^s : \text{Sym}^\bullet V^* \rightarrow \text{Sym}^\bullet V^*$$

is the unique degree- $(k-1)$  derivation of  $\text{Sym}^\bullet V^*$  determined by its action on  $\lambda \in \mathbb{R}$  and  $\alpha \in V^*$  by

$$\iota_\sigma^s \lambda := 0, \quad \iota_\sigma^s \alpha := \iota_\sigma \alpha,$$

where by  $\iota_\sigma$  we mean, for  $\varphi \in \text{Sym}^r V^*$ ,

$$(\iota_\sigma \varphi)(X_1, \dots, X_{r+k-1}) := \varphi(\sigma(X_1, \dots, X_k), X_{k+1}, \dots, X_{r+k-1}).$$

We easily see that a generalization of Lemma 2.1 is true. Namely, the map  $\iota^s : \sigma \mapsto \iota_\sigma^s$  gives an isomorphism of graded vector spaces:

$$\text{Sym}^\bullet V^* \otimes V \xrightarrow{\sim} \left\{ \begin{array}{l} \text{derivations of } \text{Sym}^\bullet V^* \text{ that} \\ \text{annihilate scalars } \text{Sym}^0 V^* = \mathbb{R} \end{array} \right\}.$$



**Lemma 3.1.** *Let  $\varphi \in \text{Sym}^r V^*$  and  $\sigma \in \text{Sym}^k V^* \otimes V$ . The symmetric contraction operator is explicitly given by*

$$\iota_\sigma^s \varphi = \frac{(r+k-1)!}{(r-1)!k!} \text{sym } \iota_\sigma \varphi.$$

*Proof.* The endomorphism on the right-hand side is clearly a degree- $(k-1)$  map and coincides with  $\iota_\sigma^s$  on scalars and 1-forms. Therefore, it is enough to check that it is a derivation of  $\text{Sym}^\bullet V^*$ , which is done analogously to Lemma 2.1.  $\square$

Globally, the symmetric contraction operator by an element of  $\Upsilon^k(M, TM)$  (in the sense of Remark 2.3) becomes an element of  $\text{Der}_{k-1}(\Upsilon^\bullet(M))$ .

**3.2. Commutator of symmetric Lie derivatives.** We can now deal with the analogue of the fifth identity.

**Proposition 3.2.** *Let  $\nabla$  be a torsion-free connection on  $M$ . For  $X, Y \in \mathfrak{X}(M)$ ,*

$$(19) \quad [L_X^s, L_Y^s] = L_{[X, Y]}^s + \iota_{2(\nabla_{\nabla_X} Y - \nabla_{\nabla_Y} X - R_\nabla(X, Y))},$$

where  $R_\nabla$  is the Riemann curvature tensor.

*Proof.* As all the terms are derivations of  $\Upsilon^\bullet(M)$ , it is enough to check the equality on functions and 1-forms. Functions are annihilated by any symmetric contraction operator, so the relation is clearly satisfied as  $L_X^s f = Xf$ . For an arbitrary 1-form  $\alpha \in \Upsilon^1(M)$  we have

$$\begin{aligned} & ([L_X^s, L_Y^s]\alpha)(Z) - (L_{[X, Y]}^s\alpha)(Z) \\ &= X(L_Y^s\alpha)(Z) - (L_Y^s\alpha)(\langle X : Z \rangle_s) - Y(L_X^s\alpha)(Z) + (L_X^s\alpha)(\langle Y : Z \rangle_s) \\ &\quad - [X, Y]\alpha(Z) + \alpha(\langle [X, Y] : Z \rangle_s) \\ &= X(Y\alpha(Z)) - X\alpha(\langle Y : Z \rangle_s) - Y\alpha(\langle X : Z \rangle_s) + \alpha(\langle Y : \langle X : Z \rangle_s \rangle_s) \\ &\quad - Y(X\alpha(Z)) + Y\alpha(\langle X : Z \rangle_s) + X\alpha(\langle Y : Z \rangle_s) - \alpha(\langle X : \langle Y : Z \rangle_s \rangle_s) \\ &\quad - [X, Y]\alpha(Z) + \alpha(\langle [X, Y] : Z \rangle_s) \\ &= \alpha(W), \end{aligned}$$

where  $W \in \mathfrak{X}(M)$  is given by  $W := \langle \langle X : Z \rangle_s : Y \rangle_s - \langle X : \langle Y : Z \rangle_s \rangle_s + \langle [X, Y] : Z \rangle_s$ . Explicitly,

$$\begin{aligned} W &= \nabla_Y \nabla_X Z + \nabla_Y \nabla_Z X + \nabla_{\nabla_X Z + \nabla_Z X} Y - \nabla_X \nabla_Y Z \\ &\quad - \nabla_X \nabla_Z Y - \nabla_{\nabla_Y Z + \nabla_Z Y} X + \nabla_{[X, Y]} Z + \nabla_Z [X, Y]. \end{aligned}$$

Using the definition of the curvature tensor and the torsion-freeness of  $\nabla$ , we get

$$W = -R_\nabla(X, Y)Z + R_\nabla(Y, Z)X + R_\nabla(Z, X)Y + 2(\nabla_{\nabla_Z X} Y - \nabla_{\nabla_Z Y} X).$$

Finally, it follows from the algebraic Bianchi identity that

$$([L_X^s, L_Y^s]\alpha)(Z) - (L_{[X, Y]}^s\alpha)(Z) = \alpha(2(\nabla_{\nabla_Z X} Y - \nabla_{\nabla_Z Y} X - R_\nabla(X, Y)Z)).$$

$\square$

**3.3. Extension of the symmetric derivative and the symmetric curvature.** Similarly as we generalized the contraction operator, we generalize also the symmetric derivative.

**Definition 3.3.** Let  $\nabla$  be a connection on  $M$  and  $A \in \Gamma(\text{End } TM)$ . We introduce the  $A$ -**symmetric derivative** of  $\nabla_A^s \in \text{Der}_1(\Upsilon^\bullet(M))$  by the formula

$$\nabla_A^s := [\iota_A^s, \nabla^s].$$

*Remark 3.4.* Note that  $\iota_{\text{id}}^s \varphi = r \varphi$  for every  $\varphi \in \Upsilon^r(M)$ . Therefore,  $\nabla^s = \nabla_{\text{id}_{TM}}^s$ .

The last piece to be introduced is a symmetric analogue of the curvature operator, whose definition will rely on the concept of **second covariant derivative**. This is, for a connection  $\nabla$  on  $M$ , the linear map  $\nabla^2 : \mathfrak{X}(M) \rightarrow \Gamma(\otimes^2 T^*M \otimes TM)$  defined by

$$(\nabla^2 X)(Y, Z) := (\nabla \nabla X)(Y, Z) = \nabla_Y \nabla_Z X - \nabla_{\nabla_Y Z} X.$$

Note that the map is not  $\mathcal{C}^\infty(M)$ -linear. However, when we take its skew-symmetric part and the connection is torsion-free, it is and we recover the Riemann curvature tensor  $R_\nabla \in \Omega^2(M, \text{End } TM)$ :

$$(2 \text{ skew } \nabla^2 X)(Y, Z) = \nabla_Y \nabla_Z X - \nabla_Z \nabla_Y X - \nabla_{[Y, Z]} X = R_\nabla(Y, Z)X.$$

**Definition 3.5.** Let  $\nabla$  be a connection on  $M$ . We introduce the **symmetric curvature operator**  $R_\nabla^s : \mathfrak{X}(M) \rightarrow \Upsilon^2(M, TM)$  by the formula

$$(R_\nabla^s X)(Y, Z) := (2 \text{ sym } \nabla^2 X)(Y, Z) = \nabla_Y \nabla_Z X + \nabla_Z \nabla_Y X - \nabla_{(Y:Z)_s} X.$$

*Remark 3.6.* If  $\nabla$  is a torsion-free connection on  $M$ , we can express the second covariant derivative in terms of the corresponding Riemannian curvature tensor and symmetric curvatures operator:

$$(\nabla^2 X)(Y, Z) = \frac{1}{2}(R_\nabla(Y, Z)X + (R_\nabla^s X)(Y, Z)).$$

A geometric interpretation of the symmetric curvature operator will be given in Remark 4.16.

**3.4. Commutator of symmetric and symmetric Lie derivatives.** We prove now the remaining identity of symmetric Cartan calculus.

**Proposition 3.7.** *Let  $\nabla$  be a torsion-free connection. For  $X \in \mathfrak{X}(M)$ ,*

$$(20) \quad [\nabla^s, L_X^s] = 2\nabla_{\nabla X}^s + \iota_{2 \text{ sym } \iota_X R_\nabla + R_\nabla^s} X.$$

*Proof.* Analogously as in Proposition 3.2, it is enough to check the relation on  $\mathcal{C}^\infty(M)$  and  $\Upsilon^1(M)$ . The relation (20) restricted on functions says

$$2d \circ \iota_X \circ d - \iota_X \circ \nabla^s \circ d = 2\iota_{\nabla X} \circ d.$$

The left-hand side acting on  $f \in \mathcal{C}^\infty(M)$  and evaluated on  $Y \in \mathfrak{X}(M)$  gives

$$\begin{aligned} 2Y(Xf) - (\nabla^s df)(X, Y) &= Y(Xf) - X(Yf) + \langle X : Y \rangle_s f \\ &= ([X, Y] + \langle X : Y \rangle_s) f = 2(\nabla_Y X) f = 2(\iota_{\nabla X} df)(Y), \end{aligned}$$

where the second equality follows from torsion-freeness of  $\nabla$ . For  $\alpha \in \Upsilon^1(M)$ ,

$$\begin{aligned} &([\nabla^s, L_X^s] \alpha)(Y, Y) - 2(\nabla_{\nabla X}^s \alpha)(Y, Y) \\ &= (\nabla^s L_X^s \alpha)(Y, Y) - (L_X^s \nabla^s \alpha)(Y, Y) - 2(\iota_{\nabla X}^s \nabla^s \alpha)(Y, Y) + 2(\nabla^s \iota_{\nabla X} \alpha)(Y, Y) \\ &= 2Y(L_X^s \alpha)(Y) - (L_X^s \alpha)(\langle Y : Y \rangle_s) - X(\nabla^s \alpha)(Y, Y) + 2(\nabla^s \alpha)(\langle X : Y \rangle_s, Y) \\ &\quad - 4(\nabla^s \alpha)(\nabla_Y X, Y) + 4Y\alpha(\nabla_Y X) - 2\alpha(\nabla_{\langle Y : Y \rangle_s} X) \\ &= 2Y(X\alpha(Y)) - 2Y\alpha(\langle X : Y \rangle_s) - X\alpha(\langle Y : Y \rangle_s) + \alpha(\langle X : \langle Y : Y \rangle_s \rangle) \\ &\quad - 2X(Y\alpha(Y)) + X\alpha(\langle Y : Y \rangle_s) + 2\langle X : Y \rangle_s \alpha(Y) + 2Y\alpha(\langle X : Y \rangle_s) \\ &\quad - 2\alpha(\langle \langle X : Y \rangle_s : Y \rangle_s) - 4(\nabla_Y X)\alpha(Y) - 4Y\alpha(\nabla_Y X) + 4\alpha(\langle \nabla_Y X : Y \rangle_s) \\ &\quad + 4Y\alpha(\nabla_Y X) - 2\alpha(\nabla_{\langle Y : Y \rangle_s} X). \end{aligned}$$

Using the fact that  $\nabla$  is torsion-free we get

$$([\nabla^s, L_X^s] \alpha)(Y, Y) - 2(\nabla_{\nabla X}^s \alpha)(Y, Y) = \alpha(W),$$

where  $W \in \mathfrak{X}(M)$  is given by

$$\begin{aligned} W &= \langle X : \langle Y : Y \rangle_s \rangle_s - 2\langle \langle X : Y \rangle_s, Y \rangle_s + 4\langle \nabla_Y X : Y \rangle_s - 2\nabla_{\langle Y : Y \rangle_s} X \\ &= 2\nabla_X \nabla_Y Y + 2\nabla_{\nabla_Y X} X - 2\nabla_{\nabla_X Y + \nabla_Y X} Y - 2\nabla_Y \nabla_X Y \\ &\quad - 2\nabla_Y \nabla_Y X + 4\nabla_{\nabla_Y X} Y + 4\nabla_Y \nabla_Y X - 4\nabla_{\nabla_Y X} X. \end{aligned}$$

It follows from the torsion-freeness of  $\nabla$  that  $W$  may be rewritten in terms of the Riemann curvature tensor and the symmetric curvature operator:

$$W = 2R_{\nabla}(X, Y)Y + (R_{\nabla}^s X)(Y, Y) = (2\text{sym } \iota_X R_{\nabla} + R_{\nabla}^s X)(Y, Y).$$

Finally, it follows from polarization that

$$([\nabla^s, L_X^s] \alpha)(Y, Z) - 2(\nabla_{\nabla X}^s \alpha)(Y, Z) = \alpha((2\text{sym } \iota_X R_{\nabla} + R_{\nabla}^s X)(Y, Z)).$$

□

Identities (19) and (20) involve rather complicated terms. We will be able to say more about them in Section 5.4, once we have described affine morphisms in Section 4.

**3.5. Dependence on the connection.** The symmetric contraction operator introduced in Section 3.1 allows us to describe how the symmetric Cartan calculus varies when we change the connection.

**Proposition 3.8.** *Let  $\nabla$  and  $\nabla'$  be two torsion-free connections on  $M$ , then*

$$\nabla'_X Y = \nabla_X Y - \frac{1}{2}\sigma(X, Y)$$

for a unique  $\sigma \in \Upsilon^2(M, TM)$ . The corresponding symmetric derivatives, symmetric Lie derivatives and symmetric brackets are related as follows:

$$\nabla'^s = \nabla^s + \iota_\sigma^s, \quad L_X'^s = L_X^s + \iota_{\sigma(X, \cdot)}^s, \quad \langle X : Y \rangle_s' = \langle X : Y \rangle_s - \sigma(X, Y).$$

*Proof.* The statement for symmetric brackets is trivial. It is enough to check the other two relations on functions and 1-forms. The equalities on functions follow from that every symmetric contraction operator annihilate functions, every symmetric derivative is geometric and  $L_X^s f = Xf = L_X'^s f$ . The equalities on 1-forms follow by straightforward calculations provided the relation between symmetric brackets.  $\square$

#### 4. AFFINE MANIFOLD MORPHISMS

We study now affine morphisms, which are the natural transformations in presence of a connection. We will see how they interact with symmetric Cartan calculus and also their infinitesimal version. Some of the results of this section are stated for arbitrary connections since they provide stronger versions.

**4.1. Definition of affine morphisms.** Given two manifolds  $M$  and  $M'$ , any diffeomorphism  $\phi : M \rightarrow M'$  intertwines their Cartan calculi in the sense that,

$$\phi_*[X, Y] = [\phi_*X, \phi_*Y], \quad \phi^* \circ d = d \circ \phi^*, \quad \phi^* \circ L_{\phi_*X} = L_X \circ \phi^*.$$

For symmetric Cartan calculus, the notion of affine diffeomorphism will play an analogous role. In this section, connections are not necessarily torsion-free.

**Definition 4.1.** Given two manifolds with connection  $(M, \nabla)$  and  $(M', \nabla')$ , a smooth map  $\phi : M \rightarrow M'$  is an **affine morphism** if  $\phi_*$  commutes with the corresponding parallel transports  $P$  and  $P'$ , that is,

$$\phi_{*\gamma(t_1)} \circ P_{t_0, t_1}^\gamma = P_{t_0, t_1}'^{\phi \circ \gamma} \circ \phi_{*\gamma(t_0)},$$

for any curve  $\gamma : I \rightarrow M$  and  $t_0, t_1 \in I$ . If, in addition,  $\phi$  is a diffeomorphism, we call it **affine diffeomorphism**. We denote the space of affine diffeomorphisms by  $\text{Aff}(\nabla, \nabla')$ , and  $\text{Aff}(\nabla)$  if  $M = M'$  and  $\nabla = \nabla'$ .

It is well known that the group of isometries of a (pseudo-)Riemannian metric is a Lie group of dimension at most  $\frac{1}{2}n(n+1)$ , where  $n := \dim M$ . A similar result is true also for the group  $\text{Aff}(\nabla)$ .

**Theorem 4.2** ([19]). *Let  $\nabla$  be a connection on  $M$  of dimension  $n \in \mathbb{N}$ . The group  $\text{Aff}(\nabla)$  is a Lie group of dimension at most  $n(n+1)$ .*

**Example 4.3.** *In the simplest case, the Euclidean connection  $\nabla^{\text{Euc}}$  on  $\mathbb{R}^n$ , we have  $\text{Aff}(\nabla^{\text{Euc}}) \cong \text{Aff}(n, \mathbb{R}) := \text{GL}(n, \mathbb{R}) \ltimes \mathbb{R}^n$ , thus attaining the maximal dimension.*

**Example 4.4.** *Every isometry between two (pseudo-)Riemannian metrics is an affine diffeomorphism between the corresponding Levi-Civita connections, see e.g. [19]. However, the converse is not true in general as it is clear from Example 4.3.*

A simpler characterization of affine morphisms will prove useful.

**Proposition 4.5.** *A map  $\phi : M \rightarrow M'$  is an affine morphism if and only if*

$$\nabla \circ \phi^* = \phi^* \circ \nabla',$$

where  $\nabla$  is seen as a map  $\Omega^1(M) \rightarrow \Omega^1(M, T^*M)$ .

*Proof.* Recall that if  $P_{t_0, t_1}^\gamma$  is the  $\nabla$ -parallel transport from  $t_0$  to  $t_1$  along  $\gamma$ , its algebraic transpose is the parallel transport from  $t_1$  to  $t_0$  along  $\gamma$  for the dual connection on  $T^*M$ , which we denote by  $P_{t_1, t_0}^{*\gamma}$ . Therefore, a map  $\phi : M \rightarrow M'$  is an affine morphism if and only if

$$(21) \quad P_{t_0, t_1}^{*\gamma} \circ \phi_{\gamma(t_0)}^* = \phi_{\gamma(t_1)}^* \circ P_{t_0, t_1}^{*\phi \circ \gamma}.$$

Consider an arbitrary vector  $u \in TM$  and  $\alpha \in \Omega^1(M')$ . We have,

$$\nabla_u \phi^* \alpha = \lim_{t \rightarrow 0} \frac{1}{t} (P_{t, 0}^{*\gamma} \phi_{\gamma(t)}^* \alpha_{\phi(\gamma(t))} - \phi_{\gamma(0)}^* \alpha_{\phi(\gamma(0))}),$$

where  $\gamma : I \rightarrow M$  is an arbitrary curve such that  $\dot{\gamma}(0) = u$ . Assume first that  $\phi$  is an affine morphism. It follows from (21) that

$$\begin{aligned} \iota_u(\nabla \phi^* \alpha) &= \nabla_u \phi^* \alpha = \lim_{t \rightarrow 0} \frac{1}{t} (\phi_{\gamma(0)}^* P_{t, 0}^{*\phi \circ \gamma} \alpha_{\phi(\gamma(t))} - \phi_{\gamma(0)}^* \alpha_{\phi(\gamma(0))}) \\ &= \phi_{\gamma(0)}^* \lim_{t \rightarrow 0} \frac{1}{t} (P_{t, 0}^{*\phi \circ \gamma} \alpha_{\phi(\gamma(t))} - \alpha_{\phi(\gamma(0))}) = \phi_{\gamma(0)}^* \nabla'_{\phi_* \gamma(0) u} \alpha = \iota_u(\phi^* \nabla' \alpha). \end{aligned}$$

Conversely, assume that  $\nabla \circ \phi^* = \phi^* \circ \nabla'$  is true. Consider an arbitrary curve  $\gamma : I \rightarrow M$ ,  $t_0, t_1 \in I$  and  $\zeta \in T_{\phi(\gamma(t_0))}^* M'$ . To prove that  $\phi$  is an affine morphism, by (21), it is enough to show that the 1-form  $\alpha \in \Gamma(\gamma^* TM)$  defined, for  $t \in I$ , by

$$\alpha(t) := \phi_{\gamma(t)}^* \beta(t), \quad \beta(t) := P_{t_0, t}^{*\phi \circ \gamma} \zeta,$$

satisfies  $\nabla_{\dot{\gamma}} \alpha = 0$ . This follows from  $\nabla'_{\phi_* \dot{\gamma}} \beta = 0$ , as

$$\nabla_{\dot{\gamma}} \alpha = \nabla_{\dot{\gamma}} \phi^* \beta = \phi^* \nabla'_{\phi_* \dot{\gamma}} \beta = 0.$$

□

For affine diffeomorphisms, we can reformulate Proposition 4.5 in terms of the original connection on  $TM$ .

**Proposition 4.6** (e.g., [20]). *A diffeomorphism  $\phi : M \rightarrow M'$  is an affine diffeomorphism if and only if, for all  $X, Y \in \mathfrak{X}(M)$ ,*

$$\phi_*(\nabla_X Y) = \nabla'_{\phi_* X} \phi_* Y$$

Given a diffeomorphism  $\phi : M \rightarrow M'$  and a connection  $\nabla'$  on  $M'$ , the **pullback connection**  $\phi^* \nabla'$  on  $M$  is given for  $X, Y \in \mathfrak{X}(M)$  by

$$(\phi^* \nabla')_X Y := \phi_*^{-1} \nabla'_{\phi_* X} \phi_* Y.$$

**Corollary 4.7.** *Let  $\phi : M \rightarrow M'$  be a diffeomorphism and  $\nabla'$  be a connection on  $M'$ . Then  $\phi \in \text{Aff}(\phi^*\nabla', \nabla')$ .*

*Remark 4.8.* Since  $\phi_*$  preserves the Lie bracket of vector fields, we get

$$T_{\phi^*\nabla'}(X, Y) = \phi_*^{-1}T_{\nabla'}(\phi_*X, \phi_*Y).$$

Therefore,  $\phi^*\nabla'$  is torsion-free if and only if  $\nabla'$  is torsion-free.

**4.2. Relation to symmetric Cartan calculus.** In case of torsion-free connections, we can rephrase Proposition 4.5 using the symmetric derivatives.

**Corollary 4.9.** *Let  $\nabla$  and  $\nabla'$  be torsion-free connections on  $M$  and  $M'$  respectively. A smooth map  $\phi : M \rightarrow M'$  is an affine morphism if and only if*

$$\nabla^s \circ \phi^* = \phi^* \circ \nabla'^s.$$

*Remark 4.10.* Note that if  $\nabla$  and  $\nabla'$  are not torsion-free, we can only claim that if  $\phi : M \rightarrow M'$  is an affine morphism, then  $\nabla^s \circ \phi^* = \phi^* \circ \nabla'^s$ .

The fact that the pullback of a smooth map intertwines two symmetric derivatives has a very clear geometrical interpretation.

**Proposition 4.11.** *Let  $\nabla$  and  $\nabla'$  be connections on  $M$  and  $M'$  respectively. A smooth map  $\phi : M \rightarrow M'$  satisfies  $\nabla^s \circ \phi^* = \phi^* \circ \nabla'^s$  if and only if  $\phi$  is geodesic preserving, that is,  $\gamma$  being a  $\nabla$ -geodesic implies that  $\phi \circ \gamma$  is a  $\nabla'$ -geodesic.*

*Proof.* Consider an arbitrary curve  $\gamma : I \rightarrow M$  and denote the curve  $\phi \circ \gamma$  on  $M'$  by  $\gamma'$ . By a straightforward calculation, for every  $\alpha \in \Upsilon^1(M')$  and  $t \in I$ , one finds

$$(22) \quad (\nabla^s \phi^* \alpha)(\dot{\gamma}(t), \dot{\gamma}(t)) - (\phi^* \nabla'^s \alpha)(\dot{\gamma}(t), \dot{\gamma}(t)) = 2((\phi^* \alpha)(\nabla_{\dot{\gamma}} \dot{\gamma}) - \alpha(\nabla'_{\dot{\gamma}'} \dot{\gamma}'))(t).$$

If the pullback  $\phi^*$  intertwines the symmetric derivatives, we get

$$(\phi^* \alpha)(\nabla_{\dot{\gamma}} \dot{\gamma}) = \alpha(\nabla'_{\dot{\gamma}'} \dot{\gamma}').$$

Clearly,  $\gamma$  being a  $\nabla$ -geodesic implies that  $\gamma' = \phi \circ \gamma$  is a  $\nabla'$ -geodesic.

Conversely, given an arbitrary  $u \in TM$ , there is a  $\nabla$ -geodesic  $\gamma$  such that  $\dot{\gamma}(0) = u$ . It follows from (22) and the fact that  $\gamma'$  is, by assumption, a  $\nabla'$ -geodesic, that

$$(\nabla^s \phi^* \alpha)(u, u) = (\phi^* \nabla'^s \alpha)(u, u).$$

Since symmetric derivatives are geometric derivations of  $\Upsilon^\bullet(M)$ , we obtain, by polarization, that  $\nabla^s \circ \phi^* = \phi^* \circ \nabla'^s$ .  $\square$

The relation between affine diffeomorphisms, geodesic preserving diffeomorphisms, and symmetric Cartan calculus is described by the following result.

**Proposition 4.12.** *Let  $\nabla$  and  $\nabla'$  be connections on  $M$  and  $M'$  and denote the associated torsion-free connections by  $\nabla^0$  and  $\nabla'^0$  respectively. Then the following statements about a diffeomorphism  $\phi : M \rightarrow M'$  are equivalent:*

- (1)  $\phi \in \text{Aff}(\nabla^0, \nabla'^0)$ ;

- (2)  $\phi$  is geodesic preserving;  
 (3)  $\nabla^s \circ \phi^* = \phi^* \circ \nabla'^s$ ;  
 (4)  $L_X^s \circ \phi^* = \phi^* \circ L_{\phi_* X}^s$  for all  $X \in \mathfrak{X}(M)$ ;  
 (5)  $\phi_* \langle X : Y \rangle_s = \langle \phi_* X : \phi_* Y \rangle'_s$  for all  $X, Y \in \mathfrak{X}(M)$ .

*Proof.* The equivalence of (1), (2) and (3) is given by Corollary 4.9 and Proposition 4.11. We see that (3) implies (4) by using the identity  $\iota_X \circ \phi^* = \phi^* \circ \iota_{\phi_* X}$ :

$$\begin{aligned} \phi^* \circ L_{\phi_* X}^s &= \phi^* \circ \iota_{\phi_* X} \circ \nabla'^s - \phi^* \circ \nabla'^s \circ \iota_{\phi_* X} = \iota_X \circ \nabla^s \circ \phi^* - \nabla^s \circ \iota_X \circ \phi^* \\ &= L_X^s \circ \phi^*. \end{aligned}$$

To see that (4) implies (5), simply consider (4) for any  $\alpha \in \Upsilon^1(M')$ , that is,

$$\begin{aligned} 0 &= (\phi^* L_{\phi_* X}^s \alpha)(Y) - (L_X^s \phi^* \alpha)(Y) = \phi^*((L_{\phi_* X}^s \alpha)(\phi_* Y)) - (L_X^s \phi^* \alpha)(Y) \\ &= \phi^*((\phi_* X)\alpha(\phi_* Y)) - \phi^*(\alpha(\langle \phi_* X : \phi_* Y \rangle'_s)) - X(\phi^* \alpha)(Y) + (\phi^* \alpha)(\langle X : Y \rangle_s) \\ &= \phi^*(\alpha(\langle \phi_* X : \phi_* Y \rangle'_s - \phi_* \langle X : Y \rangle_s)). \end{aligned}$$

The equivalence of (5) and (1) follows from Proposition 4.6 and the identities

$$\begin{aligned} \phi_* \nabla_X^0 Y &= \frac{1}{2}(\phi_* \langle X : Y \rangle_s + \phi_* [X, Y]), \\ \nabla_{\phi_* X}^0 \phi_* Y &= \frac{1}{2}(\langle \phi_* X : \phi_* Y \rangle'_s + [\phi_* X, \phi_* Y]). \end{aligned}$$

□

**4.3. Affine vector fields.** The infinitesimal version of an affine diffeomorphism from the manifold to itself is a vector field whose flow is a local 1-parameter subgroup of affine diffeomorphisms. We call such vector field an **affine vector field**. There is a useful equivalent characterization of affine vector fields.

**Proposition 4.13** ([19]). *Let  $\nabla$  be a connection on  $M$ . A vector field  $X \in \mathfrak{X}(M)$  is affine if and only if, for all  $Y, Z \in \mathfrak{X}(M)$ ,*

$$(23) \quad [L_X, \nabla_Y]Z = \nabla_{[X, Y]}Z.$$

**Example 4.14.** *Consider a (pseudo-)Riemannian manifold  $(M, g)$ . A **Killing vector field** of  $g$  is a vector field on  $M$  whose flow is a local 1-parameter subgroup of isometries. Equivalently,  $X \in \mathfrak{X}(M)$  is a Killing vector field if  $L_X g = 0$ . In particular, every Killing vector field is an affine vector field for the corresponding Levi-Civita connection. However, the converse is not true in general. For instance, the vector field  $X := x^i \partial_{x^i} \in \mathfrak{X}(\mathbb{R}^n)$  is not a Killing vector field for the Euclidean metric  $g^{\text{Euc}}$  as*

$$L_X g^{\text{Euc}} = 2g^{\text{Euc}} \neq 0,$$

*but it is an affine vector field for the Euclidean connection.*

If the connection is torsion-free, there are three more equivalent formulae characterizing affine vector fields that are closely related to symmetric Cartan calculus and will also be relevant in Section 5.4.

**Proposition 4.15.** *Let  $\nabla$  be a torsion-free connection on  $M$ . A vector field  $X \in \mathfrak{X}(M)$  is an affine vector field if and only if one of the following equivalent statements is true*

- (1)  $2 \operatorname{sym} \iota_X R_\nabla + R_\nabla^s X = 0$ ,
- (2)  $[X : \langle Y : Z \rangle_s] = \langle [X, Y] : Z \rangle_s + \langle Y : [X, Z] \rangle_s$  for all  $Y, Z \in \mathfrak{X}(M)$ .
- (3)  $[\nabla^s, L_X] = 0$ .

*Proof.* Using the fact that  $\nabla$  is torsion-free, we can rewrite the relation (23) as

$$\nabla_X \nabla_Y Z - \nabla_{\nabla_Y Z} X - \nabla_Y \nabla_X Z + \nabla_Y \nabla_Z X = \nabla_{[X, Y]} Z.$$

Taking all the terms on the left-hand side yields  $R_\nabla(X, Y)Z + (\nabla^2 X)(Y, Z) = 0$ . This is thanks to Remark 3.6 equivalent to

$$2R_\nabla(X, Y)Z + R_\nabla(Y, Z)X + (R_\nabla^s X)(Y, Z) = 0.$$

Finally, using the algebraic Bianchi identity yields

$$R_\nabla(X, Y)Z + R_\nabla(X, Z)Y + (R_\nabla^s X)(Y, Z) = 0.$$

On the other hand, by Proposition 2.17, see also Remark 2.20, we can replace all the covariant derivatives in (23) with the Lie and symmetric Lie derivatives. Therefore, (23) is satisfied if and only if

$$[L_X, L_Y]Z + [L_X, L_Y^s]Z = L_{[X, Y]}Z + L_{[X, Y]}^s Z.$$

Using the Jacobi identity for the Lie bracket of vector fields yields

$$[L_X, L_Y^s]Z = L_{[X, Y]}^s Z,$$

which can be expressed in terms of the Lie and symmetric bracket as

$$[X, \langle Y : Z \rangle_s] - \langle Y : [X, Z] \rangle_s = \langle [X, Y] : Z \rangle_s.$$

Clearly  $[\nabla^s, L_X]|_{\mathcal{C}^\infty(M)} = [d, L_X]_{\mathfrak{g}}|_{\mathcal{C}^\infty(M)} = 0$ . Moreover, for the commutator restricted on 1-forms, we have

$$\begin{aligned} & \iota_Y \circ [\nabla^s, L_X]|_{\Gamma^1(M)} \\ &= \iota_Y \circ \nabla^s \circ L_X - \iota_Y \circ L_X \circ \nabla^s \\ &= L_Y^s \circ L_X + \nabla^s \circ \iota_Y \circ L_X + \iota_{[X, Y]} \circ \nabla^s - L_X \circ \iota_Y \circ \nabla^s \\ &= L_Y^s \circ L_X - L_X \circ L_Y^s + L_{[X, Y]}^s + \nabla^s \circ \iota_{[X, Y]} + \nabla^s \circ \iota_Y \circ L_X - L_X \circ \nabla^s \circ \iota_Y. \end{aligned}$$

Since  $\nabla^s$  is geometric, we can replace it with the exterior derivative  $d$ . Using the identities  $[d, L_X]_{\mathfrak{g}} = 0$  and  $[L_X, \iota_Y]_{\mathfrak{g}} = \iota_{[X, Y]}$  yields

$$(\iota_Y \circ [\nabla^s, L_X])|_{\Gamma^1(M)} = L_{[X, Y]}^s - [L_X, L_Y^s].$$

By a straightforward calculation one finds

$$([\nabla^s, L_X]\alpha)(Y, Z) = \alpha([X, \langle Y : Z \rangle_s] - \langle Y : [X, Z] \rangle_s - \langle [X, Y] : Z \rangle_s).$$

□



*Remark 4.16.* If, in particular,  $\nabla$  is flat and torsion-free,  $X \in \mathfrak{X}(M)$  is an affine vector field if and only if  $R_{\nabla}^s X = 0$ , which gives a geometric interpretation to symmetric curvature operator.

Note that the expression on the left-hand side of (1) in Proposition 4.15 already appeared in Proposition 3.7. Therefore, we have the following.

**Corollary 4.17.** *Let  $\nabla$  be a torsion-free connection on  $M$ , we have that  $X \in \mathfrak{X}(M)$  is an affine vector field if and only if*

$$[\nabla^s, L_X^s] = 2\nabla_{\nabla X}^s.$$

## 5. THE PATTERSON-WALKER METRIC

The complicated relations obtained in Section 3 conceal a link of symmetric Cartan calculus with the so-called Patterson-Walker metric (originally called the Riemannian extension in [22]). We first reintroduce this metric from the viewpoint of connections, then express it in terms of symmetric Cartan calculus and finally spell out the link to the relations of Section 3. Although the statements of the main results are neat, the proofs require some heavy lifting.

**5.1. Definition of the Patterson-Walker metric.** For an arbitrary connection  $\nabla$  on  $M$ , the induced connection on the cotangent bundle  $\text{pr} : T^*M \rightarrow M$  gives the decomposition

$$T(T^*M) = \mathcal{H}_{\nabla} \oplus \mathcal{V} \cong \text{pr}^*TM \oplus \text{pr}^*T^*M = \text{pr}^*(TM \oplus T^*M)$$

and the  $\mathcal{C}^\infty(M)$ -module morphism  $\phi_{\nabla} : \Gamma(TM \oplus T^*M) \rightarrow \mathfrak{X}(T^*M)$ .

*Remark 5.1.* The structure of  $\mathcal{C}^\infty(M)$ -module on  $\mathfrak{X}(T^*M)$  is given, for  $f \in \mathcal{C}^\infty(M)$  and  $\mathcal{X} \in \mathfrak{X}(T^*M)$ , by  $f \cdot \mathcal{X} := (\text{pr}^*f) \mathcal{X}$ . The morphism  $\phi_{\nabla}$  is given by pullback: for  $\zeta \in T^*M$  and  $a \in \Gamma(TM \oplus T^*M)$ ,

$$(\phi_{\nabla} a)(\zeta) := a(\text{pr}(\zeta)).$$

Explicitly,  $\phi_{\nabla}|_{\mathfrak{X}(M)}$  is the horizontal lift, which we will denote  $X^h := \phi_{\nabla} X$ , and  $\phi_{\nabla}|_{\Omega^1(M)}$  is the vertical lift, which we will denote  $\alpha^v := \phi_{\nabla} \alpha$ . More details about lifts to the cotangent bundle are found in Appendix B.

The bundle  $TM \oplus T^*M$  comes equipped with a canonical symmetric pairing

$$(24) \quad \langle X + \alpha, Y + \beta \rangle_+ := \alpha(Y) + \beta(X).$$

Since pullback sections locally generate the entire space of sections, the image of  $\phi_{\nabla}$  generates  $\mathfrak{X}(T^*M)$  locally. Therefore, we can bring  $\langle \cdot, \cdot \rangle_+$  to  $T^*M$ :

**Definition 5.2.** Let  $\nabla$  be a connection on  $M$ . **The Patterson-Walker metric**  $g_{\nabla}$  is the split-signature metric on  $T^*M$  determined by

$$g_{\nabla}(\phi_{\nabla} a, \phi_{\nabla} b) := \text{pr}^* \langle a, b \rangle_+.$$

Different connections can induce different Patterson-Walker metrics. Let us show when the Patterson-Walker metrics of two different connections coincide. First, we need to prove two lemmas.

**Lemma 5.3.** *Given two connections on  $M$  that differ by  $\tau \in \Gamma(\otimes^2 T^*M \otimes TM)$ , their corresponding horizontal lifts  $(\ )^h$  and  $(\ )^{h'}$  are related as follows:*

$$X^{h'} = X^h + (\iota_X \tau)^v,$$

where  $v$  denotes the vertical lift of sections of  $\text{End } TM$  (see Appendix B).

*Proof.* By Lemma B.7, we have

$$X^{h'} Y^v = (\nabla'_{X'} Y)^v = (\nabla_X Y + \tau(X, Y))^v = (X^h + (\iota_X \tau)^v) Y^v$$

for any  $Y \in \mathfrak{X}(M)$ . Therefore, by Proposition B.2,  $X^{h'} = X^h + (\iota_X \tau)^v$ .  $\square$

**Lemma 5.4.** *Let  $\nabla$  be a connection on  $M$ . For  $A \in \Gamma(\text{End } TM)$ , we have*

$$g_{\nabla}(A^v, \alpha^v) = 0, \quad g_{\nabla}(A^v, X^h) = (AX)^v.$$

*Proof.* The first identity follows easily from the fact that vertical subbundle is isotropic. For the second one, in natural coordinates  $(T^*U, \{x^i\} \cup \{p_j\})$ , we have

$$A|_U = A_j^i dx^j \otimes \partial_{x^i}, \quad A^v|_{T^*U} = p_i A_j^i \partial_{p_j} = p_i (\iota_{dx^i} A)^v.$$

Therefore,  $g_{\nabla}(A^v, X^h)|_{T^*U} = p_i dx^i (AX) = (AX)^v|_{T^*U}$ .  $\square$

**Proposition 5.5.** *Given two connections on  $M$ , their Patterson-Walker metrics coincide if and only if the associated torsion-free connections are the same.*

*Proof.* Let  $\nabla$  and  $\nabla'$  be two connections on  $M$  related to each other by the difference tensor field  $\tau \in \Gamma(\otimes^2 T^*M \otimes TM)$ . Clearly,  $g_{\nabla}|_{\text{Sym}^2 \mathcal{V}} = 0 = g_{\nabla'}|_{\text{Sym}^2 \mathcal{V}}$ . Thanks to Lemma 5.3 and that  $\mathcal{V}$  is isotropic, we find

$$g_{\nabla}(X^{h'}, \beta^v) = g_{\nabla}(X^h + (\iota_X \tau)^v, \beta^v) = \text{pr}^* \beta(X) = g_{\nabla'}(X^{h'}, \beta^v).$$

Taking into account the fact that also the horizontal subbundle is isotropic,

$$g_{\nabla}(X^{h'}, Y^{h'}) = g_{\nabla}(X^h + (\iota_X \tau)^v, Y^h + (\iota_Y \tau)^v) = g_{\nabla}(X^h, (\iota_Y \tau)^v) + g_{\nabla}((\iota_X \tau)^v, Y^h).$$

It follows from Lemma 5.4 that

$$g_{\nabla}(X^{h'}, Y^{h'}) = (\tau(Y, X) + \tau(X, Y))^v,$$

which vanishes (i.e., it is equal to  $g_{\nabla'}(X^{h'}, Y^{h'})$ ) if and only if  $\tau \in \Omega^2(M, TM)$ . Equivalently,  $\nabla'^0 = \nabla^0$  because

$$\nabla_X^0 Y - \nabla'_X Y = \frac{1}{2}(\langle X : Y \rangle'_s - \langle X : Y \rangle_s) = \frac{1}{2}(\tau(X, Y) + \tau(Y, X)).$$

$\square$

Thanks to Proposition 5.5, we can, without loss of generality, consider only torsion-free connections, as we do for symmetric Cartan calculus.

*Remark 5.6.* In natural local coordinates  $(T^*U, \{x^j\} \cup \{p_j\})$  on  $T^*M$ , Definition 5.2 gives that the Patterson-Walker metric takes the form

$$(25) \quad g_\nabla|_{T^*U} = dp_i \odot dx^i - p_k(\text{pr}^*\Gamma_{ij}^k) dx^i \odot dx^j,$$

where  $\{\Gamma_{ij}^k\}$  are the Christoffel symbols of  $\nabla$  in the chart  $(U, \{x^j\})$ . This recovers the original definition in [22].

**5.2. Reformulation using the symmetric Cartan calculus.** We use a natural lift of the connection  $\nabla$  from  $M$ : the connection  $\hat{\nabla}$  on the manifold  $T^*M$  that is uniquely determined by

$$(26) \quad \begin{aligned} \hat{\nabla}_{X^h} Y^h &:= (\nabla_X Y)^h, & \hat{\nabla}_{\alpha^v} X^h &:= 0, \\ \hat{\nabla}_{X^h} \alpha^v &:= (\nabla_X \alpha)^v, & \hat{\nabla}_{\alpha^v} \beta^v &:= 0. \end{aligned}$$

With the lift  $\hat{\nabla}$  we can prove the following.

**Theorem 5.7.** *For any connection  $\nabla$  on  $M$ , there holds*

$$(27) \quad \hat{\nabla}^s \alpha_{\text{can}} = g_\nabla,$$

where  $\alpha_{\text{can}}$  is the canonical 1-form on  $T^*M$ .

*Proof.* We use the properties of the lifts that are proved in Appendix B. By Lemmas B.6 and B.7, we find

$$\begin{aligned} X^h \alpha_{\text{can}}(Y^h) &= X^h Y^v = (\nabla_X Y)^v = \alpha_{\text{can}}((\nabla_X Y)^h), \\ \beta^v \alpha_{\text{can}}(X^h) &= \beta^v X^v = \text{pr}^* \beta(X). \end{aligned}$$

Using these together with the fact that  $\alpha_{\text{can}}$  annihilates  $\mathcal{V}$ , we get

$$\begin{aligned} (\hat{\nabla}^s \alpha_{\text{can}})(X^h, Y^h) &= \alpha_{\text{can}}(\langle X : Y \rangle_s^h - \langle X^h : Y^h \rangle_{\hat{\nabla}^s}), \\ (\hat{\nabla}^s \alpha_{\text{can}})(\alpha^v, \beta^v) &= -\alpha_{\text{can}}(\langle \alpha^v : \beta^v \rangle_{\hat{\nabla}^s}), \\ (\hat{\nabla}^s \alpha_{\text{can}})(X^h, \beta^v) &= \text{pr}^* \beta(X) - \alpha_{\text{can}}(\langle X^h : \beta^v \rangle_{\hat{\nabla}^s}), \end{aligned}$$

where  $\langle : \rangle_{\hat{\nabla}^s}$  stands for the symmetric bracket corresponding to  $\hat{\nabla}$ . The terms with  $\alpha_{\text{can}}$  on the right-hand sides of the above equations vanish by the definition of the connection  $\hat{\nabla}$  and the result follows.  $\square$

Note that the connection  $\hat{\nabla}$  defined by (26) is not the unique connection on  $T^*M$  satisfying (27). To start with, we can take any other connection on  $T^*M$  with the same associated torsion-free affine connection. However, there is actually much more freedom.

**Lemma 5.8.** *A connection  $\mathcal{D}$  on  $T^*M$  satisfies (27) if and only if*

$$(28) \quad \langle X^h + \alpha^v : Y^h + \beta^v \rangle_{\mathcal{D}^s} - \langle X : Y \rangle_s^h \in \Gamma(\mathcal{V}),$$

where  $\langle : \rangle_{\mathcal{D}^s}$  stands for the symmetric bracket corresponding to  $\mathcal{D}$ .

*Proof.* If we follow the same approach as in the proof of Theorem 5.7, we find that the claim is true because the kernel of  $\alpha_{\text{can}}$  is precisely the vertical subbundle.  $\square$

As both  $g_{\nabla}$  and  $\hat{\nabla}$  are naturally associated to the connection  $\nabla$  on  $M$ , it is reasonable to wonder what the relation between  $\hat{\nabla}$  and the Levi-Civita connection of  $g_{\nabla}$  is. We first compute the torsion of the connection  $\hat{\nabla}$ .

**Lemma 5.9.** *Let  $\nabla$  be a connection on  $M$ . For the corresponding connection  $\hat{\nabla}$  on  $T^*M$ , we have that  $\hat{\nabla}g_{\nabla} = 0$ ,  $T_{\hat{\nabla}}|_{\mathcal{H}_{\nabla}\wedge\nabla} = 0$ ,  $T_{\hat{\nabla}}|_{\mathcal{V}\wedge\nabla} = 0$  and*

$$T_{\hat{\nabla}}(X^{\text{h}}, Y^{\text{h}}) = T_{\nabla}(X, Y)^{\text{h}} - R_{\nabla}(X, Y)^{\text{v}}.$$

*Proof.* The claim for the torsion follows directly from Lemma B.8.

For  $\hat{\nabla}g_{\nabla} = 0$ , we use the shorthand notation  $a, b, c \in \Gamma(TM \oplus T^*M)$  and

$$(29) \quad \nabla_{X+\alpha}^{\text{gen}}(Y + \beta) := \nabla_X Y + \nabla_X \beta.$$

The superscript ‘gen’ will be explained in Section 5.3. We also use  $\text{pr}_1$  and  $\text{pr}_2$  for the projections to  $TM$  and  $T^*M$ , respectively.

With this notation, we have the following for  $\nabla^{\text{gen}}$ :

$$(\text{pr}_1 a)\langle b, c \rangle_+ - \langle \nabla_a^{\text{gen}} b, c \rangle_+ - \langle b, \nabla_a^{\text{gen}} c \rangle_+ = 0.$$

Therefore, it follows from (32) that

$$\begin{aligned} & (\hat{\nabla}g_{\nabla})(\phi_{\nabla}a, \phi_{\nabla}b, \phi_{\nabla}c) \\ &= (\phi_{\nabla}a)g_{\nabla}(\phi_{\nabla}b, \phi_{\nabla}c) - g_{\nabla}(\hat{\nabla}_{\phi_{\nabla}a}\phi_{\nabla}b, \phi_{\nabla}c) - g_{\nabla}(\phi_{\nabla}b, \hat{\nabla}_{\phi_{\nabla}a}\phi_{\nabla}c) \\ &= ((\text{pr}_1 a)^{\text{h}} + (\text{pr}_2 a)^{\text{v}})\text{pr}^*\langle b, c \rangle_+ - g_{\nabla}(\phi_{\nabla}\nabla_a^{\text{gen}}b, \phi_{\nabla}c) - g_{\nabla}(\phi_{\nabla}b, \phi_{\nabla}\nabla_a^{\text{gen}}c), \end{aligned}$$

By Lemma B.6,

$$(\hat{\nabla}g_{\nabla})(\phi_{\nabla}a, \phi_{\nabla}b, \phi_{\nabla}c) = \text{pr}^*((\text{pr}_1 a)\langle b, c \rangle_+ - \langle \nabla_a^{\text{gen}} b, c \rangle_+ - \langle b, \nabla_a^{\text{gen}} c \rangle_+) = 0.$$

$\square$

We can now slightly modify  $\hat{\nabla}$  to recover the Levi-Civita connection of  $g_{\nabla}$ . Recall that we can restrict to the torsion-free case without loss of generality.

**Proposition 5.10.** *Let  $\nabla$  be a torsion-free connection on  $M$ . Then the connection  $\bar{\nabla}$  on  $T^*M$  given by*

$$\bar{\nabla}_{\phi_{\nabla}a}\phi_{\nabla}b := \hat{\nabla}_{\phi_{\nabla}a}\phi_{\nabla}b - (R_{\nabla}(\text{pr}_1 b, \cdot)(\text{pr}_1 a))^{\text{v}}.$$

*is the Levi-Civita connection of the Patterson-Walker metric  $g_{\nabla}$ .*

*Proof.* It follows directly from Lemma 5.9 that  $T_{\bar{\nabla}}|_{\mathcal{H}_{\nabla}\wedge\nabla} = 0$ ,  $T_{\bar{\nabla}}|_{\mathcal{V}\wedge\nabla} = 0$  and

$$T_{\bar{\nabla}}(X^{\text{h}}, Y^{\text{h}}) = T_{\nabla}(X, Y)^{\text{h}} - R_{\nabla}(X, Y)^{\text{v}} - (R_{\nabla}(Y, \cdot)X)^{\text{v}} + (R_{\nabla}(X, \cdot)Y)^{\text{v}},$$

which vanishes by the torsion-freeness of  $\nabla$  and the algebraic Bianchi identity. Using  $\hat{\nabla}g_\nabla = 0$ , see Lemma 5.9, and that  $\mathcal{V}$  is isotropic, we find

$$\begin{aligned} & (\bar{\nabla}g_\nabla)(\phi_\nabla a, \phi_\nabla b, \phi_\nabla c) \\ &= g_\nabla((R_\nabla(\text{pr}_1 b, \text{pr}_1 a))^v, (\text{pr}_1 c)^h) + g_\nabla((\text{pr}_1 b)^h, (R_\nabla(\text{pr}_1 c, \text{pr}_1 a))^v). \end{aligned}$$

Finally, using Lemma 5.4, we find

$$(\bar{\nabla}g_\nabla)(\phi_\nabla a, \phi_\nabla b, \phi_\nabla c) = (R_\nabla(\text{pr}_1 b, \text{pr}_1 c)\text{pr}_1 a + R_\nabla(\text{pr}_1 c, \text{pr}_1 b)\text{pr}_1 a)^v = 0.$$

□

A subtle point here is that the connection  $\bar{\nabla} = {}^{g_\nabla}\nabla$  is not the associated torsion-free connection to  $\hat{\nabla}$ . But it clearly satisfies condition (28) in Lemma 5.8, so

$$\bar{\nabla}^s \alpha_{\text{can}} = g_\nabla.$$

*Remark 5.11.* Another reformulation of the Patterson-Walker metric using the symmetric bracket can be directly deduced from [27, §17], but it involves the notions of vertical lift  $X^v \in \mathcal{C}^\infty(T^*M)$  and complete lift  $X^c \in \mathfrak{X}(T^*M)$  of a vector field  $X \in \mathfrak{X}(M)$ . They can be defined by  $X^v(\zeta) := \zeta(X_{\text{pr}(\zeta)})$  for  $\zeta \in T^*M$  and  $X^c = -\text{Ham } X^v$  (see Appendix B for more details). With these notions, one has that  $g_\nabla$  is determined (thanks to Proposition B.3) by

$$(30) \quad g_\nabla(X^c, Y^c) = -\langle X : Y \rangle_s^v.$$

Note the striking similarity between the Patterson-Walker metric and the canonical 2-form  $\omega_{\text{can}}$ :

$$\omega_{\text{can}}|_{T^*U} = dp_i \wedge dx^i, \quad d\alpha_{\text{can}} = \omega_{\text{can}}, \quad \omega_{\text{can}}(X^c, Y^c) = -[X, Y]^v.$$

This analogy is explained by the fact that  $\omega_{\text{can}}$  may be alternatively defined by

$$\omega_{\text{can}}(\phi_\nabla a, \phi_\nabla b) = \text{pr}^* \langle a, b \rangle_-,$$

where  $\langle \cdot, \cdot \rangle_-$  is the canonical skew-symmetric pairing on  $TM \oplus T^*M$ :

$$(31) \quad \langle X + \alpha, Y + \beta \rangle_- := \alpha(Y) - \beta(X).$$

Unlike the Patterson-Walker metric, the canonical 2-form is independent of the choice of torsion-free connection on  $M$ . However, considering connections with non-vanishing torsion gives rise to different non-degenerate 2-forms on  $T^*M$ , which are closed if and only if the connection is torsion-free.

**5.3. Relation to generalized geometry.** We briefly hint the relation to generalized geometry (in the sense of [17, 14]), which will be explored in forthcoming works. This section can be skipped on a first reading.

To start with, the bundle  $TM \oplus T^*M$  and the canonical symmetric pairing (24) is the starting point of generalized geometry, sometimes referred to as the generalized tangent bundle. The skew-symmetric pairing (31) is equally canonical, although its role has remained largely unexplored (we will actually explore it in future work).

Using the language of generalized geometry, we can justify the naturality of the connection  $\hat{\nabla}$  defined by (26). The key concept is that of a generalized connection (see, e.g., [15], although this concept goes back to the unpublished notes [2]). For  $E := TM \oplus T^*M$ , a generalized connection is a linear operator

$$D : \Gamma(E) \rightarrow \Gamma(E^* \otimes E)$$

such that, with the notation  $a, b \in \Gamma(E)$  and  $f \in C^\infty(M)$ ,

$$D(fa) = fDa + df \otimes a, \quad d\langle a, b \rangle_+ = \langle Da, b \rangle_+ + \langle a, Db \rangle_+.$$

For any connection  $\nabla$  on  $M$ , the connection  $\nabla^{\text{gen}}$  defined by (29) in the proof of Lemma 5.9 is a natural generalized connection (and hence the superscript ‘gen’) associated to the connection  $\nabla$ .

With this language,  $\hat{\nabla}$  is then fully determined by the natural formula

$$(32) \quad \hat{\nabla}_{\phi_{\nabla} a} \phi_{\nabla} b = \phi_{\nabla} (\nabla_a^{\text{gen}} b).$$

*Remark 5.12.* Note that this construction can be easily generalized by replacing  $\nabla^{\text{gen}}$  with an arbitrary generalized connection, which would lead, through Theorem 5.7, to a generalization of the Patterson-Walker metric.

**5.4. Killing vector fields for the Patterson-Walker metric.** As we have seen, symmetric Cartan calculus and the Patterson-Walker metrics are closely intertwined. Therefore, given a torsion-free connection, we expect that the symmetries of the Patterson-Walker metric also become symmetries of the corresponding symmetric Cartan calculus. To explore this, we will use the various lifts to the cotangent bundle that we have introduced (again, we refer to Appendix B for more details).

**Definition 5.13.** Let  $g$  be a (pseudo-)Riemannian metric on  $M$ . A vector field  $X \in \mathfrak{X}(M)$  is called a **gradient Killing vector field** of  $g$  if there is  $f \in C^\infty(M)$  such that  $X = \text{grad}_g f$  and, moreover, it is a Killing vector field for  $g$ .

**Lemma 5.14.** Let  $\nabla$  be a connection on  $M$  and denote by  $\text{grad}_{\nabla}$  the gradient map of  $g_{\nabla}$ . The vector field  $\text{grad}_{\nabla} \xi$  is vertical for a function  $\xi \in C^\infty(T^*M)$  if and only if  $\xi = \text{pr}^* f$  for some  $f \in C^\infty(M)$ .

*Proof.* Given a function  $\xi \in C^\infty(T^*M)$ , the vector field  $\text{grad}_{\nabla} \xi$  is vertical if and only if, for any  $\alpha^v$ ,  $0 = g_{\nabla}(\text{grad}_{\nabla} \xi, \alpha^v) = \alpha^v \xi$ , which is equivalent to that  $\xi$  does not depend on fibre coordinates, hence  $\xi = \text{pr}^* f$  for some  $f \in C^\infty(M)$ .  $\square$

**Proposition 5.15.** Let  $\nabla$  be a connection on  $M$ . The vertical lift of a 1-form  $\alpha \in \Upsilon^1(M)$  is a Killing vector field for  $g_{\nabla}$  if and only if  $\nabla^s \alpha = 0$ . It is, in addition, a gradient Killing vector field if and only if  $\alpha$  is exact.

*Proof.* A 1-form  $\alpha \in \Omega^1(M)$  induces a Killing vector field for  $g_{\nabla}$  through the vertical lift if and only if all the components  $(L_{\alpha^v} g_{\nabla})|_{\text{Sym}^2 \mathcal{H}_{\nabla}}$ ,  $(L_{\alpha^v} g_{\nabla})|_{\mathcal{H}_{\nabla} \otimes \mathcal{V}}$  and  $(L_{\alpha^v} g_{\nabla})|_{\text{Sym}^2 \mathcal{V}}$ , vanish. The latter vanishes simply because the vertical subbundle

is isotropic and its space of sections is an ideal for the Lie bracket of vector fields. The mixed component vanishes for the same reason and by Lemma B.6. However, for the horizontal component we find

$$(L_{\alpha^v}g_{\nabla})(X^h, Y^h) = \alpha^v g_{\nabla}(X^h, Y^h) - g_{\nabla}([\alpha^v, X^h], Y^h) - g_{\nabla}(X^h, [\alpha^v, Y^h]).$$

Since the horizontal subbundle is isotropic and  $[X^h, \alpha^v] = (\nabla_X \alpha)^v$ , we get

$$(L_{\alpha^v}g_{\nabla})(X^h, Y^h) = \text{pr}^*((\nabla_X \alpha)(Y) + (\nabla_Y \alpha)(X)) = \text{pr}^*((\nabla^s \alpha)(X, Y)).$$

Hence,  $\alpha^v$  is a Killing vector field for  $g_{\nabla}$  if and only if  $\nabla^s \alpha = 0$ . It follows from Lemma 5.14 that the vertical lift of a 1-form is a gradient vector field if and only if it is equal to  $\text{grad}_{\nabla} \text{pr}^* f$  for some  $f \in \mathcal{C}^\infty(M)$ . By Lemma B.6, for  $X \in \mathfrak{X}(M)$ ,

$$g_{\nabla}(\text{grad}_{\nabla} \text{pr}^* f, X^h) = X^h \text{pr}^* f = \text{pr}^* X f = \text{pr}^*(df)(X),$$

hence  $\text{grad}_{\nabla} \text{pr}^* f = (df)^v$ . So,  $\alpha^v$  is equal to a gradient if and only if  $\alpha$  is exact.  $\square$

*Remark 5.16.* The condition  $\nabla^s \alpha = 0$  means that  $\alpha \in \Omega^1(M)$  is a Killing 1-tensor in the sense of Definition 6.1 below.

**Proposition 5.17.** *Let  $\nabla$  be a torsion-free connection on  $M$ . The horizontal lift of a vector field  $X \in \mathfrak{X}(M)$  is a Killing vector field for  $g_{\nabla}$  if and only if  $\text{sym } \iota_X R_{\nabla} = 0$  and  $X$  is parallel. When this happens, it is always gradient Killing vector field.*

*Proof.* Similarly as in Proposition 5.15,  $X^h$  is a Killing vector field for  $g_{\nabla}$  for some  $X \in \mathfrak{X}(M)$  if and only if all of the three components of  $L_{X^h}g_{\nabla}$  vanish. Let us start with the horizontal one,

$$(L_{X^h}g_{\nabla})(Y^h, Z^h) = X^h g_{\nabla}(Y^h, Z^h) - g_{\nabla}([X^h, Y^h], Z^h) - g_{\nabla}(Y^h, [X^h, Z^h]).$$

As the horizontal subbundle is isotropic and  $[X^h, Y^h] = [X, Y]^h + R_{\nabla}(X, Y)^v$ ,

$$(L_{X^h}g_{\nabla})(Y^h, Z^h) = -(R_{\nabla}(X, Y)Z + R_{\nabla}(X, Z)Y)^v = -((\text{sym } \iota_X R_{\nabla})(Y, Z))^v,$$

which, by the continuity, vanishes if and only if  $\text{sym } \iota_X R_{\nabla} = 0$ . For the mixed component, we have

$$\begin{aligned} (L_{X^h}g_{\nabla})(Y^h, \alpha^v) &= X^h g_{\nabla}(Y^h, \alpha^v) - g_{\nabla}([X^h, Y^h], \alpha^v) - g_{\nabla}(Y^h, [X^h, \alpha^v]) \\ &= X^h \text{pr}^* \alpha(Y) - \text{pr}^* \alpha([X, Y]) - \text{pr}^*(\nabla_X \alpha)(Y) \\ &= \text{pr}^* \alpha(\nabla_Y X). \end{aligned}$$

Therefore, the mixed component vanishes if and only if  $\nabla X = 0$ , that is,  $X$  is parallel. Finally, the vertical component vanishes because the vertical subbundle is isotropic and its space of sections is an ideal for the Lie bracket of vector fields. By Lemma B.7, we find that

$$g_{\nabla}(\text{grad}_{\nabla} X^v, Y^h) = Y^h X^v = (\nabla_Y X)^v, \quad g_{\nabla}(\text{grad}_{\nabla} X^v, \alpha^v) = \alpha^v X^v = \text{pr}^* \alpha(X),$$

that is,  $\text{grad}_{\nabla} X^v = X^h + (\nabla X)^v$  and the result follows.  $\square$

**Lemma 5.18.** *Let  $\nabla$  be a torsion-free connection on  $M$ . The complete lift of a vector field  $X \in \mathfrak{X}(M)$  is a Killing vector field for  $g_\nabla$  if and only if  $X$  is an affine vector field. It is a gradient Killing vector field if and only if  $\text{sym } \iota_X R_\nabla = 0$  and  $X$  is parallel.*

*Proof.* Given  $X \in \mathfrak{X}(M)$ , by Proposition B.3, we have that  $X^c \in \mathfrak{X}(T^*M)$  is a Killing vector field for  $g_\nabla$  if and only if  $(L_{X^c} g_\nabla)(Y^c, Z^c) = 0$  for all  $Y, Z \in \mathfrak{X}(M)$ . Using (30) and the properties of the complete lift, see Appendix B, we find

$$\begin{aligned} (L_{X^c} g_\nabla)(Y^c, Z^c) &= X^c g_\nabla(Y^c, Z^c) - g_\nabla([X^c, Y^c], Z^c) - g_\nabla(Y^c, [X^c, Z^c]) \\ &= -X^c \langle Y : Z \rangle_s^v - g_\nabla([X, Y]^c, Z^c) - g_\nabla(Y^c, [X, Z]^c) \\ &= (-[X, \langle Y : Z \rangle_s] + \langle [X, Y] : Z \rangle_s + \langle Y : [X, Z] \rangle_s)^v, \end{aligned}$$

which, by the continuity, vanishes if and only if

$$[X, \langle Y : Z \rangle_s] = \langle [X, Y] : Z \rangle_s + \langle Y : [X, Z] \rangle_s.$$

This is, thanks to Proposition 4.15, equivalent to  $X$  being an affine vector field. Using the fact that  $X^c = X^h - (\nabla X)^v$  (see Lemma B.6) yields that  $X^c$  is a gradient if and only if there is a function  $\xi \in C^\infty(T^*M)$  such that

$$g_\nabla(X^h) = d\xi|_{\Gamma(\mathcal{V})}, \quad g_\nabla(-(\nabla X)^v) = d\xi|_{\Gamma(\mathcal{H}_\nabla)}.$$

Equivalently, by Lemmas 5.4 and B.7, for any  $\alpha \in \Omega^1(M)$  and  $Y \in \mathfrak{X}(M)$ ,

$$\alpha^v(X^v - \xi) = 0, \quad Y^h(X^v + \xi) = 0.$$

It follows from the first equation that  $\xi = X^v + \text{pr}^* f$  for some  $f \in C^\infty(M)$ . Employing this to the second equation, we obtain

$$2(\nabla_Y X)^v + \text{pr}^*(Yf) = 0$$

As the first term depends on the fibre coordinates and the second one does not, necessarily both of them vanish, that is,  $X$  is parallel and  $f$  is constant. Since  $X$  being parallel implies  $R_\nabla^s X = 0$ , the result follows.  $\square$

We can now prove the final result of this section.

**Theorem 5.19.** *Let  $\nabla$  be a torsion-free connection on  $M$  and  $X, Y \in \mathfrak{X}(M)$ . All of the six relations*

$$\begin{aligned} [\iota_X, \iota_Y] &= 0, & [\nabla^s, \nabla^s] &= 0, & [\iota_X, \nabla^s] &= L_X^s, \\ [L_X^s, \iota_Y] &= \iota_{\langle X:Y \rangle_s}, & [L_X^s, L_Y^s] &= 0, & [\nabla^s, L_X^s] &= 0 \end{aligned}$$

*are satisfied if and only if  $X^c$  and  $Y^c$  are gradient Killing vector fields for  $g_\nabla$ .*

*Proof.* Note that the four commutators

$$[\iota_X, \iota_Y] = 0, \quad [\nabla^s, \nabla^s] = 0, \quad [\iota_X, \nabla^s] = L_X^s, \quad [L_X^s, \iota_Y] = \iota_{\langle X:Y \rangle_s}.$$

are true for any vector fields. From Proposition 3.7, we know that, in general,

$$[\nabla^s, L_X^s] = 2\nabla_{\nabla X}^s + \iota_{2\text{sym } \iota_X R_\nabla + R_\nabla^s X}.$$



We clearly have that  $[\nabla^s, L_X^s]|_{\mathcal{C}^\infty(M)} = 0$  if and only if  $(\nabla_{\nabla X}^s)|_{\mathcal{C}^\infty(M)} = 0$ , which is equivalent to  $X$  being parallel. For a parallel vector field  $X$  (hence, also  $R_{\nabla}^s X = 0$ ), we have that  $[\nabla^s, L_X^s]|_{\Upsilon^1(M)} = 0$  if and only if  $\text{sym } \iota_X R_{\nabla} = 0$ . Altogether, by Lemma 5.18, gives that

$$[\nabla^s, L_X^s] = 0$$

if and only if  $X^c$  is a gradient Killing vector field for  $g_{\nabla}$ .

We know from Proposition 3.2 that, in general,

$$[L_X^s, L_Y^s] = L_{[X, Y]}^s + \iota_{2(\nabla_{\nabla X} Y - \nabla_{\nabla Y} X - R_{\nabla}(X, Y))}.$$

If  $X$  and  $Y$  are parallel, the identity

$$[L_X^s, L_Y^s] = L_{[X, Y]}^s$$

is satisfied if and only if  $R_{\nabla}(X, Y) = 0$ . Using the algebraic Bianchi identity, this is equivalent to  $R_{\nabla}(Y, \cdot)X + R_{\nabla}(\cdot, X)Y = 0$ , but this is clearly satisfied as  $X$  and  $Y$  are parallel.  $\square$

*Remark 5.20.* Note that if  $X$  and  $Y$  are parallel, then,  $\langle X : Y \rangle_s = 0$ . Moreover, by the torsion-freeness of  $\nabla$ , also  $[X, Y] = 0$ . Hence,  $0 = [L_X^s, L_Y^s] = L_{[X, Y]}^s = L_{\langle X : Y \rangle_s}^s$ .

The commutation relations of symmetric Cartan calculus in Theorem 5.19 are completely analogous to those of classical Cartan calculus, but they require an extra condition on the vector fields  $X, Y$ . At first glance, this may seem as an asymmetry between the two theories. However, having in mind the analogy between  $g_{\nabla}$  and  $\omega_{\text{can}}$ , the condition of the ‘complete lift being gradient Killing’ translates into the ‘complete lift being Hamiltonian’, which, by (47) in Appendix B, is trivially satisfied.

*Remark 5.21.* Note that the analogy between gradient Killing and Hamiltonian is complete. In the same way that, for a (pseudo-)Riemannian metric  $g$ , any gradient Killing vector field is automatically Killing ( $L_X g = 0$ ), for a symplectic form  $\omega$ , any Hamiltonian vector field is symplectic ( $L_X \omega = 0$ ).

## 6. SYMMETRIC COHOMOLOGY

Just as the exterior derivative gives rise to de Rham cohomology, we introduce here the analogue for the symmetric derivative  $\nabla^s$ .

**6.1. Killing tensors.** Symmetric tensors that are  $\nabla^s$ -closed have been considered before under the name of Killing tensors. We give some examples and recall their geometric significance.

**Definition 6.1.** Given a connection  $\nabla$  on  $M$ , a **Killing tensor** for  $\nabla$  is a symmetric form  $K \in \Upsilon^\bullet(M)$  such that

$$\nabla^s K = 0.$$

We denote the Killing tensors of homogeneous degree  $r$  by  $\text{Kill}_\nabla^r(M)$ . They are a vector subspace, but not a  $\mathcal{C}^\infty(M)$ -submodule, of the space of symmetric forms  $\Upsilon^r(M)$ . Together with the symmetric product, they form a graded subalgebra

$$\text{Kill}_\nabla^\bullet(M) := \bigoplus_{r \in \mathbb{N}} \text{Kill}_\nabla^r(M) \subseteq \Upsilon^\bullet(M),$$

which we call the **Killing tensor algebra** of  $\nabla$ .

**Example 6.2** (Killing 0-tensors). *As the symmetric derivative is geometric (Definition 2.4), the space of Killing 0-tensors for any connection  $\nabla$  is the space of locally constant functions.*

**Example 6.3.** *For the Levi-Civita connection of a (pseudo-)Riemannian metric  $g$ , by Proposition 2.18,  $g^{-1}$  identifies  $\text{Kill}_g^1(M)$  with the Killing vector fields for  $g$  (see Example 4.14). Moreover,  $g \in \text{Kill}_g^2(M)$ .*

Let us recall that the geometric significance of Killing tensors is that they induce conserved quantities along geodesics. For this, we use an equivalent definition of the geodesic spray  $X_\nabla \in \mathfrak{X}(TM)$ , which was introduced in Section 2.3.

**Lemma 6.4.** *Let  $\nabla$  be a connection on  $M$ . The geodesic spray is the unique vector field whose integral curve starting at  $u \in TM$  is a curve  $\dot{\gamma} : I \rightarrow TM$ , where  $\gamma : I \rightarrow M$  is a  $\nabla$ -geodesic satisfying  $\dot{\gamma}(0) = u$ . In particular, for any  $\nabla$ -geodesic  $\gamma$  on  $M$  and  $\xi \in \mathcal{C}^\infty(TM)$  there holds*

$$(X_\nabla \xi) \circ \dot{\gamma} = \frac{d}{dt}(\xi \circ \dot{\gamma}).$$

*Proof.* Consider a curve  $V : I \rightarrow TM$ . In natural coordinates  $(TU, \{x^i\} \cup \{v^j\})$ ,

$$(X_\nabla)_{V(t)} = V^i(t) \partial_{x^i}|_{V(t)} - V^i(t)V^j(t) \Gamma_{ij}^k(\gamma(t)) \partial_{v^k}|_{V(t)},$$

where  $V^i := v^i \circ V$  and  $\gamma := \text{pr} \circ V$ . Using the notation  $\gamma^i := x^i \circ \gamma$ , the equation for the integral curve of  $X_\nabla$  starting at  $u \in TM$  is locally given by

$$\dot{\gamma}^i(t) = V^i(t), \quad \dot{V}^i(t) = -V^i(t)V^j(t) \Gamma_{ij}^k(\gamma(t)), \quad V(0) = u.$$

Using the first equation on the second one, we recover the geodesic equation

$$0 = \nabla_{\dot{\gamma}} \dot{\gamma}|_U = \ddot{\gamma}^k(t) + \dot{\gamma}^i(t) \dot{\gamma}^j(t) \Gamma_{ij}^k(\gamma(t)), \quad \dot{\gamma}(0) = u.$$

The converse follows simply from the fact that every vector field is uniquely characterized by its integral curves.  $\square$

**Proposition 6.5.** *Let  $\nabla$  be a connection on  $M$ . The assignment  $K \mapsto \tilde{K}$  gives an isomorphism of unital graded algebras*

$$\text{Kill}_\nabla^\bullet(M) \xrightarrow{\sim} \left\{ \begin{array}{l} \text{polynomials in velocities on } M \\ \text{constant along } \nabla\text{-geodesics} \end{array} \right\}.$$

*Proof.* By the correspondence (8) and Proposition 2.14, we have  $K \in \text{Kill}_\nabla^\bullet(M)$  if and only if  $X_\nabla \tilde{K} = 0$ . In particular, we have  $(X_\nabla \tilde{K}) \circ \dot{\gamma} = 0$  for every  $\nabla$ -geodesic  $\gamma$  if  $K \in \text{Kill}_\nabla^\bullet(M)$ , which, by Lemma 6.4, means that  $\tilde{K} \circ \dot{\gamma}$  is constant. Conversely, assume that  $\tilde{K}$  is constant along  $\nabla$ -geodesics, hence  $(X_\nabla \tilde{K}) \circ \dot{\gamma} = 0$ . Since there is a  $\nabla$ -geodesic  $\gamma$  such that  $\dot{\gamma}(0) = u$  for every  $u \in TM$ , we get  $X_\nabla \tilde{K} = 0$ .  $\square$

We can also describe Killing tensors that are in the image of  $\nabla^s$ , which we will call  $\nabla^s$ -**exact** tensors.

**Corollary 6.6.** *Given  $K \in \text{Kill}_\nabla^r(M) \cap \text{Im } \nabla^s$  for  $r \neq 0$ , there exists  $\varphi \in \Upsilon^{r-1}(M)$  such that for every  $\nabla$ -geodesic  $\gamma$ , we have, for some constant  $\lambda \in \mathbb{R}$ ,*

$$\tilde{\varphi}(\dot{\gamma}(t)) = \tilde{K}(\dot{\gamma}(t))t + \lambda.$$

**6.2. Killing 1-tensors.** Recall that the maximum number of linearly independent Killing vector fields on a connected (pseudo-)Riemannian manifold of dimension  $n$  is  $\frac{1}{2}n(n+1)$ . In this section, we prove an analogous result concerning Killing 1-tensors.

**Lemma 6.7.** *Let  $\nabla$  be a torsion-free connection on  $M$ . For  $\alpha \in \text{Kill}_\nabla^1(M)$ ,*

$$\nabla\alpha \in \Omega^2(M), \quad \text{and} \quad \nabla\nabla\alpha = \iota_\alpha R_\nabla.$$

*Proof.* The first part is easy to see as being a Killing 1-tensor literally means that the symmetric part of  $\nabla\alpha$  vanishes. By a straightforward calculation we find, for all  $X, Y, Z \in \mathfrak{X}(M)$ , that

$$(\nabla\nabla\alpha)(X, Y, Z) - (\nabla\nabla\alpha)(Y, X, Z) = -\alpha(R_\nabla(X, Y)Z).$$

Taking the skew-symmetric projection of this relation and using the algebraic Bianchi identity together with that  $\nabla\alpha \in \Omega^2(M)$  yields

$$(\nabla\nabla\alpha)(X, Y, Z) = \alpha(R_\nabla(Y, Z)X) = R_\nabla(\alpha, X, Y, Z) = (\iota_\alpha R_\nabla)(X, Y, Z).$$

$\square$

Using Lemma 6.7, we find that every Killing 1-tensor is fully determined by linear algebraic data at a point.

**Lemma 6.8.** *Let  $M$  be connected and  $m \in M$  be an arbitrary point. Every  $\alpha \in \text{Kill}_\nabla^1(M)$  is fully determined by the following data*

$$\alpha_m \in T_m^*M, \quad (\nabla\alpha)_m \in \wedge^2 T_m^*M.$$

*Proof.* We show that having  $\alpha \in \text{Kill}_\nabla^1(M)$  such that  $\alpha_m$  and  $(\nabla\alpha)_m$  vanish at some point  $m \in M$  already implies that  $\alpha$  vanishes as a 1-form. The result then follows from the linearity. By hypothesis, the set

$$N(\alpha) := \{m \in M \mid \alpha_m = 0 \text{ and } (\nabla\alpha)_m = 0\}$$

is non-empty. It is also closed, since it can be seen as  $\alpha^{-1}(0) \cap (\nabla\alpha)^{-1}(0)$ , where 0 denotes the image of the zero section. We will prove that it is also open. Consider the vector bundle connection  $\nabla^R$  on  $TM \oplus \wedge^2 T^*M$  given by

$$\nabla_X^R(\alpha + B) := (\nabla_X\alpha - \iota_X B) + (\nabla_X B - \iota_X \iota_\alpha R_\nabla).$$

It follows from Lemma 6.7 that the section  $\alpha + \nabla\alpha \in \Gamma(T^*M \oplus \wedge^2 T^*M)$  is parallel with respect to  $\nabla^R$  for every  $\alpha \in \text{Kill}_\nabla^1(M)$ . Take an arbitrary point  $m' \in N(\alpha)$  and an open connected neighbourhood  $U \subseteq M$ . Any  $m'' \in U$  is connected to  $m'$  by a smooth curve. Since  $\alpha + \nabla\alpha$  is  $\nabla^R$ -parallel along this curve, it is necessary that  $\alpha_{m''} = 0$  and  $(\nabla\alpha)_{m''} = 0$ , that is,  $m'' \in N(\alpha)$ , which proves that  $N(\alpha)$  is open, hence  $N(\alpha) = M$ , and finally,  $\alpha = 0$ .  $\square$

**Proposition 6.9.** *Let  $M$  be connected. The maximum number of linearly independent Killing 1-tensors on  $(M, \nabla)$  is equal to  $\frac{1}{2}n(n+1)$ , where  $n := \dim M$ .*

*Proof.* By Lemma 6.8, there is an injective linear morphism from  $\text{Kill}_\nabla^1(M)$  to  $T_m^*M \oplus \wedge^2 T_m^*M$  for any  $m \in M$ . Since  $\dim(T_m^*M \oplus \wedge^2 T_m^*M) = \frac{1}{2}n(n+1)$ , the result follows.  $\square$

**6.3. Definition of symmetric cohomology.** We have looked at Killing tensors as analogues of closed forms. Since the symmetric derivative does not square to zero, we need to adjust the definition of cohomology groups: we take as  $\nabla^s$ -exact tensors the Killing tensors in the image of  $\nabla^s$ . We can then consider the quotient.

**Definition 6.10.** The  $r$ th-symmetric cohomology group of  $(M, \nabla)$  is given by

$$(33) \quad H_\nabla^r(M) := \frac{\text{Kill}_\nabla^r(M)}{\text{Kill}_\nabla^r(M) \cap \text{Im } \nabla^s|_{\mathcal{K}^{r-1}(M)}}.$$

As the symmetric derivative is geometric, that is  $\nabla^s|_{\mathcal{K}^0(M)} = d|_{C^\infty(M)}$ , we can be more specific for  $r = 0$  and  $r = 1$ .

**Proposition 6.11.** *For  $r = 0$ , the symmetric cohomology group  $H_\nabla^0(M)$  coincides with the de Rham cohomology group  $H_{dR}^0(M)$ , that is, locally constant functions.*

**Proposition 6.12.** *For  $r = 1$ , we have*

$$H_\nabla^1(M) = \frac{\text{Kill}_\nabla^1(M)}{\text{Kill}_\nabla^1(M) \cap \Omega_{\text{exact}}^1(M)},$$

that is,  $H_\nabla^1$  describes Killing 1-tensors modulo exact 1-forms. Moreover, for  $M$  connected with  $n := \dim M$ , we have

$$(34) \quad \dim H_\nabla^1(M) \leq \frac{1}{2}n(n+1).$$

*Proof.* The first claim is direct, the second follows from the bound on the dimension of the space of Killing 1-tensors given by Proposition 6.9.  $\square$

If  $\nabla$  is the Levi-Civita connection for some (pseudo-)Riemannian metric, we can rephrase  $H_{\nabla}^1(M)$  in terms of vector fields.

**Proposition 6.13.** *Let  $g$  be a (pseudo-)Riemannian metric on  $M$ .*

$$H_{g\nabla}^1(M) \cong \frac{\{\text{Killing vector fields for } g\}}{\{\text{gradient Killing vector fields for } g\}}.$$

*Proof.* The space  $\text{Kill}_{g\nabla}^1(M)$  is identified by  $g^{-1}$  with the space of Killing vector fields for  $g$ , see Example 6.3. On the other hand, since by definition  $\text{grad}_g := g^{-1} \circ \text{d}$ , exact 1-forms on  $M$  are identified by  $g^{-1}$  with gradient Killing vector fields.  $\square$

Rephrasing the symmetric cohomology in terms of polynomials in velocities on  $M$  and the geodesic spray gives a geometric interpretation for arbitrary  $r$ .

**Corollary 6.14.** *The assignment  $[K] \mapsto [\tilde{K}]$  gives an isomorphism of vector spaces*

$$H_{\nabla}^r(M) \xrightarrow{\sim} \left\{ \begin{array}{l} \text{degree-}r \text{ polynomials in velocities} \\ \text{on } M \text{ constant along } \nabla\text{-geodesics} \\ \text{modulo } \text{Im } X_{\nabla} \end{array} \right\}.$$

*Proof.* It follows directly from Proposition 2.14 and Proposition 6.5.  $\square$

Given two manifolds with connections  $(M, \nabla)$ ,  $(M', \nabla')$ , and  $\phi \in \text{Aff}(\nabla, \nabla')$ . It follows easily from Remark 4.10 that

$$K \in \text{Kill}_{\nabla'}^{\bullet}(M') \Leftrightarrow \phi^* K \in \text{Kill}_{\nabla}^{\bullet}(M), \quad \text{and} \quad K \in \text{Im } \nabla'^s \Leftrightarrow \phi^* K \in \text{Im } \nabla^s.$$

As an immediate consequence we get the next proposition.

**Proposition 6.15.** *If  $(M, \nabla)$  and  $(M', \nabla')$  are affine diffeomorphic, then*

$$\begin{aligned} \phi^* : H_{\nabla'}^r(M') &\rightarrow H_{\nabla}^r(M) \\ [K] &\mapsto [\phi^* K]. \end{aligned}$$

*is a vector space isomorphism. Consequently,  $\dim H_{\nabla}^r(M) = \dim H_{\nabla'}^r(M')$ .*

The space of connections on a manifold  $M$ , which we denote by  $\mathcal{C}(M)$ , is endowed with the natural equivalence relation defined by

$$\nabla \sim \nabla' \quad \Leftrightarrow \quad \text{Aff}(\nabla, \nabla') \neq \emptyset.$$

Equivalently,  $\nabla \sim \nabla'$  if there exists  $\phi \in \text{Diff}(M)$  such that  $\nabla = \phi^* \nabla'$ . We can thus talk about the space of connections modulo diffeomorphisms. Proposition 6.15 states that the symmetric cohomology groups are invariants in the quotient (or moduli) space.

*Remark 6.16.* Note that, as invariants, they are actually more suitable for torsion-free connections, since they cannot see the difference between a connection with torsion  $\nabla$  and its associated torsion-free connection  $\nabla^0$ , whereas it follows easily from Remark 4.8 that  $[[\nabla]] \neq [[\nabla^0]]$  in  $\mathcal{C}(M)/\text{Diff}(M)$ .

As the space of Killing tensors forms an algebra with the symmetric product, we get a *symmetric cup product* of cohomology classes, which endows the direct sum of the cohomology groups with a commutative algebra structure.

**6.4. The cartesian space  $\mathbb{R}^n$ .** Consider  $\mathbb{R}^n$  with the Euclidean connection  $\nabla^{\text{Euc}}$ . The condition for  $\alpha := \alpha_i dx^i \in \Upsilon^1(\mathbb{R}^n)$  to be a Killing tensor reads as

$$(35) \quad \frac{\partial \alpha_j}{\partial x^i} + \frac{\partial \alpha_i}{\partial x^j} = 0.$$

It is easy to see that all coordinate 1-forms  $dx^i$  are solutions, and moreover, all 1-forms of the form  $x^i dx^j - x^j dx^i$  solves the equation (35). Since all of these solutions are linearly independent and there are  $\frac{1}{2}n(n+1)$  of them, Lemma 6.8 implies that

$$\text{Kill}_{\nabla^{\text{Euc}}}^1(\mathbb{R}^n) = \text{span}(\{dx^i\}_{i=1}^n \cup \{x^i dx^j - x^j dx^i\}_{i < j}).$$

As  $dx^i$  are clearly exact and  $x^i dx^j - x^j dx^i$  are not, we conclude

$$H_{\nabla^{\text{Euc}}}^1(\mathbb{R}^n) \cong \text{span}\{x^i dx^j - x^j dx^i\}_{i < j} \cong \mathbb{R}^{\frac{n(n-1)}{2}}.$$

*Remark 6.17.* The identification of Proposition 6.13 can be understood as  $H_{\nabla^{\text{Euc}}}^1(\mathbb{R}^n)$  being, infinitesimally, symmetries modulo translational symmetries (spanned by  $\{dx^i\}_{i=1}^n$ ), which results in *rotational symmetries*, spanned by  $\{x^i dx^j - x^j dx^i\}_{i < j}$ . Thus, symmetric cohomology can be understood as a generalization of this picture.

For  $n = 1$ , we can easily calculate the symmetric cohomology for a generic connection  $\nabla$  on  $\mathbb{R}$ . Because of the dimension of the manifold we have  $\Upsilon^1(\mathbb{R}) = \Omega_{\text{cl}}^1(\mathbb{R})$  (closed 1-forms). It follows from *Poincaré lemma* that every 1-form on  $\mathbb{R}$  is exact. In conclusion,

$$H_{\nabla}^1(\mathbb{R}) = \frac{\text{Kill}_{\nabla}^1(\mathbb{R})}{\text{Kill}_{\nabla}^1(\mathbb{R}) \cap \Upsilon^1(\mathbb{R})} = \frac{\text{Kill}_{\nabla}^1(\mathbb{R})}{\text{Kill}_{\nabla}^1(\mathbb{R})} = \{0\}.$$

For  $n = 2$ , the situation is much more complicated. A generic torsion-free connection  $\nabla$  is determined by 6 smooth functions  $f_j \in \mathcal{C}^\infty(\mathbb{R}^2)$  as follows:

$$\nabla_{\partial_x} \partial_x = f_1 \partial_x + f_2 \partial_y, \quad \nabla_{\partial_x} \partial_y = \frac{1}{2}(f_3 \partial_x + f_4 \partial_y), \quad \nabla_{\partial_y} \partial_y = f_5 \partial_x + f_6 \partial_y.$$

A 1-form  $\alpha = \alpha_1 dx + \alpha_2 dy$  is Killing if and only if

$$(36) \quad \begin{aligned} 0 &= \frac{\partial \alpha_1}{\partial x} - \alpha_1 f_1 - \alpha_2 f_2, \\ 0 &= \frac{\partial \alpha_1}{\partial y} + \frac{\partial \alpha_2}{\partial x} - \alpha_1 f_3 - \alpha_2 f_4, \\ 0 &= \frac{\partial \alpha_2}{\partial y} - \alpha_1 f_5 - \alpha_2 f_6. \end{aligned}$$

Note that  $\nabla^{\text{Euc}}$  corresponds to the distinguished choice  $f_j = 0$ . Unlike the case  $n = 1$ , the symmetric cohomology of a generic connection is pretty wild. In particular, the first symmetric cohomology group can be of any dimension up to the upper bound (34). The computations supporting Examples 6.18, 6.20 and 6.21 can be found in Appendix C.

**Example 6.18** (Trivial cohomology group). *Consider a connection  $\nabla$  on  $\mathbb{R}^2$  determined by the choice*

$$f_3(x, y) := xy, \quad f_4(x, y) := y$$

and  $f_j = 0$  otherwise. One finds that the system of PDE's (36) has only the trivial solutions  $\alpha_1 = 0 = \alpha_2$ . Therefore,

$$\text{Kill}_{\nabla}^1(\mathbb{R}^2) = \{0\}, \quad H_{\nabla}^1(\mathbb{R}^2) = \{0\}.$$

**Example 6.19** (1-dimensional cohomology group). *As we have shown above, for the Euclidean connection, we have  $H_{\nabla^{\text{Euc}}}^1(\mathbb{R}^2) \cong \mathbb{R}$ .*

**Example 6.20** (2-dimensional cohomology group). *Given an arbitrary nonzero constants  $\lambda_1, \lambda_2 \in \mathbb{R}$ . We consider a connection  $\nabla$  on  $\mathbb{R}^2$  given by*

$$f_3(x, y) := \lambda_1, \quad f_4(x, y) := \lambda_2$$

and  $f_j = 0$  otherwise. One finds that

$$\text{Kill}_{\nabla}^1(\mathbb{R}^2) = \text{span} \left\{ e^{\lambda_1 y} dx, e^{\lambda_2 x} dy, \frac{1}{\lambda_1} dx - \frac{1}{\lambda_2} dy \right\}, \quad H_{\nabla}^1(\mathbb{R}^2) \cong \mathbb{R}^2.$$

**Example 6.21** (3-dimensional cohomology group). *Finally, let us have a connection  $\nabla$  on  $\mathbb{R}^2$  defined by*

$$f_3(x, y) := -2y \left( 1 + \frac{2}{1 + 2y^2} \right), \quad f_4(x, y) := -2x \left( 1 + \frac{2}{1 + 2x^2} \right)$$

and  $f_j = 0$  otherwise. Solving (36) yields

$$\text{Kill}_{\nabla}^1(\mathbb{R}^2) = \text{span} \left\{ \frac{e^{-y^2}}{1 + 2y^2} dx, \frac{e^{-x^2}}{1 + 2x^2} dy, \frac{y}{1 + 2y^2} dx - \frac{x}{1 + 2x^2} dy \right\},$$

hence  $H_{\nabla}^1(\mathbb{R}^2) \cong \mathbb{R}^3$ .

Following the discussion at the end of Section 6.3, we immediately see that the connections from Examples 6.18 to 6.21 define four different classes in the moduli space  $\mathcal{C}(\mathbb{R}^2)/\text{Diff}(\mathbb{R}^2)$ .

6.5. **The  $n$ -sphere  $S^n$ .** Consider the  $n$ -sphere  $S^n$  equipped with the round metric  $g^\circ$ . We compute its first symmetric cohomology group using Proposition 6.13.

It is a well-known fact (see, for instance [23]) that there are  $\frac{1}{2}n(n+1)$  linearly independent Killing vector fields for  $g^\circ$ . We review an important fact concerning gradient Killing vector fields on a compact oriented Riemannian manifold (see, e.g., [25]).

**Proposition 6.22.** *The only gradient Killing vector field on a compact orientable Riemannian manifold is the zero vector field.*

*Proof.* Let  $f$  be a smooth function on a compact oriented Riemannian manifold  $(M, g)$  such that  $X := \text{grad}_g f$  is a Killing vector field for  $g$ . It means that

$$0 = {}^g\nabla^s df = 2{}^g\nabla df \quad \Leftrightarrow \quad 0 = {}^g\nabla X \quad \Rightarrow \quad 0 = \text{div}_g X,$$

where  $\text{div}_g := \text{tr} \circ {}^g\nabla : \mathfrak{X}(M) \rightarrow \mathcal{C}^\infty(M)$  is the divergence operator of  $g$  and  $\text{tr} : \Gamma(\text{End } TM) \rightarrow \mathcal{C}^\infty(M)$  is the trace operator. Therefore,

$$\text{div}_g(\text{grad}_g f^2) = \text{div}_g(2fX) = 2Xf + 2f \text{div}_g X = 2\|X\|_g^2.$$

For  $Y \in \mathfrak{X}(M)$ , we have  $(\text{div}_g Y) \text{vol}_g = L_Y \text{vol}_g = d\iota_Y \text{vol}_g$ . By Stokes' theorem,

$$0 = \int_M d\iota_{\text{grad}_g f^2} \text{vol}_g = \int_M (\text{div}_g \text{grad}_g f^2) \text{vol}_g = 2 \int_M \|X\|_g^2 \text{vol}_g.$$

As  $M$  is compact, we have  $\|X\|_g^2 = 0$ , hence  $X = 0$  because  $g$  is positive definite.  $\square$

As  $(S^n, g^\circ)$  is a compact orientable Riemannian manifold, we find

$$H_{\nabla^\circ}^1(S^n) \cong \frac{\{\text{Killing vector fields for the round metric}\}}{\{0\}} \cong \mathbb{R}^{\frac{n(n+1)}{2}}.$$

*Remark 6.23.* The interpretation of the symmetric cohomology from Remark 6.17 fits nicely with the case of the round sphere  $(S^n, \nabla^\circ)$ , where we have  $\frac{1}{2}n(n+1)$  rotational symmetries but no translational ones.

For  $n = 1$ , every connection is fully determined by a function  $f \in \mathcal{C}^\infty(S^1)$ ,

$$\nabla_{\partial_\theta}^f \partial_\theta := f \partial_\theta,$$

where  $\partial_\theta$  is the standard unit (with respect to the round metric) vector field on  $S^1$ . The round connection corresponds to the distinguished choice  $f = 0$ . A 1-form  $\alpha = \alpha_1 d\theta$  (with  $d\theta$  the dual 1-form to  $\partial_\theta$ ) is a Killing 1-tensor if and only if

$$(37) \quad \partial_\theta \alpha_1 = f \alpha_1.$$

This is only possible when  $\int_{S^1} f d\theta = 0$ , in which case  $\alpha_1 = \lambda e^F$  for  $\lambda \in \mathbb{R}$  an arbitrary constant and  $F(\theta) := \int_0^\theta f d\theta$ . Therefore,

$$(38) \quad \text{Kill}_{\nabla^f}^1(S^1) \cong \begin{cases} \mathbb{R} & \text{when } \int_{S^1} f d\theta = 0, \\ \{0\} & \text{when } \int_{S^1} f d\theta \neq 0. \end{cases}$$



The 1-form  $\lambda e^F d\theta$  is never exact since  $\int_{S^1} e^F d\theta \neq 0$ . In conclusion, there are no exact Killing 1-tensors for any connection on  $S^1$ , and hence

$$(39) \quad H_{\nabla^f}^1(S^1) \cong \text{Kill}_{\nabla^f}^1(S^1).$$

Let us now show that the condition  $\int_{S^1} f d\theta = 0$  for the connection  $\nabla^f$  has a very clear geometric meaning.

**Lemma 6.24.** *Let  $f \in C^\infty(S^1)$ , the connection  $\nabla^f$  is the Levi-Civita connection for some metric on  $S^1$  if and only if  $\int_{S^1} f d\theta = 0$ .*

*Proof.* Since all connections on a 1-dimensional manifold are torsion-free, the connection  $\nabla^f$  is Levi-Civita for some metric  $g$  if and only if  $\nabla^f g = 0$ . This is equivalent, by writing  $g = g_1 d\theta \odot d\theta$  for  $g_1 \in C^\infty(S^1)$ , to the equation  $\partial_\theta g_1 = 2fg_1$ . As we saw above, this admits a nowhere vanishing solution  $g_1$  if and only if  $\int_{S^1} f d\theta = 0$ .  $\square$

*Remark 6.25.* Note that an analogous proof to that of Lemma 6.24 gives that all connections on  $\mathbb{R}^1$  are Levi-Civita for some metric.

By (38), (39) and Lemma 6.24, we can distinguish Levi-Civita connections on  $S^1$  from all other connections by using Killing tensors or just symmetric cohomology.

**Proposition 6.26.** *The only connections  $\nabla$  on  $S^1$  such that  $H_{\nabla}^1(S^1) \neq \{0\}$  (and hence  $\mathbb{R}$ ) are the Levi-Civita connections for any choice of metric on  $S^1$ .*

Since we have seen in Section 6.4 that  $H_{\nabla}^1(\mathbb{R}^1) = \{0\}$  for any connection  $\nabla$ , we have that symmetric cohomology distinguishes between  $\mathbb{R}^1$  and  $S^1$ , just as de Rham cohomology also does.

We would like to find more powerful applications of symmetric cohomology in the future. On the one hand, we can study the 'symmetric cohomology spectrum' (symmetric cohomology groups of all possible connections) to distinguish between two smooth manifolds, as we did in the case of  $\mathbb{R}^1$  and  $S^1$ . Note that even partial information about the spectrum may serve this purpose. On the other hand, the dimension of a symmetric cohomology group is invariant under the action of diffeomorphisms on the space of connections on a manifold. Therefore, it may be used to show that two connections represent different classes in the moduli space.

## 7. FURTHER TECHNIQUES FOR SYMMETRIC COHOMOLOGY

We explore now products, on which we define the Künneth subalgebra, the relation with the Patterson-Walker, which allows us to prove the Patterson-Walker bound, and left-invariant connections, which reveal the relation to Lie-admissible algebras.

**7.1. The Künneth subalgebra.** Given two manifolds with connection  $(M_1, \nabla^1)$  and  $(M_2, \nabla^2)$ , there is a natural notion of the **product connection**  $\nabla$  on the manifold  $M_1 \times M_2$ :

$$(40) \quad \nabla_{\text{pr}_1^*X_1 + \text{pr}_2^*X_2}(\text{pr}_1^*Y_1 + \text{pr}_2^*Y_2) = \text{pr}_1^*\nabla_{X_1}^1 Y_1 + \text{pr}_2^*\nabla_{X_2}^2 Y_2,$$

where  $\text{pr}_j : M_1 \times M_2 \rightarrow M_j$  are the natural projections and  $X_j, Y_j \in \mathfrak{X}(M_j)$ . Since the tangent bundle  $T(M_1 \times M_2)$  is canonically isomorphic to  $\text{pr}_1^*TM_1 \oplus \text{pr}_2^*TM_2$ , the space of vector fields  $\mathfrak{X}(M_1 \times M_2)$  is locally generated by pullbacks of vector fields on  $M_1$  and  $M_2$ , hence formula (40) completely determines a connection.

**Lemma 7.1.** *Let  $M_1$  and  $M_2$  be manifolds equipped with connections  $\nabla^1$  and  $\nabla^2$  respectively, and denote the product connection on  $M_1 \times M_2$  by  $\nabla$ . Then*

$$(41) \quad \nabla^s \circ \text{pr}_j^* = \text{pr}_j^* \circ \nabla^{j^s}.$$

*Proof.* As both sides of relation (41) are geometric derivations of  $\Upsilon^\bullet(M_j)$ , it is enough to check the relation on 1-forms. For  $X_j \in \mathfrak{X}(M_j)$  and  $\alpha \in \Upsilon^1(M_j)$ ,

$$\begin{aligned} & \frac{1}{2}(\nabla^s \text{pr}_j^* \alpha)(\text{pr}_1^*X_1 + \text{pr}_2^*X_2, \text{pr}_1^*X_1 + \text{pr}_2^*X_2) \\ &= (\text{pr}_1^*X_1 + \text{pr}_2^*X_2)((\text{pr}_j^* \alpha)(\text{pr}_1^*X_1 + \text{pr}_2^*X_2)) - (\text{pr}_j^* \alpha)(\text{pr}_1^*\nabla_{X_1}^1 X_1 + \text{pr}_2^*\nabla_{X_2}^2 X_2) \\ &= \text{pr}_j^*(X_j \alpha(X_j)) - \text{pr}_j^*(\alpha(\nabla_{X_j}^j X_j)) = \frac{1}{2} \text{pr}_j^*((\nabla^{j^s} \alpha)(X_j, X_j)) \\ &= \frac{1}{2}(\text{pr}_j^* \nabla^s \alpha)(\text{pr}_1^*X_1 + \text{pr}_2^*X_2, \text{pr}_1^*X_1 + \text{pr}_2^*X_2). \end{aligned}$$

The result follows by polarization.  $\square$

We use Lemma 7.1, and the idea behind the Künneth formula, to show the existence of a subspace in the symmetric cohomology of a product.

**Proposition 7.2.** *Let  $(M_1, \nabla^1)$  and  $(M_2, \nabla^2)$  be two manifolds with connection and denote by  $\nabla$  the product connection. The vector space*

$$(42) \quad \bigoplus_{\substack{i,j=0, \\ i+j=r}}^r H_{\nabla^1}^i(M_1) \otimes H_{\nabla^2}^j(M_2).$$

*is isomorphic to a subspace of  $H_{\nabla}^r(M_1 \times M_2)$ , which we call the Künneth subspace.*

*Proof.* It follows immediately from Lemma 7.1 that

$$\text{pr}_j^*(\text{Kill}_{\nabla^j}^r(M_j)) \leq \text{Kill}_{\nabla}^r(M_1 \times M_2), \quad \text{and} \quad \text{pr}_j^*(\text{Im } \nabla^{j^s}) \leq \text{Im } \nabla^s.$$

Moreover, by Lemma 7.1, for  $\varphi_1 \in \Upsilon^i(M_1)$  and  $\varphi_2 \in \Upsilon^j(M_2)$ , we have

$$\begin{aligned} \nabla^s(\text{pr}_1^*\varphi_1 \odot \text{pr}_2^*\varphi_2) &= \nabla^s \text{pr}_1^*\varphi_1 \odot \text{pr}_2^*\varphi_2 + \text{pr}_1^*\varphi_1 \odot \nabla^s \text{pr}_2^*\varphi_2 \\ &= \text{pr}_1^*\nabla^{1^s} \varphi_1 \odot \text{pr}_2^*\varphi_2 + \text{pr}_1^*\varphi_1 \odot \text{pr}_2^*\nabla^{2^s} \varphi_2. \end{aligned}$$

Therefore, we clearly have  $\text{pr}_1^*\varphi_1 \odot \text{pr}_2^*\varphi_2 \in \text{Kill}_{\nabla}^{i+j}(M_1 \times M_2)$  if  $\varphi_1 \in \text{Kill}_{\nabla_1}^i(M_1)$  and  $\varphi_2 \in \text{Kill}_{\nabla_2}^j(M_2)$ . Assume now that we have  $\varphi_1 \in \text{Kill}_{\nabla_1}^i(M_1) \cap \text{Im } \nabla^{1^s}$  and  $\varphi_2 \in \text{Kill}_{\nabla_2}^j(M_2) \cap \text{Im } \nabla^{2^s}$ . By a straightforward calculation one finds

$$\text{pr}_1^*\varphi_1 \odot \text{pr}_2^*\varphi_2 = \frac{1}{2}\nabla^s(\text{pr}_1^*\psi_1 \odot \text{pr}_2^*\varphi_2 + \text{pr}_1^*\varphi_1 \odot \text{pr}_2^*\psi_2),$$

where  $\psi_1 \in \Upsilon^{i-1}(M_1)$  and  $\psi_2 \in \Upsilon^{j-1}(M_2)$  satisfy  $\varphi_j = \nabla^{j^s}\psi_j$ . The result is an immediate consequence of the surjectivity of  $\text{pr}_j$ .  $\square$

In particular, for connected manifolds  $M_1$  and  $M_2$ , we have that the vector space  $H_{\nabla_1}^1(M_1) \oplus H_{\nabla_2}^1(M_2)$  determines the Künneth subspace of  $H_{\nabla}^1(M_1 \times M_2)$ .

**Example 7.3** (The Euclidean plane). *Consider two Euclidean lines  $(\mathbb{R}^1, \nabla^{\text{Euc}})$ , it is not hard to see that the Euclidean plan  $(\mathbb{R}^2, \nabla^{\text{Euc}})$  can be seen as the product manifold equipped with the product connection. Recall the result of Section 6.4,*

$$H_{\nabla^{\text{Euc}}}^1(\mathbb{R}^1) = \{0\} \quad \text{and} \quad H_{\nabla^{\text{Euc}}}^1(\mathbb{R}^2) \cong \mathbb{R}^1.$$

*So, the Künneth subspace and  $H_{\nabla}^r(M_1 \times M_2)$  are not isomorphic in general.*

**Example 7.4** (The flat cylinder). *The flat cylinder can be seen as the product manifold  $S^1 \times \mathbb{R}^1$  equipped with the product connection  $\nabla^{\text{flat}}$  of the round connection  $\nabla^\circ$  on  $S^1$  and the Euclidean connection  $\nabla^{\text{Euc}}$  on  $\mathbb{R}^1$ . It follows from the results of Sections 6.4, 6.5 and Lemma 7.1 that*

$$\text{span}\{d\theta, dx\} \leq \text{Kill}_{\nabla^{\text{flat}}}^1(S^1 \times \mathbb{R}^1).$$

*We check if there are other Killing 1-tensors. A 1-form  $\alpha \in \Upsilon^1(S^1 \times \mathbb{R}^1)$  is uniquely characterized by smooth functions  $\alpha_\theta, \alpha_x \in \mathcal{C}^\infty(S^1 \times \mathbb{R}^1)$  such that  $\alpha = \alpha_\theta d\theta + \alpha_x dx$ . It is a Killing 1-tensor if and only if*

$$0 = \partial_\theta \alpha_\theta, \quad 0 = \partial_x \alpha_x, \quad 0 = \partial_x \alpha_\theta + \partial_\theta \alpha_x.$$

*The first two equations are equivalent to  $\alpha_\theta = \text{pr}_{\mathbb{R}^1}^* f$  and  $\alpha_x = \text{pr}_{S^1}^* h$  for some  $f \in \mathcal{C}^\infty(\mathbb{R}^1)$  and  $h \in \mathcal{C}^\infty(S^1)$ . Using this in the last equation, we have that  $\partial_\theta h$  is a constant function on  $S^1$ , which is possible if and only if  $\partial_\theta h$  vanishes, hence*

$$\text{Kill}_{\nabla^{\text{flat}}}^1(S^1 \times \mathbb{R}^1) = \text{span}\{d\theta, dx\} \quad \text{and} \quad H_{\nabla^{\text{flat}}}^1(S^1 \times \mathbb{R}^1) \cong \mathbb{R}^1.$$

*Therefore, the Künneth subspace and  $H_{\nabla}^1(M_1 \times M_2)$  are isomorphic in this case.*

Note that the direct sum of Künneth subspaces forms a subalgebra with respect to the symmetric cup product mentioned at the end of Section 6.3. We call this subalgebra the **Künneth subalgebra**.

**7.2. The Patterson-Walker bound.** By Proposition 5.10, every connection  $\nabla$  on  $M$  determines a connection  $\bar{\nabla}$  on  $T^*M$ , which is the Levi-Civita connection of the corresponding Patterson-Walker metric. It is natural to ask whether the symmetric cohomology groups of these connections are related. By Proposition 6.13,

$$H_{\bar{\nabla}}^1(T^*M) \cong \frac{\{\text{Killing vector fields for } g_{\nabla}\}}{\{\text{gradient Killing vector fields for } g_{\nabla}\}}.$$

Using the relation between the vertical lift of Killing 1-tensors on  $M$  and the Killing vector fields for the Patterson-Walker metric (Proposition 5.15) and the analogous result for the complete lift of affine vector fields on  $M$  (Lemma 5.18), we obtain an immediate consequence.

**Proposition 7.5.** *Let  $\nabla$  be a torsion-free connection on  $M$  and  $\bar{\nabla}$  be the Levi-Civita connection of  $g_{\nabla}$ . The vector space*

$$\{\text{affine vector fields on } (M, \nabla)\} \oplus \text{Kill}_{\bar{\nabla}}^1(M)$$

*is isomorphic to a subspace of  $\text{Kill}_{\bar{\nabla}}^1(T^*M)$ , and moreover, the vector space*

$$(43) \quad \frac{\{\text{affine vector fields on } (M, \nabla)\}}{\{X \in \mathfrak{X}(M) \mid X \text{ is parallel and } \text{sym } \iota_X R_{\nabla} = 0\}} \oplus H_{\bar{\nabla}}^1(M)$$

*is isomorphic to a subspace of  $H_{\bar{\nabla}}^1(T^*M)$ .*

This proposition gives a bound on the dimension of  $H_{\bar{\nabla}}^1(T^*M)$ , which we refer to as **the Patterson-Walker bound**.

**Example 7.6** (The Euclidean plane). *Consider the Euclidean plane  $(\mathbb{R}^2, \nabla^{\text{Euc}})$ . By a straightforward calculation, one finds*

$$\begin{aligned} \{\text{affine vector fields on } (\mathbb{R}^2, \nabla^{\text{Euc}})\} &= \text{span}\{\partial_x, \partial_y, x\partial_x, y\partial_x, x\partial_y, y\partial_y\} \cong \mathbb{R}^6, \\ \{\text{parallel vector fields on } (\mathbb{R}^2, \nabla^{\text{Euc}})\} &= \text{span}\{\partial_x, \partial_y\} \cong \mathbb{R}^2, \\ \text{Kill}_{\nabla^{\text{Euc}}}^1(\mathbb{R}^2) &= \text{span}\{dx, dy, xdy - ydx\} \cong \mathbb{R}^3, \\ \text{Kill}_{\nabla^{\text{Euc}}}^1(\mathbb{R}^2) \cap \Omega_{\text{exact}}^1(\mathbb{R}^2) &= \text{span}\{dx, dy\} \cong \mathbb{R}^2. \end{aligned}$$

*By Proposition 7.5, we get  $\dim H_{\bar{\nabla}^{\text{Euc}}}^1(T^*\mathbb{R}^2) \geq 5$ . Indeed, by the canonical identification  $(T^*\mathbb{R}^2, \bar{\nabla}^{\text{Euc}}) \cong (\mathbb{R}^4, \nabla^{\text{Euc}})$  and Section 6.4, we have  $H_{\bar{\nabla}^{\text{Euc}}}^1(\mathbb{R}^4) \cong \mathbb{R}^6$ .*

In some cases, the Patterson-Walker bound will be attained and Proposition 7.5 will allow us to compute  $H_{\bar{\nabla}}^1(T^*M)$ . The next example shows this for  $T^*\mathbb{R}^1$ , where the computations for a large class of connections are significantly simplified: instead of a system of PDEs, only a system of ODEs needs to be solved.

**Example 7.7** (The cartesian line). *Consider an arbitrary connection on  $\mathbb{R}^1$ . It is fully determined by  $f \in C^\infty(\mathbb{R}^1)$  via  $\nabla_{\partial_x}^f \partial_x := f\partial_x$ .*

A vector field  $h\partial_x \in \mathfrak{X}(\mathbb{R}^1)$ , where  $h \in C^\infty(\mathbb{R}^1)$ , is affine if and only if

$$0 = \frac{1}{2}(R_{\nabla^f}^s h\partial_x)(\partial_x, \partial_x) = (h'' + (fh)')\partial_x,$$

see Proposition 4.15 (note that  $R_{\nabla^f} = 0$  because  $\mathbb{R}^1$  is 1-dimensional). Function  $h$  solves the ODE

$$0 = h'' + (fh)'$$

if and only if there is a constant  $\lambda \in \mathbb{R}$  such that  $\lambda = h' + fh$ . This is a first-order non-homogeneous linear ODE with variable coefficients. Its general solution, see e.g. [5], is

$$h(x) = \mu e^{-F(x)} + \lambda e^{-F(x)} \int_0^x e^{F(x)} dx,$$

where  $F(x) := \int_0^x f(x)dx$  and  $\mu \in \mathbb{R}$  is an arbitrary constant. Moreover, the vector field  $h\partial_x$  is parallel if and only if

$$0 = \nabla_{\partial_x}^f(h\partial_x) = (h' + hf)\partial_x,$$

which is satisfied if and only if

$$h(x) = \mu e^{-F(x)},$$

where  $\mu \in \mathbb{R}$  is an arbitrary constant.

On the other hand, one easily finds that

$$\text{Kill}_{\nabla^f}^1(\mathbb{R}^1) \cong \text{span}\{e^{F(x)} dx\},$$

hence

$$\{\text{affine vector fields on } (\mathbb{R}^1, \nabla^f)\} \oplus \text{Kill}_{\nabla^f}^1(\mathbb{R}^1) \cong \mathbb{R}^3.$$

By combining Proposition 7.5 with the bound on Killing 1-tensors on  $\mathbb{R}^2$  given by Proposition 6.9, we have  $H_{\nabla^f}^1(T^*\mathbb{R}^1) \cong \mathbb{R}^1$ . Finally, by the canonical identification  $T^*\mathbb{R}^1 \cong \mathbb{R}^2$  and Proposition 5.10, we have proved that

$$H_{\nabla^f}^1(\mathbb{R}^2) \cong \mathbb{R}^1$$

for the torsion-free connections  $\bar{\nabla}^f$  on  $\mathbb{R}^2$  given by

$$\bar{\nabla}_{\partial_x}^f \partial_x = f(x)\partial_x + y(2f^2(x) - f'(x))\partial_y, \quad \bar{\nabla}_{\partial_x}^f \partial_y = -f(x)\partial_y, \quad \bar{\nabla}_{\partial_y}^f \partial_y = 0.$$

*Remark 7.8.* Note that there is an analogue of the Patterson-Walker bound for de Rham cohomology. It is well known that for any symplectic manifold  $(N, \omega)$

$$(44) \quad H_{\text{dR}}^1(N) \cong \frac{\{\text{symplectic vector fields on } (N, \omega)\}}{\{\text{Hamiltonian vector fields on } (N, \omega)\}}.$$

In particular, this applies for  $(T^*M, \omega_{\text{can}})$ . For  $\alpha \in \Omega^1(M)$ , we have that the vertical lift  $\alpha^\vee \in \mathfrak{X}(T^*M)$  is symplectic if and only if  $\alpha$  is closed and, moreover, it is Hamiltonian if and only if  $\alpha$  is exact. Therefore,

$$\dim H_{\text{dR}}^1(T^*M) \geq \dim H_{\text{dR}}^1(M).$$

Since the complete lift of a vector field is always Hamiltonian, there is no contribution from vector fields as opposed to the case of symmetric cohomology, where the bound is finer because of the left term of (43).

**7.3. Symmetric cohomology of left-invariant connections.** In this section we focus on Lie groups. Consider a Lie group  $G$ , a connection  $\nabla$  on  $G$  is called **left-invariant** if, for every  $g \in G$  and  $X, Y \in \mathfrak{X}(G)$ ,

$$L_{g*}\nabla_X Y = \nabla_{L_{g*}X} L_{g*}Y,$$

where  $L_g \in \text{Diff}(G)$  is given by  $L_g h = gh$  for every  $h \in G$ .

Left-invariant connections on a Lie group that are torsion-free are closely related to the notion of a **Lie-admissible algebra** [1], that is, an algebra  $(V, \cdot)$  such that

$$[u, v] := u \cdot v - v \cdot u.$$

makes  $(V, [ \ , \ ])$  a Lie algebra. Equivalently, it is an algebra  $(V, \cdot)$  satisfying

$$\text{skew assoc} = 0,$$

where  $\text{assoc} : \otimes^3 V \rightarrow V$  is the **associator** of  $(V, \cdot)$ . Explicitly, for  $u, v, w \in V$ ,

$$\text{assoc}(u, v, w) := u \cdot (v \cdot w) - (u \cdot v) \cdot w.$$

**Example 7.9.** *Every algebra up to dimension 2 is necessarily Lie-admissible. Moreover, any associative algebra, abelian algebra or Lie algebra is Lie-admissible. Another class of Lie-admissible algebras is called **left-symmetric algebras** [8], that is, an algebra  $(V, \cdot)$  satisfying*

$$\text{assoc}(u, v, w) = \text{assoc}(v, u, w).$$

We recall the proof of a known result (see, e.g., [18, 7] for the flat case) that will play an important role.

**Proposition 7.10.** *Let  $G$  be a connected Lie group and  $(\mathfrak{g}, [ \ , \ ]_{\mathfrak{g}})$  be its Lie algebra. There is the canonical one-to-one correspondence:*

$$\left\{ \begin{array}{l} \text{left-invariant torsion-free} \\ \text{connections on } G \end{array} \right\} \xleftrightarrow{\sim} \left\{ \begin{array}{l} \text{Lie-admissible algebras } (\mathfrak{g}, \cdot) \\ \text{such that } u \cdot v - v \cdot u = [u, v]_{\mathfrak{g}} \end{array} \right\}.$$

*Moreover, a Lie-admissible algebra is, in addition, left-symmetric if and only if the corresponding left-invariant torsion-free connection is flat.*

*Proof.* Let us start with a left-invariant torsion-free connection  $\nabla$  on  $G$ . We define an algebra structure  $\cdot$  on  $\mathfrak{g}$  by

$$u \cdot v := \nabla_{u^L} v^L \Big|_e,$$

where  $u^L \in \mathfrak{X}(G)$  is the left-invariant vector field corresponding to  $u \in \mathfrak{g}$  and  $e \in G$  is the unit element. As  $\nabla$  is torsion-free, we get

$$u \cdot v - v \cdot u = (\nabla_{u^L} v^L - \nabla_{v^L} u^L) \Big|_e = [u^L, v^L] \Big|_e = [u, v]_{\mathfrak{g}}.$$

Conversely, if  $\cdot$  is a Lie-admissible algebra structure on  $\mathfrak{g}$ , whose skew-symmetric projection gives the Lie algebra structure, we define a map on the left-invariant vector fields  $\nabla : \mathfrak{X}_L(G) \rightarrow \text{End}(\mathfrak{X}_L(G))$  by

$$\nabla_X Y := (X_e \cdot Y_e)^L.$$

Since left-invariant vector fields are global generators for  $\mathfrak{X}(G)$ , the map  $\nabla$  determines a connection on  $G$ . For the same reason, it is enough to check the left-invariance and torsion-freeness of  $\nabla$  only on left-invariant vector fields, but this is clearly satisfied. The result follows because the two constructions are inverse to each other. The statement about flatness follows from that, for  $X, Y, Z \in \mathfrak{X}_L(G)$ , we have

$$\begin{aligned} R_\nabla(X, Y)Z &= (X_e \cdot (Y_e \cdot Z_e))^L - (Y_e \cdot (X_e \cdot Z_e))^L - ([X_e, Y_e]_{\mathfrak{g}} \cdot Z)^L \\ &= (X_e \cdot (Y_e \cdot Z_e) - Y_e \cdot (X_e \cdot Z_e) - (X_e \cdot Y_e - Y_e \cdot X_e) \cdot Z)^L \\ &= \text{assoc}(X_e, Y_e, Z_e) - \text{assoc}(Y_e, X_e, Z_e). \end{aligned}$$

□

Given a Lie algebra  $(\mathfrak{g}, [ \cdot, \cdot ]_{\mathfrak{g}})$ , there is always an underlying Lie-admissible algebra. To show this, consider an arbitrary linear Riemannian metric  $\langle \cdot, \cdot \rangle$  on  $\mathfrak{g}$  and define the algebra structure  $\cdot$  by

$$\langle u \cdot v, w \rangle := \frac{1}{2}(\langle [u, v]_{\mathfrak{g}}, w \rangle - \langle [v, w]_{\mathfrak{g}}, u \rangle - \langle [u, w]_{\mathfrak{g}}, v \rangle).$$

On the level of torsion-free left-invariant connections on a Lie group, this corresponds to the fact that there is always a left-invariant Riemannian metric on a Lie group and its Levi-Civita connection is also left-invariant.

For every Lie-admissible algebra  $(V, \cdot)$ , we define a canonical derivation  $d^s \in \text{Der}_1(\text{Sym}^\bullet V^*)$  completely determined by its action on the elements of  $V^*$ :

$$(45) \quad (d^s \zeta)(u, v) := -\zeta(u \cdot v + v \cdot u).$$

It vanishes on scalars, as all derivations of  $\text{Sym}^\bullet V^*$  do, and one can check by a straightforward calculation that, for  $\zeta \in \text{Sym}^r V^*$  and  $r > 0$ , it is given by

$$(d^s \zeta)(u_1, \dots, u_{r+1}) = - \sum_{\substack{i, j=1, \\ i < j}}^{r+1} \zeta(u_i \cdot u_j + u_j \cdot u_i, u_1, \dots, \hat{u}_i, \dots, \hat{u}_j, \dots, u_{r+1}).$$

We can then define its associated cohomology.

**Definition 7.11.** Let  $(V, \cdot)$  be a Lie admissible algebra, we define the  $r$ th-**Lie-admissible symmetric cohomology group**:

$$H^r(V, \cdot) := \frac{\ker d^s|_{\text{Sym}^r V^*}}{\ker d^s|_{\text{Sym}^r V^*} \cap \text{Im } d^s|_{\text{Sym}^{r-1} V^*}}.$$

As  $d^s$  vanishes on scalars, we have that, for any real Lie-admissible algebra  $(V, \cdot)$ ,

$$H^0(V, \cdot) = \mathbb{R}^1, \quad H^1(V, \cdot) \cong \ker d^s|_{V^*}.$$

**Example 7.12.** *On every real finite-dimensional vector space  $V$ , there is always the trivial Lie-admissible algebra structure:*

$$u \cdot_{\text{triv}} v := 0$$

for every  $u, v \in V$ . Then  $d^s = 0$  and  $H^r(V, \cdot_{\text{triv}}) \cong \text{Sym}^r V^*$ . Clearly, the maximum possible dimension of  $H^r(V, \cdot)$  is attained in this case.

**Example 7.13.** *Consider a 1-dimensional real vector space  $V$ . Besides the trivial Lie-admissible algebra structure, there is (up to an isomorphism) only one more algebra structure on  $V$ . It can be represented, for some nonzero  $u \in V$ , by*

$$u \cdot u := u.$$

It is straightforward to check that  $\text{Im } d^s|_{\text{Sym}^r V^*} = \text{Sym}^{r+1} V^*$  for  $r > 0$ , hence

$$H^r(V, \cdot) = \begin{cases} \mathbb{R}^1 & \text{for } r = 0, \\ \{0\} & \text{for } r > 0. \end{cases}$$

Let us now translate the Lie-admissible symmetric cohomology to the language of left-invariant torsion-free connections. We say that a symmetric form  $\varphi \in \Upsilon^\bullet(G)$  is **left-invariant** if  $L_g^* \varphi = \varphi$  and denote the space of such forms by  $\Upsilon_L^\bullet(G)$ . Analogously, the symbol  $\text{Kill}_{L, \nabla}^r(G)$  denotes the space of left-invariant Killing  $r$ -tensors. If  $\nabla$  is a left-invariant connection on  $G$ , we have, by Remark 4.10, that  $\nabla^s(\Upsilon_L^\bullet(G)) \subseteq \Upsilon_L^\bullet(G)$ . Therefore, we define the  **$r$ th-left-invariant symmetric cohomology group** of  $(G, \nabla)$  by

$$H_{L, \nabla}^r(G) := \frac{\text{Kill}_{L, \nabla}^r(G)}{\text{Kill}_{L, \nabla}^r(G) \cap \text{Im } \nabla^s|_{\Upsilon_L^{r-1}(G)}}.$$

Let  $G$  be a connected Lie group integrating a Lie algebra  $\mathfrak{g}$ . For  $\varphi \in \Upsilon_L^\bullet(G)$ , the assignment  $\varphi \mapsto \varphi_e$  gives an isomorphism of unital graded algebras

$$\Upsilon_L^\bullet(G) \cong \text{Sym}^\bullet \mathfrak{g}^*.$$

The inverse, for  $\zeta \in \text{Sym}^\bullet \mathfrak{g}^*$ , is given by  $\zeta \mapsto \zeta^L$ , where  $\zeta_g^L = L_{g^{-1}}^* \zeta$  for  $g \in G$ .

**Lemma 7.14.** *Let  $(G, \nabla)$  be a connected Lie group with a left-invariant torsion-free connection and  $(\mathfrak{g}, \cdot)$  be the corresponding Lie-admissible algebra. Then*

$$H_{L, \nabla}^r(G) \cong H^r(\mathfrak{g}, \cdot).$$

*Proof.* It is enough to check that the isomorphism between  $\Upsilon_L^\bullet(G)$  and  $\text{Sym}^\bullet \mathfrak{g}^*$  intertwines the operators  $\nabla^s$  and  $d^s$ . For every  $\varphi \in \Upsilon_L^r(G)$ ,



$$\begin{aligned}
 (d^s \varphi_e)(u_1, \dots, u_{r+1}) &= - \sum_{\substack{i,j=1, \\ i < j}}^{r+1} \varphi_e(u_i \cdot u_j + u_j \cdot u_i, u_1, \dots, \hat{u}_i, \dots, \hat{u}_j, \dots, u_{r+1}) \\
 &= - \sum_{\substack{i,j=1, \\ i < j}}^{r+1} \varphi((u_i \cdot u_j + u_j \cdot u_i)^L, u_1^L, \dots, \hat{u}_i^L, \dots, \hat{u}_j^L, \dots, u_{r+1}^L) \\
 &= - \sum_{\substack{i,j=1, \\ i < j}}^{r+1} \varphi(\langle u_i^L : u_j^L \rangle_s, u_1^L, \dots, \hat{u}_i^L, \dots, \hat{u}_j^L, \dots, u_{r+1}^L) \\
 &= (\nabla^s \varphi)(u_1^L, \dots, u_{r+1}^L) = (\nabla^\varphi)_e(u_1, \dots, u_{r+1}).
 \end{aligned}$$

□

Both  $\mathbb{R}^1$  and  $S^1$  are connected Lie groups integrating the 1-dimensional abelian Lie algebra. Left-invariant connections on  $\mathbb{R}^1$  and  $S^1$  are determined by the choice of a real constant  $\lambda \in \mathbb{R}$ :

$$\nabla_{\partial_x}^\lambda \partial_x := \lambda \partial_x, \quad \nabla_{\partial_\theta}^\lambda \partial_\theta := \lambda \partial_\theta.$$

In particular, for  $\lambda = 0$ , we have the Euclidean connection  $\nabla^{\text{Euc}}$  on  $\mathbb{R}^1$  and the round connection  $\nabla^\circ$  on  $S^1$ . By Lemma 7.14 and Examples 7.12, 7.13, we get

$$\begin{aligned}
 H_{L, \nabla^{\text{Euc}}}^r(\mathbb{R}^1) &\cong H_{L, \nabla^\circ}^r(S^1) \cong \mathbb{R}^1, \\
 H_{L, \nabla^\lambda}^r(\mathbb{R}^1) &\cong H_{L, \nabla^\lambda}^r(S^1) \cong \begin{cases} \mathbb{R}^1 & \text{for } r = 0, \\ \{0\} & \text{for } r > 0. \end{cases} \quad (\text{for } \lambda \neq 0)
 \end{aligned}$$

Comparing these with the results of Sections 6.4 and 6.5, we see that, for  $r = 1$ , the left-invariant symmetric cohomology group is isomorphic to the symmetric cohomology group for  $S^1$ , whereas for  $\mathbb{R}^1$  this is not the case.

The following result is a general statement about the compact case.

**Theorem 7.15.** *Let  $(G, \nabla)$  be a connected compact Lie group with a left-invariant torsion-free connection and  $(\mathfrak{g}, \cdot)$  be the corresponding Lie-admissible algebra. Then we have an injective vector space morphism*

$$H^r(\mathfrak{g}, \cdot) \rightarrow H_{\nabla}^r(G).$$

*Proof.* By Lemma 7.14, the inclusion map  $i : \text{Sym}^\bullet \mathfrak{g}^* \cong \Upsilon_L^\bullet(G) \hookrightarrow \Upsilon^\bullet(G)$  intertwines  $d^s$  and  $\nabla^s$ , hence it descends to the map  $i : H^r(\mathfrak{g}, \cdot) \rightarrow H_{\nabla}^r(G)$ . We show that this map is an injective vector space morphism by finding a left-inverse to it. We introduce the map  $\mathfrak{L} : \Upsilon^\bullet(G) \rightarrow \text{Sym}^\bullet \mathfrak{g}^*$ , for every  $\varphi \in \Upsilon^r(G)$  and  $u_j \in \mathfrak{g}$ , by

$$(46) \quad (\mathfrak{L}\varphi)(u_1, \dots, u_r) := \int_G (L_g^* \varphi)_e(u_1, \dots, u_r) \text{vol}_G,$$

where  $\text{vol}_G \in \det(G)$  is the *normalized left-invariant Haar measure*. Note that the map  $\mathfrak{L}$  is well defined because of the compactness of  $G$ . Using the left-invariance of  $\nabla$ , for every  $u \in \mathfrak{g}$ , yields

$$\begin{aligned} & (\mathfrak{L}\nabla^s\varphi)(u, \dots, u) \\ &= \int_G (\mathbb{L}_g^*\nabla^s\varphi)_e(u, \dots, u)\text{vol}_G = \int_G (\nabla^s\mathbb{L}_g^*\varphi)_e(u, \dots, u)\text{vol}_G \\ &= (r+1) \int_G (u^L\varphi(u^L))(g)\text{vol}_G - \frac{r(r+1)}{2} \int_G (\mathbb{L}_g^*\varphi)(\langle u^L : u^L \rangle_s, u^L, \dots, u^L)(e)\text{vol}_G, \end{aligned}$$

where we used the shorthand notation  $\varphi(u^L) := \varphi(u^L, \dots, u^L)$ . On the other hand, it follows from the correspondence of Proposition 7.10 that

$$\begin{aligned} (d^s\mathfrak{L}\varphi)(u, \dots, u) &= -\frac{r(r+1)}{2} \int_G (\mathbb{L}_g^*\varphi)_e(2u \cdot u, u, \dots, u)\text{vol}_G \\ &= -\frac{r(r+1)}{2} \int_G ((\mathbb{L}_g^*\varphi)(\langle u^L : u^L \rangle_s, u^L, \dots, u^L))(e)\text{vol}_G. \end{aligned}$$

To prove that  $\mathfrak{L}$  intertwines  $d^s$  and  $\nabla^s$ , it remains to show that the integral  $\int_G (u^L\varphi(u^L))\text{vol}_G$  vanishes. We have  $L_{u^L}\text{vol}_G = 0$  for the Lie derivative because both  $\text{vol}_G$  and  $u^L$  are left-invariant. Therefore,

$$(u^L\varphi(u^L))\text{vol}_G = (L_{u^L}\varphi(u^L))\text{vol}_G = L_{u^L}(\varphi(u^L)\text{vol}_G) = d_{u^L}(\varphi(u^L)\text{vol}_G),$$

hence, by *Stokes's theorem*, the integral  $\int_G (u^L\varphi(u^L))\text{vol}_G$  vanishes. It means that the map (46) descends to a map  $\mathfrak{L} : H^r(G) \rightarrow H^r(\mathfrak{g}, \cdot)$  and the result follows from  $\mathfrak{L} \circ i = \text{id}$ .  $\square$

As we have seen above (for the 1st-symmetric cohomology group of the circle with a left-invariant connection), we may have an isomorphism between  $H^r(\mathfrak{g}, \cdot) \cong$  and  $H^r(G)$ . Let us show that this is not always the case but that we will generally obtain a lower bound in the dimension.

**Example 7.16** (The round 3-sphere). *The 3-sphere  $S^3$  is isomorphic to  $SU(2)$ , it is thus a connected compact 3-dimensional Lie group. The round metric is left-invariant, hence its Levi-Civita connection is also left-invariant. Recall from Section 6.5 that  $H_{\nabla^\circ}^1(S^3) \cong \mathbb{R}^6$ . However, by Example 7.12, the dimension of  $H^1(\mathfrak{su}(2), \cdot)$  is bounded from above by  $\dim \mathfrak{su}(2) = 3$ . Therefore,  $H^1(\mathfrak{su}(2), \cdot)$  and  $H_{\nabla^\circ}^1(S^3)$  are not isomorphic in this case.*

**Example 7.17** (The flat  $n$ -torus). *The  $n$ -torus  $\mathbb{T}^n$ , for  $n \geq 1$ , is a connected compact Lie group whose Lie algebra  $\mathfrak{g}$  is abelian. The trivial Lie-admissible algebra structure on  $\mathfrak{g}$  corresponds to the choice of the flat connection  $\nabla^{\text{flat}}$  on  $\mathbb{T}^n$ , that is, the product connection of the round connections on  $S^1$ . By Theorem 7.15, we get*

$$\dim H_{\nabla^{\text{flat}}}^r(\mathbb{T}^n) \geq \dim H^r(\mathfrak{g}, \cdot_{\text{triv}}) = \binom{n+r-1}{r}.$$

## APPENDIX A. NON-GEOMETRIC DERIVATIONS

We have taken the symmetric derivative  $\nabla^s$  to be the starting point of symmetric Cartan calculus. Although we have explained why it is natural that  $\nabla^s$  does not square to zero, this resulted in an extra adjustment to define symmetric cohomology. A natural question is whether there are derivations of  $\Upsilon^\bullet(M)$  of degree 1, not necessarily geometric, that square to zero.

The derivations of  $\Upsilon^\bullet(M)$  were classified in [16]. In particular, derivations of degree 1 are parametrized by pairs  $(A, \sigma)$  consisting of  $A \in \Gamma(\text{End } TM)$  and  $\sigma \in \Upsilon^2(M, TM)$ . Explicitly, any  $D \in \text{Der}_1(\Upsilon^\bullet(M))$  is of the form

$$D = [\iota_A^s, \nabla^s] + \iota_\sigma^s = \nabla_A^s + \iota_\sigma^s$$

for some auxiliary symmetric derivative  $\nabla^s$ . The geometric derivations are precisely those with  $A = \text{id}_{TM}$ , see Remark 3.4 and Proposition 3.8.

**Proposition A.1.** *The only degree-1 derivation of  $\Upsilon^\bullet(M)$  that squares to zero is the trivial one,  $D = 0$ .*

*Proof.* Consider a derivation  $D \in \text{Der}_1(\Upsilon^\bullet(M))$  described by a pair  $(A, \sigma)$  as above. Assume that  $D \circ D = 0$ . We then have

$$0 = D(Df^2) = D(2fDf) = 2Df \odot Df,$$

hence  $D|_{\mathcal{C}^\infty(M)} = 0$ . In terms of the corresponding  $(A, \sigma)$ , it means  $(AX)f = 0$  for all  $X \in \mathfrak{X}(M)$  and  $f \in \mathcal{C}^\infty(M)$ . Therefore,  $A = 0$  and  $D = \iota_\sigma^s$ . In addition,

$$0 = (D(D\alpha))(X, X, X) = 3(\iota_\sigma^s \alpha)(\sigma(X, X), X) = 3\alpha(\sigma(\sigma(X, X), X))$$

for all  $\alpha \in \Upsilon^1(M)$ , so  $\sigma(\sigma(X, X), X) = 0$ . Finally, for every  $C \in \Upsilon^2(M)$ ,

$$\begin{aligned} 0 &= (D(DC))(X, X, X, X) = (3!)(\iota_\sigma^s C)(\sigma(X, X), X, X) \\ &= 12C(\sigma(\sigma(X, X), X), X) + 6C(\sigma(X, X), \sigma(X, X)). \end{aligned}$$

Since  $D \circ D|_{\Upsilon^1(M)} = 0$ , we get  $C(\sigma(X, X), \sigma(X, X)) = 0$ . If we choose  $C$  to be a Riemannian metric, this yields  $\sigma(X, X) = 0$ . By polarization, we get  $\sigma = 0$  and thus  $D = 0$ .  $\square$

This proposition reinforces the claim that the symmetric derivative is indeed the natural analogue of the exterior derivative.

## APPENDIX B. VARIOUS LIFTS TO THE COTANGENT BUNDLE

In this section, we recall several lifts from the base manifold to the total space of the cotangent bundle  $\text{pr} : T^*M \rightarrow M$  that are used in Section 5.

The **vertical subbundle**  $\mathcal{V} = \ker \text{pr}_* \subseteq T(T^*M)$  is a subbundle of rank  $\dim M$  canonically isomorphic to  $\text{pr}^*T^*M$ . The **vertical lift** of a 1-form is defined as

$$\begin{aligned} \Omega^1(M) &\rightarrow \Gamma(\mathcal{V}) \subseteq \mathfrak{X}(T^*M) \\ \alpha &\mapsto \alpha^v := \text{pr}^*\alpha, \end{aligned}$$

where  $\text{pr}^*\alpha \in \Gamma(\text{pr}^*T^*M)$  denotes the pullback section.

A connection  $\nabla$  on  $M$  induces, through the dual connection, the **horizontal subbundle**  $\mathcal{H}_\nabla \leq T(T^*M)$ , of rank  $\dim M$  and satisfying  $T(T^*M) = \mathcal{H}_\nabla \oplus \mathcal{V}$ . It is canonically isomorphic to  $\text{pr}^*TM$ , which gives the **horizontal lift**

$$\begin{aligned} \mathfrak{X}(M) &\rightarrow \Gamma(\mathcal{H}_\nabla) \subseteq \mathfrak{X}(T^*M) \\ X &\mapsto X^h := \text{pr}^*X. \end{aligned}$$

*Remark B.1.* In natural local coordinates  $(T^*U, \{x^i\} \cup \{p_j\})$  on  $T^*M$ , we have

$$\begin{aligned} \alpha|_U &= \alpha_j dx^j, & \alpha^v|_{T^*U} &= (\text{pr}^*\alpha_j)\partial_{p_j}, \\ X|_U &= X^i \partial_{x^i}, & X^h|_{T^*U} &= (\text{pr}^*X^i)(\partial_{x^i} + p_k(\text{pr}^*\Gamma_{ji}^k)\partial_{p_j}), \end{aligned}$$

where  $\{\Gamma_{ij}^k\}$  is the set of Christoffel symbols of  $\nabla$  corresponding to  $(U, \{x^j\})$ .

We recall now three more lifts and their properties from [27]. The **vertical lift of a vector field** is a map  $(\ )^v : \mathfrak{X}(M) \rightarrow \mathcal{C}^\infty(T^*M)$ . It is defined by the formula

$$X^v(\zeta) := \zeta(X_{\text{pr}(\zeta)})$$

for all  $\zeta \in T^*M$ . The importance of the vertical lift of a vector field can be seen through the following proposition.

**Proposition B.2** ([27]). *A vector field on  $T^*M$  is fully determined by its action on vertical lifts of all vector fields on  $M$ .*

This proposition allows us to give an elegant definition to the **vertical lift of a field of endomorphisms**, which is the map  $(\ )^v : \Gamma(\text{End } TM) \rightarrow \mathfrak{X}(T^*M)$  fully determined by the relation

$$A^v X^v := (AX)^v.$$

In addition to the vertical and horizontal lifts, one can consider also the **complete lift**. It is the map  $(\ )^c : \mathfrak{X}(M) \rightarrow \mathfrak{X}(T^*M)$  determined by the relation

$$X^c Y^v := [X, Y]^v.$$

Alternatively, the complete lift of  $X \in \mathfrak{X}(M)$  can be defined (up to sign) as the Hamiltonian vector field of the vertical lift of  $X$ :

$$(47) \quad X^c = -\omega_{\text{can}}^{-1}(dX^v) = -\text{Ham } X^v.$$

Let us recall two important properties of the complete lift.

**Proposition B.3** ([27]). *A tensor field that is a section of  $\otimes^r T^*M \otimes TM$  or  $\otimes^r T^*M$  is fully determined by its values on complete lifts of vector fields on  $M$ .*

In particular, every element of  $\Upsilon^\bullet(M)$  and  $\Upsilon^\bullet(M, TM)$  is fully determined by its values on complete lifts.

**Proposition B.4** ([27]). *For every  $X, Y \in \mathfrak{X}(M)$ , we have  $[X^c, Y^c] = [X, Y]^c$ .*

*Remark B.5.* In natural local coordinates, for  $X \in \mathfrak{X}(M)$  and  $A \in \Gamma(\text{End } TM)$ ,

$$\begin{aligned} X|_U &= X^i \partial_{x^i}, & X^v|_{T^*U} &= p_i(\text{pr}^* X^i), \\ X^c|_{T^*U} &= (\text{pr}^* X^i) \partial_{x^i} - p_i \left( \text{pr}^* \frac{\partial X^i}{\partial x^j} \right) \partial_{p_j}, \\ A|_U &= A_j^i dx^j \otimes \partial_{x^i}, & A^v|_{T^*U} &= p_i(\text{pr}^* A_j^i) \partial_{p_j}. \end{aligned}$$

Finally, we can state the lemmas that allow us to prove the results of Section 5.

**Lemma B.6.** *Let  $\nabla$  be a connection on  $M$ . For  $X \in \mathfrak{X}(M)$ ,  $\alpha \in \Omega^1(M)$  and  $f \in \mathcal{C}^\infty(M)$ ,*

$$\alpha_{\text{can}}(X^h) = X^v, \quad X^c = X^h - (\nabla X)^v, \quad X^h(\text{pr}^* f) = \text{pr}^*(Xf), \quad \alpha^v(\text{pr}^* f) = 0.$$

*Proof.* It follows by a straightforward calculation in natural coordinates on  $T^*M$ .  $\square$

Taking into account Proposition B.2, we can write explicit formulas that fully determine the horizontal lift of a vector field and the vertical lift of a 1-form.

**Lemma B.7.** *Let  $\nabla$  be a connection on  $M$ . For  $X, Y \in \mathfrak{X}(M)$  and  $\alpha \in \Omega^1(M)$ ,*

$$X^h Y^v = (\nabla_X Y)^v, \quad \alpha^v Y^v = \text{pr}^* \alpha(Y).$$

*Proof.* In natural coordinates, using Lemma B.6, we have

$$\begin{aligned} (X^h Y^v)|_{T^*U} &= X^h(p_l(\text{pr}^* Y^l)) = p_l(\text{pr}^*(X(Y^l))) + (\text{pr}^* X^i)(\text{pr}^* Y^j)(\text{pr}^* \Gamma_{ji}^l) \\ &= (\nabla_X Y)^v|_{T^*U}, \end{aligned}$$

where  $X|_U = X^i \partial_{x^i}$ . On the other hand,

$$(\alpha^v Y^v)|_{T^*U} = (\text{pr}^* \alpha_j) \partial_{p_j}(p_l(\text{pr}^* Y^l)) = (\text{pr}^* \alpha_j)(\text{pr}^* Y^j) = \text{pr}^* \alpha(Y)|_{T^*U},$$

where  $\alpha|_U = \alpha_j dx^j$ . Thus, also globally,  $X^h Y^v = (\nabla_X Y)^v$  and  $\alpha^v Y^v = \text{pr}^* \alpha(Y)$ .  $\square$

We finish by describing the Lie brackets between lifts.

**Lemma B.8.** *Let  $\nabla$  be a connection on  $M$ . For  $X, Y \in \mathfrak{X}(M)$  and  $\alpha, \beta \in \Omega^1(M)$ ,*

$$[X^h, Y^h] = [X, Y]^h + R_\nabla(X, Y)^v, \quad [X^h, \alpha^v] = (\nabla_X \alpha)^v, \quad [\alpha^v, \beta^v] = 0.$$

*Proof.* It follows from Lemmas B.6 and B.7, for  $Z \in \mathfrak{X}(M)$ , that

$$\begin{aligned} [X^h, Y^h]Z^v &= (\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z)^v = (\nabla_{[X, Y]} Z + R_\nabla(X, Y)Z)^v \\ &= ([X, Y]^h + R_\nabla(X, Y)^v)Z^v \\ [X^h, \alpha^v]Z^v &= X^h \text{pr}^* \alpha(Z) - \text{pr}^* \alpha(\nabla_X Z) = \text{pr}^*(\nabla_X \alpha)(Z) = (\nabla_X \alpha)^v Z^v \\ [\alpha^v, \beta^v]Z^v &= \alpha^v \text{pr}^* \beta(Z) - \beta^v \text{pr}^* \alpha(Z) = 0. \end{aligned}$$

The claim then follows from Proposition B.2.  $\square$

APPENDIX C. COMPUTATIONS FOR  $H_{\nabla}^1(\mathbb{R}^2)$ 

In this section, we enclose the computations supporting our claims in Examples 6.18, 6.20 and 6.21. In all of them, the system of PDEs (36) reduces to

$$0 = \frac{\partial \alpha_1}{\partial x}, \quad 0 = \frac{\partial \alpha_1}{\partial y} + \frac{\partial \alpha_2}{\partial x} - f_3 \alpha_1 - f_4 \alpha_2, \quad 0 = \frac{\partial \alpha_2}{\partial y}.$$

From the first and the third equation,  $\alpha_1$  and  $\alpha_2$  can be seen as smooth functions of, respectively, the single variable  $y$  and  $x$ . The second equation then becomes

$$(48) \quad 0 = \alpha_1'(y) + \alpha_2'(x) - f_3(x, y)\alpha_1(y) - f_4(x, y)\alpha_2(x).$$

*Example 6.18.* If  $f_3(x, y) = xy$  and  $f_4(x, y) = y$ , equation (48) reads as

$$(49) \quad 0 = \alpha_1'(y) + \alpha_2'(x) - xy\alpha_1(y) - y\alpha_2(x).$$

This has to be satisfied for all  $x, y \in \mathbb{R}$ . In particular, we get

$$\begin{aligned} 0 &= \alpha_1'(y) + \alpha_2'(0) - y\alpha_2(0) && \text{for } x = 0, \\ 0 &= \alpha_1'(0) + \alpha_2'(x) && \text{for } y = 0. \end{aligned}$$

We can easily solve these ODEs as

$$\alpha_1(y) = \alpha_2(0)\frac{y^2}{2} - \alpha_2'(0)y + \lambda_1, \quad \alpha_2(x) = -\alpha_1'(0)x + \lambda_2$$

for arbitrary constants  $\lambda_1, \lambda_2 \in \mathbb{R}$ . Replacing these expressions back in (49),

$$0 = \alpha_2(0)y - \alpha_2'(0) - \alpha_1'(0) - \alpha_2(0)\frac{xy^3}{2} + \alpha_2'(0)xy^2 - \lambda_1xy + \alpha_1'(0)xy - \lambda_2y.$$

Therefore,  $\alpha_2(0) = \alpha_2'(0) = \lambda_1 = \alpha_1'(0) = \lambda_2 = 0$ , which means  $\alpha_1 = 0 = \alpha_2$ .

In the case of Examples 6.20 and 6.21, we have that  $f_3$  and  $f_4$  depends only on  $y$  and  $x$  respectively. So, (48) is equivalently rewritten as

$$\alpha_1'(y) - f_3(y)\alpha_1(y) = -\alpha_2'(x) + f_4(x)\alpha_2(x).$$

This is satisfied if and only if there is a constant  $C \in \mathbb{R}$  such that

$$\alpha_1'(y) - f_3(y)\alpha_1(y) = C, \quad \alpha_2'(x) - f_4(x)\alpha_2(x) = -C.$$

These are *first-order non-homogeneous linear ODEs with variable coefficients*. It is well-known how to solve such differential equations in general, see e.g. [5].

## APPENDIX D. COMPARISON WITH CLASSICAL CARTAN CALCULUS

We show in this table a comparison between classical and symmetric Cartan calculus.

Classical Cartan calculus	Symmetric Cartan calculus
algebraic features	
graded-commutative algebra $(\Omega^\bullet(M), \wedge)$ graded derivations $\text{gDer}(\Omega^\bullet(M))$ graded commutator $[\ , \ ]_g$ $[\iota_X, \iota_Y]_g = 0$	commutative algebra $(\Upsilon^\bullet(M), \odot)$ derivations $\text{Der}(\Upsilon^\bullet(M))$ commutator $[\ , \ ]$ $[\iota_X, \iota_Y] = 0$
differentials	
exterior derivative $d$ canonical $(df)(X) = Xf$ $d \in \text{gDer}_1(\Omega^\bullet(M))$ $[d, d]_g = 2d \circ d = 0$ (non-trivial condition)	symmetric derivative $\nabla^s$ depending on the choice of $\nabla$ $(\nabla^s f)(X) = Xf$ $\nabla^s \in \text{Der}_1(\Upsilon^\bullet(M))$ $[\nabla^s, \nabla^s] = 0$ (trivially satisfied)
Lie derivatives	
Lie derivative $L_X := [\iota_X, d]_g$ $(L_X \varphi)_m = \frac{d}{dt} \Big _{t=0} (\Psi_t^X)^* \varphi_{\Psi_t^X(m)}$	symmetric Lie derivative $L_X^s := [\iota_X, \nabla^s]$ $(L_X^s \varphi)_m = \frac{d}{dt} \Big _{t=0} P_{2t,0}^\gamma (\Psi_{-t}^X)^*_{\Psi_{2t}^X(m)} \varphi_{\Psi_t^X(m)}$
brackets	
Lie bracket $[X, Y] := X \circ Y - Y \circ X$ $\iota_{[X,Y]} = [L_X, \iota_Y]_g, \quad [X, Y] = L_X Y$ foliations	symmetric bracket $\langle X : Y \rangle_s := \nabla_X Y + \nabla_Y X$ $\iota_{\langle X:Y \rangle_s} = [L_X^s, \iota_Y], \quad \langle X : Y \rangle_s = L_X^s Y$ geodesically invariant distributions
(graded-)commutation relations	
$[L_X, L_Y]_g = L_{[X,Y]}$ $[d, L_X]_g = 0$	$[L_X^s, L_Y^s] = L_{[X,Y]}^s + \iota_{2(\nabla_{\nabla X} Y - \nabla_{\nabla Y} X - R_{\nabla}(X,Y))}$ $[\nabla^s, L_X^s] = 2\nabla_{\nabla X}^s + \iota_{2\text{sym} \iota_X R_{\nabla} + R_{\nabla}^s X}$

Classical Cartan calculus	Symmetric Cartan calculus
isomorphisms	
diffeomorphisms	affine diffeomorphisms
induced geometry on the cotangent bundle	
symplectic geometry canonical 2-form $\omega_{\text{can}}$ $\omega_{\text{can}} _{T^*U} = dp_j \wedge dx^j$ $\omega_{\text{can}} = d\alpha_{\text{can}}$ $\omega_{\text{can}}(X^c, Y^c) = -[X, Y]^v$ $\omega_{\text{can}}(\phi_{\nabla}a, \phi_{\nabla}b) = \text{pr}^*\langle a, b \rangle_-$	geometry of split signature metrics Patterson-Walker metric $g_{\nabla}$ $g_{\nabla} _{T^*U} = dp_j \odot dx^j - p_k \Gamma_{ij}^k dx^i \odot dx^j$ $g_{\nabla} = \hat{\nabla}^s \alpha_{\text{can}}$ $g_{\nabla}(X^c, Y^c) = -\langle X : Y \rangle_s^v$ $g_{\nabla}(\phi_{\nabla}a, \phi_{\nabla}b) = \text{pr}^*\langle a, b \rangle_+$
simplified (graded-)commutation relations	
$[L_X, L_Y]_{\text{g}} = L_{[X, Y]}, \quad [d, L_X]_{\text{g}} = 0$ $\Leftrightarrow$ $X^c, Y^c$ are Hamiltonian for $\omega_{\text{can}}$ (trivially satisfied)	$[L_X^s, L_Y^s] = 0 (= L_{\langle X : Y \rangle_s^s}^s), \quad [\nabla^s, L_X^s] = 0$ $\Leftrightarrow$ $X^c, Y^c$ are gradient Killing for $g_{\nabla}$ (non-trivial condition)
cohomology	
de Rham cohomology $\ker d = \Omega_{\text{cl}}^{\bullet}(M)$ (closed forms) $\text{Im } d = \Omega_{\text{exact}}^{\bullet}(M)$ (exact forms)	symmetric cohomology $\ker \nabla^s = \text{Kill}_{\nabla}^{\bullet}(M)$ (Killing tensors) $\text{Im } \nabla^s \cap \text{Kill}_{\nabla}^{\bullet}(M)$ ( $\nabla^s$ -exact tensors)



## REFERENCES

- [1] A. A. Albert. Power-associative rings. *Trans. Amer. Math. Soc.*, 64:552–593, 1948.
- [2] A. Alekseev and P. Xu. Derived brackets and Courant algebroids. <https://web.archive.org/web/20160705150153/http://www.math.psu.edu/ping/anton-final.pdf>. Unpublished notes.
- [3] María Barbero-Liñán and Andrew D. Lewis. Geometric interpretations of the symmetric product in affine differential geometry and applications. *Int. J. Geom. Methods Mod. Phys.*, 9(8):1250073, 33, 2012.
- [4] Sergio Benenti. Separability in Riemannian manifolds. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 12:Paper No. 013, 21, 2016.
- [5] W. E. Boyce, R. C. DiPrima, and D. B. Meade. *Elementary Differential Equations and Boundary Value Problems*. Wiley, 2017.
- [6] Francesco Bullo and Andrew D. Lewis. *Geometric control of mechanical systems*, volume 49 of *Texts in Applied Mathematics*. Springer-Verlag, New York, 2005. Modeling, analysis, and design for simple mechanical control systems.
- [7] Dietrich Burde. Affine structures on nilmanifolds. *Internat. J. Math.*, 7(5):599–616, 1996.
- [8] Dietrich Burde. Left-symmetric algebras, or pre-Lie algebras in geometry and physics. *Cent. Eur. J. Math.*, 4(3):323–357, 2006.
- [9] B. Carter. Global structure of the Kerr family of gravitational fields. *Physical Review*, 174(5):1559–1571, 1968.
- [10] Brandon Carter. Hamilton-Jacobi and Schrödinger separable solutions of Einstein’s equations. *Comm. Math. Phys.*, 10:280–310, 1968.
- [11] Ted Courant. Tangent Lie algebroids. *J. Phys. A*, 27(13):4527–4536, 1994.
- [12] P. E. Crouch. Geometric structures in systems theory. *Proc. IEE-D*, 128(5):242–252, 1981.
- [13] Michael Eastwood. Higher symmetries of the Laplacian. *Ann. of Math. (2)*, 161(3):1645–1665, 2005.
- [14] Marco Gualtieri. Generalized complex geometry. *Oxford University doctoral thesis*, [arXiv:0401221], 2004.
- [15] Marco Gualtieri. Branes on Poisson varieties. In *The many facets of geometry*, pages 368–394. Oxford Univ. Press, Oxford, 2010.
- [16] A. Heydari, N. Boroojerdian, and E. Peyghan. A description of derivations of the algebra of symmetric tensors. *Arch. Math. (Brno)*, 42(2):175–184, 2006.
- [17] Nigel Hitchin. Generalized Calabi-Yau manifolds. *Q. J. Math.*, 54(3):281–308, 2003.
- [18] Hyuk Kim. Complete left-invariant affine structures on nilpotent Lie groups. *J. Differential Geom.*, 24(3):373–394, 1986.
- [19] Shoshichi Kobayashi. *Transformation groups in differential geometry*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1972 edition.
- [20] Shoshichi Kobayashi and Katsumi Nomizu. *Foundations of differential geometry. Vol. I*. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1996. Reprint of the 1963 original, A Wiley-Interscience Publication.
- [21] Andrew D. Lewis. Affine connections and distributions with applications to nonholonomic mechanics. *Rep. Math. Phys.*, 42(1-2):135–164, 1998. Pacific Institute of Mathematical Sciences Workshop on Nonholonomic Constraints in Dynamics (Calgary, AB, 1997).
- [22] E. M. Patterson and A. G. Walker. Riemann extensions. *Quart. J. Math. Oxford Ser. (2)*, 3:19–28, 1952.
- [23] Peter Petersen. *Riemannian geometry*, volume 171 of *Graduate Texts in Mathematics*. Springer, Cham, third edition, 2016.

- [24] M. D. Staic. From 3-algebras to  $\Delta$ -groups and symmetric cohomology. *J. Algebra*, 322(4):1360–1378, 2009.
- [25] I. Terek. Notes on Killing fields. <https://web.williams.edu/Mathematics/it3/texts/killing.pdf>. Unpublished notes.
- [26] Martin Walker and Roger Penrose. On quadratic first integrals of the geodesic equations for type {22} spacetimes. *Comm. Math. Phys.*, 18:265–274, 1970.
- [27] K. Yano and E. M. Patterson. Vertical and complete lifts from a manifold to its cotangent bundle. *J. Math. Soc. Japan*, 19:91–113, 1967.

UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BARCELONA, SPAIN;  
FACULTY OF NUCLEAR SCIENCES AND PHYSICAL ENGINEERING, CZECH TECHNICAL  
UNIVERSITY IN PRAGUE, 115 19 PRAGUE 1, CZECH REPUBLIC  
*Email address:* filip.moucka@autonoma.cat; mouckfil@cvut.cz

UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BARCELONA, SPAIN  
*Email address:* roberto.rubio@uab.es