

# Constrained mean-field control with singular control: Existence, stochastic maximum principle and constrained FBSDE

Lijun Bo <sup>\*</sup>      Jingfei Wang <sup>†</sup>      Xiang Yu <sup>‡</sup>

## Abstract

This paper studies some mean-field control (MFC) problems with singular control under general dynamic state-control-law constraints. We first propose a customized relaxed control formulation to cope with the dynamic mixed constraints and establish the existence of an optimal control using some compactification arguments in the proper canonical spaces to accommodate the singular control. To characterize the optimal pair of regular and singular controls, we treat the controlled McKean-Vlasov process as an infinite-dimensional equality constraint and recast the MFC problem as an optimization problem on canonical spaces with constraints on Banach space, allowing us to derive the stochastic maximum principle (SMP) and a constrained BSDE using a novel Lagrange multipliers method. In addition, we further investigate the uniqueness and the stability result of the solution to the constrained FBSDE associated to the constrained MFC problem with singular control.

**Keywords:** Mean-field control with singular control, dynamic state-control-law constraints, relaxed control, compactification, stochastic maximum principle, Lagrange multipliers method, constrained FBSDE

**2020 MSC:** 60H30, 49N80, 46N10, 60H10

## 1 Introduction

To overcome the high complexity in large stochastic system caused by the mutual influence among agents, mean-field game (MFG) problems (see [22] and [26]) has been proposed and popularized in academic studies, which is to find a Nash equilibrium in a competitive game formulation where each agent's impact on the aggregate distribution is negligible. Recently, MFC problems has also gained considerable attention, which concerns the cooperative population where all agents share the same goal to attain the social optimum. The present paper targets at the MFC problem, which is often interpreted as an optimal control of McKean-Vlasov dynamics by a social planner. Some pioneer studies on this topic, just to name a few, can be found in [2], [7], [10], [1], [5] for the method of stochastic maximum principle (SMP); [25] and [15] for the limit theories; [27] and [30] for the dynamic programming principle (DPP) and the HJB equation on the Wasserstein space of probability measures; [4], [31] and [32] for the study of viscosity solution and comparison principle of the HJB equation. However, it is notable that most existing studies on MFC are limited to regular controls.

---

<sup>\*</sup>Email: lijunbo@ustc.edu.cn, School of Mathematics and Statistics, Xidian University, Xi'an, 710126, China.

<sup>†</sup>Email: wjf2104296@mail.ustc.edu.cn, School of Mathematical Sciences, University of Science and Technology of China, Hefei, 230026, China.

<sup>‡</sup>Email: xiang.yu@polyu.edu.hk, Department of Applied Mathematics, The Hong Kong Polytechnic University, Kowloon, Hong Kong, China.

Singular control, also called bounded variation stochastic control, allows the displacement of the states to be discontinuous. In single agent's models, optimal singular control problems have been extensively investigated. For instance, we refer to [29] for early discussions on smooth fit principle; [20] for the existence of an optimal singular control based on the relaxed control formulation; [21] for the proof of DPP to account for singular controls; [3] for the stochastic maximum principle tailor-made for singular controls; [19] for some general theories on smooth fit principle. In the competitive game framework, there are a handful of studies on singular controls, see [11] for the n-player game with singular control as well as [17], [8] and [9] for some MFG problems with singular control. However, theoretical studies for MFC problems with singular control are underexplored in the literature, and only a recent paper [14] studies the derivation of DPP and the quasi-variational inequality using two-layer parameterisations of stochastic processes.

The goal of the present paper is to contribute to the study of MFC problems by featuring both the singular control and the joint dynamic state-control-law constraints. Motivated by practical risk regulations, liquidity requirements, or feasibility restraints, different types of constraints have been considered in stochastic control problems, but predominantly in single agent's model. Recently, in mean-field models with only regular controls, there are some emerging research to incorporate state and/or law constraints, see [18] for the level-set approach to tackle the law constraint in MFC problems; [12] for the analysis of mean-field limits under state constraints; [13] for the formulation of controlled Fokker-Planck equation under state constraints; [6] for the optimization formulation of the constrained MFC problem and the derivation of the stochastic maximum principle using the generalized Lagrange multipliers method.

It is evident that the allowance of singular control in the mean-field model will entangle with the dynamic mixed constraints, which renders some existing methods for constrained MFC with regular controls not applicable in our framework. For example, it is difficult to generalize the level set approach in [18] to account for both control constraints and the singular control. Due to the possible discontinuity of the state process and its law under the singular control, characterizing the associated Fokker-Planck equation is still an open problem in the literature, and hence the approach in [13] is also not suitable to handle the singular control. On the other hand, some conventional methods that have been employed in unconstrained MFG and MFC problems with singular control such as DPP and compactification arguments also need to be carefully and substantially modified in the setting when pathwise constraints are imposed jointly on the state process and its distribution flow.

The present paper fills the gap in the literature by advancing some theoretical foundations for constrained MFC problems with singular control. The first contribution of the paper is to address the existence of an optimal pair of regular and singular controls by developing the customized relaxed control formulation and martingale problem to embed the dynamic mixed constraints. That is, we embed the dynamic mixed constraints in item (iv) of the admissible control rules in Definition 3.1. We then present in Lemma 3.5 a martingale problem characterization associated to the admissible control rules and establish in Lemma 3.7 the closedness property of the set of probability measures fulfilling the constraints. By developing some technical compactification arguments to accommodate the singular control and combining the previous lemmas using the embedded constraints, we are able to show that the set of approximately optimal control rules is compact (see Lemma 3.8) and hence the set of optimal control rules is compact and convex that yields the existence of an optimal relaxed control (see Theorem 3.9). By some standard arguments, we also obtain the existence of an optimal strict control for the constrained MFC problem with singular control.

Our second contribution is to further characterize the optimal controls by establishing the SMP for the constrained MFC problem with singular control and derive the adjoint constrained BSDE. However, some conventional techniques such as the first order variation or the spike variation of controls may fail in our setting as the mixed constraints might be violated. Inspired

by our recent study [6] for constrained MFC with regular controls, we propose to regard the controlled McKean-Vlasov dynamics with singular control as an infinite-dimensional equality constraint and recast the MFC problem as an optimization problem on some proper canonical spaces with general constraints on Banach space (see the formulation (15)-(16)). Similar to [6], we are allowed to derive the SMP by using the generalized Fritz-John (FJ) optimality condition and the Lagrange multipliers method. To be more precise, when the forward McKean-Vlasov SDE is taken as an equality constraint, the FJ condition for the control variable gives the SMP, and the Lagrange multipliers in the space of Itô processes induced by the FJ condition on the state variable yields the adjoint processes as the solution to the constrained BSDE, see Theorem 4.4.

As the third contribution of the present paper, we also analyze the non-standard constrained FBSDE in (26)-(27) associated to our constrained MFC problem with singular control. It is worth noting that the Lagrange multiplier  $\eta$  in Theorem 4.4 is treated as part of the solution to the constrained BSDE problem. We then verify the constraints qualification and obtain the KKT optimality condition in Lemma 5.1. As a result, we further establish the uniqueness of the solution to the nonstandard constrained FBSDE in (26)-(27), see Theorem 5.3. Furthermore, under some enhanced assumptions, we also prove the stability result of the solution in the sense of (31) in Theorem 5.5.

The rest of the paper is organized as follows. Section 2 introduces the constrained MFC problem with singular control and some standing assumptions imposed on the underlying model. In Section 3, we introduce the relaxed control formulation and establish the existence of optimal control to the constrained MFC problem by applying some compactification arguments. Section 4 formulates the equivalent optimization problem with constraints in Banach space, for which we develop the SMP and a joint constrained BSDE using Lagrange multipliers method. Finally, the uniqueness and the stability result of the solution to the constrained FBSDEs associated to the constrained MFC problem with singular control are discussed in Section 5.

**Notations.** For the reader's convenience, we list below some notations that will be frequently used throughout the paper:

|   |  |
|---|--|
| $(E^*, \ \cdot\ _{E^*})$                      | The dual space of a Banach space $(E, \ \cdot\ _E)$  |
| $\langle \cdot, \cdot \rangle_{E, E^*}$       | The pairing between Banach space $(E, \ \cdot\ _E)$ and its dual   |
| $a \geq_K 0$ ( $a >_K 0$ )                    | $a \in K$ ( $a \in K^\circ$ ) with $K$ being a closed (linear) cone  |
| $Df$  | The Fréchet derivative of $f : E \mapsto \mathbb{R}$   |
| $\nabla_x f$                                  | Gradient of $f : \mathbb{R}^n \mapsto \mathbb{R}$  |
| $D_X f$                                       | The partial Fréchet derivative of $f$ with respect to $X$  |
| $D_\alpha f$                                  | The partial Fréchet derivative of $f$ with respect to $\alpha$   |
| $L^p((A, \mathcal{B}(A), \lambda_A); E)$      | Set of $L^p$ -integrable $E$ -valued mapping defined on measure space $(A, \mathcal{B}(A), \lambda_A)$ . Write $L^p(A; E)$ for short                             |
| $L^p((A, \mathcal{B}(A), \lambda_A); E)/\sim$ | The quotient set of $L^p(A; E)$ w.r.t. equivalence relation $\sim$ : $f \sim g$ iff $f = g$ except for a $\lambda_A$ -null set. Write $L^p(A; E)/\sim$ for short |
| $\mathbb{E}$ ( $\mathbb{E}'$ )                | Expectation operator under probability measure $\mathbb{P}$ ( $\mathbb{P}'$ )  |
| $\mathcal{P}_p(E)$                            | Set of probability measures on $E$ with finite $p$ -order moments  |
| $\mathcal{M}[0, T]$ ( $\mathcal{M}^+[0, T]$ ) | Set of signed (non-negative) Radon measures on $[0, T]$  |
| $\text{BV}^k[0, T]$                           | Set of functions $f : [0, T] \rightarrow \mathbb{R}^k$ with bounded variation  |
| $D^k[0, T]$                                   | Set of functions $f : [0, T] \rightarrow \mathbb{R}^k$ that are càdlàg.  |

## 2 Problem Formulation

### 2.1 MFC with singular control and mixed constraints

Let  $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$  be a filtered probability space with the filtration  $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$  satisfying the usual conditions. For  $r, l, k \in \mathbb{N}$  and  $p > 2$ , let  $W = (W_t)_{t \in [0, T]}$  be a standard  $r$ -dimensional  $(\mathbb{P}, \mathbb{F})$ -Brownian motion and  $U \subset \mathbb{R}^l$  be a closed and convex (regular) set as the policy space. Define the space of regular controls  $\mathcal{U}[0, T]$  as the set of  $\mathbb{F}$ -progressively measurable processes such that  $\mathbb{E}[\int_0^T |\alpha_t|^p dt] < \infty$ . For the space of singular controls, define  $\mathcal{A}^k[0, T]$  as the set of functions  $a : [0, T] \mapsto \mathbb{R}^k$  such that  $a = (a_t^i)_{t \in [0, T]}$  is càdlàg and non-decreasing with  $a_0^i = 0$  for  $i = 1, \dots, k$ . Let  $\mathcal{A}^k[0, T]$  be the set of  $\mathbb{F}$ -progressively measurable processes  $\zeta = (\zeta_t)_{t \in [0, T]}$  with paths in  $\mathcal{A}^k[0, T]$  and  $\mathbb{E}[|\zeta_T|^p] < \infty$ .

Let  $(b, \sigma) : [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l \mapsto \mathbb{R}^n \times \mathbb{R}^{n \times r}$  be measurable mappings and  $G : [0, T] \mapsto \mathbb{R}^{n \times k}$  be a continuous function. For  $(\alpha, \zeta) = (\alpha_t, \zeta_t)_{t \in [0, T]} \in \mathcal{U}[0, T] \times \mathcal{A}^k[0, T]$ , we consider the controlled McKean-Vlasov SDE that, for  $t \in (0, T]$ ,

$$dX_t^{\alpha, \zeta} = b(t, X_t^{\alpha, \zeta}, \mathcal{L}(X_t^{\alpha, \zeta}), \alpha_t)dt + \sigma(t, X_t^{\alpha, \zeta}, \mathcal{L}(X_t^{\alpha, \zeta}), \alpha_t)dW_t + G(t)d\zeta_t, \quad (1)$$

and  $X_0^{\alpha, \zeta} = \xi \in L^p((\Omega, \mathcal{F}_0, \mathbb{P}); \mathbb{R}^n)$  with the law of  $\xi$  under  $\mathbb{P}$  being  $\mathcal{L}(\xi) = \nu \in \mathcal{P}_p(\mathbb{R}^n)$ .

For the above controlled McKean-Vlasov dynamics, we shall impose some dynamic state-control-law constraints. To this end, let  $d \in \mathbb{N}$  be the number of inequality constraints. For  $i \in I := \{1, \dots, d\}$ , let us consider  $\phi^i : [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l \mapsto \mathbb{R}$  with  $i \in I$  such that the dynamic state-control-law inequality constraints are described by

$$\phi^i(t, X_t^{\alpha, \zeta}, \mathcal{L}(X_t^{\alpha, \zeta}), \alpha_t) \geq 0, \quad dt \times d\mathbb{P}\text{-a.s.}, \quad (2)$$

where  $X^{\alpha, \zeta} = (X_t^{\alpha, \zeta})_{t \in [0, T]}$  obeys the McKean-Vlasov SDE (1). We denote  $\mathcal{U}_{\text{ad}}[0, T]$  as the admissible control set of processes  $(\alpha, \zeta) \in \mathcal{U}[0, T] \times \mathcal{A}^k[0, T]$  such that there exists a càdlàg  $\mathbb{F}$ -adapted process  $\mathbb{R}^n$ -valued  $X^{\alpha, \zeta} = (X_t^{\alpha, \zeta})_{t \in [0, T]}$  satisfying (1) and (2).

The goal of the MFC problem is to minimize the following cost functional that, for  $(\alpha, \zeta) = (\alpha_t, \zeta_t)_{t \in [0, T]} \in \mathcal{U}_{\text{ad}}[0, T] \neq \emptyset$ ,

$$\inf_{(\alpha, \zeta) \in \mathcal{U}_{\text{ad}}[0, T]} J(\alpha, \zeta) := \inf_{(\alpha, \zeta) \in \mathcal{U}_{\text{ad}}[0, T]} \mathbb{E} \left[ \int_0^T f(t, X_t^{\alpha, \zeta}, \mathcal{L}(X_t^{\alpha, \zeta}), \alpha_t)dt + \int_0^T c(t)d\zeta_t \right], \quad (3)$$

where  $f : [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l \mapsto \mathbb{R}$  and  $c : [0, T] \mapsto \mathbb{R}^k$  are respectively the running cost function and the cost function associated with singular controls.

### 2.2 Standing assumptions

Let us first unpack some standing assumptions imposed on the model coefficients, which will be used in different sections.

**Assumption 1.** (A1) *The mappings  $b, \sigma, f$  and  $\phi^i$  for  $i \in I$  are Borel measurable,  $b, \sigma$  are continuous and  $f$  is lower semi-continuous (l.s.c.) in  $(x, \mu, u) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l$ .*

(A2) *For  $i \in I$ ,  $\phi^i$  is upper semi-continuous (u.s.c.) in  $(x, \mu, u) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l$  and there exists a constant  $M > 0$  such that, for all  $i \in I$  and  $(x, u) \in \mathbb{R}^n \times \mathbb{R}^l$ ,*

$$|\phi^i(t, x, \mu, u)| \leq M(1 + |x| + M_2(\mu) + |u|)$$

*with  $M_2(\mu) := \int_{\mathbb{R}^n} |x|^2 \mu(dx)$ . Moreover,  $\phi^i$  is uniformly u.s.c. in  $\mu \in \mathcal{P}_2(\mathbb{R}^n)$  with respect to  $(t, x, u) \in [0, T] \times \mathbb{R}^n \times \mathbb{R}^l$ .*

(A3) The mappings  $b(t, x, \mu, u)$  and  $\sigma(t, x, \mu, u)$  are uniformly Lipschitz continuous in  $(x, \mu) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$  in the sense that, there exists a constant  $M > 0$  independent of  $u \in \mathbb{R}^l$  such that, for all  $(x, \mu), (x', \mu') \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$  and  $(t, u) \in [0, T] \times \mathbb{R}^l$ ,

$$|b(t, x', \mu', u) - b(t, x, \mu, u)| + |\sigma(t, x', \mu', u) - \sigma(t, x, \mu, u)| \leq M(|x - x'| + \mathcal{W}_{2, \mathbb{R}^n}(\mu, \mu')),$$

where  $\mathcal{W}_{2, \mathbb{R}^n}$  is a 2-Wasserstein metric on  $\mathcal{P}_2(\mathbb{R}^n)$ .

(A4) There exists a constant  $M > 0$  such that, for all  $(t, x, \mu, u) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l$ ,

$$|b(t, x, \mu, u)| + |\sigma(t, x, \mu, u)| \leq M(1 + |x| + M_2(\mu) + |u|).$$

(A5)  $G : [0, T] \mapsto \mathbb{R}^{n \times k}$  and  $c : [0, T] \mapsto [0, \infty)^k$  are deterministic continuous functions. Moreover,  $c(t) = (c^1(t), \dots, c^k(t))$  satisfies  $c^i(t) \geq c_0 > 0$  for all  $t \in [0, T]$ .

(A6) There exist  $M, M_1 > 0$  such that, for all  $(t, x, \mu, u) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l$ ,

$$-M(1 + |x|^p + M_2(\mu)) + M_1|u|^p \leq f(t, x, \mu, u) \leq M(1 + |x|^p + M_2(\mu) + |u|^p).$$

**Assumption 2.** (B1) For  $i \in I$ , the mappings  $b, \sigma, f, g$  and  $\phi^i$  are Borel measurable and continuously differentiable w.r.t.  $x \in \mathbb{R}^n$ . Moreover,  $\phi^i$  is jointly continuous in  $(t, x, \mu) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$ .

(B2) There exists a constant  $M > 0$  such that, for all  $(t, x, \mu, u) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l$ ,

$$\sum_{i \in I} (|\nabla_x \phi^i(t, x, \mu, u)| + |\nabla_u \phi^i(t, x, \mu, u)|) \leq M.$$

(B3) The condition (A3) in [Assumption 1](#) holds.

(B4) There exists a constant  $M > 0$  such that, for all  $(t, x, \mu, u) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l$ ,  $|\delta(t, x, \mu, u)| \leq M(1 + |x| + |u| + M_2(\mu))$ , where  $\delta \in \{b, \sigma, \nabla_u b(\cdot), \nabla_u \sigma(\cdot), \nabla_x f(\cdot), \nabla_u f(\cdot)\}$ .

(B5) For  $i \in I$ , the mappings  $b(t, x, \mu, u), \sigma(t, x, \mu, u), f(t, x, \mu, u)$  and  $\phi^i(t, x, \mu, u)$  are continuously L-differentiable w.r.t.  $\mu \in \mathcal{P}_2(\mathbb{R}^n)$ . Moreover, there exists a constant  $M > 0$  such that, for all  $(t, x, \mu, u) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l$ ,

$$\left( \int_{\mathbb{R}^n \times \mathbb{R}^l} |\partial_\mu f(t, x, \mu, u)(x', u')|^2 \mu(dx', du') \right)^{\frac{1}{2}} \leq M(1 + |x| + |u| + M_2(\mu)),$$

and moreover

$$\sum_{i \in I} \left( \int_{\mathbb{R}^n \times \mathbb{R}^l} |\partial_\mu \phi^i(t, x, \mu, u)(x', u')|^2 \mu(dx', du') \right)^{\frac{1}{2}} \leq M.$$

(B6)  $G : [0, T] \mapsto \mathbb{R}^{n \times k}$  and  $c : [0, T] \mapsto [0, \infty)^k$  are deterministic continuous functions.

**Assumption 3.** (C1) The constrained function  $\phi$  is independent of the control variable and has the linear form of

$$\phi(t, x, \mu) = A(t)x + B(t) \int_{\mathbb{R}^n} x \mu(dx) + C(t), \quad \forall (t, x, \mu) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n),$$

where  $A = (A(t))_{t \in [0, T]}$ ,  $B = (B(t))_{t \in [0, T]}$  and  $C = (C(t))_{t \in [0, T]} \in C([0, T]; \mathbb{R}^n)$  and it holds that  $|A(t)| > 0$  and  $|A(t) + B(t)| > 0$  for all  $t \in [0, T]$ .

(C2) The Hamiltonian  $H$  defined by  $H(t, x, \mu, u, y, z) := b(t, x, \mu, u)^\top y + \text{tr}(\sigma(t, x, \mu, u)^\top z) + f(t, x, \mu, u)$  is strictly convex in  $u \in U$ . There exists a measurable mapping  $\hat{u} : [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^n \times \mathbb{R}^{n \times r} \mapsto U \subset \mathbb{R}^l$  with at most quadratic growth in  $(x, \mu, y, z) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^n \times \mathbb{R}^{n \times r}$  such that  $\hat{u}(t, x, \mu, y, z) \in \arg \min_{u \in U} H(t, x, \mu, u, y, z)$  and  $\hat{u}$  is Lipschitz continuous in the sense that there exists a constant  $M > 0$  such that, for all  $(x_i, \mu_i, y_i, z_i) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^n \times \mathbb{R}^{n \times r}$  with  $i = 1, 2$ ,

$$\begin{aligned} & |\hat{u}(t, x_1, \mu_1, y_1, z_1) - \hat{u}(t, x_2, \mu_2, y_2, z_2)| \\ & \leq M(|x_1 - x_2| + \mathcal{W}_{2, \mathbb{R}^n}(\mu_1, \mu_2) + |y_1 - y_2| + |z_1 - z_2|). \end{aligned}$$

We then set  $\hat{b}(t, x, \mu, y, z) := b(t, x, \mu, \hat{u}(t, x, \mu, y, z))$ ,  $\hat{\sigma}(t, x, \mu, y, z) := \sigma(t, x, \mu, \hat{u}(t, x, \mu, y, z))$  and  $\hat{f}(t, x, \mu, y, z) := f(t, x, \mu, \hat{u}(t, x, \mu, y, z))$ .

(C3) Let us define

$$F(t, x, \mu, u, y, z) := \nabla_x H(t, x, \mu, u, y, z) + \int_{\mathbb{R}^n} \partial_\mu H(t, x', \mu, u, y, z)(x) \mu(dx'),$$

and set

$$\begin{aligned} \hat{F}(t, x, \mu, y, z) &:= F(t, x, \mu, \hat{u}(t, x, \mu, y, z), y, z), \\ \hat{\Phi}(t, x, \mu, y, z) &:= (-\hat{F}, \hat{b}, \hat{\sigma})(t, x, \mu, y, z). \end{aligned}$$

The following monotone condition holds that

$$\begin{aligned} & (\hat{\Phi}(t, x_1, \mu, y_1, z_1) - \hat{\Phi}(t, x_2, \mu, y_2, z_2))(x_1 - x_2, y_1 - y_2, z_1 - z_2) \\ & + \beta(|x_1 - x_2|^2 + |y_1 - y_2|^2 + |z_1 - z_2|^2) \leq 0, \end{aligned}$$

where  $\beta > \frac{M}{2}$  with  $M$  being given in (A3) and the inner product is read as

$$\begin{aligned} & (\hat{\Phi}(t, x_1, \mu, y_1, z_1) - \hat{\Phi}(t, x_2, \mu, y_2, z_2))(x_1 - x_2, y_1 - y_2, z_1 - z_2) \\ & = -(\hat{F}(t, x_1, \mu, y_1, z_1) - \hat{F}(t, x_2, \mu, y_2, z_2))^\top (x_1 - x_2) \\ & + (\hat{b}(t, x_1, \mu, y_1, z_1) - \hat{b}(t, x_2, \mu, y_2, z_2))^\top (y_1 - y_2) \\ & + \text{tr} \left( (\hat{\sigma}(t, x_1, \mu, y_1, z_1) - \hat{\sigma}(t, x_2, \mu, y_2, z_2))^\top (z_1 - z_2) \right). \end{aligned}$$


(C4)  $\hat{b}$ ,  $\hat{\sigma}$ ,  $\hat{F}$  satisfy the Lipschitz condition in the sense that, there exists a constant  $M > 0$  such that, for all  $(x_i, \mu_i, y_i, z_i) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^n \times \mathbb{R}^{n \times r}$  with  $i = 1, 2$ ,

$$|\delta(t, x_1, \mu_1, y_1, z_1) - \delta(t, x_2, \mu_2, y_2, z_2)| \leq M(|x_1 - x_2| + \mathcal{W}_{2, \mathbb{R}^n}(\mu_1, \mu_2) + |y_1 - y_2| + |z_1 - z_2|),$$

for  $\delta \in \{\hat{b}, \hat{\sigma}, \hat{f}, \nabla_x b, \partial_\mu b(\cdot)(x), \nabla_x \sigma, \partial_\mu \sigma(\cdot)(x), \nabla_x f, \partial_\mu f(\cdot)(x)\}$  and  $x \in \mathbb{R}^n$ .

Note that [Assumption 2](#) strengthens [Assumption 1](#) by requiring the differentiability of the coefficients and imposing growth conditions on their derivatives, while it is weaker than [Assumption 1](#) by relaxing the growth condition on the running cost function  $f$  and omitting the requirement for a uniform positive constant  $c_0$  of the cost function  $c$ . [Assumption 3](#) introduces the monotonicity to the coefficients and guarantees the existence of a Lipschitz continuous minimizer for the Hamiltonian. The following flowchart illustrates how the above three assumptions are applied throughout the paper.





### 3 Existence of Optimal Controls

#### 3.1 Relaxed control formulation

To confirm the existence of an optimal pair of regular and singular controls to the problem (3), we first reformulate the problem in a relaxed control framework such that the compactification method can be utilized. To this end, let us introduce the following basic spaces:

- The space  $D^n[0, T]$  is endowed with the Skorokhod metric  $d_D$  and the corresponding Borel  $\sigma$ -algebra denoted by  $\mathcal{F}^D$ . Let  $\mathcal{F}_t^D$  be the Borel  $\sigma$ -algebra up to time  $t$  (c.f. Haussmann [20]) for  $t \in [0, T]$ .
- The space  $\mathcal{Q}$  of relaxed controls is defined as the set of measures  $q$  in  $[0, T] \times U$  with the first marginal equal to the Lebesgue measure and  $\int_{[0, T] \times U} |u|^p q(dt, du) < \infty$ . We endow the space  $\mathcal{Q}$  with the 2-Wasserstein metric on  $\mathcal{P}_2([0, T] \times U)$  given by  $d_{\mathcal{Q}}(q^1, q^2) = \mathcal{W}_{2, [0, T] \times U} \left( \frac{q^1}{T}, \frac{q^2}{T} \right)$ , where the metric on  $[0, T] \times U$  is given by  $((t_1, u_1), (t_2, u_2)) \mapsto |t_2 - t_1| + |u_2 - u_1|$ . Note that each  $q \in \mathcal{Q}$  can be identified with a measurable function  $[0, T] \ni t \mapsto q_t \in \mathcal{P}_2(U)$ , defined uniquely up to a.s. by  $q(dt, du) = q_t(du)dt$ . In the sequel, we will always refer to the measurable mapping  $q = (q_t)_{t \in [0, T]}$  to a relaxed control in  $\mathcal{Q}$ . Let  $\mathcal{F}^{\mathcal{Q}}$  be the Borel  $\sigma$ -algebra of  $\mathcal{Q}$  and  $\mathcal{F}_t^{\mathcal{Q}}$  be the  $\sigma$ -algebra generated by the maps  $q \mapsto q([0, s] \times V)$  with  $s \in [0, t]$  and Borel measurable  $V \subset U$ .
- The space  $A^k[0, T]$  is a subset of  $\mathcal{V} \subset \text{BV}^k[0, T]$ , where the set  $\mathcal{V} \subset \text{BV}^k[0, T]$  is the collection of mappings  $a : [0, T] \rightarrow \mathbb{R}^k$  such that  $a$  is càdlàg and  $a(0) = 0$ . As in [20], there is a natural one-to-one correspondence between  $\mathcal{V}$  and  $(\mathcal{M}[0, T])^k$ , that is, for  $a = (a^1, \dots, a^k) \in \mathcal{V}$  and  $i = 1, \dots, k$ , define  $\nu_i((s, t]) = a^i(t) - a^i(s)$  for  $0 \leq s \leq t \leq T$ . Then,  $\mathcal{V}$  is the dual space of  $(C[0, T]^*)^k$ , and hence we endow  $\mathcal{V}$  with the corresponding weak\* topology (which is also equivalent to the Fortet-Mourier norm on  $(\mathcal{M}[0, T])^k$ , c.f. [5]). Thus,  $A^k[0, T]$  is a closed subset of  $(\mathcal{M}[0, T])^k$  and can also be viewed as  $(\mathcal{M}^+[0, T])^k$ . Let  $\mathcal{F}^A$  be the Borel  $\sigma$ -algebra and  $\mathcal{F}_t^A$  be the Borel  $\sigma$ -algebra defined up to time  $t \in [0, T]$  (cf. [20]).

Now, we introduce the sample space considered in the paper, which is given by

$$\Omega := \widehat{\Omega} \times A^k[0, T] = D^n[0, T] \times \mathcal{Q} \times A^k[0, T]. \quad (4)$$

Equip  $\widehat{\Omega}$  and  $\Omega$  with the respective product metrics  $d_{\widehat{\Omega}}(\widehat{\omega}^1, \widehat{\omega}^2) = d_D(\psi^1, \psi^2) + d_{\mathcal{Q}}(q^1, q^2)$  and  $d_{\Omega}(\omega^1, \omega^2) = d_{\widehat{\Omega}}(\widehat{\omega}^1, \widehat{\omega}^2) + \sum_{i=1}^k d_{\text{FM}}(\nu_i^1, \nu_i^2)$  for  $\omega^i = (\widehat{\omega}^i, \zeta^i) = (\psi^i, q^i, \zeta^i) \in \Omega$  with  $i = 1, 2$ . Here,  $\nu^i$  is the correspondence of  $\zeta^i$  in  $\mathcal{V}$  and  $d_{\text{FM}}$  is the Fortet-Mourier metric on  $\mathcal{M}[0, T]$ . Consequently, both  $(\widehat{\Omega}, d_{\widehat{\Omega}})$  and  $(\Omega, d_{\Omega})$  are Polish spaces.

Set  $\mathcal{F} = \mathcal{F}^D \otimes \mathcal{F}^{\mathcal{Q}} \otimes \mathcal{F}^A$  as the product  $\sigma$ -algebra and define the filtration  $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$  by  $\mathcal{F}_t = \mathcal{F}_t^D \otimes \mathcal{F}_t^{\mathcal{Q}} \otimes \mathcal{F}_t^A$  for  $t \in [0, T]$ . For any  $\omega = (\psi, q, \zeta) \in \Omega$ , Let  $X_t(\omega) = \psi_t$ ,  $q_t(\omega) = q_t$ , and  $\zeta_t(\omega) = \zeta_t$  for  $t \in [0, T]$  be the coordinate process. Denote by  $\mathcal{P}_2(\Omega; \widehat{\Omega})$  the set of probability measures  $R$  on  $(\Omega, \mathcal{F})$  such that  $R|_{\widehat{\Omega}} \in \mathcal{P}_2(\widehat{\Omega})$ . Introduce the corresponding metric on  $\mathcal{P}_2(\Omega; \widehat{\Omega})$  by, for all  $R_1, R_2 \in \mathcal{P}_2(\widehat{\Omega})$ ,

$$d(R_1, R_2) = \mathcal{W}_{2, \widehat{\Omega}}(R_1|_{\widehat{\Omega}}, R_2|_{\widehat{\Omega}}) + d_{\text{FM}, \Omega}(R_1, R_2),$$

where  $d_{\text{FM}, \Omega}$  is the Fortet-Mourier metric on  $\mathcal{P}(\Omega)$ . As a result,  $R_n \rightarrow R$  in  $\mathcal{P}_2(\Omega; \widehat{\Omega})$  if and only if  $R_n|_{\widehat{\Omega}} \rightarrow R|_{\widehat{\Omega}}$  in  $\mathcal{P}_2(\widehat{\Omega})$  and  $R_n$  weakly converges to  $R$  as  $n \rightarrow \infty$ .

Next, we give the definition of relaxed controls. In particular, one important task is to guarantee that the dynamic state-control-law constraints are fulfilled when applying the compactification arguments. For this purpose, we propose to embed the constraints into the definition of admissible control rules in the following sense. Recall that  $\nu \in \mathcal{P}_p(\mathbb{R}^n)$  is the law of initial state in SDE (1).

In the sequel, we will always consider the weak formulation, i.e., we will take  $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$  introduced in last section to be the canonical space.

**Definition 3.1.** *An admissible control rule is a probability measure  $R \in \mathcal{P}_2(\Omega; \widehat{\Omega})$  on  $(\Omega, \mathcal{F})$  such that*

- (i)  $R \circ X_0^{-1} = \nu$ ;
- (ii)  $\mathbb{E}^R \left[ \int_0^T \int_U |u|^p q_t(du) dt + |\zeta_T|^p \right] < \infty$ ;
- (iii) for all  $g \in C_b^2(\mathbb{R}^n)$ ,  $M^R g = (M_t^R g)_{t \in [0, T]}$  is an  $(R, \mathbb{F})$ -martingale, where

$$\begin{aligned} M_t^R g &= g(X_t) - \int_0^t \int_U \mathbb{G}g(s, X_s, R^X(s), u) q_s(du) ds - \int_0^t \nabla_x g(X_{s-}) G(s) d\zeta_s \\ &\quad - \sum_{0 < s \leq t} (g(X_s) - g(X_{s-}) - \nabla_x g(X_{s-}) \Delta X_s) \end{aligned}$$

with  $R^X(s) = R \circ X_s^{-1}$ ,  $\Delta X_s = X_s - X_{s-}$  and

$$\mathbb{G}g(t, x, \mu, u) = b(t, x, \mu, u) \nabla_x g(x) + \frac{1}{2} \text{tr} \left( \sigma \sigma^\top(t, x, \mu, u) \nabla_{xx}^2 g(x) \right).$$

- (iv) for  $i \in I$ ,  $m \times R(A_i^R) = T$  with the admissible set  $A_i^R$  defined by

$$A_i^R = \left\{ (t, \omega) \in [0, T] \times \Omega; \int_U \phi^i(t, X_t(\omega), R^X(t), u) q_t(\omega, du) \geq 0 \right\},$$

where  $m \times R$  denotes the product measure of Lebesgue measure  $m$  and  $R$ .



### 3.2 Existence of optimal relaxed controls

For  $\nu \in \mathcal{P}_p(\mathbb{R}^n)$ , let  $\mathcal{R}_{\text{ad}}(\nu)$  be the set of admissible rules  $R$  in Definition 3.1 in the previous subsection. For  $R \in \mathcal{R}_{\text{ad}}(\nu)$ , we have the following martingale measure characterization:

**Lemma 3.2.** *We have  $R \in \mathcal{R}_{\text{ad}}(\nu)$  if and only if there is a filtered probability space  $(\Omega', \mathcal{F}', \mathbb{F}' = (\mathcal{F}'_t)_{t \in [0, T]}, P')$  supporting an  $n$ -dimensional  $\mathbb{F}'$ -adapted process  $X = (X_t)_{t \in [0, T]}$ ,  $r$ -dimensional orthogonal  $\mathbb{F}'$ -martingale measures  $N = (N^1, \dots, N^r)$  on  $U \times [0, T]$ , each with  $\mathcal{F}'_t$ -progressively measurable intensity  $q_t(du)dt$  and a  $k$ -dimensional  $\mathbb{F}'$ -progressively measurable nondecreasing process  $\zeta = (\zeta_t)_{t \in [0, T]}$  such that  $R = P' \circ (X, q, \zeta)^{-1}$ , and the following conditions hold:*

- (i)  $P' \circ X_0^{-1} = \nu$ ;
- (ii)  $\mathbb{E}^{P'} \left[ \int_0^T \int_U |u|^p q_t(du)dt + |\zeta_T|^p \right] < \infty$ ;
- (iii) the state dynamics obeys that,  $P'$ -a.s.,

$$dX_t = \int_U b(t, X_t, \mathcal{L}^{P'}(X_t), u) q_t(du)dt + \int_U \sigma(t, X_t, \mathcal{L}^{P'}(X_t), u) N(du, dt) + G(t) d\zeta_t,$$

where  $\mathcal{L}^{P'}(X_t)$  denotes the law of  $X_t$  under  $P'$ .

- (iv) the constraints hold that, for  $m \times P'$ -a.s.,  $\int_U \phi^i(t, X_t, \mathcal{L}^{P'}(X_t), u) q_t(du) \geq 0$  for  $i \in I$ .

Moreover, there exists a constant  $C > 0$  depending on  $M, \nu, p, T$  such that  $\mathbb{E}^{P'}[\sup_{t \in [0, T]} |X_t|^p] \leq C \{1 + \mathbb{E}[|\xi|^p]\}$  with  $M$  stated in Assumption 1.

*Proof.* The first assertion follows from El Karoui [16], and the second assertion can be easily proved by basic moment estimations under Assumption 1.  $\square$

Let us define the cost functional on  $\mathcal{R}_{\text{ad}}(\nu)$  by

$$\Gamma(R) := \mathbb{E}^R \left[ \int_0^T \int_U f(t, X_t, R^X(t), u) q_t(du)dt + \int_0^T c(t) d\zeta_t \right]. \quad (5)$$

Then, it follows from Assumption 1 that  $v^* := \inf_{R \in \mathcal{R}_{\text{ad}}(\nu)} \Gamma(R) \in \mathbb{R}$ . Consequently, the optimal control rule set is given by

$$\mathcal{R}_{\text{opt}}(\nu) = \inf\{R \in \mathcal{R}_{\text{ad}}(\nu); \Gamma(R) = v^*\}. \quad (6)$$

**Remark 3.3.** *For any strict admissible control  $(\alpha, \zeta) \in \mathcal{U}_{\text{ad}}[0, T]$ , if let  $X^{\alpha, \zeta}$  be the controlled state process obeyed by (1) and  $R$  be the joint law of  $(X^{\alpha, \zeta}, \delta_\alpha(du)dt, \zeta)$ , then  $R$  can be easily verified to be an admissible control rule and  $J(\alpha, \zeta) = \Gamma(R)$  with  $J(\alpha, \zeta)$  defined in (3). This yields that*

$$v^* \leq \inf_{(\alpha, \zeta) \in \mathcal{U}_{\text{ad}}[0, T]} J(\alpha, \zeta).$$

To arrive at the reverse inequality in Remark 3.3, for any  $(t, x, \nu) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$ , we define a subset of  $\mathbb{R}^n \times \mathbb{R}^{n \times n} \times \mathbb{R} \times \mathbb{R}^m$  as follows:

$$K(t, x, \nu) := \left\{ (b(t, x, \nu, u), \sigma \sigma^\top(t, x, \nu, u), z, w); z \geq f(t, x, \nu, u), w_i \leq \phi^i(t, x, \nu, u), i \in I, u \in U) \right\}. \quad (7)$$

We make the following assumption on the set  $K(t, x, \nu)$ :

**Assumption 4.** *For any  $(t, x, \nu) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$ , the set  $K(t, x, \nu)$  is convex.*

Then, we have

**Lemma 3.4.** *Let Assumption 4 hold. For any admissible (relaxed) control rule  $R \in \mathcal{R}_{\text{ad}}(\nu)$ , there exists an admissible control  $(\alpha, \zeta) \in \mathcal{U}_{\text{ad}}[0, T]$  such that  $J(\alpha, \zeta) \leq \Gamma(R)$ .*

*Proof.* For any  $\omega \in \Omega$ , define the set  $K(t, \omega) := K(t, X_t(\omega), \mathcal{L}(X_t^{\alpha, \zeta}))$  with  $(t, \omega) \in [0, T] \times \Omega$ . Since  $K(t, \omega)$  is convex and closed, we have

$$\left( \int_U b(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du), \int_U \sigma \sigma^\top(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du), \int_U f(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du), \int_U \phi^i(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du), i \in I \right) \in K(t, \omega),$$

which is equivalent to

$$U(t, \omega) := \left\{ u \in U; \int_U b(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du) = b(t, X_t(\omega), \mathcal{L}^R(X_t), u), \int_U \sigma \sigma^\top(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du) = \sigma \sigma^\top(t, X_t(\omega), \mathcal{L}^R(X_t), u), \int_U f(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du) \geq f(t, X_t(\omega), \mathcal{L}^R(X_t), u), \int_U \phi^i(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du) \leq \phi^i(t, X_t(\omega), \mathcal{L}^R(X_t), u), i \in I \right\} \neq \emptyset.$$

For  $N \geq 1$ , define  $U^N(t, \omega) := \{u \in U(t, \omega); |u| \leq N\}$ . Then, for each  $(t, \omega) \in [0, T] \times \Omega$ ,  $U^N(t, \omega)$  increases to  $U(t, \omega)$ . We also introduce  $B^N = \{(t, \omega) \in [0, T] \times \Omega; U^N(t, \omega) \neq \emptyset\}$ . Since  $U(t, \omega)$  is bounded and closed, consequently, it is also a compact subset of  $U \subset \mathbb{R}^l$ . Then, there exists a measurable selection  $\alpha^N : B^N \rightarrow U$  such that  $\alpha_t^N(\omega) \in U^N(t, \omega)$  for  $(t, \omega) \in [0, T] \times \Omega$  and  $\alpha^N$  is  $\mathcal{B}([0, t]) \otimes \mathcal{F}_t/\mathcal{B}(\mathbb{R}^l)$  measurable due to Leese [28]. Hence, we can define an  $\mathbb{F}$ -progressively measurable process  $\alpha = (\alpha_t)_{t \in [0, T]}$  by  $\alpha_t(\omega) = \sum_{N=1}^{\infty} \alpha_t^N(\omega) \mathbf{1}_{B^N \setminus B^{N-1}}(t, \omega)$ .

Next, we show that  $\alpha = (\alpha_t)_{t \in [0, T]}$  is an admissible (strict) control. Using the growth condition imposed on  $f$  in Assumption 1-(A6), we have from the above construction of  $\alpha$  that

$$\begin{aligned} +\infty &> \mathbb{E}^R \left[ \int_0^T \int_U f(t, X_t, \mathcal{L}(X_t), u) q_t(du) dt \right] \geq \mathbb{E}^R \left[ \int_0^T f(t, X_t, \mathcal{L}(X_t), \alpha_t) dt \right] \\ &\geq - \int_0^T M(1 + \mathbb{E}^R[|X_t|^p] + M_2(R^X(t))) dt + M_1 \mathbb{E}^R \left[ \int_0^T |\alpha_t|^p dt \right]. \end{aligned}$$

This, together with Lemma 3.2, indicates that  $\mathbb{E}^R[\int_0^T |\alpha_t|^p dt] < \infty$ . On the other hand, using the construction of  $\alpha$  again, we deduce that, for every  $g \in C_b^2(\mathbb{R}^n)$  and  $(t, \omega) \in [0, T] \times \Omega$ ,

$$\begin{aligned} g(X_t(\omega)) &- \int_0^t \mathbb{G}g(s, X_s(\omega), \mathcal{L}^R(X_s), \alpha_s(\omega)) ds - \int_0^t \nabla_x g(X_s(\omega)) G(s) d\zeta_s(\omega) \\ &- \sum_{0 < s \leq t} (g(X_s(\omega)) - g(X_{s-}(\omega)) - \nabla_x g(X_{s-}(\omega)) \Delta X_s(\omega)) = M_t^R g(\omega) \end{aligned}$$

is an  $(R, \mathbb{F})$ -martingale as  $R$  is a control rule in Definition 3.1. Moreover, we have, for  $m \times R$ -a.s.  $(t, \omega) \in [0, T] \times \Omega$ ,

$$\phi^i(t, X_t(\omega), \mathcal{L}^R(X_t), \alpha_t(\omega)) \geq \int_U \phi^i(t, X_t(\omega), \mathcal{L}^R(X_t), u) q_t(\omega, du) \geq 0, \quad \forall i \in I.$$

This yields that  $\alpha = (\alpha_t)_{t \in [0, T]}$  is admissible. Furthermore, by the construction of  $\alpha$ , it holds that

$$J(\alpha, \zeta) = \mathbb{E}^R \left[ \int_0^T f(t, X_t, \mathcal{L}^R(X_t), \alpha_t) dt + \int_0^T c(t) d\zeta_t \right]$$

$$\leq \mathbb{E}^R \left[ \int_0^T \int_U f(t, X_t, R^X(t), u) q_t(du) dt + \int_0^T c(t) d\zeta_t \right] = \Gamma(R).$$

Thus, we complete the proof of the lemma.  $\square$

We next study the existence of optimal relaxed controls. Consider the McKean-Vlasov SDE (1) and define the process by  $Y_t := X_t^{\alpha, \zeta} - \int_0^t G(s) d\zeta_s$  for  $t \in [0, T]$ . Then, it follows from the dynamics (1) that

$$dY_t = b(t, X_t^{\alpha, \zeta}, \mathcal{L}(X_t^{\alpha, \zeta}), \alpha_t) dt + \sigma(t, X_t^{\alpha, \zeta}, \mathcal{L}(X_t^{\alpha, \zeta}), \alpha_t) dW_t, \quad Y_0 = \xi.$$

Motivated by this expression, we have the next result whose proof is omitted because the necessary part is obvious and the sufficiency can be shown by a classical martingale argument.

**Lemma 3.5.** *Let  $R \in \mathcal{P}_2(\Omega; \widehat{\Omega})$ . Then,  $R \in \mathcal{R}_{ad}(\nu)$  if and only if there exists an  $\mathbb{F}$ -adapted process  $Y = (Y_t)_{t \in [0, T]}$  such that*

- (i)  $Y$  has continuous trajectories and  $R(X_\cdot = Y_\cdot + \int_0^\cdot G(s) d\zeta_s) = 1$ ;
- (ii)  $R \circ Y_0^{-1} = \nu$ ;
- (iii)  $\mathbb{E}^R \left[ \int_0^T \int_U |u|^p q_t(du) dt + |\zeta_T|^p \right] < \infty$ ;
- (iv) for any  $g \in C_b^2(\mathbb{R}^n)$ ,  $\tilde{M}_t^R g$  is a  $(R, \mathbb{F})$ -martingale, where

$$\tilde{M}_t^R g = g(Y_t) - \int_0^t \int_U \tilde{\mathbb{G}}g(s, X_s, R^X(s), u)(Y_s) q_s(du) ds$$

$$\text{with the operator } \tilde{\mathbb{G}}g(t, x, \mu, u)(y) := b(t, x, \mu, u) \nabla_x g(y) + \frac{1}{2} \text{tr}(\sigma \sigma^\top(t, x, \mu, u) \nabla_{xx}^2 g(y));$$

- (v) for  $i \in I$ ,  $m \times R(A_i^R) = T$ .

Moreover, we have

**Lemma 3.6.** *The objective functional  $\Gamma(R)$  defined by (5) is l.s.c. in  $R \in \mathcal{R}_{ad}(\nu)$ .*

*Proof.* Let us define the mapping  $\Lambda(R, \omega) : \mathcal{P}_2(\Omega; \widehat{\Omega}) \times \Omega \mapsto \mathbb{R}$  by

$$\Lambda(R, \omega) = \int_0^T \int_U f(t, X_t(\omega), R^X(t), u) q_t(\omega, du) dt + \int_0^T c(t) d\zeta_t(\omega).$$

Then, for  $R \in \mathcal{P}_2(\Omega; \widehat{\Omega})$ , we get that  $\Lambda(R, \cdot)$  is l.s.c. by virtue of Lemma 3.5 in Hausmann [20]. Furthermore,  $\Lambda(R, \omega)$  is uniformly l.s.c. in  $R \in \mathcal{P}_2(\Omega; \widehat{\Omega})$  with respect to  $\omega \in \Omega$  as  $f$  is uniformly l.s.c. in  $\mu \in \mathcal{P}_2(\mathbb{R}^n)$  with respect to  $(t, x, u) \in [0, T] \times \mathbb{R}^n \times \mathbb{R}^l$ . Thus, if  $R_\ell \rightarrow R$  in  $\mathcal{P}_2(\Omega; \widehat{\Omega})$  as  $\ell \rightarrow \infty$  with  $R_\ell, R \in \mathcal{R}_{ad}(\nu)$ , then  $R_\ell \rightarrow R$  in  $\mathcal{P}_2(\Omega; \widehat{\Omega})$  as  $\ell \rightarrow \infty$ , and hence  $R_\ell \rightarrow R$  as  $\ell \rightarrow \infty$  with  $R_\ell, R \in \mathcal{R}_{ad}(\nu)$ . It then follows that

$$\begin{aligned} \liminf_{\ell \rightarrow \infty} \Gamma(R_\ell) &= \liminf_{\ell \rightarrow \infty} \mathbb{E}^{R_\ell}[\Lambda(R_\ell, \omega)] \\ &\geq \liminf_{\ell \rightarrow \infty} \{ \mathbb{E}^{R_\ell}[\Lambda(R_\ell, \omega)] - \mathbb{E}^{R_\ell}[\Lambda(R, \omega)] \} + \liminf_{\ell \rightarrow \infty} \mathbb{E}^{R_\ell}[\Lambda(R, \omega)] \\ &\geq \mathbb{E}^R[\Lambda(R, \omega)]. \end{aligned}$$

This yields the desired result that the objective functional  $\Gamma$  is l.s.c. on  $\mathcal{R}_{ad}(\nu)$ .  $\square$

The next result also holds.

**Lemma 3.7.** *The set  $\{R \in \mathcal{P}_2(\Omega; \widehat{\Omega}); m \times R(A_i^R) = T, \forall i \in I\}$  is closed in  $\mathcal{P}_2(\Omega; \widehat{\Omega})$ .*

*Proof.* For  $R \in \mathcal{P}_2(\Omega; \Omega')$  and  $i \in I$ , let us define a subset  $\tilde{A}_i^R$  of  $\Omega$  by

$$\tilde{A}_i^R := \left\{ \omega \in \Omega; \int_{t_1}^{t_2} \int_U \phi^i(t, X_t(\omega), R^X(t), u) q_t(\omega, du) dt \geq 0, \forall 0 \leq t_1 < t_2 \leq T \right\}.$$

We first show that  $m \times R(A_i^R) = T$  if and only if  $R(\tilde{A}_i^R) = 1$ .

*Necessity.* For any  $\omega \in \Omega$ , we define the following  $\omega$ -section that

$$A_i^R(\omega) = \left\{ t \in [0, T]; \int_U \phi^i(t, X_t(\omega), R^X(t), u) q_t(\omega, du) \geq 0 \right\}.$$

Then, it follows from Fubini's Theorem that  $T = m \times R(A_i^R) = \int_{\Omega} m(A_i^R(\omega)) R(d\omega)$ . This implies that there exists some  $R$ -null set  $N \in \mathcal{F}$  such that  $m(A_i^R(\omega)) = T$  for all  $\omega \in \Omega \setminus N$ . For such  $\omega$  and  $0 \leq t_1 < t_2 \leq T$ , it holds that

$$\int_{t_1}^{t_2} \int_U \phi^i(t, X_t(\omega), R^X(t), u) q_t(\omega, du) dt = \int_{(t_1, t_2] \cap A_i^R(\omega)} \int_U \phi^i(t, X_t(\omega), R^X(t), u) q_t(\omega, du) dt \geq 0.$$

Consequently,  $\omega \in \tilde{A}_i^R$ , or more precisely,  $\Omega \setminus N \subset \tilde{A}_i^R$ , and hence  $R(\tilde{A}_i^R) = 1$ .

*Sufficiency.* For  $\omega \in \tilde{A}_i^R$ , we have  $m(A_i^R(\omega)) = T$ , and Fubini's Theorem results in  $m \times R(A_i^R) = \int_{\Omega} m(A_i^R(\omega)) R(d\omega) = T$ .

Now, we show that  $\tilde{A}_i^R$  is closed for any  $R \in \mathcal{P}_2(\Omega; \hat{\Omega})$ . Let  $\omega^\ell = (\psi^\ell, q^\ell, \zeta^\ell) \rightarrow \omega = (\psi, q, \zeta)$  in  $\Omega$  as  $\ell \rightarrow \infty$  and  $\omega^\ell \in \tilde{A}_i^R$ . Then, following the proof of Lemma 3.5 in Hausmann [20], we arrive at

$$\int_{t_1}^{t_2} \int_U \phi^i(t, \psi_t(\omega), R^X(t), u) q_t(\omega, du) \geq \limsup_{\ell \rightarrow \infty} \int_{t_1}^{t_2} \int_U \phi^i(t, \psi_t^\ell(\omega), R^X(t), u) q_t^\ell(\omega, du) \geq 0,$$

where we have utilized the fact that  $\phi^i$  is upper semi-continuous in  $(x, \mu, u) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l$  imposed in [Assumption 1](#)-(A2). Hence, it holds that  $\omega \in \tilde{A}_i^R$ , which yields the closedness of  $\tilde{A}_i^R$ .

Let  $R_\ell \rightarrow R$  in  $\mathcal{P}_2(\Omega; \hat{\Omega})$  as  $\ell \rightarrow \infty$  with  $R_\ell$  satisfying  $R_\ell(\tilde{A}_i^{R_\ell}) = 1$  for all  $\ell \geq 1$ . Hence,  $R_\ell(\tilde{A}_i^{R_\ell}) - R_\ell(\tilde{A}_i^R) = R_\ell(\tilde{A}_i^{R_\ell} \setminus \tilde{A}_i^R)$ . However, we have

$$\begin{aligned} \tilde{A}_i^{R_\ell} \setminus \tilde{A}_i^R \subset \left\{ \omega \in \Omega; \int_{t_1}^{t_2} \int_U \phi^i(t, \psi_t(\omega), R_\ell^X(t), u) q_t(\omega, du) \geq 0, \right. \\ \left. \int_U \phi^i(t, \psi_t(\omega), R^X(t), u) q_t(\omega, du) < 0, \exists 0 \leq t_1 < t_2 \leq T \right\} := B_\ell. \end{aligned}$$

Note that  $\phi^i$  is uniformly u.s.c. in  $\mu$  with respect to  $(t, x, u) \in [0, T] \times \mathbb{R}^n \times \mathbb{R}^l$ , which implies that

$$\phi^i(t, \psi_t(\omega), R^X(t), u) q_t(\omega, du) \geq \limsup_{\ell \rightarrow \infty} \phi^i(t, \psi_t(\omega), R_\ell^X(t), u) q_t(\omega, du),$$

uniformly with respect to  $(t, \omega, u) \in [0, T] \times \Omega \times \mathbb{R}^l$ , and hence

$$\int_{t_1}^{t_2} \int_U \phi^i(t, X_t(\omega), R^X(t), u) q_t(\omega, du) \geq \limsup_{\ell \rightarrow \infty} \int_{t_1}^{t_2} \int_U \phi^i(t, X_t(\omega), R_\ell^X(t), u) q_t(\omega, du),$$

uniformly with respect to  $\omega \in \Omega$ . Consequently,  $B_\ell$  is an empty set whenever  $\ell$  is sufficiently large. Thus, it holds that

$$\limsup_{\ell \rightarrow \infty} R_\ell(\tilde{A}_i^{R_\ell}) \leq \limsup_{\ell \rightarrow \infty} \left( R_\ell(\tilde{A}_i^{R_\ell}) - R_\ell(\tilde{A}_i^R) \right) + \limsup_{\ell \rightarrow \infty} R_\ell(\tilde{A}_i^R)$$

$$= \limsup_{\ell \rightarrow \infty} R_\ell(\tilde{A}_i^{R_\ell} \setminus \tilde{A}_i^R) + \limsup_{\ell \rightarrow \infty} R_\ell(\tilde{A}_i^R) \leq R(\tilde{A}_i^R),$$

where, for the last inequality, we have used the fact that  $\tilde{A}_i^R$  is closed and  $R_\ell$  converges weakly to  $R$  as  $\ell \rightarrow \infty$ . This yields that  $R(\tilde{A}_i^R) = 1$ , and hence  $m \times R(\tilde{A}_i^R) = T$ , which completes the proof.  $\square$

**Lemma 3.8.** *For any  $\epsilon > 0$ , let us define the  $\epsilon$ -optimal set  $\mathcal{R}_{\text{opt}}^\epsilon(\nu) = \{R \in \mathcal{R}_{\text{ad}}(\nu); \Gamma(R) \leq v^* + \epsilon\}$ . Then, the set  $\mathcal{R}_{\text{opt}}^\epsilon(\nu)$  is compact in  $\mathcal{P}_2(\Omega; \hat{\Omega})$ .*

*Proof.* We first show that  $\mathcal{R}_{\text{opt}}^\epsilon(\nu)$  is relatively compact in  $\mathcal{P}_2(\Omega; \hat{\Omega})$ . Let  $(R_\ell)_{\ell \geq 1} \subset \mathcal{R}_{\text{opt}}^\epsilon$  be a sequence of admissible control rules. We follow the proof of Theorem 3.7 in [20] to introduce the auxiliary space  $\tilde{\Omega} = \Omega \times C([0, T]; \mathbb{R}^n)$ ,  $\tilde{\mathcal{F}} = \mathcal{F} \otimes \mathcal{F}^C$  and  $\tilde{\mathcal{F}}_t = \mathcal{F}_t \otimes \mathcal{F}_t^C$  for  $t \in [0, T]$ . Here,  $\mathcal{F}^C$  is the Borel  $\sigma$ -algebra on  $C([0, T]; \mathbb{R}^n)$  and  $\mathcal{F}_t^C = \sigma(y(s); s \leq t, y \in C([0, T]; \mathbb{R}^n))$ . Denote by  $\tilde{\omega} = (\psi, q, \zeta, y)$  the generic element of  $\tilde{\Omega}$ . Define the extended probability measure  $\tilde{R}_\ell$  on  $(\tilde{\Omega}, \tilde{\mathcal{F}})$  by

$$\tilde{R}_\ell(F \times C) = \int_F \mathbf{1}_C(y(\omega)) R_\ell(d\omega), \quad \forall F \in \mathcal{F}, C \in \mathcal{F}^C \quad (8)$$

with  $y(\omega)$  being defined by  $y(\omega) = \psi(\omega) - \int_0^\cdot G(s) d\zeta_s(\omega)$ .

By virtue of Lemma 3.5-(i),  $y$  has continuous trajectories under probability  $R_\ell$ , and hence  $\tilde{R}_\ell$  is well-defined. On the other hand, for  $R \in \mathcal{R}_{\text{opt}}^\epsilon(\nu)$ , Assumption 1-(A5) and (A6) yield that

$$\begin{aligned} v^* + \epsilon &\geq \mathbb{E}^R \left[ \int_0^T \int_U f(t, X_t, R^X(t), u) q_t(du) dt + \int_0^T c(t) d\zeta_t \right] \\ &\geq \mathbb{E}^R \left[ - \int_0^T M(1 + |X_t|^p + M_2(\mu)) dt + M_1 \int_0^T \int_U |u|^p q_t(du) dt + c_0 |\zeta_T| \right]. \end{aligned}$$

Consequently, we have

$$\sup_{R \in \mathcal{R}_{\text{opt}}^\epsilon(\nu)} \mathbb{E}^R \left[ \int_0^T \int_U |u|^p q_t(du) dt + |\zeta_T| \right] < +\infty. \quad (9)$$

Then, it follows from Corollary 2.9 in [20] that  $(\tilde{R}_\ell|_{\mathcal{A}^k[0, T]})_{\ell \geq 1}$  is tight. Moreover, according to Proposition B.3 in Lacker [24] and Kolmogorov's tightness criterion, we can deduce that  $(\tilde{R}_\ell|_{\mathcal{Q} \times C([0, T]; \mathbb{R}^n)})_{\ell \geq 1}$  is relatively compact in  $\mathcal{P}_2(\mathcal{Q} \times C([0, T]; \mathbb{R}^n))$ .

We next prove that  $(\tilde{R}_\ell)_{\ell \geq 1}$  is tight. As both  $(\tilde{R}_\ell|_{\mathcal{Q} \times C([0, T]; \mathbb{R}^n)})_{\ell \geq 1}$  and  $(\tilde{R}_\ell|_{\mathcal{A}^k[0, T]})_{\ell \geq 1}$  are tight, for any  $\epsilon > 0$ , there exist some compact subsets  $K \subset \mathcal{Q}$ ,  $V \subset \mathcal{A}^k[0, T]$  and  $C \subset C([0, T]; \mathbb{R}^n)$  such that  $\sup_{\ell \geq 1} \tilde{R}_\ell(D^n[0, T] \times K \times V \times C) \geq 1 - \epsilon$ . Let  $G(V) = \{\int_0^\cdot G(s) d\zeta; \zeta \in V\}$  and  $D = C + G(V) = \{y + g; y \in C, g \in G(V)\}$ . Note that  $C$  and  $G(V)$  are compact in  $C([0, T]; \mathbb{R}^n)$  and  $\mathcal{A}^n[0, T]$ , respectively. Then, they are both compact in  $D^n[0, T]$  and so is  $D$ . Moreover,  $(\tilde{R}_\ell)_{\ell \geq 1}$  is supported on  $\{x = y + \int_0^\cdot G(s) d\zeta\}$ , and thus

$$\sup_{\ell \geq 1} \tilde{R}_\ell(D \times K \times V \times C) = \sup_{\ell \geq 1} \tilde{R}_\ell(D^n[0, T] \times K \times V \times C) \geq 1 - \epsilon.$$

Here,  $D \times K \times V \times C$  is compact in  $\Omega$ , and hence  $(R_\ell)_{\ell \geq 1}$  is tight. Furthermore, it follows from (9) that

$$\sup_{R \in \mathcal{R}_{\text{opt}}^\epsilon(\nu)} \mathbb{E}^R \left[ \sup_{t \in [0, T]} |X_t|^p + \int_0^T |u|^p q_t(du) dt + \sup_{t \in [0, T]} |Y_t|^p \right] < +\infty.$$

Thanks to Corollary B.2 in Lacker [24], we have that  $(\tilde{R}_\ell|_{\Omega' \times C([0, T]; \mathbb{R}^n)})_{\ell \geq 1}$  is relatively compact in  $\mathcal{P}_2(\widehat{\Omega} \times C([0, T]; \mathbb{R}^n))$ . As  $(R_\ell)_{\ell \geq 1}$  is the  $\Omega$ -marginal in the terminology of Jacod [23],  $(R_\ell)_{\ell \geq 1}$  is thus relatively compact in  $\mathcal{P}_2(\Omega; \widehat{\Omega})$ , and hence so is  $\mathcal{R}_{\text{opt}}^\epsilon(\nu)$ .

Lastly, we show that  $\mathcal{R}_{\text{opt}}^\epsilon(\nu)$  is closed. Let  $(R_\ell)_{\ell \geq 1} \subset \mathcal{R}_{\text{opt}}^\epsilon(\nu)$  satisfy  $R_\ell \rightarrow R$  in  $\mathcal{P}_2(\Omega; \widehat{\Omega})$  as  $\ell \rightarrow \infty$ . Then, we prove that the limit point  $R \in \mathcal{R}_{\text{ad}}(\nu)$ . In light of Lemma 3.5 and Lemma 3.7, it suffices to verify claims (i) and (iv) of Lemma 3.5 for the limit  $R$ . To do it, let  $\tilde{R}_\ell$  and  $\tilde{R}$  be the extension measure on  $(\tilde{\Omega}, \tilde{\mathcal{F}})$  as in (8). Then, by construction, we have  $\tilde{R}_\ell \xrightarrow{w} \tilde{R}$  as  $\ell \rightarrow \infty$ . Since the set  $\{\tilde{\omega} \in \tilde{\Omega}; y = x + \int_0^\cdot G(s) d\zeta_s\}$  is closed in  $\tilde{\Omega}$ , it holds that

$$\tilde{R} \left( \left\{ \tilde{\omega} \in \tilde{\Omega}; y = x + \int_0^\cdot G(s) d\zeta_s \right\} \right) \geq \limsup_{\ell \rightarrow \infty} \tilde{R}_\ell \left( \left\{ \tilde{\omega} \in \tilde{\Omega}; y = x + \int_0^\cdot G(s) d\zeta_s \right\} \right) = 1,$$

which proves claim (i) of Lemma 3.5. For the martingale argument, because  $(b, \sigma)$  are uniformly continuous in  $\mu \in \mathcal{P}_p(\mathbb{R}^n)$  (actually there exists a uniform Lipschitz constant) with respect to  $(t, x, u) \in [0, T] \times \mathbb{R}^n \times \mathbb{R}^l$ , for any  $g \in C_b^2(\mathbb{R}^n)$ , it holds that

$$\tilde{M}_t^{R_\ell} g(\omega) \rightarrow \tilde{M}_t^R g(\omega), \quad \ell \rightarrow \infty \quad (10)$$

uniformly with respect to  $(t, \omega) \in [0, T] \times \Omega$ . Moreover, for any bounded  $\mathcal{F}_s$ -measurable random variable  $\eta$  with  $0 \leq s < t \leq T$ , we have

$$\begin{aligned} & \left| \mathbb{E}^R \left[ (\tilde{M}_t^R g(\omega) - \tilde{M}_s^R g(\omega)) \eta(\omega) \right] \right| \leq \mathbb{E}^{R_\ell} \left[ \left| (\tilde{M}_t^{R_\ell} g(\omega) - \tilde{M}_s^{R_\ell} g(\omega)) \eta(\omega) \right| \right] \\ & + \mathbb{E}^{R_\ell} \left[ \left| (\tilde{M}_s^{R_\ell} g(\omega) - \tilde{M}_s^R g(\omega)) \eta(\omega) \right| \right] + \left| \mathbb{E}^R \left[ \tilde{M}_t^R g(\omega) \eta(\omega) \right] - \mathbb{E}^{R_\ell} \left[ \tilde{M}_t^{R_\ell} g(\omega) \eta(\omega) \right] \right| \\ & + \left| \mathbb{E}^R \left[ \tilde{M}_s^R g(\omega) \eta(\omega) \right] - \mathbb{E}^{R_\ell} \left[ \tilde{M}_s^{R_\ell} g(\omega) \eta(\omega) \right] \right| + \left| \mathbb{E}^{R_\ell} \left[ (\tilde{M}_t^{R_\ell} g(\omega) - \tilde{M}_s^{R_\ell} g(\omega)) \eta(\omega) \right] \right| \\ & =: I_1 + I_2 + I_3 + I_4 + I_5, \end{aligned}$$

where the term  $I_5$  equals to 0 as  $R_\ell \in \mathcal{R}_{\text{ad}}(\nu)$ ; while terms  $I_1$  and  $I_2$  converge to 0 as  $\ell \rightarrow \infty$  due to the uniform convergence (10). Lastly, terms  $I_3$  and  $I_4$  converge to 0 because  $\tilde{M}_t^R g(\omega) \eta(\omega)$  is continuous with at most quadratic growth in  $(x, q)$  and  $R_\ell \rightarrow R$  in  $\mathcal{P}_2(\Omega; \widehat{\Omega})$  as  $\ell \rightarrow \infty$ . Hence, we can conclude that  $\mathbb{E}^R[(\tilde{M}_t^R g(\omega) - \tilde{M}_s^R g(\omega)) \eta(\omega)] = 0$ . This implies claim (iv) of Lemma 3.5. Thus, we have completed the proof.  $\square$

We are ready to present the main result of this section:

**Theorem 3.9.** *Let Assumption 1 hold. Then, the set of optimal relaxed control rules  $\mathcal{R}_{\text{opt}}(\nu)$  defined by (6) is a (nonempty) compact and convex subset of  $\mathcal{P}_2(\Omega; \widehat{\Omega})$ . Consequently, there exists an optimal relaxed control to the problem (5).*

*Proof.* First of all, we note that

$$v^* = \inf_{R \in \mathcal{R}_{\text{ad}}(\nu)} \Gamma(R) = \inf_{R \in \mathcal{R}_{\text{opt}}^\epsilon(\nu)} \Gamma(R).$$

Then, it follows from Lemma 3.6 and Lemma 3.8 that  $\mathcal{R}_{\text{opt}}$  is nonempty. It also can be easily checked to be convex and closed. As a closed subset of the compact set  $\mathcal{R}_{\text{opt}}^\epsilon(\nu)$ , it is hence compact.  $\square$

We also have the following corollary as a direct consequence of Lemma 3.4 and Theorem 3.9:

**Corollary 3.10.** *Under Assumption 1 and Assumption 4, there exists an optimal strict control  $(\alpha, \zeta)$  to the constrained MFC problem (3).*

## 4 SMP by Lagrange Multipliers Method

In this section, the goal is to characterize the optimal control by establishing the stochastic maximum principle (SMP) of the constrained MFC problem (1)-(3) with singular control.

To this end, let us first introduce some basic spaces of functions and processes. Let  $\mathcal{X}$  be the space  $L^2([0, T]; \mathbb{R}^l)/\sim$  equipped with the inner product  $\langle \gamma^1, \gamma^2 \rangle_{\mathcal{X}} = \int_0^T \langle \gamma_t^1, \gamma_t^2 \rangle dt$ . Then  $(\mathcal{X}, \langle \cdot, \cdot \rangle_{\mathcal{X}})$  is a Hilbert space. Let  $\mathbb{L}_{\mathbb{F}}^2$  be the set of  $\mathbb{F}$ -adapted processes  $\beta = (\beta_t)_{t \in [0, T]}$  taking values in  $\mathbb{R}^l$  such that  $\mathbb{E}[\int_0^T |\beta_t|^2 dt] < \infty$ . As a result,  $\mathbb{L}_{\mathbb{F}}^2$  is a Hilbert space equipped with the inner product  $\langle \beta^1, \beta^2 \rangle_{\mathbb{L}_{\mathbb{F}}^2} = \mathbb{E}[\int_0^T \langle \beta_t^1, \beta_t^2 \rangle dt]$  for  $\beta^1, \beta^2 \in \mathbb{L}_{\mathbb{F}}^2$ . It can be easily verified that  $\mathcal{W}[0, T]$  is closed and convex in  $(\mathbb{L}_{\mathbb{F}}^2, \langle \cdot, \cdot \rangle_{\mathbb{L}_{\mathbb{F}}^2})$ . Let us define  $\mathbb{M}^2 (\subset \mathbb{L}_{\mathbb{F}}^2)$  as the set of  $\mathbb{F}$ -adapted (continuous)  $\mathbb{R}^n$ -valued process  $\eta = (\eta_t)_{t \in [0, T]}$  satisfying the Itô's representation that

$$\eta_t = \eta_0 + \int_0^t \eta_s^1 ds + \int_0^t \eta_s^2 dW_s, \quad (11)$$

where  $\eta_0 \in L^2(\Omega; \mathbb{R}^n)$  is  $\mathcal{F}_0$ -measurable,  $\eta^1 = (\eta_t^1)_{t \in [0, T]}$  and  $\eta^2 = (\eta_t^2)_{t \in [0, T]}$  take values respectively in  $\mathbb{R}^n$  and  $\mathbb{R}^{n \times r}$ , which are  $\mathbb{F}$ -adapted processes such that  $\mathbb{E}[\int_0^T (|\eta_t^1|^2 + |\eta_t^2|^2) dt] < \infty$ . Thus,  $\mathbb{M}^2$  is a Hilbert space under the inner product  $\langle \cdot, \cdot \rangle_{\mathbb{M}^2}$  defined by

$$\langle \eta, \delta \rangle_{\mathbb{M}^2} := \mathbb{E} \left[ \langle \eta_0, \delta_0 \rangle + \int_0^T \left( \langle \eta_t^1, \delta_t^1 \rangle + \text{tr} \left( \eta_t^2 (\delta_t^2)^\top \right) \right) dt \right], \quad \forall \eta, \delta \in \mathbb{M}^2. \quad (12)$$

We use  $\mathbb{L}_{\mathbb{F}, \mathbb{R}}^2$  to represent the set of adapted processes in  $L^2([0, T] \times \Omega; \mathbb{R})/\sim$ . Then,  $\mathbb{L}_{\mathbb{F}, \mathbb{R}}^2$  is a Hilbert space with the inner product  $\langle \beta^1, \beta^2 \rangle_{\mathbb{L}_{\mathbb{F}, \mathbb{R}}^2} := \mathbb{E}[\int_0^T \beta_t^1 \beta_t^2 dt]$  for  $\beta^1, \beta^2 \in \mathbb{L}_{\mathbb{F}, \mathbb{R}}^2$ .

As a technical preparation, we give the definition of the  $L$ -derivative as below, see Carmona and Delarue [10].

**Definition 4.1.** *A function  $u$  on  $\mathcal{P}_2(\mathbb{R}^n)$  is said to be  $L$ -differentiable at  $\mu_0 \in \mathcal{P}_2(\mathbb{R}^n)$  if there exists a random variable  $X_0$  with law  $\mu_0$ , in other words satisfying  $\mathcal{L}(X_0)$ , such that the lifted function  $\tilde{u}$  is Fréchet differentiable at  $X_0$ .*

For a continuously  $L$ -differentiable mapping  $u : \mathcal{P}_2(\mathbb{R}^n) \rightarrow \mathbb{R}$ , by Section 5.2.1 in Carmona and Delarue [10], for any  $\mu \in \mathcal{P}_2(\mathbb{R}^n)$  there exists a measurable mapping  $\partial_\mu u(\mu) : \mathbb{R}^n \times \mathbb{R}^n$  such that  $\partial_\mu u(\mu)(X) = D\tilde{u}(X)$  for all  $X \in L^2(\Omega; \mathbb{R}^n)$  with  $\mathcal{L}(X) = \mu$ . As a result of Definition 4.1, the following lemma holds.

**Lemma 4.2.** *Let  $u$  be continuously  $L$ -differentiable and denote  $\mu := \mathcal{L}(\tilde{X})$  for an  $\mathbb{R}^n$ -valued r.v.  $\tilde{X}$  on some probability space  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ . Then, for any  $\mathbb{R}^n$ -valued r.v.  $(\tilde{X}')$ , it holds that*

$$w(\mathcal{L}(\tilde{X}')) = w(\mu) + \mathbb{E}^{\tilde{P}} \left[ \langle \partial_\mu w(\mu)(\tilde{X}), \tilde{X}' - \tilde{X} \rangle \right] + o \left( \sqrt{\mathbb{E}^{\tilde{P}}[|\tilde{X}' - \tilde{X}|^2]} \right),$$

when  $\tilde{X}'$  is close to  $\tilde{X}$  in  $L^2((\tilde{\Omega}; \mathbb{R}^n))$ . Here,  $\mathbb{E}^{\tilde{P}}$  denotes the expectation operator under  $\tilde{P}$ , and the above  $\langle \cdot, \cdot \rangle$  is the inner product of  $L^2(\tilde{\Omega}; \mathbb{R}^n)$ .

To adopt the methodology introduced in Bo et al. [6], we propose to embed the state process into a suitable Banach space. Without loss of generality, we can assume that for any singular control  $\zeta \in \mathcal{A}^k[0, T]$ , the associated radon measures  $\mu_i$  for  $i \in I$  of the trajectory  $t \mapsto \zeta_t(\omega)$  is singular w.r.t. Lebesgue measure. Otherwise, we can decompose  $\mu_i$  using the Lebesgue decomposition w.r.t. the Lebesgue measure and absorb the absolutely continuous (regular) term to the drift  $b$ . Consider the following space:

$$\mathcal{X} := \left\{ X = (X_t)_{t \in [0, T]}; X \text{ is } \mathbb{F}\text{-adapted process with } X_0 \in L^2(\Omega; \mathbb{R}^n) \text{ and} \right. \quad (13)$$



$$X = X_0 + \int_0^\cdot X_s^1 ds + \int_0^\cdot X_s^2 dW_s + \int_0^\cdot G(s) dX_s^3 \text{ with } (X^1, X^2, X^3) \in \mathbb{L}_{\mathbb{F}}^2 \times \mathbb{L}_{\mathbb{F}}^2 \times \mathcal{A}^k[0, T].$$

Consequently,  $\mathcal{X}$  is a Banach space under the norm  $\|\cdot\|_{\mathcal{X}}$  given by

$$\|X\|_{\mathcal{X}} := \|X^1\|_{\mathbb{L}_{\mathbb{F}}^2} + \|X^2\|_{\mathbb{L}_{\mathbb{F}}^2} + \sqrt{\mathbb{E} \left[ \left| \int_0^T G(s) dX_s^3 \right|^2 \right]}, \quad \forall X \in \mathcal{X}.$$

By the assumption that, for any  $X \in \mathcal{X}$ , its 3rd component  $X^3$  of the decomposition in  $\mathcal{X}$  is singular w.r.t. to Lebesgue measure, one can conclude that  $\mathcal{X}$  is isometric to the product space  $\mathbb{M}^2 \times \mathbb{G}^2$  under the product norm, where  $(\mathbb{G}^2, \|\cdot\|_{\mathbb{G}^2})$  is a Banach space defined by

$$\mathbb{G}^2 := \left\{ S = (S_t)_{t \in [0, T]}; S = \int_0^\cdot G(s) d\eta_s \text{ for } \eta = (\eta_t)_{t \in [0, T]} \in \mathcal{A}^k[0, T] \right\},$$

$$\|S\|_{\mathbb{G}^2} := \sqrt{\mathbb{E} \left[ \left| \int_0^T G(s) d\eta_s \right|^2 \right]}.$$

As  $\int_0^\cdot G(s) d\eta_s \in (\mathcal{M}[0, T])^n$  for  $\eta \in \mathcal{A}^k[0, T]$ , we can identify  $\mathbb{G}^2$  as a subset of  $L^2(\Omega; (\mathcal{M}[0, T])^n)$  with  $(\mathcal{M}[0, T])^n$  equipped with the norm  $\|\mu\|_{(\mathcal{M}[0, T])^n} = \sqrt{\sum_{i=1}^n \|\mu_i\|_{\text{TV}}^2}$ . Consequently,  $\mathcal{X}$  can be viewed as a subset of  $\mathbb{T}^2 := \mathbb{M}^2 \times L^2(\Omega; (\mathbb{M}[0, T])^n)$  under the product norm. For any  $\lambda = (\lambda^1, \lambda^2) \in \mathbb{T}^{2,*} \simeq \mathbb{M}^2 \times L^2(\Omega; (C([0, T]; \mathbb{R}^{**})^n)$ , the dual pair for  $X = L + S \in \mathcal{X}$  with  $(L, S) \in \mathbb{M}^2 \times \mathbb{G}^2$  is defined by

$$\langle X, \lambda \rangle_{\mathbb{T}^2, \mathbb{T}^{2,*}} = \langle L, \lambda^1 \rangle_{\mathbb{M}^2} + \langle S, \lambda^2 \rangle_{L^2(\Omega; (\mathbb{M}[0, T])^k), L^2(\Omega; (C([0, T]; \mathbb{R}^{**})^k)}.$$

Following Theorem 2.7 in Bo et al. [6], let us introduce

$$\mathbf{X} = \mathbb{T}^2 \times \mathbb{L}_{\mathbb{F}}^2 \times L^2(\Omega; (\mathbb{M}[0, T])^n), \quad \mathbf{Y}_i = \mathbb{L}_{\mathbb{F}, \mathbb{R}}^2, \quad \forall i \in I, \quad \mathbf{H} = \mathbb{T}^2. \quad (14)$$

The norm for  $\mathbf{X}$  is defined to be the (default) product norm and we take  $\|\cdot\|_{\mathbf{Y}_i}$  ( $i \in I$ ) as the norm  $\|\cdot\|_{\mathbb{L}_{\mathbb{F}, \mathbb{R}}^2}$ , respectively. Moreover, we define  $J : \mathbf{X} \rightarrow \mathbb{R}$ ,  $\Phi_i : \mathbf{X} \rightarrow \mathbf{Y}_j$  ( $i \in I$ ) and  $F : \mathbf{X} \rightarrow \mathbf{H}$  as follows, for  $(X, \alpha, \zeta) \in \mathbf{X}$ ,

$$\left\{ \begin{array}{l} J(X, \alpha, \zeta) := \mathbb{E} \left[ \int_0^T f(t, X_t, \mathcal{L}(X_t), \alpha_t) + \int_0^T c(t) d\zeta_t \right], \\ \Phi_i(X, \alpha, \zeta)(t) := -\phi^i(t, X_t, \mathcal{L}(X_t), \alpha_t), \quad \forall t \in [0, T], \quad i \in I, \\ F(X, \alpha, \zeta)(\cdot) := X - \xi - \int_0^\cdot b(s, X_s, \mathcal{L}(X_s), \alpha_s) ds - \int_0^\cdot \sigma(s, X_s, \mathcal{L}(X_s), \alpha_s) dW_s \\ \quad - \int_0^\cdot G(s) d\zeta_s. \end{array} \right. \quad (15)$$

The well-posedness of the mappings  $\Phi_i$  for  $i \in I$  is guaranteed by [Assumption 1-\(A2\)](#). Thus, we can formulate the following optimization problem (equivalent to the constrained MFC problem (3)) with constraints that

$$\left\{ \begin{array}{l} \inf J(X, \alpha, \zeta) \\ \text{s.t. } (X, \alpha, \zeta) \in C := \mathcal{X} \times \mathcal{U}[0, T] \times \mathcal{A}^k[0, T], \\ \Phi_i(X, \alpha, \zeta) \leq_K 0, \quad \forall i \in I, \\ F(X, \alpha, \zeta) = 0, \end{array} \right. \quad (16)$$

where  $C = \mathcal{X} \times \mathcal{U}[0, T] \times \mathcal{A}^k[0, T]$  is closed and convex in  $\mathbf{X}$ , and the set  $K$  is a closed and convex cone given by

$$K := \{f \in \mathbb{L}_{\mathbb{F}, \mathbb{R}}^2; f(t, \omega) \geq 0, \forall (t, \omega) \in [0, T] \times \Omega\} =: \mathbb{L}_{\mathbb{F}, \mathbb{R}^+}^2. \quad (17)$$

Then, in view of Theorem 3.12 in [6], the FJ optimality condition holds for the constrained optimization problem (16) in the following sense.

**Lemma 4.3** (FJ optimality condition for problem (16)). *Let  $(\widehat{X}, \widehat{\alpha}, \widehat{\zeta}) \in C$  be an optimal pair to problem (16). Then, there exist (nontrivial) multipliers  $r_0 \geq 0$ ,  $\eta^i \in \mathbb{L}_{\mathbb{F}, \mathbb{R}^+}^2$  for  $i \in I$  and  $\lambda = (\lambda_1, \lambda_2) \in \mathbb{M}^2 \times L^2(\Omega; (C([0, T], \mathbb{R})^{**})^n)$ , which are not all zeros, such that*

$$\left\{ \begin{array}{l} \mathbb{E} \left[ \int_0^T \Phi_i(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(t) \eta_t^i dt \right] = 0, \quad \forall i \in I, \\ r_0 + \sum_{i \in I} \|\eta^i\|_{\mathbb{L}_{\mathbb{F}, \mathbb{R}}^2} + \|\lambda\|_{\mathbb{M}^2} = 1, \\ -\xi_1 = r_0 D_X J(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot) + \sum_{i \in I} \mathbb{E} \left[ \int_0^T D_X \Phi_i(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot)(t) \eta_t^i dt \right] - \left\langle \lambda, D_X F(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot) \right\rangle_{\mathbb{H}^*, \mathbb{H}} \quad \text{in } \mathbb{T}^*, \\ -\xi_2 = r_0 D_\alpha J(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot) + \sum_{i \in I} \mathbb{E} \left[ \int_0^T D_\alpha \Phi_i(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot)(t) \eta_t^i dt \right] - \left\langle \lambda, D_\alpha F(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot) \right\rangle_{\mathbb{H}^*, \mathbb{H}} \quad \text{in } \mathbb{L}_{\mathbb{F}}^2, \\ -\xi_3 = r_0 D_\zeta J(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot) + \sum_{i \in I} \mathbb{E} \left[ \int_0^T D_\zeta \Phi_i(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot)(t) \eta_t^i dt \right] \\ \quad - \left\langle \lambda, D_\zeta F(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})(\cdot) \right\rangle_{\mathbb{H}^*, \mathbb{H}} \quad \text{in } L^2(\Omega; (C([0, T]; \mathbb{R})^{**})^n), \end{array} \right.$$

where  $\xi = (\xi_1, \xi_2, \xi_3) \in N_C(\widehat{X}, \widehat{\alpha}, \widehat{\zeta}) \subset \mathbf{X}^*$  and  $\|\cdot\|_{\text{TV}}$  is the total variation norm on the space of (signed) Radon measures.

Then, we have the following main result in this section.

**Theorem 4.4** (SMP for constrained MFC problem with singular control (3)). *Let  $(\widehat{X}, \widehat{\alpha}, \widehat{\zeta}) \in C$  be an optimal solution to problem (3). There exist some  $r_0 \geq 0$  and  $\eta^i \in \mathbb{L}_{\mathbb{F}, \mathbb{R}^+}^2$  for  $i \in I$  such that*

- (i) *for  $i \in I$ ,  $\eta_t^i \geq 0$ ,  $dt \times d\mathbb{P}$ -a.s. and  $\eta^i \in \mathbb{L}_{\mathbb{F}, \mathbb{R}^+}^2$ , let  $A^j = (A_t^j)_{t \in [0, T]}$  be the (continuous) increasing process that  $A_t^i = \int_0^t \eta_s^i ds$ . Then, for  $\mathbb{P}$ -a.s.  $\omega$ , the process  $A^i(\omega)$  only increases on the set  $\{t \in [0, T]; \phi^j(t, \widehat{X}_t(\omega), \widehat{\mu}_t, \widehat{\alpha}_t(\omega)) = 0\}$ ,  $dt$ -a.s., with  $\widehat{\mu}_t = \mathcal{L}(\widehat{X}_t)$ ;*
- (ii) *it holds that  $r_0 + \sum_{i \in I} \|\eta^i\|_{\mathbb{L}_{\mathbb{F}, \mathbb{R}}^2} = 1$ ;*
- (iii)  *$\mathbb{P}(r_0 c_i(t) + G_i(t)^\top Y_t \geq 0, \forall (t, i) \in [0, T] \times \{1, \dots, k\}) = 1$  with  $c_i(t)$  and  $G_i(t)$  being the  $i$ -th row of  $c(t)$  and  $G(t)$ , respectively. Moreover, it holds that*

$$\mathbb{P} \left( \sum_{i=1}^k \mathbf{1}_{\{r_0 c_i(t) + G_i(t)^\top Y_t > 0\}} d\widehat{\zeta}_t^i = 0 \right) = 1. \quad (18)$$

Furthermore, the stochastic (first-order) minimum condition holds that, for any  $u \in U$  and  $dt \times d\mathbb{P}$ -a.s.,

$$\left\langle \nabla_u H^{r_0}(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t, Y_t, Z_t) - \sum_{i \in I} \nabla_u \phi^i(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) \eta_t^i, u - \widehat{\alpha}_t \right\rangle \geq 0, \quad \forall u \in U. \quad (19)$$

The Hamiltonian  $H^{r_0} : [0, T] \times \mathbb{R}^n \times \mathbb{R}^l \times \mathbb{R}^n \times \mathbb{R}^{n \times r} \mapsto \mathbb{R}$  is defined by

$$H^{r_0}(t, x, u, \mu, y, z) := \langle b(t, x, \mu, u), y \rangle + \text{tr} \left( \sigma(t, x, \mu, u) z^\top \right) + r_0 f(t, x, \mu, u), \quad (20)$$

and the process pair  $(Y, Z) = (Y_t, Z_t)_{t \in [0, T]}$  taking values in  $\mathbb{R}^n \times \mathbb{R}^{n \times r}$  is the unique solution to the constrained BSDE:

$$\begin{aligned} Y_t = & \int_t^T \nabla_x H^{r_0}(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s, Y_s, Z_s) ds + \int_t^T \mathbb{E}' \left[ \partial_\mu H(s, \widehat{X}'_s, \widehat{\mu}_s, \widehat{\alpha}'_s, Y'_s, Z'_s)(\widehat{X}_s) \right] ds \\ & - \sum_{i \in I} \int_t^T \nabla_x \phi^i(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s) \eta_s^i ds - \sum_{i \in I} \int_t^T \mathbb{E}' \left[ \partial_\mu \phi^i(s, \widehat{X}'_s, \widehat{\mu}_s, \widehat{\alpha}'_s)(\widehat{X}_s) \eta_s^{i'} \right] ds - \int_t^T Z_s dW_s. \end{aligned} \quad (21)$$

Here,  $(\widehat{X}', \widehat{\alpha}', Y', Z', \eta^{i'})$  is an independent copy of  $(\widehat{X}, \widehat{\alpha}, Y, Z, \eta^i)$  defined on the probability space  $(\Omega', \mathcal{F}', P')$ .

*Proof.* Firstly,  $r_0$  and  $(\eta^i)_{i \in I}$  can't be zeros simultaneously (see the first steps in the proof of Theorem 3.12 in [6]). Hence we can scale the multipliers such that (iii) holds. By using definition of the normal cone  $N_C(\widehat{X}, \widehat{\alpha}, \widehat{\zeta})$ , it follows that for all  $(X, \alpha, \zeta)$  and  $\xi = (\xi_1, \xi_2, \xi_3) \in N_C(\widehat{X}, \widehat{\alpha})$ ,

$$\langle \xi_1, \widehat{X} - X \rangle_{\mathbb{T}^2} + \langle \xi_2, \widehat{\alpha} - \alpha \rangle_{\mathbb{L}_{\mathbb{F}}^2} + \langle \xi_3, \zeta - \widehat{\zeta} \rangle_{(C([0, T]; \mathbb{R}))^{**n}, (\mathcal{M}[0, T])^n} \geq 0.$$

Taking  $\alpha = \widehat{\alpha}, \zeta = \widehat{\zeta}$ , we have that  $\xi_1 = 0, \langle \xi_2, \widehat{\alpha} - \alpha \rangle_{\mathbb{L}_{\mathbb{F}}^2} \geq 0, \langle \xi_3, \zeta - \widehat{\zeta} \rangle_{(C([0, T]; \mathbb{R}))^{**n}, (\mathcal{M}[0, T])^n} \geq 0$  similar as [6] (there is a slight change in the proof of Lemma 3.10 to fit into our setting here)  $\xi_1 = 0$  in Lemma 4.3 and Theorem 4.4-(iii) hold true. Then, we can conclude from Lemma 4.2 and Lemma 4.3 that

$$\begin{aligned} 0 = & r_0 \mathbb{E} \left[ \int_0^T \langle \nabla_x f(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t), L_t \rangle dt + \mathbb{E}' \left[ \int_0^T \langle \partial_\mu f(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t)(\widehat{X}'_t), L'_t \rangle dt \right] \right] \\ & - \sum_{i \in I} \mathbb{E} \left[ \int_0^T \left\langle \nabla_x \phi^i(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) + \mathbb{E}' \left[ \partial_\mu \phi^i(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t)(\widehat{X}'_t) \right], L'_t \right\rangle \eta_t^i dt \right] \\ & - \mathbb{E} \left[ \langle \lambda_1(0), L_0 \rangle + \int_0^T \langle \lambda_1^1(t), L_t^1 - \nabla_x b(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) L_t \rangle dt + \int_0^T \text{tr} \left( \lambda_1^2(t) (L_t^2 - \nabla_x \sigma(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) L_t)^\top \right) dt \right. \\ & \left. + \int_0^T \left\langle \lambda_1^1(t), \mathbb{E}' \left[ \partial_\mu b(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t)(\widehat{X}'_t) L'_t \right] \right\rangle dt + \int_0^T \text{tr} \left( \lambda_1^2(t) \left( \mathbb{E}' \left[ \partial_\mu \sigma(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t)(\widehat{X}'_t) L'_t \right] \right)^\top \right) dt \right] \\ & + \langle \lambda_2, \nu \rangle_{(C([0, T]; \mathbb{R}))^{**n}, (\mathcal{M}[0, T])^n}, \end{aligned} \quad (22)$$

where  $\lambda_1 \in \mathbb{M}^2$  has the decomposition  $\lambda_1(\cdot) = \lambda_1(0) + \int_0^\cdot \lambda_1^1(s) ds + \int_0^\cdot \lambda_1^2(s) dW_s$ ,  $L \in \mathbb{T}^2$  has the representation  $L = L_0 + \int_0^\cdot L_s^1 ds + \int_0^\cdot L_s^2 dW_s + \int_0^\cdot G(s) dL_s^3$ , and  $\nu$  is the associated radon measure of the bounded variation mapping  $\int_0^\cdot G(s) dL_s^3 \in \text{BV}^n[0, T]$ . Define an element  $p = (p_t)_{t \in [0, T]}$  in  $\mathbb{M}^2$  that, for  $t \in [0, T]$ ,

$$\begin{aligned} p_t = & - \sum_{i \in I} \int_t^T \left( \nabla_x \phi^i(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s) + \mathbb{E}' \left[ \partial_\mu \phi^i(s, \widehat{X}'_s, \widehat{\mu}_s, \widehat{\alpha}'_s)(\widehat{X}_s) \eta_s^{i'} \right] \right) ds \\ & + \int_t^T \left[ r_0 \nabla_x f(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s) + \nabla_x b(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s) \lambda_1^1(s) + \nabla_x \sigma(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s) \lambda_1^2(s) \right] ds \\ & + \int_t^T \mathbb{E}' \left[ r_0 \partial_\mu f(s, \widehat{X}'_s, \widehat{\mu}_s, \widehat{\alpha}'_s)(\widehat{X}_s) + \partial_\mu b(s, \widehat{X}'_s, \widehat{\mu}_s, \widehat{\alpha}'_s)(\widehat{X}_s) \lambda_1^1(s)' \right. \\ & \left. + \partial_\mu \sigma(s, \widehat{X}'_s, \widehat{\mu}_s, \widehat{\alpha}'_s)(\widehat{X}_s) \lambda_1^2(s)' \right] ds - \int_t^T \lambda_1^2(s) dW_s, \end{aligned} \quad (23)$$

where  $MN := \sum_{i=1}^r M_i L_i \in \mathbb{R}^n$  for any  $M = (M_1, \dots, M_r) \in \mathbb{R}^{n \times n \times r}$  and  $L = (L_1, \dots, L_r) \in \mathbb{R}^{n \times r}$ . By construction of  $L$ , we have

$$\langle \lambda_1^1(t), L_t^1 \rangle dt = \langle \lambda_1^1(t) - p(t), dL_t \rangle + \langle p(t), dL_t \rangle - \langle \lambda_1^1, \nu \rangle_{(C([0,T]; \mathbb{R})^{**})^n, (\mathcal{M}[0,T])^n}. \quad (24)$$

Noting (23) and applying Itô's formula, we can conclude that

$$\begin{aligned} \mathbb{E} \left[ \int_0^T \langle p_t, dL_t \rangle \right] &= -\mathbb{E} \left[ \int_0^T \langle L_t, dp_t \rangle dt \right] - \mathbb{E} [\langle p, L \rangle_T] \\ &= \mathbb{E} \left[ \int_0^T \left\langle r_0 \nabla_x f(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t), L_t \right\rangle dt + \int_0^T \left\langle r_0 \mathbb{E}' \left[ \partial_\mu f(t, \widehat{X}_t', \widehat{\mu}_t, \widehat{\alpha}_t')(\widehat{X}_t) \right], L_t \right\rangle dt \right] \\ &\quad + \mathbb{E} \left[ \int_0^T \left( \langle \lambda_1^1(t), \nabla_x b(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) L_t \rangle + \text{tr} \left( \lambda_1^2(t) \left( \nabla_x \sigma(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) L_t \right)^\top \right) \right) \right] \\ &\quad + \mathbb{E} \left[ \int_0^T \left\langle \lambda_1^1(t), \mathbb{E}' \left[ \partial_\mu b(t, \widehat{X}_t', \widehat{\mu}_t, \widehat{\alpha}_t')(\widehat{X}_t) L_t \right] \right\rangle dt - \langle p_0, L_0 \rangle \right] - \mathbb{E} \left[ \text{tr} \left( \lambda_1^2(t) L_t^2 \right) dt \right] \\ &\quad + \mathbb{E} \left[ \int_0^T \text{tr} \left( \lambda_1^2(t) \left( \mathbb{E}' \left[ \partial_\mu \sigma(t, \widehat{X}_t', \widehat{\mu}_t, \widehat{\alpha}_t')(\widehat{X}_t) L_t \right] \right)^\top \right) dt \right] \\ &\quad - \sum_{i \in I} \mathbb{E} \left[ \int_0^T \left\langle \nabla_x \phi^i(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t), L_t \right\rangle dt + \int_0^T \left\langle \mathbb{E}' \left[ \partial_\mu \phi^i(t, \widehat{X}_t', \widehat{\mu}_t, \widehat{\alpha}_t')(\widehat{X}_t) \right], L_t \right\rangle dt \right]. \end{aligned}$$

Inserting (24) together with the above equality into (22), we derive that

$$\mathbb{E} \left[ \langle p_0 - \lambda_1(0), L_0 \rangle + \int_0^T \langle p_t - \lambda_1^1(t), dL_t \rangle + \langle \lambda_2 - \lambda_1^1, \nu \rangle_{(C([0,T]; \mathbb{R})^{**})^n, (\mathcal{M}[0,T])^n} \right] = 0.$$

We can thus deduce that  $p_0 = \lambda_1(0)$  and  $p$  is indistinguishable from  $\lambda_1^1$  by the arbitrariness of  $L$ . Then  $\lambda_1^1$  solves (30). Moreover, we have  $G^\top \lambda_2 = G^\top \lambda_1^1$  in  $(C([0, T]; \mathbb{R})^{**})^k$ , and hence  $G^\top \lambda_2 \in C([0, T]; \mathbb{R}^k)$ . Here,  $G$  can be viewed a mapping in  $\mathcal{L}((C([0, T]; \mathbb{R})^{**})^n, ((C([0, T]; \mathbb{R})^{**})^k))$ . By the optimality condition for  $\widehat{\alpha} \in \mathcal{U}[0, T]$ , we have

$$\begin{aligned} 0 &\leq \mathbb{E} \left[ \int_0^T \left\langle r_0 \nabla_u f(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) + \nabla_u b(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t)^\top \lambda_t^1 + \nabla_u \sigma(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t)^\top \lambda_t^2, \alpha_t - \widehat{\alpha}_t \right\rangle dt \right. \\ &\quad \left. - \sum_{i \in I} \int_0^T \left\langle \nabla_u \phi^i(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) \eta_t^i, \alpha_t - \widehat{\alpha}_t \right\rangle dt \right]. \quad (25) \end{aligned}$$

Exploiting the fact that  $(\lambda_1^1, \lambda_1^2)$  solves (30), we then can derive that (25) indicates (29) by the arbitrariness of  $\alpha \in \mathcal{U}[0, T]$ . By the optimality condition for  $\widehat{\zeta} \in \mathcal{A}^k[0, T]$ , we arrive at

$$\begin{aligned} 0 &\leq -\mathbb{E} \left[ \langle \xi_3, \eta - \widehat{\zeta} \rangle_{(C([0,T]; \mathbb{R})^{**})^n, (\mathcal{M}[0,T])^n} \right] \\ &= \mathbb{E} \left[ r_0 \int_0^T c(t) d(\eta_t - \widehat{\zeta}_t) + \left\langle \lambda^2, \int_0^T G(t) d(\eta_t - \widehat{\zeta}_t) \right\rangle_{(C([0,T]; \mathbb{R})^{**})^n, (\mathcal{M}[0,T])^n} \right] \\ &= \mathbb{E} \left[ r_0 \int_0^T c(t) d(\eta_t - \widehat{\zeta}_t) + \int_0^T G(t)^\top \lambda_1^1(t) d(\eta_t - \widehat{\zeta}_t) \right], \end{aligned}$$

where for the last equality, we have used the fact that  $G^\top \lambda_2 = G^\top \lambda_1^1$ . Taking  $\eta_t = \widehat{\zeta}_t + \delta_t \in \mathcal{A}^k[0, T]$  for any  $\delta = (\delta_t)_{t \in [0, T]} \in \mathcal{A}^k[0, T]$ , we derive that

$$\mathbb{P} \left( r_0 c_i(t) + G_i(t)^\top \lambda_1^1(t) \geq 0, \forall (t, i) \in [0, T] \times \{1, \dots, k\} \right) = 1.$$

Moreover, taking  $\eta_t \equiv 0$  results in (28). Thus, the proof of the theorem is complete by just noting that (i) follows from Lemma 4.3.  $\square$

**Remark 4.5.** When there are no constraints (i.e.  $m = 0$ ) and there is no mean-field term, [Theorem 4.4](#) together with the CQ condition derived in [\[6\]](#) will reduce to the stochastic maximum principle as in [Bahali \[3\]](#).

## 5 Uniqueness and Stability of Solution to Constrained FBSDE

The aim of this section is to study the uniqueness and stability of solutions to the following generalized type of constrained FBSDE derived in [Theorem 4.4](#). Without loss of generality, we assume that  $d = 1$ :

$$\left\{ \begin{array}{l} \widehat{X}_t = \xi + \int_0^t b(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s) ds + \int_0^t \sigma(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s) dW_s + \int_0^t G(s) d\widehat{\zeta}_s, \\ Y_t = \int_t^T \nabla_x H^{r_0}(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s, Y_s, Z_s) ds \\ \quad + \int_t^T \mathbb{E}' \left[ \partial_\mu H^{r_0}(s, \widehat{X}'_s, \widehat{\mu}'_s, \widehat{\alpha}'_s, Y'_s, Z'_s)(\widehat{X}_s) \right] ds - \int_t^T \nabla_x \phi(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s) \eta_s ds \\ \quad - \int_t^T \mathbb{E}' \left[ \partial_\mu \phi(s, \widehat{X}'_s, \widehat{\mu}'_s, \widehat{\alpha}'_s)(\widehat{X}_s) \eta'_s \right] ds - \int_t^T Z_s dW_s, \end{array} \right. \quad (26)$$

subject to  $|r_0| + \|\eta\|_{\mathbb{L}_{\mathbb{F}, \mathbb{R}}} = 1$ ,  $\widehat{\mu}_t = \mathcal{L}(\widehat{X}_t)$ ,  $\mathbb{P}(G(t)Y_t + c(t) \geq 0, \forall t \in [0, T]) = 1$  and

$$\left\{ \begin{array}{l} \left\langle \nabla_u H^{r_0}(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t, Y_t, Z_t) - \sum_{i \in I} \nabla_u \phi^i(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) \eta_t^i, u - \widehat{\alpha}_t \right\rangle \geq 0, \forall u \in U, \\ \int_0^T \phi(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) \eta_t dt = 0, \phi(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t) \geq 0, \eta_t \geq 0, dt \times d\mathbb{P}\text{-a.s.}, \\ \mathbb{P} \left( \sum_{i=1}^k \mathbf{1}_{\{r_0 c_i(t) + G_i(t)^\top Y_t > 0\}} d\widehat{\zeta}_t^i = 0 \right) = 1, \end{array} \right. \quad (27)$$

which characterizes the necessary optimality condition of the constrained MFC problem [\(3\)](#) with singular control. To prove the uniqueness of solution  $(\widehat{X}, \widehat{\alpha}, \widehat{\zeta}, Y, Z, \eta) = (\widehat{X}_t, \widehat{\alpha}_t, \widehat{\zeta}_t, Y_t, Z_t, \eta_t)_{t \in [0, T]}$  satisfying [\(26\)](#)-[\(27\)](#), we first give the definition of uniqueness of solutions to [Eq. \(26\)](#)-[\(27\)](#):

**Definition 5.1** (Uniqueness of solutions to FBSDE [\(26\)](#)-[\(27\)](#)). Let  $(\widehat{X}^i, \widehat{\alpha}^i, \widehat{\zeta}^i, Y^i, Z^i, \eta^i) = (\widehat{X}_t^i, \widehat{\alpha}_t^i, \widehat{\zeta}_t^i, Y_t^i, Z_t^i, \eta_t^i)_{t \in [0, T]}$  for  $i = 1, 2$  be  $\mathbb{F}$ -adapted processes valued in  $\mathbb{R}^n \times U \times \mathbb{R}_+^k \times \mathbb{R}^n \times \mathbb{R}^{n \times r} \times \mathbb{R}_+$  satisfying the coupling constrained FBSDE [\(26\)](#)-[\(27\)](#). The solution of FBSDE [\(26\)](#)-[\(27\)](#) is said to be unique if  $\widehat{X}^1 = \widehat{X}^2$ ,  $\widehat{\alpha}^1 = \widehat{\alpha}^2$ ,  $Y^1 = Y^2$ ,  $Z^1 = Z^2$ ,  $\eta^1 = \eta^2$  and  $\int_0^\cdot G(s) d\widehat{\zeta}_s^1 = \int_0^\cdot G(s) d\widehat{\zeta}_s^2$ .

Note that, we only require  $\int_0^\cdot G(s) d\widehat{\zeta}_s^1 = \int_0^\cdot G(s) d\widehat{\zeta}_s^2$  rather than  $\widehat{\zeta}^1 = \widehat{\zeta}^2$  in [Definition 5.1](#). Once the existence of optimal triple  $(\widehat{X}, \widehat{\alpha}, \widehat{\zeta}) = (\widehat{X}_t, \widehat{\alpha}_t, \widehat{\zeta}_t)_{t \in [0, T]}$  is guaranteed, the FBSDE [\(26\)](#)-[\(27\)](#) has a solution with the backward component  $(Y, Z, \eta) = (Y_t, Z_t, \eta_t)_{t \in [0, T]}$  being the multipliers given in the last section. Then, by using [Corollary 3.10](#), we can conclude that, under [Assumption 1](#) and [Assumption 4](#), there exists a solution to the generalized type constrained FBSDE [\(26\)](#)-[\(27\)](#).

We now verify the constraints qualification condition as in (1.4) of [Zowe and Kurcyusz \[33\]](#) to establish the KKT optimality condition (i.e., we verify  $r_0 = 1$  in [Lemma 4.3](#)).

**Lemma 5.2.** Under the condition (C1) of [Assumption 3](#), the KKT optimality condition holds. More precisely, if  $(\widehat{X}, \widehat{\alpha}, \widehat{\zeta}) \in C := \mathbb{M}^2 \times \mathcal{U}[0, T] \times \mathcal{A}^k[0, T]$  is an optimal solution to the constrained MFC problem [\(3\)](#), there exists some  $\eta \in \mathbb{L}_{\mathbb{F}, \mathbb{R}_+}^2$  such that

$$(i) \mathbb{E} \left[ \int_0^T \phi(t, \widehat{X}_t, \mathcal{L}(\widehat{X}_t)) \eta_t dt \right] = 0;$$

(ii)  $\mathbb{P}(c_i(t) + G_i(t)^\top Y_t \geq 0, \forall (i, t) \in \{1, \dots, k\} \times [0, T]) = 1$ , where  $c_i(t)$  and  $G_i(t)$  are the  $i$ -th row of  $c(t)$  and  $G(t)$ , respectively. Moreover, it holds that

$$\mathbb{P} \left( \sum_{i=1}^k \mathbf{1}_{\{c_i(t) + G_i(t)^\top Y_t > 0\}} d\widehat{\zeta}_t^i = 0 \right) = 1. \quad (28)$$

Furthermore, the stochastic (first-order) minimum condition holds that, for any  $u \in U$  and  $dt \times d\mathbb{P}$ -a.s.,

$$\left\langle \nabla_u H(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t, Y_t, Z_t), u - \widehat{\alpha}_t \right\rangle \geq 0, \quad \forall u \in U. \quad (29)$$

Here, the Hamiltonian  $H(t, x, \mu, u, y, z) = H^1(t, x, \mu, u, y, z)$  for  $(t, x, \mu, u, y, z) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^l \times \mathbb{R}^n \times \mathbb{R}^{n \times r}$ , and the process pair  $(Y, Z) = (Y_t, Z_t)_{t \in [0, T]}$  taking values in  $\mathbb{R}^n \times \mathbb{R}^{n \times r}$  is the unique solution to the following general type of constrained BSDE:

$$\begin{aligned} Y_t &= \int_t^T \nabla_x H^{r_0}(s, \widehat{X}_s, \widehat{\mu}_s, \widehat{\alpha}_s, Y_s, Z_s) ds + \int_t^T \mathbb{E}' \left[ \partial_\mu H(s, \widehat{X}'_s, \widehat{\mu}'_s, \widehat{\alpha}'_s, Y'_s, Z'_s)(\widehat{X}_s) \right] ds \\ &\quad - \int_t^T \nabla_x \phi(s, \widehat{X}_s, \widehat{\mu}_s) \eta_s ds - \int_t^T \mathbb{E}' \left[ \partial_\mu \phi(s, \widehat{X}'_s, \widehat{\mu}'_s)(\widehat{X}_s) \eta'_s \right] ds - \int_t^T Z_s dW_s. \end{aligned} \quad (30)$$

*Proof.* This is a direct consequence of Lemma 3.9 in [6] and Theorem 3.1 of Zowe and Jurcysz [33].  $\square$

The strict convexity imposed on the Hamiltonian  $H$  w.r.t. the control variable  $u \in U$  given in (C3) of [Assumption 3](#) yields the result that once  $\langle \nabla_u H(t, \widehat{X}_t, \widehat{\mu}_t, \widehat{\alpha}_t, Y_t, Z_t), u - \widehat{\alpha}_t \rangle \geq 0$  for all  $u \in U$ , we must have  $\widehat{\alpha}_t = \hat{u}(t, \widehat{X}_t, \widehat{\mu}_t, Y_t, Z_t)$ . Then, we have

**Theorem 5.3** (Uniqueness of solutions to FBSDE). *Let [Assumption 2](#) and [Assumption 3](#) hold. The constrained FBSDE (26)-(27) has a unique solution in the sense of [Definition 5.1](#).*

*Proof.* Let  $(X_t^i, \zeta_t^i, Y_t^i, Z_t^i, \eta_t^i)_{t \in [0, T]}$  ( $i = 1, 2$ ) be two solutions to the FBSDE (26)-(27) satisfying  $X_0^1 = X_0^2$  and  $Y_T^1 = Y_T^2$  and set  $\bar{X}_t := X_t^1 - X_t^2$ ,  $\bar{\zeta}_t := \zeta_t^1 - \zeta_t^2$ ,  $\bar{Y}_t := Y_t^1 - Y_t^2$ ,  $\bar{Z}_t := Z_t^1 - Z_t^2$  and  $\bar{\eta}_t := \eta_t^1 - \eta_t^2$  for  $t \in [0, T]$ . Applying Itô's formula to  $\bar{X}_t \bar{Y}_t$ , we arrive at

$$\begin{aligned} \bar{X}_T \bar{Y}_T &= \int_0^T \bar{X}_t d\bar{Y}_t + \int_0^T \bar{Y}_t d\bar{X}_t + \langle \bar{X}, \bar{Y} \rangle_T \\ &= \int_0^T \{ -\Delta \bar{F}(t) \bar{X}_t + \bar{Y}_t \Delta \bar{b}(t) + \text{tr}(\bar{Z}_t \Delta \bar{\sigma}(t)) \} dt + \int_0^T (\bar{X}_t \bar{Z}_t + \bar{Y}_t \Delta \sigma(t)) dW_t \\ &\quad + \int_0^T (A(t) \bar{\eta}_t + B(t) \mathbb{E}[\bar{\eta}_t]) \bar{X}_t dt + \int_0^T G(t) \bar{Y}_t d\bar{\zeta}_t, \end{aligned}$$

where  $\Delta \bar{F}(t) := \bar{F}(t, \widehat{X}_t^1, \mathcal{L}(\widehat{X}_t^1), Y_t^1, Z_t^1) - \bar{F}(t, \widehat{X}_t^2, \mathcal{L}(\widehat{X}_t^2), Y_t^2, Z_t^2)$ ,  $\Delta \bar{b}(t) := \bar{b}(t, \widehat{X}_t^1, \mathcal{L}(\widehat{X}_t^1), Y_t^1, Z_t^1) - \bar{b}(t, \widehat{X}_t^2, \mathcal{L}(\widehat{X}_t^2), Y_t^2, Z_t^2)$  and  $\Delta \bar{\sigma}(t) := \bar{\sigma}(t, X_t^1, \mathcal{L}(X_t^1), Y_t^1, Z_t^1) - \bar{\sigma}(t, X_t^2, \mathcal{L}(X_t^2), Y_t^2, Z_t^2)$ . Taking expectations on both sides and noticing  $\bar{Y}_T = 0$ , we get

$$\begin{aligned} 0 &= \mathbb{E} \left[ \int_0^T \{ (\Phi(t, X_t^1, \mathcal{L}(X_t^1), Y_t^1, Z_t^1) - \Phi(t, X_t^2, \mathcal{L}(X_t^1), Y_t^2, Z_t^2))(X_t^1 - X_t^2, Y_t^1 - Y_t^2, Z_t^1 - Z_t^2) \right. \\ &\quad \left. + (\Phi(t, X_t^2, \mathcal{L}(X_t^1), Y_t^2, Z_t^2) - \Phi(t, X_t^2, \mathcal{L}(X_t^2), Y_t^2, Z_t^2))(X_t^1 - X_t^2, Y_t^1 - Y_t^2, Z_t^1 - Z_t^2) \right\} dt \\ &\quad \left. + \int_0^T \bar{\eta}_t \{ a(t) \bar{X}_t + b(t) \mathbb{E}[\bar{X}_t] \} dt + \int_0^T G(t) \bar{Y}_t d\bar{\zeta}_t \right]. \end{aligned}$$

Here, we have applied Fubini's theorem in the last line. Note that, it holds that

$$\begin{aligned} \mathbb{E} \left[ \int_0^T \bar{\eta}_t \{A(t)\bar{X}_t + B(t)\mathbb{E}[\bar{X}_t]\} dt \right] &= \mathbb{E} \left[ \int_0^T \bar{\eta}_t \{ \phi(t, X_t^1, \mathcal{L}(X_t^1)) - \phi(t, X_t^2, \mathcal{L}(X_t^2)) \} dt \right] \\ &= -\mathbb{E} \left[ \int_0^T \{ \phi(t, X_t^1, \mathcal{L}(X_t^1))\eta_t^2 + \phi(t, X_t^2, \mathcal{L}(X_t^2))\eta_t^1 \} dt \right] \leq 0, \end{aligned}$$

where we have used the fact that  $\mathbb{E}[\int_0^T \phi(t, X_t^i, \mathcal{L}(X_t^i))\eta_t^i dt] = 0$ ,  $\phi_i(t, X_t^i, \mathcal{L}(X_t^i)) \geq 0$  and  $\eta_t^i \geq 0$  for  $i = 1, 2$ . Similarly, we have  $\mathbb{E}[\int_0^T G(t)\bar{Y}_t d\bar{\zeta}_t] \leq 0$ . Exploiting [Assumption 3](#), it holds that

$$\begin{aligned} 0 \leq \mathbb{E} \left[ (\Phi(t, X_t^2, \mathcal{L}(X_t^1), Y_t^2, Z_t^2) - \Phi(t, X_t^2, \mathcal{L}(X_t^2), Y_t^2, Z_t^2)) (\bar{X}_t, \bar{Y}_t, \bar{Z}_t) \right. \\ \left. - \beta (|\bar{X}_t|^2 + |\bar{Y}_t|^2 + |\bar{Z}_t|^2) \right]. \end{aligned}$$

However, it follows from (A3) in [Assumption 1](#) that

$$\begin{aligned} &(\Phi(t, X_t^2, \mathcal{L}(X_t^1), Y_t^2, Z_t^2) - \Phi(t, X_t^2, \mathcal{L}(X_t^2), Y_t^2, Z_t^2)) (\bar{X}_t, \bar{Y}_t, \bar{Z}_t) \\ &\leq \frac{M}{2} \mathcal{W}_2(\mathcal{L}(X_t^1), \mathcal{L}(X_t^2)) (|X_t^1 - X_t^2| + |Y_t^1 - Y_t^2| + |Z_t^1 - Z_t^2|) \\ &\leq \frac{M}{2} (|X_t^1 - X_t^2|^2 + |Y_t^1 - Y_t^2|^2 + |Z_t^1 - Z_t^2|^2). \end{aligned}$$

Hence, combining the above two inequalities leads to  $\bar{X} = 0, \bar{Y} = 0$  and  $\bar{Z} = 0$ . It then follows that  $\bar{\alpha} = 0$  and  $\int_0^T G(s)d\bar{\zeta}_s = 0$ . Moreover, we have

$$\int_0^t A(s)\bar{\eta}_s ds + \int_0^t B(s)\mathbb{E}[\bar{\eta}_s] ds = 0, \quad \forall t \in [0, T].$$

Taking expectations on both sides of the above result, together with (C1) in [Assumption 3](#), we conclude that  $\mathbb{E}[\bar{\eta}_t] = 0$ , for  $m$ -a.e.  $t \in [0, T]$ , and hence

$$\int_0^t A(s)\bar{\eta}_s ds = 0, \quad \forall m\text{-a.e. } t \in [0, T],$$

which indicates that  $\bar{\eta} = 0$  in  $\mathbb{L}_{\mathbb{F}}^2$  by utilizing (C1) of [Assumption 3](#).  $\square$

**Remark 5.4.** Here, we only consider the case when  $Y_T = g(X_T, \mu_T)$  with  $g = 0$  in the constrained FBSDE (26)-(27). However, the uniqueness result still holds if we additionally assume:

(C5) Set  $\Psi(x, \mu) := \nabla_x g(x, \mu) + \int_{\mathbb{R}^n} \partial_\mu g(x', \mu)(x)\mu(dx')$  for  $(x, \mu) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$ . Then,  $\Psi$  satisfies the monotone condition  $(\Psi(x_1, \mu) - \Psi(x_2, \mu))(x_1 - x_2) \geq 0$  and  $\int_{\mathbb{R}^n} (\Psi(x, \mu_1) - \Psi(x, \mu_2))x'(\mu_1 - \mu_2)(dx') \geq 0$  for  $x, x_1, x_2 \in \mathbb{R}^n$  and  $\mu, \mu_1, \mu_2 \in \mathcal{P}_2(\mathbb{R}^n)$ .

Finally, we examine the stability of the constrained FBSDE (26)-(27). To this end, we introduce the following enhanced conditions:

(C1') The constrained function  $\phi$  is independent of the control variable and admits the linear form that, for  $(t, x, \mu) \in [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$ ,

$$\phi(t, x, \mu) = A(t)x + B(t) \int_{\mathbb{R}^n} x\mu(dx) + C(t),$$

where the coefficients  $A(t), B(t), C(t) \in C([0, T], \mathbb{R}^n)$  satisfying  $|A(t)| \geq c > 0$  and  $|A(t) + B(t)| \geq c > 0$  for some  $c > 0$ .

Then, we have



**Theorem 5.5.** Let *Assumption 2*, (C1'), (C3)-(C4) of *Assumption 3* hold. Assume that  $\xi_1, \xi_2 \in L^2((\Omega, \mathcal{F}_0, \mathbb{P}); \mathbb{R}^n)$  and denote by  $(\bar{X}^i, \bar{\alpha}_t^i, \bar{\zeta}_t^i, Y^i, Z^i, \eta^i)$  the unique solutions to the FBSDE (26)-(27) with respective initial data  $\xi_i$ . Set  $\bar{X}_t := X_t^1 - X_t^2$ ,  $\bar{\zeta}_t := \zeta_t^1 - \zeta_t^2$ ,  $\bar{Y}_t := Y_t^1 - Y_t^2$ ,  $\bar{Z}_t := Z_t^1 - Z_t^2$  and  $\bar{\eta}_t := \eta_t^1 - \eta_t^2$  for  $t \in [0, T]$ . Then, there exists a constant  $C > 0$  depending on  $(T, M, \alpha, \beta)$  only such that

$$\mathbb{E} \left[ |\bar{X}_T|^2 + \int_0^T (|\bar{X}_t|^2 + |\bar{\alpha}_t|^2 + |\bar{Y}_t|^2 + |\bar{Z}_t|^2 + |\bar{\eta}_t|^2) dt + \left| \int_0^T G(t) d\bar{\zeta}_t \right|^2 \right] \leq \mathbb{E} [|\xi_1 - \xi_2|^2] \quad (31)$$

*Proof.* Recall notations  $\Delta b(t), \Delta \sigma(t), \Delta F(t)$  defined in the proof of [Theorem 5.3](#). Then, one has

$$\int_0^t G(s) d\bar{\zeta}_s = \bar{X}_t - (\xi_1 - \xi_2) - \int_0^t \Delta b(s) ds - \int_0^t \Delta \sigma(s) ds. \quad (32)$$

On the other hand, applying Itô's formula to  $|\bar{X}_t|^2$  leads to

$$d|\bar{X}_t|^2 = 2 \left( \Delta b(t)^\top \bar{X}_t + \text{tr} \left( \Delta \sigma \Delta \sigma^\top(t) \right) \right) dt + 2\bar{X}_t^\top \Delta \sigma(t) dW_t + 2\bar{X}_t^\top G(t) d\bar{\zeta}_t. \quad (33)$$

It follows from the Young's inequality that, for any  $\epsilon \in (0, 1)$ ,

$$\left| \int_0^t 2\bar{X}_s^\top G(s) d\bar{\zeta}_s \right| \leq \epsilon \sup_{0 \leq s \leq t} |\bar{X}_s|^2 + \frac{1}{\epsilon} \left| \int_0^t G(s) d\bar{\zeta}_s \right|^2.$$

Plugging the above inequality into the equality (33), and taking expectations on both sides, we arrive at

$$\begin{aligned} \mathbb{E} \left[ \sup_{0 \leq s \leq t} |\bar{X}_s|^2 \right] &\leq \mathbb{E} \left[ \sup_{0 \leq s \leq t} \int_0^s 2 \left( \Delta b(s)^\top \bar{X}_s + \text{tr} \left( \Delta \sigma \Delta \sigma^\top(s) \right) \right) ds \right] + \mathbb{E} \left[ \sup_{0 \leq s \leq t} \int_0^s 2\bar{X}_s^\top \Delta \sigma(s) dW_s \right] \\ &\quad + \epsilon \mathbb{E} \left[ \sup_{0 \leq s \leq t} |\bar{X}_s|^2 \right] + \frac{1}{\epsilon} \mathbb{E} \left[ \left| \int_0^t G(s) d\bar{\zeta}_s \right|^2 \right]. \end{aligned}$$

In view of (C4) of [Assumption 3](#), the equality (32), the BDG's inequality and the Gronwall's inequality, we deduce the existence of a constant  $C > 0$  depending on  $M, \epsilon, T$  only such that

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} |\bar{X}_t|^2 \right] \leq C \mathbb{E} \left[ |\xi^1 - \xi^2|^2 + \int_0^T (|\bar{Y}_t|^2 + |\bar{Z}_t|^2) dt \right]. \quad (34)$$

With the help of the BSDE in (26), we have that, for all  $t \in [0, T]$ ,

$$\int_t^T A(s) \bar{\eta}_s ds + \int_t^T B(s) \mathbb{E} [\bar{\eta}_s] ds = \bar{Y}_T - Y_t + \int_t^T \Delta F(s) ds - \int_t^T \bar{Z}_s dW_s. \quad (35)$$

Applying the Itô's formula to  $|\bar{Y}_t|^2$ , we deduce that, for all  $t \in [0, T]$ ,

$$\begin{aligned} |\bar{Y}_t|^2 + \int_t^T |\bar{Z}_s|^2 ds &= 2 \int_t^T \bar{Y}_s^\top \Delta F(s) ds - 2 \int_t^T \bar{Y}_s \bar{Z}_s dW_s \\ &\quad - 2 \int_t^T \bar{\eta}_s A(s)^\top \bar{Y}_s ds - 2 \int_t^T \mathbb{E} [\bar{\eta}_s] B(s)^\top \bar{Y}_s ds. \end{aligned}$$

It follows from the Young's inequality again that, for all  $t \in [0, T]$ ,

$$\left| 2 \int_t^T \bar{\eta}_s A(s)^\top \bar{Y}_s ds \right| \leq \epsilon \sup_{t \leq s \leq T} |\bar{Y}_s|^2 + \frac{1}{\epsilon} \left| \int_t^T A(s)^\top \bar{\eta}_s ds \right|^2,$$

$$\left| 2 \int_t^T \mathbb{E} [\bar{\eta}_s] B(s)^\top \bar{Y}_s ds \right| \leq \epsilon \sup_{t \leq s \leq T} |\bar{Y}_s|^2 + \frac{1}{\epsilon} \left| \int_t^T B(s)^\top \mathbb{E} [\bar{\eta}_s] ds \right|^2.$$

In view of the condition (C4) of [Assumption 3](#), the condition (C1'), the BDG's inequality and the Gronwall's inequality again, we can similarly derive that

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} |\bar{Y}_t|^2 + \int_0^T |\bar{Z}_t|^2 dt \right] \leq C \mathbb{E} \left[ |\bar{X}_T|^2 + \int_0^T |\bar{X}_t|^2 dt \right], \quad (36)$$

for some constant  $C > 0$  depending on  $M, \epsilon, T$  only. Applying integration by parts to  $\bar{X}_t^\top \bar{Y}_t$ , we have

$$\begin{aligned} \bar{X}_T \bar{Y}_T &= \int_0^T (-\Delta \bar{F}(t) \bar{X}_t + \bar{Y} \Delta \bar{b}(t) + \text{tr}(\bar{Z}_t \Delta \bar{\sigma}(t))) dt + \int_0^T (\bar{X}_t \bar{Z}_t + \bar{Y}_t \Delta \sigma) dW_t \\ &\quad + \int_0^T \{A(t) \bar{\eta}_t + B(t) \mathbb{E}[\bar{\eta}_t]\} \bar{X}_t + \int_0^T G(t) \bar{Y}_t d\bar{\zeta}_t. \end{aligned}$$

Taking expectations on both sides of the above equality and following the proof of [Theorem 5.3](#), we deduce that

$$-\mathbb{E} \left[ \bar{Y}_0^\top (\xi^1 - \xi^2) \right] + \beta \mathbb{E} \left[ \int_0^T (|\bar{X}_t|^2 + |\bar{Y}_t|^2 + |\bar{Z}_t|^2) dt \right] \leq 0. \quad (37)$$

As a result, the inequality (37) leads to

$$\begin{aligned} \alpha \mathbb{E} \left[ |\bar{X}_T|^2 \right] + \beta \mathbb{E} \left[ \int_0^T (|\bar{X}_t|^2 + |\bar{Y}_t|^2 + |\bar{Z}_t|^2) dt \right] &\leq \mathbb{E} \left[ \bar{Y}_0^\top (\xi^1 - \xi^2) \right] \\ &\leq \frac{2}{\epsilon} \mathbb{E} \left[ |\bar{Y}_0|^2 \right] + \frac{1}{2\epsilon} \mathbb{E} \left[ |\xi^1 - \xi^2|^2 \right]. \end{aligned}$$

Choosing  $\epsilon > 0$  small enough and taking into account of (34) and (36), we can find a constant  $C > 0$  depending on  $M, T, \alpha, \beta$  only such that

$$\mathbb{E} \left[ |\bar{X}_T|^2 + \int_0^T (|\bar{X}_t|^2 + |\bar{Y}_t|^2 + |\bar{Z}_t|^2) dt \right] \leq C \mathbb{E} \left[ |\xi^1 - \xi^2|^2 \right].$$

Combining the Lipschitz continuity of  $\hat{u} : [0, T] \times \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^n \times \mathbb{R}^{n \times r} \mapsto U \subset \mathbb{R}^l$  in the condition (C3) of [Assumption 3](#), (32), (35) and the condition (C1'), we conclude the stability (31) from (38) above.  $\square$

**Remark 5.6.** When considering  $Y_T = g(X_T, \mu_T)$ , we further assume

(C5') Set  $\Psi(x, \mu) := \nabla_x g(x, \mu) + \int_{\mathbb{R}^n} \partial_\mu g(y, \mu)(x) \mu(dy)$  for  $(x, \mu) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$ . The function  $\Psi$  is Lipschitz continuous and satisfies the coercivity condition that, for some  $\alpha, M > 0$  and  $(x^1, \mu^1, X^1), (x^2, \mu^2, X^2) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times L^2((\Omega, \mathcal{F}_T, \mathbb{P}), \mathbb{R}^n)$ ,

$$\begin{aligned} |\Psi(x^1, \mu^1) - \Psi(x^2, \mu^2)| &\leq M(|x^1 - x^2| + \mathcal{W}_{2, \mathbb{R}^n}(\mu^1, \mu^2)), \\ \mathbb{E} [\langle \Psi(X^1, \mathcal{L}(X^1)) - \Psi(X^2, \mathcal{L}(X^2)), X^1 - X^2 \rangle] &\geq \alpha \mathbb{E} [|X^1 - X^2|^2]. \end{aligned}$$

In addition to the setting in [Theorem 5.5](#), we let  $Y^i$  possess terminal value  $g^i(X_T^i, \mu_T^i)$  and  $g^1, g^2 \in C^{1;1}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n); \mathbb{R})$  with  $g^1$  satisfying the condition (C5'). Then, it holds that

$$\begin{aligned} \mathbb{E} \left[ |\bar{X}_T|^2 + \int_0^T (|\bar{X}_t|^2 + |\bar{\alpha}_t|^2 + |\bar{Y}_t|^2 + |\bar{Z}_t|^2 + |\bar{\eta}_t|^2) dt + \left| \int_0^T G(t) d\bar{\zeta}_t \right|^2 \right] \\ \leq C \left\{ \mathbb{E} [|\xi_1 - \xi_2|^2] + \sup_{(x, \mu) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)} |\Psi^1(x, \mu) - \Psi^2(x, \mu)|^2 \right\} \quad (38) \end{aligned}$$

with  $\Psi^i(x, \mu) := \nabla_x g^i(x, \mu) + \int_{\mathbb{R}^n} \partial_\mu g^i(y, \mu)(x) \mu(dy)$  for  $i = 1, 2$ .

**Acknowledgements** L. Bo and J. Wang are supported by National Natural Science of Foundation of China (No. 12471451), Natural Science Basic Research Program of Shaanxi (No. 2023-JC-JQ-05), Shaanxi Fundamental Science Research Project for Mathematics and Physics (No. 23JSZ010). X. Yu and J. Wang are supported by the Hong Kong Polytechnic University research grant under No. P0045654.

## References

- [1] B. Acciaio, J. Backhoff-Veraguas, and R. Carmona (2017): Extended mean field control problems: stochastic maximum principle and transport perspective. *SIAM J. Contr. Optim.* 57(6): 3666-3693.
- [2] D. Andersson, and B. Djehiche (2010): A maximum principle for SDEs of mean-field type. *Appl. Math. Optim.* 63: 341-356.
- [3] S. Bahlali, and B. Mezerdi (2005): A general stochastic maximum principle for singular control problems. *Electron. J. Probab.* 10: 988-1004.
- [4] E. Bayraktar, A. Cosso, and H. Pham (2018): Randomized dynamic programming principle and Feynman-Kac representation for optimal control of McKean-Vlasov dynamics. *Trans. AMS* 370(3): 2115-60.
- [5] L. Bo, J. Wang, X. Wei, and X. Yu (2024): Extended mean-field control problems with Poissonian common noise: Stochastic maximum principle and Hamiltonian-Jacobi-Bellman equation. Preprint available at <https://arxiv.org/abs/2407.05356>
- [6] L. Bo, J. Wang, and X. Yu (2024): Extended mean field control problems with constraints: The generalized Fritz-John conditions and Lagrangian method. Preprint available at: <https://arxiv.org/abs/2408.06865>
- [7] R. Buckdahn, B. Djehiche, and J. Li (2011): A general maximum principle for SDEs of mean-field type. *Appl. Math. Optim.* 64(2): 197-216.
- [8] L. Campi, T. De Angelis, M. Ghio, and G. Livieri (2022): Mean-field games of finite-fuel capacity expansion with singular controls. *Ann. Appl. Probab.* 32(5): 3674-3717.
- [9] H. Cao, J. Dianetti and G. Ferrari (2023) Stationary discounted and ergodic mean field games with singular controls. *Math. Oper. Res.* 48(4):1871-1898.
- [10] R. Carmona, and F. Delarue (2015): Forward-backward stochastic differential equations and controlled McKean-Vlasov dynamics. *Ann. Probab.* 43(5): 2647-2700.
- [11] M. Chiarolla, G. Ferrari, and F. Riedel (2013): Generalized Kuhn-Tucker conditions for N-firm stochastic irreversible investment under limited resources. *SIAM J. Contr. Optim.* 51(5): 3863-3885.
- [12] S. Daudin (2023a): Mean-field limit for stochastic control problems under state constraint. Preprint available at: <https://arxiv.org/abs/2306.00949>
- [13] S. Daudin (2023b): Optimal control of the Fokker-Planck equation under state constraints in the Wasserstein space. *J. Math. Pures Appl.* 175: 37-75.
- [14] R. Denkert, and U. Horst (2023): Extended mean-field control problems with multi-dimensional singular controls. Preprint available at <https://arxiv.org/abs/2308.04378>

- [15] M.F. Djete, D. Possamaï, and X. Tan (2022): McKean-Vlasov optimal control: limit theory and equivalence between different formulations. *Math. Oper. Res.* 47(4): 2891-930.
- [16] N. El Karoui, and S. Méléard (1990): Martingale measures and stochastic calculus. *Probab. Theor. Rel. Fields* 84: 83-101.
- [17] G. Fu, and U. Horst (2017): Mean field games with singular controls. *SIAM J. Contr. Optim.* 55(6): 3833-3868.
- [18] M. Germain, H. Pham, and X. Warin (2023): A level-set approach to the control of state-constrained McKean-Vlasov equations: Applications to renewable energy storage and portfolio selection. *Numer. Algebra Contr. Optim.* 13: 555-582.
- [19] X. Guo, and P. Tomecek (2009): A class of singular control problems and the smooth fit principle. *SIAM J. Contr. Optim.* 47(6): 3076-3099.
- [20] U.G. Haussmann, and W. Suo (1995): Singular optimal stochastic controls I: Existence. *SIAM J. Contr. Optim.* 33: 916-936.
- [21] U.G. Haussmann, and W. Suo (1995b): Singular optimal stochastic controls II: Dynamic programming. *SIAM J. Contr. Optim.* 33(3): 937-959
- [22] M. Huang, R.P. Malhamé, and P.E. Caines (2006): Large population stochastic dynamic games closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle. *Commun. Inform. Syst.* 6(3): 221-252.
- [23] J. Jacod, and J. Mémin (1981): Sur un type de convergence intermédiaire entre la convergence en loi et la convergence en probabilité, *Séminaire de probabilités de Strasbourg* 15: 529-546.
- [24] D. Lacker (2015): Mean field games via controlled martingale problems: Existence of Markovian equilibria. *Stoch. Process. Appl.* 125: 2856-2894.
- [25] D. Lacker (2017): Limit theory for controlled McKean-Vlasov dynamics. *SIAM J. Contr. Optim.* 55(3): 1641-1672.
- [26] J.M. Lasry, and P.L. Lions (2007): Mean field games. *Jpn. J. Math.* 2(1): 229-260.
- [27] M. Laurière, and O. Pironneau (2014): Dynamic programming for mean-field type control. *Comptes Rendus Math.* 352(9): 707-713.
- [28] S.J. Leese (1978): Measurable selections and the uniformization of Souslin sets. *Amer. J. Math.* 100: 19-41.
- [29] J. Ma (1992): On the principle of smooth fit for a class of singular stochastic control problems for diffusions. *SIAM J. Contr. Optim.* 30(4): 975-999.
- [30] H. Pham, and X. Wei (2017): Dynamic programming for optimal control of stochastic McKean-Vlasov dynamics. *SIAM J. Contr. Optim.* 55(2): 1069-1101.
- [31] C. Wu, and J. Zhang (2020): Viscosity solutions to parabolic master equations and McKean-Vlasov SDEs with closed-loop controls. *Ann. Appl. Probab.* 30(2): 936-986.
- [32] J. Zhou, N. Touzi, and J. Zhang (2024): Viscosity solutions for HJB equations on the process space: Application to mean field control with common noise. Preprint available at: <https://arxiv.org/abs/2401.04920>
- [33] J. Zowe, and S. Kurcyusz (1979): Regularity and stability for the mathematical programming problem in Banach spaces. *Appl. Math. Optim.* 5: 49-62.