

Singular leaning coefficients and efficiency in learning theory

Miki Aoyagi

College of Science & Technology, Nihon University, 1-8-14, Surugadai, Kanda, Chiyoda-ku,
Tokyo 101-8308 Japan. (aoyagi.miki@nihon-u.ac.jp)

Abstract :

Singular learning models with non-positive Fisher information matrices include neural networks, reduced-rank regression, Boltzmann machines, normal mixture models, and others. These models have been widely used in the development of learning machines. However, theoretical analysis is still in its early stages.

In this paper, we examine learning coefficients, which indicate the general learning efficiency of deep linear learning models and three-layer neural network models with ReLU units. Finally, we extend the results to include the case of the Softmax function.

Keyword : resolution map, singular learning theory, multiple-layered neural networks with linear units, ReLU units, algebraic geometry.

1 Introduction

In learning theory, the log canonical threshold in algebraic geometry plays an important role in analyzing the efficiency of learning machines[19].

Assume that each sample (x_i, y_i) is drawn from a probability density function $q(x, y)$ and $(x, y)^n := \{(x_i, y_i)\}_{i=1}^n$ are n training samples selected independently and identically from $q(x, y)$. To estimate the true probability density function $q(x, y)$ using $(x, y)^n$ within the framework of Bayesian estimation, we consider a learning model expressed in probabilistic form as $p(x, y|w)$, along with an *a priori* probability density function $\psi(w)$ on a compact parameter set W , where $w \in W \subset \mathbf{R}^d$ is a parameter. The *a posteriori* probability density function $p(w|(x, y)^n)$ is then given by:

$$p(w|(x, y)^n) = \frac{1}{Z_n(\beta)} \psi(w) \prod_{i=1}^n p(x_i, y_i|w)^\beta,$$

where

$$Z_n(\beta) = \int_W \psi(w) \prod_{i=1}^n p(x_i, y_i|w)^\beta dw,$$

with β representing an inverse temperature.

The Kullback-Leibler divergence $D(p_1 | p_2) = \int p_1(z) \log \frac{p_1(z)}{p_2(z)} dz$ is a pseudo-distance between arbitrary probability density functions $p_1(z)$ and $p_2(z)$.

The optimal parameter w_0 minimizes

$$\begin{aligned} L(w) &= -E_{x,y}[\log p(x, y|w)] = \int q(x, y) \log \frac{q(x, y)}{p(x, y|w)} dx dy - \int q(x, y) \log q(x, y) dx dy \\ &= D(q(x, y) | p(x, y|w)) - \int q(x, y) \log q(x, y) dx dy \end{aligned}$$

and thus $D(q(x, y) | p(x, y|w))$ at $w = w_0$, and let

$$W_0 = \{w \in W | L(w) = \min_{w' \in W} L(w')\}.$$

Assume that its log likelihood function has relatively finite variance,

$$E_{x,y}[\log \frac{p(x, y|w_0)}{p(x, y|w)}] \geq c E_{x,y}[(\log \frac{p(x, y|w_0)}{p(x, y|w)})^2], \quad w_0 \in W_0, w \in W,$$

for a constant $c > 0$. Then, we have a unique probability density function $p_0(x, y) = p(x, y|w_0)$ for all $w_0 \in W_0$.

Let

$$f(x, y|w) = \log \frac{p_0(x, y)}{p(x, y|w)}$$

and define its average error function as

$$K(w) = E_{x,y}[f(x, y|w)].$$

It is clear that $K(w_0) = 0$ for all $w_0 \in W_0$.

By applying Hironaka's Theorem [13] to the function $K(w)$ at w_0 , we obtain a proper analytic map π from a manifold Y to a neighborhood of $w_0 \in W_0$:

$$K(\pi(u)) = u_1^{2k_1} u_2^{2k_2} \cdots u_d^{2k_d},$$

where (u_1, \dots, u_d) is a local analytic coordinate system on $U \subset Y$. Furthermore, there exist analytic functions $a(x, y|u)$ and $b(u) \neq 0$ such that:

$$f(x, y|\pi(u)) = u_1^{k_1} u_2^{k_2} \cdots u_d^{k_d} a(x, y|u),$$

and

$$\pi'(u)\psi(\pi(u)) = u_1^{h_1} u_2^{h_2} \cdots u_d^{h_d} b(u).$$

Let

$$\xi_n(u) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \{u_1^{k_1} u_2^{k_2} \cdots u_d^{k_d} - a(x_i, y_i|\pi(u))\},$$

then, we have an empirical process $K_n(\pi(u))$ such that

$$\begin{aligned} nK_n(\pi(u)) &= \sum_{i=1}^n f(x_i, y_i|\pi(u)) \\ &= n u_1^{2k_1} u_2^{2k_2} \cdots u_d^{2k_d} - \sqrt{n} u_1^{k_1} u_2^{k_2} \cdots u_d^{k_d} \xi_n(u). \end{aligned}$$

We introduce learning coefficients

$$\lambda(w_0) = \min_u \min_{1 \leq j \leq d} \frac{h_j + 1}{2k_j},$$

and its order

$$\theta(w_0) = \max_u \text{Card}(\{j : \frac{h_j + 1}{2k_j} = \lambda\}),$$

where $\text{Card}(S)$ denotes the cardinality of a set S . Without loss of generality, we can assume that

$$\lambda(w_0) = \frac{h_1 + 1}{2k_1} = \frac{h_2 + 1}{2k_2} = \cdots = \frac{h_\theta + 1}{2k_\theta} < \frac{h_j + 1}{2k_j} \quad (\theta + 1 \leq j \leq d).$$

In Bayesian estimation, the predictive probability density function of $(x, y)^n$ is given by:

$$p((x, y)^n) = Z_n(1) = \int \prod_{i=1}^n p(x_i, y_i | w) \varphi(w) dw.$$

According to Watanabe [19], for $w_0 \in W_0$, we have

$$\begin{aligned} & p((x, y)^n) \\ &= \prod_{i=1}^n p(x_i, y_i | w_0) \int \prod_{i=1}^n \frac{p(x_i, y_i | w)}{p(x_i, y_i | w_0)} \varphi(w) dw \\ &= \prod_{i=1}^n p_0(x_i, y_i) \left(\frac{(\log n)^{\theta(w_0)-1}}{n^{\lambda(w_0)}} \int_0^\infty dt \ t^{\lambda(w_0)-1} \exp(-t + \sqrt{t} \xi_n(u)) du^* + o_p\left(\frac{(\log n)^{\theta(w_0)-1}}{n^{\lambda(w_0)}}\right) \right), \end{aligned}$$

where $\mu_j = -2\lambda k_j + h_j$,

$$du^* = \frac{\prod_{i=1}^\theta \delta(u_i) \prod_{j=\theta+1}^d u_j^{\mu_j}}{(\theta-1)! \prod_{i=1}^\theta (2k_i)} b(u) du,$$

and $\delta(u)$ is Dirac's delta function.

This indicates that the most efficient parameter $w_0 \in W_0$ is the one with the smallest $\lambda(w_0)$ and the largest $\theta(w_0)$, where the Kullback-Leibler divergence between $q((x, y)^n) = \prod_{i=1}^n q(x_i, y_i)$ and $p((x, y)^n)$ is minimized:

$$\begin{aligned} D(q((x, y)^n) | p((x, y)^n)) &= \int q((x, y)^n) \log \frac{q((x, y)^n)}{p((x, y)^n)} \prod_{i=1}^n dx_i dy_i \\ &= \int q((x, y)^n) \log q((x, y)^n) \prod_{i=1}^n dx_i dy_i - \int q((x, y)^n) \log \prod_{i=1}^n p(x_i, y_i | w_0) \prod_{i=1}^n dx_i dy_i \\ &\quad + \int q((x, y)^n) \log \frac{\prod_{i=1}^n p(x_i, y_i | w_0)}{p((x, y)^n)} \prod_{i=1}^n dx_i dy_i \\ &= \int q((x, y)^n) \log q((x, y)^n) \prod_{i=1}^n dx_i dy_i - \int q((x, y)^n) \log \prod_{i=1}^n p_0(x_i, y_i) \prod_{i=1}^n dx_i dy_i \\ &\quad + \lambda(w_0) \log(n) - (\theta(w_0) - 1) \log \log(n) + O_p(1). \end{aligned}$$

Using these relations, we derive two model-selection methods: the "widely applicable Bayesian information criterion" (WBIC) [20] and the "singular Bayesian information criterion" (sBIC) [9].

The learning coefficients are known as log canonical thresholds in algebraic geometry. Theoretically, their values can be obtained using Hironaka's Theorem. However, these thresholds have been studied primarily over the complex field or algebraically closed fields in algebraic geometry and algebraic analysis [15, 16, 14]. There are significant differences between the real and complex fields. For instance, log canonical thresholds over the complex field are always less than one, while those over the real field are not necessarily so. Obtaining these thresholds for learning models is challenging due to several factors, such as degeneration with respect to their Newton polyhedra and the non-isolation of singularities [11]. As a result, determining these thresholds is a topic of interest across various disciplines, including mathematics.

Our purpose in this paper is to obtain λ and θ for deep-layered linear neural networks, and three-layer neural network models with ReLU units. Finally, we extend the results to include the case of the Softmax function. In recent studies, we obtained exact values or bounded values of the learning coefficients for Vandermonde matrix-type singularities, which are related to the three-layered neural networks and normal mixture models, among others [8, 1, 3, 5, 6]. We have also exact values for the restricted Boltzmann

machine [4]. Additionally, Rusakov and Geiger [17, 18] and Zwiernik [21], respectively, obtained the learning coefficients for naive Bayesian networks and directed tree models with hidden variables. Drton et al. [10] considered these coefficients for the Gaussian latent tree and forest models. The paper [12] empirically developed a method to obtain the local learning coefficient for deep linear networks.

2 Log canonical threshold

Definition 1 We denote constants by superscript $*$, for example, a^* , b^* , and w^* .

Let f be an analytic function in neighborhood U of w^* , and ψ be a C^∞ function on U that is also analytic in a neighborhood of w^* with compact support. Define the log canonical threshold

$$c_{w^*}(f, \psi) = \sup\{c : |f|^{-c} \text{ is locally } L^2 \text{ in a neighborhood of } w^*\}$$

over the complex field \mathbf{C} and

$$c_{w^*}(f, \psi) = \sup\{c : |f|^{-c} \text{ is locally } L^1 \text{ in a neighborhood of } w^*\}$$

over the real field \mathbf{R} .

Let $\theta_{w^*}(f, \psi)$ be its order.

If $\psi(w^*) \neq 0$, then denote $c_{w^*}(f) = c_{w^*}(f, \psi)$ and $\theta_{w^*}(f) = \theta_{w^*}(f, \psi)$ because the log canonical threshold and its order are independent of ψ .

For ideal I , generated by real analytic functions f_1, \dots, f_m in a neighborhood of w^* , define $c_{w^*}(I) = c_{w^*}(f_1^2 + \dots + f_m^2)$.

Here, $c_{w^*}(I)$ for ideal I is well-defined by Lemma 1.

Lemma 1 ([2]) Let U be a neighborhood of $w^* \in \mathbf{R}^d$. Consider the ring of analytic functions on U . Let \mathcal{J} be the ideal generated by f_1, \dots, f_n , which are analytic functions defined on U . If g_1, \dots, g_m generate ideal \mathcal{J} , then

$$c_{w^*}(f_1^2 + \dots + f_n^2) = c_{w^*}(g_1^2 + \dots + g_m^2).$$

The learning coefficient $\lambda(w^*)$ is the log canonical threshold of the average error function over the real field.

Define the norm of a matrix $C = (c_{ij})$ as $\|C\| = \sqrt{\sum_{i,j} |c_{ij}|^2}$.

Definition 2 For a matrix C , let $\langle C \rangle$ be the ideal generated by all elements of C .

Theorem 1 Let

$$h(x, w) = \sum_{0 \leq i_1, \dots, i_N \leq H} \tilde{h}_{i_1, \dots, i_N}(w) x_1^{i_1} \dots x_N^{i_N}$$

be a polynomial with variables x_1, \dots, x_N and $\tilde{h}_{i_1, \dots, i_N}(w)$'s are continuous functions. Let $q(x)$ be a continuous positive function on $X \subset \mathbf{R}^N$ with $\text{Vol}(X) = \int_X q(x) dx > 0$. Then, we have for some positive constants $\alpha_1, \alpha_2 > 0$ such that

$$\alpha_1 \sum_{0 \leq i_1, \dots, i_N \leq H} \tilde{h}_{i_1, \dots, i_N}^2(w) \leq K(w) = \int_X h^2(x, w) q(x) dx \leq \alpha_2 \sum_{0 \leq i_1, \dots, i_N \leq H} \tilde{h}_{i_1, \dots, i_N}^2(w).$$

(Proof)

Let \mathbf{v} be the vector whose elements are $\tilde{h}_{0,0,\dots,0}(w), \tilde{h}_{1,0,\dots,0}(w), \tilde{h}_{0,1,\dots,0}(w), \dots, \tilde{h}_{H,H,\dots,H}(w)$. Let C be the matrix whose (i, j) elements are $x_1^{i_1} x_2^{i_2} \dots x_N^{i_N} \cdot x_1^{j_1} x_2^{j_2} \dots x_N^{j_N}$, where the i th element of \mathbf{v} is $\tilde{h}_{i_1, \dots, i_N}(w)$ and the j th element of \mathbf{v} is $\tilde{h}_{j_1, \dots, j_N}(w)$.

We have $h^2(x, w) = \mathbf{v}^t C \mathbf{v} \geq 0$, and $K(w) = \int_X h^2(x, w) q(x) dx = \mathbf{v}^t \int_X C q(x) dx \mathbf{v} \geq 0$. Also we have $K(w) = 0$ if and only if $\mathbf{v} = 0$, and therefore $\int_X C q(x) dx$ is positive definite. By setting α_1 and α_2 as the maximum and minimum eigenvalues of C , respectively, we complete the proof.

(Q.E.D.)

In the above theorem, we assume that $h(x, w)$ is a polynomial in the variables x_1, \dots, x_N . However, by applying the Noetherian ring property, we can extend the result to analytic functions $h(x, w)$.

3 Multiple neural network with linear units.

In the paper [7], the learning coefficients for multiple-layered neural networks with linear units were obtained. In this section, we introduce the linear case with thresholds, which differs slightly from the case without thresholds.

Define matrices $A^{(s)}$ of size $H^{(s)} \times H^{(s+1)}$ and vectors $B^{(s)}$ of dimension $H^{(s)}$, for $s = 1, \dots, L$,

$$\begin{aligned} A^{(s)} &= (a_{ij}^{(s)}), & (1 \leq i \leq H^{(s)}, 1 \leq j \leq H^{(s+1)}), \\ B^{(s)} &= (b_j^{(s)}), & (1 \leq i \leq H^{(s)}). \end{aligned}$$

Definition 3 Let r be a natural number and $M^{(s)} = H^{(s)} - r$ for $s = 1, \dots, L + 1$. Define $\mathcal{M} \subset \{1, \dots, L + 1\}$ such that

$$\begin{aligned} \ell &= \text{Card}(\mathcal{M}) - 1, & \mathcal{M} &= \{S_1, \dots, S_{\ell+1}\}, \\ M^{(S_j)} &< M^{(s)} & \text{for } S_j \in \mathcal{M} \text{ and } s \notin \mathcal{M}, \\ \sum_{k=1}^{\ell+1} M^{(S_k)} &\geq \ell M^{(s)} & \text{for } s \in \mathcal{M}, \sum_{k=1}^{\ell+1} M^{(S_k)} < \ell M^{(s)} & \text{for } s \notin \mathcal{M}. \end{aligned}$$

Let M be the integer such that $M - 1 < \frac{\sum_{k=1}^{\ell+1} M^{(S_k)}}{\ell} \leq M$, and $a = \sum_{k=1}^{\ell+1} M^{(S_k)} - (M - 1)\ell$.

Let

$$\begin{aligned} &\lambda(H^{(1)}, H^{(2)}, \dots, H^{(L+1)}, r) \\ &= \frac{-r^2 + r(H^{(1)} + H^{(L+1)})}{2} + \frac{a(\ell - a)}{4\ell} + \frac{1}{4\ell} \left(\sum_{j=1}^{\ell+1} M^{(S_j)} \right)^2 - \frac{1}{4} \sum_{j=1}^{\ell+1} (M^{(S_j)})^2 \\ &= \frac{-r^2 + r(H^{(1)} + H^{(L+1)})}{2} + \frac{Ma + (M - 1) \sum_{j=1}^{\ell+1} M^{(S_j)}}{4} - \frac{1}{4} \sum_{j=1}^{\ell+1} (M^{(S_j)})^2 \end{aligned}$$

and $\theta(H^{(1)}, H^{(2)}, \dots, H^{(L+1)}, r) = a(\ell - a) + 1$.

Remark 1 If $\text{Card}\{s \mid M^{(s)} = 0\} \geq 1$, then we have $\lambda(H^{(1)}, H^{(2)}, \dots, H^{(L+1)}, r) = \frac{-r^2 + r(H^{(1)} + H^{(L+1)})}{2}$ and $\theta(H^{(1)}, H^{(2)}, \dots, H^{(L+1)}, r) = 1$.

Let W be the set of parameters

$$W = \{\{A^{(s)}\}_{1 \leq s \leq L}, \{B^{(s)}\}_{1 \leq s \leq L}\}.$$

Let $F^{(s)}(x) = A^{(s)}x + B^{(s)}$ be a function from $\mathbf{R}^{H^{(s+1)}}$ to $\mathbf{R}^{H^{(s)}}$.

Denote the input value by $x \in \mathbf{R}^{H^{(L+1)}}$ with probability density function $q(x)$ and output value $y \in \mathbf{R}^{H^{(1)}}$ for the multiple-layered neural network with linear units, which is given by

$$\begin{aligned} h(x, A, B) &= F^{(1)} \circ F^{(2)} \circ \dots \circ F^{(L)}(x) \\ &= \prod_{s=1}^L A^{(s)}x + \sum_{S=2}^L \prod_{s=1}^{S-1} A^{(s)}B^{(S)} + B^{(1)}. \end{aligned}$$

Consider the statistical model with Gaussian noise,

$$p(y|x, w) = \frac{1}{(\sqrt{2\pi})^{H^{(1)}}} \exp\left(-\frac{1}{2}\|y - h(x, A, B)\|^2\right),$$

$$p(x, y|w) = p(y|x, w)q(x).$$

The model has $H^{(1)}$ input units, $H^{(L+1)}$ output units, and $H^{(s)}$ hidden units in each hidden layer.

Define the average log loss function $L(w)$ by $L(w) = -E_{X,Y}[\log p(X, Y|w)]$ and assume the set of optimal parameters W_0 by

$$\begin{aligned} W_0 &= \{w_0 \in W | L(w_0) = \min_{w' \in W} L(w')\} \\ &= \{\{A^{(s)}\}_{1 \leq s \leq L}, \{B^{(s)}\}_{1 \leq s \leq L} | h(x, A, B) = h(x, A^*, B^*)\}. \end{aligned}$$

We have $p_0(x, y) = p(x, y|w_0)$ for all $w_0 \in W_0$.

Moreover, assume that the *a priori* probability density function $\varphi(w)$ is a C^∞ -function with compact support W , satisfying $\varphi(w^*) > 0$.

Let $K(w) = \int_X (h(x, A, B) - h(x, A^*, B^*))^2 q(x) dx$. Then, the log canonical threshold λ and its order θ are as follows.

Theorem 2 *Let r be the rank of $\prod_{s=1}^L A^{*(s)}$. We have*

$$\lambda = \frac{H^{(1)}}{2} + \lambda(H^{(1)}, H^{(2)}, \dots, H^{(L+1)}, r)$$

and

$$\theta = \theta(H^{(1)}, H^{(2)}, \dots, H^{(L+1)}, r).$$

(Proof)

By Theorem 1, we obtain

$$c_{w^*}(K(w)) = c_{w^*}(\|\prod_{s=1}^L A^{(s)} - \prod_{s=1}^L A^{*(s)}\|^2 + \|\sum_{S=2}^L \prod_{s=1}^{S-1} A^{(s)}B^{(S)} + B^{(1)} - \sum_{S=2}^L \prod_{s=1}^{S-1} A^{*(s)}B^{*(S)} - B^{*(1)}\|^2).$$

Since we can change the variables by

$$B'^{(1)} = \sum_{S=2}^L \prod_{s=1}^{S-1} A^{(s)}B^{(S)} + B^{(1)},$$

we obtain the theorem from Lemma 1 (2) and the Main Theorem in the paper [7].

(Q.E.D.)

4 Multiple-layered neural networks with activation function ReLU (Rectified Linear Unit function)

In this paper, we consider the case in multiple-layered neural networks with activation function ReLU (Rectified Linear Unit function).

For a matrix $A = (a_{ij})$, $a_{ij} \in \mathbf{R}$, define

$$A_+ = (\max\{0, a_{ij}\}).$$

Definition 4 *Let*

$$V^{(L)}(F_{i_1}^{(L)}, \dots, F_{i_k}^{(L)}) = \{x \mid F_{i_1}^{(L)}(x) > 0, \dots, F_{i_k}^{(L)}(x) > 0, \quad F_j^{(L)}(x) \leq 0 \text{ for } j \neq i_1, \dots, i_k\}$$

and

$$\Omega^{(L)} = \{V \mid V = V^{(L)}(F_{i_1}^{(L)}, \dots, F_{i_k}^{(L)}) \text{ for some } 1 \leq i_1, i_2, \dots, i_k \leq H^{(L)}\}.$$

Inductively, we define

$$\begin{aligned} V^{(s)}(F_{i_1}^{(s)}, \dots, F_{i_k}^{(s)}) &= \{x \in V^{(s+1)} \in \Omega^{(s+1)} \mid F_{i_1}^{(s)} \circ F^{(s+1)} \circ \dots \circ F^{(L)}(x) > 0, \\ &\quad \dots, F_{i_k}^{(s)} \circ F^{(s+1)} \circ \dots \circ F^{(L)}(x) > 0, \\ &\quad F_j^{(s)} \circ F^{(s+1)} \circ \dots \circ F^{(L)}(x) \leq 0 \text{ for } j \neq i_1, \dots, i_k\} \end{aligned}$$

and

$$\Omega^{(s)} = \{V \mid V = V^{(s)}(F_{i_1}^{(s)}, \dots, F_{i_k}^{(s)}) \text{ for some } 1 \leq i_1, i_2, \dots, i_k \leq H^{(s)}\}.$$

By the definition, we have $V_1 \cap V_2 = \emptyset$ for $V_1, V_2 \in \Omega^{(s)}$.

Let $h_+(x, A, B) = F_+^{(1)} \circ F_+^{(2)} \circ \dots \circ F_+^{(L)}(x)$.

If for all $x \in X$, the i th element of $F_+^{(s)} \circ \dots \circ F_+^{(L)}(x)$ is zero in a neighborhood of

$$\{A^{*(s)}, A^{*(s+1)}, \dots, A^{*(L)}, B^{*(s)}, B^{*(s+1)}, \dots, B^{*(L)}\},$$

then, we can remove the i th row of $A^{(s)}$, the i th element of $B^{(s)}$, and the i th column of $A^{(s-1)}$. This removal corresponds to eliminating the i th unit in the s th hidden layer.

Lemma 2 *Let C_1, C_2, C_3, C_4, C_5 be matrices with $C_5 = C_1(C_2, C_3)C_4$. Then we have*

$$\langle C_5, C_1(C_2, O)C_4 \rangle = \langle C_1(C_2, O)C_4, C_1(O, C_3)C_4 \rangle,$$

where O is the zero matrix.

We have

$$\begin{aligned} K(w) &= \int_X \|h_+(x, A, B) - h_+(x, A^*, B^*)\|^2 q(x) dx \\ &= \sum_{V \in \Omega^{(1)}: \text{Vol}(V) > 0} \int_V \|h_+(x, A, B) - h_+(x, A^*, B^*)\|^2 q(x) dx. \end{aligned}$$

By Theorem 1 and by Lemma 2, we have

$$c_0(K(w)) = c_0(\langle F_{i(1)}^{(1)} \circ \dots \circ F_{i(L)}^{(L)} \rangle),$$

where $i^{(s)} = \{i_1^{(s)}, i_2^{(s)}, \dots, i_{k^{(s)}}^{(s)}\} \subset \{1, 2, \dots, H^{(s)}\}$, $A_{i^{(s)}} = \begin{pmatrix} a_{i_1^{(s)}, i_1^{(s+1)}} & \cdots & a_{i_1^{(s)}, i_{k^{(s+1)}}^{(s+1)}} \\ \vdots & \cdots & \vdots \\ a_{i_{k^{(s)}}, i_1^{(s+1)}} & \cdots & a_{i_{k^{(s)}}, i_{k^{(s+1)}}^{(s+1)}} \end{pmatrix}$

and

$$B_{i^{(s)}} = \begin{pmatrix} b_{i_1^{(s)}} \\ \vdots \\ b_{i_{k^{(s)}}^{(s)}} \end{pmatrix}.$$

In this paper, we consider the case in **three-layered** neural networks with activation function ReLU (Rectified Linear Unit function).

Theorem 3 For $\alpha = 1, \dots, m$, let $f^{(\alpha)}(u^{(\alpha)})$ and $g^{(\alpha)}(u^{(\alpha)})$ be analytic functions. Also, let

$$c_{w^{*(\alpha)}} \left((f^{(\alpha)})^2, g^{(\alpha)} \right) = \lambda^{(\alpha)} \quad \text{and} \quad \theta_0 \left((f^{(\alpha)})^2, g^{(\alpha)} \right) = \theta^{(\alpha)}.$$

Then we have

$$c_{(w^{*(1)}, \dots, w^{*(m)})} \left(\langle f^{(1)}, \dots, f^{(m)} \rangle, \prod_{\alpha=1}^m g^{(\alpha)} \right) = \sum_{\alpha=1}^m \lambda^{(\alpha)},$$

and

$$\theta_{(w^{*(1)}, \dots, w^{*(m)})} \left(\langle f^{(1)}, \dots, f^{(m)} \rangle, \prod_{\alpha=1}^m g^{(\alpha)} \right) = \sum_{\alpha=1}^m (\theta^{(\alpha)} - 1) + 1.$$

(Proof)

Without loss of generality, we can assume that

$$f^{(\alpha)} = (u_1^{(\alpha)})^{k_1^{(\alpha)}} (u_2^{(\alpha)})^{k_2^{(\alpha)}} \cdots (u_{d^{(\alpha)}}^{(\alpha)})^{k_{d^{(\alpha)}}^{(\alpha)}},$$

$$g^{(\alpha)} = (u_1^{(\alpha)})^{h_1^{(\alpha)}} (u_2^{(\alpha)})^{h_2^{(\alpha)}} \cdots (u_{d^{(\alpha)}}^{(\alpha)})^{h_{d^{(\alpha)}}^{(\alpha)}}.$$

Let $u = (u_1^{(1)}, \dots, u_{d^{(1)}}^{(1)}, \dots, u_1^{(m)}, \dots, u_{d^{(m)}}^{(m)})$ and $d = \sum_{\alpha=1}^m d^{(\alpha)}$. Assume for $j \geq \theta^{(\alpha)} + 1$

$$\lambda^{(\alpha)} = \frac{h_1^{(\alpha)} + 1}{2k_1^{(\alpha)}} = \cdots = \frac{h_{\theta^{(\alpha)}}^{(\alpha)} + 1}{2k_{\theta^{(\alpha)}}^{(\alpha)}} < \frac{h_j^{(\alpha)} + 1}{2k_j^{(\alpha)}}.$$

By resolution of singularities, set $u_j^{(\alpha)} = u_1^{1,j^{(\alpha)}} \cdots u_d^{d,j^{(\alpha)}}$. For the matrix L given by

$$L = \begin{pmatrix} \ell_{1,1}^{(1)} & \ell_{2,1}^{(1)} & \cdots & \ell_{d,1}^{(1)} \\ \vdots & \vdots & \vdots & \vdots \\ \ell_{1,d^{(1)}}^{(1)} & \ell_{2,d^{(1)}}^{(1)} & \cdots & \ell_{d,d^{(1)}}^{(1)} \\ \vdots & \vdots & \vdots & \vdots \\ \ell_{1,1}^{(m)} & \ell_{2,1}^{(m)} & \cdots & \ell_{d,1}^{(m)} \\ \vdots & \vdots & \vdots & \vdots \\ \ell_{1,d^{(m)}}^{(m)} & \ell_{2,d^{(m)}}^{(m)} & \cdots & \ell_{d,d^{(m)}}^{(m)} \end{pmatrix},$$

the determinant of Jacobian $\left| \frac{\partial(u^{(1)}, \dots, u^{(m)})}{\partial u} \right|$ is

$$\prod_{s=1}^d u_s^{-1 + \sum_{\alpha=1}^m \sum_{j=1}^{d^{(\alpha)}} \ell_{s,j}^{(\alpha)}} |L|.$$

We have $\prod_{\alpha=1}^m g^{(\alpha)} = \prod_{s=1}^d u_s^{\sum_{\alpha=1}^m \sum_{j=1}^{d^{(\alpha)}} \ell_{s,j}^{(\alpha)} h_j^{(\alpha)}}$.

Since $f^{(1)^2} + \dots + f^{(m)^2}$ is normal crossing, there exists α_0 such that $f^{(\alpha_0)^2} \leq f^{(\alpha)^2}$ ($1 \leq \alpha \leq m$). Therefore, we have

$$\sum_{j=1}^{d^{(\alpha_0)}} \ell_{s,j}^{(\alpha_0)} 2k_j^{(\alpha_0)} \leq \sum_{j=1}^{d^{(\alpha)}} \ell_{s,j}^{(\alpha)} 2k_j^{(\alpha)}.$$

The possible candidates for the log canonical threshold are given by

$$\frac{\sum_{\alpha=1}^m \sum_{j=1}^{d^{(\alpha)}} (\ell_{s,j}^{(\alpha)} h_j^{(\alpha)} + \ell_{s,j}^{(\alpha)})}{\sum_{j=1}^{d^{(\alpha_0)}} \ell_{s,j}^{(\alpha_0)} 2k_j^{(\alpha_0)}} \geq \sum_{\alpha=1}^m \frac{\sum_{j=1}^{d^{(\alpha)}} (\ell_{s,j}^{(\alpha)} h_j^{(\alpha)} + \ell_{s,j}^{(\alpha)})}{\sum_{j=1}^{d^{(\alpha)}} \ell_{s,j}^{(\alpha)} 2k_j^{(\alpha)}} \geq \sum_{\alpha=1}^m \lambda^{(\alpha)},$$

where we use the inequality $\frac{s_3+s_4}{s_1+s_2} \geq \min\left\{\frac{s_3}{s_1}, \frac{s_4}{s_2}\right\}$ for $s_1, s_2, s_3, s_4 \geq 0$.

On the other hand, to obtain

$$\sum_{\alpha=1}^m \lambda^{(\alpha)} = \frac{\sum_{\alpha=1}^m \sum_{j=1}^{d^{(\alpha)}} (\ell_{s,j}^{(\alpha)} h_j^{(\alpha)} + \ell_{s,j}^{(\alpha)})}{\sum_{j=1}^{d^{(\alpha_0)}} \ell_{s,j}^{(\alpha_0)} 2k_j^{(\alpha_0)}},$$

requires $\sum_{j=1}^{\theta^{(\alpha_0)}} \ell_{s,j}^{(\alpha_0)} 2k_j^{(\alpha_0)} = \sum_{j=1}^{\theta^{(\alpha)}} \ell_{s,j}^{(\alpha)} 2k_j^{(\alpha)}$, and $\ell_{s,j}^{(\alpha)} = 0$ for $j > \theta^{(\alpha)}$. The dimension of the vectors satisfying such conditions is $\sum_{\alpha=1}^m (\theta^{(\alpha)} - 1) + 1$.

Finally, the Jacobian is nonzero if all vectors in L are linearly independent. These conditions complete the proof of the theorem.

(Q.E.D.)

Using Theorem 3, we have the following theorem.

Theorem 4 *Let*

$$h_+(x, A, B) = A^{(1)}(A^{(2)}x + B^{(2)})_+$$

The model has $H^{(3)}$ input units, $H^{(1)}$ output units, and $H^{(2)}$ hidden units in one hidden layer. Let us divide the hidden layer into $k^{(2)}$ groups:

$$H_1^{(2)} + H_2^{(2)} + \dots + H_{k^{(2)}}^{(2)} = H'^{(2)}.$$

Let $H_i'^{(1)}$ and $H_i'^{(2)}$ be the number such that $H^{(1)} - H_i'^{(1)}$ and $H^{(2)} - H_i'^{(2)}$ represents the number of removed neurons for $1 \leq i \leq k^{(2)}$, respectively.

Then, λ and θ for the model corresponding to the log canonical threshold

$$\lambda = \sum_{i=1}^{k^{(2)}} \lambda(H_i'^{(1)}, H_i'^{(2)} + 1, H^{(3)}, r_i)$$

and

$$\theta = \sum_{i=1}^{k^{(2)}} (\theta(H_i'^{(1)}, H_i'^{(2)} + 1, H^{(3)}, r_i) - 1) + 1,$$

where r_i is the rank corresponding the group i .

5 Softmax function

Next we consider the Softmax function. Let W be the set of parameters. Denote the input value by $x \in \mathbf{R}^{H(L+1)}$ with probability density function $q(x)$ and output value $y, z \in \mathbf{R}^{H(1)}$ with $y = h(x, w)$ and

$$z = \text{Softmax}(y|x, w) = \left(\frac{e^{y_1}}{\sum_i e^{y_i}}, \dots, \frac{e^{y_{H(1)}}}{\sum_i e^{y_i}} \right).$$

Consider the statistical model with Gaussian noise,

$$p(y|x, w) = \frac{1}{(\sqrt{2\pi})^{H(1)}} \exp\left(-\frac{1}{2} \|z - \text{Softmax}(y|x, w)\|^2\right).$$

Assume the set of optimal parameters W_0 by

$$\begin{aligned} W_0 &= \{w_0 \in W | L(w_0) = \min_{w' \in W} L(w')\} \\ &= \{w | \text{Softmax}(y|x, w) = \text{Softmax}(y|x, w_0)\}, \end{aligned}$$

and consider for $w^* \in W_0$,

$$c_{w^*}(\|\text{Softmax}(y|x, w) - \text{Softmax}(y|x, w^*)\|^2).$$

Theorem 5 *We have*

$$\begin{aligned} &\langle \text{Softmax}(y|x, w) - \text{Softmax}(y|x, w_0) \rangle \\ &= \langle y_2(w) - y_1(w) - (y_2(w_0) - y_1(w_0)), \dots, y_{H(1)}(w) - y_1(w) - (y_{H(1)}(w_0) - y_1(w_0)) \rangle \end{aligned}$$

(Proof)

Since

$$\begin{aligned} \left\langle \frac{1}{f} - \frac{1}{f'}, \frac{g}{f} - \frac{g'}{f'} \right\rangle &= \left\langle \frac{1}{f} - \frac{1}{f'}, \frac{g}{f} - \frac{g}{f'} + \frac{g}{f'} - \frac{g'}{f'} \right\rangle \\ &= \left\langle \frac{1}{f} - \frac{1}{f'}, g \left(\frac{1}{f} - \frac{1}{f'} \right) + \frac{g - g'}{f'} \right\rangle = \left\langle \frac{1}{f} - \frac{1}{f'}, g - g' \right\rangle, \end{aligned}$$

we have

$$\begin{aligned} &\langle \text{Softmax}(y|x, w) - \text{Softmax}(y|x, w_0) \rangle \\ &= \left\langle \left(\frac{e^{y_1(w)}}{\sum_i e^{y_i(w)}}, \dots, \frac{e^{y_{H(1)}(w)}}{\sum_i e^{y_i(w)}} \right) - \left(\frac{e^{y_1(w_0)}}{\sum_i e^{y_i(w_0)}}, \dots, \frac{e^{y_{H(1)}(w_0)}}{\sum_i e^{y_i(w_0)}} \right) \right\rangle \\ &= \left\langle \left(\frac{e^{y_1(w) - y_1(w)}}{\sum_i e^{y_i(w) - y_1(w)}}, \dots, \frac{e^{y_{H(1)}(w) - y_1(w)}}{\sum_i e^{y_i(w) - y_1(w)}} \right) - \left(\frac{e^{y_1(w_0) - y_1(w_0)}}{\sum_i e^{y_i(w_0) - y_1(w_0)}}, \dots, \frac{e^{y_{H(1)}(w_0) - y_1(w_0)}}{\sum_i e^{y_i(w_0) - y_1(w_0)}} \right) \right\rangle \\ &= \left\langle \frac{1}{\sum_i e^{y_i(w) - y_1(w)}} - \frac{1}{\sum_i e^{y_i(w_0) - y_1(w_0)}}, \right. \\ &\quad \left. e^{y_2(w) - y_1(w)} - e^{y_2(w_0) - y_1(w_0)}, \dots, e^{y_{H(1)}(w) - y_1(w)} - e^{y_{H(1)}(w_0) - y_1(w_0)} \right\rangle \\ &= \langle e^{y_2(w) - y_1(w)} - e^{y_2(w_0) - y_1(w_0)}, \dots, e^{y_{H(1)}(w) - y_1(w)} - e^{y_{H(1)}(w_0) - y_1(w_0)} \rangle \\ &= \langle e^{y_2(w) - y_1(w) - (y_2(w_0) - y_1(w_0))} - 1, \dots, e^{y_{H(1)}(w) - y_1(w) - (y_{H(1)}(w_0) - y_1(w_0))} - 1 \rangle \\ &= \langle y_2(w) - y_1(w) - (y_2(w_0) - y_1(w_0)), \dots, y_{H(1)}(w) - y_1(w) - (y_{H(1)}(w_0) - y_1(w_0)) \rangle. \end{aligned}$$

(Q.E.D.)

References

- [1] M. Aoyagi. The zeta function of learning theory and generalization error of three layered neural perceptron. *RIMS Kokyuroku, Recent Topics on Real and Complex Singularities*, 1501:153–167, 2006.
- [2] M. Aoyagi. A Bayesian learning coefficient of generalization error and Vandermonde matrix-type singularities. *Communications in Statistics - Theory and Methods*, 39(15):2667–2687, 2010.
- [3] M. Aoyagi. Consideration on singularities in learning theory and the learning coefficient. *Entropy*, 15(9):3714–3733, 2013.
- [4] M. Aoyagi. Learning coefficient in Bayesian estimation of restricted Boltzmann machine. *Journal of Algebraic Statistics*, 4(1):30–57, 2013.
- [5] M. Aoyagi. Learning coefficient of Vandermonde matrix-type singularities in model selection. *Entropy (Information Theory, Probability and Statistics)*, 21(6-561):1–12, 2019.
- [6] M. Aoyagi. Learning coefficients and information criteria. *Frontiers in Artificial Intelligence and Applications*, pages 351–362, 2019.
- [7] M. Aoyagi. Consideration on the learning efficiency of multiple-layered neural networks with linear units. *Neural Networks*, 172-106132:1–11, 2024.
- [8] M. Aoyagi and S. Watanabe. Resolution of singularities and the generalization error with Bayesian estimation for layered neural network. *IEICE Trans. J88-D-II*, 10:2112–2124, 2005a.
- [9] M. Drton. Conference lecture: Bayesian information criterion for singular models. *Algebraic Statistics 2012 in the Alleghenies at The Pennsylvania State University*, <http://jasonmorton.com/aspsu2012/>, 2012.
- [10] M. Drton, S. Lin, L. Weihs, and P. Zwiernik. Marginal likelihood and model selection for Gaussian latent tree and forest models. *Bernoulli*, 23(2):1202–1232, 2017.
- [11] W. Fulton. *Introduction to toric varieties, Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, USA, 1993.
- [12] Zach Furman and Edmund Lau. Estimating the local learning coefficient at scale. *ArXiv*, abs/2402.03698, 2024.
- [13] H. Hironaka. Resolution of singularities of an algebraic variety over a field of characteristic zero. *Annals of Math*, 79:109–326, 1964.
- [14] M. Kashiwara. B-functions and holonomic systems. *Inventiones Math.*, 38:33–53, 1976.
- [15] J. Kollár. Singularities of pairs. *Algebraic geometry-Santa Cruz 1995, Proc. Symp. Pure Math., American Mathematical Society, Providence, RI*, 62:221–287, 1997.
- [16] M. Mustata. Singularities of pairs via jet schemes. *J. Amer. Math. Soc.*, 15:599–615, 2002.
- [17] D. Rusakov and D. Geiger. Asymptotic model selection for naive Bayesian networks. *Proceedings of the Eighteenth Conference on Uncertainty in Artificial Intelligence*, pages 438–445, 2002.
- [18] D. Rusakov and D. Geiger. Asymptotic model selection for naive Bayesian networks. *Journal of Machine Learning Research*, 6:1–35, 2005.

- [19] S. Watanabe. *Algebraic Geometry and Statistical Learning Theory*, volume 25. Cambridge University Press, New York, USA, 2009.
- [20] S. Watanabe. A widely applicable bayesian information criterion. *Journal of Machine Learning Research*, (14):867–897, 2013.
- [21] P. Zwiernik. An asymptotic behavior of the marginal likelihood for general Markov models. *Journal of Machine Learning Research*, 12:3283–3310, 2011.