ON A PROBLEM OF KAC CONCERNING ANISOTROPIC LACUNARY SUMS

LORENZ FRUEHWIRTH AND MANUEL HAUKE

ABSTRACT. Given a lacunary sequence $(n_k)_{k \in \mathbb{N}}$, arbitrary positive weights $(c_k)_{k \in \mathbb{N}}$ that satisfy a Lindeberg-Feller condition, and a function $f : \mathbb{T} \to \mathbb{R}$ whose Fourier coefficients \hat{f}_k decay at rate $\frac{1}{k^{1/2+\epsilon}}$, we prove central limit theorems for $\sum_{k \le N} c_k f(n_k x)$, provided $(n_k)_{k \in \mathbb{N}}$ satisfies a Diophantine condition that is necessary in general. This addresses a question raised by M. Kac [Ann. of Math., 1946].

1. INTRODUCTION AND MAIN RESULTS

Lacunary sums are classical objects at the crossroads of number theory, dynamical systems, and the asymptotic theory of weakly dependent random variables. In this setup, one considers an integer sequence $(n_k)_{k\in\mathbb{N}}$ satisfying the Hadamard-gap condition

(1)
$$\liminf_{k \to \infty} \frac{n_{k+1}}{n_k} \ge q > 1,$$

and for a sufficiently regular function $f : \mathbb{T} = \mathbb{R} / \mathbb{Z} \to \mathbb{R}$ with mean 0, we set

$$S_N(x) := \sum_{k \le N} f(n_k x), \quad N \in \mathbb{N}, \ x \in \mathbb{T}.$$

For over a century, mathematicians have been interested in the limiting behavior of these sums as $N \to \infty$. Early works trace back to Borel [10] who studied sums $\sum_{k \le N} f(2^k x)$, generated by certain indicator functions f, in order to understand normal number properties, and to Weyl [23], who was interested in the uniform distribution properties of sequences of the form $(n_k x)_{k \in \mathbb{N}}$. Researchers soon recognized profound connections to other fields and started to investigate lacunary sums as an independent topic. In 1924, Kolmogorov proved in his article [17] that

$$\sum_{k\leq N} a_k \cos(2\pi n_k x)$$

converges for almost all $x \in \mathbb{T}$ if one assumes square summability of the coefficients $(a_k)_{k \in \mathbb{N}}$. A few years passed before researchers succeeded in establishing the first results on the scale of the central limit theorem. In his groundbreaking work [15], Kac proved in 1946 that for any mean 0 function $f : \mathbb{T} \to \mathbb{R}$ whose Fourier coefficients decay sufficiently fast, we have

$$\frac{1}{\sqrt{N}}\sum_{k\leq N}f(2^kx)\xrightarrow{w}\mathcal{N}(0,\sigma^2)$$

for some suitable $\sigma > 0$. Naively one would expect the variance σ^2 to equal $\int_{\mathbb{T}} f(x)^2 dx$ which would be in accordance with true independent behaviour; however it turned out that the resulting (asymptotic) variance is given by

$$\sigma^{2} = \int_{\mathbb{T}} f(x)^{2} dx + 2 \sum_{k=1}^{\infty} \int_{\mathbb{T}} f(2^{k} x) f(x) dx.$$

This deviation from the i.i.d. random case arises from an intriguing correlation structure, which, as later discovered, is inherent to certain dynamical systems. In the same article as well as in his later work [16], Kac also studied super-lacunary sequences, i.e. where

(2)
$$\lim_{k \to \infty} \frac{n_{k+1}}{n_k} = \infty$$

In this case, a central limit theorem was established which perfectly aligns with the classical one for a sequence of independently, uniformly distributed random variables in [0, 1), where the asymptotic

²⁰¹⁰ Mathematics Subject Classification. Primary 42A55, 60F05; Secondary 11D04, 11D45.

variance is given by $\int_{\mathbb{T}} f(x)^2 dx$. In the spirit of Kolmogorov, Kac was even able to deal with anisotropic lacunary sums of the form

$$\sum_{k\leq N} c_k f(n_k x),$$

where the sequence $(c_k)_{k \in \mathbb{N}}$ satisfies a Lindeberg-Feller-type condition, i.e. he assumed

(3)
$$\max_{k \in \mathbb{N}} c_k < \infty \quad \text{and} \quad \sum_{k \le N} c_k^2 \xrightarrow{N \to \infty} \infty.$$

In his seminal article [15], following a communication with Paul Erdős, Kac expressed his desire to obtain central limit theorems for

(4)
$$\frac{\sum_{k \le N} c_k f(n_k x)}{\left\|\sum_{k \le N} c_k f(n_k x)\right\|_2},$$

where $(n_k)_{k \in \mathbb{N}}$ is an *arbitrary* lacunary sequence only satisfying Hadamard's condition (1). It soon became clear that this goal was not attainable in full generality, even when the generating function $f: \mathbb{T} \to \mathbb{R}$ was restricted to being a trigonometric polynomial. The famous example of Erdős and Fortet (see [16, p.646]) demonstrates that choosing $f(x) = \cos(2\pi x) + \cos(4\pi x)$ and $n_k := 2^k - 1$ results in a lacunary sum $S_N = \sum_{k \le N} f(n_k x)$ that satisfies a non-degenerate distributional limit theorem, with the asymptotic distribution being non-Gaussian. However, Salem and Zygmund [21] proved a central limit theorem for $\sum_{k \le N} c_k \cos(n_k x)$, where $(n_k)_{k \in \mathbb{N}}$ is arbitrary and the weights $(c_k)_{k \in \mathbb{N}}$ are assumed to only satisfy the minimal condition (3). These examples illustrate that the asymptotic behavior of $S_N = S_N(f, (n_k)_{k \in \mathbb{N}})$ is determined not only by the function f but also by the number-theoretic properties of the sequence $(n_k)_{k \in \mathbb{N}}$, as well as by a subtle interplay between these two entities. For a more comprehensive discussion on the foundations of lacunary sums and their connections to various topics in modern mathematics, we refer the interested reader to the excellent recent survey article [5].

Over the years mathematicians tried to establish central limit theorems of the type in (4) from various directions. The *q*-adic lacunary sequences $n_k = q^k$, where $q \in \mathbb{N}_{\geq 2}$ play a special role since one can relate lacunary sums of the form $\sum_{k \in \mathbb{N}} f(q^k x)$ to the theory of dynamical systems via interpreting $q^k x$ as a *k*-fold composition of the *q*-adic shift-operator $\mathcal{T}_q : \mathbb{T} \to \mathbb{T}$ defined by $x \mapsto qx$. In recent years, it was shown in [9] that $\mathcal{T}_q \circ \ldots \circ \mathcal{T}_q$ can be replaced by a composition of potentially different operators $\mathcal{T}^1 \circ \mathcal{T}^2 \circ \ldots$, and additionally the function *f* can be replaced by a sequence of functions f_1, f_2, \ldots under some reasonably mild additional conditions: There, a central limit theorem is established for random variables of the form

$$\frac{\sum_{k \le N} f_k(\mathcal{T}_k \circ \dots \circ \mathcal{T}_1(x))}{\left|\left|\sum_{k \le N} f_k(\mathcal{T}_k \circ \dots \circ \mathcal{T}_1(x))\right|\right|_2}, \qquad x \in \mathbb{T}, \quad N \in \mathbb{N}.$$

In particular, the findings in [9] imply a central limit theorem for sequences of the form

$$\frac{\sum_{k\leq N} c_k f(q^k x)}{\left|\left|\sum_{k\leq N} c_k f(q^k x)\right|\right|_2},$$

as it was claimed¹ for q = 2 in [15, p.49].

For quite some time, the picture for arbitrary lacunary sequences $(n_k)_{k\in\mathbb{N}}$ remained much less clear, as it was necessary to identify requirements on $(n_k)_{k\in\mathbb{N}}$ to prevent anomalies like those arising in the Erdős-Fortet example. In his famous article, Gaposhkin [12] was able to show that, if for any fixed choice of $a, b, c \in \mathbb{N}$, there are only finitely many solutions (in k and ℓ) to

$$an_k - bn_\ell = c$$

then $S_N = \sum_{k \le N} f(n_k x)$ satisfies a central limit theorem with asymptotic variance $\int_{\mathbb{T}} f^2(x) dx$, where f can be taken from a reasonably large class of functions. It took about 40 years until Aistleitner and

¹Formally, Kac only states that such a central limit theorem can be obtained without significant changes to the arguments in his article, but does not provide a proof for that claim.

Berkes were able to reduce Gaposhkin's requirements (in the anisotropic case) to a minimal level. In their remarkable article [4], they demonstrated that under the conditions

(6)
$$\sup_{c \in \mathbb{N}} \# \{k, \ell \le N : jn_k - j'n_\ell = c\} + \# \{k, \ell \le N, k \ne \ell : jn_k - j'n_\ell = 0\} = o_d(N), \quad 1 \le j, j' \le d \in \mathbb{N},$$

and

(7)
$$\left\| \sum_{k \le N} f(n_k x) \right\|_2 \ge C\sqrt{N},$$

for some C > 0, we have

$$\frac{\sum_{k \le N} f(n_k x)}{\sqrt{N} ||f||_2} \xrightarrow{w} \mathcal{N}(0, 1),$$

whenever $f : \mathbb{T} \to \mathbb{R}$ is an arbitrary mean-zero function of bounded variation. In [4, Theorem 1.3], the authors also showed that the condition (6) is minimal in order to guarantee the existence of a central limit theorem. Diophantine conditions as the one in (6) were also heavily investigated on the scale of the law of the iterated logarithm, see e.g. [2, 6]. In particular, we highlight the article [22] by Weiss who considered anisotropic lacunary trigonometric sums. In [1], similar results were obtained for a more general class of generating functions *f* making progress in a question raised by Philipp in his famous article [19].

In recent years the study of lacunary sums has been extended even further to the scale of moderate and large deviations revealing an unexpected asymptotic behavior not visible on the scales of the central limit theorem and the law of the iterated logarithm (see [7, 11, 20]).

Returning to the question of the existence of central limit theorems, we remark that in case of anisotropic sums $S_N = \sum_{k \le N} c_k f(n_k x)$, progress appears to have been much slower and more limited, unless a particular structure such as $n_k = q^k$ is assumed. The existing isolated results primarily address the special cases where $f(x) = \cos(2\pi x)$ (see [21]) or $f(x) = \operatorname{sign}(\cos(2\pi x))$. The latter is related to the famous Walsh system for which central limit theorems are obtained in, e.g., [18].

The objective of this article is to establish central limit theorems for a broad class of anisotropic lacunary sums under suitable Diophantine conditions. In the article [13], central limit theorems were proven for $\sum_{k \le N} c_{k,N} f(n_k x)$, where $(c_{k,N})_{k \le N}$ is a triangular scheme of weights satisfying an analogous condition to (3) and f can be taken from a very large class of functions. However, the sequence $(n_k)_{k \in \mathbb{N}}$ needs to satisfy restrictive Diophantine conditions.

Our main results (Theorems 1 and 2) represent significant progress on the original question posed by Kac in [15], as the Diophantine condition we impose in (3) can be viewed as minimal (see the discussion in Remark 1.2).

1.1. **Main Results.** Before presenting our main results, we will introduce and discuss the assumptions required to establish Theorem 1 and Theorem 2.

Assumption 1. Let $f : \mathbb{T} \to \mathbb{R}$ be an $\mathbb{L}_2(\mathbb{T})$ -function with

$$\int_{\mathbb{T}} f(x) \, \mathrm{d}x = 0, \qquad \int_{\mathbb{T}} f^2(x) \, \mathrm{d}x =: \sigma^2 \in (0, \infty).$$

Writing $f(x) = \sum_{j=1}^{\infty} [a_j \cos(2\pi j x) + b_j \sin(2\pi j x)]$, we shall assume that there are absolute constants M > 0 and $\rho > \frac{1}{2}$ such that

$$|a_j| + |b_j| \le \frac{M}{j^{\rho}}, \quad j \in \mathbb{N}.$$

Remark 1.1. The assumption of $f : \mathbb{T} \to \mathbb{R}$ having a finite second moment is necessary in order to establish a central limit theorem. The requirement for its Fourier coefficients to decay at a certain rate is the same as in [15] and represents a slight improvement over [4], since, in particular, every function of bounded variation satisfies the above. Moreover, any Hölder-continuous function with exponent $\rho > \frac{1}{2}$ has Fourier-coefficients tending to 0 at speed $O(j^{-\rho})$.

Our methods are not restricted to a fixed sequence of weights $(c_k)_{k \in \mathbb{N}}$, but also accommodate more general triangular schemes $(c_{k,N})_{k \leq N}$, with $N \in \mathbb{N}$. The following assumption represents the natural generalization to the Lindeberg-type condition from [15] for triangular arrays of weights.

Assumption 2. The array of weights $(c_{k,N})_{k \le N, N \in \mathbb{N}}$ shall satisfy

(8)
$$\sup_{N \in \mathbb{N}} \sup_{k \le N} c_{k,N} \le 1, \quad h(N) := \sum_{k \le N} c_{k,N}^2 \longrightarrow \infty \quad as \quad N \to \infty.$$

The results in the works [2, 4, 6] suggest that it is necessary to limit the number of solutions to certain Diophantine equations in order to obtain a central limit theorem. The following assumption not only imposes a limit on the number of Diophantine solutions to equations

$$an_k - bn_\ell = c_k$$

but also captures the subtle interplay between the weights $(c_{k,N})$ and the given lacunary sequence $(n_k)_{k\in\mathbb{N}}$. Heuristically speaking, if there are not too many *inhomogeneous* (i.e. $c \neq 0$) solutions to the equation above (and some additional lower bound on the variance exists), then the clt holds with a non-degenerate variance. We remark that this bound on the inhomogeneous solutions is not satisfied in the Erdős-Fortet example and can be seen as the reason that the clt fails in this instance. If additionally the number of solutions to the *homogeneous* equation (i.e. where c = 0) is not too large, then we expect the clt to hold with asymptotic variance $\int_0^1 f^2(x) dx$. We remark that this phenomenon was accurately described in [3] on the scale of the law of the iterated logarithm. The bounds on the number of solutions in the homogeneous and inhomogeneous cases assumed in this article are as follows:

Assumption 3. For any fixed $d \in \mathbb{N}$, let us assume that the array of weights $(c_{k,N})_{k \leq N}$, $N \in \mathbb{N}$ and the sequence $(n_k)_{k \in \mathbb{N}}$ satisfy

$$L(N,d) := \sup_{c>0} \sum_{k,\ell \le N} \sum_{1 \le j,j' \le d} c_{k,N} c_{\ell,N} \mathbb{1}_{[jn_k - j'n_\ell = c]} = o(h(N)),$$

and

(9)
$$\left\| \sum_{k \le N} c_{k,N} f(n_k x) \right\|_2 \ge \delta \sqrt{h(N)}$$

for some absolute $\delta > 0$.

Assumption 4. For any $d \in \mathbb{N}$ fixed, we assume

$$L^{*}(N,d) := L(N,d) + \sum_{\substack{1 \le k,\ell \le N \\ k \ne \ell}} \sum_{\substack{j,j' \le d}} c_{k,N} c_{\ell,N} \mathbb{1}_{[jn_{k}-j'n_{\ell}=0]} = o(h(N)).$$

With these assumptions at hand, we present the main results of our article.

Theorem 1. Let $(n_k)_{k \in \mathbb{N}}$ be a lacunary sequence, let $(c_{k,N})_{k \leq N}$ be a triangular array of non-negative reals, and let $f : \mathbb{T} \to \mathbb{R}$ be such that Assumptions 1, 2 and 3 are satisfied. Then we have

$$\frac{\sum_{k \le N} c_{k,N} f(n_k x)}{\left|\left|\sum_{k \le N} c_{k,N} f(n_k x)\right|\right|_2} \xrightarrow{w} \mathcal{N}(0,1).$$

Theorem 2. Let $(n_k)_{k \in \mathbb{N}}$ be a lacunary sequence, let $(c_{k,N})_{k \leq N}$ be a triangular array of non-negative reals, and let $f : \mathbb{T} \to \mathbb{R}$ be such that Assumptions 1, 2 and 4 are satisfied. Then we have

$$\frac{\sum_{k \le N} c_{k,N} f(n_k x)}{||f||_2 \sqrt{h(N)}} \xrightarrow{w} \mathcal{N}(0,1).$$

Remark 1.2. • Since Assumption 2 imposed in Theorem 1 and Theorem 2 is equivalent to

$$\max_{k \le N} \frac{c_{k,N}}{\sqrt{\sum_{\ell \le N} c_{\ell,N}^2}} \longrightarrow 0$$

our main results can be viewed as central limit theorems under a Lindeberg-Feller condition. This is clearly a necessary requirement to obtain a clt. • Even without Assumptions 3 or 4, it follows directly from Lemma 2.7 that $||S_N||_2 \ll ||f||_2 \sqrt{h(N)}$. On the other hand, under Assumption 3, we have

$$0 < \frac{||S_N||_2}{\sqrt{h(N)}||f||_2} \ll 1,$$

where the positivity is guaranteed by (9). We emphasize that analogous assumptions to (9) were also employed to obtain central limit theorems in [4, 12, 13, 15].

• Both the conditions with Diophantine solutions to (5) for c = 0 as well as for c > 0 are in general necessary to deduce that $||S_N||_2 = ||f||\sqrt{h(N)}(1+o(1))$: In the q-adic isotropic setting, we obtain many Diophantine solutions for c = 0 which implies $||S_N||_2 = \sigma(q)\sqrt{h(N)}(1+o(1))$ with $\sigma(q) \neq ||f||_2$ in general. If some c > 0 has so many Diophantine solutions that Assumption 3 fails, [4, Theorem 1.3] implies that

$$\frac{\sum_{k \le N} f(n_k x)}{\|\sum_{k \le N} f(n_k x)\|_2}$$

does, in general, not converge to a normal distribution.

• The sharpness of the Diophantine condition in Assumption 3 in the isotropic case (i.e. where $c_{k,N} = 1$ for all $k \le N$ and for all $N \in \mathbb{N}$) shows that our Assumptions 3 resp. 4 can in general not be omitted. Naturally, the question arises whether a counterexample to a clt can be constructed for any triangular scheme of weights. Specifically, given a sequence $(c_{k,N})_{k\le N,N\in\mathbb{N}}$ satisfying Assumption 2, are there f and $(n_k)_{k\in\mathbb{N}}$ satisfying Assumption 1 and the Hadamard gap condition such that $\sum_{k\le N} c_{k,N} f(n_k x)$ fails to satisfy a clt, regardless of the choice of normalization? However, such general example cannot be constructed for obvious reasons: considering the sequence $(c_1, c_2, ...) = (1, 0, 1, 0, 0, 1, 0, 0, 1, ...)$, then independent of the choice of $(n_k)_{k\in\mathbb{N}}$, the associated lacunary sum S_N satisfies

$$S_N \asymp \sum_{k \leq \log(N)} f(\tilde{n}_k x),$$

where $(\tilde{n}_k)_{k \in \mathbb{N}}$ is a super-lacunary sequence (see also Corollary 1.4). For the latter, the Assumption 3 is satisfied (even with the number of Diophantine solutions L(N,d) being uniformly bounded in N). Such lacunary sums always satisfy the clt with asymptotic variance $\int_{\mathbb{T}} f(x)^2 dx$.

Corollary 1.3. In the isotropic case $c_{k,N} := 1$ for all $N \in \mathbb{N}$ and $k \leq N$, we recover [4, Theorem 1.1 and Theorem 1.2]. In this setting, Assumption 3 coincides with the requirement in [4, Theorem 1.1], while Assumption 4 ensures that the conclusions of [4, Theorem 1.2] hold.

Corollary 1.4. One of the few instances where a clt is known to hold for anisotropic lacunary sums is demonstrated in [16]. There, any super-lacunary sequence $(n_k)_{k \in \mathbb{N}}$ generates a lacunary sum S_N for which the clt with asymptotic variance $\int_{\mathbb{T}} f(x)^2 dx$ is satisfied. Theorem 2 directly implies the findings in [16] since any super-lacunary sequence $(n_k)_{k \in \mathbb{N}}$ satisfies

$$L^*(N,d) = O_d(1), \qquad d \in \mathbb{N}.$$

1.2. **Notation.** In this article we deal with random variables S_N , $N \in \mathbb{N}$ which are defined on the probability space $(\mathbb{T}, \mathscr{B}(\mathbb{T}), \lambda)$, where \mathscr{B} denotes the Borel- σ -algebra, \mathbb{T} is the univariate torus and λ denotes the normalized Haar-measure on \mathbb{T} . Depending on the context we sometimes also use the notation $\mathbb{P} = \lambda$ and we write $\mathbb{E}[S_N]$ for the expected value

$$\mathbb{E}[S_N] = \int_{\mathbb{T}} S_N(x) d\lambda(x) = \int_{\mathbb{T}} S_N(x) dx.$$

We denote convergence in distribution by \xrightarrow{w} and convergence in probability by $\xrightarrow{\mathbb{P}}$. When dealing with a deterministic sequence we will just write \longrightarrow to indicate convergence. For a random variable S_N on $(\mathbb{T}, \mathscr{B}(\mathbb{T}), \lambda)$ we write

$$||S_N||_2 := \left(\int_{\mathbb{T}} S_N^2(x) \, \mathrm{d}x\right)^{1/2}.$$

and $||S_N||_{\infty} = \sup_{x \in \mathbb{T}} |S_N(x)|$. The collection of random variables *X* on $(\mathbb{T}, \mathscr{B}(\mathbb{T}), \lambda)$ with $||X||_2 < \infty$ is denoted by $\mathbb{L}_2(\mathbb{T})$. Let $\mathscr{F} \subseteq \mathscr{B}(\mathbb{T})$ be another σ -algebra, then we denote the conditional expectation

of a random variable *X* with respect to \mathscr{F} by $\mathbb{E}[X|\mathscr{F}]$. We denote the standard normal distribution function as Φ , i.e., we have

$$\Phi(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{t} e^{-\frac{x^2}{2}} \mathrm{d}x.$$

For sequences $(a_N)_{N \in \mathbb{N}}$ and $(b_N)_{N \in \mathbb{N}}$ we write

$$a_N = O(b_N)$$
 or $a_N \ll b_N$,

if there exists a constant c > 0 such that $|a_N| \le cb_N$ for all but finitely many $N \in \mathbb{N}$. If $\lim_{N\to\infty} \frac{a_N}{b_N} = 0$, we write $a_N = o(b_N)$, and if $a_N = O(b_N)$ and $b_N = O(a_N)$, we write $a_N \asymp b_N$. Let $A \subseteq \mathbb{Z}$ be a finite set, then we denote the number of elements in A by #A.

1.3. Acknowledgments. LF is partially supported by the DFG project *Limit theorems for the volume of random projections of* ℓ_p *-balls* (project number 516672205). A significant part of this work was carried out during visits of LF at NTNU Trondheim respectively MH at the University of Passau. We thank both universities for their support and hospitality. We would also like to thank Christoph Aistleitner and Joscha Prochno for various helpful comments on an earlier version of this manuscript.

2. Proofs

2.1. Heuristic ideas and novelties of the proof. The main challenge is obtaining a central limit theorem for lacunary sums $S_N = S_N(f)$ when f is a trigonometric polynomial of fixed degree d. The coarse strategy to achieve this is similar to the approach in [4] where a filtered probability space and a martingale, which approximates S_N reasonably well, are constructed. Since central limit theorems for such adapted processes are well understood (see Proposition 2.1), this allows us to obtain a central limit theorem for S_N being generated by a trigonometric polynomial of finite degree. The transition to arbitrary functions with Fourier decay as in Assumption 1 then follows from standard arguments.

In order to achieve the clt for trigonometric polynomials, one partitions the set of integers [1, N] into two types of blocks. On the one hand, there are long blocks Δ_i (of polynomial length in N) that should contain the majority of the "mass" (in terms of the weights c_k^2). On the other hand, between Δ_i and Δ_{i+1} for each *i*, there are short "buffer blocks" Δ'_i (of logarithmic length) ensuring that elements of distinct long blocks behave almost independently. This allows us to establish a clt on the long blocks which implies a clt on the entire lacunary sum, provided the contribution from the Δ'_i is negligible. The latter is from the technical point of view one of the key obstructions to move from the isotropic case of [4] to the anisotropic case considered here. A crucial part of the arguments used in [4] is that in the isotropic case, the "mass" of each block Δ_i is proportional to its number of elements, i.e.

$$\sum_{k\leq N} 1 = \sum_{k\leq N} c_k = \sum_{k\leq N} c_k^2,$$

which is implicitly used at many positions of the proof, and most importantly in the above-mentioned buffer block construction. In the general anisotropic case, a direct copy of this approach is doomed to fail: A buffer block needs to contain (at least) about $\log N$ many *elements*, but this does not tell us anything about the mass $\sum_{k \in \Delta'_i} c_k^2$ in comparison to $\sum_{k \le N} c_k^2$: For arbitrary coefficients c_k , it might be that a positive proportion of $\sum_{k \le N} c_k^2$ is contained in even one single buffer block, and the whole construction breaks down. To address this issue, we partition the c_k into different sets according to their size, ensuring that all c_k within a given set do not differ significantly from one another. More specifically, we focus on the case where $c_k \ge N^{-\beta}$ for some fixed $\beta > \frac{1}{4}$. Under this assumption, it is ensured that the mass of any buffer block is negligible when compared to the entire sum (which now has polynomially growing "mass" $\sum_{k \le N} c_k^2$). With the aid of additional refined estimates, which are required to change between the 1-norm and the 2-norm of the sequence $(c_k)_{k \le N}$ in the subsequent proofs, we establish a clt for the case where the weights c_k are of similar size. This result is proved in Section 2.3.

Having a central limit theorem settled for those c_k with $c_k \ge N^{-\beta}$, a central limit theorem for weights

 c_k with $N^{-1/2} \le c_k \le N^{-\beta}$ can then be obtained immediately by renormalizing with $N^{-\beta}$. This gives us a marginal central limit theorem for

(10)
$$\left(\sum_{k\in A} c_k f(n_k x), \sum_{k\in B} c_k f(n_k x)\right),$$

where *A* contains the indices with $c_k \ge N^{-\beta}$ and *B* the ones with $N^{-1/2} \le c_k \le N^{-\beta}$. The remaining challenge lies in proving that the vector above is asymptotically uncorrelated, and thus independent. By a straight-forward pigeonhole argument, there exists a parameter $\beta \approx 1/4$ such that $\sum_{c_k \approx N^{-\beta}} c_k^2$ is negligibly small. Removing those *k* from the above sums does not destroy the marginal central limit theorems, but ensures a gap between the sizes of the coefficients, i.e. $\frac{\min_{k \in A} c_k}{\max_{k \in B} c_k} \longrightarrow 0$ at a certain speed. This suffices to show that the components of (10) are asymptotically uncorrelated. The required argument is given in the proof of Lemma 2.10 in Section 2.4.

2.2. **Prerequisites.** We establish the central limit theorem in Proposition 2.3 using methods from (discrete) martingale theory. This will need some preparation which is done in the upcoming subsections. For the moment, we will just state our main auxiliary result below.

Proposition 2.1. (*Theorem* 1 *in* [14] *for* $\delta = 1$)

Let $(X_M)_{M \in \mathbb{N}}$ with $X_M = \sum_{i=1}^M Y_i$ be a martingale adapted to a filtration $(\mathscr{F}_M)_{M \in \mathbb{N}}$ such that all Y_i have finite fourth moments. We set $V_M := \sum_{i=1}^M \mathbb{E}[Y_i^2|\mathscr{F}_{i-1}]$ and define the sequence $(s_M)_{M \in \mathbb{N}}$ via $s_M := \sum_{i=1}^M \mathbb{E}[Y_i^2]$. Then

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}\left[\frac{Y_1 + \ldots + Y_M}{\sqrt{s_M}} \le t \right] - \Phi(t) \right| \ll \left(\frac{\sum_{i=1}^M \mathbb{E}\left[Y_i^4 \right] + \mathbb{E}\left[(V_M - s_M)^2 \right]}{s_M^2} \right)^{1/5}$$

where the implicit constant is absolute and Φ denotes the standard normal distribution function.

Lemma 2.2 (Lemma 2.2 in [4]). Let $f : \mathbb{T} \to \mathbb{R}$ satisfy Assumption 1. Then for any $a, b \in \mathbb{T}$ we have

$$\int_{[a,b]} f(\lambda x) \, \mathrm{d}x \le \frac{1}{\lambda} \int_{\mathbb{T}} |f(x)| \, \mathrm{d}x \le \frac{1}{\lambda} \|f\|_{\infty}.$$

2.3. An anisotropic clt for trigonometric polynomials under weight restrictions. In this section, we focus on the case where f = p is an even trigonometric polynomial of fixed degree $d \in \mathbb{N}$. Later we will approximate $f \in \mathbb{L}_2(\mathbb{T})$ by suitably chosen trigonometric polynomials. The following proposition establishes a clt under the assumption that our weights $c_{k,N}$ are roughly of the same size on a logarithmic scale.

Proposition 2.3. Let $(n_k)_{k \in \mathbb{N}}$ be a lacunary sequence satisfying the Hadamard-gap condition for some q > 1, let $p(x) = \sum_{j=1}^{d} a_j \cos(2\pi j x)$ be a mean 0 even trigonometric polynomial of degree d and let $(c_{k,N})_{k \le N, N \in \mathbb{N}}$ satisfy Assumption 2.

Additionally, let $\beta = \frac{1}{4} + \delta$ for some $0 < \delta \le \frac{1}{12}$, and assume that $N^{-\beta} < c_{k,N} \le 1$ for all $k \le N$. Then we have the following:

(i) If Assumption 3 is satisfied (with p = f in (9)), then

$$\frac{\sum_{k \le N} c_{k,N} p(n_k x)}{\left|\left|\sum_{k \le N} c_{k,N} p(n_k x)\right|\right|_2} \xrightarrow{w} \mathcal{N}(0,1).$$

(ii) If Assumption 4 is satisfied, then

$$\frac{\sum_{k\leq N} c_{k,N} p(n_k x)}{\sqrt{h(N)} ||p||_2} \xrightarrow{w} \mathcal{N}(0,1).$$

The proof of Proposition 2.3 is the first key ingredient in order to establish Theorem 2 and will take up several pages. In the following, we will always deal with an array of weights $(c_{k,N})_{k \le N, N \in \mathbb{N}}$ satisfying Assumption 2 and we additionally assume

$$N^{-\beta} \le c_{k,N} \le 1, \quad \forall k \le N.$$

In order to keep the notational burden low, we will omit in the following the additional index N, writing c_k instead. Further, we will omit the dependence of implied constants arising on the fixed polynomial p or the variable q from the Hadamard condition. We start with the construction of a filtration and an adapted process, with the notation defined below being frequently used in the subsequent auxiliary results.

Construction of the filtration and an adapted process. As sketched in Section 2.1, given a large integer $N \in \mathbb{N}$, we partition the interval $[1, N] \subseteq \mathbb{N}$ using a finite sequence of blocks $\Delta_1, \Delta'_1, \Delta_2, \ldots$, where the Δ_i are "dense", i.e. their union contains most of the elements of [1, N] while the Δ'_i act as "buffer blocks" making sure that for $k \in \Delta_i, \ell \in \Delta_j, i \neq j$, the random variables $f(n_k x), f(n_\ell x)$ behave sufficiently "independently". In order to obtain the desired construction, we proceed algorithmically. Let $h(N) = \sum_{k \leq N} c_k^2$ and fix $0 < \gamma < 1/2$. For i = 1, we take $A_1 = 0$ and B_1 such that

$$\sum_{A_1 \leq k \leq B_1} c_k^2 \in [h(N)^{\gamma}, h(N)^{\gamma} + 1].$$

This is possible since $\sum_{k \le N} c_k^2 = h(N) \longrightarrow \infty$ and $c_k \le 1$. We then set $\Delta_1 := [A_1, B_1]$. Next, we choose $\Delta'_1 = [A'_1, B'_1]$ with $A'_1 := B_1 + 1$, $B'_1 = A'_1 + \lceil K \log_q(h(N)) \rceil$ where K > 0 is a fixed but large constant. Now we proceed inductively, assuming that we have constructed

$$A_1, B_1, A'_1, B'_1, \dots, A_{i-1}, B_{i-1}, A'_{i-1}, B'_{i-1}.$$

We choose $A_i := B'_{i-1} + 1$ and B_i such that $\sum_{A_i \le k \le B_i} c_k^2 \in [h(N)^{\gamma}, h(N)^{\gamma} + 1]$, and set $\Delta_i := [A_i, B_i]$. Further, we define $\Delta'_i := [A'_i, B'_i]$ where $A'_i := B_i + 1$, $B'_i := A'_i + [K \log_q(h(N))]$. We do this until one of these blocks exceeds N (for formal reasons, we set $c_k := 1, k > N$), which happens at some point $M = M(N) \in \mathbb{N}$ while carrying out the described construction. At this point, our algorithm terminates and we have obtained the following covering:

$$[1,N] \subseteq \bigcup_{i=1}^{M} \Delta_i \cup \Delta'_i$$

or, in other words, $N \le \max{\{\Delta'_M\}}$.

Having this construction set up, we highlight the following important properties: For all i = 1, ..., M, we have

(11)
$$\sum_{k \in \Delta_i} c_k^2 = h(N)^{\gamma} + O(1), \quad \sum_{k \in \Delta_i'} c_k^2 \ll_K \log h(N), \quad M = M(N) \approx \frac{h(N)}{h(N)^{\gamma}} = h(N)^{1-\gamma/2}$$

Based on our lacunary sequence $(n_k)_{k \in \mathbb{N}}$, we will construct a filtration $(\mathscr{F}_i)_{i \in \mathbb{N}}$ and a sequence of adapted random variables $(\varphi_k)_{k \in \Delta_i}$ which will approximate $(p(n_k x))_{k \in \Delta_i}$ with

$$p(x) = \sum_{j=1}^{d} a_j \cos(2\pi j x)$$

sufficiently well.

Lemma 2.4. Given $(\Delta_i)_{i \le M}$ as in Subsection 2.3. For $k \in \Delta_i$, we define $m(k) := \lceil \log_2 n_k + \frac{K}{2} \log_2 h(N) \rceil$. Further, we set

$$\mathscr{G}_k := \sigma\left(\left\{\left[\frac{\nu}{2^{m(k)}}, \frac{\nu+1}{2^{m(k)}}\right] : \nu = 0, \dots, 2^{m(k)} - 1\right\}\right), \quad \text{as well as} \quad \mathscr{F}_i := \mathscr{G}_{B_i}$$

Then for every $k \in \Delta_i$, there exists $\varphi_k : \mathbb{T} \to \mathbb{R}$ such that:

- (i) $\varphi_k(x)$ is constant on $I_{v,k} := \left[\frac{v}{2^{m(k)}}, \frac{v+1}{2^{m(k)}}\right]$ for any $v = 0, ..., 2^{m(k)} 1$.
- (*ii*) $\|\varphi_k p(n_k \cdot)\|_{\infty} \ll \frac{1}{h(N)^{K/2}}$.
- (*iii*) $\mathbb{E}[\varphi_k | \mathscr{F}_{i-1}] \equiv 0.$

Proof. We define $\hat{\varphi}_k := \mathbb{E}[p(n_k \cdot)|\mathscr{G}_k]$. For any $x, x' \in I_{v, B_{i-1}}$, the Lipschitz continuity of p implies

$$|p(n_k x) - p(n_k x')| \ll \frac{n_k}{2^{m(k)}} \ll \frac{1}{h(N)^{K/2}}.$$

This shows properties (*i*) and (*ii*) for the approximations $\hat{\varphi}_k$. For $x \in I_{v,k}$, we set

$$\varphi_k(x) := \hat{\varphi}_k(x) - \frac{1}{I_{\nu,k}} \int_{I_{\nu,k}} \hat{\varphi}_k(t) dt$$

Then (*i*) holds for φ_k without further work, and by construction we have $\mathbb{E}[\varphi_k | \mathscr{F}_{i-1}] = 0$, which settles (*iii*). In order to prove (*ii*), we observe

$$\begin{split} ||\varphi_{k} - p(n_{k} \cdot)||_{\infty} &\ll \max_{\nu = 0, \dots, 2^{m(B_{i-1})} - 1} \frac{1}{|I_{\nu,k}|} \left| \int_{I_{\nu,k}} \hat{\varphi}_{k}(t) dt \right| + \frac{1}{h(N)^{K/2}} \\ &\ll \max_{\nu = 0, \dots, 2^{m(B_{i-1})} - 1} \frac{1}{|I_{\nu,k}|} \left| \int_{I_{\nu,k}} p(n_{k}t) dt \right| + \frac{1}{h(N)^{K/2}} \\ &\ll \frac{2^{m(B_{i-1})}}{n_{k}} + \frac{1}{h(N)^{K/2}} \\ &\ll \frac{1}{h(N)^{K/2}}, \end{split}$$

where we used Lemma 2.2 in the second to last line and the definition of Δ'_{i-1} in the last step.

Next, we use those quantities $(\varphi_k)_{k \in \Delta_i}$ to construct a suitable martingale. Let $1 \le i \le M = M(N)$ and define

(12)

$$Y_i(x) = \sum_{k \in \Delta_i} c_k \varphi_k(x), \quad T_i(x) = \sum_{k \in \Delta_i} c_k p(n_k x), \quad T'_i(x) = \sum_{k \in \Delta'_i} c_k p(n_k x), \quad V_M(x) = \sum_{i=1}^M \mathbb{E}[Y_i^2 | \mathscr{F}_{i-1}], \quad x \in \mathbb{T}.$$

By construction, $(Y_i)_{i \in \mathbb{N}}$ is adapted to $(\mathscr{F}_i)_{i \in \mathbb{N}}$, and since $\mathbb{E}[Y_i|\mathscr{F}_{i-1}] = 0$, the process $(Y_i)_{i \in \mathbb{N}}$ is a martingale difference sequence. This means the sequence $X_M = \sum_{i=1}^M Y_i$ for $M \in \mathbb{N}$ is a martingale which puts us in the position to verify the other assumptions of Proposition 2.1. Further, let

(13)
$$w_i = \int_{\mathbb{T}} \left(\sum_{k \in \Delta_i} c_k p(n_k x) \right)^2 \mathrm{d}x, \quad s_M = \left(\sum_{i=1}^M w_i \right)^{1/2}.$$

Note that the functions φ_k as well as the filtration $(\mathscr{F}_i)_{i \in \mathbb{N}}$ (and therefore also Y_i, V_M) depend on the value of *K* chosen in Lemma 2.4. In various estimates below, we will assume implicitly that *K* is sufficiently large. We remark that the finally chosen value of *K* is absolute and could be determined explicitly. With this in mind, we can prove the following estimate:

Lemma 2.5. For any $1 \le i \le M$ and for any C > 0, we can find a K > 0 such that for $Y_i = Y_i(K)$ we have

$$||T_i^2 - Y_i^2||_{\infty} \ll \frac{1}{h(N)^C}.$$

Proof. We write

$$\begin{split} \|T_i^2 - Y_i^2\|_{\infty} &= \sup_{x \in \mathbb{T}} \left(\left(\sum_{k \in \Delta_i} \left(c_k(p(n_k x) - \varphi_k(x)) \right) \right) \left(\sum_{k \in \Delta_i} c_k(p(n_k x) + \varphi_k(x)) \right) \right) \\ &\ll \#\Delta_i \sum_{k \in \Delta_i} c_k \sup_{x \in \mathbb{T}} |p(n_k x) - \varphi_k(x)| \\ &\ll \frac{\#\Delta_i^2}{h(N)^{K/2}} \leq \frac{N^2}{h(N)^{K/2}} \leq \frac{1}{h(N)^C}. \end{split}$$

Here we used (ii) of Lemma 2.4 (for a sufficiently large K > 0) together with $h(N) \ge N^{1-2\beta}$. The following lemma will be the key ingredient in order to apply Proposition 2.1.

Lemma 2.6. Let $(V_M)_{M \in \mathbb{N}}$ and $(s_M)_{M \in \mathbb{N}}$ be defined as in (12) and (13). Then we have

(14)
$$\|V_M - s_M^2\|_2^2 = o(h(N)^2)$$

where the implied constant only depends on p.

Proof. Using Lemma 2.5 we obtain

(15)
$$\|V_{M} - s_{M}^{2}\|_{2} = \left\|\sum_{i=1}^{M} \mathbb{E}[Y_{i}^{2}|\mathscr{F}_{i-1}] - s_{M}^{2}\right\|_{2}$$
$$\leq \left\|\sum_{i=1}^{M} \mathbb{E}[T_{i}^{2}|\mathscr{F}_{i-1}] - s_{M}^{2}\right\|_{2} + \frac{M}{h(N)}$$
$$= \left\|\sum_{i=1}^{M} \mathbb{E}[T_{i}^{2} - w_{i}|\mathscr{F}_{i-1}]\right\|_{2} + \frac{M}{h(N)}.$$

Since (11) ensures $\frac{M}{h(N)} = o(h(N))$, it remains to examine $T_i^2 - w_i$ more closely. For any *i*, we observe

$$\begin{split} T_i^2(x) - w_i &= \left(\sum_{k \in \Delta_i} c_k \sum_{j=1}^d a_j \cos(2\pi j n_k x)\right)^2 - w_i \\ &\ll \sum_{\substack{k \le k' \in \Delta_i, 1 \le j, j' \le d \\ 0 < |jn_k - j'n_{k'}| < h(N)^{-K} n_{B_{i-1}}}} c_k c_{k'} a_j a_{j'} \cos(2\pi (jn_k - j'n_{k'}) x) \\ &+ \sum_{\substack{k \le k' \in \Delta_i, 1 \le j, j' \le d \\ h(N)^{-K} n_{B_{i-1}} \le |jn_k - j'n_{k'}| < n_{A_i}}} c_k c_{k'} a_j a_{j'} \cos(2\pi (jn_k - j'n_{k'}) x) \\ &+ \sum_{\substack{k \le k' \in \Delta_i, 1 \le j, j' \le d \\ n_{A_i} \le |jn_k - j'n_{k'}|}} c_k c_{k'} a_j a_{j'} \cos(2\pi (jn_k - j'n_{k'}) x) \\ &=: U_i(x) + W_i(x) + R_i(x). \end{split}$$

This means, for the quantity in (15), we get

(16)
$$\|V_M - s_M^2\|_2 \le \|\sum_{i=1}^M \mathbb{E}[U_i|\mathscr{F}_{i-1}]\|_2 + \|\sum_{i=1}^M \mathbb{E}[W_i|\mathscr{F}_{i-1}]\|_2 + \|\sum_{i=1}^M \mathbb{E}[R_i|\mathscr{F}_{i-1}]\|_2 + \frac{M}{h(N)}.$$

Starting with the estimate for R_i , we see that there are at most $2d^2 |\Delta_i|^2$ many such terms and thus we get by Lemma 2.2

(17)
$$\|\mathbb{E}[R_i|\mathscr{F}_{i-1}]\|_{\infty} \le 4d^2 |\Delta_i|^2 \frac{2^{m(B_{i-1})}}{n_{A_i}} \ll \frac{1}{N},$$

where the last estimate follows from our choice of *K* in the definition of $m(B_{i-1})$ (recall the definition of m(k) in Lemma 2.4). Using the triangle inequality and the monotonicity of the *p*-norms, this implies

$$\left\|\sum_{i=1}^{M} \mathbb{E}[R_i|\mathscr{F}_{i-1}]\right\|_2 \ll \frac{M}{N}$$

Next, we examine $\|\sum_{i=1}^{M} \mathbb{E}[U_i|\mathcal{F}_{i-1}]\|_2$. Since each U_i is a sum of cosines with frequencies ranging from 1 to $h(N)^{-K} n_{B_{i-1}}$, we can write

$$U_i(x) = \sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} c_{u,i} \cos(2\pi u x)$$

where

$$c_{u,i} = \sum_{\substack{j,j' \leq d \\ k,k' \in \Delta_i}} c_k c'_k a_j a'_j \mathbb{1}_{[jn_k - j'n_{k'} = u]}.$$

Thus, expressing

$$\sum_{i=1}^{M} U_i(x) = \sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} d_u \cos(2\pi u x)$$

where $d_u = \sum_{i=1}^{M} c_{u,i}$, we can apply Parseval's identity to derive

$$\left\|\sum_{i=1}^{M} U_{i}\right\|_{2}^{2} = \sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} d_{u}^{2} \le \max_{\nu > 0} |d_{\nu}| \sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} |d_{u}|.$$

In the following, we first establish an estimate for $\max_{v>0} |d_v|$ and then for the sum over all $|d_u|$. Indeed, we get

$$\begin{split} |d_{\nu}| &\leq \sum_{j,j' \leq d} |a_{j}a'_{j}| \sum_{k,k' \leq N} c_{k}c'_{k} \mathbb{1}_{[jn_{k}-j'n_{k'}=\nu]} \\ &\leq 2||p||_{2}^{2} \sum_{j,j' \leq d} \sum_{k,k' \leq N} c_{k}c'_{k} \mathbb{1}_{[jn_{k}-j'n_{k'}=\nu]} \\ &= o(h(N)), \end{split}$$

where we have used the elementary estimate $|a_j a_{j'}| \le 2||p||_2^2$ and the Diophantine condition from Assumption 3 (that is also included in Assumption 4) in the last step. From that we infer

(18)
$$\max_{v \ge 0} |d_v| = o(h(N)).$$

Next, we get (using $h(N)^{-K} n_{B_{i-1}} \le n_{A_i}$, which holds by the very definition of A_i and B_{i-1})

$$\sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} |d_u| \leq \sum_{i=1}^{M} \sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} |c_{u,i}|$$

$$\leq \sum_{i=1}^{M} \sum_{j,j' \leq d} |a_j a_{j'}| \sum_{\substack{k,k' \in \Delta_i \\ |j n_k - j' n_{k'}| \leq n_{A_i}}} c_k c_{k'}$$

$$\ll \sum_{i=1}^{M} \sum_{j,j' \leq d} \sum_{\substack{k \leq k' \in \Delta_i \\ |j n_k - j' n_{k'}| \leq n_{A_i}}} c_k c_{k'}.$$

Observe that if $k \le k'$ then $j'n_{k'} - jn_k \ge n_{k'} - dn_k \ge n_k q^{k'-k} - dn_k$. Thus, for $k' \ge k + \log_q(2d)$, we have $|j'n_{k'} - jn_k| \ge 2dn_k - dn_k > n_{A_i}$. Therefore it follows that

(19)
$$\sum_{\substack{k \le k' \in \Delta_i \\ |jn_k - j'n_{k'}| \le n_{A_i}}} c_k c_{k'} \le \sum_{k \in \Delta_i} \sum_{k \le k' \le k + \log_q(2d)} c_k c_{k'} \ll \sum_{k \in \Delta_i} c_k^2.$$

Thus,

$$\sum_{u=1}^{\lfloor h(N)^{-K}} n_{B_{i-1}} \rfloor d_u | \leq \sum_{i=1}^{M} \sum_{u=1}^{\lfloor h(N)^{-K}} n_{B_{i-1}} \rfloor |c_{u,i}| \leq 2d ||p||_{\infty} \sum_{i=1}^{M} \sum_{k \in \Delta_i} c_k^2 \ll h(N).$$

Combining the estimate above with (18) now yields

$$\left\|\sum_{i=1}^{M} U_{i}\right\|_{2}^{2} \leq \sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} d_{u}^{2} \leq \max_{u} |d_{u}| \sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} |d_{u}| = o_{d}(h(N)^{2})$$

As a byproduct of the previous argument, it follows that for any i = 1, ..., M, we have $\sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} |c_{u,i}| \ll h(N)$. In order to estimate for $\|\sum_{i=1}^{M} \mathbb{E}[U_i | \mathcal{F}_{i-1}]\|_2^2$, we observe that the fluctuations of U_i over an arbitrary atom of \mathcal{F}_{i-1} are bounded from above by

$$\sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} |c_{u,i}| \left| 2\pi u 2^{-m(B_{i-1})} \right| \ll \frac{n_{B_{i-1}}}{h(N)^{K}} 2^{-m(B_{i-1})} \sum_{u=1}^{\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor} |c_{u,i}|$$

$$\ll rac{n_{B_{i-1}}}{h(N)^{K-1}2^{m(B_{i-1})}} \ \ll rac{1}{N},$$

where the estimate in the last line uses the definition of $m(B_{i-1})$ together with the choice of *K*. This means we have established

(20)
$$\left\|\sum_{i=1}^{M} \mathbb{E}[U_i|\mathscr{F}_{i-1}]\right\|_2 \ll \left\|\sum_{i=1}^{M} U_i\right\|_2 + \frac{1}{N} = o_d(h(N)).$$

Next, we estimate $\left\|\sum_{i=1}^{M} \mathbb{E}[W_i | \mathcal{F}_{i-1}]\right\|_2$. We note that

$$\left\|\sum_{i=1}^{M} \mathbb{E}[W_i|\mathscr{F}_{i-1}]\right\|_2^2 \leq 2 \mathbb{E}\left[\sum_{1 \leq i, i' \leq M} \mathbb{E}[W_i|\mathscr{F}_{i-1}] \mathbb{E}[W_{i'}|\mathscr{F}_{i'-1}]\right].$$

The diagonal terms where i = i' are equal to

$$\mathbb{E}\left[\sum_{i=1}^{M} \mathbb{E}[W_i|\mathcal{F}_{i-1}]^2\right].$$

Since $W_i(x) = \sum_{u=\lfloor h(N)^{-K} n_{B_{i-1}} \rfloor+1}^{n_{A_i}} c_u \cos(2\pi u x)$, an analogous calculation as in (19), combined with Jensen's inequality for the conditional expectation, reveals that

(21)

$$\mathbb{E}\left[\sum_{i=1}^{M} \mathbb{E}[W_i|\mathscr{F}_{i-1}]^2\right] \leq \sum_{i=1}^{M} ||W_i||_2^2$$

$$\ll \sum_{i=1}^{M} \left(\sum_{k \in \Delta_i} c_k^2\right)^2$$

$$\ll h(N)^{\gamma} \sum_{i=1}^{M} \sum_{k \in \Delta_i} c_k^2$$

$$\ll h(N)^{3/2},$$

where we have used (11) in the second to last line. So we are left to treat the case where i < i'. Since $\mathbb{E}[W_i | \mathcal{F}_{i-1}]$ is \mathcal{F}_{i-1} -measurable, we get

$$\begin{split} \left| \mathbb{E} \left[\mathbb{E}[W_i | \mathcal{F}_{i-1}] \mathbb{E}[W_{i'} | \mathcal{F}_{i'-1}] | \mathcal{F}_{i-1} \right] \right| &= |\mathbb{E}[W_i | \mathcal{F}_{i-1}] \mathbb{E}[W_{i'} | \mathcal{F}_{i-1}]| \\ &\leq ||W_i||_{\infty} |\mathbb{E}[W_{i'} | \mathcal{F}_{i-1}]| \\ &\leq \left(\sum_{k \in \Delta_i} c_k^2 \right) |\mathbb{E}[W_{i'} | \mathcal{F}_{i-1}]| \\ &\ll \sqrt{h(N)} |\mathbb{E}[W_{i'} | \mathcal{F}_{i-1}]|. \end{split}$$

Again, the second to last inequality utilizes an analogous argument as in (19) whereas the last estimate uses (11). Let $W_{i'}(x) = \sum_{u=\lfloor h(N)^{-K} n_{B_{i'-1}} \rfloor + 1}^{n_{A_{i'}}} c_u \cos(2\pi u x)$ and recall that

$$\sum_{u=\lfloor h(N)^{-K}n_{B_{i'-1}}\rfloor+1}^{n_{A_{i'}}}|c_u|\ll \sum_{k\in\Delta_{i'}}c_k^2\ll\sqrt{h(N)}.$$

By Lemma 2.2 for $f(x) = \cos(2\pi x)$ together with the estimate above, we obtain

$$\begin{split} |\mathbb{E}[W_{i'}|\mathscr{F}_{i-1}]| &\leq \sum_{u=\lfloor h(N)^{-K}n_{B_{i'-1}}\rfloor+1}^{n_{A_{i'}}} |c_u|u^{-1}2^{m(B_{i-1})} \\ &\leq \frac{2^{m(B_{i-1})}}{h(N)^{-K}n_{B_{i'-1}}} \sum_{u} |c_u| \\ &\ll \frac{2^{m(B_{i-1})}}{h(N)^{-K}n_{B_{i'-1}}} h(N) \\ &\leq h(N)^{K+1}\frac{n_{B_{i-1}}}{n_{B_{i'-1}}} \\ &\ll \frac{1}{h(N)^{K}}. \end{split}$$

In the second-to-last step, we applied the definition of $m(B_{i-1})$ and completed our estimate using i' > i, which implies $\frac{n_{B_{i-1}}}{n_{B_{i'-1}}} \le q^{-h(N)^{\gamma}}$ which goes faster to zero than any polynomial power in h(N) can grow.

Combining (21) and (22) we obtain

(22)

$$\begin{split} \left\|\sum_{i=1}^{M} \mathbb{E}[W_i|\mathscr{F}_{i-1}]\right\|_2^2 &\leq \mathbb{E}\left[\sum_{1 \leq i \leq M} \mathbb{E}[W_i|\mathscr{F}_{i-1}]^2\right] + 2\mathbb{E}\left[\sum_{1 \leq i < i' \leq M} \mathbb{E}[W_i|\mathscr{F}_{i-1}]\mathbb{E}[W_{i'}|\mathscr{F}_{i'-1}]\right] \\ &\qquad \ll h(N)^{3/2} + \sum_{1 \leq i < i' \leq M} \frac{1}{h(N)^K} \\ &\ll h(N)^{3/2}, \end{split}$$

where we used (11), $\gamma < 1/2$ and that K > 0 is sufficiently large. This finishes our analysis of the terms $(W_i)_{i=1}^M$.

Aggregating the previous estimate on the W_i and (20), for the expression in (16), we finally end up with

$$\|V_M - s_M^2\|_2^2 = o(h(N)^2) + h(N)^{3/2},$$

which proves the claim.

The following lemma is used several times in the sequel.

Lemma 2.7. Let $1 \le j, j' \le d$, then we have

$$\sup_{c\in\mathbb{N}}\sum_{k,k'\in\Delta_i}c_kc_{k'}\mathbbm{1}_{[jn_k-j'n_{k'}=c]}\leq \sum_{k\in\Delta_i}c_k^2.$$

Proof. For fixed $1 \le j, j' \le d, c \in \mathbb{N}$ and for fixed $1 \le k \le N$, there is at most one k' such that $jn_k - j'n_{k'} = c$ and clearly this k' is distinct for different k. Thus this defines an injection k' = k'(c, j, j') from $A_i = A_i(c, j, j') := \{k \in \Delta_i : \exists k' \in \Delta_i : jn_k - j'n_{k'} = c\}$ to Δ_i . We obtain

$$\sum_{k,k'\in\Delta_{i}} c_{k}c_{k'}\mathbb{1}_{[jn_{k}-j'n_{k'}=c]} = \sum_{k\in A_{i}} \underbrace{c_{k}c_{k'(k)}}_{\leq \frac{1}{2}(c_{k}^{2}+c_{k'(k)}^{2})} \underbrace{\mathbb{1}_{[jn_{k}-j'n_{k'(k)}=c]}}_{\leq 1}$$
$$\leq \sum_{k\in\Delta_{i}} c_{k}^{2}.$$

Since the estimate is uniform in $c \in \mathbb{N}$, the claim follows.

Proposition 2.1 requires a bound on $\sum_{i=1}^{M} \mathbb{E}[Y_i^4]$. This is accomplished in the following axillary result. We mention that similar estimates have been obtained in [1] and [8, Lemma 2.2] under slightly more restrictive assumptions.

Proposition 2.8. Let $(Y_i)_{i=1,...,M}$ be the sequence of random variables constructed at the beginning of Section 2.3. Then there exists $\varepsilon > 0$ such that

$$\sum_{i=1}^{M} \mathbb{E}[Y_i^4] \ll h(N)^{2-\varepsilon}$$

where the implied constant only depends on p.

Proof. Since

$$\mathbb{E}[Y_i^4] = \mathbb{E}[Y_i^4 - T_i^4] + \mathbb{E}[T_i^4],$$

we can use Lemma 2.5 to obtain

$$\mathbb{E}[Y_i^4 - T_i^4] \le \|Y_i^2 - T_i^2\|_{\infty} \left(\|T_i^2\|_{\infty} + \|Y_i^2\|_{\infty}\right) \ll \frac{N}{h(N)^C}.$$

We remind ourselves that C > 0 can be chosen sufficiently large and by (11), we have $h(N) \ge N^{1-2\beta}$. This means the term in the previous display is negligibly small. So we are left to provide an upper bound for $\mathbb{E}[T_i^4]$. We recall that

$$T_i(x) = \sum_{k \in \Delta_i} c_k p(n_k x),$$

where $p(n_k x) = \sum_{j=1}^d a_j \cos(2\pi j n_k x)$. This means if we take the 4-th power of T_i and we use trigonometric identities such as $\cos(\alpha + \beta) = \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta))$, we end up with a large sum over cosines with respective frequencies of the form $\pm j_1 n_{k_1} \pm \ldots \pm j_4 n_{k_4}$, where $1 \le j_\ell \le d$ and $k_\ell \in \Delta_i$. Since we are only interested in $\mathbb{E} [T_i^4]$, solely those frequencies $\pm j_1 n_{k_1} \pm \ldots \pm j_4 n_{k_4}$ remain which are 0. This observation reveals

$$\mathbb{E}[T_i^4] \leq \sum_{k_1, k_2, k_3, k_4 \in \Delta_i} c_{k_1} c_{k_2} c_{k_3} c_{k_4} \sum_{1 \leq j_1, j_2, j_3, j_4 \leq d} |a_{j_1} a_{j_2} a_{j_3} a_{j_4}| \mathbb{1}_{[\pm j_1 n_{k_1} \pm j_2 n_{k_2} \pm j_3 n_{k_3} \pm j_4 n_{k_4} = 0]} \\ \ll \sum_{1 \leq j_1, j_2, j_3, j_4 \leq d} |a_{j_1} a_{j_2} a_{j_3} a_{j_4}| \sum_{k_1 \leq k_2 \leq k_3 \leq k_4 \in \Delta_i} c_{k_1} c_{k_2} c_{k_3} c_{k_4} \mathbb{1}_{[\pm j_1 n_{k_1} \pm j_2 n_{k_2} \pm j_3 n_{k_3} \pm j_4 n_{k_4} = 0]}.$$

An expression of the form $\pm j_1 n_{k_1} \pm j_2 n_{k_2} \pm j_3 n_{k_3} \pm j_4 n_{k_4}$ can only then be 0, if at least one sign in front of the j_i is positive; without loss of generality let it be the sign in front of j_4 . Thus we get $0 = j_4 n_{k_4} \pm j_1 n_{k_1} \pm j_2 n_{k_2} \pm j_3 n_{k_3} \ge n_{k_4} - 3dn_{k_3}$. This implies $n_{k_3} \ge \frac{n_{k_4}}{3d}$ and thus by the Hadamard-gap condition we have $k_3 \ge k_4 - \lceil \log_q(3d) \rceil$. Furthermore, for fixed (k_3, k_4) , for every k_2 , there exists at most one k_1 such that the equation $\pm j_1 n_{k_1} \pm j_2 n_{k_2} \pm j_3 n_{k_3} \pm j_4 n_{k_4} = 0$ is satisfied, and such a k_1 is distinct for different k_2 . This allows us to construct an injection $k_2 \mapsto k_1 k_{3,k_4}(k_2)$ which further implies

$$\begin{split} \sum_{k_1 \le k_2 \le k_3 \le k_4 \in \Delta_i} c_{k_1} c_{k_2} c_{k_3} c_{k_4} \mathbbm{1}_{[\pm j_1 n_{k_1} \pm j_2 n_{k_2} \pm j_3 n_{k_3} \pm j_4 n_{k_4} = 0]} \le \sum_{k_4 \in \Delta_i} c_{k_4} \sum_{k_4 - \lceil \log_q(3d) \rfloor \le k_3 \le k_4} c_{k_3} \sum_{k_2 \le k_3} c_{k_2} c_{k_{1k_3, k_4}(k_2)} \\ \ll \sum_{k_4 \in \Delta_i} c_{k_4} \sum_{k_4 - \lceil \log_q(3d) \rfloor \le k_3 \le k_4} c_{k_3} \sum_{k_2 \in \Delta_i} c_{k_2}^2 \\ \ll_d \left(\sum_{k \in \Delta_i} c_k^2 \right)^2 \\ = h(N)^{2\gamma}. \end{split}$$

Summing over *M* and using $M \simeq h(N)^{1-\gamma}$ (see (11)), we obtain

$$\sum_{i=1}^{M} \mathbb{E}[Y_i^4] \ll Mh(N)^{2\gamma} \asymp h(N)^{1+\gamma}$$

Since $\gamma < 1$, this finishes the proof.

We have now gathered all auxiliary results in order to finally prove Proposition 2.3.

Proof of Proposition 2.3. We provide a complete proof of Proposition 2.3 under Assumption 3. The corresponding claim under Assumption 4 follows mutatis mutandis. Let $N \in \mathbb{N}$ be given and choose M = M(N) such that $N \in \Delta_M \cup \Delta'_M$. Using the quantities from (12), we may write

$$\sum_{k=1}^{N} c_k p(n_k x) = \sum_{i=1}^{M} Y_i + \sum_{i=1}^{M} (T_i - Y_i) + \sum_{i=1}^{M} T'_i - \sum_{k=N+1}^{\hat{N}} c_k p(n_k x),$$

where $\hat{N} = \max \Delta'_M$. We shall now establish two properties. On the one hand, we show that under Assumption 3, the 2-norms of the quantities

$$\sum_{i=1}^{M} (T_i - Y_i), \quad \sum_{i=1}^{M} T'_i, \text{ and } \sum_{k=N+1}^{\hat{N}} c_k p(n_k x)$$

are all of order $o(||S_N||_2)$. On the other hand, we identify an asymptotically precise formula for s_M as $N \longrightarrow \infty$, which will imply in particular that s_M and $||S_N||_2$ are asymptotically equivalent. Starting with the latter, we note that for fixed $d \in \mathbb{N}$, $i, i' \in \mathbb{N}$ sufficiently large and $k \in \Delta_i, k' \in \Delta'_i, i \neq i'$, we have for all $1 \le j, j' \le d$ that $jn_k - j'n_{k'} \ne 0$. Hence, for sufficiently large $N \in \mathbb{N}$ and fixed $1 \le j, j' \le d$. d, we obtain

$$\begin{split} \sum_{i=1}^{M} \sum_{k,k' \in \Delta_{i}} c_{k} c_{k'} \mathbb{1}_{[jn_{k}-j'n_{k'}=0]} &= \sum_{k,k' \in \bigcup_{1 \le i \le M} \Delta_{i}} c_{k} c_{k'} \mathbb{1}_{[jn_{k}-j'n_{k'}=0]} \\ &= \sum_{k,k' \le N} c_{k} c_{k'} \mathbb{1}_{[jn_{k}-j'n_{k'}=0]} - \sum_{k \in \bigcup_{1 \le i \le M} \Delta_{i}'} \sum_{k' \ne k} c_{k} c_{k'} \mathbb{1}_{[jn_{k}-j'n_{k'}=0]} \\ &+ \sum_{k \in \Delta_{M}, k > N} \sum_{k' \ne k} c_{k} c_{k'} \mathbb{1}_{[jn_{k}-j'n_{k'}=0]}. \end{split}$$

Since $c_k \le 1$ and for fixed j, j', k, there is at most one k' satisfying $jn_k - j'n_{k'} = 0$, we obtain

. .

$$\sum_{k \in \bigcup_{1 \le i \le M} \Delta'_i} \sum_{k' \ne k} c_k c_{k'} \mathbb{1}_{[jn_k - j'n_{k'} = 0]} + \sum_{k \in \Delta_M, k > N} \sum_{k' \ne k} c_k c_{k'} \mathbb{1}_{[jn_k - j'n_{k'} = 0]} \ll \sum_{1 \le i \le M} |\Delta'_i| + |\Delta_M| = O(M \log N + h(N)^{\gamma}) = o(h(N)),$$

where we have used (11) and the property that each buffer block Δ'_i has at most logarithmic length. Thus, under Assumption 3, we have

$$s_{M}^{2} = \sum_{i=1}^{M} \int_{\mathbb{T}} \left(\sum_{k \in \Delta_{i}} c_{k} p(n_{k} x) \right)^{2} dx$$

$$= \sum_{i=1}^{M} \sum_{k,k' \in \Delta_{i}} c_{k} c_{k'} \sum_{j,j' \leq d} \frac{a_{j} a_{j'}}{2} \mathbb{I}_{[jn_{k}-j'n_{k'}=0]}$$

$$= \sum_{j,j' \leq d} \frac{a_{j} a_{j'}}{2} \sum_{i=1}^{M} \sum_{k,k' \in \Delta_{i}} c_{k} c_{k'} \mathbb{I}_{[jn_{k}-j'n_{k'}=0]} + o(h(N))$$

$$= \int_{\mathbb{T}} \left(\sum_{k \leq N} c_{k} p(n_{k} x) \right)^{2} dx + o(h(N))$$

$$= \left\| \left\| \sum_{k \leq N} c_{k} p(n_{k} x) \right\|_{2}^{2} (1 + o(1)).$$

where we used the estimate (9) from Assumption 3 in the last line. This shows that s_M and $||S_N||_2$ are asymptotically equivalent as $N \rightarrow \infty$. Next, we note that

$$\left\|\sum_{k=N+1}^{\hat{N}} c_k p(n_k x)\right\|_2 \ll \left(\sum_{k\in\Delta_M\cup\Delta'_M} c_k^2\right)^{1/2} \ll h(N)^{\gamma/2} = o\left(\sqrt{h(N)}\right).$$

We get using Parseval's identity (recall that $p(x) = \sum_{j=1}^{d} a_j \cos(2\pi j n_k x)$)

$$\begin{split} \left\| \sum_{i=1}^{M} T_i' \right\|_2 &= \left\| \sum_{i=1}^{M} \sum_{k \in \Delta_i'} c_k p(n_k x) \right\|_2 \\ &\leq \sum_{j=1}^{d} |a_j| \left\| \sum_{i=1}^{M} \sum_{k \in \Delta_i'} c_k \cos(2\pi j n_k) \right\|_2 \\ &\ll \left(\sum_{i=1}^{M} \sum_{k \in \Delta_i'} c_k^2 \right)^{1/2} \\ &= o(\sqrt{h(N)}). \end{split}$$

Lastly, using Lemma 2.4 (ii), we obtain

$$\begin{split} \left\| \sum_{i=1}^{M} Y_i - T_i \right\|_2 &\leq \sum_{i=1}^{M} \sum_{k \in \Delta_i} c_k \left\| p(n_k \cdot) - \varphi_k(\cdot) \right\|_2 \\ &\leq \frac{N}{h(N)^C} \\ &= o(\sqrt{h(N)}). \end{split}$$

Using the previous estimates on s_M together with Lemma 2.6 and Proposition 2.8, we have

$$\frac{\sum_{i=1}^{M} \mathbb{E}\left[Y_i^4\right] + \mathbb{E}\left[(V_M - s_M)^2\right]}{s_M^2} \longrightarrow 0, \quad \text{as } M \to \infty.$$

By Proposition 2.1 this implies

$$\frac{Y_1 + \ldots + Y_M}{\sqrt{s_M}} \xrightarrow{w} \mathcal{N}(0, 1), \quad \text{as } M \to \infty.$$

Taking all the previous estimates together, we have shown that under Assumption 3

$$\frac{\sum_{k \le N} c_k p(n_k x)}{\left|\left|\sum_{k \le N} c_k p(n_k x)\right|\right|_2} = \frac{\sum_{i=1}^M Y_i}{s_M} \cdot \frac{s_M}{\left|\left|\sum_{k \le N} c_k p(n_k x)\right|\right|_2} + r_N,$$

where r_N is a quantity with $||r_N||_2 \longrightarrow 0$. By (23), we have
$$\frac{s_M}{1 + \frac{s_M}{1 + \frac{s_M}$$

$$\frac{|\Sigma_{k\leq N} c_k p(n_k x)||_2}{||\Sigma_{k\leq N} c_k p(n_k x)||_2} \longrightarrow 1$$

which completes the proof of (i) by virtue of Slutsky's Theorem.

Corollary 2.9. Let the same assumptions as in Proposition 2.3 hold and let $A \subseteq [1, N]$ be a set such that we have ...

(i) ... Assumption 3 and

$$\liminf_{N\to\infty}\frac{\|\sum_{k\in A}c_kp(n_kx)\|_2}{\|\sum_{k\leq N}c_kp(n_kx)\|_2}>0.$$

(24)

(25)

Then

$$\frac{1}{\left|\left|\sum_{k\leq N} c_k p(n_k x)\right|\right|_2} \sum_{k\in A} c_k p(n_k x) \xrightarrow{w} \mathcal{N}(0, 1),$$

(ii) ... Assumption 4 and

$$\liminf_{N\to\infty}\frac{\sum_{k\in A}c_k^2}{\sum_{k\leq N}c_k^2}>0.$$

Then we have

$$\frac{1}{||p||_2 \sqrt{\sum_{k \in A} c_k^2}} \sum_{k \in A} c_k p(n_k x) \xrightarrow{w} \mathcal{N}(0, 1).$$

Proof. By restricting to $k \in A$, we take a subsequence of n_k which thus in particular satisfies the Hadamard gap condition. By (24) and (25), the respective Assumptions 3 and 4 also hold in this restricted setup. Therefore, an application of Proposition 2.3 proves the claim.

2.4. The clt in the general case. In this section, we stepwise extend the central limit theorem from Proposition 2.3 to the more general settings of our main results, Theorems 1 and 2. We begin by removing the (asymptotic) lower bound on c_k .

Lemma 2.10. Let $p(x) = \sum_{j=1}^{d} \cos(2\pi j x)$ be a mean zero even trigonometric polynomial and, for $N \in \mathbb{N}$, let $(c_k)_{k \leq N}$ be a scheme of weights satisfying Assumption 2. Then we have the following:

(i) If Assumption 3 is satisfied (with f = p), then

$$\frac{\sum_{k \le N} c_k p(n_k x)}{\left|\left|\sum_{k \le N} c_k p(n_k x)\right|\right|_2} \xrightarrow{w} \mathcal{N}(0, 1).$$

(ii) If Assumption 4 is satisfied, then

$$\frac{\sum_{k \le N} c_k p(n_k x)}{\sqrt{h(N)} ||p||_2} \xrightarrow{w} \mathcal{N}(0, 1).$$

Proof. We can assume without loss of generality that $c_k \ge N^{-1/2}$ for all k: This holds since we get by Lemma 2.7

$$\left\|\sum_{\substack{k\leq N\\c_k\leq N^{-1/2}}} c_k p(n_k x)\right\|_2^2 \leq \|p\|_2^2 \sum_{\substack{k\leq N\\c_k\leq N^{-1/2}}} c_k^2 \leq \|p\|_2^2$$

and thus after dividing by the diverging quantity $\sqrt{h(N)}$, this term converges to 0 in probability. We define

$$C = \left\{ k \le N : N^{-1/4 - \delta} \le c_k \le N^{-1/4} \right\}$$

where $\delta < 1/12$. Further, we define $L = L(N) = \lfloor \sqrt{\log N} \rfloor$ and partition C into $C = \bigcup_{\ell=0}^{L-1} C_{\ell}$ where

$$C_{\ell} := \left\{ k \in C : N^{-1/4 - \delta(\ell+1)/L} < c_k \le N^{-1/4 - \delta(\ell)/L} \right\}, \quad \ell = 0, \dots, L - 1$$

Since, trivially $\sum_{k \in C} c_k^2 \le h(N)$, by the pigeonhole principle there is some $\ell_0 \in \{0, \dots, L-1\}$ such that

$$\sum_{k \in C_{\ell_0}} c_k^2 \le \frac{h(N)}{L}$$

Now let $\beta_{\ell_0} = \beta_{\ell_0}(N) = \frac{1}{4} + \delta \frac{\ell_0}{L}$ and define

$$A = \left\{ k \le N : N^{-1/2} \le c_k \le N^{-\beta_{\ell_0} - \frac{\delta}{L}} \right\}, \quad B = \left\{ k \le N : c_k \ge N^{-\beta_{\ell_0}} \right\}.$$

We can now write

$$\sum_{k=1}^{N} c_k p(n_k x) = \sum_{k \in A} c_k p(n_k x) + \sum_{k \in B} c_k p(n_k x) + \sum_{k \in C_{\ell_0}} c_k p(n_k x)$$
$$=: S_N^A(x) + S_N^B(x) + \sum_{k \in C_{\ell_0}} c_k p(n_k x).$$

By Lemma 2.7, we have

$$\left\|\sum_{k\in C_{\ell_0}} c_k p(n_k x)\right\|_2^2 \le ||p||_2^2 \left\|\sum_{k\in C_{\ell_0}} c_k^2\right\|_2^2 \le \frac{||p||_2^2 h(N)}{L} = o(h(N)).$$

This shows that under Assumption 3 we have the following limit in probability:

$$\lim_{N \to \infty} \frac{\sum_{k \in C_{\ell_0}} c_k p(n_k x)}{\|\sum_{k \le N} c_k p(n_k x)\|_2} = 0.$$

Similarly, under Assumption 4 we obtain

$$\lim_{N \to \infty} \frac{\sum_{k \in C_{\ell_0}} c_k p(n_k x)}{||p||_2 \sqrt{h(N)}} = 0.$$

Thus, we are left to analyze the asymptotic stochastic behavior of S_N^A and S_N^B as $N \to \infty$. In the following, we provide a proof under Assumption 4. The claim under Assumption 3 can be shown under minimal changes. We introduce the following quantities

$$\sigma_N^A = \|p\|_2 \sqrt{\sum_{k \in A} c_k^2}, \quad \sigma_N^B = \|p\|_2 \sqrt{\sum_{k \in B} c_k^2}, \quad \sigma_N = \sqrt{(\sigma_N^A)^2 + (\sigma_N^B)^2}.$$

Let us assume that $\frac{S_N}{||p||_2 \sqrt{h(N)}}$ does not converge to the normal distribution as $N \longrightarrow \infty$. Then there exists a sequence $(N_k)_{k \in \mathbb{N}}$ and some $t \in \mathbb{R}$ such that

$$\lim_{k \to \infty} \mathbb{P}\left[\frac{S_{N_k}}{||p||_2 \sqrt{h(N_k)}} \le t\right] \ne \Phi(t),$$

where Φ denotes the standard normal distribution function. Let $\lambda_N := \frac{\sigma_N^A}{\sigma_N}$ which satisfies $0 \le \lambda_N \le 1$. Then there exists a subsequence $(N_{k_\ell})_{\ell \in \mathbb{N}} \subseteq (N_k)_{k \in \mathbb{N}}$ and some $\bar{\lambda} \in [0, 1]$ such that $\lambda_{N_{k_\ell}} \to \bar{\lambda}$. We distinguish the following cases:

Case 1: $\bar{\lambda} = 0$. In that case, we have $\sigma^B_{N_{k_\ell}} / \sigma_{N_{k_\ell}} \longrightarrow 1$ and using Lemma 2.7 shows

$$\frac{S_{N_{k_{\ell}}^{A}}}{\sigma_{N_{k_{\ell}}}} = \Lambda_{N_{k_{\ell}}} \frac{S_{N_{k_{\ell}}}^{A}}{\sigma_{N_{k_{\ell}}}^{A}} \stackrel{\mathbb{P}}{\longrightarrow} 0.$$

Consequently, this leads to

$$\frac{S_{N_{k_{\ell}}}^B}{\sigma_{N_{k_{\ell}}}} = (1 - o(1)) \frac{S_{N_{k_{\ell}}}^B}{\sigma_{N_{k_{\ell}}}^B}$$

By Corollary 2.9, we now have

$$\frac{S_{N_{k_{\ell}}}}{\sigma_{N_{k_{\ell}}}} = \frac{S_{N_{k_{\ell}}}^A}{\sigma_{N_{k_{\ell}}}} + \frac{S_{N_{k_{\ell}}}^B}{\sigma_{N_{k_{\ell}}}^B} (1 - o(1)) \xrightarrow{w} \mathcal{N}(0, 1),$$

where we have employed Slutsky's theorem. This contradicts our assumption of

$$\lim_{k \to \infty} \mathbb{P}\left[\frac{S_{N_k}}{||p||_2 \sqrt{h(N_k)}} \le t\right] \neq \Phi(t).$$

Case 2: $\bar{\lambda} = 1$. Analogously to before, we get $\frac{\sigma_{N_{k_{\ell}}^A}}{\sigma_{N_{k_{\ell}}}} \rightarrow 1$ and thus

$$\lim_{\ell \to \infty} \frac{S_{N_{k_{\ell}}}}{\sigma_{N_{k_{\ell}}}} \stackrel{w}{=} \lim_{\ell \to \infty} \frac{S_{N_{k_{\ell}}}^A}{\sigma_{N_{k_{\ell}}}^A},$$

provided the latter exists. Let $c_k \in A$, then $\tilde{c}_k := N^{1/4} c_k$ satisfies $N^{-1/4} \leq \tilde{c}_k \leq 1$. We have

$$\frac{S_{N_{k_{\ell}}}^{A}}{\sigma_{N_{k_{\ell}}}^{A}} = \frac{1}{||p||_{2}\sqrt{\sum_{k \in A} c_{k}^{2}}} \sum_{\substack{k=1\\k \in A}}^{N_{k_{\ell}}} c_{k}p(n_{k}x) = \frac{1}{||p||_{2}\sqrt{\sum_{k \in A} \tilde{c}_{k}^{2}}} \sum_{\substack{k=1\\k \in A}}^{N_{k_{\ell}}} \tilde{c}_{k}p(n_{k}x)$$

The right-hand side above converges to a standard normal distributed random variable by Corollary 2.9, a contradiction.

Case 3: $0 < \overline{\lambda} < 1$. Then there exists $\rho > 0$ such that for all sufficiently large $\ell \in \mathbb{N}$, we have

$$\frac{\sigma_{N_{k_{\ell}}}^{A}}{\sigma_{N_{k_{\ell}}}} > \rho, \quad \frac{\sigma_{N_{k_{\ell}}}^{B}}{\sigma_{N_{k_{\ell}}}} > \rho$$

By rescaling the c_k in A as in Case 2 and applying Corollary 2.9, we thus have

$$\frac{S^B_{N_{k_{\ell}}}}{\sigma^B_{N_{k_{\ell}}}} \xrightarrow{w} \mathcal{N}(0,1), \qquad \frac{S^A_{N_{k_{\ell}}}}{\sigma^A_{N_{k_{\ell}}}} \xrightarrow{w} \mathcal{N}(0,1).$$

In general, it is not clear that the sum of both random variables above converges to a sum of normal distributed random variables. In order to ensure such a limit, we need to show that $S^A_{N_{k_\ell}}$ and $S^B_{N_{k_\ell}}$ are asymptotically uncorrelated (this is the whole point in the construction of *C* to separate *A* and *B*).

Let $1 \le j, j' \le d$ and recall that, by lacunarity, $n_k j - n_{k'} j' = 0$ is only possible if $|k - k'| \le \log_q(d) \le d$. This implies

$$\begin{split} \mathbb{E}[S^{A}_{N_{k_{\ell}}}S^{B}_{N_{k_{\ell}}}] &= \sum_{j,j' \leq d} a_{j}a_{j'}\sum_{k \in A, k' \in B} c_{k}c_{k'}\mathbbm{1}_{[n_{k}j-n_{k'}j'=0]} \\ &\leq \sum_{j,j' \leq d} a_{j}a_{j'}\sum_{k' \in B} c_{k'}\sum_{\substack{k \in A \\ |k-k'| \leq d}} c_{k} \\ &\leq d\sum_{j,j' \leq d} a_{j}a_{j'}N_{k_{\ell}}^{-\beta_{\ell_{0}}-\frac{\delta}{L}}\sum_{k' \in B} c_{k'}. \end{split}$$

Since all $c_{k'} \in B$ satisfy $c_{k'} \ge N_{k_{\ell}}^{-\beta_{\ell_0}}$, we have

$$\sum_{k' \in B} c_{k'}^2 \ge \left(N_{k_\ell}^{-\beta_{\ell_0}} \right)^2 |B| \iff |B| \le \frac{1}{\left(N_{k_\ell}^{-\beta_{\ell_0}} \right)^2} \cdot \sum_{k' \in B} c_{k'}^2.$$

Therefore, by the Cauchy-Schwarz inequality

$$\sum_{\substack{k'\in B}} c_{k'} \leq \sqrt{|B| \cdot \sum_{k'\in B} c_{k'}^2} \leq N_{k_\ell}^{\beta_{\ell_0}} \sum_{k'\in B} c_{k'}^2 \leq N_{k_\ell}^{\beta_{\ell_0}} \cdot h(N_{k_\ell})$$

Consequently, we obtain

(26)
$$\mathbb{E}\left[S_{N_{k_{\ell}}}^{A}S_{N_{k_{\ell}}}^{B}\right] \leq d\sum_{j,j'\leq d} a_{j}a_{j'}h(N_{k_{\ell}})N_{k_{\ell}}^{-\delta/L} \ll_{d} h(N_{k_{\ell}})N_{k_{\ell}}^{-\delta/L} = o(\sigma_{N_{k_{\ell}}}^{2}).$$

Since $\sigma^A_{N_{k_\ell}}\sigma^B_{N_{k_\ell}} \ge \rho^2\sigma^2_{N_{k_\ell}}$, this implies

(27)
$$\lim_{\ell \to \infty} \frac{\mathbb{E}\left[S_{N_{k_{\ell}}}^{A} S_{N_{k_{\ell}}}^{B}\right]}{\sigma_{N_{k_{\ell}}}^{A} \sigma_{N_{k_{\ell}}}^{B}} = 0$$

This shows that $S^A_{N_{k_\ell}}$ and $S^B_{N_{k_\ell}}$ are asymptotically uncorrelated, and thus we get

$$\frac{S_{N_{k_\ell}}}{\sigma_{N_{k_\ell}}} \xrightarrow{w} \mathcal{N}(0, \bar{\lambda} + (1 - \bar{\lambda})) = \mathcal{N}(0, 1).$$

This contradicts our assumption and thus proves the claim in Lemma 2.10 (ii).

Having the central limit theorem settled for lacunary sums generated by an even trigonometric polynomial, we can now move on and prove Theorem 2.

Proof of Theorem 2. Let us start with the case, where f(x) is a general mean 0 trigonometric polynomial, i.e.

$$f(x) = \sum_{j=1}^{d} \left[a_j \cos(2\pi j x) + b_j \sin(2\pi j x) \right] = \sum_{j=1}^{d} a_j \cos(2\pi j x) + \sum_{j=1}^{d} b_j \sin(2\pi j x) = f_1(x) + f_2(x),$$

and hence

$$S_N = \sum_{k=1}^N c_k f_1(n_k x) + \sum_{k=1}^N c_k f_2(n_k x) =: S_N^{(1)} + S_N^{(2)}.$$

The summands $S_N^{(i)}$, i = 1, 2 are uncorrelated and hence on the scale of the central limit theorem, we can look at each $S_N^{(i)}$ separately. By Lemma 2.10, it immediately follows that $S_N^{(1)}$ satisfies a clt, when divided by its standard deviation. The clt for $S_N^{(2)}$ can be obtained under insignificant changes and thus $\frac{S_N}{||f||_2\sqrt{h(N)}}$ satisfies a clt.

Now let us consider a general $f \in \mathbb{L}_2(\mathbb{T})$ satisfying Assumption 1. By renormalizing, we can assume without loss of generality that we can write

$$f(x) = \sum_{j=1}^{\infty} \left[a_j \cos(2\pi j x) + b_j \sin(2\pi j x) \right]$$

with $\max\{|a_j|, |b_j|\} \le \frac{1}{j^{\rho}}$ for all $j \in \mathbb{N}$. Let $p(x) = \sum_{j=1}^d \left[a_j \cos(2\pi j x) + b_j \sin(2\pi j x)\right]$ and $r(x) = \sum_{j=d+1}^\infty \left[a_j \cos(2\pi j x) + b_j \sin(2\pi j x)\right]$ and let us define

$$S_{N} = \sum_{k=1}^{N} c_{k} p(n_{k} x) + \sum_{k=1}^{N} c_{k} r(n_{k} x)$$

=: $S_{N}^{p} + S_{N}^{r}$.

We write $S_N^r = S_{N,e}^r + S_{N,o}^r$ where

$$S_{N,e}^{r} = \sum_{k \le N} c_k \sum_{j=d+1}^{\infty} \left[a_j \cos(2\pi j x) \right], \quad S_{N,o}^{r} = \sum_{k \le N} c_k \sum_{j=d+1}^{\infty} \left[b_j \sin(2\pi j x) \right].$$

Clearly, by orthogonality, $||S_N^r||_2 = ||S_{N,e}^r||_2 + ||S_{N,o}^r||_2$; We will only provide an estimate for $||S_{N,e}^r||_2$, the estimate for $||S_{N,o}^r||_2$ works completely analogously.

We observe that

$$\begin{split} \|S_{N,e}^{r}\|_{2}^{2} &= \sum_{1 \leq k,k' \leq N} c_{k}c_{k'} \sum_{j,j' > d} a_{j}a_{j'} \mathbb{1}_{[jn_{k}-j'n_{k'}=0]} \\ &= \sum_{1 \leq k \leq N} c_{k}^{2} \sum_{j > d} a_{j}^{2} + 2 \sum_{1 \leq k < k' \leq N} c_{k}c_{k'} \sum_{j,j' > d} a_{j}a_{j'} \mathbb{1}_{[jn_{k}-j'n_{k'}=0]} \\ &\ll \sum_{j > d} \frac{1}{j^{2\rho}} h(N)^{2} + \sum_{1 \leq k < k' \leq N} c_{k}c_{k'} \sum_{j > d} a_{j} \sum_{j' > d} a_{j'} \mathbb{1}_{[jn_{k}-j'n_{k'}=0]}, \end{split}$$

where the last estimate employs $|a_j| \le j^{-\rho}$ for some $\rho > \frac{1}{2}$. Note that $jn_k - j'n_{k'} = 0$ and thus j' = 0 $\frac{jn_k}{n_{k'}} < jq^{k-k'}$. This means we obtain (with the formal definition of $c_k = 0$ for k > N) that

$$\begin{split} \sum_{1 \le k < k' \le N} c_k c_{k'} \sum_{j > d} a_j \sum_{j' > d} a_{j'} \mathbb{1}_{[jn_k - j'n_{k'} = 0]} \le \sum_{1 \le k < k' \le N} c_k c_{k'} \sum_{j > d} \frac{1}{j^{\rho}} \frac{1}{j^{\rho}} q^{k-k'} \\ &= \sum_{j > d} \frac{1}{j^{2\rho}} \sum_{1 \le k \le N} c_k \sum_{e \le N} c_{k+e} q^e \\ &= \sum_{j > d} \frac{1}{j^{2\rho}} \sum_{e \le N} q^e \sum_{1 \le k \le N} c_k c_{k+e} \\ &\le \sum_{j > d} \frac{1}{j^{2\rho}} \sum_{e \le N} q^e \sum_{1 \le k \le N} c_k^2 \\ &\ll_q h(N) \sum_{j > d} \frac{1}{j^{2\rho}}. \end{split}$$

For any given $\varepsilon > 0$, we can choose $d \in \mathbb{N}$ sufficiently large such that $\sum_{j>d} \frac{1}{j^{2\rho}} \le \varepsilon$ showing that $\|S_N^r\|_2^2 \le \varepsilon h(N).$

We now define

$$\sigma_N^f := \sqrt{h(N)} ||f||_2.$$

Then by construction, we have

$$(1-\varepsilon)\,\sigma_N^p \le \sigma_N^f \le (1+\varepsilon)\,\sigma_N^p$$

for any given $\varepsilon > 0$, provided the degree $d \in \mathbb{N}$ of p is sufficiently large. We now get (using the elementary estimate $\mathbb{P}[A \cap B] \ge \mathbb{P}[A] - \mathbb{P}[B^c]$)

$$\mathbb{P}\left[\frac{S_N}{\sigma_N^f} \ge t\right] \ge \mathbb{P}\left[\frac{S_N^p}{\sigma_N^f} \ge t - \sqrt[4]{\varepsilon}\right] - \mathbb{P}\left[\frac{|S_N^r|}{\sigma_N^f} \ge \sqrt[4]{\varepsilon}\right] \\ \ge \mathbb{P}\left[\frac{S_N^p}{\sigma_N^f} \ge t - \sqrt[4]{\varepsilon}\right] - \frac{\mathbb{V}\left[S_N^r\right]}{\sigma_N^f\sqrt{\varepsilon}} \\ \ge \mathbb{P}\left[\frac{S_N^p}{(1+\varepsilon)\sigma_N^p} \ge t - \sqrt[4]{\varepsilon}\right] - \frac{\sqrt{\varepsilon}}{\delta},$$

where, in the last step, we used that $\mathbb{V}[S_N^r] \leq \varepsilon \sigma_N^r$ and $\sigma_N^f = ||f||_2 \sqrt{h(N)}$. If we take the limit as $N \longrightarrow \infty$ on both sides, by Lemma 2.10 we thus obtain

$$\liminf_{N \to \infty} \mathbb{P}\left[\frac{S_N}{\sigma_N^f} \ge t\right] \ge 1 - \Phi\left(\frac{t - \sqrt[4]{\varepsilon}}{1 + \varepsilon}\right) - \sqrt{\varepsilon},$$

where Φ denotes the standard normal distribution function. Now we can take the limit as $\varepsilon \longrightarrow 0$ to get

$$\liminf_{N \to \infty} \mathbb{P}\left[\frac{S_N}{\sigma_N^f} \ge t\right] \ge 1 - \Phi(t).$$

The converse, i.e.,

$$\limsup_{N \to \infty} \mathbb{P}\left[\frac{S_N}{\sigma_N^f} \ge t\right] \le 1 - \Phi(t)$$

works analogously. In total, we have shown that

$$\frac{S_N}{\sigma_N^f} \xrightarrow{w} \mathcal{N}(0,1), \quad \text{as} \quad N \to \infty.$$

Proof of Theorem 1. The proof follows analogously to the one of Theorem 2.

References

- [1] C. Aistleitner. The law of the iterated logarithm for $\Sigma c_k f(n_k x)$. Dependence in Probability, Analysis and Number Theory. A Volume in Memory of Walter Philipp, pages 21–34, 2010.
- [2] C. Aistleitner. On the law of the iterated logarithm for the discrepancy of lacunary sequences. <u>Trans. Amer. Math. Soc.</u>, 362(11):5967–5982, 2010.
- [3] C. Aistleitner. On the law of the iterated logarithm for the discrepancy of lacunary sequences II. <u>Trans. Amer. Math.</u> Soc., 365(7):3713–3728, 2013.
- [4] C. Aistleitner and I. Berkes. On the central limit theorem for $f(n_k x)$. Probab. Theory Related Fields, 146(1-2):267–289, 2010.
- [5] C. Aistleitner, I. Berkes, and R. Tichy. Lacunary sequences in analysis, probability and number theory. <u>Societe</u> Mathematique de France, Panoramas et Synthèses, 62, 2024.
- [6] C. Aistleitner, L. Frühwirth, and J. Prochno. Diophantine conditions in the law of the iterated logarithm for lacunary systems. <u>Prob. Theory Related Fields</u>, pages 1–30, 2024.
- [7] C. Aistleitner, N. Gantert, Z. Kabluchko, J. Prochno, and K. Ramanan. Large deviation principles for lacunary sums. <u>Trans. Amer. Math. Soc.</u>, 376(01):507–553, 2023.
- [8] I. Berkes and W. Philipp. An as invariance principle for lacunary series f (nkx). <u>Acta Mathematica Academiae</u> <u>Scientiarum Hungarica</u>, 34:141–155, 1979.
- [9] J.-P. Conze and A. Raugi. Limit theorems for sequential expanding dynamical systems. In Ergodic Theory and Related Fields: 2004-2006 Chapel Hill Workshops on Probability and Ergodic Theory, University of North Carolina Chapel Hill, North Carolina, volume 430, page 89. American Mathematical Soc., 2007.
- [10] M. Émile Borel. Les probabilités dénombrables et leurs applications arithmétiques. <u>Rendiconti del Circolo</u> <u>Matematico di Palermo (1884-1940)</u>, 27(1):247–271, 1909.
- [11] L. Frühwirth, M. Juhos, and J. Prochno. The large deviation behavior of lacunary sums. Monatshefte für Mathematik, 199(1):113–133, 2022.
- [12] V. F. Gaposhkin. Lacunary series and independent functions. <u>Russian Mathematical Surveys</u>, 21(6):1, 1966.

- [13] V. F. Gaposhkin. The central limit theorem for some weakly dependent sequences. <u>Theory Probab. Appl.</u>, 15:649–666, 1971.
- [14] C. Heyde and B. Brown. On the departure from normality of a certain class of martingales. In <u>Selected Works of CC</u> <u>Heyde</u>, pages 129–133. Springer, 2010.
- [15] M. Kac. On the distribution of values of sums of the type $\sum f(2^k t)$. Ann. Math., 47(1):33–49, 1946.
- [16] M. Kac. Probability methods in some problems of analysis and number theory. <u>Bull. Amer. Math. Soc.</u>, 55:641–665, 1949.
- [17] A. Kolmogorov. Une contribution à l'étude de la convergence des séries de fourier. <u>Fundamenta Mathematicae</u>, 1(5):96–97, 1924.
- [18] G. W. Morgenthaler. On Walsh-Fourier series. <u>Trans. Amer. Math. Soc.</u>, 84(2):472–507, 1957.
- [19] W. Philipp. Limit theorems for lacunary series and uniform distribution mod 1. <u>Acta Arithmetica</u>, 26:241–251, 1975.
- [20] J. Prochno and M. Strzelecka. Moderate deviation principles and mod-gaussian convergence for lacunary trigonometric sums. <u>arXiv preprint arXiv:2312.09137</u>, 2023.
- [21] R. Salem and A. Zygmund. On lacunary trigonometric series. Proc. Natl. Acad. Sci., 33(11):333–338, 1947.
- [22] M. Weiss. The law of the iterated logarithm for lacunary trigonometric series. <u>Trans. Amer. Math. Soc.</u>, 91(3):444–469, 1959.
- [23] H. Weyl. Über die Gleichverteilung von Zahlen mod. Eins. <u>Math. Ann.</u>, 77(3):313–352, 1916.

Lorenz Frühwirth:

Faculty of Computer Science and Mathematics, University of Passau, Dr.-Hans-Kapfinger-Strasse 30, Passau, Germany

E-mail: lorenz.fruehwirth@uni-passau.de

Manuel Hauke:

Department of Mathematical Sciences, NTNU Trondheim, Sentralbygg 2 Gløshaugen,

Trondheim, Norway

E-mail: manuel.hauke@ntnu.no