V-REPRESENTABILITY AND HOHENBERG-KOHN THEOREM FOR NON-INTERACTING SCHRÖDINGER OPERATORS WITH DISTRIBUTIONAL POTENTIALS IN THE ONE-DIMENSIONAL TORUS

THIAGO CARVALHO CORSO

ABSTRACT. In this paper, we show that the ground-state density of any non-interacting Schrödinger operator on the onedimensional torus with potentials in a certain class of distributions is strictly positive. This result together with recent results from [SPR+24] provides a complete characterization of the set of non-interacting v-representable densities on the torus. Moreover, we prove that, for said class of non-interacting Schrödinger operators with distributional potentials, the Hohenberg-Kohn theorem holds, i.e., the external potential is uniquely determined by the ground-state density. In particular, the density-to-potential Kohn-Sham map is single-valued, and the non-interacting Lieb functional is differentiable at every point in this space of *v*-representable densities. These results contribute to establishing a solid mathematical foundation for the Kohn-Sham scheme in this simplified setting.

1. INTRODUCTION

Density functional theory (DFT) has become a cornerstone of quantum chemistry and materials science; by reformulating the complex many-electron problem in terms of the electron density rather than the wavefunction, DFT offers a practical method to study the electronic structure of many-body quantum systems. A central piece to the outstanding success of DFT is the celebrated Kohn-Sham [KS65] scheme, which seeks to reproduce the ground-state density of an interacting system of electrons via a fictitious system of non-interacting electrons.

However, the existence of such a fictitious non-interacting system is not well understood. This existence question is known as the v-representability problem and is a longstanding problem in the formulation of DFT. Despite its relevance (see, e.g. [WAR+23, THS+22]), in the arguably most relevant case of continuous systems in three-dimensional space, a solution to the *v*-representability problem remains elusive. Nevertheless, in simplified cases such as lattice systems [CCR85, PL21] and one-dimensional systems [AS88, CS91, CS93, SPR+24] significant progress has been made.

Of special interest to us here is the recent paper by Sutter et al [SPR+24], where the authors established sufficient conditions for a density to be ensemble *v*-representable on the one-dimensional torus $\mathbb{T} = \mathbb{R}/(2\pi\mathbb{Z})$. More precisely, they showed that, for a fixed interaction potential *w* satisfying suitable but rather general assumptions, any function $\rho : \mathbb{T} \to \mathbb{R}$ satisfying

$$\sqrt{\rho} \in \mathrm{H}^{1}(\mathbb{T}), \quad \int_{\mathbb{T}} \rho(x) \mathrm{d}x = N, \quad \text{and} \quad \rho(x) > 0 \quad \text{for all } x \in \mathbb{T},$$
 (1.1)

can be realized as the density of a (possibly mixed) ground-state of a Hamiltonian of the form

$$H_N(v,w) = -\Delta + \sum_{i\neq j}^N w(r_i - r_j) + \sum_{j=1}^N v(r_i) \quad \text{acting on} \quad \mathcal{H}_N = \bigwedge^N L^2(\mathbb{T}),$$

where v is a (distributional) potential in $H^{-1}(\mathbb{T})$. Here $H^{1}(\mathbb{T})$ stands for the Sobolev space of square integrable functions with square integrable weak derivatives of first order, and $H^{-1}(\mathbb{T})$ is the associated dual space (see Section 2.1 for precise definitions).

This result is rather remarkable as, to the best of the author's knowledge, it is the first rigorous sufficient criterion for both interacting and non-interacting *v*-representability in an infinite-dimensional and continuous system. However, as discussed in the conclusion of [SPR+24], many interesting questions regarding the *v*-representability problem on the one-dimensional torus remain unanswered; among them, the most important ones are perhaps the following:

- (1) Is the condition (2.1) also necessary, i.e., do any ground-state densities of $H_N(v, w)$ for arbitrary $v \in H^{-1}(\mathbb{T})$ satisfy (2.1)?
- (2) Is the class of potentials $v \in H^{-1}(\mathbb{T})$ too large and merely a mathematical artifact, i.e., is there a smaller class of "more reasonable" potentials that suffices to represent all densities satisfying (2.1)?

In this paper, we answer these two questions in the case of non-interacting systems. More precisely, the main contribution of this work can be summarized as follows.

Date: January 28, 2025.

²⁰²⁰ Mathematics Subject Classification. Primary: 81Q10 Secondary: 81V74, 34L40.

Key words and phrases. Density functional theory, ensemble v-representability, pure state v-representability, Hohenberg-Kohn theorem, Kohn-Sham map, exchange-correlation potential, Schrödinger equation, distributional potentials.

Funding information: DFG – Project-ID 442047500 – SFB 1481.

^{© 2024} by the authors. Faithful reproduction of this article, in its entirety, by any means is permitted for noncommercial purposes.

- (i) We show that, for non-interacting systems (i.e., w = 0), the conditions in (1.1) are not only sufficient but also necessary for *v*-representability. In particular, this gives a complete characterization of the set of non-interacting *v*-representable densities on \mathbb{T} .
- (ii) We prove the Hohenberg-Kohn theorem for distributional potentials in the case of non-interacting systems, i.e., if $H_N(v, 0)$ and $H_N(v', 0)$, with $v, v' \in H^{-1}(\mathbb{T})$, generate the same ground-state density, then v = v' up to an additive constant. In particular, this implies that the Kohn-Sham density-to-potential map is a well-defined single-valued map from the space of densities satisfying (1.1) to the space of potentials in $H^{-1}(\mathbb{T})$.
- (iii) We show that the ground-state of any single-particle Schrödinger operator with potential in $H^{-1}(\mathbb{T})$ is nondegenerate. In particular, for single-particle systems (N = 1), the conditions in (1.1) are in fact necessary and sufficient for *pure-state v-representability* and the Kohn-Sham density-to-potential map is a well-defined *smooth* and bijective map.
- (iv) On the other hand, we show that the second Hohenberg-Kohn theorem does not hold for excited states, i.e., there exists infinitely many (distinct up to an additive constant) potentials in $\mathbb{H}^{-1}(\mathbb{T})$ whose Hamiltonian has the same excited state wave-function.

2. MAIN RESULTS

To state our main results precisely, let us first introduce some notation.

2.1. Notation. Throughout this paper, we let $\mathbb{T} = \mathbb{R}/(2\pi\mathbb{Z})$ be the one-dimensional torus, and denote by $L^2(\mathbb{T})$ the standard space of (equivalent classes) of measurable functions that are square integrable with respect to the Lebesgue measure, i.e.,

$$f \in L^2(\mathbb{T})$$
 if and only if $||f||_2^2 := \langle f, f \rangle = \int_{\mathbb{T}} |f(x)|^2 dx < \infty$.

Moreover, we denote by \mathcal{H}_N the space of fermionic (or electronic) wave-functions on \mathbb{T}^N , i.e., the anti-symmetric tensor product space

$$\mathcal{H}_N \coloneqq \bigwedge^N \mathrm{L}^2(\mathbb{T}).$$

For a given wave-function $\Psi \in \mathcal{H}_N$, its (single-particle) density is defined as

$$\rho_{\Psi}(x) \coloneqq N \int_{\mathbb{T}^{N-1}} |\Psi(x, x_2, ..., x_N)|^2 \mathrm{d}x_2 ... \mathrm{d}x_N.$$

We let $H^1(\mathbb{T})$ denote the classical Sobolev space of functions $f \in L^2(\mathbb{T})$ with weak derivative $\nabla f \in L^2(\mathbb{T})$ endowed with the standard Hilbert norm, and we denote by $H^{-1}(\mathbb{T})$ the associated dual space of $H^1(\mathbb{T})$, i.e., the space of continuous linear functionals $v : H^1(\mathbb{T}) \to \mathbb{C}$ endowed with the operator norm.

Let us also introduce the following space of densities and potentials. For any $N \in \mathbb{N}$, we define \mathcal{D}_N as the space

$$\mathcal{D}_N \coloneqq \left\{ \rho : \mathbb{T} \to \mathbb{R} \text{ such that } \sqrt{\rho} \in \mathrm{H}^1(\mathbb{T}), \quad \int_{\mathbb{T}} \rho(x) \mathrm{d}x = N, \quad \text{and} \quad \rho > 0 \right\}$$
(2.1)

and \mathcal{V} as the space

 $\mathcal{V} \coloneqq \{ v \in \mathrm{H}^{-1}(\mathbb{T}) : v(f) \in \mathbb{R} \text{ for any real-valued function } f \in \mathrm{H}^{1}(\mathbb{T}) \}.$ (2.2)

Moreover, for any $v \in \mathcal{V}$, we shall denote by h(v) the self-adjoint realization of the operator

$$h(v) = -\Delta + v$$

given as a form-perturbation of the Laplacian on the torus (periodic Laplacian). More precisely, h(v) is the unique semi-bounded self-adjoint operator associated to the sesquilinear form

$$q_{h(v)}: \mathrm{H}^{1}(\mathbb{T}) \times \mathrm{H}^{1}(\mathbb{T}) \to \mathbb{C}, \quad q_{h(v)}(\varphi, \psi) = \int_{\mathbb{T}} \overline{\nabla \varphi(x)} \cdot \nabla \psi(x) \mathrm{d}x + v(\overline{\varphi}\psi).$$
(2.3)

For more details of this construction, we refer to Section 3. For $N \in \mathbb{N}$, we define the associated non-interacting *N*-particles Hamiltonian $H_N(v)$ as

$$H_N(v) = \sum_{j=1}^N 1 \otimes \dots \otimes \overbrace{h(v)}^{j^{th} position} \otimes \dots \otimes 1 \quad \text{acting on} \quad \mathcal{H}_N.$$
(2.4)

2.2. Main results. The first theorem we present here concerns the non-degeneracy of the ground-state for single-particle Hamiltonians with potentials in \mathcal{V} . This result plays a key role in the proof of the subsequent results and can be stated as follows.

Theorem 2.1 (Non-degenerate single-particle ground-state). Let $v \in V$ and $h(v) = -\Delta + v$ be the single-particle operator defined in (2.3), then the ground-state of h(v) is non-degenerate, and the unique (up to a global phase) normalized ground-state wave-function $\varphi_v \in H^1(\mathbb{T})$ is strictly positive everywhere, i.e., there exists a constant c = c(v) > 0 such that

$$\varphi_v(x) > c$$
 for any $x \in \mathbb{T}$.

Remark 2.2 (Existence of ground-state). Since the quadratic form domain of h(v) is $H^1(\mathbb{T})$, which is compactly embedded in $L^2(\mathbb{T})$, the operator h(v) has compact resolvent and therefore discrete spectrum. In particular, ground-state wave-functions of h(v) and its *N*-particles version $H_N(v)$ are guaranteed to exist.

As a consequence of Theorem 2.1, we obtain the following necessary conditions for non-interacting \mathcal{V} -representability¹.

Corollary 2.3 (Necessary conditions for non-interacting \mathcal{V} -representability). Let $N \in \mathbb{N}$ and $H_N(v)$ denote the N-particles non-interacting Hamiltonian defined in (2.4). Then the density of any mixed ground-state Γ ,

$$\rho_{\Gamma}(x) = N \int_{\mathbb{T}^{N-1}} \Gamma(x, x_2, ..., x_N, x, x_2, ..., x_N) dx_2 ... dx_N$$

satisfies

$$\sqrt{\rho_{\Gamma}} \in \mathrm{H}^{1}(\mathbb{T}), \quad \int_{\mathbb{T}} \rho_{\Gamma}(x) \mathrm{d}x = N \quad and \quad \rho_{\Gamma}(x) > 0 \quad for \ any \ x \in \mathbb{T}.$$

Combining the above result with [SPR+24, Theorem 1], we obtain a complete characterization of the set of non-interacting \mathcal{V} -representable densities on \mathbb{T} .

Theorem 2.4 (Characterization of \mathcal{V} -representable densities). A function $\rho : \mathbb{T} \to \mathbb{R}$ is the ground-state density of a *N*-particles non-interacting Hamiltonian $H_N(v)$ for some potential $v \in \mathcal{V}$ if and only if $\rho \in \mathcal{D}_N$ with \mathcal{D}_N defined according to (2.1).

The second natural question that was left open in [SPR+24] is whether one can recover the potential v from the ground-state density ρ . This question is not only natural but also specially relevant for (Kohn-Sham) DFT because it implies that the Kohn-Sham density-to-potential map $\rho \mapsto v^{\text{KS}}(\rho)$ is single-valued. Moreover, as highlighted in [SPR+24, Corollary 19], the existence of a unique potential v is also directly connected to the differentiability of the convex Lieb functional, which plays an important role in approximate schemes. Our next result provides an affirmative answer to this inverse problem for the case of non-interacting systems.

Theorem 2.5 (Hohenberg-Kohn theorem). Let $N \in \mathbb{N}$ and suppose that $\rho \in \mathcal{D}_N$ is a ground-state density of $H_N(v)$ and $H_N(v')$ for $v, v' \in \mathcal{V}$. Then v and v' are equal up to a constant, i.e., v = v' + c for some $c \in \mathbb{R}$.

Remark 2.6. Note that Theorem 2.5 is not a special case of previous results [Lie83, Zho12, Gar18, Lam18, Gar19, LBP20] because the class of potentials investigated here is rather large and include distributions such as the Dirac delta distribution. In fact, our results do not rely on the usual unique continuation for the *N*-particles wave-function [Geo79, SS80, Gar18, Kur97] as such results only guarantee that the wave-function does not vanish on a set of Lebesgue measure zero on \mathbb{T}^N , which may correspond to the support of delta-type distributions.

Theorem 2.5 guarantees that the *N*-particles Kohn-Sham (KS) density-to-potential map is a well-defined and singlevalued map from \mathcal{D}_N to the quotient space

$$\mathcal{V}/\{1\} \coloneqq \{[v]: v, v' \in [v], \text{ if } v - v' = \text{constant}\},\$$

i.e., if we identify potentials that differ only by an additive constant. Moreover, the existence of a ground-state (see Remark 2.2) implies that the KS map $v_N^{\text{KS}} : \mathcal{D}_N \to \mathcal{V}/\{1\}$ is also surjective. In particular, the fibers of v_N^{KS} , which corresponds to the set of ensemble ground-state densities of $H_N(v)$,

$$(v_N^{KS})^{-1}(\{[v]\}) = \mathcal{D}_N^{\text{ens}}(v) \coloneqq \left\{ \rho = \sum_{j=1}^m t_j \rho_{\Psi_j} : 0 \le t_j \le 1, \sum_{j=1}^m t_j = 1, \text{ and } \Psi_j \text{ ground-state of } H_N(v) \right\},$$

are all non-empty and disjoint, hence form a partition of the set \mathcal{D}_N . From this observation, we conclude that the space of potentials \mathcal{V} is not only sufficient but also necessary to represent all the densities in \mathcal{D}_N .

However, we note that, due to possible degeneracies of the ground-state, not every density in \mathcal{D}_N might be pure-state \mathcal{V} -representable. For instance, if we can find a potential $v \in \mathcal{V}$ such that the set of pure ground-state densities of $H_N(v)$,

$$\mathcal{D}_N^{\text{pure}}(v) \coloneqq \{ \rho_{\Psi} : \Psi \text{ ground-state of } H_N(v) \}$$

is not convex, then

$$\operatorname{conv}(\mathcal{D}_{N}^{\operatorname{pure}}(v)) \setminus \mathcal{D}_{N}^{\operatorname{pure}}(v) = \mathcal{D}_{N}^{\operatorname{ens}}(v) \setminus \mathcal{D}_{N}^{\operatorname{pure}}(v) \neq \emptyset$$

¹Here we use the term \mathcal{V} -representability instead of *v*-representability to emphasize the class of potentials under consideration.

and any density in this set is not pure-state \mathcal{V} -representable. In fact, as the subsets $\mathcal{D}_N^{\text{ens}}(v)$ form a partition of \mathcal{D}_N , the lack of convexity of $\mathcal{D}_N^{\text{pure}}(v)$ for some $v \in \mathcal{V}$ is the only obstruction to *pure-state* (non-interacting) \mathcal{V} -representability. In particular, as the set $\mathcal{D}_1^{\text{pure}}(v)$ for any $v \in \mathcal{V}$ consists of a single density by Theorem 2.1, we obtain a complete

In particular, as the set $\mathcal{D}_1^{rm}(v)$ for any $v \in \mathcal{V}$ consists of a single density by Theorem 2.1, we obtain a complete solution to the *pure-state* \mathcal{V} -representability problem in the case of a single-particle (N = 1). Moreover, the proof of this result does not rely on the results from [SPR+24].

Theorem 2.7 (Pure-state \mathcal{V} -representability for N = 1). Every density in \mathcal{D}_1 is pure-state \mathcal{V} -representable by a unique (up to an additive constant) potential $v \in \mathcal{V}$. In particular, the unique Kohn-Sham density-to-potential map is given by

$$v_1^{\text{KS}}: \mathcal{D}_1 \to \mathcal{V}/\{1\}, \qquad \rho \mapsto v_1^{\text{KS}}(\rho) = \frac{\Delta \sqrt{\rho}}{\sqrt{\rho}}.$$
 (2.5)

Moreover, this map is smooth² and bijective.

As a last result, we show that, while the second part of the Hohenberg-Kohn theorem (cf. [PTC+23, HK2 Theorem]) still holds for ground-state wave-functions in the non-interacting case, i.e., two different potentials with the same ground-state wave-function can only differ by an additive constant, the same is not true for excited-states. This is a drawback of extending the class of admissible potentials to include distributions as the Dirac delta.

Theorem 2.8 (No Hohenberg-Kohn for excited states). For any $v \in V$ and any real-valued excited state φ_k of h(v) with $k \ge 2$, there exist (uncountable many) potentials $v' \in V$ such that v - v' is not constant and φ_k is also an excited state of h(v').

2.3. **Outline of the paper**. Let us now outline the key steps in the proof of our results, and how they are distributed in the next sections.

The proof of Theorem 2.1 consists of three steps and is carried out in Section 4. In the first step, we use classical results from the book by Reed and Simon [RS78] (cf. Theorems XVIII.43 and 45) to show that the ground-state of h(v) is non-degenerate and strictly positive almost everywhere. In the second step, we apply Courant's nodal domain theorem to conclude that the ground-state can not vanish in more than one point. If we were dealing with an interval *I* with Dirichlet boundary conditions, these two steps would be enough to prove Theorem 2.1 because $I \setminus \{x_0\}$ consists of two connected components. In the torus, however, this is no longer true and we need a third step. This last step uses a gluing argument to obtain a contradiction with Courant's nodal domain theorem. More precisely, we use a gluing argument to construct ground-state densities that vanish on finitely many points, provided that a ground-state density vanishing on a single-point exists.

The proof of the Hohenberg-Kohn Theorem 2.5 is presented in Section 5. This proof consists of two main steps. The first step is the standard Hohenberg-Kohn argument, which shows that, if $H_N(v)$ and $H_N(v')$ have the same ground-state density, they must have a mutual ground-state wave-function. After this step, the usual argument used in previous proofs of the Hohenberg-Kohn theorem [HK64, Lie83, PTC+23, Gar18] consists in dividing the Schrödinger equation

$$(H_N(v) - H_N(v')) \Psi = 0$$
(2.6)

by this mutual ground-state wave-function Ψ , which is possible in a (almost everywhere) pointwise sense by unique continuation results. In our setting, this is no longer possible³ because we are dealing with distributional potentials which are not pointwise defined and whose support may have Lebesgue measure zero. To overcome this difficulty, we appeal to the fact that h(v) has discrete spectrum, and combine a spectral representation of the dual space $H^{-1}(\mathbb{T})$ (Lemma 3.5) with Theorem 2.1. More precisely, we show that, if Ψ satisfies (2.6) in a weak sense, has finitely many natural orbitals, and one of these orbitals can be chosen strictly positive, then the difference of potentials (v - v') is much more regular than expected, namely, belongs to $H^1(\mathbb{T})$. Combining this extra regularity with some linear algebra arguments, we can then show that (v - v') must be constant, which completes the proof of Theorem 2.5.

The proofs of Theorems 2.7 and 2.8 are also presented in Section 5. The former follows from the existence, strict positivity, and non-degeneracy of the ground-state, plus some fairly standard arguments to show smoothness of a map. The proof of the latter relies on the simple observation that any (real-valued) excited state φ_k must vanish at a point, and therefore, adding a Dirac's delta at that point still preserves φ_k as an excited state.

For the sake of completeness, we present the necessary mathematical background for our proofs in some detail in Section 3. These comprises the precise definitions and a few well-known properties of the Sobolev spaces $H^1(\mathbb{T})$ and $H^{-1}(\mathbb{T})$, the quadratic form construction of h(v), and the definition and simple properties of the natural orbital decomposition of a wave-function. In Section 6 we conclude with a brief discussion on some possible extensions of our main results and some natural open questions.

3. MATHEMATICAL BACKGROUND

In this section we briefly review the mathematical background necessary for the proofs of our main results.

²Note that \mathcal{D}_1 is a smooth manifold, as it is an open set of the closed subspace $\{f \in H^1(\mathbb{T}; \mathbb{R}) : \int_{\mathbb{T}} f(x) dx = 1\}$ of the Banach space $H^1(\mathbb{T}; \mathbb{R})$. Hence, differentiability for maps in \mathcal{D}_1 has a well-defined meaning.

³In fact, in the single-particle case, the division by the ground-state argument is still possible (in an operator sense) thanks to the strict positivity of the ground-state (cf. Theorem 2.1) and the algebra property of $H^1(\mathbb{T})$ (Lemma 3.2), see, e.g., the proof of Theorem 2.7. For $N \ge 2$, however, this argument does not apply as we have no control over the zero set of the N-particles ground-state Ψ .

3.1. Sobolev and dual spaces on the torus. We begin by recalling the definitions of $H^1(\mathbb{T})$ and $H^{-1}(\mathbb{T})$.

Definition 3.1 (Sobolev spaces). We denote by $H^1(\mathbb{T})$ the closure of the space $C^{\infty}(\mathbb{T})^4$ with respect to the norm

$$\|\varphi\|_{\mathrm{H}^{1}}^{2} \coloneqq \|\varphi\|_{\mathrm{L}^{2}}^{2} + \|\nabla\varphi\|_{\mathrm{L}^{2}}^{2}.$$
(3.1)

The space $H^{-1}(\mathbb{T})$ is the set of continuous linear functionals on $H^{1}(\mathbb{T})$ endowed with the operator norm

$$\|v\|_{\mathrm{H}^{-1}} = \sup_{f \in \mathrm{H}^{1}(\mathbb{T}) \setminus \{0\}} \frac{|v(f)|}{\|f\|_{\mathrm{H}^{1}}}.$$
(3.2)

As usual, we identify measurable functions in $f: \mathbb{T} \to \mathbb{C}$ with linear functionals on $H^1(\mathbb{T})$ via the Riesz mapping

$$g \in \mathrm{H}^{1}(\mathbb{T}) \mapsto f(g) = \langle f, g \rangle = \int_{\mathbb{T}} \overline{f(x)} g(x) \mathrm{d}x.$$
 (3.3)

That such a functional is well-defined and continuous for any function $f \in L^1(\mathbb{T})$ is a consequence of Hölder's inequality and the Gagliardo-Nirenberg-Sobolev (GNS) inequality stated below (see (3.5)). In fact, the GNS inequality implies that $H^1(\mathbb{T})$ is an algebra of functions. More precisely, we have

Lemma 3.2 (Algebra property of $H^1(\mathbb{T})$). Let $\varphi, \psi \in H^1(\mathbb{T})$, then $\varphi \psi \in H^1(\mathbb{T})$ and the following estimate holds

$$\|\varphi\psi\|_{\mathrm{H}^{1}(\mathbb{T})} \lesssim \|\varphi\|_{\mathrm{L}^{2}}^{\frac{1}{2}} \|\varphi\|_{\mathrm{H}^{1}}^{\frac{1}{2}} \|\psi\|_{\mathrm{H}^{1}}^{\frac{1}{2}} + \|\psi\|_{\mathrm{L}^{2}}^{\frac{1}{2}} \|\psi\|_{\mathrm{H}^{1}}^{\frac{1}{2}} \|\varphi\|_{\mathrm{H}^{1}}.$$
(3.4)

In particular, the operator of multiplication by $\varphi, \psi \mapsto M_{\varphi}(\psi) = \varphi \psi$, is bounded in $H^1(\mathbb{T})$. Moreover, if $|\varphi(x)| > 0$ for every $x \in \mathbb{T}$, then this operator is invertible with inverse given by $M_{1/\varphi}$.

Proof. The proof of (3.4) is straightforward from the product rule, Hölder's inequality, and the well-known Gagliardo-Nirenberg-Sobolev inequality, which shows that any $\varphi \in H^1(\mathbb{T})$ is continuous and satisfies the bound

$$\|\varphi\|_{L^{\infty}(\mathbb{T})} \lesssim \|\nabla\varphi\|_{L^{2}}^{\frac{1}{2}} \|\varphi\|_{L^{2}}^{\frac{1}{2}} + \|\varphi\|_{L^{2}} \le 2\|\varphi\|_{L^{2}}^{\frac{1}{2}} \|\varphi\|_{H^{1}}^{\frac{1}{2}}.$$
(3.5)

The following result is a simple consequence of the above lemma and will be useful to show that the KS map for a single-particle in (2.5) is smooth.

Lemma 3.3 (Differentiability of push-forward). Let $g \in C^{\infty}(\mathbb{R}; \mathbb{R})$, then the (nonlinear) push-forward map $g^{\#} : H^{1}(\mathbb{T}; \mathbb{R}) \to H^{1}(\mathbb{T}; \mathbb{R})$ given by

$$(g^{\#}\psi)(x) = g(\psi(x)), \quad x \in \mathbb{T}.$$

is smooth in $H^1(\mathbb{T}; \mathbb{R})$.

Proof. Let $\mathcal{B}(H^1(\mathbb{T}))$ denote the Banach space of bounded linear operators from $H^1(\mathbb{T})$ to $H^1(\mathbb{T})$. Then by Lemma 3.2, the map $M : H^1(\mathbb{T}) \to \mathcal{B}(H^1(\mathbb{T}))$, given by

$$\psi \in \mathrm{H}^{1}(\mathbb{T}) \to M_{\psi} \in \mathcal{B}(\mathrm{H}^{1}(\mathbb{T})) \quad \text{where} \quad M_{\psi}(\varphi) = \psi\varphi,$$

$$(3.6)$$

is continuous. Since this map is also linear, it is smooth.

Next, we claim that for any $g \in C^3(\mathbb{R}; \mathbb{R})$, the map $g^{\#}$ belongs to $C^1(H^1(\mathbb{T}; \mathbb{R}), H^1(\mathbb{T}; \mathbb{R}))$. To see this, first note that, since g is locally Lipschitz, the chain rule yields

$$\nabla(g^{\#}\psi) = \dot{g}(\psi)\nabla\psi, \quad \text{and therefore,} \quad \|\nabla(g^{\#}\psi)\|_{2} \le \|\dot{g}\|_{L^{\infty}(-C,C)}\|\nabla\psi\|_{2},$$

where $C = \|\psi\|_{L^{\infty}} \leq \|\psi\|_{H^1}$ and \dot{g} denotes the derivative of g. Hence, $g^{\#}$ maps $H^1(\mathbb{T}; \mathbb{R})$ to $H^1(\mathbb{T}; \mathbb{R})$. Furthermore, from the mean value inequality we find that

$$\begin{split} \|\nabla \left(g^{\#}(\psi+\delta) - g^{\#}(\psi) - M_{\dot{g}^{\#}(\psi)}\delta\right)\|_{2}^{2} &= \int_{\mathbb{T}} |\dot{g}(\psi+\delta)\nabla(\psi+\delta) - \dot{g}(\psi)\nabla\psi - \ddot{g}(\psi)\delta\nabla\psi - \dot{g}(\psi)\nabla\delta|^{2} dx \\ &= \int_{\mathbb{T}} |(\dot{g}(\psi+\delta) - \dot{g}(\psi))\nabla\delta + (\dot{g}(\psi+\delta) - \dot{g}(\psi) - \ddot{g}(\psi)\delta)\nabla\psi|^{2} dx \\ &\leq \|\ddot{g}\|_{L^{\infty}(-C,C)}^{2} \|\delta\|_{H^{1}}^{4} + \|g^{(3)}\|_{L^{\infty}(-C,C)}^{2} \|\psi\|_{H^{1}}^{2} \|\delta\|_{H^{1}}^{4}, \end{split}$$

where $C = \sup_{t \in [0,1]} \|\psi + t\delta\|_{L^{\infty}} \leq \|\psi\|_{H^1} + \|\delta\|_{H^1}$, and $g^{(3)}$ denotes the third derivative of g. Thus, $g^{\#}$ is indeed C^1 and the derivative at $\psi \in H^1(\mathbb{T}; \mathbb{R})$ is

$$\mathrm{d}_{\psi}g^{\#}(\delta) = \dot{g}(\psi)\delta = M_{\dot{g}^{\#}\psi}\delta.$$

As this map is a composition of the *M* map in (3.6) and the map $\dot{g}^{\#}$, we conclude that $g^{\#} \in \mathbb{C}^2$, provided that $g \in C^4$. A bootstrap (induction) argument then shows that $g^{\#} \in \mathbb{C}^{\infty}$ for $g \in C^{\infty}$, which concludes the proof.

⁴Alternatively, one can define $H^1(\mathbb{T})$ as the closure of C^1 functions on the interval $(0, 2\pi)$ that are continuous up to the boundary, have L^2 integrable derivative, and satisfy $f(0) = f(2\pi)$.

T. CARVALHO CORSO

Remark 3.4 (Lie group structure of density spaces). The GNS inequality implies that the set $\mathcal{D} := \{f \in H^1(\mathbb{T}; \mathbb{R}) : f > 0\}$

$$D := \{ f \in H^1(\mathbb{T}; \mathbb{R}) : f > 0 \}$$
(3.7)

is an open subset of the Banach space $H^1(\mathbb{T}; \mathbb{R})$. This observation plays a key role in the proof of the results of [SPR+24]. Moreover, together with Lemma 3.2, this implies that \mathcal{D} is a Lie group⁵ with respect to pointwise multiplication. In particular, \mathcal{D}_N defined in (2.1) is a submanifold of codimension 1, and the tangent space at each point $f \in \mathcal{D}_N$ can be identified with the set

$$T_f \mathcal{D}_N = \left\{ \delta \in \mathrm{H}^1(\mathbb{T}; \mathbb{R}); \int_{\mathbb{T}} \delta(x) \mathrm{d}x = 0 \right\}$$

3.2. Quadratic form definition of Hamiltonian. As a consequence of estimate (3.4) and Young's inequality, the following bound holds:

$$v(|\varphi|^{2}) \leq \|v\|_{H^{-1}} \|\varphi\|_{H^{1}}^{\frac{3}{2}} \|\varphi\|^{\frac{1}{2}} \leq \|v\|_{H^{-1}} \left(\epsilon \|\varphi\|_{H^{1}}^{2} + \frac{C}{\epsilon} \|\varphi\|_{L^{2}}^{2}\right) \quad \text{for any } \epsilon > 0.$$
(3.8)

In particular, the quadratic form induced by any $v \in H^{-1}(\mathbb{T})$ is Δ -form bounded with relative bound 0. Therefore, by the KLMN theorem [RS75, Theorem X.17], the operator h(v) is a well-defined self-adjoint operator with quadratic form

$$q_{h(v)}(\varphi,\psi) = \int_{\mathbb{T}} \overline{\nabla \varphi(x)} \nabla \psi(x) dx + v(\overline{\varphi}\psi), \quad \text{for any } \psi, \varphi \in Q(h(v)) \coloneqq H^1(\mathbb{T}).$$

(For more details on this application of the KLMN theorem, we refer to [RS75, Her89, SPR+24].)

An useful consequence of this construction is that the quadratic form domain of h(v) is the Sobolev space $H^1(\mathbb{T})$. As this space is compactly embedded in $L^2(\mathbb{T})$ by the standard Sobolev embedding theorem, the resolvent operator of h(v) is a bounded operator form $L^2(\mathbb{T})$ to $H^1(\mathbb{T})$. Consequently, the resolvent is a compact operator and, by the spectral theorem, the spectrum of h(v) is purely discrete, i.e., there exists a non-decreasing sequence $\{\lambda_k\}_{k\in\mathbb{N}}$ and an L^2 -orthonormal basis $\{\varphi_k\}_{k\in\mathbb{N}}$ such that

$$h(v)\varphi_k = \lambda_k \varphi_k$$
 and $\lambda_k \uparrow \infty$.

The above spectral decomposition of h(v) will be useful to prove the Hohenberg-Kohn theorem 2.5 in Section 5. To be more precise, there we shall use the following representation of distributions in $H^{-1}(\mathbb{T})$.

Lemma 3.5 (Spectral representation of dual Sobolev space). Let $v \in H^{-1}(\mathbb{T})$ and $\{\varphi_k\}_{k \in \mathbb{N}}$ be an orthonormal basis of eigenfunctions of h(v) with corresponding eigenvalues $\{\lambda_k\}_{k \geq 1}$. Then a distribution $f \in \mathcal{D}'(\mathbb{T})$ belongs to $H^{-1}(\mathbb{T})$ if and only if there exists a sequence $\{c_j\}_{j \in \mathbb{N}}$ such that

$$\sum_{j\in\mathbb{N}} (1-\lambda_1+\lambda_j)^{-1} |c_j|^2 < \infty \quad and \quad f(\varphi) = \sum_{j=1}^{\infty} c_j \langle \varphi_j, \varphi \rangle, \quad for \ any \ \varphi \in \mathrm{H}^1(\mathbb{T}).$$
(3.9)

Moreover, in this case $c_i = f(\varphi_i)$ *.*

Proof. The proof is rather standard, but we sketch the arguments for convenience of the reader. First, note that any $\varphi \in L^2(\mathbb{T})$, can be written as $\varphi = \sum_{j\geq 1} a_j \varphi_j$ where $a_j := \langle \varphi_j, \varphi \rangle$. Since h(v) has quadratic form domain $H^1(\mathbb{T})$ (see also (3.8)), we find that

$$\|\varphi\|_{\mathrm{H}^1(\mathbb{T})}^2 \sim \langle \varphi, (h(v) - \lambda_1 + 1) \varphi \rangle = \sum_{j=1}^{\infty} |a_j|^2 (\lambda_j - \lambda_1 + 1),$$

where ~ denotes equivalence of norms. Hence, $\varphi \in H^1(\mathbb{T})$ if and only if $\sum |a_j|^2 (1 - \lambda_1 + \lambda_j) < \infty$. Therefore, any f of the form (3.9) belongs to $H^{-1}(\mathbb{T})$ by Cauchy-Schwarz, i.e.,

$$f(\varphi) = \sum_{j=1}^{\infty} c_j a_j \le \left(\sum_{j=1}^{\infty} \frac{|c_j|^2}{(1+\lambda_j - \lambda_1)} \right)^{\frac{1}{2}} \left(\sum_{j\ge 1} |a_j|^2 (1+\lambda_j - \lambda_1) \right)^{\frac{1}{2}} \le \|\varphi\|_{\mathrm{H}^1(\mathbb{T})}$$

Conversely, if $f \in H^{-1}(\mathbb{T})$, then we can define $c_j \coloneqq f(\varphi_j)$, and conclude by using the simple fact that

$$\left|\sum_{j=1}^{\infty} c_j a_j\right| \lesssim \left(\sum_{j=1}^{\infty} |a_j|^2 (1+\lambda_j-\lambda_1)\right)^{\frac{1}{2}}, \quad \text{for any sequence } \{a_j\}_{j\in\mathbb{N}},$$

if and only if $\{c_j\}_{j \in \mathbb{N}}$ satisfies (3.9).

⁵For precise definitions of (infinite dimensional) manifolds and Lie groups, we refer, e.g., to [KM97].

3.3. Single-particle density matrix and natural orbitals. Let us now recall the definition of the single-particle density matrix and its natural orbital decomposition.

Definition 3.6 (Single-particle density matrix). Let $N \in \mathbb{N}$, then for any $\Psi \in \mathcal{H}_N$ we define its single-particle density matrix as the function

$$\gamma_{\Psi}(x,y) \coloneqq N \int_{\mathbb{T}^{N-1}} \overline{\Psi(x,x_2,...,x_N)} \Psi(y,x_2,...,x_N) \mathrm{d}x_2...\mathrm{d}x_N.$$

Alternatively, one can see the single-particle density matrix as the integral kernel of the operator

$$\langle \varphi, \gamma_{\Psi} \psi \rangle = \langle a(\varphi) \Psi, a(\psi) \Psi \rangle_{\mathcal{H}_{N-1}}, \text{ for } \varphi, \psi \in \mathcal{H}_1,$$

where $a(\varphi)$ denotes the usual fermionic annihilation operator,

$$a(\varphi): \mathcal{H}_N \to \mathcal{H}_{N-1}, \quad (a(\varphi)\Psi)(x_1, ..., x_{N-1}) = \sqrt{N} \int_{\mathbb{T}} \overline{\varphi(x)} \Psi(x, x_1, ..., x_{N-1}) \mathrm{d}x.$$
(3.10)

It is well-known that the single-particle density matrix of any *N*-particles wave-function is a bounded, positive and trace-class operator satisfying

$$\|\gamma_{\Psi}\|_{\mathcal{H}_1\to\mathcal{H}_1} = \|\Psi\|_{\mathcal{H}_N}^2 \quad \text{and} \quad \mathrm{tr}\gamma_{\Psi} = N\|\Psi\|_{\mathcal{H}_N}^2.$$

Hence, γ_{Ψ} is a compact self-adjoint operator on \mathcal{H}_1 and, by the spectral theorem, there exists an orthonormal family of eigenfunctions $\{\varphi_i\}_{i \leq M}$ and eigenvalues $\{n_k\}_{k \leq M}$ such that

$$0 < n_k \le \|\Psi\|_{\mathcal{H}_N}^2, \quad \sum_{k=1}^M n_k = N \|\Psi\|_{\mathcal{H}_N}^2, \quad \text{and} \quad \gamma_{\Psi}(x, y) = \sum_{k=1}^M n_k \overline{\varphi_k(x)} \varphi_k(y).$$
(3.11)

Here $M \in \mathbb{N} \cup \{+\infty\}$, i.e., M does not need to be finite. The eigenfunctions φ_k are called the natural orbitals of Ψ , and n_k are called the associated (natural) occupation numbers.

Remark 3.7 (Non-uniqueness of natural orbitals). Note that the natural orbitals are not uniquely defined, as any linear combination of the orbitals with the same occupation number is again a natural orbital.

For our proofs, we shall need two properties of the natural orbitals. The first one is the general fact that the natural orbitals of eigenfunctions of non-interacting operators can be chosen as eigenfunctions of the single-particle operator.

Lemma 3.8 (Natural orbital decomposition of non-interacting ground-state). Let h be a semibounded self-adjoint operator on $L^2(\mathbb{T})$ with purely discrete spectrum. Let $N \in \mathbb{N}$, then for any ground-state $\Psi \in \mathcal{H}_N$ of the non-interacting N-particles Hamiltonian

$$H_N := \sum_{j=1}^N 1 \otimes 1 \dots \otimes \stackrel{j^{tn} position}{h} \otimes 1 \dots \otimes 1, \quad acting \ on \quad \mathcal{H}_N$$

there exists an orthonormal basis $\{\varphi_j\}_{j \in \mathbb{N}}$ of eigenfunctions of h ordered in non-decreasing order, i.e., $\langle \varphi_k, h, \varphi_k \rangle \leq \langle \varphi_{k+1}h, \varphi_{k+1} \rangle$, and a finite number $M \in \mathbb{N}$ such that $\{\varphi_j\}_{j \leq M}$ are the natural orbitals of Ψ .

Proof. The proof follows from the fact that, since the spectrum of h(v) is discrete (see previous subsection), there exists an orthonormal basis of eigenfunctions $\{\varphi_j\}_{j \in \mathbb{N}}$, and therefore, the Slater determinants

$$\Phi_{j_1,...,j_N} := a(\varphi_{j_1})^* a(\varphi_{j_2})^* \dots a(\varphi_{j_N})^* 1, \quad 1 \le j_1 < j_2 < \dots < j_N \in \mathbb{N},$$

where $1 \in \mathbb{C} = \mathcal{H}_0$ is the vacuum state and $a(\varphi)^* : \mathcal{H}_{N-1} \to \mathcal{H}_N$ is the creation operator

$$(a(\varphi)^*\Psi)(x_1,...,x_N) = \frac{1}{\sqrt{N}} \sum_{i=1}^N (-1)^{1+i} \varphi(x_i) \Psi(x_1,...,x_{i-1},x_{i+1},...,x_N),$$
(3.12)

form an eigenbasis of $H_N(v)$ for the *N*-particles space \mathcal{H}_N .

The second result we shall need is the following simple formula for the overlapping density of a wave-function with its single-excitations.

Lemma 3.9 (Overlapping density with single-excitations). Let $\Psi \in \mathcal{H}_N$ and $\{\varphi_j\}_{j=1}^M$ be its natural orbitals. Then for $\Phi = a(\psi)^* a(\varphi_k) \Psi$, where $\psi \perp \{\varphi_j\}_{j \leq M}$ and $a(\psi)^*$ and $a(\varphi_k)$ are the creation and annihilation operators defined in (3.12) and (3.10), we have

$$\rho_{\Psi,\Phi}(x) = \int_{\mathbb{T}^{N-1}} \overline{\Psi(x, x_2, ..., x_N)} \Phi(x, x_2, ..., x_N) dx_2 ... dx_N = n_k \overline{\varphi_k(x)} \psi(x), \tag{3.13}$$

where $n_k > 0$ is the occupation number of φ_k .

Proof. First note that since $\psi \perp \{\varphi_j : j \leq M\}$ can be assumed to be normalized, we can extend $\{\varphi_j\}_{j \leq M}$ to an orthornomal basis $\{\varphi_j\}_{j \in \mathbb{N}}$ such that $\varphi_{M+1} = \psi$. Now consider $v \in L^{\infty}(\mathbb{T})$ and denote by V_N the *N*-particle operator of multiplication by

$$V_N(x_1, ..., x_N) = \sum_{j=1}^N v(x_j).$$

Then from the second quantization representation

$$V_N = \sum_{i,j\geq 1} v_{ij} a_i^* a_j, \text{ where } a_i = a(\varphi_i), a_j^* = a(\varphi_j)^*, \text{ and } v_{ij} = \langle \varphi_i, v\varphi_j \rangle,$$

and the canonical anti-commutation relations (CAR)

$$a_j a_k = -a_k a_j$$
, and $a_k a_j^* = \delta_{kj} - a_j^* a_k$, where δ_{ij} is the Kronecker delta,

we find that

$$\langle \Psi, V_N a(\psi)^* a(\varphi_k) \Psi \rangle = \sum_{i,j} v_{ij} \langle \Psi, a_i^* a_j a_{M+1}^* a_k \Psi \rangle = \sum_i v_{iM+1} \langle a_i \Psi, a_k \Psi \rangle + \sum_{i,j} v_{ij} \langle a_i a_{M+1} \Psi, a_j a_k \Psi \rangle$$

We can now use that $a_k \Psi = 0$ for any k > M (since $\varphi_k \perp \{\varphi_j\}_{j \le M}$ which are the natural orbitals of Ψ) to obtain

$$\langle \Psi, V_N \Phi \rangle = \sum_{i=1}^{M} v_{iM+1} \langle \varphi_i, \gamma_{\Psi} \varphi_k \rangle = n_k v_{kM+1} = \int_{\mathbb{T}} v(x) n_k \overline{\varphi}_k(x) \psi(x) dx.$$
(3.14)

On the other hand, we have

$$\langle \Psi, V_N \Phi \rangle = \int_{\mathbb{T}} v(x) \rho_{\Psi, \Phi}(x) \mathrm{d}x.$$
 (3.15)

As (3.14) and (3.15) holds for any $v \in L^{\infty}(\mathbb{T})$, we obtain (3.13).

4. Non-degeneracy of the ground-state

In this section, our goal is to prove Theorem 2.1 and Corollary 2.3. To this end, we shall need two additional lemmas. The first lemma shows that the ground-state of h(v) is non-degenerate and almost everywhere strictly positive. This kind of result is sometimes called Perron–Frobenius theorem; it can be understood as a unique continuation property but applies only to the ground-state of the single-particle system.

Lemma 4.1 (Non-degenerate ground-state). Let $v \in \mathbb{H}^{-1}(\mathbb{T})$, then the ground-state of h(v) is non-degenerate and (up to a global phase) satisfies

$$\varphi_v(x) > 0$$
 for almost every $x \in \mathbb{T}$.

Proof. For the proof, we shall use two results from the book by Reed and Simon [RS78]. To simplify the presentation, we combine the statement of these two results in a single lemma.

Lemma 4.2 (Theorem XIII.43 and XIII.45). Let h_0 and h be two semibounded self-adjoint operators in $L^2(\Omega, \mu)$, where (Ω, μ) is a sigma-finite measure space. Suppose that

- (i) h_0 has a simple and almost everywhere strictly positive ground-state.
- (ii) e^{-th_0} is positivity preserving, i.e., it maps non-negative functions to non-negative functions.

Then, if there exists a sequence of bounded functions $\{v_n\}_{n\in\mathbb{N}} \subset L^{\infty}(\Omega,\mu)$ such that $h_0 + v_n$ converges in the strong resolvent sense to h, then the ground-state of h (if existing) is also simple and strictly positive almost everywhere.

Thus, in order to prove Lemma 4.1, it suffices to find an operator h_0 satisfying (i) and (ii) and a sequence of bounded functions v_n such that $h_0 + v_n$ converges in the strong resolvent sense to h(v). As usual in the applications of such results, we choose h_0 as the free (periodic) Laplacian.

So first, note that the ground-state of $h_0 = -\Delta$ is the constant function, and therefore strictly positive; thus assumption (i) holds for h_0 . Next, recall that the heat propagator of the periodic Laplacian is given by

$$(e^{-th_0}f)(x) = (p_t * f)(x) = \int_0^{2\pi} p_t(x-y)f(y)dy,$$

where p_t is the periodization of the heat kernel of the Laplacian in \mathbb{R} . As the latter is nothing but the standard Gaussian, we find that

$$p_t(x) = \sum_{k \in \mathbb{Z}} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x+2\pi k)^2}{4t}} \ge 0 \quad \text{for any } x \in [0, 2\pi].$$

As convolution with a non-negative function is a positivity preserving operator, assumption (ii) also holds.

To conclude, note that, since $L^2(\mathbb{T})$ is dense⁶ in $H^{-1}(\mathbb{T})$ (by Lemma 3.5) and $L^{\infty}(\mathbb{T})$ is dense in $L^2(\mathbb{T})$, there exists a sequence of bounded functions $v_n \in L^{\infty}(\mathbb{T})$ such that

$$\lim_{n \to \infty} \|v - v_n\|_{\mathbf{H}^{-1}(\mathbb{T})} = 0.$$

Consequently, by estimate (3.4) in Lemma 3.2,

$$|\langle \varphi, h(v)\psi\rangle - \langle \varphi, h_0 + v_n, \psi\rangle| = |\langle \varphi, (v - v_n)\psi\rangle| = |(v - v_n)(\overline{\varphi}\psi)| \lesssim \|v - v_n\|_{\mathrm{H}^{-1}} \|\psi\|_{\mathrm{H}^{1}} \|\varphi\|_{\mathrm{H}^{1}}$$

Therefore, $h_0 + v_n$ converges to h(v) in the norm topology of operators in $\mathcal{B}(H^1(\mathbb{T}), H^{-1}(\mathbb{T}))$. Using the first resolvent formula, we then conclude that $h_0 + v_n$ converges to h(v) in the strong resolvent sense (see, e.g., [RS80, Theorem VIII.25.(c)] for a similar argument). Therefore Lemma 4.2 applies and the proof is complete.

The previous lemma only guarantees that the ground-state φ_v does not vanish in a set of measure zero. To obtain the stronger strict positivity everywhere, we shall use the following version of the classical Courant nodal domain theorem [CH89]. The proof of this result is exactly the same as in the case of Schrödinger operators with standard multiplication potentials. Yet, for the sake of completeness, we briefly sketch the main steps below.

Lemma 4.3 (Courant nodal domain theorem). Let $\varphi \in H^1(\mathbb{T})$ be an eigenfunction of h(v) with eigenvalue λ , then the number of nodal domains of φ , i.e., the number of connected components of the (open) set $\{x \in \mathbb{T} : \varphi(x) \neq 0\}$ is less or equal than $n(\lambda) := \sum_{\mu \leq \lambda} \dim \ker (h(v) - \mu)$.

Proof. First, let $\varphi \in H^1(\mathbb{T})$ be an eigenstate of h(v) with eigenvalue λ . Let $\{O_j\}_{j \le M}$ denote the connected components of $\{\varphi \neq 0\}$, and define $\varphi_j := \varphi|_{O_j}$. Since φ is continuous by the GNS inequality (3.5), each of the sets O_j is an open interval. Moreover, as $\varphi = 0$ at the ends of O_j , the extension of φ_j by zero to the whole torus \mathbb{T} belongs to $H^1(\mathbb{T})$. Since $\overline{\varphi_j}\varphi_k = 0$ everywhere for $j \neq k$, we have

$$q_{h(v)}(\varphi_j,\varphi_k) = \int_{\mathbb{T}} \overline{\nabla \varphi_j(x)} \nabla \varphi_k(x) dx + v(\overline{\varphi_j}\varphi_k) = \int_{\mathbb{T}} \overline{\nabla \varphi_j(x)} \nabla \varphi_k(x) = 0 \quad \text{if } j \neq k.$$

In particular, as $\varphi = \sum \varphi_k$ is an eigenfunction of h(v) with eigenvalue λ we have

$$\lambda \|\varphi_j\|_{\mathrm{L}^2}^2 = \lambda \langle \varphi, \varphi_j \rangle_{\mathcal{H}_1} = \langle h(v)\varphi, \varphi_j \rangle_{\mathcal{H}_1} = \int_{\mathbb{T}} \overline{\nabla \varphi(x)} \nabla \varphi_j(x) \mathrm{d}x + v(\overline{\varphi}\varphi_j) = \int_{\mathbb{T}} |\nabla \varphi_j(x)|^2 \mathrm{d}x + v(|\varphi_j|^2).$$

Therefore, for any $\psi \in \text{span}\{\varphi_1, ..., \varphi_M\}$, i.e., $\psi = \sum_{j=1}^M c_j \varphi_j$ for some $\{c_j\}_{j \le M} \subset \mathbb{C}$, we have

$$q_{h(v)}(\psi,\psi) = \sum_{j=1}^{M} |c_j|^2 q_{h(v)}(\varphi_j,\varphi_j) = \lambda \sum_{j=1}^{M} |c_j|^2 ||\varphi_j||_{L^2}^2 = \lambda ||\psi||_{L^2}^2.$$

As dim span{ $\varphi_1, ..., \varphi_M$ } = *M*, the result now follows from the (Courant-Fischer-Weyl) min-max principle.

We are now ready to prove Theorem 2.1.

Proof of Theorem 2.1. By Lemma 4.1 and Lemma 4.3, we see that $\varphi_v(x)$ can vanish on at most one point in \mathbb{T} . Indeed, if there were at least two points where $\varphi_v(x) = 0$, there would be at least two connected components of $\{x \in \mathbb{T} : \varphi_v(x) > 0\}$ which contradicts the non-degeneracy of the ground-state by Lemma 4.3. However, to complete the proof we need to show that $\varphi_v(x)$ can not vanish at a single point.

To this end, we shall argue by contradiction. More precisely, we first assume that there exists a ground-state vanishing at a single-point, and then, via a gluing argument, we construct a ground-state of a different Hamiltonian that vanishes at two distinct points. By the preceding paragraph, this gives a contradiction and suffices to complete the proof.

So suppose that there exists a ground-state $\varphi_v \ge 0$ of h(v) for some $v \in \mathbb{H}^{-1}(\mathbb{T})$ such that $\varphi_v(x) = 0$ only at x = 0. Without loss of generality, we also assume that the ground-state energy is 0. Then, we define

$$\widetilde{\varphi}(x) \coloneqq \varphi_v(x)\eta(x) + \eta(x-\pi/2) + \varphi_v(x-\pi)\eta(x-\pi) + \eta(x-3\pi/2)$$

where $\eta(x) = \eta(x \mod 2\pi)$ for some $\eta \in \mathbb{C}^{\infty}_{c}((-\pi/2, \pi/2); \mathbb{R})$ satisfying

$$\eta(x) = 0$$
 for $|x| \ge \pi/3$, $\eta(x) > 0$ for $|x| < \pi/3$, and $\eta(x) = 1$, for $|x| \le \pi/8$.

Here we denote by $x \mod 2\pi$ the unique $y \in (-\pi, \pi]$ such that $y - x \in 2\pi\mathbb{Z}$. Then, clearly, $\tilde{\varphi} \in H^1(\mathbb{T})$. Moreover, since the support of η is contained in $[-\pi/3, \pi/3]$, we have

$$\begin{cases} \widetilde{\varphi}(x) = \varphi_{v}(x) & \text{for } |x \mod 2\pi| \le \pi/8, \\ \widetilde{\varphi}(x) = \varphi_{v}(x - \pi), & \text{for } |x - \pi| \le \pi/8, \\ \widetilde{\varphi}(x) > 0, & \text{otherwise.} \end{cases}$$

$$(4.1)$$

In particular, $\tilde{\varphi}_v(x) \ge 0$ and $\tilde{\varphi}_v(x) = 0$ if and only if x = 0 or $x = \pi$.

⁶Recall that the inclusion $L^2(\mathbb{T}) \subset H^{-1}(\mathbb{T})$ is given by the Riesz mapping introduced in (3.3).

Next, let $\{p_j\}_{j=1}^3$ be a partition of the unity⁷ subordinate to the following open cover of \mathbb{T} :

$$I_1 := (-\pi/8, \pi/8) \mod 2\pi, \quad I_2 := (7\pi/8, 9\pi/8) \quad \text{and} \quad I_3 := \{x \in \mathbb{T} : |x \mod 2\pi| > \pi/16, \quad |x - \pi| > \pi/16\}.$$

In other words, each of the p_j satisfies $p_j \in C^{\infty}(\mathbb{T})$, $p_j(x) = 0$ for $x \notin I_j$, $p_j(x) \ge 0$, and together they satisfy

$$\sum_{j=1}^{3} p_j(x) = 1, \quad \text{for any } x \in \mathbb{T}.$$
(4.2)

Then, we define the distribution $\widetilde{v}\in \mathcal{D}'(\mathbb{T})$ as

$$\begin{split} \widetilde{v}(\psi) &= v(p_1\psi) + v\left(\tau_{\pi}(p_2\psi)\right) - \int_{\mathbb{T}} \nabla \widetilde{\varphi}(x) \nabla \left(\frac{p_3\psi}{\widetilde{\varphi}}\right)(x) \mathrm{d}x \\ &= v(p_1\psi) + v\left(\tau_{\pi}(p_2\psi)\right) - \int_{I_3} \frac{\nabla \widetilde{\varphi}(x)}{\widetilde{\varphi}(x)} \left(\nabla p_3(x)\psi(x) + p_3(x)\nabla\psi(x)\right) - \frac{\nabla \widetilde{\varphi}(x)^2}{\widetilde{\varphi}(x)^2} p_3(x)\psi(x) \mathrm{d}x, \end{split}$$

where $(\tau_{\pi}\psi)(x) = \psi(x - \pi)$. As $p_j \in C^{\infty}(\mathbb{T})$, the first and second term clearly define distributions in $\mathrm{H}^{-1}(\mathbb{T})$. Moreover, since $\tilde{\varphi} > c$ inside I_3 , by the GNS inequality (3.5), we have

$$|\widetilde{v}(\psi)| \lesssim \left(\|v\|_{\mathrm{H}^{-1}} + 1 + \frac{\|\widetilde{\varphi}\|_{\mathrm{H}^{1}}^{2}}{c^{2}} \right) \|\psi\|_{\mathrm{H}^{1}(\mathbb{T})}, \quad \text{for any } \psi \in \mathrm{H}^{1}(\mathbb{T}).$$

Hence $\tilde{v} \in H^{-1}(\mathbb{T})$. We now observe that, since each p_i is supported on I_i , from (4.1) and the ground-state identity

$$\langle h(v)\varphi_v,\psi\rangle = \int_{\mathbb{T}} \nabla \varphi_v(x)\nabla \psi(x)dx + v(\varphi_v\psi) = 0, \quad \text{for any } \psi \in \mathrm{H}^1(\mathbb{T}),$$

we have

$$\begin{split} -\Delta \widetilde{\varphi}(p_1 \psi) &= \int_{I_1} \nabla \widetilde{\varphi}(x) \nabla (p_1 \psi)(x) dx = \int_{\mathbb{T}} \nabla \varphi_v(x) \nabla (p_1 \psi)(x) dx = -v(p_1 \varphi_v \psi) = -v(p_1 \widetilde{\varphi} \psi), \quad \text{for any } \psi \in \mathrm{H}^1(\mathbb{T}), \\ -\Delta \widetilde{\varphi}(p_2 \psi) &= \int_{I_2} \nabla \widetilde{\varphi}(x) \nabla (p_2 \psi)(x) dx = \int_{\mathbb{T}} \nabla \varphi_v(x - \pi) \nabla (p_2 \psi)(x) dx = \int_{\mathbb{T}} \nabla \varphi_v(x) \nabla (\tau_\pi p_2 \psi)(x) \\ &= -v\left(\varphi_v \tau_\pi(p_2 \psi)\right) = -v\left(\tau_\pi(p_2 \widetilde{\varphi} \psi)\right), \quad \text{for any } \psi \in \mathrm{H}^1(\mathbb{T}), \end{split}$$

and

$$-\Delta \widetilde{\varphi}(p_3 \psi) = \int_{I_3} \nabla \widetilde{\varphi}(x) \nabla (p_3 \psi)(x) dx = \int_{\mathbb{T}} \nabla \widetilde{\varphi}(x) \nabla \left(\frac{p_3}{\widetilde{\varphi}} \widetilde{\varphi} \psi\right)(x) dx, \quad \text{for any } \psi \in \mathrm{H}^1(\mathbb{T}).$$

Hence, by the partition of the unity property (4.2) and the definition of \tilde{v} we have

$$\begin{split} -\Delta \widetilde{\varphi}(\psi) &= -\Delta \widetilde{\varphi}\left(\sum_{j=1}^{3} p_{j}\psi\right) = \sum_{j=1}^{3} -\Delta \widetilde{\varphi}(p_{j}\psi) \\ &= -v(p_{1}\widetilde{\varphi}\psi) - v\left(\tau_{\pi}(p_{2}\widetilde{\varphi}\psi)\right) + \int_{\mathbb{T}} \nabla \widetilde{\varphi}(x) \nabla \left(\frac{p_{3}\widetilde{\varphi}\psi}{\widetilde{\varphi}}\right)(x) \mathrm{d}x = -\widetilde{v}(\widetilde{\varphi}\psi), \quad \text{for any } \psi \in \mathrm{H}^{1}(\mathbb{T}). \end{split}$$

Therefore $\tilde{\varphi}$ is an eigenfunction of $h(\tilde{v})$. Since both $\tilde{\varphi}$ and the ground-state of $h(\tilde{v})$ are almost everywhere strictly positive (by Lemma 4.1), they can not be orthogonal to each other. Therefore, by the non-degeneracy of the ground-state of $h(\tilde{v})$ (see Lemma 4.1), the function $\tilde{\varphi}$ must be the ground-state of $h(\tilde{v})$, which yields the desired contradiction and concludes the proof.

We can now prove Corollary 2.3.

Proof of Corollary 2.3. Let $v \in V$ and denote by φ_v the ground-state of h(v). Since this ground-state is non-degenerate, it follows from Lemma 3.8 that any normalized ground-state of $H_N(v)$ must have φ_v as a natural orbital with occupation number 1. Thus, by Theorem 2.1 and (3.11) we have

$$\rho_{\Psi}(x) = \gamma_{\Psi}(x, x) \ge |\varphi_v(x)|^2 > 0$$
 for any $x \in \mathbb{T}$.

Hence, any pure ground-state has nowhere vanishing density. As the density of a mixed ground-state is a convex combination of the density of pure ground-states, we conclude that any mixed ground-state has nowhere vanishing density.

The integral constraint $\int \rho_{\Psi} = N$ comes from the normalization of Ψ and the regularity condition $\sqrt{\rho_{\Psi}} \in H^1(\mathbb{T})$ comes from the fact that Ψ has finite kinetic energy (and follows from standard arguments, see, [Lie8₃, SPR+24]). This completes the proof.

⁷That such partition of unity exists is a well-known fact in differential geometry. In the simple case of a torus they can be constructed explicitly from any function η as above.

5. The non-interacting Hohenberg-Kohn theorem for distributional potentials

We now turn to the proof of Theorems 2.5 and 2.7. We begin with Theorem 2.7.

Proof of Theorem 2.7. First, note that the map $v_1^{\text{KS}}(\rho) = \frac{\Delta \sqrt{\rho}}{\sqrt{\rho}}$ should be understood in the sense

$$v_{1}^{\mathrm{KS}}(\rho)(\varphi) = \frac{\Delta\sqrt{\rho}}{\sqrt{\rho}}(\varphi) = \Delta\sqrt{\rho}\left(\frac{\varphi}{\sqrt{\rho}}\right) = -\int_{\mathbb{T}} \nabla\sqrt{\rho}(x)\nabla\left(\frac{\varphi}{\sqrt{\rho}}\right)(x)\mathrm{d}x = \Delta\sqrt{\rho}\left(M_{1/\sqrt{\rho}}\varphi\right),$$

where $M_{1/\sqrt{\rho}}$ is the multiplication operator

$$\varphi \mapsto M_{1/\sqrt{\rho}}(\varphi) = \frac{\varphi}{\sqrt{\rho}}$$

Since, for any $\rho \in \mathcal{D}_1$ we have $\rho \ge c$ for some c > 0, we have $\sqrt{\rho} \in H^1(\mathbb{T})$. Thus, from Lemma 3.2, the multiplication operator $M_{1/\sqrt{\rho}}$ is bounded in $H^1(\mathbb{T})$ and therefore $v_1^{\text{KS}}(\rho) \in \mathcal{V}$ for any $\rho \in \mathcal{D}_1$. Moreover, by construction we have

$$-\Delta\sqrt{\rho}(\varphi) + v(\sqrt{\rho}\varphi) = 0$$
, for any $\varphi \in \mathrm{H}^{1}(\mathbb{T})$, where $v = v_{1}^{\mathrm{KS}}(\rho)$.

Therefore, $\sqrt{\rho}$ is an eigenfunction of h(v). As $\sqrt{\rho}$ is strictly positive, it must be the ground-state of h(v) by Theorem 2.1. Hence, v_1^{KS} is indeed the Kohn-Sham map.

To prove that $\rho \mapsto v_1^{\text{KS}}(\rho)$ is smooth, we shall show that it is a composition of smooth maps. So first, note that the map $\psi \in H^1(\mathbb{T}) \mapsto f(\psi) = \Delta \psi \in H^{-1}(\mathbb{T})$ is linear and continuous, hence smooth. Moreover, in the proof of Lemma 3.3, we have shown that the M map $\psi \mapsto M_{\psi}$ is smooth from $H^1(\mathbb{T})$ to $\mathcal{B}(H^1(\mathbb{T}))$. Furthermore, the map $h: H^{-1}(\mathbb{T}) \times \mathcal{B}(H^1(\mathbb{T})) \to H^{-1}(\mathbb{T})$ given by

$$(v,T) \to h(v,T) = v \circ T \in \mathrm{H}^{-1}(\mathbb{T}), \text{ where } v \circ T(\varphi) = v(T\varphi), \text{ for } \varphi \in \mathrm{H}^{1}(\mathbb{T}),$$

is bilinear and continuous, hence, also smooth. Since

$$v_1^{\mathrm{KS}}(\rho) = h\left(f(\sqrt{\rho}), M_{1/\sqrt{\rho}}\right),$$

it suffices to show that the maps $\rho \in \mathcal{D}_1 \mapsto \sqrt{\rho}$ and $\rho \mapsto 1/\sqrt{\rho} \in \mathcal{D}_1$ are also smooth. Moreover, as \mathcal{D}_1 is a submanifold of the set \mathcal{D} introduced in (3.7) (see Remark 3.4), we only need to show that these maps are smooth on \mathcal{D} . To this end, we observe that, since any $\rho \in \mathcal{D}$ satisfies $1/c \leq \rho \leq c$ for some $c = c(\rho) > 0$, by the GNS inequality (3.5) we can find a neighborhood $U_\rho \subset \mathcal{D}$ of $\rho \in \mathcal{D}$ such that $1/(2c(\rho)) \leq \rho' \leq 2c(\rho)$ for any $\rho' \in U_\rho$. In particular, we can find $g_1, g_2 \in C^{\infty}(\mathbb{R}; \mathbb{R})$ such that $g_1(x) = \sqrt{x}$ and $g_2(x) = 1/\sqrt{x}$ for any $1/(2c(\rho)) \leq x \leq 2c(\rho)$. Therefore, by Lemma 3.3, the maps $\rho' \in U_\rho \mapsto g_1(\rho') = \sqrt{\rho'}$ and $\rho' \in U_\rho \mapsto g_2(\rho') = 1/\sqrt{\rho'}$ are smooth. As smoothness is a local property, this shows that $\rho \mapsto \sigma_1^{\mathrm{KS}}(\rho)$ is smooth.

To complete the proof, we need to show that $v_1^{\text{KS}}(\rho)$ is the unique potential generating ρ . So suppose that $\rho \in \mathcal{D}_1$ is the ground-state density of h(v) and h(v') for $v, v' \in H^{-1}(\mathbb{T})$. Without loss of generality, we also assume that both ground-state energies are 0. Since the ground-state is strictly positive and non-degenerate by Theorem 2.1, both operators have the same ground-state wave-function $\varphi = \sqrt{\rho} \in H^1(\mathbb{T})$. Hence, we have

$$0 = \langle \varphi, h(v)\psi \rangle - \langle \varphi, h(v')\psi \rangle = (v - v')(\overline{\varphi}\psi), \quad \text{for any } \psi \in \mathrm{H}^{1}(\mathbb{T}).$$
(5.1)

As φ is strictly positive, the operator of multiplication by $\overline{\varphi}$ is an isomorphism in H¹(T) by Lemma 3.2. Equation (5.1) thus implies that $(v - v') = (v - v') \circ M_{\overline{\varphi}} \circ M_{1/\overline{\varphi}} = 0$, which concludes the proof.

We can now proceed to the proof of Theorem 2.5.

Proof of Theorem 2.5. Let $v, v' \in \mathcal{V}$ and let $N \in \mathbb{N}$. Suppose that Ψ and Ψ' are ground-state wave-functions of $H_N(v)$ and $H_N(v')$ with the same density $\rho \in \mathcal{D}_N$. Moreover, without loss of generality, let us assume that both ground-state energies are zero. We now use the standard Hohenberg-Kohn argument to show that Ψ is also a ground-state of $H_N(v')$. Precisely, we note that, since Ψ is a ground-state of $H_N(v)$ we have

$$0 = \langle \Psi', H_N(v')\Psi' \rangle \leq \langle \Psi, H_N(v')\Psi \rangle = \langle \Psi, H_N(v)\Psi \rangle + (v'-v)(\rho) = (v'-v)(\rho).$$

As the same argument holds when we exchange v and v', we conclude that $(v' - v)(\rho) = 0$. In particular, the above is an equality and Ψ is also a ground-state of $H_N(v')$. Moreover, we note that, if ρ is only representable by a mixed ground-state, i.e., $\rho = \sum_j \lambda_j \rho_{\Psi_j}$ for some $\lambda_j \ge 0$ and Ψ_j ground-states of $H_N(v)$, the same argument shows that each of the Ψ_j is a mutual ground-state of $H_N(v)$ and $H_N(v')$ (see, e.g., [PTC+23, Theorem 1] for more details). So without loss of generality, we denote by Ψ (one of) the mutual ground-states of $H_N(v)$ and $H_N(v')$.

Since Ψ is a simultaneous ground-state of both $H_N(v')$ and $H_N(v)$, we must have

$$\langle (H_N(v) - H_N(v')) \Psi, \Phi \rangle = (v - v')(\rho_{\Psi\Phi}) = 0, \text{ for any } \Phi \in \mathcal{H}_N \cap \mathrm{H}^1(\mathbb{T}^N),$$

where $\rho_{\Psi\Phi}$ is the overlapping density

$$\rho_{\Psi\Phi}(x) = N \int_{\mathbb{T}^{N-1}} \overline{\Psi(x, x_2, ..., x_N)} \Phi(x, x_2, ..., x_N) \mathrm{d}x_2 ... \mathrm{d}x_N.$$

As Ψ is the ground-state of $H_N(v)$ and h(v) has discrete spectrum, by Lemma 3.8 we can choose a basis of eigenfunctions $\{\varphi_j\}_{j\in\mathbb{N}}$ such that $\{\varphi_j\}_{j\leq M}$ for some $M < \infty$ are the natural orbitals (with non-zero occupation number) of Ψ . Thus, if we choose trial states of the form $\Phi = a(\varphi_\ell)^* a(\varphi_k) \Psi$ for $k \leq M$ and $\ell \geq M + 1$, from the formula in Lemma 3.9 we have

$$n_k(v - v')(\overline{\varphi}_k \varphi_\ell) = 0, \quad \text{for any } \ell \ge M + 1 \text{ and } 1 \le k \le M$$
 (5.2)

We can now use the representation in Lemma 3.5 to show that (v - v') is more regular than expected. Precisely, note that the operator $M_{\overline{\varphi_k}}$ of multiplication by $\overline{\varphi_k}$ is a bounded operator in $H^1(\mathbb{T})$ (see Lemma 3.2). Therefore, it follows from (5.2), the fact that $n_j > 0$ for any $j \le M$, and the representation in Lemma 3.5 that

$$(v - v') \circ M_{\overline{\varphi_k}}(\psi) = \sum_{j=1}^M A_{kj} \langle \varphi_j, \psi \rangle \quad \text{for some } \{A_{kj}\}_{k,j \le M} \in \mathbb{C}^{M \times M} \text{ and } \psi \in \mathrm{H}^1(\mathbb{T}).$$
(5.3)

As the ground-state φ_1 is strictly positive by Theorem 2.1, the operator M_{1/φ_1} is bounded in $H^1(\mathbb{T})$ by Lemma 3.2; hence,

$$(v - v')(\psi) = (v - v') \circ M_{\overline{\varphi_1}} \circ M_{1/\overline{\varphi_1}}(\psi) = \sum_{j=1}^M A_{1j}\left(\frac{\varphi_j}{\varphi_1}, \psi\right), \quad \text{for any } \psi \in \mathrm{H}^1(\mathbb{T})$$

In other words, the distribution (v - v') can be identified (via the Riesz map (3.3)) with the function

$$(v - v')(x) = \sum_{j=1}^{M} A_{1j} \frac{\varphi_j(x)}{\varphi_1(x)}.$$
(5.4)

In particular $(v - v') \in H^1(\mathbb{T})$. Hence, to conclude the proof, it suffices to show that (v - v') is piecewise constant. Indeed, if (v - v') was piecewise constant but not constant, it would have a jump, which is not possible for H^1 functions.

To see that (v - v') is piecewise constant, note that by (5.3) we have

$$(v-v')(x)\frac{\varphi_k(x)}{\varphi_1(x)} = \sum_{j=1}^N A_{kj}\frac{\varphi_j(x)}{\varphi_1(x)}, \text{ for any } 1 \le k \le M \text{ and (almost) every } x \in \mathbb{T}.$$

In other words, the above equation shows that the vector

$$\vec{\Phi}(x) \coloneqq \left(1, \frac{\varphi_2(x)}{\varphi_1(x)}, ..., \frac{\varphi_M(x)}{\varphi_1(x)}\right)^T \neq 0$$

is an eigenvector of the matrix $A = \{A_{kj}\} \in \mathbb{C}^{M \times M}$ with corresponding eigenvalue (v - v')(x) for (almost) every $x \in \mathbb{T}$. As $A \in \mathbb{C}^{M \times M}$ can have at most M distinct eigenvalues, we conclude that (v - v')(x) is piecewise constant, which completes the proof.

We now prove Theorem 2.8.

Proof of Theorem 2.8. Let φ_k be a real-valued excited state of h(v) for some $v \in H^{-1}(\mathbb{T})$. Since the ground-state φ_1 is non-degenerate and strictly positive by Theorem 2.1, there must exist some $x_0 \in \mathbb{T}$ such that $\varphi_k(x_0) = 0$, as otherwise φ_k would not change sign and the overlap $\langle \varphi_k, \varphi_1 \rangle$ would be non-zero, contradicting the orthogonality with φ_1 . We can now consider perturbations of v of the form $v_\alpha = v + \alpha \delta_{x_0}$ for $\alpha \in \mathbb{R}$. Indeed, as φ_k is an eigenfunction of h(v) with $\varphi_k(x_0) = 0$, it follows that

$$q_{h(v)}(\varphi_k,\psi) = \langle h(v)\varphi_k,\psi\rangle = \langle h(v)\varphi_k,\psi\rangle + \alpha\varphi_k(x_0)\psi(x_0) = \langle h(v_\alpha)\varphi_k,\psi\rangle, \quad \text{for any } \psi \in \mathrm{H}^1(\mathbb{T})$$

Thus φ_k is also an excited state of $h(v_\alpha)$ for any $\alpha \in \mathbb{R}$. Moreover, as the spectrum of h(v) is discrete, for $\alpha > 0$ small enough, φ_k must be the k^{th} excited state⁸ of $h(v_\alpha)$ by standard perturbation theory.

6. Concluding remarks

In this paper, we obtained a complete characterization of the set of densities that are representable by non-interacting Schrödinger operators with a certain class of distributional potentials in the one-dimensional torus. Moreover, we proved a Hohenberg-Kohn theorem for such operators, thereby establishing the uniqueness of the Kohn-Sham density-to-potential map and, thanks to [SPR+24, Corollary 19], the differentiability of the non-interacting convex Lieb functional. In particular, our results show that, in the non-interacting case, the class of distributions in $H^{-1}(\mathbb{T})$ is not only sufficient but also necessary to represent all strictly positive densities coming from wave-functions with finite kinetic energy.

Let us now comment on possible extensions of these results and further open questions.

(i) First, we emphasize that no necessary conditions for *interacting* \mathcal{V} -representability were established here. Thus an immediate open question is how to extend Theorems 2.4 and 2.5 to the case of interacting systems on \mathbb{T} .

⁸Assuming $k - 1 \in \mathbb{N}$ is the previous closed shell in the case of degeneracies, i.e., $\lambda_{k-1} < \lambda_k$.

- (ii) Second, we note that the unique continuation property restricted to ground-states in Lemma 4.1 relies on powerful abstract results of Reed and Simon that can be applied to a much more general setting such as higher-dimensional Schrödinger operators. While these results are not applicable to interacting systems due to Fermi statistics (see the comments after [RS78, Theorem X.III.46]), they can still be used to establish a Hohenberg-Kohn theorem for non-interacting systems, and therefore, the uniqueness of the Kohn-Sham density to potential map in a much broader context (e.g., for bosons).
- (iii) Third, we note that Theorems 2.1 and 2.5 can be extended to the case of an interval $I = [0, 2\pi]$ with Dirichlet boundary conditions. However, it is not true that all densities that are strictly positive inside I, and come from wave-functions with finite kinetic energy, are representable by potentials in the dual space of $H_0^1(I)$. This can be seen, for instance, by considering densities like $\sqrt{\rho}(x) = x^2$ for x close to 0, in the case of a single-particle. Indeed, in this case, $\Delta\sqrt{\rho}/\sqrt{\rho}(x) = 2/x^2$, which does not define a continuous functional in $H_0^1(I)$. Therefore, it would be interesting to systematically study the properties of the single-particle Kohn-Sham map (2.5) for functions in $H_0^1(I)$. This could lead to further insights into natural necessary and sufficient conditions for v-representability in non-compact spaces such as the line \mathbb{R} , or even to higher dimensional spaces.

Acknowledgements

The author is grateful to Markus Penz, Sarina Sutter, and Michael Ruggenthaler for helpful discussions, useful remarks on the first draft of this paper, and inspiring presentations during the workshop on Foundations and Extensions of DFT in Oslo. The author is also grateful to Andre Laestadius and Vebjørn Bakkestuen for organizing the aforementioned workshop which led to the ideas for the current paper. Special thanks also to Asbjørn Bækgaard Lauritsen for calling attention to Remark 3.7 and spotting a mistake in the previous proof of Theorem 2.5.

T.C. Corso acknowledges funding by the *Deutsche Forschungsgemeinschaft* (DFG, German Research Foundation) - Project number 442047500 through the Collaborative Research Center "Sparsity and Singular Structures" (SFB 1481).

DATA AVAILABILITY

No datasets were generated or analysed during the current study.

Competing interests

The authors have no competing interests to declare that are relevant to the content of this article.

References

| [AS88] | F. ARYASETIAWAN and J. M. STOTT, Effective potentials in density-functional theory, <i>Physical Review B</i> 38 (1988), no.5, pp. 2974–2987. |
|----------|---|
| [CCR85] | J. T. CHAYES, L. CHAYES, and M. B. RUSKAI, Density functional approach to quantum lattice systems, Journal of Statistical Physics 38 (1985), |
| [OTTA] | no.3-4, pp. 497-518. MR 788430. Zbl o632.46072. |
| [CH89] | R. COURANT and D. HILBERT, Methods of mathematical physics. Volume 1, John Wiley & Sons, (1989). MR 65391. Zbl 0729.00007. |
| [CS91] | J. CHEN and M. J. STOTT, v-representability for systems with low degeneracy, <i>Physical Review A</i> 44, no. 5, pp. 2816–2822. |
| [CS93] | J. CHEN and M. J. STOTT, v-representability for noninteracting systems, <i>Physical Review A</i> 47 (1993), no. 1, pp. 153–160. |
| [Gar18] | L. GARRIGUE, Unique continuation for many body Schrödinger operators and the Hohenberg-Kohn theorem, <i>Mathematical Physics, Analysis</i> and <i>Geometry</i> 21 (2018), no.3, Paper No. 27, 11. MR 3854406. Zbl 1400.81217. |
| [Gar19] | L. GARRIGUE, Hohenberg-Kohn theorems for interactions, spin and temperature, Journal of Statistical Physics 177 (2019), no.3, pp. 415-437. |
| [| MR 4026653. Zbl 1429.81057. |
| [Ge079] | V. GEORGESCU, On the unique continuation property for Schrödinger Hamiltonians, <i>Helvetica Physica Acta</i> 52 (1979), no,5-6, pp. 655–670. |
| [Her89] | J. HERCZYŃSKI, JAN, On Schrödinger operators with distributional potentials, <i>Journal of Operator Theory</i> 21 (1989), no.2, pp. 273–295. |
| | MR 1023316. |
| [HK64] | Р. Нонемвегд and W. Kohn, Inhomogeneous electron gas, Physical Review. Series II 136 (1964), B864–B871. MR 180312. Zbl . |
| [KM97] | A. KRIEGL and P. W. MICHOR, The convenient setting of global analysis, American Mathematical Society (1997). MR 1471480. Zbl 0889.58001. |
| [Koh83] | W. Конм, v-Representability and Density Functional Theory, Physical Review Letters 51 (1983), no. 51, pp. 1596–1598. |
| [KS65] | W. KOHN and L. J. SHAM, Self-consistent equations including exchange and correlation effects, <i>Physical Review. Series II</i> 140 (1965), A1132-A1138 MR 180732 |
| [Kuro7] | K KURATA A unique continuation theorem for the Schrödinger equation with singular magnetic field <i>Proceedings of the American</i> |
| [//] | Mathematical Society 125 (1997), no.3, pp. 853–860. MR 1363173. Zbl 0887.35026 |
| [Lam18] | P. E. LAMMERT, In search of the Hohenberg-Kohn theorem. Journal of Mathematical Physics 59 (2018), no.4, pp.042110, 19, MR 3780875, |
| [| Zbl 1386.81157. |
| [LBP20] | A. LAESTADIUS, M. BENEDICKS, and M. PENZ, Unique continuation for the magnetic Schrödinger equation, International Journal of Quantum |
| | Chemistry 120 (2020). |
| [Lie83] | E. H. LIEB, Density functionals for Coulomb systems, International Journal of Quantum Chemistry 24 (1983), pp. 243–277. |
| [PL21] | M. PENZ, and R. v. LEEUWEN, Density-functional theory on graphs, The Journal of Chemical Physics 155 (2021), no. 24, pp. 244111. |
| [PTC+23] | M. PENZ, E. I. TELLGREN, M. A. CSIRIK, M. RUGGENTHALER, and A. LAESTADIUS, The Structure of Density-Potential Mapping. Part I: Standard |
| | Density-Functional Theory, American Chemical Society (2023), 3(4), 334-347. |
| [Rego1] | R. REGBAOUI, Unique continuation from sets of positive measure, In: Carleman estimates and applications to uniqueness and control theory |
| | 46 (2001), 179–190. MR 1839175 Zbl 1165.35332. |
| [RS75] | M. REED and B. SIMON, Methods of modern mathematical physics. II. Fourier analysis, self-adjointness, Academic Press (1975). MR 493420. |
| [RS78] | M. REED and B. SIMON, Methods of modern mathematical physics. IV. Analysis of operators, Academic Press (1978). MR 493421. |
| [RS80] | M. REED and B. SIMON, Methods of modern mathematical physics. I. Functional Analysis, Academic Press (1980). MR 751959 |

T. CARVALHO CORSO

- [SPR+24] S. M. SUTTER, M. PENZ, M. RUGGENTHALER, R. VAN LEUWEEN, and K. J. H. GIESBERTZ, Solution of the v-representability problem on a one-dimensional torus, Journal of Physics. A. Mathematical and Theoretical 57 (2024), no 47. MR 4840211. Zbl 07945786.
- [SS80] M. SCHECHTER and B. SIMON, Unique continuation for Schrödinger operators with unbounded potentials, *Journal of Mathematical Analysis and Applications* 77 (1980), no.2, pp. 482–492. MR 593229. Zbl 0458.35024.
- [THS+22] A. M. TEALE, T. HELGAKER, A. SAVIN, C. ADAMO, B. ARADI, ET AL, DFT exchange: sharing perspectives on the workhorse of quantum chemistry and materials science *Physical chemistry chemical physics* 24 (2022), no. 47, pp. 28700-28781.
- [WAR+23] J. WRIGHTON, A. ALBAVERA-MATA, H. F. RODRÍGUEZ, T. S. TAN, A. C. CANCIO, J. W. DUFTY, and S. B. TRICKEY, Some problems in density functional theory, *Letters in Mathematical Physics* 113 (2023), no.2. MR 4572242. Zbl 1515.81243.
- [Zh012] A. ZHOU, Hohenberg-Kohn theorem for Coulomb type systems and its generalization, *Journal of Mathematical Chemistry* 50 (2012), no.10, pp. 2746–2754. MR 2989094. Zbl 1308.81197.

(T. Carvalho Corso) Institute of Applied Analysis and Numerical Simulation, University of Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany

Email address: thiago.carvalho-corso@mathematik.uni-stuttgart.de

14