Communication-Efficient Stochastic Distributed Learning

Xiaoxing Ren¹, Nicola Bastianello^{2*}, Karl H. Johansson², Thomas Parisini^{1,3,4}

Abstract—We address distributed learning problems, both nonconvex and convex, over undirected networks. In particular, we design a novel algorithm based on the distributed *Alternating Direction Method of Multipliers* (ADMM) to address the challenges of high communication costs, and large datasets. Our design tackles these challenges i) by enabling the agents to perform multiple local training steps between each round of communications; and ii) by allowing the agents to employ stochastic gradients while carrying out local computations. We show that the proposed algorithm converges to a neighborhood of a stationary point, for nonconvex problems, and of an optimal point, for convex problems. We also propose a variant of the algorithm to incorporate variance reduction thus achieving exact convergence. We show that the resulting algorithm indeed converges to a stationary (or optimal) point, and moreover that local training accelerates convergence. We thoroughly compare the proposed algorithms with the state of the art, both theoretically and through numerical results.

Index Terms—Distributed learning; Stochastic optimization; Variance reduction; Local training.

I. INTRODUCTION

Recent technological advances have enabled the widespread adoption of devices with computational and communication capabilities in many fields, for instance, power grids [\[1\]](#page-14-0), robotics [\[2\]](#page-14-1), [\[3\]](#page-14-2), transportation networks [\[4\]](#page-14-3), and sensor networks [\[5\]](#page-14-4). These devices connect with each other, forming multi-agent systems that cooperate to collect and process data [\[6\]](#page-14-5). As a result, there is a growing need for algorithms that enable efficient and accurate cooperative learning.

In specific terms, the objective in distributed learning is to train a model (*e.g.*, a neural network) with parameters $x \in \mathbb{R}^n$ *cooperatively* across a network of N agents. Each agent has access to a local dataset which defines the local cost as

$$
f_i(x) = \frac{1}{m_i} \sum_{h=1}^{m_i} f_{i,h}(x), \qquad (1)
$$

The work of X.R. and T.P. was supported by European Union's Horizon 2020 Research and Innovation programme under grant agreement no. 739551 (KIOS CoE).

The work of N.B. and K.H.J. was partially supported by the European Union's Horizon Research and Innovation Actions programme under grant agreement No. 101070162, and partially by Swedish Research Council Distinguished Professor Grant 2017-01078 Knut and Alice Wallenberg Foundation Wallenberg Scholar Grant.

¹Department of Electrical and Electronic Engineering, Imperial College London, London, United Kingdom.

²School of Electrical Engineering and Computer Science, and Digital Futures, KTH Royal Institute of Technology, Stockholm, Sweden.

³Department of Electronic Systems, Aalborg University, Denmark.

⁴Department of Engineering and Architecture, University of Trieste, Trieste, Italy.

[⋆]Corresponding author. Email: nicolba@kth.se.

with $f_{i,h} : \mathbb{R}^n \to \mathbb{R}$ being the loss function associated to data point $h \in \{1, \ldots, m_i\}$. Thus, the goal is for the agents to solve the following constrained problem [\[7\]](#page-14-6), [\[8\]](#page-14-7):

$$
\min_{x_i \in \mathbb{R}^n, i \in V} \quad \frac{1}{N} \sum_{i=1}^N f_i(x_i) \quad \text{s.t.} \quad x_1 = x_2 = \dots = x_N \,, \tag{2}
$$

where the objective is the sum of local costs [\(1\)](#page-0-0) to pool together the agents' data. Moreover, each agent is assigned a set x_i of local model parameters, and the consensus constraints $x_1 = x_2 = \ldots = x_N$ ensure that the agents will asymptotically agree on a shared trained model.

To effectively tackle this problem, especially when dealing with large datasets that involve sensitive information, distributed methods have become increasingly important. These techniques offer significant robustness advantages over federated learning algorithms [\[6\]](#page-14-5), as they do not rely on a central coordinator and thus, for example, have not a single point of failure. In particular, both distributed gradient-based algorithms [\[17\]](#page-14-8), [\[18\]](#page-14-9), [\[19\]](#page-14-10), [\[20\]](#page-14-11), and distributed Alternating Direction Method of Multipliers (ADMM) [\[21\]](#page-14-12), [\[22\]](#page-14-13), [\[23\]](#page-14-14), [\[24\]](#page-14-15) have proven to be effective strategies for solving such problems.

However, many learning applications face the challenges of: high communication costs, especially when training large models, and large datasets. In this paper, we jointly address these challenges with the following approach. First, we guarantee the communication efficiency of our algorithm by adopting the paradigm of *local training*, which reduces the frequency of communications. In other terms, the agents perform multiple training steps between communication rounds. We tackle the second challenge by locally incorporating *stochastic gradients*. The idea is to allow the agents to estimate local gradients by employing only a (random) subset of the available data, thus avoiding the computational burden of full gradient evaluations on large datasets.

Our main contributions are as follows:

- We propose two algorithms based on distributed ADMM, with one round of communication between multiple local update steps. The first algorithm, Local Training ADMM (LT-ADMM), uses stochastic gradient descent (SGD) for the local updates, while the second algorithm, LT-ADMM with Variance Reduction (LT-ADMM-VR), uses a variance-reduced SGD method [\[25\]](#page-14-16).
- We establish the convergence properties of LT-ADMM for both nonconvex and convex (not strongly convex) learning problems. In particular, we show almost-sure and mean-squared convergence of LT-ADMM to a neighbor-

Algorithm [Ref.]	variance reduction	grad. steps \div comm.	# stored variables [†]	comm. $size$ ^{\ddagger}	# $\nabla f_{i,j}$ evaluations per iteration	assumpt. \star	convergence
$K-GT$ [9]	x	$\tau \div 1$	$\overline{2}$	$2 \mathcal{N}_i $		n.c.	sub-linear, $\propto \sigma^2$
LED [10]	x	$\tau \div 1$	$\overline{2}$	$ \mathcal{N}_i $	1	n.c. s.c.	sub-linear, $\propto \sigma^2$ linear, $\propto \sigma^2$
RandCom [11]	X	$\left \frac{1}{p}\right \div 1$ (in mean)	$\overline{2}$	$ \mathcal{N}_i $	$\mathbf{1}$	n.c. s.c.	sub-linear, $\propto \sigma^2$ linear, $\propto \sigma^2$
VR-EXTRA/DIGing [12], GT-VR [13]		$1 \div 1$	3	$2 \mathcal{N}_i $	$ \mathcal{B} $, m_i every $\left \frac{1}{p} \right $	n.c. s.c.	sub-linear, $\rightarrow 0$ linear, $\rightarrow 0$
GT-SAGA [14], [15]		$1 \div 1$	3	$2 \mathcal{N}_i $	$\mathbf{1}$	n.c. s.c.	sub-linear, $\rightarrow 0$ linear, $\rightarrow 0$
GT-SARAH [16]		$1 \div 1$	3	$2 \mathcal{N}_i $	$ \mathcal{B} $, m_i every τ	n.c.	sub-linear, $\rightarrow 0$
GT-SVRG [14]		$1 \div 1$	3	$2 \mathcal{N}_i $	1, m_i every τ	s.c.	linear, $\rightarrow 0$
LT-ADMM [this work]	x	$\tau \div 1$	$ \mathcal{N}_i +1$	$ \mathcal{N}_i $	$ \mathcal{B} $	n.c.	sub-linear, $\propto \sigma^2$
LT-ADMM-VR [this work]		$\tau \div 1$	$ \mathcal{N}_i $ $+1$	$ {\cal N}_i $	$ \mathcal{B} $, m_i every τ	n.c.	sub-linear, $\rightarrow 0$

TABLE I COMPARISON WITH THE STATE OF THE ART IN STOCHASTIC DISTRIBUTED OPTIMIZATION.

[†] number of vectors in \mathbb{R}^n stored by each agent between iterations (disregarding temporary variables)

‡ number of messages sent by each agent during a communication round

 \star n.c. and s.c. stand for (non)convex and strongly convex

hood of the stationary point in the nonconvex case, and to a neighborhood of an optimum in the convex case. The radius of the neighborhood depends on specific properties of the problem and on tunable parameters. We prove that the algorithm achieves a convergence rate of $\mathcal{O}(\frac{1}{K\tau})$, where K is the number of iterations, and τ the number of local training steps.

- For LT-ADMM-VR, we prove *exact convergence* to a stationary point in the nonconvex case, and to an optimum in the convex case. The algorithm has a $\mathcal{O}(\frac{1}{K_{\tau}})$ rate of convergence, which is faster than $\mathcal{O}(\frac{1}{K})$ obtained by related algorithms [\[16\]](#page-14-24), [\[15\]](#page-14-23), [\[13\]](#page-14-21).
- We provide extensive numerical evaluations comparing the proposed algorithms with the state of the art. The results validate the communication efficiency of the algorithms. Indeed, LT-ADMM and LT-ADMM-VR outperform alternative methods when communications are expensive.

A. Comparison with the State of the Art

We compare our proposed algorithms – LT-ADMM and LT-ADMM-VR– with the state of the art. The comparison is holistically summarized in Table [I.](#page-1-0)

Decentralized learning algorithms, as first highlighted in the seminal paper [\[26\]](#page-14-25) on federated learning, face the fundamental challenge of high communication costs. The authors of [\[26\]](#page-14-25) address this challenge by designing a communication-efficient algorithms which allows the agents to perform multiple local training steps before each round of communication with the coordinator. However, the accuracy of the algorithm in [\[26\]](#page-14-25) degrades significantly when the agents have heterogeneous data. Since then, alternative federated learning algorithms, *e.g.*, [\[27\]](#page-14-26), [\[28\]](#page-14-27), [\[29\]](#page-14-28), [\[30\]](#page-14-29), have been designed to employ local training without compromising accuracy. The interest for communication-efficient algorithms has more recently extended to the distributed set-up, where agents rely on peerto-peer communications rather than on a coordinator, as in federated learning. Distributed algorithms with local training have been proposed in [\[31\]](#page-14-30), [\[9\]](#page-14-17), [\[10\]](#page-14-18), [\[11\]](#page-14-19). In particular, [\[31\]](#page-14-30), [\[9\]](#page-14-17), [\[10\]](#page-14-18) present gradient tracking methods which allow each agent to perform a fixed number of local updates between each communication round. The algorithm in [\[11\]](#page-14-19), which builds on [\[28\]](#page-14-27), instead triggers communication rounds according to a given probability distribution, resulting in a time-varying number of local training steps. Another related algorithm is that of [\[32\]](#page-14-31), which allows for both multiple consensus and gradient steps in each iteration. However, this algorithm requires a monotonically increasing number of communication rounds in order to guarantee exact convergence. A stochastic version of [\[32\]](#page-14-31) was then studied in [\[33\]](#page-14-32). The algorithm has inexact gradient evaluations, but only allows for multiple consensus steps.

When the agents employ stochastic gradients in the algorithms of [\[9\]](#page-14-17), [\[10\]](#page-14-18), [\[11\]](#page-14-19), they only converge to a neighborhood of a stationary point, whose radius is proportional to the stochastic gradient variance. Different *variance reduction* techniques are available to improve the accuracy of (centralized) algorithms relying on stochastic gradients, *e.g.*, [\[34\]](#page-14-33), [\[25\]](#page-14-16), [\[35\]](#page-14-34). Then, these methods have been applied to distributed optimization by combining them with widely used gradient tracking algorithms [\[12\]](#page-14-20), [\[13\]](#page-14-21), [\[14\]](#page-14-22), [\[15\]](#page-14-23), [\[16\]](#page-14-24). The resulting algorithms succeed in guaranteeing exact convergence to a stationary point despite the presence of gradient noise. However, they are not communication-efficient, as they only allow one gradient update per communication round.

We conclude by providing a summary of the features of the algorithms compared so far in Table [I.](#page-1-0) First, we classify them based on whether they use or not variance reduction and local training. For the latter, we report the ratio of gradient steps to communication rounds that characterizes each algorithm, with a ratio of $1 \div 1$ signifying that no local training is used. Notice that LT-ADMM-VR is the only

algorithm to use both variance reduction and local training, while the other only use one technique. We then compare the number of variables stored by the agents when they deploy each algorithm (disregarding temporary variables). We notice that the variable storage of LT-ADMM and LT-ADMM-VR, differently from the alternatives, scales with the size of an agent's neighborhood; this is due to the use of distributed ADMM as the foundation of our proposed algorithms [\[21\]](#page-14-12). We see that [\[10\]](#page-14-18), [\[11\]](#page-14-19), LT-ADMM, and LT-ADMM-VR require one communication per neighbor, while the other methods require two communications per neighbor. We also compare the algorithms by the computational complexity of the gradient estimators they employ, namely, the number of component gradient evaluations needed per local training iteration. The algorithms of [\[9\]](#page-14-17), [\[10\]](#page-14-18), [\[11\]](#page-14-19), [\[14\]](#page-14-22) use a single data point to estimate the gradient, while [\[12\]](#page-14-20), [\[16\]](#page-14-24), LT-ADMM, LT-ADMM-VR can apply mini-batch estimators that use a subset B of the local data points. The use of mini-batches yields more precise gradient estimates and increased flexibility. However, we remark that the gradient estimators used in [\[12\]](#page-14-20), [\[16\]](#page-14-24), [\[14\]](#page-14-22), LT-ADMM, LT-ADMM-VR require a registry of component gradient evaluations, which needs to be refreshed entirely at fixed intervals. This coincides with the evaluation of a full gradient, and thus requires m_i component gradient evaluations. Finally, we compare the algorithms' convergence. We notice that all algorithms, except for [\[14\]](#page-14-22), provide (sub-linear) convergence guarantees for convex and nonconvex problems. Additionally, some works show linear convergence for strongly convex problems. We further distinguish between algorithms which achieve exact convergence due to the use of variance reduction, or inexact convergence with an error proportional to the stochastic gradient variance ($\propto \sigma^2$).

Outline: The outline of the paper is as follows. Section [II](#page-2-0) formulates the problem at hand, and presents the proposed algorithms design. Section [III](#page-3-0) analyzes their convergence, and discusses the results. Section [IV](#page-4-0) reports and discusses numerical results comparing the proposed algorithms with the state of the art. Section [V](#page-5-0) presents some concluding remarks.

Notation: ∇f denotes the gradient of a differentiable function f. Given a matrix $A \in \mathbb{R}^{n \times n}$, $\lambda_{\min}(A)$ and $\lambda_{\max}(A)$ denotes the smallest and largest eigenvalue of A, respectively. $A > 0$ represents that matrix A is positive definite. With $n \in \mathbb{N}$, we let $\mathbf{1}_n \in \mathbb{R}^n$ be the vector with all elements equal to 1, $I \in \mathbb{R}^{n \times n}$ the identity matrix and $0 \in \mathbb{R}^{n \times n}$ the zero matrix. $\langle x, y \rangle = \sum_{h=1}^{n} x_h y_h$ represents the standard inner product of two vectors $x, y \in \mathbb{R}^n$. $\|\cdot\|$ denotes the Euclidean norm of a vector and the matrix-induced 2-norm of a matrix. The proximal of a cost f, with penalty $\rho > 0$, is defined as $\mathrm{prox}_f^{\rho}(y) = \arg \min_{y \in \mathbb{R}^n} \left\{ f(y) + \frac{1}{2\rho} ||y - x||^2 \right\}.$

II. PROBLEM FORMULATION AND ALGORITHM DESIGN

In this section, we formulate the problem at hand and present our proposed algorithms.

A. Problem Formulation

We target the solution of [\(2\)](#page-0-1) over a (undirected) graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$, where $\mathcal{V} = \{1, ..., N\}$ is the set of N agents, and $\mathcal{E} \subset \mathcal{V} \times \mathcal{V}$ is the set of edges (i, j) , $i, j \in \mathcal{V}$. In particular, we assume that the local costs $f_i : \mathbb{R}^n \to \mathbb{R}$ are in the *empirical risk minimization* form [\(1\)](#page-0-0). We make the following assumptions for [\(2\)](#page-0-1), which commonly underpin the convergence analysis of distributed algorithms.

Assumption 1. $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ *is a connected, undirected graph.*

Assumption 2. *The cost function* f_i *of each agent* $i \in V$ *is* L-smooth. That is, there exists $l > 0$ such that $\|\nabla f_i(x) \nabla f_i(y)$ || ≤ L||x – y||, $\forall x, y \in \mathbb{R}^n$. Moreover, f_i is proper: $f_i(x) > -\infty, \forall x \in \mathbb{R}^n$.

When, in the following, we specialize our results to convex scenarios, we resort to the further assumption below.

Assumption 3. *Each function* f_i , $i \in V$, *is convex.*

B. Algorithm Design

We start our design from the distributed ADMM, character-ized by the updates^{[1](#page-2-1)} [\[21\]](#page-14-12):

$$
x_{i,k+1} = \text{prox}_{f_i}^{1/\rho|\mathcal{N}_i|} \left(\frac{1}{\rho |\mathcal{N}_i|} \sum_{j \in \mathcal{N}_i} z_{ij,k} \right), \quad (3a)
$$

$$
z_{ij,k+1} = \frac{1}{2} \left(z_{ij,k} - z_{ji,k} - 2\rho x_{j,k+1} \right) , \qquad (3b)
$$

where $\mathcal{N}_i = \{j \in \mathcal{V} \mid (i,j) \in \mathcal{E}\}\$ denotes the neighbors of agent i, $\rho > 0$ is a penalty parameter, and $z_{ij} \in \mathbb{R}^n$ are auxiliary variables, one for each neighbor of agent i . This algorithm converges in a wide range of scenarios, and, differently from most gradient tracking approaches, shows robustness to many challenges (asynchrony, limited communications, *etc*) [\[21\]](#page-14-12), [\[22\]](#page-14-13). However, the drawback of [\(3\)](#page-2-2) is that the agents need to solve an optimization problem to update x_i , which in general does not have a closed-form solution. Therefore, in practice, the agents need to compute an approximate update [\(3a\)](#page-2-3), which can lead to inexact convergence [\[22\]](#page-14-13).

In this paper, we modify [\(3\)](#page-2-2) to use approximate local updates, *while ensuring that this choice does not compromise exact convergence*. In particular, we allow the agents to use $\tau \in \mathbb{N}$ iterations of a gradient-based solver to approximate [\(3a\)](#page-2-3), which yields the update:

$$
\phi_{i,k}^{0} = x_{i,k},
$$
\n
$$
\phi_{i,k}^{t+1} = \phi_i^t - \gamma \left(g_i(\phi_{i,k}^t) + \rho |\mathcal{N}_i| \phi_{i,k}^t - \sum_{j \in \mathcal{N}_i} z_{ij,k} \right),
$$
\n
$$
t = 0, \dots, \tau - 1,
$$
\n
$$
x_{i,k+1} = \phi_{i,k}^{\tau},
$$
\n(4)

where $g_i(\phi_{i,k}^{\ell})$ is an estimate of the gradient ∇f_i . The resulting algorithm is a distributed gradient method, with the difference that each communication round [\(3b\)](#page-2-4) is preceded by $\tau > 1$ local gradient evaluations. This is an application of the *local training* paradigm [\[10\]](#page-14-18). We remark that the convergence of the proposed algorithm rests on the initialization $\phi_{i,k}^0 = x_{i,k}$,

¹We remark that, more precisely, [\(3\)](#page-2-2) corresponds to the algorithm in [\[21\]](#page-14-12) with relaxation parameter $\alpha = 1/2$.

which enacts a feedback loop on the local training. In general, without this initialization, exact convergence cannot be achieved [\[22\]](#page-14-13).

The local training [\(4\)](#page-2-5) requires a local gradient evaluation or at least its estimate. In the following, we introduce two different estimator options. Notice that the gradient of the penalty term, $\rho |N_i| \phi_{i,k}^t - \sum_{j \in \mathcal{N}_i} z_{ij,k}$, is exactly known and does not need an estimator. The most straightforward idea is to simply employ a local gradient $g_i(\phi) = \nabla f_i(\phi)$. However, in learning applications, the agents may store large datasets $(m_i \gg 1)$. Therefore, computing $\nabla f_i(\phi)$ becomes computationally expensive. To remedy this, the agents can instead use *stochastic gradients*, choosing

$$
g_i(\phi) = \frac{1}{|\mathcal{B}_i|} \sum_{h \in \mathcal{B}_i} \nabla f_{i,h}(\phi), \qquad (5)
$$

where \mathcal{B}_i are randomly drawn indices from $\{1, \ldots, m_i\}$, with $|\mathcal{B}_i| < m_i$. While reducing the computational complexity of the local training iterations, the use of stochastic gradients results in inexact convergence. The idea therefore is to employ a gradient estimator based on a *variance reduction* scheme. In particular, we use the scheme proposed in [\[25\]](#page-14-16), which demonstrates faster convergence in practice and is characterized by the following procedure. Each agent maintains a table of component gradients $\{\nabla f_{i,h}(r_{i,h,k}^t)\}, h = 1, \ldots, m_i,$ where $r_{i,h,k}^t$ is the most recent iterate at which the component gradient was evaluated. This table is reset at the beginning of every new local training (that is, for any $k \in \mathbb{N}$ when $t = 0$). Using the table, the agents then estimate their local gradients as

$$
g_i\left(\phi_{i,k}^t\right) = \frac{1}{|\mathcal{B}_i|} \sum_{h \in \mathcal{B}_i} \left(\nabla f_{i,h}\left(\phi_{i,k}^t\right) - \nabla f_{i,h}\left(r_{i,h,k}^t\right)\right) \\ + \frac{1}{m_i} \sum_{h=1}^{m_i} \nabla f_{i,h}(r_{i,h,k}^t). \tag{6}
$$

The gradient estimate is then used to update $\phi_{i,k}^{t+1}$ according to [\(4\)](#page-2-5); afterwards, the agents update their local memory by setting $r_{i,h,k}^{t+1} = \phi_{i,k}^{t+1}$ if $h \in \mathcal{B}_i$, and $r_{i,h,k}^{t+1} = r_{i,h,k}^t$ otherwise. Notice that this update requires a full gradient evaluation at the beginning of each local training, to populate the memory with $\{\nabla f_{i,h}(r_{i,h,k}^0) = \nabla f_{i,h}(\phi_{i,k}^0)\}, h = 1, \ldots, m_i$. In the following steps $(t > 0)$, each agent only computes $|\mathcal{B}_i|$ component gradients.

The resulting algorithms are reported in Algorithm [1.](#page-3-1)

III. CONVERGENCE ANALYSIS AND DISCUSSION

In this section, we analyze the convergence rate of Algorithm [1](#page-3-1) in both nonconvex and convex scenarios. Throughout, we will employ the following metric of convergence

$$
\mathcal{D}_k = \mathbb{E}\left[\left\|\nabla F\left(\bar{x}_k\right)\right\|^2 + \frac{1}{\tau} \sum_{t=0}^{\tau-1} \left\|\frac{1}{N} \sum_{i=1}^N \nabla f_i\left(\phi_{i,k}^t\right)\right\|^2\right],\tag{7}
$$

where $F(x) = \frac{1}{N} \sum_{i=1}^{N} f_i(x)$ and $\bar{x}_k = \frac{1}{N} \sum_{i=1}^{N} x_i$. If the agents' converge to a stationary point of [\(2\)](#page-0-1), then $\mathcal{D}_k \to 0$.

Algorithm 1 LT-ADMM and LT-ADMM-VR

- **Input:** For each node *i*, initialize $x_{i,0} = z_{ij,0}, j \in \mathcal{N}_i$. Set the penalty ρ , the number of local training steps τ , and the local step-size γ .
- 1: for $k = 0, 1, \ldots$ every agent i do // local training 2: $\phi_{i,k}^0 = x_{i,k}, \ r_{i,h,k}^0 = x_{i,k}$, for all $h \in \{1, \ldots, m_i\}$ 3: **for** $t = 0, 1, \ldots, \tau - 1$ **do**
4: **draw** the batch B_i unit 4: draw the batch B_i uniformly at random
5: **update the gradient estimator according** update the gradient estimator according to (5) 6: update the gradient estimator according to [\(6\)](#page-3-3) 7: **update** $\phi_{i,k}$ according to [\(4\)](#page-2-5) 8: if $h \in \mathcal{B}_i$ update $r_{i,h,k}^{t+1} = \phi_{i,k}^{t+1}$, else $r_{i,h,k}^{t+1} = r_{i,h,k}^t$ 9: end for 10: $x_{i,k+1} = \phi_{i,k}^{\tau}$ // communication 11: transmit $z_{ij,k} - 2\rho x_{i,k+1}$ to each neighbor $j \in \mathcal{N}_i$, and receive the corresponding transmissions

// auxiliary update

12: update $z_{i,j,k+1}$ according to [\(3b\)](#page-2-4)

13: end for

A. Convergence with SGD

We start by characterizing the convergence of Algorithm [1](#page-3-1) when the agents use SGD during local training. To this end, we make the following standard assumption on the variance of the gradient estimators.

Assumption 4. *For all* $\phi \in \mathbb{R}^n$ *the gradient estimators* $g_i(\phi)$ *,* i ∈ V*, in* [\(5\)](#page-3-2) *are unbiased and their variance is bounded by some* $\sigma^2 > 0$ *:*

$$
\mathbb{E}\left[g_i(\phi) - \nabla f_i(\phi)\right] = 0
$$

$$
\mathbb{E}\left[\|g_i(\phi) - \nabla f_i(\phi)\|^2\right] \leq \sigma^2
$$

We are now ready to state our convergence results. All the proofs are deferred to the Appendix.

Theorem 1 (Nonconvex case). *Let Assumptions [1,](#page-2-6) [2,](#page-2-7) and [4](#page-3-4) hold. If the local step-size* γ *is chosen sufficiently small (see* [\(10\)](#page-5-1) *for the precise bound), then the output of LT-ADMM satisfies:*

$$
\frac{1}{K}\sum_{k=0}^{K-1} \mathcal{D}_k \leq \mathcal{O}\left(\frac{F(\bar{x}_0) - F(x^*)}{K\gamma\tau}\right) + \mathcal{O}\left(\gamma\tau\sigma^2\right) + \mathcal{O}\left(\frac{\|\widehat{\mathbf{d}}_0\|^2}{\rho K N}\right),\tag{8}
$$

where x^* *is a stationary point of* [\(2\)](#page-0-1)*, and* $||\mathbf{d}_0||$ *is related to the initial conditions (see* [\(26\)](#page-7-0)*).*

Theorem [1](#page-3-5) shows that LT-ADMM converges to a neighborhood of a stationary point x^* as $K \to \infty$. The radius of this neighborhood is proportional to the step-size γ , to the number of local training epochs τ , and to the stochastic gradient variance σ^2 . The result can then be particularized to the convex case as follows.

Corollary 1 (Convex case). *In the setting of Theorem [1,](#page-3-5) with the additional Assumption [3,](#page-2-8) then the output of LT-ADMM converges to a neighborhood of an optimal solution* x ∗ *characterized by* [\(8\)](#page-3-6)*.*

Remark 1 (Exact convergence). *Clearly, if we employ full gradients (and thus* $\sigma = 0$), then these results prove exact *convergence to a stationary/optimal point. This verifies that our algorithm design achieves convergence despite the use of approximate local updates.*

B. Convergence with Variance Reduction

The results of the previous sections show that only inexact convergence can be achieved when employing SGD. The following results show how Algorithm [1](#page-3-1) achieves exact convergence when using variance reduction.

Theorem 2 (Nonconvex case). *Let Assumptions [1,](#page-2-6) [2,](#page-2-7) and [4](#page-3-4) hold. If the local step-size* γ *is chosen sufficiently small (see* [\(11\)](#page-6-0) *for the precise bound), then the output of LT-ADMM converges to a stationary point* x [∗] *of* [\(2\)](#page-0-1)*, and in particular it holds:*

$$
\frac{1}{K} \sum_{k=0}^{K-1} \mathcal{D}_k \le \mathcal{O}\left(\frac{F\left(\bar{x}_0\right) - F(x^*)}{K\gamma\tau}\right) + \mathcal{O}\left(\frac{\|\widehat{\mathbf{d}}_0\|^2}{\rho^2 K N}\right) \tag{9}
$$

Corollary 2 (Convex case). *In the setting of Theorem [2,](#page-4-1) with the additional Assumption [3,](#page-2-8) then the output of LT-ADMM-VR converges to an optimal solution* x ∗ *, with rate characterized by* [\(9\)](#page-4-2)*.*

C. Discussion

1) Choice of step-size: The upper bounds to the stepsizes of LT-ADMM and LT-ADMM-VR ([\(10\)](#page-5-1) and [\(11\)](#page-6-0) in Appendix [A-A\)](#page-5-2), highlight a dependence on several *features of the problem*. In particular, the step-size bounds decrease as the smoothness constant L increases, as is usually the case for gradient-based algorithms. Moreover, the bounds are proportional to the network connectivity, represented by the second-largest eigenvalue of G's Laplacian ($|\lambda_{\text{max}}|$). Thus, less connected graphs (smaller $|\lambda_{\text{max}}|$) result in smaller bounds. Finally, we remark that the step-size bound for LT-ADMM-VR is proportional to $\frac{m_l}{m_u} = \frac{\min_{i \in \mathcal{V}} m_i}{\max_{i \in \mathcal{V}} m_i}$ $\frac{\min_{i \in \mathcal{V}} m_i}{\max_{i \in \mathcal{V}} m_i}$, where m_i is the number of data points available to agent i (see [\(1\)](#page-0-0)). This ratio can be viewed as a measure of heterogeneity between the agents. Smaller values of $\frac{m_l}{m_u}$ highlight higher imbalance in the amount of data available to the agents. The step-size bound thus is smaller for less balanced scenarios.

The step-size bounds also depend on the *tunable parameters* τ , the number of local updates, and ρ , the penalty parameter. Therefore, these two parameters can be tuned in order to increase the step-size bounds, which translates in faster convergence.

5

2) Convergence rates comparison: As discussed in section [I-A,](#page-1-1) various distributed algorithms with variance reduction have been recently proposed, for example, [\[12\]](#page-14-20), [\[14\]](#page-14-22) for strongly convex problems, and [\[16\]](#page-14-24), [\[15\]](#page-14-23), [\[13\]](#page-14-21) for nonconvex problems. Focusing on [\[16\]](#page-14-24), [\[15\]](#page-14-23), [\[13\]](#page-14-21), we notice that their convergence rate is $\mathcal{O}(\frac{1}{K})$, while Theorem [2](#page-4-1) shows that LT-ADMM-VR has rate of $\mathcal{O}(\frac{1}{\tau K})$. This shows that *employing local training accelerates convergence*.

Similarly to LT-ADMM-VR, [\[16\]](#page-14-24), [\[12\]](#page-14-20) also use batch gradient computations, *i.e.*, they update a subset of components to estimate the gradient (see [\(6\)](#page-3-3)). Interestingly, the step-size upper bound and, hence, the convergence rate in [\[16\]](#page-14-24), [\[12\]](#page-14-20) depend on the batch size. On the other hand, our theoretical results are not affected by the batch size, since we use a different variance reduction technique.

3) Choice of variance reduction mechanism: In variance reduction, we distinguish two main classes of algorithms: those that need to periodically (or randomly) perform a full gradient evaluation (SARAH-type [\[35\]](#page-14-34)), and those that do not (SAGAtype [\[25\]](#page-14-16)). In distributed learning, SARAH-type algorithms were proposed in *e.g.*, [\[16\]](#page-14-24), [\[14\]](#page-14-22), while SAGA-type algorithms in *e.g.* [\[14\]](#page-14-22). Also the proposed LT-ADMM-VR requires a periodic full gradient evaluation, as the agents re-initialize their local gradient memory at the start of each local training (since they set $r_{i,h,k}^0 = x_{i,k}$). Clearly, periodically computing a full gradient significantly increases the computational complexity of the algorithm. Thus, one can design a SAGA-type variant of LT-ADMM-VR by removing the gradient memory re-initialization at the start of local training (choosing now $r^0_{i,h,k} = r^{\tau}_{i,h,k-1}$). This variant is computationally cheaper and shows promising empirical performance, see the results for LT-ADMM-VR v2 in section [IV.](#page-4-0) However, using the outdated gradient memory leads to a more complex theoretical analysis, which we leave for future work.

4) Uncoordinated parameters: In principle, the agents could use *uncoordinated* parameters, based on the resources available to them. For example, different agents could employ different local solvers and batch sizes. Alternatively, they could use the same solver but with different step-sizes tailored to the smoothness of the local costs.

IV. NUMERICAL RESULTS

In this section we compare the proposed algorithms with the state of the art, applying them to a classification problem with nonconvex regularization, characterized by [\[10\]](#page-14-18):

$$
f_i(x) = \frac{1}{m_i} \sum_{h=1}^{m_i} \log \left(1 + \exp \left(-b_{i,h} a_{i,h}^\top x \right) \right) + \epsilon \sum_{\ell=1}^n \frac{[x]_{\ell}^2}{1 + [x]_{\ell}^2}
$$

where $[x]_l$ is the l -th component of $x \in \mathbb{R}^n$, and $a_{i,h} \in \mathbb{R}^n$ and $b_{i,h} \in \{-1,1\}$ are the pairs of feature vector and label, randomly generated with scikit-learn. We choose a ring graph with $N = 10$, and set $n = 5$, $m_i = 100$, $\epsilon = 0.01$; the initial conditions are randomly chosen as $x_{i,0} \sim \mathcal{N}(0, 100I_n)$. We use stochastic gradients with a batch of $|\mathcal{B}| = 1$. All results are averaged over 100 Monte Carlo iterations. For the algorithms with local training we select $\tau = 5$. We also tune the step-sizes of all algorithms to ensure best performance.

Finally, as performance metric we employ $\|\nabla F(\bar{x}_k)\|^2$, which is zero if the agents have reached a solution of [\(2\)](#page-0-1).

We compare LT-ADMM and LT-ADMM-VR with local training algorithms LED [\[10\]](#page-14-18) and K-GT [\[9\]](#page-14-17), as well as variance reduction algorithms GT-SARAH [\[16\]](#page-14-24), and GT-SAGA [\[14\]](#page-14-22). We also compare with the alternative version LT-ADMM-VR v2 discussed in section [III-C3.](#page-4-3) We account for the computation time (in unspecified units) of each algorithm when scaling the x-axis. This choice deviates from the common approach of using the iteration count for the x-axis, since we are interested in simulating the real performance of these algorithms. In particular, letting t_G be the time for a component gradient evaluation ($\nabla f_{i,h}$), and t_{C} the time for a round of communications, Table [II](#page-5-3) reports the computation time incurred by each algorithm over the course of τ iterations.

TABLE II COMPUTATION TIME OF THE ALGORITHMS OVER τ iterations.

Algorithm [Ref.]	Time			
LED [10] & K-GT [9]	$\tau t_{\rm G}+2t_{\rm C}$			
GT-SARAH [16]	$(m_i + \tau - 1)t_G + 2\tau t_G$			
GT-SAGA [14]	$\tau (t_{\rm G} + 2t_{\rm C})$			
LT-ADMM & LT-ADMM-VR v2	$\tau t_{\rm G}+t_{\rm C}$			
I.T.ADMM-VR	$(m_i + \tau - 1)t_{\rm G} + t_{\rm C}$			

Figure [1](#page-5-4) depicts the comparison of the algorithms in Table [II](#page-5-3) in terms of $\|\nabla F(\bar{x}_k)\|^2$. The x-axis for each algorithm is scaled to account for the individual computation time, and we plot the results for three choice of the t_G/t_C ratio. First of

Fig. 1. Algorithms' comparison for different t_G/t_C ratios. The x-axis is in unspecified computation time units to account for the algorithms' computational complexity.

all, we notice that the algorithms without variance reduction (LED, K-GT, LT-ADMM) as expected only converge to a neighborhood of the optimal solution, as stochastic gradients are employed. The asymptotic errors attained by the algorithms are: LED 8.169×10^{-4} , K-GT 7.552×10^{-4} , LT-ADMM 6.327×10^{-4} .

Turning to the algorithms with variance reduction, we see that all of them converge to the optimal solution; however, depending on the value of t_G/t_C , their relative speed of convergence changes. When gradient computations are cheaper than communications ($t_{\text{G}}/t_{\text{C}} = 0.1$), the proposed algorithm LT-ADMM-VR (and LT-ADMM-VR v2) outperform both GT-SARAH and GT-SAGA, since the latter do not employ local training. *This testifies to the benefit of employing local training in scenarios where communications are expensive*. As the ratio $t_{\rm G}/t_{\rm C}$ increases to 1 and then 10, we see how LT-ADMM-VR and GT-SARAH, on the one hand, and LT-ADMM-VR v2 and GT-SAGA, on the other hand, tend to align in terms of performance, as the bulk of the computation time is now due to gradient evaluations, of which the two pairs of algorithms have a similar number (see Table [II\)](#page-5-3).

We conclude evaluating the convergence speed of LT-ADMM-VR for different choices of the number of local training epochs τ . Figure [2](#page-5-5) reports the computation time (scaled according to Table [II\)](#page-5-3) to reach $\|\nabla F(\bar{x}_k)\|^2 < 10^{-9}$. Interestingly, it appears that there is a finite optimal value

Fig. 2. Computation time needed by LT-ADMM-VR to reach $\|\nabla F(\bar{x}_k)\|^2 <$ 10⁻⁹ for different numbers of local training epochs τ.

 $(\tau = 8)$, while smaller and larger values lead to slower convergence.

V. CONCLUDING REMARKS

In this paper, we considered (non)convex distributed learning problems. In particular, to address the challenge of expensive communication, we proposed two communicationefficient algorithms, LT-ADMM and LT-ADMM-VR, that use local training. The algorithms employ SGD and SGD with variance reduction, respectively. We have shown that LT-ADMM converges to a neighborhood of a stationary point, while LT-ADMM-VR converges exactly. We have thoroughly compared our algorithms with the state of the art, both theoretically and in simulations. Future research will focus on analyzing convergence for strongly convex problems, and extending our algorithm design to asynchronous scenarios.

APPENDIX A

PRELIMINARY ANALYSIS

In this section we summarize the step-size bounds, and present preliminary results underpinning Theorems [1](#page-3-5) and [2.](#page-4-1)

A. Step-size bounds

The step-size upper bounds for LT-ADMM and LT-ADMM-VR are, respectively:

$$
\bar{\gamma}_{sgd} := \min_{i=1,2,\dots,7} \bar{\gamma}_i,\tag{10}
$$

$$
\bar{\gamma}_{\text{sarah}} := \min_{i=1,8,9,\dots,17} \bar{\gamma}_i,\tag{11}
$$

,

where:

$$
\bar{\gamma}_{1} := \min\left\{1, \frac{2}{|\tilde{\lambda}_{\min}|\rho\tau}\right\}, \quad \bar{\gamma}_{2} := \frac{1}{4\sqrt{\tau(\tau-1)(L^{2}+\rho^{2}d_{u}^{2})}}
$$
\n
$$
\bar{\gamma}_{3} := \frac{3}{16L\tau}, \quad \bar{\gamma}_{4} := \frac{1}{8\tau\sqrt{L^{2}+4L\rho^{2}d_{u}^{2}\tau+4\rho^{2}d_{u}^{2}}},
$$
\n
$$
\bar{\gamma}_{5} := \frac{|\tilde{\lambda}_{\max}|^{2}\rho^{2}}{4(\beta_{0}\kappa_{1}+\beta_{1})}, \bar{\gamma}_{6} := \frac{|\tilde{\lambda}_{\max}|^{2}\rho^{2}}{8(\beta_{0}\kappa_{1}+\beta_{1})},
$$
\n
$$
\bar{\gamma}_{7} := \max\left\{\frac{|\tilde{\lambda}_{\max}|^{2}\rho^{2}}{256\kappa_{4}\tau^{2}N\beta_{0}}, \frac{|\tilde{\lambda}_{\max}|^{2}\rho^{2}\tau^{2}}{48L^{2}\|\hat{\mathbf{V}}^{-1}\|^{2}N}\right\}
$$
\n
$$
\bar{\gamma}_{8} := \frac{m_{l}}{8L^{2}m_{u}}, \quad \bar{\gamma}_{9} := \min\left\{\frac{1}{2(4L^{2}+2\rho^{2}d_{u}^{2})}, \frac{1}{2}\right\}
$$
\n
$$
\bar{\gamma}_{10} := \frac{\sqrt{m_{l}}}{\tau L\sqrt{384m_{u}}}, \quad \bar{\gamma}_{11} := \frac{1}{2\sqrt{6\tau(\tau-1)(3L^{2}+\rho^{2}d_{u}^{2})}}
$$
\n
$$
\bar{\gamma}_{12} := \frac{1}{8\sqrt{2\alpha_{0}\tau^{2}N+(12\alpha_{0}\tau L^{2}+\frac{L^{3}}{N})\tilde{s}_{2}}}, \quad \bar{\gamma}_{13} := \frac{3}{16L\tau}
$$
\n
$$
\bar{\gamma}_{14} := \frac{|\tilde{\lambda}_{\max}|^{2}\rho^{2}}{4(\tilde{\beta}_{0}\kappa_{1}+\tilde{\beta}_{1}+(96\tau^{2}L^{2}\tilde{\beta}_{0}+2\tilde{\beta}_{3})(\tilde{s}_{0}+\tilde{s}_{1}))}
$$
\n
$$
\bar{\gamma}_{15} := \frac{|\til
$$

These bounds depend on the following quantities. d_u = $\max\{|\mathcal{N}_i|\}_{i\in\mathcal{V}}$ denotes the maximum agents' degree. V is defined in [\(25\)](#page-7-1). $\lambda_{\min} = \lambda_{\min}(\mathbf{Q}^T \mathbf{L} \mathbf{Q})$ and λ_{\max} = $\lambda_{\text{max}}(\mathbf{Q}^T \mathbf{L} \mathbf{Q})$, with $-\mathbf{L}$ being the graph Laplacian (see [\(15\)](#page-6-1)), and \hat{Q} is being defined at the beginning of Appendix [A-C.](#page-7-2) We denote $m_u = \max_{i=1,\dots,N} m_i$ and $m_l = \min_{i=1,\dots,N} m_i$, where m_i is the number of local data points of agent i. Additionally, we have the following definitions, used both in the upper bound above and throughout the convergence analysis:

$$
\beta_0 := 4(1 + 2\rho^2 \|\tilde{\mathbf{L}}\|^2) \|\hat{\mathbf{V}}^{-1}\|^2 (L^2 + \rho^2 d_u^2) \n+ 12L^2 \|\hat{\mathbf{V}}^{-1}\|^2 \rho^2 d_u^2 \n\beta_1 := (1 + 2\rho^2 \|\tilde{\mathbf{L}}\|^2) \|\hat{\mathbf{V}}^{-1}\|^2 \tau \rho d_u^2 72 / |\tilde{\lambda}_{\text{max}}| \n+ L^2 \|\hat{\mathbf{V}}^{-1}\|^2 18 \rho d_u^2 \tau 216 / |\tilde{\lambda}_{\text{max}}| \n\kappa_1 := \frac{72\tau}{|\tilde{\lambda}_{\text{max}}|\rho} + (18 + \frac{36\tau \rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho}) 16\tau^2 \n\kappa_4 := \frac{8}{N} \left((\frac{72\tau}{|\tilde{\lambda}_{\text{max}}|\rho} + (18 + \frac{36\tau \rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho}) 16\tau^2) \right) \n\times (\frac{L^2}{2} + 2L\rho^2 d_u^2 \tau + 2\rho^2 d_u^2) \n+ \frac{2\tau}{|\tilde{\lambda}_{\text{max}}|\rho} (18\rho^2 d_u^2 + 18\rho^2 d_u^2 \tau L) \right), \n\tilde{s}_0 := \frac{36m_u}{|\tilde{\lambda}_{\text{max}}|\rho} + \frac{16m_u}{m_l} \left(18 + \frac{36\tau \rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho} \right)
$$

$$
\tilde{s}_1 := \left(\frac{72\tau}{|\tilde{\lambda}_{\text{max}}|\rho} + (18 + \frac{36\tau\rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho})16\tau^2\right) \frac{4m_u}{m_l}
$$
\n
$$
\tilde{s}_2 := \frac{64m_u}{m_l} \tau^2 N + \frac{16m_u}{m_l} N,
$$
\n
$$
\tilde{\beta}_0 := (1 + 2\rho^2 ||\tilde{\mathbf{L}}||^2) ||\hat{\mathbf{V}}^{-1}||^2 4(3L^2 + \rho^2 d_u^2)
$$
\n
$$
+ 6L^2 ||\hat{\mathbf{V}}^{-1}||^2 (2\rho^2 d_u^2 + 4L^2)
$$
\n
$$
\tilde{\beta}_1 := (1 + 2\rho^2 ||\tilde{\mathbf{L}}||^2) ||\hat{\mathbf{V}}^{-1}||^2 \tau \rho d_u^2 72/|\tilde{\lambda}_{\text{max}}|
$$
\n
$$
+ L^2 ||\hat{\mathbf{V}}^{-1}||^2 18\rho d_u^2 \tau 216/|\tilde{\lambda}_{\text{max}}|
$$
\n
$$
\tilde{\beta}_3 := (1 + 2\rho^2 ||\tilde{\mathbf{L}}||^2) ||\hat{\mathbf{V}}^{-1}||^2 8L^2 + 6L^2 ||\hat{\mathbf{V}}^{-1}||^2 4L^2,
$$
\n
$$
\alpha_0 := \frac{L^2}{2N} + \frac{2L\rho^2 d_u^2 \tau + 2\rho^2 d_u^2 + 4\tau L^3}{N}
$$
\n
$$
\alpha_1 := \frac{72\tau^2}{|\tilde{\lambda}_{\text{max}}|\rho} + (18 + \frac{36\tau\rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho})16\tau^3
$$
\n
$$
\alpha_5 := (36\rho d_u^2 \tau^2 + 36\rho d_u^2 \tau^3 L) / (N|\tilde{\lambda}_{\text{max}}|)
$$
\n
$$
\mu := 32(\alpha_5 + \alpha_0 \alpha_1 + 2(48\tau^2 L^2 \alpha_0 + 4\tau L^3/N)(\tilde{s}_0 + \tilde{s}_1)).
$$

B. Preliminary transformation

We start by rewriting the algorithm in a compact form. To this end, we introduce the following auxiliary variables: $\mathbf{Z} = \text{col}\{z_{ij}\}_{i,j\in\mathcal{E}}, \ \ \mathbf{\Phi}_k^t$ $\begin{array}{lll} t_k & = & \mathrm{col}\{\phi_{1,k}^t, \phi_{2,k}^t, ..., \phi_{N,k}^t\}, \end{array}$ $G(\Phi_k^t)$ = col $\{g_1(\phi_{1,k}^t), g_2(\phi_{2,k}^t), ..., g_N(\phi_{N,k}^t)\},$ $F(X) = \text{col}\{\frac{1}{N}f_1(x_1), \frac{1}{N}f_2(x_2), ..., \frac{1}{N}f_N(x_N)\}.$ Define $\mathbf{A} = \text{blk diag}\{\mathbf{1}_{d_i}\}_{i \in \mathcal{V}} \otimes \mathbf{I}_n \in \mathbb{R}^{Mn \times Nn}$, where $d_i = |\mathcal{N}_i|$ is the degree of node i, and $M = \sum_i |\mathcal{N}_i|$. $\mathbf{P} \in \mathbb{R}^{Mn \times Mn}$ is a permutation matrix that swaps e_{ij} with e_{ji} . If there is an edge between nodes i, j, then $A^T[i, :] PA[:, j] = 1$, otherwise $A^{T}[i, :] PA[:, j] = 0$. Therefore $\mathbf{A}^{T} \mathbf{P} \mathbf{A} = \mathbf{A}$ is the adjacency matrix.

The compact form of LT-ADMM and LT-ADMM-VR then is:

$$
\mathbf{X}_{k+1} = \mathbf{X}_k - \gamma \sum_{t=0}^{\tau-1} (\nabla G(\boldsymbol{\Phi}_k^t) + \rho \mathbf{A}^T \mathbf{A} \boldsymbol{\Phi}_k^t - \mathbf{A}^T \mathbf{Z}_k) \tag{12a}
$$

$$
\mathbf{Z}_{k+1} = \frac{1}{2} \mathbf{Z}_k - \frac{1}{2} \mathbf{P} \mathbf{Z}_k + \rho \mathbf{P} \mathbf{A} \mathbf{X}_{k+1}. \tag{12b}
$$

Moreover, we introduce the following useful variables

$$
\mathbf{Y}_k = \mathbf{A}^T \mathbf{Z}_k - \nabla F(\bar{\mathbf{X}}_k) - \rho \mathbf{D} \mathbf{X}_k \n\tilde{\mathbf{Y}}_k = \mathbf{A}^T \mathbf{P} \mathbf{Z}_k + \nabla F(\bar{\mathbf{X}}_k) - \rho \mathbf{D} \mathbf{X}_k,
$$
\n(13)

where $\bar{\mathbf{X}}_k = \mathbf{1}_N \otimes \bar{x}_k$, with $\bar{x}_k = \frac{1}{N} \mathbf{1}^T \mathbf{X}_k$, and $\mathbf{D} = \mathbf{A}^T \mathbf{A} =$ diag $\{d_i\mathbf{I}_n\}_{i\in\mathcal{V}}$ is the degree matrix.

Multiplying both sides of [\(12b\)](#page-6-2) by 1^T , and using the initial condition, we obtain $\mathbf{1}^T \mathbf{A}^T \mathbf{Z}_{k+1} = \rho \mathbf{1}^T \mathbf{D} \mathbf{X}_{k+1}$ for all $k \in \mathbb{N}$. As a consequence $\overline{Y}_k = 1 \otimes \frac{1}{N} 1^T \nabla F(\overline{X}_k) =$ $1 \otimes \frac{1}{N} \sum_i \nabla f_i(\bar{x}_k)$, and [\(12\)](#page-6-3) can be further rewritten as

$$
\begin{bmatrix} \mathbf{X}_{k+1} \\ \mathbf{Y}_{k+1} \\ \tilde{\mathbf{Y}}_{k+1} \end{bmatrix} = \begin{bmatrix} \mathbf{I} & \gamma \tau \mathbf{I} & \mathbf{0} \\ \rho \tilde{\mathbf{L}} & \rho \tilde{\mathbf{L}} \gamma \tau + \frac{1}{2} \mathbf{I} & -\frac{1}{2} \mathbf{I} \\ \mathbf{0} & -\frac{1}{2} \mathbf{I} & \frac{1}{2} \mathbf{I} \end{bmatrix} \otimes \mathbf{I}_n \begin{bmatrix} \mathbf{X}_k \\ \mathbf{Y}_k \\ \tilde{\mathbf{Y}}_k \end{bmatrix} - \mathbf{h}_k, \tag{14}
$$

where

$$
\tilde{\mathbf{L}} = \tilde{\mathbf{A}} - \mathbf{D} \tag{15}
$$

and

$$
\mathbf{h}_{k} = [\gamma \sum_{t=0}^{\tau-1} (\nabla G(\boldsymbol{\Phi}_{k}^{t}) - \nabla F(\bar{\mathbf{X}}_{k}) + \rho D \boldsymbol{\Phi}_{k}^{t} - \rho \mathbf{D} \mathbf{X}_{k});
$$

$$
\gamma \rho \tilde{\mathbf{L}} \sum_{t=0}^{\tau-1} (\nabla G(\boldsymbol{\Phi}_{k}^{t}) - \nabla F(\bar{\mathbf{X}}_{k}) + \rho \mathbf{D} \boldsymbol{\Phi}_{k}^{t} - \rho \mathbf{D} \mathbf{X}_{k})
$$

$$
+ \nabla F(\bar{\mathbf{X}}_{k+1}) - \nabla F(\bar{\mathbf{X}}_{k}); -\nabla F(\bar{\mathbf{X}}_{k+1}) + \nabla F(\bar{\mathbf{X}}_{k}).
$$

We remark that (14) can be interpreted as a linear dynamical system, with the non-linearity of the gradients as input in h_k .

C. Deviation from the average

The following lemma illustrates how far the states deviate from the average and will be used later in the proofs of Lemmas [2](#page-8-0) and [6.](#page-12-0)

Lemma [1](#page-2-6). Let Assumption 1 hold, when $\gamma < \frac{2}{\tau |\tilde{\lambda}_{\min}|\rho}$,

$$
\|\bar{\mathbf{X}}_k - \mathbf{X}_k\|^2 \le \frac{18\gamma\tau}{|\tilde{\lambda}_{\max}|\rho} \|\widehat{\mathbf{d}}_k\|^2, \quad \|\bar{\mathbf{Y}}_k - \mathbf{Y}_k\|^2 \le 9\|\widehat{\mathbf{d}}_k\|^2,
$$
\n(16)

and

$$
\|\hat{\mathbf{d}}_{k+1}\|^2 \le \delta \|\hat{\mathbf{d}}_k\|^2 + \frac{1}{1-\delta} \|\hat{\mathbf{h}}_k\|^2 \tag{17}
$$

where $\delta = 1 - |\tilde{\lambda}_{\text{max}}| \rho \tau \gamma / 2 < 1$.

Proof. By Assumption [1,](#page-2-6) graph G is undirected and connected, hence its Laplacian −L is symmetric; moreover, it has one zero eigenvalue with eigenvector 1, with all eigenvalues being positive. Denote by $\widehat{Q} \in \mathbb{R}^{N \times (N-1)}$ the matrix satisfying $\widehat{\mathbf{Q}}\widehat{\mathbf{Q}}^T = \mathbf{I}_N - \frac{1}{N}\mathbf{1}\mathbf{1}^T$, $\widehat{\mathbf{Q}}^T\widehat{\mathbf{Q}} = \mathbf{I}_{N-1}$ and $\mathbf{1}^T\widehat{\mathbf{Q}} = 0$, $\widehat{\mathbf{Q}}^T\mathbf{1} =$ 0. We have that

$$
\widehat{\mathbf{Q}}^T \widetilde{\mathbf{L}} = \widehat{\mathbf{Q}}^T \widetilde{\mathbf{L}} (\mathbf{I}_N - \frac{1}{N} \mathbf{1} \mathbf{1}^T) = \widehat{\mathbf{Q}}^T \widetilde{\mathbf{L}} \widehat{\mathbf{Q}} \widehat{\mathbf{Q}}^T.
$$
 (18)

Additionally, it holds that $\|\widehat{\mathbf{Q}}^T \mathbf{X}_k\|^2 = \mathbf{X}_k^T \widehat{\mathbf{Q}} \widehat{\mathbf{Q}}^T \widehat{\mathbf{Q}} \widehat{\mathbf{Q}}^T \mathbf{X}_k = \left\|\widehat{\mathbf{Q}} \widehat{\mathbf{Q}}^T \mathbf{X}_k\right\|^2 = \left\|\mathbf{X}_k - \bar{\mathbf{X}}_k\right\|^2$, and $\|\widehat{\mathbf{Q}}\| = 1$. Multiplying $\mathbf{Z}^2 \; = \; \left\| \mathbf{X}_k - \bar{\mathbf{X}}_k \right\|$ ², and $\|\hat{\mathbf{Q}}\| = 1$. Multiplying both sides of [\(14\)](#page-6-4) by $\hat{\mathbf{Q}}^T$ and using [\(18\)](#page-7-3) yields:

$$
\begin{bmatrix} \widehat{\mathbf{Q}}^T \mathbf{X}_{k+1} \\ \widehat{\mathbf{Q}}^T \mathbf{Y}_{k+1} \\ \widehat{\mathbf{Q}}^T \widehat{\mathbf{Y}}_{k+1} \end{bmatrix} = (\mathbf{\Theta} \otimes \mathbf{I}_n) \begin{bmatrix} \widehat{\mathbf{Q}}^T \mathbf{X}_k \\ \widehat{\mathbf{Q}}^T \mathbf{Y}_k \\ \widehat{\mathbf{Q}}^T \widehat{\mathbf{Y}}_k \end{bmatrix} - \widehat{\mathbf{Q}}^T \mathbf{h}_k \quad (19)
$$

where $\Theta =$ \lceil $\overline{1}$ ${\bf I} \hspace{1.5cm} \gamma \tau {\bf I} \hspace{1.5cm} {\bf 0}$ $\rho \widehat{\mathbf{Q}}^T \tilde{\mathbf{L}} \widehat{\mathbf{Q}} - \rho \widehat{\mathbf{Q}}^T \tilde{\mathbf{L}} \widehat{\mathbf{Q}} \gamma \tau \mathbf{I} + \frac{1}{2} \mathbf{I} \quad -\frac{1}{2} \mathbf{I}$ 0 $\frac{1}{2}$ **I** $\frac{1}{2}$ **I** 1 \cdot

The next step is to show that $\hat{Q}^T \tilde{L} \hat{Q}$ is negative definite by contradiction. Let $x \in R^{N-1}$ be an arbitrary vector, since $-\tilde{L}$ is the positive semi-definite Laplacian matrix, the quadratic form $x^T \widehat{Q}^T \widetilde{L} \widehat{Q} x = (\widehat{Q}_x)^T \widetilde{L} \widehat{Q} x \leq 0$. Moreover, if $(\widehat{\mathbf{Q}}x)^T (\widetilde{\mathbf{A}} - \mathbf{D})\widehat{\mathbf{Q}}x = 0$, we have $\widehat{\mathbf{Q}}x = 1$. Now, the properties of \dot{Q} imply that $\dot{Q}^T \dot{Q} x = x = \dot{Q}^T 1 = 0$. Therefore, for all non-zero vectors x, the quadratic form $x^T \hat{\mathbf{Q}}^T \tilde{\mathbf{L}} \hat{\mathbf{Q}} x < 0$, thus $\mathbf{Q}^T \mathbf{L} \mathbf{Q}$ is a symmetric negative-definite matrix.

We proceed now to diagonalize each block of Θ with $\phi \in$ $\mathbb{R}^{(N-1)\times(N-1)}$:

$$
\tilde{\Theta} = \phi \Theta \phi^T = \begin{bmatrix} \phi & 0 & 0 \\ 0 & \phi & 0 \\ 0 & 0 & \phi \end{bmatrix} \Theta \begin{bmatrix} \phi^T & 0 & 0 \\ 0 & \phi^T & 0 \\ 0 & 0 & \phi^T \end{bmatrix}
$$

$$
= \begin{bmatrix} \mathbf{I} & \gamma \tau & 0 \\ \rho \phi \hat{\mathbf{Q}}^T \tilde{\mathbf{L}} \hat{\mathbf{Q}} \phi^T & \rho \phi \hat{\mathbf{Q}}^T \tilde{\mathbf{L}} \hat{\mathbf{Q}} \phi^T \gamma \tau + \frac{1}{2} \mathbf{I} & -\frac{1}{2} \mathbf{I} \\ 0 & -\frac{1}{2} \mathbf{I} & \frac{1}{2} \mathbf{I} \end{bmatrix}.
$$

We denote $\phi \widehat{Q}^T \widehat{\mathbf{L}} \widehat{Q} \phi^T = \text{diag}\{\widetilde{\lambda}_i\}_{i=2,\dots,N}$, where $\widetilde{\lambda}_i < 0$ is the eigenvalue of $\hat{\mathbf{Q}}^T \tilde{\mathbf{L}} \hat{\mathbf{Q}}$, $\tilde{\lambda}_{\min} = \lambda_{\min}(\hat{\mathbf{Q}}^T \tilde{\mathbf{L}} \hat{\mathbf{Q}})$, and $\tilde{\lambda}_{\max} =$ $\lambda_{\text{max}}(\mathbf{Q}^T \mathbf{L} \mathbf{Q})$. Since each block of Θ is a diagonal matrix, there exists a permutation matrix P_0 such that $P_0 \tilde{\Theta} P_0^T =$ $\mathbf{P}_0 \boldsymbol{\phi} \boldsymbol{\Theta} \boldsymbol{\phi}^T \mathbf{P}_0^T \stackrel{\sim}{=}$ blkdiag $\{\mathbf{D}_i\}_{i=2}^N$, where

$$
\mathbf{D}_{i} = \begin{bmatrix} 1 & \gamma \tau & 0 \\ \rho \tilde{\lambda}_{i} & \rho \tilde{\lambda}_{i} \gamma \tau + 0.5 & -0.5 \\ 0 & -0.5 & 0.5 \end{bmatrix} . \tag{20}
$$

We diagonalize $D_i = V_i \Delta_i V_i^{-1}$, where Δ_i is the diagonal matrix of D_i 's eigenvalues, and

$$
\mathbf{V}_{i} = \begin{bmatrix} -\gamma \tau & d_{12} & d_{13} \\ 1 & d_{22} & d_{23} \\ 1 & 1 & 1 \end{bmatrix}
$$
 (21)

with $d_{12} = -\gamma \tau + ((\gamma \tilde{\lambda}_i \rho \tau (\gamma \tilde{\lambda}_i \rho \tau + 2))^{0.5})/(\tilde{\lambda}_i \rho), d_{13} =$ $-\gamma\tau - \left((\gamma\tilde{\lambda}_i\rho\tau(\gamma\tilde{\lambda}_i\rho\tau+2))^{0.5}\right)/(\tilde{\lambda}_i\rho), \ d_{22} = \tilde{\lambda_i}\rho d_{12} - 1,$ $d_{23} = \lambda_i \rho d_{13} - 1$. The nonzero eigenvalues λ of D_i , $i =$ 2, ..., N, satisfy $2\lambda^2 + (-2\tilde{\lambda}_i \rho \tau \gamma - 4)\lambda + \tilde{\lambda}_i \rho \tau \gamma + 2 = 0$, which can be written in the form:

$$
2\lambda^2 - 2t\lambda + t = 0\tag{22}
$$

where $t = \tilde{\lambda}_i \rho \tau \gamma + 2$. The modulus of the roots of [\(22\)](#page-7-4) is 1 – $\frac{|\tilde{\lambda}| \rho \tau \gamma}{2}$ when $-2 < \tilde{\lambda} \rho \tau \gamma < 0$. We conclude that we can write $\Theta = (\mathbf{P}_0 \phi)^T \mathbf{V} \Delta \mathbf{V}^{-1} (\mathbf{P}_0 \phi)$ where $\mathbf{V} = \text{blkdiag} \{V_i\}_{i=2}^N$ and

$$
\Delta = \text{blkdiag}\left\{\Delta_i\right\}_{i=2}^N. \tag{23}
$$

Moreover, $\|\Delta\| = 1 - |\tilde{\lambda}_{\text{max}}| \rho \tau \gamma/2$ when

$$
|\tilde{\lambda}_{\min}|\rho \tau \gamma < 2. \tag{24}
$$

Then, left multiplying both sides of [\(19\)](#page-7-5) by the inverse of

$$
\widehat{\mathbf{V}} = (P_0 \boldsymbol{\phi})^T \mathbf{V},\tag{25}
$$

which is given by $\hat{\mathbf{V}}^{-1} = \mathbf{V}^{-1}(\mathbf{P}_0 \phi)$, yields

$$
\widehat{\mathbf{d}}_{k+1} = \Delta \widehat{\mathbf{d}}_k - \widehat{\mathbf{h}}_k, \tag{26}
$$

where $\hat{\mathbf{d}}_k = \hat{\mathbf{V}}^{-1} \left[\hat{\mathbf{Q}}^T \mathbf{X}_k; \hat{\mathbf{Q}}^T \mathbf{Y}_k; \hat{\mathbf{Q}}^T \tilde{\mathbf{Y}}_k \right], \quad \hat{\mathbf{h}}_k =$ $\widehat{\mathbf{V}}^{-1}\widehat{\mathbf{Q}}^T\mathbf{h}_k$, and $\begin{bmatrix} \widehat{\mathbf{Q}}^T\mathbf{X}_k;\widehat{\mathbf{Q}}^T\mathbf{Y}_k;\widehat{\mathbf{Q}}^T\tilde{\mathbf{Y}}_k \end{bmatrix}$ = $\widehat{\mathbf{V}}\widehat{\mathbf{d}}_k$ = $\phi^T \mathbf{P}_0^T \mathbf{V} \hat{\mathbf{d}}_k = \phi^T \mathbf{P}_0^T \mathbf{V} \mathbf{P}_0 \mathbf{P}_0^T \hat{\mathbf{d}}_k$. As a consequence, from [\(19\)](#page-7-5) it holds that

$$
\begin{bmatrix}\n\widehat{\mathbf{Q}}^T \mathbf{X}_k \\
\widehat{\mathbf{Q}}^T \mathbf{Y}_k\n\end{bmatrix} = \boldsymbol{\phi}^T \begin{bmatrix}\n-\gamma \tau \mathbf{I} & d_{12} \mathbf{I} & d_{13} \mathbf{I} \\
\mathbf{I} & d_{22} \mathbf{I} & d_{23} \mathbf{I} \\
\mathbf{I} & \mathbf{I} & \mathbf{I}\n\end{bmatrix} \mathbf{P}_0^T \widehat{\mathbf{d}}_k
$$
\n
$$
= \boldsymbol{\phi}^T \begin{bmatrix}\n-\gamma \tau \mathbf{I} \mathbf{P}_0^T [1] + d_{12} \mathbf{P}_0^T [2] + d_{13} \mathbf{P}_0^T [3] \\
\mathbf{P}_0^T [1] + d_{22} \mathbf{P}_0^T [2] + d_{23} \mathbf{P}_0^T [3] \\
\mathbf{P}_0^T [1] + \mathbf{P}_0^T [2] + \mathbf{P}_0^T [3]\n\end{bmatrix} \widehat{\mathbf{d}}_k,
$$

where $\mathbf{P}_0^T[1], \mathbf{P}_0^T[2], \mathbf{P}_0^T[3]$ are the top, middle and bottom where \mathbf{r}_0 [1], \mathbf{r}_0 [2], \mathbf{r}_0 [5] are the top, findite and bottom
blocks of \mathbf{P}_0^T respectively. Moreover, we have $|d_{12}|^2$ = $|d_{13}|^2 \le \frac{2\gamma\tau}{|\bar{\lambda}_{\max}|\rho|}, \ |d_{22}| = |d_{23}| = 1.$ Now, if we let $\gamma \tau \leq \frac{2}{|\bar{\lambda}_{\max}|\rho|}$, and using $\|\phi\| = 1$, $\|\mathbf{P}_0^T[i]\| = 1$, $i = 1, 2, 3$, we derive that

$$
\begin{split} \|\bar{\mathbf{X}}_k - \mathbf{X}_k\|^2 &= \|\widehat{\mathbf{Q}}^T \mathbf{X}_k\|^2 \\ &= \|\phi^T(-\gamma \tau \mathbf{IP}_0^T[1] + d_{12} \mathbf{P}_0^T[2] + d_{13} \mathbf{P}_0^T[3])\widehat{\mathbf{d}}_k\|^2 \\ &\leq 3(\gamma^2 \tau^2 + \frac{4\gamma \tau}{|\widetilde{\lambda}_{\text{max}}|\rho}) \|\widehat{\mathbf{d}}_k\|^2 \leq \frac{18\gamma \tau}{|\widetilde{\lambda}_{\text{max}}|\rho} \|\widehat{\mathbf{d}}_k\|^2. \end{split}
$$

Applying the same manipulations to $\|\bar{\mathbf{Y}}_k - \mathbf{Y}_k\|^2$, we obtain [\(16\)](#page-7-6) holds. Denote now $\|\mathbf{\Phi}_k\|$ ² $=$ $\sum_{i=1}^{N} \sum_{t=0}^{\tau-1} \left\| \phi_{i,k}^{t} - \bar{x}_{k} \right\|$ 2 = $\sum_{t=0}^{\tau-1} ||\Phi_k^t - \bar{X}_k||$ 2 . Using Assumption [2](#page-2-7) we derive that

$$
\|\sum_{t=0}^{\tau-1} (G(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D} \Phi_k^t - \rho \mathbf{D} \mathbf{X}_k)\|^2
$$

\n
$$
= \|\sum_{t=0}^{\tau-1} (G(\Phi_k^t) - \nabla F(\Phi_k^t) + \nabla F(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k)
$$

\n
$$
+ \rho \mathbf{D}(\Phi_k^t - \bar{\mathbf{X}}_k + \bar{\mathbf{X}}_k - \mathbf{X}_k))\|^2
$$

\n
$$
\leq 4(\tau L^2 + \tau \rho^2 d_u^2) \|\hat{\Phi}_k\|^2 + 4\tau^2 \rho^2 d_u^2 \|\bar{\mathbf{X}}_k - \mathbf{X}_k\|^2
$$

\n
$$
+ 4\tau \sum_{t=0}^{\tau-1} \|G(\Phi_k^t) - \nabla F(\Phi_k^t)\|^2.
$$

Denoting $\epsilon_k = \sum_{t=0}^{\tau-1} (\rho \mathbf{1}^T \mathbf{D} \Phi_k^t - \mathbf{1}^T \mathbf{A}^T \mathbf{Z}_k) =$ $\rho \sum_{t=0}^{\tau-1} \sum_{i=1}^{N} (d_i \phi_{i,k}^t - \overline{d_i x_k} + d_i \overline{x}_k - d_i x_{i,k}),$ we have

$$
\epsilon_k^2 \le 2\rho^2 d_u^2 \tau N(\|\widehat{\mathbf{\Phi}}_k\|^2 + \tau \|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2). \tag{27}
$$

Now, denoting $\overline{G}(\Phi_k^t)$ = $\frac{1}{N} \sum_i g_i(\phi_{i,k}^t)$, $\begin{array}{l} \text{then } \quad \Vert \sum_t \overline{G}(\boldsymbol{\Phi}_k^t) \Vert^2 \quad \leq \quad \quad 2 \Vert \sum_t \overline{\nabla F}(\boldsymbol{\Phi}_k^t) \Vert^2 \quad + \\ \frac{2\tau}{N} \sum_t \sum_i \Vert \nabla f_i(\phi_{i,k}^t) - g_i(\phi_{i,k}^t) \Vert^2, \text{ where } \Vert \sum_t \overline{\nabla F}(\boldsymbol{\Phi}_k^t) \Vert^2 = \end{array}$ $\|\frac{1}{N}\sum_i\sum_t \nabla f_i\left(\phi_{i,k}^t\right)\|^2 \leq \frac{2\tau}{N}\sum_i\sum_t \|\nabla f_i(\phi_{i,k}^t)\|$ $\nabla f_i(\bar{x}_k) \|^2 + 2\tau^2 \|\frac{1}{N} \sum_i \nabla f_i(\bar{x}_k)\|^2$. We also have

$$
\|\nabla F(\bar{\mathbf{X}}_{k+1}) - \nabla F(\bar{\mathbf{X}}_k)\|^2 = N \|\nabla F(\bar{x}_{k+1}) - \nabla F(\bar{x}_k)\|^2
$$

\n
$$
\leq NL^2 \|\frac{\gamma}{N} \sum_{t=0}^{\tau-1} \sum_{i=1}^N g_i(\phi_k^{t,i}) + \frac{\gamma}{N} \sum_{t=0}^{\tau-1} (\rho \mathbf{1}^T \mathbf{D} \Phi_k^t - \mathbf{1}^T \mathbf{A}^T \mathbf{Z}_k)\|^2
$$

\n
$$
\leq 2L^2 \gamma^2 \left(2\tau \sum_t \sum_i \|\nabla f_i(\phi_{i,k}^t) - g_i(\phi_{i,k}^t)\|^2 + 2N \|\sum_t \overline{\nabla F}(\Phi_k^t)\|^2 + 2\rho^2 d_u^2 \tau (\|\widehat{\Phi}_k\|^2 + \tau \|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2) \right).
$$

Using $\|\widehat{\mathbf{Q}}^T \| = 1$ it further holds that:

$$
\begin{aligned}\n\|\widehat{\mathbf{h}}_k\|^2 &\leq \gamma^2 (1 + 2\rho^2 \|\widetilde{\mathbf{L}}\|^2) \|\widehat{\mathbf{V}}^{-1}\|^2. \\
&\cdot \|\sum_{t=0}^{\tau-1} (G(\Phi_k^t) - \nabla \mathbf{F}(\bar{\mathbf{X}}_k) + \rho \mathbf{D} \Phi_k^t - \rho \mathbf{D} \mathbf{X}_k)\|^2 \\
&\quad + 3\|\widehat{\mathbf{V}}^{-1}\|^2 \|\nabla F(\bar{\mathbf{X}}_{k+1}) - \nabla F(\bar{\mathbf{X}}_k)\|^2 \\
&\leq \gamma^2 (1 + 2\rho^2 \|\widetilde{\mathbf{L}}\|^2) \|\widehat{\mathbf{V}}^{-1}\|^2 \left(4(\tau L^2 + \tau \rho^2 d_u^2) \|\widehat{\mathbf{\Phi}}_k\|^2\right)\n\end{aligned}
$$

$$
+4\tau^{2}\rho^{2}d_{u}^{2}\left\|\bar{\mathbf{X}}_{k}-\mathbf{X}_{k}\right\|^{2}+4\tau\sum_{t=0}^{\tau-1}\|G(\Phi_{k}^{t})-\nabla F(\Phi_{k}^{t})\|^{2}\right) +6L^{2}\gamma^{2}\|\hat{\mathbf{V}}^{-1}\|^{2}\cdot\left(2\tau\sum_{t}\sum_{i}\|\nabla f_{i}\left(\phi_{i,k}^{t}\right)-g_{i}\left(\phi_{i,k}^{t}\right)\|^{2} +2N\|\sum_{t}\overline{\nabla F}\left(\Phi_{k}^{t}\right)\|^{2}+2\rho^{2}d_{u}^{2}\tau(\|\hat{\Phi}_{k}\|^{2}+\tau\|\mathbf{X}_{k}-\bar{\mathbf{X}}_{k}\|^{2})\right)
$$
(28)

Recalling [\(26\)](#page-7-0), using Jensen's inequality $||\mathbf{d}_{k+1}|| \frac{1}{||\mathbf{\Delta}||} ||\mathbf{\Delta}||^2 ||\mathbf{\hat{d}}_k||^2 + \frac{1}{1-||\mathbf{\Delta}||} ||\mathbf{\hat{h}}_k||^2$ yields [\(17\)](#page-7-7). $\frac{2}{\Box}$

APPENDIX B CONVERGENCE ANALYSIS FOR LT-ADMM

A. Key bounds

Lemma 2. *Let Assumptions [1,](#page-2-6) [2,](#page-2-7) and [4](#page-3-4) hold; when* $\gamma < \frac{2}{\tau |\tilde{\lambda}_{\text{max}}|\rho}$ *and* $4\gamma^2 \tau (L^2 + \rho^2 d_u^2) < \frac{1/4}{\tau - 1}$ *, we have* $rac{1/4}{7-1}$, we have $\mathbb{E}\left[\lVert \widehat{\mathbf{\Phi}}_k \rVert^2 \right]$ ≤ \int 72 $\gamma \tau^2$ $|\tilde{\lambda}_{\text{max}}| \rho$ + $(18 + \frac{36\gamma\tau\rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho})16\tau^3\gamma^2$ $\mathbb{E}\left[\|\hat{\mathbf{d}}_k\|^2\right]$ (29) + $16\tau^3\gamma^2 N\mathbb{E} \|\nabla F(\bar{x}_k)\|^2 + 4N\tau^2\gamma^2\sigma^2$.

Proof. From [\(12\)](#page-6-3) we can derive that

$$
\bar{x}_{k+1} - x^* = \bar{x}_k - x^* - \frac{\gamma}{N} \sum_{t=0}^{\tau-1} \sum_{i=1}^N g_i(\phi_{i,k}^t) - \frac{\gamma}{N} \left(\sum_{t=0}^{\tau-1} (\rho \mathbf{1}^T \mathbf{D} \Phi_k^t - \mathbf{1}^T \mathbf{A}^T \mathbf{Z}_k) \right)
$$
\n(30)

and

$$
\Phi_k^{t+1} = \Phi_k^t + \gamma \mathbf{Y}_k - \gamma (G(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D} \Phi_k^t - \rho \mathbf{D} \mathbf{X}_k)
$$
\n(31)

Recall that by Assumption [4,](#page-3-4) $||G(\Phi_k^t) - \nabla F(\Phi_k^t)||^2 \leq N\sigma^2$. Now, suppose that $\tau \geq 2$, using Jensen's inequality we obtain

$$
\mathbb{E}[\left\|\Phi_k^{t+1} - \bar{\mathbf{X}}_k\right\|^2]
$$
\n
$$
\geq \mathbb{E}[\left\|\Phi_k^t - \bar{\mathbf{X}}_k + \gamma \mathbf{Y}_k - \gamma(\nabla F(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D}\Phi_k^t - \rho \mathbf{D}\mathbf{X}_k)\right\|^2] + N\gamma^2 \sigma^2
$$
\n
$$
\leq (1 + \frac{1}{\tau - 1})\mathbb{E}[\left\|\Phi_k^t - \bar{\mathbf{X}}_k\right\|^2] + N\gamma^2 \sigma^2 + \sigma^2 + \tau\gamma^2 \mathbb{E}[\|\mathbf{Y}_k - (\nabla F(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D}\Phi_k^t - \rho \mathbf{D}\mathbf{X}_k)\right\|^2]
$$
\n
$$
\leq (1 + \frac{1}{\tau - 1})\mathbb{E}[\|\Phi_k^t - \bar{\mathbf{X}}_k\|^2] + N\gamma^2 \sigma^2 + \sigma^2 + \sigma^2 + \sigma^2 + \sigma^2 \mathbb{E}[\|\nabla F(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D}\Phi_k^t - \rho \mathbf{D}\bar{\mathbf{X}}_k\|^2] + \sigma^2 + 2\tau\gamma^2 \mathbb{E}[\|\mathbf{Y}_k - \bar{\mathbf{Y}}_k + \bar{\mathbf{Y}}_k + (\rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k)\|^2]
$$
\n
$$
\leq (1 + \frac{1}{\tau - 1} + 4\gamma^2 \tau (L^2 + \rho^2 d_u^2))\mathbb{E}[\|\Phi_k^t - \bar{\mathbf{X}}_k\|^2 + \sigma^2 + 2\tau\gamma^2 \|\mathbf{Y}_k - \bar{\mathbf{Y}}_k + \bar{\mathbf{Y}}_k + (\rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k)\|^2] + N\gamma^2 \sigma^2
$$
\n
$$
\leq (1 + \frac{5/4}{\tau - 1})\mathbb{E}[\|\Phi_k^t - \bar{\math
$$

.

where $\bar{\mathbf{Y}}_k = \mathbf{1} \otimes \frac{1}{N} \mathbf{1}^T \nabla F(\bar{\mathbf{X}}_k) = \mathbf{1} \otimes \frac{1}{N} \sum_i \nabla f_i(\bar{x}_k)$, and the last inequality holds when

$$
4\gamma^2 \tau (L^2 + \rho^2 d_u^2) < \frac{1/4}{\tau - 1}.\tag{32}
$$

Iterating the above inequality for $t = 0, ..., \tau - 1$

$$
\mathbb{E}[\|\Phi_k^{t+1} - \bar{\mathbf{X}}_k\|^2] \leq \left(1 + \frac{5/4}{\tau - 1}\right)^t \mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] +
$$

+ $2\tau\gamma^2 \sum_{l=0}^t \left(1 + \frac{5/4}{\tau - 1}\right)^l$

$$
\mathbb{E}[\|\mathbf{Y}_k - \bar{\mathbf{Y}}_k + \bar{\mathbf{Y}}_k + (\rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k)\|^2] +
$$

+ $N\gamma^2 \sigma^2 \sum_{l=0}^t \left(1 + \frac{5/4}{\tau - 1}\right)^l$

$$
\leq 4\mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] + 4\tau N\gamma^2 \sigma^2 +
$$

+ $8\tau^2 \gamma^2 \mathbb{E}[\|\mathbf{Y}_k - \bar{\mathbf{Y}}_k + \bar{\mathbf{Y}}_k + (\rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k)\|^2],$

where the last inequality holds by $(1 + \frac{a}{\tau-1})^t \leq \exp(\frac{at}{\tau-1}) \leq$ $\exp(a)$ for $t \leq \tau - 1$ and $a = 5/4$.

Summing over t , it follows that

$$
\mathbb{E}[\|\widehat{\Phi}_k\|^2] \le 4\tau \mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] + 16\tau^3 \gamma^2 N \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] + 4N\tau^2 \gamma^2 \sigma^2 + 16\tau^3 \gamma^2 \mathbb{E}[\|\mathbf{Y}_k - \bar{\mathbf{Y}}_k + (\rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k)\|^2];
$$
\n(33)

moreover, it is easy to verify that [\(33\)](#page-9-0) also holds for $\tau = 1$. As a consequence of [\(16\)](#page-7-6), we obtain

$$
\|\mathbf{Y}_{k} - \bar{\mathbf{Y}}_{k} + (\rho \mathbf{D}\bar{\mathbf{X}}_{k} - \rho \mathbf{D}\mathbf{X}_{k})\|^{2} \leq (18 + \frac{36\gamma\tau\rho^{2}d_{u}^{2}}{|\tilde{\lambda}_{\max}|\rho)}\|\hat{\mathbf{d}}_{k}\|^{2},
$$
\n(34)

and using this fact into [\(33\)](#page-9-0) concludes the proof.

Lemma 3. Let Assumptions [1,](#page-2-6) [2,](#page-2-7) and [4](#page-3-4) hold; when $\gamma \leq$ $\min\{\bar{\gamma}_1, \bar{\gamma}_2\}$ *, it holds that*

$$
\mathbb{E}[\|\hat{\mathbf{d}}_{k+1}\|^2] \leq (\delta + \frac{q_1}{1-\delta})\mathbb{E}[\|\hat{\mathbf{d}}_k\|^2] + + \frac{q_2}{1-\delta}\mathbb{E}[\|\sum_t \overline{\nabla F}(\mathbf{\Phi}_k^t)\|^2] + \frac{q_3}{1-\delta}\mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] + q_4\sigma^2
$$
\n(35)

where

$$
\delta := 1 - \frac{|\tilde{\lambda}_{\text{max}}| \rho \tau \gamma}{2}, q_4 := \frac{4N\tau^2 \gamma^2 c_0 + c_3}{1 - \delta}
$$

\n
$$
c_0 := \gamma^2 \tau \beta_0, c_1 := \gamma^3 \tau^2 \beta_1, c_2 := 12L^2 \gamma^2 ||\hat{\mathbf{V}}^{-1}||^2 N,
$$

\n
$$
c_3 := 4\gamma^2 (1 + 2\rho^2 ||\tilde{\mathbf{L}}||^2) ||\hat{\mathbf{V}}^{-1}||^2 \tau^2 N + 12L^2 \gamma^2 ||\hat{\mathbf{V}}^{-1}||^2 \tau^2 N,
$$

\n
$$
q_1 := c_0 \kappa_1 \gamma \tau + c_1, q_2 := c_2, q_3 := 16\tau^3 \gamma^2 N c_0.
$$

Proof. When $\gamma \leq \min{\bar{\gamma}_1, \bar{\gamma}_2}$, it follows that

$$
\gamma \tau \le \frac{2}{|\tilde{\lambda}_{\text{max}}|\rho}, \quad |\tilde{\lambda}_{\text{min}}|\rho \tau \gamma < 2, \quad \gamma^2 \tau (L^2 + \rho^2 d_u^2) < \frac{1/16}{\tau - 1}.
$$
\n(36)

Using [\(28\)](#page-8-1) and Assumption [4,](#page-3-4) we have

$$
\|\widehat{\mathbf{h}}_k\|^2 \le c_0 \|\widehat{\mathbf{\Phi}}_k\|^2 + c_1 \|\widehat{\mathbf{d}}_k\|^2 + c_2 \|\sum_t \overline{\nabla F}(\mathbf{\Phi}_k^t)\|^2 + c_3 \sigma^2;
$$

using this fact together with [\(29\)](#page-8-2) and [\(17\)](#page-7-7), and by γ < 1, we can then derive that [\(35\)](#page-9-1) holds. П

B. Theorem [1](#page-3-5)

 \Box

We start our proof by recalling that the following inequality holds for and L-smooth function f , $\forall y, z \in \mathbb{R}^n$ [\[36\]](#page-15-0):

$$
f(y) \le f(z) + \langle \nabla f(z), y - z \rangle + (L/2) \|y - z\|^2 \qquad (37)
$$

Based on [\(30\)](#page-8-3), substituting $y = \bar{x}_{k+1}$ and $z = \bar{x}_k$ into [\(37\)](#page-9-2), using Assumption [4](#page-3-4) and [\(27\)](#page-8-4), we get

$$
\mathbb{E}[F(\bar{x}_{k+1})]
$$
\n
$$
\leq \mathbb{E}[F(\bar{x}_k)] - \gamma \mathbb{E}[\langle \nabla F(\bar{x}_k), \frac{1}{N} (\sum_t \sum_i \nabla f_i (\phi_{i,k}^t) + \epsilon_k) \rangle]
$$
\n
$$
+ \frac{\gamma^2 L}{2} \mathbb{E}[\|\frac{1}{N} (\sum_t \sum_i g_i(\phi_{i,k}^t) + \epsilon_k)\|^2]
$$
\n
$$
\leq \mathbb{E}[F(\bar{x}_k)] - \gamma [\langle \nabla F(\bar{x}_k), \frac{1}{N} (\sum_t \sum_i \nabla f_i(\phi_{i,k}^t) + \epsilon_k) \rangle]
$$
\n
$$
+ 2\gamma^2 \tau L \mathbb{E}[\sum_t \|\frac{1}{N} \sum_i \nabla f_i (\phi_{i,k}^t) \|^2] + 2\gamma^2 \tau^2 L \sigma^2
$$
\n
$$
+ \frac{2\gamma^2 L \rho^2 d_u^2 \tau \mathbb{E}[\|\widehat{\Phi}_k\|^2 + \tau \|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2]}{N}.
$$

Using now $2\langle a, b \rangle = ||a||^2 + ||b||^2 - ||a - b||^2$, we have $- \langle \nabla F\left(\bar{x}_k\right), \frac{1}{N}\rangle$ N \sum t \sum $\sum_i \nabla f_i \left(\phi_{i,k}^t \right) + \frac{1}{N}$ $\frac{1}{N} \epsilon_k$ $\leq -\sum_t \langle \nabla F\left(\bar{x}_k\right), \frac{1}{N} \rangle$ N \sum $\sum_i \nabla f_i(\phi_{i,k}^t)\rangle + \frac{\tau}{4}$ $\frac{4}{4} \left\| \nabla F\left(\bar{x}_k\right) \right\|^2$ $+\frac{1}{\lambda^{10}}$ $\frac{1}{N^2\tau} \epsilon_k^2$ $=-\frac{\tau}{2}$ $\frac{\tau}{2} \left\| \nabla F\left(\bar{x}_k\right) \right\|^2 - \frac{1}{2}$ 2 \sum $\frac{1}{t}$ ∥ 1 N \sum $\sum\limits_i \nabla f_i\left(\phi_{i,k}^t\right) \|^2$ $+\frac{1}{2}$ 2 \sum $\frac{1}{t}$ ∥ 1 N \sum $\sum\limits_i \nabla f_i\left(\phi_{i,k}^t\right) - \nabla F\left(\bar{x}_k\right) \|^2$ $+\frac{7}{4}$ $\frac{\tau}{4} \left\| \nabla F\left(\bar{x}_k \right) \right\|^2 + \frac{1}{N^2}$ $\frac{1}{N^2\tau}\epsilon_k^2$ \leq $$ τ $\frac{\tau}{4} \left\| \nabla F\left(\bar{x}_k\right) \right\|^2 - \frac{1}{2}$ 2 \sum $\frac{}{t}$ ∥ 1 N \sum $\sum\limits_i \nabla f_i\left(\phi_{i,k}^t\right) \|^2$ $+\big(\frac{L^2}{2}$ $rac{L^2}{2N} + \frac{2\rho^2 d_u^2}{N}$ $\frac{\partial^2 d_u^2}{N}$) $\|\widehat{\mathbf{\Phi}}_k\|^2 + \frac{2\rho^2 d_u^2 \tau}{N}$ $\frac{a_{u}\tau}{N}\|\mathbf{X}_k-\bar{\mathbf{X}}_k\|^2$

Now, combining the two equations above, and using [\(16\)](#page-7-6), yields

$$
\mathbb{E}[F(\bar{x}_{k+1})] \leq \mathbb{E}[F(\bar{x}_k)] - \frac{\gamma \tau}{4} \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] \n- \frac{\gamma}{2} (1 - 4\gamma \tau L) \sum_t \mathbb{E}[\|\frac{1}{N} \sum_i \nabla f_i(\phi_{i,k}^t)\|^2] \n+ \left(\frac{\gamma L^2}{2N} + \frac{2\gamma^2 L \rho^2 d_u^2 \tau + 2\gamma \rho^2 d_u^2}{N}\right) \mathbb{E}[\|\hat{\Phi}_k\|^2] \n+ \frac{2\gamma \tau}{|\tilde{\lambda}_{\text{max}}| \rho} \frac{18\gamma \rho^2 d_u^2 \tau + 18\gamma^2 \rho^2 d_u^2 \tau^2 L}{N} \mathbb{E}[\|\hat{\mathbf{d}}_k\|^2] + 2\gamma^2 \tau^2 L \sigma^2.
$$

Substituting [\(29\)](#page-8-2) into the above inequality yields

$$
\mathbb{E}[F(\bar{x}_{k+1})] \leq \mathbb{E}[F(\bar{x}_k)] +
$$

$$
-\frac{\gamma \tau}{4} \left(1 - 64\gamma \tau^2 (\frac{\gamma L^2}{2} + 2\gamma^2 L\rho^2 d_u^2 \tau + 2\gamma \rho^2 d_u^2)\right).
$$

$$
\begin{aligned} &\cdot\mathbb{E}[\|\nabla F\left(\bar{x}_k\right)\|^2]\\&-\frac{\gamma}{2}(1-4\gamma L\tau)\sum_{t}\mathbb{E}[\|\frac{1}{N}\sum_{i}\nabla f_i\left(\phi_{i,k}^t\right)\|^2]\\&+\frac{1}{N}\left((\frac{72\gamma\tau^2}{|\tilde{\lambda}_{\rm max}|\rho}+(18+\frac{36\gamma\tau\rho^2d_u^2}{|\tilde{\lambda}_{\rm max}|\rho})16\tau^3\gamma^2)\cdot\right.\\&\left. \cdot(\frac{\gamma L^2}{2}+2\gamma^2L\rho^2d_u^2\tau+2\gamma\rho^2d_u^2\right)\\&+\frac{2\gamma\tau}{|\tilde{\lambda}_{\rm max}|\rho}(18\gamma\rho^2d_u^2\tau+18\gamma^2\rho^2d_u^2\tau^2L)\right)\mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2]\\&+2\gamma^2\tau^2L\sigma^2+\left(\frac{\gamma L^2}{2}+2\gamma^2L\rho^2d_u^2\tau+2\gamma\rho^2d_u^2\right)4\tau^2\gamma^2\sigma^2\end{aligned}
$$

When $\gamma \leq \min\{\bar{\gamma}_3, \bar{\gamma}_4\}$, then

$$
\gamma \tau^2 \left(\frac{\gamma L^2}{2} + 2\gamma^2 L \rho^2 d_u^2 \tau + 2\gamma \rho^2 d_u^2\right) < \frac{1}{128}, \quad \gamma L \tau < \frac{3}{16},
$$
\n(38)

and we can upper bound the previous inequality by

$$
\mathbb{E}[F(\bar{x}_{k+1})] \leq \mathbb{E}[F(\bar{x}_k)] - \frac{\gamma \tau}{8} \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] +
$$

\n
$$
- \frac{\gamma}{8} \sum_{t} \mathbb{E}[\|\frac{1}{N} \sum_{i} \nabla f_i(\phi_{i,k}^t)\|^2]
$$

\n
$$
+ \frac{1}{N} \left((\frac{72\gamma \tau^2}{|\tilde{\lambda}_{\text{max}}|\rho} + (18 + \frac{36\gamma \tau \rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho}) 16\tau^3 \gamma^2) \cdot \right.
$$

\n
$$
\cdot (\frac{\gamma L^2}{2} + 2\gamma^2 L\rho^2 d_u^2 \tau + 2\gamma \rho^2 d_u^2)
$$

\n
$$
+ \frac{2\gamma \tau}{|\tilde{\lambda}_{\text{max}}|\rho} (18\gamma \rho^2 d_u^2 \tau + 18\gamma^2 \rho^2 d_u^2 \tau^2 L) \right) \mathbb{E}[\|\hat{\mathbf{d}}_k\|^2]
$$

\n
$$
+ 2\gamma^2 \tau^2 L\sigma^2 + \left(\frac{\gamma L^2}{2} + 2\gamma^2 L\rho^2 d_u^2 \tau + 2\gamma \rho^2 d_u^2\right) 4\tau^2 \gamma^2 \sigma^2
$$

Rearranging the above relation we get

$$
\mathcal{D}_{k} \leq \frac{8}{\gamma\tau}\mathbb{E}\left[\left(\tilde{F}\left(\bar{x}_{k}\right)-\tilde{F}\left(\bar{x}_{k+1}\right)\right)\right] \n+ \frac{8}{N\gamma\tau}\left(\left(\frac{72\gamma\tau^{2}}{|\tilde{\lambda}_{\max}|\rho}+(18+\frac{36\gamma\tau\rho^{2}d_{u}^{2}}{|\tilde{\lambda}_{\max}|\rho})16\tau^{3}\gamma^{2}\right)\right) \n\cdot\left(\frac{\gamma L^{2}}{2}+2\gamma^{2}L\rho^{2}d_{u}^{2}\tau+2\gamma\rho^{2}d_{u}^{2}\right) \n+ \frac{2\gamma\tau}{|\tilde{\lambda}_{\max}|\rho}(18\gamma\rho^{2}d_{u}^{2}\tau+18\gamma^{2}\rho^{2}d_{u}^{2}\tau^{2}L)\right) \n\cdot\mathbb{E}[\|\hat{\mathbf{d}}_{k}\|^{2}]+16\left(\gamma L^{2}+4\gamma^{2}L\rho^{2}d_{u}^{2}\tau+4\gamma\rho^{2}d_{u}^{2}\right)\tau\gamma\sigma^{2} \n+16\gamma\tau L\sigma^{2},
$$

where \mathcal{D}_k is defined in [\(7\)](#page-3-7), and $\tilde{F}(\bar{x}_k) = F(\bar{x}_k) - F(x^*)$. Summing over $k = 0, 1, \ldots, K - 1$, using $-\tilde{F}(\bar{x}_k) \leq 0$ and $\gamma \leq 1$, it holds that

$$
\sum_{k=0}^{K-1} \mathcal{D}_k \le \frac{8\tilde{F}\left(\bar{x}^0\right)}{\gamma \tau} + c_4 \sum_{k=0}^{K-1} \mathbb{E}[\|\hat{\mathbf{d}}_k\|^2] + K c_5 \sigma^2 \qquad (39)
$$

where c_4 := $\gamma \kappa_4$, c_5 := $16\gamma \tau L$ + $16\left(\gamma L^2 + 4\gamma^2 L\rho^2 d_u^2 \tau + 4\gamma \rho^2 d_u^2\right) \tau \gamma.$

We now bound the term $\sum_{k=0}^{K-1} ||\hat{\mathbf{d}}_k||^2$. From [\(35\)](#page-9-1), we have $\mathbb{E}[\|\widehat{\mathbf{d}}_{k+1}\|^2]$

$$
\leq (\delta + \frac{q_1}{1-\delta}) \mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] + \frac{q_2}{1-\delta} \mathbb{E}[\|\sum_t \overline{\nabla F}(\boldsymbol{\Phi}_k^t)\|^2] \tag{40}
$$

$$
+\frac{q_3}{1-\delta}\mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] + q_4\sigma^2 \le \bar{\delta}\mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] + q_4\sigma^2 + K\mathcal{D}_k
$$
\n(41)

where

.

.

$$
R := \max\{\frac{q_3}{1-\delta}, \frac{q_2\tau^2}{1-\delta}\}.
$$
 (42)

Moreover, letting $\gamma \leq \min\{\bar{\gamma}_1, \bar{\gamma}_5\}$, we have

$$
\bar{\delta} = \delta + \frac{q_1}{1 - \delta} < 1. \tag{43}
$$

Iterating [\(41\)](#page-10-0) now gives

$$
\mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] \le \bar{\delta}_k \mathbb{E}[\|\widehat{\mathbf{d}}_0\|^2] + R \sum_{\ell=0}^{k-1} \bar{\delta}^{k-1-\ell} \mathcal{D}_\ell + \frac{q_4 \sigma^2}{1-\bar{\delta}}
$$

and summing this inequality over $k = 0, \ldots, K-1$, it follows that

$$
\sum_{k=0}^{K-1} \mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] \le \frac{\|\widehat{\mathbf{d}}_0\|^2}{1-\bar{\delta}} + \frac{R}{1-\bar{\delta}} \sum_{k=0}^{K-1} \mathcal{D}_k + \frac{q_4 \sigma^2 K}{1-\bar{\delta}}. \quad (44)
$$

Substituting [\(44\)](#page-10-1) into [\(39\)](#page-10-2) and rearranging, we obtain

$$
(1 - q_5) \sum_{k=0}^{K-1} \mathcal{D}_k \le \frac{8\tilde{F}(\bar{x}^0)}{\gamma \tau} + q_6 \|\hat{\mathbf{d}}_0\|^2 + K q_7 \sigma^2,
$$

where

$$
q_5 := \frac{c_4 R}{1 - \bar{\delta}}, \quad q_6 := \frac{c_4}{1 - \bar{\delta}} \quad q_7 := \frac{c_4 q_4}{1 - \bar{\delta}} + c_5. \tag{45}
$$

When $\gamma \leq \frac{|\tilde{\lambda}_{\text{max}}|^2 \rho^2 \tau^2}{8(\beta_0 \kappa_1 + \beta_1)}$ $\frac{|\tilde{\lambda}_{\max}|^2 \rho^2 \tau^2}{8(\beta_0 \kappa_1 + \beta_1)},$ we have $1 - \bar{\delta} \ge \frac{|\tilde{\lambda}_{\max}| \rho \tau \gamma}{4},$ and $\frac{c_4}{1-\delta} \leq \frac{4\kappa_4}{|\bar{\lambda}_{\text{max}}|\rho\tau|}$. Using this fact together with [\(42\)](#page-10-3) and $\gamma \leq 1$, we have $q_5 \leq \max\left\{\frac{128\kappa_4\tau^2 N \beta_0 \gamma}{|\tilde{\lambda}_{\max}|^2 \rho^2 \tau}\right\}$ $\frac{28 \kappa_4 \tau^2 N \beta_0 \gamma}{|\tilde{\lambda}_{\text{max}}|^2 \rho^2 \tau}, \frac{24 L^2 \|\hat{\mathbf{V}}^{-1}\|^2 N \gamma}{|\tilde{\lambda}_{\text{max}}|^2 \rho^2 \tau^2}$ $\frac{L^2 ||\mathbf{V} - \mathbf{V}||^2 N \gamma}{|\tilde{\lambda}_{\text{max}}|^2 \rho^2 \tau^2}$. Therefore, when $\gamma \le \min\{1, \bar{\gamma}_6, \bar{\gamma}_7\}$, we have

$$
q_5 \le \frac{1}{2},\tag{46}
$$

and it follows that

$$
\frac{1}{K} \sum_{k=0}^{K-1} \mathcal{D}_k \le \frac{16\tilde{F}\left(\bar{x}^0\right)}{\gamma \tau K} + \frac{2q_6}{K} \|\hat{\mathbf{d}}_0\|^2 + 2q_7\sigma^2. \tag{47}
$$

By collecting all step-size conditions, if the step-size γ < $\bar{\gamma}_{sgd} := \min_{i=1,2,...,7} \bar{\gamma}_i$, then [\(36\)](#page-9-3), [\(38\)](#page-10-4), [\(43\)](#page-10-5) and [\(46\)](#page-10-6) hold, the states $\{X_k\}$ generated by LT-ADMM converge to the neighborhood of the stationary point, concluding the proof.

APPENDIX C CONVERGENCE ANALYSIS FOR LT-ADMM-VR

A. Key bounds

We start by deriving an upper bound for the variance of the gradient estimator $\mathbb{E}\left[\|g_i\left(\phi_{i,k}^t\right)-\nabla f_i\left(\phi_{i,k}^t\right)\|^2\right]$. Define t_i^k as the averaged consensus gap of the auxiliary variables of ${r_{i,h,k}^k}_{h=1}^m$ at node *i*:

$$
\begin{aligned} & t_{i,k}^t = \frac{1}{m_i} \sum_{h=1}^{m_i} \|r_{i,h,k}^t - \bar{x}_k\|^2, \\ & t_k^t = \sum_{i=1}^N t_{i,k}^t = \frac{1}{m_i} \sum_{h=1}^{m_i} \|\mathbf{r}_{h,k}^t - \bar{\mathbf{X}}_k\|^2, \end{aligned}
$$

$$
t_k = \sum_{t=0}^{\tau-1} t_k^t = \sum_{t=0}^{\tau-1} \sum_{i=1}^N t_{i,k}^t.
$$

By the updates of $g_i(\phi_{i,k}^t)$ in LT-ADMM-VR,

$$
\mathbb{E} [\|g_i(\phi_{i,k}^t) - \nabla f_i (\phi_{i,k}^t) \|^2]
$$
\n
$$
= \mathbb{E}[\| \frac{1}{|\mathcal{B}_i|} \sum_{h \in \mathcal{B}_i} \nabla f_{i,h} (\phi_{i,k}^t) - \frac{1}{|\mathcal{B}_i|} \sum_{h \in \mathcal{B}_i} \nabla f_{i,h} (r_{i,h,k}^t)
$$
\n
$$
- (\nabla f_i (\phi_{i,k}^t) - \frac{1}{m_i} \sum_{h=1}^{m_j} \nabla f_{i,h} (r_{i,h,k}^k)) \|^2]
$$
\n
$$
\leq \mathbb{E} \left[\|\frac{1}{|\mathcal{B}_i|} \sum_{h \in \mathcal{B}_i} \nabla f_{i,h} (\phi_{i,k}^t) - \frac{1}{|\mathcal{B}_i|} \sum_{h \in \mathcal{B}_i} \nabla f_{i,h} (r_{i,h,k}^t) \|^2 \right]
$$
\n
$$
\leq \mathbb{E} \left[\frac{1}{|\mathcal{B}_i|} \sum_{h \in \mathcal{B}_i} \|\nabla f_{i,h} (\phi_{i,k}^t) - \nabla f_{i,h} (r_{i,h,k}^t) \|^2 \right]
$$
\n
$$
= \frac{1}{|\mathcal{B}_i|} \sum_{h \in \mathcal{B}_i} \mathbb{E} \left[\|\nabla f_{i,h} (\phi_{i,k}^t) - \nabla f_{i,h} (r_{i,h,k}^t) \|^2 \right]
$$
\n
$$
\leq 2L^2 \|\phi_{i,k}^t - \overline{x}_k\|^2 + 2L^2 \mathbb{E}[t_{i,k}^t],
$$

where in the first inequality we use $\mathbb{E}[\|a - \mathbb{E}[a]\|^2] \leq \mathbb{E}[\|a\|^2]$ with $a = \nabla f_{i,h}(\phi_{i,k}^t) - \nabla f_{i,h}(r_{i,h,k}^t)$; and in the second inequality we use the smoothness of the costs. As a consequence, we have

$$
\mathbb{E}[\sum_{i}\sum_{t}||g_i(\phi_{i,k}^t) - \nabla f_i(\phi_{i,k}^t)||^2] \le 2L^2 \|\widehat{\Phi}_k\|^2 + 2L^2 \mathbb{E}[t_k].
$$
\n(48)

Lemma [4](#page-3-4). *Let Assumptions [1,](#page-2-6)* [2](#page-2-7) *and* 4 *hold*; when $\gamma \tau \leq \frac{2}{|\bar{\lambda}_{\max}| \rho}$ *and* $6\gamma^2 \tau (3L^2 + \rho^2 d_u^2) < \frac{1/4}{\tau - 1}$ *, we have* $rac{1/4}{7-1}$, we have $\mathbb{E}[\|\widehat{\mathbf{\Phi}}_k\|^2] \leq \left(\frac{72\gamma\tau^2}{|\widetilde{\lambda}_{\text{max}}|} \right)$ $|\tilde{\lambda}_{\text{max}}| \rho$ $+ (18 + \frac{36\gamma\tau\rho^2d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho})16\tau^3\gamma^2\bigg)\,\mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2]$ + $16\tau^3\gamma^2 N \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] + 48\tau^2\gamma^2 L^2 \mathbb{E}[t_k]$ (49)

Proof. Suppose that $\tau \geq 2$, using [\(31\)](#page-8-5) and [\(48\)](#page-11-0) we have

$$
\mathbb{E}[\|\Phi_k^{t+1} - \bar{\mathbf{X}}_k\|^2] \n= \mathbb{E}[\|\Phi_k^t - \bar{\mathbf{X}}_k + \gamma \mathbf{Y}_k - \gamma(G(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D}\Phi_k^t - \rho \mathbf{D}\mathbf{X}_k)\|^2] \n\leq \left(1 + \frac{1}{\tau - 1}\right) \mathbb{E}[\|\Phi_k^t - \bar{\mathbf{X}}_k\|^2] + \frac{\tau \gamma^2 \mathbb{E}[\|Y_k - (G(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D}\Phi_k^t - \rho \mathbf{D}\mathbf{X}_k)\|^2] \n\leq \left(1 + \frac{1}{\tau - 1}\right) \mathbb{E}[\|\Phi_k^t - \bar{\mathbf{X}}_k\|^2] + \frac{2\tau \gamma^2 \mathbb{E}[\|(G(\Phi_k^t) - \nabla F(\Phi_k^t) + \nabla F(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D}\Phi_k^t - \rho D\bar{\mathbf{X}}_k\|^2] + \frac{2\tau \gamma^2 \mathbb{E}[\|\mathbf{Y}_k + \rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k\|^2] \n\leq \left(1 + \frac{1}{\tau - 1} + 6\gamma^2 \tau (2L^2 + L^2 + \rho^2 d_u^2)\right) \mathbb{E}[\|\Phi_k^t - \bar{\mathbf{X}}_k\|^2] + 2\tau \gamma^2 \mathbb{E}[\|\mathbf{Y}_k + \rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k\|^2] + 6\tau \gamma^2 (2L^2 \mathbb{E}[t_k^t]) \n\leq \left(1 + \frac{5/4}{\tau - 1}\right) \mathbb{E}[\|\Phi_k^t - \bar{\mathbf{X}}_k\|^2] + 12\tau \gamma^2 L^2 \mathbb{E}[t_k^t] + \frac{1}{\tau - 1} \mathbb{E}[\|\Phi_k^t - \bar{\mathbf{X}}_k\|^2] + 12\tau \gamma^2
$$

$$
+ 2\tau \gamma^2 \mathbb{E}[\|\mathbf{Y}_k + \rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k\|^2],\tag{50}
$$

where the last inequality holds when

$$
6\gamma^2 \tau (3L^2 + \rho^2 d_u^2) \le \frac{1/4}{\tau - 1}.
$$
 (51)

Iterating [\(50\)](#page-11-1) for $t = 0, ..., \tau - 1$ yields

$$
\mathbb{E}[\|\Phi_k^{t+1} - \bar{\mathbf{X}}_k\|^2] \leq \left(1 + \frac{5/4}{\tau - 1}\right)^t \mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] + + 2\tau\gamma^2 \sum_{l=0}^t \left(1 + \frac{5/4}{\tau - 1}\right)^l \mathbb{E}[\|\mathbf{Y}_k + \rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k\|^2] + + \sum_{l=0}^t \left(1 + \frac{5/4}{\tau - 1}\right)^{t-l} 12\tau\gamma^2 L^2 \mathbb{E}[t_k^l] \leq 4\mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] + 8\tau^2\gamma^2 \mathbb{E}[\|\mathbf{Y}_k + (\rho \mathbf{D}\bar{\mathbf{X}}_k - \rho \mathbf{D}\mathbf{X}_k)\|^2] + 48\tau^2\gamma^2 L^2 \mathbb{E}[t_k^t],
$$
\n(52)

where in the last inequality we used $(1+\frac{a}{\tau-1})^t \leq \exp(\frac{at}{\tau-1}) \leq$ $\exp(a)$ for $t \leq \tau - 1$. It is easy to verify that [\(52\)](#page-11-2) also holds for $\tau = 1$.

Summing (52) over t, and using (34) , it follows that

$$
\mathbb{E}[\|\hat{\Phi}_k\|^2] \leq 4\tau \mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] + 8\tau^3\gamma^2 \mathbb{E}[\|\mathbf{Y}_k - \bar{\mathbf{Y}}_k + \bar{\mathbf{Y}}_k + (\rho \mathbf{D} \bar{\mathbf{X}}_k - \rho \mathbf{D} \mathbf{X}_k)\|^2] + 48\tau^2\gamma^2 L^2 \mathbb{E}[t_k] \n\leq 4\tau \mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] + 16\tau^3\gamma^2 \mathbb{E}[\|\bar{\mathbf{Y}}_k\|^2] \n+ 16\tau^3\gamma^2 \mathbb{E}[\|\mathbf{Y}_k - \bar{\mathbf{Y}}_k + (\rho D \bar{\mathbf{X}}_k - \rho D \mathbf{X}_k)\|^2] \n+ 48\tau^2\gamma^2 L^2 \mathbb{E}[t_k] \n\leq 4\tau \mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] + 16\tau^3\gamma^2 N \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] \n+ 16\tau^3\gamma^2 \mathbb{E}[\|\mathbf{Y}_k - \bar{\mathbf{Y}}_k + (\rho D \bar{\mathbf{X}}_k - \rho D \mathbf{X}_k)\|^2] \n+ 48\tau^2\gamma^2 L^2 \mathbb{E}[t_k] \n\leq \left(\frac{72\gamma\tau^2}{|\bar{\lambda}_{\text{max}}|\rho} + (18 + \frac{36\gamma\tau\rho^2 d_w^2}{|\bar{\lambda}_{\text{max}}|\rho})16\tau^3\gamma^2\right) \mathbb{E}[\|\hat{\mathbf{d}}_k\|^2] \n+ 16\tau^3\gamma^2 N \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] + 48\tau^2\gamma^2 L^2 \mathbb{E}[t_k],
$$
\nwhich concludes the proof.

which concludes the proof.

The following lemma provides the bound on t_k .

Lemma 5. Let $\{t_k\}$ be the iterates generated by LT-ADMM-*VR. If* $\gamma \leq \min\{\bar{\gamma}_8, \bar{\gamma}_9, \bar{\gamma}_{10}\}$ *, we have for all* $k \in \mathbb{N}$ *:*

$$
\mathbb{E}[t_k] \le 2(s_0 + s_1)\mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] + 2s_2 \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2], \quad (53)
$$

where

$$
s_0 := \frac{36m_u\gamma\tau}{|\tilde{\lambda}_{\text{max}}|\rho} + \frac{4\tau}{m_l} \left(18 + \frac{36\gamma\tau\rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho}\right) (\gamma^2 + \gamma) 2m_u
$$

\n
$$
s_1 := \left(\frac{72\gamma\tau^2}{|\tilde{\lambda}_{\text{max}}|\rho} + (18 + \frac{36\gamma\tau\rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}|\rho}) 16\tau^3 \gamma^2\right) \frac{4m_u}{m_l} \quad (54)
$$

\n
$$
s_2 := \frac{64m_u}{m_l} \tau^3 \gamma^2 N + \frac{8m_u}{m_l} (\gamma^2 + \gamma) \tau N.
$$

Proof. From Algorithm [1,](#page-3-1) $\forall k, r_{i,h,k}^{t+1} = r_{i,h,k}^k$ with probability $\frac{1}{k!}$ and $r_{i,h,k}^{t+1} = r_{i,h,k}^{t+1}$ with probability $\frac{1}{k!}$ Therefore, we $1 - \frac{1}{m_i}$ and $r_{i,h,k}^{t+1} = \phi_{i,k}^{t+1}$ with probability $\frac{1}{m_i}$. Therefore, we have

$$
\mathbb{E}\left[t_{k}^{t+1}\right]=\frac{1}{m_{i}}\sum_{h=1}^{m_{i}}\mathbb{E}\left[\left\Vert \mathbf{r}_{h,k}^{t+1}-\bar{\mathbf{X}}_{k}\right\Vert ^{2}\right]
$$

$$
= \frac{1}{m_i} \sum_{h=1}^{m_i} \mathbb{E}[(1 - \frac{1}{m_i}) ||\mathbf{r}_{h,k}^t - \bar{\mathbf{X}}_k||^2 + \frac{1}{m_i} ||\Phi_k^{t+1} - \bar{\mathbf{X}}_k||^2]
$$

= $\left(1 - \frac{1}{m_i}\right) \frac{1}{m_i} \sum_{h=1}^{m_i} \mathbb{E} [||\mathbf{r}_{h,k}^t - \bar{\mathbf{X}}_k||^2]$
+ $\frac{1}{m_i} \mathbb{E} [\|\Phi_k^{t+1} - \bar{\mathbf{X}}_k\|^2].$

Now, denoting $q_k^t = \mathbf{Y}_k - (G(\Phi_k^t) - \nabla F(\bar{\mathbf{X}}_k) + \rho \mathbf{D} \Phi_k^t \rho$ **DX**_k), we have $\overline{2}$ $\overline{2}$

$$
\|\Phi_k^{t+1} - \bar{X}_k\|^2 = \|\Phi_k^{t+1} - \Phi_k^t + \Phi_k^t - \bar{X}_k\|^2 \n= \|\Phi_k^t - \bar{X}_k\|^2 + 2\gamma \langle \Phi_k^t - \bar{X}_k, q_k^t \rangle + \gamma^2 \|q_k^t\|^2 \n\le (1 + \gamma) \|\Phi_k^t - \bar{X}_k\|^2 + (\gamma^2 + \gamma) \|q_k^t\|^2,
$$

and using [\(48\)](#page-11-0) yields

$$
\mathbb{E}\left[\left\|\boldsymbol{q}_{k}^{t}\right\|^{2}\right] \n= \mathbb{E}[\left\|\mathbf{Y}_{k} - (\nabla F(\Phi_{k}^{t}) - \nabla F(\bar{\mathbf{X}}_{k}) + \rho \mathbf{D} \Phi_{k}^{t} - \rho \mathbf{D} \mathbf{X}_{k})\right\|^{2}] \n+ \mathbb{E}\left[\left\|\boldsymbol{G}(\Phi_{k}^{t}) - \nabla F(\Phi_{k}^{t})\right\|^{2}\right] \n\leq \mathbb{E}\left[\left\|\boldsymbol{G}(\Phi_{k}^{t}) - \nabla F(\Phi_{k}^{t})\right\|^{2}\right] \n+ 2 \mathbb{E}[\left\|\nabla F(\Phi_{k}^{t}) - \nabla F(\bar{\mathbf{X}}_{k}) + \rho \mathbf{D} \Phi_{k}^{t} - \rho \mathbf{D} \bar{\mathbf{X}}_{k}\right\|^{2}] \n+ 2 \mathbb{E}[\left\|\mathbf{Y}_{k} - \bar{\mathbf{Y}}_{k} + \bar{\mathbf{Y}}_{k} + (\rho \mathbf{D} \bar{\mathbf{X}}_{k} - \rho \mathbf{D} \mathbf{X}_{k})\right\|^{2}] \n\leq \mathbb{E}\left[\left\|\boldsymbol{G}(\Phi_{k}^{t}) - \nabla F(\Phi_{k}^{t})\right\|^{2}\right] + 2(L^{2} + \rho^{2} d_{u}^{2}) \mathbb{E}[\left\|\Phi_{k}^{t} - \bar{\mathbf{X}}_{k}\right\|^{2} \n+ 4\left(18 + \frac{36\gamma\tau\rho^{2}d_{u}^{2}}{|\tilde{\lambda}_{\max}|\rho|}\right) \|\hat{\mathbf{d}}_{k}\|^{2} + 4N\|\nabla F(\bar{\mathbf{x}}_{k})\|^{2} \n\leq (4L^{2} + 2\rho^{2} d_{u}^{2}) \mathbb{E}[\|\Phi_{k}^{t} - \bar{\mathbf{X}}_{k}\|^{2} + 4\left(18 + \frac{36\gamma\tau\rho^{2}d_{u}^{2}}{|\tilde{\lambda}_{\max}|\rho|}\right) \|\hat{\mathbf{d}}_{k}\|^{2} \n+ 2L^{2} \mathbb{E}[t_{k}^{t}] + 4N\
$$

Therefore we have

$$
\mathbb{E}\left[t_{k}^{t+1} | \mathcal{F}^{k}\right] = \left(1 - \frac{1}{m_{i}}\right) \frac{1}{m_{i}} \sum_{h=1}^{m_{i}} ||\mathbf{r}_{h,k}^{t} - \bar{X}_{k}||^{2} \n+ \frac{1}{m_{i}} ||\Phi_{k}^{t+1} - \bar{X}_{k}||^{2} \n\leq (1 - \frac{1}{m_{i}})t_{k}^{t} + \frac{1}{m_{i}} ((1 + \gamma) ||\Phi_{k}^{t} - \bar{X}_{k}||^{2} + (\gamma^{2} + \gamma) ||q_{k}^{t}||^{2}) \n\leq \left(1 - \frac{1}{m_{u}} + \frac{2L^{2}}{m_{l}} (\gamma^{2} + \gamma)\right) \mathbb{E}[t_{k}^{t}] + \n+ \left(\frac{1}{m_{l}}(1 + \gamma) + \frac{4L^{2} + 2\rho^{2}d_{u}^{2}}{m_{l}} (\gamma^{2} + \gamma)\right) \mathbb{E}[\|\Phi_{k}^{t} - \bar{\mathbf{X}}_{k}||^{2} + \n+ \frac{4}{m_{l}} \left(18 + \frac{36\gamma\tau\rho^{2}d_{u}^{2}}{|\tilde{\lambda}_{\max}|\rho|}\right) (\gamma^{2} + \gamma) ||\hat{\mathbf{d}}_{k}||^{2} + \n+ \frac{4N}{m_{l}} (\gamma^{2} + \gamma) ||\nabla F(\bar{x}_{k})||^{2} \n\leq \left(1 - \frac{1}{2m_{u}}\right) \mathbb{E}[t_{k}^{t}] + \frac{2}{m_{l}} \mathbb{E}[\|\Phi_{k}^{t} - \bar{\mathbf{X}}_{k} \|^{2} + \n+ \frac{4}{m_{l}} \left(18 + \frac{36\gamma\tau\rho^{2}d_{u}^{2}}{|\tilde{\lambda}_{\max}|\rho|}\right) (\gamma^{2} + \gamma) ||\hat{\mathbf{d}}_{k}||^{2} + \n+ \frac{4N}{m_{l}} (\gamma^{2} + \gamma) ||\nabla F(\bar{x}_{k})||^{2}
$$
\n(55)

where the last inequality holds when

$$
\frac{2L^2}{m_l}(\gamma^2 + \gamma) < \frac{1}{2m_u}, \quad \gamma + (4L^2 + 2\rho^2 d_u^2)(\gamma^2 + \gamma) < 1,\tag{56}
$$

which can be satisfied by choosing $\gamma < \min\{\bar{\gamma}_8, \bar{\gamma}_9\}$. Iterating [\(55\)](#page-12-1) for $t = 0, ..., \tau - 1$ then yields:

$$
\mathbb{E}[t_k^t] \leq (1 - \frac{1}{2m_u})^t \mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2] \n+ \frac{4}{m_l} \left(18 + \frac{36\gamma\tau\rho^2 d_u^2}{|\bar{\lambda}_{\text{max}}|\rho|}\right) (\gamma^2 + \gamma) \sum_{l=0}^{t-1} (1 - \frac{1}{2m_u})^l \|\hat{\mathbf{d}}_k\|^2 \n+ \frac{4N}{m_l} (\gamma^2 + \gamma) \sum_{l=0}^{t-1} (1 - \frac{1}{2m_u})^l \|\nabla F(\bar{x}_k)\|^2 \n+ \frac{2}{m_l} \sum_{l=0}^{t-1} (1 - \frac{1}{2m_u})^{t-1-l} \mathbb{E}[\|\Phi_k^l - \bar{\mathbf{X}}_k\|^2 \n\leq 9 \frac{2\gamma\tau}{|\bar{\lambda}_{\text{max}}|\rho} (1 - \frac{1}{2m_u})^t \mathbb{E}[\|\hat{\mathbf{d}}_k\|^2] \n+ \frac{4N}{m_l} (\gamma^2 + \gamma) \sum_{l=0}^{t-1} (1 - \frac{1}{2m_u})^l \|\nabla F(\bar{x}_k)\|^2 \n+ \frac{2}{m_l} \sum_{l=0}^{t-1} (1 - \frac{1}{2m_u})^{t-1-l} \mathbb{E}[\|\Phi_k^l - \bar{\mathbf{X}}_k\|^2 \n+ \frac{4}{m_l} \left(18 + \frac{36\gamma\tau\rho^2 d_u^2}{|\bar{\lambda}_{\text{max}}|\rho|}\right) (\gamma^2 + \gamma) 2m_u \|\hat{\mathbf{d}}_k\|^2.
$$

Summing the above relation over $t = 0, 1, ..., \tau - 1$ we get:

$$
\mathbb{E}[t_k]
$$
\n
$$
e_2 \leq \left(\frac{36m_u\gamma\tau}{|\tilde{\lambda}_{\max}|\rho} + \frac{4\tau}{m_l}(18 + \frac{36\gamma\tau\rho^2d_u^2}{|\tilde{\lambda}_{\max}|\rho}) (\gamma^2 + \gamma)2m_u\right) \mathbb{E}[\|\hat{\mathbf{d}}_k\|^2]
$$
\n
$$
+ \frac{4m_u}{m_l} \|\hat{\mathbf{\Phi}}_k\|^2 + \frac{4N}{m_l} (\gamma^2 + \gamma)2m_u\tau \|\nabla F(\bar{x}_k)\|^2,
$$

and using [\(49\)](#page-11-3) then yields

$$
\mathbb{E}[t_k] \le (s_0 + s_1)\mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] + s_2 \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] + + \frac{192m_u}{m_l} \tau^2 \gamma^2 L^2 \mathbb{E}[t_k],
$$

where s_0 , s_1 and s_2 are defined in [\(54\)](#page-11-4).

Letting $\gamma < \bar{\gamma}_{10}$, it follows that

$$
\frac{192m_u}{m_l}\tau^2\gamma^2L^2 < \frac{1}{2},\tag{57}
$$

and thus $\mathbb{E}[t_k] \leq 2(s_0+s_1)\mathbb{E}[||\widehat{\mathbf{d}}_k||^2] + 2s_2\mathbb{E}[||\nabla F(\bar{x}_k)||^2].$

Lemma 6. *Let Assumptions [1,](#page-2-6) [2](#page-2-7) and [4](#page-3-4) hold; when* γ < $\min\{\bar{\gamma}_1, \bar{\gamma}_{11}\}$ *, it holds that* $\forall k \geq 0$

$$
\mathbb{E}[\|\hat{\mathbf{d}}_{k+1}\|^2] \leq (\delta + \frac{\tilde{q}_1}{1-\delta})\mathbb{E}[\|\hat{\mathbf{d}}_k\|^2] + \frac{\tilde{q}_2}{1-\delta}\mathbb{E}[\|\sum_t \overline{\nabla F}(\boldsymbol{\Phi}_k^t)\|^2] + \frac{\tilde{q}_3}{1-\delta}\mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] + \tilde{q}_4\mathbb{E}[t_k],
$$
\n(58)

where

$$
\delta := 1 - \frac{|\tilde{\lambda}_{\text{max}}| \rho \tau \gamma}{2}, \tilde{c}_0 := \gamma^2 \tau \tilde{\beta}_0, \tilde{c}_1 := \gamma^3 \tau^2 \tilde{\beta}_1,
$$

\n
$$
\tilde{c}_2 := 12L^2 \gamma^2 \|\hat{\mathbf{V}}^{-1}\|^2 N, \tilde{c}_3 := \gamma^2 \tau \tilde{\beta}_3,
$$

\n
$$
\tilde{q}_1 := \tilde{c}_0 \gamma \tau \kappa_1 + \tilde{c}_1, \tilde{q}_2 := \tilde{c}_2, \quad \tilde{q}_3 := 16\tau^3 \gamma^2 N \tilde{c}_0,
$$

\n
$$
\tilde{q}_4 := (48\tau^2 \gamma^2 L^2 \tilde{c}_0 + \tilde{c}_3) / (1 - \delta).
$$

]

Proof. When γ < min $\{\bar{\gamma}_1, \bar{\gamma}_{11}\}$ we have

$$
\gamma \tau \le \frac{2}{|\tilde{\lambda}_{\text{max}}|\rho}, \quad |\tilde{\lambda}_{\text{min}}|\rho \tau \gamma < 2,
$$

6 $\gamma^2 \tau (3L^2 + \rho^2 d_u^2) < \frac{1/4}{\tau - 1}.$ (59)

Substituting [\(48\)](#page-11-0) into [\(28\)](#page-8-1) then yields

$$
\|\widehat{\mathbf{h}}_k\|^2 \leq \tilde{c}_0 \|\widehat{\mathbf{\Phi}}_k\|^2 + \tilde{c}_1 \|\widehat{\mathbf{d}}_k\|^2 + \tilde{c}_2 \|\sum_t \overline{\nabla F}(\mathbf{\Phi}_k^t)\|^2 + \tilde{c}_3 t_k,
$$

and together with [\(49\)](#page-11-3) and [\(17\)](#page-7-7), it proves that [\(58\)](#page-12-2) holds. \Box

B. Theorem [2](#page-4-1)

Based on [\(30\)](#page-8-3), substituting $y = \bar{x}_{k+1}$ and $z = \bar{x}_k$ into [\(37\)](#page-9-2) and using (27) , (48) , we get

$$
\mathbb{E}[F(\bar{x}_{k+1})]
$$
\n
$$
\leq \mathbb{E}[F(\bar{x}_k)] - \gamma \mathbb{E}[\langle \nabla F(\bar{x}_k), \frac{1}{N} (\sum_t \sum_i \nabla f_i (\phi_{i,k}^t) + \epsilon_k) \rangle]
$$
\n
$$
+ 2\gamma^2 \tau L \mathbb{E}[\sum_t ||\frac{1}{N} \sum_i \nabla f_i (\phi_{i,k}^t) ||^2]
$$
\n
$$
+ \frac{2\gamma^2 \tau L}{N} (2L^2 \mathbb{E}[||\hat{\Phi}_k||^2] + 2L^2 \mathbb{E}[t_k])
$$
\n
$$
+ \frac{2\gamma^2 L \rho^2 d_u^2 \tau(\mathbb{E}[||\hat{\Phi}_k||^2] + \tau \mathbb{E}[||\mathbf{X}_k - \bar{\mathbf{X}}_k||^2])}{N}.
$$

Similarly to Appendix [B-B,](#page-9-5) using $2\langle a, b \rangle = ||a||^2 + ||b||^2$ – $||a - b||^2$ we have

$$
\begin{split}\n&-\mathbb{E}[\langle \nabla F\left(\bar{x}_k\right), \frac{1}{N} \sum_{t} \sum_{i} \nabla f_i \left(\phi_{i,k}^t\right) + \frac{\epsilon_k}{N} \rangle] \\
&\leq -\frac{\tau}{4} \mathbb{E}[\|\nabla F\left(\bar{x}_k\right)\|^2] - \frac{1}{2} \mathbb{E}[\sum_{t} \|\frac{1}{N} \sum_{i} \nabla f_i \left(\phi_{i,k}^t\right)\|^2] \\
&+ \left(\frac{L^2}{2N} + \frac{2\rho^2 d_u^2}{N}\right) \mathbb{E}[\widehat{\Phi}_k\|^2] + \frac{2\rho^2 d_u^2 \tau}{N} \mathbb{E}[\|\mathbf{X}_k - \bar{\mathbf{X}}_k\|^2].\n\end{split}
$$

Combining the two inequalities above, and using [\(16\)](#page-7-6), yields

$$
\mathbb{E}[F(\bar{x}_{k+1})] \leq \mathbb{E}[F(\bar{x}_k)] - \frac{\gamma \tau}{4} \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2]
$$

\n
$$
- \frac{\gamma}{2} (1 - 4\gamma \tau L) \mathbb{E} \left[\sum_t \|\frac{1}{N} \sum_i \nabla f_i(\phi_{i,k}^t) \|^2 \right]
$$

\n
$$
+ \left(\frac{\gamma L^2}{2N} + \frac{2\gamma^2 L \rho^2 d_u^2 \tau + 2\gamma \rho^2 d_u^2 + 4\gamma^2 \tau L^3}{N} \right) \mathbb{E}[\|\widehat{\Phi}_k\|^2]
$$

\n
$$
+ \frac{18\gamma \rho^2 d_u^2 \tau + 18\gamma^2 \rho^2 d_u^2 \tau^2 L}{N} \frac{2\gamma \tau}{|\widetilde{\lambda}_{\max}|\rho} \mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2]
$$

\n
$$
+ \frac{4\gamma^2 \tau L^3 \mathbb{E}[t_k]}{N}.
$$

\n(60)

Combining [\(60\)](#page-13-0) together with [\(49\)](#page-11-3) and [\(53\)](#page-11-5), and using $\gamma \leq$ 1, yields

$$
\mathbb{E}[F(\bar{x}_{k+1})] \leq \mathbb{E}[F(\bar{x}_k)] - \frac{\gamma \tau}{4} \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2]
$$

$$
-\frac{\gamma}{2} (1 - 4\gamma \tau L) \mathbb{E}[\sum_t \|\frac{1}{N} \sum_i \nabla f_i(\phi_{i,k}^t) \|^2]
$$

$$
+ (k_0 k_2 + 2(k_3 k_0 + k_4) s_2) \mathbb{E}[\|\nabla F(x)\|^2]
$$

 $+ (k_5 + k_0 k_1 + 2(k_3 k_0 + k_4)(s_0 + s_1)) \mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2],$

where

$$
k_0 := \alpha_0 \gamma, k_1 := \frac{72\gamma \tau^2}{|\tilde{\lambda}_{\text{max}}| \rho} + (18 + \frac{36\gamma \tau \rho^2 d_u^2}{|\tilde{\lambda}_{\text{max}}| \rho}) 16\tau^3 \gamma^2
$$

\n
$$
k_2 := 16\tau^3 \gamma^2 N, k_3 := 48\tau^2 \gamma^2 L^2
$$

\n
$$
k_4 := \frac{4\gamma^2 \tau L^3}{N}, k_5 := \frac{36\gamma^2 \rho d_u^2 \tau^2 + 36\gamma^3 \rho d_u^2 \tau^3 L}{N|\tilde{\lambda}_{\text{max}}|}.
$$

Letting $\gamma \leq \min\{1, \bar{\gamma}_{12}, \bar{\gamma}_{13}\}\$, then

$$
k_0 k_2 + 2(k_3 k_0 + k_4) s_2 \le \frac{\gamma \tau}{8}, \quad 4\gamma \tau L \le \frac{3}{4}, \tag{61}
$$

and we can upper bound the previous inequality by

$$
\mathbb{E}[F(\bar{x}_{k+1})] \leq \mathbb{E}[F(\bar{x}_k)] - \frac{\gamma \tau}{8} \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] \n- \frac{\gamma}{8} \mathbb{E}[\sum_t \|\frac{1}{N} \sum_i \nabla f_i(\phi_{i,k}^t)\|^2] \n+ (k_5 + k_0 k_1 + 2(k_3 k_0 + k_4)(s_0 + s_1)) \mathbb{E}[\|\hat{\mathbf{d}}_k\|^2].
$$

Rearranging we get

$$
\mathcal{D}_k \leq \frac{8}{\gamma \tau} (\mathbb{E}[\tilde{F}(\bar{x}_k)] - \mathbb{E}[\tilde{F}(\bar{x}_{k+1})]) + \frac{8}{\gamma \tau} (k_5 + k_0 k_1 + 2(k_3 k_0 + k_4)(s_0 + s_1)) \mathbb{E}[\|\hat{\mathbf{d}}_k\|^2],
$$
\n(62)

where \mathcal{D}_k is defined in [\(7\)](#page-3-7), and $\tilde{F}(\bar{x}_k) = F(\bar{x}_k) - F(x^*)$. Summing [\(62\)](#page-13-1) over $k = 0, 1, ..., K - 1$ and using $-\tilde{F}(\bar{x}_k) \le$ 0, it holds that

$$
\sum_{r=0}^{K-1} \mathcal{D}_k \le \frac{8\tilde{F}(\bar{x}_0)}{\gamma \tau}
$$
\n
$$
+ \frac{8(k_5 + k_0k_1 + 2(k_3k_0 + k_4)(s_0 + s_1))}{\gamma \tau} \sum_{k=0}^{K-1} ||\mathbb{E}[\hat{\mathbf{d}}_k||^2].
$$
\n(63)

According to [\(58\)](#page-12-2) and [\(53\)](#page-11-5), we derive that $\forall k \geq 0$,

$$
\mathbb{E}[\|\widehat{\mathbf{d}}_{k+1}\|^2] \leq \left(\delta + \frac{\tilde{q}_1}{1-\delta} + 2\tilde{q}_4(s_0 + s_1)\right) \mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] \n+ \frac{\tilde{q}_2}{1-\delta} \mathbb{E}[\|\sum_t \overline{\nabla F}(\boldsymbol{\Phi}_k^t)\|^2] + \left(\frac{\tilde{q}_3}{1-\delta} + 2\tilde{q}_4 s_2\right) \mathbb{E}[\|\nabla F(\bar{x}_k)\|^2] < \tilde{\delta} \mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] + \tilde{R} \mathcal{D}_k,
$$
\n(64)

where

$$
\tilde{R} = \max \left\{ \frac{\tilde{q}_2 \tau^2}{1 - \delta}, \frac{\tilde{q}_3}{1 - \delta} + 2\tilde{q}_4 s_2 \right\}
$$
\n
$$
= \max \left\{ \frac{2\tilde{q}_2 \tau}{\rho \gamma |\tilde{\lambda}_{max}|}, \frac{2(\tilde{q}_3 + 2(48\tau^2 \gamma^2 L^2 \tilde{c}_0 + \tilde{c}_3) s_2)}{\tau \rho \gamma |\tilde{\lambda}_{max}|} \right\}.
$$

Letting $\gamma \leq \min\{\bar{\gamma}_1, \bar{\gamma}_{14}\}\$, then

$$
\tilde{\delta} = \delta + \frac{\tilde{q}_1}{1 - \delta} + 2\tilde{q}_4(s_0 + s_1) < 1. \tag{65}
$$

Iterating [\(64\)](#page-13-2) yields $\forall k \geq 1$, $\mathbb{E}[\|\hat{\mathbf{d}}_k\|^2] \leq \delta^k \mathbb{E}[\|\hat{\mathbf{d}}_0\|^2] +$ $\tilde{R} \sum_{\ell=0}^{k-1} \tilde{\delta}^{k-1-\ell} \mathcal{D}_{\ell}$, and summing over $k = 0, \ldots, K-1$ it holds that

$$
\sum_{k=0}^{K-1} \mathbb{E}[\|\widehat{\mathbf{d}}_k\|^2] \le \frac{1}{1-\tilde{\delta}}\|\widehat{\mathbf{d}}_0\|^2 + \sum_{k=0}^{K-1} \frac{\tilde{R}}{1-\tilde{\delta}} \mathcal{D}_k. \tag{66}
$$

Substituting [\(66\)](#page-13-3) into [\(63\)](#page-13-4), and rearranging, we obtain

$$
\left(1 - \frac{\tilde{R}}{1 - \tilde{\delta}} \frac{8}{\gamma \tau} (k_5 + k_0 k_1 + 2(k_3 k_0 + k_4)(s_0 + s_1))\right) \sum_{k=0}^{K-1} \mathcal{D}_k
$$

$$
\left(8\tilde{F}(\bar{x}_0) - 8(k_5 + k_0 k_1 + 2(k_3 k_0 + k_4)(s_0 + s_1))\right) \sum_{k=0}^{K-1} \mathcal{D}_k
$$

$$
\leq \frac{\sigma r \left(x_0\right)}{\gamma \tau} + \frac{\sigma \left(x_0 + x_0\right) \tau + 2 \left(x_3 \right) \tau + 4 \left(x_0 + x_1\right) \tau}{\gamma \tau (1 - \tilde{\delta})} ||\tilde{\mathbf{d}}_0||^2.
$$

Similarly to the previous analysis, when $\gamma \leq \frac{\gamma_{14}}{2} = \bar{\gamma}_{15}$, $1-\tilde{\delta} \geq \frac{|\tilde{\lambda}_{\max}|\rho\tau\gamma}{4}$. Let $\gamma \leq \min{\{\bar{\gamma}_{16}, \bar{\gamma}_{17}\}}$, then

$$
\frac{\tilde{R}}{1-\tilde{\delta}}\frac{8}{\gamma\tau}\left(k_5+k_0k_1+2(k_3k_0+k_4)(s_0+s_1)\right)\leq\frac{1}{2},
$$

and therefore it follows that

$$
\frac{1}{K} \sum_{k=0}^{K-1} \mathcal{D}_k \le \frac{16\tilde{F}(\bar{x}_0)}{K\gamma\tau} \n+ \frac{16}{K\gamma\tau} (k_5 + k_0k_1 + 2(k_3k_0 + k_4)(s_0 + s_1)) \frac{1}{1 - \tilde{\delta}} \|\hat{\mathbf{d}}_0\|^2.
$$

By collecting all step-size conditions, if the step-size γ satisfies [\(51\)](#page-11-6), [\(56\)](#page-12-3), [\(57\)](#page-12-4), [\(59\)](#page-13-5), [\(61\)](#page-13-6), [\(65\)](#page-13-7) and [\(67\)](#page-14-35), then the states $\{X_k\}$ generated by LT-ADMM-VR converge to the stationary point, concluding the proof.

REFERENCES

- [1] D. K. Molzahn, F. Dorfler, H. Sandberg, S. H. Low, S. Chakrabarti, R. Baldick, and J. Lavaei, "A Survey of Distributed Optimization and Control Algorithms for Electric Power Systems," *IEEE Transactions on Smart Grid*, vol. 8, no. 6, pp. 2941–2962, Nov. 2017.
- [2] O. Shorinwa, T. Halsted, J. Yu, and M. Schwager, "Distributed optimization methods for multi-robot systems: Part 1—a tutorial," *IEEE Robotics & Automation Magazine*, 2024.
- [3] ——, "Distributed optimization methods for multi-robot systems: Part 2—a survey," *IEEE Robotics & Automation Magazine*, 2024.
- [4] R. Mohebifard and A. Hajbabaie, "Distributed optimization and coordination algorithms for dynamic traffic metering in urban street networks," *IEEE Transactions on Intelligent Transportation Systems*, vol. 20, no. 5, pp. 1930–1941, 2018.
- [5] A. Nedić and J. Liu, "Distributed Optimization for Control," Annual *Review of Control, Robotics, and Autonomous Systems*, vol. 1, no. 1, pp. 77–103, May 2018.
- [6] T. Gafni, N. Shlezinger, K. Cohen, Y. C. Eldar, and H. V. Poor, "Federated Learning: A signal processing perspective," *IEEE Signal Processing Magazine*, vol. 39, no. 3, pp. 14–41, May 2022.
- [7] L. Bottou, F. E. Curtis, and J. Nocedal, "Optimization methods for largescale machine learning," *SIAM review*, vol. 60, no. 2, pp. 223–311, 2018.
- [8] S. A. Alghunaim and K. Yuan, "A unified and refined convergence analysis for non-convex decentralized learning," *IEEE Transactions on Signal Processing*, vol. 70, pp. 3264–3279, 2022.
- [9] Y. Liu, T. Lin, A. Koloskova, and S. U. Stich, "Decentralized Gradient Tracking with Local Steps," Jan. 2023.
- [10] S. A. Alghunaim, "Local Exact-Diffusion for Decentralized Optimization and Learning," Feb. 2023.
- [11] L. Guo, S. A. Alghunaim, K. Yuan, L. Condat, and J. Cao, "RandCom: Random Communication Skipping Method for Decentralized Stochastic Optimization," Oct. 2023.
- [12] H. Li, Z. Lin, and Y. Fang, "Variance Reduced EXTRA and DIGing and Their Optimal Acceleration for Strongly Convex Decentralized Optimization," *Journal of Machine Learning Research*, vol. 23, no. 222, pp. 1–41, 2022.
- [13] X. Jiang, X. Zeng, J. Sun, and J. Chen, "Distributed Stochastic Gradient Tracking Algorithm With Variance Reduction for Non-Convex
Optimization." IEEE Transactions on Neural Networks and Learning IEEE Transactions on Neural Networks and Learning *Systems*, vol. 34, no. 9, pp. 5310–5321, Sep. 2023.
- [14] R. Xin, U. A. Khan, and S. Kar, "Variance-Reduced Decentralized Stochastic Optimization With Accelerated Convergence," *IEEE Transactions on Signal Processing*, vol. 68, pp. 6255–6271, 2020.
- [15] - , "A Fast Randomized Incremental Gradient Method for Decentralized Nonconvex Optimization," *IEEE Transactions on Automatic Control*, vol. 67, no. 10, pp. 5150–5165, 2022.
- $_{k}$ [16] -, "Fast Decentralized Nonconvex Finite-Sum Optimization with Recursive Variance Reduction," *SIAM Journal on Optimization*, vol. 32, no. 1, pp. 1–28, Mar. 2022.
- [17] W. Shi, Q. Ling, G. Wu, and W. Yin, "Extra: An exact first-order algorithm for decentralized consensus optimization," *SIAM Journal on Optimization*, vol. 25, no. 2, pp. 944–966, 2015.
- [18] A. Nedic, A. Olshevsky, and W. Shi, "Achieving geometric convergence for distributed optimization over time-varying graphs," *SIAM Journal on Optimization*, vol. 27, no. 4, pp. 2597–2633, 2017.
- [19] F. Saadatniaki, R. Xin, and U. A. Khan, "Decentralized optimization over time-varying directed graphs with row and column-stochastic matrices," *IEEE Transactions on Automatic Control*, vol. 65, no. 11, pp. 4769–4780, 2020.
- [20] X. Ren, D. Li, Y. Xi, and H. Shao, "An accelerated distributed gradient method with local memory," *Automatica*, vol. 146, p. 110260, 2022.
- [21] N. Bastianello, R. Carli, L. Schenato, and M. Todescato, "Asynchronous Distributed Optimization Over Lossy Networks via Relaxed ADMM: Stability and Linear Convergence," *IEEE Transactions on Automatic Control*, vol. 66, no. 6, pp. 2620–2635, Jun. 2021.
- [22] N. Bastianello, D. Deplano, M. Franceschelli, and K. H. Johansson, "Robust Online Learning over Networks," *IEEE Transactions on Automatic Control*, pp. 1–14, 2024.
- [23] A. Makhdoumi and A. Ozdaglar, "Convergence rate of distributed admm over networks," *IEEE Transactions on Automatic Control*, vol. 62, no. 10, pp. 5082–5095, 2017.
- [24] V. Khatana and M. V. Salapaka, "Dc-distadmm: Admm algorithm for constrained optimization over directed graphs," *IEEE Transactions on Automatic Control*, vol. 68, no. 9, pp. 5365–5380, 2022.
- [25] A. Defazio, F. Bach, and S. Lacoste-Julien, "Saga: A fast incremental gradient method with support for non-strongly convex composite objectives," *Advances in neural information processing systems*, vol. 27, 2014.
- [26] B. McMahan, E. Moore, D. Ramage, S. Hampson, and B. A. y. Arcas, "Communication-Efficient Learning of Deep Networks from Decentralized Data," in *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics*, ser. Proceedings of Machine Learning Research, A. Singh and J. Zhu, Eds., vol. 54. Fort Lauderdale, FL, USA: PMLR, Apr. 2017, pp. 1273–1282.
- [27] X. Zhang, M. Hong, S. Dhople, W. Yin, and Y. Liu, "FedPD: A Federated Learning Framework With Adaptivity to Non-IID Data," *IEEE Transactions on Signal Processing*, vol. 69, pp. 6055–6070, 2021.
- [28] K. Mishchenko, G. Malinovsky, S. Stich, and P. Richtarik, "ProxSkip: Yes! Local Gradient Steps Provably Lead to Communication Acceleration! Finally!" in *Proceedings of the 39th International Conference on Machine Learning*, ser. Proceedings of Machine Learning Research, K. Chaudhuri, S. Jegelka, L. Song, C. Szepesvari, G. Niu, and S. Sabato, Eds., vol. 162. PMLR, Jul. 2022, pp. 15 750–15 769.
- [29] A. Mitra, R. Jaafar, G. J. Pappas, and H. Hassani, "Linear Convergence in Federated Learning: Tackling Client Heterogeneity and Sparse Gradients," in *Advances in Neural Information Processing Systems*, M. Ranzato, A. Beygelzimer, Y. Dauphin, P. S. Liang, and J. W. Vaughan, Eds., vol. 34. Curran Associates, Inc., 2021, pp. 14 606– 14 619.
- [30] L. Condat, I. Agarský, G. Malinovsky, and P. Richtárik, "TAMUNA: Doubly Accelerated Federated Learning with Local Training, Compression, and Partial Participation," May 2023.
- [31] E. D. Hien Nguyen, S. A. Alghunaim, K. Yuan, and C. A. Uribe, "On the Performance of Gradient Tracking with Local Updates," in *2023 62nd IEEE Conference on Decision and Control (CDC)*. Singapore, Singapore: IEEE, Dec. 2023, pp. 4309–4313.
- [32] A. S. Berahas, R. Bollapragada, and E. Wei, "On the Convergence of Nested Decentralized Gradient Methods With Multiple Consensus and Gradient Steps," *IEEE Transactions on Signal Processing*, vol. 69, pp. 4192–4203, 2021.
- [33] C. Iakovidou and E. Wei, "S-NEAR-DGD: A Flexible Distributed Stochastic Gradient Method for Inexact Communication," *IEEE Transactions on Automatic Control*, vol. 68, no. 2, pp. 1281–1287, Feb. 2023.
- [34] R. Johnson and T. Zhang, "Accelerating stochastic gradient descent using predictive variance reduction," *Advances in neural information processing systems*, vol. 26, 2013.
- [35] L. M. Nguyen, J. Liu, K. Scheinberg, and M. Takáč, "Sarah: A novel method for machine learning problems using stochastic recursive gradient," in *International conference on machine learning*. PMLR, 2017, pp. 2613–2621.

[36] Y. Nesterov, *Introductory lectures on convex optimization: A basic course*. Springer Science & Business Media, 2013, vol. 87.