

JIGGLING: AN H-PRINCIPLE WITHOUT HOMOTOPICAL ASSUMPTIONS

ANNA FOKMA, ÁLVARO DEL PINO, AND LAURAN TOUSSAINT

ABSTRACT. The jiggling lemma of Thurston shows that any triangulation can be jiggled (read: subdivided and then perturbed) to be in general position with respect to a distribution.

Our main result is a generalization of Thurston’s lemma. It states that piecewise smooth solutions of a given open and fiberwise dense differential relation $\mathcal{R} \subset J^1(E)$ of first order can be constructed by jiggling arbitrary sections of E . Our statement also holds in parametric and relative form. We understand this as an h-principle without homotopical assumptions for piecewise smooth solutions of \mathcal{R} .

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Subdivisions	10
4. Linearization of piecewise maps	16
5. Slope perturbations of piecewise linear maps	19
6. Jiggling	21
7. Jiggling as an h-principle	25
8. Applications	27
References	32

1. INTRODUCTION

In [29, 30], Thurston established the h-principle for foliations. One of the ideas that goes into the proof is to decompose the manifold using a triangulation whose simplices are in general position with respect to a given starting distribution; see Definition 8.8. To construct such triangulations, Thurston states what is now known as the ‘jiggling lemma’:

Lemma (Thurston’s jiggling lemma [30]). *Consider a manifold M endowed with a distribution ξ . Any smooth triangulation of M can be subdivided and subsequently perturbed to be in general position with respect to ξ , over any given compact subset. Moreover, the perturbation can be assumed to be C^1 -small.*

This result has found numerous applications in the construction of geometric structures on manifolds, beyond foliations. These include Haefliger structures with transverse geometry [19], contact structures [8, 31], and Engel structures [6, 25].

Date: January 24, 2025.

2020 Mathematics Subject Classification. Primary: 57R45. Secondary: 57Q65, 57R05, 57R35.

Key words and phrases. h-principle, jiggling, piecewise smooth, differential relations.

1.1. Jiggling sections. The main contribution of the present paper is the following observation: general h-principles in the piecewise smooth category can be established using (a suitable generalization of) jiggling. Concretely, we prove a generalization of Thurston’s lemma in which, instead of jiggling a triangulation so that it is in general position, we jiggle a section of a fiber bundle $E \rightarrow M$ to yield a piecewise smooth solution of a given differential relation $\mathcal{R} \subset J^1(E)$.

We assume a metric has been fixed on $J^1(E)$ in order to talk about C^1 -distances. This allows us to quantitatively control of the size of the jiggling:

Definition 1.1. Let $E \rightarrow M$ be a fiber bundle and let $s, s' : M \rightarrow E$ be piecewise smooth sections with respect to triangulations T and T' of M , respectively. We say that (s', T') is an ε -**jiggling** of (s, T) if T' is a subdivision of T and $d_{C^1}(s, s') < \varepsilon$, where ε is a positive constant if M is compact, and a positive function otherwise.

In order for the jiggling process to produce solutions of $\mathcal{R} \subset J^1(E)$, we need to impose two conditions on the differential relation. First of all, as is common for h-principles, we want the relation \mathcal{R} to be open as a subset of $J^1(E)$. Secondly, we want it to be fiberwise dense; this means that any jet $\sigma \in J^1(E)$ can be made to lie in \mathcal{R} using an arbitrarily small perturbation of its slope. We formalize this as follows, where $\pi_f : J^1(E) \rightarrow E$ is the front projection:

Definition 1.2. A relation $\mathcal{R} \subset J^1(E)$ is **fiberwise dense** if, for all $x \in E$, the restriction of \mathcal{R} to the fiber of π_f over x is dense.

We now state our main theorem:

Theorem 6.3. *Let M be a manifold, possibly open, endowed with a smooth triangulation $T : K \rightarrow M$. Consider a fiber bundle $E \rightarrow M$ and an open and fiberwise dense differential relation $\mathcal{R} \subset J^1(E)$. Then, given*

- a continuous function $\varepsilon : M \rightarrow \mathbb{R}_+$,
- a piecewise smooth section $s : M \rightarrow E$, and
- a subcomplex $Q \subset M$ of T such that $s|_{\mathcal{O}_p(Q)}$ is a solution of \mathcal{R} ,

there exists an ε -jiggling (s', T') of (s, T) such that

- s' is a piecewise solution of \mathcal{R} with respect to T' , and
- $s'|_Q = s|_Q$.

The idea of the proof is as follows: The conditions we pose on the relation \mathcal{R} (being open and fiberwise dense) imply that the desired solutions exist over any sufficiently small simplex. As such, after subdividing the triangulation, we have local solutions over all simplices, and it remains to glue them together to a global solution. This is slightly delicate, so we assign colors to the top-dimensional simplices in M such that a simplex and all its adjacent simplices are of a different color. Then, by induction on the color of a simplex, we jiggle the section over all simplices of the same color. If our jiggling is small enough, the solution we constructed over simplices of previous colors is only modified slightly and remains a solution.

The idea of jiggling ‘one color at a time’ was inspired by the coloring argument used by Donaldson in [9], which he used to achieve quantitative transversality in the asymptotically holomorphic setting. Coloring allows us to avoid various size and distance estimates in the Grassmannian appearing in Thurston’s original proof. It is in this simplification that our argument deviates strongly from his, and therefore it can be more readily generalized.

1.2. Jiggling as an h-principle. Theorem 6.3 naturally implies a weak equivalence between the spaces of piecewise smooth solutions of \mathcal{R} and continuous sections of E , with one subtlety. Since different sections are piecewise with respect to different triangulations, the space of triangulations is lurking in the background. However, this is a (semi-)simplicial set, not a topological space.

This leads us to define the simplicial set $\text{sSol}_{\text{PS}}(\mathcal{R})$ of piecewise smooth solutions of \mathcal{R} by letting $(\text{sSol}_{\text{PS}}(\mathcal{R}))_n$ be the set of piecewise smooth sections $s : M \times \Delta^n \rightarrow E$ that solve \mathcal{R} . We compare this simplicial set to the singular complex $\text{Sing}(\Gamma^0(E))$ of the space $\Gamma^0(E)$ of continuous sections of E .

Theorem 7.3. *Let $\mathcal{R} \subset J^1(E)$ be an open and fiberwise dense differential relation. The inclusion $\text{sSol}_{\text{PS}}(\mathcal{R}) \hookrightarrow \text{Sing}(\Gamma^0(E))$ is weak homotopy equivalence of simplicial sets.*

We pose a couple of interesting open questions: Can a version of Theorem 7.3 be established for higher order jets? Moreover, can the assumption of fiberwise density be replaced by some form of ampleness (as is required in convex integration)?

We remark that Theorem 7.3 is an h-principle without homotopical assumptions. Concretely: since our solutions $s : M \rightarrow E$ are meant to be piecewise, their first jet extensions $j^1s : M \rightarrow \mathcal{R}$ can have discontinuities along the triangulation. As such, they do not define a homotopy class of a map $M \rightarrow \mathcal{R}$. This is why no formal data appears in the statement.

The best known example of an h-principle without homotopical assumptions is the arborealization program in symplectic/contact topology [3, 1, 2]. One of its goals, which is similar in flavor to Thurston's jiggling, is to show that front projections of legendrians can be simplified so that their singularities are of so-called arboreal type. This can then be used to simplify the singularities of skeleta of Weinstein manifolds.

A key feature of arboreal singularities is that there are only finitely many models in each dimension. This is not true for singularities of triangulation type. This motivates a few more open questions:

- Can one establish a version of Theorem 7.3 for solutions with arboreal-type singularities?
- What about solutions whose germs along said arboreal singularities are modelled on some finite/countable list?
- Similarly, in the triangulated setting, can one establish a version of Theorem 7.3 in which finitely many models over each simplex are allowed?

1.3. Applications. We now state some consequences of our main theorems.

1.3.1. General position. The condition that a triangulation is in general position with respect to a distribution ξ is, of course, triangulation dependent. Because of this we have to do a bit of work to phrase it using an open and fiberwise dense relation. We do this in Section 8.3.2, establishing Thurston's jiggling lemma (Corollary 8.12). Something that is new in this direction is that we establish the result for manifolds that are not necessarily compact, which was not addressed by Thurston [30]. In fact, from a technical standpoint, an interesting feature of the proof is that we have to develop some theory regarding crystalline subdivision in the non-compact setting, which we do in Section 3.5.

Another application is the construction of triangulations that are in general position with respect to multiple distributions at the same time; a statement that has appeared already in applications, e.g. [31, 25].

1.3.2. Piecewise maps of full rank. According to Thurston's jiggling lemma, any map between manifolds can be jiggled to be piecewise transverse to a distribution or to be piecewise of maximal rank (immersive/submersive). These statements can be upgraded to full h-principles without homotopical assumptions, for the corresponding differential relations, using our results:

Corollary 8.3. *The following inclusions of simplicial sets are weak homotopy equivalences:*

$$\begin{aligned} \text{sSol}_{\text{PS}}(\mathcal{R}_{\text{transv}}) &\hookrightarrow \text{Sing}(C^0(M, N)), \\ \text{sSol}_{\text{PS}}(\mathcal{R}_{\text{imm}}) &\hookrightarrow \text{Sing}(C^0(M, N)) \text{ if } \dim M < \dim N, \text{ and} \\ \text{sSol}_{\text{PS}}(\mathcal{R}_{\text{subm}}) &\hookrightarrow \text{Sing}(C^0(M, N)) \text{ if } \dim M > \dim N. \end{aligned}$$

1.3.3. *Piecewise differentiable geometric structures.* Theorem 6.3 tells us that every non-vanishing 1-form is homotopic to a piecewise smooth contact form. Denoting by $\mathcal{R}_{\text{cont}}$ the relation for contact forms, we obtain the following h-principle:

Corollary 8.5. *The inclusion $\text{sSol}_{\text{PS}}(\mathcal{R}_{\text{cont}}) \hookrightarrow \text{Sing}(\Gamma^0(\Lambda^1(T^*M) \setminus \{0\}))$ is a weak homotopy equivalence.*

This complete flexibility in the piecewise smooth setting should be compared to the rigidity that appears in the study of other geometric structures of low regularity. Examples include C^0 -contact structures [23, 10, 26], C^0 -symplectic structures [24, 16], Lipschitz symplectic structures [18], and PL symplectic structures [5].

With that said, one may wonder whether one can “rigidify” the study of piecewise differentiable contact structures as follows. In dimension 3, one could study instead the differential relation \mathcal{R}^+ defining positive contact structures (e.g. compatible with a given orientation of the ambient manifold). This relation falls outside the scope of jiggling, since it is not fiberwise dense. We conjecture that there is rigidity in this setting. Moreover, it is unclear to us what to expect in the higher dimensional case.

1.4. **Outline of the paper.** In Section 2 we discuss the preliminaries of the paper, including jet bundles, triangulations and simplicial sets.

Sections 3 through 5 contain the ground work for jiggling. In Section 3 we discuss crystalline subdivisions of triangulations, as well as some other subdivision schemes required to work in the relative setting. In Section 4 we show how we can (locally) approximate a section by its linearization, reducing the jiggling of general sections to the jiggling of piecewise linear maps. In Section 5 we show that for open and fiberwise dense relations we can tilt any section to yield a local solution over any sufficiently small simplex.

The proof of the main theorem (Theorem 6.3) is contained in Section 6. The h-principle without homotopical assumptions that follows is discussed in Section 7. Applications are discussed in Section 8.

1.5. **Acknowledgements.** The authors want to thank Miguel Barata for reading a draft of this paper.

The second author is funded by the NWO (Dutch Research Council) project VI.Vidi.223.118. The third author is funded by the NWO project “proper Fredholm homotopy theory” (project number OCENW.M20.195) of the research programme Open Competition ENW M20-3.

2. PRELIMINARIES

In this section we cover some basics and fix notation. We discuss jet spaces and h-principles in Section 2.1. We continue with triangulations in Section 2.3, after which we end with a brief account of simplicial sets in Section 2.4.

2.1. **Jet spaces and h-principles.** In this section we briefly recall the standard terminology on (first order) jet spaces and h-principles. For a more elaborate overview we refer to [7, 13].

Given a fiber bundle $E \rightarrow M$, its **first jet bundle** $J^1(E) \rightarrow M$ is the bundle of first order Taylor polynomials of sections of E . The projection is defined by sending a Taylor polynomial at $x \in M$ to the point x . We observe that such a Taylor polynomial at x can be represented by a section locally defined around x . Vice versa, every section $s : M \rightarrow E$ induces a section $j^1s : M \rightarrow J^1(E)$ which we call its first jet. Since not every section of $J^1(E)$ comes from a section of E , we call sections which do arise in this manner **holonomic**. When interested in maps instead of sections, we define the first jet bundle $J^1(M, N) \rightarrow M$ of maps $M \rightarrow N$ in an analogous manner.

The terminology of jet spaces has proven to be a natural language to generalize the notion of a partial differential *equation* (PDE) to a partial differential *relation*.

Definition 2.1. A (first order) **partial differential relation** is a subset $\mathcal{R} \subset J^1(E)$.

We distinguish two notions of solutions of a relation \mathcal{R} . Firstly, there exists a space $\text{FSol}(\mathcal{R})$ of **formal solutions** whose elements are sections of $J^1(E)$ with image in \mathcal{R} . We endow this space with the weak C^0 -topology. Secondly, there exists a space $\text{Sol}(\mathcal{R}) \subset \Gamma^0(E)$ of **solutions** whose elements are those sections of E whose jet is a formal solution. We endow the space $\text{Sol}(\mathcal{R})$ with the weak C^1 -topology.

There is a natural inclusion,

$$\text{Sol}(\mathcal{R}) \hookrightarrow \text{FSol}(\mathcal{R}), \quad s \mapsto j^1 s,$$

which allows us to compare the space of solutions $\text{Sol}(\mathcal{R})$ to the space of formal solutions $\text{FSol}(\mathcal{R})$, which is more algebraic in nature. The theory of h-principles concerns itself with the question whether the above inclusion is a weak homotopy equivalence. If it is, we say that \mathcal{R} satisfies the **(complete) h-principle**. Gromov's method of flexible sheaves [14] shows that the h-principle holds for a large class of relations (open and Diff-invariant) on open manifolds. Partial and negative results have also been proven, such as [4, 12, 28].

Multiple variations of the complete h-principle exist. We mention two. We follow the convention that $\mathcal{O}p(C)$ denotes an unspecified arbitrarily small open neighborhood of C .

Definition 2.2. The h-principle for \mathcal{R} holds

- **relatively** if given a section $s \in \text{FSol}(\mathcal{R})$ such that $s|_C$ is holonomic, there exists a holonomic section $s' \in \text{Sol}(\mathcal{R})$, which is homotopic to s relative $\mathcal{O}p(C)$, for every closed set $C \subset M$.
- **parametrically** if given a continuous family $S : P \rightarrow \text{FSol}(\mathcal{R})$, there exists a continuous family $S' : P \rightarrow \text{Sol}(\mathcal{R})$ homotopic to S . Here P is a compact manifold acting as parameter space.

2.2. Simplicial complexes. In this section we discuss simplicial complexes and linear polyhedra. It is worth pointing out that many of the arguments in the paper are indeed linear in nature. Using them in the manifold setting will involve working chart by chart and taking into account various estimates to bound the non-linearity of the transition functions between them.

2.2.1. Simplicial complexes. We define the standard m -dimensional simplex $\Delta^m \subset \mathbb{R}^m$ as

$$\Delta^m = \{(t_1, \dots, t_m) \in \mathbb{R}^m \mid \sum_{i=1}^m t_i \leq 1 \text{ and } t_i \geq 0 \text{ for all } i\}.$$

A **linear simplex** is then any subset of Euclidean space that is affinely isomorphic to a standard simplex. By a **face** of a linear simplex we refer to a subsimplex of any dimension.

Simplices can be glued along their codimension-1 faces if they form a so-called simplicial complex:

Definition 2.3. A **simplicial complex** is a locally finite set K of linear simplices in an Euclidean space such that

- if $\sigma \in K$ then its face is also in K , and
- if $\sigma, \sigma' \in K$ then $\sigma \cap \sigma'$ is either empty or an element of K .

We denote by $K^{(\text{top})}$ the set of top dimensional simplices of a simplicial complex K , and can also define subcomplexes in an obvious manner.

A linear polyhedron is the topological space that is spanned by a simplicial complex. We introduce the word “linear” to make a distinction with polyhedra living in arbitrary manifolds, to be considered later.

Definition 2.4. A **linear polyhedron** P is the union of the linear simplices in a simplicial complex K . That is, $P = \cup_{\Delta \in K} \Delta$. In this case, we call K a **triangulation** of P and write $P = |K|$.

Due to local finiteness, our simplicial complexes will consist of countably many linear simplices. In particular, the corresponding polyhedra may be non-compact. Moreover, for our purposes it is enough to consider simplicial complexes in which the dimension of the simplices is bounded above. Furthermore, we will often work with simplicial complexes of **pure** dimension. That is, every simplex should be contained in a simplex of top dimension.

It proves useful to be able to refer to the maximal simplicial complex contained in a subset. Hence we introduce the following terminology.

Definition 2.5. Let K be a simplicial complex in \mathbb{R}^N and let $U \subset \mathbb{R}^N$ be a subset. We define $K \cap U$ as the **maximal simplicial subcomplex of K contained in U** . That is, $K \cap U$ consists of those $\Delta \in K$ such that $|\Delta| \subset U$.

2.2.2. *Adjacency.* The appropriate notion of a neighborhood of a linear simplex in a simplicial complex is that of a star, which is defined using the concept of adjacency. Related is the notion of a ring.

Definition 2.6. Consider a simplicial complex K including a linear simplex (or more generally, a subcomplex) Q .

- Two linear simplices are **adjacent** if they share a face.
- The **star** $\text{star } Q$ of Q is the set of all its adjacent simplices and their faces.
- The **closure** $\text{Cl}(A)$ of subset A of a simplicial complex is the smallest subcomplex containing the subset A .
- We define the **ring** around Q as $\text{ring } Q = \text{Cl}(\text{star}(Q) \setminus Q)$.

We write $\text{star}(Q, K)$ and $\text{ring}(Q, K)$ if we want to emphasize the simplicial complex we are working with. However, to simplify notation, we only indicate K once in expressions in which it occurs multiple times. For instance, we write $\text{ring}(\text{star}(Q, K))$ instead of $\text{ring}(\text{star}(Q, K), K)$. We also write star^n for the n -fold iteration of star.

2.2.3. *Subdivisions and isomorphisms.* The most restrictive notion of equivalence between simplicial complexes reads:

Definition 2.7. Two simplicial complexes K and K' are **isomorphic** if there exists a homeomorphism $F : |K| \rightarrow |K'|$, inducing a bijection between simplices, such that for each $\Delta \in K$ the restriction $F|_{\Delta}$ is linear.

A more “local” version for two simplicial complexes is the following:

Definition 2.8. Two simplicial complexes K and K' **agree** on a subset $Q \subset \mathbb{R}^N$ if there exist subcomplexes K_1 and K_2 of respectively K and K' such that

- K_1 and K_2 are isomorphic, and
- $Q = |K_1| = |K_2|$.

Alternatively, we can subdivide any simplicial complex to obtain a new simplicial complex triangulating the same polyhedron.

Definition 2.9. A **subdivision of a simplicial complex K** is a simplicial complex K' such that each $\Delta \in K$ satisfies $\Delta = \cup_{i \in I} \Delta'_i$ for a finite collection of $\Delta'_i \in K'$.

We remark that any two triangulations of a given polyhedron have a common subdivision.

2.2.4. *Piecewise linear maps.* Between polyhedra, the natural class of maps to consider is:

Definition 2.10. Let P be a linear polyhedron. A map $f : P \rightarrow \mathbb{R}^n$ is **piecewise linear** if:

- the map f is continuous, and
- the restriction of f to each simplex of K is affine, where K is some triangulation of P .

If we want to emphasize the role of K , we will write that (f, K) is a piecewise linear map or that f is piecewise linear with respect to K . We denote the set of piecewise linear maps by $C_{\text{PL}}(P, \mathbb{R}^n)$. We topologize it below.

We can thus speak of piecewise linear maps between linear polyhedra, yielding the category of linear polyhedra. Isomorphic simplicial complexes, as defined above, yield isomorphic polyhedra. Conversely, two polyhedra are isomorphic if they admit triangulations that are isomorphic.

2.2.5. *Piecewise smooth maps.* We also consider the more general class of maps:

Definition 2.11. Let P be a linear polyhedron and N a manifold. A map $f : P \rightarrow N$ is **piecewise smooth** if it is continuous and, for some triangulation K , the maps $f|_{\Delta}$ are smooth for all $\Delta \in K$.

The set of piecewise smooth maps will be denoted by $C_{\text{PS}}(P, N)$. We note that $C_{\text{PS}}(P, \mathbb{R}^n)$ contains $C_{\text{PL}}(P, \mathbb{R}^n)$. The set $C_{\text{PS}}(P, N)$ can readily be endowed with the C^0 -topology (either the weak or the strong version), by interpreting it as a subset of $C^0(P, N)$.

To define the C^r -topology in $C_{\text{PS}}(P, N)$, it is convenient to assume that P is of pure dimension. We fix an auxiliary triangulation K of P to make sense of $J^r(\Delta, N)$ for each top simplex $\Delta \in P$. On each $J^r(\Delta, N)$ we fix a metric, so that we can make sense of the C^r -metric on $C_{\text{PS}}(\Delta, N)$. Then we define:

Definition 2.12. Consider a pair of maps $f_1, f_2 \in C_{\text{PS}}(P, N)$, each piecewise smooth with respect to a triangulation K_i of P , for $i = 1, 2$. Consider a triangulation K' subdividing K_1, K_2 and K . We write $d_{C^r}(f_1, f_2) < \varepsilon$ if for every top dimensional simplex $\Delta \in K'$ we have $d_{C^r}(f_1|_{\Delta}, f_2|_{\Delta}) < \varepsilon$. Here we let ε be a positive constant if P is compact, and a positive function $P \rightarrow \mathbb{R}_+$ if P is not compact.

We observe that the above distance on $C_{\text{PS}}(P, N)$ depends on our choice of metrics on each $J^r(\Delta, N)$ and therefore it also depends on K . It does however not depend on the choice of K' . The underlying topologies (known as respectively the weak and strong) do not depend on any of these choices.

Analogous to Definition 1.1, we can now also define an ε -jiggling of a piecewise smooth map defined on a linear polyhedron.

Definition 2.13. Consider a pair of maps $f_1, f_2 \in C_{\text{PS}}(P, N)$, each compatible with a triangulation K_i of P , for $i = 1, 2$. Assume that the simplicial complex K_2 subdivides K_1 , and that $d_{C^1}(f_1, f_2) < \varepsilon$. Then the map (f_2, K_2) is an ε -**jiggling** of (f_1, K_1) .

2.2.6. *Jet spaces over linear polyhedra.* We can also set up the language of jet spaces and differential relations from Section 2.1 for linear polyhedra of pure dimension. We fix a finite simplicial complex K of pure dimension and consider piecewise smooth maps $|K| \rightarrow \mathbb{R}^n$. Having fixed K we can consider the jet spaces $J^r(\Delta, \mathbb{R}^n)$ for each $\Delta \in K^{(\text{top})}$. To keep notation light we will write $J^r(|K|, \mathbb{R}^n)$ for their union.

Similarly, in each $J^1(\Delta, \mathbb{R}^n)$ we can define a differential relation \mathcal{R}_{Δ} . We then write \mathcal{R} for their union. We refer to a relation \mathcal{R} as fiberwise dense and open, if each \mathcal{R}_{Δ} is. Whenever we subdivide K we can pullback these differential relations to the simplices in the subdivision.

2.2.7. *Ordered simplicial complexes.* Later on (Section 3) we will consider the crystalline subdivision method for simplicial complexes; it will be defined first for the standard simplex and then be generalized to arbitrary simplicial complexes. In order to do so, we will need to make use of an ordering of the vertices. As such, we introduce:

Definition 2.14. An **ordered simplicial complex** is a pair consisting of a simplicial complex and a total order in its set of vertices.

Observe that, in an ordered simplicial complex, each simplex can be uniquely parametrized by the standard simplex in a manner that is affine and order preserving (where the vertices of the standard simplex are ordered using the standard basis in \mathbb{R}^m).

2.2.8. *Colored simplicial complexes.* Our main construction will involve arguing over multiple simplices at once, but to do so we need them not to interact. This is possible (Section 5.2) as long as they are far apart, which we formalize by requiring that their stars are disjoint. To keep track of this we will assign a “color” to each simplex so that simplices in a star never have the same color.

For the following definition we use the notation $[C]$ with $C \in \mathbb{N}$ to denote the ordered set $\{0, \dots, C\}$.

Definition 2.15. Let K be a simplicial complex of pure dimension. A C -**coloring** of K is a map

$$c : K^{(\text{top})} \rightarrow [C]$$

such that $c(\Delta) \neq c(\Delta')$ whenever $\text{star}(\Delta) \cap \text{star}(\Delta') \neq \emptyset$.

We refer to C as the **size** of the coloring c . We write $K^{(\text{top}, i)} \subset K^{(\text{top})}$ for the top simplices of K of color i .

2.3. **Triangulations.** We now recall the notion of a triangulation of a manifold; i.e. a decomposition of the manifold into smooth simplices. This combinatorial description of the manifold is particularly useful for local arguments and, as we shall see, will turn our arguments into arguments about linear polyhedra. In this section we fix our notation and conventions on the topic. For a more detailed account of triangulations we refer to [33, Ch. IV.B] and [20].

2.3.1. *Polyhedra.* Instead of defining triangulations directly, we recall the more general notion:

Definition 2.16. A **polyhedron** $T : |K| \rightarrow M$ consists of a simplicial complex K and a family of smooth embeddings $(T_\Delta : \Delta \rightarrow M)_{\Delta \in K}$ that glue to a homeomorphism $T : |K| \rightarrow T(|K|)$.

Equivalently, a polyhedron is an embedding of a linear polyhedron $|K|$ into M that is piecewise smooth with respect to K . A subset $Q \subset M$ is a **subpolyhedron** of T if there exists a subcomplex K'' of a subdivision K' of K such that $T|_{|K''|} : |K''| \rightarrow Q$ is a homeomorphism.

Given a polyhedron $T : |K| \rightarrow M$ we define a **simplex** Δ' of T as the image of a linear simplex $\Delta \in K$ under T . With a slight abuse of notation we will write $\Delta' \in T$. Keeping K in mind, we still speak of faces, stars and rings.

Definition 2.17. A **triangulation** of a manifold M is a polyhedron $T : |K| \rightarrow M$ with $T(|K|) = M$.

It is a result of Whitehead [32] that any smooth manifold can be triangulated, uniquely up to subdivision. The name “Whitehead triangulation” is sometimes used to emphasize the compatibility between the triangulation and the smooth structure of M .

2.3.2. *Piecewise smooth sections.* Given a bundle over a triangulated manifold, we consider:

Definition 2.18. A section $s : M \rightarrow E$ of a fiber bundle E is **piecewise smooth** if there exists a triangulation $T : |K| \rightarrow M$ such that $s \circ T$ is piecewise smooth.

We denote the space of such sections by $\Gamma_{\text{PS}}(E)$. As before, if we want to emphasize the role of T , we may say that (s, T) is a piecewise smooth section.

To define the C^r -topology on $\Gamma_{\text{PS}}(E)$, we again need to fix a metric on the jet space $J^r(E)$. Compared to the setting of maps over linear polyhedra from Definition 2.12, we however do not need to fix a triangulation to make sense of jet bundles over M . Then we obtain:

Definition 2.19. Given a pair of sections $(s_1, T_1), (s_2, T_2) \in \Gamma_{\text{PS}}(E)$ we write $d_{C^r}(s_1, s_2) < \varepsilon$ if there exists a common subdivision $T : |K| \rightarrow M$ of T_1 and T_2 such that for every simplex $\Delta \in T$ we have $d_{C^r}(s_1|_\Delta, s_2|_\Delta) < \varepsilon$. Here we let ε be a positive constant if P is compact, and a positive function $P \rightarrow \mathbb{R}_+$ if P is not compact.

We point out that, even though the distances defined above will generally differ from those defined in Definition 2.12, the induced topologies on $\Gamma_{\text{PS}}(E)$ do coincide with those in Definition 2.12.

2.4. Simplicial sets. In this section we briefly recall the necessary theory of simplicial sets. We emphasize that this is a very short overview which does not give a general account of the theory, but only covers the minimum to reach what we need. In particular, we work our way up to the specific characterization of weak homotopy equivalences of simplicial sets appearing in Definition 2.25. This section is mostly based on [15, Ch. 2], to which we also refer for more details.

We start by defining the simplex category $\mathbf{\Delta}$ which has as objects $[n] = \{0, 1, \dots, n\}$. Using the natural ordering $0 \leq 1 \leq \dots \leq n$ we define morphisms in $\mathbf{\Delta}$ as non-decreasing functions. These are generated by the face maps $\delta^i : [n-1] \rightarrow [n]$ and the degeneracy maps $\sigma^i : [n+1] \rightarrow [n]$, which are defined by either skipping or doubling the element i , respectively. This allows us to define:

Definition 2.20. A **simplicial set** is a functor $X : \mathbf{\Delta}^{\text{op}} \rightarrow \mathbf{Set}$.

Unpacking the definition we see that a simplicial set X is completely determined by the sets $X_n := X([n])$, the face maps $\delta_i := X(\delta^i) : X_n \rightarrow X_{n-1}$ and the degeneracy maps $\sigma_i := X(\sigma^i) : X_n \rightarrow X_{n+1}$.

To every topological space we can associate a simplicial set by considering its singular complex. Simplicial sets still allow us to talk about weak homotopy type, but are often easier to work with than topological spaces, due to their combinatorial nature.

Definition 2.21. The functor $\text{Sing} : \mathbf{Top} \rightarrow \mathbf{sSet}$ is defined by $\text{Sing}(M)_n = \text{Hom}_{\mathbf{Top}}(\Delta^n, M)$ for all $M \in \mathbf{Top}$. We refer to $\text{Sing}(M)$ as the **singular complex** of M .

The geometric realization $|\cdot| : \mathbf{sSet} \rightarrow \mathbf{Top}$ functor is also relevant to our discussion. We do not want to define it precisely, but we do want to point out that it can be seen as the inverse of the singular complex in two ways. Firstly, it holds that there exists a bijection

$$\text{Hom}_{\mathbf{Top}}(|X|, M) \rightarrow \text{Hom}_{\mathbf{sSet}}(X, \text{Sing}(M))$$

which is natural in $X \in \mathbf{sSet}$ and $M \in \mathbf{Top}$. This can be phrased by saying that the singular complex is the right adjoint to the geometric realization. Secondly, we have the following statement:

Lemma 2.22. *Let $M \in \mathbf{Top}$, then there exists a natural map $|\text{Sing}(M)| \rightarrow M$ which is a weak homotopy equivalence. We denote this as $M \simeq |\text{Sing}(M)|$.*

2.4.1. Weak homotopy equivalences. In the category of topological spaces a weak homotopy equivalence can be defined as the vanishing of the relative homotopy groups. We can use a similar definition for simplicial sets, but for this we need simplicial analogues of the disc and its boundary. These roles are played by the standard n -simplex and its boundary. We use the notation $\Delta[n]$ for the standard n -simplex as a simplicial set, and the notation Δ^n as a topological space.

Definition 2.23. The **standard n -simplex** $\Delta[n]$ is the simplicial set

$$\Delta[n] = \text{Hom}_{\mathbf{\Delta}}(\cdot, [n]).$$

Its **boundary** $\partial\Delta[n]$ is defined by

$$(\partial\Delta[n])_m = \{f \in \text{Hom}_{\mathbf{\Delta}}([m], [n]) \mid f \text{ is not surjective}\}.$$

We have two observations to make concerning the above definition. First, note that $\Delta[n]$ and $\text{Sing}(\Delta^n)$ are not the same. One could show for instance that the latter is a so-called Kan complex whereas the former is not. The geometric realization of $\Delta[n]$ is however the standard simplex Δ^n , which explains the notation. Secondly, we note that the Yoneda lemma gives us a bijection $\text{Hom}_{\mathbf{sSet}}(\Delta[n], X) \rightarrow X_n$ for all $X \in \mathbf{sSet}$. This provides a useful interpretation of morphisms with domain $\Delta[n]$ and clarifies its role in the theory.

We can now make sense of homotopies in the category of simplicial sets (which strongly resembles the definition for topological spaces). In the following definition the product of two simplicial sets is just their product as functors. That is, $(X \times Y)_n = X_n \times Y_n$ for any given $X, Y \in \mathbf{sSet}$.

Definition 2.24. Two morphisms $f, g : X \rightarrow Y$ of simplicial sets are **homotopic** if there exists a map $\eta : X \times \Delta[1] \rightarrow Y$ such that the diagram

$$\begin{array}{ccccccc} X & \xrightarrow{\cong} & X \times \Delta[0] & \xrightarrow{\text{id} \times \delta^1} & X \times \Delta[1] & \xleftarrow{\text{id} \times \delta^0} & X \times \Delta[0] \xleftarrow{\cong} X \\ & & & \searrow f & \downarrow \eta & \swarrow g & \\ & & & & Y & & \end{array}$$

commutes.

The definition of weak homotopy equivalence for simplicial sets stated below is not the standard one but equivalent to it. It is based on a characterization of Jardine [17]. It generalizes the notion for spaces via the singular complex and the geometric realization functor.

Definition 2.25. A morphism $f : X \rightarrow Y$ of simplicial sets is a **weak homotopy equivalence** if and only if for all $n \in \mathbb{N}$ and all commutative diagrams

$$\begin{array}{ccc} \partial\Delta[n] & \longrightarrow & X \\ \downarrow & \nearrow \sigma & \downarrow f \\ \Delta[n] & \longrightarrow & Y, \end{array}$$

where $\partial\Delta[n] \rightarrow \Delta[n]$ is the standard inclusion, there exists a lift $\sigma : \Delta[n] \rightarrow X$ such that the upper triangle commutes and the lower triangle commutes up to homotopy constant along $\partial\Delta[n] \hookrightarrow \Delta[n]$.

3. SUBDIVISIONS

Various methods exist to subdivide a triangulation into smaller simplices, each with their own properties. We focus on a method called crystalline subdivision (Section 3.1), whose main benefit is that simplices do not get too distorted when subdividing (Section 3.2).

We then discuss how one covers a polyhedron with nice subpolyhedra; we need this to ultimately establish (in Section 6.1) a version of jiggling that is relative in the domain. The ideas needed for the case of compact polyhedra are explained in Section 3.3; they are based on the use of crystalline subdivision.

In the non-compact case one needs to introduce new subdivision schemes, localized over subpolyhedra. The scheme required for our main result (Theorem 6.3) can be implemented rather naively. We call it “barycentric cone off” and we explain it in Section 3.4. For other applications (e.g. general position) we need a scheme with better quantitative control over the shape of our simplices. We call it “generalized crystalline subdivision” and we describe it in Section 3.5.

3.1. Crystalline subdivision. There are various ways of defining crystalline subdivision, but here we follow the definition of Thurston from [30, p. 227].

Crystalline subdivision as described below is based on the observation that we know how to subdivide a cube into smaller cubes. We illustrate the procedure in Figure 1.

Definition 3.1. Applying **crystalline subdivision** to the standard m -simplex Δ^m with $m > 0$ is done using the following steps:

- (1) Include Δ^m into the standard m -cube I^m .
- (2) Subdivide I^m into 2^m smaller m -cubes of size $1/2$.
- (3) Subdivide each of the smaller m -cubes into $m!$ smaller linear m -simplices. Each linear simplex corresponds to a permutation π of $\{0, 1, \dots, m\}$ by identifying a permutation π with the subset $\{(x_1, \dots, x_m) \in \mathbb{R}^m \mid 0 \leq x_{\pi(0)} \leq x_{\pi(1)} \leq \dots \leq x_{\pi(m)} \leq 1\}$.

We define the crystalline subdivision of Δ^0 as itself.




FIGURE 1. The second crystalline subdivision of the standard 2-simplex Δ^2 .

This procedure makes use of the ordering of the vertices in step 1. Indeed, for $m \geq 3$, the resulting subdivision would be different if we applied an affine isomorphism that permutes the vertices.

This procedure can be iterated but it turns out to be easier to define finer crystalline subdivisions directly. Hence we define the ℓ th crystalline subdivision of the standard m -simplex Δ^m by instead subdividing the m -cube into $2^{\ell m}$ smaller m -cubes of size $2^{-\ell}$ in item 2 from Definition 3.1.

To generalize to a simplicial complex K , we choose an ordering on K such that the resulting subdivision is well-defined.

Definition 3.2. Let K be an ordered simplicial complex. Its ℓ th crystalline subdivision K_ℓ is obtained as follows: Given a simplex $\Delta \in K$, consider the unique order-preserving, affine identification ϕ between Δ and the standard simplex Δ^m . Use ϕ to transfer the ℓ th crystalline subdivision of Δ^m to Δ .

Given a polyhedron $T : |K| \rightarrow M$ associated to a specified simplicial complex K , we can also speak of its ℓ th crystalline subdivision T_ℓ , seen as the collection of embeddings $(T_\Delta : \Delta \rightarrow M)_{\Delta \in K_\ell}$ given by restricting T .

3.2. Properties of crystalline subdivision. We discuss some of the properties of crystalline subdivision next. They all have the same flavour: since crystalline does not distort simplices, we are able to obtain various quantitative bounds independent of the order of subdivision ℓ .

3.2.1. Coloring. The first advantage is that we are able to bound the maximum number of colors we need to color a simplicial complex and all its crystalline subdivisions.

Lemma 3.3. *Let K be a finite and ordered simplicial complex. Then, there exists a $C \in \mathbb{N}$ such that every crystalline subdivision K_ℓ admits a coloring of size C .*

Proof. Consider the standard m -cube I^m subdivided into smaller m -cubes of length $2^{-\ell}$ for some $\ell \in \mathbb{N}$. Let $Q \subset I^m$ be one of those smaller cubes. Then Q is adjacent to at most $3^m - 1$ cubes. In the procedure of crystalline subdivision each of these cubes is subdivided into $m!$ smaller m -simplices. If A is the number of simplices in $K^{(\text{top})}$, then each simplex $\Delta \in K_\ell^{(\text{top})}$ is adjacent to at most $(3^m - 1) \cdot (m)! \cdot A$ simplices, which means that there exists a $((3^m - 1) \cdot (m)! \cdot A)$ -coloring of K_ℓ . \square

3.2.2. Model simplices. A second property of crystalline subdivision is that the resulting simplices are equivalent to the standard simplex, up to scaling and translation. Model simplices appear already in Thurston's work [30], albeit in a different formulation.

Lemma 3.4. *Let K be a finite and ordered simplicial complex in \mathbb{R}^N of pure dimension. Then, there exists a finite collection of **model simplices** $\mathcal{C} = \{\Delta_i \subset \mathbb{R}^N \mid i = 0, \dots, I\}$ with the following property: for any $\ell \in \mathbb{N}$ and $\Delta \in K_\ell^{(\text{top})}$ there exists*

- a model simplex $\Delta_i \in \mathcal{C}$, and

- a map $t : \mathbb{R}^N \rightarrow \mathbb{R}^N$ which is a composition of a translation and a scaling by 2^ℓ

such that $\Delta_i = t(\Delta)$.

Proof. Let $\Delta \in K^{(\text{top})}$ be a top dimensional simplex, then there exists an affine isomorphism $\phi : \Delta \rightarrow \Delta^m \subset \mathbb{R}^m$, where m is the dimension of Δ . We define \mathcal{C}'_Δ to be the first crystalline subdivision of Δ^m . We note that any simplex in $(\Delta^m)_\ell$ is by construction a translated and scaled copy of a unique simplex in $(\Delta^m)_1$. Using ϕ we now push forward \mathcal{C}'_Δ to a collection of model simplices \mathcal{C}_Δ . We define \mathcal{C} to be the union over all \mathcal{C}_Δ for $\Delta \in K^{(\text{top})}$. \square

3.2.3. Shape of simplices. Given a simplex, we are interested in the maximal (r_{\max}) and minimal (r_{\min}) distances between its vertices. The former quantity controls for instance how well a map is approximated by its linearization with respect to a given triangulation (see Section 4). Their ratio on the other hand tells us how degenerate the simplex is. We define these quantities for $(m+1)$ -tuples of points in \mathbb{R}^N , but we can also speak of r_{\max} and r_{\min} of a linear simplex Δ by identifying Δ with its vertices.

Definition 3.5. The continuous functions $r_{\min}, r_{\max} : \mathbb{R}^N \times \cdots \times \mathbb{R}^N = \mathbb{R}^{N(m+1)} \rightarrow \mathbb{R}$ are defined by

$$\begin{aligned} r_{\min}(p_0, \dots, p_m) &= \min_{i,j \in [m]} d(p_i, p_j) \quad \text{and} \\ r_{\max}(p_0, \dots, p_m) &= \max_{i,j \in [m]} d(p_i, p_j). \end{aligned}$$

A priori we cannot bound these quantities and their ratio among all subdivisions of a linear polyhedron, since the space of all linear simplices in \mathbb{R}^N is not compact. However, due to the existence of the model simplices (from Lemma 3.4), we can bound these quantities when considering crystalline subdivisions:

Lemma 3.6. *Let K be simplicial complex that is ordered and finite. Then, there exists $C, D \in \mathbb{R}_+$ such that for all $\ell \in \mathbb{N}$:*

$$\max_{\Delta \in K_\ell} r_{\max}(\Delta) = C \cdot 2^{-\ell} \quad \text{and} \quad \frac{\max_{\Delta \in K_\ell} r_{\max}(\Delta)}{\min_{\Delta \in K_\ell} r_{\min}(\Delta)} = D.$$

3.3. Nice covers of polyhedra. Our arguments often have to be localized to subpolyhedra. To this end, it is important for us to be able to cover a given polyhedron by subpolyhedra that are nice. We explain how to do this now.

3.3.1. Nice subcomplexes. Before we get to the key definition, recall that the join between two subsets A, B of \mathbb{R}^N is the set

$$\text{join}(A, B) = \{ta + (1-t)b \mid a \in A, b \in B \text{ and } t \in [0, 1]\}.$$

In particular, when we take the join of a (suitable) pair of linear simplices Δ_1, Δ_2 we end up with a higher-dimensional simplex $\Delta := \text{join}(\Delta_1, \Delta_2)$ having the two original simplices as opposing faces.

Definition 3.7. Let K be a simplicial complex and let K' be a subcomplex. We will say that K' is **nice** if for each simplex $\Delta \in \text{star}(K')$ the subcomplex $\Delta \cap K'$ is a face of Δ .

The meaning of this is that any Δ in the ring of K' can be thus seen as the join of two faces, A and B , with $A \in K'$ and B disjoint from K' . This is useful in order to interpolate from a map/section given over A to a map/section given over B (as in Section 4.2). The following provides a useful criterion for niceness.

Lemma 3.8. *The following properties are equivalent for a subcomplex $K' \subset K$:*

- K' is nice.
- Every $\Delta \in K$ that equals the convex hull of $\Delta \cap K'$ is an element of K' .




FIGURE 2. We illustrate here subcomplexes K' and K'' of a simplicial complex K , where K' is not nice and K'' is.

Proof. Suppose K' is nice. Let $\Delta \in K$ equal the convex hull of $\Delta \cap K'$. Then $\Delta \in \text{star}(K')$ and hence by assumption $\Delta \cap K'$ is a face of Δ . Since a face is convex, it follows that $\Delta = \Delta \cap K'$ and hence $\Delta \in K'$.

Conversely, given $\Delta \in \text{star}(K')$ we consider the vertices in $\Delta \cap K'$. They span a face F of Δ , which a priori does not need to be an element of K' . For niceness, it suffices to show that $\Delta \cap K' = F$. Note that F is convex and therefore by assumption is an element of K' . Hence $F \subset \Delta \cap K'$ and since we cannot add any extra vertex in $\Delta \cap K'$ to F , we have the desired equality. \square

3.3.2. Nice covers via convexity. The question now is how to cover a simplicial complex by nice subcomplexes. To achieve this we exploit the usual notion of convexity in Euclidean space. We recall from Definition 2.5 that $K \cap U$ is the maximal simplicial subcomplex of K contained in a given subset U .

Lemma 3.9. *Let K be a simplicial complex in \mathbb{R}^N and let $U \subset \mathbb{R}^N$ be a convex subset. Then the intersection $K \cap U$ is a nice subcomplex.*

Proof. We check the second condition of Lemma 3.8. Let $\Delta \in K$ equal the convex hull of $\Delta \cap (K \cap U)$. Then the vertices of Δ are in $K \cap U$ and hence also its span Δ . \square

Naively, we could now consider locally finite covers $\{U_i\}$ of \mathbb{R}^N by convex opens and look at the nice subcomplexes $\{K \cap U_i\}$. The issue is that these may not cover K if the top simplices of K are too large compared to the chosen opens. However, there is an easy fix:

Lemma 3.10. *Let K be a finite simplicial complex in \mathbb{R}^N . Let $\{U_i\}$ be a locally finite cover of \mathbb{R}^N by convex opens. Then, $\{K_\ell \cap U_i\}$ is a cover of K_ℓ by nice subcomplexes, if ℓ is large enough.*

Proof. Since K is finite, the set $|K|$ is compact and hence there exists some strictly positive Lebesgue's number δ associated to the cover $\{U_i \cap |K|\}$. We now take ℓ large enough such that $r_{\max}(\Delta) < \delta$ for every $\Delta \in K_\ell$. \square

3.4. Barycentric cone off. In light of Lemma 3.10, the question remains how to handle simplicial complexes that are not finite. Indeed, without finiteness, we have no bound on the number of subdivisions ℓ required to construct a cover by nice subcomplexes.

To address this, we must work relatively. Concretely, we need a scheme capable of subdividing the simplicial complex K over a given region, while keeping it the same elsewhere. This will allow us to keep subdividing K as we deal with an infinite cover of $|K|$. It turns out that we can do this rather naively:

Definition 3.11. Let $\Delta \subset \mathbb{R}^N$ be a linear simplex. Suppose we are given a subdivision S of its boundary. The **barycentric cone off** of Δ with respect to S is obtained by taking the join of the barycenter of Δ with S .

Hence given a (nice) subcomplex K' of K and a subdivision K'' of K' , we can apply the barycentric cone off in each simplex in the ring around K' . This results in a subdivision of the whole of K that leaves the complement of $\text{star}(K')$ unchanged. This is depicted in Figure 3.




FIGURE 3. The subcomplex $K \setminus \text{star}(K')$ of K is subdivided in a crystalline manner, but K' not. Hence we apply the barycentric cone off within $\text{ring}(K')$ with respect to an arbitrary crystalline subdivision of $K \setminus \text{star}(K')$.

3.5. Generalized crystalline subdivision. The barycentric coning off procedure described in Definition 3.11 is enough for us to establish our main jiggling result, also in the case of non-compact manifolds (Section 6.2). However, it turns out not to be well suited to achieving general position (as in Thurston's jiggling lemma) over non-compact manifolds. The reason is that for this application we need some control (in terms of model simplices) as we keep coning off.

Hence we now present a way of coning off where model simplices are preserved, in the sense that we can construct model simplices for a subcomplex $K' \subset K$ before knowing what crystalline subdivision we want outside of K' . However, this means we do need a region in which we cone off, which in general will be larger than just a ring. We illustrate the construction in Figure 4.

For the following statement, to ease the notation, we will write $B(A, r)$ for the r -neighborhood of a subset $A \subset \mathbb{R}^N$.




FIGURE 4. A generalized crystalline subdivision of a polyhedron K . We have indicated below the crystalline subdivisions of K and in-between we use the barycentric cone off which we have indicated in a lighter color.

Proposition 3.12. *Let K be a simplicial complex in \mathbb{R}^N , let $\delta > 0$ be given and let $A \subset \mathbb{R}^n$ be a compact subset. Then, there exists an $L \in \mathbb{N}$ such that for all $\ell_0 \geq L$ there exist*

- a finite collection \mathcal{C} of model simplices, and
- a natural number $C \in \mathbb{N}$,

such that for all $\ell_1 \geq \ell_0$ there exists a subdivision K' of K satisfying

- K' agrees with K_{ℓ_0} on A ,
- K' agrees with K_{ℓ_1} on $\mathbb{R}^N \setminus B(A, \delta)$,
- the simplices of $K' \cap B(A, \delta)$ are modeled by \mathcal{C} , that is, they equal a model simplex in \mathcal{C} up to translation and scaling by 2^ℓ where $\ell \in [\ell_0, \ell_1]$, and
- the subcomplex $K' \cap B(A, \delta)$ can be colored with at most C colors.

Proof. Constructing the control data. The results in Section 3.2 yield a maximum number of colors C' and a finite collection of model simplices \mathcal{C}' for all crystalline subdivisions of the subcomplex of K consisting of simplices with non-empty intersection with $B(A, \delta)$.

We enlarge the collection of model simplices \mathcal{C}' to account for the barycentric cone off of Definition 3.11 as follows: Given a model simplex $\Delta \in \mathcal{C}'$, we apply crystalline subdivision to subdivide one of its faces. Now using the barycentric cone off, we obtain a subdivision of Δ and we add the resulting simplices to \mathcal{C}' . We repeat these steps for each face of Δ and each model simplex $\Delta \in \mathcal{C}'$. This defines a finite collection \mathcal{C} of model simplices.

The construction. Choose L to be large enough such that the simplices of $K_L \cap B(A, \delta)$ have a maximal radius of at most $s = \delta/4$. Let natural numbers ℓ_1 and ℓ_0 be given such that $\ell_1 \geq \ell_0 \geq L$.

Define $K^0 := K$ and let $B^{(i)}$ denote a neighborhood of A of size $\sum_{j=0}^i 2^{-j}s$. Then by induction we define a sequence of subdivisions $K^{(i)}$ as follows. Assume we have already constructed $K^{(i)}$, then we define its subdivision $K^{(i+1)}$ as the simplicial complex that

- agrees with $K^{(i)} \cap B^{(i)}$,
- is the barycentric cone off on the ring of $K^{(i)} \cap B^{(i)}$, and
- agrees with K_{ℓ_0+i+1} outside of it

We note that the double star of $K^{(i)} \cap B^{(i)}$ is contained in $B^{(i+1)}$.

We define $K' := K^{(\ell_1 - \ell_0)}$. We see that the coning off takes place in $B^{(\ell_1 - \ell_0)}$, which is contained in neighborhood of A of radius

$$\sum_{j=0}^{\ell_1 - \ell_0} 2^{-j}s < \sum_{j=0}^{\infty} 2^{-j}s = 2s < \delta.$$

Wrapping up the proof. The claimed properties follow from the fact that the simplices in K' are either simplices in K_ℓ for some $\ell \in [\ell_0, \ell_1]$, or are obtained from such a simplex by a single barycentric cone off. Hence they are indeed modeled by an element of \mathcal{C} .

We bound the maximum number of colors C as follows. Each simplex is involved in at most one coning off process. Hence we can take $C = C' \cdot A$, where we defined C' at the beginning of this proof, and A is the resulting number of top-dimensional subsimplices after applying the barycentric cone off to a simplex Δ . We can bound A by $m - 1 + 2^{m-1}$, where m is the dimension of Δ . To see this, let F be the face of Δ subdivided once in a crystalline manner. Then the subdivision of F contains 2^{m-1} simplices, which are each coned off to the barycenter. All the other faces of Δ , of which there are $m - 1$, are also coned off to the barycenter. \square

We note that by the existence of the model simplices we get bounds on the minimal and maximal distance and their ratios, similar to Lemma 3.6.

We will refer to a subdivision K' of K as a **generalized crystalline subdivision**, if each simplex in K' is an element of a crystalline subdivision of K or is obtained from such a simplex by a single barycentric cone off where the faces are subdivided at most once in a crystalline manner. The subdivision K' from Proposition 3.12 is in particular an example, since each $K^{(i)} \cap B^{(i)}$ from the proof is a nice subcomplex. Hence we refer again to Figure 4 for an example.

4. LINEARIZATION OF PIECEWISE MAPS

In this section we focus on the following notion:

Definition 4.1. Let K be a simplicial complex and let $|K|$ be the corresponding linear polyhedron. Let $s : |K| \rightarrow \mathbb{R}^n$ be a map that is piecewise smooth with respect to K . The **linearization** $s^{\text{lin}} : |K| \rightarrow \mathbb{R}^n$ of s with respect to K is the unique piecewise linear map such that s^{lin} and s agree on the vertices of K .

We observe that s and s^{lin} are homotopic through piecewise smooth maps, thanks to linear interpolation.

We now motivate this definition. We recall that our ultimate goal is to study sections of bundles E over triangulated manifolds $T : |K| \rightarrow M$. In this setting we will not be able to speak of linear maps into E . Nonetheless, we can consider linear maps with respect to specified fibered charts, but the transition functions between them will introduce non-linearity. Hence our strategy relies on the following observations:

- Linearizing a section can be done in a C^1 -small manner if we subdivide K enough.
- Linearizing changes higher derivatives drastically, but this is of no consequence to us, since we are studying first order differential relations.
- Once we have linearized, the jiggling argument is purely linear in nature.

As such, every time we move to a new fibered chart, we will linearize our data with respect to it.

4.1. The linearization statement. Fix a simplicial complex K of pure dimension. Let $s : |K| \rightarrow \mathbb{R}^n$ be a map that is piecewise smooth with respect to K . We introduce the notation s_ℓ^{lin} for the linearization of s with respect to the ℓ th crystalline subdivision K_ℓ . Then:

Lemma 4.2. *Let $s : |K| \rightarrow \mathbb{R}^n$ be a piecewise smooth map with respect to a finite simplicial complex K of pure dimension. Then, as $\ell \rightarrow \infty$ we have:*

- $d_{C^0}(s, s_\ell^{\text{lin}}) = O(2^{-2\ell})$, and
- $d_{C^1}(s, s_\ell^{\text{lin}}) = O(2^{-\ell})$.

We recall from Section 2.2.5 that the C^0 and C^1 -distances used here are independent of ℓ ; they are computed using a fixed collection of metrics on jet spaces over the simplices of K .

Proof. It suffices to prove the claim in the case where K consists of one top-dimensional linear simplex, because we can repeat the following argument for every simplex, invoking finiteness of K . Moreover, after choosing an ordering on K , we can assume that we are working with the standard simplex $\Delta^m \subset \mathbb{R}^m$ up to an affine change of coordinates. We write $\text{Hom}(\mathbb{R}^m, \mathbb{R}^n)$ for the space of linear maps $\mathbb{R}^m \rightarrow \mathbb{R}^n$ and we endow it with its standard Euclidean distance.

Consider an m -simplex $\Delta \in (\Delta^m)_\ell$. Let $\mathcal{D} \subset \text{Hom}(\mathbb{R}^m, \mathbb{R}^n)$ be the subspace of all derivatives $d_y s : \mathbb{R}^m \rightarrow \mathbb{R}^n$ with $y \in \Delta$. Then by Taylor's approximation formula, it holds that the maximum distance d_{\max} between two elements in \mathcal{D} is $O(r_{\max}(\Delta))$. This maximum exists because Δ and hence \mathcal{D} is compact.

We also fix a vertex $v \in \Delta$ and $e_1, \dots, e_m \in \mathbb{R}^m$ such that the other vertices of Δ are $\{v + e_i\}_i$. Then $s_\ell^{\text{lin}}|_\Delta$ is the map that sends $x \in \Delta$ to $s(v) + A(x - v)$ where A is the matrix with columns $s(v + e_i) - s(v)$. By the mean value theorem there exist elements $y_{ij} \in \Delta$ such that $A_{ij} = (d_{y_{ij}} s)_j e_i$. Hence we see that for the j th row of A we obtain $A_j x = (d_v s_j) x + O(d_{\max} |x|)$ for any $x \in \mathbb{R}^m$.

For the C^0 -distance between s_ℓ^{lin} and s we see that for their j th components we have

$$\begin{aligned} d_{C^0}(s_j(x), (s_\ell^{\text{lin}})_j(x)) &= |s_j(x) - s_j(v) - A_j(x - v)| \\ &\leq |s_j(x) - s_j(v) - (d_v s_j)(x - v)| + O(d_{\max} |x - v|) \end{aligned}$$

which we bound as $O(r_{\max}^2(\Delta))$, applying Taylor's approximation to the first term.

For the C^1 -distance, we consider the matrices $d_x s$ and $d_x s_\ell^{\text{lin}}$ for some $x \in \Delta$, where we point out that the latter is in fact the matrix A from earlier. The difference between their (ij) th elements is $(d_x s_j - d_{y_{ij}} s_j)e_i$, which we bound using Taylor's approximation by $O(|x - y_{ij}|) = O(r_{\max}(\Delta))$.

We now conclude the proof by recalling from Lemma 3.6 that $r_{\max}(\Delta) = O(2^{-\ell})$. \square

We observe that the proof presented does not rely specifically on the fact that we are subdividing in a crystalline fashion. The only property it uses is that the diameter r_{\max} of the simplices goes to zero as ℓ goes to infinity from Lemma 3.6.

4.2. Modifying maps on a face. As explained above, we linearize in order to work locally in a linear manner. This means that we need to explain how our local arguments globalize. To this end, we introduce an interpolation procedure between sections.

Definition 4.3. Let Δ be a linear simplex given as the join of two opposing faces A and B . Let $t : \Delta \rightarrow [0, 1]$ be the join parameter: the affine function that is 0 over A and 1 over B . Suppose that $s_A, s_B : \Delta \rightarrow \mathbb{R}^n$ are two smooth maps. We define the **interpolation** of s_A and s_B over Δ with respect to A and B , denoted as $\text{interpolate}_\Delta(A, B, s_A, s_B)$, to be the map $ts_A + (1 - t)s_B$.

The result of interpolation between sections is controlled by the shape of the simplex and the difference between the two sections that serve as input:

Lemma 4.4. *Fix a linear m -simplex Δ spanned by two opposing faces A, B . Consider moreover three maps $u_1, u_2 : A \rightarrow \mathbb{R}^n, v : B \rightarrow \mathbb{R}^n$, and the corresponding interpolations $s_i = \text{interpolate}_\Delta(A, B, u_i, v)$. Then the following bounds hold:*

- $d_{C^0}(s_1, s_2) = O(d_{C^0}(u_1, u_2))$, and
- $d_{C^1}(s_1, s_2) = d_{C^0}(s_1, s_2) + O\left(\frac{d_{C^0}(u_1, u_2)}{r_{\min}(\Delta)} + d_{C^1}(u_1, u_2)\right)$.

Here we can use the usual Euclidean C^0 and C^1 -distances, although any other choices are equivalent up to a constant.

Proof. The statement regarding the C^0 -distance is clear. For the C^1 -claim we argue as follows: The join parameter $t : \Delta \rightarrow [0, 1]$ is a smooth function whose partial derivatives are bounded above by $1/r_{\min}(\Delta)$. The interpolation sections s_i can be written as $tu_i + (1 - t)v$. Hence we obtain:

$$\begin{aligned} |\partial_j(s_1 - s_2)| &= |\partial_j(t \cdot u_1 - t \cdot u_2)| \\ &\leq |\partial_j(t)| |u_1 - u_2| + |t| |\partial_j(u_1 - u_2)| \\ &\leq \frac{1}{r_{\min}(\Delta)} d_{C^0}(u_1, u_2) + d_{C^1}(u_1, u_2). \end{aligned} \quad \square$$

4.2.1. The linear case. We note that when we interpolate sections that are linear over a simplex Δ , the result is in general not linear. Hence we define the following variation of Definition 4.3, which only depends on the sections restricted to the faces.

Definition 4.5. Let Δ be a linear simplex given as the join of two opposing faces A and B . Suppose that $s_A : A \rightarrow \mathbb{R}^n$ and $s_B : B \rightarrow \mathbb{R}^n$ are two affine maps. We define the **join** of s_A and s_B over Δ , denoted as $\text{join}_\Delta(A, B, s_A, s_B)$, to be the unique affine map $\Delta \rightarrow \mathbb{R}^n$ agreeing with s_A over A and agreeing with s_B over B .

For the join we now obtain slightly simpler bounds when perturbing, than for the interpolation in Lemma 4.4:

Lemma 4.6. *Fix a linear m -simplex Δ spanned by two opposing faces A, B . Consider moreover three linear maps $u_1, u_2 : A \rightarrow \mathbb{R}^n, v : B \rightarrow \mathbb{R}^n$, and the corresponding joins $s_i = \text{join}_\Delta(A, B, v, u_i)$. Then the following bounds hold:*

- $d_{C^0}(s_1, s_2) = O(d_{C^0}(u_1, u_2))$, and
- $d_{C^1}(s_1, s_2) = d_{C^0}(s_1, s_2) + O\left(\frac{d_{C^0}(u_1, u_2)}{r_{\min}(\Delta)}\right)$.

Proof. The statement regarding the C^0 -distance is again clear, so we only argue the C^1 -claim. We label the vertices of Δ by v_0, \dots, v_m , and assume without loss of generality that $v_0 = 0$. Additionally we assume that A is spanned by v_0, \dots, v_ℓ for some $\ell < m$.

We define affine sections $\tilde{s}_i : \Delta^m \rightarrow \mathbb{R}^n$ by requiring that $\tilde{s}_i(0) = s_i(0)$ and $\tilde{s}_i(e_i) = s_i(v_i)$, where the vectors e_i are the unit vectors in \mathbb{R}^m . If we let T be the linear transformation $\Delta \rightarrow \Delta^m$ sending v_i to e_i , we see that $s_i = \tilde{s}_i \circ T$. We also observe that the operator norm of T is $\|T\| = \frac{1}{r_{\min}(\Delta)}$.

We denote the derivative of \tilde{s}_i by the matrix D_i . The j th column of D_i equals $v(v_j) - v(0)$ if $j \in \{1, \dots, \ell\}$ and $u_i(v_j) - v(0)$ if $j \in \{\ell + 1, \dots, m\}$. The j th column of the matrix $D_1 - D_2$ consists of zero for the first ℓ columns, and $u_1(v_j) - u_2(v_j)$ if $j \in \{\ell + 1, \dots, m\}$. Hence the difference $|D_1 - D_2|$ can be bounded as $O(d_{C^0}(u_1, u_2))$.

We now revert our attention back to the s_i and obtain:

$$d_{C^0}(ds_1, ds_2) = \|D_1 T - D_2 T\| \leq \|D_1 - D_2\| \|T\| = O\left(d_{C^0}(u_1, u_2) \cdot \frac{1}{r_{\min}(\Delta)}\right). \quad \square$$

4.3. Relative linearization. We now put together Lemmas 4.2 and 4.4 in order to yield a linearization statement that applies over a subcomplex, relative to the complement of small neighborhood. We point out that in the proof below, it is the niceness of the subcomplexes (recall Definition 3.7), that allows us on the ring of the nice subcomplexes to interpolate between sections.

Corollary 4.7. *Let K be a simplicial complex in \mathbb{R}^N . Let $s : |K| \rightarrow \mathbb{R}^n$ be a map that is piecewise smooth with respect to K . Let $K' \subset K$ be a finite subcomplex and consider a neighborhood $\mathcal{O}p(|K'|)$.*

Then, for each sufficiently large ℓ there exists a map $s'_\ell : |K| \rightarrow \mathbb{R}^n$ that is piecewise smooth with respect to K_ℓ and moreover:

- s'_ℓ is piecewise linear over $|K'|$,
- $s'_\ell = s$ outside of $\mathcal{O}p(|K'|)$,
- $d_{C^0}(s'_\ell, s) = O(2^{-2\ell})$, and
- $d_{C^1}(s'_\ell, s) = O(2^{-\ell})$.

Proof. Given an open neighborhood $\mathcal{O}p(|K'|)$ we can find a finite collection of convex opens $\{U_i\}_{i=1}^I$ in \mathbb{R}^N so that $\cup_i U_i$ intersected with $|K|$ is a neighborhood of $|K'|$ contained in $\mathcal{O}p(|K'|)$. We fix such a collection and assume that ℓ is large enough so that each simplex in K_ℓ is either contained in one of the U_i or is disjoint from $|K'|$. By Lemma 3.9 it follows that each subcomplex $K^i := U_i \cap K$ is nice.

We now construct maps s_i by induction on i which each satisfy the requirements of the statement on $\cup_{j=0}^i U_j \cap K'$ instead of the whole of K' . We set $s_0 := s$, so that the base case $i = 1$ follows the same proof as the inductive step. Hence assume s_{i-1} is constructed. Over K^i we use Lemma 4.2 to linearize s_{i-1} , yielding some map \bar{s}_i . Over the ring of K^i we apply Lemma 4.4 to interpolate between \bar{s}_i and s_{i-1} ; this is what we call s_i . We define s'_ℓ as the end result of the induction, that is $s'_\ell = s_I$.

We observe that the map s'_ℓ is piecewise smooth: Smoothness on each simplex is clear from the interpolation formula. Continuity over K follows because the join parameter t is continuous over the star of each K^i and the interpolation over two adjacent simplices agrees on their intersection.

To conclude we observe that, at each stage i , the claimed bounds hold for s_i by combining Lemma 4.2 (on K^j) and Lemma 4.4 (on $\text{star}(K^i)$). As such, the bounds apply to s'_ℓ , completing the proof. \square

We will refer to s'_ℓ from Corollary 4.7 as a linearization of s over $|K'|$.

5. SLOPE PERTURBATIONS OF PIECEWISE LINEAR MAPS

Let $E \rightarrow M$ be a fiber bundle over a smooth triangulated manifold M . We are going to explain next the idea of perturbing a section $s : M \rightarrow E$ by tilting it slightly simplex by simplex. Doing this with some quantitative control is the other main ingredient behind jiggling (the first ingredient being crystalline subdivision).

First, in Section 5.1, we discuss how to approximate a 1-jet $\sigma \in \pi_f^{-1}(x) \subset J^1(E)$ by a locally defined holonomic solution of a given relation $\mathcal{R} \subset J^1(E)$, where we recall that $\pi_f : J^1(E) \rightarrow E$ is the front projection. In order to do this, we need the relation to be fiberwise dense (Definition 1.2) and open.

This idea is then applied in Section 5.2 to construct a solution over an individual simplex. This solution must then be joined to the unperturbed map away from the simplex. This will allow us to introduce such tilts colorwise, in an estimated manner.

Even though our ultimate goal is to make statements about sections of bundles over manifolds, we continue working (mostly) in the linear setting. We will explain in detail in Section 6 how one makes the transition to sections defined on manifolds, but the main idea is to work “chart by chart”.

5.1. Linear extensions of jets. To approximate a 1-jet $\sigma \in \pi_f^{-1}(x) \subset J^1(E)$ by a local solution of \mathcal{R} , we use a two step process. We first change the slope of σ such that it lies in the relation, then we extend it to a holonomic section of \mathcal{R} . We emphasize that in the first step we do not change the zeroth order information of the jet, only its slope, so that the jet remains in the fiber $\pi_f^{-1}(x)$.

5.1.1. Density of solution jets. We now fix metrics on M , E , and $J^1(E)$. We use the notation $B_f(x, r)$ for a ball around $x \in J^1(E)$ of radius r intersected with the π_f -fiber through x , that is $B_f(x, r) = B(x, r) \cap \pi_f^{-1}(\pi_f(x))$. The two requirements we pose on \mathcal{R} immediately imply the following, which realizes the first step:

Lemma 5.1. *Let $\mathcal{R} \subset J^1(E)$ be an open and fiberwise dense relation. Then, for all $\varepsilon > 0$ and $\sigma \in J^1(E)$, there exists $\sigma' \in B_f(\sigma, \varepsilon)$ and $\delta > 0$ such that $B(\sigma', \delta) \subset \mathcal{R}$.*

In fact, if we restrict ourselves to σ in a compact set and use that δ depends continuously on σ , we see that δ can be chosen such that it does no longer depend on σ . This leads to the following:

Corollary 5.2. *Let $\mathcal{R} \subset J^1(E)$ be an open and fiberwise dense relation and let $A \subset J^1(E)$ be a compact subset. For all $\varepsilon > 0$ there exists $\delta > 0$ such that: For every $\sigma \in A$, there exists $\sigma' \in B_f(\sigma, \varepsilon)$ with $B(\sigma', \delta) \subset \mathcal{R}$.*

5.1.2. Density of linear extensions. To extend the jets in $B(\sigma', \delta)$ to holonomic sections of \mathcal{R} , we make use of linear extensions. In order to make sense of linear, we have to assume here that $M = \mathbb{R}^m$ and E is the trivial bundle $\underline{\mathbb{R}^n}$.

Definition 5.3. Let $U \subset \mathbb{R}^m$ be a subset. We say that $s : U \rightarrow \mathbb{R}^n$ is the **linear extension** of $\sigma \in J^1(\mathbb{R}^m, \mathbb{R}^n)$ if $j^1 s(\pi(\sigma)) = \sigma$ and s is linear. We write in this case $s = \text{LinExt}(\sigma)$ or, when we want to emphasize the domain, $s = \text{LinExt}_U(\sigma)$.

With this definition we see that Lemma 5.1 tells us that linear extensions of the jets in $B(\sigma', \delta)$ are actually local solutions. Here we write π for the projection $J^1(\mathbb{R}^m, \mathbb{R}^n) \rightarrow \mathbb{R}^m$.

Corollary 5.4. *Let $\mathcal{R} \subset J^1(\mathbb{R}^m, \mathbb{R}^n)$ be an open and fiberwise dense relation, and let $\varepsilon > 0$ and $\sigma \in J^1(\mathbb{R}^m, \mathbb{R}^n)$ be given. Then there exists $\sigma' \in B_f(\sigma, \varepsilon)$ and $\delta > 0$ such that for all $\sigma'' \in B(\sigma', \delta)$ we have that $\text{LinExt}(\sigma'')$ is a solution of \mathcal{R} over $B(\pi(\sigma''), \delta) \subset \mathbb{R}^m$.*

As in Corollary 5.2, we now obtain:

Corollary 5.5. *Let $\mathcal{R} \subset J^1(\mathbb{R}^m, \mathbb{R}^n)$ be an open and fiberwise dense relation, let $\varepsilon > 0$ be given and consider a compact subset $A \subset J^1(\mathbb{R}^m, \mathbb{R}^n)$. Then there exists $\delta > 0$ such that: for all $\sigma \in A$ there exists $\sigma' \in B_f(\sigma, \varepsilon)$ such that for all $\sigma'' \in B(\sigma', \delta)$ we have that $\text{LinExt}(\sigma'')$ is a solution of \mathcal{R} over $B(\pi(\sigma''), \delta)$.*

5.2. Slope-perturbing. We now use the linear extensions of the previous subsection to tilt maps in a piecewise manner. The upcoming statements are quantitative and use the standard C^r -distance given by the ambient Euclidean space the simplices (or simplicial complexes) live in. By compactness of simplices, this is equivalent to any other C^r -distance we could choose.

5.2.1. Over a single simplex. Consider a linear simplex Δ . The reasoning from the previous subsection states that, given $\varepsilon > 0$ and a 1-jet σ_1 , we can choose $\sigma_2 \in B_f(\sigma_1, \varepsilon)$ and consider its linear extension s_1 over Δ . Naturally, if we had chosen some other $\sigma_2 \in B_f(\sigma, \varepsilon)$ instead of σ_1 , we would have obtained a different linear extension s_2 . However, the difference between s_1 and s_2 is bounded by ε and the size of Δ . The following is then immediate.

Lemma 5.6. *Consider*

- a positive number ε ,
- a linear simplex $\Delta \subset \mathbb{R}^N$,
- a pair of jets $\sigma_1, \sigma_2 \in J^1(\Delta, \mathbb{R}^n)$ satisfying $\sigma_2 \in B_f(\sigma_1, \varepsilon)$, and
- their linear extensions $s_i = \text{LinExt}_\Delta(\sigma_i)$.

Then the following bounds hold:

- $d_{C^0}(s_1, s_2) = O(\varepsilon r_{\max}(\Delta))$, and
- $d_{C^1}(s_1, s_2) = O(\varepsilon + \varepsilon r_{\max}(\Delta))$.

We observe that the second claim would still hold if σ_1 and σ_2 had lain in the same *fiber* over $x \in M$, but the first one would not. To emphasize that σ_1 and σ_2 lie in the same *front fiber*, we refer to s_2 as an ε -**slope-perturbation** of s_1 .

5.2.2. Over a color. We now extend this procedure to all the simplex of a given color. We package this into a definition:

Definition 5.7. Let $\varepsilon > 0$ and let K be a finite, colored, simplicial complex of pure dimension. Fix a piecewise linear map $s : |K| \rightarrow \mathbb{R}^n$ and a color i . We say that $s' : |K| \rightarrow \mathbb{R}^n$ is the result of **slope-perturbing s by ε over the i th color** if s' is obtained from s by

- first ε -slope-perturbing over all simplices of color i (as in Lemma 5.6), and
- then applying the join construction (from Definition 4.5) in the star of all such simplices.

We depict this in Figure 5.




FIGURE 5. An example of the result s' of slope-perturbing s over a simplex Δ in K . We use the join over $\text{star}(\Delta^1)$.

The following statement is immediate from the bounds established in Lemmas 4.6 and 5.6:

Lemma 5.8. *Let K be a colored simplicial complex of pure dimension m . Let K' be the collection of simplices of color i , assume it is finite. Suppose $s' : |K| \rightarrow \mathbb{R}^n$ is the result of slope-perturbing a piecewise linear map $s : |K| \rightarrow \mathbb{R}^n$ by ε over the i th color. Let r_{\max} be the maximum value of $r_{\max}(\Delta)$ over $\text{star}(K')$ and let r_{\min} be the minimum value of $r_{\min}(\Delta)$ over $\text{star}(K')$. Then, the following bounds hold:*

- $d_{C^0}(s, s') = O(\varepsilon r_{\max})$, and
- $d_{C^1}(s, s') = O\left(\varepsilon r_{\max} \left(1 + \frac{r_{\max}}{r_{\min}}\right)\right)$.

6. JIGGLING

In this section we establish our main jiggling results. We first tackle jiggling in the linear setting (Section 6.1), for maps of linear polyhedra into Euclidean space. We then address jiggling for sections of bundles over manifolds (Section 6.2).

6.1. Jiggling in the linear setting. We can formulate our first jiggling theorem, using the terminology for differential relations over polyhedra from Section 2.2.6.

Theorem 6.1. *Consider a finite simplicial complex K of pure dimension, and an open and fiberwise dense differential relation $\mathcal{R} := \{\mathcal{R}_\Delta\}_{\Delta \in K^{(\text{top})}}$. Then, given*

- $\varepsilon > 0$, and
- a piecewise smooth map $s : |K| \rightarrow \mathbb{R}^n$,

there exists an ε -jiggling (s', K_ℓ) of (s, K) such that

- s' is piecewise linear with respect to the crystalline subdivision K_ℓ of K , and
- s' is a piecewise solution of \mathcal{R} .

Proof. We write C for the number of colors needed to color all the K_ℓ (Lemma 3.3). We write $K_\ell^{(\text{top}, i)}$ for the collection of top dimensional simplices in K_ℓ of color i .

Strategy of the proof. The strategy we employ is the following: we linearize s with respect to K_ℓ and then, inductively on the color $i \in [C]$, we slope-perturb by a constant ε_i so that the resulting section $s_\ell^{(i)}$ is a piecewise smooth solution of \mathcal{R} over the simplices of color $j \leq i$. The key points to pay attention to are:

- ℓ has to be large enough so that the linearization is a good approximation.
- ℓ has to be large enough to guarantee that solutions can be constructed via slope-perturbation.
- each $j^1 s_\ell^{(i)}$ has a δ_i neighborhood lying in \mathcal{R} , which determines how large future perturbations and hence ε_{i+1} can be, so that the perturbed section is still a solution.

Choosing a good collection of constants. We now choose the constants $\{\varepsilon_i\}_{i \in [C]}$, $\{\delta_i\}_{i \in [C]}$ and ℓ . A first requirement is that ℓ must be large enough so that $d_{C^1}(s, s_\ell^{\text{lin}}) < \varepsilon/2$, which can be achieved using Lemma 4.2.

According to Lemma 5.8 we can find $\varepsilon_0 > 0$ such that, for any subdivision ℓ and color i , the C^1 -distance between s and the result of slope-perturbing s by ε_0 over the i th color in K_ℓ is at most $\frac{\varepsilon}{2C}$. In fact, we can assume that this ε_0 works for any piecewise smooth map $h : |K| \rightarrow \mathbb{R}^n$ that is ε -close to s in the C^1 -distance. To see this, two comments are relevant: First we note that this works for s because according to Lemma 3.6, we can bound the size and the ratio of the size of any two simplices independently of ℓ . Next, we point out that the 1-jet of h is contained in the ε -neighborhood of $j^1 s$ in $J^1(|K|, \mathbb{R}^n)$, which is compact, which allows us to find an ε_0 that works for all such h .

Corollary 5.5 then yields the constant $\delta_0 > 0$. Namely, it is chosen so that any 1-jet $\sigma \in \overline{B(j^1 s(|K|), \varepsilon)}$ admits an ε_0 -slope-perturbation σ' such that the δ_0 -ball around it lies in \mathcal{R} . Moreover, the linear extension of each $\sigma'' \in B(\sigma, \delta_0)$ over a δ_0 -ball is a solution of \mathcal{R} .

We then keep going, arguing similarly, inductively on $i \in [C]$. Lemma 5.8 yields $\varepsilon_i > 0$ small enough so that a ε_i -slope-perturbation over all simplices of the same color causes at most a $\delta_{i-1}/4$ perturbation over adjacent simplices, in a C^1 -sense. Our choice of ε_i then yields a $\delta_i > 0$ using Corollary 5.5. We can assume without loss of generality that $\{\varepsilon_i\}_{i \in [C]}$ is a decreasing sequence and that $2\delta_{i+1} < \delta_i$ for all $i \in [C]$.

Lastly, we ask ℓ to be large enough so that $\max(1, \|s\|_{C^1} + 2\varepsilon) r_{\max}(\Delta) < \delta_C/2$ for every $\Delta \in K_\ell$. That this is possible follows from Lemma 3.6.

Induction hypothesis. We can now construct the requested jiggling s' by induction on $i \in [C]$. The induction hypothesis is that there exists a continuous and piecewise linear map $s_\ell^{(i)} : |K| \rightarrow \mathbb{R}^n$ such that

- (IH1) for all $\Delta \in \cup_{j=0}^i K_\ell^{(\text{top}, j)}$, all $x \in \Delta$ and $\sigma'' \in B(j^1 s_\ell^{(i)}|_\Delta(x), \delta_i/2)$, the linear extension of σ'' over Δ is a solution of \mathcal{R} , and
- (IH2) $d_{C^1}(s_\ell^{(i-1)}, s_\ell^{(i)}) < \frac{\varepsilon}{2C}$ where $s_\ell^{(-1)} = s_\ell^{\text{lin}}$.

Since $\cup_{j=0}^{-1} K_\ell^{(\text{top}, j)} = \emptyset$, the base case follows from the induction step and hence we only prove the latter. Consider therefore $s_\ell^{(i)}$ as given. We want to construct $s_\ell^{(i+1)}$.

Construction. We construct $s_\ell^{(i+1)}$ by slope-perturbing $s_\ell^{(i)}$ by ε_{i+1} over all simplices in $K_\ell^{(\text{top}, i+1)}$, as follows. Consider a simplex $\Delta \in K_\ell^{(\text{top}, i+1)}$ with a vertex v . By definition of δ_{i+1} , we know there exists $\sigma' \in B_f(j^1 s_\ell^{(i)}(v), \varepsilon_{i+1})$ such that for all $\sigma'' \in B(\sigma', \delta_{i+1})$ we have that $\text{LinExt}(\sigma'')$ over $B(\pi(\sigma''), \delta_{i+1})$ is a solution. According to our assumptions on ℓ , we have that $r_{\max}(\Delta) < \delta_C/2$. As such, we can define $s_\ell^{(i+1)}|_\Delta$ as $\text{LinExt}(\sigma')$. According to Lemma 5.8, we can do this for all simplices $\Delta \in K_\ell^{(\text{top}, i+1)}$ at the same time, applying the join construction over the star.

By construction, item (IH2) is immediately satisfied. To see that item (IH1) is satisfied on each simplex $\Delta \in \cup_{j=0}^i K_\ell(i+1)$, we distinguish two cases: Δ is of color $i+1$ or of a previous color.

Simplices of color $i+1$. Take $\sigma'' \in B(j^1 s_\ell^{(i+1)}|_\Delta(x), \delta_{i+1}/2)$ for some $x \in \Delta$. Recall that v is the vertex of Δ over which we chose the 1-jet σ' we linearly extended and that hence we have $j^1 s_\ell^{(i+1)}(v) = \sigma'$. Then we see that

$$\begin{aligned} d_{C^0}(\sigma'', j^1 s_\ell^{(i+1)}(v)) &\leq d_{C^0}(\sigma'', j^1 s_\ell^{(i+1)}(x)) + d_{C^0}(j^1 s_\ell^{(i+1)}(x), j^1 s_\ell^{(i+1)}(v)) \\ &< \delta_{i+1}/2 + d_{C^0}(j^1 s_\ell^{(i+1)}(x), j^1 s_\ell^{(i+1)}(v)), \end{aligned}$$

where the latter term is bounded by $\|s_\ell^{(i+1)}\|_{C^1} r_{\max}(\Delta)$. Our assumptions on ℓ thus imply that $\sigma'' \in B(j^1 s_\ell^{(i+1)}(v), \delta_{i+1})$. Hence, the section $\text{LinExt}(\sigma'')$ is a solution over Δ .

Simplices of a previous color. Let now Δ be in $K_\ell^{(\text{top}, j)}$ for some $j < 1+i$ and take $\sigma'' \in B(j^1 s_\ell^{(i+1)}(x), \delta_{i+1}/2)$ with $x \in \Delta$. By definition of ε_{i+1} it follows in particular that $d_{C^0}(j^1 s_\ell^{(i)}, j^1 s_\ell^{(i+1)}) < \delta_i/2$, and hence we see that

$$\begin{aligned} d_{C^0}(\sigma'', j^1 s_\ell^{(i)}(x)) &\leq d_{C^0}(\sigma'', j^1 s_\ell^{(i+1)}(x)) + d_{C^0}(j^1 s_\ell^{(i+1)}(x), j^1 s_\ell^{(i)}(x)) \\ &< \delta_{i+1}/2 + \delta_i/4 < \delta_i/2, \end{aligned}$$

where the last step follows since we assumed that $2\delta_{i+1} < \delta_i$. Therefore $\text{LinExt}(\sigma'')$ over Δ is a solution of \mathcal{R} and item (IH1) is still satisfied over Δ .

End of induction. We define $s' = s_\ell^{(C)}$. The inductive hypotheses imply that s' is a piecewise linear solution approximating s . \square

With minimal adaptations, the proof we have given also provides us with the relative case. We will refer to s' below as the result of jiggling s relative to $|K'|$ and $|K''|$.

Corollary 6.2. *Suppose K , \mathcal{R} , ε , and s are given as in Theorem 6.1. Suppose moreover that we are given disjoint subcomplexes $K', K'' \subset K$ (of full dimension) with disjoint neighborhoods $\mathcal{O}_p(|K'|), \mathcal{O}_p(|K''|) \subset |K|$ such that $s|_{\mathcal{O}_p(|K'|)}$ is a solution of \mathcal{R} . Then, there exists an ε -jiggling $(s' : |K| \rightarrow \mathbb{R}^n, K_\ell)$ such that*

- s' is piecewise linear on $|K| \setminus (\mathcal{O}_p(|K'|) \cup \mathcal{O}_p(|K''|))$ with respect to K_ℓ ,
- s' is a piecewise solution of \mathcal{R} on $|K| \setminus \mathcal{O}_p(|K''|)$, and
- $s' = s$ on $|K' \cup K''|$.

Proof. Let $L \in \mathbb{N}$ be such that $|\text{star}^2(K' \cup K'', K_L)| \subset \mathcal{O}p(|K'|) \cup \mathcal{O}p(|K''|)$, implying that s restricted to $|\text{star}^2(K', K_L)|$ is a solution. Then since K' is finite, there exists $\delta > 0$ such that

$$B(j^1 s|_{|\text{star}^2(K', K_L)|}, \delta) \subset \mathcal{R}.$$

We take the number of subdivisions ℓ to be at least L .

Let $\{U_i\}_{i \in I}$ be a finite collection of convex opens in \mathbb{R}^N so that their union $\cup_i U_i$ intersected with $|K|$ covers $M \setminus (\mathcal{O}p(|K'|) \cup \mathcal{O}p(|K''|))$, but is disjoint from $|K' \cup K''|$. We define $K^i = K_\ell \cap U_i$ and note that they are nice subcomplexes of K so that we can interpolate over their ring as in Definition 4.3. We assume that ℓ is large enough so that $\cup_i |K^i|$ covers $M \setminus (\mathcal{O}p(|K'|) \cup \mathcal{O}p(|K''|))$ and so that $\text{star}(K^i) \cap (K' \cup K'') = \emptyset$.

Define $s^{(-1)} = s$. Then by induction on i we can construct maps $s^{(i)} : |K| \rightarrow \mathbb{R}^n$ which are a solution over $\cup_{j=0}^i K^j$. We do so by applying the proof of Theorem 6.1 to $s^{(i-1)}$ restricted to K^i , that is: we color the simplices in K^i and slope-perturb over those simplices. When subdivision is needed for jiggling, we subdivide the entirety of K . We define the map s^i as

- the result of jiggling on $|K^i|$,
- $s^{(i-1)}$ on $|\text{Cl}(K \setminus \text{star}(K^i))|$, and
- the result of interpolation on each simplex in $\text{ring}(K^i)$.

Note that the second item achieves in particular that $s^{(i)}$ equals s over $|K' \cup K''|$.

To ensure that the map remains a solution over $|\mathcal{O}p(K')|$, we need to make sure that, all together, the map s is perturbed less than δ . This can be achieved by choosing: 1) ℓ large enough so that the linearization is a good enough approximation, and 2) the constant ε_0 small enough, which controls the size of all slope-perturbations in the proof of Theorem 6.1. For the second point we additionally use Lemma 4.4 to bound the perturbations over simplices where we interpolate. \square

6.2. Jiggling over manifolds. We now move to the general case of jiggling sections of arbitrary fiber bundles $E \rightarrow M$. We will work chart by chart, relative to previous charts, using Corollary 6.2. We note that if, in the statement below, M is compact, the subdivision T' is a crystalline subdivision of T , but not if M is non-compact.

Theorem 6.3. *Let M be a manifold, possibly open, endowed with a smooth triangulation $T : K \rightarrow M$. Consider a fiber bundle $E \rightarrow M$ and an open and fiberwise dense differential relation $\mathcal{R} \subset J^1(E)$. Then, given*

- a continuous function $\varepsilon : M \rightarrow \mathbb{R}_+$,
- a piecewise smooth section $s : M \rightarrow E$, and
- a subcomplex $Q \subset M$ of T such that $s|_{\mathcal{O}p(Q)}$ is a solution of \mathcal{R} ,

there exists an ε -jiggling (s', T') of (s, T) such that

- s' is a piecewise solution of \mathcal{R} with respect to T' , and
- $s'|_Q = s|_Q$.

Proof. Suppose first that M is compact and Q is empty. Since the set $B(s(M), \varepsilon) \subset E$ is compact, it can be covered by finitely many fibered charts $\{\phi_i : U_i \times W_i \rightarrow \mathbb{R}^m \times \mathbb{R}^n\}_{i \in [I]}$, where U_i is a chart of M and W_i is a chart of the fiber. We can assume that ℓ is large enough such that we can cover K_ℓ by finitely many subcomplexes $\{K^{(i)}\}_{i \in [I]}$ (not necessarily nice) with $T(\text{star}^2 K^{(i)}) \subset U_i$.

The argument now amounts to doing induction on i . Starting with $s = s^{(-1)}$, we produce a sequence of sections $s^{(i)}$ via subsequent applications of Corollary 6.2. We note that linear extensions and slope-perturbations work equally well for sections as they do for maps. Concretely, in the i th step, using Corollary 6.2, we jiggle the section $\phi_i \circ s^{(i-1)} : T(|\text{star}^2(K^{(i)})|) \rightarrow \mathbb{R}^n$ relative to $|\cup_{j=0}^{i-1} K^{(j)}|$ and $|\text{ring}(\text{star}(K^{(i)}))|$. We extend the resulting section $s^{(i)}|_{|\text{star}^2(K^{(i)})|}$ to M , by requiring that $s^{(i)}$

equals $s^{(i-1)}$ over $M \setminus |\text{star}^2(K^{(i)})|$. Whenever subdivision is needed for the jiggling, we do not just subdivide $K^{(i)}$, but the entirety of K . In finitely many steps the proof is complete.

The relative case. In the relative case where $Q \neq 0$ we apply jiggling relative to Q throughout the proof.

The non-compact case. If M is non-compact, we take an exhaustion $\{C_k\}_{k \in \mathbb{N}}$ by compacts. Inductively on k , we jiggle the section s over the annulus $C_k \setminus C_{k-1}$ without changing the triangulation of C_{k-2} . In what follows we provide the details.

We set again $s^{(-1)} = s$ and $K^{(-1)} = K$ and produce a sequence of sections $s^{(k)}$ that are each a piecewise solution of \mathcal{R} on a neighborhood of C_i with respect to the triangulation $T : |K^{(k)}| \rightarrow M$. We can assume, by possibly removing some compacts from the exhaustion, that $|\text{star}^3(K \cap C_k, K)| \subset C_{k+1}$.

Assume now that $s^{(k-1)}$ and $K^{(k-1)}$ have been constructed, then the strategy is illustrated in Figure 6. We define the subcomplex $L^{(k)} = K^{(k-1)} \cap (C_k \setminus C_{k-1})$ of $K^{(k-1)}$, and whenever we take its star or ring in the following we do so as subcomplex of $K^{(k-1)}$. Using the compact case, we then jiggle $s^{(k-1)}$ over $|\text{star}^2(L^{(k)})|$ relative to $|\text{ring}(\text{star}(L^{(k)}))|$. We note that we jiggle $s^{(k-1)}$ over a manifold that generally will have a boundary, but that this does not pose any issues.

Jiggling produces a section $s^{(k)}$ over $|\text{star}^2(L^{(k)})|$ that is piecewise smooth with respect to $T : |(\text{star}^2(L^{(k)}))_\ell| \rightarrow M$ for some $\ell \in \mathbb{N}$. In particular $s^{(k)}$ is now a solution over a neighborhood of $C_k \setminus C_{k-1}$. We extend $s^{(k)}$ to M by requiring it equals $s^{(k-1)}$ on $M \setminus |\text{star}^2(L^{(k)})|$. We can assume that the jiggling was small enough so that $s^{(k)}$ restricted to $|\text{star}^2(L^{(k)})| \cap C_{k-1}$ remains a solution. Then $s^{(k)}$ is a solution over a neighborhood of C_k .

We extend the ℓ th crystalline subdivision of $\text{star}^2(L^{(k)})$ to a subdivision $K^{(k)}$ of K defined as

- the ℓ th crystalline subdivision of $\text{star}^2(L^{(k)})$,
- the 0th crystalline subdivision of $K^{(k-1)} \setminus (\text{star}^3(L^{(k)}))$, and
- the barycentric cone off in each simplex in $\text{ring}(\text{star}^2(L^{(k)}))$.

The section $s^{(k)}$ is then indeed piecewise smooth with respect to $T : K^{(k)} \rightarrow M$.

Since we assumed that $|\text{star}^3(K \cap C_k, K)| \subset C_{k+1}$, we see that $K^{(k)}$ and $K^{(k-1)}$ agree over C_{k-2} . In the limit we hence obtain a well-defined triangulation. \square

Alternatively, in the proof of Theorem 6.3 for the non-compact case, we could have used Proposition 3.12. This makes sure that the resulting subdivision of K is a generalized crystalline subdivision, which proves useful in Section 8.3.

Corollary 6.4. *Consider the assumptions from Theorem 6.3. Then, there exists an ε -jiggling (s', T') such that*

- s' is a piecewise solution of \mathcal{R} with respect to T' ,
- $s'|_Q = s|_Q$, and
- the subdivision of K underlying T' is a generalized crystalline subdivision of K .

Proof. The section s' can be constructed as in the non-compact case of Theorem 6.3. For the subdivision K' we have to make some changes to the proof, which we now indicate:

Recall that $\{C_k\}_{k \in \mathbb{N}}$ is an exhaustion by compacts of M . We define δ_k such that the closed δ_k -neighborhood $\overline{B}(C_k, \delta)$ of C_k is contained in $|\text{star}(K \cap C_k, K)|$.

We now start the induction and assume that $s^{(k-1)}$ and $K^{(k-1)}$ have been constructed. We recall that the section $s^{(k)}$ is constructed such that it is piecewise smooth with respect to $T : |(\text{star}^2(L^{(k)}))_\ell| \rightarrow M$, where $L^{(k)} = K^{(k)} \cap (C_k \setminus C_{k-1})$ is a subcomplex of $K^{(k)}$. We assume that ℓ is large enough that we can cone off $K_\ell^{(k-1)} \cap C_k$ within a δ_k -neighborhood in M to any finer crystalline subdivision of K as in Proposition 3.12. If we do so for every k (and in particular have done so for $k-2$), we can then define the subdivision $K^{(k)}$ of K as the result of Proposition 3.12 such that $K^{(k)}$ equals




FIGURE 6. A sketch of the proof of Theorem 6.3 in the non-compact case when about to construct $s^{(k)}$. The compacts C_{k-1} and C_k lie left of the respective orange lines. We have drawn $K^{(k-1)}$ and indicated the subcomplex $L^{(k)}$ and its stars. On top we indicate how to obtain $s^{(k)}$ from $s^{(k-1)}$ and on the bottom how to obtain $K^{(k)}$ from $K^{(k-1)}$.

- the 0th crystalline subdivision of $K^{(k-1)} \cap C_{k-2}$, and
- the ℓ th crystalline subdivision of $K^{(k-1)}$ in $M \setminus B(C_{k-2}, \delta_{k-2})$.

We see that $K^{(k)}$ and $K^{(k-1)}$ agree over C_{k-2} . In the limit we hence obtain a well-defined triangulation. The underlying crystalline subdivision is indeed a generalized crystalline subdivision as each simplex is only subdivided a finite number of times using crystalline subdivision, except when it lies in a $B(C_k, \delta) \setminus C_k$, then also the barycentric cone off is applied once, before it is no longer changed.

We observe that the section $s^{(k)}$ is then indeed piecewise smooth with respect to $T : K^{(k)} \rightarrow M$. \square

7. JIGGLING AS AN H-PRINCIPLE

In this section we discuss how jiggling can be interpreted as an h-principle without homotopical assumptions for piecewise solutions. As we will see in Section 7.2 (and as foreshadowed in Section 2.4), simplicial sets provide the natural setting for such an statement.

Even though the setting of topological spaces is less natural for our purposes (see Section 7.1 for an informal discussion), we do want to make the following observation: at this point we have already proven the π_0 -surjectivity at of the map $\text{Sol}_{\text{PS}}(\mathcal{R}) \hookrightarrow \Gamma_{\text{PS}}(E)$ between *topological* spaces. Here we denote by $\Gamma_{\text{PS}}(E)$ the space of piecewise smooth sections of E and by $\text{Sol}_{\text{PS}}(\mathcal{R})$ the subspace of piecewise smooth solutions of \mathcal{R} .

Corollary 7.1. *Let $\mathcal{R} \subset J^1(E)$ be an open, fiberwise dense, differential relation. Then, every section $s : M \rightarrow E$ is homotopic, through piecewise smooth sections, to a piecewise smooth solution $s' : M \rightarrow E$.*

Proof. We can ε -jiggle s using Theorem 6.3 to end up with a piecewise smooth solution $s' : M \rightarrow E$ with respect to a triangulation T' of M . If we take ε small enough, the sections s and s' are homotopic as piecewise smooth maps by linearly interpolating (with respect to some fibered charts chosen a priori) over T' . \square

7.1. A weak homotopy equivalence between topological spaces. We now make a handful of remarks about the map $\iota : \text{Sol}_{\text{PS}}(\mathcal{R}) \hookrightarrow \Gamma_{\text{PS}}(E)$. We do this to motivate the simplicial approach and pose a couple of open questions. This subsection contains no results and can otherwise be skipped.

Suppose we are interested in proving π_k -injectivity of ι . To do so we start off with a family of sections $s : (\mathbb{D}^n, \mathbb{S}^{n-1}) \rightarrow (\Gamma_{\text{PS}}(E), \text{Sol}_{\text{PS}}(\mathcal{R}))$ representing a relative homotopy class. What we would like to do is consider the induced section $\tilde{s} : M \times \mathbb{D}^n \rightarrow E$ and jiggle it. For each $t \in \mathbb{S}^{n-1}$, the sections $\tilde{s}_t : M \times \{t\} \rightarrow E$ are piecewise smooth, but with respect to different triangulations T_t of $M \times \{t\}$. However, without further assumptions, these triangulations need not be compatible in any way. In particular, even though the sections \tilde{s}_t are piecewise smooth, the section \tilde{s} need not be. It follows that we cannot apply jiggling.

Conversely, consider a section $s : M \times \mathbb{D}^n \rightarrow E$ that is piecewise smooth with respect to a triangulation T of $M \times \mathbb{D}^n$. We can then wonder whether the sections $s_t : M \times \{t\} \rightarrow E$ are piecewise smooth. However, the restriction of T to a fiber $M \times \{t\}$ need not be a triangulation. As such, the sections s_t may not be piecewise smooth.

These remarks suggest the following approach, which we leave for the reader to explore. Suppose that we consider a triangulation T of $M \times \mathbb{D}^n$ that is *in general position* with respect to the fibers $M \times \{t\}$ (recall Definition 8.8). This may be enough to deduce that T induces a triangulation on each fiber. Moreover, perhaps this can be used to show that every section that is piecewise with respect to T defines piecewise smooth sections fiberwise. This ingredient may be enough to address the connectivity of the map ι .

Instead of pursuing this, we explain in the next section how a weak homotopy equivalence of simplicial sets follows almost immediately from our earlier results.

7.2. A weak homotopy equivalence of simplicial sets. We now turn our attention to the inclusion $\text{sSol}_{\text{PS}}(\mathcal{R}) \hookrightarrow \text{Sing}(\Gamma^0(E))$ of the simplicial set of piecewise solutions into the simplicial set of continuous sections E .

We denote by $\Gamma^0(E)$ the space of continuous sections of E , endowed with the C^0 -topology. We recall that $\text{Sing}(\Gamma^0(E))$ is defined by $(\text{Sing}(\Gamma^0(E)))_n = \{\Delta^n \rightarrow \Gamma^0(E)\}$, where the latter set is in bijection with $\{s : M \times \Delta^n \rightarrow q_n^*(E) \mid s \text{ continuous section}\}$ and $q_n : M \times \Delta^n \rightarrow M$ denotes the projection onto the first factor. The simplicial set $\text{sSol}_{\text{PS}}(\mathcal{R})$ of piecewise solutions we define as the following.

Definition 7.2. $\text{sSol}_{\text{PS}}(\mathcal{R}) \in \mathbf{sSet}$ is the simplicial set with n -simplices

$$(\text{sSol}_{\text{PS}}(\mathcal{R}))_n = \{s : M \times \Delta^n \rightarrow q_n^*(E) \mid s \text{ is a piecewise smooth solution of } q_n^*(\mathcal{R})\}.$$

The face map $\delta_i : (\text{sSol}_{\text{PS}}(\mathcal{R}))_n \rightarrow (\text{sSol}_{\text{PS}}(\mathcal{R}))_{n-1}$ is given by restricting the solution to $M \times F_i$ where F_i is the face of Δ^n excluding the i th vertex, and by identifying this face F_i with Δ^{n-1} . The degeneracy map $\sigma_i : (\text{sSol}_{\text{PS}}(\mathcal{R}))_n \rightarrow (\text{sSol}_{\text{PS}}(\mathcal{R}))_{n+1}$ is induced by the pullback along the retraction $(\text{id}, r_i) : M \times \Delta^{n+1} \rightarrow M \times \Delta^n$ where r_i is the affine map collapsing the edge $\langle i, i+1 \rangle$.

Here it is worth remarking that we consider triangulations of $M \times \Delta^n$ that are compatible with the stratification of $M \times \Delta^n$ induced by the stratification of Δ^n by its faces. Concretely, we ask that the triangulation restricts to a triangulation of each $M \times F$, where F is a face. We will use the fact that any triangulation of $M \times \partial\Delta^n$ can be extended to a triangulation of $M \times \Delta^n$; this follows from the contractibility of the space of triangulations [20].

From the perspective of jiggling it does not matter that we now consider solutions of the pullback relation $q_n^*(\mathcal{R})$. If $\mathcal{R} \subset J^1(E)$ is an open and fiberwise dense relation, the relation $q_n^*(\mathcal{R}) \subset J^1(q_n^*(E))$ is as well.

To establish a weak homotopy equivalence between $\text{sSol}_{\text{PS}}(\mathcal{R})$ and $\text{Sing}(\Gamma^0(E))$, we just need to unpack what this entails in terms of sections of the bundle E . The statement will then follow almost immediately from jiggling as in Theorem 6.3. The reason is that the parametric version of jiggling, in this simplicial sense, is just a specific case of the non-parametric case.

Theorem 7.3. *Let $\mathcal{R} \subset J^1(E)$ be an open and fiberwise dense differential relation. The inclusion $\text{sSol}_{\text{PS}}(\mathcal{R}) \hookrightarrow \text{Sing}(\Gamma^0(E))$ is weak homotopy equivalence of simplicial sets.*

Proof. According to Definition 2.25, we have to show that, given a diagram of the form

$$\begin{array}{ccc} \partial\Delta[n] & \longrightarrow & \text{sSol}_{\text{PS}}(\mathcal{R}) \\ \downarrow & \nearrow \text{dashed} & \downarrow \\ \Delta[n] & \longrightarrow & \text{Sing}(\Gamma^0(E)), \end{array}$$

there exists a lift $\Delta[n] \rightarrow \text{sSol}_{\text{PS}}(\mathcal{R})$ such that the upper triangle commutes and the lower triangle commutes up to homotopy.

Hence we assume that we are given a continuous section $F : M \times \Delta^n \rightarrow q_n^*(E)$ such that F restricted to $M \times \partial\Delta^n$ is a piecewise smooth solution of $q_n^*(\mathcal{R})$. Then we need to find a section $G : M \times \Delta^n \rightarrow q_n^*(E)$ such that

- G is a piecewise smooth solution of $q_n^*(\mathcal{R})$, and
- G is homotopic to F via continuous sections relative to $M \times \partial\Delta^n$.

The construction of G consists of two steps. Firstly, we approximate F by a section F' that is piecewise smooth such that F' is a solution on $\mathcal{O}p(M \times \partial\Delta^n)$ and such that F' agrees with F on $M \times \partial\Delta^n$. Secondly, we apply jiggling of Theorem 6.3 to F' relative to $M \times \partial\Delta^n$ to end up with G . If F' and F , and in turn G and F' are sufficiently close, they are all homotopic via continuous sections. \square

Remark 7.4. We want to remark that it is also an option to define $\text{sSol}_{\text{PS}}(\mathcal{R})$ as consisting of pairs (s, T) where additionally s is piecewise smooth with respect to the triangulation T of $M \times \Delta^n$. The face map can be defined as the restriction. One way of defining the degeneracy maps is noting that the n -simplices and the face maps form a semi-simplicial set that is Kan, which can hence be endowed with degeneracy maps (and degenerate simplices) which make it into a simplicial set $\text{sSol}_{\text{PS}}^{\text{T}}(\mathcal{R})$ [27]. More concretely, the simplicial set can be described as the left Kan extension of the semi-simplicial set. With minor adaptations the proof of Theorem 7.3 follows through, proving that the inclusion $\text{sSol}_{\text{PS}}^{\text{T}}(\mathcal{R}) \hookrightarrow \text{Sing}(\Gamma^0(E))$ is a weak homotopy equivalence. In particular this recovers that the fibers of the forgetful functor $\text{sSol}_{\text{PS}}^{\text{T}}(\mathcal{R}) \rightarrow \text{sSol}_{\text{PS}}(\mathcal{R})$ are contractible.

8. APPLICATIONS

In this section we discuss two applications of jiggling. We start with maps that are transverse to a distribution in Section 8.1, which has applications to both immersions and submersions. After that we discuss contact forms in Section 8.2. Finally, in Section 8.3.1 we discuss how jiggling can be applied to triangulation to become in general position to a distribution.

8.1. Transversality. As a first application we show that using jiggling every map can be made to be piecewise transverse to a distribution. For compact (subsets of) manifolds this follows from Thurston's jiggling lemma, but here we also deal with the non-compact case.

Corollary 8.1. *Let $\varepsilon : M \rightarrow \mathbb{R}_+$ be given, let $f : M \rightarrow N$ be a piecewise smooth map and ξ a distribution of constant rank on N . Then there exists a map $g : M \rightarrow N$ that is transverse to ξ and satisfies $d_{C^1}(f, g) < \varepsilon$. In particular we can choose f and g to be homotopic as piecewise smooth maps.*

Proof. A map $M \rightarrow N$ induces a section $M \rightarrow M \times N$, which we can jiggle using Theorem 6.3 to be transverse to ξ , since this is an open and fiberwise dense relation on $J^1(M \times N)$. \square

In particular this result can be applied to maps of maximal rank by the observation that a map $M \rightarrow N$ is of maximal rank if and only if the induced section $M \rightarrow M \times N$ is transverse to the horizontal distribution.

Corollary 8.2. *Let $\varepsilon : M \rightarrow \mathbb{R}_+$ and let $f : M \rightarrow N$ be a smooth map. Then there exists a piecewise smooth map $g : M \rightarrow N$ such that g is a piecewise embedding of maximal rank and $d_{C^1}(g, f) < \varepsilon$. In particular we can choose f and g to be homotopic as piecewise smooth maps.*

Proof. We define the distribution $\xi_H : M \times N \rightarrow T(M \times N)$ as the horizontal distribution $\xi_H(x, y) = T_y N$ and we denote by $\tilde{f} : M \rightarrow M \times N$ the section induced by f . Then we observe that f has maximal rank if and only if \tilde{f} is transverse to ξ_H . Hence we apply Theorem 6.3 to f which results in a piecewise smooth map $\tilde{g} : M \rightarrow M \times N$ which is transverse to ξ_H . We can assume \tilde{g} to be a section if we take g and f to be close enough. The requested map g is then the one induced by \tilde{g} . \square

In the above examples we also obtain a weak homotopy equivalence using Theorem 7.3. Here we denote the relation that a map $M \rightarrow N$ is transverse to a given distribution by $\mathcal{R}_{\text{transv}}$, the relation that the map is an immersion by \mathcal{R}_{imm} and that it is a submersion by $\mathcal{R}_{\text{subm}}$.

Corollary 8.3. *The following inclusions of simplicial sets are weak homotopy equivalences:*

$$\begin{aligned} \text{sSol}_{\text{PS}}(\mathcal{R}_{\text{transv}}) &\hookrightarrow \text{Sing}(C^0(M, N)), \\ \text{sSol}_{\text{PS}}(\mathcal{R}_{\text{imm}}) &\hookrightarrow \text{Sing}(C^0(M, N)) \text{ if } \dim M < \dim N, \text{ and} \\ \text{sSol}_{\text{PS}}(\mathcal{R}_{\text{subm}}) &\hookrightarrow \text{Sing}(C^0(M, N)) \text{ if } \dim M > \dim N. \end{aligned}$$

8.2. Contact forms. Throughout this section we assume that M is a $(2n+1)$ -dimensional manifold. We recall that a contact form¹ is a 1-form $\alpha \in \Omega^1(M)$ such that $\alpha \wedge (d\alpha)^n \neq 0$. Equivalently, a contact form is a section $\alpha : M \rightarrow \Lambda^1(T^*M)$ whose 1-jet satisfies the contact relation $\mathcal{R}_{\text{cont}} \subset J^1(\Lambda^1(T^*M) \setminus \{0\})$. Note that we restrict to $\Lambda^1(T^*M) \setminus \{0\}$ since the contact relation is not fiberwise dense when considering also vanishing 1-forms.

Corollary 8.4. *Let $\varepsilon : M \rightarrow \mathbb{R}_+$ be given and let M be a manifold of odd dimension with a non-vanishing 1-form $\alpha \in \Omega^1(M)$. Then there exists a piecewise smooth contact form $\tilde{\alpha} \in \Omega^1(M)$ such that $d_{C^1}(\alpha, \tilde{\alpha}) < \varepsilon$. In particular, we can choose α and $\tilde{\alpha}$ to be homotopic as piecewise smooth 1-forms.*

We compare the above result to h-principles *with* homotopical assumptions for contact forms, that is, h-principles which compare the space of solutions (here: contact forms) to the space of formal solutions (here: formal contact forms). Recall that the space of formal contact forms consists of pairs $(\alpha, \beta) \in \Omega^1(M) \times \Omega^2(M)$ such that $\alpha \wedge \beta^n \neq 0$.

Gromov [14] proved that if M is an open manifold the complete h-principle holds. It tells us in particular, if we restrict ourselves to the case of π_0 -surjectivity, that any formal contact form (α, β) is homotopic to a contact form $(\tilde{\alpha}, d\tilde{\alpha})$ via formal contact forms in an open manifold. If we compare with Corollary 8.4, we see that jiggling only requires a non-degenerate 1-form α instead of also a 2-form β such that together they form a formal contact form. However, the output of jiggling is also just a piecewise smooth contact form instead of a smooth one.

For closed manifolds π_0 -surjectivity was shown to hold by Martinet [22] and Lutz [21], but π_0 -injectivity was proven to be false by Bennequin [4]. Hence for closed manifolds the full h-principle does not hold, unless we restrict ourselves to *overtwisted* contact structures as shown by Eliashberg [11].

Hence we see that the main strength of jiggling is that it also applies to closed manifolds. The full h-principle we prove, albeit without homotopical assumptions and for simplicial sets, holds true both for open and closed manifolds.

Corollary 8.5. *The inclusion $\text{sSol}_{\text{PS}}(\mathcal{R}_{\text{cont}}) \hookrightarrow \text{Sing}(\Gamma^0(\Lambda^1(T^*M) \setminus \{0\}))$ is a weak homotopy equivalence.*

¹Results for contact structures analogous to the ones we state here for contact forms also hold by taking the projectivization.

8.3. General position via jiggling. Our method of jiggling can also be used to construct triangulations in general position with respect to a distribution, which was Thurston’s original statement. However, this claim does not follow directly from the statement of Theorem 6.3. The difficulty is that, even though the differential relation encoding general position lives in the first jet space, it depends on the triangulation.

8.3.1. *General position.* We first recall:


Definition 8.6. A smooth map $f : M \rightarrow N$ is **transverse to a distribution** ξ of rank r on N if for all $x \in M$ the space $(df)_x(TM) + \xi_{f(x)}$ is maximal. That is, we require that

$$\dim((df)_x(TM) + \xi_{f(x)}) = \min\{n, m + r\},$$

where m and n denote the dimensions of M and N respectively.

If f is piecewise smooth, we require that the above holds on every top-dimensional simplex.

Observe that this does not imply that the map restricted to a lower dimensional simplex is also transverse to the distribution. Hence we consider a stronger notion than transversality, which is known as general position. This requires not just the top-dimensional simplices to be transverse but also all lower dimensional ones. Consequently, it also requires that the distribution does not vary too much within a simplex.



(A) Here the simplex is not in general position, because the edge σ is not transverse to $\mathcal{F}(\xi_{x_i})$.

(B) Here the simplex is in general position with respect to ξ , because the three edges are transverse to $\mathcal{F}(\xi_x)$ for each $x \in \Delta^2$.

FIGURE 7. We give two examples of a distribution ξ on Δ^2 such that Δ^2 is (not) in general position with respect to ξ .

In the following definition we identify a top-dimensional linear simplex Δ in a simplicial complex K with the standard simplex $\Delta^m \subset \mathbb{R}^m$. We also use the fact that any plane V in a tangent space $T_x\mathbb{R}^m$ can be extended, using the Euclidean parallel transport, to a foliation over the whole of \mathbb{R}^m . We will denote this foliation by $\mathcal{F}(V)$. We illustrate the definition in Figure 7.

Definition 8.7. Let $f : M \rightarrow N$ be a map that is piecewise smooth with respect to a triangulation $T : |K| \rightarrow M$. Let ξ be a distribution of constant rank on N . Then f is **in general position** with respect to ξ if:

- f is transverse to ξ , and
- for each top-dimensional simplex $\Delta \in K$ and each face Δ' , we have that $\Delta' \cap \mathcal{F}((f \circ T)^*\xi_x)$ for all $x \in (f \circ T)(\Delta)$.

As a special case of the above definition we can also define what it means for a triangulation T to be in general position.

Definition 8.8. Let $T : |K| \rightarrow M$ be a triangulation and let ξ be a distribution of constant rank on M . Then T is in general position with respect to ξ if $\text{id} : M \rightarrow M$ is in general position with respect to ξ and T .

The general position condition constrains maps $M \rightarrow N$ in a manner that depends on the chosen triangulation T . In particular, a map in general position may stop being so upon subdivision (see Figure 8). Jiggling on the other hand relies on being able to take sufficiently fine crystalline subdivisions T_ℓ of T . Hence this is something that we must address.




FIGURE 8. Example of a triangulation in general position with respect to the horizontal distribution ξ whose crystalline subdivision is not.

8.3.2. *Jiggling.* We now fit general position into our jiggling framework. We are given a map $f : M \rightarrow N$ that is piecewise smooth with respect to a triangulation $T : K \rightarrow M$ and we fix a distribution ξ of constant rank on N . In order to jiggle, we introduce the notation \mathcal{C}_Δ for the collection of model simplices for a single top-dimensional simplex $\Delta \in K$ under generalized crystalline subdivisions of K . That is, each simplex in a generalized crystalline subdivision of K that is a subset of Δ equals a model simplex in \mathcal{C}_Δ up to translation and scaling. The scaling factor is at this moment of no importance.

The collection \mathcal{C}_Δ is finite for each Δ , as we can construct it as in the proof of Proposition 3.12: by Lemma 3.4 there exists a finite set of model simplices \mathcal{C}'_Δ for crystalline subdivisions of Δ . Now given a model simplex $\Delta' \in \mathcal{C}'_\Delta$, we apply crystalline subdivision at most once to each of its faces. We then use the barycentric cone off to extend the subdivision of the boundary of Δ' to Δ' itself. We add each of the resulting simplices to \mathcal{C}'_Δ , to define the finite collection \mathcal{C}_Δ . This motivates the following definition.

Definition 8.9. Let M and N be smooth manifolds and fix a triangulation $T : K \rightarrow M$. We define the general position relation $\mathcal{R}_{\text{GenPos}} \subset J^1(M, N)$ as follows. A jet $\sigma \in J^1(M, N)$ with front projection $(x, y) = \pi_f(\sigma) \in M \times N$ belongs to $\mathcal{R}_{\text{GenPos}}$ if

- for every $\Delta \in K$ with $x \in T(\Delta)$, and
- every simplex Δ' in \mathcal{C}_Δ ,

it holds that Δ' is transverse to $\mathcal{F}((\sigma \circ T)^*\xi_y)$.

Here we use the fact that the pullback of a plane field, at a point, only depends on the plane field and the first jet of the map at said point. We now observe the following.

Lemma 8.10. Fix M , N , K , and ξ as above, and fix a generalized crystalline subdivision K' of K . Assume that \mathcal{C}_Δ is finite for each top dimensional simplex $\Delta \in K$. Then the following hold:

- The relation $\mathcal{R}_{\text{GenPos}}$ is open and fiberwise dense.

- If $s' : M \rightarrow N$ is a piecewise smooth solution of $\mathcal{R}_{\text{GenPos}}$ with respect to K' , then it is in general position with respect to all generalized crystalline subdivisions K'' of K that also subdivide K' .

Proof. Openness and fiberwise density follow immediately from the finiteness of all C_Δ and the local finiteness of K . The second claim follows from the definition of C_Δ as the collection of model simplices for $\Delta \in K^{(\text{top})}$ under generalized crystalline subdivisions of K . \square

The following is now a consequence of Corollary 6.4. We point out that Theorem 6.3 cannot be applied to obtain the following result, since the relation $\mathcal{R}_{\text{GenPos}}$ depends on the model simplices for top-dimensional simplices in K , which change throughout the proof of Theorem 6.3 due to the coning off of arbitrarily many subdivisions of the boundary.

Corollary 8.11. *Let M and N be manifolds, possibly open, where M is endowed with a smooth triangulation $T : K \rightarrow M$ and N with a distribution ξ of constant rank. Then, given*

- a continuous function $\varepsilon : M \rightarrow \mathbb{R}_+$,
- a map $s : M \rightarrow N$, and
- a subcomplex $Q \subset M$ of T such that $s|_{\mathcal{O}_p(Q)}$ is a solution of \mathcal{R} ,

there exists an ε -jiggling (s', T') of (s, T) such that

- s' is in general position with respect to ξ ,
- $s'|_Q = s|_Q$, and
- T' is a generalized crystalline subdivision of T .

Proof. We have already observed that C_Δ is finite for each top-dimensional simplex $\Delta \in K$ and hence by the first item of Lemma 8.10, the relation $\mathcal{R}_{\text{GenPos}}$ is open and fiberwise dense. This allows us to jiggle (s, T) using Corollary 6.4. If we check the proof of Corollary 6.4, we see that it produces a sequence of sections $s^{(k)}$ that are piecewise smooth with respect to a triangulation $T : |K^{(k)}| \rightarrow M$, and are a piecewise smooth solution over C_k , where $\{C_k\}_{k \in K}$ is an exhaustion by compacts of M . Moreover, the functions $s^{(k)}$ and $s^{(k-1)}$ agree over C_{k-2} , and $K^{(k)}$ is a generalized subdivision of K and a subdivision of $K^{(k-1)}$. Hence by the second item of Lemma 8.10, we see that the pair (s', T') produced by Corollary 6.4 satisfies the current statement. \square

It is also immediate that the same statement applies for general position with respect to a finite number of distributions.

8.3.3. Triangulations in general position. The above result can then be applied to the identity map $\text{id} : M \rightarrow M$, which is the setup for Thurston's jiggling. The resulting jiggled map will be a piecewise immersion by construction. The fact that it is a homeomorphism follows from degree theory and is immediate if we take a small enough jiggling [20, Lecture 4]. Thus:

Corollary 8.12. *Let M be a manifold triangulated by $T : |K| \rightarrow M$ and endowed with a distribution $\xi \subset TM$. Then given*

- a continuous function $\varepsilon : M \rightarrow \mathbb{R}_+$, and
- a subcomplex $K' \subset K$ such that $T|_{\mathcal{O}_p(|K'|)}$ is in general position,

there exists an ε -jiggling T' of T such that

- T' is in general position with respect to ξ , and
- $T'|_{|K'|} = T|_{|K'|}$.

REFERENCES

- [1] D. Álvarez-Gavela, Y. Eliashberg, and D. Nadler. “Arboreal models and their stability”. In: *Journal of Symplectic Geometry* 21.2 (2023), pp. 331–381.
- [2] D. Álvarez-Gavela, Y. Eliashberg, and D. Nadler. “Geomorphology of Lagrangian ridges”. In: *Journal of Topology* 15.2 (2022), pp. 844–877.
- [3] D. Álvarez-Gavela, Y. Eliashberg, and D. Nadler. *Positive arborealization of polarized Weinstein manifolds*. 2022. arXiv: [2011.08962](https://arxiv.org/abs/2011.08962) [math.SG].
- [4] D. Bennequin. “Entrelacements et équations de Pfaff”. In: *Asterisque* 107–108 (1983), pp. 87–161.
- [5] M. Bertelson and J. Distexhe. *PL approximations of symplectic manifolds*. 2024. arXiv: [2112.10118](https://arxiv.org/abs/2112.10118) [math.DG].
- [6] R. Casals et al. “Existence h-Principle for Engel structures”. In: *Invent. Math.* 210 (2017), pp. 417–451.
- [7] K. Cieliebak, Y. Eliashberg, and N. Mishachev. *Introduction to the h-principle*. Vol. 239. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2023.
- [8] V. Colin. “Stabilité topologique des structures de contact en dimension 3”. In: *Duke Mathematical Journal* 99.2 (1999), pp. 329–351.
- [9] S.K. Donaldson. “Symplectic submanifolds and almost-complex geometry”. In: *J. Differential Geom.* 44.4 (1996), pp. 666–705.
- [10] Y. Eliashberg. “A theorem on the structure of wave fronts and its applications in symplectic topology”. In: *Functional Analysis and Its Applications* 21.3 (1987), pp. 227–232.
- [11] Y. Eliashberg. “Classification of overtwisted contact structures on 3-manifolds”. In: *Invent. Math.* 98.3 (1989), pp. 623–637.
- [12] Y. Eliashberg. “Contact 3-manifolds twenty years since J. Martinet’s work”. In: *Ann. Inst. Fourier* 42 (1992), pp. 165–192.
- [13] M. Gromov. *Partial differential relations*. Vol. 9. Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1986, pp. x+363.
- [14] M. Gromov. “Stable mappings of foliations into manifolds”. In: *Mathematics of the USSR-Izvestiya* 3.4 (1969), p. 671.
- [15] G. Heuts and I. Moerdijk. *Simplicial and dendroidal homotopy theory*. Springer Nature, 2022.
- [16] V. Humilière, R. Leclercq, and S. Seyfardini. “Coisotropic rigidity and C^0 -symplectic geometry”. In: *Duke Mathematical Journal* 164.4 (2015), pp. 767–799.
- [17] J. Jardine. “Simplicial presheaves”. In: *Journal of Pure and Applied Algebra* 47.1 (1987), pp. 35–87.
- [18] D. Joksimović. “No symplectic Lipschitz structures on $S^{2n \geq 4}$ ”. In: *International Journal of Mathematics* 33.05 (2022), pp. 1–10.
- [19] F. Laudenbach and G. Meigniez. “Haefliger structures and symplectic/contact structures”. In: *Journal de l’École polytechnique - Mathématiques* 3 (2016), pp. 1–19.
- [20] J. Lurie. *Topics in geometric topology*. <https://www.math.ias.edu/~lurie/937.html>. 2009.
- [21] R. Lutz. “Sur l’existence de certaines formes différentielles remarquables sur la sphère S^3 ”. In: *C.R. Acad. Sci. Paris* 270 (1970), pp. 1597–1599.
- [22] J. Martinet. “Formes de contact sur les variétés de dimension 3”. In: *Proc. Liverpool Sing. Symp. II, Lect. Notes in Math.* 209 (1971), pp. 142–163.
- [23] S. Müller and P. Spaeth. “Gromov’s alternative, Eliashberg’s shape invariant, and C^0 -rigidity of contact diffeomorphisms”. In: *International Journal of Mathematics* 25.14 (2014), p. 1450124.
- [24] E. Opshtein. “ C^0 -rigidity of characteristics in symplectic geometry”. In: *Annales scientifiques de l’École normale supérieure* 42.5 (2009), pp. 857–864.
- [25] Á. del Pino and T. Vogel. “The Engel–Lutz twist and overtwisted Engel structures”. In: *Geometry & Topology* 24.5 (2020), pp. 2471–2546.
- [26] D. Rosen and J. Zhang. “Chekanov’s dichotomy in contact topology”. In: *Mathematical Research Letters* 27.4 (2020), pp. 1165–1193.
- [27] C.P. Rourke and B.J. Sanderson. “ Δ -sets I: Homotopy theory”. In: *The Quarterly Journal of Mathematics* 22.3 (1971), pp. 321–338.

- [28] C. H. Taubes. “The Seiberg-Witten invariants and symplectic forms”. In: *Mathematical Research Letters* 1.6 (1994), pp. 809–822.
- [29] W.P. Thurston. “Existence of codimension-one foliations”. In: *Ann. Math.* 104.2 (1976), pp. 249–268.
- [30] W.P. Thurston. “The theory of foliations of codimension greater than one”. In: *Comm. Math. Helv.* 87 (1974), pp. 214–231.
- [31] T. Vogel. “On the uniqueness of the contact structure approximating a foliation”. In: *Geometry & Topology* 20.5 (2016), pp. 2439–2573.
- [32] J.H.C. Whitehead. “On C^1 -complexes”. In: *Ann. of Math.* 41.2 (1940), pp. 809–824.
- [33] H. Whitney. *Geometric Integration Theory*. Princeton University Press, 1957.

UTRECHT UNIVERSITY, DEPARTMENT OF MATHEMATICS, BUDAPESTLAAN 6, 3584 CD UTRECHT, THE NETHERLANDS

Email address: `s.j.fokma@uu.nl`

UTRECHT UNIVERSITY, DEPARTMENT OF MATHEMATICS, BUDAPESTLAAN 6, 3584 CD UTRECHT, THE NETHERLANDS

Email address: `a.delpinogomez@uu.nl`

VU AMSTERDAM, DEPARTMENT OF MATHEMATICS, DE BOELELAAN 1111, 1081 HV AMSTERDAM, THE NETHERLANDS

Email address: `l.e.toussaint@vu.nl`