

Towards a parameter-free analysis of the QCD chiral phase transition and its universal critical behavior

Sabarnya Mitra¹ and Frithjof Karsch¹

Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany
 smitra@physik.uni-bielefeld.de, karsch@physik.uni-bielefeld.de

Abstract. To quantify the universal properties of chiral phase transition in (2+1)-flavor QCD, we use an improved, renormalized order parameter for the chiral symmetry breaking. We construct ratios of this divergence-free order parameter from its values for different pairs of light quark masses. From this, we determine in a parameter-independent manner, the chiral phase transition temperature T_c and the associated critical exponent δ of the universality class. We present first numerical results of these calculations on $N_\tau = 8$ lattices, with staggered fermions.

Keywords: Chiral phase transition, critical behavior, universality class

1 Introduction

With the order of Quantum Chromodynamics (QCD) chiral phase transition and associated universal properties still controversially poised [1,2], the extent of its influence on respective physical QCD thermodynamics and QCD axial anomaly constitute questions of profound importance. In this work, we try addressing these challenges by obtaining a first principle numerical analysis of the universal, critical behaviour of (2+1)-flavor QCD on lattice, with physical values of strange quark masses. Besides extending earlier studies of magnetic QCD equation of state [3,4,5], this work also attempts to quantify the relevant scaling regimes for various physical observables [6,7], which is pivotal for the interpretation of heavy-ion collision data in terms of chiral critical behavior.

2 Theory

With light quark mass m_ℓ , the chief ingredients in this work are the 2-flavor light quark chiral condensate, M_ℓ and the corresponding chiral susceptibility χ_ℓ ,

$$M_\ell = \frac{m_s}{f_K^4} \langle \bar{\psi} \psi \rangle, \quad \chi_\ell = m_s \frac{\partial M_\ell}{\partial m_\ell}, \quad (1)$$

where the strange quark mass m_s and kaon decay constant f_K are normalising factors. To attain a well-defined order parameter M in the continuum limit for

studying the chiral symmetry breaking, the ultraviolet divergent contributions to M_ℓ are eliminated by deducting from it, a suitable fraction of χ_ℓ :

$$M(T, H) = M_\ell(T, H) - H \chi_\ell(T, H) , \quad (2)$$

where $H \equiv m_\ell/m_s$. Thus in chiral limit, M in Eq.(2) receives in the leading order, $\mathcal{O}(H^3)$ light quark mass corrections. This improved order parameter has been introduced in [9] and used previously to probe critical behavior in lattice QCD [5,6,10]. Close to the chiral critical point ($T = T_c, H = 0$), the T and m_ℓ dependence of M is largely governed by the scaling function $f_{G\chi}(z)$ like,

$$M(T, H) = h^{1/\delta} f_{G\chi}(z) + M_{sub-lead}(T, H) , \quad (3)$$

where $f_{G\chi}(z) = f_G(z) - f_\chi(z)$. The scaling functions f_G, f_χ control the quantities M_ℓ, χ_ℓ respectively in a manner, similar to Eq.(3). Corrections to the leading, so-called singular part of the order parameter arise from *i*) non-analytic, universal corrections-to-scaling and *ii*) analytic, non-universal regular terms. They are summarized in $M_{sub-lead}$ and have a leading H -dependence of higher order in H . The scaling variable z , used in Eq.(3) is given as,

$$z = t h^{-1/\beta\delta}, \quad \text{where} \quad t = \frac{1}{t_0} \left[\frac{T}{T_c} - 1 \right], \quad h = \frac{H}{h_0} , \quad (4)$$

with non-universal constants t_0, h_0 , and β, δ being critical exponents of the related universality class. Using properties of f_G and f_χ [11], one can easily find $f_{G\chi}(0) = 1 - 1/\delta$. When plotted as a function of T for different values of H , this manifests a unique intersection point at $z = 0$, which directly provides values of T_c and δ (see left plot of Fig. 1). Exploiting this property, we construct the re-scaled order parameter (resc. in H)

$$M(T, H)/H^{1/\delta} = h_0^{-1/\delta} f_{G\chi}(z) + H^{-1/\delta} M_{sub-lead}(T, H) , \quad (5)$$

and the quantity B from the ratio of M for two different values of H as,

$$B(T, H, c) = \frac{\ln R(T, H, c)}{\ln(c)}, \quad \text{with} \quad R(T, H, c) = \frac{M(T, cH)}{M(T, H)} . \quad (6)$$

In chiral limit, $H^{-1/\delta} M_{sub-lead} \rightarrow 0$ which automatically gives $B(T_c, 0, c) = 1/\delta$.

3 Results

Here in this work, we have used HISQ and Symanzik-improved gauge actions for all the lattice computations. The relevant T and H data are taken from [5,12]. In Fig. 2 (left), we show results for $M/H^{1/\delta}$ using $\delta = 4.7798$, its value in the 3-dimensional $O(2)$ universality class as HISQ preserve only the $O(2)$ part out of full $O(4)$ symmetry. A unique intersection point is apparent from Fig. 2 (left). Its location agrees well with the previous determination of T_c *i.e.* $T_c^{N_\tau=8} =$




Fig. 1. *Left:* Scaling function $f_{G\chi}$ vs T/T_c for different values of H . Shown is $f_{G\chi}$ for the 3-d, $O(2)$ universality class using the non-universal parameter $z_0 = 1.42$ appropriate for (2+1)-flavor QCD in calculations with the HISQ action on discrete lattices of temporal extent $N_\tau = 8$. *Right:* Same as left hand figure but for the observable $B(T, H, 2)$.




Fig. 2. *Left:* The rescaled order parameter, $M(T, H)/H^{1/\delta}$ vs. T for several H values. Shown are results obtained in lattice QCD calculations with the HISQ action on lattices with temporal extent $N_\tau = 8$. *Right:* The observable $B(T, H, c)$ vs. T for different values of c and H . The dashed and dotted lines give the values for $1/\delta$ in $O(2)$ and $U(2) \times U(2)$ [2] universality classes, respectively.

143.7(2) MeV [5], and $h_0^{-1/\delta}$ (Eq.(5)); $h_0^{-1/\delta} = 39.2(4)$ [5]. For a parameter-independent analysis, we evaluate $B(T, H, c)$ as shown in Fig. 2. This plot also features a distinct intersection point, giving T_c, δ but only for small enough $m_\ell \leq m_s/40$, thereby indicating that contributions from $M_{sub-lead}$ may become significant for $H > 1/40$, at least for the present working level of precision.

4 Conclusion and Outlook

We have demonstrated in this work, that as a function of T , the re-scaled order parameter M exhibits a unique intersection point for different values of light quark mass m_ℓ , or equivalently H for small enough values of m_ℓ where the sub-leading contributions are negligible. We also showed that evaluating its ratios allows determining T_c and critical exponent δ , without having the need to make any a priori assumption about the universality class. However, we still need to

improve computational precision by incorporating more T, H data points close to critical point, besides checking the finite-volume effects and taking the continuum limit. This is to estimate δ with accuracy good enough to distinguish between $U(2) \times U(2)$ and $O(4)$ universality classes.

Acknowledgments

This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Proj. No. 315477589-TRR 211. Numerical computations have been performed on the GPU-cluster at Bielefeld University.

References

1. Pisarski, R. D. , Wilczek, F. : Remarks on the Chiral Phase Transition in Chromodynamics. Phys. Rev. D **29**, 338 (1984). doi:10.1103/PhysRevD.29.338
2. Pelissetto, A. , Vicari, E. : Relevance of the axial anomaly at the finite-temperature chiral transition in QCD. Phys. Rev. D **88**, 105018 (2013). doi:10.1103/PhysRevD.88.105018
3. Ejiri, S. , Karsch, F. , Laermann, E. , Miao, C. , Mukherjee, S. , Petreczky, P. , Schmidt, C. , Soeldner, W. , Unger, W. : Magnetic equation of state in (2+1)-flavor QCD. Phys. Rev. D **80**, 094505 (2009). doi:10.1103/PhysRevD.80.094505
4. Bazavov, A. , Bhattacharya, T. , Cheng, M. , DeTar, C. , Ding, H. T. , Gottlieb, S. , Gupta, R. , Hegde, P. , Heller, U. M. , Karsch, F. *et al.* : The chiral and deconfinement aspects of the QCD transition. Phys. Rev. D **85**, 054503 (2012). doi:10.1103/PhysRevD.85.054503
5. Ding, H. T. , Kaczmarek, O. , Karsch, F. , Petreczky, P. , Sarkar, M. , Schmidt, C. , Sharma, S. : Curvature of the chiral phase transition line from the magnetic equation of state of (2+1)-flavor QCD. Phys. Rev. D **109**, 114516 (2024). doi:10.1103/PhysRevD.109.114516
6. Kotov, A. Y. , Lombardo, M. P. , Trunin, A. : QCD transition at the physical point, and its scaling window from twisted mass Wilson fermions. Phys. Lett. B **823**, 136749 (2021). doi:10.1016/j.physletb.2021.136749
7. Braun, J. , Chen, Y. r. , Fu, W. j. , Gao, F. , Huang, C. , Ihssen, F. , Pawłowski, J. M. , Rennecke, F. , Sattler, F. R. , Tan, Y. y. *et al.* : Soft modes in hot QCD matter. arXiv:2310.19853 [hep-ph]. <https://arxiv.org/abs/2310.19853>
8. Bazavov, A. *et al.* [HotQCD]: Equation of state in (2+1)-flavor QCD. Phys. Rev. D **90**, 094503 (2014). doi:10.1103/PhysRevD.90.094503
9. Unger, W. : The chiral phase transition of QCD with 2+1 flavors : a lattice study on Goldstone modes and universal scaling. PhD thesis (2010) Bielefeld University. <https://pub.uni-bielefeld.de/record/2302381>
10. Dini, L. , Hegde, P. , Karsch, F. , Lahiri, A. , Schmidt, C. , Sharma, S. : Chiral phase transition in three-flavor QCD from lattice QCD. Phys. Rev. D **105**, 034510 (2022). doi:10.1103/PhysRevD.105.034510
11. Karsch, F. , Neumann, M. , Sarkar, M. : Scaling functions of the three-dimensional Z(2), O(2), and O(4) models and their finite-size dependence in an external field. Phys. Rev. D **108**, 014505 (2023). doi:10.1103/PhysRevD.108.014505
12. Ding, H. T. *et al.* [HotQCD]: Chiral Phase Transition Temperature in (2+1)-Flavor QCD. Phys. Rev. Lett. **123**, 062002 (2019). doi:10.1103/PhysRevLett.123.062002