Mesoscopic Edge Universality of Orthogonal Polynomial Ensembles

Wenkui Liu*

Abstract

In this paper, we study the mesoscopic fluctuations at edges of orthogonal polynomial ensembles with both continuous and discrete measures. Our main result is a Central limit Theorem (CLT) for linear statistics at mesoscopic scales. We show that if the recurrence coefficients for the associated orthogonal polynomials are slowly varying, a universal CLT holds. Our primary tool is the resolvent for the truncated Jacobi matrices associated with the orthogonal polynomials. While the Combes-Thomas estimate has been successful in obtaining bulk mesoscopic fluctuations in the literature, it is too rough at the edges. Instead, we prove an estimate for the resolvent of Jacobi matrices with slowly varying entries. Particular examples to which our CLT applies are Jacobi, Laguerre and Gaussian unitary ensembles as well as discrete ensembles from random tilings.

Contents

Supported by the Swedish Research Council (VR), grant no. 2021-06015, and the European Research Council (ERC), Grant Agreement No. 101002013.

1 Introduction

Let μ be a Borel measure on $\mathbb R$ with finite moments, i.e., $\int_{\mathbb R} |x|^k d\mu(x) < \infty$ for all $k \in \mathbb N$. The orthogonal polynomial ensemble (OPE) of order $n \in \mathbb N$ associated to μ is a probability measure on $\mathbb R^n$ polynomial ensemble (OPE) of order $n \in \mathbb{N}$ associated to μ is a probability measure on \mathbb{R}^n proportional to

$$
\prod_{1 \le i < j \le n} (x_i - x_j)^2 d\mu(x_1) \cdots d\mu(x_n). \tag{1}
$$

In this paper, we will be interested in studying the behaviour of this ensemble for large *n*.

OPEs arise naturally in many models of statistical mechanics, probability theory, combinatorics, and random matrix theory. For some surveys, we refer to [\[1,](#page-80-1) [2,](#page-80-2) [3,](#page-80-3) [4\]](#page-80-4). An important and well-known source of examples where OPEs appear are the eigenvalues of random Hermitian matrices, where the probability measure is invariant under the conjugation with unitary matrices. Classical examples are the Jacobi, Laguerre and Gaussian unitary ensembles. In these cases, the measure μ is absolutely continuous with an analytic density. The support of the measure can be compact, for example, the (modified) Jacobi unitary ensemble, or unbounded, for example the Gaussian and Laguerre unitary ensembles. When the support of the measure is unbounded, it is natural to rescale the measure with *n*, so that the eigenvalues accumulate on finitely many intervals with high probability as $n \to \infty$. For this reason, we will allow the measure μ to depend on an *n* and write $\mu = \mu_n$.

OPEs with discrete measures are also natural objects to be considered. For instance, uniformly distributed random lozenge tilings of a hexagon give rise to (extended) OPEs associated to the Hahn measure, see [\[5\]](#page-80-5). Similarly, uniformly distributed random domino tilings of an Aztec diamond give rise to (extended) OPEs associated with the Krawtchouk measure, see [\[6\]](#page-80-6). In such cases, the measure μ is discrete and its parameters depend on the size of the tiling, i.e., $\mu = \mu_n$. This is another reason that we allow the measure to be *n*-dependent.

We will study the asymptotic behaviour as $n \to \infty$ and explore cases of μ (or μ_n) both continuous and discrete in this paper. There are three regimes that one can look into, i.e., macroscopic, mesoscopic and microscopic regimes. In the macroscopic regime, one studies the global behaviour of the point processes, whereas in the microscopic regime, one zooms in near a point and studies the nearest neighbour interactions between points. This paper studies the mesoscopic regime which is intermediate between the other two and we will focus on the edges.

Let us, for clarity, start with the example of the the Laguerre unitary ensemble, whose measure is given by $d\mu_n(x) = x^\gamma e^{-nx} dx$ on [0, +∞), for some parameter $\gamma > -1$. It describes the eigenvalue distribution of a Wishart matrix. It is a classical example that has both a hard and soft edge and will therefore he of a Wishart matrix. It is a classical example that has both a hard and soft edge and will, therefore, be an interesting example for the techniques developed in this paper. For the Laguerre unitary ensemble, the eigenvalues accumulate on the interval [0, 4] almost surely as $n \to \infty$ and have the limiting distribution $d\rho_{\mu}(x) = \frac{1}{2\pi} \sqrt{\frac{4-x}{x}} dx$, see Figure [1.](#page-3-0) In the bulk, i.e., any point in the interval (0, 4), the typical distance ω_{μ} (x) = ω_{π} \sqrt{x} ax, see Figure 1. In the bunk, i.e., any point in the interval (0, +), the typical distance between neighbouring points is of order ~ n^{-1} . Near the edges points, the scaling is different. is a hard edge of the interval, since no eigenvalues can be negative. Near the origin, the limiting density behaves as $d\rho_{\mu}(x) \sim 1/\sqrt{x}dx$, which is typical at hard edges, and the neighbouring points are of distance n^{-2} . The right-end point of the interval is a typical example of soft edge where the density vanishes $\sim n^{-2}$. The right-end point of the interval is a typical example of soft edge where the density vanishes as square root $d\rho_{\mu}(x) \sim \sqrt{4 - x} dx$ and the distance between neighbours is ~ $n^{-\frac{2}{3}}$. It is well-known that the microscopic process of this Laguerre unitary ensemble converges to the Bessel point process at the hard edge, sine point process in bulk and Airy point process at the soft edge. These scaling limits on the microscopic scale are universal and observed in a wide variety of models, see [\[7,](#page-80-7) [8,](#page-80-8) [9,](#page-80-9) [10\]](#page-80-10). We also point out that [\[11\]](#page-80-11) offers an overview of universality.

We will study the mesoscopic scales at edges using the scaled linear statistics of the following form $X^{(n)}_{f, \alpha, x_0}$, where the test function *f* is a compactly supported real-valued function, the location is $x_0 \in \mathbb{R}$, and f_{f,α,x_0} , where the test random
the scale is n^{α} for some $\alpha > 0$,

$$
X_{f,\alpha,x_0}^{(n)} := \sum_{i=1}^n f(n^{\alpha}(x_i - x_0)).
$$
 (2)

This scaled linear statistics [\(2\)](#page-2-0) is to examine the points that are at a distance at most of the order of $n^{-\alpha}$ around x_0 . Given that the test function f is compactly supported, the points $\{x_j\}_j$, whose distance between *x*₀ are of order far larger than $n^{-\alpha}$, will eventually fall outside the support of *f* as $n \to \infty$ and only points within a distance of x_0 of order at most $n^{-\alpha}$ fall within the support. Note that if $\alpha = 0$, the scaled linear statistics is reduced to the global linear statistics and "sees" all points simultaneously. If α is h statistics is reduced to the global linear statistics and "sees" all points simultaneously. If α is big enough such that n^{α} is the microscopic scale around the point x_0 , then we are at the local (or microscopic) regime,

Figure 1: Equilibrium measure of the Laguerre unitary ensemble. Hard edge *x* = 0, soft edge *x* = 4.

and the scaled linear statistics will only "see" finitely many points in the limit. Hence, the linear statistics [\(2\)](#page-2-0) lives in the scale of order n^{α} , which we call the mesoscopic scale. Typically, in the bulk $\alpha \in (0, 1)$, at the hard edge $\alpha \in (0, 2)$ and at the soft edge $\alpha \in (0, 2/3)$ hard edge $\alpha \in (0, 2)$, and at the soft edge $\alpha \in (0, 2/3)$.

At this mesoscopic regime in literature, the limiting fluctuations are studied by [\[12\]](#page-80-12) for the classical compact groups. Later, the mesoscopic fluctuations in the bulk and at the soft edges for deformed Wigner matrix are studied by [\[13\]](#page-80-13), which includes GUE. The bulk universal mesoscopic fluctuations for OPEs that can be approximated by modified Jacobi ensembles are considered in [\[14\]](#page-81-0). The bulk mesoscopic fluctuations for the sparsely perturbed Jacobi unitary ensembles are considered in [\[15\]](#page-81-1) . For Wigner matrices and βensembles, the bulk mesoscopic limit is showed in [\[16\]](#page-81-2). The bulk mesoscopic fluctuations for OPEs that priorly have a sine universality are considered in [\[17\]](#page-81-3). The bulk mesoscopic fluctuations of the linear statistics with smooth test functions ($f \in C_0^6$)for β ensembles with smooth potentials ($V \in C^7$ as defined in [\(7\)](#page-4-2)) are considered in [\[18\]](#page-81-4), by the loop equation technique. The mesoscopic fluctuations for circular orthogonal polynomial ensembles are considered in [\[19\]](#page-81-5). By far, a substantial portion of research is concentrated about the bulk. Most methods designed for the bulk break down at edges, which makes the analysis more difficult. Hence, non-trivial modifications to these methods are needed to study the edge behaviour. This paper aims to address the gap in understanding the mesoscopic limit at the edges. It is dedicated to establishing general conditions for such behaviour at the mesoscopic scale for orthogonal polynomial ensembles at the edges.

Here, we present our first two pedagogical results, which are consequences of our general theorems.

Non-varying Weights

A canonical example of a compactly supported non-varying measure is the modified Jacobi Unitary Ensemble

$$
d\mu(x) = (1 - x)^{\gamma_1} (1 + x)^{\gamma_2} h(x) dx, \quad x \in [-1, 1],
$$
\n(3)

where $\gamma_1, \gamma_2 > -1$ are parameters and *h* is an analytic function in a neighbourhood of [−1, 1] and strictly positive on $[-1, 1]$. The limiting distribution for the point process is the arcsine measure $d\rho_\mu(x) = \frac{1}{\pi \sqrt{1-x^2}} dx$ on [−1, 1]. In the bulk, its mesoscopic limit has been studied by [\[14\]](#page-81-0), whose result directly implies that, for $f \in C^{1}(\mathbb{R})$, $x \in (-1, 1)$ and $\alpha \in (0, 1)$, as $r \to \infty$, the following converges in distribution $f \in C_c^1(\mathbb{R})$, $x_0 \in (-1, 1)$ and $\alpha \in (0, 1)$, as $n \to \infty$, the following converges in distribution

$$
X_{f,\alpha,x_0}^{(n)} - \mathbb{E}\left[X_{f,\alpha,x_0}^{(n)}\right] \to \mathcal{N}\left(0, \frac{1}{4\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x) - f(y)}{x - y}\right)^2 dx dy\right). \tag{4}
$$

Our result is at the (hard) edges $x_0 = 1$ or -1 .

Theorem 1.1. Let μ be the modified Jacobi weight as [\(3\)](#page-3-1), given $\gamma_1, \gamma_2 > -1$. Then we have that, for all $f \in C_c^1(\mathbb{R})$ *and* $\alpha \in (0, 2)$ *, the following converges in distribution as* $n \to \infty$

$$
X_{f,\alpha,x_0}^{(n)} - \mathbb{E}\left[X_{f,\alpha,x_0}^{(n)}\right] \to \mathcal{N}\left(0,\sigma_f^2\right) \tag{5}
$$

$$
\sigma_f^2 = \begin{cases} \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dxdy, & x_0 = 1\\ \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dxdy, & x_0 = -1. \end{cases}
$$
(6)

Theorem [1.1](#page-4-1) is a consequence of more general results Theorem [2.1](#page-6-1) or Theorem [2.3,](#page-8-0) which we will discuss in Subsection [2.3.](#page-7-0) The proof of Theorem [1.1](#page-4-1) is presented in Section [8.](#page-62-0)

Varying Weights

Let us consider the case where there exists a real potential V on $\mathbb R$ such that

$$
d\mu_n(x) = e^{-nV(x)}dx, \quad x \in \mathbb{R},\tag{7}
$$

$$
\frac{V(x)}{\log(x^2 + 1)} \to +\infty \quad \text{as } |x| \to \infty.
$$
 (8)

Condition [\(8\)](#page-4-3) is to ensure all finite moments of μ_n . This is a classical case where we are expecting soft edges, since the measure μ_n is supported on the real line. The equilibrium measure with respect to the potential *V* is defined as the unique probability measure ρ_{μ} that minimises the potential

$$
I_V(v) := -\int \int \log|x - y| d\mathsf{v}(x) d\mathsf{v}(y) + \int V(x) d\mathsf{v}(x).
$$
 (9)

The existence and uniqueness of the infinitum $I_V(\rho_\mu)$ is achieved via potential theory. For *V* being analytic and strictly convex, it is also known that ρ_{μ} is supported in a single interval. For more about the potential theory, one may refer to [\[20\]](#page-81-6).

Theorem 1.2. Assume $d\mu_n(x) = e^{-nV(x)}dx$ to be supported on the real line. Let V be an analytic and strictly convex potential satisfying (8). Without loss of generality up to some pormalization, assume the *strictly convex potential satisfying* [\(8\)](#page-4-3)*. Without loss of generality, up to some normalization, assume the equilibrium measure, which is the minimizer of* [\(9\)](#page-4-4)*, to be supported in a single interval* [−1, 1]*. Then the same asymptotic result as* [\(5\)](#page-4-5) *and* [\(6\)](#page-4-6) *in Theorem* [1.1](#page-4-1) *follows for all* $f \in C_c^1(\mathbb{R})$ *and* $\alpha \in (0, \frac{2}{3})$ $\frac{2}{3}$), as $n \to \infty$.

Theorem [1.2](#page-4-0) is a consequence of a general result Theorem [2.2,](#page-8-1) which we will discuss in Subsection [2.3.](#page-7-0) Analyticity and convexity are not the necessary conditions of the general results in this paper. However, they ensure the equilibrium measure to be supported on a single interval, see [\[21\]](#page-81-7). The proof of Theorem [1.2](#page-4-0) is presented in Section [8.](#page-62-0)

The conclusion of the results above holds significance in several aspects. First of all, the limiting variance [\(6\)](#page-4-6) is scaling invariant. Set $g(x) = f(a^2x)$ and we have $\sigma_f^2 = \sigma_g^2$. This gives a heuristic explanation of the independence of α in the limit. However, compared with the bulk mesoscopic limit in [\(4\)](#page-3-2), σ_f^2 is not translation invariant in f translation invariant in *f* .

Finally, the variances for both the soft and the hard edges share the same formula [\(6\)](#page-4-6) as indicated in the two theorems above, though microscopically, the OPEs may have different limiting processes as discussed. Our mesoscopic theorems do not require any knowledge about microscopic information a priori. It is, therefore, reasonable to anticipate that there is a broad range of OPEs for which the linear statistics should yield limiting mesoscopic fluctuations at the edges.

Our purpose of this paper is to extend Theorems [1.1](#page-4-1) and [1.2](#page-4-0) to a wider range of μ and μ_n . Analyticity and convexity of the potential *V* are not necessary. We include cases where μ or μ_n is a discrete measure, for example, Hahn polynomials (from the uniform measure on all lozenge tilings of a hexagon), Krawtchouk polynomials (from the uniform measure on all domino tilings on the Aztec diamond) and Tricomi-Carlitz polynomials given in Subsection [10.5.](#page-77-1) Our starting point is the three-term recurrence relation for the orthogonal polynomials. We give conditions on the coefficients that imply a mesoscopic CLT. Our criteria will be satisfied by the classical (properly scaled) hypergeometric orthogonal polynomials where the recurrence coefficients are explicitly known, cf. [\[22\]](#page-81-8). Our method is inspired by [\[14,](#page-81-0) [23\]](#page-81-9), which show that the recurrence coefficients can fully characterise the moments of the linear statistics of an OPE. The difficulty of adapting the methods of [\[14\]](#page-81-0) is that the standard Combes-Thomas estimate to prove mesoscopic limit in bulk is unsuitable for the analysis at the edges (cf. Sections [3.4\)](#page-18-0). In this paper, we develop estimates at the edges in place of the Combes-Thomas estimate (cf. Propositions [4.1](#page-24-0) and [4.3\)](#page-26-0). Moreover, our estimates work for the varying, as well as non-varying, weights, while Breuer and Duits only focus on the non-varying measure in [\[14\]](#page-81-0). Last but not the least, we analyse the mesoscopic linear statistics directly without implementing the microscopic information.

2 Statement of Results

In this section, we will state our main results. The proofs are given in the later sections. The general results are described in terms of the three-term recurrence relations of orthogonal polynomials, which we will introduce first.

2.1 Orthonormal Polynomials and Recurrence Relations

Given a measure μ_n on R, with finite moments, we let $\{p_{i,n}(x)\}_{i\geq 0}$ be the family of polynomial polynomials with respect to μ_n . The second subscript *n* emphasises that the measure μ_n may be varying in *n* as pointed out by the examples from [\(7\)](#page-4-2). Precisely, such $p_{j,n}$ is the unique polynomial of degree *j* with a positive leading coefficient such that

$$
\int p_{j,n}(x)p_{k,n}(x)d\mu_n(x) = \delta_{j,k}.
$$
\n(10)

It is well-known that orthonormal polynomials satisfy a three-term recurrence relation

$$
xp_{0,n}(x) = a_{1,n}p_{1,n}(x) + b_{0,n}p_{0,n}(x),
$$
\n(11)

$$
xp_{j,n}(x) = a_{j+1,n}p_{j+1,n}(x) + b_{j,n}p_{j,n}(x) + a_{j,n}p_{j-1,n}(x),
$$
\n(12)

for $j = 1, 2, \ldots$ and some coefficients $a_{j,n} > 0$ and $b_{j,n} \in \mathbb{R}$. These coefficients uniquely determine the orthonormal polynomials and are fundamental to our analysis. However, the distribution of linear statistics [\(2\)](#page-2-0) is fully determined by the recurrence coefficients, only if the the measure μ_n is determined by its moments. We will not assume the moment problem is determined in this paper. For the non-varying weight μ , the above discussion follows in the same way.

The asymptotic results of linear statistics of OPEs can be obtained by studying the recurrence coefficients. Among them, Breuer and Duits study the the global fluctuations [\[23\]](#page-81-9). One case of [\[23\]](#page-81-9) is that, given the existence of the limit

$$
\lim_{n \to \infty} a_{n+k,n} =: a, \quad \lim_{n \to \infty} b_{n+k,n} =: b,
$$
\n(13)

for all $k \in \mathbb{Z}_{\geq 0}$, the following converges in distribution, as $n \to \infty$,

$$
X_f^{(n)} - \mathbb{E}[X_f^{(n)}] \to \mathcal{N}\left(0, \sum_{k=1}^{\infty} k \left| \frac{1}{2\pi} \int_0^{2\pi} f(2a\cos(\theta) + b)e^{ik\theta}d\theta \right|^2\right).
$$
 (14)

For the non-varying weights the assumption [\(13\)](#page-5-2) can be simplified to $\lim_{n\to\infty} a_n = a$ and $\lim_{n\to\infty} b_n = b$. Compared with the mesoscopic limit in Theorems [1.1](#page-4-1) and [1.2,](#page-4-0) which are universal results, the variance of the limiting global fluctuations depends on the limits of the recurrence coefficients.

Consequently, the edges of the global fluctuations in this case are $b - 2a$ and $b + 2a$. One should be aware that though the edges of fluctuations coincide with the boundary of the equilibrium measure as in Theorem [1.1](#page-4-1) and [1.2,](#page-4-0) this is not the case in general. For example, the Tricomi Carlitz Polynomials, given in Subsection [10.5,](#page-77-1) have the equilibrium measure supported on \mathbb{R} , while the fluctuations of the associated OPE take place on [−2, 2] only.

Another important result about the limiting fluctuations of mesoscopic linear statistics in the bulk is obtained by [\[14\]](#page-81-0) i.e., [\(4\)](#page-3-2) holds, whenever the recurrence coefficients are such that

$$
a_j = a + O(j^{-1}), b_j = b + O(j^{-1}), \text{ as } j \to \infty,
$$
 (15)

for some $a > 0$ and $b \in \mathbb{R}$. The recurrence relations also play a central role in the study of mesoscopic fluctuations of Circular OPEs and sparse perturbations of JUE in [\[19,](#page-81-5) [15\]](#page-81-1) respectively.

2.2 Results on Non-Varying Weights

Consider an OPE [\(1\)](#page-1-1) with respect to the real Borel measure μ with finite moments, where the associated recurrence coefficients be a_j and b_j , given by [\(11\)](#page-5-3) and [\(12\)](#page-5-4). Note that the recurrence coefficients do not have a second subscript, since they come from the measure μ , non-varying in *n*.
Recall that $X^{(n)}_{f, \alpha, v}$ is the mesoscopic linear statistics of an OPE for a test fun

 $f_{\alpha,x_0}^{(n)}$ is the mesoscopic linear statistics of an OPE for a test function *f* around the point *x*₀
(b) Δx (2) Now we are going to state our general theorems for non-varying weights which is defined by [\(2\)](#page-2-0). Now we are going to state our general theorems for non-varying weights.

Theorem 2.1. *Consider* $0 < \alpha < 2$ *. Suppose there exist* $a > 0$ *and* $b \in \mathbb{R}$ *such that the recurrence coefficients associated with the OPE satisfy*

$$
a_j = a + O(j^{-\alpha - \varepsilon}), \quad b_j = b + O(j^{-\alpha - \varepsilon})
$$
 (16)

for some $\varepsilon \in (0, 1 - \alpha/2)$ *small, as* $j \to \infty$ *. Then, for any* $f \in C_c^1(\mathbb{R})$ *, as* $n \to \infty$ *, the following converges in distribution distribution*

$$
X_{f,\alpha,x_0}^{(n)} - \mathbb{E}\left[X_{f,\alpha,x_0}^{(n)}\right] \to \mathcal{N}(0,\sigma_f^2),\tag{17}
$$

where

$$
\sigma_f^2 = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy, \quad \text{for } x_0 = b + 2a + o(n^{-\alpha}), \tag{18}
$$

$$
\sigma_f^2 = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy, \quad \text{for } x_0 = b - 2a + o(n^{-\alpha}). \tag{19}
$$

The assumption [\(16\)](#page-6-2) is natural. Note that the Chebyshev polynomials of the second kind have constant recurrence coefficients, cf. Subsection [10.3,](#page-76-0) which comes from the Jacobi unitary ensemble from the random matrix theories. Thus the result applies. More generally, the modified Jacobi polynomials also satisfies [\(16\)](#page-6-2) (cf. [\(343\)](#page-62-1) due to [\[24\]](#page-81-10)). Hence, Theorem [1.1](#page-4-1) is a direct consequence of Theorem [2.1.](#page-6-1) This is elaborated in Section [8.](#page-62-0) Another interesting example is the OPE with a logarithm weight studied by [\[25\]](#page-81-11). The assumption [\(16\)](#page-6-2) is also satisfied in this case, see Subsection [10.6.](#page-78-0) Moreover, the assumption [\(16\)](#page-6-2) with exact the same rate of convergence is proposed by [\[14\]](#page-81-0) to show the bulk mesoscopic universality, i.e., [\(4\)](#page-3-2), in their work.

Also note that the exact edges are at $b + 2a$ and $b - 2a$. Theorem [2.1](#page-6-1) allows x_0 to be close to these edges as long as they have a distance of order at most $o(n^{-\alpha})$. This is also natural, since the mesoscopic linear statistics [\(2\)](#page-2-0) is scaled as n^{α} and any perturbations smaller than the window size should not change the result.

Another remark is that, Theorem [2.1](#page-6-1) also holds for an OPE associated with a varying weight μ_n with varying recurrence coefficients $a_{j,n}$ and $b_{j,n}$ such that

$$
\sup_{j\geq n-n^{\frac{\alpha}{2}+\varepsilon}}|a_{j,n}-a|=O(n^{-\alpha-\varepsilon}), \quad \sup_{j\geq n-n^{\frac{\alpha}{2}+\varepsilon}}|b_{j,n}-b|=O(n^{-\alpha-\varepsilon}).
$$
\n(20)

Though assumption [\(20\)](#page-7-1) is slightly weaker than the assumption [\(16\)](#page-6-2), in most reasonable examples of orthogonal polynomial ensembles, the measures of such are of bounded supports. Typically, there is no scaling with *n* in such cases. Hence, Theorem [2.1](#page-6-1) is stated for non-varying cases. To maintain the consistency of the proofs in this paper, we will use [\(20\)](#page-7-1) in the proof of Theorem [2.1](#page-6-1) in Section [9.](#page-64-0)

2.3 Results on Varying Weights

Now, we state the results about varying weights. Given an OPE [\(1\)](#page-1-1) with respect to the real Borel measure μ_n with finite moments, let the recurrence coefficients associated with measure μ_n be $a_{j,n}$ and $b_{j,n}$ defined by [\(11\)](#page-5-3) and [\(12\)](#page-5-4).

Let $0 < \alpha < 2$ and $\varepsilon \in (0, 1 - \frac{\alpha}{2})$ small. Define $I_n^{(\alpha,\varepsilon)}$ to be an indexing set

$$
I_n^{(\alpha,\varepsilon)} := \{ j \in \mathbb{N} : n - n^{\frac{\alpha}{2} + \varepsilon} \le j \le n + n^{\frac{\alpha}{2} + \varepsilon} \},\tag{21}
$$

In Theorems [2.2](#page-8-1) and [2.3,](#page-8-0) we will assume the recurrence coefficients are slowly varying as the follows.

Condition 2.1. *There exist absolute constants* $c_0, c_1 > 0$ *such that, for all* $j \in I_n^{(\alpha,\varepsilon)}$ *(where* $I_n^{(\alpha,\varepsilon)}$ *is defined* $as (21)$) and $n \in \mathbb{N}$ *as* [\(21\)](#page-7-2)*)* and $n \in \mathbb{N}$,

$$
c_0 < |a_{j,n}| < c_1, \quad |b_{j,n}| < c_1,\tag{22}
$$

$$
|a_{j,n} - a_{j-1,n}| \le \frac{c_1}{n}, \quad |b_{j,n} - b_{j-1,n}| \le \frac{c_1}{n}.
$$
 (23)

The parameter α is the same as the "scaling" parameter in the mesoscopic linear statistics. Condition [2.1](#page-7-3) means that we only assume the recurrence coefficients of order around *n* with a window size of $n^{\alpha/2+\epsilon}$ are bounded and slowly varying. As we will show, to prove the mesoscopic limit of the linear statistics, it is enough to consider recurrence coefficients of orders only inside this window. Many interesting examples in random matrix theories satisfy this condition and it is not hard to verify this. The recurrence coefficients from Theorem [1.2](#page-4-0) will satisfy this condition, and so are those from the classical (hypergeometric) orthogonal polynomials along the Askey scheme when scaled (cf. [\[22\]](#page-81-8)).

We will zoom in around a point x_0 , that may depend n , such that

$$
x_0 = b_{n-1,n} - 2\sqrt{a_{n,n}a_{n-1,n}} + o(n^{-\alpha})
$$
 on the left, or $x_0 = b_{n-1,n} + 2\sqrt{a_{n,n}a_{n-1,n}} + o(n^{-\alpha})$, on the right. (24)

Hence, for varying weights, instead of *b* ± 2*a* in Theorem [2.1,](#page-6-1) we centre x_0 around the points $b_{n-1,n} \pm 2\sqrt{a_n^2 + a_n^2}$ 2 $\sqrt{a_{n,n}a_{n-1,n}}$.

Now we are going to state our general theorems for varying weights. Recall that $X_{f,q}^{(n)}$ $f_{, \alpha, x_0}^{(n)}$, defined in [\(2\)](#page-2-0), is the mesoscopic linear statistics of an OPE for a test function f around the point x_0 .

Theorem 2.2. *Consider* $0 < \alpha < \frac{2}{3}$ *. Assume there is an* $\varepsilon \in (0, 1 - \frac{\alpha}{2})$ *such that Condition* [2.1](#page-7-3) *is satisfied for* $all \, j \in I_n^{(\alpha,\varepsilon)}$ and $n \in \mathbb{N}$. Then, for any $f \in C_c^1(\mathbb{R})$, the following converges in distribution as $n \to \infty$

$$
X_{f,\alpha,x_0}^{(n)} - \mathbb{E}\left[X_{f,\alpha,x_0}^{(n)}\right] \to \mathcal{N}(0,\sigma_f^2),\tag{25}
$$

where

$$
\tau_f^2 = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy, \quad \text{for } x_0 = b_{n-1,n} + 2\sqrt{a_{n,n} a_{n-1,n}} + o(n^{-\alpha}), \tag{26}
$$

$$
\sigma_f^2 = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy, \quad \text{for } x_0 = b_{n-1,n} - 2\sqrt{a_{n,n} a_{n-1,n}} + o(n^{-\alpha}). \tag{27}
$$

Theorem [2.2](#page-8-1) only deals with the case $0 < \alpha < \frac{2}{3}$. For soft edges with square root decay, this is optimal. But for hard edges, the result should also hold for $0 < \alpha < 2$. We will show that this is true under additional assumptions. Indeed, Theorems [1.1,](#page-4-1) [1.2](#page-4-0) and [2.2](#page-8-1) are special cases of the following.

Theorem 2.3. *Consider* $0 < \alpha < 2$. Assume there is an $\varepsilon \in (0, 1 - \frac{\alpha}{2})$ such that the Condition [2.1](#page-7-3) is satisfied *for all* $j \in I_n^{(\alpha,\varepsilon)}$ *and* $n \in \mathbb{N}$ *. Also assume the following holds, as* $n \to \infty$

$$
\max_{j \in I_n^{(\alpha,\varepsilon)}} \left| a_{j,n} a_{j-2,n} - a_{j-1,n}^2 \right| = o(n^{-\alpha-\varepsilon}),\tag{28}
$$

$$
\max_{j \in I_n^{(\alpha,\varepsilon)}} \left| (b_{j-1,n} - x_0 - a_{j,n}) a_{j-2,n} - (b_{j-2,n} - x_0 - a_{j-1,n}) a_{j-1,n} \right| = o(n^{-\frac{3\alpha}{2} - \varepsilon}).
$$
\n(29)

Then, for any $f \in C_c^1(\mathbb{R})$ *, as* $n \to \infty$ *, the following converges in distribution*

$$
X_{f,\alpha,x_0}^{(n)} - \mathbb{E}\left[X_{f,\alpha,x_0}^{(n)}\right] \to \mathcal{N}(0,\sigma_f^2),\tag{30}
$$

where σ_f^2 *is given by* [\(26\)](#page-8-2) *and* [\(27\)](#page-8-3)*.*

Note that the Condition [2.1](#page-7-3) implies that [\(28\)](#page-8-4) and [\(29\)](#page-8-5) hold for all $\alpha \in (0, \frac{2}{3})$
ly take effect when $\alpha > \frac{2}{3}$. In other words, Theorem 2.2 is a direct consequent $\frac{2}{3}$). Hence, these two assumptions only take effect when $\alpha > \frac{2}{3}$. In other words, Theorem [2.2](#page-8-1) is a direct consequence of Theorem [2.3.](#page-8-0)
These conditions are technical appearing in the proof. To understand where they come from mo

These conditions are technical appearing in the proof. To understand where they come from, more background is needed. We will postpone the explanations to the end of Subsection [3.5.](#page-20-0)

There are two examples that motivate conditions [\(28\)](#page-8-4) and [\(29\)](#page-8-5). First, it is shown in [\[24\]](#page-81-10) that the recurrence coefficients of the modified Jacobi polynomials with measure [\(3\)](#page-3-1) satisfies a stronger sense of slowly varying, i.e.,

$$
|a_j - a_{j-1}| = O(j^{-3}),
$$
 $|b_j - b_{j-1}| = O(j^{-3}),$ as $j \to \infty.$

Note that there is no scaling in this model, and we remove the second subscription *n* from the notations. Hence, the conditions [\(28\)](#page-8-4) and [\(29\)](#page-8-5) are satisfied for all $0 < \alpha < 2$. The limit of mesoscopic fluctuations holds for all $\alpha \in (0, 2)$, .i.e., Theorem [1.1.](#page-4-1) For details, see proof of Theorem [1.1](#page-4-1) in Section [8.](#page-62-0)

Another motivation of the conditions is the Laguerre ensemble, where there is a hard edge on the left (at 0). In this case, we have $a_{j,n} = \sqrt{j(j + \gamma)}/n$, $b_{j,n} = (2j + \gamma + 1)/n$ and $x_0 = 0$. Taking $\gamma = 0$, it is easy to check that assumption (28) holds for all $0 \le \alpha \le 2$ and the left hand side of 29 vanishes for all *i*. Hence check that assumption [\(28\)](#page-8-4) holds for all $0 < \alpha < 2$ and the left-hand side of [29](#page-8-5) vanishes for all *j*. Hence, Theorem [2.1](#page-6-1) applies and we obtain the the limit of mesoscopic fluctuations for all $\alpha \in (0, 2)$ at the hard (left) edge. As is observed in the Laguerre case, assumptions in Theorem [2.3](#page-8-0) reveals that although the recurrence coefficients are not slowly varying in a stronger sense, cancellation exists that leads to an order reduction. For general $\gamma > 0$, Theorem [2.3](#page-8-0) also applies, see Subsection [10.1](#page-75-0) for details.

2.4 Overview of the Rest of the Paper

The rest of the paper is organised as the following.

In the Preliminaries, Section [3,](#page-9-1) we introduce the cumulant expansion of OPEs. We will express it in terms of recurrence coefficients. Special care is needed in the case where the moment problem is indeterminate. We will also present the main tool of analysis, an estimate of a three-diagonal matrix with slowly varying entries and explain why the usual Combes-Thomas estimate is insufficient for our setup.

The varying weights are more difficult to prove than the non-varying ones. Hence, most of the content of this paper is to develop techniques for the varying weights, the proof of which we will show first.

In Section [4,](#page-23-0) we provide the proof of Theorem [2.3.](#page-8-0) We illustrate the essence of proof by demonstrating several key propositions. The proofs of these key propositions are deferred in later Sections [5](#page-30-0) [6](#page-44-0) and [7.](#page-54-0)

In Section [8,](#page-62-0) we explore the conditions of Theorem [2.3](#page-8-0) and subsequently prove Theorems [1.1,](#page-4-1) [2.2](#page-8-1) and [1.2,](#page-4-0) in that specific order.

Theorem [2.1](#page-6-1) requires a different strategy of proof than the other theorems, since it is essentially a result of non-varying weights. It is explained and proved in Section [9.](#page-64-0)

In Section [10,](#page-74-0) we give several examples that Theorems [2.1](#page-6-1) [2.2](#page-8-1) [2.3](#page-8-0) are applicable. There includes examples of continuous and discrete weights.

3 Preliminaries

In this section, we will introduce some notations and recall some preliminary facts regarding cumulants for linear statistics. We will follow the approach proposed by Breuer and Duits, [\[14,](#page-81-0) [23\]](#page-81-9). They are the first to discover that the cumulants of the linear statistics can be expressed in terms of the Jacobi (semi-finite) matrix corresponding to the measure μ_n that defines the OPE. This discovery successfully leads to several results about linear statistics of OPEs, regrading both global and mesoscopic scales. This approach can also be applied to extended-OPEs that have extra parameters indicating the time transition, that rises naturally in many random tiling models, see [\[26,](#page-81-12) [27\]](#page-81-13).

We will also introduce the Combes-Thomas estimate, which is an essential element proving the mesoscopic limit in the bulk, [\[14\]](#page-81-0). However, as we will explain in this section, at the edges, it is too rough and its improvement is required. To this end, we will introduce a formula for symmetric tri-diagonal matrices, that is well-known, but a key element for what follows.

3.1 Some Notations

We start this section by some notations that we will use, for a general reference see [\[28\]](#page-81-14). For a compact operator *A* on a (separable) Hilbert space, we denote the singular values by $\sigma_i(A)$, which are the square roots of the eigenvalues of the compact self-adjoint operator *A* [∗]*A*. Then we define

- 1. $||A||_{\infty}$ ≔ sup_{*j*} σ *j*(*A*) to be the operator norm,
- 2. $||A||_1 := \sum_j \sigma_j(A)$ to be the trace norm,
- 3. $||A||_2 := \sqrt{\sum_j \sigma_j(A)^2}$ to be the Hilbert-Schmidt norm.

Then we have the following inequalities

1. For $j = 1, 2, \infty$,

$$
||AB||_j \le ||A||_j ||B||_{\infty}, \quad ||AB||_j \le ||A||_{\infty} ||B||_j,
$$

$$
||AB||_1 \le ||A||_2 ||B||_2.
$$

2. If *A* is trace class,

 $| \text{Tr} A | \leq ||A||_1.$

We will frequently view a semi-finite matrix $A = ((A)_{i,j})$ *i*, *j*∈N with entries $(A)_{i,j}$ ∈ ℂ, as an operator on l^2 (N). Then

$$
||A||_{\infty} \le \sum_{j=-\infty}^{\infty} \sup_{k} |(A)_{k,k+j}|,
$$

$$
||A||_{1} \le \sum_{i,j=1}^{\infty} |(A)_{i,j}|,
$$

$$
||A||_{2} = \left(\sum_{i,j=1}^{\infty} |(A)_{i,j}|^{2}\right)^{1/2}.
$$

If *A* is further Hermitian, $\eta \in \mathbb{C}$ with Im $\eta \neq 0$ and *Id* is the identity operator, the following holds

$$
||(A - \eta Id)^{-1}||_{\infty} \le \frac{1}{|\operatorname{Im} \eta|}.
$$
 (31)

Note that among many trace norm inequalities, the one above may not be the optimal nor elegant one. However, in our cases, it is sufficient.

3.2 Bounded and Unbounded Jacobi Operators

Let the Jacobi matrix $\mathcal J$ be an semi-finite tri-diagonal matrix associated with the measure μ_n with entries being the recurrence coefficients defined as [\(11\)](#page-5-3) and [\(12\)](#page-5-4), i.e.,

$$
\mathcal{J} := \begin{pmatrix} b_{0,n} & a_{1,n} \\ a_{1,n} & b_{1,n} & a_{2,n} \\ a_{2,n} & b_{2,n} & a_{3,n} \\ a_{3,n} & b_{3,n} & a_{4,n} \\ \vdots & \vdots & \ddots & \vdots \end{pmatrix}.
$$
 (32)

Such $\mathcal I$ is also called the Jacobi operator associated with the measure μ_n , viewed as a linear operator from $l^2(\mathbb{N})$ to $l^2(\mathbb{N})$ acting on the space of finite sequences. It is a bounded linear operator only if both sequences ${a_{j,n}}$ *j* and ${b_{j,n}}$ *j* are bounded. For example, J is bounded for the Chebyshev polynomials where the recurrence coefficients are constants. However, in many cases $\mathcal T$ is not necessarily a bounded operator. For example, \mathcal{J} is unbounded for the Hermite and Laguerre polynomials where $a_{i,n}$ tends to infinity as $j \to \infty$. These examples are included in Section [10.](#page-74-0) Breuer and Duits studied the resolvent of bounded \mathcal{J} and they successfully obtained the mesoscopic fluctuations in the bulk for the non-varying cases [\[14\]](#page-81-0). However, though $\mathcal J$ is symmetric with all entries real numbers, its resolvent is only well-defined if $\mathcal J$ is a self-adjoint operator. However, $\mathcal J$ is essentially self-adjoint, if and only if the measure μ_n is fully characterized by its moments, see Theorems 6.10. and 6.16 in [\[29\]](#page-81-15). In terms of recurrence coefficients, that is if $\sum_{j\geq 1} a_{j,n}^{-1}$ $j_{n}^{-1} = +\infty$,

the moment problem is determinant. If $\{b_{j,n}\}_j$ is bounded, $a_{j-1,n}a_{j+1,n} \le a_{j,n}^2$ for all $j \ge j_0$ for some $j_0 \in \mathbb{N}$ and $\sum_{j\geq 1} a_{j,n}^{-1} < +\infty$, the moment problem is indeterminate. See Corollary 6.19. in [\[29\]](#page-81-15) and Theorem 1.5 in *j*,*n* [\[30\]](#page-81-16). For example, for Freud weight $d\mu_n(x) = e^{-n|x|^{\gamma}} dx$, $\gamma > 0$, shown in [\[31,](#page-82-0) [32\]](#page-82-1), the recurrence coefficients have the asymptotics as

$$
a_{j,n} = c_\gamma \left(\frac{j}{n}\right)^{\frac{1}{\gamma}} (1 + O(j^{-1})), \text{ as } j \to \infty, \text{ and } b_{j,n} = 0,
$$
 (33)

where c_y is an universal constant only depending on γ . For a precise statement, see Section [10.4.](#page-77-0) Hence, its Jacobi operator is essentially self-adjoint if and only if $\gamma \geq 1$.

In the rest of the paper, we will avoid calling a general J an operator and only treat it as an infinite tri-diagonal matrix, which is always well-defined as [\(32\)](#page-10-1). We will construct a bounded linear operator from $l^2(\mathbb{N})$ to $l^2(\mathbb{N})$ associated with \mathcal{J} , that will be suitable for our further analysis.

Let $P(x)$ be the column vector of the orthonormal polynomials

$$
P(x) = (p_{0,n}(x), p_{1,n}(x), \cdots)^T,
$$
\n(34)

where the superscript T is the transpose of a vector. We define a semi-finite matrix

$$
G_z := \int_{\mathbb{R}} (x - z)^{-1} P(x) P(x)^T d\mu_n(x), \quad z \in \mathbb{C}, \quad \text{Im } z \neq 0.
$$
 (35)

Note that each entry of G_z is well-defined and bounded, i.e.,

$$
\left| \int_{\mathbb{R}} (x - z)^{-1} p_{j,n}(x) p_{k,n}(x) d\mu(x) \right| \le \frac{1}{|\operatorname{Im} z|} \left| \int_{\mathbb{R}} p_{j,n}(x) p_{k,n}(x) d\mu(x) \right|
$$

$$
\le \frac{1}{|\operatorname{Im} z|} \left| \int_{\mathbb{R}} p_{j,n}(x)^{2} d\mu_{n}(x) \int_{\mathbb{R}} p_{k,n}(x)^{2} d\mu_{n}(x) \right|^{2} = \frac{1}{|\operatorname{Im} z|}, \quad (36)
$$

where we use the fact $|(x-z)^{-1}| \le \frac{1}{|\ln z|}$ for the first inequality, the Cauchy-Schwartz inequality for the second inequality and the normality of the orthonormal polynomials for the last equality.

Lemma 3.1. For any $z \in \mathbb{C}$ with $\text{Im } z \neq 0$, G_z represents a bounded linear operator from $l^2(\mathbb{N})$ to $l^2(\mathbb{N})$, *given the canonical basis in l*² (N)*. Its operator norm is bounded by*

$$
||G_z||_{\infty} \le |\operatorname{Im} z|^{-1}.\tag{37}
$$

Proof. Note that in the case where $\mu_n = \mu$ is non-varying or generally μ_n is varying with its moment problem being determinant, J give a self-adjoint operator and its resolvent is well defined. By the recurrence relations, we have $G_z = (\mathcal{J} - z)^{-1}$. The lemma holds directly.

However, in the case where moment problem is indeterminant, and $\mathcal J$ does not admits an unique selfadjoint extension, a more careful treatment is required.

Define the multiplication operator $M: L^2(\mu_n) \to L^2(\mu_n)$ via $Mh(x) = xh(x)$, for $h \in Dom(M) =: \{h \in L^2(\mu_n): Mh \in L^2(\mu_n)\}$. Hence $(M-z)^{-1}$ exists in particular $(M-z)^{-1}h(x) = (x-z)^{-1}h(x)$ for $h \in L^2(\mu_n)$. $L^2(\mu_n)$: $Mh \in L^2(\mu_n)$. Hence, $(M-z)^{-1}$ exists, in particular, $(M-z)^{-1}h(x) = (x-z)^{-1}h(x)$, for $h \in L^2(\mu_n)$.
The multiplication operator M is self-adjoint and hence its operator porm is bounded by The multiplication operator M is self-adjoint and hence its operator norm is bounded by

$$
\left\| (M - z)^{-1} \right\|_{\infty} \leq |{\rm Im} \, z|^{-1} \,. \tag{38}
$$

Let $e_j = (0, 0, \ldots, 1, 0, 0, \ldots)^T$ be the *j*-th vector of the canonical basis for $l^2(\mathbb{N})$, i.e., the sequence with
ly the *i*-th element to be 1 and the rests are zeros. Then each entry of *G*, equals to only the *j*-th element to be 1 and the rests are zeros. Then, each entry of G_z equals to

$$
(e_j, G_z e_k)_l^2 = \int_{\mathbb{R}} (x - z)^{-1} p_{j-1}(x) p_{k-1}(x) d\mu_n(x)
$$

=
$$
\int_{\mathbb{R}} (\mathcal{M} - z)^{-1} p_{j-1}(x) p_{k-1}(x) d\mu_n(x) = ((\mathcal{M} - z)^{-1} p_{j-1}, p_{k-1})_{L^2(\mu_n)}.
$$
 (39)

Let the subspace $\mathcal{H}_{\mu_n} := \overline{span\{p_{j,n} : j \ge 0\}}^{L^2(\mu_n)}$ be the closure of the the linear space spanned by the or-
thenormal palynomials. Closuly, $L^2(\mu_n) = \mathcal{H} \oplus \mathcal{H}^1$, Let $\mathcal{D} : L^2(\mu_n) \mapsto \mathcal{H}$, he the ortho thonormal polynomials. Clearly, $L^2(\mu_n) = \mathcal{H}_{\mu_n} \oplus \mathcal{H}_{\mu_n}^{\perp}$. Let $\mathcal{P}: L^2(\mu_n) \mapsto \mathcal{H}_{\mu_n}$ be the orthogonal projection operator onto \mathcal{H}_{μ_n} . Define an unitary operator $\mathcal{U}: \mathcal{H}_{\mu_n} \mapsto l^2(\mathbb{N})$ such that $\mathcal{U}_{p,n} = e_{j+1}$. Then [\(39\)](#page-12-0) shows that

$$
G_z = \mathcal{UP}\left(\mathcal{M} - z\right)^{-1} \mathcal{PU}^{-1}.
$$
\n⁽⁴⁰⁾

Therefore, *G^z* is a bounded operator and

$$
||G_z||_{\infty} \le ||(\mathcal{M} - z)^{-1}||_{\infty}.
$$
\n(41)

Hence, $||G_z||_{\infty} \leq |\operatorname{Im} z|^{-1}$. □

Note that the equality of [\(41\)](#page-12-1) holds only if the space $L^2(\mu_n)$ can be spanned by the polynomials $\{p_{j,n}\}\$.
By the three term recurrence relation (11) and (12) G as an infinite matrix defined in (35) is the algebra

By the three-term recurrence relation [\(11\)](#page-5-3) and [\(12\)](#page-5-4), G_z as an infinite matrix defined in [\(35\)](#page-11-0) is the algebraic inverse of \mathcal{J} – *z*, i.e., the following holds entry-wise as an infinite matrix,

$$
((\mathcal{J} - z)G_z)_{j,k} = (G_z(\mathcal{J} - z))_{j,k} = \begin{cases} 1, & j = k, \\ 0, & j \neq k. \end{cases}
$$
 (42)

For any bounded operator $A: l^2 \mapsto l^2$ whose inverse A^{-1} exists and is also a bounded linear operator, the following holds algebraically, i.e., the following holds entry-wise as an infinite matrix,

$$
G_z - A^{-1} = G_z(A - \mathcal{J} + z)A^{-1}.
$$
\n(43)

Note that $\mathcal J$ is a tri-diagonal matrix, and hence $\mathcal JA^{-1}$ is always well-defined as a semi-finite matrix. If further \mathcal{J} − *z* − *A* represents a bounded linear operator from $l^2(\mathbb{N})$ to $l^2(\mathbb{N})$, the right-hand side of [\(43\)](#page-12-2) can be viewed as the operator compositions. In such case, with an abuse of terminology, we call [\(43\)](#page-12-2) to be the resolvent identity.

The bounded operator G_z agrees with the resolvent of $\mathcal J$, whenever $\mathcal J$ is of self-adjoint. In such case, $G_z = (\mathcal{J} - z)^{-1}$.

Corollary 3.1. *Recall that we define* G_z *as* [\(35\)](#page-11-0). *Then, for* $z \in \mathbb{C}$ *with* $\text{Im } z \neq 0$, e^{G_z} *is well defined and admits a formula*

$$
e^{G_z} = \int_{\mathbb{R}} e^{(x-z)^{-1}} P(x) P(x)^T d\mu_n(x).
$$
 (44)

Proof. We will first prove for any $k \in \mathbb{N}$, by induction

$$
G_z^k = \int_{\mathbb{R}} (x - z)^{-k} P(x) P(x)^T d\mu_n(x).
$$
 (45)

Note that by Lemma [3.1,](#page-11-1) [\(45\)](#page-12-3) holds for $k = 1$. Moreover, G_z is a bounded linear operator and so is G_z^k . Then by the uniqueness of the representation it is sufficient to show each entry coincides. Note that by the three-term recurrence relation we have $(x-z)P(x)^T = P(x)^T (J-z)$ and [\(42\)](#page-12-4) algebraically (i.e., to understand the equation entry-wise as infinite matrices). Use the induction hypothesis to obtain the following, which holds algebraically, (i.e., to understand the following equations entry-wise as semi-finite matrices),

$$
G_z^{k+1} = \int_{\mathbb{R}} (x - z)^{-k} P(x) P(x)^T d\mu_n(x) G_z = \int_{\mathbb{R}} (x - z)^{-k-1} P(x) (x - z) P(x)^T d\mu_n(x) G_z
$$

=
$$
\int_{\mathbb{R}} (x - z)^{-k-1} P(x) P(x)^T d\mu_n(x) (J - z) G_z = \int_{\mathbb{R}} (x - z)^{-k-1} P(x) P(x)^T d\mu_n(x).
$$
 (46)

Recall that G_z^{k+1} is a bounded linear operator and the matrix representation of a bounded linear operator is unique. Hence, we conclude [\(45\)](#page-12-3) by induction.

Now we have algebraically, (i.e., to understand the following entry-wise as semi-finite matrices),

$$
e^{G_z} = \sum_{k\geq 0} \frac{G_z^k}{k!} = \sum_{k\geq 0} \frac{1}{k!} \int_{\mathbb{R}} (x - z)^{-k} P(x) P(x)^T d\mu_n(x) = \sum_{k\geq 0} \frac{1}{k!} \int_{\mathbb{R}} (x - z)^{-k} P(x) P(x)^T d\mu_n(x)
$$

$$
= \int_{\mathbb{R}} \sum_{k\geq 0} \frac{1}{k!} (x - z)^{-k} P(x) P(x)^T d\mu_n(x) = \int_{\mathbb{R}} e^{(x - z)^{-1}} P(x) P(x)^T d\mu_n(x). \quad (47)
$$

Note that the fourth equality above is valid due to $\int_{\mathbb{R}} |(x - z)^{-k} p_j(x) p_k(x)| d\mu_n(x) \le |\text{Im } z|^{-k}$ by the same argument as [\(36\)](#page-11-2). By Lemma [3.1,](#page-11-1) e^{G_z} is a bounded linear operator, and we conclude the corollary. \Box

3.3 Cumulants via Recurrence Relations

For a real-valued random variable *X*, whose moments are all finite, the cumulants C*m*(*X*) are uniquely defined by the cumulant generating function, for $t \in \mathbb{C}$,

$$
\log \mathbb{E}[e^{tX}] = \sum_{m \ge 1} \frac{t^m}{m!} C_m(X).
$$

From this expression, the moments of *X* can be recovered from its cumulants and vice versa. For example, the mean and variance of *X* are given by $C_1(X)$ and $C_2(X)$ respectively. Note also that, if *X* follows a Gaussian distribution, we have $\log \mathbb{E}[e^{tX}] = tC_1(X) + t^2C_2(X)/2$.
Hence, to show that the linear statistics $X_{f, \alpha, r}^{(n)}$ of an OPE as defined as [\(2\)](#page-2-0) converges to a Gaussian distri-

bution for a suitable class of test functions *f*, it is sufficient for us to study the asymptotic behaviour of its bution for a suitable class of test functions *f*, it is sufficient for us to study the asymptotic behavi cumulants. First, we consider the test function *f* to be

$$
f(x) = \operatorname{Im} \sum_{r=1}^{M} d_r \frac{1}{x - \lambda_r}
$$
\n(48)

where $M \in \mathbb{N}$, $d_r \in \mathbb{R}$, Im takes the imaginary part of a complex number, and Im $\lambda_r > 0$. We point out that any compactly supported and continuously differentiable function can be well approximated by such *f* in a certain Lipschitz space. This claim will be explained in Lemma [4.1,](#page-28-0) which is a result in [\[14\]](#page-81-0).

Denote λ_r to be the complex conjugate of λ_j . Then *f* can be rewritten as

$$
f(x) = \sum_{r=1}^{2M} c_r \frac{1}{x - \eta_r},
$$

where $c_r := \begin{cases} \frac{d_r}{2i} & r = 1, ..., M \\ -\frac{d_{r-M}}{2i} & r = 1 + M, ..., 2M \end{cases}$ $\eta_r := \begin{cases} \lambda_r & r = 1, ..., M \\ \overline{\lambda}_{r-M} & r = 1 + M, ..., 2M \end{cases}$ (49)

and, for some $x_0 \in \mathbb{R}$, define

$$
f_{\alpha,x_0}^{(n)}(x) \coloneqq f(n^{\alpha}(x - x_0)). \tag{50}
$$

$$
f_{\alpha,x_0}^{(n)}(\mathcal{J}) \coloneqq \int_{\mathbb{R}} f_{\alpha,x_0}^{(n)}(x) P(x) P(x)^T d\mu_n(x) = \sum_{r=1}^{2M} \frac{c_r}{n^{\alpha}} G_{x_0 + \frac{n_r}{n^{\alpha}}},\tag{51}
$$

where $G_{x_0+\frac{\eta_r}{n^a}}$ is defined as [\(35\)](#page-11-0). Note that $f_{\alpha,x_0}^{(n)}(\mathcal{J})$ is not necessarily a self-adjoint operator on $l^2(\mathbb{N})$. However, this is true if and only if the measure μ_n is determined by its moments. Nevertheless, similar to the properties of G in general, the following holds for $f^{(n)}$ (σ) the properties of G_z , in general, the following holds for $f_{\alpha,x_0}^{(n)}(\mathcal{J})$.

Corollary 3.2. $f_{\alpha,x_0}^{(n)}(\mathcal{J})$ *is a well defined infinite matrix and entries are real and symmetric, i.e.,*

$$
\left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)_{j,k} = \left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)_{k,j} = \overline{\left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)_{k,j}}, \quad j,k \in \mathbb{N},\tag{52}
$$

where the overline is the complex conjugate. Moreover, $f_{\alpha,x_0}^{(n)}(\mathcal{J})$ *represents a bounded linear operator from*
 $l^2(\mathbb{N})$ *to* $l^2(\mathbb{N})$. Its aparator norm is estimated by $l^2(\mathbb{N})$ *to* $l^2(\mathbb{N})$ *. Its operator norm is estimated by*

$$
\left\|f_{\alpha,x_0}^{(n)}(\mathcal{J})\right\|_{\infty} \le \sum_{r=1}^{2M} \left|\frac{c_r}{\text{Im}\,\eta_r}\right|.
$$
\n(53)

Furthermore, the bounded linear operator $e^{f_{\alpha,x_0}^{(n)}(\mathcal{J})}$ is also well-defined with matrix representation, under the *canonical basis of* $l^2(\mathbb{N})$,

$$
e^{f_{\alpha,x_0}^{(n)}}(\mathcal{J}) = \int_{\mathbb{R}} e^{f_{\alpha,x_0}^{(n)}(x)} P(x) P(x)^T d\mu_n(x).
$$
 (54)

Proof. By definition [\(48\)](#page-13-1) and [\(51\)](#page-14-0), it is clear that $f_{\alpha,x_0}^{(n)}(\mathcal{J})$ has real and symmetric entries. By Lemma [3.1](#page-11-1) *rroof.* By definition (48) and [\(51\)](#page-14-0), it is clear that $f_{\alpha,x_0}(J)$ has real and symmetric entries. By Lemma 5.1 and the definition (51), $f_{\alpha,x_0}^{(n)}(J) : l^2(\mathbb{N}) \mapsto l^2(\mathbb{N})$ is a bounded linear functional, due to line and the definition (51), $f_{\alpha,x_0}(J)$: ι (18) $\mapsto \iota$ (18) is a bounded linear functional, due to linearity. The setimate [\(53\)](#page-14-1) follows form triangle inequality and Lemma [3.1.](#page-11-1) The second statement [\(54\)](#page-14-2) follows from th same argument as Corollary [3.1.](#page-12-5) □

Breuer and Duits obtained a beautiful formula of the cumulant generating function of the linear statistics in terms of J , see the preliminaries of [\[14\]](#page-81-0). They showed the formula for J being a bounded infinite matrix. By Corollary [3.2,](#page-14-3) we now extend their result to general J , which can be unbounded.

Lemma 3.2 (Breuer-Duits). *The cumulant generating function for the linear statistics* $X_{f,a}^{(n)}$ *^f*,α,*x*⁰ *of an OPE has the following determinantal structure,*

$$
\log \mathbb{E}\left[\exp\left(tX_{f,\alpha,x_0}^{(n)}\right)\right] = \log \det\left(id + P_n\left(e^{tf_{\alpha,x_0}^{(n)}(\mathcal{J})} - Id\right)P_n\right),\tag{55}
$$

where $f_{\alpha,x_0}^{(n)}(\mathcal{J})$ *is defined in* [\(51\)](#page-14-0), *Id is the identity operator and* P_n *is a cardinal projection onto the first n*
coordinates *coordinates.*

Proof. Note that the Vandermonde determinant admits the following formula

$$
\prod_{1 \le i < j \le n} (x_i - x_j) = \frac{1}{\prod_{j=1}^n \gamma_{j,n}} \det(p_{j-1,n}(x_i))_{i,j=1}^n \tag{56}
$$

where $p_{j,n}$ is the orthonormal polynomial and $\gamma_{j,n}$ is its leading coefficients.

Hence, use [\(1\)](#page-1-1) to obtain that

$$
\mathbb{E}\left[\exp\left(tX_{f,\alpha,x_0}^{(n)}\right)\right] = \frac{1}{Z_n}\int\cdots\int\prod_{i=n}^n e^{tf_{\alpha,x_0}^{(n)}(x_i)} \left(\det(p_{j-1,n}(x_i))_{i,j=1}^n\right)^2 \mu_n(x_1)\cdots d\mu_n(x_n)
$$

$$
= \frac{1}{Z_n}\int\cdots\int \det(e^{tf_{\alpha,x_0}^{(n)}(x_i)} p_{j-1,n}(x_i))_{i,j=1}^n \det(p_{j-1,n}(x_i))_{i,j=1}^n \mu_n(x_1)\cdots d\mu_n(x_n), \quad (57)
$$

where $Z_n > 0$ is some normalising constant. Recall the Andreief's identity, for a measure μ and measurable functions $f_j, g_j \in L^2(\mu)$ for $j = 1, ..., n$,

$$
\int \cdots \int \det \left(f_j(x_k) \right)_{j,k=1}^n \det \left(g_j(x_k) \right)_{j,k=1}^n d\mu(x_1) \cdots d\mu(x_n) = n! \det \left(\int f_j(x) g_k(x) d\mu(x) \right)_{j,k=1}^n. \tag{58}
$$

Then,

$$
\mathbb{E}\left[\exp\left(tX_{f,\alpha,x_0}^{(n)}\right)\right] = \frac{n!}{Z_n} \det\left(\int e^{tf_{\alpha,x_0}^{(n)}(x)} p_{j-1,n}(x) p_{i-1,n}(x) d\mu_n(x)\right)_{j,k=1}^n.
$$
 (59)

Rewrite it in the matrix form and use Corollary [3.2](#page-14-3) to obtain

$$
\mathbb{E}\left[\exp\left(tX_{f,\alpha,x_0}^{(n)}\right)\right] = \frac{n!}{Z_n} \det\left(P_n e^{tf_{\alpha,x_0}^{(n)}(\mathcal{J})} P_n + Q_n\right). \tag{60}
$$

where P_n is the canonical projection onto the first *n* coordinates.

In particular, take $t = 0$ to obtain $\mathbb{E}[1] = \frac{n!}{\mathbb{Z}}$ $\frac{n!}{Z_n}$ det (*P_n* + *Q_n*). Hence $Z_n = n!$. This concludes the Lemma. □

Breuer and Duits also computed the cumulants in Sections 2.2 and 2.3 of [\[14\]](#page-81-0), which we extend and formulate as the following lemma. We also give the proof for completeness.

Lemma 3.3 (Breuer-Duits). *The cumulants for the linear statistics* $X_{f,a}^{(n)}$ *^f*,α,*x*⁰ *of an OPE can be expressed to be*

$$
C_1(X_{f,\alpha,x_0}^{(n)}) = \mathbb{E}[X_{f,\alpha,x_0}^{(n)}] = \text{Tr}\left(P_n f_{\alpha,x_0}^{(n)}(\mathcal{J}) P_n\right). \tag{61}
$$

For $m \geq 2$

$$
C_m(X_{f,\alpha,x_0}^{(n)}) =
$$
\n
$$
m! \sum_{j=2}^m \frac{(-1)^{j+1}}{j} \sum_{l_1+\dots+l_j=m, l_i\geq 1} \frac{\text{Tr}\left(P_n\left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)^{l_1} P_n\left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)^{l_2} P_n \dots \left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)^{l_j} P_n\right) - \text{Tr}\left(\left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)^m P_n\right)}{l_1! \cdots l_j!}.
$$
\n(62)

In particular,

$$
C_2(X_{f,\alpha,x_0}^{(n)}) = \text{Var}[X_{f,\alpha,x_0}^{(n)}] = \text{Tr}\left(P_n f_{\alpha,x_0}^{(n)}(\mathcal{J})Q_n f_{\alpha,x_0}^{(n)}(\mathcal{J})P_n\right). \tag{63}
$$

Proof. Note that for any non-singular square matrix *A*, we have log det *A* = Tr log *A*. Similarly,

$$
\log \det \left(Id + P_n \left(e^{tf_{\alpha,x_0}^{(n)}} \mathcal{J} - Id \right) P_n \right) = \operatorname{Tr} \log \left(Id + P_n \left(e^{tf_{\alpha,x_0}^{(n)}} \mathcal{J} - Id \right) P_n \right). \tag{64}
$$

Then for *t* sufficiently small, we expand the log and exp and reorder the summations to obtain,

$$
\log \mathbb{E} \left[\exp \left(t X_{f,\alpha,x_0}^{(n)} \right) \right] = \text{Tr} \sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{j} \left(P_n \left(e^{t f_{\alpha,x_0}^{(n)}(\mathcal{J})} - Id \right) P_n \right)^j
$$
\n
$$
= \text{Tr} \sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{j} \sum_{l_1 \ge 1, l_2 \ge 1, \dots, l_j \ge 1} \frac{t^{l_1 + l_2 + \dots + l_j}}{l_1! l_2! \dots l_j!} P_n \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_1} P_n \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_2} P_n \dots \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_j} P_n
$$
\n
$$
= \sum_{j=1}^{\infty} \sum_{m=j}^{\infty} t^m \frac{(-1)^{j+1}}{j} \sum_{l_1 + \dots + l_j = m, l_j \ge 1} \frac{\text{Tr} \left(P_n \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_1} P_n \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_2} P_n \dots \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_j} P_n \right)}{l_1! l_2! \dots l_j!}
$$
\n
$$
= \sum_{j=1}^{\infty} \sum_{m=j}^{\infty} t^m \frac{(-1)^{j+1}}{j} \sum_{l_1 + \dots + l_j = m, l_j \ge 1} \frac{\text{Tr} \left(P_n \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_1} P_n \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_2} P_n \dots \left(f_{\alpha,x_0}^{(n)}(\mathcal{J}) \right)^{l_j} P_n \right)}{l_1! l_2! \dots l_j!}
$$
\n
$$
= \sum_{m=1}^{\infty} t^m \sum_{j=1}^m \frac{(-1)^{j+1}}{j} \sum_{l_1 + \dots
$$

Then the cumulants of the linear statistics can be written in the following way, by expanding the right-hand side of [\(65\)](#page-16-0)

$$
C_m(X_{f,\alpha,x_0}^{(n)}) = m! \sum_{j=1}^m \frac{(-1)^{j+1}}{j} \sum_{l_1 + \dots + l_j = m, l_i \ge 1} \frac{\text{Tr}\left(P_n\left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)^{l_1} P_n\left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)^{l_2} P_n \dots \left(f_{\alpha,x_0}^{(n)}(\mathcal{J})\right)^{l_j} P_n\right)}{l_1! \cdots l_j!}.
$$
 (66)

Similarly to [\(65\)](#page-16-0), by expanding the logarithm and exponential and reordering the summations, we obtain

$$
\log(1 + (e^x - 1)) = \sum_{j=1}^{\infty} \frac{(-1)^j}{j} \sum_{l_1 \ge 1, l_2 \ge 1, \dots, l_j \ge 1} \frac{x^{l_1 + \dots + l_j}}{l_1! \cdots l_j!} = \sum_{j=1}^{\infty} \sum_{m=j}^{\infty} \frac{(-1)^j}{j} \sum_{l_1 + \dots + l_j = m, l_i \ge 1} \frac{x^m}{l_1! \cdots l_j!}
$$

$$
= \sum_{m=1}^{\infty} x^m \sum_{j=1}^m \frac{(-1)^j}{j} \sum_{l_1 + \dots + l_j = m, l_i \ge 1} \frac{1}{l_1! \cdots l_j!},
$$

for all $x \in \mathbb{R}$. Note that $log(1 + (e^x - 1)) = x$. Hence, $\sum_{j=1}^{m}$ $(-1)^{j}$ $\frac{1}{j} \sum_{l_1 + \dots + l_j = m, l_i \ge 1} \frac{1}{l_1! \dots l_j!} = 0$ for all $m \ge 2$. Then, for $m \ge 2$, [\(66\)](#page-16-1) can be rewritten to be [\(62\)](#page-15-0) □

For any linear operator A , we define

$$
C_m^{(n)}(\mathcal{A}) := m! \sum_{j=2}^m \frac{(-1)^{j+1}}{j} \sum_{l_1 + \dots + l_j = m, l_i \ge 1} \frac{\text{Tr}(\mathcal{A})^{l_1} P_n \dots (\mathcal{A})^{l_j} P_n - \text{Tr}(\mathcal{A}^m P_n)}{l_1! \dots l_j!}.
$$
 (67)

The extra term $Tr(\mathcal{A}^m P_n)$ enables us to have good estimates of the cumulants. We will use this extensively to show that the cumulants of the linear statistics of an OPE only depends on the recurrence coefficients of order around *n*, given assumptions of Theorem [2.3](#page-8-0) (cf. Proposition [4.4\)](#page-27-1). Another consequence is the following lemma, which is a by-product of Lemma 2.2 in [\[33\]](#page-82-2). We will state it in terms of linear operators and give a proof for completeness.

Lemma 3.4 (Breuer-Duits). *For a bounded linear operator* A*, whose matrix representations under the carnonial basis of l*² (N) *is such that*

$$
(\mathcal{A})_{j,k} = \overline{(\mathcal{A})_{j,k}}, \quad j,k \in \mathbb{N}.
$$

Then we have for $m \geq 2$

$$
\left| C_m^{(n)}(\mathcal{A}) \right| \leq \sqrt{\frac{2}{\pi}} m! m^{3/2} \|\mathcal{A}\|_{\infty}^{m-2} e^m C_2^{(n)}(\mathcal{A}). \tag{69}
$$

In particular, for $|t| < (e \|\mathcal{A}\|_{\infty})^{-1}$, there exists some constant $c > 0$ such that

$$
\sum_{m\geq 2} \left| \frac{t^m}{m!} C_m^{(n)}(\mathcal{A}) \right| \leq ct^2 C_2^{(n)}(\mathcal{A}).\tag{70}
$$

Note that in the original work, this lemma holds for A to be self-adjoint. However, it is clear in the proof that we can relax it to the condition [\(68\)](#page-17-0), which we will show below.

Proof. Recall the definition of a commutator of two operators $[A, B] = AB - BA$.

For $m = 2$ the estimate [\(69\)](#page-17-1) holds trivially.

For $m \geq 3$ Let $Q_n = Id - P_n$. We write

$$
\mathcal{A}^m P_n = \mathcal{A}^{l_1}(P_n + Q_n)\mathcal{A}^{l_2}(P_n + Q_n)\cdots \mathcal{A}^{l_j}P_n.
$$

By expanding the formula above, we find

$$
\mathcal{A}^{l_1} P_n \cdots \mathcal{A}^{l_j} P_n - \mathcal{A}^m P_n = -\mathcal{A}^{l_1} Q_n \mathcal{A}^{l_2} P_n \cdots \mathcal{A}^{l_j} P_n - \mathcal{A}^{l_1 + l_2} Q_n \mathcal{A}^{l_3} P_n \cdots \mathcal{A}^{l_j} P_n
$$

$$
- \mathcal{A}^{l_1 + l_2 + l_3} Q_n \mathcal{A}^{l_4} P_n \cdots \mathcal{A}^{l_j} P_n - \cdots - \mathcal{A}^{l_1 + \cdots + l_{j-1}} Q_n \mathcal{A}^{l_j} P_n. \tag{71}
$$

Using the cyclic property of the trace, we get

$$
\operatorname{Tr}\left(\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{A}^m P_n\right)=-\sum_{k=2}^j\operatorname{Tr}\left(\mathcal{A}^{l_{k+1}}P_n\cdots P_n\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\mathcal{A}^{l_k}P_n\right).
$$
(72)

Note that since $P_nP_n = P_n$ and $Q_n = Id - P_n$, we have, by the definition of a commutator, $BQ_nAP_n =$ [−][*B*, *^Pn*][*A*, *^Pn*]*Pn*. Therefore,

$$
P_n \mathcal{A}^{l_1 + \dots + l_{k-1}} Q_n \mathcal{A}^{l_k} P_n = -[\mathcal{A}^{l_1 + \dots + l_{k-1}}, P_n] [\mathcal{A}^{l_k}, P_n] P_n. \tag{73}
$$

Plug [\(73\)](#page-17-2) into [\(72\)](#page-17-3), use the trace norm inequality $||ABC||_1 \le ||A||_{\infty} ||B||_2 ||C||_2$ to obtain

$$
\left|\operatorname{Tr}\left(\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{A}^m P_n\right)\right|\leq\sum_{k=2}^j\left\|\mathcal{A}\right\|_{\infty}^{l_{k+1}+\cdots+l_j}\left\|\left[\mathcal{A}^{l_1+\cdots+l_{k-1}},P_n\right]\right\|_2\left\|\left[\mathcal{A}^{l_k},P_n\right]\right\|_2.\tag{74}
$$

For any $l \in \mathbb{N}$, by writing

$$
[\mathcal{A}^l, P_n] = \sum_{j=1}^l \mathcal{A}^{l-j}[\mathcal{A}, P_n] \mathcal{A}^{j-1},\tag{75}
$$

we see that $\left\| \left[\mathcal{A}^l, P_n \right] \right\|_2 \leq l \left\| \mathcal{A} \right\|_{\infty}^{l-1} \left\| \left[\mathcal{A}, P_n \right] \right\|_2$. Given $l_1 + \cdots + l_j = m$, we have, from [\(74\)](#page-17-4),

$$
\left|\operatorname{Tr}\left(\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{A}^m P_n\right)\right|\leq (j-1)m^2\|\mathcal{A}\|_{\infty}^{m-2}\left\|\left[\mathcal{A},P_n\right]\right\|_{2}^{2}.
$$
\n(76)

Hence, we have for $m \geq 3$

$$
\left| C_m^{(n)}(\mathcal{A}) \right| \le m! m^2 ||\mathcal{A}||_{\infty}^{m-2} ||[\mathcal{A}, P_n]||_2^2 \sum_{j=2}^m \sum_{l_1 + \dots + l_j = m, l_i \ge 1} \frac{1}{l_1! \dots l_j!}.
$$
 (77)

Now, by expanding $m^m = (1 + \cdots + 1)^m$ and Stirling's approximation

$$
\sum_{j=2}^{m} \sum_{l_1 + \dots + l_j = m, l_i \ge 1} \frac{1}{l_1! \dots l_j!} < \frac{m^m}{m!} \le \frac{e^m}{\sqrt{2\pi m}}.\tag{78}
$$

Note that by the definition of commutator,

$$
C_2^{(n)}(\mathcal{A}) = \text{Tr}\,\mathcal{A}Q_n\mathcal{A}P_n = \frac{1}{2}\,\text{Tr}[\mathcal{A}, P_n][P_n, \mathcal{A}].\tag{79}
$$

Since A satisfies [\(68\)](#page-17-0), $P_n \mathcal{A}^2 P_n$ and $P_n \mathcal{A} P_n \mathcal{A} P_n$ are self-adjoint. We have $C_2^{(n)}$ $_{2}^{(n)}(\mathcal{A}) = \frac{1}{2}$ $\frac{1}{2}$ ||[\mathcal{A}, P_n]|| $\frac{2}{2}$. There-fore, plug [\(78\)](#page-18-1) into [\(77\)](#page-18-2) to obtain that, for $m \geq 3$,

$$
C_m^{(n)}(\mathcal{A}) \le \sqrt{\frac{2}{\pi}} m! m^{3/2} \|\mathcal{A}\|_{\infty}^{m-2} e^m C_2^{(n)}(\mathcal{A}).
$$
\n(80)

Note that, $\sum_{m\geq 2} m^{3/2} ||\mathcal{A}||_{\infty}^{m-2} e^m t^{m-2} < \infty$, for $|t| < (e ||\mathcal{A}||_{\infty})^{-1}$. We conclude this lemma. □

Rather than to study the series directly, Lemma [3.4](#page-17-5) allows us to study the asymptotics of each cumulant as $n \to \infty$ and to apply the dominated convergent theorem to the series.

Let us define $F := n^{\alpha} f_{\alpha, x_0}^{(n)}(\mathcal{J})$. Then, by definition [\(51\)](#page-14-0), $F = \sum_{r=1}^{2M} c_r G_{x_0 + \frac{\eta_r}{n^{\alpha}}}$, which is a bounded linear operator due to Corollary [3.2.](#page-14-3) Then, for $m \geq 2$,

$$
C_m(X_{f,\alpha,x_0}^{(n)}) = C_m^{(n)}(n^{-\alpha}F) = n^{-m\alpha}C_m^{(n)}(F),
$$
\n(81)

where $C_m^{(n)}$ on the right-hand side is given by [\(67\)](#page-16-2). The study of the cumulants of the mesoscopic linear statistics for large *n*, then, boils down to the asymptotic behaviour of $G_{x_0 + \frac{\eta r}{n^{\alpha}}}$, defined in [\(35\)](#page-11-0). In the case *n*α that *J* is self-adjoint, $G_{x_0 + \frac{\eta_r}{n^{\alpha}}}$ is the equivalent to the resolvent of the Jacobi operators, i.e., $(\mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}})^{-1}$.

3.4 Combes–Thomas Estimate and its Limitation

In our analysis, we will need to estimate entries of the resolvent that are far away from the main diagonal. One way of doing that (which is successful in [\[14\]](#page-81-0)) is by the Combes-Thomas estimate [\[34\]](#page-82-3). However, for our purpose, this Combes-Thomas estimate is not sufficient, and we will use conditions in Theorem [2.3](#page-8-0) to improve the estimate. Before doing so, we will recall the Combes-Thomas estimate and show its limitations for our purposes.

Combes–Thomas Estimate Given $\eta \in \mathbb{C}$ with Im $\eta \neq 0$. Consider

$$
J = \begin{pmatrix} b_0 & a_1 \\ a_1 & b_1 & a_2 \\ & a_2 & b_2 & a_3 \\ & & \ddots & \ddots & \ddots \end{pmatrix} - \left(x_0 + \frac{\eta}{n^{\alpha}}\right) Id,
$$
 (82)

where $b_j, x_0 \in \mathbb{R}, 0 < a_j \le c$ for some constant $c > 0$ and *Id* is the identity operator. Then for any $j, k \in \mathbb{N}$ we have

$$
\left| \left(J^{-1} \right)_{j,k} \right| \le \frac{2n^{\alpha}}{|\operatorname{Im} \eta|} e^{-\min\{1, \frac{|\operatorname{Im} \eta|}{4\epsilon cn^{\alpha}}\}|j-k|}.
$$
\n(83)

For a proof of the Combes–Thomas estimate, see [\[34\]](#page-82-3) or Proposition 2.3. in [\[14\]](#page-81-0).

Note that the Combes-Thomas estimate works for both varying $a_{j,n}$ and $b_{j,n}$ and non-varying a_j and b_j . Here we omit the second subscript for the convenience of following discussions.

The Combes-Thomas estimate is a useful result and is successful in studying the bulk universality for OPEs, see [\[14\]](#page-81-0). However, in our set up, it has some major limitations. First of all, a_j or $a_{j,n}$ can be unbounded. In case of the scaled Hermite polynomials (corresponding to the Gaussian unitary ensemble of the random matrix theory) we have $a_{j,n} = \sqrt{\frac{j}{n}}$ $\frac{d}{n}$ and $b_{j,n} = 0$. However, in the cumulant expansion [\(62\)](#page-15-0), one only needs to estimate $J^{-1}P_n$. The following argument enables us to replace *J* by a truncated version. Take *l* = 1 for example. Take $N = n + \delta$ for some $\delta > 0$ and define

$$
J_N:=P_NJP_N+Q_N.
$$

By the resolvent identity and the trace norm inequality, we have

$$
||(J_N^{-1} - J^{-1})P_n||_1 \le ||J^{-1}||_{\infty}||(J - J_N)J_N^{-1}P_n||_1.
$$
\n(84)

Recall that J_N is a block matrix and $N > n$. Hence, $J_N^{-1}P_n = P_N J_N^{-1} P_n$. Also, recall that J and J_N are tri-diagonal matrices. Hence, the only pop-zero entry of $(I - J_N)P_N$ is the $N + 1$, N -th entry whose value is tri-diagonal matrices. Hence, the only non-zero entry of $(J - J_N)P_N$ is the $N + 1$, *N*-th entry whose value is *aN*. Now we estimate the trace norm with entries, i.e.,

$$
\| (J - J_N) J_N^{-1} P_n \|_1 = \| (J - J_N) P_N J_N^{-1} P_n \|_1 \le \sum_{k=1}^n \left| a_N \left(J_N^{-1} \right)_{N,k} \right|.
$$
 (85)

Then, we only need to estimate $(J_N^{-1})_{j,k}$ for $j = N$ and $k = 1, 2, ..., n$. We use the Combes-Thomas estimate *j*,*k* for *J_N* to conclude that [\(85\)](#page-19-0) is exponentially small if $\delta \gg n^{\alpha}$. Note that $||J^{-1}||_{\infty} \le \frac{n^{\alpha}}{|\text{Im }\eta|}$, and hence [\(84\)](#page-19-1) is exponentially small as well. Becall that the cumulant formula given by (67). This implie Exponentially small as well. Recall that the cumulant formula given by [\(67\)](#page-16-2). This implies that the difference exponentially small as well. Recall that the cumulant formula given by (67). This implies that the difference of the second cumulants $C_2^{(n)}$ $\binom{n}{2}(J^{-1})$ and $C_2^{(n)}$ $\binom{n}{2}(J_N^{-1})$ is exponentially small. The cumulants of higher order can also be compared in a similar way. The truncation J_N can be a good estimate of J in the cumulant.

However, this reveals a second limitation. This argument works well for $\delta = n^{\beta}$ and $0 < \alpha < \beta < 1$, i.e. the model of the want which is not good enough near the edge. In some examples, for instance, the modified Jacobi case, we want to zoom in further and need an estimate that holds for any $0 < \alpha < 2$.

Note that if *J* is a Toeplitz operator, the Combes-Thomas estimate can be improved near the edges. For example, take $b_{j,n} = 0$ and $a_{j,n} = 1$, which are the recurrence coefficients for Chebyshev polynomials. Take $x_0 = 2$. One can use Wiener-Hopf factorization to compute the resolvent explicitly, that is

$$
\left(J^{-1}\right)_{j,k} = \frac{\omega(\eta)}{1 - \omega(\eta)^2} \left(\frac{1}{\omega(\eta)^{|j-k|}} + \frac{1}{\omega(\eta)^{j+k}}\right),\tag{86}
$$

where

$$
\omega(\eta) := \frac{2 + \frac{\eta}{n^{\alpha}} + \left(\left(2 + \frac{\eta}{n^{\alpha}} \right)^2 - 4 \right)^{\frac{1}{2}}}{2}.
$$

We take the principle square root so that $|\omega(\eta)| > 1$. Hence, we estimate,

$$
\left| \left(J^{-1} \right)_{j,k} \right| \le d_1 n^{\frac{\alpha}{2}} e^{-d_2 n^{-\frac{\alpha}{2}} |j-k|},\tag{87}
$$

for some constant $d_1, d_2 > 0$. Hence, we tell that for $|j - k| > n^{\beta}$ with $0 < \frac{\alpha}{2} < \beta < 1$, $|(J^{-1})_{j,k}|$ is exponentially small. One should note that $\frac{\alpha}{2} < \beta$ is possible since the spectral density of *J* vanishes at the spectrum as a square root. If $x_0 \in (-2, 2)$ is chosen, we will still get the region of exponential edge of the spectrum as a square root. If $x_0 \in (-2, 2)$ is chosen, we will still get the region of exponential decay to be $0 < \alpha < \beta < 1$.

Therefore, these limitations suggest a finer estimation about J_N^{-1} , which should resemble [\(87\)](#page-20-1), for some special class of *J*. In a nutshell, the Combes-Thomas estimate works for general Jacobi matrices, but can be improved under further conditions.

3.5 Inversion of a Symmetric Tri-diagonal Matrix

Recall that J_N is a tri-diagonal symmetric matrix. To find the inverse of such a matrix is a classical algebraic question and can be reduced to compute the first and last columns of J_N^{-1} . For a review, see [\[35\]](#page-82-4). Here we only present the expressions needed.

Let us look at some general properties of an $N \times N$ non singular symmetric tri-diagonal matrix with $a_j \in \mathbb{R}, b_j \in \mathbb{R}, N \in \mathbb{N}$, and $z \in \mathbb{C}$,

$$
J_N = \begin{pmatrix} b_0 & a_1 & & & & \\ a_1 & b_1 & a_2 & & & \\ & a_2 & b_2 & a_3 & & \\ & & \ddots & \ddots & \ddots & \\ & & & a_{N-2} & b_{N-2} & a_{N-1} \\ & & & & a_{N-1} & b_{N-1} \end{pmatrix} - zId,
$$
 (88)

where *Id* is the identity matrix.

Then for $j < k$

$$
(J_N^{-1})_{j,k} = (-1)^{k-j} a_j a_{j+1} \cdots a_{k-1} \frac{d_{k+1} \cdots d_N}{\delta_j \cdots \delta_N},
$$
\n(89)

$$
(J_N^{-1})_{j,j} = \frac{d_{j+1} \cdots d_N}{\delta_j \cdots \delta_N},\tag{90}
$$

where $d_N = b_{N-1} - z$ and, for all $j = 1, \dots N - 1$, d_j are solutions to the following difference equation

$$
d_j = b_{j-1} - z - \frac{a_j^2}{d_{j+1}}.\tag{91}
$$

We also have $\delta_1 = b_0 - z$ and, for all $j = 2, \dots N$, δ_j are solutions to the following difference equation

$$
\delta_j = b_{j-1} - z - \frac{a_{j-1}^2}{\delta_{j-1}}.\tag{92}
$$

Note that J_N^{-1} is symmetric and we have $(J_N^{-1})_{k,j} = (J_N^{-1})_{j,k}$. Hence, we have an expression for each entry.

In the following we will linearize both difference equations above and rewrite the inverse formula in different ways that are suitable for our further analysis.

Linearization of the difference equation I

Recursively define ^β*^j* by

$$
\beta_N = a_{N-1}, \quad \beta_{N-1} = b_{N-1} - z, \quad \frac{\beta_j}{\beta_{j-1}} = \frac{a_{j-1}}{d_j}, \quad \text{for } j = 1, ..., N.
$$

Hence, β_j is the solution to the following linear difference equation, by [\(91\)](#page-20-2),

$$
\begin{pmatrix} \beta_j \\ \beta_{j-1} \end{pmatrix} = A_j \begin{pmatrix} \beta_{j+1} \\ \beta_j \end{pmatrix}, \quad \text{where } A_j := \begin{pmatrix} 0 & 1 \\ -\frac{a_j}{a_{j-1}} & \left(b_{j-1} - z \right) \frac{1}{a_{j-1}} \end{pmatrix}, \tag{93}
$$

for $j = 2, \ldots, N - 1$. Iteratively,

$$
\begin{pmatrix} \beta_j \\ \beta_{j-1} \end{pmatrix} = A_j A_{j+1} \cdots A_{N-1} \begin{pmatrix} \beta_N \\ \beta_{N-1} \end{pmatrix}.
$$
\n(94)

Let *N*⁰ \in N and *N*⁰ \leq *N*. Take the ratio of (J_N^{-1}) *j*,*k* and $\left(J_{N}^{-1}\right)$ j_{j,N_0} which are found by [\(89\)](#page-20-3) or [\(90\)](#page-20-4), and we rewrite the inversion formula for $j \leq N_0 < k$ to be

$$
\left(J_N^{-1}\right)_{j,k} = (-1)^{k-N_0} \frac{\beta_k}{\beta_{N_0}} \left(J_N^{-1}\right)_{j,N_0}.
$$
\n(95)

One purpose of the coming Section [5](#page-30-0) is to show that in the case $b_j = b_{j,n} - x_0$, $a_j = a_{j,n}$, $z = \eta/n^{\alpha}$, $N = n + 2mn^{\alpha/2 + \varepsilon/3}$ and $N_0 = n - 2mn^{\alpha/2 + \varepsilon/3}$ for $m \in \mathbb{N}$, given assumptions of Theorem [2.3,](#page-8-0) we have $|\beta_k/\beta_{N_0}| \le e^{-d_2(k-N_0)n^{-\alpha/2}}$, for some constant $d_2 > 0$. The same is also true for the case $b_j = b_{j+n-2mn^{\alpha/2+\epsilon/3},n}$ –
re $a_j = a_{j+n-2m}$ $a_{j+n-2m} = a_{j+n-2m}$ $a_{j+n-2m} = a_{j+n-2m}$ x_0 , $a_j = a_{j+n-2mn^{\alpha/2+\epsilon/3},n}$, $z = \eta/n^{\alpha}$, $N = 4mn^{\alpha/2+\epsilon/3}$ and $N_0 = 2mn^{\alpha/2+\epsilon/3}$ for $m \in \mathbb{N}$. Hence, we have an estimate as in Proposition 4.1, which is an analogue of the exponential part of (87) estimate as in Proposition [4.1,](#page-24-0) which is an analogue of the exponential part of [\(87\)](#page-20-1).

To this end, the following quantities are essential in this estimate. The eigenvalues of *A^j* are

$$
\omega_j^+ := \frac{b_{j-1} - z + \left((b_{j-1} - z)^2 - 4a_j a_{j-1} \right)^{\frac{1}{2}}}{2a_{j-1}}, \quad \omega_j^- := \frac{b_{j-1} - z - \left((b_{j-1} - z)^2 - 4a_j a_{j-1} \right)^{\frac{1}{2}}}{2a_{j-1}}.
$$
(96)

The square root is taken with respect to the principle branch, so that $|\omega_j^+| > |\omega_j^-|$
Note that A, can be diagonalized as $\frac{1}{j}$, whenever Re(*b*_{*j*} − *z*) > 0. Note that A_j can be diagonalized as

$$
A_j = V_j \Omega_j V_j^{-1}, \quad \text{where } V_j := \begin{pmatrix} 1 & 1 \\ \omega_j^+ & \omega_j^- \end{pmatrix}, \quad \Omega_j := \begin{pmatrix} \omega_j^+ & 0 \\ 0 & \omega_j^- \end{pmatrix}.
$$

Also of importance is

$$
M_j := V_j^{-1} V_{j+1} - Id = \frac{1}{\omega_j^- - \omega_j^+} \begin{pmatrix} \omega_j^+ - \omega_{j+1}^+ & \omega_j^- - \omega_{j+1}^- \\ \omega_{j+1}^+ - \omega_j^+ & \omega_{j+1}^- - \omega_j^- \end{pmatrix},
$$
(97)

where *Id* is the identity matrix.

Note that Proposition [4.1](#page-24-0) only matches the exponential part of (87) . To match the $n^{\alpha/2}$ factor of (87) , we need the following ingredients as well.

Linearization of the difference equation II

Similarly, we recursively define ^γ*^j* by

$$
\gamma_1 = a_1, \quad \gamma_2 = b_0 - z, \quad \frac{\gamma_j}{\gamma_{j+1}} = \frac{a_j}{\delta_j}, \quad \text{for } j = 1, \dots, N.
$$

Then γ_j also admits a recursive formula by equation [\(92\)](#page-21-0)

$$
\begin{pmatrix} \gamma_j \\ \gamma_{j+1} \end{pmatrix} = B_j \begin{pmatrix} \gamma_{j-1} \\ \gamma_j \end{pmatrix}, \quad \text{where } B_j := \begin{pmatrix} 0 & 1 \\ -\frac{a_{j-1}}{a_j} & \left(b_{j-1} - z \right) \frac{1}{a_j} \end{pmatrix}, \tag{98}
$$

for all $j = 2, ..., N - 1$. The matrix B_j is called the transfer matrix for γ_j . Iteratively

$$
\begin{pmatrix} \gamma_j \\ \gamma_{j+1} \end{pmatrix} = B_j B_{j-1} \dots B_2 \begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix}.
$$

Choose any $a_N \neq 0$ (e.g. One can pick $a_N = a_{N-1}$), rewrite [\(89\)](#page-20-3) and [\(90\)](#page-20-4) by β_j and γ_j , and we have other inversion formula for any $i \leq k$ another inversion formula for any $j \leq k$

$$
(J_N^{-1})_{j,k} = \frac{(-1)^{k-j} \gamma_j \beta_k}{\beta_N a_N \gamma_{N+1}}.
$$
\n(99)

Note that the inverse formula is independent on the choice of a_N since $a_N\gamma_{N+1} = \gamma_N\delta_N$ and γ_N and δ_N are independent of a_N . Also note that I_N is symmetric, and hence we have an expression for all entries independent of a_N . Also note that J_N is symmetric, and hence we have an expression for all entries.

One purpose of the coming Section [6](#page-44-0) is to show that in the case $b_j = b_{j+n-2mn\alpha/2+\epsilon/3,n} - x_0$, $a_j =$
 $a_j = \pi - n/n^{\alpha} N - 4mn^{\alpha/2+\epsilon/3}$ for $m \in \mathbb{N}$, we have a procise asymptotics of β_1/β_2 , and α_2/α_3 . $a_{j+n-2mn}a_{j+2}a_{j+n}$, $z = \eta/n^{\alpha}$, $N = 4mn^{\alpha/2+\epsilon/3}$ for $m \in \mathbb{N}$, we have a precise asymptotics of β_k/β_{N_0} and γ_j/γ_{N+1} as $n \to \infty$ whenever the assumptions of Theorem [2.3](#page-8-0) are satisfied. Hence, we will have a complete analogue to [\(86\)](#page-20-5), cf. Proposition [4.3](#page-26-0)

To this end, the following quantities are essential in this estimate. The eigenvalues of B_j are

$$
\lambda_j^+ := \frac{b_{j-1} - z + \left((b_{j-1} - z)^2 - 4a_j a_{j-1} \right)^{\frac{1}{2}}}{2a_j}, \quad \lambda_j^- := \frac{b_{j-1} - z - \left((b_{j-1} - z)^2 - 4a_j a_{j-1} \right)^{\frac{1}{2}}}{2a_j}.
$$
(100)

The square root is taken with respect to the principle branch, such that $|\lambda_j^+| > |\lambda_j^-|$
Note that R_j can be diagonalized as $\frac{1}{j}$, whenever Re(*b*_{*j*} − *z*) > 0. Note that B_j can be diagonalized as

$$
B_j = W_j \Lambda_j W_j^{-1}
$$
, where $W_j := \begin{pmatrix} 1 & 1 \\ \lambda_j^+ & \lambda_j^- \end{pmatrix}$, $\Lambda_j := \begin{pmatrix} \lambda_j^+ & 0 \\ 0 & \lambda_j^- \end{pmatrix}$.

Similarly we define E_j , which is an essential element for the further analysis, to be

$$
E_j := W_j^{-1} W_{j-1} - Id = \frac{1}{\lambda_j^- - \lambda_j^+} \begin{pmatrix} \lambda_j^+ - \lambda_{j+1}^+ & \lambda_j^- - \lambda_{j+1}^- \\ \lambda_{j+1}^+ - \lambda_j^+ & \lambda_{j+1}^- - \lambda_j^- \end{pmatrix} . \tag{101}
$$

The formulae [\(95\)](#page-21-1) and [\(99\)](#page-22-0) are the corner stones of key Propositions [4.1](#page-24-0) and [4.3](#page-26-0) respectively.

4 Proof of Theorem [2.3](#page-8-0)

In this section, we decompose the proof of Theorem [2.3](#page-8-0) into a number of propositions, which we will prove in the upcoming sections, cf. Sections [5,](#page-30-0) [6,](#page-44-0) and [7.](#page-54-0)

Our strategy is to first prove Theorem [2.3](#page-8-0) for some special test functions as described in [\(49\)](#page-14-4) , i.e.,

$$
f(x) = \sum_{r=1}^{2M} c_r \frac{1}{x - \eta_r}
$$

for some $\eta_r \in \{x + iy | x \in \mathbb{R}, y \neq 0\}$, in Subsection [4.1.](#page-23-1) Then, extend this result to compactly supported and continuously differentiable test functions in Subsection [4.2.](#page-27-0)

4.1 Proof of Theorem [2.3](#page-8-0) for Special Class of Test Functions

Recall that $\mathcal J$ defined in [\(32\)](#page-10-1) is the Jacobi matrix associated with an OPE and $f_{\alpha,x_0}^{(n)}(\mathcal J)$ is defined as [\(51\)](#page-14-0). Let us further define

$$
J^{(r)} := \mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}}, \quad F := n^{\alpha} f_{\alpha, x_0}^{(n)}(\mathcal{J}). \tag{102}
$$

Recall that we say $x_0 \in \mathbb{R}$ is near the edges if one of the conditions in [\(24\)](#page-7-4) is satisfied. Recall, from Section [3.3,](#page-13-0) that the *m*-th cumulant with $m \ge 2$ for the mesoscopic linear statistics is given, via [\(62\)](#page-15-0), by

$$
C_m(X_{f,\alpha,x_0}^{(n)}) = \frac{m!}{n^{\alpha m}} \sum_{j=2}^m \frac{(-1)^{j+1}}{j} \sum_{l_1 + l_1 = m, l_i \ge 1} \frac{\operatorname{Tr}\left(F^{l_1} P_n \cdots F^{l_j} P_n\right) - \operatorname{Tr}\left(F^m P_n\right)}{l_1! \cdots l_j!}.
$$
 (103)

That is $C_m(X_{f, o}^{(n)})$ $f_{f,\alpha,x_0}^{(n)}$ = $C_m^{(n)}(n^{-\alpha}F)$ or equivalently $n^{\alpha m}C_m(X_{f,\alpha}^{(n)})$ $f_{f,\alpha,x_0}^{(n)}$ = $C_m^{(n)}(F)$, as in the relation [\(81\)](#page-18-3). To show that the limiting fluctuations of $X_{f, \alpha, n}^{(n)}$ are Gaussian, it is sufficient to show that the second cumulant converges to a positive number and all cumulants of order $m \geq 3$ converge to zero, as indicated by Lemma [3.4.](#page-17-5)

It turns out that the asymptotics of the *m*-th cumulant ($m \in \mathbb{N}$) only depends on the recurrence coefficients of order around *ⁿ*, i.e., *^aj*,*n*, *^bj*,*ⁿ* for all *^j* [∼] *ⁿ* as *ⁿ* → ∞. The precise window depends on the scale considered. This has been observed in various setups as in [\[14,](#page-81-0) [23,](#page-81-9) [27\]](#page-81-13). We will show that this is the case in our setup as well. To this end, we will truncate the (semi-finite) Jacobi matrix $\mathcal T$ into the block around the n, n -th entry. For technical reasons, we will conduct a two-step truncation. To be precise, let $\beta = \frac{\alpha}{2} + \frac{\epsilon}{3}$ such that $0 \le \frac{\alpha}{2} \le \beta \le \frac{\alpha+1}{2} \le 1$. Define $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$. Define

$$
J_{n+2mn\beta}^{(r)} := P_{n+2mn\beta} J^{(r)} P_{n+2mn\beta} + Q_{n+2mn\beta}, \qquad F_{n+2mn\beta} := \sum_{r=1}^{2M} c_r \left(J_{n+2mn\beta}^{(r)} \right)^{-1}, \tag{104}
$$

$$
J_{n\pm 2mn\beta}^{(r)} := P_{n+2mn\beta} Q_{n-2mn\beta} J^{(r)} Q_{n-2mn\beta} P_{n+2mn\beta} + P_{n-2mn\beta} + Q_{n+2mn\beta},
$$
(105)

$$
F_{n\pm 2mn\beta} := \sum_{r=1}^{2M} c_r \left(J_{n\pm 2mn\beta}^{(r)} \right)^{-1} . \tag{106}
$$

The subscript *n*±2*mn*^β emphasises that the relevant entries are those with indices ranging between *n*−2*mn*^β and $n + 2mn^{\beta}$.

The following is our first finding about these truncated matrices, which is an alternative to the Combes-Thomas estimates for $J_{n+1}^{(r)}$ $\int_{n+2mn\beta}^{(r)}$ and $J_{n\pm}^{(r)}$ *n*±2*mn*β .

Figure 2: Illustration of Proposition [4.1.](#page-24-0) Given the entries of $J_{n}^{(r)}$ $J_{n+2mn\beta}^{(r)}$ and $J_{n\pm 2mn\beta}^{(r)}$ with both indices ranging between $n - 2mn^{\beta}$ and $n + 2mn^{\beta}$ are well behaved, we have that the entries of $\left(J_{n+1}^{(r)}\right)$ *n*+2*mn*β \int_0^{-1} and $\left(J_{n+1}^{(r)}\right)$ *n*±2*mn*β 1^{-1} are of exponentially small in the shaded areas respectively. Note that the matrices are block matrices and we ignore the trivial blocks in the illustration.

Proposition 4.1. *Let* $\beta = \frac{\alpha}{2} + \frac{\varepsilon}{3}$ *be such that* $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$ *. Consider J described above whose* entries are (a_1, b_1) , *Let* $m \in \mathbb{N}$ *Assume conditions in Theorem* 2.3 are satisfied *entries are* $(a_{l,n}, b_{l,n})_{l \geq 1}$ *. Let m* $\in \mathbb{N}$ *. Assume conditions in Theorem [2.3](#page-8-0) are satisfied. Consider* x_0 *to be near the edges, i.e.,* [\(24\)](#page-7-4)*. Then there exist constants* $C_0 > 0$, $d_0 > 0$ $n_0 \in \mathbb{N}$ *such that, for any n* > n_0 , *j, k with* $|j - k| \ge n^{\beta}$ *and* $\max\{j, k\} \ge n - (2m - 1)n^{\beta}$,

$$
\left| \left(\left(J_{n+2mn\beta}^{(r)} \right)^{-1} \right)_{j,k} \right| \leq C_0 n^{\alpha} e^{-d_0 n^{\beta - \frac{\alpha}{2}}},\tag{107}
$$

$$
\left| \left(\left(J_{n \pm 2m n^{\beta}}^{(r)} \right)^{-1} \right)_{j,k} \right| \leq C_0 n^{\alpha} e^{-d_0 n^{\beta - \frac{\alpha}{2}}}.
$$
\n(108)

Consequently,

$$
\left| \left(F_{n+2mn\beta} \right)_{j,k} \right| \leq C_0 n^{\alpha} e^{-d_0 n^{\beta - \frac{\alpha}{2}}},\tag{109}
$$

$$
\left| \left(F_{n \pm 2mn^{\beta}} \right)_{j,k} \right| \leq C_0 n^{\alpha} e^{-d_0 n^{\beta - \frac{\alpha}{2}}}.
$$
\n(110)

For an illustration see Figure [2.](#page-24-1) For the proof of Proposition [4.1,](#page-24-0) see Section [5.2.](#page-32-0)

Remark that the factor n^{α} is not the optimal in the estimates $\left(J_{n+1}^{(r)}\right)$ *n*±2*mn*β \int^{-1} or $F_{n+1}^{(r)}$ $\lim_{n \pm 2mn\beta}$. These are improved in Proposition [4.3.](#page-26-0) However, this is enough for the next step, which is the following proposition lying in the core of the proof of Theorem [2.3.](#page-8-0)

Proposition 4.2. *Let* $\beta = \frac{\alpha}{2} + \frac{\varepsilon}{3}$ *such that* $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$ *. Let* $m \in \mathbb{N}$ *and* $m \ge 2$ *. Assume the*
conditions in Theorem 2.3 are satisfied and x is near the edges i.e. (24). Then there *conditions in Theorem* [2.3](#page-8-0) *are satisfied and* x_0 *is near the edges, i.e.,* [\(24\)](#page-7-4)*. Then there exist constants* $d' > 0$ *,* $n_0 \in \mathbb{N}$, and $C'_m > 0$ that only depend on m and M such that for all $n > n_0$

$$
\left|n^{\alpha m} C_m(X_{f,\alpha,x_0}^{(n)}) - C_m^{(n)}(F_{n \pm 2mn\beta})\right| \le C_m' e^{-d'n^{\beta - \frac{\alpha}{2}}}.
$$
\n(111)

We give the proof for $m = 2$ here. The proof for general m follows the same principles, but is rather technical and we postpone the details to Section [5.4.](#page-41-0)

Proof of Propsition [4.2](#page-25-0) . Consider the case *m* = 2.

For any linear operator A , the second cumulant is

$$
C_2^{(n)}(\mathcal{A}) = \text{Tr}(\mathcal{A}P_n\mathcal{A}P_n) - \text{Tr}(\mathcal{A}^2P_n).
$$

Write $Q_n \coloneqq Id - P_n$ and

$$
C_2^{(n)}(\mathcal{A})=-\mathrm{Tr}\left(\mathcal{A}Q_n\mathcal{A}P_n\right).
$$

For any linear operators $\mathcal A$ and $\mathcal B$, we can write the difference of the cumulants by adding and subtracting extra terms to be

$$
C_2^{(n)}(\mathcal{A}) - C_2^{(n)}(\mathcal{B}) = -\operatorname{Tr}\left((\mathcal{A} - \mathcal{B})\mathcal{Q}_n\mathcal{A}P_n\right) - \operatorname{Tr}\left(\mathcal{B}\mathcal{Q}_n\left(\mathcal{A} - \mathcal{B}\right)P_n\right). \tag{112}
$$

Assume A and B are symmetric (not necessarily Hermitian). Use the trace norm inequality $||AB||_1 \le$ ∥*A*∥1∥*B*∥[∞] to obtain

$$
\left| C_2^{(n)}(\mathcal{A}) - C_2^{(n)}(\mathcal{B}) \right| \leq ||P_n(\mathcal{A} - \mathcal{B})Q_n||_1 (||\mathcal{A}||_{\infty} + ||\mathcal{B}||_{\infty}). \tag{113}
$$

Then take $\mathcal{A} = F$ as [\(102\)](#page-23-2) and $\mathcal{B} = F_{n+2mn\beta}$ as [\(104\)](#page-23-3). We have

$$
P_n(\mathcal{A} - \mathcal{B}) = \sum_r c_r P_n \left(G_{x_0 + \frac{\eta_r}{n^{\alpha}}} - \left(J_{n+2mn\beta}^{(r)} \right)^{-1} \right). \tag{114}
$$

Note that $n + 2mn^{\beta} > n$, and we have $P_n\left(J_{n+1}^{(r)}\right)$ *n*+2*mn*β $\int_{0}^{-1} = P_n \left(J_{n+1}^{(r)} \right)$ *n*+2*mn*β $\int_{0}^{-1} P_{n+2mn\beta}$. Define the linear operator *E* with all but two entries being zeros, i.e.,

$$
\begin{cases}\n(E)_{i,j} = (E)_{j,i} = a_{n+2mn}\beta, & \text{for } i = n+2mn\beta \text{ and } j = n+2mn\beta+1, \\
(E)_{i,j} = 0, & \text{otherwise.} \n\end{cases}
$$
\n(115)

Hence,

$$
P_n\left(J_{n+2mn\beta}^{(r)}\right)^{-1}\left(J^{(r)}-J_{n+2mn\beta}^{(r)}\right)=P_n\left(J_{n+2mn\beta}^{(r)}\right)^{-1}E.
$$
 (116)

Note that the entries of matrix [\(116\)](#page-25-1) are zeros except for those at the first *n* rows of the last column. Then apply the resolvent identity [\(43\)](#page-12-2) (also see the comment below [\(43\)](#page-12-2)) to [\(114\)](#page-25-2) to obtain

$$
P_n(\mathcal{A} - \mathcal{B}) = -\sum_{r} c_r P_n \left(J_{n+2mn\beta}^{(r)} \right)^{-1} \left(J^{(r)} - J_{n+2mn\beta}^{(r)} \right) G_{x_0 + \frac{\eta_r}{n^{\alpha}}},\tag{117}
$$

where $G_{x_0 + \frac{\eta r}{n^{\alpha}}}$ is given by [\(35\)](#page-11-0). Hence, plug [\(117\)](#page-25-3) and [\(116\)](#page-25-1) into [\(113\)](#page-25-4), use the triangle inequality to obtain

$$
\left| C_2^{(n)}(F) - C_2^{(n)}(F_{n+2mn\beta}) \right| \leq \sum_r |c_r| \left\| P_n \left(J_{n+2mn\beta}^{(r)} \right)^{-1} E \right\|_1 \left\| G_{x_0 + \frac{\eta_r}{n^{\alpha}}} \right\|_{\infty} \left(\|F\|_{\infty} + \|F_{n+2mn\beta}\|_{\infty} \right). \tag{118}
$$

The trace norm above is experientially small by [\(107\)](#page-24-2) of Proposition [4.1.](#page-24-0) All the operator normals are of order $O(n^{\alpha})$, since $\mathcal J$ is real symmetric and Im(η_r) $\neq 0$. Hence, [\(120\)](#page-26-1) is exponentially small.
Next, we take $\mathcal A = F$, as (100) and $\mathcal B = F$, as (106). Similarly we use the re-

Next, we take $\mathcal{A} = F_{n+2mn\beta}$ as [\(104\)](#page-23-3) and $\mathcal{B} = F_{n+2mn\beta}$ as [\(106\)](#page-23-4). Similarly, we use the resolvent identity and the fact that $(J_{n+1}^{(r)})$ *n*±2*mn*β $\int_{0}^{-1} Q_n = Q_{n-2mn} \int_{0}^{r} (f_{n\pm}^{(r)})^2$ *n*±2*mn*β $\int_0^{-1} Q_n$ to obtain

$$
\left(\left(J_{n+2mn\beta}^{(r)}\right)^{-1} - \left(J_{n\pm 2mn\beta}^{(r)}\right)^{-1}\right)Q_n = -\left(J_{n+2mn\beta}^{(r)}\right)^{-1}\tilde{E}\left(J_{n\pm 2mn\beta}^{(r)}\right)^{-1}Q_n,
$$

where \tilde{E} only has two entries that are not zeros, i.e.,

$$
\begin{cases}\n(\tilde{E})_{i,j} = (\tilde{E})_{j,i} = a_{n-2mn}\beta, & \text{for } i = n - 2mn\beta \text{ and } j = n - 2mn\beta - 1, \\
(\tilde{E})_{i,j} = 0, & \text{otherwise.}\n\end{cases}
$$
\n(119)

Note that the entries of matrix $\tilde{E} (J_{n+1}^{(r)})$ *n*±2*mn*β $\int_0^{-1} Q_n$ are zeros except for those at the last $2mn^{\beta}$ columns of the first row. Hence,

$$
\left| C_2^{(n)}(F_{n+2m n^{\beta}}) - C_2^{(n)}(F_{n+2m n^{\beta}}) \right| \leq \sum_r |c_r| \left\| \left(J_{n+2m n^{\beta}}^{(r)} \right)^{-1} \right\|_{\infty} \left\| \tilde{E} \left(J_{n+2m n^{\beta}}^{(r)} \right)^{-1} Q_n \right\|_1 (\|F_{n+2m n^{\beta}}\|_{\infty} + \|F_{n+2m n^{\beta}}\|_{\infty}).
$$
\n(120)

Similarly, by [\(108\)](#page-24-3) of Proposition [4.1,](#page-24-0) we have that [\(120\)](#page-26-1) is of exponentially small.

The estimates [\(118\)](#page-25-5) and [\(120\)](#page-26-1) show that $\left|C_m(F) - C_m(F_{n \pm 2mn\beta})\right|$ is of exponentially small for $m = 2$.

For the general case $m > 2$, the algebra of the difference of the cumulants is more complicated than [\(112\)](#page-25-6). However, the strategy is similar. For details, see the complete proof of Proposition [4.2](#page-25-0) in Section [5.4.](#page-41-0)

□

Proposition [4.2](#page-25-0) implies that it is sufficient to study the truncated operator $F_{n\pm 2mn\beta}$ as $n \to \infty$. By the definition of $F_{n\pm 2mn\beta}$ in [\(106\)](#page-23-4), essentially, we are left to study the resolvent $\left(J_{n\pm}^{(r)}\right)$ *n*±2*mn*β \int_{0}^{π} defined in [\(105\)](#page-23-5).

Proposition 4.3. Let $\beta = \frac{\alpha}{2} + \frac{\varepsilon}{3}$ such that $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$. Assume conditions in Theorem [2.3](#page-8-0) are satisfied. Consider x_0 to be near the edges, i.e., [\(24\)](#page-7-4). Then there exists a decomposition of $\left(J_{n+1}^{(r)}\right)$ *n*±2*mn*β \int^{-1} *such that*

$$
\left(J_{n\pm 2mn\beta}^{(r)}\right)^{-1} = T_{n\pm 2mn\beta}(\eta_r) + H_{n\pm 2mn\beta}(\eta_r)
$$
 (121)

where $T_{n\pm 2mn\beta}(\eta_r)$ *and* $H_{n\pm 2mn\beta}(\eta_r)$ *are such that as* $n \to \infty$ *, for all j, k* = $n - 2mn^{\beta}, \ldots, n + 2mn^{\beta}$ *,*

$$
(T_{n\pm 2mn^{\beta}}(\eta_r))_{j,k} = \frac{(-1)^{|k-j|}}{2a_{n,n}\left(\frac{-sgn(b_{n-1,n}-x_0)\eta_r}{a_{n,n}n^{\alpha}}\right)^{\frac{1}{2}}}\prod_{l=\min\{j,k\}}^{\max\{j,k\}-1} \left(1-\left(\frac{-sgn(b_{n-1,n}-x_0)\eta_r}{a_{n,n}n^{\alpha}}+\xi_{l}^{(r)}\right)^{\frac{1}{2}}\right)(1+o(1)), \quad (122)
$$

where $\max_{l=n-2mn\beta,...,n+2mn\beta} |\xi_l^{(r)}|$ $\binom{f}{l}$ = $o(n^{-\alpha/2})$ *and by convention* $\prod_{l=j}^{j-1}$ ≡ 1*. We also have as n* → ∞*, for all* $j, k, j + l, k + l = n - 2mn^{\beta}, \ldots, n + 2mn^{\beta},$

$$
\frac{(T_{n\pm 2mn^{\beta}}(\eta))_{j,k}}{(T_{n\pm 2mn^{\beta}}(\eta))_{j+l,k+l}} = 1 + o\left(n^{-\beta + \frac{\alpha}{2}}\right),\tag{123}
$$

$$
\left| (H_{n\pm 2mn\beta}(\eta_r))_{j,k} \right| \le C n^{\frac{\alpha}{2}} \left(e^{-dn^{-\frac{\alpha}{2}} (\max\{j,k\} - n + 2mn^{\beta})} + e^{-dn^{-\frac{\alpha}{2}} (n + 2mn^{\beta} - \min\{j,k\})} \right),\tag{124}
$$

for some constant $C, d > 0$ *.*

 $\overline{}$

For the proof of Proposition [4.3,](#page-26-0) see Section [6.2.](#page-49-0)

Compared with [\(86\)](#page-20-5), [\(121\)](#page-26-2) can be regarded as an analogue of the inverse formula of a Toeplitz operator. Formula [\(122\)](#page-26-3) implies that the entries along the same diagonals of $T_{n\pm 2mn\beta}$ are almost the same. The error is controlled by [\(123\)](#page-26-4).

Note that the trace norm of $H_{n\pm 2mn\beta}(\eta_r)$ is much smaller than that of the $T_{n\pm 2mn\beta}(\eta_r)$ by [\(124\)](#page-26-5). We will also show that its contribution to the cumulant is exponentially small in the following proposition.

Proposition 4.4. Let $\beta = \frac{\alpha}{2} + \frac{\varepsilon}{3}$ such that $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$. Let $m \in \mathbb{N}$ and $m \ge 2$. Assume conditions in Theorem 2.3 are satisfied. Consider x_0 to be at the edges. Then there exists a cons *in Theorem [2.3](#page-8-0) are satisfied. Consider x*⁰ *to be at the edges. Then there exists a constant C*′ *^m* > ⁰ *that only depends on m and M such that*

$$
\left| n^{\alpha m} C_m(X_{f,\alpha,x_0}^{(n)}) - C_m^{(n)} \left(\sum_r c_r T_{n \pm 2m n^{\beta}} (\eta_r) \right) \right| \le C_m' e^{-d' n^{\beta - \frac{\alpha}{2}}},\tag{125}
$$

for all n > n_0 *for some constant* $n_0 \in \mathbb{N}$ *.*

For the proof of Proposition [4.4,](#page-27-1) see Section [6.3.](#page-50-0)

Proposition [4.2](#page-25-0) implies that the asymptotics of the cumulants of the mesoscopic linear statistics is reduced to studying the operator $T_{n\pm 2mn\beta}$ (η_r). To this end, we will show the cumulant generating function of $\sum_r c_r T_{n\pm 2m\theta}(\eta_r)$ converges to that of a Gaussian. That is an analogue of the strong Szego's limit theorem to $\sum_r c_r T_{n\pm 2m\theta}(\eta_r)$ converges to that of a Gaussian. That is an analogue of the strong Szego's limit th this operator whose non-trivial block is close to a Toeplitz matrix but entries vary slowly along the diagonals.

Proposition 4.5. *Let* $\beta = \frac{\alpha}{2} + \frac{\epsilon}{3}$ *such that* $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$ *. Assume conditions in Theorem [2.3](#page-8-0) are*
satisfied. Consider x to be at the adoes. The test function f is defined as (40). Then for any *satisfied. Consider* x_0 *to be at the edges. The test function f is defined as* [\(49\)](#page-14-4). Then for any $t > 0$ *small,*

$$
\lim_{n \to \infty} \log \det \left(1 + P_n \left(e^{tn^{-\alpha} \sum_r c_r T_{n \pm 2mn} \beta(\eta_r)} - 1 \right) P_n \right) e^{-t \operatorname{Tr} P_n n^{-\alpha} \sum_r c_r T_{n \pm 2mn} \beta(\eta_r)} = t^2 \sigma_f^2,
$$
\nwhere, $\sigma_f^2 := \begin{cases} \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy, & \text{for } x_0 = b_{n-1,n} - 2 \sqrt{a_{n,n} a_{n-1,n}} + o(n^{-\alpha}), \\ \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy, & \text{for } x_0 = b_{n-1,n} + 2 \sqrt{a_{n,n} a_{n-1,n}} + o(n^{-\alpha}). \end{cases}$ \n(126)

Consequently, Theorem [2.3](#page-8-0) holds for such test functions f as defined in [\(49\)](#page-14-4)*.*

For the proof of Proposition [4.5,](#page-27-2) see Section [7.2.](#page-61-0)

The last step of the proof of Theorem [2.3](#page-8-0) is to extend such *f* to compactly supported functions following a standard procedure, for example, [\[14,](#page-81-0) [36\]](#page-82-5). This is elaborated in Section [4.2.](#page-27-0)

4.2 Proof of Theorem [2.3](#page-8-0)

To finish the proof of Theorem [2.3,](#page-8-0) we need to extend the result of Proposition [4.5](#page-27-2) to the test functions $f \in C_c^1$. An important role of extension is played by the following space of functions.

Definition 4.1. *Let* \mathcal{L}_w *be a space of functions of* $f : \mathbb{R} \to \mathbb{R}$ *such that* $\lim_{x \to \pm \infty} f(x) = 0$ *and*

$$
||f||_{\mathcal{L}_w} := \sup_{x,y \in \mathbb{R}} \sqrt{1 + x^2} \sqrt{1 + y^2} \left| \frac{f(x) - f(y)}{x - y} \right| < \infty.
$$
 (127)

 \mathcal{L}_w is a normed space with the weighted Lipschitz norm $||f||_{\mathcal{L}_w}$. Note that $C_c^1 \subset \mathcal{L}_w$.

Proposition 4.6 (Proposition 5.1 in[\[14\]](#page-81-0)). *Let* $g(x) = \frac{1}{x-i}$. *Then for any* $f \in \mathcal{L}_w$ *and* $n \in \mathbb{N}$ *we have*

$$
\text{Var}\left(X_{f,\alpha,x_0}^{(n)}\right) \le ||f||_{\mathcal{L}_w}^2 \left(\text{Var}\left(X_{\text{Im }g,\alpha,x_0}^{(n)}\right) + \text{Var}\left(X_{\text{Re }g,\alpha,x_0}^{(n)}\right)\right). \tag{128}
$$

A consequence of Proposition [4.6](#page-28-1) is the following corollary.

Corollary 4.1. Assume all conditions of Theorem [2.3](#page-8-0) are satisfied. Then for any $f \in \mathcal{L}_w$ we have

$$
\limsup_{n \to \infty} \text{Var}\left(X_{f,\alpha,x_0}^{(n)}\right) \le \frac{1}{8} ||f||_{\mathcal{L}_w}^2. \tag{129}
$$

Proof. We are to estimate the right-hand side of [\(128\)](#page-28-2). Note that for $g(x) = \frac{1}{x-i}$ we have

Im
$$
g(x) = \frac{1}{2i} \left(\frac{1}{x - i} - \frac{1}{x + i} \right)
$$
, Re $g(x) = \frac{1}{2} \left(\frac{1}{x - i} + \frac{1}{x + i} \right)$.

Using Proposition [4.5](#page-27-2) for these two functions, we find that at the right edge

$$
\lim_{n \to \infty} \text{Var}\left(X_{\text{Im }g,\alpha,x_0}^{(n)}\right) = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{\text{Im }g(-x^2) - \text{Im }g(-y^2)}{x - y}\right)^2 dxdy = \frac{3}{32},
$$

$$
\lim_{n \to \infty} \text{Var}\left(X_{\text{Re }g,\alpha,x_0}^{(n)}\right) = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{\text{Re }g(-x^2) - \text{Re }g(-y^2)}{x - y}\right)^2 dxdy = \frac{1}{32}.
$$

Similarly, we also obtain that at the left edge

$$
\lim_{n \to \infty} \text{Var}\left(X_{\text{Im }g,\alpha,x_0}^{(n)}\right) = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{\text{Im } g(x^2) - \text{Im } g(y^2)}{x - y}\right)^2 dxdy = \frac{3}{32},
$$

$$
\lim_{n \to \infty} \text{Var}\left(X_{\text{Re } g,\alpha,x_0}^{(n)}\right) = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{\text{Re } g(x^2) - \text{Re } g(y^2)}{x - y}\right)^2 dxdy = \frac{1}{32}.
$$

These imply that at both edges

$$
\limsup_{n\to\infty}\left(\text{Var}\left(X_{\text{Im }g,\alpha,x_0}^{(n)}\right)+\text{Var}\left(X_{\text{Re }g,\alpha,x_0}^{(n)}\right)\right)\leq\frac{1}{8}.
$$

Then combine with Proposition [4.6](#page-28-1) and we conclude this corollary. □

Lemma 4.1 (Lemma 5.3 in [\[14\]](#page-81-0)). *Let* $f \in C_c^1(\mathbb{R})$ *. For any* $\varepsilon > 0$ *, there exists a* $M \in \mathbb{N}$ *,* $d_r \in \mathbb{R}$ *, and* $\lambda_r \in \mathbb{C}$
with $\text{Im}(\lambda) > 0$ for all $r = 1$ *M* such that *with* $\text{Im}(\lambda_r) > 0$ *for all r* = 1, ..., *M such that,*

$$
\left\| f(x) - \operatorname{Im} \sum_{r=1}^{M} \frac{d_r}{x - \lambda_r} \right\|_{\mathcal{L}_w} < \varepsilon,\tag{130}
$$

where the weighted Lipschitz norm is defined in Definition [4.1.](#page-27-3)

Now we are ready to prove Theorem [2.3.](#page-8-0)

Proof of Theorem [2.3](#page-8-0). Recall that we define $f_{\alpha,x_0}(x) := f(n^{\alpha}(x - x_0))$. By the inequality $|1 - e^{ix}| \le |x|$ for all gashes the following all real *x*, we have $|e^{ix} - e^{iy}| \le |x - y|$ for any *x*, *y* real. Then use Jensen's inequality to deduce the following bound for any real-valued functions f h and real number t bound, for any real-valued functions *^f*, *^h* and real number *^t*,

$$
\left| \mathbb{E}\left[e^{itX_{f,\alpha,x_0}^{(n)}}\right]e^{-it\mathbb{E}\left[X_{f,\alpha,x_0}^{(n)}\right]} - \mathbb{E}\left[e^{itX_{h,\alpha,x_0}^{(n)}}\right]e^{-it\mathbb{E}\left[X_{h,\alpha,x_0}^{(n)}\right]}\right| \leq t^2 \operatorname{Var}\left(X_{f-h,\alpha,x_0}^{(n)}\right). \tag{131}
$$

Moreover, define

$$
\sigma_{f,L}^2 := \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy, \qquad \sigma_{f,R}^2 := \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy. \tag{132}
$$

We have for both $j \in \{L, R\}$, by Definition [4.1,](#page-27-3)

$$
\sigma_{f,j}^2 \le ||f||_{\mathcal{L}_w}^2 \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{x+y}{\sqrt{x^4+1} \sqrt{y^4+1}} \right)^2 dx dy. \tag{133}
$$

Use Cauchy residual theorem to compute the double integral $\int \int_{\mathbb{R}^2}$ $\int \frac{x+y}{\sqrt{x^4+1}\sqrt{y^4+1}}$ χ^2 $dx dy = \pi^2$. Hence,

$$
\sigma_{f,j}^2 \le \frac{1}{8} \|f\|_{\mathcal{L}_w}^2.
$$
\n(134)

Moreover, use the Cauchy-Schwartz inequality to obtain for both $j \in \{L, R\}$

$$
\left|\sigma_{f,j}^2 - \sigma_{h,j}^2\right| \le \sigma_{f+h,j}^2 \sigma_{f-g,j}^2 \le \frac{1}{16} \|f - h\|_{\mathcal{L}_w}^2 \|f + h\|_{\mathcal{L}_w}^2.
$$
\n(135)

Then use triangle inequality, [\(131\)](#page-29-0), [\(134\)](#page-29-1), [\(135\)](#page-29-2) and Corollary [4.1,](#page-28-3) to obtain the following, as $n \to \infty$,

$$
\left| \mathbb{E} \left[e^{itX_{f,\alpha,x_0}^{(n)}} \right] e^{-it\mathbb{E} \left[X_{f,\alpha,x_0}^{(n)} \right]} - e^{-\frac{t^2}{2}\sigma_{f,j}^2} \right| \leq \left| \mathbb{E} \left[e^{itX_{f,\alpha,x_0}^{(n)}} \right] e^{-it\mathbb{E} \left[X_{f,\alpha,x_0}^{(n)}} \right] - \mathbb{E} \left[e^{itX_{h,\alpha,x_0}^{(n)}} \right] e^{-it\mathbb{E} \left[X_{h,\alpha,x_0}^{(n)}} \right] \right|
$$
\n
$$
+ \left| e^{-\frac{t^2}{2}\sigma_{f,j}^2} - e^{-\frac{t^2}{2}\sigma_{h,j}^2} \right| + \left| \mathbb{E} \left[e^{itX_{h,\alpha,x_0}^{(n)}} \right] e^{-it\mathbb{E} \left[X_{h,\alpha,x_0}^{(n)}} \right] - e^{-\frac{t^2}{2}\sigma_{h,j}^2} \right|
$$
\n
$$
\leq t^2 \left(\frac{1}{8} ||f - h||_{\mathcal{L}_w}^2 + o(1) \right) + \frac{t^2}{32} ||f - h||_{\mathcal{L}_w}^2 ||f + h||_{\mathcal{L}_w}^2 + \left| \mathbb{E} \left[e^{itX_{h,\alpha,x_0}^{(n)}} \right] e^{-it\mathbb{E} \left[X_{h,\alpha,x_0}^{(n)}} \right] - e^{-\frac{t^2}{2}\sigma_{h,j}^2} \right|.
$$
\n(136)

Let $f \in C_c^1(\mathbb{R})$ and any $\varepsilon > 0$. By Lemma [4.1,](#page-28-0) there exists

$$
h = \operatorname{Im} \sum_{r=1}^{M} \frac{d_r}{x - \lambda_r}
$$

such that

$$
||f - h||_{\mathcal{L}_w} < \varepsilon. \tag{137}
$$

Moreover, since $f \in C_c^1(\mathbb{R})$, we have $||f||_{\mathcal{L}_w} < \infty$. Using triangle inequality we have

$$
\|f+h\|_{\mathcal{L}_w}<2\,\|f\|_{\mathcal{L}_w}+\varepsilon<\infty.
$$

Apply Proposition [4.5](#page-27-2) to *h* and we have

$$
\lim_{n \to \infty} \left| \mathbb{E} \left[e^{itX_{h,\alpha,x_0}^{(n)}} \right] e^{-it \mathbb{E} \left[X_{h,\alpha,x_0}^{(n)} \right]} - e^{-\frac{t^2}{2} \sigma_{h,j}^2} \right| = 0. \tag{138}
$$

Plug [\(137\)](#page-29-3) and [\(138\)](#page-30-2) into [\(136\)](#page-29-4) and we get, for any $j = L$ or R, and any $\varepsilon > 0$,

$$
\limsup_{n\to\infty}\left|\mathbb{E}\left[e^{itX_{f,\alpha,x_0}^{(n)}}\right]e^{-it\mathbb{E}\left[X_{f,\alpha,x_0}^{(n)}\right]}-e^{-\frac{t^2}{2}\sigma_{f,j}^2}\right|\leq\frac{t^2\varepsilon^2}{8}\left(1+\left(\|f\|_{\mathcal{L}_w}+\frac{\varepsilon}{2}\right)^2\right).
$$

Hence

$$
\lim_{n\to\infty}\left|\mathbb{E}\left[e^{itX^{(n)}_{f,\alpha,x_0}}\right]e^{-it\mathbb{E}\left[X^{(n)}_{f,\alpha,x_0}\right]}-e^{-\frac{t^2}{2}\sigma^2_{f,j}}\right|=0.
$$

This completes the proof of Theorem [2.3](#page-8-0) for any $f \in C_c^1(\mathbb{R})$ at both left and right edges. \Box

Note that the proof of Propositions [4.1,](#page-24-0) [4.2,](#page-25-0) [4.4](#page-27-1) and [4.3](#page-26-0) is postponed to Sections [5](#page-30-0) and [6](#page-44-0) below.

5 Proof of Propositions [4.1](#page-24-0) and [4.2](#page-25-0)

In this section we are to prove Propositions [4.1](#page-24-0) and [4.2.](#page-25-0)

In Subsection [5.1,](#page-30-1) we continue the discussion in Subsection [3.5](#page-20-0) and derive Proposition [5.1,](#page-30-3) a general theory of tri-diagonal matrices that will be used to prove Propositions [4.1](#page-24-0) and [4.2.](#page-25-0)

In Subsection [5.2,](#page-32-0) we estimate the resolvent of both $J_{n+1}^{(r)}$ (n) and the middle block of $J_{n\pm 2mn\beta}^{(r)}$ defined in [\(104\)](#page-23-3) and [\(105\)](#page-23-5) by Proposition [5.1.](#page-30-3) Then, complete the proof of Propositions [4.1](#page-24-0) and [4.2.](#page-25-0)

5.1 Inverse of a Tri-diagonal Matrix: Part I

The goal of this subsection is to estimate the inverse of the tri-diagonal symmetric matrix J_N defined in [\(88\)](#page-20-6) entry-wise.

Proposition 5.1. *Consider an N* \times *N non singular tri-diagonal symmetric matrix* J_N *defined as [\(88\)](#page-20-6) with entries* $a_j, b_j \in \mathbb{R}$, $N_1 \in \mathbb{N}$ and $z \in \mathbb{C}$ with Im $z \neq 0$. Let $N_1 \in \mathbb{N}$ with $N_1 < N$. Assume $\text{Re}(b_j - z) > 0$ for all $i > N_0$. $j \geq N_0$.

Let ω_j^+ , ω_j^- be defined as [\(96\)](#page-21-2) and M_j be defined as [\(97\)](#page-22-1). Denote

$$
c = \left| \frac{-\omega_{N-1}^+ a_{N-1} + b_{N-1} - z}{\omega_{N-1}^- a_{N-1} - b_{N-1} + z} \right|,
$$
\n(139)

$$
\varepsilon_1 = (N - N_1) \max_{l \ge N_1} ||M_l||_{\infty}.
$$
 (140)

and

$$
\varepsilon_2 = \prod_{l=N_1}^{N-1} \left| \frac{\omega_l^-}{\omega_l^+} \right|.
$$
\n(141)

Assume that $\varepsilon_1, \varepsilon_2 \in (0, 1)$ *and* $4(1 + c)\varepsilon_1 + c\varepsilon_2 < 1$ *. Then for all* $j \leq N_1 < k$ we have

$$
\left| \left(J_N^{-1} \right)_{j,k} \right| \le \frac{1 + 4(1 + c)\varepsilon_1 + c}{1 - 4(1 + c)\varepsilon_1 - c\varepsilon_2} \left\| J_N^{-1} \right\|_{\infty} \prod_{l=N_1}^{k-1} \left| \omega_l^+ \right|^{-1} . \tag{142}
$$

Proof. Apply the inverse formula for tri-diagonal matrix [\(95\)](#page-21-1) to obtain that for $j \leq N_1 < k$

$$
\left(J_N^{-1}\right)_{j,k} = (-1)^{k-N_1} \frac{\beta_k}{\beta_{N_1}} \left(J_N^{-1}\right)_{j,N_1}.
$$
\n(143)

Recall that β_j is defined recursively by [\(94\)](#page-21-3), i.e.,

$$
\begin{pmatrix} \beta_k \\ \beta_{k-1} \end{pmatrix} = A_k A_{k+1} \cdots A_{N-1} \begin{pmatrix} \beta_N \\ \beta_{N-1} \end{pmatrix}, \quad \beta_N = a_{N-1}, \quad \beta_{N-1} = b_{N-1} - z,
$$
 (144)

where the matrix A_k is defined as [\(93\)](#page-21-4). Recall that the eigenvalues ω_k^+ and $\omega_k^ \overline{k}$ of A_k are defined as [\(96\)](#page-21-2). Then *A_k* can be diagonalized as $A_k = V_k \Omega_k V_k^{-1}$ where, $V_k := \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$ + *k* ω − *k* $\Big)$ and $\Omega_k \coloneqq \Big($ ω + *k* 0 $\int_0^{\pi} \omega_k^$ *k* ! . Further define

$$
C_k \coloneqq V_k^{-1} \left(A_k A_{k+1} \dots A_{N-1} - V_k \left(\prod_{l=k}^{N-1} \Omega_l \right) V_{N-1}^{-1} \right) V_{N-1}, \quad k = 1, 2, \dots, N-1, \quad \text{and } C_N \coloneqq 0. \tag{145}
$$

Rewrite [\(144\)](#page-31-0) for $k = 1, 2, ..., N − 1$ as

$$
\begin{pmatrix} \beta_k \\ \beta_{k-1} \end{pmatrix} = V_k \left(C_k + \prod_{l=k}^{N-1} \Omega_l \right) V_{N-1}^{-1} \begin{pmatrix} \beta_N \\ \beta_{N-1} \end{pmatrix} . \tag{146}
$$

Denote $\tilde{\beta}_1 = \omega_{N-1}^- a_{N-1} - b_{N-1} + z$, $\tilde{\beta}_2 = -\omega_{N-1}^+ a_{N-1} + b_{N-1} - z$ such that $\frac{1}{\omega_{N-1}^-}$ $\omega_{N-1}^{-} - \omega_{N-1}^{+}$ $\begin{pmatrix} \tilde{\beta}_1 \ \tilde{\beta}_2 \end{pmatrix}$! $= V_{N-1}^{-1}$ $\begin{pmatrix} \beta_N \\ \beta_N \end{pmatrix}$ ^β*N*−¹ ! . Hence, we have for $k = 1, \ldots, N - 1$

$$
\beta_k = \frac{\left((C_k)_{1,1} + (C_k)_{2,1} + \prod_{l=k}^{N-1} \omega_l^+\right) \tilde{\beta}_1 + \left((C_k)_{1,2} + (C_k)_{2,2} + \prod_{l=k}^{N-1} \omega_l^-\right) \tilde{\beta}_2}{\omega_{N-1}^- - \omega_{N-1}^+}.
$$
\n(147)

By convention, let $C_N \equiv 0$ and $\prod_{l=N}^{N-1} \equiv 1$. Then we extend [\(147\)](#page-31-1) to $k = 1, ..., N$, since $\beta_N = a_{N-1}$. Use the construction of M_j in [\(97\)](#page-22-1) to write $V_k^{-1}V_{k+1} = M_k + Id$. Then rewrite C_k via the following equality

$$
A_k A_{k+1} \dots A_{N-1} = V_k \Omega_k (Id + M_k) \Omega_{k+1} (Id + M_{k+1}) \dots \Omega_{N-2} (Id + M_{N-2}) \Omega_{N-1} V_{N-1}^{-1}.
$$
 (148)

By assumption $\text{Re}(b_l - z) > 0$, we have $|\omega_l^+| \ge |\omega_l^-|$ $\frac{1}{l}$. Hence, $||\Omega_l||_{\infty} < |ω_l^+|$. Plug [\(148\)](#page-31-2) into [\(145\)](#page-31-3) to estimate

$$
||C_k||_{\infty} \le \left(\prod_{l=k}^{N-2} \left(1 + ||M_l||_{\infty}\right) - 1\right) \prod_{l=k}^{N-1} |\omega_l^+|.
$$
 (149)

Then by [\(140\)](#page-30-4), $0 < \varepsilon_1 < 1$ and the fact that $(1 + x)^y < e^{xy} < 1 + 2xy$ for all $x, y > 0$ with $0 < xy < 1$, we have

$$
||C_k||_{\infty} \le 2\varepsilon_1 \prod_{l=k}^{N-1} |\omega_l^+|.
$$
 (150)

Denote $(C_k)_{j,l}$ to be the entry of matrix C_k at the $j - th$ row and $l - th$ column. Then let

$$
C(k) := ((C_k)_{1,1} + (C_k)_{2,1}) \prod_{l=k}^{N-1} (\omega_l^+)^{-1} + ((C_k)_{1,2} + (C_k)_{2,2}) \frac{\tilde{\beta}_2}{\tilde{\beta}_1} \prod_{l=k}^{N-1} (\omega_l^+)^{-1}.
$$
 (151)

By [\(139\)](#page-30-5) we have \vert $\frac{\tilde{\beta}_2}{\tilde{\beta}_1}$ $= c$ and thus $|C(k)| \le 4(1+c)\varepsilon_1$. Moreover, use [\(151\)](#page-31-4) to rewrite [\(147\)](#page-31-1) as

$$
\beta_k = \frac{\tilde{\beta}_1 \prod_{l=k}^{N-1} \omega_l^+}{\omega_{N-1}^- - \omega_{N-1}^+} \left(1 + \frac{\tilde{\beta}_2}{\tilde{\beta}_1} \prod_{l=k}^{N-1} \frac{\omega_l^-}{\omega_l^+} + C(k) \right). \tag{152}
$$

Then plug [\(152\)](#page-32-1) into [\(143\)](#page-31-5) to obtain that for $j \leq N_1 < k$

$$
\left(J_N^{-1}\right)_{j,k} = (-1)^{k-N_1} \left(J_N^{-1}\right)_{j,N_1} \left(\prod_{l=N_1}^{k-1} \omega_l^+\right)^{-1} \frac{1 + \frac{\tilde{\beta}_2}{\tilde{\beta}_1} \prod_{l=k}^{N-1} \frac{\omega_l^-}{\omega_l^+} + C(k)}{1 + \frac{\tilde{\beta}_2}{\tilde{\beta}_1} \prod_{l=N_1}^{N-1} \frac{\omega_l^-}{\omega_l^+} + C(N_1)}.
$$
\n(153)

Note that $\left| \frac{\omega_l^-}{\omega_l^+} \right|$ $|$ < 1. Thus whenever $4(1 + c)\varepsilon_1 + c\varepsilon_2$ < 1, we have, by triangle inequality,

$$
\left| \left(J_N^{-1} \right)_{j,k} \right| \le \frac{1 + 4(1 + c)\varepsilon_1 + c}{1 - 4(1 + c)\varepsilon_1 - c\varepsilon_2} \left\| J_N^{-1} \right\|_{\infty} \prod_{l=N_1}^{k-1} \left| \omega_l^+ \right|^{-1} \tag{154}
$$

5.2 Proof of Propositions [4.1](#page-24-0)

In this subsection, we use Proposition [5.1](#page-30-3) to prove Propositions [4.1,](#page-24-0) whose assumptions are the same as Theorem [2.3.](#page-8-0)

Let $\beta = \frac{\alpha}{2} + \frac{\epsilon}{3}$ such that $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$ and $m \in \mathbb{N}$. We consider the tri-diagonal matrix J_N , where

$$
N = n + 2mn^{\beta}, \quad b_j = b_{j,n} - x_0, \quad a_j = a_{j,n}, \quad z = \frac{\eta}{n^{\alpha}},
$$
 (155)

where x_0 is given to be near the edge, i.e., [\(24\)](#page-7-4) is assumed. Indeed, we are looking at the case where $J_N = J_{n+1}^{(r)}$ $n_{n+2mn\beta}^{(r)}$ as defined as [\(102\)](#page-23-2). Since the subscript *r* of η_r will not play any role in what follows, we surpass it in this section. To consider the case $J_{n+1}^{(r)}$ $\sum_{n=2mn\beta}^{(r)}$, we will point out in the end of the proof of Propositions [4.1](#page-24-0) that the proof is the same upto relabelling of the indexing.

Now let

$$
I_{n,m}^{(\beta)} := \{ j \in \mathbb{N} : N_0 \le j \le N \}, \quad \text{where } N_0 = n - 2mn^{\beta}, N = n + 2mn^{\beta}.
$$
 (156)

Note that $I_{n,m}^{(\beta)} \subset I_n^{(\alpha,\varepsilon)}$, where $I_n^{(\alpha,\varepsilon)}$ is defined as [\(21\)](#page-7-2). The size of the indexing set is denoted by $\Big|$ *I* (β) *n*,*m* and is of order $O(n^{\beta})$.

In Theorem [2.3,](#page-8-0) our main assumptions are about the coefficients $a_{j,n} \in \mathbb{R}$ and $b_{j,n} \in \mathbb{R}$. For reader's convenience, we list them in terms of a_j and b_j (under the setup [\(155\)](#page-32-2)) as the following.

Condition 5.1 (Theorem [2.3\)](#page-8-0). *There exist some constants* c_0 , $c_1 > 0$ *such that for all j* $\in I_{n,m}^{(\beta)}$, *where* $I_{n,m}^{(\beta)}$ *is defined as* (156), we have *defined as* [\(156\)](#page-32-3)*, we have*

$$
c_0 < |a_j| < c_1, \quad |b_j| < c_1,\tag{157}
$$

$$
|a_j - a_{j-1}| \le \frac{c_1}{n}, \quad |b_j - b_{j-1}| \le \frac{c_1}{n}.
$$
 (158)

Condition 5.2 (Theorem [2.3\)](#page-8-0). As $n \to \infty$,

$$
\max_{j \in I_{n,m}^{(\beta)}} \left| a_j a_{j-2} - a_{j-1}^2 \right| = o(n^{-3\beta + \frac{\alpha}{2}})
$$
\n(159)

$$
\max_{j \in I_{n,m}^{(\beta)}} |(b_{j-1} - a_j)a_{j-2} - (b_{j-2} - a_{j-1})a_{j-1}| = o(n^{-3\beta}). \tag{160}
$$

$$
\max_{j \in I_{n,m}^{(\beta)}} |b_{n-1} - 2\sqrt{a_n a_{n-1}}| = o(n^{-\alpha}), \quad \text{for the left edge }.
$$
 (161)

For the right edge [\(161\)](#page-33-0) should be replaced by $b_{n-1} = -2\sqrt{a_n a_{n-1}} + o(n^{-\alpha})$. In the rest of this section, we will only consider the left edge. The discussion about the right edge of J is the same as that of the left edge of −J, whose diagonals are [−]*aj*,*ⁿ* which is the only reason we do not assume *^aj*,*ⁿ* > 0 in this paper.

Also note that the indexing set $I_{n,m}^{(\beta)}$ here is a subset of that in the Theorem [2.3](#page-8-0) and hence we make a subset of the statement on the statement and the statement of the statement of the statement of the statement of slightly weaker assumption. This difference is only to reduce the notations both in the statement and the proof.

We continue to explain the implications of these conditions by the following lemmas.

Lemma 5.1. *Assume conditions of Theorem [2.3](#page-8-0) are satisfied for the left edge, i.e., given Conditions [5.1](#page-32-4)* and [5.2.](#page-33-1) Then given $0 < \frac{\alpha}{2} < \beta < 1$, we have,

$$
\liminf_{n \to \infty} \min_{j \in I_{n,m}^{(\beta)}} b_j > 0,
$$
\n(162)

and as $n \to \infty$

$$
\max_{j \in I_{n,m}^{(\beta)}} \left| b_{j-1}^2 - 4a_j a_{j-1} \right| = o(n^{-\alpha}).
$$
\n(163)

Proof. Recall that *N* and N_0 are define in [\(156\)](#page-32-3). Then the size of the set $|I_{n,m}^{(\beta)}| = N - N_0 = 4mn^{\beta}$, with $0 < \beta < 1$. By [\(158\)](#page-32-5), we have

$$
\min_{j \in I_{n,m}^{(\beta)}} b_j \ge b_{n-1} - 4c_1 m n^{\beta - 1}.
$$
\n(164)

By (161) , we have

$$
b_{n-1} = 2\sqrt{a_{n-1}a_n} + o(n^{-\alpha}), \quad \text{as } n \to \infty.
$$
 (165)

Moreover, by [\(161\)](#page-33-0) and the lower bound of $|a_j|$ in [\(157\)](#page-32-6), we have $\sqrt{a_{n-1}a_n} > 2c_0 > 0$. Consequently, we have (162) have [\(162\)](#page-33-2).

Now we prove the second statement. By [\(160\)](#page-33-3), together with the lower bound of a_j in Condition [5.2,](#page-33-1) we have that for all $j \in I_{n,m}^{(\beta)}$

$$
\left(\frac{b_{j-1} - a_j}{a_{j-1}}\right) - \left(\frac{b_{j-2} - a_{j-1}}{a_{j-2}}\right) = o(n^{-3\beta}), \quad \text{as } n \to \infty.
$$
 (166)

Note that the size of the index set $|I_{n,m}^{(\beta)}| = O(n^{\beta})$. Then for all $j \in I_{n,m}^{(\beta)}$

$$
\left(\frac{b_{j-1} - a_j}{a_{j-1}}\right) - \left(\frac{b_{n-1} - a_n}{a_{n-1}}\right) = o\left(n^{-2\beta}\right), \quad \text{as } n \to \infty.
$$
 (167)

Using the fact that $\left(\sqrt{\frac{a_j}{a_{j-1}}} - 1\right)^2 = \frac{a_j}{a_{j-1}}$ $\frac{a_j}{a_{j-1}} + 1 - 2\sqrt{\frac{a_j}{a_{j-1}}}$ $\frac{a_j}{a_{j-1}}$, we can rewrite for *j* ∈ *I*_{*n*,*m*}

$$
\frac{b_{j-1} - a_j}{a_{j-1}} = \frac{b_{j-1}}{a_{j-1}} - \left(\sqrt{\frac{a_j}{a_{j-2}}} - 1\right)^2 - 2\sqrt{\frac{a_j}{a_{j-1}}} + 1.
$$
 (168)

Note that by the Condition [5.1](#page-32-4) we have $\left(\sqrt{\frac{a_j}{a_{j-1}}}-1\right)^2 = O(n^{-2})$. Also note that [\(168\)](#page-34-0) holds for all $j \in I_{n,m}^{(\beta)}$ including the special case $j = n$. Hence plugging [\(168\)](#page-34-0) into [\(167\)](#page-33-4) yields

$$
\frac{b_{j-1}}{a_{j-1}} - 2\sqrt{\frac{a_j}{a_{j-1}}} - \frac{b_{n-1}}{a_{n-1}} + 2\sqrt{\frac{a_n}{a_{n-1}}} = o(n^{-2\beta}) + O(n^{-2}), \quad \text{as } n \to \infty.
$$
 (169)

By the assumption [\(161\)](#page-33-0) in Condition [5.2,](#page-33-1) we have $\frac{b_{n-1}}{a_{n-1}} - 2\sqrt{\frac{a_n}{a_{n-1}}}$ $\frac{a_n}{a_{n-1}} = o(n^{-\alpha})$, as $n \to \infty$. Further, by the assumption $0 < \frac{\alpha}{2} < \beta < 1$, we estimate [\(169\)](#page-34-1) to be

$$
\frac{b_{j-1}}{a_{j-1}} - 2\sqrt{\frac{a_j}{a_{j-1}}} = o(n^{-\alpha}), \quad \text{as } n \to \infty.
$$
 (170)

Multiply $\frac{b_{j-1}}{a_{j-1}} + 2\sqrt{\frac{a_j}{a_{j-1}}}$ $\frac{a_j}{a_{j-1}}$ on both sides of [\(170\)](#page-34-2) to obtain that for all *j* ∈ *I*^(β),*m*

$$
\left(\frac{b_{j-1}}{a_{j-1}}\right)^2 = \frac{4a_j}{a_{j-1}} + o(n^{-\alpha}), \quad \text{as } n \to \infty,
$$
\n(171)

which completes the proof. \Box

Note that [\(162\)](#page-33-2) implies that $|\omega_j^-|$ $|\vec{v}_j| \le |\omega_j^+|$, since we are taking the principle square root. However, for the right edge, by the same argument, we have $b_j < 0$ for all $j \in I_{n,m}^{(\beta)}$. In this case, we can swap the definition of ω^+ and ω^- and all conclusions in this section follow with the same argument. Alternatively, for of ω_j^+ and $\omega_j^$ *j* and all conclusions in this section follow with the same argument. Alternatively, for the right edge, we can simply consider $-J_N$ whose diagonals are $-b_j + \frac{\eta}{n^{\frac{\alpha}{2}}}$ and off diagonals are $-a_j$. Hence, all discussions in this section can be directly applied to $(-J_N)^{-1}$, since $-b_j > 0$ for the right edge.

Lemma [5.2](#page-33-1). *Consider* $0 < \frac{\alpha}{2} < \beta < 1$ *. Assume the Conditions* [5.1](#page-32-4) *and* 5.2 *are satisfied. Then we have*

$$
\max_{j \in I_{n,m}^{(\beta)}} \left| \omega_j^- - \left(1 - \left(\frac{-\eta}{n^{\alpha} |a_{N-1}|} \right)^{\frac{1}{2}} \right) sgn(a_{N-1}) \right| = o(n^{-\frac{\alpha}{2}}),\tag{172}
$$

$$
\max_{j \in I_{n,m}^{(\beta)}} \left| \omega_j^+ - \left(1 + \left(\frac{-\eta}{n^{\alpha} |a_{N-1}|} \right)^{\frac{1}{2}} \right) sgn(a_{N-1}) \right| = o(n^{-\frac{\alpha}{2}}). \tag{173}
$$

*The choice of a*_{*N*−1} *in the formulas above is arbitrary. It can be replaced by any* a_j *for* $j \in I_{n,m}^{(\beta)}$ *. Proof.* Recall that $\omega_j^ \frac{1}{i}$ is given by

$$
\omega_j^- = \frac{b_{j-1} - \frac{\eta}{n^{\alpha}} - \left(\left(b_{j-1} - \frac{\eta}{n^{\alpha}} \right)^2 - 4a_{j-1} a_j \right)^{\frac{1}{2}}}{2a_{j-1}}.
$$
\n(174)

By Condition [5.1](#page-32-4) we have $\frac{a_l}{a_{l-1}} = 1 + O(n^{-1})$. Further, by Lemma [5.1,](#page-33-5) we obtain

$$
\left(\frac{b_{j-1}}{2a_{j-1}}\right)^2 - \frac{a_j}{a_{j-1}} = o(n^{-\alpha}), \quad \frac{b_{j-1}}{2|a_{j-1}|} = 1 + o(n^{-\alpha}) + O(n^{-1}), \quad \text{as } n \to \infty.
$$
 (175)

Condition [5.1](#page-32-4) also implies $a_{j-1} = a_{N-1} + O(n^{\beta-1})$. We can then further estimate as $n \to \infty$

$$
\left(\left(\frac{b_{j-1} - \frac{\eta}{n^{\alpha}}}{2a_{j-1}} \right)^2 - \frac{a_j}{a_{j-1}} \right)^{\frac{1}{2}} = \left(\left(\frac{b_{j-1}}{2a_{j-1}} \right)^2 - \frac{a_j}{a_{j-1}} - \frac{b_{j-1}}{2a_{j-1}} \frac{\eta}{n^{\alpha} a_{j-1}} + \left(\frac{\eta}{2n^{\alpha} a_{j-1}} \right)^2 \right)^{\frac{1}{2}}
$$

= $\left(o(n^{-\alpha}) - \left(1 + o(n^{-\alpha}) + O(n^{-1}) \right) \frac{\eta}{2n^{\alpha} \left(|a_{N-1}| + O(n^{\beta - 1}) \right)} + O(n^{2\alpha}) \right)^{\frac{1}{2}} = \left(\frac{-\eta}{n^{\alpha} |a_{N-1}|} \right)^{\frac{1}{2}} + o(n^{-\frac{\alpha}{2}}).$

Hence we proved [\(172\)](#page-34-3) for $\omega_j^ \overline{j}$. The proof for ω_j^+ follows from the same argument. □

Note that for the right edge where $b_j < 0$, we will get the following by the same argument,

$$
\omega_j^- = \left(-1 - \left(\frac{\eta}{n^{\alpha}|a_{N-1}|}\right)^{\frac{1}{2}}\right) sgn(a_{N-1}) + o(n^{-\frac{\alpha}{2}}),\tag{176}
$$

$$
\omega_j^+ = \left(-1 + \left(\frac{\eta}{n^{\alpha}|a_{N-1}|}\right)^{\frac{1}{2}}\right) sgn(a_{N-1}) + o(n^{-\frac{\alpha}{2}}). \tag{177}
$$

Lemma 5.3. *Let* $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$. Assume Conditions [5.1,](#page-32-4) and [5.2](#page-33-1) are satisfied. Then we have as $n \to \infty$ $n \rightarrow \infty$

$$
\max_{j \in I_{n,m}^{(\beta)}} \left| \omega_j^+ - \omega_{j-1}^+ \right| = o(n^{-3\beta + \frac{\alpha}{2}}), \quad \max_{j \in I_{n,m}^{(\beta)}} \left| \omega_j^- - \omega_{j-1}^- \right| = o(n^{-3\beta + \frac{\alpha}{2}}).
$$
\n(178)

Consequently we also have as $n \rightarrow \infty$

$$
\max_{j \in I_{n,m}^{(\beta)}} \|M_j\|_{\infty} = o(n^{-3\beta + \alpha}).
$$
\n(179)

Proof. First note that the difference of equation [\(159\)](#page-33-6) and [\(160\)](#page-33-3) amounts to

$$
\max_{j \in I_{n,m}^{(\beta)}} |b_{j-1}a_{j-2} - b_{j-2}a_{j-1}| = o(n^{-3\beta + \frac{\alpha}{2}}), \quad \text{as } n \to \infty.
$$
 (180)

Recall the definition of ω_j^+ and $\omega_j^ \frac{1}{i}$ in [\(96\)](#page-21-2) and we have $\omega_j^+ - \omega_j^ \frac{1}{j}$ = $\frac{\left((b_{j-1}-\frac{\eta}{n^{\alpha}})^2-4a_ja_{j-1}\right)^{\frac{1}{2}}}{a_{j-1}}$ $\frac{a_{j-1}}{a_{j-1}}$, where the square root is taken as the principle branch. By Lemma [5.1](#page-33-5) and [\(157\)](#page-32-6) we have $|\omega_j^+ - \omega_j^-|$
Moreover $j_j^{-1} = O(n^{\frac{\alpha}{2}})$ for all $j \in I_{n,m}^{(\beta)}$. Moreover,

$$
\omega_j^+ - \omega_{j-1}^+ = \frac{b_{j-1}}{2a_{j-1}} - \frac{b_{j-2}}{2a_{j-2}} - \frac{\eta}{2n^{\alpha}} \left(\frac{1}{a_{j-1}} - \frac{1}{a_{j-2}} \right) + \left(\left(\frac{b_{j-1} - \frac{\eta}{n^{\alpha}}}{2a_{j-1}} \right)^2 - \frac{a_j}{a_{j-1}} \right)^{\frac{1}{2}} - \left(\left(\frac{b_{j-2} - \frac{\eta}{n^{\alpha}}}{2a_{j-2}} \right)^2 - \frac{a_{j-1}}{a_{j-2}} \right)^{\frac{1}{2}}
$$
The first part can be estimated as $\frac{b_{j-1}}{2a_{j-1}} - \frac{b_{j-2}}{2a_{j-1}}$ $\frac{b_{j-2}}{2a_{j-2}} = o(n^{-3\beta + \frac{\alpha}{2}})$, by [\(180\)](#page-35-0). The second part can be estimated as $\frac{1}{2n^{\alpha}}$ $\left(\frac{1}{a_{j-1}} - \frac{1}{a_{j-2}}\right)$ $= O(n^{-\alpha-1})$, by Condition [5.1.](#page-32-0) For the third part, we have

$$
\left(\left(\frac{b_{j-1}-\frac{\eta}{n^{\alpha}}}{2a_{j-1}}\right)^2-\frac{a_j}{a_{j-1}}\right)^{\frac{1}{2}}-\left(\left(\frac{b_{j-2}-\frac{\eta}{n^{\alpha}}}{2a_{j-2}}\right)^2-\frac{a_{j-1}}{a_{j-2}}\right)^{\frac{1}{2}}=\frac{\left(\frac{b_{j-1}-\frac{\eta}{n^{\alpha}}}{2a_{j-1}}-\frac{b_{j-2}-\frac{\eta}{n^{\alpha}}}{2a_{j-2}}\right)\left(\frac{b_{j-1}-\frac{\eta}{n^{\alpha}}}{2a_{j-1}}+\frac{b_{j-2}-\frac{\eta}{n^{\alpha}}}{2a_{j-2}}\right)-\frac{a_j}{a_{j-1}}+\frac{a_{j-1}}{a_{j-2}}}{\left(\left(\frac{b_{j-1}-\frac{\eta}{n^{\alpha}}}{2a_{j-1}}\right)^2-\frac{a_j}{a_{j-1}}\right)^{\frac{1}{2}}+\left(\left(\frac{b_{j-2}-\frac{\eta}{n^{\alpha}}}{2a_{j-2}}\right)^2-\frac{a_{j-1}}{a_{j-2}}\right)^{\frac{1}{2}}}.
$$
\n(181)

By the same argument as before, we have 1 over the denominator is also of order $O(n^{\frac{3}{2}})$. For the numerator, Lemma [5.1](#page-33-0) together with [\(157\)](#page-32-1) implies that $\frac{b_{j-1}-\frac{\eta}{n^{\alpha}}}{2a_{j-1}} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & a_{j-1} & a_{j-1} \\ 0 & a_{j-1} & a_{j-1} \end{bmatrix}$ *aj aj*−¹ $\frac{1}{2}$ + $O(n^{-\alpha})$. Moreover, Condition [5.1](#page-32-0) implies *aj* $\frac{a_j}{a_{j-1}} = 1 + O(n^{-1})$. Since $\alpha \in (0, 2)$, we have $\frac{b_{j-1} - \frac{\eta}{n^{\alpha}}}{2a_{j-1}} + \frac{b_{j-2} - \frac{\eta}{n^{\alpha}}}{2a_{j-2}} = 2 + O(n^{-\frac{\alpha}{2}})$. The numerator of [\(181\)](#page-36-0) now becomes

$$
\frac{b_{j-1} - a_j}{a_{j-1}} - \frac{b_{j-2} - a_{j-1}}{a_{j-2}} - \frac{\eta}{n^{\alpha}} \left(\frac{1}{a_{j-1}} - \frac{1}{a_{j-2}} \right) + \left(\frac{b_{j-1}}{a_{j-1}} - \frac{b_{j-2}}{a_{j-2}} \right) O(n^{-\frac{\alpha}{2}}).
$$
(182)

The sum of the first two terms is $o(n^{-3\beta})$ by [\(160\)](#page-33-1). The third term is order $O(n^{-\alpha-1})$, by Condition [5.1.](#page-32-0) For the rest term, we have $\frac{b_{j-1}}{a_{j-1}} - \frac{b_{j-2}}{a_{j-2}}$ $\frac{b_{j-2}}{a_{j-2}} = o(n^{-3\beta + \frac{\alpha}{2}})$, by [\(180\)](#page-35-0). Then [\(182\)](#page-36-1) is of order $o(n^{3\beta})$, by the assumption $\beta < \frac{\alpha+1}{3}$. Combing the arguments above, we have $\omega_j^+ - \omega_{j-1}^+ = o(n^{-3\beta + \frac{\alpha}{2}})$. Similarly, we also have $\omega_j^ \bar{j} - \omega_{j-1}^- =$ $o(n^{-3\beta+\frac{\alpha}{2}})$. Then by the definition of *M_j* [\(97\)](#page-22-0), we have $||M_j||_{\infty} = o(n^{-3\beta+\alpha})$ \Box

Now we are ready to verify the Proposition [4.1.](#page-24-0) Recall that \mathcal{J} is the Jacobi matrix with entries $\{a_{j,n}, b_{j,n}\}$. The truncated operators $J_{n+1}^{(r)}$ $J_{n+2mn\beta}^{(r)}$, $J_{n\pm}^{(r)}$ $F_{n+2mn\beta}$, $F_{n+2mn\beta}$ and $F_{n\pm 2mn\beta}$ are defined in [\(104\)](#page-23-0), [\(105\)](#page-23-1) and [\(106\)](#page-23-2).

Proof of Proposition [4.1.](#page-24-0) First, consider

$$
J_N = J_{n+2mn\beta}^{(r)}
$$

It is sufficient to verify the conditions in Proposition [5.1.](#page-30-0)

For all $l \in I_{n,m}^{(\beta)}$, by Lemma [5.1](#page-33-0) we have $b_l > 0$. By Lemma [5.2](#page-34-0) we have $\omega_l^ \bar{l}$ = 1 + $O(n^{-\frac{\alpha}{2}})$ and − $\frac{1}{l} = 1 + O(n^{-\frac{\alpha}{2}})$. Hence

$$
\left| \frac{-\omega_{n+2m n\beta-1}^{+} a_{n+2m n\beta-1} + b_{n+2m n\beta-1} - z}{\omega_{n+2m n\beta-1}^{-} a_{n+2m n\beta-1} - b_{n+2m n\beta-1} + z} \right| = 1 + O(n^{-\frac{\alpha}{2}}), \text{ as } n \to \infty.
$$
 (183)

This shows [\(139\)](#page-30-1) is bounded.

Let *M*^{*j*} be defined as [\(97\)](#page-22-0). By Lemma [5.3](#page-35-1) we have as $n \to \infty$

$$
\left| I_{n,m}^{(\beta)} \right| \max_{l \in I_{n,m}^{(\beta)}} ||M_l||_{\infty} = o(n^{-2\beta + \alpha}) \tag{184}
$$

This shows [\(140\)](#page-30-2) to be $\varepsilon_1 = o(n^{-2\beta+\alpha})$. Moreover, use Lemma [5.2](#page-34-0) to estimate that there exist constants $d > 0$ and $n_0 \in \mathbb{N}$ such that for all $n > n_0$ and all integer N_1 with $N_0 \le N_1 \le N$ *d* > 0 and *n*⁰ ∈ N such that for all *n* > *n*⁰ and all integer *N*₁ with *N*⁰ ≤ *N*₁ < *N*

$$
\prod_{l=N_1}^{N-1} \left| \frac{\omega_l^-}{\omega_l^+} \right| \le \left(1 - \frac{d}{n^{\frac{\alpha}{2}}} \right)^{N-N_1} \le e^{-dn^{-\frac{\alpha}{2}}(N-N_1)}.
$$
\n(185)

This shows [\(141\)](#page-30-3) to be $\varepsilon_2 = O(e^{-d'n^{\beta - \frac{\alpha}{2}}})$ which is exponentially small.

Then all assumptions of Proposition [5.1](#page-30-0) are cleared. Hence, for all $j \leq N_1 < k$ we have

$$
\left| \left(J_N^{-1} \right)_{j,k} \right| \le c \left\| J_N^{-1} \right\|_{\infty} e^{-dn^{-\frac{\alpha}{2}} (k - N_1)}.
$$
\n(186)

for some constant *c* > 0. Then take $N_1 = k - n^{\beta} \ge N_0$ to obtain [\(107\)](#page-24-1).
Note that above argument is taken as $h_1 > 0$ as assumed in (162).

Note that above argument is taken as $b_j > 0$ as assumed in [\(162\)](#page-33-2). For $b_j < 0$, we consider $-J_N$ whose diagonals are $-b_j + \frac{\eta}{n^{\frac{\alpha}{2}}}$ and off diagonals are $-a_j$. Apply the result above, and we reach the same estimate for $-J_N⁻¹$.

Note that the entry $(J_{n+1}^{(r)})$ *n*±2*mn*β Í *j*,*k* is the same as $(J_{n+1}^{(r)})$ *n*+2*mn*β Í *j*,*k* for all *j*, $k \in I_{n,m}^{(\beta)}$ and the rest of the entries are in the trivial blocks. Hence, the result [\(108\)](#page-24-2) for $J_N = J_{n+1}^{(r)}$ (n) ^(r) follows the same approach by relabelling the indices appropriately.

Then, the estimates [\(109\)](#page-24-3) and [\(110\)](#page-24-4) follow from the definition of $F_{n+2mn\beta}$ and $F_{n\pm 2mn\beta}$ in [\(104\)](#page-23-0) and [\(106\)](#page-23-2) and triangle inequality and triangle inequality.

5.3 Comparison in the Trace Norm

The following lemma is essential to prove Proposition [4.2.](#page-25-0)

Lemma 5.4. *Let* $0 < \frac{\alpha}{2} < \beta < 1$ *. Let* $m \in \mathbb{N}$ *. Assume conditions in Proposition [4.2](#page-25-0) are satisfied. Then for* $am \leq k$ and $n > m$ for some constant $n \in \mathbb{N}$ we have *any* l ≤ *m and n* > *n*₀ *for some constant n*₀ ∈ $\mathbb N$ *we have*

$$
\left\| Q_{n+ln\beta} F_{n+2mn\beta}^l P_n \right\|_1 \leq C_m e^{-d' n^{\beta - \frac{\alpha}{2}}},\tag{187}
$$

$$
\left\| Q_{n+ln\beta} F^l_{n\pm 2mn\beta} P_n \right\|_1 \leq C_m e^{-d'n\beta - \frac{\alpha}{2}},\tag{188}
$$

$$
\left\| \left(F^{l} - F^{l}_{n+2mn\beta} \right) P_{n} \right\|_{1} \leq C_{m} e^{-d' n^{\beta - \frac{\alpha}{2}}}, \tag{189}
$$

where C^m > ⁰ *is a constant depends on m and d*′ > ⁰ *is also an universal constant.*

Proof. To shorten the notation in the proof, we take

 $N = n + 2mn^{\beta}$.

Recall the trace norm inequality $||A||_1 \leq \sum_{k_1,k_2} |(A)_{k_1,k_2}|$ for any matrix *A*. By Proposition [4.1,](#page-24-0) we have

$$
\left\|Q_{n+n^{\beta}}F_NP_n\right\|_1 \le \sum_{j=n+n^{\beta}+1}^N \sum_{k=1}^n C_0 n^{1+\alpha} e^{-d_0 n^{\beta-\frac{\alpha}{2}}} = C_0 (2m-1) n^{2+\alpha+\beta} e^{-d_0 n^{\beta-\frac{\alpha}{2}}},\tag{190}
$$

similarly,

$$
\left\|Q_{n+(l+1)n^{\beta}}F_{N}P_{n+ln^{\beta}}\right\|_{1} \leq \sum_{j=n+(l+1)n^{\beta}+1}^{N} \sum_{k=1}^{n+ln^{\beta}} C_{0}n^{1+\alpha}e^{-d_{0}n^{\beta-\frac{\alpha}{2}}} = C_{0}(2m-l-1)n^{1+\alpha+\beta}(n+ln^{\beta})e^{-d_{0}n^{\beta-\frac{\alpha}{2}}}.
$$
 (191)

We will prove the following statement for all $l \leq m$ by induction:

$$
\left\|Q_{n+ln\beta}F_N^lP_n\right\|_1 \leq C_0(2m-1)ln^{1+\alpha+\beta}(n+mn^{\beta})\|F_N\|_{\infty}^{l-1}e^{-d_0n^{\beta-\frac{\alpha}{2}}}. \tag{192}
$$

For $l = 1$ the statement follows by [\(190\)](#page-37-0). Then write $F_N^{l+1} = F_N(P_{n+ln\beta} + Q_{n+ln\beta})F_N^l$. With the triangle inequality and trace norm inequality $||AB||_1 \leq ||A||_1 ||B||_{\infty}$, we get

$$
\left\|Q_{n+(l+1)n^{\beta}}F_N^{l+1}P_n\right\|_1 \le \left\|Q_{n+(l+1)n^{\beta}}F_NP_{n+ln^{\beta}}\right\|_1 \left\|F_N\right\|_{\infty}^l + \left\|F_N\right\|_{\infty} \left\|Q_{n+ln^{\beta}}F_N^lP_n\right\|_1. \tag{193}
$$

Then apply [\(191\)](#page-37-1) to the first trace norm and the induction hypothesis [\(192\)](#page-37-2) for the second trace norm above, and we obtain

$$
\left\|Q_{n+(l+1)n^{\beta}}F_N^{l+1}P_n\right\|_1 \le C_0(2m-1)(l+1)n^{1+\alpha+\beta}(n+mn^{\beta})\|F_N\|_{\infty}^l e^{-d_0n^{\beta-\frac{\alpha}{2}}}.
$$
 (194)

This concludes the induction.

Moreover since \mathcal{J} is real symmetric and Im(η)/ $n^{\alpha} \neq 0$, we have \mathbb{I} \mathbb{I} \mathbb{I} $\left(J_N^{(r)}\right)$ $\binom{r}{N}^{-1}\Big|_{\infty}$ $\leq \frac{n^{\alpha}}{|\text{Im}(\eta_r)|}$. Hence $||F_N||_{\infty} \leq$ \sum_{r} *cr* $\frac{1}{2}$ n^{α} . By inserting this in [\(192\)](#page-37-2), we see [\(187\)](#page-37-3) holds.

Im(η*r*) The second statement [\(187\)](#page-37-3) follows in the same way, since Proposition [4.1](#page-24-0) also holds for $F_{n\pm 2mn\beta}$.

For the third statement [\(189\)](#page-37-4), recall that $J^{(r)}$ is defined in [\(102\)](#page-23-3) and we define

$$
E := J^{(r)} - P_N J^{(r)} P_N - Q_N J^{(r)} Q_N.
$$
\n(195)

Note that *E* is independent of *r*. It is because *E* is a semi-infinite matrix with only two entries non zero, i.e. $(E)_{N+1,N} = (E)_{N,N+1} = a_N$. This also implies $E = E P_{N+1} Q_{N-1}$.

Hence by the resolvent identity, i.e., [\(43\)](#page-12-0) and the comment thereafter, we have

$$
F - F_N = \sum_r c_r G_{x_0 + \frac{\eta r}{n^{\alpha}}} \left(Q_N - E P_{N+1} Q_{N-1} - Q_N J^{(r)} Q_N \right) \left(J_N^{(r)} \right)^{-1} . \tag{196}
$$

Further by telescopic sum, we have

$$
\left(F^{l} - F_{N}^{l}\right)P_{n} = \sum_{j=0}^{l-1} F^{j} \left(F - F_{N}\right) F_{N}^{l-j-1} P_{n}
$$
\n
$$
= \sum_{j=0}^{l-1} \sum_{r} F^{j} c_{r} G_{x_{0} + \frac{\eta_{r}}{n^{q}}} \left(Q_{N} - E P_{N+1} Q_{N-1} - Q_{N} J^{(r)} Q_{N}\right) \left(J_{N}^{(r)}\right)^{-1} F_{N}^{l-j-1} P_{n}.\tag{197}
$$

Note that $Q_N(J_N^{(r)})$ $\binom{r}{N}^{-1} F_N^{l-j-1}$ N^{l-j-1} = $(\sum_{r=1}^{2M} c_r)^{l-j-1}Q_N$, by definition $J_N^{(r)}$ N_N and F_N in [\(104\)](#page-23-0). Also, $\left(J_N^{(r)}\right)$ $\binom{r}{N}^{-1} F_N^{l-j-1}$ $P_N^{l-j-1}P_n = P_N(f_N^{(r)})$ $\binom{r}{N}^{-1} F_N^{l-j-1}$ $\sum_{N}^{N-1} P_n$, since *N* > *n*. Hence, by $Q_N P_N = 0$,

$$
\left(F^{l} - F_{N}^{l}\right)P_{n} = -\sum_{j=0}^{l-1} \sum_{r} F^{j} c_{r} G_{x_{0} + \frac{\eta_{r}}{n^{\alpha}}} E P_{N+1} Q_{N-1} \left(J_{N}^{(r)}\right)^{-1} F_{N}^{l-j-1} P_{n}.
$$
\n(198)

By the triangle inequality and the trace norm inequality, we find

$$
\left\| \left(F^{l} - F_{N}^{l} \right) P_{n} \right\|_{1} \leq \sum_{j=0}^{l-1} \sum_{r} |c_{r}| \| F \|_{\infty}^{j} \left\| G_{x_{0} + \frac{\eta r}{n^{\alpha}}} \right\|_{\infty} \| E \|_{\infty} \left\| P_{N+1} Q_{N-1} \left(J_{N}^{(r)} \right)^{-1} F_{N}^{l-j-1} P_{n} \right\|_{1}.
$$
 (199)

Write
$$
(J_N^{(r)})^{-1} = (J_N^{(r)})^{-1} (P_{n+(l-j-1)n\beta} + Q_{n+(l-j-1)n\beta})
$$
. Define, for $j < l - 1$,

$$
R_{l-j-1} := \left\| (J_N^{(r)})^{-1} \right\|_{\infty} \left\| Q_{n+(l-j-1)n\beta} F_N^{l-j-1} P_n \right\|_1, \quad R_0 \equiv 0.
$$

Use the triangle inequality and the trace norm inequality $||AB||_1 \leq ||A||_{\infty}||B||_1$ and we find

$$
\left\| P_{N+1} Q_{N-1} \left(J_N^{(r)} \right)^{-1} F_N^{l-j-1} P_n \right\|_1 \le \left\| P_{N+1} Q_{N-1} \left(J_N^{(r)} \right)^{-1} P_{n+(l-j-1)n\beta} \right\|_1 \left\| F_N \right\|_{\infty}^{l-j-1} + R_{l-j-1},\tag{200}
$$

For *^j* < *^l* [−] 1 we can estimate *^Rl*−*j*−¹ via [\(192\)](#page-37-2) to be exponentially small for large *ⁿ* i.e.,

$$
R_{l-j-1} \leq C_0 (2m-1)(l-j-1)n^{1+\alpha+\beta} (n+mn^{\beta}) \left\| \left(J_N^{(r)} \right)^{-1} \right\|_{\infty} ||F_N||_{\infty}^{l-j-2} e^{-d_0 n^{\beta-\frac{\alpha}{2}}}.
$$
 (201)

Recall that we let $N = n + 2mn^{\beta}$ and $l \leq m$. Using the trace norm inequality $||A||_1 \leq \sum_{k_1,k_2} |(A)_{k_1,k_2}|$ a matrix A and estimating the entries via Proposition 4.1 we have *A* and estimating the entries via Proposition [4.1,](#page-24-0) we have

$$
\left\| P_{N+1} Q_{N-1} \left(J_N^{(r)} \right)^{-1} P_{n+(l-j-1)n^{\beta}} \right\|_1 \leq \sum_{j=N}^{N+1} \sum_{k=1}^{n+(l-j-1)n^{\beta}} \left| \left(\left(J_N^{(r)} \right)^{-1} \right)_{j,k} \right|
$$

$$
\leq \sum_{j=N}^{N+1} \sum_{k=1}^{n+(l-j-1)n^{\beta}} C_0^{(r)} n^{1+\alpha} e^{-d_0 n^{\beta - \frac{\alpha}{2}}} \leq 2 C_0^{(r)} (n+ln^{\beta}) n^{1+\alpha} e^{-d_0 n^{\beta - \frac{\alpha}{2}}}, \quad (202)
$$

where $C_0^{(r)}$ $\binom{r}{0} > 0$ is a constant.

Note that the sum over r is finite and independent of n . Use the estimates [\(200\)](#page-39-0), [\(201\)](#page-39-1), and [\(202\)](#page-39-2) into [\(199\)](#page-38-0) to obtain that for all $l \leq m$

$$
\left\| \left(F^{l} - F_{N}^{l} \right) P_{n} \right\|_{1} \leq |a_{N}| C'_{m} e^{-d' n^{\beta - \frac{\alpha}{2}}},
$$

where $d' \in (0, d_0)$ and $C'_m > 0$ are some universal constants. Note that $||E||_{\infty} = |a_N| < c_1$ by Condition [5.1.](#page-32-0)
The lemma is concluded The lemma is concluded. □ □

Lemma 5.5. *Let* $0 < \frac{\alpha}{2} < \beta < 1$ *. Let* $m \in \mathbb{N}$ *. Then for any* $l \leq m$ *we have*

$$
\left\| P_{n-ln^{\beta}} F_{n\pm 2mn^{\beta}}^l Q_n \right\|_1 \leq C_m e^{-d'n^{\beta - \frac{\alpha}{2}}} \tag{203}
$$

$$
\left\| \left(F_{n+2mn\beta}^l - F_{n+2mn\beta}^l \right) Q_{n-mn\beta} \right\|_1 \leq C_m e^{-d' n^{\beta - \frac{\alpha}{2}}} \tag{204}
$$

where a_{n-2mn} *is the* $n-2mn^{\beta}+1$, $n-2mn^{\beta}$ -th entry of \mathcal{J} , $C_m > 0$ *is a constant depends on m and d'* > 0 *is* also a universal constant *also a universal constant.*

Proof. This proof follows the same recipe as the one of Lemma [5.4.](#page-37-5) Using the trace norm inequality $||A||_1$ ≤ $\sum_{j,k} |(A)_{j,k}|$ for any matrix *A* and applying the second statement of Proposition [4.1,](#page-24-0) we obtain

$$
\left\|P_{n-n^{\beta}}F_{n\pm 2mn^{\beta}}Q_n\right\|_{1} \le \sum_{j=n-2mn^{\beta}+1}^{n-(1+m)n^{\beta}} \sum_{k=n-mn^{\beta}+1}^{n+2mn^{\beta}} C_0 n^{1+\alpha} e^{-d_0 n^{\beta-\frac{\alpha}{2}}} = C_0 (2m-1) 2mn^{1+\alpha+2\beta} e^{-d_0 n^{\beta-\frac{\alpha}{2}}},\qquad(205)
$$

similarly,

$$
||P_{n-(l+1)n^{\beta}}F_{n\pm 2mn^{\beta}}Q_{n-ln^{\beta}}||_1 \leq \sum_{j=n-2mn^{\beta}+1}^{n-(l+1)n^{\beta}} \sum_{k=n-ln^{\beta}+1}^{n+2mn^{\beta}} C_0 n^{1+\alpha} e^{-d_0 n^{\beta-\frac{\alpha}{2}}} \\ = C_0 (2m-l-1)(2m+l)n^{1+\alpha+2\beta} e^{-d_0 n^{\beta-\frac{\alpha}{2}}}.
$$
 (206)

We will prove the following statement for all $l \leq m$ by induction:

$$
||P_{n-ln} \mathcal{F}_{n\pm 2mn}^l \mathcal{Q}_n||_1 \le 3C_0(2m-1)mn^{1+\alpha+2\beta}l||F_{n\pm mn\beta}||_{\infty}^{l-1}e^{-d_0n^{\beta-\frac{\alpha}{2}}}.
$$
 (207)

For $l = 1$ the statement follows by [\(205\)](#page-39-3). Then write $F_{l+1}^{l+1} = F_{n\pm 2mn\beta}(P_{n+ln\beta} + Q_{n+ln\beta})F_{l,n}^{l}$ *n*±2*mn*β . With the triangle inequality and trace norm inequality $||AB||_1 \le ||A||_1 ||B||_{\infty}$, we get

$$
\|P_{n-(l+1)n^{\beta}}F_{n\pm 2mn^{\beta}}^{l+1}Q_{n}\|_{1} \leq \|P_{n-(l+1)n^{\beta}}F_{n\pm 2mn^{\beta}}Q_{n-ln^{\beta}}\|_{1} \|F_{n\pm 2mn^{\beta}}\|_{\infty}^{l} + \|F_{n\pm 2mn^{\beta}}\|_{\infty} \|P_{n-ln^{\beta}}F_{n\pm 2mn^{\beta}}Q_{n}\|_{1} \quad (208)
$$

Then apply [\(206\)](#page-39-4) to the first trace norm and the induction hypothesis [\(207\)](#page-40-0) for the second trace norm above, and we obtain

$$
\left\| P_{n-(l+1)n^{\beta}} F_{n\pm 2mn^{\beta}}^{l+1} Q_n \right\|_1 \le 3C_0 (2m-1)mn^{1+\alpha+2\beta} (l+1) \|F_{n\pm mn^{\beta}}\|_{\infty}^l e^{-d_0 n^{\beta-\frac{\alpha}{2}}}.
$$
 (209)

This concludes the induction.

For the second statement, to shorten the notation, let us denote

$$
\tilde{N}=n-2mn^{\beta}.
$$

Let

$$
E := J_{n+2mn\beta}^{(r)} - P_{\tilde{N}} J_{n+2mn\beta}^{(r)} P_{\tilde{N}} - Q_{\tilde{N}} J_{n+2mn\beta}^{(r)} Q_{\tilde{N}}.
$$
 (210)

Note that *E* is independent of *r*. It is because that *E* is a semi-infinite matrix with only two entries non zero, i.e. $(E)_{\tilde{N}+1,\tilde{N}} = (E)_{\tilde{N},\tilde{N}+1} = a_{\tilde{N}}$. This also implies that $E = EP_{\tilde{N}+1}Q_{\tilde{N}-1}$. Moreover, $\left(J_{n\pm}^{(r)}\right)$ *n*±2*mn*β \int ^{-*l*} Q _{*n*−*mn*^β =} $(J_{n+2mn\beta}^{(r)} - E)^{-l} Q_{n-mn\beta}$. Hence

$$
F_{n+2mn\beta} - F_{n\pm 2mn\beta} = \sum_{r} c_r \left(J_{n+2mn\beta}^{(r)} \right)^{-1} \left(-EP_{\tilde{N}+1} Q_{\tilde{N}-1} - P_{\tilde{N}} J_{n+2mn\beta}^{(r)} P_{\tilde{N}} \right) \left(J_{n\pm 2mn\beta}^{(r)} \right)^{-1}
$$

Further by telescopic sum, we have

$$
\left(F_{n+2m n\beta}^l - F_{n\pm 2m n\beta}^l\right)Q_{n-m n\beta} = -\sum_{j=0}^{l-1}\sum_r F_{n+2m n\beta}^j c_r \left(J_{n+2m n\beta}^{(r)}\right)^{-1} E P_{\tilde{N}+1} Q_{\tilde{N}-1} \left(J_{n\pm 2m n\beta}^{(r)}\right)^{-1} F_{n\pm 2m n\beta}^{l-j-1} Q_{n-m n\beta},
$$

which follows from $P_{\tilde{N}}\left(J_{n+1}^{(r)}\right)$ *n*±2*mn*β ich follows from $P_{\tilde{N}}\left(J_{n\pm 2mn\beta}^{(r)}\right)^{-1}F_{n\pm 2mn\beta}^{l-j-1}Q_{n-mn\beta} = 0.$
By triangle inequality and trace norm inequality, we have

$$
\left\| \left(F_{n+2mn^{\beta}}^{l} - F_{n+2mn^{\beta}}^{l} \right) Q_{n-mn^{\beta}} \right\|_{1} \leq \sum_{j=0}^{l-1} \sum_{r} |c_{r}| \| F_{n+2mn^{\beta}} \|_{\infty}^{j} \left\| \left(J_{n+2mn^{\beta}}^{(r)} \right)^{-1} \right\|_{\infty} \|E\|_{\infty} \left\| P_{\tilde{N}+1} Q_{\tilde{N}-1} \left(J_{n\pm 2mn^{\beta}}^{(r)} \right)^{-1} F_{n\pm 2mn^{\beta}}^{l-j-1} Q_{n-mn^{\beta}} \right\|_{1} . \tag{211}
$$

Write $(J_{n+1}^{(r)}$ *n*±2*mn*β $J^{-1} = (J^{(r)}_{n+1})$ *n*±2*mn*β $\int_{0}^{-1} (P_{n-(l-j-1+m)n^{\beta}} + Q_{n-(l-j-1+m)n^{\beta}})$, use triangle inequality and we have

$$
\left\| P_{\tilde{N}+1} Q_{\tilde{N}-1} \left(J_{n\pm 2mn^{\beta}}^{(r)} \right)^{-1} F_{n\pm 2mn^{\beta}}^{l-j-1} Q_{n-mn^{\beta}} \right\|_{1} \leq \left\| P_{\tilde{N}+1} Q_{\tilde{N}-1} \left(J_{n\pm 2mn^{\beta}}^{(r)} \right)^{-1} Q_{n-(l-j-1+m)n^{\beta}} \right\|_{1} \left\| F_{n+2mn^{\beta}} \right\|_{\infty}^{l-j-1} + \tilde{R}_{l-j-1}
$$

where $\tilde{R}_{l-j-1} = \|\cdot\|$ $\left(J_{n}^{(r)}\right)$ *n*±2*mn*β $\Big)^{-1} \Big\|_{\infty}$ $\left\| P_{n-(l-j-1+m)n^{\beta}} F_{n\pm 2mn^{\beta}}^{l-j-1} Q_{n-mn^{\beta}} \right\|_{1}$. Note that $\tilde{R}_0 = 0$. For $j < l-1$ we can estimate \tilde{R}_{l-j-1} via [\(207\)](#page-40-0) to be

$$
\tilde{R}_{l-j-1} \le 4C_0(m-1)mn^{1+\alpha+2\beta}(l-j-1)\|J_{n\pm mn\beta}^{-1}\|_{\infty}^{l-j-1}e^{-d_0n^{\beta-\frac{\alpha}{2}}}
$$

Further using the trace norm inequality $||A||_1 \le \sum_{j,k} |(A)_{j,k}|$ for any matrix *A* and estimating the entries via
Proposition 4.1, we have Proposition [4.1,](#page-24-0) we have

$$
||P_{\tilde{N}+1}Q_{\tilde{N}-1}J_{n\pm 2mn\beta}^{-1}Q_{n-(l-j-1+m)n\beta}||_1 \leq \sum_{k=n-(l-j-1+m)n\beta+1}^{n+2mn\beta} C_0 n^{1+\alpha} n^{-d_0 n^{\beta-\frac{\alpha}{2}}} \leq C_0 (3m+l)n^{1+\alpha+\beta} e^{-d_0 n^{\beta-\frac{\alpha}{2}}}
$$

Assembling the above estimates into [\(211\)](#page-40-1), we have for all $l \leq m$

$$
\left\| \left(F_{n+2mn^{\beta}}^{l} - F_{n\pm 2mn^{\beta}}^{l} \right) Q_{n-mn^{\beta}} \right\|_{1} \leq \| E \|_{\infty} C'_{m} e^{-d' n^{\beta - \frac{\alpha}{2}}}
$$

where $d' \in (0, d_0)$ and $C'_m > 0$ some universal constant. Note that $||E||_{\infty} = |a_{\tilde{N}}| < c_1$. This concludes the lamma l emma. $□$

5.4 Proof of Proposition [4.2](#page-25-0)

Now we are ready to prove the Proposition [4.2](#page-25-0) by Lemmas [5.4](#page-37-5) and [5.5.](#page-39-5)

Proof of Proposition [4.2.](#page-25-0) Recall that for any linear operator $\mathcal A$ we write by [\(67\)](#page-16-0)

$$
C_m^{(n)}(\mathcal{A})=m!\sum_{j=2}^m\frac{(-1)^{j+1}}{j}\sum_{l_1+\cdots+l_j=m,l_i\geq 1}\frac{\mathrm{Tr}(\mathcal{A})^{l_1}P_n\cdots(\mathcal{A})^{l_j}P_n-\mathrm{Tr}(\mathcal{A}^mP_n)}{l_1!\cdots l_j!}.
$$

Let $m \ge 2$, $j = 2, \ldots, m$ and $l_i \ge 1$ with $l_1 + \cdots + l_j = m$. We can write

$$
\mathcal{A}^m P_n = \mathcal{A}^{l_1}(P_n + Q_n)\mathcal{A}^{l_2}(P_n + Q_n)\cdots \mathcal{A}^{l_j}P_n.
$$

By expanding the formula above, we find

$$
\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n-\mathcal{A}^m P_n=-\mathcal{A}^{l_1}Q_n\mathcal{A}^{l_2}P_n\cdots\mathcal{A}^{l_j}P_n-\mathcal{A}^{l_1+l_2}Q_n\mathcal{A}^{l_3}P_n\cdots\mathcal{A}^{l_j}P_n-\cdots-\mathcal{A}^{l_1+\cdots+l_{j-1}}Q_n\mathcal{A}^{l_j}P_n.
$$
\n(212)

Using the cyclic property of the trace, we get

$$
\operatorname{Tr}\left(\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{A}^m P_n\right)=-\sum_{k=2}^j\operatorname{Tr}\left(\mathcal{A}^{l_k}P_n\cdots P_n\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\right).
$$
 (213)

Similarly, using the telescoping sum and cyclic property of the trace operator, we also get

$$
\left(\operatorname{Tr}\left(\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{A}^{m}P_n\right)\right)-\left(\operatorname{Tr}\left(\mathcal{B}^{l_1}P_n\cdots\mathcal{B}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{B}^{m}P_n\right)\right)
$$
\n
$$
=-\sum_{k=2}^{j}Tr\left(\mathcal{A}^{l_k}P_n\cdots P_n\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\right)+\operatorname{Tr}\left(\mathcal{B}^{l_k}P_n\cdots P_n\mathcal{B}^{l_j}P_n\mathcal{B}^{l_1+\cdots+l_{k-1}}Q_n\right)
$$
\n
$$
=-\sum_{k=2}^{j-1}\left(\operatorname{Tr}\left((\mathcal{A}^{l_k}-\mathcal{B}^{l_k})P_n\mathcal{A}^{l_{k+1}}\cdots P_n\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\right)\right)
$$
\n
$$
+\sum_{i=k}^{j-1}\operatorname{Tr}\left(\mathcal{B}^{l_k}P_n\cdots P_n\mathcal{B}^{l_i}P_n(\mathcal{A}^{l_{i+1}}-\mathcal{B}^{l_{i+1}})P_n\cdots\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\right)
$$
\n
$$
+\operatorname{Tr}\left(\mathcal{B}^{l_k}P_n\cdots P_n\mathcal{B}^{l_j}P_n(\mathcal{A}^{l_1+\cdots+l_{k-1}}-\mathcal{B}^{l_1+\cdots+l_{k-1}})Q_n\right),\quad(214)
$$

here we take the convention that $\sum_{j=a}^{b} \equiv 0$ for any integers *b* < *a*.

Step 1 Consider $\mathcal{A} = F$ and $\mathcal{B} = F_{n+2mn\beta}$

By Lemma [5.4](#page-37-5) we have for any $l \le m$ we have \parallel $(\mathcal{A}^l - \mathcal{B}^l) P_n \Big\|_1 \leq C_m e^{-d'n^{\beta - \frac{\alpha}{2}}}$. Note that in this case \mathcal{A} and B are symmetric. Hence we also have $||P_n(\mathcal{A}^l - \mathcal{B}^l)||_1 \leq C_m e^{-d'n^{\beta - \frac{\alpha}{2}}}$. Note that we have $||\mathcal{A}||_{\infty}$, $||\mathcal{B}||_{\infty} \leq$ \sum_{r} *cr* $Im(\eta_r)$ $\begin{array}{c} \n\end{array}$ *n*^{α} for this choice. Then use the trace norm inequality $|Tr(ABC)| \le ||A||_{\infty} ||B||_1 ||C||_{\infty}$ we get

$$
\left|\text{Tr}(\mathcal{A})^{l_1}P_n\cdots(\mathcal{A})^{l_j}P_n-\text{Tr}(\mathcal{A}^m P_n)-\text{Tr}(\mathcal{B})^{l_1}P_n\cdots(\mathcal{B})^{l_j}P_n+\text{Tr}(\mathcal{B}^m P_n)\right|\leq C''e^{-d''n^{\beta-\frac{\alpha}{2}}},\qquad(215)
$$

for some constant $C'' > 0$, $d'' > 0$. Hence plug the estimate [\(215\)](#page-42-0) into the cumulant formula [\(67\)](#page-16-0) to get

$$
\left| C_m^{(n)}(F) - C_m^{(n)}(F_{n+2mn\beta}) \right| \le C_m'' e^{-d''n^{\beta - \frac{\alpha}{2}}}.
$$
\n(216)

Step 2 Consider $\mathcal{A} = F_{n \pm 2mn\beta}$ and $\mathcal{B} = F_{n+2mn\beta}$

Note that we have $||\mathcal{A}||_{\infty}$, $||\mathcal{B}||_{\infty} \leq \sum_{r}$ *cr* $\text{Im}(\eta_r)$ $\begin{array}{c} \n\end{array}$ *n*^{α} for this choice. That is, there exists a constant $C_{op} > 0$ such that

$$
\|\mathcal{A}\|_{\infty}, \|\mathcal{B}\|_{\infty} \le n^{\alpha} C_{op}.\tag{217}
$$

We are going to estimate each summand of (214) .

Recall that $l_1 + \cdots + l_j = m$. Similar to [\(212\)](#page-41-0), we write $P_n = Id - Q_n$, use the telescopic sum, and get

$$
(\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}})P_{n}\mathcal{A}^{l_{k+1}} \cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n} = -(\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}})Q_{n}\mathcal{A}^{l_{k+1}} \cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n}
$$

$$
- (\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}})\mathcal{A}^{l_{k+1}}Q_{n}\mathcal{A}^{l_{k+2}}P_{n} \cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n}
$$

$$
- (\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}})\mathcal{A}^{l_{k+1}+l_{k+2}}Q_{n}\mathcal{A}^{l_{k+3}}P_{n} \cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n}
$$

$$
- \cdots - (\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}})\mathcal{A}^{l_{k+1}+l_{k+2}+\cdots+l_{j}}Q_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n} + (\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}})\mathcal{A}^{m-l_{k}}Q_{n}.
$$
 (218)

Use the trace norm inequality $||AB||_1 \le ||A||_1 ||B||_{\infty}$ and the operator norm inequality $||AB||_{\infty} \le ||A||_{\infty} ||B||_{\infty}$ to obtain

$$
\left| \text{Tr}\left((\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}}) P_{n} \mathcal{A}^{l_{k+1}} \cdots P_{n} \mathcal{A}^{l_{j}} P_{n} \mathcal{A}^{l_{1}+\cdots+l_{k-1}} Q_{n} \right) \right|
$$

\n
$$
\leq \left\| \left(\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}} \right) Q_{n} \right\|_{1} \left(\sum_{r} \left| \frac{c_{r}}{\text{Im}(\eta_{r})} \right| n^{\alpha} \right)^{m-l_{k}} + \sum_{i=k+1}^{j} \left\| \left(\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}} \right) \mathcal{A}^{l_{k+1}+\cdots+l_{i}} Q_{n} \right\|_{1} \left(\sum_{r} \left| \frac{c_{r}}{\text{Im}(\eta_{r})} \right| n^{\alpha} \right)^{m-l_{k}-l_{k+1}-\cdots-l_{i}} + \left\| \left(\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}} \right) \mathcal{A}^{m-l_{k}} Q_{n} \right\|_{1} . \quad (219)
$$

To apply Lemma [5.5,](#page-39-5) let us define an error term, which is exponentially small for large *n*,

$$
R_n \coloneqq 2C_m \left(\sum_r \left| \frac{c_r}{\text{Im}(\eta_r)} \right| n^{\alpha} \right)^m e^{-d'n^{\beta - \frac{\alpha}{2}}}.
$$
 (220)

Writing $\mathcal{A}^{l_k} - \mathcal{B}^{l_k} = \sum_{i=0}^{l_k-1} \mathcal{B}^{l_k-1-i}(\mathcal{A} - \mathcal{B})\mathcal{A}^i$ applying [\(203\)](#page-39-6) in Lemma [5.5,](#page-39-5) we have for any $l \in \mathbb{N}$,

$$
\left\| \left(\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}} \right) \mathcal{A}^{l} \mathcal{Q}_{n} \right\|_{1} \leq \sum_{i=0}^{l_{k}-1} \left\| \mathcal{B}^{l_{k}-1-i} (\mathcal{A} - \mathcal{B}) \mathcal{A}^{i+l} \mathcal{Q}_{n} \right\|_{1}
$$
\n
$$
\leq \sum_{i=0}^{l_{k}-1} \left(\left\| \mathcal{B}^{l_{k}-1-i} \right\|_{\infty} \left\| (\mathcal{A} - \mathcal{B}) \mathcal{Q}_{n-(i+l)n\beta} \mathcal{A}^{i+l} \mathcal{Q}_{n} \right\|_{1} + \left\| \mathcal{B}^{l_{k}-1-i} (\mathcal{A} - \mathcal{B}) \right\|_{\infty} C_{m} e^{-d'n^{\beta - \frac{\alpha}{2}}} \right)
$$
\n
$$
\leq l_{k} \left(\sum_{r} \left| \frac{c_{r}}{\text{Im}(\eta_{r})} \right| n^{\alpha} \right)^{l_{k}-1+l} \left\| (\mathcal{A} - \mathcal{B}) \mathcal{Q}_{n-mn\beta} \right\|_{1} + l_{k} R_{n}, \quad (221)
$$

where in the last inequality we use the fact that $Q_{n-(i+l)n\beta} = Q_{n-mn\beta}Q_{n-(i+l)n\beta}$ and the trace norm inequality $||AB||_1 \le ||A||_1 ||B||_{\infty}$. Then [\(219\)](#page-43-0) can be further estimated as

$$
\left|\operatorname{Tr}\left((\mathcal{A}^{l_{k}}-\mathcal{B}^{l_{k}})P_{n}\mathcal{A}^{l_{k+1}}\cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n}\right)\right| \leq (j-k)l_{k}\left(\left(\sum_{r}\left|\frac{c_{r}}{\operatorname{Im}(\eta_{r})}\right|n^{\alpha}\right)^{m-1}\left\|(\mathcal{A}-\mathcal{B})Q_{n-mn^{\beta}}\right\|_{1}+R_{n}\right).
$$
\n(222)

Similarly for $i = k, \ldots, j - 1$

$$
\left| \operatorname{Tr} \left(\mathcal{B}^{l_k} P_n \cdots P_n \mathcal{B}^{l_i} P_n \left(\mathcal{A}^{l_{i+1}} - \mathcal{B}^{l_{i+1}} \right) P_n \cdots \mathcal{A}^{l_j} P_n \mathcal{A}^{l_1 + \cdots + l_{k-1}} Q_n \right) \right|
$$

$$
\leq (j - i) l_{i+1} \left(\left(\sum_r \left| \frac{c_r}{\operatorname{Im}(\eta_r)} \right| n^{\alpha} \right)^{m-1} \left\| (\mathcal{A} - \mathcal{B}) Q_{n - m n^{\beta}} \right\|_1 + R_n \right). \tag{223}
$$

Lastly,

$$
\left| \operatorname{Tr} \left(\mathcal{B}^{l_k} P_n \cdots P_n \mathcal{B}^{l_j} P_n (\mathcal{A}^{l_1 + \cdots + l_{k-1}} - \mathcal{B}^{l_1 + \cdots + l_{k-1}}) Q_n) \right| \right|
$$

$$
\leq (l_1 + \cdots + l_{k-1}) \left(\left| \sum_r \left| \frac{c_r}{\operatorname{Im}(\eta_r)} \right| n^{\alpha} \right|^{m-1} \left| \left(\mathcal{A} - \mathcal{B} \right) Q_{n - m n^{\beta}} \right| \right|_1 + R_n \right). \tag{224}
$$

Hence,

$$
\left| \text{Tr}(\mathcal{A})^{l_1} P_n \cdots (\mathcal{A})^{l_j} P_n - \text{Tr}(\mathcal{A}^m P_n) - \text{Tr}(\mathcal{B})^{l_1} P_n \cdots (\mathcal{B})^{l_j} P_n + \text{Tr}(\mathcal{B}^m P_n) \right|
$$

$$
\leq m j^2 \left(\left(\sum_r \left| \frac{c_r}{\text{Im}(\eta_r)} \right| n^{\alpha} \right)^{m-1} \left| \left(\mathcal{A} - \mathcal{B} \right) Q_{n-mn^{\beta}} \right| \left| 1 + R_n \right. \right). \tag{225}
$$

Then by [\(204\)](#page-39-7), we have

$$
\left|\text{Tr}(\mathcal{A})^{l_1}P_n\cdots(\mathcal{A})^{l_j}P_n-\text{Tr}(\mathcal{A}^m P_n)-\text{Tr}(\mathcal{B})^{l_1}P_n\cdots(\mathcal{B})^{l_j}P_n+\text{Tr}(\mathcal{B}^m P_n)\right|\leq C''e^{-d''n^{\beta-\frac{\alpha}{2}}},\qquad(226)
$$

for some constant $C'' > 0$, $d'' > 0$. Hence plug the estimate [\(226\)](#page-44-0) into the definition [\(67\)](#page-16-0) and we have

$$
\left| C_m^{(n)}(F_{n+2mn\beta}) - C_m^{(n)}(F_{n+2mn\beta}) \right| \le C_m'' e^{-d''n^{\beta - \frac{\alpha}{2}}}.
$$
\n(227)

Recall that by definition of *F*, we have $n^{\alpha m} C_m(X_{f, \alpha}^{(n)})$ *f*, a _{*f*, a , x ₀) = $C_m(F)$. Combining [\(216\)](#page-42-2) from Step 1 and [\(227\)](#page-44-1)} from Step 2, we conclude [\(111\)](#page-25-1) in Proposition [4.2.](#page-25-0) \Box

6 Proof of Propositions [4.3](#page-26-0) and [4.4](#page-27-0)

In this section, we prove Propositions [4.3](#page-26-0) and [4.4.](#page-27-0) Essentially, we are studying the inverse of the blocked operator $J_{n\pm 2mn\beta}^{(r)}$ defined in [\(105\)](#page-23-1), with $\beta = \frac{\alpha}{2} + \frac{\varepsilon}{3}$. It is the middle block, $P_{n+2mn\beta}Q_{n-2mn\beta}J^{(r)}Q_{n-2mn\beta}P_{n+2mn\beta}$, that is non-trivial and matters in the analysis.

In Section [6.1,](#page-44-2) we will compute the resolvent for a general $N \times N$ matrix with $N \in \mathbb{N}$, (cf.Proposition [6.1\)](#page-44-3). Later, in Sections [6.2](#page-49-0) and [6.3,](#page-50-0) we will apply Propositio[n6.1](#page-44-3) to the middle block the resolvent of $J_{n}^{(r)}$ $\lim_{n\pm 2mn\beta}$, by taking the size of the matrix to be $4mn^{\beta}$ with relabelling of the indexes.

6.1 Inverse of a Tri-diagonal Matrix: Part II

Consider an $N \times N$ non singular symmetric matrix with $a_j \in \mathbb{R}, b_j \in \mathbb{R}, N \in \mathbb{N}$ and $z \in \mathbb{C}$ with $\text{Im } z \neq 0$.

$$
J_N = \begin{pmatrix} b_0 & a_1 & 0 & 0 & 0 & , \cdots, & 0 \\ a_1 & b_1 & a_2 & 0 & 0 & , \cdots, & 0 \\ 0 & a_2 & b_2 & a_3 & 0 & , \cdots, & 0 \\ \cdots & & & & & & \\ 0 & 0 & 0 & 0 & a_{N-2} & b_{N-2} & a_{N-1} \\ 0 & 0 & 0 & 0 & 0 & a_{N-1} & b_{N-1} \end{pmatrix} - zId
$$
(228)

The goal of this subsection is to continue to develop a theory to estimate the inverse of J_N entry-wise by imposing stronger assumptions than in Section [5.](#page-30-4)

Proposition 6.1 (Almost Toeplitz). *Consider an N* × *N non singular symmetric matrix J_N with* $a_j \in \mathbb{R}, b_j$ *–* $\text{Re } z > 0$ and $N \in \mathbb{N}$ as defined as [\(228\)](#page-44-4). Let ω_j^+, ω_j^- as defined as [\(96\)](#page-21-0). Let M_j as defined as [\(97\)](#page-22-0). Define

$$
c_0 = \min_j |a_j|, \quad c_1 = \max \left\{ \max_j |a_j|, \max_j |b_{j-1} - z| \right\},\tag{229}
$$

$$
c_2 = \max\left\{1, \left|\frac{-a_1 + \omega_2^-(b_0 - z)}{a_1 - \omega_2^+(b_0 - z)}\right|, \left|\frac{-\omega_{N-1}^+ a_{N-1} + b_{N-1} - z}{\omega_{N-1}^- a_{N-1} - b_{N-1} + z}\right|\right\},\tag{230}
$$

$$
\varepsilon_1 = N \max_j ||M_j||_{\infty},\tag{231}
$$

$$
\varepsilon_2 = \prod_{l=2}^{N-1} \left| \frac{\omega_l^-}{\omega_l^+} \right|.
$$
\n(232)

*Assume that c*₀ > 0*, and both* ε_1 ε_1 *,* ε_2 *are sufficiently small.* ¹ *Then we have*

$$
J_N^{-1} = T_N(\eta) + H_N(\eta),
$$
\n(233)

such that for all $j \leq k$

$$
\left| (T_N(\eta))_{j,k} \right| \leq Constant \left| \frac{\prod_{l=j}^{k-1} \omega_l^-}{a_{k-1} \omega_j^- \left(\omega_{N-1}^- - \omega_{N-1}^+ \right)} \right|,\tag{234}
$$

$$
\left| (H_N(\eta))_{j,k} \right| \leq Constant \left| \frac{\prod_{l=j}^{k-1} \omega_l^{-} \left(\prod_{l=2}^j \frac{\omega_l^{-}}{\omega_l^{+}} + \prod_{l=k}^{N-1} \frac{\omega_l^{-}}{\omega_l^{+}} \right)}{a_{k-1} \omega_j^{-} \left(\omega_{N-1}^{-} - \omega_{N-1}^{+} \right)} \right|,
$$
\n(235)

where Constant is given by [\(267\)](#page-48-0) *that only depends on* c_0 , c_1 , c_2 , ε_1 *and* ε_2 *and remains bounded as* ε_1 , ε_2 \rightarrow 0*.*

Further we have for any j, k, l, as ε_1 , $\varepsilon_2 \rightarrow 0$,

$$
(T_N(\eta))_{j,k} = \frac{(-1)^{k-j}\omega_{N-1}^-}{\omega_{N-1}^+ - \omega_{N-1}^-} \frac{\prod_{m=j}^{k-1} \omega_m^-}{a_{k-1}\omega_j^-} (1 + O(\varepsilon_1 + \varepsilon_2))
$$
(236)

and

$$
\frac{(T_N(\eta))_{j,k}}{(T_N(\eta))_{j+l,k+l}} = \left(\prod_{m=j}^{k-1} \frac{\omega_m^-}{\omega_{m+l}^-}\right) \frac{\omega_{l+j}^- a_{k+l-1}}{\omega_j^- a_{k-1}} \left(1 + O(\varepsilon_1)\right). \tag{237}
$$

By convention we take $\prod_{l=k}^{k-1}$ \equiv 1. Note that J_N is symmetric and so is J_N^{-1} . Therefore we have the approxi*mation of each entry of the inverse.*

Proof. Applying the inverse formula for a tri-diagonal matrix [\(99\)](#page-22-1), we have for $j \leq k$

$$
(J_N^{-1})_{j,k} = \frac{(-1)^{k-j} \gamma_j \beta_k}{\beta_N a_N \gamma_{N+1}}.
$$
\n(238)

Choose any $a_N \neq 0$ (e.g. pick $a_N = a_{N-1}$) and Note that the inverse formula is independent on the choice of *a*_{*N*}. Recall that in the proof of Proposition [5.1](#page-30-0) we have computed β_k to be [\(152\)](#page-32-2), i.e., for $k = 1, ..., N$

$$
\beta_k = \frac{\tilde{\beta}_1 \prod_{l=k}^{N-1} \omega_l^+}{\omega_{N-1}^- - \omega_{N-1}^+} \left(1 + \frac{\tilde{\beta}_2}{\tilde{\beta}_1} \prod_{l=k}^{N-1} \frac{\omega_l^-}{\omega_l^+} + C(k) \right)
$$
(239)

¹More precisely, we assume $0 < \varepsilon_1 < 3^{-1} \left(\frac{(1+\sqrt{5})c_1}{2c_0} \right)$ \int_{0}^{2} , 0 < ε_2 and 12 $\left(1 + \frac{(1+\sqrt{5})c_1}{2c_0}\right)$ $\int_0^2 c_2^2$ $\left\{\varepsilon_1 + c_2 \varepsilon_2 < \left(\frac{(1+\sqrt{5})c_1}{2c_0}\right)\right\}$ \int_{0}^{-2} . where *C*(*k*) is defined in [\(151\)](#page-31-0) with $|C(k)| \le 4(1 + c_2)\varepsilon_1$ and

$$
\tilde{\beta}_1 := \omega_{N-1}^- a_{N-1} - b_{N-1} + z, \qquad \tilde{\beta}_2 := -\omega_{N-1}^+ a_{N-1} + b_{N-1} - z. \tag{240}
$$

Note that we have $\beta_N = a_{N-1}$, $\beta_N = a_{N-1}$ $\frac{\tilde{B}_2}{\tilde{B}_1} \prod_{l=k}^{N-1} \frac{\omega_l^-}{\omega_l^+}$ $\overline{\omega}_l^+$ $\leq c_2 \varepsilon_2.$

Recall that transfer matrices B_j is defined as [\(98\)](#page-22-2), and γ_j is defined recursively to be

$$
\begin{pmatrix} \gamma_j \\ \gamma_{j+1} \end{pmatrix} = B_j B_{j-1} \dots B_2 \begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix}, \quad \gamma_1 = a_1, \quad \gamma_2 = b_0 - z. \tag{241}
$$

Recall that the eigenvalues λ_j^+ and $\lambda_j^ \frac{1}{j}$ of *B*_{*j*} are defined as [\(96\)](#page-21-0). *B*_{*j*} can be diagonalized as *B*_{*j*} = *W*_{*j*}</sub> $\Lambda_j W_j^{-1}$ where, $W_j := \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ λ $\frac{1}{j}$ λ − *j* $\left(\begin{array}{c} \lambda \\ \lambda \end{array} \right)$ and $\Lambda_j \coloneqq \left(\begin{array}{c} \lambda \\ \lambda \end{array} \right)$ + *j* 0 $\dot{0}$ $\lambda_j^$ *j* . Further, define

$$
D_j := W_j^{-1} \left(B_j B_{j-1} \dots B_2 - W_j \left(\prod_{l=2}^j \Lambda_l \right) W_2^{-1} \right) W_2, \quad j = 2, 3, \dots, N, \text{ and } D_1 := 0. \tag{242}
$$

We can rewrite [\(241\)](#page-46-0) for all $j = 2, 3, \ldots, N$ to be

$$
\begin{pmatrix} \gamma_j \\ \gamma_{j+1} \end{pmatrix} = W_j \left(D_j + \prod_{l=2}^j \Lambda_l \right) W_2^{-1} \begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix} . \tag{243}
$$

Let $\tilde{\gamma}_1 \coloneqq \lambda_2^ \frac{1}{2}a_1-b_0+z, \tilde{\gamma}_2 \coloneqq -\lambda_2^{\dagger}a_1+b_0-z$ such that $\frac{1}{\lambda_2^{\dagger}-1}$ $\lambda_2^- - \lambda_2^+$ $\begin{pmatrix} \tilde{\gamma}_1 \\ \tilde{\gamma}_2 \end{pmatrix}$ $\tilde{\gamma}_2$ $= W_2^{-1}$ $\begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix}$ γ_2 ! . Then, we have for $j = 2, ..., N$

$$
\gamma_j = \frac{\left((D_j)_{1,1} + (D_j)_{2,1} + \prod_{l=2}^j \lambda_l^+ \right) \tilde{\gamma}_1 + \left((D_j)_{1,2} + (D_j)_{2,2} + \prod_{l=2}^j \lambda_l^- \right) \tilde{\gamma}_2}{\lambda_2^- - \lambda_2^+}.
$$
\n(244)

By convention, let $D_1 \equiv 0$ and $\prod_{l=2}^1 \equiv 1$ and we can allow $j = 1, ..., N$ for the formula γ_j above.

For γ_{N+1} , we keep B_N unchanged, estimate from B_{N-1} and write

$$
\begin{pmatrix} \gamma_N \\ \gamma_{N+1} \end{pmatrix} = B_N W_{N-1} \left(D_{N-1} + \prod_{l=2}^{N-1} \Lambda_l \right) W_2^{-1} \begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix} . \tag{245}
$$

Note that

$$
B_N W_{N-1} = \begin{pmatrix} \lambda_{N-1}^+ & \lambda_{N-1}^- \\ \frac{(b_{N-1}-z)\lambda_{N-1}^+ - a_{N-1}}{a_N} & \frac{(b_{N-1}-z)\lambda_{N-1}^- - a_{N-1}}{a_N} \end{pmatrix} .
$$
 (246)

Denote

$$
\tilde{\delta}_1 := (b_{N-1} - z) \lambda_{N-1}^+ - a_{N-1}, \qquad \tilde{\delta}_2 := (b_{N-1} - z) \lambda_{N-1}^- - a_{N-1}.
$$
\n(247)

Compute the second entry of [\(245\)](#page-46-1) to obtain

$$
\gamma_{N+1} = \frac{\tilde{\delta}_1((D_{N-1})_{1,1} + (D_{N-1})_{2,1} + \prod_{l=2}^{N-1} \lambda_l^+)\tilde{\gamma}_1 + \tilde{\delta}_2((D_{N-1})_{1,2} + (D_{N-1})_{2,2} + \prod_{l=2}^{N-1} \lambda_l^-)\tilde{\gamma}_2}{a_N(\lambda_2^- - \lambda_2^+)}.
$$
 (248)

By convention we let $D_1 \equiv 0$ and $\prod_{l=2}^1 \equiv 1$ and we can allow $j = 1, ..., N$ for all above. Moreover, by the construction of E_j in [\(101\)](#page-22-3), we have $E_j \equiv W_j^{-1}W_{j-1} - Id$. Now we estimate D_j via the expansion

$$
B_j B_{j-1} \dots B_2 = W_j \Lambda_j (Id + E_j) \Lambda_{j-1} (Id + E_{j-1}) \dots \Lambda_3 (Id + E_3) \Lambda_2 W_2^{-1}.
$$
 (249)

Also, note that by assumption $b_j > 0$, we have $|\lambda_j^+| > |\lambda_j^-|$ $j[−]$ |. Hence, $||W_j||_{\infty} \le |\lambda_j^+|$. Then, we have

$$
||D_j||_{\infty} \le \left(\prod_{l=2}^{j} (1 + ||E_l||_{\infty}) - 1\right) \prod_{l=2}^{j} |\lambda_l^+|.
$$
 (250)

Now we are about to estimate E_j . We will do so by revealing some connections between λ_j^{\pm} $\frac{1}{j}$ and ω_j^{\pm} *j* and between E_j and M_j .

By [\(229\)](#page-44-5),

$$
|\omega_j^+| \le \frac{|b_{j-1} - z| + (|b_{j-1} - z|^2 + 4|a_{j-1}a_j|)^{\frac{1}{2}}}{2|a_{j-1}|} \le \frac{(1 + \sqrt{5})c_1}{2c_0},
$$
\n(251)

$$
|\omega_j^-| \ge \frac{2|a_{j|}}{|b_{j-1} - z| + (|b_{j-1} - z|^2 + 4|a_{j-1}a_j|)^{\frac{1}{2}}} \ge \frac{2c_0}{(1 + \sqrt{5})c_1}.
$$
 (252)

Note that each entry of the 2 by 2 matrix E_j is bounded by its operator norm. Since λ_j^2 $\overline{j}\omega_j^+ = \lambda_j^+ \omega_j^+$ − $\frac{1}{j} = 1$, by [\(251\)](#page-47-0) and [\(252\)](#page-47-1), we have each entry of E_j is bounded by $\left(\frac{(1+\sqrt{5})c_1}{2c_0}\right)$ 2*c*⁰ $\int_0^2 ||M_j||_{\infty}$. Hence, $||E_j||_{\infty}$ ≤ $3\left(\frac{(1+\sqrt{5})c_1}{2c_2}\right)$ 2*c*⁰ $\int_0^2 ||M_j||_{\infty}$. By [\(231\)](#page-45-1) and 0 < 3 $\left(\frac{(1+\sqrt{5})c_1}{2c_0}\right)$ 2*c*⁰ χ^2 ε_1 < 1, by the same engagement as in [\(150\)](#page-31-1), we further estimate [\(250\)](#page-47-2) to be √

$$
||D_j||_{\infty} \le 6 \left(\frac{(1 + \sqrt{5})c_1}{2c_0} \right)^2 \varepsilon_1 \prod_{l=2}^j |\lambda_l^+|.
$$
 (253)

Then, let

$$
D(j) := ((D_j)_{1,1} + (D_j)_{2,1}) \prod_{l=2}^{j} (\lambda_l^+)^{-1} + ((D_j)_{1,2} + (D_j)_{2,2}) \frac{\tilde{\gamma}_2}{\tilde{\gamma}_1} \prod_{l=2}^{j} (\lambda_l^+)^{-1},
$$
\n(254)

$$
\tilde{D}(N-1) := ((D_{N-1})_{1,1} + (D_{N-1})_{2,1}) \prod_{l=2}^{N} (\lambda_l^+)^{-1} + \frac{\tilde{\delta}_2}{\tilde{\delta}_1} ((D_{N-1})_{1,2} + (D_{N-1})_{2,2}) \frac{\tilde{\gamma}_2}{\tilde{\gamma}_1} \prod_{l=2}^{N-1} (\lambda_l^+)^{-1}.
$$
 (255)

Hence, by[\(253\)](#page-47-3)

$$
|D(j)| \le 12\left(1 + \left|\frac{\tilde{\gamma}_2}{\tilde{\gamma}_1}\right|\right) \left(\frac{(1+\sqrt{5})c_1}{2c_0}\right)^2 \varepsilon_1, \quad |\tilde{D}(N-1)| \le 12\left(1 + \left|\frac{\tilde{\delta}_2\tilde{\gamma}_2}{\tilde{\delta}_1\tilde{\gamma}_1}\right|\right) \left(\frac{(1+\sqrt{5})c_1}{2c_0}\right)^2 \varepsilon_1.
$$
 (256)

Note that $\lambda_j^ \overline{j} \omega_j^+ = \lambda_j^+ \omega_j^+$ − $\frac{1}{\delta}$ = 1 and we have $\frac{\delta_2}{\delta_1} = \frac{(b_{N-1}-z)\lambda_{N-1}^2 - a_{N-1}}{(b_{N-1}-z)\lambda_{N-1}^2 - a_{N-1}}$ $\frac{(b_{N-1}-z)\lambda_{N-1}^- - a_{N-1}}{(b_{N-1}-z)\lambda_{N-1}^+ - a_{N-1}} = \frac{\omega_{N-1}^-}{\omega_{N-1}^+}$ $\frac{\omega_{N-1}^{-}}{\omega_{N-1}^{+}} \frac{\omega_{N-1}^{+} a_{N-1}-b_{N-1}+z}{\omega_{N-1}^{-} a_{N-1}-b_{N-1}+z}$ $\frac{\omega_{N-1}^T a_{N-1} - b_{N-1} + z}{\omega_{N-1}^- a_{N-1} - b_{N-1} + z}$ and $\frac{\tilde{\gamma}_2}{\tilde{\gamma}_1}$ $\frac{\tilde{\gamma}_2}{\tilde{\gamma}_1} = \frac{-\lambda_2^+ a_1 + b_0 - z}{\lambda_2^- a_1 - b_0 + z}$ $\frac{\lambda_2 a_1 + b_0 - z}{\lambda_2^2 a_1 - b_0 + z} =$ $\frac{\omega_2^+}{\omega_-^-}$ $\frac{\omega_2^+}{\omega_2^-} \frac{-a_1 + \omega_2^-(b_0 - z)}{a_1 - \omega_2^+(b_0 - z)}$ $\frac{a_1 + a_2 + b_0 - z_1}{a_1 - a_2 + b_0 - z_1}$. Then by [\(251\)](#page-47-0) [\(252\)](#page-47-1), we estimate,

$$
\left|\frac{\tilde{\delta}_2}{\tilde{\delta}_1}\right| \le c_2, \quad \left|\frac{\tilde{\gamma}_2}{\tilde{\gamma}_1}\right| \le \left(\frac{(1+\sqrt{5})c_1}{2c_0}\right)^2 c_2.
$$
\n(257)

Recall that $c_2 \geq 1$. Further define

$$
\tilde{\varepsilon}_1 \coloneqq 12 \left(1 + \left(\frac{(1 + \sqrt{5})c_1}{2c_0} \right)^2 c_2^2 \right) \left(\frac{(1 + \sqrt{5})c_1}{2c_0} \right)^2 \varepsilon_1. \tag{258}
$$

Then by [\(253\)](#page-47-3) we have $|C(k)|$, $|D(j)|$, $|\tilde{D}(N-1)| < \tilde{\varepsilon}_1$ for all *j*, *k*. Also, note that by definition we have the following handy relations

$$
\lambda_j^- = (\omega_j^+)^{-1}, \qquad \lambda_j^+ = (\omega_j^-)^{-1}, \qquad \omega_j^+ \omega_j^- = \frac{a_j}{a_{j-1}}, \qquad \frac{\tilde{\beta}_1}{\tilde{\delta}_1} = -\omega_{N-1}^-.
$$
 (259)

Hence for $j \leq k$

$$
(J_N^{-1})_{j,k} = \frac{(-1)^{k-j}\beta_k \gamma_j}{\beta_N a_N \gamma_{N+1}}
$$
\n
$$
= \frac{(-1)^{k-j}\beta_k}{a_{N-1}a_N} \frac{\left((D_j)_{1,1} + (D_j)_{2,1} + \prod_{l=2}^j \lambda_l^+\right)\tilde{\gamma}_1 + \left((D_j)_{1,2} + (D_j)_{2,2} + \prod_{l=2}^j \lambda_l^-\right)\tilde{\gamma}_2}{\delta_1 \left((D_{N-1})_{1,1} + (D_{N-1})_{2,1} + \prod_{l=2}^N \lambda_l^+\right)\tilde{\gamma}_1 + \tilde{\delta}_1 \left((D_{N-1})_{1,2} + (D_{N-1})_{2,2} + \prod_{l=2}^N \lambda_l^-\right)\tilde{\gamma}_2}
$$
\n(261)

$$
=\frac{(-1)^{k-j}\omega_{N-1}^{-}}{a_{k-1}}\frac{\prod_{l=j}^{k-1}\omega_{l}^{-}}{(\omega_{N-1}^{+}-\omega_{N-1}^{-})\omega_{j}^{-}}\frac{\left(1+\frac{\tilde{\gamma}_{2}}{\tilde{\gamma}_{1}}\prod_{l=2}^{j}\frac{\omega_{l}^{-}}{\omega_{l}^{+}}+D(j)\right)\left(1+\frac{\tilde{\beta}_{2}}{\tilde{\beta}_{1}}\prod_{l=k}^{N-1}\frac{\omega_{l}^{-}}{\omega_{l}^{+}}+C(k)\right)}{\left(1+\frac{\tilde{\gamma}_{2}}{\tilde{\gamma}_{1}}\prod_{l=2}^{N-1}\frac{\omega_{l}^{-}}{\omega_{l}^{+}}+\tilde{D}(N-1)\right)}.
$$
(262)

Let

$$
(T_N(\eta))_{j,k} := \frac{(-1)^{k-j}\omega_{N-1}^-}{a_{k-1}} \frac{\prod_{l=j}^{k-1} \omega_l^-}{\left(\omega_{N-1}^+ - \omega_{N-1}^-\right)\omega_j^-} \frac{(1+D(j))(1+C(k))}{\left(1+\frac{\tilde{\gamma}_2}{\tilde{\gamma}_1}\prod_{l=2}^{N-1} \frac{\omega_l^-}{\omega_l^+}+\tilde{D}(N-1)\right)},\tag{263}
$$

$$
H_N(\eta) := J_N^{-1} - T_N(\eta). \tag{264}
$$

Hence as $\varepsilon_1 \to 0$ we have

$$
\frac{(T_N(\eta))_{j,k}}{(T_N(\eta))_{j+l,k+l}} = \left(\prod_{m=j}^{k-1} \frac{\omega_m^-}{\omega_{m+l}^-}\right) \frac{\omega_{l+j}^- a_{k+l-1}}{\omega_j^- a_{k-1}} \frac{(1+D(j))(1+C(k))}{(1+D(j+l))(1+C(k+l))}
$$
(265)

$$
= \left(\prod_{m=j}^{k-1} \frac{\omega_m^-}{\omega_{m+l}^-} \right) \frac{\omega_{l+j}^- a_{k+l-1}}{\omega_j^- a_{k-1}} \left(1 + O(\varepsilon_1)\right). \tag{266}
$$

Hence, whenever $\tilde{\varepsilon}_1 + \left(\frac{(1+\sqrt{5})c_1}{2c_0}\right)$ 2*c*⁰ $\int_0^2 c_2 \varepsilon_2 < 1$

$$
\left| \frac{\omega_{N-1}^{-}(1+D(j))(1+C(k))}{\left(1+\frac{\tilde{\gamma}_{2}}{\tilde{\gamma}_{1}}\prod_{l=2}^{N-1}\frac{\omega_{l}^{-}}{\omega_{l}^{+}}+\tilde{D}(N-1)\right)}\right| \leq \frac{\frac{(1+\sqrt{5})c_{1}}{2c_{0}}(1+\tilde{\varepsilon}_{1})^{2}}{1-\left(\frac{(1+\sqrt{5})c_{1}}{2c_{0}}\right)^{2}c_{2}\varepsilon_{2}-\tilde{\varepsilon}_{1}} =:Constant'.\tag{267}
$$

Hence,

$$
\left| (T_N(\eta))_{j,k} \right| \leq Constant' \left| \frac{\prod_{l=j}^{k-1} \omega_l^-}{a_{k-1} \omega_j^- \left(\omega_{N-1}^- - \omega_{N-1}^+ \right)} \right|,
$$
\n(268)

$$
\left| (H_N(\eta))_{j,k} \right| \le 2Constant' \left| \frac{\left(\prod_{l=j}^{k-1} \omega_l^{-} \right) \left(\prod_{l=2}^j \frac{\omega_l^{-}}{\omega_l^{+}} + \prod_{l=k}^{N-1} \frac{\omega_l^{-}}{\omega_l^{+}} \right)}{a_{k-1} \left(\omega_{N-1}^{-} - \omega_{N-1}^{+} \right)} \right|.
$$
 (269)

Take the *Constant* = $2Constant'$ and we conclude the proof of Proposition [6.1.](#page-44-3)

6.2 Proof of Proposition [4.3](#page-26-0)

Proof of Proposition [4.3](#page-26-0). Recall that we define $J_{n+1}^{(r)}$ $n_{n\pm 2mn\beta}^{(r)}$ as [\(105\)](#page-23-1), which can be regarded as a three-block diagonal matrix. Recall we define the indexing set $I_{n,m}^{(\beta)} = \left\{ n - 2mn^{\beta}, n - 2mn^{\beta} + 1, \dots, n + 2mn^{\beta} \right\}$. Hence, the non-trivial block of $J_{n}^{(r)}$ $\lim_{n\pm 2mn\beta}$ is where the entries have the indices both belonging to this set $I_{n,m}^{(\beta)}$, i.e., for all $j, k \in I_{n,n}^{(\beta)}$ *ⁿ*,*^m*

$$
\left(J_{n\pm 2mn\beta}^{(r)}\right)_{j,k} = \left(P_{n+2mn\beta}Q_{n-2mn\beta}J^{(r)}Q_{n-2mn\beta}P_{n+2mn\beta}\right)_{j,k}.
$$
 (270)

In the following, we will apply Proposition [6.1](#page-44-3) to this non-trivial block of $J_{n+1}^{(r)}$ $\lim_{n\pm 2mn\beta}$, i.e., [\(270\)](#page-49-1). To short the notation, let

$$
b_j = b_{j,n} - x_0
$$
, $a_j = a_{j,n}$, and $z = \frac{\eta}{n^{\alpha}}$. (271)

Then, at the left edge, for all $j \in I_{n,m}^{(\beta)}$, a_j and b_j satisfy Conditions [5.1](#page-32-0) and [5.2.](#page-33-3) Hence, the quantities in [\(229\)](#page-44-5) are bounded below and above as desired are bounded below and above as desired.

By Lemma [5.1,](#page-33-0) we have $b_l > 0$, for all $l \in I_{n,m}^{(\beta)}$. By Lemma [5.2,](#page-34-0) we have as $n \to \infty$

$$
\max_{l \in I_{n,m}^{(\beta)}} |\omega_l^- - 1| = O(n^{-\frac{\alpha}{2}}), \quad \max_{l \in I_{n,m}^{(\beta)}} |\omega_l^+ - 1| = O(n^{-\frac{\alpha}{2}})
$$
\n(272)

Hence, as $n \to \infty$

$$
\left|\frac{-a_{n-2mn\beta}+\omega_{n-2mn\beta+1}^{-}(b_{n-2mn\beta-1}-z)}{a_{n-2mn\beta}-\omega_{n-2mn\beta+1}^{+}(b_{n-2mn\beta-1}-z)}\right|=1+O(n^{-\frac{\alpha}{2}}),\quad \left|\frac{-\omega_{n+2mn\beta-1}^{+}a_{n+2mn\beta-1}+b_{n+2mn\beta-1}-z}{\omega_{n+2mn\beta-1}^{-}a_{n+2mn\beta-1}-b_{n+2mn\beta-1}+z}\right|=1+O(n^{-\frac{\alpha}{2}}).
$$

This shows $c_2 = 1 + O(n^{\alpha/2})$ in [\(230\)](#page-45-2).

Let *M*^{*j*} be defined as [\(97\)](#page-22-0). By Lemma [5.3](#page-35-1) we have as $n \to \infty$

$$
4mn^{\beta} \max_{l \in I_{n,m}^{(\beta)}} ||M_l||_{\infty} = o(n^{-2\beta + \alpha}).
$$
\n(273)

This shows $\varepsilon_1 = o(n^{-2\beta + \alpha})$ in [\(231\)](#page-45-1). Moreover, use Lemma [5.2](#page-34-0) again to obtain that there exists a constant $d > 0$ such that for all $n > n_0$ $d > 0$ such that for all $n > n_0$

$$
\prod_{l=n-2mn^{\beta}+2}^{n+2mn^{\beta}-1} \left| \frac{\omega_l^-}{\omega_l^+} \right| \le \left(1 - \frac{d}{n^{\frac{\alpha}{2}}}\right)^{4mn^{\beta}-2} \le e^{-dn^{-\frac{\alpha}{2}}(4mn^{\beta}-2)}.
$$
\n(274)

This shows $\varepsilon_2 = O(e^{-d'n^{\beta - \frac{\alpha}{2}}})$ in [\(141\)](#page-30-3), which is exponentially small.
Hence the assumption of Proposition 6.1 is verified for the mid-

Hence, the assumption of Proposition [6.1](#page-44-3) is verified for the middle block of $J_{n+1}^{(r)}$ $\lim_{n \pm 2mn\beta}$. Hence, we obtain the estimate of $T_{n\pm 2mn\beta}(\eta_r)$ to be [\(236\)](#page-45-3). Further, by $a_j = a_n + O(n^{\beta-1})$, (236) and Lemma [5.2,](#page-34-0) we have as $n \to \infty$

$$
(T_{n\pm 2mn\beta}(\eta_r))_{j,k} = \frac{(-1)^{k-j}}{2a_n\left(\frac{-\eta_r}{n^{\alpha}|a_n|}\right)^{\frac{1}{2}}} \prod_{l=j}^{k-1} \left(1 - \left(\frac{-\eta_r}{n^{\alpha}|a_n|}\right)^{\frac{1}{2}} + \xi_l\right) (1 + o(1)),\tag{275}
$$

where $\xi_l \coloneqq \omega_l^ \frac{1}{l} - \left(1 - \left(\frac{-\eta}{n^{\alpha}|a_{N-1}|}\right)^{\frac{1}{2}}\right)$ $sgn(a_{N-1}) = o\left(n^{-\frac{\alpha}{2}}\right).$

To prove [\(123\)](#page-26-1). Note that since each entry of M_l is bounded by the operator norm, we have for all $l = n - 2mn^{\beta}, \ldots, n + 2mn^{\beta},$

$$
|\omega_l^- - \omega_{l+1}^-| \leq ||M_l||_{\infty} |\omega_l^- - \omega_l^+|.
$$

By Lemma [5.3](#page-35-1) we have $||M_l||_{\infty} = o(n^{-3\beta + \alpha})$, given that $|\omega_l^ \frac{1}{l} - \omega_l^+$ is of order $n^{-\frac{\alpha}{2}}$. Hence we have

$$
\max_{l \in I_{n,m}^{(\beta)}} |\omega_l^- - \omega_{l+1}^-| = o(n^{-3\beta + \frac{\alpha}{2}}), \quad \text{as } n \to \infty.
$$
 (276)

Lemma [5.2](#page-34-0) says that ω_m^- is bounded away from zero, hence we estimate [\(237\)](#page-45-4) to be

$$
\frac{(T_{n\pm 2mn\beta}(\eta_r))_{j,k}}{(T_{n\pm 2mn\beta}(\eta_r))_{j+l,k+l}} = \left(1 + o(n^{-3\beta + \frac{\alpha}{2}})\right)^{(k-j)l} \left(1 + O(n^{\beta - 1})\right) (1 + O(\varepsilon_1)), \quad \text{as } n \to \infty. \tag{277}
$$

Note that $1 + O(n^{\beta-1}) = \frac{a_{k+l-1}}{a_{k-1}}$ *a_{k−1}* comes from Condition [5.1.](#page-32-0) Also note that $\varepsilon_1 = o(n^{-2\beta+\alpha})$. Moreover, $(k - j)l = O(n^{2\beta})$, since $1 \le k - j, l \le 4mn^{\beta}$. Note that all the of the big-*Os* and small-*o* are uniform in the indices by assumptions. Together with the assumption $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{2} < 1$, we have indices by assumptions. Together with the assumption $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$, we have

$$
\frac{(T_{n\pm 2mn^{\beta}}(\eta_r))_{j,k}}{(T_{n\pm 2mn^{\beta}}(\eta_r))_{j+l,k+l}} = 1 + o(n^{-\beta + \frac{\alpha}{2}}), \quad \text{as } n \to \infty.
$$

This shows the statement [\(123\)](#page-26-1).

Note that above argument is taken at the left edge i.e., $b_j > 0$ as assumed in [\(162\)](#page-33-2). For the right edge, we have *b*_{*j*} < 0. Then we consider $-J_{n\pm 2mn\beta}$ whose diagonals are $-b_j + \frac{\eta_r}{n^2}$ and off diagonals are $-a_j$. Apply the result above by replacing a_n by $-a_n$ and η_r by $-\eta_r$ to estimate entries of $\left(-J_{n\pm}^{(r)}\right)$ *n*±2*mn*β \int^{-1} as in the statement. Then the result at the right edge is obtained by the simple fact $(J_{n+1}^{(r)})$ *n*±2*mn*β $\Big)^{-1} = -\Bigl(-J_{n+1}^{(r)}\Bigr)$ *n*±2*mn*β $\big)^{-1}$. This completes the proof. \Box

6.3 Proof of Proposition [4.4](#page-27-0)

Now we are ready to prove the Proposition [4.4](#page-27-0) by Lemmas [5.4](#page-37-5) and [5.5](#page-39-5) and Proposition [4.3.](#page-26-0)

Proof of Proposition [4.4.](#page-27-0) Recall that for any linear operator \mathcal{A} we write by [\(67\)](#page-16-0)

$$
C_m^{(n)}(\mathcal{A})=m!\sum_{j=2}^m\frac{(-1)^{j+1}}{j}\sum_{l_1+\cdots+l_j=m,l_i\geq 1}\frac{\mathrm{Tr}(\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n)-\mathrm{Tr}(\mathcal{A}^mP_n)}{l_1!\cdots l_j!}.
$$

Hence, consider $m \ge 2$, $j = 2, ..., m$ and $l_i \ge 1$ with $l_1 + \cdots + l_j = m$. We can write $P_n = -Q_n + Id$ and

$$
\mathcal{A}^{l_1} P_n \cdots \mathcal{A}^{l_j} P_n = -\mathcal{A}^{l_1} Q_n \cdots \mathcal{A}^{l_j} P_n + \mathcal{A}^{l_1+l_2} P_n \cdots \mathcal{A}^{l_j} P_n
$$

=
$$
-\mathcal{A}^{l_1} Q_n \cdots \mathcal{A}^{l_j} P_n - \mathcal{A}^{l_1+l_2} Q_n \cdots \mathcal{A}^{l_j} P_n + \mathcal{A}^{l_1+l_2+l_3} P_n \cdots \mathcal{A}^{l_j} P_n
$$
 (278)

Continue this argument *j* times and we get

$$
\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n-\mathcal{A}^m P_n=-\mathcal{A}^{l_1}Q_n\mathcal{A}^{l_2}P_n\cdots\mathcal{A}^{l_j}P_n-\mathcal{A}^{l_1+l_2}Q_n\mathcal{A}^{l_3}P_n\cdots\mathcal{A}^{l_j}P_n-\ldots-\mathcal{A}^{l_1+\cdots+l_{j-1}}Q_n\mathcal{A}^{l_j}P_n.
$$
\n(279)

Using the cyclic property of the trace, we get

$$
\operatorname{Tr}\left(\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{A}^m P_n\right)=-\sum_{k=2}^j\operatorname{Tr}\left(\mathcal{A}^{l_k}P_n\cdots P_n\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\right).
$$
 (280)

Similarly, using the telescoping sum and cyclic property of the trace operator, we also get

$$
\begin{split}\n\left(\operatorname{Tr}\left(\mathcal{A}^{l_1}P_n\cdots\mathcal{A}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{A}^{m}P_n\right)\right)-\left(\operatorname{Tr}\left(\mathcal{B}^{l_1}P_n\cdots\mathcal{B}^{l_j}P_n\right)-\operatorname{Tr}\left(\mathcal{B}^{m}P_n\right)\right) \\
&= \sum_{k=2}^{j}-\operatorname{Tr}\left(\mathcal{A}^{l_k}P_n\cdots P_n\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\right)+\operatorname{Tr}\left(\mathcal{B}^{l_k}P_n\cdots P_n\mathcal{B}^{l_j}P_n\mathcal{B}^{l_1+\cdots+l_{k-1}}Q_n\right) \\
&= -\sum_{k=2}^{j-1}\left(\operatorname{Tr}\left((\mathcal{A}^{l_k}-\mathcal{B}^{l_k})P_n\mathcal{A}^{l_{k+1}}\cdots P_n\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\right)\right. \\
&\left.+\sum_{i=k}^{j-1}\operatorname{Tr}\left(\mathcal{B}^{l_k}P_n\cdots P_n\mathcal{B}^{l_i}P_n(\mathcal{A}^{l_{i+1}}-\mathcal{B}^{l_{i+1}})P_n\cdots\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\right)\right. \\
&\left.+\operatorname{Tr}\left(\mathcal{B}^{l_k}P_n\cdots P_n\mathcal{B}^{l_j}P_n(\mathcal{A}^{l_1+\cdots+l_{k-1}}-\mathcal{B}^{l_1+\cdots+l_{k-1}})Q_n\right)\right),\n\end{split} \tag{281}
$$

here we take the convention that $\sum_{j=a}^{b} \equiv 0$ for any integers *b* < *a*.

Consider $\mathcal{A} = F_{n \pm 2mn\beta}$ and $\mathcal{B} = \sum_r c_r T_{n \pm 2mn\beta} (\eta_r)$

Recall that the operator norm of a linear operator can be estimated by entries, i.e., $||\mathcal{B}||_{\infty} \leq$ $\sum_{j=-\infty}^{\infty} \sup_k |(\mathcal{B})_{k,k+j}|$. Together with [\(122\)](#page-26-2) in Proposition [4.3,](#page-26-0) we compute that $n^{-\alpha}||\mathcal{B}||_{\infty}$ is uniformly bounded for all *n*. That is there exists a constant $C_{op} > 0$ such that

$$
\max\{\|\mathcal{A}\|_{\infty}, \|\mathcal{B}\|_{\infty}\} \le n^{\alpha}C_{op}.\tag{282}
$$

Without loss of generality we will assume $C_{op} > 1$.

Apply [\(188\)](#page-37-6) in Lemma [5.4,](#page-37-5) and apply the trace norm inequality, ∥*AB*∥¹ ≤ ∥*A*∥∞∥*B*∥1, and the fact that, for *ⁱ* < *^m* to obtain

$$
\|Q_{n+mn}\beta \mathcal{A}^i P_n\|_1 = \|Q_{n+mn}\beta Q_{n+in}\beta \mathcal{A}^i P_n\|_1 \le \|Q_{n+mn}\beta\|_{\infty} \|Q_{n+in}\beta \mathcal{A}^i P_n\|_1 \le C_m e^{-d'n\beta - \frac{\alpha}{2}}.
$$
 (283)

Use the fact $\mathcal{A}^{l_k} - \mathcal{B}^{l_k} = \sum_{i=0}^{l_k-1} \mathcal{B}^{l_k-1-i}(\mathcal{A} - \mathcal{B})\mathcal{A}^i$, together with [\(283\)](#page-51-0), to obtain

$$
\left\| \left(\mathcal{A}^{l_{k}} - \mathcal{B}^{l_{k}} \right) P_{n} - \sum_{i=0}^{l_{k}-1} \mathcal{B}^{l_{k}-1-i} (\mathcal{A} - \mathcal{B}) P_{n+mn\beta} \mathcal{A}^{i} P_{n} \right\|_{1} = \left\| \sum_{i=0}^{l_{k}-1} \mathcal{B}^{l_{k}-1-i} (\mathcal{A} - \mathcal{B}) Q_{n+mn\beta} \mathcal{A}^{i} P_{n} \right\|_{1}
$$

$$
\leq \sum_{i=0}^{l_{k}-1} \left\| \mathcal{B}^{l_{k}-1-i} (\mathcal{A} - \mathcal{B}) \right\|_{\infty} \left\| Q_{n+mn\beta} \mathcal{A}^{i} P_{n} \right\|_{1} \leq 2 l_{k} \left(n^{\alpha} C_{op} \right)^{l_{k}-i} C_{m} e^{-d' n^{\beta - \frac{\alpha}{2}}}.
$$
 (284)

Let us define an error term, which is exponentially small for large *n*,

$$
R_n \coloneqq 2m C_m \left(C_{op} \right)^m n^{\alpha m} e^{-d' n^{\beta - \frac{\alpha}{2}}}.
$$
\n
$$
(285)
$$

Then, for $k = 2, \ldots, j - 1$, use [\(284\)](#page-52-0) to estimate the first summand in [\(281\)](#page-51-1) to be

$$
\left| \text{Tr} \left((\mathcal{A}^{l_k} - \mathcal{B}^{l_k}) P_n \mathcal{A}^{l_{k+1}} \cdots P_n \mathcal{A}^{l_j} P_n \mathcal{A}^{l_1 + \cdots + l_{k-1}} Q_n \right) \right|
$$

$$
\leq \left\| \sum_{i=0}^{l_{k-1}} \mathcal{B}^{l_k - 1 - i} (\mathcal{A} - \mathcal{B}) P_{n + m n^{\beta}} \mathcal{A}^i P_n \mathcal{A}^{l_{k+1}} \cdots P_n \mathcal{A}^{l_j} P_n \mathcal{A}^{l_1 + \cdots + l_{k-1}} Q_n \right\|_1 + R_n. (286)
$$

We will commute Q_n from the right to the left of [\(286\)](#page-52-1). To this end, similarly to [\(218\)](#page-42-3) in the proof of Proposition [4.2,](#page-25-0) write $P_n = Id - Q_n$, use the telescopic sum to get

$$
\mathcal{A}^{i}P_{n}\mathcal{A}^{l_{k+1}}\cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n} = -\mathcal{A}^{i}Q_{n}\mathcal{A}^{l_{k+1}}\cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n}
$$

$$
-\mathcal{A}^{i+l_{k+1}}Q_{n}\mathcal{A}^{l_{k+2}}P_{n}\cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n} - \mathcal{A}^{i+l_{k+1}+l_{k+2}}Q_{n}\mathcal{A}^{l_{k+3}}P_{n}\cdots P_{n}\mathcal{A}^{l_{j}}P_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n}
$$

$$
-\cdots - \mathcal{A}^{i+l_{k+1}+l_{k+2}+\cdots+l_{j}}Q_{n}\mathcal{A}^{l_{1}+\cdots+l_{k-1}}Q_{n} + \mathcal{A}^{i+m-l_{k}}Q_{n}.
$$
 (287)

Note that there are $j - k + 2$ terms on the right-hand side of [\(287\)](#page-52-2) and $j - k + 2 \le m$.

Applying [\(203\)](#page-39-6) in Lemma [5.5](#page-39-5) to [\(279\)](#page-51-2), we have, for any $l \in \mathbb{N}$ with $l \leq m$,

$$
\|\mathcal{A}^l Q_n - Q_{n-mn}\beta \mathcal{A}^l Q_n\|_1 = \|P_{n-mn}\beta P_{n-ln}\beta \mathcal{A}^l Q_n\|_1 \le C_m e^{-d'n^{\beta - \frac{\alpha}{2}}}.
$$
\n(288)

Hence, estimate [\(287\)](#page-52-2) by [\(288\)](#page-52-3) to obtain

$$
\left\| \mathcal{A}^i P_n \mathcal{A}^{l_{k+1}} \cdots P_n \mathcal{A}^{l_j} P_n \mathcal{A}^{l_1 + \cdots + l_{k-1}} Q_n - Q_{n-mn} \mathcal{A}^i P_n \mathcal{A}^{l_{k+1}} \cdots P_n \mathcal{A}^{l_j} P_n \mathcal{A}^{l_1 + \cdots + l_{k-1}} Q_n \right\|_1
$$

\$\leq (j - k + 2) C_m \left(n^{\alpha} C_{op} \right)^{m - l_k - i} e^{-d' n^{\beta - \frac{\alpha}{2}}} .\tag{289}

Plug [\(289\)](#page-52-4) into the right-hand side of [\(286\)](#page-52-1), , and we get

$$
\|\mathcal{B}^{l_k-1-i}(\mathcal{A}-\mathcal{B})P_{n+mn\beta}\mathcal{A}^i P_n\mathcal{A}^{l_{k+1}}\cdots P_n\mathcal{A}^{l_j}P_n\mathcal{A}^{l_1+\cdots+l_{k-1}}Q_n\|_1
$$

$$
\leq (n^{\alpha}C_{op})^{m-1}\|(\mathcal{A}-\mathcal{B})P_{n+mn\beta}Q_{n-mn\beta}\|_1 + R_n. \quad (290)
$$

Plug [\(290\)](#page-52-5) into the right-hand side of [\(286\)](#page-52-1), and we get

$$
\left| \operatorname{Tr} \left((\mathcal{A}^{l_k} - \mathcal{B}^{l_k}) P_n \mathcal{A}^{l_{k+1}} \cdots P_n \mathcal{A}^{l_j} P_n \mathcal{A}^{l_1 + \cdots + l_{k-1}} Q_n \right) \right|
$$

\$\leq l_k (n^{\alpha} C_{op})^{m-1} || (\mathcal{A} - \mathcal{B}) P_{n+mn^{\beta}} Q_{n-mn^{\beta}} ||_1 + (l_k + 1) R_n. (291)

Now we estimate the second summand in [\(281\)](#page-51-1). For $i = k, \ldots, j - 1$, similarly,

$$
\left| \text{Tr} \left(\mathcal{B}^{l_k} P_n \cdots P_n \mathcal{B}^{l_i} P_n (\mathcal{A}^{l_{i+1}} - \mathcal{B}^{l_{i+1}}) P_n \cdots \mathcal{A}^{l_j} P_n \mathcal{A}^{l_1 + \cdots + l_{k-1}} Q_n) \right| \right|
$$

\$\leq l_{i+1} (n^{\alpha} C_{op})^{m-1} || (\mathcal{A} - \mathcal{B}) P_{n+mn^{\beta}} Q_{n-mn^{\beta}} ||_1 + (l_{i+1} + 1) R_n. (292)\$

Lastly, we turn to estimate the last summand in [\(281\)](#page-51-1). By [\(122\)](#page-26-2) in Proposition [4.3,](#page-26-0) we have for all $|j - k| \geq n^{\beta}$

$$
\left| \left(\mathcal{B} \right)_{j,k} \right| \leq C_0 n^{\frac{\alpha}{2}} e^{-d_0 n^{\beta - \frac{\alpha}{2}}}.
$$
\n(293)

Then use the same proof as in Lemma [5.4,](#page-37-5) we have for any $l \in \mathbb{N}$ with $l \leq m$,

$$
||P_n \mathcal{B}^l - P_n \mathcal{B}^l P_{n+mn} \mathcal{B}||_1 \le C_m e^{-d'n^{\beta - \frac{\alpha}{2}}}.
$$
\n(294)

Hence

$$
\left| \text{Tr} \left(\mathcal{B}^{l_k} P_n \cdots P_n \mathcal{B}^{l_j} P_n (\mathcal{A}^{l_1 + \cdots + l_{k-1}} - \mathcal{B}^{l_1 + \cdots + l_{k-1}}) Q_n) \right| \right|
$$

$$
\leq \left| \text{Tr} \left(\mathcal{B}^{l_k} P_n \cdots P_n \mathcal{B}^{l_j} P_n (\mathcal{A}^{l_1 + \cdots + l_{k-1}} - \mathcal{B}^{l_1 + \cdots + l_{k-1}}) P_{n + m n^{\beta}} Q_n) \right| + R_n.
$$
 (295)

Repeat the argument similar to [\(291\)](#page-52-6) to obtain

$$
\left| \text{Tr} \left(\mathcal{B}^{l_k} P_n \cdots P_n \mathcal{B}^{l_j} P_n (\mathcal{A}^{l_1 + \cdots + l_{k-1}} - \mathcal{B}^{l_1 + \cdots + l_{k-1}}) Q_n) \right| \right|
$$

$$
\leq (l_1 + \cdots + l_{k-1}) (n^{\alpha} C_{op})^{m-1} ||(\mathcal{A} - \mathcal{B}) P_{n + m n^{\beta}} Q_{n - m n^{\beta}} ||_1 + ((l_1 + \cdots + l_{k-1}) + 1) R_n. (296)
$$

Hence, plugging three estimates [\(291\)](#page-52-6), [\(292\)](#page-53-0) and [\(296\)](#page-53-1) into the formula [\(281\)](#page-51-1), we get

$$
\left| \text{Tr}(\mathcal{A})^{l_1} P_n \cdots (\mathcal{A})^{l_j} P_n - \text{Tr}(\mathcal{A}^m P_n) - \text{Tr}(\mathcal{B})^{l_1} P_n \cdots (\mathcal{B})^{l_j} P_n + \text{Tr}(\mathcal{B}^m P_n) \right|
$$

$$
\leq (m+1) \left(\left(n^{\alpha} C_{op} \right)^{m-1} ||(\mathcal{A} - \mathcal{B}) P_{n+mn^{\beta}} Q_{n-mn^{\beta}} ||_1 + R_n \right). \quad (297)
$$

Note that,

$$
\|(\mathcal{A} - \mathcal{B})P_{n+mn\beta}Q_{n-mn\beta}\|_1 \leq \sum_r |c_r| \|H_{n\pm 2mn\beta}(\eta_r)P_{n+mn\beta}Q_{n-mn\beta}\|_1,
$$

where $H_{n\pm 2mn\beta}(\eta_r)$ is defined in [\(264\)](#page-48-1). Thus by Proposition [4.3,](#page-26-0) we have

$$
||H_{n\pm 2mn^{\beta}}(\eta_r)P_{n+mn^{\beta}}Q_{n-mn^{\beta}}||_1
$$

\n
$$
\leq \sum_{j=n-2mn^{\beta}+1}^{n+2mn^{\beta}} \sum_{k=n-mn^{\beta}+1}^{n+mn^{\beta}} Constant \quad n^{\frac{\alpha}{2}} \left(\left(1 - \frac{d}{n^{\frac{\alpha}{2}}} \right)^{\max\{j,k\}-n+2mn^{\beta}} + \left(1 - \frac{d}{n^{\frac{\alpha}{2}}} \right)^{n+2mn^{\beta}-\min\{j,k\}} \right)
$$

\n
$$
\leq Constant \quad 16mn^{2\beta+\frac{\alpha}{2}}e^{-dmn^{\beta-\frac{\alpha}{2}}}.
$$
 (298)

This shows that [\(297\)](#page-53-2) is exponentially small. Therefore we conclude [\(125\)](#page-27-1).

□

7 Proof of Proposition [4.5](#page-27-2)

Before we prove the Proposition [4.5,](#page-27-2) we need to study the limiting behaviour of the following

$$
\log \det \left(1 + P_n \left(e^{tn^{-\alpha} \sum_r c_r T_{n \pm 2mn} \beta(\eta_r)} - 1 \right) P_n \right) e^{-t \operatorname{Tr} P_n n^{-\alpha} \sum_r c_r T_{n \pm 2mn} \beta(\eta_r)}, \tag{299}
$$

where $T_{n\pm 2mn\beta}$ comes from Proposition [4.3.](#page-26-0)

7.1 Limiting Behaviour of Fredholm Determinants

Suppose *A* and *B* are bounded operators such that $[A, B]$ is trace class. Then [\[37\]](#page-82-0) shows the following principle

$$
e^{-A}e^{A+B}e^{-B} - Id \text{ is trace class,}
$$
 (300)

and

$$
\det e^{-A} e^{A+B} e^{-B} = e^{-\frac{1}{2} \text{Tr}[A,B]}.
$$
\n(301)

This is a generalisation of the Helton-Howe-Pincus formula. It is useful to study generating functions for some Gaussian random variables, see for example [\[14\]](#page-81-0). We will also use this principle in our setup following [\[14\]](#page-81-0), which is a direct consequence of Lemmas [7.1](#page-54-0) and [7.2.](#page-54-1)

Lemma 7.1. *For any bounded operators A*, *B we have*

$$
e^{-A}e^{A+B}e^{-B} - Id = \sum_{m_1,m_2,m_3=0}^{\infty} \sum_{j=0}^{m_2-1} \frac{(-1)^{m_1+m_3}A^{m_1}(A+B)^j[A,B](A+B)^{m_2-j-1}B^{m_3}}{m_1!m_2!m_3!(m_1+m_2+m_3+1)}.
$$
(302)

Moreover, if [*A, B*] *is trace class then* $e^{-A}e^{A+B}e^{-B} - Id$ *is trace class.*

Proof. See Lemma 4.3 in [\[14\]](#page-81-0). □

Lemma 7.2. *Let Tn*±2*mn*^β (η*r*)[−] *be the strict upper triangle part of Tn*±2*mn*^β (η*r*) *and Tn*±2*mn*^β (η*r*)⁺ *be the lower triangle part of* $T_{n+2mn\beta}$ (η_r) *. Then we have*

$$
\det \left(Id + P_n \left(e^{tn^{-\alpha} \sum_r c_r T_{n \pm 2m\beta}(\eta_r)} - Id \right) P_n \right) e^{-tn^{-\alpha} \text{Tr } P_n \sum_r c_r T_{n \pm 2m\beta}(\eta_r)} \\
= e^{-\frac{t^2}{2n^{2\alpha}} \text{Tr} \left[\sum_r c_r T_{n \pm 2m\beta}(\eta_r)_{+}, \sum_r c_r T_{n \pm 2m\beta}(\eta_r)_{-} \right]} \det \left(Id + Q_n (R(t, \eta)^{-1} - Id) \right), \quad (303)
$$

where

$$
R(t,\eta) := e^{-tn^{-\alpha} \sum_r c_r T_{n\pm 2m\theta}(\eta_r)} + e^{tn^{-\alpha} \sum_r c_r T_{n\pm 2m\theta}(\eta_r)} e^{-tn^{-\alpha} \sum_r c_r T_{n\pm 2m\theta}(\eta_r)}.
$$
(304)

Proof. See the proof of Lemmas 4.2 and 4.4 in [\[14\]](#page-81-0). \square

Lemma 7.3. *Set*

$$
T \coloneqq \sum_{r} c_{r} T_{n \pm 2mn^{\beta}} \left(\eta_{r} \right).
$$

Let T[−] *be the strict upper triangle part of T and T*⁺ *be the lower triangle part of T. Then there exists a constant C*′ > ⁰ *such that*

$$
n^{-2\alpha} \left\| [T_+, T_-] \right\|_1 < C',\tag{305}
$$

$$
\lim_{n \to \infty} n^{-2\alpha} ||Q_n[T_+, T_-]||_1 = 0,
$$
\n(306)

$$
\lim_{n \to \infty} n^{-2\alpha} \left(\text{Tr}[T_+, T_-] + \sum_{j=n-2mn^{\beta}+1}^{n} \sum_{l=2j-n+2mn^{\beta}}^{n+2mn^{\beta}} (T_{j,l})^2 \right) = 0. \tag{307}
$$

Proof. Then by Proposition [4.3](#page-26-0) we have for all *r*, *j*, *k*, *l* with $n - 2mn^{\beta} \le j$, $j + l$, k , $k + l \le n + 2mn^{\beta}$,

$$
\frac{\left(T_{n\pm 2mn^{\beta}}\left(\eta_{r}\right)\right)_{j,k}}{\left(T_{n\pm 2mn^{\beta}}\left(\eta_{r}\right)\right)_{j+l,k+l}}=1+o\left(n^{-\beta+\frac{\alpha}{2}}\right),\quad\left|\left(T_{n\pm 2mn^{\beta}}\left(\eta_{r}\right)\right)_{j,k}\right|\leq Cn^{\frac{\alpha}{2}}\left|1-\frac{d}{n^{\frac{\alpha}{2}}}\right|^{k-j}.\tag{308}
$$

This implies that

$$
\frac{(T)_{j,k}}{(T)_{j+l,k+l}} = 1 + o\left(n^{-\beta + \frac{\alpha}{2}}\right), \quad \left|(T)_{j,k}\right| \le Cn^{\frac{\alpha}{2}} \left|1 - \frac{d}{n^{\frac{\alpha}{2}}}\right|^{k-j}.\tag{309}
$$

Note that *T*[−] = *Pn*+2*nm*β*Qn*−2*mn*β*T*[−] = *T*−*Pn*+2*nm*β*Qn*−2*mn*^β . Hence the commutator

$$
[T_+, T_-] = T_+ P_{n+2nm\beta} Q_{n-2mn\beta} T_- - T_- P_{n+2nm\beta} Q_{n-2mn\beta} T_+.
$$

This implies that for $j, k = n - 2mn^{\beta} + 1, ..., n + 2mn^{\beta}$

$$
([T_{+}, T_{-}])_{j,k} = \sum_{l=n-2mn^{\beta}+1}^{min\{j,k+1\}} (T)_{j,l} (T)_{l,k} - \sum_{l=\max\{j-1,k\}}^{n+2mn^{\beta}} (T)_{j,l} (T)_{l,k}
$$

$$
= \sum_{l=n-2mn^{\beta}+1}^{min\{j,k+1\}} ((T)_{j,l} (T)_{l,k} - (T)_{j,j+k-l} (T)_{j+k-l,k}) - \sum_{l=j+k-n+2mn^{\beta}}^{n+2mn^{\beta}} (T)_{j,l} (T)_{l,k} \chi_{j+k \le 2n}
$$

$$
= \sum_{l=n-2mn^{\beta}+1}^{min\{j,k+1\}} o(n^{-\beta+\frac{\alpha}{2}}) (T)_{j,l} (T)_{l,k} - \sum_{l=j+k-n+2mn^{\beta}}^{n+2mn^{\beta}} (T)_{j,l} (T)_{l,k} \chi_{j+k \le 2n}, \quad (310)
$$

where $\chi_{j+k\leq 2n} = 1$ if $j + k \leq 2n$ and zero otherwise. For other values of *j*, *k* we have $([T_+, T_-])_{jk} = 0$. Note that though the actual formula of $o(n^{-\beta+\frac{\alpha}{2}})$ above depends on *j*, *k*, *l*, it has an uniform limiting behaviour as $n \to \infty$. Denote \overline{T} to be the linear operator such that each entries

$$
\left(\widetilde{T}\right)_{j,k} := \left|(T)_{j,k}\right|.
$$

Also define

$$
(H)_{j,k} := \begin{cases} Cn^{\frac{\alpha}{2}} \left(1 - \frac{d}{n^{\frac{\alpha}{2}}}\right)^{j+k} & \text{for } n - 2mn^{\beta} < j, k \le n + 2mn^{\beta}, \\ 0 & \text{otherwise.} \end{cases}
$$

Then we estimate

$$
\| [T_+, T_-] \|_1 \le o\left(n^{-\beta + \frac{\alpha}{2}} \right) \|\widetilde{T_+}\|_2 \|\widetilde{T_-}\|_2 + \|H\|_2^2.
$$

Using the matrix norm relation $||AB||_1 \le ||A||_2 ||B||_2$ and $||T||_2 \le ||\widetilde{T}||_2$. Further using $||T||_2^2 = \sum_{j,k} |(T)_{j,k}|^2$, we have for any *n* with $n^{\frac{1}{2}} > d$

$$
\|\widetilde{T}_{+}\|_{2}\|\widetilde{T}_{-}\|_{2} \leq C^{2} n^{\alpha} \sum_{l=0}^{N} (N-l) \left|1 - \frac{d}{n^{\frac{\alpha}{2}}} \right|^{2l} \leq \frac{C^{2} c'}{d} n^{\beta + \frac{3\alpha}{2}},\tag{311}
$$

$$
||H||_2^2 \le n^{\alpha} \sum_{j,k=1}^N \left| 1 - \frac{d}{n^{\frac{\alpha}{2}}} \right|^{2j+2k} \le \frac{n^{2\alpha}}{d^2},\tag{312}
$$

$$
||Q_{n^{\beta}}H||_{2}^{2} \leq n^{\alpha} \sum_{j=n^{\beta}}^{N} \sum_{k=1}^{N} \left|1 - \frac{d}{n^{\frac{\alpha}{2}}}\right|^{2j+2k} \leq \frac{n^{2\alpha}}{d^{2}} e^{-2dn^{\beta-\frac{\alpha}{2}}}.
$$
 (313)

Together with the assumption $\frac{\alpha}{2} < \beta < \frac{\alpha+1}{3}$, this shows

$$
\frac{1}{n^{2\alpha}}\left\|[T_+,T_-\]\right\|_1 < \infty, \quad \frac{1}{n^{2\alpha}}\left\|Q_{n^\beta}[T_+,T_-\]\right\|_1 \to 0. \tag{314}
$$

Combine [\(310\)](#page-55-0) and [\(311\)](#page-55-1) and we conclude

$$
\lim_{n \to \infty} n^{-2\alpha} \left(\text{Tr}[T_+, T_-] + \sum_{j=n-2mn^{\beta}+1}^{n} \sum_{l=2j-n+2mn^{\beta}}^{n+2mn^{\beta}} (T_{j,l})^2 \right) = 0.
$$

Lemma 7.4. *Recall that* $R(t, \eta)^{-1}$ *is defined as* [\(304\)](#page-54-2)*. For any* β *such that* $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$ *, we have*

$$
\lim_{n \to \infty} ||Q_n(R(t, \eta)^{-1} - Id)||_1 = 0. \tag{315}
$$

Proof. Set

$$
T \coloneqq \sum_r c_r T_{n \pm 2mn^\beta}(\eta_r).
$$

Let *T*[−] be the strict upper triangle part of *T* and *T*⁺ be the lower triangle part of *T*. Then

$$
R(t,\eta)^{-1} = e^{tn^{-\alpha}T} - e^{-tn^{-\alpha}T} e^{tn^{-\alpha}T}.
$$
\n(316)

Use the expansion formula in Lemma [7.1](#page-54-0) to obtain

$$
R(t,\eta)^{-1} - Id = \sum_{m_1,m_2,m_3=0}^{\infty} (-1)^{m_2-1} (tn^{-\alpha})^{m_1+m_2+m_3+1} \sum_{j=0}^{m_2-1} \frac{(T-1)^{m_1}(T-1+1)^j [T-1+1] (T-1+1)^{m_2-j-1} (T+1)^{m_3}}{m_1! m_2! m_3! (m_1+m_2+m_3+1)}.
$$

By the operator norm inequality $||A||_{\infty} \le \sum_{j=-\infty}^{\infty} \sup_k |(A)_{k,k+j}|$ and the second estimate of [\(309\)](#page-55-2) we have

$$
\max\{\|T_{+}\|_{\infty}, \|T_{-}\|_{\infty}\} \le \sum_{r} |c_{r}| \sum_{l=0}^{\infty} C n^{\frac{\alpha}{2}} \left|1 - \frac{d}{n^{\frac{\alpha}{2}}} \right|^{l} = \frac{C \sum_{r} |c_{r}|}{d} n^{\alpha}.
$$
 (317)

By Lemma [7.3,](#page-54-3) there exists a constant $C' > 0$ such that

$$
n^{-2\alpha} \left\| [T_-, T_+] \right\|_1 < C', \tag{318}
$$

$$
n^{-2\alpha} \|\mathcal{Q}_n[T_-, T_+]\|_1 \to 0. \tag{319}
$$

Then by the equalities $Q_nT_+ = Q_nT_+Q_n$ and $Q_nT_- = Q_nT_-Q_n$, and by the dominated convergence theorem we have lim *n*→∞ $\left| R(t, \eta)^{-1} - Id \right|_1 = 0.$

Lemma 7.5. *Consider c* > 0 *and a sequence* $\xi_i \in \mathbb{C}$ *such that* $\sup_{i \in \mathbb{N}} |\xi_i| = o(n^{-\frac{\alpha}{2}})$ *as* $n \to \infty$ *. Then we have*

$$
\lim_{n \to \infty} \frac{1}{n^{\frac{\alpha}{2}}} \sum_{k=0}^{\infty} \left| \prod_{i=1}^{k} \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} + \xi_i \right) - \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} \right)^k \right| = 0.
$$
 (320)

Proof. Use the telescoping sum to obtain for $k \geq 1$

$$
\prod_{i=1}^{k} \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} + \xi_i \right) - \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} \right)^k = \sum_{l=1}^{k} \xi_l \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} \right)^{l-1} \prod_{i=l}^{k} \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} + \xi_i \right).
$$
(321)

Hence as $n \to \infty$,

$$
\left| \prod_{i=1}^{k} \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} + \xi_i \right) - \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} \right)^k \right| \le k \left(1 - \frac{c}{2n^{\frac{\alpha}{2}}} \right)^{k-1} o(n^{-\frac{\alpha}{2}}).
$$
 (322)

By the fact that

$$
\sum_{k=1}^{\infty} k \left(1 - \frac{c}{2n^{\frac{\alpha}{2}}} \right)^{k-1} = 4c^{-2}n^{\alpha},\tag{323}
$$

we conclude

$$
\lim_{n \to \infty} \frac{1}{n^{\frac{\alpha}{2}}} \sum_{k=0}^{\infty} \left| \prod_{i=1}^{k} \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} + \xi_i \right) - \left(1 - \frac{c}{n^{\frac{\alpha}{2}}} \right)^k \right| = 0.
$$
 (324)

□

Lemma 7.6. *Set*

$$
T := \sum_{r=1}^{2M} c_r T_{n \pm 2mn^{\beta}} \left(\eta_r \right).
$$

Consider the test function $f(x) = \sum_{r=1}^{2M} \frac{c_r}{x-r}$ *x*^{*x*} *while the test function* $f(x) = \sum_{r=1}^{2M} \frac{c_r}{x-\eta_r}$ *with* c_j , η_r *defined as* [\(49\)](#page-14-0)*. Then, we have at the left edge*

$$
\lim_{n \to \infty} \frac{1}{n^{2\alpha}} \operatorname{Tr}[T_+, T_-] = \frac{-1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy. \tag{325}
$$

At the right edge, we have

$$
\lim_{n \to \infty} \frac{1}{n^{2\alpha}} \operatorname{Tr}[T_+, T_-] = \frac{-1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy. \tag{326}
$$

Proof. By the limit [\(307\)](#page-54-4) of Lemma [7.3,](#page-54-3) it is sufficient to compute the following limit

$$
\lim_{n \to \infty} \frac{1}{n^{2\alpha}} \sum_{j=n-2mn^{\beta}+1}^{n} \sum_{l=2j-n+2mn^{\beta}}^{n+2mn^{\beta}} (T_{j,l})^{2}.
$$
 (327)

We estimate each entry of the fourth-fold sum [\(327\)](#page-57-0), by Proposition [4.3,](#page-26-0) that is for any *^r*, *^s* and *^j*, *^l*

$$
(T_{n\pm 2mn^{\beta}}(\eta_r))_{j,l}(T_{n\pm 2mn^{\beta}}(\eta_s))_{j,l} = \frac{n^{\alpha} \prod_{i=j}^{l-1} \left(1 - \left(-\frac{\eta_r}{n^{\alpha}|a_{n,n}|}\right)^{\frac{1}{2}} - \left(-\frac{\eta_s}{n^{\alpha}|a_{n,n}|}\right)^{\frac{1}{2}} + \xi_i^{(r,s)}\right)(1+o(1))}{4a_n^2 \left(\frac{-\eta_r}{|a_{n,n}|}\right)^{\frac{1}{2}} \left(\frac{-\eta_s}{|a_{n,n}|}\right)^{\frac{1}{2}}},\qquad(328)
$$

as $n \to \infty$, where $\xi_i^{(r,s)} \coloneqq \xi_i^{(r)}$ $\zeta_i^{(r)} + \xi_i^{(s)}$ $\zeta_i^{(s)} + \xi_i^{(r)}$ *i* ξ (*s*) $\zeta_i^{(s)} + \xi_i^{(r)}$ $\int_{i}^{(r)} \left(-\frac{\eta_s}{n^{\alpha} |a_{n,n}|} \right)$ $\int_0^{\frac{1}{2}} + \xi_i^{(s)}$ $\int_{i}^{(s)} \left(-\frac{\eta_r}{n^{\alpha}|a_{n,n}|} \right)$ $\big)^{\frac{1}{2}} = o(n^{-\frac{\alpha}{2}}).$ Relabel the indices with respect to the summations to obtain, as $n \to \infty$,

$$
\frac{1}{n^{2\alpha}} \sum_{j=n-2mn^{\beta}+1}^{n+2mn^{\beta}} \sum_{l=2j-n+2mn^{\beta}}^{n+2mn^{\beta}} (T_{j,l})^2
$$
\n
$$
= \frac{1}{n^{\alpha}} \sum_{r,s=1}^{2M} c_r c_s \sum_{j=n-2mn^{\beta}+1}^{n+2mn^{\beta}-j} \sum_{k=j-n+2mn^{\beta}}^{n+2mn^{\beta}-j} \frac{\prod_{i=j}^{k+j-1} \left(1 - \left(-\frac{\eta_r}{|a_{n,n}|}\right)^{\frac{1}{2}} - \left(-\frac{\eta_s}{n^{\alpha}|a_{n,n}|}\right)^{\frac{1}{2}} + \xi_i^{(r,s)}\right) (1+o(1))}{4a_n^2 \left(\frac{-\eta_r}{|a_{n,n}|}\right)^{\frac{1}{2}} \left(\frac{-\eta_s}{|a_{n,n}|}\right)^{\frac{1}{2}}}
$$
\n
$$
= \frac{1}{n^{\alpha}} \sum_{r,s=1}^{2M} c_r c_s \sum_{j=1}^{2mn^{\beta}+mn^{\beta}-j} \sum_{k=j}^{n+2mn^{\beta}-1} \frac{\prod_{i=j+n-2mn^{\beta}}^{n+2mn^{\beta}-1} \left(1 - \left(-\frac{\eta_r}{|a_{n,n}|}\right)^{\frac{1}{2}} - \left(-\frac{\eta_s}{n^{\alpha}|a_{n,n}|}\right)^{\frac{1}{2}} + \xi_i^{(r,s)}\right) (1+o(1))}{4a_n^2 \left(\frac{-\eta_r}{|a_{n,n}|}\right)^{\frac{1}{2}} \left(\frac{-\eta_s}{|a_{n,n}|}\right)^{\frac{1}{2}}}
$$
\n(329)

Note that for all $0 < \frac{\alpha}{2} < \beta$ we have $\left(1 - \frac{c}{n^{\frac{1}{2}}} \right)$ *n* α 2 \int_{0}^{π} = $O(e^{-n^{\beta-\frac{\alpha}{2}}})$ for any $c \neq 0$, which is exponentially small. Hence, adding exponentially small terms does not change the sum in the limit, i.e., as $n \to \infty$,

$$
\frac{1}{n^{2\alpha}} \sum_{j=n-2mn^{\beta}+1}^{n} \sum_{l=2j-n+2mn^{\beta}}^{n+2mn^{\beta}} (T_{j,l})^{2}
$$
\n
$$
= \frac{1}{n^{\alpha}} \sum_{r,s=1}^{2M} c_{r} c_{s} \sum_{j=1}^{2mn^{\beta}} \sum_{k=j}^{\infty} \frac{\prod_{i=j+n-2mn^{\beta}-1}^{k+j+n-2mn^{\beta}-1} \left(1 - \left(-\frac{\eta_{r}}{|a_{n,n}|}\right)^{\frac{1}{2}} - \left(-\frac{\eta_{s}}{n^{\alpha}|a_{n,n}|}\right)^{\frac{1}{2}} + \xi_{i}^{(r,s)}\right) (1+o(1))}{4a_{n}^{2} \left(\frac{-\eta_{r}}{|a_{n,n}|}\right)^{\frac{1}{2}} \left(\frac{-\eta_{s}}{|a_{n,n}|}\right)^{\frac{1}{2}}} + o(1). \quad (330)
$$

Then by Lemma [7.5](#page-56-0) we estimate the power sum [\(330\)](#page-58-0) to be, as $n \to \infty$,

$$
\frac{1}{n^{2\alpha}} \sum_{j=n-2mn\beta+1}^{n} \sum_{l=2j-n+2mn\beta}^{n+2mn\beta} \left(T_{j,l}\right)^2 = \frac{1}{n^{\alpha}} \sum_{r,s=1}^{2M} c_r c_s \sum_{j=1}^{2mn\beta} \sum_{k=j}^{\infty} \frac{\left(1 - \left(-\frac{\eta_r}{|a_{n,n}|}\right)^{\frac{1}{2}} - \left(-\frac{\eta_s}{n^{\alpha}|a_{n,n}|}\right)^{\frac{1}{2}}\right)^k (1+o(1))}{4a_n^2 \left(\frac{-\eta_r}{|a_{n,n}|}\right)^{\frac{1}{2}} \left(\frac{-\eta_s}{|a_{n,n}|}\right)^{\frac{1}{2}}} + o(1)
$$
\n
$$
= \sum_{r,s=1}^{2M} c_r c_s \frac{1}{4\left(-\eta_r\right)^{\frac{1}{2}} \left(-\eta_s\right)^{\frac{1}{2}} \left(-\eta_s\right)^{\frac{1}{2}} + \left(-\eta_s\right)^{\frac{1}{2}}} + o(1). \quad (331)
$$

Recall that c_j and η_j are defined as in [\(49\)](#page-14-0). Hence we have, as $n \to \infty$,

$$
\frac{1}{n^{2\alpha}} \sum_{j=n-2mn^{\beta}+1}^{n} \sum_{l=2j-n+2mn^{\beta}}^{n+2mn^{\beta}} (T_{j,l})^{2}
$$
\n
$$
= \frac{1}{2} \text{Re} \left(\sum_{r,s=1}^{M} \frac{-d_{r}d_{s}}{(-\lambda_{r})^{\frac{1}{2}} (-\lambda_{s})^{\frac{1}{2}} \left((-\lambda_{r})^{\frac{1}{2}} + (-\lambda_{s})^{\frac{1}{2}} \right)^{2}} + \frac{d_{r}d_{s}}{(-\overline{\lambda_{r}})^{\frac{1}{2}} (-\lambda_{s})^{\frac{1}{2}} \left((-\overline{\lambda_{r}})^{\frac{1}{2}} + (-\lambda_{s})^{\frac{1}{2}} \right)^{2}} \right) + o(1). \quad (332)
$$

From here the rest is to use the residual theorem to recover the double integral formula. Recall the formulation of *f* and we have

$$
\int\int_{\mathbb{R}^2}\left(\frac{f(x^2)-f(y^2)}{x-y}\right)^2dxdy=\sum_{r,s}c_r c_s\int\int_{\mathbb{R}^2}\left(\frac{\frac{1}{x^2-\eta_r}-\frac{1}{y^2-\eta_r}}{x-y}\right)\left(\frac{\frac{1}{x^2-\eta_s}-\frac{1}{y^2-\eta_s}}{x-y}\right)dxdy.
$$

Each of the double integrals can then be rewritten in the following way

$$
\begin{split} \int\int_{\mathbb{R}^2}\Bigg&\left(\frac{\frac{1}{x^2-\eta_r}-\frac{1}{y^2-\eta_r}}{x-y}\right)\Bigg(\frac{\frac{1}{x^2-\eta_s}-\frac{1}{y^2-\eta_s}}{x-y}\Bigg)dxdy\\ &=2\int_{\mathbb{R}}\frac{x^2}{(x^2-\eta_r)(x^2-\eta_s)}dx\int_{\mathbb{R}}\frac{1}{(y^2-\eta_r)(y^2-\eta_s)}dy+2\left(\int_{\mathbb{R}}\frac{x}{(x^2-\eta_r)(x^2-\eta_s)}dx\right)^2. \end{split}
$$

Now we can apply the residual theorem to each of the three integrals above. For either $\text{Im}(\eta_r) > 0$ or Im(η_r) < 0, take the principle square root, we have Im($i(-\eta_r)^{\frac{1}{2}}$) > 0. So each integral there are two residuals to consider i.e. i(, $n^{\frac{1}{2}}$ and $i(-n)^{\frac{1}{2}}$. Using residuals to consider i.e. $i(-\eta_r)^{\frac{1}{2}}$ and $i(-\eta_s)^{\frac{1}{2}}$. Hence,

$$
\frac{1}{2\pi i} \int_{\mathbb{R}} \frac{x^2}{(x^2 - \eta_r)(x^2 - \eta_s)} dx = \frac{1}{2i((-\eta_r)^{\frac{1}{2}} + (-\eta_s)^{\frac{1}{2}})},
$$
\n
$$
\frac{1}{2\pi i} \int_{\mathbb{R}} \frac{1}{(y^2 - \eta_r)(y^2 - \eta_s)} dy = \frac{1}{2i(-\eta_r)^{\frac{1}{2}} (-\eta_s)^{\frac{1}{2}} ((-\eta_r)^{\frac{1}{2}} + (-\eta_s)^{\frac{1}{2}})};
$$
\n
$$
\frac{1}{2\pi i} \int_{\mathbb{R}} \frac{x}{(x^2 - \eta_r)(x^2 - \eta_s)} dx = 0.
$$

Together we have,

$$
\frac{1}{(2\pi i)^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy = \sum_{r,s=1}^{2M} c_r c_s \frac{-1}{2(-\eta_r)^{\frac{1}{2}} (-\eta_s)^{\frac{1}{2}} \left((-\eta_r)^{\frac{1}{2}} + (-\eta_s)^{\frac{1}{2}} \right)^2}
$$

=
$$
-Re \left(\sum_{r,s=1}^M \frac{-d_r d_s}{(-\lambda_r)^{\frac{1}{2}} (-\lambda_s)^{\frac{1}{2}} \left((-\lambda_r)^{\frac{1}{2}} + (-\lambda_s)^{\frac{1}{2}} \right)^2} + \frac{d_r d_s}{(-\overline{\lambda_r})^{\frac{1}{2}} (-\lambda_s)^{\frac{1}{2}} \left((-\overline{\lambda_r})^{\frac{1}{2}} + (-\lambda_s)^{\frac{1}{2}} \right)^2} \right).
$$

Combing [\(332\)](#page-58-1) and Lemma [5.2,](#page-34-0) we conclude that, at the left edge,

$$
\lim_{n \to \infty} \frac{1}{n^{2\alpha}} \operatorname{Tr}[T_+, T_-] = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy.
$$
 (333)

Note that at the right edge instead of [\(174\)](#page-34-1) we have

$$
\omega_{l}^{(r)} = \frac{x_0 - b_{l,n} + \frac{\eta_r}{n^{\alpha}} - \left(\left(x_0 - b_{l,n} + \frac{\eta_r}{n^{\alpha}}\right)^2 - 4a_{l-1,n}a_{l,n}\right)^{\frac{1}{2}}}{-2a_{l-1,n}}.
$$

Then by the same proof of Lemma [5.2](#page-34-0) we get as $n \to \infty$, for $l \in \{n - 2mn^{\beta}, n - 2mn^{\beta} + 1, ..., n + 2mn^{\beta}\}\,$

$$
\omega_l^{(r)-} = \left(1 - \left(\frac{\eta_r}{n^{\alpha}|a_{n,n}|}\right)^{\frac{1}{2}}\right)sgn(-a_{n,n}) + o(n^{-\frac{\alpha}{2}}). \tag{334}
$$

We can now follow the same computation and obtain that

$$
\lim_{n \to \infty} \frac{1}{n^{2\alpha}} \sum_{j=n-2mn^{\beta}+1}^{n} \sum_{l=2j-n+2mn^{\beta}}^{n+2mn^{\beta}} \left(T_{j,l}\right)^2 = \sum_{r,s=1}^{2M} c_r c_s \frac{1}{4 \left(\eta_r\right)^{\frac{1}{2}} \left(\eta_s\right)^{\frac{1}{2}} \left(\eta_r\right)^{\frac{1}{2}} + \left(\eta_s\right)^{\frac{1}{2}}\right)^2}.
$$
(335)

From here the rest is to use the residual theorem to obtain the double integral formula. Recall the formulation of *f* and we have

$$
\int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy = \sum_{r,s} c_r c_s \int \int_{\mathbb{R}^2} \left(\frac{\frac{1}{x^2 + \eta_r} - \frac{1}{y^2 + \eta_r}}{x - y} \right) \left(\frac{\frac{1}{x^2 + \eta_s} - \frac{1}{y^2 + \eta_s}}{x - y} \right) dx dy.
$$

Each of the double integrals can then be rewritten in the following way

$$
\begin{split} \int\int_{\mathbb{R}^2}\Bigg&\left(\frac{\frac{1}{x^2+\eta_r}-\frac{1}{y^2+\eta_r}}{x-y}\right)\Bigg(\frac{\frac{1}{x^2+\eta_s}-\frac{1}{y^2+\eta_s}}{x-y}\Bigg)dxdy\\ &=2\int_{\mathbb{R}}\frac{x^2}{(x^2+\eta_r)(x^2+\eta_s)}dx\int_{\mathbb{R}}\frac{1}{(y^2+\eta_r)(y^2+\eta_s)}dy+2\left(\int_{\mathbb{R}}\frac{x}{(x^2+\eta_r)(x^2+\eta_s)}dx\right)^2. \end{split}
$$

Now we can apply the residual theorem to each of the three integrals above. For either $\text{Im}(\eta_r) > 0$ or Im(η_r) < 0, take the principle square root, we have Im($i(\eta_r)^{\frac{1}{2}}$) > 0. So each integral there are two residuals in the upper helf plane to consider i.e. $i(n)^{\frac{1}{2}}$ and $i(n)^{\frac{1}{2}}$. Upper in the upper half plane to consider i.e. $i(\eta_r)^{\frac{1}{2}}$ and $i(\eta_s)^{\frac{1}{2}}$. Hence,

$$
\frac{1}{2\pi i} \int_{\mathbb{R}} \frac{x^2}{(x^2 + \eta_r)(x^2 + \eta_s)} dx = \frac{1}{2i((\eta_r)^{\frac{1}{2}} + (\eta_s)^{\frac{1}{2}})},
$$
\n
$$
\frac{1}{2\pi i} \int_{\mathbb{R}} \frac{1}{(y^2 + \eta_r)(y^2 + \eta_s)} dy = \frac{1}{2i(\eta_r)^{\frac{1}{2}} (\eta_s)^{\frac{1}{2}} ((\eta_r)^{\frac{1}{2}} + (\eta_s)^{\frac{1}{2}})},
$$
\n
$$
\frac{1}{2\pi i} \int_{\mathbb{R}} \frac{x}{(x^2 + \eta_r)(x^2 + \eta_s)} dx = 0.
$$

Together we have,

$$
\frac{1}{(2\pi i)^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy = \sum_{r,s=1}^{2M} c_r c_s \frac{-1}{2(\eta_r)^{\frac{1}{2}} (\eta_s)^{\frac{1}{2}} (\eta_r)^{\frac{1}{2}} + (\eta_s)^{\frac{1}{2}} \right)^2}
$$

Combing [\(332\)](#page-58-1) and Lemma [5.2,](#page-34-0) we get

$$
\lim_{n \to \infty} \frac{1}{n^{2\alpha}} \operatorname{Tr}[T_+, T_-] = \frac{1}{8\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy.
$$
 (336)

□

7.2 Proof of Proposition [4.5](#page-27-2)

We are now ready to prove Proposition [4.5.](#page-27-2)

Proof of Proposition [4.5.](#page-27-2) Recall that we consider $f(x) = \text{Im} \sum_{r=1}^{M} d_r \frac{1}{x - \lambda_j} = \sum_{r=1}^{2M} \frac{c_r}{x - \lambda_r}$ $\frac{c_r}{x-\eta_r}$ with c_j, η_r defined as [\(49\)](#page-14-0). We also define $F := \sum_{r=1}^{2M} c_r \left(\mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}} \right)^{-1}$.

Proposition [4.2](#page-25-0) shows that the first *m* coefficients in the expansion around small *t* of below

$$
log \mathbb{E}[\exp\left(tX_{f,\alpha,x_0}-t\mathbb{E}[X_{f,\alpha,x_0}]\right)] = \log\left(\det\left(1+P_n\left(e^{tn^{-\alpha}F}-1\right)P_n\right)e^{-tn^{-\alpha}\operatorname{Tr}(P_nF)}\right) \tag{337}
$$

are asymptotically the same as those of the following

$$
\log \det \left(1 + P_n \left(e^{tn^{-\alpha} \sum_r c_r T_{n \pm 2mn\beta}(\eta_r)} - 1 \right) P_n \right) e^{-tn^{-\alpha} \operatorname{Tr} P_n \sum_r c_r T_{n \pm 2mn\beta}(\eta_r)} = \sum_{k \ge 2} \frac{t^k}{k!} C_k^{(n)} \left(\sum_r c_r n^{-\alpha} T_{n \pm 2mn\beta}(\eta_r) \right). \tag{338}
$$

By Lemmas [7.2,](#page-54-1) [7.4](#page-56-1) and [7.6,](#page-57-1) we see that, as $n \to \infty$

$$
\lim_{n \to \infty} \log \det \left(1 + P_n \left(e^{tn^{-\alpha} \sum_r c_r T_{n \pm 2mn} \beta(\eta_r)} - 1 \right) P_n \right) e^{-t \operatorname{Tr} P_n n^{-\alpha} \sum_r c_r T_{n \pm 2mn} \beta(\eta_r)} \\
= \frac{t^2}{16\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy. \tag{339}
$$

The consequence of Lemma [3.4](#page-17-0) implies that the expansions of both of the right-hand sides of [\(337\)](#page-61-0) and [\(338\)](#page-61-1) are absolutely convergent. We conclude that, via the dominated convergent theorem, for any 0 < $\frac{\alpha}{2}$ < β < $\frac{\alpha+1}{3}$ around the left edge, i.e. $x_0 = b_{n-1,n} - 2(a_{n,n}a_{n-1,n})^{\frac{1}{2}} + o(n^{-\alpha})$

$$
\lim_{n \to \infty} \log \mathbb{E} \left[\exp \left(t X_{f, \alpha, x_0}^{(n)} - t \mathbb{E} [X_{f, \alpha, x_0}^{(n)}] \right) \right] = \frac{t^2}{16\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(x^2) - f(y^2)}{x - y} \right)^2 dx dy.
$$
 (340)

At the right edge, i.e. $x_0 = b_{n-1,n} + 2\sqrt{a_{n,n}a_{n-1,n}} + o(n^{-\alpha})$, we consider $-f$. Hence, all the arguments of the left edge apply and we obtain

$$
\lim_{n \to \infty} \log \mathbb{E} \left[\exp \left(t X_{-f, \alpha, x_0}^{(n)} - t \mathbb{E} [X_{-f, \alpha, x_0}^{(n)}] \right) \right] = \frac{t^2}{16\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy.
$$
 (341)

Then by the symmetry of centred Gaussian distribution, we conclude that at the right edge

$$
\lim_{n \to \infty} \log \mathbb{E} \left[\exp \left(t X_{f,\alpha,x_0}^{(n)} - t \mathbb{E} [X_{f,\alpha,x_0}^{(n)}] \right) \right] = \frac{t^2}{16\pi^2} \int \int_{\mathbb{R}^2} \left(\frac{f(-x^2) - f(-y^2)}{x - y} \right)^2 dx dy. \tag{342}
$$

Then by the moments method we obtain $X_n(f_{\alpha,x_0}) - \mathbb{E}[X_n(f_{\alpha,x_0})]$ has the Gaussian fluctuations in the \blacksquare

Hence, together with Subsection [4.2](#page-27-3) and we complete the proof of Theorem [2.3.](#page-8-0)

8 Proof of Theorems [1.2](#page-4-0) [1.1](#page-4-1) [2.2](#page-8-1)

We will use Theorem [2.3](#page-8-0) to prove Theorems [1.1](#page-4-1) and [2.2.](#page-8-1) We will prove Theorem [1.2](#page-4-0) by Theorem [2.2.](#page-8-1)

Proof of Theorem [2.2](#page-8-1). For any $0 < \alpha < \frac{2}{3}$, condition [\(2.1\)](#page-7-0) implies that for all $j \in I_n^{(\alpha,\varepsilon)}$

$$
a_{j,n}a_{j-2,n} - a_{j-1,n}^2 = O(n^{-1}), \quad (b_{j-1,n} - x_0 - a_{j,n})a_{j-2,n} - (b_{j-2,n} - x_0 - a_{j-1,n})a_{j-1,n} = O(n^{-1}).
$$

Hence the conditions in Theorem [2.3](#page-8-0) are satisfied. Then apply Theorem [2.3](#page-8-0) and we conclude the proof of Theorem [2.2.](#page-8-1) \Box

Proof of Theorem [1.1](#page-4-1) . The asymptotics of recurrence coefficients of the modified Jacobi polynomials with measure [\(3\)](#page-3-0) are given by the Riemann-Hilbert method in [\[24\]](#page-81-1) to be

$$
a_j = \frac{1}{2} - \left(\frac{4\gamma_1^2 - 1}{32} + \frac{4\gamma_2^2 - 1}{32}\right) \frac{1}{j^2} + O(j^{-3}), \quad b_j = \frac{\gamma_2^2 - \gamma_1^2}{4j^2} + O(j^{-3}), \qquad \text{as } j \to \infty,
$$
 (343)

where γ_1 and γ_2 are some constants explicitly given in [\[24\]](#page-81-1). Note that there is no scaling in this model and, hence, we remove the second subscription *n* in the notations. Hence,

$$
|a_j - a_{j-1}| = O(j^{-3}),
$$
 $|b_j - b_{j-1}| = O(j^{-3}),$ as $j \to \infty.$

Then the conditions [\(28\)](#page-8-2) and [\(29\)](#page-8-3) are satisfied for all $0 < \frac{\alpha}{2} < \beta < 1$. Hence the limit of mesoscopic fluctuations holds for all $\alpha \in (0, 2)$. Then apply Theorem 2.3, take $\beta \to 1$ and we conclude the proof of fluctuations holds for all $\alpha \in (0, 2)$. Then apply Theorem [2.3,](#page-8-0) take $\beta \to 1$ and we conclude the proof of Theorem 1.1. Theorem [1.1.](#page-4-1) \square

Proof of Theorem [1.2.](#page-4-0) Consider $I = \{j \in \mathbb{N} : \frac{j}{n} \to 1, \text{ as } n \to \infty\}$. Then for all $j \in I_{n,m}^{(\beta)}$, let $c = \frac{j}{n}$ $\frac{f}{n}$ then we can rewrite

$$
e^{-nV(x)}dx = e^{-\frac{i}{c}V(x)}dx.
$$
 (344)

Then the conditions on *V* and positivity about *c* implies the existence and uniqueness of equilibrium measure of $c^{-1}V$, denoted by $\mu_{c^{-1}V}$ by the potential theory. By the convexity of $c^{-1}V$, the support of $\mu_{c^{-1}V}$ is a single
interval say $[\alpha_1, \alpha_2]$ which is determined by the following set of equations interval say $[\alpha_1, \alpha_2]$, which is determined by the following set of equations,

$$
\int_{\alpha_1}^{\alpha_2} \frac{V'(s)}{\sqrt{(s-\alpha_1)(\alpha_2-s)}} ds = 0, \quad \int_{\alpha_1}^{\alpha_2} \frac{sV'(s)}{\sqrt{(s-\alpha_1)(\alpha_2-s)}} ds = 2\pi c. \tag{345}
$$

In literature α_1, α_2 are called the Mhaskar-Rakhmanov-Saff numbers, (cf. [\[20\]](#page-81-2) pages 203-234) Let $F : \mathbb{R}^3 \to$ \mathbb{R}^2 be a real analytic function such that $F(x, y, c) := \begin{pmatrix} \phi(x, y) \\ \varphi(x, y, c) \end{pmatrix}$ $\varphi(x, y, c)$! where

$$
\phi(x, y) := \int_{-1}^{1} V' \left(\frac{(y - x)t + x + y}{2} \right) \frac{dt}{\sqrt{1 - t^2}},
$$

$$
\varphi(x, y, c) := \int_{-1}^{1} \frac{(y - x)t}{2} V' \left(\frac{(y - x)t + x + y}{2} \right) \frac{dt}{\sqrt{1 - t^2}} - 2\pi c.
$$

Then we have $F(\alpha_1, \alpha_2, c) = 0$ by [\(345\)](#page-62-0) and a change of variable argument. We further define

$$
d\mathcal{F}(t) \coloneqq V''\left(\frac{(\alpha_2 - \alpha_1)t + \alpha_1 + \alpha_2}{2}\right)\frac{dt}{\sqrt{1 - t^2}}.
$$

Then by convexity of *V*, we have $V'' > 0$ and thus $\mathcal{F}(t)$ defines a positive measure. Now computing the lacobian of *F* at (α, α, c) , we get Jacobian of *F* at (α_1, α_2, c) , we get

$$
D_{(x,y)}F(\alpha_1, \alpha_2, c) = \begin{pmatrix} \int_{-1}^{1} (1-t) d\mathcal{F}(t), & \int_{-1}^{1} (1+t) d\mathcal{F}(t) \\ -\frac{2\pi c}{\alpha_2 - \alpha_1} + (\alpha_2 - \alpha_1) \int_{-1}^{1} t (1-t) d\mathcal{F}(t), & \frac{2\pi c}{\alpha_2 - \alpha_1} + (\alpha_2 - \alpha_1) \int_{-1}^{1} t (1+t) d\mathcal{F}(t) \end{pmatrix}.
$$
 (346)

The determinant of [\(346\)](#page-63-0) can be computed as

$$
\frac{4\pi c}{\alpha_2 - \alpha_1} \int_{-1}^{1} d\mathcal{F}(t) + (\alpha_2 - \alpha_1) \left(\int_{-1}^{1} (1 - t) d\mathcal{F}(t) \int_{-1}^{1} t (1 + t) d\mathcal{F}(t) - \int_{-1}^{1} (1 + t) d\mathcal{F}(t) \int_{-1}^{1} t (1 - t) d\mathcal{F}(t) \right)
$$
\n
$$
= \frac{4\pi c}{\alpha_2 - \alpha_1} \int_{-1}^{1} d\mathcal{F}(t) + 2 (\alpha_2 - \alpha_1) \left(\int_{-1}^{1} d\mathcal{F}(t) \int_{-1}^{1} t^2 d\mathcal{F}(t) - \left(\int_{-1}^{1} t d\mathcal{F}(t) \right)^2 \right).
$$

Use Cauchy-Schwartz inequality and we deduce $\left(\int_{-1}^{1} t d\mathcal{F}(t)\right)^{2} \leq \int_{-1}^{1} t^{2} d\mathcal{F}(t) \int_{-1}^{1} d\mathcal{F}(t)$. Subsequently, the Jacobian matrix under consideration is strictly positive and hence non-singular. Then apply the Implicit Function Theorem to obtain that α_1 and α_2 are continuously differentiable functions of *c*, denoted by

$$
\alpha_1 = \alpha_1(c), \quad \alpha_2 = \alpha_2(c).
$$

Moreover, one can apply the same analysis in [\[21\]](#page-81-3) for $c^{-1}V$ and obtain a similar result in the Application 1 (1.64) (1.65) in [\[21\]](#page-81-3) for the regular case (also refer to the proof Theorem 2.1 in [\[38\]](#page-82-1)). Note that in their notation in the one-cut case $\tilde{M}^{(\infty)}(z)$ found as (4.71) of [\[21\]](#page-81-3) is reduced to

$$
\tilde{M}^{(\infty)}(z) = \frac{1}{2} \begin{pmatrix} \gamma(z) + \gamma(z)^{-1}, & i(\gamma(z)^{-1} - \gamma(z)) \\ i(\gamma(z) - \gamma(z)^{-1}), & \gamma(z) + \gamma(z)^{-1} \end{pmatrix},
$$

which is equivalent to *N*(*z*) defined in (6.13-6.16) of [\[38\]](#page-82-1) after rescaling the support from $[\alpha_1, \alpha_2]$ to $[-1, 1]$.
This means that the recurrence coefficients can be estimated as This means that the recurrence coefficients can be estimated as

$$
a_{j,n} = \frac{\alpha_2(c) - \alpha_1(c)}{4} + O(n^{-1}), \quad b_{j,n} = \frac{\alpha_2(c) + \alpha_1(c)}{2} + O(n^{-1}).
$$
 (347)

Recall that $c = \frac{j}{n}$ $\frac{d}{dx}$ and $\alpha_1(c)$, $\alpha_2(c)$ are continuously differentiable functions. Then for all $j \in I$ we have

$$
|a_{j,n} - a_{j-1,n}| = O(n^{-1}), \quad |b_{j,n} - b_{j-1,n}| = O(n^{-1}).
$$

Moreover, by the normalisation we have $\alpha_1(1) = -1$, $\alpha_2(1) = 1$. Hence,

$$
\lim_{j/n \to 1} a_{j,n} = \frac{1}{2}, \quad \lim_{j/n \to 1} b_{j,n} = 0.
$$

From here we obtain that the edges of the fluctuations are at $x_0 = -1$ or 1. Now we can employ Theorem [2.2](#page-8-1) and conclude Theorem [1.2.](#page-4-0) □

Note that the strict convexity is not a necessary condition for the Jacobian to be non-singular nor the equilibrium measure to be supported on a single interval. Though it can be relaxed, convexity simplifies the argument. Furthermore, analyticity is only needed for the Riemann-Hilbert technique to obtain the asymptotics of recurrence coefficients. However, it is highly possible that [\(347\)](#page-63-1) still holds for *V* having some degree of smoothness, without assuming analyticity. In the literature, the Riemann–Hilbert– $\bar{\partial}$ method is able to deal with V with two Lipschitz continuous derivatives, see [\[39\]](#page-82-2). Their result Theorem 1(1) indicates that the recurrence coefficients should be slowly varying with $O(n^{-\frac{1}{3}} \log n)$, in a more general setup. Here we believe that Theorem [1.2](#page-4-0) should still hold for *V* being a convex (or the equilibrium measure being supported in a single interval) and $C^{2+\epsilon}(\mathbb{R})$ (for some $\epsilon > 0$) potential satisfying [\(8\)](#page-4-2). This requires a more detailed study about the potential theory and we leave it open.

9 Proof of Theorem [2.1](#page-6-0)

Assumption [\(20\)](#page-7-1) allows us to to approximate the Jacobi matrix $\mathcal J$ by a Toeplitz matrix. Compared with Theorem [2.3,](#page-8-0) we have more control of the tails of the diagonals in this case. Hence, a direct comparison with a Toeplitz operator is possible by exploring the algebra of the cumulant formula. In this section we are to prove Theorem [2.1](#page-6-0) by fulfilling such idea.

For clarity, throughout this section we let

$$
\beta = \frac{\alpha + \varepsilon}{2}, \quad \text{for some } \varepsilon > 0 \quad \text{such that } 0 < \frac{\alpha}{2} < \beta < 1. \tag{348}
$$

As mentioned in the remark of Theorem [2.1,](#page-6-0) we will give a proof with a weaker assumption [\(20\)](#page-7-1). Without loss of generality we further assume

$$
\sup_{j\geq n-n^{\beta+\varepsilon/2}}|a_{j,n}-1|=O(n^{-2\beta}), \quad \sup_{j\geq n-n^{\beta+\varepsilon/2}}|b_{j,n}|=O(n^{-2\beta}).\tag{349}
$$

The general case follows by translating and scaling the point process by $(x - b)/a$, where $a := \lim_{n \to \infty} a_{n,n}$ and $b \coloneqq \lim_{n} b_{n-1,n}$.

9.1 Resolvent of a Toeplitz Operator

Let's first consider the Chebyshev polynomial (of the second kind) ensemble whose measure is given by

$$
d\tilde{\mu}(x) = \frac{\sqrt{4 - x^2}}{2\pi} dx, \quad \text{supported on } [-2, 2].
$$

Its recurrence coefficients are

$$
\tilde{a}_{j,n}=1,\quad \tilde{b}_{j,n}=0.
$$

Hence its Jacobi matrix is a (semi-finite) Toeplitz matrix, i.e, entries remains constants along diagonals. This is also called the free Jacobi operator,

$$
\widetilde{\mathcal{J}} := \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & \cdots \\ 1 & 0 & 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & 1 & 0 & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots \end{pmatrix}.
$$
 (350)

Take $x_0 = -2$ and define

$$
\omega_{+} := \frac{2 - \frac{\eta}{n^{\alpha}} + \left(\left(2 - \frac{\eta}{n^{\alpha}} \right)^{2} - 4 \right)^{\frac{1}{2}}}{2}, \quad \omega_{-} := \frac{2 - \frac{\eta}{n^{\alpha}} - \left(\left(2 - \frac{\eta}{n^{\alpha}} \right)^{2} - 4 \right)^{\frac{1}{2}}}{2},\tag{351}
$$

where the square root is taken at the principle branch so that $|\omega_{+}| > |\omega_{-}|$. The right edge (*x*₀ = 2) can be done in the exact same way by swapping the definitions of ω_+ and ω_- .

Since this $\tilde{\mathcal{J}}$ is a Toeplitz operator, we can use the Wiener-Hopf factorization to obtain the exact inverse formula entry-wise,

$$
\left(\left(\widetilde{\mathcal{J}} - \left(x_0 + \frac{\eta}{n^{\alpha}} \right) I d \right)^{-1} \right)_{j,k} = \frac{(-\omega_+)^{-|j-k|} - (-\omega_-)^{j+k}}{\omega_+ - \omega_-}.
$$
\n(352)

One should compare this formula with Proposition [4.3.](#page-26-0)

Note that

$$
\omega_{+} = 1 + \left(-\frac{\eta}{n^{\alpha}}\right)^{\frac{1}{2}} + O(n^{-\alpha}), \quad \omega_{-} = 1 - \left(-\frac{\eta}{n^{\alpha}}\right)^{\frac{1}{2}} + O(n^{-\alpha}).
$$
\n(353)

From [\(352\)](#page-64-0) and [\(353\)](#page-65-0), we see that for all j, k

$$
\left| \left| \left(\widetilde{\mathcal{J}} - \left(x_0 + \frac{\eta}{n^{\alpha}} \right) I d \right)^{-1} \right|_{j,k} \right| \le C n^{\frac{\alpha}{2}} e^{-d n^{-\frac{\alpha}{2}} |j - k|}.
$$
\n(354)

One convenient consequence of this estimate is the following lemma.

Lemma 9.1. *There exist a constant* $C_0 > 0$ *, such that for any* $m_1, m_2 \in \mathbb{N}$ *, we have*

$$
\left\|P_{m_1}Q_{m_2}\left(\widetilde{\mathcal{J}}-\left(x_0+\frac{\eta}{n^{\alpha}}\right)Id\right)^{-1}\right\|_2 \leq \sqrt{C_0n^{\frac{3\alpha}{2}}|m_1-m_2|}.\tag{355}
$$

Proof. For $m_1 \le m_2$, $P_{m_1}Q_{m_2} = 0$ and the norm on left-hand side is zero. The inequality holds trivially. We now consider the case where $m_1 > m_2$.

Recall for any linear Hilbert-Schmidt operator *A*, the Hilbert-Schmidt norm $||A||_2^2 = \sum_{j,k} |(A)_{j,k}|^2$. Then we can estimate by [\(354\)](#page-65-1)

$$
\left\| P_{m_1} Q_{m_2} \left(\widetilde{\mathcal{J}} - \left(x_0 + \frac{\eta}{n^{\alpha}} \right) I d \right)^{-1} \right\|_2^2 \le \sum_{j=m_2+1}^{m_1} \sum_{k=1}^{\infty} C^2 n^{\alpha} e^{-2dn^{-\frac{\alpha}{2}} |j-k|} \\ \le \sum_{j=m_2+1}^{m_1} \sum_{l \in \mathbb{Z}} C^2 n^{\alpha} e^{-2dn^{-\frac{\alpha}{2}} |l|} \le 2 \sum_{j=m_2+1}^{m_1} \sum_{l=0}^{\infty} C^2 n^{\alpha} e^{-2dn^{-\frac{\alpha}{2}} l} = \frac{2C^2 n^{\alpha} (m_1 - m_2)}{1 - e^{-2dn^{-\frac{\alpha}{2}}}}, \quad (356)
$$

where the second line is by adding more terms in the sum to make the series to be a power series. Note that 1 − e^{-x} ≥ $e^{-1}x$ for all $x \in [0, 1]$, we can further estimate [\(356\)](#page-65-2) to be

$$
\left\| P_{m_1} Q_{m_2} \left(\widetilde{\mathcal{J}} - \left(x_0 + \frac{\eta}{n^{\alpha}} \right) I d \right)^{-1} \right\|_2^2 \leq 4e^{2} d n^{\frac{3\alpha}{2}} (m_1 - m_2). \tag{357}
$$

This completes the proof. \Box

9.2 Lemmas of Proof of Theorem [2.1](#page-6-0)

We consider test function $f(x) = \sum_{r=1}^{2M} c_r \frac{1}{x-\eta_r}$ as described in [\(49\)](#page-14-0) for some $\eta \in \{x + iy | x \in \mathbb{R}, y \neq 0\}$. Let us define define

$$
\widetilde{F} \coloneqq \sum_{r=1}^{2M} c_r \left(\widetilde{\mathcal{J}} - x_0 - \frac{\eta_r}{n^{\alpha}} \right)^{-1} . \tag{358}
$$

Denote the Jacobi matrix of the OPE with recurrence coefficients with $\{a_{j,n}, b_{j,n}\}$ satisfies [\(349\)](#page-64-1) to be \mathcal{J} . Define

$$
F := \sum_{r=1}^{2M} c_r \left(\mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}} \right)^{-1} . \tag{359}
$$

Lemma 9.2. *Let* $0 < \frac{\alpha}{2} < \beta < 1$ *. Let* $l \in \mathbb{N}$ *. Let* $m_1 \in \mathbb{N}$ *such that* $m_1 > ln^{\beta}$ *. Then there exists a* $C_l > 0$ *such* that *that*

$$
\left\|Q_{m_1+ln\beta}\widetilde{F}^1P_{m_1}\right\|_1 \le C_l e^{-d'n^{\beta-\frac{\alpha}{2}}},\tag{360}
$$

$$
\left\|P_{m_1 - \ln^{\beta}} \widetilde{F}^1 Q_{m_1}\right\|_1 \le C_l e^{-d' n^{\beta - \frac{\alpha}{2}}}
$$
\n(361)

Proof. Repeat the proof of [\(187\)](#page-37-3) in Lemma [5.4](#page-37-5) taking $N = \infty$, using [\(354\)](#page-65-1) and one gets [\(360\)](#page-66-0). Repeat the proof of [\(203\)](#page-39-6) in Lemma [5.5](#page-39-5) summing *j* from 1 and *k* up to ∞ , using [\(354\)](#page-65-1) and one gets [\(361\)](#page-66-1). □

Lemma 9.3. *Let* $0 < \frac{\alpha}{2} < \beta < 1$. Assume [\(349\)](#page-64-1) *holds for all j* > $m_2 - 2n^{\beta}$. For any $m_1, m_2 \in \mathbb{N}$ such that $2n^{\beta} < m_2 < m_1$ and $m_1 - m_2 = O(n^{\beta})$ we have as $n \to \infty$. $2n^{\beta} < m_2 < m_1$ and $m_1 - m_2 = O(n^{\beta})$, we have, as $n \to \infty$

$$
\left\| P_{m_1} \left(\widetilde{F} - F \right) Q_{m_2} \right\|_1 = O \left(n^{\frac{3\alpha}{2} - \beta} \right). \tag{362}
$$

Proof. Let us define

$$
\widetilde{F}^{(r)} := \left(\widetilde{\mathcal{J}} - x_0 - \frac{\eta_r}{n^{\alpha}}\right)^{-1} \tag{363}
$$

$$
F^{(r)} \coloneqq \left(\mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}}\right)^{-1}.\tag{364}
$$

By triangle inequality we have

$$
\left\|P_{m_1}\left(\widetilde{F}-F\right)Q_{m_2}\right\|_1 \leq \sum_r |c_r| \left\|P_{m_1}\left(\widetilde{F}^{(r)}-F^{(r)}\right)Q_{m_2}\right\|_1. \tag{365}
$$

Now we are going to study the trace norm for any *r*. For a cleaner notation, let

$$
\mathcal{A} \equiv \widetilde{\mathcal{J}} - x_0 - \frac{\eta_r}{n^{\alpha}}, \quad \mathcal{B} \equiv \mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}}.
$$

It is sufficient to estimate $||P_{m_1}(\mathcal{A}^{-1} - \mathcal{B}^{-1}) Q_{m_2}||_1$. Note that estimates about \mathcal{A}^{-1} are discussed in Section [9.1.](#page-64-2)

We use resolvent identity and rewrite $\mathcal{B}^{-1} = \mathcal{A}^{-1} + (\mathcal{B}^{-1} - \mathcal{A}^{-1})$ to get

$$
P_{m_1}(\mathcal{A}^{-1} - \mathcal{B}^{-1}) Q_{m_2} = P_{m_1} \mathcal{B}^{-1} (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} Q_{m_2}
$$

= $P_{m_1} \mathcal{A}^{-1} (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} Q_{m_2} + P_{m_1} (\mathcal{B}^{-1} - \mathcal{A}^{-1}) (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} Q_{m_2}.$ (366)

Note that, by the fact that $Id = P_{m_2} + Q_{m_2}$ the second summand on the right-hand side can be written as

$$
P_{m_1}\left(\mathcal{B}^{-1}-\mathcal{A}^{-1}\right)P_{m_2}\left(\mathcal{B}-\mathcal{A}\right)\mathcal{A}^{-1}Q_{m_2}+P_{m_1}\left(\mathcal{B}^{-1}-\mathcal{A}^{-1}\right)Q_{m_2}\left(\mathcal{B}-\mathcal{A}\right)\mathcal{A}^{-1}Q_{m_2}.
$$

Rearrange the formula [\(366\)](#page-66-2) and we get,

$$
P_{m_1}(\mathcal{A}^{-1} - \mathcal{B}^{-1}) Q_{m_2} (Id + (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} Q_{m_2})
$$

= $P_{m_1} \mathcal{A}^{-1} (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} Q_{m_2} + P_{m_1} (\mathcal{B}^{-1} - \mathcal{A}^{-1}) P_{m_2} (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} Q_{m_2}.$ (367)

Note that, by Lemma [9.2,](#page-66-3) we have $\|\mathcal{A}^{-1}Q_{m_2} - Q_{m_2-n\beta}\mathcal{A}^{-1}Q_{m_2}\|_1 = O(e^{-d'n^{\beta-\frac{\alpha}{2}}})$. Moreover, since $\mathcal{B} - \mathcal{A}$ is three diagonal, we have $(B - A)Q_{m_2-n} = Q_{m_2-n}P_{-1}(B - A)Q_{m_2-n}P$. Also by Lemma [9.2,](#page-66-3) we have

 $\|\mathcal{A}^{-1}Q_{m_2-n\beta-1}-Q_{m_2-2n\beta-1}\mathcal{A}^{-1}Q_{m_2-n\beta-1}\|_1 = O(e^{-d'n^{\beta-\frac{\alpha}{2}}})$. Also note that $\|\mathcal{A}^{-1}\|_{\infty} = O(n^{\alpha}), \|\mathcal{A}\|_{\infty} = O(1)$ and ∥B∥[∞] = *O*(1). Hence, the first summand on the right-hand side of [\(367\)](#page-66-4) can be estimated to be

$$
\|P_{m_1}\mathcal{A}^{-1}(\mathcal{B}-\mathcal{A})\mathcal{A}^{-1}Q_{m_2}\|_1
$$

= $\left\|P_{m_1}Q_{m_2-2n^{\beta}-1}\mathcal{A}^{-1}Q_{m_2-n^{\beta}-1}(\mathcal{B}-\mathcal{A})Q_{m_2-n^{\beta}}\mathcal{A}^{-1}Q_{m_2}\right\|_1 + O\left(n^{\alpha}e^{-d'n^{\beta-\frac{\alpha}{2}}}\right).$ (368)

Similarly, we commute P_{m_1} from left to right on the right-hand side of [\(368\)](#page-67-0) with exponentially small error. Precisely speaking, by Lemma [9.2,](#page-66-3) we have $||P_{m_1}Q_{m_2-2n\beta-1}\mathcal{A}^{-1}-P_{m_1}Q_{m_2-2n\beta-1}\mathcal{A}^{-1}P_{m_1+n\beta}||_1 =$ $O(e^{-d'n^{\beta-\frac{\alpha}{2}}})$. Since $B - A$ is three diagonal, we have $P_{m_1+n^{\beta}}(\mathcal{B} - A) = P_{m_1+n^{\beta}}(\mathcal{B} - A) P_{m_1+n^{\beta+1}}$. Also by Lemma [9.2,](#page-66-3) we have $||P_{m_1+n\beta+1}Q_{m_2-n\beta}A^{-1}-P_{m_1+n\beta+1}Q_{m_2-n\beta}A^{-1}P_{m_1+2n\beta+1}||_1 = O(e^{-d'n\beta-\frac{\alpha}{2}})$. Note that $\|\mathcal{A}^{-1}\|_{\infty} = O(n^{\alpha})$, $\|\mathcal{A}\|_{\infty} = O(1)$ and $\|\mathcal{B}\|_{\infty} = O(1)$. Also note that the projections are commutative, i.e., $P_{m_1}Q_{m_2} = Q_{m_2}P_{m_1}$. Hence, [\(368\)](#page-67-0) can be further estimated to be as $n \to \infty$

$$
||P_{m_1}\mathcal{A}^{-1}(\mathcal{B}-\mathcal{A})\mathcal{A}^{-1}Q_{m_2}||_1
$$

=
$$
||P_{m_1}Q_{m_2-2n\beta-1}\mathcal{A}^{-1}P_{m_1+n\beta}Q_{m_2-n\beta-1}(\mathcal{B}-\mathcal{A})P_{m_1+n\beta+1}Q_{m_2-n\beta}\mathcal{A}^{-1}P_{m_1+2n\beta+1}Q_{m_2}||_1 + O\left(n^{\alpha}e^{-d'n^{\beta-\frac{\alpha}{2}}}\right).
$$
 (369)

Now use the trace norm inequality $||ABC||_1 \leq ||A||_2||B||_{\infty}||C||_2$ and we get as $n \to \infty$

$$
\|P_{m_1}\mathcal{A}^{-1}(\mathcal{B}-\mathcal{A})\mathcal{A}^{-1}Q_{m_2}\|_1
$$

\n
$$
\leq \|P_{m_1}Q_{m_2-2n^{\beta}-1}\mathcal{A}^{-1}\|_2 \|P_{m_1+n^{\beta}}Q_{m_2-n^{\beta}-1}(\mathcal{B}-\mathcal{A})P_{m_1+n^{\beta}+1}Q_{m_2-n^{\beta}}\|_{\infty} \|\mathcal{A}^{-1}P_{m_1+2n^{\beta}+1}Q_{m_2}\|_2 + O\left(n^{\alpha}e^{-d'n^{\beta-\frac{\alpha}{2}}}\right).
$$
\n(370)

Recall that we assume $m_1 - m_2 = O(n^{\beta})$. By Lemma [9.1,](#page-65-3) we have as $n \to \infty$

$$
\left\|P_{m_1} Q_{m_2 - 2n^{\beta} - 1} \mathcal{A}^{-1}\right\|_2 = O\left(n^{\frac{3\alpha}{4} + \frac{\beta}{2}}\right)
$$
\n(371)

$$
\left\| \mathcal{A}^{-1} P_{m_1 + 2n^{\beta} + 1} Q_{m_2} \right\|_2 = O\left(n^{\frac{3\alpha}{4} + \frac{\beta}{2}} \right). \tag{372}
$$

Since [\(349\)](#page-64-1) holds for all $j > m_2 - 2n^{\beta}$ by assumption, we have as $n \to \infty$

$$
||P_{m_1+n^\beta} Q_{m_2-n^\beta-1} (\mathcal{B} - \mathcal{A}) P_{m_1+n^\beta+1} Q_{m_2-n^\beta}||_{\infty} = O(n^{-2\beta}).
$$
\n(373)

Plug [\(371\)](#page-67-1), [\(372\)](#page-67-2) and [\(373\)](#page-67-3) into [\(370\)](#page-67-4) to get as *n* → ∞

$$
\left\|P_{m_1}\mathcal{A}^{-1}\left(\mathcal{B}-\mathcal{A}\right)\mathcal{A}^{-1}Q_{m_2}\right\|_1 = O\left(n^{\frac{3\alpha}{2}-\beta}\right).
$$
\n(374)

For the second summand of the right-hand side of (367) , we first use the resolvent identity and then continue with the same argument as [\(369\)](#page-67-5) to commute Q_{m_2} from right to left to obtain that as $n \to \infty$

$$
\|P_{m_1}\left(\mathcal{B}^{-1}-\mathcal{A}^{-1}\right)P_{m_2}\left(\mathcal{B}-\mathcal{A}\right)\mathcal{A}^{-1}Q_{m_2}\Big\|_1 = \|P_{m_1}\mathcal{B}^{-1}\left(\mathcal{A}-\mathcal{B}\right)\mathcal{A}^{-1}P_{m_2}\left(\mathcal{B}-\mathcal{A}\right)\mathcal{A}^{-1}Q_{m_2}\Big\|_1
$$

\n
$$
= \left\|P_{m_1}\mathcal{B}^{-1}\left(\mathcal{A}-\mathcal{B}\right)Q_{m_2-2n\beta-1}\mathcal{A}^{-1}P_{m_2}Q_{m_2-n\beta-1}\left(\mathcal{B}-\mathcal{A}\right)P_{m_2+1}Q_{m_2-n\beta}\mathcal{A}^{-1}Q_{m_2}\right\|_1 + O\left(n^{2\alpha}e^{-d'n^{\beta-\frac{\alpha}{2}}}\right).
$$
\n(375)

Use the trace norm inequality $||ABCD||_1 \leq ||A||_{\infty} ||B||_2 ||C||_{\infty} ||D||_2$ to get

$$
\|P_{m_1}\mathcal{B}^{-1}(\mathcal{A}-\mathcal{B})\mathcal{Q}_{m_2-2n^{\beta}-1}\mathcal{A}^{-1}P_{m_2}\mathcal{Q}_{m_2-n^{\beta}-1}(\mathcal{B}-\mathcal{A})P_{m_2+1}\mathcal{Q}_{m_2-n^{\beta}}\mathcal{A}^{-1}\mathcal{Q}_{m_2}\|_1
$$

\n
$$
\leq \|P_{m_1}\mathcal{B}^{-1}(\mathcal{A}-\mathcal{B})\mathcal{Q}_{m_2-2n^{\beta}-1}\|_{\infty}\|\mathcal{A}^{-1}P_{m_2}\mathcal{Q}_{m_2-n^{\beta}-1}\|_2\|P_{m_2}\mathcal{Q}_{m_2-n^{\beta}-1}(\mathcal{B}-\mathcal{A})\|_{\infty}\|P_{m_2+1}\mathcal{Q}_{m_2-n^{\beta}}\mathcal{A}^{-1}\mathcal{Q}_{m_2}\|_2
$$
\n(376)

In [\(376\)](#page-68-0), two operator norms are of order $O(n^{\alpha-2\beta})$ and $O(n^{-2\beta})$ respectively by the assumption that [\(349\)](#page-64-1) holds for all $j > m_2 - 2n^{\beta}$; two Hilbert-Schmidt norms are both of order $O(n^{\frac{3\alpha}{4} + \frac{\beta}{2}})$ by Lemma [9.1.](#page-65-3) Hence, as $n \to \infty$

$$
\left\|P_{m_1}\mathcal{B}^{-1}\left(\mathcal{A}-\mathcal{B}\right)Q_{m_2-2n^{\beta}-1}\mathcal{A}^{-1}P_{m_2}Q_{m_2-n^{\beta}-1}\left(\mathcal{B}-\mathcal{A}\right)P_{m_2+1}Q_{m_2-n^{\beta}}\mathcal{A}^{-1}Q_{m_2}\right\|_1 = O\left(n^{\frac{5\alpha}{2}-3\beta}\right).
$$
 (377)

Plug [\(377\)](#page-68-1) into [\(375\)](#page-67-6) to get

$$
\left\| P_{m_1} \left(\mathcal{B}^{-1} - \mathcal{A}^{-1} \right) P_{m_2} \left(\mathcal{B} - \mathcal{A} \right) \mathcal{A}^{-1} Q_{m_2} \right\|_1 = O \left(n^{\frac{5\alpha}{2} - 3\beta} \right).
$$
 (378)

We now use [\(374\)](#page-67-7) and [\(378\)](#page-68-2) to estimate the trace norm of [\(367\)](#page-66-4), and we have as $n \to \infty$

$$
\left\|P_{m_1}\left(\mathcal{A}^{-1}-\mathcal{B}^{-1}\right)Q_{m_2}\left(Id + \left(\mathcal{B}-\mathcal{A}\right)\mathcal{A}^{-1}Q_{m_2}\right)\right\|_1 = O\left(n^{\frac{3\alpha}{2}-\beta}\right).
$$
\n(379)

Note that we use the assumption $0 < \frac{\alpha}{2} < \beta < 1$ to determine $\frac{3\alpha}{2} - \beta > \frac{5\alpha}{2} - 3\beta$.

By Lemma [9.2,](#page-66-3)

$$
\left\| \left(\mathcal{B} - \mathcal{A}\right) \mathcal{A}^{-1} \mathcal{Q}_{m_2} \right\|_{\infty} = \left\| \left(\mathcal{B} - \mathcal{A}\right) \mathcal{Q}_{m_2 - n^{\beta}} \mathcal{A}^{-1} \mathcal{Q}_{m_2} \right\|_{\infty} + O\left(e^{-d' n^{\beta - \frac{\alpha}{2}}}\right).
$$
 (380)

Using the operator norm inequality we have

$$
\left\| \left(\mathcal{B} - \mathcal{A}\right) Q_{m_2 - n^{\beta}} \mathcal{A}^{-1} Q_{m_2} \right\|_{\infty} \le \left\| \left(\mathcal{B} - \mathcal{A}\right) Q_{m_2 - n^{\beta}} \right\|_{\infty} \left\| \mathcal{A}^{-1} Q_{m_2} \right\|_{\infty} . \tag{381}
$$

By the assumption that [\(349\)](#page-64-1) holds for all $j > m_2 - 2n^{\beta}$, we have as $n \to \infty \left\| (\mathcal{B} - \mathcal{A}) Q_{m_2 - n^{\beta}} \right\|_{\infty} = O(n^{-2\beta})$. Recall that $\|\mathcal{A}^{-1}\|_{\infty} = O(n^{\alpha})$. Now we have

$$
\left\| \left(\mathcal{B} - \mathcal{A}\right) \mathcal{A}^{-1} \mathcal{Q}_{m_2} \right\|_{\infty} = O\left(n^{\alpha - 2\beta}\right). \tag{382}
$$

Recall $0 < \frac{\alpha}{2} < \beta$, and hence [\(382\)](#page-68-3) is of order *o*(1). Then for large *n*,

$$
\left\| \left(Id + (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} \mathcal{Q}_{m_2} \right)^{-1} \right\|_{\infty} \le \frac{1}{1 - \left\| (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} \mathcal{Q}_{m_2} \right\|_{\infty}}.
$$
\n(383)

Moreover, by trace norm inequality $||AB||_1 \leq ||A||_1 ||B||_{\infty}$, we have

$$
\|P_{m_1}(\mathcal{A}^{-1} - \mathcal{B}^{-1}) Q_{m_2}\|_1
$$

\$\leq \|P_{m_1}(\mathcal{A}^{-1} - \mathcal{B}^{-1}) Q_{m_2} (Id + (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} Q_{m_2})\|_1 \| (Id + (\mathcal{B} - \mathcal{A}) \mathcal{A}^{-1} Q_{m_2})^{-1}\|_{\infty}.\$ (384)

Plug[\(379\)](#page-68-4),[\(382\)](#page-68-3) and [\(383\)](#page-68-5) into [\(384\)](#page-68-6) to obtain as $n \to \infty$

$$
\left\|P_{m_1}\left(\mathcal{A}^{-1}-\mathcal{B}^{-1}\right)Q_{m_2}\right\|_1 = O\left(n^{\frac{3\alpha}{2}-\beta}\right). \tag{385}
$$

Plug this estimate back into [\(365\)](#page-66-5) and we conclude as $n \to \infty$

$$
\left\| P_{m_1} \left(\widetilde{F} - F \right) Q_{m_2} \right\|_1 = O \left(n^{\frac{3\alpha}{2} - \beta} \right). \tag{386}
$$

Lemma above also implies the following by induction.

Lemma 9.4. *Let* $0 < \frac{\alpha}{2} < \beta < 1$. *Let* $m \in \mathbb{N}$. *Assume* [\(349\)](#page-64-1) *holds for all* $j > n - 2n^{\beta}$. *We have, for any* $k = 0, 1, 2, \ldots$ $k = 0, 1, 2, \ldots, m$

$$
\left\| P_{n+mn\beta} \left(\widetilde{F} - F \right) F^k Q_n \right\|_1 = O\left(n^{k\alpha + \frac{3\alpha}{2} - \beta} \right). \tag{387}
$$

Proof. First note that the following operator norms are bounded

$$
\sup_{n>0} \{ n^{-\alpha} ||\widetilde{F}||_{\infty}, n^{-\alpha} ||F||_{\infty} \} < \infty. \tag{388}
$$

Take $m_1 \equiv n + mn^{\beta}$ and $m_2 \equiv n$ to shorten the notation.

Consider the induction argument. For $k = 0$, [\(387\)](#page-69-0) is reduced to Lemma [9.3.](#page-66-6) For any $k \ge 0$, we use triangle inequality to obtain

$$
\left\| P_{m_1} \left(\widetilde{F} - F \right) F^{k+1} Q_{m_2} \right\|_1 \le \left\| P_{m_1} \left(\widetilde{F} - F \right) \widetilde{F}^{k+1} Q_{m_2} \right\|_1 + \left\| P_{m_1} \left(\widetilde{F} - F \right) \left(F^{k+1} - \widetilde{F}^{k+1} \right) Q_{m_2} \right\|_1. \tag{389}
$$

Use Lemma [9.2](#page-66-3) and we estimate the first summand on the right-hand side of [\(389\)](#page-69-1) to be, as $n \to \infty$,

$$
\left\|P_{m_1}\left(\widetilde{F}-F\right)\widetilde{F}^{k+1}Q_{m_2}\right\|_1=\left\|P_{m_1}\left(\widetilde{F}-F\right)Q_{m_2-(k+1)n^\beta}\widetilde{F}^{k+1}Q_{m_2}\right\|_1+O\left(n^\alpha e^{-d'n^{\beta-\frac{\alpha}{2}}}\right).
$$
 (390)

By the trace norm inequality $||AB||_1 \le ||A||_1 ||B||_{\infty}$, we have

$$
\left\| P_{m_1} \left(\widetilde{F} - F \right) Q_{m_2 - (k+1)n^{\beta}} \widetilde{F}^{k+1} Q_{m_2} \right\|_1 \le \left\| P_{m_1} \left(\widetilde{F} - F \right) Q_{m_2 - (k+1)n^{\beta}} \right\|_1 \left\| \widetilde{F}^{k+1} Q_{m_2} \right\|_{\infty} . \tag{391}
$$

Then use Lemma [9.3](#page-66-6) to estimate the trace norm above to be of order $O(n^{3\alpha/2-\beta})$. By [\(388\)](#page-69-2), the operator norm is of order $O(n^{(k+1)\alpha})$. Plug the estimate of [\(391\)](#page-69-3) back to [\(390\)](#page-69-4) to obtain, as $n \to \infty$

$$
\left\| P_{m_1}\left(\widetilde{F} - F\right) \widetilde{F}^{k+1} Q_{m_2} \right\|_1 = O\left(n^{(k+1)\alpha + \frac{3\alpha}{2} - \beta} \right). \tag{392}
$$

For the second summand on the right-hand side of [\(389\)](#page-69-1) we use the telescopic sum,

$$
F^{k+1} - \widetilde{F}^{k+1} = \sum_{l=0}^{k} F^{l} \left(F - \widetilde{F} \right) \widetilde{F}^{k-l},
$$

the fact that $Q_{m_2} + P_{m_2} = Id$, and triangle inequality to obtain

$$
\|P_{m_1}(\widetilde{F} - F)(F^{k+1} - \widetilde{F}^{k+1})Q_{m_2}\|_1 \leq \sum_{l=0}^k \left\|P_{m_1}(\widetilde{F} - F)F^{l}(Q_{m_2} + P_{m_2})(F - \widetilde{F})\widetilde{F}^{k-l}Q_{m_2}\right\|_1
$$

$$
\leq \sum_{l=0}^k \left(\left\|P_{m_1}(\widetilde{F} - F)F^{l}Q_{m_2}(F - \widetilde{F})\widetilde{F}^{k-l}Q_{m_2}\right\|_1 + \left\|P_{m_1}(\widetilde{F} - F)F^{l}P_{m_2}(F - \widetilde{F})\widetilde{F}^{k-l}Q_{m_2}\right\|_1\right). \quad (393)
$$

Use the trace norm inequality $||ABC||_1 \le ||A||_1 ||B||_{\infty} ||C||_{\infty}$ to obtain an estimate of the first term on the righthand side of [\(393\)](#page-69-5),

$$
\left\|P_{m_1}\left(\widetilde{F}-F\right)F^{l}Q_{m_2}\left(F-\widetilde{F}\right)\widetilde{F}^{k-l}Q_{m_2}\right\|_1 \leq \left\|P_{m_1}\left(\widetilde{F}-F\right)F^{l}Q_{m_2}\right\|_1 \left\|Q_{m_2}\left(F-\widetilde{F}\right)\right\|_{\infty} \left\|\widetilde{F}^{k-l}Q_{m_2}\right\|_{\infty}.\tag{394}
$$

The induction hypothesis for all $l = 0, \ldots, k$ implies that the first term on the right-hand side of [\(394\)](#page-70-0) to be

$$
\left\| P_{m_1} \left(\widetilde{F} - F \right) F^l Q_{m_2} \right\|_1 = O \left(n^{l \alpha + \frac{3\alpha}{2} - \beta} \right). \tag{395}
$$

For the second term on the right-hand side of [\(394\)](#page-70-0), let us define, Let us define

$$
\widetilde{F}^{(r)} := \left(\widetilde{\mathcal{J}} - x_0 - \frac{\eta_r}{n^{\alpha}}\right)^{-1} \tag{396}
$$

$$
F^{(r)} \coloneqq \left(\mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}}\right)^{-1}.\tag{397}
$$

Recall the definition of \tilde{F} and *F* in [\(358\)](#page-65-4) and [\(359\)](#page-65-5). We apply the triangle inequality and the resolvent identity to obtain

$$
\left\|Q_{m_2}\left(F-\widetilde{F}\right)\right\|_{\infty} \leq \sum_{r} |c_r| \left\|Q_{m_2}\left(F^{(r)}-\widetilde{F}^{(r)}\right)\right\|_{\infty} = \sum_{r} |c_r| \left\|Q_{m_2}\widetilde{F}^{(r)}\left(\widetilde{\mathcal{J}}-\mathcal{J}\right)F^{(r)}\right\|_{\infty}.
$$
 (398)

Use Lemma [9.2,](#page-66-3) and we get

$$
\left\| Q_{m_2} \widetilde{F}^{(r)} \left(\widetilde{\mathcal{J}} - \mathcal{J} \right) F^{(r)} \right\|_{\infty} = \left\| Q_{m_2} \widetilde{F}^{(r)} Q_{m_2 - n^{\beta}} \left(\widetilde{\mathcal{J}} - \mathcal{J} \right) F^{(r)} \right\|_{\infty} + O\left(n^{\alpha} e^{-d' n^{\beta - \frac{\alpha}{2}}} \right). \tag{399}
$$

By the operator norm inequality $||AB||_{\infty} \le ||A||_{\infty} ||B||_{\infty}$ we obtain,

$$
\left\| Q_{m_2} \widetilde{F}^{(r)} Q_{m_2 - n^{\beta}} \left(\widetilde{\mathcal{J}} - \mathcal{J} \right) F^{(r)} \right\|_{\infty} \leq \left\| Q_{m_2} \widetilde{F}^{(r)} \right\|_{\infty} \left\| Q_{m_2 - n^{\beta}} \left(\widetilde{\mathcal{J}} - \mathcal{J} \right) \right\|_{\infty} \left\| F^{(r)} \right\|_{\infty} . \tag{400}
$$

By the assumption that [\(349\)](#page-64-1) holds for all $j > m_2 - 2n^{\beta}$, we get $||Q_{m_2-n^{\beta}}(\overline{\mathcal{J}} - \mathcal{J})||_{\infty} = O(n^{-2\beta})$. Together \mathbf{H} with the fact that $||F^{(r)}||_{\infty} = O(n^{\alpha})$ and $||\overline{F}^{(r)}||_{\infty} = O(n^{\alpha})$, we estimate [\(400\)](#page-70-1) to be of order $O(n^{2\alpha-2\beta})$. Plugging it into [\(399\)](#page-70-2) and further back into [\(398\)](#page-70-3), we get

$$
\left\| Q_{m_2} \left(F - \widetilde{F} \right) \right\|_{\infty} = O \left(n^{2\alpha - 2\beta} \right). \tag{401}
$$

The last term on the right-hand side of [\(394\)](#page-70-0) is estimated to be $\|\widetilde{F}^{k-l}Q_{m_2}\|_{\infty} = O(n^{(k-l)\alpha})$. Hence, together with (395) and (401) , (394) is estimated to be

$$
\left\|P_{m_1}\left(\widetilde{F}-F\right)F^{l}Q_{m_2}\left(F-\widetilde{F}\right)\widetilde{F}^{k-l}Q_{m_2}\right\|_1=O\left(n^{(k+2)\alpha+\frac{3\alpha}{2}-3\beta}\right).
$$
\n(402)

Now we turn to the second term on the right-hand side of [\(393\)](#page-69-5). Use Lemma [9.2,](#page-66-3)

$$
\|P_{m_1}(\widetilde{F} - F) F^l P_{m_2} (F - \widetilde{F}) \widetilde{F}^{k-l} Q_{m_2}\|_1
$$

=
$$
\left\| P_{m_1}(\widetilde{F} - F) F^l P_{m_2} (F - \widetilde{F}) Q_{m_2 - (k-l)n} \widetilde{F}^{k-l} Q_{m_2}\right\|_1 + O\left(n^{(l+2)\alpha} e^{-d'n^{\beta - \frac{\alpha}{2}}}\right).
$$
 (403)

Use the operator norm inequality $||ABC||_1 \leq ||A||_{\infty} ||B||_1 ||C||_{\infty}$ to obtain

$$
\|P_{m_1}(\widetilde{F} - F) F^l P_{m_2} (F - \widetilde{F}) Q_{m_2 - (k-l)n\beta} \widetilde{F}^{k-l} Q_{m_2} \|_1
$$

\$\leq\$
$$
\left\| P_{m_1}(\widetilde{F} - F) F^l \right\|_{\infty} \left\| P_{m_2} (F - \widetilde{F}) Q_{m_2 - (k-l)n\beta} \right\|_1 \left\| \widetilde{F}^{k-l} Q_{m_2} \right\|_{\infty}.
$$
 (404)

The operator norms are estimated to be $||P_{m_1}(\overline{F} - F)F^l||_{\infty} = O(n^{(l+1)\alpha})$ and $||\overline{F}^{k-l}Q_{m_2}||_{\infty} = O(n^{(k-l)\alpha})$. The trace norm can be estimated by Lemma [9.3](#page-66-6) to be $\left\| P_{m_2} (F - \widetilde{F}) Q_{m_2 - (k-l)n^{\beta}} \right\|$ estimates into [\(404\)](#page-71-0). Then we get an estimate of [\(403\)](#page-70-6) to be $\|_1 = O\left(n^{\frac{3\alpha}{2}-\beta}\right)$. Plug these three

$$
\left\| P_{m_1}\left(\widetilde{F} - F\right) F^l P_{m_2}\left(F - \widetilde{F}\right) \widetilde{F}^{k-l} Q_{m_2} \right\|_1 = O\left(n^{(k+1)\alpha + \frac{3\alpha}{2} - \beta}\right).
$$
\n(405)

Plug [\(402\)](#page-70-7) and [\(405\)](#page-71-1) back into [\(393\)](#page-69-5), and we get an estimate of the second summand on the right-hand side of [\(389\)](#page-69-1) to be

$$
\left\|P_{m_1}\left(\widetilde{F}-F\right)\left(F^{k+1}-\widetilde{F}^{k+1}\right)Q_{m_2}\right\|_1 = O\left(n^{(k+1)\alpha+\frac{3\alpha}{2}-\beta}\right).
$$
\n(406)

Here we use the assumption $0 < \frac{\alpha}{2} < \beta < 1$ to deduce $O(n^{(k+2)\alpha + \frac{3\alpha}{2} - 3\beta}) = o(n^{(k+1)\alpha + \frac{3\alpha}{2} - \beta})$.
Plugging (392) and (406) into (389), we obtain

Plugging [\(392\)](#page-69-6) and [\(406\)](#page-71-2) into (389) , we obtain

$$
\left\| P_{m_1} \left(\widetilde{F} - F \right) F^{k+1} Q_{m_2} \right\|_1 = O \left(n^{(k+1)\alpha + \frac{3\alpha}{2} - \beta} \right). \tag{407}
$$

This concludes the induction argument. □

9.3 Proof of Theorem [2.1](#page-6-0)

The essential step of Theorem [2.1](#page-6-0) is the following proposition. Recall that $C_m^{(n)}(\mathcal{A})$ is the *m*-th cumulant for a linear operator A , which has the formula [\(67\)](#page-16-0).

Proposition 9.1. *Assume that the assumptions in Theorem [2.1](#page-6-0) are satisfied. Then we have*

$$
\lim_{n \to \infty} n^{-ma} \left| C_m^{(n)}(\widetilde{F}) - C_m^{(n)}(F) \right| = 0.
$$
\n(408)

Proof. First note that the following operator norms are bounded

$$
\sup_{n>0} \{ n^{-\alpha} ||\widetilde{F}||_{\infty}, n^{-\alpha} ||F||_{\infty} \} =: C_{op} < \infty.
$$
\n(409)

Since both sums are finite in the cumulant formula [\(67\)](#page-16-0), it is sufficient to show that

$$
\operatorname{Tr}(\widetilde{F})^{l_1} P_n \dots (\widetilde{F})^{l_j} P_n - \operatorname{Tr}(\widetilde{F}^m P_n) - \operatorname{Tr}(F)^{l_1} P_n \dots (F)^{l_j} P_n + \operatorname{Tr}(F^m P_n) = o(n^{m\alpha})
$$
(410)

Similar to [\(214\)](#page-42-1), using the telescoping sum twice and cyclic property of the trace operator, we can rewrite

the left-hand side of [\(410\)](#page-71-0) to be

$$
\sum_{k=2}^{j} Tr \left(\widetilde{F}^{l_1 + \dots + l_{k-1}} Q_n \widetilde{F}^{l_k} P_n \dots P_n \widetilde{F}^{l_j} P_n \right) - Tr \left(F^{l_1 + \dots + l_{k-1}} Q_n F^{l_k} P_n \dots P_n F^{l_j} P_n \right)
$$
\n
$$
= \sum_{k=2}^{j} Tr \left(\widetilde{F}^{l_k} P_n \dots P_n \widetilde{F}^{l_j} P_n \widetilde{F}^{l_1 + \dots + l_{k-1}} Q_n \right) - Tr \left(F^{l_k} P_n \dots P_n F^{l_j} P_n F^{l_1 + \dots + l_{k-1}} Q_n \right)
$$
\n
$$
= - \sum_{k=2}^{j-1} \left(Tr \left((\widetilde{F}^{l_k} - F^{l_k}) P_n \widetilde{F}^{l_{k+1}} \dots P_n \widetilde{F}^{l_j} P_n \widetilde{F}^{l_1 + \dots + l_{k-1}} Q_n \right) \right)
$$
\n
$$
+ \sum_{i=k}^{j-1} Tr \left(F^{l_k} P_n \dots P_n F^{l_i} P_n (\widetilde{F}^{l_{i+1}} - F^{l_{i+1}}) P_n \dots \widetilde{F}^{l_j} P_n \widetilde{F}^{l_1 + \dots + l_{k-1}} Q_n \right)
$$
\n
$$
+ Tr \left(F^{l_k} P_n \dots P_n F^{l_j} P_n (\widetilde{F}^{l_1 + \dots + l_{k-1}} - F^{l_1 + \dots + l_{k-1}}) Q_n \right)
$$

Then writing $\widetilde{F}^l - F^l = \sum_{k=0}^{l-1} F^{l-1-k} (\widetilde{F} - F) \widetilde{F}^k$, by Lemma [9.2,](#page-66-0) we commute Q_n from the right to left to get

$$
\left|\operatorname{Tr}\left((\widetilde{F}^{l_k}-F^{l_k})P_n\widetilde{F}^{l_{k+1}}\dots P_n\widetilde{F}^{l_j}P_n\widetilde{F}^{l_1+\dots+l_{k-1}}Q_n\right)\right|\leq l_kC_{op}^{m-1}n^{(m-1)\alpha}\|(\widetilde{F}-F)P_nQ_{n-mn^\beta}\|_1+R_n,\qquad(411)
$$

$$
\left| \text{Tr} \left(F^{l_k} P_n \dots P_n F^{l_i} P_n (\widetilde{F}^{l_{i+1}} - F^{l_{i+1}}) P_n \dots \widetilde{F}^{l_j} P_n \widetilde{F}^{l_1 + \dots + l_{k-1}} Q_n \right) \right|
$$

$$
\leq l_{i+1} C_{op}^{m-1} n^{(m-1)\alpha} ||(\widetilde{F} - F) P_n Q_{n-mn^{\beta}}||_1 + R_n, \quad (412)
$$

$$
\left| \text{Tr} \left(F^{l_k} P_n \dots P_n F^{l_j} P_n (\widetilde{F}^{l_1 + \dots + l_{k-1}} - F^{l_1 + \dots + l_{k-1}}) Q_n) \right| \right|
$$

$$
\leq C_{op}^{m - (l_1 + \dots + l_{k-1})} n^{(m - (l_1 + \dots + l_{k-1}))\alpha} ||P_n(\widetilde{F}^{l_1 + \dots + l_{k-1}} - F^{l_1 + \dots + l_{k-1}}) Q_n ||_1 + R_n, \quad (413)
$$

where $R_n = O(n^{m\alpha}e^{-d_0n^{\beta-\frac{\alpha}{2}}})$ is exponentially small. Define

$$
\widetilde{F}^{(r)} := \left(\widetilde{\mathcal{J}} - x_0 - \frac{\eta_r}{n^{\alpha}}\right)^{-1}, \quad F^{(r)} := \left(\mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}}\right)^{-1}.
$$
\n(414)

Use the triangle inequality to obtain

$$
\|(\widetilde{F} - F)P_n Q_{n-mn\beta}\|_1 \le \sum_{r=1}^{2M} |c_r| \left\| (\widetilde{F}^{(r)} - F^{(r)})P_n Q_{n-mn\beta} \right\|_1.
$$
 (415)

Recall that for any linear operator $\mathcal A$ and $\mathcal B$, we have the resolvent identity $\mathcal A^{-1} - \mathcal B^{-1} = \mathcal B^{-1}(\mathcal B - \mathcal A)\mathcal A^{-1}$. Further, we also have $\mathcal{B}^{-1} = (Id - \mathcal{B}^{-1}(\mathcal{B} - \mathcal{A})) \mathcal{A}^{-1}$. Combine these two formulas to get,

$$
\mathcal{A}^{-1} - \mathcal{B}^{-1} = (Id - \mathcal{B}^{-1}(\mathcal{B} - \mathcal{A}))\mathcal{A}^{-1}(\mathcal{B} - \mathcal{A})\mathcal{A}^{-1}.
$$

Take $\mathcal{A} = \widetilde{\mathcal{J}} - x_0 - \frac{\eta_r}{n^{\alpha}} = (\widetilde{F}^{(r)})^{-1}$ and $\mathcal{B} = \mathcal{J} - x_0 - \frac{\eta_r}{n^{\alpha}} = (F^{(r)})^{-1}$. Rewrite each summand of the right-hand side of [\(415\)](#page-72-0) to be

$$
\left\| (\widetilde{F}^{(r)} - F^{(r)}) P_n Q_{n-mn\beta} \right\|_1 = \left\| \left(Id - F^{(r)} \left(\mathcal{J} - \widetilde{\mathcal{J}} \right) \right) \widetilde{F}^{(r)} \left(\mathcal{J} - \widetilde{\mathcal{J}} \right) \widetilde{F}^{(r)} P_n Q_{n-mn\beta} \right\|_1. \tag{416}
$$

Recall that $\mathcal{J} - \widetilde{\mathcal{J}}$ is tri-diagonal and we have for any integer m_1

$$
\left(\mathcal{J}-\widetilde{\mathcal{J}}\right)P_{m_1}=P_{m_1+1}\left(\mathcal{J}-\widetilde{\mathcal{J}}\right)P_{m_1}.\tag{417}
$$

$$
\left(\mathcal{J}-\widetilde{\mathcal{J}}\right)Q_{m_1}=Q_{m_1-1}\left(\mathcal{J}-\widetilde{\mathcal{J}}\right)Q_{m_1}.
$$
\n(418)

Use Lemma [9.2](#page-66-0) to get that $\widetilde{F}^{(r)}$ commutes with the projection operator P_{m_1} and Q_{m_1} with an exponential small error. That is

$$
\left\| \left(\widetilde{F}^{(r)} - F^{(r)} \right) P_n Q_{n-mn\beta} \right\|_1
$$

\n
$$
\leq \left\| \left(Id - F^{(r)} \left(\mathcal{J} - \widetilde{\mathcal{J}} \right) \right) P_{n+2n\beta+1} Q_{n-(m+2)n\beta-1} \widetilde{F}^{(r)} P_{n+n\beta+1} Q_{n-(m+1)n\beta-1} \left(\widetilde{\mathcal{J}} - \mathcal{J} \right) P_{n+n\beta} Q_{n-(m+1)n\beta} \widetilde{F}^{(r)} P_n Q_{n-mn\beta} \right\|_1 + R_n,
$$
\n(419)

where $R_n = O(n^{m\alpha}e^{-d_0n^{\beta-\frac{\alpha}{2}}})$ is exponentially small. Use the trace norm inequality $||ABCD||_1 \le$ ∥*A*∥∞∥*B*∥2∥*C*∥∞∥*D*∥² and [\(416\)](#page-72-1) to obtain

$$
\left\| (\widetilde{F}^{(r)} - F^{(r)}) P_n Q_{n-mn^{\beta}} \right\|_1 \leq \| (Id - F^{(r)} \left(\mathcal{J} - \widetilde{\mathcal{J}} \right) P_{n+2n^{\beta}+1} Q_{n-(m+2)n^{\beta}-1} \|_{\infty} \| \widetilde{F}^{(r)} P_{n+n^{\beta}+1} Q_{n-(m+1)n^{\beta}-1} \|_2
$$

$$
\cdot \| \left(\widetilde{\mathcal{J}} - \mathcal{J} \right) P_{n+n^{\beta}} Q_{n-(m+1)n^{\beta}} \|_{\infty} \| \widetilde{F}^{(r)} P_n Q_{n-mn^{\beta}} \|_2 + R_n, \quad (420)
$$

where $R_n = O\left(n^{\alpha+2\beta}e^{-d_0n^{\beta-\frac{\alpha}{2}}}\right)$ is exponentially small. Then by the assumption [\(349\)](#page-64-0) and the fact that $||F^{(r)}||_{\infty} = O(n^{\alpha})$ we have

$$
\left\| \left(\widetilde{\mathcal{J}} - \mathcal{J} \right) P_{n+n\beta} Q_{n-(m+1)n\beta} \right\|_{\infty} = O(n^{-2\beta}), \tag{421}
$$

$$
\left\| F^{(r)}\left(\mathcal{J} - \widetilde{\mathcal{J}}\right) P_{n+2n^{\beta}+1} Q_{n-(m+2)n^{\beta}-1} \right\|_{\infty} = O(n^{\alpha-2\beta}). \tag{422}
$$

Use Lemma [9.1](#page-65-0) to obtain

$$
\|\widetilde{F}^{(r)}P_nQ_{n-mn^{\beta}}\|_2^2 = O(n^{\beta + \frac{3\alpha}{2}}). \tag{423}
$$

Recall that R_n is of exponentially small for *n* large. Plug [\(421\)](#page-73-0), [\(422\)](#page-73-1) and [\(423\)](#page-73-2) into [\(420\)](#page-73-3) to obtain, as $n \rightarrow \infty$,

$$
\left\| (\widetilde{F}^{(r)} - F^{(r)}) P_n Q_{n-mn^{\beta}} \right\|_1 = O(n^{-\beta + \frac{3\alpha}{2}}). \tag{424}
$$

The estimate [\(424\)](#page-73-4), together with [\(415\)](#page-72-0), implies that [\(411\)](#page-72-2) and [\(412\)](#page-72-3) are both of order $O(n^{m\alpha-\beta+\frac{\alpha}{2}})$.

For [\(413\)](#page-72-4), consider any $l = 1, 2, \ldots, m$. Recall the telescoping sum

$$
\widetilde{F}^l - F^l = \sum_{k=0}^{l-1} \widetilde{F}^{l-k-1} \left(\widetilde{F} - F \right) F^k \tag{425}
$$

Apply Lemma [9.2](#page-66-0) to obtain

$$
\left\| P_n(\widetilde{F}^l - F^l) Q_n \right\|_1 \le \sum_{k=0}^{l-1} (n^{\alpha} C_{op})^{l-k-1} \left\| P_{n+(l-k-1)n^{\beta}}(\widetilde{F} - F)F^k Q_n \right\|_1 + R_n,
$$
\n(426)

where $R_n = O\left(n^{\alpha+2\beta}e^{-d_0n^{\beta-\frac{\alpha}{2}}}\right)$ is exponentially small. We estimate [\(426\)](#page-73-5), by Lemma [9.4,](#page-69-0) to be

$$
||P_n(\widetilde{F}^l - F^l)Q_n||_1 = O\left(n^{(l-1)\alpha + \frac{3\alpha}{2} - \beta}\right)
$$
\n(427)

This shows that [\(413\)](#page-72-4) is of order $O(n^{m\alpha + \frac{\alpha}{2} - \beta})$.

In summary we have shown that [\(411\)](#page-72-2), [\(412\)](#page-72-3) and [\(413\)](#page-72-4) are all of order of $O(n^{m\alpha-\beta+\frac{\alpha}{2}})$. Hence, by assumption $0 < \frac{\alpha}{2} < \beta < 2$, we conclude that [\(410\)](#page-71-0) holds. Plug this into the cumulant formula [\(67\)](#page-16-0) and we complete the proof complete the proof. □

Now we are ready to prove Theorem [2.1.](#page-6-0)

Proof of Theorem [2.1](#page-6-0) . First consider test functions like $f(x) = \sum_{r=1}^{M} d_r \operatorname{Im}(x - \lambda_r)^{-1}$ for $d_r \in \mathbb{R}$ and $\operatorname{Im}(\lambda_r) > 0$, $f(x) = \sum_{r=1}^{M} d_r \operatorname{Im}(x - \lambda_r)^{-1}$ for $d_r \in \mathbb{R}$ and $\operatorname{Im}(\lambda_r) > 0$. 0,i.e., [\(49\)](#page-14-0). Let $X_n(f_{\alpha,x_0})$ be the mesoscopic linear statistics of the OPE given by the Chebyshev polynomial of the general line of the order of described in hosinging of Subsection 0.1. Agaly Theorem 2.2 and get the of the second kind at the edge as described in beginning of Subsection [9.1.](#page-64-1) Apply Theorem [2.3](#page-8-0) and get the following convergence in distribution as $n \to \infty$

$$
\widetilde{X}_n(f_{\alpha,x_0}) - \mathbb{E}[\widetilde{X}_n(f_{\alpha,x_0})] \to \mathcal{N}(0,\sigma_f^2),\tag{428}
$$

where σ_f^2 is the variance in Theorem [2.3.](#page-8-0) This is equivalent to the convergence of the cumulants

$$
\lim_{n \to \infty} n^{-m\alpha} C_m(\widetilde{F}) = \begin{cases} \widetilde{\sigma_f}^2, & m = 2\\ 0, & m > 2. \end{cases} \tag{429}
$$

Then Proposition [9.1](#page-71-1) implies that

$$
\lim_{n \to \infty} n^{-m\alpha} C_m(F) = \begin{cases} \widetilde{\sigma_f}^2, & m = 2\\ 0, & m > 2. \end{cases} \tag{430}
$$

This is equivalent to

$$
X_{f_{\alpha,x_0}}^{(n)} - \mathbb{E}[X_{f_{\alpha,x_0}}^{(n)}] \to \mathcal{N}(0, \sigma_f^2),
$$
\n(431)

where $X_f^{(n)}$ f_{α} , is the mesoscopic linear statistics of for OPEs whose recurrence coefficients satisfy [\(349\)](#page-64-0).

Now recall the argument in Section [4.2.](#page-27-0) Use the exact same proof of the last step of Theorem [2.3](#page-8-0) in Section [4.2](#page-27-0) and we extend [\(431\)](#page-74-0) to the test functions $f \in C_c^1$. □

10 Examples

Theorems [2.1,](#page-6-0) [2.2](#page-8-1) and [2.3](#page-8-0) can be used to obtain the asymptotics of mesoscopic fluctuations at the edges of many classes of OPEs, that are known in literature. Here we present some of the interesting examples with both continuous and discrete measures. We include classical examples as well as uncommon but still popular ones.

10.1 Laguerre Unitary Ensemble

The orthogonality measure of scaled Laguerre polynomials is given by

$$
d\mu(x) = x^{\gamma} e^{-nx} dx, \quad x \ge 0,
$$

for some parameter $\gamma > -1$. The recurrence relation reads

$$
xp_j(x) = \frac{\sqrt{(j+1)(j+1+\gamma)}}{n}p_{j+1}(x) + \frac{2j+\gamma+1}{n}p_j(x) + \frac{\sqrt{j(j+\gamma)}}{n}p_{j-1}(x)
$$

and hence the recurrence coefficients are

$$
a_{j,n} = \frac{\sqrt{j(j+\gamma)}}{n} \quad , \quad b_{j,n} = \frac{2j+\gamma+1}{n}
$$
 (432)

Clearly, Condition [2.1](#page-7-0) is satisfied for all recurrence coefficients with indices in the following set *I*,

$$
I = \left\{ j \in \mathbb{N} : \frac{j}{n} \to 1 \right\}.
$$
\n(433)

At the left edge, $b_{n-1,n} - 2\sqrt{a_{n,n}a_{n-1,n}} = \frac{\gamma^2 + 1}{4n^2}$ $\frac{d^2+1}{4n^2} + O(n^{-3})$, by Taylor approximation. Take $x_0 = 0$. We compute that for all $j \in I_{n,m}^{(\beta)}$

$$
a_{j,n}a_{j-2,n} - a_{j-1,n}^2 = -\frac{1}{n^2} + O(n^{-4}),
$$
\n(434)

$$
(b_{j-1,n} - x_0 - a_{j,n})a_{j-2,n} - (b_{j-2,n} - x_0 - a_{j-1,n})a_{j-1,n} = -\frac{\gamma^2}{2n^3} + O(n^{\beta - 4}).
$$
\n(435)

By Theorem [2.3](#page-8-0) we conclude that the mesoscopic CLT holds for all $0 < \frac{\alpha}{2} < \beta < \frac{\alpha+1}{3} < 1$. Taking $\beta \to \frac{\alpha}{2}$. We conclude that the mesoscopic CLT holds for all $0 < \alpha < 2$.

At the right edge, $b_{n-1,n} + 2\sqrt{a_{n,n}a_{n-1,n}} = 4 + (2\gamma - 2)n^{-1} + O(n^{-2})$, by Taylor approximation. Take $x_0 = 4$, by Theorem 2.2 and we conclude that the asymptotics of the edge fluctuations holds for all $\alpha \in (0, \frac{2}{3})$ apply Theorem [2.2](#page-8-1) and we conclude that the asymptotics of the edge fluctuations holds for all $\alpha \in (0, \frac{2}{3})$ $\frac{2}{3}$).

10.2 Gaussian Unitary Ensemble

The orthogonality measure of scaled Hermite polynomials is given by

$$
d\mu(x) = e^{-nx^2/2}dx, \quad x \in \mathbb{R}.
$$

The recurrence coefficients read

$$
a_{j,n} = \sqrt{\frac{j}{n}} \quad , \quad b_{j,n} = 0. \tag{436}
$$

Clearly, Condition [2.1](#page-7-0) is satisfied for all recurrence coefficients with indices in the following set *I*,

$$
I = \left\{ j \in \mathbb{N} : \frac{j}{n} \to 1 \right\}.
$$
\n(437)

Note that $b_{n-1,n} \pm 2\sqrt{a_{n,n}a_{n-1,n}} = \pm 2 + O(n^{-1})$. Take $x_0 = -2$ or 2, apply Theorem [2.2](#page-8-1) and we conclude that the asymptotics of the edge fluctuations holds for all $\alpha \in (0, \frac{2}{3})$ $\frac{2}{3}$).

10.3 Jacobi Unitary Ensemble

The orthogonality measure of Jacobi polynomials is given by

$$
d\mu(x) = (x-2)^{\gamma_1}(x+2)^{\gamma_2}dx, \quad x \in [-2,2],
$$

for some parameter $\gamma_1, \gamma_2 > -1$. The recurrence coefficients read

$$
a_j = \sqrt{\frac{16j(j+\gamma_1+\gamma_2)(j+\gamma_1)(j+\gamma_2)}{(2j+\gamma_1+\gamma_2-1)(2j+\gamma_1+\gamma_2)^2(2j+\gamma_1+\gamma_2+1)}}, \quad b_j = \frac{2(\gamma_2^2 - \gamma_1^2)}{(2j+\gamma_1+\gamma_2)(2j+\gamma_1+\gamma_2+2)}.
$$
\n(438)

Note that as $j \to \infty$,

$$
a_j = 1 + \frac{1 - 2\gamma_1^2 - 2\gamma_2^2}{8j^2} + O(j^{-3}), \quad b_j = \frac{\gamma_2^2 - \gamma_1^2}{2j^2} + O(j^{-3}).
$$

The equilibrium measure is given the the following figure.

The edges of fluctuations are at $x_0 = 2$ or -2 . Hence, by Theorem [2.3](#page-8-0) or [2.1](#page-6-0) we have the mesoscopic fluctuations at the edges hold for all $\alpha \in (0, 2)$. Note that for the Chebyshev polynomial of the second kind we have $\gamma_1 = \gamma_2 = -1/2$, $a_j = 1$ and $b_j = 0$.

10.4 Freud Weight

The orthogonality measure for Freud polynomials is given by Freud weight

$$
d\mu_n(x) = e^{-n|x|^\gamma} dx, \quad \gamma > 0, \quad x \in \mathbb{R}.
$$

The recurrence is shown in [\[31,](#page-82-0) [32\]](#page-82-1) that

$$
a_{j,n} = \frac{1}{2} \left(\frac{\Gamma(\frac{\gamma}{2}) \Gamma(\frac{1}{2})}{\Gamma(\frac{\gamma+1}{2})} \right)^{\frac{1}{\gamma}} \left(\frac{j}{n} \right)^{\frac{1}{\gamma}} (1 + res_{\gamma}(j)) \text{ and } b_{j,n} = 0,
$$
 (439)

where Γ is the Gamma function and

$$
res_{\gamma}(j) = \begin{cases} O(j^{-2}), & \gamma \ge 2 \text{ or } 0 < \gamma \le \frac{1}{2}, \\ O(j^{-\gamma}), & 1 < \gamma \le 2, \\ O(j^{-1}(\log j)^{-2}), & \gamma = 1, \\ O(j^{-1/\gamma}), & \gamma \ge 2 \text{ or } \frac{1}{2} < \gamma < 1 \end{cases}
$$
 (440)

Note that its Jacobi operator is essentially self-adjoint if and only if $\gamma \geq 1$. In such case, the moment problem is determinant. Note that for $0 < \gamma < 1$, the set of Freud orthonormal polynomials is not dense in $L^2(\mu_n)$.
The moment problem is indeterminant The moment problem is indeterminant.

Clearly, Condition [2.1](#page-7-0) is satisfied for all recurrence coefficients with indices in the following set *I*,

$$
I = \left\{ j \in \mathbb{N} : \frac{j}{n} \to 1 \right\}.
$$
\n(441)

Note that $b_{n-1,n} \pm 2\sqrt{a_{n,n}a_{n-1,n}} = \pm \left(\frac{\Gamma(\frac{\gamma}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{\gamma+1}{2})}\right)$ $\Gamma(\frac{\gamma+1}{2})$ $\int_{\gamma}^{\frac{1}{\gamma}} + O(n^{-1})$. For both edges $x_0 = -\left(\frac{\Gamma(\frac{\gamma}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{\gamma+1}{2})}\right)$ $\Gamma(\frac{\gamma+1}{2})$ $\int_{\gamma}^{\frac{1}{\gamma}}$ or $\left(\frac{\Gamma(\frac{\gamma}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{\gamma+1}{2})}\right)$ $\Gamma(\frac{\gamma+1}{2})$ $\big)^{\frac{1}{\gamma}},$ apply Theorem [2.2](#page-8-1) and we conclude that the asymptotics of the edge fluctuations holds for all $\alpha \in (0, \frac{2}{3})$
In particular for $\alpha = 2$, the Fraud weight is reduced to be the Hermite case. $\frac{2}{3}$).

In particular, for $\gamma = 2$, the Freud weight is reduced to be the Hermite case.

10.5 Tricomi-Carlitz Polynomial Ensemble

Next, we consider the Tricomi-Carlitz polynomials. Their zero distributions were first studied by [\[40,](#page-82-2) [41\]](#page-82-3). We mention this example to emphasis that the edge of fluctuations may not coincide with the edge of the equilibrium measure. Take $\gamma > 1$. The orthogonal measure $v^{(\gamma)}$ is a step function with jumps at the points

$$
v^{(\gamma)}(x) = \frac{(k+\gamma)^{k-1}e^{-k}}{k!}
$$
 at $x = \pm (k+\gamma)^{-\frac{1}{2}}$, $k = 0, 1,$

It gives a discrete polynomial $f_n^{(\gamma)}$. To obtain a reasonable limit we need to rescale *x* by $n^{-1/2}$, see examples 4.7 in [\[42\]](#page-82-4). That is we have a scaled measure

$$
\mu^{(\gamma)}(x) := \nu^{(\gamma)}(\sqrt{n}x) = \frac{(k+\gamma)^{k-1}e^{-k}}{k!} \quad \text{at } x = \pm (k+\gamma)^{-\frac{1}{2}}\sqrt{n}, \quad k = 0, 1, \dots.
$$

The scaled polynomial is given by $f_n^{(\alpha)}(n^{-1/2}x)$. However, its recurrence coefficients admits a simpler form

$$
a_{j,n} = \sqrt{\frac{jn}{(j+\gamma-1)(j+\gamma)}} \quad , \quad b_{j,n} = 0 \tag{442}
$$

Clearly, Condition [2.1](#page-7-0) is satisfied for all recurrence coefficients with indices in the following set *I*,

$$
I = \left\{ j \in \mathbb{N} : \frac{j}{n} \to 1 \right\}.
$$
\n(443)

For both edges $b_{n-1,n} \pm 2\sqrt{a_{n,n}a_{n-1,n}} = \pm 2 + O(n^{-1})$, by Taylor approximation. Take $x_0 = -2$ or 2, apply Theorem [2.2](#page-8-1) and we conclude that the asymptotics of the edge fluctuations holds for all $\alpha \in (0, \frac{2}{3})$ $\frac{2}{3}$).

10.6 Weight with Logarithm-Singularity

Deift and Piorkowski in the work [\[25\]](#page-81-0) consider orthogonal polynomials with orthogonality measure $d\mu(x)$ by

$$
d\mu(x) = \log\left(\frac{2}{1-x}\right), \qquad x \in [-1, 1).
$$

They show the asymptotics of its recurrence coefficients to be

$$
a_{k,n} = \frac{1}{2} - \frac{1}{16k^2} - \frac{3}{32k^2 \log^2(k)} + O\left(\frac{1}{k^2 \log^3(k)}\right), \qquad b_{k,n} = \frac{1}{4k^2} - \frac{3}{16k^2 \log^2(n)} + O\left(\frac{1}{k^2 \log^3(k)}\right).
$$

Hence for both edges, $b_{n-1,n} \pm 2\sqrt{a_{n,n}a_{n-1,n}} = \pm 1 + O(n^{-2})$. Take $x_0 = -1$ or 1, apply Theorem [2.1](#page-6-0) and we conclude that the example time of the edge fluctuations helds for all ≈ 6 (0.2). conclude that the asymptotics of the edge fluctuations holds for all $\alpha \in (0, 2)$.

10.7 Krawtchouk Polynomial Ensemble

Given $K \in \mathbb{N}$ and $p \in (0, 1)$. The Krawtchouk polynomial $k_i(x; p, K)$ is a discrete polynomial on $\{0, 1, \ldots, K\}$. The orthogonality weight *v* is

$$
v(x) = {K \choose x} p^{x} (1-p)^{K-x}, \qquad x = 0, 1, ..., K.
$$

The recurrence coefficients of the Krawtchouk polynomials are

$$
a_j = \sqrt{(n-j+1)jp(1-p)},
$$
 $b_j = (n-j)p + j(1-p).$

The OPE with the Krawtchouk weight describes uniformly distributed domino tilings of the Aztec diamond, see [\[6\]](#page-80-0). Here K is related to the size of the diamond and it is particularly interesting to let *K* to infinity. Following [\[6\]](#page-80-0), we consider the case where $\frac{K}{n} \to t$ for $t \ge 2$ as $n \to \infty$. Of interests are the scaled polynomials k_j (nx ; p , N). Then the scaled orthogonal weight ν is

$$
\mu(x) = {K \choose nx} p^{nx} (1-p)^{K-nx}, \qquad x = 0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{K}{n}.
$$

The scaled Krawtchouk polynomials have the recurrence coefficients,

$$
a_{j,n} = \sqrt{\frac{(K-j+1)jp(1-p)}{n^2}}, \quad b_j = \frac{(K-j)p+j(1-p)}{n}.
$$

Clearly, Condition [2.1](#page-7-0) is satisfied for all recurrence coefficients with indices in the following set *I*,

$$
I = \left\{ j \in \mathbb{N} : \frac{j}{n} \to 1 \right\}.
$$
 (444)

For both edges $b_{n-1,n} \pm 2\sqrt{a_{n,n}a_{n-1,n}} = (t-2)p + 1 \pm (t-1)\sqrt{p(1-p)} + O(n^{-1})$, by Taylor approximation. Take $x_0 = (t-2)p + 1 \pm (t-1)\sqrt{p(1-p)}$, apply Theorem [2.2](#page-8-1) and we conclude that the asymptotics of the edge fluctuations holds for all $\alpha \in (0, \frac{2}{3})$ $\frac{2}{3}$).

10.8 Hahn Polynomial Ensemble

Given $a, b > -1$ and $a, b, N \in \mathbb{N}$. The Hahn orthonormal weight is defined as

$$
v_N^{(a,b)}(x) = \binom{a+x}{x} \binom{b+N-x}{N-x}, \quad x = 0, 1, \dots, N. \tag{445}
$$

The Hahn ensemble appears in the lozenge tilings of a hexagon with uniform weights. The parameters *^a*, *^b*, *^N* are related to the size of the hexagon. To study the large hexagon, of interested when the sizes grows linearly in *n* as $n \rightarrow \infty$. Let

$$
\frac{a}{n} \to t_1, \quad \frac{b}{n} \to t_2, \quad \frac{N}{n} \to t_3, \quad \text{for some } t_1, t_2 > 0, t_3 \ge 1.
$$

The scaled Hahn weight is

$$
\mu_N^{(a,b)}(x) = \binom{a + nx}{nx} \binom{b + N - nx}{N - nx}, \quad x = 0, \frac{1}{n}, \dots, \frac{N}{n}.
$$
\n(446)

Then the scaled Hahn polynomials have the recurrence coefficients,

$$
a_{j,n} = \frac{j(j+a+b+N+1)(j+b)}{N(2j+a+b)(2j+a+b+1)} \sqrt{\frac{(N-j)(j+a+b)(a+j)(2j+a+b+1)}{j(j+a+b+N+1)(b+j)(2j+a+b-1)}},
$$

$$
b_{j,n} = \frac{(N-j)(j+a+b+1)(j+a+1)}{N(2j+a+b+N+1)(2j+a+b+2)}.
$$

Clearly, Condition [2.1](#page-7-0) is satisfied for all recurrence coefficients with indices in the following set *I*,

$$
I = \left\{ j \in \mathbb{N} : \frac{j}{n} \to 1 \right\}.
$$
 (447)

For both edges, clearly, the limits $\lim_{n} a_{n,n} = a_0$ and $\lim_{n} b_{n-1,n} = b_0$ exist, with the rate of convergence at most $O(n^{-1})$. Take $x_0 = b_0 \pm 2a_0$, apply Theorem [2.2](#page-8-1) and we conclude that the asymptotics of the edge fluctuations holds for all $\alpha \in (0, \frac{2}{3})$ $\frac{2}{3}$).

Acknowledgements

The author would like to express his deepest gratitude to Maurice Duits, for proposing this research question and for his invaluable guidance and supports throughout this work. The author is also very grateful to Grzegorz Świderski for bringing the Freud polynomials and the moment problem to the author's attention; to Mateusz Piorkowski for bringing the logarithm singularity example which motivates the development of ´ Theorem [2.1.](#page-6-0)

References

- [1] Alexei Borodin. Determinantal point processes. *The Oxford handbook of random matrix theory*, pages 231–249, 2011.
- [2] J. Ben Hough, Manjunath Krishnapur, Yuval Peres, and Bálint Virág. Determinantal processes and independence. 2006.
- [3] Kurt Johansson. Random matrices and determinantal processes. In *Les Houches*, volume 83, pages 1–56. Elsevier, 2006.
- [4] Wolfgang König. Orthogonal polynomial ensembles in probability theory. Probability Surveys, 2:385– 447, 2005.
- [5] Kurt Johansson. Random matrices and determinantal processes. *arXiv preprint math-ph*/*0510038*, 2005.
- [6] Kurt Johansson. Non-intersecting paths, random tilings and random matrices. *Probability theory and related fields*, 123(2):225–280, 2002.
- [7] Percy Deift. Orthogonal polynomials and random matrices: a riemann–hilbert approach, courant lect. *Notes Math*, 3, 1999.
- [8] Leonid Andreevich Pastur and Mariya Shcherbina. *Eigenvalue distribution of large random matrices*. Number 171. American Mathematical Soc., 2011.
- [9] Eli Levin and Doron S Lubinsky. Universality limits in the bulk for varying measures. *Advances in Mathematics*, 219(3):743–779, 2008.
- [10] Percy Deift and Dimitri Gioev. Universality at the edge of the spectrum for unitary, orthogonal, and symplectic ensembles of random matrices. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences*, 60(6):867–910, 2007.
- [11] Arno BJ Kuijlaars. Chapter 6 universality. *The Oxford Handbook on Random Matrix Theory*, pages 103–134, 2011.
- [12] Alexander Soshnikov. The central limit theorem for local linear statistics in classical compact groups and related combinatorial identities. *Annals of probability*, pages 1353–1370, 2000.
- [13] Yiting Li, Kevin Schnelli, and Yuanyuan Xu. Central limit theorem for mesoscopic eigenvalue statistics of deformed wigner matrices and sample covariance matrices. In *Annales de l'Institut Henri Poincar´e, Probabilit´es et Statistiques*, volume 57, pages 506–546. Institut Henri Poincare, 2021. ´
- [14] Jonathan Breuer and Maurice Duits. Universality of mesoscopic fluctuations for orthogonal polynomial ensembles. *Communications in Mathematical Physics*, 342(2):491–531, 2016.
- [15] Daniel Ofner. Stability of mesoscopic fluctuations of orthogonal polynomial ensembles under sparse decaying perturbations. *arXiv preprint arXiv:2410.07699*, 2024.
- [16] Benjamin Landon and Philippe Sosoe. Applications of mesoscopic clts in random matrix theory. *The Annals of Applied Probability*, 30(6):2769–2795, 2020.
- [17] Gaultier Lambert. Mesoscopic fluctuations for unitary invariant ensembles. 2018.
- [18] Florent Bekerman and Asad Lodhia. Mesoscopic central limit theorem for general beta-ensembles. *Annales de L'Institut Henri Poincare Section (B) Probability and Statistics*, 54(4):1917–1938, 2018.
- [19] Jonathan Breuer and Daniel Ofner. Mesoscopic universality for circular orthogonal polynomial ensembles. *arXiv preprint arXiv:2409.09803*, 2024.
- [20] Edward B Saff and Vilmos Totik. *Logarithmic potentials with external fields*, volume 316. Springer Science & Business Media, 2013.
- [21] Percy Deift, Thomas Kriecherbauer, K T-R McLaughlin, Stephanos Venakides, and Xin Zhou. Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences*, 52(11):1335–1425, 1999.
- [22] Roelof Koekoek, Peter A Lesky, René F Swarttouw, Roelof Koekoek, Peter A Lesky, and René F Swarttouw. *Hypergeometric orthogonal polynomials*. Springer, 2010.
- [23] Jonathan Breuer and Maurice Duits. Central limit theorems for biorthogonal ensembles and asymptotics of recurrence coefficients. *Journal of the American Mathematical Society*, 30(1):27–66, 2017.
- [24] Arno BJ Kuijlaars, KT-R McLaughlin, Walter Van Assche, and Maarten Vanlessen. The riemann– hilbert approach to strong asymptotics for orthogonal polynomials on [-1, 1]. *Advances in mathematics*, 188(2):337–398, 2004.
- [25] Percy Deift and Mateusz Piorkowski. Recurrence coefficients for orthogonal polynomials with a logarithmic weight function. *SIGMA. Symmetry, Integrability and Geometry: Methods and Applications*, 20:004, 2024.
- [26] Maurice Duits. On global fluctuations for non-colliding processes. *The Annals of Probability*, 46(3):1279–1350, 2018.
- [27] Maurice Duits, Erik Duse, and Wenkui Liu. Lozenge tilings of a hexagon and q-racah ensembles. *Journal of Physics A: Mathematical and Theoretical*, 57(40):405202, 2024.
- [28] Barry Simon. *Trace ideals and their applications*. Number 120. American Mathematical Soc., 2005.
- [29] Konrad Schmüdgen. *The moment problem*, volume 9. Springer, 2017.
- [30] Ju. Makarovich Berezansky. *Expansions in eigenfunctions of selfadjoint operators*, volume 17. American Mathematical Society, 1968.
- [31] Doron S Lubinsky, Hrushikesh N Mhaskar, and Edward B Saff. A proof of freud's conjecture for exponential weights. *Constructive Approximation*, 4:65–83, 1988.
- [32] Thomas Kriecherbauer and K McLaughlin. Strong asymptotics of polynomials orthogonal with respect to freud weights. *IMRN: International Mathematics Research Notices*, 1999(6), 1999.
- [33] Jonathan Breuer and Maurice Duits. The nevai condition and a local law of large numbers for orthogonal polynomial ensembles. *Advances in Mathematics*, 265:441–484, 2014.
- [34] Jean-Michel Combes and L Thomas. Asymptotic behaviour of eigenfunctions for multiparticle schrödinger operators. 1973.
- [35] Gérard Meurant. A review on the inverse of symmetric tridiagonal and block tridiagonal matrices. *SIAM Journal on Matrix Analysis and Applications*, 13(3):707–728, 1992.
- [36] Maurice Duits and Kurt Johansson. On mesoscopic equilibrium for linear statistics in dyson's brownian motion. 255(1222), 2018.
- [37] Torsten Ehrhardt. A generalization of pincus' formula and toeplitz operator determinants. *Archiv der Mathematik*, 80(3):302–309, 2003.
- [38] Percy Deift, Thomas Kriecherbauer, K T-R McLaughlin, Stephanos Venakides, and Xin Zhou. Strong asymptotics of orthogonal polynomials with respect to exponential weights. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences*, 52(12):1491–1552, 1999.
- [39] KT-R McLaughlin and PD Miller. The steepest descent method for orthogonal polynomials on the real line with varying weights. *International Mathematics Research Notices*, 2008:rnn075, 2008.
- [40] William MY Goh and Jet Wimp. On the asymptotics of the tricomi–carlitz polynomials and their zero distribution (i). *SIAM Journal on Mathematical Analysis*, 25(2):420–428, 1994.
- [41] William MY Goh and Jet Wimp. The zero distribution of the tricomi-carlitz polynomials. *Computers* & *Mathematics with Applications*, 33(1-2):119–127, 1997.
- [42] Arno BJ Kuijlaars and Walter Van Assche. The asymptotic zero distribution of orthogonal polynomials with varying recurrence coefficients. *Journal of approximation theory*, 99(1):167–197, 1999.