## *p*-ADIC SYMPLECTIC GEOMETRY OF INTEGRABLE SYSTEMS AND WEIERSTRASS-WILLIAMSON THEORY

LUIS CRESPO ÁLVARO PELAYO

In memory of Professor Vladimir Voevodsky (1966–2017).

ABSTRACT. We establish the foundations of the local linear symplectic geometry of *p*-adic integrable systems on *p*-adic analytic symplectic 4-dimensional manifolds, by classifying all their possible local linear models. In order to do this we develop a new approach, of independent interest, to the theory of Weierstrass and Williamson concerning the diagonalization of real matrices by real symplectic matrices, which we show can be generalized to p-adic matrices, leading to a classification of real (2n)-by-(2n) matrices and of p-adic 2-by-2 and 4-by-4 matrix normal forms. A combination of these results and the Hardy-Ramanujan formula shows that both the number of padic matrix normal forms and the number of local linear models of p-adic integrable systems grow almost exponentially with their dimensions, in strong contrast with the real case. Although the paper deals mostly with matrices and systems whose singularities are non-degenerate, we include several classifications in the degenerate p-adic case, for which the literature is limited even in the real case. The paper also includes a number of results concerning symplectic linear algebra over arbitrary fields in arbitrary dimensions as well as applications to p-adic mechanical systems and singularity theory for p-adic analytic maps on 4-manifolds. These results fit in a program, proposed a decade ago by Voevodsky, Warren and the second author, to develop a *p*-adic theory of integrable systems with the goal of later implementing it using proof assistants.

#### CONTENTS

1. Introduction and main results about integrable systems	2
1.1. <i>p</i> -adic geometry, symplectic geometry and integrable systems	2
1.2. The two (interrelated) goals of this paper	3
1.3. Main results about integrable systems: Theorems A, B and C	4
Structure of the paper	10
2. Main results concerning matrices: Theorems D–I	10
2.1. The real Weierstrass-Williamson classification	10
2.2. The <i>p</i> -adic Weierstrass-Williamson classification	11
3. Symplectic linear algebra over algebraically closed fields	17
4. <i>p</i> -adic matrix classification in dimension 2	27
4.1. General preparatory results	27
4.2. Proof of Theorems D and E	31
5. General preparatory results for fields of characteristic different from 2	35
6. <i>p</i> -adic matrix classification in dimension 4	39
6.1. Case $p \neq 2$	40
6.2. Case $p = 2$	43
6.3. Proof of Theorem F	52
6.4. Proof of Theorem G	58
6.5. Proof of Theorem H	60
7. Matrix classification in the real case, for any dimension	60

7.1. The general case	60
7.2. Example of application of our method for matrices of arbitrary order	65
7.3. Positive-definite case	66
8. Comments on the <i>p</i> -adic classification in higher dimensions and proof of	
Theorem I	67
8.1. Preparatory lemmas	67
8.2. Proof of Theorem I	68
8.3. Remarks and applications	68
9. Application to normal forms of <i>p</i> -adic singularities	69
10. Application to normal forms of singularities of integrable systems	74
10.1. Non-degenerate critical points of integrable systems	74
10.2. Degenerate critical points in the real case	75
10.3. Proof of Theorems A, B and C	77
10.4. Degenerate critical points in the <i>p</i> -adic case	78
10.5. Symplectic dynamics of integrable systems and their level sets	80
11. Application to classical mechanical systems	84
12. Examples	87
12.1. Examples with matrices	87
12.2. Examples with functions and integrable systems	90
13. Circular symmetries of the <i>p</i> -adic models	91
Appendix A. Basic properties of <i>p</i> -adic numbers	93
A.1. Definition of $\mathbb{Q}_p$	93
A.2. Critical points of <i>p</i> -adic analytic functions	95
Acknowledgments	95
References	96

#### 1. INTRODUCTION AND MAIN RESULTS ABOUT INTEGRABLE SYSTEMS

1.1. *p*-adic geometry, symplectic geometry and integrable systems. In a recent lecture [38] Lurie commented that "roughly speaking *p*-adic geometry, or rigid analytic geometry, is a version of the theory of complex manifolds where instead of using complex numbers you use something like *p*-adic numbers". In the present paper we attempt to start studying this direction for the closely related class of symplectic manifolds: we will concentrate on the local linear theory of integrable systems with *p*-adic coefficients, on *p*-adic analytic symplectic manifolds.<sup>1</sup>

More concretely, for any prime number p, the p-adic numbers  $\mathbb{Q}_p$  form an extension field of the rational numbers  $\mathbb{Q}$  which plays a prominent role in various parts of geometry, as seen for instance in the aforementioned recent lecture by Lurie and Scholze-Weinstein's lectures [55].

The present paper takes a first step in introducing *p*-adic methods in symplectic geometry of integrable systems. We focus on *p*-adic matrix theory and *p*-adic linear symplectic geometry and applying them to fully describe the local linear theory of *p*-adic integrable systems  $F = (f_1, f_2) : (M, \omega) \to (\mathbb{Q}_p)^2$  on *p*-adic analytic symplectic 4-manifolds  $(M, \omega)$ , as part of a general approach to this new field proposed ten years ago by Voevodsky,

<sup>&</sup>lt;sup>1</sup>The relation between complex and symplectic structures on manifolds already appears implicitly in the pioneering work of Kodaira [35] and in a well known paper by Thurston [57], as well as in many other contributions including for instance [6, 14, 20, 21, 27].

Warren and the second author [46, Section 7]. Our techniques are primarily algebraic and rely on the theory of p-adic extension fields.

When we started this paper we did not expect the local linear theory of p-adic integrable systems to be so rich: the possibilities in the real case represent, in comparison, only a tiny proportion of what occurs in the p-adic case (see Table 2 which for example tells us that there are exactly 211 local linear models in the case of 2-adic symplectic 4mainfolds, while there are exactly 4 such models with real coefficients). Based on what we have found in this paper and [11], we believe that p-adic symplectic geometry and its application to p-adic integrable systems are promising research directions. Another example of this richness can be seen in the p-adic version of the real Jaynes-Cummings model from quantum optics, treated in our previous paper [11].

We expect that the results of this paper can be formalized using the proof assistant Coq, in the setting of homotopy type theory and Voevodsky's Univalent Foundations, which was the original motivation for Voevodsky, Warren and the second author to formalize the construction of the *p*-adic numbers in this setting in [46]. We refer to [4, 51, 52] for an introduction to the Voevodsky's univalence axiom and homotopy type theory.

1.2. The two (interrelated) goals of this paper. This paper has two closely related goals. One of the goals is to fully describe the local symplectic geometry of p-adic integrable systems on symplectic 4-manifolds, that is, we explicitly classify their local models and give a concrete list of their formulas. In order to this, we first need to extend the seminal theory of Weierstrass [63] and Williamson [66] concerning the diagonalization of real symmetric matrices by means of symplectic matrices, to p-adic 4-by-4 matrices, which is the other goal of the paper.

More concretely, by the work of Weierstrass [63] any real symmetric positive definite matrix is diagonalizable by a symplectic matrix. This was generalized in an influential paper [66] by Williamson from 1936, where he shows that any symmetric matrix is reducible to a normal form by a symplectic matrix, and gives a classification of all matrix normal forms. In Williamson's approach, matrix reductions take place in the base field. In this paper we recover the real Wierstrass-Williamson classification with a genuinely different strategy: we lift the problem to suitable extension fields where the solution is simpler.

Our methods extend to *p*-adic matrices in all dimensions where the problem can be expected to be solved, and we do this for *p*-adic 2-by-2 and 4-by-4 matrices, hence achieving one of the aforementioned main goals.

Our method generalizes to 6-by-6 and 8-by-8 matrices, because polynomials of degree 3 and 4 are solvable by radicals, although we do not do this in this paper. One cannot expect to deal with this problem in orders higher than 8 because this solvability is not available.

All of the above results concerning matrices are stated in Section 2 ("Main results concerning matrices: Theorems D-I") of the paper. These results can be used as stepping stones to derive a complete classification of the local linear models of *p*-adic analytic integrable systems in dimension 4, this being the second main goal of this paper.

As an application of our results and the Hardy-Ramanujan formula [30] (obtained also by Uspensky [58]) in number theory, we confirm that the number of *p*-adic (2*n*)-by-(2*n*) matrix normal forms grows at least with  $e^{\pi\sqrt{2n/3}}/4\sqrt{3n}$ , which in particular implies that the number of local linear normal forms of *p*-adic integrable systems on 2*n*-dimensional symplectic manifolds at a rank 0 critical point grows in the same way. This is in strong contrast with the real case, where the number of normal forms of integrable systems at a rank 0 critical point is quadratic in the dimension.

1.3. Main results about integrable systems: Theorems A, B and C. The notions of *p*-adic analytic function and of critical point of a *p*-adic analytic function are reviewed in Appendix A.2. For the case of integrable systems the notion of non-degenerate critical point is an extension of the definition for the case of  $\mathbb{Q}_p$ -valued functions (this is explained in Definition 9.4).

In order to state our classifications we need to define the following special sets of numbers. Recall that a *quadratic residue modulo* p is an integer which is congruent to a perfect square modulo p; if this does not hold, then the integer is called a *quadratic non-residue*<sup>2</sup> modulo p. Quadratic non-residues play a crucial role in our main theorems, for which we will need the following definition.

**Definition 1.1** (Non-residue sets and coefficient functions). Let p be a prime number. If  $p \equiv 1 \mod 4$ , let  $c_0$  be the smallest quadratic non-residue modulo p. We define the *non-residue sets* 

$$X_{p} = \begin{cases} \{1, c_{0}, p, c_{0}p, c_{0}^{2}p, c_{0}^{3}p, c_{0}p^{2}\} & \text{if } p \equiv 1 \mod 4; \\ \{1, -1, p, -p, p^{2}\} & \text{if } p \equiv 3 \mod 4; \\ \{1, -1, 2, -2, 3, -3, 6, -6, 12, -18, 24\} & \text{if } p = 2. \end{cases}$$

$$Y_{p} = \begin{cases} \{c_{0}, p, c_{0}p\} & \text{if } p \equiv 1 \mod 4; \\ \{-1, p, -p\} & \text{if } p \equiv 3 \mod 4; \\ \{-1, 2, -2, 3, -3, 6, -6\} & \text{if } p = 2. \end{cases}$$

We also define the *coefficient functions*  $C_i^k : Y_p \times (\mathbb{Q}_p)^4 \to \mathbb{Q}_p$  and  $\mathcal{D}_i^k : Y_p \times (\mathbb{Q}_p)^4 \to \mathbb{Q}_p$ , for  $k \in \{1, 2\}, i \in \{0, 1, 2\}$ , by

$$\mathcal{C}_{0}^{1}(c,t_{1},t_{2},a,b) = \frac{ac}{2(c-b^{2})}, \ \mathcal{C}_{1}^{1}(c,t_{1},t_{2},a,b) = \frac{b}{b^{2}-c}, \ \mathcal{C}_{2}^{1}(c,t_{1},t_{2},a,b) = \frac{1}{2a(c-b^{2})}, \\ \mathcal{C}_{0}^{2}(c,t_{1},t_{2},a,b) = \frac{abc}{2(b^{2}-c)}, \ \mathcal{C}_{1}^{2}(c,t_{1},t_{2},a,b) = \frac{c}{c-b^{2}}, \ \mathcal{C}_{2}^{2}(c,t_{1},t_{2},a,b) = \frac{b}{2a(b^{2}-c)}, \\ \mathcal{D}_{0}^{1}(c,t_{1},t_{2},a,b) = -\frac{t_{1}+bt_{2}}{2a}, \ \mathcal{D}_{1}^{1}(c,t_{1},t_{2},a,b) = -bt_{1}-ct_{2}, \ \mathcal{D}_{2}^{1}(c,t_{1},t_{2},a,b) = -\frac{ac(t_{1}+bt_{2})}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2a}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -c(t_{1}+bt_{2}), \ \mathcal{D}_{2}^{2}(c,t_{1},t_{2},a,b) = -\frac{ac(bt_{1}+ct_{2})}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2a}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -c(t_{1}+bt_{2}), \ \mathcal{D}_{2}^{2}(c,t_{1},t_{2},a,b) = -\frac{ac(bt_{1}+ct_{2})}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2a}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -c(t_{1}+bt_{2}), \ \mathcal{D}_{2}^{2}(c,t_{1},t_{2},a,b) = -\frac{ac(bt_{1}+ct_{2})}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2a}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -c(t_{1}+bt_{2}), \ \mathcal{D}_{2}^{2}(c,t_{1},t_{2},a,b) = -\frac{ac(bt_{1}+ct_{2})}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2a}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \ \mathcal{D}_{1}^{2}(c,t_{1},t_{2},a,b) = -\frac{bt_{1}+ct_{2}}{2}, \\ \mathcal{D}_{0}^{2}(c,t_{1},t_{2},a,b) = -\frac{$$

The choice of  $c_0$  as the least quadratic residue is not essential, in the sense that the normal forms that appear in the upcoming forms are "equivalent" (in the precise sense of Propositions 4.9 and 6.11).

Now we state our main results about integrable systems, which concern non-degenerate systems (for results about degenerate systems, which are more technical, see Sections 10.2 and 10.4). We refer to Figures 2 and 3 for a depiction of some of the cases covered by Theorem A and to Figure 1 for an illustration of the real case.

Let  $(M, \omega)$  be a *p*-adic analytic symplectic 4-dimensional manifold. In the results below we use the following terminology. By *linear symplectic coordinates*  $(x, \xi, y, \eta)$  with the origin at a point  $m \in M$  we mean coordinates given by a local linear symplectomorphism  $\phi : ((\mathbb{Q}_p)^4, \omega_0) \to (M, \omega)$ , centered at (0, 0, 0, 0) (that is, such that  $\phi(0, 0, 0, 0) = m$  and

<sup>&</sup>lt;sup>2</sup>For any prime p > 2, the number of quadratic non-residues modulo p is (p-1)/2. For example, if p = 17, the quadratic non-residues modulo 17 are 3, 5, 6, 7, 10, 11, 12 and 14.




FIGURE 1. Normal forms of regular and critical points of elliptic-elliptic, focus-focus and elliptic-regular type of an integrable system  $F : \mathbb{R}^4 \to \mathbb{R}^2$ . Some of these can be normal forms of Theorem A (see Remark 1.5).

 $\phi^*\omega_m = \omega_0$ ), where  $\omega_0$  is the standard symplectic form on  $(\mathbb{Q}_p)^4$ . In terms of matrices this means that  $S^T\Omega S = \Omega_0$ , where S,  $\Omega$  and  $\Omega_0$  are the matrices of  $\phi$ ,  $\omega_m$  and the standard symplectic form  $\omega_0$  (see Definition 9.1 for details). For the notion of critical point of p-adic integrable system and its rank see Definition 10.1.

**Theorem A** (*p*-adic integrable local linear models in dimension 4). Let *p* be a prime number. Let  $X_p, Y_p, C_i^k, \mathcal{D}_i^k$  be the non-residue sets and coefficient functions in Definition 1.1. Let  $(M, \omega)$  be a *p*-adic analytic symplectic manifold of dimension 4 and let *F* :  $(M, \omega) \to (\mathbb{Q}_p)^2$  be a *p*-adic analytic integrable system. Let *m* be a non-degenerate critical point of *F*. Then there exist linear symplectic coordinates  $(x, \xi, y, \eta)$  with the origin at *m* and an invertible matrix  $B \in \mathcal{M}_2(\mathbb{Q}_p)$  such that in these coordinates we have:

(1.1) 
$$B \circ (F - F(m)) = (g_1, g_2) + \mathcal{O}(3),$$

where the expression of  $(g_1, g_2)$  depends on the rank of  $m \in \{0, 1\}$ . If m is a rank 0 critical point then one of the following situations occurs:

- (1) There exist  $c_1, c_2 \in X_p$  such that  $g_1(x, \xi, y, \eta) = x^2 + c_1 \xi^2, g_2(x, \xi, y, \eta) = y^2 + c_2 \eta^2;$
- (2) There exists  $c \in Y_p$  such that  $g_1(x,\xi,y,\eta) = x\eta + cy\xi, g_2(x,\xi,y,\eta) = x\xi + y\eta;$
- (3) There exist  $c, t_1$  and  $t_2$  corresponding to one row of Table 1 and  $(a, b) \in \{(1, 0), (a_1, b_1)\}$ , where  $(a_1, b_1)$  is given in the row in question, such that

$$g_k(x,\xi,y,\eta) = \sum_{i=0}^2 \mathcal{C}_i^k(c,t_1,t_2,a,b) x^i y^{2-i} + \sum_{i=0}^2 \mathcal{D}_i^k(c,t_1,t_2,a,b) \xi^i \eta^{2-i},$$

for  $k \in \{1, 2\}$ .

Otherwise, if m is a rank 1 point, then there exists  $c \in X_p$  such that  $g_1(x, \xi, y, \eta) = x^2 + c\xi^2$ and  $g_2(x, \xi, y, \eta) = \eta$ .

Furthermore, if there are two sets of linear symplectic coordinates in which F has one of these forms, then the pair  $(g_1, g_2)$  corresponding to the first set of linear symplectic coordinates and the pair  $(g'_1, g'_2)$  corresponding to the second set of linear symplectic coordinates are in the same case; if it is case (2) or (3), or a rank 1 point, they coincide, and in case (1) they coincide up to ordering.

1 1 4							
	$p \equiv$	≡l m	od 4				
c	$t_1$	$t_2$	$a_1$	$b_1$			
$c_0$	p	0	p	1/p			
	0	1	p	0			
	0	p	p	0			
p	$c_0$	0	1	1			
	0	1	$c_0$	0			
	0	$c_0$	$c_0$	0			
$c_0 p$	$c_0$	0	1	1			
	0	1	$c_0$	0			
	0	$c_0$	$c_0$	0			
	$p \equiv$	= 3 m	1004				
c	$t_1$	$t_2$	$a_1$	$b_1$			
-1	p	0	$b_0$	$a_0/b_0$			
	$a_0$	$b_0$	p	0			
	$pa_0$	$pb_0$	p	0			
p	-1	0	1	1			
	0	1	-1	0			
-p	-1	0	1	1			
	0	1	-1	0			

p =	$p = 2 \wedge c = -1$							
$t_1$	$t_2$	$a_1$	$b_1$					
2	0	1	2					
3	0	1	1					
6	0	1	1					
1	1	3	0					
3	3	3	0					
1	2	2	0					
2	4	2	0					
-1	3	2	0					
-2	6	2	0					

p	$p = 2 \land c = 2$							
$t_1$	$t_2$	$a_1$	$b_1$					
-1	0	1	2					
3	0	1	1					
-3	0	1	1					
0	1	-1	0					
0	3	-1	0					
1	1	-1	0					
3	3	-1	0					
2	1	3	0					
-2	-1	3	0					
6	3	3	0					
-6	-3	3	0					

$p = 2 \wedge c = -2$							
$t_1$	$t_2$	$a_1$	$b_1$				
-1	0	1	1				
3	0	1	-2				
-3	0	1	1				
0	1	3	0				
0	3	3	0				
1	1	-1	0				
-1	-1	-1	0				
-2	1	-1	0				
2	-1	-1	0				

 $\frac{b_1}{1}$   $\frac{1}{6}$  1

$p = 2 \land c = 3$							
$t_1$	$t_2$	$a_1$	$b_1$				
-1	0	1	1				
2	0	1	1				
-2	0	1	3				
0	1	2	0				
0	2	2	0				
1	1	-1	0				
-1	-1	-1	0				
3	1	-1	0				
-3	-1	-1	0				

$p = 2 \wedge c = -3$						
$t_1$	$t_2$	$a_1$	$b_1$			
-1	0	2	1/2			
2	0	2	1/2			
-2	0	1	-6			
0	1	-1	0			
0	2	-1	0			
1	2	2	0			
-1	-2	2	0			
2	4	2	0			
-2	-4	2	0			
-6	1	-1	0			
-12	2	-1	0			

$p = 2 \land c = 6$						
$t_1$	$t_2$	$a_1$				
-1	0	1				
3	0	1				
-3	0	1				
0	1	3				
0	3	3				
1	1	-1				
-1	-1	-1				
6	1	-1				
-6	-1	-1				

$p = 2 \wedge c = -6$						
$t_1$	$t_2$	$a_1$	$b_1$			
-1	0	1	-6			
3	0	1	1			
-3	0	1	1			
0	1	-1	0			
0	3	-1	0			
1	1	-1	0			
3	3	-1	0			
-6	1	3	0			
6	-1	3	0			
-18	3	3	0			
18	-3	3	0			

TABLE 1. Parameters for the normal form (3) of Theorem A. In the table, for  $p \equiv 1 \mod 4$ ,  $c_0$  is the smallest quadratic non-residue modulo p. For  $p \equiv 3 \mod 4$ ,  $a_0$  and  $b_0$  are such that  $a_0^2 + b_0^2 \equiv -1 \mod p$ . For p = 2there are many more possible parameters, and they are separated by the value of c.

**Remark 1.2.** There are many more local linear models of p-adic integrable systems than real ones, so from a physical viewpoint they should be able to model physical or other phenomena beyond the applications with real coefficients, see for example [29, 33, 34] and the references therein. For applications in biology see [3, 18].

**Remark 1.3.** Identifying a Hessian with its quadratic form, the formula (1.1) would be written  $d^2F = (g_1, g_2)$ . In the expression for  $g_1$  and  $g_2$  in any of the cases above, if we change the values of the parameters, the resulting functions still form an integrable system, but they are not a normal form. We can apply the theorem to these new functions, resulting in a new normal form  $(g'_1, g'_2)$  linearly symplectomorphic to  $(g_1, g_2)$ , in the same or a different case.

**Theorem B** (Number of *p*-adic integrable local linear models, in dimension 4). Let *p* be a prime number. Let  $X_p, Y_p, C_i^k, \mathcal{D}_i^k$  be the non-residue sets and coefficient functions in Definition 1.1. Let  $(M, \omega)$  be a *p*-adic analytic symplectic 4-manifold. Then the following statements hold:

- (1) If  $p \equiv 1 \mod 4$ , there are exactly 49 normal forms for a rank 0 non-degenerate critical point, and exactly 7 normal forms of a rank 1 non-degenerate critical point, of a p-adic analytic integrable system  $F: (M, \omega) \to (\mathbb{Q}_p)^2$  up to local linear symplectomorphisms centered at the critical point;
- (2) If  $p \equiv 3 \mod 4$ , there are exactly 32 normal forms for a rank 0 non-degenerate critical point, and exactly 5 normal forms of a rank 1 non-degenerate critical point, of a p-adic analytic integrable system  $F : (M, \omega) \to (\mathbb{Q}_p)^2$  up to local linear symplectomorphisms centered at the critical point;
- (3) If p = 2, there are exactly 211 normal forms for a rank 0 non-degenerate critical point, and exactly 11 normal forms of a rank 1 non-degenerate critical point, of a p-adic analytic integrable system  $F : (M, \omega) \to (\mathbb{Q}_p)^2$  up to local linear symplectomorphisms centered at the critical point.

In the three cases above, the normal forms for a rank 0 point are given by

$$\left\{ (x^2 + c_1 \xi^2, y^2 + c_2 \eta^2) : c_1, c_2 \in X_p \right\} \cup \left\{ (x\eta + cy\xi, x\xi + y\eta) : c \in Y_p \right\} \\ \cup \left\{ \left( \sum_{i=0}^2 \mathcal{C}_i^1(c, t_1, t_2, a, b) x^i y^{2-i} + \sum_{i=0}^2 \mathcal{D}_i^1(c, t_1, t_2, a, b) \xi^i \eta^{2-i}, \right. \\ \left. \sum_{i=0}^2 \mathcal{C}_i^2(c, t_1, t_2, a, b) x^i y^{2-i} + \sum_{i=0}^2 \mathcal{D}_i^2(c, t_1, t_2, a, b) \xi^i \eta^{2-i} \right) : \\ \left. (a, b) \in \left\{ (1, 0), (a_1, b_1) \right\}, c, t_1, t_2, a_1, b_1 \text{ in one row of Table 1} \right\}$$

and those for a rank 1 point are given by

$$\Big\{(x^2+c\xi^2,\eta):c\in X_p\Big\}.$$

**Remark 1.4.** In the real case, there are exactly 4 normal forms for a rank 0 nondegenerate critical point:

$$\left\{ (x^2 + \xi^2, y^2 + \eta^2), (x^2 + \xi^2, y\eta), (x\xi, y\eta), (x\eta - y\xi, x\xi + y\eta) \right\}$$

and exactly 2 normal forms for a rank 1 non-degenerate critical point:

$$\left\{ (x^2 + \xi^2, \eta), (x\xi, \eta) \right\}.$$




FIGURE 2. Symbolic representation of 2-dimensional fiber of focus-focus model if  $p \equiv 1 \mod 4$ , as a case of point (1) of Theorem A, which coincides with the elliptic-elliptic model. The four "cones" are 2-dimensional planes in 4-dimensional space.




FIGURE 3. Symbolic representation of 2-dimensional fiber of focus-focus model if  $p \not\equiv 1 \mod 4$ , as a case of point (2) of Theorem A, which coincides with the fiber in the real case. The two "cones" are actually 2-dimensional planes in 4-dimensional space that meet at a point.

**Remark 1.5.** In the *p*-adic category, the elliptic-elliptic and elliptic-regular points are normal forms of Theorem A: the former is a rank 0 point in case (1) with  $c_1 = c_2 = 1$ , and the latter is a rank 1 point with c = 1. The focus-focus point may also appear as a normal form, in case (2) with c = -1, but only if  $-1 \in Y_p$ , which happens if  $p \neq 1$ mod 4. Actually, if  $p \equiv 1 \mod 4$ , the normal form of a focus-focus point is ellipticelliptic; hence, the focus-focus and elliptic-elliptic points are linearly symplectomorphic if and only if  $p \equiv 1 \mod 4$ .

**Definition 1.6.** Let p be a prime number and let n be a positive integer. To each partition  $P = (a_1, \ldots, a_k)$  of n we associate the function

$$f_{P,p}(x_1,\xi_1,\ldots,x_n,\xi_n) = \sum_{i=1}^k \left( \frac{x_{b_{i-1}+1}^2}{2} + \sum_{j=b_{i-1}+1}^{b_i-1} \xi_j x_{j+1} + \frac{p\xi_{b_i}^2}{2} \right),$$

where  $b_0 = 0$  and  $b_i = \sum_{j=0}^{i} a_j$ , for  $i \in \{1, ..., k\}$ .

**Theorem C** (Number of p-adic integrable local linear models, arbitrary dimension). Let n be a positive integer. Let p be a prime number. The number of local linear normal forms of p-adic analytic integrable systems on 2n-dimensional p-adic analytic symplectic manifolds at a rank 0 non-degenerate critical point, up to local linear symplectomorphisms centered at the critical point, grows at least with

$$\frac{\mathrm{e}^{\pi\sqrt{2n/3}}}{4n\sqrt{3}}$$

Explicitly, for any two partitions P and Q of n, any two p-adic analytic integrable systems on a 2n-dimensional p-adic analytic manifold containing  $f_{P,p}$  and  $f_{Q,p}$ , respectively, as a component of their corresponding local linear normal forms, are not equivalent by local linear symplectomorphisms centered at the origin, where  $f_{P,p}$  is as given in Definition 1.6.

2n	$\mathbb{R}$	$\mathbb{Q}_2$	$\mathbb{Q}_3$	$\mathbb{Q}_5$	$\mathbb{Q}_7$
2	2	11	5	7	5
4	4	211	32	49	32
6	6	1883	123	234	129
8	9	21179	495	1054	525
10	12	161343	1595	4021	1787

TABLE 2. The number of families of normal forms of integrable systems on  $\mathbb{R}^{2n}$  and  $(\mathbb{Q}_p)^{2n}$  at a rank 0 critical point. Data is extracted from Table 3. For the *p*-adic case in dimension greater than 4, the numbers are only lower bounds. The actual number of forms might be even larger.

**Remark 1.7.** These results indicate that a global theory of *p*-adic integrable systems, which will probably be based on gluing local models, will include a large number of phenomena which do not occur in the real case. We have computed explicit lower bounds of the number of normal forms in Table 2.

The field of *p*-adic geometry is extensive, see [54, 55] and the references therein. *p*-adic geometry is also fundamental in mathematical physics and the theory of integrable systems, see for example [11, 15, 16, 17, 29, 46, 60]. For an introduction to different aspects of symplectic geometry, including its relations to mechanics and Poisson geometry, we refer to the survey articles [24, 45, 64, 65] and the books [5, 9, 31, 39, 40, 42].

The Weierstrass-Williamson theory of matrices has crucial applications in many areas including the theory of quantum states in quantum physics [12, 56, 62], hence this paper provides a new tool to further explore p-adic analogues of these applications in symplectic geometry and beyond.

Structure of the paper. In Section 2 we state the other main results of the paper, which concern matrices. Section 3 studies the classification problem in an algebraically closed field, where it is much simpler to give a classification. Sections 4 and 6 solve the classification problem over  $\mathbb{Q}_p$ , for 2-by-2 and 4-by-4 matrices, respectively. For 4-by-4 matrices there are first some general results in Section 5, then the case  $p \neq 2$  is studied in Section 6.1, then p = 2 in Section 6.2. In Section 7 we use the strategy introduced in previous sections to give a new proof of the general case of the Weierstrass-Williamson classification. The proof is given in Section 7.1, then there is an example in Section 7.2and some comments about the positive-definite case in Section 7.3. Section 8 discusses the *p*-adic classification in higher dimensions. Section 9 formulates the previous theorems concerning matrices in terms of non-degeneracy of critical points of functions. Section 10 applies the work of the previous sections to integrable systems. Section 11 applies the results about integrable systems to the concrete case of the p-adic Jaynes-Cummings system. Section 12 contains some examples and remarks. Section 13 discusses the circle actions generated by the integrable systems of Theorem A. We conclude the paper with an appendix which recalls the basic definitions and results concerning the *p*-adic numbers.

### 2. Main results concerning matrices: Theorems D-I

In this section we state our main classification results concerning normal forms of padic matrices in dimensions 2 and 4: Theorems D-I. We will use these results as stepping stones to prove the results concerning integrable systems (stated in Section 1), but they are also of independent interest and they can be read independently of all the material concerning *p*-adic analytic integrable systems and *p*-adic analytic functions.

2.1. The real Weierstrass-Williamson classification. Let us call  $\Omega_0$  the matrix of the standard symplectic form  $\omega_0$  on  $\mathbb{R}^{2n}$ , that is, a block-diagonal matrix of size 2n with all blocks equal to

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

**Definition 2.1** (Symplectic matrix). A matrix  $S \in \mathcal{M}_{2n}(\mathbb{R})$  is symplectic if  $S^T \Omega_0 S = \Omega_0$ , that is, it leaves invariant the standard symplectic form (the same definition applies if we replace  $\mathbb{R}$  by an arbitrary field F).

For a symmetric matrix  $M \in \mathcal{M}_{2n}(\mathbb{R})$  such that the eigenvalues of  $\Omega_0^{-1}M$  are pairwise distinct, the Weierstrass-Williamson classification says that there is a symplectic matrix S such that  $S^T M S = N$ , where N is a block-diagonal matrix with each block equal to

$$\begin{pmatrix} r_i & 0\\ 0 & r_i \end{pmatrix}, \begin{pmatrix} 0 & r_i\\ r_i & 0 \end{pmatrix} \text{ or } \begin{pmatrix} 0 & r_{i+1} & 0 & r_i\\ r_{i+1} & 0 & -r_i & 0\\ 0 & -r_i & 0 & r_{i+1}\\ r_i & 0 & r_{i+1} & 0 \end{pmatrix},$$

for some  $r_i \in \mathbb{R}$ ,  $1 \leq i \leq n$ , which are called *elliptic block*, hyperbolic block and focus-focus block.

Quite often the Weierstrass-Williamson classification is stated only for positive-definite matrices: in fact the condition that all eigenvalues of  $\Omega_0^{-1}M$  are pairwise distinct is

implied by M being positive definite, and in this case only the elliptic block appears. (For applications to integrable systems the condition on  $\Omega_0^{-1}M$  "translates" to the notion of a critical point being *non-degenerate*; we'll see this later.) In fact, this is the particular case of what is often called Williamson's theorem (that is, what we call the Weierstrass-Williamson classification) which is due to Weierstrass [63] (we learned this fact from the book by Hofer and Zehnder [31, Theorem 8]).

Of course, the case which Williamson treats is much more complicated and interesting: he is able to deal with the completely general situation in which the eigenvalues are not necessarily pairwise distinct. The problem with this case is that it is not feasible to write all the possibilities, for arbitrary dimension, in a compact form because the size of the blocks which are needed can increase without bound. However Williamson does provide a complete list of 4-by-4 matrix normal forms at the end of his paper [66]. These are the possibilities, expressed in a different way to align with the conventions of our paper:

$$\begin{pmatrix} 0 & r & 0 & 0 \\ r & 0 & 0 & 0 \\ 0 & 0 & 0 & s \\ 0 & 0 & s & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & 0 & s \\ 0 & 0 & s & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & s \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & r \\ 0 & 0 & r & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & r \\ 0 & 0 & 0 & a \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & a \end{pmatrix}, \begin{pmatrix} 0 & r & 0 & 0 \\ r & 0 & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & s \end{pmatrix}, \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & r & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & s \end{pmatrix}, \begin{pmatrix} a & 0 & 0 & r \\ 0 & 0 & -r & 0 \\ 0 & -r & a & 0 \\ r & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & s & 0 & r \\ s & 0 & -r & 0 \\ 0 & -r & 0 & s \\ r & 0 & s & 0 \end{pmatrix}$$

where  $r, s \in \mathbb{R}$  and  $a, b \in \{1, -1\}$ . Since in the present paper we only provide a classification of *p*-adic 4-by-4 matrices (also of *p*-adic 2-by-2 matrices, but this case is simpler), it is precisely the list above which is most relevant to us.

2.2. The *p*-adic Weierstrass-Williamson classification. In this article, we will give such a classification in the case when the real field  $\mathbb{R}$  is replaced by the *p*-adic field  $\mathbb{Q}_p$ , where *p* is an arbitrary prime number, and *V* has dimension 2 or 4. Recall that the field of *p*-adic numbers is defined as follows: it is the completion of  $\mathbb{Q}$  with respect to a non-archimedean absolute value defined as

$$|x|_p = p^{-\operatorname{ord}_p(x)},$$

where  $\operatorname{ord}_p(x)$  is the greatest power of p that divides x. We review the construction of basic properties of the field of p-adic numbers in Appendix A.1.

The classification is completely different for p = 2 than  $p \neq 2$ ; for the latter case, in turn, it depends on the class of p modulo 4.

Our main results are the following classifications of *p*-adic symmetric matrices.

**Theorem D** (p-adic classification, 2-by-2 case). Let p be a prime number. Let  $M \in \mathcal{M}_2(\mathbb{Q}_p)$  be a symmetric matrix. Let  $X_p, Y_p$  be the non-residue sets in Definition 1.1. Then, there exists a symplectic matrix  $S \in \mathcal{M}_2(\mathbb{Q}_p)$  and either  $c \in X_p$  and  $r \in \mathbb{Q}_p$ , or c = 0 and  $r \in Y_p \cup \{1\}$ , such that

$$S^T M S = \begin{pmatrix} r & 0\\ 0 & cr \end{pmatrix}.$$

Furthermore, if two symplectic matrices S and S' reduce M to the normal form of the right hand side of the equality above, then the two normal forms are equal.

In Theorem D we note that we are not saying that the value of (c, r) is unique (which essentially is, but not quite), but that the canonical matrix obtained at the right-hand side is unique.

In the statement below, and also Theorem H, we use the following terminology.

**Definition 2.2.** Let *n* be a positive integer. Let *p* be a prime number. We say that two (2n)-by-(2n) matrices *M* and *M'* with coefficients in  $\mathbb{Q}_p$  are equal up to multiplication by a symplectic matrix if there exists a (2n)-by-(2n) symplectic matrix *S* with coefficients in  $\mathbb{Q}_p$  such that  $S^TMS = M'$  (same definition works for arbitrary fields).

**Theorem E** (Number of inequivalent *p*-adic 2-by-2 matrix normal forms). Let *p* be a prime number. Let  $X_p, Y_p$  be the non-residue sets in definition 1.1. Then the following statements hold.

(1) If  $p \equiv 1 \mod 4$ , there are exactly 7 infinite families of normal forms of 2-by-2 *p*-adic matrices with one degree of freedom up to multiplication by a symplectic matrix:

$$\Big\{\Big\{r\begin{pmatrix}1&0\\0&c\end{pmatrix}:r\in\mathbb{Q}_p\Big\}:c\in X_p\Big\},$$

and exactly 4 isolated normal forms, which correspond to c = 0:

$$\Big\{ \begin{pmatrix} r & 0 \\ 0 & 0 \end{pmatrix} : r \in Y_p \cup \{1\} \Big\}.$$

- (2) If  $p \equiv 3 \mod 4$ , there are exactly 5 infinite families of normal forms of 2-by-2 p-adic matrices with one degree of freedom up to multiplication by a symplectic matrix, with the same formula as above, and exactly 4 isolated normal forms.
- (3) If p = 2, there are exactly 11 infinite families of normal forms of 2-by-2 p-adic matrices with one degree of freedom up to multiplication by a symplectic matrix, also with the same formula, and exactly 8 isolated normal forms.

This is in contrast with the real case, where there are exactly 2 families, elliptic and hyperbolic, and 2 isolated normal forms. Here by "infinite family" we mean a family of normal forms of the form  $r_1M_1 + r_2M_2 + \ldots + r_kM_k$ , where  $r_i \in \mathbb{Q}_p$  are parameters and k is the number of degrees of freedom.

Hence, already in dimension 2, the *p*-adic situation is much richer than its real counterpart. The situation is even more surprising in dimension 4. This is the classification in the case where  $\Omega_0^{-1}M$  has all eigenvalues distinct, where  $\Omega_0$  is the same matrix as before for dimension 4:

$$\Omega_0 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}$$

**Theorem F** (*p*-adic classification, 4-by-4, non-degenerate case). Let *p* be a prime number. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $(\mathbb{Q}_p)^4$ . Let  $X_p, Y_p$  be the nonresidue sets in Definition 1.1. Let  $M \in \mathcal{M}_4(\mathbb{Q}_p)$  be a symmetric matrix such that all the eigenvalues of  $\Omega_0^{-1}M$  are distinct. Then there exists a symplectic matrix  $S \in \mathcal{M}_4(\mathbb{Q}_p)$ and  $r, s \in \mathbb{Q}_p$  such that one of the following three possibilities holds: (1) There exist  $c_1, c_2 \in X_p$  such that

$$S^T M S = \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & c_1 r & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & c_2 s \end{pmatrix}.$$

(2) There exists  $c \in Y_p$  such that

$$S^{T}MS = \begin{pmatrix} 0 & s & 0 & r \\ s & 0 & cr & 0 \\ 0 & cr & 0 & s \\ r & 0 & s & 0 \end{pmatrix}$$

(3) There exist  $c, t_1$  and  $t_2$  corresponding to one row of the Table 1 such that  $S^T M S$  is equal to the matrix

$$\begin{pmatrix} \frac{ac(r-bs)}{c-b^2} & 0 & \frac{sc-rb}{c-b^2} & 0\\ 0 & \frac{-r(t_1+bt_2)-s(bt_1+ct_2)}{a} & 0 & -r(bt_1+ct_2)-sc(t_1+bt_2)\\ \frac{sc-rb}{c-b^2} & 0 & \frac{r-bs}{a(c-b^2)} & 0\\ 0 & -r(bt_1+ct_2)-sc(t_1+bt_2) & 0 & ac(-r(t_1+bt_2)-s(bt_1+ct_2)) \end{pmatrix}$$

where (a, b) are either (1, 0) or  $(a_1, b_1)$  of the corresponding row.

Furthermore, if two matrices S and S' reduce M to one of the normal forms in the righthand side of the three equalities above, then the two normal forms are in the same case; if it is case (2) or (3), they coincide, and in case (1) they coincide up to exchanging the 2 by 2 diagonal blocks.

See Figure 4 for a diagram of the classes in the statement of Theorem F.


**Definition 2.3** (Non-residue function). Let p be a prime number. If  $p \equiv 1 \mod 4$ , let  $c_0$  be the smallest quadratic non-residue modulo p. We define the *non-residue function*:

$$h_p: Y_p \to \mathbb{Q}_p \text{ given by } \begin{cases} h_p(c_0) = p, h_p(p) = h_p(c_0p) = c_0 & \text{if } p \equiv 1 \mod 4; \\ h_p(-1) = p, h_p(p) = h_p(-p) = -1 & \text{if } p \equiv 3 \mod 4; \\ h_p(-1) = h_p(-2) = h_p(3) = h_p(6) = -1, \\ h_p(-3) = h_p(-6) = 2, h_p(2) = 3 & \text{if } p = 2. \end{cases}$$

For the cases where the eigenvalues of  $\Omega_0^{-1}M$  are not pairwise distinct, we have:

**Theorem G** (p-adic classification, 4-by-4, degenerate case). Let p be a prime number. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $(\mathbb{Q}_p)^4$ . Let  $X_p, Y_p$  be the nonresidue sets in Definition 1.1. Let  $h_p: Y_p \to \mathbb{Q}_p$  be the non-residue function in Definition 2.3. Let  $M \in \mathcal{M}_4(\mathbb{Q}_p)$  a symmetric matrix such that  $\Omega_0^{-1}M$  has at least one multiple eigenvalue. Then there exists a symplectic matrix  $S \in \mathcal{M}_4(\mathbb{Q}_p)$  such that one of the following three possibilities holds:

(1) There exist  $r, s \in \mathbb{Q}_p$  and  $c_1, c_2 \in X_p \cup \{0\}$  such that  $S^T M S$  has the form in the case (1) of Theorem F. Moreover, if  $c_1 = 0$  then  $r \in Y_p \cup \{1\}$ , and if  $c_2 = 0$  then  $s \in Y_p \cup \{1\}$ .



1

-1 p


1


1


-1

p

-p










FIGURE 4. A diagram of the normal forms of Theorem F for  $p \equiv 1 \mod 4$  (first row),  $p \equiv 3 \mod 4$  (second row) and p = 2 (third to fifth row). Each point (if it is in a single cell) or line (if it is in two cells) represents a normal form. The numbers below the tables represent the first extension of  $\mathbb{Q}_p$  (the one containing the squares of the eigenvalues of  $\Omega_0^{-1}M$ ) and those in the rows and columns represent the second extension (for the eigenvalues themselves). In the first table in each block, there are two such extensions corresponding to the row and the column; in the other ones, there is one extension obtained multiplying the numbers in the row and column.

(2) There exists  $r \in \mathbb{Q}_p$  such that

$$S^T M S = \begin{pmatrix} 0 & r & 0 & 0 \\ r & 0 & 1 & 0 \\ 0 & 1 & 0 & r \\ 0 & 0 & r & 0 \end{pmatrix}$$

(3) There exist  $r \in \mathbb{Q}_p$ ,  $c \in Y_p$  and  $a \in \{1, h_p(c)\}$  such that

$$S^T M S = \begin{pmatrix} a & 0 & 0 & r \\ 0 & 0 & cr & 0 \\ 0 & cr & a & 0 \\ r & 0 & 0 & 0 \end{pmatrix}.$$

(4) There exists  $c \in Y_p \cup \{1\}$  such that

$$S^T M S = \begin{pmatrix} c & 0 & 0 & 0 \\ 0 & 0 & 0 & -c \\ 0 & 0 & 0 & 0 \\ 0 & -c & 0 & 0 \end{pmatrix}.$$

Furthermore, if two matrices S and S' reduce M to one of these normal forms on the right-hand side of the three equalities above, then the two normal forms are in the same case; if it is case (2), (3) or (4), they coincide completely, and in case (1) they coincide up to exchanging the 2 by 2 diagonal blocks.

Our proof method is different from Williamson's method and in particular gives another proof (self-contained, while Williamson's is not, as his proof relies on some applications of other substantial works which he cites in his paper [66]) of the classical Weierstrass-Williamson classification in any dimension. We carry this out in Section 7.

**Theorem H** (Number of 4-by-4 *p*-adic matrix normal forms). Let *p* be a prime number. Let  $X_p, Y_p$  be the non-residue sets in Definition 1.1. Let  $h_p : Y_p \to \mathbb{Q}_p$  be the non-residue function in Definition 2.3.

- (1) If  $p \equiv 1 \mod 4$ , there are exactly 49 infinite families of normal forms of p-adic 4-by-4 matrices with two degrees of freedom, exactly 35 infinite families with one degree of freedom, and exactly 20 isolated normal forms, up to multiplication by a symplectic matrix.
- (2) If p ≡ 3 mod 4, there are exactly 32 infinite families of normal forms of p-adic 4-by-4 matrices with two degrees of freedom, exactly 27 infinite families with one degree of freedom, and exactly 20 isolated normal forms, up to multiplication by a symplectic matrix.
- (3) If p = 2, there are exactly 211 infinite families of normal forms of p-adic 4by-4 matrices with two degrees of freedom, exactly 103 infinite families with one degree of freedom, and exactly 72 isolated normal forms, up to multiplication by a symplectic matrix.

In the three cases above, the infinite families with two degrees of freedom are given by

$$\left\{ \left\{ \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & c_1 r & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & c_2 s \end{pmatrix} : r, s \in \mathbb{Q}_p \right\} : c_1, c_2 \in X_p \right\} \cup \left\{ \left\{ \begin{pmatrix} 0 & s & 0 & r \\ s & 0 & cr & 0 \\ 0 & cr & 0 & s \\ r & 0 & s & 0 \end{pmatrix} : r, s \in \mathbb{Q}_p \right\} : c \in Y_p \right\}$$

$$\cup \Big\{ \Big\{ \begin{cases} \frac{ac(r-bs)}{c-b^2} & 0 & \frac{sc-rb}{c-b^2} & 0 \\ 0 & \frac{-r(t_1+bt_2)-s(bt_1+ct_2)}{a} & 0 & -r(bt_1+ct_2)-sc(t_1+bt_2) \\ \frac{sc-rb}{c-b^2} & 0 & \frac{r-bs}{a(c-b^2)} & 0 \\ 0 & -r(bt_1+ct_2)-sc(t_1+bt_2) & 0 & ac(-r(t_1+bt_2)-s(bt_1+ct_2)) \\ r,s \in \mathbb{Q}_p \Big\} : (a,b) \in \Big\{ (1,0), (a_1,b_1) \Big\}, c, t_1, t_2, a_1, b_1 \text{ in one row of Table 1} \Big\},$$

those with one degree of freedom are

$$\left\{ \left\{ \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & c_1 r & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} : r \in \mathbb{Q}_p \right\} : c_1 \in X_p, s \in Y_p \cup \{1\} \right\} \cup \left\{ \left\{ \begin{pmatrix} 0 & r & 0 & 0 \\ r & 0 & 1 & 0 \\ 0 & 1 & 0 & r \\ 0 & 0 & r & 0 \end{pmatrix} : r \in \mathbb{Q}_p \right\} \right\}$$
$$\cup \left\{ \left\{ \begin{pmatrix} a & 0 & 0 & r \\ 0 & 0 & cr & 0 \\ 0 & cr & a & 0 \\ r & 0 & 0 & 0 \end{pmatrix} : r \in \mathbb{Q}_p \right\} : c \in Y_p, a \in \{1, h_p(c)\} \right\},$$

and the isolated forms are

$$\Big\{ \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} : r, s \in Y_p \cup \{1\} \Big\} \cup \Big\{ \begin{pmatrix} c & 0 & 0 & 0 \\ 0 & 0 & 0 & -c \\ 0 & 0 & 0 & 0 \\ 0 & -c & 0 & 0 \end{pmatrix} : c \in Y_p \cup \{1\} \Big\}.$$

This is in contrast with the real case, where there are exactly 4 infinite families with two degrees of freedom, exactly 7 infinite families with one degree of freedom and exactly 6 isolated normal forms. Here by "infinite family" we mean a family of normal forms of the form  $r_1M_1 + r_2M_2 + \ldots + r_kM_k$ , where  $r_i$  are parameters in  $\mathbb{Q}_p$  and k is the number of degrees of freedom, and by "isolated" we mean a form that is not part of any family.

**Remark 2.4.** Note that Theorems E and H refer to infinite families of matrices, but Theorem B does not mention families. This is because the equivalence between integrable systems we consider allows for a matrix B to appear, as in equation (1.1).


We can give a lower bound for the number of blocks of size 2n that can appear in the normal forms of the matrices. In the real case, taking into account only the "nondegenerate case" where the eigenvalues of  $\Omega_0^{-1}M$  are pairwise distinct, there are two infinite families of blocks with size two (each one with one degree of freedom), one family with size four (with two degrees of freedom) and no blocks with size greater than four: this is due to the fact that all irreducible polynomials in  $\mathbb{R}$  have degree at most two, and has as a consequence that the number of families of normal forms of size 2n is quadratic in n. For the p-adic case, however, one has:

**Definition 2.5.** Let p be a prime number and let n be a positive integer. For each partition  $P = (a_1, \ldots, a_k)$  of n, we define  $M(P, p) \in \mathcal{M}_{2n}(\mathbb{Q}_p)$  as the block diagonal

2n	$\mathbb{R}$	2		$\mathbb{Q}_2$	Ç.	$\mathbb{P}_3$	(	$Q_5$	Ç.	27
	blocks	forms	blocks	forms	blocks	forms	blocks	forms	blocks	forms
2	2	2	11	11	5	5	7	7	5	5
4	1	4	145	211	17	32	21	49	17	32
6	0	6	2	1883	3	123	3	234	9	129
8	0	9	1	21179	2	495	4	1054	2	525
10	0	12	2	161343	3	1595	3	4021	3	1787
12	0	16	1	1374427	2	5111	4	14493	6	5874
14	0	20	2	9232171	3	14491	3	47462	3	17586
16	0	25	1	65570626	2	40244	4	148087	2	50614
18	0	30	2	397086458	3	103484	3	433330	9	137311
20	0	36	1	2469098766	2	259712	4	1217761	2	359463

TABLE 3. The number of families of real and p-adic normal forms of matrices of order 2n. The numbers in the real case are exact, and come from Remark 8.4. The numbers for the p-adic case, for dimensions 2 and 4, come from Theorems E and H. For the rest of dimensions, the numbers of blocks are only lower bounds coming from Lemma 8.3, and the numbers of forms are obtained summing over the partitions of n. The actual number of forms might be even larger.

matrix whose blocks have sizes  $(2a_1, \ldots, 2a_k)$  and each block has the form



**Theorem I** (Number of (2n)-by-(2n) *p*-adic matrix normal forms). Let *p* be a prime number. Let *n* be a positive integer. The number of *p*-adic families of non-degenerate normal forms of (2n)-by-(2n) matrices up to multiplication by a symplectic matrix, each family being of the form  $r_1M_1 + \ldots + r_nM_n$ , where  $r_i$  are parameters in  $\mathbb{Q}_p$ , grows at least with

$$\frac{\mathrm{e}^{\pi\sqrt{2n/3}}}{4n\sqrt{3}}$$

Explicitly, if P and P' are distinct partitions of n then the matrices M(P,p) and M(P',p)in Definition 2.5 are not equivalent by multiplication by a symplectic matrix.

We will later prove this result using the Hardy-Ramanujan formula [30]. We have computed explicit lower bounds for the number of families in Table 3.

#### 3. Symplectic linear algebra over algebraically closed fields

We start by studying the classification problem of normal forms of matrices in an algebraically closed field. This case is an essential ingredient of our strategy to obtain general classifications in the real and *p*-adic cases later on. As we will see, the problem in this case reduces to an equality of eigenvalues, and to an equality of normal forms if there are multiple eigenvalues.

**Definition 3.1.** Let *n* be a positive integer. Let *F* be a field with multiplicative identity element 1. We define  $\Omega_0$  as the (2n)-by-(2n) matrix whose blocks are all

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

that is,

$$\Omega_0 = \begin{pmatrix} 0 & 1 & & & & \\ -1 & 0 & & & & \\ & 0 & 1 & & & \\ & & -1 & 0 & & \\ & & & \ddots & & \\ & & & & 0 & 1 \\ & & & & -1 & 0 \end{pmatrix}$$

(It is also common to take  $\Omega_0$  to have the blocks in the "other" diagonal.)  $\Omega_0$  is called the standard symplectic form on  $F^{2n}$ .

**Proposition 3.2.** Let n be a positive integer. Let F be an algebraically closed field. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^{2n}$ . Let  $M_1, M_2 \in \mathcal{M}_{2n}(F)$  be symmetric matrices,

$$A_i = \Omega_0^{-1} M_i,$$

 $J_i$  the Jordan form of  $A_i$ , and let  $\Psi_i$  be such that  $\Psi_i^{-1}A_i\Psi_i = J_i$ , for  $i \in \{1, 2\}$ . Then there exists a symplectic matrix S such that  $S^T M_1 S = M_2$  if and only if  $J_1 = J_2$ . Moreover, in that case S must have the form  $\Psi_1 D \Psi_2^{-1}$ , where D is a matrix that commutes with  $J_1 = J_2$  and satisfies  $D^T \Psi_1^T \Omega_0 \Psi_1 D = \Psi_2^T \Omega_0 \Psi_2$ .

*Proof.* Let  $A_i = \Omega_0^{-1} M_i$ . Suppose first that such a S exists. Then,

$$S^{-1}A_1S = S^{-1}\Omega_0^{-1}M_1S = S^{-1}\Omega_0^{-1}(S^T)^{-1}S^TM_1S = \Omega_0^{-1}M_2 = A_2$$

hence  $A_1$  and  $A_2$  are equivalent, and  $J_1 = J_2$ .

Let  $D = \Psi_1^{-1} S \Psi_2$ . We have that  $S^T \Omega_0 S = \Omega_0$ , which implies

$$D^T \Psi_1^T \Omega_0 \Psi_1 D = \Psi_2^T \Omega_0 \Psi_2.$$

Also,

$$J_1D = \Psi_1^{-1}A_1\Psi_1D = \Psi_1^{-1}A_1S\Psi_2 = \Psi^{-1}SA_2\Psi_2 = D\Psi_2^{-1}A_2\Psi_2 = DJ_2 = DJ_1.$$

Now suppose that  $J_1 = J_2$ , let D be a matrix which satisfies the conditions and let  $S = \Psi_1 D \Psi_2^{-1}$ . The condition  $D^T \Psi_1^T \Omega_0 \Psi_1 D = \Psi_2^T \Omega_0 \Psi_2$  implies that  $S^T \Omega_0 S = \Omega_0$ , that is, S is symplectic. Moreover,

$$S^{-1}A_1S = \Psi_2 D^{-1}\Psi_1^{-1}A_1\Psi_1 D\Psi_2^{-1} = \Psi_2 D^{-1}J_1 D\Psi_2^{-1} = \Psi_2 J_1 \Psi_2^{-1} = A_2$$

which implies

$$S^{T}M_{1}S = S^{T}\Omega_{0}A_{1}S = S^{T}\Omega_{0}SS^{-1}A_{1}S = \Omega_{0}A_{2} = M_{2}$$

as we wanted.

In the case where the eigenvalues are pairwise distinct, the proposition above can be simplified:

**Lemma 3.3.** Let n be a positive integer. Let F be an algebraically closed field with characteristic different from 2. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^{2n}$  and let  $M \in \mathcal{M}_{2n}(F)$  be a symmetric matrix such that the eigenvalues of  $\Omega_0^{-1}M$  are pairwise distinct. Then there exists a basis  $\{u_1, v_1, \ldots, u_n, v_n\}$  of  $F^{2n}$  such that  $u_i$  and  $v_i$  are eigenvectors of  $\Omega_0^{-1}M$  with opposite eigenvalues and in which  $\Omega_0$  is block-diagonal with blocks of size two.

*Proof.* For ease of notation, we write the proof for n = 2, but the proof is the same for any n.

Let  $A = \Omega_0^{-1} M$ . We have that

$$det(\lambda I - A) = det(\lambda I - \Omega_0^{-1}M)$$
  
=  $det(\lambda I - (\Omega_0^{-1}M)^T)$   
=  $det(\lambda I + M\Omega_0^{-1})$   
=  $det(\lambda \Omega_0 + M) det(\Omega_0^{-1})$   
=  $det(\lambda I + \Omega_0^{-1}M)$   
=  $det(\lambda I + A).$ 

This implies that the eigenvalues of A must come in pairs, that is, if  $\lambda$  is an eigenvalue,  $-\lambda$  also is. In particular, 0 is not an eigenvalue, because it would be at least double, contradicting the hypothesis. So the eigenvalues are  $\lambda, -\lambda, \mu, -\mu$ , for  $\lambda, \mu \in F^*$ .

Now let  $w_1$  and  $w_2$  be two eigenvectors of A with eigenvalues  $\alpha_1$  and  $\alpha_2$ , such that  $\alpha_1 \neq -\alpha_2$ . Then

$$\alpha_1 w_1^T \Omega_0 w_2 = (\Omega_0^{-1} M w_1)^T \Omega_0 w_2$$
  
=  $w_1^T M (-\Omega_0^{-1}) \Omega_0 w_2$   
=  $-w_1^T M w_2$   
=  $-w_1^T \Omega_0 \Omega_0^{-1} M w_2$   
=  $-\alpha_2 w_1^T \Omega_0 w_2.$ 

As  $\alpha_1 \neq -\alpha_2$ , this implies  $w_1^T \Omega_0 w_2 = 0$ .

Let  $u_1, v_1, u_2, v_2$  be the eigenvectors of  $\lambda, -\lambda, \mu, -\mu$ , respectively. By the previous result,  $w_1^T \Omega_0 w_2 = 0$  for any two vectors  $w_1, w_2 \in \{u_1, v_1, u_2, v_2\}$  which are not a pair  $u_i, v_i$ . We call  $\Psi$  the matrix with  $(u_1, v_1, u_2, v_2)$  as columns. Then,

$$\begin{split} \Psi^T \Omega_0 \Psi &= \begin{pmatrix} 0 & u_1^T \Omega_0 v_1 & 0 & 0 \\ v_1^T \Omega_0 u_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & u_2^T \Omega_0 v_2 \\ 0 & 0 & v_2^T \Omega_0 u_2 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & u_1^T \Omega_0 v_1 & 0 & 0 \\ -u_1^T \Omega_0 v_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & u_2^T \Omega_0 v_2 \\ 0 & 0 & -u_2^T \Omega_0 v_2 & 0 \end{pmatrix}, \end{split}$$

as we wanted.

The basis in Lemma 3.3 is almost but not quite a symplectic basis:

**Definition 3.4.** Let *n* be a positive integer. Let *F* be a field. We say that a basis  $\{u_1, v_1, \ldots, u_n, v_n\}$  of  $F^{2n}$  is symplectic if, for any two vectors  $w_1, w_2$  in the basis,

 $w_1^T \Omega_0 w_2 = 1$  if  $w_1 = u_i$  and  $w_2 = v_i$  for some *i* with  $1 \leq i \leq n$ , and otherwise  $w_1^T \Omega_0 w_2 = 0$ . This condition is equivalent to saying that the matrix in  $\mathcal{M}_{2n}(F)$  with  $u_1, v_1, \ldots, u_n, v_n$  as columns is symplectic.

Actually, we can rescale the vectors  $v_i$  such that the basis becomes symplectic. But this may break the structure of the eigenvectors: for example, if  $-\lambda = \overline{\lambda}$  for some definition of the conjugate, we can take  $v_1 = \overline{u}_1$ , which will no more hold after rescaling  $v_1$ . We leave the lemma as such because we do not need that rescaling.

**Corollary 3.5.** Let n be a positive integer. Let F be an algebraically closed field with characteristic different from 2. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^{2n}$ . Given symmetric matrices  $M_1, M_2 \in \mathcal{M}_{2n}(F)$  such that  $\Omega_0^{-1}M_i$  has pairwise distinct eigenvalues for  $i \in \{1, 2\}$ , there is a symplectic matrix S such that  $S^T M_1 S = M_2$  if and only if  $\Omega_0^{-1}M_1$  and  $\Omega_0^{-1}M_2$  have the same eigenvalues. Moreover, in this case S must have the form  $\Psi_1 D \Psi_2^{-1}$ , where D is a diagonal matrix such that

(3.1) 
$$d_{2i-1,2i-1}d_{2i,2i} = \frac{(u_i^2)^T \Omega_0 v_i^2}{(u_i^1)^T \Omega_0 v_i^1},$$

 $u_i^1$  and  $v_i^1$  are those of Lemma 3.3 for the first form,  $u_i^2$  and  $v_i^2$  for the second form, and for  $j \in \{1, 2\}$ ,

$$\Psi_j = \begin{pmatrix} u_1^j & v_1^j & \dots & u_n^j & v_n^j \end{pmatrix}$$

*Proof.* By Lemma 3.3, there are  $\Psi_1$  and  $\Psi_2$  such that  $\Psi_i^{-1}A_i\Psi_i = J_i$ , where  $J_i$  is the matrix in Proposition 3.2, and  $\Psi_i^T \Omega_0 \Psi_i$  has all elements zero except those of the form (2i - 1, 2i) and (2i, 2i - 1). Moreover, in this case matrices  $J_1$  and  $J_2$  are diagonal and with all elements in the diagonal different. They are equal if and only if  $A_1$  and  $A_2$  have the same eigenvalues.

A matrix D that commutes with  $J_1 = J_2$  is necessarily diagonal, and the relation

$$D^T \Psi_1^T \Omega_0 \Psi_1 D = \Psi_2^T \Omega_0 \Psi_2$$

in this case reduces to (3.1).

In the case where the eigenvalues are not all different, the situation is not so simple, but we can do something similar to Lemma 3.3.

**Lemma 3.6.** Let n be a positive integer. Let F be an algebraically closed field with characteristic different from 2 and let  $M \in \mathcal{M}_{2n}(F)$  be a symmetric matrix. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^{2n}$ . Let  $A = \Omega_0^{-1}M$ . Then, the number of nonzero eigenvalues of A is even, that is, 2m for some integer m with  $0 \leq m \leq n$ , and there exists a set

$$\left\{u_1, v_1, \dots, u_m, v_m\right\} \subset F^{2n}$$

which satisfies the following properties:

- $Au_i = \lambda_i u_i + \mu_i u_{i-1}$  and  $Av_i = -\lambda_i v_i + \mu_{i+1} v_{i+1}$  for  $1 \leq i \leq m$ , where  $\lambda_i \in F$ ,  $\mu_i = 0$  or 1,  $\mu_1 = \mu_{m+1} = 0$ , and  $\mu_i = 1$  only if  $\lambda_i = \lambda_{i-1}$ . (That is to say, the vectors are a "Jordan basis".)
- The vectors can be completed to a symplectic basis: given  $w_1, w_2$  in the set,  $w_1^T \Omega_0 w_2 = 1$  if  $w_1 = u_i, w_2 = v_i$  for some i with  $1 \leq i \leq m$ , and otherwise  $w_1^T \Omega_0 w_2 = 0.$

*Proof.* Let J be the Jordan form of A and  $\Psi$  such that  $\Psi^{-1}A\Psi = J$ . We have that

$$J = \Psi^{-1} A \Psi$$
  
=  $\Psi^{-1} \Omega_0^{-1} M \Psi$   
=  $\Psi^{-1} \Omega_0^{-1} M \Omega_0^{-1} \Omega_0 \Psi$   
=  $\Psi^{-1} \Omega_0^{-1} (-\Omega_0^{-1} M)^T \Omega_0 \Psi$   
=  $\Psi^{-1} \Omega_0^{-1} (-\Psi J \Psi^{-1})^T \Omega_0 \Psi$   
=  $(\Psi^T \Omega_0 \Psi)^{-1} (-J^T) \Psi^T \Omega_0 \Psi$ 

But J can only be similar to  $-J^T$  if for each block having  $\lambda$  in the diagonal there is another having  $-\lambda$  in the diagonal, and with the same size. We can split the blocks in three parts, that is, there is a  $\Phi$  such that

(3.2) 
$$\Phi^{-1}A\Phi = \begin{pmatrix} J_+ & 0 & 0\\ 0 & J_- & 0\\ 0 & 0 & J_0 \end{pmatrix},$$

for some matrices  $J_+$ ,  $J_-$  and  $J_0$  in Jordan form, such that  $J_+$  and  $J_-$  have opposite eigenvalues and  $J_0$  has only 0 as eigenvalue. Let m be the size of  $J_+$  and  $J_-$ . Now 2m is the number of nonzero eigenvalues.

Let the first m columns of  $\Phi$  be  $u_1, \ldots, u_m$ . Because of (3.2), we have

$$Au_i = \lambda_i u_i + \mu_i u_{i-1},$$

for adequate  $\lambda_i$  and  $\mu_i$ , with  $\mu_i = 0$  or 1.

Now we change sign and transpose, getting

(3.3) 
$$-(\Phi^{-1}A\Phi)^T = \begin{pmatrix} -J_+^T & 0 & 0\\ 0 & -J_-^T & 0\\ 0 & 0 & -J_0^T \end{pmatrix}$$

The left-hand side equals

$$\begin{split} -\Phi^T A^T (\Phi^{-1})^T &= \Phi^T M \Omega_0^{-1} (\Phi^{-1})^T \\ &= \Phi^T \Omega_0 A \Omega_0^{-1} (\Phi^{-1})^T \\ &= \Phi^T \Omega_0 A (\Phi^T \Omega_0)^{-1}. \end{split}$$

Again, let the first m columns of  $(\Phi^T \Omega_0)^{-1}$  be  $v_1, \ldots, v_m$ . By (3.3), we have

$$Av_i = -\lambda_i v_i - \mu_{i+1} v_{i+1}$$

It is left to prove that the 2m vectors form a partial symplectic basis. To do this, first we see that  $u_i^T \Omega_0 u_j = 0$  for any i, j, by induction on i + j. The base case is when i = j = 1, which is trivial. Supposing it true for (i - 1, j) and (i, j - 1), we prove it for (i, j):

$$\lambda_i u_i^T \Omega_0 u_j = (\Omega_0^{-1} M u_i - \mu_i u_{i-1})^T \Omega_0 u_j$$
  
$$= u_i^T M (-\Omega_0^{-1}) \Omega_0 u_j - \mu_i \cdot 0$$
  
$$= -u_i^T M u_j$$
  
$$= -u_i^T \Omega_0 \Omega_0^{-1} M u_j$$
  
$$= -u_i^T \Omega_0 (\lambda_j u_j + \mu_j u_{j-1})$$
  
$$= -\lambda_j u_i^T \Omega_0 u_j$$

which implies that  $u_i^T \Omega_0 u_j = 0$  because  $\lambda_i \neq -\lambda_j$  (the opposites of the eigenvalues in the part  $J_+$  are all in  $J_-$ ). Analogously we prove that  $v_i^T \Omega_0 v_j = 0$ , making the induction backwards.

Finally,  $u_i^T \Omega_0 v_j$  is the element in the position (i, j) of  $\Phi^T \Omega_0 (\Phi^T \Omega_0)^{-1}$ , which is 1 if i = j and 0 otherwise, so the vectors are a partial symplectic basis.

Lemma 3.6 allows us to separate the part of a matrix with nonzero eigenvalues from the part with zero eigenvalues. Choosing a symplectic basis for the latter to complete the partial basis in a way that we obtain a complete "Jordan basis" is more complicated, but it can also be done, as the following theorem shows.

**Definition 3.7.** A tuple  $K = (k_1, k_2, ..., k_t)$  of natural numbers is called *good* if the numbers are in non-increasing order, that is,  $k_i \ge k_{i+1}$  for every  $1 \le i \le t-1$ , and each odd number appears an even number of times in K. We define  $f_K : \{1, ..., t\} \to \{1, ..., t\}$  as follows:

• If  $k_i$  is even,  $f_K(i) = i$ .

• If  $k_i$  is odd, let  $i_0$  be the first index such that  $k_{i_0} = k_i$ . Then,  $f_K(i) = i + (-1)^{i-i_0}$ .

If K is good,  $f_K$  is an involution that fixes the indices of even elements and pairs those of odd elements.

**Theorem 3.8.** Let n be a positive integer. Let F be a field with characteristic different from 2 and let  $M \in \mathcal{M}_{2n}(F)$  be a symmetric matrix. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^{2n}$ . Let  $A = \Omega_0^{-1}M$ . Suppose that the eigenvalues of A are all zero. Then, there is an integer  $t \ge 1$ , a good tuple  $K = (k_1, \ldots, k_t)$  and a basis of  $F^{2n}$  of the form

(3.4) 
$$\left\{u_{11}, u_{12}, \dots, u_{1k_1}, \dots, u_{t1}, u_{t2}, \dots, u_{tk_t}\right\}$$

such that the following two conditions hold:

- For every  $1 \leq i \leq t, 1 \leq j \leq k_t$ ,  $Au_{ij} = u_{i,j-1}$ , where  $u_{i0} = 0$ .
- Given  $u_{ij}, u_{i'j'}$  in this basis,  $u_{ij}^T \Omega_0 u_{i'j'} \neq 0$  if and only if  $i' = f_K(i)$  and  $j + j' = k_i + 1$ .

See Table 4, left, for an example of such a basis.

To prove this theorem, we need several intermediate results.

**Definition 3.9.** Given a tuple  $K = (k_1, \ldots, k_t)$  such that  $k_i \ge k_{i+1}$  for  $1 \le i \le t-1$ , we call  $R_K$  the set of tuples of the form  $(\ell, i, j)$  where  $1 \le \ell \le t, \ell \le i \le t, 1 \le j \le k_i$ .

Given  $R \subset R_K$ , we say that a basis is *R*-acceptable if, for each  $(\ell, i, j) \in R$ , one of these alternatives holds:

•  $j = 1, i = f_K(\ell), \ell = f_K(i), \text{ and } u_{\ell k \ell}^T \Omega_0 u_{ij} \neq 0;$ 

• 
$$j \neq 1$$
 or  $i \neq f_K(\ell)$ , and  $u_{\ell k_\ell}^T \Omega_0 u_{ij} = 0$ .

(Note that, if there exists  $(\ell, i, j) \in R$  with j = 1,  $i = f_K(\ell)$  but  $\ell \neq f_K(i)$ , no basis is *R*-acceptable.)

Given  $(\ell, i, j), (\ell', i', j') \in R_K$ , we say that  $(\ell, i, j) < (\ell', i', j')$  if

- $\ell < \ell'$ ,
- $\ell = \ell'$  and j < j', or
- $\ell = \ell', j = j'$  and i < i'.

Note that < is a total order in  $R_K$ .

For each s integer with  $0 \leq s \leq |R_K|$ , we call  $R_{K,s}$  the set of the first s elements of  $R_K$  according to the order <.

$u_{16}$							$u_{16}$					
$u_{15}$	$u_{25}$	$u_{35}$					$u_{15}$	$u_{25}$	$u_{35}$			
$u_{14}$	$u_{24}$	$u_{34}$	$u_{44}$				$u_{14}$	$u_{24}$	$u_{34}$	$u_{44} + cu_{23}$		
$u_{13}$	$u_{23}$	$u_{33}$	$u_{43}$	$u_{53}$	$u_{63}$	$\rightarrow$	$u_{13}$	$u_{23}$	$u_{33}$	$u_{43} + cu_{22}$	$u_{53}$	$u_{63}$
$u_{12}$	$u_{22}$	$u_{32}$	$u_{42}$	$u_{52}$	$u_{62}$		$u_{12}$	$u_{22}$	$u_{32}$	$u_{42} + cu_{21}$	$u_{52}$	$u_{62}$
$u_{11}$	$u_{21}$	$u_{31}$	$u_{41}$	$u_{51}$	$u_{61}$		$u_{11}$	$u_{21}$	$u_{31}$	$u_{41}$	$u_{51}$	$u_{61}$

TABLE 4. Left: a basis B of  $F^{26}$  in the form (3.4) for K = (6, 5, 5, 4, 3, 3). Right: the basis B[2, 4, 2, c], for  $c \in F$ .

Let J be the Jordan form of A. Since all eigenvalues of A are zero, J has all entries equal to zero except some in the first diagonal above the main diagonal, which are 1. That is, J is a block-diagonal matrix. Let  $k_1, \ldots, k_t$  be the sizes of the blocks, with  $k_i \ge k_{i+1}$ , and  $K = (k_1, \ldots, k_t)$ . For this K, any Jordan basis satisfies the first condition.

**Lemma 3.10.** If  $\{u_{ij}\}$  is a Jordan basis for A in the form (3.4),  $u_{ij}^T \Omega_0 u_{i'j'} = -u_{i,j-1}^T \Omega_0 u_{i',j'+1}$ . That is, the product  $u_{ij}^T \Omega_0 u_{i'j'}$  only depends on j + j', except for the sign.

*Proof.* Using that M is symmetric and  $\Omega_0$  is antisymmetric,

$$u_{ij}^{T}\Omega_{0}u_{i'j'} = u_{ij}^{T}\Omega_{0}Au_{i',j'+1} = u_{ij}^{T}Mu_{i',j'+1} = -u_{ij}^{T}A^{T}\Omega_{0}u_{i',j'+1} = -u_{i,j-1}^{T}\Omega_{0}u_{i',j'+1}.$$

**Corollary 3.11.** Let us assume the conditions of Lemma 3.10. If  $j + j' \leq \max\{k_i, k_{i'}\}$ ,  $u_{ij}^T \Omega_0 u_{i'j'} = 0$ .

*Proof.* Without loss of generality suppose that  $k_i \leq k_{i'}$ . Applying the previous lemma j times, we have

$$u_{ij}^{T}\Omega_{0}u_{i'j'} = u_{i0}^{T}\Omega_{0}u_{i',j+j'} = 0.$$

**Corollary 3.12.** If a basis  $\{u_{ij}\}$  of the form (3.4) is *R*-acceptable for some *R* which contains  $(\ell, f_K(\ell), 1)$  or  $(f_K(\ell), \ell, 1)$ , then  $u_{\ell k_\ell}^T \Omega_0 u_{f_K(\ell), 1} \neq 0$ .

Proof. If  $(\ell, f_K(\ell), 1) \in R$ , then the conclusion holds by definition. If  $(f_K(\ell), \ell, 1) \in R$ , then  $f_K(\ell) \leq \ell$ , which implies  $k_\ell = f_K(\ell)$ , and  $u_{\ell k_\ell}^T \Omega_0 u_{f_K(\ell), 1} = (-1)^{k_\ell - 1} u_{\ell 1}^T \Omega_0 u_{f_K(\ell), k_\ell} \neq 0$ .

We will need this operation to change a basis:

**Definition 3.13.** Let *B* be a basis of  $F^{2n}$  of the form (3.4),  $1 \leq \ell \leq t, 1 \leq i \leq t, 1 \leq j \leq k_i$  and  $c \in F$ . We call  $B[\ell, i, j, c]$  the basis

$$\left\{u'_{11}, u'_{12}, \dots, u'_{1k_1}, \dots, u'_{t1}, u'_{t2}, \dots, u'_{tk_t}\right\}$$

where

$$u'_{i'j'} = \begin{cases} u_{i'j'} & \text{if } i' \neq i, \\ u_{ij'} & \text{if } i' = i, j' < j, \\ u_{ij'} + cu_{\ell,j'-j+1} & \text{if } i' = i, j' \ge j. \end{cases}$$

See Table 4 for an example.

**Lemma 3.14.** (1) If B is a Jordan basis for A,  $B[\ell, i, j, c]$  is also a Jordan basis.

- (2) The only products of the form  $u_{\ell'k_{\ell'}}^T \Omega_0 u_{i'j'}$  which may be different in B and  $B[\ell, i, j, c]$ are those with  $i' = i \neq \ell'$  and  $j' \geq j$ , which vary in  $cu_{\ell'k_{\ell'}}^T \Omega_0 u_{\ell,j'-j+1}$ , those with  $\ell' = i \neq i'$ , which vary in  $cu_{\ell,k_i-j+1}^T \Omega_0 u_{i'j'}$ , and those with  $\ell' = i' = i$ , which vary in  $cu_{ik_i}^T \Omega_0 u_{\ell,j'-j+1} + cu_{\ell,k_i-j+1}^T \Omega_0 u_{ij'} + c^2 u_{\ell,k_i-j+1}^T \Omega_0 u_{\ell,j'-j+1}$  (understanding  $u_{ij} = 0$ if  $j \leq 0$  in the last equality).
- *Proof.* (1) We have that  $Au_{i'j'} = u_{i',j'-1}$  and we need to see that  $Au'_{i'j'} = u'_{i',j'-1}$  for all indices  $1 \leq i' \leq t, 1 \leq j' \leq k_{i'}$ . If  $i' \neq i$ , the conclusion follows because the vectors do not change. If i' = i and j' < j, the same happens. Otherwise, i' = i and  $j' \geq j$ , and

$$Au'_{ij'} = A(u_{ij'} + cu_{\ell,j'-j+1}) = u_{i,j'-1} + cu_{\ell,j'-j} = u'_{i,j'-1}$$

where the last equality also holds if j' = j and  $u_{\ell,j'-j} = 0$ .

(2) This follows from the definition of the new basis: to change the product we need to change any of the two vectors. □

**Lemma 3.15.** For every s with  $0 \leq s \leq |R_K|$ , there is a Jordan basis for A which is  $R_{K,s}$ -acceptable.

*Proof.* We prove this by induction in s. For s = 0,  $R_{K,0}$  is empty and the problem reduces to the existence of a Jordan basis.

Supposing it is true for s, we prove it for s + 1. Let  $\{(\ell, i, j)\} = R_{K,s+1} \setminus R_{K,s}$ , that is,  $(\ell, i, j)$  is the (s + 1)-th element of  $R_K$  according to the order <. Let  $B = \{u_{ij}\}$  be the  $R_{K,s}$ -acceptable basis given by the inductive hypothesis.

There are several cases to consider.

(1)  $i = \ell$  and  $k_{\ell} - j$  is even. By Lemma 3.10, we have that

$$u_{\ell k_{\ell}}^{T} \Omega_{0} u_{\ell j} = u_{\ell, (k_{\ell}+j)/2}^{T} \Omega_{0} u_{\ell, (k_{\ell}+j)/2} = 0$$

Either j > 1 and we want a zero, or j = 1 with  $k_{\ell}$  odd and we also want a zero; in any case B itself is  $R_{K,s+1}$ -acceptable.

(2)  $i = f_K(\ell), j = 1$ . This means that i is  $\ell$  if  $k_\ell$  is even and  $\ell + 1$  if  $k_\ell$  is odd (it cannot be  $\ell - 1$  because  $(\ell, i, j) \in R_K$ ). If  $u_{\ell k_\ell}^T \Omega_0 u_{i1} \neq 0$ , B is  $R_{K,s+1}$ -acceptable and we are done. Otherwise, since B is a basis, there are  $1 \leq i_1 \leq t, 1 \leq j_1 \leq k_{i_1}$  such that  $u_{\ell 1}^T \Omega_0 u_{i_1 j_1} \neq 0$ . By Corollary 3.11,  $j_1 + 1 > \max\{k_\ell, k_{i_1}\}$  or equivalently  $j_1 \geq \max\{k_\ell, k_{i_1}\}$ . But this implies  $k_{i_1} \geq j_1 \geq \max\{k_\ell, k_{i_1}\} \geq k_\ell$ , so  $j_1 \geq k_{i_1}$ , which implies  $j_1 = k_{i_1}$  and  $u_{\ell 1}^T \Omega_0 u_{i_1 k_{i_1}} \neq 0$ .

By Lemma 3.10,  $u_{\ell 1}^T \Omega_0 u_{ik_i} = (-1)^{k_i - 1} u_{\ell k_\ell}^T \Omega_0 u_{i1} = 0$ , so  $i \neq i_1$ . If  $i_1 < i$ , either  $i_1 < \ell$  or  $i_1 = \ell$  with  $i = \ell + 1$ . In any case  $(i_1, \ell, 1) < (\ell, i, 1)$ , and since B is  $R_{K,s}$ -acceptable,  $u_{\ell 1}^T \Omega_0 u_{i_1 k_{i_1}} = 0$ , a contradiction. Hence,  $i_1 > i \ge \ell$ . As the sequence K is non-increasing and  $k_{i_1} \ge k_\ell$ , we must have  $k_{i_1} = k_i = k_\ell$ , and again by Lemma 3.10,  $u_{\ell k_\ell}^T \Omega_0 u_{i_1 1} \neq 0$ . If  $i = \ell + 1$ ,  $k_i = k_\ell$  together with  $f_K(\ell) = i$  implies that  $f_K(i) = \ell$ .

Let  $B' = \{u'_{ij}\} = B[i_1, i, 1, 1]$ . This basis is  $R_{K,s}$ -acceptable: by Lemma 3.14, the conditions in  $R_{K,s}$  that may break with this change are  $(\ell', i, j')$  with  $\ell' < \ell$  (the only condition which may be in  $R_{K,s}$  with  $\ell' = \ell$  is  $(\ell, \ell, 1)$ ). For these tuples,

$$(u'_{\ell'k_{\ell'}})^T \Omega_0 u'_{ij'} = u^T_{\ell'k_{\ell'}} \Omega_0 u_{ij'} + u^T_{\ell'k_{\ell'}} \Omega_0 u_{i_1j'} = 0$$

because  $(\ell', i, j')$  and  $(\ell', i_1, j')$  are in  $R_{K,s}$ . Moreover, if  $i = \ell + 1$ ,

$$(u_{\ell k_{\ell}}')^{T} \Omega_{0} u_{i1}' = u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i1} + u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} = u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} \neq 0,$$

so B' is  $R_{K,s+1}$ -acceptable, as we wanted.

If  $i = \ell$ , we also define  $B'' = \{u''_{ij}\} = B[i_1, i, 1, -1]$ . Analogously to what we said for B', B'' is  $R_{K,s}$ -acceptable. Now we have

$$(u_{\ell k_{\ell}}')^{T} \Omega_{0} u_{i1}' = u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i1} + u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i1} + u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i1} + u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} = 2u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} + u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i_{1}1}$$
and

 $(u_{\ell k_{\ell}}'')^{T} \Omega_{0} u_{i1}'' = u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i1} - u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i11} - u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i1} + u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} = -2u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} + u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} + u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} + u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} = -2u_{\ell k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} + u_{i_{1}k_{\ell}}^{T} \Omega_{0} u_{i_{1}1} + u_{i_{1}k_{\ell$ where the second equality in each line is due to Lemma 3.10. If both results were zero at the same time, that would imply  $u_{\ell k_{\ell}}^T \Omega_0 u_{i_1 1} = 0$ , so one of them must be nonzero, and one of B' and B'' must be  $R_{K,s+1}$ -acceptable.

(3)  $i = \ell, j > 1$  and  $k_{\ell} - j$  is odd. We set

$$c = -\frac{u_{\ell k_{\ell}}^T \Omega_0 u_{\ell j}}{2u_{\ell k_{\ell}}^T \Omega_0 u_{f_K(\ell),1}}$$

and  $B' = B[f_K(\ell), \ell, j, c]$  (the denominator is not zero by Corollary 3.12, because  $(\ell, f_K(\ell), 1) \in R_{K,s}$  or  $(f_K(\ell), \ell, 1) \in R_{K,s})$ . By Lemma 3.14, the conditions that can break are  $(\ell, i', j')$  with  $\ell \leq i' \leq t$  and  $1 \leq j' \leq j - 1$ , and  $(\ell', \ell, j')$  with  $1 \leq \ell' \leq \ell - 1$  and  $j \leq j' \leq k_i$ . For the first type, we see that

$$(u'_{\ell k_{\ell}})^{T} \Omega_{0} u'_{i'j'} = u^{T}_{\ell k_{\ell}} \Omega_{0} u_{i'j'} + c u^{T}_{f_{K}(\ell), k_{\ell}-j+1} \Omega_{0} u_{i'j'}$$

The second term is 0 because  $k_{\ell} - j + 1 + j' \leq k_{\ell} = k_{f_K(\ell)}$  (we are using here that  $f_K(f_K(\ell)) = \ell$  because the basis is acceptable). For the second type,

$$(u_{\ell'k_{\ell'}}')^T \Omega_0 u_{\ell j'}' = u_{\ell'k_{\ell'}}^T \Omega_0 u_{\ell j'} + c u_{\ell'k_{\ell'}}^T \Omega_0 u_{f_K(\ell),j'-j+1}$$

Both terms are zero because  $(\ell', \ell, j')$  and  $(\ell', f_K(\ell), j'-j+1)$  are in  $R_{K,s}, f_K(\ell') \neq j$  $f_K(\ell)$  and j' > 1. It is only left to show that the new condition is satisfied:

 $(u_{\ell k_{\ell}}')^{T} \Omega_{0} u_{\ell j}' = u_{\ell k_{\ell}}^{T} \Omega_{0} u_{\ell j} + c u_{\ell k_{\ell}}^{T} \Omega_{0} u_{f_{K}(\ell),1} + c u_{f_{K}(\ell),k_{\ell}-j+1}^{T} \Omega_{0} u_{\ell j} + c^{2} u_{f_{K}(\ell),k_{\ell}-j+1}^{T} \Omega_{0} u_{f_{K}(\ell),1}$ The last term is zero because  $k_{\ell} - j + 1 + 1 \leq k_{\ell}$ . The second and the third are equal because, by Lemma 3.10,

$$u_{f_{K}(\ell),k_{\ell}-j+1}^{T}\Omega_{0}u_{\ell j} = -u_{\ell j}^{T}\Omega_{0}u_{f_{K}(\ell),k_{\ell}-j+1} = (-1)^{k_{\ell}-j+1}u_{\ell k_{\ell}}^{T}\Omega_{0}u_{f_{K}(\ell),1} = u_{\ell k_{\ell}}^{T}\Omega_{0}u_{f_{K}(\ell),1}.$$
Hence

пепсе

$$(u_{\ell k_{\ell}}')^{T} \Omega_{0} u_{\ell j}' = u_{\ell k_{\ell}}^{T} \Omega_{0} u_{\ell j} + 2c u_{\ell k_{\ell}}^{T} \Omega_{0} u_{f_{K}(\ell),1} = 0$$

as we wanted.

(4)  $i > \ell$  and either j > 1 or  $i \neq f_K(\ell)$ . We set

$$c = -\frac{u_{\ell k_{\ell}}^T \Omega_0 u_{ij}}{u_{\ell k_{\ell}}^T \Omega_0 u_{f_K(\ell),1}}$$

and  $B' = B[f_k(\ell), i, j, c]$  (the denominator is not zero by Corollary 3.12, because  $(\ell, f_K(\ell), 1) \in R_{K,s}$  or  $(f_K(\ell), \ell, 1) \in R_{K,s})$ . By Lemma 3.14, the conditions that can break are  $(\ell', i, j')$  with  $1 \leq \ell' \leq \ell - 1$  and  $j \leq j' \leq k_i$ . In these cases we have

$$(u'_{\ell'k_{\ell'}})^T \Omega_0 u'_{ij'} = u^T_{\ell'k_{\ell'}} \Omega_0 u_{ij'} + c u^T_{\ell'k_{\ell'}} \Omega_0 u_{f_K(\ell),j'-j+1}$$

Both terms are zero because  $(\ell', i, j')$  and  $(\ell', f_K(\ell), j'-j+1)$  are in  $R_{K,s}, f_K(\ell') \neq 0$  $f_K(\ell)$ , and either j' > 1 or  $i \neq f_K(\ell)$ . It is left to show the new condition:

$$(u'_{\ell k_{\ell}})^{T} \Omega_{0} u'_{ij} = u^{T}_{\ell k_{\ell}} \Omega_{0} u_{ij} + c u^{T}_{\ell k_{\ell}} \Omega_{0} u_{f_{K}(\ell),1} = 0.$$

With these results, we are ready to prove Theorem 3.8.

Proof of Theorem 3.8. We apply Lemma 3.15 with  $s = |R_K|$  to obtain a basis B which is  $R_K$ -acceptable. Now we show that this B is the basis we want. The first condition holds because B is a Jordan basis. For the second, if  $j + j' \leq \max\{k_i, k_{i'}\}$ , the product is zero by Corollary 3.11, as we want. Otherwise, suppose  $i \leq i'$  and  $k_i \geq k_{i'}$ . By Lemma 3.10,

$$u_{ij}^T \Omega_0 u_{i'j'} = u_{ik_i}^T \Omega_0 u_{i',j+j'-k_i} = 0,$$

and the second condition also holds.

It is left to show that K is good, but this is an immediate consequence of B being  $R_K$ -acceptable: by definition, if  $f_K(l) = i$  but  $f_K(i) \neq l$  for some  $(l, i, j) \in R_K$ , there would not be any  $R_K$ -acceptable basis.

We can use Lemma 3.6 and Theorem 3.8 to give a classification in the case where the field is algebraically closed. In particular, we can apply this to  $\mathbb{C}$ .

**Theorem 3.16.** Let n be a positive integer. Let F be an algebraically closed field with characteristic different from 2 and let  $M \in \mathcal{M}_{2n}(F)$  be a symmetric matrix. There exists a positive integer k,  $r \in F$ ,  $a \in \{0, 1\}$ , with a = 1 only if r = 0, and a symplectic matrix  $S \in \mathcal{M}_{2n}(F)$  such that  $S^T M S$  is a block-diagonal matrix whose blocks are of the form

$$M_{\rm h}(k,r,a) = \begin{pmatrix} r & & & & \\ r & 1 & & & \\ & 1 & r & & \\ & r & \ddots & & \\ & & \ddots & & 1 & \\ & & & 1 & r \\ & & & & r & a \end{pmatrix}$$

with 2k rows. Furthermore, if there are two matrices S and S' which reduce M to this form, then the two forms only differ in the order in which the blocks are arranged.

*Proof.* (a) First we prove existence. We start applying Lemma 3.6. This gives us a partial symplectic basis

$$\left\{u_1, v_1, \ldots, u_m, v_m\right\},\$$

which is also a partial Jordan basis of A corresponding to the nonzero eigenvalues, and  $u_i$  and  $v_i$  correspond to opposite eigenvalues  $\lambda_i$  and  $-\lambda_i$ . If  $\{u_1, v_1, \ldots, u_k, v_k\}$ are the vectors of a block with eigenvalues r and -r, for  $r \in F$ , these same vectors taken as columns of S give the matrix  $M_h(k, r, 0)$ .

For the other part of the Jordan form, we apply Theorem 3.8 to the eigenspace of 0, resulting in a good multiset  $K = \{k_1, \ldots, k_t\}$  and a basis

$$\left\{u_{11}, u_{12}, \ldots, u_{1k_1}, \ldots, u_{t1}, u_{t2}, \ldots, u_{tk_t}\right\}$$

This is not necessarily a symplectic basis as such, but it allows us to construct one with the required properties:

• If  $k_i = 2\ell_i$  is even, we have that

$$u_{ij}^T \Omega_0 u_{i,k_i+1-j} = (-1)^{j+1} c_i$$

for some  $c_i \in F$ . Let

$$c_i = b_i^2$$

for  $b_i \in F$ . After dividing all the chain by  $b_i$ , we can assume that  $c_i = (-1)^{\ell_i}$ . Now

$$\left\{u_{i1},(-1)^{\ell_i}u_{i,2\ell_i},-u_{i2},(-1)^{\ell_i}u_{i,2\ell_i-1},\ldots,(-1)^{\ell_i-1}u_{i\ell_i},(-1)^{\ell_i}u_{i,\ell_i+1}\right\}$$

is a partial symplectic basis which gives the form  $M_{\rm h}(\ell_i, 0, 1)$ .

• If  $k_i$  is odd and  $f_K(i) = i + 1$ , we have instead

$$u_{ij}^T \Omega_0 u_{i+1,k_i+1-j} = (-1)^{j+1} c_i$$

for some  $c_i \in F$ . After dividing the elements of the second chain by  $c_i$ , we can assume that  $c_i = 1$ . Now

$$\left\{u_{i1}, u_{i+1,k_i}, -u_{i2}, u_{i+1,k_i-1}, \dots, (-1)^{k_i-1}u_{ik_i}, u_{i+1,1}\right\}$$

is a partial symplectic basis which gives the form  $M_{\rm h}(k_i, 0, 0)$ .

(b) Uniqueness follows from Proposition 3.2, because two matrices in normal form have the same Jordan form if and only if they differ in the order of the blocks.  $\Box$ 

In the cases of greatest interest to us the matrices in the statement of Theorem 3.16 have coefficients in  $\mathbb{Q}_p$ . We can take  $F = \mathbb{C}_p$ , but the resulting matrix S will have the entries in  $\mathbb{C}_p$  and not necessarily in  $\mathbb{Q}_p$ , and we want a symplectomorphism of  $(\mathbb{Q}_p)^n$ , which is given by a symplectic matrix with entries in  $\mathbb{Q}_p$ . To avoid this, we need to manipulate adequately the symplectic basis, which translates to adjusting the matrix Din Proposition 3.2. This problem also happens in the real case, but, as we will see, the matrix S can always be adapted to have the entries in  $\mathbb{R}$  instead of  $\mathbb{C}$ .

### 4. *p*-ADIC MATRIX CLASSIFICATION IN DIMENSION 2

In this section we prove Theorems D and E, that is, the *p*-adic version of the Weierstrass-Williamson matrix classification in dimension 2. The strategy consists of lifting the problem, when needed, to an extension field of order 2, and then using the theorems in Section 3 to go back to the *p*-adic field.

4.1. General preparatory results. In dimension 2, the eigenvalues of  $\Omega_0^{-1}M$  are the roots of a degree 2 polynomial with coefficients in F, so they have the form  $\lambda$  and  $-\lambda$ . The case where the eigenvalues are in the base field is the easiest one.

**Proposition 4.1.** Let F be a field of characteristic different from 2. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^2$ . Let  $M \in \mathcal{M}_2(F)$  symmetric and invertible such that the eigenvalues of  $\Omega_0^{-1}M$  are in F. Then there is a symplectic matrix  $S \in \mathcal{M}_2(F)$  and  $a \in F$  such that

$$S^T M S = \begin{pmatrix} 0 & a \\ a & 0 \end{pmatrix}.$$

*Proof.* If the eigenvalues  $\pm \lambda$  of  $\Omega_0^{-1}M$  are in F, its eigenvectors are also in F, and the matrix  $\Psi_1$  of Corollary 3.5 is in F. In order to have the same eigenvalues for

$$\Omega_0^{-1} \begin{pmatrix} 0 & a \\ a & 0 \end{pmatrix}$$

we need  $a = \lambda$ . The matrix S that we need is precisely  $\Psi_1$ .

If the matrix is not invertible, the case of the null matrix is already covered by the previous result, with a = 0. The other case is solved in the next proposition.

**Proposition 4.2.** Let F be a field of characteristic different from 2. Let

$$M = \begin{pmatrix} a & b \\ b & c \end{pmatrix} \in \mathcal{M}_2(F)$$

where  $b^2 = ac$ . Let  $d \in F$ . There exists a symplectic matrix S such that

$$S^T M S = \begin{pmatrix} d & 0 \\ 0 & 0 \end{pmatrix}$$

if and only if ad is a square.

*Proof.* The second column of S must be (kb, -ka) for some  $k \in F$ . Let the first column be (x, y), for  $x, y \in F$ . Then

$$d = \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = ax^2 + cy^2 + 2bxy$$

and

$$ad = a^{2}x^{2} + acy^{2} + 2abxy = a^{2}x^{2} + b^{2}y^{2} + 2abxy$$

is a square. Conversely, if ad is a square, we can take

$$S = \begin{pmatrix} -kd & kb \\ 0 & -ka \end{pmatrix}$$

where k is chosen so that  $k^2ad = 1$ .

For the cases where  $\lambda \notin F$ , we need some definitions.

**Definition 4.3.** Given an Abelian group G, we call Sq(G) the subgroup formed by the squares in G.

**Definition 4.4.** Given a field F with additive identity 0 and  $c \in F$ , we call

$$DSq(F,c) = \left\{ x^2 + cy^2 : x, y \in F \right\} \setminus \{0\}$$

and

$$\overline{\mathrm{DSq}}(F,c) = \mathrm{DSq}(F,c)/\operatorname{Sq}(F^*).$$

**Lemma 4.5.** Let F be a field and  $c \in F$ . Then DSq(F,c) is a group with respect to multiplication in F.

*Proof.* We just need to see that the product of two elements of DSq(F, c) is in DSq(F, c) and the inverse of an element of DSq(F, c) is in DSq(F, c):

$$(x_1^2 + cy_1^2)(x_2^2 + cy_2^2) = (x_1x_2 + c^2y_1y_2)^2 + c(x_1y_2 - x_2y_1)^2$$

and

$$\frac{1}{x^2 + cy^2} = \left(\frac{x}{x^2 + cy^2}\right)^2 + c\left(\frac{y}{x^2 + cy^2}\right)^2.$$

The group DSq(F, c) can also be defined in terms of the Hilbert symbol:

$$(a,b)_p = \begin{cases} 1 & \text{if } ax^2 + by^2 = 1 \text{ for some } x, y \in \mathbb{Q}_p; \\ -1 & \text{otherwise.} \end{cases}$$

With this definition, we have that

$$\mathrm{DSq}(F,c) = \Big\{ a \in \mathbb{Q}_p : (a,-c)_p = 1 \Big\}.$$

Lemma 4.5 is a consequence of the multiplicativity of the Hilbert symbol.

We have that

$$F^*/\mathrm{DSq}(F,c) \cong (F^*/\mathrm{Sq}(F^*))/(\mathrm{DSq}(F,c)/\mathrm{Sq}(F^*))$$
$$= (F^*/\mathrm{Sq}(F^*))/\overline{\mathrm{DSq}}(F,c),$$

that is, we have a group isomorphism in the rightmost part of the commutative diagram

By definition,  $\overline{\text{DSq}}(F, c)$  is the subset of the classes in  $F^*/\text{Sq}(F^*)$  which contain elements of the form  $x^2 + c$ , for  $x \in F$ . Note also that DSq(F, c), and hence  $\overline{\text{DSq}}(F, c)$ , only depend on the class of c modulo  $\text{Sq}(F^*)$ .

Now we can give a necessary and sufficient condition for a matrix to be symplectomorphic to a normal form.

**Proposition 4.6.** Let F be a field of characteristic different from 2. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^2$ . Let  $M \in \mathcal{M}_2(F)$  be a symmetric matrix such that the eigenvalues of  $\Omega_0^{-1}M$  are of the form  $\pm \lambda$  with  $\lambda \notin F$  but  $\lambda^2 \in F$ . Let u be the eigenvector of value  $\lambda$  in  $\Omega_0^{-1}M$ . Then for any  $a, b \in F$  there is a symplectic matrix Ssuch that

$$S^T M S = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$$

if and only if  $\lambda^2 = -ab$  and

$$\frac{2\lambda a}{u^T \Omega_0 \bar{u}} \in \mathrm{DSq}(F, -\lambda^2).$$

*Proof.* The eigenvalues of

$$\Omega_0^{-1} \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} = \begin{pmatrix} 0 & -b \\ a & 0 \end{pmatrix}$$

are  $\pm \sqrt{-ab}$ , so we must have  $\lambda^2 = -ab$ .

The matrix  $\Psi_2$  of Corollary 3.5 has the form

$$\begin{pmatrix} \lambda & -\lambda \\ a & a \end{pmatrix}.$$

The formula for S gives that

$$S = \Psi_1 \begin{pmatrix} d_1 & 0\\ 0 & d_2 \end{pmatrix} \begin{pmatrix} \frac{1}{2\lambda} & \frac{1}{2a}\\ -\frac{1}{2\lambda} & \frac{1}{2a} \end{pmatrix}.$$

The two columns of  $\Psi_1$  are the eigenvectors of  $\lambda$  and  $-\lambda$ . The first is u, and the second is the conjugate  $\bar{u}$  (or, more precisely, can be taken as the conjugate).

If S has entries in F, let  $c_1$  and  $c_2$  be its columns. We get

$$c_1 = \frac{d_1 u - d_2 \bar{u}}{2\lambda} \in F^2;$$

$$c_2 = \frac{d_1 u + d_2 \bar{u}}{2a} \in F^2;$$

$$d_1 u = ac_2 + \lambda c_1, d_2 \bar{u} = ac_2 - \lambda c_1 = \overline{d_1 u} \Rightarrow d_2 = \overline{d_1}.$$

The numbers  $d_1$  and  $d_2$  must also satisfy (3.1), that is

(4.1) 
$$d_1 d_2 = \frac{(\lambda, a)\Omega_0(-\lambda, a)^T}{u^T \Omega_0 \bar{u}} = \frac{2\lambda a}{u^T \Omega_0 \bar{u}}.$$

Taking into account that  $d_2 = \bar{d}_1$ , we can substitute  $d_1 = r + s\lambda$  and  $d_2 = r - s\lambda$  in 4.1, giving

$$r^2 - s^2 \lambda^2 = \frac{2\lambda a}{u^T \Omega_0 \bar{u}}$$

so this is in  $DSq(F, -\lambda^2)$ .

We can apply Proposition 4.6 to the real elliptic case, that is, the case of a matrix in  $\mathcal{M}_2(\mathbb{R})$  whose eigenvalues are  $\pm \lambda = \pm ri$ . There, we want to achieve a = b.  $\lambda$  is purely imaginary, so  $\lambda^2 = -ab = -a^2$  determines exactly |a| = r. DSq $(\mathbb{R}, -\lambda^2)$  consists of all positive reals, so there is always a solution for a (we know |a| and can take the adequate sign to make  $2\lambda a/u^T \Omega_0 \bar{u}$  positive). This is why the real Weierstrass-Williamson classification in dimension 2 has just two cases.

In the *p*-adic case, the two results until now are still enough to achieve a complete Weierstrass-Williamson classification in dimension 2, but it is more complicated than the real case. First we need to compute  $DSq(\mathbb{Q}_p, c)$  for all values of *c*. Of course, "all values" means "all classes modulo  $Sq(\mathbb{Q}_p^*)$ ": by Corollary A.4, this quotient is

$$\Big\{1,c_0,p,c_0p\Big\},$$

where  $c_0$  is a quadratic non-residue modulo p, except if p = 2, in which case the quotient is

$$\{1, -1, 2, -2, 3, -3, 6, -6\}.$$

We use the notation  $\operatorname{digit}_i(x)$  for the digit in the *p*-adic expansion of *x* which is *i* positions to the left of the leading digit, that is, the digit of order  $\operatorname{ord}(x) + i$ . The value of  $\operatorname{DSq}(\mathbb{Q}_p, c)$  can also be deduced from known facts about the Hilbert symbol; however, this does not seem simpler than a direct proof.

**Proposition 4.7.** Let p be a prime number such that  $p \neq 2$  and  $c \in \mathbb{Q}_p^*$ . Then  $DSq(\mathbb{Q}_p, c)$  is given as follows (see Figure 5):

- (1) If  $p \equiv 3 \mod 4$ , then  $DSq(\mathbb{Q}_p, 1) = \{u \in \mathbb{Q}_p : \operatorname{ord}_p(u) \equiv 0 \mod 2\}$  and  $DSq(\mathbb{Q}_p, 1) = \mathbb{Q}_p$  otherwise;
- (2) if  $p \equiv 1 \mod 4$ , then  $\mathrm{DSq}(\mathbb{Q}_p, c_0) = \{u \in \mathbb{Q}_p : \mathrm{ord}_p(u) \equiv 0 \mod 2\}$  and  $\mathrm{DSq}(\mathbb{Q}_p, c_0) = \mathbb{Q}_p$  otherwise;
- (3) for any value of the prime p,  $DSq(\mathbb{Q}_p, p) = \{u \in \mathbb{Q}_p : digit_0(u) \in Sq(\mathbb{F}_p^*)\};$
- (4) for any value of the prime p,  $DSq(\mathbb{Q}_p, c_0 p) = \{u \in \mathbb{Q}_p : \operatorname{ord}_p(u) \equiv 0 \mod 2, \operatorname{digit}_0(u) \in Sq(\mathbb{F}_p^*)\} \cup \{u \in \mathbb{Q}_p : \operatorname{ord}_p(u) \neq 0 \mod 2, \operatorname{digit}_0(u) \notin Sq(\mathbb{F}_p^*)\}.$

*Proof.* For the first point, we need to look at the possibilities modulo  $\operatorname{Sq}(\mathbb{Q}_p^*)$  of numbers of the form  $x^2 + 1$ . We can immediately get 1 and  $c_0$ , the first by taking x with high order and the second by taking it with order 0 and an adequate leading digit. We can only get p and  $c_0 p$  if  $x^2$  can have -1 as a leading digit, which only happens if  $p \equiv 1 \mod 4$ .

For the second point, we need, analogously, to look at numbers of the form  $x^2 + c_0$ . We immediately get the classes 1 and  $c_0$ . We can only get p and  $c_0p$  if  $-c_0$  is a square modulo p, that is, if  $p \equiv 3 \mod 4$ .

For the last two points, numbers of the form  $x^2 + p$  are in the classes 1 and p, depending only in the order of x, and those of the form  $x^2 + c_0 p$  are in 1 and  $c_0 p$ .

**Proposition 4.8.**  $DSq(\mathbb{Q}_2, c)$  is given as follows (see Figure 6):

- (1)  $DSq(\mathbb{Q}_2, 1) = \{ u \in \mathbb{Q}_2 : digit_1(u) = 0 \};$
- (2)  $\mathrm{DSq}(\mathbb{Q}_2, -1) = \mathbb{Q}_2;$




FIGURE 5. Top:  $DSq(\mathbb{Q}_p, c)$  for  $c \in \mathbb{Q}_p$  and  $p \neq 2$ . In each group of four circles, the upper circles represent even order numbers and the lower circles odd order, and the right circles represent square leading digits and the left circles non-square digits. Bottom: these four classes depicted for p = 3. Each circle "contains" the points with the same color and the black point at the right is 0.

 $\begin{array}{ll} (3) \ \mathrm{DSq}(\mathbb{Q}_{2},2) = \{ u \in \mathbb{Q}_{2} : \mathrm{digit}_{2}(u) = 0 \}; \\ (4) \ \mathrm{DSq}(\mathbb{Q}_{2},-2) = \{ u \in \mathbb{Q}_{2} : \mathrm{digit}_{1}(u) = \mathrm{digit}_{2}(u) \}; \\ (5) \ \mathrm{DSq}(\mathbb{Q}_{2},3) = \{ u \in \mathbb{Q}_{2} : \mathrm{ord}_{2}(u) \equiv 0 \mod 2 \}; \\ (6) \ \mathrm{DSq}(\mathbb{Q}_{2},-3) = \{ u \in \mathbb{Q}_{2} : \mathrm{ord}_{2}(u) + \mathrm{digit}_{1}(u) \equiv 0 \mod 2 \}; \\ (7) \ \mathrm{DSq}(\mathbb{Q}_{2},6) = \{ u \in \mathbb{Q}_{2} : \mathrm{ord}_{2}(u) + \mathrm{digit}_{1}(u) + \mathrm{digit}_{2}(u) \equiv 0 \mod 2 \}; \\ (8) \ \mathrm{DSq}(\mathbb{Q}_{2},-6) = \{ u \in \mathbb{Q}_{2} : \mathrm{ord}_{2}(u) + \mathrm{digit}_{2}(u) \equiv 0 \mod 2 \}. \end{array}$ 

*Proof.* Table 5 indicates the leading digits and the parity of the order of  $a^2 + cb^2$  depending on c and the difference  $\operatorname{ord}_2(b) - \operatorname{ord}_2(a)$ . The result follows by putting together the cases in the same row of the table. Note that a case such as "011 even" covers *all* 2-adic numbers with even order and ending in 011.

4.2. **Proof of Theorems D and E.** First, note that Theorem E follows directly from Theorem D: the isolated normal forms correspond to the different values of r for c = 0, and each family of normal forms corresponds to a value of  $c \in X_p$  with  $c \neq 0$ . Now we prove Theorem D.

(a) First we prove existence. The characteristic polynomial of  $\Omega_0^{-1}M$  has two opposite roots, which may or may not be in  $\mathbb{Q}_p$ . If they are in  $\mathbb{Q}_p$  and are not 0, Proposition

		$\operatorname{ord}_2(b) - \operatorname{ord}_2(a)$								
c	$cb^2$	$\leq -2$	-1	0	1	$\geqslant 2$				
1	001  even	001  even	101 even	01  odd	101  even	001 even				
-1	111 even	111 even	011  even	anything	101  even	001 even				
2	001  odd	001  odd	011  odd	011  even	001  even	001 even				
-2	111 odd	111 odd	001  odd	111 even	001  even	001 even				
3	011  even	011  even	111 even	1 even	101  even	001 even				
-3	101  even	101 even	001  even	11  odd	101  even	001 even				
6	011 odd	011 odd	101 odd	111 even	001  even	001 even				
-6	101  odd	101 odd	111 odd	011  even	001  even	001 even				

TABLE 5. Leading digits and parity of the order of  $a^2 + cb^2$  depending on c and the difference  $\operatorname{ord}_2(b) - \operatorname{ord}_2(a)$ . The number  $a^2$  is always described as 001 even,  $cb^2$  depends exclusively on c, and the result of the addition of both terms depends in the offset between these digits. Note that the leading 1's will add up to 0 if the offset is 0, hence making the second digit the leading one in the cases "01 odd" and "11 odd" (in these cases the order increases in 1), the third in the case "1 even" (the order increases in 2), and giving any possible result when adding 001 and 111 at the same position.

4.1 implies that M can be converted to

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

except by a constant factor. But the matrix

$$N = \begin{pmatrix} r & 0\\ 0 & cr \end{pmatrix}$$

in Theorem D, where c = 1 if  $p \equiv 1 \mod 4$  and c = -1 otherwise, has also two eigenvalues in  $\mathbb{Q}_p$ , so it can be converted to the same matrix.

If the eigenvalues are 0, either the matrix is zero, and we are in the same situation but with r = 0, or they are not zero, and Proposition 4.2 gives the same matrix but with c = 0. In this case r must be such that ar is a square, where a is one of the coefficients. There is one and only one  $r \in Y_p \cup \{1\}$  such that this happens.

Now suppose that the roots of the characteristic polynomial of  $\Omega_0^{-1}M$  are  $\pm \lambda$  for  $\lambda \notin \mathbb{Q}_p$ . In this case, we must have  $\lambda^2 \in \mathbb{Q}_p$ . We have  $N = S^T M S$  for a symplectic matrix S and some r and c if the two conditions of Proposition 4.6 hold for some a and b = ac. The first condition reads

$$\lambda^2 = -a^2 c \Rightarrow a^2 = -\frac{\lambda^2}{c}.$$

We must now split the proof into several cases.

•  $\lambda^2$  has even order. In order for  $-\lambda^2/c$  to be a square, we need c of even order. We also know that  $\lambda^2$  is not a square, so -c must not be a square. The elements of  $X_p$  which satisfy these conditions are  $\{c_0, c_0p^2\}$  if  $p \equiv 1 \mod 4$ ,  $\{1, p^2\}$  if  $p \equiv 3 \mod 4$ , and  $\{1, 3, -3, 12\}$  if p = 2.

For p = 2, we also need that  $-\lambda^2/c$  ends in 001, so the three last digits in c must agree with those in  $-\lambda^2$ , which will be 001, 101 or 011 (not 111, which




FIGURE 6. Top:  $DSq(\mathbb{Q}_2, c)$  for  $c \in \mathbb{Q}_2$ . In each group of eight circles, the upper circles represent even order numbers and the lower circles odd order, the two rightmost circles in the row represent a 0 as second digit and the two leftmost circles a 1, and in each pair of circles, the rightmost one has 0 as third digit and the leftmost one has 1. Bottom: a depiction of the eight classes. Each circle "contains" the points with the same color and the black point in the lower left is 0.

would make  $\lambda^2$  a square). This narrows further down the options to  $\{1\}$ ,  $\{-3\}$  or  $\{3, 12\}$ , respectively.

Let C be the current set of options for c, which contains only the singleton  $\{c_1\}$  or the set of two values  $\{c_1, c_1p^2\}$ . All of them satisfy that  $-\lambda^2/c$  is a square. We still need to apply the second condition, that is, we need to choose a such that

$$\frac{2\lambda a}{u^T \Omega_0 \bar{u}} \in \mathrm{DSq}(\mathbb{Q}_p, -\lambda^2) = \mathrm{DSq}(\mathbb{Q}_p, c_1).$$

Let

$$a_1 = \sqrt{\frac{-\lambda^2}{c_1}}.$$

In the two cases where  $C = \{c_1\}$ , by Proposition 4.8(1) and (6), for any  $x \in \mathbb{Q}_p$ , either x or -x is in  $\mathrm{DSq}(\mathbb{Q}_p, c_1)$ . So, either  $a = a_1$  or  $a = -a_1$  satisfies the condition, and  $c = c_1$  is valid. If

$$C = \{c_1, c_1 p^2\},\$$

by Proposition 4.7(1) and (2) and Proposition 4.8(5),

$$\mathrm{DSq}(\mathbb{Q}_p, c_1) = \Big\{ u \in \mathbb{Q}_p : \mathrm{ord}_p(u) \equiv 0 \mod 2 \Big\}.$$

This implies that either  $a = a_1$  or  $a = a_1/p$  satisfies the condition. Hence, only one between  $c_1$  and  $c_1p^2$  is a valid value of c.

- $\lambda^2$  has odd order. Now we need c to have odd order instead of even. What happens next depends on p.
  - If  $p \equiv 1 \mod 4$ , the values of c with odd order are  $c_0^k p$  for k = 0, 1, 2, 3. The first condition implies that  $-\lambda^2/c_0^k p$  is a square, which is true for k = 0 and 2 or for k = 1 or 3, depending on the leading digit of  $-\lambda^2/p$ . Let  $c_1$  be the value which satisfies this between p and  $c_0 p$ . Now the two candidates for c are  $c_1$  and  $c_1 c_0^2$ .

We define again  $a_1 = \sqrt{-\lambda^2/c_1}$ . In this case, by Proposition 4.7(3) and (4), for any  $x \in \mathbb{Q}_p$ , either x or  $c_0 x$  is in  $DSq(\mathbb{Q}_p, c_1)$ . Hence, either  $a = a_1$  or  $a = a_1/c_0$  satisfies the condition, and either  $c = c_1$  or  $c = c_1c_0^2$  is valid.

- If  $p \equiv 3 \mod 4$ , the values of c with odd order are p and -p. As -1 is not a square, only one will make  $-\lambda^2/c$  a square. For this value of c, we set  $a_1 = \sqrt{-\lambda^2/c}$ . By Proposition 4.7(3) and (4), for any  $x \in \mathbb{Q}_p$ , either x or -x is in  $DSq(\mathbb{Q}_p, c_1)$ . Hence, either  $a = a_1$  or  $a = -a_1$  is valid, and c is valid in any case.
- If p = 2, the values of c with odd order are 2, -2, 6, -6, -18 and 24.  $-\lambda^2/c$  must end in 001, so c must agree with  $-\lambda^2$  in the last three digits, that in this case can have all possible values: 001, 101, 011 and 111. The valid c's in each case are, respectively,

$$\{2\}, \{-6\}, \{6, 24\}, \{-2, -18\}.$$

Let C be this set,  $c_1$  the element of less absolute value (in the real sense) in C, and  $a_1 = \sqrt{-\lambda^2/c_1}$ .

If C has only one element, by Proposition 4.8(3) and (8), for any  $x \in \mathbb{Q}_p$ , either x or -x is in  $DSq(\mathbb{Q}_p, c_1)$ , and again  $c_1$  is valid in any case.

If  $C = \{-2, -18\}$ , by Proposition 4.8(4), for any  $x \in \mathbb{Q}_p$ , either x or 3x is in  $DSq(\mathbb{Q}_p, -2)$  (because digit<sub>1</sub>(x) = 1 - digit<sub>1</sub>(3x) and digit<sub>2</sub>(x) = digit<sub>2</sub>(3x)). Hence, either  $a = a_1$  or  $a = a_1/3$  works and c = -2 or c = -18, respectively, works.

Finally, if  $C = \{6, 24\}$ , by Proposition 4.8(7), for any  $x \in \mathbb{Q}_p$ , either x or 2x is in DSq( $\mathbb{Q}_p, 6$ ). So either  $a = a_1$  or  $a = a_1/2$  works and c = 6 or c = 24, respectively, works.

(b) Now we prove uniqueness. In the case where the roots of the characteristic polynomial are in  $\mathbb{Q}_p$ , the rest of c's in the lists do not lead to a matrix with the eigenvalues in  $\mathbb{Q}_p$ , because their opposites are not squares. In all the other cases, we have seen that there is one and only one valid value of c. If two matrices S and S' bring M to normal form, c must coincide, hence r also coincides because the eigenvalues of the normal forms must be the same.

Recall that, in the real case, there are two normal forms: elliptic and hyperbolic. To give them the form

$$\begin{pmatrix} r & 0 \\ 0 & cr \end{pmatrix},$$

we need c = 1 and c = -1, respectively. In the *p*-adic case, these two matrices are equivalent by multiplication by a symplectic matrix if and only if  $p \equiv 1 \mod 4$ : the list for this case is the only one that does not contain -1.

**Proposition 4.9.** All choices of quadratic residue in Definition 1.1 (including the least of all, which is used in the definition) lead to the same normal form in Theorem D, up to multiplication by a symplectic matrix. That is, if  $c_0$ ,  $c'_0$  are two quadratic residues modulo p then the normal forms corresponding to  $c_0$  and the normal forms corresponding to  $c'_0$  are equivalent by multiplication by a symplectic matrix. (The order of the forms, however, may vary: for example, taking  $c'_0 \equiv c_0^3 \mod p$  results in the new form with  $c = c'_0 p$  being taken as the one which had previously  $c = c_0^3 p$ .)

*Proof.* By Theorem D, the normal forms of matrices in the first set are equivalent to one and only one normal form in the second set, so the two sets are different representatives of the same classes.  $\Box$ 

# 5. General preparatory results for fields of characteristic different from 2

In this section we prove some results which are not specific to the reals or the *p*-adics, but which are needed to prove the *p*-adic matrix classification in dimension 4 and the real classification in any dimension. They have to do with finding normal forms for 4-by-4 matrices in terms of their eigenvalues.

**Proposition 5.1.** Let F be a field of characteristic different from 2. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^4$ . Let  $M \in \mathcal{M}_4(F)$  be a symmetric matrix such that the eigenvalues of  $\Omega_0^{-1}M$  are of the form  $\pm \lambda, \pm \mu$  where  $\lambda, \mu \notin F$  but all of them are in a degree 2 extension  $F[\alpha]$  for some  $\alpha$ . Then there are  $r, s \in F$  and a symplectic matrix  $S \in \mathcal{M}_4(F)$  such that  $S^TMS$  has the form of Theorem F(2) for  $c = \alpha^2$ .

*Proof.* The condition in the statement is equivalent to say that  $\mu = \overline{\lambda}$  and both are in  $F[\alpha]$ . Let  $\lambda = s + r\alpha$  and  $\mu = s - r\alpha$ , for  $r, s \in F$ .

Let

$$N = \begin{pmatrix} 0 & s & 0 & r \\ s & 0 & r\alpha^2 & 0 \\ 0 & r\alpha^2 & 0 & s \\ r & 0 & s & 0 \end{pmatrix},$$

which is the matrix in Theorem F(2) for  $c = \alpha^2$ . The eigenvalues of

$$\Omega_0^{-1}N = \begin{pmatrix} -s & 0 & -r\alpha^2 & 0\\ 0 & s & 0 & r\\ -r & 0 & -s & 0\\ 0 & r\alpha^2 & 0 & s \end{pmatrix}$$

are precisely  $\pm s \pm r\alpha$ , that is,  $\pm \lambda$  and  $\pm \mu$ , so we can apply Corollary 3.5. The matrix  $\Psi_2$  has the form

$$\begin{pmatrix} 0 & \alpha & 0 & -\alpha \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ \alpha & 0 & -\alpha & 0 \\ & 35 \end{pmatrix}$$

with the values in the order  $(\lambda, -\lambda, \overline{\lambda}, -\overline{\lambda})$  as needed by Corollary 3.5, and the resulting matrix S is

$$S = \Psi_1 \begin{pmatrix} d_1 & 0 & 0 & 0\\ 0 & d_2 & 0 & 0\\ 0 & 0 & d_3 & 0\\ 0 & 0 & 0 & d_4 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 & \frac{1}{\alpha}\\ \frac{1}{\alpha} & 0 & 1 & 0\\ 0 & 1 & 0 & -\frac{1}{\alpha}\\ -\frac{1}{\alpha} & 0 & 1 & 0 \end{pmatrix}$$

The condition (3.1) of Corollary 3.5 implies that

(5.1) 
$$d_1 d_2 = \frac{(0, 1, 0, \alpha) \Omega_0(\alpha, 0, 1, 0)^T}{u^T \Omega_0 v} = \frac{-2\alpha}{u^T \Omega_0 v}$$

and

(5.2) 
$$d_3 d_4 = \frac{(0, 1, 0, -\alpha)\Omega_0(-\alpha, 0, 1, 0)^T}{\bar{u}^T \Omega_0 \bar{v}} = \frac{2\alpha}{\bar{u}^T \Omega_0 \bar{v}},$$

where u and v are the eigenvectors for  $\lambda$  and  $-\lambda$  respectively.

We also want S to have entries in F. Let  $c_1, c_2, c_3, c_4$  be its columns, which should be vectors in  $F^4$ :

$$\begin{cases} c_1 = \frac{d_2 v - d_4 \bar{v}}{\alpha}; \\ c_2 = d_1 u + d_3 \bar{u}; \\ c_3 = d_2 v + d_4 \bar{v}; \\ c_4 = \frac{d_1 u - d_3 \bar{u}}{\alpha}. \end{cases}$$

These expressions imply that

$$2d_1u = c_2 + \alpha c_4, 2d_3\bar{u} = c_2 - \alpha c_4 = \overline{2d_1u} \Rightarrow d_3 = \bar{d}_1$$

and

$$2d_2v = c_3 + \alpha c_1, 2d_4\bar{v} = c_3 - \alpha c_1 = 2d_2v \Rightarrow d_4 = d_2.$$

Now we can take  $d_1$  and  $d_2$  arbitrary such that (5.1) holds, and (5.2) will hold automatically because  $d_3d_4 = \overline{d_1d_2}$ .

In the real case, this is enough to complete the Weierstrass-Williamson classification in all dimensions if the eigenvalues of  $\Omega_0^{-1}M$  are pairwise distinct. Indeed, these eigenvalues can be associated in pairs of the form  $\{a, -a\}$  or  $\{ia, -ia\}$  and quadruples of the form

$$\Big\{a+\mathrm{i}b,a-\mathrm{i}b,-a+\mathrm{i}b,-a-\mathrm{i}b\Big\}.$$

We can apply respectively Propositions 4.1, 4.6 (we already explained why this is always possible) and 5.1, giving the hyperbolic, elliptic and focus-focus normal forms in Section 2.

In the *p*-adic case, such a classification is still not complete, even for 4-by-4 matrices. The reason for this difference is that, if  $\alpha \notin \mathbb{R}$  with  $\alpha^2 \in \mathbb{R}$ , this means that  $\alpha$  is an imaginary number and  $\mathbb{R}[\alpha] = \mathbb{C}$ , which is algebraically closed. But if  $\alpha \notin \mathbb{Q}_p$ with  $\alpha^2 \in \mathbb{Q}_p$ ,  $\mathbb{Q}_p[\alpha]$  is not algebraically closed. So it is possible that  $\lambda^2 \notin \mathbb{Q}_p$  and simultaneously  $\lambda \notin \mathbb{Q}_p[\lambda^2]$ .

To fix this issue, consider a degree four polynomial of the form  $t^4 + At^2 + B$  (at the moment in an arbitrary field F). Its roots are of the form  $\lambda, -\lambda, \mu, -\mu$ . If  $\lambda^2$  and  $\mu^2$  are not in F, they are conjugate in some degree 2 extension, that is,  $\lambda^2 = a + b\alpha$  and  $\mu^2 = a - b\alpha$  for some  $\alpha \in F[\lambda^2]$ . In turn, if  $\lambda$  and  $\mu$  are not in  $F[\lambda^2]$ , we have a hierarchy of fields:

$$F[\lambda, \mu] \\ | 2 \\ F[\alpha] = F[\lambda^2] = F[\mu^2] \\ | 2 \\ F$$

There is an automorphism of  $F[\lambda, \mu]$  that fixes F and moves  $\alpha$  to  $-\alpha$  (an extension of the conjugation in  $F[\alpha]$ ). We will denote this as  $x \mapsto \bar{x}$ .  $\bar{\lambda}$  must be  $\mu$  or  $-\mu$ , without loss of generality we suppose that it is  $\mu$ . There is another automorphism of  $F[\lambda, \mu]$  that fixes  $F[\alpha]$  and changes  $\lambda$  to  $-\lambda$  and  $\mu$  to  $-\mu$ , which we will denote as  $x \mapsto \hat{x}$ .

**Proposition 5.2.** Let F be a field with characteristic different from 2. Let  $F[\alpha]$  be a degree two extension of F and let  $F[\gamma, \overline{\gamma}]$  be an extension of  $F[\alpha]$  such that  $\gamma^2 \in F[\alpha]$ . Let  $\Omega_0$  be the matrix of the standard symplectic form on  $F^4$ . Let  $t_1, t_2 \in F$  such that

$$\begin{cases} \gamma^2 = t_1 + t_2 \alpha; \\ \bar{\gamma}^2 = t_1 - t_2 \alpha. \end{cases}$$

Let  $M \in \mathcal{M}_4(F)$  be a symmetric matrix such that the eigenvalues of  $\Omega_0^{-1}M$  are of the form  $\pm \lambda, \pm \mu$  with

and

$$\lambda = (r + s\alpha)\gamma$$
$$\mu = (r - s\alpha)\bar{\gamma},$$

for  $r, s \in F$ . Let  $a, b \in F$ . Let u be the eigenvector of  $\lambda$ . Then, there is a symplectic matrix  $S \in \mathcal{M}_4(F)$  such that  $S^T M S$  has the form of Theorem F(3) with  $c = \alpha^2$  if and only if

$$\frac{a\alpha\gamma(b+\alpha)}{u^T\Omega_0\hat{u}}\in \mathrm{DSq}(F[\alpha],-\gamma^2).$$

*Proof.* Let N be the matrix

$$\begin{pmatrix} \frac{a\alpha^{2}(r-bs)}{\alpha^{2}-b^{2}} & 0 & \frac{s\alpha^{2}-rb}{\alpha^{2}-b^{2}} & 0 \\ 0 & \frac{-r(t_{1}+bt_{2})-s(bt_{1}+\alpha^{2}t_{2})}{a} & 0 & -r(bt_{1}+\alpha^{2}t_{2})-s\alpha^{2}(t_{1}+bt_{2}) \\ \frac{s\alpha^{2}-rb}{\alpha^{2}-b^{2}} & 0 & \frac{r-bs}{a(\alpha^{2}-b^{2})} & 0 \\ 0 & -r(bt_{1}+\alpha^{2}t_{2})-s\alpha^{2}(t_{1}+bt_{2}) & 0 & a\alpha^{2}(-r(t_{1}+bt_{2})-s(bt_{1}+\alpha^{2}t_{2})) \end{pmatrix}$$

We have that  $\Omega_0^{-1}N$  is equal to

$$\begin{pmatrix} 0 & \frac{r(t_1+bt_2)+s(bt_1+\alpha^2 t_2)}{a} & 0 & r(bt_1+\alpha^2 t_2)+s\alpha^2(t_1+bt_2) \\ \frac{a\alpha^2(r-bs)}{\alpha^2-b^2} & 0 & \frac{s\alpha^2-br}{\alpha^2-b^2} & 0 \\ 0 & r(bt_1+\alpha^2 t_2)+s\alpha^2(t_1+bt_2) & 0 & a\alpha^2(r(t_1+bt_2)+s(bt_1+\alpha^2 t_2)) \\ \frac{s\alpha^2-br}{\alpha^2-b^2} & 0 & \frac{r-bs}{a(\alpha^2-b^2)} & 0 \end{pmatrix}$$

which has as set of eigenvalues

$$\left\{\pm (r+s\alpha)\gamma, \pm (r-s\alpha)\bar{\gamma}\right\} = \left\{\lambda, -\lambda, \mu, -\mu\right\},\$$

and the condition of Corollary 3.5 is satisfied. The matrix

$$\Psi_{2} = \begin{pmatrix} (b+\alpha)\gamma & -(b+\alpha)\gamma & (b-\alpha)\bar{\gamma} & -(b-\alpha)\bar{\gamma} \\ a\alpha & a\alpha & -a\alpha & -a\alpha \\ a\alpha\gamma(b+\alpha) & -a\alpha\gamma(b+\alpha) & -a\alpha\bar{\gamma}(b-\alpha) & a\alpha\bar{\gamma}(b-\alpha) \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

and the columns of  $\Psi_1$  are the eigenvectors of  $\lambda, -\lambda, \mu$  and  $-\mu$  in that order, which means that they are of the form  $u, \hat{u}, \bar{u}$  and  $\hat{\bar{u}}$ . Let  $c_1, c_2, c_3$  and  $c_4$  be the columns of S, which we want to be in F. We have that  $\Psi_2 S = \Psi_1 D$ , that is,

$$\Psi_2 \begin{pmatrix} c_1 & c_2 & c_3 & c_4 \end{pmatrix} = \begin{pmatrix} u & \hat{u} & \bar{u} & \hat{u} \end{pmatrix} \begin{pmatrix} d_1 & 0 & 0 & 0 \\ 0 & d_2 & 0 & 0 \\ 0 & 0 & d_3 & 0 \\ 0 & 0 & 0 & d_4 \end{pmatrix} = \begin{pmatrix} d_1 u & d_2 \hat{u} & d_3 \bar{u} & d_4 \hat{u} \end{pmatrix}$$

which expands to

$$d_1 u = (b+\alpha)\gamma c_1 + a\alpha c_2 + a\alpha\gamma(b+\alpha)c_3 + c_4;$$
  

$$d_2 \hat{u} = -(b+\alpha)\gamma c_1 + a\alpha c_2 - a\alpha\gamma(b+\alpha)c_3 + c_4;$$
  

$$d_3 \bar{u} = (b-\alpha)\bar{\gamma}c_1 - a\alpha c_2 - a\alpha\bar{\gamma}(b-\alpha)c_3 + c_4;$$
  

$$d_4 \hat{\bar{u}} = -(b-\alpha)\bar{\gamma}c_1 - a\alpha c_2 + a\alpha\bar{\gamma}(b-\alpha)c_3 + c_4;$$

that is

$$d_2\hat{u} = d_1u \Rightarrow d_2 = d_1;$$
  

$$d_3\bar{u} = \overline{d_1u} \Rightarrow d_3 = \bar{d}_1;$$
  

$$d_4\hat{\bar{u}} = \widehat{d_3\bar{u}} \Rightarrow d_4 = \hat{d}_3 = \hat{d}_1.$$

It only remains to apply the condition about D in Corollary 3.5. The result is

$$d_1 d_2 = d_1 \hat{d}_1 = \frac{4a\alpha\gamma(b+\alpha)}{u^T \Omega_0 \hat{u}};$$
  
$$d_3 d_4 = \bar{d}_1 \hat{\bar{d}}_1 = \frac{-4a\alpha\bar{\gamma}(b-\alpha)}{\bar{u}^T \Omega_0 \hat{\bar{u}}}$$

Obviously, both conditions are equivalent. Putting  $d_1 = 2(z_1 + z_2\gamma)$  with  $z_1, z_2 \in F[\alpha]$ , the first condition becomes

$$z_1^2 - z_2^2 \gamma^2 = \frac{a\alpha\gamma(b+\alpha)}{u^T \Omega_0 \hat{u}}$$

so this must be in  $DSq(F[\alpha], -\gamma^2)$ .

Proposition 5.2 reduces the problem of classifying this type of matrices to classifying (adequate) degree 4 extensions of F. In our case  $F = \mathbb{Q}_p$ .

The values of  $\alpha^2$  are given by

$$\mathbb{Q}_p^*/\operatorname{Sq}(\mathbb{Q}_p^*),$$

because two values whose quotient is a square are equivalent. The squares in  $\mathbb{Q}_p$  are the numbers with even order and with a leading digit in  $\operatorname{Sq}(\mathbb{F}_p)$ , so the quotient is  $\{1, c_0, p, c_0 p\}$ , except if p = 2, where squares have even order and end in 001, and the quotient is  $\{1, -1, 2, -2, 3, -3, 6, -6\}$ . The possible values of  $\alpha^2$  are all except 1 (because  $\alpha \notin \mathbb{Q}_p$ ). In each case, to find the normal form of M we still need to know  $\gamma$ , a and b.

In turn, the values of  $\gamma$  are given by the quotient

$$\mathbb{Q}_p[\alpha]^* / \operatorname{Sq}(\mathbb{Q}_p[\alpha]^*).$$

So we need to determine which numbers are squares in  $\mathbb{Q}_p[\alpha]$ .

**Lemma 5.3.** Let F be a field with characteristic different from 2. Let  $F[\alpha]$  be a degree 2 extension of F. Let  $a, b \in F$ . Then  $a + b\alpha$  is a square in  $F[\alpha]$  if and only if  $a^2 - b^2\alpha^2$  is a square in F and one of the numbers

$$\frac{a \pm \sqrt{a^2 - b^2 \alpha^2}}{2}$$

is also a square in F.

*Proof.* Suppose that  $a + b\alpha = (r + s\alpha)^2$ . Then also  $a - b\alpha = (r - s\alpha)^2$ , multiplying  $a^2 - b^2\alpha^2 = (r^2 - s^2\alpha^2)^2$ , and finally

$$\frac{a \pm \sqrt{a^2 - b^2 \alpha^2}}{2} = \frac{r^2 + s^2 \alpha^2 + r^2 - s^2 \alpha^2}{2} = r^2$$

Reciprocally, if both numbers are squares, they give r and s such that  $a + b\alpha = (r + s\alpha)^2$ : we get r from the previous formula, and then s from b = 2rs.

By Proposition 5.2, for fixed values of  $\alpha$  and  $\gamma$ , two normal forms for (a, b) and (a', b') are equivalent if and only if the quotient between  $a(b + \alpha)$  and  $a'(b' + \alpha)$  is an element of  $DSq(\mathbb{Q}_p[\alpha], -\gamma^2)$ . Hence, the normal forms for fixed  $\alpha$  and  $\gamma$  are given by the classes in

$$\mathbb{Q}_p[\alpha]^* / \mathrm{DSq}(\mathbb{Q}_p[\alpha], -\gamma^2),$$

or equivalently in

$$(\mathbb{Q}_p[\alpha]^*/\operatorname{Sq}(\mathbb{Q}_p[\alpha]^*))/\overline{\operatorname{DSq}}(\mathbb{Q}_p[\alpha], -\gamma^2).$$

That is, we have reduced the problem of determining subgroups of  $\mathbb{Q}_p[\alpha]$  to determining the subgroup

$$\overline{\mathrm{DSq}}(\mathbb{Q}_p[\alpha], -\gamma^2)$$

which is easier because this is a subgroup of a finite group (and in all cases of interest, isomorphic to  $\mathbb{F}_2^n$  for some n).

### 6. p-adic matrix classification in dimension 4

We will now provide the *p*-adic classification of 4-by-4 matrices. Some cases are a consequence of the previous results: the characteristic polynomial of  $\Omega_0^{-1}M$  has the form

$$At^4 + Bt^2 + C$$

and the roots are  $\lambda$ ,  $-\lambda$ ,  $\mu$  and  $-\mu$ . If  $\lambda^2$  is in  $\mathbb{Q}_p$ , then  $\mu^2$  is also in  $\mathbb{Q}_p$ . If  $\lambda \neq \mu$ , we can multiply M by a symplectic matrix to separate it into two components, one with the eigenvalues  $\lambda$  and  $-\lambda$ , and the other with  $\mu$  and  $-\mu$ , and apply to each component Theorem D. In the real case this results in three rank 0 normal forms: elliptic-elliptic, elliptic-hyperbolic and hyperbolic-hyperbolic. In the *p*-adic case, we have to combine analogously the normal forms for dimension 2, getting a total of  $\binom{8}{2} = 28$  forms for  $p \equiv 1 \mod 4$ ,  $\binom{6}{2} = 15$  for  $p \equiv 3 \mod 4$ , and  $\binom{12}{2} = 66$  for p = 2.

mod 4,  $\binom{6}{2} = 15$  for  $p \equiv 3 \mod 4$ , and  $\binom{12}{2} = 66$  for p = 2. The other case is when  $\lambda^2 \notin \mathbb{Q}_p$ . In this case,  $\lambda^2$  and  $\mu^2$  are conjugate roots in a degree two extension L of  $\mathbb{Q}_p$ . If  $\lambda^2$  is a square in L, that is,  $\lambda \in L$ , we have  $\mu^2 = \overline{\lambda^2}$  and  $\mu = \overline{\lambda}$  is also in L. We will now see that, in this case, the necessary condition of having the same eigenvalues is also sufficient to be linearly symplectomorphic. 6.1. Case  $p \neq 2$ . We start with the case  $p \equiv 1 \mod 4$ . This subdivides in three cases, depending on whether  $\alpha^2$  is  $c_0$ , p or  $c_0p$ . The following result gives a characterization of the squares in  $\mathbb{Q}_p[\alpha]$  in each case.

**Proposition 6.1.** Let p be a prime number such that  $p \equiv 1 \mod 4$ . Let  $c_0$  be a quadratic non-residue modulo p. Then the following statements hold.

- (1)  $\operatorname{Sq}(\mathbb{Q}_p[\sqrt{c_0}]^*) = \{a + b\sqrt{c_0} : a, b \in \mathbb{Q}_p, \operatorname{ord}_p(a) \leq \operatorname{ord}_p(b), \operatorname{ord}_p(a) \equiv 0 \mod 2, a^2 b^2 c_0 \in \operatorname{Sq}(\mathbb{Q}_p^*)\}.$
- (2) Sq( $\mathbb{Q}_p[\sqrt{p}]^*$ ) = { $a + b\sqrt{p} : a, b \in \mathbb{Q}_p, \operatorname{ord}_p(a) \leq \operatorname{ord}_p(b), \operatorname{digit}_0(a) \in \operatorname{Sq}(\mathbb{F}_p^*)$ }.
- (3)  $\operatorname{Sq}(\mathbb{Q}_p[\sqrt{c_0p}]^*) = \{a + b\sqrt{c_0p} : a, b \in \mathbb{Q}_p, \operatorname{ord}_p(a) \leq \operatorname{ord}_p(b), \operatorname{ord}_p(a) \equiv 0 \mod 2, \operatorname{digit}_0(a) \in \operatorname{Sq}(\mathbb{F}_p^*)\} \cup \{a + b\sqrt{c_0p} : \operatorname{ord}_p(a) \leq \operatorname{ord}_p(b), \operatorname{ord}_p(a) \not\equiv 0 \mod 2, \operatorname{digit}_0(a) \notin \operatorname{Sq}(\mathbb{F}_p^*)\}.$

*Proof.* (1) Suppose that  $a + b\sqrt{c_0}$  is a square in  $\mathbb{Q}_p$ . By Lemma 5.3,

$$a^2 - b^2 c_0 = (r^2 - s^2 c_0)^2$$

for some  $r, s \in \mathbb{Q}_p$ . In particular,  $a^2 - b^2 c_0$  is a square in  $\mathbb{Q}_p$ . If  $\operatorname{ord}(a)$  was higher than  $\operatorname{ord}(b)$ , that would make

$$\operatorname{digit}_0(a^2 - b^2 c_0) = \operatorname{digit}_0(-b^2 c_0) \notin \operatorname{Sq}(\mathbb{F}_p^*)$$

So  $\operatorname{ord}(a) \leq \operatorname{ord}(b)$ . We also have that  $a = r^2 + s^2 c_0$ , and by Proposition 4.7(2),  $\operatorname{ord}(a)$  is even.

Suppose now that a and b satisfy the three conditions. Then the first condition in Lemma 5.3 is satisfied. Let  $t_1$  and  $t_2$  be the two candidates for  $r^2$ . Note that  $t_1t_2 = b^2c_0/4$ .

The leading terms cannot cancel simultaneously in  $a + \sqrt{a^2 - b^2 c_0}$  and  $a - \sqrt{a^2 - b^2 c_0}$ . Without loss of generality, suppose that  $t_1$  has no cancellation. Then  $\operatorname{ord}(t_1) = \operatorname{ord}(a)$ , which is even, and  $\operatorname{ord}(t_2) = \operatorname{ord}(b^2 c_0/4t_1)$  is also even. But their product  $t_1 t_2 = b^2 c_0/4$  is a non-square, hence one of  $t_1$  and  $t_2$  is a square (because the product of two even-order non-squares is a square).

(2) Suppose that  $a + b\sqrt{p}$  is a square in  $\mathbb{Q}_p$ . By Lemma 5.3,

$$a^2 - b^2 p = (r^2 - s^2 p)^2$$

for some  $r, s \in \mathbb{Q}_p$ . This implies that  $\operatorname{ord}(a^2 - b^2 p)$  is even, so  $\operatorname{ord}(a) \leq \operatorname{ord}(b)$ . Here  $a = r^2 + s^2 p$ , so by Proposition 4.7(3),  $\operatorname{digit}_0(a) \in \operatorname{Sq}(\mathbb{F}_p^*)$ .

Reciprocally, if a and b satisfy the conditions,  $a^2 - b^2 p$  is a square because  $a^2$  is.  $t_1 = (a + \sqrt{a^2 - b^2 p})/2$  has the same order and leading digit than a, so if a is a square,  $t_1$  is also a square. Otherwise, a is p times a square and the same applies to  $t_1$ , and  $t_2 = b^2 p/4t_1$  is a square.

(3) Suppose that  $a + b\sqrt{c_0 p}$  is a square in  $\mathbb{Q}_p$ . By Lemma 5.3,

$$a^2 - b^2 c_0 p = (r^2 - s^2 c_0 p)^2$$

for some  $r, s \in \mathbb{Q}_p$ . This implies that  $\operatorname{ord}(a^2 - b^2 c_0 p)$  is even, so  $\operatorname{ord}(a) \leq \operatorname{ord}(b)$ . Now we have  $a = r^2 + s^2 c_0 p$ , which by Proposition 4.7(4) implies that either the order of a is even and its leading digit is square, or the order is odd and the leading digit is non-square.

Reciprocally, if a and b satisfy the conditions,  $a^2 - b^2 c_0 p$  is a square because  $a^2$  is. In the first case

$$t_1 = \frac{a + \sqrt{a^2 - b^2 c_0 p}}{40}$$

is a square. In the second case, a is p times an even order non-square,  $t_1$  is the same, and

 $t_2 = \frac{b^2 c_0 p}{4t_1}$ 

is a square.

**Corollary 6.2.** Let p be a prime number such that  $p \equiv 1 \mod 4$ . Let  $c_0$  be a quadratic non-residue modulo p. Then the following statements hold.

- (1)  $\mathbb{Q}_p[\sqrt{c_0}]^* / \operatorname{Sq}(\mathbb{Q}_p[\sqrt{c_0}]^*) = \{1, p, \sqrt{c_0}, p\sqrt{c_0}\}.$ (2)  $\mathbb{Q}_p[\sqrt{p}]^* / \operatorname{Sq}(\mathbb{Q}_p[\sqrt{p}]^*) = \{1, c_0, \sqrt{p}, c_0\sqrt{p}\}.$ (3)  $\mathbb{Q}_p[\sqrt{c_0p}]^* / \operatorname{Sq}(\mathbb{Q}_p[\sqrt{c_0p}]^*) = \{1, c_0, \sqrt{c_0p}, c_0\sqrt{c_0p}\}.$
- *Proof.* (1) Given an element  $a + b\sqrt{c_0}$  in  $\mathbb{Q}_p[\sqrt{c_0}]^*$ , if  $\operatorname{ord}(a) > \operatorname{ord}(b)$  or they are equal and  $a^2 b^2 c_0$  is not a square, we multiply it by  $\sqrt{c_0}$ . This guarantees that

$$\operatorname{ord}(a) \leq \operatorname{ord}(b)$$
 and  $a^2 - b^2 c_0$  is a square, because

$$\sqrt{c_0}(a+b\sqrt{c_0}) = a\sqrt{c_0} + bc_0$$

and

$$b^2 c_0^2 - a^2 c_0 = c_0 (a^2 - b^2 c_0)$$

so if  $a^2 - b^2 c_0$  was non-square, it is now square. Hence, if the order of a is odd, we multiply the element by p, and we obtain a square.

- (2) Given an element  $a + b\sqrt{p}$  in  $\mathbb{Q}_p[\sqrt{p}]^*$ , if  $\operatorname{ord}(a) > \operatorname{ord}(b)$ , we multiply it by  $\sqrt{p}$ , so that it has  $\operatorname{ord}(a) \leq \operatorname{ord}(b)$ . Now, multiplying it by  $c_0$  if needed, we ensure that  $\operatorname{digit}_0(a)$  is a square.
- (3) Given an element  $a + b\sqrt{c_0p}$  in  $\mathbb{Q}_p[\sqrt{c_0p}]^*$ , if  $\operatorname{ord}(a) > \operatorname{ord}(b)$ , we multiply it by  $\sqrt{c_0p}$ , so that it has  $\operatorname{ord}(a) \leq \operatorname{ord}(b)$ . Now, multiplying it by  $c_0$  if needed, we ensure that  $\operatorname{digit}_0(a)$  is a square or a non-square, depending on the order.  $\Box$

The element  $\gamma$  is the square root of an element in this set, but different from 1, which would lead to the case of Proposition 5.1. So there are three possible  $\gamma$ 's for each  $\alpha$ . Also, note that  $\gamma^2$  always is in  $\mathbb{Q}_p$  or  $\alpha$  times an element of  $\mathbb{Q}_p$ : this means that  $\bar{\gamma}^2$  is  $\gamma^2$  or  $-\gamma^2$ , that is,  $\bar{\gamma}$  is  $\gamma$  or i $\gamma$ . In any case,  $\bar{\gamma} \in \mathbb{Q}_p[\gamma]$  (here it is important that  $p \equiv 1 \mod 4$ so that  $i \in \mathbb{Q}_p$ ), or in other words, the extension  $\mathbb{Q}_p[\gamma, \bar{\gamma}]$  is the same as  $\mathbb{Q}_p[\gamma]$ , which is different for each  $\gamma$ .

The next step is to determine

$$\overline{\mathrm{DSq}}(\mathbb{Q}_p[\alpha], -\gamma^2),$$

or equivalently

$$\overline{\mathrm{DSq}}(\mathbb{Q}_p[\alpha], \gamma^2),$$

because -1 is a square. This consists of seeing which classes of  $\mathbb{Q}_p[\alpha]^*$  modulo a square are attainable by elements of the form  $x^2 + \gamma^2$  for different x. Once this is done, the quotient of  $\mathbb{Q}_p[\alpha]^*/\operatorname{Sq}(\mathbb{Q}_p[\alpha]^*)$  by this subgroup will give us the necessary a and b. The computations are shown in Table 6.

Now we make the analogous treatment with  $p \equiv 3 \mod 4$ . The values of  $\alpha$  are the same as in the previous case, but now we can take  $c_0 = -1$  to simplify the formulas, so we have  $\alpha = i, \sqrt{p}$  or  $i\sqrt{p}$ .

**Proposition 6.3.** Let p be a prime number such that  $p \equiv 3 \mod 4$ . Then the following statements hold.

(1) Sq( $\mathbb{Q}_p[\mathbf{i}]^*$ ) = { $a + \mathbf{i}b : a, b \in \mathbb{Q}_p, \min\{\operatorname{ord}_p(a), \operatorname{ord}_p(b)\} \equiv 0 \mod 2, a^2 + b^2 \in \operatorname{Sq}(\mathbb{Q}_p^*)$ }.

$\alpha^2$	all classes	$\gamma^2$	attainable classes	a	b	$a(b+\alpha)$	$[a(b+\alpha)]$
		,				· · · ·	
$c_0$	$1, p, \sqrt{c_0}, p\sqrt{c_0}$	p	1, p	1	0	$\sqrt{c_0}$	$\sqrt{c_0}$
				p	1/p	$1 + p\sqrt{c_0}$	1
		$\sqrt{c_0}$	$1, \sqrt{c_0}$	1	0	$\sqrt{c_0}$	$\sqrt{c_0}$
				p	0	$p\sqrt{c_0}$	$p\sqrt{c_0}$
		$p\sqrt{c_0}$	$1, p\sqrt{c_0}$	1	0	$\sqrt{c_0}$	$\sqrt{c_0}$
				p	0	$p\sqrt{c_0}$	$p\sqrt{c_0}$
p	$1, c_0, \sqrt{p}, c_0\sqrt{p}$	$c_0$	$1, c_0$	1	0	$\sqrt{p}$	$\sqrt{p}$
				1	1	$1 + \sqrt{p}$	1
		$\sqrt{p}$	$1,\sqrt{p}$	1	0	$\sqrt{p}$	$\sqrt{p}$
				$c_0$	0	$c_0\sqrt{p}$	$c_0\sqrt{p}$
		$c_0\sqrt{p}$	$1, c_0\sqrt{p}$	1	0	$\sqrt{p}$	$\sqrt{p}$
				$c_0$	0	$c_0\sqrt{p}$	$c_0\sqrt{p}$
$c_0 p$	$1, c_0, \sqrt{c_0 p}, c_0 \sqrt{c_0 p}$	$c_0$	$1, c_0$	1	0	$\sqrt{c_0 p}$	$\sqrt{c_0 p}$
				1	1	$1 + \sqrt{c_0 p}$	1
		$\sqrt{c_0 p}$	$1, \sqrt{c_0 p}$	1	0	$\sqrt{c_0 p}$	$\sqrt{c_0 p}$
				$c_0$	0	$c_0\sqrt{c_0p}$	$c_0\sqrt{c_0p}$
		$c_0\sqrt{c_0p}$	$1, c_0 \sqrt{c_0 p}$	1	0	$\sqrt{c_0 p}$	$\sqrt{c_0 p}$
				$c_0$	0	$c_0\sqrt{c_0p}$	$c_0\sqrt{c_0p}$

TABLE 6. Values of a and b for Proposition 5.2 with  $F = \mathbb{Q}_p$ ,  $p \equiv 1 \mod 4$ . The second column shows the classes of  $\mathbb{Q}_p[\alpha]$  modulo a square, the fourth shows the classes attainable as  $x^2 + \gamma^2$ , the fifth and sixth show values of a and b, the seventh shows the resulting  $a(b + \alpha)$ , and the eighth shows its class. These classes, multiplied by the "attainable classes", should cover the set of "all classes". (It is interesting that the attainable classes are always 1 and  $\gamma^2$ . This may have to do with the field being non-archimedean.)

- (2)  $\operatorname{Sq}(\mathbb{Q}_p[\sqrt{p}]^*) = \{a + b\sqrt{p} : a, b \in \mathbb{Q}_p, \operatorname{ord}_p(a) \leqslant \operatorname{ord}_p(b), \operatorname{digit}_0(a) \in \operatorname{Sq}(\mathbb{F}_p^*)\}.$
- (3)  $\operatorname{Sq}(\mathbb{Q}_p[\operatorname{i}\sqrt{p}]^*) = \{a + \operatorname{i}b\sqrt{p} : a, b \in \mathbb{Q}_p, \operatorname{ord}_p(a) \leq \operatorname{ord}_p(b), \operatorname{ord}_p(a) \equiv 0 \mod 2, \operatorname{digit}_0(a) \in \operatorname{Sq}(\mathbb{F}_p^*)\} \cup \{a + \operatorname{i}b\sqrt{p} : \operatorname{ord}_p(a) \leq \operatorname{ord}_p(b), \operatorname{ord}_p(a) \not\equiv 0 \mod 2, \operatorname{digit}_0(a) \notin \operatorname{Sq}(\mathbb{F}_p^*)\}.$

*Proof.* Parts (2) and (3) have the same proof as the corresponding parts of Proposition 6.1, so we focus on part (1).

Suppose that a + ib is a square. By Lemma 5.3  $a^2 + b^2 = (r^2 + s^2)^2$  for  $r, s \in \mathbb{Q}_p$ . In particular,  $a^2 + b^2$  is a square. By Proposition 4.7(1),  $r^2 + s^2$  has even order, so  $4 \mid \operatorname{ord}(a^2 + b^2)$ . As  $p \equiv 3 \mod 4$ , we cannot have a cancellation in  $a^2 + b^2$ , so

$$\operatorname{ord}(a^2 + b^2) = \min\left\{\operatorname{ord}(a^2), \operatorname{ord}(b^2)\right\} = 2\min\left\{\operatorname{ord}(a), \operatorname{ord}(b)\right\}.$$

As this is a multiple of 4,  $\min{\operatorname{ord}(a), \operatorname{ord}(b)}$  is even.

Now suppose that a and b satisfy the conditions. We have the first condition in Lemma 5.3. To check the second, let  $t_1$  and  $t_2$  be the two candidates for  $r^2$  and suppose, without loss of generality, that  $t_1 = (a + \sqrt{a^2 + b^2})/2$  does not cancel the leading terms. Then

$$\operatorname{ord}(\sqrt{a^2+b^2}) = \min\left\{\operatorname{ord}(a), \operatorname{ord}(b)\right\}$$

is even. The order of a is either higher than this (if b has lower order) or the same, and in any case  $\operatorname{ord}(t_1)$  is even.  $t_2 = -b^2/4t_1$  has also even order, and their product  $-b^2/4$  is not a square (because  $p \equiv 3 \mod 4$ ). This implies that either  $t_1$  or  $t_2$  is a square.  $\Box$ 

**Corollary 6.4.** Let p be a prime number such that  $p \equiv 3 \mod 4$ . Let  $a_0, b_0 \in \mathbb{Z}_p$  such that  $a_0^2 + b_0^2 \equiv -1 \mod p$ . (This pair exists by Proposition 4.7(1).) Then the following statements hold.

$\alpha^2$	all classes	$\gamma^2$	attainable classes	a	b	$a(b+\alpha)$	$\left[a(b+\alpha)\right]$
-1	$1, p, a_0 + ib_0, p(a_0 + ib_0)$	p	1, p	1	0	i	1
				$b_0$	$a_0/b_0$	$a_0 + ib_0$	$a_0 + ib_0$
		$a_0 + \mathrm{i}b_0$	$1, a_0 + ib_0$	1	0	i	1
				p	0	ip	p
		$p(a_0 + \mathrm{i}b_0)$	$1, p(a_0 + \mathrm{i}b_0)$	1	0	i	1
				p	0	ip	p
p	$1, -1, \sqrt{p}, -\sqrt{p}$	-1	1, -1	1	0	$\sqrt{p}$	$\sqrt{p}$
				1	1	$1 + \sqrt{p}$	1
		$\sqrt{p}$	$1, -\sqrt{p}$	1	0	$\sqrt{p}$	$\sqrt{p}$
				-1	0	$-\sqrt{p}$	$-\sqrt{p}$
-p	$1, -1, i\sqrt{p}, -i\sqrt{p}$	-1	1, -1	1	0	$i\sqrt{p}$	$i\sqrt{p}$
				1	1	$1 + i\sqrt{p}$	1
		$i\sqrt{p}$	$1, -i\sqrt{p}$	1	0	$i\sqrt{p}$	$i\sqrt{p}$
				-1	0	$-i\sqrt{p}$	$-i\sqrt{p}$

TABLE 7. Values of a and b for Proposition 5.2 with  $F = \mathbb{Q}_p, p \equiv 3 \mod 4$ . The second column shows the classes of  $\mathbb{Q}_p[\alpha]$  modulo a square, the fourth shows the classes attainable as  $x^2 - \gamma^2$ , the fifth and sixth show values of a and b, the seventh shows the resulting  $a(b+\alpha)$ , and the eighth shows its class.

- (1)  $\mathbb{Q}_p[\mathbf{i}]^* / \operatorname{Sq}(\mathbb{Q}_p[\mathbf{i}]^*) = \{1, p, a_0 + \mathbf{i}b_0, p(a_0 + \mathbf{i}b_0)\}.$
- (2)  $\mathbb{Q}_p[\sqrt{p}]^*/\operatorname{Sq}(\mathbb{Q}_p[\sqrt{p}]^*) = \{1, -1, \sqrt{p}, -\sqrt{p}\}.$
- (3)  $\mathbb{Q}_p[\mathrm{i}\sqrt{p}]^*/\operatorname{Sq}(\mathbb{Q}_p[\mathrm{i}\sqrt{p}]^*) = \{1, -1, \mathrm{i}\sqrt{p}, -\mathrm{i}\sqrt{p}\}.$

*Proof.* Again, parts (2) and (3) are similar to the corresponding ones in Corollary 6.2, so we focus on part (1).

Given a number in the form a + ib, we first ensure that  $a^2 + b^2$  is a square multiplying by  $a_0 + ib_0$  if needed (this changes  $a^2 + b^2 \mod p$  to the opposite). Then we have to ensure that  $\min{\operatorname{ord}(a), \operatorname{ord}(b)}$  is even, multiplying by p if needed, and we have a square because multiplying by p multiplies  $a^2 + b^2$  by  $p^2$  and it will still be a square. 

Now we have determined the possible  $\gamma$ 's. In this case, it is not always true that  $\bar{\gamma} \in \mathbb{Q}_p[\gamma]$ :

- If  $\gamma^2 = -1$  or p, then  $\bar{\gamma} = \gamma$ .
- If  $\gamma^2 = a_0 + ib_0$  or  $p(a_0 + ib_0)$ , the product  $\gamma^2 \bar{\gamma}^2$  is  $a_0^2 + b_0^2$  or  $p^2(a_0^2 + b_0^2)$  respectively. But  $a_0^2 + b_0^2 \equiv -1 \mod p$  implies that  $-\gamma^2 \bar{\gamma}^2$  is a square in  $\mathbb{Q}_p$ , so  $\gamma \bar{\gamma}$  is i times an element of  $\mathbb{Q}_p$ , and in this case we also have  $\bar{\gamma} \in \mathbb{Q}_p[\gamma]$ .
- Otherwise,  $\bar{\gamma} = -\gamma$ , which is a different class.

Hence, the five cases  $p, -1, a_0 + ib_0$  and  $p(a_0 + ib_0)$  give different extensions  $\mathbb{Q}_p[\gamma, \bar{\gamma}]$  and the other four cases give only two extensions, one for  $\pm \sqrt{p}$  and the other for  $\pm i\sqrt{p}$ .

The next step is to determine  $\overline{\mathrm{DSq}}(\mathbb{Q}_p[\alpha], -\gamma^2)$  for each possible  $\alpha$  and  $\gamma$ , and make the quotients. The computation is in Table 7.

6.2. Case p = 2. It only remains to make the analysis for p = 2. This case is different from the rest because we now have seven values of  $\alpha^2$  instead of three: -1, 2, -2, 3, -3, 6and -6.

**Lemma 6.5.** Let  $k, \ell \in \mathbb{N}$  with  $k \ge \ell$ . Let  $a, b, r \in \mathbb{Z}_2$  such that  $\operatorname{ord}_2(2r - a) = \ell$ . Then we have that

$$\frac{a \pm \sqrt{a^2 - b^2 \alpha^2}}{2} \equiv r \mod 2^k$$

if and only if

$$\left(\frac{b}{2}\right)^2 \alpha^2 \equiv r(a-r) \mod 2^{k+\ell}.$$

*Proof.* The first equation is equivalent to  $a \pm \sqrt{a^2 - b^2 \alpha^2} \equiv 2r \mod 2^{k+1}$ , which itself is equivalent to

$$\frac{\pm\sqrt{a^2 - b^2\alpha^2}}{2^{\ell}} \equiv \frac{2r - a}{2^{\ell}} \mod 2^{k+1-\ell}.$$

Since the right-hand side is odd, this is equivalent to all the following identities:

$$\frac{a^2 - b^2 \alpha^2}{2^{2\ell}} \equiv \frac{(2r - a)^2}{2^{2\ell}} \mod 2^{k+2-\ell} \Leftrightarrow$$

$$a^2 - b^2 \alpha^2 \equiv (2r - a)^2 \mod 2^{k+\ell+2} \Leftrightarrow$$

$$-b^2 \alpha^2 \equiv 4r^2 - 4ra \mod 2^{k+\ell+2} \Leftrightarrow$$

$$\left(\frac{b}{2}\right)^2 \alpha^2 \equiv r(a - r) \mod 2^{k+\ell},$$

as we wanted.

**Proposition 6.6.** The following statements hold.

- (1)  $\operatorname{Sq}(\mathbb{Q}_{2}[i]^{*}) = \{a + ib : a, b \in \mathbb{Q}_{p}, \operatorname{ord}_{2}(b) \operatorname{ord}_{2}(a) \ge 2, \operatorname{ord}_{2}(a) \equiv 0 \mod 2, b/4a + \operatorname{digit}_{1}(a) + \operatorname{digit}_{2}(a) \equiv 0 \mod 2\} \cup \{a + ib : a, b \in \mathbb{Q}_{p}, \operatorname{ord}_{2}(a) \operatorname{ord}_{2}(b) \ge 2, \operatorname{ord}_{2}(b) \equiv 1 \mod 2, a/4b + \operatorname{digit}_{1}(b) + \operatorname{digit}_{2}(b) \equiv 0 \mod 2\}.$
- (2)  $\operatorname{Sq}(\mathbb{Q}_2[\sqrt{2}]^*) = \{a + b\sqrt{2} : a, b \in \mathbb{Q}_p, \operatorname{ord}_2(b) \operatorname{ord}_2(a) \ge 1, \operatorname{digit}_2(a) = 0, b/2a + \operatorname{digit}_1(a) \equiv 0 \mod 2\}.$
- (3)  $\operatorname{Sq}(\mathbb{Q}_2[i\sqrt{2}]^*) = \{a + ib\sqrt{2} : a, b \in \mathbb{Q}_p, \operatorname{ord}_2(b) \operatorname{ord}_2(a) \ge 1, \operatorname{digit}_1(a) = \operatorname{digit}_2(a), b/2a + \operatorname{ord}_2(a) + \operatorname{digit}_1(a) \equiv 0 \mod 2\}.$
- (4)  $\operatorname{Sq}(\mathbb{Q}_{2}[\sqrt{3}]^{*}) = \{a+b\sqrt{3} : a, b \in \mathbb{Q}_{p}, \operatorname{ord}_{2}(b) \operatorname{ord}_{2}(a) \ge 2, \operatorname{ord}_{2}(a) \equiv 0 \mod 2, b/4a + \operatorname{digit}_{2}(a) \equiv 0 \mod 2\} \cup \{a+b\sqrt{3} : a, b \in \mathbb{Q}_{p}, \operatorname{ord}_{2}(a) \operatorname{ord}_{2}(b) = 1, \operatorname{ord}_{2}(a) \equiv 0 \mod 2, \operatorname{digit}_{1}(a) + \operatorname{digit}_{1}(b) + \operatorname{digit}_{2}(b) \equiv 0 \mod 2\}.$
- (5)  $\operatorname{Sq}(\mathbb{Q}_2[\operatorname{i}\sqrt{3}]^*) = \{a + \operatorname{i}b\sqrt{3} : a, b \in \mathbb{Q}_p, \operatorname{ord}_2(b) \operatorname{ord}_2(a) \ge 2, \operatorname{ord}_2(a) \equiv 0 \\ \operatorname{mod} 2, \operatorname{digit}_1(a) = 0\} \cup \{a + \operatorname{i}b\sqrt{3} : a, b \in \mathbb{Q}_p, \operatorname{ord}_2(a) = \operatorname{ord}_2(b), \operatorname{ord}_2(a) \equiv 1 \\ \operatorname{mod} 2, \operatorname{digit}_1(a) = 1, a^2 + 3b^2 \in \operatorname{Sq}(\mathbb{Q}_2^*)\}.$
- (6)  $\operatorname{Sq}(\mathbb{Q}_2[\sqrt{6}]^*) = \{a + b\sqrt{6} : a, b \in \mathbb{Q}_p, \operatorname{ord}_2(b) \operatorname{ord}_2(a) \ge 1, \operatorname{ord}_2(a) + \operatorname{digit}_1(a) + \operatorname{digit}_2(a) \equiv 0 \mod 2, b/2a + \operatorname{digit}_2(a) \equiv 0 \mod 2\}.$
- (7) Sq( $\mathbb{Q}_2[i\sqrt{6}]^*$ ) = { $a+ib\sqrt{6}: a, b \in \mathbb{Q}_p, \operatorname{ord}_2(b) \operatorname{ord}_2(a) \ge 1, \operatorname{ord}_2(a) + \operatorname{digit}_2(a) \equiv 0 \mod 2, b/2a + \operatorname{digit}_1(a) \equiv 0 \mod 2$ }.

*Proof.* The first condition in Lemma 5.3 implies that  $a^2 - \alpha^2 b^2$  is a square, so it has even order and ends in 001. This depends on  $\alpha^2 \in \{-1, 2, -2, 3, -3, 6, -6\}$  as well as in the difference  $\operatorname{ord}_2(b) - \operatorname{ord}_2(a)$ , in the way described in Table 5 (where  $c = -\alpha^2$ ). The valid values are as follows:

(i)  $\alpha^2$  is odd and  $\operatorname{ord}_2(b) - \operatorname{ord}_2(a) \ge 2$ . Then  $\sqrt{a^2 - b^2 \alpha^2}$  has the same order than a, and without loss of generality we suppose that  $\operatorname{digit}_1(\sqrt{a^2 - b^2 \alpha^2}) = \operatorname{digit}_1(a)$  (otherwise choose the other square root).

We have that  $t_1$  or  $t_2$  is a square, so it has even order, and  $t_1t_2 = b^2\alpha^2/4$ , which has even order, so both  $t_1$  and  $t_2$  have even order. But, as  $\operatorname{ord}_2(\sqrt{a^2 - b^2\alpha^2}) = \operatorname{ord}_2(a)$  and  $\operatorname{digit}_1(\sqrt{a^2 - b^2\alpha^2}) = \operatorname{digit}_1(a)$ ,

$$\operatorname{ord}_2(2t_1) = \operatorname{ord}_2(a + \sqrt{a^2 - b^2 \alpha^2}) = \operatorname{ord}_2(a) + 1$$
  
44

Г		
L .		
L	_	

which implies  $\operatorname{ord}_2(t_1) = \operatorname{ord}_2(a)$ . As this is even, a has even order.

To simplify the computation, we divide a and b by some power of 4 so that  $\operatorname{ord}_2(a) = 0$ . (Obviously, dividing by 4 does not affect being a square.) Now  $\operatorname{ord}_2(t_1) = 0.$ 

If  $t_1$  is a square, it must be 1 modulo 8. By Lemma 6.5, using that in this case  $\operatorname{ord}_2(2-a) = 0$ , this is equivalent to

$$\left(\frac{b}{2}\right)^2 \alpha^2 \equiv a - 1 \mod 8.$$

If  $\operatorname{ord}_2(b) = 2$ , this implies  $4\alpha^2 \equiv a-1 \mod 8$ , and, using that  $\alpha$  is odd,  $4 \equiv a-1$ mod 8, so  $a \equiv 5 \mod 8$ . Otherwise,  $\operatorname{ord}_2(b) \ge 3$  and  $a \equiv 1 \mod 8$ . In any case,  $t_1$  is square if and only if  $\operatorname{digit}_1(a) = 0$  and  $b/4a + \operatorname{digit}_2(a)$  is even.

If  $t_2$  is a square, as  $t_2 = b^2 \alpha^2 / 4t_1$ ,  $\alpha^2 / t_1$  is also a square and has order 0, so it must be 1 modulo 8 and  $t_1 \equiv \alpha^2 \mod 8$ . Now  $\operatorname{ord}_2(2\alpha^2 - a) = 0$  again, so this is equivalent to

$$\left(\frac{b}{2}\right)^2 \alpha^2 \equiv \alpha^2 (a - \alpha^2) \mod 8,$$

that is

$$\left(\frac{b}{2}\right)^2 \equiv a - \alpha^2 \mod 8.$$

The left-hand side is equivalent to 4 modulo 8 if  $\operatorname{ord}_2(b) = 2$  and 0 otherwise. If  $\alpha^2 = -1$ , a must be 3 or 7 respectively, so digit<sub>1</sub>(a) = 1 and  $b/4a + \text{digit}_2(a)$  is odd. If  $\alpha^2 = 3$ , a is 7 or 3 respectively, so digit<sub>1</sub>(a) = 1 and  $b/4a + \text{digit}_2(a)$  is even. Finally, if  $\alpha^2 = -3$ , a is 5 or 1 respectively, digit<sub>1</sub>(a) = 0 and  $b/4a + \text{digit}_2(a)$  is odd. Putting this together with the results for  $t_1$ , we obtain the first set in the cases (1), (4) and (5).

(ii)  $\alpha^2 = -1$  and  $\operatorname{ord}_2(b) - \operatorname{ord}_2(a) \leq -2$ . Now  $a^2 + b^2$  has the order of  $b^2$ , and  $a \pm \sqrt{a^2 + b^2}$  has the same order as b, so  $\operatorname{ord}_2(t_1) = \operatorname{ord}_2(t_2) = \operatorname{ord}_2(b) - 1$  and  $\operatorname{ord}_2(b)$  is odd. By dividing a and b by a power of 4, we assume that  $\operatorname{ord}_2(b) = 1$ and  $\operatorname{ord}_2(t_1) = \operatorname{ord}_2(t_2) = 0$ .

If  $t_i$  is square,  $t_i \equiv 1 \mod 8$ . Now  $\operatorname{ord}_2(2-a) = 1$  (because a is multiple of 4) and Lemma 6.5 implies that this is equivalent to

$$\left(\frac{b}{2}\right)^2 \equiv 1 - a \mod 16.$$

The left-hand side is 1 if digit<sub>3</sub>( $b^2$ ) is 0, that is, if digit<sub>1</sub>(b) + digit<sub>2</sub>(b) is even, and 9 otherwise, so a is 0 or 8 modulo 16, respectively. Putting this together, we obtain the second set in case (1).

(iii)  $\alpha^2 = 3$  and  $\operatorname{ord}_2(b) - \operatorname{ord}_2(a) = -1$ .  $a^2 - 3b^2$  has the order of  $b^2$ , and  $a \pm \sqrt{a^2 - 3b^2}$ has the order of b, so again  $\operatorname{ord}_2(t_1) = \operatorname{ord}_2(t_2) = \operatorname{ord}_2(b) - 1$  and  $\operatorname{ord}_2(b)$  is odd. By dividing a and b by a power of 4, we assume that  $\operatorname{ord}_2(b) = 1$ ,  $\operatorname{ord}_2(a) = 2$  and  $\operatorname{ord}_2(t_1) = \operatorname{ord}_2(t_2) = 0.$ 

If  $t_i$  is square,  $t_i \equiv 1 \mod 8$ . Now  $\operatorname{ord}_2(2-a) = 1$  again, and this is equivalent to

$$3\left(\frac{b}{2}\right)^2 \equiv a - 1 \mod 16$$

The left-hand side is 3 if  $\operatorname{digit}_3(b^2)$  is 0, that is, if  $\operatorname{digit}_1(b) + \operatorname{digit}_2(b)$  is even, and 11 otherwise, so a is 4 or 12 modulo 16, respectively. Putting this together, we obtain the second set in case (4).

(iv)  $\alpha^2 = -3$  and  $\operatorname{ord}_2(b) = \operatorname{ord}_2(a)$ . In this case  $\operatorname{ord}_2(\sqrt{a^2 + 3b^2}) = \operatorname{ord}_2(a) + 1$  and  $\operatorname{ord}_2(t_1) = \operatorname{ord}_2(t_2) = \operatorname{ord}_2(a) - 1$ , so  $\operatorname{ord}_2(a)$  is odd. By dividing a and b by a power of 4, we assume that  $\operatorname{ord}_2(a) = \operatorname{ord}_2(b) = 1$  and  $\operatorname{ord}_2(t_1) = \operatorname{ord}_2(t_2) = 0$ .

If  $t_i$  is square,  $t_i \equiv 1 \mod 8$ . In this case, as  $\operatorname{ord}_2(a) = 1$ ,  $\operatorname{ord}_2(2-a)$  is at least 2, and

$$3\left(\frac{b}{2}\right)^2 \equiv 1 - a \mod 32.$$

The left-hand side is 3, 27, 11 or 19 if b is  $\pm 2$ ,  $\pm 6$ ,  $\pm 10$  or  $\pm 14$  modulo 32 respectively, so a is 30, 6, 22 or 14 modulo 32. These are exactly the cases where  $a^2 + 3b^2$  is a square and digit<sub>1</sub>(a) = 1 (the opposite remainder modulo 32 for a also makes  $a^2 + 3b^2$  a square, but has digit<sub>1</sub>(a) = 0). So we obtain the second set in case (5). (v)  $\alpha^2$  is even and  $\operatorname{ord}_2(b) - \operatorname{ord}_2(a) \ge 1$ .  $\sqrt{a^2 - b^2 \alpha^2}$  has the same order than a, and without loss of generality we suppose that digit<sub>1</sub>( $\sqrt{a^2 - b^2 \alpha^2}$ ) = digit<sub>1</sub>(a)

(otherwise choose the other square root).

We have that  $t_1t_2 = b^2\alpha^2/4$ , which has odd order, so one of  $t_1$  and  $t_2$  has even order and the other has odd order. But, as  $\operatorname{ord}_2(\sqrt{a^2 - b^2\alpha^2}) = \operatorname{ord}_2(a)$  and  $\operatorname{digit}_1(\sqrt{a^2 - b^2\alpha^2}) = \operatorname{digit}_1(a)$ ,

$$\operatorname{ord}_2(2t_1) = \operatorname{ord}_2(a + \sqrt{a^2 - b^2 \alpha^2}) = \operatorname{ord}_2(a) + 1$$

which implies  $\operatorname{ord}_2(t_1) = \operatorname{ord}_2(a)$ . That is, if a has even order  $t_1$  is a square and otherwise  $t_2$  is a square.

To simplify the computation, we divide a and b by some power of 4 so that  $\operatorname{ord}_2(a)$  is 0 or 1.

If  $t_1$  is a square, then  $\operatorname{ord}_2(a) = \operatorname{ord}_2(t_1) = 0$ , so  $t_1 \equiv 1 \mod 8$ , and  $\operatorname{ord}_2(2-a) = 0$ , so

$$\left(\frac{b}{2}\right)^2 \alpha^2 \equiv a - 1 \mod 8.$$

If  $\operatorname{ord}_2(b) = 1$ , this implies  $\alpha^2 \equiv a - 1 \mod 8$ : for  $\alpha^2 = 2$  or -6,  $a \equiv 3 \mod 8$ , and otherwise  $a \equiv 7 \mod 8$ . If  $\operatorname{ord}_2(b) \ge 2$ , as  $\alpha$  is even, we get  $a \equiv 1 \mod 8$ . In any case,  $t_1$  is square if and only if  $\operatorname{digit}_2(a) = 0$  and  $b/2a + \operatorname{digit}_1(a)$  is even, if  $\alpha^2 = 2$  or -6, or  $\operatorname{digit}_1(a) = \operatorname{digit}_2(a)$  and  $b/2a + \operatorname{digit}_1(a)$  is even, otherwise.

If  $t_2$  is a square, then  $\operatorname{ord}_2(a) = \operatorname{ord}_2(t_1) = 1$ ,  $t_2 = b^2 \alpha^2 / 4t_1$ ,  $\alpha^2 / t_1$  is also a square and has order 0, so it must be 1 modulo 8 and  $t_1 \equiv \alpha^2 \mod 16$ . Now  $\operatorname{ord}_2(2\alpha^2 - a) = 1$ , so

$$\left(\frac{b}{2}\right)^2 \alpha^2 \equiv \alpha^2 (a - \alpha^2) \mod 32,$$

that is,

$$\left(\frac{b}{2}\right)^2 \equiv a - \alpha^2 \mod 16.$$

The left-hand side is equivalent to 4 modulo 16 if  $\operatorname{ord}_2(b) = 2$  and 0 otherwise. If  $\alpha^2 = 2$ , *a* must be 6 or 2 respectively, so  $\operatorname{digit}_2(a) = 0$  and  $b/2a + \operatorname{digit}_1(a)$  is even. If  $\alpha^2 = -2$ , *a* must be 2 or 14 respectively, so  $\operatorname{digit}_1(a) = \operatorname{digit}_2(a)$  and  $b/2a + \operatorname{digit}_1(a)$  is odd. If  $\alpha^2 = 6$ , *a* is 10 or 6 respectively, so  $\operatorname{digit}_1(a) \neq \operatorname{digit}_2(a)$ and  $b/2a + \operatorname{digit}_2(a)$  is even. Finally, if  $\alpha^2 = -6$ , *a* is 14 or 10 respectively, digit<sub>2</sub>(a) = 1 and b/2a + digit<sub>1</sub>(a) is even. Putting this together with the results for  $t_1$ , we obtain the result for the cases (2), (3), (6) and (7).

Now we need the analogous result to Corollaries 6.2 and 6.4. As it turns out, the quotient group has now 16 elements, instead of 4 like for the other primes, so it will be given as the list of generators. The notation  $G = \langle g_1, \ldots, g_n \rangle$  means that G is generated by the elements  $g_1, \ldots, g_n \in G$ . For example, the group  $\{1, c_0, p, c_0p\}$  can be described as  $\langle c_0, p \rangle$ .

Corollary 6.7. The following statements hold.

- (1)  $\mathbb{Q}_2[i]^* / \operatorname{Sq}(\mathbb{Q}_2[i]^*) = \langle 2, 3, 1+i, 1+2i \rangle.$
- (2)  $\mathbb{Q}_2[\sqrt{3}]^* / \operatorname{Sq}(\mathbb{Q}_2[\sqrt{3}]^*) = \langle -1, 2, \sqrt{3}, 1 + \sqrt{3} \rangle.$
- (3)  $\mathbb{Q}_2[i\sqrt{3}]^*/\operatorname{Sq}(\mathbb{Q}_2[i\sqrt{3}]^*) = \langle -1, 2, i\sqrt{3}, 1+2i\sqrt{3} \rangle.$
- (4)  $\mathbb{Q}_2[\alpha]^*/\operatorname{Sq}(\mathbb{Q}_2[\alpha]^*) = \langle -1, 3, \alpha, 1+\alpha \rangle$ , for  $\alpha^2 \in \{2, -2, 6, -6\}$ .

*Proof.* In all cases, the quotient can be computed, as with other primes, by starting with an arbitrary number in  $\mathbb{Q}_p[\alpha]$  and proving that it can be multiplied by some generators to make it a square.

- (1) First we ensure that  $\operatorname{ord}_2(a) \neq \operatorname{ord}_2(b)$  multiplying by 1 + i if needed. Then we ensure that the orders are not consecutive, multiplying by 1 + 2i if they are (such operation will increment the highest order). Then we ensure that the order of a is even, if  $\operatorname{ord}_2(a) < \operatorname{ord}_2(b)$ , and that the order of b is odd, otherwise, multiplying by 2 if needed. Finally, we ensure the condition on the digits of a or b, multiplying by 3 if needed (in general, multiplying a 2-adic number x by 3 preserves digit<sub>2</sub>(x) and inverts digit<sub>1</sub>(x)).
- (2) First we ensure that  $\operatorname{ord}_2(a) \neq \operatorname{ord}_2(b)$  multiplying by  $1 + \sqrt{3}$  if needed. Then we ensure that the difference  $\operatorname{ord}_2(b) \operatorname{ord}_2(a)$  is correct (-1 or at least 2), multiplying by  $\sqrt{3}$  if needed (this inverts the difference). Then we multiply by 2 if needed so that the order of a is even. Finally, we ensure the condition on digits by multiplying by -1: both conditions involve an odd number of digits, so they will invert on multiplication by -1.
- (3) This will be split in two cases.

(a) If  $\operatorname{ord}_2(a) = \operatorname{ord}_2(b)$ , we first make  $\operatorname{digit}_2(a^2+3b^2) = 0$  multiplying by  $1+2i\sqrt{3}$ :

$$(1+2i\sqrt{3})(a+bi\sqrt{3}) = a - 6b + (b+2a)\sqrt{3}$$

and

$$(a-6b)^2 + 3(b+2a)^2 = 13(a^2+3b^2).$$

As  $13 \equiv 5 \mod 8$ , this inverts  $\operatorname{digit}_2(a^2 + 3b^2)$ . Now we make  $\operatorname{digit}_1(a^2 + 3b^2) = 0$  multiplying by  $i\sqrt{3}$ , which will analogously multiply  $a^2 + 3b^2$  by 3. So now  $a^2 + 3b^2$  is a square. Next we multiply by 2 to make the order odd, and by -1 to make  $\operatorname{digit}_1(a) = 1$ , all without affecting  $a^2 + 3b^2$ .

- (b) If  $\operatorname{ord}_2(a) \neq \operatorname{ord}_2(b)$ , we first make  $\operatorname{ord}_2(a) < \operatorname{ord}_2(b)$  multiplying by  $i\sqrt{3}$  if needed, then  $\operatorname{ord}_2(b) \operatorname{ord}_2(a) \ge 2$  multiplying by  $1 + 2i\sqrt{3}$ , then by 2 to make  $\operatorname{ord}_2(a)$  even, and finally by -1 to make  $\operatorname{digit}_1(a) = 0$ .
- (4) First we ensure  $\operatorname{ord}_2(b) \ge \operatorname{ord}_2(a)$  multiplying by  $\alpha$  and then  $\operatorname{ord}_2(b) > \operatorname{ord}_2(a)$ multiplying by  $1 + \alpha$ . In each of the four cases, there are two conditions left, both related to digit<sub>1</sub>(a) and digit<sub>2</sub>(a). We set digit<sub>2</sub>(a) to the required value, multiplying by -1 if needed, and finally digit<sub>1</sub>(a), multiplying by 3.

A depiction of the 16 classes can be found at Figure 7 for  $\alpha^2 = -1$ , at Figure 8 for  $\alpha^2 = 2$ , at Figure 9 for  $\alpha^2 = -2$ , at Figure 10 for  $\alpha^2 = 3$ , at Figure 11 for  $\alpha^2 = -3$ , at Figure 12 for  $\alpha^2 = 6$ , and at Figure 13 for  $\alpha^2 = -6$ .

Now we have to compute which of the classes are "paired" in the sense of being the classes of  $\gamma^2$  and  $\bar{\gamma}^2$ , so that they give the same extension  $\mathbb{Q}_p[\gamma, \bar{\gamma}]$ . In general, if  $\gamma = t_1 + t_2 \alpha$ ,

$$\gamma^2 \bar{\gamma}^2 = (t_1 + t_2 \alpha)(t_1 - t_2 \alpha) = t_1^2 - t_2^2 \alpha^2$$

which is always in  $\mathbb{Q}_2$ , so two paired classes differ in a factor in  $\mathbb{Q}_2$ . In the last column of Tables 8 to 14 we give the pair of each class. After identifying the paired classes, if  $\alpha^2 \in \{-1, -2, 3, 6\}$ , 9 classes remain (not counting 1), and if  $\alpha^2 \in \{2, -3, -6\}$ , 11 classes remain.

The next step in order to achieve the classification is to compute the classes of

 $\overline{\mathrm{DSq}}(\mathbb{Q}_2[\alpha], -\gamma^2)$ 

for each possible  $\alpha$  and  $\gamma$ . For the other primes this meant three different  $\alpha$ 's and two or three  $\gamma$ 's for each one, but here we need seven  $\alpha$ 's and nine or eleven  $\gamma$ 's for each one. To simplify what would otherwise be a very long and error-prone computation, we will now use the Hilbert symbol for  $\mathbb{Q}_2[\alpha]$ .

**Lemma 6.8.** The Hilbert symbol  $(a,b)_F$  in any field F (concretely  $F = \mathbb{Q}_2[\alpha]$ ) has the following properties:

- (1)  $(1, u)_F = (u, -u)_F = 1$  for any u.
- (2)  $(u, v)_F = (v, u)_F$ .
- (3)  $(u, v)_F = 1$  if and only if  $v \in \overline{\text{DSq}}(F, -u)$ .
- (4)  $(u, v_1v_2)_F = (u, v_1)_F (u, v_2)_F$ .

We define a subset

$$S_{\alpha} \subset (\mathbb{Q}_2[\alpha]^* / \operatorname{Sq}(\mathbb{Q}_2[\alpha]^*))^2$$

for different values of  $\alpha$ :  $S_i$  is defined in Table 8,  $S_{\sqrt{2}}$  in Table 9,  $S_{i\sqrt{2}}$  in Table 10,  $S_{\sqrt{3}}$  in Table 11,  $S_{i\sqrt{3}}$  in Table 12,  $S_{\sqrt{6}}$  in Table 13, and  $S_{\sqrt{-6}}$  in Table 14.

**Lemma 6.9.** Let  $F = \mathbb{Q}_2[\alpha]$  be a degree 2 extension of  $\mathbb{Q}_2$ . If  $(u, v)_F = 1$  for all  $(u, v) \in S_\alpha$  and there exists  $(u, v) \in (\mathbb{Q}_2[\alpha]^*)^2$  such that  $(u, v)_F = -1$ , then  $(u, v)_F = -1$  for all  $(u, v) \notin S_\alpha$ .

*Proof.* In all cases, the set  $\{v : (u, v) \in S_{\alpha}\}$  for a fixed  $u \neq 1$  is a multiplicative subgroup of the quotient  $\mathbb{Q}_2[\alpha]^*/\operatorname{Sq}(\mathbb{Q}_2[\alpha]^*)$  with eight elements. If all them have  $(u, v)_F = 1$  and other v has  $(u, v)_F = -1$ , then by multiplicativity all the other v have  $(u, v)_F = -1$ .  $\Box$ 

**Proposition 6.10.** For all degree 2 extensions  $F = \mathbb{Q}_2[\alpha]$ ,  $(u, v)_F = 1$  if and only if  $(u, v) \in S_{\alpha}$ .

*Proof.* We use Lemma 6.9 for each possible  $\alpha$ . For some values  $(u, v) \in S_{\alpha}$ , it can be easily computed that they have  $(u, v)_F = 1$ , and this can be deduced for the rest of  $S_{\alpha}$  by Lemma 6.8. Then we just need to prove that there is (u, v) such that  $(u, v)_F = -1$ , and we are done.

• Case  $\alpha = i$ : since we have that 3 - 2 = 1, 2(1 + 2i) - 2(1 + i) = 2i, 6(-1 + 3i) + 6(1 + 2i) = 30i, (2 + 2i) - 2 = 2i and 4(1 + 2i) - 3 = 1 + 8i, all of which are squares, all elements of  $S_i$  have  $(u, v)_F = 1$ . Now we prove that  $(2, 1 + 2i)_F = -1$ , for which we have to see that  $x^2 - 2$  will never be 1 + 2i times a square, for any x. Suppose this happens. Then, if  $x^2 = a + bi$ , the orders of a and b must differ

in at least 2 and the orders of a - 2 and b differ in 1. This implies that a - 2 and a have different order. There are two possibilities:

- $-\operatorname{ord}_2(a-2) = 1$  and  $\operatorname{ord}_2(a) > 1$ . Then, b has order 0 or 2. Since a + bi is a square,  $\operatorname{ord}_2(a)$  is at most 0, a contradiction.
- $\operatorname{ord}_2(a-2) > 1$  and  $\operatorname{ord}_2(a) = 1$ . Since a + bi is a square, b has odd order at most -1, but it should differ in 1 with  $\operatorname{ord}_2(a-2)$ , also a contradiction.
- Case  $\alpha = \sqrt{2}$ : now the pairs that add up to a square are 3-2=1, 5-4=1,  $(1+\sqrt{2})-\sqrt{2}=1$ ,  $(2+\sqrt{2})-(1+\sqrt{2})=1$ ,  $4(1+\sqrt{2})-3=1+4\sqrt{2}$  and  $3+2\sqrt{2}=3+2\sqrt{2}$ . We prove that  $(-1,\sqrt{2})_F=-1$ , for which we have to see that  $x^2+1$  will never be  $\sqrt{2}$  times a square. Suppose it is. Let  $x^2=a+b\sqrt{2}$ . We have  $\operatorname{ord}_2(a) < \operatorname{ord}_2(b) < \operatorname{ord}_2(a+1)$ , which is possible only if a has order 0 and ends in 11. But then b must have order 1 and a ends in 011, so a+1 has order 2, and  $a+1+b\sqrt{2}$  cannot be  $\sqrt{2}$  times a square (the difference in order between a+1 and b should be at least 2).
- Case  $\alpha = i\sqrt{2}$ : now the pairs that add up to a square are 2 1 = 1, 5 4 = 1,  $(1 + i\sqrt{2}) i\sqrt{2} = 1$ ,  $3(1 + i\sqrt{2}) 3(-2 + i\sqrt{2}) = 9$ ,  $-1 + 2i\sqrt{2} = -1 + 2i\sqrt{2}$  and  $2(1 + i\sqrt{2}) 3 = -1 + 2i\sqrt{2}$ . We prove that  $(-1, 1 + i\sqrt{2})_F = -1$ , for which we have to see that  $x^2 + 1$  will never be  $1 + i\sqrt{2}$  times a square. Suppose it is. Let  $x^2 = a + bi\sqrt{2}$ . We have  $\operatorname{ord}_2(a) < \operatorname{ord}_2(b) = \operatorname{ord}_2(a + 1)$ , which is possible only if  $\operatorname{ord}_2(a) = 0$ . Also,  $b/2a + \operatorname{digit}_1(a)$  is odd, so that  $\operatorname{ord}_2(a + 1) = \operatorname{ord}_2(b)$ . This makes  $a + bi\sqrt{2}$  not a square.
- Case  $\alpha = \sqrt{3}$ : now the pairs that add up to a square are 2 1 = 1, 5 4 = 1,  $(1 + \sqrt{3}) \sqrt{3} = 1$ ,  $2(3 + \sqrt{3}) 2(1 + \sqrt{3}) = 4$ ,  $2(3 + \sqrt{3}) 2 = 4 + 2\sqrt{3}$  and  $-1 + \sqrt{3}(4 + 2\sqrt{3}) = 5 + 4\sqrt{3}$ . We prove that  $(-1, 1 + \sqrt{3})_F = -1$ , for which we have to see that  $x^2 + 1$  will never be  $1 + \sqrt{3}$  times a square. Suppose it is. Let  $x^2 = a + b\sqrt{3}$ .

- If  $\operatorname{ord}_2(b) - \operatorname{ord}_2(a) \ge 2$ ,  $\operatorname{ord}_2(a) = 0$  and a must end in 11 so that  $\operatorname{ord}_2(a+1) = \operatorname{ord}_2(b)$ . Also,  $b/4a + \operatorname{digit}_2(a)$  is odd. This makes  $a + b\sqrt{3}$  not a square.

- If  $\operatorname{ord}_2(a) \operatorname{ord}_2(b) = 1$ ,  $\operatorname{ord}_2(a)$  must be even, so it is impossible that a + 1 and b have the same order.
- Case  $\alpha = i\sqrt{3}$ : now the pairs that add up to a square are 2 1 = 1, 3 2 = 1,  $(1 + 2i\sqrt{3}) 2i\sqrt{3} = 1$ ,  $(1 + 2i\sqrt{3}) 2(-6 + i\sqrt{3}) = 13$ ,  $2 + 2i\sqrt{3} = 2 + 2i\sqrt{3}$  and  $2(-6 + i\sqrt{3}) + 14 = 2 + 2i\sqrt{3}$ . We prove that  $(-1, i\sqrt{3})_F = -1$ , for which we have to see that  $x^2 + 1$  will never be  $i\sqrt{3}$  times a square. Suppose it is. Let  $x^2 = a + bi\sqrt{3}$ .
  - If  $\operatorname{ord}_2(b) \operatorname{ord}_2(a) \ge 2$ ,  $\operatorname{ord}_2(a) = 0$  and a must end in 11 so that  $\operatorname{ord}_2(a+1) \ge \operatorname{ord}_2(b)$ , but then  $a + bi\sqrt{3}$  is not a square.
  - If  $\operatorname{ord}_2(a) = \operatorname{ord}_2(b)$ , they must be odd, and  $\operatorname{ord}_2(a+1)$  cannot be greater than  $\operatorname{ord}_2(b)$ , so they must be equal, and  $\operatorname{ord}_2(a) = \operatorname{ord}_2(a+1) = \operatorname{ord}_2(b) < 0$ . If the order is -3 or less,

$$(a+1)^2 + 3b^2 = a^2 + 3b^2 + 2a + 1$$

must be three times a square in  $\mathbb{Q}_2$ , but  $\operatorname{ord}_2(a^2 + 3b^2) \leq -4$  and  $\operatorname{digit}_1((a + 1)^2 + 3b^2) = \operatorname{digit}_1(a^2 + 3b^2) = 0$ , so it is impossible.

If the order is -1, let  $a_1 = 4a$ ,  $b_1 = 4b$ . We want  $a_1 + 4 + b_1 i\sqrt{3}$  to be  $i\sqrt{3}$  times a square. Let  $a_2 + b_2 i\sqrt{3}$  be this square. All of  $a_1$ ,  $b_1$ ,  $a_2$  and  $b_2$  have order 1, and

$$a_1 + 4 + b_1 i\sqrt{3} = i\sqrt{3}(a_2 + b_2 i\sqrt{3})$$
  
49

		1				1 -	+ i			1 -	+2i			-:	1 +	3i				
		1	2	3	6	1	2	3	6	1	2	3	6	1	2	3	6	$a_1$	$b_1$	pair
1	1	•	•	•	•	•	•	٠	٠	•	•	•	•	•	•	٠	٠			1
	2	•	•	•	•	•	•	•	•									1	2	2
	3	•	•	•	•					•	•	•	•					1	1	3
	6	•	•	•	•									•	•	٠	•	1	1	6
1 + i	1	•	•			•	•					•	•			•	•	3	0	2
	2	•	•			•	•			٠	•			•	•			3	0	1
	3	•	•					٠	٠			•	•	•	•			3	0	6
	6	•	•					٠	٠	٠	•					٠	•	3	0	3
1 + 2i	1	•		•			•		•	•		•			•		•	2	0	3
	2	•		•			•		٠		•		•	•		٠		2	0	6
	3	•		•		•		•		•		•		•		•		2	0	1
	6	•		•		•		•			•		•		•		•	2	0	2
-1 + 3i	1	•			•		•	•			•	•		•			•	2	0	6
	2	•			•		•	•		٠			•		•	•		2	0	3
	3	•			•	•			•		٠	٠			•	•		2	0	2
	6	•			•	•			•	•			•	•			•	2	0	1

TABLE 8.  $S_i$  as a subset of  $(\mathbb{Q}_2[i]^*/\operatorname{Sq}(\mathbb{Q}_2[i]^*))^2$ . For each row except the first, we need two normal forms to cover all the classes: a = 1, b = 0 and  $a = a_1, b = b_1$ . The last column indicates the second index of the class that pairs with each class; the first index is always the same.

implies that  $a_1 + 4 = -3b_2$  and  $b_1 = a_2$ . Hence,  $\operatorname{digit}_1(b_1) = \operatorname{digit}_1(a_2) = 1$ .  $a_1$  is 30, 6, 22 or 14 modulo 32 exactly when  $b_1$  can take those remainders, so  $a_1 \equiv b_1 \mod 32$ , and

$$-3b_2 = a_1 + 4 \equiv b_1 + 4 = a_2 + 4 \mod 32$$

But now  $a_2$  is 30, 6, 22 and 14 when  $b_2$  is  $\pm 2, \pm 6, \pm 10$  and  $\pm 14$  respectively, and the equation does not hold in any case.

- Case  $\alpha = \sqrt{6}$ : now the pairs that add up to a square are 2 1 = 1, 5 4 = 1,  $(1 + \sqrt{6}) \sqrt{6} = 1$ ,  $3(1 + \sqrt{6}) 3(6 + \sqrt{6}) = -15$ ,  $-1 + 2\sqrt{6} = -1 + 2\sqrt{6}$  and  $2(1 + \sqrt{6}) 3 = -1 + 2\sqrt{6}$ . We prove that  $(-1, 1 + \sqrt{6})_F = -1$ , for which we have to see that  $x^2 + 1$  will never be  $1 + \sqrt{6}$  times a square. Suppose it is. Let  $x^2 = a + b\sqrt{6}$ . We have  $\operatorname{ord}_2(a) < \operatorname{ord}_2(b) = \operatorname{ord}_2(a + 1)$ , which is possible only if  $\operatorname{ord}_2(a) = 0$ . Also,  $b/2a + \operatorname{digit}_1(a)$  is odd, so that  $\operatorname{ord}_2(a + 1) = \operatorname{ord}_2(b)$ . This makes  $a + b\sqrt{6}$  not a square.
- Case  $\alpha = i\sqrt{6}$ : now the pairs that add up to a square are 2 1 = 1, 3 2 = 1,  $(1 + i\sqrt{6}) i\sqrt{6} = 1$ ,  $(-6 + i\sqrt{6}) (1 + i\sqrt{6}) = -7$ ,  $3 + 2i\sqrt{6} = 3 + 2i\sqrt{6}$  and  $6(1 + i\sqrt{6}) 3 = 3 + 6i\sqrt{6}$ . We prove that  $(-1, i\sqrt{6})_F = -1$ , for which we have to see that  $x^2 + 1$  will never be  $i\sqrt{6}$  times a square. Suppose it is. Let  $x^2 = a + bi\sqrt{6}$ . We have  $\operatorname{ord}_2(a) < \operatorname{ord}_2(b) < \operatorname{ord}_2(a + 1)$ , which is possible only if a has order 0 and ends in 11. But then b must have order 1 and a ends in 011, so a + 1 has order 2, and  $a + 1 + bi\sqrt{6}$  cannot be  $i\sqrt{6}$  times a square (the difference in order between a + 1 and b should be at least 2).

After finding the values of the Hilbert symbol for each  $\alpha$ , we have  $\overline{\text{DSq}}(\mathbb{Q}_2[\alpha], -\gamma^2)$ : it is the row indexed by  $\gamma^2$  of the corresponding table. It always contains eight classes, so we need two pairs (a, b) to cover all classes: we can always take one of them (1, 0), and the other is  $(a_1, b_1)$  such that the class of  $a_1(b_1 + \alpha)$  is marked (in the row of  $\gamma^2$ ) if and

:	:	:	:	•	2		1		•	•	•	:	•	•	•	:
	•	•	•	•					•	•					•	•
	•								•	•		:			٠	٠
														•		
					2.1				1	1	1			•	1	1
•																
	:											:			:	
•	•		•	•	*		•		•	•	•	•	•	•	•	•
:	:	ļ,	1	•	•	1	:		2	1	:	:	2	1	1	ļ
:	:	:	1	•	•	:	•		-	:	1	:	ł	:	1	1
:	:	:		•	•	•	:		:	:	:	:	-	:	:	:
:	:	:		•	•	:	:		:	:	:	:		:	:	:
:	:		:	•	•	•	•		:	:	:	:		:		:
:	:		:	•	•	•	•		:	:	:	:		:		:

FIGURE 7. The 16 classes of Table 8. Each class contains the points x + yi with a given symbol, where x and y are the horizontal and vertical coordinates. The circles, triangles, squares and diamonds correspond to the four values of the first index (here 1, 1 + i, 1 + 2i and -1 + 3i), and the colors red, green, purple and blue to the four values of the second index (here 1, 2, 3 and 6).

		1				$ \sqrt{2}$	$\overline{2}$			1 -	$+\sqrt{2}$			2 -	$+\sqrt{2}$					
		1	-1	3	-3	1	-1	3	-3	1	-1	3	-3	1	-1	3	-3	$a_1$	$b_1$	pair
1	1	•	•	٠	•	•	•	•	•	•	•	•	•	٠	•	•	•			1
	-1	•	•	٠	•									٠	•	•	•	1	2	-1
	3	•	•	•	•	•	•	٠	•									1	1	3
	-3	•	•	•	•					٠	•	•	•					1	1	-3
$\sqrt{2}$	1	•		•			•		•		•		•	•		•		-1	0	-1
	-1	•		٠		•		•		•		•		٠		•		-1	0	1
	3	•		٠			•		•	٠		•			•		•	-1	0	-3
	-3	•		•		•		•			•		•		•		•	-1	0	3
$1+\sqrt{2}$	1	•			•		•	•			•	•		•			•	-1	0	-1
	-1	•			•	•			•	٠			•	٠			•	-1	0	1
	3	•			•		•	•		٠			•		•	•		-1	0	-3
	-3	•			•	•			•		•	•			•	•		-1	0	3
$2+\sqrt{2}$	1	•	•			•	•			•	•			•	•			3	0	1
	-1	•	•					•	•			•	•	٠	•			3	0	-1
	3	•	•			•	•					•	•			•	•	3	0	3
	-3	•	•					•	•	٠	•					•	•	3	0	-3

TABLE 9.  $S_{\sqrt{2}}$  as a subset of  $(\mathbb{Q}_2[\sqrt{2}]^*/\operatorname{Sq}(\mathbb{Q}_2[\sqrt{2}]^*))^2$ .

:	•	:	•	:	:	•	•	:		1		1	:	
	•					•	•	:					:	1
1	•	:	•		:	ļ,	:	:	•		:	:		
			2					:				:		

FIGURE 8. The 16 classes of Table 9.




TABLE 10.  $S_{i\sqrt{2}}$  as a subset of  $(\mathbb{Q}_2[i\sqrt{2}]^*/\operatorname{Sq}(\mathbb{Q}_2[i\sqrt{2}]^*))^2$ .

only if that of  $\alpha$  is unmarked. A possibility is included at the right of the corresponding row.

## 6.3. Proof of Theorem F.

(a) First we prove existence. Let  $\lambda, -\lambda, \mu, -\mu$  be the eigenvalues of  $\Omega_0^{-1}M$ . If  $\lambda^2$  is in  $\mathbb{Q}_p$ ,  $\mu^2$  is also in  $\mathbb{Q}_p$ . Let  $\{u_1, v_1, u_2, v_2\}$  be the associated basis. By Corollary 52




FIGURE 9. The 16 classes of Table 10.




TABLE 11.  $S_{\sqrt{3}}$  as a subset of  $(\mathbb{Q}_2[\sqrt{3}]^*/\operatorname{Sq}(\mathbb{Q}_2[\sqrt{3}]^*))^2$ .

3.5, there is a matrix S with entries in  $\mathbb{Q}_p[\lambda,\mu]$  such that  $S^T\Omega_0S=\Omega_0$  and

$$S^{T}MS = \begin{pmatrix} r & 0 & 0 & 0\\ 0 & c_{1}r & 0 & 0\\ 0 & 0 & s & 0\\ 0 & 0 & 0 & c_{2}s \end{pmatrix}$$




FIGURE 10. The 16 classes of Table 11.




TABLE 12.  $S_{i\sqrt{3}}$  as a subset of  $(\mathbb{Q}_2[i\sqrt{3}]^*/\operatorname{Sq}(\mathbb{Q}_2[i\sqrt{3}]^*))^2$ .

if and only if  $\lambda = r\sqrt{-c_1}$  and  $\mu = s\sqrt{-c_2}$ . There are always  $r, s \in \mathbb{Q}_p$  and  $c_1, c_2 \in X_p$  such that this is possible; moreover, there may be two valid values of  $c_1$  or  $c_2$ .

	•••••••••••••••••••••••••••••••••••••••			:	*	•				•	••••••	•	•	•
--	---	--	--	---	---	---	--	--	--	---	--------	---	---	---

FIGURE 11. The 16 classes of Table 12.

		1					$\overline{6}$			1 -	$+\sqrt{6}$			6 -	$+\sqrt{6}$					
		1	-1	3	-3	1	-1	3	-3	1	-1	3	-3	1	-1	3	-3	$a_1$	$b_1$	pair
1	1	•	•	٠	•	•	•	•	•	•	•	•	•	٠	٠	•	•			1
	-1	•	•	•	•	•	•	•	•									1	1	-1
	3	•	•	•	•									•	•	٠	٠	1	6	3
	-3	•	•	٠	•					•	•	•	•					1	1	-3
$\sqrt{6}$	1	•	•			•	•					•	•			•	•	3	0	-1
	-1	•	•			•	•			•	•			٠	•			3	0	1
	3	•	•					•	•	•	•					٠	•	3	0	-3
	-3	•	•					•	•			•	•	•	•			3	0	3
$1 + \sqrt{6}$	1	•			•		•	•			•	•		٠			•	-1	0	3
	-1	•			•		•	•		•			•		•	٠		-1	0	-3
	3	•			•	•			•	•			•	٠			•	-1	0	1
	-3	•			•	•			•		•	•			•	٠		-1	0	-1
$6 + \sqrt{6}$	1	•		•			•		•	•		•			•		•	-1	0	-3
	-1	•		٠			•		•		•		•	٠		٠		-1	0	3
	3	•		•		•		•			•		•		•		٠	-1	0	-1
	-3	•		•		•		•		•		•		٠		٠		-1	0	1

TABLE 13.  $S_{\sqrt{6}}$  as a subset of  $(\mathbb{Q}_2[\sqrt{6}]^*/\operatorname{Sq}(\mathbb{Q}_2[\sqrt{6}]^*))^2$ .

A matrix S with this property must have the form  $\Psi_1 D \Psi_2^{-1}$ , where

$$\Psi_1 = \begin{pmatrix} u_1 & v_1 & u_2 & v_2 \end{pmatrix}, \Psi_2 = \begin{pmatrix} \lambda & -\lambda & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & \mu & -\mu \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

and D is a diagonal matrix. As  $\Psi_2^{-1}$  is "box-diagonal", the first two columns of S come from the first two of  $\Psi_1$  and the last two of S come from the last two of

	•	:	•	:	:	:	:				:	:	:
	•			:	:	:	:			1		Ξ.	
•	•	٠	•	1.			-	•	• •	•	•	•	•
		•	•		•		-				•	•	•
				1.1	•		-	•	• •		•	•	•
	•			•				•		•	•		•
								•	 		-		
•	•	•	•	•	•	•	•	-	•	•	•	•	•
	:			•								:	

FIGURE 12. The 16 classes of Table 13.




TABLE 14.  $S_{i\sqrt{6}}$  as a subset of  $(\mathbb{Q}_2[i\sqrt{6}]^*/\operatorname{Sq}(\mathbb{Q}_2[i\sqrt{6}]^*))^2$ .

 $\Psi_1$ . That is to say, the existence of a D for which  $\Psi_1 D \Psi_2^{-1}$  has entries in  $\mathbb{Q}_p$  is equivalent to the existence of  $D_1$  and  $D_2$  such that

$$\begin{pmatrix} u_1 & v_1 \end{pmatrix} D_1 \begin{pmatrix} \lambda & -\lambda \\ 1 & 1 \end{pmatrix}$$
 and  $\begin{pmatrix} u_2 & v_2 \end{pmatrix} D_2 \begin{pmatrix} \mu & -\mu \\ 1 & 1 \end{pmatrix}$ 

have entries in  $\mathbb{Q}_p$ .

By Theorem D, there are always  $c_1$  and  $c_2$  in  $X_p$  for which  $D_1$  and  $D_2$  exist, so this leads to case (1).

|--|

FIGURE 13. The 16 classes of Table 14.

Now suppose that  $\lambda^2 \notin \mathbb{Q}_p$ . If  $\lambda \in \mathbb{Q}_p[\lambda^2]$ , we are in the situation of Proposition 5.1.  $\lambda$  and  $\mu$  are in a degree 2 extension  $\mathbb{Q}_p[\alpha]$ , and M is equivalent by multiplication by a symplectic matrix to the matrix of case (2) for some  $r, s \in \mathbb{Q}_p$ . The possible values of  $\alpha^2$  are the classes of  $\mathbb{Q}_p^*$  modulo squares, that is, precisely the elements of  $Y_p$ , and we have case (2).

Finally, if  $\lambda \notin \mathbb{Q}_p[\lambda^2]$ , we are in the situation of Proposition 5.2: we have a hierarchy of extensions

$$\mathbb{Q}_p \subsetneq \mathbb{Q}_p[\alpha] \subsetneq \mathbb{Q}_p[\gamma, \bar{\gamma}],$$

and M is equivalent by multiplication by a symplectic matrix to the matrix in case (3), for some  $r, s \in \mathbb{Q}_p$ , which depends on the parameters  $\alpha^2$ ,  $t_1$ ,  $t_2$ , a and b. The only ones that are not fixed by the extension are a and b: a choice of them is valid if and only if

$$\frac{a\alpha\gamma(b+\alpha)}{u^T\Omega_0\hat{u}} \in \mathrm{DSq}(F[\alpha], -\gamma^2).$$

The denominator is a constant, which implies that the valid values form a class modulo  $DSq(F[\alpha], -\gamma^2)$  in  $F[\alpha]$ . These classes are as described in Tables 6 to 14, depending on p and  $\alpha^2$ . After substituting  $c = \alpha^2$  and extracting  $t_1, t_2, a_1$  and  $b_1$  from these tables, we obtain case (3) of the theorem.

- (b) Finally we prove uniqueness. Let N and N' be the two normal forms. The case (1), (2) or (3) of the normal form is determined uniquely by the eigenvalues of A = Ω<sub>0</sub><sup>-1</sup>M, so both N and N' are in the same case. Now we split between the three cases.
  - In case (1), there are two eigenvalues of A composing the first block of the normal form and two eigenvalues composing the second one. Hence, by Theorem D, the normal form is unique up to changing the order of the blocks.

- In case (2), the extension which contains the eigenvalues is different for each c, hence N = N'.
- In case (3), analogously, the extension is different for each c,  $t_1$  and  $t_2$ , so these parameters must coincide. If a and b do not coincide, we have a number in  $\mathbb{Q}_p[\alpha]$  which is in  $\mathrm{DSq}(\mathbb{Q}_p[\alpha], -\gamma^2)$  and which cannot be there by previous results (Tables 6 to 14), so a and b must also coincide and N = N'.

#### 6.4. Proof of Theorem G.

(a) First we prove existence. Suppose first that the eigenvalues of A are  $\lambda$ ,  $\lambda$ ,  $-\lambda$  and  $-\lambda$ , with  $\lambda \neq 0$ . If A is diagonalizable, Lemma 3.6 implies that there is a symplectic basis  $\{u_1, v_1, u_2, v_2\}$  such that  $Au_i = \lambda u_i$  and  $Av_i = -\lambda v_i$ . This means we are in the first case of Theorem F, that is, case (1) of this theorem, and we can proceed from there.

If A is not diagonalizable, we can also apply Lemma 3.6, getting a symplectic basis  $\{u_1, v_1, u_2, v_2\}$ , or equivalently a symplectic matrix  $\Psi_1$ , such that

$$\Psi_1^{-1}A\Psi_1 = J = \begin{pmatrix} \lambda & 0 & 1 & 0\\ 0 & -\lambda & 0 & 0\\ 0 & 0 & \lambda & 0\\ 0 & -1 & 0 & -\lambda \end{pmatrix}.$$

If  $\lambda \in \mathbb{Q}_p$ , we can rearrange the coordinates to make J equal to  $\Omega_0^{-1}M_2$ , where  $M_2$  is the matrix in case (2) with  $r = \lambda$ . As  $\Psi_1$  is symplectic, rearranging its columns in the same way gives the S we need.

Otherwise, we can write  $\lambda = r\alpha$ , with  $\alpha = \sqrt{c}$  for some  $c \in Y_p$ . Let  $M_2$  be the matrix in case (3) and  $A_2 = \Omega_0^{-1} M_2$ . We have  $\Psi_2^{-1} A_2 \Psi_2 = J$ , where

$$\Psi_2 = \begin{pmatrix} 0 & \alpha z_1 & -\alpha z_1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & z_1 & z_1 & 0 \\ \alpha & -t_1 & t_1 & -\alpha \end{pmatrix},$$

where

$$z_1 = \frac{2\alpha}{a(1-\alpha^2)}, t_1 = \frac{1+\alpha^2}{r(1-\alpha^2)}.$$

A matrix that commutes with J has the form

$$D = \begin{pmatrix} d_1 & 0 & d_2 & 0 \\ 0 & d_3 & 0 & 0 \\ 0 & 0 & d_1 & 0 \\ 0 & d_4 & 0 & d_3 \end{pmatrix}$$

We apply the condition  $D^T \Psi_1^T \Omega_0 \Psi_1 D = \Psi_2^T \Omega_0 \Psi_2$  of Proposition 3.2. As  $\Psi_1$  is symplectic,  $\Psi_1^T \Omega_0 \Psi_1 = \Omega_0$  and the condition becomes  $d_1 d_3 = -2\alpha z_1$  and  $d_1 d_4 + d_2 d_3 = 0$ .

We also want that S has the entries in  $\mathbb{Q}_p$ . The first and fourth columns of  $\Psi_1$  are the eigenvectors of A with value  $\lambda$  and  $-\lambda$ , which are conjugate up to a multiplicative constant: we call them u and  $k\bar{u}$ . The second and third columns correspond to v and v' such that  $Av = \lambda v + u$  and  $Av' = -\lambda v' - k\bar{u}$ . This implies

 $A\bar{v} = -\lambda\bar{v} + \bar{u}$ , that is,  $v' = -k\bar{v}$ , and

$$S\Psi_2 = \Psi_1 D$$
  
=  $\begin{pmatrix} u & -k\bar{v} & v & k\bar{u} \end{pmatrix} D$   
=  $\begin{pmatrix} d_1 u & -d_3 k\bar{v} + d_4 k\bar{u} & d_2 u + d_1 v & d_3 k\bar{u} \end{pmatrix}.$ 

If we call  $c_i$  the *i*-th column of S, we have

$$c_{2} + \alpha c_{4} = d_{1}u, c_{2} - \alpha c_{4} = d_{3}k\bar{u} \Rightarrow d_{3}k = d_{1};$$
  

$$\alpha z_{1}c_{1} + z_{1}c_{3} - t_{1}c_{4} = -d_{3}k\bar{v} + d_{4}k\bar{u} = \bar{d}_{1}\bar{v} + d_{4}k\bar{u};$$
  

$$-\alpha z_{1}c_{1} + z_{1}c_{3} + t_{1}c_{4} = d_{2}u + d_{1}v.$$

Changing sign and conjugating

$$\alpha z_1 c_1 + z_1 c_3 - t_1 c_4 = d_1 \bar{v} - d_2 \bar{u},$$

so we have  $d_4k = -\bar{d}_2$ . We can take  $d_2 = d_4 = 0$ , and the condition reduces to find  $d_1$  such that

(6.1) 
$$d_1 \bar{d}_1 = \frac{-4\alpha^2 k}{a(1-\alpha^2)}.$$

So we need (6.1) to be in

$$\mathrm{DSq}(\mathbb{Q}_p, -\alpha^2) = \mathrm{DSq}(\mathbb{Q}_p, -c),$$

which is possible for a value of a in  $\mathbb{Q}_p^*/\mathrm{DSq}(\mathbb{Q}_p, -c)$ . This quotient is exactly the set called  $\{1, h_p(c)\}$  in the statement.

Now suppose that the eigenvalues are  $\lambda, -\lambda, 0, 0$  for  $\lambda \neq 0$ . By Lemma 3.6, we can choose  $u_1$  and  $v_1$  as eigenvectors with values  $\lambda$  and  $-\lambda$  such that  $u_1^T \Omega_0 v_1 = 1$  and they are  $\Omega_0$ -complementary to the kernel of A. We then complete to a symplectic basis  $\{u_1, v_1, u_2, v_2\}$ , with  $Au_2 = Av_2 = 0$ . At this point we are again in case (1) of Theorem F, with  $c_2 = 0$ .

The only case left is that all the eigenvalues of A are 0. Then Theorem 3.8 gives a good tuple K with sum 4 and a basis. The possible cases for K are (4), (2, 2), (2, 1, 1) or (1, 1, 1, 1).

- If K = (1, 1, 1, 1), M = 0 and the result follows trivially.
- If K = (2, 1, 1), the basis is  $\{u_{11}, u_{12}, u_{21}, u_{31}\}$ . We can multiply  $u_{11}$  and  $u_{12}$  by a constant so that  $u_{11}^T \Omega_0 u_{12} = 1/r$  for  $r \in Y_p \cup \{1\}$ , and  $u_{31}$  so that  $u_{21}^T \Omega_0 u_{31} = 1$ . Taking as S the matrix with the columns  $\{ru_{12}, u_{11}, u_{21}, u_{31}\}$ , we are in case (1) of Theorem F, with  $c_1 = c_2 = 0$ .
- If K = (2, 2), the basis is  $\{u_{11}, u_{12}, u_{21}, u_{22}\}$ . We multiply  $u_{11}$  and  $u_{12}$  by a constant and  $u_{21}$  and  $u_{22}$  by another constant so that  $u_{i1}^T \Omega_0 u_{i2} = 1/r_i$  for  $r_i \in Y_p \cup \{1\}, i = 1, 2$ . Taking as S the matrix with columns  $\{r_1 u_{12}, u_{11}, r_2 u_{22}, u_{21}\}$ , we are in the same case as before.
- If K = (4), the basis is  $\{u_1, u_2, u_3, u_4\}$ . Let  $k = u_1^T \Omega_0 u_4 = -u_2^T \Omega_0 u_3$ . We can multiply the four vectors by a constant so that  $1/k \in Y_p \cup \{1\}$ . Taking S with the columns  $\{u_3/k, u_2, ku_1, u_4/k^2\}$ , we are in case (4) of this theorem, with c = 1/k.
- (b) Finally we prove uniqueness. If there are two normal forms N and N', by Proposition 3.2, they must have the same Jordan form. The matrices in each case have different Jordan forms, so N and N' are in the same case.
  - If it is case (1), by Theorem D we have N = N', except perhaps for the order of the blocks.

- If it is case (2), the equality of eigenvalues implies N = N'.
- If it is case (3), each c corresponds to a different extension, so the equality of eigenvalues implies c = c' and r = r'. It is left to prove a = a'. Suppose on the contrary that a = 1 and  $a' = h_p(c)$ . Applying the proof of existence, we have that

$$\frac{-4\alpha^2 k}{1-\alpha^2} \in \mathrm{DSq}(\mathbb{Q}_p, -c) \text{ and } \frac{-4\alpha^2 k}{h_p(c)(1-\alpha^2)} \in \mathrm{DSq}(\mathbb{Q}_p, -c)$$

As  $DSq(\mathbb{Q}_p, -c)$  is a group, we also have  $h_p(c) \in DSq(\mathbb{Q}_p, -c)$ , which is a contradiction.

• If it is case (4), again by equality of eigenvalues we have c = c'.

**Proposition 6.11.** Proposition 4.9 also holds for dimension 4, that is, the choice of  $c_0$  only affects the choice of representatives of each class of matrices up to multiplication by a symplectic matrix. The same happens for  $a_0$  and  $b_0$ .

*Proof.* Applying Theorems  $\mathbf{F}$  and  $\mathbf{G}$  to the normal forms of one set gives for each one and only one form of the other set which is equivalent.

6.5. **Proof of Theorem H.** From Theorem F, if  $p \equiv 1 \mod 4$ , case (1) leads to  $\binom{8}{2} = 28$  normal forms (there are seven possible values for  $c_1$  and  $c_2$ ), case (2) to 3 normal forms, and case (3) has 9 possibilities for  $c, t_1$  and  $t_2$ , each one with two possible a and b. Hence, there is a total of 49 normal forms if  $p \equiv 1 \mod 4$ . Analogously, there is a total of  $\binom{6}{2} + 3 + 7 \cdot 2 = 32$  normal forms if  $p \equiv 3 \mod 4$ , and a total of  $\binom{12}{2} + 7 + (9 + 11 + 9 + 9 + 11 + 9 + 11) \cdot 2 = 211$  normal forms if p = 2 (7 possibilities for c, some of them with 9 options for  $t_1$  and  $t_2$  and others with 11, and 2 for a and b).


From Theorem G, case (1) produces  $7 \cdot 4 = 28$  families of normal forms with one degree of freedom if  $p \equiv 1 \mod 4$ ,  $5 \cdot 4 = 20$  if  $p \equiv 3 \mod 4$  and  $11 \cdot 8 = 88$  if p = 2, case (2) produces one such family and case (3) produces 6, 6 and 14 families, respectively. Case (1) produces 16, 16 and 64 isolated forms, and case (4) produces 4, 4 and 8 such forms.

#### 7. MATRIX CLASSIFICATION IN THE REAL CASE, FOR ANY DIMENSION

In this section we give a new proof of the most general case of the Weierstrass-Williamson classification theorem using the new strategy introduced in the previous sections of this paper. In the simplest case, that is, for positive definite symmetric matrices, the proof reduces only to a few lines.

7.1. The general case. In the real case, as already explained after Proposition 5.1, in the case where the eigenvalues of  $\Omega_0^{-1}M$  are different, the blocks up to dimension 4 are enough to classify the matrix. Actually, a weaker condition is sufficient: see Theorem 7.2. If the matrix is not diagonalizable or not invertible, the size of the blocks is not limited to 2 or 4, but instead can grow indefinitely: see Theorem 7.3. See Figure 14 for a hierarchy of properties of the decomposition.

The explicit form of the blocks is as follows:



Blocks may be repeated

FIGURE 14. Hierarchy of degeneracy levels of real matrices, according to the properties of the block decomposition of their normal forms (Theorems 7.2 and 7.3).

Definition 7.1. A diagonal block of hyperbolic type is any matrix of the form

$$M_{\rm h}(k,r,a) = \begin{pmatrix} r & & & & \\ r & 1 & & & \\ & 1 & r & & \\ & r & \ddots & & \\ & & \ddots & & 1 & \\ & & & 1 & r \\ & & & & r & a \end{pmatrix},$$

for some positive integer  $k, r \in \mathbb{R}$  and  $a \in \{-1, 0, 1\}$  with a = 0 if  $r \neq 0$ , and which has a total of 2k rows. A *diagonal block of elliptic type* is any matrix of the form

$$M_{\rm e}(k,r,a) = \begin{pmatrix} M_{\rm e1}(r) & M_{\rm e2}(1,a) & & \\ M_{\rm e2}(1,a) & M_{\rm e1}(r) & M_{\rm e2}(2,a) & & \\ & & M_{\rm e2}(2,a) & \ddots & & \\ & & & & M_{\rm e1}(r) & M_{\rm e2}'(\ell,a) \\ & & & & M_{\rm e2}'(\ell,a)^T & M_{\rm e1}'(r) \end{pmatrix}$$

if  $k = 2\ell + 1$  is odd, and

$$M_{\rm e}(k,r,a) = \begin{pmatrix} M_{\rm e1}(r) & M_{\rm e2}(1,a) \\ M_{\rm e2}(1,a) & M_{\rm e1}(r) & M_{\rm e2}(2,a) \\ & & M_{\rm e2}(2,a) & \ddots \\ & & & & M_{\rm e1}(r) & M_{\rm e2}(\ell-1,a) \\ & & & & M_{\rm e2}(\ell-1,a) & M_{\rm e1}(r) + M_{\rm e2}(\ell,a) \end{pmatrix}$$

if  $k = 2\ell$  is even, for some positive integer  $k, r \in \mathbb{R}$  and  $a \in \{-1, 1\}$ , and which has a total of 2k rows. A *diagonal block of focus-focus type* is any matrix of the form

$$M_{\rm ff}(k,r,s) = \begin{pmatrix} M_{\rm ff1}(r,s) & M_{\rm e2}(1,1) \\ M_{\rm e2}(1,1) & M_{\rm ff1}(r,s) & M_{\rm e2}(1,1) \\ & & M_{\rm e2}(1,1) & \ddots \\ & & & & M_{\rm ff1}(r,s) & M_{\rm e2}(1,1) \\ & & & & M_{\rm e2}(1,1) & M_{\rm ff1}(r,s) \end{pmatrix},$$

for some positive integer k and  $r, s \in \mathbb{R}$ , and which has a total of 4k rows. In the previous blocks the following sub-blocks are used:

$$M_{e1}(r) = \begin{pmatrix} 0 & 0 & 0 & r \\ 0 & 0 & -r & 0 \\ 0 & -r & 0 & 0 \\ r & 0 & 0 & 0 \end{pmatrix}, M_{e1}'(r) = \begin{pmatrix} r & 0 \\ 0 & r \end{pmatrix}, M_{f1}(r,s) = \begin{pmatrix} 0 & s & 0 & r \\ s & 0 & -r & 0 \\ 0 & -r & 0 & s \\ r & 0 & s & 0 \end{pmatrix},$$
$$M_{e2}(j,a) = \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & 0 & a & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \text{ if } j \text{ is odd}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & 0 & a \\ 0 & 0 & 0 & a \end{pmatrix} \text{ if } j \text{ is odd}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & 0 & a \\ 0 & 0 & 0 & a \end{pmatrix} \text{ if } j \text{ is oven},$$
$$M_{e2}'(j,a) = \begin{pmatrix} a & 0 \\ 0 & a \\ 0 & 0 \\ 0 & a \\ 0 & 0 \end{pmatrix} \text{ if } j \text{ is odd}, \begin{pmatrix} 0 & 0 \\ a & 0 \\ 0 & 0 \\ 0 & a \\ 0 & 0 \end{pmatrix} \text{ if } j \text{ is even}.$$

**Theorem 7.2.** Let n be a positive integer. Let  $\Omega_0$  be the matrix of the standard symplectic form in  $\mathbb{R}^{2n}$ . Let  $M \in \mathcal{M}_{2n}(\mathbb{R})$  be a symmetric and invertible matrix such that  $\Omega_0^{-1}M$ is diagonalizable. Then, there exists a symplectic matrix  $S \in \mathcal{M}_{2n}(\mathbb{R})$  such that  $S^T MS$ is a block-diagonal matrix with blocks of hyperbolic, elliptic type or focus-focus type with k = 1.

*Proof.* Applying Lemma 3.6, we can take a symplectic basis  $\{u_1, v_1, \ldots, u_n, v_n\}$  of  $\mathbb{R}^{2n}$  such that all the vectors in the basis are eigenvectors of A, and  $u_i$  and  $v_i$  have opposite eigenvalues  $\lambda_i$  and  $-\lambda_i$ . We can sort these vectors in such a way that two  $\lambda_i$ 's which are conjugate appear with consecutive indices.

Taking as  $\Psi_1$  the matrix with these vectors as columns, the problem decomposes into finding normal forms for each block of columns associated to eigenvalues of the form  $\{r, -r\}$ ,  $\{ir, -ir\}$  or

$$\{r+\mathrm{i}s, -r-\mathrm{i}s, r-\mathrm{i}s, -r+\mathrm{i}s\},\$$

for  $r, s \in \mathbb{R}^*$ . The first block, by Proposition 4.1, gives the hyperbolic block. The second block, by Proposition 4.6 with a = b = r or a = b = -r (one of them will always work), gives the elliptic block. The third block, by Proposition 5.1, gives the focus-focus block.

**Theorem 7.3.** Let n be a positive integer and let  $M \in \mathcal{M}_{2n}(\mathbb{R})$  be a symmetric matrix. Then, there exists a symplectic matrix  $S \in \mathcal{M}_{2n}(\mathbb{R})$  such that  $S^TMS$  is a block diagonal matrix with each of the diagonal blocks being of hyperbolic, elliptic or focus-focus type, as in Definition 7.1.

Furthermore, if there are two matrices S and S' such that  $N = S^T M S$  and  $N' = S'^T M S'$  are normal forms, then N = N' except by the order of the blocks.

**Proof.** (a) First we prove existence. The proof starts as in Theorems 3.16 and 7.2, applying Lemma 3.6. This gives us a partial symplectic basis  $\{u_1, v_1, \ldots, u_m, v_m\}$ . However, unlike Theorem 3.16, these vectors only work for the hyperbolic blocks, giving  $M_h(k, r, 0)$ ; for the rest, the blocks would not be real, so we need to recombine the vectors. We can sort the blocks of the Jordan form in such a way that the blocks of two  $\lambda_i$ 's which are conjugate appear with consecutive indices.

For an elliptic block,  $\{u_1, v_1, \ldots, u_k, v_k\}$  are the vectors corresponding to the values ir and -ir, for  $r \in \mathbb{R}$ . We see that  $M_e(k, r, 1)$  and  $M_e(k, r, -1)$  have this block as Jordan form, so we can apply Proposition 3.2. The columns of  $\Psi_1$  are  $\{u_1, v_1, \ldots, u_k, v_k\}$ , which are part of a symplectic basis, and  $u_1$  and  $v_k$  are eigenvectors with values ir and -ir, so  $v_k = c\bar{u}_1$  for some  $c \in \mathbb{C}$ . Using that  $Au_j = iru_j + u_{j-1}$  and  $Av_j = -irv_j - v_{j+1}$ , we deduce that  $v_{k+1-j} = (-1)^{j-1}c\bar{u}_j$ . Concretely,

$$cu_1^T \Omega_0 \bar{u}_k = u_1^T \Omega_0 v_1$$
  
= 1  
=  $\overline{u_k^T \Omega_0 v_k}$   
=  $\overline{(-1)^{k-1} cu_k^T \Omega_0 \bar{u}_1}$   
=  $(-1)^{k-1} \bar{c} \bar{u}_k^T \Omega_0 u_1$ 

which implies  $c = (-1)^k \bar{c}$ , that is, c is real if k is even and imaginary if k is odd. The columns of  $\Psi_2$  have two nonzero entries, of the form

$$(\ldots, \pm 1, 0, \pm i, \ldots)$$

or

$$(\ldots, \pm a, 0, \pm ia, \ldots),$$

except the two central ones if k is odd, which are  $(\ldots, 1, i)$  or a similar form. In any case, we have  $u_i^T \Omega_0 v_i' = 2a$  if k is even, and 2ia if k is odd.

We can take as D a diagonal matrix with the entries alternating between  $d_1$  and  $d_2$ , that commutes with J. The condition

$$D^T \Omega_0 D = D^T \Psi_1^T \Omega_0 \Psi_1 D = \Psi_2 \Omega_0 \Psi_2$$

implies that  $d_1d_2 = 2a$  for k even, and 2ia for k odd. As S must be a real matrix, in  $S\Psi_2 = \Psi_1 D$  the first and last columns are conjugate and we get  $\bar{d}_1 = cd_2$ , that is,  $d_1\bar{d}_1 = 2ac$  for k even and 2iac for k odd. We just need to take  $a \in \{1, -1\}$  so that this is positive: note that a is unique.

For the focus-focus case, we have in the symplectic basis a block of vectors

$$\{u_1, v_1, \ldots, u_k, v_k, u'_1, v'_1, \ldots, u'_k, v'_k\},\$$

where  $u_1$ ,  $v_k$ ,  $u'_1$  and  $v'_k$  are eigenvectors for  $\lambda, -\lambda, \bar{\lambda}$  and  $-\bar{\lambda}$  respectively. If  $\lambda = s + ir$ , the matrix  $M_f(k, r, s)$  has the same Jordan form.

The columns of  $\Psi_1$  are the vectors  $u_i, v_i, u'_i$  and  $v'_i$ ; we have that  $u'_1 = c\bar{u}_1$  for some  $c \in \mathbb{C}$ , which implies  $u'_i = c\bar{u}_i$  for all *i* and, from  $u_i^T \Omega_0 v_i = u'_i^T \Omega_0 v'_i = 1$ , we deduce  $v'_i = \bar{v}_i/c$  for all *i*. The columns of  $\Psi_2$  have now the form (0, 1, 0, i, ...),  $(i, 0, 1, 0, \ldots), (\ldots, 0, 1, 0, i, \ldots)$ , and so on, for a total of 2k, followed by their conjugates.

We can take as D a diagonal matrix with the values

$$d_1, d_2, \ldots, d_1, d_2, d_3, d_4, \ldots, d_3, d_4,$$

which commutes with J. The condition

$$D^T \Omega_0 D = D^T \Psi_1^T \Omega_0 \Psi_1 D = \Psi_2 \Omega_0 \Psi_2$$

implies that  $d_1d_2 = -2i$  and  $d_3d_4 = 2i$ . Using that S is a real matrix, the left and right halves of  $S\Psi_2 = \Psi_1 D$  are conjugate, which implies  $d_1 = cd_3$  and  $cd_2 = d_4$ , so the condition  $d_3d_4 = 2i$  reduces to a consequence of  $d_1d_2 = -2i$ , and we can take for example  $d_1 = 1$  and  $d_2 = -2i$ .

This finishes the treatment of the nonzero eigenspaces. For the other part, we can use the same treatment as in Theorem 3.16, but in the even case we cannot always make  $c_i = 1$ ; instead we make  $c_i = a_i$ . Now

$$\left\{u_{i1}, a_i u_{i,2\ell_i}, -u_{i2}, a_i u_{i,2\ell_i-1}, \dots, (-1)^{\ell_i-1} u_{i\ell_i}, a_i u_{i,\ell_i+1}\right\}$$

is a partial symplectic basis which gives the form  $M_{\rm h}(\ell_i, 0, (-1)^{\ell_i}a_i)$ .

(b) Finally we prove uniqueness. If N and N' are two normal forms which are equivalent, by Proposition 3.2, they have the same Jordan form. This means that the set of blocks is the same except for the values of  $a_i$ . But in the elliptic case we already saw that a is unique. This only leaves the hyperbolic case with r = 0.

In this case, if a can be 1 and -1 at the same time, there is a chain

$$\left\{u_1, u_2, \dots, u_{2k}\right\}$$
such that  $Au_i = u_{i-1}$ ,  $Au_1 = 0$ , and  $u_i \Omega_0 u_{2k+1-i} = (-1)^i$ , and another one

$$\left\{u_1', u_2', \dots, u_{2k}'\right\}$$

with the same properties except that

$$u_i'\Omega_0 u_{2k+1-i}' = (-1)^{i+1}.$$

In the space generated by these vectors there is only one vector in the kernel, so  $u'_1 = ku_1$  for some  $k \in \mathbb{R}$ . As  $Au_i = u_{i-1}$  and  $Au'_i = u'_{i-1}$ , we have that  $u'_i = ku_i$ for all i. This together implies that

$$(-1)^{i+1} = u'_i \Omega_0 u'_{2k+1-i} = k^2 u_i \Omega_0 u_{2k+1-i} = k^2 (-1)^i$$
  
= -1, a contradiction.

and  $k^2 =$ 

The matrix  $\Psi_1$  in the previous proof gives a complex symplectic basis in which M has the block diagonal form of Theorem 3.16. The relation between this and the final matrix S can be written in terms of vectors. In the hyperbolic case, they are the same matrix. In the elliptic case, we first multiply the vectors by the corresponding  $d_i$ :  $u_i := \sqrt{|2c|u_i|}$  $v_i := \sqrt{2/|c|} v_i$ . Then, the matrix  $\Psi_2$  indicates how the vectors in the final basis relate to these  $u_j$  and  $v_j$ : each column has a  $\pm 1$  entry, a  $\pm i$  entry and the rest are 0, so we have  $\pm u'_h + \pm i u'_\ell = u_i$ , for some indices h and  $\ell$ . As the new vectors  $u'_h$  and  $u'_\ell$  must be real, this has a unique solution, and the new vectors are the real and imaginary parts of the old ones (maybe with the sign changed).

In the focus-focus case, we also start multiplying the vectors by the  $d_i$ :

$$v_j := -2iv_j, u'_j := u'_j/c = \bar{u}_j, v'_j := 2icv'_j = \bar{v}_j.$$

Then we apply  $\Psi_2^{-1}$ : now each column of the left half of  $\Psi_2$  has a 1 and an i, and the right half has a 1 and a -i in the same positions. For every j with  $1 \leq j \leq k$ , the equations are  $v_{2j-1}'' + iv_{2j}'' = u_j$  and  $iu_{2j-1}'' + u_{2j}'' = v_j$  for the first half and their conjugates for the second half, where  $u_j''$  and  $v_j''$  are the new vectors. The solution consists of taking the real and imaginary parts of the new vectors.

**Corollary 7.4.** Let  $M \in \mathcal{M}_4(\mathbb{R})$  be a symmetric matrix. Then there exists  $r, s \in \mathbb{R}$ ,  $a, b \in \{-1, 1\}$ , and a symplectic matrix  $S \in \mathcal{M}_4(\mathbb{R})$  such that  $S^T M S$  is one of the following ten matrices:

$$\begin{pmatrix} 0 & r & 0 & 0 \\ r & 0 & 0 & 0 \\ 0 & 0 & 0 & s \\ 0 & 0 & s & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & 0 & s \\ 0 & 0 & s & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & s \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & r \\ 0 & 0 & r & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & a \end{pmatrix}, \begin{pmatrix} 0 & r & 0 & 0 \\ r & 0 & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & s \end{pmatrix}, \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & r & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & s \end{pmatrix}, \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & r & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & s \end{pmatrix}, \begin{pmatrix} a & 0 & 0 & r \\ 0 & 0 & -r & 0 \\ 0 & -r & a & 0 \\ r & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & s & 0 & r \\ s & 0 & -r & 0 \\ 0 & -r & 0 & s \\ r & 0 & s & 0 \end{pmatrix},$$

Furthermore, if there are two matrices of this form equivalent to M, they are both in the first or in the eighth case swapping r and s, or in the third case swapping a and b.

*Proof.* They are in this order: two hyperbolic blocks with k = 1 and a = 0, one with a = 0 and one with  $a \neq 0$ , two with  $a \neq 0$ , one hyperbolic block with k = 2 and a = 0, the same with  $a \neq 0$ , one hyperbolic with a = 0 and one elliptic, the same with  $a \neq 0$ , two elliptic blocks with k = 1, one elliptic block with k = 2, and one focus-focus block.  $\Box$ 

These are the same (up to symplectic transformations) that Williamson gives in his paper [66, page 24] and which we gave in Section 2.

7.2. Example of application of our method for matrices of arbitrary order. Consider the matrix

$$M = \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & 1 & & & \\ & & 1 & & \\ & & 1 & \ddots & \\ & & & \ddots & & 1 \\ & & & & 1 & \\ & & & & & 1 \end{pmatrix}$$

We have that

$$\Omega_0^{-1}M = \begin{pmatrix} & -1 & & & \\ 1 & & & & \\ & & -1 & & \\ 1 & & & \ddots & \\ & & & -1 & \\ & & 1 & & \\ & & & \ddots & & -1 \\ & & & & 1 & \end{pmatrix}$$

whose characteristic polynomial is  $\lambda^{2n} + (-1)^n$ , and the eigenvalues are  $\lambda = e^{\pi i k/n}$  for  $0 \leq k \leq 2n-1$ , if n is odd, and  $\lambda = e^{\pi i (2k+1)/2n}$  for  $0 \leq k \leq 2n-1$ , if n is even.

If n = 2m + 1 is odd, the normal form of this matrix contains a hyperbolic block, which corresponds to the eigenvalues  $\{1, -1\}$ , and *m* focus-focus blocks, for the eigenvalues

$$\left\{\mathrm{e}^{\pi\mathrm{i}k/n}, \mathrm{e}^{\pi\mathrm{i}(n-k)/n}, \mathrm{e}^{\pi\mathrm{i}(n+k)/n}, \mathrm{e}^{\pi\mathrm{i}(2n-k)/n}\right\},\,$$

for  $1 \leq k \leq m$ :

$$S^{T}MS = \begin{pmatrix} 0 & 1 & & \\ 1 & 0 & & \\ & & N(1) & \\ & & \ddots & \\ & & & N(m) \end{pmatrix}, N(k) = \begin{pmatrix} 0 & \cos\frac{\pi k}{n} & 0 & \sin\frac{\pi k}{n} \\ \cos\frac{\pi k}{n} & 0 & -\sin\frac{\pi k}{n} & 0 \\ 0 & -\sin\frac{\pi k}{n} & 0 & \cos\frac{\pi k}{n} \\ \sin\frac{\pi k}{n} & 0 & \cos\frac{\pi k}{n} & 0 \end{pmatrix}.$$

On the other hand, if n = 2m is even, the normal form contains only m focus-focus blocks N(k + 1/2), for  $0 \le k \le m - 1$ . In this case, if we change any 1 entry in M to -1, the normal form contains again the hyperbolic block, the focus-focus blocks N(k) for  $1 \le k \le m - 1$  and also an elliptic block, corresponding to the eigenvalues i and -i:

$$S^T M S = \begin{pmatrix} 0 & 1 & & & \\ 1 & 0 & & & \\ & 1 & 0 & & \\ & 0 & 1 & & \\ & & N(1) & & \\ & & & \ddots & \\ & & & & N(m-1) \end{pmatrix}$$

7.3. **Positive-definite case.** If the matrix is positive-definite, only the elliptic case may appear: this happens because, if u is an eigenvector of A with value  $\lambda$ ,

$$\bar{u}^T M u = \bar{u}^T \Omega_0 A u = \lambda \bar{u}^T \Omega_0 u$$

The left-hand side is a positive real, and for  $\bar{u}^T \Omega_0 u$  we have

$$\overline{\bar{u}^T \Omega_0 u} = u^T \Omega_0 \bar{u} = -\bar{u}^T \Omega_0 u,$$

so it is imaginary, and  $\lambda$  must be imaginary.

Also, it is impossible to obtain for these matrices the elliptic blocks with size greater than 2: if such a block appeared, there would be an element in the diagonal of  $S^T M S$  equal to 0. Let k be its index and u the k-th column of S. Then  $u^T M u = 0$ , which contradicts M being positive definite. This means that any positive definite symmetric matrix can be diagonalized by a symplectic matrix, which is the result most often referred to as "Williamson theorem".

# 8. Comments on the *p*-adic classification in higher dimensions and proof of Theorem I

Our strategy for the 4-dimensional case extends to dimension 6 and 8, using the fact that in these dimensions all polynomials are solvable by radicals (though it would be needed to take higher order radicals) but for brevity we do not deal with those cases (we expect hundreds or even thousands of possibilities for the model matrices, see Table 3). In dimension 10 or higher one cannot expect an explicit classification because polynomials of degree 5 or higher are not in general solvable by radicals.

8.1. Preparatory lemmas. In order to prove Theorem I, we need some lemmas.

**Lemma 8.1.** Let p be a prime number and let n be a positive integer. The polynomial

 $P(x) = x^n - ap,$ 

where  $\operatorname{ord}_p(a) = 0$ , is irreducible in  $\mathbb{Q}_p$ .

*Proof.* The roots of P have order 1/n. If P was reducible, a factor should have a subset of the roots whose product has integer order, but this would need all the n roots.

**Lemma 8.2.** Let p be a prime number and let n be a positive integer. Let  $a, b \in \mathbb{Q}_p$  such that  $\operatorname{ord}_p(a) = 0$  and  $\operatorname{ord}_p(b) = 1$ . If a is an n-th power in  $\mathbb{Q}_p[b^{1/n}]$ , then it is an n-th power in  $\mathbb{F}_p$ .

*Proof.* Suppose that  $a = c^n$  for  $c \in \mathbb{Q}_p[b^{1/n}]$ . We can write

$$c = c_0 + c_1 b^{\frac{1}{n}} + \ldots + c_{n-1} b^{\frac{n-1}{n}}$$

where  $c_i \in \mathbb{Q}_p$  for all *i*. Raising this to the *n*-th power, we have  $c^n = a$  at the left, and  $c_0^n$  plus terms of positive order at the right. Then,  $a - c_0^n$  has positive order, and as it is in  $\mathbb{Q}_p$  the order must be at least 1, and  $a \equiv c_0^n \mod p$ , as we wanted.

**Lemma 8.3.** Let p be a prime number and let n be a positive integer. There are at least gcd(2n, p-1) + gcd(n, p-1), if n is odd, and gcd(2n, p-1), if n is even, infinite families of blocks of size 2n in the normal form of a matrix up to multiplication by a symplectic matrix, where each family is of the form  $r_1M_1 + \ldots + r_nM_n$ .

Proof. Consider the polynomial  $P(x) = x^{2n} - ap$  where  $\operatorname{ord}_p(a) = 0$ . This is irreducible by Lemma 8.1, so it will give a block of size 2n in the normal form. This block may not be unique up to multiplication by a symplectic matrix (as happens in Propositions 4.6 and 5.2), but, in analogy with the proofs of those results, two blocks corresponding to different a will be in the same family only if the roots of the polynomials are in the same extension of  $\mathbb{Q}_p$ . Suppose that this happens for  $a_1$  and  $a_2$ . In particular,  $(a_1p)^{1/2n}$  and  $(a_2p)^{1/2n}$  are in the same extension, that is,

$$\left(\frac{a_2}{a_1}\right)^{\frac{1}{2n}} \in \mathbb{Q}_p[(a_1p)^{\frac{1}{2n}}]$$

By Lemma 8.2,  $a_2/a_1$  must be a 2*n*-th power in  $\mathbb{F}_p$ . This implies that the number of families of blocks is at least the cardinality of  $\mathbb{F}_p^*$  modulo 2*n*-th powers, which is gcd(2n, p-1), because that group is cyclic of order p-1.

If n is odd, we also consider

$$Q(x) = x^{2n} - a^2 p^2 = (x^n + ap)(x^n - ap).$$
<sub>67</sub>

The two factors are again irreducible and it also gives a block of size 2n (one factor comes from changing the sign of x in the other). Two blocks for  $a_1$  and  $a_2$  are in the same family only if  $(a_1p)^{1/n}$  and  $(a_2p)^{1/n}$  are in the same extension, that is,

$$\left(\frac{a_2}{a_1}\right)^{\frac{1}{n}} \in \mathbb{Q}_p[(a_1p)^{\frac{1}{n}}]$$

(note that choosing -ap instead of ap gives the same extension because  $(-1)^{1/n} = -1$ ). Again by Lemma 8.2,  $a_2/a_1$  is an *n*-th power in  $\mathbb{F}_p$ . So the number of families is now the cardinality of  $\mathbb{F}_p^*$  modulo *n*-th powers, which is gcd(n, p-1).

**Remark 8.4.** Concerning the number of families of normal forms (instead of just blocks), in the real case, supposing that there are k focus-focus blocks, there are 2n - 4k variables left, which can be distributed between hyperbolic and elliptic blocks in n - 2k + 1 ways. The total number of forms is

$$\sum_{k=0}^{m} 2m - 2k + 1 = 2m^2 - m(m+1) + m + 1 = m^2 + 1$$
  
if  $n = 2m$  and  
$$\sum_{k=0}^{m} 2m + 1 - 2k + 1 = m(2m+1) - m(m+1) + m + 1 = m^2 + m + 1$$

if n = 2m + 1.

8.2. **Proof of Theorem I.** Lemma 8.3 tells us that there is at least one block with each even size. Hence, the number of normal forms is at least the number of partitions of n in positive integers, that grows with  $e^{\pi\sqrt{2n/3}}/4n\sqrt{3}$  by the Hardy-Ramanujan formula [30]. In order to find the exact formulas of the matrices, we need to devise, for each partition, a matrix in  $\mathcal{M}_{2n}(\mathbb{Q}_p)$  with the product of the corresponding factors as characteristic polynomial. This can be done with the same strategy as in Section 7.2, and gives the matrix  $\mathcal{M}(P, p)$  for each partition P.

8.3. Remarks and applications. Theorem I could be strengthened by using that there is not only *one* block of each size, but this would imply making a sum over the partitions. We do not know how to make that for general n, but we can do it for small n, obtaining the results in Table 3.

From the point of view of symplectic geometry and topology of integrable systems, which is the main motivation of the authors to write this paper, currently the only known global *symplectic* classifications of integrable systems which include physically intriguing local models (that is, essentially non-elliptic models) concern dimension 4 [43, 47, 48] in the real case. These real classifications include for example the coupled angular momentum [37] and the Jaynes-Cummings model [50]. Hence, in the *p*-adic case, with hundreds of local models (Theorem B), we expect that the 4-dimensional case is already extremely complicated and that the 2*n*-dimensional case,  $n \ge 3$ , is out of reach (since it is out of reach in the real case with only a very small proportion of local models in comparison, see Theorem C).

In dimension 4 the authors analyzed one of these systems, the *p*-adic Jaynes-Cummings model [11], whose treatment is very extensive compared to its real counterpart, as expected. Although as we said, a classification of *p*-adic integrable systems in dimension 4, extending [43, 47, 48], seems out of reach, the present paper settles completely the first step: understanding explicitly *p*-adic local models. The proofs of [43, 47, 48] are based on gluing local models.

#### 9. Application to normal forms of *p*-adic singularities

The Weierstrass-Williamson's classification of matrices can be used to classify critical points of *p*-adic analytic functions.

**Definition 9.1.** Let n be a positive integer.

- Given a field F, a symplectic vector space over F is a pair  $(V, \omega)$  where V is a 2ndimensional vector space over F and  $\omega : V \times V \to F$  is a non-degenerate bilinear map such that  $\omega(v, v) = 0$  for all  $v \in V$ . We say that  $\omega$  is a linear symplectic form. If the characteristic of F is not 2, the last condition is equivalent to  $\omega$  being antisymmetric:  $\omega(v, w) = -\omega(w, v)$  for all  $v, w \in V$ .
- Let p be a prime number. A p-adic analytic symplectic manifold is a pair  $(M, \omega)$  where M is a p-adic analytic manifold and  $\omega$  is a closed non-degenerate 2-form on M. We say that  $\omega$  is a symplectic form. At each point,  $\omega$  gives a linear symplectic form.
- Given a field F, a linear symplectomorphism between two symplectic vector spaces  $(V_1, \omega_1)$  and  $(V_2, \omega_2)$  over F is a linear isomorphism  $\phi : V_1 \to V_2$  that preserves the symplectic form, that is,  $\phi^*\omega_2 = \omega_1$ . In this case we say that  $\omega_1$  and  $\omega_2$  are linearly symplectomorphic.
- Let  $(M_1, \omega_1)$  and  $(M_2, \omega_2)$  be *p*-adic analytic symplectic manifolds. Let  $m \in M$ . A local linear symplectomorphism  $\phi : U_1 \to U_2$  centered at *m* is a *p*-adic analytic diffeomorphism between some open sets  $U_1 \subset M_1$  and  $U_2 \subset M_2$ , such that  $m \in U_1$ , and which yields a linear symplectomorphism  $T_m \phi : T_m M_1 \to T_{\phi(m)} M_2$ .
- Let  $(M, \omega)$  be a *p*-adic analytic symplectic manifold. By *linear symplectic coordi*nates  $(x_1, \xi_1, ..., x_n, \xi_n)$  with the origin a point  $m \in M$  we mean coordinates given by a local linear symplectomorphism centered at m, that is,  $\phi^* \omega_m = \omega_0$ . In terms of matrices this last condition can be formulated as  $S^T \Omega S = \Omega_0$ , where S,  $\Omega$  and  $\Omega_0$  are the matrices of  $\phi$ ,  $\omega_m$  and  $\omega_0$ .

**Remark 9.2.** By definition, an automorphism of a symplectic space is a linear symplectomorphism if and only if its matrix is a symplectic matrix.

**Theorem 9.3.** Let F be a field and n be a positive integer. Every symplectic form on  $F^{2n}$  is linearly symplectomorphic to the form  $\omega_0$  which has as matrix

Hence, every two linear symplectic forms on  $F^{2n}$  are actually symplectomorphic.

*Proof.* What we want to prove is that there is a basis with respect to which  $\omega$  has the matrix  $\Omega_0$ . Then, the symplectomorphism that sends this basis to the canonical one will send  $\omega$  to  $\omega_0$ .

We go by induction on n. As  $\omega$  is non-degenerate, there are  $u_1$  and  $v_1$  with  $\omega(u_1, v_1) = 1$ . Of course,  $\langle u_1, v_1 \rangle$  is symplectic, so its complement  $\langle u_1, v_1 \rangle^{\omega}$  is also symplectic. Applying induction to this complement, we get a basis  $\{u_2, v_2, \ldots, u_n, v_n\}$ . Now,  $\{u_1, v_1, \ldots, u_n, v_n\}$  is the basis we are looking for. We refer to appendix A.2 for the definition of analytic function and critical point of a function on a *p*-adic manifold. It does not make sense to talk about the rank of such a critical point, because there is only one function and consequently only one differential form.

**Definition 9.4** (Non-degenerate critical point of *p*-adic analytic function, symplectic sense). Let *n* be a positive integer. Let *p* be a prime number. Let  $(M, \omega)$  be a 2*n*-dimensional *p*-adic analytic symplectic manifold. Let  $f: M \to \mathbb{Q}_p$  be a *p*-adic analytic function and let  $m \in M$  be a critical point of f (i.e. df(m) = 0). Let  $\Omega$  be the matrix of  $\omega$ . We say that *m* is *non-degenerate* if the eigenvalues of  $\Omega^{-1}d^2f(m)$  are all distinct.

This is not the usual notion of non-degenerate critical point (which states that the Hessian of f is invertible), but the two are related as the following proposition shows.

**Proposition 9.5.** Let n be a positive integer. Let p be a prime number. Let M be a p-adic analytic 2n-dimensional manifold,  $f: M \to \mathbb{Q}_p$  a p-adic analytic function, and m a critical point of f. Then the following are equivalent:

- (1) m is a non-degenerate critical point of f in the usual sense;
- (2) There exists a linear symplectic form  $\omega$  such that m is a non-degenerate critical point of  $f: (M, \omega) \to \mathbb{Q}_p$  in the symplectic sense (Definition 9.4).
- (3) There exist infinitely many linear symplectic forms  $\omega$  such that m is a nondegenerate critical point of  $f: (M, \omega) \to \mathbb{Q}_p$  in the symplectic sense.

Proof. Suppose (2) holds. Let  $\Omega$  be the matrix of  $\omega$ . Then  $\Omega^{-1}d^2f(m)$  has all eigenvalues distinct. Applying Lemma 3.3 to the Hessian, if zero was an eigenvalue, it would be at least double, contradicting (2). So  $\Omega^{-1}d^2f(m)$  is invertible, which implies  $d^2f(m)$  is invertible and (1) holds.

Now suppose (1) holds. Let  $H = d^2 f(m)$ . We first solve the problem for H diagonal and then the general case.

If H is diagonal, (1) means that all diagonal elements are nonzero. Let  $h_i$  be the *i*-th diagonal element. We will take  $\Omega^{-1}$  with  $a_i$  in the row 2i - 1 and column 2i,  $-a_i$  in the row 2i and column 2i - 1, and 0 the rest. After multiplying  $\Omega^{-1}$  by H,  $a_i$  becomes  $a_i h_{2i-1}$  and  $-a_i$  becomes  $-a_i h_{2i}$ . The eigenvalues are

$$\pm a_i \sqrt{-h_{2i-1}h_{2i}}$$
 for  $1 \leq i \leq n$ .

Since  $\mathbb{Q}_p$  is infinite, it is always possible to choose  $a_i$  so that these values are all different, independently of whether they are in  $\mathbb{Q}_p$  or not (just choose each  $a_i$  in turn, and there will always be a possible value).

In the general case, we can apply Gram-Schmidt orthogonalization to the canonical basis to obtain a basis  $\{v_1, \ldots, v_n\}$  such that  $v_i^T H v_j \neq 0$  if and only if i = j. Taking these vectors as columns, we have a matrix M such that  $M^T H M$  is diagonal. Applying the diagonal case to this matrix, we obtain an antisymmetric  $\Omega$  such that  $\Omega^{-1}M^T H M$  has all eigenvalues different. This matrix is similar to  $M\Omega^{-1}M^T H$ , so this has also all eigenvalues different, and  $M\Omega^{-1}M^T$  is the matrix we want.

That (3) implies (2) is trivial. If (2) holds,  $\Omega^{-1}M$  has all eigenvalues different and the same happens for any perturbation of  $\Omega$ , hence (3) holds.

In the following when we speak of non-degenerate critical points of a function we always de it in the symplectic sense of Definition 9.4.

In the real case, as a consequence of the Weierstrass-Williamson classification, it is always possible to choose linear symplectic coordinates  $(x_1, \xi_1, \ldots, x_n, \xi_n)$  with the origin at any  $m \in M$  such that

$$f = \sum_{i=1}^{n} r_i g_i + \mathcal{O}(3)$$

for some  $r_i \in \mathbb{R}$ , where  $g_i : V \to \mathbb{R}$  has one of the following forms:  $(x_i^2 + \xi_i^2)/2$  (elliptic component),  $x_i\xi_i$  (hyperbolic component), or  $x_i\xi_{i+1} - x_{i+1}\xi_i$  with the next function equal to  $x_i\xi_i + x_{i+1}\xi_{i+1}$  (focus-focus component). In the p-adic case we are also able to make this conclusion.

**Lemma 9.6.** Let n be a positive integer. Let p be a prime number. Let M be a p-adic analytic manifold of dimension 2n and let m be a non-degenerate critical point of a padic analytic function  $f: M \to \mathbb{Q}_p$ . Then there exists an open neighborhood U of m and coordinates  $(x_1, \ldots, x_n)$  with the origin at m such that the restriction of f to U, that is,  $f|_U: U \to \mathbb{Q}_p$  is given by a power series

$$\sum_{\in\mathbb{N}^n, i_1+\ldots+i_n\geqslant 2} a_I x_1^{i_1}\ldots x_n^{i_n}.$$

T

Moreover, the matrix in  $\mathcal{M}_{2n}(\mathbb{Q}_p)$  with the coefficient of  $x_i x_j$  in the row *i* and column *j*, for  $i \neq j$ , and the coefficient of  $x_i^2$  multiplied by 2 in the row and column *i*, is exactly the Hessian of *f* at *m* in the coordinates  $x_1, \ldots, x_n$ .

Proof. By definition, f is given by a power series converging in some open set U which contains m. The center of the power series can be arbitrarily chosen in U, which means that we can choose m as center. The degree 1 terms are 0 because m is a critical point, and the degree 2 terms are of the form  $x^T H x/2$  for some matrix  $H \in \mathcal{M}_{2n}(\mathbb{Q}_p)$ . Differentiating twice, we get that the Hessian of f is precisely H.

**Corollary 9.7** (Normal form of critical points in dimension 2). Let p be a prime number. Let  $(M, \omega)$  be a p-adic symplectic manifold of dimension 2 and let  $f : M \to \mathbb{Q}_p$  be a p-adic analytic function. Let  $m \in M$  be a critical point of f. Then there are linear symplectic coordinates  $(x, \xi)$  with the origin at m such that f - f(m) coincides with  $r(x^2 + c\xi^2)$  up to order 2, for some  $r \in \mathbb{Q}_p$  and  $c \in X_p$ , or  $r \in Y_p \cup \{1\}$  and c = 0. Furthermore, if f - f(m) has this form for two different linear symplectic coordinates with the origin at m, then the two forms coincide.

*Proof.* By Theorem 9.3, we can assume without loss of generality that  $\omega_m = \omega_0$ .

Applying Theorem D to  $d^2 f$ , we get a symplectic matrix S, which is the matrix of a linear symplectomorphism  $\phi$ , such that

$$d^{2}(\phi^{*}f) = S^{T}d^{2}fS = \begin{pmatrix} r & 0\\ 0 & cr \end{pmatrix} = d^{2}(rx^{2} + cr\xi^{2}).$$

By Lemma 9.6, f has the desired form. Uniqueness follows from the same theorem.  $\Box$ 

**Corollary 9.8.** Let p be a prime number. Let  $(M, \omega)$  be a p-adic analytic symplectic 2-manifold. Let  $X_p, Y_p$  be the non-residue sets in Definition 1.1. Then the following statements hold.

(1) If  $p \equiv 1 \mod 4$ , there are exactly 7 families of local linear normal forms for a nondegenerate critical point of a p-adic analytic analytic function  $f: (M, \omega) \to \mathbb{Q}_p$  up to local linear symplectomorphisms centered at the critical point, where the normal forms in each family differ by multiplication by a constant r, and exactly 4 normal forms for a degenerate critical point which only differ in the constant r.

- (2) If  $p \equiv 3 \mod 4$ , there are exactly 5 families of local linear normal forms for a nondegenerate critical point of a p-adic analytic analytic function  $f : (M, \omega) \to \mathbb{Q}_p$  up to local linear symplectomorphisms centered at the critical point, where the normal forms in each family differ by multiplication by a constant r, and exactly 4 normal forms for a degenerate critical point which only differ in the constant r.
- (3) If p = 2, there are exactly 11 families of local linear normal forms for a nondegenerate critical point of a p-adic analytic analytic function f : (M,ω) → Q<sub>p</sub> up to local linear symplectomorphisms centered at the critical point, where the normal forms in each family differ by multiplication by a constant r, and exactly 8 normal forms for a degenerate critical point which only differ in the constant r.

In the three cases the above normal forms for a non-degenerate point are given by

$$\left\{\left\{r(x^2+c\xi^2):r\in\mathbb{Q}_p\right\}:c\in X_p\right\}$$

and those for a degenerate point are given by

$$\Big\{ rx^2 : r \in Y_p \cup \{1\} \Big\}.$$

*Proof.* This follows from Theorem E and Corollary 9.7.

**Corollary 9.9** (Normal form of non-degenerate critical points in dimension 4). Let p be a prime number. Let  $X_p, Y_p, C_i^k, \mathcal{D}_i^k$  be the non-residue sets and coefficient functions in Definition 1.1. Let  $(M, \omega)$  be a p-adic symplectic manifold of dimension 4 and let  $f: M \to \mathbb{Q}_p$  be a p-adic analytic function. Let  $m \in M$  be a non-degenerate critical point of f. Then there are linear symplectic coordinates  $(x, \xi, y, \eta)$  with the origin at m such that in these cordinates we have:

$$f - f(m) = rg_1 + sg_2 + \mathcal{O}(3),$$

where  $g_1$  and  $g_2$  have one of the following forms:

- (1)  $g_1(x,\xi,y,\eta) = x^2 + c_1\xi^2$  and  $g_2(x,\xi,y,\eta) = y^2 + c_2\eta^2$ , for  $c_1, c_2 \in X_p$ .
- (2)  $g_1(x,\xi,y,\eta) = x\eta + cy\xi$  and  $g_2(x,\xi,y,\eta) = x\xi + y\eta$ , for  $c \in Y_p$ .
- (3)

$$g_k(x,\xi,y,\eta) = \sum_{i=0}^2 \mathcal{C}_i^k(c,t_1,t_2,a,b) x^i y^{2-i} + \sum_{i=0}^2 \mathcal{D}_i^k(c,t_1,t_2,a,b) \xi^i \eta^{2-i},$$

for  $k \in \{1, 2\}$ , where  $c, t_1$  and  $t_2$  correspond to one row of Table 1 and (a, b) are either (1, 0) or  $(a_1, b_1)$  of the corresponding row.

If f - f(m) has this form for two different linear symplectic coordinates, then the two forms coincide, except perhaps for swapping  $g_1$  and  $g_2$  at point (1).

*Proof.* Analogously to proof of Corollary 9.7, we apply Theorem F to the Hessian of f, and by Lemma 9.6, f has the desired form. The three cases (1), (2) and (3) for the resulting matrix correspond to the three cases of this corollary, because

$$d^{2}(\phi^{*}f) = S^{T}d^{2}fS = rd^{2}g_{1} + sd^{2}g_{2} = d^{2}(rg_{1} + sg_{2}).$$

**Corollary 9.10** (Normal form of degenerate critical points in dimension 4). Let p be a prime number. let  $X_p, Y_p$  be the non-residue sets in Definition 1.1. Let  $h_p: Y_p \to \mathbb{Q}_p$  be the non-residue function in Definition 2.3. Let  $(M, \omega)$  be a p-adic symplectic manifold of dimension 4 and let  $f: M \to \mathbb{Q}_p$  be a p-adic analytic function. Let  $m \in M$  be a degenerate critical point of f. Then there are linear symplectic coordinates  $(x, \xi, y, \eta)$  with the origin at m such that f - f(m) coincides with one of the following forms up to order 2:

- (1)  $r(x^2+c_1\xi^2)/2+s(y^2+c_2\eta^2)/2$ , for some  $c_1, c_2 \in X_p \cup \{0\}$  and  $r, s \in \mathbb{Q}_p$ . If  $c_1 = 0$ , r can be taken in  $Y_p \cup \{1\}$ , and if  $c_2 = 0$ , s can be taken in  $Y_p \cup \{1\}$ .
- (2)  $r(x\xi + y\eta) + y\xi$ , for some  $r \in \mathbb{Q}_p$ .
- (3)  $r(x\eta + cy\xi) + a(x^2 + y^2)/2$ , for some  $r \in \mathbb{Q}_p, c \in Y_p, a \in \{1, h_p(c)\}.$
- (4)  $c(x^2/2 + \xi\eta)$ , for some  $c \in Y_p \cup \{1\}$ .

Furthermore, if f - f(m) has this form for two different linear symplectic coordinates, then the two forms coincide, except perhaps for swapping  $(r, c_1)$  and  $(s, c_2)$  at point (1).

*Proof.* It is analogous the previous two proofs, but with Theorem G.

**Corollary 9.11.** Let p be a prime number. Let  $(M, \omega)$  be a p-adic analytic symplectic 4manifold. Let  $X_p, Y_p, \mathcal{C}_i^k, \mathcal{D}_i^k$  be the non-residue sets and coefficient functions in Definition 1.1. Let  $h_p: Y_p \to \mathbb{Q}_p$  be the non-residue function in Definition 2.3. Then the following statements hold.

- (1) If  $p \equiv 1 \mod 4$ , there are exactly 49 infinite families of local linear normal forms with two degrees of freedom for a critical point of a p-adic analytic function on a 4-dimensional p-adic symplectic manifold  $f : (M, \omega) \to \mathbb{Q}_p$  up to local linear symplectomorphisms centered at the critical point, exactly 35 infinite families with one degree of freedom, and exactly 20 isolated normal forms.
- (2) If  $p \equiv 3 \mod 4$ , there are exactly 32 infinite families of local linear normal forms with two degrees of freedom for a critical point of a p-adic analytic function on a 4-dimensional p-adic symplectic manifold  $f : (M, \omega) \to \mathbb{Q}_p$  up to local linear symplectomorphisms centered at the critical point, exactly 27 infinite families with one degree of freedom, and exactly 20 isolated normal forms.
- (3) If p = 2, there are exactly 211 infinite families of local linear normal forms with two degrees of freedom for a critical point of a p-adic analytic function on a 4dimensional p-adic symplectic manifold f : (M,ω) → Q<sub>p</sub> up to local linear symplectomorphisms centered at the critical point, exactly 103 infinite families with one degree of freedom, and exactly 72 isolated normal forms.
- In the three cases above, the infinite families with two degrees of freedom are given as

$$\begin{cases} \left\{ r(x^{2}+c_{1}\xi^{2})+s(y^{2}+c_{2}\eta^{2}):r,s\in\mathbb{Q}_{p}\right\}:c_{1},c_{2}\in X_{p} \\ \\ \cup\left\{ \left\{ r(x\eta+cy\xi)+s(x\xi+y\eta):r,s\in\mathbb{Q}_{p}\right\}:c\in Y_{p} \right\} \\ \\ \cup\left\{ \left\{ r\left(\sum_{i=0}^{2}\mathcal{C}_{i}^{1}(c,t_{1},t_{2},a,b)x^{i}y^{2-i}+\sum_{i=0}^{2}\mathcal{D}_{i}^{1}(c,t_{1},t_{2},a,b)\xi^{i}\eta^{2-i}\right) \\ \\ +s\left(\sum_{i=0}^{2}\mathcal{C}_{i}^{2}(c,t_{1},t_{2},a,b)x^{i}y^{2-i}+\sum_{i=0}^{2}\mathcal{D}_{i}^{2}(c,t_{1},t_{2},a,b)\xi^{i}\eta^{2-i}\right) \\ \\ r,s\in\mathbb{Q}_{p} \right\}:(a,b)\in\left\{ (1,0),(a_{1},b_{1})\right\},c,t_{1},t_{2},a_{1},b_{1} \text{ in one row of Table 1} \right\},\end{cases}$$

those with one degree of freedom are

$$\left\{\left\{r(x^{2}+c_{1}\xi^{2})+sy^{2}/2:r\in\mathbb{Q}_{p}\right\}:c_{1}\in X_{p},s\in Y_{p}\cup\{1\}\right\}\cup\left\{\left\{r(x\xi+y\eta)+y\xi:r\in\mathbb{Q}_{p}\right\}\right\}$$
$$\cup\left\{\left\{r(x\eta+cy\xi)+a(x^{2}+y^{2})/2:r\in\mathbb{Q}_{p}\right\}:c\in Y_{p},a\in\{1,h_{p}(c)\}\right\},$$

and the isolated forms are

$$\left\{ (rx^2 + sy^2)/2 : r, s \in Y_p \cup \{1\} \right\}_{73} \cup \left\{ c(x^2/2 + \xi\eta) : c \in Y_p \cup \{1\} \right\}_{73}$$

Here by "infinite family" we mean a family of normal forms of the form  $r_1f_1 + r_2f_2 + \ldots + r_kf_k$ , where  $r_i$  are parameters and k is the number of degrees of freedom, and by "isolated" we mean a form that is not part of any family.

### 10. Application to normal forms of singularities of integrable systems

The Weierstrass-Williamson classification is one of the foundational results used in the symplectic theory of integrable systems (in particular in Eliasson's linearization theorems [25, 26]). A consequence of the Weierstrass-Williamson classification states that, given an integrable system  $F = (f_1, \ldots, f_n) : (M, \omega) \to \mathbb{R}^n$  and a *non-degenerate* critical point m of F (in a precise sense which we will define shortly), it is always possible to choose linear symplectic coordinates  $(x_1, \xi_1, \ldots, x_n, \xi_n)$  with the origin at m such that in these coordinates

$$B \circ (F - F(m)) = (g_1, \ldots, g_n) + \mathcal{O}(3),$$

where B is a n-by-n matrix of reals and each  $g_i, i \in \{1, ..., n\}$  has one of the following forms:  $\xi_i$  (regular component),  $(x_i^2 + \xi_i^2)/2$  (elliptic component),  $x_i\xi_i$  (hyperbolic component), or  $x_i\xi_{i+1} - x_{i+1}\xi_i$  with the next function equal to  $x_i\xi_i + x_{i+1}\xi_{i+1}$  (focus-focus component). See Figure 1 for a representation.

10.1. Non-degenerate critical points of integrable systems. As we see next, the classification theorems for critical points of functions on symplectic manifolds can be applied to classify critical points of integrable systems. In order to do this, first we recall the notion of non-degeneracy for a critical point of an integrable system  $F : (M, \omega) \rightarrow (\mathbb{Q}_p)^n$  on a *p*-adic analytic symplectic manifold which we use in the paper, and which in the real case is equivalent to the usual definition in Vey's paper [59], see for example [49, Section 4.2.1] and [22, Lemma 2.5].

For a concise review of the basic concepts of p-adic integrable systems and p-adic symplectic geometry see [11, Section 3]. Pelayo-Voevodsky-Warren [46, Definition 7.1] defined p-adic integrable systems as follows: a p-adic analytic map on a p-adic analytic symplectic manifold of dimension 2n:

$$F := (f_1, \ldots, f_n) : (M, \omega) \to (\mathbb{Q}_p)^n$$

is a *p*-adic analytic integrable system if two conditions hold:

- The functions  $f_1, \ldots, f_n$  satisfy  $\{f_i, f_j\} = 0$  for all  $1 \le i \le j \le n$ ;
- The set where the *n* differential 1-forms  $df_1, \ldots, df_n$  are linearly dependent has *p*-adic measure zero.

In view of some examples, including the Jaynes-Cummings model treated in our previous paper [11], and which have come to our attention since [46] was written, we believe it is convenient to replace the second condition, which is too restrictive in the *p*-adic case (unlike in the real case, which is the usual formulation one gives) by the condition that the set where the *n* differential 1-forms  $df_1, \ldots, df_n$  are linearly independent is dense in M. We use this updated definition through the paper.

**Definition 10.1.** Let *n* be a positive integer. Let *p* be a prime number. Let  $(M, \omega)$  be a *p*-adic analytic symplectic manifold of dimension 2n. Let  $F = (f_1, \ldots, f_n) : (M, \omega) \to (\mathbb{Q}_p)^n$  be a *p*-adic analytic integrable system. A point  $m \in M$  is a *critical point* of *m* if the 1-forms  $df_1(m), \ldots, df_n(m)$  are linearly dependent. The number of linearly independent forms among  $df_1(m), \ldots, df_n(m)$  is called the *rank* of the critical point.

In the following definition, a subspace U of a symplectic vector space  $(V, \omega)$  is said to be *isotropic* if  $\omega(u, v) = 0$  for any  $u, v \in U$ .

**Definition 10.2** (Non-degenerate critical point of *p*-adic analytic integrable system). Let *n* be a positive integer. Let *p* be a prime number. Let  $(M, \omega)$  be a *p*-adic analytic symplectic manifold. Let  $\Omega$  be the matrix of the linear symplectic form  $\omega_m$  on the vector space  $T_m M$ . A rank 0 critical point *m* of a *p*-adic analytic integrable system  $F = (f_1, \ldots, f_n) : (M, \omega) \to (\mathbb{Q}_p)^n$  is non-degenerate if the Hessians evaluated at *m*:

$$\mathrm{d}^2 f_1(m),\ldots,\mathrm{d}^2 f_n(m)$$

are linearly independent and if there exist  $a_1, \ldots, a_n \in \mathbb{Q}_p$  such that the matrix

$$\Omega^{-1} \sum_{i=1}^{n} a_i \mathrm{d}^2 f_i(m)$$

has n different eigenvalues.

If *m* has rank *r*, then the vectors  $X_{f_1}(m), \ldots, X_{f_n}(m)$  obtained by evaluating the Hamiltonian vector fields  $X_{f_1}, \ldots, X_{f_n}$  of  $f_1, \ldots, f_n$  at *m*, form an isotropic linear subspace *L* of  $T_m M$ , whose dimension is *r*; suppose that  $X_{f_1}(m), \ldots, X_{f_r}(m)$  are linearly independent. Then  $df_{r+1}, \ldots, df_n$  descend to  $L^{\omega}/L$  in such a way that the origin is a rank 0 critical point of the integrable system induced by *F* on  $L^{\omega}/L$ . We say that the point is *non-degenerate* if the origin is a non-degenerate critical point of this induced integrable system.

**Remark 10.3.** Definition 10.2 in the *p*-adic case is motivated by the fact that in the real case the notion of being non-degenerate for a critical point on an integrable system can also be defined in this way. Indeed, the usual definition is given in terms of Cartan subalgebras as follows: if  $(M, \omega)$  is a real symplectic manifold of dimension 2n and  $F = (f_1, \ldots, f_n) : M \to \mathbb{R}^n$  is an integrable system, a critical point *m* of *F* of rank 0 is non-degenerate if the Hessians  $d^2f_1(m), \ldots, d^2f_n(m)$  span a Cartan subalgebra of the symplectic Lie algebra of quadratic forms on the tangent space  $(T_mM, \omega_m)$ .

A *p*-adic analytic integrable system  $(f_1, \ldots, f_n) : (M, \omega) \to (\mathbb{Q}_p)^n$  is non-degenerate if all of its critical points are non-degenerate.

10.2. **Degenerate critical points in the real case.** In the real case, very little is known about *degenerate* singularities. In the literature only some results are available for some special kinds of degenerate points, see for instance [67, 68]. We give here a partial classification for real integrable systems (which must be smooth but need not be analytic).

**Theorem 10.4.** Let  $(M, \omega)$  be a real symplectic 4-manifold and let  $F = (f_1, f_2)$ :  $(M, \omega) \rightarrow \mathbb{R}^2$  be a real integrable system. Let  $m \in M$  be a degenerate critical point of F. Then one of the following two statements holds:

- (1) There exist  $a_1, a_2 \in \mathbb{R}$  and a linear combination  $g = a_1 f_1 + a_2 f_2$ , such that the four eigenvalues of  $\Omega^{-1} d^2 g(m)$  are zero.
- (2) For all linear combinations  $g = a_1f_1 + a_2f_2$ , where  $a_1, a_2 \in \mathbb{R}$ , two of the eigenvalues of  $\Omega^{-1}d^2g(m)$  are zero.

*Proof.* As m is a degenerate critical point,  $\Omega^{-1} d^2 f_1(m)$  has a multiple eigenvalue. There are three cases to consider:

- All the eigenvalues of  $\Omega^{-1}d^2f_1(m)$  are zero. We are immediately in case (1).
- Two eigenvalues of  $\Omega^{-1} d^2 f_1(m)$  are zero. Let  $g_k = f_1 + k f_2$ . The eigenvalues of  $\Omega^{-1} d^2 g_k(m)$  vary continuously with k, and they must include a multiple eigenvalue for any k. For k = 0, two eigenvalues are zero and the other two are opposite

and nonzero. This implies that, for |k| sufficiently small,  $\Omega^{-1} d^2 g_k(m)$  has also two zero eigenvalues. The values of k for which this happens are either finitely many or all  $\mathbb{R}$ , hence, it must happen for all  $k \in \mathbb{R}$ .

This means that the condition of case (2) holds when  $a_1 = 1$ , which trivially implies the same for all  $a_1 \neq 0$ . For  $a_1 = 0$ , the result follows by continuity.

•  $\Omega^{-1} d^2 f_1(m)$  is invertible. By Lemma 3.6, the eigenvalues of  $\Omega^{-1} d^2 f_1(m)$  are either of the form

$$\{r, r, -r, -r\}$$

or

$$\{r\mathbf{i}, r\mathbf{i}, -r\mathbf{i}, -r\mathbf{i}\},\$$

for some  $r \in \mathbb{R}$  with  $r \neq 0$ , and the matrix can be brought by a transformation  $\Psi$  to Jordan form, which is one of the forms

$$\begin{pmatrix} \lambda & 0 & 0 & 0 \\ 0 & -\lambda & 0 & 0 \\ 0 & 0 & \lambda & 0 \\ 0 & 0 & 0 & -\lambda \end{pmatrix} \text{ or } \begin{pmatrix} \lambda & 0 & 1 & 0 \\ 0 & -\lambda & 0 & 0 \\ 0 & 0 & \lambda & 0 \\ 0 & 1 & 0 & -\lambda \end{pmatrix},$$

where  $\lambda$  is r or ri. We call this matrix  $D_1$ . The same  $\Psi$  must bring  $\Omega^{-1} d^2 f_1(m)$  to a form that commutes with  $D_1$ : we call it  $D_2$ . We have that

$$D_2 = \begin{pmatrix} \mu_1 & 0 & \mu_2 & 0 \\ 0 & \mu_5 & 0 & \mu_6 \\ \mu_3 & 0 & \mu_4 & 0 \\ 0 & \mu_7 & 0 & \mu_8 \end{pmatrix}$$

and

$$\Psi^{T} d^{2} f_{2} \Psi = \Psi^{T} \Omega_{0} \Psi \Psi^{-1} \Omega_{0}^{-1} d^{2} f_{2} \Psi = \begin{pmatrix} 0 & a & 0 & 0 \\ -a & 0 & 0 & 0 \\ 0 & 0 & 0 & b \\ 0 & 0 & -b & 0 \end{pmatrix} D_{2}$$

for some  $a, b \in \mathbb{C}$ , must be a symmetric matrix, which implies  $\mu_1 = -\mu_5$ ,  $\mu_4 = -\mu_8$ ,  $a\mu_2 = -b\mu_7$  and  $b\mu_3 = -a\mu_6$ . The last two equalities together imply  $\mu_2\mu_3 = \mu_6\mu_7$ .

If  $D_1$  has the non-diagonal form,  $D_1D_2 = D_2D_1$  implies that  $\mu_3 = \mu_6 = 0$ ,  $\mu_1 = \mu_4$  and  $\mu_5 = \mu_8$ . We can make a linear combination of the two matrices where the diagonal is 0, and whose eigenvalues will all be zero, and we are done. Now we suppose that  $D_1$  is diagonal.

If  $D_1$  is real, the matrix  $\Psi$  is and consequently  $D_2$  are real. Otherwise, we have  $\Omega_0^{-1} d^2 f_2 \Psi = \Psi D_2$  and the columns of  $\Psi$  have the form  $(u, \bar{u}, v, \bar{v})$ , this gives  $\Omega_0^{-1} d^2 f_2 (u \ \bar{u} \ v \ \bar{v}) = (\mu_1 u + \mu_3 v \ \mu_5 \bar{u} + \mu_7 \bar{v} \ \mu_2 u + \mu_4 v \ \mu_6 \bar{u} + \mu_8 \bar{v})$  which implies that

$$\overline{\mu_1 u + \mu_3 v} = \mu_5 \bar{u} + \mu_7 \bar{v}$$

and

$$\overline{\mu_2 u + \mu_4 v} = \mu_6 \overline{u} + \mu_8 \overline{v},$$

and as u and v are linearly independent, we must have  $\bar{\mu}_1 = \mu_5 = -\mu_1$  and  $\bar{\mu}_4 = \mu_8 = -\mu_4$ . That is, the diagonal of  $D_2$  is real if  $D_1$  is real, and imaginary if  $D_1$  is imaginary.

Let  $\mu_1 = r_1 \alpha$  and  $\mu_4 = r_2 \alpha$ , for  $r_1, r_2$  in  $\mathbb{R}$  and  $\alpha \in \{1, i\}$ . By adding a multiple of  $D_1$ , we can make  $r_2 = -r_1$ , so that  $D_2$  has the form

$$\begin{pmatrix} r_1 \alpha & 0 & \mu_2 & 0 \\ 0 & -r_1 \alpha & 0 & \mu_6 \\ \mu_3 & 0 & -r_1 \alpha & 0 \\ 0 & \mu_7 & 0 & r_1 \alpha \end{pmatrix}$$

and its characteristic polynomial is

$$(t^2 - r_1^2 \alpha^2 - \mu_2 \mu_3)(t^2 - r_1^2 \alpha^2 - \mu_6 \mu_7) = (t^2 - r_1^2 \alpha^2 - \mu_2 \mu_3)^2.$$

We now prove that  $r_1^2 \alpha^2 + \mu_2 \mu_3$  must be 0, that is, the characteristic polynomial is  $t^4$  and the conclusion follows. Suppose, for a contradiction, that it is not 0. Then  $D_2$  has double eigenvalues different from zero, in the form

$$\{s, s, -s, -s\}$$

or

$$\{si, si, -si, -si\}$$

and each pair of double eigenvalues has an element coming from each factor.

Consider the matrix  $D_3 = D_2 + kD_1$ , where k is a real number close to zero. The characteristic polynomial of  $D_3$  is still a product of two factors, but now they are different. For small enough k, the only way to have multiple eigenvalues is to have the same form as in  $D_2$ , that is, two real or imaginary pairs with an element from each factor. But this is not possible if the two factors are different. 

10.3. Proof of Theorems A, B and C. By Theorem 9.3, we may assume without loss of generality that  $M = (\mathbb{Q}_p)^4$  and  $\omega_m = \omega_0$ .

For a rank 1 critical point, we can take  $d\eta$  in the direction of the nonzero differential, and the problem reduces to classify the critical point in  $L^{\omega_0}/L$ . This is a system with one function (the linear combination of  $f_1$  and  $f_2$  with differential 0) in dimension 2, so we apply Corollary 9.7 and get the result.

For a rank 0 critical point, first we prove existence. The fact that  $f_1$  and  $f_2$  form an integrable system implies that  $\{f_1, f_2\} = 0$ , that is,

$$(\mathrm{d}f_1)^T \Omega_0^{-1} \mathrm{d}f_2 = 0.$$

Differentiating this twice, evaluating at m and using that  $df_1(m) = df_2(m) = 0$ ,

$$d^{2}f_{1}(m)\Omega_{0}^{-1}d^{2}f_{2}(m) = d^{2}f_{2}(m)\Omega_{0}^{-1}d^{2}f_{1}(m)$$

We define  $A_i = \Omega_0^{-1} d^2 f_i(m)$ . The previous expression implies that  $A_1$  and  $A_2$  commute.

Let  $u \in (\mathbb{C}_p)^4$  be an eigenvector of  $A_1$ , where  $\mathbb{C}_p$  is the field of complex *p*-adic numbers. Then

$$A_1 A_2 u = A_2 A_1 u = \lambda A_2 u$$

for  $\lambda \in \mathbb{C}_p$ . This implies that  $A_2 u$  is also an eigenvector of  $A_1$  with value  $\lambda$ . But the critical point is non-degenerate, which means that the only eigenvector with value  $\lambda$  is u. Hence,  $A_2 u = \mu u$  for some  $\mu \in \mathbb{C}_p$ , and u is also eigenvector of  $A_2$ . So  $A_1$  and  $A_2$  have the same eigenvectors.

In the proof of Theorem F (Section 6.3), the case in which the Hessian of f falls and the values of the parameters  $(c_1, c_2, c, a, \text{etc.})$  in the normal form are determined by the eigenvectors. This means that S is the same matrix for  $f_1$  than for  $f_2$ , and the resulting normal forms are  $r_1g_1 + s_1g_2$  and  $r_2g_1 + s_2g_2$ , for some  $r_1, r_2, s_1, s_2 \in \mathbb{Q}_p$  and  $g_1$  and  $g_2$ 77 are among the possibilities of Corollary 9.9. As the Hessians are linearly independent, the matrix

$$\begin{pmatrix} r_1 & s_1 \\ r_2 & s_2 \end{pmatrix}$$

changing  $(g_1, g_2)$  to  $(f_1, f_2)$  can be inverted, giving the matrix B that we need, and the proof of existence of Theorem A is complete. Uniqueness follows directly from Corollary 9.9.

Theorems **B** and **C** follow from applying Theorems **H** and **I**, respectively, to the Hessians of the components of the system: each normal form of the matrix gives a normal form of the integrable system, and the Hessian of  $f_{P,p}$  is exactly the matrix M(P, p).

10.4. Degenerate critical points in the *p*-adic case. Based on Theorem G, a version of Theorem A can also be deduced for degenerate singularities (a topic of growing interest in real symplectic geometry; see for instance [22, 23, 32] and the references therein) but the statement will be more complicated than that of Theorem A. The reason is that, while for non-degenerate singularities the types (in the sense of Corollary 9.9) of  $f_1$  and  $f_2$  must coincide if these functions Poisson-commute, this does not happen for degenerate singularities. For example, the function of type (1) in Corollary 9.10 with r = s = 1 and  $c_1 = c_2 = -1$  Poisson-commutes with the one of type (2) in the same list with r = 1. Hence, a full classification will need many more cases. We can give a partial result analogous to Theorem 10.4:

**Theorem 10.5.** Let p be a prime number. Let  $(M, \omega)$  be a 4-dimensional p-adic analytic symplectic manifold. Let  $F = (f_1, f_2) : (M, \omega) \to (\mathbb{Q}_p)^2$  be a p-adic analytic integrable system and let  $m \in M$  be a degenerate rank 0 critical point of F. Then one of the following two statements holds:

- (1) There are  $a_1, a_2 \in \mathbb{Q}_p$  and a linear combination  $g = a_1 f_1 + a_2 f_2$ , such that all four eigenvalues of  $\Omega^{-1} d^2 g$  are zero;
- (2) For all linear combinations  $g = a_1 f_1 + a_2 f_2$ , where  $a_1, a_2 \in \mathbb{Q}_p$ , two eigenvalues of  $\Omega^{-1} d^2 g$  are zero.

*Proof.* This proof follows the same strategy as in the real case (Theorem 10.4), we include it here for completeness.

As m is degenerate, for every linear combination  $g = a_1 f_1 + a_2 f_2$ , m is a critical point of g with repeated eigenvalues. Concretely this happens for  $f_1$  and  $f_2$  themselves. There are three cases to consider:

- $\Omega^{-1} d^2 f_1$  has all eigenvalues zero, and we are in case (1).
- $\Omega^{-1} d^2 f_1$  has exactly two eigenvalues zero. Since the other two are nonzero, they are opposites.

Consider  $g_k = f_1 + kf_2$ , for  $k \in \mathbb{Q}_p$ . Since *m* is degenerate,  $\Omega^{-1} d^2 g_k$  has a multiple eigenvalue for any *k*. If *k* is small enough, the only way to have a multiple eigenvalue is that the same two that coincide for k = 0 keep coinciding. But as they must be opposites, they must still coincide at zero. This means that  $\Omega^{-1} d^2 g_k$  has two eigenvalues equal to zero for all *k* small enough. But the values of *k* for which this happens are either finitely many or all  $\mathbb{Q}_p$ , so it must happen for all *k*.

It follows that case (2) holds when  $a_1 = 1$ , and can be extended trivially to all  $a_1 \neq 0$ . For  $a_1 = 0$ , it must also hold by continuity.

•  $\Omega^{-1} d^2 f_1$  is invertible. By Corollary 9.10, we can bring  $f_1$  to a normal form where  $\omega_m = \omega_0$ . The ones with an invertible Hessian are in case (1) with  $c_1 \neq 0$  and

 $c_2 \neq 0$ , case (2) with  $r \neq 0$ , and case (3) with  $r \neq 0$ . In the three cases, the eigenvalues are of the form  $\{\lambda, \lambda, -\lambda, -\lambda\}$ , where  $\lambda^2 \in \mathbb{Q}_p$  (in some of them  $\lambda \in \mathbb{Q}_p$  and in others it is in a degree two extension).

By Lemma 3.6, we can bring the matrix  $\Omega_0^{-1} d^2 f_1$  to Jordan form by a transformation  $\Psi$ , which is one of the two forms

$$\begin{pmatrix} \lambda & 0 & 0 & 0 \\ 0 & -\lambda & 0 & 0 \\ 0 & 0 & \lambda & 0 \\ 0 & 0 & 0 & -\lambda \end{pmatrix} \text{ or } \begin{pmatrix} \lambda & 0 & 1 & 0 \\ 0 & -\lambda & 0 & 0 \\ 0 & 0 & \lambda & 0 \\ 0 & 1 & 0 & -\lambda \end{pmatrix}$$

We call this matrix  $D_1$ . The matrix  $\Omega_0^{-1} d^2 f_2$  will become, by the same transformation  $\Psi$ , a matrix  $D_2$  that commutes with  $D_1$ , so it has the form

$$\begin{pmatrix} \mu_1 & 0 & \mu_2 & 0 \\ 0 & \mu_5 & 0 & \mu_6 \\ \mu_3 & 0 & \mu_4 & 0 \\ 0 & \mu_7 & 0 & \mu_8 \end{pmatrix}$$

We have that  $\Psi^T d^2 f_2 \Psi$  is symmetric, which leads to  $\mu_1 = -\mu_5$ ,  $\mu_4 = -\mu_8$ , and  $\mu_2 \mu_3 = \mu_6 \mu_7$ .

If  $D_1$  has the second form above,  $D_1D_2 = D_2D_1$  also implies  $\mu_3 = \mu_6 = 0$ ,  $\mu_1 = \mu_4$  and  $\mu_5 = \mu_8$ . Adding an adequate multiple of  $D_1$  to  $D_2$ , we can make  $\mu_1 = \mu_4 = \mu_5 = \mu_8 = 0$ , so that this new matrix has all eigenvalues zero, and we are done. Now we suppose that  $D_1$  has the first form.

If  $\lambda \in \mathbb{Q}_p$ , the matrix  $\Psi$  is in  $\mathbb{Q}_p$ , so  $D_2$  is also in  $\mathbb{Q}_p$ . If  $\lambda \notin \mathbb{Q}_p$ , we have  $\Omega_0^{-1} d^2 f_2 \Psi = \Psi D_2$ . Taking the columns of  $\Psi$  in the form  $(u, \bar{u}, v, \bar{v})$ , as happens in the real case but now  $\bar{u}$  means the conjugate in  $\mathbb{Q}_p[\lambda]$ , we arrive at  $\bar{\mu}_1 = \mu_5 = -\mu_1$  and  $\bar{\mu}_4 = \mu_8 = -\mu_4$ . This means that the elements in the diagonal of  $D_2$  are multiples of  $\lambda$  with coefficients in  $\mathbb{Q}_p$ , independently of whether  $\lambda$  is in  $\mathbb{Q}_p$  or not.

Let  $\mu_1 = r_1 \lambda$  and  $\mu_4 = r_2 \lambda$ , for  $r_1, r_2$  in  $\mathbb{Q}_p$ . By adding a multiple of  $D_1$ , we can make  $r_2 = -r_1$ , and  $D_2$  has now the form

$$egin{pmatrix} r_1\lambda & 0 & \mu_2 & 0 \ 0 & -r_1\lambda & 0 & \mu_6 \ \mu_3 & 0 & -r_1\lambda & 0 \ 0 & \mu_7 & 0 & r_1\lambda \end{pmatrix}.$$

The characteristic polynomial of this matrix is

$$(t^2 - r_1^2 \lambda^2 - \mu_2 \mu_3)(t^2 - r_1^2 \lambda^2 - \mu_6 \mu_7) = (t^2 - r_1^2 \lambda^2 - \mu_2 \mu_3)^2.$$

We now prove that  $r_1^2 \lambda^2 + \mu_2 \mu_3$  must be 0, from which the conclusion will follow. Suppose, for a contradiction, that it is not 0. Then  $D_2$  has double eigenvalues different from zero, and each pair of double eigenvalues has an element coming from each factor.

Consider the matrix  $D_3 = D_2 + kD_1$ , where k is a small p-adic number. The characteristic polynomial of  $D_3$  is still a product of two factors. Since the point m is degenerate, for small enough k we must have the same behavior of the eigenvalues as in  $D_2$ , that is, two pairs with an element from each factor. But this cannot hold because the two factors are not equal.

**Corollary 10.6.** Let p be a prime number. Let  $X_p, Y_p$  be the non-residue sets in Definition 1.1. If the p-adic analytic integrable system  $F : (M, \omega) \to (\mathbb{Q}_p)^2$  falls in the first case of

Theorem 10.5, then the linear combination  $g = a_1f_1 + a_2f_2$  therein can be brought to one of the following forms by a linear symplectomorphism:

- (1)  $(rx^2 + sy^2)/2 + \mathcal{O}(3)$ , for  $r, s \in Y_p \cup \{0, 1\}$ .
- (2)  $y\xi + \mathcal{O}(3)$ .
- (3)  $c(x^2/2 + \xi\eta) + \mathcal{O}(3)$ , for  $c \in Y_n \cup \{1\}$ .

*Proof.* It follows directly from Theorem 10.5 and Corollary 9.10. The selected forms are precisely those having all eigenvalues zero. 

In principle, Corollary 10.6 can be a first step towards making a classification analogous to Theorem A for degenerate critical points, but such a classification will be more complicated than the non-degenerate one, so we will not follow this direction in the paper.

**Remark 10.7.** There is an extensive theory of quadratic forms over different types of fields, we refer to the classical treatment [41] and the more recent works by Alsina-Bayer [2], Bhargava [7], Casselman [10] and Lam [36] and the references therein. In Theorem A and Theorem 10.5 we have presented a list of local normal forms of integrable systems up to linear symplectic transformations, given by sums of binary quadratic forms, but we have not carried out a further analysis of the structure/properties of these forms since this does not appear to us as applicable in our context of symplectic geometry of integrable systems.

10.5. Symplectic dynamics of integrable systems and their level sets. We now calculate the vector fields generated by the integrable systems of Theorem A.

**Proposition 10.8.** The vector fields generated by the integrable systems of Theorem A are as follows:

(1) 
$$X_{g_1} = (2c_1\xi, -2x, 0, 0), X_{g_2} = (0, 0, 2c_2\eta, -2y).$$
  
(2)  $X_{g_1} = (cy, -\eta, x, -c\xi), X_{g_2} = (x, -\xi, y, -\eta).$   
(3)  
 $X_{g_1} = \left(-\frac{t_1 + bt_2}{a}\xi - (bt_1 + ct_2)\eta, \frac{acx - by}{b^2 - c}, -ac(t_1 + bt_2)\eta - (bt_1 + ct_2)\xi, \frac{y - abx}{a(b^2 - c)}\right),$   
 $X_{g_2} = \left(-\frac{bt_1 + ct_2}{a}\xi - c(t_1 + bt_2)\eta, \frac{cy - abcx}{b^2 - c}, -ac(bt_1 + ct_2)\eta - c(t_1 + bt_2)\xi, \frac{acx - by}{a(b^2 - c)}\right).$   
*Proof.* All fields are calculated applying directly the equation  $i_{X_t}\omega_0 = df.$ 

*Proof.* All fields are calculated applying directly the equation  $i_{X_f}\omega_0 = df$ .

We can check that, in each system,  $g_1$  and  $g_2$  Poisson commute, which is equivalent to checking that, if  $A_i = \Omega_0^{-1} d^2 g_i$ , we have  $A_1 A_2 = A_2 A_1$ . This matrix is zero in case (1),

(0	0	c	$0 \rangle$
0	0	0	1
1	0	0	0
$\int 0$	c	0	0/

in case (2), and

$$\begin{pmatrix} ct_2 & 0 & \frac{t_1}{a} & 0\\ 0 & ct_2 & 0 & act_1\\ act_1 & 0 & ct_2 & 0\\ 0 & \frac{t_1}{a} & 0 & ct_2 \end{pmatrix}$$

in case (3).

We can also calculate the fibers of the systems.

**Proposition 10.9.** The fibers  $F^{-1}(0,0)$  of the integrable systems of Theorem A with a critical point of rank 0 are as follows:

- (1)  $\{(\pm d_1\xi, \xi, \pm d_2\eta, \eta) : \xi, \eta \in \mathbb{Q}_p\}$  if  $-c_1 = d_1^2$  and  $-c_2 = d_2^2$ ,  $\{(\pm d_1\xi, \xi, 0, 0) : \xi \in \mathbb{Q}_p\}$  if  $-c_1 = d_1^2$  and  $-c_2$  is not a square, and  $\{(0, 0, 0, 0)\}$  if  $-c_1$  and  $-c_2$  are not squares.
- (2)  $\{(x, 0, y, 0) : x, y \in \mathbb{Q}_p\} \cup \{(0, \xi, 0, \eta) : \xi, \eta \in \mathbb{Q}_p\}.$
- $(3) \{(0,0,0,0)\}.$

For those in which the origin has rank 1, the fibers are  $\{(\pm d\xi, \xi, y, 0) : \xi, y \in \mathbb{Q}_p\}$  if  $-c = d^2$  and  $\{(0, 0, y, 0) : y \in \mathbb{Q}_p\}$  otherwise.

*Proof.* The part about rank 1 is immediate from the formula  $(x^2 + c\xi^2, \eta)$ . For the rank 0 part, we have:

- (1) This follows from making  $x^2 + c_1\xi^2 = y^2 + c_2\eta^2 = 0$ .
- (2) We make  $x\eta + cy\xi = x\xi + y\eta = 0$ . Considering this a system in  $(\xi, \eta)$ , we have two possible cases:
  - The determinant of the coefficient matrix is 0. Then  $cy^2 x^2 = 0$ . Since c is not a square, x = y = 0.
  - The determinant of the coefficient matrix is not 0. Then  $\xi = \eta = 0$ .
- (3) We consider the coordinate change given by  $\Psi_2$  in Proposition 5.2:

$$\begin{pmatrix} x\\\xi\\y\\\eta \end{pmatrix} = \Psi_2 \begin{pmatrix} x'\\\xi'\\y'\\\eta' \end{pmatrix}.$$

We have that the first column of  $\Psi_2$  is the hat-conjugate of the second, the same happens for the third and fourth, and the original coordinates are all in  $\mathbb{Q}_p$  and they are their own conjugates, hence

$$\begin{pmatrix} x\\ \xi\\ y\\ \eta \end{pmatrix} = \begin{pmatrix} \hat{x}\\ \hat{\xi}\\ \hat{y}\\ \hat{\eta} \end{pmatrix} = \hat{\Psi}_2 \begin{pmatrix} \hat{x}'\\ \hat{\xi}'\\ \hat{y}'\\ \hat{\eta}' \end{pmatrix} = \Psi_2 \begin{pmatrix} \hat{\xi}'\\ \hat{x}'\\ \hat{\eta}'\\ \hat{\eta}' \end{pmatrix}$$

which implies  $\xi' = \hat{x}'$  and  $\eta' = \hat{y}'$ . Now, if  $(x, \xi, y, \eta)$  is a point in the fiber,

$$0 = \begin{pmatrix} x & \xi & y & \eta \end{pmatrix} M \begin{pmatrix} x \\ \xi \\ y \\ \eta \end{pmatrix} = \begin{pmatrix} x' & \xi' & y' & \eta' \end{pmatrix} \Psi_2^T M \Psi_2 \begin{pmatrix} x' \\ \xi' \\ y' \\ \eta' \end{pmatrix}$$

where M is the matrix of the normal form. We know that  $\Psi_2$  diagonalizes  $\Omega_0^{-1}M$ , so

$$\begin{split} \Psi_{2}^{\circ} M \Psi_{2} &= \Psi_{2}^{\circ} \Omega_{0} \Psi_{2} \Psi_{2}^{\circ} \Omega_{0}^{\circ} M \Psi_{2} \\ &= \begin{pmatrix} 0 & 4a\alpha \gamma(b+\alpha) & 0 & 0 \\ -4a\alpha \gamma(b+\alpha) & 0 & 0 & 0 \\ 0 & 0 & 0 & -4a\alpha \bar{\gamma}(b-\alpha) \\ 0 & 0 & 4a\alpha \bar{\gamma}(b-\alpha) & 0 \end{pmatrix} \begin{pmatrix} \lambda & 0 & 0 & 0 \\ 0 & -\lambda & 0 & 0 \\ 0 & 0 & \mu & 0 \\ 0 & 0 & 0 & -\mu \end{pmatrix} \\ &= -4a\alpha \begin{pmatrix} 0 & \gamma(b+\alpha)\lambda & 0 & 0 & 0 \\ \gamma(b+\alpha)\lambda & 0 & 0 & 0 \\ 0 & 0 & 0 & \bar{\gamma}(b-\alpha)\mu \\ 0 & 0 & \bar{\gamma}(b-\alpha)\mu \\ &= 81 \end{pmatrix}. \end{split}$$

Putting this together, we get

$$0 = \gamma(b+\alpha)\lambda x'\xi' + \bar{\gamma}(b-\alpha)\mu y'\eta' = \gamma^2(b+\alpha)(r+s\alpha)x'\hat{x}' + \bar{\gamma}^2(b-\alpha)(r-s\alpha)y'\hat{y}$$

This must hold for all  $r, s \in \mathbb{Q}_p$ . Putting (r, s) = (1, 0) and (0, 1),

$$\gamma^2(b+\alpha)x'\hat{x}' + \bar{\gamma}^2(b-\alpha)y'\hat{y}' = \alpha\gamma^2(b+\alpha)x'\hat{x}' - \alpha\bar{\gamma}^2(b-\alpha)y'\hat{y}' = 0$$

These two equations imply  $x'\hat{x}' = 0$  and  $y'\hat{y}' = 0$ , that is, x' = y' = 0, which in turn implies  $\xi' = \eta' = 0$  and the vector is zero.

In analogy with the real case, a submanifold N of a 2n-dimensional p-adic analytic symplectic manifold  $(M, \omega)$  is said to be *isotropic* if the tangent space at each point of N is an isotropic subspace of the tangent space of M, that is, if  $\omega(u, v) = 0$  for any two vectors  $u, v \in T_m N$ . It is called *Lagrangian* if it is isotropic and with dimension n.

The fibers of regular points of real integrable systems are Lagrangian, and homeomorphic to tori (for this reason, it is called a singular Lagrangian torus fibration). For p-adic integrable systems the situation is more complicated to describe in general, even for concrete examples (such as the Jaynes-Cummings model treated in our paper [11]). However, there is a common point:

**Proposition 10.10.** Let n be a positive integer. Let p be a prime number. Let  $(M, \omega)$  be a 2n-dimensional p-adic symplectic manifold. Let  $F : (M, \omega) \to (\mathbb{Q}_p)^n$  be a p-adic analytic integrable system. Suppose that the components of F are either regular components  $\xi_i$  or given by one of the normal forms of Theorem A. Then the fiber  $F^{-1}(0)$  is a union of isotropic subspaces intersecting at the origin, and if all the components are regular, then the fiber is a Lagrangian subspace.

*Proof.* For the regular case, where the system is  $(\xi_1, \ldots, \xi_n)$ , clearly the fiber of 0 is a Lagrangian subspace. Otherwise, it is enough to prove the statement for the dimension 2 and 4 normal forms, and the conclusion follows by multiplying, because the symplectic form  $\omega_0$  does not mix the coordinates of different components. For the normal forms, the result reduces to checking that

$$\omega_0((\pm d_1, 1, 0, 0), (0, 0, \pm d_2, 1)) = \omega_0((1, 0, 0, 0), (0, 0, 1, 0))$$
  
=  $\omega_0((\pm d, 1, 0, 0), (0, 0, 1, 0)) = 0.$ 

From the point of view of integrable systems and symplectic singularity theory, the fibers and images of the local models in the p-adic case are very interesting and include for example the images displayed in Figures 15 and 16 and the fibers displayed in Figures 2 and 3.

We could define a *p*-adic version of the "Williamson type" defined for the real case at [8, p. 41]. In the real case, it consists of a tuple of integers  $(k_{\rm e}, k_{\rm h}, k_{\rm f})$  that count the number of elliptic, hyperbolic and focus-focus components of the normal form. The problem with this approach is that the components of the normal forms are associated to blocks in the normal forms of matrices, which in the real case take only three possible forms. In the *p*-adic case, by Lemma 8.3, there can appear countably many different blocks, so the Williamson type will be a sequence (with a finite number of elements different from zero) instead of a tuple.

For example, for p = 2, the first 11 elements of the sequence count the number of components with each possible c in Corollary 9.7 (associated to blocks of size two), the next 145 elements count the number of pairs of components with each possible form in parts (2) and (3) of Theorem A (associated to blocks of size four), after which come the counts of trios of components associated to blocks of size six, and so on.




FIGURE 15. Images of some normal forms for the real case and the p-adic case with  $p \not\equiv 1 \mod 4$ . In the real case, the positive and negative sides of the axes represent, as usual, positive and negative numbers; if p = 2, the "positive" and "negative" sides represent numbers whose second digit is 0 and 1, respectively; finally, if  $p \equiv 3 \mod 4$ , the "positive" and "negative" sides represent even-order and odd-order numbers, respectively. (The points on the axes themselves have, as usual, a zero coordinate.) In each drawing, the green region represents regular values, the blue points are rank 1 critical values, and the red points are rank 0 critical values. An elliptic component has c = 1 in the notation of Corollary 9.7, a hyperbolic one has c = -1, and a focus-focus one has c = -1 in part (2) of Theorem A.




FIGURE 16. Images of the same normal forms for  $p \equiv 1 \mod 4$ . The slopes of the blue lines in the last image are i and -i. Note that the images of the systems with the same rank coincide, except perhaps for a coordinate change, because now the singular components of the systems belong to the same class: part (1) of Theorem A, with  $c_1 = c_2 = 1$ .

We close this section with a mention of the Eliasson-Vey's linearization theorem [25, 26, 53, 59, 61], which in the real case states that any *smooth* integrable system can be brought to its Williamson normal form by a symplectomorphism. The *analytic* case of this theorem is due to Rüßmann [53] for two degrees of freedom and Vey [59] in arbitrary dimension. In the real case Eliasson's Theorem (assuming that there are no hyperbolic components) says that there is a local diffeomorphism  $\varphi$  and symplectic coordinates  $\phi^{-1} = (x, \xi, y, \eta)$  such that  $F \circ \phi = \varphi(g_1, g_2)$ , where  $g_i$  is one of the elliptic, real or focus-focus models. The *p*-adic equivalent of this theorem is well beyond the scope of this paper, and we state it as a question.

Question 10.11 (A *p*-adic Eliasson-Vey's theorem?). Let *n* be a positive integer. Let *p* be a prime number. Given a 2*n*-dimensional *p*-adic analytic symplectic manifold  $(M, \omega)$ , an integrable system  $F : (M, \omega) \to (\mathbb{Q}_p)^n$  and a non-degenerate critical point *m* of *F*, determine under which conditions on the Williamson type of the critical point *m* there are open sets  $U \subset M$  and  $V \subset (\mathbb{Q}_p)^{2n}$ , a *p*-adic analytic symplectomorphism  $\phi : V \to U$  and a local diffeomorphism  $\varphi$  of  $(\mathbb{Q}_p)^n$  such that  $\phi(0) = m$  and

$$(F - F(m)) \circ \phi = \varphi \circ (g_1, \ldots, g_n),$$

where  $(g_1, \ldots, g_n)$  is the Williamson normal form of F in m. (In the real case it is enough that there are no hyperbolic blocks.)

**Remark 10.12.** It is important to understand that the results of this section do provide a normal form of the functions involved in a neighborhood of the critical point, but do not give a normal form of the symplectic form in the entire neighborhood, only at the tangent space at the point. The question of whether such a normal form exists in the entire neighborhood and what shapes it may take is a completely different non-linear problem which we have not attempted to answer in this paper (this would be a Darboux type theorem for *p*-adic analytic symplectic manifolds, which would be a preliminary step to answer Question 10.11).

### 11. Application to classical mechanical systems

In this section we explain how Theorem A can be applied to further study the p-adic Jaynes-Cummings model introduced and studied in [11]. We recommend the books by Abraham-Marsden [1] and de León-Rodrigues [13] for an introduction to the mathematical study of mechanics and its connections to symplectic/differential geometry.

In our paper [11] we studied the Jaynes-Cummings model with *p*-adic coefficients (see Figure 17 for the fibers in the real case and Figure 18 for a fiber in the *p*-adic case). The system was defined therein, in analogy with the real case, as follows. For any number p, first we consider the product *p*-adic analytic manifold  $S_p^2 \times (\mathbb{Q}_p)^2$  with the *p*-adic symplectic form  $\omega_{S_p^2} + du \wedge dv$ . Here we recall that

$$S_p^2 = \left\{ (x, y, z) \in \mathbb{Q}_p^2 : x^2 + y^2 + z^2 = 1 \right\}$$

and  $\omega_{S_p^2}$  is the area form in the sphere given by

$$\omega_{\mathbf{S}_p^2} = -\frac{1}{z} \mathrm{d}x \wedge \mathrm{d}y = \frac{1}{y} \mathrm{d}x \wedge \mathrm{d}z = -\frac{1}{x} \mathrm{d}y \wedge \mathrm{d}z.$$

The *p*-adic Jaynes-Cummings model is given by the *p*-adic analytic map

$$F = (J, H) : \mathrm{S}_p^2 \times (\mathbb{Q}_p)^2 \to (\mathbb{Q}_p)^2,$$
<sup>84</sup>




FIGURE 17. Image and fibers of the real Jaynes-Cummings model. The red curve consists of rank 1 critical points, and the two blue points are rank 0. The Jaynes-Cummings model is an example of a class of integrable systems called *semitoric systems*. The fibers of this system are a point, circles, 2-tori (generic fiber) and a pinched torus.

where

$$\begin{cases} J(x, y, z, u, v) = \frac{u^2 + v^2}{2} + z; \\ H(x, y, z, u, v) = \frac{ux + vy}{2}, \end{cases}$$

where  $(x, y, z) \in S_p^2$  and  $(u, v) \in (\mathbb{Q}_p)^2$ .

By [11, Proposition 2.5], at  $m_1 = (0, 0, -1, 0, 0)$ , there is a *p*-adic linear symplectomorphism changing the local coordinates to  $(x, \xi, y, \eta)$  in which the *p*-adic symplectic form is given by  $\omega = (dx \wedge d\xi + dy \wedge d\eta)/2$  and

(11.1) 
$$F_1(x,\xi,y,\eta) = \frac{1}{2}(x^2 + \xi^2, y^2 + \eta^2) + \mathcal{O}((x,\xi,y,\eta)^3).$$

Here  $F_1 = B \circ (F - F(0, 0, -1, 0, 0))$  with

$$B = \begin{pmatrix} 1 & 2 \\ 1 & -2 \end{pmatrix}.$$

At  $m_2 = (0, 0, 1, 0, 0)$ , after the change, the *p*-adic symplectic form is also given by  $\omega = (dx \wedge d\xi + dy \wedge d\eta)/2$  and

(11.2) 
$$F_2(x,\xi,y,\eta) = (x\eta - y\xi, x\xi + y\eta) + \mathcal{O}((x,\xi,y,\eta)^3).$$




FIGURE 18. Fiber  $F^{-1}(72, 1)$  of the 5-adic Jaynes-Cummings model. The blue points are values of z for which the coordinates (x, y, u, v) form two p-adic circles, at the green points they only form one circle, at the purple point z = j = 72 they have dimension 1 but they are not a circle, and the values of z which appear in grey are not in the fiber.

Here  $F_2 = B \circ (F - F(0, 0, 1, 0, 0))$  with

$$B = \begin{pmatrix} 2 & 0 \\ 0 & 4 \end{pmatrix}.$$

Hence we have the following consequence of Theorem A.

**Corollary 11.1.** Let p be a prime number. Then there exist open sets  $U_1$  and  $U_2$  such that  $m_1 \in U_1$  and  $m_2 \in U_2$  and a local linear symplectomorphism  $\phi : U_1 \to U_2$  centered at  $m_1$ , such that

$$F_2(\phi(x,\xi,y,\eta)) = F_1(x,\xi,y,\eta) + \mathcal{O}((x,\xi,y,\eta)^3)$$

for  $(x, \xi, y, \eta) \in U_1$ , if and only if  $p \equiv 1 \mod 4$ , where  $F_1$  and  $F_2$  are as described in (11.1) and (11.2).

*Proof.* By Theorem A,  $F_1$  and  $F_2$  are linearly symplectomorphic to one of the possibilities listed in its statement, so it is enough to see that it is the same for  $p \equiv 1 \mod 4$  and different otherwise. For the first normal form  $F_1$ , we have that

$$\Omega_0^{-1} d^2 (rJ_1 + sH_1) = \begin{pmatrix} 0 & -2 & 0 & 0 \\ 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 \\ 0 & 0 & 2 & 0 \end{pmatrix} \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & r & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & s \end{pmatrix} = \begin{pmatrix} 0 & -2r & 0 & 0 \\ 2r & 0 & 0 & 0 \\ 0 & 0 & 0 & -2s \\ 0 & 0 & 2s & 0 \end{pmatrix}$$

whose eigenvalues are  $\pm 2ir$  and  $\pm 2is$ . If  $p \equiv 1 \mod 4$ , we are in the situation of Proposition 4.1, so this is in case (1) of Theorem A linearly symplectomorphic to  $x^2 + \xi^2$ .

Otherwise, we are in the situation of Proposition 4.6, and for  $\lambda = 2ir$ ,

$$u^{T} \Omega_{0} \bar{u} = \begin{pmatrix} i & 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & \frac{1}{2} & 0 & 0 \\ -\frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2} \\ 0 & 0 & -\frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} -i \\ 1 \\ 0 \\ 0 \end{pmatrix} = i$$

and

$$\frac{2\lambda a}{u^T \Omega_0 \bar{u}} = 4ra$$

We need to find a and b such that  $ab = 4r^2$  and  $4ra \in DSq(\mathbb{Q}_p, 1)$ , or equivalently  $r/a \in DSq(\mathbb{Q}_p, 1)$ . Taking b = ac, we have that  $c = 4r^2/a^2$  is a square; moreover, if  $p \equiv 3 \mod 4$ , r/a has even order, hence  $4 \mid \operatorname{ord}_p(c)$ , and in the set

$$\Big\{1,-1,p,-p,p^2\Big\},$$

the c that we need is 1. If p = 2, the only c that is square is 1. The same reasoning holds for s instead of r, so this critical point is in case (1) with  $c_1 = c_2 = 1$ .

Respecting to  $F_2$ , we get

$$\Omega_0^{-1} d^2 (rJ_2 + sH_2) = \begin{pmatrix} 0 & -2 & 0 & 0 \\ 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 \\ 0 & 0 & 2 & 0 \end{pmatrix} \begin{pmatrix} 0 & s & 0 & r \\ s & 0 & -r & 0 \\ 0 & -r & 0 & s \\ r & 0 & s & 0 \end{pmatrix} = \begin{pmatrix} -2s & 0 & 2r & 0 \\ 0 & 2s & 0 & 2r \\ -2r & 0 & -2s & 0 \\ 0 & -2r & 0 & 2s \end{pmatrix}$$

has as eigenvalues  $\pm 2s \pm 2ir$ . If  $p \equiv 1 \mod 4$ , this is again in case (1) with  $c_1 = c_2 = 1$ , otherwise it is in case (2) with c = -1.

More information about the Jaynes-Cummings model and other models of interest in physics can be found at [1, 13].

### 12. Examples

In this section we show examples which illustrate our theorems, so that they can be understood more concretely.

## 12.1. Examples with matrices.

**Example 12.1.** This example follows the method given in the proof of Theorem F in order to find the normal form of a symmetric matrix. Let M be the following symmetric matrix:

$$M = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 5 & 6 & 7 \\ 3 & 6 & 8 & 9 \\ 4 & 7 & 9 & 10 \end{pmatrix}$$

The characteristic polynomial of  $A = \Omega_0^{-1}M$  is  $t^4 - 6t^2 - 2$ . We start with p = 2. In order to classify M, we need to find two things: the family of the normal form, and the normal form itself. We first calculate

$$\lambda^2 = \frac{6 \pm \sqrt{36 + 8}}{2} = 3 \pm \sqrt{11}.$$

As  $11 \equiv 3 \mod 8$ ,  $\lambda^2 \notin \mathbb{Q}_2$  and we are in case (2) or (3) of Theorem **F** with c = 3. In order to find which one, we need to check whether  $3 \pm \sqrt{11}$  is a square in  $\mathbb{Q}_2[\sqrt{3}]$ : it can

be written as  $3 \pm \sqrt{11/3}\sqrt{3}$ , where  $\sqrt{11/3} \in \mathbb{Q}_2$ . Applying the criterion on Proposition 6.6(4), we see that

$$ord(3) = ord(\sqrt{11/3}) = 0,$$

so it is not a square and we are in case (3).

The next step is to find  $t_1$  and  $t_2$ : this is the class of  $3 + \sqrt{11}$  in  $\mathbb{Q}_2[\sqrt{3}]$  modulo squares. This can be found with the procedure in the proof of Corollary 6.7(2): we need to multiply by  $1 + \sqrt{3}$  to make  $\operatorname{ord}(a) \neq \operatorname{ord}(b)$ , which gives

$$(3 + \sqrt{\frac{11}{3}}\sqrt{3})(1 + \sqrt{3}) = 3 + \sqrt{33} + \left(3 + \sqrt{\frac{11}{3}}\right)\sqrt{3}.$$

Writing this again in the form  $a + b\sqrt{3}$ , it leads to  $\operatorname{ord}(b) - \operatorname{ord}(a) = 2$ ,  $\operatorname{ord}(a) \equiv 1 \mod 2$ (so we have to multiply by 2) and  $b/4a + \operatorname{digit}_2(a) \equiv 1 \mod 2$  (so we have to change sign). The class is  $-2(1 + \sqrt{3})$ , that is,  $t_1 = t_2 = -2$ . This is not a class in the table, so we need to take its pair,  $t_1 = 1$  and  $t_2 = 1$ .

It is only left to find a and b. The two possible classes have b = 0, but one has a = 1 and the other a = -1. To find the correct one, we go to the formula in Proposition 5.2.

$$\frac{a\alpha\gamma(b+\alpha)}{u^T\Omega_0\hat{u}} = \frac{a\alpha^2\gamma}{u^T\Omega_0\hat{u}} = \frac{3a\sqrt{-2(1+\sqrt{3})}}{u^T\Omega_0\hat{u}} \in \mathrm{DSq}(\mathbb{Q}_2[\sqrt{3}], 2(1+\sqrt{3})).$$

The class of this number, with a = 1, is  $-(1 + \sqrt{3})$ . As we see in Table 11, the position in row  $\gamma^2 = -2(1 + \sqrt{3})$  and column  $-(1 + \sqrt{3})$  is unmarked, which means that the number is not in

$$DSq(\mathbb{Q}_2[\sqrt{3}], 2(1+\sqrt{3})),$$

and we need to take a = -1. This finishes the classification of the family of the normal form:

$$r\begin{pmatrix} -1 & 0 & 0 & 0\\ 0 & 1 & 0 & -3\\ 0 & 0 & -\frac{1}{3} & 0\\ 0 & -3 & 0 & 3 \end{pmatrix} + s \begin{pmatrix} 0 & 0 & 1 & 0\\ 0 & 3 & 0 & -3\\ 1 & 0 & 0 & 0\\ 0 & -3 & 0 & -9 \end{pmatrix}.$$

To find the concrete normal form, we need to put  $\lambda$  in the form  $(r + s\alpha)\gamma$ , and the result is

$$r = \frac{1}{2} \left( \sqrt{\frac{-3 + \sqrt{11} + 3\sqrt{3} - \sqrt{33}}{2}} + \sqrt{\frac{-3 - \sqrt{11} - 3\sqrt{3} - \sqrt{33}}{2}} \right)$$
$$s = \frac{1}{2\sqrt{3}} \left( \sqrt{\frac{-3 + \sqrt{11} + 3\sqrt{3} - \sqrt{33}}{2}} - \sqrt{\frac{-3 - \sqrt{11} - 3\sqrt{3} - \sqrt{33}}{2}} \right)$$

This is a common pattern in this example and the following two: the family of normal forms is very simple, but the concrete normal form is much more complicated.

**Example 12.2.** Now we classify the same matrix with p = 3. 11 is still not a square in  $\mathbb{Q}_3$ , so we are in case (2) or (3) with c = -1. We write  $\lambda^2$  as  $3 + i\sqrt{-11}$ , where  $\sqrt{-11} \in \mathbb{Q}_3$ , and we need to check whether this is a square in  $\mathbb{Q}_3[i]$ . We use Proposition 6.3(1): min{ord(a), ord(b)} \equiv 0 \mod 2 and  $a^2 + b^2 = -2$  is a square in  $\mathbb{Q}_3$ , so  $\lambda^2$  is a

square in  $\mathbb{Q}_3[i]$  and we are in case (2). In this case c = -1 is the only parameter we need. This is the focus-focus family

$$\begin{pmatrix} 0 & s & 0 & r \\ s & 0 & -r & 0 \\ 0 & -r & 0 & s \\ r & 0 & s & 0 \end{pmatrix}$$

with

$$r = \frac{\sqrt{3 + \sqrt{11}} - \sqrt{3 - \sqrt{11}}}{2i}, s = \frac{\sqrt{3 + \sqrt{11}} + \sqrt{3 - \sqrt{11}}}{2}$$

**Example 12.3.** For the same matrix and p = 5, we see that  $\lambda^2 = 3 + \sqrt{11} \in \mathbb{Q}_5$ , so we are in case (1).  $c_1$  is given by the class of  $\lambda^2$  modulo a square in  $\mathbb{Q}_5$ .

$$\lambda^2 = 3 + \sqrt{11} \equiv 3 + 1 = 4 \mod 5$$

is a square, so  $c_1 = 1$  (actually,  $c_1$  is in the class of  $-\lambda^2$ , but as  $p \equiv 1 \mod 4$  it is equivalent to take  $\lambda^2$ ). Analogously

$$\mu^2 = 3 - \sqrt{11} \equiv 3 - 1 = 2 \mod 5$$

so  $c_2 = 2$  (which is  $c_0$  for p = 5) or  $c_2 = 2p^2 = 50$ . To know which one, we go to Proposition 4.6:

$$\frac{2\mu a}{u^T \Omega_0 \bar{u}} = \dots 203033 a \in \mathrm{DSq}(\mathbb{Q}_5, 2)$$

which means, by Proposition 4.7(2), that a has even order, and since  $\operatorname{ord}(ab) = 0$ ,

$$\operatorname{ord}(c_2) = \operatorname{ord}(b) - \operatorname{ord}(a) = -2 \operatorname{ord}(a)$$

is multiple of 4, so this leaves  $c_2 = 2$ . The normal form is

$$\begin{pmatrix} r & 0 & 0 & 0 \\ 0 & r & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & 2s \end{pmatrix}$$

with

$$r = \sqrt{3 + \sqrt{11}}, s = \sqrt{\frac{\sqrt{11} - 3}{2}}.$$

**Example 12.4.** Let M be the following matrix:

$$M = \begin{pmatrix} 2 & 6 & -2 & -3 \\ 6 & 11 & 1 & -5 \\ -2 & 1 & -6 & -2 \\ -3 & -5 & -2 & 3 \end{pmatrix}$$

The characteristic polynomial of  $A = \Omega_0^{-1} M$  is  $(t^2 - 5)^2$ , so this matrix has repeated eigenvalues:

$$\{\sqrt{5}, \sqrt{5}, -\sqrt{5}, -\sqrt{5}\}.$$

For p = 5,  $\sqrt{5} \notin \mathbb{Q}_5$ , hence it is in case (3) of Theorem G. We need to write  $\lambda = r\sqrt{c}$  for  $c \in Y_p$ : this leads to c = 5 and r = 1. The value of a must be 1 or 2, and from (6.1)

we obtain that a = 1 gives the result in  $DSq(\mathbb{Q}_5, -5)$ . So the normal form is

$$\begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 5 & 0 \\ 0 & 5 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

**Example 12.5.** The same matrix with p = 11: now  $\sqrt{5} \in \mathbb{Q}_{11}$ , so it is in case (2) of Theorem G, with  $r = \lambda = \sqrt{5}$ :

$$\begin{pmatrix} 0 & \sqrt{5} & 0 & 0\\ \sqrt{5} & 0 & 1 & 0\\ 0 & 1 & 0 & \sqrt{5}\\ 0 & 0 & \sqrt{5} & 0 \end{pmatrix}$$

# 12.2. Examples with functions and integrable systems.

**Example 12.6.** The matrices from Examples 12.1 and 12.4 can be translated to functions with a critical point in the origin (the first one is non-degenerate and the second one is degenerate):

$$f_1(x,\xi,y,\eta) = \frac{x^2}{2} + 2x\xi + 3xy + 4x\eta + \frac{5}{2}\xi^2 + 6\xi y + 7\xi\eta + 4y^2 + 9y\eta + 5\eta^2$$
$$f_2(x,\xi,y,\eta) = x^2 + 6x\xi - 2xy - 3x\eta + \frac{11}{2}\xi^2 + \xi y - 5\xi\eta - 3y^2 - 2y\eta + 3\eta^2$$

The class of the critical point of  $f_1$  is that of Corollary 9.9(3), with  $c = 3, t_1 = t_2 =$ -2, a = -1, b = 0, for p = 2, that of part (2) with c = -1 for p = 3, and that of part (1) with  $c_1 = 1$  and  $c_2 = 2$  for p = 5.

The class of the critical point of  $f_2$  is that of Corollary 9.10(3) with c = 5 for p = 5, and that of part (2) for p = 11.

**Example 12.7.** Let p be a prime number such that  $p \equiv 1 \mod 4$ . Then the functions

$$F_{\text{ee}}, F_{\text{eh}}, F_{\text{hh}}, F_{\text{ff}} : ((\mathbb{Q}_p)^4, \omega_0) \to (\mathbb{Q}_p)^2$$

and the symplectic form  $\omega_0 = \mathrm{d}x \wedge \mathrm{d}\xi + \mathrm{d}y \wedge \mathrm{d}\eta$  on  $\mathbb{Q}_p^4$ :

- Elliptic-elliptic:  $F_{ee}(x,\xi,y,\eta) = (\frac{x^2+\xi^2}{2},\frac{y^2+\eta^2}{2});$  Elliptic-hyperbolic:  $F_{eh}(x,\xi,y,\eta) = (\frac{x^2+\xi^2}{2},y\eta);$
- Hyperbolic-hyperbolic:  $F_{\rm hh}(x,\xi,y,\eta) = (x\xi,y\eta);$
- Focus-focus:  $F_{\rm ff}(x,\xi,y,\eta) = (x\eta y\xi, x\xi + y\eta),$

are non-degenerate p-adic analytic integrable systems. Furthermore, all four systems are *p*-adically linearly symplectomorphic. This follows from Theorem A.

**Example 12.8.** Let p be a prime number such that  $p \not\equiv 1 \mod 4$ . Then any two distinct systems among those four are not linearly symplectomorphic. This follows from Theorem А.

**Remark 12.9.** In Corollary 9.7, the elliptic function corresponds to c = 1. The hyperbolic one, by Proposition 4.1, corresponds to c = 1 for  $p \equiv 1 \mod 4$  and c = -1otherwise. That is, six of the seven forms for  $p \equiv 1 \mod 4$ , three of the five for  $p \equiv 3$ mod 4, and nine of the eleven for p = 2, have no real equivalent.

In Theorem A, the elliptic-elliptic model corresponds to (1) with  $c_1 = c_2 = 1$ . Changing elliptic components to hyperbolic results in changing the corresponding  $c_i$  to -1, except if  $p \equiv 1 \mod 4$ , where there is no change. The focus-focus model is the same for  $p \equiv 1$ 

mod 4, and otherwise it is (2) with c = -1. The vast majority of p-adic normal forms, including all those in point (3), have no real equivalent.

We now comment on the  $S_p^1$ -actions on  $\mathbb{Q}_p^2$  induced by those systems (we refer to [11, Appendix C] for a review of the concept of *p*-adic action), where we recall that  $S_p^1$  is defined by

$$S_p^1 = \Big\{ (x, y) \in \mathbb{Q}_p^2 : x^2 + y^2 = 1 \Big\}.$$

**Remark 12.10.** Let p be a prime number.

(1) The elliptic function  $f_{\rm e}(x,\xi) = \frac{x^2+\xi^2}{2}$  induces on  $\mathbb{Q}_p^2$  an  $\mathrm{S}_p^1$ -action given by

$$(u,v) \cdot (x,\xi) = \begin{pmatrix} u & v \\ -v & u \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix},$$

for  $(u, v) \in S_p^1$  and  $(x, \xi) \in \mathbb{Q}_p^2$ .

(2) The first component  $f_1(x, \xi, y, \eta) = x\eta - y\xi$  of the focus-focus system  $F_{\rm ff}$  induces on  $\mathbb{Q}_p^4$  an action with a similar formula to the previous one, simultaneously on the plane (x, y) and the plane  $(\xi, \eta)$ :

$$(u,v) \cdot (x,\xi,y,\eta) = \begin{pmatrix} u & 0 & -v & 0 \\ 0 & u & 0 & -v \\ v & 0 & u & 0 \\ 0 & v & 0 & u \end{pmatrix} \begin{pmatrix} x \\ \xi \\ y \\ \eta \end{pmatrix},$$

for  $(u, v) \in \mathcal{S}_p^1$  and  $(x, \xi, y, \eta) \in \mathbb{Q}_p^4$ .

Indeed, by [11, Corollary 4.5], there is a subgroup of  $S_p^1$  isomorphic to  $p\mathbb{Z}_p$  by the correspondence  $t \mapsto (\cos t, \sin t)$  that contains all elements near the origin. We need to prove that, if  $\psi_t(x,\xi) = (\cos t, \sin t) \cdot (x,\xi)$ , the vector field  $X_t$  of this flow (in the sense that  $\frac{d}{dt}\psi_t(x,\xi) = X_t(\psi_t(x,\xi))$ ) satisfies Hamilton's equations  $i_{X_t}\omega_0 = df_e$ .

We have

$$\frac{\mathrm{d}}{\mathrm{d}t}\psi_t(x,\xi) = \frac{\mathrm{d}}{\mathrm{d}t} \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} -\sin t & \cos t \\ -\cos t & -\sin t \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix}$$

so  $X_t(x,\xi) = (\xi, -x)$ , and

$$i_{X_t}\omega_0 = x\mathrm{d}x + \xi\mathrm{d}\xi = \mathrm{d}f_\mathrm{e},$$

as we wanted.

For the focus-focus action, we have analogously that  $X_t(x,\xi,y,\eta) = (-y,-\eta,x,\xi)$ , and

$$i_{X_t}\omega_0 = \eta \mathrm{d}x - y\mathrm{d}\xi - \xi\mathrm{d}y + x\mathrm{d}\eta = \mathrm{d}f_1,$$

as we wanted.

#### 13. Circular symmetries of the p-adic models

Here we generalize the content of Remark 12.10 and analyze the problem of existence of circle actions for arbitrary models. In the real case, for a fixed symplectic space  $(V, \omega)$ , most multiples of a Hamiltonian that admits a circle action do not admit it. This happens because a smooth circle action over V is defined by a smooth map  $h : S^1 \times V \to V$ , satisfying  $h(g_1, h(g_2, m)) = h(g_1g_2, m)$  and h(1, m) = m, for  $g_1, g_2 \in S^1$  and  $m \in V$ . Concretely, considering  $g : \mathbb{R} \to S^1$  given by  $g(t) = (\cos t, \sin t)$ , we must have

$$h(g(t + t'), m) = h(g(t)g(t'), m) = h(g(t), h(g(t'), m)$$
91

)

and h(1,m) = m, for  $t, t' \in \mathbb{R}$  and  $m \in V$ . The induced vector field for this action, that is,  $X_h(t)$  given by

$$\frac{\mathrm{d}}{\mathrm{d}t}h(g(t),m) = X_h(t)(h(g(t),m)),$$

must coincide with  $X_f$  induced by the Hamiltonian f via the Hamilton equation  $i_{X_f}\omega_0 = df$ . When f is multiplied by a constant k, that is, f' = kf, we also have  $X_{f'} = kX_f$ , so  $X_h$  must also be multiplied by k:

$$\frac{\mathrm{d}}{\mathrm{d}t}h'(g(t),m) = X_{h'}(t)(h'(g(t),m))$$
$$= kX_h(t)(h'(g(t),m))$$
$$= kX_h(kt)(h'(g(t),m))$$

(the last equality happens because  $X_h(t) = X_f$  is independent of t). This equation is solved by h'(g(t), m) = h(g(kt), m), that is, the action is accelerated by a factor of k. As we must have  $h'(g(2\pi), m) = h'(1, m) = m$ , in general this is only possible if k is integer.

This will not happen in the *p*-adic case, because the *p*-adic circle is not closed (there is no  $t \neq 0$  such that g(t) = 1). So getting a circle action is much easier in this case, and actually any "small enough" multiple of a Hamiltonian admits a circle action.

**Proposition 13.1.** Let n be a positive integer and let p be a prime number. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $(\mathbb{Q}_p)^{2n}$ . Given a p-adic analytic Hamiltonian  $f: (\mathbb{Q}_p)^{2n} \to \mathbb{Q}_p$  such that  $f(m) = m^T Mm/2$ , for a matrix  $M \in \mathcal{M}_{2n}(\mathbb{Q}_p)$ , f admits a p-adic analytic  $S_p^1$ -action (that is, there exists  $h: S_p^1 \times (\mathbb{Q}_p)^{2n} \to (\mathbb{Q}_p)^{2n}$  analytic such that  $h(g_1, h(g_2, m)) = h(g_1g_2, m)$  and h(1, m) = m, for  $g_1, g_2 \in S_p^1$  and  $m \in (\mathbb{Q}_p)^{2n}$ ) if and only if  $\operatorname{ord}(\lambda) \geq 0$  for all  $\lambda$  which is an eigenvalue of  $\Omega_0^{-1}M$ .

*Proof.* Suppose that f admits a circle action h and let  $\psi(t, v) = h(g(t), v)$  and  $A = \Omega_0^{-1} M$ . Hamilton's equation  $i_{X_f} \omega_0 = df$  results in

$$X_f(m)^T \Omega_0 v = \mathrm{d}f(m)(v) = m^T M v \Rightarrow X_f(m) = -\Omega_0^{-1} M m = -Am.$$

Substituting in the flow equation,

$$\frac{\mathrm{d}}{\mathrm{d}t}\psi(t,m) = -A\psi(t,m)$$

and  $\psi(0,m) = m$ , which solves as

$$\psi(t,m) = \exp(-tA)m$$

where exp denotes the matrix exponential. This must exist for all t in the domain of  $g(t) = (\cos t, \sin t)$ , that is, such that  $|t|_p \leq k_p$ , where  $k_p = 1/p$  for  $p \neq 2$  and  $k_2 = 1/4$ . The exponential of -tA exists if and only if the eigenvalues  $\mu$  of -tA satisfy  $|\mu|_p \leq k_p$ , which implies

$$|-t\lambda|_p = |t|_p |\lambda|_p \leqslant k_p$$

As it must exist when  $|t|_p = k_p$ , we have  $|\lambda|_p \leq 1$ .

Conversely, suppose that  $|\lambda|_p \leq 1$  for all eigenvalues of A. Then, for all t in the domain of g,

$$|-t\lambda|_p = |t|_p |\lambda|_p \leqslant k_p$$

and  $\exp(-tA)$  exists. This means that  $\psi(t,m)$  is well defined, and  $h(g_0,m)$  exists for  $g_0 \in \operatorname{Im}(g)$ . By [11, Corollary 4.5], the quotient  $\operatorname{S}_p^1/\operatorname{Im}(g)$  is a discrete group, so we can define its action arbitrarily without affecting the flow equation, and we have an action of  $\operatorname{S}_p^1$ .

**Remark 13.2.** Often symplectic classifications of group actions in equivariant symplectic geometry include the assumption that the action is effective, as it is the case for example in Delzant's classification, Duistermaat-Pelayo [19] and Pelayo [44]. With this restriction (being effective), no multiple of a Hamiltonian which admits a circle action also admits an action. However, as Proposition 13.1 shows, in the *p*-adic case all small multiples of any Hamiltonian admit an effective action.

However, if we want the action to have the form h(g, m) = h((u, v), m) = (uI + vB)m, like the actions in Remark 12.10, the situation changes completely.

**Proposition 13.3.** Let n be a positive integer and let p be a prime number. Let  $\Omega_0$  be the matrix of the standard symplectic form on  $(\mathbb{Q}_p)^{2n}$ . Given a p-adic analytic Hamiltonian  $f: (\mathbb{Q}_p)^{2n} \to \mathbb{Q}_p$  such that  $f(m) = m^T Mm/2$ , for a matrix  $M \in \mathcal{M}_{2n}(\mathbb{Q}_p)$ , f admits a p-adic analytic  $S_p^1$ -action of the form h((u, v), m) = (uI + vB)m if and only if the eigenvalues of  $\Omega_0^{-1}M$  are i and -i, both with multiplicity n. In that case,  $B = -\Omega_0^{-1}M$ .

Proof. The flow equation implies

$$\frac{\mathrm{d}}{\mathrm{d}t}h((\cos t,\sin t),m) = -Ah((\cos t,\sin t),m),$$

that is

$$\frac{\mathrm{d}}{\mathrm{d}t}(\cos tI + \sin tB)m = (-\sin tI + \cos tB)m = -A(\cos tI + \sin tB)m,$$

which implies AB = I and B = -A. Hence, the action exists if and only if  $A^2 = -I$ . If this happens, the only possible eigenvalues are i and -i, and since they must come in opposite pairs, each one appears n times. Conversely, if the eigenvalues of A are i and -i, those of  $A^2$  are -1, which implies that  $A^2 = -I$ .

Of the normal forms in Theorem A, those with a circle action of this form are the following:

- At point (1), only the elliptic component gives eigenvalues i and -i. The remaining ones have different eigenvalues. Hence, we recover Remark 12.10(1).
- At point (2), only the focus-focus component, if  $p \not\equiv 1 \mod 4$ , has eigenvalues (i, i, -i, -i). We recover Remark 12.10(2).
- At point (3),  $t_1 + t_2\sqrt{c}$  is the square of an eigenvalue, so we must have  $t_1 = -1$  and  $t_2 = 0$  to have this kind of circle action.

# Appendix A. Basic properties of p-adic numbers

A.1. Definition of  $\mathbb{Q}_p$ . The field  $\mathbb{R}$  is defined as a completion of  $\mathbb{Q}$  with respect to the normal absolute value, that is,  $|x| = \max\{x, -x\}$ . Analogously, we define the *p*-adic absolute value as  $|x|_p = p^{-\operatorname{ord}_p(x)}$ , where  $\operatorname{ord}_p(x)$  is defined as the highest power of *p* dividing *x*, if *x* is integer, and as  $\operatorname{ord}_p(m) - \operatorname{ord}_p(n)$ , if x = m/n is rational.

**Definition A.1** ([28, Theorem 3.2.13]).  $\mathbb{Q}_p$  is defined as the completion of  $\mathbb{Q}$  with respect to the absolute value  $|\cdot|_p$ . This means that

- (1)  $\mathbb{Q}$  is a subset of  $\mathbb{Q}_p$ , and the absolute value  $|\cdot|_p$  on  $\mathbb{Q}$  extends to  $\mathbb{Q}_p$ .
- (2)  $\mathbb{Q}$  is dense in  $\mathbb{Q}_p$  with respect to the absolute value  $|\cdot|_p$ .
- (3)  $\mathbb{Q}_p$  is complete with respect to the absolute value  $|\cdot|_p$ .

The field  $\mathbb{Q}_p$  satisfying (1), (2) and (3) is unique up to isomorphism of fields preserving the absolute values.

The ring of *p*-adic integers is the topological closure of  $\mathbb{Z}$  in  $\mathbb{Q}_p$ , which coincides with the ball centered in 0 of radius 1 [28, Definition 3.3.3]:

$$\mathbb{Z}_p = \Big\{ x \in \mathbb{Q}_p \mid |x|_p \leqslant 1 \Big\}.$$

By [28, Proposition 3.3.4], for any  $a \in \mathbb{Q}_p$ , there exists a sequence  $(a_n)_{n \ge n_0}$  with  $n_0 \in \mathbb{Z}$ and  $0 \le a_n \le p-1$  such that

$$a = \sum_{n=n_0}^{\infty} a_n p^n.$$

This  $n_0$  is denoted as  $\operatorname{ord}_p(a)$ . If  $a \in \mathbb{Q}$ , it coincides with the previous definition of  $\operatorname{ord}_p(a)$ .

We also need some properties about squares in  $\mathbb{Q}_p$ . We start with a result known as Hensel's lifting:

**Theorem A.2** (Hensel's lifting, [28, Theorem 3.4.1 and Problem 112]). Let f be a polynomial in  $\mathbb{Z}_p[x]$ . Let  $\alpha_1$  be a p-adic integer,  $r = \operatorname{ord}(f(\alpha_1))$  and  $s = \operatorname{ord}(f'(\alpha_1))$ . If r > 2s, there exists  $\alpha \in \mathbb{Z}_p$  such that  $\operatorname{ord}(\alpha - \alpha_1) \ge r - s$  and  $f(\alpha) = 0$ .

A consequence which is useful for us is the following:

**Corollary A.3.** For  $a, b \in \mathbb{Z}_2$ , such that  $2 \nmid a, b, a \equiv b \mod 2^n$  or  $a \equiv -b \mod 2^n$  if and only if  $a^2 \equiv b^2 \mod 2^{n+1}$ .

*Proof.* If  $a \equiv \pm b \mod 2^n$ ,  $a = \pm b + 2^n t$  for some  $t \in \mathbb{Z}_2$ , and

$$a^2 = b^2 \pm 2^{n+1}bt + 2^{2n}t^2 \equiv b^2 \mod 2^{n+1}.$$

Suppose now that  $a^2 \equiv b^2 \mod 2^{n+1}$ . We apply Hensel's lifting to  $f(x) = x^2 - a^2$  and  $\alpha_1 = b$ . We have

$$r = \operatorname{ord}(b^2 - a^2) \ge n + 1$$

and

$$s = \operatorname{ord}(2b) = 1,$$

so there is  $\alpha$  with  $\operatorname{ord}(\alpha-b) \ge n$  and  $\alpha^2 - a^2 = 0$ . This implies  $\alpha = \pm a$ , so  $\operatorname{ord}(\pm a - b) \ge n$ , as we wanted.

Another consequence is the characterization of squares in  $\mathbb{Q}_p$ :

- **Corollary A.4.** (1) If  $p \neq 2$ ,  $a \in \mathbb{Q}_p$  is a square if and only if  $\operatorname{ord}(a)$  is even and the digit of x at the position  $\operatorname{ord}(a)$  is a square modulo p.
  - (2) If p = 2,  $a \in \mathbb{Q}_p$  is a square if and only if  $\operatorname{ord}(a)$  is even and a ends in 001 (that is,  $a/2^{\operatorname{ord}(a)} \equiv 1 \mod 8$ ).
- *Proof.* (1) For the implication to the right: if  $a = b^2$ ,  $\operatorname{ord}(a) = 2 \operatorname{ord}(b)$ , and the leading digit of a is the square of that of b modulo p.

For the implication to the left: If  $\operatorname{ord}(a)$  is even and the leading digit is square, let  $a_0 = a/p^{\operatorname{ord}(a)}$ . Let  $c \in \mathbb{Z}$  such that  $c^2 \equiv a_0 \mod p$ . We apply Hensel's lifting to  $x^2 - a_0$  with  $\alpha = c$ . We have  $r = \operatorname{ord}(c^2 - a_0) > 0$ , because the leading digits of  $c^2$  and  $a_0$  coincide, and  $s = \operatorname{ord}(2c) = 0$ . Then, there is  $b_0 \in \mathbb{Z}_p$  such that  $b_0^2 = a_0$ . Taking  $b = b_0 p^{\operatorname{ord}(a)/2}$ , we have  $b^2 = a_0 p^{\operatorname{ord}(a)} = a$ .

(2) If  $a = b^2$ , we have again  $\operatorname{ord}(a) = 2 \operatorname{ord}(b)$ . Let  $t = \operatorname{ord}(b)$ . Then  $b = 2^t(1+2c)$  for some  $c \in \mathbb{Z}_2$  and

$$a = b^{2} = 2^{2t}(1 + 4c + 4c^{2}) = 2^{2t} + 2^{2t+2}c(c+1)$$

As c is integer, c(c+1) is even, and this ends in 001.

If  $\operatorname{ord}(a)$  is even and a ends in 001, let  $a_0 = a/2^{\operatorname{ord}(a)}$ . We apply Hensel's lifting to  $x^2 - a_0$  with  $\alpha = 1$ . We have  $r = \operatorname{ord}(1 - a_0) \ge 3$ ,  $s = \operatorname{ord}(2) = 1$ , and there is  $b_0 \in \mathbb{Z}_p$  such that  $b_0^2 = a_0$ . Taking  $b = b_0 2^{\operatorname{ord}(a)/2}$ , we have  $b^2 = a_0 2^{\operatorname{ord}(a)} = a$ .  $\Box$ 

A.2. Critical points of *p*-adic analytic functions. A *power series* in  $(\mathbb{Q}_p)^n$  is given by

$$f(x) = \sum_{I \in \mathbb{N}^n} a_I (x - x_0)^I$$

where  $x^{I}$  means  $x_{1}^{i_{1}} \dots x_{n}^{i_{n}}$  and  $a_{I}$  are coefficients in  $\mathbb{Q}_{p}$ . A function

$$f:(\mathbb{Q}_p)^n\to\mathbb{Q}_p$$

is (*p*-adic) analytic if it can be expressed as a power series. (Sometimes the term is used for functions that are expressed as a power series only piecewise, but for our purposes here this distinction does not matter because we are interested only in the local behavior of analytic functions.)

We define df(m) as the vector whose coordinates are the partial derivatives of f, defined in the usual sense:

$$\mathrm{d}f(m)_i = \frac{\partial f(m)}{\partial x_i} := \lim_{t \to 0} \frac{f(x + t\mathrm{e}_i) - f(x)}{t}$$

where the definition of limit is the usual one from metric spaces (the limit of f at a point  $x_0 \in \mathbb{Q}_p^n$  is equal to  $y_0 \in \mathbb{Q}_p$  if for any  $\epsilon > 0$  there is  $\delta > 0$  such that  $|f(x) - f(x_0)|_p < \epsilon$  whenever  $|x_i - (x_0)_i|_p < \delta$  for  $1 \leq i \leq n$ ). This limit always exists for a power series, and it is equal to

$$\frac{\partial f}{\partial x_i} = \sum_{I \in \mathbb{N}^n} a_I i_j (x - x_0)^{I_j},$$

where, for  $I = (i_1, ..., i_n)$ ,  $I_j$  is defined as  $(i_1, ..., i_j - 1, ..., i_n)$ .

In the same way, we define the Hessian of f, which we denote as  $d^2f$ , the matrix with the second derivatives of f as entries:

$$(\mathrm{d}^2 f)_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}.$$

In general, if M is a p-adic analytic manifold, there are charts  $\phi$  from  $U \subset (\mathbb{Q}_p)^n$  to  $V \subset M$  for some n, and we can define *analytic function* in V as a function  $f: V \to \mathbb{Q}_p$  such that  $f \circ \phi$  is given by a power series, or more generally *analytic function* in M as  $f: M \to \mathbb{Q}_p$  given piecewise by power series. The differential of f is defined as  $df = d(f \circ \phi)$ , and its Hessian is  $d^2 f = d^2(f \circ \phi)$ . See Appendix B of [11] for the precise definitions of p-adic manifolds and analytic functions.

A critical point of  $f: M \to \mathbb{Q}_p$  is a point  $m \in M$  such that df(m) = 0.

Acknowledgments. We thank Enrique Arrondo for discussions and suggestions which have improved the paper.

The first author is funded by grants PID2019-106188GB-I00 and PID2022-137283NB-C21 of MCIN/AEI/10.13039/501100011033 / FEDER, UE and by project CLaPPo (21.SI03.64658) of Universidad de Cantabria and Banco Santander.

The second author is funded by a FBBVA (Bank Bilbao Vizcaya Argentaria Foundation) Grant for Scientific Research Projects with title *From Integrability to Randomness in Symplectic and Quantum Geometry.* He thanks the Dean of the School of Mathematical Sciences Antonio Brú and the Chair of the Department of Algebra, Geometry and Topology at the Complutense University of Madrid, Rutwig Campoamor, for their support and excellent resources he is being provided with to carry out the FBBVA project.

Part of this paper was written during a visit of the second author to the Universidad de Cantabria and the Universidad Internacional Menéndez Pelayo in the summer of 2024 and he thanks both institutions for their hospitality. Another part of this paper was written while the first author was visiting the Complutense University of Madrid in September-October 2024 and he thanks this institution.

#### References

- R. Abraham, J.E. Marsden: Foundation of Mechanics, Second edition, revised and enlarged. With the assistance of Tudor Ratiu and Richard Cushman. Benjamin/Cummings Publishing Co., Inc., Advanced Book Program, Reading, Mass., 1978.
- [2] M. Alsina, P. Bayer: Quaternion Orders, Quadratic Forms, and Shimura Curves. American Mathematical Society, CRM Monograph Series 22, 2004.
- [3] V. Avetisov, A. Bikulov: Protein ultrametricity and spectral diffusion in deeply frozen proteins. Biophys. Rev. Lett. 3 (2008), 387–396.
- [4] S. Awodey, A. Pelayo, M. A. Warren: Voevodsky's univalence axiom in homotopy type theory. Notices Amer. Math. Soc. 60(9) (2013), 1164–1167.
- [5] S. Bates, A. Weinstein: Lectures on the Geometry of Quantization. Berkeley Mathematics Lecture Notes 8, AMS-CPAM, 1997.
- [6] G. Bazzoni, V. Muñoz: Manifolds which are complex and symplectic but not Kähler. Essays in Mathematics and its Applications, Springer, 2016, 49–69.
- M. Bhargava: Higher composition laws and applications. Proceedings of the International Congress of Mathematicians Madrid, August 22–30, 2006, Volume II. European Mathematical Society, 2007, 271–294.
- [8] A. V. Bolsinov, A. T. Fomenko: Integrable Hamiltonian Systems: Geometry, Topology, Classification, CRC Press, 2004.
- [9] A. Cannas da Silva: Lectures on Symplectic Geometry. Lecture Notes in Math., 1764, Springer-Verlag, Berlin, 2001.
- [10] B. Casselman: Introduction to quadratic forms. Available at https://personal.math.ubc.ca/ cass/research/pdf/QForms.pdf
- [11] L. Crespo, A. Pelayo: The p-adic Jaynes-Cummings model in symplectic geometry, preprint, arXiv:2406.18415.
- [12] M. A. de Gosson: Gaussian quantum states can be disentangled using symplectic rotations. Lect. Math. Phys. 111 (2021), article 73.
- [13] M. de León, P. R. Rodrigues: Methods of Differential Geometry in Analytical Mechanics, North-Holland Mathematics Studies, 158, North-Holland, Amsterdam, 1989.
- [14] T. Delzant: Hamiltoniens périodiques et images convexes de l'application moment. Bull. Soc. Math. France 116(3) (1988), 315–339.
- [15] B. Dragovich: p-Adic and Adelic Quantum Mechanics. Proceedings of the First Int. Conf. on p-Adic Mathematical Physics, 1-5 October 2003, Moscow.
- B. Dragovich: Adelic harmonic oscillator. International Journal of Modern Physics A 10(16) (1995), 2349–2365.
- [17] B. Dragovich, A. Yu. Khrennikov, S. V. Kozyrev, I. V. Volovich: On p-adic mathematical physics. P-Adic Numbers, Ultrametric Analysis, and Applications 1 (2009), 1–17.
- [18] B. Dragovich, N. Ž. Mišić: *p*-adic hierarchical properties of the genetic code. *Biosystems* (2019).
- [19] J.J. Duistermaat, A. Pelayo: Symplectic torus actions with coisotropic principal orbits, Annales de l'Institut Fourier 57 (2007), 2239–2327.
- [20] J.J. Duistermaat, A. Pelayo: Reduced phase space and toric variety coordinatizations of Delzant spaces. Math. Proc. Cambridge Philos. Soc. 146(3) (2009), 695–718.
- [21] J.J. Duistermaat, A. Pelayo: Complex structures on four-manifolds with symplectic two-torus actions. Internat. J. Math. 22(3) (2011), 449–463.
- [22] H. R. Dullin, A. Pelayo: Generating hyperbolic singularities in semitoric systems via Hopf bifurcations. J. Nonlinear Sci. 26 (2016), 787–811.

- [23] K. Efstathiou, S. Hohloch, P. Santos: On the affine invariant of hypersemitoric systems. Preprint 2024, arXiv:2411.17509.
- [24] Y. Eliashberg: Recent advances in symplectic flexibility. Bull. Amer. Math. Soc. 52(1) (2015), 1–26.
- [25] L. H. Eliasson: Hamiltonian Systems with Poisson Commuting Integrals, PhD thesis, University of Stockholm, 1984.
- [26] L. H. Eliasson: Normal forms for Hamiltonian systems with Poisson commuting integrals elliptic case. Comment. Math. Helv. 65 (1990), 4–35.
- [27] M. Fernández, M. G. Gotay, A. Gray: Compact parallelizable four-dimensional symplectic and complex manifolds. Proc. Amer. Math. Soc. 103(4) (1988), 1209–1212.
- [28] F. Q. Gouvea: *p-adic Numbers*, Springer-Verlag 1993.
- [29] S. S. Gubser, J. Knaute, S. Parikh, A. Samberg, P. Witaszczyk: *p*-adic AdS/CFT. Comm. Math. Phys. 352 (2017), 1019–1059.
- [30] G. H. Hardy, S. Ramanujan: Asymptotic formulae in combinatory analysis. Proc. LMS, Second Series, 17, 75–115. Reprinted in Collected papers of Srinivasa Ramanujan, Amer. Math. Soc. (2000), 276–309.
- [31] H. Hofer, E. Zehnder: Symplectic invariants and Hamiltonian dynamics, Birkhäuser 1994.
- [32] S. Hohloch, J. Palmer, Extending compact Hamiltonian S<sup>1</sup>-spaces to integrable systems with mild degeneracies in dimension four. Preprint 2021, arXiv:2105.00523.
- [33] H. Hua, L. Hovestadt: p-adic numbers encode complex networks. Sci. Rep. 11(17) (2021).
- [34] A. Khrennikov, K. Oleschko, M. d. J. C. López: Applications of p-adic wavelets to model reactiondiffusion dynamics in random porous media. J. Fourier Anal. Appl. 22 (2016), 809–822.
- [35] K. Kodaira: On the structure of compact complex analytic surfaces, I. Amer. J. Math. 86(4) (1964), 751–798.
- [36] T. Y. Lam: Introduction to Quadratic Forms over Fields, American Mathematical Society, Graduate Studies in Mathematics 67, 2005.
- [37] Y. Le Floch, A. Pelayo: Symplectic geometry and spectral properties of classical and quantum coupled angular momenta. J. Nonlinear Sci. 29 (2016), 655–708.
- [38] J. Lurie: What is... *p*-adic geometry? Members Colloquium at IAS Princeton, October 9, 2023. Link: https://www.ias.edu/video/what-p-adic-geometry
- [39] J. E. Marsden, T. S. Ratiu: Introduction to Mechanics and Symmetry. Texts in Applied Mathematics, 17, Springer 1999.
- [40] D. McDuff, D. Salamon: Introduction to Symplectic Topology. Oxf. Grad. Texts Math. Oxford University Press, Oxford, 2017.
- [41] O. T. O'Meara: Introduction to Quadratic Forms, Springer 2013.
- [42] J.-P. Ortega, T. S. Ratiu: Momentum Maps and Hamiltonian Reduction, Progress in Mathematics, 222, Springer 2004.
- [43] J. Palmer, A. Pelayo, X. Tang: Semitoric systems of non-simple type, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas 118, 161 (2024).
- [44] A. Pelayo: Symplectic actions of 2-tori on 4-manifolds, Memoirs of the American Mathematical Society 204 (2010) no. 959, viii+ 91p.
- [45] A. Pelayo: Hamiltonian and symplectic symmetries: an introduction. Bull. Amer. Math. Soc. 54(3) (2017), 383–436.
- [46] Á. Pelayo, V. Voevodsky, M. A. Warren: A univalent formalization of the p-adic numbers. Math. Struct. in Comp. Science 25 (2015), 1147–1171.
- [47] Á. Pelayo, S. Vũ Ngọc: Semitoric integrable systems on symplectic 4-manifolds. Invent. Math. 177 (2009), 571–597.
- [48] Á. Pelayo, S. Vũ Ngọc: Constructing integrable systems of semitoric type. Acta Math. 206 (2011), 93–125.
- [49] A. Pelayo, S. Vũ Ngọc: Symplectic theory of completely integrable Hamiltonian systems. Bull. Amer. Math. Soc. 48(3) (2011), 409–455.
- [50] Á. Pelayo, S. Vũ Ngọc: Hamiltonian dynamics and spectral theory for spin-oscillators. Comm. Math. Phys. 309 (2012) 123–154.
- [51] Á. Pelayo, M. A. Warren: Homotopy type theory. Gazette des Mathématiciens Société Mathématique de France 42 (2014), 95–110.
- [52] A. Pelayo, M. A. Warren: Homotopy type theory and Voevodsky's univalent foundations. Bull. Amer. Math. Soc. 51 (2014), 597–648.

- [53] H. Rüssmann: Über das Verhalten analytischer Hamiltonscher Differentialgleichungen in der Nähe einer Gleichgewichtlösung. Math. Ann., 154 (1964), 285–300.
- [54] P. Schneider: *p-adic Lie Groups*, Springer-Verlag 2011.
- [55] P. Scholze, J. Weinstein: Berkeley lectures in p-adic geometry. Annals of Mathematics Studies, book 389.
- [56] R. Simon, E. C. G. Sudarshan, N. Mukunda: Gaussian pure states in quantum mechanics and the symplectic group. *Phys. Rev. A* 37 (1988), article 3028.
- [57] W. Thurston: Some simple examples of symplectic manifolds. Proc. Amer. Math. Soc. 55(2) (1976), 467–468. Appeared in Collected works of William P. Thurston with commentary. Vol. I. Foliations, surfaces and differential geometry, American Mathematical Society, Providence, RI, 2022, 617–618.
- [58] J. V. Uspensky: Les expressions asymptotiques des fonctions numériques, coïncidant dans les problèmes de dislocation des nombres en composés, Bulletin de l'Académie des Sciences de Russie. VI série, 14 (1920), 199–218.
- [59] J. Vey: Sur certains systèmes dynamiques séparables. Amer. J. Math., 100 (1978), 591-614.
- [60] V. S. Vladimirov, I. V. Volovich: p-adic quantum mechanics. Comm. Math. Phys. 123 (1989), 659–676.
- [61] S. Vũ Ngọc, C. Wacheux: Smooth normal forms for integrable hamiltonian systems near a focusfocus singularity. arXiv:1103.3282.
- [62] X.-B. Wang, T. Hiroshima, A. Tomita, M. Hayashi: Quantum information with Gaussian states. *Phys. Rep.* 448(1) (2007), 1–111.
- [63] K. Weierstrass: Mathematische Werke, 1858, Berlin, Band I: 233–246, Band II: 19–44, Nachtrag: 139–148.
- [64] A. Weinstein: Symplectic geometry. Bull. Amer. Math. Soc. 5(1) (1981), 1–13.
- [65] A. Weinstein: Poisson geometry. Diff. Geom. Appl. 9 (1998), 213–238.
- [66] J. Williamson: On the algebraic problem concerning the normal forms of linear dynamical systems. American Journal of Mathematics 58(1) (1936), 141–163.
- [67] N. T. Zung: A note on degenerate corank-one singularities of integrable Hamiltonian systems. Commentarii Mathematici Helvetici 75(2) (2000), 271–283.
- [68] N. T. Zung: Another note on focus-focus singularities. Letters Math. Phys. 60(1) (2002), 87–99.

LUIS CRESPO, DEPARTAMENTO DE MATEMÁTICAS, ESTADÍSTICA Y COMPUTACIÓN, UNIVERSIDAD DE CANTABRIA, AV. DE LOS CASTROS 48, 39005 SANTANDER, SPAIN *Email address*: luis.cresporuiz@unican.es

ÁLVARO PELAYO, FACULTAD DE CIENCIAS MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, 28040 MADRID, SPAIN, AND REAL ACADEMIA DE CIENCIAS EXACTAS, FÍSICAS Y NATURALES, SPAIN Email address: alvpel01@ucm.es