#### A COUNTABLE BOOLEAN ALGEBRA THAT IS REICHENBACH'S COMMON CAUSE COMPLETE

DOMINIKA BUREŠOVÁ

ABSTRACT. The common cause completeness (CCC) is a philosophical principle that asserts that if we consider two positively correlated events then it evokes a common cause. The principle is due to H. Reichenbach and has been largely studied in Boolean algebras and elsewhere. The results published so far bring about a question whether there is a small (countable) Boolean algebra with CCC. In this note we construct such an example.

# 1 Introduction

This note contributes to the previous work [1, 3, 5, 6]. In the investigation so far, the authors found that the CCC is guaranteed if the Boolean algebra is  $\sigma$ -complete, has the Darboux property, and the probability measure on it is non-atomic [3]. Of course, one would like to construct as simple CCC example as possible. We present a countable example showing that the above conditions are not necessary and the published examples of CCC are far from being the simplest.

## 2 Basic notions

The basic notions as well as their philosophical meaning are taken from [3, 6]. Standardly, the symbol  $p(a \mid b) = \frac{p(a \land b)}{p(b)}$  denotes the conditional probability.

**Definition 2.1.** Let a, b in B. We define  $cov_p(a, b) = p(a \wedge b) - p(a) \cdot p(b)$ .

**Definition 2.2.** Let B be a Boolean algebra and let p be a finitely additive probability measure on it. Suppose that a, b, c are distinct elements of B. Then we say that c is a *common cause* of a, b if the following four conditions

Department of Cybernetics, Faculty of Electrical Engineering, Czech Technical University in Prague, Czech Republic, *e-mail*: buresdo2@fel.cvut.cz.

are fulfilled:

(RCCP1)  $p(a \wedge b \mid c) = p(a \mid c) p(b \mid c),$ 

(RCCP2)  $p(a \land b \mid c') = p(a \mid c') p(b \mid c'),$ 

 $(\text{RCCP3}) \qquad \qquad p(a \mid c) > p(a \mid c'),$ 

$$(\text{RCCP4}) \qquad \qquad p(b \mid c) > p(b \mid c').$$

If each pair  $a, b \in B$  with positive covariance (i.e., positive correlation) has a common cause c, we say that (B, p) is common cause complete (CCC).

Recall that if a, b have a common cause then their correlation  $cov_p(a, b)$  has to be positive [6]. Observe that if  $cov_p(a, b) > 0$  then p(a) < 1 and p(b) < 1.

**Definition 2.3.** If there is no positively correlated pair of events, then conditions RCCP1, RCCP2, RCCP3, RCCP4 are automatically satisfied. Such Boolean algebras are called **trivially common cause complete** [3].

### 3 Results

The technical part of our study slightly overlaps with [3]. We simplified and clarified its formulations for the purpose of finding the most direct way in our construction.

**Proposition 3.1.** Let B be a Boolean algebra and p be a finitely additive probability measure on B. Let  $a, b, c \in B$ ,  $c \leq a \wedge b$ ,  $\operatorname{cov}_p(a, b) > 0$ , and p(c) > 0. Then a, b, c fulfill the conditions RCCP1, RCCP3 and RCCP4.

*Proof.* Since  $c \leq a \wedge b$  then the condition RCCP1 reduces on 1 = 1. Next, the assumptions we are to work with are  $\operatorname{cov}_p(a, b) > 0$ , p(a) < 1, p(b) < 1,  $c \leq a \wedge b \leq a$ . In order to verify RCCP3, it suffices to realize that  $\frac{p(a \wedge c)}{p(c)} = \frac{p(c)}{p(c')} = 1$  whereas  $\frac{p(a \wedge c')}{p(c')} = \frac{p(c') - p(a')}{p(c')} < 1$ . The condition RCCP4 is proved analogously.

**Proposition 3.2.** Let *B* a Boolean algebra and let *p* be a probability measure on *B*. Let  $cov_p(a, b) > 0$ . Then  $0 < \frac{p(a \land b) - p(a) \cdot p(b)}{1 + p(a \land b) - p(a) - p(b)} \le p(a \land b)$ .

Proof. Since  $(1 - p(a)) \cdot (1 - p(b)) > 0$ , we see that  $p(a \wedge b) - p(a) \cdot p(b) < 1 + p(a \wedge b) - p(a) - p(b)$ 

 $\mathbf{2}$ 

and therefore

$$\frac{p(a \wedge b) - p(a) \cdot p(b)}{1 + p(a \wedge b) - p(a) - p(b)} > 0.$$

In order to prove the other inequality, observe first the inequality of basic probability theory,

$$p(a) \cdot p(b) \ge p(a \lor b) \cdot p(a \land b).$$

When we adjust the desired inequality  $\frac{p(a \wedge b) - p(a) \cdot p(b)}{1 + p(a \wedge b) - p(a) - p(b)} \leq p(a \wedge b)$ , by multiplying with the denominator we obtain

 $p(a \wedge b) + p(a \wedge b)^2 - p(a \wedge b) \cdot p(a) - p(a \wedge b) \wedge p(b) \ge p(a \wedge b) - p(a) \cdot p(b).$ 

Manipulating with this inequality, we obtain

$$p(a \wedge b)^2 - p(a \wedge b) \cdot p(a) - p(a \wedge b) \cdot p(b) + p(a) \cdot p(b) \ge 0.$$

This is equivalent to

$$(p(a) - p(a \wedge b)) \cdot (p(b) - p(a \wedge b)) \ge 0.$$

which is obviously true. The proof is complete.

Note 3.3. The inequality in Proposition 3.2 was dealt with in the book [3], however the proof presented there was erroneous in the fundamental statement (4.92). We present a correct and new proof.

**Proposition 3.4.** Let  $cov_p(a,b) > 0, c \le a \land b$ . If c is such that p(c) = $\frac{p(a\wedge b)-p(a)\cdot p(b)}{1+p(a\wedge b)-p(a)-p(b)} \ then \ c \ is \ the \ common \ cause \ of \ a, b.$ 

*Proof.* We have to check RCCP2. Our assumption is

$$p(c) - p(c) \cdot p(a) - p(c) \cdot p(b) + p(c) \cdot p(a \wedge b) = p(a \wedge b) - p(a) \cdot p(b).$$

Adding  $p(c)^2$  to both sides of the equation and reorganizing it, we get:

$$p(c)^2 + p(a) \cdot p(b) - p(a) \cdot p(c) - p(b) \cdot p(c)$$
  
=  $p(c)^2 + p(a \wedge b) - p(c) - p(c) \cdot p(a \wedge b)$ .

This implies that

$$\underbrace{\left(p(a) - p(c)\right)}_{p(a\wedge c')} \cdot \underbrace{\left(p(b) - p(c)\right)}_{p(b\wedge c')} = \underbrace{\left(1 - p(c)\right)}_{p(c')} \cdot \underbrace{\left(p(a \wedge b) - p(c)\right)}_{p(a\wedge b\wedge c')}$$
  
s is RCCP2.

and this is RCCP2.

Let us note that finite Boolean algebras can be only trivially common cause complete (Definition 2.3). Hence a meaningful case for looking for the CCC begins with a countable cardinality of B.

**Proposition 3.5.** There is a countable Boolean algebra B with a finitely additive probability measure p such that (B, p) is non-trivially common cause complete.

#### D. BUREŠOVÁ

**Construction 3.6.** Take the interval (0, 1] and the collection of finite disjoint unions of sub-intervals of (0, 1] that have rational endpoints and that are left-open and right-closed. This collection forms a Boolean algebra. Indeed, this collection is closed under the formation of unions, intersections, and complements in (0, 1]. Also, this Boolean algebra is countable since it has the same cardinality as  $\bigcup_{i \in \mathbb{N}} \mathbb{Q}^i$  ( $\mathbb{Q}$  denotes the rational numbers of (0, 1]). Take the probability measure p as the restriction of the Lebesgue measure to B. By the construction, p is common cause complete. Indeed, consider the value  $v = \frac{p(a \wedge b) - p(a) \cdot p(b)}{1 + p(a \wedge b) - p(a) - p(b)} > 0$ . So v is a rational number with  $v \leq p(a \wedge b)$ . By the definition of v and p it is easily seen that there is an element c such that  $c \leq a \vee b$  and p(c) = v. This completes the proof.

Let us shortly comment on the result and its proof. Firstly, the same method gives us also an uncountable Boolean algebra that is non-trivially common cause complete. Indeed, the classic results of the division ring theory assert that there is a division ring  $\tilde{R}$  in the real numbers  $\mathbb{R}$  and  $\tilde{R}$  is uncountable and dense in  $\mathbb{R}$ . In analogy with Construction 3.5, one takes for B the collection of disjoint unions of intervals in  $\tilde{R} \cap (0, 1]$ . It suffices to use the Lebesgue measure again.

Secondly, though we mainly addressed the Boolean line of CCC, if we stepped into quantum logics (see [3, 2] for the adequate notions on quantum logics), we see that the situation is analogous there (including the fact that finite quantum logics cannot be non-trivially CCC). We can easily construct "finitely additive" non-trivially CCC quantum logics, too. We achieve this goal by a construction called a horizontal sum (see [4] for the adequate notion in quantum logics) of arbitrarily many copies of the example constructed in Proposition 3.5. Hence in quantum logics we can easily use our Proposition 3.5 for constructing examples of an arbitrarily large (infinite) cardinality.

Acknowledgements. The author would like to thank her colleagues Pavel Pták, Mirko Navara, and Miroslav Korbelář for professional suggestions that improved the text.

This work was supported by the CTU institutional support RVO13000.

Data availability statement. The manuscript has no associated data.

**Conflict of interest statement.** The author states that there is no conflict of interest.

#### References

 Burešová, D., Houšková, K., Navara, M., Pták, P., Slouka, M., and Ševic, J.: Reichenbach's causal completeness of quantum probability spaces. Submitted.

- [2] Gudder, S.P.: Stochastic Methods in Quantum Mechanics. North Holland, New York (1979).
- [3] Hofer-Szabó, G., Rédei, M., and Szabó, L.E.: The Principle of the Common Cause. Cambridge University Press, Cambridge (2013).
- [4] Kalmbach, G.: Orthomodular Lattices. Academic Press, London (1983).
- [5] Kitajima, Y.: Reichenbach's common cause in an atomless and complete orthomodular lattice. *Internat. J. Theoret. Phys.* 47 (2008), 511–519. DOI 10.1007/s10773-007-9475-2
- [6] Reichenbach, H.: The Direction of Time. University of California Press, Berkeley (1956).