Super-Hamiltonians for super-Macdonald polynomials

Dmitry Galakhov^{1,2,3†}, Alexei Morozov^{1,2,3‡} and Nikita Tselousov^{1,2,3§}

¹MIPT, 141701, Dolgoprudny, Russia ²NRC "Kurchatov Institute", 123182, Moscow, Russia ³IITP RAS, 127051, Moscow, Russia ⁴ITEP, Moscow, Russia

ABSTRACT

The Macdonald finite-difference Hamiltonian is lifted to a super-generalization. In addition to canonical bosonic time variables p_k new Grassmann time variables θ_k are introduced, and the Hamiltonian is represented as a differential operator acting on a space of functions of both types of variables p_k and θ_k . Eigenfunctions for this Hamiltonian are a suitable generalization of Macdonald polynomials to super-Macdonald polynomials discussed earlier in the literature. Peculiarities of the construction in comparison to the canonical bosonic case are discussed.

1 Introduction

The extension of Macdonald polynomials [1] to the super-case [2,3] and [4,5] appears to be applicable to studies of super-Yangians like $Y(\widehat{\mathfrak{gl}}_{1,1})$ [6–8] and their super-DIM generalizations [5,9,10]. Surely, a modification (introduction of Grassmann variables) of the original Macdonald construction sheds some light on peculiarities of the original construction itself: we might try to question whether an appearance of algebraic and combinatorial structures accompanying Macdonald polynomials is accidental and may be naturally extended to other similar constructions.

In approaches used in [2,4,5] the super-generalization was based on a variety of intuitive arguments. among which the major role was played by a generalization of Cauchy formulas and triangularity property. This is the old idea behind the definition of the far more general Kerov functions [11–14], where Macdonalds are distinguished by the independence of the ordering of Young diagrams – which makes triangularity a well-defined notion.

However, the most conventional approach defines Macdonald polynomials as eigenfunctions of Ruijsenaars Hamiltonians [15–17]. In this paper we also exploit this idea and construct a set of commuting super-Hamiltonians whose eigenfunctions turn out to coincide with canonically defined

 $^{^{\}dagger}\text{e-mail: galakhov@itep.ru}$

[‡]e-mail: morozov@itep.ru

[§]e-mail: tselousov.ns@phystech.edu

super-Macdonald polynomials. We discuss special properties of these Hamiltonians that are relevant for the representation theories of infinite dimensional algebras – such as Yangians [18–22] and DIM [23–28] algebras where these operators play a role of Cartan generators. Super-Hamiltonian approach to super-Macdonald polynomials was also considered in [3, 29]. Another approach to Macdonald polynomials recently discussed in [30].

The paper consists of three parts.

Section 2 is a brief reminder of the well-known Hamiltonians for Schur polynomials and a discussion of the "sum over boxes" property for eigenvalues and its connection to Pierri rules. In Sections 3 and 4 we provide basic facts about super-Schur and Macdonald cases. Section 5 is devoted to super-generalizations of Macdonald Hamiltonians – the main part of this paper. Section 6 contains a short conclusion and discussion of the future problems.

2 Hamiltonians for Schur polynomials

2.1 The simplest Hamiltonians

The grading (dilatation) operator is the simplest Hamiltonian for Schur polynomials:

$$\hat{\mathbf{G}}_{\square} = \sum_{k=1}^{\infty} k \, p_k \frac{\partial}{\partial p_k} \tag{1}$$

Its eigenvalues count the number of boxes in the Young diagram:

$$\hat{\mathbf{G}}_{\square} S_{\lambda} = \left(\sum_{\square \in \lambda} 1\right) \cdot S_{\lambda} = |\lambda| \cdot S_{\lambda} \tag{2}$$

there is a whole tower of W-operators starting with the cut-and-join operator [31,32]:

$$\hat{W} = \frac{1}{2} \sum_{a,b=1}^{\infty} \left(ab \cdot p_{a+b} \frac{\partial^2}{\partial p_a \partial p_b} + (a+b) \cdot p_a p_b \frac{\partial}{\partial p_{a+b}} \right)$$
(3)

with the eigenvalues κ_{λ} :

$$\hat{W} S_{\lambda} = \kappa_{\lambda} \cdot S_{\lambda} \qquad \qquad \boxed{\kappa_{\lambda} = \sum_{\square \in \lambda} (j_{\square} - i_{\square})}$$
(4)

Note that the eigenvalues are given by the *sum over boxes* in the Young diagram. We denote by i_{\square} and j_{\square} the vertical and horizontal coordinates respectively of a box \square in the Young diagram. Grading operators are represented as sums of units over the boxes, and higher Hamiltonians are sums of coordinate dependent functions like ω_{\square} . "Sum over boxes" property of Hamiltonians will be important in construction of higher Hamiltonians.

Another relevant for us property of Schur polynomials is the Pierri rules:

$$p_1 \cdot S_{\lambda} = \sum_{\square \in \text{Add}(\lambda)} S_{\lambda+\square} \tag{5}$$

$$\boxed{\frac{\partial}{\partial p_1} \cdot S_{\lambda} = \sum_{\square \in \text{Rem}(\lambda)} S_{\lambda - \square}}$$
(6)




Figure 1: Sets $Add(\star)$ and $Rem(\star)$.

We denote by $Add(\lambda)$ a subset of border places in diagram λ such that one can add a box to that place and the resulting diagram remains a Young diagram. Rem(λ) is the set of boxes in diagram λ that one can remove and the resulting diagram remains a Young diagram. An explanatory picture is presented on Fig.1.

2.2 Higher Hamiltonians

By combining two properties – the sum over boxes in eigenvalues (4) and Pierri rules (5), (6) – we could construct a set of nontrivial commuting operators - Hamiltonians for Schur polynomials. At the first step we define the following operators, that generalize the Pierri rule:

$$\hat{E}_0 := p_1 \tag{7}$$

$$\hat{E}_1 := \left[\hat{W}, \hat{E}_0\right] = \sum_{a=1}^{\infty} a \, p_{a+1} \frac{\partial}{\partial p_a} \tag{8}$$

$$\dots$$
 (9)

$$\hat{E}_k := \left[\hat{W}, \hat{E}_{k-1} \right] \tag{10}$$

All these operators add one box to the Young diagram yet with different coefficients:

$$\hat{E}_k S_{\lambda} = \sum_{\square \in \text{Add}(\lambda)} (\omega_{\square})^k \cdot S_{\lambda+\square}$$
(11)

where we simplify the notation by introducing $\omega_{\square} := j_{\square} - i_{\square}$. The above formula could be derived easily step by step in k. We demonstrate explicitly only the first step:

$$\hat{E}_{1} S_{\lambda} = \left[\hat{W}, \hat{E}_{0}\right] S_{\lambda} = \hat{W} \left(\sum_{\square \in \text{Add}(\lambda)} S_{\lambda+\square}\right) - \hat{E}_{0} \left(\kappa_{\lambda} \cdot S_{\lambda}\right) =$$

$$= \sum_{\square \in \text{Add}(\lambda)} \kappa_{\lambda+\square} \cdot S_{\lambda+\square} - \kappa_{\lambda} \cdot \sum_{\square \in \text{Add}(\lambda)} S_{\lambda+\square} = \sum_{\square \in \text{Add}(\lambda)} \left(\kappa_{\lambda+\square} - \kappa_{\lambda}\right) \cdot S_{\lambda+\square} = \sum_{\square \in \text{Add}(\lambda)} \omega_{\square} \cdot S_{\lambda+\square}$$

$$(12)$$

In the same way we define box-removing operators:

$$\hat{F}_0 := \frac{\partial}{\partial n_1} \tag{13}$$

$$\hat{F}_1 := -\left[\hat{W}, \hat{F}_0\right] = \sum_{a=1}^{\infty} (a+1) \, p_a \frac{\partial}{\partial p_{a+1}}$$
 (14)

$$\dots$$
 (15)

$$\hat{F}_k := -\left[\hat{W}, \hat{F}_{k-1}\right] \tag{16}$$

Operators F_k generalize the second Pierri rule (6):

$$\hat{F}_k S_{\lambda} = \sum_{\square \in \text{Rem}(\lambda)} (\omega_{\square})^k \cdot S_{\lambda - \square}$$
(17)

On the second step we commute operators E_k and F_k in order to obtain Hamiltonians:

$$\hat{H}_{a+b} := \left[\hat{E}_a, \hat{F}_b \right] \tag{18}$$

There are two nontrivial consequences:

- operators $\left[\hat{E}_a, \hat{F}_b\right]$ depend only on the sum a+b
- operators \hat{H}_k commute

$$\left[\hat{H}_k, \hat{H}_l\right] = 0 \tag{19}$$

The above properties follow directly from relations (11) and (17). One could easily prove both properties by showing that Schur polynomials are eigenfunctions of $\left[\hat{E}_a, \hat{F}_b\right]$:

For demonstrative purposes we colored the pieces into blue and red colors, if they correspond to contributions of \hat{E}_a and \hat{F}_a respectively.

The first part of the above relation represents the case when F_b removes box \square from λ , and then E_a adds the same (already removed) box \square to its place thus the resulting diagram remains λ as it was in the beginning. The second part is the same as the first part with the difference that E_a add a box to the λ and F_b removes it. Therefore, the first and the second parts contribute to the diagonal part of the operator $\left[\hat{E}_a, \hat{F}_b\right]$.

In contrast, the third and the fourth parts contribute to the off-diagonal part of the operator. In these cases a box is added to the diagram λ and the *other* box is removed from the other place of the diagram. In the formula we emphasized $\square \in \text{Rem}(\lambda)$ and $\square \in \text{Add}(\lambda)$ meaning $\square \neq \square$. In particular, adding and removing different boxes is a commutative operation, therefore:

$$\lambda - \square + \square = \lambda + \square - \square, \quad \forall \lambda \neq \emptyset, \square \tag{21}$$

In general situation cancellation of off-diagonal terms provided by the following condition:

$$\left[\hat{E}_{0},\hat{F}_{0}\right] = \text{diagonal operator}$$
 (22)

In this particular case $\left[p_1, \frac{\partial}{\partial p_1}\right] = 1$. On the other hand we have unit coefficients in Pierri rules (5) and (6), this fact directly explains absence of off-diagonal terms.

The above relation explains mutual cancellation of the third and fourth part of (20). Finally, there is the only diagonal part that depends solely on the sum of indices a+b, thus Schur polynomials are the common eigenfunctions, and H_k all commute.

The subalgebra of Hamiltonians closes due to the following relations:

$$\hat{H}_3 = -6 \cdot \hat{W} \tag{23}$$

$$\hat{H}_2 = -2 \cdot \hat{G}_{\square} \tag{24}$$

that is provided by the peculiar properties of eigenvalues and Young diagrams:

$$\sum_{\square \in \text{Rem}(\lambda)} (\omega_{\square})^3 - \sum_{\square \in \text{Add}(\lambda)} (\omega_{\square})^3 = -6 \cdot \sum_{\square \in \lambda} \omega_{\square}$$
 (25)

$$\sum_{\square \in \text{Rem}(\lambda)} (\omega_{\square})^2 - \sum_{\square \in \text{Add}(\lambda)} (\omega_{\square})^2 = -2 \cdot \sum_{\square \in \lambda} 1$$
 (26)

that are valid for any Young diagram λ .

One of the main observations in this paper: the above construction of a commuting system of Hamiltonians can be applied to another family of polynomials that is enumerated by other diagrams. The minimum piece of data that we need:

- an operator that adds boxes to the diagrams (analog of (5));
- an operator that removes boxes from the diagrams (analog of (6));
- a Hamiltonian, whose eigenvalues are given by the sum over boxes (analog of (4)).

2.3 Additional properties

At the end of this section we would like to mention two intriguing properties of Schur Hamiltonians:

1. Operators \hat{G}_{\square} and \hat{W} can be considered as the first two non-trivial terms of the following expansion:

$$\sum_{n=0}^{\infty} S_{[n]} \left\{ k p_k \right\} S_{[n]} \left\{ k^2 \frac{\partial}{\partial p_k} \right\} = 1 + \underbrace{\hat{G}_{\square}}_{\text{spin } 2} + \underbrace{\hat{W}}_{\text{spin } 3} + \dots$$
 (27)

where the omitted terms with higher spin (in our notation spin is defined as a total degree in p_k and $\frac{\partial}{\partial p_k}$ in a normal ordered form), that are no longer made from the Hamiltonians (Schur polynomials are not their eigenfunctions). The powers of k in the arguments of symmetric Schur functions might seem unfamiliar, however they will become apparent after a consideration of the Macdonald deformation.

2. All the constructed Hamiltonians H_k can be expressed in terms of the Schur polynomials themselves and their conjugates:

$$\hat{H}_k = \sum_{|\lambda| = |\mu|} A_{\lambda,\mu} \cdot S_\lambda \, \hat{S}_\mu \tag{28}$$

where we define:

$$\hat{S}_{\mu} := S_{\mu} \left(p_k = k \frac{\partial}{\partial p_k} \right) \tag{29}$$

There is an *intriguing* property of Hamiltonians H_k that their expansion in terms of the Schur functions include only a sector of single-hook Young diagrams [32]. For example, the forms of the grading operator and the cut-and-join operator read:

$$\hat{G}_{\square} = \sum_{a=1}^{\infty} a \, p_a \frac{\partial}{\partial p_a} = \sum_{n=1}^{\infty} \sum_{i,j=0}^{n-1} (-)^{i+j} \, S_{[n-i,1^i]} \hat{S}_{[n-j,1^j]}$$
(30)


$$\hat{W} = -\frac{H_3}{6} = \frac{1}{2} \sum_{a,b=1}^{\infty} (a+b) \cdot p_a p_b \frac{\partial}{\partial p_{a+b}} + ab \cdot p_{a+b} \frac{\partial}{\partial p_a} \frac{\partial}{\partial p_b} =$$

$$= \sum_{n=1}^{\infty} \sum_{i,j=0}^{n-1} (-)^{i+j} (n-1-i-j) S_{[n-i,1^i]} \hat{S}_{[n-j,1^j]}$$
(31)

where we denoted a single hook diagram as $[n-i, 1^i]$, where i is the height of the first column and n is the total number of boxes. The higher W-operators could be obtained as proper combinations of H_k [32].

3 Hamiltonians for super-Schur polynomials

Super-diagrams [2,6,9] are half-integer partitions $\lambda = [\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_{l(\lambda)}], \lambda_k \in \mathbb{N}/2$ with decreasing rule $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots \geq \lambda_{l(\lambda)}$. If values λ_k and λ_{k+1} are both half-integer then inequality necessarily becomes strict $\lambda_k > \lambda_{k+1}$. Usual Young diagrams are the subset of super-Young diagrams. We depict super-partitions as the following diagrams:



The number of super-Young diagrams p(n,m) with n full boxes and m half-boxes can be extracted from the following generating function:

$$\prod_{k=1} \frac{1+y \, x^{k-1}}{1-x^k} = \sum_{n,m=0} p(n,m) \, x^n y^m = 1 + y + x + 2xy + 2x^2 + xy^2 + 4x^2 y + \dots \tag{32}$$

If we set y = 0 the above generating function becomes generating function for number of usual Young diagrams p(n):

$$\prod_{k=1} \frac{1}{1-x^k} = \sum_{n=0} p(n) x^n = 1 + x + 2x^2 + 3x^3 + 5x^4 + \dots$$
 (33)

3.1 The simplest Hamiltonians

In the case of super-Schur polynomials ordinary time variables p_a are supplemented by Grassmann variables θ_a . Therefore the number of grading operators is increased. There is an operator that counts the number of fermions:

$$\hat{G}_{\nabla - \Delta} = \frac{1}{2} \sum_{k=1}^{\infty} \theta_k \frac{\partial}{\partial \theta_k}$$
 (34)

with eigenvalues that are given by the sum over half-boxes:

$$\hat{G}_{\mathbf{P}-\mathbf{\Delta}}\mathcal{S}_{\lambda} = \left(\sum_{\mathbf{P}\in\lambda} \frac{1}{2} - \sum_{\mathbf{\Delta}\in\lambda} \frac{1}{2}\right) \cdot \mathcal{S}_{\lambda} \tag{35}$$

Note that we treat the full box as a combination of two half-boxes:

$$\Box = \Box = \mathcal{V} + \Delta \tag{36}$$

Therefore the sum in (35) effectively runs over non-paired half-boxes \mathbf{r} on the border of the diagram. Next grading operator counts half-boxes \mathbf{r} and $\mathbf{\Delta}$ on equal footing:

$$\hat{G}_{\mathbf{P}+\mathbf{\Delta}} = \sum_{k=1}^{\infty} \left(k \, p_k \frac{\partial}{\partial p_k} + \left(k - \frac{1}{2} \right) \theta_k \frac{\partial}{\partial \theta_k} \right) \tag{37}$$

Eigenvalues of this operator have the following form:

$$\hat{G}_{\mathbf{P}+\mathbf{\Delta}}\mathcal{S}_{\lambda} = \left(\sum_{\mathbf{P}\in\lambda} \frac{1}{2} + \sum_{\mathbf{\Delta}\in\lambda} \frac{1}{2}\right) \cdot \mathcal{S}_{\lambda} \tag{38}$$

From these operators we can construct grading operators that counts only half-boxes of particular kind:

$$\hat{G}_{\mathbf{P}} = \hat{G}_{\mathbf{P}+\mathbf{\Delta}} + \hat{G}_{\mathbf{P}-\mathbf{\Delta}} = \sum_{k=1}^{\infty} \left(k \, p_k \frac{\partial}{\partial p_k} + k \, \theta_k \frac{\partial}{\partial \theta_k} \right) \tag{39}$$

$$\hat{G}_{\mathbf{\Delta}} = \hat{G}_{\mathbf{P}+\mathbf{\Delta}} - \hat{G}_{\mathbf{P}-\mathbf{\Delta}} = \sum_{k=1}^{\infty} \left(k \, p_k \frac{\partial}{\partial p_k} + (k-1) \, \theta_k \frac{\partial}{\partial \theta_k} \right) \tag{40}$$

where eigenvalues:

$$\hat{G}_{\mathbf{r}} \, \mathcal{S}_{\lambda} = \left(\sum_{\mathbf{r} \in \lambda} 1 \right) \cdot \mathcal{S}_{\lambda} \tag{41}$$

$$\hat{G}_{\Delta} S_{\lambda} = \left(\sum_{\Delta \in \lambda} 1\right) \cdot S_{\lambda} \tag{42}$$

Next we provide an explicit form of the first few non-trivial Hamiltonians:

$$\hat{\mathcal{W}}_{\nabla-\Delta} = \sum_{a,b=1}^{\infty} \left(a\theta_{a+b} \frac{\partial^2}{\partial p_a \partial \theta_b} + p_a \theta_b \frac{\partial}{\partial \theta_{a+b}} \right) \tag{43}$$

$$\hat{\mathcal{W}}_{\mathbf{r}} = \frac{1}{2} \sum_{a,b=1}^{\infty} \left(abp_{a+b} \frac{\partial^2}{\partial p_a \partial p_b} + (a+b)p_a p_b \frac{\partial}{\partial p_{a+b}} \right) + \sum_{a,b=1}^{\infty} b \left(a\theta_{a+b} \frac{\partial^2}{\partial p_a \partial \theta_b} + p_a \theta_b \frac{\partial}{\partial \theta_{a+b}} \right)$$
(44)

with the eigenvalues

$$\hat{\mathcal{W}}_{\mathbf{P}-\mathbf{\Delta}}\mathcal{S}_{\lambda} = \left(\sum_{\mathbf{P}\in\lambda} (j_{\mathbf{P}} - i_{\mathbf{P}}) - \sum_{\mathbf{\Delta}\in\lambda} (j_{\mathbf{\Delta}} - i_{\mathbf{\Delta}})\right) \cdot \mathcal{S}_{\lambda}$$
(45)

$$\hat{\mathcal{W}}_{\mathbf{P}} \, \mathcal{S}_{\lambda} = \left(\sum_{\mathbf{P} \in \lambda} (j_{\mathbf{P}} - i_{\mathbf{P}}) \right) \cdot \mathcal{S}_{\lambda} \tag{46}$$

A construction of Hamiltonians for only lower-half-boxes is also apparent:

$$\hat{\mathcal{W}}_{\Delta} = \hat{\mathcal{W}}_{\nabla} - \hat{\mathcal{W}}_{\nabla - \Delta} \tag{47}$$

$$\hat{\mathcal{W}}_{\Delta} \mathcal{S}_{\lambda} = \left(\sum_{\Delta \in \lambda} (j_{\Delta} - i_{\Delta}) \right) \cdot \mathcal{S}_{\lambda}$$
 (48)

There are similar Pierri rules for super-Schur polynomials. We define four operators

$$\hat{E}_0^{\mathbf{r}} := \theta_1 \qquad \qquad \hat{E}_0^{\mathbf{z}} := \sum_{k=1}^{\infty} p_k \frac{\partial}{\partial \theta_k}$$
 (49)

$$\hat{F}_0^{\mathbf{r}} := \frac{\partial}{\partial \theta_1} \qquad \qquad \hat{F}_0^{\mathbf{z}} := \sum_{k=1}^{\infty} k \, \theta_k \frac{\partial}{\partial p_k} \tag{50}$$

that add/remove half-boxes to/from the super-diagrams:

$$\hat{E}_0^{\nabla} \mathcal{S}_{\lambda} = \sum_{\nabla \in \text{Add}(\lambda)} C_{\lambda, \lambda + \nabla} \mathcal{S}_{\lambda + \nabla} \qquad \qquad \hat{E}_0^{\Delta} \mathcal{S}_{\lambda} = \sum_{\Delta \in \text{Add}(\lambda)} C_{\lambda, \lambda + \Delta} \mathcal{S}_{\lambda + \Delta}$$
 (51)

$$\hat{F}_0^{\nabla} \mathcal{S}_{\lambda} = \sum_{\nabla \in \text{Rem}(\lambda)} C_{\lambda, \lambda - \nabla} \mathcal{S}_{\lambda - \nabla} \qquad \qquad \hat{F}_0^{\Delta} \mathcal{S}_{\lambda} = \sum_{\Delta \in \text{Rem}(\lambda)} C_{\lambda, \lambda - \Delta} \mathcal{S}_{\lambda - \Delta}$$
 (52)




Figure 2: Sets $Add(\star)$ and $Rem(\star)$.

where $C_{\lambda,\lambda\pm \mathbf{P}}$ and $C_{\lambda,\lambda\pm \mathbf{d}}$ are some constants. The definition of sets $\mathrm{Add}(\lambda)$ and $\mathrm{Rem}(\lambda)$ is clear from the Fig.2.

Higher Hamiltonians 3.2

To construct higher Hamiltonians for the super-case we follow a procedure similar to the one presented in Sect.2. At the first step we define higher operators recursively:

$$\hat{E}_{k}^{\mathbf{r}} := \left[\hat{\mathcal{W}}_{\mathbf{r}}, \hat{E}_{k-1}^{\mathbf{r}}\right] \qquad \qquad \hat{E}_{k}^{\mathbf{\Delta}} := \left[\hat{\mathcal{W}}_{\mathbf{\Delta}}, \hat{E}_{k-1}^{\mathbf{\Delta}}\right] \tag{53}$$

$$\hat{F}_{k}^{\mathbf{r}} := -\left[\hat{\mathcal{W}}_{\mathbf{r}}, \hat{F}_{k-1}^{\mathbf{r}}\right] \qquad \qquad \hat{F}_{k}^{\mathbf{\Delta}} := -\left[\hat{\mathcal{W}}_{\mathbf{\Delta}}, \hat{F}_{k-1}^{\mathbf{\Delta}}\right] \qquad (54)$$

At the second step the higher Hamiltonians are defined as follows:

$$\hat{H}_{a+b}^{\mathbf{\Delta}} := \left\{ \hat{E}_{a}^{\mathbf{p}}, \hat{F}_{b}^{\mathbf{p}} \right\} \qquad \qquad \hat{H}_{a+b}^{\mathbf{p}} := \left\{ \hat{E}_{a}^{\mathbf{\Delta}}, \hat{F}_{b}^{\mathbf{\Delta}} \right\}$$
 (55)

Off-diagonal terms are absent due to relations (22):

$$\left\{\hat{E}_0^{\mathbf{p}}, \hat{F}_0^{\mathbf{p}}\right\} = 1\tag{56}$$

$$\left\{\hat{E}_{0}^{\mathbf{\Delta}}, \hat{F}_{0}^{\mathbf{\Delta}}\right\} = \sum_{k=1}^{\infty} \left(k \, p_{k} \frac{\partial}{\partial p_{k}} + k \, \theta_{k} \frac{\partial}{\partial \theta_{k}}\right) \tag{57}$$

Here anti-commutators are implemented rather than the usual commutators due to Grassmann odd nature of $\hat{E}_k^{\nabla,\Delta}$ and $\hat{F}_k^{\nabla,\Delta}$ operators. Note that the labels of operators in the r.h.s. are different from the labels of Hamiltonians in the l.h.s. These Hamiltonians form a commuting set:

$$\left[\hat{H}_k^{\mathbf{p}}, \hat{H}_l^{\mathbf{p}}\right] = 0 \tag{58}$$

$$\left[\hat{H}_k^{\mathbf{\Delta}}, \hat{H}_l^{\mathbf{\Delta}}\right] = 0 \tag{59}$$

$$\left[\hat{H}_{k}^{\mathbf{r}}, \hat{H}_{l}^{\mathbf{\Delta}}\right] = 0 \tag{60}$$

This a bit awkwardly looking notation for Hamiltonians is determined by the following relations that make the subalgebra closed:

$$\hat{H}_{2}^{\mathcal{\Delta}} = \hat{G}_{\mathcal{\Delta}} \qquad \qquad \hat{H}_{3}^{\mathcal{\Delta}} = 2 \cdot \hat{\mathcal{W}}_{\mathcal{\Delta}} \qquad (61)$$

$$\hat{H}_{0}^{\mathcal{P}} = \hat{G}_{\mathcal{P}} \qquad \qquad \hat{H}_{1}^{\mathcal{P}} = 2 \cdot \hat{\mathcal{W}}_{\mathcal{P}} \qquad (62)$$

$$\hat{H}_0^{\mathbf{r}} = \hat{G}_{\mathbf{r}} \qquad \qquad \hat{H}_1^{\mathbf{r}} = 2 \cdot \hat{\mathcal{W}}_{\mathbf{r}} \tag{62}$$

3.3 Additional properties

1. Since super-Schurs for integer diagrams (not containing unpaired upper-half-boxes) coincide with the ordinary Schurs (do not depend on θ_k) from the following combinations we obtain usual grading operator and cut-and-join operator:

$$\sum_{n=0}^{\infty} \mathcal{S}_{[n]} \left\{ k p_k \right\} \mathcal{S}_{[n]} \left\{ k^2 \frac{\partial}{\partial p_k} \right\} = 1 + \hat{G}_{\square} + \hat{W} + \dots$$
 (63)

However, a similar combination for half-integer diagrams gives the following result:

$$\sum_{n \text{ odd}} \mathcal{S}_{[n/2]} \left\{ k p_k, k \theta_k \right\} \mathcal{S}_{[n/2]} \left\{ k^2 \frac{\partial}{\partial p_k}, k \frac{\partial}{\partial \theta_k} \right\} = \sum_{k=1}^{\infty} k \theta_k \frac{\partial}{\partial \theta_k} + \sum_{a,b=1}^{\infty} b \left(a \theta_{a+b} \frac{\partial^2}{\partial p_a \partial \theta_b} + p_a \theta_b \frac{\partial}{\partial \theta_{a+b}} \right) + \dots$$
(64)

(note that $\frac{\partial}{\partial \theta_k}$ in the argument is multiplied by k, not by k^2). Operators as being split in brackets in the r.h.s. are *not* grading operators, and super-Schurs are *not* eigenfunctions of each of them separately. Yet the r.h.s. *is* a combination of the Hamiltonians:

$$\sum_{n=0}^{\infty} \mathcal{S}_{[n/2]} \left\{ k p_k, k \theta_k \right\} \mathcal{S}_{[n/2]} \left\{ k^2 \frac{\partial}{\partial p_k}, k \frac{\partial}{\partial \theta_k} \right\} = 1 + \hat{G}_{\nabla} + \hat{\mathcal{W}}_{\nabla} + \dots$$
 (65)

Operators \hat{G}_{Δ} and \hat{W}_{Δ} in bilinear combination of super-Schurs remain to be found.

2. If we expand this operators in terms of super-Schur polynomials and their conjugates:

$$\hat{\mathcal{S}}_{\lambda} := \mathcal{S}_{\lambda} \left(p_k = k \frac{\partial}{\partial p_k}, \theta_k = \frac{\partial}{\partial \theta_k} \right)$$
 (66)

we observe the same miraculous property as in (31): only single-hook diagrams contribute.

However, in this case there are some peculiarities. We do not know yet any natural normalization conditions for the super-Schur polynomials. We choose the normalization with unit coefficients in front of monomials p_1^n and $p_1^n\theta_1$ respectively. Therefore, the orthogonality conditions have the following form:

$$\hat{\mathcal{S}}_{\lambda} \, \mathcal{S}_{\mu} = \delta_{\lambda,\mu} \cdot ||\mathcal{S}_{\lambda}||^2 \tag{67}$$

with some coefficients $||\mathcal{S}_{\lambda}||^2$.

Moreover, in case of super-diagrams we have four types of single-hook diagrams. Two types on even levels $|\lambda| = 2m$ (B and F stand for boson and fermion respectively):

FF type:
$$\left[\frac{k}{2}, 1^l, \frac{1}{2}\right]$$
 BB type: $[k, 1^l]$ (68)

For example, $|\lambda| = 5$:

$$\left[\frac{7}{2}, 1, \frac{1}{2}\right] =$$
 $\left[3, 1^2\right] =$ $\left[69\right]$

And two types on odd levels $|\lambda| = 2m + 1$:

FB type:
$$\left[\frac{k}{2}, 1^l\right]$$
 BF type: $\left[k, 1^l, \frac{1}{2}\right]$ (70)

For example, $|\lambda| = \frac{9}{2}$:

$$\left[\frac{5}{2}, 1^2\right] = \left[2, 1^2, \frac{1}{2}\right] = \left[7, 1^2\right]$$

Finally, the following single-hook expansion holds for super-Schur Hamiltonians:

$$\hat{\mathcal{W}}_{\mathbf{P}} = \hat{\mathcal{W}}_{\mathbf{P}}^{(1)} + \hat{\mathcal{W}}_{\mathbf{P}}^{(2)} + \hat{\mathcal{W}}_{\mathbf{P}}^{(3)} + \hat{\mathcal{W}}_{\mathbf{P}}^{(4)} + \hat{\mathcal{W}}_{\mathbf{P}}^{(5)}$$
(72)

 $BB \cdot \hat{BB}$:

$$\hat{\mathcal{W}}_{\mathbf{p}}^{(1)} = \sum_{n \text{ even }} \sum_{i,j=0}^{\frac{n}{2}-1} (-)^{i+j} \left(\frac{n-2}{2} - i - j \right) \cdot \frac{C_{\frac{n-1}{2}}^{j}}{C_{\frac{n-1}{2}}^{i}} \cdot \frac{\mathcal{S}_{\left[\frac{n}{2}-i,1^{i}\right]} \hat{\mathcal{S}}_{\left[\frac{n}{2}-j,1^{j}\right]}}{||\mathcal{S}_{\left[\frac{n}{2}-i,1^{i}\right]}||^{2}}$$

 $FB \cdot \hat{FB}$:

$$\hat{\mathcal{W}}_{\mathbf{r}}^{(2)} = \sum_{n \text{ odd}} \sum_{i,j=0}^{\frac{n-3}{2}} (-)^{i+j} \left(\frac{n-1}{2} - i - j \right) \cdot \frac{C_{\frac{n-3}{2}}^{j}}{C_{\frac{n-1}{2}}^{i}} \cdot \frac{\mathcal{S}_{\left[\frac{n}{2}-i,1^{i}\right]} \hat{\mathcal{S}}_{\left[\frac{n}{2}-j,1^{j}\right]}}{||\mathcal{S}_{\left[\frac{n}{2}-i,1^{i}\right]}||^{2}}$$

 $BF \cdot \hat{BF}$:

$$\hat{\mathcal{W}}_{\mathbf{p}}^{(3)} = \sum_{n \text{ odd }} \sum_{i,j=0}^{\frac{n-3}{2}} (-)^{i+j} \left(\frac{n-5}{2} - i - j \right) \cdot \frac{C_{\frac{n-3}{2}}^{j}}{C_{\frac{n-1}{2}}^{i+1}} \cdot \frac{\mathcal{S}_{\left[\frac{n-1}{2} - i, 1^{i}, \frac{1}{2}\right]} \hat{\mathcal{S}}_{\left[\frac{n-1}{2} - j, 1^{j}, \frac{1}{2}\right]}}{||\mathcal{S}_{\left[\frac{n-1}{2} - i, 1^{i}, \frac{1}{2}\right]}||^{2}}$$
(73)

 $FB \cdot \hat{BF}$:

$$\hat{\mathcal{W}}_{\mathbf{p}}^{(4)} = \sum_{n \text{ odd }} \sum_{i,j=0}^{\frac{n-3}{2}} (-)^{i+j+1} \left(\frac{n-3}{2} - i - j \right) \cdot \frac{C_{\frac{n-3}{2}}^{j}}{C_{\frac{n-1}{2}}^{i}} \cdot \frac{\mathcal{S}_{\left[\frac{n}{2} - i, 1^{i}\right]} \hat{\mathcal{S}}_{\left[\frac{n-1}{2} - j, 1^{j}, \frac{1}{2}\right]}}{||\mathcal{S}_{\left[\frac{n}{2} - i, 1^{i}\right]}||^{2}}$$

 $BF \cdot \hat{FB}$:

$$\hat{\mathcal{W}}_{\mathbf{p}}^{(5)} = \sum_{n \text{ odd }} \sum_{i,j=0}^{\frac{n-3}{2}} (-)^{i+j+1} \left(\frac{n-3}{2} - i - j \right) \cdot \frac{C_{\frac{n-3}{2}}^{j}}{C_{\frac{n-1}{2}}^{i+1}} \cdot \frac{\mathcal{S}_{\left[\frac{n-1}{2} - i, 1^{i}, \frac{1}{2}\right]} \hat{\mathcal{S}}_{\left[\frac{n}{2} - j, 1^{j}\right]}}{||\mathcal{S}_{\left[\frac{n-1}{2} - i, 1^{i}, \frac{1}{2}\right]}||^{2}}$$

Note that for this Hamiltonian sector FF never enters. The other super-Hamiltonian $\hat{\mathcal{W}}_{\Delta}$ for super-Schurs also have single-hook expansion, although we do not present it here. In general, higher super-Hamiltonians $\hat{H}_k^{\nabla,\Delta}$ do not have neither single-hook expansion nor "sum over boxes" property for eigenvalues. However, special combinations of $\hat{H}_k^{\nabla,\Delta}$ may have these properties, we postpone this question for future research.

4 Hamiltonians for Macdonald polynomials

4.1 Differential grading operator and finite-difference/shift cut-and-join

Now in addition to the grading operator appeared in the story of the Schur polynomials in Sec. 2, which remains a differential operator,

$$\hat{G}_{\square} = \sum_{k=1}^{\infty} k \, p_k \frac{\partial}{\partial p_k} \tag{74}$$

$$\hat{\mathbf{G}}_{\square} M_{\lambda}^{q,t} = \left(\sum_{\square \in \lambda} 1\right) \cdot M_{\lambda}^{q,t} \tag{75}$$

we have non-local (shift) Hamiltonians [17], the first and the simplest one has the following vertex operator form

$$\hat{\mathcal{H}}_{\square}^{+} = \oint \frac{dz}{z} \exp\left(\sum_{k=1}^{\infty} \frac{(1 - t^{-2k})}{k} p_k z^k\right) \exp\left(\sum_{k=1}^{\infty} \frac{(q^{2k} - 1)}{z^k} \frac{\partial}{\partial p_k}\right)$$
(76)

This single Hamiltonian turns out to be sufficient to determine all the Macdonald polynomials $M_{\lambda}^{q,t}$ as eigenfunctions with distinct eigenvalues:

$$\hat{\mathcal{H}}_{\square}^{+} M_{\lambda}^{q,t} = \mathcal{E}_{\lambda}^{q,t} M_{\lambda}^{q,t} \tag{77}$$

An expression for eigenvalue $\mathcal{E}_{\lambda}^{\,q,t}$ has a form of a sum over all boxes in the Young diagram:

$$\mathcal{E}_{\lambda}^{q,t} = 1 + (q^2 - 1)(1 - t^{-2}) \cdot \sum_{\Box \in \lambda} q^{2j_{\Box}} t^{-2i_{\Box}}$$
 (78)

where the coordinate of the initial box in the Young diagram reads $(i_{\square}, j_{\square}) = (0, 0)$ and i_{\square}, j_{\square} are vertical and horizontal coordinates correspondingly.

4.2 Single-hook Schur decomposition of Macdonald Hamiltonian

With the help of reduced Cauchy formula

$$\sum_{n=0}^{\infty} S_{[n]}\{p\} \cdot z^n = \exp\left(\sum_k \frac{p_k z^k}{k}\right) \tag{79}$$

this Hamiltonian can be rewritten as a much healthier analogue of (27):

$$\hat{\mathcal{H}}_{\Box}^{+} = \sum_{n=0}^{\infty} S_{[n]} \left\{ (1 - t^{-2k}) \cdot p_k \right\} \cdot S_{[n]} \left\{ k \cdot (q^{2k} - 1) \cdot \frac{\partial}{\partial p_k} \right\}$$
 (80)

where we can further substitute the single-hook expansion [?]

$$S_{[n]}\left\{ (1 - t^{-2k}) \cdot p_k \right\} = (1 - t^{-2}) \sum_{i=0}^{n-1} (-)^i \cdot t^{-2i} \cdot S_{[n-i,1^i]} \{ p_k \}$$
(81)

to arrive to the following expression:

$$\hat{\mathcal{H}}_{\square}^{+} = 1 + (q^{2} - 1)(1 - t^{-2}) \sum_{n=1}^{\infty} \sum_{i,j=0}^{n-1} (-)^{i+j} \cdot t^{-2i} q^{2(n-1-j)} \cdot S_{[n-i,1^{i}]} \cdot \hat{S}_{[n-j,1^{j}]}$$
(82)

Note that symmetric and single-hook decompositions (80) and (82) are exact, all terms in the sums should be taken into account – not like it was in the case of (27).

4.3 q, t-inversion symmetry

The Macdonald polynomials have symmetry with respect to an inversion of q and t parameters: $M_{\lambda}^{q,t} = M_{\lambda}^{q^{-1},t^{-1}}$. Even more surprisingly, this property is not reflected by the Hamiltonian (76), which implies there should be another Hamiltonian obtained by this transformation with the same eigenfunctions:

$$\hat{\mathcal{H}}_{\square}^{-} = \oint \frac{dz}{z} \exp\left(\sum_{k=1} \frac{(1-t^{2k})}{k} p_k z^k\right) \exp\left(\sum_{k=1} \frac{(q^{-2k}-1)}{z^k} \frac{\partial}{\partial p_k}\right)$$
(83)

$$\hat{\mathcal{H}}_{\square}^{-} M_{\lambda}^{q,t} = \tilde{\mathcal{E}}_{\lambda}^{q,t} M_{\lambda}^{q,t} \tag{84}$$

$$\mathcal{H}_{\Box} M_{\lambda}^{-} = \mathcal{E}_{\lambda}^{-} M_{\lambda}^{-}$$

$$\tilde{\mathcal{E}}_{\lambda}^{q,t} = \mathcal{E}_{\lambda}^{q^{-1}, t^{-1}} = 1 + (q^{-2} - 1)(1 - t^{2}) \cdot \sum_{\Box \in \lambda} q^{-2j_{\Box}} t^{2i_{\Box}},$$
(85)

4.4 Higher Hamiltonians

Operators p_1 and $\frac{\partial}{\partial p_1}$ add and remove boxes in Macdonald polynomials:

$$p_1 \cdot M_{\lambda}^{q,t} = \sum_{\square \in \text{Add}(\lambda)} C_{\lambda,\lambda+\square}^{q,t} \ M_{\lambda+\square}^{q,t}$$
(86)

$$\frac{\partial}{\partial p_1} M_{\lambda}^{q,t} = \sum_{\square \in \text{Rem}(\lambda)} C_{\lambda,\lambda-\square}^{q,t} M_{\lambda-\square}^{q,t}$$
(87)

With the help of these two Hamiltonians one can define operators \hat{E}_k , \hat{F}_k recursively for all integer values $k \in \mathbb{Z}$:

$$\hat{E}_{k+1} := \left[\hat{\mathcal{H}}_{\Box}^{+}, \hat{E}_{k} \right] \qquad \hat{F}_{k+1} := -\left[\hat{\mathcal{H}}_{\Box}^{+}, \hat{F}_{k} \right]$$
 (88)

$$\hat{E}_{k-1} := \left[\hat{\mathcal{H}}_{\square}^-, \hat{E}_k\right] \qquad \hat{F}_{k-1} := -\left[\hat{\mathcal{H}}_{\square}^-, \hat{F}_k\right] \tag{89}$$

Initial conditions for these recursion $\hat{E}_0 = p_1$ and $\hat{F}_0 = \frac{\partial}{\partial p_1}$. Action of these operators on Macdonald polynomials reads:

$$\hat{E}_k \ M_{\lambda}^{q,t} = \sum_{\square \in \text{Add}(\lambda)} \left(q^{2j\square} t^{-2i\square} \right)^k \ C_{\lambda,\lambda+\square}^{q,t} \ M_{\lambda+\square}^{q,t}$$

$$\tag{90}$$

$$\hat{F}_k M_{\lambda}^{q,t} = \sum_{\square \in \text{Rem}(\lambda)} \left(q^{2j_{\square}} t^{-2i_{\square}} \right)^k C_{\lambda,\lambda-\square}^{q,t} M_{\lambda-\square}^{q,t}$$

$$\tag{91}$$

Then Hamiltonians reads:

$$\hat{\mathcal{H}}'_{a+b} := \left[\hat{E}_a, \hat{F}_b \right] \tag{92}$$

that form commuting family:

$$\left[\hat{\mathcal{H}}_a', \hat{\mathcal{H}}_b'\right] = 0 \tag{93}$$

Higher Hamiltonians from [17] should be expressed as proper combinations of our $\hat{\mathcal{H}}'_a$. We leave it question for future papers.

5 Hamiltonians for super-Macdonald polynomials

Super-Macdonald polynomials are q, t-deformations of super-Schur polynomials therefore the two grading operators are just the same:

$$\hat{G}_{\mathbf{P}} \mathcal{M}_{\lambda}^{q,t} = \left(\sum_{\mathbf{P} \in \lambda} 1\right) \cdot \mathcal{M}_{\lambda}^{q,t} \tag{94}$$

$$\hat{\mathbf{G}}_{\mathbf{\Delta}} \mathcal{M}_{\lambda}^{q,t} = \left(\sum_{\mathbf{\Delta} \in \lambda} 1\right) \cdot \mathcal{M}_{\lambda}^{q,t} \tag{95}$$

In complete analogy with the case of canonical Macdonald polynomials where the Hamiltonian (76) have eigenvalues:

$$\mathcal{E}_{\lambda}^{q,t} = 1 + (q^2 - 1)(1 - t^{-2}) \cdot \sum_{\Box \in \lambda} q^{2j_{\Box}} t^{-2i_{\Box}}$$
(96)

we define new super-Hamiltonians for super-Macdonald polynomials:

$$\hat{\mathcal{H}}^{\nabla,+}\mathcal{M}_{\lambda}^{q,t} = \left[1 + (q^2 - 1)(1 - t^{-2}) \left(\sum_{\nabla \in \lambda} q^{2j\nabla} t^{-2i\nabla}\right)\right] \mathcal{M}_{\lambda}^{q,t}
\hat{\mathcal{H}}^{\Delta,+}\mathcal{M}_{\lambda}^{q,t} = \left[1 + (q^2 - 1)(1 - t^{-2}) \left(\sum_{\Delta \in \lambda} q^{2j\Delta} t^{-2i\Delta}\right)\right] \mathcal{M}_{\lambda}^{q,t}
\hat{\mathcal{H}}^{\nabla,-}\mathcal{M}_{\lambda}^{q,t} = \left[1 + (q^{-2} - 1)(1 - t^2) \left(\sum_{\nabla \in \lambda} q^{-2j\nabla} t^{2i\nabla}\right)\right] \mathcal{M}_{\lambda}^{q,t}
\hat{\mathcal{H}}^{\Delta,-}\mathcal{M}_{\lambda}^{q,t} = \left[1 + (q^{-2} - 1)(1 - t^2) \left(\sum_{\Delta \in \lambda} t^{2x\Delta} q^{-2y\Delta}\right)\right] \mathcal{M}_{\lambda}^{q,t}$$
(97)

For even (ordinary) diagrams the eigenvalues of the first two Hamiltonians coincide, whereas those of the last two are obtained by a switch $q = q^{-1}$, $t = t^{-1}$. However, for a generic odd diagram all the four eigenvalues are different and independent.

For the first pair of Hamiltonians we present the following closed expression:

$$\hat{\mathcal{H}}^{\nabla,+} = \oint \frac{dw}{w} \langle \varnothing | \left[\exp \left\{ \sum_{k=1}^{\infty} \left(\frac{1 - t^{-2k}}{k} p_k w^{2k} + (1 - t^{-2}) t^{-2(k-1)} w^{2k-1} \theta_k \sum_{m=0}^{k-1} t^{2m} \psi_m \right) \right\} \times \\
\times \exp \left\{ \sum_{k=1}^{\infty} \left((q^{2k} - 1) \frac{\partial}{\partial p_k} w^{-2k} + (q^2 - 1) q^{2(k-1)} w^{-2k+1} \frac{\partial}{\partial \theta_k} \sum_{m=0}^{k-1} q^{-2m} \psi_m^{\dagger} \right) \right\} \right] | \varnothing \rangle_F ,
\hat{\mathcal{H}}^{\Delta,+} = \oint \frac{dw}{w} \langle \varnothing | \left[\exp \left\{ \sum_{k=1}^{\infty} \frac{1 - t^{-2k}}{k} p_k w^{2k} + \sum_{k=2}^{\infty} \left((1 - t^{-2}) t^{-2(k-1)} w^{2k-1} \theta_k \sum_{m=0}^{k-2} t^{2m} \psi_m \right) \right\} \\
\times \exp \left\{ \sum_{k=1}^{\infty} (q^{2k} - 1) \frac{\partial}{\partial p_k} w^{-2k} + \sum_{k=2}^{\infty} \left((q^2 - 1) q^{2(k-2)} w^{-2k+1} \frac{\partial}{\partial \theta_k} \sum_{m=0}^{k-2} q^{-2m} \psi_m^{\dagger} \right) \right\} \right] | \varnothing \rangle_F , \tag{98}$$

where fermionic correlator are defined by:

$$\left\{\psi_i, \psi_j^{\dagger}\right\} = \delta_{ij}, \quad \psi_i |\varnothing\rangle_F = 0, \quad \langle\varnothing|\varnothing\rangle_F = 1$$
 (99)

For the second pair of Hamiltonians we propose the following expressions:

$$\hat{\mathcal{H}}^{\mathcal{V},-} = \oint \frac{dw}{w} \langle \varnothing | \left[\exp \left\{ \sum_{k=1}^{\infty} \left(\frac{1 - t^{2k}}{k} p_k w^{2k} s^{-k} + (1 - t^2) w^{2k-1} \theta_k \nu \right) \right\} \times \\ \times \exp \left\{ \sum_{k=1}^{\infty} \left(\frac{q^{-2k} - 1}{1} \frac{\partial}{\partial p_k} w^{-2k} + (q^{-2} - 1) w^{-2k+1} \frac{\partial}{\partial \theta_k} \nu^{\dagger} s^{k-1} \right) \right\} \right] | \varnothing \rangle_B ,$$

$$\hat{\mathcal{H}}^{\mathcal{A},-} = \oint \frac{dw}{w} \langle \varnothing | \left[\exp \left\{ \sum_{k=1}^{\infty} \left(\frac{1 - t^{2k}}{k} p_k w^{2k} s^{-k} + (1 - t^2) w^{2k-1} \theta_k \nu \right) \right\} \times \\ \times \exp \left\{ \sum_{k=1}^{\infty} \left(\frac{q^{-2k} - 1}{1} \frac{\partial}{\partial p_k} w^{-2k} + (q^{-2} - 1) w^{-2k+1} \frac{\partial}{\partial \theta_k} \nu^{\dagger} s^{k-2} \right) \right\} \right] | | \varnothing \rangle_B$$

$$(100)$$

where s is an ordinary number, whereas ν, ν^{\dagger} anti-commutative. For their correlators we define:

$$\langle \varnothing | s^{a} | \varnothing \rangle_{B} = 1, \quad \langle \varnothing | \nu \nu^{\dagger} s^{b} | \varnothing \rangle_{B} = \begin{cases} \frac{1 - (qt)^{2b+2}}{q^{2b}(1 - (qt)^{2})}, & \text{if } b \geq 0; \\ 0, & \text{otherwise} \end{cases}$$
(101)

Note that there is a single fermionic pair ν, ν^{\dagger} , therefore these two Hamiltonians contain only terms with at most the first power of θ and θ -derivative. The primary role of s is to exclude terms like $p_{k_1} \dots p_{k_n} \frac{\partial}{\partial \theta_{m-\frac{1}{2}}}$ with $\sum k_a > m-1$ from $\hat{\mathcal{H}}^{\nabla,-}$ or $\sum k_a > m-2$ from $\hat{\mathcal{H}}^{\mathcal{A},-}$.

The above four Hamiltonians by the definition have "sum over boxes" property, however we observe that in the expansion in terms of super-Schurs S_{λ} and conjugates \hat{S}_{μ} in the spirit of (73) enters diagrams with two and more hooks. We leave this intriguing observation for further investigation elsewhere.

According to the general reasoning from Section 2 we can construct a set of commuting Hamiltonians. We provide the following example, that is based on the Pierri rules for super-Macdonald polynomials:

$$\theta_1 \cdot \mathcal{M}_{\lambda}^{q,t} = \sum_{\varphi \in \text{Add}(\lambda)} C_{\lambda,\lambda+\varphi}^{q,t} \cdot \mathcal{M}_{\lambda+\varphi}^{q,t}$$

$$\tag{102}$$

$$\frac{\partial}{\partial \theta_1} \, \mathcal{M}_{\lambda}^{q,t} = \sum_{\mathcal{V} \in \text{Rem}(\lambda)} C_{\lambda, \lambda - \mathcal{V}}^{q,t} \cdot \mathcal{M}_{\lambda - \mathcal{V}}^{q,t} \tag{103}$$

Then we define the following auxiliary operators:

$$\hat{E}_{k\pm 1}^{\nabla} = \left[\hat{\mathcal{H}}^{\nabla, \pm}, \hat{E}_{k}^{\nabla} \right] \tag{104}$$

$$\hat{F}_{k\pm 1}^{\nabla} = -\left[\hat{\mathcal{H}}^{\nabla,\pm}, \hat{F}_{k}^{\nabla}\right] \tag{105}$$

These two relations allow increase and decrease index k to define operators for any integr values of k using $\hat{E}_0^{\nabla} = \theta_1$ and $\hat{F}_0^{\nabla} = \frac{\partial}{\partial \theta_1}$ as the initial conditions of this recursive procedure. Then the commuting set of Hamiltonians read:

$$\hat{\mathcal{H}'}_{a+b} := \left\{ \hat{E}_a^{\nabla}, \hat{F}_b^{\nabla} \right\} \tag{106}$$

$$\left[\hat{\mathcal{H}'}_a, \hat{\mathcal{H}'}_b\right] = 0 \tag{107}$$

We consider here only one possible commuting family of super-Hamiltonians that is related to $\hat{\mathcal{H}}^{\nabla,+}$ and $\hat{\mathcal{H}}^{\nabla,-}$. However, we need to construct a pair of operators that add and remove semi-boxes Δ for constructing the remaining commuting family, related to $\hat{\mathcal{H}}^{\Delta,+}$ and $\hat{\mathcal{H}}^{\Delta,-}$. We leave it for future research. Of course, these two commuting families commute with each other.

6 Conclusion

This paper is devoted to the analysis of Hamiltonians for polynomials of the Schur-Macdonald family and their supersymmetric extensions. These polynomials are naturally enumerated by Young-like diagrams. We argue that the "sum over boxes" property of eigenvalues equipped with the Pierri rule for adding/removing boxes provides integrability – commutativity of an infinite set of Hamiltonians.

In the case of Schur and super-Schur polynomials each box is weighted with content $\omega = j - i$ that is given by a difference of horizontal and vertical coordinates of the box. Furthermore for the Macdonald case each box is weighted with exponentiated weights $\omega^{q,t} = q^{2j}t^{-2i}$ where q,t are free

parameters. Therefore our main idea in constructing super-Hamiltonians is to enforce the special form of eigenvalues: for an eigen function (i.e. a super-Macdonald polynomial) that is labeled by the super-Young diagram the corresponding eigenvalue is given by a sum over all semi-boxes of these diagram with respective exponent weights (97).

Super-Hamiltonians are labeled by two indices. The first index labels two different types of semi-boxes ∇ and Δ according to the "sum over boxes" property of corresponding eigenvalues. The second index encodes positive or negative powers of free parameters q,t in eigenvalues. Therefore the total number of independent super-Hamiltonians is equal to four. However, only two (e.g. two positive) super-Hamiltonians are sufficient to compute all the super-Macdonald polynomials as eigen functions with distinct eigen values.

There are plenty of details to comment on in this approach – and each one raises an intriguing question:

- 1. There is spontaneous breaking of symmetry $q \to 1/q$, $t \to 1/t$. Macdonald polynomials are invariant with respect to this transformation, whereas Ruijsenaars Hamiltonians are not (this is true about the basic ones and continues to hold for the higher Hamiltonians, see [17]). Surprisingly, this property disappears after super-generalization super-Macdonalds are no longer invariant.
- 2. This symmetry violation leads to an occurrence of extra Hamiltonians what could be partly related to the apparent difference between Yangians and DIM: there are commuting rays in the former case. These rays are promoted to entire Heisenberg *lines* in the latter [33].
- 3. As it was observed in a variety of examples the "sum over boxes" property of eigenvalues is somehow related to a single-hook expansion of the Hamiltonian in terms of eigen functions and their conjugates [32]. However, this connection is lost in case of the Macdonald super-Hamiltonians, since they do not have a naive single-hook expansion in terms of super-Schur polynomials.
- 4. The projections can be implemented in super-Hamiltonians with the help of additional Grassmannian variables with their help the formulas acquire almost the same form as the Ruijsenaars finite-difference operators (vertex-form operators).

Acknowledgments

The work was partially funded within the state assignment of the Institute for Information Transmission Problems of RAS. Our work is partly supported by the grants of the Foundation for the Advancement of Theoretical Physics and Mathematics "BASIS" (A.M. and N.T.).

References

- [1] I. G. Macdonald, Symmetric Functions and Hall Polynomials. Oxford Mathematical Monographs, 1998.
- [2] O. Blondeau-Fournier, P. Desrosiers, L. Lapointe, and P. Mathieu, "Macdonald polynomials in superspace: conjectural definition and positivity conjectures," *Lett. Math. Phys.* 101 (2012) 27–47, arXiv:1112.5188 [math-ph].

- [3] O. Blondeau-Fournier, P. Desrosiers, L. Lapointe, and P. Mathieu, "Macdonald polynomials in superspace as eigenfunctions of commuting operators," *J. Comb.* **3** (2012) 495–561, arXiv:1202.3922 [math-ph].
- [4] D. Galakhov, A. Morozov, and N. Tselousov, "Macdonald polynomials for super-partitions," *Phys. Lett. B* **856** (2024) 138911, arXiv:2407.03301 [hep-th].
- [5] D. Galakhov, A. Morozov, and N. Tselousov, "Supersymmetric polynomials and algebro-combinatorial duality," *SciPost Phys.* 17 no. 4, (2024) 119, arXiv:2407.04810 [hep-th].
- [6] D. Galakhov, A. Morozov, and N. Tselousov, "Super-Schur polynomials for Affine Super Yangian Y($\hat{\mathfrak{gl}}_{1|1}$)," *JHEP* **08** (2023) 049, arXiv:2307.03150 [hep-th].
- [7] D. Galakhov, A. Morozov, and N. Tselousov, "Wall-crossing effects on quiver BPS algebras," JHEP 05 (2024) 118, arXiv:2403.14600 [hep-th].
- [8] D. Galakhov, A. Gavshin, A. Morozov, and N. Tselousov, "Algorithms for representations of quiver Yangian algebras," *JHEP* **08** (2024) 209, arXiv:2406.20074 [hep-th].
- [9] G. Noshita and A. Watanabe, "Shifted quiver quantum toroidal algebra and subcrystal representations," *JHEP* **05** (2022) 122, arXiv:2109.02045 [hep-th].
- [10] D. Galakhov, W. Li, and M. Yamazaki, "Toroidal and elliptic quiver BPS algebras and beyond," JHEP 02 (2022) 024, arXiv:2108.10286 [hep-th].
- [11] S. V. Kerov, "Hall-littlewood functions and orthogonal polynomials," Functional Analysis and Its Applications 25 no. 1, (1991) 65–66.
- [12] A. Mironov and A. Morozov, "Kerov functions revisited," Journal of Geometry and Physics 150 (Apr., 2020) 103608.
- [13] A. Mironov and A. Morozov, "On Hamiltonians for Kerov functions," Eur. Phys. J. C 80 no. 3, (2020) 277, arXiv:1908.05176 [hep-th].
- [14] A. Mironov and A. Morozov, "Kerov functions for composite representations and Macdonald ideal," *Nucl. Phys. B* **944** (2019) 114641, arXiv:1903.00773 [hep-th].
- [15] S. Ruijsenaars, "Complete integrability of relativistic calogero-moser systems and elliptic function identities," Communications in Mathematical Physics 110 (1987) 191–213.
- [16] S. Ruijsenaars and H. Schneider, "A new class of integrable systems and its relation to solitons," *Annals of Physics* **170** no. 2, (1986) 370–405.
- [17] A. Mironov and A. Morozov, "On generalized Macdonald polynomials," *JHEP* **01** (2020) 110, arXiv:1907.05410 [hep-th].
- [18] T. Procházka, "W-symmetry, topological vertex and affine Yangian," JHEP 10 (2016) 077, arXiv:1512.07178 [hep-th].
- [19] A. Tsymbaliuk, "The affine yangian of gl1 revisited," Advances in Mathematics **304** (2017) 583–645.

- [20] D. Galakhov, A. Morozov, and N. Tselousov, "Simple representations of BPS algebras: the case of $Y(\widehat{\mathfrak{gl}}_2)$," Eur. Phys. J. C 84 no. 6, (2024) 604, arXiv:2402.05920 [hep-th].
- [21] A. Morozov and N. Tselousov, "3-Schurs from explicit representation of Yangian Y $(\hat{\mathfrak{gl}}_1)$. Levels 1–5," *JHEP* 11 (2023) 165, arXiv:2305.12282 [hep-th].
- [22] A. Morozov and N. Tselousov, "Hunt for 3-Schur polynomials," Phys. Lett. B 840 (2023) 137887, arXiv:2211.14956 [hep-th].
- [23] J. Ding, J.-t. Ding, and K. Iohara, "Generalization and deformation of Drinfeld quantum affine algebras," *Lett. Math. Phys.* **41** (1997) 181–193, arXiv:q-alg/9608002.
- [24] K. Miki, "A (q,γ) analog of the W1+ ∞ algebra," J. Math. Phys. 48 no. 12, (2007) 123520.
- [25] H. Awata, H. Kanno, A. Mironov, A. Morozov, K. Suetake, and Y. Zenkevich, "(q,t)-KZ equations for quantum toroidal algebra and Nekrasov partition functions on ALE spaces," *JHEP* 03 (2018) 192, arXiv:1712.08016 [hep-th].
- [26] A. Mironov, A. Morozov, and A. Popolitov, "Commutative families in DIM algebra, integrable many-body systems and q,t matrix models," arXiv:2406.16688 [hep-th].
- [27] B. Feigin, M. Jimbo, T. Miwa, and E. Mukhin, "Quantum toroidal gl1-algebra: Plane partitions," *Kyoto Journal of Mathematics* **52** no. 3, (2012) 621 659.
- [28] H. Awata, H. Kanno, A. Mironov, A. Morozov, K. Suetake, and Y. Zenkevich, "The MacMahon R-matrix," JHEP 04 (2019) 097, arXiv:1810.07676 [hep-th].
- [29] O. Blondeau-Fournier, P. Desrosiers, and P. Mathieu, "Supersymmetric Ruijsenaars-Schneider Model," *Phys. Rev. Lett.* 114 (2015) 121602, arXiv:1403.4667 [hep-th].
- [30] A. Mironov, A. Morozov, and A. Popolitov, "A basic triad in Macdonald theory," arXiv:2411.16517 [hep-th].
- [31] A. Mironov, A. Morozov, and S. Natanzon, "Complete Set of Cut-and-Join Operators in Hurwitz-Kontsevich Theory," *Theor. Math. Phys.* **166** (2011) 1–22, arXiv:0904.4227 [hep-th].
- [32] A. Mironov and A. Morozov, "Hook variables: Cut-and-join operators and τ -functions," Phys. Lett. B 804 (2020) 135362, arXiv:1912.00635 [hep-th].
- [33] A. Mironov, V. Mishnyakov, A. Morozov, and A. Popolitov, "Commutative families in W_{∞} , integrable many-body systems and hypergeometric τ -functions," *JHEP* **23** (2020) 065, arXiv:2306.06623 [hep-th].