# Decision Making in Changing Environments: Robustness, Query-Based Learning, and Differential Privacy

Fan Chen fanchen@mit.edu Alexander Rakhlin rakhlin@mit.edu

January 28, 2025

#### Abstract

We study the problem of interactive decision making in which the underlying environment changes over time subject to given constraints. We propose a framework, which we call *hybrid Decision Making with Structured Observations* (hybrid DMSO), that provides an interpolation between the stochastic and adversarial settings of decision making. Within this framework, we can analyze local differentially private (LDP) decision making, query-based learning (in particular, SQ learning), and robust and smooth decision making under the same umbrella, deriving upper and lower bounds based on variants of the Decision-Estimation Coefficient (DEC). We further establish strong connections between the DEC's behavior, the SQ dimension, local minimax complexity, learnability, and joint differential privacy. To showcase the framework's power, we provide new results for contextual bandits under the LDP constraint.

# **1** Introduction

The Decision-Estimation Coefficient (DEC) [Foster et al., 2021, 2023b] has been recently shown to capture the difficulty of exploration in a wide range of problems in which a learning agent interacts with an unknown environment by making decisions and observing outcomes. Such problems include structured bandits, contextual bandits, and reinforcement learning, among others. The interaction protocol, termed *Decision Making with Structured Observations* (DMSO) in [Foster et al., 2021], assumes that the unknown model is fixed over the length of the interaction, i.e. the learning agent faces a stationary environment. This is often referred to as a *stochastic setting*, or *stochastic DMSO*. In contrast, the *adversarial DMSO*, studied in [Foster et al., 2022b], is a more complex task where the model may change arbitrarily between the rounds of the interaction.

In this paper, we study a setting that interpolates between the stochastic and adversarial DMSO. This interpolation is achieved by placing constraints on the way the model may change over time. Within the constraint set, the model is allowed to change arbitrarily, and we refer to the setting as that of *constrained adversaries*, or *hybrid DMSO*. In parallel with such constraints on the adversary, we additionally study constraints placed on the information received by the decision-maker, for instance due to privacy requirements or a specific oracle model of computation. The specification of constraints allows us to study—under the same umbrella—decision making with Statistical Queries (SQ) [Kearns, 1998], local differential privacy (LDP) [Kasiviswanathan et al., 2011, Duchi et al., 2013], robustness with respect to model corruption [Huber, 1965, Huber and Ronchetti, 2011], and smooth decision making [Rakhlin et al., 2011]. For example, in SQ learning, the decision-maker obtains information by issuing queries; since the response to these queries is only approximately

correct, it is natural to model it as a response of an adversary that has limited power in providing misleading information. Similarly, for robust decision making, we can model corruption (for instance, as in Huber's contamination model [Huber, 1965]) or mis-specification (as in agnostic learning) directly as a constraint on the environment to be close to a ground-truth model. In turn, the local privacy constraint can be formulated as a restriction on the decision-maker to only observe information through differentially private channels.

Our approach begins with the *hybrid DEC* formulation that yields both lower and upper bounds for PAC learning and no-regret learning under hybrid DMSO. Then, by investigating the specific information structures imposed by the constraints on the adversary and the decision-maker, we derive the corresponding DECs and the statistical guarantees for the aforementioned (and seemingly disparate) settings. As such, the unified viewpoint leads to a systematic "recipe" for analyzing new problems under the hybrid DMSO setting; this is illustrated on numerous examples throughout the paper. What is perhaps even more surprising, all the upper bounds are achieved by only two algorithmic approaches: a generalization of the Exploration-by-Optimization Algorithm [Lattimore and Szepesvári, 2020, Lattimore and Gyorgy, 2021, Foster et al., 2022b] and a variant of the Estimation-to-Decision Algorithm [Foster et al., 2021, 2023b].

The fact that DMSO provides such a unified viewpoint on disparate problems is a testament to the power of the framework, with DEC as the central notion of inherent problem complexity.

# 1.1 Contributions

We formulate decision making in the setting of hybrid DMSO, generalizing the Decision-Estimation Coefficient framework [Foster et al., 2021, 2023b]. Our proposed notion of hybrid DEC allows us to understand, under the same umbrella, minimax behavior of statistical estimation and interactive decision making under such seemingly different settings as local differential privacy, query-based learning (in particular, statistical queries), robust learning, and smoothness. In particular, hybrid DECs for PAC learning and no-regret learning yield both lower and upper bounds for the corresponding learning goals. Our upper bounds are achieved by the unified Exploration-by-Optimization Algorithm ( $ExO^+$ , cf. Lattimore and Szepesvári [2020], Lattimore and Gyorgy [2021], Foster et al. [2022b]).

As instantiations of our framework, we derive the hybrid DECs and corresponding upper and lower bounds for query-based learning (Section 2.2), locally private learning (Section 2.3), robust decision making (Section 2.4), and decision making against smooth adversaries (Section 2.5.1). The problem of contextual bandits with adversarial contexts also naturally falls under our hybrid formulation (Section 5.5), and we provide novel results for this setting as well.

Our primary goal is to understand the complexity of learning problems at some level of generality, rather than specific examples. Still, as a concrete application, our framework provides a near-optimal  $\sqrt{T}$ -regret for linear contextual bandits with local privacy (without well-conditioned assumptions), settling the open problem of the optimal regret in this setting [Zheng et al., 2020, Han et al., 2021, Li et al., 2024].

In addition, we make the following connections to other previously studied notions:

• SQ dimension. The SQ dimension proposed by Feldman [2017] provides both lower and upper bounds for the optimal query complexity of SQ learning of distribution search problems. Not surprisingly, we show that there is quantitative equivalence between the SQ dimension

and our SQ DEC (Section 4.2). Therefore, our results extend the characterizations of Feldman [2017] to general query-based learning problems.

- Local-minimax optimality under LDP. We show that our lower and upper bounds for LDP learning can be specialized to provide a tight characterization of the local-minimax complexity (Section 6.1). In particular, for functional estimation, our results recover (up to logarithmic factors) the characterization of Duchi and Ruan [2024] through the modulus of continuity.
- LDP learnability. We show that for any problem class, the fractional covering number [Chen et al., 2024] characterizes the finite-time LDP learnability (Section 6.2). In Section 6.3, we also relate fractional covering number to the learnability under joint differential privacy (JDP) and the representation dimension [Beimel et al., 2013a].

# 1.2 Related work

**Decision-Estimation Coefficient Framework.** Towards a unifying framework for interactive decision making, Foster et al. [2021] propose Decision-Estimation Coefficient (DEC), which provides both lower and upper bounds for any decision making problem. An active line of research [Foster et al., 2022b, Chen et al., 2022, Foster et al., 2023b,a, Glasgow and Rakhlin, 2023, Chen et al., 2024] has extended the DEC framework to various more general learning goals, including adversarial decision making [Foster et al., 2022b], PAC decision making [Chen et al., 2022, Foster et al., 2022b], reward-free learning and preference-based learning [Chen et al., 2022], multi-agent decision making and partial monitoring [Foster et al., 2023a], and interactive estimation [Chen et al., 2022, 2024]. The present work further extends the DEC framework to handle changing environments and constraints on the decision maker, and our results heavily draw on the techniques developed in these previous papers.

**Exploration-by-Optimization.** The *Exploration-by-Optimization* technique is powerful machinery developed in Lattimore and Szepesvári [2020], Lattimore and Gyorgy [2021] for partial monitoring in adversarial environments and later extended by Foster et al. [2022b] to decision making in adversarial environments, achieving upper bounds in terms of the generalized Information Ratio [Russo and Van Roy, 2014, 2018, Lattimore and Gyorgy, 2021] or the DEC [Foster et al., 2021, 2022b]. In the present work, we further extend this technique by incorporating the notion of *information sets*, allowing a more granular quantification of the information and model equivalences that the decision-maker can take advantage of. The idea of using information sets in the context of posterior sampling was proposed by Dylan Foster back in 2022, and was considered by the authors of [Foster et al., 2022a] as a way of improving DEC-based results for reinforcement learning.

Local differential privacy. The notion of local differential privacy (LDP) was formalized by Kasiviswanathan et al. [2011], Duchi et al. [2013], with some earlier work on this subject dating back to Warner [1965]. A line of research has been investigating the statistical complexity of locally private learning for various statistical estimation problems [Duchi et al., 2013, 2018, Duchi and Rogers, 2019], including mean estimation [Asi et al., 2022, 2024], functional estimation [Rohde and Steinberger, 2020, Butucea and Issartel, 2021, Butucea et al., 2023, Duchi and Ruan, 2024], hypothesis testing [Berrett and Butucea, 2020, Li et al., 2023] and selection [Gopi et al., 2020, Pour et al., 2024], and regression [Wang and Xu, 2019, Berrett et al., 2021], to name a few. Beyond the setting of statistical estimation, recent research studies the complexity of interactive decision making with local privacy constraints, including contextual bandits [Zheng et al., 2020, Han et al.,

2021, Li et al., 2024] and episodic RL [Garcelon et al., 2021, Liao et al., 2023]. Notably, these works mainly focus on specific problems and adopt problem-tailored approaches.

The role of interaction in LDP learning. It has long been known that there is a statistical separation between non-interactive private channels and sequential private channels [Kasiviswanathan et al., 2011]. As is surveyed by Butucea et al. [2023], the separation of sample complexity between non-interactive and interactive channels is identified for certain problems of testing [Berrett and Butucea, 2020] and functional estimation [Butucea and Issartel, 2021, Butucea et al., 2023]. Therefore, even for statistical problems (where samples are being generated i.i.d), interactive learning is generally necessary to achieve optimal sample complexity under LDP constraints. As the DEC framework characterizes the complexity of exploration of interactive decision making, it is suitable for quantifying the complexity of interactive LDP learning.

**Statistical Queries.** The Statistical Query (SQ) model was introduced by Kearns [1998] as a restricted PAC learning model, and it turns out to be a powerful tool for understanding the computational complexity of a wide range of algorithms and problems [Feldman et al., 2015, 2017, Diakonikolas et al., 2017, Brennan et al., 2020]. Variants of SQ model have also been studied [Bshouty and Feldman, 2002, Feldman, 2017, Joshi et al., 2024]. The connection between local DP and SQ learning has been identified by Kasiviswanathan et al. [2011]. For distributional search problems, Feldman [2017] characterized the SQ query complexity in terms of the *SQ dimension*, which turns out to be recovered by the SQ DEC (when specialized to this case).

**Robust statistics.** The *robustness* of a statistical procedure refers to the ability to adapt to model mis-specification or perturbation. In robust statistics, the contamination model of Huber [1965] has been extensively studied, where the data are assumed to be sampled i.i.d from a distribution that is  $\beta$ -contaminated from the ground-truth distribution. A recent line of work [Diakonikolas et al., 2019, Diakonikolas and Kane, 2019, Liu and Moitra, 2021, Diakonikolas and Kane, 2023, Canonne et al., 2023], among others, studied stronger contamination models, where the adversary is allowed to maliciously corrupt  $\beta$ -fraction of the whole dataset (detailed discussion in Appendix A.3). The connection between robustness and differential privacy is also studied by Georgiev and Hopkins [2022], Hopkins et al. [2023], Asi et al. [2023].

# 2 Overview of Results

We start this section by formulating the *hybrid DMSO* framework (Section 2.1), a generalization of the Decision Making with Structured Observation (DMSO) framework proposed by Foster et al. [2021]. We then show how this generalization encompasses query-based learning (Section 2.2), locally differentially private learning (Section 2.3), and robust decision making (Section 2.4). For each setting, we formulate a corresponding variant of DMSO, the corresponding DEC, and the ensuing PAC guarantees. We also present regret guarantees for hybrid DMSO (Section 2.5), with application to smooth learning (Section 2.5.1).

# 2.1 Hybrid DMSO

In the DMSO formulation, studied in [Foster et al., 2021], the learner (or, the decision maker) interacts for T rounds with the environment described by an underlying model  $M^*$ , unknown to the learner (detailed discussion in Appendix A.1). While the DMSO formulation is general enough

to capture various learning tasks and problem classes, it is restricted to the *stochastic* setting, where the underlying environment is stationary (specified by the model  $M^*$ ). However, in many applications, the environment is best described as non-stationary and changing according to the previous history of interaction, while at the same time satisfying certain constraints. Inspired by Foster et al. [2022b], who consider an arbitrarily changing environment, we propose the following *hybrid DMSO* formulation. We will reserve the term "stochastic DMSO" for the original DMSO setting of Foster et al. [2021].

In the hybrid DMSO setting studied here, the environment is allowed to be (adaptively) adversarial with certain constraints, while the learner has to interact with the environment through a given class  $\Phi$  of *measurements*. Specifically, let  $\mathbf{\Pi} = \mathbf{\Pi} \times \Phi$  be the joint decision space, and let ( $\mathbf{\Pi} \rightarrow \Delta(\mathcal{O})$ ) be the set of all *models*, with each model being a conditional distribution of observation given a (decision, measurement) pair. A *constraint* for the adversary will be modeled by a subset  $\mathcal{P} \subseteq (\mathbf{\Pi} \rightarrow \Delta(\mathcal{O}))$ , and a collection of constraints—as a set  $\mathscr{P}$  of such subsets. We consider the following *T*-round interaction protocol between the environment and the learner:

- 1. Before the interaction, the environment (or, the adversary) selects a constraint  $\mathcal{P}^* \subseteq (\Pi \to \Delta(\mathcal{O}))$ , without revealing it to the learner.
- 2. For each  $t = 1, \cdots, T$ :
  - The environment selects  $M^t \in \mathcal{P}^*$ , and the learner selects a decision  $\pi_t = (\pi_t, \phi_t) \in \Pi$ .
  - The learner observes  $o_t \in \mathcal{O}$ , sampled according to  $o_t \sim M^t(\pi_t, \phi_t)$ .

The set  $\mathcal{P}^*$  restricts the power of the adversary, and we assume the learner has access to a collection  $\mathscr{P}$  of constraints that contains  $\mathcal{P}^*$ . In other words,  $\mathscr{P}$  reflects prior knowledge of the possible constraints on the adversary. We formalize this assumption as follows.

# Assumption 1 (Constraint realizability). The given class $\mathscr{P}$ contains $\mathcal{P}^{\star}$ .

For some of the settings studied in this paper, the prior knowledge is additionally reflected in a more succinct model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{O}))$ , and the constraint class  $\mathscr{P}$  will reflect this choice.

The general formulation of constraints interpolates between

- stochastic DMSO framework [Foster et al., 2021], where the environment is stochastic, i.e.,  $M^1 = \cdots = M^T = M^* \in \mathcal{M}$ , and it can be specified by constraint  $\mathcal{P}^* = \{M^*\}$  and  $\mathscr{P}_{sto} = \{\{M^*\}: M^* \in \mathcal{M}\}$ , and
- adversarial DMSO framework [Foster et al., 2022b] (detailed in Appendix A.2), where the environment is fully adversarial, i.e., the constraint is  $\mathcal{P}^{\star} = \mathcal{M}$  and  $\mathscr{P}_{\mathsf{adv}} = \{\mathcal{M}\}$ .

Further examples of hybrid DMSO include SQ DMSO (Section 2.2), where the environment is allowed to respond to queries with values that are  $\tau$ -correct with respect to a ground truth model  $M^*$ , and robust DMSO (Section 2.4), where the environment is allowed to perturb the observation generated by a ground truth model  $M^*$  with a fixed probability  $\beta$ .

In addition to the constraints on the way the environment may change, the class  $\Phi$  of measurements encodes constraints on the learner, affecting the information the learner observes. For instance, in the examples studied in this paper, the measurements will take the form of allowed queries (Section 2.2) or differentially private channels (Section 2.3). Of course, the case of  $\mathcal{P}^* \subseteq (\Pi \to \Delta(\mathcal{O}))$  may be regarded as the trivial choice  $\Phi = \{id\}$  of identity measurement. **Learning objective.** In PAC learning, the goal of the learner is to select an *output decision*  $\pi_{T+1} \in \Pi$  after T rounds of interaction, with the performance measured by

$$\mathbf{Risk}_{\mathsf{DM}}(T) := \mathbb{E}_{\pi_{T+1} \sim \widehat{p}}[L(\mathcal{P}^{\star}, \pi_{T+1})], \tag{1}$$

where  $\pi_{T+1} \sim \hat{p}$  is the randomized decision of the learner,  $L: \mathscr{P} \times \Pi \to \mathbb{R}$  is a known loss function.

To simplify the presentation in this section, we mainly focus on the PAC formulation, deferring the study of regret to Section 2.5. Further, we present all the results in terms of a *metric-based* loss function, which is specified by a certain pseudo-metric structure over the decision space  $\Pi$ .

**Definition 1** (Metric-based loss function). A loss function  $L : \mathscr{P} \times \Pi \to \mathbb{R}$  is induced by a metric (or simply metric-based) if the decision space  $\Pi$  can be equipped with a pseudo-metric  $\rho$  such that  $L(\mathcal{P}, \pi) = \rho(\pi^{\mathcal{P}}, \pi)$ , where  $\mathcal{P} \mapsto \pi^{\mathcal{P}}$  is a map from  $\mathscr{P}$  to  $\Pi$ .

For many applications in statistics, the loss function is naturally metric-based, e.g., hypothesis testing and estimation [Casella and Berger, 2002].

**PAC hybrid DEC and guarantees.** For any hybrid DMSO problem specified by the constraint class  $\mathscr{P}$ , we define the hybrid DEC of  $\mathscr{P}$  with respect to a reference model  $\overline{M}$  as

$$\mathsf{p-dec}^{\mathsf{H}}_{\varepsilon}(\mathscr{P},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{\mathcal{P} \in \mathscr{P}} \left\{ \mathbb{E}_{\pi \sim p} L(\mathcal{P},\pi) \mid \inf_{M \in \operatorname{co}(\mathcal{P})} \mathbb{E}_{\pi \sim q} D^{2}_{\mathrm{H}}\left(M(\pi),\overline{M}(\pi)\right) \leq \varepsilon^{2} \right\}, \quad (2)$$

and  $\mathbf{p}-\mathsf{dec}_{\varepsilon}^{\mathsf{H}}(\mathscr{P}) = \sup_{\overline{M} \in \mathcal{M}^{+}} \mathbf{p}-\mathsf{dec}_{\varepsilon}^{\mathsf{H}}(\mathscr{P}, \overline{M})$ , where the supremum is taken over the class of reference models  $\mathcal{M}^{+} := \operatorname{co}(\cup_{\mathcal{P} \in \mathscr{P}} \mathcal{P})$ .

We now present the first result, which states that under the hybrid DMSO framework, hybrid DEC provides both lower and upper bounds for the minimax risk. The minimax risk quantifies the fundamental limit of learning, as it measures the best possible performance of an algorithm in the face of a worst-case environment constrained by  $\mathscr{P}$  (see Section 3 for details).

**Theorem 1** (PAC lower and upper bounds; Informal). Let  $T \ge 1$ , and L be metric-like. Under mild growth assumption, the following holds:

$$\mathrm{p-dec}^{\mathrm{H}}_{\underline{\varepsilon}(T)}(\mathscr{P}) \lesssim \inf_{\mathsf{Alg}} \sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \lesssim \mathrm{p-dec}^{\mathrm{H}}_{\bar{\varepsilon}(T)}(\mathscr{P}),$$

where  $\inf_{\mathsf{Alg}}$  is taken over all *T*-round algorithms  $\mathsf{Alg}$ ,  $\sup_{\mathsf{Env}}$  is taken over all environments  $\mathsf{Env}$  constrained by  $\mathscr{P}, \, \underline{\varepsilon}(T) \asymp \frac{1}{\sqrt{T}}, \, \overline{\varepsilon}(T) \asymp \sqrt{\frac{\log|\mathscr{P}|}{T}}, \, and \, we \, omit \, poly-logarithmic \, factors.$ 

We note that the lower bound applies to the *stationary* adversaries, while the upper bound (achieved by  $ExO^+$ ) applies to arbitrary (adaptive) adversarial environments.

Let us now discuss the qualitative behavior of  $p-dec_{\varepsilon}^{\mathsf{H}}(\mathscr{P})$  with respect to the constraint class  $\mathscr{P}$ . To start, consider stochastic DMSO, where each constraint is given by a singleton  $\mathcal{P} = \{M\}$ . In this case, the infimum over  $M \in \operatorname{co}(\mathcal{P})$  disappears, recovering the definition of the original PAC DEC in Foster et al. [2023b] (see also Eq. (22)). As constraints become less stringent (informally,  $\mathcal{P}$ 's become larger), the value of the DEC increases as the Hellinger-based constraint becomes easier to satisfy. Similarly, constraints on the learner are also reflected in the Hellinger term through the amount of information the measurements provide, as will be evident in the forthcoming calculations.

In the rest of this section, we detail how both types of constraints result in the corresponding measures of complexity and the guarantees for the settings of query-based learning (Section 2.2), locally differentially private learning (Section 2.3), and robust decision making (Section 2.4).

# 2.2 Query-based learning

In query-based learning, the environment responds to the learner's measurements (or, queries) with answers that are close to the answer under the ground-truth model  $M^* : \Pi \times \Phi \to \mathcal{V}$ , and we recall that we denote  $\mathbf{\Pi} := \Pi \times \Phi$ .

We formulate the interaction protocol of ( $\tau$ -correct) SQ DMSO as follows. For each  $t = 1, \dots, T$ :

- The learner selects a decision  $\pi_t \in \Pi$  and a measurement  $\phi_t \in \Phi$ .
- The environment selects (possibly adversarially)  $v_t \in \mathcal{V}$  such that  $||v_t M^*(\pi_t, \phi_t)|| \leq \tau$  and reveals  $v_t$  to the learner, where  $\mathcal{V}$  is a fixed normed vector space, and  $\tau \geq 0$  is a known tolerance parameter.

In SQ DMSO, the underlying model  $M^*$  is a *deterministic* map  $\Pi \times \Phi \to \mathcal{V}$ , and the learner is assumed to have access to a known model class  $\mathcal{M} \subseteq (\Pi \times \Phi \to \mathcal{V})$  that contains  $M^{*,1}$  After T rounds of interaction, the learner selects an output decision  $\pi_{T+1} \sim \hat{p}$  and incurs the PAC risk

$$\mathbf{Risk}_{\mathsf{DM}}(T) := \mathbb{E}_{\widehat{p}}[L(M^{\star}, \pi_{T+1})], \tag{3}$$

where  $L: \mathcal{M} \times \Pi \to \mathbb{R}$  is a given loss function. This formulation encompasses the commonly studied *Statistical Query* (SQ) learning [Kearns, 1998] and its various variants [Bshouty and Feldman, 2002, Feldman, 2017, etc.]. Further examples are detailed in Section 4.

The setting we just described combines constraints on both the learner and the adversary. Indeed, the class  $\Phi$  represents constraints on the decision maker, limiting the information it receives. Since answers to the measurements may be imprecise (up to the tolerance level  $\tau$ ), the interaction can be modeled as decision making with a constrained adversary. Before we discuss the details of specializing the hybrid DMSO framework, we first present the definition of the DEC specific to query-based learning and its main guarantees.

**SQ DEC.** For a given model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \mathcal{V})$  and a (randomized) reference model  $\overline{M} \in \mathcal{M}^+$ , we define the SQ DEC at  $\overline{M}$  as

$$\mathsf{p-dec}_{\varepsilon}^{\tau-\mathsf{SQ}}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M,\pi)] \mid \mathbb{P}_{\pi \sim q, \bar{v} \sim \overline{M}(\pi)}(\|M(\pi) - \bar{v}\| > \tau) \le \varepsilon^2 \right\}.$$
(4)

We further define the SQ DEC of  $\mathcal{M}$  as  $\mathsf{p-dec}_{\varepsilon}^{\tau-\mathsf{SQ}}(\mathcal{M}) = \sup_{\overline{M}} \mathsf{p-dec}_{\varepsilon}^{\tau-\mathsf{SQ}}(\mathcal{M}, \overline{M})$ , where the supremum is taken over all randomized reference models  $\overline{M} : \Phi \to \Delta(\mathcal{V})$ .

For query-based learning, our main result is given by the following theorem:

**Theorem 2** (SQ DEC lower and upper bounds; Informal). Let  $T \ge 1$ ,  $\mathcal{M}$  be a given model class, and the loss function L be metric-based. Then under certain growth conditions, it holds that

$$\mathsf{p}\text{-}\mathsf{dec}_{\underline{\varepsilon}(T)}^{\tau\text{-}\mathsf{SQ}}(\mathcal{M}) \lesssim \inf_{\mathsf{Alg}} \sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \lesssim \mathsf{p}\text{-}\mathsf{dec}_{\overline{\varepsilon}(T)}^{\tau\text{-}\mathsf{SQ}}(\mathcal{M}),$$

where  $\sup_{\mathsf{Env}}$  is taken over all environments satisfying query correctness with tolerance  $\tau$  for a model  $M^{\star} \in \mathcal{M}, \, \underline{\varepsilon}(T) \asymp \frac{1}{\sqrt{T}}, \, \overline{\varepsilon}(T) \asymp \sqrt{\frac{\log |\mathcal{M}|}{T}}.$ 

<sup>&</sup>lt;sup>1</sup>The class of all stochastic models is given by  $(\Pi \to \Delta(\mathcal{V}))$ , corresponding to noisy responses. We regard  $\mathcal{M} \subset (\Pi \to \Delta(\mathcal{V}))$ .

From hybrid DMSO to SQ DMSO. To frame the ( $\tau$ -correct) SQ DMSO within hybrid DMSO, we can consider the constraint  $\mathcal{P}_{M^*}$  specified by a model  $M^* \in \mathcal{M}$ :

$$\mathcal{P}_{M^{\star}} := \{ M \in \mathcal{M}^+ : \forall \boldsymbol{\pi} \in \boldsymbol{\Pi}, \forall v \in \operatorname{supp}(M(\boldsymbol{\pi})), \| v - M^{\star}(\boldsymbol{\pi}) \| \le \tau \},$$
(5)

and the constraint class corresponding to  $\mathcal{M}$  is given by  $\mathscr{P}_{\tau\text{-query}} = \{\mathcal{P}_{M^{\star}} : M^{\star} \in \mathcal{M}\}$ , with loss function  $L(\mathcal{P}_{M^{\star}}, \pi) := L(M^{\star}, \pi)$ .

While our characterization of query-based learning (Theorem 2) is derived by a direct proof (cf. Appendix H), we can also obtain it by applying Theorem 1. Specifically, under the above choice (5), for any model  $M \in \mathcal{M}$ , we have

$$\inf_{M'\in\operatorname{co}(\mathcal{P}_M)} \mathbb{E}_{\boldsymbol{\pi}\sim q} D_{\mathrm{H}}^2\left(M'(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})\right) \asymp \mathbb{P}_{\boldsymbol{\pi}\sim q, v \sim \overline{M}(\boldsymbol{\pi})}(\|M(\boldsymbol{\pi}) - v\| > \tau),$$

where  $\approx$  here means lower and upper bounds up to constant factors (cf. Lemma E.4). Hence,

$$\mathrm{p\text{-}dec}_{\varepsilon/2}^{\tau\operatorname{-}\mathrm{SQ}}(\mathcal{M}) \leq \mathrm{p\text{-}dec}_{\varepsilon}^{\mathrm{H}}(\mathscr{P}_{\tau\operatorname{-}\mathsf{query}}) \leq \mathrm{p\text{-}dec}_{\varepsilon}^{\tau\operatorname{-}\mathrm{SQ}}(\mathcal{M}).$$

Therefore, under SQ DMSO, the hybrid DEC is equivalent to the SQ DEC, and the general guarantees of Theorem 1 apply. Details are postponed to Appendix E.3.1.

### 2.3 Locally differentially private learning

The second example of hybrid DMSO is locally differentially private (LDP) learning. We first define the differentially private (DP) channels as follows.

**Definition 2** (Differentially private channels). For the latent observation space Z and the noisy observation space  $\mathcal{O}$ , a channel Q is a (measurable) map from  $Z \to \Delta(\mathcal{O})$ . A channel Q is  $\alpha$ -DP if for  $z, z' \in Z$  and any measurable set  $E \subseteq \mathcal{O}$ ,

$$\mathsf{Q}(E|z) \le e^{\alpha} \mathsf{Q}(E|z').$$

For a fixed pair  $(\mathcal{Z}, \mathcal{O})$  of spaces, we denote by  $\mathcal{Q}_{\alpha}$  the class of all  $\alpha$ -DP channels. To simplify the presentation, we assume that  $\alpha \leq \alpha_0$  for a pre-specified universal constant  $\alpha_0$ , and we will hide dependence on  $\alpha_0$ . We also assume the observation space  $\mathcal{O}$  is non-trivial, i.e.,  $|\mathcal{O}| \geq 2$ .

**DMSO with local privacy constraint (Private DMSO).** We consider the following private variant of the DMSO framework, with the local privacy constraint formalized by a class of private channels Q. For each round t = 1, ..., T:

- The learner selects a decision  $\pi_t \in \Pi$  and a private channel  $Q_t \in \mathcal{Q}$ , where  $\Pi$  is the decision space.
- The environment generates  $z_t \in \mathcal{Z}$  sampled via  $z_t \sim M^*(\pi_t)$ , where  $\mathcal{Z}$  is the observation space.
- The learner receives a noisy observation  $o_t \in \mathcal{O}$  sampled via  $o_t \sim Q_t(\cdot | z_t)$ .

In private DMSO, the environment is stationary and specified by an underlying model  $M^* : \Pi \to \Delta(\mathcal{Z})$ , and the learner is assumed to have access to a known model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  that contains  $M^*$ . As such, private DMSO is encompassed by the stochastic DMSO framework.

In this paper, we focus on  $Q = Q_{\alpha}$ , the class of  $\alpha$ -DP channels. We call a *T*-round algorithm as preserving  $\alpha$ -LDP (or simply  $\alpha$ -LDP) if it is a learner in the above sense. This formulation is equivalent to the commonly studied model of *sequential LDP channel* [Duchi et al., 2018]. Detailed discussion is deferred to Appendix B.1. **Private PAC-DEC.** Let  $\mathcal{L} = (\mathcal{Z} \to [0, 1])$  be the class of functions from  $\mathcal{Z}$  to [0, 1]. For any  $\ell \in \mathcal{L}$ , we define the  $\ell$ -divergence between distributions  $P, Q \in \Delta(\mathcal{Z})$  as

$$\mathsf{D}_{\ell}(P,Q) := \left| \mathbb{E}_{z \sim P}[\ell(z)] - \mathbb{E}_{z \sim Q}[\ell(z)] \right|.$$
(6)

For a model class  $\mathcal{M}$  and a reference model  $\overline{M} \in co(\mathcal{M})$ , the convex hull of  $\mathcal{M}$ , we define private PAC-DEC at  $\overline{M}$  as

$$\mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi \times \mathcal{L})}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M,\pi)] \mid \mathbb{E}_{(\pi,\ell) \sim q} \mathsf{D}_{\ell}^{2}(M(\pi),\overline{M}(\pi)) \le \varepsilon^{2} \right\}, \quad (7)$$

and the private PAC-DEC of  $\mathcal{M}$  as  $\mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}) = \sup_{\overline{M} \in \mathsf{co}(\mathcal{M})} \mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}, \overline{M})$ . The  $\ell$ -divergence is a measure of closeness of two distributions that is weaker than the Hellinger distance from the DEC framework for non-private learning (cf. Eq. (22)). This divergence is closely connected to the notion of *statistical queries* (SQ), but we postpone this discussion until Section 4.3.

For learning with LDP constraints, the private PAC-DEC provides both lower and upper bounds for the expected risk, as stated in the following theorem.

**Theorem 3** (Private PAC-DEC lower and upper bounds; Informal). Let  $T \ge 1$ . If the loss function L is reward-based or metric-based, the following holds:

$$\mathsf{p}\text{-}\mathsf{dec}^{\mathsf{LDP}}_{\underline{\varepsilon}(T)}(\mathcal{M}) \lesssim \inf_{\mathsf{Alg}} \sup_{M \in \mathcal{M}} \mathbb{E}^{M,\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \lesssim \mathsf{p}\text{-}\mathsf{dec}^{\mathsf{LDP}}_{\bar{\varepsilon}(T)}(\mathcal{M}),$$

where  $\inf_{Alg}$  is taken over all T-round  $\alpha$ -LDP algorithms,  $\underline{\varepsilon}(T) \simeq \frac{1}{\sqrt{\alpha^2 T}}$ ,  $\overline{\varepsilon}(T) \simeq \sqrt{\frac{\log |\mathcal{M}|}{\alpha^2 T}}$ , and we omit poly-logarithmic factors.

**Applications.** By further specializing the above result, we provide concrete guarantees for various locally-private learning tasks, including regression (Section 5.2) and particularly linear regression (Section 5.3). Our lower and upper bounds also provide a tight characterization of the localminimax complexity under LDP (Section 6.1), recovering the characterization in Duchi and Ruan [2024]. We also provide regret guarantees under LDP constraint, with applications to contextual bandits (Section 5.5), where the contexts can be chosen adversarially by the environment. In particular, we derive a near-optimal  $\sqrt{T}$ -regret for linear contextual bandits with local privacy through the private DEC theory, settling the open problem of the optimal regret in this setting [Zheng et al., 2020, Han et al., 2021, Li et al., 2024].

From hybrid DMSO to private DMSO. For each model  $M : \Pi \to \Delta(\mathcal{Z}), M$  induces a map  $M^{\sharp} : \Pi \times \mathcal{Q} \to \Delta(\mathcal{O})$  given by  $M^{\sharp}(\pi, \mathbb{Q}) = \mathbb{Q} \circ M(\pi)$ , where for any channel  $\mathbb{Q} \in \mathcal{Q}$  and any distribution  $P \in \Delta(\mathcal{Z})$ , we denote  $\mathbb{Q} \circ P \in \Delta(\mathcal{O})$  to be the marginal distribution of o under  $z \sim P, o \sim \mathbb{Q}(\cdot|z)$ . Therefore, the private DMSO is encompassed by the hybrid DMSO with measurement class  $\Phi = \mathcal{Q}$  and constraint class  $\mathscr{P}_{\mathsf{LDP}} = \{\{M^{\sharp}\} : M \in \mathcal{M}\}$  induced by  $\mathcal{M}$ . Using the strong data-processing inequality (Proposition 20), for any distribution  $q \in \Delta(\Pi \times \mathcal{Q})$ , there exists a distribution  $q' \in \Delta(\Pi \times \mathcal{L})$ , such that

$$\mathbb{E}_{\boldsymbol{\pi} \sim q} D_{\mathrm{H}}^2 \left( M^{\sharp}(\boldsymbol{\pi}), \overline{M}^{\sharp}(\boldsymbol{\pi}) \right) \asymp \alpha^2 \mathbb{E}_{(\boldsymbol{\pi}, \ell) \sim q'} \mathsf{D}_{\ell}^2(M(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})),$$

where  $\approx$  denotes equivalence up to constant factors. Therefore, it holds that

$$\mathsf{p}\text{-}\mathsf{dec}^{\mathsf{LDP}}_{c_0\alpha\varepsilon}(\mathcal{M}) \leq \mathsf{p}\text{-}\mathsf{dec}^{\mathsf{H}}_{\varepsilon}(\mathscr{P}_{\mathsf{LDP}}) \leq \mathsf{p}\text{-}\mathsf{dec}^{\mathsf{LDP}}_{c_1\alpha\varepsilon}(\mathcal{M}),$$

where  $c_0, c_1 > 0$  are absolute constants. Details are deferred to Appendix E.3.2.

#### 2.4 Robust decision making

We now introduce the following formulation of decision making in the presence of adversarial contamination (or, *robust decision making*). We mainly focus on Huber's contamination model [Huber, 1965, Huber and Ronchetti, 2011], as the application to other types of contamination (e.g. model mis-specifications) is analogous.

**Robust DMSO.** Let  $\beta \in [0, 1]$  be a fixed rate of contamination. In robust DMSO, the interaction protocol is as follows. For each round  $t = 1, 2, \dots, T$ :

- The learner selects a decision  $\pi_t \in \Pi$  from the joint decision space.
- The environment generates  $o_t^{\star} \in \mathcal{O}$  sampled via  $o_t^{\star} \sim M^{\star}(\boldsymbol{\pi}_t)$ .
- With probability  $1 \beta$ , the environment reveals  $o_t = o_t^*$  to the learner. Otherwise, the environment selects  $o_t \in \mathcal{O}$  arbitrarily (potentially depending on the interactions up to round t).

Similar to private DMSO, we assume the ground truth model  $M^*$  belongs to a given model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \Delta(\mathcal{O}))$ . In the formulation above, the environment is allowed to be adaptive, making the learning task harder than the Huber contamination model [Huber, 1965, 1992], where the environment is *stationary*, i.e.,  $M^1 = \cdots = M^T = (1 - \beta)M^* + \beta M'$  for an arbitrary but fixed contamination model M' (cf. Definition 4). Indeed, the environment under the Huber contamination model falls within the purview of the stochastic DMSO framework. Further discussion is deferred to Appendix A.3.

To frame the above setting within hybrid DMSO, we can consider the constraint specified by a model  $M^* \in \mathcal{M}$ :

$$\mathcal{P}_{M^{\star}} := \left\{ (1 - \beta)M^{\star} + \beta M' : M' \in (\mathbf{\Pi} \to \Delta(\mathcal{O})) \right\},\tag{8}$$

and the constraint class (induced by  $\mathcal{M}$ ) as given by  $\mathscr{P}_{\beta\text{-Huber}} := \{\mathcal{P}_{M^*} : M^* \in \mathcal{M}\}$ , with loss function  $L(\mathcal{P}_{M^*}, \pi) = L(M^*, \pi)$ . Then, the robust DMSO described above is exactly hybrid DMSO with constraint class  $\mathscr{P}_{\beta\text{-Huber}}$ . By instantiating the general theory in Section 2.1, we arrive at the following (simpler) DEC formulation for robust decision making.

**Robust DEC.** For  $\beta \in [0,1]$  and distributions  $P, Q \in \Delta(\mathcal{O})$ , we consider the  $\beta$ -perturbed Hellinger divergence

$$D^{2}_{\beta-\mathsf{Huber}}(P,Q) := \inf_{P' \in \Delta(\mathcal{O})} D^{2}_{\mathrm{H}}\left((1-\beta)P + \beta P',Q\right).$$
(9)

For a model class  $\mathcal{M}$  and a reference model  $\overline{M}$ , we define robust DEC at  $\overline{M}$  as

$$\mathsf{p-dec}_{\varepsilon}^{\mathsf{R}}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M,\pi)] \mid \mathbb{E}_{\pi \sim q} D^{2}_{\beta-\mathsf{Huber}}(M(\pi),\overline{M}(\pi)) \leq \varepsilon^{2} \right\},$$
(10)

and the robust DEC of  $\mathcal{M}$  is then defined as  $\mathsf{p-dec}_{\varepsilon}^{\mathsf{R}}(\mathcal{M}) = \sup_{\overline{\mathcal{M}} \in \mathcal{M}^+} \mathsf{p-dec}_{\varepsilon}^{\mathsf{R}}(\mathcal{M}, \overline{\mathcal{M}}).$ 

In the definition of the robust DEC, we replace the Hellinger distance by the perturbed divergence (9), reflecting the fact that for a ground truth model M, the environment can vary M by a probability mass  $\beta$ . By definition, we know  $p-dec_{\varepsilon}^{\mathsf{R}}(\mathcal{M}) = p-dec_{\varepsilon}^{\mathsf{H}}(\mathscr{P}_{\beta-\mathsf{Huber}})$  for  $\varepsilon \in [0,1]$  (detailed in Appendix A.3). Therefore, as a direct corollary of Theorem 1, the robust DEC provides both lower and upper bounds for robust PAC learning. **Theorem 4** (Robust risk bounds; Informal). Let  $T \ge 1$ ,  $\mathcal{M}$  be a given model class, and the loss function L be metric-based. Then under certain growth conditions, it holds that

$$\mathrm{p-dec}^{\mathrm{R}}_{\underline{\varepsilon}(T)}(\mathcal{M}) \lesssim \inf_{\mathsf{Alg}} \sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \lesssim \mathrm{p-dec}^{\mathrm{R}}_{\bar{\varepsilon}(T)}(\mathcal{M})$$

where  $\sup_{\mathsf{Env}}$  is taken over all environments that are  $\beta$ -contaminated from a model  $M^* \in \mathcal{M}$ ,  $\underline{\varepsilon}(T) \asymp \frac{1}{\sqrt{T}}, \ \overline{\varepsilon}(T) \asymp \sqrt{\frac{\log |\mathcal{M}|}{T}}.$ 

The details and regret guarantees are presented in Appendix A.3.

## 2.5 Regret guarantees for hybrid DMSO

In this section, we study the no-regret learning goal under hybrid DMSO, and present the general regret guarantees and its application to smooth environments.

In the no-regret learning task, the performance of the learner is measured by the following notion of *regret*:

$$\operatorname{\mathbf{Reg}}_{\mathsf{DM}}(T) := \max_{\pi^{\star} \in \Pi} \sum_{t=1}^{T} V^{M^{t}}(\pi^{\star}) - \mathbb{E}_{\pi_{t} \sim q_{t}} V^{M^{t}}(\pi_{t}),$$
(11)

where for each model  $M, V^M : \Pi \to \mathbb{R}$  is an associated value function. In words,  $\operatorname{Reg}_{\mathsf{DM}}(T)$  measure the performance of the learner compared to the best decision in the hindsight. Note that due to the adversarial nature of the environment, the PAC risk (1) cannot be directly reduced from the regret (11) by the *online-to-batch* conversion.

We first extend the regret DEC [Foster et al., 2023b] to hybrid DMSO. For any model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \Delta(\mathcal{O}))$ , reference model  $\overline{M}$ , we define

$$\mathsf{r}\mathsf{-dec}^{c}_{\varepsilon}(\mathcal{M},\overline{M}) := \inf_{p \in \Delta(\mathbf{\Pi})} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[V^{M}(\pi^{M}) - V^{M}(\pi)] \mid \mathbb{E}_{\pi \sim p}D^{2}_{\mathrm{H}}\left(M(\pi),\overline{M}(\pi)\right) \leq \varepsilon^{2} \right\}, \quad (12)$$

where for each model M,  $\pi^{M} := \arg \max_{\pi \in \Pi} V^{M}(\pi)$  is an optimal decision under M. The regret DEC of  $\mathcal{M}$  is then defined as  $\mathsf{r-dec}^{c}_{\varepsilon}(\mathcal{M}) = \sup_{\overline{M} \in \mathrm{co}(\mathcal{M})} \mathsf{r-dec}^{c}_{\varepsilon}(\mathcal{M} \cup \{\overline{M}\}, \overline{M})$ .

Next, to define the regret DEC of a constraint class  $\mathscr{P}$ , we define

$$\mathcal{M}_{\mathscr{P}} := \bigcup_{\mathcal{P} \in \mathscr{P}} \operatorname{co}(\mathcal{P}), \qquad \mathsf{r-dec}_{\varepsilon}^{\mathsf{H}}(\mathscr{P}) := \mathsf{r-dec}_{\varepsilon}^{\mathsf{c}}(\mathcal{M}_{\mathscr{P}}).$$

We show that the regret DEC of  $\mathscr{P}$  provides both lower and upper bound for the minimax regret.

**Theorem 5** (Regret lower and upper bounds; Informal). Let  $T \ge 1$ . Under assumptions on the value function and the growth of the DEC, the following holds:

$$T \cdot \mathsf{r-dec}_{\underline{\varepsilon}(T)}^{c}(\mathcal{M}_{\mathscr{P}}) \lesssim \inf_{\mathsf{Alg Env}} \sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \lesssim T \cdot \mathsf{r-dec}_{\overline{\varepsilon}(T)}^{c}(\mathcal{M}_{\mathscr{P}}),$$

where  $\inf_{\mathsf{Alg}}$  is taken over all *T*-round algorithms  $\mathsf{Alg}$ ,  $\sup_{\mathsf{Env}}$  is taken over all environments  $\mathsf{Env}$  constrained by  $\mathscr{P}, \underline{\varepsilon}(T) \asymp \frac{1}{\sqrt{T}}, \ \overline{\varepsilon}(T) \asymp \sqrt{\frac{\log|\mathscr{P}| + \log|\Pi|}{T}}, \ and \ we \ omit \ poly-logarithmic \ factors.$ 

In particular, when the environment is fully adversarial,  $\mathscr{P} = \{\mathcal{M}\}$  is a singleton,  $\mathcal{M}_{\mathscr{P}} = \operatorname{co}(\mathcal{M})$ , and we recover the results of Foster et al. [2022b]. Furthermore, the log  $|\Pi|$  factor in our upper bound can further be tightened by the fractional covering number [Chen et al., 2024] (cf. Section 3.2).

As an application, we consider no-regret learning against smooth adversaries. The results for robust no-regret learning are deferred to Appendix A.3.

#### 2.5.1 Example: Smooth adversaries

Within the hybrid DMSO framework, we can also consider decision making with a smooth adversary. In this setting, we focus on the case where  $\Phi = {id}$ , i.e., only the identity measurement is considered.

For any two distributions  $P, Q \in \Delta(\mathcal{Z})$ , we define the density ratio between P, Q as

$$D_{\infty}(P \parallel Q) := \operatorname{ess\,sup} \frac{dP}{dQ}$$

We say P is  $\sigma_{sm}$ -smooth with respect to Q if  $D_{\infty}(P \parallel Q) \leq \frac{1}{\sigma_{sm}}$ .

For the setting of smooth adversary, we assume there is a known subclass  $\mathcal{M}_{\mathsf{base}} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ , such that the adversary is constrained to fix a *base model*  $M_{\mathsf{base}}$  ahead of the interaction and without revealing it to the learner, and then choose each  $M^t$  that is  $\sigma_{\mathsf{sm}}$ -smooth with respect to  $M_{\mathsf{base}}$ . Specifically, for each *base model*  $M_{\mathsf{base}} \in \mathcal{M}_{\mathsf{base}}$ , the constraint specified by  $M_{\mathsf{base}}$  is

$$\mathcal{P}_{M_{\mathsf{base}}} := \left\{ M \in \mathcal{M} : \sup_{\pi \in \Pi} D_{\infty}(M(\pi) \parallel M_{\mathsf{base}}(\pi)) \le \frac{1}{\sigma_{\mathsf{sm}}} \right\},\tag{13}$$

which is the class of all models that are  $\sigma_{sm}$ -smooth with respect to the base model  $M_{base}$ . Specifying the hybrid DMSO framework with  $\mathscr{P} = \{\mathcal{P}_{M_{base}} : M_{base} \in \mathcal{M}_{base}\}$ , we generalize the standard smooth online learning setting to interactive decision making.

Note that for each  $M_{\text{base}}$ , the class  $\mathcal{P}_{M_{\text{base}}}$  is convex, and hence we let

$$\mathcal{M}_{\mathsf{sm}} \mathrel{\mathop:}= \mathcal{M}_{\mathscr{P}} = \bigcup_{M_{\mathsf{base}} \in \mathcal{M}_{\mathsf{base}}} \mathcal{P}_{M_{\mathsf{base}}}$$

It is a direct corollary of Theorem 5 that the regret DEC of  $\mathcal{M}_{sm}$  provides both lower and upper bounds for no-regret learning against smooth adversaries.

**Theorem 6** (Regret bounds against smooth adversaries; Informal). Let  $T \ge 1$ . Under assumptions on the value function and the growth of the DEC, the following holds:

$$T \cdot \mathsf{r-dec}_{\underline{\varepsilon}(T)}^{\mathrm{c}}(\mathcal{M}_{\mathsf{sm}}) \lesssim \inf_{\mathsf{Alg }} \sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \lesssim T \cdot \mathsf{r-dec}_{\overline{\varepsilon}(T)}^{\mathrm{c}}(\mathcal{M}_{\mathsf{sm}}),$$

where  $\sup_{\mathsf{Env}}$  is taken over all environments  $\mathsf{Env}$  constrained to be  $\sigma_{\mathsf{sm}}$ -smooth with respect to a base model  $M_{\mathsf{base}} \in \mathcal{M}_{\mathsf{base}}, \, \underline{\varepsilon}(T) \asymp \frac{1}{\sqrt{T}}, \, \overline{\varepsilon}(T) \asymp \sqrt{\frac{\log |\mathcal{M}_{\mathsf{base}}| + \log |\Pi|}{T}}.$ 

# **3** DEC Theory for Hybrid DMSO

In this section, we present the details of the DEC theory for hybrid DMSO. Before proceeding to the main results, we rigorously formulate the notion of *algorithms* and *environments*.

A *T*-round algorithm Alg is specified by a sequence of mappings  $\{q_t\}_{t\in[T]} \cup \{\hat{p}\}$ , where the *t*-th mapping  $q_t(\cdot \mid \mathcal{H}_{t-1})$  specifies the distribution of  $\pi_t = (\pi_t, \phi_t)$  based on the history  $\mathcal{H}_{t-1} = (\pi_s, o_s)_{s \leq t-1}$ , and the final map  $\hat{p}(\cdot \mid \mathcal{H}_T)$  specifies the distribution of the *output decision*  $\pi_{T+1}$  based on  $\mathcal{H}_T$ . Similarly, a *T*-round *adaptive* environment Env is specified by a sequence of mappings  $\{\mu^t\}_{t\in[T]}$ , where the *t*-th mapping  $\mu^t(\cdot \mid \mathcal{H}'_{t-1})$  specifies the distribution of the model  $M^t$  based on the full-information history  $\mathcal{H}'_{t-1} = (M^s, \pi_s, o_s)_{s \leq t-1}$ . An environment is constrained by  $\mathscr{P}$  if

there exists  $\mathcal{P}^{\star} \in \mathscr{P}$  such that  $\mu^{t}(\cdot \mid \mathcal{H}'_{t-1})$  is always supported on  $\mathcal{P}^{\star}$  for all  $t \in [T]$ . As already discussed, each model  $M \in (\mathbf{\Pi} \to \Delta(\mathcal{O}))$  corresponds to a stationary environment, which chooses  $M^{1} = \cdots = M^{T} = M$  deterministically.

For any algorithm Alg and environment Env, we let  $\mathbb{P}^{\mathsf{Env},\mathsf{Alg}}(\cdot)$  to be the distribution of  $(\mathcal{H}_T, \pi_{T+1})$ generated by the algorithm Alg under the environment Env, and let  $\mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\cdot]$  to be the corresponding expectation. In particular, for any model M, we let  $\mathbb{P}^{M,\mathsf{Alg}}(\cdot)$  to be the distribution of  $(\mathcal{H}_T, \pi_{T+1})$ generated by the algorithm Alg under the stationary environment that chooses  $M^t = M$  for  $t \in [T]$ , and let  $\mathbb{E}^{M,\mathsf{Alg}}[\cdot]$  to be the corresponding expectation.

**Miscellaneous notation.** For a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{O}))$ , a finite subset  $\mathcal{M}_0 \subseteq \mathcal{M}$  is an  $\varepsilon$ -covering of  $\mathcal{M}$  if for any model  $M \in \mathcal{M}$ , there exists  $M' \in \mathcal{M}_0$  such that  $D_{\mathrm{H}}(M(\pi), M'(\pi)) \leq \varepsilon, \forall \pi \in \Pi$ . We define  $N(\mathcal{M}, \mathcal{M})$ , the  $\varepsilon$ -covering number of  $\mathcal{M}$ , to be the minimal cardinality of the  $\varepsilon$ -coverings of  $\mathcal{M}$ .

For the upper bounds in this section, we assume the model class  $\mathcal{M} = \mathcal{M}_{\mathscr{P}}$  admits finite  $\varepsilon$ -covering for any  $\varepsilon > 0$ , ensuring that the Minimax theorem can be applied.

Assumption 2 (Compactness of the model class). For any  $\Delta > 0$ , the covering number  $N(\mathcal{M}, \Delta)$  is finite.

Further, to simplify the presentation, we consider the following growth condition (following Foster et al. [2023b], Chen et al. [2024]).

**Definition 3** (Moderate decay). A function  $d : [0,1] \to \mathbb{R}_{\geq 0}$  is of moderate decay if there exists a constant  $c \geq 1$  such that  $c \frac{d(\varepsilon)}{\varepsilon} \geq \frac{d(\varepsilon')}{\varepsilon'}$  for all  $\varepsilon' \geq \varepsilon$ .

For many problems of interest, the DECs grow as  $C\varepsilon^{\rho}$  with  $\rho \leq 1$  and is automatically of moderate decay (for details, see e.g. Foster et al. [2023b]).

## 3.1 Guarantees for PAC learning

**PAC DEC lower bounds.** To better illustrate the key observation for the hybrid DEC lower bounds, we first introduce the notion of the *stationary* adversary.

**Definition 4.** For an environment Env constrained by  $\mathscr{P}$ , Env is stationary if there exists  $\mathcal{P}^* \in \mathscr{P}$ and  $\mu \in \Delta(\mathcal{P}^*)$  such that for each step  $t \in [T]$ , the model  $M^t$  is chosen as  $M^t \sim \mu$ .

In other words, in an stationary environment, the model  $M^t \sim \mu$  is chosen independently of prior interactions. The key observation of Foster et al. [2022b] is that lower bounds for adversarial DMSO can implied by the stochastic lower bounds [Foster et al., 2021, 2023b] by considering stationary environments, as stationary environments can be described by stochastic DMSO. This argument also applies to hybrid DMSO, implying the following lower bounds. The proof is deferred to Appendix E.1.

**Theorem 7** (Hybrid DEC lower bound for PAC risk). Suppose that L is metric-like. Then, for any T-round algorithm,

$$\sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \ge \frac{1}{8}\mathsf{p-dec}_{\underline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P}), \tag{14}$$

where  $\underline{\varepsilon}(T) = \frac{1}{20\sqrt{T}}$  and the supremum is taken over stationary environments.

Furthermore, for general loss function  $L : \mathscr{P} \times \Pi \to [0,1]$ , any T-round algorithm Alg, parameter  $\delta \in (0,1]$ , it holds that

$$\sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \ge \mathsf{p-dec}_{\underline{\varepsilon}_{\delta}(T)}^{\mathsf{H}}(\mathscr{P}) - \delta, \tag{15}$$

where  $\underline{\varepsilon}_{\delta}(T) := \frac{1}{13} \sqrt{\frac{\delta}{T}}$ .

We now briefly discuss the two lower bounds in Theorem 7. Eq. (14) is stated for metric-based loss, and it nearly matches the upper bound provided in Theorem 8 (with a  $\log |\mathscr{P}|$ -gap). On the other hand, Eq. (15) is stated for *any* general loss function (without requiring metric structure) and it is looser. It can be further re-written as

$$\sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \ge \frac{1}{2}\mathsf{p}-\mathsf{dec}_{\underline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P}), \qquad \text{where } T_{\underline{\varepsilon}}(T)^{2} \asymp \mathsf{p}-\mathsf{dec}_{\underline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P}). \tag{16}$$

For a problem with  $\mathbf{p}\text{-dec}_{\varepsilon}^{\mathsf{H}}(\mathscr{P}) \simeq \varepsilon$ , Eq. (16) gives a lower bound of  $\Omega\left(\frac{1}{T}\right)$ . While this is worse than the  $\Omega\left(\frac{1}{\sqrt{T}}\right)$  lower bound provided by (14) under metric-based loss, such a worse lower bound can be tight for certain problems (as shown in Foster et al. [2023a]). We also note that under stochastic DMSO and *reward-based* loss function, a tighter lower bound similar to (14) can also be derived [Foster et al., 2023b] (see also Appendix E.1).

**PAC DEC upper bounds.** Next, we present the upper bound provided by  $ExO^+$  (Algorithm 1) as follows. The description of  $ExO^+$  is deferred to Appendix F. For the simplicity of presentation, we still assume that the loss function is metric-like. While  $ExO^+$  is able to handle more general problems (and in particular reward-based loss function), we defer these details to Appendix F.

**Theorem 8** (Hybrid DEC upper bound for PAC risk). Let  $T \ge 1$ ,  $\delta \in (0,1)$ , and  $\mathscr{P}$  be given. Suppose that L is metric-like,  $\mathcal{M}_{\mathscr{P}}$  is compact (Assumption 2), and the hybrid DEC  $p-dec_{\varepsilon}^{\mathsf{H}}(\mathscr{P})$ is of moderate decay (Definition 3). Then  $\mathsf{ExO}^+$  can be suitably instantiated (as detailed in Appendix F.4.1), such that in any environment constrained by  $\mathscr{P}$ ,  $\mathsf{ExO}^+$  achieves with probability at least  $1 - \delta$  that

$$\mathbf{Risk}_{\mathsf{DM}}(T) \lesssim \mathsf{p-dec}_{\overline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P}),$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log(|\mathscr{P}|/\delta)}{T}}$ .

Furthermore, for any  $\Delta > 0$ , any general loss function L bounded in [0,1],  $\mathsf{ExO}^+$  can be suitably instantiated so that in any environment constrained by  $\mathscr{P}$ ,  $\mathsf{ExO}^+$  achieves with probability at least  $1 - \delta$  that

$$\mathbf{Risk}_{\mathsf{DM}}(T) \lesssim \mathsf{p-dec}_{\bar{\varepsilon}(\Lambda T)}^{\mathsf{H}}(\mathscr{P}) + \Delta.$$
(17)

#### 3.2 Guarantees for no-regret learning

In this section, we consider no-regret learning in hybrid DMSO. To present the DEC theory in its simplest form, we make the following assumption, which essentially requires that the value of any decision can be estimated from observations.

**Assumption 3** (Observability). For any decision  $\pi \in \Pi$ , the map  $M \mapsto V^M(\pi) \in [0, V_{\max}]$  is linear over  $\mathcal{M}_{\mathscr{P}}$ , and there exists a measurement  $\phi_{\pi} \in \Phi$ , such that for  $\pi = (\pi, \phi_{\pi}) \in \Pi$ ,

$$\left|V^{M}(\pi) - V^{\overline{M}}(\pi)\right| \leq C_{V} D_{\mathrm{H}}\left(M(\pi), \overline{M}(\pi)\right), \qquad \forall M \in \mathcal{M}_{\mathscr{P}}, \overline{M} \in \mathrm{co}(\mathcal{M}_{\mathscr{P}}).$$
(18)

Assumption 3 can also be regarded as a continuity assumption on the value function.

To better illustrate Assumption 3, we consider the example of identity measurement and rewardbased value function. This setting is extensively studied in Foster et al. [2021, 2022b, 2023b, etc.].

**Example 1** (Reward-based learning). Suppose that  $\mathscr{P}$  is induced by a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ , where  $\mathcal{Z} \subseteq \mathcal{O}$  and the measurement class is  $\Phi = \{\mathrm{id}\}$ , i.e. we overload the notation and write  $M(\pi, \mathrm{id}) = M(\pi)$  for each model  $M \in \mathcal{M}$ . In this setting, the value function V is reward-based, if there is a known reward function  $R : \mathcal{Z} \times \Pi \to [0, 1]$  such that  $V^{\mathcal{M}}(\pi) = \mathbb{E}^{M, \pi}[R(z, \pi)]$ .

This formulation encompasses many learning settings of interest, including bandits and contextual bandits, online control, reinforcement learning, etc. (for examples, see e.g. Foster et al. [2021]). In this setting, Assumption 3 holds with  $C_V = \sqrt{2}$ . We also note that for reward-based LDP learning, Assumption 3 holds with  $C_V = O(\frac{1}{\alpha})$  (as detailed in Appendix E.3.2).

**Regret lower bound.** With Assumption 3, we now present the main regret lower bound.

**Theorem 9** (Hybrid DEC lower bound for regret). Let  $T \ge 1$ ,  $\mathscr{P}$  be a given constraint class. Suppose that Assumption 3 holds for the value function V. Then, for any T-round algorithm Alg,

$$\sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \ge \sup_{\mathsf{Env}} \sup_{\pi^{\star} \in \Pi} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}\left[\sum_{t=1}^{T} V^{M^{t}}(\pi^{\star}) - V^{M^{t}}(\pi_{t})\right]$$
(19)

$$\geq \frac{T}{8} \left( \mathsf{r-dec}^{c}_{\underline{\varepsilon}(T)}(\mathcal{M}_{\mathscr{P}}) - 6C_{V\underline{\varepsilon}}(T) - \frac{V_{\max}}{T} \right)$$
(20)

where the supremum is taken over stationary environments  $\mathsf{Env}$  constrained by  $\mathscr{P}$ , and  $\underline{\varepsilon}(T) = \frac{1}{24\sqrt{T}}$ .

Similar to Theorem 7, the above regret lower bound is also proven through a reduction to the stochastic setting by considering stationary environments (detailed in Appendix E.2).

**Regret upper bound.** Before presenting the upper bound, we first introduce the notion of the *fractional covering number* [Chen et al., 2024], which captures the complexity of the decision space  $\Pi$  with respect to the class of models  $\mathcal{M}$ .

**Definition 5** (Fractional covering number). For a learning problem  $(\mathcal{M}, \Pi, L)$  and parameter  $\Delta \geq 0$ , we define the fractional covering number as

$$N_{\mathsf{frac}}(\mathcal{M}, \Delta) := \inf_{p \in \Delta(\Pi)} \sup_{M \in \mathcal{M}} \frac{1}{p(\pi : L(M, \pi) \le \Delta)}.$$
(21)

We show that the regret of  $ExO^+$  can be upper bounded in terms of the regret DEC, the fractional covering number, and log  $|\mathscr{P}|$ .

**Theorem 10** (Hybrid DEC upper bound for regret). Let  $T \ge 1$ ,  $\delta \in (0,1)$ , and  $\mathscr{P}$  be given. Suppose that  $\mathcal{M}_{\mathscr{P}}$  is compact, Assumption 3 holds for the value function V, and the regret DEC  $\mathsf{r}\operatorname{-dec}^{\mathsf{c}}_{\varepsilon}(\mathcal{M}_{\mathscr{P}})$  is of moderate decay. Then, in any environment constrained by  $\mathscr{P}$ ,  $\mathsf{ExO}^+$  (instantiated as detailed in Appendix F.4.2) achieves with probability at least  $1 - \delta$  that

$$\frac{1}{T} \operatorname{Reg}_{\mathsf{DM}}(T) \leq \Delta + O(\sqrt{\log T}) \cdot \Big[ \operatorname{r-dec}_{\bar{\varepsilon}(T)}^{c}(\mathcal{M}_{\mathscr{P}}) + C_{V}\bar{\varepsilon}(T) \Big],$$
  
where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log(|\mathscr{P}|/\delta) + \log N_{\operatorname{frac}}(\mathcal{M}_{\mathscr{P}}, \Delta)}{T}}.$ 

Finally, we remark that both our lower and upper bounds extend beyond Assumption 3, as detailed in Appendix E.2 and Appendix F.

### 3.3 Implication: Tighter bounds for convex classes

Our results for hybrid DMSO also have interesting implications for stochastic DMSO. To begin with, we recall that for a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{O}))$  under stochastic DMSO, the PAC DEC is defined as

$$\mathsf{p-dec}^{\mathsf{c}}_{\varepsilon}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M,\pi)] \mid \mathbb{E}_{\pi \sim q} D^{2}_{\mathrm{H}}\left(M(\pi),\overline{M}(\pi)\right) \leq \varepsilon^{2} \right\},$$
(22)

and  $p-dec_{\varepsilon}^{c}(\mathcal{M}) := \sup_{\overline{M} \in co(\mathcal{M})} p-dec_{\varepsilon}^{c}(\mathcal{M}, \overline{M})$ . DEC theory [Foster et al., 2021, 2023b] provides the following characterization (omitting logarithmic factors):

$$\mathsf{p-dec}^{c}_{\underline{\varepsilon}(T)}(\mathcal{M}) \lesssim \inf_{\mathsf{Alg}} \sup_{M \in \mathcal{M}} \mathbb{E}^{M,\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \lesssim \mathsf{p-dec}^{c}_{\overline{\varepsilon}(T)}(\mathcal{M}), \tag{23}$$

under certain regularity assumptions on the loss function, where  $\underline{\varepsilon}(T) \approx \frac{1}{\sqrt{T}}$ ,  $\overline{\varepsilon}(T) \approx \sqrt{\frac{\log |\mathcal{M}|}{T}}$ . Therefore, a log  $|\mathcal{M}|$ -gap remains between the known DEC lower and upper bounds, corresponding to the complexity of *estimation*, as noted by Chen et al. [2024]. The log  $|\mathcal{M}|$  factor can be undesirable for many applications beyond model-based learning.

Interestingly, it turns out the log  $|\mathcal{M}|$  factor can be replaced by a smaller quantity, potentially at the price of degradation in the DEC term. To illustrate this, we start with the *hypothesis selection* problem, which is a generalization of the standard, non-interactive hypothesis testing problem. For example, the setting below encompasses LDP hypothesis selection, where  $\Phi = \mathcal{Q}$  is the class of  $\alpha$ -DP channels, and  $\mathcal{M}$  is induced by a class of distributions over  $\Delta(\mathcal{Z})$ .

**Example 2** (Interactive hypothesis selection). Given a DMSO model class  $\mathcal{M} \subseteq (\Phi \to \Delta(\mathcal{O}))$ , a hypothesis selection problem is described by a partition

$$\mathcal{M} = \bigsqcup_{i=1}^{m} \mathcal{M}^{(i)},$$

where  $\mathcal{M}^{(1)}, \dots, \mathcal{M}^{(m)}$  are disjoint subclasses. The decision space is  $\Pi = [m]$ , and for each  $M \in \mathcal{M}$ ,  $\pi \in \Pi$ , the loss function is given by  $L(M, \pi) = \mathbf{1} \{ \pi \neq \pi^M \}$ , where  $\pi^M$  is the unique index  $i \in [m]$  such that  $M \in \mathcal{M}^{(i)}$ .

While we can frame the hypothesis selection problem within stochastic DMSO (with  $\mathscr{P}_{sto}$  corresponding to  $\mathcal{M}$ ), the upper bound provided by DEC theory scales with  $\log |\mathcal{M}|$ , the complexity of model class, which is undesirable. On the other hand, when the subclasses  $\mathcal{M}^{(1)}, \dots, \mathcal{M}^{(m)}$  are convex, we can alternative frame this problem within hybrid DMSO, with  $\mathscr{P}_m = \{\mathcal{M}^{(1)}, \dots, \mathcal{M}^{(m)}\}$  and loss function  $L(\mathcal{M}^{(i)}, \pi) = \mathbf{1} \{\pi \neq i\}$ . With such specifications, we allow the environment to be adaptive (within a fixed underlying model class  $\mathcal{M}^{(i)}$ ), while

$$\operatorname{p-dec}_{\varepsilon}^{\operatorname{H}}(\mathscr{P}_m) = \operatorname{p-dec}_{\varepsilon}^{\operatorname{c}}(\mathcal{M}), \quad \forall \varepsilon > 0.$$

Therefore, Theorem 8 implies the following tighter upper bound for hypothesis selection.

**Proposition 11** (ExO<sup>+</sup> for convex hypothesis selection). Let  $T \ge 1$ ,  $\delta \in (0,1)$ . In Example 2, suppose that  $\mathcal{M}$  is compact,  $\mathcal{M}^{(1)}, \dots, \mathcal{M}^{(m)}$  are convex, and

$$\mathsf{p-dec}^{\mathrm{c}}_{\bar{\varepsilon}(T)}(\mathcal{M}) \leq \frac{1}{3}, \qquad \bar{\varepsilon}(T) = 8\sqrt{\frac{\log(m/\delta)}{T}}.$$

Then  $\mathsf{E} \times \mathsf{O}^+$  can be suitably instantiated (on the constraint class  $\mathscr{P}_m$ , as detailed in Appendix F.4.3), so that under any model  $M^* \in \mathcal{M}_{i^*}$ , the algorithm returns  $\pi_{T+1} = i^*$  with probability at least  $1 - \delta$ .

In the above example, Theorem 8 naturally provides a tighter bound by considering hybrid DMSO and replacing the log  $|\mathcal{M}|$ -factor by log m. In general, such conversion will result in a degradation in the DEC term, if the model class is non-convex. In the following, we will make this trade-off precise.

Bounds for interactive estimation. In the interactive estimation task, the decision space  $\Pi$  is equipped with a pseudo-metric  $\rho$ , and a map  $M \mapsto \pi^M$  is given such that  $L(M, \pi) = \rho(\pi^M, \pi)$ . To apply the idea described above, we fix a parameter  $\Delta \geq 0$  and consider the constraint set specified by a  $\pi \in \Pi$ :

$$\mathcal{M}_{\pi} := \{ M : \rho(\pi^M, \pi) \le \Delta \},\$$

and the corresponding constraint class is  $\mathscr{P} = \{\mathcal{M}_{\pi} : \pi \in \Pi_{\Delta}\}$ , with  $\Pi_{\Delta}$  being a  $\Delta$ -covering of the set  $\Pi_{\mathcal{M}} := \{\pi^{M} : \pi \in \Pi\}$ . Then,  $\mathcal{M} = \bigcup_{\mathcal{P} \in \mathscr{P}} \mathcal{P}$ , and hence we can apply Theorem 8. Furthermore, assuming that  $\mathcal{M}$  is convex and  $M \mapsto \rho(\pi^{M}, \pi)$  quasi-convex for any  $\pi \in \Pi$ , then we can show that

$$\mathsf{p}\operatorname{-dec}^{\mathsf{H}}_{\varepsilon}(\mathscr{P}) \leq \Delta + \mathsf{p}\operatorname{-dec}^{\mathsf{c}}_{\varepsilon}(\mathcal{M}).$$

Therefore, Theorem 8 implies the following guarantee for interactive estimation (for affine functionals).

**Proposition 12** (ExO<sup>+</sup> for interactive estimation). Let  $T \ge 1$ ,  $\delta \in (0,1)$ ,  $\Delta \ge 0$ . Suppose that  $\mathcal{M}$  is convex,  $\Pi$  is a subset of a normed vector space, and an affine map  $\mathcal{M} \mapsto \pi^{\mathcal{M}}$  is given such that  $L(\mathcal{M}, \pi) = \|\pi^{\mathcal{M}} - \pi\|$ . Further assume that the DEC p-dec<sup>c</sup><sub> $\varepsilon$ </sub>( $\mathcal{M}$ ) is of moderate decay. Then ExO<sup>+</sup> can be suitably instantiated so that with probability at least  $1 - \delta$ , it returns  $\pi_{T+1}$  with

$$\mathbf{Risk}_{\mathsf{DM}}(T) = \rho(\pi^M, \pi_{T+1}) \lesssim \Delta + \mathsf{p-dec}^{\mathsf{c}}_{\bar{\varepsilon}(T)}(\mathcal{M}),$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log N_{\|\cdot\|}(\Pi, \Delta) + \log(1/\delta)}{T}}$ , and  $N_{\|\cdot\|}(\Pi, \Delta)$  is the  $\Delta$ -covering number of  $\Pi$  under the norm  $\|\cdot\|$ .

In particular, for bounded functional estimation,  $\Pi = [0, 1]$ , we have  $\log N_{|\cdot|}(\Pi, \Delta) = \log(1/\Delta) + O(1)$ , and hence the minimax risk of interactive functional estimation is characterized by the DEC up to logarithmic factors. This upper bound generalizes the results of Polyanskiy and Wu [2019] for *non-interactive* linear functional estimation with a convex model class.

Bounds for reward-based learning. Generalizing the above idea, we consider the rewardbased no-regret learning task (as per Example 1) in stochastic DMSO and frame this task in hybrid DMSO. Fix a parameter  $\Delta > 0$  of sub-optimality, we can consider the following "relaxed" constraint for each  $\pi \in \Pi$ :

$$\mathcal{M}_{\pi} := \{ M \in \mathcal{M} : V^{M}(\pi^{M}) - V^{M}(\pi) \le \Delta \},$$
(24)

and the corresponding "relaxed" constraint class  $\mathscr{P} := \{\mathcal{M}_{\pi} : \pi \in \Pi\}$ . For clarity, we write  $\mathcal{M}_{\Pi} := \bigcup_{\pi \in \Pi} \operatorname{co}(\mathcal{M}_{\pi})$ . Then, Theorem 10 implies that  $\mathsf{ExO}^+$  can achieve an upper bound in terms of the regret DEC of  $\mathcal{M}_{\Pi}$ . Following this idea and using a slightly more careful instantiation of  $\mathsf{ExO}^+$ , we have the following upper bounds.

**Proposition 13.** Let  $T \ge 1, \delta \in (0, 1), \Delta \ge 0$ , and we consider the reward-based no-regret learning task (Example 1) with a model class  $\mathcal{M}$ . Suppose that  $\mathcal{M}$  is compact (Assumption 2), and the regret  $DEC \operatorname{r-dec}_{\varepsilon}^{c}(\mathcal{M}_{\Pi})$ , as a function of  $\varepsilon$ , is of moderate decay. Then  $\operatorname{ExO}^{+}$  can be suitably instantiated (as detailed in Appendix F.4.4) to achieve with probability at least  $1 - \delta$  that

$$\frac{1}{T} \operatorname{\mathbf{Reg}}_{\mathsf{DM}}(T) \leq \Delta + O(\sqrt{\log T}) \cdot \Big[ \operatorname{\mathsf{r-dec}}_{\bar{\varepsilon}(T)}^{\mathsf{c}}(\mathcal{M}_{\Pi}) + \bar{\varepsilon}(T) \Big],$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log N_{\text{frac}}(\mathcal{M}, \Delta) + \log(1/\delta)}{T}}.$ 

We note that  $\mathcal{M}_{\Pi} \subseteq \operatorname{co}(\mathcal{M})$ , and hence when the model class  $\mathcal{M}$  is convex, the above upper bound in fact scales with the regret-DEC and fractional covering number of  $\mathcal{M}$ . We also note that Proposition 13 is not immediately implied by Theorem 10, because the latter also involves a term  $\log |\mathscr{P}| = \log |\Pi|$ , which can be much larger than  $\log N_{\operatorname{frac}}(\mathcal{M}, \Delta)$ . However, only slight adaptions specific to stochastic DMSO are needed (as detailed in Appendix F).

While guarantees of this form were first obtained by Chen et al. [2024], their bounds are directly reduced from Foster et al. [2022b] and scale with the DEC of  $co(\mathcal{M})$  (corresponding to the fully adversarial setting). In contrast, our framework provides finer upper bounds and has broader applicability, including convex hypothesis selection (Proposition 11), interactive estimation (Proposition 12), and also private regression (Proposition 24).

# 4 Query-Based Learning

In this section, we employ our framework to provide characterization for any query-based learning problem (Section 4.1). In particular, for learning under the *Statistical Queries* (SQ) [Kearns, 1998], the corresponding DEC recovers the *SQ dimension* of Feldman [2017], which is shown to provide both lower and upper bounds for the distributional search problems (Section 4.2). We also discuss the connection between SQ learning and LDP learning through the lens of our DEC formulation.

**Background on SQ learning.** The commonly studied setting of SQ learning is the distributional search problem (see e.g. Feldman [2017]), where a class  $\mathcal{M}^{\mathsf{d}} \subseteq \Delta(\mathcal{Z})$  of distributions is given, and each  $M \in \mathcal{M}^{\mathsf{d}}$  is associated with a set  $\Pi_M \subseteq \Pi$  of solutions, so that the loss function is specified as  $L(M, \pi) = \mathbf{1} \{ \pi \notin \Pi_M \}$ . The goal of an SQ algorithm is to find a decision  $\pi \in \Pi_M$  through adaptively querying the SQ oracle for any model  $M \in \mathcal{M}^{\mathsf{d}}$  (defined below).

**Definition 6** (SQ oracle). For a model  $M \in \Delta(\mathcal{Z})$ , tolerance parameter  $\tau > 0$ , an Statistical Query (SQ) oracle STAT<sup> $\tau$ </sup><sub>M</sub> is an oracle that, given any input  $\phi : \mathcal{Z} \to [0,1]$ , returns a value v such that  $|v - \mathbb{E}_{z \sim M} \phi(z)| \leq \tau$ .

To frame the problem of learning with SQ oracles, we consider the measurement class  $\Phi = (\mathcal{Z} \to [0,1])$ , and we note that each distribution  $M \in \mathcal{M}^{\mathsf{d}}$  induces a map  $\Pi \to \mathbb{R}$  given by  $M(\pi, \phi) = \mathbb{E}_{z \sim M}[\phi(z)]$ , i.e., the decision does not affect the response. Therefore, we may—with slight abuse of notation—write  $\mathcal{M}^{\mathsf{d}} \subseteq (\Pi \to \mathbb{R})$ , and for any  $M \in \mathcal{M}^{\mathsf{d}}$ , an SQ oracle  $\mathsf{STAT}_M^{\tau}$  corresponds to a constrained environment under the SQ DMSO. Conversely, under the specification above, any

constrained environment under the SQ DMSO corresponds to an (adaptive) SQ oracle. Therefore, our results for SQ DMSO naturally imply guarantees for SQ learning, as we discuss in Section 4.1 and Section 4.2.

# 4.1 General query oracles and DEC theory for query-based learning

Extending our discussion on SQ learning, we can formulate any SQ DMSO problem as a learning problem under certain query oracles. Specifically, given a measurement class  $\Phi$  and a model class  $\mathcal{M} \subseteq (\Pi \to \mathcal{V})$ , we define *general query* oracle as follows.

**Definition 7** (General Query). For a model  $M \in \mathcal{M}$  and tolerance parameter  $\tau > 0$ , a General Query (GQ) oracle  $\mathsf{GQ}_M^{\tau}$  is an oracle that, given any input decision  $\pi \in \Pi$  and measurement  $\phi \in \Phi$ , returns a value  $v \in \mathcal{V}$  such that  $||v - M(\pi, \phi)|| \leq \tau$ .

Clearly, there is an correspondence between the constrained environments under the SQ DMSO and general query oracles. Further, the formulation allows us to consider variants of SQ oracles, and, in particular, the standard SQ oracle and the VSTAT oracle. These are obtained below by suitably choosing the form of interaction between query and model.

**Example 3** (Symmetrized VSTAT oracle). For a distributional search problem, we can also consider learning under the VSTAT oracles. For any distribution  $M \in \Delta(\mathcal{Z})$ , tolerance parameter  $\tau \geq 0$ , a symmetrized VSTAT oracle  $\mathsf{VSTAT}$  oracles are equivalent to the standard VSTAT oracles. Clearly, a symmetrized VSTAT oracle is a GQ oracle with measurement class  $\Phi = (\mathcal{Z} \to [0, 1])$  and  $M(\phi) = \sqrt{\mathbb{E}_{z \sim M}\phi(z)}$ .

**Example 4** (Interactive SQ learning). In interactive SQ learning, the measurement class is  $\Phi = (\mathcal{Z} \to [0,1])$ , and each model  $M \in (\Pi \to \Delta(\mathcal{Z}))$  induces a map  $M : \Pi \times \Phi \to \mathbb{R}$  given by  $M(\pi, \phi) = \mathbb{E}_{z \sim M(\pi)}[\phi(z)]$ . This is a natural generalization of SQ learning to interactive decision making.

More generally, our formulation also allows us to consider other query-based learning settings, e.g., Correlation Statistical Queries [Bshouty and Feldman, 2002], Differentiable Learning Queries [Joshi et al., 2024], and the *batch* SQ learning, where at each round the learner can select a batch of queries  $\phi = (\phi^1, \dots, \phi^n) \in \mathcal{L}^n$ .

**SQ DEC lower and upper bounds.** Now, we present the SQ DEC lower and upper bounds implied by our framework. We begin with the lower bound for metric-based loss.

**Theorem 14** (Query-based lower bound). Let  $T \ge 1$ , model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \mathcal{V})$ , and the loss function L is metric-based. Suppose that Alg is a T-round query-based algorithm. Then there exists a model  $M \in \mathcal{M}$  and a GQ oracle  $\mathsf{GQ}_M^{\tau}$  such that under this oracle, the expected risk of Alg is lower bounded as

$$\mathbb{E}^{\mathrm{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \geq \frac{1}{8} \mathrm{p}\text{-}\mathsf{dec}_{\underline{\varepsilon}(T)}^{\tau\text{-}\mathsf{SQ}}(\mathcal{M}),$$

where  $\underline{\varepsilon}(T) = \frac{1}{2\sqrt{T}}$ .

Further, for general loss function  $L : \mathcal{M} \times \Pi \to [0,1]$  and  $\delta \in (0,1)$ , there exists a model  $M \in \mathcal{M}$ and a GQ oracle  $\mathsf{GQ}_M^{\tau}$  such that under this oracle, the expected risk of Alg is lower bounded as

$$\mathbb{E}^{\operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\operatorname{\mathsf{DM}}}(T)] \ge \operatorname{p-dec}_{\underline{\varepsilon}_{\delta}(T)}^{\tau-\operatorname{SQ}}(\mathcal{M}) - \delta,$$

where  $\underline{\varepsilon}_{\delta}(T) = \sqrt{\frac{\delta}{T}}$ .

Though Theorem 14 is a direct corollary of Theorem 7, we provide a more direct and simpler proof of Theorem 14 in Appendix H.1 as an illustration.

For upper bound, we propose SQ-E2D, an adaption of the E2D algorithm [Foster et al., 2023b] for SQ DMSO, which achieves an upper bound in SQ DEC with minimal assumptions (Appendix G.2). By instantiating Theorem 8, we also have the upper bound of  $ExO^+$ .

**Theorem 15** (Query-based upper bound). Let  $T \ge 1, \delta \in (0,1)$ , model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \mathcal{V})$ . Then, for any model  $M \in \mathcal{M}^{\mathsf{d}}$  and given access to any (possibly adaptive) GQ oracle  $\mathsf{GQ}_{M}^{\tau}$  of M, the SQ-E2D (Algorithm 4) achieves with probability at least  $1 - \delta$  that

$$\mathbf{Risk}_{\mathsf{DM}}(T) \lesssim \mathsf{p}\text{-}\mathsf{dec}_{\bar{\varepsilon}(T)}^{2\tau\text{-}\mathsf{SQ}}(\mathcal{M}),$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log(|\mathcal{M}|/\delta)}{T}}$ .

Further, suppose that the loss function L is metric-based, and the SQ DEC  $p-dec_{\varepsilon}^{\tau-SQ}(\mathcal{M})$  is of moderate decay. Then, for any model  $M \in \mathcal{M}^{d}$  and given access to any (possibly adaptive) GQoracle  $GQ_{M}^{\tau}$  of M,  $ExO^{+}$  (instantiated on  $\mathscr{P}_{\tau-query}$ , following Theorem 8) achieves with probability at least  $1-\delta$ 

$$\mathbf{Risk}_{\mathsf{DM}}(T) \lesssim \mathsf{p-dec}_{\overline{\varepsilon}(T)}^{\tau-\mathsf{SQ}}(\mathcal{M}).$$

Note that the upper bound of SQ-E2D scales with the SQ DEC at the correctness level  $2\tau$ . In contrast, the upper bound of  $ExO^+$  eliminates this factor of 2 under additional assumptions. We note that for  $ExO^+$ , the assumptions on the loss function and the regularity of the SQ DEC can both be relaxed (similar to Eq. (17)).

## 4.2 Connection to the SQ dimension

For a distributional search problem, Feldman [2017] studies the optimal query complexity to arbitrary SQ oracle with correctness  $\tau$ . Recall that in the distributional search problem, a class  $\mathcal{M}^{\mathsf{d}} \subseteq \Delta(\mathcal{Z})$  of distributions is given, and each  $M \in \mathcal{M}^{\mathsf{d}}$  is associated with a set  $\Pi_M \subseteq \Pi$  of solutions. Then, for success probability  $\beta$  and correctness  $\tau \geq 0$ , the optimal query complexity is the minimum number of rounds required to return a solution  $\pi \in \Pi_{M^*}$  with success probability at least  $\beta$ , given access to any SQ oracle  $\mathsf{STAT}_{M^*}^{\tau}$  for any  $M^* \in \mathcal{M}^{\mathsf{d}}$ .

More generally, for any query-based model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \Delta(\mathcal{V}))$ , we define the *T*-round minimax risk as

$$\mathfrak{M}_{T}^{\tau-\mathrm{SQ}}(\mathcal{M}) = \inf_{\mathsf{Alg Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)],$$

where the supremum is taken over all environments satisfying query correctness with tolerance  $\tau$  for a model  $M^* \in \mathcal{M}$ . Then, the minimax query complexity for achieving  $\Delta$ -risk is defined as

$$\mathfrak{C}_{\Delta}^{\tau-\mathsf{SQ}}(\mathcal{M}) := \inf \left\{ T : \mathfrak{M}_{T}^{\tau-\mathsf{SQ}}(\mathcal{M}) \leq \Delta \right\}.$$

For distributional search problems, achieving success probability  $\beta$  is equivalent to achieving  $(1-\beta)$ -risk. Hence, in the following, we state the results of Feldman [2017] in terms of  $\mathfrak{C}_{1-\beta}^{\tau-SQ}(\mathcal{M}^{\mathsf{d}})$ .<sup>2</sup>

<sup>&</sup>lt;sup>2</sup>Recall that we identify  $\mathcal{M}^{\mathsf{d}} \subseteq (\mathbf{\Pi} \to \mathbb{R})$  by regarding each model  $M \in \Delta(\mathcal{Z})$  as a map  $(\pi, \phi) \mapsto \mathbb{E}_{z \sim M} \phi(z)$ .

Characterization by SQ dimension. In the following, we first discuss the notion of SQ dimension and the results of Feldman [2017] in detail.

**Definition 8** (SQ dimension). In distributional search problems, given a model class  $\mathcal{M}^{\mathsf{d}} \subseteq \Delta(\mathcal{Z})$ , parameter  $\tau > 0$ , success probability  $\beta \in [0, 1]$ , the SQ dimension with the reference model  $\overline{M} \in \Delta(\mathcal{Z})$  is defined as

$$\mathsf{SQDim}^{\tau}_{\beta}(\mathcal{M}^{\mathsf{d}},\overline{M}) = \inf_{p \in \Delta(\Pi)} \sup_{\mu \in \Delta(\mathcal{M}^{\mathsf{d}}_{p,\beta})} \inf_{\phi: \mathcal{Z} \to [0,1]} \frac{1}{\mathbb{P}_{M \sim \mu}(\left|M(\phi) - \overline{M}(\phi)\right| > \tau)},$$

where  $\mathcal{M}_{p,\beta}^{\mathsf{d}} := \{ M \in \mathcal{M}^{\mathsf{d}} : p(\Pi_M) < \beta \}$ . The SQ dimension of  $\mathcal{M}^{\mathsf{d}}$  is then defined as  $\mathsf{SQDim}_{\beta}^{\tau}(\mathcal{M}^{\mathsf{d}}) := \sup_{\overline{M} \in \Delta(\mathcal{Z})} \mathsf{SQDim}_{\beta}^{\tau}(\mathcal{M}^{\mathsf{d}}, \overline{M})$ .

In terms of the SQ dimension defined above, Feldman [2017] provides the following lower and upper bounds on  $\mathfrak{C}^{\tau-SQ}_{\Delta}(\mathcal{M}^d)$  for any distribution search problem with a model class  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ .

**Proposition 16** (SQ dimension characterization of the query complexity, Feldman [2017]). For success probability  $\beta \in [0, 1]$ , parameter  $\delta \in (0, 1 - \beta]$ , it holds that

$$\delta \cdot \mathsf{SQDim}_{\beta-\delta}^{\tau}(\mathcal{M}^{\mathsf{d}}) \leq \mathfrak{C}_{1-\beta}^{\tau-\mathsf{SQ}}(\mathcal{M}^{\mathsf{d}}) \leq \tilde{O}\left(\mathsf{SQDim}_{\beta+\delta}^{3\tau}(\mathcal{M}^{\mathsf{d}}) \cdot \frac{C_{\mathrm{KL}}(\mathcal{M}^{\mathsf{d}})}{\tau^{2}} \log(1/\delta)\right),$$
(25)

where  $C_{\mathrm{KL}}(\mathcal{M}^{\mathsf{d}}) := \inf_{\overline{M} \in \Delta(\mathcal{Z})} \sup_{M \in \mathcal{M}^{\mathsf{d}}} D_{\mathrm{KL}}(M \parallel \overline{M})$  is the KL radius of  $\mathcal{M}^{\mathsf{d}}$ .

Comparison to the SQ DEC characterization. To compare our results with the above characterization, we first show that the SQ dimension is quantitatively equivalent to the SQ DEC of  $\mathcal{M}$ , as long as the Minimax theorem applies.

**Proposition 17.** Suppose that  $\mathcal{Z}$  is finite, and  $\mathcal{M}^{\mathsf{d}} \subseteq \Delta(\mathcal{Z})$  is a distribution class. Then for any success probability  $\beta \in [0, 1]$ , reference model  $\overline{M} \in \Delta(\mathcal{Z})$ , we have

$$\mathrm{p\text{-}dec}_{\varepsilon}^{\tau\mathrm{-}\mathrm{SQ}}(\mathcal{M}^{\mathsf{d}},\overline{M})>1-\beta\quad\Leftrightarrow\quad\varepsilon^{-2}\leq\mathrm{SQDim}_{\beta}^{\tau}(\mathcal{M}^{\mathsf{d}},\overline{M}).$$

Proof can be found in Appendix H.3. Therefore, SQ DEC can be viewed as a generalization of the SQ dimension to general query-based learning.

To have a clearer comparison, for any model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \mathcal{V})$ , we define the DEC-induced SQ dimension as<sup>3</sup>

$$\underline{\mathsf{SQDim}}^{\tau}_{\beta}(\mathcal{M}) := \min\{\varepsilon^{-2} : \mathsf{p-dec}^{\tau-\mathsf{SQ}}_{\varepsilon}(\mathcal{M}) \le 1-\beta\}.$$

Then, for any query-based learning problem with loss bounded in [0, 1], our results imply the following characterization

$$\delta \cdot \underline{\mathsf{SQDim}}_{\beta-\delta}^{\tau}(\mathcal{M}) \leq \mathfrak{C}_{1-\beta}^{\tau-\mathsf{SQ}}(\mathcal{M}) \lesssim \underline{\mathsf{SQDim}}_{\beta+\delta}^{2\tau}(\mathcal{M}) \cdot \log(|\mathcal{M}|/\delta),$$
(26)

for any success probability  $\beta \in [0, 1]$  and any parameter  $\delta \in (0, 1 - \beta]$ . We note that for metricbased loss, the  $2\tau$ -factor in the upper bound can be improved to  $\tau$  under the assumption that the SQ DEC is of moderate decay (Theorem 15).

<sup>&</sup>lt;sup>3</sup>This is slightly different from the original SQ dimension (cf. Definition 8), because in the definition (4) of SQ DEC, the supremum is taken over all *randomized* reference models  $\overline{M} \in (\mathbf{\Pi} \to \Delta(\mathcal{V}))$ .

Compared to Eq. (25), our characterization (when specialized to SQ learning in distributional search problems) does not incur the  $\tau^{-2}$ -gap between lower and upper bounds, but its upper bound scales with  $\log |\mathcal{M}^{\mathsf{d}}|$ , the complexity of the class  $\mathcal{M}^{\mathsf{d}}$ . Although it can be replaced by the log-covering number of  $\mathcal{M}^{\mathsf{d}}$ , this dependence might still be much larger than the  $C_{\mathrm{KL}}$ -factor in Eq. (25). While the dependence on  $\log |\mathcal{M}|$  can be unavoidable beyond this setting, the upper bound of Algorithm 4 for such problems can also be improved to take advantage of bounded  $C_{\mathrm{KL}}$  (see our discussion in Appendix G.2).

### 4.3 Relation between SQ learning and LDP learning

It is well known that for PAC learning, there is a (polynomial) equivalence between LDP algorithms and SQ algorithms [Kasiviswanathan et al., 2011]. We show that such an equivalence also holds between LDP DEC and SQ DEC. This is expected, since the DECs capture the complexity of the corresponding learning task. In greater generality, we state this equivalence for *interactive SQ* learning (Example 4), a generalization of SQ learning.

**Lemma 18.** Let  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{O}))$ . Then, for interactive SQ learning (Example 4), the SQ DEC can be bounded as

$$\mathsf{p}\mathsf{-dec}_{\tau+\varepsilon}^{\mathsf{LDP}}(\mathcal{M},\overline{M}) \le \mathsf{p}\mathsf{-dec}_{\varepsilon}^{\tau\mathsf{-SQ}}(\mathcal{M},\overline{M}) \le \mathsf{p}\mathsf{-dec}_{\varepsilon/\tau}^{\mathsf{LDP}}(\mathcal{M},\overline{M}), \qquad \forall \overline{M}.$$
(27)

Proof is presented in Appendix H.4. From Eq. (27), it is clear that a comparison between the DECs would typically lead to *loose* rates. This can be explained by the difference between SQ learning (where the response can be perturbed adversarially) and LDP learning (where the observations are stochastic).

In view of the relationship between LDP algorithms and SQ algorithms, Kasiviswanathan et al. [2011] established a lower bound for LDP *learning parity* by reduction. In Appendix B.4, we show that DEC theory provides a more direct LDP lower bound for learning parity through lower bounding the private PAC-DEC.

# 5 Locally Private Learning

In this section, we employ the DEC formulation to analyze private DMSO and characterize the complexity of LDP learning.

**Problems encompassed by private DMSO.** Before diving into details, we first discuss several common settings of private learning that are encompassed by private DMSO (page 8). Recall that in this setting, the learner selects, on round t, a decision  $\pi_t \in \Pi$  and a private channel  $Q_t \in Q$ , the environment generates latent observation  $z_t \sim M^*(\pi_t)$ , and the learner observes  $o_t \sim Q_t(\cdot|z_t)$ . The ground truth model  $M^*$  is known to belong to a given model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ .

In this section, one of our primary foci is the setting of *reward-based* learning [Foster et al., 2021, 2023b, Chen et al., 2024], where the goal of the learner is to maximize the expected reward of the decision, or equivalently, minimize its sub-optimality.

**Definition 9** (Reward-based value and loss function). Given a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ , we call the value function V reward-based, if there is a known reward function  $R : \mathcal{Z} \times \Pi \to [0,1]$ such that  $V^{M}(\pi) = \mathbb{E}^{M,\pi}[R(z,\pi)]$  is the expected cumulative reward of  $\pi$  under M. We also denote  $\pi^{M} := \arg \max_{\pi \in \Pi} V^{M}(\pi)$  to be the optimal decision for M (under the value function). A loss function  $L : \mathcal{M} \times \Pi \to \mathbb{R}$  is reward-based if it is specified by a reward-based value function V as

$$L(M,\pi) = V^{M}(\pi^{M}) - V^{M}(\pi).$$
(28)

Loss functions of the above form appear in many LDP learning problems of interest, including classification and regression, online learning, bandits and contextual bandits, and Reinforcement Learning (RL).

We also consider examples of *statistical tasks*, where  $z_1, \dots, z_T \sim M^*$  are independent and identically distributed, i.e., the latent observation is independent of the decision. Nonetheless, here the learner is actively choosing channels  $Q_t$ , affecting the amount of information received, and the performance is assessed by the final decision  $\pi_{T+1}$ .

**Definition 10** (Statistical task). We call the model class  $\mathcal{M}$  a statistical model class if for each model  $M \in \mathcal{M}$ ,  $M(\pi) = M \in \Delta(\mathcal{Z})$  is independent of  $\pi \in \Pi$ , i.e., we may regard  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ .

Examples of statistical tasks include hypothesis testing, hypothesis selection, classification and regression, functional estimation, and density estimation, among others. For statistical tasks, our definition of  $\alpha$ -LDP algorithms agrees with the notion of sequential private channels [Duchi et al., 2013, 2018] (as detailed in Appendix B.1).

# 5.1 DEC theory for private PAC learning

We start with the private PAC-DEC lower bounds for reward-based loss and metric-based loss.

**Theorem 19** (Private PAC-DEC lower bound). Let  $T \ge 1$ , Alg be a T-round  $\alpha$ -LDP algorithm.

(1) Suppose that the loss function L is metric-based. Then it holds that

$$\sup_{M \in \mathcal{M}} \mathbb{E}^{M, \operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] \geq \frac{1}{8} \operatorname{p-dec}_{\underline{\varepsilon}(T)}^{\operatorname{LDP}}(\mathcal{M}),$$

where  $\underline{\varepsilon}(T) = \frac{c}{\sqrt{\alpha^2 T}}$ , and c is a universal constant.

(2) Suppose that the loss function L is reward-based. Then

$$\sup_{M \in \mathcal{M}} \mathbb{E}^{M, \operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] \geq \frac{1}{4} \Big( \operatorname{p-dec}_{\underline{\varepsilon}(T)}^{\operatorname{LDP}}(\mathcal{M}) - 6\underline{\varepsilon}(T) \Big).$$

The proof of Theorem 19 is deferred to Appendix I.2 and is based on the strong data-processing inequality stated below (Proposition 20). We note that Theorem 19 (1) can also be proven directly by combining the hybrid DEC lower bound (Theorem 7) with Proposition 20. Finally, we also note that fractional covering number also provides a lower bound (Theorem 34), which is complementary to the private PAC-DEC lower bounds above.

Key ingredients for the lower bound. As we have discussed in Section 2.3, private PAC-DEC can be viewed as a special case of the hybrid DEC, based on the following characterization of the data-processing under DP channels. We recall that for any channel  $Q \in Q$  and any distribution  $P \in \Delta(\mathcal{Z})$ , we denote  $Q \circ P$  to be the marginal distribution of o under  $z \sim P, o \sim Q(\cdot|z)$ . The proof of the following result is presented in Appendix I.1. **Proposition 20** (Strong data-processing inequality). Suppose that Q is an  $\alpha$ -LDP channel. Then there exists a distribution  $q := q_{Q} \in \Delta(\mathcal{L})$ , such that for any two distributions  $P_{1}, P_{2} \in \Delta(\mathcal{Z})$  over  $\mathcal{Z}$ , it holds that

$$\frac{(e^{\alpha}-1)^2}{8e^{2\alpha}}\mathbb{E}_{\ell\sim q}\mathsf{D}_{\ell}^2(P_1,P_2) \le D_{\mathrm{H}}^2(\mathsf{Q}\circ P_1,\mathsf{Q}\circ P_2) \le \frac{(e^{\alpha}-1)^2}{8}\mathbb{E}_{\ell\sim q}\mathsf{D}_{\ell}^2(P_1,P_2).$$
(29)

Furthermore,

$$D_{\mathrm{KL}}(\mathbf{Q} \circ P_1 \| \mathbf{Q} \circ P_2) \le D_{\chi^2}(\mathbf{Q} \circ P_1 \| \mathbf{Q} \circ P_2) \le (e^{\alpha} - 1)^2 \mathbb{E}_{\ell \sim q} \mathsf{D}_{\ell}^2(P_1, P_2).$$
(30)

In particular, Eq. (30) recovers the strong data-processing inequality of Duchi et al. [2018], as the  $\ell$ -divergences are always upper bounded by TV distance.

An interpretation of the characterization in Proposition 20 is that, in terms of divergences, any private channel can be expressed in terms of a distribution over the *binary channels*.

**Example 5** (Binary channel). Perhaps the simplest nontrivial channel is the binary channel, defined as follows. For any map  $\ell : \mathcal{Z} \to [0,1]$ , the binary channel  $Q_{\ell}$  associated with  $\ell$  is given by

$$Q_{\ell}(+1|z) = \frac{1 + c_{\alpha}\ell(z)}{2}, \qquad Q_{\ell}(-1|z) = \frac{1 - c_{\alpha}\ell(z)}{2},$$

where  $c_{\alpha} = 1 - e^{-\alpha}$  and  $\mathcal{O} = \{-1, 1\}$ . It can be verified that this channel is indeed  $\alpha$ -DP. We define  $\mathcal{Q}_{\alpha,\text{bin}}$  to be the class of all binary channels described above, i.e.,  $\mathcal{Q}_{\alpha,\text{bin}} := \{\mathbf{Q}_{\ell} : \ell \in \mathcal{L}\}.$ 

It is clear that for any map  $\ell : \mathcal{Z} \to [0,1]$ , we have  $D_{\mathrm{H}}^2(\mathsf{Q} \circ P_1, \mathsf{Q} \circ P_2) \simeq \alpha^2 \mathsf{D}_{\ell}^2(P_1, P_2)$  (up to absolute constants).

**Private PAC-DEC upper bounds.** We propose LDP-E2D, an extension of the E2D algorithm of Foster et al. [2023b] to the LDP setting, providing the following upper bound for PAC learning with any problem class  $\mathcal{M}$ .

**Theorem 21** (Private PAC-DEC upper bound via E2D). For any model class  $\mathcal{M}$ , the LDP-E2D algorithm (Algorithm 2) preserves  $\alpha$ -LDP and achieves with probability at least  $1 - \delta$  that

$$\mathbf{Risk}_{\mathsf{DM}}(T) \le \mathsf{p}\text{-}\mathsf{dec}_{\bar{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}),$$

where  $\bar{\varepsilon}(T) = C \sqrt{\frac{\log(|\mathcal{M}|/\delta)\log(1/\delta)}{\alpha^2 T}}.$ 

We note that under certain assumptions,  $ExO^+$  can also be instantiated to achieve a similar upper bound, and we call the obtained algorithm LDP-ExO (detailed in Appendix F.5). In the next result, we derive an upper bound of LDP-ExO scaling with the fractional covering number of  $\mathcal{M}$ , following Proposition 13.

**Theorem 22.** Let  $T \ge 1$ ,  $\delta \in (0,1)$ , model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{O}))$ , and the loss function L be reward-based. Suppose that  $\mathcal{M}$  is compact (Assumption 2), and the private PAC-DEC p-dec<sup>LDP</sup><sub> $\varepsilon$ </sub>( $\mathcal{M}_{\Pi}$ ), as a function of  $\varepsilon$ , is of moderate decay. Then LDP-ExO (instantiated as in Appendix F.5.2) preserves  $\alpha$ -LDP and achieves with probability at least  $1 - \delta$  that

$$\mathbf{Risk}_{\mathsf{DM}}(T) \leq \Delta + O(1) \cdot \Big[ \mathsf{p-dec}_{\bar{\varepsilon}(T)}^{\mathsf{LDP}}(\operatorname{co}(\mathcal{M})) + \bar{\varepsilon}(T) \Big],$$
  
where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log N_{\mathsf{frac}}(\mathcal{M}, \Delta) + \log(1/\delta)}{\alpha^2 T}}.$ 

#### 5.2 Application: private regression

In this section, we consider the task of proper regression under LDP.

**Example 6** (Regression). In the regression task,  $\mathcal{X}$  is a given covariate space,  $\mathcal{F} \subseteq (\mathcal{X} \rightarrow [-1, 1])$  is a given function class, and  $L(\cdot, \cdot) : [-1, 1]^2 \rightarrow [0, 1]$  is a given loss. The observation space is  $\mathcal{Z} = \mathcal{X} \times [-1, 1]$ , and the loss function L is then given by

$$L(M, f) = \mathbb{E}_{(x,y) \sim M} \mathcal{L}(y, f(x)) - \min_{f^{\star} \in \mathcal{F}} \mathbb{E}_{(x,y) \sim M} \mathcal{L}(y, f^{\star}(x)).$$

Regression is a statistical task, in the sense of Definition 10, as the model class  $\mathcal{M}$  is a subset of  $\Delta(\mathcal{Z})$ . The loss function for this task is reward-based, in the sense of Definition 9, if we set the reward function as R((x, y), f) = 1 - L(y, f(x)).

The choices of loss function L of interest include (1) squared loss:  $L_{sq}(y, y') = (y - y')^2$ , and (2) absolute loss:  $L_{abs}(y, y') = |y - y'|$ . We also note that the classification task is a special case of the regression problem described above, by specializing  $\mathcal{F} \subseteq (\mathcal{X} \to \{0,1\}), \ \mathcal{M} \subseteq \Delta(\mathcal{X} \times \{0,1\})$  and  $L_{cl}(y, y') = \mathbf{1} \{y \neq y'\}$ .

In the literature, both *agnostic* regression and *well-specified* regression are studied, where the model class  $\mathcal{M}$  is specified as follows:

- Agnostic regression: the model class is  $\mathcal{M}_{agnostic} = \Delta(\mathcal{Z})$ , i.e., there is no prior knowledge of the underlying environment.
- Well-specified regression: the model class  $\mathcal{M}_{\mathcal{F}}$  consists of all models  $M \in \Delta(\mathcal{Z})$  such that there exists  $f^M \in \mathcal{F}$ , such that  $y|x \sim \operatorname{Rad}(f^M(x))$  under M.<sup>4</sup>

Notice that for agnostic regression, the model class  $\mathcal{M}_{agnostic} = \Delta(\mathcal{Z})$  is convex, and hence Theorem 22 applies immediately. In the following, we state the guarantees for agnostic regression and realizable regression. To avoid measure-theoretic issues, we assume that  $\mathcal{X}$  is finite.

**Proposition 23** (Agnostic regression). Let  $T \ge 1, \delta \in (0, 1), \Delta > 0$ . Suppose that the private PAC-DEC p-dec<sup>LDP</sup><sub> $\varepsilon$ </sub>( $\mathcal{M}_{agnostic}$ ) is of moderate decay as a function of  $\varepsilon$ . Then, LDP-ExO can be instantiated (following Theorem 22) to achieve with probability at least  $1 - \delta$ 

$$\mathbf{Risk}_{\mathsf{DM}}(T) \leq \Delta + O(1) \cdot \Big[\mathsf{p-dec}_{\bar{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}_{\mathsf{agnostic}}) + \bar{\varepsilon}(T)\Big],$$

$$\overline{_{\mathsf{frac}}(\mathcal{M}_{\mathsf{agnostic}}, \Delta) + \log(1/\delta)}_{2T}.$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log N_{\text{frac}}(\mathcal{M}_{\text{agnostic}}, \Delta) + \log(1/\delta)}{\alpha^2 T}}$ 

For well-specified regression, a similar guarantee also applies.

**Proposition 24** (Well-specified regression). Let  $T \ge 1, \delta \in (0,1), \Delta \ge 0$ . Suppose that the private PAC-DEC  $p-dec_{\varepsilon}^{LDP}(\mathcal{M}_{\mathcal{F}})$  is of moderate decay as a function of  $\varepsilon$ . Then, LDP-ExO can be instantiated (as detailed in Appendix F.5.3) to achieve with probability at least  $1 - \delta$ 

$$\mathbf{Risk}_{\mathsf{DM}}(T) \lesssim \mathsf{p-dec}_{\bar{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}_{\mathcal{F}}) + \bar{\varepsilon}(T),$$

where  $\bar{\varepsilon}(T) = \Delta + \sqrt{\frac{\log N_{\text{frac}}(\mathcal{F}, \Delta) + \log(1/\delta)}{\alpha^2 T}}$ , and the fractional covering number of  $\mathcal{F}$  is defined as

$$N_{\mathsf{frac}}(\mathcal{F}, \Delta) := \inf_{p \in \Delta(\mathcal{F})} \sup_{\mu \in \Delta(\mathcal{X}), f^{\star} \in \mathcal{F}} \frac{1}{p(f : \mathbb{E}_{x \sim \mu} | f(x) - f^{\star}(x)| \le \Delta)}.$$
(31)

<sup>&</sup>lt;sup>4</sup>For simplicity, we assume that  $y \in \{-1, 1\}$  in this case without loss of generality.

A detailed discussion of the fractional covering number  $N_{\text{frac}}(\mathcal{F}, \Delta)$  is deferred to Appendix C.2. In Appendix B.3, we also consider the online regression task (where the (x, y) pair is chosen adversarially by the environment).

#### 5.3 Application: private linear regression

In this section, we investigate LDP regression in *linear models*.

**Example 7** (Linear models). Suppose that  $\mathcal{X} \subseteq \mathbb{B}^d(1)$ , the linear function class  $\mathcal{F}_{\mathsf{Lin}}$  is given by

$$\mathcal{F}_{\mathsf{Lin}} := \{ f_{\theta}(x) = \langle \theta, x \rangle \}_{\theta \in \mathbb{R}^{d}(1)}$$

and let  $\mathcal{M}_{\mathsf{Lin}}$  be the induced class of well-specified models, i.e., each model  $M \in \mathcal{M}$  is associated with a covariate distribution  $\nu_M$  and a parameter  $\theta^M$ , such that  $(x, y) \sim M$  is generated as  $x \sim \nu_M, y \sim \operatorname{Rad}(\langle x, \theta^M \rangle)$ .

In linear models, we consider decision space  $\Pi = \mathbb{B}^d(1)$  (the space of estimators). For an estimator  $\theta \in \mathbb{B}^d(1)$ , we consider the following loss functions that measure the  $L_1(L_2)$  estimation error:

$$L_1(M,\theta) = \mathbb{E}_{x \sim M} |\langle x, \theta - \theta^M \rangle|, \qquad L_2(M,\theta) = \mathbb{E}_{x \sim M} \langle x, \theta - \theta^\star \rangle^2.$$

Note that the  $L_2$  error agrees with the squared loss of the function  $f_{\theta}(x) = \langle \theta, x \rangle$  considered in Section 5.2. However, we note that the  $L_1$  loss here measures the error of the estimator  $\theta$  with respect to the ground-truth parameter  $\theta^M$ , which is different from the absolute-loss regression considered in Section 5.2.

**Rates for**  $L_2$  regression. For LDP linear regression, to achieve the standard  $T^{-1}$ -rate under  $L_2$  risk, it is necessary to require the covariance matrix to be well-conditioned [Duchi et al., 2018, Duchi and Ruan, 2024]. Otherwise, the convergence rate can degrade to  $T^{-1/2}$  in the worst case, as indicated by the following folklore lower bound [Duchi and Ruan, 2024]. Li et al., 2024].

**Lemma 25.** Suppose that d = 1, and  $\nu$  is a given distribution over [-1,1]. Then for any *T*-round  $\alpha$ -LDP algorithm Alg with output estimator  $\hat{\theta}$ , there exists a model  $M^*$  with covariate distribution  $\nu$  and parameter  $\theta^* \in [-1,1]$ , such that

$$\mathbb{E}^{M^{\star},\operatorname{Alg}}L_{2}(M^{\star},\hat{\theta}) \gtrsim \mathbb{E}_{\nu}|x|^{2} \cdot \min\left\{\frac{1}{\alpha^{2}T(\mathbb{E}_{\nu}|x|)^{2}},1\right\}.$$

In particular, for any  $T \ge 1$ , there exists a "worst-case" covariate distribution  $\nu_T$  with  $\nu_T(0) = 1 - \frac{1}{\sqrt{\alpha^2 T}}$  and  $\nu_T(1) = \frac{1}{\sqrt{\alpha^2 T}}$ , such that any  $\alpha$ -LDP algorithm incurs an  $L_2$  loss of  $\Omega\left(\frac{1}{\sqrt{\alpha^2 T}}\right)$ .

**Rates for**  $L_1$  regression. In contrast, we show that a  $T^{-1/2}$ -rate under  $\ell_1$ -loss can still be achieved. Note that in the upper bound below, we do *not* assume the covariate distribution is known. Details are deferred to Appendix I.4.

**Theorem 26.** Let the loss function  $L = L_1$  be given by the  $L_1$  error. Then it holds that

$$\mathsf{p}\operatorname{-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}_{\mathsf{Lin}}) \leq O\Big(\sqrt{d}\varepsilon\Big).$$

Further, LDP-ExO can be instantiated to output  $\hat{\theta} \in \mathbb{B}^{d}(1)$  so that with probability at least  $1 - \delta$ ,

$$L_1(M^\star, \hat{\theta}) \le \tilde{O}\left(\sqrt{\frac{d^2 \log(1/\delta)}{\alpha^2 T}}\right),$$

which is minimax-optimal up to logarithmic factors (cf. the minimax lower bound in Corollary I.8).

To the best of our knowledge, such a assumption-free  $T^{-1/2}$ -rate is new for LDP linear regression under  $L_1$  error. More specifically, previous works mostly focus on  $L_2$  loss regression, and hence when converted to  $L_1$  loss, the results either have a  $T^{-1/4}$ -rate or need extra assumptions, e.g. a bounded condition number of the covariance matrix  $\Sigma = \mathbb{E}[xx^{\top}]$  [Duchi et al., 2018, Wang and Xu, 2019, etc.]. We note that  $L_1$  error, while less well-studied, can be of interest for a broad range of applications, including offline policy evaluation with linear function approximation.

In Section 5.5.1, we apply a similar technique to provide a near-optimal regret for learning linear contextual bandits.

#### 5.4 DEC theory for private no-regret learning

In this section, we present the private regret-DEC and the guarantees for private no-regret learning. We focus on the reward-based setting.

**Private regret-DEC.** For a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  and a value function V, we define the private regret-DEC of  $\mathcal{M}$  with respect to a reference model  $\overline{M} \in co(\mathcal{M})$  as

$$\mathsf{r}\text{-}\mathsf{dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M},\overline{M}) := \inf_{p \in \Delta(\Pi \times \mathcal{L})} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[V^{M}(\pi^{M}) - V^{M}(\pi)] \mid \mathbb{E}_{(\pi,\ell) \sim p}\mathsf{D}_{\ell}^{2}(M(\pi),\overline{M}(\pi)) \leq \varepsilon^{2} \right\},\tag{32}$$

and we define the private regret-DEC of  $\mathcal{M}$  as

$$\operatorname{\mathsf{r-dec}}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M}) := \sup_{\overline{M} \in \operatorname{co}(\mathcal{M})} \operatorname{\mathsf{r-dec}}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M} \cup \{\overline{M}\}, \overline{M}).$$
(33)

Similar to the private PAC-DEC, the private regret-DEC can also be viewed as a specification of the hybrid DEC. By instantiating Theorem 9 and Theorem 10, we have the following regret bounds.

**Theorem 27** (Private regret-DEC lower bound). Let  $T \ge 1$ . Suppose that the value function V is reward-based (Definition 9). Then, for any T-round  $\alpha$ -LDP algorithm Alg, it holds that

$$\sup_{M \in \mathcal{M}} \mathbb{E}^{M,\mathrm{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \geq \frac{T}{4} \Big( \mathrm{r-dec}_{\underline{\varepsilon}(T)}^{\mathrm{LDP}}(\mathcal{M}) - C\underline{\varepsilon}(T) \Big) - 1,$$

where  $\underline{\varepsilon}(T) = \frac{c}{\sqrt{\alpha^2 T}}$ , and c, C are universal constants.

**Theorem 28** (Private regret-DEC upper bounds). Let  $T \ge 1, \delta \in (0, 1)$ . Suppose that the model class  $\mathcal{M}$  is compact, the value function V is reward-based, and the private regret-DEC  $\operatorname{r-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M})$ is of moderate decay as a function of  $\varepsilon$ . Then, a suitable instantiation of LDP-ExO (as detailed in Appendix F.5.1) achieves with probability at least  $1 - \delta$  that

$$\frac{1}{T} \operatorname{\mathbf{Reg}}_{\mathsf{DM}}(T) \le O(\sqrt{\log T}) \cdot \Big[ \operatorname{\mathsf{r-dec}}_{\bar{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}) + \bar{\varepsilon}(T) \Big],$$
(34)

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log(|\mathcal{M}|/\delta)}{\alpha^2 T}}$ .

where  $\bar{\varepsilon}'$ 

Further, suppose that the private regret-DEC r-dec<sup>LDP</sup><sub> $\varepsilon$ </sub>(co( $\mathcal{M}$ )) is of moderate decay. Then an alternative instantiation of LDP-ExO (as detailed in Appendix F.5.2) achieves with probability at least  $1 - \delta$ 

$$\frac{1}{T} \operatorname{Reg}_{\mathsf{DM}}(T) \leq \Delta + O(\sqrt{\log T}) \cdot \Big[ \mathsf{r} - \operatorname{dec}_{\bar{\varepsilon}'(T)}^{\mathsf{LDP}}(\operatorname{co}(\mathcal{M})) + \bar{\varepsilon}'(T) \Big], \tag{35}$$
$$(T) = \sqrt{\frac{\log N_{\mathsf{frac}}(\mathcal{M}, \Delta) + \log(1/\delta)}{\alpha^2 T}}.$$

We note that under reward-based value function, the algorithms of Foster et al. [2023b], Glasgow and Rakhlin [2023] may also be adapted to achieve a regret bound similar to Eq. (34), under a weaker regularity assumption on the private regret-DEC r-dec<sup>LDP</sup><sub> $\varepsilon$ </sub>( $\mathcal{M}$ ). We state the upper bound Eq. (34) with LDP-ExO as it is more flexible.

**Applications.** As a main application of the private regret-DEC theory, in Section 5.5, we present the DEC theory for LDP learning in contextual bandits. We do not present the implications for bandits (which our framework subsumes easily) because it is already encompassed by non-private DEC framework for bandits [Foster et al., 2021, 2023b, Chen et al., 2024]: it is well-known that LDP bandits learning can be directly reduced to the standard bandits learning by adding additive noises (Laplace noise or Gaussian noise) to the random rewards.

# 5.5 Application: Contextual bandits

In this section, we focus on no-regret learning in contextual bandits, where the contexts can be adversarially chosen. Specifically, we introduce the (private) contextual DMSO framework: For each  $t = 1, \dots, T$ :

- The learner selects a decision  $\pi_t : \mathcal{X} \to \mathcal{A}$  and a private channel  $Q_t \in \mathcal{Q}$ .
- The environment selects context  $x_t \in \mathcal{X}$  and receives  $(\pi_t, Q_t)$ .
- The environment selects the action  $a_t = \pi_t(x_t)$  according to  $\pi_t$ , receives the reward  $r_t \sim \text{Rad}(f^*(x_t, a_t))$ ,<sup>5</sup> generates a noisy observation  $o_t \in \mathcal{O}$  via  $o_t \sim \mathsf{Q}_t(\cdot|x_t, a_t, r_t)$  and reveals it to the learner.

Here, we go beyond the private DMSO in that we do not assume the context of each user is stochastic; Instead, we allow  $x_t$  to depend on the history prior to step t, i.e., the context  $x_t$  can be chosen in an adversarial manner. The underlying reward function  $f^* : \mathcal{X} \times \mathcal{A} \to [-1, 1]$  encodes the mean reward value of the underlying environment, and we assume that the learner has access to a known reward function class  $\mathcal{F} \subseteq (\mathcal{X} \times \mathcal{A} \to [-1, 1])$  containing  $f^*$ . The decision space  $\Pi = (\mathcal{X} \to \mathcal{A})$  consists of all maps (policies) from the context space to the action space.

In contextual bandits, the regret of the learner is measured by

$$\operatorname{\mathbf{Reg}}_{\mathsf{DM}}(T) = \sum_{t=1}^{T} f^{\star}(x_t, \pi^{\star}(x_t)) - \mathbb{E}_{\pi_t \sim q_t} f^{\star}(x_t, \pi_t(x_t)),$$

where  $\pi^*$  is an optimal policy under the reward function  $f^*$ , i.e.,  $\pi^*(x) = \arg \max_{a \in \mathcal{A}} f^*(x, a)$  for  $x \in \mathcal{X}$ , and the expectation is with respect to  $\pi_t \sim q_t$ , the randomness of the choice of  $\pi_t$  at the *t*-th step.

<sup>&</sup>lt;sup>5</sup>For simplicity, we assume the reward is a binary random variable without loss of any generality.

Formulation in hybrid DMSO. We first briefly discuss how to frame this problem within hybrid DMSO. For  $\nu \in \Delta(\mathcal{X})$  and  $f \in \mathcal{F}$ , we define the contextual bandit model  $M_{\nu,f} : \Pi \to \Delta(\mathcal{O})$  as

$$(x, a, r) \sim M_{\nu, f}(\pi)$$
:  $x \sim \nu, a = \pi(x), r \sim \operatorname{Rad}(f(x, a)).$ 

We then consider the model class  $\mathcal{M}_{\mathcal{F},\mathsf{CB}} = \{M_{\nu,f} : \nu \in \Delta(\mathcal{X}), f \in \mathcal{F}\}$ , which is the model class of contextual bandits with stochastic context and mean reward function in  $\mathcal{F}$ . For each  $f \in \mathcal{F}$ , fspecifies a constraint  $\mathcal{P}_f$  as

$$\mathcal{P}_f := \{ M_{\nu,f}^{\sharp} : \nu \in \Delta(\mathcal{X}) \}, \tag{36}$$

i.e.,  $\mathcal{P}_f$  consists of all private (that is,  $M_{\nu,f}^{\sharp}$  includes the private channel choice) contextual bandit instances with mean reward function f, and we let  $\mathscr{P}_{\mathsf{cxt}} := \{\mathcal{P}_f : f \in \mathcal{F}\}$ . Then, the contextual bandits problem with function class  $\mathcal{F}$  can be framed within hybrid DMSO with constraint class  $\mathscr{P}_{\mathsf{cxt}}$ .

**Regret guarantees.** We show that LDP-ExO achieves a regret bound scaling with the private regret-DEC of  $\mathcal{M}_{\mathcal{F},CB}$ . Similar to Section 5.2, we assume that  $\mathcal{X}$  and  $\mathcal{A}$  are both finite throughout this section, mainly to avoid measure theoretic issues (our results do not have any dependence on  $|\mathcal{X}|$ ).

**Proposition 29.** Let  $T \ge 1, \delta \in (0, 1)$ . Suppose that  $\mathcal{X}$  and  $\mathcal{A}$  are finite, and the private regret-DEC r-dec<sup>LDP</sup><sub> $\varepsilon$ </sub> $(\mathcal{M}_{\mathcal{F},CB})$  is of moderate decay as a function of  $\varepsilon$ . Then, LDP-ExO (instantiated as in Appendix F.5.4) achieves with probability at least  $1 - \delta$ :

$$\frac{1}{T} \operatorname{Reg}_{\mathsf{DM}}(T) \leq O(\sqrt{\log T}) \cdot \left[ \operatorname{\mathsf{r-dec}}_{\bar{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}_{\mathcal{F},\mathsf{CB}}) + \bar{\varepsilon}(T) \right],$$
  
where  $\bar{\varepsilon}(T) = \inf_{\Delta \geq 0} \left( \Delta + \sqrt{\frac{\log N_{\infty}(\mathcal{F}, \Delta) + \log(1/\delta)}{\alpha^2 T}} \right)$ , and  $N_{\infty}(\mathcal{F}, \Delta)$  is the  $\Delta$ -covering number of  $\mathcal{F}$   
under  $L_{\infty}$ -norm (cf. Definition 20).

Therefore, up to a gap of the log-covering number of  $\mathcal{F}$ , the complexity of no-regret learning is characterized by the private regret-DEC of  $\mathcal{M}_{\mathcal{F},CB}$ . It is worth noting that our upper bound scales with the DEC of the *stochastic* contextual bandits, while it applies to any environment that generates contexts adversarially. Therefore, within the DEC framework, contextual decision making with (potentially) adversarial contexts is no more difficult than stochastic contexts.

This result is somewhat surprising, because with the LDP constraint, the learner can never directly observe the contexts. Indeed, this makes it challenging to estimate the ground truth mean reward function  $f^*$ , and previous works typically had to adopt problem-specific estimation methods. In contrast, Proposition 29 allows us to derive regret bounds by directly studying the DEC.

In the following, we apply our frameworks to derive near-optimal regret guarantees for linear contextual bandits and Lipschitz contextual bandits.

### 5.5.1 Linear contextual bandits

In the linear contextual bandits setting, we are given a bounded feature map  $\phi : \mathcal{X} \times \mathcal{A} \to \mathbf{B}^d(1)$ . The linear value function class  $\mathcal{F}_{\mathsf{Lin}}$  is given by

$$\mathcal{F}_{\mathsf{Lin}} = \{ f_{\theta} : f_{\theta}(x, a) = \langle \theta, \phi(x, a) \rangle \}_{\theta \in \mathbf{B}^{d}(1)},$$

Let  $\mathcal{M}_{\text{Lin-CB}}$  be the corresponding contextual bandits model class. In the following, we bound the private regret-DEC of  $\mathcal{M}_{\text{Lin-CB}}$  and provide a near-optimal guarantee for learning linear contextual bandits. Proof is presented in Appendix I.6.

**Theorem 30** (Near-optimal regret for linear contextual bandits). For the model class  $\mathcal{M}_{\text{Lin-CB}}$ , it holds that

$$\operatorname{r-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M}_{\operatorname{Lin-CB}}) \lesssim d\varepsilon.$$

Therefore, LDP-ExO achieves the following regret bound in linear contextual bandits with probability at least  $1 - \delta$ :

$$\operatorname{\mathbf{Reg}}_{\mathsf{DM}}(T) \le O\left(\frac{\sqrt{d^3T\log(T/\delta)}}{\alpha}\right).$$

The above regret bound of LDP-ExO is only a  $\tilde{O}(\sqrt{d})$  factor larger than the regret lower bound of  $\Omega\left(\sqrt{d^2T}/\alpha\right)$  for linear contextual bandits (detailed in Appendix I.8).

Our upper bound nearly settles the optimal regret for linear contextual bandits with LDP constraints. Previous works either suffer a  $T^{3/4}$  rate [Zheng et al., 2020], a  $\log^d(T) \cdot \sqrt{T}$  rate [Li et al., 2024], or require a strong assumption that the covariance matrix under *any* linear policy is well-conditioned [Han et al., 2021]. The benefit of our DEC framework is that it provides a systematic approach to obtain regret bounds, which reduces the problem to studying the private regret-DEC. We expect our techniques can be applied to a broader setting, e.g., RL with linear function approximation.

#### 5.5.2 Lipschitz contextual bandits with finite arms

As the next example, we consider a standard non-parametric contextual bandit problem: Lipschitz contextual bandits, with  $\mathcal{X}$  equipped with a metric  $\rho$ . The reward function class is

 $\mathcal{F}_{\mathsf{Lip}} = \{ f : \text{for any } a \in \mathcal{A}, f(\cdot, a) \text{ is a 1-Lipschitz function w.r.t. } \rho \},\$ 

and let  $\mathcal{M}_{\mathsf{Lip-CB}}$  be the corresponding contextual bandits model class. In the following proposition, we provide both upper and lower bounds for learning contextual bandits with  $\mathcal{F}_{\mathsf{Lip}}$ . We define  $N_{\rho}(\mathcal{X}, \Delta)$  to be the  $\Delta$ -covering number of  $\mathcal{X}$  under  $\rho$ . Details are deferred to Appendix I.7.

**Proposition 31.** For the model class  $\mathcal{M}_{Lip-CB}$ , it holds that

$$\operatorname{r-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M}_{\operatorname{Lip-CB}}) \lesssim \inf_{\Delta > 0} \left( \Delta + \sqrt{N_{\rho}(\mathcal{X}, \Delta) |\mathcal{A}|} \varepsilon \right)$$

For contextual bandits with mean reward function  $f^* \in \mathcal{F}_{Lip}$ , LDP-ExO (suitably instantiated as in Appendix I.7) achieves with probability at least  $1 - \delta$ 

$$\operatorname{\mathbf{Reg}}_{\mathsf{DM}}(T) \lesssim \inf_{\Delta > 0} \Big( T\Delta + N_{\rho}(\mathcal{X}, \Delta) \sqrt{\alpha^{-2} |\mathcal{A}| T \log(|\mathcal{A}|/\delta)} \Big).$$

On the other hand, for any  $\Delta \in (0, 1]$ , to learn an  $\Delta$ -optimal policy for  $\mathcal{M}_{\mathsf{Lip-CB}}$ , and  $\alpha$ -LDP algorithm must require T-round of interactions with  $T \gtrsim \frac{N_{\rho}(\mathcal{X}, 8\Delta)^2}{\alpha^2 \Delta^2}$  (cf. Appendix I.8).

In particular, when  $N_{\rho}(\mathcal{X}, \Delta) \simeq \Delta^{-d}$  (e.g.  $\mathcal{X}$  is a bounded domain in  $\mathbb{R}^{d}$ ), the minimax-optimal regret of privately learning  $\mathcal{F}_{\mathsf{Lip}}$  is  $\tilde{\Theta}(\alpha^{-\frac{1}{d+1}}T^{\frac{2d+1}{2d+2}})$ , up to a polynomial factor of  $|\mathcal{A}|$ .

#### 5.5.3 Concave-Lipschitz contextual bandits

Our final example is a generalization of the Lipschitz contextual bandits to continuously many arms. Assume that  $\mathcal{X}$  is equipped with a metric  $\rho$ ,  $\mathcal{A} \subset \mathbb{R}^{K}$  is a bounded convex domain, and

 $\mathcal{F}_{\mathsf{LC}} = (f : 1\text{-Lipschitz function in } (x, a) \in \mathcal{X} \times \mathcal{A}, \text{ concave in } a \in \mathcal{A}),$ 

Let  $\mathcal{M}_{LC-CB}$  be the corresponding contextual bandits model class. Similar to the Lipschitz contextual bandits, we have the following upper bound.

**Proposition 32.** For the model class  $\mathcal{M}_{LC-CB}$ , it holds that

$$\operatorname{r-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M}_{\operatorname{LC-CB}}) \leq \inf_{\Delta > 0} \left( \Delta + \tilde{O}(1) \sqrt{N_{\rho}(\mathcal{X}, \Delta) K^4} \varepsilon \right),$$

where we hide poly-logarithmic factors of the diameter of  $\mathcal{A}$ . For contextual bandits with mean reward function  $f^* \in \mathcal{F}_{LC}$ , LDP-ExO (suitably instantiated as in Appendix I.7) achieves with probability at least  $1 - \delta$ ,

$$\operatorname{Reg}_{\mathsf{DM}}(T) \leq \inf_{\Delta > 0} \Big( T\Delta + \tilde{O}\Big( N_{\rho}(\mathcal{X}, \Delta) \sqrt{\alpha^{-2} K^{5} T} \Big) \Big).$$

The upper bound above is derived by (1) reducing the contextual concave bandits to the concave bandits (without contexts) by bounding the corresponding DECs, and then (2) applying the results of Lattimore [2020]. This streamlined approach demonstrates again the advantage of the DEC framework, without which the reduction may not be easy, and we may instead need to repeat the analysis of Lattimore [2020].

Note that the lower bound of Proposition 31 also applies here (cf. Appendix I.8). Therefore, when  $N_{\rho}(\mathcal{X}, \Delta) \simeq \Delta^{-d}$ , the minimax-optimal regret of privately learning  $\mathcal{F}_{\mathsf{LC}}$  is also  $\tilde{\Theta}(\alpha^{-\frac{1}{d+1}}T^{\frac{2d+1}{2d+2}})$ , up to a polynomial factor of K.

# 6 Local Minimaxity, Learnability, and Joint Privacy

In this section, we still focus on locally private learning, and discuss how our framework relates various other notions, including local-minimax complexity, learnability, and joint differential privacy.

## 6.1 Local-minimax optimality

In this section, we demonstrate that the private PAC-DEC framework also applies to local-minimax statistical estimation under LDP, recovering the existing results in Duchi and Ruan [2024] and also providing new insights.

**Local-minimax risk.** For any learning problem given by  $\mathcal{M}$  and a model  $M_0 \in \mathcal{M}$ , we define the  $\alpha$ -LDP *local-minimax* risk at  $M_0$  as

$$\mathfrak{M}_{T}^{\mathrm{loc}}(\mathcal{M}, M_{0}) := \sup_{M_{1} \in \mathcal{M}} \inf_{\mathsf{Alg}} \sup_{M \in \{M_{0}, M_{1}\}} \mathbb{E}^{M, \mathrm{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)],$$
(37)

where the  $\inf_{Alg}$  is taken over all possible *T*-round  $\alpha$ -LDP algorithms. In words, the local minimax risk measures the best performance the algorithm can achieve when it is given the knowledge two

possible models. This risk is called local because it measures the difficulty of a particular model  $M_0$  against a *single* worst-case alternative  $M_1 \in \mathcal{M}$ .

Modulus of continuity is a commonly studied complexity measure in statistical estimation and is shown to capture the complexity of various problem classes [Donoho and Liu, 1991, Juditsky and Nemirovski, 2009, Polyanskiy and Wu, 2019]. Under local privacy constraints, Duchi and Ruan [2024] show that the following TV modulus of continuity captures the difficulty of *local* minimaxoptimal statistical estimation: They show that, for functional estimation, the minimax risk is characterized by the following TV variant of modulus of continuity:

$$w_{\varepsilon}(\mathcal{M}, M_0) := \sup_{M_1 \in \mathcal{M}} \left\{ \left| \pi^{M_1} - \pi^{M_0} \right| \mid D_{\mathrm{TV}}\left(M_1, M_0\right) \le \varepsilon \right\}.$$
(38)

We note that under LDP, the TV modulus of continuity also characterizes the complexity of linear functional estimation with a convex model class, as shown in Rohde and Steinberger [2020].

In the following, we study the local-minimax complexity of any LDP PAC learning problem (not necessarily limited to statistical tasks as per Definition 10).

**Local DEC theory.** We show that the local-minimax risk of any LDP PAC learning problem is tightly captured by the following *local* DEC:

$$\mathsf{p-dec}_{\varepsilon}^{\mathrm{loc}}(\mathcal{M}, M_0) = \sup_{M \in \mathcal{M}} \left\{ \inf_{\pi \in \Pi} L(M_1, \pi) + L(M_0, \pi) \; \middle| \; \sup_{\pi \in \Pi} D_{\mathrm{TV}}\left(M_1(\pi), M_0(\pi)\right) \le \varepsilon \right\}.$$
(39)

In particular, for functional estimation problems (where  $\Pi = \mathbb{R}$ , and  $L(M, \pi) = |\pi^M - \pi|$ ), the definition above *exactly* recovers the modulus of continuity (38). Moreover, for stochastic convex optimization, local DEC also agrees with the modulus of continuity considered in Duchi et al. [2016]. Therefore, local DEC can be regarded as the natural generalization of the modulus of continuity to any local-minimax PAC learning problem.

As an corollary of the private PAC-DEC lower and upper bounds (Appendix I.2 and Theorem 21), local DEC provides the following nearly-optimal characterization of the local-minimax risk. Details are presented in Appendix J.1.

**Theorem 33.** Let  $T \ge 1$ , model class  $\mathcal{M}$  be given. Suppose that the loss function L is bounded in  $[0, L_{\max}]$ , and for any model  $M \in \mathcal{M}$ , we have  $\min_{\pi} L(M, \pi) = 0$ . Then, the local-minimax risk at a model  $M_0 \in \mathcal{M}$  is bounded as

$$\frac{1}{8}\mathsf{p-dec}_{\underline{\varepsilon}(T)}^{\mathrm{loc}}(\mathcal{M}, M_0) \le \mathfrak{M}_T^{\mathrm{loc}}(\mathcal{M}, M_0) \le \inf_{\delta > 0} \Big(\mathsf{p-dec}_{\overline{\varepsilon}_{\delta}(T)}^{\mathrm{loc}}(\mathcal{M}, M_0) + \delta L_{\mathrm{max}}\Big),$$

where  $\underline{\varepsilon}(T) = \frac{c_0}{\sqrt{T}}$  and  $\overline{\varepsilon}_{\delta}(T) = \frac{c_1 \log(1/\delta)}{\sqrt{T}}$ .

Therefore, the local-minimax risk of interactive learning under LDP is tightly captured by the local DEC. For the particular case of functional estimation, local DEC is equivalent to the TV modulus of continuity. Hence, up to logarithmic factors, we recover the characterization of the LDP local-minimax risk of Duchi and Ruan [2024], assuming certain growth conditions. The fact that such a characterization extends to statistical estimation tasks with interaction and general loss function is a testament to the unifying power of the DEC framework.

Furthermore, from the definition of local DEC (39), we can gain some quantitative insights into how locality reduces the difficulty of learning. More specifically, with locality, the algorithm only needs

to distinguish between two models  $\{M_1, M_0\}$ , and hence avoids (1) the complexity of estimation, e.g. the log-cardinality of the model class or the function class (cf. Theorem 21), and (2) the complexity of exploration, because it suffices to pick the best distinguishing decision  $\pi$  that maximizes  $D_{\text{TV}}(M_1(\pi), M_0(\pi))$ . Hence, even though the local-minimax formulation avoids the undesirable worst-case behavior of the global-minimax LDP learning, it may be too restrictive as it trivializes the difficulty of both interaction (exploration) and estimation.

#### 6.2 Finite-time learnability under LDP

In learning theory, a central task is to investigate complexity measures that characterize the *finite-time learnability* of certain problem classes, e.g., VC dimension for binary classification, Littlestone dimension [Littlestone, 1988] for online classification [Ben-David et al., 2009], and their real-valued analogues for regression and online learning (see e.g. Rakhlin and Sridharan [2014]). Further, Bun et al. [2020], Alon et al. [2022] show that *jointly private* classification is possible if and only if the Littlestone dimension is finite. Recently, the notion of fractional covering number Definition 5 was proposed by Chen et al. [2024] and shown to characterize the non-private learnability of any stochastic bandits problems.

Following this line of work, in this section, we characterize the LDP learnability of any learning problem with reward-based loss through its fractional covering number, generalizing the results of Chen et al. [2024]. To rigorously formulate the notion of learnability, we introduce the following minimax sample complexity under LDP: For a model class  $\mathcal{M} \subset (\Pi \to \Delta(\mathcal{Z}))$ , risk level  $\Delta > 0$ , we define<sup>6</sup>

$$\mathfrak{C}_{\Delta}^{\mathsf{LDP}}(\mathcal{M}) := \min\left\{T : \exists T \text{-round } \alpha \text{-LDP algorithm Alg s.t. } \sup_{M \in \mathcal{M}} \mathbb{E}^{M,\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \leq \Delta\right\}.$$
(40)

A model class  $\mathcal{M}$  is  $\alpha$ -LDP learnable if for all risk levels  $\Delta > 0$ ,  $\mathfrak{C}_{\Delta}^{\mathsf{LDP}}(\mathcal{M}) < +\infty$ , i.e., there is an  $\alpha$ -LDP algorithm that achieves  $\Delta$ -risk in finite number of rounds.

We first show that fractional covering number provides a lower bound for any LDP learning problem, following the approach of Chen et al. [2024].

**Theorem 34.** Let  $T \ge 1$ ,  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  be a model class. Suppose that there is a T-round  $\alpha$ -LDP algorithm Alg that achieves that for all  $M \in \mathcal{M}$ ,  $\mathbf{Risk}_{\mathsf{DM}}(T) \le \Delta$  with probability at least  $\frac{1}{2}$  under  $\mathbb{P}^{M,\mathsf{Alg}}$ . Then it holds that

$$T \ge \frac{\log N_{\mathsf{frac}}(\mathcal{M}, \Delta) - 2}{2(e^{\alpha} - 1)^2}.$$

This result differs from the fractional covering number lower bound for non-private learning [Chen et al., 2024], which additionally involves the KL radius of  $\mathcal{M}$ :

$$C_{\mathrm{KL}}(\mathcal{M}) = \inf_{\overline{M}} \sup_{M \in \mathcal{M}, \pi \in \Pi} D_{\mathrm{KL}}(M(\pi) \parallel \overline{M}(\pi)).$$

In non-private learning, the dependence on  $C_{\text{KL}}^{-1}$  in the lower bound can be unavoidable (e.g., for binary classification, see also our discussion in Section 6.3.1). By contrast, Theorem 34 applies to LDP learning for *any* problem class, even when  $C_{\text{KL}} = +\infty$ .

<sup>&</sup>lt;sup>6</sup>We note that both the minimax sample complexity  $\mathfrak{C}_{\Delta}^{LDP}(\mathcal{M})$  and the fractional covering number  $N_{\text{frac}}(\mathcal{M}, \Delta)$  depend on the loss function L implicitly.

**Fractional covering number upper bound.** When the loss function is reward-based, we show that fractional covering number also provides a "brute-force" upper bound.

**Proposition 35.** Let  $T \ge 1$ ,  $\delta \in (0, 1)$ , and  $\mathcal{M}$  be a model class. Suppose that the loss function is reward-based, then there is a "brute-force" algorithm (Algorithm 5) such that with probability at least  $1 - \delta$ ,

$$\mathbf{Risk}_{\mathsf{DM}}(T) \le \Delta + O(\log(T/\delta))\sqrt{\frac{N_{\mathsf{frac}}(\mathcal{M}, \Delta)}{\alpha^2 T}}.$$

Combining the above upper bound with the lower bound of  $N_{\text{frac}}(\mathcal{M}, \Delta)$ , we have shown that  $N_{\text{frac}}(\mathcal{M}, \Delta)$  characterizes the sample complexity of LDP learning the model class, up to an exponential gap:

$$\frac{\log N_{\mathsf{frac}}(\mathcal{M}, 2\Delta)}{\alpha^2} \lesssim \mathfrak{C}_{\Delta}^{\mathsf{LDP}}(\mathcal{M}) \lesssim \frac{N_{\mathsf{frac}}(\mathcal{M}, \Delta/2)}{\alpha^2 \Delta^2},\tag{41}$$

where we omit poly-logarithmic factors. We remark that the gap between the lower and upper bounds cannot be improved in terms of fractional covering number alone:

- For classification with the parity class *F*<sub>parity</sub>, a lower bound scaling linearly with N<sub>frac</sub>(*F*<sub>parity</sub>, Δ/2) = |*F*<sub>parity</sub>| can be obtained (Proposition B.6), meaning the upper bound can be tight even for the *statistical* tasks (as per Definition 10).
- For the problem of Multi-Armed Bandits, we also have  $N_{\mathsf{frac}}(\mathcal{M}, 1/2) = |\mathcal{A}|$ , while  $\Omega\left(\frac{|\mathcal{A}|}{\alpha^2 \varepsilon^2}\right)$  samples are necessary to learn an  $\varepsilon$ -optimal policy.
- For linear bandits,  $\log N_{\text{frac}}(\mathcal{M}, 1/2) = \Omega(d)$ , and it is known that  $\tilde{O}\left(\frac{d^2}{\alpha^2 \varepsilon^2}\right)$  samples are sufficient to learn an  $\varepsilon$ -optimal policy, meaning that the lower bound can also be (nearly) tight.

While the exponential gap in Eq. (41) is unavoidable solely with fractional covering number, we have shown that the upper bound can be improved with DEC (at least for convex model classes, cf. Theorem 22).

A direct implication of Eq. (41) is that the finiteness of fractional covering number characterizes the finite-time learnability under LDP, as long as the loss function is reward-based.

**Theorem 36** (LDP learnability). Under reward-based loss, the problem class is LDP learnable if and only if  $N_{\text{frac}}(\mathcal{M}, \Delta) < \infty$  for all  $\Delta > 0$ .

The learnability characterization above is similar to the bandit learnability characterization in Chen et al. [2024]. However, we do show that fractional covering number characterizes the learnability under LDP for *any* model class  $\mathcal{M}$ , while for non-private learning fractional covering number only characterizes the learnability of model class with a bounded  $C_{\rm KL}$ .

As an application of Theorem 36, in Appendix C.2 we discuss how the fractional covering number provides insights into the LDP learnability of *regression*.

#### 6.3 Learnability under joint differential privacy

Parallel to the concept of local differential privacy (LDP), there is a notion of *joint differential* privacy (JDP) [Dwork et al., 2006].<sup>7</sup> For simplicity, in the following discussion, we focus on the notion of pure JDP for statistical problems. Detailed discussion for interactive decision making is deferred to Appendix C.1.

In this setting, the learner (algorithm) is given a dataset  $\mathcal{H}_T = (z_1, \cdots, z_T)$  consisting of i.i.d observations, i.e.,  $z_1, \cdots, z_T \sim M^*$  for a model  $M^* \in \Delta(\mathcal{Z})$ . As always, we assume the learner is given a model class  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$  that contains  $M^*$ .

For this setting, an algorithm (learner) is simply a map  $Alg : \mathbb{Z}^T \to \Delta(\Pi)$ . In the following, we define  $\alpha$ -JDP algorithms.

**Definition 11** (Pure JDP for statistical problems). For two sequence of observations  $\mathcal{H}_T = (z_1, \dots, z_T), \ \mathcal{H}'_T = (z'_1, \dots, z'_T) \in \mathcal{Z}^T$ , they are neighbored if there is at most one index  $t \in [T]$  such that  $z_t \neq z'_t$ . An algorithm Alg preserves  $\alpha$ -JDP if for any neighbored dataset  $\mathcal{H}_T, \mathcal{H}'_T$  and any measurable set  $E \subseteq \Pi$ ,

$$\mathbb{P}^{\mathsf{Alg}}(\pi_{T+1} \in E | \mathcal{H}_T) \le e^{\alpha} \cdot \mathbb{P}^{\mathsf{Alg}}(\pi_{T+1} \in E | \mathcal{H}'_T).$$

Similar to Theorem 34, we show that the fractional covering number also provides a lower bound for JDP learning.

**Proposition 37** (Fractional covering number lower bound for JDP learning). Let  $T \ge 1$ , model class  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$  be given. Suppose that Alg is a *T*-round  $\alpha$ -JDP algorithm, such that it achieves  $\operatorname{\mathbf{Risk}}_{DM}(T) \le \Delta$  with probability at least  $\frac{1}{2}$  under  $\mathbb{P}^{M,\operatorname{Alg}}$  for any  $M \in \mathcal{M}$ . Then it holds that

$$T \ge \frac{\log N_{\mathsf{frac}}(\mathcal{M}, \Delta) - \log 2}{\alpha}.$$

For binary classification under pure JDP, Beimel et al. [2013a] provide both lower and upper bounds of the sample complexity in terms of the *representation dimension*. As we discuss in Section 6.3.1, for binary classification, fractional covering number is equivalent to the representation dimension (up to an additive constant, Proposition 39).

**Pure JDP learnability**  $\equiv$  **LDP learnability.** It is clear that if an algorithm preserves  $\alpha$ -LDP, then it also preserves  $\alpha$ -JDP. Therefore, when the loss function is reward-based, as the finiteness of fractional covering number characterizes the LDP learnability, it also characterizes the JDP learnability.<sup>8</sup>

**Theorem 38.** Let privacy parameter  $\alpha > 0$ , model class  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ , and the reward-based loss function L be given. Then the following statements are equivalent:

- (1)  $\mathcal{M}$  is  $\alpha$ -LDP learnable,
- (2)  $\mathcal{M}$  is  $\alpha$ -JDP learnable, and
- (3)  $N_{\mathsf{frac}}(\mathcal{M}, \Delta) < +\infty \text{ for all } \Delta > 0.$

<sup>&</sup>lt;sup>7</sup>This notion is often referred to simply as "differential privacy." To distinguish it from local differential privacy, we use the term "joint differential privacy," as it preserves the privacy of the data points in a dataset jointly.

<sup>&</sup>lt;sup>8</sup>We note that for JDP learning in statistical problems, the exponential mechanism achieves a better upper bound scaling with log  $N_{\text{frac}}(\mathcal{M}, \Delta)$  (see e.g. Beimel et al. [2013b]). However, for interactive learning (with or without JDP), an upper bound scaling linearly with  $N_{\text{frac}}(\mathcal{M}, \Delta)$  can be necessary in general [Chen et al., 2024].

We note that a similar argument also applies to interactive decision making problems, as the fractional covering number also provides a lower bound for interactive learning under JDP (Appendix C.1).

### 6.3.1 Connection to representation dimension and Littlestone dimension

In this section, we discuss the connection between fractional covering number and two well-studied complexity measures for binary classification: representation dimension [Beimel et al., 2013a] and Littlestone's dimension [Littlestone, 1988].

**Representation dimension.** It has been known that for JDP binary classification with a function class  $\mathcal{F} \subseteq (\mathcal{X} \to \{0, 1\})$ , the sample complexity of (proper or improper) learning is tightly characterized by the following *representation dimension* [Beimel et al., 2013a]. For the simplicity of presentation, we focus on proper learning.

**Definition 12.** A distribution  $\mathscr{H}$  over finite subsets of  $\mathcal{F}$  is an  $\varepsilon$ -probabilistic representation of  $\mathcal{F}$  if for any distribution  $\nu \in \Delta(\mathscr{X})$  and  $f \in \mathcal{F}$ , with probability at least  $\frac{3}{4}$  over  $\mathcal{H} \sim \mathscr{H}$ , there exists  $h \in \mathcal{H}$  such that

$$\mathbb{P}_{x \sim \nu}(h(x) \neq f(x)) \le \varepsilon.$$

The size of  $\mathscr{H}$  is defined as  $\operatorname{size}(\mathscr{H}) = \sup_{\mathcal{H} \in \operatorname{supp}(\mathscr{H})} \log |\mathcal{H}|$ . The representation dimension of  $\mathcal{F}$  is then defined as

$$\mathsf{RDim}_{\varepsilon}(\mathcal{F}) := \inf_{\mathscr{H}} \ \mathrm{size}(\mathscr{H}),$$

where  $\inf_{\mathscr{H}}$  is taken over all  $\varepsilon$ -probabilistic representations of  $\mathcal{F}$ .

We show that for binary classification, the fractional covering number is equivalent to the representation dimension. Recall that for binary classification, the loss function (implicit in the definition of the fractional covering number, cf. Section 5.2) is given by

$$L(M,f) := \mathbb{P}_{(x,y)\sim M}(f(x) \neq y) - \inf_{f^{\star} \in \mathcal{F}} \mathbb{P}_{(x,y)\sim M}(f^{\star}(x) \neq y).$$

**Proposition 39.** For any  $\varepsilon \in [0, 1]$ , it holds that

$$|\mathsf{RDim}_{\varepsilon}(\mathcal{F}) - \log N_{\mathsf{frac}}(\mathcal{F}, \Delta)| \leq 2.$$

The details are postponed to Appendix J.7. This equivalence also agrees with the fact that both representation dimension and fractional covering number characterizes the JDP learnability of classification.

**Littlestone dimension.** It is known that for binary class,  $\mathsf{RDim}(\mathcal{F}) \geq \Omega(\mathsf{LDim}(\mathcal{F}))$  [Feldman and Xiao, 2014], and there exists classes with  $\mathsf{LDim}(\mathcal{F}) = 2$  while  $\mathsf{RDim}(\mathcal{F})$  arbitrary large. Hence, LDP learnability is a stronger notion of complexity of a class than online learnability.

It is also well-known that for binary classification, there is an equivalence between learnability under *approximate* JDP and online learnability [Bun et al., 2020, Alon et al., 2022]. For regression, joint DP learnability can be achieved under a certain growth condition on the sequential fat-shattering dimension [Golowich, 2021]. However, to learn a binary class  $\mathcal{F}$  under approximate JDP, it is only known that  $\log^*(\mathsf{LDim}(\mathcal{F}))$  samples are necessary [Bun et al., 2020].
# 7 Conclusion

We presented a systematic approach to analyzing problems of decision making with a changing environment and constraints on the amount of information received by the learner. While this approach yields upper and lower bounds on minimax performance, the question of efficient algorithms is entirely open.

### Acknowledgments

We acknowledge support from ARO through award W911NF-21-1-0328, as well as Simons Foundation and the NSF through awards DMS-2031883 and PHY-2019786.

### References

- N. Alon, M. Bun, R. Livni, M. Malliaris, and S. Moran. Private and online learnability are equivalent. ACM Journal of the ACM (JACM), 69(4):1–34, 2022.
- H. Asi, V. Feldman, and K. Talwar. Optimal algorithms for mean estimation under local differential privacy. In *International Conference on Machine Learning*, pages 1046–1056. PMLR, 2022.
- H. Asi, J. Ullman, and L. Zakynthinou. From robustness to privacy and back. In International Conference on Machine Learning, pages 1121–1146. PMLR, 2023.
- H. Asi, V. Feldman, J. Nelson, H. Nguyen, and K. Talwar. Fast optimal locally private mean estimation via random projections. *Advances in Neural Information Processing Systems*, 36, 2024.
- A. Beimel, K. Nissim, and U. Stemmer. Characterizing the sample complexity of private learners. In Proceedings of the 4th conference on Innovations in Theoretical Computer Science, pages 97–110, 2013a.
- A. Beimel, K. Nissim, and U. Stemmer. Private learning and sanitization: Pure vs. approximate differential privacy. In *International Workshop on Approximation Algorithms for Combinatorial Optimization*, pages 363–378. Springer, 2013b.
- S. Ben-David, D. Pal, and S. Shalev-Shwartz. Agnostic online learning. In *Proceedings of the 22th* Annual Conference on Learning Theory, 2009.
- T. Berrett and C. Butucea. Locally private non-asymptotic testing of discrete distributions is faster using interactive mechanisms. *Advances in Neural Information Processing Systems*, 33: 3164–3173, 2020.
- T. B. Berrett, L. Györfi, and H. Walk. Strongly universally consistent nonparametric regression and classification with privatised data. 2021.
- A. Blum, M. Furst, J. Jackson, M. Kearns, Y. Mansour, and S. Rudich. Weakly learning dnf and characterizing statistical query learning using fourier analysis. In *Proceedings of the twenty-sixth* annual ACM symposium on Theory of computing, pages 253–262, 1994.
- M. Brennan, G. Bresler, S. B. Hopkins, J. Li, and T. Schramm. Statistical query algorithms and low-degree tests are almost equivalent. arXiv preprint arXiv:2009.06107, 2020.

- N. H. Bshouty and V. Feldman. On using extended statistical queries to avoid membership queries. Journal of Machine Learning Research, 2(Feb):359–395, 2002.
- S. Bubeck, J. Ding, R. Eldan, and M. Z. Rácz. Testing for high-dimensional geometry in random graphs. *Random Structures & Algorithms*, 49(3):503–532, 2016.
- M. Bun, R. Livni, and S. Moran. An equivalence between private classification and online prediction. In 2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS), pages 389–402. IEEE, 2020.
- C. Butucea and Y. Issartel. Locally differentially private estimation of functionals of discrete distributions. Advances in Neural Information Processing Systems, 34:24753–24764, 2021.
- C. Butucea, A. Rohde, and L. Steinberger. Interactive versus noninteractive locally differentially private estimation: Two elbows for the quadratic functional. *The Annals of Statistics*, 51(2): 464–486, 2023.
- C. Canonne, S. B. Hopkins, J. Li, A. Liu, and S. Narayanan. The full landscape of robust mean testing: Sharp separations between oblivious and adaptive contamination. In 2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS), pages 2159–2168. IEEE, 2023.
- G. Casella and R. Berger. *Statistical Inference*. Duxbury advanced series in statistics and decision sciences. Thomson Learning, 2002.
- F. Chen, S. Mei, and Y. Bai. Unified algorithms for rl with decision-estimation coefficients: pac, reward-free, preference-based learning, and beyond. arXiv preprint arXiv:2209.11745, 2022.
- F. Chen, D. J. Foster, Y. Han, J. Qian, A. Rakhlin, and Y. Xu. Assouad, fano, and le cam with interaction: A unifying lower bound framework and characterization for bandit learnability. arXiv preprint arXiv:2410.05117, 2024.
- I. Diakonikolas and D. M. Kane. Recent advances in algorithmic high-dimensional robust statistics. arXiv preprint arXiv:1911.05911, 2019.
- I. Diakonikolas and D. M. Kane. Algorithmic high-dimensional robust statistics. Cambridge university press, 2023.
- I. Diakonikolas, D. M. Kane, and A. Stewart. Statistical query lower bounds for robust estimation of high-dimensional gaussians and gaussian mixtures. In 2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS), pages 73–84. IEEE, 2017.
- I. Diakonikolas, G. Kamath, D. Kane, J. Li, A. Moitra, and A. Stewart. Robust estimators in high-dimensions without the computational intractability. *SIAM Journal on Computing*, 48(2): 742–864, 2019.
- D. L. Donoho and R. C. Liu. Geometrizing rates of convergence, II. *The Annals of Statistics*, pages 633–667, 1991.
- J. Duchi and R. Rogers. Lower bounds for locally private estimation via communication complexity. In *Conference on Learning Theory*, pages 1161–1191. PMLR, 2019.
- J. C. Duchi. Lecture notes on statistics and information theory. 2023.
- J. C. Duchi and F. Ruan. The right complexity measure in locally private estimation: It is not the fisher information. *The Annals of Statistics*, 52(1):1–51, 2024.

- J. C. Duchi, M. I. Jordan, and M. J. Wainwright. Local privacy and statistical minimax rates. In 2013 IEEE 54th annual symposium on foundations of computer science, pages 429–438. IEEE, 2013.
- J. C. Duchi, J. Lafferty, Y. Zhu, et al. Local minimax complexity of stochastic convex optimization. Advances in Neural Information Processing Systems, 29, 2016.
- J. C. Duchi, M. I. Jordan, and M. J. Wainwright. Minimax optimal procedures for locally private estimation. *Journal of the American Statistical Association*, 113(521):182–201, 2018.
- C. Dwork, F. McSherry, K. Nissim, and A. Smith. Calibrating noise to sensitivity in private data analysis. In Theory of Cryptography: Third Theory of Cryptography Conference, TCC 2006, New York, NY, USA, March 4-7, 2006. Proceedings 3, pages 265–284. Springer, 2006.
- K. Fan. Minimax theorems. Proceedings of the National Academy of Sciences, 39(1):42-47, 1953.
- V. Feldman. A general characterization of the statistical query complexity. In Conference on learning theory, pages 785–830. PMLR, 2017.
- V. Feldman and D. Xiao. Sample complexity bounds on differentially private learning via communication complexity. In *Conference on Learning Theory*, pages 1000–1019. PMLR, 2014.
- V. Feldman, W. Perkins, and S. Vempala. On the complexity of random satisfiability problems with planted solutions. In *Proceedings of the forty-seventh annual ACM symposium on Theory* of Computing, pages 77–86, 2015.
- V. Feldman, E. Grigorescu, L. Reyzin, S. S. Vempala, and Y. Xiao. Statistical algorithms and a lower bound for detecting planted cliques. *Journal of the ACM (JACM)*, 64(2):1–37, 2017.
- D. Foster, D. J. Foster, N. Golowich, and A. Rakhlin. On the complexity of multi-agent decision making: From learning in games to partial monitoring. In *The Thirty Sixth Annual Conference* on Learning Theory, pages 2678–2792. PMLR, 2023a.
- D. J. Foster, S. M. Kakade, J. Qian, and A. Rakhlin. The statistical complexity of interactive decision making. arXiv preprint arXiv:2112.13487, 2021.
- D. J. Foster, N. Golowich, J. Qian, A. Rakhlin, and A. Sekhari. A note on model-free reinforcement learning with the decision-estimation coefficient. arXiv preprint arXiv:2211.14250, 2022a.
- D. J. Foster, A. Rakhlin, A. Sekhari, and K. Sridharan. On the complexity of adversarial decision making. Advances in Neural Information Processing Systems, 35:35404–35417, 2022b.
- D. J. Foster, N. Golowich, and Y. Han. Tight guarantees for interactive decision making with the decision-estimation coefficient. In *The Thirty Sixth Annual Conference on Learning Theory*, pages 3969–4043. PMLR, 2023b.
- D. J. Foster, Y. Han, J. Qian, and A. Rakhlin. Online estimation via offline estimation: An information-theoretic framework. arXiv preprint arXiv:2404.10122, 2024.
- E. Garcelon, V. Perchet, C. Pike-Burke, and M. Pirotta. Local differential privacy for regret minimization in reinforcement learning. Advances in Neural Information Processing Systems, 34: 10561–10573, 2021.
- K. Georgiev and S. Hopkins. Privacy induces robustness: Information-computation gaps and sparse mean estimation. Advances in neural information processing systems, 35:6829–6842, 2022.

- M. Glasgow and A. Rakhlin. Tight bounds for  $\gamma$ -regret via the decision-estimation coefficient. arXiv preprint arXiv:2303.03327, 2023.
- N. Golowich. Differentially private nonparametric regression under a growth condition. In *Conference on Learning Theory*, pages 2149–2192. PMLR, 2021.
- S. Gopi, G. Kamath, J. Kulkarni, A. Nikolov, Z. S. Wu, and H. Zhang. Locally private hypothesis selection. In *Conference on Learning Theory*, pages 1785–1816. PMLR, 2020.
- Y. Han, Z. Liang, Y. Wang, and J. Zhang. Generalized linear bandits with local differential privacy. Advances in Neural Information Processing Systems, 34:26511–26522, 2021.
- S. Hanneke, R. Livni, and S. Moran. Online learning with simple predictors and a combinatorial characterization of minimax in 0/1 games. In *Conference on Learning Theory*, pages 2289–2314. PMLR, 2021.
- E. Hazan, T. Koren, R. Livni, and Y. Mansour. Online learning with low rank experts. In 29th Annual Conference on Learning Theory, pages 1096–1114, 2016.
- J. He, J. Zhang, and R. Q. Zhang. A reduction from linear contextual bandits lower bounds to estimations lower bounds. In *International Conference on Machine Learning*, pages 8660–8677. PMLR, 2022.
- S. B. Hopkins, G. Kamath, M. Majid, and S. Narayanan. Robustness implies privacy in statistical estimation. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing*, pages 497–506, 2023.
- P. J. Huber. A robust version of the probability ratio test. The Annals of Mathematical Statistics, pages 1753–1758, 1965.
- P. J. Huber. Robust estimation of a location parameter. In *Breakthroughs in statistics: Methodology* and distribution, pages 492–518. Springer, 1992.
- P. J. Huber and E. M. Ronchetti. Robust statistics. John Wiley & Sons, 2011.
- T. Jayram. Hellinger strikes back: A note on the multi-party information complexity of and. In International Workshop on Approximation Algorithms for Combinatorial Optimization, pages 562–573. Springer, 2009.
- N. Joshi, T. Misiakiewicz, and N. Srebro. On the complexity of learning sparse functions with statistical and gradient queries. arXiv preprint arXiv:2407.05622, 2024.
- A. B. Juditsky and A. S. Nemirovski. Nonparametric estimation by convex programming. *The Annals of Statistics*, 37(5A):2278 – 2300, 2009. doi: 10.1214/08-AOS654. URL https://doi.org/10.1214/08-AOS654.
- S. P. Kasiviswanathan, H. K. Lee, K. Nissim, S. Raskhodnikova, and A. Smith. What can we learn privately? *SIAM Journal on Computing*, 40(3):793–826, 2011.
- M. Kearns. Efficient noise-tolerant learning from statistical queries. *Journal of the ACM (JACM)*, 45(6):983–1006, 1998.
- T. Lattimore. Improved regret for zeroth-order adversarial bandit convex optimisation. *Mathematical Statistics and Learning*, 2(3):311–334, 2020.

- T. Lattimore and A. Gyorgy. Mirror descent and the information ratio. In *Conference on Learning Theory*, pages 2965–2992. PMLR, 2021.
- T. Lattimore and C. Szepesvári. Exploration by optimisation in partial monitoring. In *Conference* on *Learning Theory*, pages 2488–2515. PMLR, 2020.
- G. Li, P. Kamath, D. J. Foster, and N. Srebro. Understanding the eluder dimension. Advances in Neural Information Processing Systems, 35:23737–23750, 2022.
- J. Li, D. Simchi-Levi, and Y. Wang. On the optimal regret of locally private linear contextual bandit. arXiv preprint arXiv:2404.09413, 2024.
- M. Li, T. B. Berrett, and Y. Yu. On robustness and local differential privacy. The Annals of Statistics, 51(2):717–737, 2023.
- C. Liao, J. He, and Q. Gu. Locally differentially private reinforcement learning for linear mixture markov decision processes. In Asian Conference on Machine Learning, pages 627–642. PMLR, 2023.
- N. Littlestone. Learning quickly when irrelevant attributes abound: A new linear-threshold algorithm. Machine learning, 2(4):285–318, 1988.
- A. Liu and A. Moitra. Settling the robust learnability of mixtures of gaussians. In Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing, pages 518–531, 2021.
- M. J. Osborne and A. Rubinstein. A course in game theory. MIT press, 1994.
- Y. Polyanskiy and Y. Wu. Dualizing le cam's method for functional estimation, with applications to estimating the unseens. arXiv preprint arXiv:1902.05616, 2019.
- A. F. Pour, H. Ashtiani, and S. Asoodeh. Sample-optimal locally private hypothesis selection and the provable benefits of interactivity. In *The Thirty Seventh Annual Conference on Learning Theory*, pages 4240–4275. PMLR, 2024.
- A. Rakhlin and K. Sridharan. Online nonparametric regression. In Conference on Learning Theory, 2014.
- A. Rakhlin, K. Sridharan, and A. Tewari. Online learning: Stochastic, constrained, and smoothed adversaries. *Advances in neural information processing systems*, 24, 2011.
- A. Rohde and L. Steinberger. Geometrizing rates of convergence under local differential privacy constraints. The Annals of Statistics, 48(5):2646–2670, 2020.
- D. Russo and B. Van Roy. Learning to optimize via posterior sampling. Mathematics of Operations Research, 39(4):1221–1243, 2014.
- D. Russo and B. Van Roy. Learning to optimize via information-directed sampling. *Operations Research*, 66(1):230–252, 2018.
- R. Shariff and O. Sheffet. Differentially private contextual linear bandits. Advances in Neural Information Processing Systems, 31, 2018.
- G. Vietri, B. Balle, A. Krishnamurthy, and S. Wu. Private reinforcement learning with pac and regret guarantees. In *International Conference on Machine Learning*, pages 9754–9764. PMLR, 2020.

- D. Wang and J. Xu. On sparse linear regression in the local differential privacy model. In *Interna*tional Conference on Machine Learning, pages 6628–6637. PMLR, 2019.
- S. L. Warner. Randomized response: A survey technique for eliminating evasive answer bias. Journal of the American statistical association, 60(309):63–69, 1965.
- K. Zheng, T. Cai, W. Huang, Z. Li, and L. Wang. Locally differentially private (contextual) bandits learning. Advances in Neural Information Processing Systems, 33:12300–12310, 2020.

# Contents

1	Introduction         1.1       Contributions	$\frac{1}{2}$
	1.2 Related work	3
2	Overview of Results         2.1       Hybrid DMSO         2.2       Query-based learning         2.3       Locally differentially private learning         2.4       Robust decision making         2.5       Regret guarantees for hybrid DMSO         2.5.1       Example: Smooth adversaries	4 7 8 10 11 12
3	DEC Theory for Hybrid DMSO         3.1 Guarantees for PAC learning         3.2 Guarantees for no-regret learning         3.3 Implication: Tighter bounds for convex classes	<b>12</b> 13 14 16
4	Query-Based Learning4.1General query oracles and DEC theory for query-based learning4.2Connection to the SQ dimension4.3Relation between SQ learning and LDP learning	<b>18</b> 19 20 22
5	Locally Private Learning5.1DEC theory for private PAC learning5.2Application: private regression5.3Application: private linear regression5.4DEC theory for private no-regret learning5.5Application: Contextual bandits5.5.1Linear contextual bandits5.5.2Lipschitz contextual bandits with finite arms5.5.3Concave-Lipschitz contextual bandits	<ul> <li>22</li> <li>23</li> <li>25</li> <li>26</li> <li>27</li> <li>28</li> <li>29</li> <li>30</li> <li>31</li> </ul>
6	<ul> <li>Local Minimaxity, Learnability, and Joint Privacy</li> <li>6.1 Local-minimax optimality</li></ul>	<b>31</b> 33 35 36
7	Conclusion	37
A	Additional Discussions and Results from Section 2A.1Stochastic DMSOA.2Adversarial DMSOA.3Robust DMSO	<b>45</b> 45 46 48
в	Additional Discussions and Results from Section 5         B.1 Sequential private channel	<b>50</b> 50

	B.2 B.3 B.4	Approximate DP channels       Additional examples         Additional examples       Additional examples         LDP lower bounds via SQ lower bounds       Additional examples	50 51 53
С	<b>Add</b> C.1 C.2	litional Discussions and Results from Section 6       Image:	<b>54</b> 54 54
D	Tecl	hnical Tools	56
$\mathbf{E}$	Pro	ofs for Lower Bounds	56
	E.1	Proof of Theorem 7	56
	E.2	Proof of Theorem 9	58
	E.3	Instantiations	59
		E.3.1 Query-based learning: Proof of Theorem 14	59
	<b>D</b> 4	E.3.2 LDP learning: Proof of Theorem 19 (1) and Theorem 27	60 60
	E.4	Proof of Proposition E.I.	62 62
	E.5 E.C	Proof of Lemma E.2 $\dots$	03 C4
	E.0	Proof of Theorem E.3	04
$\mathbf{F}$	Exp	loration-by-Optimization Algorithm and Guarantees	66
	F.1	Information set structure	68
	F.2	Exploration-by-Optimization algorithm $\ldots \ldots \ldots$	68
	F.3	Guarantees of the $ExO^+$ algorithm	70 70
	F.4	Proofs for upper bounds in Section 3	72
		F.4.1 Proof of Theorem 8	72
		F.4.2 Proof of Theorem 9	13
		F.4.3 Proof of Proposition 11	13
	ΓF	F.4.4 Proof of Proposition 15	74 74
	г.э	E 5.1 Model based learning	74 76
		F.5.2 Policy-based learning	77
		F 5.3 Value-based learning	78
		F 5.4 Contextual Bandits	79
	F.6	Proof of Theorem F.1	80
	-	F.6.1 Proof of Lemma F.13	81
	F.7	Proof of Theorem F.2	82
	F.8	Proof of Theorem F.3	82
	F.9	Proof of Lemma F.7	86
G	Esti	mation-to-Decision Algorithm and Guarantees	87
~	G.1	LDP-E2D Algorithm	87
		G.1.1 Online estimation oracle	88
		G.1.2 LDP-E2D Algorithm and its guarantees	88
		G.1.3 Proof of Proposition G.2.	91
	G.2	Query-based E2D algorithm	92
		G.2.1 Proof of Theorem 15	94

# H Proofs from Section 4

	H.1	Proof of Theorem 14
	H.2	Proof of Theorem 15
	H.3	Proof of Proposition 17
	H.4	Proof of Lemma 18
Ι	Ren	naining Proofs from Section 5 and Appendix B 98
	I.1	Proof of Proposition 20
	I.2	Proof of Theorem 19
		I.2.1 Proof of Proposition I.1
	I.3	Proof of Lemma 25
	I.4	Proof of Theorem 26
		I.4.1 Proof of Lemma I.5
		I.4.2 Proof of Lemma I.6
	I.5	Lower bound for LDP learning linear models
	I.6	Proof of Theorem 30
		I.6.1 Proof of Lemma I.10
		I.6.2 Proof of Lemma I.11
		I.6.3 Proof of Lemma I.12
	I.7	Proof of Proposition 31 and Proposition 32
		I.7.1 Proof of Theorem I.13
	I.8	Lower bounds for structured contextual bandits
	I.9	Proof of Proposition B.5
	I.10	Proof of Proposition B.6
	I.11	Proof of Proposition B.2
J	Pro	ofs from Section 6 and Appendix C 121
	J.1	Proof of Theorem 33
	J.2	Proof of Theorem 34
	<b>J</b> .3	Proof of Proposition C.1 and Proposition 37
	J.4	Proof of Proposition 35
	J.5	Proof of Lemma C.2 and Lemma C.3
	J.6	Proof of Proposition C.4
	J.7	Proof of Proposition 39

# A Additional Discussions and Results from Section 2

### A.1 Stochastic DMSO

In this section, we briefly review the original DMSO formulation of [Foster et al., 2021], which we call "stochastic DMSO" for clarity. In this setting, the learner (or, the decision maker) interacts for T rounds with the environment described by an underlying model  $M^*$ , unknown to the learner. On each round t = 1, ..., T:

- The learner selects a decision  $\pi_t \in \Pi$ , where  $\Pi$  is the decision space.
- The learner observes  $o_t \in \mathcal{O}$  sampled via  $o_t \sim M^*(\pi_t)$ , where  $\mathcal{O}$  is the observation space.

Formally speaking, the underlying model  $M^*$  is a conditional distribution, and the learner is given a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{O}))$  that contains  $M^*$ . To frame stochastic DMSO in our hybrid DMSO framework, we can consider the constraint  $\mathcal{P}^* = \{M^*\}$  and the constraint class  $\mathscr{P}_{sto} = \{\{M^*\}: M^* \in \mathcal{M}\}.$ 

Stochastic DMSO captures a number of decision making tasks, including reward-based learning [Foster et al., 2021, 2023b], interactive estimation and preference-based learning [Chen et al., 2022], multi-agent decision making and partial monitoring [Foster et al., 2023a].

Constrained DEC and hybrid DEC. Extending Foster et al. [2021], Foster et al. [2023b] propose the constrained PAC-DEC (regret-DEC) and derive lower and upper bounds for reward-based PAC learning (no-regret learning). Recall that constrained PAC-DEC is defined in Eq. (22) and the constrained regret-DEC is defined in Eq. (12). For stochastic DMSO, (with the constraint class being  $\mathscr{P}_{sto} = \{\{M\} : M \in \mathcal{M}\}$ ), and clearly

$$\mathsf{p}\mathsf{-dec}^{\mathsf{H}}_{\varepsilon}(\mathscr{P}_{\mathsf{sto}}) = \mathsf{p}\mathsf{-dec}^{\mathsf{c}}_{\varepsilon}(\mathcal{M}), \qquad \mathsf{r}\mathsf{-dec}^{\mathsf{H}}_{\varepsilon}(\mathscr{P}_{\mathsf{sto}}) = \mathsf{r}\mathsf{-dec}^{\mathsf{c}}_{\varepsilon}(\mathcal{M}), \qquad \forall \varepsilon \ge 0. \tag{42}$$

Therefore, the hybrid DEC can be regarded as a generalization of the constrained DECs.

#### A.2 Adversarial DMSO

In this section, we consider decision making against an adaptive adversary and instantiate the hybrid DEC theory developed in Section 2.5. For simplicity, we focus on the setting of Example 1, where  $\Phi = \{id\}$  and the value function is reward-based. In particular, our results tighten Foster et al. [2022b].

Adversarial DMSO. In the adversarial DMSO framework [Foster et al., 2022b], we consider the following protocol for T rounds. For each  $t = 1, \dots, T$ :

- The environment selects a model  $M^t \in \mathcal{M}$  (potentially depends on the interactions up to step t), and the learner selects a decision  $\pi_t \in \mathbf{\Pi}$ .
- The learner observes a noisy observation  $o_t$  via  $o_t \sim M^t(\pi_t)$ .

In the protocol above, the model  $M^t \in \mathcal{M}$  at step t can adaptively selected, i.e., it may depend on the history  $\mathcal{H}_{t-1}$  prior to step t. The regret of the learner is measured against the best decision in hindsight:

$$\operatorname{\mathbf{Reg}}_{\mathsf{DM}}(T) := \max_{\pi^{\star} \in \Pi} \sum_{t=1}^{T} V^{M^{t}}(\pi^{\star}) - \mathbb{E}_{\pi_{t} \sim q_{t}} V^{M^{t}}(\pi_{t}),$$
(43)

where the expectation of  $\pi_t \sim q_t$  is taken over the randomness of the learner at step t, and  $V^M(\pi) = \mathbb{E}^{M,\pi} R(o,\pi)$  is specified by a known reward function  $R : \mathcal{O} \times \Pi \to [0,1]$ .

It is clear that adversarial DMSO can be framed within hybrid DMSO framework with the constraint class  $\mathscr{P}_{adv} = \{\mathcal{M}\}$ , i.e., the constraint is always  $\mathcal{P}^* = \mathcal{M}$ . Therefore, we can directly apply Theorem 10, as follows.

**Theorem A.1** (No-regret learning against an adversary). Let  $T \ge 1, \delta \in (0, 1)$ , model class  $\mathcal{M}$ , and a reward function  $R \in [0, 1]$  be given. Suppose that  $\mathcal{M}$  is compact (Assumption 2), and the regret DEC r-dec<sup>c</sup><sub> $\varepsilon$ </sub>(co( $\mathcal{M}$ )) is of moderate decay as a function of  $\varepsilon$ . Then, ExO<sup>+</sup> (instantiated on  $\mathscr{P}_{adv}$ , following Theorem 10) achieves with probability at least  $1 - \delta$  that

$$\frac{1}{T} \operatorname{Reg}_{\mathsf{DM}}(T) \leq \Delta + O(\sqrt{\log T}) \cdot \Big[ \operatorname{\mathsf{r-dec}}_{\bar{\varepsilon}(T)}^{\mathsf{c}}(\operatorname{co}(\mathcal{M})) + \bar{\varepsilon}(T) \Big],$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log N_{\text{frac}}(\operatorname{co}(\mathcal{M}), \Delta) + \log(1/\delta)}{T}}.$ 

The above upper bound scales with the regret DEC of  $co(\mathcal{M})$  and the fractional covering number of  $co(\mathcal{M})$ , which is tighter than Foster et al. [2022b]: the latter involves a log  $|\Pi|$  factor, whereas it always holds that log  $N_{\text{frac}}(co(\mathcal{M}), \Delta) \leq \log |\Pi|$ .

Lower bounds. A direct instantiation of Theorem 9 recovers the lower bound of Foster et al. [2022b].

**Proposition A.2** (Regret lower bound with stationary adversary). Let  $T \ge 1$ ,  $\mathcal{M}$  be a given model class. Then, for any T-round algorithm Alg,

$$\sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \ge \frac{T}{8} \Big( \mathsf{r} - \mathsf{dec}_{\underline{\varepsilon}(T)}^{\mathsf{c}}(\mathsf{co}(\mathcal{M})) - 8\underline{\varepsilon}(T) \Big) - 1, \tag{44}$$

where the supremum is taken over stationary environments  $\mathsf{Env}$  specified by a distribution  $\mu \in \Delta(\mathcal{M})$ , and  $\underline{\varepsilon}(T) = \frac{1}{24\sqrt{T}}$ .

In addition to the regret DEC lower bound, we can show that fractional covering number of  $co(\mathcal{M})$  also provides a lower bound. Proposition A.3 below is a direct corollary of the fractional covering number lower bound of Chen et al. [2024] (see also Appendix J.2). Thus, we omit its proof for succinctness.

**Proposition A.3** (Fractional covering number lower bound). Let  $T \ge 1, \Delta \ge 0$ . Suppose that  $\mathcal{M}$  is a given model class, and Alg is a T-round algorithm that achieves  $\mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\operatorname{Reg}_{\mathsf{DM}}(T)] \le T\Delta$  for any stationary environment  $\mathsf{Env}$  specified by a distribution  $\mu \in \Delta(\mathcal{M})$ . Then it holds that

$$T \ge \frac{\log N_{\mathsf{frac}}(\operatorname{co}(\mathcal{M}), \Delta/2) - 2}{2C_{\mathrm{KL}}(\operatorname{co}(\mathcal{M}))}$$

A nearly "complete" characterization of the minimax regret. For no-regret learning in hybrid DMSO, the minimax regret is defined as

$$\mathfrak{M}_{T}(\mathscr{P}) := \inf_{\mathsf{Alg Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)],$$

where the supremum is taken over all environments constrained by  $\mathscr{P}$ . We also consider the following notion of minimax regret and sample complexity:

$$\mathfrak{C}_{\Delta}(\mathscr{P}) := \min \left\{ T : \mathfrak{M}_{T}(\mathscr{P}) \leq T\Delta \right\},$$

i.e.,  $\mathfrak{C}_{\Lambda}(\mathscr{P})$  is the minimum of T such that an T-round algorithm may achieve  $T\Delta$ -regret.

Under the above notation, we can translate the lower and upper bounds in this section into the following characterization of  $\mathfrak{C}_{\Delta}(\mathscr{P}_{\mathsf{adv}})$  (with  $\mathscr{P}_{\mathsf{adv}} = \{\mathcal{M}\}$ ):

$$\max\left\{\mathfrak{C}_{\Delta}^{\mathsf{dec}}(\mathrm{co}(\mathcal{M})), \frac{\log N_{\mathsf{frac}}(\mathrm{co}(\mathcal{M}), 2\Delta)}{C_{\mathrm{KL}}(\mathrm{co}(\mathcal{M}))}\right\} \lesssim \mathfrak{C}_{\Delta}(\mathscr{P}_{\mathsf{adv}}) \lesssim \mathfrak{C}_{\Delta}^{\mathsf{dec}}(\mathrm{co}(\mathcal{M})) \cdot \log N_{\mathsf{frac}}(\mathrm{co}(\mathcal{M}), \Delta/\Delta),$$

$$\tag{45}$$

where  $\mathfrak{C}_{\Delta}^{\mathsf{dec}}(\mathrm{co}(\mathcal{M})) := \min \{\varepsilon^{-2} : \mathsf{r}\operatorname{-}\mathsf{dec}_{\varepsilon}^{c}(\mathrm{co}(\mathcal{M})) \leq \Delta\}$ , and we omit logarithmic factors and assume suitable growth conditions on the regret DEC of  $\mathrm{co}(\mathcal{M})$ . Note that the lower and upper bounds of Eq. (45) match up to squaring and a factor of  $C_{\mathrm{KL}}(\mathrm{co}(\mathcal{M}))$ . In particular, for a model class  $\mathcal{M}$  with  $C_{\mathrm{KL}}(\mathrm{co}(\mathcal{M})) = O(1)$ , the DEC and fractional covering number together characterize the minimax sample complexity  $\mathfrak{C}_{\Delta}(\mathscr{P}_{\mathsf{adv}})$  (polynomially).

### A.3 Robust DMSO

In this section, we discuss the relationship between our formulation of robust DMSO and other contamination models, and present the PAC and no-regret guarantees for robust decision making.

Recall that in robust DMSO (Section 2.4), the constraint set is

$$\mathcal{P}_{M^{\star}} := \left\{ (1-\beta)M^{\star} + \beta M' : M' \in (\mathbf{\Pi} \to \Delta(\mathcal{O})) \right\},\$$

and the constraint class (induced by  $\mathcal{M}$ ) as given by  $\mathscr{P}_{\beta\text{-Huber}} := \{\mathcal{P}_{M^*} : M^* \in \mathcal{M}\}$ . To ease the notational burden, we define

$$\mathcal{M}_{\beta-\mathsf{Huber}} := \mathcal{M}_{\mathscr{P}_{\beta-\mathsf{Huber}}} = \{(1-\beta)M^{\star} + \beta M' : M^{\star} \in \mathcal{M}, M' \in (\Pi \to \Delta(\mathcal{O}))\},\$$

consisting of all stationary environments that are  $\beta$ -contaminated from a ground-truth model  $M^* \in \mathcal{M}$ .

Contamination models in robust statistics. In Huber contamination model [Huber, 1965, Huber and Ronchetti, 2011], the environment is stationary and specified by  $(1 - \beta)M^{\star} + \beta M'$ . where  $M^* \in \mathcal{M}$  is the "true model", and M' is an arbitrary contamination model. Clearly, Huber's contamination model is encompassed by stochastic DMSO (with model class  $\mathcal{M}_{\beta-\text{Huber}}$ ). Recently, for statistical estimation, the *adaptive* and *oblivious* contamination models were studied by Diakonikolas et al., 2019, Diakonikolas and Kane, 2019, Liu and Moitra, 2021, Diakonikolas and Kane, 2023, Canonne et al., 2023], among others. In these contamination models, after the i.i.d. samples  $z_1, \dots, z_T \sim M^*$  is generated, the adversary may arbitrarily corrupt  $\beta T$  many samples. The adversary is adaptive if it can choose the  $\beta T$  corrupted samples based on the whole sequence. Otherwise, the adversary is called *oblivious*. For statistical tasks, the adaptive adversary (in the above sense) can be stronger than the constrained environment in hybrid DMSO, as it is allowed to inspect the whole sequence of samples before contaminating it. On the other hand, the oblivious adversary can be much weaker. Finally, we note that both definitions of the adaptive and oblivious adversary are specialized to the statistical estimation (where the samples  $z_1, \dots, z_T$  are i.i.d). For general interactive decision making tasks, we believe the robust DMSO is a natural choice of contamination model.

**PAC lower and upper bounds.** To apply the results of hybrid DMSO, we only need to show that  $p-dec_{\varepsilon}^{\mathsf{H}}(\mathscr{P}_{\beta-\mathsf{Huber}}) = p-dec_{\varepsilon}^{\mathsf{R}}(\mathcal{M})$ . By definition, for any  $M^{\star} \in \mathcal{M}$ , reference model  $\overline{M}$ , we have

$$\inf_{M \in \operatorname{co}(\mathcal{P}_{M^{\star}})} \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^{2} \left( M(\pi), \overline{M}(\pi) \right) = \inf_{M': \mathbf{\Pi} \to \Delta(\mathcal{O})} \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^{2} \left( (1-\beta) M^{\star}(\pi) + \beta M'(\pi), \overline{M}(\pi) \right)$$
$$= \mathbb{E}_{\pi \sim q} \inf_{P' \in \Delta(\mathcal{O})} D_{\mathrm{H}}^{2} \left( (1-\beta) M^{\star}(\pi) + \beta P', \overline{M}(\pi) \right)$$
$$= \mathbb{E}_{\pi \sim q} D_{\beta-\operatorname{Huber}}^{2} \left( M^{\star}(\pi), \overline{M}(\pi) \right).$$

Therefore, for any reference model  $\overline{M}$  and  $\varepsilon \in [0, 1]$ , it holds that

$$\mathsf{p-dec}_{\varepsilon}^{\mathsf{H}}(\mathscr{P}_{\beta-\mathsf{Huber}},\overline{M}) = \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{\mathcal{P}_{M^{\star}} \in \mathscr{P}_{\beta-\mathsf{Huber}}} \left\{ \mathbb{E}_{\pi \sim p} L(\mathcal{P}_{M^{\star}},\pi) \mid \inf_{\substack{M \in \mathrm{co}(\mathcal{P}_{M^{\star}})}} \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^{2}\left(M(\pi),\overline{M}(\pi)\right) \le \varepsilon^{2} \right\}$$
$$= \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{M^{\star} \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p} L(M^{\star},\pi) \mid \mathbb{E}_{\pi \sim q} D_{\beta-\mathsf{Huber}}^{2}\left(M^{\star}(\pi),\overline{M}(\pi)\right) \le \varepsilon^{2} \right\}$$

$$= \mathsf{p-dec}_{\varepsilon}^{\mathsf{R}}(\mathcal{M}, \overline{M}).$$

Therefore, we have proven  $p-dec_{\varepsilon}^{\mathsf{H}}(\mathscr{P}_{\beta-\mathsf{Huber}}) = p-dec_{\varepsilon}^{\mathsf{R}}(\mathcal{M})$  for  $\varepsilon \in [0,1]$ . By instantiating Theorem 7 and Theorem 8, we have the following bounds.

**Theorem A.4** (PAC bounds for robust decision making). Let  $T \ge 1, \beta \in [0, 1]$ , model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \Delta(\mathcal{O}))$  be given, and the loss function L is metric-based.

(1) Lower bound: Let Alg be a T-round algorithm. Then there exists  $M^* \in \mathcal{M}$  and a stationary environment Env that is specified by  $M = (1 - \beta)M^* + \beta M'$ , such that the expected risk of Alg under Env is lower bounded as

$$\mathbb{E}^{\operatorname{Env},\operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\operatorname{DM}}(T)] \geq \frac{1}{8} \operatorname{p-dec}_{\underline{\varepsilon}(T)}^{\operatorname{R}}(\mathcal{M}),$$

where  $\underline{\varepsilon}(T) = \frac{1}{2\sqrt{T}}$ .

(2) Upper bound: Suppose the robust DEC  $p-dec_{\varepsilon}^{\mathsf{R}}(\mathcal{M})$  is of moderate decay. Then  $\mathsf{ExO}^+$  (instantiated on  $\mathscr{P}_{\beta-\mathsf{Huber}}$ , following Theorem 8) achieves, in any  $\beta$ -contaminated environment, that with probability at least  $1-\delta$ 

$$\mathbf{Risk}_{\mathsf{DM}}(T) \lesssim \mathsf{p-dec}_{\overline{\varepsilon}(T)}^{\mathsf{R}}(\mathcal{M}),$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log(|\mathcal{M}|/\delta)}{T}}$ .

**Regret lower and upper bounds.** In robust DMSO, we may also consider the no-regret learning goal (specified by Eq. (11)). For simplicity, we present the regret bounds in the setting of Example 1, i.e., the measurement class  $\Phi = \{id\}$  consists of the identity measurement, and the value function is reward-based. Then, by instantiating Theorem 9 and Theorem 10, we have the following bounds in terms of the regret DEC of  $\mathcal{M}_{\beta-Huber} = \mathcal{M}_{\mathscr{P}_{\beta-Huber}}$ .

**Theorem A.5** (Regret bounds for robust decision making). Let  $T \ge 1, \beta \in [0, 1], \Phi = \{id\}, model class <math>\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{O}))$ , and the value function V is reward-based (Example 1).

(1) Lower bound: Let Alg be a T-round algorithm. Then there exists  $M^* \in \mathcal{M}$  and a stationary environment Env that is specified by  $M = (1 - \beta)M^* + \beta M' \in \mathcal{M}_{\beta-\text{Huber}}$ , such that the expected regret of Alg under Env is lower bounded as

$$\mathbb{E}^{\operatorname{Env},\operatorname{Alg}}[\operatorname{\mathbf{Reg}}_{\operatorname{\mathsf{DM}}}(T)] \geq \frac{T}{8} \Big( \operatorname{\mathsf{r-dec}}_{\underline{\varepsilon}(T)}^{\operatorname{c}}(\mathcal{M}_{\beta\operatorname{\mathsf{-Huber}}}) - 8\underline{\varepsilon}(T) \Big) - 1,$$

where  $\underline{\varepsilon}(T) = \frac{1}{10\sqrt{T}}$ .

(2) Upper bound: Suppose that  $\mathcal{M}_{\beta-\text{Huber}}$  is compact, and the robust DEC r-dec<sup>c</sup><sub> $\varepsilon$ </sub>( $\mathcal{M}_{\beta-\text{Huber}}$ ) is of moderate decay. Then ExO<sup>+</sup> (instantiated on  $\mathscr{P}_{\beta-\text{Huber}}$ , following Theorem 10) achieves, in any  $\beta$ -contaminated environment, that with probability at least  $1 - \delta$ 

$$\frac{1}{T} \operatorname{Reg}_{\mathsf{DM}}(T) \leq \Delta + O(\sqrt{\log T}) \cdot \Big[ \operatorname{\mathsf{r-dec}}_{\bar{\varepsilon}(T)}^{c} (\mathcal{M}_{\beta-\mathsf{Huber}}) + \bar{\varepsilon}(T) \Big],$$
  
where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log(|\mathcal{M}|/\delta) + \log N_{\mathsf{frac}}(\mathcal{M}_{\beta-\mathsf{Huber}}, \Delta)}{T}}.$ 

### **B** Additional Discussions and Results from Section 5

### **B.1** Sequential private channel

The work of Duchi et al. [2013, 2018] formalizes the problem of sequential private channel selection for statistical tasks (cf. Definition 10). We rephrase its definition as follows.

**Definition 13.** A sequential channel Q from the data space  $\mathcal{Z}$  to the privatized data space  $\mathcal{O}$  is specified by a class of conditional distributions

$$\{Q(o_t = \cdot | z_t = \cdot, o_1 = \cdot, \cdots, o_{t-1} = \cdot)\}_{t \in [T]}$$

A sequential channel Q is  $\alpha$ -private if for any  $t \in [T]$ , any  $\overline{z}_t, \overline{z}_t \in \mathbb{Z}$ , any  $\overline{o}_1, \dots, \overline{o}_{t-1} \in \mathcal{O}$ , we have

$$\frac{Q(o_t \in E | z_t = \bar{z}_t, o_1 = \bar{o}_1, \cdots, o_{t-1} = \bar{o}_{t-1})}{Q(o_t \in E | z_t = \bar{z}_t, o_1 = \bar{o}_1, \cdots, o_{t-1} = \bar{o}_{t-1})} \le e^{\alpha}, \qquad \forall \text{ measurable } E \subseteq \mathcal{O}.$$

Clearly, in statistical tasks, any  $\alpha$ -private sequential channel Q induces an  $\alpha$ -LDP algorithm, which at each step  $t \in [T]$  selects the  $\alpha$ -LDP channel  $Q^t$  given by

$$Q^t(o|z) = Q(o_t = o|z_t = z, o_1, \cdots, o_{t-1}),$$

based on the history  $\mathcal{H}^{(t-1)} = (o_1, \cdots, o_{t-1})$ . Conversely, an  $\alpha$ -LDP algorithm also induces a sequential  $\alpha$ -private channel. Therefore, sequential  $\alpha$ -private channels are equivalent to the  $\alpha$ -LDP algorithms in private DMSO.

A similar argument also shows that for interactive decision making, our formulation in Appendix A.1 recovers the commonly studied interactive private channels (see e.g., Zheng et al. [2020], Garcelon et al. [2021]).

#### **B.2** Approximate DP channels

We first recall the definition of approximate DP channels.

**Definition 14** (Approximate DP channels). A channel Q (from latent observation space  $\mathcal{Z}$  to observation space  $\mathcal{O}$ ) is  $(\alpha, \beta)$ -DP if for  $z, z' \in \mathcal{Z}$  and any measurable set  $E \subseteq \mathcal{O}$ ,

$$\mathsf{Q}(E|z) \le e^{\alpha} \mathsf{Q}(E|z') + \beta.$$

The equivalence between approximate DP and pure DP under local privacy model is known [Duchi and Rogers, 2019, Duchi and Ruan, 2024]. In this section, we formalize such an equivalence in the general context of interactive decision making.

In the following, we assume  $\mathcal{O}$  is countable. The following lemma from Duchi and Rogers [2019, Lemma 25] shows that any  $(\alpha, \beta)$ -LDP channel is close to an  $\alpha$ -LDP channel.

**Lemma B.1.** For any  $(\alpha, \beta)$ -LDP channel Q, there exists an  $\alpha$ -LDP channel Q<sub>pure</sub> such that

$$\sup_{z \in \mathcal{Z}} D_{\mathrm{TV}}\left(\mathsf{Q}(\cdot|z), \mathsf{Q}_{\mathsf{pure}}(\cdot|z)\right) \leq \frac{\beta}{1 + e^{\alpha} - \beta}.$$

As a corollary, we can show that any algorithm that preserves  $(\alpha, \beta)$ -LDP is close to an algorithm that preserves  $\alpha$ -LDP. Proof is presented in Appendix I.11.

**Proposition B.2.** Suppose that  $\beta \leq \frac{1}{2}$  and Alg is a *T*-round algorithm that preserves  $(\alpha, \beta)$ -LDP. Then there is a *T*-round algorithm Alg<sub>pure</sub> that preserves  $\alpha$ -LDP, such that for any model *M*,

$$D_{\mathrm{TV}}\left(\mathbb{P}^{M,\mathsf{Alg}}(\mathcal{H}_{\pi}=\cdot),\mathbb{P}^{M,\mathsf{Alg}_{\mathsf{pure}}}(\mathcal{H}_{\pi}=\cdot)\right) \leq 2T\beta,$$

where the TV distance is taken between the distribution of the trajectory of the decisions  $\mathcal{H}_{\pi} = (\pi_1, \cdots, \pi_T, \pi_{T+1})$ . In particular, when the loss function is bounded in [0,1], it holds that for any model M,

$$\mathbb{E}^{M,\operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] \geq \mathbb{E}^{M,\operatorname{Alg}_{\mathsf{pure}}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] - 2T\beta.$$

Hence, as long as  $\beta = \frac{1}{\text{poly}(T)}$ , there is *essentially no gain* of allowing the algorithms to be  $(\alpha, \beta)$ -LDP.

#### **B.3** Additional examples

Recall that in Theorem 22, we show that LDP-ExO provides an upper bound scaling with the private PAC-DEC of  $co(\mathcal{M})$  and the fractional covering number of  $\mathcal{M}$ . To draw a clearer comparison between this upper bound and the lower bounds, we re-state our lower and upper bounds in terms of the minimax sample complexity (40). Define

$$\mathfrak{C}_{\Delta}^{\operatorname{dec}}(\mathcal{M}) := \min\{\varepsilon^{-2} : \operatorname{p-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M}) \leq \Delta\}.$$

Then, under the assumption of Theorem 22, we have the following characterization of  $\mathfrak{C}_{\Lambda}(\mathcal{M})$ :

$$\max\{\mathfrak{C}_{\Delta}^{\mathsf{dec}}(\mathcal{M}), \log N_{\mathsf{frac}}(\mathcal{M}, 2\Delta)\} \lesssim \alpha^2 \cdot \mathfrak{C}_{\Delta}(\mathcal{M}) \lesssim \mathfrak{C}_{\Delta}^{\mathsf{dec}}(\operatorname{co}(\mathcal{M})) \cdot \log N_{\mathsf{frac}}(\mathcal{M}, \Delta/2).$$
(46)

In particular, for a convex model class  $\mathcal{M}$ , under mild assumption on the growth of the private PAC-DEC and fractional covering number, the lower and upper bounds match up to squaring. We note that Eq. (46) is analogous to the observations of Chen et al. [2024] for non-private learning.

In the following, we discuss similar characterizations for convex hypothesis selection and online regression.

**Convex hypothesis selection.** As an application of Proposition 11, we consider the LDP hypothesis selection problem, which is a statistical task (Definition 10).

**Example 8.** Given a model class  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ , a hypothesis selection problem is described by a partition

$$\mathcal{M} = \bigsqcup_{i=1}^{m} \mathcal{M}^{(i)},$$

where  $\mathcal{M}^{(1)}, \dots, \mathcal{M}^{(m)}$  are disjoint subclasses. The decision space is  $\Pi = [m]$ , and for each  $M \in \mathcal{M}$ ,  $\pi \in \Pi$ , the loss function is given by  $L(M, \pi) = \mathbf{1} \{ \pi \neq \pi^M \}$ , where  $\pi^M$  is the unique index  $i \in [m]$  such that  $M \in \mathcal{M}^{(i)}$ .

Note that the LDP hypothesis selection problem can be regarded as a special case of Example 2 (with the measurement class  $\Phi = Q_{\alpha}$  the class of all  $\alpha$ -DP channels). Therefore, we summarize the lower and upper bounds for this problem, as follows.

**Proposition B.3** (Private hypothesis selection). Let  $T \ge 1$ ,  $\delta \in (0, 1)$ .

(1) Lower bound: For any  $\alpha$ -LDP algorithm Alg, it holds that

$$\sup_{M^{\star}\in\mathcal{M}}\mathbb{P}^{M^{\star},\operatorname{Alg}}\left(\pi_{T+1}\neq\pi^{M^{\star}}\right)\geq\frac{1}{8}\mathrm{p}\operatorname{-dec}_{\underline{\hat{\varepsilon}}(T)}^{\operatorname{LDP}}(\mathcal{M}),$$

where  $\underline{\varepsilon}(T) = \frac{c}{\sqrt{\alpha^2 T}}$ .

(2) Upper bound: Suppose that  $\mathcal{M}$  is compact,  $\mathcal{M}^{(1)}, \cdots, \mathcal{M}^{(m)}$  are convex, and

$$\mathsf{p-dec}^{\mathsf{LDP}}_{\bar{\varepsilon}(T)}(\mathcal{M}) \leq \frac{1}{3}, \qquad \bar{\varepsilon}(T) = C \sqrt{\frac{\log(m/\delta)}{\alpha^2 T}}$$

Then LDP-ExO can be suitably instantiated to preserve  $\alpha$ -LDP, so that under any model  $M^* \in \mathcal{M}_{i^*}$ , the algorithm returns  $\pi_{T+1} = i^*$  with probability at least  $1 - \delta$ .

In terms of the sample complexity, assuming that  $\mathcal{M}^{(1)}, \cdots, \mathcal{M}^{(m)}$  are convex, we have

$$\mathfrak{C}^{\mathsf{dec}}_{1/3}(\mathcal{M}) \lesssim \alpha^2 \cdot \mathfrak{C}_{\Delta}(\mathcal{M}) \lesssim \mathfrak{C}^{\mathsf{dec}}_{1/3}(\mathcal{M}) \cdot \log(m/\Delta),$$

for all  $\Delta \in [0, 0.05]$ . Therefore, up to the factor of  $\log(m/\Delta)$ , the sample complexity of private convex hypothesis selection is completely characterized by the private PAC-DEC.

**Online regression.** We consider the online variant of the regression task (Section 5.2). In the setting of *online regression*, for every step  $t \in [T]$ , the environment selects a pair  $(x_t, y_t)$  (potentially depends on the history prior to step t), and the learner has to pick a (randomized) prediction function  $f_t \in \mathcal{F}$ . The regret of the learner is measured by

$$\mathbf{Reg}_{\mathsf{DM}}(T) := \sum_{t=1}^{T} \mathcal{L}(y_t, f_t(x_t)) - \inf_{f^{\star} \in \mathcal{F}} \sum_{t=1}^{T} \mathcal{L}(y_t, f^{\star}(x_t)),$$

where  $L: [-1,1]^2 \rightarrow [0,1]$  is a given loss.

Clearly, online regression is encompassed by adversarial DMSO (Appendix A.2), with the constraint being  $\mathcal{P}^{\star} = \{\mathcal{M}_{agnostic}\}$ . As a corollary of Theorem A.1, we have the following regret bound for online regression.<sup>9</sup>

**Proposition B.4.** Let  $T \ge 1$ ,  $\delta \in (0,1)$ . Suppose that  $\mathcal{X}$  is finite, and  $p-dec_{\varepsilon}^{LDP}(\mathcal{M}_{agnostic})$  is of moderate decay as a function of  $\varepsilon$ . For online regression, LDP-ExO achieves the following regret bound with probability at least  $1 - \delta$ :

$$\frac{1}{T} \operatorname{\mathbf{Reg}_{\mathsf{DM}}}(T) \leq \Delta + O(\sqrt{\log T}) \cdot \operatorname{\mathbf{p-dec}}_{\bar{\varepsilon}(T)}^{\operatorname{LDP}}(\mathcal{M}_{\operatorname{agnostic}})$$
where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log N_{\operatorname{frac}}(\mathcal{M}_{\operatorname{agnostic}}, \Delta) + \log(1/\delta)}{\alpha^2 T}}.$ 

This also recovers the risk bound of Proposition 23 when the data are drawn i.i.d from a  $M^* \in \mathcal{M}_{agnostic}$ . Therefore, in this sense, online private regression is *no* more difficult than the agnostic private regression (with potential degradation of the rate of the regret), because the private PAC-DEC p-dec\_{\varepsilon}^{\text{LDP}}(\mathcal{M}\_{agnostic}) and the fractional covering number  $N_{\text{frac}}(\mathcal{M}_{agnostic}, \Delta)$  also provide lower bounds (similar to Eq. (46)). This is in sharp contrast to the non-private setting, where there is a separation between the complexity of regression and online regression.

<sup>&</sup>lt;sup>9</sup>For regression (a statistical task), we have  $\mathsf{r-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}_{\mathsf{agnostic}}) = \mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}_{\mathsf{agnostic}})$  because the decision  $f \in \mathcal{F}$  does not affect the distribution of the observation.

### B.4 LDP lower bounds via SQ lower bounds

For a more general demonstration of the power of private PAC-DEC, we consider the following variant of the commonly used SQ lower bound methods [Blum et al., 1994, Feldman et al., 2017, Brennan et al., 2020, etc.]. We focus on the statistical tasks (Definition 10, where  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ ).

**Definition 15** (Minimum correlation). For distributions  $\mathcal{D}_1, \mathcal{D}_2, \mathcal{D} \in \Delta(\mathcal{Z})$ , we define the pairwise correlation as

$$\rho_{\mathcal{D}}(\mathcal{D}_1, \mathcal{D}_2) = \mathbb{E}_{z \sim \mathcal{D}}\left(\frac{d\mathcal{D}_1(z)}{d\mathcal{D}(z)} - 1\right) \left(\frac{d\mathcal{D}_2(z)}{d\mathcal{D}(z)} - 1\right).$$

We say a set of m distributions  $\{\mathcal{D}_1, \cdots, \mathcal{D}_m\}$  is  $\varepsilon$ -correlated relative to  $\mathcal{D}$  if

$$\forall i, j, \qquad |\rho_{\mathcal{D}}(\mathcal{D}_i, \mathcal{D}_j)| \le \begin{cases} \varepsilon^2, & i \neq j, \\ m\varepsilon^2, & i = j. \end{cases}$$

Suppose  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ . For any  $\Delta$ , we define the minimum correlation  $\operatorname{cor}(\mathcal{M}, \Delta)$  to be the minimum of  $\varepsilon$  such that there exists a reference model  $\overline{M}$  and a set of models  $\{M_1, \dots, M_m\} \subseteq \mathcal{M}$ , such that (1)  $\{M_1, \dots, M_m\}$  is  $\varepsilon$ -correlated relative to  $\overline{M}$ ; (2) for any  $\pi \in \Pi$ , there is at most m/2 indices  $i \in [m]$  such that  $L(M_i, \pi) \leq \Delta$ .

In the following, we show that  $cor(\mathcal{M}, \Delta)$  provides a lower bound of private PAC-DEC of  $\mathcal{M}$ , and hence it also provides a lower bound for learning  $\mathcal{M}$  under LDP.

**Proposition B.5.** For any  $\Delta > 0$ , it holds that

$$\mathrm{p\text{-}dec}_{\varepsilon}^{\mathrm{LDP}}(\mathcal{M}) \geq \frac{\Delta}{4}, \qquad \forall \varepsilon \leq \mathrm{cor}(\mathcal{M}, \Delta).$$

In terms of the sample complexity, any  $\alpha$ -LDP algorithm requires  $\Omega\left(\frac{1}{\alpha^2 \cdot \operatorname{cor}(\mathcal{M}, 4\Delta)^2}\right)$  samples to learn a  $\Delta$ -optimal decision in  $\mathcal{M}$ .

Proof can be found in Appendix I.9.

Hardness of LDP learning parity. It has been shown that learning parity under LDP is hard [Kasiviswanathan et al., 2011], in the sense that there is a  $2^{\Omega(d)}$  lower bound on the sample complexity (where d is the dimension). In the following, we apply Proposition B.5 to recover the exponential lower bound and discuss its implication. Proof in Appendix I.10.

**Proposition B.6** (Learning parity). Let  $d \ge 2$ ,  $\varepsilon \in [0,1]$ , and  $\mathcal{X} = \{0,1\}^d$ , and  $\mathcal{F}_{\mathsf{parity}} = \{f_S\}_{S \subseteq [d]}$ , where for each subset  $S \subseteq [d]$ , the function  $f_S : \mathcal{X} \to \{-1,1\}$  is defined as

$$f_S(x) = (-1)^{\sum_{i \in S} x_i}, \quad \forall x \in \{0, 1\}^d.$$

Then, there exists a distribution  $\mu \in \Delta(\mathcal{X})$ , such that for  $\mathcal{M}_{\text{parity}}$  the class of all realizable models with the covariate distribution  $\mu$ , it holds that

$$\mathsf{p}\text{-}\mathsf{dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}_{\mathsf{parity}}) \geq \Omega(\sqrt{2^d}\varepsilon)$$

This implies a lower bound of  $\sup_{M \in \mathcal{M}_{\text{parity}}} \mathbb{E}^{M,\text{Alg}}[\operatorname{\mathbf{Risk}_{DM}}(T)] \geq \Omega\left(\sqrt{\frac{2^d}{T}}\right)$  for any T-round algorithm Alg.

Notice that for the parity function class, we have  $|\mathcal{F}_{parity}| = 2^d$ , and hence the lower bound above is in sharp contrast to the non-private setting, where the ERM can achieve a risk bound of  $\sqrt{\frac{\log |\mathcal{F}_{parity}|}{T}}$ .

# C Additional Discussions and Results from Section 6

### C.1 Joint DP in interactive learning

Generalizing the notion of JDP for non-interactive learning, Shariff and Sheffet [2018] propose a definition of JDP for contextual bandits, which is later extended to reinforcement learning by Vietri et al. [2020]. In the following, we formalize the notion of JDP for general interactive decision problems.

Recall that a *T*-round algorithm Alg (without LDP constraints) is specified by a sequence of mappings  $\{q_t\}_{t\in[T]} \cup \{\widehat{p}\}\)$ , where the *t*-th mapping  $q_t(\cdot \mid \mathcal{H}_{t-1})$  specifies the distribution of  $\pi_t$  based on the history  $\mathcal{H}_{t-1} = (\pi_s, z_s)_{s \leq t-1}$ , and the final map  $\widehat{p}(\cdot \mid \mathcal{H}_T)$  specifies the distribution of the *output* policy  $\pi_{T+1}$  based on  $\mathcal{H}_T$ .

**Definition 16** (Interactive JDP). For sequences of observations  $\mathcal{H}_{z,T} = (z_1, \dots, z_T)$  and  $\mathcal{H}'_{z,T} = (z'_1, \dots, z'_T)$ , we say  $\mathcal{H}_{z,T}$  and  $\mathcal{H}'_{z,T}$  are neighbored if there is at most one index  $t \in [T]$  such that  $z_t \neq z'_t$ .

The algorithm Alg preserves  $\alpha$ -JDP if for any two neighbored sets of observations  $\mathcal{H}_{z,T} = (z_1, \cdots, z_T)$ and  $\mathcal{H}'_{z,T} = (z'_1, \cdots, z'_T)$ , it holds that

$$\mathbb{P}^{\mathsf{Alg}}((\pi_1,\cdots,\pi_T,\pi_{T+1})\in E|\mathcal{H}_{z,T})\leq e^{\alpha}\mathbb{P}^{\mathsf{Alg}}((\pi_1,\cdots,\pi_T,\pi_{T+1})\in E|\mathcal{H}'_{z,T}),$$

for any measurable set  $E \subseteq \Pi$ , where  $\mathbb{P}^{Alg}$  is taken over the randomness of the algorithm, i.e.,

$$\mathbb{P}^{\mathsf{Alg}}((\pi_1,\cdots,\pi_T,\pi_{T+1})=\cdot|z_1,\cdots,z_T)=\prod_{t=1}^{T+1}q_t(\pi_t=\cdot|\pi_{1:t-1},z_{1:t-1}),$$

where we regard  $q_{T+1} := \hat{p}$ .

For statistical estimation problems, the definition above clearly recovers Definition 11. It also recovers the definition of interactive JDP considered by Shariff and Sheffet [2018], Vietri et al. [2020], He et al. [2022].

Similar to Proposition 37, we show that fractional covering number provides a lower bound for interactive learning under JDP.

**Proposition C.1** (Fractional covering number lower bound for JDP learning). Let  $T \ge 1$ , and Alg is a weak  $\alpha$ -JDP algorithm. Suppose that with T-round of interactions, Alg achieves  $\operatorname{Risk}_{DM}(T) \le \Delta$  with probability at least  $\frac{1}{2}$  under  $\mathbb{P}^{M,\operatorname{Alg}}$  for any  $M \in \mathcal{M}$ . Then it holds that

$$T \ge \frac{\log N_{\mathsf{frac}}(\mathcal{M}, \Delta) - \log 2}{\alpha}.$$

Note that any  $\alpha$ -LDP algorithm preserves  $\alpha$ -JDP. Hence, Theorem 38 naturally extends to interactive learning.

### C.2 Learnability of regression

In this section, we consider the learnability of the regression task, continuing Section 6.2. Recall that in Section 5.2, we study *proper* regression. More generally, in this section, we also consider the problem of *improper* regression with a function class  $\mathcal{F}^+$  not necessarily equal to  $\mathcal{F}$ .

In improper regression, the decision space is  $\Pi = \mathcal{F}^+$ , and the loss function is defined as

$$L(M, f) = \mathbb{E}_{(x,y) \sim M} \mathcal{L}(y, f(x)) - \min_{f^{\star} \in \mathcal{F}} \mathbb{E}_{(x,y) \sim M} \mathcal{L}(y, f^{\star}(x)), \qquad \forall f \in \mathcal{F}^+.$$

Define the fractional covering number of the pair  $(\mathcal{F},\mathcal{F}^+)$  as

$$N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta) := \inf_{p \in \Delta(\mathcal{F}^+)} \sup_{\mu \in \Delta(\mathcal{X}), f^\star \in \mathcal{F}} \frac{1}{p(f : \mathbb{E}_{x \sim \mu} |f(x) - f^\star(x)| \le \Delta)}.$$
(47)

When  $\mathcal{F}^+ = \mathcal{F}$ , this definition recovers the definition (31) of the fractional covering number of  $\mathcal{F}$ . We first relate  $N_{\text{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$  to the fractional covering number of  $\mathcal{M}_{\text{agnostic}}$  under the absolute loss  $L_{\text{abs}}(y, y') = |y - y'|$ .

**Lemma C.2.** Recall that  $\mathcal{M}_{\text{agnostic}} = \Delta(\mathcal{Z})$  is the class of all agnostic models. Then, under the absolute loss  $L_{\text{abs}}$  and decision space  $\Pi = \mathcal{F}^+$ , it holds that

$$N_{\text{frac}}(\mathcal{M}_{\text{agnostic}},\Delta) = N_{\text{frac}}(\mathcal{F},\mathcal{F}^+,\Delta), \qquad \forall \Delta > 0.$$

More generally, for any 1-Lipschitz loss, we have  $N_{\mathsf{frac}}(\mathcal{M}_{\mathsf{agnostic}}, \Delta) \leq N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$ .

In particular, under absolute loss, the agnostic learnability with  $(\mathcal{F}, \mathcal{F}^+)$  is characterized by the finiteness of the complexity measure  $N_{\text{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$ .

**Realizable regression.** We consider the "easier" task of realizable regression. Given the function class  $\mathcal{F}$ , a model  $M \in \Delta(\mathcal{Z})$  is realizable if there exists  $f^M \in \mathcal{F}$  such that for  $(x, y) \sim M$ ,  $y = f^M(x)$  with probability 1. Let  $\mathcal{M}_{\mathcal{F}, \text{realizable}}$  be the class of all realizable models.

**Lemma C.3.** Under the absolute loss  $L_{abs}$ , it holds that  $N_{frac}(\mathcal{M}_{\mathcal{F},realizable}, \Delta) = N_{frac}(\mathcal{F}, \mathcal{F}^+, \Delta)$ for  $\Delta > 0$ .

Therefore, under absolute loss, the learnability of realizable regression is also characterized by the finiteness of the fractional covering number  $N_{\text{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$ . In particular, the agnostic learnability is equivalent to the realizable learnability. A similar argument also applies to the squared loss, where we can show that  $N_{\text{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$  simultaneously characterizes the learnability of agnostic regression, well-specified regression (Section 5.2), and realizable regression.

Separation between proper learning and improper learning. We show that, for highdimensional linear model, there is a separation between proper and improper learning under LDP. More specifically, we consider  $\mathcal{X} := \{x \in \mathbb{R}^d : ||x|| \leq 1\}$ , and the function class  $\mathcal{F}$  given by

$$\mathcal{F}_{\mathsf{Lin}} := \{ f_{\theta}(x) = \langle \theta, x \rangle \}_{\theta : \|\theta\| \le 1}.$$

**Proposition C.4.** Let  $\mathcal{F}^+ := \{f_{\theta}(x) = \langle \theta, x \rangle\}_{\theta \in \mathbb{R}^d}$  be the class of unbounded linear functions. Then it holds that

$$\log N_{\mathsf{frac}}(\mathcal{F}, \Delta) \ge \Omega\left(d\right), \qquad \log N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta) \le \tilde{O}\left(\frac{1}{\Delta^2}\right).$$

Therefore, d-dimensional proper linear regression is infeasible when d is unbounded, while improper learning is still tractable as  $d \to \infty$ . Proof appears in Appendix J.6.

### D Technical Tools

The following lemma can be regarded as a "chain rule" of Hellinger distance [Jayram, 2009] (see also Duchi [2023, Lemma 11.5.3] or Foster et al. [2024, Lemma D.2]).

**Lemma D.1** (Sub-additivity for squared Hellinger distance). Let  $(\mathcal{X}^1, \mathfrak{F}^1), \ldots, (\mathcal{X}^T, \mathfrak{F}^T)$  be a sequence of measurable spaces, and let  $\mathcal{X}_t = \prod_{i=1}^t \mathcal{X}^i$  and  $\mathfrak{F}_t = \bigotimes_{i=1}^t \mathfrak{F}^i$ . For each t, let  $\mathbb{P}_t(\cdot | \cdot)$  and  $\mathbb{Q}_t(\cdot | \cdot)$  be probability kernels from  $(\mathcal{X}_{t-1}, \mathfrak{F}_{t-1})$  to  $(\mathcal{X}^t, \mathfrak{F}^t)$ .

Let  $\mathbb{P}$  and  $\mathbb{Q}$  be the laws of  $X_1, \ldots, X_T$  under  $X_t \sim \mathbb{P}_t(\cdot \mid X_{1:t-1})$  and  $X_t \sim \mathbb{Q}_t(\cdot \mid X_{1:t-1})$  respectively. Then it holds that

$$D_{\mathrm{H}}^{2}(\mathbb{P},\mathbb{Q}) \leq 7 \mathbb{E}_{\mathbb{P}}\left[\sum_{t=1}^{T} D_{\mathrm{H}}^{2}(\mathbb{P}_{t}(\cdot \mid X_{1:t-1}), \mathbb{Q}_{t}(\cdot \mid X_{1:t-1}))\right].$$

We also invoke the Minimax theorem.

**Theorem D.2** (Ky Fan's minimax theorem, Fan [1953]). Let X be a compact Hausdorff space and Y an arbitrary set (not topologized). Let f be a real-valued function on  $X \times Y$  such that, for every  $y \in Y$ ,  $f(\cdot, y)$  is continuous over X.

Then, if f is convex-like on X and concave-like on Y, then

$$\min_{x \in X} \sup_{y \in Y} f(x, y) = \sup_{y \in Y} \min_{x \in X} f(x, y).$$

Therefore, if f is instead concave-like on X and convex-like on Y, then we can apply Theorem D.2 to -f to obtain

$$\max_{x \in X} \inf_{y \in Y} f(x, y) = \inf_{y \in Y} \max_{x \in X} f(x, y).$$

**Theorem D.3** (Kakutani's fixed point theorem, Osborne and Rubinstein [1994, Lemma 20.1]). Let X be a compact convex subset of  $\mathbb{R}^n$ , and let  $F: X \to \mathscr{P}(X)$  be a set-valued function for which

- 1. for all  $x \in X$ , the set F(x) is nonempty and convex, and
- 2. F is upper hemicontinuous (i.e. for all sequences  $x_n$  and  $y_n$  such that  $y_n \in F(x_n)$  for all n,  $x_n \to x, y_n \to y$ , then we have  $y \in F(x)$ ).

Then, there exists  $x \in X$  such that  $x \in F(x)$ .

### **E** Proofs for Lower Bounds

#### E.1 Proof of Theorem 7

In this section, we prove a more general version of Theorem 7 through the approach developed in Chen et al. [2024], which applies to any loss function L.

Given model class  $\mathcal{M}$ , for each  $\varepsilon > 0$  and  $\delta \in [0, 1]$ , we define the quantile-based PAC DEC as

$$\mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{H}}(\mathscr{P},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{\mathcal{P} \in \mathscr{P}} \left\{ \widehat{L}_{\delta}(\mathcal{P},p) \mid \inf_{M \in \mathrm{co}(\mathcal{P})} \mathbb{E}_{\boldsymbol{\pi} \sim q} D_{\mathrm{H}}^{2}\left(M(\boldsymbol{\pi}),\overline{M}(\boldsymbol{\pi})\right) \leq \varepsilon^{2} \right\},$$
(48)

where  $\widehat{L}_{\delta}(\mathcal{P}, p)$  is the  $\delta$ -quantile loss of p, defined as

$$\widehat{L}_{\delta}(\mathcal{P}, p) = \sup_{\Delta \ge 0} \{ \Delta : \mathbb{P}_{\pi \sim p}(L(\mathcal{P}, \pi) \ge \Delta) \ge \delta \}.$$

We also denote  $\mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{H}}(\mathscr{P}) := \sup_{\overline{M} \in \mathrm{co}(\mathcal{M})} \mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{H}}(\mathscr{P},\overline{M})$ . By definition, the quantile-based PAC DEC is always bounded by the original hybrid PAC DEC:

$$\mathsf{p-dec}^{\mathsf{H}}_{\varepsilon}(\mathscr{P},\overline{M}) - \delta \le \mathsf{p-dec}^{\mathsf{q},\mathsf{H}}_{\varepsilon,\delta}(\mathscr{P},\overline{M}) \le \delta^{-1}\mathsf{p-dec}^{\mathsf{H}}_{\varepsilon}(\mathscr{P},\overline{M}), \tag{49}$$

as long as the loss function is bounded in [0, 1]. However, such a conversion can be loose in general.

The advantage of considering the quantile private PAC-DEC is that it provides the following unified lower bound for PAC learning under hybrid DMSO. Proof is presented in Appendix E.4.

**Proposition E.1** (Quantile-based hybrid DEC lower bound). For any  $T \ge 1$  and constant  $\delta \in [0,1)$ , we denote  $\underline{\varepsilon}_{\delta}(T) := \frac{1}{13}\sqrt{\frac{\delta}{T}}$ . Then, under hybrid DMSO, for any T-round algorithm Alg, there exists  $\mathcal{P}^{\star} \in \mathscr{P}$  and a distribution  $\mu^{\star} \in \Delta(\mathcal{P}^{\star})$ , such that for the stationary environment Env specified by  $\mu^{\star}$ ,

$$L(\mathcal{P}^{\star}, \pi_{T+1}) \geq \sup_{\overline{M}} \mathsf{p-dec}_{\underline{\varepsilon}_{\delta}(T), \delta}^{\mathbf{q}, \mathbf{H}}(\mathscr{P}, \overline{M}), \qquad \text{with probability at least } \delta/2 \text{ under } \mathbb{P}^{\mathsf{Env}, \mathsf{Alg}},$$

where the supremum  $\sup_{\overline{M}}$  is taken over all reference models  $\overline{M} \in (\Pi \to \Delta(\mathcal{O}))$ .

Subsequently, we specify the above lower bound to metric-based loss and any general loss function.

**Application: metric-based loss function.** When the loss function is metric-based (Definition 1), we can show that the quantile-based hybrid DEC can be lower bounded by the original hybrid DEC. More specifically, we prove the following lemma.

**Lemma E.2.** Suppose that for some constant  $C_1, C_2$ , it holds that for any models  $\mathcal{P}, \mathcal{P}' \in \mathscr{P}$ , any decision  $\pi \in \Pi$ ,

$$L(\mathcal{P}',\pi) \le C_1 L(\mathcal{P},\pi) + C_2 \inf_{\pi'} \left( L(\mathcal{P},\pi') + L(\mathcal{P}',\pi') \right).$$
(50)

Then for any  $\delta \in [0, \frac{1}{2})$  and any reference model  $\overline{M}$ , it holds that

$$\mathrm{p-dec}^{\mathrm{q},\mathrm{H}}_{\varepsilon,\delta}(\mathscr{P},\overline{M})\geq \frac{1}{2C_2}\mathrm{p-dec}^{\mathrm{H}}_{\varepsilon}(\mathscr{P},\overline{M}).$$

For example, when  $L(\mathcal{P},\pi) = \rho(\pi^{\mathcal{P}},\pi)$  for certain pseudo-metric  $\rho$  over  $\Pi$ , Eq. (50) holds with  $C_1 = C_2 = 1$ . Therefore, Eq. (50) can be viewed as a *generalized* metric structure on the loss function L. In particular, Eq. (14) of Theorem 7 follows immediately from Proposition E.1 and Lemma E.2.

**Proof of Theorem 7: Eq. (14).** Suppose that the loss function L is metric-based. Then, Lemma E.2 implies that  $p-\operatorname{dec}_{\varepsilon,\delta}^{q,H}(\mathscr{P}) \geq \frac{1}{2}p-\operatorname{dec}_{\varepsilon}^{H}(\mathscr{P})$  for any  $\delta < \frac{1}{2}$ . Thus, applying Proposition E.1 yields

$$\sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \geq \frac{\delta}{2} \sup_{\overline{M}} \mathsf{p-dec}_{\underline{\varepsilon}_{\delta}(T),\delta}^{\mathsf{q},\mathsf{H}}(\mathscr{P}) \geq \frac{\delta}{4} \sup_{\overline{M}} \mathsf{p-dec}_{\underline{\varepsilon}_{\delta}(T)}^{\mathsf{H}}(\mathscr{P},\overline{M}).$$

Letting  $\delta \to \frac{1}{2}$  gives the desired lower bound:

$$\sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \ge \frac{1}{8} \sup_{\overline{M}} \mathsf{p-dec}_{\underline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P},\overline{M}) \ge \frac{1}{8}\mathsf{p-dec}_{\underline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P}).$$
(51)

Similarly, we can apply Proposition E.1 to general loss function.

**Proof of Theorem 7: Eq. (15).** By Eq. (49), we have

$$\operatorname{p-dec}_{\varepsilon}^{\operatorname{q},\operatorname{H}}(\mathscr{P}) \geq \operatorname{p-dec}_{\varepsilon}^{\operatorname{H}}(\mathscr{P}) - \delta.$$

Hence, Eq. (15) is a direct corollary of Proposition E.1.

As a final remark, we note that under stochastic DMSO, if the loss function is reward-based (Example 1), the quantile DEC can also be lower bounded by the constrained DEC (see Chen et al. [2024] and also Appendix I.2).

### E.2 Proof of Theorem 9

In this section, we prove Theorem 9 by first reducing to stochastic DMSO, and then apply the lower bound for stochastic DMSO (Theorem E.3).

**Reduction from hybrid DMSO to stochastic DMSO.** We first argue that for any problem under hybrid DMSO, the minimax regret can always be lower bounded by a corresponding stochastic DMSO problem. The idea follows from the observation of Foster et al. [2022b].

For any stationary environment Env constrained by  $\mathscr{P}$ , Env is specified by a constraint  $\mathcal{P} \in \mathscr{P}$  and  $\mu \in \Delta(\mathcal{P})$ . Then, for each round  $t \in [T]$ , the model  $M^t \sim \mu$  independently, and hence conditional on  $(\mathcal{H}_{t-1}, \pi_t)$ , the observation  $o_t \sim M_{\mu}(\pi_t)$ , where  $M_{\mu} = \mathbb{E}_{M' \sim \mu}[M'] \in \operatorname{co}(\mathcal{P})$ . Therefore, for any T-round algorithm Alg, the marginal distribution of  $\mathcal{H}_T$  generated by Alg under Env agrees with the distribution of  $\mathcal{H}_T$  generated by Alg under the model  $M_{\mu}$ , i.e.,

$$\mathbb{P}^{\mathsf{Env},\mathsf{Alg}}(\mathcal{H}_T=\cdot)=\mathbb{P}^{M_{\mu},\mathsf{Alg}}(\mathcal{H}_T=\cdot).$$

In particular, using the linearity of the value function, we have

$$\mathbb{E}^{\operatorname{Env},\operatorname{Alg}}\left[\sum_{t=1}^{T} V^{M^{t}}(\pi_{t})\right] = \mathbb{E}^{M_{\mu},\operatorname{Alg}}\left[\sum_{t=1}^{T} V^{M_{\mu}}(\pi_{t})\right],$$

and hence

$$\mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \geq \max_{\pi^{\star} \in \Pi} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}} \left[ \sum_{t=1}^{T} V^{M^{t}}(\pi^{\star}) - V^{M^{t}}(\pi_{t}) \right]$$
$$= \max_{\pi^{\star} \in \Pi} \mathbb{E}^{M_{\mu},\mathsf{Alg}} \left[ \sum_{t=1}^{T} V^{M_{\mu}}(\pi^{\star}) - V^{M_{\mu}}(\pi_{t}) \right] = \mathbb{E}^{M_{\mu},\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)]$$

Note that for any  $M \in \mathcal{M}_{\mathscr{P}}$ , there exists  $\mathcal{P} \in \mathscr{P}$  and  $\mu \in \Delta(\mathcal{P})$  such that  $M = \mathbb{E}_{M' \sim \mu}[M']$ , and hence there exists a corresponding stationary environment. Therefore, for any algorithm Alg, it holds that

$$\sup_{\text{stationary Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \ge \sup_{M \in \mathcal{M}_{\mathscr{P}}} \mathbb{E}^{M,\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)],$$
(52)

where  $\sup_{\mathsf{Env}}$  is taken over all stationary environments  $\mathsf{Env}$  constrained by  $\mathscr{P}$ .

**Reduction to the regret DEC lower bound.** Then, we invoke the following lower bound, which is strengthened from Foster et al. [2023b], Glasgow and Rakhlin [2023], Chen et al. [2024]. The proof is deferred to Appendix E.6.

**Theorem E.3** (Constrained DEC lower bounds for stochastic DMSO). Let  $T \ge 1$ , and  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \Delta(\mathcal{O}))$  be a given model class. Suppose that V is a value function such that  $V^{M}(\pi) \in [0, V_{\max}]$ , and for any  $\pi \in \Pi$ , there exists  $\phi_{\pi} \in \Phi$ , such that

$$\left|V^{M}(\pi) - V^{\overline{M}}(\pi)\right| \le C_{V} D_{\mathrm{H}} \left(M(\pi, \phi_{\pi}), \overline{M}(\pi, \phi_{\pi})\right), \qquad \forall M \in \mathcal{M}, \overline{M} \in \mathrm{co}(\mathcal{M}).$$
(53)

Then for any T-round algorithm Alg, it holds that

$$\sup_{M \in \mathcal{M}} \mathbb{E}^{M, \mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \geq \frac{T}{8} \left( \mathsf{r-dec}_{\underline{\varepsilon}(T)}^{\mathsf{c}}(\mathcal{M}) - 6C_{V\underline{\varepsilon}}(T) - \frac{V_{\max}}{T} \right)$$

where  $\underline{\varepsilon}(T) = \frac{1}{24\sqrt{T}}$ .

Theorem 9 is then proven by combining Eq. (52) and Theorem E.3.

E.3 Instantiations

In the following, we extend the discussion in Section 2 and apply Theorem 7 and Theorem 9 to prove the lower bounds for query-based learning and LDP learning.

#### E.3.1 Query-based learning: Proof of Theorem 14

In this section, we formalize the discussion in Section 2.2 and prove that the SQ DEC can be derived from the hybrid DEC with  $\mathscr{P} = \mathscr{P}_{\tau\text{-query}}$ . In particular, we derive Theorem 14 from Theorem 7. Alternatively, a direct proof of Theorem 7 is presented in Appendix H.1.

**From hybrid DEC to SQ DEC.** The key observation is the following lemma, which relates the squared Hellinger distance to the "error probability"-style quantity in the definition of SQ DEC (4).

**Lemma E.4.** Suppose that  $P \in \Delta(\mathcal{O})$ , and  $\mathcal{O}_0 \subseteq \mathcal{O}$  is a measurable subset of  $\mathcal{O}$ . Then it holds that

$$\frac{1}{2}P(\mathcal{O}_0^c) \le \inf_{P': \operatorname{supp}(P') \subseteq \mathcal{O}_0} D_{\mathrm{H}}^2(P', P) \le P(\mathcal{O}_0^c).$$

Note that  $\mathcal{P}_M$  consists of all models M' such that  $\operatorname{supp}(M'(\pi)) \subseteq \{v : \|M(\pi) - v\| \leq \tau\}$  for all  $\pi \in \Pi$ , and particularly,  $\mathcal{P}_M$  is convex. Therefore, we can bound the quantity

$$\inf_{M'\in\mathcal{P}_M} \mathbb{E}_{\boldsymbol{\pi}\sim q} D_{\mathrm{H}}^2\left(M'(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})\right) = \mathbb{E}_{\boldsymbol{\pi}\sim q} \inf_{M'\in\mathcal{P}_M} D_{\mathrm{H}}^2\left(M'(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})\right)$$

using Lemma E.4:

$$\frac{1}{2}\mathbb{P}_{\boldsymbol{\pi}\sim q, v\sim\overline{M}(\boldsymbol{\pi})}(\|M(\boldsymbol{\pi})-v\|>\tau) \leq \inf_{M'\in\mathcal{P}_M}\mathbb{E}_{\boldsymbol{\pi}\sim q}D_{\mathrm{H}}^2\left(M'(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})\right) \leq \mathbb{P}_{\boldsymbol{\pi}\sim q, v\sim\overline{M}(\boldsymbol{\pi})}(\|M(\boldsymbol{\pi})-v\|>\tau).$$

Therefore, we have proven the following lemma.

**Lemma E.5.** Suppose that  $\tau \geq 0$ ,  $\mathscr{P}_{\tau\text{-query}}$  is specified by the model class  $\mathcal{M} \subseteq (\mathbf{\Pi} \to \mathcal{V})$ . Then, for any reference model  $\overline{M} : \mathbf{\Pi} \to \Delta(\mathcal{V})$ , it holds that

$$\mathsf{p}\text{-}\mathsf{dec}_{\varepsilon/2}^{\tau\operatorname{-}\mathsf{SQ}}(\mathcal{M},\overline{M}) \leq \mathsf{p}\text{-}\mathsf{dec}_{\varepsilon}^{\mathsf{H}}(\mathscr{P}_{\tau\operatorname{-}\mathsf{query}},\overline{M}) \leq \mathsf{p}\text{-}\mathsf{dec}_{\varepsilon}^{\tau\operatorname{-}\mathsf{SQ}}(\mathcal{M},\overline{M}), \qquad \forall \varepsilon \geq 0.$$

In particular, we have  $p-dec_{\varepsilon}^{H}(\mathscr{P}_{\tau-query}) \leq p-dec_{\varepsilon}^{\tau-sQ}(\mathcal{M}).^{10}$ 

**Proof of Theorem 14.** For metric-based loss L, we can apply Eq. (51) with  $\mathscr{P}_{\tau-\mathsf{query}}$ :

$$\sup_{\mathsf{Env}} \mathbb{E}^{\mathsf{Env},\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \geq \frac{1}{8} \sup_{\overline{M}} \mathsf{p-dec}_{\underline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P},\overline{M}) \geq \frac{1}{8}\mathsf{p-dec}_{\underline{\varepsilon}(T)/2}^{\tau-\mathsf{SQ}}(\mathcal{M}),$$

where the supremum is taken over all environments specified by a GQ oracle  $\mathsf{GQ}_M^{\tau}$  with respect to a model  $M \in \mathcal{M}$ , and the second inequality follows from Lemma E.5. Similarly, for more general loss L, a lower bound in terms of  $\mathsf{p-dec}_{\varepsilon}^{\tau-\mathsf{SQ}}(\mathcal{M})$  also follows from Eq. (15) of Theorem 7.

**Proof of Lemma E.4.** We first consider the distribution  $P_0 = P(\cdot | o \in \mathcal{O}_0)$ . Clearly,  $supp(P_0) \subseteq \mathcal{O}_0$ , and

$$D_{\rm H}^2(P_0, P) = \frac{1}{2} \left[ P(\mathcal{O}_0^c) + \left(1 - \sqrt{P(\mathcal{O}_0)}\right)^2 \right] \le D_{\rm TV}(P_0, P) = P(\mathcal{O}_0^c).$$

Hence, the upper bound is proven.

Next, we proceed to prove the lower bound. For any  $P' \in \Delta(\mathcal{O})$  such that  $\operatorname{supp}(P') \subseteq \mathcal{O}_0$ , we fix a base measure  $\mu$ , and then

$$D_{\mathrm{H}}^{2}(P_{0},P) = 1 - \int_{\mathcal{O}} \sqrt{\frac{dP}{d\mu} \cdot \frac{dP'}{d\mu}} \mu(do)$$
  
$$= 1 - \sqrt{P(\mathcal{O}_{0})} \int_{\mathcal{O}_{0}} \sqrt{\frac{dP_{0}}{d\mu} \cdot \frac{dP'}{d\mu}} \mu(do)$$
  
$$= 1 - \sqrt{P(\mathcal{O}_{0})} + \sqrt{P(\mathcal{O}_{0})} D_{\mathrm{H}}^{2}(P',P_{0})$$
  
$$\geq 1 - \sqrt{P(\mathcal{O}_{0})} \geq \frac{1}{2} P(\mathcal{O}_{0}^{c}).$$

This gives the desired lower bound.

#### E.3.2 LDP learning: Proof of Theorem 19 (1) and Theorem 27

We first recall the discussion in Section 2.3: Given a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  and the class  $\mathcal{Q} = \mathcal{Q}_{\alpha}$  of all  $\alpha$ -DP channels (from  $\mathcal{Z}$  to  $\mathcal{O}$ ), each model  $M \in \mathcal{M}$  induces a map  $M^{\sharp} : \Pi \times \mathcal{Q} \to \Delta(\mathcal{O})$  by  $M^{\sharp}(\pi, \mathbb{Q}) = \mathbb{Q} \circ M(\pi)$  for all  $\pi \in \Pi$ ,  $\mathbb{Q} \in \mathcal{Q}$ . Therefore,  $\mathcal{M}$  induces a model class  $\mathcal{M}^{\sharp}$  under hybrid DMSO:

$$\mathcal{M}^{\sharp} := \left\{ M^{\sharp} : M \in \mathcal{M} \right\} \subseteq (\Pi \times \mathcal{Q} \to \Delta(\mathcal{O}))$$
(54)

Then, a direct application of Proposition 20 yields the following lemma.

<sup>&</sup>lt;sup>10</sup>The converse might not hold, because in our definition (4) of SQ DEC, the supremum is taken over all reference models  $\overline{M} : \mathbf{\Pi} \to \Delta(\mathcal{V})$ .

**Lemma E.6.** Let the model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  be given. For the corresponding constraint class  $\mathscr{P}_{LDP} = \{M^{\sharp} : M \in \mathcal{M}\}, \text{ it holds that}$ 

$$\mathrm{p\text{-}dec}^{\mathrm{LDP}}_{c_0\varepsilon/\alpha}(\mathcal{M}) \leq \mathrm{p\text{-}dec}^{\mathrm{H}}_{\varepsilon}(\mathscr{P}_{\mathrm{LDP}}) = \mathrm{p\text{-}dec}^{\mathrm{c}}_{\varepsilon}(\mathcal{M}^{\sharp}) \leq \mathrm{p\text{-}dec}^{\mathrm{LDP}}_{c_1\varepsilon/\alpha}(\mathcal{M}), \quad \forall \varepsilon > 0,$$

where  $c_0, c_1 > 0$  are universal constants. Similarly, we also have

$$\operatorname{\mathsf{r-dec}}_{c_0\varepsilon/\alpha}^{\operatorname{\mathsf{LDP}}}(\mathcal{M}) \leq \operatorname{\mathsf{r-dec}}_{\varepsilon}^{\operatorname{\mathsf{H}}}(\mathscr{P}_{\operatorname{\mathsf{LDP}}}) = \operatorname{\mathsf{r-dec}}_{\varepsilon}^{\operatorname{c}}(\mathcal{M}^{\sharp}) \leq \operatorname{\mathsf{r-dec}}_{c_1\varepsilon/\alpha}^{\operatorname{\mathsf{LDP}}}(\mathcal{M}), \quad \forall \varepsilon > 0.$$

Therefore, there is an equivalence between the hybrid DECs and the private DECs. Based on such an equivalence, we apply the hybrid DEC lower bounds (Theorem 8 and Theorem 9) to prove Theorem 19 (1) and Theorem 27. The proof of Theorem 19 (2) is deferred to Appendix I.2, as it involves the specific properties of reward-based loss.

**Proof of Theorem 19 (1).** Fix a *T*-round  $\alpha$ -LDP algorithm Alg. Then, by Theorem 7, it holds that

$$\sup_{M \in \mathcal{M}} \mathbb{E}^{M, \mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \geq \frac{1}{8} \mathsf{p}\text{-}\mathsf{dec}_{\underline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P}_{\mathsf{LDP}}) \geq \frac{1}{8} \mathsf{p}\text{-}\mathsf{dec}_{c_0\underline{\varepsilon}(T)/\alpha}^{\mathsf{LDP}}(\mathcal{M}),$$

where  $\underline{\varepsilon}(T) = \frac{1}{20\sqrt{T}}$ , and the second inequality follows from Lemma E.6.

**Proof of Theorem 27.** We only need to verify Assumption 3. For any decision  $\pi \in \Pi$ , we consider the binary channel  $Q_{\pi} \in Q_{\alpha}$  given by

$$Q_{\pi}(+1|z) = \frac{1 + c_{\alpha}R(z,\pi)}{2}, \qquad Q_{\pi}(-1|z) = \frac{1 - c_{\alpha}R(z,\pi)}{2},$$

where  $c_{\alpha} = 1 - e^{-\alpha}$  ensures that  $Q_{\pi}$  is  $\alpha$ -DP (cf. Example 5), and we assume without loss of generality that  $\{-1, 1\} \subseteq \mathcal{O}$ . Then, by definition, it holds that

$$c_{\alpha}\left|V^{M}(\pi)-V^{\overline{M}}(\pi)\right| \leq D_{\mathrm{TV}}\left(M(\pi,\mathsf{Q}_{\pi}),\overline{M}(\pi,\mathsf{Q}_{\pi})\right) \leq \sqrt{2}D_{\mathrm{H}}\left(M(\pi,\mathsf{Q}_{\pi}),\overline{M}(\pi,\mathsf{Q}_{\pi})\right).$$

Therefore, Assumption 3 holds with  $C_V = \frac{\sqrt{2}}{c_{\alpha}} = O(\frac{1}{\alpha})$ . Hence, for any  $\alpha$ -LDP algorithm Alg, Theorem 9 yields

$$\begin{split} \sup_{M \in \mathcal{M}} \mathbb{E}^{M,\mathrm{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] &\geq \frac{T}{8} \Big( \mathsf{p}\text{-}\mathsf{dec}_{\underline{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P}_{\mathrm{LDP}}) - 6C_{V\underline{\varepsilon}}(T) \Big) - 1 \\ &\geq \frac{T}{8} \Big( \mathsf{r}\text{-}\mathsf{dec}_{c_{0\underline{\varepsilon}}(T)/\alpha}^{\mathrm{LDP}}(\mathcal{M}) - \frac{6\sqrt{2}\underline{\varepsilon}(T)}{c_{\alpha}} \Big) - 1, \end{split}$$

where  $\underline{\varepsilon}(T) = \frac{1}{24\sqrt{T}}$ , and the second inequality follows from Lemma E.6. This gives the desired lower bound.

**Proof of Lemma E.6.** We begin with the first inequality for private PAC-DEC. By Proposition 20, for any  $\alpha$ -DP channel  $Q \in Q$ , there exists a distribution  $\tilde{q}_Q \in \Delta(\mathcal{L})$ , such that

$$D_{\mathrm{H}}^{2}\left(M^{\sharp}(\pi, \mathbb{Q}), \overline{M}^{\sharp}(\pi, \mathbb{Q})\right) \leq \frac{(e^{\alpha} - 1)^{2}}{8} \mathbb{E}_{\ell \sim \tilde{q}_{\mathbb{Q}}} \mathsf{D}_{\ell}^{2}(M(\pi), \overline{M}(\pi)), \qquad \forall \pi \in \Pi, M \in \mathcal{M}, \overline{M} \in \mathrm{co}(\mathcal{M}).$$

Therefore, for any  $q \in \Delta(\Pi \times Q)$ , there exists  $\tilde{q} \in \Delta(\Pi \times \mathcal{L})$  such that

$$\left\{ M \in \mathcal{M} : \mathbb{E}_{(\pi,\ell) \sim \tilde{q}} \mathsf{D}^2_{\ell}(M(\pi), \overline{M}(\pi)) \le \left(c_0 \varepsilon / \alpha\right)^2 \right\} \subseteq \left\{ M \in \mathcal{M} : \mathbb{E}_{\pi \sim q} D^2_{\mathrm{H}}\left(M^{\sharp}(\pi), \overline{M}^{\sharp}(\pi)\right) \le \varepsilon^2 \right\},$$

where  $c_0 > 0$  is a lower bound of  $\frac{4\alpha}{e^{\alpha}-1}$  that only depends on  $\alpha_0$ . Then, by the definition of private PAC-DEC, we know

$$\mathrm{p\text{-}dec}^{\mathrm{c}}_{\varepsilon}(\mathcal{M}^{\sharp},\overline{M}^{\sharp})\geq \mathrm{p\text{-}dec}^{\mathrm{LDP}}_{c_{0}\varepsilon/\alpha}(\mathcal{M},\overline{M}), \qquad \forall \overline{M}\in\mathrm{co}(\mathcal{M}).$$

Note that  $\operatorname{co}(\mathcal{M}^{\sharp}) = \operatorname{co}(\mathcal{M})^{\sharp}$ , and hence we have  $\operatorname{\mathsf{p-dec}}_{\varepsilon}^{\operatorname{c}}(\mathcal{M}^{\sharp}) \geq \operatorname{\mathsf{p-dec}}_{c_0\varepsilon/\alpha}^{\operatorname{\mathsf{LDP}}}(\mathcal{M})$ .

Next, we prove the second inequality for the private PAC-DEC. Recall that for any  $\ell \in \mathcal{L}$ , there is a corresponding binary channel  $Q_{\ell}$ , such that

$$\begin{aligned} D_{\mathrm{H}}^{2}\left(\mathsf{Q}_{\ell}\circ P_{1},\mathsf{Q}_{\ell}\circ P_{2}\right) &\geq \frac{1}{2}D_{\mathrm{TV}}\left(\mathsf{Q}_{\ell}\circ P_{1},\mathsf{Q}_{\ell}\circ P_{2}\right)^{2} \\ &= \frac{1}{2}D_{\mathrm{TV}}\left(\mathrm{Bern}\left(\frac{1+c_{\alpha}\mathbb{E}_{P_{1}}\ell(z)}{2}\right),\mathrm{Bern}\left(\frac{1+c_{\alpha}\mathbb{E}_{P_{2}}\ell(z)}{2}\right)\right)^{2} \\ &= \frac{c_{\alpha}^{2}}{8}\left|\mathbb{E}_{P_{1}}\ell(z)-\mathbb{E}_{P_{2}}\ell(z)\right|^{2} \geq (\alpha/c_{1})^{2}\mathsf{D}_{\ell}^{2}(P_{1},P_{2}),\end{aligned}$$

where  $c_1 > 0$  is a upper bound of  $\frac{4\alpha}{c_{\alpha}}$  that only depend on  $\alpha_0$ . Therefore, for any  $q \in \Delta(\Pi \times \mathcal{L})$ , there exists  $q' \in \Delta(\Pi \times \mathcal{Q})$ , such that

$$\left\{ M \in \mathcal{M} : \mathbb{E}_{\boldsymbol{\pi} \sim q'} D_{\mathrm{H}}^2 \left( M^{\sharp}(\boldsymbol{\pi}), \overline{M}^{\sharp}(\boldsymbol{\pi}) \right) \leq \varepsilon^2 \right\} \subseteq \left\{ M \in \mathcal{M} : \mathbb{E}_{(\boldsymbol{\pi}, \ell) \sim q} \mathsf{D}_{\ell}^2(M(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})) \leq (c_1 \varepsilon / \alpha)^2 \right\}.$$

Then, by the definition of private PAC-DEC, we know

$$\mathsf{p}\text{-}\mathsf{dec}^{\mathsf{c}}_{\varepsilon}(\mathcal{M}^{\sharp},\overline{M}^{\sharp}) \leq \mathsf{p}\text{-}\mathsf{dec}^{\mathsf{LDP}}_{c_{1}\varepsilon/\alpha}(\mathcal{M},\overline{M}), \qquad \forall \overline{M} \in \mathrm{co}(\mathcal{M}),$$

and hence  $\mathsf{p-dec}^{c}_{\varepsilon}(\mathcal{M}^{\sharp}) \leq \mathsf{p-dec}^{\mathsf{LDP}}_{c_{1}\varepsilon/\alpha}(\mathcal{M}).$ 

The bounds for private regret-DEC can be proven analogously, and we omit the proof for succinctness.  $\hfill \square$ 

### E.4 Proof of Proposition E.1

Following Foster et al. [2021], we first introduce some notations.

Recall that an algorithm  $\mathsf{Alg} = \{q_t\}_{t \in [T]} \cup \{p\}$  in hybrid DMSO is specified by a sequence of mappings, where the *t*-th mapping  $q_t(\cdot \mid \mathcal{H}_{t-1})$  specifies the distribution of  $\pi_t = (\pi_t, \phi_t)$  based on the history  $\mathcal{H}_{t-1}$ , and the final map  $p(\cdot \mid \mathcal{H}_T)$  specifies the distribution of  $\pi_{T+1}$  based on  $\mathcal{H}_T$ . Therefore, for any model  $M : \mathbf{\Pi} \to \Delta(\mathcal{O})$ , we define

$$q_{M,\mathsf{Alg}} = \mathbb{E}^{M,\mathsf{Alg}} \left[ \frac{1}{T} \sum_{t=1}^{T} q_t(\cdot | \mathcal{H}_{t-1}) \right] \in \Delta(\mathbf{\Pi}), \quad p_{M,\mathsf{Alg}} = \mathbb{E}^{M,\mathsf{Alg}}[p(\mathcal{H}_T)] \in \Delta(\Pi), \tag{55}$$

The distribution  $q_{M,\text{Alg}}$  is the expected distribution of the average profile  $(\pi_1, \dots, \pi_T)$ , and  $p_{M,\text{Alg}}$  is the expected distribution of the output decision  $\pi_{T+1}$ .

Using the sub-additivity of the squared Hellinger divergence (by Lemma D.1, see e.g., Chen et al. [2024, Section 3.2]), for any model  $M, \overline{M}$ , it holds that

$$D_{\mathrm{H}}^{2}\left(\mathbb{P}^{M,\mathsf{Alg}},\mathbb{P}^{\overline{M},\mathsf{Alg}}\right) \leq 7\mathbb{E}^{\overline{M},\mathsf{Alg}}\left[\sum_{t=1}^{T} D_{\mathrm{H}}^{2}\left(M(\boldsymbol{\pi}_{t}),\overline{M}(\boldsymbol{\pi}_{t})\right)\right]$$
(56)

$$= 7T \cdot \mathbb{E}_{\boldsymbol{\pi} \sim q_{\overline{M}, \mathsf{Alg}}} D_{\mathrm{H}}^{2} \left( M(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi}) \right).$$
(57)

With Eq. (56), we now present the proof of Proposition E.1 (which is essentially following the analysis in Chen et al. [2024]).

**Proof of Proposition E.1.** We abbreviate  $\varepsilon = \underline{\varepsilon}(T)$ . Fix a  $\Delta < \sup_{\overline{M}} \mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{H}}(\mathscr{P},\overline{M})$ , and then there exists  $\overline{M}$  such that  $\Delta < \mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{H}}(\mathscr{P},\overline{M})$ . Hence, by the definition (48), we know that

$$\Delta < \sup_{\mathcal{P}\in\mathscr{P}} \left\{ \widehat{L}_{\delta}(\mathcal{P}, p_{\overline{M}, \mathsf{Alg}}) \mid \inf_{M \in \operatorname{co}(\mathcal{P})} \mathbb{E}_{\boldsymbol{\pi} \sim q_{\overline{M}, \mathsf{Alg}}} D_{\mathrm{H}}^{2}\left(M(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})\right) \leq \varepsilon^{2} \right\}.$$

Therefore, there exists  $\mathcal{P}^{\star} \in \mathscr{P}$  and  $M^{\star} \in \operatorname{co}(\mathcal{P}^{\star})$  such that

 $\mathbb{E}_{\boldsymbol{\pi} \sim q_{\overline{M}, \mathsf{Alg}}} D^2_{\mathrm{H}} \left( M^{\star}(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi}) \right) \leq \varepsilon^2, \qquad p_{\overline{M}, \mathsf{Alg}}(\boldsymbol{\pi} : L(\mathcal{P}^{\star}, \boldsymbol{\pi}) \geq \Delta) \geq \delta.$ 

By Eq. (56), we know

$$D^2_{\mathrm{H}}\left(\mathbb{P}^{{}^{M^\star,\mathrm{Alg}}},\mathbb{P}^{\overline{M},\mathrm{Alg}}\right) \leq 7T\varepsilon^2.$$

Because  $M^* \in co(\mathcal{P}^*)$ , there exists a distribution  $\mu^* \in \Delta(\mathcal{P}^*)$  such that  $M^* = \mathbb{E}_{M \sim \mu^*}[M]$ . Then, for the stationary environment Env specified by  $\mu^*$  (i.e., it selects  $M^t \sim \mu^*$  independently), it holds that  $\mathbb{P}^{\mathsf{Env},\mathsf{Alg}}(\mathcal{H}_T = \cdot) = \mathbb{P}^{M^*,\mathsf{Alg}}(\mathcal{H}_T = \cdot)$ . Therefore, by data-processing inequality, we have

$$\frac{1}{2} \Big( \sqrt{\mathbb{P}^{\mathsf{Env},\mathsf{Alg}}(L(\mathcal{P}^{\star},\pi_{T+1}) \geq \Delta)} - \sqrt{\mathbb{P}^{\overline{M},\mathsf{Alg}}(L(\mathcal{P}^{\star},\pi_{T+1}) \geq \Delta)} \Big)^2 \leq D_{\mathrm{H}}^2 \left( \mathbb{P}^{\mathsf{Env},\mathsf{Alg}}, \mathbb{P}^{\overline{M},\mathsf{Alg}} \right) \leq 7T\varepsilon^2.$$

Therefore, combining the inequalities above, we have

$$\mathbb{P}^{\mathsf{Env},\mathsf{Alg}}(L(\mathcal{P}^{\star},\pi_{T+1}) \geq \Delta) \geq \left(\sqrt{p_{\overline{M}},\mathsf{Alg}}(\pi:L(\mathcal{P}^{\star},\pi) \geq \Delta) - \sqrt{14T\varepsilon^2}\right)^2 \geq \frac{\delta}{2},$$

where we use  $p_{\overline{M},\mathsf{Alg}}(\pi: L(\mathcal{P}^{\star},\pi) \geq \Delta) \geq \delta$  and  $\sqrt{14T\varepsilon^2} \leq (1-\frac{1}{\sqrt{2}})\sqrt{\delta}$ .

Letting  $\Delta \to \mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{H}}(\mathscr{P})$  completes the proof.

### E.5 Proof of Lemma E.2

Fix a reference model  $\overline{M}$  and let  $\Delta_0 > 0 \lor \mathsf{p-dec}_{\varepsilon,\delta}^q(\mathscr{P},\overline{M})$ . Then there exists  $p \in \Delta(\Pi), q \in \Delta(\Pi)$  such that

$$\sup_{\mathcal{P}\in\mathscr{P}}\left\{ \widehat{L}_{\delta}(\mathcal{P},p) \mid \inf_{M\in\operatorname{co}(\mathcal{P})} \mathbb{E}_{\boldsymbol{\pi}\sim q} D_{\mathrm{H}}^{2}\left(M(\boldsymbol{\pi}),\overline{M}(\boldsymbol{\pi})\right) \leq \varepsilon^{2} \right\} < \Delta_{0}.$$

Therefore, we denote

$$\mathscr{P}_{q,\varepsilon}(\overline{M}) := \{ \mathcal{P} \in \mathscr{P} : \inf_{M \in \operatorname{co}(\mathcal{P})} \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^2 \left( M(\pi), \overline{M}(\pi) \right) \le \varepsilon^2 \},$$

and it holds that

$$\mathbb{P}_{\pi \sim q}(L(\mathcal{P},\pi) \geq \Delta_0) < \delta, \qquad \forall \mathcal{P} \in \mathscr{P}_{q,\varepsilon}(\overline{M}).$$

If the constrained set  $\mathscr{P}_{q,\varepsilon}(\overline{M})$  is empty, then we immediately have  $\mathsf{p-dec}^{\mathsf{c}}_{\varepsilon}(\mathscr{P},\overline{M}) = -\infty < \Delta_0$ , and the proof is completed. Therefore, in the following we may assume  $\mathscr{P}_{q,\varepsilon}(\overline{M})$  is non-empty, and fix a model  $\mathcal{P}_0 \in \mathscr{P}_{q,\varepsilon}(\overline{M})$ .

Notice that for any model  $\mathcal{P} \in \mathscr{P}_{q,\varepsilon}(\overline{M})$ , we have

$$\mathbb{P}_{\pi \sim q}(L(\mathcal{P},\pi) < \Delta_0, L(\mathcal{P}_0,\pi) < \Delta_0) \ge 1 - 2\delta > 0,$$

and hence

$$\inf_{\pi} \left( L(\mathcal{P}, \pi) + L(\mathcal{P}_0, \pi) \right) \le 2\Delta_0.$$

Therefore, Eq. (50) implies that

$$L(\mathcal{P},\pi) \leq C_1 L(\mathcal{P}_0,\pi) + 2C_2 \Delta_0, \qquad \forall \mathcal{P} \in \mathscr{P}_{q,\varepsilon}(\overline{M}).$$

Hence, we can take any  $\pi^*$  such that  $L(\mathcal{P}_0, \pi^*) = 0$ , and let  $p \in \Delta(\Pi)$  be supported on  $\pi^*$ . Then, (p,q) certifies that

$$\mathsf{p-dec}_{\varepsilon}^{\mathsf{H}}(\mathscr{P},\overline{M}) \leq \sup_{\mathcal{P}\in\mathscr{P}} \left\{ L(\mathcal{P},\pi^{\star}) \mid \inf_{M \in \mathrm{co}(\mathcal{P})} \mathbb{E}_{\boldsymbol{\pi} \sim q} D_{\mathrm{H}}^{2}\left(M(\boldsymbol{\pi}),\overline{M}(\boldsymbol{\pi})\right) \leq \varepsilon^{2} \right\} \leq 2C_{2}\Delta_{0}.$$

Letting  $\Delta_0 \to \mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{H}}(\mathscr{P},\overline{M})$  yields  $\mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathscr{P},\overline{M}) \leq 2C_2 \cdot \mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{H}}(\mathscr{P},\overline{M})$ , which is the desired result.

### E.6 Proof of Theorem E.3

Fix a *T*-round algorithm Alg and a reference model  $\overline{M} \in co(\mathcal{M})$ . Denote  $\varepsilon := \underline{\varepsilon}(T)$  and  $\Delta := r-dec_{\varepsilon}^{c}(\mathcal{M} \cup \{\overline{M}\}, \overline{M})$ . It remains to prove the following claim.

Claim. It holds that

$$\sup_{M \in \mathcal{M}} \mathbb{E}^{M, \mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \ge \frac{T}{8} \left( \Delta - 6C_V \varepsilon - \frac{V_{\max}}{T} \right).$$
(58)

**Proof of the claim.** We set  $\Delta_0 := \frac{1}{2} \left( \Delta - \sqrt{2}C_V \varepsilon - \frac{V_{\text{max}}}{T} \right)$ . If  $\Delta_0 \leq 0$ , then the claim is vacuous. In the following, we focus on the case  $\Delta_0 > 0$ .

Fix an arbitrary  $\phi_0 \in \Phi$ . For each decision  $\pi \in \Pi$ , we let  $\phi_{\pi} \in \Pi$  be an associated measurement such that Eq. (18) holds.

Consider a modified algorithm  $\operatorname{Alg}'$ : for  $t = 1, \dots, T$ , and history  $\mathcal{H}_{t-1}$ , we set  $q'_t(\cdot | \mathcal{H}_{t-1}) = q_t(\cdot | \mathcal{H}_{t-1})$  if the quantity  $G_{t-1} := \sum_{s=1}^{t-1} \left[ V^{\overline{M}}(\pi^{\overline{M}}) - \mathbb{E}_{\pi \sim q_s} V^{\overline{M}}(\pi) \right] < T\Delta_0$ , and set  $q'_t(\cdot | \mathcal{H}_{t-1})$  be supported on  $(\pi^{\overline{M}}, \phi_0)$  if otherwise. By our construction, it holds that under  $\operatorname{Alg}'$ ,

$$G_T = \sum_{s=1}^T \left[ V^{\overline{M}}(\pi^{\overline{M}}) - \mathbb{E}_{\pi \sim q_s} V^{\overline{M}}(\pi) \right] < T\Delta_0 + V_{\max}$$

Furthermore, we can define the stopping time

$$\tau = \max \{ t \in [T] : G_{t-1} < T\Delta_0 \}.$$

If  $\tau < T$ , then it holds that  $G_T = G_\tau \ge T\Delta_0$ .

Now, we consider  $p_0 := \mathbb{E}^{\overline{M}, \operatorname{Alg}'} \left[ \frac{1}{T} \sum_{t=1}^{T} q_t(\cdot | \mathcal{H}_{t-1}) \right] \in \Delta(\mathbf{\Pi})$  (following Appendix E.4). We let  $p'_0 \in \Delta(\Pi)$  be the marginal distribution of  $\pi$  under  $(\pi, \phi) \sim p_0$ , and  $p_1$  be the distribution of  $(\pi, \phi_{\pi})$  with  $\pi \sim p'_0$ . We set  $p = \frac{1}{2}(p_0 + p_1)$ .

Note that  $p_0'$  is the marginal distribution of  $\pi \sim p$ . Thus,

$$\begin{split} \mathbb{E}_{\pi \sim p} \left[ V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi) \right] &= \mathbb{E}_{\pi \sim p_0'} \left[ V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi) \right] \\ &= \frac{1}{T} \mathbb{E}^{\overline{M}, \mathsf{Alg}'} \left[ \sum_{t=1}^T V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi_t) \right] \\ &= \frac{1}{T} \mathbb{E}^{\overline{M}, \mathsf{Alg}'} [G_T] < \Delta_0 + \frac{V_{\max}}{T} \leq \Delta. \end{split}$$

Therefore, by the definition of  $\mathbf{r}-\mathbf{dec}^{c}_{\varepsilon}(\mathcal{M}\cup\{\overline{M}\},\overline{M})$ , there exists  $M\in\mathcal{M}$  such that

$$\mathbb{E}_{\pi \sim p}[V^{M}(\pi^{M}) - V^{M}(\pi)] \geq \Delta, \qquad \mathbb{E}_{\pi \sim p}D_{\mathrm{H}}^{2}\left(M(\pi), \overline{M}(\pi)\right) \leq \varepsilon^{2}.$$

We also have

$$\begin{aligned} \mathbb{E}_{\pi \sim p'_0} \left| V^M(\pi) - V^{\overline{M}}(\pi) \right|^2 &\leq C_V^2 \mathbb{E}_{\pi \sim p'_0} D_{\mathrm{H}}^2 \left( M(\pi, \phi_\pi), \overline{M}(\pi, \phi_\pi) \right) \\ &= C_V^2 \mathbb{E}_{\pi \sim p_1} D_{\mathrm{H}} \left( M(\pi), \overline{M}(\pi) \right) \\ &\leq 2 C_V^2 \varepsilon^2. \end{aligned}$$

Therefore, we have

$$\begin{aligned} V^{M}(\pi^{M}) - V^{\overline{M}}(\pi^{\overline{M}}) &= \left[ V^{M}(\pi^{M}) - \mathbb{E}_{\pi \sim p} V^{M}(\pi) \right] + \mathbb{E}_{\pi \sim p} \left[ V^{M}(\pi) - V^{\overline{M}}(\pi) \right] - \left[ V^{\overline{M}}(\pi^{\overline{M}}) - \mathbb{E}_{\pi \sim p} V^{\overline{M}}(\pi) \right] \\ &\geq \Delta - \sqrt{2} C_{V} \varepsilon - \left( \Delta_{0} + \frac{V_{\max}}{T} \right) \geq \Delta_{0}. \end{aligned}$$

In the following, we proceed to lower bound  $\operatorname{Reg}_{\mathsf{DM}}(\tau)$  under model M and algorithm  $\operatorname{Alg}'$ . Consider the random variable

$$X = \sum_{t=1}^{T} \mathbb{E}_{\pi \sim q_t} \left| V^M(\pi) - V^{\overline{M}}(\pi) \right|.$$

We then bound

$$\begin{aligned} \mathbf{Reg}_{\mathsf{DM}}(\tau) &= \sum_{t=1}^{\tau} \left[ V^{M}(\pi^{M}) - \mathbb{E}_{\pi \sim q_{t}} V^{M}(\pi) \right] \\ &= \tau (V^{M}(\pi^{M}) - V^{\overline{M}}(\pi^{\overline{M}})) + \sum_{t=1}^{\tau} \left[ V^{\overline{M}}(\pi^{\overline{M}}) - \mathbb{E}_{\pi \sim q_{t}} V^{\overline{M}}(\pi) \right] + \sum_{t=1}^{\tau} \left[ \mathbb{E}_{\pi \sim q_{t}} V^{\overline{M}}(\pi) - \mathbb{E}_{\pi \sim q_{t}} V^{M}(\pi) \right] \\ &\geq \tau (V^{M}(\pi^{M}) - V^{\overline{M}}(\pi^{\overline{M}})) + G_{\tau} - X, \end{aligned}$$

where the last line follows from the definition of  $G_{\tau}$  and X. Note that if  $\tau < T$ , we have  $G_{\tau} \ge T\Delta_0$ . Otherwise, we have  $\tau = T$  and  $G_{\tau} \ge 0$ . Therefore, under model M, it holds that (almost surely)

 $\operatorname{Reg}_{\mathsf{DM}}(\tau) \ge \min \left\{ T(V^{M}(\pi^{M}) - V^{\overline{M}}(\pi^{\overline{M}})), T\Delta_{0} \right\} - X \ge T\Delta_{0} - X.$ 

Consider the event  $\mathcal{E} := \{X > TCC_V \varepsilon\}$ . By Markov's inequality,

$$\begin{split} \mathbb{P}^{\overline{M},\mathsf{Alg}'}(\mathcal{E}) &\leq \frac{\mathbb{E}^{\overline{M},\mathsf{Alg}'}X^2}{(TCC_V\varepsilon)^2} \\ &\leq \frac{1}{T(CC_V\varepsilon)^2} \mathbb{E}^{\overline{M},\mathsf{Alg}'} \left[ \sum_{t=1}^T \mathbb{E}_{\pi \sim q_t} \left| V^M(\pi) - V^{\overline{M}}(\pi) \right|^2 \right] \\ &= \frac{1}{(CC_V\varepsilon)^2} \mathbb{E}_{\pi \sim p'_0} \Big[ \left| V^M(\pi) - V^{\overline{M}}(\pi) \right|^2 \Big] \leq \frac{2}{C^2}. \end{split}$$

Further, by Eq. (56), we have

$$D_{\mathrm{H}}^{2}\left(\mathbb{P}^{M,\mathrm{Alg}'},\mathbb{P}^{\overline{M},\mathrm{Alg}'}\right) \leq 7T \cdot \mathbb{E}_{\boldsymbol{\pi} \sim p_{0}} D_{\mathrm{H}}^{2}\left(M(\boldsymbol{\pi}),\overline{M}(\boldsymbol{\pi})\right) \leq 14T\varepsilon^{2}.$$

Therefore, by data-processing inequality, it holds that

$$\left|\mathbb{P}^{M,\operatorname{Alg}'}(\mathcal{E}) - \mathbb{P}^{\overline{M},\operatorname{Alg}'}(\mathcal{E})\right| \leq D_{\operatorname{TV}}\left(\mathbb{P}^{M,\operatorname{Alg}'}, \mathbb{P}^{\overline{M},\operatorname{Alg}'}\right) \leq \sqrt{28T\varepsilon^2},$$

which gives  $\mathbb{P}^{M,\operatorname{Alg}'}(\mathcal{E}) \leq \frac{2}{C^2} + \sqrt{28T\varepsilon^2}.$ 

Note that under the event  $\mathcal{E}^c$ , we have  $X \leq TCC_V \varepsilon$ . Therefore, we can lower bound

$$\mathbb{E}^{M,\operatorname{Alg}'}[\operatorname{\mathbf{Reg}_{\mathsf{DM}}}(\tau)] \geq \mathbb{E}^{M,\operatorname{Alg}'}[\mathbf{1}\left\{\mathcal{E}^{c}\right\}\operatorname{\mathbf{Reg}_{\mathsf{DM}}}(\tau)]$$
  
$$\geq \mathbb{E}^{M,\operatorname{Alg}'}[\mathbf{1}\left\{\mathcal{E}^{c}\right\}(T\Delta_{0}-X)]$$
  
$$\geq \mathbb{P}^{M,\operatorname{Alg}'}(\mathcal{E}^{c})\cdot T(\Delta_{0}-CC_{V}\varepsilon)$$
  
$$\geq \left(1-\frac{2}{C^{2}}-\sqrt{28T\varepsilon^{2}}\right)\cdot T(\Delta_{0}-CC_{V}\varepsilon)$$

In particular, we can choose C = 2, and by the choice  $\varepsilon = \frac{1}{24\sqrt{T}}$ , we have  $\mathbb{E}^{M,\operatorname{Alg}'}[\operatorname{Reg}_{\mathsf{DM}}(\tau)] \geq \frac{T}{4}(\Delta_0 - 2C_V\varepsilon)$ . Then, we can conclude that

$$\mathbb{E}^{M,\operatorname{Alg}}[\operatorname{\mathbf{Reg}}_{\mathsf{DM}}(T)] \geq \mathbb{E}^{M,\operatorname{Alg}}[\operatorname{\mathbf{Reg}}_{\mathsf{DM}}(\tau)] = \mathbb{E}^{M,\operatorname{Alg}'}[\operatorname{\mathbf{Reg}}_{\mathsf{DM}}(\tau)] \geq \frac{T}{4}(\Delta_0 - 2C_V\varepsilon).$$

This gives the desired lower bound.

### F Exploration-by-Optimization Algorithm and Guarantees

In this section, we present a generalization of the Exploration-by-Optimization Algorithm (ExO<sup>+</sup>) developed by Foster et al. [2022b], which is built upon Lattimore and Szepesvári [2020], Lattimore and Gyorgy [2021] and is later extended by Chen et al. [2024]. The ExO<sup>+</sup> algorithm of Foster et al. [2022b] has an *adversarial* regret guarantee for any model class  $\mathcal{M}$ , scaling with the *offset DEC* of the convexified model class  $co(\mathcal{M})$  and  $\log |\Pi|$ . For our purpose, we adapt it by incorporating certain measurement class  $\Phi$  and *information set structure*  $\Psi$ , so that it (1) handles any hybrid DMSO problem, and also (2) adapts to the structure of the decision space (e.g. capable of achieving an upper bound that scales with fractional covering number, Appendix F.4.4).

We organize this section as follows:

- In Appendix F.1, we introduce the notion of *information set structure*.
- In Appendix F.2, we present the detailed description of  $ExO^+$  algorithm based on a given information set structure  $\Psi$ .
- In Appendix F.3, we bound the risk (regret) of  $\mathsf{ExO}^+$  in terms of the offset DEC and the fractional covering number associated with  $(\mathscr{P}, \Psi)$ .
- In Appendix F.4, we instantiate  $ExO^+$  to prove the upper bounds of Section 3.
- In Appendix F.5, we apply ExO<sup>+</sup> to private DMSO to obtain the LDP-ExO algorithm the upper bounds of Section 5. Specifically, we instantiate LDP-ExO with the following information set structure:
  - Model-based information sets (Appendix F.5.1), where we prove Theorem 28 (1).
  - Policy-based information sets (Appendix F.5.2), where we prove Theorem 22 and Theorem 28 (2).
  - Value-based information sets (Appendix F.5.3), where we prove Proposition 24.
  - Contextual bandits (Appendix F.5.4), where we prove Proposition 29.
- The remaining subsections contain the proofs of the results of this section.

**Offset DECs.** For a model class  $\mathcal{M} \subseteq (\Pi \times \Phi \to \Delta(\mathcal{O}))$  under hybrid DMSO, we define the offset DECs [Foster et al., 2021] for each  $\gamma > 0$  as

$$\mathsf{p-dec}^{\mathrm{o}}_{\gamma}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\mathbf{\Pi})}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M,\pi)] - \gamma \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^{2}\left(M(\pi),\overline{M}(\pi)\right) \right\},\tag{59}$$

$$\mathsf{r-dec}^{\mathrm{o}}_{\gamma}(\mathcal{M},\overline{M}) := \inf_{p \in \Delta(\mathbf{\Pi})} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[V^{M}(\pi^{M}) - V^{M}(\pi)] - \gamma \mathbb{E}_{\pi \sim p}D_{\mathrm{H}}^{2}\left(M(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})\right) \right\}, \quad (60)$$

and we let

$$\mathsf{p-dec}^{\mathrm{o}}_{\gamma}(\mathcal{M}) = \sup_{\overline{M} \in \mathrm{co}(\mathcal{M})} \mathsf{p-dec}^{\mathrm{o}}_{\gamma}(\mathcal{M}, \overline{M}), \quad \mathsf{r-dec}^{\mathrm{o}}_{\gamma}(\mathcal{M}) = \sup_{\overline{M} \in \mathrm{co}(\mathcal{M})} \mathsf{r-dec}^{\mathrm{o}}_{\gamma}(\mathcal{M}, \overline{M}).$$
(61)

More generally, for any constraint set  $\mathscr{P}$  under hybrid DMSO, we define the offset hybrid DEC as

$$\mathsf{p-dec}_{\gamma}^{\mathrm{o},\mathrm{H}}(\mathscr{P},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{\substack{\mathcal{P} \in \mathscr{P} \\ M \in \mathrm{co}(\mathcal{P})}} \left\{ \mathbb{E}_{\pi \sim p}[L(\mathcal{P},\pi)] - \gamma \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^{2}\left(M(\pi),\overline{M}(\pi)\right) \right\}, \tag{62}$$

and let  $\mathsf{p-dec}^{\mathrm{o},\mathsf{H}}_{\gamma}(\mathscr{P}) \coloneqq \sup_{\overline{M} \in \mathrm{co}(\mathcal{M})} \mathsf{p-dec}^{\mathrm{o},\mathsf{H}}_{\gamma}(\mathscr{P},\overline{M}).$ 

As a remark, we note that when the loss function is bounded in [0, 1], it holds that

$$\mathsf{p-dec}^{\mathrm{o}}_{\varepsilon^{-2}}(\mathcal{M},\overline{M}) \le \mathsf{p-dec}^{\mathrm{c}}_{\varepsilon}(\mathcal{M},\overline{M}) \le \mathsf{p-dec}^{\mathrm{o}}_{\gamma}(\mathcal{M},\overline{M}) + \gamma \varepsilon^{2}, \tag{63}$$

and analogous conversions also hold for the regret-DECs and the hybrid DECs. The first inequality in Eq. (63) can be loose in general, and a tighter conversion is possible under reward-based loss function (Proposition F.10).

### F.1 Information set structure

Recall that in Section 2.1, we consider both PAC risk (1) (in terms of the loss function L) and the regret (11) (in terms of the value function V).

To present the  $ExO^+$  algorithm in a unified form, we first introduce the notion of *information set* structure. We consider two types of information set structure: Type 1 information set structure is introduced to handle "value-based" learning (cf. discussion below), generalizing Foster et al. [2022b]; Type 2 information set structure is for general PAC learning under hybrid DMSO.

**Type 1 information set structure.** We introduce the Type 1 information set structure primarily for no-regret learning in hybrid DMSO.

**Definition 17** (Type 1 information set structure). Given a constraint class  $\mathscr{P}$  under hybrid DMSO and a value function V, a Type 1 information set structure is a class  $\Psi$ , where each  $\psi \in \Psi$  is associated with a model class  $\mathcal{M}_{\psi} \subseteq (\mathbf{\Pi} \to \Delta(\mathcal{O}))$  and a decision  $\pi_{\psi} \in \Pi$ , such that the following holds:

(1) For each  $\mathcal{P} \in \mathscr{P}$ , there exists  $\psi \in \Psi$  such that  $\mathcal{P} \subseteq \mathcal{M}_{\psi}$ .

(2) The value  $M \mapsto V^M(\pi)$  is linear over  $\mathcal{M}_{\Psi} := \bigcup_{\psi \in \Psi} \operatorname{co}(\mathcal{M}_{\psi})$  for any  $\pi \in \Pi$ . We also denote  $L_{\psi}(M,\pi) = V^M(\pi_{\psi}) - V^M(\pi)$  for each  $\psi \in \Psi, M \in \mathcal{M}_{\psi}, \pi \in \Pi$ .

For no-regret learning in hybrid DMSO, the simplest Type 1 information set structure is given by  $\Psi = \mathscr{P} \times \Pi$ , and for each  $\psi = (\mathcal{P}^{\psi}, \pi^{\psi}) \in \Psi$ , we assign  $\mathcal{M}_{\psi} = \mathcal{P}^{\psi}, \pi_{\psi} = \pi^{\psi}$ . Then, the loss  $L_{\psi}(M, \pi)$  measures the sub-optimality of a decision  $\pi$  compared to the decision  $\pi_{\psi}$  (for the information set  $\psi$ ) under the model M.

Another example of information set structure is the "policy-based" one (cf. Section 3.3):  $\Psi = \Pi$ , where for each  $\pi \in \Psi$ ,  $\mathcal{M}_{\pi} = \{M : V^{M}(\pi^{M}) - V^{M}(\pi) \leq \Delta\}$ . In this example,  $\pi$  is a near-optimal decision for models in  $\mathcal{M}_{\pi}$ . With such an information set structure, we can derive an upper bound scaling with the fractional covering number of  $\mathcal{M}$  and the DEC of  $\mathcal{M}_{\Pi}$  (see Appendix F.4.4).

The notion of Type 1 information set structure can be viewed as an abstraction of the ideas of Foster et al. [2022b]. The idea of using information sets in the context of posterior sampling (and then AIR) was conveyed to the authors by Dylan Foster back in 2022.

In addition to no-regret learning in hybrid DMSO, Type 1 information set structure can also be applied to the "value-based" PAC learning under stochastic DMSO, as long as the loss function L is specified by the value function V as  $L(M,\pi) = V^M(\pi^M) - V^M(\pi)$ , where  $\pi^M = \arg \max_{\pi \in \Pi} V^M(\pi)$ .

**Type 2 information set structure.** For PAC learning under hybrid DMSO, we consider  $\mathscr{P}$  itself as an information set structure.

**Definition 18** (Type 2 information set structure for PAC learning in hybrid DMSO). Given a problem class  $(\mathcal{M}, \mathscr{P})$  under hybrid DMSO, we say that  $\Psi = \mathscr{P}$  is a Type 2 information set structure. To be consistent with Type 1 information set structure, we write  $\mathcal{M}_{\psi} = \psi$  and  $L_{\psi}(\mathcal{M}, \pi) := L(\psi, \pi)$ (i.e., the loss of a decision  $\pi$  only depends on the information set  $\psi \in \mathscr{P}$ ).

### F.2 Exploration-by-Optimization algorithm

The algorithm,  $ExO^+$ , is stated in Algorithm 1. It has two options: pac for PAC learning and reg for no-regret learning. For these two tasks, we specify different spaces S of distributions for

exploration-exploitation:

$$\mathbb{S}^{\mathsf{pac}} \coloneqq \Delta(\Pi) \times \Delta(\Pi), \qquad \mathbb{S}^{\mathsf{reg}} \coloneqq \{(q|_{\Pi}, q) : q \in \Delta(\Pi)\} \subset \Delta(\Pi) \times \Delta(\Pi),$$

where we recall that  $\mathbf{\Pi} := \mathbf{\Pi} \times \Phi$ , and for any distribution  $q \in \Delta(\mathbf{\Pi})$ ,  $q|_{\mathbf{\Pi}} \in \Delta(\mathbf{\Pi})$  is the marginal distribution of  $\pi$  under  $(\pi, \phi) \sim q$ . We note that for Type 2 information set structure, only the option pac applies.

At each round t, the algorithm maintains a reference distribution  $w_t \in \Delta(\Psi)$ , and uses it to obtain a joint exploration-exploitation distribution  $(p_t, q_t) \in \mathbb{S}$  and a weight function  $\xi_t \in \Xi := (\Psi \times \Pi \times \mathcal{O} \to \mathbb{R})$ ,<sup>11</sup> by solving a joint minimax optimization problem based on the exploration-by-optimization objective: Defining

$$\Gamma_{w,\gamma}(p,q,\xi;M,\psi) = \mathbb{E}_{\pi\sim p}[L_{\psi}(M,\pi)] - \gamma \mathbb{E}_{\pi\sim q} \mathbb{E}_{o\sim M(\pi)} \mathbb{E}_{\psi'\sim w} \left[1 - \exp\left(\xi(\psi';\pi,o) - \xi(\psi;\pi,o)\right)\right],$$
(64)

and

$$\Gamma_{w,\gamma}(p,q,\xi) = \sup_{(M,\psi):M\in\mathcal{M}_{\psi}} \Gamma_{w,\gamma}(p,q,\xi;M,\psi),$$
(65)

the algorithm solves

$$(p_t, q_t, \xi_t) \leftarrow \operatorname*{arg\,min}_{(p,q)\in\mathbb{S},\xi\in\Xi} \Gamma_{w_t,\gamma}(p,q,\xi).$$

The algorithm then samples  $\pi_t = (\pi_t, \phi_t) \sim q_t$  from the exploration distribution, executes  $\pi_t$  and observes  $o_t$  from the environment. Finally, the algorithm updates the reference distribution by performing the exponential weight update (67) with weight function  $\xi_t(\cdot; \pi_t, o_t)$ .

At the end of the interactions, the algorithm may also output  $\hat{p} = \frac{1}{T} \sum_{t=1}^{T} p_t \in \Delta(\Pi)$  as the distribution of  $\pi_{T+1}$ , which is the mixture of the per-step exploitation distributions.

Algorithm 1 Exploration-by-Optimization with information set structure  $(ExO^+)$ 

- **Input:** Decision space  $\Pi$ , measurement class  $\Phi$ , information set structure  $\Psi$ , prior  $w_1 \in \Delta(\Psi)$ , parameter  $T \ge 1, \gamma > 0$ .
- 1: For option pac, set  $\mathbb{S} = \mathbb{S}^{pac}$ ; for option reg, set  $\mathbb{S} = \mathbb{S}^{reg}$ .
- 2: for  $t = 1, \dots, T$  do
- 3: Solve the *exploration-by-optimization* objective:

$$(p_t, q_t, \xi_t) \leftarrow \underset{(p,q) \in \mathbb{S}, \xi \in \Xi}{\operatorname{arg\,min}} \Gamma_{w_t, \gamma}(p, q, \xi)$$
(66)

4: Sample  $\boldsymbol{\pi}_t = (\pi_t, \phi_t) \sim q_t$  and observe  $o_t \sim M^t(\boldsymbol{\pi}_t)$ 

5: Perform exponential-weight update:

$$w_{t+1}(\psi) \propto_{\psi} w_t(\psi) \exp(\xi_t(\psi; \boldsymbol{\pi}_t, o_t))$$
(67)

**Output:**  $\widehat{p} = \frac{1}{T} \sum_{t=1}^{T} p_t \in \Delta(\Pi)$ 

<sup>&</sup>lt;sup>11</sup>Formally, for infinite  $\Pi$  or  $\mathcal{O}, \Xi$  is the class of measurable, uniformly bounded functions over  $\Psi \times \Pi \times \mathcal{O}$ .

Following Foster et al. [2022b], we define

$$\exp_{\gamma}(\Psi, w) := \inf_{(p,q) \in \mathbb{S}, \xi \in \Xi} \Gamma_{w,\gamma}(p,q,\xi), \tag{68}$$

and  $\exp_{\gamma}(\Psi) = \sup_{q \in \Delta(\Psi)} \exp_{\gamma}(\Psi, w)$ . Note that  $\exp_{\gamma}(\Psi)$  implicitly depends on the space S.

Now, we present the primary guarantees of  $E \times O^+$ .

Bounds for Type 1 information set structure. Suppose that the algorithm  $ExO^+$  is instantiated with a Type 1 information set structure  $\Psi$  (with respect to the constraint class  $\mathscr{P}$ ), and the environment is constrained by  $\mathcal{P}^* \in \mathscr{P}$ . Define  $\overline{M}^* = \frac{1}{T} \sum_{t=1}^T M^t \in \operatorname{co}(\mathcal{P}^*)$  and consider the set

$$\mathcal{E}^{\star}_{\Delta} := \{ \psi : \mathcal{P}^{\star} \subseteq \mathcal{M}_{\psi}, V^{\overline{M}^{\star}}(\pi^{\overline{M}^{\star}}) - V^{\overline{M}^{\star}}(\pi_{\psi}) \leq \Delta \}.$$

Note that  $\mathcal{E}^{\star}_{\Delta} \subseteq \Psi$  depends on  $M^1, \dots, M^T$ , i.e.  $\mathcal{E}^{\star}_{\Delta}$  depends on the *T*-round interactions between the environment and the  $\mathsf{ExO}^+$  algorithm. We present an upper bound scaling with  $\log(1/w_1(\mathcal{E}^{\star}_{\Delta}))$ .

**Theorem F.1** (ExO<sup>+</sup> upper bound; Type 1). Let  $T \ge 1$ , the constraint class  $\mathscr{P}$  and the value function V be given, and  $\Psi$  be a Type 1 information set structure. Suppose that the environment is constrained by  $\mathscr{P}$ . Then the algorithm  $\mathsf{ExO}^+$  achieves that with probability at least  $1 - \delta$ ,

$$\max_{\pi \in \Pi} \sum_{t=1}^{T} V^{M^{t}}(\pi) - \mathbb{E}_{\pi_{t} \sim p_{t}} \left[ V^{M^{t}}(\pi_{t}) \right] \leq T \cdot \left[ \Delta + \exp_{\gamma}(\Psi) \right] + 2\gamma \cdot \left[ \log(1/w_{1}(\mathcal{E}_{\Delta}^{\star})) + \log(1/\delta) \right].$$

The proof of Theorem F.1 is deferred to Appendix F.6. It is based on bounding the performance of the exponential weight update (67), and then relating it to the performance of  $\mathsf{ExO}^+$  using the definition of  $\mathsf{exo}_{\gamma}(\Psi)$ . Different from the analysis in Foster et al. [2022b], Chen et al. [2024], the proof here has to carefully deal with  $\mathcal{E}^{\star}_{\Delta}$ , which is an event that depends on the *T*-round interactions.

Bounds for Type 2 information set structure. Similarly, for Type 2 information set structure  $\Psi = \mathscr{P}$ , we have the following guarantee of  $\mathsf{ExO}^+$ .

**Theorem F.2** (ExO<sup>+</sup> upper bound for PAC learning; Type 2). For PAC learning under hybrid DMSO, suppose that the algorithm  $ExO^+$  is instantiated with the Type 2 information set structure  $\Psi = \mathscr{P}$ , and  $w^1 = \text{Unif}(\mathscr{P})$ . Then for any environment constrained by  $\mathscr{P}$ ,  $ExO^+$  achieves with probability at least  $1 - \delta$ 

$$\mathbf{Risk}_{\mathsf{DM}}(T) = \mathbb{E}_{\pi_{T+1} \sim \widehat{p}} L(\mathcal{P}^{\star}, \pi_{T+1}) \le \exp_{\gamma}(\Psi) + \frac{2\gamma}{T} \cdot [\log |\mathscr{P}| + 2\log(1/\delta)].$$

The proof is postponed to Appendix F.7.

### F.3 Guarantees of the $ExO^+$ algorithm

In this section, we simplify the upper bound of Theorem F.1 and Theorem F.2. In the following, we bound the term  $\exp_{\gamma}(\Psi)$  and  $\log(1/w_1(\mathcal{E}^*_{\Delta}))$  separately.

Bounding ExO coefficient. We relate  $\exp_{\gamma}(\Psi)$  to the offset DECs, following Foster et al. [2022b, Theorem 3.1 and 3.2].

**Theorem F.3.** Suppose that the model class  $\mathcal{M}_{\Psi}$  is compact (Assumption 2). Then, the following holds:

(1) Suppose  $\Psi$  is a Type 1 information set structure and the value function V is uniformly continuous over  $\mathcal{M}_{\Psi}$ . Then, for PAC learning (option pac,  $\mathbb{S} = \mathbb{S}^{pac}$ ), we have

$$extsf{exo}_\gamma(\Psi) \leq extsf{p-dec}_{\gamma/4}^{\mathrm{o}}(\mathcal{M}_\Psi), \qquad orall \gamma > 0.$$

Analogously, for no-regret learning (option reg,  $\mathbb{S} = \mathbb{S}^{reg}$ ), we have

$$\exp_{\gamma}(\Psi) \leq \operatorname{r-dec}_{\gamma/4}^{o}(\mathcal{M}_{\Psi}), \qquad \forall \gamma > 0.$$

(2) If  $\Psi = \mathscr{P}$  is a Type 2 information set structure, then

$$\mathrm{exo}_{\gamma}(\Psi) \leq \mathrm{p}\text{-}\mathrm{dec}_{\gamma/4}^{\mathrm{o},\mathrm{H}}(\mathscr{P}), \qquad \forall \gamma > 0.$$

The proof of Theorem F.3 is a generalization of the analysis in Foster et al. [2022b] and is deferred to Appendix F.8.

**Bounding**  $w_1(\mathcal{E}^{\star}_{\Delta})$ . For Type 1 information set structure  $\Psi$ , we also need to provide a uniform upper bound on the quantity  $w_1(\mathcal{E}^{\star}_{\Delta})$ . Following Definition 5, we consider the fractional covering number of  $\mathscr{P}$  under an information set structure  $\Psi$ :

$$N_{\mathsf{frac}}(\mathscr{P}, \Psi; \Delta) := \inf_{w \in \Delta(\Psi)} \sup_{(\mathcal{P}, \overline{M})} \frac{1}{\mathbb{P}_{\psi \sim w}(\psi : \mathcal{P} \subseteq \mathcal{M}_{\psi}, V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi_{\psi}) \leq \Delta))},$$
(69)

where the supremum  $\sup_{(\mathcal{P},\overline{M})}$  is taken over all possible pair  $(\mathcal{P},\overline{M})$  with  $\mathcal{P} \in \mathscr{P}$  and  $\overline{M} \in \operatorname{co}(\mathcal{P})$ . Then, the optimal  $w_1^*$  is given by

$$w_{1}^{\star} := \underset{w \in \Delta(\Psi)}{\operatorname{arg\,min}} \sup_{(\mathcal{P},\overline{M})} \frac{1}{\mathbb{P}_{\psi \sim w}(\psi : \mathcal{P} \subseteq \mathcal{M}_{\psi}, V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi_{\psi}) \leq \Delta))}.$$
(70)

By definition, it holds that  $w_1^{\star}(\mathcal{E}_{\Delta}^{\star}) \geq \frac{1}{N_{\mathsf{frac}}(\mathscr{P}, \Psi; \Delta)}$  for any possible  $\mathcal{E}_{\Delta}^{\star}$ .

Putting these pieces together, we derive the following guarantees of  $ExO^+$  for PAC learning and no-regret learning under hybrid DMSO.

**Theorem F.4** (Guarantees of  $\text{ExO}^+$ ; Type 1). Let  $T \ge 1$ , parameter  $\gamma, \Delta > 0, \delta \in (0, 1)$ , constraint class  $\mathscr{P}$ , value function V be given. Suppose that  $\Psi$  is a Type 1 information set structure, and  $\mathcal{M}_{\Psi}$  is compact (Assumption 2). We instantiate  $\text{ExO}^+$  on  $\Psi$  and choose  $w_1 \in \Delta(\Psi)$  according to Eq. (70).

(1) With the option reg,  $ExO^+$  achieves with probability at least  $1 - \delta$ 

$$\begin{aligned} \mathbf{Reg}_{\mathsf{DM}}(T) &= \max_{\pi \in \Pi} \sum_{t=1}^{T} V^{M^{t}}(\pi) - \mathbb{E}_{\pi_{t} \sim q_{t}} \Big[ V^{M^{t}}(\pi_{t}) \Big] \\ &\leq T \cdot \Big[ \Delta + \mathsf{r-dec}_{\gamma/4}^{\mathrm{o}}(\mathcal{M}_{\Psi}) \Big] + 2\gamma [\log N_{\mathsf{frac}}(\mathscr{P}, \Psi; \Delta) + \log(1/\delta)]. \end{aligned}$$

(2) When  $\mathscr{P} = \mathscr{P}_{sto}$  (stochastic DMSO),  $\mathsf{ExO}^+$  with option pac achieves with probability at least  $1 - \delta$  that

$$\mathbb{E}_{\pi_{T+1}\sim\hat{p}} \Big[ V^{M^{\star}}(\pi^{M^{\star}}) - V^{M^{\star}}(\pi_{T+1}) \Big] \leq \Delta + \mathsf{p-dec}_{\gamma/4}^{\mathsf{o}}(\mathcal{M}_{\Psi}) + \frac{2\gamma}{T} [\log N_{\mathsf{frac}}(\mathscr{P}_{\mathsf{sto}}, \Psi; \Delta) + \log(1/\delta)].$$

Guarantees for Type 2 information set structure. Similarly, when  $E \times O^+$  is instantiated with Type 2 information set structure, we have a similar upper bound by simply choosing  $w_1 = \text{Unif}(\Psi)$ .

**Theorem F.5** (Guarantees of  $\mathsf{ExO}^+$ ; Type 2). Let  $T \ge 1, \gamma > 0, \delta \in (0, 1)$ , constraint class  $\mathscr{P}$  be given. Suppose that  $\mathcal{M}$  is compact (Assumption 2), and  $\mathsf{ExO}^+$  is instantiated with the Type 2 information set structure  $\Psi = \mathscr{P}$ ,  $w_1 = \mathrm{Unif}(\mathscr{P})$ , and option pac. Then with probability at least  $1 - \delta$ ,

$$\mathbb{E}_{\pi_{T+1}\sim \hat{p}}L(\mathcal{P}^{\star},\pi_{T+1}) \leq \mathsf{p-dec}_{\gamma/4}^{\mathsf{o},\mathsf{H}}(\mathscr{P}) + \frac{2\gamma \log(|\mathscr{P}|/\delta)}{T}.$$

**Remark F.6.** We assume that  $\mathcal{M}$  admits finite covering to ensure the Minimax theorem can be applied in Theorem F.3. Alternatively, we can assume that (1) the decision space  $\Pi$  is finite, (2) the latent observation space  $\mathcal{Z}$  is a compact metric space under a certain metric  $\rho$ , and (3) the value function is given by a reward function R (cf. Definition 9) with  $R(z,\pi)$  being Lipschitz with respect to z. This is indeed the case for agnostic regression task (Section 5.2).

In these assumptions, we can consider a finite  $\varepsilon$ -covering  $\mathcal{Z}_{\varepsilon}$  of  $\mathcal{Z}$ , and take  $\mathcal{M}_{\varepsilon} \subseteq (\Pi \to \Delta(\mathcal{Z}_{\varepsilon}))$  to be the model class induced by  $\mathcal{M}$ . Apply Theorem F.4 to the model class  $\mathcal{M}_{\varepsilon}$  with a sufficiently small  $\varepsilon$  yields the same bound on **Risk**<sub>DM</sub>(T) (or **Reg**<sub>DM</sub>(T), respectively).

### F.4 Proofs for upper bounds in Section 3

In the following, we instantiate Theorem F.4 and Theorem F.5 to prove the upper bounds in Section 3.

#### F.4.1 Proof of Theorem 8

For Theorem 8, we instantiate  $E \times O^+$  as in Theorem F.5, taking the Type 2 information set structure  $\Psi = \mathscr{P}$ . It remains to upper bound the offset hybrid DEC of  $\mathscr{P}$  by the hybrid DEC, and we invoke the following lemma. Its proof largely mimics Foster et al. [2023b] and is postponed to Appendix F.9.

**Lemma F.7.** Suppose that the loss function L is metric-based. Then it holds that

$$\inf_{\gamma>0} \left(\mathsf{p}\text{-}\mathsf{dec}_{\gamma}^{\mathrm{o},\mathrm{H}}(\mathscr{P}) + \gamma\varepsilon^2\right) \leq 8\varepsilon \cdot \sup_{\varepsilon' \in [\varepsilon,1]} \frac{\mathsf{p}\text{-}\mathsf{dec}_{\varepsilon'}^{\mathrm{H}}(\mathscr{P})}{\varepsilon'}.$$

In particular, under the assumption that the hybrid DEC of  $\mathscr{P}$  is of moderate decay with constant  $c_{\text{reg}}$  (Definition 3), we have

$$\inf_{\gamma>0} \left( \mathsf{p-dec}_{\gamma}^{\mathrm{o},\mathrm{H}}(\mathscr{P}) + \gamma \varepsilon^2 \right) \leq 10 c_{\mathrm{reg}} \mathsf{p-dec}_{\varepsilon}^{\mathrm{H}}(\mathscr{P}).$$

Hence, with an optimally tune parameter  $\gamma$ ,  $ExO^+$  (as instantiated in Theorem F.5) achieves

$$\mathbf{Risk}_{\mathsf{DM}}(T) \leq \Delta + \inf_{\gamma > 0} \left\{ \mathsf{p-dec}_{\gamma/4}^{\mathrm{o},\mathsf{H}}(\mathscr{P}) + \frac{2\gamma \log(|\mathscr{P}|/\delta)}{T} \right\}$$
$$\leq \Delta + 80c_{\mathrm{reg}} \cdot \mathsf{p-dec}_{\bar{\varepsilon}(T)}^{\mathsf{H}}(\mathscr{P}),$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log(|\mathscr{P}|/\delta)}{T}}$ .

#### F.4.2 Proof of Theorem 9

For no-regret learning, the most natural information set structure is given by  $\Psi := \Psi_{\mathscr{P},\Pi} = \mathscr{P} \times \Pi$ , such that for each  $\psi = (\mathcal{P}, \pi) \in \Psi$ , we may specify  $M_{\psi} = \mathcal{P}$  and  $\pi_{\psi} = \pi$ . With such a construction, it is direct to verify that  $\mathcal{M}_{\Psi} = \mathcal{M}_{\mathscr{P}}$  and

$$N_{\mathsf{frac}}(\mathscr{P}, \Psi; \Delta) \leq |\mathscr{P}| \cdot N_{\mathsf{frac}}(\mathcal{M}_{\mathscr{P}}, \Delta).$$

Therefore, it remains to upper bound the offset DEC of  $\mathcal{M}_{\mathscr{P}}$  with the constrained DEC of  $\mathcal{M}_{\mathscr{P}}$ . We invoke the following conversion result, which follows from Chen et al. [2024, Theorem G.5].

**Proposition F.8.** Suppose that Eq. (53) holds for the value function  $V^M$  over the model class  $\mathcal{M}$ . Then it holds that

$$\inf_{\gamma>0} \left( \mathsf{r-dec}_{\gamma}^{\mathrm{o}}(\mathcal{M}) + \gamma \varepsilon^2 \right) \leq C \sqrt{\log(1/\varepsilon)} \cdot \left( \sup_{\varepsilon' \in [\varepsilon,1]} \frac{\mathsf{r-dec}_{\varepsilon'}^{\mathrm{c}}(\mathcal{M})}{\varepsilon'} + C_V \right) \cdot \varepsilon,$$

where C is a universal constant.

Similarly, for PAC learning with loss function  $L(M,\pi) = V^M(\pi^M) - V^M(\pi)$ , it holds that

$$\inf_{\gamma>0} \left( \mathsf{p-dec}_{\gamma}^{\mathrm{o}}(\mathcal{M}) + \gamma \varepsilon^{2} \right) \leq C' \left( \sup_{\varepsilon' \in [\varepsilon, 1]} \frac{\mathsf{p-dec}_{\varepsilon'}^{\mathrm{c}}(\mathcal{M})}{\varepsilon'} + C_{V} \right) \cdot \varepsilon,$$

where C' is a universal constant.

Note that when the value function is reward-based (Example 1), Eq. (53) holds with  $C_V = \sqrt{2}$ .

Hence, under the assumptions of Theorem 10, if we instantiate  $ExO^+$  with the information set structure  $\Psi = \Psi_{\mathscr{P}}$ , option reg, and choose  $\gamma > 0$  optimally, then  $ExO^+$  achieves with probability at least  $1 - \delta$  that

$$\frac{1}{T} \operatorname{Reg}_{\mathsf{DM}}(T) \leq \Delta + \inf_{\gamma > 0} \left\{ \operatorname{r-dec}_{\gamma/4}^{o}(\mathcal{M}_{\Psi}) + \frac{2\gamma}{T} [\log N_{\mathsf{frac}}(\mathscr{P}, \Psi; \Delta) + \log(1/\delta)] \right\}$$
$$\leq \Delta + O\left(\sqrt{\log T}\right) \cdot \left[\operatorname{r-dec}_{\bar{\varepsilon}(T)}^{c}(\mathcal{M}_{\mathscr{P}}) + \bar{\varepsilon}(T)\right],$$

where  $\bar{\varepsilon}(T) = C\sqrt{\frac{\log(|\mathscr{P}|/\delta) + \log N_{\text{frac}}(\mathcal{M}_{\mathscr{P}}, \Delta)}{T}}$ . This gives Theorem 10 immediately.

## 

#### F.4.3 Proof of Proposition 11

As we have discussed in Section 3.3, for the convex hypothesis selection problem, we can consider the "relaxed" constraint class  $\mathscr{P}_m = \{\mathcal{M}_1, \cdots, \mathcal{M}_m\}$ . Then, by Theorem F.5, under any environment specified by a model  $M^* \in \mathcal{M}$ ,  $\mathsf{ExO}^+$  (when instantiated on  $\mathscr{P}_m$  and  $\gamma = \frac{4}{\overline{\varepsilon}(T)^2}$ ) achieves with probability at least  $1 - \delta$  that

$$\mathbb{E}_{\pi_{T+1} \sim \hat{p}} L(M^{\star}, \pi_{T+1}) \leq \mathsf{p-dec}_{\gamma/4}^{\mathrm{o}, \mathsf{H}}(\mathscr{P}_m) + \frac{2\gamma \log(m/\delta)}{T}.$$

Because  $\mathcal{M}_1, \cdots, \mathcal{M}_m$  are convex, we have

$$\operatorname{p-dec}_{\gamma/4}^{\mathrm{o},\mathrm{H}}(\mathscr{P}_m) = \operatorname{p-dec}_{\gamma/4}^{\mathrm{o}}(\mathcal{M}) \leq \operatorname{p-dec}_{2/\sqrt{\gamma}}^{\mathrm{c}}(\mathcal{M}) = \operatorname{p-dec}_{\bar{\varepsilon}(T)}^{\mathrm{c}}(\mathcal{M}),$$

where the inequality follows from Eq. (63). Thus, with probability at least  $1 - \delta$ , we have

$$\mathbb{E}_{\pi_{T+1}\sim\hat{p}}L(M^{\star},\pi_{T+1}) \leq \mathsf{p-dec}^{\mathsf{c}}_{\bar{\varepsilon}(T)}(\mathcal{M}) + \frac{8\log(m/\delta)}{\bar{\varepsilon}(T)^{2}T} \leq \frac{1}{3} + \frac{1}{8} < \frac{1}{2}.$$

Note that for  $M^* \in \mathcal{M}_{i^*}$ , we have  $L(M^*, \pi_{T+1}) = \mathbf{1} \{ \pi_{T+1} \neq i^* \}$ , and hence

$$\mathbb{E}_{\pi_{T+1}\sim\hat{p}}L(M^{\star},\pi_{T+1})=\mathbb{P}_{\pi\sim\hat{p}}(\pi\neq i^{\star}).$$

Therefore, we may modify  $\mathsf{ExO}^+$  so that it outputs  $\pi_{T+1} = \arg\min_{i \in [m]} \mathbb{P}_{\pi \sim \hat{p}}(\pi \neq i)$ . Then, with probability at least  $1 - \delta$ , we have  $\pi_{T+1} = i^*$ .

#### F.4.4 Proof of Proposition 13

Recall that the constraint class for stochastic DMSO is  $\mathscr{P}_{sto}$ . Therefore, fix the parameter  $\Delta \geq 0$ , we may consider the "policy-based" information set structure  $\Psi = \Pi$  (specified as in Eq. (24)):

$$\mathcal{M}_{\psi} = \{ M \in \mathcal{M} : V^{M}(\pi^{M}) - V^{M}(\pi) \le \Delta \}, \qquad \pi_{\psi} = \psi, \qquad \forall \psi \in \Psi.$$
(71)

With such an information set structure, it is clear that  $\mathcal{M}_{\Psi} = \bigcup_{\pi} \operatorname{co}(\mathcal{M}_{\pi}) = \mathcal{M}_{\Pi}$ , and

$$N_{\mathsf{frac}}(\mathscr{P}_{\mathsf{sto}}, \Psi; \Delta) = N_{\mathsf{frac}}(\mathcal{M}, \Delta).$$

Therefore, under the assumptions of Proposition 13, if we instantiate  $ExO^+$  with the information set structure  $\Psi = \Pi$  specified above, option reg, and choose  $\gamma > 0$  optimally, then  $ExO^+$  achieves with probability at least  $1 - \delta$  that

$$\begin{split} \frac{1}{T} \mathbf{Reg}_{\mathsf{DM}}(T) &\leq \Delta + \inf_{\gamma > 0} \left\{ \mathsf{r}\text{-}\mathsf{dec}^{\mathrm{o}}_{\gamma/4}(\mathcal{M}_{\Pi}) + \frac{2\gamma}{T} [\log N_{\mathsf{frac}}(\mathcal{M}, \Delta) + \log(1/\delta)] \right\} \\ &\leq \Delta + O\Big(\sqrt{\log T}\Big) \cdot \Big[\mathsf{r}\text{-}\mathsf{dec}^{\mathrm{c}}_{\bar{\varepsilon}(T)}(\mathcal{M}_{\Pi}) + \bar{\varepsilon}(T)\Big], \end{split}$$

where  $\bar{\varepsilon}(T) = \sqrt{\frac{\log N_{\text{frac}}(\mathcal{M}, \Delta) + \log(1/\delta)}{T}}$  and the second inequality follows from Proposition F.8. This gives the desired regret bound.

As a remark, for reward-based PAC learning, we can similarly obtain an upper bound scaling with the PAC-DEC of  $\mathcal{M}_{\Pi}$  and the fractional covering number.

# F.5 Instantiations of ExO<sup>+</sup> to LDP learning

In this section, we turn our focus to private DMSO (Section 2.3), where the learner is given a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ .

Let observation space be  $\mathcal{O} = \{-1, 1\}$  (i.e., only binary channels are considered), and let  $\Phi = \mathcal{Q}_{\alpha}$  be the class of all  $\alpha$ -DP channels from  $\mathcal{Z}$  to  $\{-1, 1\}$ . When  $\mathsf{ExO}^+$  is instantiated with the measurement class  $\Phi = \mathcal{Q}_{\alpha}$ , we will call the obtained algorithm LDP-ExO, because it naturally preserves  $\alpha$ -LDP.

Recall that in private DMSO, the corresponding constraint class is  $\mathscr{P}_{LDP} = \{\mathcal{M}^{\sharp} : M \in \mathcal{M}\}$ , where for each model  $M \in \mathcal{M}$ , the model  $M^{\sharp} : \Pi \times \Phi \to \Delta(\mathcal{O})$  is specified by  $M^{\sharp}(\pi, \mathsf{Q}) = \mathsf{Q} \circ M(\pi)$  and  $V^{M^{\sharp}}(\pi) = V^{M}(\pi)$  for all  $\pi \in \Pi$ ,  $\mathbb{Q} \in \Phi$ . For simplicity, we focus on the setting of rewardbased learning (Definition 9), where there is a reward function R such that the value function is given by  $V^{M}(\pi) = \mathbb{E}^{M,\pi}R(z,\pi)$ , and the loss function  $L(M,\pi) = V^{M}(\pi^{M}) - V^{M}(\pi)$ , where  $\pi^{M} = \arg \max_{\pi \in \Pi} V^{M}(\pi)$ .

For private DMSO with a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ , we restate the definition of information set structure structure as follows. Here, we focus on Type 1 information set structure, and (with slight abuse of notation) we regard  $\mathcal{M}$  as a subset of  $(\Pi \times \Phi \to \Delta(\mathcal{O}))$  by identifying each model  $M \in \mathcal{M}$  with  $M^{\sharp}$ .

**Definition 19** (Information set structure for private DMSO). Given a model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ , an information set structure is a class  $\Psi$ , where each  $\psi \in \Psi$  is associated with a model class  $\mathcal{M}_{\psi} \subseteq (\Pi \to \Delta(\mathcal{O}))$  and a decision  $\pi_{\psi} \in \Pi$ , such that for each  $M \in \mathcal{M}$ , there exists  $\psi \in \Psi$  such that  $M \in \mathcal{M}_{\psi}$ . We denote  $\mathcal{M}_{\Psi} := \bigcup_{\psi \in \Psi} \operatorname{co}(\mathcal{M}_{\psi})$ .

**Private offset-DECs.** To state the upper bounds of LDP-ExO with minimal assumptions, we introduce the offset private PAC-DEC/private regret-DEC as follows. For any model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  and a reference model  $\overline{M} \in \operatorname{co}(\mathcal{M})$ , we let

$$\mathsf{p-dec}_{\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi \times \mathcal{L})}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M,\pi)] - \gamma \mathbb{E}_{(\pi,\ell) \sim q} \mathsf{D}_{\ell}^{2}(M(\pi),\overline{M}(\pi)) \right\}, \tag{72}$$

$$\mathsf{r}\mathsf{-dec}_{\gamma}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M},\overline{M}) \coloneqq \inf_{p \in \Delta(\Pi \times \mathcal{L})} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[V^{M}(\pi^{M}) - V^{M}(\pi)] - \gamma \mathbb{E}_{(\pi,\ell) \sim q} \mathsf{D}_{\ell}^{2}(M(\pi),\overline{M}(\pi)) \right\},$$
(73)

and we define

$$\mathsf{p-dec}_{\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M}) = \sup_{\overline{M} \in \mathrm{co}(\mathcal{M})} \mathsf{p-dec}_{\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M},\overline{M}), \quad \mathsf{r-dec}_{\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M}) = \sup_{\overline{M} \in \mathrm{co}(\mathcal{M})} \mathsf{r-dec}_{\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M},\overline{M}).$$
(74)

By the data-processing inequality (Proposition 20), we can relate the offset private PAC-DEC (regret-DEC) of  $\mathcal{M}$  to the offset PAC-DEC (regret-DEC) of the induced model class  $\mathcal{M}^{\sharp}$ :

$$\mathsf{p}\mathsf{-dec}^{\mathrm{o},\mathsf{LDP}}_{c_0\alpha^2\gamma}(\mathcal{M}) \le \mathsf{p}\mathsf{-dec}^{\mathrm{o}}_{\gamma}(\mathcal{M}^{\sharp}) \le \mathsf{p}\mathsf{-dec}^{\mathrm{o},\mathsf{LDP}}_{c_1\alpha^2\gamma}(\mathcal{M}).$$
(75)

$$\operatorname{\mathsf{r-dec}}_{c_0\alpha^2\gamma}^{\mathrm{o,LDP}}(\mathcal{M}) \leq \operatorname{\mathsf{r-dec}}_{\gamma}^{\mathrm{o}}(\mathcal{M}^{\sharp}) \leq \operatorname{\mathsf{r-dec}}_{c_1\alpha^2\gamma}^{\mathrm{o,LDP}}(\mathcal{M}).$$
(76)

The proof is essentially the same as Appendix E.3.2 and hence omitted.

Guarantees of LDP-ExO. For simplicity, we denote (cf. Eq. (69))

$$N_{\mathsf{frac}}(\mathcal{M}, \Psi; \Delta) := N_{\mathsf{frac}}(\mathscr{P}_{\mathsf{LDP}}, \Psi; \Delta) = \inf_{w \in \Delta(\Psi)} \sup_{M \in \mathcal{M}} \frac{1}{\mathbb{P}_{\psi \sim w}(\psi : M \in \mathcal{M}_{\psi}, V^{M}(\pi^{M}) - V^{M}(\pi_{\psi}) \leq \Delta))}$$
(77)

With above notation, we state the guarantee of LDP-ExO as follows.

**Theorem F.9** (LDP-ExO for private DMSO). Let  $T \ge 1$ , parameter  $\gamma, \Delta > 0, \delta \in (0, 1)$ , model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  and value function V be given. Suppose that  $\Psi$  is an information set structure with respect to the model class  $\mathcal{M}$ , and  $\mathcal{M}_{\Psi}$  is compact. We instantiate  $\mathsf{ExO}^+$  on  $\Psi$  and choose  $w_1 \in \Delta(\Psi)$  according to Eq. (70). (1) With the option reg,  $ExO^+$  achieves with probability at least  $1 - \delta$ 

$$\begin{split} \mathbf{Reg}_{\mathsf{DM}}(T) &= \sum_{t=1}^{T} V^{M^{\star}}(\pi^{M^{\star}}) - \mathbb{E}_{\pi_{t} \sim q_{t}} \left[ V^{M^{\star}}(\pi_{t}) \right] \\ &\leq T \cdot \left[ \Delta + \mathsf{r-dec}_{c\alpha^{2}\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M}_{\Psi}) \right] + 2\gamma [\log N_{\mathsf{frac}}(\mathcal{M},\Psi;\Delta) + \log(1/\delta)]. \end{split}$$

(2) With the option pac,  $ExO^+$  achieves with probability at least  $1 - \delta$  that

$$\mathbb{E}_{\pi_{T+1}\sim\hat{p}}\left[V^{M^{\star}}(\pi^{M^{\star}}) - V^{M^{\star}}(\pi_{T+1})\right] \leq \Delta + \mathsf{p-dec}_{c\alpha^{2}\gamma}^{o}(\mathcal{M}_{\Psi}) + \frac{2\gamma}{T}[\log N_{\mathsf{frac}}(\mathcal{M},\Psi;\Delta) + \log(1/\delta)].$$

In the following, we provide detailed specifications of the information set structure and guarantees for various settings. To obtain upper bounds in private PAC-DEC (regret-DEC), we will frequently invoke the following conversion lemma.

**Proposition F.10.** Let  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  be a given model class. Then the following holds.

(1) No-regret learning: If the value function V is reward-based (Definition 9), then

$$\inf_{\gamma>0} \left( \mathsf{r-dec}_{\gamma}^{\mathrm{o},\mathrm{LDP}}(\mathcal{M}) + \gamma \varepsilon^2 \right) \leq C \sqrt{\log(1/\varepsilon)} \cdot \left( \sup_{\varepsilon' \in [\varepsilon,1]} \frac{\mathsf{r-dec}_{\varepsilon'}^{\mathrm{LDP}}(\mathcal{M})}{\varepsilon'} + 1 \right) \cdot \varepsilon,$$

where C is a universal constant.

(2) PAC learning: If the loss function is reward-based, then

$$\inf_{\gamma>0} \left( \mathsf{p-dec}_{\gamma}^{\mathrm{o},\mathrm{LDP}}(\mathcal{M}) + \gamma \varepsilon^2 \right) \leq C \left( \sup_{\varepsilon' \in [\varepsilon,1]} \frac{\mathsf{p-dec}_{\varepsilon'}^{\mathrm{LDP}}(\mathcal{M})}{\varepsilon'} + 1 \right) \cdot \varepsilon,$$

where C' is a universal constant.

Proposition F.10 follows immediately from Chen et al. [2024, Theorem E.7] (see also Foster et al. [2023b, Proposition 4.2]).

#### F.5.1 Model-based learning

Perhaps the most natural information set structure is the *model-based* information set structure  $\Psi_{mod}$ , given by

$$\Psi_{\text{mod}} = \mathcal{M}, \qquad \mathcal{M}_{\psi} = \{\psi\}, \qquad \forall \psi \in \Psi_{\text{mod}}, \tag{78}$$

i.e., each information set  $\psi \in \Psi_{\mathsf{mod}}$  corresponds to a model  $M \in \mathcal{M}$ .

By definition, we know that  $\mathcal{M}_{\Psi_{mod}} = \mathcal{M}$  and  $\log N_{frac}(\mathcal{M}, \Psi_{mod}; 0) = \log |\mathcal{M}|$ , achieving at the prior  $w_1 = \text{Unif}(\mathcal{M})$ .<sup>12</sup> We instantiate LDP-ExO on  $\Psi_{mod}$  to obtain the upper bound Eq. (34) in Theorem 28.

<sup>&</sup>lt;sup>12</sup>When  $\mathcal{M}$  is infinite, we may instead take  $\Psi$  to be a covering of  $\mathcal{M}$ , and our results still hold with  $\log |\mathcal{M}|$  replace by the logarithmic covering number.

**Proof of Theorem 28 (1).** Let  $\Delta = 0$  and LDP-ExO be instantiated on the information set structure  $\Psi_{mod}$ . Then, by Theorem F.9, LDP-ExO with an optimally-chosen parameter  $\gamma > 0$  achieves with probability at least  $1 - \delta$ 

$$\frac{1}{T}\mathbf{Reg}_{\mathsf{DM}}(T) \leq \inf_{\gamma > 0} \left( \mathsf{r-dec}_{c\alpha^2\gamma}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M}) + \frac{2\gamma \log(|\mathcal{M}|/\delta)}{T} \right) \leq O(\sqrt{\log T}) \cdot \Big[ \mathsf{r-dec}_{\bar{\varepsilon}(T)}^{\mathrm{LDP}}(\mathcal{M}) + \bar{\varepsilon}(T) \Big],$$

where the second inequality uses Proposition F.10 and the assumption that  $\mathsf{r-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M})$  is of moderate decay.

As a remark, we note that for reward-based PAC learning, the upper bound of Theorem 21 can also be obtained in this way.

## F.5.2 Policy-based learning

Following Chen et al. [2024] (see also Section 3.3), we consider the decision-based (or, "policybased") information set structure  $\Psi = \Psi_{pol}$  given by

$$\Psi_{\mathsf{pol}} = \Pi, \qquad \mathcal{M}_{\pi} = \{ M : V^{M}(\pi^{M}) - V^{M}(\pi) \le \Delta \}, \qquad \forall \pi \in \Pi.$$
(79)

By definition,

$$\mathcal{M}_{\Psi_{\mathsf{mod}}} = \bigcup_{\pi \in \Pi} \operatorname{co}(\mathcal{M}_{\pi}) = \mathcal{M}_{\Pi}, \qquad N_{\mathsf{frac}}(\mathcal{M}, \Psi_{\mathsf{pol}}; \Delta) = N_{\mathsf{frac}}(\mathcal{M}, \Delta).$$

Therefore, we may instantiate LDP-ExO with  $\Psi_{pol}$  to obtain the following upper bounds, which are direct implied by Theorem F.9.

**Proposition F.11** (Policy-based LDP-ExO for private PAC learning). Let  $T \ge 1$ ,  $\gamma > 0, \Delta > 0$ , model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$  be given. Suppose that  $\mathcal{M}$  is compact, and LDP-ExO is instantiated with the information set structure  $\Psi_{pol}$ . Then, the following holds.

(1) With option pac, it holds that with probability at least  $1 - \delta$ 

$$\mathbf{Risk}_{\mathsf{DM}}(T) = \mathbb{E}_{\pi_{T+1} \sim \hat{p}} L(M^{\star}, \pi_{T+1}) \leq \Delta + \mathsf{p-dec}_{c\alpha^{2}\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M}_{\Pi}) + \frac{2\gamma}{T} [\log N_{\mathsf{frac}}(\mathcal{M}, \Delta) + \log(1/\delta)].$$

(2) With option reg, it holds that with probability at least  $1 - \delta$ 

$$\frac{1}{T} \mathbf{Reg}_{\mathsf{DM}}(T) \leq \Delta + \mathsf{r-dec}_{c\alpha^2\gamma}^{\mathrm{o,LDP}}(\mathcal{M}_{\Pi}) + \frac{2\gamma}{T} [\log N_{\mathsf{frac}}(\mathcal{M}, \Delta) + \log(1/\delta)].$$

**Proof of Theorem 22.** Note that  $\mathcal{M}_{\Pi} \subseteq \operatorname{co}(\mathcal{M})$ . Thus, Theorem 22 follows immediately by choosing the optimal parameter  $\gamma > 0$  in the upper bound of Proposition F.11 (1) and then applying Proposition F.10.

**Proof of Theorem 28 (2).** Similarly, Eq. (35) of Theorem 28 follows immediately from Proposition F.11 (2) and Proposition F.10.

#### F.5.3 Value-based learning

For the well-specified regression task with a function class  $\mathcal{F} \subseteq (\mathcal{X} \to [-1, 1])$  (Section 5.2), we can employ the *value-based* information set structure  $\Psi_{\mathsf{val}} = \mathcal{F}$  (for a fixed parameter  $\Delta \geq 0$ ):

$$\mathcal{M}_f = \{ M : \mathbb{E}_{x \sim M} \left| f^M(x) - f(x) \right| \le \Delta \}, \qquad \pi_f = f, \qquad \forall f \in \Psi_{\mathsf{val}}.$$
(80)

This clearly gives a valid information set structure  $\Psi_{val}$ , and we have

$$N_{\mathsf{frac}}(\mathcal{M}_{\mathcal{F}}, \Psi_{\mathsf{val}}; \Delta) = N_{\mathsf{frac}}(\mathcal{F}, \Delta).$$

Therefore, we may instantiate LDP-ExO with such an information set structure, and it remains to upper bound the offset DEC of  $\mathcal{M}_{\Psi}$  as follows.

**Lemma F.12.** Suppose that  $\mathcal{M} = \mathcal{M}_{\mathcal{F}}$  is the class of well-specified models,  $\Psi = \Psi_{val}$ . Then it holds that

$$\mathsf{p}\text{-}\mathsf{dec}_{\gamma}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M}_{\Psi}) \leq \mathsf{p}\text{-}\mathsf{dec}_{\gamma/2}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M}) + \gamma\Delta^2 + 2\Delta.$$

**Proof of Proposition 24.** Let LDP-ExO be instantiated on the information set structure  $\Psi = \Psi_{\text{val}}$ . Then, by Theorem F.9, LDP-ExO with an optimally-chosen parameter  $\gamma > 0$  achieves with probability at least  $1 - \delta$ 

$$\begin{split} \mathbf{Risk}_{\mathsf{DM}}(T) &\leq \inf_{\gamma > 0} \left( \Delta + \mathsf{p-dec}_{c\alpha^2 \gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M}_{\Psi}) + \frac{2\gamma(\log N_{\mathsf{frac}}(\mathcal{F}, \Delta) + \log(1/\delta))}{T} \right) \\ &\leq \inf_{\gamma > 0} \left( 3\Delta + \mathsf{p-dec}_{c\alpha^2 \gamma/2}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M}) + \gamma\Delta^2 + \frac{2\gamma(\log N_{\mathsf{frac}}(\mathcal{F}, \Delta) + \log(1/\delta))}{T} \right) \\ &\leq O(1) \cdot \left( \mathsf{p-dec}_{\bar{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}) + \bar{\varepsilon}(T) \right). \end{split}$$

where the last inequality uses Proposition F.10 and the assumption that  $p-dec_{\varepsilon}^{LDP}(\mathcal{M})$  is of moderate decay.

**Proof of Lemma F.12.** By definition,  $\mathcal{M}_{\Psi} = \bigcup_{f \in \mathcal{F}} \operatorname{co}(\mathcal{M}_f)$ . We first prove the following claim.

**Claim.** For any  $f \in \mathcal{F}$  and  $\overline{M} \in co(\mathcal{M}_f)$ , there exists a model  $M' \in \mathcal{M}$  with  $f^{M'} = f$  and  $D_{\text{TV}}(M', \overline{M}) \leq \Delta$ .

Suppose  $\overline{M} \in co(\mathcal{M}_f)$  is given by  $\overline{M} = \mathbb{E}_{M \sim \mu}[M]$  with  $\mu \in \Delta(\mathcal{M}_f)$ . For each  $M \in \mathcal{M}$ , we denote  $\nu_M$  to be the distribution of x under M, and we denote  $\overline{\nu} = \mathbb{E}_{M \sim \mu} \nu_M$  to be the distribution of x under  $\overline{M}$ . Further, we know that under  $(x, y) \sim \overline{M}$ ,

$$y|x \sim \operatorname{Rad}(\bar{f}(x)), \text{ where } \bar{f}(x) = \mathbb{E}_{M|x} f^M(x),$$

where the conditional expectation is taken over  $M \sim \mu, x \sim \nu_M$ . Therefore, we have

$$\begin{aligned} \mathbb{E}_{x \sim \overline{\nu}} \left| f(x) - f(x) \right| &= \mathbb{E}_{x \sim \overline{\nu}} \left| \mathbb{E}_{M|x} f^{M}(x) - f(x) \right| \\ &\leq \mathbb{E}_{x \sim \overline{\nu}} \mathbb{E}_{M|x} \left| f^{M}(x) - f(x) \right| \\ &= \mathbb{E}_{M \sim \mu} \mathbb{E}_{x \sim \nu_{M}} \left| f^{M}(x) - f(x) \right| \leq \Delta \end{aligned}$$

Therefore, we can take M' to be the model with covariate distribution  $\overline{\nu}$  and  $f^{M'} = f$ , and we have

$$D_{\mathrm{TV}}\left(M',\overline{M}\right) = \mathbb{E}_{x \sim \overline{\nu}}\left|\overline{f}(x) - f(x)\right| \leq \Delta.$$

The proof of the claim is hence completed.

Now, with the above claim, for any reference model  $\overline{M} \in co(\mathcal{M})$ , we can bound

$$\begin{split} & \operatorname{\mathsf{p-dec}}_{\gamma}^{\operatorname{o},\operatorname{LDP}}(\mathcal{M}_{\Psi},\overline{M}) \\ &= \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi \times \mathcal{L})}} \sup_{M \in \mathcal{M}_{\Psi}} \left\{ \mathbb{E}_{\pi \sim p}[L(M,\pi)] - \gamma \mathbb{E}_{\ell \sim q} \mathsf{D}_{\ell}^{2}(M,\overline{M}) \right\} \\ &\leq \inf_{p \in \Delta(\Pi \times \mathcal{L})} \sup_{M \in \mathcal{M}_{\Psi}} \inf_{M' \in \mathcal{M}} \mathbb{E}_{\pi \sim p}[L(M',\pi)] + 2D_{\operatorname{TV}}\left(M',M\right) - \gamma \mathbb{E}_{\ell \sim q}\left[\frac{1}{2}\mathsf{D}_{\ell}^{2}(M',\overline{M}) - \mathsf{D}_{\ell}^{2}(M',M)\right] \\ &\leq \inf_{p \in \Delta(\Pi \times \mathcal{L})} \sup_{M' \in \mathcal{M}} \mathbb{E}_{\pi \sim p}[L(M',\pi)] + 2\Delta - \frac{\gamma}{2} \mathbb{E}_{\ell \sim q}\left[\mathsf{D}_{\ell}^{2}(M',\overline{M})\right] + \gamma \Delta^{2} \\ &= \operatorname{\mathsf{p-dec}}_{\gamma/2}^{\operatorname{o},\operatorname{LDP}}(\mathcal{M}) + 2\Delta + \gamma \Delta^{2}, \end{split}$$

where the second line follows from the fact that  $|L(M, \pi) - L(M', \pi)| \le 2D_{\text{TV}}(M', M)$  (because L is reward-based) and

$$\mathsf{D}^2_{\ell}(M',\overline{M}) \le 2\mathsf{D}^2_{\ell}(M,\overline{M}) + 2\mathsf{D}^2_{\ell}(M',M).$$

Taking supremum over  $\overline{M} \in co(\mathcal{M})$  gives the desired result.

## F.5.4 Contextual Bandits

In this section, we work with contextual DMSO (introduced in Section 5.5). Note that contextual DMSO is not encompassed by private DMSO, because the distribution of contexts can be changing throughout T rounds of interactions. However, the idea of Appendix F.5.3 can still be applied, and we frame it through the notation of Type 1 information set structure (with respect to the constraint class  $\mathscr{P}_{cxt}$ , defined in Eq. (36)).

We first recall the definition of the  $L_{\infty}$ -covering number.

**Definition 20.** For a function class  $\mathcal{F} \subseteq (\mathcal{X} \times \mathcal{A} \rightarrow [-1, 1])$  and parameter  $\Delta \geq 0$ , a  $\Delta$ -covering of  $\mathcal{F}$  is a subset  $\mathcal{F}' \subseteq \mathcal{F}$  such that for any  $f \in \mathcal{F}$ , there exists  $f' \in \mathcal{F}'$  with  $\sup_{x,a} |f(x, a) - f'(x, a)| \leq \Delta$ .

We define the  $\Delta$ -covering number of  $\mathcal{F}$  as  $N_{\infty}(\mathcal{F}, \Delta) := \inf \{ |\mathcal{F}'| : \mathcal{F}' \text{ is a } \Delta$ -covering of  $\mathcal{F} \}.$ 

Now, we define an Type 1 information set structure  $\Psi = \Psi_{\mathsf{cxt}}$  for the constraint class  $\mathscr{P}_{\mathsf{cxt}}$  by taking a minimal  $\Delta$ -covering  $\mathcal{F}_{\Delta}$  of  $\mathcal{F}$ , and let

$$\Psi_{\mathsf{cxt}} = \mathcal{F}_{\Delta}, \qquad \mathcal{M}_{\psi} = \left\{ M_{\nu,f} : \sup_{x,a} |f(x,a) - \psi(x,a)| \le \Delta \right\}, \qquad \forall \psi \in \Psi_{\mathsf{cxt}}, \tag{81}$$

and we set  $\pi_{\psi} \in \Pi$  be  $\pi_{\psi}(x) = \arg \max_{a \in \mathcal{A}} \psi(x, a)$ . Then by definition,  $\log N_{\mathsf{frac}}(\mathscr{P}_{\mathsf{cxt}}, \Psi_{\mathsf{cxt}}; 2\Delta) \leq \log |\Psi_{\mathsf{cxt}}| = \log N_{\infty}(\mathcal{F}, \Delta)$ .

**Proof of Proposition 29.** Similar to Lemma F.12, we can show that with  $\mathcal{M} = \mathcal{M}_{\mathcal{F},\mathsf{CB}}$  and  $\Psi = \Psi_{\mathsf{cxt}}$ ,

$$\mathrm{r\text{-}dec}_{\gamma}^{\mathrm{o},\mathrm{LDP}}(\mathcal{M}_{\Psi}) \leq \mathrm{r\text{-}dec}_{\gamma/2}^{\mathrm{o},\mathrm{LDP}}(\mathcal{M}) + \gamma\Delta^2 + 2\Delta.$$

Therefore, when LDP-ExO is instantiated with  $\Psi = \Psi_{cxt}$ , with probability at least  $1 - \delta$ ,

$$\frac{1}{T}\mathbf{Reg}_{\mathsf{DM}}(T) \le 4\Delta + \mathsf{r-dec}_{c\alpha^{2}\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M}) + 2\gamma \left[\Delta^{2} + \frac{\log N_{\infty}(\mathcal{F},\Delta) + \log(1/\delta)}{T}\right]$$

Taking a suitable  $\Delta \ge 0$  and  $\gamma > 0$  according to Proposition F.10 completes the proof of Proposition 29.

# F.6 Proof of Theorem F.1

We first invoke the following lemma, which requires careful analysis due to the adversarial nature (in particular,  $\mathcal{E}^{\star}_{\Delta}$  may depend on the full history). The proof of Lemma F.13 is deferred to the end of this section.

#### Lemma F.13. Denote

$$\operatorname{Err}(p,q,\xi;w,M^{\star},\psi) := -\log \mathbb{E}_{\boldsymbol{\pi}\sim q} \mathbb{E}_{o\sim M^{\star}(\boldsymbol{\pi})} \mathbb{E}_{\psi'\sim w} \left[ \exp\left(\xi(\psi';\boldsymbol{\pi},o) - \xi(\psi;\boldsymbol{\pi},o)\right) \right].$$

Then with probability at least  $1 - \delta$ , it holds that

$$\min_{\psi^{\star} \in \mathcal{E}_{\Delta}^{\star}} \sum_{t=1}^{I} \operatorname{Err}(p_t, q_t, \xi_t; w_t, M^t, \psi^{\star}) \le 2\log(1/w_1(\mathcal{E}_{\Delta}^{\star})) + 2\log(1/\delta)$$

Under the success event of Lemma F.13, there exists a  $\psi^{\star} \in \mathcal{E}^{\star}_{\Delta}$  such that

$$\sum_{t=1}^{T} \operatorname{Err}(p_t, q_t, \xi_t; w_t, M^t, \psi^*) \le 2 \log(1/w_1(\mathcal{E}^*_{\Delta})) + 2 \log(1/\delta)$$

Notice that  $\psi^* \in \mathcal{E}^*_{\Delta}$  implies  $M^1, \cdots, M^T \in \mathcal{M}_{\psi^*}$ , and  $V^{\overline{M}^*}(\pi_{\overline{M}^*}) - V^{\overline{M}^*}(\pi_{\psi^*}) \leq \Delta$ , and in particular,

$$\max_{\pi} \frac{1}{T} \sum_{t=1}^{T} \left( V^{M^t}(\pi) - V^{M^t}(\pi_{\psi^\star}) \right) \leq \Delta.$$

Hence,

$$\begin{split} &\max_{\pi \in \Pi} \sum_{t=1}^{T} V^{M^{t}}(\pi) - \mathbb{E}_{\pi_{t} \sim p_{t}} \left[ V^{M^{t}}(\pi_{t}) \right] \\ &\leq T\Delta + \sum_{t=1}^{T} \mathbb{E}_{\pi_{t} \sim p_{t}} \left[ V^{M^{t}}(\pi_{\psi^{\star}}) - V^{M^{t}}(\pi_{t}) \right] \\ &= T\Delta + \gamma \sum_{t=1}^{T} \operatorname{Err}(p_{t}, q_{t}, \xi_{t}; w_{t}, M^{t}, \psi^{\star}) \\ &+ \sum_{t=1}^{T} \underbrace{ \left[ \mathbb{E}_{\pi_{t} \sim p_{t}} \left[ V^{M^{t}}(\pi_{\psi^{\star}}) - V^{M^{t}}(\pi_{t}) \right] - \gamma \operatorname{Err}(p_{t}, q_{t}, \xi_{t}; w_{t}, M^{t}, \psi^{\star}) \right] }_{\leq \Gamma_{w_{t}, \gamma}(p_{t}, q_{t}, \xi_{t}; M^{t}, \psi^{\star})} \\ &\leq T\Delta + 2\gamma [\log(1/w_{1}(\mathcal{E}^{\star}_{\Delta})) + \log(1/\delta)] + \sum_{t=1}^{T} \Gamma_{w_{t}, \gamma}(p_{t}, q_{t}, \xi_{t}) \\ &\leq T(\Delta + \exp_{\gamma}(\Psi)) + 2\gamma [\log(1/w_{1}(\mathcal{E}^{\star}_{\Delta})) + \log(1/\delta)], \end{split}$$

where the second inequality uses  $M^t \in \mathcal{M}_{\psi^*}$ . This is the desired upper bound.

## F.6.1 Proof of Lemma F.13

For simplicity of presentation, we only consider the case where  $\Psi$  is countable. By definition,

$$w_t(\psi) = \frac{w_1(\psi) \exp\left(\sum_{s=1}^t \xi^s(\psi; \pi^s, o^s)\right)}{\sum_{\psi' \in \Psi} w_1(\psi') \exp\left(\sum_{s=1}^{t-1} \xi^s(\psi'; \pi^s, o^s)\right)}$$

and hence

$$\log \mathbb{E}_{\psi \sim w_t} [\exp\left(\xi_t(\psi; \boldsymbol{\pi}_t, o_t)\right)] = \log \mathbb{E}_{\psi \sim w_1} \exp\left(\sum_{s=1}^t \xi^s(\psi; \boldsymbol{\pi}^s, o^s)\right) - \log \mathbb{E}_{\psi \sim w_1} \exp\left(\sum_{s=1}^{t-1} \xi^s(\psi; \boldsymbol{\pi}^s, o^s)\right).$$

Therefore, taking summation over  $t = 1, \dots, T$ , we have

$$-\sum_{t=1}^{T} \log \mathbb{E}_{\psi \sim w_t} [\exp\left(\xi_t(\psi; \boldsymbol{\pi}_t, o_t)\right)] = -\log \mathbb{E}_{\psi \sim w_1} \left[\exp\left(\sum_{t=1}^{T} \xi_t(\psi; \boldsymbol{\pi}_t, o_t)\right)\right].$$
(82)

Thus, we define

$$X_t(\psi; \boldsymbol{\pi}_t, o_t) := -\xi_t(\psi; \boldsymbol{\pi}_t, o_t) + \log \mathbb{E}_{\psi \sim w_t}[\exp\left(\xi_t(\psi; \boldsymbol{\pi}_t, o_t)\right)]$$

and Eq. (82) implies (deterministically)

$$\mathbb{E}_{\psi \sim w_1} \exp\left(-\sum_{t=1}^T X_t(\psi; \boldsymbol{\pi}_t, o_t)\right) = 1.$$

Notice that for any  $\psi \in \Psi$ , we also have

$$\mathbb{E}^{\mathsf{ExO}^+} \exp\left(\sum_{t=1}^T X_t(\psi; \boldsymbol{\pi}_t, o_t) - \log \mathbb{E}_{t-1}[\exp\left(X_t(\psi; \boldsymbol{\pi}_t, o_t)\right)]\right) = 1,$$

where the expectation  $\mathbb{E}^{E \times O^+}$  is taken over the randomness of the interaction between  $E \times O^+$  algorithm and the environment.

Further, by the definition of  $X_t$  and Err, it holds that for any fixed  $\psi,$ 

$$-\log \mathbb{E}_{t-1}[\exp \left(X_t(\psi; \boldsymbol{\pi}_t, o_t)\right)] = \operatorname{Err}(p_t, q_t, \xi_t; w_t, M^t, \psi).$$

Combining the equations above and applying Cauchy inequality, we now have

$$\mathbb{E}_{\psi \sim w_1} \mathbb{E}^{\mathsf{E} \mathsf{x} \mathsf{O}^+} \exp\left(\frac{1}{2} \sum_{t=1}^T \operatorname{Err}(p_t, q_t, \xi_t; w_t, M^t, \psi)\right) \le 1.$$

Notice that  $\psi \sim w_1$  is independent of the randomness of the *T*-round interactions under  $\mathsf{ExO}^+$ . Therefore, we know

$$\mathbb{E}^{\mathsf{E} \times \mathsf{O}^+} \left[ w_1(\mathcal{E}^{\star}_{\Delta}) \exp\left(\frac{1}{2} \min_{\psi^{\star} \in \mathcal{E}^{\star}_{\Delta}} \sum_{t=1}^T \operatorname{Err}(p_t, q_t, \xi_t; w_t, M^t, \psi^{\star})\right) \right] \le 1.$$

Applying Markov's inequality completes the proof.

# F.7 Proof of Theorem F.2

We follow the notations of Appendix F.6, and the proof is essentially analogous.

For Type 2 information set structure  $\Psi = \mathscr{P}$ , there exists  $\psi^* \in \Psi$  such that  $\mathcal{P}^* = \mathcal{M}_{\psi^*}$ . Then, a direct adaption of Lemma F.13 yields

$$\sum_{t=1}^{T} \operatorname{Err}(p_t, q_t, \xi_t; w_t, M^t, \psi^\star) \le 2 \log |\mathscr{P}| + 2 \log(1/\delta),$$

as  $w^1 = \text{Unif}(\mathscr{P})$ .

Therefore,

$$\begin{split} \mathbf{Risk}_{\mathsf{DM}}(T) &= \mathbb{E}_{\pi_{T+1} \sim \widehat{p}} L(\mathcal{P}^{\star}, \pi) \\ &= \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}_{\pi_t \sim p_t} [L(\mathcal{P}^{\star}, \pi_t)] \\ &= \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}_{\pi_t \sim p_t} \left[ L_{\psi^{\star}}(M^t, \pi_t) \right] \\ &= \frac{\gamma}{T} \sum_{t=1}^{T} \operatorname{Err}(p_t, q_t, \xi_t; w_t, M^t, \psi^{\star}) \\ &\quad + \frac{1}{T} \sum_{t=1}^{T} \underbrace{\left[ \mathbb{E}_{\pi_t \sim p_t} \left[ L_{\psi^{\star}}(M^t, \pi_t) \right] - \gamma \operatorname{Err}(p_t, q_t, \xi_t; w_t, M^t, \psi^{\star}) \right]}_{\leq \Gamma_{w_t, \gamma}(p_t, q_t, \xi_t; M^t, \psi^{\star})} \\ &\leq \frac{2\gamma}{T} [\log |\mathscr{P}| + \log(1/\delta)] + \frac{1}{T} \sum_{t=1}^{T} \Gamma_{w_t, \gamma}(p_t, q_t, \xi_t) \\ &\leq \operatorname{exo}_{\gamma}(\Psi) + \frac{2\gamma}{T} [\log |\mathscr{P}| + \log(1/\delta)], \end{split}$$

where the first inequality uses the fact that  $M^t \in \mathcal{P}^* = \mathcal{M}_{\psi^*}$ . This is the desired result.

# F.8 Proof of Theorem F.3

The analysis below essentially follows the ideas of Foster et al. [2022b].

Let  $\mathcal{M}_0 := \bigcup_{\psi \in \Psi} \mathcal{M}_{\psi}$ . Below, we prove Theorem F.3 for finite  $\mathcal{M}_0$  (and  $\Psi$  is then automatically finite). The result for the general case then follows immediately by a covering argument (for details, see Remark F.16).

To proceed, we fix  $\gamma \geq 0, w \in \Delta(\Psi)$  and denote

$$\mathcal{I} := \{ (M, \psi) : \psi \in \Psi, M \in \mathcal{M}_{\psi} \}.$$

We also fix a parameter A > 0, and we define

$$\Xi_A := \{\xi \in \Xi : \|\xi\|_\infty \le A\}$$

Then,

$$\exp_{\gamma}(\Psi, w) \leq \inf_{(p,q) \in \mathbb{S}, \xi \in \Xi_A} \sup_{(M,\psi) \in \mathcal{I}} \Gamma_{w,\gamma}(p,q,\xi;M,\psi)$$

$$= \inf_{(p,q)\in\mathbb{S},\xi\in\Xi_A} \sup_{\mu\in\Delta(\mathcal{I})} \mathbb{E}_{(M,\psi)\sim\mu}\Gamma_{w,\gamma}(p,q,\xi;M,\psi).$$

To proceed, we apply Ky Fan's minimax theorem [Fan, 1953] (Theorem D.2). Note that  $\Delta(\mathcal{I})$  is a compact and convex subset of the Euclidean space  $\mathbb{R}^{\mathcal{I}}$  (because  $\mathcal{I}$  is finite), and  $\Xi$  is a vector space. Thus, we consider the following function

$$F_{p,q}(\xi,\mu) := \mathbb{E}_{(M,\psi)\sim\mu}\Gamma_{w,\gamma}(p,q,\xi;M,\psi),$$

and by definition,  $F_{p,q}$  is a bilinear function, and for any fixed  $\xi \in \Xi_A$ ,  $F_{p,q}(\xi, \cdot)$  is a concave, continuous function of  $\mu \in \Delta(\mathcal{I})$  (the continuity follows from the fact that  $\xi$  is uniformly bounded by A). Therefore, Ky Fan's minimax theorem (Theorem D.2) gives

$$\inf_{\xi \in \Xi_A} \max_{\mu \in \Delta(\mathcal{I})} F_{p,q}(\xi,\mu) = \max_{\mu \in \Delta(\mathcal{I})} \inf_{\xi \in \Xi_A} F_{p,q}(\xi,\mu).$$

Next, we compute  $G(p,q;\mu) := \inf_{\xi \in \Xi_A} F_{p,q}(\xi,\mu)$ . It is equivalent to compute

$$G_0(p,q;\mu) := \inf_{\xi \in \Xi_A} \mathbb{E}_{\boldsymbol{\pi} \sim q} \mathbb{E}_{o \sim M(\boldsymbol{\pi})} \mathbb{E}_{\psi' \sim w} \Big[ \exp\left(\xi(\psi';\boldsymbol{\pi},o) - \xi(\psi;\boldsymbol{\pi},o)\right) \Big].$$
(83)

For any  $\mu \in \Delta(\mathcal{I})$  and  $\pi \in \mathbf{\Pi}$ , we define  $\mathbb{P}_{\mu,\pi}$  to be the distribution of  $(\pi, o, \psi)$  generated by  $(M, \psi) \sim \mu, \pi \sim q, o \sim M(\pi)$ . Then, by Lemma F.15,

$$G_{0}(p,q;\mu) = \inf_{\xi \in \Xi_{A}} \mathbb{E}_{\pi \sim q, o \sim \mathbb{P}_{\mu,\pi}} \left\{ \mathbb{E}_{\psi \sim \mathbb{P}_{\mu,\pi}(\cdot|o)} [\exp\left(-\xi(\psi;\pi,o)\right)] \cdot \mathbb{E}_{\psi' \sim w} \left[\exp\left(\xi(\psi';\pi,o)\right)\right] \right\}$$
$$\leq \mathbb{E}_{\pi \sim q, o \sim \mathbb{P}_{\mu,\pi}} \left(1 - D_{\mathrm{H}}^{2} \left(\mathbb{P}_{\mu,\pi}(\cdot|o),w\right)\right)^{2} + 3e^{-A},$$

and hence

$$G(p,q;\mu) \leq \mathbb{E}_{(M,\psi)\sim\mu,\pi\sim p}[L_{\psi}(M,\pi)] + \gamma \mathbb{E}_{\pi\sim q,o\sim\mathbb{P}_{\mu,\pi}} \left(1 - D_{\mathrm{H}}^{2}\left(\mathbb{P}_{\mu,\pi}(\cdot|o),w\right)\right)^{2} - \gamma + 3e^{-A}.$$

Notice that for any fixed  $(p,q) \in S$ ,  $G(p,q;\mu)$  is a convex, continuous function of  $\mu$  (by definition). For any fixed  $\mu \in \Delta(\mathcal{I})$ ,  $G(p,q;\mu)$  is a linear function of  $(p,q) \in S$  and hence convex-like. Therefore, applying Ky Fan's minimax theorem (Theorem D.2) again gives

$$\inf_{(p,q)\in\mathbb{S}}\max_{\mu\in\Delta(\mathcal{I})}G(p,q;\mu)=\max_{\mu\in\Delta(\mathcal{I})}\inf_{(p,q)\in\mathbb{S}}G(p,q;\mu).$$

Finally, we proceed to bound  $G(p,q;\mu)$ . Using the fact that  $1-(1-x)^2 \ge x$  for  $x \in [0,1]$ , we have

$$G(p,q;\mu) \leq \mathbb{E}_{(M,\psi)\sim\mu,\pi\sim p}[L_{\psi}(M,\pi)] - \gamma \mathbb{E}_{\pi\sim q,o\sim\mathbb{P}_{\mu,\pi}}D_{\mathrm{H}}^{2}\left(\mathbb{P}_{\mu,\pi}(\cdot|o),w\right)$$

We then invoke the following lemma:

**Lemma F.14.** For any  $\mu \in \Delta(\mathcal{I})$  and  $\psi \in \Psi$ , we denote

$$M_{\psi|\mu} = \mathbb{E}_{M \sim \mu(\cdot|\psi)}[M] \in \operatorname{co}(\mathcal{M}_{\psi}), \qquad \overline{M}_{\mu} = \mathbb{E}_{M \sim \mu}[M] \in \operatorname{co}(\mathcal{M}_{0}).$$

Then it holds that for any  $\pi \in \Pi$ ,  $w \in \Delta(\Psi)$ ,

$$4 \cdot \mathbb{E}_{o \sim \mathbb{P}_{\mu, \pi}} D_{\mathrm{H}}^{2} \left( \mathbb{P}_{\mu, \pi}(\cdot | o), w \right) \geq \mathbb{E}_{\psi \sim \mu} D_{\mathrm{H}}^{2} \left( M_{\psi | \mu}(\pi), \overline{M}_{\mu}(\pi) \right)$$

Therefore, using Lemma F.14 and the fact that  $L_{\psi}(M,\pi)$  is affine over M, it holds that

$$G(p,q;\mu) \leq \mathbb{E}_{\psi \sim \mu, \pi \sim p} \left[ L_{\psi}(M_{\psi|\mu},\pi) \right] + 3e^{-A} - \frac{\gamma}{4} \mathbb{E}_{\psi \sim \mu, \pi \sim q} D_{\mathrm{H}}^{2} \left( M_{\psi|\mu}(\pi), \overline{M}_{\mu}(\pi) \right)$$

Hence, we have

$$\max_{\mu \in \Delta(\mathcal{I})} \inf_{(p,q) \in \mathbb{S}} G(p,q;\mu) - 3e^{-A} \\
\leq \max_{\mu \in \Delta(\mathcal{I})} \inf_{(p,q) \in \mathbb{S}} \mathbb{E}_{\psi \sim \mu, \pi \sim p} \left[ L_{\psi}(M_{\psi|\mu}, \pi) \right] - \frac{\gamma}{4} \mathbb{E}_{\psi \sim \mu, \pi \sim q} D_{\mathrm{H}}^{2} \left( M_{\psi|\mu}(\pi), \overline{M}_{\mu}(\pi) \right) \\
\leq \max_{\overline{M} \in \operatorname{co}(\mathcal{M}_{0})} \max_{\mu' \in \Delta(\mathcal{M}_{\Psi})} \inf_{(p,q) \in \mathbb{S}} \mathbb{E}_{M' \sim \mu', \pi \sim p} \left[ L_{\psi}(M', \pi) \right] - \frac{\gamma}{4} \mathbb{E}_{M' \sim \mu', \pi \sim q} D_{\mathrm{H}}^{2} \left( M'(\pi), \overline{M}(\pi) \right) \\
\leq \max_{\overline{M} \in \operatorname{co}(\mathcal{M}_{0})} \inf_{(p,q) \in \mathbb{S}} \max_{(M,\psi): M \in \mathcal{M}_{\psi}} \mathbb{E}_{\pi \sim p} \left[ L_{\psi}(M, \pi) \right] - \frac{\gamma}{4} \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^{2} \left( M(\pi), \overline{M}(\pi) \right),$$

where the last line follows again from the weak duality.

To finalize the proof, we notice that by the arbitrariness of  $w \in \Delta(\Psi)$ , we have already proven

$$\exp_{\gamma}(\Psi) \leq 3e^{-A} + \max_{\overline{M} \in \operatorname{co}(\mathcal{M}_0)} \inf_{(p,q) \in \mathbb{S}} \max_{(M,\psi): M \in \mathcal{M}_{\psi}} \mathbb{E}_{\pi \sim p}[L_{\psi}(M,\pi)] - \frac{\gamma}{4} \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^2\left(M(\pi), \overline{M}(\pi)\right).$$
(84)

Then, taking  $A \to +\infty$ , we obtain the following results (note that  $co(\mathcal{M}_0) = co(\mathcal{M}_{\mathscr{P}}) = \mathcal{M}^+$ ): (1) If  $\Psi$  is Type 1 information set structure, we have  $L_{\psi}(M, \pi) \leq L(M, \pi)$ . Hence, with option pac,  $\mathbb{S} = \mathbb{S}^{\mathsf{pac}} = \Delta(\Pi) \times \Delta(\Pi)$ , and hence in this case

$$\exp_{\gamma}(\Psi) \leq \operatorname{p-dec}_{\gamma/4}^{\mathrm{o}}(\mathcal{M}_{\Psi}).$$

Similarly, with option reg,  $\mathbb{S} = \mathbb{S}^{\mathsf{reg}} = \{(p|_{\Pi}, p) : p \in \Delta(\Pi)\}$ , and hence

$$\exp_{\gamma}(\Psi) \leq \operatorname{r-dec}_{\gamma/4}^{\mathrm{o}}(\mathcal{M}_{\Psi}).$$

(2) If  $\Psi = \mathscr{P}$  is the Type 2 information set structure, we have  $L_{\psi}(M, \pi) = L(\psi, \pi)$ , and hence

$$\mathrm{exo}_\gamma(\Psi) \leq \mathrm{p\text{-}dec}_{\gamma/4}^{\mathrm{o},\mathrm{H}}(\mathscr{P}).$$

**Proof of Lemma F.14.** Our proof essentially follows Foster et al. [2022b, Appendix C.2]. For simplicity of presentation, we abbreviate  $\mathbb{P} = \mathbb{P}_{\mu,\pi}$ . By the convexity of the squared Hellinger distance, we have

$$\mathbb{E}_{o\sim\mathbb{P}}D_{\mathrm{H}}^2\left(\mathbb{P}(\psi=\cdot|o),w\right) \ge D_{\mathrm{H}}^2\left(\mathbb{P}(\psi=\cdot),w\right).$$

Therefore, using the triangle inequality,

$$\begin{split} 4\mathbb{E}_{o\sim\mathbb{P}}D_{\mathrm{H}}^{2}\left(\mathbb{P}(\cdot|o),w\right) &\geq \mathbb{E}_{o\sim\mathbb{P}}\left[2D_{\mathrm{H}}^{2}\left(\mathbb{P}(\psi=\cdot|o),w\right) + 2D_{\mathrm{H}}^{2}\left(\mathbb{P}(\psi=\cdot),w\right)\right] \\ &\geq \mathbb{E}_{o\sim\mathbb{P}}D_{\mathrm{H}}^{2}\left(\mathbb{P}(\psi=\cdot|o),\mathbb{P}(\psi=\cdot)\right) \\ &= \mathbb{E}_{\psi\sim\mathbb{P}}D_{\mathrm{H}}^{2}\left(\mathbb{P}(o=\cdot|\psi),\mathbb{P}(o=\cdot)\right), \end{split}$$

where the last equality is because squared Hellinger distance is a f-divergence.

Recall that  $\mathbb{P} = \mathbb{P}_{\mu,\pi}$  generated  $(\psi, o)$  as  $(M, \psi) \sim \mu, o \sim M(\pi)$ . Therefore, for any  $\psi$ ,  $\mathbb{P}(o = \cdot | \psi)$  is the distribution of o generated as  $M \sim \mu(\cdot | \psi), o \sim M(\pi)$ , i.e.,  $o \sim M_{\psi|\mu}(\pi)$ . Hence,

$$D_{\mathrm{H}}^{2}\left(\mathbb{P}(o=\cdot|\psi),\mathbb{P}(o=\cdot)\right) = \mathbb{E}_{\boldsymbol{\pi}\sim q} D_{\mathrm{H}}^{2}\left(M_{\psi|\mu}(\boldsymbol{\pi}),\overline{M}_{\mu}(\boldsymbol{\pi})\right).$$

Combining the equations above completes the proof.

**Lemma F.15.** For any distribution  $\mathbb{P}, \mathbb{Q} \in \Delta(\Psi)$ , we denote

$$I_{\rm B}(\mathbb{P},\mathbb{Q}) = 1 - D_{\rm H}^2(\mathbb{P},\mathbb{Q}) = \sum_{\psi \in \Psi} \sqrt{\mathbb{P}(\psi)\mathbb{Q}(\psi)}.$$
(85)

Then for A > 0, it holds that

$$I_{\mathcal{B}}(\mathbb{P},\mathbb{Q})^{2} \leq \inf_{f \in (\Psi \to \mathbb{R}): \|f\|_{\infty} \leq A} \mathbb{E}_{\mathbb{P}}[e^{f(x)}] \mathbb{E}_{\mathbb{Q}}[e^{-f(x)}] \leq I_{\mathcal{B}}(\mathbb{P},\mathbb{Q})^{2} + 3e^{-A}.$$

*Proof.* The lower bound follows immediately from Cauchy inequality. In the following, we proceed to prove the upper bound.

Consider the function  $f = f_{\mathbb{P};\mathbb{Q}}$  given by

$$f_{\mathbb{P};\mathbb{Q}}(\psi) = \frac{1}{2} \mathsf{clip}_{[-A,A]}\left(\log \frac{\mathbb{Q}(\psi)}{\mathbb{P}(\psi)}\right).$$

Then, by definition,

$$\mathbb{E}_{\mathbb{P}}[e^{f(x)}] \le \mathbb{E}_{\mathbb{P}} \exp\left(\frac{1}{2}\log\frac{\mathbb{Q}(\psi)}{\mathbb{P}(\psi)}\right) + e^{-A} = \sum_{\psi \in \Psi} \sqrt{\mathbb{P}(\psi)\mathbb{Q}(\psi)} + e^{-A},$$

and similarly,

$$\mathbb{E}_{\mathbb{Q}}[e^{f(x)}] \le \mathbb{E}_{\mathbb{Q}} \exp\left(\frac{1}{2}\log\frac{\mathbb{P}(\psi)}{\mathbb{Q}(\psi)}\right) + e^{-A} = \sum_{\psi \in \Psi} \sqrt{\mathbb{P}(\psi)\mathbb{Q}(\psi)} + e^{-A}.$$

Therefore, for such a choice of f ensures

$$\mathbb{E}_{\mathbb{P}}[e^{f(x)}]\mathbb{E}_{\mathbb{Q}}[e^{-f(x)}] \le \left(I_{\mathrm{B}}(\mathbb{P},\mathbb{Q}) + e^{-A}\right)^2 \le I_{\mathrm{B}}(\mathbb{P},\mathbb{Q})^2 + 3e^{-A},$$

where the last inequality uses  $I_{\mathrm{B}}(\mathbb{P}, \mathbb{Q}) \leq 1$ .

**Remark F.16** (Covering argument). In the following, we briefly discuss how our analysis applies to an infinite  $\mathcal{M}_0$  with a covering argument. It is easy to deal with Type 1 information set structure, so we focus on Type 1 information set structure.

Fix a parameter  $\varepsilon \in (0, 1]$ . We take a finite subset  $\mathcal{M}' \subseteq \mathcal{M}_0$ , so that for any  $M \in \mathcal{M}_0$ , there exists  $M' \in \mathcal{M}'$ , such that

$$\sup_{\boldsymbol{\pi}} D_{\mathrm{TV}}\left(M(\boldsymbol{\pi}), M'(\boldsymbol{\pi})\right) \leq \varepsilon, \qquad \sup_{\boldsymbol{\pi}} \left|V^{\scriptscriptstyle M}(\boldsymbol{\pi}) - V^{\scriptscriptstyle M'}(\boldsymbol{\pi})\right| \leq \varepsilon.$$

Then,  $\Psi$  induces an information set structure over  $\mathcal{M}'$ , given by

$$\mathcal{M}'_{\psi} := \{ M \in \mathcal{M}' : M \in \mathcal{M}_{\psi} \}, \qquad \forall \psi \in \Psi.$$

Because  $\mathcal{M}'$  is finite, the set  $\{\mathcal{M}'_{\psi} : \psi \in \Psi\}$  is also finite. Therefore, there exists a finite subset  $\Psi' \subseteq \Psi$ , such that for any  $\psi \in \Psi, M \in \mathcal{M}_{\psi}$ , there exists  $[\psi] \in \Psi', M' \in \mathcal{M}'_{[\psi]}$ , so that

$$\sup_{\boldsymbol{\pi}} D_{\mathrm{TV}}\left(M(\boldsymbol{\pi}), M'(\boldsymbol{\pi})\right) \leq \varepsilon, \qquad \sup_{\boldsymbol{\pi}} \left| V^{\scriptscriptstyle M}(\boldsymbol{\pi}) - V^{\scriptscriptstyle M'}(\boldsymbol{\pi}) \right| \leq \varepsilon, \qquad V^{\scriptscriptstyle M}(\boldsymbol{\pi}_{\psi}) \leq V^{\scriptscriptstyle M'}(\boldsymbol{\pi}_{[\psi]}) + 2\varepsilon.$$

Then, we can bound

$$\begin{split} \exp_{\gamma}(\Psi, w) &\leq \inf_{(p,q) \in \mathbb{S}, \xi \in \Xi_{A}} \sup_{\psi \in \Psi, M \in \mathcal{M}_{\psi}} \Gamma_{w,\gamma}(p,q,\xi;M,\psi) \\ &\leq \inf_{(p,q) \in \mathbb{S}, \xi' \in \Xi'_{A}} \sup_{\psi \in \Psi', M \in \mathcal{M}'_{\psi}} \Gamma_{w',\gamma}(p,q,\xi';M,\psi) + (2+2\gamma+e^{2A})\varepsilon, \end{split}$$

where we let  $w' \in \Delta(\Psi')$  to be given by  $w'(\psi') = w(\psi : [\psi] = \psi')$  for all  $\psi' \in \Psi'$ , and the second inequality because for any map  $\xi' \in \Xi'_A := (\Psi \times \Pi \times \mathcal{O} \to [-A, A])$ , we can consider the induced map  $\xi \in \Xi_A$  given by  $\xi(\psi; \pi, o) = \xi'([\psi]; \pi, o)$  for any  $\psi \in \Psi$ .

Using Eq. (84), we have for PAC learning,

$$\exp_{\gamma}(\Psi) \leq \mathsf{p-dec}^{\mathrm{o}}_{\gamma/4}(\mathcal{M}'_{\Psi'}) + 3e^{-A} + (2+2\gamma+e^{2A})\varepsilon.$$

Note that  $\mathbf{p}-\mathbf{dec}_{\gamma/4}^{\mathbf{o}}(\mathcal{M}'_{\Psi'}) \leq \mathbf{p}-\mathbf{dec}_{\gamma/4}^{\mathbf{o}}(\mathcal{M}_{\Psi})$ , and hence first letting  $\varepsilon \to 0$  and then letting  $A \to \infty$  gives the desired result. A similar argument also applies to no-regret learning.

## F.9 Proof of Lemma F.7

Denote  $D := \sup_{\varepsilon' \in [\varepsilon, 1]} \frac{\mathsf{p-dec}_{\varepsilon'}^{\mathsf{H}}(\mathscr{P})}{\varepsilon'}$ . We consider  $\gamma = \frac{6D}{\varepsilon}$ .

We fix an arbitrary reference model  $\overline{M}$ . For each  $j \geq 0$ , we define  $\varepsilon_j = 2^{-j}$ , and let  $d_j := p-\mathsf{dec}_{\varepsilon_j}^{\mathsf{H}}(\mathscr{P}, \overline{M})$ ,

$$(p_j, q_j) := \underset{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}}{\operatorname{arg\,min}} \sup_{\mathcal{P} \in \mathscr{P}} \left\{ \mathbb{E}_{\pi \sim p} L(\mathcal{P}, \pi) \mid \inf_{M \in \operatorname{co}(\mathcal{P})} \mathbb{E}_{\pi \sim q} D_{\mathrm{H}}^2 \left( M(\pi), \overline{M}(\pi) \right) \leq \varepsilon_j^2 \right\}.$$

We define

$$\mathscr{P}_j := \left\{ \mathcal{P} : \inf_{M \in \operatorname{co}(\mathcal{P})} \mathbb{E}_{\pi \sim q_j} D_{\mathrm{H}}^2 \left( M(\pi), \overline{M}(\pi) \right) \le \varepsilon_j^2 \right\}.$$

We first claim that if for j < k,  $\mathscr{P}_j \cap \mathscr{P}_k$  is empty, then  $\mathscr{P}_{k+1}$  is empty. This is because if  $\mathscr{P}_j \cap \mathscr{P}_k = \emptyset$ , then for  $q = \frac{q_j + q_k}{2}$ , the set

$$\left\{ \mathcal{P} : \inf_{M \in \operatorname{co}(\mathcal{P})} \mathbb{E}_{\boldsymbol{\pi} \sim q} D_{\mathrm{H}}^2 \left( M(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi}) \right) \le \varepsilon_{k+1}^2 \right\}$$

must be empty, which certifies  $\mathbf{p}-\mathsf{dec}_{\varepsilon_{k+1}}^{\mathsf{H}}(\mathscr{P},\overline{M}) = -\infty$ , and hence by the optimality of  $(p_{k+1},q_{k+1})$ ,  $\mathscr{P}_{k+1}$  must be empty.

Therefore, we define  $K_0$  be the minimum integer k such that  $\mathbf{p}-\mathbf{dec}_{\varepsilon_k}^{\mathsf{H}}(\mathscr{P},\overline{M}) = -\infty$  (if such k does not exist, we write  $K_0 = \infty$ ). We further define  $K = \min\{\lfloor \log_2(1/\varepsilon) \rfloor, K_0 - 2\}$ .

For every  $j \leq K$ , by definition, for any  $\mathcal{P} \in \mathscr{P}_j$ , we have  $\mathbb{E}_{\pi \sim p_j} \rho(\pi^{\mathcal{P}}, \pi) \leq d_j$ . Thus, we can take  $\pi_j = \arg \min_{\pi' \in \Pi} \mathbb{E}_{\pi \sim p_j} \rho(\pi', \pi)$ , and then for any  $\mathcal{P} \in \mathscr{P}_j$ , we have  $\rho(\pi^{\mathcal{P}}, \pi_j) \leq 2d_j$ . Further, because  $\mathscr{P}_{K+1}$  is not empty,  $\mathscr{P}_j \cap \mathscr{P}_K$  is also not empty, and hence  $\rho(\pi_j, \pi_K) \leq 2d_j + 2d_K$ .

In the following, we choose  $\lambda_j = 2^{j-K-1}$  for  $j = 1, 2, \dots, K$  and  $\lambda_0 = 2^{-K}$ , and we set  $q = \sum_{j=0}^{K} \lambda_j q_j$ . For any  $\mathcal{P} \in \mathscr{P}$ , we proceed to bound the quantity

$$F(P) := \rho(\pi^{\mathcal{P}}, \pi_K) - \inf_{M \in \operatorname{co}(\mathcal{P})} \mathbb{E}_{\boldsymbol{\pi} \sim q} D_{\mathrm{H}}^2(M(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi})).$$

We let  $j \ge 0$  to be the largest integer such that  $\mathcal{P} \in \mathscr{P}_j$  (note that  $\mathcal{P} \in \mathscr{P}_0$  always). If j = K, then we have

$$F(P) \le \rho(\pi^{\mathcal{P}}, \pi_K) \le 2d_K$$

If j < K, then we have  $\mathcal{P} \notin \mathscr{P}_{j+1}$ , and hence

$$F(P) \leq \rho(\pi^{\mathcal{P}}, \pi_{K}) - \gamma \cdot \lambda_{j+1} \inf_{M \in \operatorname{co}(\mathcal{P})} \mathbb{E}_{\pi \sim q_{j+1}} D_{\mathrm{H}}^{2} \left( M(\boldsymbol{\pi}), \overline{M}(\boldsymbol{\pi}) \right)$$
$$\leq \rho(\pi^{\mathcal{P}}, \pi_{j}) + \rho(\pi_{j}, \pi_{K}) - \gamma \lambda_{j+1} \varepsilon_{j+1}^{2}$$
$$\leq 3d_{j} + 2d_{K} - \gamma \lambda_{j+1} \varepsilon_{j+1}^{2} \leq 2d_{K},$$

where the last line uses the fact that  $d_j \leq D\varepsilon_j$  and  $\lambda_{j+1}\varepsilon_{j+1}^2 = 2^{-K-1}\varepsilon_j \geq \frac{\varepsilon\varepsilon_j}{2}$ . Therefore, we can conclude that

$$F(\mathcal{P}) \leq 2d_K \leq 2D\varepsilon_K \leq 2D\varepsilon, \quad \forall \mathcal{P} \in \mathscr{P}.$$

This immediately implies  $\mathsf{p-dec}_{\gamma}^{\mathsf{o},\mathsf{H}}(\mathscr{P},\overline{M}) \leq 2D\varepsilon$ , and the desired upper bound follows by taking supremum over all  $\overline{M}$ .

# G Estimation-to-Decision Algorithm and Guarantees

In this section, we present the extensions of the PAC E2D algorithm [Foster et al., 2021, 2023b] to LDP learning and query-based learning.

## G.1 LDP-E2D Algorithm

In the following, we present LDP-E2D, the LDP extension of the Estimation-to-Decision algorithm [Foster et al., 2021, 2023b], for PAC learning in private DMSO. In the following, we assume without loss of generality that  $\mathcal{O} = \{-1, 1\}$ .

The LDP-E2D algorithm is based on the *binary* channels (Example 5). Specifically, LDP-E2D adopts the following protocol: For  $t = 1, \dots, T$ :

- The algorithm selects a distribution  $q_t \in \Delta(\Pi \times \mathcal{L})$  (based on the history), sample  $(\pi_t, \ell_t) \sim q_t$ .
- The environment generates a noisy observation  $o_t \sim Q_{\ell_t} \circ M^*(\pi_t)$ , and reveals  $o_t$  to the algorithm.

Note that this protocol automatically ensures the algorithm preserves  $\alpha$ -LDP. Furthermore, conditional on  $(\mathcal{H}_{t-1}, \pi_t, \ell_t)$ , the noisy observation is generated as

$$o_t \sim \operatorname{Rad}(c_{\alpha} \cdot \mathbb{E}_{z \sim M^{\star}(\pi_t)}[\ell_t(z)]).$$

For simplicity of presentation, we denote  $M(\pi)[\ell] := \mathbb{E}_{z \sim M(\pi)}[\ell(z)]$  in the following.

#### G.1.1 Online estimation oracle

The general DEC framework [Foster et al., 2021, 2023b] uses the primitive of an online estimation oracle, denoted by  $\mathbf{Alg}_{\mathbf{Est}}$ , which is an algorithm that produce estimates of the underlying model  $M^*$  at each step based on the prior observations. For LDP-E2D, an estimation oracle at each round t, given the history  $\mathcal{H}_{t-1} = (\pi_i, \ell_i, o_i)_{i=1}^{t-1}$ , returns an estimator

$$\widehat{M}_t = \mathbf{Alg}_{\mathbf{Est}}(\mathcal{H}_{t-1})$$

for the true model  $M^*$ . Here, the oracle's estimation performance is measured by cumulative squared error under each functional  $\ell_t$ , which is different from the non-private setting [Foster et al., 2021, 2023b] where the performance is measured in terms of the squared Hellinger error.

Assumption G.1 (Estimation oracle for  $\mathcal{M}$ ). At each time  $t \in [T]$ , an online estimation oracle  $Alg_{Est}$  for  $\mathcal{M}$  returns, given

$$\mathcal{H}_{t-1} = (\pi_1, \ell_1, o_1), \dots, (\pi_{t-1}, \ell_{t-1}, o_{t-1})$$

with  $(\pi_i, \ell_i) \sim p_i$  and  $o_i \sim \operatorname{Rad}(M^{\star}(\pi_i)[\ell_i])$ , an estimator  $\widehat{M}^t \in (\Pi \to \Delta(\mathcal{Z}))$  such that whenever  $M^{\star} \in \mathcal{M}$ ,

$$\mathbf{Est}_{\mathbf{sq}} := \sum_{t=1}^{T} \mathbb{E}_{(\pi_t, \ell_t) \sim q_t} \mathsf{D}^2_{\ell_t} \Big( M^{\star}(\pi_t), \widehat{M}^t(\pi_t) \Big) \le \mathbf{Est}_{\mathbf{sq}}(T, \delta),$$
(86)

with probability at least  $1 - \delta$ , where  $\mathbf{Est}_{sq}(T, \delta)$  is a known upper bound that we assumed to be a non-decreasing function in  $(T, \delta^{-1})$ .

Oracles satisfying Assumption G.1 can be obtained via online linear regression algorithms, the estimation rate  $\mathbf{Est_{sq}}(T, \delta)$  will typically reflect the statistical complexity of the class  $\mathcal{M}$ . Standard examples include Vovk's Aggregation (Proposition G.1) and Online Mirror Descent (Proposition G.2). For further background, see e.g. Foster et al. [2021, Section 4].

**Proposition G.1** (Vovk's Aggregation). Suppose that  $\mathcal{M}$  is finite. Then the Vovk's aggregation algorithm achieves

$$\mathbf{Est}_{\mathbf{sq}}(T,\delta) \lesssim \frac{1}{\alpha^2} \cdot \log(|\mathcal{M}|/\delta).$$

Furthermore, for each round  $t \in [T]$ ,  $\widehat{M}^t \in co(\mathcal{M})$ .

**Proposition G.2** (Online Mirror Descent). Suppose that  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ . Then the Online Mirror Descent (Algorithm 3) achieves

$$\mathbf{Est}_{\mathbf{sq}}(T,\delta) \lesssim \frac{1}{\alpha} \sqrt{C_{\mathrm{KL}}(\mathcal{M}) \cdot T} + \frac{\log(1/\delta)}{\alpha^2}$$

where  $C_{\mathrm{KL}}(\mathcal{M})$  is defined in Proposition 16.

### G.1.2 LDP-E2D Algorithm and its guarantees

With an online estimation oracle  $Alg_{Est}$ , we present the LDP-E2D algorithm (Appendix G), which generalizes the E2D algorithm of Foster et al. [2023b] to LDP learning. LDP-E2D algorithm consists of two phases: the exploration phase and the refining phase.

# Algorithm 2 LDP Estimation-to-Decision Algorithm for PAC learning (LDP-E2D)

**Input:** Round  $T \ge 1$ , error probability  $\delta > 0$ , model class  $\mathcal{M}$ , estimation oracle  $\mathbf{Alg}_{\mathbf{Est}}$ .

- 1: Define  $K := \lceil \log 2/\delta \rceil$ ,  $N := \frac{T}{K+1}$ , and  $\overline{\mathbf{Est}}_{\mathbf{sq}} := \mathbf{Est}_{\mathbf{sq}}(N, \frac{\delta}{4K})$ .
- 2: Set  $\overline{\varepsilon}(T) := 8\sqrt{\frac{1}{N} \cdot \overline{\mathbf{Est}}_{\mathbf{sq}}}.$
- 3: /\* Exploration phase \*/
- 4: for  $t = 1, 2, \cdots, N$  do
- 5: Compute estimator  $\widehat{M}_t = \mathbf{Alg}_{\mathbf{Est}} ((\pi_i, \ell_i, o_i)_{i=1}^{t-1}).$
- 6: Compute

$$(p_t, q_t) := \underset{\substack{p \in \Delta(\Pi)\\q \in \Delta(\Pi \times \mathcal{L})}}{\arg \min} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M, \pi)] \mid \mathbb{E}_{(\pi, \ell) \sim q} \mathsf{D}_{\ell}^2(M(\pi), \widehat{M}^t(\pi)) \le \bar{\varepsilon}(T)^2 \right\}$$

- 7: Sample decision  $(\pi_t, \ell_t) \sim q_t$ .
- 8: Receive  $o_t \sim \mathsf{Q}_{\ell_t} \circ M^{\star}(\pi_t)$  from the environment.
- 9: /\* Refining phase \*/
- 10: Sample K indices  $t_1, \ldots, t_K \sim \text{Unif}([N])$  independently.

11: for 
$$k = 1, 2, \cdots, K$$
 do

- 12: Set  $q^{(k)} := q_{t_k}$ .
- 13: **for**  $t = kN + 1, \dots, (k+1)N$  **do**

14: Compute estimator  $\widehat{M}_t = \mathbf{Alg}_{\mathbf{Est}} ((\pi_i, \ell_i, o_i)_{i=kN+1}^{t-1}).$ 

15: Sample  $(\pi_t, \ell_t) \sim q^{(k)}$ , and receive  $o_t \sim \mathsf{Q}_{\ell_t} \circ M^*(\pi_t)$  from the environment.

16: Compute  $M^{(k)} := \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \widehat{M}^t$ .

Estimation-to-Decision objective:

17: Set 
$$\widehat{k} := \arg\min_{k \in [K]} \mathbb{E}_{(\pi,\ell) \sim q^{(k)}} \mathsf{D}_{\ell}^{2} \left( \widehat{M}^{t_{k}}, M^{(k)} \right)$$
  
**Output:**  $\widehat{p} := p^{(k)}$  and  $\pi_{T+1} \sim \widehat{p}$ 

**Exploration phase.** At each round  $t \in [N]$  in this phase, the algorithm uses  $\mathbf{Alg}_{\mathbf{Est}}$  to compute an estimator  $\widehat{M}^t = \mathbf{Alg}_{\mathbf{Est}}(\mathcal{H}_{t-1})$  based on the history  $\mathcal{H}_{t-1} = (\pi_i, \ell_i, o_i)_{i=1}^{t-1}$ . Then, based on  $\widehat{M}^t$ , the algorithm computes a joint *exploration-exploitation* distribution  $(p_t, q_t)$  by solving the following

$$(p_t, q_t) := \underset{\substack{p \in \Delta(\Pi)\\q \in \Delta(\Pi \times \mathcal{L})}}{\operatorname{arg\,min}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M, \pi)] \mid \mathbb{E}_{(\pi, \ell) \sim q} \mathsf{D}^2_{\ell}(M(\pi), \widehat{M}^t(\pi)) \le \bar{\varepsilon}(T)^2 \right\}.$$
(87)

Note that the value of this minimax optimization problem is always bounded by  $\mathsf{p-dec}_{\overline{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}, \widehat{M}^t)$ . The algorithm then samples  $(\pi_t, \ell_t) \sim q_t$  from the exploitation distribution, sends it to the *t*-th user, and receives the noisy observation  $o_t \sim \operatorname{Rad}(M^*(\pi_t)[\ell_t])$  according to the interaction protocol. After the exploration phase, the goal of the algorithm is to select an index  $t \in [N]$  such that the distribution  $p_t$  achieves low risk. Note that in general, the risk of  $p_t$  may not be estimated from samples. However, if we can certify that  $\mathbb{E}_{(\pi,\ell)\sim q_t} D^2_{\ell}(M^*(\pi), \widehat{M}^t(\pi)) \leq \overline{\varepsilon}(T)^2$ , then the risk  $\mathbb{E}_{\pi\sim p^t} L(M^*, \pi) \leq \mathbf{p}\operatorname{-dec}_{\overline{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M})$  is bounded automatically. Notice that by our assumption on  $\operatorname{Alg}_{\mathsf{Est}}$ (Assumption G.1), with probability at least  $1 - \frac{\delta}{4K}$ ,

$$\sum_{t=1}^{N} \mathbb{E}_{(\pi_t,\ell_t)\sim q_t} \mathsf{D}^2_{\ell_t} \Big( M^{\star}(\pi_t), \widehat{M}^t(\pi_t) \Big) \leq \overline{\mathbf{Est}}_{\mathbf{sq}},$$

and hence there are at least N/2 indices t such that  $\mathbb{E}_{(\pi,\ell)\sim q_t} \mathsf{D}^2_{\ell}(M^{\star}(\pi), \widehat{M}^t(\pi)) \leq \frac{1}{16}\bar{\varepsilon}(T)^2$ . Therefore, in the refining phase, the algorithm proceeds as follows to identify an index t such that  $\widehat{M}^t$  achieves a small estimation error.

**Refining phase.** At the start of this phase, the algorithm randomly samples  $t_1, \dots, t_K \sim \text{Unif}([N])$ . Then, with probability at least  $1 - \frac{3}{4}\delta$ ,

there exists 
$$k \in [K]$$
 such that  $\mathbb{E}_{(\pi,\ell)\sim q_{t_k}} \mathsf{D}^2_\ell(M^\star(\pi), \widehat{M}^{t_k}(\pi)) \le \frac{1}{16}\bar{\varepsilon}(T)^2$ , (88)

as we have argued above. Thus, for each batch  $k \in [K]$ , the algorithm uses N rounds to obtain an estimator  $M^{(k)}$  of the ground-truth model  $M^*$  under the distribution  $q^{(k)} := q_{t_k}$ :

For each round  $t \in [kN + 1, (k + 1)N]$  in the k-th batch, the algorithm samples  $(\pi_t, \ell_t) \sim q^{(k)}$  and sends the pair to the learner. By running an instance of  $\mathbf{Alg}_{\mathbf{Est}}$  within the batch, it is guaranteed that with probability at least  $1 - \frac{\delta}{4K}$ 

$$\sum_{t=kN+1}^{(k+1)N} \mathbb{E}_{(\pi,\ell)\sim q^{(k)}} \mathsf{D}_{\ell}^2\Big(M^{\star}(\pi), \widehat{M}^t(\pi)\Big) \leq \overline{\mathbf{Est}}_{\mathbf{sq}}$$

Hence, by the convexity of the divergence  $\mathsf{D}^2_\ell$ , we have

$$\mathbb{E}_{(\pi,\ell)\sim q^{(k)}}\mathsf{D}^2_\ell\Big(M^\star(\pi), M^{(k)}(\pi)\Big) \le \frac{\mathbf{Est}_{\mathbf{sq}}}{N} = \frac{1}{16}\bar{\varepsilon}(T)^2.$$
(89)

Therefore, taking the union bound, Eq. (88) and Eq. (89) (for each  $k \in [K]$ ) hold simultaneously with probability at least  $1 - \delta$ . Therefore, under this success event, we know

$$\min_{k \in [K]} \mathbb{E}_{(\pi,\ell) \sim q^{(k)}} \mathsf{D}_{\ell}^2 \Big( \widehat{M}^{t_k}, M^{(k)} \Big) \le \frac{1}{4} \bar{\varepsilon}(T)^2,$$

and hence by triangle inequality,

$$\mathbb{E}_{(\pi,\ell)\sim q^{(\widehat{k})}} \mathsf{D}^2_{\ell} \Big( M^{\star}(\pi), \widehat{M}^{t_{\widehat{k}}}(\pi) \Big) \\ \leq \mathbb{E}_{(\pi,\ell)\sim q^{(\widehat{k})}} \mathsf{D}^2_{\ell} \Big( M^{\star}(\pi), M^{(\widehat{k})}(\pi) \Big) + \mathbb{E}_{(\pi,\ell)\sim q^{(\widehat{k})}} \mathsf{D}^2_{\ell} \Big( M^{(\widehat{k})}(\pi), \widehat{M}^{t_{\widehat{k}}}(\pi) \Big) \leq \bar{\varepsilon}(T)^2.$$

Therefore, for  $\widehat{p} = p_{t_{\widehat{k}}}$ , we have

$$\mathbb{E}_{\pi_{T+1}\sim \widehat{p}}L(M^{\star},\pi_{T+1}) \leq \mathsf{p-dec}_{\overline{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M},\widehat{M}^{t_{\widehat{k}}}).$$

The argument above immediately yields the following guarantee of LDP-E2D (Algorithm 2).

**Theorem G.3.** LDP-E2D (Algorithm 2) preserves  $\alpha$ -LDP, and with probability at least  $1 - \delta$ , it holds that

$$\mathbf{Risk}_{\mathsf{DM}}(T) = \mathbb{E}_{\widehat{p}}L(M^*, \pi_{T+1}) \le \max_{t \in [T]} \mathsf{p-dec}_{\overline{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}, \widehat{M}^t).$$

Theorem 21 is then a direct corollary by instantiating  $\mathbf{Alg}_{\mathbf{Est}}$  with Vovk's aggregation (Proposition G.1, where  $\widehat{M}^t \in \operatorname{co}(\mathcal{M})$  for all  $t \in [T]$ ). For statistical problems, we may also instantiate  $\mathbf{Alg}_{\mathbf{Est}}$  with Online Mirror Descent (Proposition G.2) which gives with probability at least  $1 - \delta$ ,

$$\mathbf{Risk}_{\mathsf{DM}}(T) \leq \sup_{\overline{M} \in \Delta(\mathcal{Z})} \mathsf{p-dec}_{\overline{\varepsilon}(T)}^{\mathsf{LDP}}(\mathcal{M}, \overline{M}), \qquad \text{where } \overline{\varepsilon}(T) \asymp \sqrt{\frac{C_{\mathrm{KL}}(\mathcal{M})}{\alpha^2 T}} + \frac{\log(1/\delta)}{\alpha^2 T}.$$

#### G.1.3 Proof of Proposition G.2

We present the specifications of Online Mirror Descent for online estimation in Algorithm 3, which is inspired by Feldman [2017].

Algorithm 3 Online Mirror Descent

**Input:** History  $\mathcal{H}_{t-1} = (\ell_i, o_i)_{i=1}^{t-1}$ , number of total rounds N

1: Parameters: Initial reference  $\overline{M}$  and stepsize  $\eta = \sqrt{\frac{C_{\text{KL}}}{16N}}$ .

2: Compute

$$\widehat{M}^{t}[z] \propto_{z} \overline{M}[z] \cdot \exp\left(-\eta \sum_{s=1}^{t-1} \left(c_{\alpha} \langle \ell^{s}, \widehat{M}^{s} \rangle - o^{s}\right) \ell^{s}(z)\right)$$

**Output:** Output  $\widehat{M}^t \in \Delta(\mathcal{Z})$ .

**Proof of Proposition G.2.** Consider the loss function sequence

$$L^{t}(M) = \frac{1}{2c_{\alpha}} (c_{\alpha} \langle \ell_{t}, M \rangle - o_{t})^{2}, \qquad t \in [N].$$

Then, Algorithm 3 implements the online mirror descent with regularizer  $R(M) = D_{\text{KL}}(M \parallel \overline{M})$ and stepsize  $\eta$ . Using the well-known guarantee of mirror descent (see e.g. Hazan et al. [2016]), we have

$$\sum_{t=1}^{N} \langle \nabla L^{t}(\widehat{M}^{t}), \widehat{M}^{t} - M \rangle \leq \eta \sum_{t=1}^{N} \|\nabla L^{t}(\widehat{M}^{t})\|_{\infty}^{2} + \frac{D_{\mathrm{KL}}(M \parallel \overline{M})}{\eta}, \qquad \forall M \in \Delta(\mathcal{Z}).$$

Notice that  $\|\nabla L^t(M)\|_{\infty} \leq 2$  for any  $M \in \Delta(\mathcal{Z})$ . Therefore, using the upper bound  $D_{\mathrm{KL}}(M^* \| \overline{M}) \leq C_{\mathrm{KL}}$  and our choice of  $\eta$ , we know

$$\sum_{t=1}^{N} \langle \nabla L^{t}(\widehat{M}^{t}), \widehat{M}^{t} - M \rangle \leq 16\eta N + \frac{C_{\mathrm{KL}}}{\eta} = 4\sqrt{NC_{\mathrm{KL}}}.$$

Notice that  $\nabla L^t(\widehat{M}^t) = \left(c_\alpha \langle \ell_t, \widehat{M}^t \rangle - o_t\right) \cdot \ell_t$ , and hence

$$\langle \nabla L^t(\widehat{M}^t), \widehat{M}^t - M \rangle = (M^*[\ell_t] - o_t) \cdot \left(\widehat{M}^t[\ell_t] - M^*[\ell_t]\right) + c_\alpha \left(\widehat{M}^t[\ell_t] - M^*[\ell_t]\right)^2.$$

Therefore, we denote  $X_t := \widehat{M}^t[\ell_t] - M^{\star}[\ell_t]$  and  $Z_t := o_t - M^{\star}[\ell_t]$ , and it holds that

$$\mathbb{E}[Z_t|\mathcal{H}_{t-1},\ell_t] = 0, \quad \mathbb{E}[X_t^2|\mathcal{H}_{t-1}] = \mathbb{E}_{\ell \sim q_t} \mathsf{D}_{\ell}^2 \Big(\widehat{M}^t, M^\star\Big),$$

where we recall that  $o_t \sim \text{Rad}(M^*[\ell_t])$ . In particular, by Hoeffding's inequality, for any fixed parameter  $\lambda > 0$ , with probability at least  $1 - \delta$ ,

$$\lambda \sum_{t=1}^{N} Z_t X_t - \frac{\lambda^2}{2} \sum_{t=1}^{N} X_t^2 \le \log(1/\delta).$$

Further, by Freedman's inequality and the fact that  $X_t^2 \in [0, 1]$ , with probability at least  $1 - \delta$ ,

$$\sum_{t=1}^{N} \mathbb{E}[X_t^2 | \mathcal{H}_{t-1}] \le \frac{3}{2} \sum_{t=1}^{N} X_t^2 + 5 \log(1/\delta).$$

Therefore, we may choose  $\lambda = \frac{c_{\alpha}}{2}$ , and then with probability at least  $1 - 2\delta$ ,

$$4\sqrt{NC_{\mathrm{KL}}} \geq \sum_{t=1}^{N} c_{\alpha} X_{t}^{2} - X_{t} Z_{t}$$
  
$$\geq \frac{3c_{\alpha}}{4} \sum_{t=1}^{N} X_{t}^{2} - \frac{2\log(1/\delta)}{c_{\alpha}}$$
  
$$\geq \frac{c_{\alpha}}{2} \sum_{t=1}^{N} \mathbb{E}[X_{t}^{2}|\mathcal{H}_{t-1}] - 3c_{\alpha}\log(1/\delta) - \frac{2\log(1/\delta)}{c_{\alpha}}.$$

Using the fact  $\mathbb{E}[X_t^2|\mathcal{H}_{t-1}] = \mathbb{E}_{\ell \sim q_t} \mathsf{D}_{\ell}^2(\widehat{M}^t, M^\star)$  gives the desired upper bound.

## G.2 Query-based E2D algorithm

In the following, we present the E2D algorithm (SQ-E2D, Algorithm 4) for SQ DMSO.

## Algorithm 4 Query-based Estimation-to-Decisions (SQ-E2D)

**Input:** Round  $T \ge 1$ , error probability  $\delta > 0$ , model class  $\mathcal{M} \subseteq (\Pi \to \mathcal{V})$ , GQ oracle **0**.

- 1: Define  $K := \lceil \log 2/\delta \rceil$ ,  $T_0 = \frac{T}{2}$ ,  $N := \frac{T}{2K}$ .
- 2: Set  $\bar{\gamma} := C_0 \max\left\{\frac{\log |\mathcal{M}|}{T}, \frac{\log(1/\delta)}{N}\right\}$  for a large absolute constant  $C_0$ .
- 3: /\* Exploration phase \*/
- 4: for  $t = 1, 2, \cdots, T_0$  do
- 5: Compute  $\mu^t = \text{Unif}(\widehat{\mathcal{M}}^t)$ , where

$$\widehat{\mathcal{M}}^t := \{ M \in \mathcal{M} : \| M(\pi^s) - v^s \| \le \tau, \forall s < t \},$$
(90)

#### 6: Compute

$$(p_t, q_t) := \underset{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}}{\arg \min} \sup_{\substack{M \in \mathcal{M} \\ q \in \Delta(\Pi)}} \left\{ \mathbb{E}_{\pi \sim p}[L(M, \pi)] \mid \mathbb{P}_{\pi \sim q, \overline{M} \sim \mu^t} \left( \left\| M(\pi) - \overline{M}(\pi) \right\| > 2\tau \right) \le \bar{\gamma} \right\}.$$

- 7: Sample  $\pi_t \sim q_t$ , query  $\pi_t$ , and receive  $v_t$  from the oracle O.
- 8: /\* Refining phase \*/
- 9: Sample K indices  $t_1, \ldots, t_K \sim \text{Unif}([T_0])$  independently.
- 10: Set  $k^* = 1$ .
- 11: for  $k = 1, 2, \cdots, K$  do
- 12: Set  $q^{(k)} := q^{t_k}$  and batch  $\mathcal{T}_k := \{T_0 + (k-1)N + 1, \cdots, T_0 + kN\}.$
- 13: for  $t \in \mathcal{T}_k$  do
- 14: Sample  $\pi_t \sim q_t$ , query  $\pi_t$ , and receive  $v_t$  from the oracle O.

15: Compute 
$$\hat{e}^{(k)} := \frac{1}{N} \sum_{t \in \mathcal{T}_k} \mathbb{P}_{M \sim \mu_t}(\|M(\boldsymbol{\pi}_t) - v_t\| > \tau).$$

- 16: **if**  $\hat{e}^{(k)} < \bar{\gamma}$  **then**
- 17: Set  $k^* = k$  and **break**.

**Output:**  $\widehat{p} := p^{(k^{\star})}$  and  $\pi_{T+1} \sim \widehat{p}$ 

We state the following guarantee of SQ-E2D.

**Theorem G.4.** For any model class  $\mathcal{M}$ , SQ-E2D (Algorithm 4) achieves that given access to any GSQ oracle  $\mathsf{GQ}_M^{\tau}$ , with probability at least  $1 - \delta$ ,

$$\mathbf{Risk}_{\mathsf{DM}}(T) \leq \mathsf{p-dec}_{\bar{\varepsilon}(T)}^{2\tau-\mathsf{SQ}}(\mathcal{M}),$$

where  $\bar{\varepsilon}(T) = C\sqrt{\frac{\log |\mathcal{M}| + \log^2(1/\delta)}{T}}$ .

**Remark G.5.** We note that the log  $|\mathcal{M}|$ -factor above can be necessary for more general setting (e.g., interactive SQ learning). However, under SQ setting and distributional search problem (i.e.,  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ ), Feldman [2017] derives an upper bound (25) scaling with the SQ dimension (cf. Section 4.2) and  $C_{\text{KL}}$ . When specialized to this setting, our upper bound above does not involve extra  $\tau^{-1}$ -factors, but the log  $|\mathcal{M}|$ -factor can be much larger than  $C_{\text{KL}}$ . However, if we replace the model elimination subroutine (90) with the Online Mirror Descent subroutine (Algorithm 3), then the obtained algorithm is essentially an analog of the one of Feldman [2017] and achieves an upper bound scaling with SQ DEC and  $C_{\text{KL}}$ .

## G.2.1 Proof of Theorem 15

The proof is analogous to the analysis in Appendix G.1.2. We first invoke the following lemma.

Lemma G.6. It holds that

$$\sum_{t=1}^{T_0} \mathbb{P}_{M \sim \mu^t}(\|M(\boldsymbol{\pi}_t) - v_t\| > \tau) \le \log |\mathcal{M}|.$$

Then, by Freedman's inequality, with probability at least  $1 - \frac{\delta}{4}$ , it holds that

$$\sum_{t=1}^{T_0} \mathbb{P}_{\pi \sim q_t, M \sim \mu^t} (\|M(\pi) - M^*(\pi)\| > 2\tau) \le 2 \sum_{t=1}^{T_0} \mathbb{P}_{M \sim \mu^t} (\|M(\pi_t) - M^*(\pi_t)\| > 2\tau) + 4 \log(4/\delta)$$
$$\le 2 \sum_{t=1}^{T_0} \mathbb{P}_{M \sim \mu^t} (\|M(\pi_t) - v_t\| > \tau) + 4 \log(4/\delta)$$
$$\le 2 \log |\mathcal{M}| + 4 \log(4/\delta).$$

In the following, we denote

$$e(t) := \mathbb{P}_{\boldsymbol{\pi} \sim q_t, M \sim \mu^t}(\|M(\boldsymbol{\pi}) - M^{\star}(\boldsymbol{\pi})\| > 2\tau).$$

Therefore, conditional on this success event, for at least  $\frac{T_0}{2}$  many  $t \in [T_0]$ , t belongs to the set

$$\mathcal{B} := \left\{ t \in [T_0] : \mathbb{P}_{\pi \sim q_t, M \sim \mu^t} (\|M(\pi) - M^*(\pi)\| > 2\tau) \le \frac{2(2\log|\mathcal{M}| + 4\log(2/\delta))}{T_0} \le \frac{\bar{\gamma}}{16} \right\}.$$

In particular, with probability at least  $1 - \frac{\delta}{2}$ , there exists  $k \in [K]$  such that  $t_k \in \mathcal{B}$ .

In the following, we denote  $e^{(k)} := e(t_k)$ . Then, by Freedman's inequality, with probability at least  $1 - \frac{\delta}{2}$ , the following holds for all  $k \in [K]$ :

$$\hat{e}^{(k)} \le 2e^{(k)} + \frac{4\log(4K/\delta)}{N}, \qquad e^{(k)} \le 2\hat{e}^{(k)} + \frac{4\log(4K/\delta)}{N}.$$

Therefore, conditional on the all the success events, we know that there exists  $k \in [K]$  such that  $t_k \in \mathcal{B}$ , which implies  $\hat{e}^{(k)} \leq \frac{\bar{\gamma}}{4}$ . Hence, it is ensured that  $\hat{e}^{(k^*)} \leq \frac{\bar{\gamma}}{4}$ , which in terms implies  $e^{(k^*)} \leq \bar{\gamma}$ . Therefore, for  $t^* = t_{k^*}$ , it holds that

$$\mathbb{P}_{\boldsymbol{\pi} \sim q^{t^{\star}}, \overline{M} \sim \mu^{t^{\star}}} \left( \left\| \overline{M}(\boldsymbol{\pi}) - M^{\star}(\boldsymbol{\pi}) \right\| > 2\tau \right) \leq \bar{\gamma}.$$

Thus,

$$\begin{split} \mathbb{E}_{\pi \sim p^{t^{\star}}} L(M^{\star}, \pi) &\leq \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi)}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M, \pi)] \mid \mathbb{P}_{\pi \sim q, \overline{M} \sim \mu^{t^{\star}}} \left( \left\| M(\pi) - \overline{M}(\pi) \right\| > 2\tau \right) \leq \bar{\gamma} \right\} \\ &\leq \mathsf{p-dec}_{2\bar{\gamma}}^{2\tau - \mathsf{SQ}}(\mathcal{M}). \end{split}$$

The proof of Theorem 15 is hence completed.

**Proof of Lemma G.6.** Let  $U_t = |\widehat{\mathcal{M}}^t|$ , and then

$$\mathbb{P}_{M \sim \mu^t}(\|M(\boldsymbol{\pi}_t) - v_t\| > \tau) = \frac{|\widehat{\mathcal{M}}^t \setminus \widehat{\mathcal{M}}^{t+1}|}{|\widehat{\mathcal{M}}^t|} = \frac{U_t - U_{t+1}}{U_t} \le \log U_t - \log U_{t+1}$$

Taking summation and using  $U_1 = |\mathcal{M}|$  completes the proof.

# H Proofs from Section 4

## H.1 Proof of Theorem 14

Fix  $T \ge 1, \tau \ge 0$ , reference model  $\overline{M}$ . We first consider the case L is metric-based, i.e., it is given by  $L(M,\pi) = \rho(\pi^M,\pi)$  for a pseudo-metric  $\rho$  over  $\Pi$ . We denote  $\underline{\varepsilon} := \frac{1}{2\sqrt{T}}$  and  $\Delta := \mathsf{p-dec}_{\underline{\varepsilon}}^{\tau-\mathsf{sq}}(\mathcal{M},\overline{M})$ .

We first describe any *T*-round query-based algorithm in the following way (cf. Section 3). A *T*-round algorithm  $Alg = \{q_t\}_{t \in [T]} \cup \{p\}$  is specified by a sequence of mappings, where the *t*-th mapping  $q_t(\cdot \mid \mathcal{H}_{t-1})$  specifies the distribution of  $\pi_t$  based on the history  $\mathcal{H}_{t-1} = (\pi^s, v^s)_{s \leq t-1}$ , and the final map  $p(\cdot \mid \mathcal{H}_T)$  specifies the distribution of the *output policy*  $\pi_{T+1}$  based on  $\mathcal{H}_T$ .

Next, we fix an arbitrary, randomized reference model  $\overline{M} : \mathbf{\Pi} \to \Delta(\mathcal{V})$ , and we construct a GQ oracle for each model  $M \in \mathcal{M}$  as follows. For  $M \in \mathcal{M}$ , we let  $\mathsf{GQ}_M^{\tau}$  be an oracle that response to any decision  $\pi$  as

$$\mathsf{GQ}_M^{\tau}(\boldsymbol{\pi}) = \begin{cases} v, & \text{if } \|M(\boldsymbol{\pi}) - v\| \leq \tau, \\ M(\boldsymbol{\pi}), & \text{otherwise.} \end{cases} \quad \text{where } v \sim \overline{M}(\boldsymbol{\pi}).$$

For any model M, we let  $\mathbb{P}^{M,\operatorname{Alg}}(\cdot)$  to be the distribution of  $(\mathcal{H}^T, \pi_{T+1})$  generated by the algorithm Alg under the oracle  $\mathsf{GQ}_M^{\tau}$ , and let  $\mathbb{E}^{M,\operatorname{Alg}}[\cdot]$  to be the corresponding expectation. We also define  $\overline{\mathbb{P}}$ to be the distribution of  $(\mathcal{H}^T, \pi_{T+1})$  by the algorithm Alg under the oracle  $\mathsf{GQ}_{\overline{M}}^{\tau}$  that always return  $v \sim \overline{M}(\pi)$ , and let  $\overline{\mathbb{E}}$  be the corresponding expectation.

Following Eq. (55), we define

$$q = \frac{1}{T} \sum_{t=1}^{T} \overline{\mathbb{P}}(\boldsymbol{\pi}_t = \cdot) \in \Delta(\boldsymbol{\Pi}),$$
(91)

and

$$\mathcal{M}_{q,\underline{\varepsilon}}(\overline{M}) := \{ M \in \mathcal{M} : \mathbb{P}_{\boldsymbol{\pi} \sim q, v \sim \overline{M}(\boldsymbol{\pi})} \big( \left\| M(\boldsymbol{\pi}) - \overline{M}(\boldsymbol{\pi}) \right\| > \tau \big) \le \underline{\varepsilon}^2 \}.$$
(92)

By definition, for any distribution  $p \in \Delta(\Pi)$ , there exists  $M \in \mathcal{M}_{q,\underline{\varepsilon}}(\overline{M})$  such that

 $\mathbb{E}_{\pi \sim p}[L(M,\pi)] \ge \Delta.$ 

In particular,  $\mathcal{M}_{q,\underline{\varepsilon}}(\overline{M})$  is non-empty, and we fix a model  $M_0 \in \mathcal{M}_{q,\underline{\varepsilon}}(\overline{M})$  and let  $\pi_0 := \pi^{M_0}$ . Then there exists  $M_1$  such that  $L(M_1, \pi_0) \ge \Delta$ , i.e.,  $\rho(\pi^{M_1}, \pi_0) \ge \Delta$ . We denote  $\pi_1 := \pi^{M_1}$ 

Now, using the chain rule of TV distance, it holds that for any  $M \in \mathcal{M}_{q,\underline{\varepsilon}}(\overline{M})$ 

$$D_{\mathrm{TV}}\left(\mathbb{P}^{M,\mathrm{Alg}},\bar{\mathbb{P}}\right) \leq \sum_{t=1}^{T} \bar{\mathbb{E}}\left[D_{\mathrm{TV}}\left(\mathbb{P}^{M,\mathrm{Alg}}(v_{t}=\cdot|\mathcal{H}_{t-1},\boldsymbol{\pi}_{t}),\bar{\mathbb{P}}(v_{t}=\cdot|\mathcal{H}_{t-1},\boldsymbol{\pi}_{t})\right)\right]$$
$$= \sum_{t=1}^{T} \bar{\mathbb{E}}\left[D_{\mathrm{TV}}\left(\mathsf{GQ}_{M}^{\tau}(\boldsymbol{\pi}_{t}),\mathsf{GQ}_{\overline{M}}^{\tau}(\boldsymbol{\pi}_{t})\right)\right]$$
$$= \sum_{t=1}^{T} \bar{\mathbb{E}}\left[\mathbb{P}_{v\sim\overline{M}(\boldsymbol{\pi}_{t})}(\|M(\boldsymbol{\pi}_{t})-v\|>\tau)\right]$$
$$= T \cdot \mathbb{E}_{\boldsymbol{\pi}\sim q}\left[\mathbb{P}_{v\sim\overline{M}(\boldsymbol{\pi})}(\|M(\boldsymbol{\pi})-v\|>\tau)\right]$$
$$= T \cdot \mathbb{P}_{\boldsymbol{\pi}\sim q, v\sim\overline{M}(\boldsymbol{\pi})}(\|M(\boldsymbol{\pi})-v\|>\tau) \leq T\underline{\varepsilon}^{2},$$

where the third line follows from the definition of  $\mathsf{GQ}_M^{\tau}$ . Therefore, by triangle inequality, we have

$$D_{\mathrm{TV}}\left(\mathbb{P}^{M_0,\mathsf{Alg}},\mathbb{P}^{M_1,\mathsf{Alg}}\right) \leq 2T\underline{\varepsilon}^2.$$

Hence, it holds that

$$\begin{split} & \mathbb{E}^{M_{0},\mathsf{Alg}}[\rho(\pi_{T+1},\pi_{0})] + \mathbb{E}^{M_{1},\mathsf{Alg}}[\rho(\pi_{T+1},\pi_{1})] \\ & \geq \frac{\Delta}{2} \bigg[ \mathbb{P}^{M_{0},\mathsf{Alg}} \bigg( \rho(\pi_{T+1},\pi_{0}) > \frac{\Delta}{2} \bigg) + \mathbb{P}^{M_{1},\mathsf{Alg}} \bigg( \rho(\pi_{T+1},\pi_{1}) > \frac{\Delta}{2} \bigg) \bigg] \\ & \geq \frac{\Delta}{2} \bigg[ 1 - D_{\mathrm{TV}} \left( \mathbb{P}^{M_{0},\mathsf{Alg}}, \mathbb{P}^{M_{1},\mathsf{Alg}} \right) \bigg] \geq \frac{\Delta}{4}, \end{split}$$

where the second inequality follows from  $\rho(\pi_0, \pi_1) \geq \Delta$ . Therefore,

$$\max_{M \in \{M_0, M_1\}} \mathbb{E}^{M, \mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \geq \frac{\Delta}{8}.$$

Taking the supremum over all reference models  $\overline{M}$  completes the proof for metric-based L.

For a general loss L, we may choose  $\underline{\varepsilon} := \sqrt{\frac{\delta}{T}}$ ,  $\Delta := \mathsf{p-dec}_{\underline{\varepsilon}}^{\tau-\mathsf{SQ}}(\mathcal{M}, \overline{M})$ , and we let q as in Eq. (91), and  $p = \overline{\mathbb{P}}(\pi_{T+1} = \cdot) \in \Delta(\Pi)$ . Then, we can pick a model  $M \in \mathcal{M}_{q,\underline{\varepsilon}}(\overline{M})$  such that  $\mathbb{E}_{\pi \sim p}[L(M, \pi)] \geq \Delta$ . Then, using the fact that  $D_{\mathrm{TV}}\left(\mathbb{P}^{M,\operatorname{Alg}}, \overline{\mathbb{P}}\right) \leq T\underline{\varepsilon}^2 = \delta$ , we can lower bound

$$\mathbb{E}^{M,\operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] \ge \overline{\mathbb{E}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] - D_{\operatorname{TV}}\left(\mathbb{P}^{M,\operatorname{Alg}},\overline{\mathbb{P}}\right) \ge \Delta - \delta.$$

## H.2 Proof of Theorem 15

The first upper bound of Theorem 15 is proven in Appendix G.2 (cf. Theorem G.4), and the second upper bound follows immediately from combining Theorem 8 and Lemma E.5.  $\Box$ 

## H.3 Proof of Proposition 17

We recall that for any model  $M \in \Delta(\mathcal{Z})$ , it induces a map  $\phi \mapsto M(\phi) = \mathbb{E}_{z \sim M}[\phi(z)]$ , and hence we can regard  $M : \Phi \to \mathbb{R}$ .

By definition,

 $\begin{aligned} \mathsf{p}\text{-}\mathsf{dec}_{\varepsilon}^{\tau\text{-}\mathsf{SQ}}(\mathcal{M}^{\mathsf{d}},\overline{M}) &> 1-\beta \\ \Leftrightarrow \quad \forall p \in \Delta(\Pi), \forall q \in \Delta(\mathcal{L}), \exists M \in \mathcal{M}^{\mathsf{d}}, \text{ such that } 1-p(\Pi_{M}) > 1-\beta, \mathbb{P}_{\ell \sim q}\big(\mathsf{D}_{\ell}(M,\overline{M}) > \tau\big) \leq \varepsilon^{2} \\ \Leftrightarrow \quad \forall p \in \Delta(\Pi), \forall q \in \Delta(\mathcal{L}), \exists M \in \mathcal{M}_{p,\beta}^{\mathsf{d}}, \text{ such that } \mathbb{P}_{\ell \sim q}\big(\mathsf{D}_{\ell}(M,\overline{M}) > \tau\big) \leq \varepsilon^{2} \\ \Leftrightarrow \quad \forall p \in \Delta(\Pi), \sup_{q \in \Delta(\mathcal{L})} \inf_{M \in \mathcal{M}_{p,\beta}^{\mathsf{d}}} \mathbb{P}_{\ell \sim q}\big(\mathsf{D}_{\ell}(M,\overline{M}) > \tau\big) \leq \varepsilon^{2}. \end{aligned}$ 

Further, using the Minimax theorem, we know

$$\sup_{q\in\Delta(\mathcal{L})}\inf_{M\in\mathcal{M}_{p,\beta}^{\mathsf{d}}}\mathbb{P}_{\ell\sim q}\big(\mathsf{D}_{\ell}(M,\overline{M})>\tau\big)=\inf_{\mu\in\Delta(\mathcal{M}_{p,\beta}^{\mathsf{d}})}\sup_{\ell\in\mathcal{L}}\mathbb{P}_{M\sim\mu}\big(\mathsf{D}_{\ell}(M,\overline{M})>\tau\big).$$

The Minimax theorem can be applied here because as long as  $\tau > 0$  and  $|\mathcal{Z}|$  is finite, the function class  $\{\ell \mapsto \mathbf{1} \{ \mathsf{D}_{\ell}(M, \overline{M}) > \tau \} \}_{M \in \mathcal{M}}$  admits finite eluder dimension<sup>13</sup> and hence finite threshold dimension [Li et al., 2022], and hence the Minimax theorem holds true [Hanneke et al., 2021].

Therefore, we have

$$\begin{split} & \mathsf{p}\text{-}\mathsf{dec}_{\varepsilon}^{\tau\text{-}\mathsf{SQ}}(\mathcal{M}^{\mathsf{d}},\overline{M}) > 1 - \beta \\ \Leftrightarrow & \forall p \in \Delta(\Pi), \inf_{\mu \in \Delta(\mathcal{M}_{p,\beta}^{\mathsf{d}})} \sup_{\ell \in \mathcal{L}} \mathbb{P}_{M \sim \mu} \left( \mathsf{D}_{\ell}(M,\overline{M}) > \tau \right) \leq \varepsilon^{2} \\ \Leftrightarrow & \forall p \in \Delta(\Pi), \sup_{\mu \in \Delta(\mathcal{M}_{p,\beta}^{\mathsf{d}})} \inf_{\ell \in \mathcal{L}} \frac{1}{\mathbb{P}_{M \sim \mu} \left( \mathsf{D}_{\ell}(M,\overline{M}) > \tau \right)} \geq \varepsilon^{-2} \\ \Leftrightarrow & \inf_{p \in \Delta(\Pi)} \sup_{\mu \in \Delta(\mathcal{M}_{p,\beta}^{\mathsf{d}})} \inf_{\ell \in \mathcal{L}} \frac{1}{\mathbb{P}_{M \sim \mu} \left( \mathsf{D}_{\ell}(M,\overline{M}) > \tau \right)} \geq \varepsilon^{-2} \\ \Leftrightarrow & \mathsf{SQDim}_{\beta}^{\tau}(\mathcal{M}^{\mathsf{d}},\overline{M}) \geq \varepsilon^{-2}. \end{split}$$

This is the desired result.

## H.4 Proof of Lemma 18

By definition, in interactive SQ learning, the measurement class is  $\Phi = (\mathcal{Z} \to [0, 1])$ , and for model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ , we regard  $\mathcal{M} \subseteq (\Pi \times \Phi \to \mathbb{R})$  by  $M(\pi, \phi) = \mathbb{E}_{z \sim M(\pi)} \phi(z)$ . Thus, the SQ DEC can be written as

$$\mathsf{p-dec}_{\varepsilon}^{\tau-\mathsf{SQ}}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi \times \mathcal{L})}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L(M,\pi)] \mid \mathbb{P}_{(\pi,\ell) \sim q}\left(\mathsf{D}_{\ell}(M(\pi),\overline{M}(\pi)) > \tau\right) \le \varepsilon^{2} \right\}.$$

For any  $q \in \Delta(\Pi \times \mathcal{L})$ , using Markov's inequality, we have

$$\mathbb{P}_{(\pi,\ell)\sim q}\left(\mathsf{D}_{\ell}(M(\pi),\overline{M}(\pi)) > \tau\right) \leq \frac{1}{\tau^2} \mathbb{E}_{(\pi,\ell)\sim q} \mathsf{D}_{\ell}^2(M(\pi),\overline{M}(\pi)).$$

_	

<sup>&</sup>lt;sup>13</sup>By regarding  $\mathcal{L}, \mathcal{M} \subseteq \mathbb{R}^{\mathbb{Z}}$ , we can write  $\mathsf{D}_{\ell}(M, \overline{M}) = \left| \langle \ell, M - \overline{M} \rangle \right|$  and apply the standard elliptical potential argument.

Conversely, we also have

$$\mathbb{E}_{(\pi,\ell)\sim q}\mathsf{D}^2_\ell(M(\pi),\overline{M}(\pi)) \leq \tau^2 + \mathbb{P}_{(\pi,\ell)\sim q}\big(\mathsf{D}_\ell(M(\pi),\overline{M}(\pi)) > \tau\big).$$

Combining the inequalities above completes the proof of Eq. (27).

# I Remaining Proofs from Section 5 and Appendix B

We note that we have presented the proof of the following results in the previous sections:

- Proof of Theorem 21: Appendix G.1, and see also Appendix F.5.1.
- Proof of Theorem 22 and Proposition 23: Appendix F.5.2.
- Proof of Proposition 24: Appendix F.5.3.
- Proof of Theorem 27: Appendix E.3.2, where we also provide a proof of Theorem 19 (1).
- Proof of Theorem 28: Appendix F.5.1 and Appendix F.5.2.
- Proof of Proposition 29: Appendix F.5.4.

In the subsequent subsections, we present the remaining proofs from Section 5.

## I.1 Proof of Proposition 20

Fix an  $\alpha$ -DP channel Q. By definition, the class of distributions  $\{Q(\cdot|z)\}$  admits a common base measure  $\mu$ , and hence in the following we slightly abuse notations and write a distribution P and its density  $dP/d\mu$  interchangeably. We also denote  $\mathbb{P}' = \mathbb{Q} \circ \mathbb{P}, \mathbb{Q}' = \mathbb{Q} \circ \mathbb{Q}$ .

Define  $p(o) = \inf_{z \in \mathcal{Z}} \mathbb{Q}(o = \cdot | z)$  for any  $o \in \mathcal{O}$ . Then, by definition, for any  $z \in \mathcal{Z}$ ,

$$p(o) \le \mathsf{Q}(o|z) \le e^{\alpha} p(o).$$

Therefore, we define

$$\ell_o(z) = \frac{1}{e^{\alpha} - 1} \left( \frac{\mathsf{Q}(o|z)}{p(o)} - 1 \right) \in [0, 1].$$

Then, for each  $o \in \mathcal{O}$ , it holds that

$$\mathbb{P}'(o) = \mathbb{E}_{z \sim \mathbb{P}} \mathbb{Q}(o|z) = \mathbb{E}_{z \sim \mathbb{P}}[(e^{\alpha} - 1)p(o)\ell_o(z) + p(o)],$$

and hence we know  $\mathbb{P}'(o) \in [p(o), e^{\alpha}p(o)]$ , and similarly  $\mathbb{Q}'(o) \in [p(o), e^{\alpha}p(o)]$ . Further, we also have

$$\left|\mathbb{P}'(o) - \mathbb{Q}'(o)\right| = (e^{\alpha} - 1)p(o)\left|\mathbb{P}[\ell_o] - \mathbb{Q}[\ell_o]\right|,$$

Now, by definition,

$$D_{\mathrm{H}}^{2}\left(\mathbb{P}',\mathbb{Q}'\right) = \frac{1}{2} \int \left(\sqrt{\mathbb{P}'(o)} - \sqrt{\mathbb{Q}'(o)}\right)^{2} do = \frac{1}{2} \int \frac{|\mathbb{P}'(o) - \mathbb{Q}'(o)|^{2}}{\left(\sqrt{\mathbb{P}'(o)} + \sqrt{\mathbb{Q}'(o)}\right)^{2}} do.$$

Hence, it holds that

$$\frac{(e^{\alpha}-1)^2}{8e^{\alpha}}\int |\mathbb{P}[\ell_o] - \mathbb{Q}[\ell_o]|^2 p(o)do \le D_{\mathrm{H}}^2\left(\mathbb{P}', \mathbb{Q}'\right) \le \frac{(e^{\alpha}-1)^2}{8}\int |\mathbb{P}[\ell_o] - \mathbb{Q}[\ell_o]|^2 p(o)do$$

Notice that  $\int p(o)do \in [e^{-\alpha}, 1]$ , and hence we can normalize p to a distribution  $\bar{p}$  over  $\mathcal{O}$ . The proof of Eq. (29) is hence completed, and Eq. (30) follows similarly:

$$D_{\chi^2}(\mathbb{P}' \parallel \mathbb{Q}') = \int \frac{|\mathbb{P}'(o) - \mathbb{Q}'(o)|^2}{\mathbb{Q}'(o)} do$$
  
$$\leq (e^{\alpha} - 1)^2 \int |\mathbb{P}[\ell_o] - \mathbb{Q}[\ell_o]|^2 p(o) do$$
  
$$\leq (e^{\alpha} - 1)^2 \mathbb{E}_{o \sim \bar{p}} |\mathbb{P}[\ell_o] - \mathbb{Q}[\ell_o]|^2.$$

# I.2 Proof of Theorem 19

In this section, we provide a self-contained proof of Theorem 19, following the approach of Chen et al. [2024] (see also Appendix E.1). The proof is based on the following quantile-based private PAC-DEC.

Quantile-based private PAC-DEC. Given model class  $\mathcal{M} \subseteq (\Pi \to \Delta(\mathcal{Z}))$ , for each  $\varepsilon > 0$  and  $\delta \in [0, 1]$ , we define the quantile-based private PAC-DEC as (slightly abusing the notation)

$$\mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{LDP}}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\Pi \times \mathcal{L})}} \sup_{M \in \mathcal{M}} \left\{ \widehat{L}_{\delta}(M,p) \mid \mathbb{E}_{(\pi,\ell) \sim q} \mathsf{D}_{\ell}^{2}(M(\pi),\overline{M}(\pi)) \leq \varepsilon^{2} \right\},$$
(93)

where  $\widehat{L}_{\delta}(M, p)$  is the  $\delta$ -quantile loss of p, defined as

$$\widehat{L}_{\delta}(M,p) = \sup_{\Delta \ge 0} \{ \Delta : \mathbb{P}_{\pi \sim q}(L(M,\pi) \ge \Delta) \ge \delta \}.$$

We also denote  $\mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{LDP}}(\mathcal{M}) := \sup_{\overline{\mathcal{M}} \in \mathsf{co}(\mathcal{M})} \mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{LDP}}(\mathcal{M},\overline{\mathcal{M}})$ . By definition, the quantile-based private PAC-DEC is always bounded by the original private PAC-DEC:

$$\mathsf{p-dec}_{\varepsilon}^{\mathsf{q},\mathsf{LDP}}(\mathcal{M},\overline{M}) - \delta \le \mathsf{p-dec}_{\varepsilon,\delta}^{\mathsf{q},\mathsf{LDP}}(\mathcal{M},\overline{M}) \le \delta^{-1}\mathsf{p-dec}_{\varepsilon}^{\mathsf{q},\mathsf{LDP}}(\mathcal{M},\overline{M}).$$
(94)

However, such a conversion can be loose in general.

**Quantile lower bound.** Similar to Appendix E.1, we show that the quantile-based private PAC-DEC provides a lower bound regardless of the structure of the loss function.

**Proposition I.1** (Quantile-based private PAC-DEC lower bound). For any  $T \ge 1$  and constant  $\delta \in [0,1)$ , we denote  $\underline{\varepsilon}(T) := \frac{1}{(e^{\alpha}-1)} \sqrt{\frac{\delta}{2T}}$ . Then, for any T-round  $\alpha$ -LDP algorithm Alg, there exists  $M^{\star} \in \mathcal{M}$  such that under  $\mathbb{P}^{M^{\star}, \text{Alg}}$ ,

$$\mathbf{Risk}_{\mathsf{DM}}(T) \geq \mathsf{p-dec}_{\underline{\varepsilon}(T),\delta}^{\mathsf{q},\mathsf{LDP}}(\mathcal{M}), \qquad with \ probability \ at \ least \ \delta/2.$$

Further, for reward-based loss function L, we can relate quantile-based private PAC-DEC to the original private PAC-DEC (following Chen et al. [2024, Proposition E.1]).

**Lemma I.2.** Suppose that the loss function L is reward-based. Then, for any parameter  $\varepsilon > 0, \delta \in [0,1)$ , it holds that

$$\mathsf{p}\text{-}\mathsf{dec}_{\sqrt{2}\varepsilon,\delta}^{\mathrm{q},\mathrm{LDP}}(\mathcal{M}) \geq \mathsf{p}\text{-}\mathsf{dec}_{\varepsilon}^{\mathrm{LDP}}(\mathcal{M}) - \frac{2\sqrt{2}\varepsilon}{1-\delta}.$$

Similarly, for metric-based loss function, we have the following lemma (following Lemma E.2).

**Lemma I.3.** Suppose that the loss function L is metric-based. Then, for any parameter  $\varepsilon > 0, \delta \in [0, \frac{1}{2})$ , it holds that

$$\mathrm{p\text{-}dec}_{\varepsilon,\delta}^{\mathrm{q},\mathrm{LDP}}(\mathcal{M}) \geq \frac{1}{2}\mathrm{p\text{-}dec}_{\varepsilon}^{\mathrm{LDP}}(\mathcal{M}).$$

Therefore, the proof of Theorem 19 is completed by combining Proposition I.1 with Lemma I.2 / Lemma I.3.  $\hfill \Box$ 

#### I.2.1 Proof of Proposition I.1

We follow the strategy of Appendix E.4.

Recall that an  $\alpha$ -LDP algorithm  $\mathsf{Alg} = \{q_t\}_{t \in [T]} \cup \{p\}$  is specified by a sequence of mappings, where the *t*-th mapping  $q_t(\cdot \mid \mathcal{H}_{t-1})$  specifies the distribution of  $(\pi_t, \mathsf{Q}_t)$  based on the history  $\mathcal{H}_{t-1}$ , and the final map  $p(\cdot \mid \mathcal{H}_T)$  specifies the distribution of the  $\pi_{T+1}$  based on  $\mathcal{H}_T$ . Therefore, for any model M, we define

$$q_{M,\mathsf{Alg}} = \mathbb{E}^{M,\mathsf{Alg}} \left[ \frac{1}{T} \sum_{t=1}^{T} q_t(\cdot | \mathcal{H}_{t-1}) \right] \in \Delta(\Pi \times \mathcal{Q}), \quad p_{M,\mathsf{Alg}} = \mathbb{E}^{M,\mathsf{Alg}}[p(\mathcal{H}_T)] \in \Delta(\Pi), \tag{95}$$

The distribution  $q_{M,\text{Alg}}$  is the expected distribution of the average profile  $(\pi_1, \mathbf{Q}_1, \cdots, \pi_T, \mathbf{Q}_T)$ , and  $p_{M,\text{Alg}}$  is the expected distribution of the output policy  $\pi_{T+1}$ .

Using the chain rule of KL divergence, for any model  $M, \overline{M}$ ,

$$D_{\mathrm{KL}}(\mathbb{P}^{\overline{M},\mathrm{Alg}} \parallel \mathbb{P}^{M,\mathrm{Alg}}) = \mathbb{E}^{\overline{M},\mathrm{Alg}} \left[ \sum_{t=1}^{T} D_{\mathrm{KL}}(\mathsf{Q}_{t} \circ M(\pi_{t}) \parallel \mathsf{Q}_{t} \circ M(\pi_{t})) \right]$$
$$= T \cdot \mathbb{E}_{(\pi,\mathsf{Q}) \sim q_{\overline{M}},\mathrm{Alg}} D_{\mathrm{KL}}(\mathsf{Q} \circ \overline{M}(\pi) \parallel \mathsf{Q} \circ M(\pi)).$$

Further, by Proposition 20, for any  $\alpha$ -LDP channel Q, there exists a distribution  $\tilde{q}_{Q} \in \Delta(\mathcal{L})$  such that

$$D_{\mathrm{KL}}(\mathsf{Q} \circ \mathbb{P}_1 \parallel \mathsf{Q} \circ \mathbb{P}_2) \le (e^{\alpha} - 1)^2 \mathbb{E}_{\ell \sim \tilde{q}_{\mathsf{Q}}} \mathsf{D}_{\ell}^2(\mathbb{P}_1, \mathbb{P}_2)$$

Therefore, for any model  $M \in \mathcal{M}$ , we define  $\tilde{q}_{M,\mathsf{Alg}} \in \Delta(\Pi \times \mathcal{L})$  to be the distribution of  $(\pi, \ell)$ , where  $(\pi, \mathsf{Q}) \sim q_{M,\mathsf{Alg}}$ , and  $\ell \sim \tilde{q}_{\mathsf{Q}}$ . Then, our argument above gives

$$D_{\mathrm{KL}}(\mathbb{P}^{\overline{M},\mathrm{Alg}} \parallel \mathbb{P}^{M,\mathrm{Alg}}) \le (e^{\alpha} - 1)^2 T \cdot \mathbb{E}_{(\pi,\ell) \sim \tilde{q}_{\overline{M},\mathrm{Alg}}} \mathsf{D}^2_{\ell}(M(\pi),\overline{M}(\pi)).$$
(96)

With this chain rule, we now present the proof of Proposition I.1 (which is essentially following the analysis in Chen et al. [2024]).

**Proof of Proposition I.1.** We abbreviate  $\varepsilon = \underline{\varepsilon}(T)$ . Fix a  $\Delta < \mathsf{p-dec}_{\varepsilon,\delta}^{q,\mathsf{LDP}}(\mathcal{M})$ , and then there exists  $\overline{M}$  such that  $\Delta < \mathsf{p-dec}_{\varepsilon,\delta}^{q}(\mathcal{M},\overline{M})$ . Hence, by the definition (93), we know that

$$\Delta < \sup_{M \in \mathcal{M}} \left\{ \widehat{L}_{\delta}(M, p_{\overline{M}, \mathsf{Alg}}) \mid \mathbb{E}_{(\pi, \ell) \sim \widetilde{q}_{\overline{M}, \mathsf{Alg}}} \mathsf{D}_{\ell}^{2}(M(\pi), \overline{M}(\pi)) \leq \varepsilon^{2} \right\}.$$

Therefore, there exists  $M \in \mathcal{M}$  such that

$$\mathbb{E}_{(\pi,\ell)\sim \tilde{q}_{\overline{M},\mathsf{Alg}}} \mathsf{D}^2_{\ell}(M(\pi),\overline{M}(\pi)) \leq \varepsilon^2, \qquad \mathbb{P}_{\pi\sim p_{\overline{M}},\mathsf{Alg}}(L(M,\pi) > \Delta) \geq \delta.$$

By (96), we know

$$D_{\mathrm{KL}}(\mathbb{P}^{\overline{M},\mathrm{Alg}} \parallel \mathbb{P}^{M,\mathrm{Alg}}) \leq (e^{\alpha} - 1)^2 T \varepsilon^2.$$

By data-processing inequality, we have

$$D_{\mathrm{TV}}\left(p_{\overline{M},\mathsf{Alg}},p_{M,\mathsf{Alg}}\right) \leq D_{\mathrm{TV}}\left(\mathbb{P}^{\overline{M},\mathsf{Alg}},\mathbb{P}^{M,\mathsf{Alg}}\right) \leq \sqrt{\frac{1}{2}}(e^{\alpha}-1)^{2}T\varepsilon^{2}.$$

Therefore, combining the inequalities above, we have

$$p_{M,\mathsf{Alg}}(\pi: L(M,\pi) > \Delta) \ge p_{\overline{M},\mathsf{Alg}}(\pi: L(M,\pi) > \Delta) - \sqrt{\frac{1}{2}(e^{\alpha} - 1)^2 T \varepsilon^2} \ge \frac{\delta}{2}.$$

By the definition of  $p_{M,\text{Alg}}$ , this gives  $\mathbb{P}^{M,\text{Alg}}(L(M,\pi_{T+1}) > \Delta) \geq \frac{\delta}{2}$ . Letting  $\Delta \to \text{p-dec}_{\varepsilon,\delta}^{q}(\mathcal{M})$  completes the proof.

### I.3 Proof of Lemma 25

For each  $\theta \in [-1, 1]$ , we denote  $M_{\theta} \in \Delta(\mathcal{Z})$  to be the model given by

$$(x,y) \sim M_{\theta}$$
:  $x \sim \nu, y \sim \operatorname{Rad}(\theta x)$ .

Then it holds that

$$D_{\mathrm{TV}}(M_{\theta}, M_0) \leq \mathbb{E}_{x \sim \nu} |\theta x| = |\theta| \mathbb{E} |x|,$$

and

$$L(M_{\theta},\pi) = \mathbb{E}_{x \sim p} |x(\theta - \pi)|^2 = |\theta - \pi|^2 \cdot \mathbb{E}|x|^2.$$

Therefore, for the model class  $\mathcal{M} = \{M_{\theta} : \theta \in [-1, 1]\}$ , we can consider  $\theta = \min\left\{\frac{\varepsilon}{\mathbb{E}|x|}, 1\right\}$ , which gives

$$\mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}, M_0) \ge \mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\{M_{\theta}, M_0\}, M_0) \ge \frac{\mathbb{E}|x|^2}{4} \min\left\{\frac{\varepsilon^2}{(\mathbb{E}|x|)^2}, 1\right\}.$$

Applying Theorem 19 gives the desired lower bound.

### I.4 Proof of Theorem 26

Following Appendix F.5.3 (Lemma F.12), we consider the following information set structure  $\Psi = \Theta$ :

$$\mathcal{M}_{\theta} = \{ M \in \mathcal{M}_{\mathsf{Lin}} : \theta^{M} = \theta \}, \qquad \pi_{\theta} = \theta, \qquad \theta \in \Psi.$$

Then,  $\Psi$  is a information set structure with respect to the model class  $\mathcal{M}_{\text{Lin}}$  and value function  $V^{M}(\pi) = -L_{1}(M, \theta)$ . It is clear that  $\mathcal{M}_{\Psi} = \mathcal{M}_{\text{Lin}}$ , and hence we have the following guarantee of LDP-ExO (by Theorem F.4).

**Proposition I.4.** Let  $T \ge 1, \gamma > 0$ . Then, for linear regression under  $L_1$  loss, LDP-ExO (instantiated on  $\Psi$  defined above) achieves with probability at least  $1 - \delta$ 

$$\mathbb{E}_{\hat{\theta} \sim \hat{p}} L_1(M^\star, \hat{\theta}) \leq \mathsf{p-dec}_{c\alpha^2 \gamma}^{\mathrm{o}, \mathsf{LDP}}(\mathcal{M}_{\mathsf{Lin}}) + \frac{2\gamma \log(|\Theta|/\delta)}{T}.$$

Note that for simplicity, we assume  $\Theta$  is finite. By applying the argument on a covering of  $\Theta \subset \mathbb{B}^d(1)$ , we can regard  $\log |\Theta| \leq \tilde{O}(d)$ .

In the following, we denote  $\mathcal{M} := \mathcal{M}_{\mathsf{Lin}}$ , and it remains to upper bound  $\mathsf{p-dec}_{\gamma}^{\mathsf{o},\mathsf{LDP}}(\mathcal{M})$ . For simplicity of presentation, we assume that  $\mathcal{X} \subseteq \mathbb{B}^d(1) \setminus \{0\}$  (without loss of generality).

Fix a reference model  $\overline{M} \in \Delta(\mathcal{Z})$ . Let  $\overline{\nu} \in \Delta(\mathcal{X})$  be the marginal distribution of x under  $(x, y) \sim \overline{M}$ , and let  $f^{\overline{M}}(x) = \mathbb{E}^{\overline{M}}[y|x]$  for  $x \in \mathcal{X}$ . Note that  $f^{\overline{M}}$  is not necessarily a linear function. Further, for any  $M \in \mathcal{M}$ , we let  $\theta^M \in \mathbb{B}^d(1)$  be the associated parameter so that  $\mathbb{E}^M[y|x] = \langle \theta^M, x \rangle$ . In the following, we proceed to upper bound the offset private PAC-DEC (59) of  $\mathcal{M}$  with respect to  $\overline{M}$ , which is defined as

$$\mathsf{p}\text{-}\mathsf{dec}_{\gamma}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M},\overline{M}) := \inf_{\substack{p \in \Delta(\Pi) \\ q \in \Delta(\mathcal{L})}} \sup_{M \in \mathcal{M}} \left\{ \mathbb{E}_{\pi \sim p}[L_1(M,\pi)] - \gamma \mathbb{E}_{\ell \sim q} \mathsf{D}_{\ell}^2(M,\overline{M}) \right\}.$$

**Construction of** (p,q). The key observation is the following lemma.

**Lemma I.5.** Suppose that  $\lambda_0 > 0$  and  $\overline{\nu} \in \Delta(\mathcal{X})$  are given. Then there exists a PSD matrix  $U \in \mathbb{R}^{d \times d}$  satisfies the following equation:

$$\mathbb{E}_{x\sim\overline{\nu}}\frac{Uxx^{\top}U}{\|Ux\|} + \lambda_0 U = I_d.$$
(97)

In particular, by taking trace, it holds that  $\mathbb{E}_{x \sim \overline{\nu}} \| Ux \| \leq d$ .

We fix a  $\lambda_0 > 0$  and invoke Lemma I.5 to obtain a PSD matrix U satisfies (97). Based on the matrix U, we define the normalization map  $\mathbf{n} : \mathbb{R}^d \to \mathbb{B}^d(1)$  as  $\mathbf{n}(v) = \frac{Uv}{\|Uv\|}$  for any vector  $v \neq 0$ . Then, Eq. (97) ensures that

$$\mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x) x^{\top} + \lambda_0 I = U^{-1}.$$
(98)

To construct a distribution  $q \in \Delta(\mathcal{L})$ , we invoke the following lemma.

**Lemma I.6.** Suppose that  $\mathbf{v} : \mathcal{Z} \to \mathbb{B}^D(1)$ . Then there exists a distribution  $Q(\mathbf{v})$  over  $\mathcal{L} = (\mathcal{Z} \to [0,1])$ , such that for any  $\mathbb{P}, \mathbb{Q} \in \Delta(\mathcal{Z})$ , it holds

$$\mathbb{E}_{\ell \sim Q(\mathbf{v})} \mathsf{D}_{\ell}^{2}(\mathbb{P}, \mathbb{Q}) \geq \|\mathbb{P}[\mathbf{v}] - \mathbb{Q}[\mathbf{v}]\|^{2},$$

where we denote  $\mathbb{P}[\mathbf{v}] := \mathbb{E}_{z \sim \mathbb{P}} \mathbf{v}(z)$ .

To apply Lemma I.6, we define maps

$$\mathbf{v}_1(z) = \mathbf{n}(x) \cdot y, \quad \mathbf{v}_2(z) = \operatorname{vec}(\mathbf{n}(x)x^{\top}), \quad \mathbf{v}_3(z) = \left[\mathbf{1}\left\{\|Ux\| < \gamma\right\}\gamma^{-1}\|Ux\|, \mathbf{1}\left\{\|Ux\| \ge \gamma\right\}\right],$$

and then by Lemma I.6, there exists a distribution  $q \in \Delta(\mathcal{L})$  such that for any model  $M \in \mathcal{M}$ ,

$$\mathbb{E}_{\ell \sim q} \mathsf{D}_{\ell}^{2}(M, \overline{M}) \geq \frac{1}{3} \sum_{i=1}^{3} \|\mathbb{E}_{z \sim M} \mathbf{v}_{i}(z) - \mathbb{E}_{z \sim \overline{M}} \mathbf{v}_{i}(z)\|^{2}.$$

For notational simplicity, in the following, we denote  $\varepsilon_{M,i}^2 := \|M[\mathbf{v}_i] - \overline{M}[\mathbf{v}_i]\|^2$  for each  $i \in \{1, 2, 3\}$  and each  $M \in \mathcal{M}$ . Then, by definition, we know

$$\begin{split} & \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x) \left( \langle \theta^{M}, x \rangle - f^{\overline{M}}(x) \right) \right\| \\ & \leq \left\| \mathbb{E}_{x \sim \nu_{M}} \mathsf{n}(x) \langle \theta^{M}, x \rangle - \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x) \langle \theta^{M}, x \rangle \right\| + \left\| \mathbb{E}_{x \sim \nu_{M}} \mathsf{n}(x) \langle \theta^{M}, x \rangle - \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x) f^{\overline{M}}(x) \right\| \\ & = \left\| \langle \mathbb{E}_{x \sim \nu_{M}} \mathsf{n}(x) x - \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x) x, \theta^{M} \rangle \right\| + \left\| \mathbb{E}_{(x,y) \sim M} \mathsf{n}(x) y - \mathbb{E}_{(x,y) \sim \overline{M}} \mathsf{n}(x) y \right\| \\ & \leq \left\| \mathbb{E}_{x \sim \nu_{M}} \mathsf{n}(x) x - \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x) x \right\|_{F} + \left\| \mathbb{E}_{(x,y) \sim M} \mathsf{n}(x) y - \mathbb{E}_{(x,y) \sim \overline{M}} \mathsf{n}(x) y \right\| \\ & = \varepsilon_{M,2} + \varepsilon_{M,1}. \end{split}$$

Therefore, we define

$$\bar{\theta} = \underset{\theta \in \mathbb{B}^{d}(1)}{\arg\min} \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x) \left( \langle \theta, x \rangle - f^{\overline{M}}(x) \right) \right\|^{2}.$$

Notice that the objective function above is a quadratic function of  $\theta$ , we know that for any  $M \in \mathcal{M}$ ,

$$\left\|\mathbb{E}_{x\sim\overline{\nu}}\mathsf{n}(x)\langle\theta^{M}-\bar{\theta},x\rangle\right\|\leq\left\|\mathbb{E}_{x\sim\overline{\nu}}\mathsf{n}(x)\big(\langle\theta^{M},x\rangle-f^{\overline{M}}(x)\big)\right\|\leq\varepsilon_{M,1}+\varepsilon_{M,2}$$

Using Eq. (98), we then have

$$\left\| U^{-1}(\theta^{M} - \bar{\theta}) \right\| = \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x) \langle \theta^{M} - \bar{\theta}, x \rangle + \lambda_{0}(\theta^{M} - \bar{\theta}) \right\| \le \varepsilon_{M,1} + \varepsilon_{M,2} + 2\lambda_{0}.$$

We let p be supported on  $\bar{\theta}$ .

Bounding the offset DEC risk. Define  $\mathcal{X}_0 := \{x : ||Ux|| < \gamma\}$ . For any  $M \in \mathcal{M}$ , we bound

$$L_{1}(M,\bar{\theta}) = \mathbb{E}_{x \sim \nu_{M}} \left| \langle x,\bar{\theta}-\theta^{M} \rangle \right|$$
  

$$\leq \mathbb{E}_{x \sim \nu_{M}} \left[ 2 \cdot \mathbf{1} \left\{ x \notin \mathcal{X}_{0} \right\} + \mathbf{1} \left\{ x \in \mathcal{X}_{0} \right\} \left| \langle x,\bar{\theta}-\theta^{M} \rangle \right| \right]$$
  

$$\leq 2\mathbb{E}_{x \sim \nu_{M}} \mathbf{1} \left\{ x \notin \mathcal{X}_{0} \right\} + \left(\mathbb{E}_{x \sim \nu_{M}} \mathbf{1} \left\{ x \in \mathcal{X}_{0} \right\} \|Ux\|\right)^{1/2} \left(\mathbb{E}_{x \sim \nu_{M}} \frac{\langle x,\bar{\theta}-\theta^{M} \rangle^{2}}{\|Ux\|}\right)^{1/2}$$

First, notice that  $\mathbb{E}_{x \sim \overline{\nu}} \mathbf{1} \{ x \notin \mathcal{X}_0 \} \leq \frac{1}{\gamma} \mathbb{E}_{x \sim \overline{\nu}} \| U x \|$ , and hence

$$\mathbb{E}_{x \sim \nu_M} \mathbf{1} \{ x \notin \mathcal{X}_0 \}$$
  
$$\leq |\mathbb{E}_{x \sim \nu_M} \mathbf{1} \{ x \notin \mathcal{X}_0 \} - \mathbb{E}_{x \sim \overline{\nu}} \mathbf{1} \{ x \notin \mathcal{X}_0 \} | + \mathbb{E}_{x \sim \overline{\nu}} \mathbf{1} \{ x \notin \mathcal{X}_0 \} \leq \varepsilon_{M,3} + \frac{d}{\gamma}.$$

Similarly,

$$\mathbb{E}_{x \sim \nu_M} \mathbf{1} \{ x \in \mathcal{X}_0 \} \| U x \|$$
  
$$\leq \| \mathbb{E}_{x \sim \nu_M} \mathbf{1} \{ x \in \mathcal{X}_0 \} \| U x \| - \mathbb{E}_{x \sim \overline{\nu}} \mathbf{1} \{ x \in \mathcal{X}_0 \} \| U x \| + \mathbb{E}_{x \sim \overline{\nu}} \mathbf{1} \{ x \in \mathcal{X}_0 \} \| U x \| \leq \gamma \varepsilon_{M,3} + d.$$

Next, we denote  $H_M = \mathbb{E}_{x \sim \nu_M} \frac{xx^\top}{\|Ux\|}$ , and we bound

$$\mathbb{E}_{x \sim \nu_M} \frac{\langle x, \theta - \theta^M \rangle^2}{\|Ux\|} = \left\| \bar{\theta} - \theta^M \right\|_{H_M}^2$$
$$= \left\| \bar{\theta} - \theta^M \right\|_{H_M}^2 - \left\| \bar{\theta} - \theta^M \right\|_{H_{\overline{M}}}^2 + \left\| \bar{\theta} - \theta^M \right\|_{H_{\overline{M}}}^2$$

$$\leq \left\| (H_M - H_{\overline{M}})(\overline{\theta} - \theta^M) \right\| + \left\| U^{-1}(\overline{\theta} - \theta^M) \right\|^2,$$

where the second inequality uses  $H_{\overline{M}} = \mathbb{E}_{x \sim \overline{\nu}} \frac{xx^{\top}}{\|\overline{U}x\|} \preceq U^{-2}$  (by Eq. (97)). Notice that

$$\|U(H_M - H_{\overline{M}})\|_F = \|\mathbb{E}_{x \sim \nu_M} \mathsf{n}(x)x - \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}(x)x\|_F \le \varepsilon_{M,2},$$

and hence

$$\left\| (H_M - H_{\overline{M}})(\overline{\theta} - \theta^M) \right\| \le \left\| U(H_M - H_{\overline{M}}) \right\|_F \left\| U^{-1}(\overline{\theta} - \theta^M) \right\|.$$

Combining the inequalities above, we can conclude that

$$L_1(M,\bar{\theta}) \lesssim \varepsilon_{M,3} + \frac{d}{\gamma} + \sqrt{(d+\gamma\varepsilon_{M,3})}(\varepsilon_{M,1} + \varepsilon_{M,2} + \lambda_0).$$

Therefore, by applying the weighted AM-GM inequality, we have

$$L_1(M,\bar{\theta}) - \frac{\gamma}{3}(\varepsilon_{M,1}^2 + \varepsilon_{M,2}^2 + \varepsilon_{M,3}^2) \lesssim \frac{d}{\gamma} + \sqrt{d+\gamma}\lambda_0.$$

and hence  $\mathbf{p}-\mathbf{dec}_{\gamma}^{\mathrm{o},\mathrm{LDP}}(\mathcal{M},\overline{M}) \leq C\left(\frac{d}{\gamma} + \sqrt{d+\gamma}\lambda_0\right)$  for some absolute constant C. Taking  $\lambda_0 \to 0$  gives  $\mathbf{p}-\mathbf{dec}_{\gamma}^{\mathrm{o},\mathrm{LDP}}(\mathcal{M},\overline{M}) \leq \frac{Cd}{\gamma}$ .

Finalizing the proof. We have shown that  $p-dec_{\gamma}^{0,LDP}(\mathcal{M}) \leq C\frac{d}{\gamma}$ . In particular, this implies  $p-dec_{\varepsilon}^{LDP}(\mathcal{M}) \leq \sqrt{Cd\varepsilon}$  for any  $\varepsilon \in [0,1]$ .

Further, by Proposition I.4, LDP-ExO achieves with probability at least  $1 - \delta$  that

$$\mathbb{E}_{\pi \sim \widehat{p}} L_1(M^\star, \pi) \le O(1) \cdot \left[ \frac{d}{\alpha^2 \gamma} + \frac{2\gamma \log(|\Theta|/\delta)}{T} \right].$$

Note that  $L_1(M^*, \pi)$  is a convex function with respect to  $\pi \in \mathbb{B}^d(1)$ , and hence we can let LDP-ExO output  $\hat{\theta} = \mathbb{E}_{\pi \sim \hat{p}}[\pi] \in \mathbb{B}^d(1)$ . Then, by choosing  $\gamma > 0$  suitably, it is guaranteed that with probability at least  $1 - \delta$ 

$$L_1(M^\star, \hat{\theta}) \le \tilde{O}\left(\sqrt{\frac{d^2 \log(1/\delta)}{\alpha^2 T}}\right)$$

The proof of Theorem 26 is hence completed.

## I.4.1 Proof of Lemma I.5

Consider the compact, convex region  $\mathcal{U} \subset \mathbb{R}^{d \times d}$  given by

$$\mathcal{U} = \{ U : (1+\lambda_0)^{-1/2} I_d \leq U \leq \lambda_0^{-1/2} I_d \}.$$
(99)

Define function  $F: \mathcal{U} \to \mathbb{R}^{d \times d}$  as follows:

$$F(U) := \left( \mathbb{E}_{x \sim \overline{\nu}} \frac{U^{1/2} x x^{\top} U^{1/2}}{\|Ux\|} + \lambda_0 I_d \right)^{-1}.$$

Note that by definition, for any  $U \in \mathcal{U}$ ,  $\lambda_0 I_d \preceq F(U)^{-1} \preceq (\lambda_0 + 1)I_d$ . Therefore, F maps  $\mathcal{U}$  to itself. Further, the map  $(U, x) \mapsto \frac{U^{1/2}xx^{\top}U^{1/2}}{\|Ux\|}$  is uniformly continuous with respect to  $U \in \mathcal{U}$  and  $x \neq 0$ . Therefore, F(U) is continuous in U, and Brouwer fixed-point theorem implies that there exists  $U \in \mathcal{U}$  such that F(U) = U, i.e., U satisfies Eq. (97).

### I.4.2 Proof of Lemma I.6

Define s(t) = 1 if  $t \ge 0$ , and s(t) = 0 otherwise.

Fix the map  $\mathbf{v}: \mathcal{Z} \to \mathbb{B}^D(1)$ . For each  $w \in \mathbb{R}^D$ , we define  $\ell_w \in \mathcal{L}$  as

$$\ell_w(z) = \|\mathbf{v}(z)\| \, \mathbf{s}(\langle \mathbf{v}(z), w \rangle), \qquad \forall z \in \mathcal{Z}.$$

Then, we define  $Q(\mathbf{v}) \in \Delta(\mathcal{L})$  to be the distribution of  $\ell = \ell_w$  with  $w \sim \mathsf{N}(0, I_D)$ .

Now, for any  $x \in \mathbb{R}^D$  with ||x|| = 1 and any fixed  $z \in \mathcal{Z}$ , it holds

$$\mathbb{E}_w \ell_w(z) \langle w, x \rangle = \langle \mathbf{v}(z), x \rangle,$$

where  $\mathbb{E}_w$  is taken over  $w \sim \mathsf{N}(0, I_D)$  and the equality follows from the rotational invariance of the Gaussian distribution.

Therefore, for the distribution  $Q(\mathbf{v})$  defined above, we have

Hence, by the arbitrariness of x, we have

$$\|\mathbb{P}[\mathbf{v}] - \mathbb{Q}[\mathbf{v}]\| = \sup_{x \in \mathbb{R}^D : \|x\| = 1} \langle \mathbb{P}[\mathbf{v}] - \mathbb{Q}[\mathbf{v}], x \rangle \le \left(\mathbb{E}_{\ell \sim Q(\mathbf{v})} \mathsf{D}_{\ell}^2(\mathbb{P}, \mathbb{Q})\right)^{\frac{1}{2}}.$$

## I.5 Lower bound for LDP learning linear models

Fix  $d \geq 1$  and  $\Delta \in [0,1]$ . Let  $\Theta = \{-\Delta, \Delta\}^d$  and  $\mathcal{X} = [d]$ , and for each  $\theta \in \Theta$ , we define  $f_{\theta}$  as

$$f_{\theta}(i) = \theta_i,$$

and let  $M_{\theta} \in \Delta(\mathcal{Z})$  given by

$$(x, y) \sim M_{\theta}$$
:  $x \sim \text{Unif}(\mathcal{X}), y | x \sim \text{Rad}(f_{\theta}(x)).$ 

Then, we let  $\mathcal{F} = \{f_{\theta} : \theta \in \Theta\} \subseteq (\mathcal{X} \to [-1,1])$ , and  $\mathcal{M}_d := \{M_{\theta} : \theta \in \Theta\}$  is the class of well-specified models (with respect to  $\mathcal{F}$ ) with covariate distribution  $\text{Unif}(\mathcal{X})$ .

Recall that for such a problem class, the decision space is  $\Pi \subseteq (\mathcal{X} \to [-1, 1])$ , which can be naturally identified as a subset of  $[-1, 1]^d$ . Then, the loss function is given by

$$L(M_{\theta}, \pi) = \mathbb{E}_{x \sim \text{Unif}(\mathcal{X})} |\pi(x) - f_{\theta}(x)| = \frac{1}{d} \sum_{i=1}^{d} |\pi(i) - \theta_i|$$

**Proposition I.7.** Let  $T \ge 1$ ,  $\Delta \in (0,1]$ . Suppose that Alg is a T-round  $\alpha$ -LDP algorithm, such that  $\mathbb{E}^{M,\text{Alg}}[\operatorname{Risk}_{\mathsf{DM}}(T)] \le \frac{\Delta}{4}$  for all  $M \in \mathcal{M}_d$ . Then it holds that  $T \ge \Omega\left(\frac{d^2}{\alpha^2 \Delta^2}\right)$ .

*Proof.* For each  $\theta \in \Theta$ , we consider

$$p_{\theta} := \mathbb{P}^{M_{\theta}, \operatorname{Alg}}(\pi_{T+1} = \cdot) \in \Delta(\Pi).$$

Recall the chain rule of KL divergence and Proposition 20: for any model  $M, \overline{M}$ ,

$$D_{\mathrm{KL}}(\mathbb{P}^{M,\mathrm{Alg}} \parallel \mathbb{P}^{\overline{M},\mathrm{Alg}}) = \mathbb{E}^{M,\mathrm{Alg}} \left[ \sum_{t=1}^{T} D_{\mathrm{KL}}(\mathbb{Q}^{t} \circ M \parallel \mathbb{Q}^{t} \circ \overline{M}) \right]$$
$$\leq T(e^{\alpha} - 1)^{2} D_{\mathrm{TV}}\left(M,\overline{M}\right)^{2}.$$

Therefore, by data-processing inequality, we have

$$D_{\mathrm{TV}}(p_{\theta}, p_{\theta'}) \le (e^{\alpha} - 1)\sqrt{T} D_{\mathrm{TV}}(M_{\theta}, M_{\theta'}) = (e^{\alpha} - 1)\sqrt{T} \cdot \frac{1}{d} \left\| \theta - \theta' \right\|_{1}.$$

We further denote  $p_{\theta,i} = p_{\theta}(\pi(i) \ge 0)$ . Then it holds that

$$\mathbb{E}^{M_{\theta}, \operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] = \mathbb{E}_{\pi \sim p_{\theta}} L(M_{\theta}, \pi)$$
$$= \frac{1}{d} \sum_{i=1}^{d} \mathbb{E}_{\pi \sim p_{\theta}} |\pi(i) - \theta_{i}|$$
$$\geq \frac{\Delta}{d} \left( \sum_{i:\theta_{i}=-\Delta} p_{\theta,i} + \sum_{i:\theta_{i}=+\Delta} (1 - p_{\theta,i}) \right).$$

Thus,

$$\sum_{\theta \in \Theta} \mathbb{E}^{M_{\theta}, \operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] \geq \frac{\Delta}{d} \sum_{i=1}^{d} \left( \sum_{\theta: \theta_{i} = -\Delta} p_{\theta,i} + \sum_{\theta: \theta_{i} = +\Delta} (1 - p_{\theta,i}) \right)$$
$$= \frac{\Delta}{d} \sum_{i=1}^{d} \left( 2^{d-1} - \sum_{(\theta, \theta') \in \Theta_{i}} (p_{\theta,i} - p_{\theta',i}) \right),$$

where  $\Theta_i = \{(\theta, \theta') : \theta_i = +\Delta, \theta'_i = -\Delta, \text{and } \forall j \neq i, \theta_j = \theta'_j\}$ . Notice that for any  $(\theta, \theta') \in \Theta_i$ , we have

$$\left|p_{\theta,i} - p_{\theta',i}\right| \le D_{\mathrm{TV}}\left(p_{\theta}, p_{\theta'}\right) \le \frac{2\Delta(e^{\alpha} - 1)\sqrt{T}}{d}$$

Therefore, we have

$$\sum_{\theta \in \Theta} \mathbb{E}^{M_{\theta}, \operatorname{Alg}}[\operatorname{\mathbf{Risk}}_{\mathsf{DM}}(T)] \ge \Delta \cdot 2^{d-1} \left( 1 - \frac{2\Delta(e^{\alpha} - 1)\sqrt{T}}{d} \right),$$

which immediately implies

$$\max_{M \in \mathcal{M}} \mathbb{E}^{M_{\theta}, \mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \geq \frac{\Delta}{2} \left( 1 - \frac{2\Delta(e^{\alpha} - 1)\sqrt{T}}{d} \right)$$

Hence, we must have  $T \ge \frac{d^2}{4\Delta^2(e^{\alpha}-1)^2}$ .

.

Choosing  $\Delta = \min\left\{\frac{d}{4(e^{\alpha}-1)\sqrt{T}}, \frac{1}{\sqrt{d}}\right\}$  in the proof above (which ensures  $\Theta \subset \mathbb{B}^d(1)$ ), we have the following corollary.

**Corollary I.8.** There exists a covariate distribution  $\mu$  over  $\mathbb{B}^d(1)$ , such that for the model class  $\mathcal{M}$  consisting of the linear models with covariate distribution  $\mu$ , any *T*-round  $\alpha$ -LDP algorithm with output  $\hat{\theta}$ , it holds that

$$\sup_{M^{\star} \in \mathcal{M}} \mathbb{E}^{M^{\star}, \operatorname{Alg}} L_1(M^{\star}, \hat{\theta}) \gtrsim \min\left\{\frac{d}{\alpha\sqrt{T}}, \frac{1}{\sqrt{d}}\right\}$$

This lower bounds implies that the upper bound of Theorem 26 is nearly minimax-optimal (up to logarithmic factors).

## I.6 Proof of Theorem 30

We claim that the private regret-DEC of  $\mathcal{M}$  can be bounded as

$$\mathsf{r}\mathsf{-dec}^{\mathsf{LDP}}_{\varepsilon}(\mathcal{M}) = \sup_{\overline{M} \in \mathrm{co}(\mathcal{M})} \mathsf{r}\mathsf{-dec}^{\mathsf{LDP}}_{\varepsilon}(\mathcal{M} \cup \{\overline{M}\}, \overline{M}) \le (20d+6)\varepsilon, \qquad \forall \varepsilon \in [0,1].$$
(100)

With Eq. (100), we may directly apply Proposition 29, as  $\log N_{\infty}(\mathcal{F}_{\text{Lin}}, \Delta) \leq O(d \log(1/\Delta))$ .

In the following, it remains to prove Eq. (100). We only need to upper bound  $\operatorname{r-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M} \cup \{\overline{M}\}, \overline{M})$  for any fixed reference model  $\overline{M} \in \operatorname{co}(\mathcal{M})$ . Following the proof of Theorem 26 (Appendix I.4), we assume that  $\phi(x, a) \neq 0$  for all  $x \in \mathcal{X}$ ,  $a \in \mathcal{A}$  without loss of generality.

Fix a reference model  $\overline{M} = \mathbb{E}_{M \sim \mu}[M] \in \operatorname{co}(\mathcal{M})$  and  $\varepsilon \in [0, 1]$ . Note that  $\overline{M}$  is also associated with a mean reward function  $f^{\overline{M}}$  (not necessarily in  $\mathcal{F}_{\mathsf{Lin}}$ ) and a context distribution  $\overline{\nu} \in \Delta(\mathcal{X})$ . For any  $M \in \mathcal{M}$ , we let  $\theta^M \in \mathbb{B}^d(1)$  be the associated parameter so that the mean reward function  $f^M \in \mathcal{F}$ is given by  $f^M(x, a) = \langle \theta^M, \phi(x, a) \rangle$ .

In the following, we proceed to upper bound the private regret-DEC :

$$\mathsf{r-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M} \cup \{\overline{M}\}, \overline{M}) \\ \coloneqq \inf_{p \in \Delta(\Pi \times \mathcal{L})} \sup_{M \in \mathcal{M} \cup \{\overline{M}\}} \left\{ \mathbb{E}_{\pi \sim p}[V^{M}(\pi^{M}) - V^{M}(\pi)] \mid \mathbb{E}_{(\pi,\ell) \sim p} \mathsf{D}_{\ell}^{2}(M(\pi), \overline{M}(\pi)) \leq \varepsilon^{2} \right\}.$$

For notational simplicity, for any  $p \in \Delta(\Pi \times \mathcal{L})$ , we denote

$$\mathcal{M}_{p,\varepsilon^2}(\overline{M}) := \left\{ M \in \mathcal{M} : \mathbb{E}_{(\pi,\ell) \sim p} \mathsf{D}^2_{\ell}(M(\pi), \overline{M}(\pi)) \le \varepsilon^2 \right\}.$$

Let  $0 < \lambda_0 < \min\left\{\frac{\varepsilon^2}{100}, \frac{\varepsilon}{\log |\mathcal{A}|}\right\}$  be a sufficiently small, fixed parameter. Let  $\lambda = 5\varepsilon$ .

In the following, for notational simplicity, we denote  $\phi(x,\pi) := \phi(x,\pi(x)) \in \mathbb{R}^d$ . We first invoke the following corollary of Lemma I.5.

**Lemma I.9.** Suppose that  $\lambda_0 > 0$  and  $\overline{\nu} \in \Delta(\mathcal{X})$  are given. Then by Lemma I.5, for each  $\pi \in \Pi$ , there exists a PSD matrix  $U_{\pi} \in \mathbb{R}^{d \times d}$  satisfies the following equation:

$$\mathbb{E}_{x \sim \overline{\nu}} \frac{U_{\pi} \phi(x, \pi) \phi(x, \pi)^{\top} U_{\pi}}{\|U_{\pi} \phi(x, \pi)\|} + \lambda_0 U_{\pi} = I_d.$$
(101)

We further define  $\mathbf{n}_{\pi}(x) := \left[1; \frac{U_{\pi}\phi(x,\pi)}{\|U_{\pi}\phi(x,\pi)\|}\right]$ .

**Fixed point argument.** Our proof strategy is that, for any distribution  $P \in \Delta(\Pi)$ , we define a "refinement"  $F(P) \in \Delta(\Pi)$  of P. Then, the fixed point of F is a distribution of good properties.

(1) Define the constrained set

$$\Theta_{\overline{M}} := \{\theta \in \mathbb{B}^d(1) : \left| \mathbb{E}_{x \sim \overline{\nu}} \langle \theta, \phi(x, \pi^{\overline{M}}) \rangle - V^{\overline{M}}(\pi^{\overline{M}}) \right| \le 4\varepsilon \}.$$

Then, for each  $P \in \Delta(\Pi)$ , we define

$$\widehat{\theta}_{P} := \underset{\theta \in \Theta_{\overline{M}}}{\operatorname{arg\,min}} L_{P}(\theta) := \mathbb{E}_{\pi \sim P} \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \left( \langle \theta, \phi(x, \pi) \rangle - f^{\overline{M}}(x, \pi(x)) \right) \right\|^{2} + \lambda_{0}^{2} \|\theta\|^{2} \,. \tag{102}$$

By the strong convexity of  $L_P$ ,  $\hat{\theta}_P$  is a continuous function of  $P \in \Delta(\Pi)$ .

(2) Define  $\Sigma_P := \mathbb{E}_{\pi \sim P} U_{\pi}^{-2}$  and

$$\widehat{f}_P(x,a) := \langle \widehat{\theta}_P, \phi(x,a) \rangle + 2\min\{\lambda \| \phi(x,a) \|_{\Sigma_P^{-1}}, 1\}, \qquad \forall (x,a) \in \mathcal{X} \times \mathcal{A}.$$
(103)

(3) For each  $x \in \mathcal{X}$ , we define

$$Q_P(\cdot|x) := \underset{q \in \Delta(\mathcal{A})}{\operatorname{arg\,max}} \ \mathbb{E}_{a \sim q} \widehat{f}_P(x, a) + \lambda_0 H(q), \tag{104}$$

where  $H(q) = -\sum_{a \in \mathcal{A}} q(a) \log q(a)$  is the entropy of  $q \in \Delta(\mathcal{A})$ . Notice that the objective function is strongly concave with respect to  $q \in \Delta(\mathcal{A})$ , and hence  $Q_P$  is continuous with respect to P.

(4) Finally, define  $F(P) = \bigotimes_{x \in \mathcal{X}} Q_P(\cdot | x) \in \Delta(\Pi)$ . Formally, we define  $F(P) \in \Delta(\Pi)$  as<sup>14</sup>

$$F(P)[\pi] := \prod_{x \in \mathcal{X}} Q_P(\pi(x)|x).$$

By definition,  $F : \Delta(\Pi) \to \Delta(\Pi)$  is continuous, and hence by Theorem D.3, there exists  $P \in \Delta(\Pi)$  such that F(P) = P. In the following, we work with such a fixed-point distribution P.

We start with the following lemmas.

**Lemma I.10.** (1) For any 
$$M \in \mathcal{M}$$
 with  $\left\| \theta^M - \widehat{\theta}_P \right\|_{\Sigma_P} \leq \lambda$ , it holds that  
 $V^M(\pi^M) - \mathbb{E}_{\pi \sim P} V^M(\pi) \leq 4\mathbb{E}_{\pi \sim P} \mathbb{E}_{x \sim \nu_M} \min\{\lambda \| \phi(x, \pi) \|_{\Sigma_P^{-1}}, 1\} + \varepsilon.$ 

(2) It holds that

$$\mathbb{E}_{\pi \sim P} \mathbb{E}_{x \sim \overline{\nu}} \| \phi(x, \pi) \|_{\Sigma_{p}^{-1}} \leq d.$$

<sup>&</sup>lt;sup>14</sup>Alternatively, we can also define  $F(P) \in \Delta(\Pi)$  as the distribution of  $\pi$  generated as  $\pi(x) \sim Q_P(\cdot|x)$  independently for  $x \in \mathcal{X}$ .
**Construction of** *p*. First, we define a distribution  $\bar{p} \in \Delta(\Pi \times \mathcal{L})$  as follows. Consider the maps

$$\bar{\mathbf{v}}(z) = [r, \phi(x, \pi^{\overline{M}})],$$

and Lemma I.6 implies that there exists  $\bar{q} \in \Delta(\mathcal{L})$  such that for any model  $M \in \mathcal{M}, \pi \in \Pi$ ,

$$\mathbb{E}_{\ell \sim \bar{q}} \mathsf{D}_{\ell}^{2}(M(\pi), \overline{M}(\pi)) \geq \frac{1}{2} \left\| \mathbb{E}_{z \sim M(\pi)} \bar{\mathbf{v}}(z) - \mathbb{E}_{z \sim \overline{M}(\pi)} \bar{\mathbf{v}}(z) \right\|^{2}.$$

We then choose  $\bar{p} \in \Delta(\Pi \times \mathcal{L})$  to be the distribution of  $(\pi^{\overline{M}}, \ell)$  under  $\ell \sim \bar{q}$ .

**Lemma I.11.** Suppose that  $M \in \mathcal{M}_{\bar{p},2\varepsilon^2}(\overline{M})$ . Then  $\theta^M \in \Theta_{\overline{M}}$  and  $\left|V^M(\pi^{\overline{M}}) - V^{\overline{M}}(\pi^{\overline{M}})\right| \leq 2\varepsilon$ .

Next, we define  $p^* \in \Delta(\Pi \times \mathcal{L})$  as follows. For each policy  $\pi \in \Pi$ , we define a map  $\mathbf{v}_{\pi} : \mathcal{Z} \to \mathbb{R}^{(d+1)^2+1}$ :

$$\mathbf{v}_{\pi}(z) := [ \mathbf{n}_{\pi}(x) \cdot r; \operatorname{vec}(\mathbf{n}_{\pi}(x)\phi(x,\pi)^{\top}); \min\{\|\phi(x,\pi)\|_{\Sigma_{P}^{-1}}, 1\}],\$$

and Lemma I.6 implies that there exists  $q_{\pi} \in \Delta(\mathcal{L})$  such that for any model  $M \in \mathcal{M}, \pi \in \Pi$ ,

$$\mathbb{E}_{\ell \sim q_{\pi}} \mathsf{D}_{\ell}^{2}(M(\pi), \overline{M}(\pi)) \geq \frac{1}{5} \left\| \mathbb{E}_{z \sim M(\pi)} \mathbf{v}_{\pi}(z) - \mathbb{E}_{z \sim \overline{M}(\pi)} \mathbf{v}_{\pi}(z) \right\|^{2}.$$
 (105)

We then define  $p^* \in \Delta(\Pi \times \mathcal{L})$  to be the distribution of  $(\pi, \ell)$  under  $\pi \sim P$ ,  $\ell \sim q_{\pi}$ . We summarize the properties of  $p^*$  in the following lemma.

**Lemma I.12.** Suppose that  $M \in \mathcal{M}_{p^*, 2\varepsilon^2}(\overline{M})$ . Then it holds that

$$\mathbb{E}_{\pi \sim P} \left| V^M(\pi) - V^{\overline{M}}(\pi) \right| \le 2\varepsilon, \tag{106}$$

$$\mathbb{E}_{\pi \sim P} \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \left( \langle \theta, \phi(x, \pi) \rangle - f^{\overline{M}}(x, \pi(x)) \right) \right\|^2 \le 20\varepsilon^2, \tag{107}$$

$$\mathbb{E}_{\pi \sim P} \left| \mathbb{E}_{x \sim \nu_M} \min\{\lambda \| \phi(x, \pi) \|_{\Sigma_P^{-1}}, 1\} - \mathbb{E}_{x \sim \nu_{\overline{M}}} \min\{\lambda \| \phi(x, \pi) \|_{\Sigma_P^{-1}}, 1\} \right| \le 4\varepsilon.$$
(108)

In particular, when  $\theta^M \in \Theta_{\overline{M}}$ , it holds that  $\left\| \theta^M - \widehat{\theta}_P \right\|_{\Sigma_P} \leq 5\varepsilon = \lambda$ .

Now, we consider three cases. Define  $M_P \in \mathcal{M}$  be the model with context distribution  $\overline{\nu}$  and parameter  $\hat{\theta}_P$ , and let

$$\varepsilon_P := \mathbb{E}_{\pi \sim P} \left| V^{M_P}(\pi) - V^{\overline{M}}(\pi) \right|.$$

**Case 1:**  $\Theta_{\overline{M}} = \emptyset$ . In this case, the set  $\mathcal{M}_{\overline{p},2\varepsilon^2}(\overline{M}) = \emptyset$  by Lemma I.11. Therefore, we can set  $p = \overline{p}$  and bound

$$\mathsf{r-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}\cup\{\overline{M}\},\overline{M}) \leq \sup_{M\in\mathcal{M}_{\bar{p},\varepsilon^2}(\overline{M})\cup\{\overline{M}\}} \mathbb{E}_{\pi\sim\bar{p}}[V^M(\pi^M) - V^M(\pi)] = \mathbb{E}_{\pi\sim\bar{p}}\left[V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi)\right] = 0$$

**Case 2:**  $\Theta_{\overline{M}} \neq \emptyset$  and  $\varepsilon_P \leq 5\varepsilon$ . In this case, we set  $p = \frac{\overline{p} + p^*}{2}$ . We proceed to upper bound  $V^M(\pi^M) - \mathbb{E}_{\pi \sim p} V^M(\pi)$  for any  $M \in \mathcal{M}_{p,\varepsilon^2}(\overline{M}) \cup \{\overline{M}\}$ .

**Case 2(a):**  $M \in \mathcal{M}_{p,\varepsilon^2}(\overline{M})$ . Then, we know that  $\left\|\theta^M - \widehat{\theta}_P\right\|_{\Sigma_P} \leq 5\varepsilon = \lambda$  by Lemma I.12. Then, invoking Lemma I.10 gives

$$V^{M}(\pi^{M}) - \mathbb{E}_{\pi \sim P} V^{M}(\pi) \leq 4\mathbb{E}_{\pi \sim P} \mathbb{E}_{x \sim \nu_{M}} \min\{\lambda \| \phi(x, \pi) \|_{\Sigma_{P}^{-1}}, 1\} + \varepsilon$$

 $\leq 4\varepsilon + 5d\lambda,$ 

where the second inequality uses Eq. (108) and Lemma I.10 (2).

Therefore, it remains to upper bound  $V^{M}(\pi^{M}) - V^{M}(\pi^{\overline{M}})$ . Combining Eq. (106) and Lemma I.11 and the fact that  $V^{\overline{M}}(\pi) \leq V^{\overline{M}}(\pi^{\overline{M}})$ , we have

$$\mathbb{E}_{\pi \sim P} V^{M}(\pi) - V^{M}(\pi^{\overline{M}}) = \mathbb{E}_{\pi \sim P} [V^{M}(\pi) - V^{\overline{M}}(\pi) + V^{\overline{M}}(\pi) - V^{\overline{M}}(\pi^{\overline{M}})] + V^{\overline{M}}(\pi^{\overline{M}}) - V^{M}(\pi^{\overline{M}}) \le 5\varepsilon.$$

To conclude, we have

$$V^{M}(\pi^{M}) - \mathbb{E}_{\pi \sim p} V^{M}(\pi) = V^{M}(\pi^{M}) - \mathbb{E}_{\pi \sim P} V^{M}(\pi) + \frac{1}{2} \left[ \mathbb{E}_{\pi \sim P} V^{M}(\pi) - V^{M}(\pi^{\bar{M}}) \right] \\ \leq (20d + 6)\varepsilon.$$

**Case 2(b):**  $M = \overline{M}$ . Consider the model  $M_P \in \mathcal{M}$ . Then, Lemma I.10 implies that

$$V^{M_P}(\pi^{M_P}) - \mathbb{E}_{\pi \sim P} V^{M_P}(\pi) \le 4\mathbb{E}_{\pi \sim P} \mathbb{E}_{x \sim \overline{\nu}} \lambda \left\| \phi(x, \pi) \right\|_{\Sigma_P^{-1}} + \varepsilon \le 4d\lambda + \varepsilon.$$

Further, because  $\widehat{\theta}_P \in \Theta_{\overline{M}}$ , we also have  $|V^{M_P}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi^{\overline{M}})| \leq 2\varepsilon$ , and hence  $V^{\overline{M}}(\pi^{\overline{M}}) \leq 2\varepsilon + V^{M_P}(\pi^{\overline{M}})$ .

Therefore, combining the inequalities above, we have

$$V^{\overline{M}}(\pi^{\overline{M}}) - \mathbb{E}_{\pi \sim P} V^{\overline{M}}(\pi) \leq V^{\overline{M}}(\pi^{\overline{M}}) - V^{M_P}(\pi^{M_P}) + V^{M_P}(\pi^{M_P}) - \mathbb{E}_{\pi \sim P} V^{M_P}(\pi) + \mathbb{E}_{\pi \sim P} [V^{\overline{M}}(\pi) - V^{M_P}(\pi)]$$
  
$$\leq 3\varepsilon + 4d\lambda + \varepsilon_P.$$

Hence, using  $\varepsilon_P \leq 5\varepsilon$ ,

$$\mathbb{E}_{\pi \sim p} \left[ V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi) \right] = \frac{1}{2} \left( V^{\overline{M}}(\pi^{\overline{M}}) - \mathbb{E}_{\pi \sim P} V^{\overline{M}}(\pi) \right) \le (10d+5)\varepsilon.$$

Combining the case (a) and (b), we conclude that

$$\operatorname{r-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M}\cup\{\overline{M}\},\overline{M})\leq (20d+6)\varepsilon.$$

**Case 3:**  $\Theta_{\overline{M}} \neq \emptyset$  and  $\varepsilon_P > 5\varepsilon$ . In this case, we set  $b = \frac{5\varepsilon}{2\varepsilon_P} < \frac{1}{2}$ , and  $p = (1-b)\overline{p} + bp^{\star}$ .

We first show that  $\mathcal{M}_{p,\varepsilon^2}(\overline{M}) = \emptyset$ . Otherwise, there exists  $M \in \mathcal{M}_{p,\varepsilon^2}(\overline{M})$ , and hence  $M \in \mathcal{M}_{p^\star,\varepsilon^2/b}(\overline{M})$  and  $M \in \mathcal{M}_{\overline{p},2\varepsilon^2}(\overline{M})$ . The latter implies  $\theta^M \in \Theta_{\overline{M}}$  (by Lemma I.11), and the former implies (by Lemma I.12)

$$\mathbb{E}_{\pi \sim P} \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \left( \langle \theta^{M}, \phi(x, \pi) \rangle - f^{\overline{M}}(x, \pi(x)) \right) \right\|^{2} \leq \frac{10\varepsilon^{2}}{b},$$

and hence by the definition Eq. (102),

$$\mathbb{E}_{\pi \sim P} \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \Big( \langle \widehat{\theta}_{P}, \phi(x, \pi) \rangle - f^{\overline{M}}(x, \pi(x)) \Big) \right\|^{2} \leq \frac{10\varepsilon^{2}}{b} + \lambda_{0}.$$

Notice that the first coordinate of  $n_{\pi}(x)$  is always 1, and hence

$$\mathbb{E}_{\pi \sim P} \left| V^{M_P}(\pi) - V^{\overline{M}}(\pi) \right|^2 \le \frac{10\varepsilon^2}{b} + \lambda_0^2, \quad \Rightarrow \quad \varepsilon_P^2 \le \frac{10\varepsilon^2}{b} + \lambda_0^2.$$

By our choice of b, this is a contradiction.

Therefore, it remains to bound  $\mathbb{E}_{\pi \sim \bar{p}} \left[ V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi) \right]$ . Notice that

$$\mathbb{E}_{\pi \sim p} \left[ V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi) \right] = b \cdot \left( V^{\overline{M}}(\pi^{\overline{M}}) - \mathbb{E}_{\pi \sim P} V^{\overline{M}}(\pi) \right),$$

and our argument in Case 2(b) also applies here:

$$V^{\overline{M}}(\pi^{\overline{M}}) - \mathbb{E}_{\pi \sim P} V^{\overline{M}}(\pi) \le 3\varepsilon + 4d\lambda + \varepsilon_P.$$

Therefore, we also have

$$\mathbb{E}_{\pi \sim p} \left[ V^{\overline{M}}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi) \right] \le (10d+4)\varepsilon.$$

The proof is completed by combining the three cases above.

# I.6.1 Proof of Lemma I.10

We denote  $P_x := Q_P(\cdot|x) \in \Delta(\mathcal{A})$  for each  $x \in \mathcal{X}$ . Then, using the definition of P = F(P), we have

$$\mathbb{E}_{\pi \sim P} V^M(\pi) = \mathbb{E}_{x \sim \nu_M} \mathbb{E}_{a \sim P_x} f^M(x, a).$$

Next, for a fixed  $x \in \mathcal{X}$ , by the definition of  $Q_P$ , it holds

$$\forall a' \in \mathcal{A}, \qquad \widehat{f}_P(x, a') \le \mathbb{E}_{a \sim P_x} \widehat{f}_P(x, a) + \lambda_0 \log |\mathcal{A}|$$

Notice that  $\left\|\theta^{M} - \widehat{\theta}_{P}\right\|_{\Sigma_{P}} \leq \lambda$ , and hence

$$\left| \langle \widehat{\theta}_P, \phi(x, a) \rangle - \langle \theta^M, \phi(x, a) \rangle \right| \le \lambda \, \|\phi(x, a)\|_{\Sigma_P^{-1}} \,,$$

which implies (using  $f^M, \hat{f}_P \in [-1, 1]$ )

$$f^{M}(x,a) \leq \widehat{f}_{P}(x,a) \leq f^{M}(x,a) + 4\min\{\lambda \|\phi(x,\pi)\|_{\Sigma_{P}^{-1}},1\}$$

Therefore, we can now combine the inequalities above to obtain

$$f^{M}(x,\pi^{M}(x)) = \max_{a\in\mathcal{A}} f^{M}(x,a) \leq \max_{a\in\mathcal{A}} \widehat{f}_{P}(x,a) \leq \mathbb{E}_{a\sim P_{x}} \widehat{f}_{P}(x,a) + \varepsilon$$
$$\leq \mathbb{E}_{a\sim P_{x}} \Big[ f^{M}(x,a) + 4\min\{\lambda \|\phi(x,\pi)\|_{\Sigma_{P}^{-1}}, 1\} \Big] + \varepsilon,$$

where we use  $\lambda_0 \log |\mathcal{A}| \leq \varepsilon$ . Taking expectation over  $x \sim \nu_M$  completes the proof of (1). Now we proceed to prove (2). For any fixed  $\pi \in \Pi$ , by Cauchy inequality,

$$\mathbb{E}_{x \sim \overline{\nu}} \|\phi(x, \pi)\|_{\Sigma_{P}^{-1}} = \mathbb{E}_{x \sim \overline{\nu}} \sqrt{\|U_{\pi} \phi(x, \pi)\|} \frac{\|\phi(x, \pi)\|_{\Sigma_{P}^{-1}}^{2}}{\|U_{\pi} \phi(x, \pi)\|} \\ \leq \sqrt{\mathbb{E}_{x \sim \overline{\nu}} \|U_{\pi} \phi(x, \pi)\|} \cdot \mathbb{E}_{x \sim \overline{\nu}} \frac{\|\phi(x, \pi)\|_{\Sigma_{P}^{-1}}^{2}}{\|U_{\pi} \phi(x, \pi)\|}.$$

Notice that by the definition of  $U_{\pi}$  (Lemma I.9), it holds that  $\mathbb{E}_{x \sim \overline{\nu}} \| U_{\pi} \phi(x, \pi) \| \leq d$ , and

$$\mathbb{E}_{x \sim \overline{\nu}} \frac{\phi(x, \pi) \phi(x, \pi)^{\top}}{\|U_{\pi} \phi(x, \pi)\|} \preceq U_{\pi}^{-2}.$$

Hence,

$$\mathbb{E}_{x\sim\overline{\nu}}\frac{\|\phi(x,\pi)\|_{\Sigma_{P}^{-1}}^{2}}{\|U_{\pi}\phi(x,\pi)\|} = \mathbb{E}_{x\sim\overline{\nu}}\left\langle\Sigma_{P}^{-1}, \frac{\phi(x,\pi)\phi(x,\pi)^{\top}}{\|U_{\pi}\phi(x,\pi)\|}\right\rangle = \left\langle\Sigma_{P}^{-1}, \mathbb{E}_{x\sim\overline{\nu}}\frac{\phi(x,\pi)\phi(x,\pi)^{\top}}{\|U_{\pi}\phi(x,\pi)\|}\right\rangle \leq \left\langle\Sigma_{P}^{-1}, U_{\pi}^{-2}\right\rangle,$$

where we recall for matrix  $A, B \in \mathbb{R}^{d \times d}$ , the Frobenius inner product is defined as  $\langle A, B \rangle = tr(A^{\top}B)$ . Therefore,

$$\mathbb{E}_{\pi \sim P} \mathbb{E}_{x \sim \overline{\nu}} \| \phi(x, \pi) \|_{\Sigma_{P}^{-1}} \leq \mathbb{E}_{\pi \sim P} \sqrt{d \cdot \langle \Sigma_{P}^{-1}, U_{\pi}^{-2} \rangle}$$
$$\leq \sqrt{d \cdot \mathbb{E}_{\pi \sim P} \langle \Sigma_{P}^{-1}, U_{\pi}^{-2} \rangle}$$
$$= \sqrt{d \langle \Sigma_{P}^{-1}, \mathbb{E}_{\pi \sim P} U_{\pi}^{-2} \rangle} = d,$$

where the last line follows from  $\mathbb{E}_{\pi \sim P} U_{\pi}^{-2} = \Sigma_P$  and  $\operatorname{tr}(I_d) = d$ .

# I.6.2 Proof of Lemma I.11

Suppose  $M \in \mathcal{M}_{\bar{p}, 2\varepsilon^2}(\overline{M})$ . Then by the definition of  $\bar{p}$ , we have

$$\left|\mathbb{E}_{r\sim M(\pi^{\overline{M}})}r - \mathbb{E}_{r\sim\overline{M}(\pi^{\overline{M}})}r\right|^{2} + \left\|\mathbb{E}_{x\sim\nu_{M}}\phi(x,\pi^{\overline{M}}) - \mathbb{E}_{x\sim\nu_{M}}\phi(x,\pi^{\overline{M}})\right\|^{2} \le 4\varepsilon^{2}.$$

Notice that  $V^{M}(\pi^{\overline{M}}) = \mathbb{E}_{r \sim M(\pi^{\overline{M}})}r$ , and hence  $|V^{M}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi^{\overline{M}})| \leq 2\varepsilon$  follows immediately. Further, we also have

$$V^{M}(\pi^{\overline{M}}) = \mathbb{E}_{r \sim M(\pi^{\overline{M}})} r = \left\langle \mathbb{E}_{x \sim \nu_{M}} \phi(x, \pi^{\overline{M}}), \theta^{M} \right\rangle,$$

and hence

$$\begin{aligned} & \left| \mathbb{E}_{x \sim \overline{\nu}} \langle \theta, \phi(x, \pi^{\overline{M}}) \rangle - V^{\overline{M}}(\pi^{\overline{M}}) \right| \\ & \leq \left| V^{M}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi^{\overline{M}}) \right| + \left| \left\langle \mathbb{E}_{x \sim \nu_{M}} \phi(x, \pi^{\overline{M}}), \theta^{M} \right\rangle - \left\langle \mathbb{E}_{x \sim \overline{\nu}} \phi(x, \pi^{\overline{M}}), \theta^{M} \right\rangle \right| \\ & \leq \left| V^{M}(\pi^{\overline{M}}) - V^{\overline{M}}(\pi^{\overline{M}}) \right| + \left\| \mathbb{E}_{x \sim \nu_{M}} \phi(x, \pi^{\overline{M}}) - \mathbb{E}_{x \sim \nu_{M}} \phi(x, \pi^{\overline{M}}) \right\| \leq 4\varepsilon. \end{aligned}$$

This immediately implies  $\theta^M \in \Theta_{\overline{M}}$ .

## I.6.3 Proof of Lemma I.12

Fix any  $M \in \mathcal{M}_{p^{\star}, 2\varepsilon^2}(\overline{M})$ . Then by the definition of  $p^{\star}$ , we know

$$\mathbb{E}_{\ell \sim q_{\pi}} \mathsf{D}_{\ell}^2(M(\pi), \overline{M}(\pi)) \le 2\varepsilon^2.$$

The inequality Eq. (106) and Eq. (108) follows immediately from Eq. (105) (notice that the first coordinate of  $n_{\pi}(x)$  is 1).

The inequality Eq. (107) follows similarly from the proof of Lemma I.11 (see Appendix I.6.2):

$$\begin{aligned} & \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \left( \langle \theta^{M}, \phi(x, \pi) \rangle - f^{\overline{M}}(x, \pi(x)) \right) \right\| \\ & \leq \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \langle \theta^{M}, \phi(x, \pi) \rangle - \mathbb{E}_{x \sim \nu_{M}} \mathsf{n}_{\pi}(x) \langle \theta^{M}, \phi(x, \pi) \rangle \right\| + \left\| \mathbb{E}_{x \sim \nu_{M}} \mathsf{n}_{\pi}(x) \langle \theta^{M}, \phi(x, \pi) \rangle - \mathbb{E}_{x \sim \overline{\nu}} f^{\overline{M}}(x, \pi(x)) \right\| \\ & \leq \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \phi(x, \pi)^{\top} - \mathbb{E}_{x \sim \nu_{M}} \mathsf{n}_{\pi}(x) \phi(x, \pi)^{\top} \right\|_{F} + \left\| \mathbb{E}_{z \sim M(\pi)} r - \mathbb{E}_{z \sim \overline{M}(\pi)} r \right\|, \end{aligned}$$

where the second inequality follows from  $\|\theta^M\| \leq 1$  and the fact that  $\mathbb{E}^{M(\pi)}[r|x] = \langle \theta^M, \phi(x,\pi) \rangle$ and  $\mathbb{E}^{\overline{M}(\pi)}[r|x] = f^{\overline{M}}(x,\pi(x))$ . Therefore,

$$\left\|\mathbb{E}_{x\sim\overline{\nu}}\mathsf{n}_{\pi}(x)\big(\langle\theta^{M},\phi(x,\pi)\rangle-f^{\overline{M}}(x,\pi(x))\big)\right\|^{2}\leq 2\left\|\mathbb{E}_{z\sim M(\pi)}\mathbf{v}_{\pi}(z)-\mathbb{E}_{z\sim\overline{M}(\pi)}\mathbf{v}_{\pi}(z)\right\|^{2},$$

and Eq. (107) follows immediately.

Finally, we bound  $\|\theta^{M} - \hat{\theta}_{P}\|_{\Sigma_{P}}$  assuming  $\theta^{M} \in \Theta_{\overline{M}}$ . Using Eq. (107), we know  $L_{P}(\theta^{M}) \leq 20\varepsilon^{2} + \lambda_{0}^{2}$ , where the quadratic loss function  $L_{P}$  is defined in Eq. (102). Therefore, using  $\hat{\theta}_{P} = \arg \min_{\theta \in \Theta_{\overline{M}}} L_{P}(\theta)$ , we have

$$\frac{1}{2} \left\| \theta^{M} - \widehat{\theta}_{P} \right\|_{\nabla^{2} L_{P}}^{2} \leq L_{P}(\theta^{M}) - L_{P}(\widehat{\theta}_{P}) \leq 20\varepsilon^{2} + \lambda_{0}^{2},$$

where using the definition of  $L_P$ , we also have

$$\frac{1}{2} \left\| \theta \right\|_{\nabla^2 L_P}^2 = \mathbb{E}_{\pi \sim P} \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \phi(x, \pi)^\top \theta \right\|^2.$$

Notice that, for  $n_{\pi}$  defined as in Lemma I.9, we have

$$\mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \phi(x, \pi)^{\top} = \mathbb{E}_{x \sim \overline{\nu}} \mathbb{E}_{x \sim \overline{\nu}} \frac{U_{\pi} \phi(x, \pi) \phi(x, \pi)^{\top}}{\|U_{\pi} \phi(x, \pi)\|} = U_{\pi}^{-1} - \lambda_0 I_d.$$

Therefore, we know

$$\frac{1}{2} \left\| \theta \right\|_{\nabla^2 L_P}^2 = \mathbb{E}_{\pi \sim P} \left\| \mathbb{E}_{x \sim \overline{\nu}} \mathsf{n}_{\pi}(x) \phi(x, \pi)^\top \theta \right\|^2 \ge \mathbb{E}_{\pi \sim P} \left[ \frac{5}{6} \left\| U_{\pi}^{-1} \theta \right\|^2 - 5\lambda_0^2 \left\| \theta \right\|^2 \right],$$

where we use  $a^2 \ge \frac{5}{6}(a+b)^2 - 5b^2$  for scalar  $a, b \ge 0$ . Plugging in  $\theta = \theta^M - \hat{\theta}_P$  and re-arranging yield

$$\left\|\theta^{M} - \widehat{\theta}_{P}\right\|_{\Sigma_{P}} = \mathbb{E}_{\pi \sim P} \left\|U_{\pi}^{-1}(\theta^{M} - \widehat{\theta}_{P})\right\|^{2} \le 24\varepsilon^{2} + 20\lambda_{0}^{2} \le 25\varepsilon^{2}.$$

## I.7 Proof of Proposition 31 and Proposition 32

In this section, we prove the results of Lipschitz contextual bandits (Proposition 31 and Proposition 32). We first state the general result for any Lipschitz contextual bandits under the following conditions on the value function class  $\mathcal{F} \subseteq (\mathcal{X} \times \mathcal{A} \rightarrow [-1, 1])$ :

(1)  $\mathcal{X}$  and  $\mathcal{A}$  are both metric space, and for any  $f \in \mathcal{F}$ ,  $x \in \mathcal{X}$ ,  $a \in \mathcal{A}$ , the function  $f(\cdot, a)$  is 1-Lipschitz over  $\mathcal{X}$ , and the function  $f(x, \cdot)$  is 1-Lipschitz over  $\mathcal{A}$ .

(2) There is a convex function class  $\mathcal{F}_{\mathsf{b}} \subseteq (\mathcal{A} \to [-1, 1])$ , such that for any  $f \in \mathcal{F}, x \in \mathcal{X}$ , we have  $f(x, \cdot) \in \mathcal{F}_{\mathsf{b}}$ .

(3) The offset DEC of  $\mathcal{F}_{\mathsf{b}}$  is defined as

$$\operatorname{\mathsf{r-dec}}_{\gamma}^{\mathrm{o}}(\mathcal{F}_{\mathsf{b}}, \bar{f}) = \inf_{p \in \Delta(\mathcal{A})} \sup_{f \in \mathcal{F}_{\mathsf{b}}} \mathbb{E}_{a \sim p} \left[ \max_{a'} f(a') - f(a) - \gamma_{x} (f(a) - \bar{f}(a))^{2} \right],$$
(109)

and we define  $\mathsf{r-dec}^{o}_{\gamma}(\mathcal{F}_{\mathsf{b}}) = \sup_{\bar{f} \in \mathcal{F}_{\mathsf{b}}} \mathsf{r-dec}^{o}_{\gamma}(\mathcal{F}_{\mathsf{b}}, \bar{f})$  following Foster et al. [2021]. We assume that there is an increasing function  $d(\gamma) \geq 1$  such that

$$\mathsf{r} ext{-dec}^{\mathrm{o}}_\gamma(\mathcal{M}_\mathsf{b}) \leq rac{d(\gamma)}{\gamma}, \quad orall \gamma > 0.$$

Under the above conditions, we prove that the offset DEC of  $\mathcal{M} = \mathcal{M}_{\mathcal{F},CB}$  can be bounded, and  $ExO^+$  can be suitably instantiated to achieve the desired regret.

**Specifications of LDP-ExO.** To instantiate LDP-ExO, we consider an information set structure that is slightly different from the one considered in Appendix F.5.4.

Fix a parameter  $\Delta \geq 0$ , we denote  $N_X := N(\mathcal{X}, \Delta), N_A := N(\mathcal{A}, \Delta)$ . Let  $\mathcal{X}_\Delta \subset \mathcal{X}$  be a minimal  $\Delta$ -covering of  $\mathcal{X}$ , and for each  $x \in \mathcal{X}$ , we let  $[x] \in \mathcal{X}_\Delta$  such that  $\rho(x, [x]) \leq \Delta$ . Similarly, we take a minimal  $\Delta$ -covering  $\mathcal{A}_\Delta$  of  $\mathcal{A}$ , and for each  $a \in \mathcal{A}$ , we define  $[a] \in \mathcal{A}_\Delta$  such that  $\rho(a, [a]) \leq \Delta$ . We consider the space  $\Pi_+ := \prod_{x \in \mathcal{X}_\Delta} \mathcal{A}_\Delta$ , i.e., for each  $\bar{\pi} \in \Pi_+, \bar{\pi}$  is a  $\mathcal{A}_\Delta$ -valued vector indexed by  $x \in \mathcal{X}_\Delta$ . We also identify  $\Pi_+ \subseteq \Pi$ , by associating  $\bar{\pi}(x) = \bar{\pi}([x])$  for all  $x \in \mathcal{X}$ .

For  $\bar{\pi} \in \Pi_+$ , we let  $\mathcal{F}_{\bar{\pi}}$  be class of all reward functions f such that  $\bar{\pi}$  is a near-optimal policy:

$$\mathcal{F}_{\bar{\pi}} := \left\{ f \in \mathcal{F} : \forall x \in \mathcal{X}_{\Delta}, f(x, \bar{\pi}(x)) \ge \max_{a \in \mathcal{A}} f(x, a) - \Delta \right\}.$$

By definition,  $\mathcal{F} = \bigcup_{\bar{\pi} \in \Pi_+} \mathcal{F}_{\bar{\pi}}$ .

Then, we consider the (Type 1) information set structure  $\Psi = \Pi_+$ , with

$$\mathcal{M}_{\psi} := \{ M_{\nu, f} : \nu \in \Delta(\mathcal{X}), f \in \mathcal{F}_{\psi} \}, \qquad \pi_{\psi} = \psi, \qquad \psi \in \Psi = \Pi_{+}.$$

Clearly,  $\Psi$  is a valid information set structure for the constraint set  $\mathscr{P}_{\mathsf{cxt}}$  introduced in Section 5.5, and we have  $\log |\Psi| = N_X \log N_A$ . Therefore, it remains to upper bound the regret DEC r-dec $^{0,\mathsf{LDP}}_{\gamma}(\mathcal{M}_{\Psi})$ .

Theorem I.13 (Learning Lipschitz contextual bandits). It holds that

$$\mathsf{r}\operatorname{-dec}_{\gamma}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M}_{\mathcal{F},\mathsf{CB}}) \leq \mathsf{r}\operatorname{-dec}_{\gamma}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M}_{\Psi}) \leq \Delta + \frac{N_X(10d(\gamma) + 5)}{\gamma}$$
(110)

Further, LDP-ExO (when instantiated on  $\Psi$ ) achieve with probability at least  $1 - \delta$ 

$$\frac{1}{T} \mathbf{Reg}_{\mathsf{DM}}(T) \lesssim \Delta + N_X \left( \frac{d(\gamma)}{\alpha^2 \gamma} + \frac{\gamma \log(N_A/\delta)}{T} \right)$$

The proof of Theorem I.13 is deferred to Appendix I.7.1. Using Theorem I.13, we prove Proposition 31 and Proposition 32 as follows.

**Proof of Proposition 31 (upper bound).** By Foster et al. [2021, Proposition 5.2], for any function class  $\mathcal{F}_{b} \subseteq (\mathcal{A} \rightarrow [-1, 1])$ , we have

$$\mathsf{r-dec}^{\mathrm{o}}_{\gamma}(\mathcal{F}_{\mathsf{b}}) \leq \frac{|\mathcal{A}|}{\gamma}$$

Therefore, we have

$$\mathrm{r-dec}_{\varepsilon}^{\mathrm{LDP}}(\mathcal{M}) \leq \inf_{\gamma \geq 0} \left( \mathrm{r-dec}_{\gamma}^{\mathrm{o},\mathrm{LDP}}(\mathcal{M}) + \gamma \varepsilon^2 \right) \lesssim \Delta + \sqrt{N_X |\mathcal{A}| \varepsilon}$$

Similarly, we can suitably choose the parameter  $\gamma > 0$  such that  $\mathsf{ExO}^+$  achieves

$$\operatorname{\mathbf{Reg}}_{\mathsf{DM}} \lesssim T\Delta + N_X \sqrt{|\mathcal{A}|T \log(|\mathcal{A}|/\delta)}.$$

Proof of Proposition 32. In this case, we have

 $\mathcal{F}_{\mathsf{b}} = \{ f : \mathcal{A} \to [-1, 1] : f \text{ is concave and 1-Lipschitz over } \mathcal{A} \subset \mathbb{R}^{K} \text{ under } \| \cdot \|_{2} \}.$ 

Suppose that the diameter of  $\mathcal{A}$  is bounded by  $R \geq 1$ . Then, by Foster et al. [2021, Proposition 6.3], we have

$$\mathsf{r-dec}^{\mathrm{o}}_{\gamma}(\mathcal{F}_{\mathsf{b}}) \leq \frac{K^4}{\gamma} \cdot \operatorname{poly} \log(\gamma, R).$$

Therefore, we can bound

$$\operatorname{r-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M}) \leq \inf_{\gamma > 0} \left(\operatorname{r-dec}_{\gamma}^{\operatorname{o},\operatorname{LDP}}(\mathcal{M}) + \gamma \varepsilon^{2}\right) \lesssim \Delta + \sqrt{N_{X} K^{4} \varepsilon} \cdot \operatorname{poly} \log(\varepsilon^{-1}, R).$$

Also note that  $\log N_A \leq K \log(R/\Delta)$ . Therefore, we can suitably choose the parameter  $\gamma > 0$  such that  $\mathsf{ExO}^+$  achieves with probability at least  $1 - \delta$ 

$$\operatorname{\mathbf{Reg}}_{\mathsf{DM}} \lesssim T\Delta + N_X \sqrt{K^5 T} \cdot \operatorname{poly} \log(T, R, \delta^{-1}, \Delta^{-1}).$$

### I.7.1 Proof of Theorem I.13

Let  $\mathsf{ExO}^+$  be instantiated with the measurement class  $\Phi = \mathcal{Q}_{\alpha}$  and information set structure  $\Psi$ . Then, by Theorem F.4 and our analysis in Appendix E.3.2, it holds that with probability at least  $1 - \delta$ 

$$\frac{1}{T}\mathbf{Reg}_{\mathsf{DM}}(T) \leq 3\Delta + \mathrm{r-dec}_{c\alpha^{2}\gamma}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M}_{\Psi}) + \frac{2\gamma}{T}[\log N_{\mathsf{frac}}(\mathscr{P}_{\mathsf{cxt}},\Psi,3\Delta) + \log(1/\delta)].$$

Therefore, it remains to upper bound the fractional covering number  $\log N_{\text{frac}}(\mathscr{P}_{\text{cxt}}, \Psi, 3\Delta)$  and the private regret-DEC.

We first prove that  $\log N_{\text{frac}}(\mathscr{P}_{\text{cxt}}, \Psi, 3\Delta) \leq \log |\Psi|$ . Consider the prior  $w = \text{Unif}(\Psi)$ . Note that for any  $f \in \mathcal{F}$ , there exists  $\bar{\pi} \in \Pi_+$  such that  $f \in \mathcal{F}_{\bar{\pi}}$ . Then, for any  $x \in \mathcal{X}$ ,

$$\max_{a \in \mathcal{A}} f(x, a) \le \max_{a \in \mathcal{A}} f([x], a) + \Delta$$

r	-	

$$\leq f([x], \bar{\pi}([x])) + 2\Delta \leq f(x, \bar{\pi}(x)) + 3\Delta,$$

where we use the Lipschitzness of f and  $\bar{\pi}(x) = \bar{\pi}([x])$ . Therefore, we know  $\mathcal{P}_f \subseteq \mathcal{M}_{\bar{\pi}}$  and for any  $M \in \mathcal{P}_f$ 

$$V^{M}(\pi^{M}) - V^{M}(\bar{\pi}) = \mathbb{E}_{x \sim M} \left[ \max_{a \in \mathcal{A}} f(x, a) - f(x, \bar{\pi}(x)) \right] \le 3\Delta.$$

This implies that  $N_{\mathsf{frac}}(\mathscr{P}_{\mathsf{cxt}}, \Psi, 3\Delta) \leq |\Psi|$ .

Therefore, it remains to prove Eq. (110).

**Proof of Eq. (110).** To upper bound  $\mathsf{r-dec}_{\gamma}^{o,\mathsf{LDP}}(\mathcal{M}_{\Psi})$ , we fix a reference model  $\overline{M} \in \mathrm{co}(\mathcal{M}_{\Psi}) = \mathrm{co}(\mathcal{M}_{\mathcal{F},\mathsf{CB}}) =: \mathcal{M}^+$ .

For any model  $M \in \mathcal{M}^+$ , we consider the function

$$\tilde{f}^{M}(x_{i},a) = \mathbb{E}^{M}[r|[x] = x_{i},a].$$

Then, because  $M \in \mathcal{M}^+$ , we know  $\tilde{f}^M(x_i, \cdot)$  is a convex combination of elements in  $\mathcal{F}_{\mathsf{b}}$ , and hence  $\tilde{f}^M(x_i, \cdot) \in \mathcal{F}_{\mathsf{b}}$  for all  $x_i \in \mathcal{X}_{\Delta}$ . We also denote  $\nu_M \in \Delta(\mathcal{X}_{\Delta})$  to be distribution of  $[x], x \sim M$ .

Construction of the distribution  $p \in \Delta(\Pi \times \mathcal{L})$ : For each  $x \in \mathcal{X}_{\Delta}$ , we consider  $\bar{f}_x := \tilde{f}^{\overline{M}}(x, \cdot) \in \mathcal{F}_{\mathsf{b}}$ ,  $\gamma_x = \frac{\gamma \nu_{\overline{M}}(x)}{5}$ , and

$$p_x = \underset{p \in \Delta(\mathcal{A})}{\operatorname{arg min}} \sup_{f \in \mathcal{F}_{\mathsf{b}}} \mathbb{E}_{a \sim p} \left[ \underset{a'}{\max} f(a') - f(a) - \gamma_x (f(a) - \bar{f}_x(a))^2 \right]$$

Then we know

$$\mathbb{E}_{a \sim p_x} \left[ \max_{a'} f(a') - f(a) - \gamma_x (f(a) - \bar{f}_x(a))^2 \right] \le \frac{d(\gamma_x)}{\gamma_x}, \quad \forall f \in \mathcal{F}_{\mathsf{b}}$$

Next, we consider the following maps  $\mathbf{v}_0, \mathbf{v} : \mathcal{Z} \to \mathbb{R}^{N_X}$ :

$$\mathbf{v}_0(x,a,r) = \mathbf{e}_{[x]}, \qquad \mathbf{v}(x,a,r) = r\mathbf{e}_{[x]},$$

where  $\mathbf{e}_{x_i} \in \mathbb{R}^{N_X}$  is the vector with the *i*-th coordinate being 1 and other coordinates being 0. Then, by Lemma I.6, there exists a distribution  $Q \in \Delta(\mathcal{L})$  such that for all  $M \in \mathcal{M}^+$ ,

$$2\mathbb{E}_{\ell \sim Q} \mathsf{D}_{\ell}(M(\pi), M(\pi))^{2}$$

$$\geq \left\| \mathbb{E}_{z \sim M(\pi)}[\mathbf{v}_{0}(z)] - \mathbb{E}_{z \sim \overline{M}(\pi)}[\mathbf{v}_{0}(z)] \right\|^{2} + \left\| \mathbb{E}_{z \sim M(\pi)}[\mathbf{v}(z)] - \mathbb{E}_{z \sim \overline{M}(\pi)}[\mathbf{v}(z)] \right\|^{2}.$$

Then, we let  $p \in \Delta(\Pi \times \mathcal{L})$  be the distribution of  $(\pi, \ell)$  under  $\ell \sim Q$ ,  $\pi([x]) \sim p_{[x]}$  independently for all  $[x] \in \mathcal{X}_{\Delta}$ , and  $\pi(x) = \pi([x])$  for all  $x \in \mathcal{X}$ .

Then, by definition

$$2\mathbb{E}_{(\pi,\ell)\sim p} \mathsf{D}_{\ell}(M(\pi), \overline{M}(\pi))^{2}$$

$$= 2\mathbb{E}_{\pi\sim p} \mathbb{E}_{\ell\sim Q} \mathsf{D}_{\ell}(M(\pi), \overline{M}(\pi))^{2}$$

$$\geq \mathbb{E}_{\pi\sim p} \left[ \sum_{x\in\mathcal{X}_{\Delta}} |\nu_{M}(x) - \nu_{\overline{M}}(x)|^{2} + |\nu_{M}(x)\tilde{f}^{M}(x, \pi(x)) - \nu_{\overline{M}}(x)\tilde{f}^{\overline{M}}(x, \pi(x))|^{2} \right]$$

$$\geq \mathbb{E}_{\pi \sim p} \left[ \frac{1}{2} \sum_{x \in \mathcal{X}_{\Delta}} \nu_{\overline{M}}(x)^2 \left| \tilde{f}^M(x, \pi(x)) - \tilde{f}^{\overline{M}}(x, \pi(x)) \right|^2 \right]$$
$$= \frac{1}{2} \sum_{x \in \mathcal{X}_{\Delta}} \nu_{\overline{M}}(x)^2 \mathbb{E}_{a \sim p_x} \left| \tilde{f}^M(x, a) - \tilde{f}^{\overline{M}}(x, a) \right|^2,$$

where the last line follows from the definition of  $\pi \sim p$ , as  $\pi([x]) \sim p_{[x]}$  independently.

Next, for any  $M \in \mathcal{M}_{\Psi}$ , there exists  $\bar{\pi} \in \Pi_+$  such that  $M \in co(\mathcal{M}_{\bar{\pi}})$ , and by our argument above, we know  $V^M(\pi^M) - V^M(\bar{\pi}) \leq 3\Delta$ , and thus

$$\begin{aligned} & \mathbb{E}_{\pi \sim p} [V^{M}(\pi^{M}) - V^{M}(\pi)] \\ &\leq 3\Delta + \mathbb{E}_{\pi \sim p} [V^{M}(\bar{\pi}) - V^{M}(\pi)] \\ &= 3\Delta + \mathbb{E}_{\pi \sim p} \mathbb{E}_{x \sim \nu_{M}} \left[ \tilde{f}^{M}(x, \bar{\pi}(x)) - \tilde{f}^{M}(x, \pi(x)) \right] \\ &\leq 3\Delta + \mathbb{E}_{x \sim \nu_{M}} \mathbb{E}_{a \sim p_{[x]}} \left[ \max_{a' \in \mathcal{A}} \tilde{f}^{M}(x, a') - \tilde{f}^{M}(x, a) \right] \\ &\leq 3\Delta + \mathbb{E}_{x \sim \nu_{\overline{M}}} \mathbb{E}_{a \sim p_{[x]}} \left[ \max_{a' \in \mathcal{A}} \tilde{f}^{M}(x, a') - \tilde{f}^{M}(x, a) \right] + 2 \sum_{x \in \mathcal{X}_{\Delta}} |\nu_{M}(x) - \nu_{\overline{M}}(x)| \,. \end{aligned}$$

Combining the inequalities above with the AM-GM inequality  $2a \leq \frac{a^2}{5} + \frac{5}{\gamma}$ , we know

$$\begin{split} & \mathbb{E}_{\pi \sim p}[V^{M}(\pi^{M}) - V^{M}(\pi)] - \gamma \mathbb{E}_{(\pi,\ell) \sim p} \mathsf{D}_{\ell}(M(\pi), \overline{M}(\pi))^{2} \\ & \leq 3\Delta + \sum_{x \in \mathcal{X}_{\Delta}} \left( \nu_{\overline{M}}(x) \mathbb{E}_{a \sim p_{[x]}} \left[ \max_{a' \in \mathcal{A}} \tilde{f}^{M}(x, a') - \tilde{f}^{M}(x, a) \right] + 2 \left| \nu_{M}(x) - \nu_{\overline{M}}(x) \right| \right) \\ & - \gamma \sum_{x \in \mathcal{X}_{\Delta}} \left( \frac{1}{5} \nu_{\overline{M}}(x)^{2} \mathbb{E}_{a \sim p_{x}} \left| \tilde{f}^{M}(x, a) - \tilde{f}^{\overline{M}}(x, a) \right|^{2} + \frac{1}{10} \left| \nu_{M}(x) - \nu_{\overline{M}}(x) \right|^{2} \right) \\ & \leq 3\Delta + \frac{10}{\gamma} + \sum_{x \in \mathcal{X}_{\Delta}} \nu_{\overline{M}}(x) \mathbb{E}_{a \sim p_{[x]}} \left[ \max_{a' \in \mathcal{A}} \tilde{f}^{M}(x, a') - \tilde{f}^{M}(x, a) - \frac{\nu_{\overline{M}}(x)\gamma}{5} \left| \tilde{f}^{M}(x, a) - \tilde{f}^{\overline{M}}(x, a) \right|^{2} \right] \\ & \leq 3\Delta + \frac{10}{\gamma} + \sum_{x \in \mathcal{X}_{\Delta}} \nu_{\overline{M}}(x) \cdot \mathbf{r} \cdot \mathsf{dec}_{\gamma_{x}}^{\mathsf{O},\mathsf{LDP}}(\mathcal{F}_{\mathsf{b}}, \bar{f}_{x}) \leq 3\Delta + \frac{10}{\gamma} + N_{X} \cdot \frac{5d(\gamma)}{\gamma}, \end{split}$$

where the last line follows from the choice of  $p_x$ . This gives the desired upper bound on the offset DEC as

$$\mathsf{r-dec}_{\gamma}^{\mathrm{o},\mathsf{LDP}}(\mathcal{M},\overline{M}) \leq \sup_{M \in \mathcal{M}_{\Psi}} \mathbb{E}_{(\pi,\ell) \sim p} \left[ L(M,\pi) - \gamma \mathsf{D}_{\ell}(M(\pi),\overline{M}(\pi))^2 \right] \leq \frac{N_X(5d(\gamma) + 10)}{\gamma}.$$

Therefore, by the arbitrariness of  $\overline{M}$ , the proof of Eq. (110) is completed.

# I.8 Lower bounds for structured contextual bandits

The argument of Appendix I.5 also implies the following lower bound for contextual bandits.

**Proposition I.14.** Let  $d \ge 1, \Delta \in (0, 1]$ . Consider the contextual bandits problem with context space  $\mathcal{X} = [d]$ , action space  $\mathcal{A} = \{0, 1\}$ , reward function class

$$\mathcal{F}_d := \{ f_\theta : \forall i \in [d], f_\theta(i,0) = 0, f_\theta(i,1) = \theta_i \Delta \}_{\theta \in \Theta},$$

where  $\Theta = \{-1, 1\}^d$ . Let  $\mathcal{M}_d$  be the contextual bandits problem class with reward function in  $\mathcal{F}_d$ and context distribution  $\mu = \text{Unif}(\mathcal{X})$ . Then, for any T-round  $\alpha$ -LDP algorithm,

$$\sup_{M \in \mathcal{M}_d} \mathbb{E}^{M, \mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \geq \frac{\Delta}{4}, \qquad unless \ T \gtrsim \frac{d^2}{\alpha^2 \Delta^2}.$$

Note that when  $\Delta \leq \frac{1}{\sqrt{d}}$ ,  $\mathcal{F}_d$  is a class of linear functions, and hence Proposition I.14 immediately implies a regret lower bound for linear contextual bandits.

**Corollary I.15.** Let  $d \ge 1, T \ge 1$ . Then for any *T*-round  $\alpha$ -LDP algorithm,

$$\sup_{M \in \mathcal{M}_{\mathsf{Lin-CB}}} \mathbb{E}^{M,\mathsf{Alg}}[\mathbf{Reg}_{\mathsf{DM}}(T)] \ge c \min\left\{\frac{d\sqrt{T}}{\alpha}, \frac{T}{\sqrt{d}}\right\}.$$

Similarly, we can prove the lower bound of Proposition 31 as follows.

**Proof of Proposition 31 (lower bound).** Fix a  $\Delta > 0$ , and we set  $d = N(\mathcal{X}, 2\Delta)$ ,  $\mathcal{A} = \{0, 1\}$ . By the duality of packing and covering, there exists  $x_1, \dots, x_d \in \mathcal{X}$  such that  $\rho(x_i, x_j) \ge 2\Delta, \forall i \neq j$ .

Then, for each  $\theta \in \{-1,1\}^d$ , we define  $f_{\theta} \in \mathcal{F}_{\mathsf{Lip}}$  as follows: for any  $x \in \mathcal{X}$ , we set  $f_{\theta}(x,0) = 0$  and

$$f_{\theta}(x,1) := \sum_{i=1}^{d} \theta_i \max\{\Delta - \rho(x,x_i), 0\}.$$

By definition,  $f_{\theta}(\cdot, 1)$  is clearly 1-Lipschitz, because for any  $x \in \mathcal{X}$ , there is at most one  $i \in [d]$  such that  $\rho(x, x_i) \leq \Delta$ .

Therefore, we have an inclusion  $\iota : \mathcal{M}_d \to \mathcal{M}_{\mathsf{Lip-CB}}$ . Hence, Proposition I.14 implies that for any *T*-round  $\alpha$ -LDP algorithm Alg, we have

$$\sup_{M \in \mathcal{M}_{\mathsf{Lip-CB}}} \mathbb{E}^{M,\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)] \geq \frac{\Delta}{4}, \qquad \text{unless } T \gtrsim \frac{d^2}{\alpha^2 \Delta^2}.$$

This is the desired result.

## I.9 Proof of Proposition B.5

Fix  $\Delta > 0$  and let  $\varepsilon_0 = \operatorname{cor}(\mathcal{M}, \Delta)$ .

Then, there exists a reference model  $\overline{M}$  and a set of models  $\{M_1, \dots, M_m\} \subseteq \mathcal{M}$ , such that (1)  $\{M_1, \dots, M_m\}$  is  $\varepsilon$ -correlated relative to  $\overline{M}$ ; (2) for any  $\pi \in \Pi$ , there is at most m/2 indexes  $i \in [m]$  such that  $L(M_i, \pi) \leq \Delta$ .

In the following, we proceed to lower bound the quantile-based private PAC-DEC (cf. Appendix I.2). For any  $\ell \in \mathcal{L}$ , we have

$$\sum_{i=1}^{m} \mathsf{D}_{\ell}(M_i, \overline{M})^2 = \sum_{i=1}^{m} \left| M_i[\ell] - \overline{M}[\ell] \right|^2 = \sup_{w \in \mathbb{R}^m : ||w|| \le 1} \left| \sum_{i=1}^{m} w_i \left( M_i[\ell] - \overline{M}[\ell] \right) \right|^2.$$

 _	
	L
	н

For any fixed  $w \in \mathbb{R}^m$ , we can consider the shifted  $\overline{\ell}(z) = \ell(z) - \frac{1}{2} \in [-\frac{1}{2}, \frac{1}{2}]$  and bound

$$\begin{aligned} \left|\sum_{i=1}^{m} w_i \left(M_i[\ell] - \overline{M}[\ell]\right)\right|^2 &= \left|\sum_{i=1}^{m} w_i \left(M_i[\bar{\ell}] - \overline{M}[\bar{\ell}]\right)\right|^2 \\ &= \left|\mathbb{E}_{z \sim \overline{M}} \left[ \left(\sum_{i=1}^{m} w_i \left(\frac{M_i[z]}{\overline{M}[z]} - 1\right)\right) \bar{\ell}(z) \right] \right|^2 \\ &\leq \mathbb{E}_{z \sim \overline{M}} \left(\sum_{i=1}^{m} w_i \left(\frac{M_i[z]}{\overline{M}[z]} - 1\right)\right)^2 \cdot \mathbb{E}_{z \sim \overline{M}} \bar{\ell}(z)^2 \\ &\leq \frac{1}{4} \sum_{i,j} \rho_{\overline{M}}(M_i, M_j) w_i w_j, \end{aligned}$$

where the first inequality follows from the Cauchy inequality. Therefore,

$$\sum_{i=1}^{m} \mathsf{D}_{\ell}(M_i, \overline{M})^2 = \sup_{w \in \mathbb{R}^m : \|w\| \le 1} \left| \sum_{i=1}^{m} w_i \left( M_i[\ell] - \overline{M}[\ell] \right) \right|^2$$
$$\leq \frac{1}{4} \sup_{w \in \mathbb{R}^m : \|w\| \le 1} \sum_{i,j} |\rho_{\overline{M}}(M_i, M_j)| \cdot |w_i w_j|$$
$$\leq \frac{1}{4} \left( m \varepsilon_0^2 + \sqrt{m(m-1)} \varepsilon_0^2 \right) \le \frac{m \varepsilon_0^2}{2},$$

where the last inequality follows from the definition of  $\varepsilon$ -correlation.

Hence, we may consider  $\mu = \text{Unif}(\{M_1, \dots, M_m\}) \subseteq \Delta(\mathcal{M})$ . Then, for any  $p \in \Delta(\Pi), q \in \Delta(\mathcal{L})$ , we know that

$$\mathbb{P}_{M \sim \mu, \pi \sim p}(L(M, \pi) \ge \Delta) \ge \inf_{\pi} \mathbb{P}_{M \sim \mu}(L(M, \pi) \ge \Delta) \ge \frac{1}{2}.$$

Therefore, there must exist  $\mathcal{M}_0 \subset \mathcal{M}$  such that  $\mu(\mathcal{M}_0) \geq \frac{1}{3}$ , and

$$\mathbb{P}_{\pi \sim p}(L(M,\pi) \ge \Delta) \ge \frac{1}{4}, \quad \forall M \in \mathcal{M}_0.$$

Then, we also know

$$\mu(\mathcal{M}_0) \min_{M \in \mathcal{M}_0} \mathbb{E}_{\ell \sim q} \mathsf{D}_{\ell}(M, \overline{M})^2 \le \mathbb{E}_{M \sim \mu} \mathbb{E}_{\ell \sim q} \mathsf{D}_{\ell}(M, \overline{M})^2 \le \frac{1}{2} \varepsilon_0^2$$

and hence there exists  $M \in \mathcal{M}_0$  with  $\mathbb{E}_{\ell \sim q} \mathsf{D}_{\ell}(M, \overline{M})^2 \leq \frac{3}{2} \varepsilon_0^2$ . This gives

$$\mathsf{p}\text{-}\mathsf{dec}_{\sqrt{2}\varepsilon_0,1/4}^{\mathsf{q},\mathsf{LDP}}(\mathcal{M},\overline{M}) \geq \Delta,$$

which also implies  $\mathsf{p-dec}_{\sqrt{2}\varepsilon_0}^{\mathsf{LDP}}(\mathcal{M},\overline{\mathcal{M}}) \geq \frac{\Delta}{4}$ . Hence, the desired lower bounds on  $\mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M})$  follows for all  $\varepsilon \geq \varepsilon_0$ .

Furthermore, applying Proposition I.1 with  $p-dec_{\sqrt{2}\varepsilon_0,1/4}^{q,\text{LDP}}(\mathcal{M},\overline{M}) \geq \Delta$ , we also have the desired lower bound on sample complexity.

## I.10 Proof of Proposition B.6

Fix a parameter  $\lambda \in [0, \frac{1}{2}]$  and denote  $\mathcal{X}_+ = \mathcal{X} \setminus \{\mathbf{0}\}, \ \mu_+ = \text{Unif}(\mathcal{X}_+).$ 

Then, for each  $S \subseteq [n]$ , we consider  $M_S := M_{S,\lambda}$  the model with covariate distribution being  $\mu_{\lambda} = (1 - \lambda)\delta_0 + \lambda\mu_+$ , and  $y = f_S(x)$  for  $(x, y) \sim M_S$ . We then consider the subclass  $\mathcal{M}_{\lambda} = \{M_{S,\lambda}\}_{S \subseteq [n]} \subset \mathcal{M}$ , and let  $\overline{M} = \overline{M}_{\lambda} \in \Delta(\mathcal{Z})$  be the reference model given by  $x \sim \mu$ , and  $y \sim \text{Rad}(0)$  if  $x \neq \mathbf{0}$ , and y = 0 otherwise.

By definition, the pairwise correlation is given by

$$\rho_{\overline{M}}(M_S, M_{S'}) = \lambda \mathbb{E}_{x \sim \mu_+} f_S(x) f_{S'}(x)$$

Further, we know  $f_S(x) = (-1)^{\sum_{i \in S} x_i}$ , and hence

$$\mathbb{E}_{x \sim \operatorname{Unif}(\mathbb{Z}_2^d)} f_S(x) f_{S'}(x) = \mathbb{E}_{x \sim \operatorname{Unif}(\mathbb{Z}_2^d)} (-1)^{\sum_{i \in S \Delta S'} x_i} = \begin{cases} 0, & S \neq S', \\ 1, & S = S'. \end{cases}$$

Therefore, we have

$$\rho_{\overline{M}}(M_S, M_{S'}) = \begin{cases} -\frac{\lambda}{2^d - 1}, & S \neq S', \\ \lambda, & S = S'. \end{cases}$$

We also know that for any  $f \in (\mathcal{X} \to \{-1, 1\}),$ 

$$L(M_S, f) \ge \lambda \mathbb{P}_{x \sim \mu_+}(f(x) \neq f_S(x)) = \lambda \mathbb{E}_{x \sim \mu_+} \left[ \frac{1 - f(x) f_S(x)}{2} \right].$$

Therefore, for any  $S \neq S'$ ,

$$L(M_S, f) + L(M_{S'}, f) \ge \mathbb{P}_{x \sim \mu}(f_S(x) \neq f_{S'}(x)) \ge \frac{\lambda}{2} \left(1 - \mathbb{E}_{x \sim \mu_+} f_S(x) f_{S'}(x)\right) = \frac{\lambda}{2} \left(1 - \frac{1}{2^d - 1}\right).$$

Hence, for any  $f \in (\mathcal{X} \to \{-1, 1\})$ , there exists at most one model  $M_S$  such that  $L(M_S, f) \leq \frac{\lambda}{8}$ , and by definition of the minimum correlation (Definition 15), we know

$$\operatorname{cor}(\mathcal{M}_{\lambda},\lambda/8) \le \frac{\lambda}{2^d-1}.$$
 (111)

Note that for  $\lambda = \Theta(1)$ , Eq. (111) is enough for proving lower bound  $\Omega(2^d)$  for constant suboptimality: applying Proposition B.5 immediately yields the desired result (for sub-optimality level  $\varepsilon = \Theta(1)$ ).

In the following, we use a slightly more careful argument to show the lower bound of private PAC-DEC. Notice that Eq. (111) implies that for  $\lambda = \frac{1}{2}$  and  $\varepsilon_0 = \frac{1}{\sqrt{2^d-1}}$ ,

$$\operatorname{p-dec}_{\varepsilon_0}^{\operatorname{LDP}}(\mathcal{M}_{1/2},\overline{M}_{1/2}) \geq \frac{1}{16}.$$

Further, notice that for  $\lambda \leq \frac{1}{2}$ ,  $S \subseteq [n]$ , we have

$$L(M_{S,\lambda},f) = 2\lambda L(M_{S,1/2},f), \qquad \mathsf{D}_{\ell}(M_{S,\lambda},\overline{M}_{\lambda}) = 2\lambda \mathsf{D}_{\ell}(M_{S,1/2},\overline{M}_{1/2}).$$

Therefore, by the definition of private PAC-DEC (7),

$$\mathsf{p}\text{-}\mathsf{dec}_{2\lambda\varepsilon_0}^{\mathsf{LDP}}(\mathcal{M}_{\lambda},\overline{M}_{\lambda}) = 2\lambda\cdot\mathsf{p}\text{-}\mathsf{dec}_{\varepsilon_0}^{\mathsf{LDP}}(\mathcal{M}_{1/2},\overline{M}_{1/2}) \geq \frac{\lambda}{8},$$

and hence for any  $\varepsilon \leq \varepsilon_0$ , we con set  $\lambda(\varepsilon) = \frac{\varepsilon}{2\varepsilon_0}$ , and then

$$\mathsf{p}\text{-}\mathsf{dec}^{^{\mathrm{LDP}}}_{\varepsilon}(\mathcal{M}) \geq \mathsf{p}\text{-}\mathsf{dec}^{^{\mathrm{LDP}}}_{\varepsilon}(\mathcal{M}_{\lambda(\varepsilon)}, \overline{M}_{\lambda(\varepsilon)}) \geq \frac{\lambda(\varepsilon)}{8} = \frac{1}{16}\sqrt{2^d - 1}\varepsilon.$$

Applying Theorem 19 completes the proof, as L is a metric-based loss.

## I.11 Proof of Proposition B.2

Fix the *T*-round algorithm Alg with rules  $\{q_t\} \cup \{p\}$ , we define  $\mathsf{Alg}_{\mathsf{pure}}$  as follows: for each round  $t \in [T]$ ,

- Sample  $(\pi_t, \widetilde{\mathsf{Q}}_t) \sim q_t(\cdot | \pi_1, \widetilde{\mathsf{Q}}_1, z_1, \cdots, \pi_{t-1}, \widetilde{\mathsf{Q}}_{t-1}, z_{t-1}).$
- Set  $\pi_t = (\pi_t, \widetilde{\mathsf{Q}}_{\mathsf{pure},t})$  and observe  $z_t$ .

Now, we define  $\widetilde{\mathbb{P}}^{M,\text{pr}}$  to be the joint distribution of  $\mathcal{H} = \{(\pi_t, \widetilde{\mathsf{Q}}_t, \widetilde{\mathsf{Q}}_{\mathsf{pure},t}, z_t)\}_{t \in [T]}$  under  $\mathsf{Alg}_{\mathsf{pure}}$  and model  $M \in (\Pi \to \Delta(\mathcal{Z}))$ .

As an intermediate step of proof, we also consider  $\widetilde{\mathbb{P}}^{M}$ , the distribution of  $\mathcal{H} = \{(\pi_t, \mathsf{Q}_t, \mathsf{Q}_{\mathsf{pure},t}, z_t)\}_{t \in [T]}$ under Alg and model  $M \in \mathcal{M}$ . By data-processing inequality, we have

$$D_{\mathrm{TV}}\left(\mathbb{P}^{M,\mathrm{Alg}}(\mathcal{H}_{\pi}=\cdot),\mathbb{P}^{M,\mathrm{Alg}_{\mathsf{pure}}}(\mathcal{H}_{\pi}=\cdot)\right) \leq D_{\mathrm{TV}}\left(\widetilde{\mathbb{P}}^{M,\mathrm{pr}},\widetilde{\mathbb{P}}^{M}\right).$$

Then, we may apply the chain rule of TV distance, which gives

$$\begin{split} D_{\mathrm{TV}}\left(\widetilde{\mathbb{P}}^{M,\mathrm{pr}},\widetilde{\mathbb{P}}^{M}\right) &\leq \sum_{t=1}^{T} \mathbb{E}_{\widetilde{\mathbb{P}}^{M}} D_{\mathrm{TV}}\left(\widetilde{\mathbb{P}}^{M,\mathrm{pr}}(z_{t}=\cdot|\pi_{1:t},\mathsf{Q}_{1:t},\mathsf{Q}_{\mathsf{pure},1:t},z_{1:t-1}),\widetilde{\mathbb{P}}^{M}(z_{t}=\cdot|\pi_{1:t},\mathsf{Q}_{1:t},\mathsf{Q}_{\mathsf{pure},1:t},z_{1:t-1})\right) \\ &= \sum_{t=1}^{T} \mathbb{E}_{\widetilde{\mathbb{P}}^{M}} D_{\mathrm{TV}}\left(\widetilde{\mathbb{P}}^{M,\mathrm{pr}}(z_{t}=\cdot|\pi_{t},\mathsf{Q}_{\mathsf{pure},t}),\widetilde{\mathbb{P}}^{M}(z_{t}=\cdot|\pi_{t},\mathsf{Q}_{t})\right) \\ &= \sum_{t=1}^{T} \mathbb{E}_{\widetilde{\mathbb{P}}^{M}} D_{\mathrm{TV}}\left(\mathsf{Q}_{\mathsf{pure},t}\circ M(\pi_{t}),\mathsf{Q}_{t}\circ M(\pi_{t})\right) \\ &\leq T \cdot \sup_{\mathsf{Q}} \sup_{z\in\mathcal{Z}} D_{\mathrm{TV}}\left(\mathsf{Q}_{\mathsf{pure}}(\cdot|z),\mathsf{Q}(\cdot|z)\right) \\ &\leq \frac{T\beta}{1+e^{\alpha}-\beta}, \end{split}$$

where the expectation  $\mathbb{E}_{\widetilde{\mathbb{P}}^M}$  is taken over the trajectory  $\mathcal{H} = \{(\pi_t, \mathsf{Q}_t, \mathsf{Q}_{\mathsf{pure},t}, z_t)\}_{t \in [T]} \sim \widetilde{\mathbb{P}}^M$ . Combining the inequalities above completes the proof.

# J Proofs from Section 6 and Appendix C

# J.1 Proof of Theorem 33

For simplicity, for any model class  $\mathcal{M} \subseteq \Delta(\mathcal{Z})$ , we denote

$$\mathfrak{M}_{T}(\mathcal{M}) := \inf_{\mathsf{Alg}} \sup_{M \in \mathcal{M}} \mathbb{E}^{M,\mathsf{Alg}}[\mathbf{Risk}_{\mathsf{DM}}(T)],$$

where the  $\inf_{Alg}$  is taken over  $\alpha$ -LDP algorithms. Then we know

$$\mathfrak{M}_T^{\mathrm{loc}}(\mathcal{M}, M_0) := \sup_{M_1 \in \mathcal{M}} \mathfrak{M}_T(\{M_1, M_0\}).$$

**Proof of the upper bound.** We only need to bound Private PAC-DEC in terms of the local DEC, as follows.

**Lemma J.1.** For any 2-point model class  $\mathcal{M}_0 = \{M_1, M_0\} \subseteq \mathcal{M}$ , it holds that

$$\mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}_0) \leq \begin{cases} \inf_{\pi \in \Pi} \left( L(M_1, \pi) + L(M_0, \pi) \right), & \text{if } \sup_{\pi} D_{\mathrm{TV}} \left( M_1(\pi), M_0(\pi) \right) \leq 2\varepsilon, \\ 0, & \text{otherwise.} \end{cases}$$

With Lemma J.1, we know that

$$\sup_{M_1 \in \mathcal{M}} \operatorname{p-dec}_{\varepsilon}^{\operatorname{LDP}}(\{M_0, M_1\}) \leq \operatorname{p-dec}_{2\varepsilon}^{\operatorname{loc}}(\mathcal{M}).$$

Applying Theorem 21 gives

$$\mathfrak{M}_T(\{M_1, M_0\}) \le \mathsf{p-dec}_{\bar{\varepsilon}_{\delta}(T)}^{\mathsf{LDP}}(\{M_0, M_1\}) + \delta.$$

Therefore, we may combine the two inequalities above to obtiin

$$\mathfrak{M}_{T}^{\mathrm{loc}}(\mathcal{M}, M_{0}) = \sup_{M_{1} \in \mathcal{M}} \mathfrak{M}_{T}(\{M_{1}, M_{0}\}) \leq \mathsf{p-dec}_{2\bar{\varepsilon}_{\delta}(T)}^{\mathrm{loc}}(\mathcal{M}) + \delta.$$

This is the desired upper bound.

**Proof of the lower bound.** Similar to the proof of upper bound, we can directly lower bound  $\sup_{M_1 \in \mathcal{M}} \mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\{M_1, M_0\})$  by the local DEC  $\mathsf{p-dec}_{\varepsilon}^{\mathsf{loc}}(\mathcal{M})$ . However, the private PAC-DEC lower bound (Theorem 19) requires certain structural assumptions on the loss function L, which is in fact artificial in this case. Therefore, in the following, we utilize the quantile DEC lower bound (Appendix I.2) to obtain a better lower bound.

**Lemma J.2.** For any  $\varepsilon > 0$ , it holds

$$\sup_{M_1 \in \mathcal{M}} \mathsf{p-dec}_{\varepsilon, 1/2}^{q, \mathsf{LDP}}(\{M_1, M_0\}, M_0) \geq \frac{1}{2} \mathsf{p-dec}_{\varepsilon}^{\mathrm{loc}}(\mathcal{M}, M_0).$$

Now, applying Proposition I.1 gives

$$\mathfrak{M}_T(\{M_1,M_0\}) \geq \frac{1}{4} \mathrm{p-dec}_{\underline{\varepsilon}(T),1/2}^{\mathrm{q,LDP}}(\{M_1,M_0\}),$$

and hence Lemma J.2 yields

$$\mathfrak{M}_{T}^{\mathrm{loc}}(\mathcal{M}, M_{0}) = \sup_{M_{1} \in \mathcal{M}} \mathfrak{M}_{T}(\{M_{1}, M_{0}\}) \geq \frac{1}{4} \sup_{M_{1} \in \mathcal{M}} \mathsf{p-dec}_{\underline{\varepsilon}(T), 1/2}^{q, \mathsf{LDP}}(\{M_{1}, M_{0}\}) \geq \frac{1}{8} \mathsf{p-dec}_{\varepsilon}^{\mathrm{loc}}(\mathcal{M}, M_{0}).$$

This is the desired result.

**Proof of Lemma J.1.** Define  $\varepsilon_0 = \sup_{\pi} D_{\text{TV}}(M_1(\pi), M_0(\pi))$  and

$$\pi^{\mathsf{exp}} = \underset{\pi \in \Pi}{\operatorname{arg\,max}} D_{\mathrm{TV}} \left( M_1(\pi), M_0(\pi) \right).$$

Further, we choose  $\ell \in \mathcal{L}$  such that

$$\mathsf{D}_{\ell}(M_1(\pi^{\mathsf{exp}}), M_0(\pi^{\mathsf{exp}})) = D_{\mathrm{TV}}\left(M_1(\pi^{\mathsf{exp}}), M_0(\pi^{\mathsf{exp}})\right).$$

Then, we consider the distribution  $q \in \Delta(\Pi \times \mathcal{L})$  supported on  $(\pi^{exp}, \ell)$ , and any reference model  $\overline{M} \in co(\mathcal{M}_0)$  given by  $\overline{M} = \lambda M_0 + (1 - \lambda)M_1$  (where  $\lambda \in [0, 1]$ ). There are two cases:

(1) If  $\varepsilon_0 \leq 2\varepsilon$ , then, we choose

$$\pi^{\mathsf{out}} = \underset{\pi \in \Pi}{\operatorname{arg\,min}} \left( L(M_1, \pi) + L(M_0, \pi) \right),$$

and let  $p \in \Delta(\Pi)$  be the distribution supported on  $\pi^{out}$ . Then, p certifies that

$$\operatorname{p-dec}_{\varepsilon}^{\operatorname{LDP}}(\mathcal{M}_0, \overline{M}) \leq \inf_{\pi \in \Pi} \left( L(M_1, \pi) + L(M_0, \pi) \right).$$

(2) If  $\varepsilon_0 > 2\varepsilon$ , then using the fact that

$$\mathsf{D}_{\ell}\big(M_0(\pi^{\mathsf{exp}}), \overline{M}(\pi^{\mathsf{exp}})\big) = (1-\lambda)\mathsf{D}_{\ell}(M_1(\pi^{\mathsf{exp}}), M_0(\pi^{\mathsf{exp}})) = (1-\lambda)\varepsilon_0, \\ \mathsf{D}_{\ell}\big(M_1(\pi^{\mathsf{exp}}), \overline{M}(\pi^{\mathsf{exp}})\big) = \lambda\mathsf{D}_{\ell}(M_1(\pi^{\mathsf{exp}}), M_0(\pi^{\mathsf{exp}})) = \lambda\varepsilon_0,$$

we know there is at most one index  $i \in \{0,1\}$  such that  $\mathsf{D}_{\ell}(M_i(\pi^{\mathsf{exp}}), \overline{M}(\pi^{\mathsf{exp}})) \leq \varepsilon$ . If such an index does not exist, then we already have  $\mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}_0, \overline{M}) = 0$ . Otherwise, given such an index i, we can take a decision  $\pi^{\mathsf{out}}$  such that  $L(M_i, \pi^{\mathsf{out}}) = 0$ , which also certifies  $\mathsf{p-dec}_{\varepsilon}^{\mathsf{LDP}}(\mathcal{M}_0, \overline{M}) = 0$ .  $\Box$ 

**Proof of Lemma J.2.** We take  $0 \leq \Delta < p-dec_{\varepsilon}^{loc}(\mathcal{M}, M_0)$ . Then by definition, there exists  $M_1 \in \mathcal{M}$  such that

$$\sup_{\pi} D_{\mathrm{TV}}\left(M_1(\pi), M_0(\pi)\right) \le \varepsilon, \qquad \inf_{\pi} \left(L(M_1, \pi) + L(M_0, \pi)\right) \ge \Delta.$$

Then, for any  $p \in \Delta(\Pi)$  and  $q \in \Delta(\Pi \times \mathcal{L})$ , we have

$$\mathbb{E}_{(\pi,\ell)\sim q} \mathsf{D}^2_{\ell}(M_1(\pi), M_0(\pi)) \leq \sup_{\pi} D_{\mathrm{TV}}\left(M_1(\pi), M_0(\pi)\right)^2 \leq \varepsilon^2,$$

and we also know  $\{\pi: L(M_1, \pi) < \Delta/2\}$  and  $\{\pi: L(M_0, \pi) < \Delta/2\}$  are disjoint, which implies

$$p(\pi : L(M_1, \pi) \ge \Delta/2) + p(\pi : L(M_0, \pi) \ge \Delta/2) \ge 1.$$

Therefore, the quantile-based private PAC-DEC can be lower bounded as

$$\mathrm{p\text{-}dec}_{\varepsilon,1/2}^{\mathrm{q},\mathrm{LDP}}(\{M_1,M_0\},M_0)\geq \frac{\Delta}{2}$$

This gives the desired result by letting  $\Delta \to \mathsf{p-dec}^{\mathrm{loc}}_{\varepsilon}(\mathcal{M}, M_0)$ .

# J.2 Proof of Theorem 34

We first recall the notations and results of Appendix I.2.1. Using Eq. (96), we know that for any models  $M, \overline{M}$ ,

$$D_{\mathrm{KL}}(\mathbb{P}^{M,\mathrm{Alg}} \parallel \mathbb{P}^{\overline{M},\mathrm{Alg}}) \le (e^{\alpha} - 1)^2 T \cdot \mathbb{E}_{(\pi,\ell) \sim \tilde{q}_{M,\mathrm{Alg}}} \mathsf{D}_{\ell}^2(M(\pi),\overline{M}(\pi)) \le (e^{\alpha} - 1)^2 T.$$

On the other hand, by the definition Definition 5 of  $N_{\mathsf{frac}}(\mathcal{M}, \Delta)$ , we have that

$$\frac{1}{N_{\mathsf{frac}}(\mathcal{M},\Delta)} := \sup_{p \in \Delta(\Pi)} \inf_{M \in \mathcal{M}} p(\pi : L(M,\pi) \le \Delta).$$

Therefore, we may fix a reference model  $\overline{M} \in \mathcal{M}$ , and it holds that

$$\inf_{M \in \mathcal{M}} p_{\overline{M}, \mathsf{Alg}}(\pi : L(M, \pi) \le \Delta) \le \frac{1}{N_{\mathsf{frac}}(\mathcal{M}, \Delta)},$$

and hence there exists  $M \in \mathcal{M}$  such that

$$p_{\overline{M},\mathsf{Alg}}(\pi: L(M,\pi) \le \Delta) \le \frac{1}{N_{\mathsf{frac}}(\mathcal{M},\Delta)}.$$

On the other hand, the condition of Theorem 34 gives  $p_{M,Alg}(\pi : L(M, \pi) \leq \Delta) \geq \frac{1}{2}$ . Therefore, by data-processing inequality,

$$D_{\mathrm{KL}}(\mathbb{P}^{M,\mathrm{Alg}} \parallel \mathbb{P}^{\overline{M},\mathrm{Alg}}) \ge D_{\mathrm{KL}}(p_{M,\mathrm{Alg}} \parallel p_{\overline{M},\mathrm{Alg}}) \ge \frac{\log N_{\mathrm{frac}}(\mathcal{M},\Delta)}{2} - 1.$$

Comparing the lower and upper bounds above complete the proof.

# J.3 Proof of Proposition C.1 and Proposition 37

With the following lemma (which generalizes Beimel et al. [2013a]), the proof is essentially similar to Appendix J.2.

**Lemma J.3.** Suppose that Alg is a T-round  $\alpha$ -JDP algorithm. Then for any two models  $M, \overline{M}$ , it holds that

$$\mathbb{P}^{M,\mathsf{Alg}}(\pi_{T+1} \in E) \le e^{T\alpha} \cdot \mathbb{P}^{\overline{M},\mathsf{Alg}}(\pi_{T+1} \in E), \qquad \forall E.$$
(112)

Notice that by definition, for any model  $M \in \mathcal{M}$ ,

$$\mathbb{P}^{M,\operatorname{Alg}}(L(M,\pi_{T+1}) \le \Delta) \ge \frac{1}{2},$$

and hence

$$\mathbb{P}^{\overline{M},\mathsf{Alg}}(L(M,\pi_{T+1}) \leq \Delta) \geq e^{-T\alpha} \mathbb{P}^{M,\mathsf{Alg}}(L(M,\pi_{T+1}) \leq \Delta) \geq \frac{1}{2}e^{-T\alpha}, \qquad \forall M \in \mathcal{M}.$$

Then, by the definition of fractional covering number (Definition 5), we know for the distribution  $p = \mathbb{P}^{\overline{M}, \operatorname{Alg}}(\pi_{T+1} = \cdot) \in \Delta(\Pi)$ , it holds that

$$2e^{T\alpha} \ge \sup_{M \in \mathcal{M}} \frac{1}{p(L(M,\pi) \le \Delta)} \ge N_{\mathsf{frac}}(\mathcal{M},\Delta).$$

This gives the desired lower bound  $\alpha T \geq \log N_{\mathsf{frac}}(\mathcal{M}, \Delta) - \log 2$ .

**Proof of Lemma J.3.** We first consider the setting of statistical problems, which is easier to analyze. In this case, by definition, for any sequence of observations  $\mathcal{H}_{z,T} = (z_1, \dots, z_T)$ ,  $\mathcal{H}'_{z,T} = (z'_1, \dots, z'_T) \in \mathcal{Z}^T$ ,  $\alpha$ -JDP implies that

$$\mathbb{P}^{\mathsf{Alg}}(\pi_{T+1} \in E|\mathcal{H}_T) \le e^{T\alpha} \cdot \mathbb{P}^{\mathsf{Alg}}(\pi_{T+1} \in E|\mathcal{H}_T'), \qquad \forall E.$$
(113)

Therefore, we may take expectation over  $\mathcal{H}_{z,T} = (z_1, \cdots, z_T) \sim M$  and  $\mathcal{H}'_{z,T} = (z'_1, \cdots, z'_T) \sim \overline{M}$ , which completes the proof of Lemma J.3.

More generally, for interactive learning, for any two sequences  $\mathcal{H}_{z,T} = (z_1, \dots, z_T)$  and  $\mathcal{H}'_{z,T} = (z'_1, \dots, z'_T)$ , it holds that

$$\mathbb{P}^{\mathsf{Alg}}((\pi_1,\cdots,\pi_T,\pi_{T+1})=\cdot|\mathcal{H}_{z,T})\leq e^{T\alpha}\mathbb{P}^{\mathsf{Alg}}\big((\pi_1,\cdots,\pi_T,\pi_{T+1})=\cdot|\mathcal{H}'_{z,T}\big).$$

Therefore, for any fixed sequence  $(\pi_1, \dots, \pi_T, \pi_{T+1})$ , we may take expectation over  $z_t \sim M(\pi_t), z'_t \sim \overline{M}(\pi_t)$  independently (recursively for  $t = T, T - 1, \dots, 1$ ), which gives

$$\mathbb{P}^{M,\mathsf{Alg}}(\pi_1,\cdots,\pi_T,\pi_{T+1}) \leq e^{T\alpha}\mathbb{P}^{\overline{M},\mathsf{Alg}}(\pi_1,\cdots,\pi_T,\pi_{T+1}).$$

Hence, the proof of Lemma J.3 is completed.

## J.4 Proof of Proposition 35

We consider the private analog of the algorithm of Chen et al. [2024]. For the simplicity of presentation, we focus on PAC learning.

# Algorithm 5 "Brute-Force" Algorithm

**Input:** Model class  $\mathcal{M}$ , decision space  $\Pi$ , parameter  $\Delta, \delta > 0, T \ge 1$ . 1: Set  $N = N_{\mathsf{frac}}(\mathcal{M}, \Delta) \log(1/\delta), J = \frac{T}{N}$ , and

$$p_{\Delta}^{\star} = \arg\min_{p \in \Delta(\Pi)} \sup_{M \in \mathcal{M}} \frac{1}{p(\pi : L(M, \pi) \le \Delta)}.$$
(114)

2: for  $k = 0, 1, \dots, N-1$  do 3: Sample  $\pi_{(k)} \sim p_{\Delta}^{\star}$ . 4: Set  $\ell_{(k)} = R(\cdot, \pi_{(k)})$  and  $Q_{(k)} = Q_{\ell_{(k)}}$  be the binary channel (Example 5). 5: for  $t = kJ + 1, \dots, (k+1)J$  do 6: Select  $\pi_t = \pi_{(k)}$  and  $Q_t = Q_{(k)}$  and observes  $o_t \sim Q_t \circ M^{\star}(\pi_t)$ . 7: Compute  $\widehat{r}_{(k)} = \frac{1}{J} \sum_{t=kJ+1}^{(k+1)J} o_t$ . 8: Set  $\widehat{k} = \arg \max_{k \in [N]} \widehat{r}_{(k)}$ . Output:  $\pi_{T+1} = \pi_{(\widehat{k})}$ .

Analysis of Algorithm 5. By definition,

$$\pi_{(1)}, \cdots, \pi_{(N)} \sim p_{\Delta}^{\star}$$
 independently.

Hence,

$$\mathbb{P}\big(\forall k \in [N], L(M^{\star}, \pi_{(k)}) > \Delta\big) = p_{\Delta}^{\star}(\pi : L(M^{\star}, \pi) > \Delta)^{N}$$

$$f^{M} = \operatorname*{arg\,min}_{f \in \mathcal{F}} \mathbb{E}_{(x,y) \sim M} \mathcal{L}(y, f(x)),$$

and then for any  $f \in \mathcal{F}^+$ ,

$$L(M,f) = \mathbb{E}_{(x,y)\sim M}[\mathcal{L}(y,f(x)) - \mathcal{L}(y,f^{M}(x))] \le \mathbb{E}_{x\sim M} \left| f(x) - f^{M}(x) \right|.$$

Therefore, we have  $N_{\mathsf{frac}}(\mathcal{M}_{\mathsf{agnostic}}, \Delta) \leq N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$  for any  $\Delta > 0$ .

We next consider the absolute loss  $L_{abs}$ . By definition, for any  $M \in \mathcal{M}_{\mathcal{F}, realizable}$ , we have L(M, f) = $\mathbb{E}_{x \sim M} |f(x) - f^M(x)|$ , and hence it holds that  $N_{\mathsf{frac}}(\mathcal{M}_{\mathcal{F},\mathsf{realizable}}, \Delta) = N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$ . Notice that  $\mathcal{M}_{\mathcal{F},\mathsf{realizable}} \subseteq \mathcal{M}_{\mathsf{agnostic}}$ , and hence we have

$$N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta) = N_{\mathsf{frac}}(\mathcal{M}_{\mathcal{F}, \mathsf{realizable}}, \Delta) \le N_{\mathsf{frac}}(\mathcal{M}_{\mathsf{agnostic}}, \Delta) \le N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta).$$

This gives the desired results.

**Remark J.4.** Similarly, under the squared loss  $L_{sq}$ , we can also show that

$$N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \sqrt{2\Delta}) \le N_{\mathsf{frac}}(\mathcal{M}_{\mathcal{F}, \mathsf{realizable}}, \Delta) = N_{\mathsf{frac}}(\mathcal{M}_{\mathcal{F}}, \Delta) \le N_{\mathsf{frac}}(\mathcal{M}_{\mathsf{agnostic}}, \Delta) \le N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta).$$

$$\leq \left(1 - \frac{1}{N_{\mathsf{frac}}(\mathcal{M}, \Delta)}\right)^{N}$$
$$\leq \exp\left(-\frac{N}{N_{\mathsf{frac}}(\mathcal{M}, \Delta)}\right) \leq \delta$$

Therefore, with probability at least  $1 - \delta$ , there exists  $k \in [K]$  such that  $L(M^*, \pi_{(k)}) \leq \Delta$ .

Furthermore, by the definition of  $(\pi_{(k)}, \mathsf{Q}_{(k)})$ , we know that for  $t \in [kJ+1, (k+1)J]$ , the observation  $o_t \sim \operatorname{Rad}(c_{\alpha} V^{M^{\star}}(\pi_{(k)}))$  are generated independently. Therefore, with probability at least  $1 - \frac{\delta}{N}$ , it holds that

$$\left|\widehat{r}_{(k)} - c_{\alpha} V^{M^{\star}}(\pi_{(k)})\right| \leq \sqrt{\frac{2\log(2N/\delta)}{J}} =: \varepsilon_J.$$

Hence, taking the union bound, we know that with probability at least  $1 - 2\delta$ ,

$$c_{\alpha}V^{M^{\star}}(\pi_{(\widehat{k})}) \ge c_{\alpha} \max_{k \in [N]} V^{M^{\star}}(\pi_{(\widehat{k})}) - 2\varepsilon_{J} \ge c_{\alpha} \left( V^{M^{\star}}(\pi^{M^{\star}}) - \Delta \right) - 2\varepsilon_{J}.$$

Reorganizing yields

$$\begin{split} L(M^{\star}, \pi_{T+1}) &= V^{M^{\star}}(\pi^{M^{\star}}) - V^{M^{\star}}(\pi_{(\widehat{k})}) \leq \Delta + \frac{2}{c_{\alpha}} \sqrt{\frac{2N \log(T/\delta)}{T}} \\ &\leq \Delta + O\left(\frac{\log(T/\delta)}{\alpha} \sqrt{\frac{N_{\mathsf{frac}}(\mathcal{M}, \Delta)}{T}}\right). \end{split}$$

This is the desired result.

#### **J.5** Proof of Lemma C.2 and Lemma C.3

We first show that  $N_{\text{frac}}(\mathcal{M}_{\text{agnostic}}, \Delta) \leq N_{\text{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$ 1-Lipschitz loss L. For any  $M \in$  $\mathcal{M}_{agnostic}$ , we denote

$$f^{M} = \operatorname*{arg\,min}_{f \in \mathcal{F}} \mathbb{E}_{(x,y) \sim M} \mathcal{L}(y, f(x)),$$

$$M, f) = \mathbb{E}_{(x,y)\sim M}[\mathcal{L}(y, f(x)) - \mathcal{L}(y, f^{M}(x))] \le \mathbb{E}_{x\sim M}|f(x) - f^{M}(x)|$$

# J.6 Proof of Proposition C.4

**Proof of the lower bound.** For any parameter  $C \ge 1$ , we define

$$\mathcal{F}_{\mathsf{Lin},C} := \{ f_{\theta}(x) = \langle \theta, x \rangle \}_{\theta : \|\theta\| \le C}$$

We lower bound  $N_{\mathsf{frac}}(\mathcal{F}_{\mathsf{Lin}}, \mathcal{F}_{\mathsf{Lin},C}, \Delta)$  as follows. Denote  $\mathcal{F} := \mathcal{F}_{\mathsf{Lin}}$  and  $\mathcal{F}_C := \mathcal{F}_{\mathsf{Lin},C}$ . Fix any  $p \in \Delta(\mathcal{F}_C)$ , and we bound

$$\inf_{\mu \in \Delta(\mathcal{X}), f^{\star} \in \mathcal{F}} \mathbb{P}_{f \sim p} \left( f : \mathbb{E}_{x \sim \mu} | f(x) - f^{\star}(x) | \leq \frac{1}{2} \right) \leq \mathbb{E}_{x_0 \sim \mathrm{Unif}(\mathbb{S}^{d-1})} \mathbb{P}_{f \sim p} \left( f : | f(x_0) - 1 | \leq \frac{1}{2} \right) \\
= \mathbb{E}_{f \sim p} \mathbb{P}_{x_0 \sim \mathrm{Unif}(\mathbb{S}^{d-1})} \left( f : | f(x_0) - 1 | \leq \frac{1}{2} \right).$$

Notice that for any fix  $\theta \in \mathbb{R}^d$  and  $x_0 \sim \text{Unif}(\mathbb{S}^{d-1})$ , we have  $\langle \theta, x_0 \rangle = \|\theta\| t$ , where the random variable  $t \in [-1, 1]$  has density function

$$P(t) = \frac{\Gamma(d/2)}{\Gamma((d-1)/2)\sqrt{\pi}} (1-t^2)^{(d-3)/2},$$

see e.g. Bubeck et al. [2016, Section 2]. Therefore,

$$\mathbb{P}_{x_0 \sim \text{Unif}(\mathbb{S}^{d-1})} \left( f : |f(x_0) - 1| \le \frac{1}{2} \right) = \mathbb{P}_{t \sim P} \left( t \in \left[ \frac{1}{2 \|\theta\|}, \frac{3}{2 \|\theta\|} \right] \right)$$
$$\le \frac{1}{\|\theta\|} \cdot O\left(\sqrt{d}\right) \left( 1 - \frac{4}{\|\theta\|^2} \right)^{(d-3)/2}$$
$$\le O(1) \frac{\sqrt{d}}{C} \exp\left( -\frac{d-3}{2C^2} \right),$$

Therefore, as long as  $C \leq c_0 \sqrt{d}$ , we have

$$\inf_{\mu \in \Delta(\mathcal{X}), f^{\star} \in \mathcal{F}} \mathbb{P}_{f \sim p} \left( f : \mathbb{E}_{x \sim \mu} |f(x) - f^{\star}(x)| \le \frac{1}{2} \right) \le \exp\left(-c_1 \frac{d}{C^2}\right), \qquad \forall p \in \Delta(\mathcal{F}_C),$$

for some universal constants  $c_1, c_0 > 0$ . Therefore,

$$\log N_{\mathsf{frac}}(\mathcal{F}_{\mathsf{Lin}}, \mathcal{F}_{\mathsf{Lin}, C}, \Delta) \ge c_1 \frac{d}{C^2}, \qquad \forall C \in [1, c_0 \sqrt{d}].$$

In particular, this gives the desired lower bound by letting C = 1.

**Proof of the upper bound.** As the above lemma indicates, to upper bound  $N_{\text{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta)$ , we must choose  $p \in \Delta(\Pi)$  to be highly improper. We construct such a distribution of improper functions as follows.

Fix a parameter  $\lambda \in [0, \lambda_0]$  for some small enough universal constant  $\lambda_0$ . We set p to be the distribution of  $f_{\theta}$  with  $\theta \sim N(0, \lambda^2 I_d)$ . We proceed to lower bound the probability

$$\mathbb{P}_{f \sim p}(f : \mathbb{E}_{x \sim \mu} | f(x) - f^{\star}(x) | \le \Delta)$$

for arbitrary fixed  $f^{\star} = f_{\theta^{\star}} \in \mathcal{F}$  and distribution  $\mu \in \Delta(\mathcal{X})$ . Notice that for  $\theta \in \mathbb{R}^d$ , we have

$$\left(\mathbb{E}_{x\sim\mu}|f_{\theta}(x) - f^{\star}(x)|\right)^{2} \leq \mathbb{E}_{x\sim\mu}|f_{\theta}(x) - f^{\star}(x)|^{2} = \|\theta - \theta^{\star}\|_{\Sigma}^{2},$$

where  $\Sigma = \mathbb{E}_{x \sim \mu}[xx^{\top}]$ . By the rotational invariance, we may assume that  $\Sigma = \text{diag}(\lambda_1, \dots, \lambda_d)$ with  $\lambda_1 \geq \dots \geq \lambda_d \geq 0$ . Notice that  $\text{tr}(\Sigma) \leq 1$ , and hence we have  $\sum_{i=1}^n \lambda_i \leq 1$  and  $\lambda_k \leq \frac{1}{k}$ . Therefore, we know

$$\|\theta - \theta^{\star}\|_{\Sigma}^{2} \leq \max_{1 \leq i \leq k} (\theta_{i} - \theta_{i}^{\star})^{2} + \sum_{i=k+1}^{d} \lambda_{i} (\theta_{i} - \theta_{i}^{\star})^{2}.$$

Using the fact that  $\theta_i \sim N(0, \lambda^2)$ , we know

$$\mathbb{P}_{\theta}\left(\left(\theta_{i}-\theta_{i}^{\star}\right)^{2}\leq\lambda^{2}\right)\geq\frac{\lambda}{\sqrt{2\pi}}\exp\left(-\frac{\left(\left|\theta_{i}^{\star}\right|+\lambda\right)^{2}}{2\lambda^{2}}\right).$$

Therefore, using the independence between  $\theta_1, \cdots, \theta_k$ , we have

$$\mathbb{P}_{\theta}\Big(\forall i \in [k], (\theta_i - \theta_i^{\star})^2 \le \lambda^2\Big) \ge \left(\frac{\lambda}{\sqrt{2\pi}}\right)^k \exp\left(-\sum_{i=1}^k \frac{\left(|\theta_i^{\star}| + \lambda\right)^2}{2\lambda^2}\right) \ge \left(\frac{\lambda}{\sqrt{2\pi}e}\right)^k \exp\left(-\frac{1}{\lambda^2}\right).$$

Further, using the fact that

$$\mathbb{E}_{\theta}\left[\sum_{i=k+1}^{d} \lambda_i (\theta_i - \theta_i^{\star})^2\right] = \sum_{i=k+1}^{d} \lambda_i \left(|\theta_i^{\star}|^2 + \lambda^2\right) \le \frac{1}{k} + \lambda^2,$$

we know that

$$\mathbb{P}_{\theta}\left(\sum_{i=k+1}^{d} \lambda_i (\theta_i - \theta_i^{\star})^2 \le \frac{2}{k} + 2\lambda^2\right) \ge \frac{1}{2}.$$

Therefore, using the independence between  $\theta_1, \cdots, \theta_d$ , we have

$$\mathbb{P}_{\theta}\bigg(\|\theta - \theta^{\star}\|_{\Sigma}^{2} \le \frac{2}{k} + 3\lambda^{2}\bigg) \ge \frac{1}{2}\exp\bigg(-\frac{1}{\lambda^{2}} + k\log(\sqrt{2\pi}e/\lambda)\bigg).$$

Setting  $k = \frac{\Delta^2}{4}$  and  $\lambda^2 = \frac{\Delta^2}{6}$  gives

$$\mathbb{P}_{\theta}\Big(\|\theta - \theta^{\star}\|_{\Sigma}^{2} \leq \Delta^{2}\Big) \geq \exp\left(-\frac{C_{0}}{\Delta^{2}}\log\left(\frac{1}{\Delta}\right)\right),$$

where  $C_0$  is a large universal constant. By the arbitrariness of  $\mu$  and  $\theta^{\star}$ , we have

$$-\log \mathbb{P}_{f\sim p}(f:\mathbb{E}_{x\sim \mu}|f(x)-f^{\star}(x)| \leq \Delta) \leq \frac{C_0}{\Delta^2}\log\left(\frac{1}{\Delta}\right), \qquad \forall \mu \in \Delta(\mathcal{X}), \theta^{\star} \in \mathbb{B}^d(1).$$

Therefore, p certifies that  $\log N_{\mathsf{frac}}(\mathcal{F}, \mathcal{F}^+, \Delta) \leq \frac{C_0}{\Delta^2} \log \left(\frac{1}{\Delta}\right)$ , and the proof is hence completed.  $\Box$ 

# J.7 Proof of Proposition 39

Suppose that  $p \in \Delta(\mathcal{F})$  is given by

$$p := \arg\min_{p \in \Delta(\mathcal{F})} \sup_{\nu \in \Delta(\mathcal{X}), f^{\star} \in \mathcal{F}} \frac{1}{p(f : \mathbb{E}_{x \sim \nu} |f(x) - f^{\star}(x)| \leq \Delta)}.$$

Then, for any given  $\nu \in \Delta(\mathcal{X}), f^{\star} \in \mathcal{F}$ , we have

$$\mathbb{P}_{f \sim p}(\mathbb{P}_{x \sim \nu}(f(x) \neq f^{\star}(x)) \leq \Delta) \geq \frac{1}{N_{\mathsf{frac}}(\mathcal{F}, \Delta)}.$$

Therefore, for  $N \ge 1$ , we consider the distribution  $p_N$  over the subsets of  $\Pi$  given by

 $\mathcal{H} \sim p_N : \mathcal{H} = \{f_1, \cdots, f_N\}, \qquad f_1, \cdots, f_N \sim p \text{ independently.}$ 

Then, we can bound

$$\mathbb{P}_{\mathcal{H}\sim p_N}(\exists f \in \mathcal{H}, \mathbb{P}_{x\sim\nu}(f(x) \neq f^{\star}(x)) \leq \Delta) \geq 1 - \left(1 - \frac{1}{N_{\mathsf{frac}}(\mathcal{F}, \Delta)}\right)^N.$$

Choosing  $N \ge N_{\mathsf{frac}}(\mathcal{F}, \Delta) \log(4)$  yields that  $p_N$  is a  $\Delta$ -probabilistic representation of  $\mathcal{F}$ , and hence

$$\mathsf{RDim}_{\Delta}(\mathcal{F}) \leq \log N \leq \log N_{\mathsf{frac}}(\mathcal{F}, \Delta) + 2.$$

Conversely, suppose that  $\mathscr{H}$  is an optimal  $\varepsilon$ -probabilistic representation of  $\mathcal{F}$ , i.e.  $\mathsf{RDim}_{\varepsilon}(\mathcal{F}) = \operatorname{size}(\mathscr{H})$ . Then  $\mathscr{H}$  induces a distribution  $p_{\mathscr{H}} \in \Delta(\Pi)$  as

$$f \sim p_{\mathscr{H}} : \mathscr{H} \sim \mathscr{H}, \ f \sim \text{Unif}(\mathscr{H}).$$

Then, for any  $\nu \in \Delta(\mathcal{X}), f^{\star} \in \mathcal{F},$ 

$$\mathbb{P}_{f \sim p_{\mathscr{H}}}(f : \mathbb{E}_{x \sim \nu} | f(x) - f^{\star}(x) | \leq \varepsilon) \geq \mathbb{E}_{\mathcal{H} \sim \mathscr{H}} \left[ \frac{1}{|\mathcal{H}|} \mathbf{1} \{ \exists f \in \mathcal{H}, \mathbb{P}_{x \sim \nu}(f(x) \neq f^{\star}(x)) \leq \varepsilon \} \right]$$
$$\geq \frac{3}{4} \frac{1}{\sup_{\mathcal{H} \in \mathrm{supp}(\mathscr{H})} |\mathcal{H}|}.$$

Therefore,  $p_{\mathscr{H}}$  certifies that

$$\log N_{\mathsf{frac}}(\mathcal{F},\varepsilon) \leq \sup_{\mathcal{H} \in \mathrm{supp}(\mathscr{H})} \log |\mathcal{H}| + \log(4/3) \leq \mathsf{RDim}_{\varepsilon}(\mathcal{F}) + 1.$$

Combining the inequalities above completes the proof.