

Improved global determination of two-meson distribution amplitudes from multi-body B decays

Da-Cheng Yan¹, Hsiang-nan Li², Zhou Rui³, Zhen-Jun Xiao⁴, and Ya Li^{5*}

¹ *School of Mathematics and Physics, Changzhou University, Changzhou, Jiangsu 213164, China*

² *Institute of Physics, Academia Sinica, Taipei, Taiwan 115, Republic of China*

³ *Department of Physics, Yantai University, Yantai 264005, China*

⁴ *Department of Physics and Institute of Theoretical Physics, Nanjing Normal University, Nanjing 210023, Jiangsu, China and*

⁵ *Department of Physics, College of Sciences, Nanjing Agricultural University, Nanjing, Jiangsu 210095, China*
(Dated: January 28, 2025)

We improve the perturbative QCD (PQCD) formalism for multi-body charmless hadronic B meson decays, such as $B \rightarrow VP_3 \rightarrow P_1P_2P_3$, by resolving the possible discrepancy in parametrizing the contribution of the P -wave resonance V to the two-meson distribution amplitudes (DAs) associated with the pairs $P_1P_2 = \pi\pi, K\pi, KK$. The determination of the Gegenbauer moments in the two-meson DAs is then updated in the global fit of the improved PQCD factorization formulas at leading order in the strong coupling α_s to available data for branching ratios and polarization fractions of three- and four-body B decays. The convergence of the Gegenbauer expansion of the resultant two-meson DAs is manifest. The satisfactory quality of the fit implies the consistency of the PQCD framework for multi-body B decays and the universality of the nonperturbative two-meson DAs. In particular, the predicted longitudinal polarization fraction $f_0(B_s^0 \rightarrow K^{*0}\bar{K}^{*0}) = 28.2_{-9.5}^{+8.8}\%$ with the updated Gegenbauer moments matches well the measurement. The observable $L_{K^{*0}\bar{K}^{*0}} = 7.7_{-3.8}^{+4.9}$, defined as the ratio of the longitudinal amplitudes of the two U -spin related channels $B_s^0 \rightarrow K^{*0}\bar{K}^{*0}$ and $B^0 \rightarrow K^{*0}\bar{K}^{*0}$, accommodates the current data within errors. It is found that the direct CP asymmetries $\mathcal{A}_{CP}^{0,\parallel,\perp}$ in the polarization states of some four-body decays $B \rightarrow V_1V_2 \rightarrow (P_1P_2)(P_3P_4)$ might be large, but the destruction among them result in small net CP violation. Our predictions can be confronted with LHCb and Belle-II data in the future.

I. INTRODUCTION

Multi-body B meson decays offer one of the most promising avenues for studying involved strong and weak dynamics and for probing CP violation (CPV). CPV is a number in two-body decays, but depends on the invariant mass of a meson pair in final states of multi-body decays and varies from region to region in a Dalitz plot [1, 2]. Sizable direct CP asymmetries in localized regions of the Dalitz plots for the $B^\pm \rightarrow \pi^\pm\pi^+\pi^-$ and $B^\pm \rightarrow K^\pm K^+K^-$ decays have been experimentally well established [3–5]. The CPV originating from the interference between S - and P -wave resonances was also observed, which causes large local asymmetries [6]. Multi-body B meson decays thus deserve thorough theoretical and phenomenological investigations. Nevertheless, their analysis is considerably more challenging than that of two-body decays owing to complicated interplay between resonant and nonresonant contributions as well as possible significant final-state interactions [7–9].

A factorization formalism that describes multi-body B meson decays in entire phase space is not yet available at present. It seems reasonable to assume the factorization of three-body B meson decays as a working principle, when two final-state mesons are collimated and the bachelor meson recoils back [10–13]. The nonperturbative (collinear) dynamics responsible for the production of the meson pair, including final-state interactions between the two mesons, is absorbed into two-meson distribution amplitudes (DAs) naturally [14–20]. The formulation of three-body B meson decays is then simplified to that of quasi-two-body decays, where a Feynman diagram for hard kernels contains a single virtual gluon exchange at leading order (LO) in the strong coupling α_s . The same idea is applicable to the exploration of four-body charmless hadronic B meson decays, which proceed dominantly with two intermediate resonances.

To attain definite predictions from the above factorization theorem, nonperturbative inputs of two-meson DAs must be known to high precision. We have performed a global fit of the Gegenbauer moments in two-meson DAs to measured branching ratios and direct CP asymmetries of three-body charmless hadronic B meson decays $B \rightarrow VP_3 \rightarrow P_1P_2P_3$ in the perturbative QCD

*Corresponding author: liyakelly@163.com

(PQCD) approach, where V stands for an intermediate vector resonance, and P_i , $i = 1, 2, 3$, represent final-state pseudoscalar mesons [21]. The Gegenbauer moments in the DAs for the mesons $P_3 = \pi, K$ have been fixed in the global analysis of two-body B meson decays [22]. The leading-twist (twist-2) and subleading-twist (twist-3) DAs for the pairs $P_1 P_2 = \pi\pi, K\pi$ and KK with the intermediate vector mesons $V = \rho, K^*$ and ϕ , respectively, were then determined in Ref. [21]. It has been demonstrated that the predictions for various observables in multi-body B meson decays from the fitted Gegenbauer moments are in general consistency with data [23–30].

However, the Gegenbauer moments in the $K\pi$ and KK DAs, being slightly higher than unity [21], are not favored in view of the convergence of the Gegenbauer expansion. Hence, We introduce additional parameters to compensate the possible discrepancy between two theoretical treatments of the hadronic matrix elements for vacuum transition to meson pairs [31]; one is based on the definitions for time-like meson form factors, and another is in terms of Breit-Wigner (BW) propagators for intermediate P -wave resonances. With this new ingredient, we refine the global determination of the Gegenbauer moments, which indeed exhibit better convergence. The branching ratios, CP asymmetries and polarization fractions of numerous three- and four-body B meson decays are then predicted using the updated Gegenbauer moments. It will be shown that the derived longitudinal polarization fraction $f_0(B_s^0 \rightarrow K^{*0} \bar{K}^{*0}) = 28.2_{-9.5}^{+8.8}\%$, distinct from the previous result [23], agrees well with the measurement now. The observable $L_{K^{*0} \bar{K}^{*0}} = 7.7_{-3.8}^{+4.9}$, defined as the ratio of the longitudinal amplitudes of the two U -spin related channels $B_s^0 \rightarrow K^{*0} \bar{K}^{*0}$ and $B^0 \rightarrow K^{*0} \bar{K}^{*0}$ [32], accommodates the current data within errors. It is found that the direct CP asymmetries $\mathcal{A}_{CP}^{0,||,\perp}$ in the polarization states of the four-body decays $B \rightarrow V_1 V_2 \rightarrow (P_1 P_2)(P_3 P_4)$ might be large, but the destruction among them turn in small net CPV. Our predictions can be confronted with LHCb and Belle-II data in the future.

The rest of the paper is organized as follows. The kinematic variables for multi-body hadronic B meson decays are assigned in Sec. II. The two-meson P -wave DAs are parametrized and normalized to time-like form factors, which take the relativistic Breit-Wigner (RBW) model [33] or the Gounaris-Sakurai (GS) model [34]. Such a parametrization is allowed by the Watson theorem [35], that guarantees the factorization of elastic rescattering effects into a final-state meson pair. The extra parameters are associated with the P -wave strength in two-meson DAs. We elaborate how to perform the global fit, and present and discuss the numerical results for branching ratios, polarization fractions and direct CP asymmetries of multi-body B meson decays in Sec. III, which is followed by the Conclusion.

II. THEORETICAL FRAMEWORK

A. Kinematics

Consider the charmless B meson decay into three pseudoscalar mesons via an intermediate vector resonance, $B(p_B) \rightarrow V(p)P_3(p_3) \rightarrow P_1(p_1)P_2(p_2)P_3(p_3)$, with the momenta $p_B = p + p_3$ and $p = p_1 + p_2$. We work in the B meson rest frame and arrange the meson pair $P_1 P_2$ and the bachelor meson P_3 to move in the directions $n = (1, 0, 0_T)$ and $v = (0, 1, 0_T)$, respectively. The above momenta in the light-cone coordinates are chosen as

$$\begin{aligned} p_B &= \frac{m_B}{\sqrt{2}}(1, 1, \mathbf{0}_T), & k_B &= \left(0, x_B \frac{m_B}{\sqrt{2}}, \mathbf{k}_{BT}\right), \\ p &= \frac{m_B}{\sqrt{2}}(f_+, f_-, \mathbf{0}_T), & k &= \left(z f_+ \frac{m_B}{\sqrt{2}}, 0, \mathbf{k}_T\right), \\ p_3 &= \frac{m_B}{\sqrt{2}}(g_-, g_+, \mathbf{0}_T), & k_3 &= \left(0, x_3 g_+ \frac{m_B}{\sqrt{2}}, \mathbf{k}_{3T}\right), \end{aligned} \quad (1)$$

where m_B is the B meson mass, and k_B, k and k_3 are the valence quark momenta in the B meson, the meson pair and the bachelor meson, respectively, with the parton momentum fractions (transverse momenta) x_B, z and x_3 (k_{BT}, k_T and k_{3T}). The functions f_{\pm} and g_{\pm} in Eq. (1) read

$$\begin{aligned} f_{\pm} &= \frac{1}{2} \left(1 + \eta - r_3 \pm \sqrt{(1 - \eta)^2 - 2r_3(1 + \eta) + r_3^2}\right), \\ g_{\pm} &= \frac{1}{2} \left(1 - \eta + r_3 \pm \sqrt{(1 - \eta)^2 - 2r_3(1 + \eta) + r_3^2}\right), \end{aligned} \quad (2)$$

with the ratios $r_3 = m_{P_3}^2/m_B^2$ and $\eta = \omega^2/m_B^2$, m_{P_3} being the bachelor meson mass and $\omega^2 = p^2$ being the invariant mass squared of the meson pair. For a P -wave meson pair, we introduce the longitudinal polarization vector

$$\epsilon = \frac{1}{\sqrt{2}\eta}(f_+, -f_-, \mathbf{0}_T). \quad (3)$$

The individual meson momenta $p_{1,2}$ in the meson pair are then derived as

$$\begin{aligned} p_1 &= \left(\left(\zeta + \frac{r_1 - r_2}{2\eta} \right) f_+ \frac{m_B}{\sqrt{2}}, \left(1 - \zeta + \frac{r_1 - r_2}{2\eta} \right) f_- \frac{m_B}{\sqrt{2}}, \mathbf{p}_T \right), \\ p_2 &= \left(\left(1 - \zeta - \frac{r_1 - r_2}{2\eta} \right) f_+ \frac{m_B}{\sqrt{2}}, \left(\zeta - \frac{r_1 - r_2}{2\eta} \right) f_- \frac{m_B}{\sqrt{2}}, -\mathbf{p}_T \right), \\ p_T^2 &= \zeta(1 - \zeta)\omega^2 + \frac{(m_{P_1}^2 - m_{P_2}^2)^2}{4\omega^2} - \frac{m_{P_1}^2 + m_{P_2}^2}{2}, \end{aligned} \quad (4)$$

from the relation $p = p_1 + p_2$ and the on-shell conditions $p_{1,2}^2 = m_{P_{1,2}}^2$, $m_{P_{1,2}}$ being the $P_{1,2}$ meson masses, with the ratios $r_{1,2} = m_{P_{1,2}}^2/m_B^2$. The variable $\zeta + (r_1 - r_2)/(2\eta) = p_1^+/p^+$ bears the meaning of the meson momentum fraction up to corrections from the final state masses. Alternatively, one can define the polar angle θ of the meson P_1 in the P_1P_2 pair rest frame. The transformation between the B meson rest frame and the meson pair rest frame leads to the relation between the meson momentum fraction ζ and the polar angle θ ,

$$2\zeta - 1 = \sqrt{1 - 2\frac{r_1 + r_2}{\eta} + \frac{(r_1 - r_2)^2}{\eta^2}} \cos \theta, \quad (5)$$

with the bounds

$$\zeta_{\max,\min} = \frac{1}{2} \left[1 \pm \sqrt{1 - 2\frac{r_1 + r_2}{\eta} + \frac{(r_1 - r_2)^2}{\eta^2}} \right]. \quad (6)$$

The $B \rightarrow P_1P_2P_3$ decay amplitude \mathcal{M} is expressed, in the PQCD factorization approach, as

$$\mathcal{M} = \Phi_B \otimes H \otimes \Phi_{P_1P_2} \otimes \Phi_{P_3}, \quad (7)$$

where Φ_B (Φ_{P_3}) is the B (bachelor) meson DA, and the two-meson DA $\Phi_{P_1P_2}$ absorbs the nonperturbative dynamics responsible for the production of the meson pair P_1P_2 . The hard kernel H , containing only one virtual gluon, gathers the perturbative strong and electroweak interactions, as in the formalism for two-body decays. The symbol \otimes denotes the convolution of the above factors in the parton momenta. The corresponding branching ratio is given by [36]

$$\mathcal{B} = \frac{\tau_B m_B}{256\pi^3} \int_{(\sqrt{r_1} + \sqrt{r_2})^2}^1 d\eta \sqrt{(1 - \eta)^2 - 2r_3(1 + \eta) + r_3^2} \int_{\zeta_{\min}}^{\zeta_{\max}} d\zeta |\mathcal{M}|^2, \quad (8)$$

with the B meson lifetime τ_B . A direct CP asymmetry \mathcal{A}_{CP} is defined as

$$\mathcal{A}_{CP} = \frac{\mathcal{B}(\bar{B} \rightarrow \bar{f}) - \mathcal{B}(B \rightarrow f)}{\mathcal{B}(\bar{B} \rightarrow f) + \mathcal{B}(B \rightarrow f)}. \quad (9)$$

For the four-body decay $B \rightarrow V_1V_2 \rightarrow (P_1P_2)(P_3P_4)$, the kinematics is described in a way similar to that for a three-body decay, which has been specified in our previous work [23], and will not be presented explicitly here. There are three helicity amplitudes, A_0 for the longitudinal polarization, and A_{\parallel} and A_{\perp} for the transverse polarizations with spins being parallel and perpendicular to each other, respectively. The associated polarization fractions f_h , $h = 0, \parallel$ and \perp , the direct CP asymmetry in each polarization component and the overall asymmetry are defined as

$$f_h = \frac{\mathcal{B}_h}{\mathcal{B}_0 + \mathcal{B}_{\parallel} + \mathcal{B}_{\perp}}, \quad \mathcal{A}_{CP}^h = \frac{\bar{\mathcal{B}}_h - \mathcal{B}_h}{\bar{\mathcal{B}}_h + \mathcal{B}_h}, \quad \mathcal{A}_{CP}^{\text{dir}} = \frac{\sum_h \bar{\mathcal{B}}_h - \sum_h \mathcal{B}_h}{\sum_h \bar{\mathcal{B}}_h + \sum_h \mathcal{B}_h}, \quad (10)$$

where $\bar{\mathcal{B}}_h$ is the branching ratio of the CP -conjugate channel. It is obvious that the polarization fractions f_h obey the normalization $f_0 + f_{\parallel} + f_{\perp} = 1$.

B. Distribution Amplitudes

We focus on the leading-power contribution from the B meson wave function, which has been widely adopted in the PQCD calculations [37–42],

$$\Phi_B = \frac{i}{\sqrt{2N_c}} (\not{p}_B + m_B) \gamma_5 \phi_B(x, b), \quad (11)$$

with the impact parameter b being conjugate to the parton transverse momentum k_{BT} . The B meson DA $\phi_B(x, b)$ is parametrized as

$$\phi_B(x, b) = N_B x^2 (1-x)^2 \exp \left[-\frac{m_B^2 x^2}{2\omega_B^2} - \frac{1}{2}(\omega_B b)^2 \right], \quad (12)$$

where the constant N_B is related to the B meson decay constant f_B through the normalization condition $\int_0^1 dx \phi_B(x, b=0) = f_B/(2\sqrt{2N_c})$. The shape parameter takes the values $\omega_B = 0.40$ GeV for B^+ , B^0 mesons and $\omega_{B_s} = 0.48$ GeV [22, 37, 43, 44] for a B_s^0 meson with 10% variation in the numerical study below.

The light-cone hadronic matrix element for a B meson contains in fact two DAs ϕ_B and $\bar{\phi}_B$, which are the linear combinations of ϕ_B^+ and ϕ_B^- defined in the literature [45], $\phi_B = (\phi_B^+ + \phi_B^-)/2$ and $\bar{\phi}_B = (\phi_B^+ - \phi_B^-)/2$. It has been verified that the next-to-power contribution from $\bar{\phi}_B$ is numerically suppressed [38, 39, 46] compared with the leading-power one from ϕ_B . For instance, the former contribution to the $B \rightarrow \pi$ transition form factor $F_0^{B \rightarrow \pi}$ is about 20% of the latter as shown in the PQCD evaluation [46]. The higher-twist B meson DAs have been designated systematically in the heavy quark effective theory [47], which are decomposed according to the twist and conformal spin assignments up to twist 6. In principle, all the next-to-leading-power sources can be taken into account for a complete analysis. However, the currently available data are not sufficient for their determination.

The light-cone matrix element for a pseudoscalar meson is decomposed, up to twist 3, into [38, 39]

$$\Phi_P \equiv \frac{i}{\sqrt{2N_c}} \gamma_5 [p_3 \phi_P^A(x_3) + m_{03} \phi_P^P(x_3) + m_{03} (\not{p} - 1) \phi_P^T(x_3)], \quad (13)$$

with $P = \pi, K$ and the chiral scale m_{03} . The involved DAs ϕ_P^i are expanded, in terms of the Gegenbauer polynomials,

$$C_1^{3/2}(t) = 3t, \quad C_2^{3/2}(t) = \frac{3}{2}(5t^2 - 1), \quad C_4^{3/2}(t) = \frac{15}{8}(1 - 14t^2 + 21t^4). \quad (14)$$

as

$$\begin{aligned} \phi_\pi^A(x) &= \frac{3f_\pi}{\sqrt{6}} x(1-x) [1 + a_2^\pi C_2^{3/2}(2x-1) + a_4^\pi C_4^{3/2}(2x-1)], \\ \phi_\pi^P(x) &= \frac{f_\pi}{2\sqrt{6}} [1 + a_{2P}^\pi C_2^{1/2}(2x-1)], \\ \phi_\pi^T(x) &= \frac{f_\pi}{2\sqrt{6}} (1-2x) [1 + a_{2T}^\pi (10x^2 - 10x + 1)], \\ \phi_K^A(x) &= \frac{3f_K}{\sqrt{6}} x(1-x) [1 + a_1^K C_1^{3/2}(2x-1) + a_2^K C_2^{3/2}(2x-1) + a_4^K C_4^{3/2}(2x-1)], \\ \phi_K^P(x) &= \frac{f_K}{2\sqrt{6}} [1 + a_{2P}^K C_2^{1/2}(2x-1)], \\ \phi_K^T(x) &= -\frac{f_K}{2\sqrt{6}} [C_1^{1/2}(x) + a_{2T}^K C_3^{1/2}(x)], \end{aligned} \quad (15)$$

with the pion (kaon) decay constant f_π (f_K). The Gegenbauer coefficients in the above pion and kaon DAs have been derived at the scale 1 GeV in a recent global analysis [22] based on the LO PQCD factorization formulas, which are summarized as

$$\begin{aligned} a_2^\pi &= 0.64 \pm 0.08, & a_4^\pi &= -0.41 \pm 0.10, & a_{2P}^\pi &= 1.08 \pm 0.15, & a_{2T}^\pi &= -0.48 \pm 0.33, \\ a_1^K &= 0.33 \pm 0.08, & a_2^K &= 0.28 \pm 0.10, & a_4^K &= -0.40 \pm 0.07, & a_{2P}^K &= 0.24, & a_{2T}^K &= 0.35. \end{aligned} \quad (16)$$

Note that the twist-3 DAs ϕ_K^P and ϕ_K^T were not obtained in Ref. [22], but quoted from the sum-rule results [48].

The P -wave two-meson DAs for both longitudinal and transverse polarizations are decomposed, up to twist 3, into [49, 50]

$$\Phi_P^L(z, \zeta, \omega) = \frac{1}{\sqrt{2N_c}} \left[\omega \not{\epsilon}_P \phi_P^0(z, \omega^2) + \omega \phi_P^s(z, \omega^2) + \frac{\not{p}_1 \not{p}_2 - \not{p}_2 \not{p}_1}{\omega(2\zeta-1)} \phi_P^t(z, \omega^2) \right] (2\zeta-1), \quad (17)$$

$$\begin{aligned} \Phi_P^T(z, \zeta, \omega) &= \frac{1}{\sqrt{2N_c}} \left[\gamma_5 \not{\epsilon}_T \not{p} \phi_P^T(z, \omega^2) + \omega \gamma_5 \not{\epsilon}_T \phi_P^a(z, \omega^2) + i\omega \frac{\epsilon^{\mu\nu\rho\sigma} \gamma_\mu \epsilon_{T\nu} p_\rho n_{-\sigma}}{p \cdot n_-} \phi_P^v(z, \omega^2) \right] \\ &\cdot \sqrt{\zeta(1-\zeta) + \alpha}, \end{aligned} \quad (18)$$

with the kinematic parameter $\alpha = (r_1 - r_2)^2/(4\eta^2) - (r_1 + r_2)/(2\eta)$. The DAs ϕ_P^i are expanded as

$$\phi_{\pi\pi}^0(z, \omega^2) = \frac{3F_{\pi\pi}^{\parallel}(\omega^2)}{\sqrt{2N_c}} z(1-z) \left[1 + a_{2\rho}^0 \frac{3}{2} (5t^2 - 1) \right], \quad (19)$$

$$\phi_{\pi\pi}^s(z, \omega^2) = \frac{3F_{\pi\pi}^{\perp}(\omega^2)}{2\sqrt{2N_c}} t \left[1 + a_{2\rho}^s (10z^2 - 10z + 1) \right], \quad (20)$$

$$\phi_{\pi\pi}^t(z, \omega^2) = \frac{3F_{\pi\pi}^{\perp}(\omega^2)}{2\sqrt{2N_c}} t^2 \left[1 + a_{2\rho}^t \frac{3}{2} (5t^2 - 1) \right], \quad (21)$$

$$\phi_{\pi\pi}^T(z, \omega^2) = \frac{3F_{\pi\pi}^{\perp}(\omega^2)}{\sqrt{2N_c}} z(1-z) \left[1 + a_{2\rho}^T \frac{3}{2} (5t^2 - 1) \right], \quad (22)$$

$$\phi_{\pi\pi}^a(z, \omega^2) = \frac{3F_{\pi\pi}^{\parallel}(\omega^2)}{4\sqrt{2N_c}} t \left[1 + a_{2\rho}^a (10z^2 - 10z + 1) \right], \quad (23)$$

$$\phi_{\pi\pi}^v(z, \omega^2) = \frac{3F_{\pi\pi}^{\parallel}(\omega^2)}{8\sqrt{2N_c}} \left[1 + t^2 + a_{2\rho}^v (3t^2 - 1) \right], \quad (24)$$

$$\phi_{K\pi}^0(z, \omega^2) = \frac{3F_{K\pi}^{\parallel}(\omega^2)}{\sqrt{2N_c}} z(1-z) \left[1 + a_{1K^*}^{\parallel} 3t + a_{2K^*}^{\parallel} \frac{3}{2} (5t^2 - 1) \right], \quad (25)$$

$$\phi_{K\pi}^s(z, \omega^2) = \frac{3F_{K\pi}^{\perp}(\omega^2)}{2\sqrt{2N_c}} t, \quad (26)$$

$$\phi_{K\pi}^t(z, \omega^2) = \frac{3F_{K\pi}^{\perp}(\omega^2)}{2\sqrt{2N_c}} t^2, \quad (27)$$

$$\phi_{K\pi}^T(z, \omega^2) = \frac{3F_{K\pi}^{\perp}(\omega^2)}{\sqrt{2N_c}} z(1-z) \left[1 + a_{1K^*}^{\perp} 3t + a_{2K^*}^{\perp} \frac{3}{2} (5t^2 - 1) \right], \quad (28)$$

$$\phi_{K\pi}^a(z, \omega^2) = \frac{3F_{K\pi}^{\parallel}(\omega^2)}{4\sqrt{2N_c}} t, \quad (29)$$

$$\phi_{K\pi}^v(z, \omega^2) = \frac{3F_{K\pi}^{\parallel}(\omega^2)}{8\sqrt{2N_c}} (1 + t^2), \quad (30)$$

$$\phi_{KK}^0(z, \omega^2) = \frac{3F_{KK}^{\parallel}(\omega^2)}{\sqrt{2N_c}} z(1-z) \left[1 + a_{2\phi}^0 \frac{3}{2} (5t^2 - 1) \right], \quad (31)$$

$$\phi_{KK}^s(z, \omega^2) = \frac{3F_{KK}^{\perp}(\omega^2)}{2\sqrt{2N_c}} t, \quad (32)$$

$$\phi_{KK}^t(z, \omega^2) = \frac{3F_{KK}^{\perp}(\omega^2)}{2\sqrt{2N_c}} t^2, \quad (33)$$

$$\phi_{KK}^T(z, \omega^2) = \frac{3F_{KK}^{\perp}(\omega^2)}{\sqrt{2N_c}} z(1-z) \left[1 + a_{2\phi}^T \frac{3}{2} (5t^2 - 1) \right], \quad (34)$$

$$\phi_{KK}^a(z, \omega^2) = \frac{3F_{KK}^{\parallel}(\omega^2)}{4\sqrt{2N_c}} t, \quad (35)$$

$$\phi_{KK}^v(z, \omega^2) = \frac{3F_{KK}^{\parallel}(\omega^2)}{8\sqrt{2N_c}} (1 + t^2), \quad (36)$$

with the variable $t = 1 - 2z$ and the time-like form factors $F^{\parallel,\perp}(\omega^2)$. The twist-3 $K\pi$ DAs $\phi_{K\pi}^{a,v}$ and KK DAs $\phi_{KK}^{s,t}$ have been fixed to the asymptotic forms owing to the limited amount of data. The Gegenbauer coefficients $a_{2\rho}^{0(s,t)}$, $a_{1(2)K^*}^{\parallel}$ and $a_{2\phi}^{0(T)}$ were determined in a recent global investigation of three- and four-body B meson decays based on the PQCD formalism [21, 26, 29]. Note that the moments $a_{2\rho}^{T,a,v}$ in Eqs. (22)-(24) cannot be acquired from present fit, since only the measured transverse polarization fraction $f_{\perp}(B^0 \rightarrow \rho^0 K^{*0})$ has been reported [51] so far. They are thus set to the values $a_{2\rho}^T = 0.5 \pm 0.5$, $a_{2\rho}^a = 0.4 \pm 0.4$ and $a_{2\rho}^v = -0.5 \pm 0.5$ deduced in Ref. [50].

According to the Watson theorem [35], elastic rescattering effects in a final-state meson pair can be absorbed into the time-like

form factors $F^{\parallel,\perp}(\omega^2)$ as mentioned before. We employ the RBW line shape for the form factors

$$F_{K\pi(KK)}^{\parallel}(\omega^2) = \frac{m_{K^*(\phi)}^2}{m_{K^*(\phi)}^2 - \omega^2 - im_{K^*(\phi)}\Gamma_{K^*(\phi)}(\omega^2)}, \quad (37)$$

associated with the narrow intermediate resonances K^* and ϕ , $m_{K^*(\phi)}$ and $\Gamma_{K^*(\phi)}$ being the pole mass and width, respectively. The mass-dependent width is written as

$$\Gamma_{K^*(\phi)}(\omega^2) = \Gamma_{K^*(\phi)} \left(\frac{m_{K^*(\phi)}}{\omega} \right) \left(\frac{k(\omega)}{k(m_{K^*(\phi)})} \right)^{(2L_R+1)}, \quad (38)$$

where $k(\omega)$ is the momentum vector of the decay product measured in the resonance rest frame, and $k(m_{K^*(\phi)})$ is the value of $k(\omega)$ at $\omega = m_{K^*(\phi)}$. Its explicit expression is given by

$$k(\omega) = \frac{\sqrt{\lambda(\omega^2, m_{h_1}^2, m_{h_2}^2)}}{2\omega}, \quad (39)$$

with the Källén function $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2(ab + ac + bc)$. The orbital angular momenta $L_R = 0, 1, \dots$ correspond to the S, P, \dots partial-wave resonances in the $\pi\pi, K\pi$ and KK systems.

The contribution from a broad ρ resonance is usually parameterized as the GS model [34] based on the BW function [52] in the experimental survey of multi-body hadronic B meson decays. Taking into account the ρ - ω interference and excited state contributions, we have the form factor [49, 53, 54]

$$F_P^{\parallel}(\omega^2) = \left[\text{GS}_\rho(\omega^2, m_\rho, \Gamma_\rho) \frac{1 + c_\omega \text{BW}_\omega(\omega^2, m_\omega, \Gamma_\omega)}{1 + c_\omega} + \Sigma c_j \text{GS}_j(\omega^2, m_j, \Gamma_j) \right] (1 + \Sigma c_j)^{-1}, \quad (40)$$

where $m_{\rho,\omega,j}$ ($\Gamma_{\rho,\omega,j}$), $j = \rho'(1450), \rho''(1700)$ and $\rho'''(2254)$, are the masses (decay widths) of the series of ρ resonances, and $c_{\omega,j}$ are the corresponding weights. The function $\text{GS}_\rho(s, m_\rho, \Gamma_\rho)$ reads

$$\text{GS}_\rho(\omega^2, m_\rho, \Gamma_\rho) = \frac{m_\rho^2 [1 + d(m_\rho)\Gamma_\rho/m_\rho]}{m_\rho^2 - \omega^2 + f(\omega^2, m_\rho, \Gamma_\rho) - im_\rho\Gamma(\omega^2, m_\rho, \Gamma_\rho)}, \quad (41)$$

with the factors

$$\begin{aligned} \Gamma(s, m_\rho, \Gamma_\rho) &= \Gamma_\rho \frac{s}{m_\rho^2} \left(\frac{\beta_\pi(s)}{\beta_\pi(m_\rho^2)} \right)^3, \\ d(m) &= \frac{3}{\pi} \frac{m_\pi^2}{k^2(m^2)} \ln \left(\frac{m + 2k(m^2)}{2m_\pi} \right) + \frac{m}{2\pi k(m^2)} - \frac{m_\pi^2 m}{\pi k^3(m^2)}, \\ f(s, m, \Gamma) &= \frac{\Gamma m^2}{k^3(m^2)} [k^2(s)[h(s) - h(m^2)] + (m^2 - s)k^2(m^2)h'(m^2)], \\ k(s) &= \frac{1}{2} \sqrt{s} \beta_\pi(s), \quad h(s) = \frac{2}{\pi} \frac{k(s)}{\sqrt{s}} \ln \left(\frac{\sqrt{s} + 2k(s)}{2m_\pi} \right), \end{aligned} \quad (42)$$

and $\beta_\pi(s) = \sqrt{1 - 4m_\pi^2/s}$. For the poorly known form factor $F^\perp(\omega^2)$, we assume the relation $F^\perp(\omega^2)/F^\parallel(\omega^2) \approx f_V^T/f_V$, in which f_V and f_V^T are the decay constants of the intermediate vector resonance V .

C. P -wave Parametrization

The amplitude of the three-body decay $B(p_B) \rightarrow P_1(p_1)P_2(p_2)P_3(p_3)$ contains the factorizable contribution

$$\frac{G_F}{\sqrt{2}} \langle P_1(p_1)P_2(p_2) | J_\mu | 0 \rangle \langle P_3(p_3) | J^\mu | B(p_B) \rangle, \quad (43)$$

with the vector current operators J_μ . The $B \rightarrow P_3$ vector and scalar transition form factors $F_1^{BP_3}(p^2)$ and $F_0^{BP_3}(p^2)$, respectively, are defined via the matrix element

$$\langle P_3(p_3) | J^\mu | B(p_B) \rangle = \left[(p_B + p_3)^\mu - \frac{m_B^2 - m_{P_3}^2}{p^2} p^\mu \right] F_1^{BP_3}(p^2) + \frac{m_B^2 - m_{P_3}^2}{p^2} p^\mu F_0^{BP_3}(p^2). \quad (44)$$

Similarly, the matrix element for the transition from vacuum to the $P_1 P_2$ meson pair defines the vector (scalar) form factor $F_1^{P_1 P_2}(p^2)$ ($F_0^{P_1 P_2}(p^2)$),

$$\langle P_1(p_1)P_2(p_2)|J_\mu|0\rangle = \left[(p_1 - p_2)_\mu - \frac{m_{P_1}^2 - m_{P_2}^2}{p^2} p_\mu \right] F_1^{P_1 P_2}(p^2) + \frac{m_{P_1}^2 - m_{P_2}^2}{p^2} p_\mu F_0^{P_1 P_2}(p^2). \quad (45)$$

See Eq. (36) of Ref. [55].

The above subprocess can be described in an alternative way: the quark pair from a hard decay kernel forms the intermediate vector meson V , which propagates following the BW factor, and then proceeds with the $V \rightarrow P_1 P_2$ transition. We write

$$\begin{aligned} \langle P_1(p_1)P_2(p_2)|J_\mu|0\rangle &= \sum_{\lambda=0,\pm 1} \langle P_1(p_1)P_2(p_2)|V, \epsilon^\lambda\rangle \frac{1}{m_V^2 - s - im_V \Gamma(s)} \langle V, \epsilon^\lambda|J_\mu|0\rangle \\ &= \frac{g^{V \rightarrow P_1 P_2}}{m_V^2 - s - im_V \Gamma(s)} \sum_{\lambda=0,\pm 1} \epsilon^\lambda \cdot (p_1 - p_2) \langle V, \epsilon^\lambda|J_\mu|0\rangle \\ &= \frac{g^{V \rightarrow P_1 P_2}}{m_V^2 - s - im_V \Gamma(s)} \sum_{\lambda=0,\pm 1} \epsilon^\lambda \cdot (p_1 - p_2) \epsilon_\mu^\lambda f_V m_V, \end{aligned} \quad (46)$$

where $s = p^2 = (p_1 + p_2)^2$ is the invariant mass squared of the meson pair, ϵ^λ (m_V , f_V) represents the polarization vector (mass, decay constant) of the resonance V , and the coupling strength $g^{V \rightarrow P_1 P_2}$ can be evaluated from Eq. (4.17) in Ref. [56].

The completeness relation for the summation over the polarization states λ

$$\sum_{\lambda=0,\pm 1} \epsilon_\mu^\lambda(p) \epsilon_\nu^{\lambda*}(p) = - \left(g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right), \quad (47)$$

demands, according to Eqs. (45) and (46),

$$\frac{g^{V \rightarrow P_1 P_2}}{m_V^2 - s - im_V \Gamma(s)} \cdot f_V m_V = N_P F_1^{P_1 P_2}(p^2). \quad (48)$$

The coefficient N_P has been introduced to remedy the possible theoretical mismatch between the meson form factors and the properties of the intermediate P -wave resonance. One can roughly estimate the values of N_P for the $\pi\pi$, $K\pi$ and KK meson pairs based on Eqs. (37), (40) and (48),

$$N_{\pi\pi} \approx 1.00, \quad N_{K\pi} \approx 1.40, \quad N_{KK} \approx 1.20. \quad (49)$$

The above three coefficients will be handled as free parameters, determined in our global fit, and then compared with Eq. (49) in the next section.

III. NUMERICAL ANALYSIS

A. Global Fit

To leviate the precision of the two-meson DAs, we update the fitted Gegenbauer moments by including the additional four-body decays $B \rightarrow \rho K^* \rightarrow (\pi\pi)(K\pi)$, $B \rightarrow K^* K^* \rightarrow (K\pi)(K\pi)$ and $B \rightarrow \phi K^* \rightarrow (KK)(K\pi)$ in the global study. The moments $a_{1K^*}^\perp$ and $a_{2K^*}^\perp$ of the transverse $K\pi$ DA in Eq. (28) will be derived for the first time. We specify the inputted masses and widths (in units of GeV) [36] in the numerical analysis,

$$\begin{aligned} m_B &= 5.280, & m_{B_s} &= 5.367, & m_b &= 4.8, & m_{K^\pm} &= 0.494, \\ m_{K^0} &= 0.498, & m_{\pi^\pm} &= 0.140, & m_{\pi^0} &= 0.135, & \Gamma_\rho &= 0.1496 \\ \Gamma_{K^*} &= 0.0473, & \Gamma_\phi &= 0.00425. \end{aligned} \quad (50)$$

The decay constants (in units of GeV) and the B meson lifetimes (in units of ps) take the values [48, 53]

$$\begin{aligned} f_B &= 0.21, & f_{B_s} &= 0.24, & f_\rho &= 0.216, & f_\rho^T &= 0.184, \\ f_{\phi(1020)} &= 0.215, & f_{\phi(1020)}^T &= 0.186, & f_{K^*} &= 0.217, & f_{K^*}^T &= 0.185, \\ \tau_{B^0} &= 1.519, & \tau_{B^\pm} &= 1.638, & \tau_{B_s} &= 1.512. \end{aligned} \quad (51)$$

The Wolfenstein parameters in the CKM matrix are set to $A = 0.836 \pm 0.015$, $\lambda = 0.22453 \pm 0.00044$, $\bar{\rho} = 0.122^{+0.018}_{-0.017}$ and $\bar{\eta} = 0.355^{+0.012}_{-0.011}$ [36].

The amplitudes \mathcal{M} for the considered B meson decays can be expanded in terms of the Gegenbauer moments with the two meson DAs in Eqs. (19)-(36). Taking the $B \rightarrow K^* \phi \rightarrow (K\pi)(KK)$, $B \rightarrow \rho K^* \rightarrow (\pi\pi)(K\pi)$ and $B \rightarrow K^* K^* \rightarrow (K\pi)(K\pi)$ decays as examples, we decompose the longitudinal squared amplitudes $|\mathcal{M}_{K^* \phi}^L|^2$, $|\mathcal{M}_{\rho K^*}^L|^2$ and $|\mathcal{M}_{K^* K^*}^L|^2$ into the linear combinations of the Gegenbauer moments and their products,

$$\begin{aligned} |\mathcal{M}_{K^* \phi}^L|^2 &= M_{0K^* \phi}^L + a_{2\phi}^0 M_{1K^* \phi}^L + a_{1K^*}^{\parallel} M_{2K^* \phi}^L + a_{2K^*}^{\parallel} M_{3K^* \phi}^L + (a_{2\phi}^0)^2 M_{4K^* \phi}^L \\ &+ (a_{1K^*}^{\parallel})^2 M_{5K^* \phi}^L + (a_{2K^*}^{\parallel})^2 M_{6K^* \phi}^L + (a_{2\phi}^0 a_{1K^*}^{\parallel}) M_{7K^* \phi}^L + (a_{2\phi}^0 a_{2K^*}^{\parallel}) M_{8K^* \phi}^L \\ &+ (a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{9K^* \phi}^L + (a_{2\phi}^0)^2 a_{1K^*}^{\parallel} M_{10K^* \phi}^L + (a_{2\phi}^0)^2 a_{2K^*}^{\parallel} M_{11K^* \phi}^L \\ &+ a_{2\phi}^0 (a_{1K^*}^{\parallel})^2 M_{12K^* \phi}^L + a_{2\phi}^0 (a_{2K^*}^{\parallel})^2 M_{13K^* \phi}^L + (a_{2\phi}^0 a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{14K^* \phi}^L \\ &+ (a_{2\phi}^0 a_{1K^*}^{\parallel})^2 M_{15K^* \phi}^L + (a_{2\phi}^0 a_{2K^*}^{\parallel})^2 M_{16K^* \phi}^L + (a_{2\phi}^0)^2 a_{1K^*}^{\parallel} a_{2K^*}^{\parallel} M_{17K^* \phi}^L, \end{aligned} \quad (52)$$

$$\begin{aligned} |\mathcal{M}_{K^* K^*}^L|^2 &= M_{0K^* K^*}^L + a_{1K^*}^{\parallel} M_{1K^* K^*}^L + a_{2K^*}^{\parallel} M_{2K^* K^*}^L + (a_{1K^*}^{\parallel})^2 M_{3K^* K^*}^L + (a_{2K^*}^{\parallel})^2 M_{4K^* K^*}^L \\ &+ (a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{5K^* K^*}^L + (a_{1K^*}^{\parallel})^3 M_{6K^* K^*}^L + (a_{2K^*}^{\parallel})^3 M_{7K^* K^*}^L + (a_{1K^*}^{\parallel}) (a_{2K^*}^{\parallel})^2 M_{8K^* K^*}^L \\ &+ (a_{1K^*}^{\parallel})^2 (a_{2K^*}^{\parallel}) M_{9K^* K^*}^L + (a_{1K^*}^{\parallel})^4 M_{10K^* K^*}^L + (a_{1K^*}^{\parallel})^3 (a_{2K^*}^{\parallel}) M_{11K^* K^*}^L \\ &+ (a_{1K^*}^{\parallel})^2 (a_{2K^*}^{\parallel})^2 M_{12K^* K^*}^L + (a_{1K^*}^{\parallel}) (a_{2K^*}^{\parallel})^3 M_{13K^* K^*}^L + (a_{2K^*}^{\parallel})^4 M_{14K^* K^*}^L, \end{aligned} \quad (53)$$

$$\begin{aligned} |\mathcal{M}_{\rho K^*}^L|^2 &= M_{0\rho K^*}^L + a_{2\rho}^0 M_{1\rho K^*}^L + a_{2\rho}^s M_{2\rho K^*}^L + a_{2\rho}^t M_{3\rho K^*}^L + a_{1K^*}^{\parallel} M_{4\rho K^*}^L + a_{2K^*}^{\parallel} M_{5\rho K^*}^L \\ &+ (a_{2\rho}^0)^2 M_{6\rho K^*}^L + (a_{2\rho}^s)^2 M_{7\rho K^*}^L + (a_{2\rho}^t)^2 M_{8\rho K^*}^L + (a_{1K^*}^{\parallel})^2 M_{9\rho K^*}^L + (a_{2K^*}^{\parallel})^2 M_{10\rho K^*}^L \\ &+ (a_{2\rho}^0 a_{2\rho}^s) M_{11\rho K^*}^L + (a_{2\rho}^0 a_{2\rho}^t) M_{12\rho K^*}^L + (a_{2\rho}^0 a_{1K^*}^{\parallel}) M_{13\rho K^*}^L + (a_{2\rho}^0 a_{2K^*}^{\parallel}) M_{14\rho K^*}^L \\ &+ (a_{2\rho}^s a_{2\rho}^t) M_{15\rho K^*}^L + (a_{2\rho}^s a_{1K^*}^{\parallel}) M_{16\rho K^*}^L + (a_{2\rho}^s a_{2K^*}^{\parallel}) M_{17\rho K^*}^L + (a_{2\rho}^t a_{1K^*}^{\parallel}) M_{18\rho K^*}^L \\ &+ (a_{2\rho}^t a_{2K^*}^{\parallel}) M_{19\rho K^*}^L + (a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{20\rho K^*}^L + (a_{2\rho}^0)^2 (a_{1K^*}^{\parallel}) M_{21\rho K^*}^L + (a_{2\rho}^0)^2 (a_{2K^*}^{\parallel}) M_{22\rho K^*}^L \\ &+ (a_{2\rho}^0 a_{2\rho}^s a_{1K^*}^{\parallel}) M_{23\rho K^*}^L + (a_{2\rho}^0 a_{2\rho}^s a_{2K^*}^{\parallel}) M_{24\rho K^*}^L + (a_{2\rho}^0 a_{2\rho}^t a_{1K^*}^{\parallel}) M_{25\rho K^*}^L + (a_{2\rho}^0 a_{2\rho}^t a_{2K^*}^{\parallel}) M_{26\rho K^*}^L \\ &+ (a_{2\rho}^s)^2 (a_{1K^*}^{\parallel}) M_{27\rho K^*}^L + (a_{2\rho}^s)^2 (a_{2K^*}^{\parallel}) M_{28\rho K^*}^L + (a_{2\rho}^s a_{2\rho}^t a_{1K^*}^{\parallel}) M_{29\rho K^*}^L + (a_{2\rho}^s a_{2\rho}^t a_{2K^*}^{\parallel}) M_{30\rho K^*}^L \\ &+ (a_{2\rho}^t)^2 (a_{1K^*}^{\parallel}) M_{31\rho K^*}^L + (a_{2\rho}^t)^2 (a_{2K^*}^{\parallel}) M_{32\rho K^*}^L + (a_{2\rho}^0) (a_{1K^*}^{\parallel})^2 M_{33\rho K^*}^L + (a_{2\rho}^0 a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{34\rho K^*}^L \\ &+ (a_{2\rho}^s) (a_{1K^*}^{\parallel})^2 M_{35\rho K^*}^L + (a_{2\rho}^s a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{36\rho K^*}^L + (a_{2\rho}^t) (a_{1K^*}^{\parallel})^2 M_{37\rho K^*}^L + (a_{2\rho}^t a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{38\rho K^*}^L \\ &+ (a_{2\rho}^0) (a_{2K^*}^{\parallel})^2 M_{39\rho K^*}^L + (a_{2\rho}^s) (a_{2K^*}^{\parallel})^2 M_{40\rho K^*}^L + (a_{2\rho}^t) (a_{2K^*}^{\parallel})^2 M_{41\rho K^*}^L + (a_{2\rho}^0)^2 (a_{1K^*}^{\parallel})^2 M_{42\rho K^*}^L \\ &+ (a_{2\rho}^0)^2 (a_{2K^*}^{\parallel})^2 M_{43\rho K^*}^L + (a_{2\rho}^s)^2 (a_{1K^*}^{\parallel})^2 M_{44\rho K^*}^L + (a_{2\rho}^s)^2 (a_{2K^*}^{\parallel})^2 M_{45\rho K^*}^L + (a_{2\rho}^t)^2 (a_{1K^*}^{\parallel})^2 M_{46\rho K^*}^L \\ &+ (a_{2\rho}^t)^2 (a_{2K^*}^{\parallel})^2 M_{47\rho K^*}^L + (a_{2\rho}^0)^2 (a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{48\rho K^*}^L + (a_{2\rho}^0 a_{2\rho}^s a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{49\rho K^*}^L \\ &+ (a_{2\rho}^0 a_{2\rho}^s) (a_{1K^*}^{\parallel})^2 M_{50\rho K^*}^L + (a_{2\rho}^0 a_{2\rho}^t) (a_{1K^*}^{\parallel})^2 M_{51\rho K^*}^L + (a_{2\rho}^0 a_{2\rho}^t a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{52\rho K^*}^L \\ &+ (a_{2\rho}^0 a_{2\rho}^s) (a_{2K^*}^{\parallel})^2 M_{53\rho K^*}^L + (a_{2\rho}^0 a_{2\rho}^t) (a_{2K^*}^{\parallel})^2 M_{54\rho K^*}^L + (a_{2\rho}^s)^2 (a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{55\rho K^*}^L \\ &+ (a_{2\rho}^s a_{2\rho}^t) (a_{1K^*}^{\parallel})^2 M_{56\rho K^*}^L + (a_{2\rho}^s a_{2\rho}^t a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{57\rho K^*}^L + (a_{2\rho}^s a_{2\rho}^t) (a_{2K^*}^{\parallel})^2 M_{58\rho K^*}^L \\ &+ (a_{2\rho}^t)^2 (a_{1K^*}^{\parallel} a_{2K^*}^{\parallel}) M_{59\rho K^*}^L. \end{aligned} \quad (54)$$

The transverse squared amplitudes $|\mathcal{M}_{K^* \phi}^{N(T)}|^2$ and $|\mathcal{M}_{K^* K^*}^{N(T)}|^2$ are inferred from the above expressions through the transformations

$$a_{2\phi}^0 \rightarrow a_{2\phi}^T, \quad a_{1K^*}^{\parallel} \rightarrow a_{1K^*}^{\perp}, \quad a_{2K^*}^{\parallel} \rightarrow a_{2K^*}^{\perp}, \quad M_{iK^* \phi}^L \rightarrow M_{iK^* \phi}^{N(T)}, \quad M_{iK^* K^*}^L \rightarrow M_{iK^* K^*}^{N(T)}. \quad (55)$$

We then compute the coefficients M , which contain only the Gegenbauer polynomials, to establish the database for the global fit. The formulas of the squared amplitudes for the $B \rightarrow VP_3 \rightarrow P_1 P_2 P_3$ and $B \rightarrow \rho\rho \rightarrow (\pi\pi)(\pi\pi)$, $B \rightarrow \phi\phi \rightarrow (KK)(KK)$ decays can be found in Refs. [21, 26, 29].

Repeating the procedure in Ref. [22], we determine the Gegenbauer moments in the two-meson DAs by fitting Eqs. (52)–(54) with the Gegenbauer-moment-independent database to the measured branching ratios and polarization fractions of three- and

four-body B meson decays. The standard nonlinear least- χ^2 (lsq) method [57] is employed, in which the χ^2 function is defined for n pieces of experimental data $v_i \pm \delta v_i$ with the errors δv_i and the corresponding fitted values v_i^{th} as

$$\chi^2 = \sum_{i=1}^n \left(\frac{v_i - v_i^{\text{th}}}{\delta v_i} \right)^2. \quad (56)$$

The inclusion of more data in the fit decreases statistical uncertainties. The measurements with significance lower than 3σ , which do not impose stringent constraints, need not be taken into account. The data of those modes, which are greatly impacted by subleading contributions according to the existent PQCD calculations [58, 59], such as $B^0 \rightarrow \pi^0 \rho^0 \rightarrow \pi^0(\pi^+ \pi^-)$ and $B^0 \rightarrow \rho^0 \rho^0 \rightarrow (\pi^+ \pi^-)(\pi^+ \pi^-)$, are also excluded, even though they may have higher precision.

B. Gegenbauer Moments of Two-meson DAs

The Gegenbauer moments of the twist-2 and twist-3 DAs for the $\pi\pi$, $K\pi$ and KK pairs from a joint fit with $\chi^2 = 1.6$ are presented in Table I, whose errors mainly arise from experimental uncertainties. The fitted $N_{\pi\pi} = 1.05 \pm 0.04$, $N_{K\pi} = 1.48 \pm 0.03$ and $N_{KK} = 1.22 \pm 0.03$ are basically consistent with the expected values in Eq. (49). The latter two confirm that the discrepancy for the P -wave parametrization is remarkable for the $K\pi$ and KK pairs. The longitudinal $\pi\pi$ DAs are from the fit to the eight pieces of $B \rightarrow P(\rho \rightarrow)\pi\pi$ data and the six pieces of $B \rightarrow \rho\rho, \rho K^*$ data marked by “†” in Tables II-III. As mentioned before, it is not practical to fit the parameters $a_{2\rho}^{T,v,a}$ in the transverse $\pi\pi$ DAs because of the limited experimental inputs, which have been set to the values extracted in Ref. [50]. It is seen that our Gegenbauer moments deviate from the ones of the $\rho(770)$ meson DAs in QCD sum rules [60]; the ρ - ω mixing effect and the contributions from higher ρ resonances with finite widths have been included in the $\pi\pi$ DAs via Eq. (40), so they need not be the same as the $\rho(770)$ meson DAs. The added $B^+ \rightarrow \rho^0 K^{*+}$ and $B^+ \rightarrow \rho^+ K^{*0}$ data constrain effectively the corresponding Gegenbauer moments, leading to $a_{2\rho}^0 = 0.39 \pm 0.11$, $a_{2\rho}^s = -0.34 \pm 0.26$ and $a_{2\rho}^t = -0.13 \pm 0.04$, which are distinct from those in Ref. [29].

The moments $a_{1K^*}^{\parallel}$ and $a_{2K^*}^{\parallel}$ for the twist-2 $K\pi$ DA $\phi_{K\pi}^0$, and $a_{1K^*}^{\perp}$ and $a_{2K^*}^{\perp}$ for the twist-2 DA $\phi_{K\pi}^T$ are governed by the six pieces of $B_{(s)} \rightarrow P(K^* \rightarrow)K\pi$ data and the twelve pieces of $B_{(s)} \rightarrow K^*\rho, K^*K^*, K^*\phi$ data. We highlight that $a_{1K^*}^{\perp} = 0.61 \pm 0.21$ and $a_{2K^*}^{\perp} = 0.45 \pm 0.06$ in the transverse component $\phi_{K\pi}^T$ are attained in the global analysis for the first time, which are controlled by the $B_s^0 \rightarrow K^{*0}\bar{K}^{*0}$, $B^+ \rightarrow K^{*+}\phi$ and $B^0 \rightarrow K^{*0}\phi$ data. The fitted $B_s^0 \rightarrow K^{\pm}(K^{*\mp} \rightarrow)K\pi$ and $B_s^0 \rightarrow \bar{K}^0(\bar{K}^{*0} \rightarrow)K\pi$ branching ratios deviate from the central values of the data, in contrast with the other channels considered in the fit. This is not unexpected because of the involved large experimental errors, which do not give tight constraints actually. As a test, we remove these two modes, and collect the new outcomes in Table IV. The central values of the Gegenbauer moments change little relative to those in Table I, indicating that the measured $B_s^0 \rightarrow K^{\pm}(K^{*\mp} \rightarrow)K\pi$ and $B_s^0 \rightarrow \bar{K}^0(\bar{K}^{*0} \rightarrow)K\pi$ branching ratios indeed have minor impacts on the fit. Similarly, the obtained Gegenbauer moments for the $K\pi$ DAs vary away from those of the K^* meson DAs in QCD sum rules [60]. We point out that $a_{2K^*}^{\parallel} = 1.13 \pm 0.32$ in the previous fit [21], being greater than unity, is in fact not favored in view of the convergence of the Gegenbauer expansion. The refined result $a_{2K^*}^0 = -0.75 \pm 0.08$ in the present work is much smaller for two factors at least: the inclusions of the parameter $N_{K\pi}$ and of the data from the $B_{(s)} \rightarrow K^*\rho, K^*K^*, K^*\phi$ decays, among which the latter two dominated by the $B_{(s)} \rightarrow K^*$ transition can impose severe constraints on the $K\pi$ DAs.

The outcomes for $a_{2\phi}^0$ and $a_{2\phi}^T$ in the longitudinal and transverse twist-2 KK DAs ϕ_{KK}^0 and ϕ_{KK}^T , respectively, are controlled by the two pieces of $B \rightarrow K(\phi \rightarrow)KK$ data and the nine pieces of $B_{(s)} \rightarrow \phi K^*, \phi\phi$ data. Analogous to the $\pi\pi$ and $K\pi$ pairs, the above Gegenbauer moments differ from those of the ϕ meson DAs in QCD sum rules [60], and from $a_{2\phi}^0 = 0.40 \pm 0.06$ and $a_{2\phi}^T = 1.48 \pm 0.07$ in Ref. [26]. It is evident that the convergence of the Gegenbauer expansion for the $K\pi$ and KK DAs has been made more reliable by adding the extra parameters $N_{K\pi}$ and N_{KK} .

TABLE I: Fitted Gegenbauer moments for the twist-2 and twist-3 two-meson DAs.

	$a_{2\rho}^0$	$a_{2\rho}^s$	$a_{2\rho}^t$		
fit	0.16 ± 0.10	-0.11 ± 0.14	-0.21 ± 0.04		
	$a_{1K^*}^{\parallel}$	$a_{2K^*}^{\parallel}$	$a_{1K^*}^{\perp}$	$a_{2K^*}^{\perp}$	
fit	0.45 ± 0.11	-0.75 ± 0.08	0.61 ± 0.21	0.45 ± 0.06	
	$a_{2\phi}^0$	$a_{2\phi}^T$	$N_{\pi\pi}$	$N_{K\pi}$	N_{KK}
fit	-0.54 ± 0.14	0.77 ± 0.04	1.05 ± 0.04	1.48 ± 0.03	1.22 ± 0.03

TABLE II: CP averaged branching ratios (in units of 10^{-6}) of the three-body decays $B \rightarrow P_3(V \rightarrow P_1 P_2)$ in the PQCD approach. The data for comparison are quoted from [36]. Those data marked by “†” are included in the fit. The theoretical uncertainties are attributed to the variations of the shape parameter $\omega_{B(s)}$ in the $B_{(s)}$ meson DA, of the Gegenbauer moments in the two-meson DAs and of the hard scale t , which are then added in quadrature.

Channels	Results	Data
$B^+ \rightarrow K^+(\rho^0 \rightarrow)\pi\pi$	$2.90^{+1.51}_{-0.84}$	3.7 ± 0.5 †
$B^+ \rightarrow K^0(\rho^+ \rightarrow)\pi\pi$	$6.77^{+3.67}_{-2.30}$	$7.3^{+1.0}_{-1.2}$ †
$B^0 \rightarrow K^+(\rho^- \rightarrow)\pi\pi$	$8.31^{+4.35}_{-2.93}$	7.0 ± 0.9 †
$B^0 \rightarrow K^0(\rho^0 \rightarrow)\pi\pi$	$3.53^{+1.15}_{-0.92}$	3.4 ± 1.1 †
$B_s^0 \rightarrow K^-(\rho^+ \rightarrow)\pi\pi$	$16.9^{+4.9}_{-6.4}$...
$B_s^0 \rightarrow \bar{K}^0(\rho^0 \rightarrow)\pi\pi$	$0.18^{+0.03}_{-0.03}$...
$B^+ \rightarrow \pi^+(\rho^0 \rightarrow)\pi\pi$	$7.05^{+1.87}_{-1.42}$	8.3 ± 1.2 †
$B^+ \rightarrow \pi^0(\rho^+ \rightarrow)\pi\pi$	$10.07^{+4.23}_{-2.84}$	$10.6^{+1.2}_{-1.3}$ †
$B^0 \rightarrow \pi^0(\rho^0 \rightarrow)\pi\pi$	$0.04^{+0.02}_{-0.01}$	2.0 ± 0.5
$B^0 \rightarrow \pi^\pm(\rho^\mp \rightarrow)\pi\pi$	$30.49^{+9.33}_{-6.55}$	23.0 ± 2.3 †
$B_s^0 \rightarrow \pi^+(\rho^- \rightarrow)\pi\pi$	$0.17^{+0.06}_{-0.04}$...
$B_s^0 \rightarrow \pi^-(\rho^+ \rightarrow)\pi\pi$	$0.12^{+0.02}_{-0.02}$...
$B_s^0 \rightarrow \pi^0(\rho^0 \rightarrow)\pi\pi$	$0.16^{+0.05}_{-0.03}$...
$B^+ \rightarrow K^+(\bar{K}^{*0} \rightarrow)K\pi$	$0.59^{+0.23}_{-0.15}$	0.59 ± 0.08 †
$B^+ \rightarrow \bar{K}^0(K^{*+} \rightarrow)K\pi$	$0.21^{+0.09}_{-0.06}$...
$B^0 \rightarrow K^\pm(K^{*\mp} \rightarrow)K\pi$	$0.51^{+0.15}_{-0.11}$	< 0.4
$B^0 \rightarrow \bar{K}^0(\bar{K}^{*0} \rightarrow)K\pi$	$0.60^{+0.21}_{-0.15}$	< 0.96
$B_s^0 \rightarrow K^\pm(K^{*\mp} \rightarrow)K\pi$	$9.35^{+2.89}_{-2.01}$	19 ± 5 †
$B_s^0 \rightarrow \bar{K}^0(\bar{K}^{*0} \rightarrow)K\pi$	$9.31^{+3.14}_{-2.11}$	20 ± 6 †
$B^+ \rightarrow \pi^+(K^{*0} \rightarrow)K\pi$	$8.89^{+3.52}_{-2.54}$	10.1 ± 0.8 †
$B^+ \rightarrow \pi^0(K^{*+} \rightarrow)K\pi$	$5.84^{+2.12}_{-1.53}$	6.8 ± 0.9 †
$B^0 \rightarrow \pi^-(K^{*+} \rightarrow)K\pi$	$7.40^{+2.73}_{-1.92}$	7.5 ± 0.4 †
$B^0 \rightarrow \pi^0(K^{*0} \rightarrow)K\pi$	$2.80^{+1.09}_{-0.74}$	3.3 ± 0.6 †
$B_s^0 \rightarrow \pi^+(K^{*-} \rightarrow)K\pi$	$3.63^{+1.69}_{-1.27}$	2.9 ± 1.1
$B_s^0 \rightarrow \pi^0(\bar{K}^{*0} \rightarrow)K\pi$	$0.11^{+0.03}_{-0.02}$...
$B^+ \rightarrow K^+(\phi \rightarrow)KK$	$9.45^{+4.66}_{-3.27}$	$8.8^{+0.7}_{-0.6}$ †
$B^0 \rightarrow K^0(\phi \rightarrow)KK$	$8.63^{+4.25}_{-3.06}$	7.3 ± 0.7 †
$B_s^0 \rightarrow \bar{K}^0(\phi \rightarrow)KK$	$0.043^{+0.015}_{-0.014}$...
$B^+ \rightarrow \pi^+(\phi \rightarrow)KK$	$0.011^{+0.005}_{-0.004}$	0.032 ± 0.015
$B^0 \rightarrow \pi^0(\phi \rightarrow)KK$	$0.005^{+0.002}_{-0.002}$	< 0.015
$B_s^0 \rightarrow \pi^0(\phi \rightarrow)KK$	$0.11^{+0.04}_{-0.03}$...

C. Branching Ratios of Three-body B Meson Decays

With the fitted Gegenbauer moments in Table I, we compute the CP averaged branching ratios of all the considered three-body B meson decays in the LO PQCD formalism, and summarize the predictions in Table II. The listed theoretical uncertainties are assessed from three sources; the first one is associated with the shape parameter $\omega_B = 0.40$ GeV or $\omega_{B_s} = 0.48$ GeV with 10% variation; the second one originates from the Gegenbauer moments in the two-meson DAs; the last uncertainty is caused by tuning the hard scale t from $0.75t$ to $1.25t$ to characterize next-to-leading-order (NLO) QCD corrections. The individual uncertainties are then added in quadrature to get the total errors of the LO PQCD predictions in Table II. The errors due to the CKM matrix elements are tiny and can be safely ignored. The aforementioned three uncertainties are comparable in magnitude, and their combined effects could exceed 50%, implying that the nonperturbative parameters in the DAs of the initial and final states need to be known more precisely, and higher-order corrections to multi-body B meson decays are critical.

Most of the included $B \rightarrow P(\rho \rightarrow)\pi\pi$ data in Table II, in particular those with higher precision, are well reproduced. The central value of the fitted $B^0 \rightarrow \pi^\pm(\rho^\mp \rightarrow)\pi\pi$ branching ratio around 30×10^{-6} is a bit higher than the data. As illustrated in Ref. [59], NLO corrections can reduce the LO $B^0 \rightarrow \pi^\pm(\rho^\mp \rightarrow)\pi\pi$ branching ratio by about 40%. The disparity between the PQCD predictions and the data for three-body B meson decays can be diminished effectively by NLO contributions in principle,

TABLE III: CP averaged branching ratios \mathcal{B} (in units of 10^{-6}) and polarization fractions (in units of %) of the four-body decays $B \rightarrow V_1 V_2 \rightarrow (P_1 P_2)(P_3 P_4)$ in the PQCD approach, where the final states $(P_1 P_2)(P_3 P_4)$ are not written for simplicity. The data for comparison are quoted from [36]. Those data marked by “†” are included in the fit. The sources of theoretical errors are same as in Table II.

Channels	Results			Data		
	$\mathcal{B}(10^{-6})$	$f_0(\%)$	$f_\perp(\%)$	$\mathcal{B}(10^{-6})$	$f_0(\%)$	$f_\perp(\%)$
$B^+ \rightarrow \rho^+ \rho^0$	$12.7^{+4.5}_{-3.3}$	$97.7^{+0.8}_{-0.9}$	$1.3^{+0.5}_{-0.4}$	24 ± 1.9	95.0 ± 1.6	...
$B^0 \rightarrow \rho^+ \rho^-$	$27.0^{+10.7}_{-7.5}$	$92.2^{+3.6}_{-4.3}$	$4.5^{+2.3}_{-1.9}$	$27.7 \pm 1.9^\dagger$	$99.0^{+2.1}_{-1.9}^\dagger$...
$B^0 \rightarrow \rho^0 \rho^0$	$0.35^{+0.12}_{-0.07}$	$37.9^{+9.4}_{-3.2}$	$33.9^{+2.6}_{-5.7}$	0.96 ± 0.15	71^{+8}_{-9}	...
$B_s^0 \rightarrow \rho^+ \rho^-$	$1.35^{+0.84}_{-0.43}$	$99.4^{+0.6}_{-0.7}$	$0.3^{+0.1}_{-0.2}$
$B_s^0 \rightarrow \rho^0 \rho^0$	$0.68^{+0.42}_{-0.22}$	$99.4^{+0.6}_{-0.7}$	$0.3^{+0.1}_{-0.2}$	< 320
$B^+ \rightarrow \rho^0 K^{*+}$	$6.92^{+2.06}_{-3.21}$	$57.8^{+15.1}_{-12.8}$	$16.9^{+6.2}_{-6.9}$	$4.6 \pm 1.1^\dagger$	$78 \pm 12^\dagger$...
$B^+ \rightarrow \rho^+ K^{*0}$	$11.1^{+5.81}_{-3.82}$	$48.7^{+17.5}_{-13.4}$	$25.7^{+7.6}_{-8.5}$	$9.2 \pm 1.5^\dagger$	$48 \pm 8^\dagger$...
$B^0 \rightarrow \rho^- K^{*+}$	$9.91^{+4.84}_{-3.29}$	$48.0^{+16.5}_{-12.9}$	$26.2^{+6.3}_{-8.0}$	10.3 ± 2.6	38 ± 13	...
$B^0 \rightarrow \rho^0 K^{*0}$	$4.35^{+2.40}_{-1.59}$	$33.2^{+10.9}_{-9.9}$	$41.2^{+4.6}_{-7.8}$	3.9 ± 1.3	17.3 ± 2.6	40 ± 4
$B_s^0 \rightarrow \rho^0 \bar{K}^{*0}$	$0.35^{+0.11}_{-0.06}$	$59.4^{+9.4}_{-9.6}$	$21.5^{+4.9}_{-4.8}$	< 767
$B_s^0 \rightarrow \rho^+ K^{*-}$	$12.1^{+4.4}_{-3.9}$	$89.4^{+1.5}_{-2.3}$	$5.3^{+1.2}_{-0.9}$
$B^+ \rightarrow \rho^+ \phi$	$0.025^{+0.013}_{-0.009}$	$87.4^{+4.9}_{-7.2}$	$5.8^{+2.9}_{-1.9}$	< 3.0
$B^0 \rightarrow \rho^0 \phi$	$0.012^{+0.006}_{-0.004}$	$87.4^{+4.9}_{-7.2}$	$5.8^{+2.9}_{-1.9}$	< 0.33
$B_s^0 \rightarrow \rho^0 \phi$	$0.20^{+0.09}_{-0.06}$	$82.9^{+2.7}_{-1.9}$	$8.9^{+1.1}_{-1.4}$	0.27 ± 0.08
$B^0 \rightarrow \phi \phi$	$0.015^{+0.004}_{-0.003}$	$98.6^{+0.7}_{-2.0}$	$0.01^{+0.01}_{-0.00}$	< 0.027
$B_s^0 \rightarrow \phi \phi$	$16.6^{+6.6}_{-4.8}$	$38.7^{+10.6}_{-10.3}$	$30.9^{+5.1}_{-5.5}$	$18.5 \pm 1.4^\dagger$	$37.9 \pm 0.8^\dagger$	$31.0 \pm 0.6^\dagger$
$B^+ \rightarrow \phi K^{*+}$	$11.5^{+4.4}_{-3.9}$	$54.6^{+4.6}_{-9.1}$	$23.1^{+4.5}_{-2.2}$	$10 \pm 2^\dagger$	$55.0 \pm 5.0^\dagger$	$20.0 \pm 5.0^\dagger$
$B^0 \rightarrow \phi K^{*0}$	$10.4^{+4.6}_{-3.5}$	$51.9^{+4.3}_{-8.7}$	$24.5^{+4.2}_{-3.1}$	$10.00 \pm 0.50^\dagger$	$49.7 \pm 1.7^\dagger$	$22.4 \pm 1.5^\dagger$
$B_s^0 \rightarrow \phi \bar{K}^{*0}$	$0.29^{+0.17}_{-0.10}$	$62.3^{+11.9}_{-13.2}$	$25.2^{+8.8}_{-8.2}$	1.14 ± 0.30	51 ± 17	...
$B^+ \rightarrow K^{*+} \bar{K}^{*0}$	$0.71^{+0.34}_{-0.16}$	$83.5^{+5.0}_{-3.8}$	$8.5^{+1.5}_{-3.3}$	0.91 ± 0.29	82^{+15}_{-21}	...
$B^0 \rightarrow K^{*+} K^{*-}$	$1.24^{+0.38}_{-0.32}$	~ 100	~ 0	< 2.0
$B^0 \rightarrow K^{*0} \bar{K}^{*0}$	$0.60^{+0.22}_{-0.14}$	$81.1^{+3.2}_{-5.2}$	$9.6^{+2.7}_{-1.7}$	$0.83 \pm 0.24^\dagger$	$74 \pm 5^\dagger$...
$B_s^0 \rightarrow K^{*+} K^{*-}$	$13.7^{+5.4}_{-3.7}$	$32.3^{+10.5}_{-10.6}$	$33.9^{+5.3}_{-5.2}$
$B_s^0 \rightarrow K^{*0} \bar{K}^{*0}$	$13.2^{+5.2}_{-3.7}$	$28.2^{+8.8}_{-9.5}$	$35.7^{+4.7}_{-4.2}$	$11.1 \pm 2.7^\dagger$	$24 \pm 4^\dagger$	$38 \pm 12^\dagger$

TABLE IV: Fitted Gegenbauer moments for the twist-2 and twist-3 two-meson DAs without the $B_s^0 \rightarrow K^\pm(K^{*\mp} \rightarrow)K\pi$ and $B_s^0 \rightarrow \bar{K}^0(\bar{K}^{*0} \rightarrow)K\pi$ decays.

	$a_{2\rho}^0$	$a_{2\rho}^s$	$a_{2\rho}^t$		
fit	0.18 ± 0.10	-0.11 ± 0.14	-0.21 ± 0.04		
	$a_{1K^*}^\parallel$	$a_{2K^*}^\parallel$	$a_{1K^*}^\perp$	$a_{2K^*}^\perp$	
fit	0.43 ± 0.11	-0.76 ± 0.08	0.63 ± 0.21	0.44 ± 0.06	
	$a_{2\phi}^0$	$a_{2\phi}^T$	$N_{\pi\pi}$	$N_{K\pi}$	N_{KK}
fit	-0.57 ± 0.14	0.76 ± 0.04	1.05 ± 0.04	1.47 ± 0.03	1.22 ± 0.03

whose complete analysis in the PQCD formalism, however, goes beyond the scope of this work. The observables excluded in the fit are predicted at the LO accuracy, and compared with the data in Table II. The predicted branching ratio of the color-suppressed mode $B^0 \rightarrow \pi^0(\rho^0 \rightarrow)\pi\pi$, which suffers substantial subleading contributions, is still below the data, similar to what was obtained in the earlier publication on two-body decays [59].

Generally speaking, the distinction between the updated and previous results [21] is more notable for the $B_s \rightarrow P(K^* \rightarrow)K\pi$ branching ratios than for the $B \rightarrow P(K^* \rightarrow)K\pi$ ones. Taking the $B_s^0 \rightarrow \pi^+(K^{*-} \rightarrow)K\pi$ and $B^0 \rightarrow \pi^-(K^{*+} \rightarrow)K\pi$ decays as examples, we find that the prediction $\mathcal{B}(B_s^0 \rightarrow \pi^+(K^{*-} \rightarrow)K\pi) = (3.63^{+1.69}_{-1.27}) \times 10^{-6}$ is smaller than $(12.13^{+4.94}_{-3.85}) \times 10^{-6}$ in [21] by a factor of ~ 4 , and becomes closer to the data $(2.9 \pm 1.1) \times 10^{-6}$. For the $B^0 \rightarrow \pi^-(K^{*+} \rightarrow)K\pi$ decay, our branching ratio reveals no deviation from that in [21]. The above observations are understood easily; the former involves the $B_s \rightarrow (K^* \rightarrow)K\pi$ transition form factor, which is more sensitive to the variation of the Gegenbauer moments in the $K\pi$ DAs.




FIG. 1: Dependencies of $\mathcal{B}(B_s^0 \rightarrow K^\pm(K^{*\mp} \rightarrow)K\pi)$ on the Gegenbauer moments $a_{1K^*}^s$ and $a_{1K^*}^t$.

As stated before, the predicted branching ratios $\mathcal{B}(B_s^0 \rightarrow K^\pm(K^{*\mp} \rightarrow)K\pi)$ and $\mathcal{B}(B_s^0 \rightarrow \bar{K}^0(\bar{K}^{*0} \rightarrow)K\pi)$ are somewhat lower than the data, but consistent with the previous evaluations based on two-body decays [61]. The authors of Ref. [61] have noticed that the $B_s^0 \rightarrow K^\pm K^{*\mp}$ and $B_s^0 \rightarrow \bar{K}^0 \bar{K}^{*0}$ branching ratios can be enhanced significantly by NLO corrections, which reach $\sim 40\%$ in magnitude of the LO contributions. We have stuck to the asymptotic forms of the twist-3 $K\pi$ DAs $\phi_{K\pi}^s$ and $\phi_{K\pi}^t$ owing to the finite amount of data in the current study. Here we explore the influence on the above branching ratios from the subleading Gegenbauer moments in $\phi_{K\pi}^s$ and $\phi_{K\pi}^t$, given by

$$\phi_{K\pi}^s(z, \omega^2) = \frac{3F_{K\pi}^\perp(\omega^2)}{2\sqrt{2N_c}} [t(1 + a_{1K^*}^s t) - a_{1K^*}^s 2z(1 - z)], \quad (57)$$

$$\phi_{K\pi}^t(z, \omega^2) = \frac{3F_{K\pi}^\perp(\omega^2)}{2\sqrt{2N_c}} t[t + a_{1K^*}^t (3t^2 - 1)]. \quad (58)$$

The dependencies of $\mathcal{B}(B_s^0 \rightarrow K^\pm(K^{*\mp} \rightarrow)K\pi)$ on the Gegenbauer moments $a_{1K^*}^s$ and $a_{1K^*}^t$ are plotted in Fig. 1, which imply the sensitivity to $a_{1K^*}^s$ and $a_{1K^*}^t$ clearly. It suggests that more precise data of $\mathcal{B}(B_s^0 \rightarrow K^\pm(K^{*\mp} \rightarrow)K\pi)$ and $\mathcal{B}(B_s^0 \rightarrow \bar{K}^0(\bar{K}^{*0} \rightarrow)K\pi)$ can yield better grasp of the twist-3 $K\pi$ DAs.

With a single Gegenbauer moment $a_{2\phi}^0$ in the twist-2 KK DA ϕ_{KK}^0 , the two pieces of $B \rightarrow K(\phi \rightarrow)KK$ data included in the fit are well reproduced as indicated in Table II. The predicted $B^+ \rightarrow \pi^+(\phi \rightarrow)KK$ branching ratio shows minor deviation from, but still agrees with the data within uncertainties.

TABLE V: PQCD predictions for the $B_{(s)} \rightarrow P_1 P_2$ transition form factors $F_\perp(m_R^2, 0)$ with $R = \rho, K^*, \phi$, whose sources of theoretical errors are same as in Table II. The results from light-cone sum rules [62–68] are listed for comparison.

Decay modes	This work	LCSR [62]	LCSR [63]	LCSR [64]	LCSR [65]	LCSR [66]	LCSR [67]	LCSR [68]
$B \rightarrow \pi\pi$	39_{-10}^{+9}	29 ± 15	34 ± 11	35 ± 3	43 ± 12	34 ± 4
$B \rightarrow K\pi$	113_{-16}^{+17}	79 ± 26	93 ± 26	81 ± 10	...	98 ± 11	86 ± 43	62 ± 36
$B_s \rightarrow K\pi$	62_{-10}^{+14}	72 ± 7	...	75 ± 9
$B_s \rightarrow KK$	905_{-132}^{+128}	1080 ± 92	...	1121 ± 136

D. $B \rightarrow P_1 P_2$ Transition Form Factors

We compute the $B_{(s)} \rightarrow P_1 P_2$ transition form factors, which constitute an essential ingredient in the formulation of the semileptonic decays $B_{(s)} \rightarrow P_1 P_2 l^+ l^-$, using the fitted Gegenbauer moments. The exercise is motivated by the universality

of the two-meson DAs in the PQCD factorization. There are three (seven) form factors involved in the (S -wave) P -wave $B_{(s)} \rightarrow P_1 P_2$ transition, whose definitions can be found in Refs. [65, 68]. For simplicity, we focus only on the form factor $F_{\perp}(\omega^2, Q^2)$ defined via the vector current $\bar{q}\gamma^{\mu}b$, $q = u, d, s$, with the invariant mass ω of an intermediate resonance R and the momentum transfer Q ,

$$i\langle P_1(p_1)P_2(p_2)|\bar{q}\gamma^{\mu}b|B(p_B)\rangle = F_{\perp}(\omega^2, Q^2)k_{\perp}^{\mu}, \quad (59)$$

where the vector k_{\perp}^{μ} has been specified in Ref. [68]. The PQCD factorization formula for $F_{\perp}(\omega^2, Q^2)$ is referred to Eq. (A.59) in Ref. [23]. The predictions for $F_{\perp}(m_R^2, 0)$ with $R = \rho, K^*, \phi$ at maximal recoil are presented in Table V, and compared with those from light-cone sum rules (LCSR) in the narrow-width limit. The consistency between the two approaches is satisfactory within theoretical uncertainties, hinting that the PQCD formalism and the predictions from it are reasonable. A complete and systematic PQCD investigation of all the $B_{(s)} \rightarrow P_1 P_2$ form factors, including their ω^2 and Q^2 dependencies, will be postponed to a future work.

The $B, B_s \rightarrow K\pi$ transition form factors $F_{\perp}(m_{K^*}^2, 0)$ have been calculated in the PQCD approach [23]. We find that the updated $F_{\perp}(m_{K^*}^2, 0) = 113$ for $B^0 \rightarrow K\pi$ is larger than the previous one 88, while the updated $F_{\perp}(m_{K^*}^2, 0) = 62$ is very close to the previous one 64 for $B_s^0 \rightarrow K\pi$. It is easy to see that the destruction between the contributions from the Gegenbauer polynomials $C_1^{3/2}(t)$ and $C_2^{3/2}(t)$ in the transverse twist-2 $K\pi$ DA $\Phi_{K\pi}^T$ weakens the dependence of $F_{\perp}(m_{K^*}^2, 0)$ on the moments $a_{1K^*}^{\perp}$ and $a_{2K^*}^{\perp}$ in the latter case. For the $B^0 \rightarrow K\pi$ transition, the value of $a_{1K^*}^{\perp}$, related to $SU(3)$ symmetry breaking effects, flips sign, since the momentum fraction is associated with the \bar{s} quark in the K^{*0} meson. The contributions from the aforementioned $C_1^{3/2}(t)$ and $C_2^{3/2}(t)$ then become constructive, and the associated $F_{\perp}(m_{K^*}^2, 0)$ is more sensitive to the Gegenbauer moments.

E. Branching Ratios and Polarization Fractions of $B \rightarrow V_1 V_2$ Decays

Most of the predicted branching ratios of the four-body decays $B \rightarrow V_1 V_2 \rightarrow (P_1 P_2)(P_3 P_4)$ match the data within sizable theoretical errors as evinced in Table III. Our predictions $\mathcal{B}(B^+ \rightarrow \rho^0 \rho^+) = (12.7_{-3.3}^{+4.5}) \times 10^{-6}$ and $\mathcal{B}(B^0 \rightarrow \rho^0 \rho^0) = (0.35_{-0.07}^{+0.12}) \times 10^{-6}$ are slightly below those from other theoretical approaches [69–71] and the data [36], but close to the earlier PQCD results from the two-body-decay framework [72]. It has been observed that strong cancellations occurs to the nonfactorizable contributions in the color-suppressed mode $B^0 \rightarrow \rho^0 \rho^0$ at LO, and its branching ratio turns out to be as tiny as $\mathcal{B}(B^0 \rightarrow \rho^0 \rho^0) = (0.35_{-0.06}^{+0.10}) \times 10^{-6}$. Then the relation of $\mathcal{B}(B^0 \rightarrow \rho^+ \rho^-) \approx 2\mathcal{B}(B^+ \rightarrow \rho^+ \rho^0)$ always holds in the PQCD analysis, because of the small $\mathcal{B}(B^0 \rightarrow \rho^0 \rho^0) \sim 10^{-7}$ in the isospin triangle. On the experimental side, nevertheless, the two rates are roughly equal to each other within errors, leading to a puzzle that has persisted for a long time.

It is naively expected that the longitudinal components dominate the charmless $B \rightarrow V_1 V_2$ decays based on the counting rules for the polarization fractions [73],

$$f_0 \sim 1 - \mathcal{O}(m_V^2/m_B^2), \quad f_{\parallel} \sim f_{\perp} \sim \mathcal{O}(m_V^2/m_B^2), \quad (60)$$

with a vector meson mass m_V . The penguin-dominated modes, such as $B \rightarrow K^* \rho, B \rightarrow K^* \phi$ and $B_s^0 \rightarrow \phi \phi$ with the observed longitudinal polarization fractions being around 50% [51, 74–78], do not respect the above counting rules. To interpret the large transverse polarizations, a number of resolutions within or beyond the SM have been proposed [79–102]. In the PQCD approach, the sizable transverse contributions are accommodated by means of the hard emission diagrams and the chirally enhanced annihilation diagrams, especially the $(S - P)(S + P)$ penguin annihilation from the QCD penguin operator O_6 [81, 87].

The longitudinal polarization fraction for the color-suppressed $B^0 \rightarrow \rho^0 \rho^0$ decay is predicted to be $f_0 = (37.9_{-3.2}^{+9.4})\%$, which is much lower than the data $(71_{-9}^{+8})\%$ [36]. It has been found that the longitudinal polarization contributions from the hard emission diagrams largely cancel each other in this type of decays. The minor penguin annihilation contributions then play a role in enhancing the transverse components, such that the transverse polarization fraction f_{\perp} becomes comparable with the longitudinal one. We point out that NLO corrections to $f_0(B^0 \rightarrow \rho^0 \rho^0)$ can shrink the gap between the PQCD prediction and the data [103], which will be addressed in future studies. It is worth mentioning that the predicted longitudinal polarization fraction of the pure-penguin decay $f_0(B_s^0 \rightarrow K^{*0} \bar{K}^{*0}) = (28.2_{-9.5}^{+8.8})\%$, far below our previous result [23], is now in good agreement with the measured one $f_0(B_s^0 \rightarrow K^{*0} \bar{K}^{*0})_{\text{exp}} = (24 \pm 4)\%$ [36]. The improved consistency is mainly attributed to the updated Gegenbauer moments $a_{1K^*}^{\parallel(\perp)}$ and $a_{2K^*}^{\parallel(\perp)}$. To be specific, we display $f_0(B_s^0 \rightarrow K^{*0} \bar{K}^{*0})$ as a function of the above moments in Fig. 2, which decreases quickly as $a_{2K^*}^{\parallel}$ turns negative and as the other three increase. It is then not hard to realize the drop of $f_0(B_s^0 \rightarrow K^{*0} \bar{K}^{*0})$ by comparing $a_{1K^*}^{\parallel} = a_{1K^*}^{\perp} = 0.31 \pm 0.16$ and $a_{2K^*}^{\parallel} = a_{2K^*}^{\perp} = 1.188 \pm 0.099$ employed in [23] with the corresponding ones in Table I.

It is well known that flavor-changing neutral-current transitions are very sensitive to new physics (NP) contributions. Though the polarization anomaly associated with the $B_s^0 \rightarrow K^{*0} \bar{K}^{*0}$ decay is resolved here, we cannot rule out the possibility of NP




FIG. 2: Longitudinal polarization fraction f_0 in the $B_s^0 \rightarrow K^{*0} \bar{K}^{*0}$ decay as a function of the Gegenbauer moments $a_{1K^*}^{||(\perp)}$ and $a_{2K^*}^{||(\perp)}$. The blue dashed line corresponds to the central value of the data [36] with the shaded band representing the experimental errors.

effects. A new observable $L_{K^{*0} \bar{K}^{*0}}$ in terms of two U -spin related decays $B_s^0 \rightarrow K^{*0} \bar{K}^{*0}$ and $B^0 \rightarrow K^{*0} \bar{K}^{*0}$ has been constructed to maximize the sensitivity to NP by reducing hadronic uncertainties [32],

$$L_{K^{*0} \bar{K}^{*0}} = \frac{\mathcal{B}(B_s^0 \rightarrow K^{*0} \bar{K}^{*0})g(B^0 \rightarrow K^{*0} \bar{K}^{*0})f_L(B_s^0 \rightarrow K^{*0} \bar{K}^{*0})}{\mathcal{B}(B^0 \rightarrow K^{*0} \bar{K}^{*0})g(B_s^0 \rightarrow K^{*0} \bar{K}^{*0})f_L(B^0 \rightarrow K^{*0} \bar{K}^{*0})},$$

with the phase space factors

$$g(B_{(s)} \rightarrow K^{*0} \bar{K}^{*0}) = \frac{\tau_{B_{(s)}}}{16\pi M_{B_{(s)}}^2} \sqrt{M_{B_{(s)}}^2 - 4M_{K^{*0}}^2}. \quad (61)$$

Adopting the latest inputs from the PDG [36], one derives the experimental measurement [32]

$$L_{K^{*0} \bar{K}^{*0}}^{\text{exp}} = 4.43 \pm 0.92, \quad (62)$$

smaller than the QCDF [32] and PQCD [104] predictions

$$L_{K^{*0} \bar{K}^{*0}} = \begin{cases} 19.5_{-6.8}^{+9.3} & (\text{QCDF}), \\ 12.7_{-3.2}^{+5.6} & (\text{PQCD}). \end{cases} \quad (63)$$

The difference between Eqs. (62) and (63), being regarded as a NP signal [32], motivated the authors of Ref. [104] to consider a family non-universal Z' boson in the $b \rightarrow sq\bar{q}$ transition. We update the ratio $L_{K^{*0} \bar{K}^{*0}}$ based on Table III,

$$L_{K^{*0} \bar{K}^{*0}}^{\text{PQCD}} = 7.7_{-3.8}^{+4.9}, \quad (64)$$

which turns closer to Eq. (62). Namely, the deviation of our prediction from the data can be alleviated by improving the precision of nonperturbative parameters. We encourage experimentalists to conduct direct measurements on $L_{K^{*0} \bar{K}^{*0}}$ to uncover potential NP signals.

F. Direct CP Asymmetries of Multi-body B Decays

The direct CP asymmetries from the integration over the phase space of the considered three-body B meson decays are listed in Table VI. Those modes induced only by penguin operators are expected to have vanishing CP asymmetries in the LO PQCD framework and not shown in the table. The predicted direct CP asymmetry of the $B^+ \rightarrow \pi^+ \rho^0$ decay is found to be negative and large, but the measured value $\mathcal{A}_{CP}(B^+ \rightarrow \pi^+ \rho^0) = (-0.4 \pm 1.7 \pm 0.9)\%$ is consistent with zero [105]. This mismatch

TABLE VI: Direct CP asymmetries (in units of %) of the three-body decays $B \rightarrow VP_3 \rightarrow P_1P_2P_3$. The data for comparison are quoted from [36]. The sources of theoretical errors are same as in Table II.

Modes	Results	Data
$B^+ \rightarrow K^+(\rho^0 \rightarrow)\pi\pi$	$62.4^{+11.6}_{-13.5}$	16 ± 2
$B^+ \rightarrow K^0(\rho^+ \rightarrow)\pi\pi$	$9.3^{+1.2}_{-2.2}$	-3 ± 15
$B^0 \rightarrow K^+(\rho^- \rightarrow)\pi\pi$	$44.1^{+7.2}_{-7.3}$	20 ± 11
$B^0 \rightarrow K^0(\rho^0 \rightarrow)\pi\pi$	$1.5^{+2.4}_{-2.2}$	-4 ± 20
$B_s^0 \rightarrow K^-(\rho^+ \rightarrow)\pi\pi$	$19.6^{+4.4}_{-5.2}$...
$B_s^0 \rightarrow \bar{K}^0(\rho^0 \rightarrow)\pi\pi$	$-36.4^{+26.8}_{-17.9}$...
$B^+ \rightarrow \pi^+(\rho^0 \rightarrow)\pi\pi$	$-34.1^{+7.3}_{-9.8}$	0.9 ± 1.9
$B^+ \rightarrow \pi^0(\rho^+ \rightarrow)\pi\pi$	$23.1^{+7.0}_{-6.1}$	2 ± 11
$B^0 \rightarrow \pi^+(\rho^- \rightarrow)\pi\pi$	$-26.5^{+5.4}_{-5.7}$	-8 ± 8
$B^0 \rightarrow \pi^-(\rho^+ \rightarrow)\pi\pi$	$9.5^{+2.3}_{-2.6}$	13 ± 6
$B^0 \rightarrow \pi^0(\rho^0 \rightarrow)\pi\pi$	$16.7^{+23.5}_{-18.9}$	27 ± 24
$B_s^0 \rightarrow \pi^+(\rho^- \rightarrow)\pi\pi$	$-37.9^{+19.1}_{-13.1}$...
$B_s^0 \rightarrow \pi^-(\rho^+ \rightarrow)\pi\pi$	$-66.2^{+7.0}_{-4.5}$...
$B_s^0 \rightarrow \pi^0(\rho^0 \rightarrow)\pi\pi$	$-40.0^{+8.8}_{-4.4}$...
$B^+ \rightarrow K^+(\bar{K}^{*0} \rightarrow)K\pi$	$-4.0^{+12.2}_{-10.3}$	4 ± 5
$B^+ \rightarrow \bar{K}^0(K^{*+} \rightarrow)K\pi$	$-61.9^{+4.2}_{-19.2}$...
$B^0 \rightarrow K^+(K^{*-} \rightarrow)K\pi$	$18.4^{+8.9}_{-6.3}$...
$B^0 \rightarrow K^-(K^{*+} \rightarrow)K\pi$	$14.3^{+4.6}_{-9.9}$...
$B_s^0 \rightarrow K^+(K^{*-} \rightarrow)K\pi$	$11.5^{+7.2}_{-8.8}$...
$B_s^0 \rightarrow K^-(K^{*+} \rightarrow)K\pi$	$-7.3^{+8.0}_{-7.9}$...
$B^+ \rightarrow \pi^+(K^{*0} \rightarrow)K\pi$	$-2.3^{+1.2}_{-0.5}$	-4 ± 9
$B^+ \rightarrow \pi^0(K^{*+} \rightarrow)K\pi$	$-3.5^{+4.0}_{-3.9}$	-39 ± 21
$B^0 \rightarrow \pi^-(K^{*+} \rightarrow)K\pi$	$-15.6^{+5.9}_{-6.4}$	-27 ± 4
$B^0 \rightarrow \pi^0(K^{*0} \rightarrow)K\pi$	$-13.5^{+1.6}_{-1.8}$	-15 ± 13
$B_s^0 \rightarrow \pi^+(K^{*-} \rightarrow)K\pi$	$-18.1^{+5.7}_{-5.9}$...
$B_s^0 \rightarrow \pi^0(\bar{K}^{*0} \rightarrow)K\pi$	$-22.3^{+15.7}_{-17.8}$...
$B^+ \rightarrow K^+(\phi \rightarrow)KK$	$0.7^{+1.2}_{-1.8}$	2.4 ± 2.8
$B_s^0 \rightarrow \pi^0(\phi \rightarrow)KK$	$31.2^{+3.7}_{-3.7}$...

represents a puzzle that needs to be resolved. The CPV originating from the interference between the S - and P -wave $\pi^+\pi^-$ pairs in the $B^+ \rightarrow \pi^+\rho^0$ decay has been observed by the LHCb [6]. The interference causes significant local asymmetries, which, however, cancel in the integration over the phase space of the decay. It has been postulated [56, 106] that the nearly diminishing CPV is due to the destruction between the power corrections from the penguin annihilation and hard spectator interactions.

The LHCb Collaboration reported the first observation of the direct CP asymmetry $\mathcal{A}_{CP}(B^+ \rightarrow K^+\rho^0) = (15.0 \pm 1.9 \pm 1.1)\%$ [105] recently based on the measurements of the three-body decay $B^\pm \rightarrow K^\pm\pi^+\pi^-$. Our prediction $\mathcal{A}_{CP}(B^+ \rightarrow K^+\rho^0) = (62.4^{+11.6}_{-13.5})\%$, similar to the one from the formalism for two-body decays [107], is higher than the data. The authors of Ref. [108] noticed that NLO corrections to the scalar pion form factor are small in magnitude but generate a large strong phase, which can affect dramatically direct CP asymmetries. The discrepancy between our prediction and the data for $\mathcal{A}_{CP}(B^+ \rightarrow K^+\rho^0)$ may thus be removed, when complete NLO corrections to multi-body B decays, in particular those to nonfactorizable spectator and annihilation diagrams, are available.

CP asymmetries of the $B \rightarrow V_1V_2$ modes have not been measured at a level higher than 5σ so far, most of which exhibit no apparent deviation from zero. There are three possible spin configurations corresponding to the polarizations of the final-state vector mesons V_1 and V_2 , which are allowed by the angular momentum conservation, while there is only one helicity amplitude in B meson decays into two pseudoscalars. We will demonstrate that CP asymmetries in individual helicity final states of some $B \rightarrow V_1V_2$ decays can be as large as those in the $B \rightarrow P_1P_2$ ones, greater than 10%, but the cancellation between different polarization components results in small net CPV.

For the four-body decays $B \rightarrow V_1V_2 \rightarrow (P_1P_2)(P_3P_4)$, one can disentangle the helicities of the $V_1(\rightarrow P_1P_2)V_2(\rightarrow P_3P_4)$ final states via an angular analysis. We collect in Table VII the direct CP asymmetries in the three helicity states $\mathcal{A}_{CP}^{0,||,\perp}$ together

TABLE VII: Direct CP asymmetries (in units of %) of the four-body decays $B \rightarrow V_1 V_2 \rightarrow (P_1 P_2)(P_3 P_4)$ in the PQCD approach. The data for comparison are quoted from [36]. The sources of theoretical errors are same as in Table II. The percentage in the parentheses specifies the proportion of CPV of each helicity state to the direct CPV.

Modes	\mathcal{A}_{CP}^0	$\mathcal{A}_{CP}^{\parallel}$	\mathcal{A}_{CP}^{\perp}	$\mathcal{A}_{CP}^{\text{dir}}$
$B^+ \rightarrow \rho^+ \rho^0 \rightarrow (\pi^+ \pi^0)(\pi^+ \pi^-)$	$0.4_{-0.1}^{+0.2}$ (97.7%)	$-0.1_{-0.6}^{+0.4}$ (1.0%)	$0.5_{-0.6}^{+0.2}$ (1.3%)	$0.4_{-0.1}^{+0.2}$
Data	-5 ± 5
$B^0 \rightarrow \rho^+ \rho^- \rightarrow (\pi^+ \pi^0)(\pi^- \pi^0)$	$-3.7_{-2.0}^{+0.9}$ (92.2%)	$43.4_{-17.7}^{+11.7}$ (3.3%)	$38.4_{-16.9}^{+12.0}$ (4.5%)	$-0.3_{-2.4}^{+2.9}$
Data	0 ± 9
$B^0 \rightarrow \rho^0 \rho^0 \rightarrow (\pi^+ \pi^-)(\pi^+ \pi^-)$	$61.7_{-14.8}^{+12.3}$ (37.9%)	$64.9_{-8.4}^{+6.1}$ (28.2%)	$76.9_{-9.0}^{+8.8}$ (33.9%)	$67.8_{-9.9}^{+7.1}$
Data	20 ± 90
$B_s^0 \rightarrow \rho^+ \rho^- \rightarrow (\pi^+ \pi^0)(\pi^- \pi^0)$	$5.1_{-2.5}^{+4.3}$ (99.4%)	$4.3_{-3.9}^{+4.6}$ (0.3%)	$5.9_{-5.3}^{+8.6}$ (0.3%)	$5.1_{-2.6}^{+4.2}$
$B_s^0 \rightarrow \rho^0 \rho^0 \rightarrow (\pi^+ \pi^-)(\pi^+ \pi^-)$	$5.1_{-2.5}^{+4.3}$ (99.4%)	$4.3_{-3.9}^{+4.6}$ (0.3%)	$5.9_{-5.3}^{+8.6}$ (0.3%)	$5.1_{-2.6}^{+4.2}$
$B^+ \rightarrow \rho^+ K^{*0} \rightarrow (\pi^+ \pi^0)(K^+ \pi^-)$	$-0.2_{-3.4}^{+1.9}$ (48.7%)	$1.6_{-1.3}^{+0.9}$ (25.6%)	$1.7_{-0.6}^{+0.2}$ (25.7%)	$0.8_{-1.8}^{+0.8}$
Data	-1 ± 16
$B^+ \rightarrow \rho^0 K^{*+} \rightarrow (\pi^+ \pi^-)(K^0 \pi^+)$	$32.8_{-8.5}^{+2.7}$ (57.8%)	$0.8_{-4.6}^{+4.4}$ (25.3%)	$-56.2_{-9.4}^{+9.7}$ (16.9%)	$9.6_{-7.7}^{+7.1}$
Data	31 ± 13
$B^0 \rightarrow \rho^0 K^{*0} \rightarrow (\pi^+ \pi^-)(K^+ \pi^-)$	$0.1_{-7.3}^{+5.3}$ (33.2%)	$-34.5_{-12.2}^{+9.2}$ (25.6%)	$12.8_{-1.4}^{+0.8}$ (41.2%)	$-3.5_{-2.7}^{+2.2}$
Data	-6 ± 9
$B^0 \rightarrow \rho^- K^{*+} \rightarrow (\pi^- \pi^0)(K^0 \pi^+)$	$57.2_{-10.1}^{+5.8}$ (48.0%)	$-26.5_{-4.7}^{+5.3}$ (25.8%)	$-30.7_{-4.9}^{+4.6}$ (26.2%)	$12.6_{-9.5}^{+12.0}$
Data	21 ± 15
$B_s^0 \rightarrow \rho^+ K^{*-} \rightarrow (\pi^+ \pi^0)(\bar{K}^0 \pi^-)$	$-14.2_{-3.8}^{+3.4}$ (89.4%)	$73.7_{-15.4}^{+12.5}$ (5.3%)	$75.2_{-14.1}^{+11.5}$ (5.3%)	$-4.7_{-2.9}^{+3.1}$
$B_s^0 \rightarrow \rho^0 \bar{K}^{*0} \rightarrow (\pi^+ \pi^-)(K^- \pi^+)$	$21.2_{-16.5}^{+19.6}$ (59.4%)	$73.2_{-25.0}^{+15.4}$ (19.1%)	$81.8_{-21.8}^{+11.6}$ (21.5%)	$44.2_{-13.9}^{+11.5}$
$B_s^0 \rightarrow \rho^0 \phi \rightarrow (\pi^+ \pi^-)(K^+ K^-)$	$-2.4_{-6.9}^{+9.0}$ (82.9%)	$-18.3_{-4.2}^{+5.2}$ (8.2%)	$-16.8_{-2.9}^{+4.0}$ (8.9%)	$-5.0_{-5.8}^{+8.2}$
$B^+ \rightarrow K^{*+} \phi \rightarrow (K^0 \pi^+)(K^+ K^-)$	$-5.7_{-3.7}^{+11.2}$ (54.6%)	$2.8_{-2.9}^{+0.3}$ (22.3%)	$-2.3_{-1.0}^{+1.3}$ (23.1%)	$-3.3_{-1.6}^{+7.6}$
$B^+ \rightarrow K^{*+} \bar{K}^{*0} \rightarrow (K^0 \pi^+)(K^+ \pi^-)$	$-21.5_{-10.8}^{+8.7}$ (83.5%)	$-9.0_{-1.3}^{+1.4}$ (8.0%)	$7.9_{-1.5}^{+1.7}$ (8.5%)	$-19.4_{-9.2}^{+7.6}$
$B^0 \rightarrow K^{*+} K^{*-} \rightarrow (K^0 \pi^+)(\bar{K}^0 \pi^-)$	$19.1_{-1.5}^{+2.8}$ ($\sim 100\%$)	$-29.5_{-16.0}^{+17.3}$ (~ 0)	$10.2_{-5.1}^{+6.7}$ (~ 0)	$19.1_{-1.5}^{+2.8}$
$B_s^0 \rightarrow K^{*+} K^{*-} \rightarrow (K^0 \pi^+)(\bar{K}^0 \pi^-)$	$33.1_{-8.2}^{+7.5}$ (32.3%)	$-17.4_{-5.4}^{+4.5}$ (33.8%)	$-16.6_{-5.2}^{+4.6}$ (33.9%)	$-0.9_{-2.4}^{+1.6}$

with $\mathcal{A}_{CP}^{\text{dir}}$ from the summation over all helicity states. The total direct CP asymmetry can be well approximated by the weighted sum of the three asymmetries,

$$\mathcal{A}_{CP}^{\text{dir}} \approx f_0 \mathcal{A}_{CP}^0 + f_{\parallel} \mathcal{A}_{CP}^{\parallel} + f_{\perp} \mathcal{A}_{CP}^{\perp}, \quad (65)$$

with the coefficients f_i , $i = 0, \parallel, \perp$, being the corresponding polarization fractions. The predicted $\mathcal{A}_{CP}^{\parallel}$ for the $B^0 \rightarrow \rho^0 K^{*0} \rightarrow (\pi^+ \pi^-)(K^+ \pi^-)$ channel can reach $(-34.5_{-12.2}^{+9.2})\%$, but the cancellation among the three helicity components in Eq. (65) decreases the magnitude of $\mathcal{A}_{CP}^{\text{dir}}$, and render it compatible with the data. We mention that similar cancellation between various partial waves also appears in the analysis of the CPV in bottom baryon decays [109]. The tiny asymmetry \mathcal{A}_{CP}^0 in the longitudinal component of the $B_s^0 \rightarrow \rho^0 \phi \rightarrow (\pi^+ \pi^-)(K^+ K^-)$ decay governs the direct CPV owing to the weight of nearly unity, $f_0 \sim 80\%$, giving rise to small $\mathcal{A}_{CP}^{\text{dir}}$. The direct CP asymmetries of the other decays can be understood in the same manner.

IV. CONCLUSION

We have improved the PQCD formalism of multi-body charmless hadronic B meson decays by resolving the possible inconsistency in the parametrization for P -wave resonances in two-meson DAs. The determination of the Gegenbauer moments in two-meson DAs was then updated by performing the global fit to measured branching ratios and polarization fractions for the three-body $B \rightarrow VP_3 \rightarrow P_1 P_2 P_3$ and four-body $B \rightarrow V_1 V_2 \rightarrow (P_1 P_2)(P_3 P_4)$ decays in the LO PQCD framework. The data from the three additional four-body decays $B \rightarrow \rho K^* \rightarrow (\pi\pi)(K\pi)$, $B \rightarrow K^* K^* \rightarrow (K\pi)(K\pi)$ and $B \rightarrow \phi K^* \rightarrow (KK)(K\pi)$ were included to reduce theoretical uncertainties. The moments $a_{1K^*}^{\perp}$ and $a_{2K^*}^{\perp}$ in the transverse $K\pi$ DA were extracted from the global analysis for the first time. It has been shown that the resultant Gegenbauer expansion of two-meson DAs becomes more convergent.

With the updated two-meson DAs, the branching ratios, polarization fractions and the direct CP asymmetries of the considered multi-body B meson decays were reexamined. It has been demonstrated that most of the data involved in the fit are well

reproduced; namely, the fit quality is satisfactory. It was elaborated that the precision of the two-meson DAs does play a crucial role in accounting for the data, especially for the unexpected low longitudinal polarization fraction of the $B_s^0 \rightarrow K^{*0} \bar{K}^{*0}$ decay. We have predicted the observable $L_{K^*K^*}$ in the PQCD approach, whose confrontation with measurements can probe the presence of NP contributions. It was also found that although the direct CP asymmetry in each polarization component of some four-body B meson decays might be sizable, the strong cancellation among polarization states leads to diminishing net CPV. Therefore, it is worthwhile to search for CPV in individual polarization states experimentally.

The present work is still based on the LO PQCD factorization formulas. The precision of the two-meson DAs can be further enhanced systematically, when higher-order and/or higher-power corrections to multi-body hadronic B -meson decays are taken into account. More precise measurements, in particular those of CP asymmetries, are also necessary. If a high-precision global investigation discloses notable tensions between theoretical results and data, it may hint that NP effects are inevitable.

Acknowledgments

We thank Fu-Sheng Yu, Rui-Lin Zhu and Jia-Jie Han for valuable discussions. This work is supported by the National Natural Science Foundation of China under Grant Nos. 12005103, 12075086, 12105028 and by NSTC of R.O.C. under Grant No. NSTC-113-2112-M-001-024-MY3. ZR is supported in part by the Natural Science Foundation of Hebei Province under Grant Nos. A2019209449 and A2021209002.

-
- [1] R. H. Dalitz, *Phil. Mag. Ser. 7* **44** (1953), 1068-1080. doi:10.1080/14786441008520365
 - [2] R. H. Dalitz, *Phys. Rev.* **94** (1954), 1046-1051 doi:10.1103/PhysRev.94.1046
 - [3] R. Aaij *et al.* [LHCb], *Phys. Rev. Lett.* **111** (2013), 101801 doi:10.1103/PhysRevLett.111.101801 [arXiv:1306.1246 [hep-ex]].
 - [4] R. Aaij *et al.* [LHCb], *Phys. Rev. Lett.* **112** (2014) no.1, 011801 doi:10.1103/PhysRevLett.112.011801 [arXiv:1310.4740 [hep-ex]].
 - [5] R. Aaij *et al.* [LHCb], *Phys. Rev. D* **108** (2023) no.1, 012008 doi:10.1103/PhysRevD.108.012008 [arXiv:2206.07622 [hep-ex]].
 - [6] R. Aaij *et al.* [LHCb], *Phys. Rev. Lett.* **124** (2020) no.3, 031801 doi:10.1103/PhysRevLett.124.031801 [arXiv:1909.05211 [hep-ex]].
 - [7] I. Bediaga, T. Frederico and O. Lourenço, *Phys. Rev. D* **89** (2014) no.9, 094013 doi:10.1103/PhysRevD.89.094013 [arXiv:1307.8164 [hep-ph]].
 - [8] I. Bediaga and P. C. Magalhães, [arXiv:1512.09284 [hep-ph]].
 - [9] X. W. Kang, B. Kubis, C. Hanhart and U. G. Meißner, *Phys. Rev. D* **89** (2014), 053015 doi:10.1103/PhysRevD.89.053015 [arXiv:1312.1193 [hep-ph]].
 - [10] C. H. Chen and H. n. Li, *Phys. Lett. B* **561** (2003), 258-265 doi:10.1016/S0370-2693(03)00486-6 [arXiv:hep-ph/0209043 [hep-ph]].
 - [11] B. El-Bennich, A. Furman, R. Kaminski, L. Lesniak, B. Loiseau and B. Moussallam, *Phys. Rev. D* **79** (2009), 094005 [erratum: *Phys. Rev. D* **83** (2011), 039903] doi:10.1103/PhysRevD.83.039903 [arXiv:0902.3645 [hep-ph]].
 - [12] J. Virto, *PoS FPCP2016* (2017), 007 doi:10.22323/1.280.0007 [arXiv:1609.07430 [hep-ph]].
 - [13] S. Kränkl, T. Mannel and J. Virto, *Nucl. Phys. B* **899** (2015), 247-264 doi:10.1016/j.nuclphysb.2015.08.004 [arXiv:1505.04111 [hep-ph]].
 - [14] A. G. Grozin, *Sov. J. Nucl. Phys.* **38** (1983), 289-292
 - [15] A. G. Grozin, *Theor. Math. Phys.* **69** (1986), 1109-1121 doi:10.1007/BF01037870
 - [16] D. Müller, D. Robaschik, B. Geyer, F. M. Dittes and J. Hořejši, *Fortsch. Phys.* **42** (1994), 101-141 doi:10.1002/prop.2190420202 [arXiv:hep-ph/9812448 [hep-ph]].
 - [17] M. Diehl, T. Gousset, B. Pire and O. Teryaev, *Phys. Rev. Lett.* **81** (1998), 1782-1785 doi:10.1103/PhysRevLett.81.1782 [arXiv:hep-ph/9805380 [hep-ph]].
 - [18] M. Diehl, T. Gousset and B. Pire, *Phys. Rev. D* **62** (2000), 073014 doi:10.1103/PhysRevD.62.073014 [arXiv:hep-ph/0003233 [hep-ph]].
 - [19] B. Pire and L. Szymanowski, *Phys. Lett. B* **556** (2003), 129-134 doi:10.1016/S0370-2693(03)00134-5 [arXiv:hep-ph/0212296 [hep-ph]].
 - [20] M. V. Polyakov, *Nucl. Phys. B* **555** (1999), 231 doi:10.1016/S0550-3213(99)00314-4 [arXiv:hep-ph/9809483 [hep-ph]].
 - [21] Y. Li, D. C. Yan, J. Hua, Z. Rui and H. n. Li, *Phys. Rev. D* **104** (2021) no.9, 096014 doi:10.1103/PhysRevD.104.096014 [arXiv:2105.03899 [hep-ph]].
 - [22] J. Hua, H. n. Li, C. D. Lu, W. Wang and Z. P. Xing, *Phys. Rev. D* **104** (2021) no.1, 016025 doi:10.1103/PhysRevD.104.016025 [arXiv:2012.15074 [hep-ph]].
 - [23] Z. Rui, Y. Li and H. n. Li, *JHEP* **05**, 082 (2021) doi:10.1007/JHEP05(2021)082 [arXiv:2103.00642 [hep-ph]].
 - [24] Y. Li, D. C. Yan, Z. Rui and Z. J. Xiao, *Eur. Phys. J. C* **81**, no.9, 806 (2021) doi:10.1140/epjc/s10052-021-09608-5 [arXiv:2107.10684 [hep-ph]].
 - [25] C. Q. Zhang, J. M. Li, M. K. Jia, Y. Li and Z. Rui, *Phys. Rev. D* **105**, no.5, 053002 (2022) doi:10.1103/PhysRevD.105.053002 [arXiv:2112.10939 [hep-ph]].
 - [26] D. C. Yan, Z. Rui, Z. J. Xiao and Y. Li, *Phys. Rev. D* **105**, no.9, 093001 (2022) doi:10.1103/PhysRevD.105.093001 [arXiv:2204.01092 [hep-ph]].
 - [27] Z. Q. Zhang, Y. C. Zhao, Z. L. Guan, Z. J. Sun, Z. Y. Zhang and K. Y. He, *Chin. Phys. C* **46**, no.12, 123105 (2022) doi:10.1088/1674-1137/ac89d1 [arXiv:2207.02043 [hep-ph]].

- [28] Z. T. Zou, W. S. Fang, X. Liu and Y. Li, Eur. Phys. J. C **82**, no.11, 1076 (2022) doi:10.1140/epjc/s10052-022-11060-y [arXiv:2210.08522 [hep-ph]].
- [29] D. C. Yan, Z. Rui, Y. Yan and Y. Li, Eur. Phys. J. C **83**, no.10, 974 (2023) doi:10.1140/epjc/s10052-023-12152-z [arXiv:2308.12543 [hep-ph]].
- [30] D. C. Yan, Y. Yan and Z. Rui, Eur. Phys. J. C **84**, no.7, 754 (2024) doi:10.1140/epjc/s10052-024-13087-9 [arXiv:2404.19198 [hep-ph]].
- [31] J. P. Dedonder, A. Furman, R. Kaminski, L. Lesniak and B. Loiseau, Acta Phys. Polon. B **42** (2011), 2013 doi:10.5506/APhysPolB.42.2013 [arXiv:1011.0960 [hep-ph]].
- [32] M. Algueró, A. Crivellin, S. Descotes-Genon, J. Matias and M. Novoa-Brunet, JHEP **04**, 066 (2021) doi:10.1007/JHEP04(2021)066 [arXiv:2011.07867 [hep-ph]].
- [33] R. Aaij *et al.* [LHCb], Eur. Phys. J. C **78** (2018) no.12, 1019 doi:10.1140/epjc/s10052-018-6447-z [arXiv:1809.07416 [hep-ex]].
- [34] G. J. Gounaris and J. J. Sakurai, Phys. Rev. Lett. **21** (1968), 244-247 doi:10.1103/PhysRevLett.21.244
- [35] K. M. Watson, Phys. Rev. **88** (1952), 1163-1171 doi:10.1103/PhysRev.88.1163
- [36] S. Navas *et al.* [Particle Data Group], Phys. Rev. D **110** (2024) no.3, 030001 doi:10.1103/PhysRevD.110.030001
- [37] Y. Y. Keum, H. N. Li and A. I. Sanda, Phys. Rev. D **63** (2001), 054008 doi:10.1103/PhysRevD.63.054008 [arXiv:hep-ph/0004173 [hep-ph]].
- [38] T. Kurimoto, H. n. Li and A. I. Sanda, Phys. Rev. D **65** (2002), 014007 doi:10.1103/PhysRevD.65.014007 [arXiv:hep-ph/0105003 [hep-ph]].
- [39] C. D. Lu and M. Z. Yang, Eur. Phys. J. C **28** (2003), 515-523 doi:10.1140/epjc/s2003-01199-y [arXiv:hep-ph/0212373 [hep-ph]].
- [40] H. n. Li, Prog. Part. Nucl. Phys. **51** (2003), 85-171 doi:10.1016/S0146-6410(03)90013-5 [arXiv:hep-ph/0303116 [hep-ph]].
- [41] Z. J. Xiao, W. F. Wang and Y. y. Fan, Phys. Rev. D **85** (2012), 094003 doi:10.1103/PhysRevD.85.094003 [arXiv:1111.6264 [hep-ph]].
- [42] H. N. Li, Y. L. Shen and Y. M. Wang, JHEP **02** (2013), 008 doi:10.1007/JHEP02(2013)008 [arXiv:1210.2978 [hep-ph]].
- [43] C. D. Lu, K. Ukai and M. Z. Yang, Phys. Rev. D **63** (2001), 074009 doi:10.1103/PhysRevD.63.074009 [arXiv:hep-ph/0004213 [hep-ph]].
- [44] Y. Y. Keum, H. n. Li and A. I. Sanda, Phys. Lett. B **504** (2001), 6-14 doi:10.1016/S0370-2693(01)00247-7 [arXiv:hep-ph/0004004 [hep-ph]].
- [45] A. G. Grozin and M. Neubert, Phys. Rev. D **55** (1997), 272-290 doi:10.1103/PhysRevD.55.272 [arXiv:hep-ph/9607366 [hep-ph]].
- [46] Y. Yang, L. Lang, X. Zhao, J. Huang and J. Sun, Phys. Rev. D **103** (2021) no.5, 056006 doi:10.1103/PhysRevD.103.056006 [arXiv:2012.10581 [hep-ph]].
- [47] V. M. Braun, Y. Ji and A. N. Manashov, JHEP **05** (2017), 022 doi:10.1007/JHEP05(2017)022 [arXiv:1703.02446 [hep-ph]].
- [48] A. Ali, G. Kramer, Y. Li, C. D. Lu, Y. L. Shen, W. Wang and Y. M. Wang, Phys. Rev. D **76** (2007), 074018 doi:10.1103/PhysRevD.76.074018 [arXiv:hep-ph/0703162 [hep-ph]].
- [49] W. F. Wang and H. n. Li, Phys. Lett. B **763** (2016), 29-39 doi:10.1016/j.physletb.2016.10.026 [arXiv:1609.04614 [hep-ph]].
- [50] Z. Rui, Y. Li and H. N. Li, Phys. Rev. D **98** (2018) no.11, 113003 doi:10.1103/PhysRevD.98.113003 [arXiv:1809.04754 [hep-ph]].
- [51] R. Aaij *et al.* [LHCb], JHEP **05**, 026 (2019) doi:10.1007/JHEP05(2019)026 [arXiv:1812.07008 [hep-ex]].
- [52] G. Breit and E. Wigner, Phys. Rev. **49** (1936), 519-531 doi:10.1103/PhysRev.49.519
- [53] Y. Li, A. J. Ma, W. F. Wang and Z. J. Xiao, Phys. Rev. D **95** (2017) no.5, 056008 doi:10.1103/PhysRevD.95.056008 [arXiv:1612.05934 [hep-ph]].
- [54] J. P. Lees *et al.* [BaBar], Phys. Rev. D **86** (2012), 032013 doi:10.1103/PhysRevD.86.032013 [arXiv:1205.2228 [hep-ex]].
- [55] A. Ali, G. Kramer and C. D. Lu, Phys. Rev. D **58** (1998), 094009 doi:10.1103/PhysRevD.58.094009 [arXiv:hep-ph/9804363 [hep-ph]].
- [56] H. Y. Cheng, Phys. Rev. D **106** (2022) no.11, 113004 doi:10.1103/PhysRevD.106.113004 [arXiv:2211.03965 [hep-ph]].
- [57] P. Lepage and C. Gohlke, gplepage/lqfit: lqfit version 11.7, Zenodo. <http://doi.org/10.5281/zenodo.4037174>.
- [58] H. n. Li and S. Mishima, Phys. Rev. D **74** (2006), 094020 doi:10.1103/PhysRevD.74.094020 [arXiv:hep-ph/0608277 [hep-ph]].
- [59] Z. Rui, X. Gao and C. D. Lu, Eur. Phys. J. C **72** (2012), 1923 doi:10.1140/epjc/s10052-012-1923-3 [arXiv:1111.0181 [hep-ph]].
- [60] P. Ball, V. M. Braun, Y. Koike and K. Tanaka, Nucl. Phys. B **529** (1998), 323-382 doi:10.1016/S0550-3213(98)00356-3 [arXiv:hep-ph/9802299 [hep-ph]].
- [61] D. C. Yan, P. Yang, X. Liu and Z. J. Xiao, Nucl. Phys. B **931**, 79-104 (2018) doi:10.1016/j.nuclphysb.2018.04.007 [arXiv:1707.06043 [hep-ph]].
- [62] N. Gubernari, A. Kokulu and D. van Dyk, JHEP **01**, 150 (2019) doi:10.1007/JHEP01(2019)150 [arXiv:1811.00983 [hep-ph]].
- [63] A. Khodjamirian, T. Mannel and N. Offen, Phys. Rev. D **75**, 054013 (2007) doi:10.1103/PhysRevD.75.054013 [arXiv:hep-ph/0611193 [hep-ph]].
- [64] A. Bharucha, D. M. Straub and R. Zwicky, JHEP **08**, 098 (2016) doi:10.1007/JHEP08(2016)098 [arXiv:1503.05534 [hep-ph]].
- [65] S. Cheng, A. Khodjamirian and J. Virto, JHEP **05**, 157 (2017) doi:10.1007/JHEP05(2017)157 [arXiv:1701.01633 [hep-ph]].
- [66] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014029 (2005) doi:10.1103/PhysRevD.71.014029 [arXiv:hep-ph/0412079 [hep-ph]].
- [67] A. Khodjamirian, T. Mannel, A. A. Pivovarov and Y. M. Wang, JHEP **09**, 089 (2010) doi:10.1007/JHEP09(2010)089 [arXiv:1006.4945 [hep-ph]].
- [68] S. Descotes-Genon, A. Khodjamirian and J. Virto, JHEP **12**, 083 (2019) doi:10.1007/JHEP12(2019)083 [arXiv:1908.02267 [hep-ph]].
- [69] H. Y. Cheng and C. K. Chua, Phys. Rev. D **80** (2009), 114026 doi:10.1103/PhysRevD.80.114026 [arXiv:0910.5237 [hep-ph]].
- [70] C. Wang, S. H. Zhou, Y. Li and C. D. Lu, Phys. Rev. D **96** (2017) no.7, 073004 doi:10.1103/PhysRevD.96.073004 [arXiv:1708.04861 [hep-ph]].
- [71] C. Wang, Q. A. Zhang, Y. Li and C. D. Lu, Eur. Phys. J. C **77** (2017) no.5, 333 doi:10.1140/epjc/s10052-017-4889-3 [arXiv:1701.01300 [hep-ph]].
- [72] Z. T. Zou, A. Ali, C. D. Lu, X. Liu and Y. Li, Phys. Rev. D **91** (2015), 054033 doi:10.1103/PhysRevD.91.054033 [arXiv:1501.00784 [hep-ph]].
- [73] H. n. Li, Phys. Lett. B **622** (2005), 63-68 doi:10.1016/j.physletb.2005.06.077 [arXiv:hep-ph/0411305 [hep-ph]].

- [74] T. Aaltonen *et al.* [CDF], Phys. Rev. Lett. **107** (2011), 261802 doi:10.1103/PhysRevLett.107.261802 [arXiv:1107.4999 [hep-ex]].
- [75] R. Aaij *et al.* [LHCb], Phys. Lett. B **713** (2012), 369-377 doi:10.1016/j.physletb.2012.06.012 [arXiv:1204.2813 [hep-ex]].
- [76] K. F. Chen *et al.* [Belle], Phys. Rev. Lett. **91** (2003), 201801 doi:10.1103/PhysRevLett.91.201801 [arXiv:hep-ex/0307014 [hep-ex]].
- [77] B. Aubert *et al.* [BaBar], Phys. Rev. D **78** (2008), 092008 doi:10.1103/PhysRevD.78.092008 [arXiv:0808.3586 [hep-ex]].
- [78] J. P. Lees *et al.* [BaBar], Phys. Rev. D **85** (2012), 072005 doi:10.1103/PhysRevD.85.072005 [arXiv:1112.3896 [hep-ex]].
- [79] M. Beneke, J. Rohrer and D. Yang, Nucl. Phys. B **774** (2007), 64-101 doi:10.1016/j.nuclphysb.2007.03.020 [arXiv:hep-ph/0612290 [hep-ph]].
- [80] C. W. Bauer, D. Pirjol, I. Z. Rothstein and I. W. Stewart, Phys. Rev. D **70** (2004), 054015 doi:10.1103/PhysRevD.70.054015 [arXiv:hep-ph/0401188 [hep-ph]].
- [81] H. n. Li and S. Mishima, Phys. Rev. D **71** (2005), 054025 doi:10.1103/PhysRevD.71.054025 [arXiv:hep-ph/0411446 [hep-ph]].
- [82] H. Y. Cheng and K. C. Yang, Phys. Rev. D **78** (2008), 094001 [erratum: Phys. Rev. D **79** (2009), 039903] doi:10.1103/PhysRevD.79.039903 [arXiv:0805.0329 [hep-ph]].
- [83] Y. Grossman, Int. J. Mod. Phys. A **19** (2004), 907-917 doi:10.1142/S0217751X04018865 [arXiv:hep-ph/0310229 [hep-ph]].
- [84] P. K. Das and K. C. Yang, Phys. Rev. D **71** (2005), 094002 doi:10.1103/PhysRevD.71.094002 [arXiv:hep-ph/0412313 [hep-ph]].
- [85] C. H. Chen and C. Q. Geng, Phys. Rev. D **71** (2005), 115004 doi:10.1103/PhysRevD.71.115004 [arXiv:hep-ph/0504145 [hep-ph]].
- [86] Y. D. Yang, R. M. Wang and G. R. Lu, Phys. Rev. D **72** (2005), 015009 doi:10.1103/PhysRevD.72.015009 [arXiv:hep-ph/0411211 [hep-ph]].
- [87] A. L. Kagan, Phys. Lett. B **601** (2004), 151-163 doi:10.1016/j.physletb.2004.09.030 [arXiv:hep-ph/0405134 [hep-ph]].
- [88] M. Beneke, J. Rohrer and D. Yang, Phys. Rev. Lett. **96** (2006), 141801 doi:10.1103/PhysRevLett.96.141801 [arXiv:hep-ph/0512258 [hep-ph]].
- [89] A. Datta, A. V. Gritsan, D. London, M. Nagashima and A. Szynekman, Phys. Rev. D **76** (2007), 034015 doi:10.1103/PhysRevD.76.034015 [arXiv:0705.3915 [hep-ph]].
- [90] P. Colangelo, F. De Fazio and T. N. Pham, Phys. Lett. B **597** (2004), 291-298 doi:10.1016/j.physletb.2004.07.024 [arXiv:hep-ph/0406162 [hep-ph]].
- [91] M. Ladisa, V. Laporta, G. Nardulli and P. Santorelli, Phys. Rev. D **70** (2004), 114025 doi:10.1103/PhysRevD.70.114025 [arXiv:hep-ph/0409286 [hep-ph]].
- [92] H. Y. Cheng, C. K. Chua and A. Soni, Phys. Rev. D **71** (2005), 014030 doi:10.1103/PhysRevD.71.014030 [arXiv:hep-ph/0409317 [hep-ph]].
- [93] E. Alvarez, L. N. Epele, D. Gomez Dumm and A. Szynekman, Phys. Rev. D **70** (2004), 115014 doi:10.1103/PhysRevD.70.115014 [arXiv:hep-ph/0410096 [hep-ph]].
- [94] K. C. Yang, Phys. Rev. D **72** (2005), 034009 [erratum: Phys. Rev. D **72** (2005), 059901] doi:10.1103/PhysRevD.72.034009 [arXiv:hep-ph/0506040 [hep-ph]].
- [95] S. Baek, A. Datta, P. Hamel, O. F. Hernandez and D. London, Phys. Rev. D **72** (2005), 094008 doi:10.1103/PhysRevD.72.094008 [arXiv:hep-ph/0508149 [hep-ph]].
- [96] C. S. Huang, P. Ko, X. H. Wu and Y. D. Yang, Phys. Rev. D **73** (2006), 034026 doi:10.1103/PhysRevD.73.034026 [arXiv:hep-ph/0511129 [hep-ph]].
- [97] C. H. Chen and H. Hatanaka, Phys. Rev. D **73** (2006), 075003 doi:10.1103/PhysRevD.73.075003 [arXiv:hep-ph/0602140 [hep-ph]].
- [98] A. Faessler, T. Gutsche, J. C. Helo, S. Kovalenko and V. E. Lyubovitskij, Phys. Rev. D **75** (2007), 074029 doi:10.1103/PhysRevD.75.074029 [arXiv:hep-ph/0702020 [hep-ph]].
- [99] C. H. Chen, C. Q. Geng, Y. K. Hsiao and Z. T. Wei, Phys. Rev. D **72** (2005), 054011 doi:10.1103/PhysRevD.72.054011 [arXiv:hep-ph/0507012 [hep-ph]].
- [100] C. H. Chen and C. Q. Geng, Phys. Rev. D **75** (2007), 054010 doi:10.1103/PhysRevD.75.054010 [arXiv:hep-ph/0701023 [hep-ph]].
- [101] H. Y. Cheng and K. C. Yang, Phys. Rev. D **83** (2011), 034001 doi:10.1103/PhysRevD.83.034001 [arXiv:1010.3309 [hep-ph]].
- [102] C. Bobeth, M. Gorbahn and S. Vickers, Eur. Phys. J. C **75** (2015) no.7, 340 doi:10.1140/epjc/s10052-015-3535-1 [arXiv:1409.3252 [hep-ph]].
- [103] H. n. Li and S. Mishima, Phys. Rev. D **73**, 114014 (2006) doi:10.1103/PhysRevD.73.114014 [arXiv:hep-ph/0602214 [hep-ph]].
- [104] Y. Li, G. H. Zhao, Y. J. Sun and Z. T. Zou, Phys. Rev. D **106**, no.9, 093009 (2022) doi:10.1103/PhysRevD.106.093009 [arXiv:2209.13389 [hep-ph]].
- [105] R. Aaij *et al.* [LHCb], Phys. Rev. D **108**, no.1, 012013 (2023) doi:10.1103/PhysRevD.108.012013 [arXiv:2206.02038 [hep-ex]].
- [106] H. Y. Cheng, [arXiv:2005.06080 [hep-ph]].
- [107] J. Chai, S. Cheng, Y. h. Ju, D. C. Yan, C. D. Lü and Z. J. Xiao, Chin. Phys. C **46**, no.12, 123103 (2022) doi:10.1088/1674-1137/ac88bd [arXiv:2207.04190 [hep-ph]].
- [108] S. Cheng, Z. J. Xiao and Y. L. Zhang, Nucl. Phys. B **896**, 255-280 (2015) doi:10.1016/j.nuclphysb.2015.04.021 [arXiv:1409.5947 [hep-ph]].
- [109] J. X. Yu, J. J. Han, Y. Li, H. n. Li, Z. J. Xiao and F. S. Yu, [arXiv:2409.02821 [hep-ph]].