Stochastic Volatility under Informative Missingness

Gehui Zhang

Southwest Petroleum University & Department of Biostatistics, University of Pittsburgh Gong Tang Department of Biostatistics, University of Pittsburgh Lori N. Scott

Department of Psychiatry, University of Pittsburgh

and

Robert T. Krafty*

Department of Biostatistics and Bioinformatics, Emory University

January 28, 2025

Abstract

Stochastic volatility models that treat the variance of a time series as a stochastic process have proven to be important tools for analyzing dynamic variability. Current methods for fitting and conducting inference on stochastic volatility models are limited by the assumption that any missing data are missing at random. With a recent explosion in technology to facilitate the collection of dynamic self-response data for which mechanisms underlying missing data are inherently scientifically informative, this limitation in statistical methodology also limits scientific advancement. The goal of this article is to develop the first statistical methodology for modeling, fitting, and conducting inference on stochastic volatility with data that are missing not at random. The approach is based upon a novel imputation method derived using Tukey's representation, which utilizes the Markovian nature of stochastic volatility models to overcome unidentifiable components often faced when modeling informative missingness in other settings. This imputation method is combined with a new conditional particle filtering with ancestor sampling procedure that accounts for variability in imputation to formulate a complete particle Gibbs sampling scheme. The use of the method is illustrated through the analysis of mobile phone self-reported mood from individuals being monitored after unsuccessful suicide attempts.

Keywords: Ecological momentary assessment (EMA); Particle methods; Sequential Monte Carlo; Stochastic volatility; Time series; Tukey's representation

^{*}Corresponding author Robert T. Krafty, Department of Biostatistics and Bioinformatics, Emory University, Atlanta, GA 30322 (e-mail:rkrafty@emory.edu). This work is supported by National Institutes of Health grants R01MH115388, R01GM140476 and R01GM113243.

1 Introduction

Stochastic volatility models are nonlinear state space models where the variance or dispersion of a time series is modeled as a dynamic stochastic process. Given the presence and importance of heteroscedastic variability in derivative options, stochastic volatility models have become a popular tool for analyzing financial and econometric time series, and a sizable amount of research has been conducted on methods for their analysis. A comprehensive description of the development and use of stochastic volatility models can be found in the first chapter of Shephard (2005).

Technological advances have led to an explosion in the amount of dynamic data that are collected across a variety of fields. An important example of such data is activity and emotional data collected by mobile devices in real time, which are often referred to as ecological momentary assessment (EMA) data. Researchers have found associations between variability in mood and affect with clinical outcomes (Thompson et al. 2011, Kuppens et al. 2007, McConville & Cooper 1996, Angst et al. 2003). However, the current state of the art for evaluating variability in psychosocial and behavioral data either assumes homoscedastic variability, despite empirical evidence of heteroscedasticity, or utilizes coarse and naive windowed estimates within predefined time intervals. Stochastic volatility could be a potentially interpretable and predictive metric from EMA data that provides a more comprehensive, dynamic picture of variably. An example of such data comes from our motivating application, which is discussed in further detail in Section 7, which considers data from two young adults who experienced recent suicidal thoughts and/or behaviors (Cox et al. 2023). The participants were prompted to self-report measures of affect and mood seven times per day using a cell phone-based application. The time series of the participants' self-reported level of happiness are displayed in Figure 1. We desire an anal-



Figure 1: Self-reported happiness over time from participants who recently had suicidal ideation and/or behavior. The participants were prompted to report happiness 7 times per day over 3 weeks. One hundred (100) represents maximum happiness, and 0 represents minimum happiness.

ysis of these data that can provide estimates of stochastic emotional volatility, which can potentially be used to predict clinical outcomes, such as suicide ideation.

The observational nature of self-reported data prompted by mobile devices inherently results in nonignorable missingness. However, existing methods for estimating stochastic volatility assume any missing data are missing at random. For financial time series where the ignorability of missing data is plausible, *ad hoc* methods that assume data are missing completely at random, such as ignoring missing points, averaging, or aggregation, are commonly used to handle missing data. A technique for handling irregular sampling in data that uses the expectation-maximization (EM) algorithm within particle Gibbs sampling was proposed by Kim & Stoffer (2008), which can be used to analyze times series with missing data by assuming that missing observations do not relate to latent volatility and are missing at random. To the best of our knowledge, no methods currently exist for stochastic volatility

models with missing data that are not missing at random. The broad goal of this article is to develop the first formal method for the analysis of stochastic volatility with informative missing data.

Selection factorization (Rubin 1974) and pattern-mixture factorization (Rubin 1974, Little & Wang 1996) are arguably the two most popular approaches for addressing nonignorable missingness. However, both of these approaches involve an unidentifiable component that has to come from prior knowledge or remain unknown. Holland (1986) demonstrated that this unidentifiability is propagated into other components of these two popular factorizations. To address this challenge, as recorded by Holland (1986), John W. Tukey proposed an alternative method for decomposing the joint distribution in which all components of the representation are accessible. Tukey's representation was recently re-examined by Franks et al. (2020), who demonstrated its tractability when data are independent (i.e. not time series) and the distribution of the variable of interest is in the exponential family. One of two specific methodological contributions of our article is the development of a novel imputation procedure for the stochastic volatility model based on Tukey's representation. Although it provides a factorization with identifiable components, a potential challenge in the use of Tukey's representation is the need to estimate the conditional probability of missingness utilizing all data, both observed and unobserved. We show that the Markovian state space nature of the stochastic volatility model allows one to formulate and estimate this quantity using either a parametric or nonparametric logistic regression model. To the best of our knowledge, all past usages of the Tukey's representation have only considered parametric models for the missingness mechanism that are dictated by the natural parameter of the exponential family considered. Our method not only represents what we believe is the first use of Tukey's representation for time series, but also its first usage with

nonparametric modeling of the missing-data mechanism.

As a nonlinear state space model, there are a variety of techniques to approach Bayesian analysis for the stochastic volatility model. Carlin et al. (1992), Jacquier et al. (1994), Taylor (1994), and Kim et al. (1998) discussed classic Markov chain Monte Carlo (MCMC) methods for the stochastic volatility model. Particle Gibbs methods, also commonly known as sequential Monte Carlo, have become a popular approach to fitting stochastic volatility models. And rieu et al. (2010) proposed using the conditional particle filter (CPF) for the latent state, which guarantees that the target distribution is invariant, unlike previous methods. Later, Lindsten et al. (2014) proposed the conditional particle filter with ancestor sampling (CPF-AS) to overcome the path degeneracy common with CPF. The second specific methodological contribution of this article is the development of an imputed CPF-AS algorithm (ICPF-AS) that incorporates information from the missingness mechanism and presents a complete particle Gibbs method for the analysis of stochastic volatility models with informed missingness. To the best of our knowledge, the novel ICPF-AS is the first method for fitting stochastic volatility models with informed missingness. It should be noted that, as discussed in Section 8, although formulated for the analysis of the stochastic volatility model, the proposed method provides insights into the analysis of general nonlinear state space models with informed missingness.

The rest of this article is presented as follows. After introducing the univariate discrete time stochastic volatility model with missing data in Section 2, a likelihood decomposition is derived in Section 4.2 to facilitate and motivate the proposed methodology. The imputation method is constructed in Section 3, the new ICPF-AS to account for imputation is introduced in Section 4.2.2, while priors distributions and the complete particle Gibbs sampling scheme are discussed in Section 5. Simulation studies are presented in Section 6

to investigate the approach's empirical properties, and its use in the analysis of EMA data is illustrated in Section 7. A discussion of extensions, limitations, and further implications is provided in Section 8. Appendices provided in online supplemental materials provide details of the Gibbs sampler and additional simulation results.

2 Model

The basic univariate discrete time stochastic volatility model for a real-valued zero-mean time series $\{y_t\}_{t=1}^n$ defined as

$$y_t = \exp\left(\frac{h_t + \mu}{2}\right)\epsilon_t$$

$$h_{t+1} = \mu + \phi \left(h_t - \mu\right) + \eta_t$$
(1)

where $\epsilon_t \sim N(0,1)$, $\eta_t \sim N(0,\sigma^2)$, and $h_1 \sim N\left(\mu, \frac{\sigma^2}{1-\phi^2}\right)$, ϵ_t , η_t , j_1 are independent from each other. Here, $\{h_t\}_{t=1}^n$ is the volatility process and the primary focus of our analysis. In this article, we will utilize the notation where $h_{1:n}$ and $y_{1:n}$ represent the epochs of the stochastic processes $\{h_t\}_{t=1}^n$ and $\{y_t\}_{t=1}^n$.

Shephard (1996) and Taylor (1994) discussed theoretical details and econometric properties of this model when y_t are observed for all t = 1, ..., n. This article considers the more complicated setting where some values of $y_{1:n}$ could be missing. Define the response indicator at time t, r_t , as:

$$r_t = \begin{cases} 1 & \text{if } y_t \text{ is observed} \\ 0 & \text{if } y_t \text{ is missing.} \end{cases}$$

We assume that y_1 is always observed so that $r_1 = 1$. We are interested in the scenario where the probability of missing data at time t only depends on y_t , so that the presence of missing data is informative. We assume that r_t only depends on $h_{1:n}$ and $y_{1:n}$ through y_t , and that $\{r_t\}_{t=1}^n$ are independent conditional on $y_{1:n}$.

We consider that the basic stochastic volatility model (1) holds for the observed y_t given $r_t = 1$, and believe the conditional distribution is different when y_t is missing. Further, we consider the scenario where a model for the conditional log odds of $r_t = 1$ based on y_t can be assumed such that

$$\operatorname{logit}\left[f\left(r_{t}=1\mid y_{1:n},\theta_{R|Y}\right)\right]=g\left(y_{t}\mid\theta_{R|Y}\right)$$

$$(2)$$

for some function g of y_t that depends on a set of parameters $\theta_{R|Y}$. We consider two modeling approaches for (2) in this article. The first approach is the simple parametric case where $\theta_{R|Y} = (\beta_0, \beta_1)^T$ and $g(y_t|\theta_{R|Y}) = \beta_0 + \beta_1 y_t$, which accurately captures the hypothesized mechanism of the motivating application where lower levels of positive affect are more likely to be missing. As will be demonstrated in the next section, this setting provides for a simple, closed form for the distribution of y_t when $r_t = 0$.

The second approach is a nonparametric approach that models g as a smooth function with square integrable second derivatives through the Bayesian formulation of the cubic smoothing spline. Here, we model $g(y_t|\theta_{R|Y}) = \beta_0 + \beta_1 y_t + u(y_t)$, where u is a function with square-integrable second derivatives that is orthogonal to a space of linear functions in that $\int u(y)dy = \int u'(y)dy = 0$, and $\theta_{R|Y} = (\beta_0, \beta_1, u)$. This model decomposes g into a linear component and a component that accounts for curvature. Under our fully Bayesian approach that will be defined in detail in Section 5, weakly informative Gaussian priors are assumed for β_0 and β_1 , and u is modeled as a mean-zero Gaussian process whose covariance function (after scaling its domain between [0,1]) is $\lambda^{-1}R(x,y)$ where $R(x,y) = k_2(x)k_2(y) - k_4(|x-y|), k_2(x) = \frac{1}{2}[(x-0.5)^2 - \frac{1}{12}]$ and $k_4(x) = \frac{1}{24}[(x-0.5)^4 - \frac{(x-0.5)^2}{2} + \frac{7}{240}]$ (Gu 2013, Chapter 2). The smoothing parameter λ regularizes the roughness of g such that, as $\lambda \to 0$, u almost surely approaches a constant function of zero, and g approaches a linear function. Although any reasonable nonparametric approach could conceivably be considered, we consider the cubic smoothing spline as its decomposition into linear and nonlinear components, and the closed form under a linear model, enables for the intuitive and efficient formulation of an important sampler.

3 Implied distribution for missing data

With a slight abuse of notation that makes the dependence on h_t and Θ implicit, let

$$r(y_t) = p(r_t = 1 | y_t, \Theta, h_t)$$
$$p = p(r_t = 1 | \Theta, h_t).$$

If the positivity condition is satisfied where $P(R_t = r_t | \Theta) > 0$ and $P(Y_t = y_t | \Theta) > 0$ such that $P(R_t = r_t, Y_t = y_t | \Theta) > 0$ (Holland 1986), then Tukey's representation expresses the conditional distribution of missing y as

$$f(y_t|r_t = 0, \Theta, h_t) = \frac{p}{1-p} \frac{1-r(y_t)}{r(y_t)} f(y_t|r_t = 1, \Theta, h_t)$$
(3)

(details provided in the supplementary materials). The term $\frac{p}{1-p}$ is a normalizing constant that can be determined by normalizing the integral of $f(y_t|r_t = 0, \Theta, h_t)$ to unity. Thus, the unidentifiable $f(y_t|r_t = 0, \Theta, h_t)$ is expressed in terms of specified elements that can be estimated from observed data.

To understand each of these specified elements, we begin by expressing the density of

 y_t given $(\Theta, h_t, \mathbf{Y}^1)$ using the total probability rule as

$$f(y_t | \Theta, h_t, \mathbf{Y}^1) = pf(y_t | r_t = 1, \Theta, h_t)$$

+ $(1-p) \left[\frac{p}{1-p} \frac{1-r(y_t)}{r(y_t)} f(y_t | r_t = 1, \Theta, h_t) \right].$

Integrating over y_t , it can be found that $(1 - p)/p = \int f(y_t|r_t = 1, \Theta, h_t) \frac{1 - r(y_t)}{r(y_t)} dy_t$. In light of our assumption that the basic stochastic volatility model (1) is applicable to the observed y_t when $r_t = 1$,

$$y_t \mid r_t = 1, \Theta, h_t \sim N(0, \sigma_t^2)$$

 $\sigma_t^2 = \exp(h_t + \mu)$

Under the simple logistic-linear regression model where $\theta_{R|Y} = (\beta_0, \beta_1)'$ and $g(y_t \mid \theta_{R|Y}) = \beta_0 + \beta_1 y_t$ in Equation 2, $p/(1-p) = \exp\left(\beta_0 - \frac{\beta_1^2 \sigma_t^2}{2}\right)$. Consequently, it follows that

$$f(y_t|r_t = 0, \Theta, h_t, \sigma_t^2) = \frac{1}{\sqrt{2\pi\sigma_t^2}} \exp\left[-\frac{(y_t + \beta_1 \sigma_t^2)^2}{2\sigma_t^2}\right] \sim N(-\beta_1 \sigma_t^2, \sigma_t^2),$$

As expected, the implied distribution depends on the association of the probability of missingness, with imputed values having an expected value of zero when there is no association with $\beta_1 = 0$, a positive expected value when the odds of missingness is larger for larger values of y with $\beta < 0$, and a negative expected value when the odds of missingness are larger for smaller values of y with $\beta > 0$. Both the variance and the magnitude of the mean of the imputation distribution are proportional to σ_t^2 , whose expected magnitude depends on the current time's volatility h_t . Large positive values of current volatility, as well as large negative values of current volatility, imply larger expected values of σ_t^2 .

Under the cubic smoothing spline approach where $g(y_t|\theta_{R|Y}) = \beta_0 + \beta_1 y_t + u(y_t)$,

$$f(y_t|r_t = 0, \Theta, h_t) \propto \exp\left[-u(y_t)\right] \times N(-\beta_1 \sigma_t^2, \sigma_t^2).$$
(4)

Note that the nonlinear structure in the cubic smoothing spline alters the implied distribution, thereby affecting the estimation procedure for both volatility and imputation steps. Whereas the distribution under missingness for the parametric log-linear model is Gaussian from which observations can be directly simulated, this is not the case under the nonparametric model. However, as the imputation procedure is embedded into a larger sampler that utilizes importance sampling, this does not cause any complications in its use.

4 Particle Gibbs for SVM with missingness

4.1 Particle Gibbs for SVM

Particle filter (PF) methods (Pitt & Shephard 1999, Doucet & Johansen 2011), also known as sequential Monte Carlo methods, are a class of recursive algorithms used for filtering, smoothing, and estimating states in nonlinear and non-Gaussian dynamic systems. These methods approximate the posterior distribution using a set of N weighted random samples (particles), which are iteratively propagated through time based on a sequence of observations to provide estimates of the underlying state. PF methods blend sequential importance sampling (SIS) and resampling (Andrieu et al. 2010). In the SIS step, N particles are drawn from the proposal density and assigned important weights that account for the discrepancy between the true density and proposal density. In the resampling step, a new set of Nparticles is generated by sampling N times from the particles obtained in the SIS step.

Conditional particle filter (CPF) methods, also referred to as Particle Gibbs methods, are similar to the standard PF methods but with the difference that a reference trajectory is specified and retained throughout the sampling procedure (Andrieu et al. 2010). To address the poor mixing problem due to path degeneracy for CPF methods, Lindsten et al. (2014) proposed conditional particle with ancestor sampling (CPF-AS) methods. In CPF-AS methods, an artificial history is assigned to the last particle in an ancestor sampling step.

In the absence of any missing data, Bayesian approaches to analyzing the stochastic volatility model conduct inference based on simulated realizations from conditional posterior distributions through a Gibbs sampler that iterates between sampling $\theta_{Y|R} = (\mu, \sigma^2, \phi)$ given $(h_{1:n}, y_{1:n})$ and sampling $h_{1:n}$ given $(\theta_{Y|R}, y_{1:n})$. The particle methods can be utilized for sampling sequentially from the target distribution $f(h_{1:n} | y_{1:n}, \theta_{Y|R})$ utilizing a proposal distribution $q(h_t | h_{t-1}, y_t, \theta_{Y|R})$ based on the decomposition

$$f(h_{1:n} \mid y_{1:n}, \theta_{Y|R}) \propto f(h_{1:n}, y_{1:n} \mid \theta_{Y|R})$$

= $f(h_{1:n-1}, y_{1:n-1} \mid \theta_{Y|R}) \frac{f(h_n \mid h_{n-1}, \theta_{Y|R})f(y_n \mid h_n, \theta_{Y|R})}{q(h_n \mid h_{n-1}, y_n, \theta_{Y|R})}$
 $\times q(h_n \mid h_{n-1}, y_n, \theta_{Y|R}).$ (5)

If we consider N particles and let \tilde{h}_{t-1}^i be the resampled-derived ancestor of h_t , details of which will be provided in Algorithm 2, the weight for the *i*th particle at t is defined as

$$w_{t}^{i} = \frac{f(y_{t}|h_{t}^{i}, \theta_{Y|R})f(h_{t}^{i}|h_{t-1}^{i}, \theta_{Y|R})}{q(h_{t}^{i}|\tilde{h}_{t-1}^{i}, y_{t}, \theta_{Y|R})}$$

Although one can choose many appropriate proposal distributions, as is common for state space models, the proposal distribution can be selected to be the state equation so that $q(h_t^i|\tilde{h}_{t-1}^i, y_t, \theta_{Y|R}) = f(h_t^i|\tilde{h}_{t-1}^i, \theta_{Y|R})$, which is $N\left[\mu + \phi\left(\tilde{h}_{t-1}^i - \mu\right), \sigma^2\right]$ for t > 1, and the weight functions simplify to $w_t^i = f\left(y_t \mid h_t^i, \theta_{Y|R}\right)$.

4.2 Particle Gibbs for SVM with missingness

Let $\mathbf{Y}^1 = \{y_t : r_t = 1, t = 1...n\}$ and $\mathbf{Y}^0 = \{y_t : r_t = 0, t = 1...n\}$ be the collection of observed and missing values up to time *n* such that $y_{1:n} = \{\mathbf{Y}^1, \mathbf{Y}^0\}$, and let $\Theta = (\theta_{Y|R}, \theta_{R|Y})$ be the collection of parameters $\theta_{Y|R}$ of the stochastic volatility model and $\theta_{R|Y}$ of the logistic regression model for the probability of missingness. Our proposed Gibbs sampler that accounts for missingness-related information iterates between:

- (i) Sampling $\theta_{Y|R}$ given $(h_{1:n}, \mathbf{Y}^1, \mathbf{Y}^0, \theta_{R|Y})$.
- (ii) Sampling $\theta_{R|Y}$ given $(h_{1:n}, \mathbf{Y}^1, \mathbf{Y}^0, \theta_{Y|R})$.
- (iii) Sampling $(\mathbf{Y}^0, h_{1:n})$ given (\mathbf{Y}^1, Θ) .

Step (i) can be successfully achieved through existing methods that have been used for estimating static parameters in the stochastic volatility model without missing data, and it follows from Equation 2 that step (ii) can be achieved through logistic regression. A challenging and innovative component to solving this novel problem is in step (iii).

We propose to sample $(\mathbf{Y}^0, h_{1:n})$ given (\mathbf{Y}^1, Θ) sequentially using particle methods that are derived in Section 4.2.1. At time t,

- (i) if y_t is observed, sample h_t from proposal density $f(h_t \mid h_{t-1}, \Theta)$ and calculate the importance weights $w_t = f(y_t \mid h_t, \Theta)$, and
- (ii) if y_t is missing, sample (y_t, h_t) from proposal density $q(y_t, h_t \mid h_{t-1}, \Theta) = q(y_t \mid h_t, \Theta) f(h_t \mid h_{t-1}, \Theta)$ and compute the corresponding importance weights $w_t = 1$ for the parametric logistic linear setting, and $w_t = \exp[-u(y_t)]$ for the nonparametric smoothing spline setting.

The weights when $r_t = 1$ are the same as those for PF methods when there are no missing data and quantify the likelihood of a proposed h given the observed y. When $r_t = 0$, under the parametric logistic linear model, we can sample both h and y directly from their target distributions, so that particles have equal weighting and importance sampling is trivial. Under the nonparametric smoothing spline model, we can sample h directly from its target distribution, but cannot sample y directly from its target distribution; the weights reflect the differences between the proposal and target distributions of y.

4.2.1 Decomposition of the target distribution

While the dimensions of the proposal density differ for the proposed sequential sampler depending on $r_t = 0$, where we want to sample h_t , or $r_t = 1$, where we want to sample (y_t, h_t) , a prerequisite for basic particle methods is that the target distribution has a fixed dimension. We derive the presented sampler by embedding it into a larger problem with a fixed dimension. Consider $\{x_t\}_{t=1}^n$ defined as

$$x_t \sim \exp\left[-u(x_t)\right] \times N(-\beta_1 \sigma_t^2, \sigma_t^2) \tag{6}$$

where $\sigma_t^2 = \exp(h_t + \mu)$ and u(x) = 0 in the simple logistic-linear case. The series $\{x_t\}_{t=1}^n$ is generated under the missing distribution at all time points and is never observed. When $r_t = 0, y_t = x_t$; when $r_t = 1, x_t$ is a counterfactual that could have occurred if no data were observed. Sequentially sampling from the variable dimension $(h_{1:n}, Y_{1:n}^0 | Y_{1:n}^1, \Theta)$ can be extended to sampling from a fixed dimension $(h_{1:n}, x_{1:n} | Y_{1:n}^1, \Theta)$. At time t,

$$f(h_{1:t}, x_{1:t} \mid Y_{1:t}^{1}, \Theta) \propto f(h_{1:t}, x_{1:t}, Y_{1:t}^{1} \mid \Theta)$$

$$= f(h_{1:t-1}, x_{1:t-1}, Y_{1:t-1}^{1} \mid \Theta)$$

$$\times \left[\frac{f(x_{t} \mid h_{t}, \Theta) f(h_{t} \mid h_{t-1}, \Theta)}{q(x_{t}, h_{t} \mid h_{t-1}, \Theta)} q(x_{t}, h_{t} \mid h_{t-1}, \Theta) \right]^{1-r_{t}}$$

$$\times \left[\frac{f(x_{t} \mid h_{t}, y_{t}, \Theta) f(y_{t} \mid h_{t}, \Theta) f(h_{t} \mid h_{t-1}, \Theta)}{q(x_{t}, h_{t} \mid h_{t-1}, \Theta)} q(x_{t}, h_{t} \mid h_{t-1}, \Theta) \right]^{r_{t}}$$
(7)

where $q(x_t, h_t \mid h_{t-1}, \Theta)$ is the proposal density for the target distribution. At time t, the importance weight for each particle will be adjusted based on the value of r_t . For the *i*th particle,

$$w_t^i = W_{\Theta,t}(h_t^i, x_t^i, r_t) = \left[\frac{f(x_t^i \mid h_t^i, \Theta) f(h_t^i \mid \tilde{h}_{t-1}^i, \Theta)}{q(x_t^i, h_t^i \mid \tilde{h}_{t-1}^i, \Theta)}\right]^{1-r_t} \times \left[\frac{f(x_t^i \mid h_t^i, y_t, \Theta) f(y_t \mid h_t^i, \Theta) f(h_t^i \mid \tilde{h}_{t-1}^i, \Theta)}{q(x_t^i, h_t^i \mid \tilde{h}_{t-1}^i, \Theta)}\right]^{r_t}.$$
(8)

where \tilde{h}_{t-1}^i is the ancestor for (h_t^i, x_t^i) . However, if we choose the proposal density as $q(x_t, h_t \mid h_{t-1}, \Theta) = f(x_t \mid h_t, \Theta) f(h_t \mid h_{t-1}, \Theta)$ at time points when $r_t = 1$, the sampled x_t remains unused for importance weights or the inference of x and h series for these timepoints.

Under the cubic smoothing spline missing-data mechanism, the importance weights are given by $w_t^i = W_{\Theta,t}(h_t^i, x_t^i, r_t) = \{\exp[-u(y_t)]\}^{1-r_t}$ where at instances when $r_t = 1$, the importance weights incorporate the information from observed data, while when $r_t = 0$, the weights can account for the discrepancy between the proposal and target density. Under the simple logistic-linear missing-data mechanism, the importance weights are reduced to $w_t^i = W_{\Theta,t}(h_t^i, x_t^i, r_t) = [1]^{1-r_t} \times [f(y_t \mid h_t^i, \Theta)]^{r_t}$ where now at instances when $r_t = 0$, equal weights are assigned. This is because, in the latter case, sampling is done directly from the true density.

For the ancestor sampling step in the CPF-AS algorithm, the index for the last particle is sampled with probability

$$P(J=i) = \frac{W_{t-1}^i f_{\theta}(h'_t, x'_t | h_{t-1}^i)}{\sum_{k=1}^N W_{t-1}^k f_{\theta}(h'_t, x'_t | h_{t-1}^k)},$$

where (h'_t, x'_t) is the reference trajectory that is prespecified and retained for the CPF-AS

algorithm. The probability function can be reduced to

$$P(J=i) = \frac{W_{t-1}^{i} f_{\theta}(x_{t}' \mid h_{t}') f_{\theta}(h_{t}' \mid h_{t-1}^{i})}{\sum_{k=1}^{N} W_{t-1}^{k} f_{\theta}(x_{t}' \mid h_{t}') f_{\theta}(h_{t}' \mid h_{t-1}^{k})} = \frac{W_{t-1}^{i} f_{\theta}(h_{t}' \mid h_{t-1}^{i})}{\sum_{k=1}^{N} W_{t-1}^{k} f_{\theta}(h_{t}' \mid h_{t-1}^{k})}.$$
(9)

The first equation follows from the definition of x_t , and the second equation holds as $f_{\theta}(x'_t \mid h'_t)$ is invariant across particles by definition of the conditional particle filter methods.

We observe that the sampled x_t is not utilized in the computation of any important weights or indices. Additionally, the proposal density suggests that the sampling of h_t does not incorporate x_t , irrespective of whether $r_t = 1$ or $r_t = 0$. Therefore, we can create a sub-sequence $(Y_{1:n}^0, h_{1:n})$ from $(x_{1:n}, h_{1:n})$ in a way that excludes the unused $\{x_t : r_t = 1\}$ from the sampling process, which constitutes our proposed sampling scheme introduced in 4.2.

4.2.2 Imputed CPF-AS (ICPF-AS)

We refer to the CPF-AS methods under imputation as imputed CPF-AS (ICPF-AS). The sampling procedure of ICPF-AS involves two steps for sampling h: the initial step (t = 1)and the updating step (t = 2...n). For the initial step in Algorithm 1, particles for h_1 are sampled from the proposal density and weights computed, while Algorithm 2 sequentially utilizes re-sampling and ancestor sampling for times t > 1.

5 Prior Distributions and Sampling Scheme

The approaches discussed in Section 4.2 for sampling $(\mathbf{Y}^{\mathbf{0}}, h_{1:n})$, which is one component of the Gibbs sampler that iterates between sampling $\theta_{Y|R}$, $\theta_{R|Y}$ and $(\mathbf{Y}^{\mathbf{0}}, h_{1:n})$. A full iteration

Algorithm 1 Initialization step for ICPF-AS at t=1

Input: The reference trajectory $(h'_{1:n}, \{y_{1:n}\}')$, and Θ .

- 1: Draw $h_1^i \sim q_{\theta,t}(h_1|y_1)$, for $i = 1, \dots, N-1$.
- 2: Set $h_1^N = h_1'$.
- 3: For i = 1, ..., N, calculate the importance weights $w_1^i = W_{\Theta,1}(h_1^i, r_1)$, and get normalized weights W_1^i .
- 4: return $\{h_1^i\}_{i=1}^N, \{W_1^i\}_{i=1}^N$.

Algorithm 2 Updating ICPF-AS at time t > 1

Input: The reference trajectory $(h'_{1:n}, \{y_{1:n}\}'), \{(h_{1:t-1}, Y^0_{1:t-1})^i\}_{i=1}^N, \{W^i_{t-1}\}_{i=1}^N, \text{ and } \Theta.$

- 1: Draw with replacement from $\{(h_{1:t-1}, Y_{1:t-1}^{0})^i\}_{i=1}^N$ with probability $\{W_{t-1}^i\}_{i=1}^N$ for N-1 times to generate $\{(\tilde{h}_{1:t-1}, \tilde{Y}_{1:t-1}^0)^i\}_{i=1}^{N-1}$.
- 2: Set $(\tilde{h}_{1:t-1}, \tilde{Y}_{1:t-1}^{0})^{N} = (h_{1:t-1}, Y_{1:t-1}^{0})^{J}$, where J is drawn from 1, ..., N with probability $P(J=i) = \frac{W_{t-1}^{i} f_{\theta}(h'_{t}|h_{t-1}^{i})}{\sum_{k=1}^{N} W_{t-1}^{k} f_{\theta}(h'_{t}|h_{t-1}^{k})}$.
- 3: if $r_t = 0$ then
- Sample $(h_t, y_t^0)^i \sim q(y_t^0, h_t \mid h_{t-1}, \Theta)$ for i = 1, ..., N 1, and set $(h_t, y_t^0)^N =$ 4: $(\tilde{h}_{t-1}, \tilde{y}_{t-1}^0)^N.$
- 5: else if $r_t = 1$ then
- Sample $h_t^i \sim q(h_t \mid h_{t-1}, \Theta)$ for i = 1, ..., N-1, and set $h_t^N = \tilde{h}_{t-1}^N$. 6:
- 7: end if
- 8: For i = 1, ..., N, compute the importance weights $w_t^i = W_{\Theta,t}(h_t^i, y_t^{0i}, r_t)$ and calculate normalized weights $W_t^i = w_t^i / \sum_{j=1}^N w_t^j$. 9: **return** $\{(h_{1:t}, Y_{1:t}^0)^i\}_{i=1}^N, \{W_t^i\}_{i=1}^N$.

of the Gibbs sampler, including assumed prior distributions is given by:

(i) Sampling $\theta_{Y|R}$ given $(h_{1:n}, \mathbf{Y}^1, \mathbf{Y}^0, \theta_{R|Y})$

We assume independent priors for μ and (σ, ϕ) , which allows for their sequential sampling. Assuming a diffuse prior on μ (Kim et al. 1998), we sample μ from its posterior distribution

$$\mu | \sigma^2, \phi, h_{1:n}, y_{1:n} \sim N(\hat{\mu}, \sigma_{\mu}^2)$$

where $\hat{\mu} = \sigma_{\mu}^2 \left\{ \frac{(1-\phi^2)}{\sigma^2} h_1 + \frac{(1-\phi)}{\sigma^2} \sum_{t=1}^{n-1} (h_{t+1} - \phi h_t) \right\}$ and $\sigma_{\mu}^2 = \frac{\sigma^2}{(n-1)(1-\phi)^2 + (1-\phi^2)}$.

Although individual sampling of σ and ϕ has been commonly used in the past

(Kim et al. 1998), the mutual dependence of σ and ϕ makes the use of independent priors and condition sampling inefficient. We consider an approach that samples σ and ϕ jointly using a bivariate normal prior that accounts for this dependency and the Random Walk Metropolis-Hastings (RWMH) algorithm (Gong & Stoffer 2021). Details for the joint sampling of σ and ϕ and a brief review of individual sampling methods are provided in the supplemental materials.

- (ii) Sampling $\theta_{R|Y}$ given $(h_{1:n}, \mathbf{Y}^1, \mathbf{Y}^0, \theta_{Y|R})$.
 - (a) Under the simple logistic missing-data mechanism, the parameters β_0 , β_1 can be sampled using the Pólya-Gamma latent variable method for Bayesian logistic regression (Polson et al. 2013). For hyperparameters m_0 and B, we assume the prior where $\beta \sim N(m_0, B)$. To be consistent with logistic regression, we reparameterize and consider $m_t = 1 - r_t, t = 1, ..., n$ as the outcome variable that indicates missing in the regression. Let $X = (x_1, ..., x_n)'$ denote the design matrix, where $x_i = (1, y_i)$. We then sample β as

$$z_i | \boldsymbol{\beta} \sim PG(1, x_i^T \boldsymbol{\beta})$$

 $\boldsymbol{\beta} | m, z \sim N(\mu_z, V_z)$

where $V_z = (X^T Z X + B^{-1})^{-1}$, $\mu_z = V_z (X^T \kappa + B^{-1}b)$, $\kappa = \frac{1}{2} \mathbf{1}_n$ and Z is the $n \times n$ diagonal matrix with *i*th diagonal element z_i . The estimation of $\boldsymbol{\beta} = (\beta_0, \beta_1)$ is performed with standardized y such that the covariance matrix B can be reduced to a diagonal matrix. According to the interpretation of model parameters of a logistic regression model, we propose to choose and update the hyperparameter m_0 with the following procedure:

• Choose the first item of m_0 as $\log\left(\frac{m_{mean}}{1-m_{mean}}\right)$, where m_{mean} is the mean value

of $\{m_t\}_{t=1}^n$.

- For the second item of m_0 , we choose an arbitrary starting value for the first iteration, and for the *i*th iteration (i > 1), the timepoints are split into $T^u = t : y_t > 0.5$ and $T^l = t : y_t <= 0.5$, and update the item as $\log\left(\frac{m_{mean}^u/1-m_{mean}^u}{m_{mean}^l/1-m_{mean}^l}\right)$ where m_{mean}^u is the mean value of $\{m_t\}_{t\in T^u}$ and m_{mean}^l is the mean value of $\{m_t\}_{t\in T^l}$.
- (b) In the context of the cubic smoothing spline missing-data mechanism, we propose to employ a computationally efficient low-rank approximation. Rescalling data between [0,1], define y^{*}_t = [y_t min(y)] / [max(y) min(y)], d₁ = β₀+β₁min(y), and d₂ = β₁ [max(y) min(y)]. The right side of the missing-data mechanism can be written as g(Y^{*}) = Sd + Rc, where Y^{*} is the n dimensional vector with *i*th element y^{*}_i, S is the n × 2 matrix with *i*th row (1, y^{*}_i), R is the n × n matrix with *ij* element R(y^{*}_i, y^{*}_j), R(·, ·) is the kernel defined in Section 2, and c ~ N(0, λ⁻¹R⁻¹). The k-dimensional low-rank approximation considers a k-dimensional grid s_j = j/k, defines the k × k matrix Q as the evaluation of the kernel R over this grid, and lets Q = UDU^T be its singular value decomposition. Defining the n × k matrix Q̃ with *ij*th element R(y^{*}_i, s_j) and R̃ = Q̃U^TD^{-1/2}, the low-rank approximation considers g(Y^{*}) = Sd + R̃c where

$$(\boldsymbol{d}, \tilde{\boldsymbol{c}})^T \sim N(m_{k+2 \times 1}, B_{(k+2) \times (k+2)}),$$

 $\lambda^{-\frac{1}{2}} \sim \text{Half-}t(\nu_{\lambda}, G_{\lambda}),$

 $m_{k+2\times 1}$ is the zero vector and ν_{λ} , G_{λ} and $B_{(k+2)\times (k+2)} = \text{diag} \{\sigma_d^2, \sigma_d^2, \lambda^{-1}, \dots, \lambda^{-1}\}$ are hyperparameters. The half-t distribution is equivalent to a scale mixture of inverse Gamma distribution (Wand et al. 2011),

Algorithm 3 ICPF-AS for Sampling Y^0 , $h_{1:n}$

Input: The reference trajectory $(h'_{1:n}, \{y_{1:n}\}')$, and Θ .

- 1: Initiate the algorithm using Algorithm 1.
- 2: for t = 2 to n do
- 3: Update h_t using the Algorithm 2.
- 4: end for
- 5: Sample K from 1, ..., N with $P(K = i) = W_n^i$
- 6: return $h_{1:n}^K$.

$$\lambda^{-1} | q_{\lambda} \sim \text{Inv-}\Gamma\left(\frac{v}{2}, \frac{v}{q_{\lambda}}\right)$$
$$q_{\lambda} \sim \text{Inv-}\Gamma\left(\frac{1}{2}, \frac{1}{G_{\lambda}^{2}}\right)$$

The coefficients \boldsymbol{d} and $\boldsymbol{\tilde{c}}$ are then sampled with the Pólya-Gamma latent variable

method. The posterior distribution for λ is

$$egin{aligned} \Omega_{\lambda} \mid oldsymbol{d}, ilde{oldsymbol{c}}, \lambda &\sim \operatorname{Inv-}\Gamma\left(rac{
u_{\lambda}+1}{2}, rac{1}{G_{\lambda}^{2}} +
u_{\lambda}\lambda
ight) \ \lambda^{-1} \mid oldsymbol{d}, ilde{oldsymbol{c}}, \Omega &\sim \operatorname{Inv-}\Gamma\left(rac{
u_{\lambda}+k}{2}, rac{
u_{\lambda}}{\Omega_{\lambda}} + rac{1}{2} ilde{oldsymbol{c}}^{T} ilde{oldsymbol{c}}
ight) \end{aligned}$$

(iii) Sampling $(\mathbf{Y}^0, h_{1:n})$ given (\mathbf{Y}^1, Θ)

The parameters $(\mathbf{Y}^0, h_{1:n})$ are sampled using ICPF-AS in the last part of the Gibbs sampler utilizing methods developed in Sections 4.2, as formalized in Algorithm 3.

6 Simulation

For the stochastic volatility model 1, we set the true values of $\theta_{Y|R} = (\mu, \sigma^2, \phi)$ as (0.1, 0.25, 0.8)and the number of time points n = 100, 500. For the linear missing-data mechanism, we set the parameters for the missing-data mechanism $\beta_0 = -3$ and $\beta_1 = \ln(2.5), \ln(3), \ln(3.5)$. For example, Figure 2 shows the volatility h and time series y sequence generated from



Figure 2: The volatility h and time series y sequence generated from model 1 with sample size n = 100 and $\beta_1 = \ln(3)$.

model 1 with sample size n = 100 and $\beta_1 = \ln(3)$.

For the smoothing spline missing-data mechanism, the true missing-data mechanism is selected as $\log \left[f(r_t = 1 \mid y_{1:n}, \theta_{R|Y}) \right] = \beta_0 + \beta_1 y_t + y_t^2. \tag{10}$

We set the parameters for missing-data mechanism $\beta_0 = -2$ and $\beta_1 = \ln(2.5), \ln(3.5), \ln(4.5)$. The number of particles is N = 20. We run 500 replicates for each beta and sample size combination. For example, Figure 3 shows the volatility h and time series y sequence generated from model 1 with sample size n = 100 and $\beta_1 = \ln(4.5)$.

Regarding the hyperparameters for model parameters, we set the hyperparameters for ϕ and σ as $\mu_{\phi} = 0.875$, $\mu_{\sigma} = 0.45$, $\sigma_{\phi} = 0.075$, $\sigma_q = 0.1$ and $\rho = -0.25$. For the linear missingdata mechanism, the first hyperparameter for $\boldsymbol{\beta} = (\beta_0, \beta_1)$ is updated in each iteration with starting value $m_0 = (0, 1)$ and the second hyperparameter is selected as $B = I_{2\times 2}$. This updating procedure for m_0 can help mitigate the sensitivity of hyperparameters. Regarding B, we recommend starting with the identity matrix and considering increasing or decreasing



Figure 3: The volatility h and time series y sequence generated from model 1 with sample size n = 100 and smoothing spline missing-data mechanism with $\beta_1 = \ln(4.5)$.

the diagonal items if the algorithm fails. For smoothing spline estimation, we set k = 15.

For all simulation settings, the number of particles is N = 20. We run 500 replicates for each β and sample size combination, and we run the Gibbs sampler for 32500 iterations and discard the first 2500 iterations as burn-in. For the linear missing-data mechanism, we set the initial value for the parameters as $(\mu^{[0]}, \phi^{[0]}, \sigma^{[0]}, \beta_0^{[0]}, \beta_1^{[0]}) = (0.15, 0.9, 0.2, -1, 1)$. Figure 4 presents the true volatility h, posterior median volatility, and corresponding 95% credible interval with sample size n = 100 and $\beta_1 = \ln(3)$ for a simulated series under the linear missing-data mechanism. For the smoothing spline missing-data mechanism, we set the initial value for the parameters as $(\mu^{[0]}, \phi^{[0]}, \sigma^{[0]}) = (0.15, 0.9, 0.2)$, $d^{[0]} = (0, -1)'$, $c^{[0]} = \mathbf{0}_{k\times 1}$, and $\lambda^{[0]} = \exp(-7)$. Figure 5 presents the true volatility h, posterior median volatility, and corresponding 95% credible interval for a simulated series with sample size n = 100 and $\beta_1 = \ln(4.5)$.

We calculated the mean posterior coverage rate and width of 95% credible intervals over time, along with the average mean squared error (AMSE) over time for the stochastic



Figure 4: The true volatility h (dashed line), posterior median volatility (solid line), and corresponding 95% credible interval (shaded area) with sample size n = 100 and $\beta_1 = \ln(3)$ under the linear missing-data mechanism.



Figure 5: The true volatility h (dashed line), posterior median volatility (solid line), and corresponding 95% credible interval (shaded area) with sample size n = 100 and smoothing spline missing-data mechanism with $\beta_1 = \ln(4.5)$.

volatility $h_{1:n}$. The summarized results are presented in Table 1. According to the results, the mean coverage rates show comparability across the two sample size groups. Within the same sample size category, the mean coverage rate is lower in the group with a larger odds ratio $\exp(\beta_1)$ of missingness under the logistic missing-data mechanism. The average mean squared error is smaller when the sample size is larger, but increases when the odds ratio of missingness is larger.

Missing-data	Sample size	$\exp(\beta_1)^1$	AMSE	Mean width	Mean coverage rate
Mechanism					
		2.5	0.7598	2.3018	0.9290
	500	3	0.7962	2.3004	0.9185
Lincor		3.5	0.8055	2.2526	0.9054
Lillear		2.5	0.7811	2.3676	0.9359
	100	3	0.8130	2.3746	0.9307
		3.5	0.8400	2.3772	0.9250
		2.5	0.7368	2.3411	0.9421
	500	3.5	0.7403	2.3552	0.9442
Smoothing		4.5	0.7493	2.3763	0.9457
Spline		2.5	0.7753	2.4233	0.9495
	100	3.5	0.7756	2.4372	0.9505
		4.5	0.7847	2.4469	0.9503

Table 1: Mean posterior coverage rate and width of 95% credible interval over time and the average mean squared error(AMSE) over time for the stochastic volatility h

We also conduct a comparison between the proposed method (method P) and two commonly used imputation techniques: imputing missing y values with the mean of observed data (method A) and using the last observed data point for imputation (method B). We employ the same simulation settings. The summarized results for the stochastic volatility h are presented in Table 5. The findings clearly indicate that the proposed method outperforms the other two imputation methods concerning the estimation of volatility series, and shows comparable performance in terms of model parameter estimation.

Additional results, including the mean posterior coverage rate, the width of the 95% credible interval over time, and the mean squared error (AMSE) over time for model parameters, are provided in the supplementary materials.

				AMSE	
Missing-data Mechanism	Sample size	$\exp(\beta_1)$	Method P	Method A	Method B
		2.5	0.7598	0.9133	0.7891
	500	3	0.7836	1.1203	0.8352
Lincor		3.5	0.8055	1.4925	0.8906
Linear		2.5	0.7811	0.8738	0.7987
	100	3	0.8130	1.0103	0.8535
		3.5	0.8400	1.1687	0.8981
		2.5	0.7368	0.8005	0.7786
	500	3.5	0.7403	0.8229	0.7913
Smoothing		4.5	0.7493	0.8394	0.7961
Spline		2.5	0.7753	0.8183	0.7825
	100	3.5	0.7756	0.8219	0.7856
		4.5	0.7847	0.9061	0.8725

Table 2: Average mean squared error (AMSE) over time for the stochastic volatility h

7 Application

Suicide rates in the United States among young adults aged 18-24 have been consistently rising over the past two decades (Garnett & Curtin 2023); there is an urgent need to identify ways to understand who is at the highest risk for suicide in the near future. Technological advancements in the form of mobile devices have given clinicians and researchers the ability to monitor those at risk at fine time scales. These data can potentially be used to identify dynamic, proximal risk factors, which could be used to deploy time-sensitive interventions. Our motivating application considers data collected at the University of Pittsburgh as part of a recently completed study to identify near-term risk factors for suicidal thoughts and behavior in young adults who had suicidal ideation and/or behavior in the past 4 months. The data considered, which are displayed in Figure 1, are EMA self-response from two study participants to the question "At this moment, I feel happy" using a visual analog scale slider with a scale of 0 to 100, where 0 indicates not all and 100 indicates very much. The participants were prompted 7 times per day for 21 days, for a potential total of 147 observed responses. The mean (standard deviation) of the happiness responses are 29.43 (21.98) and 33.99 (27.92) for subjects one and two, respectively, and there 1 - p = 4.1% and 1 - p = 8.8% missing data where the participant did not respond to the prompt for subjects 1 and 2, respectively.

We analyzed the motivating data using the proposed procedure under both the simple logistic-linear missing-data mechanism and the cubic smoothing spline mechanism. Since the findings remain consistent across these two distinct missing-data mechanisms, we provide an elaborate discussion solely on the results obtained using the simple logistic-linear missing-data mechanism (Results pertaining to the cubic smoothing spline mechanism are available in the supplemental materials). The estimated posterior median (standard deviation) for β_1 was -0.059 (0.057) for subject 1 and -0.081 (0.051) for subject 2. This suggests that the odds of the participant not responding to the prompt are slightly higher for smaller values of happiness, which is consistent with disengagement during periods of low positive affect. The estimated posterior median of stochastic volatilities is displayed in Figure 7. We see a dynamic pattern in volatility, in particular a nadir between 10 and 16 days from the start of observation for subject 1 and between 3 and 7 days for subject 2.

Researchers have found variability in mood to be associated with suicidal ideation (Palmier-Clauss et al. 2013), where variability was quantified in two ways: as the mean squared successive difference between successive responses and as sample variance across the entire observation period. Both of these methods inherently assume homogeneous variability across time. To the best of our knowledge, our current analysis is the first examination of the stochastic volatility of EMA in the context of suicidal risk. To investigate the potential clinical associations and benefits of the fine-resolution time-course measure of variability provided by stochastic volatility, we consider its association with instances



Figure 6: Estimated median posterior volatility of self-reported happiness (curve) along with time points when the participant reported that they wished to be dead (dotted vertical lines).

where the participant reported that they wished to be dead, which are indicated in Figure 7 through dotted vertical lines. We estimated time varying intensity function of events of wishing to be dead by applying Gaussian kernels to each data point and aggregating their contributions using the R function density() from the stats package. For kernel bandwidth selection, we utilized the default rule of thumb method. We then computed Pearson's correlation between this intensity and estimated volatility across time for each subject. The estimated correlation between volatility of happiness and intensity of wishing to be dead is 0.4518 with a p-value less than 0.0001 for subject 1, and 0.3769 with a p-value less than 0.0001 for subject 2.

As self-reported EMA levels of happiness themselves and their means have been shown to be associated with suicide ideation (Kivela et al. 2022), it is important to compare the strengths of association between suicide ideation and volatility of happiness to its associations with reported levels of happiness and mean happiness. The estimated correlation between the intensity of the participant wishing that they were dead and self-reported happiness is 0.0377 with a p-value of 0.6570 for subject 1 and -0.0271 with a p-value of 0.7562 for subject 2. In addition, we applied a generalized cross-validated smoothing spline estimator to the self-reported time series of happiness to obtain a de-noised time-varying estimate of mean happiness. The correlation between this smoothed estimate of mean happiness and the intensity of a participant wishing that they were dead was estimated as 0.0699 with a p-value of 0.4001 for subject 1 and -0.0049 with a p-value of 0.9528 for subject 2. These results, where stochastic volatility has a stronger correlation with a measure of suicide ideation, suggest a potential use in improving monitoring and predicting suicide ideation beyond that provided by reported levels of mood.

8 Discussion

In this paper, we introduced the first method for fitting stochastic volatility models with informed missingness. The approach is built on an imputation algorithm for the stochastic volatility model with informative missingness based on Tukey's representation and extended the conditional particle filter with ancestor sampling algorithm to account for the imputation. Although we focused on the stochastic volatility model in this paper, the conditional particle filter with ancestor sampling algorithm that is the foundation of the proposed method is developed for nonlinear state space models (Lindsten et al. 2015). Thus, the proposed method can potentially be extended to other nonlinear state space models by modifying the imputation step according to the model settings.

This presents the first approach to the analysis of stochastic volatility with informed missingness and considers the simplest case of single realizations of univariate conditionally Gaussian data. It leaves many open questions that can be the focus of future work. The first line of extensions is with regards to the form of responses, including methods for the analysis of multivariate time series and for ordinal time series, which are common forms of EMA responses. Although methods for defining multivariate stochastic volatility models exist, primarily via factor models, how to account for different patterns of missingness among different variables could be challenging. A second line of important extensions is to second-level analyses from multiple subjects. This article considered the estimation from data from a single subject, and used exploratory analyses to evaluate the potential use of volatility as a clinical measure. Future work can extend the single subject model into a multilevel model for multiple subjects, as well as joint models for the simultaneous modeling of stochastic volatility and prediction of time to events.

References

- Andrieu, C., Doucet, A. & Holenstein, R. (2010), 'Particle markov chain monte carlo methods', Journal of the Royal Statistical Society. Series B (Statistical Methodology)
 72(3), 269–342.
- Angst, J., Gamma, A. & Endrass, J. (2003), 'Risk factors for the bipolar and depression spectra', Acta psychiatrica Scandinavica. Supplementum 108, 15–9.
- Carlin, B. P., Polson, N. G. & Stoffer, D. S. (1992), 'A monte carlo approach to nonnormal and nonlinear state-space modeling', *Journal of the American Statistical Association* 87(418), 493–500.
- Chib, S. & Greenberg, E. (1994), 'Bayes inference in regression models with arma (p, q) errors', *Journal of Econometrics* **64**(1), 183–206.
- Cox, R. C., Brown, S. L., Chalmers, B. N. & Scott, L. N. (2023), 'Examining sleep distur-

bance components as near-term predictors of suicide ideation in daily life', *Psychiatry Research* **326**, 115323.

- Doucet, A. & Johansen, A. M. (2011), A tutorial on particle filtering and smoothing: Fifteen years later, in D. Crisan & B. Rozovskii, eds, 'The Oxford Handbook of Nonlinear Filtering', Oxford University Press, New York, pp. 656–704.
- Franks, A. M., Airoldi, E. M. & Rubin, D. B. (2020), 'Nonstandard conditionally specified models for nonignorable missing data', *Proceedings of the National Academy of Sciences* 117(32), 19045–19053.
- Garnett, M. & Curtin, S. (2023), Suicide mortality in the united states, 2001–2021, Technical report, NCHS data brief 464, Hyattsville.
- Gelman, A. (2006), 'Prior distributions for variance parameters in hierarchical models (comment on article by Browne and Draper)', Bayesian Analysis 1(3), 515 – 534.
- Gong, C. & Stoffer, D. S. (2021), 'A note on efficient fitting of stochastic volatility models', Journal of Time Series Analysis 42(2), 186–200.
- Gu, C. (2013), Smoothing Spline ANOVA Models, Springer New York.
- Holland, P. W. (1986), Discussion 4: Mixture modeling versus selection modeling with nonignorable nonresponse, in W. Howard, ed., 'Drawing Inferences from Self-Selected Samples', Springer, New York, p. 143–149.
- Jacquier, E., Polson, N. G. & Rossi, P. E. (1994), 'Bayesian analysis of stochastic volatility models', Journal of Business & Economic Statistics 12(4), 371–389.
- Kim, J. & Stoffer, D. S. (2008), 'Fitting stochastic volatility models in the presence of

irregular sampling via particle methods and the em algorithm', *Journal of Time Series* Analysis **29**(5), 811–833.

- Kim, S., Shephard, N. & Chib, S. (1998), 'Stochastic volatility: Likelihood inference and comparison with arch models', *The Review of Economic Studies* 65(3), 361–393.
- Kivela, L., van der Does, W. A. J., Riese, H. & Antypa, N. (2022), 'Don't miss the moment: A systematic review of ecological momentary assessment in suicide research', *Frontiers in Digital Health* 4(876595).
- Kuppens, P., Mechelen, I., Nezlek, J., Dossche, D. & Timmermans, T. (2007), 'Individual differences in core affect variability and their relationship to personality and adjustment', *Emotion (Washington, D.C.)* 7, 262–74.
- Li, Z., Rosen, O., Ferrarelli, F. & Krafty, R. T. (2021), 'Adaptive bayesian spectral analysis of high-dimensional nonstationary time series', *Journal of Computational and Graphical Statistics* **30**(3), 794–807. PMID: 35936018.
- Lindsten, F., Douc, R. & Moulines, E. (2015), 'Uniform ergodicity of the particle gibbs sampler', Scandinavian Journal of Statistics 42(3), 775–797.
- Lindsten, F., Jordan, M. I. & Schön, T. B. (2014), 'Particle gibbs with ancestor sampling', J. Mach. Learn. Res. 15(1), 2145–2184.
- Little, R. J. A. & Wang, Y. (1996), 'Pattern-mixture models for multivariate incomplete data with covariates', *Biometrics* **52**(1), 98–111.
- McConville, C. & Cooper, C. (1996), 'Mood variability and the intensity of depressive states', *Curr Psychol* 14, 329–338.

- Palmier-Clauss, J., Shryane, N., Taylor, P., Lewis, S. & Drake, R. (2013), 'Mood variability predicts the course of suicidal ideation in individuals with first and second episode psychosis', *Psychiatric Research* 206(3), 240–250.
- Pitt, M. K. & Shephard, N. (1999), 'Filtering via simulation: Auxiliary particle filters', Journal of the American Statistical Association 94(446), 590–599.
- Polson, N. G., Scott, J. G. & Windle, J. (2013), 'Bayesian inference for logistic models using pólya–gamma latent variables', Journal of the American Statistical Association 108(504), 1339–1349.
- Rubin, D. B. (1974), 'Characterizing the estimation of parameters in incomplete-data problems', Journal of the American Statistical Association 69(346), 467–474.
- Shephard, N. (1996), Statistical aspects of ARCH and stochastic volatility, (edited by d.r. cox, david v. hinkley and ole e. barndorff-neilsen) edn, Chapman & Hall, London, pp. 1–67.
- Shephard, N., ed. (2005), Stochastic Volatility: Selected Readings, Oxford University Press.
- Taylor, S. J. (1994), 'Modeling stochastic volatility: A review and comparative study', Mathematical Finance 4(2), 183–204.
- Thompson, R. J., Berenbaum, H. & Bredemeier, K. (2011), 'Cross-sectional and longitudinal relations between affective instability and depression', *Journal of Affective Disorders* 130(1), 53–59.
- Wand, M. P., Ormerod, J. T., Padoan, S. A. & Frühwirth, R. (2011), 'Mean Field Variational Bayes for Elaborate Distributions', *Bayesian Analysis* 6(4), 847 – 900.

Appendix A Imputation procedure for informative missingess

Following the notations introduced in Sections 2 and 4, we provide supplemental details on the derivation of the proposed imputation procedure.

A.1 Details of the Tukey's representation

First note that

$$f(y_t|\theta_{Y|R}, h_{t-1}) = pf(y_t|r_t = 1, \theta_{Y|R}, h_{t-1}) + (1-p)f(y_t|r_t = 0, \theta_{Y|R}, h_{t-1})$$

Thus,

$$r(y_t) = P(r_t = 1 | y_t, \Theta, h_{t-1}) = \frac{pf(y_t | r_t = 1, \theta_{Y|R}, h_{t-1})}{f(y_t | \theta_{Y|R}, h_{t-1})}$$

The logit of $r(y_t)$ is given by

$$\log it \left(\frac{r(y_t)}{1 - r(y_t)}\right) = \log [r(y_t)] - \log [1 - r(y_t)]$$

= $\log \left[\frac{pf(y_t|r_t = 1, \theta_{Y|R}, h_{t-1})}{f(y_t|\theta_{Y|R}, h_{t-1})}\right] - \log \left[\frac{(1 - p)f(y_t|r_t = 0, \theta_{Y|R}, h_{t-1})}{f(y_t|\theta_{Y|R}, h_{t-1})}\right]$
= $\log \left[\frac{p}{1 - p}\right] + \log \left[f(y_t|r_t = 1, \theta_{Y|R}, h_{t-1})\right]$
 $- \log \left[f(y_t|r_t = 0, \theta_{Y|R}, h_{t-1})\right]$ (11)

The equation (11) can be rewritten as

$$f(y_t|r_t = 0, \theta_{Y|R}, h_{t-1}) = \frac{p}{1-p} \frac{1-r(y_t)}{r(y_t)} f(y_t|r_t = 1, \theta_{Y|R}, h_{t-1}).$$
(12)

A.2 Details of the implied distribution for missing data y_t

As introduced in Section 4 of the main article, under the univariate stochastic volatility model setting, we have

$$\frac{1 - r(y_t)}{r(y_t)} = \frac{1}{exp(\beta_0 + \beta_1 y_t)}$$
(13)

$$\frac{p}{1-p} = \exp\left(\beta_0 - \frac{\beta_1 \sigma_t^2}{2}\right) \tag{14}$$

$$y_t | r_t = 1, \Theta, h_{t-1} \sim N(0, \sigma_t^2).$$
 (15)

Applying Tukey's representation 12, the implied distribution for missing y_t can be derived as

$$\begin{split} f(y_t|r_t &= 0, \theta_{Y|R}, h_{t-1}) = \frac{p}{1-p} \frac{1-r(y_t)}{r(y_t)} f(y_t|r_t = 1, \theta_{Y|R}, h_{t-1}) \\ &= \exp\left(\beta_0 - \frac{\beta_1 \sigma_t^2}{2}\right) \frac{1}{exp(\beta_0 + \beta_1 y_t)} \times \frac{1}{\sqrt{2\pi\sigma_t^2}} exp(\frac{-y_t^2}{2\sigma_t^2}) \\ &= \frac{1}{\sqrt{2\pi\sigma_t^2}} exp(\frac{-(y_t + \beta_1 \sigma_t^2)^2}{2\sigma_t^2}) \\ &\sim N(-\beta_1 \sigma_t^2, \sigma_t^2). \end{split}$$

Appendix B Additional details of the Gibbs sampler

For a more comprehensive understanding of the Gibbs sampler, as originally introduced in Section 6 of the main article, we present additional details on sampling σ and ϕ in this section. We provide details for two priors and approaches for sampling σ and ϕ . The first approach assumes independent priors where We used two methods for sampling ϕ, σ^2 . The first method involves sampling ϕ and σ^2 separately.

(a) ϕ

We used a method based on the Metropolis-Hastings algorithm(Chib & Greenberg 1994, Kim et al. 1998).

• prior:

$$\frac{1+\phi}{2} \sim Beta(a,b)$$
$$\pi(\phi) \propto \{\frac{1+\phi}{2}\}^{a-1} \{\frac{1-\phi}{2}\}^{b-1}$$

where a, b are hyperparameters that satisfy $a, b > \frac{1}{2}$.

• posterior and acceptance sampling method: At *i*th iteration, given $\phi^{(i-1)}$ from i - 1th iteration, sample ϕ^* from proposal distribution $N(\hat{\phi}, V_{\phi})$, where $\hat{\phi} = \frac{\sum_{t=1}^{n-1} (h_{t+1}-\mu)(h_t-\mu)}{\sum_{t=1}^{n-1} (h_t-\mu)^2}$ and $V_{\phi} = \frac{\sigma^2}{\sum_{t=1}^{n-1} (h_t-\mu)^2}$. Then if the proposal value is in the stationary region, it will be accepted with probability $exp(g(\phi^*) - g(\phi^{(i-1)}))$, where

$$g(\phi) = \log \pi(\phi) - \frac{(h_1 - \mu)^2 (1 - \phi^2)}{2\sigma^2} + \frac{1}{2} \log(1 - \phi^2).$$

If ϕ^* is rejected, set $\phi^{(i)}$ as $\phi^{(i-1)}$.

(b) σ^2

• prior: A Half-t prior distribution (Gelman 2006, Li et al. 2021) is used for σ

$$\sigma \sim \text{Half-}t(v,G)$$

$$\pi(\sigma) \propto \{1 + \frac{1}{v}(\frac{\sigma}{G})^2\}^{-(v+1)/2}, \sigma > 0$$

where v, G are hyperparameters. This distribution is equivalent to a scale mixture of inverse Gamma distribution(IG) for σ^2 (Wand et al. 2011):

$$\sigma^2 | q_\sigma \sim IG(\frac{v}{2}, \frac{v}{q_\sigma}),$$
$$q_\sigma \sim IG(\frac{1}{2}, \frac{1}{G^2}).$$

• posterior:

$$\sigma^{2}|q_{\sigma}, h_{1:n}, y_{1:n}, \phi, \mu \sim IG(\frac{v+n}{2}, A)$$
$$q_{\sigma}|\sigma^{2}, h_{1:n}, y_{1:n}, \phi, \mu \sim IG(\frac{v+1}{2}, \frac{1}{G^{2}} + \frac{v}{\sigma^{2}})$$

where

$$A = \frac{v}{q_{\sigma}} + \frac{1}{2} \{ (1 - \phi^2)(h_1 - \mu)^2 + \sum_{t=2}^n [(h_t - \mu) - \phi(h_{t-1} - \mu)^2] \}$$

Another method involves sampling ϕ and σ^2 together from posterior distribution $f(\phi, \sigma | \mu, y_{1:T}, h_{1:T})$ using Random Walk Metropolis-Hastings(RWMH) algorithm (Gong & Stoffer 2021). The prior and posterior for (ϕ, σ) are presented below and the Random Walk Metropolis-Hastings(RWMH) algorithm is summarized in the Algorithm 4.

• prior:

$$\begin{pmatrix} \phi \\ \sigma \end{pmatrix} \sim N(\begin{pmatrix} \mu_{\phi} \\ \mu_{\sigma} \end{pmatrix}, \Sigma)$$

where

$$\Sigma = \begin{bmatrix} \sigma_{\phi}^2 & \rho \sigma_{\phi} \sigma_q \\ \\ \rho \sigma_{\phi} \sigma_q & \sigma_q^2 \end{bmatrix}$$

• posterior:

$$f(\phi, \sigma | \mu, y_{1:T}, h_{1:T}) \propto \pi(\phi, \sigma) f(h_1 | \phi, \sigma, \mu) \prod_{i=2}^T f(h_i | h_{i-1}, \phi, \sigma, \mu)$$

Algorithm 4 Random Walk Metropolis-Hastings(RWMH)

1: The initial state $\xi^0 = (\phi, \sigma)$. 2: for $r \ge 0$ do Draw $\epsilon_r \sim N(0, \lambda \Sigma_{\epsilon})$ and let $\eta^r = \xi^{r-1} + \epsilon_r$. 3: Compute the acceptance probability $\alpha = min(\frac{g(\xi)}{g(\xi^r)}, 1)$. 4: 5:Generate $U_r \sim unif(0, 1)$. 6: if $U_r < \alpha$ then Set $\xi^r = \eta^r$ 7: else if $U_r \geq \alpha$ then 8: Set $\xi^r = \xi^{r-1}$ 9: end if 10: 11: end for

where

$$g(\xi) = exp\{-\frac{(\phi - \mu_{\phi})^{2}\sigma_{q}^{2} + (\sigma - \mu_{q})^{2}\sigma_{\phi}^{2} - 2\rho\sigma_{\phi}\sigma_{q}(\phi - \mu_{\phi})(\sigma_{-}\mu_{q})}{2(1 - \rho^{2})\sigma_{\phi}^{2}\sigma_{q}^{2}}\}$$
$$\times \frac{\sqrt{1 - \phi^{2}}}{\sigma^{n}}exp\{-\frac{(1 - \phi^{2})(h_{1} - \mu)^{2} + \sum_{i=2}^{n}[(h_{i} - \mu) - \phi(h_{i-1} - \mu)]^{2}}{2\sigma^{2}}\}$$

Appendix C Additional results for simulation studies

In Section 6 of the main manuscript, we present the posterior results for the stochastic volatility series. In this section, we provide the mean posterior coverage rate and width of 95% credible intervals and mean squared error for model parameters $\Theta = (\sigma, \phi, \mu, \beta_0, \beta_1)$.

The posterior results for parameters for the proposed method and commonly used ad hoc methods are also provided in this section. The AMSEs for model parameters using the proposed method are comparable to those from ad hoc methods.

Parameter	Sample size	$\exp(\beta_1)$	Mean coverage rate	Mean width	MSE
		2.5	0.9580	0.2477	0.0076
	500	3.5	0.9900	0.2643	0.0069
_		4.5	0.9020	0.2428	0.0095
0		2.5	1.0000	0.3144	0.0098
	100	3.5	0.9979	0.3116	0.0102
		4.5	0.9979	0.3123	0.0107
		2.5	0.9920	0.1829	0.0033
	500	3.5	0.9880	0.2263	0.0061
4		4.5	0.9980	0.1931	0.0034
φ		2.5	0.9959	0.2285	0.0057
	100	3.5	1.0000	0.2320	0.0058
		4.5	0.9959	0.2351	0.0058
		2.5	0.9980	0.3995	0.0151
	500	3.5	0.9460	0.3283	0.0127
		4.5	0.9540	0.3502	0.0150
μ		2.5	1.0000	0.8928	0.0689
	100	3.5	1.0000	0.8787	0.0715
		4.5	1.0000	0.8757	0.0743
		2.5	0.8120	1.3196	0.4408
	500	3.5	0.8800	1.0254	0.1921
ß		4.5	0.8560	1.0475	0.2644
\wp_1		2.5	0.8330	1.6966	0.5985
	100	3.5	0.8274	2.1267	0.8541
		4.5	0.8066	1.6917	0.6569

Table 3: Mean posterior coverage rate and width of 95% credible intervals and mean squared error for model parameters under linear missing mechanism

Appendix D Additional results for the application

We analyzed the motivating data using the proposed procedure under the cubic smoothing spline mechanism and the findings are consistent with those under the logistic missing mechanism.

Parameter	Sample size	$\exp(\beta_1)$	Mean coverage rate	Mean width	MSE
		2.5	0.9733	0.2489	0.0067
	500	3.5	0.9868	0.2533	0.0069
σ		4.5	0.9957	0.2568	0.0066
0		2.5	1.0000	0.3216	0.0091
	100	3.5	1.0000	0.3213	0.0090
		4.5	1.0000	0.3258	0.0090
		2.5	0.9756	0.1792	0.0033
	500	3.5	0.9868	0.1797	0.0033
4		4.5	0.9762	0.1818	0.0033
ϕ		2.5	0.9931	0.2290	0.0057
	100	3.5	0.9911	0.2288	0.0057
		4.5	0.9977	0.2314	0.0055
		2.5	1.0000	0.4756	0.0200
	500	3.5	1.0000	0.4793	0.0199
μ -		4.5	1.0000	0.4897	0.0203
		2.5	1.0000	0.9629	0.1021
	100	3.5	1.0000	0.9717	0.0796
		4.5	1.0000	0.9705	0.0836

Table 4: Mean posterior coverage rate and width of 95% credible intervals and mean squared error for model parameters under cubic smoothing spline missing mechanism

				AMSE	
Parameter	Sample size	$\exp(\beta_1)$	Method P	Method A	Method B
		2.5	0.7598	0.9133	0.7891
	500	3	0.7836	1.1203	0.8352
L		3.5	0.8055	1.4925	0.8906
11		2.5	0.7811	0.8738	0.7987
	100	3	0.8130	1.0103	0.8535
		3.5	0.8400	1.1687	0.8981
		2.5	0.0076	0.0133	0.0069
	500	3	0.0069	0.0261	0.0067
_		3.5	0.0095	0.0742	0.0071
0		2.5	0.0098	0.0098	0.0095
	100	3	0.0102	0.0107	0.0096
		3.5	0.0107	0.0117	0.0098
		2.5	0.0033	0.0040	0.0032
	500	3	0.0061	0.0057	0.0030
4		3.5	0.0034	0.0081	0.0032
ϕ		2.5	0.0057	0.0051	0.0059
	100	3	0.0058	0.0049	0.0058
		3.5	0.0058	0.0049	0.0059
		2.5	0.0151	0.0219	0.0160
	500	3	0.0127	0.0431	0.0160
		3.5	0.0150	0.0919	0.0176
μ		2.5	0.0689	0.0718	0.0685
	100	3	0.0715	0.0819	0.0775
		3.5	0.0743	0.1116	0.0739

Table 5: Average mean squared error (AMSE) over time for the stochastic volatility h and model parameters under linear missing mechanism

				AMSE	
Parameter	Sample size	$\exp(\beta_1)$	Method P	Method A	Method B
		2.5	0.7368	0.8005	0.7786
	500	3.5	0.7403	0.8229	0.7913
Ь		4.5	0.7493	0.8394	0.7961
11		2.5	0.7753	0.8183	0.7825
	100	3.5	0.7756	0.8219	0.7856
		4.5	0.7847	0.9061	0.8725
		2.5	0.0067	0.0117	0.0082
	500	3.5	0.0069	0.0146	0.0090
-		4.5	0.0066	0.0166	0.0091
0		2.5	0.0091	0.0109	0.0096
	100	3.5	0.0090	0.0112	0.0094
		4.5	0.0090	0.0172	0.0141
		2.5	0.0033	0.0030	0.0030
	500	3.5	0.0033	0.0029	0.0030
4		4.5	0.0033	0.0030	0.0030
ϕ		2.5	0.0057	0.0055	0.0061
	100	3.5	0.0057	0.0054	0.0061
		4.5	0.0055	0.0051	0.0053
		2.5	0.0200	0.0179	0.0194
	500	3.5	0.0199	0.0185	0.0206
		4.5	0.0203	0.0191	0.0189
μ		2.5	0.1021	0.0855	0.1007
	100	3.5	0.0796	0.0813	0.0963
		4.5	0.0836	0.1151	0.0989

Table 6: Average mean squared error(AMSE) over time for the stochastic volatility h and model parameters under cubic smoothing spline missing mechanism

Table 7: Mean posterior coverage rate and width of 95% credible interval over time and the average mean squared error(AMSE) over time for smoothing spline function

Sample size	$\exp(\beta_1)$	AMSE	Mean width	Mean coverage rate
	2.5	19.8800	7.3030	0.8974
500	3.5	20.4429	6.9856	0.8594
	4.5	22.5088	7.4068	0.8203
	2.5	26.5555	12.3529	0.9732
100	3.5	27.0224	11.9002	0.9647
	4.5	27.6660	12.0874	0.9602



Figure 7: Estimated median posterior volatility of self-reported happiness (curve) along with time points when the participant reported that they wished to be dead (dotted vertical lines).

Table 8: Correlation and corresponding p-values for correlation test between the intensity function and estimated volatility(h), original mood series(y), and spline-smoothed mood series(\tilde{y}).

Subject	Measure	Pearson's correlation	p-value
	h	0.5257	< 0.0001
1	y	0.0377	0.6570
	${ ilde y}$	0.0699	0.4001
	h	0.5252	< 0.0001
2	y	-0.0271	0.7562
	\widetilde{y}	-0.0049	0.9528