# Separable Computation of Information Measures

Xiangxiang Xu and Lizhong Zheng Dept. EECS, MIT Cambridge, MA 02139, USA Email: {xuxx, lizhong}@mit.edu

Abstract—We study a separable design for computing information measures, where the information measure is computed from learned feature representations instead of raw data. Under mild assumptions on the feature representations, we demonstrate that a class of information measures admit such separable computation, including mutual information, *f*-information, Wyner's common information, Gács–Körner common information, and Tishby's information bottleneck. Our development establishes several new connections between information measures and the statistical dependence structure. The characterizations also provide theoretical guarantees of practical designs for estimating information measures through representation learning.

#### I. INTRODUCTION

The computation of information measures, e.g., mutual information, is a fundamental task in information theory [1] and its applications. In machine learning tasks, for example, computing information measures becomes a key step when using information-theoretic tools for algorithm analyses and designs. Due to the unknown statistical model behind practical data, information measures cannot be directly computed from the definitions. The high dimensionality and complicated structures of data also lead to an enormous computation complexity of non-parametric approaches.

Recent developments employed deep learning techniques in designing information measure estimators [2], [3], leveraging the capability of deep neural networks in effectively processing high-dimensional and structured data. For instance, [3] proposed an approach to estimate mutual information by using deep neural networks to extract features from data and then applying a non-parametric mutual information estimator on the extracted features. Despite their performance gains on empirical evaluations, such designs are often heuristic without guarantees, where the learned features may fail to carry useful information for the estimation tasks [3, Sec. 3.3].

In this paper, we formulate the problem of computing information measures from extracted features, which allows separable computation and implementation. We restrict to the bivariate case and consider computing an information measure  $\theta(X, Y)$  from a pair of random variables X and Y, where X or Y correspond to possibly high-dimensional data in practical applications. As illustrated in Fig. 1, the separable computation is conducted by first learning features s(X), t(Y) from X, Y, and then applying  $\theta$ -estimator on the extracted features. Our focus is on establishing the theoretical conditions such that the features contain the necessary information for estimating the  $\theta$ , i.e.,  $\theta(X, Y) = \theta(s(X), t(Y))$ . In particular, we assume that s(X) and t(Y) can be any sufficient statistics of X and Y, i.e.,



Fig. 1. Computing an information measure  $\theta(X, Y)$  in two steps: (1) obtain transformed variables s(X), t(Y) and (2) apply an estimator on s(X), t(Y). For high-dimensional X, Y with unknown probability structures, the first step can be implemented by data-driven approaches, e.g., deep neural network training, corresponding to the feature learning process, where s(X) and t(Y) are the learned feature representations.

we have the Markov relations X-s(X)-Y and X-t(Y)-Y. This assumption allows flexible feature learning designs. Moreover, the statistical sufficiency allows feature extractors to discard irrelevant information while capturing the statistical dependence between X and Y [4]. Under this assumption, we establish several classical information measures admitting the separable computation, including mutual information and its variants using *f*-divergence, Wyner's common information [5], Gács–Körner common information [6], and Tishby's information bottleneck [7]. Our results provide theoretical guarantees of modular designs in applying deep neural networks for information measure estimations, which can lead to more systematic designs and improve implementation efficiency.

#### **II. PRELIMINARIES AND NOTATIONS**

For a random variable Z, we denote the corresponding alphabet by  $\mathcal{Z}$ , use z to denote a specific value in  $\mathcal{Z}$ , and use  $P_Z$  to denote the probability distribution. Throughout our development, we restrict to a pair of discrete random variables X, Y on finite alphabets  $\mathcal{X}, \mathcal{Y}$ . Without loss of generality, we assume  $P_X$  and  $P_Y$  have positive probability masses, i.e.,  $P_X(x) > 0$  for all  $x \in \mathcal{X}$  and  $P_Y(y) > 0$  for all  $y \in \mathcal{Y}$ , since otherwise we can remove the zero-probability symbols from the alphabets. For  $n \geq 1$ , we denote  $[n] \triangleq \{1, 2, ..., n\}$ .

# A. Canonical Dependence Kernel and Modal Decomposition

Given  $(X, Y) \sim P_{X,Y}$  the associated *canonical dependence* kernel (CDK) function [8] is defined as a joint function  $i_{X;Y}$ :

$$\mathfrak{i}_{X;Y}(x,y) \triangleq \frac{P_{X,Y}(x,y)}{P_X(y)P_Y(y)} - 1. \tag{1}$$

By applying the singular value decomposition (SVD) in the space of joint function, we can write CDK  $i_{X;Y}$  as a superposition of rank-one singular modes, referred to as its modal decomposition [9]:

$$i_{X;Y}(x,y) = \sum_{i=1}^{K} \sigma_i \cdot f_i^*(x) g_i^*(y),$$
(2)

In particular,  $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_K > 0$  are the singular values, where  $K \ge 0$  denotes the rank of  $i_{X;Y}$ . Analogous to the orthogonality of singular vectors in matrix SVD,  $f_i^*, g_i^*$ satisfy  $\mathbb{E}\left[f_i^*(X)f_j^*(X)\right] = \mathbb{E}\left[g_i^*(Y)g_j^*(Y)\right] = \delta_{i,j}$ , for all  $i, j \in [K]$ . For convenience, we also define functions

$$f^* \triangleq (f_1^*, \dots, f_K^*)^{\mathrm{T}}, \quad g^* \triangleq (g_1^*, \dots, g_K^*)^{\mathrm{T}}. \tag{3}$$

For convenience, we further define  $\sigma_0 \triangleq 1, f_0^* \colon \mathfrak{X} \ni x \mapsto 1$ and  $g_0^* \colon \mathcal{Y} \ni y \mapsto 1$ . Then, we can write the SVD of the density ratio as [cf. (1)]:  $\frac{P_{X,Y}(x,y)}{P_X(x)P_Y(y)} = \sum_{i=0}^{K} \sigma_i \cdot f_i^*(x)g_i^*(y).$ 

# B. Sufficiency and Minimal Sufficiency

Definition 1: Given X, Y, S = s(X) is a sufficient statistic of X for inferring Y if we have the Markov relation X – s(X) - Y. We call f(X) a minimal sufficient statistic if f(X)is sufficient, and for each given sufficient statistic S of X, f(X) is a function of S.

We have the following useful characterization of sufficient statistics. A proof is provided in Section V-A.

Proposition 1: Given X, Y and S = s(X), T = t(Y), the following statements are equivalent:

- X S Y, X T Y;
- $\begin{aligned} X S T Y; \\ P_{X,Y}(x,y) \\ P_X(x)P_Y(y) \end{aligned} = \frac{P_{S,T}(s(x), t(y))}{P_S(s(x))P_T(t(y))} \text{ for all } x \in \mathfrak{X}, y \in \mathcal{Y}. \end{aligned}$

From such equivalences, the Markov relation X - S - T - Yis equivalent to X - S - Y and X - T - Y, i.e., both S and T are sufficient statistics.

In addition, the following result demonstrates a useful connection between minimal sufficiency and the modal decomposition of CDK.

*Proposition 2 ([4, Proposition 2]):* The features  $f^*(X), q^*(Y)$  as defined by (2)–(3) are minimal sufficient statistics of X and Y.

### **III. MAIN RESULTS**

Throughout our development in the paper, we will assume that S = s(X), T = t(Y) are sufficient statistics of given X, Y, i.e., X - s(X) - t(Y) - Y. We demonstrate several common examples of information measures  $\theta(X, Y)$  that can be evaluated from S and T, i.e.,  $\theta(X, Y) = \theta(S, T)$ .

# A. Mutual Information and f-Information

It is straightforward to verify that the mutual information

$$I(X;Y) \triangleq \sum_{x \in \mathfrak{X}, y \in \mathfrak{Y}} P_{X,Y}(x,y) \log \frac{I_{X,Y}(x,y)}{P_X(x)P_Y(y)}$$

satisfies I(X;Y) = I(S;T). To see this, note that we have S - X - Y - T since S and T are deterministic functions of X and Y, respectively. Then, the data processing inequality [cf. Lemma 1] implies that  $I(S;T) \leq I(X;Y)$ . Also, the sufficiency of S and T implies that X - S - T - Y and  $I(X;Y) \le I(S;T).$ 

We can extend the above result to the general class of finformation, defined as

$$I_f(X;Y) \triangleq \sum_{\substack{x \in \mathcal{X} \\ y \in \mathcal{Y}}} P_X(x) P_Y(y) \cdot f\left(\frac{P_{X,Y}(x,y)}{P_X(x)P_Y(y)}\right), \quad (4)$$

where f is a convex function defined on  $[0, \infty)$  with f(1) = 0. In particular the mutual information I(X;Y) corresponds to the specific  $f(u) = u \cdot \log u$ . Then we have the following characterization, as a straightforward corollary of Proposition 1. We omit the proof.

Corollary 1: For each f-information defined by (4), we have  $I_f(X;Y) = I_f(S;T).$ 

# B. Common Information

We consider the common information between a pair of random variables formulated by Wyner [5] and the common information introduced by Gács and Körner [6].

1) Wyner's Common Information: Given X, Y, Wyner's common information [5] between X and Y is defined as

$$C(X,Y) \triangleq \min_{P_{W|X,Y}: X-W-Y} I(W;X,Y).$$
(5)

Then, we have the following characterization. A proof is provided in Section V-B.

Theorem 1: Suppose S = s(X) and T = t(Y) are sufficient. Then, C(X, Y) = C(S, T), and the optimal W in (5) satisfies W - (S,T) - (X,Y).

2) Gács-Körner Common Information: Different from Wyner's notion, Gács-Körner common information [6] is

$$C_{\mathrm{GK}}(X,Y) \triangleq \max_{\substack{f,g: \mathbb{P}\{f(X)=g(Y)\}=1}} H(f(X)), \quad (6)$$

where  $f: \mathfrak{X} \to \mathcal{W}$  and  $g: \mathcal{Y} \to \mathcal{W}$  map X and Y to some common alphabet  $\mathcal{W}$ , and where  $H(\cdot)$  denotes Shannon entropy. Detailed analyses and properties of  $C_{GK}(X, Y)$  can also be found in, e.g., [10] and [8, Section 5.9].

Here, we establish that  $C_{GK}(X, Y)$  can be computed from sufficient statistics. A proof is provided in Section V-C.

Theorem 2: We have  $C_{GK}(X,Y) = C_{GK}(S,T) =$  $H(f_0^*(X), \ldots, f_k^*(X))$ , where we have defined  $k \triangleq \max\{0 \le 1\}$  $i \leq K$ :  $\sigma_i = 1$ , and where  $\sigma_i$ 's and K are defined by the modal decomposition (2).

#### C. Information Bottleneck

Given X, Y, the information bottleneck problem [7] investigates the description of X by a new variable U, such that U captures as much of the information about Y as possible. This can be formalized as solving the maximal I(U;Y) under the constraint  $I(U; X) \leq R$  for some  $R \geq 0$ , where U is specified by  $P_{U|X}$ . This is equivalent to computing

$$\vartheta_{X,Y}(R) \triangleq \max_{\substack{P_{U|X}: \ U-X-Y\\I(U;X) < R}} I(U;Y).$$
(7)

From the data processing inequality, we have  $\vartheta_{X,Y}(R) \leq$ I(X; Y). The equality can be achieved, e.g., for any S = s(X)satisfy X - S - Y and R = H(S).

*Remark 1:* The information bottleneck formulation is a specific example of rate-distortion theory by employing a KL divergence as the distortion function [7]. The related quantities also appear in other information-theoretic studies. For example,  $\vartheta_{X,Y}(R)$  is the optimal error exponent of a distributed hypothesis testing problem (cf. [11, Theorem 3]), where *R* corresponds to a communication rate constraint.

The Lagrangian  $I(U; X) - \beta \cdot I(U; Y)$  was introduced in [7] to characterize the tradeoff between I(U; Y) and I(U; X), where  $\beta > 0$  is the Lagrange multiplier. Specifically, let us denote its minimum value as

$$L_{\mathrm{IB}}^{*}(X,Y;\beta) \triangleq \min_{P_{U|X}: |U-X-Y|} \left[ I(U;X) - \beta \cdot I(U;Y) \right].$$
(8)

Compared with previous examples of information measures,  $\vartheta(\cdot)$  and  $L_{\mathrm{IB}}^*$  are generally not symmetric with respect to X and Y, and the results are parameterized functions instead of single scalar-valued measures. However,  $\vartheta(\cdot)$  and  $L_{\mathrm{IB}}^*$  admit the same separable computation property, as demonstrated below. A proof is provided in Section V-D.

Theorem 3: Given X, Y, suppose S = s(X), T = t(Y)are their sufficient statistics. Then, for all  $\beta > 0$ , we have  $L^*_{IB}(X,Y;\beta) = L^*_{IB}(S,T;\beta)$ , and the optimal U that achieves  $L^*_{IB}(X,Y;\beta)$  satisfies U-S-X-Y. In addition,  $\vartheta_{X,Y}(R) = \vartheta_{S,T}(R)$  for all  $R \ge 0$ .

## **IV. DISCUSSIONS**

We demonstrated the separable design for estimating information measure  $\theta$ , where  $\theta$  satisfies employing  $\theta(X, Y) = \theta(S, T)$  for sufficient statistics S and T of X and Y. This separability enables modular designs and often more efficient implementations. A practical choice of S and T can be the  $f^*(X)$  and  $g^*(Y)$  defined by the modal decomposition [cf. (2)]. The detailed design of learning  $f^*(X)$  and  $g^*(Y)$  from the X, Y samples was discussed in [9]. More general separable designs for feature-centric information processing systems, including multivariate cases, were also studied in [9].

Our developments are also deeply related to the statistical dependence. For example, the relation  $\theta(X, Y) = \theta(S, T)$ , i.e., the information measure  $\theta$  is invariance to the choices of sufficient statics S and T, also appeared as a key criterion in characterizing the statistical dependence [4]. In addition, [8] demonstrated that the features  $f^*(X), g^*(Y)$ , referred to as the universal features, are optimal for a large class of inference problems. In particular, under assumptions of weak dependence or Gaussian dependence, [8] developed analytical expressions of information measures in terms of  $f^*(X), g^*(Y)$ .

#### V. PROOFS

We first introduce several useful results. The first is the data processing inequality, a proof of which can be found in, e.g., [12, Theorem 2.8.1].

Lemma 1: For X, Y, Z with the Markov relation X - Y - Z,  $I(X; Z) = I(X; Y) - I(X; Y|Z) \le I(X; Y)$ .

Lemma 2: Given X, U taking values from  $\mathfrak{X} \times \mathfrak{U}$  with joint distribution  $P_{X,U}$ , define  $V \triangleq v(X)$  for a function v on  $\mathfrak{X}$ , and define a new random variable U' on  $\mathfrak{U}$  specified by

 $P_{U'|X}(u|x) \triangleq P_{U|V}(u|v(x))$  for all  $u \in \mathcal{U}, x \in \mathcal{X}$ . Then we have U' - V - X and  $P_{U',V} = P_{U,V}$ .

*Proof:* Since  $P_{U'|X}(u|x)$  depends on x only through v(x), we obtain U'-V-X, i.e., V is a sufficient statistic of X to infer U'. In addition, from  $P_{U'|V}(u|v(x)) = P_{U|V}(u|v(x))$ , we obtain  $P_{U'|V} = P_{U|V}$ . This implies that  $P_{U'} = P_U$  and thus  $P_{U',V} = P_{U,V}$ .

Lemma 3: If U, X, Y, Z satisfy U - X - Y, U - (X, Y) - Z, and X - Z - Y, then we have U - X - Z - Y. *Proof*: Omitted.

A. Proof of Proposition 1

First, note that "2"  $\implies$  "1" can be readily obtained by taking corresponding marginal distributions. It suffices to prove that "1"  $\implies$  "3"  $\implies$  "2".

To establish "1"  $\implies$  "3", from X - S - Y we have  $P_{X,Y}(x,y) = P_{Y|X}(y|x) = P_{Y|S}(y|s(x))$ 

$$P_X(x)P_Y(y) \stackrel{-}{\longrightarrow} P_Y(y) \stackrel{-}{\longrightarrow} P_Y(y)$$
,  
ch depends on x only through  $s(x)$ . Similarly, from

which depends on x only through s(x). Similarly, from X - T - Y we obtain

$$\frac{P_{X,Y}(x,y)}{P_X(x)P_Y(y)} = \frac{P_{X|T}(x|t(y))}{P_X(x)}$$

which depends on y only through t(y). Therefore, there exists a function  $\tau: S \times T \to \mathbb{R}$ , such that

$$\frac{P_{X,Y}(x,y)}{P_X(x)P_Y(y)} = \tau(s(x), t(y)).$$
(9)

This implies that for any  $s_0 \in S, t_0 \in T$ ,

$$P_{S,T}(s_0, t_0) = \sum_{x: \ s(x)=s_0} \sum_{y: \ t(y)=t_0} P_{X,Y}(x, y)$$
$$= P_S(s_0) P_T(t_0) \cdot \tau(s_0, t_0).$$

From (9) we obtain "3", since

$$\begin{split} \frac{\dot{P}_{X,Y}(x,y)}{P_X(x)P_Y(y)} &= \tau(s(x),t(y)) = \frac{P_{S,T}(s(x),t(y))}{P_S(s(x))P_T(t(y))}.\\ \text{To establish "3" } \implies "2", \text{ note that} \\ P_{X,Y,S,T}(x,y,s(x),t(y)) \\ &= P_{X,Y}(x,y) \\ &= P_X(x)P_Y(y) \cdot \frac{P_{S,T}(s(x),t(y))}{P_S(s(x))P_T(t(y))} \\ &= \frac{P_X(x)}{P_S(s(x))} \cdot \frac{P_Y(y)}{P_T(t(y))} \cdot P_{S,T}(s(x),t(y)) \\ &= P_{X|S}(x|s(x)) \cdot P_{S,T}(s(x),t(y)) \cdot P_{Y|T}(y|t(y)), \\ \text{which gives } X - S - T - Y. \end{split}$$

#### B. Proof of Theorem 1

We first introduce a useful lemma.

Lemma 4: Given X, Y, and suppose S = s(X) and T = t(Y) are sufficient. Then, for each random variable W with X - W - Y specified by the alphabet W and the conditional distribution  $P_{W|X,Y}$ , we can construct W' specified by  $P_{W'|X,Y}$  such that: W' takes values from W,

• 
$$(X,Y) - (S,T) - W', P_{W'|S,T} = P_{W|S,T};$$
  
•  $X - S - W' - T - Y;$   
 $I(W';X,Y) = I(W;S,T)$   
 $= I(W;X,Y) - I(W;X,Y|S,T).$  (10)

Proof of Lemma 4: We construct W' on  $\mathcal{W}$  as  $P_{W'|X,Y}(w|x,y) \triangleq P_{W|S,T}(w|s(x),t(y))$ , for all  $x \in \mathfrak{X}, y \in \mathcal{Y}$ . From Lemma 2, we obtain  $P_{W,S,T} = P_{W',S,T}$  and

$$' - (S,T) - (X,Y).$$
 (11)

We then verify that X - S - W' - T - Y, i.e.,  $P_{X,Y,S,T|W'}(x, y, s(x), t(y)|w)$ 

W

$$= P_{X|S}(x|s(x))P_{S|W'}(s(x)|w)P_{T|W'}(t(y)|w)P_{Y|T}(y|t(y))$$
  
To see this, note that

 $P_{X,Y,S,T|W'}(x, y, s(x), t(y)|w) = P_{X,Y|W'}(x, y|w) = \frac{P_{W'|X,Y}(w|x, y)P_{X,Y}(x, y)}{P_{W'}(w)}$ (12)

$$=\frac{P_{W|S,T}(w|s(x),t(y))P_{X,Y}(x,y)}{P_{W}(w)}$$
(13)

$$= P_{S,T|W}(s(x), t(y)|w) \cdot \frac{P_{X,Y}(x,y)}{P_{S,T}(s(x), t(y))}.$$
 (14)

To simplify (14), note that due to X - W - Y, we have the Markov relation S - W - T and thus

 $P_{S,T|W}(s(x),t(y)|w) = P_{S|W}(s(x)|w) \cdot P_{T|W}(t(y)|w).$  (15) In addition, from the sufficiency of S and T, we have X - S - T - Y. Thus, from Proposition 1, we have

$$\frac{P_{X,Y}(x,y)}{P_{S,T}(s(x),t(y))} = \frac{P_X(x)P_Y(y)}{P_S(s(x))P_T(t(y))} = P_{X|S}(x|s(x)) \cdot P_{Y|T}(y|t(y)).$$
(16)

Combining (15) and (16), we can rewrite (14) as

 $P_{X,Y,S,T|W'}(x,y,s(x),t(y)|w)$ 

$$= P_{S|W}(s(x)|w) \cdot P_{T|W}(t(y)|w)$$
  
 
$$\cdot P_{X|S}(x|s(x)) \cdot P_{Y|T}(y|t(y))$$
  
 
$$= \left[P_{X|S}(x|s(x)) \cdot P_{S|W'}(s(x)|w)\right]$$
  
(17)

$$\left[P_{T|W'}(t(y)|w) \cdot P_{Y|T}(y|t(y))\right], \qquad (18)$$

where the last equality follows from that due to  $P_{S,T|W} = P_{S,T|W'}$ , we have  $P_{S|W} = P_{S|W'}, P_{T|W} = P_{T|W'}$ .

Finally, (10) follows from

$$I(W'; X, Y) = I(W'; S, T)$$
(19)

$$=I(W;S,T) \tag{20}$$

$$= I(W; X, Y) - I(W; X, Y|S, T),$$
(21)

where (19) follows from (11), (20) follows from the fact that  $P_{W',S,T} = P_{W,S,T}$ , and (21) follows from the Markov relation W - (X,Y) - (S,T) and Lemma 1.

Proceeding to our proof of Theorem 1, from Lemma 4, for each W with X - W - Y, we can construct W' specified by  $P_{W'|S,T}$  with W' - (S,T) - (X,Y) and X - S - W' - T - Y (which implies X - W' - Y), such that I(W';X,Y) =I(W;S,T) = I(W;X,Y) - I(W;X,Y|S,T). Therefore, the optimal W satisfies I(W;X,Y|S,T) = 0, i.e., W - (X,Y) - (S,T), since otherwise W' is strictly better than W. Thus,

$$C(X,Y) = \min_{\substack{P_{W|X,Y}: X - W - Y \\ P_{W|S,T}: W - (S,T) - (X,Y) \\ X - S - W - T - Y}} I(W; S,T).$$
(22)

In addition, for each  $P_{W|S,T}$  such that S - W - T, we can construct  $\overline{W}$  taking values from W, such that  $P_{\overline{W}|X,Y}(w|x,y) = P_{W|S,T}(w|s(x),t(y))$ . Then, we have  $\overline{W} - (S,T) - (X,Y)$  and  $P_{\overline{W},S,T} = P_{W,S,T}$ . This implies that  $S - \overline{W} - T$  and  $I(W;S,T) = I(\overline{W};S,T)$ . Note that we also have  $X - S - \overline{W} - T - Y$  since  $P_{X,Y,C,T|\overline{W}}(x,y,s(x),t(y)|w)$ 

$$\begin{aligned} &P_{X,Y,S,T|\bar{W}}(x,y,s(x),t(y)|w) \\ &= P_{X,Y|S,T,\bar{W}}(x,y|s(x),t(y),w)P_{S,T|\bar{W}}(s(x),t(y)|w) \\ &= P_{X,Y|S,T}(x,y|s(x),t(y))P_{S,T|\bar{W}}(s(x),t(y)|w) \\ &= P_{X,Y|S,T}(x,y|s(x),t(y))P_{S,T|\bar{W}}(s(x),t(y)|w) \end{aligned}$$

 $= P_{X|S}(x|s(x))P_{Y|T}(y|t(y))P_{S|\bar{W}}(s(x)|w)P_{T|\bar{W}}(t(y)|w),$ where the second equality follows from  $\bar{W} - (S,T) - (X,Y)$ , and where the last equality follows from X - S - T - Y and  $S - \bar{W} - T$ . Therefore, we obtain

$$C(S,T) = \min_{\substack{P_{W|S,T}: S - W - T \\ P_{\bar{W}|S,T}: \bar{W} - (S,T) - (X,Y) \\ S - \bar{W} - T}} I(\bar{W}; S,T)$$

$$= \min_{\substack{P_{\bar{W}|S,T}: \bar{W} - (S,T) - (X,Y) \\ X - S - \bar{W} - T - Y}} I(\bar{W}; S,T).$$
(23)

Combining (22) and (23), we have C(X, Y) = C(S, T).

# C. Proof of Theorem 2

S

From the modal decomposition (2), we define  $\hat{f}$  and  $\hat{g}$  as  $\hat{f}(x) \triangleq (f_0^*(x), \ldots, f_k^*(x))^{\mathrm{T}}, \hat{g}(y) \triangleq (g_0^*(y), \ldots, g_k^*(y))^{\mathrm{T}}$ . We first demonstrate that, for *d*-dimensional  $f = (f_1, \ldots, f_d)^{\mathrm{T}}$  and  $g = (g_1, \ldots, g_d)^{\mathrm{T}}$ ,  $\mathbb{P}\{f(X) = g(Y)\} = 1$  if and only if  $f(x) = A\hat{f}(x)$  and  $g(y) = A\hat{g}(y)$  for some  $A \in \mathbb{R}^{d \times (k+1)}$ . Note that  $\mathbb{P}\{f(X) = g(Y)\} = 1$  is equivalent to  $\mathbb{P}\{f_i(X) = g_i(Y)\} = 1$  for all  $i = 1, \ldots, d$ . Therefore, it suffices to consider the case d = 1.

To begin, we can uniquely decompose f and g as

$$f = \sum_{i=0}^{K} \alpha_i \cdot f_i^* + r_f, \quad g = \sum_{i=0}^{K} \beta_i \cdot g_i^* + r_g,$$
  
uch that  $\mathbb{E}[r_f(X)f_i^*(X)] = \mathbb{E}[r_g(Y)g_i^*(Y)] = 0$  for all  $i = K$ . Then, it sufficient to prove that

 $0, \dots, K$ . Then, it suffices to prove that  $\alpha_i = \beta_i$  for i = 0 k (24a)

$$\alpha_i = \beta_i \quad \text{for } i = 0, \dots, n, \quad (24a)$$

$$\alpha_i = \beta_i = 0 \quad \text{for } i = k+1, \dots, K, \tag{24b}$$

$$r_f = 0, \quad r_g = 0.$$
 (24c)

To this end, note that from the orthogonality between functions,  $P \left[ \left( f(X) \right) - \left( X \right) \right]^2$ 

$$\mathbb{E}\left[ (f(X) - g(Y))^2 \right] \\= \mathbb{E}\left[ \left( \sum_{i=0}^K (\alpha_i f_i^*(Y) - \beta_i g_i^*(Y)) + r_f(X) - r_g(Y) \right)^2 \right] \\= \sum_{i=0}^K \mathbb{E}\left[ (\alpha_i \cdot f_i^*(Y) - \beta_i \cdot g_i^*(Y))^2 \right] \\+ \mathbb{E}\left[ r_f^2(X) \right] + \mathbb{E}\left[ r_g^2(Y) \right] \\= \sum_{i=0}^K (\alpha_i^2 - 2\sigma_i \cdot \alpha_i \beta_i + \beta_i^2) + \mathbb{E}\left[ r_f^2(X) \right] + \mathbb{E}\left[ r_g^2(Y) \right] \\= \sum_{i=0}^k (\alpha_i - \beta_i)^2 + \sum_{i=k+1}^K (1 - \sigma_i)(\alpha_i^2 + \beta_i^2)$$

+ 
$$\sum_{i=k+1}^{K} \sigma_i (\alpha_i - \beta_i)^2 + \mathbb{E} \left[ r_f^2(X) \right] + \mathbb{E} \left[ r_g^2(Y) \right].$$
 (25)

Since  $\mathbb{P}\left\{f(X) = g(Y)\right\} = 1$  implies  $\mathbb{E}\left[(f(X) - g(Y))^2\right] = 0$ , from (25) we obtain the conditions (24).

As a result, if  $\mathbb{P}\left\{f(X) = g(Y)\right\} = 1$ , then  $H(f(X)) = H(\hat{f}(X)) \leq H(\hat{f}(X)) = H(f_0^*(X), \dots, f_k^*(X))$ , where the equality holds if  $f = \hat{f}$ . Therefore, we obtain

$$C_{\rm GK}(X,Y) = \max_{f,g: \ \mathbb{P}\{f(X)=g(Y)\}=1} H(f(X))$$
  
=  $H(f_0^*(X), \dots, f_k^*(X)).$  (26)

From Proposition 1, we have

$$_{S;T}(s(x), t(y)) = \sum_{i=1}^{K} \sigma_i \cdot f_i^*(x) \cdot g_i^*(y),$$

which is the modal decomposition of  $i_{S,T}$ . Moreover, from the minimal sufficiency of  $f^*(X), g^*(Y)$  [cf. Proposition 2],  $f_i^*(x)$  is a function of s(x), and  $g_i^*(y)$  is a function of t(y), for each  $i = 0, \ldots, K$ . As a result, it follows from (26) that  $C_{\text{GK}}(S,T) = H(f_0^*(X), \ldots, f_k^*(X)) = C_{\text{GK}}(X,Y)$ .

# D. Proof of Theorem 3

We first introduce a useful lemma.

Lemma 5: Given X, Y, and suppose S = s(X) satisfies X - s(X) - Y. Then, for each random variable U with U - X - Y specified by the alphabet  $\mathcal{U}$  and the conditional distribution  $P_{U|X}$ , we can construct U' such that takes values from  $\mathcal{U}$  and satisfies U' - S - X - Y,  $P_{U',S} = P_{U,S}$ ,  $P_{U',Y} = P_{U,Y}$ , and I(U'; X) = I(U; X) - I(U; X|S), I(U'; Y) = I(U; Y).

Proof of Lemma 5: We construct U' on  $\mathcal{U}$  such that  $P_{U'|X,Y}(u|x,y) = P_{U|S}(u|s(x))$ . Then, it follows from Lemma 2 that U' - S - (X, Y) and  $P_{U',S} = P_{U,S}$ . Since we also have S - X - Y, we obtain  $P_{U',S,X,Y} = P_{U',S}P_{X,Y|S} = P_{U',S}P_{X|S}P_{Y|X}$ , i.e., U' - S - X - Y.

Moreover, from S - X - U and Lemma 1, we have

$$I(U;X) = I(U;X|S) + I(U;S).$$
(27)

Similarly, we have

$$I(U';X) = I(U';X|S) + I(U';S) = I(U';S),$$
(28)  
where the last equality follows from  $U' - S - X$ .

From (27), (28) and  $P_{U,S} = P_{U',S}$ , we obtain I(U';X) = I(U';S) = I(U;S) = I(U;X) - I(U;X|S).

It remains only to establish  $P_{U',Y} = P_{U,Y}$ . To this end, note that from U - X - Y, U - (X, Y) - S, and X - S - Y, from Lemma 3 we obtain the Markov relation U - X - S - Y. Therefore, we have U - S - Y and

$$\begin{split} P_{U,Y}(u,y) &= \sum_{s \in \mathbb{S}} P_{Y|S}(y|s) P_{U,S}(u,s). \\ \text{In addition, due to } U' - S - X - Y, \text{ we have } U' - S - Y \text{ and} \\ P_{U',Y}(u,y) &= \sum_{s \in \mathbb{S}} P_{Y|S}(y|s) P_{U',S}(u,s). \end{split}$$

From  $P_{U',S} = P_{U,S}$ , we have  $P_{U',Y} = P_{U,Y}$  and I(U';Y) = I(U;Y).

Proceeding to the proof of Theorem 3, we first show that

$$L_{\rm IB}^*(X, Y; \beta) = L_{\rm IB}^*(S, Y; \beta) = L_{\rm IB}^*(S, T; \beta).$$
(29)

To establish the first equality, note that the optimal U that achieves  $L_{IB}^*(X, Y; \beta)$  in (8) satisfies U - S - X. Otherwise,

I(U; X|S) > 0 and from Lemma 5, we can construct U' on the same alphabet  $\mathcal U$  with U' - S - X - Y and

$$I(U';X) = I(U;X) - I(U;X|S) < I(U;X),$$
(30a)  
$$I(U';Y) = I(U;Y).$$
(30b)

This implies that  $I(U'; X) - \beta I(U; Y) < I(U; X) - \beta I(U; Y)$ , which contradicts the optimality of U.

Note that since we also have U - X - Y and X - (U, Y) - S, from Lemma 3, U - S - X implies U - S - X - Y. Therefore,  $L_{IB}^*(X, Y; \beta) = \min_{P_{U|X}: |U - X - Y} [I(U; X) - \beta \cdot I(U; Y)]$  $= \min_{P_{U|S}: |U - S - X - Y} [I(U; S) - \beta \cdot I(U; Y)]$  $= \min_{P_{U|S}: |U - S - Y} [I(U; S) - \beta \cdot I(U; Y)]$  $= L_{IB}^*(S, Y; \beta),$ 

where the second equality follows from that the optimal U satisfies U - S - X - Y, and thus (cf. Lemma 1) I(U; X) = I(U; S) + I(U; X|S) = I(U; S). To obtain the third equality, note that for either U - S - X - Y or U - S - Y, I(U; S) and I(U; Y) are solely determined by  $P_{U|S}$ . Finally, the first and the last equalities follow from the definition (8).

To establish the second equality of (29), we note that U - S - Y implies U - S - T - Y and I(U;Y) = I(U;T). Indeed, from U - S - Y, U - (S,Y) - T, and S - T - Y (due to the statistical sufficiency of S, T), we obtain U - S - T - Yby applying Lemma 3. Therefore, we have U - T - Y and U - Y - T, which implies that I(U;Y) = I(U;T). Therefore,  $L_{\text{TB}}^*(S,Y;\beta) = \min [I(U;S) - \beta \cdot I(U;Y)]$ 

$$B(T, \beta) = \lim_{P_{U|S}: |U-S-Y|} [I(U, \beta) - \beta \cdot I(U, T)]$$
  
= 
$$\min_{P_{U|S}: |U-S-Y-T|} [I(U; S) - \beta \cdot I(U; T)]$$
  
= 
$$\min_{P_{U|S}: |U-S-T|} [I(U; S) - \beta \cdot I(U; T)],$$

where the last equality follows from the fact that I(U;S) and I(U;T) depend only on  $P_{U|S}$ .

Similarly, we obtain  

$$\vartheta_{X,Y}(R) \triangleq \max_{\substack{P_{U|X}: |U-X-Y\\I(U,Y) \leq R}} I(U;Y)$$
(31)

$$= \max_{\substack{P_{U|S}: |U-S-X-Y\\I(U:S) < R}} I(U;Y)$$
(32)

$$= \max_{\substack{P_{U|S}: U-S-Y\\I(U;S) \le R}} I(U;Y)$$
(33)

$$= \max_{\substack{P_{U|S}: |U-S-Y-T\\I(U;S) \le R}} I(U;T)$$
(34)

$$= \max_{\substack{P_{U|S}: U-S-T\\I(U:S) \leq R}} I(U;T) = \vartheta_{S,T}(R)$$
(35)

where (32) follows from that it is without loss of generality to restrict U to U - S - X - Y [cf. (30)], which satisfies I(U;X) = I(U;S). (34) follows from that U - S - Y implies I(U;Y) = I(U;T). To obtain (33) and the first equality of (35) we have dropped variables in the middle of Markov chains, as the objective functions (I(U;Y) or I(U;T)) are uniquely determined by  $P_{U|S}$ .

#### References

- [1] C. E. Shannon, "Prediction and entropy of printed english," *Bell system technical journal*, vol. 30, no. 1, pp. 50–64, 1951.
- [2] M. I. Belghazi, A. Baratin, S. Rajeshwar, S. Ozair, Y. Bengio, A. Courville, and D. Hjelm, "Mutual information neural estimation," in *International conference on machine learning*. PMLR, 2018, pp. 531–540.
- [3] G. Gowri, X. Lun, A. M. Klein, and P. Yin, "Approximating mutual information of high-dimensional variables using learned representations," in *The Thirty-eighth Annual Conference on Neural Information Processing Systems*, 2024. [Online]. Available: https://openreview.net/forum?id=HN05DQxyLl
- [4] X. Xu and L. Zheng, "Dependence induced representations," in 2024 60th Annual Allerton Conference on Communication, Control, and Computing. IEEE, 2024, pp. 1–8.
  [5] A. Wyner, "The common information of two dependent random vari-
- [5] A. Wyner, "The common information of two dependent random variables," *IEEE Transactions on Information Theory*, vol. 21, no. 2, pp. 163–179, 1975.
- [6] P. Gács, and J. Körner, "Common information is far less than mutual information." *Problems of Control and Information Theory*, vol. 2, pp. 149–162, 1973.
- [7] N. Tishby, F. C. Pereira, and W. Bialek, "The information bottleneck method," arXiv preprint physics/0004057, 2000.
- [8] S.-L. Huang, A. Makur, G. W. Wornell, and L. Zheng, "Universal features for high-dimensional learning and inference," *Foundations and Trends® in Communications and Information Theory*, vol. 21, no. 1-2, pp. 1–299, 2024. [Online]. Available: http://dx.doi.org/10.1561/0100000107
- [9] X. Xu and L. Zheng, "Neural feature learning in function space," *Journal of Machine Learning Research*, vol. 25, no. 142, pp. 1–76, 2024.
- [10] H. S. Witsenhausen, "On sequences of pairs of dependent random variables," *SIAM Journal on Applied Mathematics*, vol. 28, no. 1, pp. 100–113, 1975.
- [11] R. Ahlswede and I. Csiszár, "Hypothesis testing with communication constraints," *IEEE transactions on information theory*, vol. 32, no. 4, pp. 533–542, 1986.
- [12] T. M. Cover and J. A. Thomas, *Elements of infor*mation theory (2. ed.). Wiley, 2006. [Online]. Available: http://www.elementsofinformationtheory.com/