

Finding network effect of randomized treatment under weak assumptions for any outcome and any effect heterogeneity

Myoung-jae Lee

Dept. Economics, Korea University
145 Anam-ro, Sungbuk-gu
Seoul 02841, South Korea
myoungjae@korea.ac.kr

Dept. Finance, Accounting & Economics
University of Nottingham Ningbo China
Ningbo 315100, China
myoung-jae.lee@nottingham.edu.cn

In estimating the effects of a treatment/policy with a network, an unit is subject to two types of treatment: one is the direct treatment on the unit itself, and the other is the indirect treatment (i.e., network/spillover influence) through the treated units among the friends/neighbors of the unit. In the literature, linear models are widely used where either the number of the treated neighbors or the proportion of them among the neighbors represents the intensity of the indirect treatment. In this paper, we obtain a nonparametric network-based “causal reduced form (CRF)” that allows any outcome variable (binary, count, continuous, ...) and any effect heterogeneity. Then we assess those popular linear models through the lens of the CRF. This reveals what kind of restrictive assumptions are embedded in those models, and how the restrictions can result in biases. With the CRF, we conduct almost model-free estimation and inference for network effects.

Running Head: Network effect for any outcome and heterogeneity.

Keywords: causal reduced form, network/spillover effect, treatment intensity.

Funding Information: This research has been supported by the National Research Foundation of Korea (NRF) grant from the Korea government (MSIT) (RS-2024-00337766).

Compliance with ethical standard, no conflict of interest, and no AI usage: no human/animal subject is involved in this research, and there is no conflict of interest to disclose. Also, no generative-AI-related technology has been used for this paper.

1 Introduction

Consider N units connected through a network, and a binary randomized treatment/policy D . Let Y_i be an outcome of unit $i = 1, \dots, N$, F_i be the number of direct (“first-order”) friends/neighbors of unit i (“ F ” for friends), and T_i be the total number of the treated direct friends/neighbors subject to $T_i \leq F_i$. When treatment spillover is present, Y_i is influenced not only by its own treatment D_i but also by the treatments received by friends: two treatments (direct, and indirect through the network) appear.

Let U_{0i} and U_{1i} be error terms. Define the treated proportion among friends $R_i \equiv T_i/F_i$. Two popular network-effect linear models are: with β parameters, under $F_i > 0$,

$$Y_i = \beta_0 + \beta_d D_i + \beta_\tau T_i + \beta_f F_i + U_{0i}; \quad (\text{T-model})$$

$$Y_i = \beta_0 + \beta_d D_i + \beta_r R_i + U_{1i} = \beta_0 + \beta_d D_i + \frac{\beta_r}{F_i} T_i + U_{1i}. \quad (\text{R-model})$$

See, e.g., Miguel and Kremer (2004), Oster and Thornton (2012), Bryan et al. (2014), and Cai et al. (2015) for empirical examples using the above models (or their variants), whose theoretical justifications were provided by Manski (2013), Aronow and Samii (2017), and Leung (2020). Henceforth, we often omit the subscript i unless doing so causes confusion.

In the T- and R-models, β_d is the direct effect common across the two models, but the effect (i.e., slope) of the network treatment T (the number of treated friends) differs, due to T being taken as the network treatment in the second expression of the R-model, instead of the network treatment R in the first expression. This way of rewriting the R-model helps comparing the two models. Whereas D is binary, the network treatment T is non-binary with an integer-valued treatment intensity. Network effects can be defined and accounted for in various ways (see, e.g., Hu et al. (2022) and references therein), but the T-, and R-models postulate that it operates only through the single route T or R .

A problem with the above models is that the interaction DT is ruled out. A general model encompassing the two models while allowing for interaction effects is: still under $F > 0$, for an error term U_2 ,

$$\begin{aligned} Y_i &= \beta_0 + \beta_f F_i + \beta_d D_i + \beta_\tau T_i + \beta_r R_i + \beta_{d\tau} D_i T_i + \beta_{dr} D_i R_i + U_{2i} \\ &= \beta_0 + \beta_f F_i + \beta_d D_i + \left(\beta_\tau + \frac{\beta_r}{F_i}\right) T_i + \left(\beta_{d\tau} + \frac{\beta_{dr}}{F_i}\right) D_i T_i + U_{2i}. \quad (\text{TR-model}) \end{aligned}$$

We can see which (or neither) is correct between the T- or R-models by estimating the TR-model. The total effect of a vaccine consists of the direct effect of the unit getting vaccinated and the network effect of the unit’s friends getting vaccinated. The indirect effect would be greater if the unit is also vaccinated, which is the interaction effect of DT .

Although the TR-model is more general, there are questions for all three models. First, in general, the linear models are invalid when Y is noncontinuous, and thus for noncontinuous Y , is it all right to still employ the linear models or should a nonlinear outcome model be considered as in Oster and Thornton (2012)? Second, in reality, the network effect of T is likely heterogeneous in F (and other covariates), neither constant nor proportional to F^{-1} . Third, in view of this issue, we may want to control F to find the heterogeneous effect: how exactly then should F be controlled in this case?

The issue of controlling F may be viewed more generally as the issue of specifying the network-effect part as a function of (T, F) . The T-model adopts the additive form $\beta_r T + \beta_f F$, and the R-model adopts the ratio form $\beta_r R = \beta_r T/F$. Given that the discrepancy between two groups $D = 0, 1$ can be assessed with the linear form $E(Y|D = 1) - E(Y|D = 0)$ or the ratio form $E(Y|D = 1)/E(Y|D = 0)$, both T-model and R-models may be plausible. What is right between the additive and ratio forms might be an empirical matter. Surprisingly, however, we know the true model. This is intuitively explained in the following, using the basic framework of the effect of only D on Y .

As is usual in treatment effect analysis, let X denote covariates, and (Y^0, Y^1) be the potential outcomes for $D = 0, 1$, so that the observed outcome is $Y = Y^0 + (Y^1 - Y^0)D$. Since we deal with a randomized D , ‘ $(Y^0, Y^1) \perp\!\!\!\perp D|X$ ’ trivially holds, where $\perp\!\!\!\perp$ stands for independence. Then, take $E(\cdot|D, X)$ on $Y = Y^0 + (Y^1 - Y^0)D$ to obtain:

$$E(Y|D, X) = E(Y^0|X) + E(Y^1 - Y^0|X)D.$$

Now, defining $U \equiv Y - E(Y|D, X)$ renders a nonparametric “causal reduced form (CRF)” that holds for any Y (binary, count, continuous, ...):

$$Y = E(Y^0|X) + E(Y^1 - Y^0|X)D + U, \quad E(U|D, X) = 0. \quad (1.1)$$

This CRF is the true model that holds always, where $E(Y^0|X)$ is the X -conditional intercept and $E(Y^1 - Y^0|X)$ is the X -conditional slope of D . The name “CRF” is ap-

appropriate because it is a derived/reduced form (RF), not a structural form (SF), and yet it contains the causal parameter of interest $E(Y^1 - Y^0|X)$. Of course, we do not know the functional forms of $E(Y^0|X)$ and $E(Y^1 - Y^0|X)$, but we can see the assumptions to justify the usual linear model, say $Y = \beta'_x X + \beta_d D + error$: $E(Y^0|X) = \beta'_x X$ and $E(Y^1 - Y^0|X) = \beta_d$. If X is “rich/detailed”, then $E(Y^0|X) = \beta'_x X$ might be plausible, but it would be difficult to justify $E(Y^1 - Y^0|X) = \beta_d$ because treatment effects are almost always heterogeneous in reality—think of the Covid vaccine effects. In view of this, the CRF may be estimated by ordinary least squares (OLS) estimator under $E(Y^0|X) = \beta'_x X$ and $E(Y^1 - Y^0|X) = \beta'_{dx} X$, which yields $Y = \beta'_x X + \beta'_{dx} X D + error$.

Turning back to the network effect framework, the situation is more complex due to the two types of treatment: the binary (direct) treatment $D = 0, 1$ on each unit, and an integer (indirect) treatment $T = 0, 1 \dots F$. The goals of this paper are establishing a nonparametric CRF appropriate for this situation to show what kind of restrictions are (unknowingly) embedded in the above linear models, and explaining how to estimate all effects of interest using the CRF derived by doing analogously to deriving (1.1). In the remainder of this paper, Section 2 derives two versions of CRF. Section 3 provides a simulation study. Finally, Section 4 concludes this paper.

2 Causal Reduced Form (CRF) for Network Effect

Let $Y_i(\mathbf{d})$ be the potential outcome of unit i for the counterfactual treatment vector $\mathbf{d} \equiv (d_1, d_2 \dots d_N)$. Under some assumptions, $Y_i(\mathbf{d})$ depends solely on (d_i, t_i) where t_i is the realized value of T_i , which we maintain throughout this paper; e.g., the Assumption 2 (with $g_i = t_i$) in Forastiere et al. (2021), or more specifically, the assumption that network effect occurs only among the direct friends who are exchangeable in Leung (2020).

2.1 Unrestricted CRF

Our causal effects of interest conditional on F are:

$$\delta_t(F) \equiv E(Y^{1t} - Y^{0t}|F), \tau_{1t}(F) \equiv E(Y^{1t} - Y^{10}|F), \tau_{0t}(F) \equiv E(Y^{0t} - Y^{00}|F). \quad (2.1)$$

Here, $\delta_t(F)$ is the **direct** (thus **delta**) effect of the own treatment, when the unit has t treated among F friends in total; $\tau_{1t}(F)$ is the network effect on a treated unit of having t treated out of F ; and $\tau_{0t}(F)$ is the network effect on a control unit of having t treated.

Define the (“net”) interaction effect of $(D = 1, T = t)$ as (Choi and Lee, 2018):

$$\Delta Y_t^\pm \equiv Y^{1t} - Y^{10} - Y^{0t} + Y^{00} = Y^{1t} - Y^{00} - (Y^{10} - Y^{00}) - (Y^{0t} - Y^{00})$$

which is the “gross” interaction effect $Y^{1t} - Y^{00}$ of $(D = 1, T = t)$ minus the “partial” effects $Y^{10} - Y^{00}$ of $D = 1$ and $Y^{0t} - Y^{00}$ of $T = t$. Let $1[A] \equiv 1$ if A holds and 0 otherwise. Theorem 1 below is proved in the appendix.

THEOREM 1. *The following CRF holds for any Y (binary, count, continuous,...): with $E(U|D, T, F) = 0$ by construction,*

$$Y = E(Y^{00}|F) + \delta_0(F)D + \sum_{t=1}^F \tau_{0t}(F)1[T = t] + \sum_{t=1}^F \tau_{\pm t}(F) \cdot D1[T = t] + U,$$

$$\tau_{\pm t}(F) \equiv E(Y^{1t} - Y^{10} - Y^{0t} + Y^{00}|F) = \tau_{1t}(F) - \tau_{0t}(F) = \delta_t(F) - \delta_0(F); \quad (\text{CRF1})$$

non-zero interaction effect is $\tau_{\pm t}(F) \neq 0 \iff \tau_{1t}(F) \neq \tau_{0t}(F) \iff \delta_t(F) \neq \delta_0(F)$.

As $T = \sum_{t=1}^F t \cdot 1[T = t]$, in view of CRF1, the restrictions in the linear models are:

	$E(Y^{00} F)$	$\delta_0(F)$	$\tau_{0t}(F)$	$\tau_{\pm t}(F)$	
T-model:	$\beta_0 + \beta_f F$	β_d	$\beta_\tau t$	0	
R-model:	β_0	β_d	$(\beta_r/F)t$	0	
TR-model:	$\beta_0 + \beta_f F$	β_d	$(\beta_\tau + \beta_r/F)t$	$(\beta_{d\tau} + \beta_{dr}/F)t$	(2.2)

Some restrictions here are strong; e.g., $E(Y^{00}|F) = \beta_0$ in the R-model, and $\delta_0(F) = \beta_d$ in all three models. Also, $\tau_{0t}(F)$ and $\tau_{\pm t}(F)$ are unknown functions of (t, F) , and yet the specifications of these functions are not even linear in (t, F) in all three-models.

“CRF” may sound strange, but CRF has been fruitfully used in various contexts recently—but not for network effects yet: Lee (2018, 2021, 2024), Mao and Li (2020), Choi et al. (2023), Lee et al. (2023), and Kim and Lee (2024). Traditionally, often a tightly specified SF is used that contains a parameter of interest. Instead of this, “targeted learning” (Van der Laan and Rose, 2011) declares a causal parameter of interest first, and

then considers how to identify and estimate it. Our CRF-based approach stands on a middle ground, in the sense that it uses a nonparametric RF with causal parameters.

All effects in CRF1 are allowed to be heterogeneous in F , but since F is integer-valued, those effects can be specified as linear in the dummy variables for F without loss of generality: e.g., with $f = 0, 1, \dots, \bar{f}$ for an upper bound \bar{f} on F with $P(F = \bar{f}) > 0$,

$$\delta_0(F) = \sum_{f=0}^{\bar{f}} \delta_0(f) 1[F = f] = \delta_0(0) + \sum_{f=1}^{\bar{f}} \{\delta_0(f) - \delta_0(0)\} 1[F = f].$$

Hence, OLS of Y on $(1, D, 1[T = t], D1[T = t])$ interacted with $(1[F = 1] \dots 1[F = \bar{f}])$ provides nonparametric estimators for $\{E(Y^{00}|F), \delta_0(F), \tau_{0t}(F), \tau_{\pm t}(F)\}$, from which we can then identify $\tau_{1t}(F)$ using $\tau_{1t}(F) = \tau_{0t}(F) + \tau_{\pm t}(F)$ in Theorem 1, and $\delta_t(F)$ using $\delta_t(F) = \delta_0(F) + \tau_{\pm t}(F)$. Instead of this “long OLS”, we may as well do the “short OLS” of Y on $(1, D, 1[T = t], D1[T = t])$ separately for each subsample $F = f$.

2.2 Restricted CRF with Power-Function Specifications

The aforementioned long OLS is cumbersome if the support of F is large, and is numerically unstable if $P(F = f)$ is too small for some f . E.g., suppose $\bar{f} = 20$ and the maximum value of T is 10. Then the long OLS would have about $20 + 20 + 20 \times 10 + 20 \times 10 = 440$ parameters, where the four parts correspond to those in CRF1. Here, we reduce the long OLS dimension with power-function approximations in t and F .

ASSUMPTION 1. For a known J , and some β, δ and τ parameters,

$$E(Y^{00}|F) = \sum_{j=0}^J \beta_{0j} F^j, \quad \delta_0(F) = \sum_{j=0}^J \delta_{0j} F^j, \quad \tau_{0t}(F) = \sum_{j=0}^J \tau_{01j} F^j t, \quad \tau_{\pm t}(F) = \sum_{j=0}^J \tau_{\pm 1j} F^j t.$$

Assumption 1 specifies order- J power-functions for $E(Y^{00}|F)$ and $\delta_0(F)$, and order- J (in F) and order-1 (in t) power functions for $\tau_{0t}(F)$ and $\tau_{\pm t}(F)$. Of course, a higher-order can be used for t , if desired. Using $T = \sum_{t=1}^F t 1[T = t]$, Assumption 1 simplifies CRF1 much by turning the $1[T = t]$ and $D1[T = t]$ parts in CRF1 into $(\sum_{j=0}^J \tau_{01j} F^j)T$ and $(\sum_{j=0}^J \tau_{\pm 1j} F^j)DT$, as Theorem 2 below shows in CRF2. The interaction treatment takes the “natural” form DT in CRF2, to which OLS can be easily applied.

THEOREM 2. *Under Assumption 1, CRF1 simplifies to:*

$$Y = \sum_{j=0}^J \beta_{0j} F^j + \left(\sum_{j=0}^J \delta_{0j} F^j \right) D + \left(\sum_{j=0}^J \tau_{01j} F^j \right) T + \left(\sum_{j=0}^J \tau_{\pm 1j} F^j \right) \cdot DT + U. \quad (\text{CRF2})$$

If order-2 approximation were used for t , then this would add T^2 and DT^2 terms in CRF2. One might argue that our CRF-based estimation can also misspecify the functions of F and T as the conventional T- and R-models can. This is true, but the functions of F and T in the CRF's are of RF varieties, not SF's as in the T- and R-models, and the consequences of misspecifying RF's should be less serious than misspecifying SF's. Our CRF-based estimation rests on this premise; i.e., specify RF's, not SF's, if one has to.

3 Simulation Study

Our simulation design uses the following as the basis: for an error term U_3 ,

$$\begin{aligned} Y &= \beta_0 + \beta_f F + (\beta_d + \beta_{f2} \ln F) D + (\beta_\tau + \beta_r F^{-1} + \beta_{f2} \ln F) T \\ &\quad + (\beta_{d\tau} + \beta_{dr} F^{-1} + \beta_{f2} \ln F) DT + U_3, \\ E(D) &= 0.5, \quad U_3 \sim N(0, 1) \text{ II}(D, T, F), \quad \beta_0 = 0, \quad \beta_f = -2, \quad \beta_d = 2, \\ \beta_{f2} &= 0, 0.4, \quad \beta_\tau = 0, 0.2, \quad \beta_r = 0, 2, \quad \beta_{d\tau} = 0, 0.2, \quad \beta_{dr} = 0, 2, \\ N &= 1000, 2000 \text{ (and 5000)}, \quad 1000 \text{ repetitions.} \end{aligned} \quad (3.1)$$

The parameter values for β_τ and β_{dr} are set to small numbers to make the network effect of all treated friends not too different from the direct effect of D .

For each i , we generate its longitude and latitude from the two-dimensional uniform distribution on $[0, 1]^2$, and F_i is the number of units falling within the circle of radius 0.025 from the unit i 's location, and T_i are the number of the treated among the F_i friends. Then we select only those with $F > 0$ to estimate the effects; about 63% are retained when $N = 1000$, and about 85% when $N = 2000$. Hence, the selected sample size with $N = 2000$ is about three times that with $N = 1000$.

In the selected sample with $N = 1000$, we have $\bar{F} \simeq 2.5$ with $SD(F) \simeq 1.3$ where SD stands for standard deviation, and $\bar{T} \simeq 1.6$ with $SD(T) \simeq 0.78$; the maximum value of F

is about 7. In the selected sample with $N = 2000$, we have $\bar{F} \simeq 4.2$ with $SD(F) \simeq 1.9$, and $\bar{T} \simeq 2.3$ with $SD(T) \simeq 1.2$; the maximum value of F is about 12. Whereas the effect of D is always $\beta_d = 2$, the network effect of T and the interaction effect of DT are

$$E(\beta_\tau + \beta_r F^{-1} + \beta_{f2} \ln F) \quad \text{and} \quad E(\beta_{d\tau} + \beta_{dr} F^{-1} + \beta_{f2} \ln F).$$

Different parameter values result in different models: whereas the CRF holds always,

- (i) $\beta_\tau = 0.2$ and $\beta_{f2} = \beta_r = \beta_{d\tau} = \beta_{dr} = 0$: T- and TR-models hold;
- (ii) $\beta_r = 2$ and $\beta_{f2} = \beta_\tau = \beta_{d\tau} = \beta_{dr} = 0$: R- and TR-models hold;
- (iii) $\beta_{f2} = 0$ but $\beta_\tau, \beta_r, \beta_{d\tau}, \beta_{dr}$ are not zero : TR-model holds; (3.2)
- (iv) $\beta_{f2}, \beta_\tau, \beta_r, \beta_{d\tau}, \beta_{dr}$ are all not zero: no linear model holds.

Table 1 presents the simulation results only for $N = 2000$ —the table for $N = 1000$ is omitted to save space: *four OLS's to T-, R- and TR-model, and CRF2 with quadratic approximations in F*. We consider four data generating processes (DGP's) of (3.2) using the parameter values in (3.1), as can be seen in the column headers of Table 1. The column ‘True Effect’ shows the true effects of (D, T, DT) when each of the models (i)-(iv) in (3.2) is correct; e.g., the first three rows in the True Effect column are the true effects when (i) of (3.2) is correct, and the next three rows are the true effects when (ii) is correct.

Since the true effects differ much across the DGP's, it might be better to standardize the bias's and SD's by dividing them with the true effects, which is not done, however, because some true effects are zero. For each entry of the tables, the absolute bias and SD are presented. No SD appears for the interaction effects in the T-OLS and R-OLS, because the interaction effects are supposed (i.e., “estimated”) to be zero.

In Table 1, first, when the DGP is the T model, the T-, TR- and CRF-OLS's are unbiased, but R-OLS is biased by about 10% ($100 \times 0.02 / 0.20$) in network effect as the network effect is wrong only in R-OLS. Second, when the DGP is the R-model, the R- and TR-OLS's are mostly unbiased, but T- and CRF-OLS's are biased in network effect. The bias of 30% ($= 100 \times 0.18 / 0.61$) is understandable for T-OLS as its network effect is wrong, but the bias of about 15% ($100 \times 0.09 / 0.61$) in CRF-OLS is not small. Third, when the DGP is the TR-model, both T- and R-OLS's are highly biased, whereas TR-OLS does

best and CRF-OLS is somewhat biased in three effects by about 6~13%. Finally, when the DGP is the CRF, the T- and R-OLS's are highly biased, and TR-OLS does still best despite the misspecifications (the omitted $\ln F$), and CRF-OLS is biased by 3~6%.

Table 1. Bias & SD of Four OLS's with $N = 2000$										
	True Effect		(i) T model		(ii) R model		(iii) TR		(iv) CRF	
T-OLS	(i) D	2.00	0.00	0.05	0.00	0.05	1.60	0.06	2.95	0.09
	T	0.20	0.00	0.03	0.18	0.03	0.15	0.04	0.57	0.05
	DT	0.00	0.00	0.00	0.80	1.35
R-OLS	(ii) D	2.00	0.00	0.05	0.00	0.05	1.60	0.06	2.95	0.10
	T	0.61	0.02	0.03	0.00	0.04	0.45	0.04	0.91	0.08
	DT	0.00	0.00	0.00	0.80	1.35
TR-OLS	(iii) D	2.00	0.00	0.13	0.01	0.13	0.01	0.14	0.03	0.14
	T	0.80	0.00	0.05	0.00	0.05	0.02	0.05	0.08	0.05
	DT	0.80	0.00	0.06	0.00	0.07	0.01	0.07	0.02	0.07
CRF-OLS	(iv) D	2.52	0.00	0.14	0.01	0.14	0.11	0.14	0.07	0.14
	T	1.35	0.00	0.05	0.09	0.05	0.08	0.05	0.08	0.05
	DT	1.35	0.00	0.07	0.00	0.07	0.09	0.07	0.07	0.07

'True Effect' is the true effects under the DGP's (i)-(iv) in (3.2)

Table 2. Bias & SD of Two OLS's with $N = 5000$						
	True Effect		(iii) TR		(iv) CRF	
TR-OLS	(iii) D	2.00	0.00	0.09	0.07	0.10
	T	0.44	0.00	0.01	0.01	0.01
	DT	0.44	0.00	0.02	0.00	0.02
CRF-OLS	(iv) D	2.88	0.01	0.10	0.00	0.10
	T	1.32	0.01	0.01	0.00	0.01
	DT	1.32	0.01	0.02	0.00	0.02

In Table 1, most results are what was expected for the four OLS's. However, when the DGP is TR or CRF, TR-OLS performed better than expected whereas CRF-OLS did worse. This might cast a doubt on CRF-OLS. To dissipate the doubt, Table 2 with

$N = 5000$ presents part of the results using (iii) and (iv) for TR-OLS and CRF-OLS. When the TR model is the DGP, TR-OLS does a little better than CRF-OLS, with their SD's being almost the same. However, when CRF is the DGP, CRF-OLS outperforms TR-OLS, with their SD's being the same. These findings are remarkable, considering that CRF-OLS uses only quadratic approximations—the most basic nonlinear extension from linear—whereas the DGP has F^{-1} or $(F^{-1}, \ln F)$, not F or (F, F^2) .

In summary, despite good performances of the T- and R-model OLS in some cases, overall, we find no good reason not to use the TR-model OLS, which is almost as easy to implement as the T- and R-model OLS. The TR-model can reveal which is correct between the T- and R-models, or neither. If the sample size is large, CRF-OLS should be also applied, as it can reveal whether the TR-model holds or not.

Why did TR-OLS work well in many cases? Recalling (2.2), the TR-model specifies

$$E(Y^{00}|F) = \beta_0 + \beta_f F, \quad \tau_{0t}(F) = (\beta_\tau + \beta_r F^{-1})t, \quad \tau_{\pm t}(F) = (\beta_{d\tau} + \beta_{dr} F^{-1})t.$$

These may be taken as linear specifications to the unknown functions of F , as using $\beta_r F^{-1}$ is analogous to using $-\beta_r F$. Since linear specifications often work well in practice, this seems to be why TR-OLS performed well. CRF-OLS then used F^2 extra, whose benefits were realized only in large samples, with multicollinearity problems appearing in small samples. This explains the performances of CRF-OLS vis-à-vis TR-OLS.

4 Conclusions

Network/spillover effect is an important issue, as human beings do not live in isolation and are constantly affected through interactions with others. Given a randomized binary treatment D , one popular model (“T model”) used to find the network effect takes the number of the treated friends T as the network treatment, and another popular model (“R model”) takes the ratio T/F as the network treatment where F is the number of friends/neighbors. The two models are vastly different though: if a constant β is the effect of T in the T-model, then the effect of T in the R-model is β/F —heterogeneous in F . Hence, an important question arises: which is the right model?

This paper provided an answer, considering a general model (“TR-model”) that encompasses both models. Estimating the TR-model with OLS, which is as easy as estimating the T or R-model, we can see which is the right model, or neither is.

Since the effect heterogeneity in F may not necessarily take the form β/F , we further obtained a “causal reduced form (CRF)” that holds for any form of outcome Y and for any form of effect heterogeneity in F . The CRF contains causal parameters of interest as slopes; e.g., denoting the potential outcomes as Y^{dt} where d and t index the values of (D, T) , the effect $E(Y^{10} - Y^{00}|F)$ of D with $T = 0$ appears as the slope of D in the CRF. The CRF can be estimated by specifying the unknown functions such as $E(Y^{10} - Y^{00}|F)$. Of course, misspecifications can occur, but the functions such as $E(Y^{10} - Y^{00}|F)$ are “reduced forms (RF’s)”, not “structural forms (SF’s)” as the T- and R-models are, and specifying RF’s should be less riskier than specifying SF’s.

In our simulation study, the TR-model worked well, and only in large samples, OLS for CRF outperformed OLS for the TR-model. This was attributed to that the TR-model essentially uses a linear approximation to unknown functions such as $E(Y^{10} - Y^{00}|F)$, and OLS for CRF uses power functions of F extra. Given that the TR-model was motivated by the simple desire to encompass both T- and R-models, finding its good performance is something of a “serendipity”. Our hope is that researchers see the advantages of the TR-model and CRF-based estimation, and apply them fruitfully in their research.

Appendix: Proof of Theorem 1

Using $Y^{dT} = \sum_{t=0}^F 1[T = t] \cdot Y^{dt}$, $Y = (1 - D)Y^{0T} + DY^{1T}$ becomes

$$\begin{aligned}
Y &= (1 - D) \sum_{t=0}^F 1[T = t] \cdot Y^{0t} + D \sum_{t=0}^F 1[T = t] \cdot Y^{1t} \\
&= (1 - D) \{Y^{00} + \sum_{t=1}^F 1[T = t](Y^{0t} - Y^{00})\} + D \{Y^{10} + \sum_{t=1}^F 1[T = t](Y^{1t} - Y^{10})\} \\
&= Y^{00} + (Y^{10} - Y^{00})D + \sum_{t=1}^F \{(1 - D)(Y^{0t} - Y^{00}) + D(Y^{1t} - Y^{10})\} \cdot 1[T = t] \\
&= Y^{00} + (Y^{10} - Y^{00})D + \sum_{t=1}^F (Y^{0t} - Y^{00} + \Delta Y_t^\pm D) \cdot 1[T = t].
\end{aligned}$$

Since T is as good as randomized given $F > 0$, $E(Y^{dt}|D, T, F) = E(Y^{dt}|F)$ holds when $F > 0$. Also, $F = 0$ implies $T = 0$ to drop T from $E(Y^{dt}|D, T, F = 0)$. Hence, $E(Y^{dt}|D, T, F) = E(Y^{dt}|F)$ holds always. Take $E(\cdot|D, T, F)$ on the above Y equation:

$$E(Y|D, T, F) = E(Y^{00}|F) + E(Y^{10} - Y^{00}|F)D + \sum_{t=1}^F \{\tau_{0t}(F) + E(\Delta Y_t^\pm|F)D\} \cdot 1[T = t].$$

Then, $U \equiv Y - E(Y|D, T, F)$ renders CRF1 with $E(U|D, T, F) = 0$. Also,

$$\begin{aligned} \tau_{1t}(F) - \tau_{0t}(F) &= E(Y^{1t} - Y^{10}|F) - E(Y^{0t} - Y^{00}|F) = E(\Delta Y_t^\pm|F); \\ \delta_t(F) - \delta_0(F) &= E(Y^{1t} - Y^{0t}|F) - E(Y^{10} - Y^{00}|F) = E(\Delta Y_t^\pm|F). \end{aligned}$$

REFERENCES

Aronow, P.M. and C. Samii, 2017, Estimating average causal effects under general interference, *Annals of Applied Statistics* 11, 1912-1947.

Bryan, G., S. Chowdhury and A.M. Mobarak, 2014, Underinvestment in a profitable technology: the case of seasonal migration in Bangladesh, *Econometrica* 82, 1671-1748.

Cai, J., A. De Janvry and E. Sadoulet, 2015, Social networks and the decision to insure, *American Economic Journal: Applied Economics* 7, 81-108.

Choi, J.Y., G. Lee and M.J. Lee, 2023, Endogenous treatment effect for any response conditional on control propensity score, *Statistics and Probability Letters* 196, 109747.

Choi, J.Y. and M.J. Lee, 2018, Regression discontinuity with multiple running variables allowing partial effects, *Political Analysis* 26, 258-274.

Forastiere, L., E.M. Airoidi and F. Mealli, 2021, Identification and estimation of treatment and interference effects in observational studies on networks, *Journal of the American Statistical Association* 116, 901-918.

Hu, Y., S. Li and S. Wager, 2022, Average direct and indirect causal effects under interference, *Biometrika* 109, 1165-1172.

Kim, B.R. and M.J. Lee, 2024, Instrument-residual estimator for multi-valued instruments under full monotonicity, *Statistics and Probability Letters* 213, 110187.

Lee, G., J.Y. Choi and M.J. Lee, 2023, Minimally capturing heterogeneous complier effect of endogenous treatment for any outcome variable, *Journal of Causal Inference* 11, 20220036.

Lee, M.J., 2018, Simple least squares estimator for treatment effects using propensity score residuals, *Biometrika* 105, 149-164.

Lee, M.J., 2021, Instrument residual estimator for any response variable with endogenous binary treatment, *Journal of the Royal Statistical Society (Series B)* 83, 612-635.

Lee, M.J., 2024, Direct, indirect and interaction effects based on principal stratification with a binary mediator, *Journal of Causal Inference* 12, 20230025.

Leung, M.P., 2020, Treatment and spillover effects under network interference, *Review of Economics and Statistics* 102, 368-380.

Manski, C.F., 2013, Identification of treatment response with social interactions, *Econometrics Journal* 16, S1-S23.

Mao, H. and L. Li, 2020, Flexible regression approach to propensity score analysis and its relationship with matching and weighting, *Statistics in Medicine* 39, 2017-2034.

Miguel, E. and M. Kremer, 2004, Worms: identifying impacts on education and health in the presence of treatment externalities, *Econometrica* 72, 159-217.

Oster, E. and R. Thornton, 2012, Determinants of technology adoption: peer effects in menstrual cup take-up, *Journal of the European Economic Association* 10, 1263-1293.

Van der Laan, M.J. and S. Rose, 2011, Targeted learning: causal inference for observational and experimental data, Springer.