

# Moving toward best practice when using propensity score weighting in survey observational studies

Yukang Zeng<sup>1,2</sup>, Fan Li<sup>1,3,4</sup>, Guangyu Tong<sup>1,2,3,4</sup>

<sup>1</sup> Department of Biostatistics, Yale School of Public Health, New Haven, CT, USA

<sup>2</sup> Cardiovascular Medicine Analytics Center, Yale School of Medicine, New Haven, CT, USA

<sup>3</sup> Section of Cardiovascular Medicine, Department of Internal Medicine,  
Yale School of Medicine, New Haven, CT, USA

<sup>4</sup> Center for Methods in Implementation and Prevention Science,  
Yale School of Public Health, New Haven, CT, USA

Corresponding Author: Guangyu Tong, PhD

Email: [guangyu.tong@yale.edu](mailto:guangyu.tong@yale.edu)

January 28, 2025

## Abstract

Propensity score weighting is a common method for estimating treatment effects with survey data. The method is applied to minimize confounding using measured covariates that are often different between individuals in treatment and control. However, existing literature does not reach a consensus on the optimal use of survey weights for population-level inference in the propensity score weighting analysis. Under the balancing weights framework, we provided a unified solution for incorporating survey weights in both the propensity score of estimation and the outcome regression model. We derived estimators for different target populations, including the combined, treated, controlled, and overlap populations. We provide a unified expression of the sandwich variance estimator and demonstrate that the survey-weighted estimator is asymptotically normal, as established through the theory of M-estimators. Through an extensive series of simulation studies, we examined the performance of our derived estimators and compared the results to those of alternative methods. We further carried out two case studies to illustrate the application of the different methods of propensity score analysis with complex survey data. We concluded with a discussion of our findings and provided practical guidelines for propensity score weighting analysis of observational data from complex surveys.

**Keywords:** complex survey, propensity score weighting, survey weights, overlap weights, covariate balance, augmented weighting estimators

# 1 Introduction

The use of observational data to estimate the causal effects of exposures or interventions is prevalent in medical research, particularly when randomized trials are unavailable or unethical. In many cases, researchers resort to observational data garnered through surveys with complex sampling techniques. These techniques are designed to ensure data from a sample can accurately represent the target population. In extensive surveys, these techniques may encompass complex features such as multistage sampling and deliberate oversampling of less representative subgroups.[1] The survey weights computed based on the design are integral to recovering the population features so that the differentiated selection probabilities and non-responses are appropriately accounted for. There is consensus on the importance of incorporating the survey design into the analytical process to derive causal effect estimates that are applicable to the target population.[2, 3, 4]

Survey weights also play a crucial role in ensuring the accuracy and representativeness of the propensity score method, which is among the most commonly used tool for causal inference using observational data.[5] In the absence of randomization, systematic differences in baseline characteristics usually exist between exposed and unexposed individuals, leading to the issue of confounding. The propensity score methods rest on the treatment assignment process with observed covariates,[6] and with estimated propensity scores, differences in the observed baseline features across groups can be reduced so that the estimated treatment effect will no longer be attributable to pre-existing group differences. In practice, implementing propensity score methods involve a two-step procedure, including propensity score estimation and the subsequent treatment effect estimation. The latter step leverages the estimated propensity score through stratification,[7] matching,[8] or weighting.[9]

The appropriate use of survey weights in stratification to obtain population causal effect estimates has been previously addressed, with a recommendation that survey weights should be used when estimating treatment effects within strata but not in the estimation of the propensity scores. [10, 2] DuGoff et al. later suggested that survey weights can carry unique design features that are not captured by observed covariates, and therefore should be included as an additional covariate in the propensity score model to enhance covariate balance and address potential bias due to unmeasured covariates.[3] Similar recommendations have also been extended to propensity score matching and weighting, where the estimation of propensity score can incorporate survey weights as an additional covariate; but in those contexts, whether to perform survey-weighted analysis in the outcome model depends on the goal of the analysis (i.e., whether a sample or the underlying population defines the target of inference).[3]. Austin et al.[11] illustrated through simulations that including survey weight as a covariate in the propensity score model showed no clear advantage over a survey-weighted propensity score model. Neither approach can consistently outperform the other regarding covariate balance and bias in the treatment effect estimation across various scenarios. Dong et al. [12] evaluated multiple strategies for incorporating survey weights into both propensity score and outcome models, recommending their use at both stages, although they observed that estimates from different strategies did not yield apparent differences in their data application. Lenis et al.[13] found through simulations that incorporating survey weights in the propensity score model does not necessarily improve the estimation of population-level treatment effect using matching, especially when sufficient population-

level balance in confounders can already be achieved. They also showed through simulations that incorporating survey weights in propensity score estimation can reduce bias when the missingness in outcome is correlated with survey weights.[13]

Regarding propensity score weighting, Ridgeway et al.[4] provided a theoretical justification for the use of a survey-weighted propensity score model in obtaining a consistent estimator of the population-level average treatment effect. They proved that using survey weights in both stages leads to consistent and robust population average treatment effect estimators under various generating mechanisms (e.g., survey weights are either informative or non-informative for the treatment assignment mechanism). Salerno et al. [14] recently explored a complex survey mechanism in health disparities research, where survey weights depend on the group variable of interest under comparison, such as race. They further revealed that whether the propensity score should be survey-weighted corresponds to whether the available survey weights are tied to comparison variables such as treatment or exposure. Whereas Ridgeway et al.[4] derived the results and performed simulations for the continuous outcome scenario, Yang et al.[15] extended the results to a binary outcome and evaluated the performance of different ways to incorporate survey weights in the propensity score model and the outcome model. Both Ridgeway et al.[4] and Yang et al.[15] considered the potentially differentiated practice of incorporating survey weights when estimating the average treatment effect (ATE) and the average treatment effect of the treated (ATT).

In general, the emerging message from the existing literature is that the use of survey weights at the propensity score model stage and the outcome model stage can potentially be beneficial in reducing the bias of population-level inference. However, a consensus on the best practice and the underlying justifications for using survey weights to target different causal estimands of interest have not been fully elucidated. This paper thus aims to contribute to the literature on propensity score weighting in the following several ways. First, the target estimands in the prior development on survey observational data methods were limited to ATE and ATT, but not to other estimands (e.g., ATO, the average treatment effect among the overlap population[9]). We address this by supplying a unified framework that incorporates survey weights into the general framework of balancing weights.[9] Second, as a further extension, we detail the use of survey weights for augmented weighting estimators under the balancing weights framework, which can potentially improve efficiency over weighting alone by leveraging information from a posited outcome regression function. Third, we developed closed-form variance estimators that accommodate survey weights in each estimator we considered for computationally efficient inference. Lastly, given that no consensus has been reached on the best use of survey weights in propensity score weighting analysis under various real-world scenarios (such as varying levels of covariate overlap and model misspecification), we report on an extension simulation study that shed light on the relative comparisons between methods and provide recommendations to help practitioners enhance the reliability of causal evidence obtained from survey observational data.

The rest of this paper is organized as follows. Section 2 defines the notation and assumptions for the potential outcomes framework under the context of survey observational data. Under the balancing weight framework, this section then derives the weighting estimator and emphasizes the role of survey weights in addition to balancing weights. We provide a unified expression of the sandwich variance estimator and demonstrate that the survey-weighted estimator is asymptotically normal, as established through the theory of

M-estimators. In Section 3, we present three augmented estimators incorporating survey weights and further provide the associated sandwich variance structure for large-sample inference. In Section 4, we conduct a full series of simulations to examine the performance of different estimators incorporating propensity score weights for different estimands with data generated from multistage clustering sampling. Section 5 compares the performance of the different estimation procedures in an application using the Early Childhood Longitudinal Study, Kindergarten Class of 1998-1999 (ESCL-K). [16] In Section 6, we present our main conclusions and provide a discussion.

## 2 Methods for incorporating survey weights into propensity score weighting

### 2.1 Definitions and assumptions

In this section, we establish our study framework starting with the setup of finite sample and population under the context of survey observational data. Consider a finite sample of size  $n$  selected from a superpopulation of size  $N$ . Each individual in the sample is indexed by  $i$ , where  $i = 1, \dots, n$ . Assuming a  $p$ -dimensional vector for baseline covariates for each individual  $\mathbf{X}_i$  that includes observed pretreatment variables, and a binary exposure variable  $Z_i = z$  that indicates treatment ( $z = 1$ ) or control ( $z = 0$ ) group affiliation. Under the Stable Unit Treatment Value Assumption (SUTVA), the potential outcomes  $Y_i(z)$  for  $z \in \{0, 1\}$  are well-defined with no interference between population units where the observed outcome for each individual in the sample is  $Y_i = Z_i Y_i(1) + (1 - Z_i) Y_i(0)$ . For estimating population effects using our sample, we define the survey indicator  $S_i$  where  $S_i = 1$  denotes selected to the sample and  $S_i = 0$  denotes unselected to the sample. We assume selection positivity such that every individual in the superpopulation has a non-zero probability of selection,  $0 < p_i = p(\mathbf{x}_i) = P(S_i = 1 | \mathbf{X}_i = \mathbf{x}_i) < 1$ .

We assume conditional independence (or unconfoundedness) for the treatment assignment,  $\{Y(0), Y(1)\} \perp Z | \mathbf{X}$ , and for the sampling process after the exposure has occurred,  $\{Y(0), Y(1)\} \perp S | Z, \mathbf{X}$ , which implies that treatment assignment bias arises solely from  $\mathbf{X}$ , and selection bias from the afterwards sampling process can only be caused by  $Z$  and  $\mathbf{X}$ . These assumptions can be relaxed to weak treatment assignment exchangeability, expressed as  $E[Y(z) | \mathbf{X}] = E[Y(z) | Z = z, \mathbf{X}]$ , and weak selection exchangeability, formulated as  $E[Y(z) | Z = z, \mathbf{X}] = E[Y(z) | Z = z, \mathbf{X}, S = 1]$ . These relaxed assumptions still enable the identification of causal effects and enable the generalization from the finite sample to the superpopulation, as previously noted by Salerno et al.[14] One causal estimand of interest is the average treatment effect (ATE), denoted as  $\tau(\mathbf{x}) = E[Y(1) - Y(0) | \mathbf{X} = \mathbf{x}]$ . Imbens[17] differentiates population-level ATE (PATE)  $\tau_{\text{PATE}} = E[Y(1) - Y(0)]$  from the sample average treatment effect (SATE), which we denote as  $\tau_{\text{SATE}} = E[Y(1) - Y(0) | S = 1]$ . The population and sample level propensity scores for individual  $i$  are denoted as  $e_{\text{sp}}(\mathbf{x}_i) = P(Z_i = 1 | \mathbf{X}_i = \mathbf{x}_i)$  and  $e_{\text{fp}}(\mathbf{x}_i) = P(Z_i = 1 | \mathbf{X}_i = \mathbf{x}_i, S_i = 1)$  respectively. They will be equal only when sampling and treatment assignment are independent conditional on baseline covariates, i.e.,  $Z \perp S | \mathbf{X}$ .

In a prospective study setup where treatment assignment occurred after the sampling,

we can define the sampling probability for individual  $i$  as the probability of inclusion in the study based on observed covariates,  $p_i = P(S_i = 1 | \mathbf{X}_i = \mathbf{x}_i)$ . In contrast, for a retrospective study where sampling occurs after exposure or treatment assignment, the sampling probability can depend on both observed covariates and treatment and is denoted as  $p_{z,i} = p_z(\mathbf{x}_i) = P(S_i = 1 | Z_i = z, \mathbf{X}_i = \mathbf{x}_i)$ . To provide further clarity, we define  $p_1(\mathbf{x}_i) = P(S_i = 1 | Z_i = 1, \mathbf{X}_i = \mathbf{x}_i)$  and  $p_0(\mathbf{x}_i) = P(S_i = 1 | Z_i = 0, \mathbf{X}_i = \mathbf{x}_i)$  as the specific sampling probabilities for the selected individuals in the treatment and control groups, respectively. When we assume conditional independence between sampling and treatment assignment, represented as  $Z \perp S | \mathbf{X}$ , the sampling probabilities for both prospective and retrospective designs align conceptually with the propensity score framework. In this framework, sampling is viewed as a separate assignment mechanism that can be adjusted for in the analysis to address potential biases. Moreover, these different sampling probabilities can then be connected through a ratio,  $p_i = r_{z,i} p_{z,i}$ , where  $r_{z,i} = r_z(\mathbf{x}_i)$  captures the discrepancy between propensity scores in the sample and population levels. For a treated unit ( $z = 1$ ),  $r_1(\mathbf{x}_i)$  equals  $e_{\text{sp}}(\mathbf{x}_i)/e_{\text{fp}}(\mathbf{x}_i)$ , the ratio of propensity score at population and sample level. For a control unit ( $z = 0$ ),  $r_0(\mathbf{x}_i)$  equals  $(1 - e_{\text{sp}}(\mathbf{x}_i))/(1 - e_{\text{fp}}(\mathbf{x}_i))$ . Furthermore,  $r_z(\mathbf{x}_i)p_z(\mathbf{x}_i)$  can be used to approximate  $p_i = P(S_i = 1 | \mathbf{X}_i = \mathbf{x}_i)$  even when the marginalized survey weights  $p_i$  are not directly accessible. As Salerno et al.[14] highlighted, the distinction between prospective and retrospective studies underscore two different ways of factorizing  $P(S_i = 1, Z_i = z | \mathbf{X}_i = \mathbf{x}_i)$ : one conditional on the covariates alone (for prospective studies) and the other conditional on both treatment and covariates (for retrospective studies). To ensure the validity of causal estimates, assumptions for positive probabilities of sampling and treatment assignment must hold at both the population and sample levels, with unconfoundedness providing the necessary ignorable conditions for both mechanisms. In this paper, we primarily focus on the retrospective study setup, which commonly occurs in survey observational data; in other words, we address the scenario where the exposure is already implemented in the target population.

Regarding the setup of propensity score weighting, we follow the framework of balancing weights [9] that characterizes a general class of weighted distributions of covariates across two treatment groups that can achieve balance. Balancing weights typically leverages the propensity score to calibrate covariates of each group towards a target population with an estimand of interest. The density of the target population is represented as  $g(\mathbf{X}) = f(\mathbf{X})h(\mathbf{X})$ , where  $f(\mathbf{X})$  signifies the marginal density of covariates within the finite population, and  $h(\mathbf{X})$ , known as the tilting function, characterizes the target population and estimand. The balancing weights framework articulates the estimand as a weighted average of individual-level conditional treatment effects: [9, 18]

$$\tau_h = \frac{E[h(\mathbf{X})\tau(\mathbf{X})]}{E[h(\mathbf{X})]}. \quad (1)$$

In this equation, the expectations in both the numerator and the denominator are taken over the target population. The tilting function  $h(\mathbf{X})$  is a non-negative function of covariates and satisfies  $E[h(\mathbf{X})] > 0$ .

## 2.2 Incorporating survey weights into propensity score weighting

Under the balancing weights framework, the tilting function  $h(\mathbf{X}) = g(\mathbf{X})/f(\mathbf{X})$  serves as a bridge to tilt the covariate distribution in the population  $f(\mathbf{X})$  to that in the target population  $g(\mathbf{X})$ . Since our goal is to derive the expressions of various estimands for target populations in the superpopulation, we first derive a general form of this class of estimands  $\tau_h(\mathbf{X})$  considering the average treatment effect over the target population in the superpopulation. Under weak exchangeability in both the treatment assignment and sampling processes, Equation (1) can be expressed as,

$$\tau_h = \frac{E \left[ \frac{h(\mathbf{X})}{p_1(\mathbf{X})e_{\text{sp}}(\mathbf{X})} ZY \mid S = 1 \right]}{E \left[ \frac{h(\mathbf{X})}{p_1(\mathbf{X})e_{\text{sp}}(\mathbf{X})} Z \mid S = 1 \right]} - \frac{E \left[ \frac{h(\mathbf{X})}{p_0(\mathbf{X})(1 - e_{\text{sp}}(\mathbf{X}))} (1 - Z)Y \mid S = 1 \right]}{E \left[ \frac{h(\mathbf{X})}{p_0(\mathbf{X})(1 - e_{\text{sp}}(\mathbf{X}))} (1 - Z) \mid S = 1 \right]}. \quad (2)$$

When the expectations are replaced with sample means, a unbiased estimator of  $\tau_h$  in the form of a weighted difference in the outcomes between treatment groups in the target population can be expressed as

$$\hat{\tau}_h = \frac{\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i) Z_i Y_i}{\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i) Z_i} - \frac{\sum_{i=1}^n \omega_{0,p}(\mathbf{X}_i) (1 - Z_i) Y_i}{\sum_{i=1}^n \omega_{0,p}(\mathbf{X}_i) (1 - Z_i)}. \quad (3)$$

In this equation, the weights are given by

$$(\omega_{1,p}(\mathbf{X}_i), \omega_{0,p}(\mathbf{X}_i)) = \left( \frac{h(\mathbf{X}_i)}{p_1(\mathbf{X}_i)e_{\text{sp}}(\mathbf{X}_i)}, \frac{h(\mathbf{X}_i)}{p_0(\mathbf{X}_i)(1 - e_{\text{sp}}(\mathbf{X}_i))} \right). \quad (4)$$

Details of proof are included in Web Appendix A.1. It is important to note that the above derivation primarily applies to retrospective studies, where exposure occurs before survey sampling begins. However, this approach can also be generalized to prospective studies by defining the sampling weights as  $p(S_i = 1 \mid \mathbf{X}_i)$  for individual  $i$ . In such cases, one can factorize  $p(Z_i = 1, S_i = 1 \mid \mathbf{X}_i)$  differently as  $p(S_i = 1 \mid \mathbf{X}_i)p(Z_i = 1 \mid S_i = 1, \mathbf{X}_i)$ , where the second term represents the in-sample propensity score.[14] Further details on the corresponding estimators under prospective settings are extended and discussed in Web Appendix A.6.

The specification of  $h(\mathbf{X})$  can adopt various forms depending on specific statistical or scientific considerations. Some commonly used forms of balancing weights and their associated target populations, tilting functions, and estimands are presented in Table 1. In this table, PATE refers to the Population Average Treatment Effect, defined as  $\tau_{\text{PATE}} = E[Y(1) - Y(0)]$ , with  $g(\mathbf{X}) = f(\mathbf{X})$  and  $h(\mathbf{X}) = 1$ . PATT denotes the Population Average Treatment Effect for the Treated, expressed as  $\tau_{\text{PATT}} = E[Y(1) - Y(0) \mid Z = 1]$ , where  $g(\mathbf{X}) = f(\mathbf{X} \mid Z = 1)$  and  $h(\mathbf{X}) = e_{\text{sp}}(\mathbf{X})$ . Complimentary to PATT, PATC represents the Population Average Treatment Effect for the Control, given by  $\tau_{\text{PATC}} = E[Y(1) - Y(0) \mid Z = 0]$ , with  $g(\mathbf{X}) = f(\mathbf{X} \mid Z = 0)$  and  $h(\mathbf{X}) = 1 - e_{\text{sp}}(\mathbf{X})$ . The overlap weights defined by Li et al. [9] correspond to the Average Treatment Effect among the Overlap Population (PATO) under the tilting function of the average  $h(\mathbf{X}) = e_{\text{sp}}(\mathbf{X})(1 - e_{\text{sp}}(\mathbf{X}))$ , such that greater emphasis

is given to units with propensity scores close to 0.5 and smaller weights were assigned to those with extreme propensity scores with more deterministic treatment assignment. In the absence of survey sampling, the choice of the overlap tilting function attains the minimum asymptotic variance among all choices of the tilting functions under the homoscedasticity assumption. [9] Under the survey observational data setting, we show that this minimum asymptotic variance property continues to hold, along with the exact balance properties. We summarize the properties below and details of the proof can be found in Web Appendix A.2.

**Proposition 1.** When propensity scores are estimated using a survey-weighted logistic regression under weighted maximum likelihood estimation, where  $\hat{e}_{\text{sp}}(\mathbf{X}_i) = \text{logit}^{-1}(\hat{\boldsymbol{\beta}}_0 + \mathbf{X}_i \hat{\boldsymbol{\beta}}_{\text{sw}}^\top)$ , the application of overlap weights ensures an exact balance in the population means of any given covariate across treatment and control groups. That is,

$$\frac{\sum_{i=1}^n X_{ik} Z_i (1 - \hat{e}_{\text{sp}}(\mathbf{X}_i)) / p_1(\mathbf{X}_i)}{\sum_{i=1}^n Z_i (1 - \hat{e}_{\text{sp}}(\mathbf{X}_i)) / p_1(\mathbf{X}_i)} = \frac{\sum_{i=1}^n X_{ik} (1 - Z_i) \hat{e}_{\text{sp}}(\mathbf{X}_i) / p_0(\mathbf{X}_i)}{\sum_{i=1}^n (1 - Z_i) \hat{e}_{\text{sp}}(\mathbf{X}_i) / p_0(\mathbf{X}_i)}, \quad \text{for } k = 1, 2, \dots, K$$

**Proposition 2.** The function  $h(\mathbf{X}) \propto e_{\text{sp}}(\mathbf{X}) (1 - e_{\text{sp}}(\mathbf{X}))$  minimizes the asymptotic variance of the weighting estimator  $\hat{\tau}_h$  across all possible  $h$  functions, under the assumption of homoscedasticity for both the original outcome  $Y(z)$  and the pseudo-outcome  $Y(z)/p_z(\mathbf{X})$  across treatment groups ( $z = 1$  and  $z = 0$ ). As the sample size  $n$  approaches infinity:

$$n \min_h \{E_{\mathbf{X}} \text{Var} [\hat{\tau}_h \mid \mathbf{X}, S = 1]\} \rightarrow \frac{\sqrt{vc}}{C_{hs}^2} \int e_{\text{sp}}(\mathbf{X}) (1 - e_{\text{sp}}(\mathbf{X})) \frac{1}{p(S = 1 \mid \mathbf{X})} f(\mathbf{X} \mid S = 1) \mu(d\mathbf{X}),$$

where  $v = \text{Var}(Y(z)/p_z(\mathbf{X}) \mid \mathbf{X})$ ,  $c = \text{Var}(Y(z) \mid \mathbf{X})$ , and  $C_{hs} = \int (h(\mathbf{X})/p(S = 1 \mid \mathbf{X})) f(\mathbf{X} \mid S = 1) \mu(d\mathbf{X})$ .

Proposition 2 leads to optimal efficiency in estimating  $\hat{\tau}_h$ , minimizing the variance among all forms of balancing weights. While the homoscedasticity condition may not always hold in practice, our simulations demonstrate that using overlap weights consistently achieves smaller variances in finite survey samples. Furthermore, previous empirical work on nonsurvey-weighted outcomes has shown that overlap weights improve efficiency compared to inverse probability of treatment weighting (IPTW), regardless of the homoscedasticity condition. [19, 20, 21] Under these two propositions, overlap weights can still mitigate the covariate balancing and extreme propensity score issues under the survey sampling setup. In addition, for truncated combined, the tilting function  $h(\mathbf{X}) = 1\{\alpha < e_{\text{sp}}(\mathbf{X}) < 1 - \alpha\}$  can be used to exclude individuals with extreme propensity scores, which facilitates more reliable estimations by focusing on a population with sufficient overlap. [22] The parameter  $\alpha$ , typically chosen within the range  $(0, 0.1)$ , indicates the truncation threshold for propensity scores.

### 2.3 Asymptotic normality and the sandwich variance estimator

Under the framework of balancing weights, this section establishes the asymptotic normality of the propensity score weighting estimator and derives the robust sandwich variance

Table 1: Survey-weighted propensity score weights for different forms of balancing weights, target population, tilting function, and estimand.

Target population	$h(\mathbf{X})$	Estimand	Weight $(\omega_{1,p}, \omega_{0,p})$
Combined	1	PATE	$\left( \frac{1}{p_1(\mathbf{X}) e_{\text{sp}}(\mathbf{X})}, \frac{1}{p_0(\mathbf{X}) (1 - e_{\text{sp}}(\mathbf{X}))} \right)$
Treated	$e_{\text{sp}}(\mathbf{X})$	PATT	$\left( \frac{1}{p_1(\mathbf{X})}, \frac{1}{p_0(\mathbf{X}) (1 - e_{\text{sp}}(\mathbf{X}))} \right)$
Control	$(1 - e_{\text{sp}}(\mathbf{X}))$	PATC	$\left( \frac{1}{p_1(\mathbf{X}) (1 - e_{\text{sp}}(\mathbf{X}))}, \frac{1}{p_0(\mathbf{X})} \right)$
Overlap	$e_{\text{sp}}(\mathbf{X}) (1 - e_{\text{sp}}(\mathbf{X}))$	PATO	$\left( \frac{1}{p_1(\mathbf{X}) (1 - e_{\text{sp}}(\mathbf{X}))}, \frac{1}{p_0(\mathbf{X}) e_{\text{sp}}(\mathbf{X})} \right)$
Truncated combined	$\mathbf{1}(\alpha < e_{\text{sp}}(\mathbf{X}) < 1 - \alpha)$		$\left( \frac{1}{p_1(\mathbf{X})} \frac{\mathbf{1}(\alpha < e_{\text{sp}}(\mathbf{X}) < 1 - \alpha)}{e_{\text{sp}}(\mathbf{X})}, \frac{1}{p_0(\mathbf{X})} \frac{\mathbf{1}(\alpha < e_{\text{sp}}(\mathbf{X}) < 1 - \alpha)}{1 - e_{\text{sp}}(\mathbf{X})} \right)$

estimator in the presence of survey weights. One viable approach for inference is the bootstrap method, which could ascertain the variance of  $\hat{\tau}_h$  across various target populations. However, bootstraps can involve a substantial amount of computation, especially when the sample size gets larger. In addition, bootstrap survey data often entails additional complications, see, for example, Ackerman et al.[23] Leveraging the asymptotic normality property of the survey-weighted estimator, we provide a closed-form variance estimator via the empirical sandwich method, which serves as a feasible solution to circumvent the need for extensive computation.[24] Detailed proof can be found in Web Appendix A.3.

Assuming the propensity scores are estimated using a survey-weighted logistic regression model and in the form of  $e_{\text{sp}}(\mathbf{X}_i; \boldsymbol{\beta}) = 1 / (1 + \exp(-\mathbf{X}_i^\top \boldsymbol{\beta}))$ . We define  $\tau_1 = \frac{E[h(\mathbf{X})\mu_1(\mathbf{X})]}{E[h(\mathbf{X})]}$  and  $\tau_0 = \frac{E[h(\mathbf{X})\mu_0(\mathbf{X})]}{E[h(\mathbf{X})]}$ , where  $\mu_1(\mathbf{X}) = E[Y(1) | \mathbf{X}]$  and  $\mu_0(\mathbf{X}) = E[Y(0) | \mathbf{X}]$ , as the treatment-specific average potential outcomes in the target population, and the treatment effect estimator of target population can be written as  $\hat{\tau}_h = \hat{\tau}_1 - \hat{\tau}_0$ . The consistent sandwich-type variance estimator for  $\hat{\tau}_h$  is given by:

$$\widehat{\text{Var}}(\hat{\tau}_h) = n^{-2} \sum_{i=1}^n \left( \hat{I}_i / \hat{\nu} \right)^2, \quad (5)$$

where  $\hat{\nu} = n^{-1} \sum_{i=1}^n h(\mathbf{X}_i) / P(S_i = 1 | \mathbf{X}_i)$  represents the weighted adjustment factor that accounts for survey weights  $1/p_z(\mathbf{X}_i)$ . The influence function  $\hat{I}_i$  here, characterizing the contribution of the  $i$ -th sampled individual to the overall population-level treatment effect estimator, is defined as:

$$\hat{I}_i = Z_i \hat{\omega}_{1,p}(\mathbf{X}_i; \hat{\boldsymbol{\beta}}) (Y_i - \hat{\tau}_1) - (1 - Z_i) \hat{\omega}_{0,p}(\mathbf{X}_i; \hat{\boldsymbol{\beta}}) (Y_i - \hat{\tau}_0) + \hat{\mathbf{H}}_{\hat{\boldsymbol{\beta}}}^\top \hat{E}_{\hat{\boldsymbol{\beta}}}^{-1} \frac{1}{p_z(\mathbf{X}_i)} \left( Z_i - \hat{e}_{\text{sp}}(\mathbf{X}_i; \hat{\boldsymbol{\beta}}) \right) \mathbf{X}_i. \quad (6)$$

where the gradient of the balancing weights with respect to the logistic regression parameters



$\beta$  is:

$$\widehat{\mathbf{H}}_{\widehat{\beta}} = n^{-1} \sum_{i=1}^n [Z_i (Y_i - \widehat{\tau}_1) \omega'_{1,p,\beta} - (1 - Z_i) (Y_i - \widehat{\tau}_0) \omega'_{0,p,\beta}], \quad (7)$$

Here,  $\omega'_{1,p,\beta} = \partial \omega_{1,p}(\mathbf{X}_i; \beta) / \partial \beta$  and  $\omega'_{0,p,\beta} = \partial \omega_{0,p}(\mathbf{X}_i; \beta) / \partial \beta$  quantify how sensitive the balancing weights are to changes in propensity score model parameters. Further, the covariance matrix of the propensity score estimates  $\widehat{\mathbf{E}}_{\beta\beta}$  here is expressed as:

$$\widehat{E}_{\widehat{\beta}\widehat{\beta}} = n^{-1} \sum_{i=1}^n \frac{1}{p_z(\mathbf{X}_i)} \widehat{e}_{\text{sp}}(\mathbf{X}_i; \widehat{\beta}) \left(1 - \widehat{e}_{\text{sp}}(\mathbf{X}_i; \widehat{\beta})\right) \mathbf{X}_i \mathbf{X}_i^\top. \quad (8)$$

The sandwich variance estimator accounts for the uncertainty in the estimation of propensity scores and also adjusts for the known survey weights.

### 3 Improving the weighting estimator through outcome regression

To improve the weighting estimators, augmented weighting estimators have proven useful for causal inference and missing data analysis.[7, 25, 26] These estimators are specifically designed to leverage additional information from outcome modeling, and for inverse probability weighting (IPW), can provide consistent estimation of causal effects even if only one of the models—treatment assignment mechanism or the outcome model—is correctly specified, but not necessarily both. In the absence of survey sampling, Mao et al.[27] investigated these augmentation methods to estimate different target estimands in observational studies under the balancing weight framework.

In this section, we first develop the augmented weighting (AW) estimating equations based on the balancing weights in Section 3.1 in survey observational data. We then introduce three versions of augmented estimators adapted to in Sections 3.2 to 3.4: the moment estimator, the clever covariate estimator, and the weighted regression estimator. Each augmentation weighting method is discussed with a focus on how survey weights are integrated at various stages, clarifying their applications in the survey setting. Section 3.5 presents the closed-form sandwich variance estimation for the augmented estimators for computationally efficient inference. This section primarily focuses on the development of estimators under the retrospective case, consistent with our previous notation, while modifications required for prospective designs are outlined in the Web Appendix A.6.

#### 3.1 Augmented weighting estimating equations for survey observational data

Following the framework outlined in §13.5 of Tsiatis [28] for defining the class of influence functions for regular asymptotically linear estimators, we can extend the asymptotic properties of augmented estimators[27] that incorporate survey weights. In Web Appendix A.4, we prove the proposition below that guarantees the local efficiency of the augmented estimator after incorporating the survey weights.

**Proposition 3.** If the survey weights and the propensity scores model are known, the class of observed sample data influence functions for regular asymptotically linear estimators of population estimand is,

$$\tau_h^{\text{AW}} = \left\{ \frac{h(e_{\text{sp}}(\mathbf{X}))/P(S = 1 | \mathbf{X})}{\Psi} \left[ \frac{ZY}{e_{\text{fp}}(\mathbf{X})} - \frac{(1-Z)Y}{1 - e_{\text{fp}}(\mathbf{X})} \right] - \tau_h \right\} + \{Z - e_{\text{fp}}(\mathbf{X})\}\varphi(\mathbf{X}),$$

for any function  $\varphi(\mathbf{X})$ .

Here  $\Psi = E[h(e_{\text{sp}}(\mathbf{X}))/P(S = 1 | \mathbf{X}) | S = 1]$  is the normalization term and we use  $h(e_{\text{sp}}(\mathbf{X}))$  to explicitly highlight that the tilting function  $h(\mathbf{X})$  can depend on the population propensity score  $e_{\text{sp}}(\mathbf{X})$ , as illustrated in Table 1. Based on this proposition, one can follow the argument in Tsiatis and show that, among all estimators whose influence functions fall within this class, the augmented estimator with

$$\varphi(\mathbf{X}) = -\frac{h(e_{\text{sp}}(\mathbf{X}))/p(S = 1 | \mathbf{X})}{\Psi} \left[ \frac{E[Y | Z = 1, \mathbf{X}, S = 1]}{e_{\text{fp}}(\mathbf{X})} + \frac{E[Y | Z = 0, \mathbf{X}, S = 1]}{1 - e_{\text{fp}}(\mathbf{X})} \right] \quad (9)$$

where  $E[Y | \mathbf{X}, Z = 1, S = 1]$  and  $E[Y | \mathbf{X}, Z = 0, S = 1]$  are modeled by correctly specified outcome models, achieves the smallest asymptotic variance.[28] This theoretical guarantee allows us to obtain the augmented weighting estimating equation tailored to survey observational data:

$$\frac{1}{\Psi} \frac{1}{n} \sum_{i=1}^n \left\{ \frac{h(\mathbf{X}_i)}{P(S_i = 1 | \mathbf{X}_i)} (m_1(\mathbf{X}_i) - m_0(\mathbf{X}_i)) + \frac{Z_i h(\mathbf{X}_i)}{e_{\text{sp}}(\mathbf{X}_i) p_1(\mathbf{X}_i)} (Y_i - m_1(\mathbf{X}_i)) - \frac{(1 - Z_i) h(\mathbf{X}_i)}{(1 - e_{\text{sp}}(\mathbf{X}_i)) p_0(\mathbf{X}_i)} (Y_i - m_0(\mathbf{X}_i)) - \tau_h^{\text{AW}} \right\} = 0, \quad (10)$$

where  $m_1(\mathbf{X}) = E[Y(1) | \mathbf{X}] = E[Y | \mathbf{X}, Z = 1, S = 1]$  represents the conditional expectation of the outcome given covariates for the treatment group in the survey sample. The second term in Equation (10) relies on the weak exchangeability assumption for both  $Z$  and  $S$ , as discussed in Section 2.1. Similarly,  $m_0(\mathbf{X}) = E[Y(0) | \mathbf{X}] = E[Y | \mathbf{X}, Z = 0, S = 1]$  is the conditional expectation for the control group. This formulation enables the estimation of the treatment effect between potential outcomes in the population based on observed data under the assumption of weak exchangeability. By plugging in  $\Psi$  in Equation (3) as  $\Psi = E[Z\omega_{1,p}(\mathbf{X}) | S = 1] = E[(1 - Z_i)\omega_{0,p}(\mathbf{X}) | S = 1] = E[h(\mathbf{X})/p(S = 1 | \mathbf{X}) | S = 1]$ , the general solution to this augmented weighting estimating equation becomes,

$$\hat{\tau}_h^{\text{AW}} = \frac{\sum_{i=1}^n h(\mathbf{X}_i)/P(S_i = 1 | \mathbf{X}_i) (\hat{m}_1(\mathbf{X}_i) - \hat{m}_0(\mathbf{X}_i))}{\sum_{i=1}^n h(\mathbf{X}_i)/P(S_i = 1 | \mathbf{X}_i)} \quad (11)$$

$$+ \frac{\sum_{i=1}^n Z_i \omega_{1,p}(\mathbf{X}_i) (Y_i - \hat{m}_1(\mathbf{X}_i))}{\sum_{i=1}^n Z_i \omega_{1,p}(\mathbf{X}_i)} - \frac{\sum_{i=1}^n (1 - Z_i) \omega_{0,p}(\mathbf{X}_i) (Y_i - \hat{m}_0(\mathbf{X}_i))}{\sum_{i=1}^n (1 - Z_i) \omega_{0,p}(\mathbf{X}_i)} \quad (12)$$

where  $(\omega_{1,p}(\mathbf{X}_i), \omega_{0,p}(\mathbf{X}_i))$  are balancing weights pair for survey observational data, as defined in equation (4) in Section 2.2. In retrospective studies, when the marginalized survey weights

$P(S_i = 1 | \mathbf{X}_i)$  are not directly accessible, they can be approximated using  $r_z(\mathbf{X}_i)p_z(\mathbf{X}_i) = P(S_i = 1 | \mathbf{X}_i)$  for each individual with  $z_i = 1$  or 0.

This estimator is a locally efficient estimator for  $\tau_h$ , and the details of the proof can be found in Web Appendix A.4. The consistency of  $\hat{\tau}_h^{\text{AW}}$  is contingent on the inclusion of the population-level propensity score  $e_{\text{sp}}(\mathbf{X}; \hat{\boldsymbol{\beta}})$  in the tilting function  $h(\mathbf{X})$  in the first term of Equation (11). If the population-level propensity score model is misspecified with probability limit defined as  $e_{\text{sp}}(\mathbf{X}; \tilde{\boldsymbol{\beta}})$ , the estimator will inevitably be biased for the original weighted average treatment effect estimand but still consistent for  $\tilde{\tau}_h = E \left[ h \left( e_{\text{sp}}(\mathbf{X}; \tilde{\boldsymbol{\beta}}) \right) \tau(\mathbf{X}) \right] / E \left[ h(e_{\text{sp}}(\mathbf{X}; \tilde{\boldsymbol{\beta}})) \right]$ , which represents the average treatment effect in a shifted target population determined by the shifted tilting function  $h(e_{\text{sp}}(\mathbf{X}; \tilde{\boldsymbol{\beta}}))$ . Generalizing from Mao et al.,[27] we have the following proposition (detailed derivations in Web Appendix A.4) .

**Proposition 4.** In the survey setting, the augmented weighting estimator  $\hat{\tau}_h^{\text{AW}}$  remains consistent for  $\tau_h$  provided that the population-level propensity score model is correctly specified, irrespective of the accuracy of the outcome models. When the population-level propensity score model  $e_{\text{sp}}(\mathbf{X})$  is misspecified but the outcome models are accurate, this estimator consistently estimates  $\tilde{\tau}_h$ , which only equals to  $\tau_h$  when the tilting function  $h(\mathbf{X})$  does not depend on  $e_{\text{sp}}(\mathbf{X})$ .

In the following, we introduce three variants of the augmented weighting estimators that are applicable to the survey observational data, all connected to and motivated from Equation (11).

### 3.2 One-step moment estimator

The first augmented estimator is the moment estimator (MOM). Under the balancing weights framework, the moment estimator can be adapted for any specified target population defined by covariate density  $g(\mathbf{X})$  or tilting function  $h(\mathbf{X})$ . [27] We now extend this traditional augmented estimator to the survey setting by augmenting the propensity score weighting estimator in Equation (3) with the outcome regression model. We denote the regression model as the conditional expectation of the outcome given covariates over treatment and control group in the observed sample as  $m_1^{\text{MOM}}(\mathbf{X}; \alpha_1)$  and  $m_0^{\text{MOM}}(\mathbf{X}; \alpha_0)$ . Here,  $\alpha_1$  and  $\alpha_0$  are vectors of parameters estimated from the observed data in each arm, representing the relationship between covariates  $\mathbf{X}$  and the expected outcome in the treatment and control groups, respectively. By plugging in  $m_1^{\text{MOM}}(\mathbf{X}; \hat{\alpha}_1)$  and  $m_0^{\text{MOM}}(\mathbf{X}; \hat{\alpha}_0)$  into  $\hat{\tau}_h^{\text{AW}}$ , the moment estimator can be subsequently expressed as,

$$\begin{aligned} \hat{\tau}_h^{\text{MOM}} = & \frac{\sum_{i=1}^n h(\mathbf{X}_i) / P(S_i = 1 | \mathbf{X}_i) [m_1^{\text{MOM}}(\mathbf{X}_i; \hat{\alpha}_1) - m_0^{\text{MOM}}(\mathbf{X}_i; \hat{\alpha}_0)]}{\sum_{i=1}^n h(\mathbf{X}_i) / P(S_i = 1 | \mathbf{X}_i)} \\ & + \frac{\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i) Z_i [Y_i - m_1^{\text{MOM}}(\mathbf{X}_i; \hat{\alpha}_1)]}{\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i) Z_i} - \frac{\sum_{i=1}^n \omega_{0,p}(\mathbf{X}_i) (1 - Z_i) [Y_i - m_0^{\text{MOM}}(\mathbf{X}_i; \hat{\alpha}_0)]}{\sum_{i=1}^n \omega_{0,p}(\mathbf{X}_i) (1 - Z_i)} \end{aligned} \quad (13)$$

In this moment estimator, survey weights must be incorporated in three different places. First, survey weights are used to estimate the population-level propensity score,  $\hat{e}_{\text{sp}}(\mathbf{X}_i; \hat{\boldsymbol{\beta}})$ , via weighted logistics regression. Second, the estimated  $\hat{e}_{\text{sp}}(\mathbf{X}_i; \hat{\boldsymbol{\beta}})$  is utilized to construct

the tilting function  $h(\mathbf{X}_i) = h(\widehat{e}_{\text{sp}}(\mathbf{X}_i; \widehat{\boldsymbol{\beta}}))$ , which defines the target population estimand, as illustrated in Table 1. Third, survey weights are further incorporated into the construction of the balancing weight pair,  $\omega_{1,p}(\mathbf{X}_i) = h(\widehat{e}_{\text{sp}}(\mathbf{X}_i; \widehat{\boldsymbol{\beta}})) / (p_1(\mathbf{X}_i) \widehat{e}_{\text{sp}}(\mathbf{X}_i; \widehat{\boldsymbol{\beta}}))$  and  $\omega_{0,p}(\mathbf{X}_i) = h(\widehat{e}_{\text{sp}}(\mathbf{X}_i; \widehat{\boldsymbol{\beta}})) / (p_0(\mathbf{X}_i) (1 - \widehat{e}_{\text{sp}}(\mathbf{X}_i; \widehat{\boldsymbol{\beta}})))$ , effectively reweighting the sample to represent the target population through the tilting function  $h(\widehat{e}_{\text{sp}}(\mathbf{X}_i; \widehat{\boldsymbol{\beta}}))$  and additional multiplication with the sampling probabilities  $p_z(\mathbf{X}_i)$ . Here, the conditional expectations,  $m_1^{\text{MOM}}(\mathbf{X}_i; \alpha_1)$  and  $m_0^{\text{MOM}}(\mathbf{X}_i; \alpha_0)$ , are regression models applied within the sample to recover the missing potential outcomes. Detailed proof for the consistency of  $\widehat{\tau}_h^{\text{MOM}}$  is in Web Appendix A.5. Equation 13 can be decomposed into two components:  $\widehat{\tau}_h^{\text{MOM}} = \widehat{\mu}_1^{\text{MOM}} - \widehat{\mu}_0^{\text{MOM}}$ , where

$$\widehat{\mu}_1^{\text{MOM}} = \frac{\sum_{i=1}^n h(\mathbf{X}_i) / P(S_i = 1 | \mathbf{X}_i) m_1^{\text{MOM}}(\mathbf{X}_i; \widehat{\alpha}_1)}{\sum_{i=1}^n h(\mathbf{X}_i) / P(S_i = 1 | \mathbf{X}_i)} + \frac{\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i) Z_i [Y_i - m_1^{\text{MOM}}(\mathbf{X}_i; \widehat{\alpha}_1)]}{\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i) Z_i}, \quad (14)$$

and  $\widehat{\mu}_0^{\text{MOM}}$  is the counterpart for the control group. Each of  $\widehat{\mu}_1^{\text{MOM}}$  and  $\widehat{\mu}_0^{\text{MOM}}$  is characterized by a regression model enhanced by weighted residuals for each group. Regarding the augmentation properties of  $\widehat{\tau}_h^{\text{MOM}}$ , consider the expression for  $\widehat{\mu}_1^{\text{MOM}}$ , which has the following asymptotic approximation,

$$\widehat{\mu}_1^{\text{MOM}} \xrightarrow{p} \frac{E[h(\mathbf{X})Y(1)]}{E[h(\mathbf{X})]} + \frac{1}{E[h(\mathbf{X})]} E \left[ h(\mathbf{X}) \left( \frac{Z}{e_{\text{sp}}(\mathbf{X})} - 1 \right) (Y(1) - m_1^{\text{MOM}}(\mathbf{X}; \alpha_1)) \right] \quad (15)$$

The second term in this equation vanishes when either the postulated regression model  $m_1^{\text{MOM}}(\mathbf{X}; \alpha_1)$  or the survey-weighted propensity score model  $e_{\text{sp}}(\mathbf{X}; \boldsymbol{\beta})$  is correctly specified. When the tilting function  $h(\mathbf{X})$  is specified as a constant proportional function, such as  $h(\mathbf{X}) \propto 1$ , it corresponds to inverse probability weighting and defines the entire population as the target population. The estimand in this case is the population average treatment effect (PATE). Under this configuration, this moment estimator  $\widehat{\tau}_h^{\text{MOM}}$  exhibits double robustness, meaning that consistency is guaranteed if either the outcome model  $m_1^{\text{MOM}}(\mathbf{X}; \alpha_1)$  or the propensity score model  $e_{\text{sp}}(\mathbf{X}; \boldsymbol{\beta})$  is correctly specified. Similarly, when  $h(\mathbf{X})$  corresponds to trimming functions, it depends directly on  $\mathbf{X}$  and does not rely on the correct specification of  $e_{\text{sp}}(\mathbf{X}; \boldsymbol{\beta})$ . In such cases,  $\widehat{\tau}_h^{\text{MOM}}$  retains double robustness.

In more general cases, when the tilting function  $h(\mathbf{X})$  explicitly depends on  $e_{\text{sp}}(\mathbf{X})$ , as illustrated in Table 1, the estimator  $\widehat{\mu}_1^{\text{MOM}}$  exhibits only single robustness. In this context, consistency is ensured only if the propensity score model  $e_{\text{sp}}(\mathbf{X}; \boldsymbol{\beta})$  is correctly specified, which allows for unbiased estimation of the first term in Equation 15. However, as demonstrated by Mao et al.,[27] integrating an outcome model into the weighting estimator can improve efficiency compared to weighting alone. Even in cases where  $e_{\text{sp}}(\mathbf{X}; \boldsymbol{\beta})$  is misspecified, leveraging  $m_1^{\text{MOM}}(\mathbf{X}; \alpha_1)$  can help reduce residual variance, thereby enhancing the estimator's precision.

### 3.3 Clever covariate regression estimator

An alternative strategy for constructing an augmented estimator involves incorporating functions of estimated propensity scores into the outcome regression model as an additional covariate. Scharfstein et al.[29] introduced a specific version of this approach by adding the

inverse propensity score into outcome models. Later, Bang et al.[25] demonstrated that this estimator is essentially a solution to the augmented weighting estimating equation, making the inclusion of a single propensity-score-based covariate sufficient to achieve double robustness for augmented IPW estimators. This method is often referred to as the “clever covariate” estimator (CVR) because the additional covariate “cleverly” adjusts for potential misspecification of the outcome model by introducing a coarse but sufficient representation of all covariates, thereby ensuring robustness. Moreover, as noted by Robins et al., [26] when the CVR is structured with an identity link, it aligns with the targeted maximum likelihood estimation (TMLE) for marginal means over the target population, using, in our case, population-level survey-weighted balancing weights as covariates to re-target and correct the original naive outcome model to better estimate potential outcomes in the target population.

Under the context of survey observational data, by solving the augmented weighting estimating equation10 and replacing the conventional outcome models  $m_1(\mathbf{X}_i)$  and  $m_0(\mathbf{X}_i)$  with generalized linear regression models  $m_1^{\text{CVR}}(\mathbf{X}; \hat{\boldsymbol{\alpha}}_1^{\text{CVR}}) = g^{-1}\{m_1(\mathbf{X}; \hat{\alpha}_1) + \hat{\phi}_1 \omega_{1,p}(\mathbf{X})\}$  and  $m_0^{\text{CVR}}(\mathbf{X}; \hat{\boldsymbol{\alpha}}_0^{\text{CVR}}) = g^{-1}\{m_0(\mathbf{X}; \hat{\alpha}_0) + \hat{\phi}_0 \omega_{0,p}(\mathbf{X})\}$ , where  $g$  is the canonical link function, we can construct the clever covariate estimator adapted to survey observational data as follows:

$$\begin{aligned} & \hat{\tau}_h^{\text{CVR}} \\ &= \hat{\tau}_1^{\text{CVR}} - \hat{\tau}_0^{\text{CVR}} \tag{16} \\ &= \frac{\sum_{i=1}^n h(\mathbf{X}_i)/p(S_i = 1 | \mathbf{X}_i) m_1^{\text{CVR}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_1^{\text{CVR}})}{\sum_{i=1}^n h(\mathbf{X}_i)/p(S_i = 1 | \mathbf{X}_i)} + \frac{\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i, \hat{\boldsymbol{\beta}}) Z_i \left( Y_i - m_1^{\text{CVR}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_1^{\text{CVR}}) \right)}{\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i; \hat{\boldsymbol{\beta}}) Z_i} \\ &- \frac{\sum_{i=1}^n h(\mathbf{X}_i)/p(S_i = 1 | \mathbf{X}_i) m_0^{\text{CVR}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_0^{\text{CVR}})}{\sum_{i=1}^n h(\mathbf{X}_i)/p(S_i = 1 | \mathbf{X}_i)} - \frac{\sum_{i=1}^n \omega_{0,p}(\mathbf{X}_i, \hat{\boldsymbol{\beta}}) (1 - Z_i) \left( Y_i - m_0^{\text{CVR}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_0^{\text{CVR}}) \right)}{\sum_{i=1}^n \omega_{0,p}(\mathbf{X}_i; \hat{\boldsymbol{\beta}}) (1 - Z_i)} \end{aligned}$$

The second terms of  $\hat{\tau}_1^{\text{CVR}}$  and  $\hat{\tau}_0^{\text{CVR}}$  are essentially the score functions of  $m_1^{\text{CVR}}(\mathbf{X}; \hat{\boldsymbol{\alpha}}_1^{\text{CVR}})$  and  $m_0^{\text{CVR}}(\mathbf{X}; \hat{\boldsymbol{\alpha}}_0^{\text{CVR}})$  acting as the weighted residuals of first terms of each, and they become zero when they are correctly specified, simplifying  $\hat{\tau}_h^{\text{CVR}}$  to the following form:

$$\hat{\tau}_h^{\text{CVR}} = \frac{\sum_{i=1}^n h(\mathbf{X}_i)/p(S_i = 1 | \mathbf{X}_i) \left( m_1^{\text{CVR}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_1^{\text{CVR}}) - m_0^{\text{CVR}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_0^{\text{CVR}}) \right)}{\sum_{i=1}^n h(\mathbf{X}_i)/p(S_i = 1 | \mathbf{X}_i)} \tag{17}$$

Here, by substituting the expression of balancing weights pair  $(\omega_{1,p}(\mathbf{X}_i), \omega_{0,p}(\mathbf{X}_i))$  as the clever covariates for each arm, we have

$$\begin{aligned} m_1^{\text{CVR}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_1^{\text{CVR}}) &= g^{-1} \left\{ m_1(\mathbf{X}_i; \hat{\alpha}_1) + \hat{\phi}_1 h(\mathbf{X}_i) / (P(S_i = 1 | Z_i = 1, \mathbf{X}_i) e_{\text{sp}}(\mathbf{X}_i; \hat{\boldsymbol{\beta}})) \right\} \\ \text{and } m_0^{\text{CVR}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_0^{\text{CVR}}) &= g^{-1} \left\{ m_0(\mathbf{X}_i; \hat{\alpha}_0) + \hat{\phi}_0 h(\mathbf{X}_i) / (P(S_i = 1 | Z_i = 0, \mathbf{X}_i) (1 - e_{\text{sp}}(\mathbf{X}_i; \hat{\boldsymbol{\beta}}))) \right\}. \end{aligned}$$

In this estimator, survey weights need to be incorporated three times in the clever covariate

method: during survey-weighted propensity score regression for  $e_{\text{sp}}(\mathbf{X}_i; \hat{\boldsymbol{\beta}})$ , when formulating balancing weights as clever covariates, and in standardizing the outcome predictions toward the target estimands.

The clever covariate estimator is asymptotically equivalent to the moment estimator in that both are solutions to the augmented weighting estimating equation and share the same influence function. When the tilting function  $h(\mathbf{X}) \propto 1$ , the clever covariate method becomes equivalent to the augmented IPW estimator, which exhibits double robustness, ensuring consistency if either the outcome model (without clever covariates)  $m_z(\mathbf{X}; \hat{\boldsymbol{\alpha}}_z)$  or the propensity score model  $e_{\text{sp}}(\mathbf{X}; \hat{\boldsymbol{\beta}})$  is correctly specified. In more general cases, when  $h(\mathbf{X})$  explicitly depends on  $e_{\text{sp}}(\mathbf{X})$ , it may improve efficiency over simple weighting estimators discussed in Section 2. However, this requires the correct specification of  $e_{\text{sp}}(\mathbf{X}; \hat{\boldsymbol{\beta}})$ ; otherwise, the estimator converges to a biased weighted average treatment effect estimand  $\tilde{\tau}_h$ , where the weights depend on the misspecified tilting function  $h(e_{\text{sp}}(\mathbf{X}; \tilde{\boldsymbol{\beta}}))$ .

In the absence of survey sampling, Kang and Schafer [30] pointed out that, when the outcome models  $m_1(\mathbf{X}; \hat{\boldsymbol{\alpha}}_1)$  and  $m_0(\mathbf{X}; \hat{\boldsymbol{\alpha}}_0)$  are already correctly specified, the clever covariates as coarse summaries of the covariates may lead to slight overfitting by adding mean-zero noise through weighted residuals. When the survey-weighted propensity score model is correctly specified, the clever covariate method ensures consistent estimation of target parameters, provided that the conditional mean of the outcome varies smoothly with both the survey-weighted propensity score and the sampling probabilities. In theory, this smooth variation remains essential for maintaining the consistency and robustness of the clever covariate method, even in complex survey designs. However, incorporating survey weights twice in constructing the clever covariates may potentially exacerbate the extrapolation problem noted by Robins et al.[26] because the inflation of extreme value can lead to finite-sample bias, making the clever covariate estimator less stable compared to the weighted regression methods introduced in Section 3.4.

### 3.4 Weighted regression estimator

A third regression-based augmented estimator is the weighted regression estimator (WET)[31, 26, 32, 33] that, in our case, utilizes balancing weights during outcome regression process. The weighted regression estimator can directly incorporate both propensity scores and survey weights. It enables effective bias correction and aligns the estimated coefficients more closely with the true population parameters. [26, 30] These properties are especially advantageous in complex data environments where both models are prone to misspecification and the sampling process is not necessarily independent of exposure and treatment assignment. Adapted to our setting of survey sampling, the weighted regression estimator can be expressed as,

$$\hat{\tau}_h^{\text{WET}} = \frac{\sum_{i=1}^n h(\mathbf{X}_i)/p(S_i = 1 | \mathbf{X}_i) \left( m_1^{\text{WET}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_1^{\text{WET}}) - m_0^{\text{WET}}(\mathbf{X}_i; \hat{\boldsymbol{\alpha}}_0^{\text{WET}}) \right)}{\sum_{i=1}^n h(\mathbf{X}_i)/p(S_i = 1 | \mathbf{X}_i)} \quad (18)$$

where  $m_1^{\text{WET}}(\mathbf{X}; \hat{\boldsymbol{\alpha}}_1^{\text{WET}})$  and  $m_0^{\text{WET}}(\mathbf{X}; \hat{\boldsymbol{\alpha}}_0^{\text{WET}})$  are the outcome regression models with canonical link functions, similar to those in the clever covariate estimator, but using weighted

regression with balancing weights  $\omega_{1,p}(\mathbf{X})$  and  $\omega_{0,p}(\mathbf{X})$ . In this estimator, survey weights are incorporated in three distinct stages: during the survey-weighted propensity score regression for  $e_{\text{sp}}(\mathbf{X}_i; \hat{\boldsymbol{\beta}})$ , for the construction of the balancing weight pairs  $\omega_{1,p}(\mathbf{X}_i)$  and  $\omega_{0,p}(\mathbf{X}_i)$ , which subsequently serve as regression weights in the outcome modeling process, and in standardizing the predicted potential outcomes toward the target population. This strategy adjusts the weighted least squares coefficients  $\hat{\boldsymbol{\alpha}}_1^{\text{WET}}$  and  $\hat{\boldsymbol{\alpha}}_0^{\text{WET}}$  by accounting for the mechanism between  $\mathbf{X}$  and potential outcomes under treatment and control conditions respectively in the target population, as long as the SUTVA and weak exchangeability assumptions in Section 2.1 hold. For ease of reference, Table 2 summarizes where survey weights need to be applied for each simple weighting estimator and augmented weighting estimator.

Similar to the moment and clever covariate estimators by sharing the same influence function, the weighted regression estimator ensures consistent estimation when the survey-weighted propensity score model  $e_{\text{sp}}(\mathbf{X}; \hat{\boldsymbol{\beta}})$  is correctly specified, with additional robustness depending on the definition of the tilting function. When the tilting function  $h(\mathbf{X}) \propto 1$ , the weighted regression estimator is doubly robust, leading to consistent estimation if either the weighted outcome regression model or the survey-weighted propensity score model  $e_{\text{sp}}(\mathbf{X}; \hat{\boldsymbol{\beta}})$  is correctly specified. In cases where  $h(\mathbf{X})$  explicitly depends on  $e_{\text{sp}}(\mathbf{X}; \hat{\boldsymbol{\beta}})$ , incorporating the weighted outcome regression models can enhance efficiency relative to the simple weighting estimators discussed in Section 2. Consistency to the originally intended target estimand in this case is achieved only if  $e_{\text{sp}}(\mathbf{X}; \hat{\boldsymbol{\beta}})$  is correctly specified; otherwise, the estimator will converge to a biased target estimand  $\tilde{\tau}_h$ , determined by the misspecified tilting function  $h(e_{\text{sp}}(\mathbf{X}; \hat{\boldsymbol{\beta}}))$ .

Compared to the clever covariate estimator, the weighted regression estimator may offer better performance in scenarios involving highly variable survey weights or extreme propensity scores. As discussed by Kang et al. [30] and Robins et al., [26] the weighted regression estimator is more stable under extreme balancing weights, reducing the risk of overfitting that may occur under the clever covariate construction. However, under the IPW scheme, the weighted regression estimator remains susceptible to the impact of extreme inverse propensity scores, which can lead to inflated variance and finite-sample bias, particularly when these extreme weights dominate the regression model. We plan to examine to what extent these observations carry forward to the survey observational study settings in Section 4.

### 3.5 Sandwich variance estimation for augmented estimator under survey observational data

According to Proposition 3, all three augmented estimators are derived from the solution to the augmented weighting estimating equation 11. This allows us to construct a unified empirical sandwich variance estimator using M-estimation theory, [34] which enables computationally efficient solution for variance and confidence interval estimation under the context of survey observational data.

Specifically, the generalized form of the augmented estimator in Equation 11 can be expressed as  $\hat{\tau}_h^{\text{AW}} = \hat{v}_1 + \hat{v}_2 - \hat{v}_3$ . Adapting the empirical sandwich variance method [7, 9, 27] based on M-estimation theory, [29] we construct the estimating equation 19 for augmented estimators under the context of survey observational data. This estimating equation adjusts

Table 2: Summary of components that incorporate survey weights in propensity score weighting and augmented estimators under the context of retrospective studies using survey observational data.

Estimator	Estimating propensity scores	Constructing the balancing weights	Fitting the outcome regression model	Standardizing predicted potential outcomes to the target estimand
PSW	✓	✓	Not applicable	Not applicable
MOM	✓	✓	×	✓
CVR	✓	✓	✓ (through additional covariates)	✓
WET	✓	✓	✓ (through weighted regression)	✓

Note: For PSW, survey weights are used to estimate the propensity score  $e_{sp}(\mathbf{X}_i; \hat{\boldsymbol{\beta}})$  and construct the balancing weights  $\omega_{z,p}(\mathbf{X}_i) = h(e_{sp}(\mathbf{X}_i; \hat{\boldsymbol{\beta}})) / (p_z(\mathbf{X}_i) e_{sp}(\mathbf{X}_i; \hat{\boldsymbol{\beta}}))$ ; for MOM, survey weights are similarly used for  $e_{sp}(\mathbf{X}_i; \hat{\boldsymbol{\beta}})$  and  $\omega_{z,p}(\mathbf{X}_i)$ , but the outcome regression does not include survey weights; for CVR, balancing weights  $\omega_{z,p}(\mathbf{X}_i)$  are included as additional covariates in the outcome regression; for WET, balancing weights  $\omega_{z,p}(\mathbf{X}_i)$  are incorporated as regression weights, introducing survey weights directly into the outcome regression process.

for the uncertainty in estimating the survey-weighted propensity score and two potential outcome models by utilizing score functions corresponding to  $\boldsymbol{\theta} = (v_1, v_2, v_3, \boldsymbol{\alpha}_0^T, \boldsymbol{\alpha}_1^T, \boldsymbol{\beta}_{fp}^T, \boldsymbol{\beta}_{sp}^T)^T$ , where  $v_1$ ,  $v_2$ , and  $v_3$  are the components in true value of the general augmented estimator  $\tau_h^{AW}$ ,  $\boldsymbol{\alpha}_0$  and  $\boldsymbol{\alpha}_1$  are the parameters of the outcome models  $m_0^{AW}(\mathbf{X}; \boldsymbol{\alpha}_0)$  and  $m_1^{AW}(\mathbf{X}; \boldsymbol{\alpha}_1)$ , respectively, and  $\boldsymbol{\beta}_{fp}$  and  $\boldsymbol{\beta}_{sp}$  represent the parameters of the sample-level and population-level propensity score models, respectively. Therefore, the estimating equation for the augmented estimators under the context of survey observational data is:

$$\mathbf{0} = \sum_{i=1}^n \boldsymbol{\Psi}_{\boldsymbol{\theta}}(\mathbf{X}_i, Y_i, Z_i) = \sum_{i=1}^n \begin{bmatrix} \psi_{PS,FP} \\ \psi_{PS,SP} \\ \psi_{OR,1} \\ \psi_{OR,0} \\ \psi_{v1} \\ \psi_{v2} \\ \psi_{v3} \end{bmatrix} = \sum_{i=1}^n \begin{bmatrix} \omega_{PS,FP}(\mathbf{X}_i) \mathbf{X}_i (Z_i - e_{fp}(\mathbf{X}_i; \boldsymbol{\beta}_{fp})) \\ \omega_{PS,SP}(\mathbf{X}_i^*) \mathbf{X}_i^* (Z_i - e_{sp}(\mathbf{X}_i^*; \boldsymbol{\beta}_{sp})) \\ \omega_{OR,1}(\mathbf{X}_i^*) Z_i \mathbf{X}_i' (Y_i - m_1^{AW}(\mathbf{X}_i^*; \boldsymbol{\alpha}_1)) \\ \omega_{OR,0}(\mathbf{X}_i^*) (1 - Z_i) \mathbf{X}_i'' (Y_i - m_0^{AW}(\mathbf{X}_i^*; \boldsymbol{\alpha}_0)) \\ h_p(\mathbf{X}_i^*) (m_1^{AW}(\mathbf{X}_i^*; \boldsymbol{\alpha}_1) - m_0^{AW}(\mathbf{X}_i^*; \boldsymbol{\alpha}_0) - v_1) \\ \omega_{1,p}(\mathbf{X}_i^*) Z_i (Y_i - m_1^{AW}(\mathbf{X}_i^*; \boldsymbol{\alpha}_1) - v_2) \\ \omega_{0,p}(\mathbf{X}_i^*) (1 - Z_i) (Y_i - m_0^{AW}(\mathbf{X}_i^*; \boldsymbol{\alpha}_0) - v_3) \end{bmatrix} \quad (19)$$

The equation above includes six distinct components:  $\psi_{PS,FP}$ ,  $\psi_{PS,SP}$ ,  $\psi_{OR,1}$ ,  $\psi_{OR,0}$ ,  $\psi_{v1}$ ,  $\psi_{v2}$ , and  $\psi_{v3}$ , each representing different aspects of the augmented weighting estimation process. The first two components,  $\psi_{PS,FP}$  and  $\psi_{PS,SP}$ , correspond to the score functions associated with the estimation of the propensity score model at the sample and population levels, respectively. In retrospective settings,  $e_{fp}(\mathbf{X})$  is used exclusively to calculate  $r_z(\mathbf{X}_i)$ , which approximates  $P(S_i = 1 | \mathbf{X}_i) = r_z(\mathbf{X}_i) p_z(\mathbf{X}_i)$ , and is applied in the augmented estimator to tilt the imputed outcome difference. The term  $\omega_{PS,FP}(\mathbf{X}_i)$  represents the regression



weights for estimating the within-sample propensity score and is set to 1. Furthermore,  $\omega_{\text{PS,SP}}(\mathbf{X}_i^*)$  refers to the regression weights applied to the covariate vector  $\mathbf{X}_i^*$  in the population level propensity score model, which is either the survey weights when the regression is survey-weighted (W.PS) or set to 1 when the regression is unweighted (U.PS). Depending on the model specification,  $\mathbf{X}_i^*$  may consist of baseline covariates alone or specified original covariates with survey weights included as an additional covariate (C.PS), as recommended by DuGoff et al.[3], to capture unique design features and address latent sampling information, enhancing covariate balance and reducing bias from unmeasured confounders.

The components  $\psi_{\text{OR},1}$  and  $\psi_{\text{OR},0}$  represent the score functions for estimating the unobserved potential outcome models for the treated ( $Z_i = 1$ ) and control ( $Z_i = 0$ ) groups, respectively. The weights  $\omega_{\text{OR},1}(\mathbf{X}_i^*)$  and  $\omega_{\text{OR},0}(\mathbf{X}_i^*)$  depend on the choice of augmented estimator. For the moment estimator (MOM) and clever covariate estimator (CVR), these weights are set to 1. For the weighted regression estimator (WET), they correspond to the balancing weights  $\omega_{1,p}(\mathbf{X}_i^*)$  and  $\omega_{0,p}(\mathbf{X}_i^*)$  as specified in Section 3.4. The outcome models  $m_1^{\text{AW}}$  and  $m_0^{\text{AW}}$  reflect the specific type of augmentation used, where the superscript "AW" can denote "MOM", "CVR", or "WET", depending on the choice of estimator. For the clever covariate estimator, the outcome model covariates  $\mathbf{X}'_i$  and  $\mathbf{X}''_i$  include the clever covariates for each arm, whereas for MOM and WET, only the baseline covariates are included, i.e.,  $\mathbf{X}_i$ .

The component  $\psi_{v_1}$  pertains to the difference in predicted full outcomes between the treated and control models, adjusted by the tilting function  $h_p(\mathbf{X}_i^*) = h(e_{\text{sp}}(\mathbf{X}_i^*; \boldsymbol{\beta})/P(S_i = 1 | \mathbf{X}_i))$  and accounting for the first component  $v_1$  in the general augmented treatment effect estimation 11. Here,  $m_1^{\text{AW}}(\mathbf{X}'_i; \boldsymbol{\alpha}_1)$  and  $m_0^{\text{AW}}(\mathbf{X}''_i; \boldsymbol{\alpha}_0)$  represent the specified outcome models for the treated and control groups, respectively, using covariate vectors  $\mathbf{X}'_i$  and  $\mathbf{X}''_i$ . The components  $\psi_{v_2}$  and  $\psi_{v_3}$  represent the residual adjustments for the treated and control groups, respectively. These adjustments incorporate the balancing weights  $\omega_{1,p}(\mathbf{X}_i^*)$  and  $\omega_{0,p}(\mathbf{X}_i^*)$ , which are designed to adjust for confounding effects. These balancing weights are as described in Table 1 and are generalized for survey observational data under different model specifications.

These six components together form the complete estimating equation for a given augmented weighting approach under the context of survey observational data, capturing the uncertainties associated with the estimated propensity score, outcome models, and treatment effects in a comprehensive manner. This approach ensures robust estimation of treatment effects under complex survey designs by considering various augmentations and model specifications. Given that each component of the estimating equations represents unbiased estimating equations, under suitable regularity conditions,[24] it follows that for each augmented estimator we can achieve asymptotic normality  $\sqrt{n}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \xrightarrow{d} N(\mathbf{0}, \mathbf{V}(\boldsymbol{\theta}))$ , where the variance-covariance matrix  $\mathbf{V}(\boldsymbol{\theta})$  is defined as  $\mathbf{V}(\boldsymbol{\theta}) = \mathbf{A}(\boldsymbol{\theta})^{-1} \mathbf{B}(\boldsymbol{\theta}) \{\mathbf{A}(\boldsymbol{\theta})^{-1}\}^T$  with  $\mathbf{A}(\boldsymbol{\theta}) = E \left\{ -\frac{\partial \boldsymbol{\Psi}_{\boldsymbol{\theta}}(\mathbf{X}, Y, Z)}{\partial \boldsymbol{\theta}} \mid S = 1 \right\}$ , and  $\mathbf{B}(\boldsymbol{\theta}) = E \left\{ \boldsymbol{\Psi}_{\boldsymbol{\theta}}(\mathbf{X}, Y, Z) \boldsymbol{\Psi}_{\boldsymbol{\theta}}(\mathbf{X}, Y, Z)^T \mid S = 1 \right\}$ . The quantities  $\mathbf{A}(\boldsymbol{\theta})$  and  $\mathbf{B}(\boldsymbol{\theta})$  can be consistently estimated by replacing the expected values with their empirical counterparts,  $\hat{\mathbf{A}}(\hat{\boldsymbol{\theta}}) = -\frac{1}{n} \sum_{i=1}^n \partial \boldsymbol{\Psi}_{\hat{\boldsymbol{\theta}}}(\mathbf{X}_i, Y_i, Z_i) / \partial \boldsymbol{\theta}$  and  $\hat{\mathbf{B}}(\hat{\boldsymbol{\theta}}) = \frac{1}{n} \sum_{i=1}^n \boldsymbol{\Psi}_{\hat{\boldsymbol{\theta}}}(\mathbf{X}_i, Y_i, Z_i) \boldsymbol{\Psi}_{\hat{\boldsymbol{\theta}}}(\mathbf{X}_i, Y_i, Z_i)^T$  resulting in the empirical sandwich variance estimator  $\hat{\mathbf{V}}(\hat{\boldsymbol{\theta}}) = \hat{\mathbf{A}}(\hat{\boldsymbol{\theta}})^{-1} \hat{\mathbf{B}}(\hat{\boldsymbol{\theta}}) \{\hat{\mathbf{A}}(\hat{\boldsymbol{\theta}})^{-1}\}^T$ . The variance of the augmented estimator  $\hat{\tau}_h^{\text{AW}}$  can be

consistently estimated using the corresponding elements of  $\widehat{\mathbf{V}}(\widehat{\boldsymbol{\theta}})$ .

Table 3: Covariate vector notations used in the estimating equations for the sandwich variance estimation of augmented estimators under survey observational data.

Notatio	Definition and Context
$\mathbf{X}_i^*$	Covariates used in the propensity score model $e_{\text{sp}}(\mathbf{X}_i^*; \boldsymbol{\beta})$ . In unweighted (U.PS) and survey-weighted regression scenarios (W.PS), it includes only baseline covariates. When survey weights are incorporated as an additional covariate (C.PS), it contains both the specified original covariates and the survey weights.
$\mathbf{X}_i'$	Covariates in the treated outcome regression model $m_1^{\text{AW}}(\mathbf{X}_i'; \boldsymbol{\alpha}_1)$ . For CVR, this includes clever covariates; for MOM and WET, only baseline covariates are used.
$\mathbf{X}_i''$	Covariates in the control outcome regression model $m_0^{\text{AW}}(\mathbf{X}_i''; \boldsymbol{\alpha}_0)$ . Mirrors the definition of $\mathbf{X}_i'$ based on the estimator type.

Note: The specified original covariates used in the propensity score model may differ from those in the outcome regression models, and the covariates used in the outcome models for each arm are not necessarily the same. In MOM and WET, where both outcome models include only baseline covariates without distinct clever covariates,  $\mathbf{X}_i'$  and  $\mathbf{X}_i''$  may be identical or even unified, using one model for both arms' counterfactual outcome predictions. In cases where the same set of covariates are used across all models, all three covariate notations ( $\mathbf{X}_i^*$ ,  $\mathbf{X}_i'$  and  $\mathbf{X}_i''$ ) in MOM and WET with U.PS or W.PS could be identical.

## 4 Simulation Studies

We conducted a simulation study to evaluate the performance of these methods under the context of survey observational data in estimating various population-level estimands, particularly under different degrees of overlap and model misspecification. Below we detail our simulation design and key findings.

### 4.1 Simulating design

Our data-generating process was adapted from the multi-stage strategy employed by Austin et al. and Lenis et al. [11, 13] We simulated a population of 1,000,000 individuals, organized into 10 strata ( $j = 1, \dots, 10$ ). Each stratum was further divided into 20 clusters, resulting in a total of 200 clusters ( $k = 1, \dots, 200$ ), with 5000 individuals in each cluster ( $i = 1, \dots, 5000$ ). For each individual, we generated six baseline covariates ( $l = 1, \dots, 6$ ) from normal distributions, where the vector of covariates for the  $i$ -th individual is denoted as  $\mathbf{X}_i = \{X_{1i}, X_{2i}, X_{3i}, X_{4i}, X_{5i}, X_{6i}\}$ . The mean of these covariates were allowed to vary at both the stratum and cluster levels. For the  $l$ -th covariate, a stratum-specific random effect for the  $j$  th stratum  $\nu_{l,j}^{\text{stratum}} \sim N(0, \sigma_l^{\text{stratum}})$  was generated, along with a cluster-specific random effect for the  $k$  -th cluster  $\nu_{l,k}^{\text{cluster}} \sim N(0, \sigma_l^{\text{cluster}})$ . The  $l$ -th covariate for the  $i$ -th individual in the  $j$ -th stratum and  $k$  -th cluster was then generated as  $X_{l,ijk} \sim N(\nu_{l,j}^{\text{stratum}} + \nu_{l,k}^{\text{cluster}}, 1)$ .

We set the standard deviation associated with stratum-level means to  $\sigma^{\text{stratum}} = 0.35$  and with cluster-level means to  $\sigma^{\text{cluster}} = 0.15$ .

We then generated a binary treatment assignment for each individual using a logistic model to determine the probability of treatment:

$$\text{logit}(e_{\text{sp}}(\mathbf{X}_i)) = a_0 + \psi (\mathbf{a}_{1:6}^T \mathbf{X}_i + a_7 X_{1i} X_{2i})$$

where  $\mathbf{a}_{1:6} = (\log(1.1), \log(1.25), \log(1.5), \log(1.75), \log(2), \log(2.5))^T$ , and  $a_7 = \log(1.1)$ . Treatment status  $Z_i$  was simulated from a Bernoulli distribution:  $Z_i \sim \text{Bernoulli}(e_{\text{sp}}(\mathbf{X}_i))$ . The scalar  $\psi$  reflects the strength of confounding and dictates the level of overlap in the propensity score. [35] We set  $\psi = 0.6$  or  $2$  to mimic scenarios with good or poor overlap in the full population, as visualized in Fig. 1(a) and Fig. 1(b), respectively. For both scenarios, the overall treatment prevalence was maintained at approximately 30% by adjusting  $a_0$  to  $\log(35/80)$  for the good overlap and  $\log(20/80)$  for the poor overlap, allowing us to isolate the impact of overlap when evaluating estimator performance.




Figure 1: Distribution of Propensity Scores in Different Overlap Scenarios

We simulated continuous outcomes for each individual  $i$  in the population using a linear combination of covariates, treatment effects, and interaction terms. Specifically, the potential outcome  $Y_i$  was modeled as:

$$Y_i = b_0 + \delta_0 (\mathbf{b}^T \mathbf{X}_i + b_7 X_{1i} X_{2i}) + Z_i (\delta_1 + \delta_2 (\mathbf{b}^T \mathbf{X}_i + b_8 X_{1i} X_{2i})) + \varepsilon$$

where  $\varepsilon \sim N(0, 1)$ . We set  $b_0 = 0$  and  $\mathbf{b} = (2.5, -2, 1.75, -1.25, 1.5, 1.1)^T$  for the covariate coefficients. The average treatment effect was represented by  $\delta_1 = 1$ , while  $\delta_0 = 0.3$  and  $\delta_2 = 0.2$  introduced heterogeneity in the control and treatment conditions, respectively. Additionally, two interaction terms were included, with  $b_7 = 2.5$  and  $b_8 = 1.5$ , to capture the more complex influence of baseline covariates on outcomes across both conditions.

Once we simulated data for the superpopulation, we drew 1,000 random samples with 5000 individuals each (representing 0.5% of the overall population) using a multi-stage sampling design, where the sampling mechanism was conditionally independent of the treatment assignment mechanism. Sample sizes were allocated across the 10 strata as  $\{850, 750, 700, 650, 600,$

400, 350, 300, 250, 150}, ensuring that the observed sample would have a different distribution from the population. Within each stratum, five clusters were selected using simple random sampling, and from each selected cluster, an equal number of individuals were sampled. For each sampled individual, we calculated the true survey weights by taking the inverse of the sampling probability, representing the number of individuals in the population that the sampled individual represents.

To validate the robustness of our estimator under different sampling mechanisms, we further extended the simulation by introducing an alternative sampling mechanism dependent on both baseline covariates and treatment assignment. We used logistic regression to model the sampling probability for individual  $i$  as a function of both the treatment assignment indicator  $Z_i$  and covariates  $\mathbf{X}_i$ , allowing sampling to reflect more complex real-world dependencies. The sampling probability  $p_{z,i} = P(S_i = 1 \mid Z_i, \mathbf{X}_i)$  was computed as  $\text{logit}(p_{z,i}) = c_0 + \delta_z^{(s)} Z_i + \mathbf{c}^T \mathbf{X}_i$ , where  $c_0 = \log\left(\frac{0.005}{1-0.005}\right)$  defines the baseline sampling rate of 0.5% consistent with the above multi-stage design, and  $\delta_z^{(s)} = \log(0.9)$  slightly reduces the sampling probability for the treatment group ( $Z = 1$ ), addressing possible treatment-related biases. The vector  $\mathbf{c} = (c_1, c_2, c_3, c_4, c_5, c_6)^T$  represents the covariate coefficients with values  $c_1 = \log(1.05)$ ,  $c_2 = \log(1.10)$ ,  $c_3 = \log(1.15)$ ,  $c_4 = \log(1.10)$ ,  $c_5 = \log(1.05)$ , and  $c_6 = \log(1.10)$ , corresponding to the influences of covariates  $X_1$  through  $X_6$  on the selection probability. To avoid the impact of near-zero values leading to extreme weights in further estimation, we will perform and recommend truncation with  $\alpha = 0.001$  on the fitted propensity scores after incorporating the varied survey weights in this settings. The sampling indicator  $S_i$  was subsequently generated as a Bernoulli random variable with probability  $p_{z,i}$  for individual  $i$ , capturing the interaction between treatment assignment and covariates in determining sample inclusion. More details and results of this alternative simulation are presented in Web Appendix B.2.

For each sample within each scenario, we evaluated four types of estimators: the PSW (propensity score weighting estimator) and three augmentation methods, referred to as MOM (moment estimator), CVR (clever covariate regression estimator), and WET (Weighted Regression estimator), as previously described. To investigate the role of survey weights in estimating the propensity scores, we primarily compare two strategies for integrating survey weights in the propensity score model: as model weights (W.PS) and as an additional covariate (C.PS). The latter involves directly adjusting for survey weight as one additional single covariate in the propensity score model, which is distinct from including the balancing weight as a clever covariate in the outcome regression model. Additionally, results for unweighted propensity scores (U.PS) and survey weights applied as both model weights and covariates (CW.PS) are presented in the Web Appendix B.1. These comparisons may shed light on previous debates in the literature regarding which level of covariate balance should we achieve with propensity score and in what way.[3, 4]

In total, we constructed eight estimators by combining the four types of estimators with the two methods of incorporating survey weights into the propensity score model. These estimators were evaluated across three common estimands: the population average treatment effect (PATE), the population average treatment effect for the treated, (PATT), and the population average treatment effect among the overlap population (PATO). Additionally, we assessed their performance under four model misspecification scenarios: (i) correct propensity

score model and correct outcome model (Cor|Cor), (ii) incorrect propensity score model but correct outcome model (Mis|Cor), (iii) correct propensity score model but incorrect outcome model (Cor|Mis), and (iv) both models misspecified (Mis|Mis).

Given the complexity of the design, it is instructive to note the total number of considered scenarios. We examined two levels of overlap (good vs. poor), two primary methods for incorporating survey weights into the propensity score model (W.PS vs. C.PS), four estimators (PSW, MOM, CVR, WET), three estimands (PATE, PATT, PATO), and four model specification scenarios (Cor|Cor, Mis|Cor, Cor|Mis, Mis|Mis). This combination yields  $2 \times 2 \times 4 \times 3 \times 4 = 192$  distinct configurations in the main analysis. Additional scenarios that incorporate the unweighted and combined weighting-plus-covariate propensity score approaches and the sampling mechanism depending on treatment increase this number even further. The factorial nature of this design ensures that our conclusions are drawn from a broad range of realistic conditions, and the resulting comparisons are adequately comprehensive. Table 4 outlines these factors and their levels, providing an overview of the multiplicity of scenarios explored.

Table 4: Summary of factors and number of scenarios in the main simulation design

Factor	Levels	Count
Overlap scenario	Good vs. Poor	2
Method of incorporating survey weights into PS model	W.PS, C.PS	2
Estimator	PSW, MOM, CVR, WET	4
Estimand	PATE, PATT, PATO	3
Model specification scenario	Cor Cor, Mis Cor, Cor Mis, Mis Mis	4
Total scenarios in main analysis		192

Performance metrics included relative bias, relative efficiency, and empirical coverage of the 95% confidence intervals. Relative bias was calculated as the percentage difference between the estimated and true population treatment effects, averaged across 1,000 simulations. Relative efficiency was determined as the ratio of the Monte Carlo variance of the reference estimator (the balancing weight estimator for PATE with W.PS under correct model specification) to that of each evaluated estimator. Empirical coverage was assessed by calculating the proportion of confidence intervals, constructed using sandwich variances, that contained the true population treatment effect. The simulation code and implementations of the survey-weighted balancing weight estimator and the augmented estimators for PATE, PATT, and PATO are available online at [https://github.com/ykzeng-yale/SW\\_PSW](https://github.com/ykzeng-yale/SW_PSW), and they are implemented in the PSweight R package (Version 2.1.0).

## 4.2 Simulations results

In this section, we focus on the primary four types of estimators (PSW, MOM, CVR and WET) and two distinct methods of incorporating survey weights into propensity score models (W.PS and C.PS). Table 5 presents the simulation results under the good overlap scenario,

summarizing the relative bias, relative efficiency, and coverage of PATE, PATT, and PATO under various modeling strategies with either correctly or incorrectly specified propensity score and outcome regression models. In parallel, Table 6 presents the results when the overlap is poor. In addition to these primary results, we present more comprehensive analyses that include U.PS and CW.PS in Web Tables 3 to 5 and 6 to 8 for each estimand under good and poor overlap. Furthermore, simulation results for the extended scenario, where sampling depends on treatment assignment, are provided in Web Tables 9 and 10 for W.PS and C.PS results, and Web Tables 11 to 13 and 14 to 16 for more comprehensive analyses. To facilitate the summary of findings, we focus on the following three thorny questions; that is, how survey weights should be incorporated at each stage of the balancing weight estimator, which augmented estimators perform better, and which method is more robust (i.e., under poor overlap and model misspecification).

#### 4.2.1 How should survey weights be incorporated into the propensity score weighting estimator?

Focusing first on the results from the weighting estimator alone (PSW), our simulation results support our theoretical conclusion that incorporating survey weights in both the propensity score and outcome estimation leads to unbiased and efficient propensity score weighting estimators of the population-level treatment effect. Under good overlap (Table 5), using survey-weighted propensity score models (W.PS) consistently produces lower bias, higher relative efficiency, and coverage closer to the nominal 95% compared to using survey weights as an additional covariate (C.PS). This pattern is particularly evident for PATO, where W.PS attains near-ideal balance and performance across all specification scenarios. These observations, shown in the “Cor|Cor” columns of the first two rows of each estimand corresponding to PSW results in Table 5, indicate that W.PS excels when the propensity score model is correct and continues to outperform C.PS even when misspecified, as seen in the “Mis|Cor” columns.

Under poor overlap (Table 6), the advantage of W.PS over C.PS becomes even more pronounced. While C.PS suffers from inflated bias and poor coverage, particularly for PATE and PATT, W.PS maintains lower bias, higher efficiency, and closer-to-nominal coverage across all estimands. Particularly, inference for PATO is more robust to poor overlap, PSW with W.PS achieves the lowest biases and highest relative efficiency among all PSW methods, underscoring its ability to attain population-level covariate balance. Moreover, PSW with W.PS for PATO maintains near-nominal coverage at 93.8%, while C.PS drops below 90%. Even when the propensity score model is misspecified and interaction terms are ignored (“Mis|Cor” columns in Tables 5 and 6), W.PS consistently outperforms C.PS in terms of bias and coverage, demonstrating its desirability and robustness under model misspecification, regardless of overlap.

Additional evidence from alternative simulation results (Web Tables 9 and 10), where sampling depends on treatment assignment, confirms W.PS’s advantages. In these extended scenarios, W.PS achieves near-zero bias, while C.PS’s bias deteriorates to 25%–54% under good overlap (Web Table 9). Extreme survey weights, likely when sampling depends on treatment, exacerbate model extrapolation and misspecification issues under C.PS. In contrast, W.PS handles these conditions effectively, maintaining robust performance. Inference

Table 5: Relative bias (%), relative efficiency, and coverage of the 95% confidence intervals for the estimators for continuous outcomes under good overlap across 5000 simulations.

Estimand	Specification					Relative Bias(%)					Relative Efficiency					Coverage						
	W.PS	C.PS	PSW	MOM	CVR	WEI	Cor Cor	Mis Cor	Cor Mis	Mis Mis	Cor Cor	Mis Cor	Cor Mis	Mis Mis	Cor Cor	Mis Cor	Cor Mis	Mis Mis	Cor Cor	Mis Cor	Cor Mis	Mis Mis
PATE	✓	✓	✓				0.554	8.742	-	-	1.000	0.757	-	-	0.951	0.795	-	-	-	-	-	-
			✓				0.741	9.144	-	-	0.729	0.621	-	-	0.926	0.786	-	-	-	-	-	-
	✓	✓		✓			0.188	0.211	0.236	8.704	1.521	1.521	1.365	0.900	0.929	0.929	0.983	0.796	0.930	0.930	0.930	0.974
			✓				0.171	0.194	0.218	8.766	1.520	1.519	1.201	0.893	0.930	0.930	0.974	0.795	0.930	0.930	0.930	0.974
	✓	✓			✓		0.214	0.196	0.152	6.791	1.395	1.395	0.750	0.809	0.940	0.940	0.854	0.859	0.938	0.938	0.938	0.938
PATT	✓	✓					0.195	0.174	8.152	6.677	1.392	1.392	0.739	0.810	0.940	0.941	0.848	0.860	0.935	0.935	0.935	0.804
						✓	0.188	0.196	0.272	8.596	1.525	1.525	1.380	0.919	0.935	0.935	0.986	0.804	0.935	0.935	0.935	0.804
	✓	✓				✓	0.169	0.177	0.251	8.581	1.525	1.524	1.219	0.918	0.935	0.935	0.975	0.805	0.935	0.935	0.975	0.805
			✓				-0.222	5.515	-	-	1.220	0.882	-	-	0.937	0.805	-	-	-	-	-	-
	✓	✓					-0.034	5.792	-	-	0.886	0.725	-	-	0.898	0.773	-	-	-	-	-	-
PATO	✓	✓		✓			-0.035	-1.358	-0.175	5.436	1.425	1.456	1.368	0.954	0.913	0.907	0.935	0.829	0.923	0.923	0.923	0.828
							-0.098	-1.471	-0.226	5.346	1.469	1.491	1.268	0.974	0.911	0.902	0.923	0.828	0.923	0.923	0.923	0.828
	✓	✓			✓		-0.067	-1.390	1.778	6.163	1.649	1.687	1.470	1.114	0.936	0.931	0.931	0.826	0.931	0.931	0.931	0.826
							-0.128	-1.502	1.696	6.062	1.711	1.736	1.478	1.139	0.938	0.939	0.826	0.826	0.939	0.939	0.939	0.826
	✓	✓				✓	-0.034	-1.360	-0.128	5.423	1.427	1.457	1.374	0.959	0.940	0.932	0.863	0.863	0.932	0.932	0.932	0.863
PATO	✓	✓					-0.099	-1.475	-0.181	5.319	1.473	1.494	1.274	0.980	0.938	0.927	0.864	0.864	0.938	0.938	0.938	0.864
			✓				0.328	7.286	-	-	1.633	1.049	-	-	0.951	0.766	-	-	-	-	-	-
	✓	✓					0.481	7.602	-	-	1.042	0.817	-	-	0.914	0.755	-	-	-	-	-	-
				✓			0.286	-0.306	0.299	7.322	1.648	1.664	1.646	1.049	0.925	0.924	0.969	0.783	0.924	0.924	0.924	0.783
	✓	✓					0.252	-0.379	0.251	7.290	1.665	1.677	1.454	1.055	0.924	0.924	0.969	0.783	0.924	0.924	0.924	0.783
PATO	✓	✓					0.204	-0.393	3.974	8.687	1.709	1.725	1.363	1.061	0.938	0.930	0.748	0.748	0.938	0.938	0.938	0.748
							0.168	-0.466	3.900	8.643	1.726	1.738	1.379	1.067	0.937	0.938	0.748	0.748	0.937	0.937	0.937	0.748
	✓	✓				✓	0.288	-0.313	0.318	7.299	1.648	1.663	1.644	1.052	0.935	0.936	0.805	0.805	0.935	0.935	0.936	0.805
						✓	0.252	-0.387	0.275	7.228	1.667	1.677	1.456	1.061	0.936	0.935	0.805	0.805	0.936	0.936	0.936	0.805
	✓	✓					0.252	-0.387	0.275	7.228	1.667	1.677	1.456	1.061	0.936	0.935	0.805	0.805	0.936	0.936	0.936	0.805

Note:

- **W.PS** represents the survey-weighted propensity score model, where survey weights are directly used as regression weights in the propensity score estimation.
- **C.PS** refers to the covariate-adjusted propensity score model, where survey weights are used as an additional covariate in the propensity score estimation.
- **PSW** represents the propensity score weighting estimator without augmentation.
- **MOM** stands for the Moment Estimator.
- **CVR** stands for the Clever Covariate Estimator.
- **WET** represents the Weighted Regression Estimator.
- The four categories under **Relative Bias**, **Relative Efficiency**, and **Coverage** represent combinations of model specification correctness:
  - **Cor|Cor**: Both the propensity score and outcome models are correctly specified.
  - **Mis|Cor**: The outcome model is correctly specified, while the propensity score model is misspecified.
  - **Cor|Mis**: The propensity score model is correctly specified, while the outcome model is misspecified.
  - **Mis|Mis**: Both the propensity score and outcome models are misspecified.
- The - symbol indicates scenarios where the estimator does not involve the outcome model, hence no results are reported.

Table 6: Relative bias (%), relative efficiency, and coverage of the 95% confidence intervals for the estimators for continuous outcomes under poor overlap across 5000 simulations.

Estimand	Specification					Relative Bias(%)					Relative Efficiency					Coverage				
	W.PS	C.PS	PSW	MOM	CVR	WEI	Cor Cor	Mis Cor	Cor Mis	Mis Mis	Cor Cor	Mis Cor	Cor Mis	Mis Mis	Cor Cor	Mis Cor	Cor Mis	Mis Mis		
PATE	✓	✓	✓				8.848	36.838	-	-	1.000	1.073	-	-	0.761	0.505	-	-		
			✓				9.192	37.923	-	-	0.984	1.044	-	-	0.761	0.504	-	-		
	✓	✓		✓			0.727	1.000	2.573	33.376	4.182	4.402	1.463	0.797	0.902	0.904	0.924	0.595		
		✓		✓			0.744	1.016	2.811	33.956	4.211	4.384	1.450	0.777	0.903	0.904	0.923	0.592		
	✓	✓			✓		0.351	0.358	24.972	22.415	23.756	23.663	12.343	13.261	0.800	0.803	0.330	0.365		
PATT	✓	✓					0.340	0.334	24.947	22.298	23.733	23.656	12.257	13.270	0.788	0.790	0.292	0.356		
		✓				✓	1.020	1.012	2.234	29.742	5.760	5.818	3.657	3.223	0.859	0.862	0.882	0.543		
		✓				✓	1.022	1.011	2.308	29.743	5.761	5.806	3.592	3.226	0.858	0.861	0.873	0.542		
	✓	✓	✓				0.774	17.438	-	-	2.299	2.291	-	-	0.793	0.517	-	-		
		✓		✓			1.055	18.092	-	-	2.190	2.173	-	-	0.780	0.504	-	-		
PATO	✓	✓		✓			-0.555	-2.739	0.167	16.538	6.976	7.403	4.661	4.949	0.578	0.559	0.635	0.275		
		✓		✓			-0.567	-2.851	0.218	16.417	7.002	7.423	4.542	4.905	0.580	0.556	0.632	0.282		
	✓	✓			✓		0.022	-2.186	13.213	13.100	19.046	19.401	11.532	11.655	0.938	0.928	0.497	0.602		
		✓		✓			-0.007	-2.320	13.119	13.032	19.097	19.441	11.328	11.513	0.937	0.923	0.523	0.599		
	✓	✓			✓		-0.701	-2.886	1.059	16.278	9.262	9.386	6.606	6.052	0.886	0.874	0.888	0.575		
PATO	✓	✓				✓	-0.721	-3.011	1.084	16.148	9.321	9.442	6.431	6.118	0.885	0.870	0.888	0.576		
		✓					0.492	23.623	-	-	24.336	15.337	-	-	0.938	0.157	-	-		
	✓	✓	✓				0.544	24.283	-	-	15.168	11.695	-	-	0.892	0.191	-	-		
		✓		✓			0.423	-0.454	0.395	23.793	25.027	25.254	24.745	15.280	0.912	0.911	0.973	0.174		
	✓	✓		✓			0.402	-0.565	0.391	23.794	25.105	25.287	21.835	15.342	0.912	0.912	0.962	0.175		
PATO	✓	✓			✓		0.364	-0.510	18.058	21.367	31.746	31.867	20.604	19.124	0.938	0.939	0.347	0.216		
		✓		✓			0.344	-0.621	17.989	21.286	31.862	31.918	20.692	19.221	0.938	0.938	0.346	0.218		
	✓	✓			✓		0.423	-0.474	0.482	23.638	25.049	24.949	24.893	15.469	0.942	0.941	0.983	0.220		
		✓			✓		0.405	-0.585	0.511	23.505	25.162	25.004	22.065	15.619	0.942	0.941	0.976	0.224		
	✓	✓			✓															

**Note:**

- **W.PS** represents the survey-weighted propensity score model, where survey weights are directly used as regression weights in the propensity score estimation.
- **C.PS** refers to the covariate-adjusted propensity score model, where survey weights are used as an additional covariate in the propensity score estimation.
- **PSW** represents the propensity score weighting estimator without augmentation.
- **MOM** stands for the Moment Estimator.
- **CVR** stands for the Clever Covariate Estimator.
- **WET** represents the Weighted Regression Estimator.
- The four categories under **Relative Bias**, **Relative Efficiency**, and **Coverage** represent combinations of model specification correctness:
  - **Cor|Cor**: Both the propensity score and outcome models are correctly specified.
  - **Mis|Cor**: The outcome model is correctly specified, while the propensity score model is misspecified.
  - **Cor|Mis**: The propensity score model is correctly specified, while the outcome model is misspecified.
  - **Mis|Mis**: Both the propensity score and outcome models are misspecified.
- The - symbol indicates scenarios where the estimator does not involve the outcome model, hence no results are reported.



for PATO demonstrates outstanding robustness in the alternative sampling mechanism, especially under poor overlap (Web Table 10). PSW with W.PS for PATO achieves the smallest bias (0.5%) and the highest efficiency—nearly 30 times the reference method (PSW with W.PS for PATE)—and is the only PSW estimator maintaining nominal coverage at 94.4%. These findings align with insights from Ridgeway et al.,[4] which highlight the benefits of survey-weighted propensity scores across various sampling mechanisms, particularly when influenced by treatment assignment, though limited to PATT estimation. Our results extend this by demonstrating W.PS’s robustness across various estimands (PATE, PATT, and PATO) and under different overlap and misspecification conditions.

While initially demonstrated with simple weighting estimators (PSW), these advantages generalize to augmented estimators. In our main scenarios with multi-stage sampling setups (Tables 5 and 6), augmented estimators with W.PS or C.PS perform similarly; this indicates that augmentation provides some protective effect and confers a degree of robustness regardless of how the propensity scores are estimated. However, under treatment-dependent sampling (Web Tables 9 and 10), the differences become substantial, and W.PS wins over C.PS in both bias and coverage even with augmentation. Thus, our simulations highlight W.PS as the preferred method for incorporating survey weights into the propensity score model, regardless of whether an outcome regression component is used in the final estimation.

#### **4.2.2 Is there a universal winner among the augmented estimators when all models are correctly specified?**

Our simulation results, when both the propensity score and outcome models are correctly specified, indicate that all three augmented estimators—moment estimator (MOM), clever covariate estimator (CVR), and weighted regression estimator (WET)—consistently outperform conventional propensity score weighting estimators (PSW). While CVR demonstrate high efficiency under good overlap (Table 5), its sensitivity to different sampling mechanisms and overlap scenarios reduces its practical reliability. MOM offers consistent robustness but provides less bias control and coverage stability compared to WET across all estimands. Consequently, WET with W.PS emerges as the most reliable and efficient estimator, maintaining near-nominal coverage, lower bias, and superior performance, particularly under poor overlap (Table 6) and treatment-dependent sampling (Web Table 9 and 10).

Focusing first on the "Cor|Cor" columns under good overlap (Table 5), W.PS-based MOM, CVR, and WET outperform PSW across all estimands (PATE, PATT, and PATO). The CVR estimator achieves significant efficiency gains, notably for PATT and PATO. Meanwhile, WET consistently performs slightly better than MOM, maintaining around 93% coverage and biases below 2% across most rows.

Under poor overlap ("Cor|Cor" columns in Table 6) PSW alone produces sizable biases (e.g., 8.848% for PATE) and reduced coverage (76.1%). This highlights the well-known vulnerability of simple weighting methods under poor overlap.[36] In contrast, MOM, CVR, and WET demonstrate lower biases and significantly improved efficiency and coverage. For example, CVR with W.PS achieves nearly twenty-fold efficiency gains for PATE, reaching a relative efficiency of 23.8. Similarly, CVR boosts efficiency for PATT and PATO while maintaining better control over coverage. However, comparing across the augmented estimators, CVR and MOM show greater sensitivity under poor overlap. For instance, CVR achieves

only 80% coverage for PATE, and MOM drops to 57.8% coverage for PATT. In comparison, WET consistently maintains coverage near or above 90%, excelling particularly in PATO, where the inherent robustness of PATO protects all methods—both PSW and augmented estimators—enabling them to achieve nominal-level coverage. Among these, WET with W.PS stands out, achieving the highest nominal coverage for PATO at 94.2% and outperforming other estimators.

When sampling probabilities depend on treatment assignment (Web Tables 9 and 10), the advantage of WET becomes even more evident. Under good overlap scenarios (Web Table 9), WET with W.PS achieves minimal biases and relative efficiency exceeding 1.5 for PATE and PATT while maintaining nominal coverage around 94%, demonstrating its resilience under favorable overlap conditions. By contrast, CVR suffers from higher bias and reduced coverage, particularly for PATT, where extreme survey weights correlated with the group variable destabilize its performance. Similar patterns emerge under poor overlap (Web Table 10), where WET maintains near-nominal coverage for PATO. MOM shows better stability than CVR under poor overlap but still falls short of WET’s efficiency.

Taken together, WET with W.PS excels under complex sampling mechanisms. The weighted regression framework effectively minimizes bias and maintains stable coverage, even under extreme overlap conditions. While MOM provides consistent robustness and CVR achieves high efficiency under ideal conditions as in Table 5, both are more vulnerable to instability when survey weights are heterogeneous. Thus, WET with W.PS stands out as the most robust choice, particularly when the sampling mechanism depends on treatment and generates highly variable survey weights.

### **4.2.3 Across all misspecification scenarios, which methods are more robust and should be recommended under the complex survey setting?**

When model specifications deviate from the truth, the performance of estimators varies by estimand and overlap. Examining the “Mis|Cor,” “Cor|Mis,” and “Mis|Mis” columns in Tables 5 and 6 reveals distinct advantages of MOM and WET over CVR, particularly under poor overlap and misspecified outcome models.

When the propensity score model is misspecified but the outcome model is correct under good overlap (“Mis|Cor” columns in Table 5), all three augmented estimators achieve nominal coverage for PATE, demonstrating their doubly robust properties compared to PSW alone, which struggles to achieve even 80% coverage. MOM and WET control bias and coverage better than CVR. In poor overlap scenarios (Table 6), CVR achieves smaller bias but suffers reduced coverage, dropping to only 80%. In contrast, WET with W.PS maintains robustness, achieving 94.1% coverage for PATO, even with misspecified propensity score models.

When the propensity score model is correct but the outcome model is misspecified (“Cor|Mis” columns), CVR demonstrates significant instability. Its bias increases dramatically for PATE under good overlap (Table 5), reaching nearly 10% compared to 0.2% bias for MOM and WET, and its coverage drops below 50% under poor overlap (Table 6). This highlights CVR’s strong reliance on the correct outcome model to manage the extreme values introduced by clever covariates in poor overlap scenarios. While MOM struggles with PATT coverage, WET consistently performs best, achieving the optimal efficiency-coverage balance for PATO, despite the poor overlap and outcome model misspecification.

Under the most challenging scenario, where both the propensity score and outcome models are misspecified (“Mis|Mis” columns), all methods experience significant biases and deteriorated coverage. Despite these challenges, WET remains the most reliable option, consistently maintaining coverage around 80–85% under good overlap, while CVR and MOM frequently see coverage drop below 80%. The extent of bias varies depending on the type of misspecification. In our simulation settings, where interaction terms are omitted from both models, this misspecification strongly affects propensity score estimation and outcome regression, leading to highly biased estimates and poor coverage across all estimators.

Additional simulations (Web Tables 9 and 10), where sampling depends on treatment and extreme survey weights are present, further highlight the relative advantage of WET. WET with W.PS consistently preserves its reliability across misspecified models and overlap conditions, achieving minimal bias (less than 3%) and near-nominal coverage. CVR, by contrast, becomes highly vulnerable under outcome model misspecification, with biases escalating to beyond 70–100% and coverage collapsing to near zero for PATT under poor overlap. MOM remains more stable than CVR but falls short of WET in terms of overall consistency.

In conclusion, among the augmented estimators, while the clever covariate estimator (CVR) demonstrates potential for high efficiency with correctly specified models under good overlap, its heightened sensitivity to model misspecification and poor overlap limits its practical reliability. In contrast, both the moment estimator (MOM) and weighted regression estimator (WET) exhibit greater robustness, with WET emerging as the most resilient choice across varying survey settings, maintaining low bias and near-nominal coverage for a wide range of estimands, even under challenging conditions. Thus, WET with W.PS is strongly recommended for reliable performance in real-world applications.

## 5 Case studies

### 5.1 Application to the ECLS-K dataset

In this first data example, we present a real-world example with multi-stage sampling to demonstrate the application of the different methods for using propensity score weighting with complex survey data. The data used are from the The Early Childhood Longitudinal Study, Kindergarten class 1998-1999 (ECLS-K), [37] a nationally representative longitudinal study that tracks early school experiences of a large cohort from kindergarten through eighth grade in the United States. The study collects comprehensive information at the child, household, and school levels, with survey weights reflecting the multi-stage probability sampling design employed to ensure representativeness. [16] The participants were selected prior to the assignment of treatment. Specifically, children were sampled at the start of the study upon their entry into kindergarten, and subsequent treatment, such as the receipt of special education services, was determined in later stages, typically based on administrative records collected during follow-up data collection waves, such as in first grade or fifth grade. This example reflects a prospective study design, consistent with our main simulation setting, in which the sampling process does not depend on treatment assignment

and survey weights are explicitly known. This design simplifies the integration of survey weights into subsequent estimations by reducing the steps required. Corresponding estimators tailored for prospective studies are further detailed in the appendix. The treatment status was defined as participation in special education services, as documented in school administrative records. The outcome of interest was the IRT-scaled math achievement test score from the ECLS-K dataset. A set of 39 baseline covariates covering demographic, household, and school characteristics were included to address potential confounding, as described in Keller and Tipton.[38]

Web Figure 1 illustrates the population-level covariate balance, measured by Population-level survey-weighted Standardized Mean Differences (PSMD), which align with the weights  $(\omega_{1,p}(\mathbf{X}_i), \omega_{0,p}(\mathbf{X}_i))$  used in the effect estimation and correspond to the specifications outlined in Table 1 for each target population and estimand. The PSMD for a baseline covariate  $X$  is calculated as the absolute difference between the weighted means of  $X$  for the treated and control groups, scaled by the pooled weighted standard deviation  $s_p$ . Specifically, PSMD is defined as  $j\bar{X}_1 - \bar{X}_0j/s_p$ , where the weighted means are  $\bar{X}_1 = (\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i)X_iZ_i) / (\sum_{i=1}^n \omega_{1,p}(\mathbf{X}_i)Z_i)$  for the treated group and  $\bar{X}_0 = (\sum_{i=1}^n \omega_{0,p}(\mathbf{X}_i)X_i(1 - Z_i)) / (\sum_{i=1}^n \omega_{0,p}(\mathbf{X}_i)(1 - Z_i))$  for the control group. The pooled weighted standard deviation,  $s_p$ , reflects a composite measure of the variability of  $X$  within the treated and control groups, weighted according to the propensity score weights  $\omega_{1,p}$  and  $\omega_{0,p}$ , and accounts for the sample size in each group. The figure compares the covariate balance for each population-level estimand—PATE, PATT, and PATO—using survey-weighted regression (W.PS). The visual evidence clearly demonstrates that W.PS achieves superior covariate balance, as indicated by a larger proportion of covariates achieving PSMDs below the 0.10 threshold, a criterion for adequate balance.[39] This is further detailed in the grey-shaded cells of Web Table 17, which shows how W.PS systematically improves covariate balance across most baseline characteristics. Specifically, W.PS achieves exact balance for PATO, further supporting its appropriateness for estimands where balancing the overlap population is critical.

To estimate the causal effect of elementary school special education services on fifth-grade math achievement, we applied the Propensity Score Weighting (PSW) estimator alongside three augmented estimators: the Moment Estimator (MOM), Clever Covariate Regression (CVR), and Weighted Regression Estimator (WET). Building on theoretical results and simulation studies that consistently identify survey-weighted propensity scores (W.PS) as the most robust and reliable approach across diverse conditions, this analysis emphasizes the application of W.PS. The estimated treatment effects for PATE, PATT, and PATO using W.PS are presented in Table 7, underscoring its practical advantages in achieving reliable causal inference.

The estimated treatment effect of special education services on fifth-grade math achievement using the WET under W.PS for PATO was -5.80, with a 95% confidence interval of (-7.34, -4.26). This significant negative effect indicates that, on average, students in special education scored approximately 5.8 points lower in math compared to the general population with similar baseline characteristics. In contrast, the PSW estimator produced larger standard errors, such as  $SE = 2.80$  for PATT, and confidence intervals that included zero, highlighting its lower precision and reduced ability to confirm significant effects. The augmented estimators (MOM, CVR, and WET) consistently yielded confidence intervals that excluded zero across all estimands, demonstrating sufficient statistical power for detecting

the underlying causal effect. Among them, WET under W.PS consistently yielded more precise and robust estimates, whereas MOM and CVR exhibited greater variability, particularly under potential model misspecification in real-world analyses. Notably, the PATO estimand remained stable across all estimators, emphasizing the reliability of the overlap population for deriving comparable treatment effects. These results align with theoretical guarantees and simulation findings, further supporting the superior performance of WET when combined with W.PS. The evidence underscores WET’s practical utility and robustness in addressing complex survey designs and achieving reliable causal estimates.

Table 7: Estimated treatment effects and standard errors (SE) for PATE, PATT, and PATO using different estimators under W.PS in the ECLS-K study.

PSW	MOM	CVR	WET	PATE (SE)	PATT (SE)	PATO (SE)
✓				-3.31(2.64)	-1.12(2.80)	-5.81(1.61)
	✓			-4.47(1.54)	-3.68(1.90)	-5.96(1.61)
		✓		-3.91(1.47)	-4.38(1.28)	-4.65(1.15)
			✓	-4.86(1.43)	-4.58(1.64)	-5.80(1.53)

## 5.2 Application to the 2009 Medical Expenditure Panel Survey dataset

In our second empirical example, we demonstrate how the proposed methods perform in a retrospective survey design where the sampling mechanism depends directly on the exposure—here, race. We use data from the 2009 Medical Expenditure Panel Survey (MEPS),[?] which historically oversamples minority groups, including Hispanics and Blacks, to improve the precision of health disparity analyses. Starting in 2006, this strategy expanded to include Asian households in response to changing demographics and analytical priorities.[40] This oversampling design ensures adequate representation of minority groups but can also lead to disproportionate survey weights, particularly for subgroups with limited population overlap. Unlike Data Example 1 (ECLS-K), where participants were sampled before treatment status was realized, MEPS employs a retrospective design where individuals are sampled after their race is determined, leading to survey weights that often correlate with race and demographic strata. This underscores the importance of incorporating these survey weights into propensity score and effect estimation.

The 2009 MEPS public-release file includes 9830 non-Hispanic White adults (hereafter “Whites”), 4020 Blacks, 1446 Asians, and 5150 Hispanics, each assigned a survey weight reflecting their probability of selection and nonresponse adjustments.[41, 40] Here, we focus on two pairwise comparisons (White–Asian and White–Hispanic) to estimate racial disparities in total annual healthcare expenditures, measured in 2009 USD. Although “race” is not a manipulable treatment, the methodological goal parallels standard propensity score approaches

to balancing covariates between two groups. Following the framework for unconfounded descriptive comparisons described in Li and Li,[42] we control for a set of variables reflecting clinical need, as instructed by the Institute of Medicine (IOM),[43, 44] including body mass index (BMI), SF-12 physical and mental component summaries, health conditions, age, gender, and marital status. Additionally, we account for selected socioeconomic and functional limitation variables, ensuring alignment with established approaches to health care disparity analyses. These variable selections align with prior applications of MEPS data in disparity research,[41, 36, 45, 42] while recognizing that some SES variables, such as education and region, are not fully incorporated due to limitations in the available dataset.

Web Figures 2, 3, and 4 collectively illustrate the interplay between survey weights, propensity score distributions estimated by the survey-weighted propensity score model (W.PS), and covariate balance under different weighting methods for White–Asian and White–Hispanic comparisons. In the White–Asian comparison, MEPS’s oversampling strategy leads to highly concentrated survey weights for Asians and a broader range with extreme weights for Whites, amplifying disparities in baseline covariates like BMI. These imbalances are reflected in the propensity score distributions, estimated using the survey-weighted propensity score model (W.PS), which show limited overlap between the groups, with most Whites concentrated at high scores (above 0.9) and Asians clustering near 0.8–0.9. The love plots further demonstrate that this limited overlap results in poor covariate balance under PATE and PATT, particularly under C.PS, where many covariates exceed the 0.1 PSMD threshold. Conversely, overlap weighting (PATO), especially under W.PS, achieves exact balance by focusing on comparable subpopulations, mitigating instability caused by extreme weights. In the White–Hispanic comparison, stronger overlap and more balanced survey weights yield better covariate alignment across methods, with PATO consistently outperforming other estimands in achieving robust and interpretable results. Overall, these findings underscore the importance of selecting appropriate weighting methods. Poor balance under PATE and PATT, especially in White–Asian comparisons, will inflate future variance estimation and bias effect estimates due to skewed target populations and extreme weights. Overlap weighting for PATO, particularly under W.PS, emerges as the most effective strategy for achieving population-level covariate balance and robust inference by focusing on comparable subpopulations and minimizing instability from extreme scores.[36, 45, 40, 42]

Table 8 summarizes the estimated differences in annual healthcare expenditures (in 2009 USD) between Whites and Asians and between Whites and Hispanics for PATE, PATT, and PATO under W.PS. In the White–Asian comparison, limited overlap, particularly among high-BMI Asians, resulted in a PATE estimate using PSW of \$2263.23 (SE = 575.60), while the PATO estimate, focusing on comparable overlap subpopulations, had a notably smaller SE of 236.56 with a point estimate of \$1352.88, highlighting the stability of overlap-based estimands. Augmented estimators, especially WET, maintained consistent performance and resilience to extreme weights, producing a PATO estimate of \$1358.16 (SE = 237.55). Similarly, for the White–Hispanic comparison, the PATE estimate under PSW was \$607.78 (SE = 390.54), and the PATO estimate achieved a smaller SE of 164.71 with a point estimate of \$1317.00. Augmented estimators further reduced variability, with WET yielding a PATO estimate of \$1308.27 (SE = 162.38), reflecting robust performance across different racial comparisons under both favorable and challenging overlap conditions. Overall, these results underscore the effectiveness of W.PS in addressing survey weight imbalances across minor-

Table 8: Estimated treatment effects and standard errors (SE) for PATE, PATT, and PATO using different estimators under W.PS in the White–Asian and White–Hispanic comparisons of annual healthcare expenditures (2009 USD), in the MEPS study.

Comparison Group	PSW	MOM	CVR	WET	PATE (SE)	PATT (SE)	PATO (SE)
White–Asian	✓				2263.228 (575.601)	2327.342 (600.877)	1352.879 (236.564)
		✓			3962.495 (1913.583)	4053.586 (1958.159)	1349.055 (236.329)
			✓		1237.268 (614.993)	1231.726 (652.689)	1497.991 (283.749)
				✓	1179.610 (424.043)	1152.960 (442.362)	1358.156 (235.356)
White–Hispanic	✓				607.782 (390.542)	463.373 (447.725)	1317.004 (164.707)
		✓			1230.326 (311.024)	1206.963 (352.923)	1308.903 (161.845)
			✓		1157.681 (317.028)	1172.387 (365.968)	1288.049 (164.289)
				✓	1242.545 (241.151)	1227.124 (264.413)	1308.268 (161.411)

ity groups and demonstrate the robustness of WET in achieving precise treatment effect estimation, emphasizing their utility for analyzing racial disparities in MEPS. [36, 45]

## 6 Discussion

Our paper introduces a unified framework for incorporating survey weights into the balancing weights paradigm, enhancing their applicability to survey observational data and providing robust tools for population-level causal inference. We establish the asymptotic normality of survey-weighted estimators for retrospective designs, with extensions to prospective cases detailed in the appendix. A robust sandwich variance estimator for weighting methods is developed via M-estimation theory, which extends to a unified variance framework for all three augmented estimators— one-step moment estimator (MOM), clever covariate regression estimator (CVR), and weighted regression estimator (WET)—derived from the shared augmented weighting estimating equations for survey observational data. Simulation studies evaluate these methods under varying overlap and model misspecification conditions. Results demonstrate WET’s robustness, MOM’s consistency, and CVR’s efficiency. Practical applicability is illustrated through two data examples: estimating the effect of special education services on fifth-grade math achievement using the Early Childhood Longitudinal Study Kindergarten Class (ECLS-K) dataset under a prospective multi-stage sampling survey design, and analyzing racial disparities in healthcare expenditures using the Medical Expenditure Panel Survey (MEPS) under a retrospective descriptive comparison framework. These case studies highlight the strengths of survey-weighted propensity score models (W.PS) in achieving population-level covariate balance, WET in providing robust estimation, and the PATO estimand in enabling reliable inference within comparable subpopulations. All proposed methods are implemented in the publicly available PSweight R package.

First, incorporating survey weights throughout all phases of the propensity score weighting estimator is essential for unbiased treatment effects estimation in survey observational studies. Using a survey-weighted regression model (W.PS) for propensity score estimation

aligns sampled data covariate distributions with the target population, enhancing efficiency, coverage, and robustness across varying overlap scenarios. Consistent with Ridgeway et al.,[4] our results suggest that omitting survey weights restricts balance to within the sample,[2, 46] potentially introducing bias when generalizing to target populations. Consequently, W.PS is integral to valid population-level causal inferences.

Second, augmented estimators are instrumental in reducing bias, enhancing efficiency, and improving coverage in survey observational data, particularly when both the propensity score and outcome models are correctly specified. Simulation results highlight the strengths of MOM, CVR and WET in different scenarios. WET demonstrates superior robustness under model misspecification and consistently performs well across estimands and overlap conditions. CVR shows high efficiency under good overlap and correct models but exhibits sensitivity to outcome model misspecification and diminished reliability in poor overlap scenarios, limiting its utility in real-world applications. These findings align with prior literature [30, 26] that highlights the instability of inverse propensity-based covariate approaches, particularly under extreme values. Extrapolation issues are exacerbated when the clever covariate incorporates large survey weights, amplifying variability and undermining model performance. Reformulating the clever covariate to preserve boundedness properties[26] could enhance its robustness and practical utility, particularly in scenarios with extreme survey weights or limited overlap.

Third, WET consistently emerges as the most robust and efficient estimator across diverse overlap conditions and model misspecifications. Compared to CVR and MOM, WET demonstrates superior stability and precision, particularly in poor overlap or when models are misspecified. This aligns with Kang and Schafer[30] and Robins et al.[26], who suggest that clever covariates can supplement outcome models but fail to address misspecifications under poor overlap, leading to coverage declines as low as 30%, as shown in Table 6. Moreover, as noted by Robins et al.,[26] when the CVR is structured with an identity link, it aligns with Targeted Maximum Likelihood Estimation (TMLE) for marginal means but can perform worse than the weighted regression estimator (WET) when both models are misspecified. Simulation results and case studies further confirm WET’s robustness, showing lower bias and more reliable coverage in both prospective and retrospective survey designs. Case study analyses demonstrate WET’s capacity to mitigate the influence of extreme survey weights, particularly for PATO, delivering stable and interpretable results. Overall, WET under W.PS is recommended for complex survey settings, combining efficiency, robustness, and practical reliability.

The validity of our study’s findings hinges on critical assumptions, including the stable unit treatment value assumption (SUTVA) and treatment assignment ignorability. [47, 48] These assumptions are crucial for unbiased causal effect estimation under observational designs but may be violated in real-world settings due to issues such as latent confounders or non-overlapping covariate distributions.[49, 50] Hence, sensitivity analyses could provide critical insights into the potential impact of unmeasured confounding for weighting and augmented weighting estimator, and need to be generalized to complex survey data.[51] Second, our analysis assumes that survey weights are known and derived from well-designed surveys. If this assumption does not hold—for instance, in cases where the survey design introduces biases or weights are not appropriately generated—causal inferences may be distorted. Poor survey design, including unaddressed non-response biases or inadequate post-stratification,



can result in flawed survey weights that exacerbate selection bias or produce extreme values, destabilizing estimators and reducing efficiency.[50, 19] Future research could extend sensitivity analyses to explore the robustness of survey weights under alternative assumptions about the sampling mechanism.[49, 52] Third, while our findings focus on binary treatments, they are also readily extensible to multiple treatments. With multiple nominal or ordinal treatments, the generalized propensity scores (GPS) provide a key device for simultaneously balancing covariates across multiple treatment arms, [53, 45] and are key ingredients in developing weighting and augmented weighting estimators. Future work is needed to extend the current methods to multiple treatments in the context of survey data. Finally, our analysis used parametric models for propensity score and outcome estimation, as these are the most commonly used methods in real practice. However, these parametric models may fail to capture complex, high-dimensional, or nonlinear structures in real-world data [54]. More flexible approaches, such as Bayesian Additive Regression Trees (BART) [55, 56] and Causal Forests [57] can effectively capture non-linear relationships and heterogeneous treatment effects. These flexible modeling frameworks are particularly advantageous in high-dimensional settings or when traditional parametric models prove inadequate.[58] Advanced estimators like the Targeted Maximum Likelihood Estimation (TMLE) can achieve semi-parametric efficiency by incorporating machine learning methods to optimize both outcome and propensity score estimates.[59] Extensions of AIPW methods to double machine learning (DML) frameworks can also further enhance robustness by addressing model misspecification and overfitting challenges.[54] These data-adaptive extensions of the current estimators in the context of survey observational data represent fruitful directions for additional research.

## References

- [1] LaVange L, Koch G, Koch G, Schwartz T. Applying sample survey methods to clinical trials data. *Statistics in Medicine* 2001; 20. doi: 10.1002/sim.732
- [2] Zanutto EL. A comparison of propensity score and linear regression analysis of complex survey data. *Journal of Data Science* 2006; 4: 67–91.
- [3] DuGoff EH, Schuler M, Stuart EA. Generalizing observational study results: applying propensity score methods to complex surveys. *Health Services Research* 2014; 49(1): 284–303.
- [4] Ridgeway G, Kovalchik SA, Griffin BA, Kabeto MU. Propensity score analysis with survey weighted data. *Journal of Causal Inference* 2015; 3(2): 237–249.
- [5] Imai K, King G, Stuart E. Misunderstandings between experimentalists and observationalists about causal inference. *Journal of the Royal Statistical Society: Series A* 2008; 171(2): 481–502. doi: 10.1111/j.1467-985X.2007.00527.x
- [6] Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika* 1983; 70(1): 41–55.

- [7] Lunceford JK, Davidian M. Stratification and weighting via the propensity score in estimation of causal treatment effects: a comparative study. *Statistics in Medicine* 2004; 23(19): 2937–2960.
- [8] Stuart E. Matching methods for causal inference: A review and a look forward. *Statistical Science* 2010; 25(1): 1–21. doi: 10.1214/09-STS313
- [9] Li F, Morgan KL, Zaslavsky AM. Balancing Covariates via Propensity Score Weighting. *Journal of the American Statistical Association* 2018; 113(521): 390–400. doi: 10.1080/01621459.2016.1260466
- [10] Zanutto E, Lu B, Hornik R. Using propensity score subclassification for multiple treatment doses to evaluate a national antidrug media campaign. *Journal of Educational and Behavioral Statistics* 2005; 30(1): 59–73. doi: 10.3102/10769986030001059
- [11] Austin P, Jembere N, Chiu M. Propensity score matching and complex surveys. *Statistical Methods in Medical Research* 2018; 27(4): 1240–1257. doi: 10.1177/0962280216658920
- [12] Dong N, Stuart E, Lenis D, Quynh Nguyen T. Using propensity score analysis of survey data to estimate population average treatment effects: a case study comparing different methods. *Evaluation Review* 2020; 44(1): 84–108. doi: 10.1177/0193841X20938497
- [13] Lenis D, Nguyen T, Dong N, Stuart E. It’s all about balance: propensity score matching in the context of complex survey data. *Biostatistics* 2019; 20(1): 147–163. doi: 10.1093/biostatistics/kxx063
- [14] Salerno S, Roberts EK, Needham BL, McCormick TH, Mukherjee B, Shi X. What’s the Weight? Estimating Controlled Outcome Differences in Complex Surveys for Health Disparities Research. <https://arxiv.org/abs/2406.19597>; 2024.
- [15] Yang C, Cuerden M, Zhang Wea. Propensity score weighting with survey weighted data when outcomes are binary: a simulation study. *Health Serv Outcomes Res Method* 2023. doi: 10.1007/s10742-023-00317-y
- [16] Education Statistics fNC. Early childhood longitudinal study [United States]: Kindergarten Class of 1998–1999 Kindergarten–Eighth Grade Full Sample. Inter-university Consortium for Political and Social Research [distributor]; 2011.
- [17] Imbens GW. Nonparametric Estimation of Average Treatment Effects Under Exogeneity: A Review. *The Review of Economics and Statistics* 2004; 86(1): 4–29. doi: 10.1162/003465304323023651
- [18] Hirano K, Imbens GW, Ridder G. Efficient estimation of average treatment effects using the estimated propensity score. *Econometrica* 2003; 71(4): 1161–1189.
- [19] Li F, Morgan KL, Zaslavsky AM. Balancing covariates via propensity score weighting. *Journal of the American Statistical Association* 2018; 113(521): 390–400.

- [20] Cheng C, Li F, Thomas LE, Li FF. Addressing Extreme Propensity Scores in Estimating Counterfactual Survival Functions via the Overlap Weights. *American Journal of Epidemiology* 2022; 191(6): 1140-1151. doi: 10.1093/aje/kwac043
- [21] Zeng S, Li F, Hu L, Li F. Propensity Score Weighting Analysis of Survival Outcomes Using Pseudo-observations. *Statistica Sinica* 2023; 33(3): 2161–2184. doi: 10.5705/ss.202021.0175
- [22] Crump RK, Hotz VJ, Imbens GW, Mitnik OA. Dealing with Limited Overlap in Estimation of Average Treatment Effects. *Biometrika* 2009; 96(1): 187–199. doi: 10.1093/biomet/asn055
- [23] Ackerman B, Lesko C, Siddique J, Susukida R, Stuart E. Generalizing randomized trial findings to a target population using complex survey population data. *Statistics in Medicine* 2021; 40(5): 1101–1120. Epub 2020 Nov 26doi: 10.1002/sim.8822
- [24] Stefanski LA, Boos DD. The calculus of M-estimation. *The American Statistician* 2002; 56(1): 29–38.
- [25] Bang H, Robins JM. Doubly robust estimation in missing data and causal inference models. *Biometrics* 2005; 61(4): 962–973. doi: 10.1111/j.1541-0420.2005.00377.x
- [26] Robins JM, Sued M, Lei-Gomez Q, Rotnitzky A. Comment: Performance of double-robust estimators when "inverse probability" weights are highly variable. *Statistical Science* 2007; 22(4): 544–559.
- [27] Mao H, Li L, Greene T. Propensity score weighting analysis and treatment effect discovery. *Statistical methods in medical research* 2019; 28(8): 2439–2454.
- [28] Tsiatis AA. *Semiparametric Theory and Missing Data*. Springer Series in Statistics New York, NY: Springer . 2006
- [29] Scharfstein DO, Rotnitzky A, Robins JM. Adjusting for nonignorable drop-out using semiparametric nonresponse models. *Journal of the American Statistical Association* 1999; 94(448): 1096–1120.
- [30] Kang JDY, Schafer JL. Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data. *Statistical Science* 2007; 22(4): 523–539.
- [31] Schafer JL, Kang J. Average causal effects from nonrandomized studies: a practical guide and simulated example. *Psychological Methods* 2008; 13(4): 279–313. doi: 10.1037/a0014268
- [32] Vansteelandt S, Keiding N. Invited Commentary: G-Computation—Lost in Translation?. *American Journal of Epidemiology* 2011; 173(7): 739–742. doi: 10.1093/aje/kwq474
- [33] Gabriel EE, Sachs MC, Martinussen T, et al. Inverse probability of treatment weighting with generalized linear outcome models for doubly robust estimation. *Statistics in Medicine* 2024; 43(3): 534–547. doi: <https://doi.org/10.1002/sim.9969>

- [34] Shook-Sa BE, Hudgens MG, Knittel AK, et al. Exposure effects on count outcomes with observational data, with application to incarcerated women. *The Annals of Applied Statistics* 2024; 18(3): 2147–2165.
- [35] Hu L, Ji J, Li F. Estimating heterogeneous survival treatment effect in observational data using machine learning. *Statistics in Medicine* 2021; 40(21): 4691–4713. doi: 10.1002/sim.9090
- [36] Li F, Thomas L, Li F. Addressing extreme propensity scores via the overlap weights. *American Journal of Epidemiology* 2019; 188(1): 250–257. doi: 10.1093/aje/kwy201
- [37] Tourangeau K, Nord C, Lê T, Sorongon AG, Najarian M. Early Childhood Longitudinal Study Kindergarten Class of 1998–99 (ECLS-K): Combined User’s Manual for the ECLS-K Eighth-Grade and K-8 Full Sample Data Files and Electronic CodeBooks. NCES Report 2009-004, National Center for Education Statistics; National Center for Education Statistics, Washington, DC: 2009.
- [38] Keller B, Tipton E. Propensity score analysis in R: a software review. *Journal of Educational and Behavioral Statistics* 2016; 41(2): 326–348.
- [39] Austin PC, Stuart EA. Moving towards best practice when using inverse probability of treatment weighting (IPTW) using the propensity score to estimate causal treatment effects in observational studies. *Statistics in Medicine* 2015; 34(28): 3661–3679. Epub 2015 Aug 3doi: 10.1002/sim.6607
- [40] Chowdhury SR, Machlin SR, Gwet KL. Sample Designs of the Medical Expenditure Panel Survey Household Component, 1996–2006 and 2007–2016. Methodology Report 33, Agency for Healthcare Research and Quality; Rockville, MD: 2019.
- [41] Cook BL, McGuire TG, Lock K, Zaslavsky AM. Comparing Methods of Racial and Ethnic Disparities Measurement Across Different Settings of Mental Health Care. *Health Services Research* 2010; 45(3): 825–847. doi: 10.1111/j.1475-6773.2010.01100.x
- [42] Li F, Li F. Using propensity scores for racial disparities analysis. *Observational Studies* 2023; 9(1): 59–68. doi: 10.1353/obs.2023.0005
- [43] McGuire TG, Alegria M, Cook BL, Wells KB, Zaslavsky AM. Implementing the institute of medicine definition of disparities: An application to mental health care. *Health Services Research* 2006; 41: 1979–2005. doi: 10.1111/j.1475-6773.2006.00583.x
- [44] Cook BL, McGuire TG, Meara E, Zaslavsky AM. Adjusting for health status in non-linear models of health care disparities. *Health Services and Outcomes Research Methodology* 2009; 9: 1–21. doi: 10.1007/s10742-009-0040-2
- [45] Li F, Li F. Propensity Score Weighting for Causal Inference with Multiple Treatments. *The Annals of Applied Statistics* 2019; 13(4): 2389–2415. doi: 10.1214/19-AOAS1282

- [46] Dugoff E, Schuler M, Stuart E. Generalizing observational study results: applying propensity score methods to complex surveys. *Health Services Research* 2014; 49(1): 284–303. doi: 10.1111/1475-6773.12090
- [47] Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika* 1983; 70(1): 41–55.
- [48] Imbens GW, Rubin DB. *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press . 2015.
- [49] Rosenbaum PR, Rubin DB. Assessing sensitivity to an unobserved binary covariate in an observational study with binary outcome. *Journal of the Royal Statistical Society: Series B (Methodological)* 1983; 45(2): 212–218.
- [50] Cole SR, Stuart EA. Generalizing evidence from randomized clinical trials to target populations: The ACTG 320 trial. *American Journal of Epidemiology* 2010; 172(1): 107–115.
- [51] Rosenbaum PR. Sensitivity analysis for certain permutation inferences in matched observational studies. *Biometrika* 1987; 74(1): 13–26. doi: 10.1093/biomet/74.1.13
- [52] Hartman E, Huang M. Sensitivity Analysis for Survey Weights. *Political Analysis* 2024; 32: 1–16.
- [53] Yang S, Imbens GW, Cui Z, Faries DE, Kadziola Z. Propensity score weighting for causal inference with multiple treatments. *Statistics in Medicine* 2020; 39(11): 1718–1736.
- [54] Chernozhukov V, Chetverikov D, Demirer M, et al. Double/debiased machine learning for treatment and structural parameters: inference with machine learning. *The Econometrics Journal* 2018; 21(1): C1–C68.
- [55] Hill JL. Bayesian nonparametric modeling for causal inference. *Journal of Computational and Graphical Statistics* 2011; 20(1): 217–240.
- [56] Chen X, Harhay MO, Tong G, Li F. A Bayesian machine learning approach for estimating heterogeneous survivor causal effects: applications to a critical care trial. *The annals of applied statistics* 2024; 18(1): 350.
- [57] Wager S, Athey S. Estimation and Inference of Heterogeneous Treatment Effects using Random Forests. *Journal of the American Statistical Association* 2018; 113(523): 1228–1242.
- [58] Athey S, Wager S. Estimation and inference of heterogeneous treatment effects using random forests. *Journal of the American Statistical Association* 2018; 113(523): 1228–1242.
- [59] Laan v. dMJ, Rubin D. Targeted maximum likelihood learning. *The International Journal of Biostatistics* 2006; 2(1).