

Symmetries of 3-webs around a point

Jean Paul Dufour.

January 30, 2025

Abstract

Let W be a real analytic planar 3-web defined on a neighborhood of a point M . In this paper we consider local diffeomorphisms of the plane which preserve M and which map any foliation of W onto a (not necessarily the same) foliation. We call them "symmetry" of W around M . There is three types of such symmetries: the first one are those which preserve each foliation; the second one are those which preserve only one foliation and permute the two others; the third are those which permute circularly the three foliations.

We know that hexagonal (i.e. flat) 3-webs have always the three types of symmetries around each point. In this paper we study the non-flat case.

We give a classification of 3-webs which admit a symmetry of first or second type. The problem is more difficult for the third case but we give a method for building all 3-webs with this type of symmetry. We also give a precise example which is non-flat.

Keywords: planar 3-webs.

AMS classification : 53A60

1 Introduction

In this text we work in the real analytic framework. The results could be extended to the smooth and to the complex frameworks.

A planar 3-web is a triple of 1-dimensional foliations, mutually transverse, on an open domain of the plane. In the sequel we will work only with these 3-webs: so we forget the word *planar*. Let W be such a web and M a point of its domain.

A **symmetry of W around M** is a local diffeomorphism of the plane which preserves M and maps any foliation of W to another (possibly different) foliation.

We have to consider three different types of such symmetries: **simple symmetries** are those which preserve each foliation; **mirror symmetries** are those which preserve only one foliation and permute the two others; **circular symmetries** are those which give a circular permutation of the three foliations.

We choose local coordinates (x, y) near M , vanishing at this point, such that the foliations are given by the verticals lines $x = \text{constant}$, the horizontal lines $y = \text{constant}$ and the level sets of some function f (the sets of (x, y) such that $f(x, y) = \text{constant}$). Such a 3-web is denoted (x, y, f) .

We know that W is **flat** near M iff we can find such coordinates with $f(x, y) = x + y$. In that case it is easy to see that it has circular, mirror or non-trivial simple symmetries.

2 Normal forms and simple symmetries.

A classical result ([DJ]), based on a result of S. Sternberg ([SS]), claims that we can always choose coordinates (x, y) vanishing at M such that

$$f(x, y) = x + y + xy(x - y)g(x, y) \tag{1}$$

where g is an analytic (or smooth) function.

We say that $(x, y, x + y + xy(x - y)g(x, y))$ is a **normal form** of W near M . They are characterised by the three equations

$$f(t, 0) = t, \quad f(0, t) = t, \quad f(t, t) = 2t,$$

for any small t , and W is **non-flat** ([AH]) near M iff the function $g(x, y)$ does not vanish identically near the origin. Moreover these coordinates are unique up to a homothety $(x, y) \mapsto (\lambda x, \lambda y)$. At least this notion proves easily the following result.

Proposition 2.1 *If W admits a simple symmetry ϕ around M with has Id (identity) as tangent map at M then ϕ is the identity.*

We denote by $\sum g_{rs}x^r y^s$ the Taylor expansion of the fonction g near the origin, the above homothety maps $(x, y, f(x, y))$ to a the web $(x, y, x + y +$

$xy(x - y)G(x, y)$ with

$$G(x, y) = \sum g_{rs} \lambda^{r+s+2} x^r y^s.$$

Suppose there is at least one non-vanishing coefficient $g_{r_0 s_0}$ with $r_0 + s_0$ odd. When we choose correctly λ , we obtain that the new coefficient of $x^{r_0} y^{s_0}$ is 1. With this new constraint, the normal form is unique and the only simple symmetry is the identity (Id).

Suppose now that all the coefficients g_{rs} with $r + s$ odd vanish. Then the homothety $(x, y) \mapsto (-x, -y)$ ($-Id$) doesn't change our normal form. So we can conclude with the following

Theorem 2.2 *If the 3-web W is non-flat, either it has only Id as simple symmetry around M , or it has two simple symmetries Id and ϕ , where ϕ is a non trivial involution ($\phi^2 = I$ and the linear part of ϕ at the origin is $-Id$).*

Moreover it has exactly two simple symmetries if and only if its normal form $(x, y, x + y + xy(x - y)g(x, y))$ is such that $g(x, y) = g(-x, -y)$ for every x and y .

As far I know, this result never appeared in the classical web litterature.

Remark: The notion of normal form works also in the complex case. We have an analog of the preceeding proposition but we may have more than the above non-trivial simple symmetries: We suppose that n is the lowest integer such that they are non-zero g_{rs} with $n = r + s$. For a convenient choice of λ the diffeomorphism $\lambda.Id$ gives a new normal form but with $g_{r_0 s_0} = 1$ for some couple (r_0, s_0) with $r_0 + s_0 = n$. The only diffeomorphisms ϕ which preserve this constraint are of the form $\mu.Id$ where μ is a $(n + 2)$ -root of the unit ($\mu^{n+2} = 1$). As ϕ preserves only the monomial $x^r y^s$ with $r + s + 2 = k.(n + 2)$ for any integer k , $\mu.Id$ will give $n + 1$ non-trivial simple symmetries when G contains only these monomials in its Taylor expansion.

3 Mirror symmetries of 3-webs.

Suppose that W has a mirror symmetry ϕ around M .

We first remark that ϕ^2 is a simple symmetry; the theorem above implies that we have two possibilities : either $\phi^2 = Id$ or $\phi^4 = Id$ and the linear part of ϕ^2 at the origin is $-Id$.

We can choose local coordinates (x, y) , which give the normal form for W , and such that ϕ permutes the foliations $x = constant$ and $y = constant$.

Thus ϕ has the form $(x, y) \mapsto (X(y), Y(x))$ for two diffeomorphisms X and Y of the line (fixing the origin). The remark above proves that there are two cases: the linear part of ϕ is $(x, y) \mapsto (y, x)$ or $(x, y) \mapsto (-y, -x)$.

Since ϕ is a mirror symmetry, it must preserve the third foliation. As this last is given by the the level lines of a fonction $f(x, y) = x + y + xy(x - y)g(x, y)$, this may be written as

$$f((X(y), Y(x))) = Z(f(x, y)),$$

for some third diffeomorphism Z of the line.

When we use the relation $f(t, 0) = f(0, t) = t$ we obtain $X = Y = Z$.

We have also the obvious property $f(t, t) = 2t$. So when we write $x = y = t$ in the above formula we obtain

$$X(2t) = 2X(t),$$

which implies that X , Y and Z are the same homothety: Id or $-Id$. So we can rewrite our equation as

$$f(y, x) = f(x, y)$$

or

$$f(-y, -x) = -f(x, y)$$

This can be written as

$$g(y, x) = -g(x, y).$$

or

$$g(-y, -x) = -g(x, y).$$

Theorem 3.1 *If the 3-web W is non flat, it has zero, one or two mirror symmetries around M .*

It has a mirror symmetry ϕ which is an involution ($\phi^2 = Id$) iff it has a normal form $(x, y, x + y + xy(x - y)g(x, y))$, with a function g which satisfies $g(y, x) = -g(x, y)$. It has a mirror symmetry ϕ which $\phi^4 = Id$ iff it has a normal form with a function g which satisfies $g(y, x) = -g(-x, -y)$. It may have simultaneously the two types of mirror symmetries.

4 Circular symmetries.

Suppose that W has a circular symmetry ϕ around M . We choose local coordinates (x, y) which provide to W the form $(x, y, f(x, y))$, with

$$f(x, y) = x + y + \text{hot},$$

("hot" means higher order terms) and such that ϕ maps the foliation by vertical lines to the foliation by horizontal lines, the foliation by horizontal lines to the foliation by the level sets of f and the foliation by the level sets of f to the foliation by the vertical lines.

Remark that ϕ^3 is a simple symmetry of W . So, if W is non-flat, we have $\phi^3 = Id$ or $\phi^6 = Id$.

In the first case the linear part $\phi^{(1)}$ of ϕ at the origin has the form $(x, y) \mapsto (-x - y, x)$ and $(x, y) \mapsto (x + y, -x)$ in the second case.

In the sequel we study only the first case $\phi^3 = Id$; the second can be treated exactly as the first one.

In that case ϕ gives a local analytic action of $\mathbb{Z}/3\mathbb{Z}$ wich fixes the origin. A classical result claims that such local action is conjugate to its linear part. This means that there is a local diffeomorphism ψ (with identity as linear part at the origin) which fixes the origin and such that

$$\phi = \psi^{-1} \circ \phi^{(1)} \circ \psi$$

on a neighborhood of the origin. With notations $\psi(x, y) = (A(x, y), B(x, y))$ and $\psi^{-1}(x, y) = (U(x, y), V(x, y))$ and $\phi(x, y) = (F(x, y), G(x, y))$ we have the relations

$$U(A(x, y), B(x, y)) - x = 0, \tag{2}$$

$$V(A(x, y), B(x, y)) - y = 0, \tag{3}$$

$$F(x, y) = U(-A(x, y) - B(x, y), A(x, y)), \tag{4}$$

$$G(x, y) = V(-A(x, y) - B(x, y), A(x, y)). \tag{5}$$

The linear parts of A and U at the origin is x and the linear parts of B and V at the origin is y .

Now we remark that ϕ maps vertical lines to horizontal lines if and only if $G(x, y)$ is independant of y . So we have the following equation

$$V(-A(x, y) - B(x, y), A(x, y)) - \mu(x) = 0 \quad (6)$$

for some function $\mu = x + hot$.

When the functions μ and $V(x, y) = y + hot$ are given, the theorem of implicit functions claims that the pair of equations (6) and (3) have a suitable local solution (A, B) . Then the equation (2) has a solution U and we obtain F by (4).

It is obvious that ϕ maps level sets of the function F to vertical lines. Because $\phi^3 = Id$, it is also obvious that ϕ is a circular symmetry of the 3-web (x, y, F) . As we have $F = -x - y + hot$ we obtain $f = -F$.

This gives a way to construct all the non-flat webs which admit a circular symmetry. However this method, starting with different (μ, V) , could give isomorphic 3-webs; it can also give flat webs. In the sequel we will give a V such that, with $\mu = Id$, we obtain a non-flat 3-web in normal form.

Lemma 4.1 *We assume that $P(x, y)$ is a function invariant by $\phi^{(1)}$, i.e. such that $P(-x - y, x) = P(x, y)$, with a vanishing 1-jet at the origin. We assume also that $\theta(x, y)$ is the local solution of the equation*

$$\theta + P(x + \theta, y + \theta) = 0.$$

For $V = y + P$ and $\mu = Id$, we obtain $f(x, y) = -x - y - 3\theta$.

Proof: We write $A = x + \alpha(x, y)$, $B = y + \beta(x, y)$ and $U = x + \gamma(x, y)$ where the 1-jets at the origin of α , β and γ vanish. Equations (6) and (3) give $B + P(A, B) = y$ and $A + P(A, B) = x$; so we get $\beta = \alpha$ and $\alpha + P(x + \alpha, y + \alpha) = 0$. So we have $\alpha = \theta$.

Now equation (2) gives $\gamma = P$ and $F = -A - B + P(A, B)$ leads to $f(x, y) = -x - y - 3\theta$, which achieves our proof.

Consider the polynomial $P_0 = xy(x - y)(x + y)(2x + y)(x + 2y)$. It is invariant by $\phi^{(1)}$.

Lemma 4.2 *The equation $\theta + P_0(x + \theta, y + \theta) = 0$ have a solution of the shape $\theta = P_0(-1 + hot)$.*

To prove this lemma we impose $\theta = P_0(-1 + \eta)$ and we apply the theorem of implicit functions to our equation (considered as an equation of the shape $K(x, y, \eta) = 0$).

Now we find that F has the shape $-x - y - 3P_0(1 + \eta)$. So our web has the normal form (x, y, f) with $f = -F$ and it is non-flat.

Remark that the level set $f = 0$ is the line $y = -x$. The level set $f = t$ contains the points $(-t, 2t)$, $(0, t)$, $(t/2, t/2)$, $(t, 0)$ and $(2t, -t)$; they are all on the line $x + y = t$ but, for $t \neq 0$, this level set is different of this line.

So we have proved the following theorem.

Theorem 4.3 *There are non-flat 3-webs which have a circular symmetry around some point.*

Remarks.

Using [DJ], we could obtain analogous results for local $(p + 1)$ -webs of codimension n on a $p.n$ dimensional space.

We fall on this results trying to construct a counter-example to the celebrate "Gronwall's conjecture for planar 3-webs" first appearing in [THG] (see [SA] for the better history of this conjecture and the last results). To be more precise, we tried to construct a **linear** non-flat 3-web, with a symmetry around some point, which is not a homography. In fact, after very long calculations via Maple, it seems that there is no example of such 3-web.

References

- [SA] S.I. Agafonov, *Gronwall's conjecture for 3-webs with two pencils of lines*. Differential Geometry and its Applications, vol.91 (dec. 2023).
- [DJ] J. P. Dufour, P. Jean *Rigidity of Webs and Families of Hypersurfaces*. in Singularities and Dynamical Systems, (Iraklion, 1983), North Holland, Amsterdam, (1985), 271-283.
- [THG] T. H. Gronwall, *Sur les équations entre trois variables représentables par les nomogrammes à points alignés*. Journal de Liouville, 8, (1912),59-102.
- [AH] A. Hénaut, *Analytic web geometry*, 6-48, in Web theory and related topics. (J. Grifone and E. Salem editors) World Scientific Publishing Co. Ptc. Ltd 2001.
- [SS] S. Sternberg, *On the structure of local homeomorphisms of euclidean n space*. Ann. of Math. 80, (1958), 623-631.