More on the asymptotic behaviour of moments of branching Markov processes

Christopher B. C. Dean^{*} Emma Horton^{*}

Abstract

Consider a branching Markov process, $X = (X_t, t \ge 0)$, with non-local branching mechanism. Studying the asymptotic behaviour of the moments of X has recently received attention in the literature [6, 7] due to the importance of these results in understanding the underlying genealogical structure of X. In this article, we generalise the results of [7] to allow for a nonsimple leading eigenvalue and to also study the higher order fluctuations of the moments of X. These results will be useful for proving central limit theorems and extending well-known LLN results.

Keywords : Branching process, fluctuations, moments, non-local branching.

MSC: 60J80, 60F05.

1 Introduction

Let E be a Polish space and $\dagger \notin E$ a cemetery (absorbing) state. We consider a measure-valued stochastic process $X = (X_t, t \ge 0)$ given by

$$X_t := \sum_{i=1}^{N_t} \delta_{x_i(t)}, \qquad t \ge 0,$$

whose atoms $\{x_i(t) : i = 1, ..., N_t\}$ evolve in $E \cup \{\dagger\}$ according to the following dynamics. From an initial position $x \in E$, particles evolve independently in E according to a Markov process (ξ, \mathbf{P}_x) . When at $y \in E$, particles branch at rate $\gamma(y)$, which we assume to be bounded, at which point the particle is replaced by a random number, N, of offspring at positions x_1, \ldots, x_N in E. The law of the offspring locations and their number is denoted by \mathcal{P}_y , where y denotes the position of the parent particle at the branch time. We will also often use the notation \mathcal{E}_y for the associated expectation operator. We refer to X as a $(\mathbf{P}, \gamma, \mathcal{P})$ -branching Markov process (BMP). We will let \mathbb{P}_{μ} denote its law when initiated from $\mu \in \mathcal{M}(E)$, the space of finite measures on E. We also let B(E) denote the space of bounded complex-valued functions, $B^+(E)$ denote the space of non-negative real bounded functions and $B_1(E)$, (resp. $B_1^+(E)$) denote those functions in B(E), (resp. $B^+(E)$) that are bounded by unity.

^{*}Department of Statistics, University of Warwick, Coventry, CV4 7AL, UK. Email: {Christopher.B.C.Dean}, {Emma.Horton}@warwick.ac.uk

One way to characterise the behaviour of X is via its linear expectation semigroup,

$$\psi_t[f](x) := \mathbb{E}_{\delta_x} \left[X_t[f] \right], \quad t \ge 0, \, x \in E, \, f \in B(E), \tag{1.1}$$

where \mathbb{E}_{δ_x} denotes the expectation conditional on X_0 consisting of a single particle at x, and, for a measure μ and a function f, we have used the notation

$$\mu[f] := \int_E f(x)\mu(\mathrm{d}x)$$

In many cases, it is well known that the semigroup ψ satisfies the following Perron Frobenius-type decomposition.

(PF) There exists $\lambda_1 \in \mathbb{R}, \varphi_1 \in B^+(E)$ and a probability measure $\tilde{\varphi}_1$ on E such that

$$\psi_t[\varphi_1] = e^{\lambda_1 t} \varphi_1, \qquad \tilde{\varphi}_1[\psi_t[f]] = e^{\lambda_1 t} \tilde{\varphi}_1[f],$$

and

$$\sup_{x \in E, f \in B_1(E)} \left| e^{-\lambda_1 t} \psi_t[f](x) - \tilde{\varphi}_1[f] \varphi_1(x) \right| \to 0, \quad \text{as } t \to \infty.$$

When this decomposition holds, along with moment conditions on the offspring distribution, it has been shown in [7] that similar asymptotics hold for the moments $\mathbb{E}_{\delta_x}[X_t[f]^k]$. We also mention the previous works [3, 5, 11, 12], where asymptotics for moments of branching processes were studied under varying assumptions. Results of this kind are particularly useful in proving laws of large numbers [1, 4, 9], for setting benchmarks for Monte Carlo codes [2] and for understanding the genealogical structure of branching Markov processes [6, 8, 10]. However, these results do not give a complete description of the asymptotic behaviour of X. Indeed, they do not allow us to say anything about the fluctuations of the process via a CLT, for example. Moreover, there are many cases where (PF) is not satisfied, for example when the first eigenvalue is not simple. The aim of this article is to study the asymptotic moments of X in more depth by considering what happens when we have a more general version of (PF). These results will play a key role in proving CLTs for X and will also allow us to provide new proofs for a LLN result.

1.1 Assumptions

Before stating our main results, we first spend some time discussing the assumptions. Our main assumption generalises (PF) to the case when λ_1 is not a simple eigenvalue and where we may have more points in the discrete spectrum of ψ . Here and throughout, we will use the notation $\ker(\lambda_{\ell}) := \{f \in B(E) : \tilde{\varphi}_{i,j}^{(k)}[f] = 0, 1 \le i \le \ell, 1 \le j \le p_i, 1 \le k \le k_{i,j}\}$ for each $1 \le \ell \le m$, where the $\tilde{\varphi}_{i,j}^{(k)}$ are defined in the following assumption.

(H1) There exists an integer $m \ge 1$, a real number $\lambda_1 > 0, \lambda_2, \ldots, \lambda_m \in \mathbb{C}$, that satisfy $\lambda_1 > \operatorname{Re}\lambda_2 \ge \cdots \ge \operatorname{Re}\lambda_m$, such that, for $1 \le i \le m$, there exist bounded functions

$$\varphi_{i,1}^{(1)}, \dots, \varphi_{i,1}^{(k_{i,1})}, \dots, \varphi_{i,p_i}^{(1)}, \dots, \varphi_{i,p_i}^{(k_{i,p_i})} \in B(E)$$

bounded linear functionals

 $\tilde{\varphi}_{i,1}^{(1)},\ldots,\tilde{\varphi}_{i,1}^{(k_{i,1})},\ldots,\tilde{\varphi}_{i,p_i}^{(1)},\ldots,\tilde{\varphi}_{i,p_i}^{(k_{i,p_i})}:B(E)\to\mathbb{C},$

¹If m = 1, the list $\lambda_2 \dots, \lambda_m$ is an empty list.

such that $\tilde{\varphi}_{i,j}^{(k)}[\varphi_{i,j}^{(k)}] = 1$, $\tilde{\varphi}_{i,j}^{(k)}[\varphi_{\ell,n}^{(q)}] = 0$ for each $(i, j, k) \neq (\ell, n, q)$. There also exists and a linear operator \mathcal{N} , such that

- for each $1 \le i \le m, \ 1 \le k \le k_{i,1}, \ \mathcal{N}\varphi_{i,1}^{(k)} = 0;$
- for each $1 \leq i \leq m, 2 \leq j \leq p_i, 1 \leq k \leq k_{i,j}$ there exists² $k^* \in \{1, \ldots, k_{i,j-1}\}$ such that $\mathcal{N}\varphi_{i,j}^{(k)} = \varphi_{i,j-1}^{(k^*)};$
- for $f \in \ker(\lambda_m), \mathcal{N}f = 0$,

x

and, for any $x \in E$, $t \ge 0$ and $f \in B(E)$,

$$\psi_t[\varphi_{i,j}^{(k)}](x) = \mathrm{e}^{(\lambda_i + \mathcal{N})t}\varphi_{i,j}^{(k)}(x), \quad \text{and} \quad \tilde{\varphi}_{i,j}^{(k)}[\psi_t[f]] = \mathrm{e}^{\lambda_i t}\tilde{\varphi}_{i,j}^{(k)}[\mathrm{e}^{\mathcal{N}t}f]. \tag{1.2}$$

Moreover,

$$\sup_{e \in E, f \in B_1(E)} e^{-\operatorname{Re}\lambda_m t} \left| \psi_t[f](x) - \sum_{j=1}^{p_i} e^{(\lambda_i + \mathcal{N})t} \Phi_i[f](x) \right| \to 0, \quad \text{as } t \to \infty, \tag{1.3}$$

where

$$\Phi_{i,j}[f](x) = \sum_{k=1}^{k_{i,j}} \tilde{\varphi}_{i,j}^{(k)}[f]\varphi_{i,j}^{(k)}(x), \quad \Phi_i[f](x) = \sum_{j=1}^{p_i} \Phi_{i,j}[f](x).$$

Let us spend some time unpacking the above assumption. First note that, as mentioned above, this allows for the case that the leading eigenvalue associated with ψ is not simple. For simplicity, assume that m = 1 in (H1). In the case that $p_1 = k_{1,1} = 1$, we recover (PF). In the case that $p_1 = 1$ but $k_{1,1} > 1$, we extend (PF) to the case where λ_1 is not simple. This case implies that $k_{1,1}$ is both the algebraic and geometric multiplicity of λ_1 , with corresponding eigenfunctions given by $\varphi^{(1)}, \ldots, \varphi^{(k_{1,1})}$ (note, we have dropped the subscript for simplicity). The fact that these are eigenfunctions can be deduced from (1.2) and the first property of \mathcal{N} . The case that $p_1 = 2$, say, the functions $\varphi_{1,1}^{(1)}, \ldots, \varphi_{1,1}^{(k_{1,1})}$ are eigenfunctions, but the functions $\varphi_{1,2}^{(1)}, \ldots, \varphi_{1,2}^{(k_{1,2})}$ are generalised eigenfunctions of rank 2, meaning $\mathcal{N}\varphi_{1,2}^{(k)} \neq 0$ but $\mathcal{N}^2\varphi_{1,2}^{(k)} = 0$, which can be deduced from the first two properties of \mathcal{N} . In general, if $p_1 \geq 3$, then $\varphi_{1,j}^{(k)}$ is of rank j. Note that, in the case that m = 1 and $p_1 \geq 2$, the geometric multiplicity of λ_1 is $k_{1,1}$ and the algebraic multiplicity is $\sum_{j=1}^{p_1} k_{1,j}$. Allowing $m \geq 2$ generalises this to m eigenvalues $e^{\lambda_1 t}, \ldots, e^{\lambda_m t}$, each with eigenfunctions $\varphi_{i,1}^{(1)}, \ldots, \varphi_{i,1}^{(k_{i,1})}$ and generalised eigenfunctions $\varphi_{i,j}^{(1)}, \ldots, \varphi_{i,j}^{(k_{i,j})}$ of rank j. The asymptotic given in (1.3) stipulates that, when normalised by the smallest growth rate $e^{\operatorname{Re}\lambda_m t}$, the expectation semigroup converges.

1.2 Notation

We are interested in seeing how (H1) propagates to higher moments of X_t . We will shortly see that it will obey one of three possible dynamics depending on the growth rate of $\psi_t[f]$. However, we first introduce some notation that will be used throughout the rest of the article.

For $k \ge 1$, $A \subseteq [k] := \{1, \ldots, k\}$, and $\boldsymbol{f} \in B(E)^k$, we write

$$\psi_t^{(A)}[\boldsymbol{f}](x) = \mathbb{E}_{\delta_x}\left[\prod_{i \in A} X_t[f_i]\right], \quad t \ge 0.$$

²If $k_1 \neq k_2$ then $k^*(i, j, k_1) \neq k^*(i, j, k_2)$

If A = [k], we will write $\psi_t^{(k)}$ instead.

Similarly to [7], we will use an inductive argument to prove our results concerning the asymptotic behaviour of $\psi_t^{(k)}$. For this, we introduce the following notation.

For a set $A \subset \mathbb{N}$, we let $\mathcal{P}(A)$ denote the set of partitions of A and set $\mathcal{P}^*(A) := \mathcal{P}(A) \setminus \{A\}$. Further, if $|A| \ge 2$ and $2 \le \ell \le |A|$, we set

$$B_{\ell,A} := \{ \boldsymbol{i} = (i_1, \dots, i_\ell) \in A^\ell : i_j \neq i_k, j \neq k \}.$$

For $\sigma \in \mathcal{P}(A)$, we write $|\sigma|$ for the number of blocks in σ and σ_j for the *j*-th block. For $k \geq 1$, we write $\mathcal{P}(k)$, $\mathcal{P}^*(k)$ and $B_{\ell,k}$ in place of $\mathcal{P}([k])$, $\mathcal{P}^*([k])$ and $B_{\ell,[k]}$, respectively. Then, for $A \subset \mathbb{N}$ and a function $\boldsymbol{g} : \mathfrak{P}(A) \times E \to \mathbb{C}$, where $\mathfrak{P}(A)$ denotes the power set of A, we define

$$\zeta_A[\boldsymbol{g}](x) = \mathcal{E}_x \left[\sum_{\sigma \in \mathcal{P}^*(A)} \sum_{\boldsymbol{i} \in B_{|\sigma|,N}} \prod_{j=1}^{|\sigma|} \boldsymbol{g}(\sigma_j, x_{i_j}) \right].$$
(1.4)

The above operator will allow us to write $\psi_t^{(k)}$ in terms of the lower order moments, thus motivating inductive proofs. For similar reasons, we would like to develop asymptotics akin to that of (H1) but for the other eigenvalues. To this end, recall the notation from (H1) and define

$$\operatorname{Ei}^{*}(\lambda_{1}) = \{ f \in B(E) : \Phi_{1,j}[f] \neq 0 \text{ for some } 1 \leq j \leq p_{1} \},$$

$$\operatorname{Ei}^{*}(\lambda_{i}) = \{ f \in B(E) : \Phi_{i,j}[f] \neq 0 \text{ for some } 1 \leq j \leq p_{i} \}, \quad 2 \leq i \leq m,$$

and

$$\operatorname{Ei}(\lambda_1) = \operatorname{Ei}^*(\lambda_1), \quad \operatorname{Ei}(\lambda_i) = \operatorname{Ei}^*(\lambda_i) \setminus \bigcup_{\substack{\operatorname{Re}\lambda_j \ge \operatorname{Re}\lambda_i \\ \lambda_j \neq \lambda_i}} \operatorname{Ei}^*(\lambda_j), \quad 2 \le i \le m.$$

We will also use the notation $\text{Ei}_1(\lambda_i)$ to denote $\text{Ei}(\lambda_i)$ restricted to functions bounded by unity. Then, thanks to (1.3), for $1 \leq i \leq m$, we have

$$\sup_{x \in E, f \in \operatorname{Ei}_1(\lambda_i)} e^{-\operatorname{Re}\lambda_i t} \left| \psi_t[f](x) - e^{(\lambda_i + \mathcal{N})t} \Phi_i[f](x) \right| \to 0, \quad \text{as } t \to \infty.$$
(1.5)

For $1 \leq i \leq m$ and $f \in \text{Ei}_1(\lambda_i)$, if $2\text{Re}\lambda_i \geq \lambda_1$, then we will show that the asymptotic growth rate of the higher moments of $X_t[f]$ depends on the growth rate of the semigroup $\psi_t[f]$ given in (1.5). In this setting, we say that λ_i is *large* when $2\text{Re}\lambda_i > \lambda_1$ and *critical* when $2\text{Re}\lambda_i = \lambda_1$. If however, $2\text{Re}\lambda_i < \lambda_1$, then the asymptotic growth rate of the higher moments of $X_t[f]$ depends instead on λ_1 and p_1 . We say that λ_i is *small* when $2\text{Re}\lambda_i < \lambda_1$. For this reason, we will work with a different function space for each of these three regimes, which we now define.

When it exists, let τ be the unique index such that $2\text{Re}\lambda_{\tau-1} \geq \lambda_1$ but $2\text{Re}\lambda_{\tau} < \lambda_1$. Set

$$\operatorname{Ei}(\Lambda_L) := \bigcup_{2\operatorname{Re}\lambda_i > \lambda_1} \operatorname{Ei}(\lambda_i), \quad \operatorname{Ei}(\Lambda_C) := \bigcup_{2\operatorname{Re}\lambda_i = \lambda_1} \operatorname{Ei}(\lambda_i), \quad \operatorname{Ei}(\Lambda_S) := \ker(\lambda_{\tau-1}),$$

where ker (λ_{ℓ}) was defined just before (H1), and where if τ is not well defined (i.e. there are no small eigenvalues), we set $\text{Ei}(\Lambda_S) = \emptyset$.

For $f \in \text{Ei}(\Lambda_L) \cup \text{Ei}(\Lambda_C)$, let $\nu(f)$ be the unique index such that $f \in \text{Ei}(\lambda_{\nu(f)})$, set $\lambda(f) := \lambda_{\nu(f)}$ and define

$$p(f) := \max\{i \ge 0 : \mathcal{N}^{i-1}f \neq 0\}.$$

Then, for $f \in (\text{Ei}(\Lambda_L) \cup \text{Ei}(\Lambda_C))^k$, define

$$\tilde{\lambda}(\boldsymbol{f}) = \sum_{i=1}^{k} \lambda(f), \quad \tilde{p}(\boldsymbol{f}) = \sum_{i=1}^{k} p(f).$$

Finally, for $\ell \geq 1$ with ℓ even and $\mathbf{f} \in \text{Ei}(\Lambda_C)^{\ell}$, we say that $(\lambda(f_1), \ldots, \lambda(f_\ell))$ is a *conjugate* ℓ -tuple if one can split $(f_i)_{i=1}^{\ell}$ into $\ell/2$ pairs such that, for each pair $(f_i, f_j), \lambda(f_i) = \overline{\lambda}(f_j)$.

2 Main results

In this section, we will present our main results. We are interested in the asymptotic behaviour of $\psi_t^{(k)}$. As briefly mentioned in the previous section, this behaviour depends on whether we are in the large, critical or small regime. For this reason, we will only consider the asymptotics of $\psi_t^{(k)}[\mathbf{f}]$ for $\mathbf{f} \in \text{Ei}(\Lambda_L)^k \cup \text{Ei}(\Lambda_C)^k \cup \text{Ei}(\Lambda_S)^k$. In the case that we have $\mathbf{f} \in B_1(E)^k$, the same proof techniques can be used, however this would result in significantly longer theorems and proofs, since many more cases need to be considered.

From here on we assume that (H1) is in force.

Theorem 2.1 (Large Regime). Assume that for some $k \geq 2$,

$$\sup_{x\in E} \mathcal{E}_x[N^k] < \infty$$

For $1 \leq \ell \leq k$ and $t \geq 0$ set

$$\Delta_{\ell,t} = \sup_{x \in E, \boldsymbol{f} \in \mathrm{Ei}_1(\Lambda_L)^{\ell}} \left| \mathrm{e}^{-\tilde{\lambda}(\boldsymbol{f})t} (1+t)^{\tilde{p}(\boldsymbol{f})-\ell} \psi_t^{(\ell)}[\boldsymbol{f}](x) - \prod_{j=1}^{\ell} (p(f_j)-1)! L_{[\ell]}[\boldsymbol{f}^*](x) \right|,$$

where, for $1 \leq i \leq \ell$, $L_{\{i\}}[\boldsymbol{f}](x) = \Phi_{\nu(f_i)}[f_i](x)$, and for $A \subseteq [\ell]$ with $2 \leq |A| \leq \ell$,

$$L_A[\boldsymbol{f}](x) = \int_0^\infty e^{-\tilde{\lambda}(\boldsymbol{f})s} \psi_s \left[\gamma \zeta_A[L_{\cdot}[(e^{-\mathcal{N}s}f_1, \dots, e^{-\mathcal{N}s}f_{\ell})]] \right](x) \mathrm{d}s, \qquad (2.1)$$

and where

$$\boldsymbol{f}^* = (\mathcal{N}^{p(f_1)-1} f_1, \dots, \mathcal{N}^{p(f_\ell)-1} f_\ell).$$
(2.2)

Then, for $1 \leq \ell \leq k$,

$$\sup_{t \ge 0} \Delta_{\ell,t} < \infty \ and \ \lim_{t \to \infty} \Delta_{\ell,t} = 0$$

Theorem 2.2 (Small Regime). Assume that for some $k \ge 2$,

$$\sup_{x\in E}\mathcal{E}_x[N^k]<\infty$$

For $t \ge 0$ and $2 \le \ell \le k$, let

$$\Delta_{\ell,t} = \sup_{x \in E, \boldsymbol{f} \in \mathrm{Ei}_1(\Lambda_S)^{\ell}} \left| \mathrm{e}^{-\frac{\ell \lambda_1}{2}} (1+t)^{-\frac{\ell(p_1-1)}{2}} \psi_t^{(\ell)}[\boldsymbol{f}](x) - L_{[\ell]}[\boldsymbol{f}](x) \right|,$$
(2.3)

where λ_1 and p_1 were given in (H1), and, for $A \subseteq \{1, \ldots, \ell\}$ with |A| even, if |A| = 2, we have

 $L_{A}[\boldsymbol{f}](x) = (p_{1}-1)!\Phi_{1,1}[\mathcal{N}^{p_{1}-1}(f_{A_{1}}f_{A_{2}})](x) + (p_{1}-1)!\int_{0}^{\infty} e^{-\lambda_{1}s}\Phi_{1,1}[\mathcal{N}^{p_{1}-1}(\gamma\zeta_{A}[\psi_{s}^{(\cdot)}[\boldsymbol{f}]])](x)ds,$

and for $|A| \ge 4$, we have

$$L_A[\boldsymbol{f}](x) = \int_0^\infty e^{-\frac{\lambda_1|A|}{2}} \psi_s \left[\gamma \zeta_A[L_{\cdot}[\boldsymbol{f}]]\right](x) ds.$$

Otherwise, $L_A[\mathbf{f}](x) = 0$. Then, for $1 \le \ell \le k$,

$$\sup_{t\geq 0} \Delta_{\ell,t} < \infty \ and \ \lim_{t\to\infty} \Delta_{\ell,t} = 0.$$

Remark 2.3. Note that under a different scaling one can use the techniques presented here to obtain convergence of the odd moments in the small regime. This scaling depends on whether $\text{Re}\lambda_{\tau}$ is greater than, equal to, or less than 0, and we see a similar trichotomy as in this article. We omit these results, since it would make the presentation much more technical whilst providing little gain.

Theorem 2.4 (Critical Regime). Assume that for some $k \ge 2$,

$$\sup_{x\in E}\mathcal{E}_x[N^k] < \infty$$

For $2 \leq \ell \leq k$, let

$$\Delta_{\ell,t} = \sup_{x \in E, \boldsymbol{f} \in \mathrm{Ei}_1(\Lambda_C)^{\ell}} \left| \mathrm{e}^{-\frac{\ell \lambda_1}{2}} (1+t)^{-\left(p(\boldsymbol{f}) + \frac{\ell(p_1-2)}{2}\right)} \psi_t^{(\ell)}[\boldsymbol{f}](x) - L_{[\ell]}[\boldsymbol{f}](x) \right|, \qquad (2.4)$$

where λ_1 and p_1 were given in (H1), and, for $A \subseteq \{1, \ldots, \ell\}$ with |A| even, we have

$$L_{A}[\boldsymbol{f}](x) = \frac{\Phi_{1,1}[\mathcal{N}^{p_{1}-1}(\gamma\zeta_{[2]}[\Phi_{\lambda(\cdot),1}[\boldsymbol{f}^{*}]])](x)(\tilde{p}(\boldsymbol{f})-2)!}{(p(f_{1})-1)!(p(f_{2})-1)!(p_{1}+\tilde{p}(\boldsymbol{f})-2)!}, \quad |A|=2, \quad (f_{i}:i\in A) \text{ conjugate 2-tuple},$$

where f^* is as in (2.2), and for $|A| \ge 4$,

$$L_A[\boldsymbol{f}](x) = \int_0^\infty e^{-\frac{\lambda_1|A|}{2}} \psi_s \left[\gamma \zeta_A[L_{\boldsymbol{\cdot}}[\boldsymbol{f}]] \right](x) ds, \quad (f_i : i \in A) \ conjugate \ |A| - tuple.$$

Otherwise $L_A[\mathbf{f}] = 0$. Then, for $1 \le \ell \le k$,

$$\sup_{t\geq 0} \Delta_{\ell,t} < \infty \ and \ \lim_{t\to\infty} \Delta_{\ell,t} = 0.$$

Similarly to Remark 2.3, we can use the techniques presented here to obtain convergence of the odd moments under a different scaling. In this setting, the conjugate tuple property extends by asking for the final eigenvalue to be real and equal to $\lambda_1/2$.

3 Proofs

This section is dedicated to the proofs of the main results. The proofs will follow similar ideas to those presented in [7]. The main idea is to use an inductive argument using Lemma 4.1, with the base case given by (H1). Before starting the proofs, we introduce some notation and several useful inequalities which will be used throughout. We use C for a generic non-negative constant that may change between equations. For multiple generic constants, we use the notation C_1, C_2, \ldots . Firstly, for $\mathbf{f} \in (\text{Ei}(\Lambda_L) \cup \text{Ei}(\Lambda_C))^k$, and $A \subseteq [k]$, we extend the definitions of $\tilde{p}(\mathbf{f})$ and $\tilde{\lambda}(\mathbf{f})$ as follows:

$$\tilde{p}(\boldsymbol{f}, A) = \sum_{i \in A} p(f_i),$$
$$\tilde{\lambda}(\boldsymbol{f}, A) = \sum_{i \in A} \lambda(f_i).$$

Next, since \mathcal{N} is non-zero on a finite dimensional space, it is bounded and therefore there exists a constant C such that, for any $f \in B(E)$,

$$\|e^{\mathcal{N}t}f\|_{\infty} \le C(1+t)^{p(f)-1}\|f\|_{\infty}, \quad t \in \mathbb{R}.$$
 (3.1)

This and (H1) imply the existence of a further constant C such that, for $1 \le i \le m$ and $f \in \text{Ei}(\lambda_i)$,

$$\|\psi_t[f]\|_{\infty} \le C e^{\operatorname{Re}\lambda_i t} (1+t)^{p(f)-1} \|f\|_{\infty},$$
(3.2)

where in particular we have used (1.5). Finally, for any $k \ge 2$, we have that

$$\zeta_{[k]}[1](x) = \mathcal{E}_x \left[\sum_{\sigma \in \mathcal{P}^*(A)} \sum_{i \in B_{|\sigma|,N}} 1 \right] \le \mathcal{E}_x[N^k].$$
(3.3)

3.1 Proof of Theorem 2.1

We first prove an intermediate result that is of a similar form to Theorem 2.1.

Lemma 3.1. Assume that for some $k \geq 2$,

$$\sup_{x\in E}\mathcal{E}_x[N^k]<\infty.$$

For $1 \leq \ell \leq k$ and $t \geq 0$ set

$$\tilde{\Delta}_{\ell,t} = \sup_{x \in E, \boldsymbol{f} \in \mathrm{Ei}_1(\Lambda_L)^{\ell}} \left| \mathrm{e}^{-\tilde{\lambda}(\boldsymbol{f})t} \psi_t^{(\ell)}[(\mathrm{e}^{-\mathcal{N}t} f_1, \dots, \mathrm{e}^{-\mathcal{N}t} f_\ell)](x) - L_{[\ell]}[\boldsymbol{f}](x) \right|,$$

where the $L_{[\ell]}$ were defined in Theorem 2.1. Then, for $1 \leq \ell \leq k$,

$$\sup_{t \ge 0} \Delta_{\ell,t} < \infty \text{ and } \lim_{t \to \infty} \Delta_{\ell,t} = 0.$$

Proof. Let $k \ge 2$ and assume that Theorem 2.1 holds for $\ell \le k - 1$, where the case of k = 1 holds by (H1). Furthermore, for ease of notation, for a function $\mathbf{f} \in B(E)^k$, let

$$\boldsymbol{f}_t = (f_{t,1}, \dots, f_{t,k}), \quad f_{t,i} = e^{-\mathcal{N}t} f_i, \quad 1 \le i \le k, \quad t \in \mathbb{R}.$$
(3.4)

Firstly, by Lemma 4.1, for any $\boldsymbol{f} \in B(E)^k$, $x \in E$, and $t \ge 0$,

$$\psi_t^{(k)}[\boldsymbol{f}](x) = \psi_t \left[\prod_{i=1}^k f_i\right](x) + \int_0^t \psi_s \left[\gamma \zeta_{[k]}[\psi_{t-s}^{(\cdot)}[\boldsymbol{f}]]\right](x) \mathrm{d}s.$$
(3.5)

Since the product of any function uniformly bounded by 1 is also uniformly bounded by 1, by (3.1) and (3.2), the first term on the right-hand side satisfies

$$\lim_{t \to \infty} \sup_{x \in E, \mathbf{f} \in \mathrm{Ei}_1(\Lambda_L)^k} \mathrm{e}^{-\mathrm{Re}\tilde{\lambda}(f)t} \left| \psi_t \left[\prod_{i=1}^k f_{t,i} \right] (x) \right| = 0, \tag{3.6}$$

where we have used, for any $\mathbf{f} \in \text{Ei}(\Lambda_L)^k$, $\text{Re}\tilde{\lambda}(\mathbf{f}) > \lambda_1$. For the integral term in (3.5), first note that by the inductive hypothesis and (3.1), we have

$$\sup_{0 \le s \le t < \infty} \sup_{\boldsymbol{f} \in \operatorname{Ei}(\Lambda_L)^k} \sup_{A \subseteq [k]} (1+s)^{-(\tilde{p}(f,A)-k)} \| \mathrm{e}^{-\tilde{\lambda}(f,A)(t-s)} \psi_{t-s}^{(\cdot)}[\boldsymbol{f}_t](A,\cdot) - L_A[\boldsymbol{f}_s] \|_{\infty} < \infty,$$

where we have applied the inductive hypothesis to the functions $((1+s)^{-(p(f_i)-1)}f_{s,i})_{i\in A}$. Furthermore, again by the inductive hypothesis, the left-hand side tends to 0 as $t - s \to \infty$. This and (3.3) imply

$$\sup_{0 \le s \le t < \infty} \sup_{\boldsymbol{f} \in \text{Ei}(\Lambda_L)^k} (1+s)^{-(\tilde{p}(f,A)-k)} \| e^{-\tilde{\lambda}(f)(t-s)} \zeta_{[k]}[\psi_{t-s}^{(\cdot)}[\boldsymbol{f}_t]](x) - \zeta_{[k]}[L_{\cdot}[\boldsymbol{f}_{-s}]] \|_{\infty} < \infty,$$

and that the left-hand side tends to 0 as $t - s \rightarrow \infty$. This and (3.2) give

$$\begin{split} \sup_{x \in E, \boldsymbol{f} \in \operatorname{Ei}_{1}(\Lambda_{L})^{k}} \left| \int_{0}^{t} \psi_{s} \left[\gamma(\mathrm{e}^{-\tilde{\lambda}(f)t} \zeta_{[k]}[\psi_{t-s}^{(\cdot)}[\boldsymbol{f}_{t}]] - \mathrm{e}^{-\tilde{\lambda}(f)s} \zeta_{[k]}[L_{\cdot}[\boldsymbol{f}_{s}]]) \right](x) \mathrm{d}s \right| \\ \leq \sup_{\boldsymbol{f} \in \operatorname{Ei}_{1}(\Lambda_{L})^{k}} C_{1} \int_{0}^{t} \mathrm{e}^{(\lambda_{1} - \operatorname{Re}\tilde{\lambda}(f))s} (1+s)^{p_{1}-1} \| \mathrm{e}^{-\tilde{\lambda}(f)(t-s)} \zeta_{[k]}[\psi_{t-s}^{(\cdot)}[\boldsymbol{f}_{t}]](x) - \zeta_{[k]}[L_{\cdot}[\boldsymbol{f}_{s}]] \|_{\infty} \mathrm{d}s \\ \leq C_{2} \int_{0}^{t} \mathrm{e}^{(\lambda_{1} - \operatorname{Re}\tilde{\lambda}(f))s} (1+s)^{\tilde{p}(f) + p_{1} - (k+1)} g(t-s) \mathrm{d}s, \end{split}$$

where $g: \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ satisfies $g(t) \to 0$ as $t \to \infty$. Therefore,

$$\lim_{t \to \infty} \sup_{x \in E, \boldsymbol{f} \in \mathrm{Ei}_1(\Lambda_L)^k} \left| \int_0^t \psi_s \left[\gamma(\mathrm{e}^{-\tilde{\lambda}(f)t} \zeta_{[k]}[\psi_{t-s}^{(\cdot)}[\boldsymbol{f}_t]] - \mathrm{e}^{-\tilde{\lambda}(f)s} \zeta_{[k]}[L_{\cdot}[\boldsymbol{f}_s]]) \right](x) \mathrm{d}s \right| = 0.$$
(3.7)

Using (3.6) and (3.7) in (3.5) gives us Lemma 3.1.

Returning to the proof of the theorem, let $k \geq 1$ and note that for any $\mathbf{f} \in \text{Ei}(\Lambda_L)^k$, we have

$$\boldsymbol{f} = \mathrm{e}^{\mathcal{N}t} \boldsymbol{f}_t.$$

By definition of \mathcal{N} and (3.1),

$$\sup_{\substack{f \in \text{Ei}_{1}(\Lambda_{L})}} \lim_{t \to \infty} \|t^{-(p(f)-1)} e^{\mathcal{N}t} f - (p(f)-1)!^{-1} \mathcal{N}^{p(f)-1} f\|_{\infty} = 0,$$
$$\sup_{f \in \text{Ei}_{1}(\Lambda_{L})} \|\mathcal{N}^{p(f)-1} f\|_{\infty} \le C \|f\|_{\infty}.$$

The corollary follows from this and applying Lemma 3.1 to the functions

$$(\mathcal{N}^{p(f_1)-1}f,\ldots,\mathcal{N}^{p(f_k)-1}f), \quad f\in \mathrm{Ei}_1(\Lambda_L).$$

3.2 Proof of Theorem 2.2

We start by showing the case of k = 2. For this, we will actually prove the following stronger result.

Lemma 3.2. Assume that

$$\sup_{x\in E}\mathcal{E}_x[N^2]<\infty$$

For $t \geq 0$, let

$$\Delta_{2,t}^{*} = \sup_{x \in E, \boldsymbol{f} \in \text{Ei}_{1}(\Lambda_{S})^{2}} \left| e^{-(\lambda_{1} + \mathcal{N})t} \psi_{t}^{(2)}[\boldsymbol{f}](x) - L_{[2]}^{*}[\boldsymbol{f}](x) \right|, \qquad (3.8)$$

where

$$L_{[2]}^{*}[\boldsymbol{f}](x) = \Phi_{1}[f_{1}f_{2}](x) + \int_{0}^{\infty} e^{-(\lambda_{1}+\mathcal{N})s} \Phi_{1}[\gamma\zeta_{[2]}[\psi_{s}^{(\cdot)}[\boldsymbol{f}]]](x) ds.$$

Then

$$\sup_{t \ge 0} \Delta_{2,t}^* < \infty \text{ and } \lim_{t \to \infty} \Delta_{2,t}^* = 0$$

By Lemma 4.1, for any $\boldsymbol{f} \in B(E)^k$, $x \in E$, and $t \ge 0$,

$$\psi_t^{(2)}[\mathbf{f}](x) = \psi_t \left[f_1 f_2 \right](x) + \int_0^t \psi_{t-s} \left[\gamma \zeta_{[2]}[\psi_s^{(\cdot)}[\mathbf{f}]] \right](x) \mathrm{d}s.$$
(3.9)

For the first term on the right-hand side, by (H1), we have that

$$\lim_{t \to \infty} \sup_{x \in E, \mathbf{f} \in \text{Ei}_1(\Lambda_S)^2} \left| e^{-(\lambda_1 + \mathcal{N})t} \psi_t \left[f_1 f_2 \right](x) - \Phi_1[f_1 f_2](x) \right| = 0.$$
(3.10)

Similarly, for the integrand in (3.9), we have that

$$\lim_{t \to \infty} \sup_{s \ge 0} \sup_{x \in E, \mathbf{f} \in \text{Ei}_1(\Lambda_S)^2} \frac{\left| e^{-(\lambda_1 + \mathcal{N})t} \psi_t \left[\gamma \zeta_{[2]} [\psi_s^{(\cdot)}[\mathbf{f}]] \right] (x) - \Phi_1 [\gamma \zeta_{[2]} [\psi_s^{(\cdot)}[\mathbf{f}]] (x) \right|}{1 + \|\gamma \zeta_{[2]} [\psi_s^{(\cdot)}[\mathbf{f}]] \|_{\infty}} = 0.$$
(3.11)

Next, by (3.2) and (3.3), there exists $\varepsilon > 0$, such that

$$\sup_{\boldsymbol{f}\in \mathrm{Ei}_{1}(\Lambda_{S})^{2}} \|\gamma\zeta_{[2]}[\psi_{s}^{(\cdot)}[\boldsymbol{f}]]\|_{\infty} \leq C\mathrm{e}^{(\lambda_{1}-\varepsilon)t},$$

where we have used $2\text{Re}\lambda_{\tau} < \lambda_1$. This, (3.11), and (3.1) imply

$$\sup_{x \in E, \boldsymbol{f} \in \operatorname{Ei}_{1}(\Lambda_{S})^{2}} \left| \int_{0}^{t} e^{-(\lambda_{1}+\mathcal{N})t} \psi_{t-s} \left[\gamma \zeta_{[2]}[\psi_{s}^{(\cdot)}[\boldsymbol{f}]] \right](x) - e^{-(\lambda_{1}+\mathcal{N})s} \Phi_{1}[\gamma \zeta_{[2]}[\psi_{s}^{(\cdot)}[\boldsymbol{f}]](x) ds \right| \\
\leq C_{1} \sup_{\boldsymbol{f} \in \operatorname{Ei}_{1}(\Lambda_{S})^{2}} \int_{0}^{t} g(t-s)(1+s)^{p_{1}-1} e^{-\lambda_{1}s}(1+\|\gamma \zeta_{[2]}[\psi_{s}^{(\cdot)}[\boldsymbol{f}]\|_{\infty}) ds \\
\leq C_{2} \int_{0}^{t} g(t-s)(1+s)^{p_{1}-1} e^{-\varepsilon s} ds,$$

where $g: \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ satisfies $g(t) \to 0$ as $t \to \infty$, and to exchange the integral and operator \mathcal{N} , we have used that \mathcal{N} is continuous since it is bounded. Thus, the right-hand side tends to 0 as $t \to \infty$. Using this and (3.10) in (3.5) gives Lemma 3.2.

We return to the proof of Theorem 2.2. To obtain the case of k = 2 from Lemma 3.2, we use that, for the eigenspace with eigenvalue λ_1 , the dominating term of $e^{\mathcal{N}t}$ is $(p_1 - 1)!^{-1}t^{p_1-1}\mathcal{N}^{p_1-1}$. Furthermore, by (H1), for $f \in B(E)$,

$$\mathcal{N}^{p_1-1}\Phi_1[f] = \Phi_{1,1}[\mathcal{N}^{p_1-1}f]$$

Thus, (2.3) for k = 2 follows from this and (3.8). The inductive step that completes the proof follows an identical structure to the proof of Theorem 2.1. We thus omit the details.

3.3 Proof of Theorem 2.4

We start by showing the case of k = 2. Similarly to the proof of Theorem 2.2, we prove the following stronger result.

Lemma 3.3. Assume that

$$\sup_{x\in E}\mathcal{E}_x[N^2]<\infty$$

For $0 \leq \alpha \leq p_1 - 1$, let

$$\Delta_{2,t}^{\alpha} = \sup_{x \in E, \boldsymbol{f} \in \text{Ei}_{1}(\Lambda_{C})^{2}} \left| e^{-\frac{\ell \lambda_{1}}{2}} (1+t)^{-(\tilde{p}(\boldsymbol{f})+p_{1}-\alpha-2)} \mathcal{N}^{\alpha} \psi_{t}^{(2)}[\boldsymbol{f}](x) - L_{[2]}^{\alpha}[\boldsymbol{f}](x) \right|, \quad (3.12)$$

where λ_1 and p_1 were given in (H1), and,

$$L_{[2]}^{\alpha}[\boldsymbol{f}](x) = \frac{\Phi_{1,1}[\mathcal{N}^{p_1-1}(\gamma\zeta_{[2]}[\Phi_{\lambda(\cdot),1}[\boldsymbol{f}^*]])](x)(\tilde{p}(\boldsymbol{f})-2)!}{(p(f_1)-1)!(p(f_2)-1)!(p_1+\tilde{p}(\boldsymbol{f})-\alpha-2)!}, \quad \boldsymbol{f} \text{ conjugate 2-tuple}.$$

Otherwise $L^{\alpha}_{A}[\mathbf{f}] = 0$. Then, for $1 \leq \ell \leq k$,

$$\sup_{t\geq 0} \Delta_{2,t} < \infty \ and \ \lim_{t\to\infty} \Delta_{2,t} = 0.$$

Proof. Fix $0 \le \alpha \le p_1 - 1$. By Lemma 4.1, for any $\mathbf{f} \in B(E)^k$, $x \in E$, and $t \ge 0$,

$$\psi_t^{(2)}[\boldsymbol{f}](x) = \psi_t \left[f_1 f_2 \right](x) + \int_0^t \psi_{t-s} \left[\gamma \zeta_{[2]}[\psi_s^{(\cdot)}[\boldsymbol{f}]] \right](x) \mathrm{d}s.$$
(3.13)

For ease of notation, let $\beta(\mathbf{f}) = \tilde{p}(\mathbf{f}) + p_1 - \alpha - 2$. For the first term on the right-hand side, an identical argument to (3.10) implies

$$\lim_{t \to \infty} \sup_{x \in E, \mathbf{f} \in \operatorname{Ei}_{1}(\Lambda_{C})^{2}} \left| (1+t)^{-\beta(\mathbf{f})} \mathrm{e}^{-\lambda_{1} t} \mathcal{N}^{\alpha} \psi_{t} \left[f_{1} f_{2} \right] (x) \right| = 0.$$

We now show convergence of the integral term in (3.13). Let

$$h_{f}(x) = \frac{\gamma \zeta_{[2]}[\Phi_{\lambda(\cdot),1}[f^*]](x)}{(p(f_1) - 1)!(p(f_2) - 1)!},$$

where f^* is as in (2.2). By boundedness of \mathcal{N} and (3.3), there exists constant C, such that

$$\sup_{\boldsymbol{f} \in \mathrm{Ei}_1(\Lambda_C)^2} \|h_{\boldsymbol{f}}\|_{\infty} < C.$$
(3.14)

This, (3.3) and (H1) imply

$$\lim_{t \to \infty} \sup_{x \in E, \boldsymbol{f} \in \operatorname{Ei}_1(\Lambda_C)^2} \left| t^{-(\tilde{p}(\boldsymbol{f})-2)} \mathrm{e}^{-\tilde{\lambda}(\boldsymbol{f})t} \gamma \zeta_{[2]}[\psi_t^{(\cdot)}[\boldsymbol{f}]](x) - h_{\boldsymbol{f}}(x) \right| = 0.$$
(3.15)

Next, by (H1), we have

$$\lim_{t \to \infty} \sup_{s \ge 0} \sup_{x \in E, \boldsymbol{f} \in \text{Ei}_{1}(\Lambda_{C})^{2}} \frac{\left| e^{-\lambda_{1}t} \mathcal{N}^{\alpha} \psi_{t} \left[\gamma \zeta_{[2]} [\psi_{s}^{(\cdot)}[\boldsymbol{f}]] \right] (x) - e^{\mathcal{N}t} \mathcal{N}^{\alpha} \Phi_{1} [\gamma \zeta_{[2]} [\psi_{s}^{(\cdot)}[\boldsymbol{f}]] (x) \right|}{1 + \|\gamma \zeta_{[2]} [\psi_{s}^{(\cdot)}[\boldsymbol{f}]\|_{\infty}} = 0.$$
(3.16)

Thus, by feeding the limit (3.15) into (3.16), and using (3.14), we obtain, for $x \in E$, and $\mathbf{f} \in \text{Ei}_1(\Lambda_C)^2$,

$$\begin{aligned} (1+t)^{-\beta(f)} \left| \int_{0}^{t} e^{-\lambda_{1}t} \mathcal{N}^{\alpha} \psi_{t-s} \left[\gamma \zeta_{[2]} [\psi_{s}^{(\cdot)}[f]] \right] (x) &- e^{(\tilde{\lambda}(f) - \lambda_{1})s} s^{\tilde{p}(f) - 2} e^{\mathcal{N}(t-s)} \mathcal{N}^{\alpha} \Phi_{1}[h_{f}](x) ds \right| \\ &\leq (1+t)^{-\beta(f)} \left| \int_{0}^{t} e^{-\lambda_{1}t} \psi_{t-s} \left[\gamma \zeta_{[2]} [\psi_{s}^{(\cdot)}[f]] \right] (x) - e^{-\lambda_{1}s} e^{\mathcal{N}(t-s)} \mathcal{N}^{\alpha} \Phi_{1}[\gamma \zeta_{[2]} [\psi_{t-s}^{(\cdot)}[f]](x) ds \right| \\ &+ (1+t)^{-\beta(f)} \left| \int_{0}^{t} e^{\mathcal{N}(t-s)} \mathcal{N}^{\alpha} \Phi_{1}[\gamma \zeta_{[2]} [\psi_{t-s}^{(\cdot)}[f]] (x) - e^{(\tilde{\lambda}(f) - \lambda_{1})s} s^{\tilde{p}(f) - 2} e^{\mathcal{N}(t-s)} \mathcal{N}^{\alpha} \Phi_{1}[h_{f}](x) ds \right| \\ &\leq (1+t)^{-\beta(f)} \left| \int_{0}^{t} (1+t-s)^{p_{1}-1-\alpha} (1+s)^{\tilde{p}(f) - 2} g_{1}(t-s) ds \right| \\ &+ (1+t)^{-\beta(f)} \int_{0}^{t} (1+t-s)^{p_{1}-1-\alpha} (1+s)^{\tilde{p}(f) - 2} g_{2}(s) ds \end{aligned}$$

where $g_1, g_2 : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$, C are independent of the choice of x and f, and $g_1(t), g_2(t) \to 0$ as $t \to \infty$. Therefore, the right-hand side tends to 0 as $t \to \infty$. It is left to show convergence of

$$(1+t)^{-\beta(\boldsymbol{f})} \int_0^t \mathrm{e}^{(\tilde{\lambda}(\boldsymbol{f})-\lambda_1)s} s^{\tilde{p}(\boldsymbol{f})-2} \mathrm{e}^{\mathcal{N}(t-s)} \mathcal{N}^{\alpha} \Phi_1[h_{\boldsymbol{f}}](x) \mathrm{d}s \to \begin{cases} 0, \quad \tilde{\lambda}(\boldsymbol{f})-\lambda_1 \in \mathbb{C}, \\ \frac{\Phi_{1,1}[\mathcal{N}^{p_1-1}h_{\boldsymbol{f}}](\tilde{p}(\boldsymbol{f})-2)!}{(p_1+\tilde{p}(\boldsymbol{f})-2)!}, \quad \tilde{\lambda}(\boldsymbol{f})-\lambda_1 = 0. \end{cases}$$

as $t \to \infty$. First note that, by (H1) and (3.14), for any $0 < \varepsilon < 1$,

$$\lim_{t \to \infty} \sup_{\boldsymbol{f} \in \operatorname{Ei}_1(\Lambda_C)^2} t^{\varepsilon} \left\| t^{-(p_1 - \alpha - 1)} e^{\mathcal{N}t} \mathcal{N}^{\alpha} \Phi_1[h_{\boldsymbol{f}}] - \frac{\Phi_{1,1}[\mathcal{N}^{p_1 - 1}h_{\boldsymbol{f}}]}{(p_1 - \alpha - 1)!} \right\|_{\infty} = 0.$$

Thus, uniformly, for any $\boldsymbol{f} \in \mathrm{Ei}_1(\Lambda_C)^2, x \in E$,

$$\lim_{t \to \infty} (1+t)^{-\beta(f)} \left| \int_0^t e^{(\tilde{\lambda}(f) - \lambda_1)s} s^{\tilde{p}(f) - 2} \left(e^{\mathcal{N}(t-s)} \sum_{j=1}^{p_1} \Phi_{1,j}[h_f](x) - \frac{(t-s)^{p_1 - \alpha - 1} \Phi_{1,1}[\mathcal{N}^{p_1 - 1}h_f]}{(p_1 - \alpha - 1)!} \right) ds \right| = 0.$$

as $t \to \infty$. Finally, by standard integration techniques, we have that

$$\lim_{t \to \infty} (1+t)^{-\beta(f)} \frac{\Phi_{1,1}[\mathcal{N}^{p_1-1}h_f]}{(p_1-\alpha-1)!} \int_0^t e^{(\tilde{\lambda}(f)-\lambda_1)s} (t-s)^{p_1-\alpha-1} s^{\tilde{p}(f)-2} ds$$
$$= \begin{cases} 0, \quad \tilde{\lambda}(f) - \lambda_1 \in \mathbb{C}, \\ \frac{\Phi_{1,1}[\mathcal{N}^{p_1-1}h_f](\tilde{p}(f)-2)!}{(p_1+\tilde{p}(f)-\alpha-2)!}, \quad \lambda(f) - \lambda_1 = 0. \end{cases}$$

Returning to the proof of Theorem 2.4, the case of k = 2 is handled by Lemma 3.3 with $\alpha = 0$. The proof of $k \ge 3$ follows an identical structure to the proof of Theorem 2.1. We omit the details.

4 Appendix

In this section, we state an evolution equation relating moments of the form $\prod_{i=1}^{k} X_t[f_i]$ for $f_1, \ldots, f_k \in B(E)$ to lower order product moments.

Lemma 4.1. Fix $k \ge 1$. Assume that

$$\sup_{x \in E} \mathcal{E}_x[N^k] < \infty. \tag{4.1}$$

Then, for any $f_1, \ldots, f_k \in B(E)$, $x \in E$, and $t \ge 0$, we have that

$$\psi_t^{(k)}[\mathbf{f}](x) = \psi_t \left[f_1 \cdots f_k \right](x) + \int_0^t \psi_s \left[\gamma \zeta_{[k]}[\psi_{t-s}^{(\cdot)}[\mathbf{f}]] \right](x) \mathrm{d}s, \quad t \ge 0,$$
(4.2)

where we have used the notation introduced in Section 2.

This result is an extension of [9, Proposition 9.1]. Since the proof is extremely similar, we only sketch the steps and leave the details to the reader.

Sketch proof. The proof follows by induction. We start by assuming that $f_1, \ldots, f_k \in B^+(E)$ with the additional assumption that the functions are real-valued. The result for k = 1 holds trivially since $\zeta_{[1]}[\psi_t^{(\cdot)}[\mathbf{f}]] = 0$ for all $t \ge 0$. Assuming the result holds for $k \ge 2$, for $t \ge 0, x \in E, f \in B^+(E)$ with f real-valued, define

$$u_t[f](x) = \mathbb{E}_{\delta_x} \left[1 - e^{-X_t[f] ds} \right]$$

Then, by [9, Theorem 8.2], for $\theta_1, \ldots, \theta_k > 0$, we have that

$$u_t \left[\sum_{i=1}^k \theta_i f_i\right](x) = \psi_t \left[1 - e^{-\sum_{i=1}^k \theta_i f_i}\right](x) - \int_0^t \psi_s \left[A \left[u_{t-s} \left[\sum_{i=1}^k \theta_i f_i\right]\right]\right](x) ds, \quad (4.3)$$

where

$$A[f](x) = \gamma(x)\mathcal{E}_x\left[\prod_{i=1}^N (1 - f(x_i)) - 1 + \sum_{i=1}^N f(x_i)\right], \quad x \in E.$$

Differentiating with respect to $\theta_1, \ldots, \theta_k$ and setting $\theta_1 = \cdots = \theta_k = 0$, as in [9, Proposition 9.1], yields the result for f_1, \ldots, f_k non-negative and real-valued. To remove the restriction that f_1, \ldots, f_k are non-negative, we can write each f_i as the difference of its positive and negative parts. Some algebra, combinatorics and the result for non-negative functions then yields the result for general bounded, real-valued functions. Finally, to extend the result to complex-valued functions, we write each complex function as the sum of its real and imaginary parts and the proof then follows as in the real case.

Acknowledgements

This work was supported by the EPSRC grant MaThRad EP/W026899/1. Part of this work was completed while the first author was in receipt of a scholarship from the EPSRC Centre for Doctoral Training in Statistical Applied Mathematics at Bath (SAMBa), under the project EP/S022945/1.

References

- S. Asmussen and H. Hering. Strong limit theorems for general supercritical branching processes with applications to branching diffusions. <u>Z. Wahrscheinlichkeitstheorie und Verw. Gebiete</u>, 36(3):195–212, 1976.
- [2] Eric Dumonteil, Emma Horton, Andreas E Kyprianou, and Andrea Zola. Limit theorems for the neutron transport equation. arXiv preprint arXiv:2407.04820, 2024.
- [3] S.D. Durham. Limit theorems for a general critical branching process. <u>Journal of Applied</u> Probability, 8(1):1–16, 1971.
- [4] J. Engländer, S.C. Harris, and A. E. Kyprianou. Strong law of large numbers for branching diffusions. Ann. Inst. Henri Poincaré Probab. Stat., 46(1):279–298, 2010.
- [5] J. Fleischman. Limiting distributions for branching random fields. <u>Trans. Amer. Math. Soc.</u>, 239:353–389, 1978.
- [6] Félix Foutel-Rodier and Emmanuel Schertzer. Convergence of genealogies through spinal decomposition with an application to population genetics. <u>Probability Theory and Related</u> Fields, 187(3):697–751, 2023.
- [7] I. Gonzalez, E. Horton, and A. E. Kyprianou. Asymptotic moments of spatial branching processes. <u>Probability Theory and Related Fields</u>, 184(3-4):805–858, 2022.
- [8] Simon C Harris, Emma Horton, Andreas E Kyprianou, and Ellen Powell. Many-to-few for non-local branching markov process. Electronic Journal of Probability, 29:1–26, 2024.
- [9] E. Horton and A. E. Kyprianou. <u>Stochastic neutron transport and non-local branching Markov</u> processes. Probability and its Applications. Birkhäuser, 2023.
- [10] Emma Horton, Andreas E Kyprianou, Pedro Martín-Chávez, Ellen Powell, and Victor Rivero. Stability of (sub) critical non-local spatial branching processes with and without immigration. arXiv preprint arXiv:2407.05472, 2024.
- [11] I. Iscoe. On the supports of measure-valued critical branching Brownian motion. <u>Ann. Probab.</u>, 16(1):200-221, 1988.
- [12] A. Klenke. Multiple scale analysis of clusters in spatial branching models. <u>Ann. Probab.</u>, 25(4):1670–1711, 1997.