Interacted two-stage least squares with treatment effect heterogeneity

Anqi Zhao Peng Ding Fan Li^{*}

Abstract

Treatment effect heterogeneity with respect to covariates is common in instrumental variable (IV) analyses. An intuitive approach, which we term the interacted two-stage least squares (2SLS), is to postulate a linear working model of the outcome on the treatment, covariates, and treatment-covariate interactions, and instrument it by the IV, covariates, and IV-covariate interactions. We clarify the causal interpretation of the interacted 2SLS under the local average treatment effect (LATE) framework when the IV is valid conditional on the covariates. Our contributions are threefold. First, we show that the interacted 2SLS with centered covariates is consistent for estimating the LATE if either of the following conditions holds: (i) the IV-covariate interactions are linear in the covariates; (ii) the linear outcome model underlying the interacted 2SLS is correct. In particular, condition (i) is satisfied if either (ia) the covariates are categorical or (ib) the IV is randomly assigned. In contrast, existing 2SLS procedures with an additive second stage only recover weighted averages of the conditional LATEs that generally differ from the LATE. Second, we show that the coefficients of the treatment-covariate interactions from the interacted 2SLS are consistent for estimating treatment effect heterogeneity with regard to covariates among compliers if either condition (i) or condition (ii) holds. Moreover, we connect the 2SLS estimator with the weighting perspective in Abadie (2003) and establish the necessity of condition (i) in the absence of additional assumptions on the outcome model. Third, leveraging the consistency guarantees of the interacted 2SLS for categorical covariates, we propose a stratification strategy based on the IV propensity score to approximate the LATE and treatment effect heterogeneity with regard to the IV propensity score when neither condition (i) nor condition (ii) holds.

Keywords: Conditional average treatment effect, instrumental variable, local average treatment effect, potential outcome, treatment effect variation

^{*}Anqi Zhao: Fuqua School of Business, Duke University. Peng Ding: Department of Statistics, University of California, Berkeley. Fan Li: Department of Statistical Science, Duke University. Peng Ding was supported by the U.S. National Science Foundation (grant #1945136).

1 Introduction

Two-stage least squares (2SLS) is widely used for estimating treatment effects when instrumental variables (IVs) are available for endogenous treatments. Under the potential outcomes framework, Imbens and Angrist (1994) and Angrist et al. (1996) defined the local average treatment effect (LATE) as the average treatment effect over the subpopulation of *compliers* whose treatment status is affected by the IV, and formalized the assumptions on the IV that ensure the consistency of 2SLS for estimating the LATE.

Often, IVs only satisfy the IV assumptions after conditioning on a set of covariates. Refer to such IVs as *conditionally valid* IVs hence. A common strategy is to include the corresponding covariates as additional regressors when fitting the 2SLS. When treatment effect heterogeneity with regard to covariates is suspected, an intuitive generalization, which we term the *interacted* 2SLS, is to add the IV-covariate and treatment-covariate interactions to the first and second stages, respectively (Angrist and Pischke, 2009, Section 4.5.2).

To our knowledge, theoretical properties of the interacted 2SLS have not been discussed under the LATE framework except when the IV is randomly assigned hence *unconditionally valid* (Ding et al., 2019). Our paper closes this gap and clarifies the properties of the interacted 2SLS in the more prevalent setting where the IV is conditionally valid.

Our contributions are threefold. First, we show that the interacted 2SLS with centered covariates is consistent for estimating the LATE if either of the following conditions holds:

(i) the IV-covariate interactions are linear in the covariates;

(ii) the linear outcome model underlying the interacted 2SLS is correct.

In particular, condition (i) is satisfied if either (ia) the covariates are categorical or (ib) the IV is randomly assigned. In contrast, existing 2SLS procedures with an additive second

stage only recover weighted averages of the conditional LATEs that generally differ from the LATE (Angrist and Imbens, 1995; Sloczynski, 2022; Blandhol et al., 2022). This illustrates an advantage of including treatment-covariate interactions in the second stage.

Second, define systematic treatment effect variation as the variation in individual treatment effects that can be explained by covariates (Heckman et al., 1997; Djebbari and Smith, 2008). We show that the coefficients of the treatment-covariate interactions from the interacted 2SLS are consistent for estimating the systematic treatment effect variation among compliers if either condition (i) or condition (ii) holds. Furthermore, we connect the 2SLS estimator with the weighting perspective in Abadie (2003), and establish the necessity of condition (i) in the absence of additional assumptions on the outcome model. The results extend the literature on the connection between regression and weighting for estimating the average treatment effect and LATE (Kline, 2011; Chattopadhyay and Zubizarreta, 2023; Chattopadhyay et al., 2024; Słoczyński et al., 2025; ?).

Third, to remedy the possible biases when neither condition (i) nor condition (ii) holds, we propose stratifying based on the estimated propensity score of the IV and using the stratum dummies as the new covariates to fit the interacted 2SLS. While propensity score stratification is standard for estimating the average treatment effect (Rosenbaum and Rubin, 1983) and LATE (Cheng and Lin, 2018; Pashley et al., 2024), we highlight its value in approximating treatment effect heterogeneity with respect to the IV propensity score.

Notation. For a set of tuples $\{(u_i, v_{i1}, \ldots, v_{iL}) : i = 1, \ldots, N\}$ with $u_i \in \mathbb{R}^K$ and $v_{il} \in \mathbb{R}^{K_l}$ for $l = 1, \ldots, L$, denote by $\operatorname{Im}(u_i \sim v_{i1} + \cdots + v_{iL})$ the least squares fit of the *component-wise* linear regression of u_i on $(v_{i1}^\top, \ldots, v_{iL}^\top)^\top$. We allow each v_{il} to be a scalar or a vector and use + to denote concatenation of regressors. Throughout, we focus on the numeric outputs of least squares without invoking any assumption of the

corresponding linear model. For two random vectors $Y \in \mathbb{R}^p$ and $X \in \mathbb{R}^q$, denote by $\operatorname{proj}(Y \mid X)$ the linear projection of Y on X in that $\operatorname{proj}(Y \mid X) = BX$, where $B = \operatorname{argmin}_{b \in \mathbb{R}^{p \times q}} \mathbb{E}\{(Y - bX)^2\} = \mathbb{E}(YX^{\top})\{\mathbb{E}(XX^{\top})\}^{-1}$. Let $1(\cdot)$ denote the indicator function. Let \perp denote independence.

2 Interacted 2SLS and identifying assumptions

2.1 Interacted 2SLS

Consider a study with two treatment levels, indexed by d = 0, 1, and a study population of N units, indexed by i = 1, ..., N. For each unit, we observe a treatment status $D_i \in \{0, 1\}$, an outcome of interest $Y_i \in \mathbb{R}$, a baseline covariate vector $X_i \in \mathbb{R}^K$, and a binary IV $Z_i \in \{0, 1\}$ that is valid conditional on X_i . We assume $X_i = (1, X_{i1}, \ldots, X_{i,K-1})^{\top}$ includes the constant one as the first element unless specified otherwise, and state the IV assumptions in Section 2.2. Let $\mathbb{E}(Z_i \mid X_i) = \mathbb{P}(Z_i = 1 \mid X_i)$ denote the *IV propensity score* of unit *i* (Rosenbaum and Rubin, 1983).

Definition 1 below reviews the standard 2SLS procedure for estimating the causal effect of the treatment on outcome. We call it the *additive* 2SLS to signify the additive regression specifications in both stages.

- **Definition 1** (Additive 2SLS). (i) First stage: Fit $lm(D_i \sim Z_i + X_i)$ over i = 1, ..., N. Denote the fitted values by \hat{D}_i for i = 1, ..., N.
 - (ii) Second stage: Fit $lm(Y_i \sim \hat{D}_i + X_i)$ over i = 1, ..., N. Estimate the treatment effect by the coefficient of \hat{D}_i .

The standard 2SLS in Definition 1 is motivated by the *additive working model* $Y_i = D_i\beta_D + X_i^{\top}\beta_X + \eta_i$, where $\mathbb{E}(\eta_i \mid Z_i, X_i) = 0$. The coefficient of D_i , β_D , is the constant

treatment effect of interest, and we use Z_i to instrument the possibly endogenous D_i . We use the term *working model* to refer to models that are used to construct estimators but not necessarily correct.

When treatment effect heterogeneity with regard to covariates is suspected, an intuitive generalization is to consider the *interacted* working model

$$Y_i = D_i X_i^\top \beta_{DX} + X_i^\top \beta_X + \eta_i, \quad \text{where} \quad \mathbb{E}(\eta_i \mid Z_i, X_i) = 0, \tag{1}$$

with $X_i^{\top} \beta_{DX}$ representing the covariate-dependent treatment effect. We can then use $Z_i X_i$ to instrument the possibly endogenous $D_i X_i$; see, e.g., Angrist and Pischke (2009, Section 4.5.2). We term the resulting 2SLS procedure the *interacted* 2SLS, formalized in Definition 2 below.

- **Definition 2** (Interacted 2SLS). (i) First stage: Fit $lm(D_iX_i \sim Z_iX_i + X_i)$ over $i = 1, \ldots, N$, as the component-wise regression of D_iX_i on $(Z_iX_i^{\top}, X_i^{\top})^{\top}$. Denote the fitted values by \widehat{DX}_i for $i = 1, \ldots, N$.
 - (ii) Second stage: Fit $lm(Y_i \sim \widehat{DX}_i + X_i)$ over i = 1, ..., N. Let $\hat{\beta}_{2SLS}$ denote the coefficient vector of \widehat{DX}_i .

Of interest is the causal interpretation of the coefficient vector $\hat{\beta}_{2\text{SLS}}$.

2.2 IV assumptions and causal estimands

We formalize below the IV assumptions and corresponding causal estimands using the potential outcomes framework (Imbens and Angrist, 1994; Angrist et al., 1996).

For $z, d \in \{0, 1\}$, let $D_i(z)$ denote the potential treatment status of unit *i* if $Z_i = z$, and let $Y_i(z, d)$ denote the potential outcome of unit *i* if $Z_i = z$ and $D_i = d$. The observed treatment status and outcome satisfy $D_i = D_i(Z_i)$ and $Y_i = Y_i(Z_i, D_i) = Y_i(Z_i, D_i(Z_i))$.

Following the literature, we classify the units into four compliance types, denoted by U_i , based on the joint values of $D_i(1)$ and $D_i(0)$. We call unit *i* an *always-taker* if $D_i(1) = D_i(0) = 1$, denoted by $U_i = a$; a *complier* if $D_i(1) = 1$ and $D_i(0) = 0$, denoted by $U_i = c$; a *defier* if $D_i(1) = 0$ and $D_i(0) = 1$, denoted by $U_i = d$; and a *never-taker* if $D_i(1) = D_i(0) = 0$, denoted by $U_i = n$.

Assumption 1 below reviews the assumptions for *conditionally valid* IVs that we assume throughout the paper.

Assumption 1. $\{Y_i(z, d), D_i(z), X_i, Z_i : z = 0, 1; d = 0, 1\}$ are independent and identically distributed across i = 1, ..., N, and satisfy the following conditions:

- (i) Selection on observables: $Z_i \perp \{Y_i(z, d), D_i(z) : z = 0, 1; d = 0, 1\} \mid X_i$.
- (ii) **Overlap:** $0 < \mathbb{E}(Z_i \mid X_i) < 1.$
- (iii) Exclusion restriction: $Y_i(0, d) = Y_i(1, d) = Y_i(d)$ for d = 0, 1, where $Y_i(d)$ denotes the common value.
- (iv) **Relevance:** $\mathbb{E}\{D_i(1) D_i(0) \mid X_i\} \neq 0.$
- (v) Monotonicity: $D_i(1) \ge D_i(0)$.

Assumption 1(i) ensures Z_i is as-if randomly assigned given X_i . Assumption 1(ii) ensures positive probabilities of $Z_i = 1$ and $Z_i = 0$ at all possible values of X_i . Assumption 1(iii) ensures Z_i has no effect on the outcome once we condition on the treatment status. Assumption 1(iv) ensures Z_i has a nonzero causal effect on the treatment status at all possible values of X_i . Assumption 1(v) precludes defiers. Assumption 1(iv)–(v) together ensure positive proportion of compliers at all possible values of X_i , i.e., $\mathbb{P}(U_i = c \mid X_i) > 0$. Recall $Y_i(d)$ as the common value of $Y_i(1, d) = Y_i(0, d)$ for d = 0, 1 under Assumption 1(iii). Define $\tau_i = Y_i(1) - Y_i(0)$ as the individual treatment effect of unit *i*. Define

$$\tau_{\rm c} = \mathbb{E}(\tau_i \mid U_i = {\rm c}) \tag{2}$$

as the local average treatment effect (LATE) on compliers, and define

$$\tau_{\mathbf{c}}(X_i) = \mathbb{E}(\tau_i \mid X_i, U_i = \mathbf{c}) \tag{3}$$

as the conditional LATE given X_i . The law of iterated expectations ensures $\tau_c = \mathbb{E} \{ \tau_c(X_i) \mid U_i = c \}$.

To quantify treatment effect heterogeneity with regard to covariates, consider the linear projection of τ_i on X_i among compliers, denoted by $\operatorname{proj}_{U_i=c}(\tau_i \mid X_i) = X_i^{\top}\beta_c$. The coefficient vector equals

$$\beta_{\mathbf{c}} = \operatorname{argmin}_{b \in \mathbb{R}^{K}} \mathbb{E}\left\{ (\tau_{i} - X_{i}^{\top}b)^{2} \mid U_{i} = \mathbf{c} \right\} = \left\{ \mathbb{E}(X_{i}X_{i}^{\top} \mid U_{i} = \mathbf{c}) \right\}^{-1} \mathbb{E}(X_{i}\tau_{i} \mid U_{i} = \mathbf{c}).$$
(4)

By Angrist and Pischke (2009, Theorem 3.1.6), $X_i^{\top}\beta_c$ is also the linear projection of the conditional LATE $\tau_c(X_i)$ on X_i among compliers with $\beta_c = \operatorname{argmin}_{b \in \mathbb{R}^K} \mathbb{E}[\{\tau_c(X_i) - X_i^{\top}b\}^2 | U_i = c]$. This ensures $X_i^{\top}\beta_c$ is the best linear approximation to both τ_i and $\tau_c(X_i)$ based on X_i , generalizing the idea of systematic treatment effect variation in Djebbari and Smith (2008) to the IV setting; see also Heckman et al. (1997). We define β_c as the causal estimand for quantifying treatment effect heterogeneity among compliers that is explained by X_i .

Recall $\hat{\beta}_{2\text{SLS}}$ as the coefficient vector from the interacted 2SLS in Definition 2. Under the finite-population, design-based framework, Ding et al. (2019) showed that $\hat{\beta}_{2\text{SLS}}$ is consistent for estimating β_c when Z_i is randomly assigned hence valid without conditioning on X_i .

We establish in Section 3 the properties of $\hat{\beta}_{2\text{SLS}}$ for estimating τ_c , $\tau_c(X_i)$, and β_c in the more prevalent setting where Z_i is only conditionally valid given X_i .

2.3 Identifying assumptions

We introduce below two assumptions central to our identification results. First, Assumption 2 below restricts the functional form of the IV propensity score, requiring the product of $\mathbb{E}(Z_i \mid X_i)$ and X_i to be linear in X_i .

Assumption 2 (Linear IV-covariate interactions). $\mathbb{E}(Z_i X_i \mid X_i)$ is linear in X_i .

In contrast, Assumption 3 below imposes linearity on the outcome model.

Assumption 3 (Correct interacted outcome model). The interacted working model (1) is correct.

Proposition 1 below states three implications of Assumptions 2–3.

Proposition 1. Let K denote the dimension of the covariate vector X_i .

- (i) Assumption 2 holds if either of the following conditions holds
 - (a) Categorical covariates: X_i is categorical with K levels.
 - (b) Random assignment of IV: $Z_i \perp \!\!\!\perp X_i$.
- (ii) If Assumption 2 holds, then $\mathbb{E}(Z_i \mid X_i)$ takes at most K distinct values across all possible values of X_i . If Assumption 2 holds for all possible assignment mechanisms of Z_i , then X_i is categorical with K levels.
- (iii) Assume Assumption 1. Assumption 3 holds if the potential outcomes satisfy

 $\mathbb{E}\{Y_i(1) \mid X_i, U_i = \mathbf{a}\} = \mathbb{E}\{Y_i(1) \mid X_i, U_i = \mathbf{c}\} = X_i^\top \beta_1,$ $\mathbb{E}\{Y_i(0) \mid X_i, U_i = \mathbf{n}\} = \mathbb{E}\{Y_i(0) \mid X_i, U_i = \mathbf{c}\} = X_i^\top \beta_0$

for some fixed $\beta_0, \beta_1 \in \mathbb{R}^K$.

Proposition 1(i) provides two important special cases of Assumption 2. Together with Assumption 1, the condition of $Z_i \perp X_i$ in Proposition 1(ib) ensures the IV is valid without conditioning on X_i .

Proposition 1(ii) ensures that the IV propensity score takes at most K distinct values under Assumption 2. This substantially restricts the possible IV assignment mechanisms beyond the two special cases in Proposition 1(i).

Proposition 1(iii) provides a special case of Assumption 3 where the potential outcomes are linear in the covariates.

3 Identification by interacted 2SLS

We establish in this section the conditions for the interacted 2SLS to recover the LATE and treatment effect heterogeneity with regard to covariates in the presence of treatment effect heterogeneity.

3.1 Identification of LATE

We first establish the properties of the interacted 2SLS for estimating the LATE and unify the results with the literature. Assume throughout this subsection that $X_i = (1, X_{i1}, \ldots, X_{i,K-1})^{\top}$ has the constant one as the first element.

3.1.1 Interacted 2SLS with centered covariates

Let $\mu_k = \mathbb{E}(X_{ik} \mid U_i = c)$ denote the complier average of X_{ik} for $k = 1, \dots, K - 1$. While μ_k is generally unknown, we can consistently estimate it by the method of moments proposed by Imbens and Rubin (1997) or the weighted estimator proposed by Abadie (2003). Definition 3 below formalizes our proposed estimator of the LATE.

Definition 3. Given $X_i = (1, X_{i1}, \ldots, X_{i,K-1})^{\top}$, let $\hat{\mu}_k$ be a consistent estimate of $\mu_k = \mathbb{E}(X_{ik} \mid U_i = c)$ for $k = 1, \ldots, K - 1$. Let $\hat{\tau}_{\times \times}$ be the first element of $\hat{\beta}_{2\text{SLS}}$ from the interacted 2SLS in Definition 2 that uses $X_i^0 = (1, X_{i1} - \hat{\mu}_1, \ldots, X_{i,K-1} - \hat{\mu}_{K-1})^{\top}$ as the covariate vector.

We use the subscript $\times \times$ to signify the interacted first and second stages in the interacted 2SLS. Given $D_i X_i^0 = (D_i, D_i (X_{i1} - \hat{\mu}_1), \dots D_i (X_{i,K-1} - \hat{\mu}_{K-1}))^{\top}$, $\hat{\tau}_{\times \times}$ is the coefficient of the fitted value of D_i from the interacted second stage. Let $\tau_{\times \times}$ denote the probability limit of $\hat{\tau}_{\times \times}$ as the sample size N goes to infinity. Theorem 1 below establishes the conditions for $\tau_{\times \times}$ to identify the LATE.

Theorem 1. Assume Assumption 1. Then $\tau_{\times\times} = \tau_c$ if either Assumption 2 or Assumption 3 holds.

Theorem 1 ensures the consistency of $\hat{\tau}_{\times\times}$ for estimating the LATE when either Assumption 2 or Assumption 3 holds. From Proposition 1, this encompasses (i) categorical covariates, (ii) unconditionally valid IV, and (iii) correct interacted working model as three special cases. In particular, the regression specification of the interacted 2SLS is saturated when X_i is categorical, and is therefore a special case of the saturated 2SLS discussed by Blandhol et al. (2022).

Recall that $\hat{\tau}_{\times\times}$ is the first element of $\hat{\beta}_{2\text{SLS}}$ with centered covariates. We provide intuition for the use of centered covariates in Section 3.2 after presenting the more general theory for $\hat{\beta}_{2\text{SLS}}$, and demonstrate the sufficiency of Assumptions 2 and 3 in Section 4. See Hirano and Imbens (2001) and Lin (2013) for similar uses of centered covariates when using interacted regression to estimate the average treatment effect. Note that when computing $\hat{\tau}_{\times\times}$, we use the *estimated* complier averages to center the covariates, because the true values are unknown. To conduct inference based on $\hat{\tau}_{\times\times}$, we need to account for the uncertainty in estimating the complier means. This can be achieved by bootstrapping.

3.1.2 Unification with the literature

Theorem 1 contributes to the literature on using 2SLS to estimate the LATE in the presence of treatment effect heterogeneity. Previously, Angrist and Imbens (1995) studied a variant of the additive 2SLS in Definition 1 for categorical X_i , and showed that the coefficient of \hat{D}_i recovers a weighted average of the conditional LATEs that generally differs from τ_c . Sloczynski (2022) studied the additive 2SLS in Definition 1, and showed that when $\mathbb{E}(Z_i \mid X_i)$ is linear in X_i , the coefficient of \hat{D}_i recovers a weighted average of the conditional LATEs that generally differs from τ_c . Blandhol et al. (2022) studied the additive 2SLS and showed that the coefficient of \hat{D}_i recovers a weighted average of the conditional LATEs that generally differs from τ_c . Blandhol et al. (2022) studied the additive 2SLS and showed that the coefficient of \hat{D}_i recovers a weighted average of the conditional LATEs when the additive first stage is correctly specified, or *saturated*. In contrast, Theorem 1 ensures that the interacted 2SLS directly recovers τ_c if either (i) the covariates are categorical, or (ii) the IV is unconditionally valid, or (iii) the interacted working model (1) is correct. This illustrates an advantage of including the treatment-covariate interactions in the second stage.

We formalize the above overview in Definition 4 and Proposition 2 below. First, Definition 4 states the 2SLS procedure considered by Angrist and Imbens (1995) and Sloczynski (2022).

Definition 4 (Interacted-additive 2SLS). (i) First stage: Fit $lm(D_i \sim Z_i X_i + X_i)$ over i = 1, ..., N. Denote the fitted values by \hat{D}_i for i = 1, ..., N.

(ii) Second stage: Fit $lm(Y_i \sim \hat{D}_i + X_i)$ over i = 1, ..., N. Estimate the treatment effect by the coefficient of \hat{D}_i , denoted by $\hat{\tau}_{\times +}$.

We use the subscript \times + to signify the interacted first stage and additive second stage. The additive, interacted, and interacted-additive 2SLS in Definitions 1–2 and 4 are three variants of 2SLS discussed in the LATE literature, summarized in Table 1. The combination of additive first stage and interacted second stage leads to degenerate second stage and is hence omitted.

Let $\hat{\tau}_{++}$ denote the coefficient of \hat{D}_i from the additive 2SLS in Definition 1. Let τ_{++} and $\tau_{\times+}$ denote the probability limits of $\hat{\tau}_{++}$ and $\hat{\tau}_{\times+}$, respectively. Proposition 2 below states the conditions for τ_{++} and $\tau_{\times+}$ to identify a weighted average of $\tau_c(X_i)$, and contrasts the results with that of $\tau_{\times\times}$.

Let

$$w(X_i) = \frac{\operatorname{var}\{\mathbb{E}(D_i \mid Z_i, X_i) \mid X_i\}}{\mathbb{E}\left[\operatorname{var}\{\mathbb{E}(D_i \mid Z_i, X_i) \mid X_i\}\right]}$$

with $w(X_i) > 0$ and $\mathbb{E}\{w(X_i)\} = 1$. Let $\pi(X_i) = \mathbb{P}(U_i = c \mid X_i)$ denote the proportion of compliers given X_i . Let $\tilde{\pi}(X_i)$ denote the linear projection of $\pi(X_i)$ on X_i weighted by $\operatorname{var}(Z_i \mid X_i)$. That is, $\tilde{\pi}(X_i) = a^{\top}X_i$ with $a = \mathbb{E}\{\operatorname{var}(Z_i \mid X_i) \cdot X_i X_i^{\top}\}^{-1}\mathbb{E}\{\operatorname{var}(Z_i \mid X_i) \cdot X_i \pi(X_i)\}$.

Proposition 2. Assume Assumption 1.

(i) If $\mathbb{E}(Z_i \mid X_i)$ is linear in X_i , then $\tau_{++} = \mathbb{E}\{w_+(X_i) \cdot \tau_c(X_i)\}$, where

$$w_{+}(X_{i}) = \frac{\operatorname{var}(Z_{i} \mid X_{i}) \cdot \pi(X_{i})}{\mathbb{E}\left\{\operatorname{var}(Z_{i} \mid X_{i}) \cdot \pi(X_{i})\right\}} \quad \text{with} \quad w_{+}(X_{i}) > 0 \text{ and } \mathbb{E}\left\{w_{+}(X_{i})\right\} = 1.$$

Further assume that $\mathbb{E}(D_i \mid Z_i, X_i)$ is linear in (Z_i, X_i) , i.e., the additive first stage in

Definition 1 is correctly specified. Then $w_+(X_i) = w(X_i)$.

(ii) If Assumption 2 holds, then $\tau_{\times +} = \mathbb{E}\{w_{\times}(X_i) \cdot \tau_{c}(X_i)\}$, where

$$w_{\times}(X_i) = \frac{\operatorname{var}(Z_i \mid X_i) \cdot \tilde{\pi}(X_i) \cdot \pi(X_i)}{\mathbb{E}\left\{\operatorname{var}(Z_i \mid X_i) \cdot \tilde{\pi}(X_i)^2\right\}} \quad \text{with} \quad \mathbb{E}\left\{w_{\times}(X_i)\right\} = 1$$

Further assume that $\mathbb{E}(D_i \mid Z_i, X_i)$ is linear in $(Z_i X_i, X_i)$, i.e., the interacted first stage in Definition 4 is correctly specified. Then $w_{\times}(X_i) = w(X_i)$.

(iii) If Assumption 2 or Assumption 3 holds, then $\tau_{\times\times} = \tau_c$.

Proposition 2(i) reviews Sloczynski (2022, Corollary 3.4) and ensures that when the IV propensity score $\mathbb{E}(Z_i \mid X_i)$ is linear in X_i , $\hat{\tau}_{++}$ recovers a weighted average of $\tau_c(X_i)$ that generally differs from τ_c . We further show that $w_+(X_i)$ simplifies to $w(X_i)$ when the additive first stage is correctly specified. See Sloczynski (2022) for a detailed discussion about the deviation of τ_{++} from τ_c .

Proposition 2(ii) extends Angrist and Imbens (1995) on categorical covariates to general covariates, and ensures that when $\mathbb{E}(Z_iX_i \mid X_i)$ is linear in X_i , $\hat{\tau}_{\times +}$ recovers a weighted average of $\tau_c(X_i)$ that generally differs from τ_c . A technical subtlety is that $w_{\times}(X_i)$ may be negative due to $\tilde{\pi}(X_i)$, so that some $\tau_c(X_i)$ may be negatively weighted. Note that categorical covariates satisfy both Assumption 2 and the condition that $\mathbb{E}(D_i \mid Z_i, X_i)$ is linear in (Z_iX_i, X_i) . Proposition 2(ii) then simplifies to the special case in Angrist and Imbens (1995).

Proposition 2(iii) follows from Theorem 1 and ensures that when $\mathbb{E}(Z_i X_i \mid X_i)$ is linear in X_i , $\hat{\tau}_{\times\times}$ from the interacted 2SLS directly recovers τ_c .

Together, Proposition 2 illustrates the advantage of $\hat{\tau}_{\times\times}$ over $\hat{\tau}_{\times+}$ and $\hat{\tau}_{++}$ for estimating the LATE, especially when the covariates are categorical or the IV is unconditionally valid;

c.f. Proposition 1. See Table 1 for a summary and Section 6 for a simulated example.

	Second stage			
First stage	Additive	Interacted		
	${\tt lm}(Y_i\sim \hat{D}_i+X_i)$	${\tt lm}(Y_i\sim \widehat{DX}_i+X_i)$		
$\begin{array}{c} \text{Additive} \\ & \texttt{lm}(D_i \sim Z_i + X_i) \end{array}$	$\hat{\tau}_{++} \xrightarrow{\mathbb{P}} \mathbb{E} \{ w_+(X_i) \cdot \tau_c(X_i) \}$ if $\mathbb{E}(Z_i \mid X_i)$ is linear in X_i	n.a.		
Interacted	$\hat{\tau}_{\times +} \xrightarrow{\mathbb{P}} \mathbb{E} \left\{ w_{\times}(X_i) \cdot \tau_{\mathrm{c}}(X_i) \right\}$ if Assumption 2 holds	$\hat{\tau}_{\times\times} \xrightarrow{\mathbb{P}} \tau_{c}$ if if Assumption 2 or Assumption 3 holds		

Table 1: 2SLS for estimating the LATE.

Note. $\hat{\tau}_{++}$ denotes the coefficient of \hat{D}_i from the additive 2SLS in Definition 1. $\hat{\tau}_{\times+}$ denotes the coefficient of \hat{D}_i from the interacted-additive 2SLS in Definition 4. $\hat{\tau}_{\times\times}$ denotes the coefficient of \hat{D}_i from the interacted 2SLS with centered covariates in Definition 3. $w_+(X_i)$ and $w_{\times}(X_i)$ are defined in Proposition 2.

3.2 Identification of treatment effect heterogeneity

We establish in this subsection the properties of the interacted 2SLS for estimating $\beta_{\rm c}$ that quantifies treatment effect heterogeneity with regard to covariates. Recall $\hat{\beta}_{2\rm SLS}$ as the coefficient vector of $D_i X_i$ from the interacted 2SLS in Definition 2. Let $\beta_{2\rm SLS}$ denote the probability limit of $\hat{\beta}_{2\rm SLS}$.

3.2.1 General theory

Let $\operatorname{proj}(D_iX_i \mid Z_iX_i, X_i)$ denote the linear projection of D_iX_i on $(Z_iX_i^{\top}, X_i^{\top})^{\top}$. Let \widetilde{DX}_i denote the residual from the linear projection of $\operatorname{proj}(D_iX_i \mid Z_iX_i, X_i)$ on X_i . Let $\epsilon_i = \tau_i - X_i^{\top}\beta_c$ for compliers, as the residual from the linear projection of τ_i on X_i among compliers. Let $\Delta_i = Y_i(0)$ for compliers and never-takers, and $\Delta_i = Y_i(1) - X_i^{\top}\beta_c$ for

always-takers. Let

$$B_{1} = C_{1} \cdot \mathbb{E}\left[\left\{\mathbb{E}(Z_{i}X_{i} \mid X_{i}) - \operatorname{proj}(Z_{i}X_{i} \mid X_{i})\right\} \cdot \left\{\mathbb{E}(\Delta_{i} \mid X_{i}) - \operatorname{proj}(\Delta_{i} \mid X_{i})\right\}\right],$$

$$B_{2} = C_{1}(I_{K} - C_{2}) \cdot \mathbb{E}\left\{\mathbb{E}(Z_{i}X_{i} \mid X_{i}) \cdot \epsilon_{i} \mid U_{i} = c\right\} \cdot \mathbb{P}(U_{i} = c),$$
(5)

where C_1 is the coefficient matrix of $Z_i X_i$ in $\operatorname{proj}(D_i X_i \mid Z_i X_i, X_i)$ and C_2 is the coefficient matrix of X_i in $\operatorname{proj}(Z_i X_i \mid X_i)$.

Theorem 2. Assume Assumption 1. Then $\beta_{2\text{SLS}} = \beta_c + \{\mathbb{E}(\widetilde{DX}_i \cdot \widetilde{DX}_i^{\mathsf{T}})\}^{-1}(B_1 + B_2),$ where $B_1 = B_2 = 0$ if either Assumption 2 or Assumption 3 holds.

Theorem 2 ensures that $\beta_{2_{\text{SLS}}}$ identifies β_c if and only if $B_1 + B_2 = 0$, and establishes Assumption 2 and Assumption 3 as two sufficient conditions. We discuss the sufficiency and "almost necessity" of these two assumptions in Section 4.

Recall that Assumption 2 holds when the IV is randomly assigned. Therefore, Theorem 2 ensures $\beta_{2\text{SLS}} = \beta_c$ for randomly assigned IVs as a special case. See Ding et al. (2019, Theorem 7) for similar results in the finite-population, design-based framework.

In addition, recall that Assumption 2 holds when the covariates are categorical. Therefore, Theorem 2 ensures $\beta_{2\text{SLS}} = \beta_c$ for categorical covariates. As it turns out, the elements of β_c equal the category-specific LATEs if we define X_i as the vector of all category dummies, without the constant one. This allows us to recover the conditional LATEs as coefficients of the interacted 2SLS for categorical covariates. We formalize the intuition below.

3.2.2 The special case of categorical covariates

Assume throughout this subsection that the covariate vector X_i encodes a K-level categorical variable, represented by $X_i^* \in \{1, \dots, K\}$, using category dummies:

$$X_i = (1(X_i^* = 1), \dots, 1(X_i^* = K))^{\top}.$$
(6)

Define

$$\tau_{[k]c} = \mathbb{E}(\tau_i \mid X_i^* = k, U_i = c)$$

as the conditional LATE on compliers with $X_i^* = k$; c.f. Eq. (3). Lemma 1 below states the simplified forms of β_c and $\tau_c(X_i)$ under (6).

Lemma 1. Assume (6). Then $\beta_{c} = (\tau_{[1]c}, \ldots, \tau_{[K]c})^{\top}$ and $\tau_{c}(X_{i}) = X_{i}^{\top}\beta_{c}$.

Lemma 1 has two implications. First, when X_i is the dummy vector, the corresponding β_c has the subgroup LATEs $\tau_{[k]c}$'s as its elements, which adds interpretability. Second, with categorical X_i , the conditional LATE $\tau_c(X_i)$ is linear in X_i and equals the linear projection $X_i^{\top}\beta_c$. This is not true for general X_i , so that linear projection is only an approximation.

Corollary 1 below follows from Theorem 2 and Lemma 1, and ensures the coefficients of the interacted 2SLS are consistent for estimating the conditional LATEs.

Corollary 1. Assume Assumption 1 and (6). We have $\beta_{2sLs} = (\tau_{[1]c}, \ldots, \tau_{[K]c})^{\top}$.

3.3 Reconciliation of Theorems 1 and 2

We now clarify the connection between Theorems 1 and 2, and provide intuition for the use of centered covariates in estimating the LATE.

Recall that $\hat{\tau}_{\times\times}$ is the first element of $\hat{\beta}_{2\text{SLS}}$ with centered covariates. Theorem 2 ensures that $\hat{\beta}_{2\text{SLS}}$ is consistent for estimating β_c when either Assumption 2 or Assumption 3 holds. Therefore, a logical basis for using $\hat{\tau}_{\times\times}$ to estimate τ_c is that τ_c equals the first element of β_c . Lemma 2 below ensures that this condition is satisfied when the nonconstant covariates are centered by their respective complier averages. This establishes Theorem 1 as a direct consequence of Theorem 2, and justifies the use of centered covariates for estimating the LATE.

Lemma 2. Assume $X_i = (1, X_{i1}, \ldots, X_{i,K-1})^{\top}$. If $\mathbb{E}(X_{ik} \mid U_i = c) = 0$ for $k = 1, \ldots, K-1$, then the first element of β_c equals τ_c .

4 Discussion of Assumptions 2 and 3

We discuss in this section the sufficiency and "almost necessity" of Assumptions 2 and 3 to ensure $\beta_{2SLS} = \beta_c$ in Theorem 2.

4.1 Sufficiency and "almost necessity" of Assumption 2

We discuss below the sufficiency and "almost necessity" of Assumption 2. We first demonstrate the sufficiency of Assumption 2 to ensure $B_1 = B_2 = 0$, as a sufficient condition for $\beta_{2_{SLS}} = \beta_c$ by Theorem 2. We then adopt the weighting perspective in Abadie (2003) and establish the necessity of Assumption 2 to ensure $\beta_{2_{SLS}} = \beta_c$ in the absence of additional assumptions on the outcome model beyond Assumption 1.

4.1.1 Sufficiency

Recall the definitions of B_1 and B_2 from Eq. (5), where C_2 denotes the coefficient matrix of X_i in $\text{proj}(Z_iX_i \mid X_i)$. Assumption 2 ensures

$$\mathbb{E}(Z_i X_i \mid X_i) = \operatorname{proj}(Z_i X_i \mid X_i) = C_2 X_i$$
(7)

so that $B_1 = 0$.

In addition, recall that ϵ_i is the residual from the linear projection of τ_i on X_i among compliers. Properties of linear projection ensure $\mathbb{E}(X_i\epsilon_i \mid U_i = c) = 0$ so that the expectation term in B_2 satisfies

$$\mathbb{E}\Big\{\mathbb{E}(Z_iX_i \mid X_i) \cdot \epsilon_i \mid U_i = c\Big\} \stackrel{(7)}{=} \mathbb{E}(C_2X_i\epsilon_i \mid U_i = c) = 0.$$

This ensures $B_2 = 0$ and therefore the sufficiency of Assumption 2 to ensure $B_1 = B_2 = 0$.

4.1.2 Necessity in the absence of additional assumptions on outcome model

Proposition 3 below expresses $\beta_{2_{\text{SLS}}}$ as a weighted estimator and establishes the necessity of Assumption 2 to ensure $\beta_{2_{\text{SLS}}} = \beta_c$ in the absence of additional assumptions on the outcome model. Recall that \widetilde{DX}_i denotes the residual from the linear projection of $\operatorname{proj}(D_iX_i \mid Z_iX_i, X_i)$ on X_i .

Proposition 3. (i) $\beta_{2\text{SLS}} = \mathbb{E}(w_i Y_i)$, where $w_i = \{\mathbb{E}(\widetilde{DX}_i \cdot \widetilde{DX}_i^{\top})\}^{-1} \widetilde{DX}_i$.

(ii) Under Assumption 1, we have $\beta_{c} = \mathbb{E}(u_{i}Y_{i})$, where $u_{i} = \{\mathbb{E}(X_{i}X_{i}^{\top} \mid U_{i} = c)\}^{-1}\pi_{c}^{-1}\Delta\kappa X_{i}$ with

$$\Delta \kappa = \frac{Z_i - \mathbb{E}(Z_i \mid X_i)}{\mathbb{E}(Z_i \mid X_i) \{1 - \mathbb{E}(Z_i \mid X_i)\}}$$

(iii) If $w_i = u_i$, then Assumption 2 must hold.

Proposition 3(i) expresses β_{2SLS} as a weighted average of the outcome Y_i . The result is numeric and does not require Assumption 1.

Proposition 3(ii) follows from Abadie (2003) and expresses β_c as a weighted average of Y_i when Assumption 1 holds.

Proposition 3(iii) connects Proposition 3(i)–(ii) and establishes the necessity of Assumption 2 to ensure $w_i = u_i$.

Together, Proposition 3(i)–(iii) establish the necessity of Assumption 2 to ensure $\beta_{2SLS} = \beta_c$ in the absence of additional assumptions on the outcome model.

4.2 Sufficiency and "almost necessity" of Assumption 3

We discuss below the sufficiency and "almost necessity" of Assumption 3 to ensure $B_1 = B_2 = 0$ in the absence of additional assumptions on Z_i beyond Assumption 1.

Without additional assumptions on Z_i , the $\mathbb{E}(Z_iX_i \mid X_i)$ in the definitions of B_1 and B_2 is arbitrary. To ensure $B_1 = 0$, we generally need $\mathbb{E}(\Delta_i \mid X_i) = \text{proj}(\Delta_i \mid X_i)$, which is equivalent to

$$\mathbb{E}(\Delta_i \mid X_i) \text{ is linear in } X_i. \tag{8}$$

In addition, Lemma S4 in the Supplementary Material ensures that $C_1(I_K - C_2)$ is invertible. From Eq. (5), we have $B_2 = 0$ if and only if

$$\mathbb{E}\Big\{\mathbb{E}(Z_i X_i \mid X_i) \cdot \epsilon_i \mid U_i = c\Big\} = 0.$$
(9)

By the law of iterated expectations, the left-hand side of condition (9) equals

$$\mathbb{E}\Big\{\mathbb{E}(Z_iX_i \mid X_i) \cdot \epsilon_i \mid U_i = c\Big\} = \mathbb{E}\Big[\mathbb{E}\Big\{\mathbb{E}(Z_iX_i \mid X_i) \cdot \epsilon_i \mid X_i, U_i = c\Big\} \mid U_i = c\Big]$$
$$= \mathbb{E}\Big[\mathbb{E}(Z_iX_i \mid X_i) \cdot \mathbb{E}(\epsilon_i \mid X_i, U_i = c) \mid U_i = c\Big].$$

When $\mathbb{E}(Z_i X_i \mid X_i)$ is arbitrary, condition (9) generally requires $\mathbb{E}(\epsilon_i \mid X_i, U_i = c) = 0$, which is equivalent to

$$\tau_{\rm c}(X_i) = \mathbb{E}(\tau_i \mid X_i, U_i = {\rm c}) \text{ is linear in } X_i.$$
⁽¹⁰⁾

The above observations suggest the sufficiency and "almost necessity" of the combination of conditions (8) and (10) to ensure $B_1 = B_2 = 0$ in the absence of additional assumptions on Z_i . We further show in Lemma S5 in the Supplementary Material that under Assumption 1, Assumption 3 is equivalent to the combination of conditions (8) and (10). This demonstrates the sufficiency and "almost necessity" of Assumption 3 to ensure $B_1 = B_2 = 0$ in the absence of additional assumptions on Z_i .

Remark 1. From Theorem 2 and Lemma 2, $\tau_{\times\times} = \tau_c$ if and only if

the first dimension of
$$\{\mathbb{E}(\widetilde{DX}_i \cdot \widetilde{DX}_i^{\mathsf{T}})\}^{-1}(B_1 + B_2)$$
 equals 0, (11)

which is weaker than the condition of $B_1 + B_2 = 0$ to ensure $\beta_{2\text{SLS}} = \beta_c$. Nonetheless, we did not find intuitive relaxations of Assumptions 2 and 3 to satisfy (11), and therefore invoke the same identifying conditions for τ_c and β_c in Theorems 1 and 2.

5 Stratification based on IV propensity score

The results in Section 3 suggest that the interacted 2SLS may be inconsistent for estimating the LATE and treatment effect heterogeneity when neither Assumption 2 or Assumption 3 holds for the original covariates. When this is the case, we propose stratifying based on the IV propensity score and using the stratum dummies as the new covariates to fit the interacted 2SLS. The consistency of the interacted 2SLS for categorical covariates provides heuristic justification for using the resulting 2SLS coefficients to approximate the LATE and treatment effect heterogeneity with regard to the IV propensity score. We formalize the intuition below. See Cheng and Lin (2018) and Pashley et al. (2024) for other applications of stratification based on the IV propensity score.

Assume that X_i^* is a general covariate vector that ensures the conditional validity of Z_i . Let $e_i = \mathbb{P}(Z_i = 1 \mid X_i^*)$ denote the IV propensity score of unit *i*, and define

$$\tau_{\mathbf{c}}(e_i) = \mathbb{E}(\tau_i \mid e_i, U_i = \mathbf{c})$$

as the conditional LATE given e_i ; c.f. Eq. (3). Properties of the propensity score (Rosenbaum and Rubin, 1983) ensure $Z_i \perp \{Y_i(z, d), D_i(z) : z = 0, 1; d = 0, 1\} | e_i$ so that Z_i is also conditionally valid given e_i . This, together with the consistency guarantees of the interacted 2SLS for categorical covariates, motivates viewing e_i as a new covariate that ensures the validity of Z_i , and fitting the interacted 2SLS with a discretized version of e_i to approximate τ_c and $\tau_c(e_i)$. In most applications, e_i is unknown so we need to first estimate e_i using X_i^* . We formalize the procedure in Algorithm 1 below.

Algorithm 1. (i) Estimate $e_i = \mathbb{P}(Z_i = 1 \mid X_i^*)$ as \hat{e}_i .

(ii) Partition [0, 1] into K strata, denoted by $\{S_k : k = 1, \dots, K\}$. Define $\hat{\mathcal{I}}_i = (1, \mathcal{I}_{i1}, \dots, \mathcal{I}_{i,K-1})^\top$

and $\tilde{\mathcal{I}}_i = (\mathcal{I}_{i1}, \dots, \mathcal{I}_{iK})^{\top}$, where $\mathcal{I}_{ik} = 1(\hat{e}_i \in \mathcal{S}_k)$, as two representations of the stratum of \hat{e}_i . In practice, we can choose K and the cutoffs for \mathcal{S}_k based on the realized values of \hat{e}_i 's.

(iii) Compute $\hat{\tau}_{\times\times}^*$ by letting $X_i = \hat{\mathcal{I}}_i$ in Definition 3 to approximate the LATE. Compute $\hat{\beta}_{2\text{SLS}}^*$ by letting $X_i = \tilde{\mathcal{I}}_i$ in Definition 2. For $k = 1, \ldots, K$, use the *k*th element of $\hat{\beta}_{2\text{SLS}}^*$ to approximate $\tau_c(e_i)$ for $e_i \in \mathcal{S}_k$.

Algorithm 1 transforms arbitrary covariate vector X_i^* into categorical $\hat{\mathcal{I}}_i$ and $\tilde{\mathcal{I}}_i$ to fit the interacted 2SLS. Note that $\hat{\mathcal{I}}_i$ and $\tilde{\mathcal{I}}_i$ are discretized approximations to the actual e_i . Heuristically, an IV that is conditionally valid given e_i can be viewed as "approximately valid" given $\hat{\mathcal{I}}_i$ or $\tilde{\mathcal{I}}_i$. This, together with Theorem 1 and Corollary 1, provides the heuristic justification of the resulting $\hat{\tau}^*_{\times\times}$ and $\hat{\beta}^*_{2SLS}$ to approximate the LATE and subgroup LATES

$$\tau_{[k]c} = \mathbb{E}(\tau_i \mid e_i \in \mathcal{S}_k, U_i = c), \quad \text{where} \quad k = 1, \dots, K,$$
(12)

respectively. The subgroup LATEs in (12) further provide a piecewise approximation to $\tau_{\rm c}(e_i)$ by assuming $\tau_{\rm c}(e_i) \approx \tau_{[k]c}$ for $e_i \in \mathcal{S}_k$. This is analogous to the idea of *regressogram* in nonparametric regression (Liero, 1989; Wasserman, 2006).

6 Simulation

6.1 Interacted 2SLS for estimating LATE

We now use a simulated example to illustrate the advantage of the interacted 2SLS for estimating the LATE. Assume the following model for (X_i, Z_i, D_i, Y_i) :

(i) $X_i = (1, X_{i1})^{\top}$, where $X_{i1} \sim \text{Bernoulli}(0.5)$.




Figure 1: Distributions of $\hat{\tau}_{\times\times}$, $\hat{\tau}_{++}$, and $\hat{\tau}_{\times+}$ over 1000 replications.

- (ii) $Z_i \mid X_i \sim \text{Bernoulli}(0.5 + 0.4X_{i1}).$
- (iii) $D_i = 1(U_i = a) + Z_i \cdot 1(U_i = c)$, where $\mathbb{P}(U_i = a \mid X_i) = 0.1$ and $\mathbb{P}(U_i = c \mid X_i) = 0.7 0.5X_{i1}$.

(iv)
$$Y_i = D_i Y_i(1) + (1 - D_i) Y_i(0)$$
 with $Y_i(0) = 0$ and $Y_i(1) = -1 + 5X_{i1}$.

The data-generating process ensures $\tau_{[1]c} = \mathbb{E}(\tau_i \mid X_{i1} = 1, U_i = c) = 4, \tau_{[2]c} = \mathbb{E}(\tau_i \mid X_{i1} = 0, U_i = c) = -1$, and $\tau_c = 1/9$. For each replication, we generate N = 10000 independent realizations of (X_i, Z_i, D_i, Y_i) and compute $\hat{\tau}_{++}, \hat{\tau}_{\times\times}$, and $\hat{\tau}_{\times+}$ from the additive, interacted, and interacted-additive 2SLS in Definitions 1–4; c.f. Table 1. We use the method of moments to estimate the complier average of X_{i1} in computing $\hat{\tau}_{\times\times}$.

Figure 1 shows the distributions of the three estimators over 1000 replications. Almost all realizations of $\hat{\tau}_{++}$ and $\hat{\tau}_{\times+}$ are below -0.4 while the actual τ_c is positive. In contrast, the distribution of $\hat{\tau}_{\times\times}$ is approximately centered at the actual τ_c , which is coherent with Theorem 1 for categorical covariates.

6.2 Approximation of LATE by Algorithm 1.

We now illustrate the improvement of Algorithm 1 for estimating the LATE. Assume the following model for generating (X_i, Z_i, D_i, Y_i) .

- (i) $X_i = (1, X_{i1}, X_{i2})^{\top}$, where X_{i1} and X_{i2} are independent standard normals.
- (ii) $\mathbb{P}(Z_i \mid X_i) = \{1 + \exp(X_{i1} + X_{i2})\}^{-1}.$

(iii)
$$D_i = Z_i \cdot 1(U_i = c) + 1(U_i = a)$$
, where $\mathbb{P}(U_i = c) = 0.7$ and $\mathbb{P}(U_i = a) = 0.2$.

(iv)
$$Y_i = D_i Y_i(1) + (1 - D_i) Y_i(0)$$
 with $Y_i(0) = 0$ and $Y_i(1) = X_{1i}^2 + X_{2i}^2$

For each replication, we generate N = 1000 independent realizations of (X_i, Z_i, D_i, Y_i) and compute $\hat{\tau}^*_{\times\times}$ by Algorithm 1. We consider K = 5, 10, 15 to evaluate the impact of the number of strata on estimation. For comparison, we also compute $\hat{\tau}_{++}$, $\hat{\tau}_{\times\times}$, and $\hat{\tau}_{\times+}$ from the additive, interacted, and interacted-additive 2SLS in Definitions 1–4.

Figure 2 shows the violin plot of the deviations of the resulting estimators from the LATE over 1000 replications. The three estimators by Algorithm 1 substantially reduce the biases compared with $\hat{\tau}_{++}$, $\hat{\tau}_{\times+}$, and $\hat{\tau}_{\times\times}$. In addition, increasing the number of strata further reduces the bias but increases variability. Table 2 summarizes the means and standard deviations of the deviations.

Table 2: Biases and standard deviations of the 2SLS estimators over 1000 replications.

	â	Â	$\hat{\tau}_{\times \times}$	$\hat{\tau}^*_{\times\times}$ from Algorithm 1		
	7++	$ au_{\times +}$		K = 5	K = 10	K = 15
Bias	-0.559	-0.556	-0.557	-0.106	-0.054	-0.043
Standard dev.	0.144	0.165	0.157	0.124	0.140	0.336




Figure 2: Violin plot of the deviations of the 2SLS estimators from the LATE over 1000 replications. The first three columns correspond to $\hat{\tau}_{++}$, $\hat{\tau}_{\times+}$, and $\hat{\tau}_{\times\times}$, respectively. The last three columns correspond to $\hat{\tau}^*_{\times\times}$ by Algorithm 1 with K = 5, 10, 15.

6.3 Possible inconsistency for estimating β_c

We now illustrate the possible inconsistency of the interacted 2SLS in Theorem 2. Assume the following model for (X_i, Z_i, D_i, Y_i) :

(i)
$$X_i = (1, X_{i1})^{\top}$$
, where $X_{i1} \sim \text{Uniform}(0, 1)$.

(ii)
$$Z_i \sim \text{Bernoulli}(e_i)$$
 with $e_i = X_{i1}$.

(iii)
$$D_i = Z_i \cdot 1(U_i = c) + 1(U_i = a)$$
, where $\mathbb{P}(U_i = c) = 0.7$ and $\mathbb{P}(U_i = a) = 0.2$.

(iv)
$$Y_i = D_i Y_i(1) + (1 - D_i) Y_i(0)$$
 with $Y_i(0) = 0$ and $Y_i(1) = X_{i1}^2$.

The data-generating process ensures $\tau_i = Y_i(1) - Y_i(0) = X_{i1}^2$ and $\beta_c = (-1/6, 1)^{\top}$.

For each replication, we generate N = 1000 independent realizations of (X_i, Z_i, D_i, Y_i) and compute $\hat{\beta}_{2\text{SLS}}$ from the interacted 2SLS in Definition 2. Figure 3 shows the distribution of $\hat{\beta}_{2\text{SLS}} - \beta_c$ over 1000 replications, indicating clear empirical biases in both dimensions.

6.4 Approximation of $\tau_{\mathbf{c}}(e_i)$ by Algorithm 1.

We now illustrate the utility of Algorithm 1 for approximating the conditional LATEs given the IV propensity score. Assume the same model as in Section 6.3 for generating




Figure 3: Distributions of $\hat{\beta}_{2\text{SLS}} - \beta_c$ over 1000 replications. The labels $\hat{\beta}_{2\text{SLS},1} - \beta_{c,1}$ and $\hat{\beta}_{2\text{SLS},2} - \beta_{c,2}$ denote the first and second elements of $\hat{\beta}_{2\text{SLS}} - \beta_c$, respectively.

 (X_i, Z_i, D_i, Y_i) . The data-generating process ensures $\tau_i = X_{i1}^2 = e_i^2$ with $\tau_c(e_i) = \mathbb{E}(\tau_i \mid e_i, U_i = c) = e_i^2$.

To use Algorithm 1 to approximate $\tau_c(e_i)$, we estimate the IV propensity score by the logistic regression of Z_i on X_i , and create the strata by dividing $(\hat{e}_i)_{i=1}^N$ into K equal-sized bins. Figure 4 shows the results at K = 5, 10, 15. The first row is based on one replication to illustrate the evolution of the approximation as K increases. The second row shows the 95% confidence bands based on 3000 replications. The coverage of the confidence band improves as K increases. Note that the logistic regression for estimating e_i is misspecified in this example. The results suggest the robustness of the method to misspecification of the propensity score model with moderate number of strata.

7 Application

We now apply Algorithm 1 to the dataset that Abadie (2003) used to study the effects of 401(k) retirement programs on savings. The data contain cross-sectional observations of




Figure 4: Piecewise approximations of $\tau_c(e_i)$ by Algorithm 1. The first row is based on one replication, and the second row shows the 95% confidence band based on 3000 replications.

N = 9275 units on eligibility for and participation in 401(k) plans along with income and demographic information. The outcome of interest is net financial assets. The treatment variable is an indicator of participation in 401(k) plans, and the IV is an indicator of 401(k) eligibility. Based on Abadie (2003), the IV is conditionally valid after controlling for income, age, and marital status.

Table 3 shows the point estimates of the LATE by the additive, interacted, interactedadditive 2SLS in Definitions 1–4 and Algorithm 1, along with their respective bootstrap standard deviations. Figure 5 shows the piecewise approximations of $\tau_c(e_i)$ based on Algorithm 1 and the 95% confidence bands based on bootstrapping. The result suggests heterogeneity of conditional LATEs across different strata of the propensity score.

Table 3: Point estimates of the LATE and bootstrap standard deviations by the additive, interacted, additive-interacted 2sLs and Algorithm 1 over 1000 replications.

	â	Â	â	$\hat{\tau}^*_{\times\times}$ from Algorithm 1 K = 5 $K = 10$ $K = 15$ $K = 20$			
	7++	7×+	$\gamma_{\times \times}$	K = 5	K = 10	K = 15	K = 20
Point estimate	9.658						
Standard dev.	2.222	2.520	1.913	2.076	2.079	2.081	2.089




Figure 5: Piecewise approximations of $\tau_{\rm c}(e_i)$ by Algorithm 1 and the 95% confidence bands by bootstrapping.

8 Extensions

8.1 Extension to partially interacted 2SLS

Let X_i denote the covariate vector that ensures the conditional validity of the IV. The interacted 2SLS in Definition 2 includes interactions of Z_i and D_i with the whole set of X_i in accommodating treatment effect heterogeneity. When treatment effect heterogeneity is of interest or suspected for only a subset of X_i , denoted by $V_i \subset X_i$, an intuitive modification is to interact Z_i and D_i with only V_i in fitting the 2SLS, formalized in Definition 5 below. See Resnjanskij et al. (2024) for a recent application. See also Hirano and Imbens (2001) and Hainmueller et al. (2019) for applications of the partially interacted regression in the ordinary least squares setting.

- **Definition 5** (Partially interacted 2SLS). (i) First stage: Fit $lm(D_iV_i \sim Z_iV_i + X_i)$ over i = 1, ..., N. Denote the fitted values by $\widehat{DV_i}$ for 1, ..., N.
 - (ii) Second stage: Fit $lm(Y_i \sim \widehat{DV_i} + X_i)$ over i = 1, ..., N. Let $\hat{\beta}'_{2SLS}$ denote the coefficient vector of $\widehat{DV_i}$.

The partially interacted 2SLS in Definition 5 generalizes the interacted 2SLS, and reduces to the interacted 2SLS when $V_i = X_i$. It is the 2SLS procedure corresponding to the working model

$$Y_i = D_i V_i^{\top} \beta_{DV} + X_i^{\top} \beta_X + \eta_i, \quad \text{where} \quad \mathbb{E}(\eta_i \mid Z_i, X_i) = 0, \tag{13}$$

with $V_i^{\top}\beta_{DV}$ representing the covariate-dependent treatment effect. We establish below the causal interpretation of $\hat{\beta}'_{2SLS}$.

Causal estimand. Analogous to the definition of β_c in Eq. (4), let $\operatorname{proj}_{U_i=c}(\tau_i \mid V_i) = V_i^{\top}\beta'_c$ denote the linear projection of τ_i on V_i among compliers with

$$\beta_{\mathbf{c}}' = \operatorname{argmin}_{b \in \mathbb{R}^Q} \mathbb{E}\Big\{ (\tau_i - V_i^\top b)^2 \mid U_i = \mathbf{c} \Big\}.$$

Parallel to the discussion after Eq. (4), $V_i^{\top}\beta'_c$ is also the linear projection of $\tau_c(X_i)$ on V_i among compliers with $\beta'_c = \operatorname{argmin}_{b \in \mathbb{R}^Q} \mathbb{E}[\{\tau_c(X_i) - V_i^{\top}b\}^2 \mid U_i = c]$. This ensures $V_i^{\top}\beta'_c$ is the best linear approximation to both τ_i and $\tau_c(X_i)$ based on V_i . We define β'_c as the causal estimand for quantifying treatment effect heterogeneity with regard to V_i among compliers. **Properties of** $\hat{\beta}'_{2\text{sls}}$ for estimating β'_{c} . Let $\beta'_{2\text{sls}}$ denote the probability limit of $\hat{\beta}'_{2\text{sls}}$. Proposition 4 below generalizes Theorem 2 and states three sufficient conditions for $\beta'_{2\text{sls}}$ to identify β'_{c} .

Proposition 4. Assume $X_i = (V_i^{\top}, W_i^{\top})^{\top} \in \mathbb{R}^K$, where $V_i \in \mathbb{R}^Q$ and $W_i \in \mathbb{R}^{K-Q}$ with $Q \leq K$. Let $\widetilde{DV_i}$ denote the residual of the linear projection of $\operatorname{proj}(D_iV_i \mid Z_iV_i, X_i)$ on X_i . Let $B' = \mathbb{E}\{\widetilde{DV_i} \cdot (Y_i - D_iV_i^{\top}\beta_c')\}$. Assume Assumption 1. Then

$$\beta_{2_{\mathrm{SLS}}}' = \beta_{\mathrm{c}}' + \{\mathbb{E}(\widetilde{DV_i} \cdot \widetilde{DV_i}^{\top})\}^{-1}B',\$$

where B' = 0 if any of the following conditions holds:

- (i) **Categorical covariates:** V_i is categorical with Q levels, and satisfies $\mathbb{E}(Z_i \mid X_i) = \mathbb{E}(Z_i \mid V_i)$ and $\operatorname{proj}(Z_i V_i \mid X_i) = \operatorname{proj}(Z_i V_i \mid V_i)$.
- (ii) Random assignment of IV: $Z_i \perp \!\!\!\perp X_i$.

(iii) Correct interacted outcome model: The interacted working model (13) is correct.

Proposition 4(i) parallels the identification of β_c with categorical covariates by Theorem 2, and states a sufficient condition for β'_{2SLS} to identify β'_c when V_i is categorical. A special case is when X_i encodes two categorical variables and V_i corresponds to one of them.

Proposition 4(ii) parallels the identification of β_c with randomly assigned IV by Theorem 2, and ensures $\beta'_{2SLS} = \beta'_c$ when Z_i is unconditionally valid.

Proposition 4(iii) parallels the identification of β_c under correct interacted working model (1) by Theorem 2, and ensures $\beta'_{2SLS} = \beta'_c$ when the working model (13) underlying the partially interacted 2SLS is correct.

We relegate possible relaxations of these three sufficient conditions to Lemma S9 in the Supplementary Material.

8.2 Interacted OLS as a special case

Ordinary least squares (OLS) is a special case of 2SLS with $Z_i = D_i$. Therefore, our theory extends to the OLS regression of Y_i on (D_iX_i, X_i) when D_i is conditionally randomized. Interestingly, the discussion of interacted OLS in observational studies is largely missing except Rosenbaum and Rubin (1983). See Aronow and Samii (2016) and Chattopadhyay and Zubizarreta (2023) for the corresponding discussion of additive OLS.

Renew $X_i^0 = (1, X_{i1} - \bar{X}_1, \dots, X_{i,K-1} - \bar{X}_{K-1})^{\top}$, where $\bar{X}_k = N^{-1} \sum_{i=1}^N X_{ik}$, as a variant of X_i centered by covariate means over all units. Let $\hat{\tau}_{\text{OLS}}$ be the coefficient of D_i in $\ln(Y_i \sim D_i X_i^0 + X_i^0)$, as the OLS analog of $\hat{\tau}_{\times\times}$ when $Z_i = D_i$. Let $\hat{\beta}_{\text{OLS}}$ be the coefficient vector of $D_i X_i$ in $\ln(Y_i \sim D_i X_i + X_i)$, as the analog of $\hat{\beta}_{2\text{SLS}}$ when $Z_i = D_i$. We derive below the causal interpretation of $\hat{\tau}_{\text{OLS}}$ and $\hat{\beta}_{\text{OLS}}$.

First, letting $Z_i = D_i$ ensures that all units are compliers. The definitions of τ_c and β_c in Eqs. (2) and (4) simplify to $\tau = \mathbb{E}(\tau_i)$, as the average treatment effect, and $\beta = \operatorname{argmin}_{b \in \mathbb{R}^K} \mathbb{E}\{(\tau_i - X_i^{\top}b)^2\}$, as the coefficient vector in the linear projection of τ_i on X_i , respectively. The IV assumptions in Assumption 1 simplify to the selection on observables and overlap assumptions on D_i , summarized in Assumption 4 below.

Assumption 4. (i) Selection on observables: $D_i \perp \{Y_i(d) : d = 0, 1\} \mid X_i;$

(ii) **Overlap:** $0 < \mathbb{P}(D_i = 1 | X_i) < 1.$

Proposition 5 below follows from applying Theorems 1–2 with $Z_i = D_i$, and establishes the properties of $\hat{\tau}_{\text{OLS}}$ and $\hat{\beta}_{\text{OLS}}$ for estimating τ and β .

Proposition 5. Let τ_{OLS} and β_{OLS} denote the probability limits of $\hat{\tau}_{\text{OLS}}$ and $\hat{\beta}_{\text{OLS}}$. Assume Assumption 4. Then $\tau_{\text{OLS}} = \tau$ and $\beta_{\text{OLS}} = \beta$ if either of the following conditions holds:

(i) Linear treatment-covariate interactions: $\mathbb{E}(D_i X_i \mid X_i)$ is linear in X_i .

(ii) **Linear outcome:** $Y_i = D_i X_i^\top \beta_{ZX} + X_i^\top \beta_X + \eta_i$, where $\mathbb{E}(\eta_i \mid D_i, X_i) = 0$, for some fixed $\beta_{ZX}, \beta_X \in \mathbb{R}^K$.

Parallel to Proposition 1(i), Proposition 5(i) encompasses (a) categorical covariates and (b) randomly assigned treatment as special cases. Proposition 5(ii) implies that the interacted regression is correctly specified.

Similar results can be obtained for partially interacted OLS regression of Y_i on (D_iV_i, X_i) by letting $Z_i = D_i$ in Proposition 4. We omit the details.

9 Discussion

We formalized the interacted 2SLS as an intuitive approach for incorporate treatment effect heterogeneity in IV analyses, and established its properties for estimating the LATE and systematic treatment effect variation under the LATE framework. We showed that the coefficients from the interacted 2SLS are consistent for estimating the LATE and treatment effect heterogeneity with regard to covariates among compliers if either (i) the IV-covariate interactions are linear in the covariates or (ii) the linear outcome model underlying the interacted 2SLS is correct. Condition (i) includes categorical covariates and randomly assigned IV as special cases. When neither condition (i) nor condition (ii) holds, we propose a stratification strategy based on the IV propensity score to approximate the LATE and treatment effect heterogeneity with regard to the IV propensity score. We leave possible relaxations of the identifying conditions for the LATE to future work.

References

- Abadie, A. (2003). Semiparametric instrumental variable estimation of treatment response models. Journal of Econometrics 113, 231–263.
- Angrist, J. D. (1998). Estimating the labor market impact of voluntary military service using social security data on military applicants. *Econometrica* 66, 249–288.
- Angrist, J. D. and G. W. Imbens (1995). Two-stage least squares estimation of average causal effects in models with variable treatment intensity. *Journal of the American statistical Association 90*, 431–442.
- Angrist, J. D., G. W. Imbens, and D. B. Rubin (1996). Identification of causal effects using instrumental variables (with discussion). Journal of the American Statistical Association 91, 444–455.
- Angrist, J. D. and J.-S. Pischke (2009). Mostly Harmless Econometrics: An Empiricist's Companion. Princeton: Princeton University Press.
- Aronow, P. M. and C. Samii (2016). Does regression produce representative estimates of causal effects? *American Journal of Political Science* 60, 250–267.
- Blandhol, C., J. Bonney, M. Mogstad, and A. Torgovitsky (2022). When is tsls actually late? Technical report, National Bureau of Economic Research Cambridge, MA.
- Chattopadhyay, A., N. Greifer, and J. R. Zubizarreta (2024). lmw: Linear model weights for causal inference. *Observational Studies* 10, 33–62.
- Chattopadhyay, A. and J. R. Zubizarreta (2023). On the implied weights of linear regression for causal inference. *Biometrika* 110, 615–629.

- Cheng, J. and W. Lin (2018). Understanding causal distributional and subgroup effects with the instrumental propensity score. *American Journal of Epidemiology* 187, 614–622.
- Ding, P., A. Feller, and L. Miratrix (2019). Decomposing treatment effect variation. *Journal* of the American Statistical Association 114, 304–317.
- Djebbari, H. and J. Smith (2008). Heterogeneous impacts in progresa. Journal of Econometrics 145, 64–80.
- Hainmueller, J., J. Mummolo, and Y. Xu (2019). How much should we trust estimates from multiplicative interaction models? Simple tools to improve empirical practice. *Political Analysis 27*, 163–192.
- Heckman, J. J., J. Smith, and N. Clements (1997). Making the most out of programme evaluations and social experiments: Accounting for heterogeneity in programme impacts. *The Review of Economic Studies* 64, 487–535.
- Hirano, K. and G. W. Imbens (2001). Estimation of causal effects using propensity score weighting: An application to data on right heart catheterization. *Health Services and Outcomes Research Methodology 2*, 259–278.
- Imbens, G. W. and J. D. Angrist (1994). Identification and estimation of local average treatment effects. *Econometrica* 62, 467–475.
- Imbens, G. W. and D. B. Rubin (1997). Estimating outcome distributions for compliers in instrumental variables models. The Review of Economic Studies 64, 555–574.
- Kline, P. (2011). Oaxaca-Blinder as a reweighting estimator. American Economic Review 101, 532–537.

- Liero, H. (1989). Strong uniform consistency of nonparametric regression function estimates. Probability Theory and Related Fields 82, 587–614.
- Lin, W. (2013). Agnostic notes on regression adjustments to experimental data: Reexamining Freedman's critique. *The Annals of Applied Statistics* 7, 295–318.
- Pashley, N. E., L. Keele, and L. W. Miratrix (2024, 08). Improving instrumental variable estimators with poststratification. *Journal of the Royal Statistical Society Series A: Statistics in Society*, qnae073.
- Resnjanskij, S., J. Ruhose, S. Wiederhold, L. Woessmann, and K. Wedel (2024). Can mentoring alleviate family disadvantage in adolescence? a field experiment to improve labor market prospects. *Journal of Political Economy 132*, 000–000.
- Rosenbaum, P. R. and D. B. Rubin (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika* 70, 41–55.
- Sloczynski, T. (2022). When should we (not) interpret linear IV estimands as LATE?
- Słoczyński, T., S. D. Uysal, and J. M. Wooldridge (2025). Abadie's kappa and weighting estimators of the local average treatment effect. Journal of Business & Economic Statistics 43, 164–177.
- Strang, G. (2006). Linear Algebra and Its Applications (4 ed.). Cengage Learning.
- Wasserman, L. (2006). All of Nonparametric Statistics. New York: Springer.

Supplementary Material

Section S1 gives additional results that complement the main paper.

Section S2 gives the lemmas for the proofs. In particular, Section S2.4 gives two linear algebra results that underlie Proposition 1(ii).

Section S3 gives the proofs of the results in the main paper.

Section S4 gives the proofs of the results in Section S1.

Section S5 gives the proofs of the linear algebra results in Section S2.4.

Notation. Recall that for two random vectors $Y \in \mathbb{R}^p$ and $X \in \mathbb{R}^q$, we use $\operatorname{proj}(Y \mid X)$ to the linear projection of Y on X. Further let $\operatorname{res}(Y \mid X) = Y - \operatorname{proj}(Y \mid X)$ denote the corresponding residual.

Write

$$\widehat{DX}_i = \operatorname{proj}(D_i X_i \mid Z_i X_i, X_i) = C_1 Z_i X_i + C_0 X_i, \quad \operatorname{proj}(Z_i X_i \mid X_i) = C_2 X_i, \quad (S1)$$

where C_1 , C_0 , and C_2 denote the coefficient matrices. A useful fact for the proofs is

$$\widetilde{DX}_{i} = \operatorname{res}(\widehat{DX}_{i} \mid X_{i}) \stackrel{(\mathrm{S1})}{=} \operatorname{res}(C_{1}Z_{i}X_{i} + C_{0}X_{i} \mid X_{i})$$
$$= C_{1} \cdot \operatorname{res}(Z_{i}X_{i} \mid X_{i})$$
$$\stackrel{(\mathrm{S1})}{=} C_{1}(Z_{i}X_{i} - C_{2}X_{i}).$$
(S2)

Recall the definition of ϵ_i and Δ_i from Theorem 2. Extend the definition of ϵ_i to all units as

$$\epsilon_i = \begin{cases} \tau_i - X_i^\top \beta_c & \text{for compliers} \\ 0 & \text{for always- and never-takers} \end{cases}$$

A useful fact for the proofs is

$$Y_{i} - D_{i}X_{i}^{\top}\beta_{c} = \begin{cases} Y_{i}(1) - X_{i}^{\top}\beta_{c} & \text{for always-takers} \\ Y_{i}(0) + Z_{i}(\tau_{i} - X_{i}^{\top}\beta_{c}) & \text{for compliers} \\ Y_{i}(0) & \text{for never-takers} \end{cases} = \Delta_{i} + Z_{i}\epsilon_{i}.$$
(S3)

•

S1 Additional results

Population 2SLS estimand. Linear projection gives the population analog of leastsquares regression. Proposition S1 below formalizes the population analog of the interacted 2SLS in Definition 2.

Proposition S1. Define the *population interacted* 2SLS as the combination of the following two linear projections:

- (i) First stage: the linear projection of $D_i X_i$ on $(Z_i X_i^{\top}, X_i^{\top})^{\top}$, denoted by $\operatorname{proj}(D_i X_i \mid Z_i X_i, X_i)$.
- (ii) Second stage: the linear projection of Y_i on the concatenation of $\operatorname{proj}(D_i X_i \mid Z_i X_i, X_i)$ and X_i .
- Then β_{2SLS} equals the coefficient vector of $\operatorname{proj}(D_iX_i \mid Z_iX_i, X_i)$ from the second stage.

Invariance of the interacted 2SLS to nondegenerate linear transformations. The values of $\hat{\beta}_{2\text{SLS}}$ from the interacted 2SLS and β_c from Eq. (4) both depend on the choice of the covariate vector X_i . Let $\hat{\beta}_{2\text{SLS}}(X_i)$, $\beta_{2\text{SLS}}(X_i)$, and $\beta_c(X_i)$ denote the values of $\hat{\beta}_{2\text{SLS}}$, $\beta_{2\text{SLS}}$, and β_c corresponding to covariate vector X_i . Proposition S2 below states their invariance to nondegenerate linear transformation of X_i .

Proposition S2. For $K \times 1$ covariate vectors $\{X_i \in \mathbb{R}^K : i = 1, ..., N\}$ and an invertible $K \times K$ matrix Γ , we have

$$\hat{\beta}_{2\text{SLS}}(\Gamma X_i) = (\Gamma^{\top})^{-1} \hat{\beta}_{2\text{SLS}}(X_i), \quad \beta_{2\text{SLS}}(\Gamma X_i) = (\Gamma^{\top})^{-1} \beta_{2\text{SLS}}(X_i), \quad \beta_{\text{c}}(\Gamma X_i) = (\Gamma^{\top})^{-1} \beta_{\text{c}}(X_i).$$

Generalized additive 2SLS. Definition S1 below generalizes the additive and interactedadditive 2SLS in Definitions 1 and 4 to arbitrary first stage.

Definition S1 (Generalized additive 2SLS). Let $R_i = R(X_i, Z_i)$ be an arbitrary vector function of (X_i, Z_i) .

- (i) First stage: Fit $lm(D_i \sim R_i)$ over i = 1, ..., N. Denote the fitted values by \hat{D}_i for i = 1, ..., N.
- (ii) Second stage: Fit $lm(Y_i \sim \hat{D}_i + X_i)$ over i = 1, ..., N. Denote the coefficient of \hat{D}_i by $\hat{\tau}_{*+}$.

We use the subscript *+ to represent the arbitrary first stage and additive second stage. The additive 2SLS in Definition 1 is a special case of the generalized additive 2SLS with $R_i = (Z_i, X_i^{\top})^{\top}$. The interacted-additive 2SLS in Definition 4 is a special case with $R_i = (Z_i X_i^{\top}, X_i^{\top})^{\top}$.

Let τ_{*+} denote the probability limit of $\hat{\tau}_{*+}$. Proposition S3 below clarifies the causal interpretation of τ_{*+} .

Let $\check{D}_i = \operatorname{proj}(D_i \mid R_i)$ denote the population fitted value from the first stage of the generalized additive 2SLS. Let $\tilde{D}_i = \operatorname{res}(\check{D}_i \mid X_i)$. Let $\Delta_i = Y_i(0)$ for compliers and never-takers and $\Delta_i = Y_i(1) - \tau_c(X_i)$ for always-takers. Recall

$$w(X_i) = \frac{\operatorname{var}\{\mathbb{E}(D_i \mid Z_i, X_i) \mid X_i\}}{\mathbb{E}\left[\operatorname{var}\{\mathbb{E}(D_i \mid Z_i, X_i) \mid X_i\}\right]}.$$

Proposition S3. Assume Assumption 1. Then $\tau_{*+} = \mathbb{E} \{ \alpha(X_i) \cdot \tau_c(X_i) \} + \{\mathbb{E}(\tilde{D}_i^2)\}^{-1}B$, where

$$\alpha(X_i) = \frac{\mathbb{E}(D_i \check{D}_i \mid X_i)}{\mathbb{E}(\tilde{D}_i^2)}, \quad B = \mathbb{E}\left[\left\{\mathbb{E}(\check{D}_i \mid X_i) - \operatorname{proj}(\check{D}_i \mid X_i)\right\} \cdot \left\{\mathbb{E}(\Delta_i \mid X_i) - \operatorname{proj}(\Delta_i \mid X_i)\right\}\right]$$

In particular,

- (i) B = 0 if either of the following conditions holds:
 - (a) $\mathbb{E}(\check{D}_i \mid X_i)$ is linear in X_i .
 - (b) $\mathbb{E}(\Delta \mid X_i)$ is linear in X_i .
- (ii) $\alpha(X_i) = w(X_i)$ if both of the following conditions hold:
 - (a) $\mathbb{E}(\check{D}_i \mid X_i)$ is linear in X_i .
 - (b) $\check{D}_i = \mathbb{E}(D_i \mid Z_i, X_i)$; i.e., the arbitrary first stage is correct.

S2 Lemmas

S2.1 Projection and conditional expectation

Lemmas S1–S3 below review some standard results regarding projection and conditional expectation. We omit the proofs.

Lemma S1. For random vectors A and B,

- (i) $\mathbb{E}(A \mid B) = \operatorname{proj}(A \mid B)$ if and only if $\mathbb{E}(A \mid B)$ is linear in B;
- (ii) if $A \perp B$, then $\mathbb{E}(AB \mid B) = \operatorname{proj}(AB \mid B) = \mathbb{E}(A)B$;
- (iii) if A is binary, then $\mathbb{E}(A \mid B)$ is constant implies $A \perp B$.

Lemma S2. Let X be an $m \times 1$ random vector with $\mathbb{E}(XX^{\top})$ being invertible. Then

- (i) for any subvector of X, denoted by $V, \mathbb{E}(VV^{\top})$ is invertible.
- (ii) X takes at least m distinct values.
- (iii) If X takes exactly m distinct values, denoted by $x_1, \ldots, x_m \in \mathbb{R}^m$, then

- $\Gamma = (x_1, \ldots, x_m) \in \mathbb{R}^{m \times m}$ is invertible, and $X = \Gamma(1(X = x_1), \ldots, 1(X = x_m))^{\top}$.

- XX^{\top} is component-wise linear in X.
- for any random vector A, $\mathbb{E}(A \mid X)$ is linear in X.

Lemma S3 (Population Frisch–Waugh–Lovell (FWL)). For a random variable Y, a $p_1 \times 1$ random vector X_1 , and a $p_2 \times 1$ random vector X_2 , let $\tilde{Y} = \operatorname{res}(Y \mid X_2)$ and $\tilde{X}_1 = \operatorname{res}(X_1 \mid X_2)$ denote the residuals from the linear projections of Y and X_1 on X_2 , respectively. Then the coefficient vector of X_1 in the linear projection of Y on $(X_1^{\top}, X_2^{\top})^{\top}$ equals the coefficient vector of \tilde{X}_1 in the linear projection of Y on \tilde{X}_1 .

S2.2 Lemmas for the interacted 2SLS

Assumption S1 below states the rank condition for the interacted 2SLS, which we assume implicitly throughout the main paper. By the law of large numbers,

$$\beta_{2\text{SLS}} = \text{the first } K \text{ dimensions of } \left[\mathbb{E} \left\{ \begin{pmatrix} Z_i X_i \\ X_i \end{pmatrix} \begin{pmatrix} D_i X_i^\top, X_i^\top \end{pmatrix} \right\} \right]^{-1} \mathbb{E} \left\{ \begin{pmatrix} Z_i X_i \\ X_i \end{pmatrix} Y_i \right\}.$$

Assumption S1.
$$\mathbb{E}\left\{ \begin{pmatrix} Z_i X_i \\ X_i \end{pmatrix} \begin{pmatrix} D_i X_i^\top, X_i^\top \end{pmatrix} \right\}$$
 is invertible.

Lemma S4 below states some implications of Assumption S1 that are useful for the proofs.

Lemma S4. Recall C_0, C_1, C_2 from (S1). Assume Assumption S1. We have

(i)
$$\mathbb{E}\left\{ \begin{pmatrix} Z_i X_i \\ X_i \end{pmatrix} \begin{pmatrix} Z_i X_i^{\top}, X_i^{\top} \end{pmatrix} \right\}$$
 and $\mathbb{E}(X_i X_i^{\top})$ are both invertible with

$$(C_1, C_0) = \mathbb{E}\left\{ D_i X_i (Z_i X_i^{\top}, X_i^{\top}) \right\} \left[\mathbb{E}\left\{ \begin{pmatrix} Z_i X_i \\ X_i \end{pmatrix} (Z_i X_i^{\top}, X_i^{\top}) \right\} \right]^{-1},$$

$$C_2 = \mathbb{E}(Z_i X_i X_i^{\top}) \left\{ \mathbb{E}(X_i X_i^{\top}) \right\}^{-1}.$$

- (ii) X_i takes at least K distinct values.
- (iii) C_1 , C_2 , and $I_K C_2$ are all invertible.

Proof of Lemma S4. We verify below Lemma S4(i)–(iii), respectively. We omit the subscript i in the proof. Let

$$\Gamma_1 = \mathbb{E}\left\{ \begin{pmatrix} DX \\ X \end{pmatrix} (ZX^{\top}, X^{\top}) \right\}, \quad \Gamma_2 = \mathbb{E}\left\{ \begin{pmatrix} ZX \\ X \end{pmatrix} (ZX^{\top}, X^{\top}) \right\}$$

be shorthand notation with Γ_1 being invertible under Assumption S1.

Proof of Lemma S4(i): We prove the result by contradiction. Note that Γ_2 is positive semidefinite. If Γ_2 is degenerate, then there exists a nonzero vector a such that $a^{\top}\Gamma_2 a = 0$. This implies

$$0 = a^{\top} \mathbb{E} \left\{ \begin{pmatrix} ZX \\ X \end{pmatrix} (ZX^{\top}, X^{\top}) \right\} a = \mathbb{E} \left\{ a^{\top} \begin{pmatrix} ZX \\ X \end{pmatrix} (ZX^{\top}, X^{\top}) a \right\}$$

such that $(ZX^{\top}, X^{\top})a = 0$. As a result, we have

$$\Gamma_1 a = \mathbb{E}\left\{ \begin{pmatrix} DX \\ X \end{pmatrix} (ZX^{\top}, X^{\top}) \right\} a = \mathbb{E}\left\{ \begin{pmatrix} DX \\ X \end{pmatrix} (ZX^{\top}, X^{\top})a \right\} = 0$$

for a nonzero a. This contradicts with Γ_1 being invertible.

That $\mathbb{E}(XX^{\top})$ is invertible then follows from Lemma S2(i).

Proof of Lemma S4(ii). Given $\mathbb{E}(XX^{\top})$ is invertible as we just proved, the result follows from Lemma S2(ii).

Proof of Lemma S4(iii): Properties of linear projection ensure

$$\operatorname{proj}\left\{ \begin{pmatrix} DX\\ X \end{pmatrix} \mid ZX, X \right\} = \begin{pmatrix} C_1 & C_0\\ 0 & I_K \end{pmatrix} \begin{pmatrix} ZX\\ X \end{pmatrix},$$

with

$$\begin{pmatrix} C_1 & C_0 \\ 0 & I_K \end{pmatrix} = \mathbb{E}\left\{ \begin{pmatrix} DX \\ X \end{pmatrix} (ZX^{\top}, X^{\top}) \right\} \left[\mathbb{E}\left\{ \begin{pmatrix} ZX \\ X \end{pmatrix} (ZX^{\top}, X^{\top}) \right\} \right]^{-1} = \Gamma_1 \Gamma_2^{-1}.$$
(S4)

This ensures $\det(C_1) = \det\begin{pmatrix} C_1 & C_0 \\ 0 & I_K \end{pmatrix} = \det(\Gamma_1 \Gamma_2^{-1}) \neq 0.$

In addition, note that

$$I_{K} - C_{2} = \mathbb{E}(XX^{\top}) \cdot \left\{ \mathbb{E}(XX^{\top}) \right\}^{-1} - \mathbb{E}(ZXX^{\top}) \cdot \left\{ \mathbb{E}(XX^{\top}) \right\}^{-1}$$
$$= \mathbb{E}\left\{ (1 - Z)XX^{\top} \right\} \cdot \left\{ \mathbb{E}(XX^{\top}) \right\}^{-1}.$$

To show that C_2 and $I_K - C_2$ are invertible, it suffices to show that $\mathbb{E}(ZXX^{\top})$ and $\mathbb{E}\{(1 - Z)XX^{\top}\}$ are invertible. This is ensured by

$$0 < \det(\Gamma_2) = \det \left\{ \begin{pmatrix} \mathbb{E}(ZXX^{\top}) & \mathbb{E}(ZXX^{\top}) \\ \mathbb{E}(ZXX^{\top}) & \mathbb{E}(XX^{\top}) \end{pmatrix} \right\}$$
$$= \det \left\{ \begin{pmatrix} \mathbb{E}(ZXX^{\top}) & \mathbb{E}(ZXX^{\top}) \\ 0 & \mathbb{E}\{(1-Z)XX^{\top}\} \end{pmatrix} \right\}$$
$$= \det \left\{ \mathbb{E}(ZXX^{\top}) \right\} \cdot \det \left[\mathbb{E}\{(1-Z)XX^{\top}\} \right].$$

Lemma S5 below formalizes the sufficiency and necessity of Assumption 3 to ensure the combination of conditions (8) and (10); c.f. Section 4.2.

Lemma S5. Assume Assumption 1. Assumption 3 is both sufficient and necessary for

$$\mathbb{E}(\Delta_i \mid X_i) = X_i^{\top} \beta_{\Delta} \text{ for some } \beta_{\Delta} \in \mathbb{R}^K, \quad \tau_{c}(X_i) = \mathbb{E}(\tau_i \mid X_i, U_i = c) = X_i^{\top} \beta_{c}.$$
(S5)

Proof of Lemma S5. We verify below the sufficiency and necessity of Assumption 3,

respectively.

Sufficiency. Under Assumption 3, we have

$$\eta_{i} = Y_{i} - D_{i}X_{i}^{\top}\beta_{DX} - X_{i}^{\top}\beta_{X}$$

$$= \begin{cases} Y_{i}(1) - X_{i}^{\top}(\beta_{DX} + \beta_{X}) & \text{if } U_{i} = a \\ Y_{i}(0) - X_{i}^{\top}\beta_{X} & \text{if } U_{i} = n \\ Y_{i}(0) - X_{i}^{\top}\beta_{X} + Z_{i}(\tau_{i} - X_{i}^{\top}\beta_{DX}) & \text{if } U_{i} = c \end{cases}$$

$$= \lambda_{i} + Z_{i}\zeta_{i},$$
(S6)

where

$$\lambda_i = \begin{cases} Y_i(1) - X_i^\top (\beta_{DX} + \beta_X) & \text{if } U_i = \mathbf{a} \\ Y_i(0) - X_i^\top \beta_X & \text{if } U_i = \mathbf{n} \\ Y_i(0) - X_i^\top \beta_X & \text{if } U_i = \mathbf{c} \end{cases}, \quad \zeta_i = \begin{cases} \tau_i - X_i^\top \beta_{DX} & \text{if } U_i = \mathbf{c} \\ 0 & \text{if } U_i \neq \mathbf{c} \end{cases}.$$

The definition of ζ_i ensures

$$\mathbb{E}(\zeta_i \mid X_i) = \mathbb{E}(\zeta_i \mid X_i, U_i = c) \cdot \mathbb{P}(U_i = c \mid X_i)
= \left\{ \mathbb{E}(\tau_i \mid X_i, U_i = c) - X_i^\top \beta_{DX} \right\} \cdot \mathbb{P}(U_i = c \mid X_i)
= \left\{ \tau_c(X_i) - X_i^\top \beta_{DX} \right\} \cdot \mathbb{P}(U_i = c \mid X_i).$$
(S7)

This, together with (S6), ensures

$$\mathbb{E}(\eta_i \mid X_i, Z_i) \stackrel{(S6)}{=} \mathbb{E}(\lambda_i \mid X_i, Z_i) + Z_i \cdot \mathbb{E}(\zeta_i \mid X_i, Z_i) \\
\stackrel{\text{Assm. 1}}{=} \mathbb{E}(\lambda_i \mid X_i) + Z_i \cdot \mathbb{E}(\zeta_i \mid X_i) \\
\stackrel{(S7)}{=} \mathbb{E}(\lambda_i \mid X_i) + Z_i \cdot \left\{ \tau_c(X_i) - X_i^\top \beta_{DX} \right\} \cdot \mathbb{P}(U_i = c \mid X_i). \quad (S8)$$

Under Assumption 3, we have $\mathbb{E}(\eta_i \mid X_i, Z_i) = 0$ so that (S8) implies

$$\mathbb{E}(\lambda_i \mid X_i) \stackrel{\text{(S8)}}{=} \mathbb{E}(\eta_i \mid X_i, Z_i = 0) \stackrel{\text{Assm. 3}}{=} 0, \tag{S9}$$

$$\tau_{\mathbf{c}}(X_i) - X_i^{\top} \beta_{DX} \stackrel{(\mathrm{S8})}{=} \frac{\mathbb{E}(\eta_i \mid X_i, Z_i = 1) - \mathbb{E}(\eta_i \mid X_i, Z_i = 0)}{\mathbb{P}(U_i = \mathbf{c} \mid X_i)} \stackrel{\mathrm{Assm. 3}}{=} 0.$$
(S10)

(S10), together with Lemma S1(i), ensures

$$\tau_{\rm c}(X_i) = X_i^{\top} \beta_{DX} \quad \text{with} \quad \beta_{\rm c} = \beta_{DX}. \tag{S11}$$

Plugging (S11) in the definition of λ_i ensures $\Delta_i = \lambda_i + X_i^{\top} \beta_X$ with

$$\mathbb{E}(\Delta_i \mid X_i) = \mathbb{E}(\lambda_i \mid X_i) + X_i^{\top} \beta_X \stackrel{(S9)}{=} X_i^{\top} \beta_X.$$

This, together with (S10), verifies the sufficiency of Assumption 3 to ensure (S5).

<u>Necessity.</u> We show below that (S5) implies Assumption 3. Recall from (S3) that $Y_i - D_i X_i^\top \beta_c = \Delta_i + Z_i \epsilon_i$. When (S5) holds, we have $\mathbb{E}(\epsilon_i \mid X_i, U_i = c) = 0$ so that

$$\mathbb{E}(\epsilon_i \mid X_i) = \mathbb{E}(\epsilon_i \mid X_i, U_i = c) \cdot \mathbb{P}(U_i = c \mid X_i) = 0.$$
(S12)

This ensures

$$\mathbb{E}(Y_i - D_i X_i^\top \beta_c \mid X_i, Z_i) \stackrel{(S3)}{=} \mathbb{E}(\Delta_i \mid X_i, Z_i) + Z_i \cdot \mathbb{E}(\epsilon_i \mid X_i, Z_i) \\
\stackrel{\text{Assm. 1}}{=} \mathbb{E}(\Delta_i \mid X_i) + Z_i \cdot \mathbb{E}(\epsilon_i \mid X_i) \\
\stackrel{(S5)+(S12)}{=} X_i^\top \beta_{\Lambda}.$$
(S13)

Let $\eta'_i = Y_i - D_i X_i^\top \beta_c - X_i^\top \beta_\Delta$. We have

$$Y_{i} = D_{i}X_{i}^{\top}\beta_{c} + X_{i}^{\top}\beta_{\Delta} + \eta_{i}',$$
$$\mathbb{E}(\eta_{i}' \mid X_{i}, Z_{i}) = \mathbb{E}(Y_{i} - D_{i}X_{i}^{\top}\beta_{c} \mid X_{i}, Z_{i}) - X_{i}^{\top}\beta_{\Delta} \stackrel{(S13)}{=} 0$$

so that Assumption 3 holds.

Lemma S6 below states a numeric result that we use to prove Corollary 1.

Lemma S6. Let \widehat{DX}_i and \hat{D}_i denote the fitted values from $lm(D_iX_i \sim Z_iX_i + X_i)$ and

$$lm(D_i \sim Z_i X_i + X_i), \tag{S14}$$

respectively. Assume that X_i is categorical with K levels. Then $\widehat{DX}_i = \hat{D}_i X_i$.

Proof of Lemma S6. We verify below the result for $X_i = (1(X_i^* = 1), \dots, 1(X_i^* = K))^{\top}$, where $X_i^* \in \{1, \dots, K\}$ denotes a K-level categorical variable. The result for general categorical X_i then follows from the invariance of least squares to nondegenerate linear transformations of the regressor vector.

Given $X_i = (1(X_i^* = 1), \dots, 1(X_i^* = K))^{\top}$, for $k = 1, \dots, K$, the *k*th element of $\hat{D}_i X_i$ equals $\hat{D}_i \cdot 1(X_i^* = k)$, and the *k*th element of \widehat{DX}_i equals the fitted value from

$$lm\Big\{D_i \cdot 1(X_i^* = k) \sim Z_i X_i + X_i\Big\},\tag{S15}$$

denoted by $\widehat{DX}_i[k]$. By symmetry, to show that $\widehat{DX}_i = \hat{D}_i X_i$, it suffices to show that the first element of $\hat{D}_i X_i$ equals the first element of \widehat{DX}_i , i.e.,

$$\hat{D}_i \cdot 1(X_i^* = 1) = \widehat{DX}_i[1].$$
 (S16)

We verify (S16) below.

Left-hand side of (S16). Standard results ensure that (S14) is equivalent to K stratumwise regressions

$$lm(D_i \sim Z_i + 1)$$
 over units with $X_i^* = k$

for k = 1, ..., K, such that \hat{D}_i equals the sample mean of $\{D_i : X_i^* = k, Z_i = z\}$ for units with $\{X_i^* = k, Z_i = z\}$. This ensures

$$\hat{D}_i \cdot 1(X_i^* = 1) = \begin{cases} 0 & \text{for } \{i : X_i^* \neq 1\}; \\ \text{sample mean of } \{D_i : X_i^* = 1, Z_i = z\} & \text{for } \{i : X_i^* = 1; Z_i = z\}. \end{cases}$$
(S17)

Right-hand side of (S16). From (S15), $\widehat{DX}_i[1]$ equals the fitted value from

$$\mathrm{lm}\Big\{D_i\cdot 1(X_i^*=1)\sim Z_iX_i+X_i\Big\},$$

which is equivalent to K stratum-wise regressions

$${\rm lm}\Big\{D_i\cdot 1(X_i^*=1)\sim Z_i+1\Big\} \quad {\rm over \ units \ with } \ X_i^*=k$$

for k = 1, ..., K. This ensures that $\widehat{DX}_i[1]$ equals the sample mean of $\{D_i \cdot 1(X_i^* = 1) : X_i^* = k, Z_i = z\}$ for units with $\{X_i^* = k, Z_i = z\}$, which simplies to the follows:

$$\widehat{DX}_{i}[1] = \begin{cases} 0 & \text{for } \{i : X_{i}^{*} \neq 1\}; \\ \text{sample mean of } \{D_{i} : X_{i}^{*} = 1, Z_{i} = z\} & \text{for } \{i : X_{i}^{*} = 1; Z_{i} = z\}. \end{cases}$$

This is identical to the expression of $\hat{D}_i \cdot 1(X_i^* = 1)$ in (S17), and ensures (S16).

S2.3 Lemmas for the partially interacted 2SLS

Assumption S2 below states the rank condition for the partially interacted 2SLS.

Assumption S2.
$$\mathbb{E}\left\{ \begin{pmatrix} Z_i V_i \\ X_i \end{pmatrix} \begin{pmatrix} D_i V_i^{\top}, X_i^{\top} \end{pmatrix} \right\}$$
 is invertible.

Lemma S7 below generalizes Lemma S4 and states some implications of Assumption S2 useful for the proofs. The proof is similar to that of Lemma S4, hence omitted.

Lemma S7. Assume Assumption S2. We have

(i)
$$\mathbb{E}\left\{ \begin{pmatrix} Z_i V_i \\ X_i \end{pmatrix} \begin{pmatrix} Z_i V_i^{\top}, X_i^{\top} \end{pmatrix} \right\}$$
, $\mathbb{E}(X_i X_i^{\top})$, and $\mathbb{E}(V_i V_i^{\top})$ are all invertible with

$$\operatorname{proj}(D_i V_i \mid Z_i V_i, X_i) = C_1 Z_i V_i + C_0 X_i, \qquad \operatorname{proj}(Z_i V_i \mid V_i) = C_2 V_i$$

with

$$(C_1, C_0) = \mathbb{E}\left\{D_i V_i(Z_i V_i^{\top}, X_i^{\top})\right\} \left[\mathbb{E}\left\{\begin{pmatrix}Z_i V_i\\X_i\end{pmatrix}\left(Z_i V_i^{\top}, X_i^{\top}\right)\right\}\right]^{-1}, C_2 = \mathbb{E}(Z_i V_i V_i^{\top})\left\{\mathbb{E}(V_i V_i^{\top})\right\}^{-1}.$$

- (ii) X_i takes at least K distinct values. V_i takes at least Q distinct values.
- (iii) C_1 , C_2 , and $I_Q C_2$ are all invertible.

Lemma S8. Assume $X_i = (V_i^{\top}, W_i^{\top})^{\top} \in \mathbb{R}^K$, where $V_i \in \mathbb{R}^Q$ with $Q \leq K$ and $\mathbb{E}(X_i X_i^{\top})$ is invertible. Then $\mathbb{E}(Z_i V_i \mid X_i)$ is linear in X_i and $\mathbb{E}(Z_i \mid X_i) \cdot \{V_i - \operatorname{proj}(Z_i V_i \mid X_i)\}$ is linear in V_i if any of the following conditions holds:

- (i) V_i is categorical with Q levels; $\operatorname{proj}(Z_iV_i \mid X_i) = \operatorname{proj}(Z_iV_i \mid V_i)$ and $\mathbb{E}(Z_i \mid X_i) = \mathbb{E}(Z_i \mid V_i)$.
- (ii) $Z_i \perp \!\!\!\perp X_i$.

Proof of Lemma S8. To ensure $\mathbb{E}(Z_iV_i \mid X_i)$ is linear in X_i ,

- the sufficiency of Lemma S8(i) follows from $\mathbb{E}(Z_i V_i \mid X_i) = \mathbb{E}(Z_i V_i \mid V_i)$, where $\mathbb{E}(Z_i V_i \mid V_i)$ is linear in V_i by Lemma S2(iii).
- the sufficiency of Lemma S8(ii) follows from $\mathbb{E}(Z_i V_i \mid X_i) = \mathbb{E}(Z_i \mid X_i) \cdot V_i = \mathbb{E}(Z_i) \cdot V_i$.

We verify below the sufficiency of the conditions for ensuring that $\mathbb{E}(Z_i \mid X_i) \cdot \{V_i - \text{proj}(Z_iV_i \mid X_i)\}$ is linear in V_i . We omit the subscript *i* in the proof.

Proof of Condition (i). Given $\mathbb{E}(XX^{\top})$ is invertible, Lemma S2(i) ensures that $\mathbb{E}(VV^{\top})$ is invertible. When V is categorical with Q levels, Lemma S2(iii) ensures that VV^{\top} is component-wise linear in V and that $\mathbb{E}(Z \mid V)$ is linear in V. We can write

$$V - \operatorname{proj}(ZV \mid V) = CV, \quad \mathbb{E}(Z \mid V) = V^{\top}c, \tag{S18}$$

for $C = I_Q - \mathbb{E}(ZVV^{\top}) \{\mathbb{E}(VV^{\top})\}^{-1}$ and some constant $c \in \mathbb{R}^Q$. This, together with $\operatorname{proj}(ZV \mid X) = \operatorname{proj}(ZV \mid V)$ and $\mathbb{E}(Z \mid X) = \mathbb{E}(Z \mid V)$, ensures

$$\left\{V - \operatorname{proj}(ZV \mid X)\right\} \cdot \mathbb{E}(Z \mid X) = \left\{V - \operatorname{proj}(ZV \mid V)\right\} \cdot \mathbb{E}(Z \mid V) \stackrel{(S18)}{=} CV \cdot V^{\top}c,$$

which is linear in V when VV^{\top} is component-wise linear in V.

Proof of Condition (ii). When $Z \perp X$, we have

$$\operatorname{proj}(ZX \mid X) = \mathbb{E}(Z)X \tag{S19}$$

by Lemma S1(ii). Given $V = (I_Q, 0)X$, we have

$$\operatorname{proj}(ZV \mid X) = \operatorname{proj} \{ Z \cdot (I_Q, 0)X \mid X \}$$
$$= (I_Q, 0) \cdot \operatorname{proj}(ZX \mid X) \stackrel{(S19)}{=} (I_Q, 0) \cdot \mathbb{E}(Z)X = \mathbb{E}(Z)V$$

so that $\mathbb{E}(Z \mid X) \cdot \{V - \operatorname{proj}(ZV \mid X)\} = \mathbb{E}(Z) \cdot \{1 - \mathbb{E}(Z)\} \cdot V.$

Lemma S9 below underlies the sufficient conditions in Proposition 4. Renew

$$B_{1} = C_{1} \cdot \mathbb{E}\left[\left\{\mathbb{E}(Z_{i}V_{i} \mid X_{i}) - \operatorname{proj}(Z_{i}V_{i} \mid X_{i})\right\} \cdot \left\{\mathbb{E}(\Delta_{i} \mid X_{i}) - \operatorname{proj}(\Delta_{i} \mid X_{i})\right\}\right],$$
(S20)
$$B_{2} = C_{1} \cdot \mathbb{E}\left[\mathbb{E}(Z_{i} \mid X_{i}) \cdot \{V_{i} - \operatorname{proj}(Z_{i}V_{i} \mid X_{i})\} \cdot \epsilon_{i} \mid U_{i} = c\right] \cdot \mathbb{P}(U_{i} = c),$$

where C_1 is the coefficient matrix of $Z_i V_i$ in $\operatorname{proj}(D_i V_i \mid Z_i V_i, X_i)$. Renew

$$\epsilon_{i} = \begin{cases} \tau_{i} - V_{i}^{\top} \beta_{c}^{\prime} & \text{for compliers;} \\ 0 & \text{for always-takers and never-takers,} \end{cases}$$
$$\Delta_{i} = \begin{cases} Y_{i}(0) & \text{for compliers and never-takers;} \\ Y_{i}(1) - V_{i}^{\top} \beta_{c}^{\prime} & \text{for always-takers.} \end{cases}$$

The renewed ϵ_i , Δ_i , and C_1 reduce to their original definitions in Theorem 2 when $V_i = X_i$.

Lemma S9. Recall $B' = \mathbb{E}\{\widetilde{DV}_i \cdot (Y_i - D_i V_i^\top \beta_c')\}$ with $\widetilde{DV}_i = \operatorname{res}\{\operatorname{proj}(D_i V_i \mid Z_i X_i, X_i) \mid X_i\}$ in Proposition 4. Under Assumptions 1 and S2, we have $B' = B_1 + B_2$ with B_1 and B_2 satisfying the following:

- (i) $B_1 = B_2 = 0$ if either of Lemma S8(i)–(ii) holds.
- (ii) $B_1 = 0$ if $\mathbb{E}(\Delta_i \mid X_i)$ is linear in X_i .

$$B_2 = 0$$
 if $\tau_c(X_i) = \mathbb{E}(\tau_i \mid X_i, U_i = c)$ is linear in V_i .

Proof of Lemma S9. Observe that

$$Y_{i} - D_{i}V_{i}^{\top}\beta_{c}^{\prime} = \begin{cases} Y_{i}(1) - V_{i}^{\top}\beta_{c}^{\prime} & \text{for always-takers} \\ Y_{i}(0) + D_{i}\tau_{i} - D_{i}V_{i}^{\top}\beta_{c}^{\prime} & \text{for compliers} \\ Y_{i}(0) & \text{for never-takers} \end{cases} = \Delta_{i} + Z_{i}\epsilon_{i}.$$
(S21)

This ensures

$$B' = \mathbb{E}\left\{\widetilde{DV}_{i} \cdot (Y_{i} - D_{i}V_{i}^{\top}\beta_{c}')\right\}$$

$$\stackrel{(S21)}{=} \mathbb{E}\left\{\widetilde{DV}_{i} \cdot (\Delta_{i} + Z_{i}\epsilon_{i})\right\} = \mathbb{E}(\widetilde{DV}_{i} \cdot \Delta_{i}) + \mathbb{E}(\widetilde{DV}_{i} \cdot Z_{i}\epsilon_{i}).$$

Therefore, it suffices to show that

$$B_1 = \mathbb{E}(\widetilde{DV_i} \cdot \Delta_i), \quad B_2 = \mathbb{E}(\widetilde{DV_i} \cdot Z_i \epsilon_i)$$
(S22)

for B_1 and B_2 in Eq. (S20). To simplify the notation, write

$$\operatorname{proj}(D_i V_i \mid Z_i V_i, X_i) = C_1 Z_i V_i + C_0 X_i,$$

where C_1 and C_0 are well defined by Lemma S7 with C_1 being invertible under Assumption S2. We have

$$\widetilde{DV_i} = \operatorname{res} \left\{ \operatorname{proj}(D_i V_i \mid Z_i X_i, X_i) \mid X_i \right\}$$

$$= \operatorname{res}(C_1 Z_i V_i + C_0 X_i \mid X_i)$$

$$= C_1 \cdot \operatorname{res}(Z_i V_i \mid X_i)$$

$$= C_1 \left\{ Z_i V_i - \operatorname{proj}(Z_i V_i \mid X_i) \right\},$$

$$\widetilde{DV_i} \cdot Z_i = C_1 \left\{ Z_i V_i - Z_i \cdot \operatorname{proj}(Z_i V_i \mid X_i) \right\} = C_1 \cdot Z_i \cdot \left\{ V_i - \operatorname{proj}(Z_i V_i \mid X_i) \right\}.$$

This ensures

$$\mathbb{E}(\widetilde{DV_i} \mid X_i) = C_1 \Big\{ \mathbb{E}(Z_i V_i \mid X_i) - \operatorname{proj}(Z_i V_i \mid X_i) \Big\}, \\
\mathbb{E}(\widetilde{DV_i} \cdot Z_i \mid X_i) = C_1 \cdot \mathbb{E}(Z_i \mid X_i) \cdot \Big\{ V_i - \operatorname{proj}(Z_i V_i \mid X_i) \Big\}.$$
(S23)

Let $\xi_i = \Delta_i - \operatorname{proj}(\Delta_i \mid X_i)$ with $\mathbb{E}(\xi_i \mid X_i) = \mathbb{E}(\Delta_i \mid X_i) - \operatorname{proj}(\Delta_i \mid X_i)$. Properties of linear projection ensure $\mathbb{E}\{\widetilde{DV_i} \cdot \operatorname{proj}(\Delta_i \mid X_i)\} = 0$ such that

$$\mathbb{E}(\widetilde{DV_i} \cdot \Delta_i) = \mathbb{E}\left\{\widetilde{DV_i} \cdot \operatorname{proj}(\Delta_i \mid X_i)\right\} + \mathbb{E}(\widetilde{DV_i} \cdot \xi_i) = \mathbb{E}(\widetilde{DV_i} \cdot \xi_i).$$
(S24)

In addition, $\widetilde{DV_i}$ is a function of (X_i, Z_i) , whereas ξ_i and ϵ_i are both functions of $\{Y_i(z, d), D_i(z), X_i : z = 0, 1; d = 0, 1\}$. This ensures

$$\widetilde{DV}_i \perp \!\!\!\perp \xi_i \mid X_i, \qquad \widetilde{DV}_i \cdot Z_i \perp \!\!\!\perp \epsilon_i \mid X_i \tag{S25}$$

under Assumption 1.

Proof of $B_1 = \mathbb{E}(\widetilde{DV}_i \cdot \Delta_i)$ in (S22) and sufficient conditions of $B_1 = 0$. (S23)–(S25) ensure

$$\mathbb{E}(\widetilde{DV_i} \cdot \Delta_i) \stackrel{(S24)}{=} \mathbb{E}(\widetilde{DV_i} \cdot \xi_i)$$

$$= \mathbb{E}\left\{\mathbb{E}(\widetilde{DV}_{i} \cdot \xi_{i} \mid X_{i})\right\}$$

$$\stackrel{(S25)}{=} \mathbb{E}\left[\mathbb{E}(\widetilde{DV}_{i} \mid X_{i}) \cdot \mathbb{E}(\xi_{i} \mid X_{i})\right]$$

$$\stackrel{(S23)}{=} C_{1} \cdot \mathbb{E}\left[\left\{\mathbb{E}(Z_{i}V_{i} \mid X_{i}) - \operatorname{proj}(Z_{i}V_{i} \mid X_{i})\right\} \cdot \left\{\mathbb{E}(\Delta_{i} \mid X_{i}) - \operatorname{proj}(\Delta_{i} \mid X_{i})\right\}\right]$$

$$= B_{1}.$$

By Lemma S8, Lemma S9(i) ensures $\mathbb{E}(Z_i V_i \mid X_i)$ is linear in X_i so that $B_1 = 0$. By Lemma S1(i), Lemma S9(ii) ensures $\mathbb{E}(\Delta_i \mid X_i) = \operatorname{proj}(\Delta_i \mid X_i)$ so that $B_1 = 0$.

Proof of $B_2 = \mathbb{E}(\widetilde{DV_i} \cdot Z_i \epsilon_i)$ in (S22) and sufficient conditions of $B_2 = 0$. Let $g(X_i) = \mathbb{E}(Z_i \mid X_i) \cdot \{V_i - \operatorname{proj}(Z_i V_i \mid X_i)\}$ be a shorthand notation. (S23) and (S25) ensure

$$\begin{split} \mathbb{E}(\widetilde{DV_i} \cdot Z_i \epsilon_i) &\stackrel{(\text{S25})}{=} & \mathbb{E}\left\{ \mathbb{E}(\widetilde{DV_i} \cdot Z_i \mid X_i) \cdot \mathbb{E}(\epsilon_i \mid X_i) \right\} \\ &\stackrel{(\text{S23})}{=} & C_1 \cdot \mathbb{E}\left\{g(X_i) \cdot \mathbb{E}(\epsilon_i \mid X_i)\right\} \\ &= & C_1 \cdot \mathbb{E}\left[\mathbb{E}\left\{g(X_i) \cdot \epsilon_i \mid X_i\right\}\right] \\ &= & C_1 \cdot \mathbb{E}\left\{g(X_i) \cdot \epsilon_i\right\} \\ &= & C_1 \cdot \mathbb{E}\left\{g(X_i) \cdot \epsilon_i \mid U_i = c\right\} \cdot \mathbb{P}(U_i = c) = B_2. \end{split}$$

Given C_1 is invertible and $\mathbb{P}(U_i = c) > 0$, we have $B_2 = 0$ if and only if

$$\mathbb{E}\left\{g(X_i)\cdot\epsilon_i\mid U_i=\mathbf{c}\right\}=0.$$
(S26)

The definition of ϵ_i ensures that (S26) holds when $g(X_i)$ is linear in V_i . By Lemma S8, Lemma S9(i) ensures that $g(X_i)$ is linear in V_i .

In addition, we have

$$\mathbb{E}\{g(X_i) \cdot \epsilon_i \mid U_i = c\} = \mathbb{E}\left[\mathbb{E}\{g(X_i) \cdot \epsilon_i \mid X_i, U_i = c\} \mid U_i = c\right]$$
$$= \mathbb{E}\left\{g(X_i) \cdot \mathbb{E}\left(\epsilon_i \mid X_i, U_i = c\right) \mid U_i = c\right\}$$

so that (S26) holds if $\mathbb{E}(\epsilon_i \mid X_i, U_i = c) = 0$. This is equivalent to $\mathbb{E}(\tau_i \mid X_i, U_i = c) = V_i^{\top} \beta'_c$, which is ensured by Lemma S9(ii).

S2.4 Two linear algebra propositions

Propositions S4–S5 below state two linear algebra results that underlie Proposition 1(ii). We did not find formal statements or proofs of these results in the literature. We provide our original proofs in Section S5.

Proposition S4. Assume $X = (X_0, X_1, \ldots, X_m)^{\top}$ is an $(m+1) \times 1$ vector such that XX_0 is linear in (1, X). That is, there exists constants $\{c_i, b_i, a_{ij} \in \mathbb{R} : i = 0, \ldots, m; j = 1, \ldots, m\}$ such that

$$XX_{0} = \begin{pmatrix} X_{0} \\ X_{1} \\ \vdots \\ X_{m} \end{pmatrix} X_{0} = \begin{pmatrix} c_{0} \\ c_{1} \\ \vdots \\ c_{m} \end{pmatrix} + \begin{pmatrix} b_{0} \\ b_{1} \\ \vdots \\ b_{m} \end{pmatrix} X_{0} + \begin{pmatrix} a_{01} & \dots & a_{0m} \\ a_{11} & \dots & a_{1m} \\ \vdots \\ a_{m1} & \dots & a_{mm} \end{pmatrix} \begin{pmatrix} X_{1} \\ \vdots \\ X_{m} \end{pmatrix}.$$
(S27)

Then X_0 takes at most m + 2 distinct values across all solutions of X to (S27).

Proposition S5 below extends Proposition S4 and ensures that if XX_0, XX_1, \ldots, XX_m are all linear in (1, X), then the whole vector $X = (X_0, X_1, \ldots, X_m)^{\top}$ takes at most m + 2distinct values.

Proposition S5. Assume $X = (X_1, \ldots, X_m)^{\top}$ is an $m \times 1$ vector such that XX^{\top} is component-wise linear in (1, X). That is, there exists constants $\{c_{ij}, a_{ij[k]} \in \mathbb{R} : i, j, k = 1, \ldots, m\}$ such that

$$X_i X_j = c_{ij} + \sum_{k=1}^m a_{ij[k]} X_k$$
 for $i, j = 1, \dots, m$.

Then X takes at most m + 1 distinct values.

Corollary S1 is a direct implication of Propositions S4–S5.

Corollary S1. Let $X = (X_1, \ldots, X_m)^{\top}$ be an $m \times 1$ vector.

- (i) If there exists an $m \times 1$ constant vector $c = (c_1, \ldots, c_m)^{\top}$ such that $XX^{\top}c$ is linear in (1, X), then $X^{\top}c$ takes at most m + 1 distinct values.
- (ii) If $XX^{\top}c$ is linear in (1, X) for all $m \times 1$ constant vectors $c \in \mathbb{R}^m$, then X takes at most m + 1 distinct values.

S3 Proofs of the results in the main paper

S3.1 Proofs of the results in Section 2

Proof of Proposition 1. We verify below Proposition 1(i)–(iii), respectively.

Proof of Proposition 1(i). The sufficiency of categorical covariates follows from Lemma S1. The sufficiency of randomly assigned IV follows from $\mathbb{E}(Z_iX_i \mid X_i) = \mathbb{E}(Z_i \mid X_i)X_i = \mathbb{E}(Z_i)X_i$.

Proof of Proposition 1(ii). Let $X_{i[-1]} = (X_1, \ldots, X_{K-1})^\top$ with $X_i = (1, X_{i[-1]}^\top)^\top$. Assumption 2 requires that

$$\mathbb{E}(Z_i \mid X_i)$$
 and $\mathbb{E}(Z_i X_{i[-1]} \mid X_i)$ are both linear in X_i . (S28)

From (S28), there exists constants $a_0 \in \mathbb{R}$ and $a_X \in \mathbb{R}^{K-1}$ so that

$$\mathbb{E}(Z_i \mid X_i) = a_0 + X_{i[-1]}^{\top} a_X.$$
(S29)

This ensures

$$\mathbb{E}(Z_i X_{i[-1]} \mid X_i) = X_{i[-1]} \cdot \mathbb{E}(Z_i \mid X_i) \stackrel{(S29)}{=} X_{i[-1]} a_0 + X_{i[-1]} X_{i[-1]}^\top a_X.$$
(S30)

From (S30), for $\mathbb{E}(Z_i X_{i[-1]} | X_i)$ to be linear in $(1, X_{i[-1]})$ as suggested by (S28), we need $X_{i[-1]}X_{i[-1]}^{\top}a_X$ to be linear in $(1, X_{i[-1]})$. By Corollary S1(i), this implies that $X_{i[-1]}^{\top}a_X$ takes at most K distinct values, which in turn ensures $\mathbb{E}(Z_i | X_i)$ takes at most K distinct values given (S29).

In addition, that Assumption 2 holds for arbitrary IV assignment mechanism implies that $X_{i[-1]}X_{i[-1]}^{\top}a_X$ is linear in $(1, X_{i[-1]})$ for arbitrary a_X . By Corollary S1(ii), this implies X_i is categorical.

Proof of Proposition 1(iii). Assume that the potential outcomes satisfy

$$\mathbb{E}\{Y_{i}(1) \mid X_{i}, U_{i} = \mathbf{a}\} = \mathbb{E}\{Y_{i}(1) \mid X_{i}, U_{i} = \mathbf{c}\} = X_{i}^{\top}\beta_{1},$$

$$\mathbb{E}\{Y_{i}(0) \mid X_{i}, U_{i} = \mathbf{n}\} = \mathbb{E}\{Y_{i}(0) \mid X_{i}, U_{i} = \mathbf{c}\} = X_{i}^{\top}\beta_{0}$$
(S31)

for some fixed $\beta_0, \beta_1 \in \mathbb{R}^K$. Define

$$\eta_i = Y_i - D_i X_i^\top (\beta_1 - \beta_0) - X_i^\top \beta_0 \tag{S32}$$

with

$$\eta_{i} = \begin{cases} Y_{i}(1) - X_{i}^{\top}\beta_{1} & \text{if } U_{i} = a; \\ Y_{i}(0) - X_{i}^{\top}\beta_{0} & \text{if } U_{i} = n; \\ Y_{i}(1) - X_{i}^{\top}\beta_{1} & \text{if } U_{i} = c \text{ and } Z_{i} = 1; \\ Y_{i}(0) - X_{i}^{\top}\beta_{0} & \text{if } U_{i} = c \text{ and } Z_{i} = 0. \end{cases}$$
(S33)

Under Assumption 1, we have

$$\mathbb{E}(\eta_i \mid Z_i, X_i, U_i = \mathbf{a}) \stackrel{(S33)}{=} \mathbb{E}\left\{Y_i(1) - X_i^\top \beta_1 \mid Z_i, X_i, U_i = \mathbf{a}\right\}$$

$$\stackrel{\text{Assm. 1}}{=} \mathbb{E}\left\{Y_i(1) - X_i^\top \beta_1 \mid X_i, U_i = \mathbf{a}\right\}$$

$$\stackrel{(S31)}{=} 0,$$

$$\mathbb{E}(\eta_i \mid Z_i, X_i, U_i = \mathbf{n}) \stackrel{(S33)}{=} \mathbb{E}\left\{Y_i(0) - X_i^\top \beta_0 \mid Z_i, X_i, U_i = \mathbf{n}\right\}$$

$$\stackrel{\text{Assm. 1}}{=} \mathbb{E}\left\{Y_i(0) - X_i^\top \beta_0 \mid X_i, U_i = \mathbf{n}\right\}$$

$$\stackrel{(S31)}{=} 0,$$

and

$$\mathbb{E}(\eta_{i} \mid Z_{i} = 1, X_{i}, U_{i} = c) \stackrel{(S33)}{=} \mathbb{E}\left\{Y_{i}(1) - X_{i}^{\top}\beta_{1} \mid Z_{i} = 1, X_{i}, U_{i} = c\right\}
\stackrel{\text{Assm. 1}}{=} \mathbb{E}\left\{Y_{i}(1) - X_{i}^{\top}\beta_{1} \mid X_{i}, U_{i} = c\right\}
\stackrel{(S31)}{=} 0,
\mathbb{E}(\eta_{i} \mid Z_{i} = 0, X_{i}, U_{i} = c) \stackrel{(S33)}{=} \mathbb{E}\left\{Y_{i}(0) - X_{i}^{\top}\beta_{0} \mid Z_{i} = 0, X_{i}, U_{i} = c\right\}
\stackrel{\text{Assm. 1}}{=} \mathbb{E}\left\{Y_{i}(0) - X_{i}^{\top}\beta_{0} \mid X_{i}, U_{i} = c\right\}
\stackrel{(S31)}{=} 0.$$

This ensures $\mathbb{E}(\eta_i \mid Z_i, X_i, U_i) = 0$ so that

$$\mathbb{E}(\eta_i \mid Z_i, X_i) = \mathbb{E}\{\mathbb{E}(\eta_i \mid Z_i, X_i, U_i) \mid Z_i, X_i\} = 0.$$
(S34)

(S32) and (S34) ensure that Assumption 3 holds.

S3.2 Proofs of the results in Section 3.2

Proof of Theorem 2. With a slight abuse of notation, renew $\widehat{DX}_i = \operatorname{proj}(D_iX_i \mid Z_iX_i, X_i)$ as the *population* fitted value from the first stage of the interacted 2SLS throughout this proof. From Proposition S1, β_{2SLS} equals the coefficient vector of \widehat{DX}_i in $\operatorname{proj}(Y_i \mid \widehat{DX}_i, X_i)$. Recall $\widehat{DX}_i = \operatorname{res}(\widehat{DX}_i \mid X_i)$. The population FWL in Lemma S3 ensures

$$\beta_{2\text{SLS}} = \left\{ \mathbb{E} \left(\widetilde{DX}_i \cdot \widetilde{DX}_i^\top \right) \right\}^{-1} \mathbb{E} \left(\widetilde{DX}_i \cdot Y_i \right).$$
(S35)

Write $Y_i = (D_i X_i)^\top \beta_c + (Y_i - D_i X_i^\top \beta_c)$ with

$$\widetilde{DX}_i \cdot Y_i = \widetilde{DX}_i \cdot (D_i X_i)^\top \beta_{\rm c} + \widetilde{DX}_i \cdot (Y_i - D_i X_i^\top \beta_{\rm c}).$$

This ensures the $\mathbb{E}(\widetilde{DX}_i \cdot Y_i)$ on the right-hand side of (S35) equals

$$\mathbb{E}\left(\widetilde{DX}_{i}\cdot Y_{i}\right) = \mathbb{E}\left\{\widetilde{DX}_{i}\cdot (D_{i}X_{i})^{\top}\right\} \cdot \beta_{c} + \mathbb{E}\left\{\widetilde{DX}_{i}\cdot (Y_{i}-D_{i}X_{i}^{\top}\beta_{c})\right\} \\
= \mathbb{E}\left\{\widetilde{DX}_{i}\cdot (D_{i}X_{i})^{\top}\right\} \cdot \beta_{c} + B,$$
(S36)

where

$$B = \mathbb{E}\left\{\widetilde{DX}_i \cdot (Y_i - D_i X_i^\top \beta_c)\right\}.$$

Plugging (S36) in (S35) ensures

$$\beta_{2\text{SLS}} = \left\{ \mathbb{E}\left(\widetilde{DX}_i \cdot \widetilde{DX}_i^{\mathsf{T}}\right) \right\}^{-1} \mathbb{E}\left\{\widetilde{DX}_i \cdot (D_i X_i)^{\mathsf{T}}\right\} \cdot \beta_{\mathsf{c}} + \left\{ \mathbb{E}\left(\widetilde{DX}_i \cdot \widetilde{DX}_i^{\mathsf{T}}\right) \right\}^{-1} B.$$

Therefore, it suffices to show that

$$\mathbb{E}\left\{\widetilde{DX}_i \cdot (D_i X_i)^{\top}\right\} = \mathbb{E}\left(\widetilde{DX}_i \cdot \widetilde{DX}_i^{\top}\right), \quad B = B_1 + B_2.$$
(S37)

The sufficiency of Assumptions 2 and 3 to ensure $B_1 = B_2 = 0$ then follows from Section 4 and Lemma S5.

Proof of $\mathbb{E}\{\widetilde{DX}_i \cdot (D_i X_i)^{\top}\} = \mathbb{E}(\widetilde{DX}_i \cdot \widetilde{DX}_i^{\top})$ in (S37). Write

$$D_{i}X_{i} - \widetilde{DX}_{i} = \left(D_{i}X_{i} - \widehat{DX}_{i}\right) + \left(\widehat{DX}_{i} - \widetilde{DX}_{i}\right)$$
$$= \operatorname{res}(D_{i}X_{i} \mid Z_{i}X_{i}, X_{i}) + \operatorname{proj}(\widehat{DX}_{i} \mid X_{i}).$$
(S38)

By definition, $\widetilde{DX}_i = \widehat{DX}_i - \operatorname{proj}(\widehat{DX}_i \mid X_i)$ is a linear combination of (Z_iX_i, X_i) . Properties of linear projection ensure

$$\mathbb{E}\left\{\widetilde{DX}_{i}\cdot\operatorname{res}(D_{i}X_{i}\mid Z_{i}X_{i}, X_{i})^{\mathsf{T}}\right\} = 0, \quad \mathbb{E}\left\{\widetilde{DX}_{i}\cdot\operatorname{proj}(\widehat{DX}_{i}\mid X_{i})^{\mathsf{T}}\right\} = 0.$$
(S39)

(S38)-(S39) together ensure

$$\mathbb{E}\left\{\widetilde{DX}_{i}\cdot(D_{i}X_{i})^{\top}\right\} - \mathbb{E}\left(\widetilde{DX}_{i}\cdot\widetilde{DX}_{i}^{\top}\right)$$

$$= \mathbb{E}\left\{\widetilde{DX}_{i}\cdot(D_{i}X_{i}-\widetilde{DX}_{i})^{\top}\right\}$$

$$\stackrel{(S38)}{=} \mathbb{E}\left\{\widetilde{DX}_{i}\cdot\operatorname{res}(D_{i}X_{i}\mid Z_{i}X_{i},X_{i})^{\top}\right\} + \mathbb{E}\left\{\widetilde{DX}_{i}\cdot\operatorname{proj}(\widehat{DX}_{i}\mid X_{i})^{\top}\right\}$$

$$\stackrel{(S39)}{=} 0.$$

Proof of $B = B_1 + B_2$ in (S37). Recall from (S3) that $Y_i - D_i X_i^{\top} = \Delta_i + Z_i \epsilon_i$. We have

$$B = \mathbb{E}\left\{\widetilde{DX}_{i} \cdot (Y_{i} - D_{i}X_{i}^{\top}\beta_{c})\right\}$$

$$\stackrel{(S3)}{=} \mathbb{E}\left\{\widetilde{DX}_{i} \cdot (\Delta_{i} + Z_{i}\epsilon_{i})\right\}$$

$$= \mathbb{E}(\widetilde{DX}_{i} \cdot \Delta_{i}) + \mathbb{E}(\widetilde{DX}_{i} \cdot Z_{i}\epsilon_{i}).$$

We show below

$$B_1 = \mathbb{E}(\widetilde{DX}_i \cdot \Delta_i), \quad B_2 = \mathbb{E}(\widetilde{DX}_i \cdot Z_i \epsilon_i)$$
(S40)

to complete the proof. Recall from (S2)

$$\widetilde{DX}_{i} = C_{1} \Big\{ Z_{i} X_{i} - \operatorname{proj}(Z_{i} X_{i} \mid X_{i}) \Big\}$$
(S41)

$$\stackrel{(S2)}{=} C_1(Z_i X_i - C_2 X_i) \tag{S42}$$

so that

$$\widetilde{DX}_{i} \cdot Z_{i} \stackrel{(S42)}{=} C_{1}(Z_{i}X_{i} - C_{2}Z_{i}X_{i}) = C_{1}(I_{K} - C_{2}) \cdot Z_{i}X_{i}.$$
(S43)

Proof of $B_1 = \mathbb{E}(\widetilde{DX}_i \cdot \Delta_i)$ in (S40). Let

$$\xi_i = \Delta_i - \operatorname{proj}(\Delta_i \mid X_i) \tag{S44}$$

denote the residual from the linear projection of Δ_i on X_i . Properties of linear projection ensure $\mathbb{E}\{\widetilde{DX}_i \cdot \operatorname{proj}(\Delta_i \mid X_i)\} = 0$ so that

$$\mathbb{E}(\widetilde{DX}_i \cdot \Delta_i) \stackrel{(\text{S44})}{=} \mathbb{E}(\widetilde{DX}_i \cdot \xi_i) + \mathbb{E}\left\{\widetilde{DX}_i \cdot \operatorname{proj}(\Delta_i \mid X_i)\right\} = \mathbb{E}(\widetilde{DX}_i \cdot \xi_i).$$
(S45)

In addition, ξ_i is a function of $\{Y_i(z, d), D_i(z), X_i : z = 0, 1; d = 0, 1\}$, whereas \widetilde{DX}_i is a function of (Z_i, X_i) . This ensures

$$\xi_i \perp DX_i \mid X_i \tag{S46}$$

under Assumption 1(i). Accordingly, we have

$$\mathbb{E}(\widetilde{DX}_{i} \cdot \Delta_{i}) \stackrel{(S45)}{=} \mathbb{E}(\widetilde{DX}_{i} \cdot \xi_{i}) \\
= \mathbb{E}\left\{\mathbb{E}(\widetilde{DX}_{i} \cdot \xi_{i} \mid X_{i})\right\} \\
\stackrel{(S46)}{=} \mathbb{E}\left\{\mathbb{E}(\widetilde{DX}_{i} \mid X_{i}) \cdot \mathbb{E}(\xi_{i} \mid X_{i})\right\} \\
\stackrel{(S41)+(S44)}{=} C_{1} \cdot \mathbb{E}\left[\left\{\mathbb{E}(Z_{i}X_{i} \mid X_{i}) - \operatorname{proj}(Z_{i}X_{i} \mid X_{i})\right\} \cdot \left\{\mathbb{E}(\Delta_{i} \mid X_{i}) - \operatorname{proj}(\Delta_{i} \mid X_{i})\right\} \\
= B_{1},$$

where the second to last equality follows from

$$\mathbb{E}(\widetilde{DX}_i \mid X_i) = C_1 \cdot \left\{ \mathbb{E}(Z_i X_i \mid X_i) - \operatorname{proj}(Z_i X_i \mid X_i) \right\} \text{ by (S41)},$$
$$\mathbb{E}(\xi_i \mid X_i) = \mathbb{E}(\Delta_i \mid X_i) - \operatorname{proj}(\Delta_i \mid X_i) \text{ by (S44)}.$$

Proof of $B_2 = \mathbb{E}(\widetilde{DX}_i \cdot Z_i \epsilon_i)$ in (S40). Note that ϵ_i is a function of $\{Y_i(z, d), D_i(z), X_i : z = 0, 1; d = 0, 1\}$. Assumption 1(i) ensures

$$Z_i X_i \perp \epsilon_i \mid X_i. \tag{S47}$$

Accordingly, we have

$$\mathbb{E}(Z_{i}X_{i} \cdot \epsilon_{i}) = \mathbb{E}\left\{\mathbb{E}(Z_{i}X_{i} \cdot \epsilon_{i} \mid X_{i})\right\}$$

$$\stackrel{(S47)}{=} \mathbb{E}\left\{\mathbb{E}(Z_{i}X_{i} \mid X_{i}) \cdot \mathbb{E}(\epsilon_{i} \mid X_{i})\right\}$$

$$= \mathbb{E}\left[\mathbb{E}\left\{\mathbb{E}(Z_{i}X_{i} \mid X_{i}) \cdot \epsilon_{i} \mid X_{i}\right\}\right]$$

$$= \mathbb{E}\left\{\mathbb{E}(Z_{i}X_{i} \mid X_{i}) \cdot \epsilon_{i}\right\}$$

$$= \mathbb{E}\left\{\mathbb{E}(Z_{i}X_{i} \mid X_{i}) \cdot \epsilon_{i} \mid U_{i} = c\right\} \cdot \mathbb{P}(U_{i} = c).$$
(S48)

This, combined with (S43), ensures

$$\mathbb{E}(\widetilde{DX}_i \cdot Z_i \epsilon_i) \stackrel{(S43)}{=} C_1(I_K - C_2) \cdot \mathbb{E}(Z_i X_i \cdot \epsilon_i)$$

$$\stackrel{(S48)}{=} C_1(I_K - C_2) \cdot \mathbb{E}\left\{\mathbb{E}(Z_i X_i \mid X_i) \cdot \epsilon_i \mid U_i = c\right\} \cdot \mathbb{P}(U_i = c) = B_2.$$

Proof of Lemma 1. The expression of $\tau_{c}(X_{i})$ follows from Lemma S2(iii) and Lemma S1(i). We verify below the expression of β_{c} .

Recall from Eq. (4) that $\beta_c = \{\mathbb{E}(X_i X_i^\top \mid U_i = c)\}^{-1} \cdot \mathbb{E}(X_i \tau_i \mid U_i = c)$. When $X_i = (1(X_i^* = 1), \dots, 1(X_i^* = K))^\top$, we have

(i) $X_i X_i^{\top} = \text{diag}\{1(X_i^* = k)\}_{k=1}^K$ so that

$$\mathbb{E}(X_i X_i^\top \mid U_i = \mathbf{c}) = \operatorname{diag}(p_k)_{k=1}^K,$$
(S49)

where $p_k = \mathbb{E}\{1(X_i^* = k) \mid U_i = c\} = \mathbb{P}(X_i^* = k \mid U_i = c).$

(ii)

$$\mathbb{E}\left(X_i\tau_i \mid U_i = \mathbf{c}\right) = (a_1, \dots, a_K)^{\mathsf{T}},\tag{S50}$$

where

$$a_{k} = \mathbb{E} \Big\{ 1(X_{i}^{*} = k) \cdot \tau_{i} \mid U_{i} = c \Big\}$$

= $\mathbb{E} (\tau_{i} \mid X_{i}^{*} = k, U_{i} = c) \cdot \mathbb{P} (X_{i}^{*} = k \mid U_{i} = c) = \tau_{[k]c} \cdot p_{k}$

Eqs. (4) and (S49)–(S50) together ensure the kth element of β_c equals $p_k^{-1} \cdot a_k = \tau_{[k]c}$ for $k = 1, \ldots, K$.

Proof of Corollary 1. Let $\hat{\tau}_{\text{wald},[k]}$ denote the Wald estimator based on units with $X_i^* = k$. We show in the following that

$$\hat{\beta}_{2\text{SLS}} = (\hat{\tau}_{\text{wald},[1]}, \dots, \hat{\tau}_{\text{wald},[K]})^{\top}, \tag{S51}$$

which ensures $\beta_{2\text{SLS}} = (\tau_{\text{wald},[1]}, \ldots, \tau_{\text{wald},[K]})^{\top} = \tau_{\text{c}}.$

Recall the definition of forbidden regression from above Lemma S6. In addition, consider the following stratum-wise 2SLS based on units with $X_i^* = k$:

- (i) First stage: Fit $lm(D_i \sim 1 + Z_i)$ over $\{i : X_i^* = k\}$. Denote the fitted value by \hat{D}_i .
- (ii) Second stage: Fit $lm(Y_i \sim 1 + \hat{D}_i)$ over $\{i : X_i^* = k\}$. Denote the coefficient of \hat{D}_i by $\tilde{\beta}_{2\text{SLS},[k]}$.

When $X_i = (1(X_i^* = 1), \dots, 1(X_i^* = K))^{\top}$,

(a) properties of least squares ensure that the coefficients of $\hat{D}_i X_i$ from the forbidden regression equal $(\tilde{\beta}_{2\text{SLS},[1]}, \ldots, \tilde{\beta}_{2\text{SLS},[K]})^{\top}$;

(b) Lemma S6 ensures the coefficients of $\hat{D}_i X_i$ from the forbidden regression equal $\hat{\beta}_{2SLS}$.

These two observations ensure $\hat{\beta}_{2\text{SLS}} = (\tilde{\beta}_{2\text{SLS},[1]}, \dots, \tilde{\beta}_{2\text{SLS},[K]})^{\top}$. This, together with $\tilde{\beta}_{2\text{SLS},[k]} = \hat{\tau}_{\text{wald},[k]}$ by standard theory, ensures (S51).

S3.3 Proofs of the results in Section 3.3

Proof of Lemma 2. Recall

$$\beta_{\mathbf{c}} = \{ \mathbb{E}(X_i X_i^\top \mid U_i = \mathbf{c}) \}^{-1} \mathbb{E} (X_i \tau_i \mid U_i = \mathbf{c})$$

from Eq. (4). Let $X_{i[-1]} = (X_{i1}, \ldots, X_{i,K-1})^{\top}$ to write $X_i = (1, X_{i[-1]}^{\top})^{\top}$. We have

$$X_{i}X_{i}^{\top} = \begin{pmatrix} 1 \\ X_{i[-1]} \end{pmatrix} \begin{pmatrix} 1, X_{i[-1]}^{\top} \end{pmatrix} = \begin{pmatrix} 1 & X_{i[-1]}^{\top} \\ X_{i[-1]} & X_{i[-1]}X_{i[-1]}^{\top} \end{pmatrix}, \quad X_{i}\tau_{i} = \begin{pmatrix} 1 \\ X_{i[-1]} \end{pmatrix} \tau_{i} = \begin{pmatrix} \tau_{i} \\ X_{i[-1]} \tau_{i} \end{pmatrix}$$

with

$$\mathbb{E}(X_i X_i^\top \mid U_i = \mathbf{c}) = \begin{pmatrix} 1 & \Phi \\ \Phi^\top & D \end{pmatrix}, \quad \mathbb{E}(X_i \tau_i \mid U_i = \mathbf{c}) = \begin{pmatrix} \tau_{\mathbf{c}} \\ \mathbb{E}(X_{i[-1]} \tau_i \mid U_i = \mathbf{c}) \end{pmatrix},$$

where $\Phi = \mathbb{E}(X_{i[-1]}^{\top} \mid U_i = c)$ and $D = \mathbb{E}(X_{i[-1]}X_{i[-1]}^{\top} \mid U_i = c)$. Therefore, $\Phi = 0$ ensures that the first element of β_c equals τ_c .

S3.4 Proofs of the results in Section 3.1

Proof of Theorem 1. Let $\check{X}_i^0 = (1, X_{i1} - \mathbb{E}(X_{i1} | U_i = c), \dots, X_{i,K-1} - \mathbb{E}(X_{i,K-1} | U_i = c))^{\top}$ denote the population analog of X_i^0 . Let $\hat{\tau}_{\times\times}(X_i^0) = \hat{\tau}_{\times\times}^*$ and $\hat{\tau}_{\times\times}(\check{X}_i^0)$ denote the first elements of $\hat{\beta}_{2\text{SLS}}(X_i^0)$ and $\hat{\beta}_{2\text{SLS}}(\check{X}_i^0)$, respectively. Then $\hat{\tau}_{\times\times}(X_i^0)$ and $\hat{\tau}_{\times\times}(\check{X}_i^0)$ have the same probability limit.

In addition, Lemma 2 ensures that $\tau_{\rm c}$ equals the first element of $\beta_{\rm c}(\check{X}_i^0)$. This ensures

$$plim_{N\to\infty} \hat{\tau}_{\times\times}(X_i^0) = \tau_c$$

$$\iff plim_{N\to\infty} \hat{\tau}_{\times\times}(\check{X}_i^0) = \tau_c$$

$$\stackrel{\text{Lemma 2}}{\iff} \text{ the first element of } \beta_{2\text{SLS}}(\check{X}_i^0) \text{ equals the first element of } \beta_c(\check{X}_i^0)$$

$$\stackrel{\text{Proposition S2}}{\iff} \text{ the first element of } \beta_{2\text{SLS}}(X_i) \text{ equals the first element of } \beta_c(X_i), \quad (S52)$$

where the last equivalence follows from \check{X}_i^0 being a nondegenerate linear transformation of X_i and the invariance of 2SLS to nondegenerate linear transformations in Proposition S2. By Theorem 2, the equivalent result in (S52) is ensured by either Assumption 2 or 3.

Theorem 1 ensures Proposition 2(iii). We verify below Proposition 2(i) and (ii).

Proof of Proposition 2(i). Assume throughout this proof that $\check{D}_i = \operatorname{proj}(D_i \mid Z_i, X_i)$ denotes the population fitted value from the additive first stage $\operatorname{lm}(D_i \sim Z_i + X_i)$, and let $\check{D}_i = \check{D}_i - \operatorname{proj}(\check{D}_i \mid X_i)$ denote the residual from the linear projection of \check{D}_i on X_i . Given Proposition S3, it suffices to verify that when $\mathbb{E}(Z_i \mid X_i)$ is linear in X_i , we have

$$\mathbb{E}(\check{D}_i \mid X_i) \text{ is linear in } X_i, \quad \alpha(X_i) = \frac{\mathbb{E}(D_i \check{D}_i \mid X_i)}{\mathbb{E}(\check{D}_i^2)} = w_+(X_i).$$
(S53)

Write

$$\check{D}_i = a_+ Z_i + b^\top X_i, \quad \text{where} \quad a_+ = \frac{\mathbb{E}\left\{ \operatorname{var}(Z_i \mid X_i) \cdot \pi(X_i) \right\}}{\mathbb{E}\left\{ \operatorname{var}(Z_i \mid X_i) \right\}}$$
(S54)

by Angrist (1998); see also Sloczynski (2022, Lemma A.1).

Proof of linear $\mathbb{E}(\check{D}_i \mid X_i)$ in (S53). From (S54), we have $\mathbb{E}(\check{D}_i \mid X_i) = a_+ \cdot \mathbb{E}(Z_i \mid X_i) + b^\top X_i$, which is linear in X_i when $\mathbb{E}(Z_i \mid X_i)$ is linear in X_i .

Proof of $\alpha(X_i) = w_+(X_i)$ in (S53). When $\mathbb{E}(Z_i \mid X_i)$ is linear in X_i , we just showed that $\mathbb{E}(\check{D}_i \mid X_i)$ is linear in X_i . This ensures $\operatorname{proj}(\check{D}_i \mid X_i) = \mathbb{E}(\check{D}_i \mid X_i)$ so that

$$\tilde{D}_{i} = \check{D}_{i} - \operatorname{proj}(\check{D}_{i} \mid X_{i})
= \check{D}_{i} - \mathbb{E}(\check{D}_{i} \mid X_{i})
\stackrel{(S54)}{=} a_{+} \Big\{ Z_{i} - \mathbb{E}(Z_{i} \mid X_{i}) \Big\}.$$
(S55)

This ensures

$$\mathbb{E}(\tilde{D}_i^2 \mid X_i) \stackrel{(S55)}{=} a_+^2 \cdot \mathbb{E}\Big[\big\{Z_i - \mathbb{E}(Z_i \mid X_i)\big\}^2 \mid X_i\Big] = a_+^2 \operatorname{var}(Z_i \mid X_i),$$

and therefore

$$\mathbb{E}(\tilde{D}_i^2) = a_+^2 \cdot \mathbb{E}\left\{ \operatorname{var}(Z_i \mid X_i) \right\}.$$
(S56)

In addition, (S55) ensures

$$\mathbb{E}(D_i \tilde{D}_i \mid Z_i, X_i) = \mathbb{E}(D_i \mid Z_i, X_i) \cdot \tilde{D}_i = a_+ \cdot \mathbb{E}(D_i \mid Z_i, X_i) \cdot \{Z_i - \mathbb{E}(Z_i \mid X_i)\},\$$

which implies

$$\mathbb{E}(D_i \tilde{D}_i \mid Z_i = 1, X_i) = a_+ \cdot \mathbb{E}(D_i \mid Z_i = 1, X_i) \cdot \{1 - \mathbb{E}(Z_i \mid X_i)\}, \\
\mathbb{E}(D_i \tilde{D}_i \mid Z_i = 0, X_i) = -a_+ \cdot \mathbb{E}(D_i \mid Z_i = 0, X_i) \cdot \mathbb{E}(Z_i \mid X_i).$$
(S57)

Therefore, we have

$$\mathbb{E}(D_{i}\tilde{D}_{i} \mid X_{i}) = \mathbb{E}(D_{i}\tilde{D}_{i} \mid Z_{i} = 1, X_{i}) \cdot \mathbb{P}(Z_{i} = 1 \mid X_{i}) \\
+ \mathbb{E}(D_{i}\tilde{D}_{i} \mid Z_{i} = 0, X_{i}) \cdot \mathbb{P}(Z_{i} = 0 \mid X_{i}) \\
\stackrel{(\text{S57})}{=} a_{+} \cdot \mathbb{E}(D_{i} \mid Z_{i} = 1, X_{i}) \cdot \{1 - \mathbb{E}(Z_{i} \mid X_{i})\} \cdot \mathbb{E}(Z_{i} \mid X_{i}) \\
- a_{+} \cdot \mathbb{E}(D_{i} \mid Z_{i} = 0, X_{i}) \cdot \mathbb{E}(Z_{i} \mid X_{i}) \cdot \{1 - \mathbb{E}(Z_{i} \mid X_{i})\} \\
= a_{+} \cdot \{\mathbb{E}(D_{i} \mid Z_{i} = 1, X_{i}) - \mathbb{E}(D_{i} \mid Z_{i} = 0, X_{i})\} \cdot \mathbb{E}(Z_{i} \mid X_{i}) \cdot \{1 - \mathbb{E}(Z_{i} \mid X_{i})\} \\
= a_{+} \cdot \pi(X_{i}) \cdot \operatorname{var}(Z_{i} \mid X_{i}).$$
(S58)

Eqs. (S54), (S56), and (S58) together ensure

$$\alpha(X_i) = \frac{\mathbb{E}(D_i \tilde{D}_i \mid X_i)}{\mathbb{E}(\tilde{D}_i^2)}$$

$$\stackrel{(S56)+(S58)}{=} \frac{a_+ \cdot \pi(X_i) \cdot \operatorname{var}(Z_i \mid X_i)}{a_+^2 \cdot \mathbb{E}\left\{\operatorname{var}(Z_i \mid X_i)\right\}}$$

$$= \frac{\pi(X_i) \cdot \operatorname{var}(Z_i \mid X_i)}{a_+ \cdot \mathbb{E}\left\{\operatorname{var}(Z_i \mid X_i)\right\}}$$

$$\stackrel{(S54)}{=} \frac{\pi(X_i) \cdot \operatorname{var}(Z_i \mid X_i)}{\mathbb{E}\left\{\pi(X_i) \cdot \operatorname{var}(Z_i \mid X_i)\right\}} = w_+(X_i).$$

Proof of Proposition 2(ii). Assume throughout this proof that $\check{D}_i = \text{proj}(D_i \mid Z_i X_i, X_i)$ denotes the population fitted value from the interacted first stage $\text{lm}(D_i \sim Z_i X_i + X_i)$, and let $\tilde{D}_i = \check{D}_i - \text{proj}(\check{D}_i \mid X_i)$ denote the residual from the linear projection of \check{D}_i on X_i . Given Proposition S3, it suffices to verify that when $\mathbb{E}(Z_i X_i \mid X_i)$ is linear in X_i , we have

$$\mathbb{E}(\check{D}_i \mid X_i) \text{ is linear in } X_i, \quad \alpha(X_i) = \frac{\mathbb{E}(D_i \tilde{D}_i \mid X_i)}{\mathbb{E}(\tilde{D}_i^2)} = w_{\times}(X_i).$$
(S59)

Write

$$\check{D}_i = a^\top Z_i X_i + b^\top X_i.$$
(S60)

Proof of linear $\mathbb{E}(\check{D}_i \mid X_i)$ in (S59). From (S60), we have $\mathbb{E}(\check{D}_i \mid X_i) = a^{\top} \mathbb{E}(Z_i X_i \mid X_i) + b^{\top} X_i$, which is linear in X_i when $\mathbb{E}(Z_i X_i \mid X_i)$ is linear in X_i .

Proof of $\alpha(X_i) = w_{\times}(X_i)$ in (S59). When $\mathbb{E}(Z_iX_i \mid X_i)$ is linear in X_i , we just showed that $\mathbb{E}(\check{D}_i \mid X_i)$ is linear in X_i . This ensures $\operatorname{proj}(\check{D}_i \mid X_i) = \mathbb{E}(\check{D}_i \mid X_i)$ so that

$$\tilde{D}_{i} = \check{D}_{i} - \operatorname{proj}(\check{D}_{i} \mid X_{i})
= \check{D}_{i} - \mathbb{E}(\check{D}_{i} \mid X_{i})
\stackrel{(S60)}{=} a^{\top}Z_{i}X_{i} - \mathbb{E}(a^{\top}Z_{i}X_{i} \mid X_{i}) = (a^{\top}X_{i}) \{Z_{i} - \mathbb{E}(Z_{i} \mid X_{i})\}.$$
(S61)

Note that the last expression in (S61) parallels (S55) after replacing a_+ with $a^{\top}X_i$. Replacing a_+ by $a^{\top}X_i$ in the proof of Proposition 2(i) after (S55) ensures that the renewed $\alpha(X_i)$ in (S59) satisfies

$$\alpha(X_i) = \frac{\mathbb{E}(D_i \tilde{D}_i \mid X_i)}{\mathbb{E}(\tilde{D}_i^2)} = \frac{(a^\top X_i) \cdot \pi(X_i) \cdot \operatorname{var}(Z_i \mid X_i)}{\mathbb{E}\{(a^\top X_i)^2 \cdot \operatorname{var}(Z_i \mid X_i)\}}.$$
(S62)

We show below $a = \mathbb{E}\{\operatorname{var}(Z_i \mid X_i) \cdot X_i X_i^{\top}\}^{-1} \mathbb{E}\{\operatorname{var}(Z_i \mid X_i) \cdot X_i \pi(X_i)\}$, which completes the proof.

By the population FWL, we have

$$a = \mathbb{E}\left\{ \operatorname{res}(Z_i X_i \mid X_i) \cdot Z_i X_i^{\top} \right\}^{-1} \mathbb{E}\left\{ \operatorname{res}(Z_i X_i \mid X_i) \cdot D_i \right\}.$$

Therefore, it suffices to show that

$$\mathbb{E}\left\{ \operatorname{res}(Z_i X_i \mid X_i) \cdot Z_i X_i^{\top} \right\} = \mathbb{E}\left\{ \operatorname{var}(Z_i \mid X_i) \cdot X_i X_i^{\top} \right\},$$
(S63)

$$\mathbb{E}\left\{\operatorname{res}(Z_iX_i \mid X_i) \cdot D_i\right\} = \mathbb{E}\left\{\operatorname{var}(Z_i \mid X_i) \cdot X_i \pi(X_i)\right\}.$$
(S64)

A useful fact is that when $\mathbb{E}(Z_i X_i \mid X_i)$ is linear in X_i , we have

$$\operatorname{res}(Z_i X_i \mid X_i) = Z_i X_i - \operatorname{proj}(Z_i X_i \mid X_i)$$
$$= Z_i X_i - \mathbb{E}(Z_i X_i \mid X_i) = \{Z_i - \mathbb{E}(Z_i \mid X_i)\} \cdot X_i$$

with

$$\mathbb{E}\left\{\operatorname{res}(Z_i X_i \mid X_i) \mid Z_i = 1, X_i\right\} = \left\{1 - \mathbb{E}(Z_i \mid X_i)\right\} \cdot X_i,$$
(S65)

$$\mathbb{E}\left\{ \operatorname{res}(Z_i X_i \mid X_i) \cdot h(X_i) \right\} = 0 \quad \text{for arbitrary } h(\cdot), \tag{S66}$$

$$\operatorname{res}(Z_i X_i \mid X_i) \cdot Z_i = \{1 - \mathbb{E}(Z_i \mid X_i)\} \cdot Z_i X_i.$$
(S67)

Proof of (S63). From (S65), we have

$$\mathbb{E}\left\{ \operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot Z_{i}X_{i}^{\top} \mid X_{i} \right\} = \mathbb{E}\left\{ \operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot Z_{i} \mid X_{i} \right\} \cdot X_{i}^{\top} \\ = \mathbb{P}(Z_{i} = 1 \mid X_{i}) \cdot \mathbb{E}\left\{ \operatorname{res}(Z_{i}X_{i} \mid X_{i}) \mid Z_{i} = 1, X_{i} \right\} \cdot X_{i}^{\top} \\ \stackrel{(\text{S65)}}{=} \mathbb{E}(Z_{i} \mid X_{i}) \cdot \left\{ 1 - \mathbb{E}(Z_{i} \mid X_{i}) \right\} \cdot X_{i} \cdot X_{i}^{\top} \\ = \operatorname{var}(Z_{i} \mid X_{i}) \cdot X_{i}X_{i}^{\top}.$$

This ensures

$$\mathbb{E}\left\{\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot Z_{i}X_{i}^{\top}\right\} = \mathbb{E}\left[\mathbb{E}\left\{\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot Z_{i}X_{i}^{\top} \mid X_{i}\right\}\right] = \mathbb{E}\left\{\operatorname{var}(Z_{i} \mid X_{i}) \cdot X_{i}X_{i}^{\top}\right\}.$$
(S68)

Proof of (S64). Let $\delta_i = D_i(1) - D_i(0)$ with $D_i = D_i(0) + Z_i \delta_i$ and $\mathbb{E}(\delta_i \mid Z_i, X_i) = \mathbb{E}(\delta_i \mid X_i) = \pi(X_i)$. This ensures

$$\mathbb{E}(D_i \mid Z_i, X_i) = \mathbb{E}\{D_i(0) \mid Z_i, X_i\} + Z_i \cdot \mathbb{E}(\delta_i \mid Z_i, X_i)$$
$$= \mathbb{E}\{D_i(0) \mid X_i\} + Z_i \cdot \pi(X_i)$$
(S69)

with

$$\mathbb{E}\left\{\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot D_{i} \mid Z_{i}, X_{i}\right\} = \operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot \mathbb{E}(D_{i} \mid Z_{i}, X_{i}) \\ \stackrel{(\mathrm{S69})}{=} \operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot \mathbb{E}\{D_{i}(0) \mid X_{i}\} \\ + \operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot Z_{i} \cdot \pi(X_{i}). \quad (\mathrm{S70})$$

Note that the two terms in (S70) satisfy

$$\mathbb{E}\left[\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot \mathbb{E}\{D_{i}(0) \mid X_{i}\}\right] \stackrel{(\text{S66})}{=} 0, \\
\mathbb{E}\left\{\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot Z_{i} \cdot \pi(X_{i})\right\} \stackrel{(\text{S67})}{=} \mathbb{E}\left[\left\{1 - \mathbb{E}(Z_{i} \mid X_{i})\right\} \cdot Z_{i}X_{i} \cdot \pi(X_{i})\right] \\
= \mathbb{E}\left[\left\{1 - \mathbb{E}(Z_{i} \mid X_{i})\right\} \cdot \mathbb{E}(Z_{i} \mid X_{i}) \cdot X_{i} \cdot \pi(X_{i})\right]^{(\text{S71})} \\
= \mathbb{E}\left\{\operatorname{var}(Z_{i} \mid X_{i}) \cdot X_{i} \cdot \pi(X_{i})\right\}.$$

This ensures

$$\mathbb{E}\left\{\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot D_{i}\right\} = \mathbb{E}\left[\mathbb{E}\left\{\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot D_{i} \mid Z_{i}, X_{i}\right\}\right] \\ \stackrel{(S70)}{=} \mathbb{E}\left[\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot \mathbb{E}\left\{D_{i}(0) \mid X_{i}\right\}\right] \\ + \mathbb{E}\left\{\operatorname{res}(Z_{i}X_{i} \mid X_{i}) \cdot Z_{i} \cdot \pi(X_{i})\right\} \\ \stackrel{(S71)}{=} \mathbb{E}\left\{\operatorname{var}(Z_{i} \mid X_{i}) \cdot X_{i} \cdot \pi(X_{i})\right\}.$$

S3.5 Proofs of the results in Section 4

Proof of Proposition 3. Proposition 3(i) follows from the population FWL in Lemma S3. Proposition 3(ii) follows from Abadie (2003). We verify below Proposition 3(iii).

First, it follows from (S2) and Lemma S4 that

$$\mathbb{E}(\widetilde{DX}_i \mid X_i) = C_1 \Big\{ \mathbb{E}(Z_i X_i \mid X_i) - C_2 X_i \Big\}, \quad \text{where } C_1 \text{ is invertible.}$$
(S72)

Note that a necessary condition for $w_i = u_i$ is

$$\mathbb{E}(w_i \mid X_i) = \mathbb{E}(u_i \mid X_i), \tag{S73}$$

where it follows from $\mathbb{E}(\Delta \kappa \mid X_i) = 0$ that

$$\mathbb{E}(u_i \mid X_i) = \{\mathbb{E}(X_i X_i^\top \mid U_i = \mathbf{c})\}^{-1} \pi_{\mathbf{c}}^{-1} \cdot \mathbb{E}(\Delta \kappa \mid X_i) \cdot X_i = 0.$$

Therefore, the necessary condition in (S73) has the following equivalent forms:

$$(S73) \iff \mathbb{E}(w_i \mid X_i) = 0$$

$$\iff \mathbb{E}(DX_i \mid X_i) = 0$$

$$\stackrel{(S72)}{\iff} \mathbb{E}(Z_iX_i \mid X_i) = C_2X_i$$

This ensures the necessity of Assumption 2 to ensure $w_i = u_i$.

S3.6 Proofs of the results in Section 8

Proof of Proposition 4. Recall $\widetilde{DV_i}$ as the residual from the linear projection of $\operatorname{proj}(D_iV_i \mid Z_iV_i, X_i)$ on X_i . The population FWL in Lemma S3 ensures

$$\beta_{2_{\rm SLS}}' = \left\{ \mathbb{E} \left(\widetilde{DV_i} \cdot \widetilde{DV_i}^{\top} \right) \right\}^{-1} \mathbb{E} \left(\widetilde{DV_i} \cdot Y_i \right).$$
(S74)

Write $Y_i = (D_i V_i)^\top \beta'_{\rm c} + (Y_i - D_i V_i^\top \beta'_{\rm c})$ with

$$\widetilde{DV_i} \cdot Y_i = \widetilde{DV_i} \cdot (D_i V_i)^\top \beta_c' + \widetilde{DV_i} \cdot (Y_i - D_i V_i^\top \beta_c')$$

and

$$\mathbb{E}\left(\widetilde{DV}_{i}\cdot Y_{i}\right) = \mathbb{E}\left\{\widetilde{DV}_{i}\cdot (D_{i}V_{i})^{\top}\right\}\beta_{c}' + \mathbb{E}\left\{\widetilde{DV}_{i}\cdot (Y_{i}-D_{i}V_{i}^{\top}\beta_{c}')\right\} \\
= \mathbb{E}\left\{\widetilde{DV}_{i}\cdot (D_{i}V_{i})^{\top}\right\}\beta_{c}' + B'.$$
(S75)

Plugging (S75) in (S74) ensures

$$\beta_{2\text{SLS}}' = \left\{ \mathbb{E}\left(\widetilde{DV_i} \cdot \widetilde{DV_i}^{\mathsf{T}} \right) \right\}^{-1} \mathbb{E}\left\{ \widetilde{DV_i} \cdot (D_i V_i)^{\mathsf{T}} \right\} \beta_{\mathrm{c}}' + \left\{ \mathbb{E}\left(\widetilde{DV_i} \cdot \widetilde{DV_i}^{\mathsf{T}} \right) \right\}^{-1} B'.$$

Therefore, it suffices to show that

$$\mathbb{E}\left\{\widetilde{DV_i}\cdot(D_iV_i)^{\top}\right\} = \mathbb{E}\left(\widetilde{DV_i}\cdot\widetilde{DV_i}^{\top}\right).$$
(S76)

The sufficient conditions then follow from Lemma S9. We verify (S76) below.

Proof of (S76). With a slight abuse of notation, renew $\widehat{DV_i} = \operatorname{proj}(D_iV_i \mid Z_iV_i, X_i)$ and write

$$D_i V_i - \widetilde{DV_i} = D_i V_i - \widehat{DV_i} + \widehat{DV_i} - \widetilde{DV_i} = \operatorname{res}(D_i V_i \mid Z_i V_i, X_i) + \operatorname{proj}(\widetilde{DV_i} \mid X_i).$$
(S77)

Note that $\widetilde{DV_i}$ is by definition a linear combination of (Z_iV_i, X_i) . Properties of linear projection ensure

$$\mathbb{E}\left\{\widetilde{DV_i} \cdot \operatorname{res}(D_iV_i \mid Z_iV_i, X_i)^{\top}\right\} = 0, \quad \mathbb{E}\left\{\widetilde{DV_i} \cdot \operatorname{proj}(\widetilde{DV_i} \mid X_i)^{\top}\right\} = 0.$$
(S78)

(S77)-(S78) together ensure (S76) as follows:

$$\mathbb{E}\left\{\widetilde{DV_{i}}\cdot(D_{i}V_{i})^{\top}\right\} - \mathbb{E}\left(\widetilde{DV_{i}}\cdot\widetilde{DV_{i}}^{\top}\right)$$

$$= \mathbb{E}\left\{\widetilde{DV_{i}}\cdot(D_{i}V_{i}-\widetilde{DV_{i}})^{\top}\right\}$$

$$\stackrel{(S77)}{=} \mathbb{E}\left\{\widetilde{DV_{i}}\cdot\operatorname{res}(D_{i}V_{i}\mid Z_{i}V_{i}, X_{i})^{\top}\right\} - \mathbb{E}\left\{\widetilde{DV_{i}}\cdot\operatorname{proj}(\widetilde{DV_{i}}\mid X_{i})^{\top}\right\}$$

$$\stackrel{(S78)}{=} 0.$$

Proof of Proposition 5. The result follows from letting $Z_i = D_i$ in Theorems 1–2. \Box

S4 Proofs of the results in Section S1

Proof of Proposition S3. Recall $R_i = R(X_i, Z_i)$ as the regressor vector in the first stage of the generalized additive 2SLS in Definition S1. Recall $\check{D}_i = \text{proj}(D_i \mid R_i)$ and $\tilde{D}_i = \check{D}_i - \text{proj}(\check{D}_i \mid X_i)$. The correspondence between least squares and linear projection ensures τ_{*+} is the coefficient of \check{D}_i in $\text{proj}(Y_i \mid \check{D}_i, X_i)$. The population FWL in Lemma S3 ensures

$$\tau_{*+} = \frac{\mathbb{E}(Y_i \tilde{D}_i)}{\mathbb{E}(\tilde{D}_i^2)}.$$
(S79)

We simplify below the numerator $\mathbb{E}(Y_i \tilde{D}_i)$ in (S79).

Recall $\tau_{c}(X_{i}) = \mathbb{E}(\tau_{i} \mid U_{i} = c, X_{i})$. Let $\delta_{i} = D_{i}(1) - D_{i}(0)$ to write

$$D_i = D_i(0) + \delta_i Z_i. \tag{S80}$$

Write

$$\Delta_i = \begin{cases} Y_i(0) & \text{for compliers and never-takers} \\ Y_i(1) - \tau_{\rm c}(X_i) & \text{for always-takers} \end{cases} = Y_i(0) + \{\tau_i - \tau_{\rm c}(X_i)\} \cdot D_i(0).$$

We have

$$Y_{i} = Y_{i}(0) + \tau_{i} \cdot D_{i}$$

$$= Y_{i}(0) + \{\tau_{i} - \tau_{c}(X_{i})\} \cdot D_{i} + \tau_{c}(X_{i}) \cdot D_{i}$$

$$\stackrel{(S80)}{=} Y_{i}(0) + \{\tau_{i} - \tau_{c}(X_{i})\} \cdot \{D_{i}(0) + \delta_{i}Z_{i}\} + \tau_{c}(X_{i}) \cdot D_{i}$$

$$= Y_{i}(0) + \{\tau_{i} - \tau_{c}(X_{i})\} \cdot D_{i}(0) + \{\tau_{i} - \tau_{c}(X_{i})\} \cdot \delta_{i}Z_{i} + \tau_{c}(X_{i}) \cdot D_{i}$$

$$= \Delta_{i} + \{\tau_{i} - \tau_{c}(X_{i})\} \cdot \delta_{i}Z_{i} + \tau_{c}(X_{i}) \cdot D_{i}.$$
(S81)

(S81) ensures

$$\mathbb{E}(Y_i \tilde{D}_i) = \mathbb{E}(\Delta_i \cdot \tilde{D}_i) + \mathbb{E}\left[\left\{\tau_i - \tau_c(X_i)\right\} \delta_i \cdot Z_i \tilde{D}_i\right] + \mathbb{E}\left\{\tau_c(X_i) \cdot D_i \tilde{D}_i\right\}.$$
 (S82)

We compute below the three expectations on the right-hand side of (S82), respectively.

<u>Compute $\mathbb{E}(\Delta_i \cdot \tilde{D}_i)$ in (S82)</u>. Let $\xi_i = \Delta_i - \operatorname{proj}(\Delta_i \mid X_i)$ with

$$\mathbb{E}(\xi_i \mid X_i) = \mathbb{E}(\Delta_i \mid X_i) - \operatorname{proj}(\Delta_i \mid X_i).$$
(S83)

Properties of linear projection ensure $\mathbb{E}\{\operatorname{proj}(\Delta_i \mid X_i) \cdot \tilde{D}_i\} = 0$ so that

$$\mathbb{E}(\Delta_i \cdot \tilde{D}_i) = \mathbb{E}\left\{ \operatorname{proj}(\Delta_i \mid X_i) \cdot \tilde{D}_i \right\} + \mathbb{E}(\xi_i \cdot \tilde{D}_i) = \mathbb{E}(\xi_i \cdot \tilde{D}_i).$$
(S84)

In addition, ξ_i is a function of $\{Y_i(z,d), D_i(z), X_i : z = 0, 1; d = 0, 1\}$, whereas \tilde{D}_i is a function of (X_i, Z_i) . This ensures

$$\xi_i \perp \hspace{-0.1cm}\perp \hspace{-0.1cm} \tilde{D}_i \mid X_i \tag{S85}$$

under Assumption 1(i). (S83)–(S85) together ensure

$$\mathbb{E}(\Delta_{i} \cdot \tilde{D}_{i}) \stackrel{(\text{S84})}{=} \mathbb{E}(\xi_{i} \cdot \tilde{D}_{i} \mid X_{i}) \}$$

$$= \mathbb{E}\left\{\mathbb{E}(\xi_{i} \cdot \tilde{D}_{i} \mid X_{i})\right\}$$

$$\stackrel{(\text{S85})}{=} \mathbb{E}\left\{\mathbb{E}(\xi_{i} \mid X_{i}) \cdot \mathbb{E}(\tilde{D}_{i} \mid X_{i})\right\}$$

$$\stackrel{(\text{S83})}{=} \mathbb{E}\left[\left\{\mathbb{E}(\Delta_{i} \mid X_{i}) - \operatorname{proj}(\Delta_{i} \mid X_{i})\right\} \cdot \mathbb{E}(\tilde{D}_{i} \mid X_{i})\right]$$

$$= B,$$
(S86)

where the last equality follows from $\mathbb{E}(\tilde{D}_i \mid X_i) = \mathbb{E}(\check{D}_i \mid X_i) - \operatorname{proj}(\check{D}_i \mid X_i)$ by definition.

$$\{\tau_i - \tau_c(X_i)\}\delta_i \perp Z_i \tilde{D}_i \mid X_i$$
(S87)

under Assumption 1(i). In addition,

$$\mathbb{E}\Big[\big\{\tau_i - \tau_{\mathbf{c}}(X_i)\big\}\delta_i \mid X_i\Big] = \mathbb{E}\big\{\tau_i - \tau_{\mathbf{c}}(X_i) \mid U_i = \mathbf{c}, X_i\big\} \cdot \mathbb{P}(U_i = \mathbf{c} \mid X_i) = 0.$$
(S88)

(S87)-(S88) together ensure

$$\mathbb{E}\Big[\big\{\tau_i - \tau_{\mathbf{c}}(X_i)\big\}\delta_i \cdot Z_i \tilde{D}_i \mid X_i\Big] \stackrel{(S87)}{=} \mathbb{E}\Big[\big\{\tau_i - \tau_{\mathbf{c}}(X_i)\big\}\delta_i \mid X_i\Big] \cdot \mathbb{E}(Z_i \tilde{D}_i \mid X_i) \\ \stackrel{(S88)}{=} 0,$$

and therefore

$$\mathbb{E}\Big[\big\{\tau_i - \tau_c(X_i)\big\}\delta_i \cdot Z_i\tilde{D}_i\Big] = \mathbb{E}\Big(\mathbb{E}\Big[\big\{\tau_i - \tau_c(X_i)\big\}\delta_i \cdot Z_i\tilde{D}_i \mid X_i\Big]\Big) = 0.$$
(S89)

Compute $\mathbb{E}\{\tau_{\mathbf{c}}(X_i) \cdot D_i \tilde{D}_i\}$ in (S82). We have

$$\mathbb{E}\left\{\tau_{c}(X_{i})\cdot D_{i}\tilde{D}_{i}\mid X_{i}\right\} = \tau_{c}(X_{i})\cdot\mathbb{E}(D_{i}\tilde{D}_{i}\mid X_{i})$$
$$= \tau_{c}(X_{i})\cdot\alpha(X_{i})\cdot\mathbb{E}(\tilde{D}_{i}^{2}),$$

and therefore

$$\mathbb{E}\left\{\tau_{c}(X_{i}) \cdot D_{i}\tilde{D}_{i}\right\} = \mathbb{E}\left[\mathbb{E}\left\{\tau_{c}(X_{i}) \cdot D_{i}\tilde{D}_{i} \mid X_{i}\right\}\right] \\
= \mathbb{E}\left\{\tau_{c}(X_{i}) \cdot \alpha(X_{i})\right\} \cdot \mathbb{E}(\tilde{D}_{i}^{2}).$$
(S90)

Plugging (S86), (S89), (S90) into (S82) ensures

$$\mathbb{E}(Y_i \tilde{D}_i) = B + \mathbb{E}\left\{\tau_{c}(X_i) \cdot \alpha(X_i)\right\} \cdot \mathbb{E}(\tilde{D}_i^2),$$

which verifies the expression of $\tau_{*+} = \{\mathbb{E}(\tilde{D}_i^2)\}^{-1}B + \mathbb{E}\{\alpha(X_i) \cdot \tau_c(X_i)\}$ by (S79).

The sufficient conditions for B = 0 are straightforward. We verify below the sufficient conditions for $\alpha(X_i) = w(X_i)$.

Conditions for $\alpha(X_i) = w(X_i)$. Assume that $\mathbb{E}(\check{D}_i \mid X_i)$ is linear in X_i . We have $\mathbb{E}(\check{D}_i \mid X_i) = \operatorname{proj}(\check{D}_i \mid X_i)$ with

$$\tilde{D}_i = \check{D}_i - \mathbb{E}(\check{D}_i \mid X_i). \tag{S91}$$

This, together with $\tilde{D}_i = \check{D}_i - \text{proj}(\check{D}_i \mid X_i)$ being a function of (X_i, Z_i) , ensures

$$\mathbb{E}(\tilde{D}_{i}^{2} \mid X_{i}) \stackrel{(\text{S91})}{=} \mathbb{E}\left[\left\{\check{D}_{i} - \mathbb{E}(\check{D}_{i} \mid X_{i})\right\}^{2} \mid X_{i}\right] = \operatorname{var}(\check{D}_{i} \mid X_{i}), \quad (\text{S92})$$
$$\mathbb{E}(D_{i}\tilde{D}_{i} \mid Z_{i}, X_{i}) = \mathbb{E}(D_{i} \mid Z_{i}, X_{i}) \cdot \tilde{D}_{i}$$

$$\stackrel{(S91)}{=} \mathbb{E}(D_i \mid Z_i, X_i) \cdot \check{D}_i - \mathbb{E}(D_i \mid Z_i, X_i) \cdot \mathbb{E}(\check{D}_i \mid X_i).$$
(S93)

From (S93), we have

$$\mathbb{E}(D_{i}\tilde{D}_{i} \mid X_{i}) = \mathbb{E}\left\{\mathbb{E}(D_{i}\tilde{D}_{i} \mid Z_{i}, X_{i}) \mid X_{i}\right\} \\
= \mathbb{E}\left\{\mathbb{E}(D_{i} \mid Z_{i}, X_{i}) \cdot \check{D}_{i} \mid X_{i}\right\} - \mathbb{E}\left\{\mathbb{E}(D_{i} \mid Z_{i}, X_{i}) \cdot \mathbb{E}(\check{D}_{i} \mid X_{i}) \mid X_{i}\right\} \\
= \mathbb{E}\left\{\mathbb{E}(D_{i} \mid Z_{i}, X_{i}) \cdot \check{D}_{i} \mid X_{i}\right\} - \mathbb{E}\left\{\mathbb{E}(D_{i} \mid Z_{i}, X_{i}) \mid X_{i}\right\} \cdot \mathbb{E}(\check{D}_{i} \mid X_{i}) \\
= \operatorname{cov}\left\{\mathbb{E}(D_{i} \mid Z_{i}, X_{i}), \check{D}_{i} \mid X_{i}\right\}.$$
(S94)

(S92) and (S94) together ensure

$$\alpha(X_i) = \frac{\mathbb{E}(D_i \tilde{D}_i \mid X_i)}{\mathbb{E}(\tilde{D}_i^2)} \stackrel{(S92)+(S94)}{=} \frac{\operatorname{cov}\left\{\mathbb{E}(D_i \mid Z_i, X_i), \check{D}_i \mid X_i\right\}}{\mathbb{E}\{\operatorname{var}(\check{D}_i \mid X_i)\}}.$$
(S95)

When $\check{D}_i = \mathbb{E}(D_i \mid Z_i, X_i)$, (S95) simplifies to $\alpha(X_i) = \frac{\operatorname{var}\{\mathbb{E}(D_i \mid Z_i, X_i) \mid X_i\}}{\mathbb{E}\left[\operatorname{var}\{\mathbb{E}(D_i \mid Z_i, X_i) \mid X_i\}\right]} = w(X_i).$

S5 Proofs of the results in Section S2.4

S5.1 Lemmas

Lemma S10. Let G be an $m \times m$ matrix. Let $r = \operatorname{rank}(G)$ denote the rank of G with $0 \le r \le m$. Then there exists an invertible $m \times m$ matrix

$$\Gamma = \begin{pmatrix} \Gamma_{11} & \Gamma_{12} \\ I_{m-r} & \Gamma_{22} \end{pmatrix}_{m \times m}$$

with $\Gamma_{11} \in \mathbb{R}^{r \times (m-r)}$, $\Gamma_{12} \in \mathbb{R}^{r \times r}$, and $\Gamma_{22} \in \mathbb{R}^{(m-r) \times r}$ such that

$$\Gamma G = \begin{pmatrix} I_r & C \\ 0 & 0 \end{pmatrix}_{m \times m} \quad \text{for some } C \in \mathbb{R}^{r \times (m-r)}.$$

Proof of Lemma S10. The results for r = 0 and r = m are straightforward by letting $\Gamma = I_m$ and $\Gamma = G^{-1}$, respectively. We verify below the result for 0 < r < m.

Given rank(G) = r, the theory of reduced row echelon form (see, e.g., Strang (2006)) ensures that there exists an invertible $m \times m$ matrix Γ' such that

$$\Gamma'G = \begin{pmatrix} I_r & C\\ 0 & 0 \end{pmatrix}_{m \times m} \quad \text{for some } C \in \mathbb{R}^{r \times (m-r)}.$$
(S96)

Write

$$\Gamma' = \begin{pmatrix} \Gamma_1 \\ \Gamma'_2 \end{pmatrix} \quad \text{with } \Gamma_1 \in \mathbb{R}^{r \times m} \text{ and } \Gamma'_2 \in \mathbb{R}^{(m-r) \times m}.$$
(S97)

That Γ' is invertible ensures Γ'_2 has full row rank with $\operatorname{rank}(\Gamma'_2) = m - r$. The theory of reduced row echelon form ensures that there exists an invertible $(m - r) \times (m - r)$ matrix Γ_3 such that

$$\Gamma_3 \Gamma'_2 = (I_{m-r}, \ \Gamma_{22}) \quad \text{for some } (m-r) \times r \text{ matrix } \Gamma_{22}.$$
(S98)

Define

$$\Gamma = \begin{pmatrix} I_r & 0\\ 0 & \Gamma_3 \end{pmatrix} \Gamma'.$$
(S99)

Let $(\Gamma_{11}, \Gamma_{12})$ denote a partition of Γ_1 with $\Gamma_{11} \in \mathbb{R}^{r \times (m-r)}$ and $\Gamma_{12} \in \mathbb{R}^{r \times r}$. It follows from (S97)–(S98) that

$$\Gamma \stackrel{(S97)}{=} \begin{pmatrix} I_r & 0\\ 0 & \Gamma_3 \end{pmatrix} \begin{pmatrix} \Gamma_1\\ \Gamma_2' \end{pmatrix} = \begin{pmatrix} \Gamma_1\\ \Gamma_3\Gamma_2' \end{pmatrix} \stackrel{(S98)}{=} \begin{pmatrix} \Gamma_{11} & \Gamma_{12}\\ I_{m-r} & \Gamma_{22} \end{pmatrix}$$
(S100)

with

$$\Gamma G \stackrel{(S99)}{=} \begin{pmatrix} I_r & 0\\ 0 & \Gamma_3 \end{pmatrix} \Gamma' G \stackrel{(S96)}{=} \begin{pmatrix} I_r & 0\\ 0 & \Gamma_3 \end{pmatrix} \begin{pmatrix} I_r & C\\ 0 & 0 \end{pmatrix} = \begin{pmatrix} I_r & C\\ 0 & 0 \end{pmatrix}.$$

Lemma S11. Assume the setting of Proposition S4. Let

 $\mathcal{S} = \{ X = (X_0, X_1, \dots, X_m)^\top \in \mathbb{R}^{m+1} : X \text{ satisfies (S27)} \}$

denote the solution set of (S27). Let

$$\mathcal{S}_0 = \{X_0 : \text{there exists } X = (X_0, X_1, \dots, X_m)^\top \in \mathcal{S}\}$$

denote the set of values of X_0 across all solutions to (S27). Let $A_0 = (a_{01}, \ldots, a_{0m}) \in \mathbb{R}^{1 \times m}$ denote the coefficient vector of $(X_1, \ldots, X_m)^{\top}$ in the first equation in (S27). Let $A = (a_{ij})_{i,j=1,\ldots,m} \in \mathbb{R}^{m \times m}$ denote the coefficient matrix of $(X_1, \ldots, X_m)^{\top}$ in the last m equations in (S27). Let $\sigma(A)$ denote the set of eigenvalues of A. Let

$$\mathcal{S}_{\overline{\sigma(A)}} = \{ X = (X_0, X_1, ..., X_m)^\top \in \mathcal{S} : X_0 \notin \sigma(A) \}$$

denote the set of solutions to (S27) that have X_0 not equal to any eigenvalue of A. Then

(i)
$$|\mathcal{S}_0| \le 2$$
 if $A_0 = 0_{1 \times m}$.

(ii) Each $X_0 \in \mathcal{S}_0 \setminus \sigma(A)$ corresponds to a unique solution to (S27) with

$$|\mathcal{S}_{\overline{\sigma(A)}}| = |\mathcal{S}_0 \setminus \sigma(A)| \le \begin{cases} 2 & \text{if } A_0 = 0_{1 \times m}; \\ m+2 & \text{if } A_0 \neq 0_{1 \times m}. \end{cases}$$

(iii) Further assume that $S_0 \cap \sigma(A) \neq \emptyset$. Let $\lambda_0 \in S_0 \cap \sigma(A)$ denote an eigenvalue of A that is also in S_0 . Let $r = \operatorname{rank}(\lambda_0 I - A)$ denote the rank of $\lambda_0 I - A$. Let

$$\mathcal{S}_{\lambda_0} = \{ X = (X_0, X_1, \dots, X_m)^\top \in \mathcal{S} : X_0 = \lambda_0 \},\$$
$$\mathcal{S}_{\overline{\lambda_0}} = \{ X = (X_0, X_1, \dots, X_m)^\top \in \mathcal{S} : X_0 \neq \lambda_0 \}$$

denote a partition of S, containing solutions to (S27) that have $X_0 = \lambda_0$ and $X_0 \neq \lambda_0$, respectively.

- (a) If $A \neq \lambda_0 I$, then 0 < r < m and there exists constant $g \in \mathbb{R}^r$, $G \in \mathbb{R}^{r \times (m-r)}$, $h \in \mathbb{R}^{m-r}$, and $H \in \mathbb{R}^{(m-r) \times r}$ such that
 - for all $X = (X_0, X_1, \dots, X_m)^\top \in \mathcal{S}_{\lambda_0}$ with $X_0 = \lambda_0$, the corresponding $X_{1:r} = (X_1, \dots, X_r)^\top$ and $X_{(r+1):m} = (X_{r+1}, \dots, X_m)^\top$ satisfy

$$X_{1:r} = g + GX_{(r+1):m};$$

- for all $X = (X_0, X_1, \dots, X_m)^\top \in \mathcal{S}_{\overline{\lambda_0}}$ with $X_0 \neq \lambda_0$, the corresponding $X_{1:(m-r)} = (X_1, \dots, X_{m-r})^\top$ and $X_{(m-r+1):m} = (X_{m-r+1}, \dots, X_m)^\top$ satisfy

$$X_{1:(m-r)} = h + HX_{(m-r+1):m}$$

(b) If $A = \lambda_0 I$, then

 $- \text{ for all } X = (X_0, X_1, \dots, X_m)^\top \in \mathcal{S}_{\overline{\lambda_0}} \text{ with } X_0 \neq \lambda_0, \text{ we have } (X_1, \dots, X_m) = (b_1, \dots, b_m).$ $- |\mathcal{S}_{\overline{\lambda_0}}| = |\mathcal{S}_0 \setminus \{\lambda_0\}| \leq \begin{cases} 1 & \text{if } A_0 = 0_{1 \times m}; \\ 2 & \text{if } A_0 \neq 0_{1 \times m}. \end{cases}$ $- |\mathcal{S}_0| \leq \begin{cases} 2 & \text{if } A_0 = 0_{1 \times m}; \\ 3 & \text{if } A_0 \neq 0_{1 \times m}. \end{cases}$

Proof of Lemma S11. Let $U = (X_1, ..., X_m)^{\top}$, $c = (c_1, ..., c_m)^{\top}$, $b = (b_1, ..., b_m)^{\top}$, and

$$l(t) = c + bt = \begin{pmatrix} c_1 + b_1 t \\ \vdots \\ c_m + b_m t \end{pmatrix} \in \mathbb{R}^m$$
(S101)

to write the last m equations in (S27) as

$$UX_0 = l(X_0) + AU. (S102)$$

Write the first equation in (S27) as

$$X_0^2 - b_0 X_0 - c_0 = A_0 U. (S103)$$

We verify below Lemma S11(i)–(iii) one by one.

Proof of Lemma S11(i):

When $A_0 = 0$, (S103) reduces to $X_0^2 - b_0 X_0 - c_0 = 0$. This ensures X_0 takes at most 2 distinct values across all solutions to (S27).

Proof of Lemma S11(ii):

Write (S102) as

$$(X_0 I - A)U = l(X_0). (S104)$$

For $X_0 \in \mathcal{S}_0 \setminus \sigma(A)$ that is not an eigenvalue of A, we have $X_0I - A$ is invertible with

$$(X_0 I - A)^{-1} = \frac{1}{\det(X_0 I - A)} \cdot \operatorname{Adj}(X_0 I - A),$$
(S105)

where $det(\cdot)$ and $Adj(\cdot)$ denote the determinant and adjoint matrix, respectively. (S104)–(S105) together ensure that

for all solutions
$$X = (X_0, X_1, \dots, X_m)^{\top}$$
 to (S27) that have $X_0 \notin \sigma(A)$,
 $U = (X_1, \dots, X_m)^{\top}$ is uniquely determined by X_0 as
$$U \stackrel{(S104)}{=} (X_0I - A)^{-1} \cdot l(X_0) \stackrel{(S105)}{=} \frac{1}{\det(X_0I - A)} \cdot \operatorname{Adj}(X_0I - A) \cdot l(X_0).$$
(S106)

(S106) ensures that

each $X_0 \in \mathcal{S}_0 \setminus \sigma(A)$ corresponds to a unique solution to (S27) with $|\mathcal{S}_{\overline{\sigma(A)}}| = |\mathcal{S}_0 \setminus \sigma(A)|$.

It remains to show that

$$|\mathcal{S}_0 \setminus \sigma(A)| \le \begin{cases} 2 & \text{if } A_0 = 0_{1 \times m}; \\ m+2 & \text{if } A_0 \neq 0_{1 \times m}. \end{cases}$$
(S107)

When $A_0 = 0_{1 \times m}$, Lemma S11(i) ensures $|\mathcal{S}_0 \setminus \sigma(A)| \leq |\mathcal{S}_0| \leq 2$. When $A_0 \neq 0_{1 \times m}$, plugging (S106) in (S103) ensures

$$X_0^2 - b_0 X_0 - c_0 = \frac{1}{\det(X_0 I - A)} \cdot A_0 \cdot \operatorname{Adj}(X_0 I - A) \cdot l(X_0),$$

and therefore

$$\det(X_0I - A) \cdot (X_0^2 - b_0X_0 - c_0) = A_0 \cdot \operatorname{Adj}(X_0I - A) \cdot l(X_0).$$
(S108)

Note that $det(X_0I - A)$ is at most *m*th-degree in X_0 and the elements of $Adj(X_0I - A)$ are at most (m - 1)th-degree in X_0 . This, together with $l(X_0) = c + bX_0$, ensures (S108) is at most (m + 2)th-degree in X_0 . By the fundamental theorem of algebra, X_0 takes at most m + 2 distinct values beyond the eigenvalues of A. This verifies (S107).

Proof of Lemma S11(iii):

From (S101)–(S102), direct algebra ensures that

for all $X \in \mathcal{S}$, we have

$$(U-b)(X_0 - \lambda_0) = UX_0 - \lambda_0 U - bX_0 + b\lambda_0$$

$$\stackrel{(S101)+(S102)}{=} (c + bX_0 + AU) - \lambda_0 U - bX_0 + b\lambda_0$$

$$= (c + b\lambda_0) + AU - \lambda_0 U$$

$$\stackrel{(S101)}{=} l(\lambda_0) - (\lambda_0 I - A)U. \qquad (S109)$$

We verify below the result for $A \neq \lambda_0 I$ and $A = \lambda_0 I$, respectively.

Proof of Lemma S11(iiia) for $A \neq \lambda_0 I$:

Recall $r = \operatorname{rank}(\lambda_0 I - A)$. When $A \neq \lambda_0 I$, we have 0 < r < m. From Lemma S10, there exists an invertible $m \times m$ matrix

$$\Gamma = \begin{pmatrix} \Gamma_{11} & \Gamma_{12} \\ I_{m-r} & \Gamma_{22} \end{pmatrix} = \begin{pmatrix} \Gamma_1 \\ \Gamma_2 \end{pmatrix}$$
(S110)

with $\Gamma_{11} \in \mathbb{R}^{r \times (m-r)}$, $\Gamma_{12} \in \mathbb{R}^{r \times r}$, $\Gamma_{22} \in \mathbb{R}^{(m-r) \times r}$, $\Gamma_1 = (\Gamma_{11}, \Gamma_{12})$, and $\Gamma_2 = (I_{m-r}, \Gamma_{22})$ such that

$$\Gamma(\lambda_0 I - A) = \begin{pmatrix} I_r & C \\ 0 & 0 \end{pmatrix} \quad \text{for some } C \in \mathbb{R}^{r \times (m-r)}.$$
(S111)

Multiplying both sides of (S109) by Γ ensures that

for all
$$X \in \mathcal{S}$$
, we have

$$\Gamma(U-b)(X_0 - \lambda_0) = \Gamma \cdot l(\lambda_0) - \Gamma(\lambda_0 I - A)U.$$
(S112)

We simplify below the left- and right-hand sides of (S112), respectively.

Write

$$U = (X_1, \dots, X_m)^{\top} = (X_{1:r}^{\top}, X_{(r+1):m}^{\top})^{\top} = (X_{1:(m-r)}^{\top}, X_{(m-r+1):m}^{\top})^{\top}$$
(S113)

with

$$X_{1:r} = (X_1, \dots, X_r)^{\top}, \qquad X_{(r+1):m} = (X_{r+1}, \dots, X_m)^{\top};$$
$$X_{1:(m-r)} = (X_1, \dots, X_{m-r})^{\top}, \qquad X_{(m-r+1):m} = (X_{m-r+1}, \dots, X_m)^{\top}.$$

This, together with (S110)–(S111), ensures

$$\Gamma(\lambda_0 I - A) U \stackrel{(\text{S111})+(\text{S113})}{=} \begin{pmatrix} I_r & C \\ 0 & 0 \end{pmatrix} \begin{pmatrix} X_{1:r} \\ X_{(r+1):m} \end{pmatrix} = \begin{pmatrix} X_{1:r} + C X_{(r+1):m} \\ 0 \end{pmatrix}, \quad (\text{S114})$$

and

$$\Gamma_{2}(U-b) = \Gamma_{2}U - \Gamma_{2}b$$

$$\stackrel{(S110)+(S113)}{=} (I_{m-r}, \Gamma_{22}) \begin{pmatrix} X_{1:(m-r)} \\ X_{(m-r+1):m} \end{pmatrix} - \Gamma_{2}b$$

$$= X_{1:(m-r)} + \Gamma_{22}X_{(m-r+1):m} - \Gamma_{2}b.$$
(S115)

Given (S110)–(S115), the left-hand side of (S112) equals

$$\Gamma(U-b)(X_{0}-\lambda_{0}) \stackrel{(S110)}{=} \begin{pmatrix} \Gamma_{1} \\ \Gamma_{2} \end{pmatrix} (U-b)(X_{0}-\lambda_{0}) \\ = \begin{pmatrix} \Gamma_{1}(U-b)(X_{0}-\lambda_{0}) \\ \Gamma_{2}(U-b)(X_{0}-\lambda_{0}) \end{pmatrix} \\ \stackrel{(S115)}{=} \begin{pmatrix} \Gamma_{1}(U-b)(X_{0}-\lambda_{0}) \\ (X_{1:(m-r)}+\Gamma_{22}X_{(m-r+1):m}-\Gamma_{2}b)(X_{0}-\lambda_{0}) \end{pmatrix}, (S116)$$

and the right-hand side of (S112) equals

$$\Gamma \cdot l(\lambda_0) - \Gamma(\lambda_0 I - A) U \stackrel{(S110)+(S114)}{=} \begin{pmatrix} \Gamma_1 \\ \Gamma_2 \end{pmatrix} l(\lambda_0) - \begin{pmatrix} X_{1:r} + CX_{(r+1):m} \\ 0 \end{pmatrix}$$
$$= \begin{pmatrix} \Gamma_1 \cdot l(\lambda_0) - X_{1:r} - CX_{(r+1):m} \\ \Gamma_2 \cdot l(\lambda_0) \end{pmatrix}. \quad (S117)$$

Given (S112), comparing the first and second rows of (S116) and (S117) ensures that

for all $X \in \mathcal{S}$, we have

$$\Gamma_1(U-b)(X_0-\lambda_0) = \Gamma_1 \cdot l(\lambda_0) - X_{1:r} - CX_{(r+1):m},$$
(S118)
$$\left(X_{1:(m-r)} + \Gamma_{22}X_{(m-r+1):m} - \Gamma_2 b\right)(X_0-\lambda_0) = \Gamma_2 \cdot l(\lambda_0).$$
(S119)

The derivation so far, hence (S118)–(S119), only depends on 0 < r < m and holds regardless of whether $\lambda_0 \in S_0$. Now that we know λ_0 is indeed within S_0 , (S118)–(S119) together lead to the following:

• Letting $X_0 = \lambda_0$ in (S118) ensures

for all
$$X = (X_0, X_1, \dots, X_m)^\top \in \mathcal{S}_{\lambda_0}$$
 with $X_0 = \lambda_0$, we have

$$X_{1:r} = \Gamma_1 \cdot l(\lambda_0) - CX_{(r+1):m}.$$
(S120)

This verifies the first part of Lemma S11(iiia) with $g = \Gamma_1 \cdot l(\lambda_0)$ and G = -C.

• Letting $X_0 = \lambda_0$ in (S119) ensures

$$\Gamma_2 \cdot l(\lambda_0) = 0, \tag{S121}$$

as a constraint on the coefficients $\{c_i, b_i : i = 1, ..., m\}$; c.f. $l(\lambda_0) = c + b\lambda_0$ by (S101). Plugging (S121) back in (S119) ensures that

for all
$$X \in \mathcal{S}$$
, we have

$$\left(X_{1:(m-r)} + \Gamma_{22}X_{(m-r+1):m} - \Gamma_{2}b\right)(X_{0} - \lambda_{0}) = 0.$$
(S122)

From (S122), we have

$$X_{1:(m-r)} = \Gamma_2 b - \Gamma_{22} X_{(m-r+1):m} \quad \text{for all } X \in \mathcal{S}_{\overline{\lambda_0}} \text{ with } X_0 \neq \lambda_0.$$
(S123)

This verifies the second part of Lemma S11(iiia) with $h = \Gamma_2 b$ and $H = -\Gamma_{22}$.

Proof of Lemma S11(iiib) for $A = \lambda_0 I$:

When $A = \lambda_0 I$, (S109) simplifies to

for all
$$X \in \mathcal{S}$$
, we have $(U - b)(X_0 - \lambda_0) = l(\lambda_0)$. (S124)

Given $\lambda_0 \in \mathcal{S}_0$ by assumption, letting $X_0 = \lambda_0$ in (S124) ensures

$$l(\lambda_0) = 0, \tag{S125}$$

as a constraint on $\{c_i, b_i : i = 1, ..., m\}$ analogous to (S121). Plugging (S125) back in (S124) ensures

for all
$$X \in \mathcal{S}$$
, we have $(U - b)(X_0 - \lambda_0) = 0$, (S126)

analogous to (S122). This ensures

for all
$$X = (X_0, X_1, \dots, X_m)^{\top} \in \mathcal{S}_{\overline{\lambda_0}}$$
 with $X_0 \neq \lambda_0$, we have $U = b$, (S127)

analogous to (S123).

In addition, that $A = \lambda_0 I$ implies $\sigma(A) = \{\lambda_0\}$ with $S_{\overline{\lambda_0}} = S_{\overline{\sigma(A)}}$. This, together with Lemma S11(ii), ensures

$$|\mathcal{S}_{\overline{\lambda_0}}| = |\mathcal{S}_{\overline{\sigma(A)}}| = |\mathcal{S}_0 \setminus \sigma(A)| = |\mathcal{S}_0 \setminus \{\lambda_0\}|.$$

When $A_0 = 0_{1 \times m}$, Lemma S11(i) ensures $|\mathcal{S}_0| \leq 2$ such that $|\mathcal{S}_0 \setminus \{\lambda_0\}| = |\mathcal{S}_0| - 1 \leq 1$. When $A_0 \neq 0_{1 \times m}$, plugging (S127) in (S103) ensures that

for all
$$X = (X_0, X_1, \dots, X_m)^\top \in \mathcal{S}_{\overline{\lambda_0}}$$
 with $X_0 \neq \lambda_0$, we have $X_0^2 - b_0 X_0 - c_0 = A_0 b$.

The fundamental theorem of algebra ensures that X_0 takes at most 2 distinct values beyond λ_0 , i.e., $|\mathcal{S}_0 \setminus {\lambda_0}| \leq 2$. The bound for $|\mathcal{S}_0|$ then follows from $|\mathcal{S}_0| = |\mathcal{S}_0 \setminus {\lambda_0}| + 1$.

Remark S1. From Lemma S10, the definition of Γ in (S110) reduces to $\Gamma = I_m = \Gamma_2$ when r = 0. This underlies the correspondence between (S125)–(S127) and (S121)–(S123), and ensures that the proof for Lemma S11(iiia) implies the proof for Lemma S11(iiib) as a special case. We nonetheless provided a separate proof to add clarity.

S5.2 Proof of Proposition S4

We verify below Proposition S4 by induction.

<u>Proof of the result for m = 1.</u>

When m = 1, (S27) suggests

$$X_0^2 = c_0 + b_0 X_0 + a_{01} X_1, (S128)$$

$$X_1 X_0 = c_1 + b_1 X_0 + a_{11} X_1 \tag{S129}$$

for fixed constants $\{c_i, b_i, a_{i1} : i = 0, 1\}$. If $a_{01} = 0$, then (S128) reduces to $X_0^2 = c_0 + b_0 X_0$ so that X_0 takes at most 2 distinct values. If $a_{01} \neq 0$, then (S128) implies $X_1 = a_{01}^{-1}(X_0^2 - b_0 X_0 - c_0)$. Plugging this in (S129) yields

$$a_{01}^{-1}(X_0^2 - b_0X_0 - c_0)X_0 = c_1 + b_1X_0 + a_{11}a_{01}^{-1}(X_0^2 - b_0X_0 - c_0),$$

as a cubic equation of X_0 . The fundamental theorem of algebra ensures that X_0 takes at most 3 distinct values. This verifies the result for m = 1.

Proof of the result for $m \ge 2$.

Assume that the result holds for m = 1, ..., Q - 1. We verify below the result for m = Q, where (S27) becomes

$$\begin{pmatrix} X_0 \\ X_1 \\ \vdots \\ X_Q \end{pmatrix} X_0 = \begin{pmatrix} c_0 \\ c_1 \\ \vdots \\ c_Q \end{pmatrix} + \begin{pmatrix} b_0 \\ b_1 \\ \vdots \\ b_Q \end{pmatrix} X_0 + \begin{pmatrix} a_{01} & \dots & a_{0Q} \\ a_{11} & \dots & a_{1Q} \\ \vdots \\ a_{Q1} & \dots & a_{QQ} \end{pmatrix} \begin{pmatrix} X_1 \\ \vdots \\ X_Q \end{pmatrix}.$$
(S130)

Following Lemma S11, renew

$$\mathcal{S} = \{X = (X_0, X_1, \dots, X_Q)^\top \in \mathbb{R}^{Q+1} : X \text{ satisfies (S130)}\},\$$

$$\mathcal{S}_0 = \{X_0 : \text{there exists } X = (X_0, X_1, \dots, X_Q)^\top \in \mathcal{S}\}.$$

The goal is to show that $|\mathcal{S}_0| \leq Q+2$.

First, let
$$U = (X_1, \dots, X_Q)^{\top}$$
, $c = (c_1, \dots, c_Q)^{\top}$, $b = (b_1, \dots, b_Q)^{\top}$, $A_0 = (a_{01}, \dots, a_{0Q}) \in$

 $\mathbb{R}^{1 \times Q}$, and $A = (a_{qq'})_{q,q'=1,\dots,Q} \in \mathbb{R}^{Q \times Q}$ to write (S130) as

$$\begin{pmatrix} X_0 \\ U \end{pmatrix} X_0 = \begin{pmatrix} c_0 \\ c \end{pmatrix} + \begin{pmatrix} b_0 \\ b \end{pmatrix} X_0 + \begin{pmatrix} A_0 \\ A \end{pmatrix} U.$$
(S131)

Let $\sigma(A)$ denote the set of eigenvalues of A. We have the following observations from Lemma S11:

- If $S_0 \cap \sigma(A) = \emptyset$ so that X_0 does not equal any eigenvalue of A across all solutions to (S130), then $S_0 = S_0 \setminus \sigma(A)$, and it follows from Lemma S11(ii) that $|S_0| = |S_0 \setminus \sigma(A)| \le Q + 2$.
- If $S_0 \cap \sigma(A) \neq \emptyset$ and there exists $\lambda \in S_0 \cap \sigma(A)$ so that $A = \lambda_0 I$, then it follows from Lemma S11(iiib) that $|S_0| \leq 3 \leq Q+2$.

Therefore, it remains to show that $|\mathcal{S}_0| \leq Q + 2$ when

$$\mathcal{S}_0 \cap \sigma(A) \neq \emptyset$$
 and $A \neq \lambda I$ for all $\lambda \in \mathcal{S}_0 \cap \sigma(A)$. (S132)

Proof of $|\mathcal{S}_0| \leq Q + 2$ under (S132).

Let λ_0 denote an element in $S_0 \cap \sigma(A)$ with $A \neq \lambda_0 I$. Let $r = \operatorname{rank}(\lambda_0 I - A)$ denote the rank of $\lambda_0 I - A$ with 0 < r < Q. Let

$$X_{1:(Q-r)} = (X_1, \dots, X_{Q-r})^{\top}, \quad X_{(Q-r+1):Q} = (X_{Q-r+1}, \dots, X_Q)^{\top}$$

denote a partition of $U = (X_1, \ldots, X_Q)^{\top}$. The last r equations in (S131) equal

$$\begin{aligned} X_{(Q-r+1):Q}X_{0} &= \begin{pmatrix} X_{Q-r+1} \\ \vdots \\ X_{Q} \end{pmatrix} X_{0} \\ &= \begin{pmatrix} c_{Q-r+1} \\ \vdots \\ c_{Q} \end{pmatrix} + \begin{pmatrix} b_{Q-r+1} \\ \vdots \\ b_{Q} \end{pmatrix} X_{0} + \begin{pmatrix} a_{Q-r+1,1} & \dots & a_{Q-r+1,Q} \\ & \ddots & \\ a_{Q1} & \dots & a_{QQ} \end{pmatrix} U \\ &= c_{(Q-r+1):Q} + b_{(Q-r+1):Q}X_{0} + A_{(Q-r+1):Q}U, \end{aligned}$$
(S133)

where

$$c_{(Q-r+1):Q} = \begin{pmatrix} c_{Q-r+1} \\ \vdots \\ c_Q \end{pmatrix}, \quad b_{(Q-r+1):Q} = \begin{pmatrix} b_{Q-r+1} \\ \vdots \\ b_Q \end{pmatrix}, \quad A_{(Q-r+1):Q} = \begin{pmatrix} a_{Q-r+1,1} & \dots & a_{Q-r+1,Q} \\ & \ddots & & \\ a_{Q1} & \dots & a_{QQ} \end{pmatrix}$$

•

In addition, let

$$\mathcal{S}_{\overline{\lambda_0}} = \{ X = (X_0, X_1, \dots, X_Q)^\top \in \mathcal{S} : X_0 \neq \lambda_0 \}.$$

Lemma S11(iii) ensures that there exists constant $h \in \mathbb{R}^{Q-r}$ and $H \in \mathbb{R}^{(Q-r) \times r}$ such that

for all
$$X = (X_0, X_1, \dots, X_Q)^\top \in \mathcal{S}_{\overline{\lambda_0}}$$
 with $X_0 \neq \lambda_0$, we have

$$X_{1:(Q-r)} = h + HX_{(Q-r+1):Q}.$$
(S134)

Consequently, we have

for all
$$X = (X_0, X_1, \dots, X_Q)^\top \in \mathcal{S}_{\overline{\lambda_0}}$$
 with $X_0 \neq \lambda_0$, the corresponding U satisfies

$$U = \begin{pmatrix} X_1 \\ \vdots \\ X_Q \end{pmatrix} = \begin{pmatrix} X_{1:(Q-r)} \\ X_{(Q-r+1):Q} \end{pmatrix} \stackrel{(S134)}{=} \begin{pmatrix} h + HX_{(Q-r+1):Q} \\ X_{(Q-r+1):Q} \end{pmatrix} = \begin{pmatrix} h \\ 0 \end{pmatrix} + \begin{pmatrix} H \\ I_r \end{pmatrix} X_{(Q-r+1):Q}.$$
(S135)

(S131)-(S135) together ensure that

for all $X = (X_0, X_1, \dots, X_Q)^\top \in \mathcal{S}_{\overline{\lambda_0}}$ with $X_0 \neq \lambda_0$, we have

$$\begin{pmatrix} X_0 \\ X_{(Q-r+1):Q} \end{pmatrix} X_0 \stackrel{(S131)+(S133)}{=} \begin{pmatrix} c_0 \\ c_{(Q-r+1):Q} \end{pmatrix} + \begin{pmatrix} b_0 \\ b_{(Q-r+1):Q} \end{pmatrix} X_0 + \begin{pmatrix} A_0 \\ A_{(Q-r+1):Q} \end{pmatrix} U$$

$$\stackrel{(S135)}{=} \begin{pmatrix} c_0 \\ c_{(Q-r+1):Q} \end{pmatrix} + \begin{pmatrix} b_0 \\ b_{(Q-r+1):Q} \end{pmatrix} X_0 \qquad (S136)$$

$$+ \begin{pmatrix} A_0 \\ A_{(Q-r+1):Q} \end{pmatrix} \cdot \left\{ \begin{pmatrix} h \\ 0 \end{pmatrix} + \begin{pmatrix} H \\ I_r \end{pmatrix} X_{(Q-r+1):Q} \right\}.$$

for all solutions $X = (X_0, X_1, \ldots, X_Q)^{\top}$ to (S130) that have $X_0 \neq \lambda_0$, the corresponding $X' = (X_0, X_{(Q-r+1):Q}^{\top})^{\top} = (X_0, X_{Q-r+1}, \ldots, X_Q)^{\top}$ is an $(r+1) \times 1$ vector that satisfies $X'X_0$ is linear in (1, X') with $r \leq Q - 1$.

By induction, applying Proposition S4 to $X' = (X_0, X_{(Q-r+1):Q}^{\top})^{\top}$ at $m = r (\leq Q - 1)$ ensures that

across all solutions $X = (X_0, X_1, \dots, X_Q)^{\top}$ to (S130) that have $X_0 \neq \lambda_0$, X_0 takes at most r + 2 distinct values, where $r \leq Q - 1$.

This ensures $|\mathcal{S}_0 \setminus \{\lambda_0\}| \le r+2 \le Q+1$ so that $|\mathcal{S}_0| = |\mathcal{S}_0 \setminus \{\lambda_0\}| + 1 \le Q+2$.

S5.3 Proof of Proposition S5

We verify below Proposition S5 by induction.

<u>Proof of the result for m = 1.</u>

Assume $X = X_1 \in \mathbb{R}$ is a scalar such that $XX^{\top} = X_1^2$ is linear in $(1, X) = (1, X_1)$. Then there exists constants $a_0, a_1 \in \mathbb{R}$ such that $X_1^2 = a_0 + a_1X_1$. By the fundamental theorem of algebra, X_1 takes at most 2 distinct values.

Proof of the result for $m \ge 2$.

Assume that the result holds for m = 1, ..., Q. We verify below the result for m = Q + 1. Assume $X = (X_1, ..., X_Q, X_{Q+1})^{\top}$ is a $(Q+1) \times 1$ vector such that all elements of

$$XX^{\top} = \begin{pmatrix} X_1X_1 & \dots & X_1X_Q & X_1X_{Q+1} \\ \vdots & \vdots & \vdots \\ X_QX_1 & \dots & X_QX_Q & X_QX_{Q+1} \\ \hline X_{Q+1}X_1 & \dots & X_{Q+1}X_Q & X_{Q+1}X_{Q+1} \end{pmatrix}$$
(S137)

are linear in (1, X) with

$$X_q X_{q'} = c_{qq'} + \sum_{k=1}^{Q+1} a_{qq'[k]} X_k \quad \text{for } q, q' = 1, \dots, Q+1$$
(S138)

for constants $\{c_{qq'}, a_{qq'[k]} \in \mathbb{R} : q, q', k = 1, \dots, Q + 1\}$. Without loss of generality, assume the coefficients for $X_q X_{q'}$ and $X_{q'} X_q$ are identical with

$$c_{qq'} = c_{q'q}, \quad a_{qq'[k]} = a_{q'q[k]} \quad \text{for} \quad q, q', k = 1, \dots, Q+1.$$
 (S139)

Let

$$\mathcal{S} = \{ X = (X_1, \dots, X_{Q+1})^\top \in \mathbb{R}^{Q+1} : X \text{ satisfies (S138)} \}$$

denote the solution set of (S138). To goal is to show that $|\mathcal{S}| \leq Q + 2$.

To this end, consider the last column of XX^{\top} in (S137) that consists of $X_1X_{Q+1}, \ldots, X_QX_{Q+1}, X_{Q+1}^2$. The corresponding part in (S138) is

$$X_q X_{Q+1} = c_{q,Q+1} + \sum_{k=1}^{Q+1} a_{q,Q+1[k]} X_k$$
 for $q = 1, \dots, Q+1$,

and can be summarized in matrix form as

$$\begin{pmatrix}
X_{Q+1} \\
X_1 \\
\vdots \\
X_Q
\end{pmatrix} X_{Q+1} = \begin{pmatrix}
c_{Q+1,Q+1} \\
c_{1,Q+1} \\
\vdots \\
c_{Q,Q+1}
\end{pmatrix} + \begin{pmatrix}
a_{Q+1,Q+1[Q+1]} \\
a_{1,Q+1[Q+1]} \\
\vdots \\
a_{Q,Q+1[Q+1]}
\end{pmatrix} X_{Q+1} \\
+ \begin{pmatrix}
\frac{a_{Q+1,Q+1[1]} & \cdots & a_{Q+1,Q+1[Q]} \\
a_{1,Q+1[1]} & \cdots & a_{1,Q+1[Q]} \\
\vdots \\
a_{Q,Q+1[1]} & \cdots & a_{Q,Q+1[Q]}
\end{pmatrix} \begin{pmatrix}
X_1 \\
\vdots \\
X_Q
\end{pmatrix}.$$
(S140)

Let

$$A = \begin{pmatrix} a_{1,Q+1[1]} & \dots & a_{1,Q+1[Q]} \\ & \ddots & \\ a_{Q,Q+1[1]} & \dots & a_{Q,Q+1[Q]} \end{pmatrix} \in \mathbb{R}^{Q \times Q}$$

denote the $Q \times Q$ coefficient matrix of (X_1, \ldots, X_Q) in the last Q rows of (S140). Let $\sigma(A)$ denote the set of eigenvalues of A. Let

$$\mathcal{S}_{Q+1} = \{X_{Q+1} : \text{there exists } X = (X_1, \dots, X_{Q+1})^\top \in \mathcal{S}\}$$

denote the set of values of X_{Q+1} across all solutions to (S138). Note that all solutions to (S138) must satisfy (S140). Applying Proposition S4 and Lemma S11(ii) to (S140) with X_{Q+1} treated as X_0 ensures

- (i) $|\mathcal{S}_{Q+1}| \leq Q+2$ by Proposition S4.
- (ii) If $S_{Q+1} \cap \sigma(A) = \emptyset$ such that X_{Q+1} does not equal any eigenvalue of A across all solutions to (S138), then $S_{Q+1} = S_{Q+1} \setminus \sigma(A)$, and it follows from Lemma S11(ii) that $|S| = |S_{Q+1} \setminus \sigma(A)| = |S_{Q+1}|.$

These two observations together ensure

$$|\mathcal{S}| = |\mathcal{S}_{Q+1}| \le Q+2$$
 when $\mathcal{S}_{Q+1} \cap \sigma(A) = \emptyset$.

We show below $|\mathcal{S}| \leq Q + 2$ when $\mathcal{S}_{Q+1} \cap \sigma(A) \neq \emptyset$.

Let λ_0 denote an element in $\mathcal{S}_{Q+1} \cap \sigma(A)$, as an eigenvalue of A that X_{Q+1} takes in at least one solution to (S138). We verify below $|\mathcal{S}| \leq Q + 2$ for the cases of $A = \lambda_0 I$ and $A \neq \lambda_0 I$, respectively. Throughout, let

$$\mathcal{S}_{\lambda_0} = \{ X = (X_1, \dots, X_Q, X_{Q+1})^\top \in \mathcal{S} : X_{Q+1} = \lambda_0 \},\$$
$$\mathcal{S}_{\overline{\lambda_0}} = \{ X = (X_1, \dots, X_Q, X_{Q+1})^\top \in \mathcal{S} : X_{Q+1} \neq \lambda_0 \}$$

denote a partition of S, containing solutions to (S138) that have $X_{Q+1} = \lambda_0$ and $X_{Q+1} \neq \lambda_0$, respectively.

S5.3.1 $|\mathcal{S}| \leq Q + 2$ when $A = \lambda_0 I$.

Let q = q' = Q + 1 in (S138) to see

$$X_{Q+1}^2 = c_{Q+1,Q+1} + a_{Q+1,Q+1[Q+1]} X_{Q+1} + \sum_{k=1}^Q a_{Q+1,Q+1[k]} X_k.$$
 (S141)

Condition S1 below states the condition that the coefficients for X_1, \ldots, X_Q in (S141) all equal zero. We verify below that

$$|\mathcal{S}_{\overline{\lambda_0}}| \leq \begin{cases} 1 & \text{if Condition S1 holds;} \\ 2 & \text{otherwise,} \end{cases}$$
(S142)

$$|\mathcal{S}_{\lambda_0}| \leq \begin{cases} Q+1 & \text{if Condition S1 holds;} \\ Q & \text{otherwise.} \end{cases}$$
(S143)

These two parts together ensure $|\mathcal{S}| = |\mathcal{S}_{\lambda_0}| + |\mathcal{S}_{\overline{\lambda_0}}| \le Q + 2$.

Condition S1. $a_{Q+1,Q+1[k]} = 0$ for k = 1, ..., Q.

Part I: Proof of (S142).

Condition S1 ensures that $A_{Q+1} = (a_{Q+1,Q+1[1]}, \ldots, a_{Q+1,Q+1[Q]})$ equals zero, which corresponds to the condition of $A_0 = 0$ in Lemma S11(iiib). Inequality (S142) follows from Lemma S11(iiib) with X_{Q+1} treated as X_0 .

Part II: Proof of (S143).

First, (S138) ensures

$$X_q X_{q'} = c_{qq'} + a_{qq'[Q+1]} X_{Q+1} + \sum_{k=1}^Q a_{qq'[k]} X_k \quad \text{for } q, q' = 1, \dots, Q.$$
(S144)

Letting $X_{Q+1} = \lambda_0$ in (S144) ensures that

for all solutions $X = (X_1, \ldots, X_Q, X_{Q+1})^\top$ to (S138) that have $X_{Q+1} = \lambda_0$, the corresponding $X' = (X_1, \ldots, X_Q)^\top$ satisfies

$$X_{q}X_{q'} = c_{qq'} + a_{qq'[Q+1]}\lambda_0 + \sum_{k=1}^{Q} a_{qq'[k]}X_k \quad \text{for } q, q' = 1, \dots, Q;$$
(S145)

i.e., $X'X'^{\dagger}$ is component-wise linear in (1, X').

By induction, applying Proposition S5 to $X' = (X_1, \ldots, X_Q)^{\top}$ at m = Q ensures that

across all solutions $X = (X_1, \ldots, X_Q, X_{Q+1})^\top$ to (S138) that have $X_{Q+1} = \lambda_0$, the corresponding $X' = (X_1, \ldots, X_Q)^\top$ take at most Q + 1 distinct values.

This ensures $|S_{\lambda_0}| \leq Q + 1$, and verifies the first line of (S143) when Condition S1 holds. When Condition S1 does not hold, there exists $k \in \{1, \ldots, Q\}$ such that $a_{Q+1,Q+1[k]} \neq 0$. Without loss of generality, assume that $a_{Q+1,Q+1[Q]} \neq 0$. Letting $X_0 = \lambda_0$ in (S141) implies

for all solutions $X = (X_1, \ldots, X_Q, X_{Q+1})^\top$ to (S138) that have $X_{Q+1} = \lambda_0$, the corresponding $X' = (X_1, \ldots, X_Q)^\top$ satisfies

$$\lambda_0^2 = c_{Q+1,Q+1} + a_{Q+1,Q+1[Q+1]}\lambda_0 + \sum_{k=1}^Q a_{Q+1,Q+1[k]}X_k$$
(S146)

with

$$X_Q = a_{Q+1,Q+1[Q]}^{-1} \left(\lambda_0^2 - c_{Q+1,Q+1} - a_{Q+1,Q+1[Q+1]} \lambda_0 - \sum_{k=1}^{Q-1} a_{Q+1,Q+1[k]} X_k \right).$$

Plugging the expression of X_Q in (S146) back in (S145) ensures

for all solutions $X = (X_1, \ldots, X_Q, X_{Q+1})^\top$ to (S138) that $X_{Q+1} = \lambda_0$, the corresponding $X' = (X_1, \ldots, X_{Q-1})^\top$ satisfies

$$X_q X_{q'} = c_{qq'}^* + \sum_{k=1}^{Q-1} a_{qq'[k]}^* X_k \quad \text{for} \quad q, q' = 1, \dots, Q-1,$$

where $\{c_{qq'}^*, a_{qq'[k]}^* : q, q', k = 1, \dots, Q - 1\}$ are constants determined by λ_0 and $\{c_{qq'}, a_{qq'[k]} : q, q', k = 1, \dots, Q + 1\}$; i.e., $X'X'^{\top}$ is component-wise linear in (1, X').

By induction, applying Proposition S5 to $X' = (X_1, \ldots, X_{Q-1})^{\top}$ at m = Q-1 ensures that

across all solutions $X = (X_1, \dots, X_Q, X_{Q+1})^\top$ to (S138) that have $X_{Q+1} = \lambda_0$, the corresponding $X' = (X_1, \dots, X_{Q-1})^\top$ take at most Q distinct values. (S147)

The expression of X_Q in (S146) further ensures that

each solution $X = (X_1, \dots, X_Q, X_{Q+1})^\top$ to (S138) with $X_{Q+1} = \lambda_0$ is uniquely determined by its subvector $X' = (X_1, \dots, X_{Q-1})^\top$. (S148)

Observations (S147)–(S148) together ensure $|S_{\lambda_0}| \leq Q$ in the second line of (S143) when Condition S1 does not hold. S5.3.2 $|S| \leq Q + 2$ when $A \neq \lambda_0 I$.

We now show that $|\mathcal{S}| \leq Q + 2$ when $A \neq \lambda_0 I$. The proof parallels that for the case of $A = \lambda_0 I$ but involves more technicality. We will first reparameterize X to state a condition analogous to Condition S1, and then bound $|\mathcal{S}_{\lambda_0}|$ and $|\mathcal{S}_{\overline{\lambda_0}}|$ when the new condition holds and does not hold, respectively, analogous to (S142)–(S143).

Let $r = \operatorname{rank}(\lambda_0 I - A)$, with $1 \le r \le Q - 1$ when $A \ne \lambda_0 I$. Let

$$X_{1:r} = (X_1, \dots, X_r)^{\top}, \qquad X_{(r+1):Q} = (X_{r+1}, \dots, X_Q)^{\top},$$
$$X_{1:(Q-r)} = (X_1, \dots, X_{Q-r})^{\top}, \qquad X_{(Q-r+1):Q} = (X_{Q-r+1}, \dots, X_Q)^{\top}$$

define two partitions of $(X_1, \ldots, X_Q)^{\top}$. Applying Lemma S11(iiia) to (S140) with X_{Q+1} treated as X_0 ensures that there exists constant $g \in \mathbb{R}^r$, $G \in \mathbb{R}^{r \times (Q-r)}$, $h \in \mathbb{R}^{Q-r}$, and $H \in \mathbb{R}^{(Q-r) \times r}$ such that

$$X_{1:r} = g + GX_{(r+1):Q} \quad \text{for all } X = (X_1, \dots, X_{Q+1})^\top \in \mathcal{S}_{\lambda_0} \text{ with } X_{Q+1} = \lambda_0,$$

$$X_{1:(Q-r)} = h + HX_{(Q-r+1):Q} \quad \text{for all } X = (X_1, \dots, X_{Q+1})^\top \in \mathcal{S}_{\overline{\lambda_0}} \text{ with } X_{Q+1} \neq \lambda_0.$$
(S149)

A reparametrization.

Let $G_q \in \mathbb{R}^{1 \times (Q-r)}$ denote the *q*th row of *G* for $q = 1, \ldots, r$ with $G = (G_1, \ldots, G_r)^\top$. Motivated by the first row of (S149), define $X^* = (X_1^*, \ldots, X_{Q+1}^*)^\top = (X_{1:r}^{*\top}, X_{(r+1):Q}^{*\top}, X_{Q+1}^*)^\top$ as a reparametrization of $X = (X_1, \ldots, X_{Q+1})^\top$ with

$$X_{1:r}^* = (X_1^*, \dots, X_r^*)^\top = X_{1:r} - GX_{(r+1):Q},$$

$$X_{(r+1):Q}^* = (X_{r+1}^*, \dots, X_Q^*)^\top = X_{(r+1):Q}, \qquad X_{Q+1}^* = X_{Q+1},$$
(S150)

summarized as

$$X^* = \begin{pmatrix} X_{1:r}^* \\ X_{(r+1):Q}^* \\ X_{Q+1}^* \end{pmatrix} = \begin{pmatrix} I_r & -G & 0 \\ 0 & I_{Q-r} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} X_{1:r} \\ X_{(r+1):Q} \\ X_{Q+1} \end{pmatrix} = \Gamma_0 X, \text{ where } \Gamma_0 = \begin{pmatrix} I_r & -G & 0 \\ 0 & I_{Q-r} & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Denote by

$$X_{q}^{*}X_{q'}^{*} = c_{qq'} + \sum_{k=1}^{Q+1} a_{qq'[k]}^{*}X_{k}^{*} \quad \text{for } q, q' = 1, \dots, Q+1,$$
(S151)

the reparameterization of (S138) in terms of $X^* = (X_1^*, \ldots, X_{Q+1}^*)^\top$. The coefficients $\{a_{qq'[k]}^* : q, q', k = 1, \ldots, Q+1\}$ in (S151) are constants fully determined by $a_{qq'[k]}$'s and G, and satisfy

$$a_{qq'[k]}^* = a_{q'q[k]}^*$$
 for $q, q', k = 1, \dots, Q+1$ (S152)

under (S139). We omit their explicit forms because they are irrelevant to the proof. Let

$$\mathcal{S}^* = \{X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathbb{R}^{Q+1} : X^* \text{ satisfies (S151)}\}$$

denote the solution set of (S151). Let

$$\mathcal{S}_{\lambda_0}^* = \{ X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}^* : \ X_{Q+1}^* = \lambda_0 \},\$$
$$\mathcal{S}_{\overline{\lambda_0}}^* = \{ X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}^* : \ X_{Q+1}^* \neq \lambda_0 \}$$

denote a partition of S^* , containing solutions to (S151) that have $X_{Q+1}^* = \lambda_0$ and $X_{Q+1}^* \neq \lambda_0$, respectively. The definition of (S151) ensures

$$\begin{array}{ll}
X \in \mathcal{S}_{\lambda_0} & \iff & X^* = \Gamma_0 X \in \mathcal{S}^*_{\lambda_0} \\
X \in \mathcal{S}_{\overline{\lambda_0}} & \iff & X^* = \Gamma_0 X \in \mathcal{S}^*_{\overline{\lambda_0}}
\end{array} \quad \text{with} \quad |\mathcal{S}_{\lambda_0}| = |\mathcal{S}^*_{\lambda_0}|, \quad |\mathcal{S}_{\overline{\lambda_0}}| = |\mathcal{S}^*_{\overline{\lambda_0}}|. \quad (S153)$$

Eq. (S153) ensures that

for all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$,
the corresponding $X = (X_1, \dots, X_{Q+1})^\top = \Gamma_0^{-1} X^*$ is in \mathcal{S}_{λ_0} with $X_{Q+1} = \lambda_0$.

This, together with (S149) and $X_{1:r}^* = X_{1:r} - GX_{(r+1):Q}$ from (S150), ensures that

for all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$, the subvector
 $X_{1:r}^* = (X_1^*, \dots, X_r^*)^\top$ satisfies $X_{1:r}^* = g$ with $X_q^* = g_q$ for $q = 1, \dots, r$,
(S154)

where g_q denotes the *q*th element of *g* for $q = 1, \ldots, r$.

A condition analogous to Condition S1.

Write (S151) as

$$X_{q}^{*}X_{q'}^{*} = c_{qq'} + \sum_{k=1}^{r} a_{qq'[k]}^{*}X_{k}^{*} + \sum_{k=r+1}^{Q} a_{qq'[k]}^{*}X_{k}^{*} + a_{qq'[Q+1]}^{*}X_{Q+1}^{*} \quad \text{for } q, q' = 1, \dots, Q + (S155)$$

Plugging (S154) to the right-hand side of (S155) ensures that

for all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$, we have

$$X_{q}^{*}X_{q'}^{*} = c_{qq'} + \sum_{k=1}^{r} a_{qq'[k]}^{*}g_{k} + \sum_{k=r+1}^{Q} a_{qq'[k]}^{*}X_{k}^{*} + a_{qq'[Q+1]}^{*}\lambda_{0}$$

$$= \gamma_{qq'} + \sum_{k=r+1}^{Q} a_{qq'[k]}^{*}X_{k}^{*},$$
(S156)
where $\gamma_{qq'} = c_{qq'} + \sum_{k=1}^{r} a_{qq'[k]}^{*}g_{k} + a_{qq'[Q+1]}^{*}\lambda_{0},$

for $q, q' = 1, \dots, Q + 1$.

Divide combinations of (q, q') for $q, q' = 1, \ldots, Q + 1$ into nine blocks as follows:

	$q' \in \{1, \dots, r\}$	$q' \in \{r+1, \dots, Q\}$	q' = Q + 1	
$q \in \{1, \dots, r\}$	(i)	(ii)	(iii)	(S157)
$q \in \{r+1, \dots, Q\}$			(iv)	(5157)
q = Q + 1			(v)	

Given (S154), writing out $X_q^* = g_q$ for q = 1, ..., r on the left-hand side of (S156) implies the following for combinations of (q, q') in blocks (i)–(v) in (S157):

for all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$, we have (S158)
(i) $g_q g_{q'} = \gamma_{qq'} + \sum_{k=r+1}^Q a_{qq'[k]}^* X_k^*$ for $q, q' \in \{1, \dots, r\}$;
(ii) $g_q X_{q'}^* = \gamma_{qq'} + \sum_{k=r+1}^Q a_{qq'[k]}^* X_k^*$ for $q \in \{1, \dots, r\}$ and $q' \in \{r+1, \dots, Q\}$;
(iii) $g_q \lambda_0 = \gamma_{q,Q+1} + \sum_{k=r+1}^Q a_{q,Q+1[k]}^* X_k^*$ for $q \in \{1, \dots, r\}$ and $q' = Q+1$;
(iv) $X_q^* \lambda_0 = \gamma_{q,Q+1} + \sum_{k=r+1}^Q a_{q,Q+1[k]}^* X_k^*$ for $q \in \{r+1, \dots, Q\}$ and $q' = Q+1$;
(v) $\lambda_0^2 = \gamma_{Q+1,Q+1} + \sum_{k=r+1}^Q a_{q+1,Q+1[k]}^* X_k^*$ for $q = q' = Q+1$.

Condition S2 below parallels Condition S1, and states the condition that all coefficients of $(X_{r+1}^*, \ldots, X_Q^*)$ in (S158) are zero. We show below

$$|\mathcal{S}_{\overline{\lambda_0}}| \leq \begin{cases} r+1 & \text{if Condition S2 holds;} \\ r+2 & \text{otherwise,} \end{cases}$$
(S159)
$$|\mathcal{S}_{\lambda_0}| \leq \begin{cases} Q-r+1 & \text{if Condition S2 holds;} \\ Q-r & \text{otherwise.} \end{cases}$$
(S160)

These two parts together ensure $|\mathcal{S}| = |\mathcal{S}_{\lambda_0}| + |\mathcal{S}_{\overline{\lambda_0}}| \le Q + 2.$

Condition S2. All coefficients of $(X_{r+1}^*, \ldots, X_Q^*)$ in (S158) are zero. That is,

(i)
$$a_{qq'[k]}^* = 0$$
 for $k = r + 1, \dots, Q$ in (S158) row (i) for $q, q' \in \{1, \dots, r\}$;

(ii)
$$\begin{cases} a_{qq'[q']}^* = g_q & q \in \{1, \dots, r\}; \\ a_{qq'[k]}^* = 0 & \text{for } k \in \{r+1, \dots, Q\} \setminus \{q'\} & q' \in \{r+1, \dots, Q\}; \\ q' \in \{r+1, \dots, Q\}; \end{cases}$$

(iii)
$$a_{q,Q+1[k]}^* = 0$$
 for $k = r + 1, \dots, Q$ in (S158) row (iii) for $q \in \{1, \dots, r\}$;

(iv)
$$\begin{cases} a_{q,Q+1[q]}^* = \lambda_0 & \text{in (S158) row (iv)} & \text{for } q \in \{r+1,\dots,Q\}; \\ a_{q,Q+1[k]}^* = 0 & \text{for } k \in \{r+1,\dots,Q\} \setminus \{q\} \end{cases}$$

(v)
$$a_{Q+1,Q+1[k]}^* = 0$$
 for $k = r+1, \dots, Q$ in (S158) row (v).

Part I: Proof of (S159).

 $\frac{|\mathcal{S}_{\overline{\lambda_0}}| \leq r+1 \text{ when Condition S2 holds.}}{1 \text{ from (S152). Condition S2(i), (iii), and (v) together ensure}}$ Recall that $a_{qq'[k]}^* = a_{q'q[k]}^*$ for $q, q', k = 1, \ldots, Q+$

$$a_{qq'[r+1]}^*, a_{qq'[r+2]}^*, \dots, a_{qq'[Q]}^* = 0$$
 for all $q, q' \in \{1, \dots, r\} \cup \{Q+1\}.$

Therefore, when Condition S2 holds, the coefficients of X_{r+1}^*, \ldots, X_Q^* are all zero in the linear expansions of $\{X_q^*X_{q'}^*: q, q' = 1, \ldots, r; Q+1\}$ in (S151). The corresponding part of

(S151) reduces to

$$X_{q}^{*}X_{q'}^{*} = c_{qq'} + \sum_{k=1}^{r} a_{qq'[k]}^{*}X_{k}^{*} + a_{qq'[Q+1]}^{*}X_{Q+1}^{*} \quad \text{for} \quad q, q' = 1, \dots, r; \ Q+1.$$
(S161)

In words, (S161) ensures that

for all solutions $X^* = (X_1^*, \dots, X_{Q+1}^*)^\top$ to (S151), the subvector $X' = (X_1^*, \dots, X_r^*, X_{Q+1}^*)^\top = (X_{1:r}^{*\top}, X_{Q+1}^*)^\top$ is an $(r+1) \times 1$ vector that satisfies $X'X'^\top$ is component-wise linear in (1, X').

By induction, applying Proposition S5 to $X' = (X_1^*, \ldots, X_r^*, X_{Q+1}^*)^\top$ at $m = r + 1 (\leq Q)$ ensures that

across all solutions $X^* = (X_1^*, \dots, X_{Q+1}^*)^\top$ to (S151), the corresponding $X' = (X_{1:r}^{*\top}, X_{Q+1}^*)^\top$ take at most r+2 distinct values.

Given that $X_{Q+1}^* = \lambda_0$ in at least one solution of X^* to (S151), we have

across all solutions of $X^* = (X_1^*, \dots, X_{Q+1}^*)^\top$ to (S151) that have $X_{Q+1}^* \neq \lambda_0$, the corresponding $X' = (X_{1:r}^{*\top}, X_{Q+1}^*)^\top$ take at most r+1 distinct values. (S162)

In addition, Condition S2(iv) ensures that

for
$$q = r + 1, \dots, Q$$
, we have $\begin{aligned} a_{q,Q+1[q]}^* &= \lambda_0, \\ a_{q,Q+1[k]}^* &= 0 \quad \text{for } k \in \{r+1,\dots,Q\} \setminus \{q\}. \end{aligned}$

The linear expansions of $\{X_q^*X_{Q+1}^*: q = r+1, \ldots, Q\}$ in (S151) reduce to

$$X_q^* X_{Q+1}^* = c_{q,Q+1} + \sum_{k=1}^r a_{q,Q+1[k]}^* X_k^* + \lambda_0 X_q^* + a_{q,Q+1[Q+1]}^* X_{Q+1}^* \quad \text{for} \quad q = r+1, \dots, Q,$$

implying

$$X_{q}^{*}(X_{Q+1}^{*}-\lambda_{0}) = c_{q,Q+1} + \sum_{k=1}^{r} a_{q,Q+1[k]}^{*}X_{k}^{*} + a_{q,Q+1[Q+1]}^{*}X_{Q+1}^{*} \quad \text{for} \quad q = r+1, \dots, Q.(S163)$$

for all solutions $X^* = (X_1^*, \dots, X_{Q+1}^*)^\top$ to (S151) with $X_{Q+1}^* \neq \lambda_0$, the corresponding $X_{(r+1):Q}^* = (X_{r+1}^*, \dots, X_Q^*)^\top$ is uniquely determined by $X' = (X_1^*, \dots, X_r^*, X_{Q+1}^*)^\top = (X_{1:r}^{*\top}, X_{Q+1}^*)^\top$.

This, together with (S162), ensures

there are at most r + 1 distinct solutions of $X^* = (X_1^*, \dots, X_{Q+1}^*)^\top$ to (S151) that have $X_{Q+1}^* \neq \lambda_0$, i.e., $|\mathcal{S}_{\overline{\lambda_0}}^*| \leq r+1$.

This ensures $|\mathcal{S}_{\overline{\lambda_0}}| \leq r+1$ from (S153).

 $|\mathcal{S}_{\overline{\lambda_0}}| \leq r+2$ when Condition S2 does not hold. Recall from (S149) that

$$X_{1:(Q-r)} = h + HX_{(Q-r+1):Q} \quad \text{for all } X = (X_1, \dots, X_{Q+1})^\top \in \mathcal{S}_{\overline{\lambda_0}} \text{ with } X_{Q+1} \neq \lambda_0.$$

This ensures that

for all
$$X = (X_1, \dots, X_{Q+1})^\top \in S_{\overline{\lambda_0}}$$
 with $X_{Q+1} \neq \lambda_0$, we have

$$\sum_{k=1}^{Q-r} a_{qq'[k]} X_k = A_{qq'[1:(Q-r)]} X_{1:(Q-r)},$$
where $A_{qq'[1:(Q-r)]} = (a_{qq'[1]}, \dots, a_{qq'[Q-r]}) \in \mathbb{R}^{1 \times (Q-r)}$

$$\stackrel{(\text{S149})}{=} A_{qq'[1:(Q-r)]} \left(h + H X_{(Q-r+1):Q}\right)$$

$$= A_{qq'[1:(Q-r)]} h + A_{qq'[1:(Q-r)]} H \cdot X_{(Q-r+1):Q}$$

$$= \tilde{c}_{qq'} + \sum_{k=Q-r+1}^{Q} \tilde{a}_{qq'[k]} X_k,$$
(S164)

where $\tilde{c}_{qq'} = A_{qq'[1:(Q-r)]}h$ and $\{\tilde{a}_{qq'[k]}: k = Q - r + 1, \dots, Q\}$ are elements of $A_{qq'[1:(Q-r)]}H$. Plugging (S164) in (S138) ensures that

for all $X = (X_1, \ldots, X_{Q+1})^\top \in \mathcal{S}_{\overline{\lambda_0}}$ with $X_{Q+1} \neq \lambda_0$, we have

$$\begin{aligned} X_q X_{q'} &= c_{qq'} + \sum_{k=1}^{Q+1} a_{qq'[k]} X_k \\ &= c_{qq'} + \sum_{k=1}^{Q-r} a_{qq'[k]} X_k + \sum_{k=Q-r+1}^{Q} a_{qq'[k]} X_k + a_{qq'[Q+1]} X_{Q+1} \\ &\stackrel{(\text{S164})}{=} c_{qq'} + \left(\tilde{c}_{qq'} + \sum_{k=Q-r+1}^{Q} \tilde{a}_{qq'[k]} X_k \right) + \sum_{k=Q-r+1}^{Q} a_{qq'[k]} X_k + a_{qq'[Q+1]} X_{Q+1} \\ &= (c_{qq'} + \tilde{c}_{qq'}) + \sum_{k=Q-r+1}^{Q} (\tilde{a}_{qq'[k]} + a_{qq'[k]}) X_k + a_{qq'[Q+1]} X_{Q+1} \\ &\text{for } q, q' = Q - r + 1, \dots, Q + 1. \end{aligned}$$

In words, this ensures that

for all $X = (X_1, \ldots, X_{Q+1})^\top \in \mathcal{S}_{\overline{\lambda_0}}$ with $X_{Q+1} \neq \lambda_0$, the corresponding $X' = (X_{Q-r+1}, \ldots, X_Q, X_{Q+1})^\top$ is an $(r+1) \times 1$ vector that satisfies $X'X'^\top$ is component-wise linear in (1, X').

By induction, applying Proposition S5 to $X' = (X_{Q-r+1}, \ldots, X_Q, X_{Q+1})^{\top}$ at m = r+1 ($\leq Q$) ensures

across all $X = (X_1, \dots, X_{Q+1})^\top \in \mathcal{S}_{\overline{\lambda_0}}$ with $X_{Q+1} \neq \lambda_0$, the corresponding $X' = (X_{Q-r+1}, \dots, X_Q, X_{Q+1})^\top$ take at most r+2 distinct values. (S165)

In addition, the second row in (S149) further ensures that

all solutions $X = (X_1, \ldots, X_{Q+1})^{\top}$ to (S138) that have $X_{Q+1} \neq \lambda_0$ are uniquely determined by their corresponding $(X_{(Q-r+1):Q}, X_{Q+1})$.

This, together with (S165), ensures $|\mathcal{S}_{\overline{\lambda_0}}| \leq r+2$.

Part II: Proof of (S160).

Let $\mathcal{S}_{(r+1):Q}$ denote the set of values of $X_{(r+1):Q} = (X_{r+1}, \ldots, X_Q)^{\top}$ across all solutions $X = (X_1, \ldots, X_{Q+1})^{\top}$ to (S138) that have $X_{Q+1} = \lambda_0$. The first line in (S149) ensures that

all solutions $X = (X_1, \ldots, X_{Q+1})^\top$ to (S138) that have $X_{Q+1} = \lambda_0$ are uniquely determined by their corresponding $X_{(r+1):Q} = (X_{r+1}, \ldots, X_Q)^\top$. This ensures $|\mathcal{S}_{\lambda_0}| = |\mathcal{S}_{(r+1):Q}|$ such that it suffices to show that

$$|\mathcal{S}_{(r+1):Q}| \leq \begin{cases} Q-r+1 & \text{if Condition S2 holds;} \\ Q-r & \text{otherwise.} \end{cases}$$

A useful fact is that (S156) ensures

for all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$, we have
 $X_q^* X_{q'}^* = \gamma_{qq'} + \sum_{k=r+1}^Q a_{qq'[k]}^* X_k^*$ for $q, q' = r+1, \dots, Q$. (S166)

 $|\mathcal{S}_{(r+1):Q}| \le Q - r + 1$ when Condition S2 holds. In words, (S166) implies that

for all $X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$ with $X_{Q+1}^* = \lambda_0$, the corresponding $X_{(r+1):Q}^* = (X_{r+1}^*, \dots, X_Q^*)^\top$ is a $(Q-r) \times 1$ vector that satisfies $X_{(r+1):Q}^* X_{(r+1):Q}^{*\top}$ is component-wise linear in $(1, X_{(r+1):Q}^*)$.

By induction, applying Proposition S5 to $X^*_{(r+1):Q}$ at $m = Q - r \ (\leq Q)$ ensures that

across all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$,
the corresponding $X_{(r+1):Q}^*$ take at most $Q - r + 1$ distinct values.

This, together with the correspondence of $\mathcal{S}^*_{\lambda_0}$ and \mathcal{S}_{λ_0} from (S153), ensures that

across all $X = (X_1, \ldots, X_{Q+1})^\top \in \mathcal{S}_{\lambda_0}$ with $X_{Q+1} = \lambda_0$, the corresponding $X_{(r+1):Q}$ take at most Q - r + 1 distinct values so that $|\mathcal{S}_{(r+1):Q}| \leq Q - r + 1$.

We next improve the bound by one when Condition S2 does not hold.

 $\frac{|\mathcal{S}_{(r+1):Q}| \leq Q - r \text{ when Condition S2 does not hold.}}{\text{hold, at least one coefficient of } (X_{r+1}^*, \dots, X_Q^*) \text{ in (S158)}} \text{ When Condition S2 does not hold, at least one coefficient of } (X_{r+1}^*, \dots, X_Q^*) \text{ in (S158)} \text{ is nonzero. This ensures that there exists constants } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ such that } \{\beta_0, \beta_{r+1}, \dots, \beta_Q \in \mathbb{R} : \beta_{r+1}, \dots, \beta_Q \text{ are not all zero}\} \text{ are not all zero}\} \text{ are not all zero} \} \text{ are not all zero} \{\beta_0, \beta_1, \beta_2, \dots, \beta_Q \in \mathbb{R} : \beta_1, \dots, \beta_Q \in \mathbb{R} \} \text{ are not all zero}\} \text{ are not all zero} \} \text{ are not all zero} \text{ are not all zero} \} \text{ are not all zero} \} \text{ are not all zero} \} \text{ are not all zero} \mathbb{E} \{\beta_1, \beta_2, \dots, \beta_Q \in \mathbb{R} : \beta_1, \dots, \beta_Q \in \mathbb{R} \} \text{ are not all zero} \| \beta_1, \beta_2, \dots, \beta_Q \in \mathbb{R} \} \text{ are not all zero} \| \beta_1, \beta_2, \dots, \beta_Q \in \mathbb{R} \} \text{ are not all zero} \| \beta_1, \beta_2, \dots, \beta_Q \in \mathbb{R} \} \text{ are not all zero} \| \beta_1, \beta_2, \dots, \beta_Q \in \mathbb{R} \} \text{ are not all zero} \| \beta$

for all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$, we have $\beta_0 + \sum_{k=r+1}^Q \beta_k X_k^* = 0$.

Without loss of generality, assume that $\beta_Q \neq 0$ such that

for all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$, we have $X_Q^* = -\beta_Q^{-1} \left(\beta_0 + \sum_{k=r+1}^{Q-1} \beta_k X_k^*\right)$. (S167)

Plugging (S167) in (S166) ensures that

for all
$$X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$, we have
 $X_q^* X_{q'}^* = \gamma_{qq'} + \sum_{k=r+1}^Q a_{qq'[k]}^* X_k^*$

$$= \gamma_{qq'} + \sum_{k=r+1}^{Q-1} a_{qq'[k]}^* X_k^* + a_{qq'[Q]}^* X_Q^*$$
(S168)
 $\stackrel{(\text{S167})}{=} \gamma_{qq'} + \sum_{k=r+1}^{Q-1} a_{qq'[k]}^* X_k^* - a_{qq'[Q]}^* \beta_Q^{-1} \left(\beta_0 + \sum_{k=r+1}^{Q-1} \beta_k X_k^* \right)$
for $q, q' = r + 1, \dots, Q - 1$.

In words, (S168) ensures that

for all
$$X^* = (X_1^*, \ldots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$$
 with $X_{Q+1}^* = \lambda_0$,
the subvector $X' = (X_{r+1}^*, \ldots, X_{Q-1}^*)^\top$ is a $(Q - r - 1) \times 1$ vector that satisfies $X'X'^\top$ is component-wise linear in $(1, X')$.

By induction, applying Proposition S5 to $X' = (X_{r+1}^*, \dots, X_{Q-1}^*)^\top$ at m = Q - r - 1 (< Q) ensures that

across all $X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$ with $X_{Q+1}^* = \lambda_0$, the corresponding $X' = (X_{r+1}^*, \dots, X_{Q-1}^*)^\top$ take at most Q - r distinct values. (S169)

This, together with (S167), ensures that

across all $X^* = (X_1^*, \dots, X_{Q+1}^*)^\top \in \mathcal{S}_{\lambda_0}^*$ with $X_{Q+1}^* = \lambda_0$, the corresponding $X_{(r+1):Q}^* = (X_{r+1}^*, \dots, X_{Q-1}^*, X_Q^*)^\top$ take at most Q - r distinct values.

It then follows from (S153) that $|\mathcal{S}_{(r+1):Q}| \leq Q - r$.

S5.4 Proof of Corollary S1

We verify below Corollary S1(i)–(ii) one by one.

Proof of Corollary S1(i). The result is trivial when $c = 0_m$. When $c \neq 0_m$, define $X_0 = X^{\top}c = \sum_{j=1}^m c_j X_j$. Without loss of generality, assume $c_m \neq 0$ with $X_m = c_m^{-1}(X_0 - \sum_{j=1}^{m-1} c_i X_i)$. This ensures $X = (X_1, \ldots, X_{m-1}, X_m)^{\top}$ and $X' = (X_0, X_1, \ldots, X_{m-1})^{\top}$ are linearly equivalent. Therefore,

$$\begin{aligned} XX^{\top}c \text{ is linear in } (1,X^{\top})^{\top} & \iff XX_0 \text{ is linear in } (1,X^{\top})^{\top} \\ & \iff XX_0 \text{ is linear in } (1,X'^{\top})^{\top} \\ & \iff X'X_0 \text{ is linear in } (1,X'^{\top})^{\top}. \end{aligned}$$

The result follows from applying Proposition S4 to $X' = (X_0, X_1, \dots, X_{m-1})^\top$.

Proof of Corollary S1(ii). Let e_j denote the *j*th column of the $m \times m$ identity matrix for j = 1, ..., m. Then $XX^{\top} = (XX^{\top}e_1, XX^{\top}e_2, ..., XX^{\top}e_m)$ is component-wise linear in (1, X). The result follows from Proposition S5.