# Online Generalized Method of Moments for Time Series

Man Fung Leung <sup>\*1</sup>, Kin Wai Chan<sup>2</sup>, and Xiaofeng Shao<sup>3</sup>

<sup>1</sup>Department of Statistics, University of Illinois at Urbana-Champaign <sup>2</sup>Department of Statistics, The Chinese University of Hong Kong <sup>3</sup>Department of Statistics and Data Science, and Department of Economics, Washington University in St. Louis

#### Abstract

Online learning has gained popularity in recent years due to the urgent need to analyse large-scale streaming data, which can be collected in perpetuity and serially dependent. This motivates us to develop the online generalized method of moments (OGMM), an explicitly updated estimation and inference framework in the time series setting. The OGMM inherits many properties of offline GMM, such as its broad applicability to many problems in econometrics and statistics, natural accommodation for over-identification, and achievement of semiparametric efficiency under temporal dependence. As an online method, the key gain relative to offline GMM is the vast improvement in time complexity and memory requirement.

Building on the OGMM framework, we propose improved versions of online Sargan–Hansen and structural stability tests following recent work in econometrics and statistics. Through Monte Carlo simulations, we observe encouraging finite-sample performance in online instrumental variables regression, online overidentifying restrictions test, online quantile regression, and online anomaly detection. Interesting applications of OGMM to stochastic volatility modelling and inertial sensor calibration are presented to demonstrate the effectiveness of OGMM.

*Keywords:* Instrumental variable; Online learning; Quantile regression; Recursive estimation; Streaming data.

<sup>\*</sup>Correspondence email: mfleung2@illinois.edu

### 1 Introduction

The generalized method of moments (GMM, Hansen, 1982) is a fundamental estimation and inference framework in econometrics and statistics. It is widely used to analyse economic and financial data, and subsumes many popular statistical methods such as least squares, maximum likelihood, and instrumental variables regression (Hall, 2005). However, despite being partly motivated by situations in which asymptotically efficient estimation (e.g., maximum likelihood) is computationally burdensome (Hansen, 1982; Hall, 2005), GMM does not scale to modern data sets with millions of data points. For large-scale streaming data, which can be collected in perpetuity and serially dependent, the classical (offline) GMM is too computationally expensive and may fail due to violation of memory constraints (Chen et al., 2023).

With the emergence of streaming data collection techniques, it has become important to develop online estimation and inference procedures that can be computationally (and memory) efficient while preserving statistical properties of their offline counterparts. In the context of GMM, Chen et al. (2023) adapted the stochastic approximation (Robbins and Monro, 1951) to linear instrumental variables regression for independent data and provided plug-in and self-normalized inference procedures. They also discussed learning rate selection, multi-epoch estimation, and nonlinear GMM. Luo and Song (2020) developed renewable estimation and incremental inference for generalized linear models with independent and identically distributed (iid) data. Their proposed estimator is based on estimating equations, which correspond to GMM with exact identification. Subsequently, Luo et al. (2022) extended renewable estimation to longitudinal data analysis using the quadratic inference function (Qu et al., 2000), which is also a special case of GMM. They further discussed anomaly detection based on a modified Sargan–Hansen test. In a related work, Luo et al. (2023) focused on streaming longitudinal data with a fixed set of participants. Under a first-order autoregressive working correlation structure, they proposed a new decomposition for the quadratic inference function that allows online updates. They also considered time-varying parameter estimation by introducing an exponential smoothing factor that dynamically adjusts the weights applied to historical data batches.

The overarching goal of this article is to develop the online generalized method of moments (OGMM), a general and efficient estimation and inference framework for streaming time series, which seems lacking in the literature. Using the first-order Taylor approximation and a lagged linearization (i.e., a linearization step with the OGMM estimator obtained at the previous step plugged in), we derive an explicitly updated estimator in contrast to implicitly updated renewable estimators in the literature (Schifano et al., 2016; Luo and Song, 2020; Luo et al., 2022, 2023); see Toulis and Airoldi (2017) for the difference between explicit and implicit updates in the context of stochastic approximation. To achieve semiparametric efficiency, we employ the optimal weighting matrix by inverting the online long-run variance estimator in Leung and Chan (2025). This allows us to perform efficient estimation and inference for the parameter in a fully online fashion.

Our major contributions include:

- (a) We develop an online GMM estimator in the time series setting and show that the OGMM estimator is consistent, asymptotically normal, and achieves the same asymptotic efficiency as offline GMM and implicitly updated renewable estimators;
- (b) Our framework is autocorrelation-robust by handling the serial dependence in a nonparametric fashion. The semiparametric efficiency is thus achieved through online estimation of the optimal weighting matrix. Other than the user-chosen parameters involved in the online updates of the weighting matrix, our algorithm does not involve any tuning parameter such as the learning rate in stochastic approximation;
- (c) Following the work of Chen et al. (2023) and Luo et al. (2022), we propose improved versions of online Sargan–Hansen and structural stability tests, which do not suffer from considerable size distortion (Chen et al., 2023) nor require historical raw data (Luo et al., 2022);
- (d) Owing to the broad framework of GMM, we can include online one-step estimation, online least squares, online quantile regression and online instrumental variables regression as special cases. Numerical comparison with several existing methods shows that the finite-sample performance of our OGMM is very competitive.

The rest of this paper is organized as follows. Section 2 reviews relevant literature and introduces some notation. Section 3 formulates OGMM by solving three computational challenges faced by GMM, and shows that OGMM covers many existing statistical methods as special cases. Section 4 develops the asymptotic theory of OGMM. Section 5 compares OGMM with some existing statistical methods in Monte Carlo experiments. Section 6 presents two applications in stochastic volatility modelling and inertial sensor calibration. Section 7 discusses our findings and some future directions. All experiments are performed on Red Hat Enterprise Linux 9.4 with an Intel Xeon Gold 6148 CPU and R version 4.2.3. An R-package ogmm that implements our framework is available online. All proofs and some additional results are also deferred to the online Supplement(Leung et al., 2025).

### 2 Literature review

Consider stationary and ergodic time series data  $\{D_i = (x_{i,1}, x_{i,2}, \ldots, x_{i,n_i})\}_{i=1}^b$  arriving in batches, where  $x_{i,j} \in \mathbb{R}^d$  is the *j*-th vector of observations in the *i*-th batch of data with sample size  $n_i$ . Let  $\theta^* \in \Theta \subset \mathbb{R}^p$  be a vector of unknown parameters which are to be estimated, and  $g(\theta, x)$  be a  $q \times 1$  vector of functions that satisfies the population moment condition  $\mathbb{E}\{g(\theta^*, x_{i,j})\} = 0$  for all i, j with  $q \ge p$ . To simplify the notation, write  $G(\theta; D_i) = \sum_{j=1}^{n_i} g(\theta, x_{i,j})$  and  $N_b = \sum_{i=1}^b n_i$ . The celebrated GMM estimator of  $\theta^*$  is

$$\tilde{\theta}_{b,\mathsf{GMM}} = \operatorname*{argmin}_{\theta \in \Theta} \bar{g}_b(\theta)^{\mathsf{T}} \tilde{W} \bar{g}_b(\theta), \qquad (2.1)$$

where  $\bar{g}_b(\theta) = N_b^{-1} \sum_{i=1}^b G(\theta; D_i)$  is the sample moment and  $\tilde{W}$  is a weighting matrix.

Under suitable conditions (Hansen 1982), it is well known that (2.1) is asymptotically optimal when  $\tilde{W} = \tilde{\Sigma}^{-1}$ , where  $\tilde{\Sigma}$  is a consistent estimator of the long-run variance matrix

$$\Sigma = \lim_{N_b \to \infty} N_b \operatorname{Var}\{\bar{g}_b(\theta^*)\}.$$
(2.2)

In this case,  $\sqrt{N_b}(\tilde{\theta}_{b,\mathsf{GMM}} - \theta^*) \stackrel{\mathrm{d}}{\to} \mathrm{N}(0, (V^{\mathsf{T}}\Sigma^{-1}V)^{-1})$ , where  $V = \mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}$  is the expectation of the gradient of  $g(\theta^*, x_{i,j})$  and  $\stackrel{\mathrm{d}}{\to}$  denotes convergence in distribution. However, as  $\Sigma$  depends on the unknown  $\theta^*$ , substitutes of  $\theta^*$  are necessary for efficient estimation. Classical variants of **GMM** such as two-step, iterated and continuously updating differ in the way to substitute  $\theta^*$ ; see Hall (2005) and the references therein.

There is a rich literature on the estimation of (2.2) because the long-run variance is an important quantity in time series analysis. However, classical nonparametric estimators utilizing the overlapping batch means (Meketon and Schmeiser, 1984) or kernels/lag windows (Parzen, 1957; Andrews, 1991) cannot be updated online. Wu (2009) and Chan and Yau (2016, 2017) proposed different online alternatives based on subsample selection rules, which replace the constant batch size of a batch means estimator with a sequence of batch sizes. Leung and Chan (2025) developed a different approach by decomposing kernels, which led to estimators with higher statistical and computational efficiency.

To update GMM and obtain online estimates, the stochastic approximation (Robbins and Monro, 1951) is a natural candidate. By averaging the iterates (Polyak and Juditsky, 1992), one can further conduct inference using an online long-run variance estimator (Chen et al., 2020; Zhu et al., 2023; Leung and Chan, 2025) or a self-normalizer/random scaling (Lee et al., 2022). Chen et al. (2023) appears to be the first to extend the stochastic approximation to the GMM setting, and they developed the stochastic GMM (SGMM) for independent data. They also proposed an online Sargan–Hansen test statistic and state its asymptotic distribution under the null hypothesis. Nevertheless, their premier algorithm assumes a fixed sample size and may not be efficient after the first epoch, which is why they recommend running over multiple epochs in practice (Chen et al., 2023). The computational cost or number of epochs to obtain an efficient estimate may vary, which makes SGMM unsuitable for streaming data collected in perpetuity. On the other hand, their online Sargan–Hansen test may suffer from considerable size distortion in finite sample. Since SGMM is based on explicit stochastic approximation, its performance can also be sensitive to the choice of initial learning rate (Toulis and Airoldi, 2017). We will revisit each of these issues and show numerical evidence in Section 5.

Method	Data assumption	Over-identification	Update
SGMM	independent	allowed	explicit
Luo and Song $(2020)$	iid	not allowed	implicit
Luo et al. (2022)	independent	under working correlation	implicit
Luo et al. (2023)	first-order autoregressive	under working correlation	implicit
$OGMM\ (\mathrm{proposal})$	stationary & ergodic	allowed	$\operatorname{explicit}$

Table 1: Summary of properties of different online estimation and inference framework.

Another candidate for updating GMM is renewable estimation, a term coined in Luo and Song (2020). Essentially, one stores a finite number of statistics after first-order Taylor approximations so that the estimating equation and its solution (i.e., the renewable estimator) only depends on the stored statistics and the latest batch of data (Schifano et al., 2016; Luo and Song, 2020). Luo et al. (2022, 2023) investigated extension to longitudinal data analysis using quadratic inference function, which allows for over-identification due to the inverse working correlation matrix being approximated by a linear combination of some basis matrices. Renewable estimation is also widely applied in variants of online quantile regression (Jiang and Yu, 2022; Sun et al., 2024; Jiang and Yu, 2024). Nevertheless, renewable estimators involve solving fixed-point equations so their updates depend on the stopping criteria and are implicit in the sense of Toulis and Airoldi (2017). Explicitly updated estimators with a lower computational cost remain largely unexplored in the literature. Additionally, the anomaly detection statistic in Luo et al. (2022) requires storing a reference data batch and solving an optimization problem for every new data batch. A fully online detector remains to be developed. We will discuss renewable estimation again in Section 3 as it is closely related to our proposal. Table 1 highlights some key differences between our proposal and some existing ones.

Before ending this section, we introduce some notations. Denote the gradient operation with respect to  $\theta$  by  $\nabla_{\theta}$ . For any vector a and matrix A, denote the  $l^2$ -vector norm and Frobenius norm by  $||a||_2 = \sqrt{a^{\intercal}a}$  and  $||A||_F = \sqrt{tr}(A^{\intercal}A)$ , respectively. Denote the identity matrix of size q by  $I_q$ . We reserve the tilde, circumflex and check for offline, OGMM and other online methods (e.g.,  $\tilde{\theta}_{b,\text{GMM}}$ ,  $\hat{\theta}_{b,\text{OGMM}}$  and  $\check{\theta}_{k,\text{SGMM}}$ ), respectively. We may also suppress the subscript if the context is clear.

### 3 Methodology

In Sections 3.1 to 3.3, we formulate OGMM by solving three computational challenges involved in adapting GMM to an online setting. Then, we discuss implementation and some special cases in Section 3.4. Some online inference examples are given in Section 3.5.

#### 3.1 Challenges in moment computation

In general, the sample moment  $\bar{g}_b(\theta)$  cannot be updated online when  $g(\theta, x)$  is nonlinear in  $\theta$ . To see this, suppose we have  $\bar{g}_{b-1}(\theta_1)$  from the previous step and would like to compute  $\bar{g}_b(\theta_2)$ . We cannot apply the recursive formula for sample mean because it requires the evaluation of  $\bar{g}_{b-1}(\theta_2)$ . However, we can derive an online moment using the first-order Taylor approximation. Let  $\hat{\theta}_1$  be an initial estimate based on  $D_1$  and  $\hat{W}$  be a weighting matrix. For b = 1, 2, ..., define

$$V_b = \frac{1}{N_b} \sum_{i=1}^b \nabla G(\hat{\theta}_i; D_i) \quad \text{and} \quad U_b = \frac{1}{N_b} \sum_{i=1}^b \left\{ G(\hat{\theta}_i; D_i) - \nabla G(\hat{\theta}_i; D_i) \hat{\theta}_i \right\},$$

which are estimators of  $V = \mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}$  and  $U = \mathbb{E}\{g(\theta^*, x_{i,j})\} - V\theta^*$ , respectively. Since  $V_b$  and  $U_b$  are the latest online estimates after  $\hat{\theta}_b$  is obtained,  $U_b + V_b\theta$  is an online approximation of the sample moment  $\bar{g}_b(\theta)$ . When b = 2, we have

$$N_{2}\bar{g}_{2}(\theta) = G(\hat{\theta}_{1}; D_{1}) + \nabla G(\hat{\theta}_{1}; D_{1})(\theta - \hat{\theta}_{1}) + G(\theta; D_{2}) + O_{p}\left(n_{1} \left\|\hat{\theta}_{1} - \theta\right\|_{2}^{2}\right)$$
$$= N_{1}(U_{1} + V_{1}\theta) + G(\theta; D_{2}) + O_{p}\left(n_{1} \left\|\hat{\theta}_{1} - \theta\right\|_{2}^{2}\right).$$

By storing  $U_1$ ,  $V_1$  and  $N_1$  in advance, we can compute  $N_1(U_1 + V_1\theta) + G(\theta; D_2)$  in  $O(n_2q)$  time without using  $D_1$  explicitly. If the error term is asymptotically negligible, a reasonable online estimator can be obtained from

$$\underset{\theta \in \Theta}{\operatorname{argmin}} \frac{1}{N_2^2} \left\{ N_1(U_1 + V_1\theta) + G(\theta; D_2) \right\}^{\mathsf{T}} \hat{W} \left\{ N_1(U_1 + V_1\theta) + G(\theta; D_2) \right\}.$$
(3.1)

The derivation of (3.1) shares the same idea as renewable estimation, except that the literature (e.g., Luo and Song 2020) consider solving the first-order optimality condition

$$\frac{1}{N_2^2} \left\{ N_1 V_1 + \nabla G(\hat{\theta}_2; D_2) \right\}^{\mathsf{T}} \hat{W} \left\{ N_1 (U_1 + V_1 \hat{\theta}_2) + G(\hat{\theta}_2; D_2) \right\} = V_2^{\mathsf{T}} \hat{W} (U_2 + V_2 \hat{\theta}_2) = 0. \quad (3.2)$$

Rewriting (3.2) gives  $\hat{\theta}_2 = -(V_2 \hat{W} V_2)^{-1} V_2^{\mathsf{T}} \hat{W} U_2$ , which is a fixed-point equation because  $U_2$  and  $V_2$  depend on  $\hat{\theta}_2$ . Therefore, the Newton–Raphson method is usually used to solve (3.2) iteratively in the literature.

### 3.2 Challenges in stable computation

For time-sensitive applications, solving an optimization problem like (3.1) or a fixed-point equation like (3.2) for every new data batch may be undesirable because the time to meet the stopping criteria may vary. In light of it, we propose to estimate V and U by

$$\hat{V}_b = \frac{1}{N_b} \sum_{i=1}^b \nabla G(\hat{\theta}_{\min(i,b-1)}; D_i)$$

$$= \frac{1}{N_b} \left\{ N_{b-1} V_{b-1} + \nabla G(\hat{\theta}_{b-1}; D_b) \right\} \text{ and}$$
$$\hat{U}_b = \frac{1}{N_b} \sum_{i=1}^b \left\{ G(\hat{\theta}_{\min(i,b-1)}; D_i) - \nabla G(\hat{\theta}_{\min(i,b-1)}; D_i) \hat{\theta}_{\min(i,b-1)} \right\}$$
$$= \frac{1}{N_b} \left\{ N_{b-1} U_{b-1} + G(\hat{\theta}_{b-1}; D_b) - \nabla G(\hat{\theta}_{b-1}; D_b) \hat{\theta}_{b-1} \right\},$$

respectively. Essentially, we replace  $\hat{\theta}_b$  by  $\hat{\theta}_{b-1}$  in  $V_b$  and  $U_b$  to obtain  $\hat{V}_b$  and  $\hat{U}_b$ , respectively. Then, we can stabilize the computational time by using the lagged  $\hat{\theta}_1$  to linearize  $G(\hat{\theta}_2; D_2)$  and replacing  $\nabla G(\hat{\theta}_2; D_2)$  with  $\nabla G(\hat{\theta}_1; D_2)$  in (3.2), which leads to

$$\hat{V}_2^{\mathsf{T}}\hat{W}(\hat{U}_2 + \hat{V}_2\hat{\theta}_2) = 0.$$
(3.2')

The solution  $\hat{\theta}_2 = -(\hat{V}_2 \hat{W} \hat{V}_2)^{-1} \hat{V}_2^{\mathsf{T}} \hat{W} \hat{U}_2$  is explicit and does not involve any tuning parameter. These are also true for b > 2 with

$$\hat{\theta}_{b,\mathsf{OGMM}} = -(\hat{V}_b^{\mathsf{T}} \hat{W} \hat{V}_b)^{-1} \hat{V}_b^{\mathsf{T}} \hat{W} \hat{U}_b; \tag{3.3}$$

see Algorithm 3.1 for the recursive steps involved in the computation of  $\hat{\theta}_{b, \mathsf{OGMM}}$ .

If  $\hat{W} = I_q$  and  $g(\theta, x)$  is the score function, (3.3) is same as the implicit estimator in Luo and Song (2020) obtained via the Newton–Raphson method with one iteration. In other words, (3.3) can be interpreted as an explicit estimator, which seems largely unexplored in the renewable estimation literature. For stochastic gradient descent, Toulis and Airoldi (2017) showed that implicit and explicit procedures have identical asymptotic performance, but explicit estimators are more sensitive to the learning rate. Interestingly, (3.3) also achieves the same asymptotic efficiency as its offline and implicit online counterparts under an additional condition that  $n_i = o(N_{i-1}^2)$  for  $i = 2, \ldots, b$ ; see Theorem 2 and Remark 1 for a discussion. However, (3.3) does not require learning rate selection.

### 3.3 Challenges in weighting matrix computation

The final challenge is the choice of  $\hat{W}$ , which is important as it affects the statistical efficiency. Computationally, there are at least three estimation approaches as discussed below. (1) We compute  $\hat{\Sigma}_1$  based on  $D_1$  and fix  $\hat{W} = \hat{\Sigma}_1^{-1}$ . Many classical estimators are available and the computational cost is minimized. Nevertheless, this approach may be inefficient since it does not utilize the sequentially arrived data. (2) We assume  $g(\theta^*, x_{i,j})$  follows some parametric model (e.g., a vector autoregressive moving average model; see Section 3.5.2 of Hall 2005) and update such model recursively. Then, we can plug in the online parameter estimates to obtain  $\hat{\Sigma}_b$  via the long-run variance estimator based on the parametric model and set  $\hat{W} = \hat{\Sigma}_b^{-1}$ . This is similar to using a working correlation matrix in Luo et al. (2022, 2023), which leads to a loss of statistical efficiency in case of model misspecification. (3) We update  $\hat{W}$  using an online nonparametric estimator  $\hat{\Sigma}_b(K)$ ,

where K is a kernel function that satisfies some conditions. This approach yields both statistical efficiency and autocorrelation robustness. While it involves some smoothing parameter, there are automatic optimal selectors in the literature (Chan and Yau, 2017; Leung and Chan, 2025) so that users only need to provide problem-specific information, e.g., the strength of serial dependence and the memory constraint.

In general, we recommend the third approach. If users have a strict time budget or are certain that the data are independent, they can use a fixed weighting matrix or invert Welford's (1962) sample variance estimator, which are implemented in our R-package ogmm. However, the default choice is to invert the long-run variance estimator in Leung and Chan (2025), whose implementation allows batch updates and includes a positive definiteness adjustment. To be specific, their long-run variance estimator takes the form

$$\hat{\Sigma}_b(K) = \frac{1}{N_b} \sum_{i=1}^{N_b} \sum_{j=1}^{N_b} K_{N_b}(i,j) (X_i - \bar{X}_{N_b}) (X_j - \bar{X}_{N_b})^{\mathsf{T}}, \qquad (3.4)$$

where  $X_k$ 's are the evaluated moment functions indexed by a single subscript in our setting, i.e.,  $X_k = g(\hat{\theta}_b, x_{b,j})$  if  $k = \sum_{i=1}^{b-1} n_i + j$ , and  $\bar{X}_{N_b} = N_b^{-1} \sum_{i=1}^{N_b} X_i \equiv N_b^{-1} \sum_{j=1}^{b} G(\hat{\theta}_j; D_j)$ . Their kernel takes the form

$$K_n(i,j) = \left(1 - \frac{|i-j|^{\lambda}}{t_n^{\lambda}}\right) \mathbb{1}_{|i-j| \le s'_{i \lor j}},$$

where  $s_n = \min(\lfloor \Psi n^{\psi} \rfloor, n-1); t_n = \min(\lceil \Xi n^{\xi} \rceil, n);$ 

$$s'_{n} = \begin{cases} s'_{n-1} + 1, & \text{if } s_{n-1} \le s'_{n-1} + 1 < \phi s_{n-1}; \\ s_{n}, & \text{if } s'_{n-1} + 1 \ge \phi s_{n-1}. \end{cases}$$

 $\Psi, \Xi \in \mathbb{R}^+$ ;  $\psi, \xi \in (0, 1)$ ;  $\lambda \in \mathbb{Z}^+$ ; and  $\phi \in [1, \infty)$ . Their optimal parameters selector automatically handles  $\Psi, \Xi, \psi$  and  $\xi$  so users only need to choose  $\lambda$  and  $\phi$ . Following Leung and Chan (2025), we recommend  $\lambda = 1$  ( $\lambda = 3$ ) if the serial dependence is strong (weak), and  $\phi = 1$  ( $\phi = 2$ ) if the memory is abundant (scarce).

#### 3.4 Implementation and special cases

Algorithm 3.1 summarizes the key steps in OGMM estimation. Alternatively, let  $\hat{\theta}'_b = \hat{\theta}'_{b-1} - \{(\hat{V}'_b)^{\top}\hat{W}\hat{V}'_b\}^{-1}(\hat{V}'_b)^{\top}\hat{W}\hat{U}'_b$ , where  $V'_1 = \nabla \bar{g}_1(\hat{\theta}'_1)$ ,  $V'_b = N_b^{-1}\{N_{b-1}V'_{b-1} + \nabla G(\hat{\theta}'_b; D_b)\}$ ,  $\hat{V}'_b = N_b^{-1}\{N_{b-1}V'_{b-1} + \nabla G(\hat{\theta}'_b; D_b)\}$ ,  $U'_1 = \bar{g}_1(\hat{\theta}'_1)$ ,  $U'_b = N_b^{-1}\{N_{b-1}U'_{b-1} + N_{b-1}V'_{b-1}(\hat{\theta}'_b - \hat{\theta}'_{b-1}) + G(\hat{\theta}'_b; D_b)\}$ , and  $\hat{U}'_b = N_b^{-1}\{N_{b-1}U'_{b-1} + G(\hat{\theta}'_{b-1}; D_b)\}$ . Here,  $U'_b$  and  $\hat{U}'_b$  are estimators of  $U + V\theta^* = \mathbb{E}\{g(\theta^*, x_{i,j})\}$ . The following proposition verifies that  $\hat{\theta}'_b \equiv \hat{\theta}_b$  and  $U'_b \equiv U_b + V_b\hat{\theta}_b$ . Since fewer arithmetic operations are involved in this alternative definition, we actually implement  $\hat{\theta}'_b$  in our R-package ogmm.

**Proposition 1** (Validity of the telescoping-based formula). If  $\hat{\theta}'_1 = \hat{\theta}_1$ , then  $\hat{\theta}'_b \equiv \hat{\theta}_b$  and

Algorithm 3.1: Online Generalized Method of Moments

[1] initialization: [2] Set b = 1 and  $N_1 = n_1$ [3] Compute  $\hat{\theta}_1$  using  $D_1$  and any reasonable method (e.g., GMM) Initialize  $\hat{\Sigma}_1$  using  $\hat{\theta}_1$  and set  $\hat{W} = \hat{\Sigma}_1^{-1}$ [4] Set  $U_1 = \bar{g}_1(\hat{\theta}_1) - \nabla \bar{g}_1(\hat{\theta}_1)\hat{\theta}_1$  and  $V_1 = \nabla \bar{g}_1(\hat{\theta}_1)$ [5] begin [6] Set b = b + 1[7] Receive  $D_b$  and set  $N_b = N_{b-1} + n_b$ [8] Set  $\hat{U}_b = N_b^{-1} \{ N_{b-1} U_{b-1} + G(\hat{\theta}_{b-1}; D_b) - \nabla G(\hat{\theta}_{b-1}; D_b) \hat{\theta}_{b-1} \}$ [9] Set  $\hat{V}_b = N_b^{-1} \{ N_{b-1} V_{b-1} + \nabla G(\hat{\theta}_{b-1}; D_b) \}$ [10] Compute  $\hat{\theta}_b = -(\hat{V}_b^{\mathsf{T}}\hat{W}\hat{V}_b)^{-1}\hat{V}_b^{\mathsf{T}}\hat{W}\hat{U}_b$ [11] (Optional) update  $\hat{\Sigma}_{b-1}$  using  $\hat{\theta}_b$  and set  $\hat{W} = \hat{\Sigma}_b^{-1}$ [12] Set  $U_b = N_b^{-1} \{ N_{b-1} U_{b-1} + G(\hat{\theta}_b; D_b) - \nabla G(\hat{\theta}_b; D_b) \hat{\theta}_b \}$ [13] Set  $V_b = N_b^{-1} \{ N_{b-1} V_{b-1} + \nabla G(\hat{\theta}_b; D_b) \}$ [14]

$$U_b' \equiv U_b + V_b \hat{\theta}_b$$
 for  $b = 1, 2, \dots$ 

Besides, it is not necessary to initialize  $\hat{\theta}_1$  using GMM. Any  $\sqrt{N_1}$ -consistent  $\hat{\theta}_1$  will be sufficient for  $\hat{\theta}_2$  to achieve the same asymptotic efficiency as GMM. For example, one can compute  $\hat{\theta}_1$  using the two-stage least squares (2SLS) for linear instrumental variables estimation. The consistency of  $\hat{\theta}_1$  is important so users may want to combine several data batches to form a large  $D_1$  in practice, which is also recommended in Luo et al. (2022).

We further analyse the computational complexities of Algorithm 3.1 after initialization. For lines 7–11 and 13–14, the total time and space complexities are  $O(n_bpq + pq^2)$ and  $O(n_bq + q^2)$ , respectively. For line 12 with the long-run variance estimator in Leung and Chan (2025), the time complexity is  $O(n_bq^2 + q^3)$ , and the space complexity is  $O(N_b^{1/(1+2\lambda)}q^2\mathbb{1}_{\phi<2} + n_bq + q^2)$ . For comparison, the time complexity of offline GMM is at least  $O(N_bq)$ , which can be greater depending on the optimization method and how the weighting matrix is chosen, and the space complexity is  $O(N_bq + q^2)$ . As for general SGMM, the time complexity to update for  $n_b$  iterations is  $O(n_bpq^2)$ , and the space complexity is  $O(n_bq + q^2)$  (or  $O(N_bq + q^2)$  if multiple epochs are needed). We will compare the computational cost of GMM, SGMM and OGMM in simulation studies later.

Same as GMM, OGMM covers many statistical methods as special cases. We give a few examples below to demonstrate the generality of Algorithm 3.1.

**Example 1** (One-step estimation). Suppose  $\hat{W} = I_q$  and  $g(\theta, x)$  is the score function. Let  $\hat{\theta}_1$  be a  $\sqrt{N_1}$ -consistent estimator and  $D_2$  be empty. Then,  $\hat{\theta}_2 = -(\hat{V}_2\hat{W}\hat{V}_2)^{-1}\hat{V}_2^{\top}\hat{W}\hat{U}_2 = \hat{\theta}_1 - (V_1^{\top}I_qV_1)^{-1}V_1^{\top}I_q\bar{g}_1(\hat{\theta}_1) = \hat{\theta}_1 - V_1^{-1}\bar{g}_1(\hat{\theta}_1)$  performs Le Cam's (1956) one-step estimation provided that  $V_1$  is a consistent estimator for the Fisher information matrix. In this regard, OGMM can be viewed as a one-step estimator for GMM with online data.

**Example 2** (Online least squares). Let  $Y_i = (y_{i,1}, \ldots, y_{i,n_i})^{\mathsf{T}}$  and  $X_i = (x_{i,1}, \ldots, x_{i,n_i})^{\mathsf{T}}$ be the *i*-th batch of responses and predictors, respectively. The ordinary least squares is equivalent to GMM with  $\hat{W} = I_q$  and  $g(\theta, \{x, y\}) = x(y - x^{\mathsf{T}}\theta)$ . We can verify that  $\nabla g(\theta, \{x, y\}) = -xx^{\mathsf{T}}, \ \hat{V}_b = -N_b^{-1} \sum_{i=1}^b X_i^{\mathsf{T}} X_i$  and  $\hat{U}_b = N_b^{-1} \sum_{i=1}^b X_i^{\mathsf{T}} Y_i$ . Therefore,  $\hat{\theta}_b = (\sum_{i=1}^b X_i^{\mathsf{T}} X_i)^{-1} (\sum_{i=1}^b X_i^{\mathsf{T}} Y_i)$  exactly recovers the ordinary least squares.

**Example 3** (Online quantile regression). To simplify the discussion here, suppose we only use an initial estimate  $\hat{\theta}_{\tau,1}$  to update the summary statistics of OGMM and the online linear estimator for quantile regression (LEQR) in Chen et al. (2019), where  $\tau$  is a quantile level of interest. We will compare the actual estimators in Section 5.3. Let  $\check{U}_{b,\text{LEQR}} = N_b^{-1} \sum_{i=1}^b X_i^{\top} [H\{(Y_i - X_i\hat{\theta}_{\tau,1})/h_i\} + \tau - 1 + (Y_i/h_i)H\{(Y_i - X_i\hat{\theta}_{\tau,1})/h_i\}]$ , where  $H(\cdot)$  is a smooth approximation of the indicator function  $\mathbb{1}_{>0}$ ,  $\nabla H(\cdot)$  is the gradient of  $H(\cdot)$ , and  $\{h_i\}$  is a sequence of bandwidths. The smoothed quantile regression (de Castro et al., 2019) is equivalent to GMM with  $\hat{W} = I_q$  and  $g(\theta_{\tau}, \{x, y, h\}) = x[H\{(y - x^{\top}\theta_{\tau})/h\} + \tau - 1]$ . We can verify that  $\nabla g(\theta_{\tau}, \{x, y, h\}) = -(xx^{\top}/h)\nabla H\{(y - x^{\top}\theta_{\tau})/h\}$ ,  $\hat{V}_b = -N_b^{-1} \sum_{i=1}^b (X_i^{\top} X_i/h_i)\nabla H\{(Y_i - X_i\hat{\theta}_{\tau,1})/h_i\}$  and  $\hat{U}_b = N_b^{-1} \sum_{i=1}^b X_i^{\top} [H\{(Y_i - X_i\hat{\theta}_{\tau,1})/h_i\}]$ . The resulting estimator  $\hat{\theta}_b = \hat{V}_b^{-1}\hat{U}_b$  is almost same as  $\check{\theta}_{b,\text{LEQR}} = \hat{V}_b^{-1}\check{U}_{b,\text{LEQR}}$ , except that  $X_i\hat{\theta}_{\tau,1}/h_i$  in  $\hat{U}_b$  is replaced with  $Y_i/h_i$  in  $\check{U}_{b,\text{LEQR}}$ .

### **3.5** Online inference and applications

To conduct inference under the OGMM framework, we can simply replace offline statistics in the GMM literature with online counterparts from Algorithm 3.1. Consequently, we can improve the computational efficiency while preserving statistical properties of many classical inferential procedures.

**Example 4** (Over-identifying restrictions testing and confidence region estimation). Let  $\chi^2_{p,\alpha}$  be the  $\alpha$ -quantile of a chi-squared distribution with p degrees of freedom. The offline Sargan–Hansen test statistic is  $N_b \bar{g}_b (\tilde{\theta}_{b,\mathsf{GMM}})^{\mathsf{T}} \tilde{\Sigma}^{-1} \bar{g}_b (\tilde{\theta}_{b,\mathsf{GMM}})$ , where  $\tilde{\Sigma}$  is an offline estimator of  $\Sigma$ . Since  $U_b + V_b \theta$  is an online approximation of  $\bar{g}_b(\theta)$ , a natural online test statistic is  $T = N_b (U_b + V_b \hat{\theta}_{b,\mathsf{OGMM}})^{\mathsf{T}} \hat{\Sigma}_b^{-1} (U_b + V_b \hat{\theta}_{b,\mathsf{OGMM}})$ . We reject the null hypothesis of  $\mathbb{E}\{g(\theta^*, x_{i,j})\} = 0$  at  $100\alpha\%$  nominal level if  $T > \chi^2_{q-p,1-\alpha}$ . Similarly, an online  $100(1-\alpha)\%$  confidence region for  $\theta^*$  is  $\{\theta : N_b(\hat{\theta}_{b,\mathsf{OGMM}} - \theta)^{\mathsf{T}} (V_b^{\mathsf{T}} \hat{\Sigma}_b^{-1} V_b)^{-1} (\hat{\theta}_{b,\mathsf{OGMM}} - \theta) \leq \chi^2_{p,1-\alpha}\}$ .

**Example 5** (Anomaly detection). Consider the abnormal data batch detection setting in Luo et al. (2022), where  $D_1$  is the normal reference and  $D_b$  is possibly abnormal. We are interested in testing  $H_0: \mathbb{E}\{g(\theta^*, x_{1,j})\} = \mathbb{E}\{g(\theta^*, x_{b,h})\} = 0$  against  $H_1: \mathbb{E}\{g(\theta^*, x_{1,j})\} = 0$ , but  $\mathbb{E}\{g(\theta^*, x_{b,h})\} \neq 0$  for  $j \in \{1, \ldots, n_1\}$  and  $h \in \{1, \ldots, n_b\}$ . Luo et al. (2022) proposed

$$T_{\rm R} = n_1^{-1} G(\tilde{\theta}_{b,{\rm R}}; D_1)^{\mathsf{T}} \tilde{C}_1^{-1} G(\tilde{\theta}_{b,{\rm R}}; D_1) + n_b^{-1} G(\tilde{\theta}_{b,{\rm R}}; D_b)^{\mathsf{T}} \tilde{C}_b^{-1} G(\tilde{\theta}_{b,{\rm R}}; D_b), \quad \text{where} \quad (3.5)$$

$$\tilde{\theta}_{b,\mathrm{R}} = \underset{\theta \in \Theta}{\operatorname{argmin}} \left\{ n_1^{-1} G(\theta; D_1)^{\mathsf{T}} \tilde{C}_1(\theta)^{-1} G(\theta; D_1) + n_b^{-1} G(\theta; D_b)^{\mathsf{T}} \tilde{C}_b(\theta)^{-1} G(\theta; D_b) \right\}$$
(3.6)

is a restricted estimator, and  $\tilde{C}_i(\theta)$  is the sample variance that depends on  $\theta$  and  $D_i$  for  $i \in \{1, b\}$ . Same as (2.1), one can obtain (3.6) using two-step, iterated or continuously updating GMM. The test statistic in (3.5) is not autocorrelation-robust though and its computational complexities depend on  $D_1$ . If the normal reference contains all previous data batches, computing (3.5) will be very costly. Therefore, we propose

$$T_{\rm F} = n_1 (U_1 + V_1 \hat{\theta}_{b,\rm F})^{\mathsf{T}} \hat{\Sigma}_1^{-1} (U_1 + V_1 \hat{\theta}_{b,\rm F}) + n_b^{-1} G(\hat{\theta}_{b,\rm F}; D_b)^{\mathsf{T}} \hat{\Sigma}_1^{-1} G(\hat{\theta}_{b,\rm F}; D_b), \qquad (3.7)$$

where  $\hat{\theta}_{b,\mathrm{F}}$  is the full-sample estimator that can be obtained by an OGMM update with  $D_b$ , and  $\hat{\Sigma}_1$  is the estimated long-run variance before receiving  $D_b$  (assuming the serial dependence structure does not change over time). Under  $H_0$ , both (3.5) and (3.7) converge in distribution to  $\chi^2_{2q-p}$ . However, (3.7) does not require solving another optimization problem nor storing previous data batches.

**Example 6** (Structural stability testing). Abnormality in Example 5 corresponds to a more general behaviour termed structural instability with a known break point in the context of GMM. Indeed, replacing the restricted estimator in (3.5) with the fullsample estimator in (3.7) is inspired by a common practice in structural stability testing; see p.173–174 of Hall (2005) for additional references and discussion. Now, for  $i \in \{1, b\}$ , let  $\Sigma_i(\theta)$  and  $P_i(\theta)$  be the subsample analogues of  $\Sigma(\theta) = \lim_{N_b \to \infty} N_b \operatorname{Var}\{\bar{g}_b(\theta)\}$ and  $P(\theta) = \Sigma(\theta)^{-1/2} V(\theta) \{V(\theta)^{\top} \Sigma(\theta)^{-1} V(\theta)\}^{-1} V(\theta)^{\top} \Sigma(\theta)^{-1/2}$  for  $D_i$ , respectively, where  $V(\theta) = \mathbb{E}\{\nabla g(\theta, x_{i,j})\}$ . Hall and Sen (1999) decomposed  $H_0$  into

$$H_{0,\mathrm{I}}:P_{1}(\theta^{*})\Sigma_{1}(\theta^{*})^{-1/2}\mathbb{E}\{g(\theta^{*},x_{1,j})\} = P_{b}(\theta^{*})\Sigma_{b}(\theta^{*})^{-1/2}\mathbb{E}\{g(\theta^{*},x_{b,j})\} = 0 \text{ and} \\ H_{0,\mathrm{O}}:\{I_{q}-P_{1}(\theta^{*})\}\Sigma_{1}(\theta^{*})^{-1/2}\mathbb{E}\{g(\theta^{*},x_{1,j})\} = \{I_{q}-P_{b}(\theta^{*})\}\Sigma_{b}(\theta^{*})^{-1/2}\mathbb{E}\{g(\theta^{*},x_{b,j})\} = 0.$$

If  $H_0$  is rejected and  $\mathbb{E}\{g(\theta^*, x_{b,h})\} \neq 0$  for  $h \in \{1, \ldots, n_b\}$ , there are two possibilities. (1) There is some unique  $\theta' \in \Theta \setminus \{\theta^*\}$  that satisfies  $\mathbb{E}\{g(\theta', x_{b,h})\} = 0$ , which means that the instability is caused by a change point in parameter values. In this case, only  $H_{0,I}$ is violated. (2) There is no  $\theta \in \Theta$  that satisfies  $\mathbb{E}\{g(\theta, x_{b,h})\} = 0$ , which means that the instability is caused by a more fundamental misspecification instead of a change point in parameter value. In this case,  $H_{0,O}$  and likely  $H_{0,I}$  are both violated. The test statistics in Example 5 are not able to distinguish between these two cases. Therefore, based on Hall and Sen (1999), we propose

$$T_{\rm U} = n_1 (U_1 + V_1 \hat{\theta}_1)^{\mathsf{T}} \hat{\Sigma}_1^{-1} (U_1 + V_1 \hat{\theta}_1) + n_b^{-1} G(\tilde{\theta}_{b,\rm U}; D_b)^{\mathsf{T}} \tilde{\Sigma}_{b,\rm U}^{-1} G(\tilde{\theta}_{b,\rm U}; D_b), \qquad (3.8)$$

where  $\tilde{\theta}_{b,\mathrm{U}}$  and  $\tilde{\Sigma}_{b,\mathrm{U}}$  are the (unrestricted) GMM and long-run variance estimator computed with  $D_b$  only. Since  $U_1$ ,  $V_1$ ,  $\hat{\theta}_1$  and  $\hat{\Sigma}_1$  are available before receiving  $D_b$ , (3.8) is computationally efficient. Under the null hypothesis, (3.8) converges in distribution to  $\chi^2_{2(q-p)}$  and is asymptotically independent of (3.5) and (3.7). Therefore, given that (3.5) or (3.7) is rejected, we may believe that there is a change point in parameter value if (3.8) is not rejected, and the model is misspecified if (3.8) is rejected. The unknown break point case seems more challenging and is left for future research.

### 4 Theory

To compare with GMM on the same theoretical basis, we develop the asymptotic theory of OGMM based on the assumptions in Hall (2005). Two different cases that correspond to the scenario (S2) in Luo et al. (2022) are considered: (a)  $\min(n_1, \ldots, n_b) \rightarrow \infty$ ; (b)  $n_1 \rightarrow \infty$  and no restrictions on  $n_2, n_3, \ldots$ . We allow b to be any finite positive integer but require  $n_1 \rightarrow \infty$  to ensure the initial estimate is consistent, which is implicitly assumed in Luo and Song (2020) and many subsequent works.

#### 4.1 Consistency

Assumption 1 (Strict stationarity). The time series data  $\{x_{1,1}, \ldots, x_{1,n_1}, x_{2,1}, \ldots, x_{2,n_2}, \ldots\}$  form a strictly stationary process with sample space  $\mathcal{X} \subset \mathbb{R}^d$ .

Assumption 2 (Regularity conditions for  $g(\theta, x)$ ). The function  $g(\theta, x)$  satisfies:

- (a) it is continuous on  $\Theta$  for each  $x \in \mathcal{X}$ ;
- (b)  $\mathbb{E}\{g(\theta, x_{i,j})\}$  exists and is finite for every  $\theta \in \Theta$ ;
- (c)  $\mathbb{E}\{g(\theta, x_{i,j})\}$  is continuous on  $\Theta$ .

Assumption 3 (Population moment condition). The data  $\{x_{i,j}\}$  and the parameter  $\theta^*$  satisfy the population moment condition  $\mathbb{E}\{g(\theta^*, x_{i,j})\} = 0$ .

Assumption 4 (Global identification).  $\mathbb{E}\{g(\theta', x_{i,j})\} \neq 0 \text{ for all } \theta' \in \Theta \setminus \{\theta^*\}.$ 

Assumption 5 (Regularity conditions for  $\nabla g(\theta, x)$ ).

- (a)  $\nabla g(\theta, x)$  exists and is continuous on  $\Theta$  for each  $x \in \mathcal{X}$ ;
- (b)  $\theta^*$  is an interior point of  $\Theta$ ;
- (c)  $\mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}$  exists and is finite.

Assumption 6 (Weighting matrix).  $\hat{W}$  is a positive semi-definite matrix which converges in probability to a positive definite constant matrix W.

Assumption 7 (Ergodicity). The random process  $\{x_{i,j}\}$  is ergodic.

Assumption 8 (Compactness).  $\Theta$  is a compact set.

Assumption 9 (Domination of  $g(\theta, x)$ ).  $\mathbb{E}\{\sup_{\theta \in \Theta} \|g(\theta, x_{i,j})\|_2\} < \infty$ .

Assumption 10 (Long-run variance).

- (a)  $\mathbb{E}\{g(\theta^*, x_{i,j})g(\theta^*, x_{i,j})^{\mathsf{T}}\}\$  exists and is finite;
- (b)  $\Sigma$  defined in (2.2) exists and is a finite positive definite matrix.

Assumption 11 (Local Lipschitz continuity of  $\nabla g(\theta, x)$ ). Let  $B_{\epsilon} = \{\theta \in \Theta : \|\theta - \theta^*\|_2 \le \epsilon\}$ .

- (a) For some  $\epsilon > 0$ , there exists L > 0 such that  $\|\mathbb{E}\{\nabla g(\theta, x_{i,j})\} \mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}\|_F \le L \|\theta \theta^*\|_2$  for all  $\theta \in B_{\epsilon}$ .
- (b) For some  $\epsilon > 0$ , there exists L > 0 such that  $\|\nabla g(\theta, x) \nabla g(\theta^*, x)\|_F \le L \|\theta \theta^*\|_2$ for all  $\theta \in B_{\epsilon}$  and  $x \in \mathcal{X}$ .

Assumption 12 (Local uniform convergence to  $\mathbb{E}\{\nabla g(\theta, x_{i,j})\}$ ).

- (a) For some  $\epsilon > 0$  and i = 1, ..., b, as  $n_i \to \infty$ ,
  - (i)  $\sup_{\theta \in B_{\epsilon}} \|n_i^{-1} \nabla G(\theta; D_i) \mathbb{E}\{\nabla g(\theta, x_{i,j})\}\|_F = o_p(1)$ , which is implied by
  - (ii)  $\sup_{\theta \in B_{\epsilon}} \|n_i^{-1} \nabla G(\theta; D_i) \mathbb{E} \{ \nabla g(\theta, x_{i,j}) \} \|_F = O_p(n_i^{-1/2}).$
- (b) For some  $\epsilon > 0$ , as  $N_b \to \infty$ ,
  - (i)  $\left\|N_b^{-1}\sum_{k=1}^b \nabla G(\theta^*; D_k) \mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}\right\|_F = o_p(1)$ , which is implied by
  - (ii)  $\sup_{\theta \in B_{\epsilon}} \left\| N_b^{-1} \sum_{k=1}^b \nabla G(\theta; D_k) \mathbb{E} \{ \nabla g(\theta, x_{i,j}) \} \right\|_F = o_p(1).$

Assumptions 1–10 are identical to Assumptions 3.1–3.5 and 3.7–3.11 in Hall (2005). Assumptions 11 and 12 are required to hold for some neighbourhood  $B_{\epsilon}$  of  $\theta^*$  only so both are local assumptions. Although Assumptions 11(a) and (b) (local Lipschitz continuity) are stronger than Assumption 3.12 (continuity) in Hall (2005), they are standard in the renewable estimation literature to bound the Taylor approximation error. Compared with Condition 3 in Luo and Song (2020), Assumption 11(b) is slightly weaker because only local smoothness is required, and Assumption 11(a) is implied by Assumption 11(b). Note that Assumption 12(a)(ii) is stronger than Assumption 12(b)(ii), where the latter is identical to Assumption 3.13 in Hall (2005).

**Theorem 1** (Consistency of  $\hat{\theta}_b$ ). Suppose Assumptions 1–9 hold.

- (a) Under Assumptions 11(a) and 12(a)(i),  $\hat{\theta}_b \xrightarrow{p} \theta^*$  as  $\min(n_1, \ldots, n_b) \to \infty$ .
- (b) Under Assumptions 11(b) and 12(b)(i),  $\hat{\theta}_b \xrightarrow{p} \theta^*$  as  $n_1 \to \infty$ .

When b = 1, Theorem 1 is exactly Theorem 3.1 in Hall (2005), which states the consistency of  $\hat{\theta}_1$  if it is computed using GMM. Then, for  $b \ge 2$ , Assumptions 11 and 12(i) ensure the Taylor approximation error is negligible so that  $\hat{\theta}_b$  remains consistent.

### 4.2 Asymptotic normality

**Theorem 2** (Asymptotic normality of  $\hat{\theta}_b$ ). Recall that  $V = \mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}$ . Let  $H = (V^{\mathsf{T}}WV)^{-1}V^{\mathsf{T}}W$ . Suppose Assumptions 1–10 hold and  $n_i = o(N_{i-1}^2)$  for  $i = 2, \ldots, b$ .

- (a) Under Assumptions 11(a) and 12(a)(ii),  $\sqrt{N_b}(\hat{\theta}_b \theta^*) \stackrel{d}{\to} N(0, H\Sigma H^{\intercal})$  as  $\min(n_1, \dots, n_b) \to \infty.$
- (b) Under Assumptions 11(b) and 12(b)(ii),  $\sqrt{N_b}(\hat{\theta}_b \theta^*) \xrightarrow{d} N(0, H\Sigma H^{\intercal})$  as  $n_1 \to \infty$ .

When b = 1, Theorem 2 is exactly Theorem 3.2 in Hall (2005), which states the asymptotic normality of  $\hat{\theta}_1$  if it is computed using GMM. The asymptotic variance is minimized at  $(V^{\top}\Sigma^{-1}V)^{-1}$  when  $W = \Sigma^{-1}$ . For  $b \ge 2$ ,  $\hat{\theta}_b$  achieves the same asymptotic efficiency regardless of whether the initial estimator is GMM. As briefly discussed in Example 1, any  $\sqrt{N_1}$ -consistent  $\hat{\theta}_1$  will be sufficient for  $\hat{\theta}_2$  to achieve the same asymptotic efficiency as GMM. Therefore, users have some flexibility in choosing  $\hat{\theta}_1$  in Algorithm 3.1.

Compared with renewable estimators in the literature, OGMM also achieves the same asymptotic efficiency as its offline counterpart in Theorem 2. The condition  $n_i = o(N_{i-1}^2)$ for i = 2, ..., b appears to be new and mild, and is needed to control the Taylor approximation error of explicit updates in OGMM. It is easily seen to be satisfied when the batch size is constant, and can be fulfilled in practice by splitting a large batch of new data into smaller batches. Other assumptions are similar to the standard conditions in the GMM or renewable estimation literature.

**Remark 1** (Explicit renewable estimators). The renewable estimators (4.5) and (4.12) for unconditional quantile regression in Jiang and Yu (2024) are also explicit. Their additional condition is  $n_i = O(n_1)$  for i = 2, ..., b, which implies our condition.

**Remark 2** (Verification of Assumptions 11(b) and 12(b)(ii)). In practice, case (b) is usually of broader interest because there are no restrictions on  $n_2, n_3, \ldots$  Therefore, we give some examples where we can verify Assumptions 11(b) and 12(b)(ii).

- (a) For the ordinary least squares in Example 2, Assumption 11(b) holds because  $\nabla g(\theta, \{x, y\}) = -xx^{\intercal}$  does not depend on  $\theta$ . By the law of large numbers, Assumption 12(b)(ii) also holds if  $\mathbb{E}\{\|xx^{\intercal}\|_{F}\} < \infty$ .
- (b) The instrumental variables estimation is equivalent to GMM with  $g(\theta, \{x, y, z\}) = z(y-x^{\mathsf{T}}\theta)$ , where z is a vector of instrumental variables. Similar to the ordinary least squares, Assumption 11(b) holds because  $\nabla g(\theta, \{x, y, z\}) = -zx^{\mathsf{T}}$  does not depend on  $\theta$ , and Assumption 12(b)(ii) holds if  $\mathbb{E}\{\|zx^{\mathsf{T}}\|_F\} < \infty$ .
- (c) The maximum likelihood estimation is equivalent to GMM with  $g(\theta, x) = \nabla \ell(\theta, x)$ , where  $\ell(\theta, x)$  is the log-likelihood function. If  $\ell(\theta, x)$  is twice continuously differentiable for all  $\theta \in \Theta$  (Condition 3 in Luo and Song 2020), Assumption 11(b) holds by the mean value theorem. If  $\Theta$  is compact and  $\mathbb{E}\{\sup_{\theta \in \Theta} \|\nabla^2 \ell(\theta, x)\|_F\} < \infty$ , then Assumption 12(b)(ii) also holds by Lemma 2.4 in Newey and McFadden (1994).

### 4.3 Over-identifying restrictions test

**Theorem 3** (Asymptotic distributions of the online sample moment and online Sargan–Hansen test statistic). If the conditions in Theorem 2(a) or (b) hold, then  $\hat{W}^{1/2}\sqrt{N_b}(U_b + V_b\hat{\theta}_b) \stackrel{d}{\rightarrow} N(0, RW^{1/2}\Sigma(W^{1/2})^{\intercal}R^{\intercal})$  as (a)  $\min(n_1, \ldots, n_b) \rightarrow \infty$ ; or (b)  $n_1 \rightarrow \infty$ , where  $R = I_q - W^{1/2}V(V^{\intercal}WV)^{-1}V^{\intercal}(W^{1/2})^{\intercal}$ . Suppose further that  $\hat{\Sigma}$  is positive semi-definite and converges in probability to  $\Sigma$  in Assumption 10. Then  $N_b(U_b + V_b\hat{\theta}_b)^{\top}\hat{\Sigma}^{-1}(U_b + V_b\hat{\theta}_b) \xrightarrow{d} \chi^2_{q-p}$  as (a)  $\min(n_1, \ldots, n_b) \to \infty$ ; or (b)  $n_1 \to \infty$ .

When b = 1, Theorem 3 is equivalent to Theorems 3.3 and 5.1 in Hall (2005) if  $\hat{\theta}_1$  is computed using GMM because  $U_1 + V_1 \hat{\theta}_1 \equiv \bar{g}_1(\hat{\theta}_1)$ . For  $b \ge 2$ , the online sample moment and online Sargan–Hansen test statistic differ from their offline version as more than one  $\hat{\theta}_i$ 's are plugged in. Therefore, we have an additional first-order correction term  $N_b^{-1} \sum_{i=1}^b \nabla G(\hat{\theta}_i; D_i)(\hat{\theta}_b - \hat{\theta}_i)$  compared with Chen et al. (2023). We remark that the condition on  $\hat{\Sigma}$  is satisfied by the online long-run variance estimator in Leung and Chan (2025) under standard conditions.

### 5 Simulation results

In this section, we examine the finite-sample performance of OGMM, as compared to several competitors in the literature. Specifically, we report the results for online instrumental variables regression in Section 5.1, for online Sargan-Hansen test in Section 5.2, for online quantile regression in Section 5.3 and for online anomaly detection in Section 5.4.

Throughout Sections 5 and 6, we may lighten the notation by using a single subscript to index variables, e.g.,  $x_k = x_{b,j}$  with  $k = \sum_{i=1}^{b-1} n_i + j$ . Unless otherwise stated, we use the R-package momentfit 0.5 for offline two-step GMM and 2SLS, sandwich 3.1.1 for the Bartlett kernel estimator with a bandwidth selected by fitting a first-order autoregressive model (Andrews, 1991) and prewhite=FALSE, rlaser 0.1.1 for Leung and Chan's (2025) recursive long-run variance estimator with  $\lambda = 1$ ,  $\phi = 1$ , pilot equals to  $n_1$  and warm=FALSE, and ogmm 0.0.1 for OGMM, SGMM and LEQR. Non-default arguments will be discussed whenever they are modified. All simulations are repeated for 1000 times.

#### 5.1 Online instrumental variables regression

Before we investigate the finite-sample performance of OGMM, we describe SGMM (Chen et al., 2023), another framework for computing GMM with online updates. Let  $\eta_k = \eta_0 k^{-a}$ be the learning rate at the k-th iteration for some  $\eta_0 > 0$  and  $a \in (1/2, 1)$ . The general SGMM algorithm considers  $\check{\theta}_{k+1} = \check{\theta}_k - \eta_{k+1}(\check{V}_k^{\top}\check{W}_k\check{V}_k)^{-1}\check{V}_k^{\top}\check{W}_kg(\check{\theta}_k, x_{k+1})$ , where  $\check{V}_k$  is an online average that estimate  $V = \mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}$ , and  $\check{W}_k$  is the inverse of the second sample moment updated using the Sherman–Morrison–Woodbury formula. By updating the average  $\bar{\theta}_N = N^{-1} \sum_{k=1}^N \check{\theta}_k$ , inference on  $\theta^*$  can be performed using a plug-in approach. To conduct the Sargan–Hansen test, Chen et al. (2023) proposed the test statistic

$$N\check{g}_N^{\mathsf{T}}\check{W}_N\check{g}_N$$
, where  $\check{g}_N = \frac{1}{N}\sum_{k=1}^N g(\bar{\theta}_k, x_k).$  (5.1)

For the initial learning rate, consider using an initialization data batch  $D_0$  to compute  $\check{\theta}_0$ ,  $\check{V}_0$  and  $\check{W}_0$ . Then, Chen et al. (2023) proposed to use  $\eta_0 = 1/\Psi_0(\kappa)$ , where

$$\Psi_0(\kappa) = \operatorname{quantile}_{\kappa} \left\{ p^{-1} \left\| (\check{V}_0^{\mathsf{T}} \check{W}_0 \check{V}_0)^{-1} \check{V}_0^{\mathsf{T}} \check{W}_0 \nabla g(\check{\theta}_0, x) \right\|_S : x \in D_0 \right\},$$
(5.2)

 $\kappa$  is a predetermined quantile level and  $\|\cdot\|_S$  is the spectral norm in (5.2).

Now, consider the following linear instrumental variables regression models:

- (a) Independent: let  $y_k = x_k^{\mathsf{T}} \theta^* + 5 \exp(z_{k,q})(\nu_k + \varepsilon_k)$ , where  $\nu_k, \varepsilon_k \stackrel{\text{iid}}{\sim} \mathcal{N}(0,1)$ ,  $z_k \stackrel{\text{iid}}{\sim} \mathcal{N}_q(0, \mathcal{V})$  with  $\mathcal{V}_{i,j} = \rho^{|i-j|}$ ,  $x_{k,1} = 0.1 \sum_{j=2}^p x_{k,j} + 0.5 \sum_{j=p}^q z_{k,j} + \nu_k$ , and  $x_{k,j} = z_{k,j-1}$  for  $j = 2, \ldots, p$ . We exactly replicate a simulation model in Chen et al. (2023) by setting  $\theta^* = (1, \ldots, 1)^{\mathsf{T}}$ , p = 5, q = 20, and  $\rho = 0.5$ .
- (b) Dependent: let  $y_k = \theta_1^* y_{k-1} + \theta_2^* y_{k-2} + \phi_1^* \varepsilon_{k-1} + \phi_2^* \varepsilon_{k-2} + \varepsilon_k$ , where  $\varepsilon_k \stackrel{\text{iid}}{\sim} N(0, 1)$ . We exactly replicate a simulation model in Chaussé (2010) by setting  $\theta^* = (1.4, -0.6)^{\top}$  and  $\phi^* = (0.6, -0.3)^{\top}$ . The selected instruments are  $z_k = (y_{k-3}, \dots, y_{k-6})^{\top}$ .

We are interested in estimating  $\theta^*$  at  $N_b = 2000b$  for  $b = 1, 2, \dots, 100$  using

- (a) 2SLS: the two-stage least squares as the offline benchmark and initial estimator for model (a). The Bartlett kernel estimator is used for inference.
- (b) GMM: the two-step GMM as the offline benchmark and initial estimator for model
  (b). The Bartlett kernel estimator is used for inference and the weighting matrix. We also use θ<sub>b-1</sub> as the initial value when we optimize for θ<sub>b</sub>, and θ<sub>0</sub> = (0,0)<sup>T</sup>.
- (c) SGMM: for model (a), we consider different initial learning rates selected at  $\kappa \in \{0.5, 0.7, 0.9\}$ . For model (b), we only consider  $\kappa = 0.5$  but further try Leung and Chan's (2025) recursive long-run variance estimator for both inference and the weighting matrix. For both models, we set a = 0.501 in the learning rate  $\eta_k$ . However, the stochastic 2SLS algorithm is not used for warm up as in Chen et al. (2023) because it requires a randomly partitioned subsample with size that scales with  $N_b$ , which is not feasible in our online setting where  $N_b$  is increasing. Similarly, we do not consider the multi-epoch approach in Chen et al. (2023).
- (d) OGMM: both Welford's (1962) and Leung and Chan's (2025) estimators are considered for inference and the weighting matrix. We further compare explicit and implicit updates with the stopping criteria in Luo et al. (2022). Specifically, we stop when the number of Newton–Raphson iterations reaches 50 or  $\{(\hat{V}_{b}^{(r)})^{\top}\hat{W}\hat{U}_{b}^{(r)}\}^{\top}\{(\hat{V}_{b}^{(r)})^{\top}\hat{W}\hat{V}_{b}^{(r)}\}^{-1}(\hat{V}_{b}^{(r)})^{\top}\hat{W}\hat{U}_{b}^{(r)} < 10^{-6}, \text{ where } \hat{V}_{b}^{(r)} = N_{b}^{-1}\{N_{b-1}V'_{b-1}+\nabla G(\hat{\theta}_{b}^{(r-1)};D_{b})\}, \hat{U}_{b}^{(r)} = N_{b}^{-1}\{N_{b-1}U'_{b-1}+N_{b-1}V'_{b-1}(\hat{\theta}_{b}^{(r-1)}-\hat{\theta}'_{b-1})+G(\hat{\theta}_{b}^{(r-1)};D_{b})\}, \text{ and } \hat{\theta}_{b}^{(r)} = \hat{\theta}_{b}^{(r-1)} \{(\hat{V}_{b}^{(r)})^{\top}\hat{W}\hat{V}_{b}^{(r)}\}^{-1}(\hat{V}_{b}^{(r)})^{\top}\hat{W}\hat{U}_{b}^{(r)}.$

In Figures 1 & 2, we compare mean squared errors of the point estimates, coverage rates of the 95% confidence intervals for  $\theta_1^*$ , and computational time in seconds. Figure 1 confirms that suboptimal  $\sqrt{N_1}$ -consistent  $\hat{\theta}_1$  (such as 2SLS) is still sufficient for OGMM to be asymptotically optimal. Despite the fact that data are independent under model



Figure 1: Online instrumental variables estimation of  $\theta_1^*$  in the independent model (a). A better estimator has a lower mean squared error (MSE), a coverage rate closer to 95% and a shorter computation time.



Figure 2: Online instrumental variables estimation of  $\theta_1^*$  in the dependent model (b). The caption of Figure 1 also applies here.

(a), using a recursive long-run variance estimator for autocorrelation robustness does not affect the performance of OGMM much. On the other hand, SGMM is not sensitive to the choice of  $\kappa$  here. Its mean squared errors relative to 2SLS and coverage rates are similar to those reported in Chen et al. (2023). Nevertheless, SGMM is not as efficient as OGMM in a moderately large sample ( $N_b \leq 2 \times 10^5$ ).

Figure 2 verifies the asymptotic efficiency of OGMM relative to GMM. It also shows that OGMM and SGMM greatly reduce the computational cost of GMM. Since data are dependent under model (b), recursive long-run variance estimation improves the coverage rates of SGMM, which was originally designed for independent data. However, OGMM still outperforms SGMM when the same long-run variance estimator is used. It does not gain efficiency through implicit updates though because the moment is linear in  $\theta$ ; see Remark 2. Additional results with the same findings are deferred to the Supplement.

### 5.2 Online Sargan–Hansen test

In Section 4.3, we observe that there is a first-order correction term in our online Sargan– Hansen test statistic because we are plugging in a sequence of estimates rather than a single estimate in the offline setting. On the other hand, there is no first-order correction term in (5.1). To investigate the influence of this term, consider the following models:

- (c) Independent: let  $y_k = \theta_1^* x_k + \theta_2^* z_{k,1} + \varepsilon_k$  and  $x_k = z_{k,1} + z_{k,2} + \nu_k$ , where  $(z_{k,1}, z_{k,2}, \nu_k, \varepsilon_k)^{\intercal} \sim N_4(0, \mathcal{V})$  with  $\mathcal{V}_{i,j} = \mathbb{1}_{i=j} + 0.5\mathbb{1}_{(i,j)\in\{(1,2),(2,1),(3,4),(4,3)\}}$ . This is the simulation model in Hall (2000), and we consider  $\theta_1^* = 1$  and  $\theta_2^* \in \{0, 0.05, \dots, 0.5\}$ .
- (d) Dependent: same as model (c) but (z<sub>k,1</sub>, z<sub>k,2</sub>, ν<sub>k</sub>, ε<sub>k</sub>)<sup>T</sup> follows a vector autoregressive model of order 1 with mean (0,...,0)<sup>T</sup>, coefficient matrix diag(0.5,...,0.5) and Gaussian noise covariance matrix V in model (c).

We estimate  $\theta_1^*$  at  $N_b = (100)2^b$  for b = 1, 2, ..., 5 under the assumption that  $\mathbb{E}\{z_k(y_k - \theta_1^*x_k)\} = 0$ . Clearly, misspecification happens when  $\theta_2^* \neq 0$ . The method settings are same as those for model (b). We continue to use two-step GMM as the offline benchmark and initial estimator, and SGMM with  $\kappa = 0.5$  and a = 0.501.

Figure 3 reports the empirical size and size-adjusted power of different Sargan-Hansen tests for  $\theta_2^* \in \{0, 0.2, 0.4\}$ . Overall, the performance of OGMM is very encouraging. Although the batch size  $n_b = N_{b-1}$  is increasing, the statistical efficiency of OGMM remains comparable to that of GMM, which corroborates with our theory in Theorems 2 and 3. Recursive long-run variance estimation does not cause size distortion under model (c) and allows autocorrelation-robust online testing under model (d). On the other hand, SGMM suffers from considerable size distortion and exhibits non-monotonic size-adjusted power (with respect to the sample size) that seems counter-intuitive and cannot be explained by the variance estimator. This confirms the validity and necessity of the first-order correction term in Theorem 3 when a sequence of estimates is plugged in. The results for other values of  $\theta_2^*$  are similar and deferred to the Supplement.

### 5.3 Online quantile regression

As illustrated in Example 3, online methods for quantile regression are often based on smoothing the indicator function in the quantile loss (Chen et al., 2019; Sun et al., 2024; Jiang and Yu, 2024). There is another class of online methods based on a different kind of smoothing (Jiang and Yu, 2022). However, we continue with Example 3 to compare with Chen et al. (2019) because other estimators are based on implicit updates. Suppose observations are divided into intervals by their indices. Define  $b_0 = 0$ ,  $c_0 = -\infty$ , and  $c_{2k-1} = 2^{k-1} + 1/2$  and  $c_{2k} = 2^{k-1} + 3/4$  for  $k \ge 1$ . Let *m* be the memory constraint and



Figure 3: Online Sargan–Hansen test at 5% nominal level under the independent model (c) (upper panel) and dependent model (d) (lower panel). When  $\theta_2^* = 0$ , the model is well-specified. A size closer to 5% means that the test controls the type I error better. When  $\theta_2^* > 0$ , the model is misspecified. A high size-adjusted power means the test controls the type II error better.

 $b_l = \lfloor m^{c_{l-1}} \rfloor + 1$  be the first index in the *l*-th interval. Chen et al. (2019) elaborated that  $\{b_l\}$  was chosen in this way because there would be no improvement of online LEQR after  $n^2$  fresh observations if the previous estimate was based on *n* observations. This is similar to our condition  $n_i = o(N_{i-1}^2)$  because  $n_i$  is the number of fresh observations and  $N_{i-1}$  is previous sample size. However, apart from the minor difference in Example 3, OGMM is methodologically different from LEQR in two ways. First, OGMM includes all data in the summary statistics, whereas LEQR mainly includes data from the previous and current intervals; see Algorithm 2 in Chen et al. (2019). Second, OGMM may utilize the memory constraint of size *m* so that  $\hat{\theta}_{b-1}$  is updated frequently, whereas LEQR only updates the estimate that is used to compute summary statistics at the start of a new interval. Now, consider the following models:

(e) Independent: let  $y_k = x_k^{\mathsf{T}} \theta^* + \varepsilon_k$ , where  $x_{k,1} = 1, x_{k,2}, \ldots, x_{k,p}$  follow a uniform distribution on [0, 1] with  $\operatorname{Corr}(x_{k,i}, x_{k,j}) = 0.5^{|i-j|}$ , and  $\varepsilon_k \stackrel{\text{iid}}{\sim} \operatorname{N}(0, 1)$ . We set  $\theta^* =$ 



Figure 4: Online quantile regression under the independent model (e) (upper panel) and dependent model (f) (lower panel). A better estimator has a lower median absolute error (MAE), a coverage rate closer to 95% and a shorter computation time.

- $(1, \ldots, 1)^{\mathsf{T}}$  and p = 10, which replicates a simulation model in Chen et al. (2019).
- (f) Dependent: same as model (e) but  $(\varepsilon_k, x_{k,2}, \dots, x_{k,p})^{\mathsf{T}}$  follows a vector autoregressive model of order 1 with mean  $(0, \dots, 0)^{\mathsf{T}}$ , coefficient matrix diag $(0.5, \dots, 0.5)$  and Gaussian noise covariance matrix  $\mathcal{V}_{i,j} = 0.5^{|i-j|} \mathbb{1}_{i\neq 1} \mathbb{1}_{j\neq 1} + \mathbb{1}_{i=1} \mathbb{1}_{j=1}$  for  $1 \leq i, j \leq p$ .

We are interested in estimating  $\theta_{\tau}^*$  at  $N_b = 2000b$  for  $b = 1, \ldots, 250$ . Recall that  $\theta_{\tau}^*$  can be computed by shifting  $\varepsilon_k$  such that  $\mathbb{P}(\varepsilon_k \leq 0 \mid x_k) = \tau$  (Chen et al., 2019). For  $\tau = 0.1$ , the intercept  $\theta_{\tau 1}^*$  in models (e) and (f) are approximately -0.28 and -0.48, respectively. The method settings are close to those for model (a). The differences are that we use the default modified Barrodale and Roberts algorithm in the R-package quantreg (Koenker, 2005) for initial estimation, LEQR as the online benchmark, and SGMM with  $\kappa \in \{0.97, 0.98, 0.99\}$  and a = 0.501. For the smoothing in Example 3, we follow Chen et al. (2019) to use  $H(x) = \{1/2 + (15/16)(x - 2x^3/3 + x^5/5)\}\mathbb{1}_{|x|<1} + \mathbb{1}_{x\geq 1}$  and  $h_b = \sqrt{p/N_{b-1}}$ , where b is the current number of intervals (LEQR) or batches (SGMM/OGMM).

In Figure 4, we compare the median absolute errors of point estimates, the computa-

tional time in seconds, and the coverage rates of 95% confidence intervals for the summed coefficient  $J^{\dagger}\theta_{\tau}^{*}$ , where  $J = (1, \ldots, 1)^{\dagger}$  and  $\tau = 0.1$ . The median absolute error is considered here because SGMM with  $\kappa \in \{0.97, 0.98\}$  failed to initialize or gave unreasonable estimates in some replications. Figure 4 illustrates that OGMM's methodological differences from LEQR are beneficial. The computational cost of OGMM is slightly higher due to recursive long-run variance estimation, but it enables autocorrelation-robust inference in model (f). On the other hand, SGMM is very sensitive to the choice of  $\kappa$  here because a lot of  $\nabla g(\check{\theta}_0, x)$  in (5.2) are zero or very small. It is also computationally inefficient because we can only apply the general algorithm instead of the one specialized for instrumental variables in Chen et al. (2023). The latter one includes a Sherman–Morrison–Woodbury formula for  $(\check{V}_k^{\dagger}\check{W}_k\check{V}_k)^{-1}$ , which is not available in the general version.

#### 5.4 Online anomaly detection

Recall from Examples 5 and 6 that there are two types of abnormality or structural instability caused by a change point in parameter value and a misspecified model, respectively. Now, we extend the experiments in Section 5.2 by modifying the models there:

- (g) Independent: let  $y_k = (\theta_1^* + \theta_2^* \mathbb{1}_{k \in \Lambda_{\mathrm{I}}}) x_k + \theta_2^* \mathbb{1}_{k \in \Lambda_{\mathrm{O}}} z_{k,1} + \varepsilon_k$  and  $x_k = z_{k,1} + z_{k,2} + \nu_k$ , where  $(z_{k,1}, z_{k,2}, \nu_k, \varepsilon_k)^{\top} \stackrel{\mathrm{iid}}{\sim} \mathrm{N}_4(0, \mathcal{V})$  with  $\mathcal{V}_{i,j} = \mathbb{1}_{i=j} + 0.5\mathbb{1}_{(i,j)\in\{(1,2),(2,1),(3,4),(4,3)\}}, \Lambda_{\mathrm{I}} = \{2001, \ldots, 2500, 6001, \ldots, 6500\}$  and  $\Lambda_{\mathrm{O}} = \{4001, \ldots, 4500, 8001, \ldots, 8500\}$ . Same as model (c), we consider  $\theta_1^* = 1$  and  $\theta_2^* \in \{0, 0.05, \ldots, 0.5\}$ .
- (h) Dependent: same as model (g) but  $(z_{k,1}, z_{k,2}, \nu_k, \varepsilon_k)^{\mathsf{T}}$  follows the vector autoregressive model in model (d).

We estimate  $\theta_1^*$  at  $N_b = 500b$  for b = 1, 2, ..., 20 under the assumption that  $\mathbb{E}\{z_k(y_k - \theta_1^*x_k)\} = 0$ . There are abnormalities caused by a change point in parameter value and a misspecified model for  $b \in \{5, 13\}$  and  $b \in \{9, 17\}$ , respectively. To detect them, consider

- (a) Luo et al.: the offline detection statistic in (3.5) with 2SLS and two-step GMM as the initial estimator and the restricted estimator in (3.6), respectively. The Bartlett kernel estimator is used for inference and the weighting matrix.
- (b) OGMM I: the online detection statistic in (3.7) with 2SLS as the initial estimator. The OGMM summary statistics are based on  $D_1$  only (initial) or updated with data batches where abnormality is not detected (cumulative). Leung and Chan's (2025) estimator is used for inference and the weighting matrix.
- (c) OGMM O: same as OGMM I but (3.8) is used. We expect that OGMM O should be sensitive to model misspecification only.

We compare the empirical rejection rates and size-adjusted rejection rates of different proposals. Figure 5 reports the results for  $\theta_2^* = 0.2$  and for  $b \in \{5,9\}$ . It confirms that OGMM is able to preserve the statistical properties when we replace offline statistics with online counterparts in our framework. For example, OGMM O only rejects when



Figure 5: Online anomaly detection at 5% nominal level under the independent model (g) (upper panel) and dependent model (h) (lower panel). The dotted and dashed lines in plots (i) and (iv) are the locations of abnormality caused by a change point in parameter value and a misspecified model, respectively. For  $\theta_2^* = 0$  or  $b \notin \{5,9,13,17\}$ , there is no abnormality so the rejection rates are ideally close to 5%. For  $\theta_2^* > 0$  and  $b \in \{5,9,13,17\}$ , there is abnormality so the tests are expected to reject.

b = 9 but not when b = 5, which helps identify the two possible causes of structural instability. We can also use more normal data to improve the sensitivity of OGMM I without worrying about the computational cost. Under model (g), there are 38% (initial) and 22.6% (cumulative) reductions in averaged time compared with Luo et al. (2022). Under model (h), the reductions change to 26.7% (initial) and 10% (cumulative) due to long-run variance estimation, but it allows autocorrelation-robust online testing. The results for other values of  $\theta_2^*$  or b are similar and deferred to the Supplement.

### 6 Applications

In Section 6.1, we consider the rolling approach of stochastic volatility modelling (Pascalau and Poirier, 2023) and find that our OGMM can be a handy addition to a forecaster's toolbox. In Section 6.2, we show that OGMM can be extended to the frequency domain and offers some alternative insights into the inertial sensor calibration problem.

### 6.1 Stochastic volatility modelling

Forecasting is a central topic in econometrics and statistics. With wider availability of high frequency data, practitioners are interested in updating models quickly to generate real-time prediction. However, traditional stochastic volatility models are usually trained with daily aggregated data. In light of it, Pascalau and Poirier (2023) proposed a rolling approach to incorporate intraday information. We consider different online estimation methods via OGMM here. Our data set consists of 7976478 one-minute bar euro dollar rate (EUR/USD) from 2000 to 2023 (UTC-5), which are freely available on HistData.com. We exclude 1681 observations in the weekend (UTC+2) and compute the close-to-close log 5-minute return  $r_{t,h,m}$  for m = 1, ..., 12 in the *h*-th hour on day *t*. Define the rolling daily realized variance and realized quarticity

$$\mathrm{RV}_{t,h} = \sum_{i=h-23}^{h} \sum_{j=1}^{12} r_{t,i,j}^2 \quad \text{and} \quad \mathrm{RQ}_{t,h} = \frac{24 \times 12}{3} \sum_{i=h-23}^{h} \sum_{j=1}^{12} r_{t,i,j}^4,$$

where  $r_{t,i-\ell,j} = r_{t-1,i+24-\ell,j}$  if  $i - \ell \in (-24, 0]$ , and  $r_{t,i+\ell,j} = r_{t+1,i-24+\ell,j}$  if  $i + \ell \in (24, 48]$ . We forecast the one-day-ahead daily realized variance  $RV_{t+1,24}$  using the following approaches with all available observations to date.

(a) Traditional: the quarticity-adjusted heterogeneous autoregressive model

$$RV_{t,24} = \theta_0^* + \left(\theta_1^* + \theta_2^* \sqrt{RQ_{t-1,24}}\right) RV_{t-1,24} + \theta_3^* \sum_{\ell=1}^5 \frac{RV_{t-\ell,24}}{5} + \theta_4^* \sum_{\ell=1}^{21} \frac{RV_{t-\ell,24}}{21} + \varepsilon_{t,24}$$
(6.1)

in Bollerslev et al. (2016) is estimated using ordinary least squares and updated daily. Following Pascalau and Poirier (2023), the initial sample size is 1000.

(b) Rolling: the model in (6.1) is modified to

$$RV_{t,h} = \theta_0^* + (\theta_1^* + \theta_2^* \sqrt{RQ_{t-1,h}}) RV_{t-1,h} + \theta_3^* \sum_{\ell=1}^5 \frac{RV_{t-\ell,h}}{5} + \theta_4^* \sum_{\ell=1}^{21} \frac{RV_{t-\ell,h}}{21} + \varepsilon_{t,h}$$
(6.2)

using the rolling approach (Pascalau and Poirier, 2023). Therefore, the initial sample size becomes  $1000 \times 24$ . Other settings are same as Traditional and we focus on forecasting RV<sub>t+1,24</sub> at day t's end.

(c) OGMM: consider (6.2) with different moment conditions. (LS) least squares; see Example 2. (QR) quantile regression with  $\tau = 0.5$ ; see Section 5.3. (IV) instrumental variables regression with 1,  $\sum_{\ell=1}^{5} \text{RV}_{t-\ell,h}/5$ ,  $\sum_{\ell=1}^{21} \text{RV}_{t-\ell,h}/21$  and  $\{\sqrt{\text{RQ}_{t-i,h}}\}_{i=k}^{21}$  as instruments, and 2SLS as the initial estimator; see Section 5.1. As Bollerslev et al. (2016) suggested that the measurement error in  $\text{RV}_{t-1,h}$  is most serious, we try to

	MSE rel	E relative to Traditional		Time relative to Traditional		
	Rolling	OGMM	SGMM	Rolling	OGMM	SGMM
LS	0.997	0.997	1.088	19.221	0.118	0.368
QR	—	0.994	1.076	—	0.251	0.661
IV $(k = 1)$	—	0.990	1.015	—	0.289	0.791
IV $(k = 2)$	—	0.990	1.019	—	0.270	0.789
IV $(k = 3)$	—	0.980	1.013	—	0.275	0.780
IV $(k = 4)$	—	0.970	1.008	—	0.250	0.793
IV $(k = 5)$	—	0.960	1.004	—	0.249	0.766

Table 2: One-day-ahead forecasting performances of (6.2) estimated using different methods. The mean squared errors and computation times are relative to (6.1).

use  $\{\sqrt{RQ_{t-i,h}}\}_{i=k}^{21}$  as instruments for terms that involve  $RV_{t-1,h}$ . The weighting matrix is inverted from Welford's (1962) estimator for (LS) and (QR), and Leung and Chan's (2025) estimator for (IV). Other settings are same as Rolling.

(d) SGMM: same as OGMM. The learning rate is chosen by (5.2) with  $\kappa = 0.5$  and a = 0.501.

Table 2 reports the mean squared error and computation time relative to the traditional approach to update models and generate predictions. It confirms that intraday information can be used to deliver more accurate forecasts, which is consistent with the findings in Pascalau and Poirier (2023). Furthermore, OGMM LS vastly improves the computational efficiency of the rolling approach by reducing its computing time by approximately 160 times. Meanwhile, OGMM QR and IV open the door to statistical refinements by further reducing the relative mean squared error from 0.997 to 0.994 and then to 0.96 (with k = 5). SGMM is unable to achieve them as its mean squared error is higher than that of the traditional method and its computational cost is approximately 2.5 to 3 times that of OGMM. Moreover, SGMM is affected by the choice of learning rate. In practice, OGMM allows us to conduct online estimation and inference easily given any moment condition. Users can maintain several sets of OGMM summary statistics and ensemble forecasts in real time. The efficiency and flexibility of OGMM are thus valuable in the analysis of large-scale streaming data.

### 6.2 Inertial sensor calibration

Inertial sensors are very common in modern vehicles and mobile robots. However, they may be corrupted by stochastic errors of complex structure, e.g.,

$$y_t = \sum_{i=1}^2 z_t^{(i)} + u_t^{(3)}, \tag{6.3}$$

	GMWM	OGMM
$\rho_1 \times 10^{-1}$	$9.592 \ (9.592, \ 9.592)$	9.910 (9.910, 9.910)
$\sigma_1^2 \times 10^{-12}$	23.217 (17.620, 28.814)	$4.191 \ (3.174, \ 5.209)$
$ ho_2  imes 10^{-2}$	$7.529\ (7.529,\ 7.529)$	$7.733\ (7.733,\ 7.733)$
$\sigma_2^2 \times 10^{-7}$	3.445(3.409, 3.482)	3.350(3.281, 3.419)
$\sigma_3^2  imes 10^{-8}$	$3.234\ (2.864,\ 3.605)$	4.177 (3.499, 4.855)
Estimation time (seconds)	298.46	11.19
Total time (seconds)	594.33	175.59
Sargan-Hansen test $(p$ -value)	0.000585	0.107

Table 3: Estimated parameters with associated 95% confidence intervals for (6.3).

where  $\{y_t\}$  are the observed error signals,  $z_t^{(i)} = \rho_i z_{t-1}^{(i)} + u_t^{(i)}$  follow latent first-order autoregressive processes, and  $u_1^{(i)}, u_2^{(i)}, \ldots \stackrel{\text{iid}}{\sim} N(0, \sigma_i^2)$  and are independent across *i* (Stebler et al., 2014). To estimate  $\theta^* = (\rho_1, \sigma_1^2, \rho_2, \sigma_2^2, \sigma_3^2)^{\mathsf{T}}$ , we can apply the generalized method of wavelet moments (GMWM, Guerrier et al., 2013). Specifically, let  $w_{t,j}$  be the maximal overlap discrete wavelet transform of  $y_t$  at scale  $\tau_j = 2^{j-1}$  using the Haar filter. Then, the *j*-th element of the moment function is

$$w_{t,j}^2 - \sum_{i=1}^2 \frac{(0.5\tau_j - 3\rho_i - 0.5\tau_j\rho_i^2 + 4\rho_i^{1+0.5\tau_j} - \rho_i^{1+\tau_j})\sigma_i^2}{0.5\tau_j^2(1 - \rho_i)^2(1 - \rho_i^2)} - \frac{\sigma_3^2}{\tau_j}, \quad \text{for} \quad j = 1, \dots, q;$$

see the R-package gmwm and Table 3 in Guerrier et al. (2013). When q > 5, the Sargan–Hansen test can be used to assess the goodness of fit of (6.3).

Now, consider the y-axis gyroscope data set available through imar.gyro in the R-package imudata. 5760000 error signals were collected at 400 Hz in 4 hours. We divide them into b = 48 equal batches (each of size 120000) and fit (6.3) using q = 10 and

- (a) GMWM: the two-step GMWM implemented in the R-package gmwm with full data. The default first-step weighting matrix is diagonal, which may not be optimal. Therefore, parametric bootstraps of size 100 are used for the second-step weighting matrix in estimation and the asymptotic variance in inference separately.
- (b) OGMM: OGMM with normal data batches. Using the two-step GMWM as the initial estimator, we update OGMM summary statistics if (3.7) and (3.8) are not rejected; see the cumulative strategy in Section 5.4.

Table 3 reports the estimation results. For GMWM, the Sargan-Hansen test suggests that there are abnormal data batches, or (6.3) is a poorly fitted model. To investigate the latter possibility, we also estimate (6.3-)  $y_t = z_t^{(1)} + u_t^{(2)}$  and (6.3+)  $y_t = \sum_{i=1}^3 z_t^{(i)} + u_t^{(4)}$ , which are considered in Guerrier et al. (2013) and Stebler et al. (2014). The Sargan-Hansen test *p*-values are (6.3-)  $1.021 \times 10^{-14}$  and (6.3+)  $8.822 \times 10^{-10}$ , which indicate worse fit. In contrast, applying GMWM and OGMM to the first data batch leads to *p*-values of 0.494 and 0.173, respectively. This evidence supports structural stability testing

in OGMM. Using (3.7) and (3.8), we find that 17 batches of data (number 10, 11, 13, 14, 15, 16, 17, 19, 21, 24, 25, 29, 36, 39, 41, 45, 46) are normal, and most abnormalities are due to model misspecification because (3.8) is rejected for 28 batches of data. The resulting OGMM estimate of  $\sigma_1^2$  is much smaller than GMWM, while estimates of other parameters are closer. The OGMM *p*-value 0.107 suggests better fit than GMWM, although it is not adjusted for possible post-screening issues. In terms of total time (estimation and testing), (3.8) requires solving an optimization problem for each data batch. Nevertheless, OGMM offers online estimates and remains faster than GMWM because it does not perform bootstrap for optimal estimation and inference. Overall, this application illustrates that the OGMM framework is very general and offers an alternative approach to estimation and inference for time series models formulated in wavelet domains.

### 7 Discussion

Time series data are inherently serially dependent and sequential. Motivated by the need to analyse large streaming time series data, we develop an online version of the classical GMM, which is robust to weak serial dependence, preserves the asymptotic efficiency of offline GMM, and vastly reduces the time complexity and memory requirements. Our methodological development covers OGMM point estimation, confidence region construction, over-identifying restrictions testing and anomaly detection with rigorous theoretical justification. Our numerical simulation and data illustration convincingly demonstrate the favourable performance of OGMM-based estimation and inference as compared to several existing procedures in terms of computational time and statistical errors.

To conclude, we mention a few future research topics. Throughout this paper the dimension of the parameter is assumed fixed. It would be interesting to develop an online version of GMM that can accommodate high dimensionality of the parameter; see Belloni et al. (2018) for regularized GMM in the offline setting. Additionally, GMM has been extended to handle the estimation of a finite dimensional parameter when a nuisance function is present; see Ackerberg et al. (2014) and Chen and Liao (2015). Developing an online version of two-step GMM to deal with this semiparametric setting would be interesting. Finally, it is of interest to allow for time-varying parameter (Luo et al., 2023) and/or local stationarity of streaming time series, as stationarity and time-invariant parameter might be too strong to be satisfied for a very long streaming time series.

### Acknowledgments

Chan was supported in part by grants GRF-14306421 and GRF-14307922 provided by the Research Grants Council of HKSAR. Shao would like to acknowledge partial financial support from National Science Foundation DMS-2210002 and DMS-2412833.

### References

- Ackerberg, D., Chen, X., Hahn, J. and Liao, Z. (2014) Asymptotic efficiency of semiparametric two-step GMM. *The Review of Economic Studies*, 81, 919–943.
- Andrews, D. W. K. (1991) Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica*, **59**, 817–858.
- Belloni, A., Chernozhukov, V., Chetverikov, D., Hansen, C. and Kato, K. (2018) Highdimensional econometrics and regularized GMM. Manuscript.
- Bollerslev, T., Patton, A. J. and Quaedvlieg, R. (2016) Exploiting the errors: A simple approach for improved volatility forecasting. *Journal of Econometrics*, **192**, 1–18.
- de Castro, L., Galvao, A. F., Kaplan, D. M. and Liu, X. (2019) Smoothed GMM for quantile models. *Journal of Econometrics*, **213**, 121–144.
- Chan, K. W. and Yau, C. Y. (2016) New recursive estimators of the time-average variance constant. *Statistics and Computing*, 26, 609–627.
- (2017) Automatic optimal batch size selection for recursive estimators of time-average covariance matrix. Journal of the American Statistical Association, **112**, 1076–1089.
- Chaussé, P. (2010) Computing generalized method of moments and generalized empirical likelihood with R. *Journal of Statistical Software*, **34**, 1–35.
- Chen, X., Lee, J. D., Tong, X. T. and Zhang, Y. (2020) Statistical inference for model parameters in stochastic gradient descent. *The Annals of Statistics*, **48**, 251–273.
- Chen, X., Lee, S., Liao, Y., Seo, M. H., Shin, Y. and Song, M. (2023) SGMM: Stochastic approximation to generalized method of moments. *Journal of Financial Econometrics*.
- Chen, X. and Liao, Z. (2015) Sieve semiparametric two-step GMM under weak dependence. Journal of Econometrics, 189, 163–186.
- Chen, X., Liu, W. and Zhang, Y. (2019) Quantile regression under memory constraint. The Annals of Statistics, 47, 3244–3273.

- Guerrier, S., Skaloud, J., Stebler, Y. and Victoria-Feser, M.-P. (2013) Wavelet-variancebased estimation for composite stochastic processes. *Journal of the American Statistical* Association, 108, 1021–1030.
- Hall, A. R. (2000) Covariance matrix estimation and the power of the overidentifying restrictions test. *Econometrica*, **68**, 1517–1527.
- (2005) Generalized Method of Moments. Oxford University Press.
- Hall, A. R. and Sen, A. (1999) Structural stability testing in models estimated by generalized method of moments. *Journal of Business & Economic Statistics*, **17**, 335–348.
- Hansen, L. P. (1982) Large sample properties of generalized method of moments estimators. *Econometrica*, **50**, 1029–1054.
- Jiang, R. and Yu, K. (2022) Renewable quantile regression for streaming data sets. Neurocomputing, 508, 208–224.
- (2024) Unconditional quantile regression for streaming datasets. Journal of Business
   & Economic Statistics, to appear.
- Koenker, R. (2005) Quantile Regression. Cambridge University Press.
- Le Cam, L. (1956) On the asymptotic theory of estimation and testing hypotheses. In Proceedings of the Berkeley Symposium on Mathematical Statistics and Probability, vol. 1, 129–156. University of California Press.
- Lee, S., Liao, Y., Seo, M. H. and Shin, Y. (2022) Fast and robust online inference with stochastic gradient descent via random scaling. *Proceedings of the AAAI Conference* on Artificial Intelligence, 36, 7381–7389.
- Leung, M. F. and Chan, K. W. (2025) Principles of statistical inference in online problems. Manuscript available at https://arxiv.org/abs/2209.05399.
- Leung, M. F., Chan, K. W. and Shao, X. (2025) Supplement to "online generalized method of moments for time series".
- Luo, L. and Song, P. X.-K. (2020) Renewable estimation and incremental inference in generalized linear models with streaming data sets. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 82, 69–97.
- Luo, L., Wang, J. and Hector, E. C. (2023) Statistical inference for streamed longitudinal data. *Biometrika*, **110**, 841–858.

- Luo, L., Zhou, L. and Song, P. X.-K. (2022) Real-time regression analysis of streaming clustered data with possible abnormal data batches. *Journal of the American Statistical Association*, **118**, 2029–2044.
- Meketon, M. S. and Schmeiser, B. (1984) Overlapping batch means: something for nothing? In *Proceedings of the Winter Simulation Conference*, WSC '84, 226–230.
- Newey, W. K. and McFadden, D. (1994) Large sample estimation and hypothesis testing. vol. 4 of *Handbook of Econometrics*, chap. 36, 2111–2245. Elsevier.
- Parzen, E. (1957) On consistent estimates of the spectrum of a stationary time series. The Annals of Mathematical Statistics, 28, 329–348.
- Pascalau, R. and Poirier, R. (2023) Increasing the information content of realized volatility forecasts. *Journal of Financial Econometrics*, **21**, 1064–1098.
- Polyak, B. T. and Juditsky, A. B. (1992) Acceleration of stochastic approximation by averaging. SIAM Journal on Control and Optimization, 30, 838–855.
- Qu, A., Lindsay, B. G. and Li, B. (2000) Improving generalised estimating equations using quadratic inference functions. *Biometrika*, 87, 823–836.
- Robbins, H. and Monro, S. (1951) A stochastic approximation method. The Annals of Mathematical Statistics, 22, 400–407.
- Schifano, E. D., Wu, J., Wang, C., Yan, J. and Chen, M.-H. (2016) Online updating of statistical inference in the big data setting. *Technometrics*, 58, 393–403.
- Stebler, Y., Guerrier, S., Skaloud, J. and Victoria-Feser, M.-P. (2014) Generalized method of wavelet moments for inertial navigation filter design. *IEEE Transactions on Aerospace and Electronic Systems*, 50, 2269–2283.
- Sun, X., Wang, H., Cai, C., Yao, M. and Wang, K. (2024) Online renewable smooth quantile regression. *Computational Statistics & Data Analysis*, 185, 107781.
- Toulis, P. and Airoldi, E. M. (2017) Asymptotic and finite-sample properties of estimators based on stochastic gradients. *The Annals of Statistics*, **45**, 1694–1727.
- Welford, B. P. (1962) Note on a method for calculating corrected sums of squares and products. *Technometrics*, **4**, 419–420.
- Wu, W. B. (2009) Recursive estimation of time-average variance constants. The Annals of Applied Probability, 19, 1529–1552.
- Zhu, W., Chen, X. and Wu, W. B. (2023) Online covariance matrix estimation in stochastic gradient descent. *Journal of the American Statistical Association*, **118**, 393–404.

### A Proof of theorems

### A.1 Proof of Theorem 1

Theorem 3.1 in Hall (2005) states that  $\hat{\theta}_1 \xrightarrow{p} \theta^*$  under Assumptions 1–4 and 6–9. We do not separate the remaining proof by cases (a) and (b) because they only differ by the use of Lemma 1(a) and (b).

To prove by induction, we first consider b = 2. Recall that  $\hat{W} \xrightarrow{p} W$  under Assumption 6, and  $\hat{V}_2 \xrightarrow{p} V$  and  $\hat{U}_2 \xrightarrow{p} -V\theta^*$  by Lemma 1. Applying Slutsky's theorem gives

$$\hat{\theta}_2 = -(\hat{V}_2^{\mathsf{T}}\hat{W}\hat{V}_2)^{-1}\hat{V}_2^{\mathsf{T}}\hat{W}\hat{U}_2 \xrightarrow{\mathrm{p}} (V^{\mathsf{T}}WV)^{-1}V^{\mathsf{T}}WV\theta^* = \theta^*.$$

Now, suppose that  $\|\hat{\theta}_i - \theta^*\| = o_p(1)$  and  $\hat{\theta}_i \in B_{\epsilon}$  for  $i = 1, \ldots, b-1$ . Then, we have  $\hat{V}_b \xrightarrow{p} V$  and  $\hat{U}_b \xrightarrow{p} -V\theta^*$  by Lemma 1 again. By the same argument,

$$\hat{\theta}_b = -(\hat{V}_b^{\mathsf{T}} \hat{W} \hat{V}_b)^{-1} \hat{V}_b^{\mathsf{T}} \hat{W} \hat{U}_b \xrightarrow{\mathrm{p}} (V^{\mathsf{T}} W V)^{-1} V^{\mathsf{T}} W V \theta^* = \theta^*.$$

In the above argument, we assume that  $\hat{\theta}_i \in B_{\epsilon}$  for  $i = 1, \dots, b-1$  to simplify the proof. Technically speaking, we can partition the probability space into  $\Omega_{b-1} = \bigcup_{i=1}^{b-1} \{\hat{\theta}_i \notin B_{\epsilon}\}$  and its complement  $\Omega_{b-1}^c$ . On  $\Omega_{b-1}^c$ , all the above discussion can go through under the assumptions stated for a local neighborhood  $B_{\epsilon}$ . Hence all statements that hold in probability are still valid since  $P(\Omega_{b-1}) \leq \sum_{i=1}^{b-1} P(\hat{\theta}_i \notin B_{\epsilon}) \to 0$ ; see Section 3.5 of Newey and McFadden (1994) for an example with  $\hat{\theta}_1$  only. In the following proofs, we shall not repeat this argument but instead assume  $\hat{\theta}_i \in B_{\epsilon}$  for  $i = 1, \dots, b-1$  to simplify the presentation.

### A.2 Proof of Theorem 2

Recall that Assumptions 1–10 are identical to Assumptions 3.1–3.5 and 3.7–3.11 in Hall (2005), and both cases (a) and (b) in Assumptions 11 and 12(ii) imply Assumptions 3.12 and 3.13 in Hall (2005), respectively. By Theorem 3.2 in Hall (2005), we have  $\sqrt{N_1}(\hat{\theta}_1 - \theta^*) \xrightarrow{d} N(0, H\Sigma H^{\intercal})$ . However, an asymptotically efficient  $\hat{\theta}_1$  is not necessary in the remaining proof. Any  $\sqrt{N_1}$ -consistent  $\hat{\theta}_1$  will suffice.

(Case a). To prove by induction, we first consider b = 2. By the mean value theorem,

$$\bar{g}_2(\hat{\theta}_2) = \bar{g}_2(\theta^*) + \nabla \bar{g}_2(\bar{\theta}_2)(\hat{\theta}_2 - \theta^*),$$

where  $\bar{\theta}_b$  is a point between  $\hat{\theta}_b$  and  $\theta^*$ . Multiplying by  $\sqrt{N_2}\hat{V}_2^{\dagger}\hat{W}$  and rearranging,

$$\sqrt{N_2}(\hat{\theta}_2 - \theta^*) = \{\hat{V}_2^{\mathsf{T}} \hat{W} \nabla \bar{g}_2(\bar{\theta}_2)\}^{-1} \hat{V}_2^{\mathsf{T}} \hat{W} \sqrt{N_2} \bar{g}_2(\hat{\theta}_2) 
- \{\hat{V}_2^{\mathsf{T}} \hat{W} \nabla \bar{g}_2(\bar{\theta}_2)\}^{-1} \hat{V}_2^{\mathsf{T}} \hat{W} \sqrt{N_2} \bar{g}_2(\theta^*) 
= J_{2,1} + J_{2,2}.$$
(A.1)

Using (C.3) and the condition that  $\hat{V}_2^{\dagger}\hat{W}(\hat{U}_2 + \hat{V}_2\hat{\theta}_2) = 0$ , the first term simplifies to

$$J_{2,1} = \sqrt{N_2} \{ \hat{V}_2^{\mathsf{T}} \hat{W} \nabla \bar{g}_2(\bar{\theta}_2) \}^{-1} \hat{V}_2^{\mathsf{T}} \hat{W} \left\{ \hat{U}_2 + \hat{V}_2 \hat{\theta}_2 + O_p \left( \frac{\sqrt{n_1} + \sqrt{n_2}}{N_2} \left\| \hat{\theta}_1 - \hat{\theta}_2 \right\| + \left\| \hat{\theta}_1 - \hat{\theta}_2 \right\|^2 \right) \right\}$$
$$= \{ \hat{V}_2^{\mathsf{T}} \hat{W} \nabla \bar{g}_2(\bar{\theta}_2) \}^{-1} \hat{V}_2^{\mathsf{T}} \hat{W} O_p \left( \frac{\sqrt{n_1} + \sqrt{n_2}}{\sqrt{N_2}} \left\| \hat{\theta}_1 - \hat{\theta}_2 \right\| + \sqrt{N_2} \left\| \hat{\theta}_1 - \hat{\theta}_2 \right\|^2 \right) .$$

Check that

$$\hat{\theta}_2 = -(\hat{V}_2^{\mathsf{T}}\hat{W}\hat{V}_2)^{-1}\hat{V}_2^{\mathsf{T}}\hat{W}\hat{U}_2 = \hat{\theta}_1 - (\hat{V}_2^{\mathsf{T}}\hat{W}\hat{V}_2)^{-1}\hat{V}_2^{\mathsf{T}}\hat{W}\frac{1}{N_2}\left\{G(\hat{\theta}_1; D_1) + G(\hat{\theta}_1; D_2)\right\}.$$
 (A.2)

Using the same trick in (C.2) and Assumption 5(c), we have

$$G(\hat{\theta}_{1}; D_{1}) = G(\theta^{*}; D_{1}) + \nabla G(\theta^{*}; D_{1})(\hat{\theta}_{1} - \theta^{*}) + O_{p}\left(\sqrt{n_{1}} \|\hat{\theta}_{1} - \theta^{*}\| + n_{1} \|\hat{\theta}_{1} - \theta^{*}\|^{2}\right)$$
  
$$= G(\theta^{*}; D_{1}) + n_{1}\mathbb{E}\{\nabla g(\theta^{*}, x_{i,j})\}(\hat{\theta}_{1} - \theta^{*}) + O_{p}\left(\sqrt{n_{1}} \|\hat{\theta}_{1} - \theta^{*}\| + n_{1} \|\hat{\theta}_{1} - \theta^{*}\|^{2}\right)$$
  
$$= G(\theta^{*}; D_{1}) + O_{p}\left(n_{1} \|\hat{\theta}_{1} - \theta^{*}\|\right).$$
(A.3)

Therefore, by the central limit theorem for  $\bar{g}_2(\theta^*)$  (see, e.g., Lemma 3.2 in Hall 2005), and  $\|\hat{\theta}_1 - \theta^*\| = O_p(N_1^{-1/2})$ ,

$$\hat{\theta}_{2} - \hat{\theta}_{1} = -(\hat{V}_{2}^{\top} \hat{W} \hat{V}_{2})^{-1} \hat{V}_{2}^{\top} \hat{W} \frac{1}{N_{2}} \left\{ G(\hat{\theta}_{1}; D_{1}) + G(\hat{\theta}_{1}; D_{2}) \right\} = -(\hat{V}_{2}^{\top} \hat{W} \hat{V}_{2})^{-1} \hat{V}_{2}^{\top} \hat{W} \bar{g}_{2}(\theta^{*}) + O_{p}(\|\hat{\theta}_{1} - \theta^{*}\|) = O_{p}(N_{2}^{-1/2} + N_{1}^{-1/2}) = O_{p}(N_{1}^{-1/2}).$$
(A.4)

Using the same procedure that leads to (C.1), we have  $\nabla \bar{g}_2(\bar{\theta}_2) \xrightarrow{p} V$ . Moreover, recall that  $\hat{W} \xrightarrow{p} W$  under Assumption 6 and  $\hat{V}_2 \xrightarrow{p} V$  by Lemma 1(a). Therefore,  $J_{2,1} = o_p(1)$ if and only if  $\sqrt{N_2} = o(N_1) \Leftrightarrow n_2 = o(N_1^2)$ . For  $J_{2,2}$ , the central limit theorem for  $\bar{g}_2(\theta^*)$ and Slutsky's theorem gives  $J_{2,2} \xrightarrow{d} N(0, H\Sigma H^{\intercal})$ . Combining the results with Slutsky's theorem, we have  $\sqrt{N_2}(\hat{\theta}_2 - \theta^*) \xrightarrow{d} N(0, H\Sigma H^{\intercal})$ . Now, suppose that  $\|\hat{\theta}_i - \theta^*\| = O_p(N_i^{-1/2})$ for  $i = 1, \ldots, b - 1$ . Using the same procedure that leads to (A.1), we have

$$\sqrt{N_b}(\hat{\theta}_b - \theta^*) = \{\hat{V}_b^{\mathsf{T}} \hat{W} \nabla \bar{g}_b(\bar{\theta}_b)\}^{-1} \hat{V}_b^{\mathsf{T}} \hat{W} \sqrt{N_b} \bar{g}_b(\hat{\theta}_b) - \{\hat{V}_b^{\mathsf{T}} \hat{W} \nabla \bar{g}_b(\bar{\theta}_b)\}^{-1} \hat{V}_b^{\mathsf{T}} \hat{W} \sqrt{N_b} \bar{g}_b(\theta^*) = J_{b,1} + J_{b,2}.$$
(A.5)

By (C.3) and the condition that  $\hat{V}_b^{\top}\hat{W}(\hat{U}_b + \hat{V}_b\hat{\theta}_b) = 0$ ,

$$J_{b,1} = \{ \hat{V}_b^{\mathsf{T}} \hat{W} \nabla \bar{g}_b(\bar{\theta}_b) \}^{-1} \hat{V}_b^{\mathsf{T}} \hat{W} O_p \left( \frac{\sqrt{n_b}}{\sqrt{N_b}} \| \hat{\theta}_{b-1} - \hat{\theta}_b \| + \sum_{i=1}^{b-1} \frac{\sqrt{n_i}}{\sqrt{N_b}} \| \hat{\theta}_i - \hat{\theta}_b \| \right) \\ + \{ \hat{V}_b^{\mathsf{T}} \hat{W} \nabla \bar{g}_b(\bar{\theta}_b) \}^{-1} \hat{V}_b^{\mathsf{T}} \hat{W} O_p \left( \frac{n_b}{\sqrt{N_b}} \| \hat{\theta}_{b-1} - \hat{\theta}_b \|^2 + \sum_{i=1}^{b-1} \frac{n_i}{\sqrt{N_b}} \| \hat{\theta}_i - \hat{\theta}_b \|^2 \right).$$

For  $\|\hat{\theta}_{b-1} - \hat{\theta}_b\|$ , there is no direct formula like (A.2). Therefore, we bound the difference in another way. Using the trick in (C.2) and (A.3),

$$\nabla G(\hat{\theta}_i; D_i) = \nabla G(\theta^*; D_i) + \nabla G(\hat{\theta}_i; D_i) - \nabla G(\theta^*; D_i)$$
$$= n_i \mathbb{E} \{ \nabla g(\theta^*, x_{i,j}) \} + O_p(n_i^{1/2} + \|\hat{\theta}_i - \theta^*\|) = O_p(n_i)$$

By the triangle inequality and  $\|\hat{\theta}_i - \theta^*\| = O_p(N_i^{-1/2})$  for  $i = 1, \dots, b-1$ ,

$$\begin{split} & \left\| \hat{U}_{b} - \left[ \frac{1}{N_{b}} \left\{ \sum_{i=1}^{b-1} G(\hat{\theta}_{i}; D_{i}) + G(\hat{\theta}_{b-1}; D_{b}) \right\} - \hat{V}_{b} \hat{\theta}_{b-1} \right] \right\| \\ &= \left\| \frac{1}{N_{b}} \sum_{i=1}^{b-2} \nabla G(\hat{\theta}_{i}; D_{i}) (\hat{\theta}_{b-1} - \hat{\theta}_{i}) \right\| \\ &\leq \left\| \frac{1}{N_{b}} \sum_{i=1}^{b-2} \nabla G(\hat{\theta}_{i}; D_{i}) (\hat{\theta}_{i} - \theta^{*}) \right\| + \left\| \frac{1}{N_{b}} \sum_{i=1}^{b-2} \nabla G(\hat{\theta}_{i}; D_{i}) (\hat{\theta}_{b-1} - \theta^{*}) \right\| \\ &= O_{p} \left( \sum_{i=1}^{b-2} \frac{n_{i}}{N_{b}} \left\| \hat{\theta}_{i} - \theta^{*} \right\| \right) = O_{p} \left( \sum_{i=1}^{b-2} \frac{n_{i}}{N_{b} \sqrt{N_{i}}} \right). \end{split}$$

Applying the trick in (A.3) again and the central limit theorem for  $\bar{g}_b(\theta^*)$ ,

$$\begin{split} \hat{\theta}_{b} - \hat{\theta}_{b-1} &= -(\hat{V}_{b}^{\top} \hat{W} \hat{V}_{b})^{-1} \hat{V}_{b}^{\top} \hat{W} \hat{U}_{b} - (\hat{V}_{b}^{\top} \hat{W} \hat{V}_{b})^{-1} \hat{V}_{b}^{\top} \hat{W} \hat{V}_{b} \hat{\theta}_{b-1} \\ &= -(\hat{V}_{b}^{\top} \hat{W} \hat{V}_{b})^{-1} \hat{V}_{b}^{\top} \hat{W} \frac{1}{N_{b}} \left\{ \sum_{i=1}^{b-1} G(\hat{\theta}_{i}; D_{i}) + G(\hat{\theta}_{b-1}; D_{b}) \right\} + O_{p} \left( \sum_{i=1}^{b-2} \frac{n_{i}}{N_{b} \sqrt{N_{i}}} \right) \\ &= -(\hat{V}_{b}^{\top} \hat{W} \hat{V}_{b})^{-1} \hat{V}_{b}^{\top} \hat{W} \bar{g}_{b} (\theta^{*}) + O_{p} \left( \frac{n_{b}}{N_{b} \sqrt{N_{b-1}}} + \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b} \sqrt{N_{i}}} \right) \\ &= O_{p} \left( \frac{1}{\sqrt{N_{b}}} + \frac{n_{b}}{N_{b} \sqrt{N_{b-1}}} + \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b} \sqrt{N_{i}}} \right). \end{split}$$

By Lemma 2,

$$\frac{1}{\sqrt{N_b}} + \frac{n_b}{N_b\sqrt{N_{b-1}}} + \sum_{i=1}^{b-1} \frac{n_i}{N_b\sqrt{N_i}} \le \frac{1}{\sqrt{N_b}} + \frac{n_b}{N_b\sqrt{N_{b-1}}} + \frac{2\sqrt{N_{b-1}}}{N_b} \le \frac{3}{\sqrt{N_b}} + \frac{n_b}{N_b\sqrt{N_{b-1}}}.$$

Therefore,

$$\left\|\hat{\theta}_{b-1} - \hat{\theta}_b\right\| = O_p\left(\frac{n_b}{N_b\sqrt{N_{b-1}}}\right). \tag{A.6}$$

Using  $\|\hat{\theta}_i - \theta^*\| = O_p(N_i^{-1/2})$  for  $i = 1, \dots, b-1$ , a direct consequence is that

$$\|\hat{\theta}_{i} - \hat{\theta}_{b}\| \leq \|\hat{\theta}_{i} - \theta^{*}\| + \|\theta^{*} - \hat{\theta}_{b-1}\| + \|\hat{\theta}_{b-1} - \hat{\theta}_{b}\| = O_{p}(N_{i}^{-1/2}).$$

We are ready to bound  $J_{b,1}$ . By the triangle inequality, (A.6) and Lemma 2,

$$\frac{\sqrt{n_b}}{\sqrt{N_b}} \left\| \hat{\theta}_{b-1} - \hat{\theta}_b \right\| + \sum_{i=1}^{b-1} \frac{\sqrt{n_i}}{\sqrt{N_b}} \left\| \hat{\theta}_i - \hat{\theta}_b \right\| = O_p \left( \frac{1}{\sqrt{N_{b-1}}} + \sum_{i=1}^{b-1} \frac{\sqrt{n_i}}{\sqrt{N_i N_b}} \right)$$

$$= O_p \left\{ \frac{1}{\sqrt{N_{b-1}}} + \frac{1}{\sqrt{\min(n_1, \dots, n_b)}} \right\} = o_p(1).$$

Similarly, we can apply the triangle inequality to obtain

$$\begin{aligned} &\frac{n_b}{\sqrt{N_b}} \left\| \hat{\theta}_{b-1} - \hat{\theta}_b \right\|^2 + \sum_{i=1}^{b-1} \frac{n_i}{\sqrt{N_b}} \left\| \hat{\theta}_i - \hat{\theta}_b \right\|^2 \\ \leq &\sqrt{N_b} \left\| \hat{\theta}_{b-1} - \hat{\theta}_b \right\|^2 + 2 \left\| \hat{\theta}_{b-1} - \hat{\theta}_b \right\| \sum_{i=1}^{b-1} \frac{n_i}{\sqrt{N_b}} \left\| \hat{\theta}_i - \hat{\theta}_{b-1} \right\| + \sum_{i=1}^{b-1} \frac{n_i}{\sqrt{N_b}} \left\| \hat{\theta}_i - \hat{\theta}_{b-1} \right\|^2. \end{aligned}$$

By (A.6) and Lemma 2, we have

$$\left\|\hat{\theta}_{b-1} - \hat{\theta}_{b}\right\| \sum_{i=1}^{b-1} \frac{n_{i}}{\sqrt{N_{b}}} \left\|\hat{\theta}_{i} - \hat{\theta}_{b-1}\right\| = o_{p}(1)O_{p}\left(\sum_{i=1}^{b-1} \frac{n_{i}}{\sqrt{N_{i}N_{b}}}\right) = o_{p}(1)O_{p}\left(\frac{2\sqrt{N_{b}}}{\sqrt{N_{b}}}\right) = O_{p}(1)O_$$

and

$$\sum_{i=1}^{b-1} \frac{n_i}{\sqrt{N_b}} \left\| \hat{\theta}_i - \hat{\theta}_{b-1} \right\|^2 = O_p \left( \sum_{i=1}^{b-1} \frac{n_i}{N_i \sqrt{N_b}} \right) = O_p \left( \frac{1 + \log N_b - \log n_1}{\sqrt{N_b}} \right) = o_p(1).$$

The conclusion then follows by the same argument for b = 2 as we can verify that

$$J_{b,1} = O_p\left(\sqrt{N_b} \left\|\hat{\theta}_{b-1} - \hat{\theta}_b\right\|^2\right) = O_p\left(\frac{n_b^2}{N_b^{3/2}N_{b-1}}\right) = o_p(1)$$

when  $n_b = o(N_{b-1}^2)$ .

(Case b). The proof is very similar to case (a) so we only highlight the difference. When b = 2, we still have (A.1). Applying (C.6) and the condition that  $\hat{V}_2^{\dagger}\hat{W}(\hat{U}_2+\hat{V}_2\hat{\theta}_2) = 0$  to  $J_{2,1}$ , we have

$$J_{2,1} = \{ \hat{V}_2^{\mathsf{T}} \hat{W} \nabla \bar{g}_2(\bar{\theta}_2) \}^{-1} \hat{V}_2^{\mathsf{T}} \hat{W} O_p \left( \sqrt{N_2} \left\| \hat{\theta}_1 - \hat{\theta}_2 \right\|^2 \right).$$

The formula in (A.2) and conclusion in (A.4) remain unchanged. However, the argument for (A.3) is slightly different. By (C.5), Assumptions 5(c) and 12(b)(i), we have

$$\frac{1}{N_2} \sum_{i=1}^2 G(\hat{\theta}_1; D_i) = \bar{g}_2(\theta^*) + \nabla \bar{g}_2(\theta^*)(\hat{\theta}_1 - \theta^*) + O_p\left(\left\|\hat{\theta}_1 - \theta^*\right\|^2\right) \\ = \bar{g}_2(\theta^*) + \mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}(\hat{\theta}_1 - \theta^*) + O_p\left(\left\|\hat{\theta}_1 - \theta^*\right\| + \left\|\hat{\theta}_1 - \theta^*\right\|^2\right) \\ = \bar{g}_2(\theta^*) + O_p\left(\left\|\hat{\theta}_1 - \theta^*\right\|\right).$$

Using the same procedure that leads to (C.4), we have  $\nabla \bar{g}_2(\bar{\theta}_2) \xrightarrow{p} V$ . The remaining arguments are the same. For b > 2, check that

$$J_{b,1} = \{ \hat{V}_b^{\mathsf{T}} \hat{W} \nabla \bar{g}_b(\bar{\theta}_b) \}^{-1} \hat{V}_b^{\mathsf{T}} \hat{W} O_p \left( \frac{n_b}{\sqrt{N_b}} \left\| \hat{\theta}_{b-1} - \hat{\theta}_b \right\|^2 + \sum_{i=1}^{b-1} \frac{n_i}{\sqrt{N_b}} \left\| \hat{\theta}_i - \hat{\theta}_b \right\|^2 \right).$$

The other arguments are same as case (a) or those in the above for b = 2.

### A.3 Proof of Theorem 3

The proof of Theorem 3 is based on the results in the proof of Theorem 2. Therefore, we do not separate the proof by cases (a) and (b) but refer to the relevant results instead. By Theorem 3.3 in Hall (2005), we have  $\hat{W}^{1/2}\sqrt{N_1}(U_1 + V_1\hat{\theta}_1) = \hat{W}^{1/2}\sqrt{N_1}\bar{g}_1(\hat{\theta}_1) \xrightarrow{d} N(0, RW^{1/2}\Sigma(W^{1/2})^{\intercal}R^{\intercal})$ , where the square root of a positive semi-definite matrix A is defined as the unique positive semi-definite matrix B such that  $BB = B^{\intercal}B = A$ . For b = 2, by the trick in (C.2) and the result in (A.4),

$$\hat{W}^{1/2}\sqrt{N_2}(U_2 + V_2\hat{\theta}_2) = \hat{W}^{1/2}\sqrt{N_2}\bar{g}_2(\hat{\theta}_2) + O_p\left(\frac{n_1}{\sqrt{N_2}} \left\|\hat{\theta}_1 - \hat{\theta}_2\right\|^2\right)$$
$$= \hat{W}^{1/2}\sqrt{N_2}\bar{g}_2(\hat{\theta}_2) + o_p(1).$$

It follows from the mean value theorem and (A.1) that

$$\begin{split} \hat{W}^{1/2} \sqrt{N_2} (U_2 + V_2 \hat{\theta}_2) \\ &= \hat{W}^{1/2} \sqrt{N_2} \bar{g}_2(\theta^*) + \hat{W}^{1/2} \nabla \bar{g}_2(\bar{\theta}_2) \sqrt{N_2} (\hat{\theta}_2 - \theta^*) + o_p(1) \\ &= \left[ I_q - \hat{W}^{1/2} \nabla \bar{g}_2(\bar{\theta}_2) \{ \hat{V}_2^{\mathsf{T}} \hat{W} \nabla \bar{g}_2(\bar{\theta}_2) \}^{-1} \hat{V}_2^{\mathsf{T}} (\hat{W}^{1/2})^{\mathsf{T}} \right] \hat{W}^{1/2} \sqrt{N_2} \bar{g}_2(\theta^*) + o_p(1). \end{split}$$

Recall that  $\hat{W} \xrightarrow{p} W$ ,  $\hat{V}_2 \xrightarrow{p} V$  and  $\nabla \bar{g}_2(\bar{\theta}_2) \xrightarrow{p} V$  in the proof of Theorem 2. Therefore, the central limit theorem for  $\bar{g}_2(\theta^*)$  and Slutsky's theorem gives  $\hat{W}^{1/2}\sqrt{N_2}(U_2 + V_2\hat{\theta}_2) \xrightarrow{d} N(0, RW^{1/2}\Sigma(W^{1/2})^{\intercal}R^{\intercal})$ . For b > 2, by the trick in (C.2) and the result in (A.6),

$$\begin{split} \hat{W}^{1/2} \sqrt{N_b} (U_b + V_b \hat{\theta}_b) &= \hat{W}^{1/2} \sqrt{N_b} \bar{g}_b(\hat{\theta}_b) + O_p \left( \sum_{i=1}^{b-1} \frac{n_i}{\sqrt{N_b}} \left\| \hat{\theta}_i - \hat{\theta}_b \right\|^2 \right) \\ &= \hat{W}^{1/2} \sqrt{N_b} \bar{g}_b(\hat{\theta}_b) + o_p(1). \end{split}$$

By the mean value theorem and (A.5),

$$\begin{split} \hat{W}^{1/2} \sqrt{N_b} (U_b + V_b \hat{\theta}_b) \\ &= \hat{W}^{1/2} \sqrt{N_b} \bar{g}_b(\theta^*) + \hat{W}^{1/2} \nabla \bar{g}_b(\bar{\theta}_b) \sqrt{N_b} (\hat{\theta}_b - \theta^*) + o_p(1) \\ &= \left[ I_q - \hat{W}^{1/2} \nabla \bar{g}_b(\bar{\theta}_b) \{ \hat{V}_b^{\mathsf{T}} \hat{W} \nabla \bar{g}_b(\bar{\theta}_b) \}^{-1} \hat{V}_b^{\mathsf{T}} (\hat{W}^{1/2})^{\mathsf{T}} \right] \hat{W}^{1/2} \sqrt{N_b} \bar{g}_b(\theta^*) + o_p(1). \end{split}$$

The conclusion then follows by the same argument for b = 2. When  $\hat{\Sigma}$  is positive semidefinite and converges in probability to  $\Sigma$  in Assumption 10, the asymptotic distribution of  $N_b(U_b + V_b\hat{\theta}_b)^{\mathsf{T}}\hat{\Sigma}^{-1}(U_b + V_b\hat{\theta}_b)$  is a direct consequence of the asymptotic normality of  $\hat{W}^{1/2}\sqrt{N_b}(U_b + V_b\hat{\theta}_b)$  with  $\hat{W} = \hat{\Sigma}^{-1}$ ; see the proof of Theorem 5.1 in Hall (2005).

### **B** Proof of Proposition 1

To prove by induction, we first consider b = 2. Since  $\hat{\theta}'_1 = \hat{\theta}_1$ , we have  $\hat{V}'_2 = \hat{V}_2$  and

$$\hat{U}_{2}' = \frac{1}{N_{2}} \left\{ N_{1}U_{1}' + G(\hat{\theta}_{1}'; D_{2}) \right\} = \frac{1}{N_{2}} \left\{ G(\hat{\theta}_{1}; D_{1}) + G(\hat{\theta}_{1}; D_{2}) \right\}.$$

It follows from (A.2) that  $\hat{\theta}_2' \equiv \hat{\theta}_2$  and

$$U_{2}' = \frac{1}{N_{2}} \left\{ N_{1}U_{1}' + N_{1}V_{1}'(\hat{\theta}_{2}' - \hat{\theta}_{1}') + G(\hat{\theta}_{2}'; D_{2}) \right\} \equiv U_{2} + V_{2}\hat{\theta}_{2}.$$

Now, suppose  $\hat{\theta}'_i \equiv \hat{\theta}_i$  and  $U'_i \equiv U_i + V_i \hat{\theta}_i$  for  $i = 1, \dots, b-1$ . Check that  $\hat{V}'_b = \hat{V}_b$  and

$$\hat{U}_{b} + \hat{V}_{b}\hat{\theta}_{b-1} = \frac{1}{N_{b}} \left\{ N_{b-1}U_{b-1} + N_{b-1}V_{b-1}\hat{\theta}_{b-1} + G(\hat{\theta}_{b-1}; D_{b}) \right\}$$
$$= \frac{1}{N_{b}} \left\{ N_{b-1}U_{b-1}' + G(\hat{\theta}_{b-1}'; D_{b}) \right\} = \hat{U}_{b}'.$$

Therefore,

$$\begin{aligned} \hat{\theta}_{b} &= -(\hat{V}_{b}^{\top} \hat{W} \hat{V}_{b})^{-1} \hat{V}_{b}^{\top} \hat{W} \hat{U}_{b} \\ &= -\{(\hat{V}_{b}')^{\top} \hat{W} \hat{V}_{b}'\}^{-1} (\hat{V}_{b}')^{\top} \hat{W} (\hat{U}_{b}' - \hat{V}_{b}' \hat{\theta}_{b-1}') \\ &= \hat{\theta}_{b-1}' - \{(\hat{V}_{b}')^{\top} \hat{W} \hat{V}_{b}'\}^{-1} (\hat{V}_{b}')^{\top} \hat{W} \hat{U}_{b}' \equiv \hat{\theta}_{b}', \end{aligned}$$

and similarly  $U_b' \equiv U_b + V_b \hat{\theta}_b$ .

## C Proof of lemmas

### C.1 Lemma 1

**Lemma 1** (Convergence of  $\hat{U}_b$  and  $\hat{V}_b$ ). Suppose Assumptions 1–9 hold.

- (a) Under Assumptions 11(a) and 12(a)(i),  $\hat{V}_b \xrightarrow{p} V = \mathbb{E}\{\nabla g(\theta^*, x_{i,j})\}$  and  $\hat{U}_b \xrightarrow{p} -V\theta^*$ as  $\min(n_1, \ldots, n_b) \to \infty$ .
- (b) Under Assumptions 11(b) and 12(b)(i),  $\hat{V}_b \xrightarrow{p} V$  and  $\hat{U}_b \xrightarrow{p} -V\theta^*$  as  $n_1 \to \infty$ .

*Proof.* We can prove this lemma under the condition that  $\|\hat{\theta}_i - \theta^*\| = o_p(1)$  and  $\hat{\theta}_i \in B_\epsilon$  for  $i = 1, \ldots, b-1$ ; see the proof of Theorem 1 based on induction with the same assumptions.

(Case a). First, we follow Theorem 4.1 in Newey and McFadden (1994) to show that  $\hat{V}_b \xrightarrow{p} V$ . By the triangle inequality, Assumptions 11(a) and 12(a)(i), we have

$$\left\| \hat{V}_{b} - V \right\|$$
  
=  $\left\| \frac{1}{N_{b}} \left\{ \sum_{i=1}^{b-1} \nabla G(\hat{\theta}_{i}; D_{i}) + \nabla G(\hat{\theta}_{b-1}; D_{b}) \right\} - \mathbb{E} \{ \nabla g(\theta^{*}, x_{i,j}) \} \right\|$ 

		_
L		
L		
L		
-		_

$$\leq \left\| \frac{1}{N_{b}} \left\{ \sum_{i=1}^{b-1} \nabla G(\hat{\theta}_{i}; D_{i}) + \nabla G(\hat{\theta}_{b-1}; D_{b}) \right\} - \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \mathbb{E} \{ \nabla g(\hat{\theta}_{i}, x_{i,j}) \} - \frac{n_{b}}{N_{b}} \mathbb{E} \{ \nabla g(\hat{\theta}_{b-1}, x_{i,j}) \} \right\| \\ + \left\| \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \mathbb{E} \{ \nabla g(\hat{\theta}_{i}, x_{i,j}) \} + \frac{n_{b}}{N_{b}} \mathbb{E} \{ \nabla g(\hat{\theta}_{b-1}, x_{i,j}) \} - \sum_{i=1}^{b} \frac{n_{i}}{N_{b}} \mathbb{E} \{ \nabla g(\theta^{*}, x_{i,j}) \} \right\| \\ \leq \sum_{i=1}^{b} \frac{n_{i}}{N_{b}} \sup_{\theta \in B_{\epsilon}} \left\| \frac{1}{n_{i}} \nabla G(\theta; D_{i}) - \mathbb{E} \{ \nabla g(\theta, x_{i,j}) \} \right\| + L \left( \frac{n_{b}}{N_{b}} \left\| \hat{\theta}_{b-1} - \theta^{*} \right\| + \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \left\| \hat{\theta}_{i} - \theta^{*} \right\| \right) \\ \stackrel{\mathrm{P}}{\to} 0. \tag{C.1}$$

Next, we bound the Taylor approximation error using Assumptions 11(a) and 12(a)(ii). We will point out the difference under Assumption 12(a)(i) later. To illustrate,

$$\begin{aligned} \left\| \frac{1}{n_i} \nabla G(\bar{\theta}_i; D_i) - \frac{1}{n_i} \nabla G(\hat{\theta}_i; D_i) \right\| \\ &\leq \left\| \frac{1}{n_i} \nabla G(\bar{\theta}_i; D_i) - \mathbb{E} \{ \nabla g(\bar{\theta}_i, x_{i,j}) \} \right\| + \left\| \frac{1}{n_i} \nabla G(\hat{\theta}_i; D_i) - \mathbb{E} \{ \nabla g(\hat{\theta}_i, x_{i,j}) \} \right\| \\ &+ \left\| \mathbb{E} \{ \nabla g(\bar{\theta}_i, x_{i,j}) \} - \mathbb{E} \{ \nabla g(\hat{\theta}_i, x_{i,j}) \} \right\| \\ &\leq 2 \sup_{\theta \in B_{\epsilon}} \left\| \frac{1}{n_i} \nabla G(\theta; D_i) - \mathbb{E} \{ \nabla g(\theta, x_{i,j}) \} \right\| + L \left\| \bar{\theta}_i - \hat{\theta}_i \right\| \\ &= O_p(n_i^{-1/2} + \left\| \hat{\theta}_i - \theta^* \right\|), \end{aligned}$$

where  $\bar{\theta}_i$  is a point between  $\hat{\theta}_i$  and  $\theta^*$ . By the mean value theorem,

$$G(\theta^*; D_i) = G(\hat{\theta}_i; D_i) + \nabla G(\hat{\theta}_i; D_i)(\theta^* - \hat{\theta}_i) + \left\{ \nabla G(\bar{\theta}_i; D_i) - \nabla G(\hat{\theta}_i; D_i) \right\} (\theta^* - \hat{\theta}_i),$$

where the sub-multiplicativity of Frobenius norm gives

$$\left\|\left\{\nabla G(\bar{\theta}_i; D_i) - \nabla G(\hat{\theta}_i; D_i)\right\} (\theta^* - \hat{\theta}_i)\right\| \leq \left\|\nabla G(\bar{\theta}_i; D_i) - \nabla G(\hat{\theta}_i; D_i)\right\| \left\|\theta^* - \hat{\theta}_i\right\|$$
$$= O_p \left(n_i^{1/2} \left\|\hat{\theta}_i - \theta^*\right\| + n_i \left\|\hat{\theta}_i - \theta^*\right\|^2\right). \quad (C.2)$$

Repeating this trick, we have

$$\bar{g}_{b}(\theta^{*}) = \frac{1}{N_{b}} \sum_{i=1}^{b-1} \left\{ G(\hat{\theta}_{i}; D_{i}) + \nabla G(\hat{\theta}_{i}; D_{i})(\theta^{*} - \hat{\theta}_{i}) \right\} + \frac{1}{N_{b}} G(\theta^{*}; D_{b}) \\
+ O_{p} \left\{ \sum_{i=1}^{b-1} \left( \frac{n_{i}^{1/2}}{N_{b}} \| \hat{\theta}_{i} - \theta^{*} \| + \frac{n_{i}}{N_{b}} \| \hat{\theta}_{i} - \theta^{*} \|^{2} \right) \right\} \\
= \frac{N_{b-1}}{N_{b}} (U_{b-1} + V_{b-1}\theta^{*}) + \frac{1}{N_{b}} G(\theta^{*}; D_{b}) + O_{p} \left\{ \sum_{i=1}^{b-1} \left( \frac{n_{i}^{1/2}}{N_{b}} \| \hat{\theta}_{i} - \theta^{*} \| + \frac{n_{i}}{N_{b}} \| \hat{\theta}_{i} - \theta^{*} \|^{2} \right) \right\} \\
= \hat{U}_{b} + \hat{V}_{b}\theta^{*} + O_{p} \left( \frac{n_{b}^{1/2}}{N_{b}} \| \hat{\theta}_{b-1} - \theta^{*} \| + \sum_{i=1}^{b-1} \frac{n_{i}^{1/2}}{N_{b}} \| \hat{\theta}_{i} - \theta^{*} \|^{2} \right) \\
+ O_{p} \left( \frac{n_{b}}{N_{b}} \| \hat{\theta}_{b-1} - \theta^{*} \|^{2} + \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \| \hat{\theta}_{i} - \theta^{*} \|^{2} \right).$$
(C.3)

If we use Assumption 12(a)(i) instead, (C.3) becomes

$$\bar{g}_{b}(\theta^{*}) = \hat{U}_{b} + \hat{V}_{b}\theta^{*} + o_{p}\left(\frac{n_{b}}{N_{b}} \left\|\hat{\theta}_{b-1} - \theta^{*}\right\| + \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \left\|\hat{\theta}_{i} - \theta^{*}\right\|\right) + O_{p}\left(\frac{n_{b}}{N_{b}} \left\|\hat{\theta}_{b-1} - \theta^{*}\right\|^{2} + \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \left\|\hat{\theta}_{i} - \theta^{*}\right\|^{2}\right).$$

Since  $\bar{g}_b(\theta^*) \xrightarrow{p} 0$  by the uniform law of large numbers (see, e.g., Lemma 2.4 in Newey and McFadden 1994),  $\hat{V}_b \xrightarrow{p} V$ , and  $\|\hat{\theta}_i - \theta^*\| = o_p(1)$  for  $i = 1, \ldots, b-1$ , applying Slutsky's theorem gives  $\hat{U}_b = \bar{g}_b(\theta^*) - \hat{V}_b\theta^* + o_p(1) \xrightarrow{p} -V\theta^*$ .

(Case b). Again, we try to show that  $\hat{V}_b \xrightarrow{p} V$  first. By the triangle inequality, Assumptions 11(b) and 12(b)(i),

$$\begin{aligned} \left\| \hat{V}_{b} - V \right\| \\ &= \left\| \frac{1}{N_{b}} \left\{ \sum_{i=1}^{b-1} \nabla G(\hat{\theta}_{i}; D_{i}) + \nabla G(\hat{\theta}_{b-1}; D_{b}) \right\} - \mathbb{E} \{ \nabla g(\theta^{*}, x_{i,j}) \} \right\| \\ &\leq \left\| \frac{1}{N_{b}} \left\{ \sum_{i=1}^{b-1} \nabla G(\hat{\theta}_{i}; D_{i}) + \nabla G(\hat{\theta}_{b-1}; D_{b}) \right\} - \frac{1}{N_{b}} \sum_{i=1}^{b} \nabla G(\theta^{*}; D_{i}) \right\| \\ &+ \left\| \frac{1}{N_{b}} \sum_{i=1}^{b} \nabla G(\theta^{*}; D_{i}) - \mathbb{E} \{ \nabla g(\theta^{*}, x_{i,j}) \} \right\| \\ &\leq L \left( \frac{n_{b}}{N_{b}} \left\| \hat{\theta}_{b-1} - \theta^{*} \right\| + \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \left\| \hat{\theta}_{i} - \theta^{*} \right\| \right) + \left\| \frac{1}{N_{b}} \sum_{i=1}^{b} \nabla G(\theta^{*}; D_{i}) - \mathbb{E} \{ \nabla g(\theta^{*}, x_{i,j}) \} \right\| \\ \stackrel{\mathrm{P}}{\to} 0. \end{aligned}$$

$$(C.4)$$

Using Assumption 11(b),

$$\begin{aligned} \left\|\nabla G(\bar{\theta}_i; D_i) - \nabla G(\hat{\theta}_i; D_i)\right\| &\leq \left\|\nabla G(\bar{\theta}_i; D_i) - \nabla G(\theta^*; D_i)\right\| + \left\|\nabla G(\hat{\theta}_i; D_i) - \nabla G(\theta^*; D_i)\right\| \\ &= L \cdot n_i \left(\left\|\bar{\theta}_i - \theta^*\right\| + \left\|\hat{\theta}_i - \theta^*\right\|\right) = O_p\left(n_i \left\|\hat{\theta}_i - \theta^*\right\|\right), \end{aligned}$$
(C.5)

where  $\bar{\theta}_i$  is a point between  $\hat{\theta}_i$  and  $\theta^*$ . Repeating the mean value theorem, we have

$$\bar{g}_{b}(\theta^{*}) = \frac{1}{N_{b}} \sum_{i=1}^{b-1} \left\{ G(\hat{\theta}_{i}; D_{i}) + \nabla G(\hat{\theta}_{i}; D_{i})(\theta^{*} - \hat{\theta}_{i}) \right\} + \frac{1}{N_{b}} G(\theta^{*}; D_{b}) + O_{p} \left( \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \left\| \hat{\theta}_{i} - \theta^{*} \right\|^{2} \right) \\ = \frac{N_{b-1}}{N_{b}} (U_{b-1} + V_{b-1}\theta^{*}) + \frac{1}{N_{b}} G(\theta^{*}; D_{b}) + O_{p} \left( \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \left\| \hat{\theta}_{i} - \theta^{*} \right\|^{2} \right) \\ = \hat{U}_{b} + \hat{V}_{b}\theta^{*} + O_{p} \left( \frac{n_{b}}{N_{b}} \left\| \hat{\theta}_{b-1} - \theta^{*} \right\|^{2} + \sum_{i=1}^{b-1} \frac{n_{i}}{N_{b}} \left\| \hat{\theta}_{i} - \theta^{*} \right\|^{2} \right).$$
(C.6)

The conclusion then follows by the same argument in case (a).

### C.2 Lemma 2

**Lemma 2** (Bound for sums of sample sizes). Recall that  $n_i$  is the sample size of  $D_i$  and  $N_i = \sum_{j=1}^i n_j$ . We have

$$\sum_{i=1}^{b} \frac{n_i}{N_i} \le 1 + \log \frac{N_b}{n_1}, \quad \sum_{i=1}^{b} \frac{n_i}{\sqrt{N_i}} \le 2\sqrt{N_b} \quad and \quad \sum_{i=1}^{b} \frac{\sqrt{n_i}}{\sqrt{N_i}} \le \frac{2\sqrt{N_b}}{\sqrt{\min(n_1, \dots, n_b)}}$$

*Proof.* The first two inequalities are restated from Luo et al. (2022) for ease of reference. Their proof can be found in the supplementary material of Luo et al. (2022). The last inequality follows from

$$\sum_{i=1}^{b} \frac{\sqrt{n_i}}{\sqrt{N_i}} \le \frac{1}{\sqrt{\min(n_1, \dots, n_b)}} \sum_{i=1}^{b} \frac{n_i}{\sqrt{N_i}} \le \frac{2\sqrt{N_b}}{\sqrt{\min(n_1, \dots, n_b)}}.$$

### **D** Additional results

### D.1 Online instrumental variables regression

Figures 6 and 7 report the additional results of online instrumental variables regression in the independent model (a) and dependent model (b), respectively. The findings are same as those described in Section 5.1.

### D.2 Online Sargan–Hansen test

Figure 8 reports the additional results of online Sargan–Hansen test at 5% nominal level. It verifies the findings in Section 5.2 for other values of  $\theta_2^*$ .

### D.3 Online quantile regression

Figure 9 reports the additional results of online quantile regression. It shows that implicit updates do not improve the statistical efficiency.

### D.4 Online anomaly detection

Figure 10 reports the additional results of online anomaly detection at 5% nominal level. The findings are same as those described in Section 5.4.



Figure 6: Online instrumental variables estimation of  $\theta^*$  in the independent model (a). The caption of Figure 1 also applies here.

### **E** Implementation details

### E.1 Stochastic volatility modeling

Apart from removing observations in the weekend, we remove abnormal exchange rates at 2001-09-12 03:12:00 (due to negativity) and 2001-06-08 06:32:00 (due to an extreme jump that leads to a log 5-minute return of over 700%).

### E.2 Inertial sensor calibration

To use OGMM with wavelet moments, we extend the pyramid algorithm for maximal overlap discrete wavelet transform using the Haar filter to an online setting in Algorithm E.1. The key idea is to store the necessary data in queues so that they can be retrieved and later deleted in O(1) time. As a result, Algorithm E.1 gives identical wavelets compared with its offline counterpart. For an incoming batch of size  $n_b$ , the time and space complexities are  $O(n_bq)$  and  $O(n_b + 2^q)$ , respectively.



Figure 7: Online instrumental variables estimation of  $\theta^*$  in the dependent model (b). The caption of Figure 1 also applies here.

In OGMM estimation, we slightly modify the arguments for long-run variance estimation compared with other examples in our paper. Specially, we use Leung and Chan's (2025) recursive long-run variance estimator with  $\lambda = 1$ ,  $\phi = 1$ , pilot=0 and warm=TRUE, which is the default setting in the R-package rlaser. We also ignore  $\{w_{t,j}\}_{t=1}^{2^q-1}$  for  $j = 1, \ldots, q$  since  $w_{t,q}$  is only defined for  $t \ge 2^q$ . For structural stability testing with (3.8), we need to perform unrestricted GMM estimation. This is done using optim in the R-package stats with method="L-BFGS-B", lower = c(rep(c(0.0001, 1e-13), 2), 1e-13) and upper = c(rep(c(0.9999, 1e-5), 2), 1e-5), where the order of parameter is  $\theta^* = (\rho_1, \sigma_1^2, \rho_2, \sigma_2^2, \sigma_3^2)^{\mathsf{T}}$ .

Finally, we remark that the R-package gmwm is slightly modified in our paper; see https://github.com/hemanlmf/gmwm/tree/gmwm2-improvement. We add the possibility to specify the number of moment condition q as the original version fixes  $q = \lfloor \log_2 N \rfloor$ , where N is the sample size. We also include an argument to change the optimization algorithm, although the default conjugate gradients method is still used at the end.



Figure 8: Online Sargan–Hansen test at 5% nominal level under the independent model (c) (upper panel) and dependent model (d) (lower panel). The case  $\theta_2^* = 0$  is omitted because Figure 3 has already shown the size at all sample sizes. Therefore, a higher size-adjusted power means that the test is better here.



Figure 9: Online quantile regression under the independent model (e) (upper panel) and dependent model (f) (lower panel). The caption of Figure 1 also applies here.



Figure 10: Online anomaly detection at 5% nominal level under the independent model (g) (upper panel) and dependent model (h) (lower panel). The caption of Figure 5 also applies here.

Algorithm E.1: Online maximal overlap discrete wavelet transform

[1] initialization: [2] Receive q and  $\{y_t\}_{t=1}^n$ , where  $n > 2^q$ [3] Initialize empty queues  $\vec{z}_0, \ldots, \vec{z}_{q-1}$ [4] for j = 0 to q - 1 do Set  $\tau = 2^j$ [5] for  $t = 2^{j+1}$  to n do [6] Set  $w_{t,j} = 0.5y_t - 0.5y_{t-\tau}$  // Haar filter [7] Set  $v_t = 0.5y_t + 0.5y_{t-\tau}$  // pyramid algorithm [8] Store  $y_{n-\tau+1}, \ldots, y_n$  in  $\vec{z}_j$  // for online updates after initialization [9] Set  $\{y_t\}_{t=1}^n = \{v_t\}_{t=1}^n$ [10] [11] begin Receive  $y_{n+1}$ [12] for j = 0 to q - 1 do [13] Set  $\tau = 2^j$ [14] Pop  $y_{n-\tau+1}$  from  $\vec{z}_j$ [15] Set  $w_{n+1,j} = 0.5y_{n+1} - 0.5y_{n-\tau+1}$ [16] Push  $y_{n+1}$  into  $\vec{z}_j$ [17] Set  $y_{n+1} = 0.5y_{n+1} + 0.5y_{n-\tau+1}$ [18] Set n = n + 1[19]