

# Kaon Modification in Pion Medium

Yu. B. Ivanov<sup>1,2,\*</sup>

<sup>1</sup>*Bogoliubov Laboratory of Theoretical Physics, JINR Dubna, 141980 Dubna, Russia*

<sup>2</sup>*National Research Center "Kurchatov Institute", 123182 Moscow, Russia*

Kaon properties in pion medium are studied. The dense pion medium is produced in high-energy heavy-ion collisions. The consideration is based on the chiral nucleon-kaon-pion Lagrangian. It is found that the pion medium does not produce a substantive contribution but enhances the effect of the baryon matter. Numerical estimate shows that this enhancement factor induced by the pion medium does not exceed 11% at the freeze-out stage of heavy-ion collisions, i.e. it is small. However, the impact of this enhancement can be higher in actual nuclear collisions because the effect of the in-medium (anti)kaon modification is accumulated during the (anti)kaon evolution before the freeze-out when the pion density is higher.

It is well known that kaons are modified in the baryon medium, which attracted attention since Kaplan and Nelson suggested possibility of kaon condensation in dense matter [1]. The attractive kaon-nucleon interaction may result in condensation in the interior of neutron stars, see review papers [2–5]. Theoretical studies of kaon production from A+A collisions at moderately relativistic energies indicated that in-medium modifications of the properties of (anti)kaons are seen in the collective flow pattern as well as in the abundance and spectra of kaons and antikaons [5–7].

In all aforementioned studies the in-medium modifications of kaons were induced by the baryon medium. At higher collision energies, mesons (mostly pions) are abundantly produced. In this paper we address the question if the pionic medium can modify the kaon properties additionally to the baryon-induced modifications?

The present consideration follows the lines of Ref. [8], where the influence of the Bose-Einstein pion condensate on kaon properties was studied. In the present paper, the effect of the thermal pion medium on kaon properties is considered.

The starting point of the consideration is the conventional nucleon-kaon Lagrangian (see, e.g., Ref. [9]) supplemented by the chiral pion-kaon Lagrangian [10], which well reproduces the  $\pi K$  scattering and dimeson atomic data. This Lagrangian reads

$$\begin{aligned} \mathcal{L} = & \bar{N}(i\gamma_\mu\partial^\mu - m_N)N + \partial_\mu\bar{K} \cdot \partial^\mu K - m_K^2\bar{K}K \\ & + \frac{\Sigma_{KN}}{f_K^2}[\bar{N}N\bar{K}K - \frac{3i}{8f_K^2}\bar{N}\gamma_\mu N(\bar{K}\partial^\mu K - \partial^\mu\bar{K} \cdot K)] \\ & + \frac{1}{2}(\partial_\mu\boldsymbol{\pi} \cdot \partial^\mu\boldsymbol{\pi} - m_\pi^2\boldsymbol{\pi}^2) \\ & + \frac{1}{4f_\pi^2}[(m_\pi^2 + m_K^2)\bar{K}K\boldsymbol{\pi}^2 - \bar{K}K\partial_\mu\boldsymbol{\pi} \cdot \partial^\mu\boldsymbol{\pi} \\ & - \boldsymbol{\pi}^2\partial_\mu\bar{K} \cdot \partial^\mu K - \bar{K}\boldsymbol{\pi}\partial_\mu K \cdot \partial^\mu\boldsymbol{\pi} - (\partial_\mu\bar{K} \cdot \partial^\mu\boldsymbol{\pi})K\boldsymbol{\pi}], \end{aligned} \quad (1)$$

where  $\boldsymbol{\pi}$  is the isotopic vector of pion fields and the kaon

is the isotopic spinor

$$K = \begin{pmatrix} K^+ \\ K^0 \end{pmatrix} \quad \text{and} \quad \bar{K} = (K^- \quad \bar{K}^0), \quad (2)$$

$f_\pi = 93$  MeV is the pion decay constant,  $f_K$  is the kaon decay constant, and  $\Sigma_{KN}$  is the kaon-nucleon sigma term [6]. The term proportional to  $\Sigma_{KN}$  results from the attractive scalar interaction due to explicit chiral symmetry breaking. The  $\pi N$  interaction is omitted in this Lagrangian because it is irrelevant to our purposes. The chiral  $\pi K$  Lagrangian contains additional 1/2 factor as compared to Refs. [8, 10] because scalar  $\boldsymbol{\pi}$  fields are used here instead of charged  $\pi^+$  and  $\pi^-$  in Refs. [8, 10].

The above Lagrangian contains only  $s$ -wave kaon-nucleon interaction. Importance of  $p$ -wave interactions was indicated in Refs. [7, 11, 12]. Also there are more extended and sophisticated versions of the chiral  $\pi K$  Lagrangian, see, e.g., Ref. [13, 14]. Therefore, Lagrangian (1) can only serve as a basis for semi-quantitative estimation of the in-medium effects in (anti)kaon production, which is enough for our purposes.

From the Euler-Lagrange equation and using the mean-field approximation for the nucleon and pion fields one obtains the following Klein-Gordon equation for a kaon in the nucleon-pion medium

$$\begin{aligned} & \left(1 - \frac{\langle\boldsymbol{\pi}^2\rangle}{4f_\pi^2}\right)(\partial_\mu\partial^\mu K + m_K^2 K) - \frac{\Sigma_{KN}}{f_K^2}\rho_S K \\ & + \frac{3i}{4f_K^2}J_\mu\partial^\mu K - \frac{1}{4f_\pi^2}\langle\partial_\mu\boldsymbol{\pi}^2\rangle \cdot \partial^\mu K \\ & - \frac{1}{4f_\pi^2}[m_\pi^2\langle\boldsymbol{\pi}^2\rangle + \langle\boldsymbol{\pi}\partial_\mu\partial^\mu\boldsymbol{\pi}\rangle]K = 0, \end{aligned} \quad (3)$$

where

$$J_\mu = \langle\bar{N}\gamma_\mu N\rangle \quad (4)$$

$$\rho_S = \langle\bar{N}N\rangle \quad (5)$$

are the nucleon current and scalar density, respectively. In the homogeneous and stationary medium

$$\partial_\mu\langle\boldsymbol{\pi}^2\rangle = 0.$$

\*e-mail: yivanov@theor.jinr.ru

The term [...] in Eq. (3) can be estimated by means of the Euler-Lagrange equation for pions. Then it is

$$\frac{1}{4f_\pi^2} [m_\pi^2 \langle \pi^2 \rangle + \langle \pi \partial_\mu \partial^\mu \pi \rangle] \propto \frac{\langle \bar{K} K \rangle}{4f_\pi^2} \frac{\langle \pi^2 \rangle}{4f_\pi^2},$$

i.e. contains additional smallness

$$\frac{\langle \bar{K} K \rangle}{\langle \pi^2 \rangle} \ll 1.$$

We also could add the pion nonlinear (Weinberg) term into the Lagrangian (1) [8]

$$-\frac{1}{8f_\pi^2} (2\pi^2 \partial_\mu \pi \cdot \partial^\mu \pi - m_\pi^2 \pi^4)$$

similarly to Ref. [8]. Then it would be

$$\frac{1}{4f_\pi^2} [m_\pi^2 \langle \pi^2 \rangle + \langle \pi \partial_\mu \partial^\mu \pi \rangle] \propto \left( \frac{\langle \pi^2 \rangle}{4f_\pi^2} \right)^2,$$

i.e. it also contains extra smallness. Thus, the term [...] in Eq. (3) can safely neglected.

Finally, in the rest frame of homogeneous and stationary medium, i.e.  $\mathbf{J} = 0$ , we arrive at

$$\partial_\mu \partial^\mu K + m_K^2 K + F_\pi \left[ \frac{3i}{4f_K^2} \rho_N \partial_t K - \frac{\Sigma_{KN}}{f_K^2} \rho_S K \right] = 0, \quad (6)$$

where  $\rho_N$  is the nucleon density in the rest frame,  $J_\mu = (\rho_N, 0, 0, 0)$ , and

$$F_\pi = \left( 1 - \frac{\langle \pi^2 \rangle}{4f_\pi^2} \right)^{-1}. \quad (7)$$

Thus, we arrive at the well-known equation for the kaon in the nucleon medium [9] but with additional enhancement factor  $F_\pi$  that results from the pion medium. Similar manipulations for anti-kaons result to the same equation except for the opposite sign in front of the  $\rho_N \partial_t \bar{K}$  term.

The (anti)kaon dispersion relation in nuclear-pion matter is then given by

$$\omega^2 = m_K^2 + \mathbf{k}^2 - \frac{F_\pi \Sigma_{KN}}{f_K^2} \rho_S \pm \frac{3F_\pi}{4f_K^2} \omega \rho_N, \quad (8)$$

where the upper (lower) sign in front of the  $\omega \rho_N$  term corresponds to the (anti)kaon and  $(\omega, \mathbf{k})$  is the four-momentum of the (anti)kaon, The quantity

$$m_\pi \langle \pi^2 \rangle = \rho_S^\pi = 3m_\pi \int \frac{d^3k}{(2\pi)^3 \omega_\pi(k)} \frac{1}{\exp(\omega_\pi/T) - 1} \quad (9)$$

is the scalar pion density with  $T$  being the temperature of the medium.

From the dispersion relation, the (anti)kaon energy in medium can be obtained

$$\omega(\mathbf{k}) = \sqrt{m_K^2 + \mathbf{k}^2 - \frac{F_\pi \Sigma_{KN}}{f_K^2} \rho_S + \left( \frac{3F_\pi \rho_N}{8f_K^2} \right)^2} \pm \frac{3F_\pi \rho_N}{8f_K^2},$$

where again the upper(lower) sign refers to  $K(\bar{K})$ . The nucleon scalar and number densities,  $\rho_S$  and  $\rho_N$ , respectively, are usually replaced by respective baryon densities in actual calculations. As seen, the pion medium does not produce a substantive contribution but enhances the effect of the baryon matter.

To quantitatively estimate the enhancement factor  $F_\pi$  due to the pion medium, use is made of the freeze-out temperatures and baryon chemical potentials ( $\mu_B$ ) deduced from experimental data with the statistical model [15, 16]

$$T(\mu_B) = a - b\mu_B^2 - c\mu_B^4$$

where  $a = 0.166$  GeV,  $b = 0.139$  GeV<sup>-1</sup>,  $c = 0.053$  GeV<sup>-3</sup>, and

$$\mu_B(\sqrt{s_{NN}}) = \frac{d}{1 + e\sqrt{s_{NN}}}$$

with  $d = 1.308$  GeV and  $e = 0.273$  GeV<sup>-1</sup>. Here,  $\sqrt{s_{NN}}$  is the center-of-mass energy of the nuclear collision (in GeV).




FIG. 1: Enhancement factor  $F_\pi$  induced by the pion medium at the freeze-out stage as a function the center-of-mass energy of the nuclear collision  $\sqrt{s_{NN}}$ .

The result of numerical calculation of the enhancement factor  $F_\pi$  induced by the pion medium at the freeze-out stage as a function the center-of-mass energy of the nuclear collision is presented in Fig. 1. As seen, it does not exceed 11%, i.e. it is small. This retroactively confirms the approximations made at derivation of Eq. (10). However, the impact of this enhancement can be higher in actual nuclear collisions because the effect of the in-medium (anti)kaon modification is accumulated during the (anti)kaon evolution before the freeze-out when the pion density is higher.

Enlightening discussions with D.N. Voskresensky are gratefully acknowledged. This work was carried out using

computing resources of the federal collective usage center “Complex for simulation and data processing for mega-

science facilities” at NRC “Kurchatov Institute” [17].

- 
- [1] D. B. Kaplan and A. E. Nelson, “Strange Goings on in Dense Nucleonic Matter,” *Phys. Lett. B* **175**, 57-63 (1986), doi:10.1016/0370-2693(86)90331-X A. E. Nelson and D. B. Kaplan, “Strange Condensate Realignment in Relativistic Heavy Ion Collisions,” *Phys. Lett. B* **192**, 193 (1987), doi:10.1016/0370-2693(87)91166-X
- [2] A. L. Watts, N. Andersson, D. Chakrabarty, M. Feroci, K. Hebeler, G. Israel, F. K. Lamb, M. C. Miller, S. Morsink and F. Özel, *et al.* “Colloquium : Measuring the neutron star equation of state using x-ray timing,” *Rev. Mod. Phys.* **88**, no.2, 021001 (2016), doi:10.1103/RevModPhys.88.021001 [arXiv:1602.01081 [astro-ph.HE]].
- [3] B. A. Li, B. J. Cai, W. J. Xie and N. B. Zhang, “Progress in Constraining Nuclear Symmetry Energy Using Neutron Star Observables Since GW170817,” *Universe* **7**, no.6, 182 (2021), doi:10.3390/universe7060182 [arXiv:2105.04629 [nucl-th]].
- [4] G. F. Burgio, H. J. Schulze, I. Vidana and J. B. Wei, “Neutron stars and the nuclear equation of state,” *Prog. Part. Nucl. Phys.* **120**, 103879 (2021), doi:10.1016/j.ppnp.2021.103879 [arXiv:2105.03747 [nucl-th]].
- [5] L. Tolos and L. Fabbietti, “Strangeness in Nuclei and Neutron Stars,” *Prog. Part. Nucl. Phys.* **112**, 103770 (2020), doi:10.1016/j.ppnp.2020.103770 [arXiv:2002.09223 [nucl-ex]].
- [6] C. Hartnack, H. Oeschler, Y. Leifels, E. L. Bratkovskaya and J. Aichelin, “Strangeness Production close to Threshold in Proton-Nucleus and Heavy-Ion Collisions,” *Phys. Rept.* **510**, 119-200 (2012), doi:10.1016/j.physrep.2011.08.004 [arXiv:1106.2083 [nucl-th]].
- [7] T. Song, L. Tolos, J. Wirth, J. Aichelin and E. Bratkovskaya, “In-medium effects in strangeness production in heavy-ion collisions at (sub)threshold energies,” *Phys. Rev. C* **103**, no.4, 044901 (2021), doi:10.1103/PhysRevC.103.044901 [arXiv:2012.05589 [nucl-th]].
- [8] E. E. Kolomeitsev and D. N. Voskresensky, “Bose-Einstein condensation of pions in ultrarelativistic nucleus-nucleus collisions and the spectra of kaons,” *Phys. Atom. Nucl.* **58**, 2082-2087 (1995).
- [9] C. M. Ko and G. Q. Li, “Medium effects in high-energy heavy ion collisions,” *J. Phys. G* **22**, 1673-1725 (1996), [arXiv:nucl-th/9611027 [nucl-th]].
- [10] A. A. Belkov, D. Ebert and V. N. Pervushin, “LOW-ENERGY PREDICTIONS OF MODERN CHIRAL LAGRANGIANS BASED ON QUARK DYNAMICS,” *Sov. J. Part. Nucl.* **22**, 1-30 (1991).
- [11] E. E. Kolomeitsev, B. Kampfer and D. N. Voskresensky, “Kaon polarization in nuclear matter,” *Nucl. Phys. A* **588**, 889-917 (1995), doi:10.1016/0375-9474(95)00084-E
- [12] E. E. Kolomeitsev and D. N. Voskresensky, “Negative kaons in dense baryonic matter,” *Phys. Rev. C* **68**, 015803 (2003), doi:10.1103/PhysRevC.68.015803 [arXiv:nucl-th/0211052 [nucl-th]].
- [13] A. Roessl, “Pion kaon scattering near the threshold in chiral SU(2) perturbation theory,” *Nucl. Phys. B* **555**, 507-539 (1999), doi:10.1016/S0550-3213(99)00336-3 [arXiv:hep-ph/9904230 [hep-ph]].
- [14] J. Bijnens, P. Dhonte and P. Talavera, “pi K scattering in three flavor ChPT,” *JHEP* **05**, 036 (2004), doi:10.1088/1126-6708/2004/05/036 [arXiv:hep-ph/0404150 [hep-ph]].
- [15] J. Cleymans, H. Oeschler, K. Redlich and S. Wheaton, “Comparison of chemical freeze-out criteria in heavy-ion collisions,” *Phys. Rev. C* **73**, 034905 (2006), doi:10.1103/PhysRevC.73.034905 [arXiv:hep-ph/0511094 [hep-ph]].
- [16] F. Karsch and K. Redlich, “Probing freeze-out conditions in heavy ion collisions with moments of charge fluctuations,” *Phys. Lett. B* **695**, 136-142 (2011), doi:10.1016/j.physletb.2010.10.046 [arXiv:1007.2581 [hep-ph]].
- [17] <http://ckp.nrcki.ru/>