

Study of the mass spectra of doubly heavy Ξ_{QQ} and Ω_{QQ} baryons

Ji-Hai Pan^{1,*} and Ji-Si Pan^{†2,‡}

¹*Department of Mathematics and Physics,*

Liuzhou Institute of Technology, Liuzhou 545616, China

²*School of Physics and Information Engineering,*

Guangxi Science & Technology Normal University, Laibin 546199, China

In the paper, we enumerated the mass spectra of the radial and orbital excited states for the doubly heavy Ξ_{QQ} and Ω_{QQ} baryons using the Regge trajectory model and the scaling rules. Recently, LHCb Collaboration first observed a doubly charmed baryon Ξ_{cc}^{++} in the $\Lambda_c^+ K^- \pi^+ \pi^+$ decay with a mass of 3621.40 ± 0.78 MeV. Our studies show that Ξ_{cc}^{++} can be grouped into the $1S$ -wave state with the spin-parity quantum number $J^P = 1/2^-$. On the other hand, the mass of Ξ_{cc}^{++} with $J^P = 3/2^-$ is predicted to be 3699.69 MeV. We also predict the mass spectra of the unknown ground and excited states for the doubly heavy baryons, which provide useful references for the experimental test in the future.

I. INTRODUCTION

During the past two decades, the study of the spectroscopy of the heavy baryons is an interesting and important issue in hadronic physics. Furthermore, it enables us to gain a deeper understanding of the internal structure and strong interactions of hadrons. Doubly heavy baryon is composed of one light quark ($q = u, d, \text{ or } s$), and two heavy quarks ($QQ = cc, bc, \text{ and } bb$), where the two heavy quarks can be defined as the heavy diquark (QQ). The doubly heavy baryons have Ξ_{QQ} and Ω_{QQ} family. The Ξ_{QQ} family has up quark (u) or down quark (d), namely, Ξ_{cc} , Ξ_{bc} , and Ξ_{bb} baryons, while the Ω_{QQ} family has one light strange quark (s), namely, Ω_{cc} , Ω_{bc} , and Ω_{bb} baryons. As a component of hadron, it has been explored by various methods and aroused wide interest.

In 2017, LHCb Collaboration has observed a doubly charmed baryon Ξ_{cc}^{++} in the $\Lambda_c^+ K^- \pi^+ \pi^+$ mass spectra [1], the measured mass is 3621.40 ± 0.78 MeV. Later, the LHCb Collaboration confirmed the existence of this state in the $\Xi_c^+ \pi^+$ decay [2]. In 2020, precision measurement of the Ξ_{cc}^{++} mass is given by $3621.55 \pm 0.23 \pm 0.30$ MeV in Ref. [3]. However, the spin-parity quantum number

[†] Corresponding author

*Electronic address: Tunmnwnu@outlook.com

[‡]Electronic address: panjisi@gxstnu.edu.cn

J^P of the Ξ_{cc}^{++} state have not been determined. According to our calculation results indicated that may be regarded as a $1S$ state with $J^P = 1/2^-$. More assignments for the Ξ_{cc}^{++} state can be found in Refs [4–10].

In earlier times of 2002, the SELEX collaboration reported the first observation of a candidate for a double charmed baryon state Ξ_{cc}^+ in the charged decay mode $\Xi_{cc}^+ \rightarrow \Lambda_c^+ K^- \pi^+$ [11]. The observed mass of Ξ_{cc}^+ state is 3519 ± 1 MeV. In 2005, the SELEX collaboration reported again the doubly charmed baryon Ξ_{cc}^+ in the charged decay mode $\Xi_{cc}^+ \rightarrow p D^+ K^-$ [12], but could not be confirmed by Belle, *BABAR*, CLEO, and LHCb collaborations [13]. However, our knowledge of the ground and higher excited states for the doubly heavy baryons are still lacking.

In this paper we study the effective masses of the heavy diquark and light quark with the relative kinetic energy. We also use the Regge trajectory to complete the spin-average masses of the doubly heavy Ξ_{QQ} and Ω_{QQ} baryons. In addition, to obtain the mass shifts, we exploit the scaling relations to calculate the spin-coupling parameters. For the unknown doubly heavy baryons in experiments, the mass spectra of the radial excited and orbital excited states are calculated and predicted.

The structure of this paper is as follows. In Sec. II, we study the effective masses of heavy diquark and light quark. We analyze the spin-average masses of the doubly heavy baryons in Sec. III. We use a all- JLS coupling to compute the mass spectra expressions of doubly heavy baryons in Sec. IV. We discuss the scaling relations in Sec. V. In Sec. VI, we calculate the mass spectra of the Ξ_{QQ} baryons. In Sec. VII, a similar mass analysis is given by the scaling relations for the Ω_{QQ} baryons. Finally, we outline our conclusion in Sec. VIII.

II. THE EFFECTIVE MASS OF QUARK

Generally, the effective mass of the quark is a concept in quantum chromodynamics (QCD) that arises due to the confinement of quarks within hadrons [14, 15]. It would be more interesting to study the effective mass of the quarks in the baryon system. Accordingly, the effective mass of the quark should be different from the choices of the radial and orbital quantum number for the baryons with the current mass of quark.

Including relativistic effects, one can obtain the effective masses M_1 and M_2 of the quarks for the heavy baryons given by

$$M_1 = \frac{m_1}{\sqrt{1 - v_1^2}}, \quad (1)$$

$$M_2 = \frac{m_2}{\sqrt{1 - v_2^2}}, \quad (2)$$

where m_1, m_2 can be regarded as the current masses of the quarks, v_1 and v_2 are the velocities of the quarks for the baryons, respectively. For simplicity, we have chosen the velocity of light $c = 1$. To obtain the values of the quark velocities v_i ($i = 1, 2$) in the heavy-light system we exploit the relative kinetic energy $T_i = \frac{1}{2}m_i v_i^2$ with the current mass m_i of the quarks. Then, by taking the average of the square velocity v_i^2 , we get

$$\langle v_i^2 \rangle = \left\langle \frac{2}{m_i} T_i \right\rangle. \quad (3)$$

Considering the relative spherical coordinates r and the Coulomb potential $V = -4\alpha_s/3r$ with the running coupling constant α_s for the heavy baryons, and the Virial theorem implies

$$\langle T_i \rangle = \frac{1}{2} \langle r V' \rangle. \quad (4)$$

This together with the momentum conservation $m_1 v_1 = m_2 v_2$, the square velocity $\langle v_i^2 \rangle$ with $\langle 1/r \rangle = 1/a_B N^2$ are

$$\langle v_1^2 \rangle = \frac{1}{m_1} \frac{4\alpha_s}{3} \frac{1}{a_B N^2}, \quad (5)$$

$$\langle v_2^2 \rangle = \frac{m_1}{m_2^2} \frac{4\alpha_s}{3} \frac{1}{a_B N^2}, \quad (6)$$

where a_B is the Bohr radius related to the reduced mass $\mu = m_1 m_2 / (m_1 + m_2)$ in the baryon system as $a_B = 1/\mu$ corresponding to the masses of the quarks. Here, $N = n + L + 1$ is the principle quantum number for the heavy baryons with the radial quantum number n ($n = 0, 1, 2, 3, \dots$) and the orbital quantum number L ($L = 0, 1, 2, 3, \dots$). Using the above Eqs. (5) and (6), Eqs. (1) and (2) become

$$M_1 = \frac{m_1}{\sqrt{1 - \frac{1}{m_1} \frac{4\alpha_s}{3} \frac{\mu}{N^2}}}, \quad (7)$$

$$M_2 = \frac{m_2}{\sqrt{1 - \frac{m_1}{m_2^2} \frac{4\alpha_s}{3} \frac{\mu}{N^2}}}. \quad (8)$$

III. THE SPIN-AVERAGE MASSES OF DOUBLY HEAVY BARYONS

In the heavy-light quark picture in this work, we consider the baryon system as a bound state of three constituent quarks under the strong interaction. The doubly heavy baryon can be done by viewing a QCD rotating-string tied to the heavy diquark (QQ) with spin-1 as one end and to

the light quark (q) with spin-1/2 as the other end. Based on this model, we make an attempt to investigate the Regge trajectory behavior of the hadronic system.

For the orbital excitations of the doubly heavy baryons, the spin-average mass \bar{M} and the angular momentum L following the equations are given by Refs. [16, 17]

$$\bar{M} = \frac{m_{QQ}}{\sqrt{1-v_{QQ}^2}} + \frac{\alpha}{\omega} \int_0^{v_{QQ}} \frac{du}{\sqrt{1-u^2}} + \frac{m_q}{\sqrt{1-v_q^2}} + \frac{\alpha}{\omega} \int_0^{v_q} \frac{du}{\sqrt{1-u^2}}, \quad (9)$$

$$L = \frac{m_{QQ}v_{QQ}^2}{\sqrt{1-v_{QQ}^2}} + \frac{\alpha}{\omega^2} \int_0^{v_{QQ}} \frac{u^2 du}{\sqrt{1-u^2}} + \frac{m_q v_q^2}{\sqrt{1-v_q^2}} + \frac{\alpha}{\omega^2} \int_0^{v_q} \frac{u^2 du}{\sqrt{1-u^2}}, \quad (10)$$

where m_{QQ} and m_q can be regarded as current mass of the heavy diquark and light quark, respectively. Then α is the QCD string tension coefficient. In Ref. [18], the slope of the Regge trajectory is mainly determined by the effective mass M_{QQ} of the heavy diquark, the dependence of the Regge slopes α' on M_{QQ} has the form:

$$\alpha' = \frac{1}{2\pi\alpha} \propto \frac{1}{\sqrt{M_{QQ}}}. \quad (11)$$

By introducing the coefficient k , the coefficient α is defined as

$$\alpha = \frac{k}{2\pi} (M_{QQ})^{\frac{1}{2}}. \quad (12)$$

In addition, the velocity of the quark in the doubly heavy baryon system is defined as $v_{i'} = \omega r_{i'}$ ($i' = QQ, q$), where ω and $r_{i'}$ can be regarded as the angular velocity and the position from the centre-of-mass, respectively. Applying Eqs. (1) and (2), Eqs. (9) and (10) can be integrated to the second order give

$$\bar{M} = M_{QQ} + M_q + M_{QQ}v_{QQ}^2 + \frac{\pi\alpha}{2\omega}, \quad (13)$$

$$L = \frac{1}{\omega} (M_q + M_{QQ}v_{QQ}^2 + \frac{\pi\alpha}{4\omega}). \quad (14)$$

For the string ending at the heavy diquark we utilize the boundary condition

$$\frac{\alpha}{\omega} = \frac{m_{QQ}v_{QQ}}{1-v_{QQ}^2} \approx M_{QQ}v_{QQ}. \quad (15)$$

Substituting Eq. (15) into Eqs. (13) and (14) eliminating the angular velocity ω gives the spin-averaged mass formula for the orbital excited states,

$$(\bar{M} - M_{QQ})^2 = \pi\alpha L + (M_q + M_{QQ}v_{QQ}^2)^2. \quad (16)$$

Using Eq. (6), the spin-averaged mass formula (16) can be written as

$$(\bar{M} - M_{QQ})^2 = \pi\alpha L + \left(M_q + \frac{M_{QQ}m_q}{m_{QQ}^2} \frac{4\alpha_s}{3} \frac{1}{a_B N^2} \right)^2. \quad (17)$$

In order to obtain the spin-average masses of the radial excited states for the doubly heavy baryons, we would like to extend the Regge-like mass relation Eq. (17) by replacing $\pi\alpha L$ with $\pi\alpha(L + 1.37n + 1)$ in Ref. [19], in which α that appears in Eq. (12),

$$\bar{M} = M_{QQ} + \left(\frac{1}{2}(M_{QQ})^{\frac{1}{2}}(L + 1.37n + 1) + \left(M_q + \frac{M_{QQ}m_q}{m_{QQ}^2} \frac{4\alpha_s}{3} \frac{\mu}{(n + L + 1)^2} \right)^2 \right)^{\frac{1}{2}}. \quad (18)$$

IV. THE MASS SPECTRA EXPRESSIONS OF DOUBLY HEAVY BARYONS

For the doubly heavy baryon states, the mass spectra forms with the spin-coupling parameters are constructed. To estimate the mass splitting, we need to consider the spin-dependent Hamiltonian [20, 21],

$$H^{SD} = a_1 \mathbf{L} \cdot \mathbf{S}_{QQ} + a_2 \mathbf{L} \cdot \mathbf{S}_q + b_1 S_{12} + c_1 \mathbf{S}_{QQ} \cdot \mathbf{S}_q, \quad (19)$$

where a_1 , a_2 , b_1 , and c_1 are the spin-coupling parameters. The first two terms are the spin-orbit interactions, the third is the tensor energy, and the last is the contact interaction between the heavy diquark spin \mathbf{S}_{QQ} and light quark spin \mathbf{S}_q . Here, the tensor interaction S_{12} can be given by [22–24]

$$\begin{aligned} S_{12} &= 3(\mathbf{S}_q \cdot \hat{\mathbf{r}})(\mathbf{S}_{QQ} \cdot \hat{\mathbf{r}})/r^2 - \mathbf{S}_{QQ} \cdot \mathbf{S}_q \\ &= -\frac{3}{(2L-1)(2L+3)} [(\mathbf{L} \cdot \mathbf{S}_q)(\mathbf{L} \cdot \mathbf{S}_{QQ}) + (\mathbf{L} \cdot \mathbf{S}_{QQ})(\mathbf{L} \cdot \mathbf{S}_q) - \frac{2}{3}L(L+1)(\mathbf{S}_{QQ} \cdot \mathbf{S}_q)], \end{aligned} \quad (20)$$

where $\hat{\mathbf{r}} = \vec{\mathbf{r}}/r$ determines the unity vector of position pointing from the center of mass of the light quark relative to the heavy diquark. Obviously, it can be seen that the form of S_{12} would depend on the different value of L .

Calculated the mass spectrum for the doubly heavy baryons from the spin-average masses \bar{M} in Eq. (18) and the mass shift $\Delta M(J, j)$, the baryon mass becomes

$$M(J, j) = \bar{M} + \Delta M(J, j), \quad (21)$$

where $\mathbf{J} = \mathbf{S} + \mathbf{L}$ is the total angular momentum with the total spin $\mathbf{S} = \mathbf{S}_{QQ} + \mathbf{S}_q$, and $j = L + S_{QQ}$ is the total angular momentum of the heavy diquark.

We analyze the S -wave states of the doubly heavy baryons with $L = 0$ by using the Hamiltonian Eq. (19). In this case, only the last term survives,

$$H^{SD}(L = 0) = c_1 \mathbf{S}_{QQ} \cdot \mathbf{S}_q. \quad (22)$$

The spin of the heavy diquark $S_{QQ} = 1$ can be coupled with the light quark $S_q = 1/2$. Thus, there are two possibilities for the total spin S , one is $1/2$ and the other is $3/2$. More detailed calculations of the S -wave states are found in Appendix A. The expectation value of $\mathbf{S}_{QQ} \cdot \mathbf{S}_q$ is obtained by Eq. (A3),

$$\langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle = \begin{bmatrix} -1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}. \quad (23)$$

The set of masses are

$$\begin{aligned} M(1/2, 1) &= \bar{M} - c_1, \\ M(3/2, 1) &= \bar{M} + \frac{1}{2}c_1. \end{aligned} \quad (24)$$

For the P -wave baryon system with $L = 1$, our purpose is to diagonalize the Hamiltonian in Eq. (19) involving all terms for the doubly heavy baryons,

$$H^{SD}(L = 1) = a_1 \mathbf{L} \cdot \mathbf{S}_{QQ} + a_2 \mathbf{L} \cdot \mathbf{S}_q + b_1 S_{12} + c_1 \mathbf{S}_{QQ} \cdot \mathbf{S}_q, \quad (25)$$

as the operator. Therefore, in contrast to the scheme used in Ref. [19], we proposed a new scheme of states classification named all- JLS coupling. Coupling $S_{QQ} = 1$ and $S_q = 1/2$ in L - S coupling scheme to form $S = 1/2$ or $3/2$, and then couple S and $L = 1$ to give $J = 1/2, 3/2$ or $1/2, 3/2, 5/2$ with negative parity $P = -1$. By using the all- JLS coupling, one can obtain the five mass shifts $\Delta M(J, j)$ in Eq. (B6) for the P -wave states, and refer to Appendix B. From Eqs. (21) and (B6), we find the following mass spectra of the doubly heavy baryons,

$$\begin{aligned} M(1/2, 0) &= \bar{M} + \frac{1}{4}(-a_1 - 6a_2 - 2b_1 - c_1) \\ &\quad - \frac{1}{12}\sqrt{(9a_1 - 2a_2 + 10b_1 - 7c_1)^2 + 8(2a_2 - b_1 - 2c_1)^2}, \\ M(1/2, 1) &= \bar{M} + \frac{1}{4}(-a_1 - 6a_2 - 2b_1 - c_1) \\ &\quad + \frac{1}{12}\sqrt{(9a_1 - 2a_2 + 10b_1 - 7c_1)^2 + 8(2a_2 - b_1 - 2c_1)^2}, \\ M(3/2, 1) &= \bar{M} + \frac{1}{20}(-5a_1 + 8b_1 - 5c_1) \\ &\quad - \frac{1}{60}\sqrt{(45a_1 - 40a_2 - 16b_1 - 5c_1)^2 + 5(20a_2 - 10b_1 - 20c_1)^2}, \end{aligned}$$

$$\begin{aligned}
M(3/2, 2) &= \bar{M} + \frac{1}{20} (-5a_1 + 8b_1 - 5c_1) \\
&\quad + \frac{1}{60} \sqrt{(45a_1 - 40a_2 - 16b_1 - 5c_1)^2 + 5(20a_2 - 10b_1 - 20c_1)^2}, \\
M(5/2, 2) &= \bar{M} + \frac{a_1}{2} + a_2 - \frac{b_1}{5} + \frac{c_1}{2}.
\end{aligned} \tag{26}$$

For D -wave baryon system with $L = 2$, we further discuss the mass expressions for calculating the doubly heavy baryons. The spin $S_{QQ} = 1$ in L - S coupling scheme can be coupled with $S_q = 1/2$ to determine the total spin $S = 1/2, 3/2$. Coupling of $L = 2$ give six states with $J = 3/2, 5/2$ or $1/2, 3/2, 5/2, 7/2$ with positive parity $P = +1$. Similarly, we also analyze the mass shifts $\Delta M(J, j)$ in Eq. (C4) for the D -wave states, and the Hamiltonian H^{SD} in Eq. (19) is treated as a representation operator,

$$H^{SD}(L = 2) = a_1 \mathbf{L} \cdot \mathbf{S}_{QQ} + a_2 \mathbf{L} \cdot \mathbf{S}_q + b_1 S_{12} + c_1 \mathbf{S}_{QQ} \cdot \mathbf{S}_q. \tag{27}$$

Details of calculating the mass shifts are presented in Appendix C. The mass spectra expressions can be written as

$$\begin{aligned}
M(1/2, 1) &= \bar{M} - \frac{3a_1}{2} - 3a_2 - b_1 + \frac{c_1}{2}, \\
M(3/2, 1) &= \bar{M} + \frac{1}{4} (-a_1 - 8a_2 - c_1) \\
&\quad - \frac{1}{20} \sqrt{(25a_1 - 16a_2 + 8b_1 - 9c_1)^2 + 36(2a_2 - b_1 - 2c_1)^2}, \\
M(3/2, 2) &= \bar{M} + \frac{1}{4} (-a_1 - 8a_2 - c_1) \\
&\quad + \frac{1}{20} \sqrt{(25a_1 - 16a_2 + 8b_1 - 9c_1)^2 + 36(2a_2 - b_1 - 2c_1)^2}, \\
M(5/2, 2) &= \bar{M} + \frac{1}{28} (-7a_1 + 14a_2 + 10b_1 - 7c_1) \\
&\quad - \frac{1}{140} \sqrt{(175a_1 - 182a_2 - 342b_1 + 7c_1)^2 + 2744(2a_2 - b_1 - 2c_1)^2}, \\
M(5/2, 3) &= \bar{M} + \frac{1}{28} (-7a_1 + 14a_2 + 10b_1 - 7c_1) \\
&\quad + \frac{1}{140} \sqrt{(175a_1 - 182a_2 - 342b_1 + 7c_1)^2 + 2744(2a_2 - b_1 - 2c_1)^2}, \\
M(7/2, 3) &= \bar{M} + a_1 + 2a_2 - \frac{2}{7}b_1 + \frac{c_1}{2}.
\end{aligned} \tag{28}$$

With the above expressions (24), (26), and (28), we present the estimates of the mass spectra of the ground and excited states in the Ξ_{QQ} and Ω_{QQ} baryon system in Sec. VI and Sec. VII, respectively.

V. THE SCALING RELATIONS OF DOUBLY HEAVY BARYONS

The study of data set of the spin-coupling parameters a_1 , a_2 , b_1 , and c_1 , as well as the effective mass of the quark, is of special interest to better understand the baryon structure. Under the color configurations, we utilize the scaling relations based on the similarity between a baryon and its partner baryons to study the spin-coupling parameters.

In our previous studies on the mass spectra of the singly heavy baryons Σ_Q , Ξ'_Q , and Ω_Q ($Q = c, b$) [19], we employed the following scaling relations to describe the parameters,

$$\begin{cases} a_1(B_a, (n+1)L) = \frac{M'_Q M'_d N'_{a_1}}{M_Q M_d N_{a_1}} a_1(B'_a, (n'+1)L'), \\ a_2(B_a, (n+1)L) = \frac{M'_Q M'_d N'_{a_2}}{M_Q M_d N_{a_2}} a_2(B'_a, (n'+1)L'), \\ b_1(B_a, (n+1)L) = \frac{M'_Q M'_d N'_{b_1}}{M_Q M_d N_{b_1}} b_1(B'_a, (n'+1)L'), \\ c_1(B_a, (n+1)L) = \frac{M'_Q M'_d N'_{c_1}}{M_Q M_d N_{c_1}} c_1(B'_a, (n'+1)L'), \end{cases} \quad (29)$$

where M'_Q , M_Q are the effective masses of heavy quarks, and M'_d , M_d are the effective masses of light diquark in the singly heavy baryon system. Then, $n, n' = 0, 1, 2, \dots$, $L, L' = S, P, D, F, \dots$, and B_a, B'_a are baryons with

$$\begin{aligned} N_{a_1} &= (n+L+1)^2 = N_{a_2}, \\ N_{b_1} &= L(L+1/2)(L+1)(n+L+1)^3, \\ N_{c_1} &= (L+\lambda)(n+L+1)^3, \end{aligned} \quad (30)$$

corresponding to the similar form of N'_{a_1} , N'_{a_2} , N'_{b_1} , N'_{c_1} with L' and n' , respectively. Here, the prime denotes the quantities of the baryon B'_a obtained from experiments, distinguishing them from that of an unobserved baryon B_a .

The parameters a_1 , a_2 , b_1 , and c_1 of the $1P$ -wave states for the singly heavy Ω_c baryon in Ref. [19] are treated as the object of the scaling relations in Eq. (29) to calculate the parameters of its partner baryon states, the results are listed in Table I.

TABLE I: The spin-coupling parameters (in GeV) of singly heavy baryons.

baryon	a_1	a_2	b_1	c_1
Ω_c	26.96	25.76	13.51	4.04
Σ_c	35.86	34.27	17.97	5.37
Ξ'_c	30.64	29.28	15.35	4.59

In order to further investigate the spin-coupling parameters a_1 , a_2 , b_1 , and c_1 of the doubly heavy Ξ_{QQ} and Ω_{QQ} baryons, we need to generalize Eq. (29). As these parameters in Ref. [25] should be related to the running coupling constant α_s from the Coulomb potential $V = -4\alpha_s/3r$. Thus, the scaling relations (29) become

$$\begin{cases} a_1(B_a, (n+1)L) = \frac{M'_{QQ}M'_q N'_{a_1}}{M_{QQ}M_q N_{a_1}} \frac{\alpha_s(B_a)}{\alpha_s(B'_a)} a_1(B'_a, (n'+1)L'), \\ a_2(B_a, (n+1)L) = \frac{M'_{QQ}M'_q N'_{a_2}}{M_{QQ}M_q N_{a_2}} \frac{\alpha_s(B_a)}{\alpha_s(B'_a)} a_2(B'_a, (n'+1)L'), \\ b_1(B_a, (n+1)L) = \frac{M'_{QQ}M'_q N'_{b_1}}{M_{QQ}M_q N_{b_1}} \frac{\alpha_s(B_a)}{\alpha_s(B'_a)} b_1(B'_a, (n'+1)L'), \\ c_1(B_a, (n+1)L) = \frac{M'_{QQ}M'_q N'_{c_1}}{M_{QQ}M_q N_{c_1}} \frac{\alpha_s(B_a)}{\alpha_s(B'_a)} c_1(B'_a, (n'+1)L'). \end{cases} \quad (31)$$

In this subsection, we estimate the effective masses of the quarks for the doubly heavy baryons. In general, it is extracted from the experimental values of mass spectra for the baryons that have been discovered. Different from the case of the singly heavy baryons and singly heavy mesons, apart from the two $\Xi_{cc}^{++}(3621)$ and $\Xi_{cc}^+(3519)$ states, there are no more experimental values for the doubly heavy Ξ_{QQ} and Ω_{QQ} baryons. Hence, when describing some of the details on how to estimate the quark masses in the Ξ_{QQ} and Ω_{QQ} baryons, we adopt the current masses of the quarks from PDG [26] as follows

$$\begin{aligned} m_c &= 1.273 \text{ GeV}, \quad m_b = 4.183 \text{ GeV}, \\ m_u &= 0.00216 \text{ GeV}, \quad m_d = 0.0047 \text{ GeV}, \quad m_s = 0.0935 \text{ GeV}. \end{aligned} \quad (32)$$

With this information and the help of the relations (7) and (8), we first consider the ρ -mode excited states, the masses of the heavy quarks c and b are obtained by

$$\begin{aligned} cc &: M_c = 1381.03 \text{ MeV}, \quad m_{cc} = M_c + M_c = 2762.06 \text{ MeV}, \\ bc &: M_c = 1451.20 \text{ MeV}, \quad M_b = 4228.38 \text{ MeV}, \quad m_{bc} = M_c + M_b = 5679.58 \text{ MeV}, \\ bb &: M_b = 4537.99 \text{ MeV}, \quad m_{bb} = M_b + M_b = 9075.98 \text{ MeV}. \end{aligned} \quad (33)$$

And then the effective masses of the heavy diquark and light quark for the λ -mode excited states are calculated, with the results

$$\begin{aligned} M_{cc} &= 2763.20 \text{ MeV}, \quad M_{bc} = 5680.15 \text{ MeV}, \quad M_{bb} = 9076.33 \text{ MeV}, \\ M_u &= 4.25 \text{ MeV}, \quad M_d = 9.24 \text{ MeV}, \quad M_s = 176.18 \text{ MeV}. \end{aligned} \quad (34)$$

Note that only a part of the effective masses of the quarks for 1S-wave states are described for simplicity. In addition, in Eqs. (12) and (30), there are two parameters which should be fixed, *i.e.*,

k and λ . In the following calculations, we take $k = 1.0$ and $\lambda = 0.68$ by analyzing the experimental values. For applying Eq. (31) and the parameters in Table I, one can calculate the spin-coupling parameters for the Ξ_{QQ} baryons listed in Table II, and in Table III for the Ω_{QQ} baryons with $\alpha_s(B'_a) = 0.593$ GeV and $\alpha_s(B_a) = 0.557$ GeV corresponding to the singly heavy baryons and doubly heavy baryons, respectively.

VI. THE Ξ_{QQ} BARYONS

Let us now begin with calculating the masses of the ground and excited states for the doubly heavy Ξ_{cc} , Ξ_{bc} , and Ξ_{bb} baryons. It was a pleasant surprise that the doubly charmed baryon Ξ_{cc}^{++} is observed by the LHCb Collaboration in the $\Lambda_c^+ K^- \pi^+ \pi^+$ mass spectrum [1]. The mass reported for this resonance of Ξ_{cc}^{++} is $M(\Xi_{cc}^{++}) = 3621.40 \pm 0.78$ MeV, but its J^P value is still unknown. In addition, the LHCb Collaboration has also carried out searches for the doubly heavy baryon Ξ_{cc}^+ decaying to $D^0 p K^-$ [27], and for the doubly heavy baryons Ξ_{cc}^0 and Ω_{cc}^0 decaying to $\Lambda_c^+ \pi^-$ and $\Xi_c^+ \pi^-$ in Refs. [28, 29], these baryons have not yet been observed. Many workers try to study their properties and internal structure for the doubly heavy Ξ_{QQ} baryons, we recommend interested readers to see Refs. [30–38].

Based on the close similarity of the strong interaction binds the heavy diquark and light quark in hadron system, we obtain a relation between the mass splitting of the doubly heavy baryons and that of the singly heavy mesons ($Q\bar{q}$) in Ref. [25],

$$\Delta M(\Xi_{QQ}) = \frac{3}{2} \frac{M_Q M_{\bar{q}}}{M_{QQ} M_q} \Delta M(B, D), \quad (35)$$

with the factor $3/2$ being just the ratio of the baryon and meson spin matrix elements.

From PDG [26], the D^\pm and $D^*(2010)^\pm$ states, which are regarded as $1S$ -wave states corresponding to the masses $M(D^\pm) = 1869.66$ MeV and $M(D^*(2010)^\pm) = 2010.26$ MeV with $J^P = 1/2^-$ and $3/2^-$, respectively. Thus, the hyperfine splitting between D^\pm and $D^*(2010)^\pm$ would be $\Delta M(D) = M(D^*(2010)^\pm) - M(D^\pm) = 140.60$ MeV. Substituting this value into Eq. (35), the predicted hyperfine splitting of $1S$ -wave for the Ξ_{cc} baryon is given as follows

$$\begin{aligned} \Delta M(\Xi_{cc}) &= \frac{3}{2} \frac{M_c M_{\bar{u}}}{M_{cc} M_u} \Delta M(D) \\ &= 78.12 \text{ MeV}, \end{aligned} \quad (36)$$

with $M_c = 1273.00$ MeV, $M_{cc} = 2762.07$ MeV, $M_{\bar{u}} = 3.42$ MeV, and $M_u = 4.25$ MeV. By applying

TABLE II: Spin coupling parameters (MeV) of Ξ_{QQ} baryons.

Baryon	State	a_1	a_2	b_1	c_1
Ξ_{cc}	1S				52.08
	2S				11.57
	3S				3.64
	4S				1.56
	5S				0.81
	1P	31.27	29.88	4.64	4.68
	2P	14.75	14.10	1.46	1.47
	3P	8.46	8.08	0.63	0.63
	4P	5.46	5.22	0.32	0.33
	5P	3.81	3.64	0.19	0.19
	1D	14.75	14.10	0.29	0.92
	2D	8.46	8.08	0.13	0.40
	3D	5.46	5.22	0.06	0.21
	4D	3.81	3.64	0.04	0.12
	5D	2.81	2.68	0.02	0.08
Ξ_{bc}	1S				25.30
	2S				5.62
	3S				1.77
	4S				0.76
	5S				0.39
	1P	15.21	14.53	0.67	2.28
	2P	7.17	6.85	0.21	0.72
	3P	4.11	3.93	0.09	0.31
	4P	2.66	2.54	0.05	0.16
	5P	1.85	1.77	0.03	0.09
	1D	7.17	6.85	0.04	0.45
	2D	4.11	3.93	0.02	0.19
	3D	2.66	2.54	0.01	0.10
	4D	1.85	1.77	0.005	0.06
	5D	1.37	1.30	0.003	0.04
Ξ_{bb}	1S				15.83
	2S				3.52
	3S				1.11
	4S				0.48
	5S				0.25
	1P	9.52	9.09	0.42	1.42
	2P	4.49	4.29	0.13	0.45
	3P	2.57	2.46	0.06	0.19
	4P	1.66	1.59	0.03	0.10
	5P	1.16	1.11	0.02	0.06
	1D	4.49	4.29	0.03	0.28
	2D	2.57	2.46	0.01	0.12
	3D	1.66	1.59	0.006	0.06
	4D	1.16	1.11	0.003	0.04
	5D	0.85	0.82	0.002	0.02

TABLE III: Spin coupling parameters (MeV) of Ω_{QQ} baryons.

Baryon	State	a_1	a_2	b_1	c_1
Ω_{cc}	1S				40.39
	2S				8.62
	3S				2.70
	4S				1.16
	5S				0.60
	1P	23.28	22.24	3.46	3.49
	2P	10.96	10.47	1.08	1.09
	3P	6.28	6.00	0.47	0.47
	4P	4.05	3.87	0.24	0.24
	5P	2.83	2.70	0.14	0.14
	1D	10.96	10.47	0.22	0.69
	2D	6.28	6.00	0.09	0.29
	3D	4.05	3.87	0.05	0.15
	4D	2.83	2.70	0.03	0.09
	5D	2.08	1.99	0.02	0.06
Ω_{bc}	1S				19.22
	2S				4.19
	3S				1.31
	4S				0.57
	5S				0.29
	1P	11.30	10.80	0.50	1.69
	2P	5.33	5.09	0.16	0.53
	3P	3.05	2.92	0.07	0.23
	4P	1.97	1.88	0.03	0.12
	5P	1.37	1.31	0.02	0.07
	1D	5.33	5.09	0.03	0.33
	2D	3.05	2.92	0.01	0.14
	3D	1.97	1.88	0.007	0.07
	4D	1.37	1.31	0.004	0.04
	5D	1.01	0.97	0.003	0.03
Ω_{bb}	1S				11.93
	2S				2.62
	3S				0.82
	4S				0.35
	5S				0.18
	1P	7.07	6.75	0.31	1.06
	2P	3.33	3.18	0.10	0.33
	3P	1.91	1.82	0.04	0.14
	4P	1.23	1.18	0.02	0.07
	5P	0.86	0.82	0.01	0.04
	1D	3.33	3.18	0.02	0.21
	2D	1.91	1.82	0.01	0.09
	3D	1.23	1.18	0.004	0.05
	4D	0.86	0.82	0.003	0.03
	5D	0.63	0.61	0.002	0.02

Eq. (18), one estimates the spin-average mass of 1S-wave ($n = 0, L = 0$) for the Ξ_{cc}^{++} baryon,

$$\begin{aligned}\bar{M}(\Xi_{cc}) &= M_{cc} + \left(\frac{1}{2}(M_{cc})^{1/2} + \left(M_u + \frac{M_{cc}m_u}{m_{cc}^2} \frac{4\alpha_s}{3} \mu_{ccu} \right)^2 \right)^{\frac{1}{2}} \\ &= 3673.65 \text{ MeV},\end{aligned}\quad (37)$$

with $\mu_{ccu} = 2.158$ MeV, as well as the following rough estimate for the parameter c_1 by Eq. (31),

$$c_1(\Xi_{cc}) = \frac{M_c M_{uu}}{M_{cc} M_u} \frac{N'_{c_1}}{N_{c_1}} \frac{\alpha_s(\Xi_{cc})}{\alpha_s(\Sigma_c)} c_1(\Sigma_c, 1P) = 52.08 \text{ MeV}.\quad (38)$$

Note that so far no other experiments confirmed the existence of the Ξ_{cc}^{++} [39, 40], as they find no evidence for the Ξ_{cc}^+ baryon in $\Lambda_c^+ K^- \pi^+$ and $\Xi_c^0 \pi^+$ decay, and the Ξ_{cc}^{++} baryon in $\Lambda_c^+ K^- \pi^+ \pi^+$ and $\Xi_c^0 \pi^+ \pi^+$ decay. However, in our model, the $\Xi_{cc}^{++}(3621)$ can be grouped into the 1S state. We assign $J^P = 1/2^+$. Thus, the mass spectra of the two 1S-wave states for the Ξ_{cc} baryon are given by

$$\begin{aligned}\text{State}|J, j\rangle : & \quad |1/2, 1\rangle & \quad |3/2, 1\rangle \\ M(\Xi_{cc}^{++}, 1S): & \quad 3621.57 \text{ MeV} & \quad 3699.69 \text{ MeV}.\end{aligned}\quad (39)$$

From the predicted values of the constituent quark model in Ref. [4], our prediction for $3/2^+$ state differs only by 9.69 MeV. For the ground and excited states for the Ξ_{cc} baryon, the result of the masses are shown in Table IV by our model and compared with other models, and refer to Refs. [44–49] for more details.

In 2021, the neutral doubly heavy baryon $\Xi_{bc}^0 (bcd)$ was searched by the LHCb experiment in $\Xi_{bc}^0 \rightarrow \Xi_c^+ \pi^-$ decay [28], no significant excess is found for invariant masses between 6700 and 7300 MeV. The LHCb experiment had searched the baryon Ξ_{bc}^+ (bcd) in the charged decay mode $\Xi_{bc}^+ \rightarrow J/\psi \Xi_c^+$ [29] in 2023. The most significant peaks in the mass region considered correspond to local (global) significance of 4.3σ (2.8σ) and 4.1σ (2.4σ) at 6571 MeV and 6694 MeV. Thus, there are no evidence for the Ξ_{bc}^0 and Ξ_{bc}^+ baryons with the current data sample. According to this information for the Ξ_{bc} and Ξ_{bb} baryons in our model, the spin-average masses of the ground states are given by using the spin-independent mass formula (18) with $n = 0, L = 0, \mu_{bcu} = 2.159$ MeV, and $\mu_{bbu} = 2.160$ MeV,

$$\begin{aligned}\bar{M}(\Xi_{bc}) &= M_{bc} + \left(\frac{1}{2}(M_{bc})^{1/2} + \left(M_u + \frac{M_{bc}m_u}{m_{bc}^2} \frac{4\alpha_s}{3} \mu_{bcu} \right)^2 \right)^{\frac{1}{2}} \\ &= 6771.19 \text{ MeV},\end{aligned}\quad (40)$$

and

$$\bar{M}(\Xi_{bb}) = M_{bb} + \left(\frac{1}{2}(M_{bb})^{1/2} + \left(M_u + \frac{M_{bb}m_u}{m_{bb}^2} \frac{4\alpha_s}{3} \mu_{bbu} \right)^2 \right)^{\frac{1}{2}}$$

$$= 10303.31 \text{ MeV}. \quad (41)$$

We utilize the scaling relation (31) to calculate the spin coupling parameter c_1 for the Ξ_{bc} and Ξ_{bb} baryons, the results are obtained by

$$c_1(\Xi_{bc}) = \frac{M_{cc}M'_u}{M_{bc}M_u} \frac{N'_{c_1}}{N_{c_1}} \frac{\alpha_s(\Xi_{bc})}{\alpha_s(\Xi_{cc})} c_1(\Xi_{cc}, 1P) = 25.30 \text{ MeV}, \quad (42)$$

$$c_1(\Xi_{bb}) = \frac{M_{cc}M'_u}{M_{bb}M_u} \frac{N'_{c_1}}{N_{c_1}} \frac{\alpha_s(\Xi_{bb})}{\alpha_s(\Xi_{cc})} c_1(\Xi_{cc}, 1P) = 15.83 \text{ MeV}. \quad (43)$$

Therefore, the predicted masses of the $1S$ -wave states are

$$\begin{aligned} \text{State}|J, j\rangle : & \quad |1/2, 1\rangle & \quad |3/2, 1\rangle \\ M(\Xi_{bc}, 1S): & \quad 6745.89 \text{ MeV} & \quad 6783.84 \text{ MeV}, \end{aligned} \quad (44)$$

and

$$\begin{aligned} \text{State}|J, j\rangle : & \quad |1/2, 1\rangle & \quad |3/2, 1\rangle \\ M(\Xi_{bb}, 1S): & \quad 10287.48 \text{ MeV} & \quad 10311.22 \text{ MeV}, \end{aligned} \quad (45)$$

respectively. In our model calculations, the $1S$ -wave level-splitting for the partner Ξ_{bc} and Ξ_{bb} baryons are predicted to be $\Delta M(\Xi_{bc}) = 37.95 \text{ MeV}$ and $\Delta M(\Xi_{bb}) = 23.74 \text{ MeV}$ smaller than 47 MeV and 35 MeV in Ref. [25], respectively. For the Ξ_{bc} and Ξ_{bb} baryons, by exploiting the mass spectra expressions (24), (26), and (28), we calculate the mass spectra listed in Table V and Table VI. Calculation of the mass spectra of doubly heavy Ξ_{cc} , Ξ_{bc} , and Ξ_{bb} baryons together with a more detailed discussion can be found in Refs. [50–56], (see also [57–59]).

VII. THE Ω_{QQ} BARYONS

The doubly heavy Ω_{QQ} baryons are regarded as an important and unique part of the baryons in heavy-light quark system, as they are composed of light strange quark s . Up to now, the doubly heavy Ω_{QQ} baryons have not been reported yet by the experimental, and their nature are still unknown. Thus, the doubly heavy Ω_{QQ} baryons were studied by using various approaches, we recommend interested readers to see Refs. [59–66]. In this section, we may apply the similar methods to investigate the mass spectra of the Ω_{cc} , Ω_{bc} , and Ω_{bb} baryons. The calculation results of the masses are listed in Table VII, VIII, and IX for the Ω_{cc} , Ω_{bc} , and Ω_{bb} baryons, respectively.

For the Ω_{cc} baryon, the spin-averaged mass of the $1S$ -wave states with $\mu_{ccs} = 90.44 \text{ MeV}$ by applying Eq. (18) is predicted to be

$$\bar{M}(\Omega_{cc}) = M_{cc} + \left(\frac{1}{2}(M_{cc})^{1/2} + \left(M_s + \frac{M_{cc}m_s}{m_{cc}^2} \frac{4\alpha_s}{3} \mu_{ccs} \right)^2 \right)^{\frac{1}{2}}$$

TABLE IV: Mass spectra (MeV) of Ξ_{cc} baryons are given and compared with different quark models.

State	J^P	Ours	[5]	[41]	[42]	[43]
$1^1 S_{1/2}$	$1/2^+$	3621.57	3620	3606	3678	3627
$1^3 S_{3/2}$	$3/2^+$	3699.69	3727	3675	3752	3690
$2^1 S_{1/2}$	$1/2^+$	4153.86		4004	4311	
$2^3 S_{3/2}$	$3/2^+$	4171.21		4036	4368	
$3^1 S_{1/2}$	$1/2^+$	4521.34				
$3^3 S_{3/2}$	$3/2^+$	4526.79				
$4^1 S_{1/2}$	$1/2^+$	4821.15				
$4^3 S_{3/2}$	$3/2^+$	4823.50				
$5^1 S_{1/2}$	$1/2^+$	5081.76				
$5^3 S_{3/2}$	$3/2^+$	5082.97				
$1^2 P_{1/2}$	$1/2^-$	3972.72	4053	3998	4081	
$1^4 P_{1/2}$	$1/2^-$	4017.48	4136	3985	4073	
$1^2 P_{3/2}$	$3/2^-$	4026.94	4101	4014	4077	
$1^4 P_{3/2}$	$3/2^-$	4061.26	4196	4025	4079	
$1^4 P_{5/2}$	$5/2^-$	4098.16	4155	4050	4089	
$2^2 P_{1/2}$	$1/2^-$	4398.91				
$2^4 P_{1/2}$	$1/2^-$	4420.24				
$2^2 P_{3/2}$	$3/2^-$	4423.09				
$2^4 P_{3/2}$	$3/2^-$	4440.97				
$2^4 P_{5/2}$	$5/2^-$	4457.42				
$3^2 P_{1/2}$	$1/2^-$	4725.85				
$3^4 P_{1/2}$	$1/2^-$	4738.15				
$3^2 P_{3/2}$	$3/2^-$	4739.32				
$3^4 P_{3/2}$	$3/2^-$	4750.06				
$3^4 P_{5/2}$	$5/2^-$	4759.22				
$4^2 P_{1/2}$	$1/2^-$	5001.92				
$4^4 P_{1/2}$	$1/2^-$	5009.89				
$4^2 P_{3/2}$	$3/2^-$	5010.46				
$4^4 P_{3/2}$	$3/2^-$	5017.59				
$4^4 P_{5/2}$	$5/2^-$	5023.39				
$5^2 P_{1/2}$	$1/2^-$	5245.85				
$5^4 P_{1/2}$	$1/2^-$	5251.43				
$5^2 P_{3/2}$	$3/2^-$	5251.74				
$5^4 P_{3/2}$	$3/2^-$	5256.80				
$5^4 P_{5/2}$	$5/2^-$	5260.80				
$1^4 D_{1/2}$	$1/2^+$	4276.72				
$1^2 D_{3/2}$	$3/2^+$	4298.45				
$1^4 D_{3/2}$	$3/2^+$	4319.26				
$1^2 D_{5/2}$	$5/2^+$	4334.44				
$1^4 D_{5/2}$	$5/2^+$	4353.97				
$1^4 D_{7/2}$	$7/2^+$	4384.29				
$2^4 D_{1/2}$	$1/2^+$	4630.81				
$2^2 D_{3/2}$	$3/2^+$	4643.24				
$2^4 D_{3/2}$	$3/2^+$	4655.36				
$2^2 D_{5/2}$	$5/2^+$	4663.84				
$2^4 D_{5/2}$	$5/2^+$	4675.26				
$2^4 D_{7/2}$	$7/2^+$	4692.47				
$3^4 D_{1/2}$	$1/2^+$	4922.24				
$3^2 D_{3/2}$	$3/2^+$	4930.25				
$3^4 D_{3/2}$	$3/2^+$	4938.15				
$3^2 D_{5/2}$	$5/2^+$	4943.54				
$3^4 D_{5/2}$	$5/2^+$	4951.00				
$3^4 D_{7/2}$	$7/2^+$	4962.03				
$4^4 D_{1/2}$	$1/2^+$	5176.13				
$4^2 D_{3/2}$	$3/2^+$	5181.71				
$4^4 D_{3/2}$	$3/2^+$	5187.26				
$4^2 D_{5/2}$	$5/2^+$	5190.98				
$4^4 D_{5/2}$	$5/2^+$	5196.22				
$4^4 D_{7/2}$	$7/2^+$	5203.89				
$5^4 D_{1/2}$	$1/2^+$	5404.38				
$5^2 D_{3/2}$	$3/2^+$	5408.48				
$5^4 D_{3/2}$	$3/2^+$	5412.59				
$5^2 D_{5/2}$	$5/2^+$	5415.31				
$5^4 D_{5/2}$	$5/2^+$	5419.19				
$5^4 D_{7/2}$	$7/2^+$	5424.82				

TABLE V: Mass spectra (MeV) of Ξ_{bc} baryons are given and compared with different quark models.

State	J^P	Ours	[5]	[42]	[44]	[4]
$1^1 S_{1/2}$	$1/2^+$	6745.89	6933	7014	6904	6914
$1^3 S_{3/2}$	$3/2^+$	6783.84	6980	7064	6936	6969
$2^1 S_{1/2}$	$1/2^+$	7354.46		7634	7478	
$2^3 S_{3/2}$	$3/2^+$	7362.90		7676	7495	
$3^1 S_{1/2}$	$1/2^+$	7788.87			7904	
$3^3 S_{3/2}$	$3/2^+$	7791.53			7917	
$4^1 S_{1/2}$	$1/2^+$	8146.42				
$4^3 S_{3/2}$	$3/2^+$	8147.56				
$5^1 S_{1/2}$	$1/2^+$	8457.95				
$5^3 S_{3/2}$	$3/2^+$	8458.54				
$1^2 P_{1/2}$	$1/2^-$	7186.89		7397		
$1^4 P_{1/2}$	$1/2^-$	7206.79		7390		
$1^2 P_{3/2}$	$3/2^-$	7210.25		7392		
$1^4 P_{3/2}$	$3/2^-$	7228.23		7394		
$1^4 P_{5/2}$	$5/2^-$	7246.48		7399		
$2^2 P_{1/2}$	$1/2^-$	7666.23				
$2^4 P_{1/2}$	$1/2^-$	7676.04				
$2^2 P_{3/2}$	$3/2^-$	7677.06				
$2^4 P_{3/2}$	$3/2^-$	7686.16				
$2^4 P_{5/2}$	$5/2^-$	7694.26				
$3^2 P_{1/2}$	$1/2^-$	8046.25				
$3^4 P_{1/2}$	$1/2^-$	8051.99				
$3^2 P_{3/2}$	$3/2^-$	8052.40				
$3^4 P_{3/2}$	$3/2^-$	8057.80				
$3^4 P_{5/2}$	$5/2^-$	8062.29				
$4^2 P_{1/2}$	$1/2^-$	8371.44				
$4^4 P_{1/2}$	$1/2^-$	8375.19				
$4^2 P_{3/2}$	$3/2^-$	8375.39				
$4^4 P_{3/2}$	$3/2^-$	8378.94				
$4^4 P_{5/2}$	$5/2^-$	8381.79				
$5^2 P_{1/2}$	$1/2^-$	8660.59				
$5^4 P_{1/2}$	$1/2^-$	8663.23				
$5^2 P_{3/2}$	$3/2^-$	8663.34				
$5^4 P_{3/2}$	$3/2^-$	8665.85				
$5^4 P_{5/2}$	$5/2^-$	8667.81				
$1^4 D_{1/2}$	$1/2^+$	7539.15				
$1^2 D_{3/2}$	$3/2^+$	7549.62				
$1^4 D_{3/2}$	$3/2^+$	7559.73				
$1^2 D_{5/2}$	$5/2^+$	7567.05				
$1^4 D_{5/2}$	$5/2^+$	7576.61				
$1^4 D_{7/2}$	$7/2^+$	7591.39				
$2^4 D_{1/2}$	$1/2^+$	7943.64				
$2^2 D_{3/2}$	$3/2^+$	7949.64				
$2^4 D_{3/2}$	$3/2^+$	7955.54				
$2^2 D_{5/2}$	$5/2^+$	7959.63				
$2^4 D_{5/2}$	$5/2^+$	7965.21				
$2^4 D_{7/2}$	$7/2^+$	7973.59				
$3^4 D_{1/2}$	$1/2^+$	8283.32				
$3^2 D_{3/2}$	$3/2^+$	8287.19				
$3^4 D_{3/2}$	$3/2^+$	8291.03				
$3^2 D_{5/2}$	$5/2^+$	8293.64				
$3^4 D_{5/2}$	$5/2^+$	8297.28				
$3^4 D_{7/2}$	$7/2^+$	8302.65				
$4^4 D_{1/2}$	$1/2^+$	8582.23				
$4^2 D_{3/2}$	$3/2^+$	8584.93				
$4^4 D_{3/2}$	$3/2^+$	8587.62				
$4^2 D_{5/2}$	$5/2^+$	8589.42				
$4^4 D_{5/2}$	$5/2^+$	8591.98				
$4^4 D_{7/2}$	$7/2^+$	8595.71				
$5^4 D_{1/2}$	$1/2^+$	8852.43				
$5^2 D_{3/2}$	$3/2^+$	8854.42				
$5^4 D_{3/2}$	$3/2^+$	8856.41				
$5^2 D_{5/2}$	$5/2^+$	8857.73				
$5^4 D_{5/2}$	$5/2^+$	8859.63				
$5^4 D_{7/2}$	$7/2^+$	8862.37				

TABLE VI: Mass spectra (MeV) of Ξ_{bb} baryons are given and compared with different quark models.

State	J^P	Ours	[5]	[41]	[30]	[42]
$1^1 S_{1/2}$	$1/2^+$	10287.48	10202	10138	10171	10322
$1^3 S_{3/2}$	$3/2^+$	10311.22	10237	10169	10195	10352
$2^1 S_{1/2}$	$1/2^+$	10961.90	10832	10662	10738	10940
$2^3 S_{3/2}$	$3/2^+$	10967.18	10860	10675	10753	10972
$3^1 S_{1/2}$	$1/2^+$	11448.40				
$3^3 S_{3/2}$	$3/2^+$	11450.06				
$4^1 S_{1/2}$	$1/2^+$	11849.90				
$4^3 S_{3/2}$	$3/2^+$	11850.62				
$5^1 S_{1/2}$	$1/2^+$	12199.99				
$5^3 S_{3/2}$	$3/2^+$	12200.36				
$1^2 P_{1/2}$	$1/2^-$	10788.86	10632	10525	10593	10694
$1^4 P_{1/2}$	$1/2^-$	10801.32	10675	10504	10547	10694
$1^2 P_{3/2}$	$3/2^-$	10803.49	10647	10526	10606	10691
$1^4 P_{3/2}$	$3/2^-$	10814.73	10694	10528	10561	10692
$1^4 P_{5/2}$	$5/2^-$	10826.16	10661	10547	10560	10695
$2^2 P_{1/2}$	$1/2^-$	11318.24				
$2^4 P_{1/2}$	$1/2^-$	11324.38				
$2^2 P_{3/2}$	$3/2^-$	11325.02				
$2^4 P_{3/2}$	$3/2^-$	11330.71				
$2^4 P_{5/2}$	$5/2^-$	11335.78				
$3^2 P_{1/2}$	$1/2^-$	11741.84				
$3^4 P_{1/2}$	$1/2^-$	11745.44				
$3^2 P_{3/2}$	$3/2^-$	11745.69				
$3^4 P_{3/2}$	$3/2^-$	11749.07				
$3^4 P_{5/2}$	$5/2^-$	11751.88				
$4^2 P_{1/2}$	$1/2^-$	12105.71				
$4^4 P_{1/2}$	$1/2^-$	12108.06				
$4^2 P_{3/2}$	$3/2^-$	12108.19				
$4^4 P_{3/2}$	$3/2^-$	12110.41				
$4^4 P_{5/2}$	$5/2^-$	12112.19				
$5^2 P_{1/2}$	$1/2^-$	12429.85				
$5^4 P_{1/2}$	$1/2^-$	12431.51				
$5^2 P_{3/2}$	$3/2^-$	12431.57				
$5^4 P_{3/2}$	$3/2^-$	12433.14				
$5^4 P_{5/2}$	$5/2^-$	12434.37				
$1^4 D_{1/2}$	$1/2^+$	11182.28			10913	
$1^2 D_{3/2}$	$3/2^+$	11188.83			10918	
$1^4 D_{3/2}$	$3/2^+$	11195.16			10798	
$1^2 D_{5/2}$	$5/2^+$	11199.74			10921	
$1^4 D_{5/2}$	$5/2^+$	11205.72			10803	
$1^4 D_{7/2}$	$7/2^+$	11214.97			10805	
$2^4 D_{1/2}$	$1/2^+$	11630.45				
$2^2 D_{3/2}$	$3/2^+$	11634.20				
$2^4 D_{3/2}$	$3/2^+$	11637.89				
$2^2 D_{5/2}$	$5/2^+$	11640.46				
$2^4 D_{5/2}$	$5/2^+$	11643.95				
$2^4 D_{7/2}$	$7/2^+$	11649.19				
$3^4 D_{1/2}$	$1/2^+$	12009.20				
$3^2 D_{3/2}$	$3/2^+$	12011.63				
$3^4 D_{3/2}$	$3/2^+$	12014.03				
$3^2 D_{5/2}$	$5/2^+$	12015.66				
$3^4 D_{5/2}$	$5/2^+$	12017.94				
$3^4 D_{7/2}$	$7/2^+$	12021.30				
$4^4 D_{1/2}$	$1/2^+$	12343.54				
$4^2 D_{3/2}$	$3/2^+$	12345.22				
$4^4 D_{3/2}$	$3/2^+$	12346.91				
$4^2 D_{5/2}$	$5/2^+$	12348.04				
$4^4 D_{5/2}$	$5/2^+$	12349.64				
$4^4 D_{7/2}$	$7/2^+$	12351.98				
$5^4 D_{1/2}$	$1/2^+$	12646.28				
$5^2 D_{3/2}$	$3/2^+$	12647.52				
$5^4 D_{3/2}$	$3/2^+$	12648.77				
$5^2 D_{5/2}$	$5/2^+$	12649.59				
$5^4 D_{5/2}$	$5/2^+$	12650.78				
$5^4 D_{7/2}$	$7/2^+$	12652.49				

$$= 3692.18 \text{ MeV}. \quad (46)$$

Combining the value of the $c_1(\Omega_{cc}) = 40.39 \text{ MeV}$ show in Table III, we obtain the masses with $J^P = 1/2^+$ and $3/2^+$ for the Ω_{cc} states,

$$\begin{aligned} \text{State}|J, j\rangle : & \quad |1/2, 1\rangle & \quad |3/2, 1\rangle \\ M(\Omega_{cc}, 1S): & \quad 3651.79 \text{ MeV} & \quad 3712.37 \text{ MeV}. \end{aligned} \quad (47)$$

It can be seen that our result of $1S$ -wave level-splitting mass is about $\Delta M(\Omega_{cc}, 1S) = 60 \text{ MeV}$ smaller than the result $\Delta M(\Omega_{cc}, 1S) = 94 \text{ MeV}$ in Ref. [5]. Over the past two decades, the properties of the Ω_{cc} baryon with one strange quark can be explored experimentally. For example, in Ref. [68], the LHCb experiment had searched the doubly charmed baryon Ω_{cc}^+ in the charged decay mode $\Omega_{cc}^+ \rightarrow \Lambda_c^+ K^- \pi^+$. No significant signal is observed within the invariant mass range of 3.6 to 4.0 GeV. Additionally, we predict the spin-averaged mass of the $2S$ -wave Ω_{cc} states with $L = 0$, $n = 1$ gives

$$\bar{M}(\Omega_{cc}, 2S) = 4169.58 \text{ MeV}, \quad (48)$$

which are about 477 MeV higher than the spin-averaged mass in Eq. (46) of the $1S$ -wave states for the Ω_{cc} baryon. Thus, the results of the masses for the $2S$ -wave are

$$\begin{aligned} \text{State}|J, j\rangle : & \quad |1/2, 1\rangle & \quad |3/2, 1\rangle \\ M(\Omega_{cc}, 2S): & \quad 4160.96 \text{ MeV} & \quad 4173.89 \text{ MeV}. \end{aligned} \quad (49)$$

On the other hand, we also have calculated the spin-average of the $1P$ -wave ($L = 1$, $n = 0$) for Ω_{cc} states,

$$\begin{aligned} \bar{M}(\Omega_{cc}, 1P) &= M_{cc} + \left(\frac{1}{2}(M_{cc})^{\frac{1}{2}} \times 2 + \left(M_s + \frac{M_{cc}m_s}{m_{cc}^2} \frac{4\alpha_s}{3} \frac{\mu_{ccs}}{4} \right)^2 \right)^{\frac{1}{2}} \\ &= 4055.73 \text{ MeV}, \end{aligned} \quad (50)$$

the parameters are

$$\begin{aligned} a_1(\Omega_{cc}, 1P) &= \frac{M_c M_{ss}}{M_{bc} M_u} \frac{N'_{c_1}}{N_{c_1}} \frac{\alpha_s(\Omega_{cc})}{\alpha_s(\Omega_c)} c_1(\Omega_c, 1P) = 23.28 \text{ MeV}, \\ a_2(\Omega_{cc}, 1P) &= \frac{M_c M_{ss}}{M_{bc} M_u} \frac{N'_{c_1}}{N_{c_1}} \frac{\alpha_s(\Omega_{cc})}{\alpha_s(\Omega_c)} c_1(\Omega_c, 1P) = 22.24 \text{ MeV}, \\ b_1(\Omega_{cc}, 1P) &= \frac{M_c M_{ss}}{M_{bc} M_u} \frac{N'_{c_1}}{N_{c_1}} \frac{\alpha_s(\Omega_{cc})}{\alpha_s(\Omega_c)} c_1(\Omega_c, 1P) = 3.46 \text{ MeV}, \\ c_1(\Omega_{cc}, 1P) &= \frac{M_c M_{ss}}{M_{bc} M_u} \frac{N'_{c_1}}{N_{c_1}} \frac{\alpha_s(\Omega_{cc})}{\alpha_s(\Omega_c)} c_1(\Omega_c, 1P) = 3.49 \text{ MeV}. \end{aligned} \quad (51)$$

The obtained masses with the bases $|J, j\rangle$ are

$$\begin{aligned} \text{State}|J, j\rangle : & \quad |1/2, 0\rangle \quad |1/2, 1\rangle \quad |3/2, 1\rangle \quad |3/2, 2\rangle \quad |5/2, 2\rangle \\ M(\Omega_{cc}, 1P): & \quad 3997.28 \text{ MeV} \quad 4030.60 \text{ MeV} \quad 4037.65 \text{ MeV} \quad 4063.19 \text{ MeV} \quad 4090.66 \text{ MeV}. \end{aligned} \quad (52)$$

Thus, our level-splitting mass for $1P$ -wave Ω_{cc} states is expected to be about 50 larger than the Ξ_{cc} states in Table IV. More valuable information of the Ω_{cc} baryon can be provided to the further experimental exploration.

In Ref. [28], the LHCb experiment had searched the baryon Ω_{bc}^0 (bcs) in the $\Omega_{bc}^0 \rightarrow \Lambda_c^+ \pi^-$ decay. The search for the Ω_{bc}^0 baryon is performed in the mass range between 6700 MeV and 7300 MeV, no significant excess is found in the LHCb experiment. In our work, similar to the case for the Ω_{bc} baryon, we calculate the spin-average masses of the ground states with $\mu_{bcs} = 91.99$ MeV,

$$\begin{aligned} \bar{M}(\Omega_{bc}) &= M_{bc} + \left(\frac{1}{2}(M_{bc})^{1/2} + \left(M_s + \frac{M_{bc}m_s}{m_{bc}^2} \frac{4\alpha_s}{3} \mu_{bcs} \right)^2 \right)^{\frac{1}{2}} \\ &= 6786.72 \text{ MeV}. \end{aligned} \quad (53)$$

The predicted masses are given by

$$\begin{aligned} \text{State}|J, j\rangle : & \quad |1/2, 1\rangle \quad |3/2, 1\rangle \\ M(\Omega_{bc}, 1S): & \quad 6767.50 \text{ MeV} \quad 6796.33 \text{ MeV}. \end{aligned} \quad (54)$$

The mass difference of the $M(\Omega_{bc}, 1/2^+)$ and $M(\Omega_{bc}, 3/2^+)$ states is predicted to be 28 MeV, which is comparable with these results from the relativistic quark model Ref. [5]. The computed ground and excited states for the Ω_{bc} baryons are compared with different theoretical approaches shown in Table VIII.

We also calculate the predicted mass of the $1S$ -wave state with $\mu_{bcs} = 92.55$ MeV for the Ω_{bb} baryon,

$$\begin{aligned} \bar{M}(\Omega_{bb}) &= M_{bb} + \left(\frac{1}{2}(M_{bb})^{1/2} + \left(M_s + \frac{M_{bb}m_s}{m_{bb}^2} \frac{4\alpha_s}{3} \mu_{bbs} \right)^2 \right)^{\frac{1}{2}} \\ &= 10317.14 \text{ MeV}. \end{aligned} \quad (55)$$

The predicted masses of the Ω_{bb} baryon are

$$\begin{aligned} \text{State}|J, j\rangle : & \quad |1/2, 1\rangle \quad |3/2, 1\rangle \\ M(\Omega_{bb}, 1S): & \quad 10305.21 \text{ MeV} \quad 10323.11 \text{ MeV}. \end{aligned} \quad (56)$$

According to the analysis of our calculated values, we infer that the mass shifts of about 18 MeV between the $\Omega_{bb}(1/2^+)$ and $\Omega_{bb}(3/2^+)$ states in $1S$ -wave are relatively small due to the large mass

of heavy diquark bb . As indicated by our results in Table IX, the obtained masses of these five $1P$ -wave states with the bases $|J, j\rangle$ are

$$\begin{aligned} \text{State}|J, j\rangle : & \quad |1/2, 0\rangle & \quad |1/2, 1\rangle & \quad |3/2, 1\rangle & \quad |3/2, 2\rangle & \quad |5/2, 2\rangle \\ M(\Omega_{bb}, 1P): & 10797.92 \text{ MeV} & 10807.17 \text{ MeV} & 10808.78 \text{ MeV} & 10817.13 \text{ MeV} & 10825.61 \text{ MeV}. \end{aligned} \quad (57)$$

As can be seen from the above (57), the $\Omega_{bb}(10807.17)$ and $\Omega_{bb}(10808.78)$ states are the mass degenerate mixed states in $1P$ -wave with $J^P = 1/2^-$ and $3/2^-$, respectively. Though they are most likely appears consistent with a single resonance due to its degeneracy, is actually composed of this two states. For the excited states for the Ω_{bb} baryon, the results of the predicted masses are listed in Table IX and compared with other models, and refer to Refs. [69, 70].

VIII. CONCLUSION

In this paper, we study the effective masses of the heavy diquark and light quark in heavy-light hadron framework. We obtain the effective masses of quarks, which are different from the choices of the radial and orbital excited states with L and n for the doubly heavy baryons.

We expand the Regge trajectory formula to complete the spin-average masses \bar{M} of doubly heavy Ξ_{QQ} , Ω_{QQ} baryons. To obtain the mass shifts, we exploit a new scaling relation to calculate the spin-coupling parameters a_1 , a_2 , b_1 , and c_1 . Therefore, we obtain the mass spectra of the radial and orbital excited states for the doubly heavy baryons. In particular, for the ground states of the Ξ_{QQ} baryon, our prediction results are consistent with many theoretical models. Finally, we have also predicted the unobserved states of doubly heavy Ξ_{QQ} and Ω_{QQ} baryons, which have a good reference to be detected in the experiment.

Appendix A: S -wave

For the S -wave system, we consider the spin-dependent Hamiltonian $H^{SD}(L=0)$ in Eq. (22). The matrix elements of $\mathbf{S}_{QQ} \cdot \mathbf{S}_q$ may be evaluated by the square of the total spin $\mathbf{S} = \mathbf{S}_{QQ} + \mathbf{S}_q$,

$$\mathbf{S}_{QQ} \cdot \mathbf{S}_q = (\mathbf{S}^2 - \mathbf{S}_{QQ}^2 - \mathbf{S}_q^2)/2. \quad (A1)$$

Then, the basis states of L - S coupling with the third component S_3 can be constructed as a linear combination of the $|S_{QQ3}, S_{q3}\rangle$ states. The two basis states are

$$|^2S_{1/2}, S_3 = 1/2\rangle = \sqrt{\frac{2}{3}}|1, -\frac{1}{2}\rangle - \sqrt{\frac{1}{3}}|0, \frac{1}{2}\rangle,$$

TABLE VII: Mass spectra (MeV) of Ω_{cc} baryons are given and compared with different quark models.

State	J^P	Ours	[5]	[41]	[62]	[67]
$1^1 S_{1/2}$	$1/2^+$	3651.79	3778	3715	3650	3736
$1^3 S_{3/2}$	$3/2^+$	3712.37	3872	3772	3810	3837
$2^1 S_{1/2}$	$1/2^+$	4160.96		4118	4028	4078
$2^3 S_{3/2}$	$3/2^+$	4173.89		4142	4085	4110
$3^1 S_{1/2}$	$1/2^+$	4525.12			4317	4320
$3^3 S_{3/2}$	$3/2^+$	4529.18			4345	4334
$4^1 S_{1/2}$	$1/2^+$	4823.86			4570	4514
$4^3 S_{3/2}$	$3/2^+$	4825.61			4586	4521
$5^1 S_{1/2}$	$1/2^+$	5083.96			4801	4676
$5^3 S_{3/2}$	$3/2^+$	5084.86			4811	4680
$1^2 P_{1/2}$	$1/2^-$	3997.28	4208	4087	3964	4011
$1^4 P_{1/2}$	$1/2^-$	4030.60	4271	4081	3972	4014
$1^2 P_{3/2}$	$3/2^-$	4037.65	4252	4107	3948	4004
$1^4 P_{3/2}$	$3/2^-$	4063.19	4325	4114	3981	4007
$1^4 P_{5/2}$	$5/2^-$	4090.66	4303	4134	3935	3998
$2^2 P_{1/2}$	$1/2^-$	4411.31			4241	4253
$2^4 P_{1/2}$	$1/2^-$	4427.16			4248	4256
$2^2 P_{3/2}$	$3/2^-$	4429.28			4228	4248
$2^4 P_{3/2}$	$3/2^-$	4442.56			4234	4251
$2^4 P_{5/2}$	$5/2^-$	4454.78			4216	4244
$3^2 P_{1/2}$	$1/2^-$	4733.62			4492	4453
$3^4 P_{1/2}$	$1/2^-$	4742.75			4498	4454
$3^2 P_{3/2}$	$3/2^-$	4743.62			4479	4450
$3^4 P_{3/2}$	$3/2^-$	4751.59			4486	4451
$3^4 P_{5/2}$	$5/2^-$	4758.39			4469	4447
$4^2 P_{1/2}$	$1/2^-$	5007.43			4723	
$4^4 P_{1/2}$	$1/2^-$	5013.35			4728	
$4^2 P_{3/2}$	$3/2^-$	5013.78			4712	
$4^4 P_{3/2}$	$3/2^-$	5019.06			4717	
$4^4 P_{5/2}$	$5/2^-$	5023.37			4703	
$5^2 P_{1/2}$	$1/2^-$	5250.09			4939	
$5^4 P_{1/2}$	$1/2^-$	5254.23			4944	
$5^2 P_{3/2}$	$3/2^-$	5254.46			4929	
$5^4 P_{3/2}$	$3/2^-$	5258.21			4934	
$5^4 P_{5/2}$	$5/2^-$	5261.18			4921	
$1^4 D_{1/2}$	$1/2^+$	4296.41			4156	4101
$1^2 D_{3/2}$	$3/2^+$	4312.55			4133	4091
$1^4 D_{3/2}$	$3/2^+$	4328.01			4141	4094
$1^2 D_{5/2}$	$5/2^+$	4339.28			4113	4082
$1^4 D_{5/2}$	$5/2^+$	4353.79			4121	4086
$1^4 D_{7/2}$	$7/2^+$	4376.31			4075	4075
$2^4 D_{1/2}$	$1/2^+$	4642.80			4407	4324
$2^2 D_{3/2}$	$3/2^+$	4652.02			4389	4318
$2^4 D_{3/2}$	$3/2^+$	4661.02			4395	4320
$2^2 D_{5/2}$	$5/2^+$	4667.32			4372	4312
$2^4 D_{5/2}$	$5/2^+$	4675.80			4378	4314
$2^4 D_{7/2}$	$7/2^+$	4688.57			4358	4307
$3^4 D_{1/2}$	$1/2^+$	4930.50			4446	
$3^2 D_{3/2}$	$3/2^+$	4936.44			4425	
$3^4 D_{3/2}$	$3/2^+$	4942.30			4432	
$3^2 D_{5/2}$	$5/2^+$	4946.30			4407	
$3^4 D_{5/2}$	$5/2^+$	4951.84			4414	
$3^4 D_{7/2}$	$7/2^+$	4960.03			4391	
$4^4 D_{1/2}$	$1/2^+$	5182.30			4863	
$4^2 D_{3/2}$	$3/2^+$	5186.43			4847	
$4^4 D_{3/2}$	$3/2^+$	5190.55			4853	
$4^2 D_{5/2}$	$5/2^+$	5193.31			4833	
$4^4 D_{5/2}$	$5/2^+$	5197.20			4838	
$4^4 D_{7/2}$	$7/2^+$	5202.89			4821	
$5^4 D_{1/2}$	$1/2^+$	5409.24				
$5^2 D_{3/2}$	$3/2^+$	5412.28				
$5^4 D_{3/2}$	$3/2^+$	5415.33				
$5^2 D_{5/2}$	$5/2^+$	5417.35				
$5^4 D_{5/2}$	$5/2^+$	5420.23				
$5^4 D_{7/2}$	$7/2^+$	5424.41				

TABLE VIII: Mass spectra (MeV) of Ω_{bc} baryons are given and compared with different quark models.

State	J^P	Ours	[5]	[62]	[67]	[44]
$1^1 S_{1/2}$	$1/2^+$	6767.50	7088	7136	7079	6994
$1^3 S_{3/2}$	$3/2^+$	6796.33	7130	7187	7182	7017
$2^1 S_{1/2}$	$1/2^+$	7359.24		7473	7435	7559
$2^3 S_{3/2}$	$3/2^+$	7365.52		7490	7469	7571
$3^1 S_{1/2}$	$1/2^+$	7791.65		7753	7688	7976
$3^3 S_{3/2}$	$3/2^+$	7793.62		7761	7703	7985
$4^1 S_{1/2}$	$1/2^+$	8148.51		8004	7895	
$4^3 S_{3/2}$	$3/2^+$	8149.36		8009	7903	
$5^1 S_{1/2}$	$1/2^+$	8459.70		8236	8073	
$5^3 S_{3/2}$	$3/2^+$	8460.14		8239	8077	
$1^2 P_{1/2}$	$1/2^-$	7199.88		7375	7369	
$1^4 P_{1/2}$	$1/2^-$	7214.67		7381	7371	
$1^2 P_{3/2}$	$3/2^-$	7217.25		7363	7364	
$1^4 P_{3/2}$	$3/2^-$	7230.59		7369	7366	
$1^4 P_{5/2}$	$5/2^-$	7244.16		7353	7360	
$2^2 P_{1/2}$	$1/2^-$	7673.13		7657	7620	
$2^4 P_{1/2}$	$1/2^-$	7680.41		7662	7621	
$2^2 P_{3/2}$	$3/2^-$	7681.16		7647	7616	
$2^4 P_{3/2}$	$3/2^-$	7687.92		7652	7618	
$2^4 P_{5/2}$	$5/2^-$	7693.93		7639	7614	
$3^2 P_{1/2}$	$1/2^-$	8050.77		7912	7832	
$3^4 P_{1/2}$	$1/2^-$	8055.04		7916	7833	
$3^2 P_{3/2}$	$3/2^-$	8055.34		7903	7830	
$3^4 P_{3/2}$	$3/2^-$	8059.34		7908	7831	
$3^4 P_{5/2}$	$5/2^-$	8062.68		7896	7828	
$4^2 P_{1/2}$	$1/2^-$	8374.79		8147		
$4^4 P_{1/2}$	$1/2^-$	8377.57		8151		
$4^2 P_{3/2}$	$3/2^-$	8377.72		8140		
$4^4 P_{3/2}$	$3/2^-$	8380.35		8143		
$4^4 P_{5/2}$	$5/2^-$	8382.46		8133		
$5^2 P_{1/2}$	$1/2^-$	8663.26		8368		
$5^4 P_{1/2}$	$1/2^-$	8665.22		8372		
$5^2 P_{3/2}$	$3/2^-$	8665.30		8361		
$5^4 P_{3/2}$	$3/2^-$	8667.16		8365		
$5^4 P_{5/2}$	$5/2^-$	8668.61		8355		
$1^4 D_{1/2}$	$1/2^+$	7549.77		7562	7464	
$1^2 D_{3/2}$	$3/2^+$	7557.54		7545	7458	
$1^4 D_{3/2}$	$3/2^+$	7565.04		7551	7460	
$1^2 D_{5/2}$	$5/2^+$	7570.48		7531	7452	
$1^4 D_{5/2}$	$5/2^+$	7577.57		7536	7454	
$1^4 D_{7/2}$	$7/2^+$	7588.54		7518	7447	
$2^4 D_{1/2}$	$1/2^+$	7950.30		7821	7700	
$2^2 D_{3/2}$	$3/2^+$	7954.76		7807	7695	
$2^4 D_{3/2}$	$3/2^+$	7959.13		7812	7697	
$2^2 D_{5/2}$	$5/2^+$	7962.17		7795	7691	
$2^4 D_{5/2}$	$5/2^+$	7966.31		7799	7692	
$2^4 D_{7/2}$	$7/2^+$	7972.53		7784	7687	
$3^4 D_{1/2}$	$1/2^+$	8288.05		8060		
$3^2 D_{3/2}$	$3/2^+$	8290.92		8048		
$3^4 D_{3/2}$	$3/2^+$	8293.77		8052		
$3^2 D_{5/2}$	$5/2^+$	8295.70		8037		
$3^4 D_{5/2}$	$5/2^+$	8298.41		8041		
$3^4 D_{7/2}$	$7/2^+$	8302.39		8028		
$4^4 D_{1/2}$	$1/2^+$	8585.86		8085		
$4^2 D_{3/2}$	$3/2^+$	8587.86		8274		
$4^4 D_{3/2}$	$3/2^+$	8589.86		8277		
$4^2 D_{5/2}$	$5/2^+$	8591.20		8263		
$4^4 D_{5/2}$	$5/2^+$	8593.10		8267		
$4^4 D_{7/2}$	$7/2^+$	8595.87		8254		
$5^4 D_{1/2}$	$1/2^+$	8855.37				
$5^2 D_{3/2}$	$3/2^+$	8856.85				
$5^4 D_{3/2}$	$3/2^+$	8858.33				
$5^2 D_{5/2}$	$5/2^+$	8859.31				
$5^4 D_{5/2}$	$5/2^+$	8860.71				
$5^4 D_{7/2}$	$7/2^+$	8862.74				

TABLE IX: Mass spectra (MeV) of Ω_{bb} baryons are given and compared with different quark models.

State	J^P	Ours	[5]	[41]	[30]	[62]
$1^1 S_{1/2}$	$1/2^+$	10305.21	10359	10230	10266	10446
$1^3 S_{3/2}$	$3/2^+$	10323.11	10389	10258	10291	10467
$2^1 S_{1/2}$	$1/2^+$	10967.74	10970	10751	10816	10730
$2^3 S_{3/2}$	$3/2^+$	10969.66	10992	10763	10830	10737
$3^1 S_{1/2}$	$1/2^+$	11450.73				10973
$3^3 S_{3/2}$	$3/2^+$	11451.97				10976
$4^1 S_{1/2}$	$1/2^+$	11851.70				11191
$4^3 S_{3/2}$	$3/2^+$	11852.23				11193
$5^1 S_{1/2}$	$1/2^+$	12201.51				11393
$5^3 S_{3/2}$	$3/2^+$	12201.78				11394
$1^2 P_{1/2}$	$1/2^-$	10797.92	10771	10605	10669	10580
$1^4 P_{1/2}$	$1/2^-$	10807.17	10804	10591	10641	10581
$1^2 P_{3/2}$	$3/2^-$	10808.78	10785	10610	10681	10578
$1^4 P_{3/2}$	$3/2^-$	10817.13	10821	10611	10656	10579
$1^4 P_{5/2}$	$5/2^-$	10825.61	10798	10625	10655	10576
$2^2 P_{1/2}$	$1/2^-$	11323.18				10796
$2^4 P_{1/2}$	$1/2^-$	11327.74				10797
$2^2 P_{3/2}$	$3/2^-$	11328.21				10795
$2^4 P_{3/2}$	$3/2^-$	11332.44				10796
$2^4 P_{5/2}$	$5/2^-$	11336.20				10794
$3^2 P_{1/2}$	$1/2^-$	11745.18				10982
$3^4 P_{1/2}$	$1/2^-$	11747.85				10982
$3^2 P_{3/2}$	$3/2^-$	11748.04				10981
$3^4 P_{3/2}$	$3/2^-$	11750.54				10981
$3^4 P_{5/2}$	$5/2^-$	11752.63				10980
$4^2 P_{1/2}$	$1/2^-$	12108.25				
$4^4 P_{1/2}$	$1/2^-$	12109.99				
$4^2 P_{3/2}$	$3/2^-$	12110.08				
$4^4 P_{3/2}$	$3/2^-$	12111.73				
$4^4 P_{5/2}$	$5/2^-$	12113.05				
$5^2 P_{1/2}$	$1/2^-$	12431.92				
$5^4 P_{1/2}$	$1/2^-$	12433.14				
$5^2 P_{3/2}$	$3/2^-$	12433.19				
$5^4 P_{3/2}$	$3/2^-$	12434.36				
$5^4 P_{5/2}$	$5/2^-$	12435.27				
$1^4 D_{1/2}$	$1/2^+$	11189.58			10971	10662
$1^2 D_{3/2}$	$3/2^+$	11194.45			10975	10660
$1^4 D_{3/2}$	$3/2^+$	11199.14			10891	10661
$1^2 D_{5/2}$	$5/2^+$	11202.54			10979	10657
$1^4 D_{5/2}$	$5/2^+$	11206.98			10896	10658
$1^4 D_{7/2}$	$7/2^+$	11213.84			10898	10655
$2^4 D_{1/2}$	$1/2^+$	11635.15				10866
$2^2 D_{3/2}$	$3/2^+$	11637.93				10864
$2^4 D_{3/2}$	$3/2^+$	11640.67				10865
$2^2 D_{5/2}$	$5/2^+$	11642.57				10863
$2^4 D_{5/2}$	$5/2^+$	11645.16				10863
$2^4 D_{7/2}$	$7/2^+$	11649.05				10861
$3^4 D_{1/2}$	$1/2^+$	12012.62				
$3^2 D_{3/2}$	$3/2^+$	12014.41				
$3^4 D_{3/2}$	$3/2^+$	12016.20				
$3^2 D_{5/2}$	$5/2^+$	12017.41				
$3^4 D_{5/2}$	$5/2^+$	12019.10				
$3^4 D_{7/2}$	$7/2^+$	12021.59				
$4^4 D_{1/2}$	$1/2^+$	12346.21				
$4^2 D_{3/2}$	$3/2^+$	12347.47				
$4^4 D_{3/2}$	$3/2^+$	12348.72				
$4^2 D_{5/2}$	$5/2^+$	12349.55				
$4^4 D_{5/2}$	$5/2^+$	12350.74				
$4^4 D_{7/2}$	$7/2^+$	12352.47				
$5^4 D_{1/2}$	$1/2^+$	12648.49				
$5^2 D_{3/2}$	$3/2^+$	12649.41				
$5^4 D_{3/2}$	$3/2^+$	12650.34				
$5^2 D_{5/2}$	$5/2^+$	12650.95				
$5^4 D_{5/2}$	$5/2^+$	12651.83				
$5^4 D_{7/2}$	$7/2^+$	12653.10				

$$|^4S_{3/2}, S_3 = 3/2\rangle = |1, \frac{1}{2}\rangle. \quad (\text{A2})$$

The eigenvalues (two diagonal elements) of $\langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle$ in the basis $[^2S_{1/2}, ^4S_{3/2}]$ can be given by

$$\begin{aligned} \langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle &= \begin{bmatrix} \langle ^2S_{1/2}, S_3 = 1/2 | \mathbf{S}_{QQ} \cdot \mathbf{S}_q | ^2S_{1/2}, S_3 = 1/2 \rangle & \langle ^2S_{1/2}, S_3 = 1/2 | \mathbf{S}_{QQ} \cdot \mathbf{S}_q | ^4S_{3/2}, S_3 = 3/2 \rangle \\ \langle ^4S_{3/2}, S_3 = 3/2 | \mathbf{S}_{QQ} \cdot \mathbf{S}_q | ^2S_{1/2}, S_3 = 1/2 \rangle & \langle ^4S_{3/2}, S_3 = 3/2 | \mathbf{S}_{QQ} \cdot \mathbf{S}_q | ^4S_{3/2}, S_3 = 3/2 \rangle \end{bmatrix} \\ &= \begin{bmatrix} -1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}. \end{aligned} \quad (\text{A3})$$

Appendix B: P -wave

For the P -wave states of the doubly heavy baryons, the expectation value of $\mathbf{L} \cdot \mathbf{S}$ in any coupling scheme is

$$\langle \mathbf{L} \cdot \mathbf{S} \rangle = [J(J+1) - L(L+1) - S(S+1)]/2. \quad (\text{B1})$$

The calculation of the operator $\mathbf{L} \cdot \mathbf{S}_i$ ($i = QQ, q$) with raising and lowering operator $L_{\pm}, S_{i\pm}$ results in

$$\mathbf{L} \cdot \mathbf{S}_i = L_3 S_{i3} + (L_+ S_{i-} + L_- S_{i+})/2. \quad (\text{B2})$$

In the L - S basis can be constructed as linear combinations of the states $|S_{QQ3}, S_{q3}, L_3\rangle$ of the third components of the respective angular momenta. Thus, these five P -wave states of the doubly heavy baryons may be classified as $^{2S+1}P_J = ^2P_{1/2}, ^4P_{1/2}, ^2P_{3/2}, ^4P_{3/2}, ^4P_{5/2}$ with the third components of the total angular momenta J_3 ,

$$\begin{aligned} |^2P_{1/2}, J_3 = 1/2\rangle &= \frac{\sqrt{2}}{3}|1, -\frac{1}{2}, 0\rangle - \frac{1}{3}|0, \frac{1}{2}, 0\rangle - \frac{\sqrt{2}}{3}|0, -\frac{1}{2}, 1\rangle + \frac{2}{3}|-1, \frac{1}{2}, 1\rangle, \\ |^4P_{1/2}, J_3 = 1/2\rangle &= \frac{1}{\sqrt{2}}|1, \frac{1}{2}, -1\rangle - \frac{1}{3}|1, -\frac{1}{2}, 0\rangle - \frac{\sqrt{2}}{3}|0, \frac{1}{2}, 0\rangle + \frac{1}{3}|0, -\frac{1}{2}, 1\rangle + \frac{1}{3\sqrt{2}}|-1, \frac{1}{2}, 1\rangle, \\ |^2P_{3/2}, J_3 = 3/2\rangle &= \sqrt{\frac{2}{3}}|1, -\frac{1}{2}, 1\rangle - \sqrt{\frac{1}{3}}|0, \frac{1}{2}, 1\rangle, \\ |^4P_{3/2}, J_3 = 3/2\rangle &= \sqrt{\frac{3}{5}}|1, \frac{1}{2}, 0\rangle - \sqrt{\frac{2}{15}}|1, -\frac{1}{2}, 1\rangle - \frac{2}{\sqrt{15}}|0, \frac{1}{2}, 1\rangle, \\ |^4P_{5/2}, J_3 = 5/2\rangle &= |1, \frac{1}{2}, 1\rangle. \end{aligned} \quad (\text{B3})$$

The expectation values of $\langle \mathbf{L} \cdot \mathbf{S}_i \rangle$, $\langle S_{12} \rangle$ and $\langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle$ are given by

$$\begin{aligned} \langle \mathbf{L} \cdot \mathbf{S}_{QQ} \rangle_{J=\frac{1}{2}} &= \begin{bmatrix} -\frac{4}{3} & -\frac{\sqrt{2}}{3} \\ -\frac{\sqrt{2}}{3} & -\frac{5}{3} \end{bmatrix}, \langle \mathbf{L} \cdot \mathbf{S}_q \rangle_{J=\frac{1}{2}} = \begin{bmatrix} \frac{1}{3} & \frac{\sqrt{2}}{3} \\ \frac{\sqrt{2}}{3} & -\frac{5}{6} \end{bmatrix}, \langle S_{12} \rangle_{J=\frac{1}{2}} = \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -1 \end{bmatrix}, \\ \langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle_{J=\frac{1}{2}} &= \begin{bmatrix} -1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}, \end{aligned}$$

$$\begin{aligned}
\langle \mathbf{L} \cdot \mathbf{S}_{QQ} \rangle_{J=\frac{3}{2}} &= \begin{bmatrix} \frac{2}{3} & -\frac{\sqrt{5}}{3} \\ -\frac{\sqrt{5}}{3} & -\frac{2}{3} \end{bmatrix}, \langle \mathbf{L} \cdot \mathbf{S}_q \rangle_{J=\frac{3}{2}} = \begin{bmatrix} -\frac{1}{3} & \frac{\sqrt{5}}{3} \\ \frac{\sqrt{5}}{3} & -\frac{1}{3} \end{bmatrix}, \langle S_{12} \rangle_{J=\frac{3}{2}} = \begin{bmatrix} 0 & -\frac{\sqrt{5}}{10} \\ -\frac{\sqrt{5}}{10} & \frac{4}{5} \end{bmatrix}, \\
\langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle_{J=\frac{3}{2}} &= \begin{bmatrix} -1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}, \\
\langle \mathbf{L} \cdot \mathbf{S}_{QQ} \rangle_{J=\frac{5}{2}} &= 1, \quad \langle \mathbf{L} \cdot \mathbf{S}_q \rangle_{J=\frac{5}{2}} = \frac{1}{2}, \quad \langle S_{12} \rangle_{J=\frac{5}{2}} = -\frac{1}{5}, \quad \langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle_{J=\frac{5}{2}} = \frac{1}{2}.
\end{aligned} \tag{B4}$$

The matrix forms of these mass shifts are

$$\begin{aligned}
\Delta \mathcal{M}_{J=1/2} &= \begin{bmatrix} \frac{1}{3}(a_2 - 4a_1) - c_1 & \frac{\sqrt{2}}{3}(a_2 - a_1) + \frac{b_1}{\sqrt{2}} \\ \frac{\sqrt{2}}{3}(a_2 - a_1) + \frac{b_1}{\sqrt{2}} & -\frac{5}{3}(a_1 + \frac{1}{2}a_2) - b_1 + \frac{1}{2}c_1 \end{bmatrix}, \\
\Delta \mathcal{M}_{J=3/2} &= \begin{bmatrix} \frac{2}{3}a_1 - \frac{1}{6}a_2 - c_1 & \frac{\sqrt{5}}{3}(a_2 - a_1) - \frac{b_1}{2\sqrt{5}} \\ \frac{\sqrt{5}}{3}(a_2 - a_1) - \frac{b_1}{2\sqrt{5}} & -\frac{1}{3}(2a_1 + a_2) + \frac{4b_1}{5} + \frac{1}{2}c_1 \end{bmatrix}, \\
\Delta \mathcal{M}_{J=5/2} &= a_1 + \frac{1}{2}a_2 - \frac{b_1}{5} + \frac{c_1}{2}.
\end{aligned} \tag{B5}$$

Diagonalizing the above matrices Eq. (B5), we can obtain the mass shifts $\Delta M(J, j)$ with the $j = 0, 1, 2$,

$$\begin{aligned}
\Delta M(1/2, 0) &= \frac{1}{4}(-a_1 - 6a_2 - 2b_1 - c_1) \\
&\quad - \frac{1}{12}\sqrt{(9a_1 - 2a_2 + 10b_1 - 7c_1)^2 + 8(2a_2 - b_1 - 2c_1)^2}, \\
\Delta M(1/2, 1) &= \frac{1}{4}(-a_1 - 6a_2 - 2b_1 - c_1) \\
&\quad + \frac{1}{12}\sqrt{(9a_1 - 2a_2 + 10b_1 - 7c_1)^2 + 8(2a_2 - b_1 - 2c_1)^2}, \\
\Delta M(3/2, 1) &= \frac{1}{20}(-5a_1 + 8b_1 - 5c_1) \\
&\quad - \frac{1}{60}\sqrt{(45a_1 - 40a_2 - 16b_1 - 5c_1)^2 + 5(20a_2 - 10b_1 - 20c_1)^2}, \\
\Delta M(3/2, 2) &= \frac{1}{20}(-5a_1 + 8b_1 - 5c_1) \\
&\quad + \frac{1}{60}\sqrt{(45a_1 - 40a_2 - 16b_1 - 5c_1)^2 + 5(20a_2 - 10b_1 - 20c_1)^2}, \\
\Delta M(5/2, 2) &= \frac{a_1}{2} + a_2 - \frac{b_1}{5} + \frac{c_1}{2}.
\end{aligned} \tag{B6}$$

Appendix C: *D*-wave

For analyzing the *D*-wave system, the relevant linear combinations of six basis states ${}^{2S+1}D_J = {}^4D_{1/2}, {}^2D_{3/2}, {}^4D_{3/2}, {}^2D_{5/2}, {}^4D_{5/2}, {}^4D_{7/2}$ are given by

$$\begin{aligned}
|{}^4D_{1/2}, J_3 = 1/2\rangle &= \frac{1}{\sqrt{10}}|1, \frac{1}{2}, -1\rangle - \frac{1}{\sqrt{15}}|1, -\frac{1}{2}, 0\rangle - \sqrt{\frac{2}{15}}|0, \frac{1}{2}, 0\rangle + \frac{1}{\sqrt{5}}|0, -\frac{1}{2}, 1\rangle + \frac{1}{\sqrt{10}}|-1, \frac{1}{2}, 1\rangle \\
&\quad - \sqrt{\frac{2}{5}}|-1, -\frac{1}{2}, 2\rangle,
\end{aligned}$$

$$\begin{aligned}
|^2D_{3/2}, J_3 = 3/2\rangle &= \sqrt{\frac{2}{15}}|1, -\frac{1}{2}, 1\rangle - \frac{1}{\sqrt{15}}|0, \frac{1}{2}, 1\rangle - \frac{2}{\sqrt{15}}|0, -\frac{1}{2}, 2\rangle + \sqrt{\frac{8}{15}}|-1, \frac{1}{2}, 2\rangle, \\
|^4D_{3/2}, J_3 = 3/2\rangle &= \frac{1}{\sqrt{5}}|1, \frac{1}{2}, 0\rangle - \sqrt{\frac{2}{15}}|1, \frac{1}{2}, 1\rangle - \frac{2}{\sqrt{15}}|0, \frac{1}{2}, 1\rangle + \frac{2}{\sqrt{15}}|0, -\frac{1}{2}, 2\rangle + \sqrt{\frac{2}{15}}|-1, \frac{1}{2}, 2\rangle, \\
|^2D_{5/2}, J_3 = 5/2\rangle &= \sqrt{\frac{2}{3}}|1, -\frac{1}{2}, 2\rangle - \sqrt{\frac{1}{3}}|0, \frac{1}{2}, 2\rangle, \\
|^4D_{5/2}, J_3 = 5/2\rangle &= \frac{3}{\sqrt{21}}|1, \frac{1}{2}, 1\rangle - \frac{2}{\sqrt{21}}|1, -\frac{1}{2}, 2\rangle - \frac{2\sqrt{2}}{\sqrt{21}}|0, \frac{1}{2}, 2\rangle, \\
|^4D_{7/2}, J_3 = 7/2\rangle &= |1, \frac{1}{2}, 2\rangle.
\end{aligned} \tag{C1}$$

The expectation values of $\langle \mathbf{L} \cdot \mathbf{S}_i \rangle$ ($i = QQ, q$), $\langle S_{12} \rangle$ and $\langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle$ are

$$\begin{aligned}
\langle \mathbf{L} \cdot \mathbf{S}_{QQ} \rangle_{J=\frac{1}{2}} &= -3, \quad \langle \mathbf{L} \cdot \mathbf{S}_q \rangle_{J=\frac{1}{2}} = -\frac{3}{2}, \quad \langle S_{12} \rangle_{J=\frac{1}{2}} = -1, \quad \langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle_{J=\frac{1}{2}} = \frac{1}{2}, \\
\langle \mathbf{L} \cdot \mathbf{S}_{QQ} \rangle_{J=\frac{3}{2}} &= \begin{bmatrix} -2 & -1 \\ -1 & -2 \end{bmatrix}, \quad \langle \mathbf{L} \cdot \mathbf{S}_q \rangle_{J=\frac{3}{2}} = \begin{bmatrix} \frac{1}{2} & 1 \\ 1 & -1 \end{bmatrix}, \quad \langle S_{12} \rangle_{J=\frac{3}{2}} = \begin{bmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{bmatrix}, \\
\langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle_{J=\frac{3}{2}} &= \begin{bmatrix} -1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}, \\
\langle \mathbf{L} \cdot \mathbf{S}_{QQ} \rangle_{J=\frac{5}{2}} &= \begin{bmatrix} \frac{4}{3} & -\frac{\sqrt{14}}{3} \\ -\frac{\sqrt{14}}{3} & -\frac{1}{3} \end{bmatrix}, \quad \langle \mathbf{L} \cdot \mathbf{S}_q \rangle_{J=\frac{5}{2}} = \begin{bmatrix} -\frac{1}{3} & \frac{\sqrt{14}}{3} \\ \frac{\sqrt{14}}{3} & -\frac{1}{6} \end{bmatrix}, \quad \langle S_{12} \rangle_{J=\frac{5}{2}} = \begin{bmatrix} 0 & -\frac{\sqrt{14}}{14} \\ -\frac{\sqrt{14}}{14} & \frac{5}{7} \end{bmatrix}, \\
\langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle_{J=\frac{5}{2}} &= \begin{bmatrix} -1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}, \\
\langle \mathbf{L} \cdot \mathbf{S}_{QQ} \rangle_{J=\frac{7}{2}} &= 2, \quad \langle \mathbf{L} \cdot \mathbf{S}_q \rangle_{J=\frac{7}{2}} = 1, \quad \langle S_{12} \rangle_{J=\frac{7}{2}} = -\frac{2}{7}, \quad \langle \mathbf{S}_{QQ} \cdot \mathbf{S}_q \rangle_{J=\frac{7}{2}} = \frac{1}{2}.
\end{aligned} \tag{C2}$$

The matrix forms of these mass shifts are

$$\begin{aligned}
\Delta \mathcal{M}_{J=1/2} &= -3a_1 - \frac{3a_2}{2} - b_1 + \frac{c_1}{2}, \\
\Delta \mathcal{M}_{J=3/2} &= \begin{bmatrix} -2a_1 + \frac{1}{2}a_2 - c_1 & -a_1 + a_2 + \frac{1}{2}b_1 \\ -a_1 + a_2 + \frac{1}{2}b_1 & -2a_1 - a_2 + \frac{1}{2}c_1 \end{bmatrix}, \\
\Delta \mathcal{M}_{J=5/2} &= \begin{bmatrix} \frac{4}{3}a_1 - \frac{1}{3}a_2 - c_1 & -\frac{\sqrt{14}}{3}a_1 + \frac{\sqrt{14}}{3}a_2 - \frac{\sqrt{14}}{14}b_1 \\ -\frac{\sqrt{14}}{3}a_1 + \frac{\sqrt{14}}{3}a_2 - \frac{\sqrt{14}}{14}b_1 & -\frac{1}{3}a_1 - \frac{1}{6}a_2 + \frac{5}{7}b_1 + \frac{1}{2}c_1 \end{bmatrix}, \\
\Delta \mathcal{M}_{J=7/2} &= 2a_1 + a_2 - \frac{2}{7}b_1 + \frac{1}{2}c_1.
\end{aligned} \tag{C3}$$

Diagonalizing the above matrices Eq. (C3), we can obtain the six mass shifts $\Delta M(J, j)$ with $j = 1, 2, 3$,

$$\begin{aligned}
\Delta M(1/2, 1) &= -\frac{3a_1}{2} - 3a_2 - b_1 + \frac{c_1}{2}, \\
\Delta M(3/2, 1) &= \frac{1}{4}(-a_1 - 8a_2 - c_1) \\
&\quad - \frac{1}{20}\sqrt{(25a_1 - 16a_2 + 8b_1 - 9c_1)^2 + 36(2a_2 - b_1 - 2c_1)^2}, \\
\Delta M(3/2, 2) &= \frac{1}{4}(-a_1 - 8a_2 - c_1)
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{20} \sqrt{(25a_1 - 16a_2 + 8b_1 - 9c_1)^2 + 36(2a_2 - b_1 - 2c_1)^2}, \\
\Delta M(5/2, 2) &= \frac{1}{28} (-7a_1 + 14a_2 + 10b_1 - 7c_1) \\
& - \frac{1}{140} \sqrt{(175a_1 - 182a_2 - 342b_1 + 7c_1)^2 + 2744(2a_2 - b_1 - 2c_1)^2}, \\
\Delta M(5/2, 3) &= \frac{1}{28} (-7a_1 + 14a_2 + 10b_1 - 7c_1) \\
& + \frac{1}{140} \sqrt{(175a_1 - 182a_2 - 342b_1 + 7c_1)^2 + 2744(2a_2 - b_1 - 2c_1)^2}, \\
\Delta M(7/2, 3) &= a_1 + 2a_2 - \frac{2}{7}b_1 + \frac{c_1}{2}. \tag{C4}
\end{aligned}$$

-
- [1] R. Aaij *et al.* (LHCb Collaboration), Observation of the doubly charmed baryon Ξ_{cc}^{++} , *Phys. Rev. Lett.* **119**, 112001 (2017). [arXiv:1707.01621 [hep-ex]].
- [2] R. Aaij *et al.* (LHCb Collaboration), First observation of the doubly charmed baryon decay $\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$, *Phys. Rev. Lett.* **121**, 162002 (2018). [arXiv:1807.01919 [hep-ex]].
- [3] R. Aaij *et al.* (LHCb Collaboration), Precision measurement of the Ξ_{cc}^{++} mass, *JHEP*, **2002**, 049 (2020). [arXiv:1911.08594 [hep-ex]].
- [4] M. Karliner, J. L. Rosner, Baryons with two heavy quarks: masses, production, decays, and detection, *Phys. Rev. D* **90**, 094007 (2014). [arXiv:1408.5877 [hep-ph]].
- [5] D. Ebert, R. N. Faustov, V. O. Galkin, A. P. Martynenko, Mass spectra of doubly heavy baryons in the relativistic quark model, *Phys. Rev. D* **66**, 014008 (2002). [arXiv:0201217 [hep-ph]].
- [6] Z. Shah, A. K. Rai, Excited state mass spectra of doubly heavy Ξ baryons, *Eur. Phys. J. C* **77**, 129 (2017). [arXiv:1702.02726 [hep-ph]].
- [7] N. Mohajery, N. Salehi, H. Hassanabadi, A new model for calculating the ground and excited states masses spectra of doubly heavy Ξ baryons, *Adv. High. Energy. Phys* **2018**, 1326438 (2018) [arXiv:1807.06800 [nucl-th]].
- [8] E. Ortiz-Pacheco, R. Bijker, Masses and radiative decay widths of S - and P -wave singly, doubly, and triply heavy charm and bottom baryons, *Phys. Rev. D* **108**, 054014 (2023). [arXiv:2307.04939 [hep-ph]].
- [9] J. Soto, J. T. Castellà, Effective QCD string and doubly heavy baryons, *Phys. Rev. D* **104**, 074027 (2021). [arXiv:2108.00496 [hep-ph]].
- [10] J-R. Zhang, M-Q. Huang, Doubly heavy baryons in QCD sum rules, *Phys. Rev. D* **78**, 094007 (2008). [arXiv:0810.5396 [hep-ph]].
- [11] M. Mattson *et al.* (SELEX Collaboration), First observation of the doubly charmed baryon Ξ_{cc}^+ , *Phys. Rev. Lett.* **89**, 112001 (2002). [arXiv:0208014 [hep-ex]].
- [12] A. Ocherashvili *et al.* (SELEX Collaboration), Confirmation of the doubly charmed baryon $\Xi_{cc}^+(3520)$ via its decay to pD^+K^- , *Phys. Lett. B* **628**, 18 (2005). [arXiv:0406033 [hep-ex]].

- [13] R. Aaij *et al.* (LHCb Collaboration), Search for the doubly charmed baryon Ξ_{cc}^+ , *Sci. China-Phys. Mech. Astron* **63**, 221062 (2020). [[arXiv:1909.12273 \[hep-ex\]](#)].
- [14] D. J. Griffiths, Introduction to elementary particles. (John Wiley & Sons, Inc., Canada, 1987).
- [15] F. E. Close, An introduction to quarks and partons, Second Printing. (Academic Press, Inc., London, 1982).
- [16] D. LaCourse and M. G. Olsson, String potential model: spinless quarks, *Phys. Rev. D* **39**, 2751 (1989).
- [17] B. Chen, K-W Wei, and A. Zhang, Investigation of Λ_Q and Ξ_Q baryons in the heavy quark-light diquark picture, *Eur. Phys. J. A* **51**, 82 (2015). [[arXiv:1406.6561 \[hep-ph\]](#)].
- [18] D. Ebert, R. N. Faustov, V. O. Galkin, Spectroscopy and Regge trajectories of heavy quarkonia and B_c mesons, *Eur. Phys. J. C* **71**, 1825 (2011). [[arXiv:1111.0454 \[hep-ph\]](#)].
- [19] J-H. Pan, J. Pan, Investigation of the mass spectra of singly heavy baryons Σ_Q , Ξ'_Q , and Ω_Q ($Q = c, b$) in the Regge trajectory model, *Phys. Rev. D* **109**, 076010 (2024). [[arXiv:2308.11769 \[hep-ph\]](#)].
- [20] D. Ebert, R. N. Faustov, and V. O. Galkin, Spectroscopy and Regge trajectories of heavy baryons in the relativistic quark-diquark picture, *Phys. Rev. D* **84**, 014025 (2011). [[arXiv:1105.0583 \[hep-ph\]](#)].
- [21] M. Karliner and J. L. Rosner, Prospects for observing the lowest-lying odd-parity Σ_c and Σ_b baryons, *Phys. Rev. D* **92**, 074026 (2015). [[arXiv:1506.01702 \[hep-ph\]](#)].
- [22] L. D. Landau and E. M. Lifshitz, Quantum Mechanics (Nonrelativistic Theory), Pergamon Press, Oxford (1977).
- [23] M. Karliner and J. L. Rosner, Very narrow excited Ω_c baryons, *Phys. Rev. D* **95**, 114012 (2017). [[arXiv:1703.07774 \[hep-ph\]](#)].
- [24] A. Ali, I. Ahmed, M. J. Aslam, A. Ya. Parkhomenko, and A. Rehman, Mass spectrum of the hidden-charm pentaquarks in the compact diquark model, *JHEP*, **2019**, 256 (2019). [[arXiv:1907.06507 \[hep-ph\]](#)].
- [25] D. Ebert, R. N. Faustov, V. O. Galkin, A. P. Martynenko, Mass spectra of doubly heavy baryons in the relativistic quark model, *Phys. Rev. D* **66**, 014008 (2002). [[arXiv:0201217 \[hep-ph\]](#)].
- [26] S. Navas *et al.* (Particle Data Group), Review of Particle Physics, *Phys. Rev. D* **110**, 030001 (2024).
- [27] R. Aaij *et al.* (LHCb Collaboration), Search for the doubly heavy Ξ_{bc}^0 baryon via decays to $D^0 p K^-$, *JHEP*, **2020**, 095 (2020). [[arXiv:2009.02481 \[hep-ex\]](#)].
- [28] R. Aaij *et al.* (LHCb Collaboration), Search for the doubly heavy baryons Ω_{bc}^0 and Ξ_{bc}^0 decaying to $\Lambda_c^+ \pi^-$ and $\Xi_c^+ \pi^-$, *Chin. Phys. C* **45**, 093002 (2021). [[arXiv:2104.04759 \[hep-ex\]](#)].
- [29] R. Aaij *et al.* (LHCb Collaboration), Search for the doubly heavy baryon Ξ_{bc}^+ decaying to $J/\psi \Xi_c^+$. *Chin. Phys. C* **47**, 093001 (2023). [[arXiv:2204.09541 \[hep-ex\]](#)].
- [30] B. Chen, S-Q. Luo, K-W Wei, X. Liu, b -hadron spectroscopy study based on the similarity of double bottom baryon and bottom meson, *Phys. Rev. D* **105**, 074014 (2022). [[arXiv:2201.05728 \[hep-ph\]](#)].
- [31] J. Oudichhya, K. Gandhi, A. K. Rai, Mass-spectra of singly, doubly, and triply bottom baryons, *Phys. Rev. D* **104**, 114027 (2021). [[arXiv:2111.00236 \[hep-ph\]](#)].
- [32] A. V. Berezhnuy, A. K. Likhoded, A. V. Luchinsky, Doubly heavy baryons at LHC,

- Phys. Rev. D 98, 113004 (2018). [arXiv:1809.10058 [hep-ph]].
- [33] Z. S. Brown, W. Detmold, S. Meinel, and K. Orginos, Charmed bottom baryon spectroscopy from lattice QCD, Phys. Rev. D 90, 094507 (2014). [arXiv:1409.0497 [hep-ph]].
- [34] Y-L. Ma, M. Harada, Chiral partner structure of doubly heavy baryons with heavy quark spin-flavor symmetry, J. Phys. G: Nucl. Part. Phys 45, 075006 (2018). [arXiv:1709.09746 [hep-ph]].
- [35] D-H. He, K. Qian, Y-B. Ding, X-Q. Li, P-N. Shen, Evaluation of the spectra of baryons containing two heavy quarks in a bag model, Phys. Rev. D 70, 094004 (2004). [arXiv:0403301 [hep-ph]].
- [36] C. Albertus, E. Hernandez, J. Nieves, J. M. Verde-Velasco, Static properties and semileptonic decays of doubly heavy baryons in a nonrelativistic quark model, Eur. Phys. J. A 36, 119 (2008). [arXiv:0610030 [hep-ph]].
- [37] X-Z. Weng, X-L. Chen, W-Z. Deng, Masses of doubly heavy-quark baryons in an extended chromomagnetic model, Phys. Rev. D 97, 054008 (2018). [arXiv:1801.08644 [hep-ph]].
- [38] M. Padmanath, R. G. Edwards, N. Mathur, and M. Peardon, Spectroscopy of doubly charmed baryons from lattice QCD, Phys. Rev. D 91, 094502 (2015). [arXiv:1502.01845 [hep-ph]].
- [39] B. Aubert, *et al.* (BABAR Collaboration), Search for doubly charmed baryons Ξ_{cc}^+ and Ξ_{cc}^{++} in BABAR, Phys. Rev. D 74, 011103 (2006). [arXiv:0605075 [hep-ex]].
- [40] R. Chistov, *et al.* (BABAR Collaboration), Observation of new states decaying into $\Lambda_c^+ K^- \pi^+$ and $\Lambda_c^+ K_S^0 \pi^-$, Phys. Rev. Lett. 97, 162001 (2006). [arXiv:0606051 [hep-ex]].
- [41] Q-F. Lü, K-L. Wang, L-Y. Xiao, X-H. Zhong, Mass spectra and radiative transitions of doubly heavy baryons in a relativized quark model, Phys. Rev. D 96, 114006 (2017). [arXiv:1708.04468 [hep-ph]].
- [42] B. Eakins and W. Roberts, Symmetries and systematics of doubly heavy hadrons, Int. J. Mod. Phys. A 27, 1250039 (2012). [arXiv:1201.4885 [nucl-th]].
- [43] M. Karliner, J. L. Rosner, Strange baryons with two heavy quarks, Phys. Rev. D 97, 094006 (2018). [arXiv:1803.01657 [hep-ph]].
- [44] F. Giannuzzi, Doubly heavy baryons in a Salpeter model with AdS/QCD inspired potential, Phys. Rev. D 79, 094002 (2009). [arXiv:0902.4624 [hep-ph]].
- [45] T. Yoshida, E. Hiyama, A. Hosaka, M. Oka, and K. Sadato, Spectrum of heavy baryons in the quark model, Phys. Rev. D 92, 114029 (2015). [arXiv:1510.01067 [hep-ph]].
- [46] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, A. I. Onishchenko, Spectroscopy of doubly heavy baryons, Phys. Atom. Nucl 63, 274 (2000). [arXiv:9811212 [hep-ph]].
- [47] W. Wang, F-S. Yu, Z-X. Zhao, Weak decays of doubly heavy baryons: the $1/2 \rightarrow 1/2$ case, Eur. Phys. J. C 77, 781 (2017). [arXiv:1707.02834 [hep-ph]].
- [48] C. Alexandrou, V. Drach, K. Jansen, C. Kallidonis, G. Koutsou, Baryon spectrum with $N_f=2+1+1$ twisted mass fermions, Phys. Rev. D 90, 074501 (2014). [arXiv:1406.4310 [hep-ph]].
- [49] H-Y. Cheng, Y-L. Shi, Lifetimes of doubly charmed baryons, Phys. Rev. D 98, 113005 (2018). [arXiv:1809.08102 [hep-ph]].
- [50] K. Azizi, N. Er, Effects of a dense medium on parameters of doubly heavy baryons,

- Phys. Rev. D 100, 074004 (2019). [arXiv:1906.09087 [hep-ph]].
- [51] Z. Shah, A. K. Rai, Excited state mass spectra of doubly heavy Ξ baryons, Eur. Phys. J. C 77, 129 (2017). [arXiv:1702.02726 [hep-ph]].
- [52] J-W. Zhang, X-G. Wu, T. Zhong, Y. Yu, Z-Y. Fang, Hadronic production of the doubly heavy baryon Ξ_{cc} at LHC, Phys. Rev. D 83, 034026 (2011). [arXiv:1101.1130 [hep-ph]].
- [53] H-Y. Cheng, F. Xu, Lifetimes of doubly heavy baryons \mathfrak{B}_{bb} and \mathfrak{B}_{bc} , Phys. Rev. D 99, 073006 (2019). [arXiv:1903.08148 [hep-ph]].
- [54] V. V. Kiselev, A. K. Likhoded, A. I. Onishchenko, Lifetimes of Ξ_{bc}^+ and Ξ_{bc}^0 baryons, Eur. Phys. J. C 16, 461 (2000). [arXiv:9901224 [hep-ph]].
- [55] C-H. Chang, J-X. Wang, X-G. Wu, GENXICC2.0: an upgraded version of the generator for hadronic production of double heavy baryons Ξ_{cc} , Ξ_{bc} and Ξ_{bb} , Comput. Phys. Commun 181, 1144 (2010). [arXiv:0910.4462 [hep-ph]].
- [56] E. Ortiz-Pacheco, R. Bijker, Masses and radiative decay widths of S - and P -wave singly, doubly, and triply heavy charm and bottom baryons, Phys. Rev. D 108, 054014 (2023). [arXiv:2307.04939 [hep-ph]].
- [57] X-Z. Weng, X-L. Chen, and W-Z. Deng, Masses of doubly heavy-quark baryons in an extended chromomagnetic model, Phys. Rev. D 97, 054008 (2018). [arXiv:1801.08644 [hep-ph]].
- [58] K. U. Can, G. Erkol, B. Isildak, M. Oka, T. T. Takahashi, Electromagnetic properties of doubly charmed baryons in Lattice QCD, Phys. Lett. B 726, 703 (2013). [arXiv:1306.0731 [hep-ph]].
- [59] Z-G Wang, Analysis of the $\frac{1}{2}^+$ doubly heavy baryon states with QCD sum rules, Eur. Phys. J. A 45, 267 (2010). [arXiv:1001.4693 [hep-ph]].
- [60] H-Y. Cheng, Charmed baryon physics circa 2021, Chin. J. Phys 78, 324 (2022). [arXiv:2109.01216 [hep-ph]].
- [61] H-S. Li, W-L. Yang, Spin- $\frac{3}{2}$ doubly charmed baryon contribution to the magnetic moments of the spin- $\frac{1}{2}$ doubly charmed baryons, Phys. Rev. D 103, 056024 (2021). [arXiv:2012.14596 [hep-ph]].
- [62] Z. Shah, K. Thakkar, A. K. Rai, Excited state mass spectra of doubly heavy baryons Ω_{cc} , Ω_{bb} and Ω_{bc} , Eur. Phys. J. C 76, 530 (2016). [arXiv:1609.03030 [hep-ph]].
- [63] J. Oudichhya, K. Gandhi, A. K. Rai, Mass spectra of Ξ_{cc} , Ξ_{bc} , Ω_{cc} , and Ω_{bc} baryons in Regge Phenomenology, Phys. Scr. 97, 054001 (2022). [arXiv:2204.10045 [hep-ph]].
- [64] V. V. Kiselev, A. K. Likhoded, O. N. Pakhomova, V. A. Saleev, Mass spectra of doubly heavy $\Omega_{QQ'}$ baryons, Phys. Rev. D 66, 034030 (2002). [arXiv:0206140 [hep-ph]].
- [65] M-H. Weng, X-H. Guo, A. W. Thomas, Bethe-Salpeter equation for doubly heavy baryons in the covariant instantaneous approximation, Phys. Rev. D 83, 056006 (2011). [arXiv:1012.0082 [hep-ph]].
- [66] N. Mathur, M. Padmanath, S. Mondal, Precise predictions of charmed-bottom hadrons from lattice QCD, Phys. Rev. Lett. 121, 202002 (2018). [arXiv:1806.04151 [hep-ph]].
- [67] A. Kakadiya, A. K. Rai, Spectroscopic study of Ω_{cc} , Ω_{cb} and Ω_{bb} baryons, Few-Body Systems 64, 17 (2023). [arXiv:2303.03771 [hep-ph]].
- [68] R. Aaij *et al.* (LHCb Collaboration), Search for the doubly charmed baryon Ω_{cc}^+ ,

- Sci. China-Phys. Mech. Astron. 64, 101062 (2021). [[arXiv:2105.06841 \[hep-ex\]](#)].
- [69] K-W Wei, B. Chen, N. Liu, Q-Q Wang, X-H Guo, Spectroscopy of singly, doubly, and triply bottom baryons, *Phys. Rev. D* 95, 116005 (2017). [[arXiv:1609.02512 \[hep-ph\]](#)].
- [70] N. Salehi, Spectroscopy of Ω_{cc} , Ω_{bb} and Ω_{bc} baryons in hypercentral constituent quark model via ansatz method, *Acta. Phys. Polon. B* 50, 735 (2019).