# Light quark contributions to Higgs decays

A. I. Hernández-Juárez,<sup>1, \*</sup> R. Gaitán,<sup>1</sup> and R. Martinez<sup>2</sup>

<sup>1</sup>Departamento de Física, FES-Cuautitlán, Universidad Nacional Autónoma de México, C.P. 54770, Estado de México, México.

<sup>2</sup>Departamento de Física, Universidad Nacional de Colombia, K. 45 No. 26-85, Bogotá D.C., Colombia

(Dated: February 5, 2025)

The literature establishes that the light fermions contributions to the decays  $H \to Z\gamma$  and  $H \to \gamma\gamma$ are negligible since their coupling with the Higgs is proportional to  $m_f$ . In the present letter, we show that although such a conclusion is true for leptons, the light quark contributions are zero when we consider their non-perturbative effects.

### I. INTRODUCTION

Precision measurements of Higgs couplings at the LHC are essential for assessing whether they align with the predictions of the SM. Recently, the CMS and ATLAS collaborations announced the first measurements of the Higgs boson width, as well as the off-shell contributions of the Higgs to the production of two Z bosons [1, 2]. Additionally, both collaborations have observed an excess in the production of  $Z\gamma$  pairs through a Higgs boson [3, 4].



FIG. 1. One-loop SM contributions in the unitary gauge to the  $H \to Z\gamma$  decay.

The  $H \to \gamma \gamma$  and  $H \to Z \gamma$  decays are loop-induced in the SM through the Feynman diagrams in Fig. 1, with the vertex function for both processes given as follows [5, 6]:

$$\Gamma^{\mu\nu}_{V\gamma H} = h_1^{V\gamma} g^{\mu\nu} + \frac{1}{m_Z^2} \Big\{ h_2^{V\gamma} p_1^{\nu} p_2^{\mu} + h_3^{V\gamma} \epsilon^{\mu\nu\alpha\beta} p_{1\alpha} p_{2\beta} \Big\}, \quad (V = \gamma, Z),$$
(1)

where we follow the notation in Fig 2. The  $h_{1,2}^{V\gamma}$  form factors are *CP*-conserving, whereas  $h_3^{V\gamma}$  is associated with *CP* violation. Due to gauge invariance, the  $h_1^{V\gamma}$  and  $h_2^{V\gamma}$  form factors are not independent of each other:

$$h_2^{Z\gamma} = \frac{2 m_Z^2}{m_Z^2 - m_H^2} h_1^{Z\gamma},\tag{2}$$

$$h_2^{\gamma\gamma} = -2 \ h_1^{\gamma\gamma}. \tag{3}$$



FIG. 2.  $HZ\gamma$  coupling

<sup>\*</sup> alan.hernandez@cuautitlan.unam.mx

$$h_1^{Z\gamma}(W) = -3.4 \times 10^{-1} \text{GeV}$$
<sup>(4)</sup>

$$h_1^{\gamma\gamma}(W) = 4.8 \times 10^{-1} \text{GeV}$$
 (5)

In the literature, contributions from heavy fermions are usually emphasized, while those from light quarks and leptons are regarded as negligible. This conclusion arises from the coupling  $H\bar{f}f$ , which is proportional to the mass  $m_f$ . Consequently, the contributions to  $h_1^{ZV}$  ( $V = Z, \gamma$ ) will be proportional to  $m_f^2$  and can be neglected (see the discussion of one-loop contributions of light fermions in Refs [7, 11, 12, 14, 16–26]). Nevertheless, this argument is based on a perturbative approach, which is inadequate for light quarks due to their non-perturbative effects.

In this study, we aim to investigate the role of light quarks in the decay processes  $H \to Z\gamma$  and  $H \to \gamma\gamma$  to determine whether their contributions can be considered insignificant or zero.

# II. FERMION CONTRIBUTIONS TO $h_1^{V\gamma}$ ( $V = Z, \gamma$ )

In perturbation theory, the fermion contribution from the diagram in Fig. 1(a) can be expressed in terms of the Passarino-Veltman scalar functions as follows

$$h_{1}^{Z\gamma}(F) = \frac{N_{c}\mathcal{Q}_{f} \ g \ g_{V} \ e \ m_{f}^{2}}{8\pi^{2}c_{W}m_{W} \left(m_{H}^{2} - m_{Z}^{2}\right)} \left\{ 2m_{Z}^{2} \left[ B_{0} \left(m_{H}^{2}, m_{f}^{2}, m_{f}^{2}\right) - B_{0} \left(m_{Z}^{2}, m_{f}^{2}, m_{f}^{2}\right) \right] + \left(m_{H}^{2} - m_{Z}^{2}\right) \left( \left(-m_{H}^{2} + 4m_{f}^{2} + m_{Z}^{2}\right) C_{0} \left(0, m_{H}^{2}, m_{Z}^{2}, m_{f}^{2}, m_{f}^{2}\right) + 2 \right) \right\},$$

$$(6)$$

$$h_1^{\gamma\gamma}(F) = -\frac{e^2 \mathcal{Q}_f^2 m_f^2 N_c}{8\pi^2 m_W} \Big\{ 2 - \left( m_H^2 - 4m_f^2 \right) \mathcal{C}_0 \left( 0, 0, m_H^2, m_f^2, m_f^2, m_f^2 \right) \Big\}.$$
(7)

where  $m_f$  and  $Q_f$  denote the mass and charge of the fermion f in the loop, respectively. Additionally,  $g_V$  corresponds to the vector coupling of the Z boson with the fermions, and  $N_c$  stands for the color number. Contributions from heavy quarks and leptons can be calculated using Eqs (6) and (7). For this analysis, we adopt the masses  $m_t = 172.57$ GeV,  $m_c = 1.51$  GeV, and  $m_b = 4.18$  GeV, as recommended by the LHCHWG [27, 28]. Public codes, such as HDECAY, have implemented the masses  $\overline{m}_q(\mu)$  in the  $\overline{\text{MS}}$  scheme to assess the contributions of the bottom and charm quarks [29]. The numerical values from the top, bottom, and charm quarks, along with the total contributions from leptons, are summarized in Table. I.

The imaginary parts of the form factors  $h_1^{V\gamma}$  ( $V = Z, \gamma$ ) emerge when the fermions in the loop that are coupled to the Higgs boson can be on-shell. Notably, the contributions from the top quark are comparable in magnitude to those arising from the W-loops. The remaining contributions, however, are two to three orders of magnitude smaller than the top quark contributions and can be considered negligible. This finding is expected from Eqs (6)-(7), where we find that  $h_1^{V\gamma} \sim m_f^2$ , reinforcing the argument found in the literature to neglect the contributions from light fermions.

Contribution	$h_1^{\gamma\gamma}$	$h_1^{Z\gamma}$
Top quark	$-1.07 \times 10^{-1} \text{ GeV}$	$1.9 \times 10^{-2} \text{ GeV}$
Bottom quark	$(1.399 \times 10^{-3} - i \ 1.84 \times 10^{-3}) \ \text{GeV}$	$(-3.91 \times 10^{-4} + i \ 2.20 \times 10^{-4}) \text{ GeV}$
Charm quark	$(1.13 \times 10^{-3} - i \ 9.26 \times 10^{-4}) \ \text{GeV}$	$(-8.07 \times 10^{-5} + i \ 3.32 \times 10^{-5}) \ \text{GeV}$
Leptons	$(1.395 \times 10^{-3} - i \ 1.26 \times 10^{-3}) \ { m GeV}$	$(-1.46 \times 10^{-5} + i \ 6.3 \times 10^{-6}) \ \text{GeV}$

TABLE I. Perturbative contributions to the form factors  $h_1^{\gamma\gamma}$  and  $h_1^{Z\gamma}$  from the top, bottom, charm quarks, and leptons. For the quark masses, we use the values recommended by the LHCHWG [27, 28].

The contributions from light fermions require a different treatment, as their masses are not well-defined within a perturbative regime. One viable approach is to consider the chiral limit [30–32], defined in a non-perturbative

framework. In QCD, chiral symmetry is dynamically broken [33–35], which provides a mechanism for generating quark masses. Within context, one can define and compute one-loop corrections of the light quarks, including their chromomagnetic dipole moments [36, 37]. In Table II, we present our calculations of the light quark contributions to  $h_1^{V\gamma}$ . This analysis was conducted through a perturbative approach, assuming that custodial symmetry is preserved. For the light quark masses, we adopt  $m_u \approx m_d \approx 8$  MeV and  $m_s \approx 150$  MeV for the strange quark. We observe that the contributions from light quarks are negligible compared with those reported in Table I, as pointed out in the literature.

Contribution	$h_1^{\gamma\gamma}$	$h_1^{Z\gamma}$
Up quark	$(2.28 \times 10^{-7} - i \ 7.73 \times 10^{-8}) \ \text{GeV}$	$(-5.38 \times 10^{-9} + i \ 9.33 \times 10^{-10}) \ \text{GeV}$
Down quark	$(5.72 \times 10^{-8} - i \ 1.93 \times 10^{-8}) \ \text{GeV}$	$(-4.64 \times 10^{-9} + i \ 8.06 \times 10^{-10}) \ \text{GeV}$
Strange quark	$(9.36 \times 10^{-6} - i \ 4.73 \times 10^{-6}) \ \text{GeV}$	$(-1.1 \times 10^{-6} + i \ 2.83 \times 10^{-7}) \ \text{GeV}$

TABLE II. Perturbative contributions to the form factors  $h_1^{\gamma\gamma}$  and  $h_1^{Z\gamma}$  from the up, down and strange quarks. For the quark masses, we use  $m_u = m_d = 8$  MeV and  $m_s = 150$  MeV.

#### **III. PION CONTRIBUTIONS**



FIG. 3.  $H \to V\gamma$  ( $V = Z, \gamma$ ) decay mediated by the  $\pi^0, \eta^0$  and  $\eta'$  mesons.

At low energies, the light quarks form bound states and the  $H \to V\gamma$  ( $V = \gamma, Z$ ) decay is mediated by  $\pi^0, \eta^0$  and  $\eta'$  mesons as shown in Fig. 3. These particles are described by the states:

$$\pi^0 = \frac{\overline{u}u - \overline{d}d}{\sqrt{2}},\tag{8}$$

$$\eta^0 = \frac{\overline{u}u + \overline{d}d - 2\overline{s}s}{\sqrt{6}},\tag{9}$$

$$\eta' = \frac{\overline{u}u + \overline{d}d + \overline{s}s}{\sqrt{3}}.$$
(10)

Moreover, the coupling of the Higgs boson with the light quarks is given as

$$\mathcal{L} = -\frac{g}{2m_W} \Big( m_u \overline{u}u + m_d \overline{d}d + m_u \overline{s}s \Big) H, \tag{11}$$

Therefore, the amplitude for the decay  $H \rightarrow \gamma \gamma$  mediated by a meson can be expressed as follows

$$\mathcal{M}(H \to \gamma \gamma) = -\frac{g}{2m_W(m_H^2 - m_{P^0}^2)} \left\langle 0 \right| m_u \overline{u} u + m_d \overline{d} d + m_u \overline{s} s \left| P^0 \right\rangle A_{P^0}^{\mu\nu} \epsilon^{\mu}(p_1) \epsilon^{\nu}(p_2), \tag{12}$$

where  $P^0 = \pi^0$ ,  $\eta$ ,  $\eta'$  and  $A^{\mu\nu}_{P^0}$  correspondes to the  $P^0 \to \gamma\gamma$  amplitude. It is well known that the under parity transformations (P) that

$$P\left|P^{0}\right\rangle = -\left|P^{0}\right\rangle,\tag{13}$$

$$P\left|0\right\rangle = \left|0\right\rangle,\tag{14}$$

$$P\overline{\psi}\psi P^{-1} = \overline{\psi}\psi. \tag{15}$$

Therefore,

$$\langle 0| P^{-1}P(m_u\overline{u}u + m_d\overline{d}d + m_u\overline{s}s)P^{-1}P|\beta\rangle = -\langle 0| m_u\overline{u}u + m_d\overline{d}d + m_u\overline{s}s|\beta\rangle = 0$$
(16)

and

$$\mathcal{M}(H \to \gamma \gamma) = 0. \tag{17}$$

Therefore, the light fermions do not contribute to the  $H \to \gamma \gamma$  decay. This conclusion stands in contrast to those reached in Refs. [7, 11, 12, 14, 16–26], which suggest that the contributions from light quarks can be neglected as they are insignificant. From a similar argument, the light quarks do not contribute to the  $H \to Z\gamma$  decay either.

## IV. CONCLUSION

In this note, we demonstrated that the contributions from the bottom, charm quark, and lepton to the decays  $H \rightarrow \gamma \gamma$  and  $H \rightarrow Z \gamma$  are significantly smaller than those arising from the top quark and W-loops. Therefore, as pointed out in the literature, these contributions can be disregarded. In contrast, the contributions from the up, down, and strange quarks are zero rather than negligible.

#### ACKNOWLEDGMENTS

This work was supported by UNAM Posdoctoral Program (POSDOC). We also acknowledge support from Sistema Nacional de Investigadores (Mexico).

- A. Tumasyan *et al.* (CMS), Measurement of the Higgs boson width and evidence of its off-shell contributions to ZZ production, Nature Phys. 18, 1329 (2022), arXiv:2202.06923 [hep-ex].
- [2] G. Aad et al. (ATLAS), Evidence of off-shell Higgs boson production from ZZ leptonic decay channels and constraints on its total width with the ATLAS detector, Phys. Lett. B 846, 138223 (2023), arXiv:2304.01532 [hep-ex].
- [3] A. Tumasyan *et al.* (CMS), Search for Higgs boson decays to a Z boson and a photon in proton-proton collisions at  $\sqrt{s} = 13$  TeV, JHEP **05**, 233, arXiv:2204.12945 [hep-ex].
- [4] G. Aad et al. (ATLAS, CMS), Evidence for the Higgs Boson Decay to a Z Boson and a Photon at the LHC, Phys. Rev. Lett. 132, 021803 (2024), arXiv:2309.03501 [hep-ex].
- [5] R. Martínez, M. Pérez, and J. Toscano, The decays  $H^0 \to Z\gamma$  and  $Z' \to H^0\gamma$  in left-right symmetric models, Physics Letters B 234, 503 (1990).
- [6] R. Martinez, M. A. Perez, and J. J. Toscano, The Two Photon Decay Width of the Higgs Boson in Left-right Symmetric Theories, Phys. Rev. D 40, 1722 (1989).
- [7] J. Ellis, M. K. Gaillard, and D. Nanopoulos, A phenomenological profile of the higgs boson, Nuclear Physics B 106, 292 (1976).
- [8] M. A. Shifman, A. I. Vainshtein, M. B. Voloshin, and V. I. Zakharov, Low-Energy Theorems for Higgs Boson Couplings to Photons, Sov. J. Nucl. Phys. 30, 711 (1979).
- [9] L. B. Okun, Leptons and Quarks: Special Edition Commemorating the Discovery of the Higgs Boson (North-Holland, Amsterdam, Netherlands, 1982).
- [10] M. Gavela, G. Girardi, C. Malleville, and P. Sorba, A non-linear  $R_{\xi}$  gauge condition for the electroweak  $SU(2) \times U(1)$  model, Nuclear Physics B **193**, 257 (1981).
- [11] R. N. Cahn, M. S. Chanowitz, and N. Fleishon, Higgs particle production by  $Z \to H\gamma$ , Phys. Lett. B 82, 113 (1979).
- [12] L. Bergstrom and G. Hulth, Induced Higgs Couplings to Neutral Bosons in  $e^+e^-$  Collisions, Nucl. Phys. B **259**, 137 (1985), [Erratum: Nucl.Phys.B **276**, 744–744 (1986)].
- [13] J. F. Gunion, H. E. Haber, G. L. Kane, and S. Dawson, The Higgs Hunter's Guide, Vol. 80 (2000).

- [14] R. Decker, M. Nowakowski, and D. Woitschitzky, The  $Z^0 \to H\gamma$  decay in two Higgs doublet models, Phys. Lett. B 255, 605 (1991).
- [15] A. I. Hernández-Juárez, R. Gaitán, and R. Martinez, The  $H \to Z\gamma$  decay and CP violation, (2024), arXiv:2405.03094 [hep-ph].
- [16] P. S. Bhupal Dev, D. K. Ghosh, N. Okada, and I. Saha, 125 GeV Higgs Boson and the Type-II Seesaw Model, JHEP 03, 150, [Erratum: JHEP 05, 049 (2013)], arXiv:1301.3453 [hep-ph].
- [17] M. Spira, A. Djouadi, and P. M. Zerwas, QCD corrections to the  $HZ\gamma$  coupling, Phys. Lett. B 276, 350 (1992).
- [18] R. Bonciani, V. Del Duca, H. Frellesvig, J. M. Henn, F. Moriello, and V. A. Smirnov, Next-to-leading order QCD corrections to the decay width  $H \rightarrow Z\gamma$ , Journal of High Energy Physics **2015**, 108 (2015).
- [19] T. Gehrmann, S. Guns, and D. Kara, The rare decay  $H \to Z\gamma$  in perturbative QCD, Journal of High Energy Physics **2015**, 38 (2015).
- [20] H.-q. Zheng and D.-d. Wu, First-order QCD corrections to the decay of the Higgs boson into two photons, Phys. Rev. D 42, 3760 (1990).
- [21] S. Dawson and R. P. Kauffman, QCD corrections to the Higgs-boson decay  $H \to \gamma \gamma$ , Phys. Rev. D 47, 1264 (1993).
- [22] A. Djouadi, M. Spira, and P. Zerwas, Two-photon decay widths of Higgs particles, Physics Letters B **311**, 255 (1993).
- [23] A. Djouadi, M. Spira, J. van der Bij, and P. Zerwas, QCD corrections to  $\gamma\gamma$  decays of Higgs particles in the intermediate mass range, Physics Letters B **257**, 187 (1991).
- [24] K. Melnikov and O. Yakovlev, Higgs-two-photon interaction in the standard model. The QCD radiative correction, Physics Letters B 312, 179 (1993).
- [25] J. Fleischer, O. Tarasov, and V. Tarasov, Analytical result for the two-loop QCD correction to the decay  $H \rightarrow 2\gamma$ , Physics Letters B **584**, 294 (2004).
- [26] A. Djouadi, P. Gambino, and B. Kniehl, Two-loop electroweak heavy-fermion corrections to Higgs-boson production and decay, Nuclear Physics B 523, 17 (1998).
- [27] D. de Florian *et al.* (LHC Higgs Cross Section Working Group), Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector 2/2017, 10.23731/CYRM-2017-002 (2016), arXiv:1610.07922 [hep-ph].
- [28] R. L. Workman et al. (Particle Data Group), Review of Particle Physics, PTEP 2022, 083C01 (2022).
- [29] A. Djouadi, J. Kalinowski, M. Muhlleitner, and M. Spira, HDECAY: Twenty++ years after, Computer Physics Communications 238, 214 (2019).
- [30] H. Pagels, Chiral Symmetry Realization in the Quark Model, Phys. Rev. Lett. 28, 1482 (1972).
- [31] H. Leutwyler, Mesons in terms of quarks on a null plane, Nuclear Physics B 76, 413 (1974).
- [32] P. Maris and C. D. Roberts, π- and K-meson Bethe-Salpeter amplitudes, Physical Review C 56, 3369 (1997).
- [33] M. S. Bhagwat, M. A. Pichowsky, C. D. Roberts, and P. C. Tandy, Analysis of a quenched lattice-QCD dressed-quark propagator, Phys. Rev. C 68, 015203 (2003).
- [34] M. S. Bhagwat, Analysis of full-QCD and quenched-QCD lattice propagators, in AIP Conference Proceedings, Vol. 842 (AIP, 2006) pp. 225–227.
- [35] P. O. Bowman, U. M. Heller, D. B. Leinweber, M. B. Parappilly, A. G. Williams, and J. Zhang, Unquenched quark propagator in Landau gauge, Physical Review D 71, 10.1103/physrevd.71.054507 (2005).
- [36] L. Chang, Y.-X. Liu, and C. D. Roberts, Dressed-Quark Anomalous Magnetic Moments, Physical Review Letters 106, 10.1103/physrevlett.106.072001 (2011).
- [37] R. Bermudez, L. Albino, L. X. Gutiérrez-Guerrero, M. E. Tejeda-Yeomans, and A. Bashir, Quark-gluon vertex: A perturbation theory primer and beyond, Phys. Rev. D 95, 034041 (2017).