NONLINEAR COVARIANCE SHRINKAGE FOR HOTELLING'S T^2 IN HIGH DIMENSION *

BY BENJAMIN D. ROBINSON †,a AND VAN LATIMER ‡,b

Air Force Office of Scientific Research 875 N. Randolph Rd Arlington, VA *machine.itel@us.af.mil

Radial Research and Development 1210 E. Dayton-Yellow Springs Rd Fairborn, OH byan.latimer@radialrad.com

In this paper we study the problem of comparing the means of a single observation and a reference sample in the presence of a common data covariance matrix, where the data dimension p grows linearly with the number of samples n and p/n converges to a number between 0 and 1. The approach we take is to replace the sample covariance matrix with a nonlinear shrinkage estimator—i.e., a matrix with the same eigenvectors—in Hotelling's T^2 test. Current approaches of this sort typically assume that the data covariance matrix has a condition number or spiked rank that increases slowly with dimension. However, this assumption is ill-suited to data sets containing many strongly correlated background covariates, as often found in finance, genetics, and remote sensing. To address this problem we construct, using variational methods and new local random-matrix laws, a nonlinear covariance shrinkage method tailored to optimize detection performance across a broad range of spiked ranks and condition numbers. We then demonstrate, via both simulated and real-world data, that our method outperforms existing approaches.

1. Introduction. Comparing the means of two samples sharing a covariance matrix is a fundamental problem in multivariate statistics, and Hotelling's T^2 test is often the method of choice when the number of samples n is much larger than the data dimension p [And63, Mui09]. In the more modern high-dimensional regime where n grows linearly with p, various authors [BS96, CQ10, LAP+20b] have proposed improvements that effectively replace the sample covariance matrix S_n appearing in the test by an affine function of S_n , also known as a linear shrinkage estimator [LW04, ML05, CM14]. However, for data corrupted by many strongly correlated signals, linear shrinkage can result in low testing power, likely because its single affine parameter cannot accurately model the full covariance spectrum [RML+22]. Therefore, more flexible *nonlinear* functions of the sample covariance matrix are needed.

Such scenarios featuring many strong correlations commonly arise in genetic microarray analysis [JL09] and heavily cluttered signal processing [BT02]. As a result, the problem of nonlinearly transforming—i.e., shrinking—the spectrum of a sample covariance matrix for various applications has been a significant topic [Ste75, Ste86, DS85, Nad14, DGJ18, LAP20a, LW22]. Stein provided an early glimpse into nonlinear shrinkage theory, but work

^{*}Thanks to AFOSR Lab Tasks 18RYCOR004, 19RYCOR036, 22RYCOR006

[†]US Air Force Office of Scientific Research

[‡]Radial Research and Development, Inc.

MSC2020 subject classifications: Primary 62H15, 60B20; secondary 62G10.

Keywords and phrases: Nonlinear covariance shrinkage, Hotelling's T^2 , high-dimensional statistics, random matrix theory, Hilbert transform.

stalled for many years due to a theoretically opaque "eigenvalue-isotonization" procedure. It was not until the recent work of Ledoit and Wolf [LW20] that a computationally efficient nonlinear shrinkage method was found and proven to be asymptotically optimal compared to any oracle shrinkage method for various loss functions of interest. Since then, this type of shrinkage has been successfully applied to many domains, including financial portfolio optimization [LW17b, LW20] and radar signature detection [RMH21]. However, to our knowledge, nonlinear shrinkage has not yet been rigorously explored for Hotelling's T^2 , despite the envisioned importance of doing so [NPW21].

In this paper, we develop the theory of nonlinear covariance shrinkage for Hotelling's T^2 . To enable the use of the powerful local anisotropic law of random matrix theory [KY17], we limit our attention to monotonic, bounded, and positive limiting spectral densities with bounded positive support. Although this choice may seem restrictive at first, we note that our model accurately approximates many finite-dimensional matrices drawn from other popular ones, such as the generalized spiked model [BY12], as well as many matrices with O(p) large eigenvalues.

As a preliminary to our main results, in Theorems 3 and 5 we find a stochastic convergence rate associated with Ledoit and Wolf's nonlinear shrinkage eigenvalues from [LW20]. Our main accomplishments in this paper are then the following:

- 1. In Theorem 6, we establish significance levels for the Hotelling T^2 test modified with covariance shrinkage.
- 2. In Theorem 7, we formulate and solve a deterministic variational problem for a shrinkage function that approximately optimizes the test's detection power.
- 3. In Theorem 8, we find an empirical approximation to the above shrinkage function for finite sample sizes that has the same asymptotic power.

We present these results and their preliminaries in the following sections. In Section 2, we provide an outline of the mean-shift detection problem to be considered, shrinkage estimators, and the shrinkage-modified Hotelling T^2 statistic (SRHT). In Section 3, we give an overview of the relevant local-law results in random matrix theory and present our subgaussian data model. In Section 4, we introduce the idea of nonlinear covariance shrinkage and present our convergence-rate analysis of Ledoit and Wolf's nonlinear shrinkage eigenvalues. In Section 5, we present our three main accomplishments listed above. In Section 6, we present our empirical results, showing that our approach outperforms existing ones on various simulated and real-world data.

2. Background on covariance shrinkage and Hotelling's T^2 . Mean-shift detection in this paper will mean the problem of comparing the mean of an observation to that of a reference data set having the same covariance matrix. Formulated as a hypothesis test, the question is, given i.i.d. random vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n \sim P$ and a random vector $\mathbf{y} \sim Q$ with $cov(\mathbf{x}_1) = cov(\mathbf{y})$, to decide between $\mathcal{H}_0 : \mathbb{E}[\mathbf{x}_1] = \mathbb{E}[\mathbf{y}]$ and $\mathcal{H}_1 : \mathbb{E}[\mathbf{x}_1] \neq \mathbb{E}[\mathbf{y}]$. When the common covariance matrix Σ is known, a typical test statistic for mean-shift detection is the energy detector, which takes the form

(1)
$$(\mathbf{y} - \overline{\mathbf{x}})' \mathbf{\Sigma}^{-1} (\mathbf{y} - \overline{\mathbf{x}}) \underset{\mathcal{H}_0}{\gtrless} \tau,$$

where $\overline{\mathbf{x}}$ is the sample mean of $\{\mathbf{x}_i\}$, τ is some detection threshold, and $(\cdot)'$ denotes the transpose operation. However, unless n is large—i.e., much larger than p—it not realistic to assume that one can approximate all the p^2 entries of Σ with much accuracy. In this case, it is standard to replace Σ by some estimator $\hat{\Sigma}$. The focus of this paper is on how to choose $\hat{\Sigma}$ in the *high-dimensional regime*, where n and p go to infinity and $p/n \to \phi \in (0,1)$.

Very often $\hat{\Sigma}$ is chosen to be the sample covariance matrix of $\{x_i\}$, defined to be

(2)
$$\mathbf{S}_n := \frac{1}{n-1} \sum_{i=1}^n (\mathbf{x}_i - \overline{\mathbf{x}}) (\mathbf{x}_i - \overline{\mathbf{x}})' = \frac{1}{n-1} \mathbf{X}_n \left(1 - \frac{1}{n} \overline{\mathbf{x}} \mathbf{1}_p \right) \mathbf{X}'_n,$$

where \mathbf{X}_n is the $p \times n$ matrix $[\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n]$ and $\mathbf{1}_p$ is the $1 \times p$ vector $(1, 1, \dots, 1)$. The resulting plug-in energy detector can be identified with Hotelling's T^2 statistic [Mui09] for large n. However, although this choice minimizes asymptotic detection loss compared to (1) when $n \gg p$, there is no reason to believe it has the same property in the high-dimensional regime. Indeed, Hotelling's T^2 is inconsistent in this regime [BS96].

A natural alternative approach is to choose $\hat{\Sigma}^{-1}$ to be a *precision shrinkage estimator*, which we define to be a positive semi-definite matrix that has the same eigenvectors as \mathbf{S}_n . In this case, we call the resulting test statistic a *shrinakge-regularized Hotelling* T^2 *statistic*, or *SRHT* statistic. Regarding precision shrinkage estimators, the most common ones take the form $f(\mathbf{S}_n) = \sum_{i=1}^p f(\lambda_i) \mathbf{u}_i \mathbf{u}_i'$ for some continuous, deterministic function f, where \mathbf{S}_n is diagonalized as $\sum_{i=1}^p \lambda_i \mathbf{u}_i \mathbf{u}_i'$. However, the values taken by f could depend on random quantities, such as the full spectrum of \mathbf{S}_n [LW20]. Given a sequence of shrinkage functions f_n , we will always assume f_n are almost surely continuous and uniformly bounded for large enough n.

The most common type of shrinkage-function sequence f_n corresponds to ridge or "linear" shrinkage and takes the $f_n(x) = 1/(x+\lambda_n)$ for some scalars $\lambda_n > 0$. This choice corresponds to the regularized Hotelling T^2 test statistics of [LAP+20b], which form a one-parameter subclass of the SRHTs we consider. Linear shrinkage is extremely well-studied, but for all the effort devoted to it, virtually all procedures for finding λ_n appear to either depend on *ad hoc* rules of thumb, suffer from numerical instability, or incur high computationally complexity, calling into question their usefulness in real-time applications.

In order to find the asymptotic distribution of the SRHT

(3)
$$\tilde{T}_n^2 \equiv \tilde{T}_n^2(f_n) := (\mathbf{y}_n - \overline{\mathbf{x}}_n)' f_n(\mathbf{S}_n) (\mathbf{y}_n - \overline{\mathbf{x}}_n),$$

we will need to summarize and develop a few results in random-matrix and nonlinear-shrinkage theory. In particular, we will need to provide a summary of the theory of so-called *nonlinear* shrinkage estimators, in which $f_n(x)$ is generally not of the form $(x + \lambda)^{-1}$. This more flexible type of shrinkage estimator has been an active subject of study beginning as early as with C. Stein [Ste75], but has only in the last decade or so been put on a solid theoretical foundation by Ledoit and Wolf [LW17a] and their collaborator S. Péché [LP11]. Motivated by questions of Sharpe-ratio optimization in mathematical finance, Ledoit and Wolf showed that the optimal relative weights in an investment portfolio can be obtained by applying their precision estimator to a known vector of real asset returns. In the context of hypothesis testing, some of our earlier work [RMH21] showed that input signal-to-interference-plus-noise ratio in radar detection is mathematically equivalent to the Sharpe ratio, so that the same eigenvalues approximately optimize power when used as replacements for the sample eigenvalues in an "adaptive matched filter," as defined in [RFKN92]. In the present paper, we provide a similar analysis for SRHTs.

3. Background on random matrix theory. The starting point of nonlinear shrinkage theory is the celebrated Marčenko–Pastur Theorem [MP67, Sil95], which we will now state in one of its modern forms. We emphasize that the samples \mathbf{x}_{ni} and sample means $\overline{\mathbf{x}}_n$, both of increasing dimension, depend on n (and p) by denoting them as \mathbf{x}_{ni} and $\overline{\mathbf{x}}_n$. We write the

sample matrix again in block form as $\mathbf{X}_n = [\mathbf{x}_{n1}, \mathbf{x}_{n2}, \dots, \mathbf{x}_{nn}]$, which can be visualized as in the following diagram:

$$p\left\{ \begin{bmatrix} \overbrace{ | |$$

Given that $\mathbf{S}_n = (n-1)^{-1}\mathbf{X}_n(1-n^{-1}\overline{\mathbf{x}}_n\mathbf{1}_p)\mathbf{X}_n$, as before, the norm difference of \mathbf{S}_n and $\tilde{\mathbf{S}}_n = n^{-1}\mathbf{X}_n\mathbf{X}_n'$ goes to zero when the population has mean zero. As a result of this and shift invariance, we will frequently assume the mean vanishes and identify $\tilde{\mathbf{S}}_n$ with \mathbf{S}_n , commenting on any complications that arise in the case of nonzero mean as necessary.

Except when commenting on the nonzero-mean case, we will assume throughout the rest of the paper the following data model (H1)-(H5) for the \mathbf{x}_{ni} 's unless otherwise noted.

- (H1) \mathbf{X}_n can be expressed as $\mathbf{\Sigma}_n^{1/2}\mathbf{W}_n$, where \mathbf{W}_n is a matrix of i.i.d. random variables $\sim W$ with zero mean, unit variance, and $\mathbb{E}[e^{sW}] \leq e^{s^2/2}$ for all real s.
- (H2) Σ_n is a deterministic symmetric positive-definite $p \times p$ matrix with eigenvalues $\tau_{n1}, \tau_{n2}, \dots \tau_{np}$,
- (H3) There is $\phi \in (0,1)$ such that $|p/n \phi| = O(1/p)$ as $n, p \to \infty$,
- (H4) As $n, p \to \infty$, (a) defining δ_{τ} to be the Dirac mass at τ , the population spectral distribution $\pi_n = p^{-1} \sum_{i=1}^p \delta_{\tau_{ni}}$ of Σ_n converges weakly to some limiting distribution, π_{∞} with compact positive support, and (b) all population eigenvalues eventually lie in this bounded set.
- (H5) π_{∞} is absolutely continuous, and $d\pi_{\infty}(x)/dx$ is positive, monotonic, and bounded on its support.

The assumptions (H1)-(H4) closely mimic the assumptions of [LW20]. The moment condition in (H1), also known as *one-subgaussianity*, can easily be relaxed to a sub-Weibull-type condition, but we will not be concerned this generalization since the analysis is much the same. Similarly, our assumption in (H3) that $\phi \in (0,1)$ is weaker than Ledoit and Wolf's, which allows for the "sample-starved" case of $\phi > 1$, but we leave analysis of this extension for future work since it is longer, is more technical, and does not particularly clarify our main results.

We note that the assumption (H3) means that we will be able to use the condition "as $n \to \infty$ " as a substitute for the equivalent but more verbose condition " $n, p \to \infty$ and $p/n = \phi + O(1/p)$." Even more briefly, when we express a convergence relation, such as $\|\mathbf{\Sigma}_n - \mathbf{I}_p\| \to 0$ without identifying a limiting regime, it will be understood that this convergence occurs as n (and thus p) goes to infinity.

Our primary notion of convergence rate in probability will be that of *stochastic domination*, defined as follows as in [KY17]:

DEFINITION 1. Let $X_n(u)$ and $Y_n(u)$ be random variables, parameterized by a family of real vectors $u \in U$. We say X_n is stochastically dominated by Y_n uniformly in u if for all $\epsilon, D > 0$ we have

$$\sup_{u \in U} \Pr\left[X_n(u) < n^{\epsilon} Y_n(u) \right] \le n^{-D}$$

for all n greater than some $n_0(\epsilon, D)$. More briefly, we write $|X_n| \prec Y_n$ or $X_n = O_{\prec}(Y_n)$.

REMARK 1. The underlying set U parameterizing random variables of interest will often not be explicitly identified; instead the notation $O_{\prec}(\,\cdot\,)$ will be implicitly taken to mean uniform stochastic domination over all parameters (e.g., matrix indices or complex spectral parameters) not expressly deemed constant. As a further notational remark, $|X_n| \lesssim Y_n$ will mean $X_n = O(Y_n)$ with probability as above, and $X_n \asymp Y_n$ will mean both $|X_n| \lesssim Y_n$ and $|Y_n| \lesssim X_n$.

We note that the sample covariance matrix can be expressed just as before, except with terms indexed by n to indicate their increasing dimension and dependence on n:

$$\mathbf{S}_n = (n-1)^{-1} \sum_{i=1}^n (\mathbf{x}_{ni} - \overline{\mathbf{x}}_n) (\mathbf{x}_{ni} - \overline{\mathbf{x}}_n)'.$$

Further, we will frequently make use of the sample eigen-decomposition:

$$\mathbf{S}_n = \sum_{i=1}^p \lambda_{ni} \mathbf{u}_{ni} \mathbf{u}'_{ni},$$

where \mathbf{u}_{ni} are almost surely uniquely defined since the probability of repeated sample eigenvalues is zero. The Marčenko-Pastur Theorem, then, is essentially a statement about the limiting behavior of the resolvent $\mathbf{R}_n(z)$ of \mathbf{S}_n , given by

$$\mathbf{R}_n(z) = (\mathbf{S}_n - z\mathbf{I}_n)^{-1},$$

and the spectral measure of S_n : namely,

$$\mu_n = p^{-1} \sum_{i=1}^p \delta_{\lambda_{ni}}.$$

A strong form of the Marčenko-Pastur Theorem due to Knowles and Yin is reproduced below.

THEOREM 1 (Local Marčenko–Pastur Laws [KY17]). Assume (H1)-(H5) and that μ_n is the spectral distribution of the sample covariance matrix \mathbf{S}_n . Let

$$m_n(z) := p^{-1} \operatorname{tr}(\mathbf{R}_n(z)).$$

Then for all bounded $z = x + i\eta$ in the complex upper half plane \mathbb{C}^+ , we have

(4)
$$m_n(z) = m_{\infty}(z) + O_{\prec}\left(\frac{1}{n\eta}\right)$$

as $n, p \to \infty$, where $m_{\infty}(z)$ is the unique solution to

(5)
$$m_{\infty}(z) = \int_0^{\infty} \frac{d\pi_{\infty}(\tau)}{\tau(1 - \phi - \phi z m_{\infty}(z)) - z}.$$

Furthermore, (the "small-scale" laws) there is a nonrandom measure μ_{∞} such that

(6)
$$\sup_{(a,b)\subset\mathbb{R}} |\mu_n(a,b) - \mu_\infty(a,b)| \prec \frac{1}{n}$$

and, uniformly in $f \in C^2$,

(7)
$$\int f \, d\mu_n = \int f \, d\mu_\infty + O_{\prec} \left(\frac{\|f\|_1}{n} + \frac{\|f'\|_1}{n} + \frac{\|f''\|_1}{n^2} \right),$$

as $n, p \to \infty$.




Fig 1: Marchenko-Pastur law $w(x) = d\mu_{\infty}(x)/dx$ (red) versus histogram of eigenvalues of \mathbf{S}_n for n = 5000, p = 1000, and $\mathbf{\Sigma}_n = \mathbf{I}_p$. Due the error bounds of roughly 1/n in the small-scale law of Theorem 1, a fairly high-resolution histogram matches w(x) reasonably well.

PROOF. The first statement is the main conclusions of the fluctuating-averaging argument in the cited paper. The small-scale laws follow from the somewhat more general Corollary 1 and Lemma 6 of Appendix B. Both small-scale laws are immediate consequences of existing work by Helffer and Sjöstrand, which that appendix reproduces for convenience.

REMARK 2. The classical Marčenko–Pastur Theorems of [Sil95] hold under more general moment and spectral conditions, but only provide $o_{\rm a.s.}(1)$ convergence rates. The condition (H5) is sufficient to obtain our conclusion, but not necessary: a somewhat more general spectral criterion than (H5) is provided in [KY17, Definition 2.7]. If only convergence in probability is needed, we conjecture that the condition on $d\pi_{\infty}(x)/dx$ can be dropped completely.

Throughout this paper, we will let $F := \operatorname{supp} \mu_{\infty}$, let $w(x) := d\mu_{\infty}(x)/dx$, and let $\kappa(x)$ be defined as the distance from x to the boundary of F. Some important facts are that F is the union of finitely many disjoint closed intervals of nonzero length [BS⁺98], and that w(x) is a smooth function except near the edges of these intervals, where w(x) and $\kappa(x)^{1/2}$ are on the same order of magnitude [KY17].

REMARK 3. We note that the integral equation (5) does not have a known solution for $m_{\infty}(z)$ except in special cases, such as when $\Sigma_n = \mathbf{I}_p$ or π_{∞} is otherwise a convex combination of two Dirac masses. As a result, the same is true for w(x). The rare solvable case of $\Sigma_n = \mathbf{I}_p$, and its relation to the small-scale law of Theorem 1, is illustrated in Figure 1.

Define the Stieltjies transform of a positive measure ρ by $\mathcal{S}[d\rho](z) := \int (t-z)^{-1} d\rho(t)$, so that $m_n(z) = \mathcal{S}[d\mu_n](z)$. It turns out that μ_∞ is defined by $m_\infty(z) = \mathcal{S}[d\mu_\infty](z)$ for $z \in \mathbb{C}^+$, and that exteding $m_\infty(z)$ and related Stieltjies transforms to the real line will be of interest. In [SC⁺95], it is shown that for $x \in \mathbb{R}$ the following limit exits

$$\lim_{z \in \mathbb{C}^+ \to x} m_{\infty}(z) =: \breve{m}(x),$$

and, further $\mathrm{Im}[\check{m}(x)] = \pi w(x)$. Another fact is that the real part of $\check{m}(x)$ is given by $\pi \mathcal{H}[d\mu_{\infty}](x)$ [LW20, Supplement A.1], where the Hilbert transform $\mathcal{H}[d\rho](x)$ of a finite Borel measure ρ with $d\rho/dx \in L^q$ for $q \in (1,\infty)$ is given by the principle-value integral

$$\mathcal{H}[d\rho](x) := \text{p.v.} \frac{1}{\pi} \int \frac{d\rho(t)}{t-x},$$

which we will also define to be the Hilbert transform of $d\rho/dx$. Combining these two results about $\check{m}(x)$, we get

$$\check{m}(x) = \pi \mathcal{H}[d\mu_{\infty}](x) + i\pi w(x). \qquad (x \in \mathbb{R})$$

It follows in much the same way that if $f \in L^1(d\mu_\infty)$, we may write

(8)
$$\lim_{z \in \mathbb{C}^+ \to x} \mathcal{S}[f \, d\mu_{\infty}](z) = \pi \mathcal{H}[f \, d\mu_{\infty}](x) + i\pi f(x)w(x)$$

for all $x \in \mathbb{R}$. We emphasize that Hilbert transforms of the form $\mathcal{H}[f dx]$ will, as usual,often be written as $\mathcal{H}[f]$ or simply $\mathcal{H}f$. For example, $\mathcal{H}[f d\mu_{\infty}]$ will often be denoted by $\mathcal{H}[fw]$.

4. Nonlinear shrinkage background and refinements. In a seminal paper extending the classical Marčenko-Pastur theorem of [Sil95], Ledoit and Péché [LP11] derive an almost sure deterministic limit for generalized resolvents of the form $\Theta_n^g(z) = p^{-1} \operatorname{tr}(g(\Sigma_n) \mathbf{R}_n(z))$, where g is a bounded piece-wise continuous function applied spectrally to Σ_n . These resolvents are important in covariance and precision shrinkage, as well as in the determination of asymptotic eigenvector biases, which appear, for example, in principle-components analysis.

We present in Subsection 4.1 a refinement of the Ledoit–Péché theory that provides explicit convergence rates—analogous to Theorem 1's refinement of the classical Marčenko–Pastur theorem. Then in Subsection 4.2, we present a similar refinement of Ledoit and Wolf's work [LW20], which can be seen as the empirical version of Ledoit and Péché's law. These refinements will enable us to standardize the SRHTs of interest in Section 5.

4.1. Ledoit–Péché Theory. The functions $\Theta^g(z)$ are of particular interest when g(x)=1, just discussed in Theorem 1, and when $g(x)=\operatorname{id}(x)\equiv x$, which is the subject of this section. When $g=\operatorname{id}$, we will simplify notation by writing $\Theta_n(z)$ instead of $\Theta_n^{\operatorname{id}}(z)$. Then the following analogue of the classical Marčenko–Pastur theorem holds.

THEOREM 2 ([LP11]). Assume (H1)-(H4) and $z \in \mathbb{C}^+$. Let

$$\Theta_n(z) := p^{-1} \operatorname{tr}(\mathbf{\Sigma}_n \mathbf{R}_n(z)).$$

Then $\Theta_n(z) = \mathcal{S}[d\nu_n](z)$, where

$$\nu_n := \frac{1}{p} \sum_{i=1}^p \mathbf{u}'_{ni} \mathbf{\Sigma}_n \mathbf{u}_{ni} \delta_{\lambda_{ni}},$$

and there exists a nonrandom analytic function $\Theta_{\infty}(z)$ such that

(9)
$$\Theta_n(z) = \Theta_{\infty}(z) + o_{as}(1)$$

as $n, p \to \infty$. Furthermore, if $\mathcal{H}w(x)$ is the Hilbert transform of w(x) and the function $\delta : \mathbb{R}_+ \to \mathbb{R}_+$ is defined by

(10)
$$\delta(x) = \frac{x}{[1 - \phi - \phi \pi x \mathcal{H} w(x)]^2 + \phi^2 \pi^2 x^2 w(x)^2}$$

then

(11)
$$d\nu_n \to \delta d\mu_\infty =: d\nu_\infty$$

weakly almost surely as $n, p \to \infty$.

PROOF. The first statement follows from diagonalization of $\mathbf{R}_n(z)$ and the circular-permutation property of trace. Equation (9) follows from [LP11, Theorem 2]. Equation (11) follows from [LP11, Theorem 4].

REMARK 4. The form of (10) differs in appearance from the form given in Ledoit and Péché's paper, but defines the same function of x. The form given there is

$$\frac{1}{|1-\phi-\phi x\breve{m}(x)|^2}.$$

As we discussed the last section, the imaginary part of $\check{m}(x)$ is just $\pi w(x)$, and the real part is π times the Hilbert transform of $d\mu_{\infty}(x) = w(x) \, dx$, so the identification is complete.

The choice $g=\operatorname{id}$ studied in this section is important for several reasons. First, the coefficients of the measure ν_n resulting from $\Theta_n^g(z)$ optimize various loss functions of statistical interest, such as Frobenius loss [LP11], reverse Stein loss [LW18], the Sharpe ratio appearing in portfolio balancing [LW17b], and the Reed–Mallet–Brennan figure of merit in radar detection [RMH21], assuming an ignorant signal prior for the latter two. A potential wrinkle, as discussed in Remark 3, is that w(x) has no known closed form expression except in a handful of special cases. However, fortunately, Ledoit and Wolf [LW20] have developed observable substitutes for w(x) and $\mathcal{H}w(x)$ in $\delta(x)$ that exhibit various forms of consistency, as we will discuss in Theorem 4.

For the applications just mentioned, more machinery is not needed, but for the application in this paper, an explicit rate of convergence is needed between w(x) and its substitute. We will address this rate in the next subsection, but first we need to refine Theorem 2 so that it includes precise convergence rates.

THEOREM 3 (Local Ledoit–Péché Laws). Assume (H1)-(H5) and z is bounded in \mathbb{C}^+ . Then we have the following analogue of (4) from Theorem 1:

(12)
$$\Theta_n(z) = \Theta_{\infty}(z) + O_{\prec}\left(\frac{1}{n\eta}\right)$$

as $n, p \to \infty$. Furthermore (the "small-scale laws,") we have the following analogue of (6) from Theorem 1:

(13)
$$\sup_{(a,b)\subset\mathbb{R}} |\nu_n(a,b) - \nu_\infty(a,b)| \prec \frac{1}{n}$$

and, uniformly in $f \in C^2$,

(14)
$$\int f \, d\nu_n = \int f \, d\nu_\infty + O_{\prec} \left(\frac{\|f\|_1}{n} + \frac{\|f'\|_1}{n} + \frac{\|f''\|}{n^2} \right),$$

as $n, p \to \infty$.

PROOF. Equation (12) is proved in Appendix A, and Equations (13) and (14) are proved in Appendix B. \Box

4.2. Ledoit–Wolf Approximation. In this subsection, we present the approximations devised by Ledoit and Wolf [LW17a] for the Marčenko–Pastur density w(x) and its Hilbert transform. Upon substitution in the expression for $\delta(x)$, these approximations give rise to a shrinkage function $\tilde{d}_n(x)$ that exhibits uniform convergence in probability to $\delta(x)$ on F. As discussed, this mode of convergence is sufficient to establish asymptotic optimality in the applications of [LW20, RMH21].

Our main refinement of [LW20] is to develop an explicit rate of convergence for $\tilde{d}_n(x)$ to $\delta(x)$, although we study convergence in $L^1(d\mu_n)$ instead of in $L^\infty(d\mu_\infty)$. With this essentially more relaxed convergence mode, we are able to obtain convergence at a rate of $O_{\prec}(n^{-2/3})$, which will be fast enough for standardization and optimization of the SRHTs that will be our focus starting in the next section.

First we describe the main approximation result of [LW20]. As mentioned before, the idea is to replace w(x) and $\mathcal{H}w(x)$ in $\delta(x)$ by suitable substitutes. First, one replaces w(x) everywhere by

(15)
$$\tilde{w}_n(x) := p^{-1} \sum_{i=1}^p k_{ni}(x),$$

where

(16)
$$k_{ni}(x) := \frac{1}{n^{-1/3}\lambda_{ni}} k\left(\frac{x - \lambda_{ni}}{n^{-1/3}\lambda_{ni}}\right)$$

for a suitable bump-like kernel function k with known Hilbert transform K. Second, they replace $\mathcal{H}w(x)$ in $\delta(x)$ by

(17)
$$\mathcal{H}\tilde{w}_n(x) := p^{-1} \sum_{i=1}^p K_{ni}(x),$$

where

(18)
$$K_{ni}(x) := \frac{1}{n^{-1/3}\lambda_{ni}} K\left(\frac{x - \lambda_{ni}}{n^{-1/3}\lambda_{ni}}\right).$$

using invariance properties of \mathcal{H} .

The main theoretical result of [LW20] is then the following, which proves that if these replacements are made in $\delta(x)$, the resulting function consistently estimates $\delta(x)$.

THEOREM 4 (Theorem 4.1 of [LW20]). Assume (H1)-(H4) and $k(x) = \frac{1}{2\pi} \sqrt{(4-x^2)_+}$ and $K(x) = (2\pi)^{-1}(-x + \mathrm{sign}(x)\sqrt{(x^2-4)_+})$, where $(\cdot)_+ = \max\{\cdot, 0\}$. Then, with $\tilde{w}_n(x)$ as in (15) and $\mathcal{H}\tilde{w}_n(x)$ as in (17), the function

$$\tilde{d}_n(x) = \frac{x}{[1 - \phi - \phi \pi x \mathcal{H} \tilde{w}_n(x)]^2 + \phi^2 \pi^2 x^2 \tilde{w}_n(x)^2}$$

has the property that

$$\sup_{x \in F} |\tilde{d}_n(x) - \delta(x)|$$

converges to zero in probability as $n, p \to \infty$.

REMARK 5. The authors actually choose k(x) to be the so-called *Epanechnikov* kernel $3(1-x^2/5)_+/(4\sqrt{5})$ in the main body of their paper, but remark in [LW20, Supplement D.5] that the semicircular kernel above works equally well in theory. We choose the latter since the Epanechnikov kernel appears to require much more care in its implementation to avoid numerical instabilities.

As indicated above, we do not analyze rates of stochastic convergence in the rather strong $L^{\infty}(\mu_{\infty})$ norm that appears in the above theorem since $L^{1}(\mu_{n})$ convergence will suffice for our purposes. The following theorem provides an explicit rate for this type of convergence and is our main contribution the Ledoit and Wolf's approximation framework.

THEOREM 5 (Error Analysis of Ledoit–Wolf Eigenvalues in $L^1(\mu_n)$). Making the same assumptions as in Theorem 4, plus (H5), we have

$$p^{-1} \sum_{i=1}^{p} \left| \tilde{d}_n(\lambda_{ni}) - \delta(\lambda_{ni}) \right| \prec \frac{1}{n^{2/3}}$$

as $n, p \to \infty$.

PROOF. See Appendix C.

With this result in hand, we may now return to our application of mean-shift detection.

5. Shrinkage-regularized Hotelling T^2 statistics (SRHTs). In order to discuss the asymptotic properties of our test statistic (3), we must now introduce a model for our test datum $\mathbf{y} = \mathbf{y}_n$. This random vector must, like the \mathbf{x}_{ni} 's, of course be $p \times 1$ and increasing in dimension as n increases. We assume that $\mathbf{y}_n = \boldsymbol{\mu}_n + \boldsymbol{\Sigma}_n^{1/2} \mathbf{z}_n$ for some unknown vector (possibly random) $\boldsymbol{\mu}_n$, where the components of \mathbf{z}_n are i.i.d. $\sim W$ and independent of \mathbf{x}_{ni} . In other words, we assume $\mathbf{y}_n - \boldsymbol{\mu}_n$ and $\{\mathbf{x}_{ni}\}_{i=1}^n$ are jointly i.i.d.

In Subsection 5.1, we find significance levels for the SRHT. Then in Subsection 5.2, we find a surrogate functional for detection performance, identify an (unobservable) shrinkage function that optimizes it in the limit, and present an an empirical shrinkage function that approximates it. Throughout this section we will need the following notion of a regular sequence of functions.

DEFINITION 2. Suppose $f_n \in C^2(\mathbb{R})$ be a sequence for which $|f_n|$ is uniformly bounded for large enough n, and $|f_n'| \prec n^{1/3}$ and $|f_n''| \prec n^{2/3}$. Then we say f_n is regular.

5.1. Significance level. In this subsection, we fix a regular sequence f_n and determine a significance level for $\tilde{T}_n^2 = \tilde{T}_n^2(f_n)$. Suppose we shift and rescale by empirical quantities $\tilde{m}_n p \approx p$ and $\tilde{\sigma}_n \sqrt{p} \approx \sqrt{p}$, obtaining

(19)
$$\hat{Z}_n := \frac{\tilde{T}_n^2 - \tilde{m}_n p}{\tilde{\sigma}_n \sqrt{p}},$$

Performing a binomial expansion on \tilde{T}_n^2 , it can be shown that the mean and variance of \tilde{T}_n^2 are both almost surely of order p and that the quantity $\overline{\mathbf{x}}_n$ can be almost surely replaced by μ_n in the limit. Thus, if we define the following unobservable quantity

$$\hat{Z}_n^o := \frac{\mathbf{z}_n' \tilde{\Sigma}_n \mathbf{z}_n - \tilde{m}_n p}{\tilde{\sigma}_n \sqrt{p}},$$

then $\hat{Z}_n - \hat{Z}_n^o$ converges in distribution to δ_0 , where $\tilde{\Sigma}_n = \Sigma_n^{1/2} f_n(\mathbf{S}_n) \Sigma_n^{1/2}$. Let Z_n^o be the additional unobservable quantity defined as follows

(20)
$$Z_n^o := \frac{\mathbf{z}_n' \hat{\boldsymbol{\Sigma}}_n \mathbf{z}_n - m_n p}{\sigma_n \sqrt{p}},$$

where

$$m_n \equiv m_n(f_n) := p^{-1} \sum_{i=1}^p f_n(\lambda_{ni}) \mathbf{u}'_{ni} \mathbf{\Sigma} \mathbf{u}_{ni}$$

and

$$\sigma_n^2 \equiv \sigma_n^2(f_n) := p^{-1} \operatorname{tr}(f_n(\mathbf{S}_n) \mathbf{\Sigma}_n f_n(\mathbf{S}_n) \mathbf{\Sigma}_n).$$

In the special case where the fourth moment of W matches a standard normal's, this statistic is asymptotically normal by conditioning on \mathbf{X}_n , applying Berry-Esseen, and taking the expectation over the distribution of \mathbf{X}_n . Thus, comparing Z_n^o , \hat{Z}_n^o , and \hat{Z}_n motivates choosing \tilde{m}_n and $\tilde{\sigma}_n$ to consistently approximate m_n and σ_n . Suitable approximations will be given in Lemma 2, but first we need a lemma approximating σ_n using a deterministic limiting form.

LEMMA 1. Assume (H1)-(H5) and let $H:=L^2(F)$. Let $\sigma^2_{\infty}(f,g)$ be a quadratic form on $H\times H$ defined by

(21)
$$\sigma_{\infty}^{2}(f,g) = \int_{F} Tf(x)Tg(x)x\delta(x) d\mu_{\infty}(x),$$

where $T: H \to H$ is defined by

$$Tf(x) = f(x) - \phi \pi \left(\mathcal{H} \left[f d\nu_{\infty} \right](x) - \mathcal{H} \left[d\nu_{\infty} \right](x) f(x) \right). \qquad (x \in F)$$

Then, if f_n is regular we have

(22)
$$\sigma_n(f_n) = \sigma_{\infty}(f_n, f_n) + o_{a.s.}(1)$$

as $n, p \to \infty$.

With these formulas in mind, we now present our choices of $\tilde{\sigma}_n$ and \tilde{m}_n , as well as the resulting asymptotic equivalence of \hat{Z}_n and Z_n^o .

LEMMA 2. Assume (H1)-(H5) and that f_n is regular. Let

$$\tilde{m}_n \equiv \tilde{m}_n(f_n) := p^{-1} \sum_{i=1}^p f_n(\lambda_{ni}) \tilde{d}_n(\lambda_{ni})$$

and

(23)
$$T_n f(x) := f(x) - \frac{1}{n} \sum_{j=1}^{p} (f(\lambda_{nj}) - f(x)) \tilde{d}_n(\lambda_{nj}) K_{nj}(x)$$

and

(24)
$$\tilde{\sigma}_n^2 \equiv \tilde{\sigma}_n^2(f_n) := p^{-1} \sum_{i=1}^p \left[T_n f_n(\lambda_{ni}) \right]^2 \lambda_{ni} \tilde{d}_n(\lambda_{ni}).$$

Then, with \hat{Z}_n and Z_n^o as defined in (19) and (20), $\hat{Z}_n - Z_n^o$ converges in distribution to δ_0 as $n, p \to \infty$.

PROOF. It follows from Theorems 3 and 5 together is that we have

$$\frac{|m_n(f_n) - \tilde{m}_n|p}{\sqrt{p}} = \sqrt{p}|m_n(f_n) - \tilde{m}_n| \prec \sqrt{p} \left(\frac{||f_n||}{n} + \frac{||f_n'||}{n} + n^{-4/3}\right) + \frac{\sqrt{p}}{n^{2/3}} \prec \frac{1}{n^{1/6}}$$

as $n \to \infty$. Thus, as long as $\tilde{\sigma}_n$ is a consistent approximation of $\sigma_n(f_n)$ our choice of \tilde{m}_n is suitable. The consistency of $\tilde{\sigma}_n$ and $\sigma_n(f_n)$ follows from the formal similarity of (24) and (21) and the analysis of Appendix E.

It remains to calculate the asymptotic significance level of either Z_n^o or \hat{Z}_n . It follows from the Hanson–Wright inequality [RV13] and the fact that $\|\tilde{\Sigma}_n\| = \|\Sigma_n f_n(\mathbf{S}_n)\|$ that, almost surely,

(25)
$$\Pr\left[|\mathbf{z}_{n}'f_{n}(\mathbf{S}_{n})\mathbf{z}_{n} - m_{n}p| > \tau \mid \mathbf{X}_{n}, \mathcal{H}_{0}\right]$$

$$\leq 2 \exp\left[-c \min\left(\frac{\tau^{2}}{C^{4}\sigma_{n}^{2}p}, \frac{\tau}{C^{2} \left\|\mathbf{\Sigma}_{n}f_{n}(\mathbf{S}_{n})\right\|}\right)\right]$$

for an absolute known constant c and some C depending only on the moments of W. Scaling τ by $\sigma_n\sqrt{p}$, this upper bound can naturally be used to set a nontrivial asymptotic bound on the false-alarm rate, since the second term in the minimum then goes to infinity like \sqrt{p} , almost surely. By the defintion of Z_n^o , then, we have:

(26)
$$\Pr[Z_n^o > \tau \mid \mathbf{X}_n, \mathcal{H}_0] \le 2 \exp(-c\tau^2/C^4) + o_{a.s.}(1).$$

Taking the expectation with respect to the distribution of X_n and using Lemma 2 gives the following main result for this subsection.

THEOREM 6. Assume (H1)-(H5), that f_n is regular, and that c and C are the known constants defined above. Then

$$\Pr\left[\hat{Z}_n > \tau \mid \mathcal{H}_0\right] \le 2\exp(-c\tau^2/C^4) + o(1)$$

as $n, p \to \infty$.

REMARK 6. Consider again the case where the fourth moment of W matches that of a standard normal. In this case, Theorem 6 can be strengthened by saying that \hat{Z}_n is asymptotically standard normal under \mathcal{H}_0 , since, as mentioned earlier, the same is true of Z_n^o . For numerical evidence of this claim, see Figure 2.

5.2. Detection power and optimal shrinkage. Assume again that f_n is a regular sequence. Further, assume now that \mathcal{H}_1 is in force and consider the detection rate. By expanding quadratic forms, the asymptotic variance of \tilde{T}_n^2 conditioned on \mathbf{X}_n and $\boldsymbol{\mu}_n$ is given by

(27)
$$\mathbb{E}[W^4] \sum_{i} (\tilde{\Sigma}_n)_{ii}^2 + 3 \sum_{i \neq j} (\tilde{\Sigma}_n)_{ij} + 4 \mu'_n f_n(\mathbf{S}_n) \mathbf{\Sigma} f_n(\mathbf{S}_n) \boldsymbol{\mu}_n.$$

Since large values of $\|\boldsymbol{\mu}_n\|_2$ would trivialize the hypothesis test, we assume for now that $\|\boldsymbol{\mu}_n\|_2 = o_{\text{a.s.}}(\sqrt{p})$, which allows us to ignore the term containing $\boldsymbol{\mu}_n$ above. Thus, \tilde{T}_n^2 has the same asymptotic variance under \mathcal{H}_1 and \mathcal{H}_0 . Arguing similarly, define

(28)
$$\mathcal{U}_n \equiv \mathcal{U}_n(f_n) := \frac{\boldsymbol{\mu}'_n f_n(\mathbf{S}_n) \boldsymbol{\mu}_n}{\tilde{\sigma}_n \sqrt{p}},$$

the difference between \hat{Z}_n under \mathcal{H}_1 and \mathcal{H}_0 is given by

$$\frac{2\mathbf{z}_n'\boldsymbol{\Sigma}_n^{1/2}f_n(\mathbf{S}_n)\boldsymbol{\mu}_n}{\tilde{\sigma}_n\sqrt{p}}+\mathcal{U}_n,$$

and and the co-term of \mathcal{U}_n can be neglected due to the asymptotic assumption on $\|\boldsymbol{\mu}_n\|_2$ and another application of Hanson–Wright. Thus, the mean of \hat{Z}_n conditioned on \mathbf{X}_n and $\boldsymbol{\mu}_n$ is \mathcal{U}_n larger under \mathcal{H}_1 than under \mathcal{H}_0 .




Fig 2: A plot showing \hat{Z}_n of (19) to be roughly standard normal when both the test datum y and the training data $\{\mathbf{x}_i\}$ are a coloring matrix times a vector of uniform random variables whose first four moments match a standard Gaussian's. Here, the dimension is 200, the population covariance is the sum of a matrix with Unif[0,1] eigenvalues and rank-40 matrix with eigenvalues $10^{(40-j)/10}$ for j=0,...39.

As a result, $U_n - \hat{Z}_n$ is subgaussian under \mathcal{H}_1 , so that using Hanson-Wright again as in (25), we have

(29)
$$\Pr\left[\hat{Z}_{n} > \tau \mid \mathbf{X}_{n}, \boldsymbol{\mu}_{n}, \mathcal{H}_{1}\right]$$

$$\geq 1 - \Pr\left[\left|\mathcal{U}_{n} - \hat{Z}_{n}\right| \geq (\mathcal{U}_{n} - \tau)_{+} \mid \mathbf{X}_{n}, \boldsymbol{\mu}_{n}, \mathcal{H}_{1}\right]$$

$$\geq 1 - 2\exp\left(-c(\mathcal{U}_{n} - \tau)_{+}^{2}/C^{4}\right) + o_{a.s.}(1)$$

as $n, p \to \infty$. Moreover, taking the expectation over the distributions of μ_n and \mathbf{X}_n gives

(30)
$$\Pr\left[\hat{Z}_n > \tau \mid \mathcal{H}_1\right] \ge 1 - 2\mathbb{E}\exp\left(-c(\mathcal{U}_n - \tau)_+^2/C^4\right) + o(1).$$

Crucially, any almost-sure limit of \mathcal{U}_n appears to be a surrogate for detection performance. As a result, a reasonable goal appears to treat $\mathcal{U}_n(f_n)$ as a *utility functional* and choose f_n to almost-surely maximize it.

REMARK 7. Although the above argument only shows \mathcal{U}_n to be a meaningful surrogate when it exceeds τ , with a bit of algebra it can be shown that the same surrogate arises whenever the fourth moment of W does not exceed a standard normal's, and that this surrogate is meaningful for all values of τ in that case. In particular, the same surrogate arises when the fourth moment of W equals a standard normal's, in which case $\mathcal{U}_n - \hat{Z}_n$ is asymptotically standard normal, as before. Thus, there are cases of interest in which our detection-power surrogate is a reasonable objective functional regardless of the value of τ .

Since the direction in \mathbb{R}^p of the signal μ_n is unknown, it is not possible to analyze the utility functional \mathcal{U}_n without additional information. To rectify this shortcoming, one often assumes some sort of prior on μ_n . For example, in [LAP+20b], the authors assume $\mu_n \sim \mathcal{N}(0,\Omega_n)$ with Ω_n being some properly scaled monomial function of Σ_n . In this paper, we allow for more general dispersion matrices, but require a rate of convergence, as follows.

ASSUMPTION (Assumption (H6)). Let μ_n be a random vector independent of \mathbf{z}_n for which $\mathbb{E}[\mathbf{y}_n \mid \boldsymbol{\mu}_n] = \boldsymbol{\mu}_n$, and let $\boldsymbol{\mu}_n \sim \mathcal{N}(0, \boldsymbol{\Omega}_n)$ for some $\|\boldsymbol{\Omega}_n\| \asymp n^{1/2}$. Further, assume there is a measure ω such that $d\omega/d\mu_\infty$ is C^1 on some neighborhood of F and, for $\eta = \operatorname{Im} z > 0$,

(31)
$$p^{-3/2}\operatorname{tr}\left(\mathbf{\Omega}_{n}\mathbf{R}_{n}(z)\right) = \int \frac{d\omega(x)}{x-z} + O_{\prec}\left(\frac{1}{n\eta}\right).$$

REMARK 8. By inspection, assuming $\|\Omega_n\| \approx n^{1/2}$ ensures that $\|\mu_n\|^2 \approx n^{1/2}$ and thus that \mathcal{U}_n is of order 1, which is precisely the condition that ensures the hypothesis test is tractable but not trivially so in the limit. By Theorem 1 and 3, two suitable examples of Ω_n are $p^{1/2}\mathbf{I}_p$ and $p^{1/2}\mathbf{\Sigma}_n$. The signal prior corresponding to the former example is the ignorant prior, which is commonly applicable in cases where preexisting signal information is absent. If the example of $p^{1/2}\mathbf{\Sigma}_n$ is applicable, then the signal tends to be obscured not just by noise but by the strongest sources of background interference, which results in a more challenging detection problem.

In the following theorem, we will show that under this new assumption (H6), the utilities $\mathcal{U}_n(f_n)$, like the variances $\sigma_n(f_n)$ from the previous subsection, can be approximated using a deterministic form. Further, this deterministic form can be optimized by solving a variational problem explicitly terms of Hilbert transforms—a fact that will be highly useful for our eventual goal of approximately optimizing the utility for finite sample sizes. Making the important definitions

$$g(x) := 1 - \phi - \phi \pi x \mathcal{H} w(x)$$
 and $G(x) := -\phi \pi x w(x)$,

we make our claim precise as follows.

THEOREM 7. Assume (H1)-(H6). Then

(a) (Deterministic Limit.) Suppose f_n is regular. Then

$$\mathcal{U}_n(f_n) = \mathcal{U}(f_n, \omega) + o_{a.s.}(1),$$

as $n \to \infty$, where

$$\mathcal{U}(f,\omega) = \frac{\int f d\omega}{\sigma_{\infty}(f,f)}.$$

(b) (Optimal Limiting Shrinkage.) If $\omega'(x) = d\omega(x)/dx$ and $\Omega(x)$ is the Hilbert transform $\mathcal{H}[\omega'](x)$ and $\mathrm{id}(x) \equiv x$, then $f \in L^2(F) \mapsto \mathcal{U}(f,\omega)$ is maximized by the function

(32)
$$f^* := \frac{g^2 \omega' + gG\Omega}{\delta w \operatorname{id}} - \mathcal{H} \left[\frac{G^2 \Omega + Gg\omega'}{\delta w \operatorname{id}} \right].$$

PROOF. See Appendix F for part (a) and Appendix G for part (b).

The main thing that remains is constructing f_n as a suitable approximation to the optimizer f^* of $\mathcal{U}(\cdot,\omega)$, which is accomplished by the following theorem.

THEOREM 8. Assume (H1)-(H6). Let

$$\tilde{q}_n(x) = 1 - \phi - \phi \pi x \mathcal{H} \tilde{w}_n(x)$$

and

$$\xi_{nj} = \frac{\tilde{g}_n(\lambda_{nj})^2 \mathbf{u}'_{nj} \mathbf{\Omega}_n \mathbf{u}_{nj} - \phi \pi \lambda_{nj} \tilde{g}_n(\lambda_{nj}) p^{-1} \sum_{i=1}^p \mathbf{u}'_{ni} \mathbf{\Omega}_n \mathbf{u}_{ni} K_{ni}(x - \lambda_{ni})}{\tilde{d}_n(\lambda_{nj}) \lambda_{nj}}$$

and

$$\eta_{nj} = -\frac{\phi \pi \lambda_{nj}}{\tilde{g}_n(\lambda_{nj})} \xi_{nj}$$

Further, let $f_n(\mathbf{S}_n)$ be the precision shrinkage estimator that maps the sample eigenvalue λ_{ni} to

(33)
$$\xi_{ni} - p^{-1} \sum_{j=1}^{p} \eta_{nj} K_{nj} (\lambda_{ni} - \lambda_{nj}).$$

Then f_n is regular and satisfies

(34)
$$\lim_{n\to\infty} \mathcal{U}_n(f_n) = \mathcal{U}(f^*,\omega). \qquad (a.s.)$$

As a result, the lower bound on detection power in (30) is maximized by f_n and almost-surely asymptotically equivalent to

(35)
$$1 - 2\exp\left(-c(\mathcal{U}_n(f_n) - \tau)_+^2/C^4\right).$$

PROOF. The vectors ξ_{ni} and η_{ni} correspond to $(g^2\omega' + gG\Omega)/(\delta w \mathrm{id})$ and $G^2\Omega + Gg\omega')/(\delta w \mathrm{id})$ evaluated at λ_{ni} , respectively. See Appendix H for details. The last statement and (35) arise from (29) after taking expectations and almost-sure limits.

6. Empirical results. In this section we present a number of numerical tests of our proposed shrinkage algorithm. One set of tests is performed in the somewhat sterile environment where we choose the population covariance matrix and generate simulated data satisfying the moment conditions in (H1)-(H6). Another set of tests concerns the real-world "Crawdad umich/rss" data set [HIPS22]. In the real-world set, a sensor network collects a time series of received signal strengths during periods of activity and inactivity in a lab setting. In this case, the population covariance matrix and data model are, of course, unknown, but the truth information of when people entered and exited the lab is known.

In the simulated environment, we find that our algorithm outperforms the competitors, which we will describe below—sometimes by significant margins. For the crawdad data, we imagine that there are significant dependencies and inhomogeneities between the samples, making the number of independent samples different from the number of training samples, apparently corrupting some of the detectors. Nevertheless, our detector performs as well as the best competitor in our tests.

Before we discuss results further, we describe the alternative algorithms we chose as comparators.

6.1. Competing algorithms.

6.1.1. BS96 and CQ10. In the context of broader two-sample testing, where two samples have mean $\overline{\mathbf{x}}_1$ and $\overline{\mathbf{x}}_2$, Bai and Saranadasa [BS96] suggest replacing Hotelling's T^2 with $\|\overline{\mathbf{x}}_1 - \overline{\mathbf{x}}_2\|^2$, effectively using the trivial shrinkage estimator of $f(\mathbf{S}_n) = \mathbf{I}_p$, under the "condition-number" assumption that $\|\mathbf{\Sigma}_n\|^2 \ll \operatorname{tr}(\mathbf{\Sigma}_n^2)$ holds. This requirement does indeed hold in our model since we assume uniform spectral boundedness of $\mathbf{\Sigma}_n$; however, if the limiting condition number of $\mathbf{\Sigma}_n$ is even somewhat large, the assumption may not become

relevant until p is enormous, resulting in slow convergence. Nevertheless, the detector has the desirable qualities that, asymptotically, it has known constant false-alarm rate and predictable detection power.

Similar to BS96, but also suitable to the case of $p \gg n$, Chen and Qin [CQ10] propose a test that is asymptotically optimal given a similar constraint on the condition number of Σ_n . To be precise, the covariance matrix is required to satisfy the constraint $\operatorname{tr}(\Sigma_n^4) \ll \operatorname{tr}^2(\Sigma_n^2)$. Their test statistic for samples $\{\mathbf{x}_{1i}\}_{i=1}^{n_1}$ and $\{\mathbf{x}_{2i}\}_{i=1}^{n_2}$ is

$$\frac{\sum_{i\neq j}^{n_1} \mathbf{x}'_{1i}\mathbf{x}_{1j}}{n_1(n_1-1)} + \frac{\sum_{i\neq j}^{n_2} \mathbf{x}'_{2i}\mathbf{x}_{2j}}{n_2(n_2-1)} - \frac{2\sum_{i=1}^{n_1} \sum_{j=1}^{n_2} \mathbf{x}'_{1i}\mathbf{x}_{2j}}{n_1n_2},$$

Although this test statistic, BS96, and several other algorithms we describe are applicable to values of n_1 and n_2 that both differ from 1, we will always assume that $n_2=1$ for the sake of comparison with ours. In our simulations, we do not test the ultra-high-dimensional regime or the case where the condition number diverges, apparently making CQ10 perform equivalently to BS96. As a result, in the plots that follow we will show only CQ10 to represent both algorithms.

6.1.2. LAPPW20. Li et al. [LAP+20b] assume instead that Σ_n follows the well-known spiked covariance model of Johnstone [Joh01], i.e., that the all the eigenvalues of Σ_n are equal to one except for a fixed finite number that are larger. While this assumption may not clash with the condition-number assumptions of BS96 and CQ10 in the limit, a spiked matrix with large condition could require an extremely large value of p before those algorithms begin to exhibit the type of performance they asymptotically guarantee. In LAPPW20, the "pooled" sample covariance matrix S_n (equivalent to our S_n when $n_2 = 1$) is replaced by the linear shrinkage estimator of the form $S_n + \lambda I_p$, for some data-dependent $\lambda > 0$. The constant λ is chosen to locally maximize an asymptotic utility function similar to ours, where the signal prior is $I_p, \Sigma_n, \Sigma_n^2$, or some linear combination thereof. (Higher-order polynomials in Σ_n can be handled similarly.) The proposed method of maximization is to use a grid search from $p^{-1}\text{tr}(S_n)$ to $20 ||S_n||$ to maximize a spiked analogue of $\mathcal{U}(f,\omega)$ for $f(x) = x + \lambda$.

In our implementation, we only consider signal priors of I_p and Σ_n , and we perform a log-scale grid search on the designated interval consisting of 1×10^6 points.

Like BS96 and CQ10, LAPPW20 has a known asymptotically constant false-alarm rate and predictable detection power. Further, Li et al. remark that LAPPW20 reduces to BS96 for population covariance matrices that have condition number sufficiently close to one.

- 6.1.3. LW22. Based Ledoit and Wolf's work in [LW22], which is a refinement of earlier work [LW20], we choose to replace the sample covariance matrix with the "quadratic inverse shrinkage" estimator using code provided by the authors in their appendix. Although the algebraic form of their covariance estimator appears slightly different compared to the one in their earlier work and our Theorem 5, it continues to satisfy the conclusions of Theorem 4, differing mainly in the choice of kernel function k to be the Cauchy density $1/(\pi(1+x^2))$.
- 6.1.4. CWH11. The mean-shift detector of Chen et al. [CWH11] is applied in a similar context to ours, where two samples are tested but one sample is just a singleton. Their model is a somewhat different generalization of the Gaussian one, though. They assume a spherically-invariant-random-vectors, or elliptical, data model, rather than a separable-covariates model as we do. The Marčenko–Pastur Theorem happens to be valid in both settings, so it would not be a surprise if their work extends to our model and vice versa.

Fundamentally, CWH11 is an extension of the well-known covariance M-estimation algorithm of Tyler [Tyl87] to the sample-starved case of n < p, by formulating and iteratively

solving a fixed-point equation. (Thus, in the plots we will show, CWH11 is equivalent to the Tyler estimator.) This linear shrinkage is shown to outperform Ledoit and Wolf's *linear* shrinkage algorithm [LW04], for example, on a supervised detection task using the crawdad data set.

- 6.1.5. Proposed algorithm. The proposed algorithm is essentially just the approximation to f^* described in Theorem 8. The only modification we make is to replace every shrunken eigenvalue by its max with 0. Since f^* is asymptotically non-negative, this does not impact consistency and can only accelerate convergence.
- 6.2. Results for synthetic subgaussian data. For our first series of experiments, we tested our detector on artificially generated training and test data sets in Matlab. The data generation process was similar to that in [RML+22]. First, let p=200, choose a signal prior covariance Ω and signal magnitude γ , choose the number of training samples n, and generate a $p \times p$ covariance matrix Σ with piece-wise log-linear eigenvalues $\{10^{\kappa i/40}\}_{i=1}^{40} \cup \{10^{(i-1)/(40(p-41))}\}_{i=1}^{p-40}$ for some $\kappa \geq 1$. (See Figure 3 for a scree plot with $\kappa = 2$.) Next, perform 100 Monte Carlo trials as follows:
- Generate an i.i.d. $p \times n$ matrix \mathbf{Z} with uniformly distributed components of mean 0 and variance 1. Let $\mathbf{X} = \mathbf{\Sigma}^{1/2} \mathbf{Z}$, the training-data matrix.
- Generate a large number test observations, some signal-free ones distributed as the columns of \mathbf{X} , and some signal-containing ones, which are signal-free observations added to $\gamma \mathbf{z}/\|\mathbf{z}\|$, where $\mathbf{z} \sim \mathcal{N}(0, \mathbf{\Omega})$.
- Compute S_n as in (2) and generate detection scores for each algorithm in the last section and each test observation.

Once detection scores have been generated for all 100 trials, they are assembled into approximate receiver-operator-characteristic (ROC) curves of size versus power by counting the number of \mathcal{H}_0 and \mathcal{H}_1 observations that exceed each of a large number of thresholds.

Figures 4 and 5 show the cases of $\Omega = I$, n = 300, and $\kappa = 1, 2, 4$, and 8.

- REMARK 9. We note that essentially equivalent results can be observed in the more nearly spiked model where the trailing 160 eigenvalues of Σ are chosen to be 1, suggesting that our results may extend beyond the model of (H5) to the spiked model. Nevertheless, to demonstrate the validity of our results under the conditions we have given, we choose instead to conduct tests in which (H5) is more closely approximated.
- 6.3. Results for measured data. In addition to testing our algorithm on synthetically generated subgaussian data, we also test our algorithm on real, measured sensor-network data. The data collect to which we apply our method is the crawdad umich/rss one [HIPS22], in which 14 Mica2 sensors were distributed throughout a lab space to detect whether a person was present and/or moving there. In order to do so, the sensor network collected and recorded received signal strength (RSS) measurements for each sender-receiver pair of sensors, totaling $p=14\cdot 13=182$ measurements at each time instant. Whether or not there was activity in the room was known in the experiment and can be used to determine the performance of a detection algorithm applied to the data. RSS measurements were collected over 3127 time instants, spaced 0.5 seconds apart, with 327 time instants corresponding to activity. In addition to human activity, RSS measurements were weakened/disturbed by background interference and noise due to cellular, wifi, and radio signals, and other nuisance sources of EM radiation. Further, the data are de-trended in the same way as in the simulations of [CWH11].





Fig 3: Scree plot of population covariance matrix chosen to generate artificial data, $\kappa = 2$.

In contrast to our tests in the previous section, one does not know a priori the signal prior for the crawdad data collect. A natural guess might be the ignorant prior of $\Omega \propto I_p$. However, this choice appears to result in fairly poor performance, suggesting that the true prior distribution of signals is not completely isotropic. A more conservative assumption, accounting for a much more challenging type of signal behavior, would be the assumption that $\Omega \propto \Sigma$, the unknown population covariance matrix. This would mean that signals indicating human activity are well-masked by interfering signals, with the likelihood of a human signal in a given direction being correlated to the strength of an interfering signal in that direction. This assumption could of course be made even more conservative if replaced by $\Omega \propto \Sigma^k$ for some $k \geq 2$, for example, but for simplicity we do not concern ourselves with such assumptions here.


Assuming that $\Omega \propto \Sigma$ means that, according to (31), in the limit $\omega(x)$ should be equal to $\delta(x)$. We also calibrate LAPPW20 so that the signal prior of $\Omega \propto \Sigma$ is assumed, as in [LAP+20b].

Our experimental procedure is to pick 1000 samples of size n from the 3127-327=2800 inactive time indices, and for each sample test the remaining 3127-n time indices, generating a detection statistic for each using our algorithm and the comparators from the last section. We then generate an empirical ROC curve for each algorithm by plotting how many detection statistics for inactive and active time indices exceed a sliding threshold relative to the totals of 2800 and 327, respectively. Our results for n=200 and n=300 are shown in Figure 6.

The most noticable feature of this figure is that our algorithm roughly ties with LW22. Similar experiments with synthetic data show the same pattern, which is intriguing since the eigenvalues of LW22 differ significantly from ours—in particular, since many of ours are zero. This trend suggests that the class of nearly optimal estimators is fairly flexible in some settings. An interesting avenue for future research would be to explore whether this flexibility can be exploited to improve the mean and/or variance of our algorithm's finite-sample performance.




(a) LW22 and Proposed perform similarly, and CQ10 dominates the others due to relatively low condition number.




(b) Proposed outperforms LW22 since $\Sigma \approx I$ is now less accurate. LAPPW20 now outperforms the others, likely due to the increasing condition number and relatively more spikey nature of the spectrum. (See Figure 3.)

Fig 4: Gap between Proposed and alternatives increases with κ .



(a) Proposed dominates LW22 to an even greater degree, likely for the same reason as in Figure 4b. CWH11 outperforms the others, including LAPPW20, likely due to its stringent search requirements for a regularization factor.



(b) Proposed dominates perhaps to an even greater degree. CWH11, LAPPW20, and CQ10 all lie on the chance line, possibly due to the high condition number and/or numerical instabilities.

Fig 5: Proposed dominates for high condition number κ .




Fig 6: Our detector approximately ties with LW22 and Hotelling for best on the crawdad data set for n=300 and 200. (Hotelling not shown for n=300 due to tie.) The lower performance of CQ10 and LAPPW20 suggests that the training data are not i.i.d. and/or do not have a common spiked or well-conditioned covariance matrix.

Acknowledgments. This work was supported by the United States Air Force Sensors Directorate and AFOSR Lab Tasks 18RYCOR004, 19RYCOR03 and 24RYCOR006. However, the views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of any agency of the U.S. government. Examples of analysis performed within this article are only examples. Assumptions made within

the analysis are also not reflective of the position of any U.S. Government entity. The Public Affairs approval number of this document is AFRL-2024-6552.

APPENDIX A: CONVERGENCE RATE OF THE LEDOIT-PÉCHÉ ANALYTIC FUNCTION

First we make an important remark about notation.

REMARK 10. In this appendix and the following ones, some of the formulas will involve many key quantities that depend on n, which could result in an excessive number of subscripts. As a result, we will follow the convention of [KY17] and typically omit indices of n, so that all quantities not expressly deemed constant (as the measure π_{∞} is) may have a hidden dependence on n. Thus, we abbreviate $m_n(z)$ by m(z), Σ_n by Σ , S_n by S, the sample resolvent $R_n(z)$ by R(z), etc; and statements such as $m(z) \to m_{\infty}(z)$ (a.s.) should cause no confusion. Having made this note, however, we will still occasionally use indices of n for emphasis or clarity.

The result about $\Theta(z) \equiv \Theta_n(z)$ will follow by carefully keeping track of the estimation errors in [LP11, Lemma 2]. Their result is principally a study of $p^{-1}\text{tr}(\mathbf{SR}(z))$, due to the resolvent identity

$$p^{-1}\operatorname{tr}(\mathbf{SR}(z)) = 1 + zm(z),$$

where we have suppressed three subscripts of n. From this identity, if one could say that $p^{-1}\mathrm{tr}(\mathbf{SR}(z))$ is in some sense approximately $p^{-1}\mathrm{tr}(\mathbf{\Sigma R}(z)) = \Theta($, almost-sure convergence of the right-hand side would follow from almost-sure convergence of m(z) and the fact that Σ is spectrally bounded.

In actuality, the two normalized traces do not converge to each others' limits in our asymptotic regime, but they are functionally related. The right-hand side above is, by the averaged local Marčenko–Pastur law, the same as $1+zm_{\infty}(z)$, up to error $O_{\prec}\left(n^{-1}\eta^{-1}\right)$. Further that quantity, by [LP11, Lemma 2] can be expressed as

(36)
$$1 + z m_{\infty}(z) = \frac{\phi \Theta_{\infty}(z)}{\phi + \phi^2 \Theta_{\infty}(z)}.$$

Thus,

(37)
$$p^{-1}\operatorname{tr}(\mathbf{SR}(z)) = \frac{\phi\Theta_{\infty}(z)}{\phi + \phi^{2}\Theta_{\infty}(z)}.$$

Our approach will be to study the left-hand side, relate it to $\Theta(z)$, and use that relationship to make and explicit comparison with $\Theta_{\infty}(z)$.

First, we need a lemma.

LEMMA 3. Assume (H1)-(H5). Then there is are constants A, a > 0 such that, almost surely, for all bounded $z \in \mathbb{C}^+$ and sufficiently large n

$$(38) a \le |1 + \phi\Theta(z)| \le A.$$

PROOF. The idea is to use the fact that the limit as $\eta \to 0^+$ of (A) is $\delta(x)/x$, as defined in Theorem 2, then to use that theorem to bound it. In addition, we use the asymptotic equivalence of $\Theta(z)$ and $\Theta_{\infty}(z)$ that theorem.

By [PWY23] and the fact that the spectral density is piecewise C^{∞} , the function $m_{\infty}(x)$ is smooth except possibly at finitely many points, and $\delta(x)$ is therefore continuous and bounded.

In addition, by Theorem 2, $\delta(x)$ can only exceed $\|\Sigma\| \ge \max \mathbf{u}_i^* \Sigma \mathbf{u}_i$ on a set of measure zero. Thus, $\delta(x) \le \|\Sigma\|$ on all of F, which we recall is $\operatorname{supp} \mu_{\infty} \setminus \{0\}$. Thus, we get

$$(39) |1 - \phi - \phi x \check{m}(x)|^2 \ge \frac{\min F}{\|\Sigma\|}$$

for all $x \in E$. Using the fact that $|m_{\infty}(z) - \breve{m}(x)| \leq \eta$, we get that $|zm_{\infty}(z) - x\breve{m}(x)| \leq \eta$ and for small enough η ,

$$(40) |1 - \phi - \phi z m(z)| \ge 1/A$$

for some large enough A.

Similarly, we may upper bound the function in (39) by $(\max F) \|\Sigma^{-1}\|$, so that (40) can be augmented to

(41)
$$1/a \ge |1 - \phi - \phi z m(z)| \ge 1/A$$

for some sufficiently small a > 0.

Finally, we get the following additional identity from [LP11, Lemma 2], equivalent to (36) above:

$$1 + \phi \Theta_{\infty}(z) = \frac{1}{1 - \phi - \phi z m_{\infty}(z)}.$$

Combining with (41) gives the desired result.

It will be convenient if to define a few quantities related to the sample resolvent. First, let $\mathbf{c}_j = \mathbf{x}_j/\sqrt{n}$, where we recall that \mathbf{x}_j is standing in for \mathbf{x}_{nj} . By (H1), an alternative expression for \mathbf{c}_j is $\mathbf{\Sigma}^{1/2}\mathbf{z}_j/\sqrt{n}$, where \mathbf{z}_j is a vector of i.i.d. zero-mean, variance-one subgaussian random variables. Next, we define $\mathbf{R}^{(j)}(z)$ to be $(\mathbf{S} - z\mathbf{I} - \mathbf{c}_j\mathbf{c}_j')^{-1}$ and $\mathbf{R}^{(kj)}(z) = (\mathbf{S} - z\mathbf{I} - \mathbf{c}_j\mathbf{c}_j' - \mathbf{c}_k\mathbf{c}_k')^{-1}$. Finally, we define $\Theta^j(z) = p^{-1}\mathrm{tr}(\mathbf{\Sigma}\mathbf{R}^{(j)}(z))$.

Our next preliminary result comes from [SB95, Lemma 2.6]:

(42)
$$\frac{1}{p} \left| \operatorname{tr} \left(\mathbf{\Sigma} (\mathbf{R}^{(j)}(z) - \mathbf{R}(z)) \right) \right| \le \frac{\|\mathbf{\Sigma}\|}{p\eta}$$

We also have that $\mathbf{c}_j'\mathbf{R}^{(j)}(z)\mathbf{c}_j$ converges almost surely to $\Theta^j(z)$ by concentration of quadratic forms in subgaussian variables, which by (42), converges almost surely at a rate of $O(n^{-1}\eta^{-1})$ to $\Theta(z)$. Thus, a corollary of Lemma 3 is that there is a > 0 such that

(43)
$$|1 + \phi \Theta^{j}(z)| \ge a \qquad (\text{all } j \le n \text{ and } z \in \mathbb{C}^{+})$$

almost surely for sufficiently large n. Likewise, since $1 + \phi \mathbf{c}'_j \mathbf{R}^{(j)}(z) \mathbf{c}_j$ is one larger than a quadratic form in subgaussian random vectors, it converges almost surely to the left side of (43), and

(44)
$$|1 + \mathbf{c}_j' \mathbf{R}^{(j)}(z) \mathbf{c}_j| \ge a.$$
 (all $j \le n$ and $z \in \mathbb{C}^+$)

almost surely for sufficiently large n.

We continue to follow along with the argument in the appendix of [LP11], as prescribed above. Some algebra shows that

$$1 + zm(z) = 1 + p^{-1} \operatorname{tr}(\mathbf{SR}(z))$$
$$= p^{-1} \operatorname{tr}\left(\sum_{i=1}^{n} \mathbf{c}_{i} \mathbf{c}'_{i} \mathbf{R}(z)\right)$$
$$= p^{-1} \sum_{i=1}^{n} \mathbf{c}'_{i} \mathbf{R}(z) \mathbf{c}_{i}.$$

Using the Woodbury–Morrison identity gives

(45)
$$\mathbf{R}(z) = \mathbf{R}^{(i)}(z) - \frac{\mathbf{R}^{(i)}(z)\mathbf{c}_i\mathbf{c}_i'\mathbf{R}^{(i)}(z)}{1 + \mathbf{c}_i'\mathbf{R}^{(i)}(z)\mathbf{c}_i},$$

so that $p^{-1}\mathrm{tr}(\mathbf{SR}(z))$ becomes

(46)
$$\frac{1}{p} \sum_{i=1}^{n} \frac{\mathbf{c}_{i}' \mathbf{R}^{(i)}(z) \mathbf{c}_{i}}{1 + \mathbf{c}_{i}' \mathbf{R}^{(i)}(z) \mathbf{c}_{i}}.$$

In order to estimate the above, we define the error

$$\epsilon_i := \mathbf{c}_i' \mathbf{R}^{(i)}(z) \mathbf{c}_i - \phi \Theta^i(z).$$

Our next lemma concerns the order of magnitude of ϵ_i .

LEMMA 4. Assume (H1)-(H5). Then $|\epsilon_i| \prec 1/\sqrt{\eta n}$.

PROOF. By Hanson-Wright [RV13],

$$\Pr\left[\left|\epsilon_{i}\right| \geq t \mid \mathbf{R}^{(i)}(z)\right] \leq 2 \exp\left[-c \min\left(\frac{n^{2}t^{2}}{C^{4} \left\|\mathbf{\Sigma}^{1/2}\mathbf{R}^{(i)}(z)\mathbf{\Sigma}^{1/2}\right\|_{\mathrm{HS}}^{2}}, \frac{nt}{C^{2} \left\|\mathbf{\Sigma}\mathbf{R}^{(i)}(z)\right\|}\right)\right],$$

for some C > 0 depending only on the moments of W. Thus, we would like to estimate the Hilbert-Schmidt norm above: namely,

(47)
$$\operatorname{tr}\left(\mathbf{\Sigma}\mathbf{R}^{(i)}(z)\mathbf{\Sigma}\mathbf{R}^{(i)}(z)^{*}\right) \leq \|\mathbf{\Sigma}\|\operatorname{tr}\left(\mathbf{R}^{(i)}(z)\mathbf{\Sigma}\mathbf{R}^{i}(\overline{z})\right)\right)$$

(48)
$$\lessapprox \sum_{k=1}^{p} \frac{\mathbf{u}_{k}' \mathbf{\Sigma} \mathbf{u}_{k}}{|\tilde{\lambda}_{k} - z|^{2}}$$

$$= \frac{p}{i\eta} \left(\Theta^{i}(z) - \Theta^{i}(\overline{z}) \right),$$

where $\tilde{\lambda}_i$ are the eigenvalues of $\mathbf{S} - \mathbf{c}_i \mathbf{c}_i'$. Since, as discussed in Section 4.2, $\Theta^i(z)$ has a limiting value as $z \to x \in \mathbb{R} \setminus \{0\}$, the left-hand side of (47) has order at most n/η as $n \to \infty$, almost surely. It can also be seen that the spectral norm has order at most $1/\eta$. Thus, taking t to be $n^{\epsilon}/\sqrt{\eta n}$ and smoothing the conditional expectation gives the result.

Using Woodbury to expand $\mathbf{R}^{(j)}(z)$ as $\mathbf{R}^{(ij)}(z) = (\mathbf{R}^{(j)}(z)^{-1} - \mathbf{c}_i \mathbf{c}_i')^{-1}$ and applying the ideas of Lemma 4 and (48) again, this time to the squared Frobenius norm of $\Sigma \mathbf{R}^{(ij)}(z)\Sigma \mathbf{R}^{(ij)}(z)^*$, we get an "almost-independence" result for ϵ_i and ϵ_j , the proof of which is omitted:

LEMMA 5. Assume (H1)-(H5) and that $i \neq j$. Then $|\epsilon_i \epsilon_j| \prec n^{-2} \eta^{-2}$.

We continue with our analysis of $p^{-1}\text{tr}(\mathbf{SR}(z))$, returning to (46). Letting f(x)=(x-1)/x, we see that

$$p^{-1}\operatorname{tr}(\mathbf{SR}(z)) = p^{-1} \sum_{i=1}^{n} f(1 + \mathbf{c}_{i}'\mathbf{R}^{(i)}(z)\mathbf{c}_{i}).$$

The mean value theorem tell us $|f(x) - f(y)| \le |x - y|/a^2$ for $x, y \ge a > 0$, so that we can make the following estimate

(49)
$$\left| p^{-1} \operatorname{tr}(\mathbf{S}\mathbf{R}(z)) - p^{-1} \sum_{i=1}^{n} f\left(1 + \phi \Theta^{i}(z)\right) \right|^{2}$$

$$\leq \frac{1}{a^{4}} \left(p^{-1} \sum_{i=1}^{n} |\epsilon_{i}| \right)^{2}$$

$$\lessapprox p^{-2} \sum_{i=1}^{n} |\epsilon_{i}|^{2} + p^{-2} \sum_{i \neq j} |\epsilon_{i} \epsilon_{j}|$$

$$\prec \frac{1}{n^{2} \eta^{2}}.$$

Thus, $p^{-1}\mathrm{tr}(\mathbf{SR}(z))$ can be replaced by $p^{-1}\sum_{i=1}^n f\left(1+\phi\Theta^i(z)\right)$ in our main identity (37), committing only another error of $O_{\prec}\left(n^{-1}\eta^{-1}\right)$. Repeating the argument (49)-(50) for comparing $p^{-1}\sum_{i=1}^n f\left(1+\phi\Theta^i(z)\right)$ and $p^{-1}\sum_{i=1}^n f\left(1+\phi\Theta(z)\right)$, gives yet another error of order $O_{\prec}\left(n^{-1}\eta^{-1}\right)$. Thus, we get

(51)
$$p^{-1} \sum_{i=1}^{n} f\left(1 + \phi\Theta(z)\right) = \frac{\phi\Theta_{\infty}(z)}{\phi + \phi^{2}\Theta_{\infty}(z)}.$$

But the left-hand side above is just

$$\frac{n}{p} \frac{\phi \Theta_n(z)}{1 + \phi \Theta_n(z)},$$

where we have re-added subscripts of n for emphasis. Using $n/p - 1/\phi = O(1/p)$ gives

$$\frac{1}{1 + \phi \Theta_n(z)} = \frac{1}{1 + \phi \Theta_{\infty}(z)} + O_{\prec} \left(\frac{1}{n\eta}\right).$$

Since, by Lemma 3 again, $|1 + \phi\Theta_{\infty}(z)|$ is asymptotically almost surely bounded above and below, this identity is readily inverted to give the desired result.

APPENDIX B: HELFFER-SJÖSTRAND RESULTS

This section closely follows [BGK16, Section 8]. The main result of this section is as follows.

LEMMA 6. Assume (H1)-(H5). Then, uniformly for $f \in C^2(\mathbb{R})$, we have

(52)
$$\int f d\mu - \int f d\mu_{\infty} = O_{\prec} \left(\frac{1}{n} \|f\|_{1} + \frac{1}{n} \|f'\|_{1} + \frac{1}{n^{2}} \|f''\|_{1} \right)$$

Moreover, letting ν be the sample Ledoit-Péché measure and ν_{∞} its limit, we have the same result when μ and μ_{∞} are replaced with ν and ν_{∞}

We have stated Lemma 6 in this way to accommodate general n-dependent functions f. One consequence is the more familiar "law on small scales":

COROLLARY 1. Let $I = [a, b] \subseteq \mathbb{R}$ be an interval. Then uniformly in I,

(53)
$$\mu(I) - \mu_{\infty}(I) = O_{\prec}\left(\frac{1}{n}\right)$$

and

(54)
$$\nu(I) - \nu_{\infty}(I) = O_{\prec}\left(\frac{1}{n}\right)$$

PROOF OF COROLLARY 1. Fix $\epsilon > 0$ and let $\eta = n^{\epsilon - 1}$. Let $f : \mathbb{R} \to [0, 1]$ be C^{∞} , equal 1 on I, equal 0 on $(a - \eta, b + \eta)^c$, and satisfy $\|f'\|_{\infty} \leq C\eta^{-1}$ and $\|f''\|_{\infty} \leq C\eta^{-2}$. Thus f "just barely" dominates the indicator function of I. Lemma 6 implies that

(55)
$$\int f d\mu - \int f d\mu_{\infty} = O_{\prec} \left(n^{-1} \right)$$

so that, using the facts that the density of μ_{∞} is bounded and, by Lemma 3, the density of ν_{∞} is bounded,

$$(56) \qquad \mu(I) \ge \int f d\mu - C n^{-1+\epsilon} = \int f d\mu_{\infty} + O_{\prec} \left(n^{-1} \right) \ge \mu_{\infty}(I) + O_{\prec} \left(n^{-1} \right).$$

Repeating this logic when f is instead "just barely" dominated by the indicator function of I provides the reverse inequality.

PROOF OF LEMMA 6. We proceed under the assumption that f is compactly supported; this only strengthens the bound since the supports of μ and ν are bounded by some fixed constant with high probability.

Let $\hat{\mu} = \mu - \mu_{\infty}$, and let $\hat{m} = m - m_{\infty}$ be the Stieltjes transform of $\hat{\mu}$.

Let $\epsilon > 0$ and $\eta = n^{-1+\epsilon}$.

Let $\chi \in \mathcal{C}^{\infty}(\mathbb{R}, [0,1])$ be a smooth cutoff function with $\chi(0) > 0$. Lastly, let $h \in \mathcal{C}^{\infty}(\mathbb{R}, [0,1])$ be supported on $[a-\eta, b+\eta]$, be identically 1 on [a,b], and satisfy $\|h'\|_{\infty} \leq C\eta^{-1}$, $\|h''\|_{\infty} \leq C\eta^{-2}$.

By Helffer-Sjöstrand formula, we may write the left-hand side of equation (52) as

(57)
$$\int f(\lambda)\hat{\mu}(d\lambda) = \frac{1}{2\pi} \int dx \int dy \left(\partial_x + i\partial_y\right) \left[\left(f(x) + iyf'(x) \right) \chi(y) \right] \hat{m}(x + iy),$$

which upon expanding and using that the left-hand side is real, reads

(58)
$$\int f(\lambda)\hat{\mu}(d\lambda) = -\frac{1}{2\pi} \int dx \int dy f''(x)\chi(y)y \operatorname{Im}\hat{m}(x+iy)$$
$$-\frac{1}{2\pi} \int dx \int dy f(x)\chi'(y) \operatorname{Im}\hat{m}(x+iy)$$
$$-\frac{1}{2\pi} \int dx \int dy f'(x)\chi'(y)y \operatorname{Re}\hat{m}(x+iy)$$

The second and third terms are easiest to bound. Since $|\hat{m}| \leq \eta$ uniformly on the purely complex set $\operatorname{supp} \chi'(y) \subseteq \mathbb{C}$, we bound the second term by

(59)
$$\lessapprox \left| \int dx f(x) \int dy \chi'(y) \operatorname{Im} \hat{m}(x + iy) \right|$$
$$\lessapprox \eta \int dx |f(x)|.$$

Similarly the third term is bounded by

(60)
$$\lessapprox \left| \int dx f'(x) \int dy \chi'(y) y \operatorname{Re} \hat{m}(x + iy) \right|$$
$$\lessapprox \eta \int dx \left| f'(x) \right|.$$

Now, we must bound the first term

(61)
$$-\frac{1}{2\pi} \int dx \int dy f''(x) \chi(y) y \operatorname{Im} \hat{m}(x+iy).$$

First we bound

(62)
$$-\frac{1}{2\pi} \int dx \int_{|y| \le \eta} dy f''(x) \chi(y) y \operatorname{Im} \hat{m}(x + iy).$$

It is a general fact of Stieltjes transforms t of positive measures that

$$(63) y \mapsto y|t(x+iy)|$$

is nondecreasing for y>0 and for any x. Since $\eta|\hat{m}(x+\mathrm{i}\eta)|\leq\eta$ with high probability, and moreover $\eta|m_\infty(x+\mathrm{i}\eta)|\leq\eta$ since the density of μ_∞ is bounded, we find that $\eta|m(x+\mathrm{i}\eta)|\leq\eta$. This general fact may then be applied to both $y|m(x+\mathrm{i}y)|$ and $y|m_\infty(x+\mathrm{i}y)|$, together with $t(\overline{z})=\overline{t(z)}$, to establish

(64)
$$\max_{|y| \le \eta} |y\hat{m}(x+iy)| \le \eta$$

with high probability. Now we may bound (62) by

(65)
$$\lessapprox \int dx \left| f''(x) \right| \int_{|y| \le \eta} dy \cdot \eta \lessapprox \eta^2 \int dx \left| f''(x) \right|$$

with high probability.

Lastly we bound

(66)
$$-\frac{1}{2\pi} \int dx \int_{|y|>\eta} dy f''(x) \chi(y) y \operatorname{Im} \hat{m}(x+\mathrm{i}y).$$

This is more difficult that the previous bound because the simple bound $|f''| \le C\eta^{-2}$ is not affordable any longer. The way around this is to use that $\int dx f''(x)$ has a much tigher bound than $\int dx |f''(x)|$, through the fundamental theorem of calculus. Some care is still required, however, since we must not bound $\int dx f''(x)$ only but the larger expression (66) which has another x-dependent factor in the integrand.

We integrate by parts: first in x, noting that f' vanishes at $\pm \infty$,

$$-\frac{1}{2\pi} \int dx \int_{|y|>\eta} dy f''(x) \chi(y) y \operatorname{Im} \hat{m}(x+iy)$$

$$= \frac{1}{2\pi} \int dx \int_{|y|>\eta} dy f'(x) \chi(y) y \partial_x \operatorname{Im} \hat{m}(x+iy)$$

$$= -\frac{1}{2\pi} \int dx \int_{|y|>\eta} dy f'(x) \chi(y) y \partial_y \operatorname{Re} \hat{m}(x+iy)$$

where in the last line we used the holomorphy of \hat{m} , and then integrating over y,

$$(68)$$

$$= -\frac{1}{2\pi} \int dx \int_{|y|>\eta} dy f'(x) \chi(y) y \partial_y \operatorname{Re} \hat{m}(x+\mathrm{i}y)$$

$$= -\frac{1}{2\pi} \int dx f'(x) \left(\left[y\chi(y) \operatorname{Re} \hat{m}(x+\mathrm{i}y) \right]_{\eta}^{-\eta} - \int_{|y|>\eta} dy (\chi(y) + y\chi'(y)) \operatorname{Re} \hat{m}(x+\mathrm{i}y) \right)$$

Continuing,

$$\begin{aligned} |\cdot| &\lessapprox \int dx \left| f'(x) \right| \left| O(\eta) - \int_{|y| > \eta} dy (\chi(y) \operatorname{Re} \hat{m}(x + \mathrm{i}y) + y \chi'(y) \operatorname{Re} \hat{m}(x + \mathrm{i}y)) \right| \\ &\lessapprox \int dx \left| f'(x) \right| \left(O(\eta) + \int_{|y| > \eta} dy \chi(y) \left| \operatorname{Re} \hat{m}(x + \mathrm{i}y) \right| + \int_{|y| > 1} y \left| \operatorname{Re} \hat{m}(x + \mathrm{i}y) \right| \right) \\ &\lessapprox \int dx \left| f'(x) \right| \left(O(\eta) + \int_{\eta}^{1} dy \frac{\eta}{y} + O(\eta) \right) \\ &\lessapprox \int dx \left| f'(x) \right| \left(O(\eta) - \eta \log \eta \right) \\ &\lessapprox \eta \log \eta \int dx \left| f'(x) \right| \end{aligned}$$

with high probability, where we have used (64) in the middle step. Thus, we have established equations (66), (61), and ultimately, (52).

The proof is exactly the same when μ, μ_{∞} are replaced with ν, ν_{∞} .

APPENDIX C: L^1 ERROR IN THE LEDOIT-WOLF EIGENVALUES

In what follows, using our convention of suppressing subscripts of n, $L^1(\mu)$ will denote the n-dependent norm $L^1(\mu_n)$, and $\tilde{w}(x)$ will denote $\tilde{w}_n(x)$, for example.

The goal is to approximate $\delta(x) = x/|1 - \phi - \phi x \check{m}(x)|^2$ in the $L^1(\mu)$ -norm. Since the denominator $|1 - \phi - \phi x \check{m}(x)|^2$ is bounded away from zero for x in a neighborhood of F by Lemma 3 and $\check{m}(x) = \pi \mathcal{H}w(x) + i\pi w(x)$, it suffices to show $\|\mathcal{H}\tilde{w} - \mathcal{H}w\|_{L^1(\mu)}$ and $\|\tilde{w} - w\|_{L^1(\mu)}$ are both $O_{\prec}(n^{-2/3})$. Throughout this and later appendices, Δ will stand for $n^{-1/3}$.

For the sake of brevity, we will alternatively denote $\tilde{w}(x)$ to be the additive convolution $\varphi_{\Delta} * \mu$, where $\varphi_{\Delta}(x)$ is the even function $\Delta^{-1}k(x/\Delta)$, and μ is again the empirical spectral measure. This marks a slight departure from the multiplicative convolution with the approximation to the identity $x \mapsto \Delta^{-1}k((x-1)/\Delta)$ appearing in the theorem, but the proofs are identical to order Δ^2 after appropriate logarithmic transformations and Hilbert-transform invariances are applied. The reason the multiplicative convolution was originally proposed is a practical one: the smallest estimated eigenvalue becomes positive more quickly as $n \to \infty$.

Our first main result of this section is the following.

LEMMA 7. Assume the conditions of Theorem 5. Then \tilde{w} satisfies

(70)
$$\frac{1}{p} \sum_{i=1}^{p} |\tilde{w}(\lambda_i) - w(\lambda_i)| < n^{-2/3}.$$

(In other words, $\|\tilde{w} - w\|_{L^1(\mu)} \prec n^{-2/3}$.)

PROOF. We will use the decomposition, with $w_{\Delta} := \varphi_{\Delta} * d\mu_{\infty}$:

$$(71) |w - \tilde{w}| \le |w - w_{\Delta}| + |\varphi_{\Delta} * (d\mu_{\infty} - d\mu)|.$$

By Lemma 6, the second term of the right-hand side above is

We only consider the case where φ and φ' have no more than 3 monotonic intervals. The fundamental theorem of calculus then yields

so that

(74)
$$\frac{1}{p} \sum_{i=1}^{p} |w_{\Delta}(\lambda_i) - \tilde{w}(\lambda_i)| \prec \Delta^{-1} n^{-1}.$$

We will now focus on the first term.

We sum $|w-w_{\Delta}|(\lambda_i)$ over λ_i falling into two cases: $\kappa < 2\Delta$ and $\kappa > 2\Delta$, where $\kappa = \kappa(\lambda_i)$ is the distance from the spectral edge. We note that, as in [KY17], $w(x) \approx \kappa(x)^{1/2}$ for x near the spectral edge.

For $\kappa(x) \leq 2\Delta$, we have

(75)
$$|w(x) - w_{\Delta}(x)| \leq \int_{x-\Delta}^{x+\Delta} \varphi_{\Delta}(x-t)|w(x) - w(t)| dt$$

$$\lesssim \int_{x-\Delta}^{x+\Delta} |\varphi_{\Delta}(x-t)| \Delta^{1/2} dt$$

$$\leq \Delta^{1/2}.$$

Therefore, we have

(76)
$$\frac{1}{p} \sum_{\kappa(\lambda_i) \le 2\Delta} |w(x) - w_{\Delta}(x)| \le \left(C(2\Delta)^{3/2} + p^{\epsilon - 1} \right) \Delta^{1/2} \lessapprox \Delta^2$$

with high probability, where we estimated the number of terms in the sum using the small-scale clause of Theorem 1.

For $\kappa(x) > 2\Delta$, Taylor-expanding w(t) about t = x and using evenness of φ gives that

(77)
$$w(x) - w_{\Delta}(x) = \int_{x-\Delta}^{x+\Delta} \varphi_{\Delta}(x-t)(w(x) - w(t)) dt$$

$$= \int_{x-\Delta}^{x+\Delta} \varphi_{\Delta}(x-t)$$

$$\cdot \left(w'(x)(x-t) - \frac{w''(x)}{2}(x-t)^2 + O(\Delta^3 \max_{[x-\Delta, x+\Delta]} w''') \right) dt$$

$$\lessapprox \Delta^2(\kappa(x) - \Delta)^{-3/2}$$

$$\lessapprox \Delta^2 \kappa(x)^{-3/2}$$

where we have used [KY17] again to bound $|w^{(k)}| \lesssim \kappa^{1/2-k}$ for any fixed k.

Therefore, we have

(78)
$$\frac{1}{p} \sum_{\kappa(\lambda_i) > 2\Delta} |w(x) - w_{\Delta}(x)| \lessapprox \frac{1}{p} \sum_{\kappa(\lambda_i) > 2\Delta} \Delta^2 \kappa(\lambda_i)^{-3/2}$$

The right hand side is precisely

(79)
$$\int \widetilde{f} d\mu,$$

where $\widetilde{f}(x):=\Delta^2\kappa(x)^{-3/2}\mathbf{1}_{\kappa(x)>2\Delta}$. The only difficulty in applying Lemma 6 is that \widetilde{f} is discontinuous, so we must adjust it. Let $g:\mathbb{R}\to[0,1]$ be C^∞ , equal 0 for $x\geq 2$, equal 1 for $x\leq 1$ and satisfy $\|g'\|_\infty+\|g''\|_\infty\leq C$. Then define

(80)
$$f(x) = \widetilde{f}(x)g((2\Delta)^{-1}\kappa(x)),$$

which is now C^2 and satisfies

(81)
$$||f||_1 + ||f'||_1 + ||f''||_1 \le 1.$$

Therefore Lemma 6 gives

$$\frac{1}{p} \sum_{\kappa(\lambda_{i})>2\Delta} |w(x) - w_{\Delta}(x)| \lessapprox \frac{1}{p} \sum_{\kappa(\lambda_{i})>2\Delta} \Delta^{2} \kappa(\lambda_{i})^{3/2}$$

$$\lessapprox \int \widetilde{f}(x) d\mu(x)$$

$$= \int \widetilde{f}(x) d\mu_{\infty}(x) + O_{\prec}(n^{-1})$$

$$\lessapprox \int \widetilde{f}(x) \kappa(x)^{1/2} dx + O_{\prec}(n^{-1})$$

$$\lessapprox \Delta^{2} \int_{\kappa(x)>2\Delta} \kappa(x)^{-3/2} \kappa(x)^{1/2} dx + O_{\prec}(n^{-1})$$

$$\asymp \Delta^{2} \log |\Delta| \prec \Delta^{2}$$

with high probability.

Equations (76) and (82) together establish that

(83)
$$\frac{1}{p} \sum_{i=1}^{p} |\tilde{w}(\lambda_i) - w_{\Delta}(\lambda_i)| \prec \Delta^2,$$

which, with equation (74) gives

(84)
$$\frac{1}{p} \sum_{i=1}^{p} |w(\lambda_i) - \tilde{w}(\lambda_i)| < \Delta^2 + \Delta^{-1} n^{-1}.$$

Thus the choice of $\Delta = n^{-1/3}$ becomes clear, and we conclude the proof.

The Hilbert transform $\mathcal{H}w$ may be treated in much the same way as w.

LEMMA 8. Assume the conditions of Theorem 5. Then $\mathcal{H}\tilde{w}$ satisfies

(85)
$$\frac{1}{p} \sum_{i=1}^{p} |\mathcal{H}\tilde{w}(\lambda_i) - \mathcal{H}w(\lambda_i)| < n^{-2/3}.$$

PROOF. We decompose:

$$|\mathcal{H}w - \mathcal{H}w_{\Delta}^{\mu}| \le |\mathcal{H}w - \mathcal{H}w_{\Delta}| + |\mathcal{H}w_{\Delta} - \mathcal{H}\tilde{w}|$$

Using the relation between \mathcal{H} and convolution, the above may be written as

(87)
$$|\mathcal{H}w - \varphi_{\Delta} * (\mathcal{H}w)| + |(\mathcal{H}\varphi_{\Delta}) * \mu_{\infty} - (\mathcal{H}\varphi_{\Delta}) * \mu|$$

The latter term may be written, letting $\hat{\varphi} := \mathcal{H}\varphi$, as

(88)
$$|(\mathcal{H}\varphi_{\Delta}) * \mu_{\infty} - (\mathcal{H}\varphi_{\Delta}) * \mu| = |\hat{\varphi}_{\Delta} * \mu_{\infty} - \hat{\varphi}_{\Delta} * \mu|$$

Let $f = \hat{\varphi}_{\Delta} \cdot g$, where g is C^{∞} , equals 1 on [-C, C], equals 0 on $\mathbb{R} \setminus [-2C, 2C]$, and has bounded first and second derivatives, for some large constant C. As in the proof of Lemma 7, we use Lemma 6: uniformly in $x \leq C/2$,

(89)
$$|\hat{\varphi}_{\Delta} * \mu_{\infty}(x) - \hat{\varphi}_{\Delta} * \mu(x)| = |f * \mu_{\infty}(x) - f * \mu(x)| \\ \prec n^{-1} \|f\|_{1} + n^{-1} \|f'\|_{1} + n^{-2} \|f''\|_{1}$$

(the appearance of $||f||_1$ is why we needed to attenuate $\hat{\varphi}_{\Delta}$ by a cutoff function: $\mathcal{H}\varphi$ is not L^1 ; we may do this because μ_{∞} and μ have uniformly bounded support with high probability). Now we may conclude similarly to in the argument leading to (74). We have that $||f||_1 \lesssim$ $|\log \Delta|$, and that f and f' may be ensured to have a bounded number of monotonic intervals, so that $\|f'\|_1 \lesssim \|f\|_\infty \lesssim \Delta^{-1}$ and $\|f''\|_1 \lesssim \|f'\|_\infty \lesssim \Delta^{-2}$. Now we treat the former term

(90)
$$\|\mathcal{H}w - \varphi_{\Delta} * (\mathcal{H}w)\|$$

Now, $\mathcal{H}w$ is analytic inside the support of μ_{∞} . We have for $\mathcal{H}w$ that

(91)
$$(\mathcal{H}w)^{(k)} \lesssim \kappa(x)^{1/2-k}$$

for any fixed k just as we had for w, so that we may repeat the argument in Lemma 7.

APPENDIX D: DETERMINISTIC LIMITING VARIANCE

We choose the somewhat more general task of estimating $p^{-1} \operatorname{tr}(\mathbf{\Xi}_n f_n(\mathbf{S}_n) \mathbf{\Sigma}_n g_n(\mathbf{S}_n))$ for some matrix Ξ_n having

(92)
$$p^{-1} \sum_{i=1}^{p} \frac{\mathbf{u}'_{ni} \Xi_n \mathbf{u}_{ni}}{\lambda_{ni} - z} = \int \frac{\xi(x)}{x - z} dx + O_{\prec} \left(\frac{1}{n\eta}\right)$$

for some bounded measurable $\xi: \mathbb{R} \to \mathbb{R}$ and all $z \in \mathbb{C}^+$. By Theorem 2 and Appendix B, this is certainly true of Σ_n —the value of Ξ_n that is of most interest—in which case $\xi(x) = \delta(x)$.

The strategy is to first estimate the alternating trace above for f and g equal to resolvent functions, then use the usual limiting argument as complex arguments go the real axis to go from resolvent functions to continuous functions. For the remainder of this appendix, we will resume exercising the convention of suppressing most subscripts of n.

Assume as a first case that $f(\lambda) \equiv f_n(\lambda) = (\lambda - x_z - i\eta_z)^{-1}$ and $g(\lambda) \equiv g_n(\lambda) = (\lambda - i\eta_z)^{-1}$ $(x_v - i\eta_v)^{-1}$ for positive and distinct η_x and η_z , and let $z = x_z + i\eta_z$ and $w = x_v + i\eta_v$. Our first main goal will be to show that

$$(93) p^{-1} \operatorname{tr} (\mathbf{\Xi} f(\mathbf{S}) \mathbf{\Sigma} g(\mathbf{S})) =$$

$$p^{-1} \sum_{i=1}^{p} \frac{\lambda_{i} \mathbf{u}_{i}' \mathbf{\Xi} \mathbf{u}_{i}}{(\lambda_{i} - z)(\lambda_{i} - v)} \left(1 + \frac{1}{n} \sum_{i=1}^{p} \frac{\mathbf{u}_{i}' \mathbf{\Sigma} \mathbf{u}_{i}}{\lambda_{i} - z} \right) \left(1 + \frac{1}{n} \sum_{i=1}^{p} \frac{\mathbf{u}_{i}' \mathbf{\Sigma} \mathbf{u}_{i}}{\lambda_{i} - v} \right)$$

$$+ O_{\prec} \left(\frac{1}{n n_{r} n_{r}} \right).$$

To this end, we first observe the following:

$$p^{-1}\operatorname{tr}(\mathbf{R}(z)\mathbf{\Xi}\mathbf{R}(v)) = p^{-1}\operatorname{tr}(\mathbf{R}(v)\mathbf{R}(z)\mathbf{\Xi})$$

(94)
$$= p^{-1} \sum_{i=1}^{p} \frac{\mathbf{u}_i' \mathbf{\Xi} \mathbf{u}_i}{(\lambda_i - z)(\lambda_i - v)},$$

Consider

(95)
$$p^{-1}\operatorname{tr}(\mathbf{\Xi}\mathbf{R}(v)) + zp^{-1}\operatorname{tr}(\mathbf{R}(z)\mathbf{\Xi}\mathbf{R}(v)).$$

Since $I + z\mathbf{R}(z) = \mathbf{S}\mathbf{R}(z)$, we can write the above as

$$p^{-1}\operatorname{tr}((\mathbf{I} + z\mathbf{R}(z))\mathbf{\Xi}\mathbf{R}(v))$$
$$= p^{-1}\operatorname{tr}(\mathbf{S}\mathbf{R}(z)\mathbf{\Xi}\mathbf{R}(v)).$$

Writing $S = \sum_{k=1}^{n} c_k c'_k$, with c_k defined as in Appendix A, we get the above is

(96)
$$p^{-1} \sum_{k=1}^{n} \mathbf{c}_{k}' \mathbf{R}(z) \mathbf{\Xi} \mathbf{R}(v) \mathbf{c}_{k}.$$

Another application of the Woodbury formula (45) gives the following expansion of (96).

$$p^{-1} \sum_{k=1}^{n} \mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{\Xi} \mathbf{R}^{(k)}(v) \mathbf{c}_{k} +$$

$$- p^{-1} \sum_{k=1}^{n} \mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{\Xi} \mathbf{R}^{(k)}(v) \mathbf{c}_{k} \left(1 - \frac{1}{1 + \mathbf{c}_{k}' \mathbf{R}^{(k)}(v) \mathbf{c}_{k}} \right) +$$

$$- p^{-1} \sum_{k=1}^{n} \mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{\Xi} \mathbf{R}^{(k)}(v) \mathbf{c}_{k} \left(1 - \frac{1}{1 + \mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{c}_{k}} \right) +$$

$$+ p^{-1} \sum_{k=1}^{n} \mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{\Xi} \mathbf{R}^{(k)}(v) \mathbf{c}_{k} \frac{\mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{c}_{k}}{1 + \mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{c}_{k}} \frac{\mathbf{c}_{k}' \mathbf{R}^{(k)}(v) \mathbf{c}_{k}}{1 + \mathbf{c}_{k}' \mathbf{R}^{(k)}(v) \mathbf{c}_{k}}$$

$$= p^{-1} \sum_{k=1}^{n} \frac{\mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{\Xi} \mathbf{R}^{(k)}(v) \mathbf{c}_{k}}{(1 + \mathbf{c}_{k}' \mathbf{R}^{(k)}(z) \mathbf{c}_{k}) \left(1 + \mathbf{c}_{k}' \mathbf{R}^{(k)}(v) \mathbf{c}_{k}} \right)}.$$

$$(97)$$

Observe that

$$f^k(\mathbf{w}) = \mathbf{w}' \mathbf{\Sigma}^{1/2} \mathbf{R}^{(k)}(z) \mathbf{\Sigma}^{1/2} \mathbf{w}$$

is a quadratic form in w. Since $\Sigma^{-1/2}\mathbf{c}_k$ is a subgaussian random vector, Hanson-Wright then implies

$$\left| \mathbf{c}_k' \mathbf{R}^{(k)}(z) \mathbf{c}_k - \frac{\operatorname{tr}(\mathbf{\Sigma} \mathbf{R}^{(k)}(z))}{n} \right| \prec \left(\frac{1}{\eta_z \sqrt{p}} \right).$$

By a similar calculation,

$$\left|\mathbf{c}_k'\mathbf{R}^{(k)}(z)\mathbf{\Xi}\mathbf{R}^{(k)}(v)\mathbf{c}_k - \frac{\operatorname{tr}(\mathbf{\Sigma}\mathbf{R}^{(k)}(z)\mathbf{\Xi}\mathbf{R}^{(k)}(v))}{n}\right| \prec \left(\frac{1}{\eta_z\eta_v\sqrt{p}}\right).$$

Using uniformity in k, we could use the same bound for (97), but we can improve the factor of $1/\sqrt{p}$ to 1/p using the "almost-independence" argument identical to Lemma 5, obtaining:

(98)
$$p^{-1} \sum_{k=1}^{n} \frac{\frac{1}{n} \operatorname{tr}(\mathbf{\Sigma} \mathbf{R}^{(k)}(z) \mathbf{\Xi} \mathbf{R}^{(k)}(v))}{\left(1 + \frac{1}{n} \operatorname{tr}(\mathbf{\Sigma} \mathbf{R}^{(k)}(z))\right) \left(1 + \frac{1}{n} \operatorname{tr}(\mathbf{\Sigma} \mathbf{R}^{(k)}(v))\right)} + O_{\prec} \left(\frac{1}{\eta_{z} \eta_{v} p}\right).$$

Using the fact that $\mathbf{R}^{(k)}(z)$ can be replaced by $\mathbf{R}(z)$ with an error of only $O(n^{-1}\eta^{-1})$ (by Woodbury again), (98) becomes

$$\frac{1}{p} \frac{\operatorname{tr}(\mathbf{\Sigma}\mathbf{R}(z)\mathbf{\Xi}\mathbf{R}(v))}{\left(1 + \frac{1}{n}\operatorname{tr}(\mathbf{\Sigma}\mathbf{R}(z))\right)\left(1 + \frac{1}{n}\operatorname{tr}(\mathbf{\Sigma}\mathbf{R}(v))\right)} + O_{\prec}\left(\frac{1}{\eta_z\eta_v p}\right).$$

Comparing to (94) and (95) gives

$$\frac{1}{p} \sum_{i=1}^{p} \frac{\mathbf{u}_{i}' \mathbf{\Xi} \mathbf{u}_{i}}{\lambda_{i} - v} + z \left(p^{-1} \sum_{i=1}^{p} \frac{\mathbf{u}_{i}' \mathbf{\Xi} \mathbf{u}_{i}}{(\lambda_{i} - z)(\lambda_{i} - v)} \right)$$

$$= \frac{1}{p} \frac{\operatorname{tr}(\mathbf{\Sigma} \mathbf{R}(z) \mathbf{\Xi} \mathbf{R}(v))}{(1 + \frac{1}{n} \operatorname{tr}(\mathbf{\Sigma} \mathbf{R}(z))) (1 + \frac{1}{n} \operatorname{tr}(\mathbf{\Sigma} \mathbf{R}(v)))} + O_{\prec} \left(\frac{1}{\eta_{z} \eta_{v} p} \right)$$

so that, using the asymptotic boundedness of $1 + \frac{1}{n} tr(\Sigma \mathbf{R}(z))$, proved in Lemma 3, we have

(99)
$$= \left(1 + \frac{1}{n} \sum_{i=1}^{p} \frac{\mathbf{u}_{i}' \mathbf{\Sigma} \mathbf{u}_{i}}{\lambda_{i} - z}\right) \left(1 + \frac{1}{n} \sum_{i=1}^{p} \frac{\mathbf{u}_{i}' \mathbf{\Sigma} \mathbf{u}_{i}}{\lambda_{i} - v}\right) p^{-1} \sum_{i=1}^{p} \frac{\lambda_{i} \mathbf{u}_{i}' \mathbf{\Xi} \mathbf{u}_{i}}{(\lambda_{i} - z)(\lambda_{i} - v)} + O_{\prec} \left(\frac{1}{\eta_{z} \eta_{v} p}\right),$$

which accomplishes the goal of (93).

Applying the argument of Appendix B to the real and imaginary parts of

$$n^{-1} \sum_{i} \frac{\mathbf{u}_{i}' \mathbf{\Sigma} \mathbf{u}_{i}}{\lambda_{i} - z}$$

and using linearity allows us to replace this summation by $z\mapsto \phi\int d\nu(\lambda)/(\lambda-z)$, with a negligible error of $1/(n\min\{\eta_z,\eta_v\})$. Similarly, using partial fractions on the remaining sum and assuming $\liminf_n |\eta_z/\eta_v-1|\neq 0$ allows us to replace this sum by

$$\int \frac{\lambda \xi(\lambda)}{(\lambda - z)(\lambda - v)} d\mu_{\infty}(\lambda) + O_{\prec}\left(\frac{1}{\eta_z \eta_v p}\right).$$

Expanding the resulting product of three integral expressions and using Fubini's theorem and some algebra gives

(100)

$$p^{-1}\operatorname{tr}(\mathbf{\Xi}f(\mathbf{S})\boldsymbol{\Sigma}g(\mathbf{S}))$$

$$= \int f(x)g(x)x\xi(x)\,d\mu_{\infty}(x)$$

$$-\phi \int_{\mathbb{R}^{2}} f(x)\frac{g(y) - g(x)}{y - x}\,x\xi(x)\,d\mu_{\infty}(x)d\nu_{\infty}(y)$$

$$-\phi \int_{\mathbb{R}^{2}} g(x)\frac{f(y) - f(x)}{y - x}\,x\xi(x)\,d\mu_{\infty}(x)d\nu_{\infty}(y)$$

$$+\phi^{2} \int_{\mathbb{R}^{3}} \frac{g(u) - g(x)}{u - x}\frac{f(y) - f(x)}{y - x}\,x\xi(x)\,d\mu_{\infty}(x)d\nu_{\infty}(y)d\nu_{\infty}(u) + O_{\prec}\left(\frac{1}{\eta_{z}\eta_{v}n}\right).$$

Next, we claim that this identity holds for general regular functions $\tilde{f} \equiv \tilde{f}_n$ and $\tilde{g} \equiv \tilde{g}_n$. For this, we simply integrate against $\tilde{f}(\eta_z)$ and $\tilde{g}(\eta_v)$ and let η_z and η_v go to zero at a

rate faster than $n^{-1/3}$ but slower than $n^{-1/2}$. This results in the above equality with $f \leftarrow \tilde{f} * \varphi_{\eta_z}$ and $g \leftarrow \tilde{g} * \varphi_{\eta_v}$, where φ is the Cauchy kernel. Further, since \tilde{f} is regular, we have $\tilde{f} * \varphi_{\eta}(x) = \tilde{f}(x) + O_{\prec}(n^{2/3}\eta^2)$ uniformly in x, and so we replace these convolutions by \tilde{f} and \tilde{g} , incurring only $o_{a.s.}(1)$ error. The claim follows since our choice of η_z, η_v implies that the error of $O_{\prec}(n^{-1}\eta_z^{-1}\eta_v^{-1})$ goes to zero almost surely.

Thus, we may assume (100) holds for general regular functions f and g and the right-hand side of this identity can be factored, giving

(101)
$$\sigma_{\infty}^{2}(f,g;\xi) := \int Tf(x)Tg(x)x\xi(x) d\mu_{\infty}(x)$$

since linearity of the Hilbert transform can be used to express T as

$$Tf(x) = f(x) - \phi \int \frac{f(y) - f(x)}{y - x} d\nu_{\infty}(y).$$

In particular, (100) holds if the right-hand side is replaced by $\sigma^2_{\infty}(f,f) = \sigma^2_{\infty}(f,f;\delta)$, as desired.

APPENDIX E: CONSISTENCY OF VARIANCE ESTIMATE

The main technical step will be to prove that

(102)
$$\left\| (T_n f_n - T f_n)^{(j)} \right\|_{L^1(E)} \prec \Delta^{2-j}$$

for derivatives of order j = 0, 1, and 2, where $\Delta = n^{-1/3}$.

First we consider j=0. The terms of $T_nf_n(x)$ and $Tf_n(x)$ that depend on x can be shown to be consistent with one another in $L^1(\mu_n)$ within $O_{\prec}(\Delta^2)$ using Appendices B and C. The terms not depending on x are of the form $\mathcal{H}_w[f_n\tilde{d}_n]$ and $\tilde{\mathcal{H}}_w[f_n\tilde{d}_n]$, where

(103)
$$\mathcal{H}_{w} f \equiv \mathcal{H}_{w}[f] := \mathcal{H}[fw]$$

and

(104)
$$\tilde{\mathcal{H}}_w f(x) \equiv \tilde{\mathcal{H}}_w[f](x) := p^{-1} \sum_{i=1}^p f(\lambda_i) K_{h_{ni}}(x - \lambda_i),$$

To simplify our analysis, as before, we make the replacement $h_{ni} \leftarrow \Delta$ in $K_{h_{ni}}$ above. Similar to before, without this replacement $\tilde{\mathcal{H}}_w[f](x)$ is a convolution with respect to multiplication rather than addition, so the analyses do not significantly differ in character.

As a result of the above analysis, showing (102) for j = 0 can be reduced to the following lemma.

LEMMA 9. Assume (H1)-(H5) and that f is regular. Then we have the following estimate

(105)
$$\left\| \left(\tilde{\mathcal{H}}_w[f] - \mathcal{H}_w[f] \right) w \right\|_q^q \prec \Delta^2$$

for any finite $q \geq 1$.

PROOF. Using the notation $K_{\Delta}(x) = \Delta^{-1}K(x/\Delta)$ and, we have $\tilde{\mathcal{H}}_w[f] = K_{\Delta}*(f d\mu_n)$. Using the Helffer-Sjöstrand argument of Appendix B and the fact that

$$\|(d/d\lambda)K_{\Delta}(x-\lambda)\|_1 \lessapprox 1/\Delta \qquad \text{and} \qquad \left\|(d/d\lambda)^2K_{\Delta}(x-\lambda)\right\|_1 \lessapprox 1/\Delta^2,$$

we get that $ilde{\mathcal{H}}_w f$ can be approximated by

(106)
$$\int f(\lambda)K_{\Delta}(x-\lambda)w(\lambda)\,d\lambda = \int f(\lambda)K_{\Delta}(x-\lambda)w(\lambda)\varphi(\lambda)\,d\lambda$$

where φ is a smooth cutoff function with $\varphi' \leq \Delta^{-1}$ and $\varphi'' \leq \Delta^{-2}$ that takes the value 1 on F and 0 on $\mathbb{R} \setminus (F + \Delta)$. By Appendix B, then, the error in the above approximation of $\tilde{\mathcal{H}}_w f$ can be written as

(107)

$$O_{\prec}\left(\frac{\|fK\|_{L^{1}(F)}}{n} + \frac{\|f'K + fK'/\Delta\|_{L^{1}(F)}}{n} + \frac{\|f''K + f'K'/\Delta + fK''/\Delta^{2}\|_{L^{1}(F)}}{n^{2}}\right),$$

which is $O_{\prec}(\Delta^2)$ since f'' is bounded and K, K', and K'' are integrable on F.

We may now use the anti-self-adjointess of the Hilbert transform to obtain from the integral above

$$\mathcal{H}_w[f] * k_{\Delta}(x) = \int \mathcal{H}[fw](\lambda) k_{\Delta}(x-\lambda) d\lambda.$$

Using continuity and/or the properties of approximate identities, this function converges in various modes to the desired function $\mathcal{H}_w f$, but we have yet to establish the desired mode and rate of convergence.

Up to this point, we have shown that $\tilde{\mathcal{H}}_w[f]$ is an $O_{\prec}(\Delta^2)$ approximation to $\mathcal{H}_w[f]*k_{\Delta}$ with repect to the Lebesgue measure's infinity norm. We next show that $\mathcal{H}_w[f]*k_{\Delta}$ converges to \mathcal{H}_wf at a rate of $O_{\prec}(\Delta^{2/q})$ in the $L^q(\mu_{\infty})$ norm for finite $q \geq 1$. That is,

(108)
$$\int |\mathcal{H}_w[f] * k_{\Delta}(\lambda) - \mathcal{H}_w f(\lambda)|^q w \, dx \prec \Delta^2.$$

This estimate follows from the fact that the integrand is eventually bounded and for q=1 the left-hand side is bounded by $O_{\prec}(\Delta^2|\log\Delta|) = O_{\prec}(\Delta^2)$ in exactly the same way as in Lemma 6, since square-root behavior of w near the spectral edges implies square-root behavior of $\mathcal{H}[fw]$ near the spectral edges.

Now let us reconsider $\tilde{\sigma}_n^2(f_n) = p^{-1} \sum_{i=1}^p T_n f_n(\lambda_{ni})^2 \lambda_{ni} \tilde{d}_n(\lambda_{ni})$. The terms $\tilde{d}_n(\lambda_{ni})$ may be replaced by $\delta(\lambda_{ni})$ by the work in Appendix C, committing only an error of order $O_{\prec}(\Delta^2)$. The result is equivalent to

$$\sigma_{\infty}^{2}(f_{n}, f_{n}) = \int T f_{n}(\lambda)^{2} \lambda \delta(\lambda) d\mu_{\infty}(\lambda)$$

up to an error of order

$$O_{\prec} \left(\frac{\|g_n\|_1}{n} + \frac{\|g_n'\|_1}{n} + \frac{\|g_n''\|_1}{n^2} \right),$$

where $g_n=(T_nf_n)^2-(Tf_n)^2$. The first term is stochastically dominated by Δ^2/n by the lemma and discussion above, and the fact that f_n , and thus g_n , is uniformly bounded. For the second and third term, we need (102) with j=1 and j=2. But these cases follow by observing that $\Delta^{j+\epsilon}(T_nf_n-Tf_n)^{(j)}$ is regular for $j\in\{1,2\}$ and small positive ϵ , using linearity and the fact that $g_n^{(j)}\prec\Delta^{-j}$. The theorem is proved.

APPENDIX F: PROOF OF THEOREM 7(A)

Since Lemma 1 proves $\tilde{\sigma}_n(f_n) = \sigma_{\infty}(f_n, f_n) + o_{a.s.}(1)$, it suffices to show that

(109)
$$p^{-1/2} \boldsymbol{\mu}'_n f_n(\mathbf{S}_n) \boldsymbol{\mu}_n = \int f_n \, d\omega + o_{a.s.}(1).$$

By Hanson-Wright, the left side is almost-surely equivalent to

$$p^{-3/2}\operatorname{tr}(f_n(\mathbf{S}_n)\mathbf{\Omega}_n).$$

Thus, by following the argument of Appendix B again, we get that the error in (109) is

$$p^{-3/2}\operatorname{tr}(f_n(\mathbf{S}_n)\mathbf{\Omega}_n) = \int f_n d\omega + O_{\prec} \left(\frac{\|f_n\|_1}{n} + \frac{\|f'_n\|_1}{n} + \frac{\|f''_n\|_1}{n^2} \right),$$

which is $o_{a.s.}(1)$ by regularity, as desired.

APPENDIX G: PROOF OF THEOREM 7(B)

We prove the theorem for a slightly more general objective function. Let

(110)
$$\mathcal{U}(f,\omega,\xi) = \frac{\int f \,d\omega}{\sigma_{\xi}(f,f;\xi)},$$

where $\sigma(\cdot;\xi)$ was defined in (101). This objective function arises, for instance, in the work of Ledoit and Wolf on financial portfolio optimization, in which case $\xi \equiv 1$, though we will not pursue this case in more detail.

More suggestively, we may re-express $\mathcal{U}(f,\omega,\xi)$ as follows:

(111)
$$\frac{\langle f, \omega' \rangle}{\sqrt{\langle M_{\text{id} w\xi} T f, T f \rangle}},$$

where $\langle \cdot \rangle$ is the inner product on $H \times H$. Taking $A = T' M_{\mathrm{id} w \xi} T : H \to H$, where T' is the transpose of $T : H \to H$, Cauchy-Schwarz implies that this ratio is maximized precisely when

$$f = A^{-1}\omega'$$
.

Inversion of the operator A can be performed in two main steps. First, writing $A^{-1} = T^{-1}M_{\mathrm{id}\,w\xi}^{-1}(T')^{-1}$, we would like to invert T'. A next, minor step is to multiply by the reciprocal of the function $x \mapsto xw(x)\xi(x)$. The second major step is to invert T, which can be done by writing $T^{-1} = ((T')^{-1})'$ —just the adjoint of the previously found inverse $(T')^{-1}$.

For the first main step, observe that T takes the following form:

(112)
$$Tf(x) = b(x)f(x) - \phi \text{ p.v.} \int \frac{f(y)}{y-x} d\nu_{\infty}(y)$$

(113)
$$= b(x)f(x) - \phi \pi \mathcal{H}[f\delta w](x)$$

where

$$b(x) := 1 + \phi \pi \mathcal{H}[d\nu_{\infty}](x).$$

Thus,

(114)
$$T'f(x) = b(x)f(x) - B(x)\mathcal{H}[f](x),$$

where

(115)
$$B(x) := -\phi \pi \delta(x) w(x).$$

In order to invert T', we need a lemma.

LEMMA 10. The equation $T'f = \varphi \in H$ has the unique solution

$$(116) f = g\varphi + G\mathcal{H}[\varphi],$$

where
$$g(x) = 1 - \phi - \phi \pi x \mathcal{H} w(x)$$
 and $G(x) = -\phi \pi x w(x)$.

PROOF. We show this in two steps. First, we simplify b by considering the analytic signal b+iB and its extension to the complex upper half-plane. Second, we invert T' on H by solving a singular integral equation.

First we simplify b.

As mentioned in (36), this Stieltjies transform satisfies an identity equivalent to

$$1 + \phi \Theta_{\infty}(z) = \frac{1}{1 - \phi - \phi z m_{\infty}(z)}.$$

Using the result from Theorem 2 that for x > 0,

$$\lim_{\eta \to 0^+} \operatorname{Im} \left[\Theta_{\infty}(x + i\eta) \right] = \pi \delta(x) w(x) = \pi d\nu_{\infty}(x) / dx$$

Thus, by the properties of the Hilbert transform, $\pi \mathcal{H}[d\nu_{\infty}]$ is given by the limit of the real part of $\Theta_{\infty}(z)$ as $\mathrm{Im}z \to 0^+$, and similarly g is given by the following:

$$\begin{split} b(x) &= \lim_{\eta \to 0^+} \operatorname{Re} \left[1 + \phi \Theta_\infty(x + i \eta) \right] \\ &= \operatorname{Re} \lim_{\eta \to 0^+} \frac{1}{1 - \phi - \phi z m_\infty(z)} \\ &= \operatorname{Re} \left[\frac{1}{1 - \phi - \phi x \breve{m}(x)} \right] \\ &= \frac{1 - \phi - \phi x \operatorname{Re} [\breve{m}(x)]}{|1 - \phi - \phi x \breve{m}(x)|^2}. \end{split}$$

By the fact that $\breve{m}(x) = \pi \mathcal{H}[d\mu_{\infty}] + i\pi w(x)$, the above can be written

$$\frac{1 - \phi - \phi \pi x \mathcal{H} w(x)}{(1 - \phi - \phi \pi x \mathcal{H} w(x))^2 + \phi^2 \pi^2 x^2 w(x)^2}.$$

Assume $T'f = \varphi \in H$. Observe that we may write

$$b(x) = 1 + \phi \pi \mathcal{H}[\delta w](x).$$

Using the fact that 1 has vanishing Hilbert transform and \mathcal{H} is an anti-involution, we observe that $B(x) = \mathcal{H}[b](x)$. Thus, we may take the Hilbert transform of (114) and use the identity $\mathcal{H}[bf - B\mathcal{H}f] = Bf + b\mathcal{H}f$ of [CL77] to obtain

$$Bf + b\mathcal{H}f = \mathcal{H}\varphi$$
.

Together with (114), this yields a two-by-two linear system for f and $\mathcal{H}f$, which can be solved for f as follows:

(117)
$$(T')^{-1}\varphi = f = \frac{b\varphi + B\mathcal{H}\varphi}{h^2 + R^2}.$$

In order to simplify the coefficients $b/(b^2+B^2)$ and $B/(b^2+B^2)$, observe that

$$B(x) = -\phi \pi \delta(x) w(x) = \frac{-\phi \pi x w(x)}{(1 - \phi - \phi \pi x \mathcal{H} w(x))^2 + \phi^2 \pi^2 x^2 w(x)^2},$$

so that

(118)
$$\frac{b(x)}{b(x)^2 + B(x)^2} = 1 - \phi - \phi \pi x \mathcal{H} w(x) = g(x)$$

and

$$\frac{B(x)}{b(x)^2 + B(x)^2} = -\phi \pi x w(x) = G(x),$$

by the definitions of g and G. In other words,

$$(119) (T')^{-1}\varphi = q\varphi + G\mathcal{H}\varphi,$$

as desired.

As a result of the above lemma, we may easily apply $(T')^{-1}$ to ω' . The next step is to apply in the inverse of $M_{\mathrm{id}\,w\xi}$ to the result, obtaining

(120)
$$f_* := M_{\mathrm{id}\, w\xi}^{-1}(T')^{-1}\omega'$$

(121)
$$= \frac{1}{\operatorname{id} w\xi} \left(g\omega' + G\mathcal{H}[\omega'] \right).$$

The above lemma also enables us to easily compute $((T')^{-1})'$, which coincides with T^{-1} , the final operator we must apply. Using the anti-self-adjointness of \mathcal{H} and the self-adjointness of multiplication operators, one obtains that for $\varphi \in H$

$$T^{-1}\varphi = ((T')^{-1})'\varphi = g\varphi - \mathcal{H}[G\varphi].$$

Thus,

$$\begin{split} A^{-1}\omega' &= T^{-1}f_* \\ &= gf_* - \mathcal{H}\left[Gf_*\right] \\ &= g\frac{g\omega' + G\mathcal{H}[\omega']}{\operatorname{id} w\xi} - \mathcal{H}\left[G\frac{g\omega' + G\mathcal{H}[\omega']}{\operatorname{id} w\xi}\right], \end{split}$$

as desired.

APPENDIX H: CONSISTENCY OF PROPOSED SHRINKER

As in previous appendices, we will frequently suppress subscripts of n wherever they are unnecessary. As in Appendix C, we will use $\tilde{w}(x)$ and $\mathcal{H}\tilde{w}(x)$ to denote the additive convolutions of μ with k or K rather than the multiplicative ones, with the understanding that our results can easily be adapted to the multiplicative versions.

The main idea will then be to show almost-sure convergence of $p^{-3/2}\mathrm{tr}(f_n(\mathbf{S}_n)\mathbf{\Omega}_n)$ to $\int f^* d\omega$ and $\sigma_n^2(f_n)$ to $\sigma_\infty^2(f^*,f^*)$. For the former, Appendix B and Cauchy-Schwarz imply that the identity $\|(f_n-f^*)w\|_2 \overset{\mathrm{a.s.}}{\to} 0$ is sufficient. For the latter, by the triangle inequality, it suffices to show that $\sigma_\infty^2(f_n-f^*,f_n-f^*) \overset{\mathrm{a.s.}}{\to} 0$. By using the expression of $\sigma_\infty^2(\cdot)$ in terms of the bounded operator $T:H\to H$, we see that this also follows if $\|(f_n-f^*)w\|_2 \overset{\mathrm{a.s.}}{\to} 0$.

To establish this norm's convergence, observe that from Appendix G, f^* takes the form of $\psi_1\Psi + \psi_2\mathcal{H}_w[\psi_3\Psi] + \mathcal{H}_w[\psi_4\mathcal{H}_w[\Psi]]$, where $\Psi = \omega'/w$ is non-singular and ψ_j are non-singular functions depending on $\mathcal{H}w$, and that f_n is obtained by simply replacing each instance of \mathcal{H}_w with $\tilde{\mathcal{H}}_w$ and $\mathcal{H}w$ with $\mathcal{H}\tilde{w}$. To shorten our argument, we will analyze only the term $\mathcal{H}_w[\psi_4\mathcal{H}_w[\Psi]] - \tilde{\mathcal{H}}_w[\tilde{\psi}_4\tilde{\mathcal{H}}_w[\Psi]]$, where $\tilde{\psi}_4$ is obtained from ψ_4 by substitution of $\mathcal{H}\tilde{w}$ for $\mathcal{H}w$. The other terms can be analyzed similarly.

Thus, for brevity's sake, for the rest of this appendix, we define $f = \psi_4 \mathcal{H}_w[\Psi]$ and $\tilde{f} = \tilde{\psi}_4 \tilde{\mathcal{H}}_w[\Psi]$. Since f and \tilde{f} are regular, Lemma 9 gives

$$\left\| \left(\tilde{\mathcal{H}}_{w} \tilde{f} - \mathcal{H}_{w} f \right) w \right\|_{2}$$

$$\prec \left\| \left(\mathcal{H}_{w} \tilde{f} - \mathcal{H}_{w} f \right) w \right\|_{2},$$

This inequality can be continued as

where we have used Lemma 9 in the last step and Appendix C in the previous step, together with the fact that ψ_4 , $\tilde{\psi}_4$, and Ψ are almost surely bounded as $n \to \infty$. The proof is complete.

REFERENCES

[And63] Theodore Wilbur Anderson.

Asymptotic theory for principal component analysis.

The Annals of Mathematical Statistics, 34(1):122-148, 1963.

[BGK16] Florent Benaych-Georges and Antti Knowles.

Lectures on the local semicircle law for Wigner matrices.

arXiv preprint arXiv:1601.04055, 2016.

[BS96] Zhidong Bai and Hewa Saranadasa.

Effect of high dimension: By an example of a two sample problem.

Statistica Sinica, pages 311–329, 1996.

[BS⁺98] Zhi-Dong Bai, Jack W. Silverstein, et al.

No eigenvalues outside the support of the limiting spectral distribution of large-dimensional sample covariance matrices.

The Annals of Probability, 26(1):316-345, 1998.

[BT02] J.S. Bergin and P.M. Techau.

High-fidelity site-specific radar simulation: KASSPER'02 workshop datacube.

Information Systems Laboratories, Inc., Vienna, VA, Technical Report ISL-SCRD-TR-02-105, 2002.

[BY12] Zhidong Bai and Jianfeng Yao.

On sample eigenvalues in a generalized spiked population model.

Journal of Multivariate Analysis, 106:167–177, 2012.

[CL77] C. Carton-Lebrun.

Product properties of Hilbert transforms.

Journal of Approximation Theory, 21(4):356-360, 1977.

[CM14] Romain Couillet and Matthew McKay.

Large dimensional analysis and optimization of robust shrinkage covariance matrix estimators.

Journal of Multivariate Analysis, 131:99-120, 2014.

[CQ10] Song Xi Chen and Ying-Li Qin.

A two-sample test for high-dimensional data with applications to gene-set testing.

The Annals of Statistics, 38(2):808-835, 2010.

[CWH11] Yilun Chen, Ami Wiesel, and Alfred O. Hero.

Robust shrinkage estimation of high-dimensional covariance matrices.

IEEE Transactions on Signal Processing, 59(9):4097–4107, 2011.

[DGJ18] David L. Donoho, Matan Gavish, and Iain M. Johnstone.

Optimal shrinkage of eigenvalues in the spiked covariance model.

Annals of Statistics, 46(4):1742, 2018.

[DS85] Dipak K. Dey and C. Srinivasan.

Estimation of a covariance matrix under Stein's loss.

The Annals of Statistics, pages 1581-1591, 1985.

[HIPS22] Alfred O. Hero III, Neal Patwari, and Kumar Sricharan.

Crawdad umich/rss, 2022.

[JL09] Iain M. Johnstone and Arthur Yu Lu.

On consistency and sparsity for principal components analysis in high dimensions.

Journal of the American Statistical Association, 104(486):682-693, 2009.

[Joh01] Iain M. Johnstone.

On the distribution of the largest eigenvalue in principal components analysis.

Annals of Statistics, pages 295-327, 2001.

[KY17] Antti Knowles and Jun Yin.

Anisotropic local laws for random matrices.

Probability Theory and Related Fields, 169(1):257–352, 2017.

[LAP20a] Haoran Li, Alexander Aue, and Debashis Paul.

High-dimensional general linear hypothesis tests via non-linear spectral shrinkage.

Bernoulli, 26(4), 2020.

[LAP⁺20b] Haoran Li, Alexander Aue, Debashis Paul, Jie Peng, and Pei Wang.

An adaptable generalization of Hotelling's T^2 test in high dimension.

The Annals of Statistics, 48(3):1815-1847, 2020.

[LP11] Olivier Ledoit and Sandrine Péché.

Eigenvectors of some large sample covariance matrix ensembles.

Probability Theory and Related Fields, 151(1-2):233-264, 2011.

[LW04] Olivier Ledoit and Michael Wolf.

A well-conditioned estimator for large-dimensional covariance matrices.

Journal of Multivariate Analysis, 88(2):365-411, 2004.

[LW17a] Olivier Ledoit and Michael Wolf.

Direct nonlinear shrinkage estimation of large-dimensional covariance matrices.

Technical report, Working Paper, 2017.

[LW17b] Olivier Ledoit and Michael Wolf.

Nonlinear shrinkage of the covariance matrix for portfolio selection: Markowitz meets Goldilocks.

The Review of Financial Studies, 30(12):4349-4388, 2017.

[LW18] Olivier Ledoit and Michael Wolf.

Optimal estimation of a large-dimensional covariance matrix under Stein's loss.

Bernoulli, 24(4B):3791-3832, 2018.

[LW20] Olivier Ledoit and Michael Wolf.

Analytical nonlinear shrinkage of large-dimensional covariance matrices.

The Annals of Statistics, 48(5):3043-3065, 2020.

[LW22] Olivier Ledoit and Michael Wolf.

Quadratic shrinkage for large covariance matrices.

Bernoulli, 28(3):1519-1547, 2022.

[ML05] Xavier Mestre and Miguel Ángel Lagunas.

Finite sample size effect on minimum variance beamformers: Optimum diagonal loading factor for large arrays.

IEEE Transactions on Signal Processing, 54(1):69-82, 2005.

[MP67] Vladimir A. Marčenko and Leonid Andreevich Pastur.

Distribution of eigenvalues for some sets of random matrices.

Mathematics of the USSR-Sbornik, 1(4):457, 1967.

[Mui09] Robb J. Muirhead.

Aspects of Multivariate Statistical Theory.

John Wiley & Sons, 2009.

[Nad14] Raj Rao Nadakuditi.

Optshrink: An algorithm for improved low-rank signal matrix denoising by optimal, data-driven singular value shrinkage.

 ${\it IEEE Transactions on Information Theory}, 60 (5): 3002-3018, 2014.$

[NPW21] Jamshid Namdari, Debashis Paul, and Lili Wang.

High-dimensional linear models: A random matrix perspective.

Sankhya A, 83(2):645-695, 2021.

[PWY23] Yifei Pan, Jianfei Wang, and Yu Yan.

On the local regularity of the Hilbert transform.

arXiv preprint arXiv:2308.11947, 2023.

[RFKN92] Frank C. Robey, Daniel R. Fuhrmann, Edward J. Kelly, and Ramon Nitzberg.

A CFAR adaptive matched filter detector.

IEEE Transactions on Aerospace and Electronic Systems, 28(1):208-216, 1992.

[RMH21] Benjamin D. Robinson, Robert Malinas, and Alfred O. Hero.

Space-time adaptive detection at low sample support.

IEEE Transactions on Signal Processing, 69:2939–2954, 2021.

[RML $^+$ 22] Benjamin D. Robinson, Robert Malinas, Van Latimer, Beth Morrison, and Alfred O. Hero. An improvement on the Hotelling T^2 test using the Ledoit-Wolf nonlinear shrinkage estimator. In 2022 30th European Signal Processing Conference (EUSIPCO), pages 2106–2110, 2022.

[RV13] Mark Rudelson and Roman Vershynin.

Hanson-Wright inequality and sub-gaussian concentration.

Electron. Commun. Probab., pages 1-9, 2013.

[SB95] Jack W. Silverstein and Z. D. Bai.

On the empirical distribution of eigenvalues of a class of large dimensional random matrices. *Journal of Multivariate Analysis*, 54(2):175–192, 1995.

[SC⁺95] Jack W. Silverstein, Sang-Il Choi, et al.

Analysis of the limiting spectral distribution of large dimensional random matrices.

Journal of Multivariate Analysis, 54(2):295-309, 1995.

[Sil95] Jack W. Silverstein.

Strong convergence of the empirical distribution of eigenvalues of large dimensional random matri-

Journal of Multivariate Analysis, 55(2):331-339, 1995.

[Ste75] Charles Stein.

Estimation of a covariance matrix, Rietz lecture.

In 39th Annual Meeting IMS, Atlanta, GA, 1975, 1975.

[Ste86] Charles Stein.

Lectures on the theory of estimation of many parameters.

Journal of Soviet Mathematics, 34(1):1373–1403, 1986. [Tyl87] David E Tyler.

A distribution-free M-estimator of multivariate scatter.

The Annals of Statistics, pages 234–251, 1987.