

Sampling models for selective inference

Daniel García Rasines^{1*†} and G. Alastair Young^{2†}

^{1*}Department of Quantitative Methods, CUNEF Universidad, Calle Almansa 101, Madrid, 28040, Spain. ORCID: 0000-0002-1558-5860.

²Department of Mathematics, Imperial College London, Exhibition Road, London, SW7 2AZ, United Kingdom. ORCID: 0000-0002-8333-7981.

*Corresponding author(s). E-mail(s): daniel.garciasines@cunef.edu;
Contributing authors: alastair.young@imperial.ac.uk;

†These authors contributed equally to this work.

Abstract

This paper explores the challenges of constructing suitable inferential models in scenarios where the parameter of interest is determined in light of the data, such as regression after variable selection. Two compelling arguments for conditioning converge in this context, whose interplay can introduce ambiguity in the choice of conditioning strategy: the Conditionality Principle, from classical statistics, and the ‘condition on selection’ paradigm, central to selective inference. We discuss two general principles that can be employed to resolve this ambiguity in some recurrent contexts. The first one refers to the consideration of how information is processed at the selection stage. The second one concerns an exploration of ancillarity in the presence of selection. We demonstrate that certain notions of ancillarity are preserved after conditioning on the selection event, supporting the application of the Conditionality Principle. We illustrate these concepts through examples and provide guidance on the adequate inferential approach in some common scenarios.

Keywords: Selection, inference, conditioning, ancillarity

1 Introduction

Most statistical methodology is designed to provide error guarantees in situations where the objectives of the analysis are specified before collecting the data. In many contemporary problems, however, such idealised settings are the exception rather than

the norm. A more realistic situation is one in which the statistician performs an exploratory analysis of the data before selecting a set of relevant inferential questions to address. It is well known that failing to acknowledge this adaptivity in the subsequent inferences yields the reported error assessments invalid. For example, type I error guarantees of testing procedures are lost. This problem, commonly known as selection bias, has received considerable attention in the past few years, especially in the regression literature. The problem discussed in this paper concerns the appropriate choice of statistical model to conduct inference in the presence of selection.

To fix ideas, let us consider a standard problem concerning selection: the ‘selected mean’ problem. Suppose that Y_1, \dots, Y_n are independent normal observations with different unknown means $\theta_1, \dots, \theta_n$ and known common variance, and that we seek inference for the means of the largest k observations for some $k < n$. A problem like this can arise, for example, in a clinical trial experiment, where n treatments are tested and only the most promising ones are put forward for further investigation. In this situation, reporting standard estimates and confidence intervals for the selected parameters ignoring selection—a face-value approach—will likely lead to flawed conclusions, as the largest observations tend to overestimate their corresponding means. This particular case of selection bias is known in the literature as the winner’s curse. Besides being important in its own right, this problem provides a simple theoretical benchmark for the analysis of more complex scenarios.

A general formulation of selection problems is as follows. Suppose that we have data $Y \in \mathcal{Y}$, whose sampling distribution we model by some parametric family $\{F(y; \theta) : \theta \in \Theta\}$, where $F(y; \theta)$ is the distribution function of Y under θ . We assume throughout that \mathcal{Y} and Θ are both subsets of a Euclidean space, and that all the distributions in the family are dominated by the Lebesgue or counting measures. The corresponding Lebesgue densities or probability mass functions will be denoted by $f(y; \theta)$. In addition, we assume that there exists a set of m potential parameters of interest, $\{\psi_1(\theta), \dots, \psi_m(\theta)\}$, from which at most one is to be selected for inference after observing the data, possibly by an artificially randomised procedure. For example, in the previous problem, the set of potential parameters of interest would contain all the subsets of $\{\theta_1, \dots, \theta_n\}$ of size k . Thus, we assume that there exist functions $p_i : \mathcal{Y} \rightarrow [0, 1]$, $i = 1, \dots, m$, such that, having observed $Y = y$, $\psi_i(\theta)$ is selected for inference with probability $p_i(y)$.

Under this formulation of the problem, at most one of the parameters can be selected, so we will generally drop the subscript and write the selected parameter under consideration simply as $\psi \equiv \psi_i(\theta)$, and the corresponding selection probability as $p(y) \equiv p_i(y)$. The latter function, which encodes all the relevant information regarding the selection mechanism of ψ , will be referred to as the selection function. By randomised selection procedure we mean one for which the decision to select a parameter for inference depends not only on y but also on random noise generated by the statistician. As a consequence, $p(y)$ is not an indicator function. This is typically carried out to increase the power of the subsequent inference (Tian and Taylor, 2018). The following examples illustrate the framework.

Example 1. *File drawer problem.* Let $Y_1, \dots, Y_{n_1} \in \mathbb{R}$ be a random sample from a distribution with scalar parameter θ , and suppose that we use this sample to test

$H_0: \theta = 0$ against $H_a: \theta > 0$, because we are only interested in θ if it is positive. If the test results in rejection, we collect a second set of samples Y_{n_1+1}, \dots, Y_n to obtain more information about θ . At the end, the whole data vector is $Y = (Y_1, \dots, Y_n)^T$. If the testing procedure rejects H_0 whenever $T(Y_1, \dots, Y_{n_1}) > t$ for some statistic T and some threshold t , the selection function is $p(y) = \mathbf{1}\{T(y_1, \dots, y_{n_1}) > t\}$, where $\mathbf{1}\{\cdot\}$ is the indicator function. Within this context, a randomised procedure could be of the form $T(y_1, \dots, y_{n_1}) + W > t$, where W is generated from a known distribution independently of the data, and the resulting selection function would be $p(y) = \mathbb{P}\{T(y_1, \dots, y_{n_1}) + W > t\}$.

Now, suppose that for a certain population under study the true parameter is $\theta = 0$, and that a process like the one described before is carried out by several independent researchers. If only those analyses for which the initial test rejected the null are published, the literature concerning this parameter will be filled with results that overstate the true effect size, leading to the false belief that θ is positive, when in fact it is not. This is a stylised example of what Rosenthal (1979) termed the ‘file drawer effect’, also known as publication bias, which occurs when the decision of whether to make a scientific finding public is influenced by the outcome of the analysis. Another important family of selection problems concerns inference for a regression model after variable selection. This is discussed in greater depth in Section 3.

Example 2. *Regression model.* Let $Y \in \mathbb{R}^n$ be a response vector and $X \in \mathbb{R}^{n \times p}$ be a known, fixed design matrix containing the observed values of p covariates. Suppose that a variable-selection algorithm is used to select a non-empty subset $s \subseteq \{1, \dots, p\}$ of the covariates. We think of s as being the output of some algorithm such as the LASSO or best subset selection applied to (Y, X) . Given s , we may wish to provide inference for the best linear predictor of Y given only the selected covariates, $\psi = \{X(s)^T X(s)\}^{-1} X(s)^T \mathbb{E}_\theta[Y \mid X]$, where $X(s)$ is the submatrix of X containing the observations of the selected covariates. Denoting by $E(s) \subseteq \mathbb{R}^n$ the set of observations for which the variable-selection algorithm would have produced the same set s , we can write the selection function as $p(y) = \mathbf{1}\{y \in E(s)\}$.

Like many other fundamental problems in statistics, selection issues were first studied by R.A. Fisher in Fisher (1934a), who considered the sampling bias arising in albinism studies. The type of problem considered by Fisher is one in which independent samples Y_1, Y_2, \dots from a given population are generated and each sample is recorded or not with probability $w(y_i)$, so that the data available for inference constitutes a random sample from a weighted distribution $g(y; \theta) \propto f(y; \theta)w(y)$. There is a large body of literature dedicated to the study of these models. A good survey is given by Rao (1985).

Although these problems fit within the framework considered here, on defining the selection function $p(y_1, \dots, y_n) = \prod_{i=1}^n w(y_i)$, they are different in nature to the kind of problems studied in this work. Conceptually, the main difference is that in our problems selection refers to the decision of providing or not inference for a given parameter, while in the other case the inferential objective is decided in advance, and selection refers to an inherent bias of the sampling mechanism. Mathematically, the selection functions considered here cannot usually be broken up into marginal selection functions for the different samples: the decision of providing inference for a given

parameter depends on some summary statistic of the full dataset or a subset of it. The implications of this are important. When $p(y_1, \dots, y_n) = \prod_{i=1}^n w(y_i)$ and Y_1, \dots, Y_n are independent, the distribution induced by the selection rule can be factorised into independent components. By contrast, the selection rules considered here induce non-trivial dependencies between the observations, and such dependencies do not vanish when the sample size increases except in trivial circumstances.

As a side remark, in this work we are only concerned with situations where the selection function of the selected parameter is known, at least to a very high degree of precision. Inference with a fully unknown selection function is ill-posed, as the model becomes highly non-identifiable. This problem has been considered by some authors (e.g. Bayarri and DeGroot, 1987) in cases where the selection function is assumed to be of the form $p(y) = \prod_{i=1}^n w(y_i)$.

2 The conditional approach to selective inference

Different approaches to statistical inference lead to two opposing views as to the correct analysis of the data in the presence of selection. On the one hand, frequentist methods evaluate the accuracy of inferential procedures with respect to the sampling distribution of the data at a fixed value of the parameter. Since selection modifies the sampling distribution by favouring data points with higher selection probability, it is clear that the reported accuracy should be appropriately modified. On the other hand, Bayesians typically adopt the view that, once the data has been observed, the recognition that a different realisation could have resulted in a different inferential problem, or in no problem at all, should have no effect on the inference (Dawid, 1994). The latter position, however, has recently been challenged by some authors (Yekutieli, 2012; Harville, 2021; García Rasines and Young, 2022).

Adoption of the first viewpoint leads to adherence to the so-called conditional approach (Fithian et al., 2017), which advocates that inference for the selected ψ should be based on the conditional distribution of the data given selection. Such distribution has density or mass function

$$f_S(y; \theta) = \frac{f(y; \theta)p(y)}{\varphi(\theta)}, \quad \varphi(\theta) = \mathbb{E}_\theta[p(Y)];$$

where the normalising constant $\varphi(\theta)$ is the probability that ψ gets selected when θ is the true parameter. The motivation for basing inference on (2) is that, under repeated sampling from a fixed distribution $F(\cdot; \theta)$, if inferences are only provided for those samples that get selected, the reported error assessments are well-calibrated in a frequentist sense. For example, nominal $1 - \alpha$ confidence sets contain the true parameter at least $(1 - \alpha)\%$ of the times they are reported. In the remainder of this work, we will refer to the conditional distribution of Y given selection as the selective distribution.

An intuitive way to interpret the conditional approach is as a form of information splitting. For a given ψ , let R be the random variable that takes the value 1 if ψ gets selected and 0 otherwise. That is, $R|Y \sim \text{Bernoulli}\{p(Y)\}$. Following Fithian et al. (2017), the data-generating process of Y may be thought of as consisting of two stages.

In the first stage, the value of R , r say, is sampled from its marginal distribution, and in the second, Y is sampled from the conditional distribution $Y \mid r$. Since it is R that determines whether we are going to provide inference for ψ or not, inference based on information revealed in stage two is necessarily free of any selection bias, as it removes the information about the parameter provided by R .

In Kuffner and Young (2018) it is argued that the conditional approach is in fact a modern re-expression of well-established Fisherian arguments. Specifically, the authors consider what they term the *Fisherian proposition*, based on Fisher (1925, 1934b), which states that relevance of the inference to the observed data is achieved by conditioning on certain sample statistics. Typically, these statistics are required to be uninformative about the parameter of interest, though it is generally accepted that this condition can be relaxed in some circumstances (Barndorff-Nielsen and Cox, 1994, Chapter 2).

In a similar vein, we argue that the conditional approach follows from arguments analogous to those underpinning the Conditionality Principle (Birnbaum, 1962), as both advocate conditioning on the random events that have occurred right up to the formulation of the inferential problem. However, the nature of the conditioning differs in each case. In the selective context, it necessarily entails conditioning on statistics that are informative about the parameter of interest, as they provide evidence that the parameter is sufficiently relevant to warrant analysis.

The conditional approach has received considerable attention in recent years, particularly in the context of inference after variable selection. Interest on this approach increased notably following the realization that some widely used selection rules produce simple selection events, like affine sets, which can be handled with relative ease. This line of work began with Lockhart et al. (2014) and has been explored by numerous authors, including Lee and Taylor (2014); Loftus and Taylor (2014); Lee et al. (2016); Fithian et al. (2017); Tibshirani et al. (2018); and Panigrahi et al. (2021). Notable selection rules of this kind include the LASSO and LARS with fixed penalty parameters, stepwise procedures with a fixed number of steps, and multiplicity correction methods such as the Benjamini-Hochberg procedure (Benjamini and Hochberg, 1995). On the other hand, progress on the conditional approach has been hindered by the analytical complexity of many contemporary selection algorithms. This has led to the development of analytical approximations (Panigrahi and Taylor, 2018), simulation-based approaches (Markovic et al., 2019), and randomisation mechanisms (Tian and Taylor, 2018; García Rasines and Young, 2023; Leiner et al., 2023; Neufeld et al., 2024).

3 Sampling models

The conditional approach appears straightforward to apply from a conceptual perspective: if the selection function $p(y)$ is known and one is comfortable with a particular sampling model for the data, then the selective distribution (2) is unequivocally determined. However, in this section we argue that, just as the determination of an appropriate sampling distribution in classical statistics is subject to some degree of

ambiguity, so is, in some circumstances, the appropriate choice of the conditional distribution for selective inference. To motivate the discussion, let us consider the following example.

Example 3. *Random sample size.* Let Y_1, \dots, Y_{n_1} be a random sample from a $N(\theta, 1)$ distribution, where the sample size $n_1 \geq 1$ has been generated from some known distribution $f_{N_1}(n_1)$. Suppose that we use this sample to test whether θ is greater than zero. If we observe $y_1 + \dots + y_{n_1} > 1.96n_1^{1/2}$, say, we collect a second set of observations of fixed size n_2 , Y_{n_1+1}, \dots, Y_n , where $n = n_1 + n_2$, which we use to obtain more information about θ . In the joint model for (N_1, Y_1, \dots, Y_n) , with selection function $p(n_1, y_1, \dots, y_n) = \mathbf{1}(y_1 + \dots + y_{n_1} > 1.96n_1^{1/2})$, the conditional distribution of the full data given selection is

$$f_S(n_1, y_1, \dots, y_n; \theta) = \frac{f_{N_1}(n_1)f(y_1, \dots, y_n | n_1; \theta)p(n_1, y_1, \dots, y_n)}{\sum_{\tilde{n}_1=1}^{\infty} f_{N_1}(\tilde{n}_1)\Phi(\theta\tilde{n}_1^{1/2} - 1.96)}.$$

Note that in this model the sample size is not independent of θ , as its probability mass function is given by

$$f_S(n_1; \theta) = \frac{f_{N_1}(n_1)\Phi(\theta n_1^{1/2} - 1.96)}{\sum_{\tilde{n}_1=1}^{\infty} f_{N_1}(\tilde{n}_1)\Phi(\theta\tilde{n}_1^{1/2} - 1.96)}.$$

The intuitive explanation of this dependence is that, if the true θ is smaller than zero, we have a higher chance of falsely concluding that $\theta > 0$ if we have a less informative sample (i.e. if n_1 is small), and vice versa. Therefore, inference based on this sampling model would make use of the sample size distribution, and would arrive to a different conclusion had the sample size been decided in advance by the statistician rather than random, which appears counter-intuitive. Instead, it is more reasonable to work with the model

$$f_S(n_1, y_1, \dots, y_n; \theta) = \frac{f_{N_1}(n_1)f(y_1, \dots, y_n | n_1; \theta)p(n_1, y_1, \dots, y_n)}{\Phi(\theta n_1^{1/2} - 1.96)}.$$

This model conditions on selection after the sample size has been observed, which therefore remains independent of the parameter after conditioning. This reflects more closely the selection process, since the selection test was designed to achieve a certain significance level conditionally on the observed n_1 . That is, the information about θ in the selection step is interpreted according to the conditional model $Y_1, \dots, Y_{N_1} | N_1$, rather than to the unconditional one.

Example 3 illustrates a key issue about selection models: conditioning on selection can create artificial dependencies in the data that do not reflect the nature of the sampling process. Formally, a statistic A which is distribution constant unconditionally will typically depend on the parameter after conditioning on selection. However, it is clear that whether a sample is selected or not cannot make A informative. To avoid this, we have to ensure that the selective distribution reflects the selection process as closely as possible.

Suppose that, unconditionally, the data Y can be reduced to a minimal sufficient statistic (T, A) , where A is distribution constant. In such cases A is said to be ancillary for θ , and the Conditionality Principle, a formal re-expression of the Fisherian proposition, asserts that the information provided by Y about θ is equivalent to that provided by T given the observed value of A (Birnbbaum, 1962). If the selection process complies with this principle, so that the information about θ in the selection stage is interpreted via the conditional distribution $T | A$, as in Example 3, then it is more reasonable to define the selective distribution by

$$f_S(t, a; \theta) = \frac{f_A(a)f_{T|A}(t | a; \theta)p(t, a)}{\varphi(\theta; a)}$$

rather than

$$f_S(t, a; \theta) = \frac{f_A(a)f_{T|A}(t | a; \theta)p(t, a)}{\varphi(\theta)},$$

where $p(t, a) = \mathbb{E}[p(Y) | t, a]$ denotes the selection function in terms of $(T, A) = (t, a)$, independent of θ by sufficiency, $f_A(a)$ and $f_{T|A}(t | a; \theta)$ are the densities or probability mass functions of A and $T | A$, and $\varphi(\theta; a) = \mathbb{E}_\theta[p(T, A) | a]$ is the selection probability under θ given $A = a$. Ancillary statistics arise, for example, in models with an underlying group structure, such as location models.

Example 4. *Location model.* Let Y_1, \dots, Y_n be a random sample from a location model with density $f(y_i; \theta) = g(y_i - \theta)$, $\theta \in \mathbb{R}$. In this model, the Conditionality Principle dictates that sample evidence about θ should be interpreted via the conditional distribution of $T = \hat{\theta}$ given $A = (Y_1 - \hat{\theta}, \dots, Y_n - \hat{\theta})$, where $\hat{\theta}$ is the maximum likelihood estimator of θ and A is the configuration statistic. The density of $\hat{\theta} | a$ admits the simple expression

$$f_{T|A}(t | a; \theta) = c(\theta, a) \prod_{i=1}^n g(t + a_i - \theta),$$

where $c(\theta, a)$ is a normalising constant. Suppose that θ is selected for inference if the p -value

$$u(T, a) = c(0, a) \int_{\hat{\theta}}^{\infty} \prod_{i=1}^n g(\tilde{t} + a_i) d\tilde{t}$$

is below some level α . Then, the selective density of $(\hat{\theta}, A)$ should be

$$f_S(t, a; \theta) = \frac{f(a)c(\theta, a) \prod_{i=1}^n g(t + a_i - \theta)p(t, a)}{\mathbb{P}_\theta(u(T, a) \leq \alpha | a)},$$

where $p(t, a) = \mathbf{1}\{u(t, a) \leq \alpha\}$.

A similar case can be made about models involving nuisance parameters. Suppose that $\theta = (\psi, \chi)$, where ψ is the potential parameter of interest and χ is a nuisance parameter. In these models it is sometimes possible to identify a minimal sufficient statistic (T, A) such that the distribution of $T | A$ is independent of χ and the distribution of A is independent of ψ , or depends on it in such a way that through observation of A alone no information can be extracted about ψ (Barndorff-Nielsen and Cox, 1994,

Chapter 2). In such cases, a natural extension of the Conditionality Principle asserts that information about ψ should be interpreted via the conditional model $T \mid A$. As before, if the selection process interprets the sample information about ψ via the conditional distribution $T \mid A$, it is more natural to define the selective density as

$$f_S(t, a; \theta) = \frac{f_A(a; \theta) f_{T|A}(t \mid a; \psi) p(t, a)}{\varphi(\psi; a)}$$

rather than

$$f_S(t, a; \theta) = \frac{f_A(a; \theta) f_{T|A}(t \mid a; \psi) p(t, a)}{\varphi(\psi, \chi)},$$

where we are using the same notational conventions as before.

An important setting where this argument applies is when inference is sought for a component of the canonical parameter in a full exponential family. Let the model be given by

$$f(y; \theta) = h(y) \exp \{ \psi s_1(y) + \chi^T s_2(y) - K(\theta) \}.$$

Standard arguments show that the distribution of $s_1(Y) \mid s_2(Y)$ is independent of χ , and even though the distribution of $s_2(Y)$ usually depends on ψ , power considerations based on Neyman-Pearson theory indicate that inference for ψ should be provided via the conditional distribution $s_1(Y) \mid s_2(Y)$ (Young and Smith, 2005, Chapter 7). Moreover, $s_2(Y)$ is uninformative for ψ in a sense made precise by Jørgensen (1994), so the Conditionality Principle applies.

A second important class of examples with this structure that arise frequently in selective inference is the following.

Example 5. *Inference on winners.* Suppose that Y_1, \dots, Y_m are independent observations with densities $f(y_i; \theta_i)$, $i = 1, \dots, m$, and suppose that we want to provide inference for the parameter which produces the largest observation, $\theta_I = \arg \max \{ y_i : i = 1, \dots, m \}$. Assume without loss of generality that $I = 1$, so that $T = Y_1$ and $A = (Y_2, \dots, Y_m)^T$. We can identify two natural sampling mechanisms that are consistent with selection of the first mean. In the first one, the whole vector $(Y_1, \dots, Y_m)^T$ is sampled until Y_1 is observed to be the maximum. The corresponding density of this generative process is

$$f_S(y_1, y_2, \dots, y_m; \theta) = \frac{\prod_{i=1}^m f(y_i; \theta_i) \mathbf{1}(I = 1)}{\mathbb{P}_\theta(Y_1 > Y_i \forall i > 1)}. \quad (1)$$

The second sampling mechanism is that in which $(Y_2, \dots, Y_m)^T$ is sampled from its unconditional distribution, and conditionally on its observed value, Y_1 is sampled until it exceeds $\max\{y_2, \dots, y_m\}$. The density of the data under this model is

$$f_S(y_1, y_2, \dots, y_m; \theta) = \frac{\prod_{i=1}^m f(y_i; \theta_i) \mathbf{1}(I = 1)}{\mathbb{P}_{\theta_1}(Y_1 > Y_i \forall i > 1 \mid y_2, \dots, y_m)}. \quad (2)$$

In the first model, conditioning on selection breaks the independence structure of the data. As a consequence, the observations from the non-selected parameters depend on θ_1 , and the marginal distribution of Y_1 depends on $\theta_2, \dots, \theta_m$. This makes

manipulation of the likelihood function awkward and computationally expensive if m is large, despite the apparent simplicity of the problem. However, one could nevertheless argue that it is appropriate to condition on the observed values of Y_2, \dots, Y_m for either of the following two reasons:

1. If selection is used to determine whether θ_1 is the largest θ_i , since the observed values of Y_2, \dots, Y_m do not provide any direct information about this fact, it appears more reasonable to conduct the selection step conditionally on $a = (y_2, \dots, y_m)^T$. Thus, preference of the second inferential model over the first can be justified without resorting to computational considerations.
2. Even if the previous argument does not apply in a given problem, we might still decide to condition on a on top of the selection event on the basis that the conditional distribution of A given the selection event depends very mildly on the parameter of interest (it is non-informative, in a certain sense), and leads to simpler inferences. We formalise this notion in the following section.

Since the distribution of Y_2, \dots, Y_m depends on θ_1 given selection, conditioning on them, even if advisable on conceptual grounds, is expected to reduce inferential power. We have investigated this empirically in García Rasines and Young (2022), in a situation where the marginal distributions of the Y_i 's are Gaussian; see also Section 3 of (García Rasines and Young, 2021). The main conclusion is that inference based on (1) is slightly more powerful than inference based on the simpler model (2) (as indicated by confidence intervals 5-10% shorter). However, it requires considerably more computational power than the latter and is more sensitive to numerical approximations.

Another important area of application of selective inference involving ancillary statistics is regression after variable selection, which was briefly introduced in Example 2. In this setting, data consists of an n -dimensional response vector Y and a $n \times p$ design matrix X , containing information on p covariates. If p large, either absolutely or relative to n , it is often convenient to reduce the set of covariates to be analysed. This reduction may be conducted to increase the interpretability of the fitted model, for computational or analytical efficiency, or because it is believed that only a few covariates have explanatory power over Y , and by getting rid of the inactive ones higher inferential power can be achieved. Following the notation of Example 2, let $s \subseteq \{1, \dots, p\}$ be the subset of selected covariates and $X(s)$ be the corresponding sub-design matrix.

An ongoing matter of discussion in this context concerns the adequacy of conditioning on X in cases where it has been randomly generated. In standard regression settings, a fixed-design approach can be compellingly justified by appealing to the Conditionality Principle, provided it is reasonable to assume that the distribution of X carries no information about the regression parameter $\mu = E(Y | X)$. By contrast, incorporating a selection step into the analysis introduces additional complexity that must be accounted for. We distinguish the following two scenarios, which lead to two different treatments of the design matrix.

Scenario 1. Sometimes the goal of the selection step is to decide which joint distribution from $\{(Y, X(s)) : s \subseteq \{1, \dots, p\}\}$ is going to be analysed. That is, the selected

parameter of interest is $\psi_s = E[Y | X(s)]$. In these circumstances, conditioning on X before selection is not appropriate, as this would prevent consideration of the different joint distributions. Once we have selected the distribution for inference, the conditional approach dictates that the regression of y on $X(s)$ ought to be carried out conditionally on the selection event $E = \{(y, X) : s(y, X) = s\}$, where $s(y, X)$ is the output of the selection algorithm applied to a generic (y, X) . In general, the conditional distribution of $X(s) | E$ depends on the regression parameter ψ_s , so $X(s)$ would not be ancillary for it and a fixed-design analysis would not be justified. Conditioning on E with a random X is very challenging, as it requires modeling its distribution, and to the best of our knowledge the only viable proposal to achieve validity in this setting is data splitting (Rinaldo et al., 2019).

Scenario 2. A different type of selection problem is as follows. In some cases, we may want to assess the effect of the selected covariates $X(s)$ relative to the full set of covariates X . In this context, standard arguments justify conditioning on X before carrying out selection, thereby leading to a fixed-design approach. The selection event is then $E = \{y : s(y, X) = s\}$, where X is fixed at its observed value. This problem has been extensively studied and there are multiple analytical and computational procedures available solve it, depending on the selection algorithm and the error distribution of $Y | X$ (Lee and Taylor, 2014; Loftus and Taylor, 2014; Lee et al., 2016; Fithian et al., 2017; Tibshirani et al., 2018; Panigrahi et al., 2021; Panigrahi and Taylor, 2018; García Rasines and Young, 2023; Markovic et al., 2019). Typically, the parameter of interest in this context is the so-called projection parameter $\psi_s = [X(s)^T X(s)]^{-1} X(s)^T \mu$, which provides the best linear predictor of Y using only the variables in s , though alternative parameters have been proposed.

4 Ancillarity in selection models with nuisance parameters

In the previous section, we argued that the choice of inferential model should account for whether the information available at the selection stage was analysed conditionally on an ancillary statistic. When this is not the case, statistics that were independent of the selected parameter before selection may become dependent on it after conditioning. This raises the key question of whether conditioning on selection actually renders these statistics informative about the parameter of interest.

The most intuitive definition of ancillarity in the presence of nuisance parameters requires the marginal distribution of the ancillary statistic A to be free of ψ , which as we have just seen does not usually hold in selective distributions. Weaker definitions of ancillarity have been suggested that allow A to depend on ψ , but in such a way that if we only observed A we would not be able to extract any information about ψ . In this section we are going to show that two of these proposed notions of ancillarity are preserved after conditioning on selection, and can thus be invoked to justify conditioning on these statistics in situations not covered by the discussion in Section 3.

Throughout this section we will assume that we can write $\theta = (\psi, \chi)$, that the parameter space can be written as $\Theta = \Theta_1 \times \Theta_2$, where Θ_1 and Θ_2 are the parameter spaces of ψ and χ , and that there exists a minimal sufficient statistic (T, A) for θ

such that the distribution of $T \mid A$ does not depend on χ . Instead of modifying the selective sampling distribution according to the nature of the sampling process, as in the previous section, we now assume that the selective distribution is $f_S(y; \theta) \propto f(y; \theta)p(y)$, and justify the change of inferential distribution differently.

In the previous setting, Godambe (1976) calls A ancillary if it is complete for χ for any fixed value of ψ . This is, if, for all ψ , whenever $\mathbb{E}_{\psi, \chi}[g(A)] = 0$ for all $\chi \in \Theta_2$ and some real-valued measurable function g , it must be the case that $\mathbb{P}_{\psi, \chi}\{g(A) = 0\} = 1$ for all $\chi \in \Theta_2$. We will refer to this type of ancillarity as G -ancillarity. Godambe (1980) showed that if A is G -ancillary, then the distribution of T contains the same information about ψ as the distribution of $T \mid A$, so that the latter can be used for inference about ψ without any loss of information. It is easy to see that G -ancillarity is preserved after conditioning on selection events under minimal assumptions. We have the following result.

Proposition 1. *If the selection mechanism is such that $\varphi(\psi; a) > 0$ for all possible (ψ, a) , then A is G -ancillary in the non-selective model if and only if it is G -ancillary in the selective model.*

Proof. First of all, we have that, for any y_1, y_2 in the support of the selective distribution,

$$\frac{f_S(y_1; \theta)}{f_S(y_2; \theta)} = \frac{f(y_1; \theta) p(y_1)}{f(y_2; \theta) p(y_2)}$$

is free of θ if and only if $f(y_1; \theta)/f(y_2; \theta)$ is free of θ , which, by the minimal sufficiency of (T, A) in the non-selective regime, holds if and only if (T, A) is equal for y_1 and y_2 . This implies that (T, A) is also minimal sufficient for θ in the selective model. Secondly, we need to show that the distribution of $T \mid A$ in the selective model is free of χ . This is trivial, as

$$f_S(t \mid a; \theta) = \frac{p(t, a)}{\varphi(\psi; a)} f(t \mid a; \psi).$$

Now, for A to be G -ancillary in the non-selective model it means that the condition $\mathbb{E}_{\psi, \chi}[g(A)] = 0$ for all χ implies that $\mathbb{P}_{\psi, \chi}\{g(A) = 0\} = 1$ for all χ , given an arbitrary ψ . However, note that

$$\mathbb{E}_{\psi, \chi}[g(A) \mid S] = \frac{\mathbb{E}_{\psi, \chi}[g(A)\varphi(\psi; A)]}{\varphi(\psi, \chi)},$$

so if $\mathbb{E}_{\psi, \chi}[g(A) \mid S] = 0$ we must have that $\mathbb{P}_{\psi, \chi}\{g(A)\varphi(\psi; A) = 0\} = 1$. With the assumption that $\varphi(\psi; a)$ is positive everywhere, this implies that $\mathbb{P}_{\psi, \chi}\{g(A) = 0\} = 1$, so A is G -ancillary in the selective model. The reciprocal claim follows by the same type of argument. \square

The second notion of ancillarity that we will consider stems from the field of noninformation theory, introduced by Barndorff-Nielsen in Barndorff-Nielsen (1976) and extended by Jørgensen (1994). Several types of ancillarity can be derived from this work. In the selective inference context, the most relevant one is M -ancillarity and extensions thereof, all of which follow from the following general definition: if,

for every possible pair (ψ, a) , the model $\{f_A(a; \psi, \chi) : \chi \in \Theta_2\}$ provides a *perfect fit* for the data a , then A is said to be ancillary ψ (Jørgensen, 1994). Different types of ancillarity can then be derived by adopting different definitions of perfect fit.

The rationale behind this definition is as follows: if two models provide a perfect fit for a given observation, then both are equally good at explaining the data, and we cannot assume that one is more likely to have generated the data than the other. Thus, in the previous setting, we cannot compare two values of ψ , ψ_1 and ψ_2 , say, only through observation of $A = a$, because the models $\{f_A(a; \psi_1, \chi) : \chi \in \Theta_2\}$ and $\{f_A(a; \psi_2, \chi) : \chi \in \Theta_2\}$ explain observation of a equally well.

In the case of M -ancillarity, perfect fit is defined as follows (Barndorff-Nielsen, 1976). A family of densities or mass functions $\{g(x; \lambda) : \lambda \in \Lambda\}$ is said to provide a perfect fit for a given observation $x \in \mathcal{X}$ if there exists a $\lambda \in \Lambda$ such that x is an approximate mode of $g(x; \lambda)$. Formally, if

$$\text{for all } \varepsilon > 0 \text{ there exists a } \lambda \in \Lambda \text{ such that } g(x; \lambda) > (1 - \varepsilon) \sup_{\tilde{x} \in \mathcal{X}} g(\tilde{x}; \lambda).$$

One drawback of this definition is that it is not invariant under transformations of the data. For example, the family of exponential distributions does not provide a perfect fit for any positive observation, as all the densities have mode 0. However, one can easily show that after applying a log-transformation the family provides a perfect fit for any real observation. To remove this arbitrariness, we propose to change the definition and require instead that the previous condition holds under some change of variable. This produces a new type of ancillarity, which we term \tilde{M} -ancillarity. We will assume that the distribution of A is continuous for simplicity. Extension to the discrete case is straightforward.

Definition 1. *Let \mathcal{A} be the sample space of A and assume it is open. A is \tilde{M} -ancillary for ψ if, for all $\psi \in \Theta_1$, there exists a smooth bijection $h : \mathcal{A} \rightarrow \mathcal{A}$ such that condition (4) holds for $h(A)$.*

We then have the following proposition.

Proposition 2. *If the selection mechanism is such that $\varphi(\psi; a) > 0$ for all possible (ψ, a) , then A is \tilde{M} -ancillary in the non-selective model if and only if it is \tilde{M} -ancillary in the selective model.*

Proof. The first part of the proof is the same as in Proposition 1. The marginal density of A given selection is

$$f_S(a; \theta) = \frac{\varphi(\psi; a)}{\varphi(\theta)} f_A(a; \theta).$$

Suppose that A is \tilde{M} -ancillary for ψ in the non-selective model. Let ψ be arbitrarily fixed and define $Z_1 = h_1(A)$, and $Z_2 = h_2(Z_1)$, where h_1 is a bijection such that Z_1 satisfies the condition of Definition 1, and h_2 is a bijection with absolute Jacobian determinant $\varphi(\psi; h_1^{-1}(z_1)) |Jh_1^{-1}(z_1)|$, where $|Jh_1^{-1}(z_1)|$ denotes the absolute Jacobian determinant of $h_1^{-1}(z_1)$. By construction, the density of Z_2 given selection is proportional to $f_{Z_1}(h_2^{-1}(z_2); \psi, \chi)$. By assumption, for any z_1 there exists a $\chi, \hat{\chi}(z_1)$, such that z_1 is a mode of $f_{Z_1}(\cdot; \psi, \hat{\chi}(z_1))$. Hence, for all z_2 , $\hat{\chi}\{h_2^{-1}(z_2)\}$ is such that z_2 is a mode of $f_{S, Z_2}(\cdot; \psi, \hat{\chi}\{h_2^{-1}(z_2)\})$, which shows \tilde{M} -ancillarity of A in the selective model. The converse can be proved analogously. \square

The two forms of ancillarity considered in this section can be used to formally justify conditioning on certain sample statistics which depend on the parameter of interest. The simplest scenario is that described in Example 5, where the non-selective distribution of the data is the product of m independent random variables with different underlying parameters. In that example, if the data from the non-selected parameters, $A = (Y_2, \dots, Y_m)^T$, is either complete or is such that the set of possible modes is the whole sample space, and if $\varphi(\psi; a)$ is positive everywhere, then A is respectively G or \tilde{M} -ancillary for ψ in the selective model, and adherence to the Conditionality Principle would require providing inference for the selected parameter conditionally on the data from the non-selected parameters. This occurs, for example, if $Y_i \sim N(\theta_i, 1)$, $\theta_i \in \mathbb{R}$.

This can be generalised to more interesting situations. Consider a situation where $Y \sim N(\theta, I_n)$ follows a p -dimensional normal distribution for some $p > 1$, and inference is provided for a linear function $\psi = \eta^T \theta$. This problem arises, for example, in regression settings with Gaussian errors and a fixed design, when interest lies on the coefficients of the projection parameter $\psi_s = [X(s)^T X(s)]^{-1} X(s)^T \mu$. In this case, if P is a matrix containing $p - 1$ linearly independent vectors orthogonal to η as columns, then $A = P^T Y$ can be easily seen to be both G and \tilde{M} -ancillary for ψ without selection, and therefore also after selection. On defining $T = \eta^T Y$, one can easily show that the distribution of $T | A$ given selection is a truncated one-dimensional Gaussian distribution, which admits an exact analytical characterization provided the selection event is not too complicated. For selection events which can be expressed as a union of polyhedra, analytical characterisations are provided in Lee et al. (2016).

5 Discussion

Conditioning plays a fundamental role in all approaches to statistical inference. In selective inference, the choice of conditioning strategy is particularly important, as it can significantly impact the complexity of the problem and the validity of the results. We have discussed two ways to make this decision. The first relies on carefully assessing how information is processed during the selection stage. The second one exploits refined definitions of ancillarity that work nicely in selection problems.

Ancillarity in the presence of nuisance parameters is an intricate subject, and multiple definitions of it have been proposed, some of which are not preserved by conditioning on selection. A more complete examination of the relation between selection and ancillarity would be interesting.

References

- Barndorff-Nielsen, O.E. 1976. Noninformation. *Biometrika* 63(3): 567–571. <https://doi.org/10.1093/biomet/63.3.567> .
- Barndorff-Nielsen, O.E. and D.R. Cox. 1994. *Inference and asymptotics*. London: Chapman and Hall/CRC.

- Bayarri, M.J. and M.H. DeGroot. 1987. Bayesian analysis of selection models. *J. R. Statist. Soc. D (The Statistician)* 36(2/3): 137–146. <https://doi.org/10.2307/2348506> .
- Benjamini, Y. and Y. Hochberg. 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. R. Statist. Soc. B (Methodological)* 57(1): 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x> .
- Birnbaum, A. 1962. On the foundations of statistical inference. *J. Am. Stat. Assoc.* 57: 269–326. <https://doi.org/10.1080/01621459.1962.10480660> .
- Dawid, A.P. 1994. ‘Selection paradoxes of Bayesian inference’. In T. W. Anderson, K. T. Fang, and I. Olkin (Eds.), *Multivariate analysis and its applications*, Volume 24, United Kingdom, pp. 211–220. Institute of Mathematical Statistics, Monograph Series.
- Fisher, R.A. 1925. Theory of statistical estimation. *Proc. Camb. Phil. Soc.* 22(5): 700–725. <https://doi.org/10.1017/S0305004100009580> .
- Fisher, R.A. 1934a. The effect of methods of ascertainment upon the estimation of frequencies. *Ann. Eugen.* 6: 13–25. <https://doi.org/10.1111/j.1469-1809.1934.tb02105.x> .
- Fisher, R.A. 1934b. Two new properties of mathematical likelihood. *Proc. R. Soc. Lond. A* 144(852): 285–307. <https://doi.org/10.1098/rspa.1934.0050> .
- Fithian, W., D.L. Sun, and J.E. Taylor 2017. ‘Optimal inference after model selection’. arXiv:1410.2597v4.
- García Rasines, D. and G. Young 2021. Bayesian selective inference: non-informative priors. arXiv:2008.04584v2.
- García Rasines, D. and G.A. Young. 2022. Chapter 2 - Bayesian selective inference, In *Advancements in Bayesian Methods and Implementation*, eds. Srinivasa Rao, A.S., G.A. Young, and C. Rao, Volume 47 of *Handbook of Statistics*, 43–65. United Kingdom: Elsevier.
- García Rasines, D. and G.A. Young. 2023. Splitting strategies for post-selection inference. *Biometrika* 110(3): 597–614. <https://doi.org/10.1093/biomet/asac070> .
- Godambe, V.P. 1976. Conditional likelihood and unconditional optimum estimating equations. *Biometrika* 63(2): 277–284. <https://doi.org/10.2307/2335620> .
- Godambe, V.P. 1980. On sufficiency and ancillarity in the presence of a nuisance parameter. *Biometrika* 67(1): 155–162. <https://doi.org/10.2307/2335328> .
- Harville, D.A. 2021. Bayesian inference is unaffected by selection: fact or fiction? *Am. Stat.* 00(0): 1–7. <https://doi.org/10.1080/00031305.2020.1858963> .

- Jørgensen, B. 1994. The rules of conditional inference: is there a universal definition of non-information? *J. Ital. Statistic. Soc.* 3: 355–384. <https://doi.org/10.1007/BF02589024> .
- Kuffner, T.A. and G.A. Young 2018. ‘Principled statistical inference in data science’. In N. Adams, E. Cohen, and Y.-K. Guo (Eds.), *Statistical Data Science*, Singapore, pp. 21–36. World Scientific Publishing.
- Lee, J.D., D.L. Sun, Y. Sun, and J.E. Taylor. 2016. Exact post-selection inference, with application to the lasso. *Ann. Stat.* 44(3): 907–927. <https://doi.org/10.1214/15-AOS1371> .
- Lee, J.D. and J.E. Taylor. 2014. Exact post model selection inference for marginal screening. *Adv. Neural Inf. Process Syst.* 1: 136–144 .
- Leiner, L., B. Duan, L. Wasserman, and A. Ramdas. 2023. Data fission: Splitting a single data point. *JASA* 0(0): 1–12. <https://doi.org/10.1080/01621459.2023.2270748> .
- Lockhart, R., J.E. Taylor, R. Tibshirani, and R. Tibshirani. 2014. A significance test for the lasso. *Ann. Stat.* 42(2): 413–468. <https://doi.org/10.1214/13-AOS1175> .
- Loftus, J.R. and J.E. Taylor 2014. ‘A significance test for forward stepwise model selection’. arXiv:1405.3920v1.
- Markovic, J., J. Taylor, and J. Taylor 2019. Inference after black box selection. arXiv:1901.09973v1.
- Neufeld, A., A. Dharamshi, L. Gao, and D. Witten. 2024. Data thinning for convolution-closed distributions. *JMLR* 25(57): 1–35 .
- Panigrahi, S. and J. Taylor. 2018. Scalable methods for Bayesian selective inference. *Electron. J. Stat.* 12: 2355–2400. <https://doi.org/10.1214/18-EJS1452> .
- Panigrahi, S., J.E. Taylor, and A. Weinstein. 2021. Integrative methods for post-selection inference under convex constraints. *Ann. Stat.* 49(5): 2803 – 2824. <https://doi.org/10.1214/21-AOS2057> .
- Rao, C.R. 1985. Weighted distributions arising out of methods of ascertainment: What population does a sample represent? New York, NY, pp. 543–569. Springer.
- Rinaldo, A., L. Wasserman, and M. G’Sell. 2019. Bootstrapping and sample splitting for high-dimensional, assumption-lean inference. *Ann. Stat.* 47(6): 3438–3469. <https://doi.org/10.1214/18-AOS1784> .
- Rosenthal, R. 1979. The file drawer problem and tolerance for null results. *Psychol Bull.* 86(3): 638–641. <https://doi.org/10.1037/0033-2909.86.3.638> .

- Tian, X. and J.E. Taylor. 2018. Selective inference with a randomized response. *Ann. Stat.* 46(2): 679–710. <https://doi.org/10.1214/17-AOS1564> .
- Tibshirani, R., A. Rinaldo, R. Tibshirani, and L. Wasserman. 2018. Uniform asymptotic inference and the bootstrap after model selection. *Ann. Stat.* 46(3): 1255–1287. <https://doi.org/10.1214/17-AOS1584> .
- Yekutieli, D. 2012. Adjusted Bayesian inference for selected parameters. *J. R. Statist. Soc. B* 74(3): 515–541. <https://doi.org/10.1111/j.1467-9868.2011.01016.x> .
- Young, G.A. and R.L. Smith. 2005. *Essentials of Statistical Inference*. United Kingdom: Cambridge University Press.