

Primordial Black Hole Formation via Inverted Bubble Collapse

Kai Murai,^{1,*} Kodai Sakurai,^{1,†} and Fuminobu Takahashi^{1,‡}

¹*Department of Physics, Tohoku University, Sendai, Miyagi 980-8578, Japan*

We propose a novel mechanism of primordial black hole (PBH) formation through inverted bubble collapse. In this scenario, bubbles nucleate sparsely in an incomplete first-order phase transition, followed by a bulk phase transition in the rest of the universe that inverts these pre-existing bubbles into false vacuum regions. These spherically symmetric false-vacuum bubbles subsequently collapse to form PBHs. Unlike conventional PBH formation mechanisms associated with domain wall collapse or bubble coalescence, our inverted bubble collapse mechanism naturally ensures spherical collapse. We demonstrate that, when applied to the singlet extension of the Standard Model, this mechanism can produce highly monochromatic PBHs with masses up to $\mathcal{O}(10^{-7} - 10^{-5})M_{\odot}$, which potentially explain the microlensing events observed in the OGLE and Subaru HSC data.

Introduction.— The composition of our Universe presents one of the most profound mysteries in modern cosmology. Observational evidence indicates that approximately one-quarter of the energy density of the Universe consists of dark matter, whose nature remains unknown [1–3]. Among various candidates proposed to explain dark matter, primordial black holes (PBHs) stand out as an intriguing possibility [4–8]. While PBHs could potentially account for all of the dark matter, they might also coexist with other dark matter components. Indeed, scenarios in which PBHs constitute a non-negligible fraction or even all of the dark matter have been widely studied. [7, 9–11].

The formation of PBHs requires significant density perturbations in the early Universe. Many proposed mechanisms rely on superhorizon-scale fluctuations, with one widely studied scenario involving the generation of large curvature perturbations during inflation that later reenter the horizon and collapse into PBHs [4–6, 12–15]. However, such scenarios often require fine-tuning of the inflaton potential, particularly in creating ultra-flat regions in the case of single-field inflation models, which imposes non-trivial constraints on inflationary model building. Another possibility is to make use of the large isocurvature perturbations of baryons or dark matter. For instance, PBH formation from collapsing baryon/axion bubbles represents an interesting realization of such a mechanism [16–24].

Alternative mechanisms for generating large overdensities include those involving topological defects such as domain walls [25–35] and first-order phase transitions (FOPTs) [36–52]. However, these scenarios face fundamental challenges, as the overdense regions are expected to deviate significantly from spherical symmetry, and it remains unclear how much deviation is allowed for successful PBH formation (see Refs. [53–55] for the discussion on the PBH formation from Gaussian curvature perturbations). These uncertainties make it challenging to

predict the PBH abundance and understand the collapse process.

In this Letter, we propose a novel mechanism for PBH formation through inverted bubble collapse (IBC) in an incomplete FOPT followed by a bulk phase transition. In our scenario, bubbles sparsely nucleate during an incomplete FOPT. These bubbles are spherically symmetric when nucleated [56], and maintain this symmetry unless they collide. Crucially, before this transition is completed, a second phase transition occurs throughout the Universe except within the bubbles, transforming the bubble interiors into false vacuum regions. Consequently, the initially expanding bubbles begin to contract and eventually collapse. When sufficiently energetic bubbles collapse, they can form PBHs while naturally preserving spherical symmetry. Unlike PBH formation scenarios involving topological defect collapse or bubble coalescence, our mechanism allows for precise prediction of PBH abundance in terms of model parameters thanks to this preserved spherical symmetry.

We demonstrate that our mechanism can be implemented in the context of the electroweak phase transition (EWPT) by extending the Standard Model (SM) with singlet scalar fields. In this setup, an initial (incomplete) FOPT and the subsequent bulk phase transition occur in the singlet sector, followed by the EWPT in the Higgs direction. We show that this scenario can produce highly monochromatic PBHs with masses up to $\mathcal{O}(10^{-7} - 10^{-5})M_{\odot}$, which could potentially explain the microlensing events observed in the OGLE and Subaru HSC data [57–59] and will also be probed by the ongoing Subaru HSC searches for PBHs and other compact objects. Depending on the model parameters, it can also generate PBHs with much smaller masses that could explain all dark matter. Furthermore, if either the bulk phase transition in the singlet sector or the EWPT is of first-order, it could generate gravitational waves detectable by future space-borne interferometers such as LISA [60] (see also Refs. [61–69] for theoretical studies of gravitational waves in the singlet extension).

Lastly, we note that the expanding and shrinking bubbles have also been studied in Ref. [70], which focused on the bubble nucleation during the very early reheating

* kai.murai.e2@tohoku.ac.jp

† kodai.sakurai.e3@tohoku.ac.jp

‡ fumi@tohoku.ac.jp

stage. In their scenario, the shrinking bubbles leave small density perturbations that later collapse to PBHs, which grow via mass accretion during the inflaton matter dominated universe. In contrast, our mechanism operates in the radiation dominated (RD) universe and forms PBHs directly through bubble collapse. Furthermore, our approach allows us to calculate the PBH mass spectrum solely from the phase transition parameters, independent of specific inflation models and reheating processes.

PBH formation in the IBC mechanism.— In this section, we derive the mass distribution of PBHs formed through our IBC mechanism. We first calculate the bubble size distribution, then determine their collapse conditions and derive the mass spectrum of PBHs.

To calculate the bubble size distribution, we first need to track the evolution of bubbles in the cooling universe. We evaluate this at time $t = t_c$, when the bubble walls start to shrink after their interiors have become false vacuum regions through the bulk phase transition. Note that t_c is generically later than the bulk phase transition that occurs at $t = t_{PT}$.

Well before the bubble percolation, the number density of the bubbles, $n_b(t)$, satisfies

$$\frac{dn_b}{dt} = -3Hn_b + \Gamma(T(t)), \quad (1)$$

where $\Gamma(T)$ is the bubble nucleation rate per unit volume, which depends on the cosmic temperature T and grows exponentially as the universe cools down. The solution to this equation is

$$n_b(t)a^3(t) = e^{\int^t dt' a^3(t')\Gamma(T(t'))}. \quad (2)$$

With this solution for the bubble number density, we can derive the size distribution. A key ingredient is the relation between a bubble's nucleation time t_n and its size R at $t = t_c$. Assuming that each bubble expands with a constant velocity v after nucleation, the size at $t = t_c$ is given by

$$\begin{aligned} R(t_c; t_n) &= a(t_c) \left(\frac{R_{\text{in}}}{a(t_n)} + \int_{t_n}^{t_c} d\tilde{t} \frac{v}{a(\tilde{t})} \right) \\ &= \sqrt{\frac{t_c}{t_n}} R_{\text{in}} + 2vt_c \left[1 - \left(\frac{t_n}{t_c} \right)^{1/2} \right], \end{aligned} \quad (3)$$

where a is the scale factor, and R_{in} is the physical critical radius at the bubble nucleation. In the second equality, we used $a \propto t^{1/2}$, as we focus on PBH formation during the RD era in this Letter. As a result, the size distribution of the bubble at $t = t_c$ is given by

$$\begin{aligned} \frac{dn_b}{dR}(t_c) &= \left| \frac{dt_R}{dR(t_c; t_R)} \right| \frac{1}{a(t_c)^3} \frac{dn_b a^3}{dt}(t_R) \\ &= \frac{\Gamma(T_R) t_R^2}{v t_c^2} \left(1 + \frac{R_{\text{in}}}{2vt_R} \right)^{-1}, \end{aligned} \quad (4)$$

where t_R is the nucleation time of a bubble whose radius is R at $t = t_c$, i.e., t_R is the time satisfying $R(t_c; t_R) = R$,

and we define $T_R \equiv T(t_R)$. Typically, the contribution of R_{in} in Eq. (3) becomes negligibly small soon after bubble nucleation, so it can be dropped. Then, we obtain

$$t_R \simeq t_c \left(1 - \frac{R}{2vt_c} \right)^2, \quad (5)$$

and

$$\frac{dn_b}{dR}(t_c) \simeq \frac{\Gamma(T(t_R))}{v} \left(1 - \frac{R}{2vt_c} \right)^4. \quad (6)$$

Note that the bubble radius is bounded by the sound horizon as

$$R \lesssim 2vt_c. \quad (7)$$

Next, we discuss the PBH formation from a false vacuum bubble. Here, we denote the potential energy difference between the inside and outside of the bubble by ΔV , which we treat as constant during the relevant timescale of bubble collapse. In realistic situations, ΔV depends on temperature, as we will see later; however, for simplicity, we treat it as a constant in the following analysis.

Let us assume that the radius of the bubble is R when the bubble wall starts to shrink after its interior transitions to the false vacuum. Then, the energy of the bubble can be approximately expressed as

$$M_b \simeq \frac{4\pi}{3} R^3 \Delta V. \quad (8)$$

Under certain situations, when the energy difference is sufficiently large, the bubble wall can undergo runaway acceleration even in cases with plasma dissipation, whereas in the absence of dissipation, runaway is naturally achieved [71, 72]. Then, as the bubble shrinks, a part of the vacuum energy swept by the bubble wall is accumulated in the kinetic energy of the wall, and this energy gets progressively concentrated inside a smaller region. In such runaway cases, we introduce an efficiency parameter $\epsilon (\leq 1)$ to characterize the fraction of vacuum energy converted into the wall's kinetic energy. Then, if the bubble size becomes smaller than the Schwarzschild radius, $R_s = 2G\epsilon M_b$, the bubble collapses into a PBH. Considering that the bubble can be as small as the width of the bubble wall, δ , the condition for the PBH formation is given by¹

$$R_s = 2G\epsilon M_b \simeq \frac{\epsilon R^3 \Delta V}{3M_{\text{Pl}}^2} \gtrsim \delta. \quad (9)$$

Considering the upper bound of the bubble radius, we obtain the possible range of the PBH mass as

$$M_{\text{min}} \equiv \frac{\delta}{2G} \lesssim M \lesssim M_{\text{max}} \equiv \frac{32\pi}{3} \epsilon v^3 t_c^3 \Delta V. \quad (10)$$

¹ In fact, the bubble wall width becomes thinner due to the Lorentz contraction as it gets accelerated. The wall width δ here is assumed to include this effect.

Since the bubble size never exceeds the particle horizon, the false vacuum energy inside the bubble does not dominate the universe before collapsing into a PBH. Thus, $\Delta V \lesssim 3M_{\text{Pl}}^2 H^2$ holds until the PBH is formed. As a result, we obtain the condition for ΔV given by

$$\frac{3M_{\text{Pl}}^2 \delta}{\epsilon R^3} \lesssim \Delta V \lesssim 3M_{\text{Pl}}^2 H_c^2 = \frac{\pi^2 g_* T_c^4}{30}, \quad (11)$$

where H_c and T_c denote the Hubble parameter and temperature when $t = t_c$, respectively. In the following, we assume that M_{min} is much smaller than the mass region of interest, which is a condition typically satisfied in realistic situations.²

Taking into account the size distribution of the bubbles, we obtain the mass function of the PBHs given by

$$\begin{aligned} \frac{d\rho_{\text{PBH}}}{d \ln M}(t) &= M \frac{dn_b}{dR}(t_c) \frac{dR}{d \ln M_b} \left(\frac{a(t_c)}{a(t)} \right)^3 \Theta(M - M_{\text{min}}) \\ &= \frac{MR \Gamma(T_R)}{3} \frac{t_R^2}{v} \frac{t_c^2}{t_c^2} \left(\frac{a(t_c)}{a(t)} \right)^3 \Theta(M - M_{\text{min}}), \end{aligned} \quad (12)$$

where R is related to M_b via Eq. (8), $M = \epsilon M_b$, and t represents the time after the PBH formation.

Next, we evaluate the PBH mass function assuming the temperature dependence of the bubble nucleation rate. The bubble nucleation rate is considered to increase progressively toward the bulk phase transition that occurs at temperature $T = T_{\text{PT}} (\geq T_c)$, and then effectively drops down to zero afterwards. Thus, around the bulk phase transition, it can be parametrized as³

$$\Gamma(T) \simeq c H_{\text{PT}}^4 e^{-\alpha \frac{T - T_{\text{PT}}}{T_{\text{PT}}}} \quad \text{for } T \geq T_{\text{PT}}, \quad (13)$$

where H_{PT} is the Hubble parameter at the bulk phase transition, and c and α are constant parameters. From Eqs. (5) and (8), we obtain

$$\begin{aligned} T_R &= \sqrt{\frac{t_c}{t_R}} T_c \simeq \left(1 - \frac{R}{2vt_c} \right)^{-1} T_c \\ &\simeq \left[1 - \frac{H_c}{v} \left(\frac{3M}{4\pi\epsilon\Delta V} \right)^{1/3} \right]^{-1} T_c. \end{aligned} \quad (14)$$

² The width of the bubble wall in the rest frame is typically estimated by the inverse of the mass scale of the field driving the phase transition. For instance, if we take the mass scale to be 100 GeV, we obtain $M_{\text{min}} \simeq 6 \times 10^{-22} M_{\odot} / \gamma$ with γ being the Lorentz factor of the wall.

³ We require that the FOPT via bubble nucleation should not occur inside the bubbles, as it would spoil the spherical symmetry and reduce the energy inside them. This condition is met in the case of the singlet extension of the SM studied in the next section.

Then, we obtain the energy ratio to dark matter by

$$\begin{aligned} \frac{df_{\text{PBH}}}{d \ln M} &\equiv \frac{1}{\rho_{\text{DM}}(t)} \frac{d\rho_{\text{PBH}}}{d \ln M}(t) \\ &\simeq \frac{1}{0.44 \text{ eV}} \frac{45}{2\pi^2 g_{*s}(T_c) T_c^3} \frac{c M H_{\text{PT}}^4}{3v} \left(\frac{3M}{4\pi\epsilon\Delta V} \right)^{1/3} \\ &\quad \times \left[1 - \frac{H_c}{v} \left(\frac{3M}{4\pi\epsilon\Delta V} \right)^{1/3} \right]^4 \\ &\quad \times \exp \left[-\alpha \left(\frac{T_c}{T_{\text{PT}}} \left[1 - \frac{H_c}{v} \left(\frac{3M}{4\pi\epsilon\Delta V} \right)^{1/3} \right]^{-1} - 1 \right) \right] \end{aligned} \quad (15)$$

for M satisfying $M_{\text{min}} < M < M_{\text{max}}$ and $T_R > T_{\text{PT}}$. Here, we used $\rho_{\text{DM}}/s = 0.44 \text{ eV}$ with s being the entropy density.

In Fig. 1, we show the predicted PBH abundance in our scenario. We use the parameter sets (a) and (b) shown in Tab. I for the red and blue solid lines, respectively. These parameter sets correspond to a constant bubble nucleation rate ($\alpha = 0$), fast bubble expansion ($v = 1$), and efficient accumulation of the false vacuum energy ($\epsilon = 1$). For comparison, we also show the results for the cases with $\alpha = 10$ (red dashed line) and with $v = 0.5$ (red dot-dashed line) based on the parameter set (a). Note that v , ϵ , and ΔV appear in Eq. (15) in the form of $v(\epsilon\Delta V)^{1/3}$, and the changes of their values are degenerate in $df_{\text{PBH}}/d \ln M$. In these cases, the mass distribution follows $df_{\text{PBH}}/d \ln M \propto M^{4/3}$ for $M \ll M_{\text{max}}$ and has a cutoff at $M = M_{\text{max}}$. On the other hand, the PBH mass function for the parameter set (b) has a cutoff on smaller mass. This is because all the bubbles continue to expand at least between t_{PT} and t_c , and there is a minimal bubble radius at $t = t_c$, leading to the lower bound on the PBH mass. As we will see below, the PBH mass distribution approaches a monochromatic distribution as t_c is significantly delayed from t_{PT} . Setting $T_{\text{PT}} = T_c$ in parameter set (b) yields the same mass function for a specific choice of c , but without the cutoff for small masses (blue dashed line).

Singlet Extended Model.— In this section, we apply the IBC mechanism to the singlet extension of the SM where we introduce two real singlets S_1 and S_2 . The scalar potential is given by

$$\begin{aligned} V(\Phi, S_1, S_2) &= -m_{\Phi}^2 |\Phi|^2 + \lambda |\Phi|^4 + \mu_{\Phi 1} |\Phi|^2 S_1 \\ &\quad + \lambda_{\Phi 1} |\Phi|^2 S_1^2 + t_1 S_1 - m_1^2 S_1^2 + \mu_1 S_1^3 \\ &\quad + \lambda_1 S_1^4 - m_2^2 S_2^2 + \lambda_2 S_2^4 + \lambda_{\Phi 2} |\Phi|^2 S_2^2 \\ &\quad + \lambda_{12} S_1^2 S_2^2 + \mu_{12} S_1 S_2^2, \end{aligned} \quad (16)$$

where Φ denotes the Higgs doublet, and the potential respects the Z_2 symmetry acting on S_2 .⁴ In our IBC

⁴ The collider phenomenology in the extension of the SM with two real singlet scalars with a $Z_2 \times Z_2'$ discrete symmetry are studied in Ref. [78].

TABLE I. Parameter sets for the red and blue solid lines in Fig. 1.

Set	$T_{\text{PT}} [\text{GeV}]$	$T_c [\text{GeV}]$	$\Delta V/\rho_{\text{tot}}(T_c)$	v	ϵ	c	α
(a)	15	15	0.1	1	1	2.0×10^{-8}	0
(b)	3.0×10^4	2.8×10^4	0.1	1	1	1.7×10^{-11}	0




FIG. 1. PBH fraction in dark matter, $df_{\text{PBH}}/d \ln M$. The red and blue solid lines correspond to the parameter sets (a) and (b) in Table I, respectively. The red dashed and red dot-dashed lines correspond to modifications of parameter set (a), where we set $\alpha = 10$ and $v = 0.5$, respectively. For the blue dashed line, we set T_{PT} equal to T_c in the parameter set (b) and tune the parameter c so that it overlaps with the blue solid line for larger masses. The green dots represent the PBH fraction f_{PBH} in the singlet-extended model, with the dotted line showing its variation with the triple coupling of S_1 . The orange and light blue regions are obtained assuming the lensing event observed in the HSC M31 observation [58] and OGLE [57, 59] are due to PBHs, respectively. The purple region is the joint posterior for them [73]. The gray lines represent the observational upper bounds on the PBH abundance by EROS [74], HSC [58], OGLE-III+OGLE-IV [75], OGLE-IV high cadence survey [76]. The evaporation constraint is taken from PBH bounds [11, 77].

mechanism, PBH formation is related to phase transitions in the directions of S_1 and S_2 before the EWPT happens. Thus, we analyze the scalar potential with the CP-even component of Φ set to be $\phi = 0$. While the required late EWSB can be realized for a certain choice of the model parameters [65, 68], we focus on the phase transition dynamics in the singlet sector for our purpose.

We choose the parameters in the scalar potential such that the phase transition proceeds in two stages. First, the S_1 field undergoes a FOPT with bubble nucleation, where the S_1 field takes a larger value inside the bubbles than outside. Subsequently, S_2 in the bulk (outside the bubbles) undergoes a phase transition at time $t = t_{\text{PT}}$, before the completion of the FOPT in the S_1 direction. While the vacuum state with a non-zero vacuum expectation value (VEV) of S_2 in the bulk is not necessarily the true vacuum at t_{PT} , it evolves into the true vacuum

by $t = t_c$. At this point, the bubble interior, where the S_2 remains in its symmetric phase, becomes a false vacuum, triggering the bubbles to collapse and form PBHs within a few Hubble times. Note that the evolution of the shrinking bubble likely exhibits runaway behavior because the phase transition occurs solely in the singlet sector [72].

When t_c is significantly delayed beyond t_{PT} , the resultant PBH mass tends to be larger for two reasons: the continued bubble expansion increases the net false vacuum energy inside, and the collapse exhibits more runaway behavior, leading to a larger efficiency parameter ϵ . Additionally, as t_c is delayed, the physical size of bubbles asymptotically approaches the sound horizon size at $t = t_c$. This behavior occurs because no new bubbles form after t_{PT} , and existing bubbles gradually approach the sound horizon. As a consequence, these similarly-sized bubbles collapse to form PBHs with highly monochromatic mass distribution, resulting in a sharp mass peak.

The PBH abundance shown in Fig. 1 (represented by green dots and dotted line) corresponds to the parameter choice $v = 246.2 \text{ GeV}$, $m_h = 125.09 \text{ GeV}$, $m_{H_1} = 223.1 \text{ GeV}$, $m_{H_2} = 246 \text{ GeV}$, $v_2 = 70 \text{ GeV}$, $\theta_1 = \theta_2 = \theta_3 = 0$, $\lambda_1 = 0.335$, $\lambda_{12} = 0.5$, $\lambda_{\Phi 1} = -0.14$, $\mu_1 = [-190.49, -190.48] \text{ GeV}$, where v is the EW VEV, m_h is the mass of SM-like Higgs boson, m_{H_i} ($i = 1, 2$) is the mass of heavy Higgs bosons, and θ_j ($j = 1, 2, 3$) is the mixing angle for the CP-even Higgs bosons. Due to the extremely narrow mass spectrum in this model, we represent the PBH fraction f_{PBH} by green dots in the figure (using the same numerical scale as $df_{\text{PBH}}/d \ln M$ indicated on the vertical axis), with its variation with the singlet triple coupling shown by the dotted lines (see Appendix A for the detailed spectrum). We have used CosmoTransitions [79] to evaluate the tunneling rate and ΔV in this model. For instance, we obtain $T_{\text{PT}} \approx 221 \text{ GeV}$ and $T_c \approx 46 \text{ GeV}$ for the lower green dot in Fig. 1. The existence of such viable parameters in this simple setup suggests that our PBH formation mechanism can be naturally realized in a wide class of models with multi-step phase transitions.

Discussion.— For successful PBH formation in our IBC scenario, the scalar potential parameters must be set appropriately; the FOPT should remain incomplete, as bubble collisions would break spherical symmetry and complicate PBH formation. If PBHs constitute a significant fraction of dark matter, this parameter constraint may be explained by anthropic selection; a completed FOPT would trap our universe in a false vacuum, triggering old inflation and diluting baryons, while without

a FOPT, no PBH dark matter would form.

One of the key elements of our IBC mechanism is the runaway behavior of the collapsing bubbles. There is currently ongoing debate about establishing a definite criterion for the runaway bubble dynamics, but the bubbles in the singlet sector are likely the ones that exhibit the runaway behavior. Even if the bubble dynamics does not exhibit this runaway behavior, we may have successful PBH formation by introducing a short period of inflation [80–82] after the bubble nucleation. Then, the overdense regions in the interior of the bubbles are stretched beyond the horizon, and they could collapse into PBHs when the bubbles reenter the horizon, if the density perturbations are sizable. This is conceptually related to the PBH formation using vacuum bubbles [83, 84] (see also Ref. [85]).

Furthermore, while this work focuses on pre-EWPT phase transitions in the singlet sector, similar IBC mechanisms might operate during or after the EWPT in singlet or hidden sectors, potentially triggered by the EWPT itself. These scenarios offer additional pathways for PBH formation worthy of future investigation. Also, it is an interesting question if the baryogenesis can be incorporated in our example of the singlet-extended model, especially in a case where the bulk phase transition or the EWPT is of first order. Investigation of baryogenesis scenarios [86, 87] within this framework is left for future investigation.

ACKNOWLEDGMENTS

We thank Katsuya Hashino for useful discussions, and Wen-Yuan Ai for informing us of Ref. [70]. This work is supported by JSPS Core-to-Core Program (grant number: JPJSCCA20200002) (F.T.), JSPS KAKENHI Grant Numbers 20H01894 (F.T.), 20H05851 (F.T.), 23KJ0088 (K.M.), 24K17039 (K.M.), and 23KJ0086 (K.S.). This article is based upon work from COST Action COSMIC WISPerS CA21106, supported by COST (European Cooperation in Science and Technology).

Appendix A: Singlet Extension Model for EWPT

In this Appendix, we give a more detailed explanation of the singlet extension model given by Eq. (16). The component fields of Φ and S_i are given by

$$\Phi = \begin{bmatrix} G^+ \\ \frac{1}{\sqrt{2}}(v + \phi + iG^0) \end{bmatrix}, S_i = v_i + s_i \quad (i = 1, 2), \quad (\text{A1})$$

where $v \simeq 246$ GeV and v_i are the VEVs of Φ and S_i , respectively, and G^+ and G^0 are Nambu-Goldstone (NG) bosons that are absorbed by the weak gauge bosons. The shift of S_1 , $S_1 \rightarrow S_1 + v'_1$, does not change physics [88]. Using this shift transformation, we take $v_1 = 0$ in the following discussion. The CP-even scalar states ϕ and s_i

mix with each other. The mass matrix in the weak basis (ϕ, s_1, s_2) is given by

$$M^2 = \begin{pmatrix} M_{\phi\phi}^2 & M_{\phi s_1}^2 & M_{\phi s_2}^2 \\ M_{\phi s_1}^2 & M_{s_1 s_1}^2 & M_{s_1 s_2}^2 \\ M_{\phi s_2}^2 & M_{s_1 s_2}^2 & M_{s_2 s_2}^2 \end{pmatrix}, \quad (\text{A2})$$

$$M_{\phi\phi}^2 = 2\lambda v^2, \quad M_{\phi s_1}^2 = \mu_{\Phi 1} v, \quad M_{\phi s_2}^2 = 2\lambda_{\Phi 2} v v_2, \quad (\text{A3})$$

$$M_{s_1 s_1}^2 = -2m_S^2 + \lambda_{\Phi 1} v^2 + 2\lambda_{12} v_2^2, \quad (\text{A4})$$

$$M_{s_1 s_2}^2 = 2\mu_{12} v_2, \quad M_{s_2 s_2}^2 = 8\lambda_2 v_2^2, \quad (\text{A5})$$

This mass matrix is diagonalized by the following orthogonal transformation:

$$\begin{pmatrix} \phi \\ s_1 \\ s_2 \end{pmatrix} = R_\theta \begin{pmatrix} h \\ H_1 \\ H_2 \end{pmatrix}, \quad (\text{A6})$$

$$R_\theta = \begin{pmatrix} c_1 c_2 & -s_1 c_2 & -s_2 \\ s_1 c_3 - c_1 s_2 s_3 & c_1 c_3 + s_1 s_2 s_3 & -c_2 s_3 \\ c_1 s_2 c_3 + s_1 s_3 & c_1 s_3 - s_1 s_2 c_3 & c_2 c_3 \end{pmatrix}, \quad (\text{A7})$$

with the shorthanded notations $s_i = \sin \theta_i$, and $c_i = \cos \theta_i$ ($i = 1, 2, 3$). Thus, the masses of H_i and h are obtained by

$$\text{diag}(m_h^2, m_{H_1}^2, m_{H_2}^2) = R_\theta M^2 R_\theta^T. \quad (\text{A8})$$

Through the relations (A8) and the tadpole conditions of the scalar potential (16), the potential parameters m_{Φ}^2 , m_i^2 ($i = 1, 2$), t_1 , λ , λ_2 , $\lambda_{\Phi 2}$, $\mu_{\Phi 1}$ and μ_{12} can be expressed in terms of the physical masses and mixing parameters as

$$m_{\Phi}^2 = \lambda v^2 + \lambda_{\Phi 2} v_2^2, \quad (\text{A9})$$

$$m_1^2 = -\frac{1}{2} m_h^2 s_1^2 c_2^2 - \frac{1}{2} m_{H_1}^2 (s_1 s_2 s_3 + c_1 c_3)^2 - \frac{1}{2} m_{H_2}^2 (s_1 s_2 c_3 - c_1 s_3)^2 + \frac{\lambda_{\Phi 1} v^2}{2} + \lambda_{12} v_2^2, \quad (\text{A10})$$

$$m_2^2 = \frac{\lambda_{\Phi 2} v^2}{2} + 2\lambda_2 v_2^2, \quad (\text{A11})$$

$$t_1 = \frac{1}{2} (-\mu_{\Phi 1} v^2 - 2\mu_{12} v_2^2), \quad (\text{A12})$$

$$\lambda = \frac{m_h^2 c_1^2 c_2^2}{2v^2} + \frac{m_{H_1}^2 (s_1 c_3 - c_1 s_2 s_3)^2}{2v^2} + \frac{m_{H_2}^2 (c_1 s_2 c_3 + s_1 s_3)^2}{2v^2}, \quad (\text{A13})$$

$$\lambda_2 = \frac{m_h^2 s_2^2}{8v_2^2} + \frac{m_{H_1}^2 c_2^2 s_3^2}{8v_2^2} + \frac{m_{H_2}^2 c_2^2 c_3^2}{8v_2^2}. \quad (\text{A14})$$




FIG. 2. Contours of the effective potential in the (s_2, s_1) plane at temperatures $T = 30, 100, 220,$ and 600 GeV. The red points indicate the global minimum at each temperature.

$$\lambda_{\Phi 2} = -\frac{m_h^2 c_1 s_2 c_2}{2vv_2} + \frac{m_{H_1}^2 c_2 (c_1 s_2 s_3^2 - s_1 s_3 c_3)}{2vv_2} + \frac{m_{H_2}^2 c_2 (c_1 s_2 c_3^2 + s_1 s_3 c_3)}{2vv_2}, \quad (\text{A15})$$

$$\mu_{\Phi 1} = -\frac{m_h^2 s_1 c_1 c_2^2}{v} + \frac{m_{H_1}^2}{8v} \left[2s_1 c_1 \{2c_2^2 - (-s_2^2 + c_2^2 - 3)(c_3^2 - s_3^2)\} - 8s_2 s_3 c_3 (c_1^2 - s_1^2) \right] + \frac{m_{H_2}^2}{8v} \left[8s_2 s_3 c_3 (c_1^2 - s_1^2) + 2s_1 c_1 \{(-s_2^2 + c_2^2 - 3)(c_3^2 - s_3^2) + 2c_2^2\} \right], \quad (\text{A16})$$

$$\mu_{12} = \frac{m_h^2 s_1 s_2 c_2}{2v_2} - \frac{m_{H_1}^2 c_2 (s_1 s_2 s_3^2 + c_1 s_3 c_3)}{2v_2} + \frac{m_{H_2}^2 c_2 (c_1 s_3 c_3 - s_1 s_2 c_3^2)}{2v_2}, \quad (\text{A17})$$

We choose the following twelve parameters as model input parameters:

$$v, \quad m_h, \quad v_2, \quad m_{H_1}, \quad m_{H_2}, \quad \theta_i \quad (i = 1, 2, 3), \quad (\text{A18})$$

$$\mu_1, \quad \lambda_1, \quad \lambda_{\Phi 1}, \quad \lambda_{12},$$

where the EW VEV and the SM-like Higgs boson mass are fixed to be $v = 246.2$ GeV, $m_h = 125.09$ GeV.

From the potential (16), one can calculate the effective potential, which consists of the three contributions, i.e., $V_{\text{eff}}(\phi, s_i, T) = V_0(\phi, s_i) + V_{\text{CW}}(\phi, s_i) + V_T(\phi, s_i, T)$ ($i = 1, 2$), where $V_0(\phi, s_i)$ is the tree-level potential, $V_{\text{CW}}(\phi, s_i)$ is Coleman-Weinberg potential up to the one-loop level [89], and $V_T(\phi, s_i, T)$ is the finite-temperature effect of the effective potential [90]. The tree-level contribution V_0 can be straightforwardly obtained from the potential (16) with the replacement of $(\Phi, S_i) \rightarrow (\phi, s_i)$. The remaining parts V_{CW} and V_T depend on the field-dependent masses $\bar{m}_k^2(\phi, s_i)$ of the model. In terms of the field-dependent masses, V_{CW} and V_T are given by

$$V_{\text{CW}} = \sum_k (-1)^{F_k} \frac{n_k}{64\pi^2} \bar{m}_k^4(\phi, s_i) \times \left(\log \frac{\bar{m}_k^2(\phi, s_i)}{\mu_R^2} - c_k \right), \quad (\text{A19})$$

$$V_T = \frac{T^4}{2\pi^2} \sum_k n_k J_{\mp} \left(\frac{\bar{m}_k(\phi, s_i)}{T} \right), \quad (\text{A20})$$

where n_k is the number of degrees of freedom of the particle species and the constant c_k is $3/2$ ($5/6$) for scalar bosons and fermions (gauge bosons). The index F_k is 0 (1) for bosons (fermions). The function $J_{\mp}(x)$ is given

by

$$J_{\mp}(x) = \pm \int_0^{\infty} dy y^2 \log \left(1 \mp e^{-\sqrt{y^2+x^2}} \right), \quad (\text{A21})$$

with the minus (plus) sign being for the bosons (fermions). Renormalization of the Coleman-Weinberg potential is performed in the $\overline{\text{MS}}$ scheme, by which the renormalization scale μ_R is introduced in the effective potential.⁵ In our analysis, we set $\mu_R = m_t$.

In the following, we list field-dependent masses of the scalar bosons in the model with two real singlet scalars. For the CP-even Higgs bosons, from the tree-level effective potential $V_0(\phi, s_i)$, one obtains the field-dependent mass matrix

$$\bar{\mathcal{M}}^2 = \begin{pmatrix} \bar{\mathcal{M}}_{\phi\phi}^2 & \bar{\mathcal{M}}_{\phi 1}^2 & \bar{\mathcal{M}}_{\phi 2}^2 \\ \bar{\mathcal{M}}_{\phi 1}^2 & \bar{\mathcal{M}}_{11}^2 & \bar{\mathcal{M}}_{12}^2 \\ \bar{\mathcal{M}}_{\phi 2}^2 & \bar{\mathcal{M}}_{12}^2 & \bar{\mathcal{M}}_{22}^2 \end{pmatrix}, \quad (\text{A22})$$

$$\bar{\mathcal{M}}_{\phi\phi}^2 = 3\lambda\phi^2 - m_{\Phi}^2 + \lambda_{\Phi 1}s_1^2 + \mu_{\Phi 1}s_1 + \lambda_{\Phi 2}s_2^2, \quad (\text{A23})$$

$$\bar{\mathcal{M}}_{\phi 1}^2 = \phi(\mu_{\Phi 1} + 2\lambda_{\Phi 1}s_1), \quad \bar{\mathcal{M}}_{\phi 2}^2 = 2\lambda_{\Phi 2}s_2\phi, \quad (\text{A24})$$

$$\begin{aligned} \bar{\mathcal{M}}_{11}^2 &= \lambda_{\Phi 1}\phi^2 - 2m_S^2 + 12\lambda_1s_1^2 \\ &\quad + 6\mu_Ss_1 + 2\lambda_{12}s_2^2, \end{aligned} \quad (\text{A25})$$

$$\bar{\mathcal{M}}_{12}^2 = 2s_2(\mu_{12} + 2\lambda_{12}s_1), \quad (\text{A26})$$

$$\begin{aligned} \bar{\mathcal{M}}_{22}^2 &= \lambda_{\Phi 2}\phi^2 - 2m_2^2 + 2\lambda_{12}s_1^2 \\ &\quad + 2\mu_{12}s_1 + 12\lambda_2s_2^2. \end{aligned} \quad (\text{A27})$$

Diagonalizing it, one obtains field-dependent masses of the CP-even Higgs bosons, i.e., $\bar{m}_h^2, \bar{m}_{H_1}^2, \bar{m}_{H_2}^2$. The field-dependent masses of the NG bosons are given by

$$\bar{m}_{G^0}^2 = \lambda\phi^2 - m_{\Phi}^2 + \lambda_{\Phi 1}s_1^2 + \mu_{\Phi 1}s_1 + \lambda_{\Phi 2}s_2^2, \quad (\text{A28})$$

$$\bar{m}_{G^{\pm}}^2 = \bar{m}_{G^0}^2. \quad (\text{A29})$$

Those of weak gauge bosons and fermions are the same as the SM [93].

We implement the thermal effects of the field-dependent masses by the replacement $\bar{m}_k^2(\phi, s_i) \rightarrow \bar{M}_k^2(\phi, s_i, T)$ in the effective potential V_{eff} [94]. Thermal corrected masses $\bar{M}_k^2(\phi, s_i, T)$ involve the temperature dependent self-energy $\Pi_k(T)$. The one for the Higgs

field and singlet scalar fields are given by

$$\begin{aligned} \Pi_{\Phi}(T) &= T^2 \left[\frac{1}{16} (3g^2 + g'^2) + \frac{y_t^2}{4} \right. \\ &\quad \left. + \frac{1}{6} \left(\frac{3\lambda}{2} + \frac{\lambda_{\Phi 1}}{2} + \frac{\lambda_{\Phi 2}}{2} \right) + \frac{\lambda}{4} \right] \end{aligned} \quad (\text{A30})$$

$$\Pi_{S_1}(T) = T^2 \left[\lambda_1 + \frac{1}{6}(\lambda_{12} + 2\lambda_{\Phi 1}) \right], \quad (\text{A31})$$

$$\Pi_{S_2}(T) = T^2 \left[\frac{1}{6}(\lambda_{12} + 2\lambda_{\Phi 2}) + \lambda_2 \right], \quad (\text{A32})$$

One then obtains field-dependent mass matrix with the thermal corrections in the basis of (ϕ, s_1, s_2) : $\bar{M}^2 = \bar{\mathcal{M}}^2 + \text{diag}(\Pi_{\Phi}, \Pi_{S_1}, \Pi_{S_2})$. Diagonalization of \bar{M}^2 yields the thermal corrected masses of the CP-even Higgs bosons. The thermal corrected masses of the NG boson are obtained by $M_{G^0, G^{\pm}}^2 = \bar{m}_{G^0, G^{\pm}}^2 + \Pi_{\Phi}$. Masses of longitudinal mode of the weak gauge bosons also replaced by thermal corrected ones [93]. While the effective potential $V_{\text{eff}}(\phi, s_i, T)$ of the three dimensional fields is presented here, we analyze $V_{\text{eff}}(0, s_i, T)$ to examine phase transitions in the direction of S_1 and S_2 .

For the parameter points presented in the main text, we have checked that the parameter points satisfy the theoretical constraints for the scalar potential, i.e., perturbative unitarity [78] and the stability condition of the potential [78]. Experimental constraints such as direct searches of the heavy Higgs boson and Higgs coupling measurements for the model with a single real singlet scalar field are discussed with the Run II data of LHC in the Appendix of Ref. [69]. The above parameter points satisfy all these experimental constraints.

For the model parameters adopted in the main text, there are three relevant local minima that appear at different temperatures. At sufficiently high temperatures, there is only a unique vacuum where the Z_2 symmetry (as well as the EW symmetry) is restored. Note that the singlet S_1 has a nonzero VEV in this vacuum, since we do not impose Z_2 symmetry on S_1 . Then, as the temperature decreases, another local minimum appears along the S_1 direction. Let us call this minimum the S_1 -vacuum. We have confirmed that the bubble nucleation rate progressively increases as the universe cools down, where we have used `CosmoTransitions` [79] to evaluate the tunneling rate. Subsequently, the second PT occurs in the bulk along the S_2 -direction. Let us call this vacuum the S_2 -vacuum. For the adopted model parameters, the second PT is of second order, and completes around $T_{\text{PT}} \approx 221$ GeV. The energy difference between the S_1 -vacuum and the S_2 -vacuum, ΔV , depends on the cosmic temperature. For the adopted parameters, the S_2 -vacuum is still meta-stable soon after the second PT, and it becomes the true vacuum around $T_t \approx 60-70$ GeV. To illustrate such phase transitions, Fig. 2 shows the effective potential in the (s_2, s_1) plane at different temperatures $T = 30, 100, 220, \text{ and } 600$ GeV.

⁵ The Coleman-Weinberg potential can also be evaluated by applying another renormalization scheme, such as on shell-like schemes. The renormalization scheme difference of the effective potential is discussed in Refs. [91, 92].




FIG. 3. The highly monochromatic PBH mass spectrum in the singlet-extension model. The spectra from top to bottom are calculated with different values of the triple coupling of S_1 , $\mu_1 = -190.5, -190.49, -190.48, -190.47, -190.46$ GeV, while other parameters are fixed as in the main text. Note that the PBH fraction shown by the green dots is about two orders of magnitude smaller than the peak height due to the extremely narrow width.

We set $\mu_1 = -190.48$ GeV, and other input parameters are taken as presented in the main text. One can see that the S_1 -vacuum first appears in the S_1 -direction, followed by the appearance of the S_2 -vacuum in the S_2 -direction.

Fig. 3 shows the highly monochromatic PBH mass spectrum predicted in the singlet-extension model. It should be noted that due to the extremely narrow peak width, the PBH fraction in dark matter obtained by integrating the distribution (green dots) is about two orders of magnitude smaller than the peak height of the mass distribution. This spectrum is obtained by substituting the temperature-dependent bubble nucleation rate and ΔV calculated using *CosmoTransition* into Eq. (15). We have assumed that the bubbles start to shrink one Hubble time after the S_2 -vacuum becomes the true vacuum and used $T_c = T_t/\sqrt{2}$. Here, the spectra from top to bottom correspond to different values of the triple coupling of S_1 , $\mu_1 = -190.5, -190.49, -190.48, -190.47, -190.46$ GeV, while keeping the other parameters fixed as in the main text.

-
- [1] N. Aghanim *et al.* (Planck), *Astron. Astrophys.* **641**, A6 (2020), [Erratum: *Astron. Astrophys.* 652, C4 (2021)], [arXiv:1807.06209 \[astro-ph.CO\]](#).
- [2] V. C. Rubin and W. K. Ford, Jr., *Astrophys. J.* **159**, 379 (1970).
- [3] F. Zwicky, *Helv. Phys. Acta* **6**, 110 (1933).
- [4] S. W. Hawking, *Commun. Math. Phys.* **25**, 152 (1972).
- [5] B. J. Carr and S. W. Hawking, *Mon. Not. Roy. Astron. Soc.* **168**, 399 (1974).
- [6] B. J. Carr, *Astrophys. J.* **201**, 1 (1975).
- [7] P. H. Frampton, M. Kawasaki, F. Takahashi, and T. T. Yanagida, *JCAP* **04**, 023 (2010), [arXiv:1001.2308 \[hep-ph\]](#).
- [8] B. Carr, F. Kuhnel, and M. Sandstad, *Phys. Rev. D* **94**, 083504 (2016), [arXiv:1607.06077 \[astro-ph.CO\]](#).
- [9] B. Carr, K. Kohri, Y. Sendouda, and J. Yokoyama, *Rept. Prog. Phys.* **84**, 116902 (2021), [arXiv:2002.12778 \[astro-ph.CO\]](#).
- [10] B. Carr and F. Kuhnel, *Ann. Rev. Nucl. Part. Sci.* **70**, 355 (2020), [arXiv:2006.02838 \[astro-ph.CO\]](#).
- [11] A. M. Green and B. J. Kavanagh, *J. Phys. G* **48**, 043001 (2021), [arXiv:2007.10722 \[astro-ph.CO\]](#).
- [12] Y. B. Zel'dovich and I. D. Novikov, *Sov. Astron.* **10**, 602 (1967).
- [13] J. Yokoyama, *Astron. Astrophys.* **318**, 673 (1997), [arXiv:astro-ph/9509027](#).
- [14] J. García-Bellido, A. Linde, and D. Wands, *Phys. Rev. D* **54**, 6040 (1996), [arXiv:astro-ph/9605094 \[astro-ph\]](#).
- [15] M. Kawasaki, N. Sugiyama, and T. Yanagida, *Phys. Rev. D* **57**, 6050 (1998), [arXiv:hep-ph/9710259 \[hep-ph\]](#).
- [16] A. Dolgov and J. Silk, *Phys. Rev. D* **47**, 4244 (1993).
- [17] A. D. Dolgov, M. Kawasaki, and N. Kevlishvili, *Nucl. Phys. B* **807**, 229 (2009), [arXiv:0806.2986 \[hep-ph\]](#).
- [18] M. Kawasaki and K. Murai, *Phys. Rev. D* **100**, 103521 (2019), [arXiv:1907.02273 \[astro-ph.CO\]](#).
- [19] N. Kitajima and F. Takahashi, *JCAP* **11**, 060 (2020), [arXiv:2006.13137 \[hep-ph\]](#).
- [20] M. Kawasaki, K. Murai, and H. Nakatsuka, *JCAP* **10**, 025 (2021), [arXiv:2107.03580 \[astro-ph.CO\]](#).
- [21] K. Kasai, M. Kawasaki, and K. Murai, *JCAP* **10**, 048 (2022), [arXiv:2205.10148 \[astro-ph.CO\]](#).
- [22] K. Kasai, M. Kawasaki, N. Kitajima, K. Murai, S. Neda, and F. Takahashi, *JCAP* **10**, 049 (2023), [arXiv:2305.13023 \[astro-ph.CO\]](#).
- [23] K. Kasai, M. Kawasaki, N. Kitajima, K. Murai, S. Neda, and F. Takahashi, *JCAP* **05**, 092 (2024), [arXiv:2310.13333 \[astro-ph.CO\]](#).
- [24] K. Kasai, M. Kawasaki, K. Murai, and S. Neda, (2024), [arXiv:2405.09790 \[astro-ph.CO\]](#).
- [25] F. Ferrer, E. Masso, G. Panico, O. Pujolas, and F. Rompineve, *Phys. Rev. Lett.* **122**, 101301 (2019), [arXiv:1807.01707 \[hep-ph\]](#).
- [26] S. Ge, *Phys. Dark Univ.* **27**, 100440 (2020), [arXiv:1905.12182 \[hep-ph\]](#).
- [27] J. Liu, Z.-K. Guo, and R.-G. Cai, *Phys. Rev. D* **101**, 023513 (2020), [arXiv:1908.02662 \[astro-ph.CO\]](#).
- [28] G. B. Gelmini, J. Hyman, A. Simpson, and E. Vitagliano, *JCAP* **06**, 055 (2023), [arXiv:2303.14107 \[hep-ph\]](#).
- [29] N. Kitajima, J. Lee, K. Murai, F. Takahashi, and W. Yin, *Phys. Lett. B* **851**, 138586 (2024), [arXiv:2306.17146 \[hep-ph\]](#).
- [30] S. Ge, J. Guo, and J. Liu, *Phys. Rev. D* **109**, 123030 (2024), [arXiv:2309.01739 \[hep-ph\]](#).
- [31] Y. Gouttenoire and E. Vitagliano, *Phys. Rev. D* **110**, L061306 (2024), [arXiv:2306.17841 \[gr-qc\]](#).
- [32] Y. Gouttenoire and E. Vitagliano, *Phys. Rev. D* **109**, 123507 (2024), [arXiv:2311.07670 \[hep-ph\]](#).
- [33] R. Z. Ferreira, A. Notari, O. Pujolàs, and F. Rompineve, *JCAP* **06**, 020 (2024), [arXiv:2401.14331 \[astro-ph.CO\]](#).
- [34] D. I. Dunskey and M. Kongsore, *JHEP* **06**, 198 (2024), [arXiv:2402.03426 \[hep-ph\]](#).
- [35] Y. Gouttenoire, S. F. King, R. Roshan, X. Wang, G. White, and M. Yamazaki, (2025), [arXiv:2501.16414](#)

- [hep-ph].
- [36] K. Sato, M. Sasaki, H. Kodama, and K.-i. Maeda, *Prog. Theor. Phys.* **65**, 1443 (1981).
- [37] H. Kodama, M. Sasaki, K. Sato, and K.-i. Maeda, *Prog. Theor. Phys.* **66**, 2052 (1981).
- [38] H. Kodama, M. Sasaki, and K. Sato, *Prog. Theor. Phys.* **68**, 1979 (1982).
- [39] K.-i. Maeda, K. Sato, M. Sasaki, and H. Kodama, *Phys. Lett. B* **108**, 98 (1982).
- [40] K. Jedamzik and J. C. Niemeyer, *Phys. Rev. D* **59**, 124014 (1999), [arXiv:astro-ph/9901293](#).
- [41] J. Liu, L. Bian, R.-G. Cai, Z.-K. Guo, and S.-J. Wang, *Phys. Rev. D* **105**, L021303 (2022), [arXiv:2106.05637 \[astro-ph.CO\]](#).
- [42] K. Hashino, S. Kanemura, and T. Takahashi, *Phys. Lett. B* **833**, 137261 (2022), [arXiv:2111.13099 \[hep-ph\]](#).
- [43] K. Hashino, S. Kanemura, T. Takahashi, and M. Tanaka, *Phys. Lett. B* **838**, 137688 (2023), [arXiv:2211.16225 \[hep-ph\]](#).
- [44] K. Kawana, T. Kim, and P. Lu, *Phys. Rev. D* **108**, 103531 (2023), [arXiv:2212.14037 \[astro-ph.CO\]](#).
- [45] M. Lewicki, P. Toczek, and V. Vaskonen, *JHEP* **09**, 092 (2023), [arXiv:2305.04924 \[astro-ph.CO\]](#).
- [46] Y. Gouttenoire and T. Volansky, *Phys. Rev. D* **110**, 043514 (2024), [arXiv:2305.04942 \[hep-ph\]](#).
- [47] S. Balaji, M. Fairbairn, and M. O. Olea-Romacho, *Phys. Rev. D* **109**, 075048 (2024), [arXiv:2402.05179 \[hep-ph\]](#).
- [48] M. Lewicki, P. Toczek, and V. Vaskonen, *Phys. Rev. Lett.* **133**, 221003 (2024), [arXiv:2402.04158 \[astro-ph.CO\]](#).
- [49] M. M. Flores, A. Kusenko, and M. Sasaki, *Phys. Rev. D* **110**, 015005 (2024), [arXiv:2402.13341 \[hep-ph\]](#).
- [50] S. Kanemura, M. Tanaka, and K.-P. Xie, *JHEP* **06**, 036 (2024), [arXiv:2404.00646 \[hep-ph\]](#).
- [51] M. Lewicki, P. Toczek, and V. Vaskonen, (2024), [arXiv:2412.10366 \[astro-ph.CO\]](#).
- [52] K. Hashino, S. Kanemura, T. Takahashi, M. Tanaka, and C.-M. Yoo, (2025), [arXiv:2501.11040 \[hep-ph\]](#).
- [53] C.-M. Yoo, T. Harada, and H. Okawa, *Phys. Rev. D* **102**, 043526 (2020), [Erratum: *Phys.Rev.D* 107, 049901 (2023)], [arXiv:2004.01042 \[gr-qc\]](#).
- [54] A. Escrivà and C.-M. Yoo, (2024), [arXiv:2410.03451 \[gr-qc\]](#).
- [55] A. Escrivà and C.-M. Yoo, (2024), [arXiv:2410.03452 \[gr-qc\]](#).
- [56] S. R. Coleman, V. Glaser, and A. Martin, *Commun. Math. Phys.* **58**, 211 (1978).
- [57] P. Mroz *et al.*, *Nature* **548**, 183 (2017), [arXiv:1707.07634 \[astro-ph.EP\]](#).
- [58] H. Niikura *et al.*, *Nature Astron.* **3**, 524 (2019), [arXiv:1701.02151 \[astro-ph.CO\]](#).
- [59] H. Niikura, M. Takada, S. Yokoyama, T. Sumi, and S. Masaki, *Phys. Rev. D* **99**, 083503 (2019), [arXiv:1901.07120 \[astro-ph.CO\]](#).
- [60] K. G. Arun *et al.* (LISA), *Living Rev. Rel.* **25**, 4 (2022), [arXiv:2205.01597 \[gr-qc\]](#).
- [61] K. Hashino, M. Kakizaki, S. Kanemura, P. Ko, and T. Matsui, *Phys. Lett. B* **766**, 49 (2017), [arXiv:1609.00297 \[hep-ph\]](#).
- [62] V. Vaskonen, *Phys. Rev. D* **95**, 123515 (2017), [arXiv:1611.02073 \[hep-ph\]](#).
- [63] A. Beniwal, M. Lewicki, J. D. Wells, M. White, and A. G. Williams, *JHEP* **08**, 108 (2017), [arXiv:1702.06124 \[hep-ph\]](#).
- [64] K. Hashino, R. Jinno, M. Kakizaki, S. Kanemura, T. Takahashi, and M. Takimoto, *Phys. Rev. D* **99**, 075011 (2019), [arXiv:1809.04994 \[hep-ph\]](#).
- [65] J. Ellis, M. Lewicki, and J. M. No, *JCAP* **04**, 003 (2019), [arXiv:1809.08242 \[hep-ph\]](#).
- [66] A. Alves, T. Ghosh, H.-K. Guo, K. Sinha, and D. Vagie, *JHEP* **04**, 052 (2019), [arXiv:1812.09333 \[hep-ph\]](#).
- [67] T. Alanne, T. Hügler, M. Platscher, and K. Schmitz, *JHEP* **03**, 004 (2020), [arXiv:1909.11356 \[hep-ph\]](#).
- [68] J. Ellis, M. Lewicki, M. Merchand, J. M. No, and M. Zych, *JHEP* **01**, 093 (2023), [arXiv:2210.16305 \[hep-ph\]](#).
- [69] M. J. Ramsey-Musolf, T. V. I. Tenkanen, and V. Q. Tran, (2024), [arXiv:2409.17554 \[hep-ph\]](#).
- [70] W.-Y. Ai, L. Heurtier, and T. H. Jung, (2024), [arXiv:2409.02175 \[astro-ph.CO\]](#).
- [71] D. Bodeker and G. D. Moore, *JCAP* **05**, 009 (2009), [arXiv:0903.4099 \[hep-ph\]](#).
- [72] D. Bodeker and G. D. Moore, *JCAP* **05**, 025 (2017), [arXiv:1703.08215 \[hep-ph\]](#).
- [73] S. Sugiyama, M. Takada, and A. Kusenko, *Phys. Lett. B* **840**, 137891 (2023), [arXiv:2108.03063 \[hep-ph\]](#).
- [74] P. Tisserand *et al.* (EROS-2), *Astron. Astrophys.* **469**, 387 (2007), [arXiv:astro-ph/0607207](#).
- [75] P. Mróz *et al.*, *Nature* **632**, 749 (2024), [arXiv:2403.02386 \[astro-ph.GA\]](#).
- [76] P. Mróz *et al.*, *Astrophys. J. Lett.* **976**, L19 (2024), [arXiv:2410.06251 \[astro-ph.CO\]](#).
- [77] B. J. Kavanagh, “PBH bounds,” (2019).
- [78] T. Robens, T. Stefaniak, and J. Wittbrodt, *Eur. Phys. J. C* **80**, 151 (2020), [arXiv:1908.08554 \[hep-ph\]](#).
- [79] C. L. Wainwright, *Comput. Phys. Commun.* **183**, 2006 (2012), [arXiv:1109.4189 \[hep-ph\]](#).
- [80] K. Yamamoto, *Phys. Lett. B* **168**, 341 (1986).
- [81] D. H. Lyth and E. D. Stewart, *Phys. Rev. D* **53**, 1784 (1996), [arXiv:hep-ph/9510204](#).
- [82] N. Kitajima, S. Nakagawa, and F. Takahashi, *Phys. Rev. D* **105**, 103011 (2022), [arXiv:2111.06696 \[hep-ph\]](#).
- [83] J. Garriga, A. Vilenkin, and J. Zhang, *JCAP* **02**, 064 (2016), [arXiv:1512.01819 \[hep-th\]](#).
- [84] H. Deng and A. Vilenkin, *JCAP* **12**, 044 (2017), [arXiv:1710.02865 \[gr-qc\]](#).
- [85] A. Caravano, K. Inomata, and S. Renaux-Petel, *Phys. Rev. Lett.* **133**, 151001 (2024), [arXiv:2403.12811 \[astro-ph.CO\]](#).
- [86] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, *Phys. Lett. B* **155**, 36 (1985).
- [87] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, *Ann. Rev. Nucl. Part. Sci.* **43**, 27 (1993), [arXiv:hep-ph/9302210](#).
- [88] S. Kanemura, M. Kikuchi, and K. Yagyu, *Nucl. Phys. B* **917**, 154 (2017), [arXiv:1608.01582 \[hep-ph\]](#).
- [89] S. R. Coleman and E. J. Weinberg, *Phys. Rev. D* **7**, 1888 (1973).
- [90] L. Dolan and R. Jackiw, *Phys. Rev. D* **9**, 3320 (1974).
- [91] C.-W. Chiang, Y.-T. Li, and E. Senaha, *Phys. Lett. B* **789**, 154 (2019), [arXiv:1808.01098 \[hep-ph\]](#).
- [92] P. Athron, C. Balazs, A. Fowlie, L. Morris, G. White, and Y. Zhang, *JHEP* **01**, 050 (2023), [arXiv:2208.01319 \[hep-ph\]](#).
- [93] M. E. Carrington, *Phys. Rev. D* **45**, 2933 (1992).
- [94] R. R. Parwani, *Phys. Rev. D* **45**, 4695 (1992), [Erratum: *Phys.Rev.D* 48, 5965 (1993)], [arXiv:hep-ph/9204216](#).