On the limits of some Bayesian model evaluation statistics

Hien Duy Nguyen^{1,2}, Mayetri Gupta³, Jacob Westerhout⁴, and TrungTin Nguyen⁵

¹Department of Mathematics and Physical Science, La Trobe University, Melbourne, Australia.

²Institute of Mathematics for Industry, Kyushu University, Fukuoka, Japan.

³School of Mathematics and Statistics, University of Glasgow, Glasgow, UK.

⁴School of Mathematics and Physics, University of Queensland, Brisbane,

Australia.

⁵School of Mathematical Sciences, Queensland University of Technology, Brisbane, Australia.

February 7, 2025

Abstract

Model selection and order selection problems frequently arise in statistical practice. A popular approach to addressing these problems in the frequentist setting involves information criteria based on penalized maxima of log-likelihoods for competing models. In the Bayesian context, similar criteria are employed, replacing the maxima of log-likelihoods with their posterior expectations. Despite their popularity in applications, the large-sample behavior of these criteria—such as the deviance information criterion (DIC), Bayesian predictive information criterion (BPIC), and widely-applicable Bayesian information criterion (WBIC)—has received relatively little attention. In this work, we investigate the almost sure limits of these criteria and establish novel results on posterior and generalized posterior consistency, which are of independent interest. The utility of our theoretical findings is demonstrated via illustrative technical and numerical examples.

Keywords. Bayesian model comparison, Bayesian information criterion, deviance information criterion, Bayesian consistency, almost sure weak convergence

1 Introduction

1.1 General introduction

Bayesian statistics provides a cohesive and comprehensive set of tools for the analysis of data arising from natural and man-made phenomena. In recent decades, with advances in the availability of computational resources, Bayesian analysis has become an increasingly popular paradigm for the development of novel statistical methods in applied settings, and Bayesian statistical inference is an increasingly popular area of fundamental research and development (Ghosh et al., 2006; Robert, 2007).

As in frequentist settings, the Bayesian framework often requires the analyst to choose between various models that may each provide plausible fits to the observed data. To this end, a popular technique is to employ the method of information criteria, such as using the Akaike information criterion (AIC; Akaike, 1974) or the Bayesian information criterion (BIC; Schwarz, 1978), in order to evaluate the goodness of fit of competing models. Both the AIC and BIC are defined via the maximum of the log-likelihood function of the data. Some texts that explore these information criteria in detail are Burnham & Anderson (2002) and Konishi & Kitagawa (2008). The limiting behaviors of the AIC, BIC, and other information criteria based on the maximum of some data dependent objective has been investigated in general contexts, for example, in the works of Sin & White (1996) and Baudry (2015).

As an alternative to the maximum likelihood approach, some authors have suggested information criteria based on the posterior average of the log-likelihood function, as well as the log-likelihood function evaluated at Bayesian quantities, such as the log-likelihood evaluated at the posterior mean. Comprehensive resources for these types of criteria are Ando (2010) and Watanabe (2018) who discuss examples of such criteria, including the deviance information criterion (DIC; Spiegelhalter et al., 2002), the Bayesian predictive information criterion (BPIC; Ando, 2007), and the widely applicable Bayesian information criterion (WBIC; Watanabe, 2013).

In this work, we seek to obtain general regularity conditions under which the almost sure limits of the DIC, BPIC, and WBIC can be obtained. Our technique relies on the concept of almost sure weak convergence, otherwise known as almost sure conditional weak convergence, as espoused in the works of Sweeting (1989), Berti et al. (2006), and Grübel & Kabluchko (2016). This is the mode of convergence under which Bayesian consistency is often established, as well as limit theorems for the Bootstrap; see, for example, Ghosh & Ramamoorthi (2003, Ch. 4), Ghosal & Van der Vaart (2017, Ch. 6), and van der Vaart & Wellner (2023, Sec. 1.13 and 3.7). The almost sure convergence of posterior measures is then combined with the convergence of averaged functions of data and parameters using generalizations of the Lebesgue convergence theorems, as described in the works of Serfozo (1982), Feinberg et al. (2020a) and Feinberg et al. (2020b).

We note that convergence in mean value for the listed information criteria and related objects can be obtained, for example, via the many results of Watanabe (2018). Our almost sure convergence results can thus be seen as complementary to the approach of Watanabe (2018). We shall provide technical details in the sequel.

1.2 Technical introduction

Let $(\Omega, \mathfrak{F}, \mathbb{P})$ be a probability space with typical element ω and expectation operator \mathbb{E} , and let $(\mathbb{T}, \mathfrak{B}_{\mathbb{T}})$ be a measurable space with typical element θ , where \mathbb{T} is either an open or closed subset of \mathbb{R}^p with its Borel σ -algebra $\mathfrak{B}_{\mathbb{T}}$. For each $n \in \mathbb{N}$, let $f_n : \Omega \times \mathbb{T} \to \mathbb{R}$ be measurable with respect to $\omega \in \Omega$, for each $\theta \in \mathbb{T}$, and let $\mathbb{Q}_n : \Omega \times \mathfrak{B}_{\mathbb{T}} \to \mathbb{R}_{\geq 0}$, be a random measure in the sense that for each ω , $\mathbb{Q}_n(\omega, \cdot) \in \mathcal{P}(\mathbb{T})$, where $\mathcal{P}(\mathbb{T})$ is the space of measures on $(\mathbb{T}, \mathfrak{B}_{\mathbb{T}})$.

Following the exposition of Grübel & Kabluchko (2016), we say that the sequence $(Q_n)_{n \in \mathbb{N}}$ converges weakly to $Q \in \mathcal{P}(\mathbb{T})$, almost surely, with respect to the measure P (which we denote as $Q_n \xrightarrow[n \to \infty]{P-a.s.w.} Q$), if for each bounded and continuous $f : \mathbb{T} \to \mathbb{R}$,

$$\int_{\mathbb{T}} f(\theta) \operatorname{Q}_{n}(\omega, \mathrm{d}\theta) \xrightarrow[n \to \infty]{} \int_{\mathbb{T}} f(\theta) \operatorname{Q}(\mathrm{d}\theta), \qquad (1)$$

for every ω on a P-almost sure (P-a.s.) set (where the set may depend on f). Now suppose that the sequence of random functions $(f_n(\omega, \cdot))_{n \in \mathbb{N}}$ converges to some deterministic $(\mathbb{T}, \mathfrak{B}_{\mathbb{T}})$ -measurable function f, in some appropriate sense, on a P-a.s. set. Then, we seek to investigate integrals of the form

$$\int_{\mathbb{T}} f_n(\omega, \theta) \, \mathcal{Q}_n(\omega, \mathrm{d}\theta) \,, \tag{2}$$

and deduce sufficient conditions under which such integrals will converge to the limit

$$\int_{\mathbb{T}} f(\theta) \operatorname{Q}(\mathrm{d}\theta),$$

for each ω on a P-a.s. set.

We note that such integrals of the form (2) occur frequently in Bayesian statistics. To present some common cases, let us introduce some further notation. Let $(X_i)_{i\in[n]}$ be a sequence of random variables $X_i : \Omega \to \mathbb{X}$, for each $i \in [n] \in \{1, \ldots, n\}$, where $\mathbb{X} \subset \mathbb{R}^q$ is a real subspace with σ -algebra $\mathfrak{B}_{\mathbb{X}}$. Next, we suppose that $\Pi \in \mathcal{P}(\mathbb{T})$ is a prior probability measure on $(\mathbb{T}, \mathfrak{B}_{\mathbb{T}})$, with density function $\pi : \mathbb{T} \to \mathbb{R}_{\geq 0}$ with respect to some dominating measure \mathfrak{m} on $(\mathbb{T}, \mathfrak{B}_{\mathbb{T}})$.

Next, we suppose that $p(x_1, \ldots, x_n | \theta)$ is a likelihood function in the sense that for each fixed $\theta \in \mathbb{T}$, $p(\cdot | \theta) : \mathbb{X}^n \to \mathbb{R}_{\geq 0}$ is a probability density function with respect to the *n*-fold product of some dominating measure \mathfrak{n} on $(\mathbb{X}, \mathfrak{B}_{\mathbb{X}})$. Then, we may identify the sequence of posterior measures $(\Pi_n(\omega, \cdot))_{n \in \mathbb{N}}$ as a sequence of random measures, where

$$\Pi_n(\omega, \cdot) = \Pi\left(\cdot | X_1(\omega), \dots, X_n(\omega)\right),\tag{3}$$

is defined by its density function

$$\theta \mapsto \pi\left(\theta | X_1, \dots, X_n\right) = \frac{p\left(X_1, \dots, X_n | \theta\right) \pi\left(\theta\right)}{\int_{\mathbb{T}} p\left(X_1, \dots, X_n | \tau\right) \pi\left(\tau\right) \mathfrak{m}\left(\mathrm{d}\tau\right)}$$

Suppose further that $(\theta_n)_{n \in \mathbb{N}}$ is a sequence of estimators of θ , in the sense that $\theta_n : \mathbb{X}^n \to \mathbb{T}$, for each $n \in \mathbb{N}$. A first example of an object of the form (2) is to consider parametric loss functions of the form

 $\ell : \mathbb{T} \times \mathbb{T} \to \mathbb{R}_{\geq 0}$. Then, we may identify $\omega \mapsto \ell(\theta_n(\omega), \theta)$ as a random function and write the posterior risk with respect to ℓ , as defined in Lehmann & Casella (1998, Ch. 4), by the expression

$$\int_{\mathbb{T}} \ell\left(\theta_n\left(\omega\right), \theta\right) \Pi_n\left(\omega, \mathrm{d}\theta\right)$$

which is in the form (2). More pertinent to our investigation, we can consider more elaborate random functions f_n that are best described as utility or score functions $u_n : \mathbb{X}^n \times \mathbb{T} \to \mathbb{R}$, as per the exposition of Berger (1985) and Bernardo & Smith (2009). Examples of such utility functions include the log-likelihood function

$$u_n(x_1,\ldots,x_n;\theta) = \log p(x_1,\ldots,x_n|\theta).$$

Here, the integral of the log-likelihood with respect to the posterior measure is of particular interest as it appears in both the DIC and BPIC, while the integral with respect to a modified posterior measure, describe below, forms the definition of the WBIC.

We can also identify other Bayesian objects with sequences of measures $(Q_n(\omega, \cdot))_{n \in \mathbb{N}}$. For example, approximate Bayesian computation pseudo-posteriors can be identified as random measures (see, e.g., Bernton et al., 2019, Nguyen et al., 2020 and Forbes et al., 2022), as with asymptotic Bayesian approximations of the posterior distribution obtained by a Bernstein–von Mises Theorem (see, e.g., van der Vaart, 1998, Ch. 10 and Ghosh et al., 2006, Ch. 4). Related to such objects are data dependent measures on $(\mathbb{T}, \mathfrak{B}_{\mathbb{T}})$ often studied as fiducial distributions or confidence distributions (see e.g., Hannig et al., 2016 and Schweder & Hjort, 2016). A general method for generating sequences $(Q_n)_{n\in\mathbb{N}}$ that exhibit P-a.s. convergence, including variational Bayesian posteriors (see e.g., Zhang & Gao, 2020 and Nguyen et al., 2024), the generalized posteriors (see, e.g., Zhang, 2006 and Bissiri et al., 2016), and typical posteriors of form (3) have recently been studied by Knoblauch et al. (2022) (see also Knoblauch, 2019). Of particular interest in our study is the generalized posterior $\Pi_n^{\beta_n}$, for sequence $(\beta_n)_{n\in\mathbb{N}} \subset \mathbb{R}_{>0}$, whose density is defined by

$$\pi^{\beta_n} \left(\theta | X_1, \dots, X_n \right) = \frac{\left\{ p \left(X_1, \dots, X_n | \theta \right) \right\}^{\beta_n} \pi \left(\theta \right)}{\int_{\mathbb{T}} \left\{ p \left(X_1, \dots, X_n | \theta \right) \right\}^{\beta_n} \pi \left(\tau \right) \mathfrak{m} \left(\mathrm{d} \tau \right)}$$

which appears in the definition of the WBIC (in particular, $\beta_n = 1/\log n$).

The modes of weak convergence, almost surely and in probability, have previously been formally studied in the works of Sweeting (1989), Berti et al. (2006), Bain & Crisan (2009), and Grübel & Kabluchko (2016). Let $\delta_{\theta_0} : 2^{\mathbb{T}} \to [0,1]$ denote the delta measure that takes value 1 if $\theta_0 \in \mathbb{B}$ and 0, otherwise, for every $\mathbb{B} \in 2^{\mathbb{T}}$. Then, the usual notion of posterior consistency, with respect to a parameter $\theta_0 \in \mathbb{T}$, in Bayesian analysis, can be characterized as the weak convergence of posterior measure $\Pi_n^1 = \Pi_n$ to δ_{θ_0} , P-a.s. (cf. Ghosal & Van der Vaart, 2017, Prop. 6.2). For another example, if the sequence $(\theta_n)_{n\in\mathbb{N}}$ converges P-a.s. to θ_0 , then since \mathbb{T} is separable, we can determine weak convergence by checking condition (1) using only a countable number of bounded continuous functions (cf. Bain & Crisan, 2009, Thm. 2.18), and determine that $(\delta_{\theta_n})_{n\in\mathbb{N}}$ converges to δ_{θ_0} , P-a.s.w., by continuous mapping. Since we can write

$$f_n(\omega,\theta_n) = \int_{\mathbb{T}} f_n(\omega,\theta) \,\delta_{\theta_n}(\mathrm{d}\theta) \,, \, \mathrm{and} \, f(\theta_0) = \int_{\mathbb{T}} f(\theta) \,\delta_{\theta_0}(\mathrm{d}\theta) \,, \tag{4}$$

the convergence of $(f_n(\cdot, \theta_n))_{n \in \mathbb{N}}$ to $f(\theta_0)$, P-a.s., falls within the context of this work.

The remainder of the manuscript is organized as follows. In Section 2, we present our main results. Section 3 provides some example applications of our results, accompanied by numerical simulations, while Section 4 is dedicated to proofs and technical details. A concluding discussion follows in Section 5. The Appendix then contains some technical calculations from the examples in Section 4.

2 Main results

In the sequel, we will retain the notation and technical setup of the introduction. Furthermore, to simplify discussions, we will assume that $(X_i)_{i\in\mathbb{N}}$ is an independent and identically distributed (IID) sequence of random variables, each identical to $X : \Omega \to \mathbb{X}$.

We are largely concerned with three information criteria, which have found particular popularity in the Bayesian literature; namely, the DIC of Spiegelhalter et al. (2002), the BPIC of Ando (2010), and the WBIC of Watanabe (2013), taking the respective forms:

$$\mathrm{DIC}_{n} = -\frac{4}{n} \int_{\mathbb{T}} \sum_{i=1}^{n} \log p\left(X_{i}|\theta\right) \Pi_{n}\left(\mathrm{d}\theta\right) + \frac{2}{n} \sum_{i=1}^{n} \log p\left(X_{i}|\bar{\theta}_{n}\right),\tag{5}$$

$$BPIC_n = -\frac{2}{n} \int_{\mathbb{T}} \sum_{i=1}^n \log p\left(X_i|\theta\right) \Pi_n\left(\mathrm{d}\theta\right) + 2\frac{p}{n},\tag{6}$$

and

WBIC_n =
$$-\frac{2}{n} \int_{\mathbb{T}} \sum_{i=1}^{n} \log p(X_i|\theta) \Pi_n^{1/\log n}(\mathrm{d}\theta)$$
,

where

$$\bar{\theta}_{n} = \left(\int_{\mathbb{T}} \theta_{1} \Pi_{n} \left(\mathrm{d}\theta \right), \dots, \int_{\mathbb{T}} \theta_{j} \Pi_{n} \left(\mathrm{d}\theta \right), \dots, \int_{\mathbb{T}} \theta_{p} \Pi_{n} \left(\mathrm{d}\theta \right) \right)$$

is the posterior mean, with $\theta_j : \mathbb{R}^p \to \mathbb{R}$ denoting the *j*th coordinate projection of θ . Here, $x \mapsto p(x|\theta)$ characterizes a family of probability density functions (PDFs), indexed by $\theta \in \mathbb{T}$, with

$$p(x_1,...,x_n|\theta) = \prod_{i=1}^n p(x_i|\theta)$$

in the definitions of Π_n and $\Pi_n^{\beta_n}$.

Observe that the first expression in each of the criteria is in form (2) with

$$f_n(\omega, \theta) = -\frac{2}{n} \sum_{i=1}^n \log p(X_i | \theta), \qquad (7)$$

in each case, and where Q_n equals $\prod_n^{\beta_n}$ for the WBIC, for some choice of $(\beta_n)_{n \in \mathbb{N}}$. In particular, the BPIC and DIC take $\beta_n = 1$, whereas $\beta_n = 1/\log n$ is used in the original definition of the WBIC. The second term in the DIC also has form (2), where f_n has form (7) but Q_n equals $\delta_{\bar{\theta}_n}$.

Note that we have presented the information criteria as functions of sample averages for convenience. We have also scaled the WBIC by a factor of two to standardize the scales of the three quantities. That is, in our notation, the original characterizations of the DIC, BPIC, and WBIC present the objects in the forms: $n\text{DIC}_n$, $n\text{BPIC}_n$, and $n\text{WBIC}_n/2$, respectively.

2.1 Consistency of $\Pi_n^{\beta_n}$

Let $\mathbb{B}_{\epsilon}(\theta) = \{\tau \in \mathbb{T} : \|\theta - \tau\| < \epsilon\}$ and $\overline{\mathbb{B}}_{\epsilon}(\theta) = \{\tau \in \mathbb{T} : \|\theta - \tau\| \le \epsilon\}$ denote the open and closed balls around $\theta \in \mathbb{T}$ of radius $\epsilon > 0$, respectively. We shall also use $(\cdot)^{c}$ to denote the set complement with respect to universe \mathbb{T} . Make the following additional assumptions:

- A1 The PDF p is Caratheodory in the sense that $p(\cdot|\theta)$ is measurable for each $\theta \in \mathbb{T}$ and $p(x|\cdot)$ is continuous for each $x \in \mathbb{X}$ and $p(x|\theta) > 0$ for each $\theta \in \mathbb{T}$ and $x \in \mathbb{X}$.
- **A2** There exists a P-a.s. set on which for every compact $\mathbb{K} \subset \mathbb{T}$,

$$\sup_{\theta \in \mathbb{K}} \left| \frac{1}{n} \sum_{i=1}^{n} \log p\left(X_i | \theta \right) - \mathbb{E} \left[\log p\left(X | \theta \right) \right] \right| \xrightarrow[n \to \infty]{} 0.$$
(8)

A3 There exists a unique maximizer $\theta_0 \in \mathbb{T}$ of $\mathbb{E}[\log p(X|\theta)]$, in the sense that, for each $\theta \in \mathbb{T} \setminus \{\theta_0\}$,

$$E \left[\log p(X|\theta) \right] < E \left[\log p(X|\theta_0) \right].$$

A4 There exists a compact set $\mathbb{S} \subsetneq \mathbb{T}$ containing θ_0 , such that there is a P-a.s. set on which

$$\limsup_{n \to \infty} \sup_{\theta \in \mathbb{S}^{c}} \frac{1}{n} \sum_{i=1}^{n} \log p(X_{i}|\theta) < E\left[\log p(X|\theta_{0})\right].$$

Proposition 1. Assume that $(X_i)_{i\in\mathbb{N}}$ are IID, $\Pi(\mathbb{B}_{\rho}(\theta_0)) > 0$, for every $\rho > 0$, and A1-A4 hold. If $n\beta_n \to \infty$, then the sequence of posterior measures $(\Pi_n^{\beta_n})_{n\in\mathbb{N}}$ is consistent with respect to $\theta_0 \in \mathbb{T}$ in the sense that for every $\epsilon > 0$,

$$\Pi_{n}^{\beta_{n}}\left(\bar{\mathbb{B}}_{\epsilon}\left(\theta_{0}\right)\right) \xrightarrow[n \to \infty]{P-a.s.} 1.$$

$$\tag{9}$$

We shall establish a more general result in the next section, which implies Proposition 1. The proof of this proposition is provided in Section 4.1. Assumption A2 provides an almost sure event that guarantees the uniform convergence of the average log-likelihood on all compact sets, which we then use to guarantee the continuous convergence of the average log-likelihood over the entirety of \mathbb{T} , since it is an open or closed Euclidean subset and is therefore hemi-compact (see Beckenstein et al., 2011, Def. 2.1). A3 is an identifiability assumption regarding the maximizer θ_0 of E [log p $(X|\theta)$], requiring that E [log p $(X|\theta)$] be uniquely maximized on \mathbb{T} . Taken together with A1 and A3, A4 can also be verified if the uniform convergence on compact sets $\mathbb{K} \subset \mathbb{T}$ in A2 is instead replaced by uniform convergence over the entirety of \mathbb{T} , P-a.s. By this observation, under A1-A3, A4 is immediately satisfied if \mathbb{T} is compact. In the sequel, we further detail a useful approach, based on quasiconvexity of the log-likelihood functions. Based on Ghosal & Van der Vaart (2017, Prop. 6.2), we have the fact that (9) is equivalent to the statement that $(\Pi_n^{\beta_n})_{n \in \mathbb{N}}$ converges to δ_{θ_0} , P-a.s.w. We consider that both A1 and A3 are fundamental assumptions about the property of the PDFs p and must be inspected for each problem. However, A2 can be implied by more verifiable assumptions using a uniform strong law of large numbers (e.g., Theorem 3). Make the following additional assumptions:

A2a For each $\theta \in \mathbb{T}$,

$$\frac{1}{n} \sum_{i=1}^{n} \log p(X_i | \theta) \xrightarrow[n \to \infty]{\text{P-a.s.}} \mathbb{E} \left[\log p(X | \theta) \right],$$

where $E[|\log p(X|\theta)|] < \infty$.

A2b For each compact set $\mathbb{K} \subset \mathbb{T}$, there exists a dominating function $\Delta : \mathbb{X} \to \mathbb{R}$, such that

$$\left|\log p\left(X|\theta\right)\right| \le \Delta\left(X\right),$$

for each $\theta \in \mathbb{K}$, and $\mathbb{E}\left[\Delta(X)\right] < \infty$.

The conclusion of Proposition 1 holds with A2 replaced by A2a and A2b, above. Thus, we now have enough practical conditions to establish that $(\Pi_n^{\beta_n})_{n \in \mathbb{N}}$ converges to δ_{θ_0} , P-a.s.w.

2.2 A general convergence theorem

We now present conditions under which (2) converges to its limit, generally. To this end, we adapt the deterministic generalized Lebesgue convergence theorems of Serfozo (1982), Feinberg et al. (2020b) and Feinberg et al. (2020a).

To proceed, we require a number of definitions. First, we will say that a sequence of random functions $(f_n(\omega, \cdot))_{n \in \mathbb{N}}$ continuously converges to the deterministic function $f : \mathbb{T} \to \mathbb{R}$, P-a.s., if for every sequence $(\theta_n)_{n \in \mathbb{N}}$ converging to a limit $\theta \in \mathbb{T}$, it holds that $(f_n(\omega, \theta_n))_{n \in \mathbb{N}}$ converges to $f(\theta)$, for every ω on a P-a.s. set. Next, we say that $(f_n)_{n \in \mathbb{N}}$ is asymptotically uniformly integrable (AUI) with respect to $(Q_n)_{n \in \mathbb{N}}$, P-a.s., if

$$\lim_{\delta \to \infty} \limsup_{n \to \infty} \int_{\mathbb{T}} |f_n(\omega, \theta)| \, \chi_{\{\tau \in \mathbb{T} : |f_n(\omega, \tau)| \ge \delta\}}(\theta) \, \mathcal{Q}_n(\omega, \mathrm{d}\theta) = 0,$$

for every ω on a P-a.s. set. If each $f_n = f$, then we will instead say that f is AUI with respect to $(Q_n)_{n \in \mathbb{N}}$. Here, $\chi_{\mathbb{A}} : \mathbb{T} \to \mathbb{R}$ is the usual characteristic function for the set $\mathbb{A} \subset \mathbb{T}$. A direct modification of Feinberg et al. (2020a, Cor. 2.8) for the stochastic setting yields the following general Lebesgue dominated convergence theorem.

Theorem 1. Assume that $(Q_n)_{n \in \mathbb{N}}$ converge to Q, P-a.s.w., and $(f_n)_{n \in \mathbb{N}}$ continuously converge to f, P-a.s. If $(f_n)_{n \in \mathbb{N}}$ is AUI with respect to $(Q_n)_{n \in \mathbb{N}}$, P-a.s., then

$$\lim_{n \to \infty} \int_{\mathbb{T}} f_n(\omega, \theta) \, \mathcal{Q}_n(\omega, \mathrm{d}\theta) = \int_{\mathbb{T}} f(\theta) \, \mathcal{Q}(\mathrm{d}\theta)$$

for almost every ω , with respect to P.

To verify that $(f_n)_{n \in \mathbb{N}}$ continuously converges to f, P-a.s., we can use the equivalent notion that fis continuous on \mathbb{T} , and for almost every ω with respect to P, for every compact $\mathbb{K} \subset \mathbb{T}$, $(f_n(\omega, \cdot))_{n \in \mathbb{N}}$ converges uniformly to f on \mathbb{K} (cf. Remmert 1991, Thm. 3.1.5). At first this might seem daunting since we would have to construct an almost sure event of Ω on which uniform convergence holds for an uncountable number of compact subsets $\mathbb{K} \subset \mathbb{T}$. However, we shall use the fact that since $\mathbb{T} \subset \mathbb{R}^p$ is open or closed, it is hemi-compact, which then provides the necessary regularity to construct such an almost sure event. We shall demonstrate applications of these facts in the sequel.

The AUI property is more difficult to verify but can be implied by more practical assumptions. For instance, Feinberg et al. (2020a, Cor. 2.10) suggests that if there exists a sequence of stochastic functions $(g_n)_{n\in\mathbb{N}}$, converging continuously to $g: \mathbb{T} \to \mathbb{R}$, P-a.s., with $g_n: \Omega \times \mathbb{T} \to \mathbb{R}$, for each $n \in \mathbb{N}$, such that for almost every ω with respect to P, $|f_n(\omega, \theta)| \leq g_n(\omega, \theta)$, for every $\theta \in \mathbb{T}$, and

$$\limsup_{n \to \infty} \int_{\mathbb{T}} g_n(\omega, \theta) \, \mathcal{Q}_n(\mathrm{d}\theta) \le \int_{\mathbb{T}} g(\theta) \, \mathcal{Q}(\mathrm{d}\theta) < \infty, \tag{10}$$

then $(f_n)_{n\in\mathbb{N}}$ is AUI with respect to $(Q_n)_{n\in\mathbb{N}}$, P-a.s., if $(f_n)_{n\in\mathbb{N}}$ continuously converges to f, P-a.s. We note that a version of this criterion, when $f_n = f$ for each n, is given in van der Vaart & Wellner (2023, Thm. 1.11.3). Trivially, the condition holds if there exists a constant $b < \infty$, such that $|f_n(\omega, \theta)| \le b$, for each n, ω , and θ , i.e. $(f_n)_{n\in\mathbb{N}}$ is uniformly bounded by b.

Motivated by van der Vaart & Wellner (2023, Example 1.11.5), we have the fact that for $1 < r < q < \infty$, the sequence $(f_n^r)_{n \in \mathbb{N}}$ is AUI with respect to $(Q_n)_{n \in \mathbb{N}}$, P-a.s., if

$$\limsup_{n \to \infty} \int_{\mathbb{T}} |f_n(\omega, \theta)|^q \, \mathcal{Q}_n(\omega, \mathrm{d}\theta) < \infty,$$

holds, for every ω on a P-a.s. set. This is because, for each $\delta > 0$,

$$\limsup_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^r \chi_{\{\tau \in \mathbb{T} : |f_n(\omega, \theta)| \ge \delta\}} Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \limsup_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \mathrm{d}\theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \theta) \le \delta^{r-q} \log_{n \to \infty} \int_{\mathbb{T}} \left| f_n(\omega, \theta) \right|^q Q_n(\omega, \theta)$$

where $\lim_{\delta \to \infty} \delta^{r-q} = 0$.

2.3 Convergence of the BPIC and WBIC

We are now ready to state sufficient conditions for the convergence of the $BPIC_n$ and $WBIC_n$ to their limits. The following further assumptions are required:

A5 The sequence of average log-likelihoods $(n^{-1}\sum_{i=1}^{n} \log p(X_i|\cdot))_{n\in\mathbb{N}}$ is AUI with respect to $(\prod_{n=1}^{\beta_n})_{n\in\mathbb{N}}$, P-a.s.

A6 The sequence $\left(\Pi_{n}^{\beta_{n}}\right)_{n\in\mathbb{N}}$ converges to $\Pi_{0}\in\mathcal{P}\left(\mathbb{T}\right)$, P-a.s.w.

Proposition 2. Assume that $(X_i)_{i \in \mathbb{N}}$ are IID and that A1 and A2 hold. If A5 and A6 hold for $\beta_n = 1$, then

$$BPIC_n \xrightarrow[n \to \infty]{P-a.s.} -2 \int_{\mathbb{T}} E\left[\log p\left(X|\theta\right)\right] \Pi_0\left(\mathrm{d}\theta\right).$$
(11)

If A5 and A6 hold with $n\beta_n \to \infty$, then

WBIC_n
$$\xrightarrow[n \to \infty]{P-a.s.}{} -2 \int_{\mathbb{T}} E\left[\log p\left(X|\theta\right)\right] \Pi_0\left(\mathrm{d}\theta\right).$$
 (12)

If we replace A6 with A3 and A4, we instead obtain the following more specific outcome.

Corollary 1. Assume that $(X_i)_{i\in\mathbb{N}}$ are IID and $\Pi(\mathbb{B}_{\rho}(\theta_0)) > 0$, for every $\rho > 0$. If A1-A5 hold for $\beta_n = 1$, then

$$\operatorname{BPIC}_{n} \xrightarrow[n \to \infty]{\text{P-a.s.}} -2\operatorname{E}\left[\log \operatorname{p}\left(X|\theta_{0}\right)\right].$$

If A1-A5 hold with $n\beta_n \to \infty$, then

WBIC_n
$$\xrightarrow[n \to \infty]{\text{P-a.s.}} -2\text{E} \left[\log p\left(X|\theta_0\right)\right]$$
.

2.4 Convergence of the DIC

Finally, to obtain the limit of the DIC, we require the convergence of the limit of the second right-hand term of (5). To this end, we require the following assumption:

A7 The norm $\|\cdot\|$ is AUI with respect to $(\Pi_n)_{n\in\mathbb{N}}$, P-a.s.

Via Proposition 4, we can show that when taken together with A1, A2, and A6, A7 implies:

$$\bar{\theta}_n \xrightarrow[n \to \infty]{\text{P-a.s.}} \bar{\theta}_0 = \left(\int_{\mathbb{T}} \theta_1 \Pi_0 \left(\mathrm{d}\theta \right), \dots, \int_{\mathbb{T}} \theta_j \Pi_0 \left(\mathrm{d}\theta \right), \dots, \int_{\mathbb{T}} \theta_p \Pi_0 \left(\mathrm{d}\theta \right) \right).$$
(13)

If we replace A6 by A3 and A4, then $\bar{\theta}_0 = \theta_0$. We can establish that $\delta_{\bar{\theta}_n} \xrightarrow[n \to \infty]{n \to \infty} b_{\bar{\theta}_0}$, by the same argument as presented prior to (4). Then, A1 and A2 imply the continuous convergence of $\left(n^{-1}\sum_{i=1}^n \log p(X_i|\cdot)\right)_{n \in \mathbb{N}}$ to E [log p (X|·)], P-a.s., and thus

$$\frac{2}{n} \sum_{i=1}^{n} \log p\left(X_i | \bar{\theta}_n\right) \xrightarrow[n \to \infty]{\text{P-a.s.}} 2\text{E}\left[\log p\left(X | \bar{\theta}_0\right)\right].$$
(14)

We therefore establish the following convergence result for DIC_n .

Proposition 3. Assume that $(X_i)_{i\in\mathbb{N}}$ are IID and $\Pi(\mathbb{B}_{\rho}(\theta_0)) > 0$, for every $\rho > 0$. If A1, A2, and A5–A7 hold for $\beta_n = 1$, then

$$\operatorname{DIC}_{n} \xrightarrow[n \to \infty]{P-a.s.} -4 \int_{\mathbb{T}} \operatorname{E} \left[\log p\left(X | \theta \right) \right] \Pi_{0} \left(\mathrm{d} \theta \right) + 2 \operatorname{E} \left[\log p\left(X | \bar{\theta}_{0} \right) \right]$$

If A6 is replaced by A3 and A4, then

$$\operatorname{DIC}_n \xrightarrow[n \to \infty]{\text{P-a.s.}} -2\operatorname{E}\left[\log \operatorname{p}\left(X|\theta_0\right)\right].$$

3 Examples

We providing three technical examples, followed by numerical results to demonstrate our theoretical findings. Derivations and technical details are provided in the Appendix.

3.1 The geometric model

We begin with the simple case of the geometric model, whose probability mass function (PMF) is characterized by

$$p(x|\theta) = (1-\theta)^x \theta,$$

for $x \in \mathbb{X} = \mathbb{N} \cup \{0\}$ and $\theta \in \mathbb{T} = (0, 1)$. We will endow θ with the uniform prior measure on \mathbb{T} ; that is, Π has density $\pi(\theta) = 1$ for each $\theta \in \mathbb{T}$. With the sample mean as $\bar{X}_n = n^{-1} \sum_{i=1}^n X_i$, let $\bar{\theta}_n$ denote the posterior mean value, and suppose that $\mathbb{E}|X| < \infty$. Then, for $\beta_n > 0$ the information criteria have the forms:

$$DIC_n = -\frac{4}{n} \int_0^1 \left\{ \bar{X}_n \log \left(1 - \theta\right) + \log \theta \right\} \Pi_n \left(d\theta\right) + \frac{2}{n} \sum \left\{ \bar{X}_n \log \left(1 - \bar{\theta}_n\right) + \log \bar{\theta}_n \right\},$$
$$BPIC_n = -2 \int_0^1 \left\{ \bar{X}_n \log \left(1 - \theta\right) + \log \theta \right\} \Pi_n \left(d\theta\right) + \frac{2}{n},$$
$$WBIC_n = -2 \int_0^1 \left\{ \bar{X}_n \log \left(1 - \theta\right) + \log \theta \right\} \Pi_n^{\beta_n} \left(d\theta\right).$$

Let A1–A6 hold with $\theta_0 = \{1 + EX\}^{-1}$. Then, Corollary 1 and Proposition 3 imply that, as $n \to \infty$, with $n\beta_n \to \infty$, we have

$$\operatorname{DIC}_{n}, \operatorname{BPIC}_{n}, \operatorname{WBIC}_{n} \xrightarrow[n \to \infty]{} 2 \log \{1 + \operatorname{E}X\} - 2\operatorname{E}X \log \left\{\frac{\operatorname{E}X}{1 + \operatorname{E}X}\right\}.$$
 (15)

3.2 The normal model

We now consider the normal model, defined by the PDF:

$$p(x|\theta) = (2\pi)^{-p/2} \exp\left\{-\frac{1}{2} ||x - \theta||^2\right\},\$$

for $x \in \mathbb{X} = \mathbb{R}^p$ and $\theta \in \mathbb{T} = \mathbb{R}^p$, equipped with the normal prior measure Π , defined by the PDF

$$\pi(\theta) = (2\pi)^{-p/2} \exp\left\{-\frac{1}{2} \|\theta - \mu\|^2\right\},\$$

for some $\mu \in \mathbb{R}^p$. We will take $\bar{\theta}_n$ to be the multivariate mean of the posterior distribution, and assume that X has a finite second moment: $\mathbb{E}\left[\|X\|^2\right] < \infty$. Then, for $\beta_n > 0$,

$$DIC_{n} = p \log (2\pi) + \frac{2}{n} \int_{\mathbb{R}^{p}} \sum_{i=1}^{n} \|X_{i} - \theta\|^{2} \Pi_{n} (d\theta) - \frac{1}{n} \sum_{i=1}^{n} \|X_{i} - \bar{\theta}_{n}\|^{2},$$

$$BPIC_{n} = p \log (2\pi) + \frac{1}{n} \int_{\mathbb{R}^{p}} \sum_{i=1}^{n} \|X_{i} - \theta\|^{2} \Pi_{n} (d\theta) + \frac{2p}{n},$$
$$WBIC_{n} = p \log (2\pi) + \frac{1}{n} \int_{\mathbb{R}^{p}} \sum_{i=1}^{n} \|X_{i} - \theta\|^{2} \Pi_{n}^{\beta_{n}} (d\theta).$$
(16)

Let A1–A6 hold with $\theta_0 = EX$. Then, Corollary 1 and Proposition 3 imply that, as $n \to \infty$, with $n\beta_n \to \infty$, we have

$$\operatorname{DIC}_n, \operatorname{BPIC}_n, \operatorname{WBIC}_n \xrightarrow[n \to \infty]{\text{P-a.s.}} p \log (2\pi) + \operatorname{E} \left[\|X\|^2 \right] - \|\operatorname{E} X\|^2.$$
 (17)

3.3 The Laplace model

We conclude by considering a nonconjugate prior pair, taking a Laplace model for $X \in \mathbb{X} = \mathbb{R}$, with PDF

$$p(x|\theta) = \frac{1}{2\gamma} \exp\left\{-\frac{|x-\mu|}{\gamma}\right\},$$

where $\theta = (\mu, \gamma) \in \mathbb{T} = [-m, m] \times [1/s, s]$, with m > 0 and s > 1. We will endow θ with a prior measure Π whose density π is strictly positive on \mathbb{T} .

Write Med (X) as the median of X and assume the regularity assumptions that $E|X| < \infty$, $\mu_0 = Med(X) \in [-m,m]$ and $\gamma_0 = E|X - Med(X)| \in [s^{-1},s]$. Further write $\bar{\theta}_n = (\bar{\mu}_n, \bar{\gamma}_n)$ to be the posterior mean. Then, for $\beta_n > 0$,

$$DIC_{n} = 4 \int_{-\infty}^{\infty} \log (2\gamma) + \frac{1}{\gamma n} \sum_{i=1}^{n} |X_{i} - \mu| \Pi_{n} (d\theta) - 2 \left\{ \log (2\gamma_{0}) + \frac{1}{\gamma_{0} n} \sum_{i=1}^{n} |X_{i} - \mu_{0}| \right\},\$$
$$BPIC_{n} = 2 \int_{-\infty}^{\infty} \log (2\gamma) + \frac{1}{\gamma n} \sum_{i=1}^{n} |X_{i} - \mu| \Pi_{n} (d\theta) + \frac{4}{n},\$$
$$WBIC_{n} = 2 \int_{-\infty}^{\infty} \log (2\gamma) + \frac{1}{\gamma n} \sum_{i=1}^{n} |X_{i} - \mu| \Pi_{n}^{\beta_{n}} (d\theta).$$

Let A1–A6 hold with $\theta_0 = (\mu_0, \gamma_0)$. Then, Corollary 1 and Proposition 3 imply that, as $n \to \infty$, with $n\beta_n \to \infty$, we have

$$\operatorname{DIC}_n, \operatorname{BPIC}_n, \operatorname{WBIC}_n \xrightarrow[n \to \infty]{\text{P-a.s.}} 2 \log (2\gamma_0) + 2.$$

3.4 Numerical examples

The following numerical experiments are conducted in the R programming language. Code for all results are made available on our Github page: https://github.com/hiendn/BayesianIC.

3.4.1 DIC for the geometric model

We continue with the geometric distribution example from Section 3.1. Let us broaden the scenario by replacing the uniform prior with a beta distribution prior measure Π , with parameters $\alpha, \beta > 0$, whose

law we shall write as Beta (α, β) . Then, conjugacy implies that the posterior measure Π_n is the law Beta (a_n, b_n) , where $a_n = n + \alpha$ and $b_n = n\bar{X}_n + \beta$. Let us write $\psi : \mathbb{R}_{>0} \to \mathbb{R}_0$ to be the digamma function (defined in the Appendix) for which we have the Poincare-type approximation (Olver et al., 2010, Eq. 5.11.2): for each a > 0,

$$\psi\left(a\right) \approx \log a - \frac{1}{2a}$$

Using this fact, we derive the following large n approximation of DIC_n in the Appendix:

$$DIC_n \approx -2\log\left\{\frac{n+\alpha}{n+\alpha+n\bar{X}_n+\beta}\right\} + 2\frac{n\bar{X}_n+\beta}{(n+\alpha)\left(n+\alpha+n\bar{X}_n+\beta\right)} \\ -2\bar{X}_n\log\left\{\frac{n\bar{X}_n+\beta}{n+\alpha+n\bar{X}_n+\beta}\right\} + 2\frac{(n+\alpha)\bar{X}_n}{\left(n\bar{X}_n+\beta\right)\left(n+\alpha+n\bar{X}_n+\beta\right)}.$$

Equipped with this computation formula, we design a simulation study to check the convergence of DIC_n to its large sample limit, for a range of values of θ_0 , with various prior hyperparameters α and β , and sample sizes n. The values for θ_0 are chosen between 0.1 and 0.9, with prior hyperparameters chosen as pairs with $\alpha, \beta \in \{1, 10, 100\}$. Sample sizes are chosen in the range 10^2 to 10^7 , and each scenario is repeated 10 times. For each θ_0 , the limiting value of DIC_n is as per (15):

$$\operatorname{DIC}_{n} \xrightarrow[n \to \infty]{\text{P-a.s.}} -2\operatorname{E}\left[\log \operatorname{p}\left(X|\theta_{0}\right)\right] = -2\left\{\frac{1-\theta_{0}}{\theta_{0}}\log\left(1-\theta_{0}\right)+\log\left(\theta_{0}\right)\right\}$$

Here, we make the substitution $\theta_0 = \{1 + EX\}^{-1}$. From Figure 1, we see that for smaller *n*, DIC_n can have high variance, but as *n* increases, we indeed have convergence to the theoretical limit for each θ_0 .

3.4.2 WBIC for the normal model

We next consider the normal model in Section 3.2 with p = 1. Here, we wish to investigate the convergence of the WBIC_n for different choices of sequences β_n that satisfy the condition $n\beta_n \to \infty$ as $n \to \infty$. To this end, we employ a unit variance normal prior distribution Π with mean hyperperameter $\mu = 0$, whose power posterior $\Pi_n^{\beta_n}$ has a normal law (see the Appendix):

$$\mathcal{N}\left(\frac{n\beta_n\bar{X}_n+\mu}{n\beta_n+1},\frac{1}{n\beta_n+1}\right).$$

Using this fact, we can evaluate (16) via numerical integration, under the original power sequence $\beta_n = 1/\log n$ of Watanabe (2013) along with alternative choices $\beta_n = 1/\log \log n$, $\beta_n = 1$, and $\beta_n = 1/\sqrt{n}$. We will also consider choices of $\beta_n = 1/n$ and $\beta_n = 1/\{n \log n\}$ in order to investigate the necessity of the condition $n\beta_n \to \infty$. Figure 2 displays our results for 10 replicates of each sample sizes n ranging from 10^1 to 10^5 , with data simulated from the normal law N ($\theta_0, 1$), with mean parameter $\theta_0 = 1$.

The simulation outcome is as expected, with $WBIC_n$ appearing to converge to the predicted limit as per (17) of

$$-2\mathrm{E}\left[\log p\left(X|\theta_0\right)\right] = \log\left(2\pi\right) + 1,$$

in all cases where $n\beta_n \to \infty$. In the cases where $n\beta_n \to 0$ (i.e., Figure 2 (e, f)) we observe that WBIC_n is




Figure 1: DIC_n values for the geometric model. Points indicate the replicates of DIC_n evaluations for a range of values of θ_0 with different sample sizes n, while the lines correspond to the limiting value of DIC_n as $n \to \infty$. Both points and lines are coloured according to the true value of θ_0 in each case.




Figure 2: Simulated WBIC_n values under various choices of β_n : (a) $1/\log n$, (b) $1/\log \log n$, (c) 1, (d) $1/\sqrt{n}$, (e) 1/n, and (f) $1/\{n \log n\}$. Points indicate the replicates of WBIC_n evaluations, and the solid lines indicate the theoretical limit under Corollary 1.

either diverging or converging to a limit that is not the one predicted by Corollary 1, providing evidence towards the necessity of the condition $n\beta_n \to \infty$.

4 Proofs and technical results

4.1 Proof of Proposition 1

As promised, we shall prove a more general result that implies Proposition 1. Following Section 1, we define a sequence of data-dependent utility functions $(u_n)_{n \in \mathbb{N}}$, where, for each $\omega \in \Omega$ and $n \in \mathbb{N}$:

$$u_n(\omega,\theta) = u_n(X_1(\omega),\ldots,X_n(\omega);\theta).$$

We make the following assumptions regarding $(u_n)_{n \in \mathbb{N}}$, mirroring assumptions A1–A4.

- **B1** For each $n \in \mathbb{N}$, the utility u_n is Caratheodory in the sense that $u_n(\cdot, \theta)$ is measurable for each $\theta \in \mathbb{T}$ and $u_n(\omega, \cdot)$ is continuous for each $\omega \in \Omega$.
- **B2** There exists a deterministic function $u: \mathbb{T} \to \mathbb{R}$ and a P-a.s. set on which for every compact $\mathbb{K} \subset \mathbb{T}$,

$$\sup_{\theta \in \mathbb{K}} |u_n(\omega, \theta) - u(\theta)| \xrightarrow[n \to \infty]{} 0.$$
(18)

B3 There exists a unique maximizer $\theta_0 \in \mathbb{T}$ of $u(\theta)$, in the sense that, for every $\theta \in \mathbb{T} \setminus \{\theta_0\}$,

$$u(\theta) < u(\theta_0).$$

B4 There exists a compact set $\mathbb{S} \subseteq \mathbb{T}$ containing θ_0 , such that there is a P-a.s. set on which

$$\limsup_{n \to \infty} \sup_{\theta \in \mathbb{S}^{c}} u_{n} \left(\omega, \theta \right) < u \left(\theta_{0} \right).$$

Let $(\Pi_n)_{n\in\mathbb{N}}$ be a sequence of random measures, where $\Pi_n: \Omega \times \mathfrak{B}_{\mathbb{T}} \to \mathbb{R}_{\geq 0}$ for each $n \in \mathbb{N}$, defined by

$$\Pi_{n}(\omega,\mathbb{A}) = \frac{\int_{\mathbb{A}} \exp\left(\gamma_{n} u_{n}(\omega,\theta)\right) \Pi\left(\mathrm{d}\theta\right)}{\int_{\mathbb{T}} \exp\left(\gamma_{n} u_{n}(\omega,\theta)\right) \Pi\left(\mathrm{d}\theta\right)},$$

for each $\omega \in \Omega$, where $\mathbb{A} \in \mathfrak{B}_{\mathbb{T}}$. Here $(\gamma_n)_{n \in \mathbb{N}} \subset \mathbb{R}_{>0}$ diverges to infinity. The measures Π_n extend the generalized or Gibbs posterior measures and appear in the works of Zhang (2006), Bissiri et al. (2016), Miller (2021), and Knoblauch et al. (2022), among others. We prove the following result.

Theorem 2. If $\Pi (\mathbb{B}_{\rho}(\theta_0)) > 0$, for every $\rho > 0$, and B1–B4 hold, then the sequence $(\Pi_n)_{n \in \mathbb{N}}$ is consistent in the sense that for every ω on a P-a.s. set, for every $\epsilon > 0$,

$$\Pi_n\left(\omega, \bar{\mathbb{B}}_{\epsilon}\left(\theta_0\right)\right) \xrightarrow[n \to \infty]{} 1.$$

Proof of Theorem 2. We start by selecting an arbitrary $\epsilon > 0$. By B2 we have that u is continuous on every compact set \mathbb{K} , as u is the uniform limit of continuous functions $u_n(\omega, \cdot)$ (cf. Remmert 1991, Thm. 3.1.5). This then implies that u is continuous on $\mathbb{T} \subseteq \mathbb{R}^p$. For any $\mathbb{A}, \mathbb{B} \subset \mathbb{R}^p$ and $\epsilon > 0$, let

$$\bar{\mathbb{B}}_{\epsilon}(\mathbb{A}) = \left\{ x \in \mathbb{R}^p : \inf_{y \in \mathbb{A}} \|x - y\| \le \epsilon \right\},\$$

and define the distance $d(\mathbb{A}, \mathbb{B}) = \inf_{x \in \mathbb{A}} \inf_{y \in \mathbb{B}} \{ \|x - y\| \}$. When \mathbb{T} is open, $d(\mathbb{S}, \mathbb{T}^c) > 0$ and so there is an $\epsilon^* > 0$ such that $\overline{\mathbb{B}}_{\epsilon^*}(\mathbb{S}) \subseteq \mathbb{T}$. Let $\epsilon_0 = \sup_{\theta \in \overline{\mathbb{B}}_{\epsilon^*}(\mathbb{S})} d(\theta, \mathbb{S})$ and note that because \mathbb{S} is closed and $\mathbb{S} \neq \mathbb{T}$, $\epsilon_0 > 0$. When \mathbb{T} is closed, we can choose $\epsilon_0 > 0$, arbitrarily.

We define the maximum function $u^*: [0, \epsilon_0] \to \mathbb{R}$ given by

$$u^{*}\left(\rho\right) = \max_{\theta \in \mathbb{B}_{\rho}^{c}(\theta_{0}) \cap \bar{\mathbb{B}}_{\epsilon_{0}}(\mathbb{S})} u\left(\theta\right),$$

and observe that for $\rho \in [0, \epsilon_0]$, $\mathbb{B}_{\rho}^{c}(\theta_0) \cap \overline{\mathbb{B}}_{\epsilon_0}(\mathbb{S})$ is compact and non-empty.

By B3 we can then take $\epsilon_1 \in (0, \min\{\epsilon, \epsilon_0\})$ such that

$$u^*(\epsilon_1) < u(\theta_0),$$

where we note that this inequality holds for any $\epsilon_1 > 0$. Now define the minimum function $u_* : \mathbb{R}_{\geq 0} \to \mathbb{R}$ by

$$u_{*}\left(\rho\right) = \min_{\theta \in \bar{\mathbb{B}}_{\rho}(\theta_{0})} u\left(\theta\right).$$

Further note that the correspondence

$$\phi(\rho) = \overline{\mathbb{B}}_{\rho}(\theta_0)$$

is continuous at 0 and that u_* is decreasing. Berge's maximum theorem (see, e.g. Beer 1993, Thm. 6.3.8) then gives that u_* is continuous at 0; i.e.

$$\lim_{\rho \to 0} u_*(\rho) = u(\theta_0).$$

We can then choose an $\epsilon_{2,1} \in (0, \epsilon_1)$ so that

$$u_*(\epsilon_{2,1}) > u^*(\epsilon_1).$$
 (19)

B4 allows us to choose a $\epsilon_{2,2} > 0$ so that

$$u_*(\epsilon_{2,2}) > \limsup_{n \to \infty} \sup_{\theta \in \bar{\mathbb{B}}_{\epsilon_0}^c} u_n(\omega, \theta).$$
⁽²⁰⁾

Let $\epsilon_2 = \min{\{\epsilon_{2,1}, \epsilon_{2,2}\}}$. By the monotonicity of u_* and u^* , (19) and (20) still hold for ϵ_2 . Then take any $\delta > 0$ such that,

$$u_*(\epsilon_2) - \delta > u^*(\epsilon_1) + \delta, \tag{21}$$

and

$$u_*(\epsilon_2) - \delta > \limsup_{n \to \infty} \sup_{\theta \in \bar{\mathbb{B}}_{\epsilon_0}^c(\mathbb{S})} u_n(\omega, \theta).$$
(22)

We thus deduce the following expressions:

$$\begin{split} \Pi_{n}\left(\omega,\bar{\mathbb{B}}_{\epsilon}\left(\theta_{0}\right)\right) &\geq \Pi_{n}\left(\omega,\bar{\mathbb{B}}_{\epsilon_{1}}\left(\theta_{0}\right)\right) \\ &= \frac{\int_{\bar{\mathbb{B}}_{\epsilon_{1}}\left(\theta_{0}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}{\int_{\mathbb{T}}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)} \\ &= \left\{\frac{\int_{\mathbb{T}}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}{\int_{\bar{\mathbb{B}}_{\epsilon_{1}}\left(\theta_{0}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}\right\}^{-1} \\ &= \left\{1 + \frac{\int_{\bar{\mathbb{B}}_{\epsilon_{1}}\left(\theta_{0}\right)\cap\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}{\int_{\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}\right\}^{-1} \\ &= \left\{1 + \frac{\int_{\bar{\mathbb{B}}_{\epsilon_{1}}\left(\theta_{0}\right)\cap\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right) + \int_{\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}{\int_{\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}\right\}^{-1} \\ &\geq \left\{1 + \frac{\int_{\mathbb{B}_{\epsilon_{1}}\left(\theta_{0}\right)\cap\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right) + \int_{\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}{\int_{\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)}\exp\left(\gamma_{n}u_{n}\left(\omega,\theta\right)\right)\Pi\left(\mathrm{d}\theta\right)}\right\}^{-1}. \end{split}$$

We now seek upper bounds for the two ratios

$$a_{n}(\omega) = \frac{\int_{\mathbb{B}_{\epsilon_{1}}^{c}(\theta_{0})\cap\bar{\mathbb{B}}_{\mathbb{S}}(\theta_{0})} \exp\left(\gamma_{n}u_{n}(\omega,\theta)\right) \Pi\left(\mathrm{d}\theta\right)}{\int_{\bar{\mathbb{B}}_{\epsilon_{2}}(\theta_{0})} \exp\left(\gamma_{n}u_{n}(\omega,\theta)\right) \Pi\left(\mathrm{d}\theta\right)}$$

and

$$b_{n}(\omega) = \frac{\int_{\bar{\mathbb{B}}_{\epsilon_{0}}^{c}(\mathbb{S})} \exp\left(\gamma_{n}u_{n}(\omega,\theta)\right) \Pi\left(\mathrm{d}\theta\right)}{\int_{\bar{\mathbb{B}}_{\epsilon_{2}}(\theta_{0})} \exp\left(\gamma_{n}u_{n}(\omega,\theta)\right) \Pi\left(\mathrm{d}\theta\right)}.$$

Starting with a_n , because $\mathbb{B}_{\epsilon_1}^c(\theta_0) \cap \overline{\mathbb{B}}_{\epsilon_0}(\mathbb{S})$ is compact, B2 provides a P-a.s. set Ω_u such that for each $\omega \in \Omega_u$, there is an $n(\omega)$, such that whenever $n > n(\omega)$,

$$|u_n(\omega,\theta) - u(\theta)| < \delta,$$

for all $\theta \in \mathbb{B}_{\epsilon_1}^c(\theta_0) \cap \overline{\mathbb{B}}_{\epsilon_0}(\mathbb{S})$. That is, for each $n > n(\omega)$ and each $\theta \in \mathbb{B}_{\epsilon_1}^c(\theta_0) \cap \overline{\mathbb{B}}_{\epsilon_0}(\mathbb{S})$, it follows that

$$u(\theta) - \delta \le u_n(\omega, \theta) \le u(\theta) + \delta.$$
⁽²³⁾

We then have, for each $\omega \in \Omega_u$ and each $n > n(\omega)$,

$$\begin{split} a_{n}\left(\omega\right) &\leq \frac{\int_{\mathbb{B}_{\epsilon_{1}}^{c}\left(\theta_{0}\right)\cap\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)}\exp\left(\gamma_{n}\left\{u\left(\theta\right)+\delta\right\}\right)\Pi\left(\mathrm{d}\theta\right)}{\int_{\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)}\exp\left(\gamma_{n}\left\{u\left(\theta\right)-\delta\right\}\right)\Pi\left(\mathrm{d}\theta\right)} \\ &\leq \frac{\Pi\left(\mathbb{B}_{\epsilon_{1}}^{c}\left(\theta_{0}\right)\cap\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)\right)\exp\left(\gamma_{n}\left\{\max_{\theta\in\mathbb{B}_{\epsilon_{1}}^{c}\left(\theta_{0}\right)\cap\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)}u\left(\theta\right)+\delta\right\}\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)\right)\exp\left(\gamma_{n}\left\{\min_{\theta\in\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)}u\left(\theta\right)-\delta\right\}\right)} \\ &= \frac{\Pi\left(\mathbb{B}_{\epsilon_{1}}^{c}\left(\theta_{0}\right)\cap\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)\right)\exp\left(\gamma_{n}\left\{u^{*}\left(\epsilon_{1}\right)+\delta\right\}\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)\right)\exp\left(\gamma_{n}\left\{u^{*}\left(\epsilon_{1}\right)-u_{*}\left(\epsilon_{2}\right)+2\delta\right\}\right)} \\ &= \frac{\Pi\left(\mathbb{B}_{\epsilon_{1}}^{c}\left(\theta_{0}\right)\cap\bar{\mathbb{B}}_{\epsilon_{0}}\left(\mathbb{S}\right)\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)\right)}\exp\left(\gamma_{n}\left\{u^{*}\left(\epsilon_{1}\right)-u_{*}\left(\epsilon_{2}\right)+2\delta\right\}\right). \end{split}$$

Note that because $\Pi(\mathbb{B}_{\rho}(\theta_0)) > 0$, for all $\rho > 0$, we have that the above expressions are well defined. Similarly, (23) implies that for each $\omega \in \Omega_u$ and each $n > n(\omega)$,

$$\begin{split} b_{n}\left(\omega\right) &\leq \frac{\Pi\left(\bar{\mathbb{B}}_{\epsilon_{0}}^{c}(\mathbb{S})\right)\exp\left(\gamma_{n}\sup_{\theta\in\bar{\mathbb{B}}_{\epsilon_{0}}^{c}(\mathbb{S})}u_{n}\left(\omega,\theta\right)\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)\right)\exp\left(\gamma_{n}\left\{u_{*}\left(\epsilon_{2}\right)-\delta\right\}\right)} \\ &= \frac{\Pi\left(\bar{\mathbb{B}}_{\epsilon_{0}}^{c}(\mathbb{S})\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_{2}}\left(\theta_{0}\right)\right)}\exp\left(\gamma_{n}\left\{\sup_{\theta\in\bar{\mathbb{B}}_{\epsilon_{0}}^{c}(\mathbb{S})}u_{n}\left(\omega,\theta\right)-u_{*}\left(\epsilon_{2}\right)+\delta\right\}\right). \end{split}$$

Since $\gamma_n \xrightarrow[n \to \infty]{} \infty$, via (21) we have

$$\limsup_{n \to \infty} a_n(\omega) \leq \lim_{n \to \infty} \frac{\Pi\left(\mathbb{B}_{\epsilon_1}^{c}(\theta_0) \cap \bar{\mathbb{B}}_{\epsilon_0}(\mathbb{S})\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_2}(\theta_0)\right)} \exp\left(\gamma_n\left\{u^*\left(\epsilon_1\right) - u_*\left(\epsilon_2\right) + 2\delta\right\}\right)$$
$$= \frac{\Pi\left(\mathbb{B}_{\epsilon_1}^{c}\left(\theta_0\right) \cap \bar{\mathbb{B}}_{\epsilon_0}(\mathbb{S})\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_2}(\theta_0)\right)} \exp\left(\lim_{n \to \infty} \gamma_n\left\{u^*\left(\epsilon_1\right) - u_*\left(\epsilon_2\right) + 2\delta\right\}\right)$$
$$= 0,$$

and via (22) we have

$$\begin{split} \limsup_{n \to \infty} b_n(\omega) &\leq \lim_{n \to \infty} \frac{\Pi\left(\bar{\mathbb{B}}_{\epsilon_0}^{c}(\mathbb{S})\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_2}(\theta_0)\right)} \exp\left(\gamma_n \left\{\sup_{\theta \in \bar{\mathbb{B}}_{\epsilon_0}^{c}(\mathbb{S})} u_n(\omega, \theta) - u_*(\epsilon_2) + \delta\right\}\right) \\ &= \frac{\Pi\left(\bar{\mathbb{B}}_{\epsilon_0}^{c}(\mathbb{S})\right)}{\Pi\left(\bar{\mathbb{B}}_{\epsilon_2}(\theta_0)\right)} \exp\left(\lim_{n \to \infty} \gamma_n \left\{\sup_{\theta \in \bar{\mathbb{B}}_{\epsilon_0}^{c}(\mathbb{S})} u_n(\omega, \theta) - u_*(\epsilon_2) + \delta\right\}\right) \\ &= 0. \end{split}$$

Because $a_n, b_n \ge 0$ this gives

$$\lim_{n \to \infty} b_n(\omega) = \lim_{n \to \infty} a_n(\omega) = 0$$

Thus,

$$\lim_{n \to \infty} \Pi_n \left(\omega, \bar{\mathbb{B}}_{\epsilon} \left(\theta_0 \right) \right) \ge \left\{ 1 + \lim_{n \to \infty} a_n \left(\omega \right) + \lim_{n \to \infty} b_n \left(\omega \right) \right\}^{-1} = 1,$$

as required. Since this conclusion is obtained as an intersection of finitely many P-a.s. events, it also occurs on a P-a.s. set. $\hfill \Box$

We note that this proof follows essentially the same steps as that of Ghosh & Ramamoorthi (2003, Thm. 1.3.4) whose proof is specialized to the situation where

$$u_n(\omega,\theta) = \frac{1}{n} \sum_{i=1}^n \log p\left(X_i(\omega)|\theta\right)$$
(24)

is the log-likelihood of an IID sequence $(X_i)_{i \in [n]}$ and $\gamma_n = n$. We also point out that our result is complementary to the abstract methods of Miller (2021, Sec. 2) (Theorem 3(1,2), particularly), where the author provides sufficient conditions for the posterior consistency of non-stochastic generalized posterior measures in the case when $\gamma_n = n$. In fact our assumptions B1–B4 can be read as stochastic counterparts of Miller (2021, Thm. 3(2)), specialised to $\mathbb{T} \subset \mathbb{R}^p$. Our proof establishes the required stochastic assumptions in the case when $\mathbb{T} \subset \mathbb{R}^p$ and makes the generalization to generic sequences $(\gamma_n)_{n \in \mathbb{N}}$. To prove Proposition 1 using Theorem 2, we observe that B1–B4 are satisfied by A1–A4 when taking u_n of form (24). The result is then implied by setting $\gamma_n = n\beta_n$.

Remark 1. From the proof of Theorem 2, we see that the exponential function can be replaced by $\exp \circ \eta$, where $\eta : \mathbb{R} \to \mathbb{R}$ is monotone and satisfies the condition that if $\gamma_n \to \infty$, then

$$\eta\left(a\gamma_n\right) - \eta\left(b\gamma_n\right) \to \infty$$

for every a > b. For example, one may consider any increasing homogeneous function, such as $\eta(x) = x^{2k+1}$, for $k \in \mathbb{N} \cup \{0\}$. We note that, to the best of our knowledge, this fact has not been reported in any prior works, although we also do not know of any immediate application.

When \mathbb{T} is convex, we say that $u: \mathbb{T} \to \mathbb{R}$ is quasiconvex if for each $\theta, \tau \in \mathbb{T}$ and $\lambda \in [0, 1]$,

$$u\left(\lambda\theta + (1-\lambda)\tau\right) \le \max\left\{u\left(\theta\right), u\left(\tau\right)\right\}.$$

Furthermore, we say that u is quasiconcave if -u is quasiconvex. The class of quasiconvex functions include the convex functions, and form a generalization that preserves the property of convex level sets. Using the notion of quasiconvexity, we provide the following result for verifying B4, providing a generalization of the convexity-based result of Miller (2021, Lem. 27), and when taken together with Theorem 2, we obtain a stochastic quasiconvexity-based version of Miller (2021, Thm. 3(3)).

Lemma 1. Let $\mathbb{T} \subset \mathbb{R}^p$ be convex. Suppose that $(u_n)_{n \in \mathbb{N}}$ is a sequence of functions, where $u_n : \mathbb{T} \to \mathbb{R}$ is

continuous and quasiconcave, for each $n \in \mathbb{N}$, and

$$\sup_{\theta \in \mathbb{K}} |u_n(\theta) - u(\theta)| \underset{n \to \infty}{\longrightarrow} 0,$$
(25)

for each compact $\mathbb{K} \subset \mathbb{T}$, for some $u : \mathbb{T} \to \mathbb{R}$. Then:

- (i) The function u is continuous and quasiconcave.
- (ii) If $u(\theta) < u(\theta_0)$, for every $\theta \in \mathbb{T} \setminus \{\theta_0\}$, then, for every $\epsilon > 0$,

$$\limsup_{n \to \infty} \sup_{\theta \in \mathbb{T} \setminus \mathbb{B}_{\epsilon}(\theta_0)} u_n(\theta) < u(\theta_0).$$

Proof of Lemma 1. (i) We firstly show that $u : \mathbb{T} \to \mathbb{R}$ is quasiconcave. It suffices to show that the pointwise limit of quasi-concave functions is quasiconcave since every singleton $\{\theta\}$ is compact. Let $\theta, \tau \in \mathbb{T}$ and $\lambda \in [0, 1]$. Then,

$$-u \left(\lambda \theta + (1 - \lambda) \tau\right) = \lim_{n \to \infty} -u_n \left(\lambda \theta + (1 - \lambda) \tau\right)$$
$$\leq \lim_{n \to \infty} \max\left\{-u_n \left(\theta\right), -u_n \left(\tau\right)\right\}$$
$$= \max\left\{-u \left(\theta\right), -u \left(\tau\right)\right\},$$

as required. Then (25) implies that $(u_n)_{n \in \mathbb{N}}$ is continuously convergent to u, which must be continuous as a consequence.

(ii) For any $\epsilon > 0$, such that define the sphere of radius ϵ as

$$\mathbb{S}_{\epsilon} = \left\{ \theta \in \mathbb{T} : \|\theta - \theta_0\| = \epsilon \right\},\$$

and write

$$\alpha_n = u_n(\theta_0) - \sup_{\theta \in \mathbb{S}_{\epsilon}} u_n(\theta)$$
, and $\alpha = u(\theta_0) - \sup_{\theta \in \mathbb{S}_{\epsilon}} u(\theta)$.

Since S_{ϵ} is compact, u_n converges uniformly to u, on S_{ϵ} , and since u and u_n are continuous, for each n, we have $\lim_{n\to\infty} \alpha_n = \alpha$. Additionally, the extreme value theorem gives that these suprema are attained. Then $\alpha > 0$ because $u(\theta) < u(\theta_0)$, for every $\theta \in \mathbb{T} \setminus \{\theta_0\}$.

Let $\tau \in \mathbb{T} \setminus \mathbb{B}_{\epsilon}(\theta_0)$ and choose $\theta_1 \in \mathbb{S}_{\epsilon}$ so that $\theta_1 = \theta_0 + \lambda_1 \{\tau - \theta_0\}$, for some $\lambda_1 \in (0, 1]$. Then define

$$g_n(\lambda) = f_n(\theta_0 + \lambda \{\tau - \theta_0\}),$$

for every $\lambda \in \mathbb{T}_{\tau} = \{t \in \mathbb{R} : \theta_0 + t (\tau - \theta_0) \in \mathbb{T}\}$. Note that g_n is equal to u_n restricted to the line passing through θ_0 , θ_1 , and τ . By Cambini & Martein (2008, Thm. 2.2.10), $g_n : \mathbb{T}_{\tau} \to \mathbb{R}$ is a quasiconcave function and by Florenzano & Le Van (2001, Prop. 8.1.3), g_n is either monotonically increasing, monotonically decreasing, or there exists some $\lambda^* \in \mathbb{T}_{\tau}$, such that g_n is increasing on $(-\infty, \lambda^*) \cap \mathbb{T}_{\tau}$ and decreasing on $(\lambda^*, \infty) \cap \mathbb{T}_{\tau}$. Since $\alpha > 0$ and $\lim_{n\to\infty} \alpha_n = \alpha$, $\exists N \in \mathbb{N}$ such that for each n > N. $\alpha_n > 0$. Thus for n > N,

$$g_n\left(0\right) - g_n\left(\lambda_1\right) \ge \alpha_n > 0.$$

In particular, $g_n(0) > g_n(\lambda_1)$ implying that either g_n is decreasing on \mathbb{T}_{τ} or there exists some $\lambda^* \in (0, \lambda_1)$, where g_n switches from increasing to decreasing. In either case, g_n is decreasing on $(\lambda_1, \infty) \cap \mathbb{T}_{\tau}$, and thus $g_n(\lambda_1) \ge g_n(1)$, implying that

$$g_n(0) - g_n(1) \ge g_n(0) - g_n(\lambda_1) \ge \alpha_n,$$

That is,

$$u_n(\theta_0) - u_n(\tau) \ge u_n(\theta_0) - u_n(\theta_1) \ge \alpha_n.$$

Thus, for n > N, $u_n(\theta_0) \ge u_n(\tau) + \alpha_n$, for each $\tau \in \mathbb{T} \setminus \mathbb{B}_{\epsilon}(\theta_0)$, implying that

$$u_n\left(\theta_0\right) \ge \sup_{\theta \in \mathbb{T} \setminus \mathbb{B}_{\epsilon}\left(\theta_0\right)} u_n\left(\theta\right) + \alpha_n$$

Taking the limit of both sides then yields

$$u(\theta_{0}) \geq \limsup_{n \to \infty} \sup_{\theta \in \mathbb{T} \setminus \mathbb{B}_{\epsilon}(\theta_{0})} u_{n}(\theta) + \alpha > \limsup_{n \to \infty} \sup_{\theta \in \mathbb{T} \setminus \mathbb{B}_{\epsilon}(\theta_{0})} u_{n}(\theta).$$

as required.

Lemma 1 implies that B4 can be replaced with the following quasiconcavity assumption, since we can choose $\mathbb{S} = \mathbb{B}_{\epsilon}(\theta_0)$, for some $\epsilon > 0$.

B4a For almost every $\omega \in \Omega$, $u_n(\omega, \cdot) : \mathbb{T} \to \mathbb{R}$ is continuous and quasiconcave.

4.2 Assumptions A2a and A2b

Next, we wish to verify that A2a and A2b together imply A2. To this end, we require the following uniform strong law of large numbers, which is standard and can be found, for example, in Shapiro et al. (2021, Thm. 9.60). Make the following assumptions:

C1 $\mathbb{K} \subset \mathbb{R}^p$ is compact.

- **C2** $u_n(\omega, \theta) = n^{-1} \sum_{i=1}^n v(X_i(\omega); \theta)$, where v is Caratheodory in the sense that $v(\cdot; \theta)$ is $(\mathbb{X}, \mathfrak{B}_{\mathbb{X}})$ measurable for each $\theta \in \mathbb{T}$ and $v(x; \cdot)$ is continuous for each $x \in \mathbb{X}$.
- **C3** There exists a dominating function $\Delta : \mathbb{X} \to \mathbb{R}$ such that $|v(x;\theta)| \leq \Delta(x)$ for each $\theta \in \mathbb{T}$, such that $\operatorname{E}[\Delta(X)] < \infty$.

Theorem 3. Assume that $(X_i)_{i \in \mathbb{N}}$ are IID. If C1–C3 hold then

$$\sup_{\theta \in \mathbb{K}} \left| u_n\left(\omega, \theta\right) - u\left(\theta\right) \right| \stackrel{\text{P-a.s.}}{\underset{n \to \infty}{\longrightarrow}} 0,$$

where $u(\theta) = E[v(X; \theta)].$

Observe that for any compact set $\mathbb{K} \subset \mathbb{T}$, A1, A2a and A2b together allow us to apply Theorem 3 to conclude that (8) holds P-a.s. on \mathbb{K} . But since $\mathbb{T} \subset \mathbb{R}^p$ is either open or closed, it is hemi-compact and we can construct a countable sequence of compact sets $(\mathbb{K}_i)_{i \in \mathbb{N}}$ such that $\mathbb{T} = \bigcup_{i \in \mathbb{N}} \mathbb{K}_i$. Then, since $\mathbb{K} \subset \mathbb{K}_i$ for each compact set \mathbb{K} and some index $i \geq 1$, A2 is implied by the intersect of the events that (8) hold when $\mathbb{K} = \mathbb{K}_i$, for each $i \in \mathbb{N}$, which each hold P-a.s. and thus jointly hold on a P-a.s. set, as required.

4.3 Convergence of the BPIC and WBIC

Firstly, we wish to demonstrate that Proposition 1 is a direct consequence of Theorem 1. To see that (11) is true, we make the substitutions $f_n(\omega, \cdot) = n^{-1} \sum_{i=1}^n \log p(X_i(\omega) | \cdot)$, for each fixed ω , and $Q_n = \prod_n$. Under A1 and A2, we have the fact that $f_n \xrightarrow[n \to \infty]{\text{P-a.s.}} E[\log p(X_i | \cdot)]$, continuously, by definition. Then, A5 and A6 together with A1 and A2 imply that the conditions for Theorem 1 are met, and we therefore have

$$\lim_{n \to \infty} \int_{\mathbb{T}} \frac{1}{n} \sum_{i=1}^{n} \log p\left(X_{i}\left(\omega\right) | \cdot\right) \Pi_{n}\left(\omega, \mathrm{d}\theta\right) = \int_{\mathbb{T}} \mathrm{E}\left[\log p\left(X | \theta\right)\right] \Pi_{0}\left(\mathrm{d}\theta\right),\tag{26}$$

for almost every $\omega \in \Omega$. Thus, given expression (6), we have

$$\lim_{n \to \infty} \operatorname{BPIC}_{n} = -2 \lim_{n \to \infty} \int_{\mathbb{T}} \frac{1}{n} \sum_{i=1}^{n} \log p(X_{i}(\omega) | \cdot) \Pi_{n}(\omega, \mathrm{d}\theta) + 2 \lim_{n \to \infty} \frac{p}{n}$$
$$= -2 \int_{\mathbb{T}} \operatorname{E} \left[\log p(X|\theta) \right] \Pi_{0}(\mathrm{d}\theta) ,$$

for almost every $\omega \in \Omega$, as required. Result (12) follows similarly, using the definition of the WBIC and taking $Q_n = \prod_n^{\beta_n}$, with $n\beta_n \to \infty$.

Corollary 1 then follows by the fact that A1–A5 imply that the conditions for Proposition 1 and Theorem 1 are both satisfied, with (26) satisfied, and $\Pi_n^{\beta_n}$ converging P-a.s.w. to $\Pi_0 = \delta_{\theta_0}$. The result is obtained by observing that

$$\int_{\mathbb{T}} \operatorname{E}\left[\log p\left(X|\theta\right)\right] \delta_{\theta_0}\left(\mathrm{d}\theta\right) = \operatorname{E}\left[\log p\left(X|\theta_0\right)\right],\tag{27}$$

as per (4).

4.4 Convergence of the DIC

The following result is required to establish the convergence of the posterior mean, i.e. (13).

Proposition 4. Let $(Q_n)_{n \in \mathbb{N}}$ converge to Q, P-a.s.w., and suppose that $\|\cdot\|$ is AUI with respect to $(Q_n)_{n \in \mathbb{N}}$, P-a.s., for any choice of norm $\|\cdot\|$ on \mathbb{R}^p . Then, $\bar{\vartheta}_n : \Omega \to \mathbb{R}^p$ converges to $\bar{\vartheta}_0 \in \mathbb{R}^p$, P-a.s., where

$$\bar{\vartheta}_{n}(\omega) = \left(\int_{\mathbb{T}} \theta_{1} \mathbf{Q}_{n}(\omega, \mathrm{d}\theta), \dots, \int_{\mathbb{T}} \theta_{j} \mathbf{Q}_{n}(\omega, \mathrm{d}\theta), \dots, \int_{\mathbb{T}} \theta_{p} \mathbf{Q}_{n}(\omega, \mathrm{d}\theta)\right),$$

for each $n \in \mathbb{N}$, and

$$\bar{\vartheta}_{0} = \left(\int_{\mathbb{T}} \theta_{1} \mathbf{Q} \left(\mathrm{d} \theta \right), \dots, \int_{\mathbb{T}} \theta_{j} \mathbf{Q} \left(\mathrm{d} \theta \right), \dots, \int_{\mathbb{T}} \theta_{p} \mathbf{Q} \left(\mathrm{d} \theta \right) \right).$$

Proof of Proposition 4. Consider that

$$\left\|\bar{\vartheta}_{n}-\bar{\vartheta}_{0}\right\|_{1}=\sum_{j=1}^{p}\left|\int_{\mathbb{T}}\theta_{j}\mathbf{Q}_{n}\left(\mathrm{d}\theta\right)-\int_{\mathbb{T}}\theta_{j}\mathbf{Q}\left(\mathrm{d}\theta\right)\right|$$

and so $\|\bar{\vartheta}_n - \bar{\vartheta}_0\|_1 \xrightarrow[n \to \infty]{\text{P-a.s.}} 0$ if for each $j \in [p]$,

$$\left| \int_{\mathbb{T}} \theta_j \mathbf{Q}_n \left(\mathrm{d} \theta \right) - \int_{\mathbb{T}} \theta_j \mathbf{Q} \left(\mathrm{d} \theta \right) \right| \stackrel{\text{P-a.s.}}{\underset{n \to \infty}{\longrightarrow}} 0.$$

By Theorem 2, this holds if for each $j, \theta_j \mapsto |\theta_j|$ is AUI with respect to $(Q_n)_{n \in \mathbb{N}}$, P-a.s., which is true if $\|\cdot\|_1$ is AUI with respect to $(Q_n)_{n \in \mathbb{N}}$, P-a.s., since $|\theta_j| \leq \|\theta\|_1$. The result then follows from the equivalence of norms on \mathbb{R}^p .

By (5), the limit of DIC_n is determined by the limits of form (14) and (26). Result (14) follows from the same argument as that of Section 4.3, under assumptions A1, A2, A5, and A6. By Proposition 4, A6 and A7 imply result (13), where A1 and A2 then imply (14) by the definition of continuous convergence, as previously discussed, which establishes the first result of Proposition 3. When A6 is replaced by A3 and A4, $\bar{\theta}_0 = \theta_0$, which thus implies the second result of the proposition by the same argument as (4).

5 Discussion

In this work, we thoroughly investigate the limiting behavior of several Bayesian model selection criteria namely, the DIC, BPIC, and WBIC—as sample sizes approach infinity, under the condition that posterior distributions are consistent. We employ generalizations of the dominated convergence theorem, which have not been widely utilized in the literature, and we establish several general results regarding Bayesian posterior consistency, whose utility extends beyond the scope of this study.

It is important to note that our analysis is qualitative in nature and does not provide finite sample predictions for the behavior of the aforementioned information criteria. Specifically, our investigation does not offer comparative recommendations regarding the utility or merit of these criteria for model or variable selection.

In the frequentist setting, Sin & White (1996), as well as subsequent works by Nguyen (2024) and Westerhout et al. (2024), have explored general conditions under which information criteria based on minima of risk functions can achieve model selection consistency. This consistency is understood as the asymptotic selection of the minimal complexity model that minimizes risk as $n \to \infty$. Conversely, finite sample properties of information criteria-like, penalty-based methods have been studied in works such as those by Massart (2007) and Koltchinskii (2011), which provide broadly applicable bounds on the risk incurred by these methods. Within this context, our results provide only a minimal qualitative guarantee. Specifically, in the language of Claeskens & Hjort (2008), our findings establish that each of the criteria considered exhibits weak consistency. That is, when used to compare models, the criteria will almost surely select one among those that minimize the expected negative log-likelihood. This property, often referred to as efficiency, is analogous to the guarantee provided by the AIC or simply by comparing models using their sample log-likelihood values. Further research is necessary to derive conditions under which Bayesian model selection criteria, based on posterior integrated likelihoods rather than maximum likelihoods, can achieve consistent parsimonious model selection.

Additionally, while our primary analysis focuses on the posterior distribution in the role of the random measure Π_n and the log-likelihood function as the integrand, our methodology extends broadly to the construction of information criteria based on more general objects. For instance, our framework can be applied to analyze the behavior of the variational inference information criteria proposed by You et al. (2014), generalized posterior measures with corresponding risk functions, or scenarios in which the posterior is replaced by Bernstein–von Mises approximations or approximate Bayesian posterior measures that exhibit consistency. Thus, our work provides a flexible template for analyzing a broad class of Bayesian model selection approaches.

Finally, given our focus on information criteria for parametric models, we have focused on establishing sufficient conditions for posterior consistency within this context. However, with modifications, these conditions could be generalized to align with the deterministic metric space framework of Miller (2021). Furthermore, our analysis raises interesting questions about alternative definitions of posterior consistency, such as allowing posterior mass to converge on a set rather than a single point. Such a perspective could allow for the development of limit theorems for information criteria in non-identifiable models. We aim to extend our proofs to these settings in future work.

Appendix

A Geometric model limits

We will verify the sufficient conditions of Corollary 1 and Proposition 3 with respect to the geometric model from Section 3.1. To this end, we first recall that the geometric model has PMF

$$p(x|\theta) = (1-\theta)^x \theta$$

for $x \in \mathbb{X} = \mathbb{N} \cup \{0\}$ and $\theta \in \mathbb{T} = (0, 1)$, and we further equip \mathbb{T} with the uniform prior measure Π , characterized by density $\pi(\theta) = 1$, for each $\theta \in \mathbb{T}$. Taking the logarithm yields the log-likelihood

$$\log p(x|\theta) = X \log (1-\theta) + \log \theta$$

which is continuous in θ , for each $X \in \mathbb{X}$, and is thus Caratheodory, verifying A1. To verify A2a, we simply note that

$$E\left[\log p\left(X|\theta\right)\right] = EX\log\left(1-\theta\right) + \log\theta$$

is finite for each fixed $\theta \in (0, 1)$ as long as $EX < \infty$.

Since each compact set of \mathbb{T} is a subset of $\mathbb{A} = [a, 1 - a]$ for some $a \in (0, 1/2)$, we need only verify A2b on \mathbb{A} . Since $\log(1 - \theta)$ and $\log \theta$ are both continuous on \mathbb{A} , the Weierstrass extreme value theorem implies that

$$\left|\log p\left(X|\theta\right)\right| \le X \sup_{\theta \in \mathbb{A}} \left|\log \left(1-\theta\right)\right| + \sup_{\theta \in \mathbb{A}} \left|\log \theta\right| = \Delta\left(X\right),$$

where $E[\Delta(X)] < \infty$ whenever $EX < \infty$, which implies A2b and thus A2.

To show A3, we simply observe that

$$E\left[\log p\left(X|\theta\right)\right] = EX\log\left(1-\theta\right) + \log\theta$$
(28)

is maximized at $\theta_0 = 1/\{1 + EX\}$, by Fermat's first order condition and the fact that (28) is strictly concave. Finally, the concavity of $\theta \mapsto \log p(x|\theta)$, for each $x \in \mathbb{X}$, implies quasiconcavity, which verifies A4 via B4a. Since $\Pi ((\theta_0 - \rho, \theta_0 + \rho) > 0)$ for each $\rho > 0$, as long as EX > 0, Proposition 1 implies that the posterior sequence $(\Pi_n^{\beta_n})_{n \in \mathbb{N}}$ converges to δ_{θ_0} , P-a.s.w., whenever $n\beta_n \to \infty$. In fact, for each $n \in \mathbb{N}$, we observe that the density of $\Pi_n^{\beta_n}$ is

$$\pi_n^{\beta_n}(\theta) \propto \exp\left\{n\beta_n \bar{X}_n \log\left(1-\theta\right) + n\beta_n \log\theta\right\}.$$

We can thus identify $\prod_{n=1}^{\beta_n} \hat{X}_n$ with the law of the beta distribution Beta (a_n, b_n) , where $a_n = n\beta_n + 1$ and $b_n = n\beta_n \bar{X}_n + 1$.

To proceed, we are required to show that

$$\frac{1}{n}\sum_{i=1}^{n}\log p(X_i|\theta) = \bar{X}_n \log (1-\theta) + \log \theta$$

is AUI. To this end, we shall define

$$g_n(\theta) = \bar{X}_n \left| \log \left(1 - \theta \right) \right| + \left| \log \theta \right|,$$

and observe that

$$\left|\bar{X}_n \log\left(1-\theta\right) + \log\theta\right| \le g_n\left(\theta\right).$$

Since $g_n: \mathbb{T} \to \mathbb{R}_{\geq 0}$ is continuous, Theorem 3 implies P-a.s., g_n continuously converges to

$$g(\theta) = \mathbf{E}X \left| \log \left(1 - \theta \right) \right| + \left| \log \theta \right|$$

by definition. To verify condition (10), notice that

$$\int_{\mathbb{T}} \left| \log \left(1 - \theta \right) \right| \Pi_n^{\beta_n} \left(\mathrm{d}\theta \right) = -\psi \left(b_n \right) + \psi \left(a_n + b_n \right)$$

and

$$\int_{\mathbb{T}} \left| \log \theta \right| \Pi_n^{\beta_n} \left(\mathrm{d} \theta \right) = -\psi \left(a_n \right) + \psi \left(a_n + b_n \right),$$

where $\psi : \mathbb{R}_{>0} \to \mathbb{R}$ is the digamma function (cf. Olver et al., 2010, Eq. 5.2.2), defined as:

$$\psi(a) = \frac{\left(\mathrm{d}\Gamma(z)/\mathrm{d}z\right)(a)}{\Gamma(a)}$$

Here, for each z > 0,

$$\Gamma(z) = \int_0^\infty \exp\left\{-t\right\} t^{z-1} \mathrm{d}t$$

is the Gamma function.

Next, from Alzer (1997, Eq. (2.2)), we have the inequalities:

$$\log a - \frac{1}{a} \le \psi(a) \le \log a - \frac{1}{2a},$$

for each a > 0, which implies

$$\log\left(\frac{a_n+b_n}{a_n}\right)+o_{\text{a.s.}}\left(1\right)\leq -\psi\left(a_n\right)+\psi\left(a_n+b_n\right)\leq \log\left(\frac{a_n+b_n}{a_n}\right)+o_{\text{a.s.}}\left(1\right).$$

But since

$$\frac{a_n + b_n}{a_n} = \frac{n\beta_n + n\beta_n X_n + 2}{n\beta_n + 1} \xrightarrow[n \to \infty]{\text{P-a.s.}} 1 + \text{E}X,$$

which implies

$$\int_{\mathbb{T}} \left| \log \theta \right| \prod_{n=0}^{\beta_n} (\mathrm{d}\theta) \xrightarrow[n \to \infty]{\text{P-a.s.}} \log \left\{ 1 + \mathrm{E}X \right\}$$

by the squeeze theorem. In the same way, we have

$$\int_{\mathbb{T}} \left| \log \left(1 - \theta \right) \right| \prod_{n=1}^{\beta_n} \left(\mathrm{d}\theta \right) \xrightarrow[n \to \infty]{\text{P-a.s.}} \log \left\{ \frac{1 + \mathrm{E}X}{\mathrm{E}X} \right\}$$

and consequently,

$$\int_{\mathbb{T}} g_n\left(\theta\right) \Pi_n^{\beta_n}\left(\mathrm{d}\theta\right) \xrightarrow[n \to \infty]{\text{P-a.s.}} \mathrm{E}X \log\left\{\frac{1 + \mathrm{E}X}{\mathrm{E}X}\right\} + \log\left\{1 + \mathrm{E}X\right\} = g\left(\theta_0\right),$$

as required. We have thus verified the conditions of Corollary 1 and thus

$$BPIC_n, WBIC_n \xrightarrow[n \to \infty]{P-a.s.} -2E \left[\log p\left(X | \theta_0 \right) \right] = 2 \log \left\{ 1 + EX \right\} - 2EX \log \left\{ \frac{EX}{1 + EX} \right\}.$$

Finally, to verify that DIC_n has the same limit, we need to demonstrate that $\bar{\theta}_n \xrightarrow[n \to \infty]{n \to \infty} \theta_0$. To this end, we simply observe that $|\theta| \leq 1$ for each $\theta \in \mathbb{T}$, and is hence AUI with respect to $\left(\Pi_n^{\beta_n}\right)_{n \in \mathbb{N}}$, and the result holds by application of Proposition 4. Thus, by Proposition 3,

$$DIC_n \xrightarrow[n \to \infty]{P-a.s.} 2 \log \{1 + EX\} - 2EX \log \left\{\frac{EX}{1 + EX}\right\}$$

B Normal model limits

We now verify the conditions of Corollary 1 and Proposition 3 for the normal model in Section 3.2. We start by recalling that the normal model has PDF:

for $x \in \mathbb{X} = \mathbb{R}^p$ and $\theta \in \mathbb{T} = \mathbb{R}^p$, which we equip with the normal prior measure Π , whose PDF has the form

$$\pi(\theta) = (2\pi)^{-p/2} \exp\left\{-\frac{1}{2} \|\theta - \mu\|^2\right\},\$$

for some hyperparameter $\mu \in \mathbb{R}^p$, we observe that the log-likelihood

$$\log p(X|\theta) = -\frac{p}{2}\log(2\pi) - \frac{1}{2} ||X - \theta||^2$$

is Caratheodory, verifying A1. By further expansion, we have

$$\log p(X|\theta) = -\frac{p}{2}\log(2\pi) - \frac{1}{2}\left\{ \|X\|^2 - 2X^{\top}\theta + \|\theta\|^2 \right\},\$$

Thus, for each $\theta \in \mathbb{T}$, $E[\log p(X|\theta)]$ is finite as long as $E[||X||^2] < \infty$, thus A2a is verified.

Note that every compact subset of \mathbb{T} is contained in a closed ball $\overline{\mathbb{B}}_{\rho}(0)$, for some $\rho > 0$. Further note that $\theta \mapsto \log p(X|\theta)$ is continuous for each $X \in \mathbb{X}$, since $X^{\top}\theta$ and $\|\theta\|^2$ are continuous in θ . Thus, the Weierstrass extreme value theorem implies that

$$\log p(X|\theta) \le \frac{p}{2} \log (2\pi) + \|X\|^2 + 2|X|^{\top} \sup_{\theta \in \bar{\mathbb{B}}_{\rho}(0)} |\theta| + \sup_{\theta \in \bar{\mathbb{B}}_{\rho}(0)} \|\theta\|^2 = \Delta(X),$$

where $|\cdot|$, when applied to a vector, denotes the elementwise absolute value function. Observe that $E[\Delta(X)] < \infty$ whenever $E[||X||^2] < \infty$, and thus A2b is verified. Therefore A2 holds.

To verify A3, we observe that the derivative of

$$E[\log p(X|\theta)] = -\frac{p}{2}\log(2\pi) - \frac{1}{2}\left\{E\left[\|X\|^{2}\right] - 2E[X]^{\top}\theta + \|\theta\|^{2}\right\}$$

is $2\theta - 2EX$, which yields the maximum $\theta_0 = EX$, by Fermat's condition and by the fact that $E[\log p(X|\theta)]$ is strictly concave. Concavity implies quasiconcavity and thus A4 is also verified via B4a. Since Π is normal, it has positive mass on every open subset of \mathbb{R}^p , thus we can apply Proposition 1 to guarantee that $(\Pi_n^{\beta_n})_{n\in\mathbb{N}}$ converges to δ_{θ_0} , P-a.s., whenever $n\beta_n \to \infty$.

Like the geometric model, conjugacy implies that $\Pi_n^{\beta_n}$ has density

$$\pi_n^{\beta_n} \propto \exp\left\{-\frac{1}{2(n\beta_n+1)^{-1}} \left\|\theta - \frac{\beta_n \sum_{i=1}^n X_i + \mu}{n\beta_n+1}\right\|^2\right\},\$$

which implies that $\Pi_n^{\beta_n}$ is distributed as per the normal law N (μ_n, Σ_n) , where

$$\mu_n = \frac{n\beta_n \bar{X}_n + \mu}{n\beta_n + 1},$$

and $\Sigma_n = (n\beta_n + 1)^{-1} \mathbf{I}$, with \mathbf{I} denoting the identity matrix. Using this expression, we can now verify that

$$\frac{1}{n} \sum_{i=1}^{n} \log p(X_i | \theta) = -\frac{p}{2} \log (2\pi) - \frac{1}{2n} \sum_{i=1}^{n} \|X_i - \theta\|^2$$

is AUI. To do so, we bound it above by

$$g_n(\theta) = \frac{p}{2} \log (2\pi) + \frac{1}{2n} \sum_{i=1}^n ||X_i - \theta||^2$$

= $\frac{p}{2} \log (2\pi) + \frac{1}{2n} \left\{ \sum_{i=1}^n ||X_i||^2 - 2n \bar{X}_n^\top \theta + n ||\theta||^2 \right\}.$

Clearly, we have

$$\left| -\frac{p}{2} \log (2\pi) - \frac{1}{2n} \sum_{i=1}^{n} \|X_i - \theta\|^2 \right| \le g_n(\theta).$$

Given that $\mathbf{E}\left[\|X\|^2\right] < \infty$, by the strong law of large numbers,

$$\frac{1}{n} \sum_{i=1}^{n} \|X_i\|^2 \xrightarrow[n \to \infty]{\text{P-a.s.}} \mathbf{E}\left[\|X\|^2\right]$$

and $\bar{X}_n \xrightarrow[n \to \infty]{\text{P-a.s.}} \text{E}X$. Since g_n is continuous, we have continuous convergence of g_n to

$$g(\theta) = \frac{p}{2}\log(2\pi) + \frac{1}{2}\left\{ E\left[\|X\|^2 \right] - 2(EX)^\top \theta + \|\theta\|^2 \right\},\$$

P-a.s., by Theorem 3. Now, to verify (10), we write

$$\int_{\mathbb{R}^{p}} \|X - \theta\|^{2} \Pi_{n}^{\beta_{n}} (\mathrm{d}\theta) = \mathrm{tr} \left[(n\beta_{n} + 1)^{-1} \mathbf{I} \right] + \left\{ \frac{n\beta_{n}\bar{X}_{n} + \mu}{n\beta_{n} + 1} - X \right\}^{\top} \left\{ \frac{n\beta_{n}\bar{X}_{n} + \mu}{n\beta_{n} + 1} - X \right\}$$
$$= \frac{p}{n\beta_{n} + 1} + \left\| \frac{n\beta_{n}\bar{X}_{n} + \mu}{n\beta_{n} + 1} \right\|^{2} - 2X^{\top} \frac{n\beta_{n}\bar{X}_{n} + \mu}{n\beta_{n} + 1} + \|X\|^{2}.$$

Thus, we have

$$\frac{1}{2n} \sum_{i=1}^{n} \int_{\mathbb{R}^{p}} \|X_{i} - \theta\|^{2} \Pi_{n}^{\beta_{n}} (\mathrm{d}\theta) = \frac{1}{2} \left\{ \frac{p}{n\beta_{n} + 1} + \left\| \frac{n\beta_{n}\bar{X}_{n} + \mu}{n\beta_{n} + 1} \right\|^{2} - 2\bar{X}_{n}^{\top} \frac{n\beta_{n}\bar{X}_{n} + \mu}{n\beta_{n} + 1} + \frac{1}{n} \sum_{i=1}^{n} \|X_{i}\|^{2} \right\}$$
$$\xrightarrow{\text{P-a.s.}}_{n \to \infty} \frac{1}{2} \left\{ \mathrm{E} \left[\|X\|^{2} \right] - \|\mathrm{E}X\|^{2} \right\}.$$

Our verification is complete by noting that

$$\int_{\mathbb{T}} g\left(\theta\right) \mathrm{d}\delta_{\theta_{0}} = \frac{p}{2} \log\left(2\pi\right) + \frac{1}{2} \left\{ \mathrm{E}\left[\left\|X\right\|^{2}\right] - \left\|\mathrm{E}X\right\|^{2} \right\},\$$

upon noting that $\theta_0 = EX$. We have verified the necessary conditions to apply Corollary 1, which implies the desired result that

$$BPIC_n, WBIC_n \xrightarrow[n \to \infty]{P-a.s.} -2E \left[\log p\left(X | \theta_0 \right) \right] = -\frac{p}{2} \log \left(2\pi \right) - \frac{1}{n} \left\{ E \left[\left\| X \right\|^2 \right] - \left\| EX \right\|^2 \right\}$$

To obtain the limit for DIC_n , we must verify that $\|\cdot\| : \mathbb{T} \to \mathbb{R}_{\geq 0}$ is AUI with respect to $(\Pi_n^{\beta_n})_{n \in \mathbb{N}}$. To this end, we use the fact that $\sqrt{a} \leq x/2 + 1/2$, implying that

$$\|\theta\| \le \frac{1}{2} \|\theta\|^2 + \frac{1}{2}$$

Using the bound, we again verify condition (10) by noting that

$$\int_{\mathbb{T}} \|\theta\|^2 \Pi_n^{\beta_n} (\mathrm{d}\theta) = \mathrm{tr} \left[(n\beta_n + 1)^{-1} \mathbf{I} \right] + \left\| \frac{n\beta_n \bar{X}_n + \mu}{n\beta_n + 1} \right\|^2,$$

which converges, P-a.s., to $||EX||^2$. This is as required, since

$$\int_{\mathbb{T}} \|\theta\|^2 \,\mathrm{d}\delta_{\theta_0} = \|\theta_0\|^2 = \|\mathbf{E}X\|^2$$

We have therefore verified the sufficient conditions for Proposition 3. Thus

$$\operatorname{DIC}_{n} \xrightarrow[n \to \infty]{\text{P-a.s.}} -\frac{p}{2} \log \left(2\pi\right) - \frac{1}{n} \left\{ \operatorname{E}\left[\left\|X\right\|^{2}\right] - \left\|\operatorname{E}X\right\|^{2} \right\}.$$

C Laplace model limits

We now examine our final technical example from Section 3.3: the Laplace model. Recall that the Laplace distribution has PDF:

$$p(x|\theta) = \frac{1}{2\gamma} \exp\left\{-\frac{|x-\mu|}{\gamma}\right\},$$

for $x \in \mathbb{X} = \mathbb{R}$, where we consider only a compact support $\mathbb{T} = [-m, m] \times [s^{-1}, s]$ for $\theta = (\mu, \gamma) \in \mathbb{T}$, where m > 0 and s > 1. Since \mathbb{T} is compact, the analysis is the same for any prior distribution Π whose density π is strictly positive on \mathbb{T} .

Writing the log-likelihood as

$$\log p(X|\theta) = -\log(2\gamma) - \frac{1}{\gamma} |X - \mu|,$$

we observe that $\theta \mapsto \log p(X|\theta)$ is is continuous for each fixed $X \in \mathbb{X}$, thus it is Caratheodory and satisfies

A1. We verify A2a by first noting that by the strong law of large numbers

$$\frac{1}{n} \sum_{i=1}^{n} \log p(X_i | \theta) = -\log(2\gamma) - \frac{1}{\gamma n} \sum_{i=1}^{n} |X_i - \mu|$$

converges, P-a.s., to

$$E\left[\log p\left(X|\theta\right)\right] = -\log\left(2\gamma\right) - \frac{1}{\gamma}E\left[|X - \mu|\right],$$

as long as $E|X| < \infty$, for each fixed $\theta \in \mathbb{T}$. A2b is verified by noting that

$$\left|\log p\left(X|\theta\right)\right| \le \sup_{\gamma \in [s^{-1},s]} \left|\log\left(2\gamma\right)\right| + \left|X\right| \sup_{\gamma \in [s^{-1},s]} \frac{1}{\gamma} + \sup_{\theta \in \mathbb{T}} \frac{|\mu|}{\gamma} = \Delta\left(X\right),$$

where $E[\Delta(X)] < \infty$ if $E|X| < \infty$, by the Weierstrass extreme value theorem. Thus A2 holds. For A3, we observe that

$$\mathbf{E}\left[\log \mathbf{p}\left(X|\theta\right)\right] = -\log\left(2\gamma\right) - \frac{1}{\gamma}\mathbf{E}\left[|X - \mu|\right]$$

has a unique maximizer $\mu_0 = \text{Med}(X)$ and $\gamma_0 = \mathbb{E}|X - \mu_0|$, whenever the law of X has a continuous distribution function and whenever $\theta_0 = (\mu_0, \gamma_0) \in \mathbb{T}$. The optimality of μ_0 for any fixed γ can be proved, for example, via (Schay Thm. 6.6.3). The optimality of γ_0 given $\mu = \mu_0$ is then a consequence of Fermat's first order condition. A4 is verified by the continuity of the log-likelihood and the compactness of T. Together with the strict positivity of π implying that $\Pi(\mathbb{B}_{\rho}(\theta_0)) > 0$ for each $\rho > 0$, we have verified the sufficient conditions for Proposition 1 and can conclude that whenever $n\beta_n \to \infty$, $(\Pi_n^{\beta_n})_{n\in\mathbb{N}}$ converges to δ_{θ_0} , P-a.s..

The compactness of \mathbb{T} also makes the verification of the AUI condition simple. Observe that

$$\left|\frac{1}{n}\sum_{i=1}^{n}\log p\left(X_{i}|\theta\right)\right| \leq g_{n}\left(\theta\right) = \sup_{\gamma \in [s^{-1},s]}\left|\log\left(2\gamma\right)\right| + \sup_{\gamma \in [s^{-1},s]}\frac{1}{\gamma} \times \frac{1}{n}\sum_{i=1}^{n}\left|X_{i}\right| + \sup_{\theta \in \mathbb{T}}\frac{|\mu|}{\gamma},$$

where $g_n(\theta)$ is a constant function with respect to θ for each fixed $\omega \in \Omega$. Thus, the strong law of large numbers implies that $(g_n)_{n \in \mathbb{N}}$ converges continuously, P-a.s., to

$$g\left(\theta\right) = \sup_{\gamma \in [s^{-1},s]} \left|\log\left(2\gamma\right)\right| + \sup_{\gamma \in [s^{-1},s]} \frac{1}{\gamma} \mathbf{E}\left|X\right| + \sup_{\theta \in \mathbb{T}} \frac{|\mu|}{\gamma},$$

and thus $(g_n)_{n\in\mathbb{N}}$ is AUI with respect to $\left(\prod_{n=1}^{\beta_n}\right)_{n\in\mathbb{N}}$, since

$$\int_{\mathbb{T}} g_n \mathrm{d}\Pi_n^{\beta_n} = g_n \stackrel{\text{P-a.s.c}}{\underset{n \to \infty}{\longrightarrow}} g = \int_{\mathbb{T}} g \mathrm{d}\delta_{\theta_0}.$$

Since

$$\int_{\mathbb{T}} \frac{1}{n} \sum_{i=1}^{n} \log p(X_i | \theta) \prod_{n=1}^{\beta_n} (\mathrm{d}\theta) \xrightarrow[n \to \infty]{\text{P-a.s.}} \mathbb{E} \left[\log p(X | \theta_0) \right],$$

where

$$\operatorname{E}\left[\log p\left(X|\theta_{0}\right)\right] = -\log\left(2\gamma_{0}\right) - 1,$$

Corollary 1 implies that

$$BPIC_n, WBIC_n \xrightarrow[n \to \infty]{\text{P-a.s.}} -2E \left[\log p\left(X | \theta_0 \right) \right] = 2 \log \left(2\gamma_0 \right) + 2$$

as required. The same compactness and continuity argument verifies that $\|\cdot\|$ is AUI with respect to $\left(\prod_{n=1}^{\beta_n}\right)_{n\in\mathbb{N}}$, P-a.s., thus DIC_n has the same limit as above, by Proposition 3.

D Approximation of DIC_n for the geometric model

Under the geometric model, we recall that the PMF of X is

$$p(x|\theta) = (1-\theta)^x \theta,$$

where we now endow θ with a prior Π whose law is Beta (α, β) , $\alpha, \beta > 0$. We further note that the average log-likelihood has the form

$$\frac{1}{n}\sum_{i=1}^{n}\log p\left(X_{i}|\theta\right) = \bar{X}_{n}\log\left(1-\theta\right) + \log\theta.$$

To evaluate DIC_n , we must evaluate the two expressions

$$\int_0^1 \frac{1}{n} \sum_{i=1}^n \log p(X_i | \theta) \Pi_n(\mathrm{d}\theta) \text{ and } \frac{1}{n} \sum_{i=1}^n \log p(X_i | \bar{\theta}_n)$$

where $\bar{\theta}_n$ is the expectation of the law Π_n . Both of these evaluations can be facilitated by noting that the conjugacy between the geometric and beta distribution implies that Π_n has law Beta (a_n, b_n) , where $a_n = n + \alpha$ and $b_n = n\bar{X}_n + \beta$. Furthermore, if $Y \sim \text{Beta}(a_n, b_n)$, then we have the useful expression

$$EY = \frac{a_n}{a_n + b_n}, E \log (Y) = \psi (a_n) - \psi (a_n + b_n), \text{ and } E \log (1 - Y) = \psi (b_n) - \psi (a_n + b_n).$$

Given these facts,

$$\int_{0}^{1} \frac{1}{n} \sum_{i=1}^{n} \log p(X_{i}|\theta) \Pi_{n}(d\theta) = E\left[\bar{X}_{n} \log(1-Y) + \log Y|\bar{X}_{n}\right]$$
$$= \bar{X}_{n} \left\{\psi(b_{n}) - \psi(a_{n}+b_{n})\right\} + \psi(a_{n}) - \psi(a_{n}+b_{n}).$$

We then apply the Poincare-type approximation

$$\psi(a) \approx \log a - \frac{1}{2a}$$

to get the approximation

$$\bar{X}_n\left\{\log\left(\frac{b_n}{a_n+b_n}\right) - \frac{a_n}{2b_n\left(a_n+b_n\right)}\right\} + \left\{\log\left(\frac{a_n}{a_n+b_n}\right) - \frac{b_n}{2a_n\left(a_n+b_n\right)}\right\}.$$

By substitution of $\bar{\theta}_n = EY$, we also have

$$\frac{1}{n}\sum_{i=1}^{n}\log p\left(X_{i}|\bar{\theta}_{n}\right) = \bar{X}_{n}\log\left(\frac{b_{n}}{a_{n}+b_{n}}\right) + \log\left(\frac{a_{n}}{a_{n}+b_{n}}\right).$$

Our approximation is then obtained via the definition of DIC_n .

References

- Akaike, H. (1974). A new look at the statistical model identification. IEEE Transactions on Automatic Control, 19, 716–723. (Cited on page 2.)
- Alzer, H. (1997). On some inequalities for the gamma and psi functions. Mathematics of computation, 66(217), 373–389. (Cited on page 26.)
- Ando, T. (2007). Bayesian predictive information criterion for the evaluation of hierarchical bayesian and empirical bayes models. *Biometrika*, 94(2), 443–458. (Cited on page 2.)
- Ando, T. (2010). Bayesian Model Selection and Statistical Modeling. Boca Raton: CRC Press. (Cited on pages 2 and 5.)
- Bain, A. & Crisan, D. (2009). Fundamentals of Stochastic Filtering. New York: Springer. (Cited on page 4.)
- Baudry, J.-P. (2015). Estimation and model selection for model-based clustering with the conditional classification likelihood. *Electronic Journal of Statistics*, 9, 1041–1077. (Cited on page 2.)
- Beckenstein, E., Narici, L., & Suffel, C. (2011). Topological Algebras. Elsevier. (Cited on page 6.)
- Beer, G. (1993). Topologies on closed and closed convex sets, volume 268. Springer Science & Business Media. (Cited on page 16.)
- Berger, J. O. (1985). Statistical decision theory and Bayesian analysis. Springer Science & Business Media. (Cited on page 4.)
- Bernardo, J. M. & Smith, A. F. (2009). Bayesian theory. John Wiley & Sons. (Cited on page 4.)
- Bernton, E., Jacob, P. E., Gerber, M., & Robert, C. P. (2019). Approximate Bayesian computation with the Wasserstein distance. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 81(2), 235–269. (Cited on page 4.)
- Berti, P., Pratelli, L., & Rigo, P. (2006). Almost sure weak convergence of random probability measures. Stochastics and Stochastics Reports, 78(2), 91–97. (Cited on pages 2 and 4.)

- Bissiri, P. G., Holmes, C. C., & Walker, S. G. (2016). A general framework for updating belief distributions. Journal of the Royal Statistical Society B, 78, 1103–1130. (Cited on pages 4 and 15.)
- Burnham, K. P. & Anderson, D. R. (2002). Model Selection And Multimodel Inference: A Practical Information-theoretic Approach. New York: Springer. (Cited on page 2.)
- Cambini, A. & Martein, L. (2008). *Generalized Convexity and Optimization: Theory and Applications*. Springer. (Cited on page 20.)
- Claeskens, G. & Hjort, N. L. (2008). Model Selection And Model Averaging. Cambridge: Cambridge University Press. (Cited on page 23.)
- Feinberg, E. A., Kasyanov, P. O., & Liang, Y. (2020a). Fatou's lemma for weakly converging measures under the uniform integrability condition. *Theory of Probability and Its Applications*, 64, 615–630. (Cited on pages 2, 7, and 8.)
- Feinberg, E. A., Kasyanov, P. O., & Liang, Y. (2020b). Fatou's lemma in its classical form and Lebesgue's convergence theorems for varying measures with applications to Markov decision processes. *Theory of Probability and Its Applications*, 65, 270–291. (Cited on pages 2 and 7.)
- Florenzano, M. & Le Van, C. (2001). Finite Dimensional Convexity and Optimization. Springer. (Cited on page 20.)
- Forbes, F., Nguyen, H. D., Nguyen, T., & Arbel, J. (2022). Summary statistics and discrepancy measures for approximate Bayesian computation via surrogate posteriors. *Statistics and Computing*, 32(5), 85. (Cited on page 4.)
- Ghosal, S. & Van der Vaart, A. (2017). Fundamentals of nonparametric Bayesian inference, volume 44. Cambridge University Press. (Cited on pages 2, 4, and 7.)
- Ghosh, J. K., Delampady, M., & Samanta, T. (2006). An Introduction to Bayesian Analysis: Theory and Methods. New York: Springer. (Cited on pages 2 and 4.)
- Ghosh, J. K. & Ramamoorthi, R. V. (2003). Bayesian Nonparametrics. New York: Springer. (Cited on pages 2 and 19.)
- Grübel, R. & Kabluchko, Z. (2016). A functional central limit theorem for branching random walks, almost sure weak convergence and applications to random trees. *The Annals of Applied Probability*, 26(6), 3659–3698. (Cited on pages 2, 3, and 4.)
- Hannig, J., Iyer, H., Lai, R. C., & Lee, T. C. (2016). Generalized fiducial inference: A review and new results. *Journal of the American Statistical Association*, 111(515), 1346–1361. (Cited on page 4.)
- Knoblauch, J. (2019). Frequentist consistency of generalized variational inference. arXiv. (Cited on page 4.)

- Knoblauch, J., Jewson, J., & Damoulas, T. (2022). An optimization-centric view on Bayes' rule: reviewing and generalizing variational inference. *Journal of Machine Learning Reseach*, 23, 1–109. (Cited on pages 4 and 15.)
- Koltchinskii, V. (2011). Oracle Inequalities in Empirical Risk Minimization and Sparse Recovery Problems. Springer. (Cited on page 23.)
- Konishi, S. & Kitagawa, G. (2008). Information Criteria and Statistical Modeling. New York: Springer. (Cited on page 2.)
- Lehmann, E. L. & Casella, G. (1998). Theory Of Point Estimation. New York: Springer. (Cited on page 4.)
- Massart, P. (2007). Concentration Inequalities and Model Selection. New York: Springer. (Cited on page 23.)
- Miller, J. W. (2021). Asymptotic normality, concentration, and coverage of generalized posteriors. The Journal of Machine Learning Research, 22(1), 7598–7650. (Cited on pages 15, 19, and 24.)
- Nguyen, H. D. (2024). Panic: Consistent information criteria for general model selection problems. Australian & New Zealand Journal of Statistics. (Cited on page 23.)
- Nguyen, H. D., Arbel, J., Lü, H., & Forbes, F. (2020). Approximate Bayesian computation via the energy statistic. *IEEE Access*, 8, 131683–131698. (Cited on page 4.)
- Nguyen, T., Forbes, F., Arbel, J., & Duy Nguyen, H. (2024). Bayesian nonparametric mixture of experts for inverse problems. *Journal of Nonparametric Statistics*, (pp. 1–60). (Cited on page 4.)
- Olver, F. W. J., Lozier, D. W., Boisvert, R. F., & Clark, C. W., Eds. (2010). NIST Handbook of Mathematical Functions. Cambridge: Cambridge University Press. (Cited on pages 12 and 26.)
- Remmert, R. (1991). Theory of Complex Functions. New York: Springer. (Cited on pages 8 and 16.)
- Robert, C. P. (2007). The Bayesian choice: from decision-theoretic foundations to computational implementation. New York: Springer. (Cited on page 2.)
- Schwarz, G. (1978). Estimating the dimensions of a model. Annals of Statistics, 6, 461–464. (Cited on page 2.)
- Schweder, T. & Hjort, N. L. (2016). Confidence, likelihood, probability, volume 41. Cambridge University Press. (Cited on page 4.)
- Serfozo, R. (1982). Convergence of Lebesgue integrals with varying measures. Sankhya A, 44, 380–402. (Cited on pages 2 and 7.)
- Shapiro, A., Dentcheva, D., & Ruszczynski, A. (2021). Lectures on Stochastic Programming. SIAM. (Cited on page 21.)

- Sin, C.-Y. & White, H. (1996). Information criteria for selecting possibly misspecified parametric models. Journal of Econometrics, 71, 207–225. (Cited on pages 2 and 23.)
- Spiegelhalter, D. J., Best, N. G., Carlin, B. P., & Van Der Linde, A. (2002). Bayesian measures of model complexity and fit. Journal of the royal statistical society: Series b (statistical methodology), 64(4), 583-639. (Cited on pages 2 and 5.)
- Sweeting, T. (1989). On conditional weak convergence. *Journal of Theoretical Probability*, 2, 461–474. (Cited on pages 2 and 4.)
- van der Vaart, A. W. (1998). Asymptotic Statistics. Cambridge: Cambridge University Press. (Cited on page 4.)
- van der Vaart, A. W. & Wellner, J. A. (2023). Weak convergence and empirical processes. Cham: Springer. (Cited on pages 2 and 8.)
- Watanabe, S. (2013). A widely applicable Bayesian information criterion. Journal of Machine Learning Research, 14, 867–897. (Cited on pages 2, 5, and 12.)
- Watanabe, S. (2018). *Mathematical theory of Bayesian statistics*. Boca Raton: CRC Press. (Cited on page 2.)
- Westerhout, J., Nguyen, T., Guo, X., & Nguyen, H. D. (2024). On the asymptotic distribution of the minimum empirical risk. In Forty-first International Conference on Machine Learning. (Cited on page 23.)
- You, C., Ormerod, J. T., & Mueller, S. (2014). On variational bayes estimation and variational information criteria for linear regression models. Australian & New Zealand Journal of Statistics, 56, 73–87. (Cited on page 24.)
- Zhang, F. & Gao, C. (2020). Convergence rates of variational posterior distributions. The Annals of Statistics, 48(4), 2180 – 2207. Publisher: Institute of Mathematical Statistics. (Cited on page 4.)
- Zhang, T. (2006). Information-theoretic upper and lower bounds for statistical estimation. *IEEE Transactions on Information Theory*, 52, 1307–1321. (Cited on pages 4 and 15.)