Quasi maximum likelihood estimation of high-dimensional approximate dynamic matrix factor models via the EM algorithm

Matteo Barigozzi * Luca Trapin [†]

February 7, 2025

Abstract

This paper considers an approximate dynamic matrix factor model that accounts for the time series nature of the data by explicitly modelling the time evolution of the factors. We study Quasi Maximum Likelihood estimation of the model parameters based on the Expectation Maximization (EM) algorithm, implemented jointly with the Kalman smoother which gives estimates of the factors. This approach allows to easily handle arbitrary patterns of missing data. We establish the consistency of the estimated loadings and factor matrices as the sample size T and the matrix dimensions p_1 and p_2 diverge to infinity. The finite sample properties of the estimators are assessed through a large simulation study and an application to a financial dataset of volatility proxies.

Keywords: Matrix Factor Models; Expectation-Maximization Algorithm; Kalman Smoother; Missing Observations.

^{*}Department of Economics, University of Bologna, Piazza Scaravilli 2, Bologna, Italy E-mail: matteo.barigozzi@unibo.it

[†]Department of Statistical Sciences, University of Bologna, Via Belle Arti 41, Bologna, Italy E-mail: luca.trapin@unibo.it

1 Introduction

Matrix-variate time series data are becoming increasingly popular in economics and finance. For example, forecasts of regional economic time series exploit matrices of observations on several variables from different regions (Chernis et al., 2020), investigating the dynamics of international trade flows require time series matrices of bilateral trade (Chen and Chen, 2022), measures of financial connectedness are extracted from a sequence of networks among financial assets represented through adjacency matrices (Billio et al., 2021). This has stimulated the development of high-dimensional methods to analyze matrix time series data, including matrix autoregressive models (Chen et al., 2021; Hsu et al., 2021; Billio et al., 2023), matrix panel regression models (Kapetanios et al., 2021), and matrix factor models (Wang et al., 2019; Yu et al., 2022; Chen and Fan, 2023; Xu et al., 2024; Yu et al., 2024).

This paper considers a matrix factor model for the $p_1 \times p_2$ zero-mean matrix-valued stationary process $\{\mathbf{Y}_t\}$, with latent factors generated by a Matrix Autoregressive (MAR) model of order P. Namely, for any $t \in \mathbb{Z}$, we consider the model given by

$$\mathbf{Y}_t = \mathbf{R} \mathbf{F}_t \mathbf{C}' + \mathbf{E}_t, \tag{1}$$

$$\mathbf{F}_{t} = \sum_{i=1}^{P} \mathbf{A}_{i} \mathbf{F}_{t-i} \mathbf{B}_{i}' + \mathbf{U}_{t}.$$
(2)

In (1), \mathbf{F}_t is a $k_1 \times k_2$ matrix of latent factors with $k_1, k_2 < \min(p_1, p_2)$, \mathbf{R} and \mathbf{C} are $p_1 \times k_1$ and $p_2 \times k_2$ matrices of unknown row and column loadings, \mathbf{E}_t is a $p_1 \times p_2$ matrix of idiosyncratic components with $p_1 \times p_1$ row covariance matrix \mathbf{H} and $p_2 \times p_2$ column covariance matrix \mathbf{K} . In (2), \mathbf{A}_i and $\mathbf{B}_i, i=1,\ldots,P$, are $k_1 \times k_1$ and $k_2 \times k_2$ matrices of autoregressive parameters, and \mathbf{U}_t is a $k_1 \times k_2$ matrix of innovations from a matrix-variate distribution with $k_1 \times k_1$ row covariance matrix \mathbf{P} and $k_2 \times k_2$ column covariance matrix \mathbf{Q} . The processes $\{\mathbf{F}_t\}$ and $\{\mathbf{E}_t\}$ are assumed to be uncorrelated (at all leads and lags).

In our setting the idiosyncratic components are allowed to be correlated both across rows and columns, i.e., in general **H** and **K** are allowed to be full matrices, and we say that the model is *approximate*. Furthermore, the model is *dynamic* since the factors are autocorrelated as specified by the MAR in (2), and, moreover, we also allow the idiosyncratic components to be autocorrelated, although no explicit model for their dynamics is introduced. Therefore, we call a model defined by (1)-(2) an approximate dynamic matrix factor model (DMFM). It combines the matrix factor model, as formulated by Yu et al. (2022) and Chen and Fan (2023), and the MAR proposed by Chen et al. (2021).

In this paper, we propose Quasi Maximum Likelihood (QML) estimation of the factor loading matrices and factor matrices of the DMFM, implemented via the Expectation Maximization (EM) algorithm jointly with the Kalman smoother. We prove consistency of the estimated loadings and factors as $\min(p_1, p_2, T) \rightarrow \infty$.

Similarly to the vector case, the DMFM can be identified only in the limit $p_1, p_2 \rightarrow \infty$ due to its approximate structure. That is to say that the numbers of factors k_1 and k_2 can be consistently estimated only in a high-dimensional setting. This is what allows one to disentangle the factor driven component from the idiosyncratic one. However, this forces us to work in a high-dimensional setting. This makes joint Maximum Likelihood estimation of (1)-(2) a hard if not unfeasible task due to the large number of parameters we need to estimate, which is $O((p_1^2 + p_2^2)T)$ (all autocovariances of the factors and idiosyncratic components), and due to the lack of a closed form solution.

The estimation approach we consider has two main features which allow us to solve both problems. First, it is based on a mis-specified likelihood where the idiosyncratic components are treated as if they were uncorrelated. This reduces the number of parameters to be estimated to $O(p_1+p_2)$. Second, it is an iterative approach where, in a first step, for given parameters we estimate the factors via the Kalman smoother, and, in a second step, for given factors we estimate all parameters by maximizing the expected conditional likelihood. This allows us to derive a closed form expression for all estimators.

Our approach is the generalization of the approach proposed by Doz et al. (2012) for the vector case. However, such generalization is non-trivial, indeed, in the present matrix time series setting, we need, at each iteration of the EM algorithm, to jointly estimate the two matrices of loadings, \mathbf{R} and \mathbf{C} , which depend on each other (the same goes for the row and column idiosyncratic covariances, \mathbf{H} and \mathbf{K} , the MAR coefficients, \mathbf{A} and \mathbf{B} , and the MAR innovation covariances, \mathbf{P} and \mathbf{Q}). In order to respect this bi-linear structure requires then to modify the algorithm accordingly and makes the derivation of the asymptotic properties more challenging.

We argue that accounting for factor dynamics via the Kalman smoother and considering Maximum Likelihood estimation is particularly convenient as it allows the user to impose *a priori* restrictions on the models' parameters and/or dynamics, construct counterfactual scenarios, conditional forecasts, obtain now-casts, and deal with missing values due to different sampling frequencies or plain unavailability of the data (see, e.g., the applications in Bańbura and Modugno, 2014; Bańbura et al., 2015, in the case of vector time series).

To show the potential of the the proposed approach we apply it to the analysis of a matrix times series containing various volatility proxies for many stocks. Since not all proxies are available for all stocks we show how to adapt the EM algorithm to deal with missing values and we then produce volatility forecasts for all stocks.

The rest of the paper is organized as follows: Section 2 discusses related works; Section 3 presents the QML estimator and the EM algorithm; Section 4 presents the assumptions and the consistency results; Section 5 explains how to extend the EM algorithm to handle missing data; Section 6 studies the finite sample properties of the EM algorithm through Monte Carlo simulations; Section 7 applies the EM algorithm to estimate the latent variance of financial assets. All notation, as well as relevant results on matrix operations, are given in Appendix A.

2 Related literature

We briefly review alternative estimation approaches of large matrix factor models, possibly augmented with a MAR for the factors.

A first set of works considers only estimation of the factor model in (1), thus without explicitly accounting for the factors' dynamics. Specifically, Wang et al. (2019) introduced the class of large matrix factor models under the assumption of serially uncorrelated idiosyncratic components, and propose to estimate the loadings by means of eigenvectors of a long-run covariance matrix (see also Chen et al., 2020). Successively, Yu et al. (2022) and Chen and Fan (2023) extended this approach to the case of possibly autocorrelated idiosyncratic components, and propose two different generalizations to the matrix setting of the Principal Component (PC) estimators typically used in the vector case (Bai, 2003). Both these work consider also methods for determining the number of factors (see also He et al., 2023 and Han et al., 2022, for alternative methods). Last, Yuan et al. (2023) and Xu et al. (2024) consider two different specifications of a matrix factor model and generalize to those settings the classical QML estimation approach, where only the loadings and the idiosyncratic variances are estimated and the idiosyncratic components are treated as if they were uncorrelated (Bai and Li, 2012, 2016).

To the best of our knowledge only Yu et al. (2024) consider a DMFM as specified by (1)-(2). However, our work differs in several aspects. First, we consider joint estimation of factors and parameters of the model, while they consider a two-step approach where first the loadings and the factors are estimated and then a MAR is estimated on the factors. Second, we allow the idiosyncratic components to be serially correlated, while they impose a different factor structure with time independent idiosyncratic components. Third, we derive the asymptotic properties of the factors estimated via the Kalman smoother, while they do not study the asymptotic properties of such estimator, although entertaining the possibility of retrieving the factors via filtering. As a last difference we stress that our estimation approach can be easily adapted to make predictions also in presence of arbitrary patterns of missing data.

Our work is also related to two other strands of the literature. First, the idea of considering a misspecified likelihood in factor analysis to make its maximization more treatable dates back to Tipping and Bishop (1999) who, in a vector context, treated the idiosyncratic components as i.i.d.. This idea

was then extended by Doz et al. (2012) and Bai and Li (2016) to the case of high-dimensional vectors of time series having serially and cross-sectionally correlated idiosyncratic components. In particular, while Bai and Li (2016) do not model the dynamics of the factors, Doz et al. (2012) explicitly model the factors as generated by a VAR.

Second, there exist many factor model approaches for handling missing values in high-dimensional vector time series. On the one hand, Bańbura and Modugno (2014) propose an EM-based approach which we generalize to the matrix setting in this paper. On the other hand, there are a few approaches based on various modifications of standard PC analysis, see, e.g., the recent works by Xiong and Pelger (2023) and Cahan et al. (2023). Finally, Cen and Lam (2024) consider a PC based approach for the tensor case, which includes the matrix case.

3 Estimation

3.1 The log-likelihood

Let consider a DMFM as defined in (1)-(2), and without loss of generality assume that the MAR is of order P=1. For any $t\in\mathbb{Z}$, our data generating process is then given by:

$$\mathbf{Y}_t = \mathbf{R}\mathbf{F}_t \mathbf{C}' + \mathbf{E}_t, \tag{3}$$

$$\mathbf{F}_t = \mathbf{A}\mathbf{F}_{t-1}\mathbf{B}' + \mathbf{U}_t. \tag{4}$$

Denote as $\mathbf{y}_t = \operatorname{vec}(\mathbf{Y}_t)$, $\mathbf{f}_t = \operatorname{vec}(\mathbf{F}_t)$, $\mathbf{e}_t = \operatorname{vec}(\mathbf{E}_t)$ and $\mathbf{u}_t = \operatorname{vec}(\mathbf{U}_t)$, the vectorized versions of matrices \mathbf{Y}_t , \mathbf{F}_t , \mathbf{E}_t and \mathbf{U}_t at time t. Consider a sample of T observations, and let $\mathbf{Y}_T = (\mathbf{y}'_1, \dots, \mathbf{y}'_T)'$ and $\mathbf{E}_T = (\mathbf{e}'_1, \dots, \mathbf{e}'_T)'$ be $(p_1 p_2 T)$ -dimensional vectors containing all the observations and idiosyncratic components, respectively, and let $\mathbf{F}_T = (\mathbf{f}'_1, \dots, \mathbf{f}'_T)'$ be the $(k_1 k_2 T)$ -dimensional vector of factors. Let $\mathbf{\Omega}_T^{\mathbf{Y}} = \mathbb{E}_{\boldsymbol{\theta}}[\mathbf{Y}_T \mathbf{Y}'_T]$, $\mathbf{\Omega}_T^{\mathbf{E}} = \mathbb{E}_{\boldsymbol{\theta}}[\mathbf{E}_T \mathbf{E}'_T]$, $\mathbf{\Omega}_T^{\mathbf{F}} = \mathbb{E}_{\boldsymbol{\theta}}[\mathbf{F}_T \mathbf{F}'_T]$ be covariance matrices containing all the cross-sectional row and column covariances and all the autocovariances up to lag (T-1). Notice that $\mathbf{\Omega}_T^{\mathbf{F}}$ is fully characterized by the matrices of MAR parameter \mathbf{A} , \mathbf{B} , \mathbf{P} and \mathbf{Q} , thus, hereafter, we denote it as $\mathbf{\Omega}_{T(\mathbf{A},\mathbf{B},\mathbf{P},\mathbf{Q})}^{\mathbf{F}}$.

It follows that the DMFM is fully characterized by the covariance matrix:

$$\boldsymbol{\Omega}_{T}^{\mathsf{Y}} = (\mathbb{I}_{T} \otimes \mathbf{C} \otimes \mathbf{R}) \boldsymbol{\Omega}_{T(\mathbf{A},\mathbf{B},\mathbf{P},\mathbf{Q})}^{\mathsf{F}} (\mathbb{I}_{T} \otimes \mathbf{C} \otimes \mathbf{R})' + \boldsymbol{\Omega}_{T}^{\mathsf{E}}.$$

In an approximate DMFM as the one we consider, Ω_T^{E} is allowed to be a full-matrix, but this implies that it has $\frac{p_1p_2T(p_1p_2T+1)}{2}$ entries to be estimated while we have only p_1p_2T observations. This makes Maximum Likelihood estimation unfeasible.

A solution consists in considering a misspecified likelihood where the idiosyncratic components \mathbf{E}_t are treated as if they were serially and cross-sectionally uncorrelated, i.e. when we replace Ω_T^{E} with $\mathbb{I}_T \otimes \mathrm{dg}(\mathbf{K}) \otimes \mathrm{dg}(\mathbf{H})$. This is the approach follwed in the vector factor model case, by, e.g., Bai and Li (2012, 2016) and Doz et al. (2012). Under this misspecification, the vector of parameters to be estimated reduces to

$$\boldsymbol{\theta} = \left(\operatorname{vec}(\mathbf{R})', \operatorname{vec}(\mathbf{C})', \operatorname{vec}(\operatorname{dg}(\mathbf{H}))', \operatorname{vec}(\operatorname{dg}(\mathbf{K}))', \operatorname{vec}(\mathbf{A})', \operatorname{vec}(\mathbf{B})', \operatorname{vec}(\mathbf{P})', \operatorname{vec}(\mathbf{Q})'\right)'.$$

which has dimension now growing as p_1+p_2 , thus it can be estimated using p_1p_2T observations. Consequently, we consider a misspecified, or quasi, log-likelihood given by:

$$\ell(\mathbf{Y}_{T};\boldsymbol{\theta}) = \frac{p_{1}p_{2}T}{2}\log(2\pi) - \log\left(\left|\left(\mathbb{I}_{T}\otimes\mathbf{C}\otimes\mathbf{R}\right)\boldsymbol{\Omega}_{T(\mathbf{A},\mathbf{B},\mathbf{P},\mathbf{Q})}^{\mathsf{F}}(\mathbb{I}_{T}\otimes\mathbf{C}\otimes\mathbf{R})' + \mathbb{I}_{T}\otimes\mathrm{dg}(\mathbf{K})\otimes\mathrm{dg}(\mathbf{H})\right|\right) - \frac{1}{2}\left[\mathbf{Y}_{T}'\left(\left(\mathbb{I}_{T}\otimes\mathbf{C}\otimes\mathbf{R}\right)\boldsymbol{\Omega}_{T(\mathbf{A},\mathbf{B},\mathbf{P},\mathbf{Q})}^{\mathsf{F}}(\mathbb{I}_{T}\otimes\mathbf{C}\otimes\mathbf{R})' + \mathbb{I}_{T}\otimes\mathrm{dg}(\mathbf{K})\otimes\mathrm{dg}(\mathbf{H})\right)^{-1}\mathbf{Y}_{T}\right].$$
(5)

Due to the introduced misspecifications, we say that the maximizer of (5) is a QML estimator.

In principle, QML estimation of θ can be performed by writing the model in vectorized form and maximizing the prediction error decomposition of the Gaussian likelihood obtained from the Kalman filter (see e.g. section 7.2 in Durbin and Koopman, 2012). This approach is applicable when p_1 , and p_2 are relatively small, but it becomes quickly unfeasible in larger settings due to a lack of closed form solution. Furthermore, by vectorizing the data we lose the bilinear structure of the model. We resort to the EM algorithm instead.

3.2 EM algorithm

The EM algorithm is an iterative procedure proposed by Dempster et al. (1977) to maximize the log-likelihood in problems where missing or latent observations make the likelihood intractable. This procedure works in two steps: given a set of parameter values, the E-step computes the expectation of the log-likelihood conditional on the observed data, thus "filling" the missing observations; the M-step maximizes the expected log-likelihood with respect to the model parameters. These two steps are iterated until a convergence criterion is satisfied. As the factor process $\{\mathbf{F}_t\}$ is unobserved in our setting, the EM algorithm is a suitable option to perform QML estimation.

3.2.1 Kalman smoother

For any iteration $n \ge 0$ of the EM algorithm, and given an estimator $\widehat{\theta}^{(n)}$, we run the Kalman smoother on the vectorized approximate DMFM:

$$\mathbf{y}_t = (\mathbf{C} \otimes \mathbf{R}) \mathbf{f}_t + \mathbf{e}_t, \tag{6}$$

$$\mathbf{f}_t = (\mathbf{B} \otimes \mathbf{A}) \mathbf{f}_{t-1} + \mathbf{u}_t. \tag{7}$$

The Kalman smoother gives the linear projection $f_{t|T}^{(n)} = \operatorname{Proj}_{\widehat{\theta}^{(n)}}[f_t|\mathbf{Y}_T]$ and the associated MSE, denoted as $\mathbf{\Pi}_{t|T}^{(n)}$. Moreover, by considering the Kalman smoother for the augmented state vector $(f'_t f'_{t-1})'$, we denote the top-left $k_1k_2 \times k_1k_2$ block of the associated $2k_1k_2 \times 2k_1k_2$ MSE as $\Delta_{t|T}^{(n)}$. In particular, to run the Kalman smoother we need also to specify the covariance matrices of \mathbf{e}_t and \mathbf{u}_t . For \mathbf{e}_t , consistently with the misspecified log-likelihood (5), we use an estimator of the $p_1p_2 \times p_1p_2$ misspecified covariance matrix $dg(\mathbf{K}) \otimes dg(\mathbf{H})$, which is always invertible. While for \mathbf{u}_t we can use an estimator of the $k_1k_2 \times k_1k_2$ full-covariance matrix $\mathbf{Q} \otimes \mathbf{P}$. We refer to, e.g., Section 4.4 in Durbin and Koopman (2012), for the explicit expressions of these quantities. Note that the engineering literature proposes matrix versions of the Kalman filter (e.g. Choukroun et al., 2006) for matrix state-space models like the one in (3)-(4). However, these approaches heavily rely on the vec(\cdot) operator and offer only minor computational advantages, primarily due to algebraic simplifications. Similarly to our approach, also Yu et al. (2024) utilize the vectorized Kalman filter.

Under joint Gaussianity of F_T and Y_T , it is known that $\mathsf{f}_{t|T}^{(n)}$, $\mathbf{\Pi}_{t|T}^{(n)}$, and $\boldsymbol{\Delta}_{t|T}^{(n)}$ are estimators of the first and second conditional moments of f_t given Y_T , obtained when computing expectations using the estimated parameters $\hat{\boldsymbol{\theta}}^{(n)}$. Nevertheless, we stress here that consistency of the proposed estimators of the matrices of loadings and factors does not require any Gaussianity assumption (see the next section for more details).

3.2.2 E-step

In the E-step, we use the estimated conditional moments to compute the expected Gaussian loglikelihood of the approximate DMFM. Spcifically, given Y_T and $\hat{\theta}^{(n)}$, by Bayes' rule, we have¹

$$\ell(\mathsf{Y}_T;\boldsymbol{\theta}) = \underbrace{\mathbb{E}_{\widehat{\boldsymbol{\theta}}^{(n)}}[\ell(\mathsf{Y}_T|\mathsf{F}_T;\boldsymbol{\theta})|\mathsf{Y}_T] + \mathbb{E}_{\widehat{\boldsymbol{\theta}}^{(n)}}[\ell(\mathsf{F}_T;\boldsymbol{\theta})|\mathsf{Y}_T]}_{\mathcal{Q}(\boldsymbol{\theta},\widehat{\boldsymbol{\theta}}^{(n)})} - \mathbb{E}_{\widehat{\boldsymbol{\theta}}^{(n)}}[\ell(\mathsf{F}_T|\mathsf{Y}_T;\boldsymbol{\theta})|\mathsf{Y}_T].$$

¹Notice that conditioning on Y_T which is a vector is equivalent to conditioning on the sequence of matrices $\{Y_1, ..., Y_T\}$, likewise we can equivalently write the argument of the log-likelihood in both ways. The same applies for F_T and the sequence of matrices $\{F_1, ..., F_T\}$. Therefore, in order to avoid introducing further notation hereafter we use only Y_T and F_T as conditioning random variables and as arguments of the log-likelihoods even when the latter are expressed explicitly as function of matrix valued time series.

As proved in Dempster et al. (1977) and Wu (1983), for models in the exponential family, maximizing $\ell(\mathbf{Y}_T; \boldsymbol{\theta})$ is equivalent to maximizing $\mathcal{Q}(\boldsymbol{\theta}, \widehat{\boldsymbol{\theta}}^{(n)})$, and we thus need to compute only the latter. Specifically, we have (up to constant terms and initial conditions)

$$\mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T}|\mathbf{F}_{T};\boldsymbol{\theta})|\mathbf{Y}_{T}] = -\frac{T}{2}(p_{1}\log(|\mathbf{K}|) + p_{2}\log(|\mathbf{H}|))$$

$$-\frac{1}{2}\sum_{t=1}^{T}\mathbb{E}_{\widehat{\boldsymbol{\theta}}^{(n)}}\left[\mathbf{tr}\left(\mathbf{H}^{-1}(\mathbf{Y}_{t}-\mathbf{RF}_{t}\mathbf{C}')\mathbf{K}^{-1}(\mathbf{Y}_{t}-\mathbf{RF}_{t}\mathbf{C}')'\right)|\mathbf{Y}_{T}\right],$$

$$\mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{F}_{T};\boldsymbol{\theta})|\mathbf{Y}_{T}] = -\frac{T-1}{2}(k_{1}\log(|\mathbf{Q}|) + k_{2}\log(|\mathbf{P}|))$$

$$-\frac{1}{2}\sum_{t=1}^{T}\mathbb{E}_{\widehat{\boldsymbol{\theta}}^{(n)}}\left[\mathbf{tr}\left(\mathbf{P}^{-1}(\mathbf{F}_{t}-\mathbf{AF}_{t-1}\mathbf{B}')\mathbf{Q}^{-1}(\mathbf{F}_{t}-\mathbf{AF}_{t-1}\mathbf{B}')'\right)|\mathbf{Y}_{T}\right].$$

$$(8)$$

$$(9)$$

Notice that these log-likelihoods depend directly on the data in its matrix form (see Appendix B.1 for the derivation from the vectorized form).

3.2.3 M-step

In the M-step, we maximize (8) and (9) to obtain a new estimate of the parameters $\widehat{\theta}^{(n+1)}$. In particular, at any $n \ge 0$ iteration, the row and column loadings estimators are given by (see Appendix B.2 for the derivation):

$$\widehat{\mathbf{R}}^{(n+1)} = \left(\sum_{t=1}^{T} \mathbf{Y}_t \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \mathbf{F}_{t|T}^{(n)\prime}\right) \left(\sum_{t=1}^{T} \left(\widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)}\right) \star \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)}\right)\right)^{-1},$$
(10)

$$\widehat{\mathbf{C}}^{(n+1)} = \left(\sum_{t=1}^{T} \mathbf{Y}_{t}' \widehat{\mathbf{H}}^{(n)-1} \widehat{\mathbf{R}}^{(n+1)} \mathbf{F}_{t|T}^{(n)}\right) \left(\sum_{t=1}^{T} \left(\widehat{\mathbf{R}}^{(n+1)'} \widehat{\mathbf{H}}^{(n)-1} \widehat{\mathbf{R}}^{(n+1)}\right) \star \left(\mathbb{K}_{k_{1}k_{2}} \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)'} + \mathbf{\Pi}_{t|T}^{(n)}\right) \mathbb{K}_{k_{1}k_{2}}'\right)\right)^{-1}.$$

Clearly, the estimators of \mathbf{R} and \mathbf{C} depend on each other. Here, we choose to first estimate \mathbf{R} conditional on the previous iteration estimator of \mathbf{C} . Since, as shown in the next section, all these estimators are consistent at any iteration $n \ge 0$, we ensure, in this way, that the bilinear structure of the DMFM is preserved. We can of course equivalently choose to first estimate \mathbf{C} conditional on the previous iteration estimator of \mathbf{R} .

By, using $\widehat{\mathbf{R}}^{(n+1)}$ and $\widehat{\mathbf{C}}^{(n+1)}$ we can compute the estimators of **H** and **K** which are enforced to be diagonal matrices in agreement with the considered misspecified log-likelihood given in (5). Thus, for $i=1,\ldots,p_1$, we have:

$$\begin{split} [\widehat{\mathbf{H}}^{(n+1)}]_{ii} = & \frac{1}{Tp_2} \sum_{t=1}^{T} \left[\mathbf{Y}_t \widehat{\mathbf{K}}^{(n)-1} \mathbf{Y}'_t - \mathbf{Y}_t \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n+1)} \mathbf{F}_{t|T}^{(n)'} \widehat{\mathbf{R}}^{(n+1)'} - \widehat{\mathbf{R}}^{(n+1)} \mathbf{F}_{t|T}^{(n)} \widehat{\mathbf{C}}^{(n+1)'} \widehat{\mathbf{K}}^{(n)-1} \mathbf{Y}'_t \\ & + \left(\widehat{\mathbf{C}}^{(n+1)'} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n+1)} \right) \star \left(\left(\mathbb{I}_{k_2} \otimes \widehat{\mathbf{R}}^{(n+1)} \right) \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)'} + \mathbf{\Pi}_{t|T}^{(n)} \right) \left(\mathbb{I}_{k_2} \otimes \widehat{\mathbf{R}}^{(n+1)} \right)' \right) \right]_{ii}, \end{split}$$

and $[\widehat{\mathbf{H}}^{(n+1)}]_{ij} = 0$ if $i \neq j$. Likewise, for $i=1,\ldots,p_2$, we have:

$$\begin{split} [\widehat{\mathbf{K}}^{(n+1)}]_{ii} &= \frac{1}{Tp_1} \sum_{t=1}^{T} \left[\mathbf{Y}_t' \widehat{\mathbf{H}}^{(n+1)-1} \mathbf{Y}_t - \mathbf{Y}_t' \widehat{\mathbf{H}}^{(n+1)-1} \widehat{\mathbf{R}}^{(n+1)} \mathbf{F}_{t|T}^{(n)} \widehat{\mathbf{C}}^{(n+1)\prime} - \widehat{\mathbf{C}}^{(n+1)} \mathbf{F}_{t|T}^{(n)\prime} \widehat{\mathbf{R}}^{(n+1)\prime} \widehat{\mathbf{H}}^{(n+1)-1} \mathbf{Y}_t \\ &+ \left(\widehat{\mathbf{R}}^{(n+1)\prime} \widehat{\mathbf{H}}^{(n+1)-1} \widehat{\mathbf{R}}^{(n+1)} \right) \star \left(\left(\mathbb{I}_{k_1} \otimes \widehat{\mathbf{C}}^{(n+1)} \right) \left(\mathbb{K}_{k_1 k_2} \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)} \right) \mathbb{K}_{k_1 k_2}^{\prime} \right) \left(\mathbb{I}_{k_1} \otimes \widehat{\mathbf{C}}^{(n+1)} \right)^{\prime} \right) \right]_{ii}. \end{split}$$

and $[\widehat{\mathbf{K}}^{(n+1)}]_{ij}=0$ if $i \neq j$. As for the loadings, the estimators of \mathbf{H} and \mathbf{K} depend on each other, and, in order to preserve the bilinear structure of the model, we first estimate \mathbf{H} conditional on the previous iteration estimator of \mathbf{K} .

Finally, in this paper, we are not directly interested in the autoregressive parameters and the related innovation covariances. These are just needed to run the Kalman smoother, which, in our algorithm, is implemented on vectorized data. Therefore, we just consider the estimators obtained from the vectorized MAR:

$$\widehat{\mathbf{B} \otimes \mathbf{A}}^{(n+1)} = \left(\sum_{t=2}^{T} \mathbf{f}_{t|T}^{(n+1)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Delta}_{t|T}^{(n)}\right) \left(\sum_{t=2}^{T} \mathbf{f}_{t-1|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Pi}_{t-1|T}^{(n)}\right)^{-1},$$
$$\widehat{\mathbf{Q} \otimes \mathbf{P}}^{(n+1)} = \frac{1}{T} \sum_{t=2}^{T} \mathbf{f}_{t|T}^{(k)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)} - \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Delta}_{t|T}^{(n)}\right) \widehat{\mathbf{B} \otimes \mathbf{A}}^{(n+1)\prime}.$$

Clearly, these estimators do not satisfy the constraints imposed by the bilinear structure of the MAR (see Appendix B.2 for the expressions of the estimators $\widehat{\mathbf{A}}^{(n+1)}$, $\widehat{\mathbf{B}}^{(n+1)}$, $\widehat{\mathbf{P}}^{(n+1)}$, and $\widehat{\mathbf{Q}}^{(n+1)}$ satisfying those constraints). Nevertheless, the asymptotic properties of the estimated loadings and factor matrices are unaffected by this choice. This is also confirmed by our simulations (see Section 6).

3.2.4 Initialization

We use the projected estimator (PE) of Yu et al. (2022) to obtain initial estimates of the row and column loading matrices and the factor matrices. Let $\mathbf{M}_1 = \frac{1}{p_1 p_2 T} \sum_{t=1}^T \mathbf{Y}_t \mathbf{Y}_t'$ and $\mathbf{M}_2 = \frac{1}{p_1 p_2 T} \sum_{t=1}^T \mathbf{Y}_t' \mathbf{Y}_t$, and define $\overline{\mathbf{X}}_t = p_2^{-1} \mathbf{Y}_t \overline{\mathbf{C}}$ and $\overline{\mathbf{Z}}_t = p_1^{-1} \mathbf{Y}_t' \overline{\mathbf{R}}$, where $\overline{\mathbf{R}} = \sqrt{p_1} \Gamma^{M_1}$ and $\overline{\mathbf{C}} = \sqrt{p_2} \Gamma^{M_2}$, with Γ^{M_i} containing the k_i leading eigenvectors of \mathbf{M}_i , for i=1,2. Let $\overline{\mathbf{M}}_1 = \frac{1}{p_1 p_2 T} \sum_{t=1}^T \overline{\mathbf{X}}_t \overline{\mathbf{X}}_t'$ and $\overline{\mathbf{M}}_2 = \frac{1}{p_1 p_2 T} \sum_{t=1}^T \overline{\mathbf{Z}}_t' \overline{\mathbf{Z}}_t$. Preestimators of \mathbf{R} and \mathbf{C} are given by:

$$\widehat{\mathbf{R}}^{(0)} = \sqrt{p_1} \Gamma^{\overline{M}_1}, \qquad \widehat{\mathbf{C}}^{(0)} = \sqrt{p_2} \Gamma^{\overline{M}_2}.$$
(11)

The pre-estimator of the factor matrix is then obtained by linear projection as: $\mathbf{\tilde{F}}_t = \frac{\mathbf{\hat{R}}^{(0)'} \mathbf{Y}_t \mathbf{\hat{C}}^{(0)}}{p_1 p_2}$. Then, letting $\mathbf{\hat{E}}^{(0)} = \mathbf{Y}_t - \mathbf{\hat{R}}^{(0)} \mathbf{\tilde{F}}_t \mathbf{\hat{C}}^{(0)'}$, the pre-estimators of **H** and **K** are given by:

$$\widehat{\mathbf{K}}^{(0)} = \frac{1}{Tp_1} \sum_{t=1}^{T} \operatorname{tr}\left(\widehat{\mathbf{E}}_t^{(0)\prime} \widehat{\mathbf{E}}_t^{(0)}\right), \qquad \widehat{\mathbf{H}}^{(0)} = \frac{1}{Tp_1} \sum_{t=1}^{T} \operatorname{tr}\left(\widehat{\mathbf{E}}_t^{(0)} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{E}}_t^{(0)\prime}\right).$$
(12)

Notice that only their diagonal terms of these estimators are needed for running the EM algorithm.

Then, denoting the pre-estimator of the vectorized factors as $\tilde{f}_t = \frac{(\hat{C}^{(0)} \otimes \hat{R}^{(0)})' y_t}{p_1 p_2}$, the pre-estimators for the MAR parameters are given by:

$$\widehat{\mathbf{B} \otimes \mathbf{A}}^{(0)} = \left(\sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t} \widetilde{\mathbf{f}}_{t-1}^{\prime}\right) \left(\sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t-1} \widetilde{\mathbf{f}}_{t-1}^{\prime}\right)^{-1},$$
$$\widehat{\mathbf{Q} \otimes \mathbf{P}}^{(0)} = \left(\sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t} - \widehat{\mathbf{B} \otimes \mathbf{A}}^{(0)} \widetilde{\mathbf{f}}_{t-1}\right) \left(\sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t} - \widehat{\mathbf{B} \otimes \mathbf{A}}^{(0)} \widetilde{\mathbf{f}}_{t-1}\right)^{\prime},$$

Alternatively, we can obtain the pre-estimators $\widehat{\mathbf{A}}^{(0)}$, $\widehat{\mathbf{B}}^{(0)}$, $\widehat{\mathbf{P}}^{(0)}$ and $\widehat{\mathbf{Q}}^{(0)}$ with the projection method of Chen et al. (2021) computed when using the estimated factors $\widetilde{\mathbf{f}}_t$.

Finally, we initialize the Kalman filter by setting $\mathbf{f}_{0|0}^{(0)} = \mathbf{0}_k$ at n = 0 and $\mathbf{f}_{0|0}^{(n)} = \mathbf{f}_{0|T}^{(n-1)}$ at $n \ge 1$, and $\mathbf{\Pi}_{0|0}^{(n)} = \mathbb{I}_k$ at $n \ge 0$.

3.2.5 EM convergence

As in Doz et al. (2012), we run the EM algorithm for a finite number of iterations n_{max} and, for a given tolerance level ϵ , it is stopped at the first iteration $n^* < n_{\text{max}}$ such that

$$\Delta \mathcal{L}_{n^{*}} = \frac{\left| \mathcal{L}\left(\mathsf{Y}_{t}, \widehat{\boldsymbol{\theta}}^{(n^{*}+1)}\right) - \mathcal{L}\left(\mathsf{Y}_{t}; \widehat{\boldsymbol{\theta}}^{(n^{*})}\right) \right|}{\frac{1}{2} \left| \mathcal{L}\left(\mathsf{Y}_{T}; \widehat{\boldsymbol{\theta}}^{(n^{*}+1)}\right) + \mathcal{L}\left(\mathsf{Y}_{t}; \widehat{\boldsymbol{\theta}}^{(n^{*})}\right) \right|} < \epsilon,$$

where $\mathcal{L}(Y_T; \boldsymbol{\theta})$ is the prediction error log-likelihood, which, in turn, is given by (up to constant terms and initial conditions):

$$\begin{split} \mathcal{L}(\mathbf{Y}_T; \boldsymbol{\theta}) &= -\frac{1}{2} \sum_{t=1}^T \log \left| (\mathbf{C} \otimes \mathbf{R}) \mathbf{\Pi}_{t|t-1} (\mathbf{C} \otimes \mathbf{R})' + \mathbf{K} \otimes \mathbf{H} \right| \\ &- \frac{1}{2} \sum_{t=1}^T \left(\mathsf{y}_t - (\mathbf{C} \otimes \mathbf{R}) \mathsf{f}_{t|t-1} \right)' \left((\mathbf{C} \otimes \mathbf{R}) \mathbf{\Pi}_{t|t-1} (\mathbf{C} \otimes \mathbf{R})' + \mathbf{K} \otimes \mathbf{H} \right)^{-1} \left(\mathsf{y}_t - (\mathbf{C} \otimes \mathbf{R}) \mathsf{f}_{t|t-1} \right), \end{split}$$

where $f_{t|t-1}$ is the one-step-ahead prediction of f_t computed via the Kalman filter and $\Pi_{t|t-1}$ is the associated MSE, both computed when using θ .

Once the EM algorithm reaches convergence, we define the EM estimator of the model parameters as

 $\widehat{\boldsymbol{\theta}} = \widehat{\boldsymbol{\theta}}^{(n^*+1)}$. Finally, we obtain a final estimate of the factor matrices by running the Kalman smoother one last time, that is $\widehat{\mathbf{F}}_t \equiv \text{unvec}\left(\mathbf{f}_{t|T}^{(n^*+1)}\right)$ for any $t=1,\ldots,T$.

4 Asymptotic results

We make the following standard assumptions.

Assumption 1. (COMMON COMPONENT).

- (i) $\|\mathbf{R}\|_{\max} \leq \bar{r}$ and $\|\mathbf{C}\|_{\max} \leq \bar{c}$, for finite positive reals \bar{r} and \bar{c} , and, as $\min\{p_1, p_2\} \rightarrow \infty$, $\|p_1^{-1}\mathbf{R'R} \mathbb{I}_{k_1}\| \rightarrow 0$ and $\|p_2^{-1}\mathbf{C'C} \mathbb{I}_{k_2}\| \rightarrow 0$.
- (*ii*) For all $t \in \mathbb{Z}$, $\mathbb{E}[\mathbf{F}_t] = \mathbf{0}_{k_1,k_2}$, $\mathbb{E}||\mathbf{F}_t||^4 < \infty$, and, as $T \to \infty$, $T^{-1} \sum_{t=1}^T \mathbf{F}_t \mathbf{F}_t' \xrightarrow{p} \mathbf{\Sigma}_1$ and $T^{-1} \sum_{t=1}^T \mathbf{F}_t' \mathbf{F}_t \xrightarrow{p} \mathbf{\Sigma}_2$ where $\mathbf{\Sigma}_i$ is a $k_i \times k_i$ matrix with distinct eigenvalues and spectral decomposition $\mathbf{\Sigma}_i = \mathbf{\Gamma}_i^F \mathbf{\Lambda}_i^F \mathbf{\Gamma}_i^{F'}$, for i=1,2. The factor numbers k_1 and k_2 are finite and independent of T, p_1 , and p_2 .
- (*iii*) $\|\mathbf{B}\| < 1$ and $\|\mathbf{A}\| < 1$.
- (iv) For all $t \in \mathbb{Z}$, $\mathbb{E}[\mathbf{U}_t] = \mathbf{0}_{k_1,k_2}$, and $\mathbb{E}[\mathbf{U}_t\mathbf{U}'_t] = \mathbf{Ptr}(\mathbf{Q})$ and $\mathbb{E}[\mathbf{U}'_t\mathbf{U}_t] = \mathbf{Qtr}(\mathbf{P})$, with \mathbf{P} and \mathbf{Q} $k_1 \times k_1$ and $k_2 \times k_2$ positive definite matrices such that $C_P^{-1} \leq [\mathbf{P}]_{ii} \leq C_P$ and $C_Q^{-1} \leq [\mathbf{Q}]_{ii} \leq C_Q$, for some finite positive reals C_P and C_Q independent of i, and spectral decomposition $\mathbf{P} = \mathbf{\Gamma}^P \mathbf{\Lambda}^P \mathbf{\Gamma}^{P'}$ and $\mathbf{Q} = \mathbf{\Gamma}^Q \mathbf{\Lambda}^Q \mathbf{\Gamma}^{Q'}$. For all $t, k \in \mathbb{Z}$ with $k \neq 0$, \mathbf{U}_t and \mathbf{U}_{t-k} are independent.

Assumption 2. (IDIOSYNCRATIC COMPONENT).

(i) The process $\{\mathbf{e}_t\}$ is α -mixing, i.e. there exists $\gamma > 2$ such that $\sum_{h=1}^{\infty} \alpha(h)^{1-2/\gamma} \leq \infty$, with $\alpha(h) = \sup_{t \in \mathbb{Z}} \sup_{A \in \mathcal{F}_{-\infty}^t, B \in \mathcal{F}_{t+h}^\infty} |\Pr(A \cap B) - \Pr(A) \cap \Pr(B)|$ and \mathcal{F}_{τ}^s the σ -field generated by $\{\mathbf{e}_t: \tau \leq t \leq s\}$.

There exists a finite positive real c, independent of T, p_1 , and p_2 , such that:

(*ii*) for all t=1,...,T, $i=1,...,p_1$, $j=1,...,p_2$, $\mathbb{E}[e_{tij}]=0$, $\mathbb{E}[|e_{tij}|^4] \leq c$. Moreover

$$\mathbb{E}\left[\mathbf{E}_{t}\mathbf{E}_{t}'\right] = \mathbf{H}\mathbf{tr}(\mathbf{K}), \qquad \mathbb{E}\left[\mathbf{E}_{t}'\mathbf{E}_{t}\right] = \mathbf{K}\mathbf{tr}(\mathbf{H}),$$

with \mathbf{H} and \mathbf{K} $p_1 \times p_1$ and $p_2 \times p_2$ positive definite matrices such that $\operatorname{tr}(\mathbf{H}) = p_1$, and $C_H^{-1} \leq [\mathbf{H}]_{ii} \leq C_H$ and $C_K^{-1} \leq [\mathbf{K}]_{ii} \leq C_K$, for some finite positive reals C_H and C_K independent of i.

(iii) for all $T, p_1, p_2 \in \mathbb{N}$,

$$\frac{1}{Tp_1p_2} \sum_{t,s=1}^T \sum_{i_1,i_2=1}^{p_1} \sum_{j_1,j_2=1}^{p_2} |\mathbb{E}[e_{ti_2j_1}e_{si_1j_2}]| \le c$$

(iv) for all t=1,...,T, $i,l_1=1,...,p_1$, $j,h_1=1,...,p_2$, and all $T,p_1,p_2 \in \mathbb{N}$,

$$\sum_{s=1}^{T} \sum_{l_{2}=1}^{p_{1}} \sum_{h=1}^{p_{2}} |\mathbb{C}\mathrm{ov}[e_{tij}e_{tl_{1}j}, e_{sih}e_{sl_{2}h}]| \le c, \qquad \sum_{s=1}^{T} \sum_{l=1}^{p_{1}} \sum_{h_{2}=1}^{p_{2}} |\mathbb{C}\mathrm{ov}[e_{tij}e_{tih_{1}}, e_{sl_{2}j}e_{sih_{2}}]| \le c,$$
$$\sum_{s=1}^{T} \sum_{i,l_{2}=1}^{p_{1}} \sum_{j,h_{2}=1}^{p_{2}} (|\mathbb{C}\mathrm{ov}[e_{tij}e_{tl_{1}h_{1}}, e_{sij}e_{sl_{2}h_{2}}]| + |\mathbb{C}[e_{tl_{1}j}e_{tih_{1}}, e_{sl_{2}j}e_{sih_{2}}]|) \le c.$$

Assumption 3. (COMPONENTS DEPENDENCE). There exists a finite positive real c, independent of T, p_1 , and p_2 , such that:

(i) for any deterministic vector \mathbf{v} and \mathbf{w} satisfying $\|\mathbf{v}\|=1$ and $\|\mathbf{w}\|=1$,

$$\mathbb{E}\left[\left\|\frac{1}{\sqrt{T}}\sum_{t=1}^{T}\left(\mathbf{F}_{t}\mathbf{v}'\mathbf{E}_{t}\mathbf{w}\right)\right\|_{F}^{2}\right] \leq c;$$

(*ii*) for all $T, p_1, p_2 \in \mathbb{N}$, and all $i=1,...,p_1, j=1,...,p_2$,

$$\left\|\sum_{h=1}^{p_2} \mathbb{E}\left[\boldsymbol{\zeta}_{ij} \otimes \boldsymbol{\zeta}_{ih}\right]\right\|_{\max} \leq c, \qquad \left\|\sum_{l=1}^{p_1} \mathbb{E}\left[\boldsymbol{\zeta}_{ij} \otimes \boldsymbol{\zeta}_{lj}\right]\right\|_{\max} \leq c,$$

and all $i_1, l_1 = 1, \dots, p_1, j_1, h_1 = 1, \dots, p_2$,

$$\left\| \sum_{j_{2}=1}^{p_{2}} \sum_{h_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{1}j_{2}} \otimes \boldsymbol{\zeta}_{l_{1}h_{2}}\right] \right\|_{\max} \leq c, \qquad \left\| \sum_{i_{2}=1}^{p_{1}} \sum_{h_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{2}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{2}}\right] \right\|_{\max} \leq c, \qquad \left\| \sum_{i_{2}=1}^{p_{1}} \sum_{h_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{2}j_{1}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right] \right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{1}} \sum_{j_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{1}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right] \right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{1}} \sum_{j_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{1}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right] \right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{1}} \sum_{j_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{1}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right] \right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{1}} \sum_{j_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{1}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right] \right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{1}} \sum_{j_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{1}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{1}} \sum_{j_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{2}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{1}j_{1}} \otimes \boldsymbol{\zeta}_{l_{1}h_{1}}, \boldsymbol{\zeta}_{i_{2}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{2}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right]\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{i_{2}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{l_{2}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{l_{2}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{l_{2}j_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \otimes \boldsymbol{\zeta}_{l_{2}h_{1}}\right\|_{\max} \leq c, \qquad \left\| \sum_{l_{2}=1}^{p_{2}} \mathbb{C}\operatorname{ov}\left[\boldsymbol{\zeta}_{l_{2}j_{2}} \otimes \boldsymbol{\zeta}_{l_{$$

Assumptions 1(i)-1(ii) matches Assumptions B and C in Yu et al. (2022). Assumption 1(iii) guarantees the stationarity of the MAR(1) model. Assumption 1(iv) requires the factor innovations \mathbf{U}_t to have positive definite covariance matrix. Assumptions 2 closely matches Assumptions A and D in Yu et al. (2022). In particular, Assumption 2(i) controls serial idiosyncratic dependence by requiring them to be α -mixing (see also Chen and Fan, 2023, Assumption 1). Assumption 2(i) imposes finite absolute fourth moments and requires **H** and **K** to be positive definite matrices, and, since **H** and **K** are only determined up to a positive constant, and only their Kronecker product, $\mathbf{K} \otimes \mathbf{H}$, is uniquely defined, we impose the constraint $\text{tr}(\mathbf{H})=p_1$ (see, e.g., Viroli (2012)). Assumption 2(ii) controls cross-sectional idiosyncratic dependence across both rows and columns (see also Chen and Fan, 2023, Assumption

2), while Assumption 2(iv) bounds fourth order cumulants to allow for consistent estimation of the second order moments. Last, Assumption 3 bounds the dependence between factors and idiosyncratic components and matches Assumption E in Yu et al. (2022).

Under the above assumptions can then derive theoretical results on the convergence rates of the EM estimators for the loading and factor matrices $\widehat{\mathbf{R}}$, $\widehat{\mathbf{C}}$, and $\widehat{\mathbf{F}}_t$, defined in Section 3.2. In particular, we show that using the initialization setting discussed in Section 3.2.4, the estimators converge in probability after one iteration of the EM algorithm, i.e. $n^*=0$.

Proposition 1. Let $\widehat{\mathbf{R}} \equiv \widehat{\mathbf{R}}^{(n^*+1)}$ and $\widehat{\mathbf{C}} \equiv \widehat{\mathbf{C}}^{(n^*+1)}$ be the EM estimators of the loadings, with $n^* \ge 0$. Under Assumptions 1 through 3, there exist matrices $\widehat{\mathbf{J}}_1$ of size $k_1 \times k_1$ and $\widehat{\mathbf{J}}_2$ of size $k_2 \times k_2$ satisfying $\widehat{\mathbf{J}}_1 \widehat{\mathbf{J}}'_1 \xrightarrow{p} \mathbb{I}_{k_1}$ and $\widehat{\mathbf{J}}_2 \widehat{\mathbf{J}}'_2 \xrightarrow{p} \mathbb{I}_{k_2}$, such that, as $\min\{p_1, p_2, T\} \rightarrow \infty$,

$$\min\left(\sqrt{Tp_1}, \sqrt{Tp_2}, p_1p_2\right) \left\| \frac{\widehat{\mathbf{R}} - \widehat{\mathbf{RJ}}_1}{\sqrt{p_1}} \right\| = O_p(1), \quad \min\left(\sqrt{Tp_1}, \sqrt{Tp_2}, p_1p_2\right) \left\| \frac{\widehat{\mathbf{C}} - \widehat{\mathbf{CJ}}_2}{\sqrt{p_2}} \right\| = O_p(1),$$

and, for any given $i=1,\ldots,p_1$ and $j=1,\ldots,p_2$, as $\min\{p_1,p_2,T\} \rightarrow \infty$,

$$\min\left(\sqrt{Tp_1}, \sqrt{Tp_2}, p_1p_2\right) \left\| \widehat{\mathbf{r}}_{i} - \mathbf{r}_{i} \cdot \widehat{\mathbf{J}}_1 \right\| = O_p(1), \quad \min\left(\sqrt{Tp_1}, \sqrt{Tp_2}, p_1p_2\right) \left\| \widehat{\mathbf{c}}_{j} - \mathbf{c}_{j} \cdot \widehat{\mathbf{J}}_2 \right\| = O_p(1).$$

Proposition 2. Let $\widehat{\mathbf{F}}_t \equiv \text{unvec}(\widehat{\mathbf{f}}_{t|T}^{(n^*)})$ be the Kalman smoother estimator of the factors computed using the estimated parameters $\widehat{\boldsymbol{\theta}}^{(n^*+1)}$, with $n^* \geq 0$. Under Assumptions 1 through 3, and given $\widehat{\mathbf{J}}_1$ and $\widehat{\mathbf{J}}_2$ as defined in Proposition 1, for any given t=1,...,T, as $\min\{p_1,p_2,T\} \rightarrow \infty$,

$$\min\left(\sqrt{Tp_1}, \sqrt{Tp_2}, \sqrt{p_1p_2}\right) \left\| \widehat{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right\| = O_p(1).$$

Note that numerous applications in economics and finance might present situations where one of the dimension of the matrix \mathbf{Y}_t , say p_1 , can be assumed to be large $(p_1 \to \infty)$, while the other one, say p_2 , is fixed $(p_2 < \infty)$. For example, the number of traded assets can be assumed to be very large but the number of liquidity or volatility proxies is likely to be finite. Under these circumstances, consistency of the matrices of factors and factor loadings still holds at the rate min $\left(\frac{1}{\sqrt{T}}, \frac{1}{p_1}\right)$.

5 Extension to the case of missing data

If the data contains missing values, the estimation of the factors and their second moments with the Kalman smoother is still possible (see e.g. section 6.4 in Durbin and Koopman, 2012). Now, since $\mathbb{E}_{\widehat{\theta}_{(n)}}[\ell(\mathsf{F}_T; \theta)|\mathsf{Y}_T]$ depends only on the factors but not on the data, its expression remains unchanged. Thus, the estimators of \mathbf{A} , \mathbf{B} , \mathbf{P} , and \mathbf{Q} are also unchanged. However, $\mathbb{E}_{\widehat{\theta}_{(n)}}[\ell(\mathsf{Y}_T|\mathsf{F}_T; \theta)|\mathsf{Y}_T]$ depends

on the data and therefore its expression is affected by missing values. It follows that we need to adjust the estimators of \mathbf{R} , \mathbf{C} , \mathbf{H} and \mathbf{K} in the M-step accordingly. To this end, we extend the procedure of Bańbura and Modugno (2014) to the matrix setting.

Let \mathbf{W}_t be a $p_1 \times p_2$ matrix with (i, j) entry equal to zero if y_{tij} and equal to one otherwise. For any iteration $n \ge 0$, the estimators of the row and column loadings are then modified to:

$$\operatorname{vec}\left(\widehat{\mathbf{R}}^{(n+1)}\right) = \left(\sum_{t=1}^{T}\sum_{s=1}^{p_{1}}\sum_{q=1}^{p_{1}} \left(\left(\widehat{\mathbf{C}}^{(n)'}\mathbb{D}_{\mathbf{W}_{t}}^{[s,q]}\widehat{\mathbf{K}}^{(n)-1}\widehat{\mathbf{C}}^{(n)}\right) \star \left(\mathbf{f}_{t|T}^{(n)}\mathbf{f}_{t|T}^{(n)'} + \mathbf{\Pi}_{t|T}^{(n)}\right)\right) \otimes \left(\mathbb{E}_{p_{1},p_{1}}^{(s,q)}\widehat{\mathbf{H}}^{(n)-1}\right)\right)^{-1} \\ \times \left(\sum_{t=1}^{T}\operatorname{vec}\left(\left(\mathbf{W}_{t}\circ\widehat{\mathbf{H}}^{(n)-1}\mathbf{Y}_{t}\widehat{\mathbf{K}}^{(n)-1}\right)\widehat{\mathbf{C}}^{(n)}\mathbf{F}_{t|T}^{(n)'}\right)\right), \\ \operatorname{vec}\left(\widehat{\mathbf{C}}^{(n+1)}\right) = \left(\sum_{t=1}^{T}\sum_{k=1}^{p_{2}}\sum_{q=1}^{p_{2}}\left(\left(\widehat{\mathbf{R}}^{(n+1)'}\mathbb{D}_{\mathbf{W}_{t}^{'}}^{[s,q]}\widehat{\mathbf{H}}^{(n)-1}\widehat{\mathbf{R}}^{(n+1)}\right) \star \left(\mathbb{K}_{k_{1}k_{2}}\left(\mathbf{f}_{t|T}^{(n)}\mathbf{f}_{t|T}^{(n)'} + \mathbf{\Pi}_{t|T}^{(n)}\right)\mathbb{K}_{k_{1}k_{2}}^{(n)}\right)\right) \otimes \left(\mathbb{E}_{p_{2},p_{2}}^{(s,q)}\widehat{\mathbf{K}}^{(n)-1}\right)\right)^{-1} \\ \times \left(\sum_{t=1}^{T}\operatorname{vec}\left(\left(\mathbf{W}_{t}\circ\widehat{\mathbf{H}}^{(n)-1}\mathbf{Y}_{t}\widehat{\mathbf{K}}^{(n)-1}\right)'\widehat{\mathbf{R}}^{(n)}\mathbf{F}_{t}^{(n)}\right)\right).$$

Details on the derivations of these expressions as well as expressions for the estimators of the idiosyncratic variances are given in Appendix B.3.

6 Simulation study

We perform Monte Carlo simulations in order to assess the finite sample properties of the EM estimators. We generate observations according to the following DMFM

$$\begin{aligned} \mathbf{Y}_t = \mathbf{R} \mathbf{F}_t \mathbf{C}' + \mathbf{E}_t \\ \mathbf{F}_t = \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}' + \mathbf{U}_t, \qquad \mathbf{U}_t \sim \mathcal{MN}_{k_1, k_2}(\mathbb{O}_{k_1, k_2}, \mathbf{P}, \mathbf{Q}) \\ \mathbf{E}_t = \mathbf{D} \mathbf{E}_{t-1} \mathbf{G}' + \mathbf{V}_t, \qquad \mathbf{V}_t \sim \mathcal{MN}_{p_1, p_2}(\mathbb{O}_{p_1, p_2}, \mathbf{H}, \mathbf{K}) \end{aligned}$$

for t=1,...,T, where the loading matrices are such that $[\mathbf{R}]_{ij}, [\mathbf{C}]_{ij} \sim \mathcal{U}(-1,1)$, and both latent factors and idiosyncratic shocks follow matrix autoregressive processes with

$$[\mathbf{B}]_{ij}, [\mathbf{A}]_{ij} \sim \begin{cases} \mathcal{U}(0.7, 0.9) & i=i \\ & & \\ \mathcal{U}(0, 0.5) & i\neq j \end{cases}, \quad [\mathbf{P}]_{ij}, [\mathbf{Q}]_{ij} = \begin{cases} 1 & i=i \\ 0 & i\neq j \end{cases}$$

$$[\mathbf{D}]_{ij}, [\mathbf{G}]_{ij} = \begin{cases} \mathcal{U}(0, \delta) & i = i \\ 0 & i \neq j \end{cases}, \quad [\mathbf{H}]_{ij}, [\mathbf{K}]_{ij} = \begin{cases} \mathcal{U}(0.7, 1.2) & i = i \\ \tau^{|i-j|} & i \neq j \end{cases}$$

with τ and δ controlling the degree of cross-sectional and serial correlation in the idiosyncratic shocks, respectively.

6.1 Comparison with PE

We compare the performance of the EM estimators for the loading and factor matrices with those of the PE of Yu et al. (2022). We use the column space distance $\mathcal{D}(\mathbf{R}, \widehat{\mathbf{R}})$ and $\mathcal{D}(\mathbf{C}, \widehat{\mathbf{C}})$ to evaluate the loadings matrices estimators, with

$$\mathcal{D}(\mathbf{A},\widehat{\mathbf{A}}) = \left\| \widehat{\mathbf{A}} \left(\widehat{\mathbf{A}}' \widehat{\mathbf{A}} \right)^{-1} \widehat{\mathbf{A}}' - \mathbf{A} \left(\mathbf{A}' \mathbf{A} \right)^{-1} \mathbf{A}' \right\|$$

for any $m \times n$ matrix **A**. To compare the performance of recovering the signal $\mathbf{S}_t = \mathbf{RF}_t \mathbf{C}'_t$, we consider the mean squared error, i.e.

$$MSE_{\mathbf{S}} = \frac{1}{Tp_1p_2} \sum_{t=1}^{T} \left\| \widehat{\mathbf{S}}_t - \mathbf{S}_t \right\|_{\mathrm{F}}^2.$$

For each performance measure, $\mathcal{D}(\mathbf{R}, \widehat{\mathbf{R}})$, $\mathcal{D}(\mathbf{C}, \widehat{\mathbf{C}})$ and MSE_S, Table 1 reports summary statistics for the ratio between the performance obtained with the EM estimator and those obtained with the PE, over 100 replications. The EM algorithm improves upon PE across all the different settings.

Furthermore, in Figure 1 we show for one replication the log-likelihood as function of the number of iterations of the EM algorithm. As expected the log-likelihood increases monotonically and the first few iterations seem to be the most important ones.

| | | | | $T \!=\! 100$ | | | T = 400 | |
|-----------------------|-------|-------|---|--|-----------------------------|---|--|-----------------------------|
| | p_1 | p_2 | $\mathcal{D}(\mathbf{R}, \widehat{\mathbf{R}})$ | $\mathcal{D}(\mathbf{C},\widehat{\mathbf{C}})$ | $\mathrm{MSE}_{\mathbf{S}}$ | $\mathcal{D}(\mathbf{R}, \widehat{\mathbf{R}})$ | $\mathcal{D}(\mathbf{C},\widehat{\mathbf{C}})$ | $\mathrm{MSE}_{\mathbf{S}}$ |
| $\delta = 0 \tau = 0$ | 20 | 20 | 0.98 | 0.97 | 0.92 | 0.98 | 0.96 | 0.91 |
| 0-0,7-0 | | | (0.05) | (0.05) | (0.03) | (0.05) | (0.05) | (0.01) |
| | 10 | 30 | 0.98 | 0.96 | 0.90 | 0.96 | 0.96 | 0.90 |
| | | | (0.09) | (0.05) | (0.03) | (0.10) | (0.05) | (0.01) |
| $\delta = 0.7 = -0.5$ | 20 | 20 | 0.80 | 0.71 | 0.73 | 0.74 | 0.65 | 0.75 |
| 0-0.1,7-0.5 | | | (0.07) | (0.08) | (0.04) | (0.06) | (0.06) | (0.02) |
| | 10 | 30 | 0.87 | 0.68 | 0.70 | 0.82 | 0.63 | 0.75 |
| | | | (0.10) | (0.08) | (0.05) | (0.10) | (0.06) | (0.03) |

Table 1: Average and standard deviation (in parenthesis) of the ratio between the performance of the EM estimator and PE over 100 replications, for each of $\mathcal{D}(\mathbf{R}, \widehat{\mathbf{R}})$, $\mathcal{D}(\mathbf{C}, \widehat{\mathbf{C}})$ and MSE_S.



Figure 1: Log-likelihood as function of the EM iterations.

6.2 Dealing with missing data

We now introduce missing observations in the data generating process. In practice, datasets can present missing observations for several reasons and therefore display different missing patterns. We consider two patterns widely seen in empirical application: (i) randomly missing, when \mathbf{Y}_t displays holes occurring at random at each point in time, e.g. failures in the measurement operations. We assume a constant probability of missings through time, $\pi = \{25\%, 50\%\}$; (ii) block missing, when a fixed portion of \mathbf{Y}_t is missing for a given period of time, e.g. information on some individuals is available for a shorter period of time. We consider a portion $\pi = \{25\%, 50\%\}$ of missing at each point in time. In particular, we assume the bottom-right quarter and the right-half of \mathbf{Y}_t are missing for the first half of the time series when $\pi = 25\%$ and $\pi = 50\%$, respectively.

To fill the missing observations we use the procedure of Xiong and Pelger (2023). This step is required to obtain estimates with the PE and initialize the EM algorithm. Table 2 reports summary statistics for the ratio between the performance obtained with the EM estimator and those obtained with the PE, over 100 replications. These results confirm that estimates obtained with the EM algorithm improve upon those obtained with PE.

| | | | | T = 1 | 100 | | | T = 4 | 400 | |
|-------|-------|-------|---|--|-----------------------------|--------------------------|---|--|-----------------------------|--------------------------|
| π | p_1 | p_2 | $\mathcal{D}(\mathbf{R}, \widehat{\mathbf{R}})$ | $\mathcal{D}(\mathbf{C},\widehat{\mathbf{C}})$ | $\mathrm{MSE}_{\mathbf{S}}$ | $MSE_{\mathbf{Y}^{(0)}}$ | $\mathcal{D}(\mathbf{R}, \widehat{\mathbf{R}})$ | $\mathcal{D}(\mathbf{C},\widehat{\mathbf{C}})$ | $\mathrm{MSE}_{\mathbf{S}}$ | $MSE_{\mathbf{Y}^{(0)}}$ |
| | | | | | | Randoml | y missing | | | |
| 25% | 20 | 20 | 0.85 | 0.87 | 0.78 | 0.99 | 0.86 | 0.86 | 0.81 | 1.00 |
| 2070 | | | (0.10) | (0.12) | (0.05) | (0.00) | (0.08) | (0.09) | (0.01) | (0.00) |
| | 10 | 30 | 0.84 | 0.88 | 0.79 | 0.99 | 0.84 | 0.89 | 0.90 | 1.00 |
| | | | (0.14) | (0.08) | (0.05) | (0.00) | (0.14) | (0.08) | (0.02) | (0.00) |
| 5007 | 20 | 20 | 0.58 | 0.57 | 0.41 | 0.96 | 0.51 | 0.52 | 0.74 | 0.99 |
| 3070 | | | (0.12) | (0.13) | (0.06) | (0.01) | (0.11) | (0.10) | (0.03) | (0.00) |
| | 10 | 30 | 0.53 | 0.55 | 0.36 | 0.94 | 0.42 | 0.46 | 0.64 | 0.98 |
| | | | (0.13) | (0.11) | (0.08) | (0.02) | (0.12) | (0.1) | (0.08) | (0.01) |
| | | | | | | Block | missing | | | |
| 25.07 | 20 | 20 | 0.41 | 0.47 | 0.34 | 0.91 | 0.27 | 0.29 | 0.33 | 0.93 |
| 2070 | | | (0.11) | (0.12) | (0.09) | (0.04) | (0.06) | (0.07) | (0.04) | (0.02) |
| | 10 | 30 | 0.38 | 0.45 | 0.29 | 0.82 | 0.22 | 0.27 | 0.27 | 0.84 |
| | | | (0.12) | (0.14) | (0.09) | (0.07) | (0.08) | (0.05) | (0.04) | (0.03) |
| 50% | 20 | 20 | 0.95 | 0.23 | 0.21 | 0.82 | 0.99 | 0.17 | 0.21 | 0.79 |
| 3070 | | | (0.05) | (0.08) | (0.08) | (0.08) | (0.08) | (0.04) | (0.04) | (0.05) |
| | 10 | 30 | 0.92 | 0.31 | 0.22 | 0.78 | 0.93 | 0.19 | 0.22 | 0.82 |
| | | | (0.09) | (0.12) | (0.10) | (0.13) | (0.10) | (0.06) | (0.06) | (0.12) |

Table 2: Average and standard deviation (in parenthesis) of the ratio between the performance of α -PCA and of the EM algorithm over 100 replications, for each of $\mathcal{D}(\mathbf{R}, \widehat{\mathbf{R}})$, $\mathcal{D}(\mathbf{C}, \widehat{\mathbf{C}})$ and MSE_S.

7 Forecasting volatilities

Despite the abundant use of high-frequency data in the financial econometrics literature, their availability is often limited to major equity indices or large U.S. stocks (Bollerslev et al., 2018), limiting the chance of building high-frequency-based estimates of volatility for a large number of traded companies. Given that volatility measures tend to covary across assets (Barigozzi and Hallin, 2016; Herskovic et al., 2016), a natural question is whether high-frequency-based volatility measures on a set of assets can be used to improve volatility estimates for a set of assets for which only daily observations are available.

We collect daily returns and realized measures for 30 assets listed in the S&P500 under the Financial GICS sector. The data covers the period that goes from the beginning of 2006 to the end of 2010, covering the great financial crisis. We consider the following 10 realized measures of the daily integrated volatility: 5-min realized variance, autocorrelation-corrected 5-min realized variance (Hansen and Lunde, 2006), realized range (Christensen and Podolskij, 2007), realized kernel (Barndorff-Nielsen et al., 2008), pre-averaged realized variance (Jacod et al., 2009), maximum likelihood realized variance (Xiu, 2010), and three OHLC estimators. These measures are all available for half the stocks in the sample becasue we can access high-frequency data. For the remaining stocks we only have OHLC data, and can therefore compute just the three low-frequency variance estimators. We thus obtain a matrix time series of $p_2=10$

estimates of the daily variance on $p_1=30$ assets for T=1259 days, with a block of missing observations.

We cast our empirical problem into the DMFM framework. Let $\sigma_{i,t}^2$ be the *t*th day latent variance of the *i*th asset and define $\tilde{\sigma}_t^2 = (\tilde{\sigma}_{1,t}^2, \dots, \tilde{\sigma}_{p_1,t}^2)'$, with $\tilde{\sigma}_{i,t}^2 = \sigma_{i,t}^2/\bar{\sigma}_{i,t}^2$ and $\bar{\sigma}_i^2 = (\prod_{t=1}^T \sigma_{i,t}^2)^{\frac{1}{T}}$, for all $i=1,\dots,p_1$. We assume that the centered latent log-variances follows,

$$\log\left(\widetilde{\boldsymbol{\sigma}}_{t}^{2}\right) = \mathbf{R}\mathbf{f}_{t} + \boldsymbol{\varepsilon}_{t},$$

where **R** is a $p_1 \times k_1$ loading matrix, \mathbf{f}_t is a vector of k_1 factors evolving according to a VAR(1),

$$\mathbf{f}_t = \mathbf{A}\mathbf{f}_{t-1} + \mathbf{u}_t,$$

with \mathbf{u}_t a vector of shocks. Let $\mathbf{s}_{i,t}$ be the vector of p_2 realized measures for the *i*th asset on the *t*th day and define $\widetilde{\mathbf{s}}_{i,t} = (\widetilde{s}_{i,1,t}, \dots, \widetilde{s}_{1,p_2,t})'$, with $\widetilde{s}_{i,j,t} = s_{i,j,t}/\overline{s}_i$ and $\overline{s}_i = \left(\prod_{t=1}^T \prod_{j=1}^{p_2} s_{i,j,t}\right)^{\frac{1}{T_{p_2}}}$. Let \mathbf{c} be a p_2 loading vector, we assume that

$$\log(\widetilde{\mathbf{s}}_{i,t}) = \mathbf{c}\log(\widetilde{\sigma}_{i,t}^2) + \boldsymbol{\epsilon}_{i,t} \qquad \forall i = 1,...,p_1$$

and with $\epsilon_{i,t}$ a vector of shocks. Let \mathbf{Y}_t be a $p_1 \times p_2$ matrix stacking the vectors of log-transformed realized measures on each of its rows, i.e. $\mathbf{Y}_t = \left(\log(\mathbf{\tilde{s}}_{1,t})', \dots, \log(\mathbf{\tilde{s}}_{p_1,t})'\right)'$. Using matrix notation, we can write our system of equations as

$$\mathbf{Y}_t = \mathbf{R}\mathbf{f}_t\mathbf{c}' + \mathbf{E}_t$$

 $\mathbf{f}_t = \mathbf{A}\mathbf{f}_{t-1} + \mathbf{u}_t$

with $\mathbf{E}_t \neq p_1 \times p_2$ matrix of shocks.

To assess whether the estimation of the latent variance of an asset for which we only have OHLC data can be improved by high-frequency variance estimates on other assets, we compare the forecasts obtained with our DMFM model to those of a dynamic factor model on the low-frequency variance measures of a single asset. We define an in-sample window of 750 observations for the models estimation and leave 509 observations for the out-of-sample forecast evaluation. We estimate a DMFM with $k_1=2$ and $k_2=1$ using the EM algorithm, and obtain forecasts of $\tilde{\sigma}_{i,t}^2$ as follows

$$\widehat{\widetilde{\sigma}}_{i,t}^{2} = \exp\left(\widehat{\mathbf{R}}_{i}^{\prime} \widehat{\mathbf{f}}_{t}\right).$$

For comparison purposes, we consider the following dynamic factor model with k=1 factor for the 15



Figure 2: MSE of the forecasts obtained with the DMFM and the dynamic factor model of each stock.

assets with only OHLC data available,

$$\log(\widetilde{\mathbf{s}}_{i,t}) = \lambda_i g_{i,t} + \mathbf{v}_{i,t}$$
$$g_{i,t} = \beta g_{i,t-1} + u_{i,t}$$

and obtain forecasts of $\tilde{\sigma}_{i,t}^2$ as follows

$$\widehat{\widetilde{\sigma}}_{i,t}^2 = \exp(\widehat{g}_{i,t}).$$

We use the mean squared error (MSE) loss function to evaluate the daily foreasts over the out-ofsample period, and use the daily range as a proxy for the latent variance. Figure 2 depicts the MSE values obtained from the two models. Predictions obtained with the DMFM always outperform those of the stock-specific factor model except in one case, and differences are always statistically significant according to the Diebold-Mariano test.

References

Bai, J. Inferential theory for factor models of large dimensions. *Econometrica*, 71:135–171, 2003.

Bai, J. and Li, K. Statistical analysis of factor models of high dimension. The Annals of Statistics, 40: 436–465, 2012.

- Bai, J. and Li, K. Maximum likelihood estimation and inference for approximate factor models of high dimension. The Review of Economics and Statistics, 98:298–309, 2016.
- Bańbura, M. and Modugno, M. Maximum likelihood estimation of factor models on datasets with arbitrary pattern of missing data. *Journal of Applied Econometrics*, 29:133–160, 2014.
- Bańbura, M., Giannone, D., and Lenza, M. Conditional forecasts and scenario analysis with vector autoregressions for large cross-sections. *International Journal of forecasting*, 31(3):739–756, 2015.
- Barigozzi, M. and Hallin, M. Generalized dynamic factor models and volatilities: recovering the market volatility shocks. *The Econometrics Journal*, 19(1):C33–C60, 2016.
- Barigozzi, M. and Luciani, M. Quasi maximum likelihood estimation and inference of large approximate dynamic factor models via the EM algorithm. arXiv 1910.03821v5, 2024.
- Barndorff-Nielsen, O. E., Hansen, P. R., Lunde, A., and Shephard, N. Designing realized kernels to measure the ex post variation of equity prices in the presence of noise. *Econometrica*, 76(6):1481–1536, 2008.
- Billio, M., Casarin, R., Costola, M., and Iacopini, M. A matrix-variate t model for networks. Frontiers in Artificial Intelligence, 4:49, 2021.
- Billio, M., Casarin, R., Iacopini, M., and Kaufmann, S. Bayesian dynamic tensor regression. Journal of Business & Economic Statistics, 41:429–439, 2023.
- Bollerslev, T., Hood, B., Huss, J., and Pedersen, L. H. Risk everywhere: Modeling and managing volatility. *The Review of Financial Studies*, 31(7):2729–2773, 2018.
- Cahan, E., Bai, J., and Ng, S. Factor-based imputation of missing values and covariances in panel data of large dimensions. *Journal of Econometrics*, 233(1):113–131, 2023.
- Cen, Z. and Lam, C. Tensor time series imputation through tensor factor modelling. arXiv 2403.13153, 2024.
- Chen, E. Y. and Chen, R. Modeling dynamic transport network with matrix factor models: an application to international trade flow. *Journal of Data Science*, 21:490–507, 2022.
- Chen, E. Y. and Fan, J. Statistical inference for high-dimensional matrix-variate factor models. *Journal* of the American Statistical Association, 118:1038–1055, 2023.
- Chen, E. Y., Tsay, R. S., and Chen, R. Constrained factor models for high-dimensional matrix-variate time series. Journal of the American Statistical Association, 115:775–793, 2020.

- Chen, R., Xiao, H., and Yang, D. Autoregressive models for matrix-valued time series. Journal of Econometrics, 222:539–560, 2021.
- Chernis, T., Cheung, C., and Velasco, G. A three-frequency dynamic factor model for nowcasting canadian provincial gdp growth. *International Journal of Forecasting*, 36:851–872, 2020.
- Choukroun, D., Weiss, H., Bar-Itzhack, I. Y., and Oshman, Y. Kalman filtering for matrix estimation. *IEEE Transactions on Aerospace and Electronic Systems*, 42(1):147–159, 2006.
- Christensen, K. and Podolskij, M. Realized range-based estimation of integrated variance. Journal of Econometrics, 141(2):323–349, 2007.
- Dempster, A. P., Laird, N. M., and Rubin, D. B. Maximum likelihood from incomplete data via the em algorithm. Journal of the Royal Statistical Society: Series B, 39:1–22, 1977.
- Doz, C., Giannone, D., and Reichlin, L. A quasi-maximum likelihood approach for large, approximate dynamic factor models. *The Review of Economics and Statistics*, 94:1014–1024, 2012.
- Durbin, J. and Koopman, S. J. *Time series analysis by state space methods*. Oxford university press, 2012.
- Han, Y., Chen, R., and Zhang, C.-H. Rank determination in tensor factor model. *Electronic Journal of Statistics*, 16:1726–1803, 2022.
- Hansen, P. R. and Lunde, A. Realized variance and market microstructure noise. Journal of Business
 & Economic Statistics, 24(2):127–161, 2006.
- He, Y., Kong, X., Trapani, L., and Yu, L. One-way or two-way factor model for matrix sequences? Journal of Econometrics, 235:1981–2004, 2023.
- Herskovic, B., Kelly, B., Lustig, H., and Van Nieuwerburgh, S. The common factor in idiosyncratic volatility: Quantitative asset pricing implications. *Journal of Financial Economics*, 119(2):249–283, 2016.
- Hsu, N.-J., Huang, H.-C., and Tsay, R. S. Matrix autoregressive spatio-temporal models. Journal of Computational and Graphical Statistics, 30:1143–1155, 2021.
- Jacod, J., Li, Y., Mykland, P. A., Podolskij, M., and Vetter, M. Microstructure noise in the continuous case: the pre-averaging approach. *Stochastic processes and their applications*, 119(7):2249–2276, 2009.

- Kapetanios, G., Serlenga, L., and Shin, Y. Estimation and inference for multi-dimensional heterogeneous panel datasets with hierarchical multi-factor error structure. *Journal of Econometrics*, 220:504–531, 2021.
- Merikoski, J. K. and Kumar, R. Inequalities for spreads of matrix sums and products. Applied Mathematics E-Notes, 4(150-159):9, 2004.
- Tipping, M. E. and Bishop, C. M. Probabilistic principal component analysis. Journal of the Royal Statistical Society Series B: Statistical Methodology, 61:611–622, 1999.
- Viroli, C. On matrix-variate regression analysis. Journal of Multivariate Analysis, 111:296–309, 2012.
- Wang, D., Liu, X., and Chen, R. Factor models for matrix-valued high-dimensional time series. Journal of Econometrics, 208:231–248, 2019.
- Wu, C. J. On the convergence properties of the em algorithm. The Annals of statistics, pages 95–103, 1983.
- Xiong, R. and Pelger, M. Large dimensional latent factor modeling with missing observations and applications to causal inference. *Journal of Econometrics*, 233(1):271–301, 2023.
- Xiu, D. Quasi-maximum likelihood estimation of volatility with high frequency data. Journal of Econometrics, 159:235–250, 2010.
- Xu, S., Yuan, C., and Guo, J. Quasi maximum likelihood estimation for large-dimensional matrix factor models. *Journal of Business & Economic Statistics*, 2024. available online.
- Yu, L., He, Y., Kong, X., and Zhang, X. Projected estimation for large-dimensional matrix factor models. *Journal of Econometrics*, 229:201–217, 2022.
- Yu, R., Chen, R., Xiao, H., and Han, Y. Dynamic matrix factor models for high dimensional time series. arXiv 2407.05624, 2024.
- Yuan, C., Gao, Z., He, X., Huang, W., and Guo, J. Two-way dynamic factor models for high-dimensional matrix-valued time series. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 85:1517–1537, 2023.

Supplementary Material

A Notation and results on matrix operations

We adopt the following notation.

- The Hadamard and Kronecker product are denoted with \circ and \otimes , respectively.
- We use $vec(\cdot)$ and $unvec(\cdot)$ to denote the vectorization operation and its inverse.
- \mathbb{I}_d denotes the $d \times d$ identity matrix.
- We use $\mathbf{1}_m$ and $\mathbf{0}_m$ to denote an m dimensional vector filled with ones and zeros, and use $\mathbf{1}_{m,n}$ and $\mathbf{0}_{m,n}$ to denote $m \times n$ matrices filled with ones and zeros, respectively.
- Let **A** be an $n \times n$ matrix, the matrix $dg(\mathbf{A})$ denotes the matrix having the diagonal entries of **A** in the diagonal and zeros in the off-diagonal entries.
- Let A be an m×n matrix, we denote with D_A a mn×mn matrix stacking the columns of A on its diagonal, i.e. D_A=I_{mn}vec(A).
- $\mathbb{E}_{m,n}^{(i,j)}$ denotes a standard basis $(m \times n)$ matrix with a one in the (i,j) entry.
- \mathbb{K}_{nm} denotes an $nm \times nm$ commutation matrix, $\mathbb{K}_{nm} = \sum_{i=1}^{n} \sum_{j=1}^{m} \mathbb{E}_{n,m}^{(i,j)} \otimes \mathbb{E}_{n,m}^{(i,j)'}$.
- The generic (i,j) entry of a matrix **A** is denoted as $a_{ij} \equiv [\mathbf{A}]_{ij}$, while $\mathbf{a}_{i} \equiv [\mathbf{A}]_{i}$ and $\mathbf{a}_{j} \equiv [\mathbf{A}]_{.j}$ denote the generic *i*th row and *j*th column of a matrix **A**, respectively.
- Let **A** be a $mp \times nq$ matrix, we denote with $\mathbf{A}^{[i,j]}$ the special partition of dimension $m \times n$, $\mathbf{A}^{[i,j]} = \sum_{r=1}^{m} \sum_{s=1}^{n} a_{(rp-p+i)(sq-q+j)} \mathbb{E}_{m,n}^{(r,s)}$, for $i=1,\ldots,p$ and $j=1,\ldots,q$.
- Let **A** and **B** be $m \times n$ and $mp \times nq$ matrices such that **B** can be partitioned into mn sub-matrices of dimension $p \times q$. The star product between **A** and **B** is defined as $\mathbf{A} \star \mathbf{B} = \sum_{i=1}^{m} \sum_{j=1}^{n} a_{ij} \mathbf{B}_{ij}^{(p,q)}$, where $\mathbf{B}_{ij}^{(p,q)}$ denotes the ijth block of dimension $p \times q$ of the matrix **B**.
- We denote by $\mathcal{U}(a,b)$ the uniform distribution on the interval (a,b), $\mathcal{N}(\mathbf{A}, \Sigma)$ the normal distribution with mean \mathbf{A} and covariance matrix Σ , and by $\mathcal{MN}_{m,n}(\mathbf{A}, \Sigma, \Omega)$ the $(m \times n)$ matrix normal distribution with mean \mathbf{A} , row covariance Σ , and column covariance Ω .
- We denote as $\nu^{(k)}(\mathbf{A})$ the *k*th largest eigenvalue of a generic squared matrix \mathbf{A} . The matrix norm induced by a vector *p*-norm is denoted as $\|\mathbf{A}\|_p$, with $\|\mathbf{A}\|$ the spectral norm. The Frobenious norm is denoted $\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^n \sum_{j=1}^n a_{ij}} = \mathbf{tr}(\mathbf{A}\mathbf{A}')$. The max norm is denoted as $\|\mathbf{A}\|_{\max} = \max_{ij} a_{ij}$.

• The o_p is for convergence to zero in probability and O_p is for stochastic boundedness. For two random series, X_n and Y_n , $X_n \leq Y_n$ means that $X_n = O_p(Y_n)$, and $X_n \geq Y_n$ means that $Y_n = O_p(X_n)$. The notation $X_n \simeq Y_n$ means that $X_n \leq Y_n$ and $X_n \geq Y_n$.

We make abundant use of the following matrix results. Recall that \mathbb{K}_{nm} is the $(nm \times nm)$ commutation matrix. Let **X**, **Y**, and **Z** be $m \times n$, $n \times p$ and $p \times q$ matrices,

$$\operatorname{vec}(\mathbf{X}\mathbf{Z}\mathbf{Y}) = (\mathbf{Y}' \otimes \mathbf{X})\operatorname{vec}(\mathbf{Z}),$$
 (13)

$$\operatorname{vec}(\mathbf{X}') = \mathbb{K}_{mn}\operatorname{vec}(\mathbf{X}),$$
 (14)

$$\mathbf{XZY} = \mathbf{Z} \star \Big(\operatorname{vec}(\mathbf{X}) \operatorname{vec}(\mathbf{Y}')' \Big), \qquad (15)$$

$$\mathbf{tr}(\mathbf{X}(\mathbf{Y} \circ \mathbf{Z})) = \mathbf{tr}((\mathbf{X}' \circ \mathbf{Y})'\mathbf{Z}).$$
(16)

Let **A** be a $mp \times nq$ matrix. Recall that $\mathbf{A}^{[i,j]} = \sum_{r=1}^{m} \sum_{s=1}^{n} a_{(rp-p+i)(sq-q+j)} \mathbb{E}_{m,n}^{(r,s)}$, we can decompose **A** as follows

$$\mathbf{A} = \sum_{i=1}^{p} \sum_{j=1}^{q} \mathbf{A}^{[i,j]} \otimes \mathbb{E}_{p,q}^{(ij)}.$$
(17)

B Details on the EM algorithm

B.1 Derivation of the expected likelihoods

The expressions of the expected log-likelihoods stated in terms of the data in its original matrix form, which are given in (8) and (9) (up to constant terms and initial conditions), are obtained from the usual expressions for vectorized data as follows (here we consider expectations computed using a generic estimator of the parameters $\hat{\theta}$):

$$\begin{split} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T}|\mathbf{F}_{T};\boldsymbol{\theta})|\mathbf{Y}_{T}] &= -\frac{T}{2}\mathrm{log}(|\mathbf{K}\otimes\mathbf{H}|) - \frac{1}{2}\sum_{t=1}^{T}\mathbb{E}_{\widehat{\boldsymbol{\theta}}}\Big[(\mathbf{y}_{t} - (\mathbf{C}\otimes\mathbf{R})\mathbf{f}_{t})'(\mathbf{K}\otimes\mathbf{H})^{-1}(\mathbf{y}_{t} - (\mathbf{C}\otimes\mathbf{R})\mathbf{f}_{t})|\mathbf{Y}_{T}\Big] \\ &= -\frac{T}{2}(\mathrm{log}(|\mathbf{K}|^{p_{1}}|\mathbf{H}|^{p_{2}})) - \frac{1}{2}\sum_{t=1}^{T}\mathbb{E}_{\widehat{\boldsymbol{\theta}}}\Big[\mathrm{vec}(\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')'(\mathbf{K}^{-1}\otimes\mathbf{H}^{-1})\mathrm{vec}(\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')|\mathbf{Y}_{T}\Big] \\ &= -\frac{T}{2}(\mathrm{log}(|\mathbf{K}|^{p_{1}}|\mathbf{H}|^{p_{2}})) - \frac{1}{2}\sum_{t=1}^{T}\mathbb{E}_{\widehat{\boldsymbol{\theta}}}\Big[\mathrm{vec}(\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')'\mathrm{vec}(\mathbf{H}^{-1}(\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')\mathbf{K}^{-1})|\mathbf{Y}_{T}\Big] \\ &= -\frac{T}{2}(\mathrm{log}(|\mathbf{K}|^{p_{1}}|\mathbf{H}|^{p_{2}})) - \frac{1}{2}\sum_{t=1}^{T}\mathbb{E}_{\widehat{\boldsymbol{\theta}}}\Big[\mathrm{tr}\big(\mathbf{H}^{-1}(\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')\mathbf{K}^{-1}(\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')'\big|\mathbf{Y}_{T}\Big]. \end{split}$$

$$\begin{split} \mathbb{E}_{\widehat{\theta}}[\ell(\mathsf{F}_{T}; \widehat{\theta}) | \mathsf{Y}_{T}] &= -\frac{T-1}{2} \log(|\mathbf{Q} \otimes \mathbf{P}|) - \frac{1}{2} \sum_{t=2}^{T} \mathbb{E}_{\widehat{\theta}} \Big[(\mathsf{f}_{t} - (\mathbf{B} \otimes \mathbf{A}) \mathsf{f}_{t-1})' (\mathbf{Q} \otimes \mathbf{P})^{-1} (\mathsf{f}_{t} - (\mathbf{B} \otimes \mathbf{A}) \mathsf{f}_{t-1}) | \mathsf{Y}_{T} \Big] \\ &= -\frac{T-1}{2} \Big(\log(|\mathbf{Q}|^{k_{1}} | \mathbf{P} |^{k_{2}}) \Big) - \frac{1}{2} \sum_{t=2}^{T} \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{vec}(\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}')' (\mathbf{Q} \otimes \mathbf{P})^{-1} \operatorname{vec}(\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}') | \mathsf{Y}_{T} \Big] \\ &= -\frac{T-1}{2} \Big(\log(|\mathbf{Q}|^{k_{1}} | \mathbf{P} |^{k_{2}}) \Big) - \frac{1}{2} \sum_{t=2}^{T} \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{vec}(\mathbf{f}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}')' \operatorname{vec}(\mathbf{P}^{-1} (\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}') \mathbf{Q}^{-1}) | \mathsf{Y}_{T} \Big] \\ &= -\frac{T-1}{2} \Big(\log(|\mathbf{Q}|^{k_{1}} | \mathbf{P} |^{k_{2}}) \Big) - \frac{1}{2} \sum_{t=2}^{T} \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{tr}(\mathbf{P}^{-1} (\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}') \mathbf{Q}^{-1} (\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}')' | \mathsf{Y}_{T} \Big]. \end{split}$$

B.2 EM updates

To obtain the EM updates we first compute the derivatives of $\mathcal{Q}(\theta, \hat{\theta})$ with respect to each parameter in θ , and obtain the following

$$\begin{split} \frac{\partial \mathcal{Q}(\boldsymbol{\theta}, \widehat{\boldsymbol{\theta}})}{\partial \mathbf{R}} &= \frac{\partial \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T} | \mathbf{F}_{T}; \boldsymbol{\theta}) | \mathbf{Y}_{T}]}{\partial \mathbf{R}} \\ &= \frac{1}{2} \sum_{t=1}^{T} \left(\frac{\partial \mathbf{tr} \left(\mathbf{H}^{-1} \mathbf{Y}_{t} \mathbf{K}^{-1} \mathbf{C} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t}' | \mathbf{Y}_{T}] \mathbf{R}' \right)}{\partial \mathbf{R}} + \frac{\partial \mathbf{tr} \left(\mathbf{H}^{-1} \mathbf{R} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t} | \mathbf{Y}_{T}] \mathbf{C}' \mathbf{K}^{-1} \mathbf{Y}_{t}' \right)}{\partial \mathbf{R}} - \frac{\partial \mathbf{tr} \left(\mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[\mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{R}' | \mathbf{Y}_{T} \right] \right)}{\partial \mathbf{R}} \right) \\ &= \frac{1}{2} \sum_{t=1}^{T} \left(2 \mathbf{H}^{-1} \mathbf{Y}_{t} \mathbf{K}^{-1} \mathbf{C} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t}' | \mathbf{Y}_{T}] - 2 \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[\mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' | \mathbf{Y}_{T} \right] \right), \end{split}$$

$$\begin{split} \frac{\partial \mathcal{Q}(\boldsymbol{\theta}, \widehat{\boldsymbol{\theta}})}{\partial \mathbf{C}} &= \frac{\partial \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T} | \mathbf{F}_{T}; \boldsymbol{\theta}) | \mathbf{Y}_{T}]}{\partial \mathbf{C}} \\ &= \frac{1}{2} \sum_{t=1}^{T} \left(\frac{\partial \mathbf{tr} \left(\mathbf{H}^{-1} \mathbf{Y}_{t} \mathbf{K}^{-1} \mathbf{C} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t}' | \mathbf{Y}_{T}] \mathbf{R}' \right)}{\partial \mathbf{C}} + \frac{\partial \mathbf{tr} \left(\mathbf{H}^{-1} \mathbf{R} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t} | \mathbf{Y}_{T}] \mathbf{C}' \mathbf{K}^{-1} \mathbf{Y}_{t}' \right)}{\partial \mathbf{C}} - \frac{\partial \mathbf{tr} \left(\mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[\mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{R}' | \mathbf{Y}_{T}] \right)}{\partial \mathbf{C}} \right)}{\partial \mathbf{C}} \\ &= \frac{1}{2} \sum_{t=1}^{T} \left(2 \mathbf{K}^{-1} \mathbf{Y}_{t}' \mathbf{H}^{-1} \mathbf{R} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t} | \mathbf{Y}_{T}] - 2 \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[\mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{R}' \mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} | \mathbf{Y}_{T} \right] \right), \end{split}$$

$$\begin{aligned} \frac{\partial \mathcal{Q}(\boldsymbol{\theta}, \widehat{\boldsymbol{\theta}})}{\partial \mathbf{H}} &= \frac{\partial \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T} | \mathbf{F}_{T}; \boldsymbol{\theta}) | \mathbf{Y}_{T}]}{\partial \mathbf{H}} \\ &= -\frac{Tp_{2}}{2} \frac{\partial \log(|\mathbf{H}|)}{\partial \mathbf{H}} - \frac{1}{2} \sum_{t=1}^{T} \frac{\partial \mathbf{tr} \left(\mathbf{H}^{-1} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \mathbf{K}^{-1} (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' | \mathbf{Y}_{T} \right] \right)}{\partial \mathbf{H}} \\ &= -\frac{Tp_{2}}{2} \mathbf{H}^{-1} + \frac{1}{2} \sum_{t=1}^{T} \mathbf{H}^{-1} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \mathbf{K}^{-1} (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' | \mathbf{Y}_{T} \right]' \mathbf{H}^{-1}, \end{aligned}$$

$$\begin{split} \frac{\partial \mathcal{Q}(\boldsymbol{\theta}, \widehat{\boldsymbol{\theta}})}{\partial \mathbf{K}} &= \frac{\partial \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T} | \mathbf{F}_{T}; \boldsymbol{\theta}) | \mathbf{Y}_{T}]}{\partial \mathbf{K}} \\ &= -\frac{Tp_{1}}{2} \frac{\partial \log(|\mathbf{K}|)}{\partial \mathbf{K}} - \frac{1}{2} \sum_{t=1}^{T} \frac{\partial \mathbf{tr} \left(\mathbf{K}^{-1} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \mathbf{H}^{-1} (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') | \mathbf{Y}_{T} \right] \right)}{\partial \mathbf{K}} \\ &= -\frac{Tp_{1}}{2} \mathbf{K}^{-1} + \frac{1}{2} \sum_{t=1}^{T} \mathbf{K}^{-1} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \mathbf{H}^{-1} (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') | \mathbf{Y}_{T} \right]' \mathbf{K}^{-1}, \end{split}$$

$$\begin{aligned} \frac{\partial \mathcal{Q}(\boldsymbol{\theta}, \boldsymbol{\hat{\theta}})}{\partial \mathbf{A}} &= \frac{\partial \mathbb{E}_{\boldsymbol{\hat{\theta}}}[\ell(\mathbf{F}_{T}; \boldsymbol{\theta}) | \mathbf{Y}_{T}]}{\partial \mathbf{A}} \\ &= \frac{1}{2} \sum_{t=2}^{T} \left(\frac{\partial \mathbf{tr}(\mathbb{E}_{\boldsymbol{\hat{\theta}}}[\mathbf{P}^{-1}\mathbf{F}_{t}\mathbf{Q}^{-1}\mathbf{B}\mathbf{F}'_{t-1}\mathbf{A}' | \mathbf{Y}_{T}])}{\partial \mathbf{A}} + \frac{\partial \mathbf{tr}(\mathbb{E}_{\boldsymbol{\hat{\theta}}}[\mathbf{P}^{-1}\mathbf{A}\mathbf{F}_{t-1}\mathbf{B}'\mathbf{Q}^{-1}\mathbf{F}'_{t}| \mathbf{Y}_{T}])}{\partial \mathbf{A}} \\ &- \frac{\partial \mathbf{tr}(\mathbb{E}_{\boldsymbol{\hat{\theta}}}[\mathbf{P}^{-1}\mathbf{A}\mathbf{F}_{t-1}\mathbf{B}'\mathbf{Q}^{-1}\mathbf{B}\mathbf{F}'_{t-1}\mathbf{A}' | \mathbf{Y}_{T}])}{\partial \mathbf{A}} \right) \\ &= \frac{1}{2} \sum_{t=2}^{T} 2 \left(\mathbb{E}_{\boldsymbol{\hat{\theta}}} \Big[\mathbf{P}^{-1}\mathbf{F}_{t}\mathbf{Q}^{-1}\mathbf{B}\mathbf{F}'_{t-1} | \mathbf{Y}_{T} \Big] - 2\mathbb{E}_{\boldsymbol{\hat{\theta}}} \Big[\mathbf{P}^{-1}\mathbf{A}\mathbf{F}_{t-1}\mathbf{B}'\mathbf{Q}^{-1}\mathbf{B}\mathbf{F}'_{t-1} | \mathbf{Y}_{T} \Big] \right), \end{aligned}$$

$$\begin{aligned} \frac{\partial \mathcal{Q}(\theta, \hat{\theta})}{\partial \mathbf{B}} &= \frac{\partial \mathbb{E}_{\hat{\theta}}[\ell(\mathbf{F}_{T}; \theta)|\mathbf{Y}_{T}]}{\partial \mathbf{B}} \\ &= \frac{1}{2} \sum_{t=2}^{T} \left(\frac{\partial \mathbf{tr} \left(\mathbb{E}_{\hat{\theta}} \left[\mathbf{P}^{-1} \mathbf{F}_{t} \mathbf{Q}^{-1} \mathbf{B} \mathbf{F}'_{t-1} \mathbf{A}' | \mathbf{Y}_{T} \right] \right)}{\partial \mathbf{B}} + \frac{\partial \mathbf{tr} \left(\mathbb{E}_{\hat{\theta}} \left[\mathbf{P}^{-1} \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}' \mathbf{Q}^{-1} \mathbf{F}'_{t} | \mathbf{Y}_{T} \right] \right)}{\partial \mathbf{B}} \\ &- \frac{\partial \mathbf{tr} \left(\mathbb{E}_{\hat{\theta}} \left[\mathbf{P}^{-1} \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}' \mathbf{Q}^{-1} \mathbf{B} \mathbf{F}'_{t-1} \mathbf{A}' | \mathbf{Y}_{T} \right] \right)}{\partial \mathbf{B}} \right) \\ &= \frac{1}{2} \sum_{t=2}^{T} \left(2 \mathbb{E}_{\hat{\theta}} \left[\mathbf{Q}^{-1} \mathbf{F}'_{t} \mathbf{P}^{-1} \mathbf{A} \mathbf{F}'_{t-1} | \mathbf{Y}_{T} \right] - 2 \mathbb{E}_{\hat{\theta}} \left[\mathbf{Q}^{-1} \mathbf{B} \mathbf{F}'_{t-1} \mathbf{A}' \mathbf{P}^{-1} \mathbf{A} \mathbf{F}_{t-1} | \mathbf{Y}_{T} \right] \right), \end{aligned}$$

$$\frac{\partial \mathcal{Q}(\theta, \hat{\theta})}{\partial \mathbf{P}} &= \frac{\partial \mathbb{E}_{\hat{\theta}} [\ell(\mathbf{F}_{T}; \theta) | \mathbf{Y}_{T}]}{\partial \mathbf{P}} \\ &= -\frac{(T-1)k_{2}}{2} \frac{\partial \log(|\mathbf{P}|)}{\partial \mathbf{P}} + \frac{1}{2} \sum_{t=2}^{T} \frac{\partial \mathbf{tr} \left(\mathbf{P}^{-1} \mathbb{E}_{\hat{\theta}} \left[(\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}') \mathbf{Q}^{-1} (\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}')' | \mathbf{Y}_{T} \right] \right)}{\partial \mathbf{P}} \\ &= -\frac{(T-1)k_{2}}{2} \mathbf{P}^{-1} + \frac{1}{2} \sum_{t=2}^{T} \mathbf{P}^{-1} \mathbb{E}_{\hat{\theta}} \left[(\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}') \mathbf{Q}^{-1} (\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}')' | \mathbf{Y}_{T} \right]' \mathbf{P}^{-1}, \end{aligned}$$

$$\frac{\partial \mathcal{Q}(\boldsymbol{\theta}, \widehat{\boldsymbol{\theta}})}{\partial \mathbf{Q}} = \frac{\partial \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{F}_{T}; \boldsymbol{\theta}) | \mathbf{Y}_{T}]}{\partial \mathbf{Q}}$$

$$= -\frac{(T-1)k_{1}}{2} \frac{\partial \log(|\mathbf{Q}|)}{\partial \mathbf{Q}} + \frac{1}{2} \sum_{t=2}^{T} \frac{\partial \mathbf{tr}(\mathbf{Q}^{-1} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[(\mathbf{F}_{t} - \mathbf{A}\mathbf{F}_{t-1}\mathbf{B}')'\mathbf{P}^{-1}(\mathbf{F}_{t} - \mathbf{A}\mathbf{F}_{t-1}\mathbf{B}')|\mathbf{Y}_{T}])}{\partial \mathbf{Q}}$$

$$= -\frac{(T-1)k_{1}}{2} \mathbf{Q}^{-1} + \frac{1}{2} \sum_{t=2}^{T} \mathbf{Q}^{-1} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[(\mathbf{F}_{t} - \mathbf{A}\mathbf{F}_{t-1}\mathbf{B}')'\mathbf{P}^{-1}(\mathbf{F}_{t} - \mathbf{A}\mathbf{F}_{t-1}\mathbf{B}')|\mathbf{Y}_{T}]'\mathbf{Q}^{-1}.$$

First order conditions (FOC) then yield

$$\begin{split} \mathbf{R} &= \left(\sum_{t=1}^{T} \mathbf{Y}_{t} \mathbf{K}^{-1} \mathbf{C} \mathbb{E}_{\widehat{\theta}} \left[\mathbf{F}_{t}' | \mathbf{Y}_{T} \right] \right) \left(\sum_{t=1}^{T} \mathbb{E}_{\widehat{\theta}} \left[\mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' | \mathbf{Y}_{T} \right] \right)^{-1}, \\ \mathbf{C} &= \left(\sum_{t=1}^{T} \mathbf{Y}_{t}' \mathbf{H}^{-1} \mathbf{R} \mathbb{E}_{\widehat{\theta}} \left[\mathbf{F}_{t} | \mathbf{Y}_{T} \right] \right) \left(\sum_{t=1}^{T} \mathbb{E}_{\widehat{\theta}} \left[\mathbf{F}_{t}' \mathbf{R}' \mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} | \mathbf{Y}_{T} \right] \right)^{-1}, \\ \mathbf{H} &= \frac{1}{T p_{2}} \sum_{t=1}^{T} \mathbb{E}_{\widehat{\theta}} \left[\left(\mathbf{Y}_{t} - \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right) \mathbf{K}^{-1} \left(\mathbf{Y}_{t} - \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right)' | \mathbf{Y}_{T} \right], \\ \mathbf{K} &= \frac{1}{T p_{1}} \sum_{t=1}^{T} \mathbb{E}_{\widehat{\theta}} \left[\left(\mathbf{Y}_{t} - \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right)' \mathbf{H}^{-1} \left(\mathbf{Y}_{t} - \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right) | \mathbf{Y}_{T} \right], \\ \mathbf{A} &= \left(\sum_{t=1}^{T} \mathbb{E}_{\widehat{\theta}} \left[\mathbf{F}_{t} \mathbf{Q}^{-1} \mathbf{B} \mathbf{F}_{t-1}' | \mathbf{Y}_{T} \right] \right) \left(\sum_{t=1}^{T} \mathbb{E}_{\widehat{\theta}} \left[\mathbf{F}_{t-1} \mathbf{B}' \mathbf{Q}^{-1} \mathbf{B} \mathbf{F}_{t-1}' | \mathbf{Y}_{T} \right] \right)^{-1}, \\ \mathbf{B} &= \left(\sum_{t=1}^{T} \mathbb{E}_{\widehat{\theta}} \left[\mathbf{F}_{t}' \mathbf{P}^{-1} \mathbf{A} \mathbf{F}_{t-1} | \mathbf{Y}_{T} \right] \right) \left(\sum_{t=1}^{T} \mathbb{E}_{\widehat{\theta}} \left[\mathbf{F}_{t-1} \mathbf{A}' \mathbf{P}^{-1} \mathbf{A} \mathbf{F}_{t-1} | \mathbf{Y}_{T} \right] \right)^{-1}, \\ \mathbf{P} &= \frac{1}{(T-1)k_{2}} \sum_{t=2}^{T} \mathbb{E}_{\widehat{\theta}} \left[\left(\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}' \right) \mathbf{Q}^{-1} \left(\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}' \right) | \mathbf{Y}_{T} \right], \\ \mathbf{Q} &= \frac{1}{(T-1)k_{1}} \sum_{t=2}^{T} \mathbb{E}_{\widehat{\theta}} \left[\left(\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}' \right)' \mathbf{P}^{-1} \left(\mathbf{F}_{t} - \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}' \right) | \mathbf{Y}_{T} \right]. \end{split}$$

Using the conditional moments of the Kalman smoother recursions and (13)-(15), we obtain

$$\begin{split} \mathbb{E}_{\widehat{\theta}}[\mathbf{F}_{t}|\mathbf{Y}_{T}] = \operatorname{unvec}\left(\mathbb{E}_{\widehat{\theta}}[f_{t}|\mathbf{Y}_{T}]\right) = \operatorname{unvec}\left(f_{t|T}\right) = \mathbf{F}_{t|T}, \\ \mathbb{E}_{\widehat{\theta}}[\mathbf{F}_{t}(\mathbf{Y}_{T}]] = \operatorname{unvec}\left(\mathbb{E}_{\widehat{\theta}}[f_{t}|\mathbf{Y}_{T}]\right)' = \operatorname{unvec}\left(f_{t|T}\right)' = \mathbf{F}_{t|T}', \\ \mathbb{E}_{\widehat{\theta}}[\mathbf{F}_{t}\mathbf{C}'\mathbf{K}^{-1}\mathbf{C}\mathbf{F}_{t}'|\mathbf{Y}_{T}] = \left(\mathbf{C}'\mathbf{K}^{-1}\mathbf{C}\right) \times \mathbb{E}_{\widehat{\theta}}[\operatorname{tec}(\mathbf{F}_{t})\operatorname{vec}(\mathbf{F}_{t})'|\mathbf{Y}_{T}] \\ &= \left(\mathbf{C}'\mathbf{K}^{-1}\mathbf{C}\right) \times \mathbb{E}_{\widehat{\theta}}[f_{t}'|\mathbf{Y}_{T}] \\ &= \left(\mathbf{C}'\mathbf{K}^{-1}\mathbf{C}\right) \times \mathbb{E}_{\widehat{\theta}}[f_{t}(\mathbf{T}_{t}'|\mathbf{T}] + \mathbf{I}_{t|T}\right), \\ \mathbb{E}_{\widehat{\theta}}[\mathbf{F}_{t}'\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\mathbf{F}_{t}|\mathbf{Y}_{T}] = \left(\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\right) \times \mathbb{E}_{\widehat{\theta}}[\operatorname{vec}(\mathbf{F}_{t}')\operatorname{vec}(\mathbf{F}_{t}')'|\mathbf{Y}_{T}] \\ &= \left(\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\right) \times \left(\mathbb{K}_{k_{1}k_{2}}\mathbb{E}_{\widehat{\theta}}[f_{t}f_{t}'|\mathbf{Y}_{T}]\mathbb{K}_{k_{1}k_{2}}\right) \\ &= \left(\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\right) \times \left(\mathbb{K}_{k_{1}k_{2}}\left(f_{t|T}f_{t}'|\mathbf{T} + \mathbf{I}_{t|T}\right)\mathbb{K}_{k_{1}k_{2}}\right), \\ \mathbb{E}_{\widehat{\theta}}[\mathbf{R}\mathbf{F}_{t}\mathbf{C}'\mathbf{K}^{-1}\mathbf{C}\mathbf{F}_{t}'\mathbf{R}'|\mathbf{Y}_{T}] = \left(\mathbf{C}'\mathbf{K}^{-1}\mathbf{C}\right) \times \mathbb{E}_{\widehat{\theta}}\left[\operatorname{vec}(\mathbf{R}\mathbf{F}_{t})\operatorname{vec}(\mathbf{R}\mathbf{F}_{t}'|\mathbf{Y}_{T}]\right] \\ &= \left(\mathbf{C}'\mathbf{K}^{-1}\mathbf{C}\right) \times \left((I_{k_{2}}\otimes\mathbf{R})\mathbb{E}_{\widehat{\theta}}[f_{t}f_{t}'|\mathbf{Y}_{T}](I_{k_{2}}\otimes\mathbf{R})'\right) \\ &= \left(\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\right) \times \left((I_{k_{1}}\otimes\mathbf{C})\mathbb{E}_{k_{1}k_{2}}\mathbb{E}_{\widehat{\theta}}[f_{t}f_{t}'|\mathbf{Y}_{T}](I_{k_{2}}\otimes\mathbf{R})'\right) \\ &= \left(\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\right) \times \left((I_{k_{1}}\otimes\mathbf{C})\mathbb{E}_{\widehat{\theta}}[f_{t}f_{t}'|\mathbf{Y}_{T}]\mathbb{E}_{k_{1}k_{2}}\right)\left(I_{k_{1}}\otimes\mathbf{C}\right)'\right) \\ &= \left(\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\right) \times \left((I_{k_{1}}\otimes\mathbf{C})\mathbb{E}_{k_{1}k_{2}}\mathbb{E}_{k_{1}k_{2}}\mathbb{E}_{k_{1}k_{2}}\right)(I_{k_{1}}\otimes\mathbf{C}\right)'\right) \\ &= \left(\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\right) \times \left((I_{k_{1}}\otimes\mathbf{C})\mathbb{E}_{\widehat{\theta}}[f_{t}f_{t}'|\mathbf{Y}_{T}]\mathbb{E}_{k_{1}k_{2}}\right)\left(I_{k_{1}}\otimes\mathbf{C}\right)'\right) \\ &= \left(\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}\right) \times \left((I_{k_{1}}\otimes\mathbf{C})\mathbb{E}_{k_{1}k_{2}}\mathbb{E}_{k_{1}k_{2}k_{2}}\mathbb{E}_{k_{1}k_{2}}\mathbb{E}_{k_{1}k_{2}k_{2}}\right)\left(I_{k$$

$$\mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t}\mathbf{Q}^{-1}\mathbf{F}_{t}^{*}|\mathbf{Y}_{T}] = \mathbf{Q}^{-1} \star \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\operatorname{Vec}(\mathbf{F}_{t})\operatorname{Vec}(\mathbf{F}_{t})|\mathbf{Y}_{T}]$$

$$= \mathbf{Q}^{-1} \star \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{f}_{t}\mathbf{f}_{t}^{\prime}|\mathbf{Y}_{T}]$$

$$= \mathbf{Q}^{-1} \star \left(\mathbf{f}_{t|T}\mathbf{f}_{t|T}^{\prime} + \mathbf{\Pi}_{t|T}\right),$$

$$\mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t}^{\prime}\mathbf{P}^{-1}\mathbf{F}_{t}|\mathbf{Y}_{T}] = \mathbf{P}^{-1} \star \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\operatorname{vec}(\mathbf{F}_{t}^{\prime})\operatorname{vec}(\mathbf{F}_{t}^{\prime})^{\prime}|\mathbf{Y}_{T}]$$

$$= \mathbf{P}^{-1} \star \left(\mathbb{K}_{k_{1}k_{2}}\mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{f}_{t}\mathbf{f}_{t}^{\prime}|\mathbf{Y}_{T}]\mathbb{K}_{k_{1}k_{2}}^{\prime}\right)$$

$$= \mathbf{P}^{-1} \star \left(\mathbb{K}_{k_1 k_2} \left(\mathbf{f}_{t|T} \mathbf{f}_{t|T}' + \mathbf{\Pi}_{t|T} \right) \mathbb{K}_{k_1 k_2}' \right),$$

$$\begin{split} \mathbb{E}_{\widehat{\theta}} \Big[\mathbf{F}_{t} \mathbf{Q}^{-1} \mathbf{B} \mathbf{F}'_{t-1} | \mathbf{Y}_{T} \Big] &= (\mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{f}_{t}) \operatorname{vec}(\mathbf{F}_{t-1})' | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(f_{t-1}' | \mathbf{Y}_{T}) \Big] \\ &= (\mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) \operatorname{vec}(\mathbf{F}_{t-1})' | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) \operatorname{vec}(\mathbf{F}_{t-1})' | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \Big] \\ &= (\mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \Big] \\ &= (\mathbf{P}^{-1} \mathbf{A}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{P}^{-1} \mathbf{A}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{E}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{A}' \mathbf{P}^{-1} \mathbf{A}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{C} (\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{A}' \mathbf{P}^{-1} \mathbf{A}) * \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{fv}(\mathbf{C} (\mathbf{F}_{t-1}) | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{A}' \mathbf{P}^{-1} \mathbf{A}) * (\mathbb{E}_{k_{1}k_{2}} \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{ft}_{t-1} | \mathbf{F}_{t-1} | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{A}' \mathbf{P}^{-1} \mathbf{A}) * (\mathbb{E}_{k_{1}k_{2}} \Big[\operatorname{ft}_{t-1} \mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}'_{t-1} | | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{B}_{\widehat{\theta}} \Big[\mathbf{F}_{t-1} \mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}'_{t-1} | | \mathbf{Y}_{T} \Big] \\ &+ \mathbf{A} \mathbb{E}_{\widehat{\theta}} \Big[\mathbf{F}_{t-1} \mathbf{B}' \mathbf{Q}^{-1} \mathbf{B}_{t-1} | \mathbf{Y}_{T} \Big] \mathbf{A}', \\ \\ \\ &= \widehat{\theta}_{\widehat{\theta}} \Big[\mathbf{F}_{t-1} \mathbf{A}' \mathbf{P}^{-1} \mathbf{A} \mathbf{F}_{t-1} | \mathbf{Y}_{T} \Big] \mathbf{B}', \\ \\ &- \mathbf{B} \mathbb{E}_{\widehat{\theta}} \Big[\mathbf{F}_{t-1}' \mathbf{A}' \mathbf{P}^{-1} \mathbf{A} \mathbf{F}_{t-1} | \mathbf{Y}_{T} \Big] \mathbf{B}'. \end{aligned}$$

Combining these results together with the FOC we obtain

$$\begin{split} \mathbf{R} &= \left(\sum_{t=1}^{T} \mathbf{Y}_{t} \mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t|T}'\right) \left(\sum_{t=1}^{T} \left(\mathbf{C}' \mathbf{K}^{-1} \mathbf{C}_{t}\right) \star \left(\mathbf{f}_{t|T} \mathbf{f}_{t|T}' + \mathbf{\Pi}_{t|T}\right)\right)^{-1}, \\ \mathbf{C} &= \left(\sum_{t=1}^{T} \mathbf{Y}_{t}' \mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t|T}\right) \left(\sum_{t=1}^{T} \left(\mathbf{R}' \mathbf{H}^{-1} \mathbf{R}\right) \star \left(\mathbb{K}_{k_{1}k_{2}} \left(\mathbf{f}_{t|T} \mathbf{f}_{t|T}' + \mathbf{\Pi}_{t|T}\right) \mathbb{K}_{k_{1}k_{2}}'\right)\right)^{-1}, \\ \mathbf{H} &= \frac{1}{Tp_{2}} \sum_{t=1}^{T} \left[\mathbf{Y}_{t} \mathbf{K}^{-1} \mathbf{Y}_{t}' - \mathbf{Y}_{t} \mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t|T}' \mathbf{R}' - \mathbf{R} \mathbf{F}_{t|T} \mathbf{C}' \mathbf{K}^{-1} \mathbf{Y}_{t}' \\ &+ \left(\mathbf{C}' \mathbf{K}^{-1} \mathbf{C}\right) \star \left(\left(\mathbb{I}_{k_{2}} \otimes \mathbf{R}\right) \left(\mathbf{f}_{t|T} \mathbf{f}_{t|T}' + \mathbf{\Pi}_{t|T}\right) \left(\mathbb{I}_{k_{2}} \otimes \mathbf{R}\right)'\right)\right], \\ \mathbf{K} &= \frac{1}{Tp_{1}} \sum_{t=1}^{T} \left[\mathbf{Y}_{t}' \mathbf{H}^{-1} \mathbf{Y}_{t} - \mathbf{Y}_{t}' \mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t|T} \mathbf{C}' - \mathbf{C} \mathbf{F}_{t|T}' \mathbf{R}' \mathbf{H}^{-1} \mathbf{Y}_{t} \\ &+ \left(\mathbf{R}' \mathbf{H}^{-1} \mathbf{R}\right) \star \left(\left(\mathbb{I}_{k_{1}} \otimes \mathbf{C}\right) \left(\mathbb{K}_{k_{1}k_{2}} \left(\mathbf{f}_{t|T} \mathbf{f}_{t|T}' + \mathbf{\Pi}_{t|T}\right) \mathbb{K}_{k_{1}k_{2}}'\right) \left(\mathbb{I}_{k_{1}} \otimes \mathbf{C}\right)'\right)\right], \end{split}$$

$$\begin{split} \mathbf{A} &= \left(\sum_{t=1}^{T} \left(\mathbf{Q}^{-1}\mathbf{B}\right) \star \left(f_{t|T}f_{t-1|T}' + \boldsymbol{\Delta}_{t|T}\right)\right) \left(\sum_{t=1}^{T} \left(\mathbf{B}'\mathbf{Q}^{-1}\mathbf{B}\right) \star \left(f_{t-1|T}f_{t-1|T}' + \boldsymbol{\Pi}_{t-1|T}\right)\right)^{-1}, \\ \mathbf{B} &= \left(\sum_{t=1}^{T} \left(\mathbf{P}^{-1}\mathbf{A}\right) \star \left(\mathbb{K}_{k_{1}k_{2}}\left(f_{t|T}f_{t-1|T}' + \boldsymbol{\Delta}_{t|T}\right)\mathbb{K}_{k_{1}k_{2}}'\right)\right) \left(\sum_{t=1}^{T} \left(\mathbf{A}'\mathbf{P}^{-1}\mathbf{A}\right) \star \left(\mathbb{K}_{k_{1}k_{2}}\left(f_{t-1|T}f_{t-1|T}' + \boldsymbol{\Pi}_{t-1|T}\right)\mathbb{K}_{k_{1}k_{2}}'\right)\right)^{-1} \\ \mathbf{P} &= \frac{1}{(T-1)k_{2}}\sum_{t=2}^{T} \left\{\mathbf{Q}^{-1} \star \left(f_{t|T}f_{t|T}' + \boldsymbol{\Pi}_{t|T}\right) - \left[\left(\mathbf{Q}^{-1}\mathbf{B}\right) \star \left(f_{t|T}f_{t-1|T}' + \boldsymbol{\Delta}_{t|T}\right)\right]\mathbf{A}' - \\ \mathbf{A} \left[\left(\mathbf{Q}^{-1}\mathbf{B}\right) \star \left(f_{t|T}f_{t-1|T}' + \boldsymbol{\Delta}_{t|T}\right)\right]' + \mathbf{A} \left[\left(\mathbf{B}'\mathbf{Q}^{-1}\mathbf{B}\right) \star \left(f_{t-1|T}f_{t-1|T}' + \boldsymbol{\Pi}_{t-1|T}\right)\right]\mathbf{A}'\right\}, \\ \mathbf{Q} &= \frac{1}{(T-1)k_{1}}\sum_{t=2}^{T} \left\{\mathbf{P}^{-1} \star \left(\mathbb{K}_{k_{1}k_{2}}\left(f_{t|T}f_{t|T}' + \boldsymbol{\Pi}_{t|T}\right)\mathbb{K}_{k_{1}k_{2}}'\right) - \left[\left(\mathbf{P}^{-1}\mathbf{A}\right) \star \left(\mathbb{K}_{k_{1}k_{2}}\left(f_{t|T}f_{t-1|T}' + \boldsymbol{\Delta}_{t|T}\right)\mathbb{K}_{k_{1}k_{2}}'\right)\right]' \\ &\quad - \mathbf{B} \left[\left(\mathbf{P}^{-1}\mathbf{A}\right) \star \left(\mathbb{K}_{k_{1}k_{2}}\left(f_{t|T}f_{t-1|T}' + \boldsymbol{\Delta}_{t|T}\right)\mathbb{K}_{k_{1}k_{2}}'\right)\right]' \\ &\quad + \mathbf{B} \left[\left(\mathbf{A}'\mathbf{P}^{-1}\mathbf{A}\right) \star \left(\mathbb{K}_{k_{1}k_{2}}\left(f_{t-1|T}f_{t-1|T}' + \boldsymbol{\Pi}_{t-1|T}\right)\mathbb{K}_{k_{1}k_{2}}'\right)\right]\mathbf{B}'\right\}. \end{split}$$

For any iteration $n \ge 0$, given an estimator of the parameters $\hat{\boldsymbol{\theta}}^{(n)}$, the explicit solutions for **R**, **C**, **H**, and **K** obtained from the previous FOCs are given in Section 3.2.3. While the solutions for **A**, **B**, **P**, and **Q** are the following:

$$\begin{aligned} \widehat{\mathbf{A}}^{(n+1)} &= \left(\sum_{t=2}^{T} \left(\widehat{\mathbf{Q}}^{(n)-1} \widehat{\mathbf{B}}^{(n)} \right) \star \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)'} + \mathbf{\Delta}_{t|T}^{(n)} \right) \right) \left(\sum_{t=1}^{T} \left(\widehat{\mathbf{B}}^{(n)'} \widehat{\mathbf{Q}}^{(n)-1} \widehat{\mathbf{B}}^{(n)} \right) \star \left(\mathbf{f}_{t-1|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)'} + \mathbf{\Pi}_{t-1|T}^{(n)} \right) \right)^{-1}, \\ \widehat{\mathbf{B}}^{(n+1)} &= \left(\sum_{t=2}^{T} \left(\widehat{\mathbf{P}}^{(n)-1} \widehat{\mathbf{A}}^{(n+1)} \right) \star \left(\mathbb{K}_{k_{1}k_{2}} \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)'} + \mathbf{\Delta}_{t|T}^{(n)} \right) \mathbb{K}_{k_{1}k_{2}}^{(n)} \right) \right) \\ &\times \left(\sum_{t=1}^{T} \left(\widehat{\mathbf{A}}^{(n+1)'} \widehat{\mathbf{P}}^{(n)-1} \widehat{\mathbf{A}}^{(n+1)} \right) \star \left(\mathbb{K}_{k_{1}k_{2}} \left(\mathbf{f}_{t-1|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)'} + \mathbf{\Pi}_{t-1|T}^{(n)} \right) \mathbb{K}_{k_{1}k_{2}}^{(n)} \right) \right)^{-1}, \end{aligned}$$

$$\begin{split} \widehat{\mathbf{P}}^{(n+1)} &= \frac{1}{(T-1)k_2} \sum_{t=2}^{T} \Big[\widehat{\mathbf{Q}}^{(n)-1} \star \Big(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)} \Big) \\ &- \Big(\Big(\widehat{\mathbf{Q}}^{(n)-1} \widehat{\mathbf{B}}^{(n+1)} \Big) \star \Big(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Delta}_{t|T}^{(n)} \Big) \Big) \widehat{\mathbf{A}}^{(n+1)\prime} \\ &- \widehat{\mathbf{A}}^{(n+1)} \Big(\Big(\widehat{\mathbf{Q}}^{(n)-1} \widehat{\mathbf{B}}^{(n+1)} \Big) \star \Big(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Delta}_{t|T}^{(n)} \Big) \Big)' \\ &+ \widehat{\mathbf{A}}^{(n+1)} \Big(\Big(\widehat{\mathbf{B}}^{(n+1)\prime} \widehat{\mathbf{Q}}^{(n)-1} \widehat{\mathbf{B}}^{(n+1)} \Big) \star \Big(\mathbf{f}_{t-1|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Pi}_{t-1|T}^{(n)} \Big) \Big) \widehat{\mathbf{A}}^{(n+1)\prime} \Big], \end{split}$$

$$\begin{aligned} \widehat{\mathbf{Q}}^{(n+1)} &= \frac{1}{(T-1)k_1} \sum_{t=2}^{T} \left[\widehat{\mathbf{P}}^{(n)-1} \star \left(\mathbb{K}_{k_1 k_2} \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)} \right) \mathbb{K}'_{k_1 k_2} \right) \\ &- \left(\left(\mathbf{P}^{(n)-1} \mathbf{A}^{(n+1)} \right) \star \left(\mathbb{K}_{k_1 k_2} \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Delta}_{t|T}^{(n)} \right) \mathbb{K}'_{k_1 k_2} \right) \right) \widehat{\mathbf{B}}'_{(n+1)} \\ &- \widehat{\mathbf{B}}^{(n+1)} \left(\left(\widehat{\mathbf{P}}^{(n)-1} \widehat{\mathbf{A}}^{(n+1)} \right) \star \left(\mathbb{K}_{k_1 k_2} \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Delta}_{t|T}^{(n)} \right) \mathbb{K}'_{k_1 k_2} \right) \right)' \\ &+ \widehat{\mathbf{B}}^{(n+1)} \left(\left(\widehat{\mathbf{A}}^{(n+1)\prime} \widehat{\mathbf{P}}^{(n)-1} \widehat{\mathbf{A}}^{(n+1)} \right) \star \left(\mathbb{K}_{k_1 k_2} \left(\mathbf{f}_{t-1|T}^{(n)} \mathbf{f}_{t-1|T}^{(n)\prime} + \mathbf{\Pi}_{t-1|T}^{(n)} \right) \mathbb{K}'_{k_1 k_2} \right) \right) \widehat{\mathbf{B}}^{(n+1)\prime} \right] \end{aligned}$$

In practice, estimating the factor matrices using $\widehat{\mathbf{B}}^{(n+1)} \otimes \widehat{\mathbf{A}}^{(n+1)}$ and $\widehat{\mathbf{P}}^{(n+1)} \otimes \widehat{\mathbf{Q}}^{(n+1)}$ or using $\widehat{\mathbf{B}} \otimes \widehat{\mathbf{A}}^{(n+1)}$ and $\widehat{\mathbf{P} \otimes \mathbf{Q}}^{(n+1)}$, as given in Section 3.2.3, does not make any appreciable difference, so for ease of computation we suggest to use the latter.

B.3 EM updates with missing observations

Let \mathbf{W}_t be a $p_1 \times p_2$ matrix with ones corresponding to the non-missing entries in \mathbf{Y}_t and zeros otherwise. Decomposing \mathbf{Y}_t as follows

$$\mathbf{Y}_t = \mathbf{W}_t \circ \mathbf{Y}_t + (\mathbf{1}_{p_1, p_2} - \mathbf{W}_t) \circ \mathbf{Y}_t,$$

we can write

$$\begin{aligned} \operatorname{tr} \left(\mathbf{H}^{-1} (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}') \mathbf{K}^{-1} (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')' \right) \\ &= \operatorname{tr} \left(\mathbf{H}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}'))' \right) \\ &+ \operatorname{tr} \left(\operatorname{H}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')) \mathbf{K}^{-1} ((\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}'))' \right) \\ &+ \operatorname{tr} \left(\operatorname{H}^{-1} ((\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}'))' \right) \\ &+ \operatorname{tr} \left(\operatorname{H}^{-1} (((\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')) \mathbf{K}^{-1} ((\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}'))' \right). \end{aligned}$$
(18)

Moreover, by the law of iterated expectations, we have that

$$\mathbb{E}_{\widehat{\theta}}\left[\operatorname{tr}\left(\mathbf{H}^{-1}\left(\mathbf{Y}_{t}-\mathbf{R}\mathbf{F}_{t}\mathbf{C}'\right)\mathbf{K}^{-1}\left(\mathbf{Y}_{t}-\mathbf{R}\mathbf{F}_{t}\mathbf{C}'\right)'\right)|\mathbf{Y}_{T}\right] = \mathbb{E}_{\widehat{\theta}}\left[\mathbb{E}_{\widehat{\theta}}\left[\operatorname{tr}\left(\mathbf{H}^{-1}\left(\mathbf{Y}_{t}-\mathbf{R}\mathbf{F}_{t}\mathbf{C}'\right)\mathbf{K}^{-1}\left(\mathbf{Y}_{t}-\mathbf{R}\mathbf{F}_{t}\mathbf{C}'\right)'\right)|\mathbf{F}_{t},\mathbf{Y}_{T}\right]\right].$$
(19)

Using the properties of Hadamard product and (16), we obtain

$$\mathbb{E}_{\widehat{\theta}} \left[\operatorname{tr} \left(\mathbf{H}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')) \mathbf{K}^{-1} ((\mathbf{1}_{p_{1}, p_{2}} - \mathbf{W}_{t}) \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}'))' \right) | \mathbf{F}_{t}, \mathbf{Y}_{T} \right] \\
= \mathbb{E}_{\widehat{\theta}} \left[\operatorname{tr} \left(\left(\mathbf{W}_{t} \circ (\mathbf{H}^{-1} (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')) \mathbf{K}^{-1} \right) \left((\mathbf{1}_{p_{1}, p_{2}} - \mathbf{W}_{t})' \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \right) \right) | \mathbf{F}_{t}, \mathbf{Y}_{T} \right] \\
= \mathbb{E}_{\widehat{\theta}} \left[\operatorname{tr} \left(\left(\mathbf{W}_{t}' \circ (\mathbf{K}^{-1} (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \mathbf{H}^{-1} \right) \circ (\mathbf{1}_{p_{1}, p_{2}} - \mathbf{W}_{t})' \right) (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \right) | \mathbf{F}_{t}, \mathbf{Y}_{T} \right] \\
= \mathbb{E}_{\widehat{\theta}} \left[\operatorname{tr} \left(\left(\mathbf{0}_{p_{1}, p_{2}} \circ (\mathbf{K}^{-1} (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \mathbf{H}^{-1} \right) \right) (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \right) | \mathbf{F}_{t}, \mathbf{Y}_{T} \right] = 0, \tag{20}$$

and

$$\begin{split} & \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{tr} \big(\mathbf{H}^{-1} ((\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')) \mathbf{K}^{-1} ((\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}'))' \big) | \mathbf{F}_{t}, \mathbf{Y}_{T} \Big] \\ &= \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{tr} \big(\big((\mathbf{H}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t})' \mathbf{H}^{-1}) \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \circ (\mathbf{K}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t})'))' (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \big) | \mathbf{F}_{t}, \mathbf{Y}_{T} \Big] \\ &= \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{tr} \big(\big((\mathbf{H}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \mathbf{K}^{-1}) \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \big) (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \big) | \mathbf{F}_{t}, \mathbf{Y}_{T} \Big] \\ &= \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{tr} \big(\big((\mathbf{H}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \mathbf{K}^{-1}) \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \big) (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')' \big) | \mathbf{F}_{t}, \mathbf{Y}_{T} \Big] \\ &= \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{tr} \big(\big((\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \big) \big(\mathbf{H}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \mathbf{K}^{-1} \big) \big) | \mathbf{F}_{t}, \mathbf{Y}_{T} \Big] \\ &= \mathbb{E}_{\widehat{\theta}} \Big[\operatorname{tr} \big(\big((\mathbf{H}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \mathbf{K}^{-1} \big) \mathbb{E}_{\widehat{\theta}} \Big[\big((\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \big) | \mathbf{F}_{t}, \mathbf{Y}_{T} \Big] \\ &= \operatorname{tr} \big(\big(\mathbf{H}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \mathbf{K}^{-1} \big) \mathbb{E}_{\widehat{\theta}} \Big[\big((\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}') \big) ' | \mathbf{F}_{t}, \mathbf{Y}_{T} \Big] \big) \\ &= \operatorname{tr} \Big(\big(\mathbf{H}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t}) \mathbf{K}^{-1} \big) \Big(\widehat{\mathbf{H}} \mathbf{1}_{p_{1},p_{2}} \widehat{\mathbf{K}} \Big)' \Big). \end{split}$$

Combining (18), (19), (20) and (21) we get, up to constant terms, that

$$\begin{split} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T}|\mathbf{F}_{T};\boldsymbol{\theta})|\mathbf{Y}_{T}] &= -\frac{T}{2}(p_{1}\log(|\mathbf{K}|) + p_{2}\log(|\mathbf{H}|)) \\ &- \frac{1}{2}\sum_{t=1}^{T}\mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[\mathbf{tr}\left(\mathbf{H}^{-1}(\mathbf{W}_{t}\circ(\mathbf{Y}_{t}-\mathbf{RF}_{t}\mathbf{C}'))\mathbf{K}^{-1}(\mathbf{W}_{t}\circ(\mathbf{Y}_{t}-\mathbf{RF}_{t}\mathbf{C}'))'\right)|\mathbf{Y}_{T}\right] \\ &- \frac{1}{2}\sum_{t=1}^{T}\mathbf{tr}\left(\mathbf{H}^{-1}(\mathbf{1}_{p_{1},p_{2}}-\mathbf{W}_{t})\mathbf{K}^{-1}\left(\widehat{\mathbf{H}}\mathbf{1}_{p_{1},p_{2}}\widehat{\mathbf{K}}\right)'\right). \end{split}$$

Therefore, the EM updates must be modified accordingly. We first compute derivatives of $\mathbb{E}_{\hat{\theta}}[\ell(\mathbf{Y}_T | \mathbf{F}_T; \boldsymbol{\theta}) | \mathbf{Y}_T]$ with respect to **R**, **C**, **H**, and **K**, obtaining

$$\begin{split} \frac{\partial \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T}|\mathbf{F}_{T};\boldsymbol{\theta})|\mathbf{Y}_{T}]}{\partial \mathbf{R}} &= \frac{1}{2} \sum_{t=1}^{T} \left(2 \frac{\partial \mathbf{tr} \left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \left(\mathbf{Y}_{t}' \circ \mathbf{C} \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\mathbf{F}_{t}'|\mathbf{Y}_{T}] \mathbf{R}' \right) \right)}{\partial \mathbf{R}} - \frac{\partial \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\mathbf{tr} \left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} (\mathbf{C} \mathbf{F}_{t}' \mathbf{R}' \circ \mathbf{C} \mathbf{F}_{t}' \mathbf{R}') \right) |\mathbf{Y}_{T}]}{\partial \mathbf{R}} \right) \\ &= \frac{1}{2} \sum_{t=1}^{T} \left(2 \frac{\partial \mathbf{tr} \left(\left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{Y}_{t} \right) \mathbf{C} \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\mathbf{F}_{t}'|\mathbf{Y}_{T}] \mathbf{R}' \right)}{\partial \mathbf{R}} - \frac{\partial \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\mathbf{tr} \left(\left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right) \mathbf{C} \mathbf{F}_{t}' \mathbf{R}' \right) |\mathbf{Y}_{T}]}{\partial \mathbf{R}} \right) \\ &= \frac{1}{2} \sum_{t=1}^{T} \left(2 \left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{Y}_{t} \right) \mathbf{C} \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\mathbf{F}_{t}'|\mathbf{Y}_{T}] - 2 \mathbb{E}_{\hat{\boldsymbol{\theta}}} \left[\left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right) \mathbf{C} \mathbf{F}_{t}' |\mathbf{Y}_{T} \right] \right), \end{split}$$

$$\begin{split} \frac{\partial \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T}|\mathbf{F}_{T};\boldsymbol{\theta})|\mathbf{Y}_{T}]}{\partial \mathbf{C}} &= \frac{1}{2} \sum_{t=1}^{T} \left(2 \frac{\partial \mathbf{tr} \left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \left(\mathbf{Y}_{t}' \circ \mathbf{C} \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\mathbf{F}_{t}'|\mathbf{Y}_{T}] \mathbf{R}' \right) \right)}{\partial \mathbf{R}} - \frac{\partial \mathbb{E}_{\hat{\boldsymbol{\theta}}} \left[\mathbf{tr} \left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} (\mathbf{C} \mathbf{F}_{t}' \mathbf{R}' \circ \mathbf{C} \mathbf{F}_{t}' \mathbf{R}') \right) |\mathbf{Y}_{T} \right]}{\partial \mathbf{C}} \right) \\ &= \frac{1}{2} \sum_{t=1}^{T} \left(2 \frac{\partial \mathbf{tr} \left(\left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{Y}_{t} \right) \mathbf{C} \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\mathbf{F}_{t}'|\mathbf{Y}_{T}] \mathbf{R}' \right)}{\partial \mathbf{C}} - \frac{\partial \mathbb{E}_{\hat{\boldsymbol{\theta}}} \left[\mathbf{tr} \left(\left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right) \mathbf{C} \mathbf{F}_{t}' \mathbf{R}' \right) |\mathbf{Y}_{T} \right]}{\partial \mathbf{C}} \right) \\ &= \frac{1}{2} \sum_{t=1}^{T} \left(2 \left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{Y}_{t} \right)' \mathbf{R} \mathbb{E}_{\hat{\boldsymbol{\theta}}} \left[\mathbf{F}_{t} |\mathbf{Y}_{T} \right] - 2 \mathbb{E}_{\hat{\boldsymbol{\theta}}} \left[\left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right)' \mathbf{R} \mathbf{F}_{t} |\mathbf{Y}_{T} \right] \right), \end{split}$$

$$\begin{split} & \frac{\partial \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T}|\mathbf{F}_{T};\boldsymbol{\theta})|\mathbf{Y}_{T}]}{\partial \mathbf{H}} \\ = & -\frac{T_{p_{2}}}{2} \frac{\partial \log(|\mathbf{H}|)}{\partial \mathbf{H}} - \frac{1}{2} \sum_{t=1}^{T} \frac{\partial \mathbf{tr} \left(\mathbf{H}^{-1} \left(\mathbb{E}_{\hat{\boldsymbol{\theta}}}[(\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t}\mathbf{C}'))\mathbf{K}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t}\mathbf{C}'))'|\mathbf{Y}_{T}] + (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t})\mathbf{K}^{-1} (\hat{\mathbf{H}}\mathbf{1}_{p_{1},p_{2}}\hat{\mathbf{K}})')\right)}{\partial \mathbf{H}} \\ = & -\frac{T_{p_{2}}}{2} \mathbf{H}^{-1} + \frac{1}{2} \sum_{t=1}^{T} \mathbf{H}^{-1} \left(\mathbb{E}_{\hat{\boldsymbol{\theta}}} \left[(\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t}\mathbf{C}'))\mathbf{K}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t}\mathbf{C}'))'|\mathbf{Y}_{T} \right]' \\ & + \left(\widehat{\mathbf{H}}\mathbf{1}_{p_{1},p_{2}}\widehat{\mathbf{K}} \right) \mathbf{K}^{-1} (\mathbf{1}_{p_{1},p_{2}} - \mathbf{W}_{t})' \right) \mathbf{H}^{-1}, \end{split}$$

$$\begin{split} & \frac{\partial \mathbb{E}_{\hat{\boldsymbol{\theta}}}[\ell(\mathbf{Y}_{T}|\mathbf{F}_{T};\boldsymbol{\theta})|\mathbf{Y}_{T}]}{\partial \mathbf{K}} \\ &= -\frac{Tp_{1}}{2} \frac{\partial \log(|\mathbf{K}|)}{\partial \mathbf{K}} - \frac{1}{2} \sum_{t=1}^{T} \frac{\partial \mathbf{tr} \left(\mathbf{K}^{-1} \left(\mathbb{E}_{\hat{\boldsymbol{\theta}}} \left[(\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}'))'\mathbf{H}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}'))|\mathbf{Y}_{T} \right] + \left(\hat{\mathbf{H}} \mathbf{1}_{p_{1}, p_{2}} \hat{\mathbf{K}} \right)' \mathbf{H}^{-1} \left(\mathbf{1}_{p_{1}, p_{2}} - \mathbf{W}_{t} \right) \right) \right)}{\partial \mathbf{H}} \\ &= -\frac{Tp_{1}}{2} \mathbf{K}^{-1} + \frac{1}{2} \sum_{t=1}^{T} \mathbf{K}^{-1} \left(\mathbb{E}_{\hat{\boldsymbol{\theta}}} \left[(\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}'))'\mathbf{H}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t} \mathbf{C}')) |\mathbf{Y}_{T} \right]' \\ &+ (\mathbf{1}_{p_{1}, p_{2}} - \mathbf{W}_{t})'\mathbf{H}^{-1} \left(\hat{\mathbf{H}} \mathbf{1}_{p_{1}, p_{2}} \hat{\mathbf{K}} \right) \right) \mathbf{K}^{-1}. \end{split}$$

The FOC for ${\bf R}$ then yields

$$\sum_{t=1}^{T} \left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{Y}_{t} \right) \mathbf{C} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[\mathbf{F}_{t}' | \mathbf{Y}_{T} \right] = \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[\left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right) \mathbf{C} \mathbf{F}_{t}' | \mathbf{Y}_{T} \right],$$

from which we obtain

$$\begin{split} \sum_{t=1}^{T} \operatorname{vec}\left(\left(\mathbf{W}_{t} \circ \mathbf{H}^{-1} \mathbf{Y}_{t} \mathbf{K}^{-1}\right) \mathbf{C} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t}'|\mathbf{Y}_{T}]\right) &= \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[\operatorname{vec}\left(\left(\mathbf{W}_{t} \circ \mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1}\right) \mathbf{C} \mathbf{F}_{t}'\right)|\mathbf{Y}_{T}\right] \\ &= \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[\left(\mathbf{F}_{t} \mathbf{C}' \otimes \mathbb{I}_{p_{1}}\right) \operatorname{vec}\left(\left(\mathbf{W}_{t} \circ \mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1}\right) \mathbf{C} \mathbf{F}_{t}'\right)|\mathbf{Y}_{T}\right] \\ &= \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[\left(\mathbf{F}_{t} \mathbf{C}' \otimes \mathbb{I}_{p_{1}}\right) \mathbb{D}_{\mathbf{W}_{t}} \operatorname{vec}\left(\mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1}\right)|\mathbf{Y}_{T}\right] \\ &= \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[\left(\mathbf{F}_{t} \mathbf{C}' \otimes \mathbb{I}_{p_{1}}\right) \mathbb{D}_{\mathbf{W}_{t}}\left(\mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \otimes \mathbf{H}^{-1}\right)|\mathbf{Y}_{T}\right] \operatorname{vec}(\mathbf{R}), \end{split}$$

implying that

$$\operatorname{vec}(\mathbf{R}) = \left(\sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[(\mathbf{F}_{t} \mathbf{C}' \otimes \mathbb{I}_{p_{1}}) \mathbb{D}_{\mathbf{W}_{t}} \left(\mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \otimes \mathbf{H}^{-1} \right) | \mathbf{Y}_{T} \right] \right)^{-1} \left(\sum_{t=1}^{T} \operatorname{vec}\left(\left(\mathbf{W}_{t} \circ \mathbf{H}^{-1} \mathbf{Y}_{t} \mathbf{K}^{-1} \right) \mathbf{C} \mathbb{E}_{\widehat{\boldsymbol{\theta}}}[\mathbf{F}_{t}' | \mathbf{Y}_{T}] \right) \right).$$

The FOC for ${\bf C}$ yields

$$\sum_{t=1}^{T} \left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{Y}_{t} \right)' \mathbf{R} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} [\mathbf{F}_{t} | \mathbf{Y}_{T}] = \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[\left(\mathbf{H}^{-1} \mathbf{W}_{t} \mathbf{K}^{-1} \circ \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \right)' \mathbf{R} \mathbf{F}_{t} | \mathbf{Y}_{T} \right],$$

from which we obtain

$$\begin{split} \sum_{t=1}^{T} \operatorname{vec} \left(\left(\mathbf{W}_{t} \circ \mathbf{H}^{-1} \mathbf{Y}_{t} \mathbf{K}^{-1} \right)' \mathbf{R} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} [\mathbf{F}_{t} | \mathbf{Y}_{T}] \right) &= \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[\operatorname{vec} \left(\left(\mathbf{W}_{t} \circ \mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1} \right)' \mathbf{R} \mathbf{F}_{t} \right) | \mathbf{Y}_{T} \right] \\ &= \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{F}_{t}' \mathbf{R}' \otimes \mathbb{I}_{p_{2}}) \operatorname{vec} \left(\left(\mathbf{W}_{t} \circ \mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \mathbf{C}' \mathbf{K}^{-1} \right)' \mathbf{R} \mathbf{F}_{t} \right) | \mathbf{Y}_{T} \right] \\ &= \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{F}_{t}' \mathbf{R}' \otimes \mathbb{I}_{p_{2}}) \mathbb{D}_{\mathbf{W}_{t}'} \operatorname{vec} \left(\mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{R}' \mathbf{H}^{-1} \right) | \mathbf{Y}_{T} \right] \\ &= \sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{F}_{t}' \mathbf{R}' \otimes \mathbb{I}_{p_{2}}) \mathbb{D}_{\mathbf{W}_{t}'} \left(\mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \otimes \mathbf{K}^{-1} \right) | \mathbf{Y}_{T} \right] \operatorname{vec}(\mathbf{C}), \end{split}$$

implying that

$$\operatorname{vec}(\mathbf{C}) = \left(\sum_{t=1}^{T} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{F}_{t}' \mathbf{R}' \otimes \mathbb{I}_{p_{2}}) \mathbb{D}_{\mathbf{W}_{t}'} (\mathbf{H}^{-1} \mathbf{R} \mathbf{F}_{t} \otimes \mathbf{K}^{-1}) | \mathbf{Y}_{T} \right] \right)^{-1} \left(\sum_{t=1}^{T} \operatorname{vec} \left((\mathbf{W}_{t} \circ \mathbf{H}^{-1} \mathbf{Y}_{t} \mathbf{K}^{-1})' \mathbf{R} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} [\mathbf{F}_{t} | \mathbf{Y}_{T}] \right) \right).$$

The FOC for ${\bf H}$ yields

$$\frac{Tp_2}{2}\mathbf{H}^{-1} = \frac{1}{2}\sum_{t=1}^{T}\mathbf{H}^{-1} \Big(\mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{W}_t \circ (\mathbf{Y}_t - \mathbf{RF}_t \mathbf{C}'))\mathbf{K}^{-1} (\mathbf{W}_t \circ (\mathbf{Y}_t - \mathbf{RF}_t \mathbf{C}'))' | \mathbf{Y}_T \Big]' \\ + \Big(\widehat{\mathbf{H}} \mathbf{1}_{p_1, p_2} \widehat{\mathbf{K}} \Big) \mathbf{K}^{-1} (\mathbf{1}_{p_1, p_2} - \mathbf{W}_t)' \Big) \mathbf{H}^{-1}$$

implying that, for $i=1,\ldots,p_1$,

$$[\mathbf{H}]_{ii} = \left[\frac{\sum_{t=1}^{T} \left(\mathbb{E}_{\widehat{\theta}}\left[(\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t}\mathbf{C}'))\mathbf{K}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t}\mathbf{C}'))' | \mathbf{Y}_{T}\right]' + \left(\widehat{\mathbf{H}}\mathbf{1}_{p_{1}, p_{2}}\widehat{\mathbf{K}}\right)\mathbf{K}^{-1} \left(\mathbf{1}_{p_{1}, p_{2}} - \mathbf{W}_{t}\right)'\right)}{Tp_{2}}\right]_{ii}.$$

The FOC for ${\bf K}$ yields

$$\begin{split} \frac{Tp_1}{2}\mathbf{K}^{-1} &= \frac{1}{2}\sum_{t=1}^{T}\mathbf{K}^{-1} \left(\mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{W}_t \circ (\mathbf{Y}_t - \mathbf{RF}_t \mathbf{C}'))' \mathbf{H}^{-1} (\mathbf{W}_t \circ (\mathbf{Y}_t - \mathbf{RF}_t \mathbf{C}')) | \mathbf{Y}_T \right]' \\ &+ (\mathbf{1}_{p_1, p_2} - \mathbf{W}_t)' \mathbf{H}^{-1} \left(\widehat{\mathbf{H}} \mathbf{1}_{p_1, p_2} \widehat{\mathbf{K}} \right) \right) \mathbf{K}^{-1}, \end{split}$$

implying that, for $i=1,\ldots,p_2$,

$$[\mathbf{K}]_{ii} = \left[\frac{\sum_{t=1}^{T} \left(\mathbb{E}_{\widehat{\boldsymbol{\theta}}}\left[(\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t}\mathbf{C}'))'\mathbf{H}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{RF}_{t}\mathbf{C}'))|\mathbf{Y}_{T}\right]' + \left(\mathbf{1}_{p_{1}, p_{2}} - \mathbf{W}_{t}\right)'\mathbf{H}^{-1} \left(\widehat{\mathbf{H}}\mathbf{1}_{p_{1}, p_{2}}\widehat{\mathbf{K}}\right)\right)}{Tp_{1}}\right]_{ii}.$$

Using the conditional moments computed with the Kalman recursions and (17), we obtain

$$\begin{split} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{F}_{t} \mathbf{C}' \otimes \mathbb{I}_{p_{1}}) \mathbb{D}_{\mathbf{W}_{t}} \left(\mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \otimes \mathbf{H}^{-1} \right) | \mathbf{Y}_{T} \Big] \\ &= \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \left[(\mathbf{F}_{t} \mathbf{C}' \otimes \mathbb{I}_{p_{1}}) \left(\sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{1}} \mathbb{D}_{\mathbf{W}_{t}}^{[s,q]} \otimes \mathbb{E}_{p_{1},p_{1}}^{(s,q)} \right) \left(\mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \otimes \mathbf{H}^{-1} \right) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{1}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{F}_{t} \mathbf{C}' \otimes \mathbb{I}_{p_{1}}) \left(\mathbb{D}_{\mathbf{W}_{t}}^{[s,q]} \otimes \mathbb{E}_{p_{1},p_{1}}^{(s,q)} \right) \left(\mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \otimes \mathbf{H}^{-1} \right) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{1}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\left(\mathbf{F}_{t} \mathbf{C}' \mathbb{D}_{\mathbf{W}_{t}}^{[s,q]} \otimes \mathbb{I}_{p_{1}} \mathbb{E}_{p_{1},p_{1}}^{(s,q)} \right) \left(\mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \otimes \mathbf{H}^{-1} \right) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{1}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\left(\mathbf{F}_{t} \mathbf{C}' \mathbb{D}_{\mathbf{W}_{t}}^{[s,q]} \otimes \mathbb{I}_{p_{1}} \mathbb{E}_{p_{1},p_{1}}^{(s,q)} \mathbf{H}^{-1} \right) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{1}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\left(\mathbf{C}' \mathbb{D}_{\mathbf{W}_{t}}^{[s,q]} \mathbf{K}^{-1} \mathbf{C} \mathbf{F}_{t}' \otimes \mathbb{I}_{p_{1}} \mathbb{E}_{p_{1},p_{1}}^{(s,q)} \mathbf{H}^{-1} \right) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{1}} \sum_{q=1}^{p_{1}} \left(\left(\mathbf{C}' \mathbb{D}_{\mathbf{W}_{t}}^{[s,q]} \mathbf{K}^{-1} \mathbf{C} \right) \star \mathbb{E}_{\widehat{\boldsymbol{\theta}}} [f_{t} f_{t}' | \mathbf{Y}_{T}] \right) \otimes \left(\mathbb{I}_{p_{1}} \mathbb{E}_{p_{1},p_{1}}^{(s,q)} \mathbf{H}^{-1} \right) \\ &= \sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{1}} \left(\left(\mathbf{C}' \mathbb{D}_{\mathbf{W}_{t}}^{[s,q]} \mathbf{K}^{-1} \mathbf{C} \right) \star \left(f_{t|T} f_{t|T}' + \mathbf{\Pi}_{t|T} \right) \right) \otimes \left(\mathbb{I}_{p_{1}} \mathbb{E}_{p_{1},p_{1}}^{(s,q)} \mathbf{H}^{-1} \right) , \end{split}$$

$$\begin{split} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{F}_{t}'\mathbf{R}' \otimes \mathbb{I}_{p_{2}}) \mathbb{D}_{\mathbf{W}_{t}'} (\mathbf{H}^{-1}\mathbf{R}\mathbf{F}_{t} \otimes \mathbf{K}^{-1}) | \mathbf{Y}_{T} \Big] \\ &= \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{F}_{t}'\mathbf{R}' \otimes \mathbb{I}_{p_{2}}) \left(\sum_{s=1}^{p_{2}} \sum_{q=1}^{p_{2}} \mathbb{D}_{\mathbf{W}_{t}'}^{[s,q]} \otimes \mathbb{E}_{p_{2},p_{2}}^{(s,q)} \right) (\mathbf{H}^{-1}\mathbf{R}\mathbf{F}_{t} \otimes \mathbf{K}^{-1}) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{2}} \sum_{q=1}^{p_{2}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{F}_{t}'\mathbf{R}' \otimes \mathbb{I}_{p_{2}}) \left(\mathbb{D}_{\mathbf{W}_{t}'}^{[s,q]} \otimes \mathbb{E}_{p_{2},p_{2}}^{(s,q)} \right) (\mathbf{H}^{-1}\mathbf{R}\mathbf{F}_{t} \otimes \mathbf{K}^{-1}) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{2}} \sum_{q=1}^{p_{2}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\left(\mathbf{F}_{t}'\mathbf{R}' \mathbb{D}_{\mathbf{W}_{t}'}^{[s,q]} \otimes \mathbb{I}_{p_{2}} \mathbb{E}_{p_{2},p_{2}}^{(s,q)} \right) (\mathbf{H}^{-1}\mathbf{R}\mathbf{F}_{t} \otimes \mathbf{K}^{-1}) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{2}} \sum_{q=1}^{p_{2}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\left(\mathbf{F}_{t}'\mathbf{R}' \mathbb{D}_{\mathbf{W}_{t}'}^{[s,q]} + \mathbb{I}^{-1}\mathbf{R}\mathbf{F}_{t} \otimes \mathbb{I}_{p_{2}} \mathbb{E}_{p_{2},p_{2}}^{(s,q)} \mathbf{K}^{-1} \right) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{2}} \sum_{q=1}^{p_{2}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\left(\mathbf{F}_{t}'\mathbf{R}' \mathbb{D}_{\mathbf{W}_{t}'}^{[s,q]} + \mathbb{I}^{-1}\mathbf{R}\mathbf{F}_{t} \otimes \mathbb{I}_{p_{2}} \mathbb{E}_{p_{2},p_{2}}^{(s,q)} \mathbf{K}^{-1} \right) | \mathbf{Y}_{T} \Big] \\ &= \sum_{s=1}^{p_{2}} \sum_{q=1}^{p_{2}} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\left(\mathbf{R}' \mathbb{D}_{\mathbf{W}_{t}'}^{[s,q]} + \mathbb{I}^{-1}\mathbf{R}\mathbf{K} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\operatorname{vec}(\mathbf{F}_{t}') \operatorname{vec}(\mathbf{F}_{t}') | \mathbf{Y}_{T} \Big] \right) \otimes \Big(\mathbb{I}_{p_{2}} \mathbb{E}_{p_{2},p_{2}}^{(s,q)} \mathbf{K}^{-1} \Big) \\ &= \sum_{s=1}^{p_{2}} \sum_{q=1}^{p_{2}} \left(\mathbf{R}' \mathbb{D}_{\mathbf{W}_{t}'}^{[s,q]} + \mathbb{I}^{-1}\mathbf{R}\mathbf{K} \Big(\mathbb{K}_{k_{1}k_{2}} \Big(\mathbf{f}_{t|T}\mathbf{f}_{t|T}' + \mathbf{\Pi}_{t|T} \Big) \mathbb{K}_{k_{1}k_{2}}' \Big) \Big) \otimes \Big(\mathbb{I}_{p_{2}} \mathbb{E}_{p_{2},p_{2}}^{(s,q)} \mathbf{K}^{-1} \Big). \end{aligned}$$

Moreover,

$$\begin{split} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}') \mathbf{K}_t^{-1} (\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}')' | \mathbf{Y}_T \Big] \\ &= \mathbf{K}_t^{-1} \star \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\operatorname{vec} (\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}') \operatorname{vec} (\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}')' | \mathbf{Y}_T \Big] \\ &= \mathbf{K}_t^{-1} \star \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\mathbb{D}_{\mathbf{W}_t} \operatorname{vec} (\mathbf{R} \mathbf{F}_t \mathbf{C}') \operatorname{vec} (\mathbf{R} \mathbf{F}_t \mathbf{C}')' \mathbb{D}_{\mathbf{W}_t}' | \mathbf{Y}_T \Big] \\ &= \mathbf{K}_t^{-1} \star (\mathbb{D}_{\mathbf{W}_t} (\mathbf{C} \otimes \mathbf{R}) \mathbb{E}_{\widehat{\boldsymbol{\theta}}} [f_t f_t' | \mathbf{Y}_T] (\mathbf{C} \otimes \mathbf{R})' \mathbb{D}_{\mathbf{W}_t}') \\ &= \mathbf{K}_t^{-1} \star \Big(\mathbb{D}_{\mathbf{W}_t} (\mathbf{C} \otimes \mathbf{R}) \Big(f_{t|T} f_{t|T}' + \mathbf{\Pi}_{t|T} \Big) (\mathbf{C} \otimes \mathbf{R})' \mathbb{D}_{\mathbf{W}_t}' \Big), \end{split}$$

$$\begin{split} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}')' \mathbf{H}^{-1} (\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}') | \mathbf{Y}_T \Big] \\ &= \mathbf{H}^{-1} \star \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\operatorname{vec} \left((\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}')' \right) \operatorname{vec} \left((\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}')' \right)' | \mathbf{Y}_T \Big] \\ &= \mathbf{H}^{-1} \star \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\mathbb{D}_{\mathbf{W}'_t} \operatorname{vec} (\mathbf{C} \mathbf{F}'_t \mathbf{R}') \operatorname{vec} (\mathbf{C} \mathbf{F}'_t \mathbf{R}')' \mathbb{D}'_{\mathbf{W}'_t} | \mathbf{Y}_T \Big] \\ &= \mathbf{H}^{-1} \star \Big(\mathbb{D}_{\mathbf{W}'_t} (\mathbf{R} \otimes \mathbf{C}) \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[\operatorname{vec} (\mathbf{F}'_t) \operatorname{vec} (\mathbf{F}'_t)' | \mathbf{Y}_T \Big] (\mathbf{R} \otimes \mathbf{C})' \mathbb{D}'_{\mathbf{W}'_t} \Big) \\ &= \mathbf{H}^{-1} \star \Big(\mathbb{D}_{\mathbf{W}'_t} (\mathbf{R} \otimes \mathbf{C}) \Big(\mathbb{K}_{k_1 k_2} \Big(\mathbf{f}_{t|T} \mathbf{f}'_{t|T} + \mathbf{\Pi}_{t|T} \Big) \mathbb{K}'_{k_1 k_2} \Big) (\mathbf{R} \otimes \mathbf{C})' \mathbb{D}'_{\mathbf{W}'_t} \Big), \end{split}$$

$$\begin{split} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}')) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ (\mathbf{Y}_{t} - \mathbf{R}\mathbf{F}_{t}\mathbf{C}'))' | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{W}_{t} \circ \mathbf{Y}_{t}) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ \mathbf{Y}_{t})' - (\mathbf{W}_{t} \circ \mathbf{Y}_{t}) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ \mathbf{R}\mathbb{E}_{\widehat{\boldsymbol{\theta}}} [\mathbf{F}_{t} | \mathbf{Y}_{T}] \mathbf{C}')' \\ &- \left(\mathbf{W}_{t} \circ \mathbf{R}\mathbb{E}_{\widehat{\boldsymbol{\theta}}} [\mathbf{F}_{t} | \mathbf{Y}_{T}] \mathbf{C}' \right) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ \mathbf{Y}_{t})' \\ &+ \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{W}_{t} \circ \mathbf{R}\mathbf{F}_{t}\mathbf{C}') \mathbf{K}^{-1} (\mathbf{W}_{t} \circ \mathbf{R}\mathbf{F}_{t}\mathbf{C}')' | \mathbf{Y}_{T} \Big] \\ &= (\mathbf{W}_{t} \circ \mathbf{Y}_{t}) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ \mathbf{Y}_{t})' - (\mathbf{W}_{t} \circ \mathbf{Y}_{t}) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ \mathbf{R}\mathbf{F}_{t|T}\mathbf{C}')' \\ &- \left(\mathbf{W}_{t} \circ \mathbf{R}\mathbf{F}_{t|T}\mathbf{C}' \right) \mathbf{K}^{-1} (\mathbf{W}_{t} \circ \mathbf{Y}_{t})' \\ &+ \mathbf{K}_{t}^{-1} \star \Big(\mathbb{D}_{\mathbf{W}_{t}} (\mathbf{C} \otimes \mathbf{R}) \Big(\mathbf{f}_{t|T} \mathbf{f}_{t|T}' + \mathbf{\Pi}_{t|T} \Big) (\mathbf{C} \otimes \mathbf{R})' \mathbb{D}_{\mathbf{W}_{t}}' \Big), \end{split}$$

$$\begin{split} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{W}_t \circ (\mathbf{Y}_t - \mathbf{R} \mathbf{F}_t \mathbf{C}'))' \mathbf{H}^{-1} (\mathbf{W}_t \circ (\mathbf{Y}_t - \mathbf{R} \mathbf{F}_t \mathbf{C}')) | \mathbf{Y}_T \Big] \\ &= (\mathbf{W}_t \circ \mathbf{Y}_t)' \mathbf{H}^{-1} (\mathbf{W}_t \circ \mathbf{Y}_t) - (\mathbf{W}_t \circ \mathbf{Y}_t)' \mathbf{H}^{-1} \big(\mathbf{W}_t \circ \mathbf{R} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} [\mathbf{F}_t | \mathbf{Y}_T] \mathbf{C}' \big) \\ &- \big(\mathbf{W}_t \circ \mathbf{R} \mathbb{E}_{\widehat{\boldsymbol{\theta}}} [\mathbf{F}_t | \mathbf{Y}_T] \mathbf{C}' \big)' \mathbf{H}^{-1} (\mathbf{W}_t \circ \mathbf{Y}_t) \\ &+ \mathbb{E}_{\widehat{\boldsymbol{\theta}}} \Big[(\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}')' \mathbf{H}^{-1} (\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_t \mathbf{C}') | \mathbf{Y}_T \Big] \\ &= (\mathbf{W}_t \circ \mathbf{Y}_t)' \mathbf{H}^{-1} (\mathbf{W}_t \circ \mathbf{Y}_t) - (\mathbf{W}_t \circ \mathbf{Y}_t)' \mathbf{H}^{-1} \big(\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_{t|T} \mathbf{C}' \big) \\ &- \big(\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_{t|T} \mathbf{C}' \big)' \mathbf{H}^{-1} (\mathbf{W}_t \circ \mathbf{Y}_t) \\ &+ \mathbf{H}^{-1} \star \Big(\mathbb{D}_{\mathbf{W}_t'} (\mathbf{R} \otimes \mathbf{C}) \Big(\mathbb{K}_{k_1 k_2} \Big(\mathbf{f}_{t|T} \mathbf{f}_{t|T}' + \mathbf{\Pi}_{t|T} \Big) \mathbb{K}_{k_1 k_2}' \Big) (\mathbf{R} \otimes \mathbf{C})' \mathbb{D}_{\mathbf{W}_t'}' \Big). \end{split}$$

Combining these results with the FOC we obtain

$$\begin{aligned} \operatorname{vec}(\mathbf{R}) &= \left(\sum_{l=1}^{T} \sum_{s=1}^{p_1} \sum_{q=1}^{p_1} \left(\left(\mathbf{C}' \mathbb{D}_{\mathbf{W}_t}^{[s,q]} \mathbf{K}^{-1} \mathbf{C} \right) \star \left(\mathbf{f}_{l|T} \mathbf{f}_{l|T}' + \mathbf{H}_{l|T} \right) \right) \otimes \left(\mathbb{I}_{p_1} \mathbb{E}_{p_1,p_1}^{(s,q)} \mathbf{H}^{-1} \right) \right)^{-1} \\ &\qquad \left(\sum_{t=1}^{T} \operatorname{vec} \left(\left(\mathbf{W}_t \circ \mathbf{H}^{-1} \mathbf{Y}_t \mathbf{K}^{-1} \right) \mathbf{C} \mathbf{F}_{l|T}' \right) \right), \end{aligned} \right) \\ \operatorname{vec}(\mathbf{C}) &= \left(\sum_{t=1}^{T} \sum_{s=1}^{p_2} \sum_{q=1}^{p_2} \left(\mathbf{R}' \mathbb{D}_{\mathbf{W}_t}^{[k,q]} \mathbf{H}^{-1} \mathbf{R} \star \left(\mathbb{K}_{k_1 k_2} \left(\mathbf{f}_{l|T} \mathbf{f}_{l|T}' + \mathbf{\Pi}_{l|T} \right) \mathbb{K}_{k_1 k_2}' \right) \right) \otimes \left(\mathbb{I}_{p_2} \mathbb{E}_{p_2, p_2}^{(s,q)} \mathbf{K}^{-1} \right) \right)^{-1} \\ &\qquad \left(\sum_{t=1}^{T} \operatorname{vec} \left(\left(\mathbf{W}_t \circ \mathbf{H}^{-1} \mathbf{Y}_t \mathbf{K}^{-1} \right)' \mathbf{R} \mathbf{F}_t \right) \right), \end{aligned} \right) \end{aligned}$$

$$\left[\mathbf{H} \right]_{ii} &= \frac{1}{T_{p_2}} \sum_{t=1}^{T} \left[\left[\left((\mathbf{W}_t \circ \mathbf{Y}_t) \mathbf{K}^{-1} (\mathbf{W}_t \circ \mathbf{Y}_t)' - (\mathbf{W}_t \circ \mathbf{Y}_t) \mathbf{K}^{-1} (\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_{t|T} \mathbf{C}')' \right) \right. \\ &\qquad \left. - \left(\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_{t|T} \mathbf{C}' \right) \mathbf{K}^{-1} (\mathbf{W}_t \circ \mathbf{Y}_t)' \right. \\ \left. + \left(\widehat{\mathbf{H}}_{1p_1, p_2} \widehat{\mathbf{K}} \right) \mathbf{K}^{-1} (\mathbf{1}_{1p_1, p_2} - \mathbf{W}_t)' \right]_{ii}, \end{aligned}$$

$$\left[\mathbf{K} \right]_{ii} &= \frac{1}{T_{p_1}} \sum_{t=1}^{T} \left[\left[\left((\mathbf{W}_t \circ \mathbf{Y}_t)' \mathbf{H}^{-1} (\mathbf{W}_t \circ \mathbf{Y}_t) - (\mathbf{W}_t \circ \mathbf{R} \mathbf{F}_{t|T} \mathbf{C}' \right) \mathbf{K}^{-1} (\mathbf{W}_t \otimes \mathbf{R} \mathbf{F}_{t|T} \mathbf{C}') \right) \right. \\ &\qquad \left. + \left(\widehat{\mathbf{H}}_{1p_1, p_2} \widehat{\mathbf{K}} \right) \mathbf{K}^{-1} (\mathbf{1}_{p_1, p_2} - \mathbf{W}_t)' \right]_{ii}, \end{aligned}$$

For any iteration $n \ge 0$, given an estimator of the parameters $\widehat{\boldsymbol{\theta}}^{(n)}$, the explicit solutions for **R** and **C** obtained from the previous FOCs are given in Section 5. While the solutions for the diagonal elements

of ${\bf H}$ and ${\bf K}$ are the following:

$$\begin{split} [\widehat{\mathbf{H}}^{(n+1)}]_{ii} &= \frac{1}{Tp_2} \sum_{t=1}^{T} \left[\left((\mathbf{W}_t \circ \mathbf{Y}_t) \widehat{\mathbf{K}}^{(n)-1} (\mathbf{W}_t \circ \mathbf{Y}_t)' \right. \\ &\left. - (\mathbf{W}_t \circ \mathbf{Y}_t) \widehat{\mathbf{K}}^{(n)-1} \left(\mathbf{W}_t \circ \left(\widehat{\mathbf{R}}^{(n+1)} \mathbf{F}_{t|T}^{(n)} \widehat{\mathbf{C}}^{(k+1)\prime} \right) \right)' \right. \\ &\left. - \left(\mathbf{W}_t \circ \left(\widehat{\mathbf{R}}^{(n+1)} \mathbf{F}_{t|T}^{(n)} \widehat{\mathbf{C}}^{(k+1)\prime} \right) \right) \widehat{\mathbf{K}}^{(n)-1} (\mathbf{W}_t \circ \mathbf{Y}_t)' \right. \\ &\left. + \widehat{\mathbf{K}}^{(n)-1} \star \left(\mathbb{D}_{\mathbf{W}_t} \left(\widehat{\mathbf{C}}^{(n+1)} \otimes \widehat{\mathbf{R}}^{(n+1)} \right) \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)} \right) \left(\widehat{\mathbf{C}}^{(n+1)} \otimes \widehat{\mathbf{R}}^{(n+1)} \right)' \mathbb{D}'_{\mathbf{W}_t} \right) \right)' \\ &\left. + \left(\widehat{\mathbf{H}}^{(n)} \mathbf{1}_{p_1, p_2} \widehat{\mathbf{K}}^{(n)} \right) \widehat{\mathbf{K}}^{(n)-1} (\mathbf{1}_{p_1, p_2} - \mathbf{W}_t)' \right]_{ii}, \end{split}$$

$$\begin{split} [\widehat{\mathbf{K}}^{(n+1)}]_{ii} &= \frac{1}{Tp_1} \sum_{t=1}^{T} \Big[\Big((\mathbf{W}_t \circ \mathbf{Y}_t)' \widehat{\mathbf{H}}^{(n+1)-1} (\mathbf{W}_t \circ \mathbf{Y}_t) \\ &- (\mathbf{W}_t \circ \mathbf{Y}_t)' \widehat{\mathbf{H}}^{(n+1)-1} \Big(\mathbf{W}_t \circ \Big(\widehat{\mathbf{R}}^{(n+1)} \mathbf{F}_{t|T}^{(n)} \widehat{\mathbf{C}}^{(k+1)'} \Big) \Big) \\ &- \Big(\mathbf{W}_t \circ \Big(\widehat{\mathbf{R}}^{(n+1)} \mathbf{F}_{t|T}^{(n)} \widehat{\mathbf{C}}^{(k+1)'} \Big) \Big)' \widehat{\mathbf{H}}^{(n+1)-1} (\mathbf{W}_t \circ \mathbf{Y}_t) \\ &+ \widehat{\mathbf{H}}^{(n+1)-1} \star \Big(\mathbb{D}_{\mathbf{W}_t'} \Big(\widehat{\mathbf{R}}^{(n+1)} \otimes \widehat{\mathbf{C}}^{(n+1)} \Big) \Big(\mathbb{K}_{k_1 k_2} \Big(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)'} + \mathbf{\Pi}_{t|T}^{(n)} \Big) \mathbb{K}_{k_1 k_2}' \Big) \Big(\widehat{\mathbf{R}}^{(n+1)} \otimes \widehat{\mathbf{C}}^{(n+1)} \Big)' \mathbb{D}_{\mathbf{W}_t'}' \Big) \Big)' \\ &+ (\mathbf{1}_{p_1, p_2} - \mathbf{W}_t)' \widehat{\mathbf{H}}^{(n+1)-1} \Big(\widehat{\mathbf{H}}^{(n+1)} \mathbf{1}_{p_1, p_2} \widehat{\mathbf{K}}^{(n)} \Big) \Big]_{ii}. \end{split}$$

C Asymptotic results

As k_1 , k_2 are both fixed constants, without loss of generality, we assume $k_1 = k_2 = 1$ in some parts of the proofs as long as it simplifies the notations.

C.1 Proof of main results

Proof of Proposition 1. Recall that

$$\begin{aligned} \widehat{\mathbf{R}}^{(n+1)} &= \left(\sum_{t=1}^{T} \mathbf{Y}_{t} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \widehat{\mathbf{F}}_{t|T}^{(n)\prime}\right) \left(\left(\widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)}\right) \star \left(\sum_{t=1}^{T} \mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)}\right)\right)^{-1} \\ &= \mathbf{R} \left(\sum_{t=1}^{T} \mathbf{F}_{t} \mathbf{C} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \widehat{\mathbf{F}}_{t|T}^{(n)\prime}\right) \left(\left(\widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(n)}\right) \star \left(\sum_{t=1}^{T} \mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)}\right)\right)^{-1} \\ &+ \left(\sum_{t=1}^{T} \mathbf{E}_{t} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \widehat{\mathbf{F}}_{t|T}^{(n)\prime}\right) \left(\left(\widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)}\right) \star \left(\sum_{t=1}^{T} \mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)}\right)\right)^{-1} \end{aligned}$$

we can thus write

$$\begin{aligned} \widehat{\mathbf{R}}^{(n+1)} - \widehat{\mathbf{R}} \widehat{\mathbf{J}}_{1} &= \left\{ \widehat{\mathbf{R}} \widehat{\mathbf{J}}_{1} \left(\frac{1}{p_{2}} \widehat{\mathbf{J}}_{2}^{\prime} \mathbf{C}^{\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \left(f_{t|T}^{(n)} - \widehat{\mathbf{J}}^{-1} f_{t} \right) f_{t|T}^{(n)\prime} \right) \\ &- \widehat{\mathbf{R}} \widehat{\mathbf{J}}_{1} \left(\frac{1}{p_{2}} \left(\widehat{\mathbf{C}}^{(n)} - \widehat{\mathbf{C}} \widehat{\mathbf{J}}_{2} \right)^{\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} f_{t|T}^{(n)} f_{t|T}^{(n)\prime} \right) \\ &- \widehat{\mathbf{R}} \widehat{\mathbf{J}}_{1} \left(\frac{1}{p_{2}} \widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \mathbf{\Pi}_{t|T}^{(n)} \right) \\ &+ \frac{1}{Tp_{2}} \sum_{t=1}^{T} \widehat{\mathbf{E}}_{t} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \widehat{\mathbf{F}}_{t|T}^{(n)\prime} \right\} \\ &\times \left(\left(\frac{1}{p_{2}} \widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} f_{t|T}^{(n)} f_{t|T}^{(n)\prime} + \mathbf{\Pi}_{t|T}^{(n)} \right) \right)^{-1}. \end{aligned}$$

Now, let n=0, we have

$$\frac{1}{\sqrt{p_{1}}} \left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \left(\frac{\widehat{\mathbf{J}}_{2}' \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)}}{p_{2}} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right) \mathbf{f}_{t|T}^{(0)'} \right) \right\| \\
\leq k_{2}^{2} \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \left\| \frac{\widehat{\mathbf{J}}_{2}' \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)}}{p_{2}} \right\| \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right) \mathbf{f}_{t|T}^{(0)'} \right\| \\
\leq \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \frac{\left\| \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|}{\sqrt{p_{2}}} \left\| \frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - (\mathbf{C} \widehat{\mathbf{J}}_{2})' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_{2}}} \right\| \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right) \mathbf{f}_{t|T}^{(0)'} \right\| \\
+ \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \frac{\left\| \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|}{\sqrt{p_{2}}} \left\| \frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_{2}}} \right\| \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right) \mathbf{f}_{t|T}^{(0)'} \right\| \\
= O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right)$$
(23)

by Lemmas 1, 3(iii), 3(v), 8(iii), and 11(i),

$$\frac{1}{\sqrt{p_{1}}} \left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \left(\frac{1}{p_{2}} \left(\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right)' \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} \right) \right\| \\
\leq k_{2}^{2} \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \left\| \frac{\left(\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right)' \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right\|}{p_{2}} \right\| \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} \right\| \\
\leq \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \frac{\left\| \widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|}{\sqrt{p_{2}}} \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \frac{\left\| \widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|}{\sqrt{p_{2}}} \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} \right\| \\
+ \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \frac{\left\| \widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|}{\sqrt{p_{2}}} \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \frac{\left\| \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|}{\sqrt{p_{2}}} \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} \right\| \\
= O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{p_{1}p_{2}}, \frac{1}{Tp_{2}} \right\} \right)$$
(24)

by Lemmas 1, 3(iii), 4(i), 8(i), and since

$$\begin{aligned} \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)\prime} \right\| &\leq & \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right) \left(\mathbf{f}_{t|T}^{(0)\prime} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right) \right\| + 2 \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right) \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right\| + \left\| \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \mathbf{f}_t' \widehat{\mathbf{J}}^{-\prime} \right\| \\ &\lesssim & \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right) \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right\| + \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_t \mathbf{f}_t' \right\| \\ &= & O_p(1) \end{aligned}$$

by Assumption 1(ii) and Lemma 11(i),

$$\frac{1}{\sqrt{p_{1}}} \left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \left(\frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)}}{p_{2}} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \mathbf{\Pi}_{t|T}^{(0)} \right) \right\| \leq k_{2}^{2} \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \left\| \frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)}}{p_{2}} \right\| \max_{1 \leq t \leq T} \left\| \mathbf{\Pi}_{t|T}^{(0)} \right\| \\ \leq \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \frac{\left\| \widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|^{2}}{p_{2}} \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \max_{1 \leq t \leq T} \left\| \mathbf{\Pi}_{t|T}^{(0)} \right\| \\ + \frac{\left\| \mathbf{R} \widehat{\mathbf{J}}_{1} \right\|}{\sqrt{p_{1}}} \frac{\left\| \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|^{2}}{p_{2}} \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \max_{1 \leq t \leq T} \left\| \mathbf{\Pi}_{t|T}^{(0)} \right\| \\ = O_{p} \left(\frac{1}{p_{1} p_{2}} \right)$$
(25)

by Lemmas 1, 3(iii), 4(ii), 8(i), and Lemma D.12 in Barigozzi and Luciani (2024). Moreover,

$$\frac{1}{\sqrt{p_{1}}} \left\| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{E}_{t} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{F}}_{t|T}^{(0)\prime} \right\| \lesssim \frac{1}{\sqrt{p_{1}}} \left\| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{E}_{t} \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \left(\widehat{\mathbf{F}}_{t|T}^{(0)} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)' \right\|
= \frac{1}{\sqrt{p_{1}}} \left\| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{E}_{t} \left(\widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \right) \mathbf{F}_{t}' \right\|
= \frac{1}{\sqrt{p_{1}}} \left\| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{E}_{t} \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \mathbf{F}_{t}' \right\|
= O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right)$$
(26)

since

$$\begin{split} \frac{1}{T\sqrt{p_1p_2}} \left\| \sum_{t=1}^{T} \mathbf{E}_t \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \left(\widehat{\mathbf{F}}_{t|T}^{(0)} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \right\| &= \frac{1}{T\sqrt{p_1p_2}} \left\| \left(\mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right) \star \left(\sum_{t=1}^{T} \mathbf{e}_t \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right)' \right) \right\| \\ &= \frac{1}{T\sqrt{p_1p_2}} \left\| \sum_{i=1}^{p_2} \sum_{j=1}^{k_2} k_{ii}^{-1} c_{ij} \left[\sum_{t=1}^{T} \mathbf{e}_t \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right)' \right]_{ij}^{(p_1,k_1)} \right\| \\ &\leq \frac{\overline{c}C_K}{T\sqrt{p_1p_2}} \left\| \sum_{i=1}^{p_2} \sum_{j=1}^{k_2} \left[\sum_{t=1}^{T} \mathbf{e}_t \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right)' \right]_{ij}^{(p_1,k_1)} \right\| \\ &\leq \overline{c}C_K \left\| \frac{1}{T\sqrt{p_1p_2}} \sum_{i=1}^{p_2} \sum_{t=1}^{T} \mathbf{e}_{t-i} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right)' \right\| \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_{1p_2}} \right\} \right) \end{split}$$

by Lemma 11(iv), and

$$\begin{aligned} \frac{1}{\sqrt{p_1}} \left\| \frac{1}{Tp_2} \sum_{t=1}^T \mathbf{E}_t \left(\widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}} - \operatorname{dg}(\mathbf{K})^{-1} \mathbf{C} \right) \mathbf{F}'_t \right\| &\leq \frac{1}{\sqrt{Tp_1p_2}} \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{E}_t \otimes \mathbf{F}'_t \right\| \left\| \frac{\widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}} - \operatorname{dg}(\mathbf{K})^{-1} \mathbf{C}}{\sqrt{p_2}} \right\| \\ &= \frac{1}{\sqrt{T}} O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right) \end{aligned}$$

by Assumption 3(i) and Lemma 8(iii),

$$\frac{1}{\sqrt{p_1}} \left\| \frac{1}{Tp_2} \sum_{t=1}^T \mathbf{E}_t \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \mathbf{F}_t' \right\| = O_p \left(\frac{1}{\sqrt{Tp_2}} \right)$$

by Lemma A.1 in Yu et al. (2022). Finally, we have that

$$\begin{aligned} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} + \mathbf{\Pi}_{t|T}^{(0)} \right) - \left(\frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_{2}} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{f}_{t}' \widehat{\mathbf{J}}' \right) \right\| \\ \leq \left\| \left(\frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_{2}} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} + \mathbf{\Pi}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{f}_{t}' \widehat{\mathbf{J}}' \right) \right\| \\ + \left\| \left(\frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_{2}} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} + \mathbf{\Pi}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{f}_{t}' \widehat{\mathbf{J}}' \right) \right\| \\ + \left\| \left(\frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_{2}} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \mathbf{J}^{-1} \mathbf{f}_{t} \mathbf{f}_{t}' \widehat{\mathbf{J}}' \right) \right\| \\ = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right) \end{aligned}$$

$$(27)$$

since the first addendum is dominated by the other two terms and

$$\begin{split} \left\| \left(\frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_2} \right) \star \left(\frac{1}{T} \sum_{t=1}^T \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} + \mathbf{\Pi}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \mathbf{f}_t' \widehat{\mathbf{J}}' \right) \right\| \\ & \leq k_2^2 \left\| \frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_2}} \right\| \left\| \frac{\mathbf{C}}{\sqrt{p_2}} \right\| \left\| \left(\frac{1}{T} \sum_{t=1}^T \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} + \mathbf{\Pi}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \mathbf{f}_t' \widehat{\mathbf{J}}' \right) \right\| \\ & \leq \left\| \left(\frac{1}{T} \sum_{t=1}^T \mathbf{f}_{t|T}^{(0)} \mathbf{f}_{t|T}^{(0)'} - \frac{1}{T} \sum_{t=1}^T \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \mathbf{f}_t' \widehat{\mathbf{J}}' \right) \right\| + \max_{1 \leq t \leq T} \left\| \mathbf{\Pi}_{t|T}^{(0)} \right\| \\ & \leq \left\| \left(\frac{1}{T} \sum_{t=1}^T \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right) \mathbf{f}_{t|T}^{(0)'} \right) \right\| + \max_{1 \leq t \leq T} \left\| \mathbf{\Pi}_{t|T}^{(0)} \right\| \\ & = O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) + O_p \left(\frac{1}{p_1p_2} \right) \end{split}$$

by Lemmas 1, 3(iii), 3(v), 11(i) and Lemma D.12 in Barigozzi and Luciani (2024),

$$\begin{split} \left\| \left(\frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_2} \right) \star \left(\frac{1}{T} \sum_{t=1}^T \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \mathbf{f}_t' \widehat{\mathbf{J}}^{-\prime} \right) \right\| \\ & \leq k_2^2 \left\| \left(\frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_2} \right) \right\| \left\| \frac{1}{T} \sum_{t=1}^T \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \mathbf{f}_t' \widehat{\mathbf{J}}^{-\prime} \right\| \\ & \leq \left\| \frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - (\mathbf{C} \widehat{\mathbf{J}}_2)' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_2}} \right\| \left\| \frac{\mathbf{C} \widehat{\mathbf{J}}_2}{\sqrt{p_2}} \right\| + \left\| \frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_2}} \right\| \left\| \frac{\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2}{\sqrt{p_2}} \right\| \\ & = O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right) \end{split}$$

by Assumption 1(ii) and Lemmas 1, 3(iii), 3(v), 4(ii) and 8(iii). Moreover,

$$\left\| \left(\left(\frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_2} \right) \star \left(\frac{1}{T} \sum_{t=1}^T \mathbf{f}_t \mathbf{f}_t' \right) \right)^{-1} \right\| = \left\| \left(\frac{1}{Tp_2} \sum_{t=1}^T \mathbf{F}_t \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \mathbf{F}_t' \right)^{-1} \right\|$$

$$\leq \left\| \left(\frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_2} \right)^{-1} \right\| \left\| \left(\frac{1}{T} \sum_{t=1}^T \mathbf{F}_t \mathbf{F}_t' \right)^{-1} \right\|$$

$$= O_p(1)$$

$$(28)$$

by Lemma 3(iv) and Assumption 1(ii). Combining (22) with (23)-(26) and (27)-(28), we obtain

$$\frac{1}{\sqrt{p_1}} \left\| \widehat{\mathbf{R}}^{(1)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right).$$

Consider now n>0, the consistency result for $\widehat{\mathbf{R}}^{(n+1)}$ follows iterating the same steps but using Lemma 10 and this proposition in place of Lemmas 4 and 8. The proof for $\widehat{\mathbf{C}}$ follows analogously and it is omitted.

For the row-wise consistency note that we can use the decomposition in (22) using \mathbf{r}_i in place of \mathbf{R} ,

then (23)-(25) follows analogously as $\|\mathbf{r}_{i}\| = O_p(1)$ by Assumption 1(*i*). Then

$$\begin{aligned} \left\| \frac{1}{T_{p_2}} \sum_{t=1}^{T} \mathbf{e}'_{ti} \cdot \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{F}}^{(0)\prime}_{t|T} \right\| &\lesssim \left\| \frac{1}{T_{p_2}} \sum_{t=1}^{T} \mathbf{e}'_{ti} \cdot \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \left(\widehat{\mathbf{F}}^{(0)}_{t|T} - \widehat{\mathbf{J}}^{-1}_{1} \mathbf{F}_{t} \widehat{\mathbf{J}}^{-\prime}_{2} \right)' \right\| \\ &\left\| \frac{1}{T_{p_2}} \sum_{t=1}^{T} \mathbf{e}'_{ti} \cdot \left(\widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \right) \mathbf{F}'_{t} \right\| \\ &\left\| \frac{1}{T_{p_2}} \sum_{t=1}^{T} \mathbf{e}'_{ti} \cdot \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \mathbf{F}'_{t} \right\| \\ &= O_p \left(\max \left\{ \frac{1}{\sqrt{T_{p_1}}}, \frac{1}{\sqrt{T_{p_2}}}, \frac{1}{p_{1p_2}} \right\} \right) \end{aligned}$$
(29)

since

$$\frac{1}{Tp_2} \left\| \sum_{t=1}^{T} \mathbf{e}'_{ti.} \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \left(\widehat{\mathbf{F}}_{t|T}^{(0)} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \right\| = \frac{1}{Tp_2} \left\| \sum_{t=1}^{T} \sum_{j=1}^{p_2} \sum_{q=1}^{k_2} e_{tij} k_{jj}^{-1} c_{jq} \left[\widehat{\mathbf{F}}_{t|T}^{(0)} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right]_{\cdot q} \right\| \\ \leq \bar{c} C_K \left\| \frac{1}{Tp_2} \sum_{j=1}^{p_2} \sum_{t=1}^{T} e_{tij} \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right) \right\| \\ = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \right)$$

by Lemma 11(ii),

$$\left\| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{e}_{ti}' \left(\widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right) \mathbf{F}_t' \right\| \lesssim \frac{1}{\sqrt{T}} \frac{\left\| \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right\|}{\sqrt{p_2}} \left\| \frac{1}{\sqrt{Tp_2}} \sum_{t=1}^{T} \sum_{j=1}^{p_2} e_{tij} \mathbf{F}_t' \right\|$$
$$= \frac{1}{\sqrt{T}} O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right)$$

by Assumption 3(i) and Lemma 8(iii),

$$\left\|\frac{1}{Tp_2}\sum_{t=1}^{T}\mathbf{e}_{ti} \cdot \mathrm{dg}(\mathbf{K})^{-1}\mathbf{C}\mathbf{F}_t'\right\| \leq \frac{C_K}{\sqrt{Tp_2}} \left\|\frac{1}{\sqrt{T}}\sum_{t=1}^{T}\sum_{j=1}^{p_2} e_{tij}\frac{\mathbf{c}_{j\cdot}}{\sqrt{p_2}}\mathbf{F}_t'\right\| = O_p\left(\frac{1}{\sqrt{Tp_2}}\right)$$

follows directly from Assumption 3(i). Iterating the same steps using Lemma 10 in place of Lemma 8 yields the result for n>0. Row-wise consistency for $\hat{\mathbf{c}}_{j}$ can be established analogously.

Proof of Proposition 2. Since $\operatorname{vec}\left(\widehat{\mathbf{F}}_{t}^{(n)}\right) = \mathbf{f}_{t|T}^{(n)}$, we have

$$\begin{aligned} \left\| \widehat{\mathbf{F}}_{t}^{(n)} - \widehat{\mathbf{J}}_{t}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right\| &\leq \left\| \widehat{\mathbf{F}}_{t}^{(n)} - \widehat{\mathbf{J}}_{t}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right\|_{F} \\ &= \left\| \mathbf{f}_{t|T}^{(n)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right\| \\ &\leq \left\| \mathbf{f}_{t|T}^{(n)} - \mathbf{f}_{t|t}^{(n)} \right\| + \left\| \mathbf{f}_{t|t}^{(n)} - \mathbf{f}_{t}^{LS(n)} \right\| + \left\| \mathbf{f}_{t}^{LS(n)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right\| \end{aligned}$$

with $\widehat{\mathbf{J}} = \widehat{\mathbf{J}}_2 \otimes \widehat{\mathbf{J}}_1$ and

$$\mathbf{f}_{t}^{LS(n)} = \left(\left(\left(\widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right)^{-1} \widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(n)\prime} \widehat{\mathbf{H}}^{(n)-1} \widehat{\mathbf{R}}^{(n)} \right)^{-1} \widehat{\mathbf{R}}^{(n)\prime} \widehat{\mathbf{H}}^{(n)-1} \right) \right) \mathbf{y}_{t}.$$

Consider the case n=0. Combining Lemmas D.15 and D.16 in Barigozzi and Luciani (2024) with Lemmas 4, 6, 8, we have that

$$\left| \mathbf{f}_{t|T}^{(0)} - \mathbf{f}_t \right| \leq \left\| \mathbf{f}_t^{LS(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right\| + O_p \left(\frac{1}{p_1 p_2} \right)$$

Then,

$$\begin{split} \left\| f_{t}^{LS(0)} - \widehat{\mathbf{J}}^{-1} f_{t} \right\| &\leq \\ \left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \right) \right) \right. \\ &\times \left(\mathbf{C} \widehat{\mathbf{J}}_{2} \otimes \mathbf{R} \widehat{\mathbf{J}}_{1} - \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} \right) \left\| \left\| \widehat{\mathbf{J}}^{-1} f_{t} \right\| \right. \\ &+ \left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \right) \right) \right\| \\ &\leq \\ &\left\| \left(\left(\left(\mathbf{C}^{'} \mathbf{K}^{-1} \mathbf{C} \right)^{-1} \mathbf{C}^{'} \mathbf{K}^{-1} \right) \otimes \left(\left(\mathbf{R}^{'} \mathbf{H}^{-1} \mathbf{R} \right)^{-1} \mathbf{R}^{'} \mathbf{H}^{-1} \right) \right) \left(\mathbf{C} \widehat{\mathbf{J}}_{2} \otimes \mathbf{R} \widehat{\mathbf{J}}_{1} - \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} \right) \right\| \left\| \widehat{\mathbf{J}}^{-1} f_{t} \right\| \\ &+ \\ &\left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \right) \right) \\ &- \left(\left(\mathbf{C}^{'} \mathbf{K}^{-1} \mathbf{C}^{-1} \mathbf{C}^{'} \mathbf{K}^{-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \right) \right\| \\ &+ \\ &\left\| \left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\mathbf{R}^{'} \mathbf{H}^{-1} \mathbf{R} \right)^{-1} \right) \right\| \\ &\times \left\| \left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \otimes \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \right) \otimes \left(\mathbf{R}^{'} \mathrm{dg}(\mathbf{H})^{-1} \right) \right) \right\| \\ &+ \\ &\left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \otimes \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \right) \\ &\times \left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \otimes \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \right) - \left(\mathbf{C}^{'} \mathrm{dg}(\mathbf{K})^{-1} \right) \right) \right\| \right\| \\ \\ &+ \\ &\left\| \left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \otimes \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \right)^{-1} \right) \\ &+ \\ &\left\| \left(\left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \otimes \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \right) - \left(\mathbf{C}^{'} \mathrm{dg}(\mathbf{K})^{-1} \right) \right) \right\| \right\| \right\| \\ \\$$

Now,

$$\begin{split} I &\leq \left\| \left(\frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_2} \right)^{-1} \right\| \left\| \frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_2}} \right\| \left\| \left(\frac{\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R}}{p_1} \right)^{-1} \right\| \left\| \frac{\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1}}{\sqrt{p_2}} \right\| \left\| \frac{\mathbf{C} \widehat{\mathbf{J}}_2 \otimes \mathbf{R} \widehat{\mathbf{J}}_1 - \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)}}{\sqrt{p_1 p_2}} \right\| \left\| \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right\| \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right) \end{split}$$

by Lemmas 3(iv), 3(v), 4(iii), and since $\|\mathbf{f}_t\| = O_p(1)$ by Assumption 1(*ii*),

$$II \leq \sqrt{p_2} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - \left(\mathbf{C'} \mathbf{K}^{-1} \mathbf{C} \right)^{-1} \mathbf{C'} \mathbf{K}^{-1} \right\| \left\| \left(\frac{\mathbf{R'} \mathbf{H}^{-1} \mathbf{R}}{p_1} \right)^{-1} \right\| \frac{\| \mathbf{R'} \mathbf{H}^{-1} \|}{\sqrt{p_1}} \\ \times \frac{1}{\sqrt{p_1 p_2}} \left\| \left(\mathbf{C} \widehat{\mathbf{J}}_2 \otimes \mathbf{R} \widehat{\mathbf{J}}_1 - \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} \right) \right\| \left\| \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right\| \\ \left\| \left(\frac{\mathbf{C'} \mathbf{K}^{-1} \mathbf{C}}{p_2} \right)^{-1} \right\| \frac{\| \mathbf{C'} \mathbf{K}^{-1} \|}{\sqrt{p_2}} \sqrt{p_1} \right\| \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} - \left(\mathbf{R'} \mathbf{H}^{-1} \mathbf{R} \right)^{-1} \mathbf{R'} \mathbf{H}^{-1} \right\| \\ \times \frac{1}{\sqrt{p_1 p_2}} \left\| \left(\mathbf{C} \widehat{\mathbf{J}}_2 \otimes \mathbf{R} \widehat{\mathbf{J}}_1 - \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} \right) \right\| \left\| \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right\| \\ = o_p \left(\max \left\{ \frac{1}{\sqrt{p_1} p_1}, \frac{1}{\sqrt{p_2}}, \frac{1}{p_1 p_2} \right\} \right)$$

by Lemmas 3(iv), 3(v), 4(iii), 8(ix), 8(x), and Assumption 1(ii),

$$III \leq \left\| \left(\frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)}}{p_2} \right)^{-1} \right\| \left\| \left(\frac{\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)}}{p_1} \right)^{-1} \right\| \frac{1}{p_1 p_2} \left\| \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{E}_t \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right\|_F$$
$$= O_p \left(\frac{1}{\sqrt{p_1 p_2}} \right)$$

by Lemma 8(v), 8(vi), and since

$$\mathbb{E}\left[\left\|\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{E}_{t} \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}\right\|_{F}^{2}\right] = \mathbb{E}\left[\sum_{i=1}^{k_{1}} \sum_{j=1}^{k_{2}} \left(\mathbf{r}'_{\cdot i} \mathrm{dg}(\mathbf{H})^{-1} \mathbf{E}_{t} \mathrm{dg}(\mathbf{K})^{-1} \mathbf{c}_{\cdot j}\right)^{2}\right] \\ = \sum_{i=1}^{k_{1}} \sum_{j=1}^{k_{2}} \mathbb{E}\left[\left(\sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{2}} r_{si} h_{ss}^{-1} e_{tsq} k_{qq}^{-1} c_{qj}\right)^{2}\right] \\ \leq k \bar{r}^{2} \bar{c}^{2} C_{K}^{2} C_{H}^{2} \mathbb{E}\left[\left(\sum_{s=1}^{p_{1}} \sum_{q=1}^{p_{2}} e_{tsq}\right)^{2}\right] \\ \lesssim \sum_{s_{1}, s_{2}} \sum_{q_{1}, q_{2}}^{p_{2}} |\mathbb{E}[e_{ts_{1}q_{1}} e_{ts_{2}q_{2}}]| \\ = O_{p}(p_{1}p_{2})$$

by Assumptions 1(i), 2(ii), and 2(iii),

$$\begin{split} IV &\leq \left\| \left(\frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)}}{p_2} \right)^{-1} \right\| \left\| \left(\frac{\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)}}{p_1} \right)^{-1} \right\| \left\| \frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_2}} \right\| \left\| \frac{\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1}}{\sqrt{p_1}} \right\| \left\| \frac{\mathbf{e}_t}{\sqrt{p_1 p_2}} \right\| \\ &+ \left\| \left(\frac{\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)}}{p_2} \right)^{-1} \right\| \left\| \left(\frac{\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)}}{p_1} \right)^{-1} \right\| \left\| \frac{\widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} - \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1}}{\sqrt{p_1}} \right\| \left\| \frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_2}} \right\| \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right) \end{split}$$

by Lemma 3(v), 8(iii)-8(vi) and since

$$\mathbb{E}\left[\left\|\mathbf{e}_{t}\right\|^{2}\right] = \mathbb{E}\left[\left\|\mathbf{E}_{t}\right\|_{F}^{2}\right] = \sum_{i=1}^{p_{1}} \sum_{j=1}^{p_{2}} \mathbb{E}\left[\left|e_{tij}\right|^{2}\right] \le p_{1}p_{2}C_{k}C_{h} = O(p_{1}p_{2})$$

by Assumption 2(ii). Iterating the same steps, using Proposition 1, Proposition (a.4)-(a.5) in Barigozzi

and Luciani (2024), and Lemma 10 in place of Lemmas 4, 6, 8, we can obtain the result for n > 0.

C.2 Auxiliary lemmata

C.2.1 Preliminary results

Lemma 1. Let **A** and **B** be $m \times n$ and $mp \times nq$ matrices, respectively. We have that

$$\|\mathbf{A}\star\mathbf{B}\| \leq mn \|\mathbf{A}\|_{\max} \|\mathbf{B}\|$$

Proof.

$$\begin{aligned} \|\mathbf{A}\star\mathbf{B}\| &= \left\| \sum_{i=1}^{m} \sum_{j=1}^{n} a_{ij} \mathbf{B}_{ij}^{(p,q)} \right\| \\ &\leq \|\mathbf{A}\|_{\max} \sum_{i=1}^{m} \sum_{j=1}^{n} \left\| \mathbf{B}_{ij}^{(p,q)} \right\| \\ &\leq mn \|\mathbf{A}\|_{\max} \max_{i,j} \left\| \mathbf{B}_{ij}^{(p,q)} \right\| \\ &\leq mn \|\mathbf{A}\|_{\max} \|\mathbf{B}\| \end{aligned}$$

| Г | - | - | ٦ |
|---|---|---|---|
| | | | 1 |
| | | | 1 |
| - | | | - |

Lemma 2. For any $k_1 \times k_1$ and $k_2 \times k_2$ orthogonal matrices \mathbf{J}_1 and \mathbf{J}_2 , the DMFM in (3)-(4) is equivalent to

$$\mathbf{Y}_t = \widetilde{\mathbf{R}} \widetilde{\mathbf{F}}_t \widetilde{\mathbf{C}}' + \mathbf{E}_t \tag{30}$$

$$\widetilde{\mathbf{F}}_{t} = \widetilde{\mathbf{A}} \widetilde{\mathbf{F}}_{t-1} \widetilde{\mathbf{B}}' + \widetilde{\mathbf{U}}_{t}$$
(31)

and the vectorized form in (6)-(7) can be written as follows,

$$\mathbf{y}_t = \left(\widetilde{\mathbf{C}} \otimes \widetilde{\mathbf{R}}\right) \widetilde{\mathbf{f}}_t + \mathbf{e}_t \tag{32}$$

$$\widetilde{\mathsf{f}}_t = \left(\widetilde{\mathbf{B}} \otimes \widetilde{\mathbf{A}}\right) \widetilde{\mathsf{f}}_{t-1} + \widetilde{\mathsf{u}}_t \tag{33}$$

with $\mathbb{E}\left[\widetilde{\mathbf{U}}_{t}\widetilde{\mathbf{U}}_{t}'\right] = \widetilde{\mathbf{P}}tr\left(\widetilde{\mathbf{Q}}\right)$ and $\mathbb{E}\left[\widetilde{\mathbf{U}}_{t}'\widetilde{\mathbf{U}}_{t}\right] = \widetilde{\mathbf{Q}}tr\left(\widetilde{\mathbf{P}}\right)$ such that $\widetilde{\mathbf{P}} = \widetilde{\mathbf{\Gamma}}^{P}\widetilde{\mathbf{\Lambda}}^{P}\widetilde{\mathbf{\Gamma}}^{P'}$ and $\widetilde{\mathbf{Q}} = \widetilde{\mathbf{\Gamma}}^{Q}\widetilde{\mathbf{\Lambda}}^{Q}\widetilde{\mathbf{\Gamma}}^{Q'}$, where

$$\begin{split} \widetilde{\mathbf{R}} &= \mathbf{R} \mathbf{J}_{1}, & \widetilde{\mathbf{C}} = \mathbf{C} \mathbf{J}_{2}, & \widetilde{\mathbf{F}}_{t} = \mathbf{J}_{1}^{-1} \mathbf{F}_{t} (\mathbf{J}_{2}^{-1})', \\ \widetilde{\mathbf{A}} &= \mathbf{J}_{1}^{-1} \mathbf{A} \mathbf{J}_{1}, & \widetilde{\mathbf{B}} = \mathbf{J}_{2}^{-1} \mathbf{B} \mathbf{J}_{2}, & \widetilde{\mathbf{U}}_{t} = \mathbf{J}_{1}^{-1} \mathbf{U}_{t} (\mathbf{J}_{2}^{-1})', \\ \widetilde{\mathbf{P}} &= \mathbf{J}_{1}^{-1} \mathbf{P} \mathbf{J}_{1}, & \widetilde{\mathbf{\Gamma}}^{P} = \mathbf{J}_{1}^{-1} \mathbf{\Gamma}^{P} & \widetilde{\mathbf{A}}^{P} = \mathbf{A}^{P} \\ \widetilde{\mathbf{Q}} &= \mathbf{J}_{2}^{-1} \mathbf{Q} \mathbf{J}_{2} & \widetilde{\mathbf{\Gamma}}^{Q} = \mathbf{J}_{2}^{-1} \mathbf{\Gamma}^{Q} & \widetilde{\mathbf{A}}^{Q} = \mathbf{A}^{Q} \end{split}$$

Proof. Plug-in all the rotated ("tilde") matrices to obtain,

$$\mathbf{Y}_{t} = \widetilde{\mathbf{R}}\widetilde{\mathbf{F}}_{t}\widetilde{\mathbf{C}}' + \mathbf{E}_{t}$$
$$= \mathbf{R}\mathbf{J}_{1}\mathbf{J}_{1}^{-1}\mathbf{F}_{t}\mathbf{J}_{2}^{-1}\mathbf{J}_{2}'\mathbf{C}' + \mathbf{E}_{t}$$
$$= \mathbf{R}\mathbf{F}_{t}\mathbf{C}' + \mathbf{E}_{t}$$

$$\begin{split} \widetilde{\mathbf{F}}_t &= \widetilde{\mathbf{A}} \widetilde{\mathbf{F}}_{t-1} \widetilde{\mathbf{B}}' + \widetilde{\mathbf{U}}_t \\ \mathbf{J}_1 \widetilde{\mathbf{F}}_t \mathbf{J}_2' &= \mathbf{J}_1 \widetilde{\mathbf{A}} \widetilde{\mathbf{F}}_{t-1} \widetilde{\mathbf{B}}' \mathbf{J}_2' + \mathbf{J}_1 \widetilde{\mathbf{U}}_t \mathbf{J}_2' \\ \mathbf{J}_1 \mathbf{J}_1^{-1} \mathbf{F}_t (\mathbf{J}_2^{-1})' \mathbf{J}_2' &= \mathbf{J}_1 \mathbf{J}_1^{-1} \mathbf{A} \mathbf{J}_1 \mathbf{J}_1^{-1} \mathbf{F}_t (\mathbf{J}_2^{-1})' \mathbf{J}_2' \mathbf{B}' (\mathbf{J}_2^{-1})' \mathbf{J}_2' + \mathbf{J}_1 \mathbf{J}_1^{-1} \mathbf{U}_t (\mathbf{J}_2^{-1})' \mathbf{J}_2' \\ \mathbf{F}_t &= \mathbf{A} \mathbf{F}_{t-1} \mathbf{B}' + \mathbf{U}_t \end{split}$$

Moreover, using Assumption 1(iv), we have

$$\begin{split} \mathbb{E}\Big[\widetilde{\mathbf{U}}_t\widetilde{\mathbf{U}}_t'\Big] &= \mathbb{E}\big[\mathbf{J}_1^{-1}\mathbf{U}_t(\mathbf{J}_2^{-1})'\mathbf{J}_2^{-1}\mathbf{U}_t(\mathbf{J}_1^{-1})'\big] \\ &= \mathbf{J}_1^{-1}\mathbf{P}\mathrm{tr}(\mathbf{Q})(\mathbf{J}_1^{-1})' \\ &= \mathbf{J}_1^{-1}\mathbf{P}(\mathbf{J}_1^{-1})'\mathrm{tr}\left(\mathbf{Q}(\mathbf{J}_2^{-1})'\mathbf{J}_2^{-1}\right) \\ &= \widetilde{\mathbf{P}}\mathrm{tr}\left(\widetilde{\mathbf{Q}}\right) \end{split}$$

and analogous derivation can be obtained for $\mathbb{E}\left[\widetilde{\mathbf{U}}_{t}'\widetilde{\mathbf{U}}_{t}\right]$. The derivation of the vectorized form follows naturally.

Lemma 3. Consider the rotated system (30)-(31) defined in Lemma 2. For any $k_1 \times k_1$ and $k_2 \times k_2$ orthogonal matrices \mathbf{J}_1 and \mathbf{J}_2 , under Assumptions 1-2, we have that

(i)
$$\|\widetilde{\mathbf{A}}\| \le 1$$
, $\|\widetilde{\mathbf{B}}\| < 1$, and $\|\widetilde{\mathbf{B}} \otimes \widetilde{\mathbf{A}}\| < 1$
(ii) $\|\mathbf{H}\| = O(1)$, $\|\mathbf{K}\| = O(1)$, and $\|\mathbf{K} \otimes \mathbf{H}\| = O(1)$
(iii) $p_1^{-1/2} \|\widetilde{\mathbf{R}}\| = O(1)$, and $p_2^{-1/2} \|\widetilde{\mathbf{C}}\| = O(1)$,
(iv) $p_2 \| \left(\widetilde{\mathbf{C}}' \operatorname{dg}(\mathbf{K})^{-1} \widetilde{\mathbf{C}} \right)^{-1} \| = O(1)$ and $p_1 \| \left(\widetilde{\mathbf{R}}' \operatorname{dg}(\mathbf{H})^{-1} \widetilde{\mathbf{R}} \right)^{-1} \| = O(1)$
(v) $\frac{1}{\sqrt{p_2}} \| \widetilde{\mathbf{C}}' \operatorname{dg}(\mathbf{K})^{-1} \| = O(1)$ and $\frac{1}{\sqrt{p_1}} \| \widetilde{\mathbf{R}}' \operatorname{dg}(\mathbf{H})^{-1} \| = O(1)$

Proof. To show (i), note that by Assumption 1(iii),

$$\|\widetilde{\mathbf{A}}\| \leq \|\mathbf{J}_1^{-1}\mathbf{A}\mathbf{J}_1\|_F = \sqrt{\operatorname{tr}\left(\mathbf{J}_1^{-1}\mathbf{A}\mathbf{J}_1\mathbf{J}_1'\mathbf{A}'(\mathbf{J}_1^{-1})'\right)} = \sqrt{\operatorname{tr}\left(\mathbf{A}\mathbf{A}'\right)} = \|\mathbf{A}\| < 1$$

$$\|\widetilde{\mathbf{B}}\| = \|\mathbf{J}_2^{-1}\mathbf{B}\mathbf{J}_2\| < \|\mathbf{J}_2^{-1}\|\|\mathbf{B}\|\|\mathbf{J}_2\| = \|\mathbf{B}\| < 1$$

and $\|\widetilde{\mathbf{B}} \otimes \widetilde{\mathbf{A}}\| \leq \|\widetilde{\mathbf{B}}\| \|\widetilde{\mathbf{A}}\| < 1$. To show (ii), start noticing that since **H** and **K** are symmetric matrices then also $\mathbf{K} \otimes \mathbf{H}$ is symmetric, implying that $\|\mathbf{K} \otimes \mathbf{H}\| = \rho(\mathbf{K} \otimes \mathbf{H}) < \|\mathbf{K} \otimes \mathbf{H}\|_1$. Note that each column of $\mathbf{K} \otimes \mathbf{H}$ can be written as $\mathbf{k}_{\cdot j} \otimes \mathbf{h}_{\cdot i}$ for $j=1,...,p_2$ and $i=1,...,p_1$. By the symmetry of **K** and **H**, we have that $\mathbf{k}_{\cdot j} \otimes \mathbf{h}_{\cdot i} = \mathbf{k}_{j \cdot} \otimes \mathbf{h}_{i \cdot}$ and $\mathbf{k}_{j \cdot} \otimes \mathbf{h}_{i \cdot} = \operatorname{vec}(\mathbb{E}[\mathbf{e}_{\cdot j}\mathbf{e}'_{i \cdot}])$. We conclude $\|\mathbf{K} \otimes \mathbf{H}\|_1 = O(1)$ by Assumption 2(iii). The same conclusion follows for $\|\mathbf{H}\|$ and $\|\mathbf{K}\|$ noticing that $\|\mathbf{K} \otimes \mathbf{H}\| = \|\mathbf{K}\| \|\mathbf{H}\|$. To show (iii), note that by Assumption 1(i)

$$\frac{1}{p_1} \left\| \widetilde{\mathbf{R}} \right\|^2 \approx \frac{1}{p_1} \left\| \mathbf{R} \right\|_F^2 = \frac{1}{p_1} \sum_{i=1}^{p_1} \sum_{j=1}^{k_1} r_{ij}^2 \le k_1 \bar{r} = O(1),$$

and

$$\frac{1}{p_2} \left\| \widetilde{\mathbf{C}} \right\|^2 \approx \frac{1}{p_2} \left\| \mathbf{C} \right\|_F^2 = \frac{1}{p_2} \sum_{i=1}^{p_2} \sum_{j=1}^{k_2} c_{ij}^2 \leq k_2 \bar{c} = O(1).$$

To show (iv), note that by Theorem 1 in Merikoski and Kumar (2004), we have

$$p_{2} \left\| \left(\widetilde{\mathbf{C}}' \mathrm{dg}(\mathbf{K})^{-1} \widetilde{\mathbf{C}} \right)^{-1} \right\| = \frac{p_{2}}{\nu^{(k_{2})} \left(\widetilde{\mathbf{C}}' \mathrm{dg}(\mathbf{K})^{-1} \widetilde{\mathbf{C}} \right)} \\ \leq \frac{p_{2}}{\nu^{(k_{2})} \left(\widetilde{\mathbf{C}}' \widetilde{\mathbf{C}} \right) \nu^{(p_{2})} \left(\mathrm{dg}(\mathbf{K})^{-1} \right)} \\ \leq \frac{p_{2}}{\nu^{(k_{2})} \left(\widetilde{\mathbf{C}}' \widetilde{\mathbf{C}} \right) \left(\nu^{(1)} (\mathrm{dg}(\mathbf{K})) \right)^{-1}} \\ \lesssim \frac{1}{C_{K}^{-1}}$$

by Assumptions 1(*i*) and 2(*ii*). The proof for $p_1 \left\| \left(\widetilde{\mathbf{R}}' dg(\mathbf{H})^{-1} \widetilde{\mathbf{R}} \right)^{-1} \right\|$ follows the same steps. To show (v), note that

$$\frac{1}{\sqrt{p_2}} \left\| \widetilde{\mathbf{C}}' \mathrm{dg}(\mathbf{K})^{-1} \right\| \leq \frac{1}{\sqrt{p_2}} \left\| \widetilde{\mathbf{C}} \right\| \left\| \mathrm{dg}(\mathbf{K})^{-1} \right\|$$
$$\leq \frac{1}{\sqrt{p_2}} \left\| \widetilde{\mathbf{C}} \right\| \max_{j=1,\dots,p_2} k_{ii}^{-1}$$
$$= O(1)$$

by (iii) and Assumption 2(*ii*). The proof for $\frac{1}{\sqrt{p_1}} \left\| \widetilde{\mathbf{R}}' dg(\mathbf{H})^{-1} \right\|$ follows the same steps.

C.2.2 Results on pre-estimators

Lemma 4. Under Assumptions 1 through 3, there exist matrices $\widehat{\mathbf{J}}_1$ and $\widehat{\mathbf{J}}_2$ satisfying $\widehat{\mathbf{J}}_1 \widehat{\mathbf{J}}_1' \xrightarrow{p} \mathbb{I}_{k_1 k_1}$ and $\widehat{\mathbf{J}}_2 \widehat{\mathbf{J}}_2' \xrightarrow{p} \mathbb{I}_{k_2 k_2}$, such that as $\min\{p_1, p_2, T\} \rightarrow \infty$,

(*i*)
$$\frac{1}{\sqrt{p_1}} \left\| \widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right\| = O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2}, \frac{1}{Tp_1} \right\} \right)$$

$$(ii) \quad \frac{1}{\sqrt{p_2}} \left\| \widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1 p_2}, \frac{1}{Tp_2} \right\} \right)$$
$$(iii) \quad \frac{1}{\sqrt{p_1 p_2}} \left\| \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2 \otimes \mathbf{R} \widehat{\mathbf{J}}_1 \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right)$$

Proof. Note that (i) and (ii) follows immediately from Theorem 3.1 in Yu et al. (2022). Moreover, consider (iii) and note that

$$\begin{split} \frac{1}{\sqrt{p_1 p_2}} \left\| \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2 \otimes \mathbf{R} \widehat{\mathbf{J}}_1 \right\| &\leq \frac{1}{\sqrt{p_1 p_2}} \left\| \left(\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right) \otimes \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right) \right\| \\ &\quad + \frac{1}{\sqrt{p_1 p_2}} \left\| \left(\widehat{\mathbf{C}} \widehat{\mathbf{J}}_2 \right) \otimes \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right) \right\| \\ &\quad + \frac{1}{\sqrt{p_1 p_2}} \left\| \left(\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right) \otimes \left(\mathbf{R} \widehat{\mathbf{J}}_1 \right) \right\| \\ &\leq \left\| \frac{\mathbf{C} \widehat{\mathbf{J}}_2}{\sqrt{p_2}} \right\| \left\| \frac{\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1}{\sqrt{p_1}} \right\| + \left\| \frac{\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2}{\sqrt{p_2}} \right\| \left\| \frac{\mathbf{R} \widehat{\mathbf{J}}_1}{\sqrt{p_1}} \right\| \\ &= O_p \left(\max \left\{ \frac{1}{\sqrt{T p_2}}, \frac{1}{p_1 p_2}, \frac{1}{T p_1} \right\} \right) + O_p \left(\max \left\{ \frac{1}{\sqrt{T p_1}}, \frac{1}{p_1 p_2}, \frac{1}{T p_2} \right\} \right) \end{split}$$

by (i), (ii) and Lemma 3(iii).

Lemma 5. Under Assumptions 1 through 3, there exist matrices $\widehat{\mathbf{J}}_1$ and $\widehat{\mathbf{J}}_2$ satisfying $\widehat{\mathbf{J}}_1 \widehat{\mathbf{J}}_1' \xrightarrow{p} \mathbb{I}_{k_1 k_1}$ and $\widehat{\mathbf{J}}_2 \widehat{\mathbf{J}}_2' \xrightarrow{p} \mathbb{I}_{k_2 k_2}$, such that for $s_1, s_2 \in \{0, 1\}$ as $\min\{p_1, p_2, T\} \rightarrow \infty$,

$$(i) \quad \frac{1}{T} \sum_{t}^{T} \left(\left(\frac{\widehat{\mathbf{C}}^{(0)}}{p_{2}} \otimes \frac{\widehat{\mathbf{R}}^{(0)}}{p_{1}} \right)' \mathsf{y}_{t-s_{1}} - \widehat{\mathbf{J}}^{-1} \mathsf{f}_{t-s_{1}} \right) \mathsf{f}_{t-s_{2}}' \widehat{\mathbf{J}}^{-1\prime} = O_{p} \left(\max\left\{ \frac{1}{\sqrt{T}}, \frac{1}{p_{1}p_{2}} \right\} \right),$$
$$(ii) \quad \frac{1}{T} \sum_{t}^{T} \left(\left(\frac{\widehat{\mathbf{C}}^{(0)}}{p_{2}} \otimes \frac{\widehat{\mathbf{R}}^{(0)}}{p_{1}} \right)' \mathsf{y}_{t-s_{1}} - \widehat{\mathbf{J}}^{-1} \mathsf{f}_{t-s_{1}} \right) \mathsf{u}_{t}' = O_{p} \left(\max\left\{ \frac{1}{\sqrt{T}}, \frac{1}{p_{1}p_{2}} \right\} \right),$$
$$with \quad \widehat{\mathbf{J}} = \widehat{\mathbf{J}}_{2} \otimes \widehat{\mathbf{J}}_{1}.$$

Proof. Note that the left hand side of (i) can be written as

$$\frac{1}{T}\sum_{t=2}^{T}\left(\operatorname{vec}\left(\frac{\widehat{\mathbf{R}}^{(0)\prime}}{p_{1}}\mathbf{R}\widehat{\mathbf{J}}_{1}\widehat{\mathbf{J}}_{1}^{-1}\mathbf{F}_{t-s_{1}}\widehat{\mathbf{J}}_{2}^{-1\prime}\widehat{\mathbf{J}}_{2}^{\prime}\mathbf{C}^{\prime}\frac{\widehat{\mathbf{C}}^{(0)}}{p_{2}}+\frac{\widehat{\mathbf{R}}^{(0)\prime}}{p_{1}}\mathbf{E}_{t}\frac{\widehat{\mathbf{C}}^{(0)}}{p_{2}}\right)-\widehat{\mathbf{J}}^{-1}\mathbf{f}_{t-s_{1}}\right)\mathbf{f}_{t-s_{2}}^{\prime}\widehat{\mathbf{J}}^{-1\prime}$$
(34)

By Theorem 3.1 in Yu et al. (2022), we have that

$$p_1^{-1}\widehat{\mathbf{R}}^{(0)\prime}\mathbf{R} = \widehat{\mathbf{J}}_1' + o_p(1), \qquad p_2^{-1}\widehat{\mathbf{C}}^{(0)\prime}\mathbf{C} = \widehat{\mathbf{J}}_2' + o_p(1), \tag{35}$$

therefore (34) is asymptotically equivalent to

$$\frac{1}{T} \sum_{t=2}^{T} \left(\frac{\widehat{\mathbf{C}}^{(0)}}{p_2} \otimes \frac{\widehat{\mathbf{R}}^{(0)}}{p_1} \right)' \mathbf{e}_{t-s_1} \mathbf{f}'_{t-s_2} \widehat{\mathbf{J}}^{-1'}.$$
(36)

We can bound (36) as follows

$$\begin{split} \frac{1}{T} \sum_{t=2}^{T} \left(\frac{\widehat{\mathbf{C}}^{(0)}}{p_2} \otimes \frac{\widehat{\mathbf{R}}^{(0)}}{p_1} \right)' \mathbf{e}_t \mathbf{f}'_t \widehat{\mathbf{J}}^{-1\prime} &\lesssim \quad \frac{1}{T\sqrt{p_1 p_2}} \sum_{t=2}^{T} \left(\frac{\widehat{\mathbf{C}}^{(0)}}{\sqrt{p_2}} \otimes \frac{\widehat{\mathbf{R}}^{(0)}}{\sqrt{p_1}} - \frac{\mathbf{C}\widehat{\mathbf{J}}_2}{\sqrt{p_2}} \otimes \frac{\mathbf{R}\widehat{\mathbf{J}}_1}{\sqrt{p_1}} \right)' \mathbf{e}_t \mathbf{f}'_t \widehat{\mathbf{J}}^{-1\prime} \\ &+ \frac{1}{T\sqrt{p_1 p_2}} \sum_{t=2}^{T} \left(\frac{\mathbf{C}\widehat{\mathbf{J}}_2}{\sqrt{p_2}} \otimes \frac{\mathbf{R}\widehat{\mathbf{J}}_1}{\sqrt{p_1}} \right)' \mathbf{e}_t \mathbf{f}'_t \widehat{\mathbf{J}}^{-1\prime} \\ &\lesssim \quad O_p \left(\max\left\{ \frac{1}{\sqrt{T}}, \frac{1}{p_1 p_2} \right\} \right). \end{split}$$

by Assumption 3(i), and Lemmas 3 and 4. Consider (ii) and note that by (35), we have

$$\begin{split} \frac{1}{T} \sum_{t}^{T} \left(\left(\frac{\widehat{\mathbf{C}}^{(0)}}{p_{2}} \otimes \frac{\widehat{\mathbf{R}}^{(0)}}{p_{1}} \right)' \mathsf{y}_{t-s_{1}} - \widehat{\mathbf{J}}^{-1} \mathsf{f}_{t-s_{1}} \right) \mathsf{u}_{t}' &\lesssim \quad \frac{1}{T\sqrt{p_{1}p_{2}}} \sum_{t=2}^{T} \left(\frac{\widehat{\mathbf{C}}^{(0)}}{\sqrt{p_{2}}} \otimes \frac{\widehat{\mathbf{R}}^{(0)}}{\sqrt{p_{1}}} \right)' \mathsf{e}_{t} \mathsf{u}_{t}' \\ &\lesssim \quad O_{p} \left(\max\left\{ \frac{1}{\sqrt{T}}, \frac{1}{p_{1}p_{2}} \right\} \right). \end{split}$$

by Lemmas 3 and 4, and since We can bound (36) as follows

$$\begin{split} \mathbb{E}\left[\left\|\frac{1}{T\sqrt{p_{1}p_{2}}}\sum_{t=2}^{T}\mathsf{e}_{t}\mathsf{u}_{t}'\right\|^{2}\right] &\leq \frac{1}{T^{2}p_{1}p_{2}}\sum_{i_{1}=1}^{p_{1}}\sum_{i_{2}=1}^{p_{2}}\sum_{j_{1}=1}^{k_{1}}\sum_{j_{2}=1}^{k_{2}}\sum_{t,s=1}^{T}\mathbb{E}[e_{ti_{1}i_{2}}e_{si_{1}i_{2}}u_{tj_{1}j_{2}}u_{sj_{1}j_{2}}] \\ &\leq \frac{kC_{P}C_{Q}}{T^{2}p_{1}p_{2}}\sum_{i_{1}=1}^{p_{1}}\sum_{i_{2}=1}^{p_{2}}\sum_{t,s=1}^{T}|\mathbb{E}[e_{ti_{1}i_{2}}e_{si_{1}i_{2}}]| \\ &\lesssim O_{p}\left(\frac{1}{T}\right). \end{split}$$

by Assumptions 1(iv) and 2(iii).

Lemma 6. Under Assumptions 1 through 3, there exist matrices $\widehat{\mathbf{J}}_1$ and $\widehat{\mathbf{J}}_2$ satisfying $\widehat{\mathbf{J}}_1 \widehat{\mathbf{J}}_1' \xrightarrow{p} \mathbb{I}_{k_1}$ and $\widehat{\mathbf{J}}_2 \widehat{\mathbf{J}}_2' \xrightarrow{p} \mathbb{I}_{k_2}$, such that as $\min\{p_1, p_2, T\} \rightarrow \infty$,

(i)
$$\left\|\widehat{\mathbf{B}\otimes\mathbf{A}}^{(0)} - \widehat{\mathbf{J}}^{-1}(\mathbf{B}\otimes\mathbf{A})\widehat{\mathbf{J}}\right\| = O_p\left(\max\left\{\frac{1}{\sqrt{T}}, \frac{1}{p_1p_2}\right\}\right)$$

(ii) $\left\|\widehat{\mathbf{Q}\otimes\mathbf{P}}^{(0)} - \widehat{\mathbf{J}}^{-1}(\mathbf{Q}\otimes\mathbf{P})\widehat{\mathbf{J}}\right\| = O_p\left(\max\left\{\frac{1}{\sqrt{T}}, \frac{1}{p_1p_2}\right\}\right)$,
with $\widehat{\mathbf{J}} = \widehat{\mathbf{J}}_2 \otimes \widehat{\mathbf{J}}_1$.

Proof. Consider (i) and note that

$$\begin{split} \left\| \widehat{\mathbf{B} \otimes \mathbf{A}}^{(0)} - \widehat{\mathbf{J}}^{-1} (\mathbf{B} \otimes \mathbf{A}) \widehat{\mathbf{J}} \right\| &\leq \\ \left\| \left(\frac{1}{T} \sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t} \widetilde{\mathbf{f}}_{t-1}' - \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{f}_{t-1}' \widehat{\mathbf{J}}^{-\prime} \right) \left(\frac{1}{T} \sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t-1} \widetilde{\mathbf{f}}_{t-1}' \right)^{-1} \right\| \\ &+ \\ \left\| \widehat{\mathbf{J}}^{-1} (\mathbf{B} \otimes \mathbf{A}) \widehat{\mathbf{J}} \left(\frac{1}{T} \sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t-1} \widetilde{\mathbf{f}}_{t-1}' - \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t-1} \mathbf{f}_{t-1}' \widehat{\mathbf{J}}^{-\prime} \right) \left(\frac{1}{T} \sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t-1} \widetilde{\mathbf{f}}_{t-1}' \right)^{-1} \\ &+ \\ \\ &+ \\ \left\| \left(\frac{1}{T} \sum_{t=2}^{T} \mathbf{u}_{t} \widetilde{\mathbf{f}}_{t-1}' \right) \left(\frac{1}{T} \sum_{t=2}^{T} \widetilde{\mathbf{f}}_{t-1} \widetilde{\mathbf{f}}_{t-1}' \right)^{-1} \right\| . \end{split}$$

| | | - | |
|---|--|---|--|
| L | | | |
| 1 | | | |
| 1 | | | |
| | | | |

where $\tilde{\mathsf{f}}_t = \left(\frac{\hat{\mathbf{C}}^{(0)}}{p_2} \otimes \frac{\hat{\mathbf{R}}^{(0)}}{p_1}\right)' \mathsf{y}_t$. Now, for $s = \{0, 1\},$

$$\frac{1}{T}\sum_{t=2}^{T}\tilde{\mathbf{f}}_{t-s}\tilde{\mathbf{f}}_{t-1}' - \frac{1}{T}\sum_{t=2}^{T}\hat{\mathbf{J}}^{-1}\mathbf{f}_{t-s}\mathbf{f}_{t-1}'\hat{\mathbf{J}}^{-\prime} = \frac{1}{T}\sum_{t=2}^{T}\left(\tilde{\mathbf{f}}_{t-s}-\hat{\mathbf{J}}^{-1}\mathbf{f}_{t-s}\right)\mathbf{f}_{t-1}'\hat{\mathbf{J}}^{-\prime} + \frac{1}{T}\sum_{t=2}^{T}\hat{\mathbf{J}}^{-1}\mathbf{f}_{t-s}\left(\tilde{\mathbf{f}}_{t-1}-\hat{\mathbf{J}}^{-1}\mathbf{f}_{t-1}\right)' \\ + \frac{1}{T}\sum_{t=2}^{T}\left(\tilde{\mathbf{f}}_{t-s}-\hat{\mathbf{J}}^{-1}\mathbf{f}_{t-s}\right)\left(\tilde{\mathbf{f}}_{t-1}-\hat{\mathbf{J}}^{-1}\mathbf{f}_{t-1}\right)'.$$

Note that the first two terms dominate the third one. By Lemma 5(i) we have that

$$\left\| \frac{1}{T} \sum_{t=2}^{T} \tilde{\mathbf{f}}_{t} \tilde{\mathbf{f}}_{t-1}' - \frac{1}{T} \sum_{t=2}^{T} \mathbf{\hat{J}}^{-1} \mathbf{f}_{t} \mathbf{f}_{t-1}' \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{T}}, \frac{1}{p_{1}p_{2}} \right\} \right) \\ \left\| \frac{1}{T} \sum_{t=2}^{T} \tilde{\mathbf{f}}_{t-1} \tilde{\mathbf{f}}_{t-1}' - \frac{1}{T} \sum_{t=2}^{T} \mathbf{\hat{J}}^{-1} \mathbf{f}_{t-1} \mathbf{f}_{t-1}' \mathbf{\hat{J}}^{-\prime} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{T}}, \frac{1}{p_{1}p_{2}} \right\} \right).$$

Moreover, combining the latter with Assumption 1(ii), we have that,

$$\left(\frac{1}{T}\sum_{t=2}^{T}\widetilde{\mathsf{f}}_{t-1}\widetilde{\mathsf{f}}_{t-1}'\right)^{-1} = O_p(1).$$

Finally, note that

$$\left\|\frac{1}{T}\sum_{t=2}^{T}\mathsf{u}_{t}\widetilde{\mathsf{f}}_{t-1}'\right\| \leq \left\|\frac{1}{T}\sum_{t=2}^{T}\mathsf{u}_{t}\left(\widetilde{\mathsf{f}}_{t-1}-\mathsf{f}_{t-1}\right)\right\| + \left\|\frac{1}{T}\sum_{t=2}^{T}\mathsf{u}_{t}\mathsf{f}_{t-1}\right\| = O_{p}\left(\min\left\{\frac{1}{\sqrt{T}},\frac{1}{p_{1}p_{2}}\right\}\right),$$

by Lemma 5(ii) and Assumption 1(iv), concluding the proof. The result for (ii) follows using the same steps of (i) noting that

$$\begin{split} \widehat{\mathbf{Q} \otimes \mathbf{P}}^{(0)} &= \frac{1}{T} \sum_{t=2}^{T} \left\{ \widetilde{\mathbf{f}}_{t} - \widehat{\mathbf{B} \otimes \mathbf{A}} \widetilde{\mathbf{f}}_{t-1} \right\} \left\{ \widetilde{\mathbf{f}}_{t} - \widehat{\mathbf{B} \otimes \mathbf{A}} \widetilde{\mathbf{f}}_{t-1} \right\}' \\ &= \frac{1}{T} \sum_{t=2}^{T} \left\{ \left(\widetilde{\mathbf{f}}_{t} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right) + \left(\widehat{\mathbf{B} \otimes \mathbf{A}} - \mathbf{B} \otimes \mathbf{A} \right) \left(\widetilde{\mathbf{f}}_{t-1} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t-1} \right) \right. \\ &+ \left(\widehat{\mathbf{B} \otimes \mathbf{A}} - \mathbf{B} \otimes \mathbf{A} \right) \widetilde{\mathbf{f}}_{t-1} + \mathbf{B} \otimes \mathbf{A} \left(\widetilde{\mathbf{f}}_{t-1} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t-1} \right) + \widehat{\mathbf{J}}^{-1} \mathbf{u}_{t} \right\} \\ &\left\{ \left(\widetilde{\mathbf{f}}_{t} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right) + \left(\widehat{\mathbf{B} \otimes \mathbf{A}} - \mathbf{B} \otimes \mathbf{A} \right) \left(\widetilde{\mathbf{f}}_{t-1} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t-1} \right) \right. \\ &+ \left(\widehat{\mathbf{B} \otimes \mathbf{A}} - \mathbf{B} \otimes \mathbf{A} \right) \widetilde{\mathbf{f}}_{t-1} + \mathbf{B} \otimes \mathbf{A} \left(\widetilde{\mathbf{f}}_{t-1} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t-1} \right) \\ &+ \left(\widehat{\mathbf{B} \otimes \mathbf{A}} - \mathbf{B} \otimes \mathbf{A} \right) \widetilde{\mathbf{f}}_{t-1} + \mathbf{B} \otimes \mathbf{A} \left(\widetilde{\mathbf{f}}_{t-1} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t-1} \right) + \widehat{\mathbf{J}}^{-1} \mathbf{u}_{t} \right\}' \end{split}$$

and that

$$\left\|\frac{1}{T}\sum_{t=1}^{T}\mathsf{u}_{t}\mathsf{u}_{t}'-\mathbf{Q}\otimes\mathbf{P}\right\|=O_{p}\left(\frac{1}{\sqrt{T}}\right)$$

by Assumption 1(iv).

Lemma 7. Under Assumption (1) through (3), as $\min\{p_1, p_2, T\} \rightarrow \infty$,

$$\begin{array}{l} (i) \ \left| \hat{k}_{jj}^{(0)} - k_{jj} \right| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1 p_2}, \frac{1}{Tp_2}, \right\} \right) \ uniformly \ in \ j \\ (ii) \ \left| \hat{h}_{ii}^{(0)} - h_{ii} \right| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2}, \frac{1}{Tp_1} \right\} \right) \ uniformly \ in \ i \\ (iii) \ \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left(\hat{k}_{jj}^{(0)} - k_{jj} \right) \right| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1 p_2}, \frac{1}{Tp_2} \right\} \right) \\ (iv) \ \frac{1}{p_1} \left| \sum_{i=1}^{p_1} \left(\hat{h}_{ii}^{(0)} - h_{ii} \right) \right| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2}, \frac{1}{Tp_1} \right\} \right) \end{array}$$

Proof. Start from (i), and recall by Assumption 2(ii) we have that $tr(\mathbf{H}) = p_1$, thus $\mathbf{K} = \frac{1}{p_1} \mathbb{E}[\mathbf{E}'_t \mathbf{E}_t]$. We can then write

$$\begin{aligned} \widehat{k}_{jj}^{(0)} - k_{jj} \middle| &= \left| \frac{1}{Tp_1} \sum_{t=1}^{T} \sum_{i=1}^{p_1} (\widehat{e}_{tij}^{(0)})^2 - \frac{1}{p_1} \sum_{i=1}^{p_1} \mathbb{E}\left[e_{tij}^2\right] \right| \\ &= \left| \frac{1}{Tp_1} \sum_{t=1}^{T} \sum_{i=1}^{p_1} \left((s_{tij} - \widehat{s}_{tij}^{(0)})^2 + 2(s_{tij} - \widehat{s}_{tij}^{(0)})e_{tij} + e_{tij}^2 \right) - \frac{1}{p_1} \sum_{i=1}^{p_1} \mathbb{E}\left[e_{tij}^2\right] \right| \\ &\leq \left| \frac{1}{Tp_1} \sum_{t=1}^{T} \sum_{i=1}^{p_1} (s_{tij} - \widehat{s}_{tij}^{(0)})^2 \right| + 2 \left| \frac{1}{Tp_1} \sum_{t=1}^{T} \sum_{i=1}^{p_1} (s_{tij} - \widehat{s}_{tij}^{(0)})e_{tij} \right| \\ &+ \left| \frac{1}{Tp_1} \sum_{t=1}^{T} \sum_{i=1}^{p_1} e_{tij}^2 - \frac{1}{p_1} \sum_{i=1}^{p_1} \mathbb{E}\left[e_{tij}^2\right] \right| \end{aligned}$$

Consider the first addendum, we have

$$\begin{aligned} \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} (s_{tij} - \widehat{s}_{tij}^{(0)})^{2} \right| &= \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} (\mathbf{s}_{t\cdot j} - \widehat{\mathbf{s}}_{t\cdot j}^{(0)})^{2} \right| \\ &\lesssim \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j} \cdot \mathbf{F}_{t}' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_{1} \right)' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_{1} \right) \mathbf{F}_{t} \mathbf{c}_{j} \right| \\ &+ \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \mathbf{F}_{t}' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_{1} \right)' \mathbf{R} \left(\widetilde{\mathbf{F}}_{t} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2} \right) \mathbf{c}_{j} \right| \\ &+ \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \mathbf{F}_{t}' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_{1} \right)' \mathbf{R} \mathbf{F}_{t} \left(\widehat{\mathbf{c}}_{j} - \mathbf{c}_{j}' \cdot \widehat{\mathbf{J}}_{2} \right) \right| \\ &+ \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j} \cdot \left(\widetilde{\mathbf{F}}_{t} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2} \right)' \mathbf{R}' \mathbf{R} \left(\widetilde{\mathbf{F}}_{t} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2} \right) \mathbf{c}_{j} \right| \\ &+ \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j} \cdot \left(\widetilde{\mathbf{F}}_{t} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2} \right)' \mathbf{R}' \mathbf{R} \mathbf{F}_{t} \left(\widehat{\mathbf{c}}_{j} - \mathbf{c}_{j}' \cdot \widehat{\mathbf{J}}_{2} \right) \right| \\ &+ \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j} \cdot \left(\widetilde{\mathbf{F}}_{t} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2} \right)' \mathbf{R}' \mathbf{R} \mathbf{F}_{t} \left(\widehat{\mathbf{c}}_{j} - \mathbf{c}_{j}' \cdot \widehat{\mathbf{J}}_{2} \right) \right| \\ &+ \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \left(\widehat{\mathbf{c}}_{j} - \mathbf{c}_{j}' \cdot \widehat{\mathbf{J}}_{2} \right) \mathbf{F}_{t}' \mathbf{R}' \mathbf{R} \mathbf{F}_{t} \left(\widehat{\mathbf{c}}_{j} - \mathbf{c}_{j}' \cdot \widehat{\mathbf{J}}_{2} \right) \right| \\ &= O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}p_{2}}}, \frac{1}{\sqrt{Tp_{1}p_{2}}}, \frac{1}{\sqrt{Tp_{1}p_{2}}}, \frac{1}{Tp_{1}}, \frac{1}{Tp_{1}} \right\} \right) \end{aligned}$$

since

$$\frac{1}{Tp_1} \sum_{t=1}^{T} \mathbf{c}_{j \cdot} \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right) \mathbf{F}_t \mathbf{c}_{j \cdot} \bigg| \lesssim \frac{1}{p_1} \left\| \widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right\|^2 \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{F}_t' \mathbf{F}_t \right\|$$
$$= O_p \left(\max \left\{ \frac{1}{Tp_2}, \frac{1}{p_1^2 p_2^2}, \frac{1}{T^2 p_1^2} \right\} \right)$$

because of Lemma 4(i) and Assumption 1(ii)

$$\begin{aligned} \left| \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j\cdot}' \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{R} \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \mathbf{c}_{j\cdot} \right| &\lesssim \quad \frac{1}{p_1} \left\| \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{R} \right\| \\ &\times \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{F}_t' \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \right\| \\ &= \quad O_p \left(\frac{1}{Tp_1 p_2} \right) \end{aligned}$$

because of Lemma E.1 in Yu et al. (2022) and since

$$\begin{aligned} \widetilde{\mathbf{F}}_{t} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-1} &= \frac{1}{p_{1} p_{2}} \widehat{\mathbf{R}}^{(0)'} \left(\mathbf{R} - \widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{J}}_{1}^{-1} \right) \mathbf{F}_{t} \left(\mathbf{C} - \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{J}}_{2}^{-1} \right)' \widehat{\mathbf{C}}^{(0)} \\ &+ \frac{1}{p_{1}} \widehat{\mathbf{R}}^{(0)'} \left(\mathbf{R} - \widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{J}}_{1}^{-1} \right) \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} + \frac{1}{p_{2}} \widehat{\mathbf{J}}^{-1} \mathbf{F}_{t} \left(\mathbf{C} - \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{J}}_{2}^{-1} \right)' \widehat{\mathbf{C}}^{(0)} \\ &+ \frac{1}{p_{1} p_{2}} \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_{1} \right)' \mathbf{E}_{t} \left(\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right) \\ &+ \frac{1}{p_{1} p_{2}} \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_{1} \right)' \mathbf{E}_{t} \mathbf{C} \widehat{\mathbf{J}}_{2} + \frac{1}{p_{1} p_{2}} \widehat{\mathbf{J}}_{1}' \mathbf{R}' \mathbf{E}_{t} \left(\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right) \\ &+ \frac{1}{p_{1} p_{2}} \widehat{\mathbf{J}}_{1}' \mathbf{R}' \mathbf{E}_{t} \mathbf{C} \widehat{\mathbf{J}}_{2} \\ &\leq \frac{1}{p_{1} p_{2}} \mathbf{R}' \mathbf{E}_{t} \mathbf{C} \end{aligned}$$
(38)

we have that

$$\left|\frac{1}{T}\sum_{t=1}^{T}\mathbf{F}_{t}'\left(\widetilde{\mathbf{F}}_{t}-\widehat{\mathbf{J}}_{1}^{-1}\mathbf{F}_{t}\widehat{\mathbf{J}}_{2}\right)\right\|\lesssim\frac{1}{Tp_{1}p_{2}}\sum_{t=1}^{T}\mathbf{F}_{t}'\mathbf{R}'\mathbf{E}_{t}\mathbf{C}=O_{p}\left(\frac{1}{\sqrt{Tp_{1}p_{2}}}\right)$$
(39)

by Lemma A.1 in Yu et al. (2022) and Chebyshev inequality,

$$\begin{aligned} \left| \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}'_{j\cdot} \mathbf{F}'_t \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{R} \mathbf{F}_t \left(\widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}'_{j\cdot} \widehat{\mathbf{J}}_2 \right) \right| &\lesssim \quad \frac{1}{p_1} \left\| \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{R} \right\| \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{F}'_t \mathbf{F}_t \right\| \left\| \widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}'_{j\cdot} \widehat{\mathbf{J}}_2 \right\| \\ &= \quad O_p \left(\frac{1}{Tp_1} \frac{1}{Tp_2}, \frac{1}{p_1p_2}, \frac{1}{\sqrt{Tp_1p_2}} \right) O_p \left(\frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2} \frac{1}{Tp_2} \right) \end{aligned}$$

because of Theorem 3.1 and Lemma E.1 in Yu et al. (2022) and Assumption 1(ii),

$$\begin{aligned} \left| \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j\cdot} \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \mathbf{R}' \mathbf{R} \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \mathbf{c}_{j\cdot} \right| &\lesssim \left\| \frac{1}{Tp_1} \sum_{t=1}^T \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \right\| \\ &\lesssim \left\| \frac{1}{Tp_1^2 p_2^2} \sum_{t=1}^T \mathbf{C}' \mathbf{E}'_t \mathbf{R} \mathbf{R}' \mathbf{E}_t \mathbf{C} \right\| \\ &= \frac{1}{Tp_1^2 p_2^2} \| \mathbf{C} \| \| \mathbf{R} \| \sum_{i=1}^{p_1} \sum_{j=1}^{p_2} \left\| \sum_{t=1}^T \mathbf{C}' \mathbf{E}_{ti\cdot} \mathbf{E}'_{t\cdot j} \mathbf{R} \right\| \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1 p_2}}, \frac{1}{p_1 p_2} \right\} \right) \end{aligned}$$

because of (38) and (A.3) in Yu et al. (2022),

$$\begin{aligned} \left| \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j\cdot} \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \mathbf{R}' \mathbf{R} \mathbf{F}_t \left(\widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}'_{j\cdot} \widehat{\mathbf{J}}_2 \right) \right| &\lesssim & \left\| \frac{1}{T} \sum_{t=1}^T \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \mathbf{F}_t \right\| \left\| \widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}'_{j\cdot} \widehat{\mathbf{J}}_2 \right\| \\ & \left\| \frac{1}{Tp_1p_2} \sum_{t=1}^T \mathbf{C}' \mathbf{E}'_t \mathbf{R} \mathbf{F}_t \right\| \left\| \widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}'_{j\cdot} \widehat{\mathbf{J}}_2 \right\| \\ &= & O_p \left(\frac{1}{\sqrt{Tp_1p_2}} \right) O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right) \end{aligned}$$

because of (39) and Theorem 3.1 in Yu et al. (2022),

$$\begin{aligned} \left| \frac{1}{Tp_1} \sum_{t=1}^{T} \left(\widehat{\mathbf{c}}_{j \cdot} - \mathbf{c}_{j \cdot}' \widehat{\mathbf{J}}_2 \right) \mathbf{F}_t' \mathbf{R}' \mathbf{R} \mathbf{F}_t \left(\widehat{\mathbf{c}}_{j \cdot} - \mathbf{c}_{j \cdot}' \widehat{\mathbf{J}}_2 \right) \right| &\lesssim \left\| \widehat{\mathbf{c}}_{j \cdot} - \mathbf{c}_{j \cdot}' \widehat{\mathbf{J}}_2 \right\|^2 \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{F}_t' \mathbf{F}_t \right\| \\ &= o_p \left(\max \left\{ \frac{1}{Tp_1}, \frac{1}{p_1^2 p_2^2}, \frac{1}{T^2 p_2^2} \right\} \right) \end{aligned}$$

because of Assumption 1(ii) and Theorem 3.1 in Yu et al. (2022). Consider now the second addendum,

$$\begin{aligned} \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} (s_{tij} - \widehat{s}_{tij}^{(0)}) e_{tij} \right| &= \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} (\mathbf{s}_{t\cdot j} - \widehat{\mathbf{s}}_{t\cdot j}^{(0)}) \mathbf{e}_{t\cdot j} \right| \\ &\lesssim \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \mathbf{F}_{t}' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_{1} \right)' \mathbf{e}_{t\cdot j} \right| \\ &+ \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \left(\widetilde{\mathbf{F}}_{t} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2} \right)' \mathbf{R}' \mathbf{e}_{t\cdot j} \right| \\ &+ \left| \frac{1}{Tp_{1}} \sum_{t=1}^{T} \left(\widehat{\mathbf{c}}_{j} \cdot - \mathbf{c}_{j} \cdot \widehat{\mathbf{J}}_{2} \right)' \mathbf{F}_{t}' \mathbf{R}' \mathbf{e}_{t\cdot j} \right| \end{aligned} \tag{40}$$

since

$$\left| \frac{1}{Tp_1} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{e}_{t \cdot j} \right| \lesssim O_p \left(\max \left\{ \frac{1}{\sqrt{T}p_1}, \frac{1}{Tp_2} \right\} \right)$$

by the same steps as in the proof of Lemma B.3 in Yu et al. (2022),

$$\begin{aligned} \left| \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j\cdot}' \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \mathbf{R}' \mathbf{e}_{t\cdot j} \right| &\lesssim & \left\| \frac{1}{Tp_1^2 p_2} \sum_{t=1}^T \mathbf{C}' \mathbf{E}_t' \mathbf{R} \mathbf{R}' \mathbf{e}_{t\cdot j} \right\| \\ &= & \left\| \frac{1}{Tp_1^2 p_2} \sum_{t=1}^T \sum_{i=1}^{p_1} [\mathbf{C}' \mathbf{E}_t']_{.i} \mathbf{r}_{i\cdot}' \mathbf{R}' \mathbf{e}_{t\cdot j} \right\| \\ &\lesssim & \frac{1}{p_1} \sum_{i=1}^{p_1} \left\| \frac{1}{Tp_1 p_2} \sum_{t=1}^T \mathbf{C}' \mathbf{e}_{t.i} \mathbf{e}_{t\cdot j}' \mathbf{R} \right\| \\ &= & O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1 p_2}}, \frac{1}{p_1 p_2} \right\} \right) \end{aligned}$$

by (A.3) in Yu et al. (2022), and

$$\begin{aligned} \frac{1}{Tp_1} \sum_{t=1}^T \left(\widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}_{j\cdot} \widehat{\mathbf{J}}_2 \right)' \mathbf{F}'_t \mathbf{R}' \mathbf{e}_{t\cdot j} \middle| &\leq \left\| \widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}_{j\cdot} \widehat{\mathbf{J}}_2 \right\| \left\| \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{F}'_t \mathbf{R}' \mathbf{e}_{t\cdot j} \right\| \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right) O_p \left(\frac{1}{\sqrt{Tp_1}} \right) \end{aligned}$$

by Theorem 3.1 and assumption 3(i). Finally, note that

$$\mathbb{E}\left[\left(\frac{1}{Tp_{1}}\sum_{t=1}^{T}\sum_{i=1}^{p_{1}}e_{tij}^{2}-\mathbb{E}\left[e_{tij}^{2}\right]\right)^{2}\right] = \frac{1}{T^{2}p_{1}^{2}}\sum_{t,s}^{T}\sum_{i_{1},i_{2}}^{p_{1}}\mathbb{E}\left[\left(e_{ti_{1}j}^{2}-\mathbb{E}\left[e_{ti_{1}j}^{2}\right]\right)\left(e_{si_{2}j}^{2}-\mathbb{E}\left[e_{si_{2}j}^{2}\right]\right)\right] \\
\leq \frac{1}{T^{2}p_{1}^{2}}\sum_{t,s}^{T}\sum_{i_{1},i_{2}}^{p_{1}}\mathbb{C}\left[e_{ti_{1}j}^{2},e_{si_{2}j}^{2}\right] \\
= O\left(\frac{1}{Tp_{1}}\right)$$
(41)

by Assumption 2(iv). Therefore, for the third addendum, we have

$$\left|\frac{1}{Tp_1}\sum_{t=1}^{T}\sum_{i=1}^{p_1}e_{tij}^2 - \frac{1}{p_1}\sum_{i=1}^{p_1}\mathbb{E}\left[e_{tij}^2\right]\right| = O_p\left(\frac{1}{\sqrt{Tp_1}}\right)$$
(42)

Combining (37), (40), and (42) yields the desired result. Part (ii) follows by similar steps.

Consider (iii), we have that

$$\begin{aligned} \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left(\widehat{k}_{jj}^{(0)} - k_{jj} \right) \right| &= \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \sum_{i=1}^{p_1} (\widehat{e}_{tij}^{(0)})^2 - \frac{1}{p_1} \sum_{i=1}^{p_1} \mathbb{E} \left[e_{tij}^2 \right] \right\} \right| \\ &\leq \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \sum_{i=1}^{p_1} (s_{tij} - \widehat{s}_{tij}^{(0)})^2 \right\} \right| + \frac{2}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \sum_{i=1}^{p_1} (s_{tij} - \widehat{s}_{tij}^{(0)})^2 \right\} \right| \\ &+ \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \sum_{i=1}^{p_1} e_{tij}^2 - \frac{1}{p_1} \sum_{i=1}^{p_1} \mathbb{E} \left[e_{tij}^2 \right] \right\} \right| \end{aligned}$$

Consider the first addendum, we have that

$$\frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} (s_{tij} - \hat{s}_{tij}^{(0)})^{2} \right\} \right| = \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j} \cdot \mathbf{F}_{t}' (\hat{\mathbf{R}}^{(0)} - \mathbf{R}\hat{\mathbf{J}}_{1})' (\hat{\mathbf{R}}^{(0)} - \mathbf{R}\hat{\mathbf{J}}_{1}) \mathbf{F}_{t} \mathbf{c}_{j} \cdot \mathbf{F}_{j} \right| \\
\lesssim \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j} \cdot \mathbf{F}_{t}' (\hat{\mathbf{R}}^{(0)} - \mathbf{R}\hat{\mathbf{J}}_{1})' (\mathbf{R}^{(0)} - \mathbf{R}\hat{\mathbf{J}}_{1}) \mathbf{F}_{t} \mathbf{c}_{j} \cdot \mathbf{F}_{j} \right\} \right| \\
+ \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \mathbf{F}_{t}' (\hat{\mathbf{R}}^{(0)} - \mathbf{R}\hat{\mathbf{J}}_{1})' \mathbf{R} (\tilde{\mathbf{F}}_{t} - \hat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \hat{\mathbf{J}}_{2}) \mathbf{c}_{j} \cdot \mathbf{F}_{j} \right| \\
+ \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \mathbf{F}_{t}' (\hat{\mathbf{R}}^{(0)} - \mathbf{R}\hat{\mathbf{J}}_{1})' \mathbf{R} (\hat{\mathbf{C}}_{j} - \mathbf{c}_{j}' \cdot \hat{\mathbf{J}}_{2}) \mathbf{c}_{j} \cdot \mathbf{F}_{j} \right| \\
+ \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \mathbf{F}_{t}' (\hat{\mathbf{R}}^{(0)} - \mathbf{R}\hat{\mathbf{J}}_{1})' \mathbf{R} (\hat{\mathbf{F}}_{t} - \hat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \hat{\mathbf{J}}_{2}) \mathbf{c}_{j} \cdot \mathbf{F}_{j} \right| \\
+ \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot \mathbf{F}_{t}' (\hat{\mathbf{R}}^{(0)} - \mathbf{R}\hat{\mathbf{J}}_{1})' \mathbf{R} (\hat{\mathbf{F}}_{t} - \hat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \hat{\mathbf{J}}_{2}) \mathbf{c}_{j} \cdot \mathbf{F}_{j} \right| \\
+ \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j}' \cdot (\tilde{\mathbf{F}}_{t} - \hat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \hat{\mathbf{J}}_{2})' \mathbf{R}' \mathbf{R} \mathbf{F}_{t} (\hat{\mathbf{c}}_{j} - \mathbf{c}'_{j} \cdot \hat{\mathbf{J}}_{2}) \mathbf{c}_{j} \cdot \mathbf{F}_{j} \right| \\
+ \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} \mathbf{c}_{j} \cdot (\tilde{\mathbf{F}}_{t} - \hat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \hat{\mathbf{J}}_{2})' \mathbf{R}' \mathbf{R} \mathbf{F}_{t} (\hat{\mathbf{c}}_{j} - \mathbf{c}'_{j} \cdot \hat{\mathbf{J}}_{2}) \right\} \right| \\
+ \frac{1}{p_{2}} \left| \sum_{j=1}^{p_{2}} \left\{ \frac{1}{Tp_{1}} \sum_{t=1}^{T} (\hat{\mathbf{c}}_{j} - \mathbf{c}'_{j} \cdot \hat{\mathbf{J}}_{2}) \mathbf{F}_{t}' \mathbf{R}' \mathbf{R} \mathbf{F}_{t} (\hat{\mathbf{c}}_{j} - \mathbf{c}'_{j} \cdot \hat{\mathbf{J}}_{2}) \right\} \right| \\
= O_{p} \left(\max\{ \frac{1}{Tp_{1}} \sum_{j=1}^{T} \frac{1}{Tp_{1}} \sum_{j=1}^{T} \frac{1}{Tp_{1}} \sum_{j=1}^{T} \frac{1}{Tp_{1}} \sum_{j=1}^{T} \mathbf{R}_{j} \cdot \mathbf{R}_{j} \cdot \mathbf{R}_{j} \cdot \mathbf{R}_{j} \cdot \mathbf{R}_{j} \cdot \mathbf{R}_{j}$$

since

$$\begin{split} \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_j \cdot \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right) \mathbf{F}_t \mathbf{c}_j \cdot \right\} \right| \\ &= \frac{1}{Tp_1 p_2} \left| \operatorname{tr} \left(\mathbf{C} \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right) \mathbf{F}_t \mathbf{C}' \right) \right| \\ &\approx \frac{1}{Tp_1} \left| \operatorname{tr} \left(\mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right) \mathbf{F}_t \right) \right| \\ &\lesssim \frac{\left\| \widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right\|^2}{p_1} \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{F}_t' \mathbf{F}_t \right\| \\ &= O_p \left(\max \left\{ \frac{1}{Tp_2}, \frac{1}{p_1^2 p_2^2}, \frac{1}{T^2 p_1^2} \right\} \right) \end{split}$$

by Lemma 4(i) and Assumption 1(ii),

$$\begin{aligned} \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j}' \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{R} \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \mathbf{c}_{j} \right\} \right| &\lesssim \frac{\|\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1\|}{\sqrt{p_1}} \frac{\|\mathbf{R}\|}{\sqrt{p_1}} \\ &\times \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{F}_t' \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \right\| \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2}, \frac{1}{Tp_1} \right\} \right) O_p \left(\frac{1}{\sqrt{Tp_1p_2}} \right) \end{aligned}$$

by Lemmas 3(iii), 4(i) and (39),

$$\begin{aligned} \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j\cdot}' \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{R} \mathbf{F}_t \left(\widehat{\mathbf{c}}_{j\cdot}^{(0)} - \mathbf{c}_{j\cdot}' \widehat{\mathbf{J}}_2 \right) \right\} \right| &\lesssim \frac{\|\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1\|}{\sqrt{p_1}} \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{F}_t' \mathbf{F}_t \right\| \frac{\|(\widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2)' \mathbf{C}\|}{p_2} \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2}, \frac{1}{Tp_1} \right\} \right) \\ &\times O_p \left(\min\left\{ \frac{1}{Tp_1}, \frac{1}{Tp_2}, \frac{1}{p_1p_2}, \frac{1}{\sqrt{Tp_1p_2}} \right\} \right) \end{aligned}$$

because of Assumption 1(i) and Lemmas 3(iii), 4(i) and E.1 in Yu et al. (2022),

$$\frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j \cdot} \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \mathbf{R}' \mathbf{R} \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \mathbf{c}_{j \cdot} \right\} \right| \lesssim \left\| \frac{1}{T} \sum_{t=1}^T \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \right\| \\ \lesssim \frac{1}{Tp_1^2 p_2^2} \left\| \mathbf{C} \right\| \left\| \mathbf{R} \right\| \sum_{i=1}^{p_2} \sum_{j=1}^{p_2} \left\| \sum_{t=1}^T \mathbf{C}' \mathbf{E}_{ti \cdot} \mathbf{E}'_{t \cdot j} \mathbf{R} \right\| \\ = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1 p_2}}, \frac{1}{p_1 p_2} \right\} \right) \right)$$

because of (A.3) in Yu et al. (2022) and Lemma 3(iii),

$$\frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j \cdot} \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \mathbf{R}' \mathbf{R} \mathbf{F}_t \left(\widehat{\mathbf{c}}_{j \cdot} - \mathbf{c}'_{j \cdot} \widehat{\mathbf{J}}_2 \right) \right\} \right| \lesssim \frac{\|\mathbf{C}\|_F}{\sqrt{p_2}} \frac{\|\widehat{\mathbf{C}}^{(0)} - \mathbf{C}\widehat{\mathbf{J}}_2\|_F}{\sqrt{p_2}} \\ \times \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{F}'_t \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right) \right\|_F \\ = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right) O_p \left(\frac{1}{\sqrt{Tp_1p_2}} \right)$$

by Lemmas 3(iii), 4(ii) and (39),

$$\frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \left(\widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}_{j\cdot}' \widehat{\mathbf{J}}_2 \right) \mathbf{F}_t' \mathbf{R}' \mathbf{R} \mathbf{F}_t \left(\widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}_{j\cdot}' \widehat{\mathbf{J}}_2 \right) \right\} \right| \lesssim \frac{\|\widehat{\mathbf{C}}^{(0)} - \mathbf{C}\widehat{\mathbf{J}}_2\|_F^2}{p_2} \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{F}_t' \mathbf{F}_t \right\|_F$$
$$= O_p \left(\max\left\{ \frac{1}{Tp_1}, \frac{1}{p_1^2 p_2^2}, \frac{1}{T^2 p_2^2} \right\} \right)$$

because of Theorem 3.1 in Yu et al. (2022) and Assumption 1(ii). Consider the second addendum, we have that

$$\frac{1}{p_{2}}\sum_{j=1}^{p_{2}}\left|\left\{\frac{1}{Tp_{1}}\sum_{t=1}^{T}\sum_{i=1}^{p_{1}}(s_{tij}-\widehat{s}_{tij}^{(0)})e_{tij}\right\}\right| = \frac{1}{p_{2}}\sum_{j=1}^{p_{2}}\left|\left\{\frac{1}{Tp_{1}}\sum_{t=1}^{T}(\mathbf{s}_{t\cdot j}-\widehat{\mathbf{s}}_{t\cdot j}^{(0)})'\mathbf{e}_{t\cdot j}\right\}\right| \\
\lesssim \frac{1}{p_{2}}\left|\sum_{j=1}^{p_{2}}\left\{\frac{1}{Tp_{1}}\sum_{t=1}^{T}\mathbf{c}_{j}'.\mathbf{F}_{t}'\left(\widehat{\mathbf{R}}^{(0)}-\mathbf{R}\widehat{\mathbf{J}}_{1}\right)'\mathbf{e}_{t\cdot j}\right\}\right| \\
+ \frac{1}{p_{2}}\left|\sum_{j=1}^{p_{2}}\left\{\frac{1}{Tp_{1}}\sum_{t=1}^{T}\mathbf{c}_{j}'.\left(\widetilde{\mathbf{F}}_{t}-\widehat{\mathbf{J}}_{1}^{-1}\mathbf{F}_{t}\widehat{\mathbf{J}}_{2}\right)'\mathbf{R}'\mathbf{e}_{t\cdot j}\right\}\right| \\
+ \frac{1}{p_{2}}\left|\sum_{j=1}^{p_{2}}\left\{\frac{1}{Tp_{1}}\sum_{t=1}^{T}\mathbf{c}_{j}'.\left(\widetilde{\mathbf{F}}_{t}-\widehat{\mathbf{J}}_{1}^{-1}\mathbf{F}_{t}\widehat{\mathbf{J}}_{2}\right)'\mathbf{R}'\mathbf{e}_{t\cdot j}\right\}\right| \\
= O_{p}\left(\max\left\{\frac{1}{\sqrt{Tp_{1}p_{2}}},\frac{1}{p_{1}p_{2}},\frac{1}{\sqrt{Tp_{1}}},\frac{1}{Tp_{2}}\right\}\right)$$
(44)

since

$$\begin{aligned} \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j\cdot}' \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{e}_{t\cdot j} \right\} \right| &= \operatorname{tr} \left(\frac{1}{Tp_1p_2} \sum_{t=1}^T \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{E}_t \mathbf{C} \right) \\ &\lesssim \frac{1}{\sqrt{p_2}} \left\| \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{F}_t' \left(\widehat{\mathbf{R}}^{(0)} - \mathbf{R} \widehat{\mathbf{J}}_1 \right)' \mathbf{E}_t \right\| \frac{\|\mathbf{C}\|}{\sqrt{p_2}} \\ &= O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{Tp_2} \right\} \right) \end{aligned}$$

by Lemma 3(iii) and (B.3) in Yu et al. (2022),

$$\frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \mathbf{c}_{j\cdot}' \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2 \right)' \mathbf{R}' \mathbf{e}_{t\cdot j} \right\} \right| = \operatorname{tr} \left(\frac{1}{Tp_1 p_2} \sum_{t=1}^T \left(\widetilde{\mathbf{F}}_t - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-1} \right)' \mathbf{R}' \mathbf{E}_t \mathbf{C} \right) \\ \lesssim \left\| \frac{1}{Tp_1^2 p_2^2} \sum_{t=1}^T \mathbf{C}' \mathbf{E}_t' \mathbf{R} \mathbf{R}' \mathbf{E}_t \mathbf{C} \right\| \\ = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1 p_2}}, \frac{1}{p_1 p_2} \right\} \right)$$

by (38), Lemma 3(iii) and (A.3) in Yu et al. (2022),

$$\frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \left(\widehat{\mathbf{c}}_{j\cdot} - \mathbf{c}_{j\cdot} \widehat{\mathbf{J}}_2 \right)' \mathbf{F}'_t \mathbf{R}' \mathbf{e}_{t\cdot j} \right\} \right| = \operatorname{tr} \left(\frac{1}{Tp_1 p_2} \sum_{t=1}^T \mathbf{F}_t \mathbf{R}' \mathbf{E}_t \left(\widehat{\mathbf{C}} - \mathbf{C} \widehat{\mathbf{J}}_2 \right) \right) \\ \lesssim \frac{\|\widehat{\mathbf{C}} - \mathbf{C} \widehat{\mathbf{J}}_2\|}{\sqrt{p_2}} \left\| \frac{1}{Tp_1 \sqrt{p_2}} \sum_{t=1}^T \mathbf{F}_t \mathbf{R}' \mathbf{E}_t \right\| \\ = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1 p_2} \frac{1}{Tp_2} \right\} \right) O_p \left(\frac{1}{\sqrt{Tp_1}} \right)$$

by Lemma 4(ii) and A.1 in Yu et al. (2022). Consider the third addendum, since the result in (41) does

not depend on j, we have that

$$\frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \sum_{i=1}^{p_1} e_{tij}^2 - \frac{1}{p_1} \sum_{i=1}^{p_1} \mathbb{E}\left[e_{tij}^2\right] \right\} \right| = \max_j \left| \left\{ \frac{1}{Tp_1} \sum_{t=1}^T \sum_{i=1}^{p_1} e_{tij}^2 - \frac{1}{p_1} \sum_{i=1}^{p_1} \mathbb{E}\left[e_{tij}^2\right] \right\} \right| = O_p \left(\frac{1}{\sqrt{Tp_1}} \right).$$

$$(45)$$

Combining (43), (44) and (45) yields the desired result. The proof of (iv) follows similar steps. \Box Lemma 8. Under Assumptions 1-2, we have that as min{ T, p_1, p_2 } $\rightarrow \infty$

$$\begin{aligned} (i) & \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| = O_{p}(1) \\ (ii) & \left\| \widehat{\mathbf{H}}^{(0)-1} \right\| = O_{p}(1) \\ (iii) & \frac{1}{\sqrt{p_{2}}} \left\| \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{p_{1p_{2}}}, \frac{1}{Tp_{2}} \right\} \right) \\ (iv) & \frac{1}{\sqrt{p_{1}}} \left\| \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} - \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1p_{2}}}, \frac{1}{Tp_{1}} \right\} \right) \\ (v) & p_{2} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \right\| = O_{p}(1) \\ (vi) & p_{1} \left\| \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} - \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{p_{1p_{2}}}, \frac{1}{Tp_{2}} \right\} \right) \\ (viii) & p_{2} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} - \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1p_{2}}}, \frac{1}{Tp_{2}} \right\} \right) \\ (viii) & p_{1} \left\| \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{R}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1p_{2}}}, \frac{1}{Tp_{2}} \right\} \right) \\ (ix) & \sqrt{p_{2}} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} - \left(\mathbf{C}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1p_{2}}}, \frac{1}{Tp_{2}} \right\} \right) \\ (x) & \sqrt{p_{1}} \left\| \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} - \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{Tp_{2}}, \frac{1}{Tp_{2}} \right\} \right) \\ (x) & \sqrt{p_{1}} \left\| \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} - \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{Tp_{2}} \right\} \right) \\ (x) & \sqrt{p_{1}} \left\| \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{R}}^{(0)-1} - \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \right\| = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{Tp_{2}} \right\} \right) \\ (x) & \sqrt{p_{1}} \left\| \left(\widehat{\mathbf{$$

Proof. Consider (i) and note that

$$\begin{aligned} \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| &= \left\{ \nu^{(p_2)} \left(\widehat{\mathbf{K}}^{(0)} \right) \right\}^{-1} \\ &= \left\{ \min_{j=1,\dots,p_2} k_{jj} + \widehat{k}_{jj}^{(0)} - k_{jj} \right\}^{-1} \\ &= \left\{ \min_{j=1,\dots,p_2} k_{jj} - \min_{j=1,\dots,p_2} \left| \widehat{k}_{jj}^{(0)} - k_{jj} \right| \right\}^{-1} \\ &= \left\{ C_K^{-1} - \left| \widehat{k}_{jj}^{(0)} - k_{jj} \right| \right\}^{-1} = C_K + O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right) \end{aligned}$$

by Assumption 2(ii) and Lemma 7(i). The proof for (ii) follows the same steps. Consider (iii), we have

that

$$\begin{split} & p_{2}^{-1} \left\| \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\|^{2} = p_{2}^{-1} \left\| \sum_{i=j}^{p_{2}} \widehat{\mathbf{c}}_{j}^{(0)} \widehat{k}_{jj}^{(0)-1} - \sum_{j=1}^{p_{2}} \mathbf{c}_{j} . k_{jj}^{-1} \right\|^{2} \\ & = p_{2}^{-1} \left\| \sum_{j=1}^{p_{2}} \widehat{\mathbf{c}}_{j}^{(0)} \left(\widehat{k}_{jj}^{(0)} - k_{jj} + k_{jj} \right)^{-1} - \sum_{j=1}^{p_{2}} \mathbf{c}_{j} . k_{jj}^{-1} \right\|^{2} \\ & = p_{2}^{-1} \left\| \sum_{j=1}^{p_{2}} \widehat{\mathbf{c}}_{j}^{(0)} k_{jj}^{-1} \left(1 + (\widehat{k}_{jj}^{(0)} - k_{jj}) / k_{jj} \right)^{-1} - \sum_{j=1}^{p_{2}} \mathbf{c}_{j} . k_{jj}^{-1} \right\|^{2} \\ & \leq p_{2}^{-1} \left\| \left(1 + \min_{j=1,\dots,p_{2}} (\widehat{k}_{jj}^{(0)} - k_{jj}) / k_{jj} \right)^{-1} \sum_{j=1}^{p_{2}} \widehat{\mathbf{c}}_{j}^{(0)} k_{jj}^{-1} - \sum_{j=1}^{p_{2}} \mathbf{c}_{j} . k_{jj}^{-1} \right\|^{2} \\ & \leq p_{2}^{-1} \left\| \left(1 - \min_{j=1,\dots,p_{2}} (\widehat{k}_{jj}^{(0)} - k_{jj}) / k_{jj} + o \left(\min_{j=1,\dots,p_{2}} (\widehat{k}_{jj}^{(0)} - k_{jj}) / k_{jj} \right) \right) \sum_{j=1}^{p_{2}} \widehat{\mathbf{c}}_{j}^{(0)} k_{jj}^{-1} - \sum_{j=1}^{p_{2}} \mathbf{c}_{j} . k_{jj}^{-1} \right\|^{2} \\ & \leq p_{2}^{-1} \left\| \widehat{\mathbf{C}}^{(0)'} \mathrm{dg}(\mathbf{K})^{-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\|^{2} + p_{2}^{-1} \left(\min_{j=1,\dots,p_{2}} \left| \widehat{k}_{jj}^{(0)} - k_{jj} \right| / k_{jj} \right| \right)^{2} \left\| \widehat{\mathbf{C}}^{(0)'} \mathrm{dg}(\mathbf{K})^{-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\|^{2} + p_{2}^{-1} C_{K}^{2} \left(\min_{j=1,\dots,p_{2}} \left| \widehat{k}_{jj}^{(0)} - k_{jj} \right| \right)^{2} \left\| \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|^{2} \right\| \mathrm{dg}(\mathbf{K})^{-1} \right\|^{2} \\ & + p_{2}^{-1} C_{K}^{2} \left(\min_{j=1,\dots,p_{2}} \left| \widehat{k}_{jj}^{(0)} - k_{jj} \right| \right)^{2} \left\| \widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right\|^{2} \left\| \mathrm{dg}(\mathbf{K})^{-1} \right\|^{2} \\ & = O_{p} \left(\max\left\{ \frac{1}{T_{p_{1}}}, \frac{1}{T_{p_{1}2}^{2}}, \frac{1}{T^{2} p_{2}^{2}} \right\} \right)$$

by Lemmas 3(iii), 4, 7(iii) and Assumption 2(ii). Part (iv) follows analogously. Consider (v) and note that

$$\det\left(\widehat{\mathbf{C}}^{(0)'}\widehat{\mathbf{K}}^{(0)-1}\widehat{\mathbf{C}}^{(0)}\right)^{-1} = \prod_{j=1}^{k_2} \nu^{(j)} \left(\widehat{\mathbf{C}}^{(0)'}\widehat{\mathbf{K}}^{(0)-1}\widehat{\mathbf{C}}^{(0)}\right)$$
$$\geq \left(\nu^{(k_2)} \left(\widehat{\mathbf{C}}^{(0)'}\widehat{\mathbf{K}}^{(0)-1}\widehat{\mathbf{C}}^{(0)}\right)\right)^{k_2}$$
$$\geq \left(\nu^{(k_2)} \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}\mathbf{C}\right) - \left|\nu^{(k_2)} \left(\widehat{\mathbf{C}}^{(0)'}\widehat{\mathbf{K}}^{(0)-1}\widehat{\mathbf{C}}^{(0)}\right) - \nu^{(k_2)} \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}\mathbf{C}\right)\right|\right)^{k_2}$$

From Lemma 3(iv), we have that $\lim_{p_2 \to \infty} p_2^{-1} \nu^{(k_2)} \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right) > 0$. Moreover,

$$\begin{aligned} \frac{1}{p_2} \left| \nu^{(k_2)} \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right) - \nu^{(k_2)} \left(\mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \right) \right| &\leq \frac{1}{p_2} \left\| \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathbf{C}' \mathrm{dg}(\mathbf{K}) \mathbf{C} \right\| \\ &\lesssim \frac{1}{p_2} \left\| \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K}) \right\| \| \mathbf{C} \| \\ &\quad + \frac{1}{p_2} \left\| \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\| \left\| \widehat{\mathbf{C}}^{(0)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right\| \\ &= O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right) \end{aligned}$$

by Lemmas 3(iii), 3(v), Proposition 1 and term (iii), implying that det $\left(p_2^{-1}\widehat{\mathbf{C}}^{(0)'}\widehat{\mathbf{K}}^{(0)-1}\widehat{\mathbf{C}}^{(0)}\right)^{-1} > 0$ with probability tending to one as min{ T, p_1, p_2 } goes to infinity, i.e. $p_2 \left\| \left(\widehat{\mathbf{C}}^{(0)'}\widehat{\mathbf{K}}^{(0)-1}\widehat{\mathbf{C}}^{(0)}\right)^{-1} \right\| = O_p(1)$. The proof for (vi) follows the same steps. Consider (vii), we have that

$$p_{2} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} - \left(\mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \right)^{-1} \right\| \leq p_{2} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \right\| p_{2} \left\| \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \right\| \\ \frac{1}{p_{2}} \left\| \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right\| \\ = O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{p_{1}p_{2}}, \frac{1}{Tp_{2}} \right\} \right)$$

because of Lemma 3(iv), term (v) and since $\frac{1}{p_2} \left\| \widehat{\mathbf{C}}^{(0)\prime} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{p_1p_2}, \frac{1}{Tp_2} \right\} \right)$ by the same steps used in the proof of term (iii). Proof for (viii) follows analogously. The proofs for (ix) and (x) follow directly from (iii) and (vii), and from (iv) and (viii), respectively.

C.2.3 Results on EM estimators

Lemma 9. Under Assumption (1) through (3), for all $n \in \mathbb{N}_+$, as $\min\{p_1, p_2, T\} \rightarrow \infty$,

$$\begin{array}{l} (i) \ \left| \widehat{k}_{jj}^{(n)} - k_{jj} \right| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \text{ uniformly in } j \\ (ii) \ \left| \widehat{h}_{ii}^{(n)} - h_{ii} \right| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \text{ uniformly in } i \\ (iii) \ \frac{1}{p_2} \left| \sum_{j=1}^{p_2} \left(\widehat{k}_{jj}^{(n)} - k_{jj} \right) \right| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \\ (iv) \ \frac{1}{p_1} \left| \sum_{i=1}^{p_1} \left(\widehat{h}_{ii}^{(n)} - h_{ii} \right) \right| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \end{array}$$

Proof. Consider (ii) and recall that

$$\widehat{h}_{ii}^{(n)} = \frac{1}{Tp_2} \sum_{t=1}^{T} \left[\mathbf{Y}_t \widehat{\mathbf{K}}^{(n-1)-1} \mathbf{Y}'_t - \mathbf{Y}_t \widehat{\mathbf{K}}^{(n-1)-1} \widehat{\mathbf{C}}^{(n)} \mathbf{F}_{t|T}^{(n-1)\prime} \widehat{\mathbf{R}}^{(n)\prime} - \widehat{\mathbf{R}}^{(n)} \mathbf{F}_{t|T}^{(n-1)} \widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n-1)-1} \mathbf{Y}'_t \right. \\ \left. + \left(\widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n-1)-1} \widehat{\mathbf{C}}^{(n)} \right) \star \left(\left(\mathbb{I}_{k_2} \otimes \widehat{\mathbf{R}}^{(n)} \right) \left(\mathbf{f}_{t|T}^{(n-1)\prime} \mathbf{f}_{t|T}^{(n-1)\prime} + \mathbf{\Pi}_{t|T}^{(n-1)} \right) \left(\mathbb{I}_{k_2} \otimes \widehat{\mathbf{R}}^{(n)} \right)' \right) \right]_{ii},$$

and that $h_{ii} = \frac{1}{p_1} \mathbb{E} \left[\mathbf{e}'_{ti} \cdot \mathrm{dg}(\mathbf{K})^{-1} \mathbf{e}_{ti} \right]$. Let n = 1, we have

$$\begin{aligned} \left| \hat{h}_{ii}^{(1)} - h_{ii} \right| &= \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}_{i.}' \mathbf{F}_t \mathbf{C}' \hat{\mathbf{K}}^{(0)-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{r}_{i.} - \mathbf{r}_{i.}' \mathbf{F}_t \mathbf{C}' \hat{\mathbf{K}}^{(0)-1} \hat{\mathbf{C}}^{(1)} \mathbf{F}_{t|T}^{(0)} \hat{\mathbf{r}}_{i.}^{(1)} \right| \\ &+ \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}_{i.}' \mathbf{F}_t \mathbf{C}' \hat{\mathbf{K}}^{(0)-1} \mathbf{e}_{ti.} - \hat{\mathbf{r}}_{i.}^{(1)} \mathbf{F}_{t|T}^{(0)} \hat{\mathbf{C}}^{(1)'} \hat{\mathbf{K}}^{(0)-1} \mathbf{e}_{ti.} \right| \\ &+ \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{e}_{ti.}' \hat{\mathbf{K}}^{(0)-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{r}_{i.}' - \mathbf{e}_{ti.} \hat{\mathbf{K}}^{(0)-1} \hat{\mathbf{C}}^{(1)'} \mathbf{F}_{t|T}^{(0)'} \hat{\mathbf{r}}_{i.}^{(1)} \right| \\ &+ \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \left[\left(\hat{\mathbf{C}}^{(1)'} \hat{\mathbf{K}}^{(0)-1} \hat{\mathbf{C}}^{(1)} \right) \star \left(\left(\mathbb{I}_{k_2} \otimes \hat{\mathbf{R}}^{(1)} \right) \left(\mathbf{f}_{t|T}^{(n)} \mathbf{f}_{t|T}^{(n)'} + \mathbf{\Pi}_{t|T}^{(0)} \right) \left(\mathbb{I}_{k_2} \otimes \hat{\mathbf{R}}^{(1)} \right)' \right) \right]_{ii} \\ &- \hat{\mathbf{r}}_{i.}^{(1)} \mathbf{F}_{t|T}^{(0)} \hat{\mathbf{C}}^{(1)'} \hat{\mathbf{K}}^{(0)-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{r}_{i.} \right| \\ &+ \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{e}_{ti.}' \hat{\mathbf{K}}^{(0)-1} \mathbf{e}_{ti.} - \frac{1}{p_1} \mathbb{E} \left[\mathbf{e}_{ti.}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{e}_{ti.} \right] \right| \\ &= I + II + III + IV + V \end{aligned}$$

For term I we have that

$$\begin{split} \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{t} \cdot \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \left(\mathbf{C} \mathbf{F}'_t \mathbf{r}_{i} - \widehat{\mathbf{C}}^{(1)} \mathbf{F}^{(0)'}_{t|T} \widehat{\mathbf{r}}^{(0)}_{i} \right) \right| \\ & \leq \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{i} \cdot \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \left(\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right) \left(\mathbf{F}^{(0)}_{t|T} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \widehat{\mathbf{J}}'_1 \mathbf{r}_{i} \right| \\ & + \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{i} \cdot \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \left(\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right) \left(\widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \left(\widehat{\mathbf{r}}^{(1)}_{i} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i} \right) \right| \\ & + \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{i} \cdot \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_2 \left(\mathbf{F}^{(0)}_{t|T} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \left(\widehat{\mathbf{r}}^{(1)}_{i} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i} \right) \right| \\ & + \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{i} \cdot \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_2 \left(\mathbf{F}^{(0)}_{t|T} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \widehat{\mathbf{J}}_1' \mathbf{r}_{i} \right| \\ & + \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{i} \cdot \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_2 \left(\mathbf{F}^{(0)}_{t|T} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \widehat{\mathbf{J}}_1' \mathbf{r}_{i} \right| \\ & + \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{i} \cdot \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_2 \left(\mathbf{F}^{(0)}_{t|T} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \widehat{\mathbf{J}}_1' \mathbf{r}_{i} \right| \\ & + \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{i} \cdot \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_2 \left(\widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \left(\widehat{\mathbf{r}}^{(1)}_{i} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i} \right) \right| \\ & = O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}} , \frac{1}{\sqrt{Tp_2}} , \frac{1}{p_{1p_2}}} \right\} \right) \end{aligned}$$

since |

by Lemmas 1, 3(iii), 8(i), 11(i) and Proposition 1,

by Assumption 1(ii), Lemmas 1, 3(iii), 8(i) and Proposition 1,

by Lemmas 1, 3(iii), 8(i), 11(i) and Proposition 1,

$$\begin{aligned} \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{r}'_{t} \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \left(\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right) \left(\widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \widehat{\mathbf{J}}'_1 \mathbf{r}_{i.} \right| \\ \approx \left\| \left(\frac{\mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} (\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_2)}{p_2} \right) \star \left(\frac{1}{T} \sum_{t=1}^{T} \mathbf{f}_t \mathbf{f}'_t \right) \right\| \\ \lesssim \frac{\|\mathbf{C}\|}{\sqrt{p_2}} \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \frac{\|\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_2\|}{p_2} \\ = O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \end{aligned}$$

by Assumption 1(ii), Lemmas 1, 3(iii), 8(i) and Proposition 1,

by Lemmas 1, 3(iii), 8(i) and 11(i),

$$\begin{aligned} \left| \frac{1}{Tp_2} \sum_{t=1}^T \mathbf{r}'_{i\cdot} \mathbf{F}_t \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_2 \left(\widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \left(\widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right) \right| \\ \approx \left\| \left(\frac{\mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C}}{p_2} \right) \star \left(\frac{1}{T} \sum_{t=1}^T \mathbf{f}_t \mathbf{f}'_t \right) \right\| \left| \widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right| \\ \lesssim \frac{\|\mathbf{C}\|^2}{p_2} \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \left| \widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right| \\ = O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \end{aligned}$$

by Assumption 1(ii), Lemmas 1, 3(iii), 8(i) and Proposition 1. For terms II and III we have that

$$\begin{split} \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti.}^{\prime} \widehat{\mathbf{K}}^{(0)-1} \left(\mathbf{C} \mathbf{F}_{t}^{\prime} \mathbf{r}_{i.} - \widehat{\mathbf{C}}^{(1)} \mathbf{F}_{t|T}^{(0)} \widehat{\mathbf{r}}_{i.}^{(0)} \right) \right| \\ &\leq \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti.}^{\prime} \widehat{\mathbf{K}}^{(0)-1} \left(\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right) \left(\mathbf{F}_{t|T}^{(0)} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)^{\prime} \widehat{\mathbf{J}}_{1}^{\prime} \mathbf{r}_{i.} \right| \\ &+ \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti.}^{\prime} \widehat{\mathbf{K}}^{(0)-1} \left(\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right) \left(\widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)^{\prime} \left(\widehat{\mathbf{r}}_{i.}^{(1)} - \widehat{\mathbf{J}}_{1}^{\prime} \mathbf{r}_{i.} \right) \right| \\ &+ \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti.}^{\prime} \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \left(\mathbf{F}_{t|T}^{(0)} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)^{\prime} \left(\widehat{\mathbf{r}}_{i.}^{(1)} - \widehat{\mathbf{J}}_{1}^{\prime} \mathbf{r}_{i.} \right) \right| \\ &+ \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti.}^{\prime} \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \left(\mathbf{F}_{t|T}^{(0)} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)^{\prime} \widehat{\mathbf{J}}_{1}^{\prime} \mathbf{r}_{i.} \right| \\ &+ \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti.}^{\prime} \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \left(\mathbf{F}_{t|T}^{(0)} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)^{\prime} \widehat{\mathbf{J}}_{1}^{\prime} \mathbf{r}_{i.} \right| \\ &+ \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti.}^{\prime} \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \left(\mathbf{F}_{t|T}^{(0)} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)^{\prime} \widehat{\mathbf{J}}_{1}^{\prime} \mathbf{r}_{i.} \right| \\ &+ \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti.}^{\prime} \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \left(\widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)^{\prime} \left(\widehat{\mathbf{r}}_{i.}^{(1)} - \widehat{\mathbf{J}}_{1}^{\prime} \mathbf{r}_{i.} \right) \right| \\ &= O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right)$$

since

by Lemmas 8(i), 11(v) and Proposition 1,

$$\begin{aligned} \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{e}_{ti}' \cdot \widehat{\mathbf{K}}^{(0)-1} \left(\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right) \left(\widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \left(\widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right) \right| \\ \approx \left\| \left(\frac{\widehat{\mathbf{K}}^{(0)-1} (\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_2)}{\sqrt{p_2}} \right) \star \left(\frac{1}{T\sqrt{p_2}} \sum_{t=1}^{T} \mathbf{e}_{ti\cdot} \mathbf{f}_t' \right) \right\| \left| \widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right| \\ \lesssim \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \frac{\left\| \widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right\|}{\sqrt{p_2}} \left\| \frac{1}{T\sqrt{p_2}} \sum_{t=1}^{T} \mathbf{e}_{ti\cdot} \mathbf{f}_t' \right\| \left| \widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right| \\ = O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right) \end{aligned}$$

by Assumption 3(i), Lemma 8(i), and Proposition 1,

by Lemmas 3(iii), 8(i), 11(v) and Proposition 1,

by Assumption 1(ii), Lemmas 8(i) and Proposition 1,

$$\begin{aligned} \left| \frac{1}{Tp_2} \sum_{t=1}^T \mathbf{e}'_{ti} \cdot \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_2 \left(\mathbf{F}_{t|T}^{(0)} - \widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \widehat{\mathbf{J}}'_1 \mathbf{r}_{i} \cdot \right| \\ & \asymp \left\| \left(\frac{\widehat{\mathbf{K}}^{(0)-1} \mathbf{C}}{\sqrt{p_2}} \right) \star \left(\frac{1}{T\sqrt{p_2}} \sum_{t=1}^T \mathbf{e}_{ti} \cdot \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right)' \right) \right\| \\ & \lesssim \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \frac{\|\mathbf{C} \widehat{\mathbf{J}}_2\|}{\sqrt{p_2}} \left\| \frac{1}{T\sqrt{p_2}} \sum_{t=1}^T \mathbf{e}_{ti} \cdot \left(\mathbf{f}_{t|T}^{(0)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \right)' \right\| \\ &= O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \end{aligned}$$

by Lemma 3(iii), 8(i) and 11(v),

$$\begin{aligned} \left| \frac{1}{Tp_2} \sum_{t=1}^T \mathbf{e}'_{ti} \cdot \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_2 \left(\widehat{\mathbf{J}}_1^{-1} \mathbf{F}_t \widehat{\mathbf{J}}_2^{-\prime} \right)' \left(\widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right) \right| \\ \approx \left\| \left(\frac{\widehat{\mathbf{K}}^{(0)-1} \mathbf{C}}{\sqrt{p_2}} \right) \star \left(\frac{1}{T\sqrt{p_2}} \sum_{t=1}^T \mathbf{e}_{ti} \cdot \mathbf{f}_t' \right) \right\| \left| \widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right| \\ \lesssim \left\| \widehat{\mathbf{K}}^{(0)-1} \right\| \frac{\|\mathbf{C} \widehat{\mathbf{J}}_2\|}{\sqrt{p_2}} \left\| \left(\frac{1}{T\sqrt{p_2}} \sum_{t=1}^T \mathbf{e}_{ti} \cdot \mathbf{f}_t' \right) \right\| \left| \widehat{\mathbf{r}}_{i\cdot}^{(1)} - \widehat{\mathbf{J}}_1' \mathbf{r}_{i\cdot} \right| \end{aligned}$$

For term V we have

$$\begin{aligned} \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \mathbf{e}'_{ti} \cdot \widehat{\mathbf{K}}^{(0)-1} \mathbf{e}_{ti} - \frac{1}{p_1} \mathbb{E} \Big[\mathbf{e}'_{ti} \cdot \mathrm{dg}(\mathbf{K})^{-1}_{ti} \Big] \right| &\leq \max_j \left| \widehat{k}^{(0)-1}_{jj} - k^{-1}_{jj} \right| \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \sum_{j=1}^{p_2} e^2_{tij} \right| \\ &+ \left| \frac{1}{Tp_2} \sum_{t=1}^{T} \sum_{j=1}^{p_2} e_{tij} - \mathbb{E} \Big[e^2_{tij} \Big] \right| \left| k^{-1}_{jj} \Big| \\ &= O_p \left(\max \Big\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \Big\} \right) \end{aligned}$$

by Assumptions 2(ii), 2(iv) and Lemma 7(iii). The proof for (i) follows the same steps. Consider (iv), then

$$\begin{aligned} \frac{1}{p_{1}} \sum_{i=1}^{p_{1}} \left| \hat{h}_{ii}^{(1)} - h_{ii} \right| &= \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{r}_{i}'. \mathbf{F}_{t} \mathbf{C}' \hat{\mathbf{K}}^{(0)-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{r}_{i}. - \mathbf{r}_{i}'. \mathbf{F}_{t} \mathbf{C}' \hat{\mathbf{K}}^{(0)-1} \hat{\mathbf{C}}^{(0)} \mathbf{F}_{t|T}^{(0)} \hat{\mathbf{r}}_{i}^{(0)} \right| \\ &+ \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{r}_{i}'. \mathbf{F}_{t} \mathbf{C}' \hat{\mathbf{K}}^{(0)-1} \mathbf{e}_{ti} - \hat{\mathbf{r}}_{i}^{(0)} \mathbf{F}_{t|T}^{(0)} \hat{\mathbf{C}}^{(0)'} \hat{\mathbf{K}}^{(0)-1} \mathbf{e}_{ti} \right| \\ &+ \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{e}_{ti}'. \hat{\mathbf{K}}^{(0)-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{r}_{i}' - \mathbf{e}_{ti}. \hat{\mathbf{K}}^{(0)-1} \hat{\mathbf{C}}^{(0)'} \mathbf{F}_{t|T}^{(0)'} \hat{\mathbf{r}}_{i}^{(0)} \right| \\ &+ \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{e}_{ti}'. \hat{\mathbf{K}}^{(0)-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{r}_{i}' - \mathbf{e}_{ti}. \hat{\mathbf{K}}^{(0)-1} \hat{\mathbf{C}}^{(0)'} \mathbf{F}_{t|T}^{(0)'} \mathbf{f}_{i}^{(0)} \right| \\ &+ \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{e}_{ti}'. \hat{\mathbf{K}}^{(0)-1} \mathbf{C}^{(n+1)} \hat{\mathbf{K}}^{(0)-1} \mathbf{C}^{(n+1)} \right| \\ &- \hat{\mathbf{r}}_{i}^{(0)'} \mathbf{F}_{t|T}^{(0)} \hat{\mathbf{C}}^{(0)'} \hat{\mathbf{K}}^{(0)-1} \mathbf{C} \mathbf{F}_{t}' \mathbf{r}_{i} \right| \\ &+ \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{e}_{ti}'. \hat{\mathbf{K}}^{(0)-1} \mathbf{e}_{ti} - \frac{1}{p_{1}} \mathbb{E} \left[\mathbf{e}_{ti}'. \mathrm{dg}(\mathbf{K})^{-1} \mathbf{e}_{ti} \right] \right| \\ &= I + II + III + IIV + V \end{aligned}$$

For term I we have that

$$\begin{split} \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{r}_{i}' \mathbf{F}_{t} \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \left(\mathbf{C} \mathbf{F}_{t}' \mathbf{r}_{i} - \widehat{\mathbf{C}}^{(1)} \mathbf{F}_{t|T}^{(0)'} \widehat{\mathbf{r}}_{i}^{(0)} \right) \right| \\ \lesssim \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{r}_{i}' \mathbf{F}_{t} \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \left(\widehat{\mathbf{C}}^{(1)} - \mathbf{C} \widehat{\mathbf{J}}_{2} \right) \left(\widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)' \widehat{\mathbf{J}}_{1}' \mathbf{r}_{i} \right| \\ + \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{r}_{i}' \mathbf{F}_{t} \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \left(\mathbf{F}_{t|T}^{(0)} - \widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)' \widehat{\mathbf{J}}_{1}' \mathbf{r}_{i} \right| \\ + \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{r}_{i}' \mathbf{F}_{t} \mathbf{C}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C} \widehat{\mathbf{J}}_{2} \left(\widehat{\mathbf{J}}_{1}^{-1} \mathbf{F}_{t} \widehat{\mathbf{J}}_{2}^{-\prime} \right)' \left(\widehat{\mathbf{r}}_{i}^{(1)} - \widehat{\mathbf{J}}_{1}' \mathbf{r}_{i} \right) \right| \\ = O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right) \end{split}$$

since

by Assumption 1(ii), Lemmas 3(iii), 8(i), and Proposition 1,

by Lemmas 3(iii), 8(i), 11(i), and Proposition 1,

by Assumption 1(ii), Lemmas 3(iii), 8(i), and Proposition 1. For terms II and III we have that

$$\begin{split} \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{e}_{ti}' \widehat{\mathbf{K}}^{(0)-1} \left(\mathbf{C}\mathbf{F}_{t}'\mathbf{r}_{i} - \widehat{\mathbf{C}}^{(1)}\mathbf{F}_{t|T}^{(0)'} \widehat{\mathbf{r}}_{i}^{(0)} \right) \right| \\ \lesssim \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{e}_{ti}' \widehat{\mathbf{K}}^{(0)-1} \left(\widehat{\mathbf{C}}^{(1)} - \mathbf{C}\widehat{\mathbf{J}}_{2} \right) \left(\widehat{\mathbf{J}}_{1}^{-1}\mathbf{F}_{t}\widehat{\mathbf{J}}_{2}^{-\prime} \right)' \widehat{\mathbf{J}}_{1}' \mathbf{r}_{i} \right. \\ + \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{e}_{ti}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C}\widehat{\mathbf{J}}_{2} \left(\mathbf{F}_{t|T}^{(0)} - \widehat{\mathbf{J}}_{1}^{-1}\mathbf{F}_{t}\widehat{\mathbf{J}}_{2}^{-\prime} \right)' \widehat{\mathbf{J}}_{1}' \mathbf{r}_{i} \right. \right| \\ + \left| \frac{1}{Tp_{1}p_{2}} \sum_{t=1}^{T} \sum_{i=1}^{p_{1}} \mathbf{e}_{ti}' \widehat{\mathbf{K}}^{(0)-1} \mathbf{C}\widehat{\mathbf{J}}_{2} \left(\widehat{\mathbf{J}}_{1}^{-1}\mathbf{F}_{t}\widehat{\mathbf{J}}_{2}^{-\prime} \right)' \left(\widehat{\mathbf{r}}_{i}^{(1)} - \widehat{\mathbf{J}}_{1}' \mathbf{r}_{i} \right) \right| \\ = O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right) \end{split}$$

since

by Lemma (A.1) in Yu et al. (2022), Lemmas 8(i) and Proposition 1,

by Assumptions 1(i), 2(ii), Lemmas 3(iii), 7(iii), and 11(iv),

by Assumptions 1(i), 2(ii), 3(i), Lemma 7(iii), and Proposition 1. For term V we have

$$\begin{aligned} \frac{1}{p_{1}} \sum_{i=1}^{p_{1}} \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \mathbf{e}_{ti}' \cdot \widehat{\mathbf{K}}^{(0)-1} \mathbf{e}_{ti} - \frac{1}{p_{2}} \mathbb{E} \left[\mathbf{e}_{ti}' \cdot \mathrm{dg}(\mathbf{K})_{ti}^{-1} \mathbf{e}_{ti} \right] \right| &\leq \max_{j} \left| \widehat{k}_{jj}^{(0)-1} - k_{jj}^{-1} \right| \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \sum_{j=1}^{p_{2}} e_{tij}^{2} \right| \\ &+ \left| \frac{1}{Tp_{2}} \sum_{t=1}^{T} \sum_{j=1}^{p_{2}} e_{tij} - \mathbb{E} \left[e_{tij}^{2} \right] \right| \left| k_{jj}^{-1} \right| \\ &= O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right) \end{aligned}$$

by Assumptions 2(ii), 2(iv) and Lemma 7(i). Repeating the same steps for all $n \in \mathbb{N}_+$, replacing Lemmas 7 and 8 with Lemmas 9 and 10, respectively, completes the proof.

Lemma 10. Under Assumptions 1-2, for all $n \in \mathbb{N}_+$, we have that as $\min\{T, p_1, p_2\} \rightarrow \infty$

(i)
$$\left\|\widehat{\mathbf{K}}^{(n)-1}\right\| = O_p(1)$$

(ii) $\left\|\widehat{\mathbf{H}}^{(n)-1}\right\| = O_p(1)$
(iii) $\frac{1}{\sqrt{p_2}} \left\|\widehat{\mathbf{C}}^{(n)'}\widehat{\mathbf{K}}^{(n)-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}\right\| = O_p\left(\max\left\{\frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2}\right\}\right)$
(iv) $\frac{1}{\sqrt{p_1}} \left\|\widehat{\mathbf{R}}^{(n)'}\widehat{\mathbf{H}}^{(n)-1} - \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1}\right\| = O_p\left(\max\left\{\frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2}\right\}\right)$
(v) $p_2 \left\|\left(\widehat{\mathbf{C}}^{(n)'}\widehat{\mathbf{K}}^{(n)-1}\widehat{\mathbf{C}}^{(n)}\right)^{-1}\right\| = O_p(1)$

$$(vi) \ p_1 \left\| \left(\widehat{\mathbf{R}}^{(n)'} \widehat{\mathbf{H}}^{(n)-1} \widehat{\mathbf{R}}^{(n)} \right)^{-1} \right\| = O_p(1)$$

$$(vii) \ p_2 \left\| \left(\widehat{\mathbf{C}}^{(n)'} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right)^{-1} - \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right)$$

$$(viii) \ p_1 \left\| \left(\widehat{\mathbf{R}}^{(n)'} \widehat{\mathbf{H}}^{(n)-1} \widehat{\mathbf{R}}^{(n)} \right)^{-1} - \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right)$$

$$(ix) \ \sqrt{p_2} \left\| \left(\widehat{\mathbf{C}}^{(n)'} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right)^{-1} \widehat{\mathbf{C}}^{(n)'} \widehat{\mathbf{K}}^{(n)-1} - \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right)$$

$$(x) \ \sqrt{p_1} \left\| \left(\widehat{\mathbf{R}}^{(n)'} \widehat{\mathbf{H}}^{(n)-1} \widehat{\mathbf{R}}^{(n)} \right)^{-1} \widehat{\mathbf{R}}^{(n)'} \widehat{\mathbf{H}}^{(n)-1} - \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right)$$

Proof. Consider (i) and note that

$$\begin{aligned} \left\| \widehat{\mathbf{K}}^{(n)-1} \right\| &= \left\{ \nu^{(p_2)} \left(\widehat{\mathbf{K}}^{(n)} \right) \right\}^{-1} \\ &= \left\{ \min_{j=1,\dots,p_2} k_{jj} + \widehat{k}_{jj}^{(n)} - k_{jj} \right\}^{-1} \\ &= \left\{ \min_{j=1,\dots,p_2} k_{jj} - \min_{j=1,\dots,p_2} \left| \widehat{k}_{jj}^{(n)} - k_{jj} \right| \right\}^{-1} \\ &= \left\{ C_K^{-1} - \left| \widehat{k}_{jj}^{(n)} - k_{jj} \right| \right\}^{-1} = C_K + O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \end{aligned}$$

by Assumption 2(ii) and Lemma 9(i). The proof for (ii) follows the same steps. Consider (iii), we have that

$$\begin{aligned} \frac{1}{\sqrt{p_2}} \left\| \widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\| &\lesssim \frac{1}{\sqrt{p_2}} \left\| \widehat{\mathbf{C}}^{(n)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right\| \left\| \mathrm{dg}(\mathbf{K})^{-1} \right\| + \frac{1}{\sqrt{p_2}} \left\| \mathbf{C}' \left(\widehat{\mathbf{K}}^{(n)-1} - \mathrm{dg}(\mathbf{K})^{-1} \right) \right\| \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2} \right\} \right) \end{aligned}$$

because of Proposition 1, Assumption 2(ii), and since

$$\begin{split} \frac{1}{\sqrt{p_2}} \left\| \mathbf{C}' \left(\widehat{\mathbf{K}}^{(n)-1} - \mathrm{dg}(\mathbf{K})^{-1} \right) \right\| &\leq \frac{1}{\sqrt{p_2}} \left\{ \sum_{i=1}^{k_2} \sum_{j=1}^{p_2} \left| \left[\mathbf{C}' \left(\widehat{\mathbf{K}}^{(n)-1} - \mathrm{dg}(\mathbf{K})^{-1} \right) \right]_{ij} \right|^2 \right\}^{\frac{1}{2}} \\ &\leq \sqrt{k_2} \max_{ij} \left| \mathbf{c}'_{i} \left[\widehat{\mathbf{K}}^{(n)-1} - \mathrm{dg}(\mathbf{K})^{-1} \right]_{.j} \right| \\ &\leq \bar{c} \sqrt{k_2} \max_{j} \left| \left\{ \widehat{k}_{jj}^{(n)-1} k_{jj}^{-1} \left(\widehat{k}_{jj}^{(n)} - k_{jj} \right) \right\} \right| \\ &\leq \bar{c} \sqrt{k_2} \max_{j} \left| \left(\min_{j} \widehat{k}_{jj}^{(n)} \right)^{-1} \left(\min_{j} k_{jj} \right)^{-1} \sum_{j=1}^{p_2} \left(\widehat{k}_{jj}^{(n)} - k_{jj} \right) \right| \\ &\leq \bar{c} \sqrt{k_2} C_K^2 \left| \widehat{k}_{jj}^{(n)} - k_{jj} \right| \\ &= O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2}, \right\} \right) \end{split}$$

by Assumptions 1(i), 2(ii) and Lemma 9(i). The proof for (iv) follows the same steps. Consider (v)

and note that

$$\det\left(\widehat{\mathbf{C}}^{(n)'}\widehat{\mathbf{K}}^{(n)-1}\widehat{\mathbf{C}}^{(0)}\right)^{-1} = \prod_{j=1}^{k_2} \nu^{(j)} \left(\widehat{\mathbf{C}}^{(n)'}\widehat{\mathbf{K}}^{(n)-1}\widehat{\mathbf{C}}^{(n)}\right)$$

$$\geq \left(\nu^{(k_2)} \left(\widehat{\mathbf{C}}^{(n)'}\widehat{\mathbf{K}}^{(n)-1}\widehat{\mathbf{C}}^{(n)}\right)\right)^{k_2}$$

$$\geq \left(\nu^{(k_2)} \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}\mathbf{C}\right) - \left|\nu^{(k_2)} \left(\widehat{\mathbf{C}}^{(n)'}\widehat{\mathbf{K}}^{(n)-1}\widehat{\mathbf{C}}^{(n)}\right) - \nu^{(k_2)} \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}\mathbf{C}\right)\right|\right)^{k_2}$$

From Lemma 3(iv), we have that $\lim_{p_2 \to \infty} p_2^{-1} \nu^{(k_2)} \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right) > 0$. Moreover,

$$\begin{split} \frac{1}{p_2} \left| \nu^{(k_2)} \left(\widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right) - \nu^{(k_2)} \left(\mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \right) \right| &\leq \quad \frac{1}{p_2} \left\| \widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} - \mathbf{C}' \mathrm{dg}(\mathbf{K}) \mathbf{C} \right\| \\ &\lesssim \quad \frac{1}{p_2} \left\| \widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} - \mathbf{C}' \mathrm{dg}(\mathbf{K}) \right\| \| \mathbf{C} \| \\ &\quad + \frac{1}{p_2} \left\| \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\| \left\| \widehat{\mathbf{C}}^{(n)} - \mathbf{C} \widehat{\mathbf{J}}_2 \right\| \\ &= \quad O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right) \end{split}$$

by Lemmas 3(iii), 3(v), Proposition 1 and term (iii), implying that det $\left(p_2^{-1}\widehat{\mathbf{C}}^{(n)'}\widehat{\mathbf{K}}^{(n)-1}\widehat{\mathbf{C}}^{(n)}\right)^{-1} > 0$ with probability tending to one as min $\{T, p_1, p_2\}$ goes to infinity, i.e. $p_2 \left\| \left(\widehat{\mathbf{C}}^{(n)'}\widehat{\mathbf{K}}^{(n)-1}\widehat{\mathbf{C}}^{(n)}\right)^{-1} \right\| = O_p(1)$. The proof for (vi) follows the same steps. Consider (vii), we have that

$$p_{2} \left\| \left(\widehat{\mathbf{C}}^{(n)'} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right)^{-1} - \left(\mathbf{C}' \mathbf{K}^{-1} \mathbf{C} \right)^{-1} \right\| \leq p_{2} \left\| \left(\widehat{\mathbf{C}}^{(n)'} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right)^{-1} \right\| p_{2} \left\| \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \right\|$$
$$\frac{1}{p_{2}} \left\| \widehat{\mathbf{C}}^{(n)'} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} - \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right\|$$
$$= O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}} \frac{1}{p_{1}p_{2}} \right\} \right)$$

because of Lemma 3(iv), term (v) and since $\frac{1}{p_2} \left\| \widehat{\mathbf{C}}^{(n)'} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} - \mathbf{C'K^{-1}C} \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}} \frac{1}{p_1p_2} \right\} \right)$. Proof for (viii) follows analogously. The proofs for (ix) and (x) follow directly from (iii) and (vii), and from (iv) and (viii), respectively.

Lemma 11. Under Assumptions 1 through 3, there exist matrices $\hat{\mathbf{J}}_1$ and $\hat{\mathbf{J}}_2$ satisfying $\hat{\mathbf{J}}_1 \hat{\mathbf{J}}_1' \stackrel{p}{\to} \mathbb{I}_{k_1 k_1}$ and $\hat{\mathbf{J}}_2 \hat{\mathbf{J}}_2' \stackrel{p}{\to} \mathbb{I}_{k_2 k_2}$, such that, for all $n \in \mathbb{N}$, as $\min\{p_1, p_2, T\} \to \infty$, (i) $\left(\frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(n)} - \hat{\mathbf{J}}^{-1} \mathbf{f}_t\right) \mathbf{f}_t' \hat{\mathbf{J}}\right) = O_p \left(\max\left\{\frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2}\right\}\right)$ (ii) $\left\|\frac{1}{Tp_2} \sum_{j=1}^{p_2} \sum_{t=1}^{T} e_{tij} \left(\mathbf{f}_{t|T}^{(n)} - \hat{\mathbf{J}}^{-1} \mathbf{f}_t\right)\right\| = O_p \left(\max\left\{\frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2}\right\}\right)$ (iii) $\left\|\frac{1}{Tp_1} \sum_{i=1}^{p_1} \sum_{t=1}^{T} e_{tij} (\mathbf{f}_{t|T}^{(n)} - \hat{\mathbf{J}}^{-1} \mathbf{f}_t)\right\| = O_p \left(\max\left\{\frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2}\right\}\right)$ (iv) $\left\|\frac{1}{T\sqrt{p_1 p_2}} \sum_{i=1}^{p_2} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(n)} - \hat{\mathbf{J}}^{-1} \mathbf{f}_t\right) \mathbf{e}'_{t\cdot i}\right\| = O_p \left(\max\left\{\frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1 p_2}\right\}\right)$

(v)
$$\left\| \frac{1}{Tp_1\sqrt{p_2}} \sum_{i=1}^{p_1} \sum_{t=1}^T \mathbf{e}_{ti} \cdot (\mathbf{f}_{t|T}^{(n)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_t)' \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2} \right\} \right)$$

Proof. Let $\mathbf{x}_t = \left\{ \mathbf{f}_t \widehat{\mathbf{J}}, \frac{1}{p_2} \sum_{j=1}^{p_2} e_{tij}, \frac{1}{p_1} \sum_{i=1}^{p_1} e_{tij}, \frac{1}{\sqrt{p_1}p_2} \sum_{i=1}^{p_2} \mathbf{e}_{t\cdot i}, \frac{1}{p_1\sqrt{p_2}} \sum_{i=1}^{p_1} \mathbf{e}_{ti} \right\}$. From (B.4) in Barigozzi and Luciani (2024), we have that

$$\begin{aligned} \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t}^{(n)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right) \mathbf{x}_{t}' \right\| &\leq \\ \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|T}^{(n)} - \mathbf{f}_{t|t}^{(n)} \right) \mathbf{x}_{t}' \right\| \\ &+ \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t|t}^{(n)} - \mathbf{f}_{t}^{LS(n)} \right) \mathbf{x}_{t}' \right\| \\ &+ \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{f}_{t}^{LS(n)} - \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \right) \mathbf{x}_{t}' \right\| \\ &= I + II + III \end{aligned}$$

where

$$\mathbf{f}_{t}^{LS(n)} = \left(\left(\left(\widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \widehat{\mathbf{C}}^{(n)} \right)^{-1} \widehat{\mathbf{C}}^{(n)\prime} \widehat{\mathbf{K}}^{(n)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(n)\prime} \widehat{\mathbf{H}}^{(n)-1} \widehat{\mathbf{R}}^{(n)} \right)^{-1} \widehat{\mathbf{R}}^{(n)\prime} \widehat{\mathbf{H}}^{(n)-1} \right) \right) \mathbf{y}_{t}$$

Consider the case n=0. From Lemmas 4, 6, 8, (B.5) and (B.6) in Barigozzi and Luciani (2024), it follows that terms I and II are both $O_p\left(\frac{1}{p_1p_2}\right)$. Focusing on the third term, we have

$$\begin{split} III &\leq \left\| \frac{1}{T} \sum_{k=1}^{T} \left(\left(\left(\widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \right) \right) \\ &\times \left(\mathbf{C} \widehat{\mathbf{J}}_{2} \otimes \mathbf{R} \widehat{\mathbf{J}}_{1} - \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} \right) \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{x}_{t}^{t} \right\| \\ &+ \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\left(\left(\widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \right) \right) \right\| \\ &\leq \left\| \left(\left(\left(\mathbf{C}^{\prime} \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \mathbf{C}^{\prime} \mathbf{K}^{-1} \right) \otimes \left(\left(\mathbf{R}^{\prime} \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \mathbf{R}^{\prime} \mathrm{dg}(\mathbf{H})^{-1} \right) \right) \\ &\times \left(\mathbf{C} \widehat{\mathbf{J}}_{2} \otimes \mathbf{R} \widehat{\mathbf{J}}_{1} - \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} \right) \right\| \right\| \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{x}_{t}^{t} \right\| \\ &+ \left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \right) \right) \\ &- \left(\left(\mathbf{C}^{\prime} \mathbf{K}^{-1} \mathbf{C} \right)^{-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \right) \right) \right\| \\ &- \left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} - \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\mathbf{R}^{\prime} \mathbf{H}^{-1} \mathbf{R} \right)^{-1} \mathbf{R}^{\prime} \mathbf{H}^{-1} \right) \right\| \\ &- \left\| \left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \right)^{-1} \mathbf{R}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \right) \right) \right\| \\ &+ \left\| \left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)-1} \right) \right) \right\| \\ &+ \left\| \left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes \left(\left(\widehat{\mathbf{R}}^{(0)} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)-1} \right) \right) \right\| \\ &+ \left\| \left\| \left(\left(\left(\widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)} \widehat{\mathbf{K}}^{(0)-1} \right) \otimes$$

Since,

$$III_{a} \leq \left\| p_{2} \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \right\| \left\| \frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}}{\sqrt{p_{2}}} \right\| \left\| p_{1} \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \right\| \left\| \frac{\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1}}{\sqrt{p_{1}}} \right\|$$
$$\times \frac{1}{\sqrt{p_{1}p_{2}}} \left\| \mathbf{C} \widehat{\mathbf{J}}_{2} \otimes \mathbf{R} \widehat{\mathbf{J}}_{1} - \widehat{\mathbf{C}}^{(0)} \otimes \widehat{\mathbf{R}}^{(0)} \right\| \left\| \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{x}_{t}' \right\|$$
$$= O_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right) \left\| \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{x}_{t}' \right\|$$

by Lemmas 3(iv), 3(v), 4(iii),

$$\begin{split} III_{b} &\lesssim \sqrt{p_{2}} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\| \\ & \times \left\| \left(\frac{\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R}}{p_{1}} \right)^{-1} \right\| \frac{\|\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1}\|}{\sqrt{p_{1}}} \frac{\|\mathbf{C} \widehat{\mathbf{J}}_{2} \otimes \mathbf{R} \widehat{\mathbf{J}}_{1} - \widehat{\mathbf{C}} \otimes \widehat{\mathbf{R}} \|}{\sqrt{p_{1}p_{2}}} \right\| \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{x}_{t}' \right\| \\ & + \sqrt{p_{1}} \left\| \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} - \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \right\| \\ & \times \left\| \left(\frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_{2}} \right)^{-1} \right\| \frac{\|\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}\|}{\sqrt{p_{1}}} \left\| \frac{\mathbf{C} \widehat{\mathbf{J}}_{2} \otimes \mathbf{R} \widehat{\mathbf{J}}_{1} - \widehat{\mathbf{C}} \otimes \widehat{\mathbf{R}}}{\sqrt{p_{1}p_{2}}} \right\| \left\| \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{x}_{t}' \right\| \\ &= o_{p} \left(\max \left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right) \left\| \frac{1}{T} \sum_{t=1}^{T} \widehat{\mathbf{J}}^{-1} \mathbf{f}_{t} \mathbf{x}_{t}' \right\| \end{split}$$

by Lemmas 3(iv), 3(v), 4(iii), 8(ix) and 8(x),

$$III_{c} \leq \left\| \left(\frac{\mathbf{C}'\mathbf{K}^{-1}\mathbf{C}}{p_{2}} \right)^{-1} \right\| \left\| \left(\frac{\mathbf{R}'\mathbf{H}^{-1}\mathbf{R}}{p_{1}} \right)^{-1} \right\| \frac{1}{p_{1}p_{2}} \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{C}'\mathbf{K}^{-1} \right) \otimes \left(\mathbf{R}'\mathbf{H}^{-1} \right) \mathbf{e}_{t} \mathbf{x}_{t}' \right\|$$
$$\lesssim \frac{1}{p_{1}p_{2}} \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{C}'\mathbf{K}^{-1} \right) \otimes \left(\mathbf{R}'\mathbf{H}^{-1} \right) \mathbf{e}_{t} \mathbf{x}_{t}' \right\|$$

by Lemma 3(iv), and

$$III_{d} \lesssim \sqrt{p_{2}} \left\| \left(\widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} \widehat{\mathbf{C}}^{(0)} \right)^{-1} \widehat{\mathbf{C}}^{(0)'} \widehat{\mathbf{K}}^{(0)-1} - \left(\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C} \right)^{-1} \mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \right\| \\ \times \left\| \left(\frac{\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R}}{p_{1}} \right)^{-1} \right\| \frac{\|\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1}\|}{\sqrt{p_{1}}} \frac{1}{\sqrt{p_{1}p_{2}}} \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{e}_{t} \mathbf{x}_{t}' \right\| \\ + \sqrt{p_{1}} \left\| \left(\widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} \widehat{\mathbf{R}}^{(0)} \right)^{-1} \widehat{\mathbf{R}}^{(0)'} \widehat{\mathbf{H}}^{(0)-1} - \left(\mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \mathbf{R} \right)^{-1} \mathbf{R}' \mathrm{dg}(\mathbf{H})^{-1} \right\| \\ \times \left\| \left(\frac{\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1} \mathbf{C}}{p_{2}} \right)^{-1} \right\| \frac{\|\mathbf{C}' \mathrm{dg}(\mathbf{K})^{-1}\|}{\sqrt{p_{1}}} \left\| \frac{1}{\sqrt{p_{1}p_{2}}} \right\| \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{e}_{t} \mathbf{x}_{t}' \right\| \\ = O_{p} \left(\max\left\{ \frac{1}{\sqrt{Tp_{1}}}, \frac{1}{\sqrt{Tp_{2}}}, \frac{1}{p_{1}p_{2}} \right\} \right) \frac{1}{\sqrt{p_{1}p_{2}}} \left\| \frac{1}{T} \sum_{t=1}^{T} \mathbf{e}_{t} \mathbf{x}_{t}' \right\|$$

by Lemmas 3(iv)-(v), 4, 8(ix)-(x), we obtain

$$III = O_p \left(\max\left\{\frac{1}{\sqrt{Tp_1}}, \frac{1}{\sqrt{Tp_2}}, \frac{1}{p_1p_2}\right\} \right) \left\{ \left\| \frac{1}{T} \sum_{t=1}^T \widehat{\mathbf{J}}^{-1} \mathbf{f}_t \mathbf{x}_t' \right\| + \frac{1}{\sqrt{p_1p_2}} \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{e}_t \mathbf{x}_t' \right\| \right\} + \frac{1}{p_1p_2} \left\| \frac{1}{T} \sum_{t=1}^T \left(\mathbf{C}' \mathbf{K}^{-1} \right) \otimes \left(\mathbf{R}' \mathbf{H}^{-1} \right) \mathbf{e}_t \mathbf{x}_t' \right\|$$

From Lemma 2, we have that the stochastic behavior of $\hat{\mathbf{J}}^{-1}\mathbf{f}_t$ is equivalent to that of \mathbf{f}_t . Set $\mathbf{x}_t = \mathbf{f}_t$, we have that

$$\left\|\frac{1}{T}\sum_{t=1}^{T}\mathsf{f}_{t}\mathsf{f}_{t}'\right\| = O_{p}(1)$$

by Assumption 1(ii),

$$\frac{1}{\sqrt{Tp_1p_2}} \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^{T} \mathsf{e}_t \mathsf{f}'_t \right\| = O_p\left(\frac{1}{\sqrt{T}}\right)$$

by Assumption 3(i)

$$\frac{1}{p_1p_2} \left\| \frac{1}{T} \sum_{t=1}^{T} \left(\mathbf{C}' \mathbf{K}^{-1} \right) \otimes \left(\mathbf{R}' \mathbf{H}^{-1} \right) \mathbf{e}_t \mathbf{f}'_t \right\| = O_p \left(\max \left\{ \frac{1}{\sqrt{Tp_1p_2}} \right\} \right)$$

by Assumption 3(i) and Lemma 3(iii). This concludes the proof for (i). Let $x_t = \frac{1}{p_2} \sum_{j=1}^{p_2} e_{tij}$, we have that

$$\left\|\frac{1}{Tp_2}\sum_{j=1}^{p_2}\sum_{t=1}^{T}\mathsf{f}_t e_{tij}\right\| = O_p(1)$$

by Assumption 3(i),

$$\frac{1}{\sqrt{p_1 p_2}} \left\| \frac{1}{T p_2} \sum_{i=1}^{p_2} \sum_{t=1}^{T} \mathbf{e}_t e_{tij} \right\| = O_p\left(\frac{1}{\sqrt{T p_2}}\right)$$

since

$$\mathbb{E}\left[\frac{1}{p_{1}p_{2}}\left\|\frac{1}{Tp_{2}}\sum_{i=1}^{p_{2}}\sum_{t=1}^{T}\mathbf{e}_{t}e_{tij}\right\|_{F}^{2}\right] = \mathbb{E}\left[\frac{1}{p_{1}p_{2}}\sum_{l}\sum_{h}^{p_{1}}\sum_{h}^{p_{2}}\left|\frac{1}{Tp_{2}}\sum_{j=1}^{p_{2}}\sum_{t=1}^{T}e_{tlh}e_{tij}\right|^{2}\right]$$
$$= \max_{l}\max_{h}\mathbb{E}\left[\left|\frac{1}{Tp_{2}}\sum_{j=1}^{p_{2}}\sum_{t=1}^{T}e_{tlh}e_{tij}\right|^{2}\right]$$
$$= \max_{l}\max_{h}\frac{1}{T^{2}p_{2}^{2}}\sum_{j_{1},j_{2}=1}\sum_{s,t=1}^{T}\mathbb{E}[e_{tlh}e_{tij_{1}}e_{slh}e_{sij_{2}}]$$
$$= O_{p}\left(\frac{1}{Tp_{2}}\right)$$

by Assumption 2(iv), and

$$\frac{1}{p_1p_2} \left\| \frac{1}{Tp_2} \sum_{j=1}^{p_2} \sum_{t=1}^{T} \left(\mathbf{C}' \mathbf{K}^{-1} \right) \otimes \left(\mathbf{R}' \mathbf{H}^{-1} \right) \mathbf{e}_t e_{tij} \right\| = O_p \left(\max\left\{ \frac{1}{\sqrt{Tp_1p_2}} \right\} \right)$$

This concludes the proof for (ii). The results for (iii), (iv) and (v) can be established analogously. Repeating the same steps for all $n \in \mathbb{N}$ using Proposition 1, Proposition 1 (a.4)-(a.5) in Barigozzi and Luciani (2024), and Lemma 10 in place of Lemmas 4, 6, 8 completes the proof.