Uniqueness of generalized conformal restriction measures and Malliavin-Kontsevich-Suhov measures for $c \in (0, 1]$

Gefei Cai^{*} Yifan Gao[†]

Abstract

In this paper, we present a unified approach to establish the uniqueness of generalized conformal restriction measures with central charge $c \in (0, 1]$ in both chordal and radial cases, by relating these measures to the Brownian loop soup. Our method also applies to the uniqueness of the Malliavin-Kontsevich-Suhov loop measures for $c \in (0, 1]$, which was recently obtained in [Baverez-Jego, arXiv:2407.09080] for all $c \leq 1$ from a CFT framework of SLE loop measures. In contrast, though only valid for $c \in (0, 1]$, our approach provides additional probabilistic insights, as it directly links natural quantities of MKS measures to loop-soup observables.

Key words and phrases: conformal restriction, Malliavin-Kontsevich-Suhov measure, Brownian loop soup, Schramm-Loewner evolution.

1 Introduction

In the study of Brownian intersection exponents, Lawler and Werner [LW00] first realized that any conformally invariant process satisfying a certain restriction property would give rise to the same intersection exponents as the Brownian motion. This idea, combined with Schramm's discovery of SLE process [Sch00], finally led to the rigorous determination of the Brownian intersection exponents [LSW01a, LSW01b, LSW02b, LSW02a]. To further deepen this idea, Lawler, Schramm and Werner [LSW03] gave a complete understanding of the *conformal restriction measures* in the chordal case (which corresponds to Definition 1.1 with c = 0 below). In particular, they showed that the boundary of conformal restriction samples can be described by variants of SLE_{8/3}. Conformal restriction measures in other cases are also studied in [Wu15, Qia18].

For general SLE_{κ} with $\kappa \in (0, 4]$, there is a corresponding parameter $c \leq 1$ called *central charge* related to it via

$$c(\kappa) = 1 - 6\left(\frac{2}{\sqrt{\kappa}} - \frac{\sqrt{\kappa}}{2}\right)^2,\tag{1.1}$$

such that SLE_{κ} and its variants satisfy a generalized conformal restriction property that involves the Brownian loop mass with the corresponding c; see e.g. [LSW03, Dub05, Law09, WW13, Qia21]. For

^{*}caigefei1917@pku.edu.cn; Beijing International Center for Mathematical Research, Peking University

[†]yifangao@cityu.edu.hk; Department of Mathematics, City University of Hong Kong

 $c \in (0, 1]$, corresponding to $\kappa \in (\frac{8}{3}, 4]$, there is another simple way to obtain a general restriction sample by taking the filled union of a standard one with all loop-soup clusters it intersects from an independent Brownian loop soup of intensity $\frac{c}{2}$. However, it remains open whether these generalized restriction measures are unique.

There also exists a loop version of restriction measure, known as the Malliavin-Kontsevich-Suhov (MKS) measure [KS07]. It is an infinite measure defined on simple loops on \mathbb{C} with a parameter $c \leq 1$; see Definition 1.4 below. To construct an MKS measure, one uses the loop version of SLE_{κ} [KW16, Zha21] (see also [Wer08, BD16] for the special cases c = 0, -2). Recently, Baverez and Jego [BJ24] established the framework on the conformal field theory (CFT) of SLE, and as an application, they obtained the uniqueness of MKS measures for all $c \leq 1$. It would be interesting to explore whether their approach can be adapted to show the uniqueness of generalized restriction measures.

In this paper, we present a unified and direct approach to establish the uniqueness of all these restriction measures for $c \in (0, 1]$, based on a key observation that connects these measures to the *Brownian loop soup* or the SLE_{8/3} *loop soup*. In particular, we explicitly express various natural quantities associated with these measures in terms of loop-soup observables, which provides further probabilistic understanding of these measures.

In the following, we will state our results on generalized restriction measures and MKS measures in Sections 1.1 and 1.2, respectively, and then provide several applications in Section 1.3.

1.1 Generalized chordal restriction measures

For convenience, we focus on the unit disk $\mathbb{D} \subset \mathbb{C}$, with two marked boundary points -1 and 1. Let $\mathbb{S}^1 := \partial \mathbb{D}$. Let \mathcal{K} (resp. \mathcal{K}_-) be the collection of simply connected closed sets $K \subset \overline{\mathbb{D}}$ such that $K \cap \mathbb{S}^1 = \{-1, 1\}$ (resp. $K \cap \mathbb{S}^1 = \mathbb{S}^1 \cap \{z : \Im z \ge 0\}$). Let \mathcal{Q} be the collection of compact sets $A \subset \mathbb{D}$ such that $\mathbb{D} \setminus A$ is simply connected and $-1, 1 \notin A$. For $A \in \mathcal{Q}$, let $f_A : \mathbb{D} \setminus A \to \mathbb{D}$ be a conformal map that preserves -1, 1. We also denote the set of $A \in \mathcal{Q}$ with $A \cap \mathbb{S}^1 \subset \{z \in \mathbb{C} : \Im z < 0\}$ by \mathcal{Q}_- .

We equip \mathcal{K} (resp. \mathcal{K}_{-}) with the σ -algebra generated by the collection of events $\{K \in \mathcal{K} : K \cap A = \emptyset\}$ (resp. $\{K \in \mathcal{K}_{-} : K \cap A = \emptyset\}$) for all $A \in \mathcal{Q}$ (resp. $A \in \mathcal{Q}_{-}$).

Definition 1.1. Let $c, \alpha \in \mathbb{R}$. We say that a probability measure \mathbb{P} on \mathcal{K} (resp. \mathcal{K}_{-}) satisfies the two-sided (resp. one-sided) chordal c-restriction with exponent α , if for all $A \in \mathcal{Q}$ (resp. $A \in \mathcal{Q}_{-}$), we have

$$\frac{d\mathbb{P}(K)}{d\mathbb{P}_A(K)}\mathbf{1}_{K\cap A=\emptyset} = \mathbf{1}_{K\cap A=\emptyset}(f'_A(1)f'_A(-1))^{\alpha}\exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}}(K,A)\right),\tag{1.2}$$

where $\mathbb{P}_A = \mathbb{P} \circ f_A$, and $\Lambda_{\mathbb{D}}(K, A)$ is the total mass of loops on \mathbb{D} intersecting both K and A under the Brownian loop measure μ^{BL} , defined in [LW04]. Note that $f'_A(1)f'_A(-1)$ does not depend on the choice of conformal map f_A .

When c = 0, Definition 1.1 reduces to the *standard* chordal restriction measures in [LSW03]. The uniqueness of such measures is straightforward, as (1.2) now gives $\mathbb{P}[K \cap A = \emptyset] = (f'_A(1)f'_A(-1))^{\alpha}$.

Note that the two-sided case above has been studied in [Qia21]. In [Qia21, Proposition 6.2], the author constructed measures that satisfy the two-sided chordal *c*-restriction with exponent α in the range $c \leq 1$ (i.e., $\kappa \in (0, 4]$) and $\alpha \geq \frac{6-\kappa}{2\kappa}$, using variants of SLE_{κ}. However, it remains open whether these measures are unique or whether this range of α is maximal for which these measures exist.

The first main result of this paper is to provide positive answers to both questions for $c \in (0, 1]$. Similar result also holds for the one-sided chordal *c*-restriction measures.

Theorem 1.2. Let $c \in (0,1]$ and $\kappa \in (\frac{8}{3},4]$ be related via (1.1). Then the two-sided (resp. one-sided) chordal c-restriction with exponent α exists if and only if $\alpha \geq \frac{6-\kappa}{2\kappa}$ (resp. $\alpha > 0$). Furthermore, when $\alpha \geq \frac{6-\kappa}{2\kappa}$ (resp. $\alpha > 0$), there exists a unique probability measure on \mathcal{K} (resp. \mathcal{K}_{-}) satisfying the two-sided (resp. one-sided) chordal c-restriction with exponent α in Definition 1.1.

The main ingredient to show the uniqueness part of Theorem 1.2 is to express the non-intersection probability $\mathbb{P}[K \cap A = \emptyset]$ by some quantity only involving the Brownian loop soup. For this purpose, we let $\mathcal{L}_{\mathbb{D}}$ be the Brownian loop soup on \mathbb{D} of intensity $\frac{c}{2}$, that is, a Poisson point process with intensity measure $\frac{c}{2}\mu^{\mathrm{BL}}$ restricted to loops fully contained in \mathbb{D} . For any bounded set $B \subset \mathbb{C}$, the *filling* of B is defined as the complement of unbounded connected component of $\mathbb{C} \setminus B$. Let \widetilde{A} be the filling of the union of A and all the loop-soup clusters in $\mathcal{L}_{\mathbb{D}}$ that intersect A. We need to consider the following event

$$E := \{-1 \text{ and } 1 \text{ are in the same connected component of } \overline{\mathbb{D}} \setminus \widetilde{A} \}.$$
(1.3)

On the event E, we let $f_{\widetilde{A}}$ be the conformal map from the connected component of $\overline{\mathbb{D}} \setminus \widetilde{A}$ containing 1 to \mathbb{D} with -1 and 1 fixed. Note that E happens almost surely when $A \in \mathcal{Q}_-$. Then we have

Theorem 1.3. Let $c \in (0,1]$ and $\alpha \in \mathbb{R}$. Suppose \mathbb{P} satisfies the two-sided (resp. one-sided) chordal *c*-restriction with exponent α . Then for any $A \in \mathcal{Q}$ (resp. $A \in \mathcal{Q}_{-}$),

$$\mathbb{P}[K \cap A = \emptyset] = \mathbf{E}\left[\mathbf{1}_E\left(f'_{\widetilde{A}}(1)f_{\widetilde{A}}(-1)\right)^{\alpha}\right]$$
(1.4)

where **E** denotes the expectation with respect to the Brownian loop soup $\mathcal{L}_{\mathbb{D}}$.

The proof of (1.4) is based on a simple observation that $\exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}}(K,A)\right)$ in (1.2) is in fact the probability that there is no loop in $\mathcal{L}_{\mathbb{D}}$ that intersects both K and A given K. Hence, by interpolating extra randomness from the Brownian loop soup, we obtain the first equation that $\mathbb{P}[K \cap A = \emptyset] = \mathbf{E} \otimes \mathbb{E}_A \left[(f'_A(1)f'_A(-1))^{\alpha} \mathbf{1}_{K \cap A_1 = \emptyset} \right]$, where \mathbb{E}_A represents the expectation for a restriction sample in $\mathbb{D} \setminus A$, and A_1 is the filling of the union of A and all loops in $\mathcal{L}_{\mathbb{D}}$ that intersect A. Then, one can iterate the first equation until no loops can be added anymore, i.e. arriving at \widetilde{A} , and the limiting equation is just (1.4); see Section 2.1 for details. In fact, one can verify (1.4) straightforwardly if K is constructed by the union of a standard restriction sample with exponent α and all the clusters in $\mathcal{L}_{\mathbb{D}}$ that intersect it, which satisfies (1.2) directly.

Moreover, similar results also hold for the generalized radial and tri-chordal restriction measures; see Section 2.2 for detailed discussions. Next, we turn to the loop case.

1.2 The Malliavin-Kontsevich-Suhov measure

Let $\mu_{\mathbb{C}}$ be a Borel measure on the set of simple loops on \mathbb{C} . We say that $\mu_{\mathbb{C}}$ is *non-trivial* if for any bounded domain D and $\delta > 0$, the $\mu_{\mathbb{C}}$ -mass of simple loops ℓ with $\ell \subset D$ and diam $(\ell) > \delta$ is finite.

Definition 1.4 ([KS07]). A non-trivial measure $\mu_{\mathbb{C}}$ on the set of simple loops on \mathbb{C} is called a Malliavin-Kontsevich-Suhov measure if the following holds. For any simply connected domain $D \subset \mathbb{C}$, define μ_D by

$$\frac{d\mu_D(\eta)}{d\mu_{\mathbb{C}}(\eta)} \mathbf{1}_{\eta \subset D} = \mathbf{1}_{\eta \subset D} \exp\left(\frac{c}{2}\Lambda^*(\eta, \partial D)\right),\tag{1.5}$$

where $\Lambda^*(\eta, \partial D)$ is the total mass of loops on \mathbb{C} intersecting both η and ∂D under the normalized Brownian loop measure defined in [FL13]. Then for any two conformally equivalent domains D and D', the pushforward of μ_D under any conformal map from D to D' equals $\mu_{D'}$.

The study of loop versions of SLEs can be traced back to Werner [Wer03]. Following this, Kontsevich and Suhov wondered whether the restriction property of these loops could be alternatively expressed in the framework of CFT, which led to the formalism described in [KS07]. They also conjectured the existence and uniqueness of the measure in Definition 1.4 for all $c \leq 1$ [KS07, Conjecture 1], and related it to the unitarizing measure considered by Airault and Malliavin [AM01]. Now this measure is often known as the Malliavin-Kontsevich-Suhov measure.

Such an MKS measure was first considered by Werner [Wer08] for c = 0, where it is proved that the measure \mathcal{W} induced from μ^{BL} by taking outer boundaries of Brownian loops indeed characterizes the c = 0 MKS measure. We will call \mathcal{W} Werner's SLE_{8/3} loop measure on \mathbb{C} . A construction for the case c = -2 was given in [BD16]. Later, Kemppainen and Werner [KW16] constructed MKS measures for $c \in (0, 1]$ by taking the counting measure on loops in the full-plane CLE_{κ} configuration, denoted by SLE^{loop}_{κ} in the sequel. In particular, they studied the SLE_{8/3} loop soup on \mathbb{C} and proved that the outer and inner boundaries of SLE_{8/3} loop soup clusters are both equal to SLE^{loop}_{κ}. The construction for all $c \leq 1$ was finally completed by Zhan [Zha21]. Recently, Baverez and Jego [BJ24] rigorously derived the CFT of SLE loop measures, based on an in-depth study of its Virasoro algebra structures. As a consequence, they obtained the uniqueness of MKS measures for all $c \leq 1$. See also a contemporaneous work [GQW24] on the Virasoro representation of SLE loops.

The second main result of this paper is a direct and more "probabilistic" proof in the regime $c \in (0, 1]$, following a similar approach used in the generalized chordal restriction measures.

Theorem 1.5. Let $c \in (0,1]$. The MKS measure is unique up to a positive multiplicative constant.

Similar to Theorem 1.2, we will prove Theorem 1.5 by relating the MKS loop measure to loop-soup observables. However, since the Brownian loop soup on \mathbb{C} has only one cluster, one needs to consider the SLE_{8/3} loop soup instead, as indicated in [KW16]. This approach is feasible thanks to [CW24]; namely, we can study a related measure $\hat{\mu}_D$, which is defined in a similar fashion to (1.5) but with Λ^* replaced by Werner's SLE_{8/3} loop measure \mathcal{W} on \mathbb{C} , see (3.1). Let $\mathcal{L}_{\mathbb{C}}^{\mathcal{W}}$ be the whole-plane SLE_{8/3} loop soup of intensity $\frac{c}{2}$, that is, a Poisson point process with intensity measure $\frac{c}{2}\mathcal{W}$. Suppose D and U are simply connected domains in \mathbb{C} containing the origin, and $U \subset D$. Let $\mathcal{C}_{\partial D}$ be the union of ∂D and all clusters in $\mathcal{L}_{\mathbb{C}}^{\mathcal{W}}$ that intersect ∂D . Let D_{∞} be the connected components of $\mathbb{C} \setminus \mathcal{C}_{\partial D}$ that contains the origin. Furthermore, let $f_{D,\infty}$ be the conformal maps from $D_{\infty} \to \mathbb{D}$ such that $f_{D,\infty}(0) = 0$ and $f'_{D,\infty}(0) > 0$. Define $f_{U,\infty}$ in the same way. Then we have the following analog of Theorem 1.3.

Theorem 1.6. Let $c \in (0,1]$. Let $\mu_{\mathbb{C}}^0$ be the restriction of $\mu_{\mathbb{C}}$ to loops surrounding the origin. Then there is a constant $\lambda \in (0,\infty)$ such that for all $U \subset D$ as above,

$$\mu^0_{\mathbb{C}}[\ell \not\subset U, \ell \subset D] = \lambda \mathbf{E} \log f'_{U,\infty}(0) - \lambda \mathbf{E} \log f'_{D,\infty}(0),$$

where **E** denotes the expectation with respect to the $\text{SLE}_{8/3}$ loop soup $\mathcal{L}_{\mathbb{C}}^{\mathcal{W}}$.

Theorem 1.6 is new and it directly implies Theorem 1.5. Note that for c = 0 and $\mu_{\mathbb{C}} = \mathcal{W}$, the corresponding result of Theorem 1.6 has been provided in [Wer08, Proposition 3] with $\lambda = \frac{\pi}{5}$ [Wer08, Page 151]. Furthermore, if one views the sample of $\mu_{\mathbb{C}}$ as the outer boundary of the union of the loop from \mathcal{W} and the clusters in $\mathcal{L}_{\mathbb{C}}^{\mathcal{W}}$ intersecting it, then Theorem 1.6 is a direct consequence of its c = 0 counterpart [Wer08, Proposition 3]. Moreover, if $c \in (0, 1]$ and $\kappa = \kappa(c) \in (\frac{8}{3}, 4]$, with $\mu_{\mathbb{C}}$ chosen as $\mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ (i.e., the counting measure on loops of the full-plane CLE_{κ}), then the corresponding λ in Theorem 1.6 can be explicitly determined as $\lambda = \frac{1}{\pi}(\frac{\kappa}{4} - 1)\cot(\pi(1 - \frac{4}{\kappa}))$; see Corollary 3.4.

1.3 Applications

Here we provide several consequences of the results in Sections 1.1 and 1.2.

As mentioned to us by W. Werner, the *reversibility* and *duality* of SLE can be obtained from the uniqueness of these restriction measures, which was indeed considered (see e.g. [Wer05]) before the proofs [Zha08b] (see also [MS16b, LY21])¹ and [Zha08a, Dub09, MS16a]. This approach to the reversibility of simple SLE loop measures was completed in [BJ24, Section 7.1], as well as the $\kappa \leftrightarrow \frac{16}{\kappa}$ duality of SLE_{κ} loops [BJ24, Theorem 7.1] (this duality was also proved in a contemporaneous work [ACSW24b, Theorem 1.2]). Using similar ideas, our results on the uniqueness of general chordal restriction measures (Theorem 1.2) will provide alternate proofs in the regime $\kappa \in (\frac{8}{3}, 4]$ for the reversibility of chordal SLE_{κ} (and its variants), and the $\kappa \leftrightarrow \frac{16}{\kappa}$ duality in the chordal case.

The next application is about the SLE characterization of one-sided chordal restriction measures. Let γ be the lower boundary of an one-sided chordal 0-restriction sample with exponent $\alpha > 0$, which is a simple path from -1 to 1 on \mathbb{D} . Let Γ be an independent $\operatorname{CLE}_{\kappa}$ on \mathbb{D} with $\kappa \in (\frac{8}{3}, 4]$. Let η be the lower boundary of the union of γ and all loops of Γ that γ intersects. Then the law of η , which was characterized by $\operatorname{SLE}_{\kappa}(\rho)$ process in [WW13], now can be deduced from Theorem 1.2 directly.

Proposition 1.7 ([WW13, Theorem 1.1, Theorem 2.1]). Let $\kappa \in (\frac{8}{3}, 4]$ and $\alpha > 0$. The law of η is an SLE_{κ}(ρ) on \mathbb{D} from -1 to 1 with force point $e^{i\pi-}$, where $\rho \in (-2, \infty)$ satisfies $\alpha = \frac{(\rho+2)(\rho+6-\kappa)}{4\kappa}$.

¹We mention that the reversibility for non-simple SLE is established in [MS16c].

Proof. Since the CLE_{κ} on \mathbb{D} can be realized as the outer boundaries of outermost clusters in a Brownian loop soup on \mathbb{D} of intensity $\frac{c}{2}$ [SW12], it is straightforward to check that the connected component of $\mathbb{D} \setminus \eta$ above η satisfies the one-sided chordal *c*-restriction with exponent α as in Definition 1.1. On the other hand, as indicated in [Dub05], the upper part of $\text{SLE}_{\kappa}(\rho)$ on \mathbb{D} satisfies the one-sided chordal *c*-restriction (1.2) with exponent α as well (see [WW13, Page 10-11] for a detailed justification). Combined, the result follows from the uniqueness part of Theorem 1.2.

Analogously, the two-sided case of Theorem 1.2 immediately gives the following. A radial counterpart will be given later by Proposition 2.9.

Proposition 1.8. Let $\kappa \in (\frac{8}{3}, 4]$ and $\alpha \geq \frac{6-\kappa}{2\kappa}$. Let K be a two-sided chordal c-restriction with exponent α . Let Γ be an independent $\operatorname{CLE}_{\kappa}$ on \mathbb{D} . Let $\mathcal{C}(K)$ be the union of K and all loops of Γ that K intersects. Then the filling of $\mathcal{C}(K)$ has the same law as that constructed in [Qia21, Proposition 6.2] using variants of $\operatorname{SLE}_{\kappa}$.

Based on Theorem 1.6, for any connected compact set V separating 0 and ∞ , we can express the $\mu^0_{\mathbb{C}}$ -mass of loops intersecting V by some $\mathrm{SLE}_{8/3}$ loop-soup observables; see (1.6) below. As before, let \mathcal{C}_V be the union of V and all clusters in $\mathcal{L}^{\mathcal{W}}_{\mathbb{C}}$ that intersect V. Let $\Omega_{V,\infty}$ and $\Omega^*_{V,\infty}$ be the connected components of $\mathbb{C} \setminus \mathcal{C}_V$ containing 0 and ∞ , respectively. Let $V_{\infty} := \mathbb{C} \setminus (\Omega_{V,\infty} \cup \Omega^*_{V,\infty})$, $f_{V_{\infty}} : \Omega_{V,\infty} \to \mathbb{D}$ be conformal fixing 0, and $h_{V_{\infty}} : \Omega^*_{V,\infty} \to \mathbb{D}^* := \mathbb{C} \setminus \overline{\mathbb{D}}$ be conformal fixing ∞ .

Theorem 1.9. For $c \in (0,1]$, there is a constant $\zeta \in \mathbb{R}$ such that for any connected compact set V separating 0 and ∞ ,

$$\mu_{\mathbb{C}}^{0}[\ell \cap V \neq \emptyset] = \lambda \mathbf{E}\left[\theta(V_{\infty})\right] + \zeta, \tag{1.6}$$

where $\theta(V_{\infty}) := \log |f'_{V_{\infty}}(0)| - \log |h'_{V_{\infty}}(\infty)|$ is the so-called electrical thickness of V_{∞} [KW04], and **E** is the expectation for the SLE_{8/3} loop soup $\mathcal{L}^{\mathcal{W}}_{\mathbb{C}}$. In fact, ζ can be related to λ through (4.1).

The proof of Theorem 1.9 will be given in Section 4. Theorem 1.9 can also be straightforwardly extended to c = 0 and $\mu_{\mathbb{C}} = \mathcal{W}$, in which case $\zeta = \frac{\sqrt{6}}{15}$; see Remark 4.4. However, for general $c \in (0, 1]$ and $\mu_{\mathbb{C}} = \text{SLE}_{\kappa}^{\text{loop}}$, the exact value of ζ remains unknown.

Organization of the paper. In Section 2, we present the proof of Theorems 1.2 and 1.3, and discuss its radial counterpart. In Section 3, we prove Theorems 1.5 and 1.6. Then we show Theorem 1.9 in Section 4. Finally in Section 5, we give some further remarks and discussions.

2 Generalized chordal and radial restriction measures

2.1 The chordal case

In this section, we prove Theorem 1.3 by using a Markovian exploration process (Definition 2.1) and a recursive formula (Proposition 2.3) derived from the restriction property (1.2). Based on these results, we then finish the proof Theorem 1.2. We first focus on the one-sided case. Let $c \in (0, 1]$, $\alpha \in \mathbb{R}$, and $A \in \mathcal{Q}_-$. Let **P** and **E** be the law and the expectation with respect to the Brownian loop soup $\mathcal{L}_{\mathbb{D}}$ on \mathbb{D} (of intensity $\frac{c}{2}$ as assumed). Now, we define a natural exploration process associated with a set $A \in \mathcal{Q}_-$ by revealing the loops connected to A in $\mathcal{L}_{\mathbb{D}}$ in a Markovian way.

Definition 2.1 (Exploration process). Suppose $A \in Q_-$. Let $A_0 := A$, and for all $n \ge 0$, let A_{n+1} be the filling of the union of A_n and all loops in $\mathcal{L}_{\mathbb{D}}$ that intersect A_n .

As introduced, \widetilde{A} denotes the filling of the union of A and all clusters in $\mathcal{L}_{\mathbb{D}}$ that intersect A. For clarity, we further choose a point $w_0 \in \mathbb{S}^1 \setminus A$ other than -1, 1, and let f_{A_n} (resp. $f_{\widetilde{A}}$) be the unique conformal map from $\mathbb{D} \setminus A_n$ (resp. $\mathbb{D} \setminus \widetilde{A}$) to \mathbb{D} that fixes $-1, 1, w_0$. For sets $B_1, B_2 \subset \mathbb{C}$, the Hausdorff distance between B_1 and B_2 is given by

$$d_H(B_1, B_2) := \inf\{\varepsilon > 0 : B_1 \subset B_2^{(\varepsilon)}, B_2 \subset B_1^{(\varepsilon)}\}, \text{ where } B_i^{(\varepsilon)} := \{z : \operatorname{dist}(z, B_i) < \varepsilon\}.$$

We first show that the exploration process A_n converges to \widetilde{A} in Hausdorff distance.

Lemma 2.2. For any $A \in Q_{-}$, we have **P**-a.s. $\lim_{n\to\infty} d_H(A_n, \widetilde{A}) = 0$, and therefore, $f_{A_n}^{-1} \to f_{\widetilde{A}}^{-1}$ under the locally uniform topology.

Proof. For any $\varepsilon > 0$, by [SW12, Lemma 9.7], there are only finitely many clusters in $\mathcal{L}_{\mathbb{D}}$ with diameter larger than ε . We denote those that intersect A by $\mathcal{C}_1, \ldots, \mathcal{C}_m$, which are at positive distance from each other. Below, we use Fill to denote the filling of a set for brevity. Then, $A' := \operatorname{Fill}(A \cup \mathcal{C}_1 \cup \ldots \cup \mathcal{C}_m) \subset \widetilde{A}$, and $d_H(A', \widetilde{A}) \leq \varepsilon$. According to [vdBCL16, Theorem 4.1], every cluster \mathcal{C}_i , $1 \leq i \leq m$, can be approximated by a sequence of finite subclusters \mathcal{C}_i^N of \mathcal{C}_i (i.e., a chain of finite number of loops in \mathcal{C}_i) such that $\lim_{N\to\infty} d_H(\operatorname{Fill}(\mathcal{C}_i^N), \operatorname{Fill}(\mathcal{C}_i)) = 0$. Since each finite subcluster \mathcal{C}_i^N is contained in A_n for sufficiently large n, we conclude $\lim_{n\to\infty} d_H(A_n, \widetilde{A}) = 0$. Once $A_n \to \widetilde{A}$ in Hausdorff distance, $\{f_{A_n}^{-1}\}_{n\geq 1}$ form a normal family, and for each convergent subsequence $\{f_{A_{n_k}}^{-1}\}$, its limit is a conformal map from \mathbb{D} to \widetilde{A} , hence equals $f_{\widetilde{A}}^{-1}$, as desired.

We derive the following recursive formula for the non-intersection probability of restriction samples by the loop-soup interpolation.

Proposition 2.3. Let $c \in (0,1]$ and $\alpha \in \mathbb{R}$. Suppose \mathbb{P} satisfies the one-sided chordal c-restriction with exponent α as in Definition 1.1. For any $A \in \mathcal{Q}_{-}$, we have

$$\mathbb{P}[K \cap A = \emptyset] = \mathbf{E} \otimes \mathbb{E}\left[(f'_{A_n}(1)f'_{A_n}(-1))^{\alpha} \exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}}\left((f_{A_{n-1}} \circ f_{A_n}^{-1})(K), f_{A_{n-1}}(A_n \setminus A_{n-1})\right)\right) \right].$$

Proof. According to (1.2), it holds that

$$\mathbb{P}[K \cap A = \emptyset] = \mathbb{E}_A \left[(f'_A(1)f'_A(-1))^{\alpha} \exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}}(K,A)\right) \right]$$
$$= \mathbf{E} \otimes \mathbb{E}_A \left[(f'_A(1)f'_A(-1))^{\alpha} \mathbf{1}_{K \cap A_1 = \emptyset} \right]$$
(2.1)

$$= \mathbf{E} \otimes \mathbb{E}_{A_1} \left[(f'_{A_1}(1)f'_{A_1}(-1))^{\alpha} \exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}\setminus A}(K,A_1)\right) \right],$$
(2.2)

where we used $\mathbf{P}[K \cap A_1 = \emptyset] = \exp(-\frac{c}{2}\Lambda_{\mathbb{D}}(K, A))$ given K such that $K \cap A = \emptyset$ in (2.1), and $\frac{d\mathbb{P}_A(K)}{d\mathbb{P}_{A_1}(K)}\mathbf{1}_{K\cap A_1=\emptyset} = \mathbf{1}_{K\cap A_1=\emptyset}(\phi'(1)\phi'(-1))^{\alpha}\exp(-\frac{c}{2}\Lambda_{\mathbb{D}\setminus A}(K, A_1))$ for $\phi = f_{A_1} \circ f_A^{-1}$ in (2.2). By induction, we have for any $n \ge 1$,

$$\mathbb{P}[K \cap A = \emptyset] = \mathbf{E} \otimes \mathbb{E}_{A_n} \left[(f'_{A_n}(1)f'_{A_n}(-1))^{\alpha} \exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}\setminus A_{n-1}}(K,A_n)\right) \right].$$

Now, by the conformal invariance of Brownian loop measure,

$$\Lambda_{\mathbb{D}\setminus A_{n-1}}(K,A_n) = \Lambda_{\mathbb{D}}(f_{A_{n-1}}(K), f_{A_{n-1}}(A_n \setminus A_{n-1})).$$

Moreover, if K is sampled from \mathbb{P} , then $f_{A_n}^{-1}(K)$ is distributed as \mathbb{P}_{A_n} . The lemma then follows. \Box

Next, we give a simple lemma on the Brownian loop soup.

Lemma 2.4. Let $0 < a < \frac{1}{2}$. Suppose *B* is a connected subset of \mathbb{D} such that $\partial B \cap \mathbb{S}^1 \neq \emptyset$ and $B \cap (1-a)\mathbb{S}^1 \neq \emptyset$. Let *H* be the event that there is a loop in $\mathcal{L}_{\mathbb{D}}$ that intersects both *B* and $\frac{1}{2}\mathbb{D}$. Then, there exists a constant $\delta = \delta(c, a) > 0$ such that for all *B*,

$$\mathbf{P}[H] \ge \delta.$$

Proof. Let z be some point on $B \cap (1-a)\mathbb{S}^1$. Let L be the line segment connecting z and z/(2|z|). For any b > 0, define $L(b) := \{w : \operatorname{dist}(w, L) \le b\}$. Let $\Lambda(z)$ be the set of loops in $L(\frac{a}{10}) \setminus L(\frac{a}{20})$ that surround $L(\frac{a}{20})$. Note that B contains a curve intersecting both $\partial L(\frac{a}{10})$ and $\partial L(\frac{a}{20})$. Every loop in $\Lambda(z)$ will intersect B. Hence, $\mathbf{P}[H] \ge \mathbf{P}[\Lambda(z) \neq \emptyset]$. By rotation invariance, $\mu^{\mathrm{BL}}[\Lambda(z)] \ge \delta'(a)$ for some constant $\delta'(a)$ independent of z. This allows us to conclude the proof.

The following lemma shows that $f_{A_{n-1}}(A_n \setminus A_{n-1})$ will stay close to the boundary as $n \to \infty$.

Lemma 2.5. For any $A \in \mathcal{Q}_{-}$, let $d_n := \sup\{\operatorname{dist}(z, \mathbb{S}^1) \mid z \in f_{A_{n-1}}(A_n \setminus A_{n-1})\}$. Under $\mathbf{P}, d_n \to 0$ in probability.

Proof. Otherwise, there is $\varepsilon_0 > 0$, $\rho > 0$ and a sequence $n_i \to \infty$ such that $\mathbf{P}[d_{n_i} > \varepsilon_0] > \rho$. Then by Lemma 2.4, given $d_{n_i} > \varepsilon_0$, the probability that there is a loop in $\mathcal{L}_{\mathbb{D}}$ intersecting both $f_{A_{n_i-1}}(A_{n_i} \setminus A_{n_i-1})$ and $\frac{1}{2}\mathbb{D}$ is bounded from below by a constant $\delta(\varepsilon_0) > 0$ only depending on ε_0 . Hence, by conformal invariance of the Brownian loop soup,

$$\mathbf{P}\Big[f_{A_{n_i-1}}(A_{n_i+1} \setminus A_{n_i}) \cap \left(\frac{1}{2}\mathbb{D}\right) \neq \emptyset\Big] > \rho\delta(\varepsilon_0).$$
(2.3)

By Lemma 2.2, $f_{A_n}^{-1} \to f_{\widetilde{A}}^{-1}$ locally uniformly a.s. Then for any $\sigma > 0$ and sufficiently large n,

$$\mathbf{P}\left[f_{A_n}^{-1}\left(\frac{1}{2}\mathbb{D}\right) \not\subset f_{\widetilde{A}}^{-1}\left(\frac{1+\sigma}{2}\mathbb{D}\right)\right] < \frac{1}{2}\rho\delta(\varepsilon_0).$$
(2.4)

Combining with (2.3), we obtain that $\mathbf{P}\left[\left(A_{n_i+1} \setminus A_{n_i}\right) \cap f_{\widetilde{A}}^{-1}\left(\frac{1+\sigma}{2}\mathbb{D}\right) \neq \emptyset\right] > \frac{1}{2}\rho\delta(\varepsilon_0)$ for sufficiently large n_i , which contradicts with the fact that $A_{n_i+1} \subset \widetilde{A}$ and $\widetilde{A} \cap f_{\widetilde{A}}^{-1}\left(\frac{1+\sigma}{2}\mathbb{D}\right) = \emptyset$.

The following lemma deals with the behavior of $(f_{A_n})_{n\geq 1}$ near the boundary points -1, 1.

Lemma 2.6. P-almost surely, for any $A \in Q_{-}$, all of the followings hold. First, $f'_{A_n}(\pm 1) \downarrow f'_{\widetilde{A}}(\pm 1)$. Next, the distance between $f_{A_{n-1}}(A_n \setminus A_{n-1})$ and $\{-1,1\}$ has a (random) uniform positive lower bound. Furthermore, for any $K \in \mathcal{K}$, $(f_{A_{n-1}} \circ f^{-1}_{A_n})(K) \to K$ in Hausdorff distance.

Proof. Let $B_{\varepsilon}(1) := \{z : |z-1| < \varepsilon\}$. Let $\varepsilon < \frac{1}{100}$ be such that $B_{2\varepsilon}(1) \cap \widetilde{A} = \emptyset$ and $f_{\widetilde{A}}(B_{\varepsilon}(1)) \cap \frac{1}{2}\mathbb{D} = \emptyset$. By Schwarz reflection for circular arcs, f_{A_n} (for sufficiently large n) and $f_{\widetilde{A}}$ can be analytically extended to functions φ_n and φ defined on $B_{\varepsilon}(1)$, respectively. Then $\{\varphi_n\}$ forms a normal family and every limit in this family, when restricted to $B_{\varepsilon}(1) \cap \mathbb{D}$, is identical to $\varphi|_{B_{\varepsilon}(1)\cap\mathbb{D}} = f_{\widetilde{A}}|_{B_{\varepsilon}(1)\cap\mathbb{D}}$ by Lemma 2.2. By the uniqueness of analytic function, we then conclude $\varphi_n \to \varphi$ locally uniformly, and as an interior point in $B_{\varepsilon}(1)$, $f'_{A_n}(1) = \varphi'_n(1)$ converges to $f'_{\widetilde{A}}(1) = \varphi'(1)$. In particular, $|\varphi'_n(1)| \ge C$ for some random C > 0 and hence $\varphi_n(B_{\varepsilon}(1)) \supset B_{C\varepsilon/4}(1)$ by Koebe's 1/4 theorem. Hence $f_{A_{n-1}}(A_n \setminus A_{n-1})$, as a subset of $f_{A_{n-1}}(\widetilde{A} \setminus A_{n-1})$, has distance $\ge \frac{1}{4}C\varepsilon$ from 1. Similar results hold for the case -1. The last claim follows directly from the locally uniform convergence of f_{A_n} and φ_n .

Finally, we are ready to show Theorem 1.3.

Proof of Theorem 1.3. We first finish the proof for the one-sided case, which requires the use of Proposition 2.3 by taking $n \to \infty$. By Lemmas 2.5 and 2.6, we conclude that under $\mathbf{P} \otimes \mathbb{P}$,

$$\Lambda_{\mathbb{D}}\left((f_{A_{n-1}} \circ f_{A_n}^{-1})(K), f_{A_{n-1}}(A_n \setminus A_{n-1})\right) \to 0 \text{ in probability.}$$

Note that **P**-a.s. $f'_{A_n}(\pm 1) \in (0,1)$ and $f'_{A_n}(\pm 1) \downarrow f'_{\widetilde{A}}(\pm 1)$ by Lemma 2.6. Therefore,

$$\mathbf{E} \otimes \mathbb{E} \left[(f'_{A_n}(1)f'_{A_n}(-1))^{\alpha} \exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}}\left((f_{A_{n-1}} \circ f_{A_n}^{-1})(K), f_{A_{n-1}}(A_n \setminus A_{n-1})\right)\right) \right] \rightarrow \mathbf{E} \left[\left(f'_{\widetilde{A}}(1)f'_{\widetilde{A}}(-1)\right)^{\alpha} \right] \qquad \text{as } n \to \infty,$$

$$(2.5)$$

where one can use the monotone convergence theorem to show that \limsup of LHS is smaller than RHS, and use Fatou's lemma to show that \liminf of LHS is greater than RHS. Hence, by Proposition 2.3, we conclude $\mathbb{P}[K \cap A = \emptyset] = \mathbf{E}[(f'_{\widetilde{A}}(1)f'_{\widetilde{A}}(-1))^{\alpha}]$ and finish the one-sided case.

The two-sided case is proved in a similar way. However, compared to the one-sided case, now it could happen that -1 and 1 are separated by \widetilde{A} inside \mathbb{D} . Below, we only give the necessary modifications and omit further details. Let \mathbb{P} be the two-sided chordal *c*-restriction measure with exponent α (Definition 1.1). For $A \in \mathcal{Q}$, the exploration process $(A_n)_{n\geq 0}$ in Definition 2.1 is still valid, and **P**-a.s. $\lim_{n\to\infty} d_H(A_n, \widetilde{A}) = 0$. Furthermore, if we let E_n be the event that -1 and 1 are in the same connected component of $\overline{\mathbb{D}} \setminus A_n$, then **P**-a.s. $\mathbf{1}_{E_n} \downarrow \mathbf{1}_E$, where *E* is defined in (1.3). The same argument as in the proof of Proposition 2.3 now gives

$$\mathbb{P}[K \cap A = \emptyset] = \mathbf{E} \otimes \mathbb{E}\left[\mathbf{1}_{E_n}(f'_{A_n}(1)f'_{A_n}(-1))^{\alpha} \exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}}\left((f_{A_{n-1}} \circ f^{-1}_{A_n})(K), f_{A_{n-1}}(A_n \setminus A_{n-1})\right)\right)\right].$$

For $A \in \mathcal{Q}$ and on the event E, analogues of Lemmas 2.5 and 2.6 hold as well. Then similar to (2.5), the above converges to $\mathbf{E}[\mathbf{1}_E(f'_{\widetilde{A}}(1)f'_{\widetilde{A}}(-1))^{\alpha}]$, which implies (1.4), as desired.

Proof of Theorem 1.2. Let $c \in (0, 1]$. The existence of the two-sided chordal *c*-restriction with exponent α for $\alpha \geq \frac{6-\kappa}{2\kappa}$ is given by [Qia21, Proposition 6.2]. Note that the proof of Proposition 1.7 indeed gives the construction of the one-sided case via chordal $SLE_{\kappa}(\rho)$.

The uniqueness part of Theorem 1.2 follows immediately from Theorem 1.3. It remains to show that the two-sided (resp. one-sided) chordal *c*-restriction measure with exponent α exists if and only if $\alpha \geq \frac{6-\kappa}{2\kappa}$ (resp. $\alpha > 0$). For the one-sided case, when $\alpha \leq 0$, the right side of (1.4) is always ≥ 1 , which cannot be the case. The argument for the two-sided case is similar to [LSW03, Corollary 8.6]. Suppose that there is a probability measure \mathbb{P} on random sets $K \in \mathcal{K}$ satisfying the two-sided chordal *c*-restriction with exponent α for some $\alpha < \frac{6-\kappa}{2\kappa}$. Then the lower boundary γ of K satisfies the one-sided chordal *c*-restriction with the same exponent α ; hence $\alpha > 0$ and γ is an $\mathrm{SLE}_{\kappa}(\rho)$ for some $\rho < 0$. In particular, $\mathbb{P}[0$ is above $\gamma] < \frac{1}{2}$. However, the symmetry of \mathbb{P} implies $\mathbb{P}[0$ is above $\gamma]$ is at least $\frac{1}{2}$ (it can be strictly larger when K is of positive Lebesgue measure), a contradiction.

2.2 The radial case

In this section, we provide the radial counterpart. To lighten notation, we use the same symbols as the chordal case to denote their radial counterparts. Let \mathcal{K} be the collection of all simply connected compact sets $K \subset \overline{\mathbb{D}}$ such that $0 \in K$ and $K \cap \mathbb{S}^1 = \{1\}$. Let \mathcal{Q} be the collection of all compact sets $A \subset \overline{\mathbb{D}}$ such that $\mathbb{D} \setminus A$ is simply connected and $0, 1 \notin A$. For any $A \in \mathcal{Q}$, let $f_A : \mathbb{D} \setminus A \to \mathbb{D}$ be the conformal map that fixes 0, 1. The following definition for general $c \leq 1$ is from [Qia21, Definition 1.3], which extends the c = 0 case considered in [Wu15].

Definition 2.7. Let $c \leq 1$ and $\alpha, \beta \in \mathbb{R}$. We say that a probability measure \mathbb{P} on \mathcal{K} satisfies the radial *c*-restriction with exponents (α, β) , if for all $A \in \mathcal{Q}$,

$$\frac{d\mathbb{P}(K)}{d\mathbb{P}_A(K)}\mathbf{1}_{K\cap A=\emptyset} = \mathbf{1}_{K\cap A=\emptyset} |f'_A(0)|^{\alpha} f'_A(1)^{\beta} \exp\left(-\frac{c}{2}\Lambda_{\mathbb{D}}(K,A)\right),\tag{2.6}$$

where $\mathbb{P}_A = \mathbb{P} \circ f_A$ and $\Lambda_{\mathbb{D}}(K, A)$ is the same as in Definition 1.1.

In [Qia21, Theorem 1.6], the author used the radial hypergeometric SLE_{κ} to construct a probability measure $\mathbb{P}_{\kappa}^{\alpha,\beta}$, which satisfies the radial *c*-restriction with exponents (α,β) , in the range $\alpha \leq \eta_{\kappa}(\beta)$ and $\beta \geq \frac{6-\kappa}{2\kappa}$, where $\eta_{\kappa}(\beta)$ is the generalized disconnection exponent [Qia21, (1.7)]. The author further conjectured that for any $c \leq 1$ ($\kappa \in (0,4]$), the restriction property (2.6) uniquely characterizes the law of \mathbb{P} . Up to now, the only proved case is c = 0 [Wu15]. Using the approach similar to the chordal case, we are able to confirm the uniqueness for $c \in (0,1]$.

Theorem 2.8. Let $c \in (0,1]$ and $\alpha, \beta \in \mathbb{R}$. Suppose \mathbb{P} satisfies the radial *c*-restriction with exponents (α, β) . Then, for all $A \in \mathcal{Q}$, we have

$$\mathbb{P}[K \cap A = \emptyset] = \mathbf{E}\left[\mathbf{1}_{0 \notin \widetilde{A}} |f'_{\widetilde{A}}(0)|^{\alpha} f'_{\widetilde{A}}(1)^{\beta}\right], \qquad (2.7)$$

where \mathbf{E} denotes the expectation for the Brownian loop soup $\mathcal{L}_{\mathbb{D}}$ of intensity $\frac{c}{2}$, \widetilde{A} is the filling of the union of A and all clusters in $\mathcal{L}_{\mathbb{D}}$ that intersect A, and $f_{\widetilde{A}}$ is the conformal map from $\mathbb{D} \setminus \widetilde{A}$ to \mathbb{D}

that fixes 0,1. As a result, for any $c \in (0,1]$, the probability measure on \mathcal{K} that satisfies the radial *c*-restriction with exponents (α, β) as in Definition 2.7 is unique.

Proof. We only need to show (2.7). For any $A \in \mathcal{Q}$, the exploration process $(A_n)_{n\geq 0}$ given in Definition 2.1 is still valid, and we have **P**-a.s. $\lim_{n\to\infty} d_H(A_n, \widetilde{A}) = 0$. Moreover, **P**-a.s. $\mathbf{1}_{0\notin A_n} \downarrow \mathbf{1}_{0\notin \widetilde{A}}$. Similar to Proposition 2.3, we can use (2.6) to deduce the following formula

$$\mathbb{P}[K \cap A = \emptyset] = \mathbf{E} \otimes \mathbb{E} \left[\mathbf{1}_{0 \notin A_n} |f'_{A_n}(0)|^{\alpha} f'_{A_n}(1)^{\beta} \exp\left(-\frac{c}{2} \Lambda_{\mathbb{D}} \left((f_{A_{n-1}} \circ f_{A_n}^{-1})(K), f_{A_{n-1}}(A_n \setminus A_{n-1}) \right) \right) \right].$$
(2.8)

On the event $\{0 \notin \widetilde{A}\}$, we have $f_{A_n}^{-1} \to f_{\widetilde{A}}^{-1}$, $|f'_{A_n}(0)| \uparrow |f'_{\widetilde{A}}(0)|$, $f'_{A_n}(1) \downarrow f'_{\widetilde{A}}(1)$, and the exponential term tends to 1 as before. Letting $n \to \infty$ in (2.8) and similar to (2.5), we arrive at (2.7).

Consequently, we obtain the following result, which plays a crucial role in determining the Hausdorff dimension of certain exceptional sets arising from loop-soup clusters [GLQ22].

Proposition 2.9. Let $\kappa \in (\frac{8}{3}, 4]$, $\beta \geq \frac{5}{8}$ and $\alpha \leq \eta(\beta)$, where $\eta(\beta) = \eta_{8/3}(\beta)$ is the Brownian disconnection exponent. Let K be a standard radial restriction sample in \mathbb{D} with exponents (α, β) and marked points 0 and 1 (i.e. the c = 0 case in Definition 2.7). Let Γ be an independent CLE_{κ} on \mathbb{D} . Let $\mathcal{C}(K)$ be the union of K and all loops of Γ that K intersects. Then, the filling of $\mathcal{C}(K)$ has the same law as $\mathbb{P}_{\kappa}^{\alpha,\beta}$, which is constructed by the radial hypergeometric SLE_{κ} in [Qia21, Section 4.1].

Proof. Note that both the law of the filling of $\mathcal{C}(K)$ and $\mathbb{P}_{\kappa}^{\alpha,\beta}$ satisfy the radial *c*-restriction with exponents (α,β) ; see Theorem 1.6 and Lemma 2.1 of [Qia21], respectively. Now, Theorem 2.8 gives the equivalence between these two.

Remark 2.10. Our analysis can be also adapted to the trichordal case; see [Qia18] and [Qia21, Section 6.2.2] for related notions. We leave it to the interested reader.

3 The Malliavin-Kontsevich-Suhov measure

This section focuses on the MKS loop measure $\mu_{\mathbb{C}}$ (see Definition 1.4) on the whole-plane. We will prove Theorem 1.6 in Section 3.2, which implies Theorem 1.5 immediately.

As mentioned before, since the Brownian loop soup on \mathbb{C} a.s. has only one cluster for any c > 0, we will use the $\mathrm{SLE}_{8/3}$ loop soup on \mathbb{C} instead, as considered in [KW16]. Recall that Werner's $\mathrm{SLE}_{8/3}$ loop measure \mathcal{W} on \mathbb{C} is induced from the Brownian loop measure μ^{BL} on \mathbb{C} by taking outer boundaries. For an MKS loop measure $\mu_{\mathbb{C}}$, on any simply connected $D \subset \mathbb{C}$, define $\hat{\mu}_D$ by

$$\frac{d\widehat{\mu}_D(\eta)}{d\mu_{\mathbb{C}}(\eta)} \mathbf{1}_{\eta \subset D} = \mathbf{1}_{\eta \subset D} \exp\left(\frac{c}{2}\mathcal{W}(\eta, \partial D)\right).$$
(3.1)

We first show that the above $(\hat{\mu}_D)_{D \subset \mathbb{C}}$ is also conformally invariant as the family $(\mu_D)_{D \subset \mathbb{C}}$ defined in (1.5), based on a relation between \mathcal{W} and μ^{BL} obtained by Carfagnini and Wang [CW24]. **Lemma 3.1.** $(\widehat{\mu}_D)_{D \subset \mathbb{C}}$ defined above is conformally invariant. That is, for $\phi : D \to D'$ conformal, the pushforward of $\widehat{\mu}_D$ under ϕ equals $\widehat{\mu}_{D'}$.

Proof. Since $\mu_{\mathbb{C}}$ is an MKS measure, we have $\phi \circ \mu_D = \mu_{D'}$, and by (1.5),

$$\int_{\{\eta \subset D\}} F(\phi(\eta)) \exp\left(\frac{c}{2}\Lambda^*(\eta, \partial D)\right) \mu_{\mathbb{C}}(d\eta) = \int_{\{\eta \subset D'\}} F(\eta) \exp\left(\frac{c}{2}\Lambda^*(\eta, \partial D')\right) \mu_{\mathbb{C}}(d\eta)$$

for any positive measurable F defined on loops contained in D'. Now we choose

$$F(\eta) = \exp\left(\frac{c}{2}(\mathcal{W}(\phi^{-1}(\eta), \partial D) - \Lambda^*(\phi^{-1}(\eta), \partial D))\right) G(\eta).$$

By [CW24, Theorem 2.5] and the remark after it, we have

$$F(\eta) = \exp\left(\frac{c}{2}(\mathcal{W}(\eta, \partial D') - \Lambda^*(\eta, \partial D'))\right) G(\eta).$$

Hence we obtain

$$\int_{\{\eta \subset D\}} G(\phi(\eta)) \exp\left(\frac{c}{2} \mathcal{W}(\eta, \partial D)\right) \mu_{\mathbb{C}}(d\eta) = \int_{\{\eta \subset D'\}} F(\eta) \exp\left(\frac{c}{2} \mathcal{W}(\eta, \partial D')\right) \mu_{\mathbb{C}}(d\eta),$$

which is exactly that $\phi \circ \hat{\mu}_D = \hat{\mu}_{D'}$, as desired.

We use a superscript 0 to denote the corresponding measure restricted to loops surrounding the origin. As in Section 2, we will use the $SLE_{8/3}$ loop soup observables to express quantities of $\mu^0_{\mathbb{C}}$. Let $\mathcal{L}^{\mathcal{W}}_{\mathbb{C}}$ be the $SLE_{8/3}$ loop soup on \mathbb{C} of itensity $\frac{c}{2}$, and \mathbf{P}, \mathbf{E} be the law and the expectation with respect to $\mathcal{L}^{\mathcal{W}}_{\mathbb{C}}$. Recall the notation introduced above Theorem 1.6. In the following, for any function f defined on some $\Omega \subset \mathbb{C}$, we let $f(A) := f(A \cap \Omega)$ for any $A \subset \mathbb{C}$.

Proposition 3.2. For any 0 < r < 1, we have $\widehat{\mu}^0_{\mathbb{D}}[\ell \not\subset r\mathbb{D}] < \infty$. Furthermore, for any simply connected $D \subset \mathbb{D}$ containing the origin, we have

$$\mu^{0}_{\mathbb{C}}[\ell \not\subset r\mathbb{D}, \ell \subset D] = \mathbf{E} \otimes \widehat{\mu}^{0}_{\mathbb{D}}[\ell \not\subset f_{D,\infty}(r\mathbb{D})].$$
(3.2)

Proof. We use ideas similar to Section 2. First, we define an exploration process analogous to Definition 2.1 as follows. Let $D_0 := D$, and $D_{n+1} \subset D_n$ be the connected component containing 0 of the complement of the union of ∂D_n and all loops in $\mathcal{L}^{\mathcal{W}}_{\mathbb{C}}$ that intersect ∂D_n . Using (3.1), we obtain that

$$\mu^{0}_{\mathbb{C}}[\ell \not\subset r\mathbb{D}, \ell \subset D] = \widehat{\mu}^{0}_{D} \Big[\mathbf{1}_{\ell \not\subset r\mathbb{D}} \exp\left(-\frac{c}{2} \mathcal{W}(\ell, \partial D)\right) \Big] = \mathbf{E} \otimes \widehat{\mu}^{0}_{D}[\ell \not\subset r\mathbb{D}, \ell \subset D_{1}].$$
(3.3)

Let $f_{D,n}$ be the conformal map from D_n to \mathbb{D} such that $f_{D,n}(0) = 0$ and $f'_{D,n}(0) > 0$. Iterating (3.3), we conclude that for all n,

$$\mu^{0}_{\mathbb{C}}[\ell \not\subset r\mathbb{D}, \ell \subset D] = \mathbf{E} \otimes \widehat{\mu}^{0}_{D_{n}}[\ell \not\subset r\mathbb{D}, \ell \subset D_{n+1}] = \mathbf{E} \otimes \widehat{\mu}^{0}_{\mathbb{D}}[\ell \not\subset f_{D,n}(r\mathbb{D}), \ell \subset f_{D,n}(D_{n+1})], \quad (3.4)$$

where we used the conformal invariance of $\hat{\mu}_{\mathbb{D}}$ from Lemma 3.1. Using Lemma 2.2, together with the finiteness of clusters in $\mathcal{L}^{\mathcal{W}}_{\mathbb{C}}$ [KW16], and the inversion invariance of $\mathcal{L}^{\mathcal{W}}_{\mathbb{C}}$ [Wer08], we have $D_n \to D_{\infty}$ in Hausdorff distance, and $f_{D,n}^{-1} \to f_{D,\infty}^{-1}$ in the locally uniform topology. In particular, we have

$$\mathbf{1}_{\ell \not\subset f_{D,n}(r\mathbb{D}), \ell \subset f_{D,n}(D_{n+1})} \to \mathbf{1}_{\ell \not\subset f_{D,\infty}(r\mathbb{D})}, \quad \mathbf{E} \otimes \widehat{\mu}^0_{\mathbb{D}} - \text{a.e.}$$
(3.5)

Then, by Fatou's Lemma, taking $n \to \infty$ in (3.4), we obtain

$$\infty > \mu^0_{\mathbb{C}}[\ell \not\subset r\mathbb{D}, \ell \subset D] \ge \mathbf{E} \otimes \widehat{\mu}^0_{\mathbb{D}}[\ell \not\subset f_{D,\infty}(r\mathbb{D})].$$
(3.6)

Now, (3.6) implies that $\widehat{\mu}^0_{\mathbb{D}}[\ell \not\subset r\mathbb{D}] < \infty$. Indeed, there is a positive probability p_0 under **P** such that $\mathbb{D}_{\infty} \supset \frac{1}{2}\mathbb{D}$. Note that $\mathbb{D}_{\infty} \supset \frac{1}{2}\mathbb{D}$ implies $f_{\mathbb{D},\infty}(r\mathbb{D}) \subset 2r\mathbb{D}$ by Schwarz lemma. Then (3.6) gives

$$\infty > \mathbf{E} \otimes \widehat{\mu}^0_{\mathbb{D}}[\ell \not\subset f_{\mathbb{D},\infty}(r\mathbb{D})] \ge p_0 \,\widehat{\mu}^0_{\mathbb{D}}[\ell \not\subset 2r\mathbb{D}]. \tag{3.7}$$

It remains to show (3.2). We go back to (3.4). By Schwarz lemma again, $D_n \subset \mathbb{D}$ gives $r\mathbb{D} \subset f_{D,n}(r\mathbb{D})$. Hence, we can rewrite (3.4) as

$$\mu^{0}_{\mathbb{C}}[\ell \not\subset r\mathbb{D}, \ell \subset D] = \mathbf{E} \otimes \widehat{\mu}^{0}_{\mathbb{D}}[\ell \not\subset f_{D,n}(r\mathbb{D}), \ell \subset f_{D,n}(D_{n+1}); \ \ell \not\subset r\mathbb{D}].$$
(3.8)

Now we know from (3.7) that $\mathbf{E} \otimes \hat{\mu}^0_{\mathbb{D}}[\cdot; \ell \not\subset r\mathbb{D}]$ is a finite measure for any fixed r. Then, taking $n \to \infty$ in (3.8), by (3.5) and the dominated convergence, we finally obtain (3.2).

In the following, we first provide several basic estimates on $\mu_{\mathbb{C}}^0$ and $\hat{\mu}_{\mathbb{D}}^0$ in Section 3.1, and then finish the proof of Theorems 1.5 and 1.6 in Section 3.2.

3.1 A priori estimates on $\mu_{\mathbb{C}}^0$ and $\hat{\mu}_{\mathbb{D}}^0$

Let $\|\ell\|_{\infty} := \max_{z \in \ell} |z|$ for a simple loop $\ell \subset \mathbb{C}$.

Lemma 3.3. There is a $\lambda \in (0,\infty)$ such that for any a > b > 0, $\mu^0_{\mathbb{C}}(\|\ell\|_{\infty} \in [a,b)) = \lambda \log \frac{b}{a}$.

Proof. Let $U(a,b) := \mu_{\mathbb{C}}^0(\|\ell\|_{\infty} \in [a,b))$. Then U(a,b) + U(b,c) = U(a,c) for any 0 < a < b < c. Furthermore, the scaling invariance of $\mu_{\mathbb{C}}^0$ gives U(a,b) = U(ka,kb) for all k > 0. Then $x \mapsto U(1,x)$ is increasing with U(1,xy) = U(1,x) + U(1,y); thus $U(1,x) = \lambda \log x$ for some $\lambda > 0$.

Corollary 3.4. For $c \in (0,1]$, let $\mu_{\mathbb{C}} = \text{SLE}_{\kappa}^{\text{loop}}$, i.e. the counting measure on full-plane CLE_{κ} loops, then the λ in Lemma 3.3 is given by $\lambda = \frac{1}{\pi} (\frac{\kappa}{4} - 1) \cot(\pi(1 - \frac{4}{\kappa}))$.

Proof. Let Γ be sampled from the full-plane $\operatorname{CLE}_{\kappa}$, and N_C be the number of loop ℓ 's in Γ surrounding the origin with $\|\ell\|_{\infty} \in [1, e^C)$. By [ACSW24b, Lemma 9.2] (which is based on [SSW09]), we have $\frac{\mathbb{E}[N_C]}{C} \to \frac{1}{\pi}(\frac{\kappa}{4}-1)\cot(\pi(1-\frac{4}{\kappa}))$ as $C \to \infty$. Comparing with Lemma 3.3, the result then follows. \Box

We also mention that for c = 0 and $\mu_{\mathbb{C},c=0}$ replaced by Werner's measure \mathcal{W} , the corresponding λ in Lemma 3.3 equals $\frac{\pi}{5}$ [Wer08, Page 151].

Lemma 3.5. There is an $\alpha > 0$ such that for any $0 < \varepsilon < \frac{1}{2}$, $\mathcal{W}(\varepsilon \mathbb{D}, \mathbb{S}^1) = O(\varepsilon^{\alpha})$.

Proof. This basically follows from [NW11]. We add some details here for completeness. First, by [NW11, Lemma 2] and scaling invariance of \mathcal{W} , we have

$$\mathcal{W}(\varepsilon \mathbb{D}, \mathbb{S}^1) \le \int_0^{2\varepsilon} \frac{1}{r} \mathbb{E}(\operatorname{Area}(\{z : \operatorname{dist}(z, \gamma) \le r\})) \, dr, \tag{3.9}$$

where Area denotes the Lebesgue measure on \mathbb{R}^2 , and γ under \mathbb{E} has the law of the outer boundary of the Brownian loop of time-length 1. By the proof of [NW11, Lemma 4], the expected area of *r*-neighborhood of γ is $O(r^{\alpha})$ for some constant $\alpha > 0$. This concludes the proof from (3.9).

In fact, the optimal α in Lemma 3.5 is $\frac{2}{3}$, which can be deduced from the Brownian disconnection exponents. However, we will not need such an explicit bound.

Proposition 3.6. For any $c \in (0,1]$, there is a constant $\zeta_1 \in (0,\infty)$ such that $\hat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in [\varepsilon,1)) = \lambda |\log \varepsilon| + \zeta_1 + o(1)$ as $\varepsilon \to 0$.

Proof. By Proposition 3.2, $\hat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in [\varepsilon, 1)) < \infty$. By (3.1), Lemmas 3.3 and 3.5, we have

$$0 \le \widehat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in [\varepsilon, 2\varepsilon)) - \mu^0_{\mathbb{C}}(\|\ell\|_{\infty} \in [\varepsilon, 2\varepsilon)) \le \mu^0_{\mathbb{C}}(\|\ell\|_{\infty} \in [\varepsilon, 2\varepsilon)) \left(\exp(\frac{c}{2}\mathcal{W}(2\varepsilon\mathbb{D}, \mathbb{S}^1)) - 1\right) = O(\varepsilon^{\alpha}).$$

This implies that $\hat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in [\varepsilon, 1)) - \mu^0_{\mathbb{C}}(\|\ell\|_{\infty} \in [\varepsilon, 1))$ converges to some $\zeta_1 \in (0, \infty)$ as $\varepsilon \to 0$. \Box

Corollary 3.7. Let $\beta \in [0, \frac{1}{2}]$. We have

$$\widehat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in [\varepsilon, a\varepsilon)) = \lambda \log a + o(1) \quad as \quad \varepsilon \to 0,$$

uniformly for all $1 \le a \le \varepsilon^{-\beta}$, where o(1) is independent of β .

Proof. Write $\hat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in [\varepsilon, a\varepsilon)) = \hat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in [\varepsilon, 1)) - \hat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in [a\varepsilon, 1))$. The result then follows from Proposition 3.6.

3.2 The proof of Theorems 1.5 and 1.6

This section is dedicated to the proof of Theorem 1.6, which implies the uniqueness of the MKS measure (Theorem 1.5) immediately. By scaling invariance of MKS measures, it suffices to prove Theorem 1.6 for simply connected domains U, D such that $\{0\} \subset U \subset D \subset \mathbb{D}$. Therefore, we will assume $D \subset \mathbb{D}$ throughout this section.

We first define the following important quantity I(D) whenever the limit exists (which is the case as we will see immediately)

$$I(D) := \lim_{\varepsilon \to 0} \left(\widehat{\mu}^0_{\mathbb{D}}[\ell \not\subset \varepsilon \mathbb{D}] - \mu^0_{\mathbb{C}}[\ell \not\subset \varepsilon \mathbb{D}, \ell \subset D] \right).$$
(3.10)

Note in particular that $I(\mathbb{D}) = \zeta_1$ according to Proposition 3.6. We will show that I(D) has a nice loop-soup expression (see Proposition 3.9), which is the key to Theorem 1.6. Before going any further, we first show that the limit in (3.10) exists by establishing an equality between I(D) and the $\mu_{\mathbb{C}}^0$ -mass of loops in \mathbb{D} that are not fully in D. **Lemma 3.8.** The limit in (3.10) exists and $I(D) = \mu_{\mathbb{C}}^0[\ell \not\subset D, \ell \subset \mathbb{D}] + \zeta_1 \in (0, \infty).$

Proof. Since $D \subset \mathbb{D}$, we have

$$\begin{split} \mu^{0}_{\mathbb{C}}[\ell \not\subset D, \ell \subset \mathbb{D}] &= \lim_{\varepsilon \to 0} \left(\mu^{0}_{\mathbb{C}}[\ell \not\subset \varepsilon \mathbb{D}, \ell \subset \mathbb{D}] - \mu^{0}_{\mathbb{C}}[\ell \not\subset \varepsilon \mathbb{D}, \ell \subset D] \right) \\ &= \lim_{\varepsilon \to 0} \left(\widehat{\mu}^{0}_{\mathbb{D}}[\ell \not\subset \varepsilon \mathbb{D}] - \zeta_{1} - \mu^{0}_{\mathbb{C}}[\ell \not\subset \varepsilon \mathbb{D}, \ell \subset D] \right), \end{split}$$

where the second line is from Proposition 3.6. Hence $I(D) = \mu_{\mathbb{C}}^0[\ell \not\subset D, \ell \subset \mathbb{D}] + \zeta_1$. Note that $\mu_{\mathbb{C}}^0[\ell \not\subset D, \ell \subset \mathbb{D}] \in (0, \infty)$ thanks to the non-triviality of $\mu_{\mathbb{C}}$.

The following proposition provides a clean form of I(D) that only involves randomness from the loop soup.

Proposition 3.9. We have $I(D) = \lambda \mathbf{E} \log f'_{D,\infty}(0)$. In particular, $\zeta_1 = I(\mathbb{D}) = \lambda \mathbf{E} \log f'_{\mathbb{D},\infty}(0)$.

We first show how to derive Theorem 1.6 (and hence Theorem 1.5) from Proposition 3.9.

Proof of Theorem 1.6, assuming Proposition 3.9. Write $\mu^0_{\mathbb{C}}[\ell \not\subset U, \ell \subset D] = \mu^0_{\mathbb{C}}[\ell \not\subset U, \ell \subset \mathbb{D}] - \mu^0_{\mathbb{C}}[\ell \not\subset D, \ell \subset \mathbb{D}]$. We finish the proof by applying Lemma 3.8 and Proposition 3.9.

Proof of Theorem 1.5. From Theorem 1.6, we have determined $\mu^0_{\mathbb{C}}[\ell \not\subset U, \ell \subset D]$ for all simply connected domains U and D satisfying $U \subset D$. This indeed characterizes $\mu_{\mathbb{C}}$ by a rather straightforward measure-theoretical argument, as presented in [Wer08, Page 146].

In the remaining of this section, we aim to show Proposition 3.9. Our proof relies crucially on the following loop-soup interpolation of I(D).

Lemma 3.10. We have

$$I(D) = \lim_{\varepsilon \to 0} \mathbf{E} \otimes \widehat{\mu}^0_{\mathbb{D}} [\ell \subset f_{D,\infty}(\varepsilon \mathbb{D}) \setminus (\varepsilon \mathbb{D})].$$
(3.11)

Proof. Since $D_{\infty} \subset \mathbb{D}$, by Schwarz lemma, we have $f_{D,\infty}(\varepsilon \mathbb{D}) \supset \varepsilon \mathbb{D}$. Then by Proposition 3.2,

$$\begin{aligned} \widehat{\mu}^{0}_{\mathbb{D}}[\ell \not\subset \varepsilon \mathbb{D}] - \mu^{0}_{\mathbb{C}}[\ell \not\subset \varepsilon \mathbb{D}, \ell \subset D] &= \mathbf{E} \otimes \widehat{\mu}^{0}_{\mathbb{D}}[\ell \not\subset \varepsilon \mathbb{D}] - \mathbf{E} \otimes \widehat{\mu}^{0}_{\mathbb{D}}[\ell \not\subset f_{D,\infty}(\varepsilon \mathbb{D})] \\ &= \mathbf{E} \otimes \widehat{\mu}^{0}_{\mathbb{D}}[\ell \subset f_{D,\infty}(\varepsilon \mathbb{D}) \setminus (\varepsilon \mathbb{D})]. \end{aligned}$$

Taking $\varepsilon \to 0$, we get (3.11).

We will use the following technical lemma to control the distortion of conformal maps.

Lemma 3.11. Let D be a simply connected domain in \mathbb{D} that contains 0. Let f be the conformal map from $D \to \mathbb{D}$ with f(0) = 0 and f'(0) > 0. Suppose $\varepsilon, \beta \in (0, 1), \varepsilon^{1-\beta} \leq \frac{1}{16}$ and $f'(0) < \varepsilon^{-\beta}$. There is a universal constant C such that for any z with $|z| = \varepsilon$,

$$f'(0)\varepsilon(1-C\varepsilon^{1-\beta}) \le |f(z)| \le f'(0)\varepsilon(1+C\varepsilon^{1-\beta}).$$

Proof. By Koebe's 1/4 theorem, we have $r := \operatorname{dist}(0, \partial D) \ge \frac{1}{4}f'(0)^{-1} > \frac{1}{4}\varepsilon^{\beta}$. Consider the conformal map g(w) := f(rw)/(rf'(0)) on \mathbb{D} , which satisfies g'(0) = 1. By [Law05, Proposition 3.26], for any $|w| = \frac{\varepsilon}{r} \le 4\varepsilon^{1-\beta} \le \frac{1}{4}$, we have $|g(w) - w| \le C |w|^2$ for some universal constant C. Setting z = rw, we have $|f(z) - f'(0)z| \le Cf'(0)\varepsilon^{2-\beta}$, concluding the proof.

Lemma 3.12. For any $\alpha_0 \in (0, \frac{1}{2}]$, we have

$$\lim_{\varepsilon \to 0} \mathbf{E} \otimes \widehat{\mu}^0_{\mathbb{D}} [\ell \subset f_{D,\infty}(\varepsilon \mathbb{D}) \setminus (\varepsilon \mathbb{D}); \ f'_{D,\infty}(0) < \varepsilon^{-\alpha_0}] = \lambda \mathbf{E} \log f'_{D,\infty}(0).$$
(3.12)

Therefore,

$$\lambda \mathbf{E} \log f'_{D,\infty}(0) \le I(D) < \infty. \tag{3.13}$$

Proof. By Lemma 3.11, we have

$$\begin{aligned} \left| \mathbf{E} \otimes \widehat{\mu}_{\mathbb{D}}^{0} [\ell \subset f_{D,\infty}(\varepsilon \mathbb{D}) \setminus (\varepsilon \mathbb{D}); \ f_{D,\infty}'(0) < \varepsilon^{-\alpha_{0}}] - \mathbf{E} \otimes \widehat{\mu}_{\mathbb{D}}^{0} [\|\ell\|_{\infty} \in [\varepsilon, f_{D,\infty}'(0)\varepsilon); \ f_{D,\infty}'(0) < \varepsilon^{-\alpha_{0}}] \\ \leq \mathbf{E} \otimes \widehat{\mu}_{\mathbb{D}}^{0} [\|\ell\|_{\infty} \in [f_{D,\infty}'(0)\varepsilon(1 - C\varepsilon^{1-\alpha_{0}}), f_{D,\infty}'(0)\varepsilon(1 + C\varepsilon^{1-\alpha_{0}}); \ f_{D,\infty}'(0) < \varepsilon^{-\alpha_{0}}] \\ \leq O(\varepsilon^{1-\alpha_{0}}) \mathbf{P}(f_{D,\infty}'(0) < \varepsilon^{-\alpha_{0}}) \to 0 \quad \text{as} \quad \varepsilon \to 0, \end{aligned}$$

where we used Corollary 3.7 to estimate the $\hat{\mu}^0_{\mathbb{D}}$ -mass of ℓ in the last inequality. It follows that

$$\begin{split} &\lim_{\varepsilon \to 0} \mathbf{E} \otimes \widehat{\mu}_{\mathbb{D}}^{0} [\|\ell\|_{\infty} \in [\varepsilon, f_{D,\infty}'(0)\varepsilon); \ f_{D,\infty}'(0) < \varepsilon^{-\alpha_{0}}] \\ &= \lim_{\varepsilon \to 0} \mathbf{E}[\lambda \log f_{D,\infty}'(0) + o(1); \ f_{D,\infty}'(0) < \varepsilon^{-\alpha_{0}}] \\ &= \lambda \mathbf{E} \log f_{D,\infty}'(0) \end{split}$$
(by Corollary 3.7)
(by Corollary 3.7)

Combined, we complete the proof of (3.12). Now combining (3.12) and Lemma 3.10, we obtain that

$$\lambda \mathbf{E} \log f'_{D,\infty}(0) \le \lim_{\varepsilon \to 0} \mathbf{E} \otimes \widehat{\mu}^0_{\mathbb{D}}(\ell \subset f_{D,\infty}(\varepsilon \mathbb{D}) \setminus (\varepsilon \mathbb{D})) = I(D)$$

By Lemma 3.8, $I(D) < \infty$, and (3.13) then follows.

Finally, we turn to

Proof of Proposition 3.9. By (3.12) and Lemma 3.10, it remains to show that

$$\lim_{\varepsilon \to 0} \mathbf{E} \otimes \widehat{\mu}^0_{\mathbb{D}}[\ell \subset f_{D,\infty}(\varepsilon \mathbb{D}) \setminus (\varepsilon \mathbb{D}); \ f'_{D,\infty}(0) \ge \varepsilon^{-\alpha_0}] = 0.$$

First, by (3.13), as $\varepsilon \to 0$,

$$|\log \varepsilon| \mathbf{P}[f'_{D,\infty}(0) \ge \varepsilon^{-\alpha_0}] \le \frac{1}{\alpha_0} \mathbf{E}[\log f'_{D,\infty}(0), f'_{D,\infty}(0) > \varepsilon^{-\alpha_0}] \to 0.$$

Therefore, by Proposition 3.6, we have

$$\mathbf{E} \otimes \widehat{\mu}^{0}_{\mathbb{D}} [\ell \subset f_{D,\infty}(\varepsilon \mathbb{D}) \setminus (\varepsilon \mathbb{D}); \ f'_{D,\infty}(0) \ge \varepsilon^{-\alpha_{0}}] \le \mathbf{E} \otimes \widehat{\mu}^{0}_{\mathbb{D}} [\|\ell\|_{\infty} \in [\varepsilon, 1); \ f'_{D,\infty}(0) \ge \varepsilon^{-\alpha_{0}}]$$
$$= (\lambda |\log \varepsilon| + \zeta_{1} + o(1)) \mathbf{P} [f'_{D,\infty}(0) \ge \varepsilon^{-\alpha_{0}}],$$

which tends to 0 as $\varepsilon \to 0$. The proof is now complete.

4 Electrical thickness

This section is to establish Theorem 1.9 from Theorem 1.6. We inherit the notation from Section 3. We first show that the $\mu_{\mathbb{C}}^{0}$ -mass of loops that intersect the unit circle \mathbb{S}^{1} is finite.

Lemma 4.1. For any $c \in (0,1]$, we have $\zeta_2 := \mu^0_{\mathbb{C}}(\ell \cap \mathbb{S}^1 \neq \emptyset) < \infty$.

Proof. According to the uniqueness of the MKS measure (Theorem 1.5), it suffices to prove the statement for $\mu_{\mathbb{C}} = \operatorname{SLE}_{\kappa}^{\operatorname{loop}}$, the counting measure on the full-plane $\operatorname{CLE}_{\kappa}$ configuration Γ . First, there is a universal u > 0 such that for any simply connected domain D containing 0 with $\operatorname{dist}(0, \partial D) < 1$, with probability at least u, the outermost $\operatorname{CLE}_{\kappa}$ loop in D surrounding 0 is contained in $\frac{1}{2}\mathbb{D}$ (this is directly from the loop-soup construction of $\operatorname{CLE}_{\kappa}$ [SW12]; see e.g. [KW16, Section 3.1, Fact 4]). Then by the Markov property of $\operatorname{CLE}_{\kappa}$, the number of loops in Γ intersecting \mathbb{S}^1 is stochastically dominated by a geometric distribution with parameter u, i.e., $\mu_{\mathbb{C}}^0(\ell \cap \mathbb{S}^1 \neq \emptyset) \leq \sum_{n>1} nu(1-u)^{n-1} < \infty$.

Denote $A_{\varepsilon,R} := \{z : \varepsilon < |z| < R\}$. The following gives the $\mu^0_{\mathbb{C}}$ -mass of loops that intersect $A_{\varepsilon,R}$.

Lemma 4.2. For any $c \in (0, 1]$, let

$$\zeta := \zeta_2 - 2\zeta_1 = \mu^0_{\mathbb{C}}(\ell \cap \mathbb{S}^1 \neq \emptyset) - 2\lambda \mathbf{E} \log f'_{\mathbb{D},\infty}(0) \in (-\infty,\infty),$$
(4.1)

where ζ_1 and ζ_2 are defined in Proposition 3.6 and Lemma 4.1 respectively. Then, we have

$$\lim_{\varepsilon \to 0, R \to \infty} \left(\mu^0_{\mathbb{C}}(\ell \cap \mathbb{A}_{\varepsilon, R} \neq \emptyset) - \widehat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in (\varepsilon, 1)) - \widehat{\mu}^0_{\mathbb{D}}(\|\ell\|_{\infty} \in (R^{-1}, 1)) \right) = \zeta.$$
(4.2)

Proof. The last equation of (4.1) follows from Proposition 3.9. To show (4.2), write $\|\ell\|_{\min} := \min_{z \in \ell} |z|$, and we decompose the mass of loops as

$$\mu^{0}_{\mathbb{C}}(\ell \cap \mathbb{A}_{\varepsilon,R} \neq \emptyset) = \mu^{0}_{\mathbb{C}}(\|\ell\|_{\infty} \in (\varepsilon, 1)) + \mu^{0}_{\mathbb{C}}(\|\ell\|_{\min} \in (1, R)) + \mu^{0}_{\mathbb{C}}(\ell \cap \mathbb{S}^{1} \neq \emptyset).$$
(4.3)

Note that $\mu^0_{\mathbb{C}}(\|\ell\|_{\min} \in (1, R)) = \mu^0_{\mathbb{C}}(\|\ell\|_{\infty} \in (R^{-1}, 1))$ by inversion invariance. Using Proposition 3.6 to estimate the first two terms on the right-hand side of (4.3), we finish the proof.

Next, we consider general simply connected domain $D \subset \mathbb{C}$ containing 0 (without the assumption that $D \subset \mathbb{D}$). The definition of I(D) given in (3.10) can be extended to this general case straightforwardly. We can generalize the result in Proposition 3.9 to $D \subset \mathbb{C}$ by scaling.

Proposition 4.3. For any simply connected domain $D \subset \mathbb{C}$ containing 0, we have

$$I(D) = \lambda \mathbf{E} \log f'_{D,\infty}(0). \tag{4.4}$$

Proof. Let $a := \sup_{z \in D} |z|$ and $\widetilde{D} := a^{-1}D$. By (3.10) and scaling invariance of $\mu_{\mathbb{C}}^0$, we have $I(D) = I(\widetilde{D}) - \lambda \log a$. Further, since $\widetilde{D} \subset \mathbb{D}$, Proposition 3.9 gives $I(\widetilde{D}) = \lambda \mathbf{E} \log f'_{\widetilde{D},\infty}(0)$. Moreover, by scaling invariance of the whole-plane SLE_{8/3} loop-soup, $f'_{D,\infty}(0)$ has the same law as $a^{-1}f'_{\widetilde{D},\infty}(0)$. Combined, we obtain $I(D) = \lambda \mathbf{E} \log f'_{\widetilde{D},\infty}(0) - \lambda \log a = \lambda \mathbf{E} \log f'_{D,\infty}(0)$, as desired.

Now we complete the proof of Theorem 1.9. Let V be any connected compact set separating 0 and ∞ , and recall the notation introduced above Theorem 1.9. For clarity, we fix the conformal maps $f_{V_{\infty}}$ and $h_{V_{\infty}}$ such that $f'_{V_{\infty}}(0), h'_{V_{\infty}}(\infty) > 0$.

Proof of Theorem 1.9. By Proposition 4.3 and inversion invariance, we have

$$I(\Omega_{V,\infty}) = \lim_{\varepsilon \to 0} \left(\widehat{\mu}^0_{\mathbb{D}}(\ell \not\subset \varepsilon \mathbb{D}) - \mu^0_{\mathbb{C}}(\ell \not\subset \varepsilon \mathbb{D}, \ell \subset \Omega_{V,\infty}) \right) = \lambda \mathbf{E} \log f'_{V_{\infty}}(0), \tag{4.5}$$

$$\lim_{R \to \infty} \left(\widehat{\mu}^0_{\mathbb{D}}(\ell \not\subset R^{-1}\mathbb{D}) - \mu^0_{\mathbb{C}}(\ell \not\subset R\mathbb{D}^*, \ell \subset \Omega^*_{V,\infty}) \right) = -\lambda \mathbf{E} \log h'_{V_{\infty}}(\infty).$$
(4.6)

Moreover, for any $0 < \varepsilon < R < \infty$ such that $V \subset \mathbb{A}_{\varepsilon,R}$, we have

$$\begin{split} \mu^{0}_{\mathbb{C}}(\ell \cap V \neq \emptyset) &= \mu^{0}_{\mathbb{C}}(\ell \cap \mathbb{A}_{\varepsilon,R} \neq \emptyset) - \mu^{0}_{\mathbb{C}}(\ell \not\subset \varepsilon \mathbb{D}, \ell \subset \Omega_{V,\infty}) - \mu^{0}_{\mathbb{C}}(\ell \not\subset R \mathbb{D}^{*}, \ell \subset \Omega^{*}_{V,\infty}) \\ &= \left(\widehat{\mu}^{0}_{\mathbb{D}}(\ell \not\subset \varepsilon \mathbb{D}) - \mu^{0}_{\mathbb{C}}(\ell \not\subset \varepsilon \mathbb{D}, \ell \subset \Omega_{V,\infty})\right) + \left(\widehat{\mu}^{0}_{\mathbb{D}}(\ell \not\subset R^{-1}\mathbb{D}) - \mu^{0}_{\mathbb{C}}(\ell \not\subset R \mathbb{D}^{*}, \ell \subset \Omega^{*}_{V,\infty})\right) \\ &+ \left(\mu^{0}_{\mathbb{C}}(\ell \cap \mathbb{A}_{\varepsilon,R} \neq \emptyset) - \widehat{\mu}^{0}_{\mathbb{D}}(\ell \not\subset \varepsilon \mathbb{D}) - \widehat{\mu}^{0}_{\mathbb{D}}(\ell \not\subset R^{-1}\mathbb{D})\right). \end{split}$$

Taking $\varepsilon \to 0$ and $R \to \infty$, using (4.5), (4.6) and Lemma 4.2 to deal with the three terms on the right-hand side above respectively, we conclude that

$$\mu^0_{\mathbb{C}}(\ell \cap V \neq \emptyset) = \lambda \mathbf{E} \log f'_{V_{\infty}}(0) - \lambda \mathbf{E} \log h'_{V_{\infty}}(\infty) + \zeta = \lambda \mathbf{E} \left[\theta(V_{\infty})\right] + \zeta,$$

which finishes the proof.

Remark 4.4. Theorem 1.9 can be straightforwardly extended to the case c = 0 (i.e. $\kappa = \frac{8}{3}$). Now $\zeta = \mu_{\mathbb{C},c=0}^{0}(\ell \cap \mathbb{S}^{1} \neq \emptyset)$ by (4.1), and V_{∞} is simply V. Namely,

$$\mu^{0}_{\mathbb{C},c=0}(\ell \cap V \neq \emptyset) = \lambda \theta(V) + \mu^{0}_{\mathbb{C},c=0}(\ell \cap \mathbb{S}^{1} \neq \emptyset),$$
(4.7)

where $\theta(V)$ is the electrical thickness of V. Note that (4.7) looks similar to [WX24, Corollary 4.8], which computes the total mass of Brownian loops with winding $m \in \mathbb{Z}_+$ around 0 and hitting V.

As mentioned before, if $\mu_{\mathbb{C},c=0}$ is Werner's measure \mathcal{W} , then $\lambda = \frac{\pi}{5}$ [Wer08]. One can also determine that $\mathcal{W}^0(\ell \cap \mathbb{S}^1 \neq \emptyset) = \mathcal{W}(\ell \text{ surrounds } 0, \ \ell \cap \mathbb{S}^1 \neq \emptyset) = \frac{\sqrt{6}}{15}$.²

5 Discussions and Possible Extensions

At the end of this paper we briefly list several remarks and open questions.

General Riemann Surfaces. Both the Brownian loop measure and the SLE loop measure can be defined on a general Riemann surface Σ , see e.g. [Wer08, Law11, Zha21]. It is interesting to see how our approach in Section 3 can be extended to the SLE loop measure μ_{Σ} on Σ . For instance, one can

²Let $Z_{8/3}(\tau)$ be the total mass of \mathcal{W} on non-contractible loops in $\{z : e^{-2\pi\tau} < |z| < 1\}$. Then $\mathcal{W}(\ell$ surrounds 0, $\ell \cap \mathbb{S}^1 \neq \emptyset$) = $\lim_{\tau \to \infty} Z_{8/3}(\tau) - 2Z_{8/3}(\tau)$. The explicit formula of $Z_{8/3}(\tau)$, up to a multiplicative constant, is given by Cardy $Z_{\text{Cardy}}(\tau)$ [Car06, (5)] and proved by [ARS22]. By a forthcoming work of Liu, Qian, Sun, Wu and the first author of this paper, this multiplicative constant can be further specified, i.e., $Z_{8/3}(\tau) = \frac{3\sqrt{2}}{5}Z_{\text{Cardy}}(\tau)$.

consider the SLE_{8/3} loop soup on Σ , and relate it to the quantity $\mu_{\Sigma}[\ell \subset A \text{ and non-contractible in } A]$ for some annular region $A \subset \Sigma$.

Note that recently Wang and Xue [WX24] demonstrated that there is a deep connection between the Brownian loop measure and the length spectrum of geodesics on Σ . Hence, it is natural to ask that on Σ how the SLE loop measure relates to such geometric quantities.

The regime c < 0 or $\kappa \in (0, \frac{8}{3})$. Our approach relies on the interpretation of the loop-mass term in the generalized conformal restriction through the Brownian loop soup, which indeed utilizes the coupling from $SLE_{8/3}$ to SLE_{κ} for $\kappa > \frac{8}{3}$. It would be interesting to see whether a similar approach can be adapted to the regime c < 0. This is connected to an alternative characterization of SLE_{κ} for $\kappa < \frac{8}{3}$ as the unique simple curve with the property that, when augmented with a specific density of Brownian loops, the resulting hull coincides with the union of certain Brownian motions [Wer03].

Integrability of the SLE loop. Our results, Theorems 1.6 and 1.9, establish relationships between natural quantities of the SLE_{κ} loop measure and the $SLE_{8/3}$ loop soup. A natural follow-up question is whether these quantities can be expressed as *exact functions*, and whether further integrability for the SLE_{κ} loop measure can be obtained. To date, only a few results on the integrability of the SLE_{κ} loop measure are known, including its two- and three-point Green functions and its electrical thickness [ACSW24a, ACSW24b], by Ang, Sun, Wu and the first author of this paper.

Acknowledgement. G.C. thanks X. Sun and B. Wu for helpful comments and discussions. Y.G. thanks W. Qian for bringing his attention to this problem and helpful discussions. We thank W. Werner for pointing out the relevant work [Wer03] and his remarks on the applicability of our results to the reversibility and duality properties of SLE. G.C. is supported by National Key R&D Program of China (No. 2021YFA1002700). Y.G. is supported by National Key R&D Program of China (No. 2023YFA1010700).

References

- [ACSW24a] Morris Ang, Gefei Cai, Xin Sun, and Baojun Wu. Integrability of Conformal Loop Ensemble: Imaginary DOZZ Formula and Beyond. arXiv preprint arXiv:2107.01788, 2024.
- [ACSW24b] Morris Ang, Gefei Cai, Xin Sun, and Baojun Wu. SLE Loop Measure and Liouville Quantum Gravity. arXiv preprint arXiv:2409.16547, 2024.
- [AM01] Hélène Airault and Paul Malliavin. Unitarizing probability measures for representations of Virasoro algebra.
 J. Math. Pures Appl. (9), 80(6):627–667, 2001.
- [ARS22] Morris Ang, Guillaume Remy, and Xin Sun. The moduli of annuli in random conformal geometry. arXiv preprint arXiv:2203.12398, 2022.
- [BD16] Stéphane Benoist and Julien Dubédat. An SLE₂ loop measure. Ann. Inst. H. Poincaré Probab. Statist., 52(3):1406–1436, 2016.
- [BJ24] Guillaume Baverez and Antoine Jego. The CFT of SLE loop measures and the Kontsevich–Suhov conjecture. arXiv preprint arXiv:2407.09080, 2024.
- [Car06] John Cardy. The O(n) model on the annulus. J. Stat. Phys., 125(1):1–21, 2006.

- [CW24] Marco Carfagnini and Yilin Wang. Onsager-Machlup functional for SLE_{κ} loop measures. Comm. Math. Phys., 405(11):258, 2024.
- [Dub05] Julien Dubédat. $SLE(\kappa, \rho)$ martingales and duality. Ann. Probab., 33(1):223–243, 2005.
- [Dub09] Julien Dubédat. Duality of Schramm-Loewner evolutions. Ann. Sci. Éc. Norm. Supér. (4), 42(5):697–724, 2009.
- [FL13] Laurence S. Field and Gregory F. Lawler. Reversed radial SLE and the Brownian loop measure. J. Stat. Phys., 150(6):1030–1062, 2013.
- [GLQ22] Yifan Gao, Xinyi Li, and Wei Qian. Multiple points on the boundaries of Brownian loop-soup clusters. arXiv preprint arXiv:2205.11468, 2022.
- [GQW24] Maria Gordina, Wei Qian, and Yilin Wang. Infinitesimal conformal restriction and unitarizing measures for Virasoro algebra. arXiv preprint arXiv:2407.09426, 2024.
- [KS07] Maxim Kontsevich and Yuri Suhov. On Malliavin measures, SLE, and CFT. Tr. Mat. Inst. Steklova, 258:107–153, 2007.
- [KW04] Richard W. Kenyon and David B. Wilson. Conformal radii of loop models. 2004.
- [KW16] Antti Kemppainen and Wendelin Werner. The nested simple conformal loop ensembles in the Riemann sphere. *Probab. Theory Related Fields*, 165(3-4):835–866, 2016.
- [Law05] Gregory F. Lawler. Conformally invariant processes in the plane, volume 114 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2005.
- [Law09] Gregory F. Lawler. Partition functions, loop measure, and versions of SLE. J. Stat. Phys., 134(5-6):813– 837, 2009.
- [Law11] Gregory F. Lawler. Defining SLE in multiply connected domains with the Brownian loop measure. arXiv preprint arXiv:1108.4364, 2011.
- [LSW01a] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Values of Brownian intersection exponents. I. Half-plane exponents. Acta Math., 187(2):237–273, 2001.
- [LSW01b] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Values of Brownian intersection exponents. II. Plane exponents. Acta Math., 187(2):275–308, 2001.
- [LSW02a] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Analyticity of intersection exponents for planar Brownian motion. Acta Math., 189(2):179–201, 2002.
- [LSW02b] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Values of Brownian intersection exponents. III. Two-sided exponents. Ann. Inst. H. Poincaré Probab. Statist., 38(1):109–123, 2002.
- [LSW03] Gregory Lawler, Oded Schramm, and Wendelin Werner. Conformal restriction: the chordal case. J. Amer. Math. Soc., 16(4):917–955, 2003.
- [LW00] Gregory F. Lawler and Wendelin Werner. Universality for conformally invariant intersection exponents. J. Eur. Math. Soc., 2(4):291–328, 2000.
- [LW04] Gregory F. Lawler and Wendelin Werner. The Brownian loop soup. *Probab. Theory Related Fields*, 128(4):565–588, 2004.
- [LY21] Gregory F. Lawler and Stephen Yearwood. A new proof of reversibility of SLE_{κ} for $\kappa \leq 4$. arXiv preprint arXiv:2111.06960, 2021.
- [MS16a] Jason Miller and Scott Sheffield. Imaginary geometry I: interacting SLEs. Probab. Theory Related Fields, 164(3-4):553–705, 2016.
- [MS16b] Jason Miller and Scott Sheffield. Imaginary geometry II: reversibility of $SLE_{\kappa}(\rho_1; \rho_2)$ for $\kappa \in (0, 4)$. Ann. Probab., 44(3):1647–1722, 2016.

- [MS16c] Jason Miller and Scott Sheffield. Imaginary geometry III: reversibility of SLE_{κ} for $\kappa \in (4, 8)$. Ann. of Math. (2), 184(2):455–486, 2016.
- [NW11] Şerban Nacu and Wendelin Werner. Random soups, carpets and fractal dimensions. J. Lond. Math. Soc., 83(3):789–809, 2011.
- [Qia18] Wei Qian. Conformal restriction: the trichordal case. *Probab. Theory Related Fields*, 171(3-4):709–774, 2018.
- [Qia21] Wei Qian. Generalized disconnection exponents. Probab. Theory Related Fields, 179(1-2):117–164, 2021.
- [Sch00] Oded Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118:221–288, 2000.
- [SSW09] Oded Schramm, Scott Sheffield, and David B. Wilson. Conformal radii for conformal loop ensembles. Comm. Math. Phys., 288(1):43–53, 2009.
- [SW12] Scott Sheffield and Wendelin Werner. Conformal loop ensembles: the Markovian characterization and the loop-soup construction. Ann. of Math. (2), 176(3):1827–1917, 2012.
- [vdBCL16] Tim van de Brug, Federico Camia, and Marcin Lis. Random walk loop soups and conformal loop ensembles. Probab. Theory Related Fields, 166(1-2):553–584, 2016.
- [Wer03] Wendelin Werner. SLEs as boundaries of clusters of Brownian loops. C. R. Math. Acad. Sci. Paris, 337(7):481–486, 2003.
- [Wer05] Wendelin Werner. Conformal restriction and related questions. Probab. Surv., 2:145–190, 2005.
- [Wer08] Wendelin Werner. The conformally invariant measure on self-avoiding loops. J. Amer. Math. Soc., 21(1):137–169, 2008.
- [Wu15] Hao Wu. Conformal restriction: the radial case. *Stochastic Process. Appl.*, 125(2):552–570, 2015.
- [WW13] Wendelin Werner and Hao Wu. From $CLE(\kappa)$ to $SLE(\kappa, \rho)$'s. Electron. J. Probab., 18(36):1–20, 2013.
- [WX24] Yilin Wang and Yuhao Xue. The Brownian loop measure on Riemann surfaces and applications to length spectra. *arXiv preprint arXiv:2406.09108*, 2024.
- [Zha08a] Dapeng Zhan. Duality of chordal SLE. Invent. Math., 174(2):309–353, 2008.
- [Zha08b] Dapeng Zhan. Reversibility of chordal SLE. Ann. Probab., 36(4):1472–1494, 2008.
- [Zha21] Dapeng Zhan. SLE loop measures. Probab. Theory Related Fields, 179(1-2):345–406, 2021.