

Anytime-valid FDR control with the stopped e-BH procedure

Hongjian Wang¹, Sanjit Dandapanthula¹, and Aaditya Ramdas²

¹, ²Department of Statistics and Data Science, Carnegie Mellon University

²Machine Learning Department, Carnegie Mellon University
 {hjnwang, sanjitd, aramdas}@cmu.edu

February 13, 2025

Abstract

The recent e-Benjamini-Hochberg (e-BH) procedure for multiple hypothesis testing is known to control the false discovery rate (FDR) under arbitrary dependence between the input e-values. This paper points out an important subtlety when applying the e-BH procedure with e-processes, which are sequential generalizations of e-values (where the data are observed sequentially). Since adaptively stopped e-processes are e-values, the e-BH procedure can be repeatedly applied at every time step, and one can continuously monitor the e-processes and the rejection sets obtained. One would hope that the “stopped e-BH procedure” (se-BH) has an FDR guarantee for the rejection set obtained at any stopping time. However, while this is true if the data in different streams are independent, it is not true in full generality, because each stopped e-process is an e-value only for stopping times in its own *local* filtration, but the se-BH procedure employs a stopping time with respect to a *global* filtration. This can cause information to *leak across time*, allowing one stream to know its future by knowing past data of another stream. This paper formulates a simple causal condition under which *local e-processes* are also *global e-processes* and thus the se-BH procedure does indeed control the FDR. The condition excludes unobserved confounding from the past and is met under most reasonable scenarios including genomics.

1 Introduction

Controlling the false discovery rate (FDR) at a pre-specified level α is the central task in multiple hypothesis testing. In this paper, we consider the FDR control problem in sequential multiple hypothesis testing, where each time one observes a batch of data that corresponds to all or some hypotheses, a natural set-up in e.g. biostatistics where cell observations for the full genome are collected and used for testing sequentially.

The Benjamini-Hochberg (BH) procedure (Benjamini and Hochberg, 1995) is among the most widely used approaches to FDR control. However, the BH procedure operates directly on fixed-sample p-values, which are inherently non-sequential. Further, the BH procedure does not hold under arbitrary cross-hypothesis dependence, and there has been an on-going effort in deciding the exact dependence condition for BH.

Recently, Wang and Ramdas (2022) showed that, by applying the BH procedure to the inverses of *e-values*, the same FDR control can be extended to arbitrary dependence across hypotheses. Further, since e-values arise naturally in sequential experiments, this result which is commonly referred to as “e-BH” opens up the possibility of sequential multiple testing. The sequential eBH was later spelled out in detail by Xu et al. (2021), whose result allows bandit-like active queries into the hypotheses, each carrying an e-process (to be defined formally later), and the FDR is controlled at any *stopping time* under arbitrary dependence.

This seemingly very satisfactory result by [Xu et al. \(2021\)](#), however, bears a very crucial caveat. It is assumed that the e-processes of the hypotheses are valid under a shared “global” filtration. This indicates that one can *not* simply apply e-BH to e-processes each constructed “locally” *within* the hypotheses without further verification. As we shall elaborate soon, filtration is often an important matter in sequential statistics as using an e-process adapted to the “wrong” filtration defies the safety (type 1 error or FDR) guarantee.

In this paper, we first formally define the local-global distinction of e-processes, and identify some further conditions on which e-processes constructed within hypotheses using generic methods are not only local, but also global e-processes. We exemplify our conditions by a parametric and a nonparametric problem and also discuss when they do not hold. Additionally, we define the more generalized notion of global compound e-processes, and show that e-BH with these processes is necessary and sufficient to obtain stopped FDR control.

2 Background and problem set-up

Throughout the paper, we shall liberally adopt the notations $\mathbb{E}_P, \mathbb{E}_\theta$ etc. to denote the expected value of some random variable when the underlying data-generating distribution is P , or when the underlying parameter of interest takes the value of θ .

2.1 Multiple testing with e-values and e-BH

We first recall the standard definitions of e-values. Let \mathcal{P} be a set of distributions. A nonnegative random variable E is called an e-value for \mathcal{P} if $\mathbb{E}_P E \leq 1$ for all $P \in \mathcal{P}$. E-values have been used for safely testing the null hypothesis that the true distribution comes from \mathcal{P} by Markov’s inequality using the rejection rule that $E \geq 1/\alpha$ and are understood as the quantification of evidence against the null \mathcal{P} .

In the multiple testing set-up, we assume there is a set of all possible distributions \mathcal{M} of the model, with G subsets of null distributions $\mathcal{P}^1, \dots, \mathcal{P}^G \subseteq \mathcal{M}$. We denote by $[G]$ the set $\{1, \dots, G\}$. The underlying distribution $P \in \mathcal{M}$ may belong to arbitrary number of sets among $(\mathcal{P}^g : g \in [G])$. The g^{th} null hypothesis \mathcal{H}_0^g is true if $P \in \mathcal{P}^g$. Let us denote by $\text{eBH}_\alpha : \mathbb{R}_{\geq 0}^{[G]} \rightarrow 2^{[G]}$ the e-BH procedure at level α that takes G e-values and returns a set of rejected hypotheses ([Wang and Ramdas, 2022](#), Section 4):

$$\text{eBH}_\alpha(E_1, \dots, E_G) = \{g : E_g \geq E_{(g^*)}\} \quad \text{where } g^* = \max \left\{ g \in [G] : \frac{gE_{(g)}}{G} \geq \frac{1}{\alpha} \right\}, \quad (1)$$

and $(E_{(g)} : g \in [G])$ are $(E_g : g \in [G])$ sorted decreasingly. The FDR guarantee of e-BH on G e-values is proved by [Wang and Ramdas \(2022, Proposition 2\)](#), who also generalize it to G random variables satisfying a condition weaker than being e-values ([Wang and Ramdas, 2022, Proposition 3](#)). These were recently named *compound e-values* by [Ignatiadis et al. \(2024, Definition 1.1\)](#).

Definition 2.1 (Compound e-values). G random variables $\mathbf{E} = (E_g : g \in [G]) \in \mathbb{R}_{\geq 0}^G$ are called compound e-values for $(\mathcal{P}^g : g \in G)$ if,

$$\sup_{P \in \mathcal{M}} \sum_{g \in [G]} \mathbb{1}\{P \in \mathcal{P}^g\} \cdot \mathbb{E}_P E_g \leq G. \quad (2)$$

Clearly, if each E_g is an e-value for \mathcal{P}^g , that is, $\sup_{P \in \mathcal{P}^g} \mathbb{E}_P E_g \leq 1$, then \mathbf{E} are compound e-values. However compound e-values form a much larger class including weighted e-values. The e-BH on compound e-values controls the FDR due to [Ignatiadis et al. \(2024, Theorem 3.2\)](#).

Lemma 2.2 (FDR property of e-BH). *Let $\mathbf{E} = (E_g : g \in [G])$ be compound e-values for $(\mathcal{P}^g : g \in [G])$. Then, for any underlying $P \in \mathcal{M}$,*

$$\text{FDR} = \mathbb{E}_P \left[\frac{\sum_{g=1}^G \mathbb{1}\{P \in \mathcal{P}^g\} \cdot \mathbb{1}\{g \in \text{eBH}_\alpha(\mathbf{E})\}}{1 \vee \sum_{g=1}^G \mathbb{1}\{g \in \text{eBH}_\alpha(\mathbf{E})\}} \right] \leq \alpha. \quad (3)$$

It is worth noting that the converse to Lemma 2.2 is also true, due to Ignatiadis et al. (2024, Theorems 3.1 and 3.3). That is, a random set of rejected hypotheses controls the FDR if and only if it is the e-BH of G compound e-values. We shall revisit this equivalence after fully developing our theory in the sequential setting.

2.2 Sequential multiple testing and filtrations

We now introduce the sequential experiment setting and let us first review the terminology with a single hypothesis. Let $\{\mathcal{F}_n\}_{n \geq 0}$ be a filtration. An $\{\mathcal{F}_n\}$ -adapted nonnegative process $\{M_n\}_{n \geq 1}$ is called an e-process for \mathcal{P} on $\{\mathcal{F}_n\}$ if either of the following two equivalent conditions holds:

- For any $\{\mathcal{F}_n\}$ -stopping time τ , the stopped variable M_τ is an e-value for \mathcal{P} .
- For any $P \in \mathcal{P}$, there is a process $\{N_n^P\}$ such that $N_0^P = 1$, it is a nonnegative supermartingale under P , and $N_n^P \geq M_n$ for all n .

Therefore, one can stop at any $\{\mathcal{F}_n\}$ -stopping time τ and safely reject the null \mathcal{P} if the stopped e-value $M_\tau \geq 1/\alpha$. In particular, τ can be $\min\{n : M_n \geq 1/\alpha\}$.

With Lemma 2.2, it is tempting to consider the following sequential multiple testing scheme with e-processes: since “the stopped value of an e-process is an e-value”, if one runs an e-process for each hypothesis and computes e-BH with the stopped values of e-processes, does one achieve the sequential e-BH control at any stopping time? Here, however, lies the subtlety of the problem: filtration. Let us carefully define the sequential problem with multiple hypotheses.

We assume G null sets of distributions $\mathcal{P}^1, \dots, \mathcal{P}^G \subseteq \mathcal{M}$ as before, and that the g^{th} null \mathcal{H}_0^g states that the underlying P belongs to \mathcal{P}^g . Further, for each $g \in [G]$, there is a *local* filtration $\{\mathcal{F}_n^g\}_{n \geq 0}$ for the hypothesis $\mathcal{H}^g : P \in \mathcal{P}^g$. This represents the evolution of information gathered from testing \mathcal{H}_0^g . The combination of these G local filtrations gives rise to the *global* filtration $\{\mathcal{F}_n\}_{n \geq 0}$:

$$\mathcal{F}_n = \sigma(\mathcal{F}_n^g : g \in [G]), \quad (4)$$

representing all information gathered up to n observations from all hypotheses. The following data generating model that we shall later pay closer attention to may illustrate this local-global distinction. Suppose at time n , the n^{th} observation gathered includes a common variable X_n and an array of hypothesis-specific variables $(Y_n^g : g \in [G])$. Then \mathcal{F}_n^g is defined as $\sigma(X_i, Y_i^g : i \leq n)$ whereas \mathcal{F}_n as $\sigma(X_i, Y_i^g : i \leq n, g \in [G])$. We shall see an important variation on this choice of filtrations later. Our definitions apply to more general cases where the data sharing can happen between arbitrary subsets of hypotheses and can change over time.

The concept of local and global filtrations leads to that of the stopped FDR control which is only defined globally, and the local and global e-processes. Below, we denote the set of all stopping times on $\{\mathcal{F}_n^g\}$ by \mathbb{T}^g , and on $\{\mathcal{F}_n\}$ by \mathbb{T} .

Definition 2.3. *A $2^{[K]}$ -valued set process $\{R_n\}$ adapted to $\{\mathcal{F}_n\}$ is said to satisfy the level- α stopped FDR control if*

$$\sup_{\substack{P \in \mathcal{M} \\ \tau \in \mathbb{T}}} \mathbb{E}_P \left[\frac{\sum_{g=1}^G \mathbb{1}\{P \in \mathcal{P}^g\} \cdot \mathbb{1}\{g \in R_\tau\}}{1 \vee \sum_{g=1}^G \mathbb{1}\{g \in R_\tau\}} \right] \leq \alpha. \quad (5)$$

Definition 2.4. Let $(\{M_n^g\} : g \in [G])$ be G nonnegative processes. They are called:

1. local e-processes for $(\mathcal{P}^g : g \in [G])$, if for any $g \in [G]$, $\{M_n^g\}$ is adapted to $\{\mathcal{F}_n^g\}$ and

$$\sup_{\substack{P \in \mathcal{P}^g \\ \tau \in \mathbb{T}^g}} \mathbb{E}_P M_\tau^g \leq 1; \quad (6)$$

2. local compound e-processes for $(\mathcal{P}^g : g \in [G])$, if for any $g \in [G]$, $\{M_n^g\}$ is adapted to $\{\mathcal{F}_n^g\}$ and

$$\sup_{\substack{P \in \mathcal{M} \\ \tau^1 \in \mathbb{T}^1 \\ \vdots \\ \tau^G \in \mathbb{T}^G}} \left(\sum_{g \in [G]} \mathbb{1}\{P \in \mathcal{P}^g\} \cdot \mathbb{E}_{P^g} M_{\tau^g}^g \right) \leq G; \quad (7)$$

3. global e-processes for $(\mathcal{P}^g : g \in [G])$, if for any $g \in [G]$, $\{M_n^g\}$ is adapted to $\{\mathcal{F}_n\}$ and

$$\sup_{\substack{P \in \mathcal{P}^g \\ \tau \in \mathbb{T}}} \mathbb{E}_P M_\tau^g \leq 1; \quad (8)$$

4. global compound e-processes for $(\mathcal{P}^g : g \in [G])$, if for any $g \in [G]$, $\{M_n^g\}$ is adapted to $\{\mathcal{F}_n\}$ and

$$\sup_{\substack{P \in \mathcal{M} \\ \tau \in \mathbb{T}}} \left(\sum_{g \in [G]} \mathbb{1}\{P \in \mathcal{P}^g\} \cdot \mathbb{E}_{P^g} M_\tau^g \right) \leq G. \quad (9)$$

These definitions satisfy the clear inclusion relations $1 \subseteq 2$ and $3 \subseteq 4$. 1 and 3, for example, are not included in either direction, because while adaptivity to a larger filtration is a weaker condition, returning an e-value at any stopping time on a larger filtration is a stronger condition. Nonetheless, local and local compound e-processes are much more natural and straightforward to construct, as we shall argue soon. Our definitions lead to the following direct corollary of Lemma 2.2.

Theorem 2.5. Let $(\{M_n^g\} : g \in [G])$ be global compound e-processes. Then, the set process $\{\text{eBH}_\alpha(M_n^g : g \in [G])\}$ satisfies the level- α stopped FDR control.

We describe the stopped e-BH (se-BH) procedure following Theorem 2.5 as Algorithm 1. The “any stopping time” in the algorithm, we note, can depend on the rejection set R_n . For example, it can be $\min\{n : |R_n| \geq 100\}$, the first time when 100 hypotheses are rejected. We shall finally show in Section 6 that the converse to Theorem 2.5 holds: a procedure controls the stopped FDR if and only if it is the stopped e-BH procedure with global compound e-processes.

Algorithm 1 Stopped e-BH

Require: null sets $(\mathcal{H}_0^g : g \in [G])$; global compound e-processes $(\{M_n^g\} : g \in [G])$; FDR level $\alpha \in (0, 1)$

Ensure: stopped FDR $\leq \alpha$

$M_0^g \leftarrow 1$ for all $g \in [G]$

for $n = 1, 2, \dots$ **do**

compute M_n^g for all $g \in [G]$

$R_n \leftarrow \text{eBH}_\alpha(M_n^g : g \in [G])$

at any stopping time: **return** R_n

end for

If the local filtrations are independent, then local e-processes are global e-processes (Appendix A). However, this rarely happens in reality as it is common to have cross-hypothesis dependence. How can one construct global or global compound e-processes under dependence? By definition, the local filtration $\{\mathcal{F}_g^n\}$ describes \mathcal{H}_0^g -specific information. As most of the growing e-process literature is concerned with a single hypothesis, the construction of e-processes often does not incorporate (and more accurately, does not *allow*) external randomness. If an e-process $\{M_n^g\}$ is constructed to test \mathcal{H}_0^g , it is naturally one *for* \mathcal{P}^g *on* $\{\mathcal{F}_n^g\}$, thus a local e-process on which only stopping at a local stopping time on $\{\mathcal{F}_n^g\}$ yields an e-value. There is no guarantee whatsoever that stopping at a “wild”, outside, non- $\{\mathcal{F}_n^g\}$ stopping time, the stopped value remains an e-value. The toolbox for global e-processes, therefore, is inherently limited.

In fact, the subtlety brought by filtrations is noted in numerous recent papers on e-processes, even in the single hypothesis testing case. For example, Wang and Ramdas (2025) note that in a problem as simple as testing if an i.i.d. Gaussian sequence X_1, X_2, \dots has mean 0, there are e-processes on a sub-filtration of the canonical filtration $\{\sigma(X_i : i \leq n)\}$, which return non-e-values when stopped at stopping times on $\{\sigma(X_i : i \leq n)\}$.

Xu et al. (2021) are aware of this issue, and in their formulation of bandit e-BH framework, they indeed assume that all e-processes are e-processes with respect to a shared global filtration. It still remains to be answered under what circumstances one may construct e-processes locally within a hypothesis using any of the many existing e-processes in the literature and still enjoy the global property to stop them at a single cross-hypothesis stopping time.

3 Global e-processes by a Markovian assumption

We formulate the sequential multiple testing problem with the following set-up. Consider at time $n = 1, 2, \dots$, we observe the n^{th} observation $Z_n = (X_n, \mathbf{Y}_n)$ that includes the covariate $X_n \in \mathcal{X}$ and an array of response variables $\mathbf{Y}_n = (Y_n^1, \dots, Y_n^G) \in \mathcal{Y}_1 \times \dots \times \mathcal{Y}_G$.

We assign to each $g \in [G]$ the following model, which can be parametric or nonparametric, describing the *conditional* distribution of the response variable Y_n^g given the covariate X_n :

$$Y^g | X, \theta \sim p_g(Y^g | X, \theta) \quad (10)$$

where $\theta \in \Theta$ is the main parameter of interest, and $p_g(\cdot | x, \theta)$ is a distribution over \mathcal{Y}_g for any $x \in \mathcal{X}$ and $\theta \in \Theta$.

The conditional model (10) naturally leads to the following causal assumption.

Assumption 3.1. For any $n \geq 2$, $\mathbf{Y}_n \perp (X_1, \dots, X_{n-1}; \mathbf{Y}_1, \dots, \mathbf{Y}_{n-1}) | X_n$.

The assumption states that, given the covariate X_n , there is allowed an arbitrary dependence *between* the G response variables Y_n^1, \dots, Y_n^G ; but the array of these response variables *as a whole* is independent from everything one has previously collected. \mathbf{Y}_n depends on the past only through its covariate X_n , a property akin to that of a Markov chain. In fact, Assumption 3.1 states that the sequence

$$X_1, \mathbf{Y}_1, X_2, \mathbf{Y}_2, X_3, \mathbf{Y}_3 \dots \quad (11)$$

is Markovian at the \mathbf{Y}_n 's. Further, Assumption 3.1 poses no limitation on the covariate X_n . It can be random, deterministic, chosen or sampled adaptively depending on previous observations and inference results.

Since arbitrary dependence between Y_n^1, \dots, Y_n^G is allowed, the model (10) thus describes only the marginal distributions of Y_n^g conditioned on X_n . We therefore carefully write down the definition of the model set \mathcal{M} , null set \mathcal{P}^g , and null hypothesis \mathcal{H}_0^g for each $g \in [G]$, connecting to our previous terminology.

Definition 3.2. \mathcal{M} contains all distributions for $\{Z_n\}_{n \geq 1}$ satisfying Assumption 3.1 such that there exists a $\theta \in \Theta$, for all $n \geq 1$, the conditional distribution of the G -tuple

$$\mathbf{Y}_n | X_n \quad (12)$$

satisfies that for all $g \in [G]$, the g^{th} marginal distribution equals $p_g(\cdot | X_n, \theta)$. \mathcal{P}^g contains all such distributions such that the θ above can be chosen from Θ^g ; and when this happens, the null hypothesis \mathcal{H}_0^g holds.

The global filtration we work with differs slightly from the obvious choice $\{\sigma(Z_i : i \leq n)\}$ which served as an illustrating example in the previous section, but is instead defined as follows:

$$\mathcal{F}_n = \sigma(\mathbf{Y}_i, X_j : 1 \leq i \leq n, 1 \leq j \leq n+1). \quad (13)$$

That is, \mathcal{F}_n includes all the information after the $(n+1)^{\text{st}}$ covariate is available but before the $(n+1)^{\text{st}}$ response variables are revealed. With these definitions and assumptions above, we establish the following theorem stating that locally defined e-processes can be lifted to this global filtration.

Theorem 3.3. Let $g \in [G]$. Suppose for each $n \geq 1$, there is an \mathcal{F}_{n-1} -measurable random function $E_n : \Theta_0^g \times \mathcal{X} \times \mathcal{Y}_g \rightarrow \mathbb{R}_{\geq 0}$ that satisfies

$$\sup_{\substack{\theta \in \Theta_0^g \\ x \in \mathcal{X}}} \int_{\mathcal{Y}_g} E_n(\theta, x, y) p_g(dy | x, \theta) \leq 1. \quad (14)$$

Then, under Assumption 3.1, for every $\theta \in \Theta_0^g$ the process

$$M_n(\theta) = \prod_{i=1}^n E_i(\theta, X_i, Y_i^g) \quad (15)$$

is a nonnegative supermartingale on $\{\mathcal{F}_n\}$ under $p_g(\cdot | \cdot, \theta)$. Consequently, the process

$$U_n^g = \inf_{\theta \in \Theta_0^g} M_n(\theta) \quad (16)$$

is an e-process on $\{\mathcal{F}_n\}$ under the null Θ_0^g .

Proof. First, it is easy to see that $\{M_n(\theta)\}$ is adapted to $\{\mathcal{F}_n\}$ since $M_n(\theta)$ only depends on $\{Z_i : 1 \leq i \leq n\}$. Therefore, to see that it is a supermartingale, it suffices to show that $\mathbb{E}_{\theta^g}[E_n(\theta, X_n, Y_n^g) | \mathcal{F}_{n-1}] \leq 1$. This follows from

$$\mathbb{E}_{\theta}[E_n(\theta, X_n, Y_n^g) | \mathcal{F}_{n-1}] \quad (17)$$

$$= \mathbb{E}_{\theta}[E_n(\theta, X_n, Y_n^g) | \mathbf{Y}_i, X_j : 1 \leq i \leq n-1, 1 \leq j \leq n] \quad (18)$$

$$\text{(Assumption 3.1)} \quad (19)$$

$$= \mathbb{E}_{\theta}[E_n(\theta, X_n, Y_n^g) | X_n] = \int_{\mathcal{Y}_g} E_n(\theta, X_n, y) p_g(dy | X_n, \theta) \leq 1, \quad (20)$$

where \mathbb{E}_{θ} denotes the expectation under $p_g(\cdot | \cdot, \theta)$. \square

We remark that in the theorem above, the local e-process that tests \mathcal{H}_g^0 is computed via the infimum over a \mathcal{H}_g^0 -indexed family of nonnegative supermartingales, each incrementally updated by an e-value $E_n(\theta, X_n, Y_n^g)$ at time n that only takes X_n and Y_n^g . The assumption that E_n being \mathcal{F}_{n-1} -measurable is only for the sake of generality. In a purely local construction we often let E_n be $\sigma(X_i, Y_i^g : i \leq n-1)$ -measurable, which of course implies \mathcal{F}_{n-1} -measurability. Recalling the second equivalent *definition* of e-processes in the opening of Section 2.2, this procedure is universal as it encompasses every possible local construction of e-processes. Theorem 3.3 then states that these local e-processes are global. Combining with Theorem 2.5, we see that these global e-processes stopped at a global stopping time can produce FDR-controlled multiple testing results with e-BH.

Corollary 3.4. *The set process $\{\text{eBH}_\alpha(M_n^g : g \in [G])\}$ satisfies the level- α stopped FDR control. More generally, let $(w_g : g \in [G])$ be G nonnegative numbers such that $\sum_{g \in [G]} w_g \leq G$. The set process $\{\text{eBH}_\alpha(w_g M_n^g : g \in [G])\}$ also satisfies the level- α stopped FDR control.*

4 Global e-processes via compound adjusters

While we have previously discussed certain conditions under which local e-processes are naturally global e-processes, we now quote a complementary recent result by [Choe and Ramdas \(2024\)](#) which states that by slightly “muting” an e-process, it becomes an e-process on arbitrary refinement of the filtration. This is helpful when [Assumption 3.1](#) does not hold or is not verifiable. We first recall the definition of an *adjuster*, which traces back to [Shafer et al. \(2011\)](#).

Definition 4.1. *An adjuster is a non-decreasing function $A : \mathbb{R}_{\geq 1} \rightarrow \mathbb{R}_{\geq 0}$ such that*

$$\int_1^\infty \frac{A(x)}{x^2} dx \leq 1. \quad (21)$$

Simple examples include $x \mapsto kx^{1-k}$ for $k \in (0, 1)$ and $\sqrt{x} - 1$. Applying an adjuster to an e-process, or more generally the running maximum of an e-process is referred to as “e-lifting” by [Choe and Ramdas \(2024, Theorem 2\)](#) and yields an e-process for any finer filtrations. That is, if $\{M_n\}$ is an e-process for \mathcal{P} on $\{\mathcal{G}_n\}$, then for any adjuster A and any filtration $\{\mathcal{F}_n\} \supseteq \{\mathcal{G}_n\}$, the adjusted process $\{M_n^a\}$ where

$$M_0^a = 1, \quad M_n^a = A\left(\max_{0 \leq i \leq n} M_i\right) \quad (22)$$

is an e-process for \mathcal{P} on $\{\mathcal{F}_n\}$. This immediately leads to the following implication, beginning with a compound extension of [Definition 4.1](#).

Definition 4.2. *Let $(A^g : g \in [G])$ be a family of non-decreasing functions $\mathbb{R}_{\geq 1} \rightarrow \mathbb{R}_{\geq 0}$. They are called compound adjusters if their average is an adjuster, i.e.,*

$$\int_1^\infty \frac{\sum_{g \in [G]} A^g(x)}{x^2} dx \leq G. \quad (23)$$

That is, compound adjusters arise from decomposing an adjuster into G monotone components. For example, a vector of G adjusters are compound adjusters; and similar to compound e-values, one can construct compound adjusters by taking a weighted sum of G adjusters as long as weights sum up $\leq G$.

Corollary 4.3. *Let $(\{M_n^g\} : g \in [G])$ be local e-processes and $(A^g : g \in [G])$ be compound adjusters. Then, the set process $\{\text{eBH}_\alpha(M_n^{ga} : g \in [G])\}$ satisfies the level- α stopped FDR control, where*

$$M_0^{ga} = 1, \quad M_n^{ga} = A^g\left(\max_{0 \leq i \leq n} M_i^g\right). \quad (24)$$

5 Examples and counterexamples

5.1 Parametric regression and universal inference

Let us present an example of [Theorem 3.3](#) that involves a parametric model with computable and optimizable likelihood functions, where the e-processes are computed according to the *universal inference* method due to [Wasserman et al. \(2020\)](#). The example allows the Markov-like condition [Assumption 3.1](#) satisfied with arbitrary cross-hypothesis dependence among $\mathbf{Y}|X$.

Let $\mathcal{X} = \mathbb{R}^d$, $\mathcal{Y}^g = \mathbb{R}$. We assume the parameter θ can be split into G \mathbb{R}^m -valued, hypothesis-specific parameters ($\theta_g = \phi^g(\theta) : g \in [G]$) via G functions $\phi^g : \Theta \rightarrow \mathbb{R}^m$, with the g^{th} null region being $\Phi_0^g \subseteq \mathbb{R}^m$; that is,

$$\Theta_0^g = \{\theta \in \Theta : \phi^g(\theta) \in \Phi_0^g\}. \quad (25)$$

For example, Θ can be $\mathbb{R}^{m \times G}$ and $\phi^g(\theta)$ can be the g^{th} column of the matrix θ . But our assumption above is more flexible. With a slight abuse of notations, we assume the model (10) is specified by the conditional probability density or mass functions $p_g(y|x, \theta^g)$. Then, the stepwise e-value function E_n can be simply chosen as the likelihood ratio

$$E_n(\theta, x, y) = \frac{p_g(y|x, \tilde{\theta}_{n-1}^g)}{p_g(y|x, \theta^g)}, \quad (26)$$

where $\tilde{\theta}_{n-1}^g$ is any estimator for θ^g computed from Z_1, \dots, Z_{n-1} , thus is \mathcal{F}_{n-1} -measurable. To see that (14) holds,

$$\int_{\mathcal{Y}^g} E_n(\theta, x, y) p_g(y|x, \theta^g) dy = \int_{\mathcal{Y}^g} p(y|x, \tilde{\theta}_{n-1}^g) dy = 1. \quad (27)$$

In particular, define the maximum likelihood estimates within the null and the full models

$$\hat{\theta}_n^g = \operatorname{argmax}_{\theta \in \Theta_0^g} \prod_{i=1}^n p(Y_i^g | X_i, \theta), \quad \tilde{\theta}_n^g = \operatorname{argmax}_{\theta \in \Theta^g} \prod_{i=1}^n p(Y_i^g | X_i, \theta). \quad (28)$$

Then the e-process $\{U_n^g\}$ takes the following simple form

$$U_n^g = \prod_{i=1}^n \frac{p(Y_i^g | X_i, \tilde{\theta}_{i-1}^g)}{p(Y_i^g | X_i, \hat{\theta}_i^g)}. \quad (29)$$

Numerous generalized regression-like statistical models take this form. For example in genomics, the vector X_n consists of the d regressor variables of the n^{th} cell, and for each gene $g \in [G]$, the vector θ^g consists of the $m = d$ linear coefficients controlling its expression level. Y_n^g is the expression level of gene g in cell n , and follows a negative binomial distribution with mean $\exp(X_n^\top \theta^g)$ and dispersion γ_g estimated separately. That is, $p_g(y|x, \theta^g) = \text{nb}(y | \exp(x^\top \theta^g), \gamma_g)$ where $\text{nb}(y|\mu, \gamma)$ is the probability mass function of a negative binomial distribution with mean μ and dispersion γ , which has a closed-form expression and its maximum likelihood estimates (28) have already been computed frequently in numerous biostatistics packages.

5.2 Nonparametric heavy-tailed conditional mean testing

Our second example of Theorem 3.3 is nonparametric, where $\mathcal{Y}^g = \mathbb{R}$ and for each $g \in [G]$ we specify a function $\mu^g : \mathcal{X} \rightarrow \mathbb{R}$. We test the hypotheses

$$\mathcal{H}_0^g : \mathbb{E}[Y^g | X] \leq \mu^g(X) \quad (30)$$

under the finite conditional variance assumption $\mathbf{Var}[Y^g | X] \leq v^g(X)$. Therefore, the parameter set Θ indexes *all* conditional laws $\mathbf{Y}|X$ such that $\sup_{g \in [G]} (\mathbf{Var}[Y^g | X] - v^g(X)) \leq 0$, and Θ_0^g all those in Θ with $\mathbb{E}[Y^g | X] \leq \mu^g(X)$. Again, Assumption 3.1 is satisfied with arbitrary cross-hypothesis dependence among $\mathbf{Y}|X$. We employ the following ‘‘Catoni-style’’ e-value due to Wang and Ramdas (2023):

$$E_n(\theta, x, y) = \exp(\phi(\lambda_n(y - \mu^g(x))) - \lambda_n^2 v^g(x)/2) \quad (31)$$

where

$$\phi(x) = \begin{cases} \log(1 + x + x^2/2), & x \geq 0, \\ -\log(1 - x + x^2/2), & x < 0, \end{cases} \quad (32)$$

and $\lambda_n > 0$ is \mathcal{F}_{n-1} -measurable. Of note, this function of E does *not* depend on the “parameter” θ that now lives in an infinite-dimensional space. The corresponding e-process therefore equals the supermartingale

$$U_n^g = M_n(\theta) = \exp \left\{ \sum_{i=1}^n \phi(\lambda_i(Y_i^g - \mu^g(X_i))) - \frac{\sum_{i=1}^n \lambda_i^2 v^g(X_i)}{2} \right\}. \quad (33)$$

If the nulls are $\mathbb{E}[Y^g|X] \geq \mu^g(X)$ instead, one can simply replace ϕ with $-\phi$; and if they are $\mathbb{E}[Y^g|X] = \mu^g(X)$, use a convex combination (e.g. the average) of the e-process for $\mathbb{E}[Y^g|X] \leq \mu^g(X)$ and the e-process for $\mathbb{E}[Y^g|X] \geq \mu^g(X)$. This example thus encompasses a wide scope of nonparametric problems including heavy-tailed linear regression.

5.3 Failure of Assumption 3.1

We now illustrate a simple counterexample where the Markovian causal condition (Assumption 3.1) does not hold. Our example is akin to a recent construction by [Dandapanthula and Ramdas \(2025, Section 9.2\)](#). This involves $G = 2$ streams of variables where the second stream “peeks into the future” of the first stream.

We construct two different set-ups indexed by $d \in \{0, 1\}$. Consider empty covariates $\{X_n\}$, and the response variables for \mathcal{H}_0^1 , $\{Y_n^1\}$, an i.i.d. sequence of $\text{Ber}(\theta)$ random variables. Now, we define $Y_n^2 = Y_{n+d}^1$ for all $n \geq 1$, where $d \in \{0, 1\}$. That is, the second stream, while also an i.i.d. $\text{Ber}(\theta)$ sequence, reproduces the coin tosses of the first stream if $d = 0$; and foretells the upcoming coin tosses of the first stream if $d = 1$. The statistician tests $\mathcal{H}_0^1 : \theta = \theta_1$ using $\{Y_n^1\}$, and $\mathcal{H}_0^2 : \theta = \theta_2$ using $\{Y_n^2\}$ with two local e-processes (for example by universal inference) and the stopped e-BH.

If $d = 0$, the test is perfectly valid as Assumption 3.1, equivalent to

$$Y_n^1 \perp (Y_1^1, \dots, Y_{n-1}^1), \quad (34)$$

holds. However, the test is invalid if $d = 1$ as Assumption 3.1, now equivalent to

$$(Y_n^1, Y_{n+1}^1) \perp (Y_1^1, \dots, Y_n^1), \quad (35)$$

is not satisfied.

The scenario above does not happen as long as the time index n is a bona fide representation of the chronological evolution of the experiment, and no retrocausality is allowed. It warns, however, that in asynchronous experiments or settings where retrocausality is not excluded¹, one must beware the possible violation of Assumption 3.1 and conduct the sequential hypothesis testing with care: local e-processes may fail to be global e-processes. A remedy, we note, is the filtration-agnostic adjustment we mentioned in Section 4.

6 Universality of se-BH and global compound e-processes

We establish the following converse to Theorem 2.5.

Theorem 6.1. *Let $\{R_n\}$ be a set process adapted to $\{\mathcal{F}_n\}$ satisfying the level- α stopped FDR control. Then, there exist G global compound e-processes $(\{M_n^g\} : g \in [G])$ for $(\mathcal{P}^g : g \in [G])$ such that*

$$R_n = \text{eBH}_\alpha(M_n^g : g \in [G]) \quad \text{for all } n. \quad (36)$$

¹For example, there has been an ongoing debate in the philosophy of quantum mechanics community whether a retrocausal theory is valid ([Friederich and Evans, 2023](#)). This is much beyond the scope of the current paper.

Proof. First, at any non-random n , since R_n is a \mathcal{F}_n -measurable random set that controls the FDR at level α , it follows from [Ignatiadis et al. \(2024, Theorems 3.1 and 3.3\)](#) that

$$R_n = \text{eBH}_\alpha(M_n^g : g \in [G]) \quad (37)$$

for G compound e-values M_n^1, \dots, M_n^G defined by

$$M_n^g = \frac{G}{\alpha} \cdot \frac{\mathbb{1}\{g \in R_n\}}{|R_n| \vee 1}. \quad (38)$$

Hence these compound e-values are \mathcal{F}_n -measurable as well. We have thus defined the processes $\{M_n^g\}$ for $g \in [G]$, all globally adapted, and it remains to verify they are compound e-processes globally. Take any $\tau \in \mathbb{T}$. Since the FDR is controlled for the set R_τ , again we have

$$R_\tau = \text{eBH}_\alpha(E_\tau^g : g \in [G]) \quad (39)$$

for G compound e-values $E_\tau^1, \dots, E_\tau^G$ where

$$E_\tau^g = \frac{G}{\alpha} \cdot \frac{\mathbb{1}\{g \in R_\tau\}}{|R_\tau| \vee 1}, \quad (40)$$

which equals M_τ^g . Therefore

$$\sum_{g \in [G]} \mathbb{1}\{P \in \mathcal{P}^g\} \mathbb{E}_P M_\tau^g = \sum_{g \in [G]} \mathbb{1}\{P \in \mathcal{P}^g\} \mathbb{E}_P E_\tau^g \leq G \quad (41)$$

under any $P \in \mathcal{M}$, concluding that they are global compound e-processes. \square

7 Summary

We propose the stopped e-BH procedure for sequential, anytime-valid multiple hypothesis testing, a procedure both necessary and sufficient for strict false discovery rate control at all stopping times. We carefully distinguish local and global stopping times and e-processes: local e-processes are more commonly constructed, but global (compound) e-processes are more relevant to the FDR control objective. A crucial causal condition is identified that bridges the local-global distinction. Our work demonstrates the theoretical foundation of using e-values and e-processes for large-scale and complex scientific experiments. These experiments can have arbitrarily dependent hypotheses, sample sizes and sampling schemes for which allowing optional stopping and continuation is necessarily beneficial.

References

- Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(1):289–300, 1995.
- Y. J. Choe and A. Ramdas. Combining evidence across filtrations. *arXiv preprint arXiv:2402.09698*, 2024.
- S. Dandapanthula and A. Ramdas. Multiple testing in multi-stream sequential change detection. *arXiv e-prints*, pages arXiv–2501, 2025.
- S. Friederich and P. W. Evans. Retrocausality in Quantum Mechanics. In E. N. Zalta and U. Nodelman, editors, *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, Winter 2023 edition, 2023.

- N. Ignatiadis, R. Wang, and A. Ramdas. Asymptotic and compound e-values: multiple testing and empirical Bayes. *arXiv preprint arXiv:2409.19812*, 2024.
- G. Shafer, A. Shen, N. Vereshchagin, and V. Vovk. Test martingales, bayes factors and p-values. 2011.
- H. Wang and A. Ramdas. Catoni-style confidence sequences for heavy-tailed mean estimation. *Stochastic Processes and their Applications*, 163:168–202, 2023. ISSN 0304-4149. doi: 10.1016/j.spa.2023.05.007.
- H. Wang and A. Ramdas. Anytime-valid t-tests and confidence sequences for Gaussian means with unknown variance. *Sequential Analysis*, 0(0):1–55, 2025. doi: 10.1080/07474946.2024.2428245.
- R. Wang and A. Ramdas. False discovery rate control with e-values. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 84(3):822–852, 2022.
- L. Wasserman, A. Ramdas, and S. Balakrishnan. Universal inference. *Proceedings of the National Academy of Sciences*, 117(29):16880–16890, 2020.
- Z. Xu, R. Wang, and A. Ramdas. A unified framework for bandit multiple testing. *Advances in Neural Information Processing Systems*, 34:16833–16845, 2021.

A Independent local filtrations

In this appendix, we formally prove that when there is no cross-hypothesis dependence, local e-processes are always global e-processes.

Theorem A.1. *Following the terminology in Definition 2.4, if the local σ -algebras at infinity ($\mathcal{F}_\infty^g : g \in [G]$) are independent, local e-processes ($\{M_n^g\} : g \in [G]$) are global.*

Proof. Our proof is based on the two equivalent definitions of e-processes mentioned in Section 2.2. Fix a $g \in [G]$. Since $\{M_n^g\}$ is an e-process for \mathcal{P}^g on $\{\mathcal{F}_n^g\}$, for every $P \in \mathcal{P}^g$, it is upper bounded by a nonnegative supermartingale $\{N_n^P\}$ on $\{\mathcal{F}_n^g\}$ with $N_0^P = 1$. It suffices to prove it is a supermartingale on $\{\mathcal{F}_n\}$ as well. This follows from

$$\mathbb{E}[N_{n+1}^P | \mathcal{F}_n] = \mathbb{E}[N_{n+1}^P | \mathcal{F}_n^g; \mathcal{F}_n^h : h \in [G] \setminus \{g\}] \stackrel{(*)}{=} \mathbb{E}[N_{n+1}^P | \mathcal{F}_n^g] \leq N_n, \quad (42)$$

where the equality (*) follows from lifting Lemma A.2 to σ -algebras: $\sigma(\mathcal{F}_n^h : h \in [G] \setminus \{g\})$ being independent from $\sigma(N_{n+1}^P, \mathcal{F}_n^g)$. \square

Lemma A.2. *Let A, B, C be events in a probability space such that B is independent from $\sigma(A, C)$, and $\mathbb{P}[B], \mathbb{P}[C] > 0$. Then $\mathbb{P}[A|B \cap C] = \mathbb{P}[A|C]$.*

Proof. $\mathbb{P}[A|B \cap C] = \frac{\mathbb{P}[A \cap B \cap C]}{\mathbb{P}[B \cap C]} = \frac{\mathbb{P}[A \cap C] \mathbb{P}[B]}{\mathbb{P}[B] \mathbb{P}[C]} = \frac{\mathbb{P}[A \cap C]}{\mathbb{P}[C]} = \mathbb{P}[A|C]$. \square