STRENGTH AND PARTITION RANK UNDER LIMITS AND FIELD EXTENSIONS

ARTHUR BIK, JAN DRAISMA, AMICHAI LAMPERT, AND TAMAR ZIEGLER

ABSTRACT. The strength of a multivariate homogeneous polynomial is the minimal number of terms in an expression as a sum of products of lowerdegree homogeneous polynomials. Partition rank is the analogue for multilinear forms. Both ranks can drop under field extensions, and both can jump in a limit. We show that, for fixed degree and under mild conditions on the characteristic of the ground field, the strength is at most a polynomial in the border strength. We also establish an analogous result for partition rank. Our results control both the jump under limits and the drop under field extensions.

1. INTRODUCTION AND MAIN RESULTS

Let K be a field and let $f \in K[x_1, \ldots, x_n]_d$ be a form (i.e., homogeneous polynomial) of degree d. In [Sch85], Schmidt introduced the following measure of complexity for f:

Definition 1.0.1. The strength of f, denoted $s_K(f)$, is the minimal number r of terms in any expression

$$f = \sum_{i=1}^{r} a_i b_i$$

where the a_i and b_i are forms in $K[x_1, \ldots, x_n]$ of positive degrees. If d = 1 and f is nonzero, then we define $s_K(f) = \infty$.

1.1. Strength and field extensions. Strength, also known as Schmidt rank, was independently introduced by Green-Tao in their work on distribution of polynomials over finite fields [GT09] and by Ananyan-Hochster in their proof of Stillman's conjecture [AH20]. One of the key results in the work of both Schmidt and Ananyan-Hochster is that for algebraically closed K, the strength of a form is closely related to the codimension of its singular locus.

This relationship also plays a central role in the arithmetic applications studied by Schmidt and Green-Tao, where, however, K is not algebraically closed. It is therefore important to understand the behaviour of strength under field extensions. If L is a field extension of K, then we may also consider f as a form with coefficients in L. The inequalities $s_{\overline{K}}(f) \leq s_L(f) \leq s_K(f)$ are immediate, where \overline{K} is an algebraic closure of K. The following simple example shows that these inequalities may be strict.

JD was partially supported by Swiss National Science Foundation project grant 200021-227864.

AL was supported by NSF grants DMS-2402041 and DMS-1926686.

TZ was supported by ISF grant 2112/20.

The authors thank the Institute for Advanced Study (Princeton, USA) for excellent working conditions; this collaboration started there in Fall 2022.

Example 1.1.1. For d = 2, $s_K(f)$ is the minimal codimension of a subspace of K^n on which the quadratic form f vanishes identically. So for $f = x_1^2 + \cdots + x_n^2$ we have $s_{\mathbb{R}}(f) = n$ and $s_{\mathbb{C}}(f) = \lceil \frac{n}{2} \rceil$.

Bik-Draisma-Snowden give an example [BDS24b, Example 3.4] showing that, in general, the drop in strength can be arbitrarily large. However, for semi-perfect fields they show in [BDS24a, Theorem 1.3] that $s_K(f) \leq \Gamma(s_{\overline{K}}(f), d)$ where Γ is some non-explicit function that may depend on K.

For certain fields satisfying char(K) = 0 or char(K) > d, Lampert-Ziegler proved quantitative bounds for Γ [LZ24]. Our first theorem gives a quantitative bound for any field satisfying this condition.

Theorem 1.1.2. Suppose that $s_{\overline{K}}(f) = r$ and that $\operatorname{char}(K) = 0$ or $\operatorname{char}(K) > d$. Then $s_K(f) \ll_d r^{d-1}$ for infinite K. If K is finite then $s_K(f) \ll_d r^{d-1} \log r$.

Here, \ll_d means that the left-hand side is bounded by a constant times the right-hand side, where the constant depends only on d (and in particular not on K).

1.2. **De-bordering strength.** Theorem 1.1.2 is a consequence of a stronger result which we now describe. In the following definition, we write \mathcal{P}_d for the affine space over \mathbb{Z} with coordinates labelled by monomials in x_1, \ldots, x_n of degree d. The K-points of this scheme are the homogeneous forms of degree d in x_1, \ldots, x_n with coefficients in K.

Definition 1.2.1. The border strength $\underline{s}(f)$ of $f \in K[x_1, \ldots, x_n]_d$ is defined as the minimal r for which there exist $d_1, \ldots, d_r \in \{1, \ldots, d-1\}$ such that f is a K-point in the Zariski closure of the image of the morphism

$$\mu_{d_1,\dots,d_r}:\prod_{i=1}^r (\mathcal{P}_{d_i} \times \mathcal{P}_{d-d_i}) \to \mathcal{P}_d,$$
$$((a_1,b_1),\dots,(a_r,b_r)) \mapsto \sum_{i=1}^r a_i b_i.$$

The border strength of a nonzero linear form is defined as ∞ .

 \diamond

Write \overline{K} for an algebraic closure of K. The border strength $\underline{s}(f)$ is also the minimal r such that f lies in the Zariski closure in $\overline{K}[x_1, \ldots, x_n]_d$ of forms of strength $\leq r$. The latter set is the union of the images of the μ_{d_1,\ldots,d_r} , hence constructible by Chevalley's theorem. Hence for $K = \mathbb{C}$, since constructible sets have the same closure in the Zariski topology as in the Euclidean topology, $\underline{s}(f)$ is also the minimal r such that there exists a sequence of forms of strength $\leq r$ that converges to f in the Euclidean topology. This explains the term border strength.

Remark 1.2.2. We do not know an explicit example of a form over \mathbb{C} whose border strength is strictly lower than its strength. However, such forms do exist for d = 4 [BBOV22], and the proof there can be adapted to d > 4. For $d \leq 3$ they do not exist, since there strength is the minimal codimension of a linear space on which the form vanishes identically, and vanishing on a large subspace is a Zariski-closed condition on the form; see [DES17].

We have the following fundamental inequalities:

(1)
$$s_K(f) \ge s_{\overline{K}}(f) \ge \underline{s}(f)$$

The goal of this paper is to upper bound $s_K(f)$ as a function of $\underline{s}(f)$, as follows.

Theorem 1.2.3. Suppose that $\underline{s}(f) = r$ and $\operatorname{char}(K) = 0$ or $\operatorname{char}(K) > d$. For infinite K we have $s_K(f) \ll_d r^{d-1}$ and for finite K we have $s_K(f) \ll_d r^{d-1} \log r$.

Note that Theorem 1.1.2 follows immediately by the inequalities (1). Following the theoretical computer science literature $[DGI^+22]$, we call a result like that in Theorem 1.2.3 a *de-bordering result*.

1.3. Expressing low-strength polynomials as polynomials in derivatives. As a by-product of the proof of Theorem 1.2.3 and its generalisations below, we will establish the following result. Given $f \in K[x_1, \ldots, x_n]_d$, let D(f) be the subspace of $\bigoplus_{e=1}^{d-1} K[x_1, \ldots, x_n]_e$ spanned by all polynomials of the form

$$\frac{\partial}{\partial x_{i_1}}\cdots\frac{\partial}{\partial x_{i_l}}f$$

with $l \in \{1, \ldots, d-1\}$ and $i_1, \ldots, i_l \in \{1, \ldots, n\}$.

Theorem 1.3.1. Suppose $\underline{s}(f) = r$ and $\operatorname{char}(K) = 0$ or $\operatorname{char}(K) > d$. Then, if K is infinite, f lies in a subalgebra generated by $\ll_d r^d$ elements of D(f). For finite K, f lies in a subalgebra generated by $\ll_d r^d \log r$ elements of D(f).

1.4. Connections to the literature. Bounds for the strength of a form and for the partition rank of a tensor in terms of its strength / partition rank over the algebraic closure have been found for cubics in [AKZ21], for quartics in [KP23], and for general degree but over certain fields in [LZ24, BL24]. In [AKZ21], these results were applied to finite fields in order to bound strength/partition rank in terms of *analytic rank*. A well-known conjecture in higher order Fourier analysis is that, for fixed degree, strength is bounded by a constant multiple of analytic rank; for the currently strongest results along these lines see [CM23, MZ22]. Adiprasito-Kazhdan-Ziegler conjectured [AKZ21] that Theorem 1.1.2 holds with linear bounds, and showed that this implies linear bounds for strength in terms of analytic rank. In [MZ22] and [BL24], quasi-linear bounds were obtained for finite fields. See [CY24] for additional results related to this conjecture. We can now formulate two natural stronger versions of Theorems 1.2.3 and 1.3.1.

Question 1.4.1. Can the bounds in Theorem 1.2.3 and/or Theorem 1.3.1 be taken linear in r?

Another strand of research, motivated by questions in theoretical computer science, concerns *de-bordering* complexity measures. A recent result in this direction is that the Waring rank of a degree-*d* form over \mathbb{C} admits an upper bound that is exponential in its border Waring rank but *linear* in *d* [DGI⁺22].

Finally, our work is closely related to Karam's result that says that bounded partition rank can be recognised by looking at all subtensors of some given size [Kar22]. Indeed, a quantitative version of that statement involved degree estimations for equations vanishing on tensors of bounded partition rank similar to those we will use below; see [DK24].

1.5. **De-bordering collective strength.** Our techniques for proving Theorem 1.2.3 also apply to tuples of polynomials, to tensors, and to tuples of tensors. In this and the next few sections, we record these additional results. First, for tuples of degree-d forms, the following notion is a natural analogue of strength.

Definition 1.5.1. Let $\underline{f} = (f_1, \ldots, f_m) \in K[x_1, \ldots, x_n]_d^m$. The strength $s_K(\underline{f})$ is the minimum of $s_K(c_1f_1 + \cdots + c_mf_m)$ over all $(c_1, \ldots, c_m) \in K^m \setminus \{(0, \ldots, 0)\}$.

Thus the strength is 0 if and only if f_1, \ldots, f_m are linearly dependent.

Definition 1.5.2. The border strength $\underline{s}(\underline{f})$ is defined as the minimal r such that there exist $d_1, \ldots, d_r \in \{1, \ldots, d-1\}$ for which \underline{f} is a K-point in the image closure of the following morphism over \mathbb{Z} :

(2)
$$\mu_{m,d_1,\dots,d_r} : \mathbb{A}^{m \times m} \times (\mathcal{P}_d)^{m-1} \times \prod_{i=1}^{r} (\mathcal{P}_{d_i} \times \mathcal{P}_{d-d_i}) \to (\mathcal{P}_d)^m$$
$$(g, (h_1,\dots,h_{m-1}), ((a_1,b_1),\dots,(a_r,b_r)))$$
$$\mapsto g \cdot (h_1,\dots,h_{m-1},\mu_{d_1,\dots,d_r}((a_1,b_1),\dots,(a_r,b_r))).$$

Here the notation g· means that in the *i*-th position, we take the linear combination of the following *m* polynomials with coefficients given by the *i*-th row of *g*. \diamondsuit

On K-points, the morphism in (2) does the following: we pick m-1 arbitrary degree-*d* forms and a degree-*d* form of strength $\leq r$ (which admits a decomposition via μ_{d_1,\ldots,d_r}), and take *m* arbitrary linear combinations of these *r* forms. Again, $\underline{s(f)}$ is the minimal *r* such that \underline{f} lies in the Zariski closure in $\overline{K}[x_1,\ldots,x_n]_d^m$ of the set of *m*-tuples of collective strength $\leq r$.

Theorem 1.5.3. Suppose $\underline{s}(\underline{f}) = r$ and $\operatorname{char}(K) = 0$ or $\operatorname{char}(K) > d$. Then $s_K(\underline{f}) \ll_d m^3 r^{d-1}$ if K is infinite and $s_K(\underline{f}) \ll_d m^3 r^{d-1} \cdot \log(r+m)$ if K is finite. Consequently, $s_K(f)$ is also bounded by a polynomial in $s_{\overline{K}}(f)$.

1.6. **De-bordering partition rank.** The results that we will state now require no assumptions on char(K). Let V_1, \ldots, V_d be finite-dimensional K-vector spaces and let $t \in V_1 \otimes \cdots \otimes V_d$; unless stated otherwise, tensor products are over K. For a subset $I \subseteq [d] := \{1, \ldots, d\}$ we write $V_I := \bigotimes_{i \in I} V_i$. We have a canonical isomorphism $V_I \otimes V_{I^c} \to V_1 \otimes \cdots \otimes V_d$, where $I^c = [d] \setminus I$, and we will use this isomorphism to identify the two spaces.

Definition 1.6.1. The partition rank $prk_{K}(t)$ of t is the minimal r in any expression

$$t = \sum_{i=1}^{r} a_i \otimes b_i$$

where each $a_i \in V_{I_i}$ and $b_i \in V_{I_i^c}$ for some proper, nonempty subset $I_i \subset [d]$. The border partition rank prk(t) is defined in a similar fashion as the border strength. \diamond

Partition rank was introduced by Naslund in [Nas20b] and has found applications in higher Fourier analysis and additive combinatorics [CM23, GK22, Lov19, Mil19, MZ22, Nas20a]. As we will soon see, it is closely related to strength.

Remark 1.6.2. The partition rank of any tensor $t \in V_1 \otimes \cdots \otimes V_d$ is at most $r := \min_j \dim(V_j)$, because t admits an expression as above with $I_i = \{j\}$ for all i, where j attains the minimum, and with a_i running through a basis of V_j .

For a field extension L of K, write V_L for the L-vector space $L \otimes_K V$. We may think of $t \in V_1 \otimes \cdots \otimes V_d$ as an element of the tensor product $(V_1)_L \otimes_L \cdots \otimes_L (V_d)_L$, and then we have $\operatorname{prk}_L(t) \leq \operatorname{prk}_K(t)$. In particular, as before, we have

$$\operatorname{prk}_{K}(t) \ge \operatorname{prk}_{L}(t) \ge \operatorname{prk}_{\overline{K}}(t) \ge \operatorname{prk}(t).$$

Our de-bordering result for partition rank is as follows.

Theorem 1.6.3. Suppose that $\underline{\operatorname{prk}}(t) = r$. Then $\operatorname{prk}_K(t) \ll_d r^{d-1}$ if K is infinite and $\operatorname{prk}_K(t) \leq \ll_d r^{d-1} \cdot \log(r)$ if K is finite. In particular, $\operatorname{prk}_K(t)$ is also bounded by a polynomial in $\operatorname{prk}_{\overline{K}}(t)$.

1.7. De-bordering collective partition rank.

Definition 1.7.1. Let *m* be a positive integer. The *(collective) partition rank*, $\operatorname{prk}_{K}(\underline{t})$, of $\underline{t} = (t_{1}, \ldots, t_{m}) \in (V_{1} \otimes \cdots \otimes V_{d})^{m}$ is the minimum of $\operatorname{prk}_{K}(c_{1}t_{1} + \cdots + c_{m}t_{m})$ over all $(c_{1}, \ldots, c_{m}) \in K^{m} \setminus \{(0, \ldots, 0)\}$. The border *(collective) partition rank* of \underline{t} is defined similarly as before. \diamondsuit

Theorem 1.7.2. Suppose $\underline{\operatorname{prk}}(\underline{t}) = r$. Then $\operatorname{prk}_K(\underline{t}) \ll_d m^3 r^{d-1}$ if K is infinite and $\operatorname{prk}_K(\underline{t}) \ll_d m^3 r^{d-1} \log(r+m)$ if K is finite. In particular, $\operatorname{prk}_K(\underline{t})$ is also bounded by a polynomial in m and $\operatorname{prk}_{\overline{K}}(\underline{t})$.

1.8. **Proof overview.** We will establish de-bordering results for partition rank and strength by following the approach in [BDE19]. Indeed, there it was shown that if one has any tensor property that is preserved under linear maps, and a nontrivial polynomial equation satisfied by the tensors with that property, then one can derive an upper bound on the partition rank over K of the tensors with that property. In this paper, the tensor property is "having border partition rank $\leq r$ ". We find bounds for such a polynomial equation, both in terms of dimensions of the underlying vector spaces (this suffices for the result over infinite fields) and in terms of degree of the equation (this is needed for the result over finite fields), and use these to upper bound the partition rank over K.

We observe that Theorem 1.7.2 implies Theorem 1.6.3 and Theorem 1.5.3 implies Theorem 1.2.3. Furthermore, we will show that Theorem 1.7.2 implies Theorem 1.5.3. It then suffices to prove Theorem 1.7.2.

2. Preliminaries

2.1. Comparing strength and partition rank. Let V be a finite-dimensional vector space over K and write S^dV for the d-th symmetric power of V. Given any basis x_1, \ldots, x_n of V, the space S^dV is canonically isomorphic to $K[x_1, \ldots, x_n]_d$, and since strength is basis-invariant, we may transport the notion of strength to S^dV . We have linear maps determined by

$$\pi = \pi_d : V^{\otimes d} \to S^d V, \quad v_1 \otimes \ldots \otimes v_d \mapsto v_1 \cdots v_d \text{ and}$$
$$\iota = \iota_d : S^d V \to V^{\otimes d}, \quad v_1 \cdots v_d \mapsto \sum_{\pi \in S_d} v_{\pi(1)} \otimes \cdots \otimes v_{\pi(d)}.$$

These satisfy $\pi(\iota(f)) = d!f$. The following well-known proposition relates the partition rank of elements in $V^{\otimes d}$ with the strength of elements in $S^d V$.

Proposition 2.1.1. For $t \in V^{\otimes d}$ and $f \in S^d V$ we have

$$s_K(\pi(t)) \leq \operatorname{prk}_K(t) \text{ and } \operatorname{prk}_K(\iota(f)) \leq D \cdot s_K(f) \text{ where } D := \begin{pmatrix} d \\ \lfloor \frac{d}{2} \rfloor \end{pmatrix}.$$

Proof. Since π and ι are linear and strength/partition rank are subadditive, it suffices to prove these inequalities when $\operatorname{prk}_K(t) = 1$ and $s_K(f) = 1$, respectively. For $\emptyset \subseteq I \subseteq [d]$, $a \in V^{\otimes I}$ and $b \in V^{\otimes I^c}$ we have $\pi_d(a \otimes b) = \pi_e(a)\pi_{d-e}(b)$ (where

 π_e is short-hand for the concatenation of an isomorphism $V^{\otimes I} \to V^{\otimes e}$ induced by a bijection $I \to [e]$ and the map π_e ; and similarly for π_{d-e}). This yields the first inequality.

For $a \in S^e V$ and $b \in S^{d-e} V$ we have

$$\iota(ab) = \sum_{I \subseteq [d], |I|=e} \iota_I(a) \otimes \iota_{I^c}(b)$$

where $\iota_I : S^e V \to V^{\otimes I}$ stands for the composition of ι_e and an isomorphism $V^{\otimes e} \to V^{\otimes I}$ coming from a bijection $[e] \to I$. This shows the second inequality. \Box

Remark 2.1.2. The proposition readily implies that also $\underline{s}(\pi(t)) \leq \underline{\mathrm{prk}}(t)$ and $\mathrm{prk}(\iota(f)) \leq D \cdot \underline{s}(f)$.

Proposition 2.1.3. Fix d and m. Then Theorem 1.7.2 implies Theorem 1.5.3.

Proof. Assume Theorem 1.7.2 and consider

$$\underline{f} = (f_1, \dots, f_m) \in K[x_1, \dots, x_n]_d^m \cong (S^d V)^m.$$

Set $r := \underline{s}(\underline{f})$, so f lies in the Zariski closure of the set of m-tuples of degree-d forms over \overline{K} of collective strength $\leq r$. Applying ι over \overline{K} and using Proposition 2.1.1, we find that $\underline{t} := (\iota(f_1), \ldots, \iota(f_m))$ lies in the Zariski closure of the set of m-tuples of tensors over \overline{K} of collective partition rank $\leq Dr$. Hence $\operatorname{prk}(\underline{t}) \leq Dr$.

Under the assumptions of Theorem 1.5.3, d! is invertible in K, so that

$$s_K(f) = s_K(d!f) = s_K(\pi(\iota(f_1)), \dots, \pi(\iota(f_m))) \le \operatorname{prk}_K(\underline{t}) \le Q_m(Dr),$$

where the first inequality uses Proposition 2.1.1 (applied to the linear combination of $\iota(f_1), \ldots, \iota(f_m)$ of minimal partition rank) and the second inequality uses Theorem 1.7.2.

2.2. Behaviour under finite field extensions. We record a simple and well-known relation between the (collective) partition rank over K and that over a field extension of K.

Proposition 2.2.1. Let $\underline{t} \in (V_1 \otimes \cdots \otimes V_d)^m$ and let L be an extension of K with $e := [L : K] < \infty$. Then the collective partition rank satisfies $\operatorname{prk}_K(\underline{t}) \leq e \cdot \operatorname{prk}_L(ul_t)$.

Proof. Write

$$\sum_{k=1}^m c_k t_k = \sum_{i=1}^r a_i \otimes_L b_i$$

where $r := \operatorname{prk}_L(\underline{t})$, $a_i \in L \otimes_K V_{I_i}$, $b_i \in L \otimes_K V_{I_i^c}$, and $c_k \in L$, where the the c_k are not all zero. Assume, without loss of generality, that $c_m = 1$. Let z_1, \ldots, z_e be a basis of L over K. Then we can write $a_i = \sum_j z_j a_{ij}$ for certain tensors $a_{ij} \in V_{I_i}$. We then have

$$t_m + \sum_{k=1}^{m-1} c_k t_k = \sum_{i=1}^r \sum_{j=1}^e a_{ij} \otimes_L (z_j b_i).$$

The c_k and $z_j b_i$ are a solution over L to a certain system of linear equations with coefficients from K. This system then also has a solution over K, and such a solution witnesses $\operatorname{prk}_K(\underline{t}) \leq e \cdot r$.

Remark 2.2.2. The corresponding statement also holds for strength, with the same proof. The implication of our main theorems that the (collective) strength or partition rank over K can be bounded by the corresponding quantity over \overline{K} is therefore interesting only for fields K for which \overline{K} is not finite-dimensional over K, i.e., by the Artin-Schreier theorem, for fields that are not real closed or algebraically closed.

2.3. Subvarieties of tensor products. It is convenient, for the time being, to assume that K is infinite. We will later explain how the reasoning can be adjusted to finite fields. Let **Vec** be the category of finite-dimensional K-vector spaces, let d be a natural number, and denote by **Vec**^d the category whose objects and morphisms are d-tuples of objects and morphisms in **Vec**, respectively.

Definition 2.3.1. A functor $P : \mathbf{Vec}^d \to \mathbf{Vec}$ is polynomial of degree $\leq e$ if for any $V, W \in \mathbf{Vec}^d$ the map $P : \mathrm{Hom}(V, W) \mapsto \mathrm{Hom}(P(V), P(W))$ is polynomial of degree $\leq e$.

By a polynomial functor $\mathbf{Vec}^d \to \mathbf{Vec}$ we will always mean one that is polynomial of degree at most some e. In fact, the most interesting ones for this paper are the functors

$$T_I: \mathbf{Vec}^d \to \mathbf{Vec}, (V_1, \dots, V_d) \mapsto \bigotimes_{i \in I} V_i = V_I$$

where I is a subset of [d].

We will think of P(V) not just as a K-vector space, but also as the spectrum of the symmetric algebra of $P(V)^*$, an affine scheme of finite type over K. For $\varphi \in \operatorname{Hom}_{\operatorname{Vec}^d}(V, W)$ the linear map $P(\varphi)$ induces a morphism $P(V) \to P(W)$ of affine schemes over K.

Definition 2.3.2. A closed subvariety of a polynomial functor $P : \operatorname{Vec}^d \to \operatorname{Vec}$ is a rule X that assigns to every tuple $V \in \operatorname{Vec}^d$ a reduced, closed subscheme $X(V) \subseteq P(V)$ in such a manner that for any $\varphi : V \to W$ the map $P(\varphi)$ maps X(V) into X(W). We then write $X(\varphi)$ for the restriction $X(V) \to X(W)$ of the morphism $P(\varphi)$ to X(V).

Let X, Y be closed subvarieties of polynomial functors $\mathbf{Vec}^d \to \mathbf{Vec}$. A morphism $\alpha : X \to Y$ is the data of a morphism $\alpha_V : X(V) \to Y(V)$ of affine algebraic varieties over K for all $V \in \mathbf{Vec}^d$ in such a manner that for all $V, W \in \mathbf{Vec}^d$ and all $\varphi \in \mathrm{Hom}_{\mathbf{Vec}^d}(V, W)$ the following diagram commutes:

$$\begin{array}{c|c} X(V) & \xrightarrow{\alpha_V} & Y(V) \\ X(\varphi) & & & & \\ X(\psi) & & & & \\ X(W) & \xrightarrow{\alpha_W} & Y(W). \end{array}$$

The most important example of a closed subvariety for us lives in $T^m_{[d]}$ (*m*-tuples of tensors) and is defined by

$$X_r(V) := \{ \underline{t} \in T^m_{[d]}(V) \mid \operatorname{prk}(\underline{t}) \le r \};$$

the variety of *m*-tuples of tensors of border partition rank $\leq r$.

 \diamond

Definition 2.3.3. Given a tuple $U \in \mathbf{Vec}^d$ and a polynomial functor $P : \mathbf{Vec}^d \to \mathbf{Vec}$, we get a new polynomial functor $\mathrm{Sh}_U P : \mathbf{Vec}^d \to \mathbf{Vec}$, the *shift of* P over U, by setting

$$\operatorname{Sh}_U P(V) := P(U \oplus V) \text{ and } (\operatorname{Sh}_u P)(\varphi) := P(\operatorname{id}_U \oplus \varphi).$$

Furthermore, if X is a closed subvariety of P, then we define $\operatorname{Sh}_U X$ as $(\operatorname{Sh}_U X)(V) := X(U \oplus V) \subseteq (\operatorname{Sh}_U P)(V)$; this is a closed subvariety of $\operatorname{Sh}_U P$.

The following important example of a shift will be used intensively.

Example 2.3.4. Since tensor products distribute over direct sums, we have

$$(\operatorname{Sh}_U T_{[d]})(V) = \bigoplus_{I \subseteq [d]} U_{I^c} \otimes V_I,$$

so $\operatorname{Sh}_U T_{[d]}$ is a direct sum, over all subsets I of [d], of $\prod_{i \in I^c} (\dim(U_i))$ copies of T_I . Note that there is precisely one summand of degree d, namely, that with I = [d]; this summand is $U_{\emptyset} \otimes V_{[d]}$, which we identify with $V_{[d]} = T_{[d]}(V)$. The quotient $(\operatorname{Sh}_U T_{[d]})/T_{[d]}$ is a polynomial functor of degree d - 1.

Similarly, for any subset $I \subseteq [d]$, $\operatorname{Sh}_U T_I^m$ equals T_I^m plus a direct sum of copies of $T_{I'}^{m'}$ where $I' \subseteq I$ and where m' < m if I' = I.

Definition 2.3.5. Given a closed subvariety X of a polynomial functor P and a function $h \in K[X(0)]$, we may think of h as a function on every X(V) via the composition with the map $X(0_{V\to 0}) : X(V) \to X(0)$. We define the functor X[1/h] from Vec^d to affine K-schemes by sending V to the basic open subset of X(V) defined by the nonvanishing of h. This is a closed subset of the polynomial functor $V \mapsto K \oplus P(V)$ defined by the equations for $X(V) \subseteq P(V)$ and the equation $y \cdot h = 1$, where y is the coordinate on the affine line corresponding to the summand K.

3. Proof of Theorem 1.7.2 for infinite K

We continue to assume that K is infinite. We have

$$\Gamma^m_{[d]}(K^{n_1},\ldots,K^{n_d}) = (K^{n_1} \otimes \cdots \otimes K^{n_d})^m.$$

In this space we have the standard basis vectors

$$e_{(i_1,\ldots,i_d),\ell} := (0,\ldots,0,e_{i_1}\otimes\cdots\otimes e_{i_d},0,\ldots)$$

where the nonzero entry is in position $\ell \in [m]$ and where the e_{i_j} are the standard basis vectors in K^{n_j} . We write $x_{(i_1,\ldots,i_d),\ell}$ for the corresponding standard coordinates.

The following theorem is a summary of results proved in [BDE19, §4], but generalised from tensors to *m*-tuples of tensors. Informally, the theorem says that on an open subset where a partial derivative of a defining function for a closed subset $X \subseteq T_{[d]}^m$ is nonzero, the coordinates of the *m*-th tensor can be reconstructed from those of the remaining tensors.

Theorem 3.0.1. Let $X \subseteq T_{[d]}^m$ be a closed subvariety and assume that X is defined over the prime field of K in the following sense: for all n_1, \ldots, n_d , $X(K^{n_1}, \ldots, K^{n_d}) \subseteq (K^{n_1} \otimes \cdots \otimes K^{n_d})^m$ is the zero set of polynomials in the standard coordinates with coefficients from the prime field F of K. Let $U = (U_1, \ldots, U_d) \in \mathbf{Vec}^d$ with $U_j = K^{n_j}$ be such that $X(U) \subsetneq (T_{[d]}(U))^m$. Let f be a nonzero polynomial with coefficients in F that vanishes identically on X(U). Let x_1 be the standard coordinate $x_{(1,\ldots,1),m}$ and set

$$h := \frac{\partial f}{\partial x_1}.$$

Let M be the m-th copy of $T_{[d]}$ in $T_{[d]}^m$. It is then naturally a summand of $\operatorname{Sh}_U T_{[d]}^m$, and the projection $\operatorname{Sh}_U T_{[d]}^m \to (\operatorname{Sh}_U T_{[d]}^m)/M$ restricts to a closed embedding, defined over F,

$$\psi: (\operatorname{Sh}_U X)[1/h] \to ((\operatorname{Sh}_U T^m_{[d]})/M)[1/h].$$

Furthermore, there exists a morphism, defined over F,

$$\sigma: ((\operatorname{Sh}_U T^m_{[d]})/M)[1/h] \to (\operatorname{Sh}_U T^m_{[d]})[1/h]$$

such that $\sigma \circ \psi$ is the identity on $(\operatorname{Sh}_U X)[1/h]$.

Corollary 3.0.2. In the setting of Theorem 3.0.1, for any $V \in \mathbf{Vec}^d$ and any K-point \underline{t} of X(V) for which there exists an m-tuple $\varphi: V \to U$ of K-linear maps with $h(T^m_{\mathrm{fd}}(\varphi)\underline{t}) \neq 0$, the collective partition rank $\mathrm{prk}_K(\underline{t})$ of \underline{t} is at most

(3)
$$m \cdot \sum_{I \subseteq [d]: 1 \le \lfloor d/2 \rfloor} \prod_{j \in I^c} n_j$$

Proof. Let $\underline{\tilde{t}}$ be the image of \underline{t} in $T^m_{[d]}(U \oplus V)$ under the linear map $T^m_{[d]}(\varphi \oplus \mathrm{id}_V)$. So the *i*-th component of $\underline{\tilde{t}}$ equals $t_i \in V_{[d]}$ plus a remainder in

$$\bigoplus_{I \subsetneq [d]} U_{I^c} \otimes V_I = \sum_{i=1}^d U_i \otimes \bigotimes_{j \neq i} (U_j \oplus V_j).$$

We analyse t_m , which is the part of \tilde{t}_m in M.

We have

$$h(\tilde{t}) = h(T^m_{[d]}(\varphi)\underline{t}) \neq 0,$$

so that we may regard $\underline{\tilde{t}}$ as a *K*-point of $(\operatorname{Sh}_U X)(V)[1/h]$. Therefore, Theorem 3.0.1 yields the equality $\tilde{t} = \sigma(\psi(\tilde{t}))$. The *M*-component of σ is a morphism $((\operatorname{Sh}_U T^m_{[d]})/M)[1/h] \to M = T_{[d]}$ and hence, by basic representation theory for $\prod_i \operatorname{GL}(V_i)$, a linear combination (with coefficients in $K[T^m_{[d]}(U)][1/h]$) of morphisms that take the components in some summands T_{I_1}, \ldots, T_{I_k} for which $I_1 \sqcup \cdots \sqcup I_k = [d]$ and returns their tensor product. (See the paragraph [BDE19, §4.8] on covariants for details.)

Among these morphisms, those with k = 1 are isomorphisms from the m - 1 copies of $T_{[d]}$ in $\operatorname{Sh}_U T_{[d]}^m/M$; this is where t_1, \ldots, t_{m-1} live. So for suitable $c_1, \ldots, c_{m-1} \in K, t' := t_m + \sum_{i=1}^{m-1} c_i t_i$ is a linear combination of tensors in the images of those morphisms with $k \geq 2$. For these, the smallest I_j has cardinality at most $\lfloor \frac{d}{2} \rfloor$. So we find that t' has partition rank at most the number of (copies of) T_I with $|I| \leq \lfloor \frac{d}{2} \rfloor$ in $\operatorname{Sh}_U T_{[d]}^m$, which is

$$m \cdot \sum_{I \subseteq [d]: 1 \le |I| \le \lfloor d/2 \rfloor} \prod_{j \in I^c} n_j.$$

Thus the collective partition rank of \underline{t} is bounded by the number above.

We record the following corollary of the proof.

Definition 3.0.3. A proper contraction of $\underline{t} \in T_{[d]}(V)^m$ is any tensor of the form

$$\sum_{i=1}^{m} \xi_i(t_i) \in T_I(V),$$

 \diamond

where I is a nonempty proper subset of [d], and $\xi_i \in T_{I^c}(V^*)$ for all i.

Corollary 3.0.4. There exists an N such that for any \underline{t} as in Corollary 3.0.2 there is a K-linear combination $t' = t_m + \sum_{i=1}^{m-1} c_i t_i$ which is contained in a subalgebra of $T(V) = \bigoplus_{e=0}^{\infty} T_{[d]}(V)$ generated by $\leq \sum_{j=1}^{d} n_j + m \cdot \prod_{j \in [d]} (n_j + 1)$ proper contractions of \underline{t} .

Such a tensor product is of the form $s_1 \otimes \cdots \otimes s_e$ where $e \geq 2$, $s_i \in T_{I_i}(V)$ and I_1, \ldots, I_e form a partition of [d] into nonempty sets.

Proof. Without loss of generality, \underline{t} is *concise*: each V_j is the smallest space $V'_j \subseteq V_j$ with $\underline{t} \in T^m_{[d]}(V_1, \ldots, V_{j-1}, V'_j, V_{j+1}, \ldots, V_d)$. Then each element of V_j is a proper contraction of \underline{t} .

Let V'_j be the kernel of φ_j and let U'_j be a complement of V'_j in V_j . Then the restriction of φ to U' is injective, and we may regard this as an inclusion, so that $U'_j \subseteq U_j$ for all j and \underline{t} lies in $T^m_{[d]}(U' \oplus V') \subseteq T^m_{[d]}(U \oplus V')$. We will treat \underline{t} as we treated $\underline{\tilde{t}}$ in the proof of Corollary 3.0.2. Consider the linear combination $t' := t_m + \sum_{i=1}^{m-1} c_i t_i$ constructed in Corollary 3.0.2. We want to show that this admits a decomposition as desired.

Choose a basis of each U_j containing a basis of U'_j . Each t_i admits a unique decomposition

$$t_i = \sum_u u \otimes t_{i,u}$$

where u runs over all $\prod_{j=1}^{d} (1+n_j)$ "monomials" in the chosen bases of the U_j : each lives in a tensor product $\bigotimes_{j \in I} U_j$ for some subset $I \subseteq [d]$, while its coefficient $t_{i,u}$ lives in the tensor product $\bigotimes_{j \in I^c} V'_j$. From the fact that \underline{t} lies in $T^m_{[d]}(U' \oplus V')$ it follows that that $t_{i,u}$ is zero whenever u involves basis elements of some U_j that are not in U'_j . So in all nonzero terms above, u is contained in the subalgebra generated by the union of the bases of U'_j , which consists of $\leq \sum_{j=1}^d n_j$ proper contractions of \underline{t} .

Now consider a monomial $u \in \bigotimes_{j \in I} U'_j$ of positive degree. Then inclusion-exclusion yields:

$$t_{i,u} = \sum_{u',u'':u'=u\otimes u''} (-1)^{|u'|-|u|} \xi_{u'}(t_i) \otimes u''$$

where the sum is over all monomials u' that are multiples of u, hence $u' \in \bigotimes_{j \in I'} U'_j$ with $I' \supseteq I$, and $\xi_{u'}$ is a linear form on the space $\bigotimes_{j \in I'} (U'_j \oplus V'_j)$ that is 1 on u', zero on any other monomial in $\bigotimes_{j \in I'} U'_j$, and zero on any tensor product of vectors containing a vector in some V'_j with $j \in I'$. The number of contractions $\xi_{u'}(t_i)$ appearing in these expressions for any i and u of positive degree is $\leq m \cdot \prod_{j \in [d]} (n_j + 1)$.

10

Now, t' has a unique decomposition like the $t_{i,u}$:

$$t' = \sum_u u \otimes t'_u,$$

where each t'_{u} is a linear combination of the $t_{i,u}$, and hence is contained in the subalgebra we have so far described, as long as u has positive degree. On the other hand, in the proof of Corollary 3.0.2 we saw that $t'_u = t'_1$, where u = 1 is the unique degree-0 monomial, is a sum of tensor products of the $t_{i,u}$ for monomials uof positive degree. This completes the proof.

Proposition 3.0.5. The closed subvariety $X_r(V) := \{\underline{t} \in T^m_{[d]}(V) \mid \underline{\mathrm{prk}}(\underline{t}) \leq r\}$ of $T_{[d]}^m$ satisfies the conditions of Theorem 3.0.1 with $n_1 = \ldots = n_d = n$ where n is minimal such that

(4)
$$n^d > m^2 + r(n^{d-1} + n)$$

Proof. For any choice of r proper nonempty subsets I_1, \ldots, I_r of [d], each of size $\leq \lfloor d/2 \rfloor$, and $U := (K^n, \ldots, K^n)$ consider the parameterisation

(5)
$$K^{m \times m} \times T^{m-1}_{[d]}(U) \times \prod_{i=1}^{r} \left(T_{I_i}(U) \times T_{I_i^c}(U) \right) \to T^m_{[d]}(U),$$
$$(g, (t_1, \dots, t_{m-1}), ((a_i, b_i))_i) \mapsto g \cdot (t_1, \dots, t_{m-1}, \sum_i a_i \otimes b_i)$$

which is the tensor analogue to the map in (2). The locus of tensor tuples of collective partition rank at most r is the union of the images of these parameterisations over all r-tuples (I_1, \ldots, I_r) as above. Since the parameterisation is defined over the prime field, X is defined over the prime field. The dimension of $V_{I_i} \times V_{I_i^c}$ is at most $n^{d-1} + n$, with equality if I_i or I_i^c is a

singleton. Hence the left-hand side in the parameterisation has dimension at most

$$(m-1)n^d + m^2 + r(n^{d-1} + n) < mn^d.$$

By inequality (4), the parameterisation is not dominant into $T^m_{[d]}(U)$, hence there exists a nonzero polynomial F that vanishes on the image. Taking the product of these Fs for the finitely many choices of (I_1, \ldots, I_r) gives a nonzero polynomial vanishing on all of $X_r(U)$. \square

We note the crucial fact that for d fixed, the lower bound on n imposed by (4) is linear in $r + m^{2/d}$. We can now prove Theorem 1.7.2 in the case where K is infinite.

Proof of Theorem 1.7.2 when K is infinite. By Proposition 3.0.5, $X_r(U) \subsetneq T^m_{[d]}(U)$ for $U_i := K^n, i = 1, ..., d$ with n minimal such that (4) holds. We claim that, for any $V \in \mathbf{Vec}^d$, the collective partition rank $\operatorname{prk}_K(\underline{t})$ of any K-point \underline{t} of $X_r(V)$ is at most

(6)
$$m \cdot \sum_{e=1}^{\lfloor \frac{d}{2} \rfloor} \binom{d}{e} n^{d-e}.$$

Since n is linear in $r + m^{2/d}$, this expression is $\ll_d m^3 r^{d-1}$.

We prove the claim with X_r replaced by any \mathbf{Vec}^d -subvariety Y of X_r that is defined over the prime field F of K. This has the advantage that we can do induction on the minimal degree $\delta(Y)$ of a nonzero polynomial vanishing on Y(U): when proving it for Y, we may assume that the claim holds for all Vec^{d} -subvarieties Y' of X_r for which $\delta(Y') < \delta(Y)$.

Thus let $f \in K[T^m_{[d]}(U)]$ be a nonzero polynomial, with coefficients in F, of minimal degree that vanishes on Y. If f is constant, then Y is empty and the claim holds trivially. Otherwise, if $\operatorname{char}(K) = p$, then since f is of minimal degree, it is not a p-th power of a polynomial. Hence in some monomial in f, some variable has an exponent that is not divisible by p. Without loss of generality, this variable is the variable x_1 in Theorem 3.0.1. Construct construct h as in that theorem. By construction, h is nonzero and has smaller degree than f.

Let $Y' \subseteq Y$ be the **Vec**^d-subvariety defined by

(7)
$$Y'(V) := \{ \underline{t} \in Y(V) \mid \forall \varphi \in \operatorname{Hom}(V, U) : h(T^m_{[d]}(\varphi)\underline{t}) = 0 \}.$$

This is again defined over F. Indeed, instead of letting φ run over all of $\operatorname{Hom}(V, U)$, we can take for φ a tuple of matrices with entries that are variables, expand $h(T^m_{[d]}(\varphi)\underline{t})$ as a polynomial in these variables, and take the coefficients; these define Y'—we use here that K is infinite.

On Y' the nonzero polynomial h of strictly lower degree than $\deg(f)$ vanishes, so that $\delta(Y') < \delta(Y)$. Hence the claim holds for Y' by the induction hypothesis. On the other hand, for <u>t</u> a K-point of $Y'(V) \setminus Y(V)$ we can apply Corollary 3.0.2 to conclude that <u>t</u> has collective partition rank at most (6).

4. Proof of Theorem 1.7.2 for finite K

In this section we assume that char K = p > 0 and drop the requirement that K is infinite. We consider the affine scheme $X_r \subseteq T^m_{[d]}$ whose \overline{K} -points are the tuples of border collective partition rank $\leq r$. Let $U_i = K^n$ where n is such that (4) holds and let f be a nonzero polynomial, with coefficients in the prime field, of minimal degree in the ideal of $X_r(U)$. Assume that x_1 appears in some monomial in f and its exponent in that monomial is not divisible by p, so that $h := \frac{\partial f}{\partial x_n}$ is nonzero.

its exponent in that monomial is not divisible by p, so that $h := \frac{\partial f}{\partial x_1}$ is nonzero. If \underline{t} is a K-point of $X_r(V)$ for which there exists a d-tuple $\varphi \in \operatorname{Hom}(V, U)$ of K-linear maps such that $h(T^m_{[d]}(\varphi)\underline{t}) \neq 0$, then Corollary 3.0.2 still yields the desired upper bound on $\operatorname{prk}_K(\underline{t})$ —indeed, we may extend K to an infinite field L to apply that result, and we use that the morphisms ψ, σ from Theorem 3.0.1 are defined over the prime field, hence over K.

However, what if \underline{t} is a K-point of $X_r(V)$ such that $h(T^m_{[d]}(\varphi)\underline{t}) = 0$ for all $\varphi \in \operatorname{Hom}(V, U)$? If K is finite, then this does not a priori imply that $h(T^m_{[d]}(\varphi)\underline{t}) = 0$ holds for all $\varphi \in \operatorname{Hom}_L(L \otimes V, L \otimes U)$, and so \underline{t} is perhaps not a point in the proper subvariety $Y' \subsetneq X_r$ defined in (7).

To nevertheless obtain a bound on $\operatorname{prk}_{K}(\underline{t})$, we proceed in this case as follows. We find a field L and a $\varphi \in \operatorname{Hom}_{L}(L \otimes V, L \otimes U)$ such that $h(T^{m}_{[d]}(\varphi)\underline{t}) \neq 0$. Then Corollary 3.0.2 gives an upper bound on the collective partition rank $\operatorname{prk}_{L}(\underline{t})$. Finally, we use Proposition 2.2.1 to derive an upper bound on $\operatorname{prk}_{K}(\underline{t})$.

For this to work, we need that L is large enough. Observe that $h(T^m_{[d]}(\varphi)\underline{t})$ has degree $d \cdot \deg(h)$ in the entries of φ . If $|L| > d \cdot \deg(h)$, then a $\varphi \in \operatorname{Hom}_L(L \otimes V, L \otimes U)$ exists for which this is nonzero, unless \underline{t} is a K-point of Y', in which case we can invoke the induction hypothesis.

Hence it suffices to take for L an extension of degree greater than $\log_2(d \cdot \deg(h))$ —the base 2 rather than |K| ensures that we obtain a bound that is independent on the finite field. In what follows, log stands for \log_2 .

Proposition 4.0.1. For d fixed and r sufficiently large, $n = 4(r + m^{2/d})$ satisfies (4). Choose this n and set $U := (K^n, \ldots, K^n)$. For any fixed parametrisation (5), let $X_{I_1,\ldots,I_r} \subset T^m_{[d]}$ be the image. Then the minimal degree D of a nonzero polynomial f that vanishes identically on $X_{I_1,\ldots,I_r}(U)$ satisfies

$$\log(D) \ll_d \log(r+m).$$

Proof. The first statement is immediate. Now consider the parameterisation in (5), expanded in the standard bases of these spaces. Along the pull-back of this map, a coordinate on the space $T^m_{[d]}(U)$ is mapped to a polynomial each of whose monomials is of degree ≤ 3 in the coordinates of g and the t_i, a_i, b_i (actually, these monomials have more structure, but we ignore this). Hence a polynomial of degree $\leq 3D$ on $T^m_{[d]}(U)$ is mapped to a polynomial each of whose monomials is of degree to a polynomial each of whose monomials is of degree $\leq 3D$ in the coordinates on the left-hand side. The number of such monomials is at most

$$\binom{m^2 + (m-1)n^d + r(n^{d-1} + n) + 3D}{3D} \le \binom{(m - \frac{1}{4})n^d + 3D}{(m - \frac{1}{4})n^d} =: A$$

We determine D such that this pullback is guaranteed to have a nontrivial kernel. For this, it suffices that there are more degree- $\leq D$ monomials in $T^m_{[d]}(U)$ than the expression above. Explicitly, it suffices that

$$\binom{mn^d + D}{mn^d} > A.$$

We use the well-known bounds

$$\left(\frac{N}{k}\right)^k \le \binom{N}{k} \le \left(\frac{eN}{k}\right)^k$$

to find sufficient conditions on D. By the right-hand bound and choosing $D \ge (m - \frac{1}{4})n^d$ we find

$$A \le \left(\frac{e((m-\frac{1}{4})n^d + 3D)}{(m-\frac{1}{4})n^d}\right)^{(m-\frac{1}{4})n^d} \le \left(\frac{4eD}{(m-\frac{1}{4})n^d}\right)^{(m-\frac{1}{4})n^d}$$

On the other hand, using the lower bound for binomial coefficients, we find

$$\binom{mn^d + D}{mn^d} \ge \left(\frac{mn^d + D}{mn^d}\right)^{mn^a} \ge \left(\frac{D}{mn^d}\right)^{mn^d}.$$

Taking logarithms, we find that it suffices that

$$\log(D) \cdot \frac{1}{4}n^{d} \ge mn^{d}\log(mn^{d}) + (m - \frac{1}{4})n^{d}\log\left(\frac{4e}{(m - \frac{1}{4})n^{d}}\right).$$

This yields the sufficient condition

$$\log(D) \ge 4\log(mn^d) + 4(m - \frac{1}{4})\log\left(\frac{3e}{(m - \frac{1}{3})n^d}\right)$$

We see that we can take $\log(D)$ a suitable constant multiple of $\log(r+m)$, as desired.

Proof of Theorem 1.7.2 for finite K. Any $\underline{t} \in X_r(V)$ lies in X_{I_1,\ldots,I_r} for some choice of proper, nonempty subsets $I_1, \ldots, I_r \subset [d]$. By Proposition 4.0.1, choosing $n = 4(r + m^{2/d})$ for r sufficiently large and setting $U := (K^n, \ldots, K^n)$, the minimal degree δ of a nonzero polynomial vanishing on all of $X_{I_1,\ldots,I_r}(U)$ satisfies $\log(\delta) \ll_d \log(r + m)$. This then also holds for the **Vec**^d-subvarieties $Y \subseteq X_r$ used in the proof of Theorem 1.7.2 for infinite fields. By the discussion preceding the proof of Proposition 4.0.1, for any K-point $\underline{t} \in Y(V)$ we have the following dichotomy: either \underline{t} lies in the proper subvariety Y'(V) defined by (7), so that the induction hypothesis applies; or else there exists an extension L of K with $[L:K] \ll_d \log(r+m)$ such that $\operatorname{prk}_L(\underline{t})$ is at most the polynomial Q(r,m) from (6) (defined using $n = 4(r + m^{2/d})$). Then Proposition 2.2.1 implies that

$$\operatorname{prk}_{K}(\underline{t}) \ll_{d} m^{3} r^{d-1} \log(r+m),$$

as desired.

5. Proof of Theorem 1.3.1

We recall that Theorem 1.3.1 asks for an expression of f as a polynomial in a bounded number of elements of the space D(f).

Proof of Theorem 1.3.1. First we assume that K is infinite and let f be a K-point of the variety of forms of border strength $\leq r$ in S^dV , where V is an n-dimensional vector space. Then $t := \iota(f) \in V^{\otimes d}$ is a tensor of border partition rank $\leq Dr$, where D is as in Proposition 2.1.1. By Corollary 3.0.4, t is contained in a subalgebra generated by $\ll_d r^d$ proper contractions. This uses the same inductive reasoning as used in the proof of Theorem 1.7.2.

Given a proper contraction $s_j \in T_{I_j}(V)$, a straightforward computation shows that $h_j := \pi_{I_j}(s_j) \in S^{|I_j|}V$ is an element of D(f) and that $f = \frac{1}{d!}\pi_d(t)$ is contained in the subalgebra generated by the $\pi_{I_j}(s_j)$. This concludes the proof when K is infinite.

When K is finite, as in the proof of Theorem 1.7.2 over finite fields, we extend K to a field L such that $[L:K] \ll_d \log r$ and we find that f lies in an algebra generated by $\ll_d r^d$ elements of $L \otimes_K D(f)$. As in Proposition 2.2.1, this implies that f is in an algebra generated by $\ll_d r^d \log r$ elements of D(f).

References

- [AH20] Tigran Ananyan and Melvin Hochster. Small subalgebras of polynomial rings and Stillman's conjecture. J. Am. Math. Soc., 33(1):291–309, 2020.
- [AKZ21] Karim Adiprasito, David Kazhdan, and Tamar Ziegler. On the schmidt and analytic ranks for trilinear forms. 2021. Preprint, arXiv:2102.03659.
- [BBOV22] Edoardo Ballico, Arthur Bik, Alessandro Oneto, and Emanuele Ventura. The set of forms with bounded strength is not closed. C. R., Math., Acad. Sci. Paris, 360:371– 380, 2022.
- [BDE19] Arthur Bik, Jan Draisma, and Rob H. Eggermont. Polynomials and tensors of bounded strength. *Commun. Contemp. Math.*, 21(7), 2019. paper number 1850062 (24 pages).
- [BDS24a] Arthur Bik, Jan Draisma, and Andrew Snowden. The geometry of polynomial representations in positive characteristic. Math. Z., 2024. To appear, arXiv:2406.07415.
- [BDS24b] Arthur Bik, Jan Draisma, and Andrew Snowden. Two improvements in Brauer's theorem on forms. 2024. Preprint, arXiv:2401.02067.
- [BL24] Benjamin Baily and Amichai Lampert. Strength, partition rank and algebraic closure. 2024. Preprint, arXiv:2410.00248.

- [CM23] Alex Cohen and Guy Moshkovitz. Partition and analytic rank are equivalent over large fields. Duke Math. J., 172(12):2433–2470, 2023.
- [CY24] Qiyuan Chen and Ke Ye. Stability of ranks under field extensions. 2024. Preprint, arXiv:2409.04034.
- [DES17] Harm Derksen, Rob H. Eggermont, and Andrew Snowden. Topological noetherianity for cubic polynomials. Algebra Number Theory, 11(9):2197–2212, 2017.
- [DGI+22] Pranjal Dutta, Fulvio Gesmundo, Christian Ikenmeyer, Gorav Jindal, and Vladimir Lysikov. De-bordering and geometric complexity theory for waring rank and related models. 2022. Preprint, arXiv:2211.07055.
- [DK24] Jan Draisma and Thomas Karam. On subtensors of high partition rank. Proc. Am. Math. Soc., 152(12):5083–5093, 2024.
- [GK22] W. T. Gowers and Thomas Karam. Equidistribution of high-rank polynomials with variables restricted to subsets of \mathcal{O}_p . Preprint, arXiv:2209.04932 [math.CO] (2022), 2022.
- [GT09] Ben Green and Terence Tao. The distribution of polynomials over finite fields, with applications to the Gowers norms. *Contrib. Discrete Math.*, 4(2):1–36, 2009.
- [Kar22] Thomas Karam. High-rank subtensors of high-rank tensors. 2022. Preprint, arXiv:2207.08030.
- [KP23] David Kazhdan and Alexander Polishchuk. Schmidt rank of quartics over perfect fields. Isr. J. Math., 255(2):851–869, 2023.
- [Lov19] Shachar Lovett. The analytic rank of tensors and its applications. Discrete Anal., 2019:10, 2019. Id/No 7.
- [LZ24] Amichai Lampert and Tamar Ziegler. On rank in algebraic closure. Sel. Math., New Ser., 30, 2024. paper number 15.
- [Mil19] Luka Milićević. Polynomial bound for partition rank in terms of analytic rank. Geom. Funct. Anal., 29(5):1503–1530, 2019.
- [MZ22] Guy Moshkovitz and Daniel G. Zhu. Quasi-linear relation between partition and analytic rank. 2022. Preprint, arXiv:2211.05780.
- [Nas20a] Eric Naslund. Exponential bounds for the Erdős-Ginzburg-Ziv constant. J. Comb. Theory, Ser. A, 174:18, 2020. Id/No 105185.
- [Nas20b] Eric Naslund. The partition rank of a tensor and k-right corners in \mathbb{F}_q^n . J. Comb. Theory, Ser. A, 174:24, 2020. Id/No 105190.
- [Sch85] Wolfgang M. Schmidt. The density of integer points on homogeneous varieties. Acta Math., 154:243–296, 1985.

D.E. Shaw & Co., Floor 52, Two Manhattan West, 375 Ninth Avenue New York, 10001 NY, USA

Email address: arthur.bik@deshaw.com

Mathematical Institute, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzer-Land; and Department of Mathematics and Computer Science, Eindhoven University of Technology, P.O. Box 513, 5600MB, Eindhoven, the Netherlands

Email address: jan.draisma@unibe.ch

Mathematics Department, University of Michigan, 530 Church Street, Ann Arbor, MI 48109-1043, USA

Email address: amichai@umich.edu

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT RAM. JERUSALEM, 9190401, ISRAEL

Email address: tamar.ziegler@mail.huji.ac.il