

HIGHER SCHWARZIAN, QUASIMODULAR FORMS AND EQUIVARIANT FUNCTIONS

HICHAM SABER AND ABDELLAH SEBBAR

ABSTRACT. The Schwarzian derivative plays a fundamental role in complex analysis, differential equations, and modular forms. In this paper, we investigate its higher-order generalizations, known as higher Schwarzians, and their connections to quasimodular forms and equivariant functions. We prove that a meromorphic function is equivariant if and only if its higher Schwarzians are quasimodular forms of prescribed weight and depth, thereby extending classical results and linking projective differential operators to the structure of modular and quasimodular forms.

1. INTRODUCTION

The Schwarzian derivative defined by

$$\{f, z\} = \frac{f'''(z)}{f'(z)} - \frac{3}{2} \left(\frac{f''(z)}{f'(z)} \right)^2$$

is a fundamental operator in complex analysis and differential equations, with deep connections to projective geometry, univalent functions, and modular forms. It plays a crucial role in the study of conformal mappings, Teichmüller theory, and the monodromy of second-order differential equations. Over time, various generalizations of the Schwarzian derivative, known as higher Schwarzians, have been introduced to extend its applicability to broader mathematical contexts.

In this work, We study the Aharonov invariants of a meromorphic function on a simply connected domain, denoted $S_n[f]$, $n \geq 1$, which are referred to as higher Schwarzians. Notably, the first two cases recover well-known operators: $2S_1[f] = f''/f'$ known as the pre-Schwarzian, and $6S_2[f] = \{f, z\}$, the classical Schwarzian derivative.

Parallel to this, the theory of modular and quasimodular forms has become central in number theory, arithmetic geometry, and mathematical physics. Quasimodular

2010 *Mathematics Subject Classification.* 11F03, 11F11, 34M05.

Key words and phrases. Schwarz derivative, Modular forms, Quasimodular forms, Equivariant functions, representations of the modular group.

forms, introduced by Kaneko and Zagier, extend classical modular forms by incorporating derivatives, leading to a richer algebraic structure that remains closed under differentiation.

A key perspective on modular forms involves their transformation properties under discrete subgroups of $\mathrm{PSL}_2(\mathbb{R})$. Similarly, equivariant functions satisfy specific transformation rules under discrete group actions. Given a discrete group Γ acting on the upper half-plane \mathbb{H} , a meromorphic function is equivariant if it commutes with the action of Γ . More generally, if f is a meromorphic function on \mathbb{H} , and $\rho : \Gamma \rightarrow \mathrm{GL}_2(\mathbb{C})$ is a two-dimensional complex representation, then f is called ρ -equivariant for if

$$f(\gamma z) = \rho(\gamma)f(z), \quad \text{for all } \gamma \in \Gamma \text{ and all } z \in \mathbb{H}.$$

It turns out that a function f is ρ -equivariant for some representation ρ of Γ if and only if the Schwarzian derivative $\{f, z\}$ is a weight 4 meromorphic modular form for Γ . Equivariant functions naturally arise in the study of monodromy representations, vector bundles over modular curves, and modular differential equations, providing a unifying framework for understanding the role of higher Schwarzians in relation to projective differential operators and modular forms.

In this paper, we explore the deep connections between higher Schwarzians, quasimodular forms, and equivariant functions. We show that higher Schwarzians inherently exhibit quasimodular behavior, extending classical results on the Schwarzian derivative. Specifically, we prove that a meromorphic function on \mathbb{H} is ϱ -equivariant for some representation ϱ of a discrete group Γ if and only if its higher Schwarzians $S_n[f]$, for $n \geq 2$, are quasimodular forms for Γ of weight $2n$ and depth $n - 2$.

The paper is organized as follows: In Section 1, we review the classical Schwarzian derivative and introduce its higher-order generalizations following Aharonov's construction. Section 2 provides an overview of quasimodular forms and their fundamental properties, emphasizing their role as a generalization of modular forms. In Section 3, we introduce equivariant functions and explore their connections to modular transformations and representation theory. Finally, we present our main results on the quasimodular nature of higher Schwarzians.

2. HIGHER SCHWARZIAN

Let D be a domain in the extended complex plane $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. If f is a meromorphic function on D , the Schwarz derivative (or the Schwarzian) of f is defined as

$$(2.1) \quad \{f, z\} = \left(\frac{f''(z)}{f'(z)} \right)' - \frac{1}{2} \left(\frac{f''(z)}{f'(z)} \right)^2 = \frac{f'''(z)}{f'(z)} - \frac{3}{2} \left(\frac{f''(z)}{f'(z)} \right)^2.$$

This is the lowest-order differential operator that is projectively invariant. Specifically, if $g(z) = \frac{af(z)+b}{cf(z)+d}$ with $ad - bc \neq 0$, then $\{f, z\} = \{g, z\}$. This operator was named

after Schwarz by Cayley; however, Schwarz himself credited its discovery to Lagrange in 1781. It also appeared in a paper by Kummer in 1836 [12].

The Schwarz derivative is fundamental in various areas of mathematics, including the study of the complex projective line, univalent functions, conformal mappings, Teichmüller spaces, and the theories of automorphic and hypergeometric functions [2, 3, 5, 10, 11, 14]. Furthermore, it has a profound connection to the theory of ordinary differential equations, as outlined below.

Let $R(z)$ be a meromorphic function on D and consider the second order differential equation

$$(2.2) \quad y'' + R(z)y = 0$$

with two linearly independent solutions y_1 and y_2 . Then $f = y_1/y_2$ satisfies the Schwarzian differential equation

$$(2.3) \quad \{f, z\} = 2R(z).$$

This relationship with second-order ordinary differential equations underpins many key properties of the Schwarz derivative which we now give:

- (1) Cocycle property: If w is a function of z , then

$$\{f, z\} = \{f, w\}(dw/dz)^2 + \{w, z\},$$

which says that $\{f, z\}$ behaves as a quadratic differential

$$\{f, z\}dz^2 = \{f, w\}dw^2 + \{w, z\}dz^2.$$

- (2) We have

$$\{f, z\} = 0 \text{ if and only if } f(z) = \frac{az + b}{cz + d} \text{ for some } a, b, c, d \in \mathbb{C}$$

- (3) If $w = \frac{az + b}{cz + d}$ with $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{C})$, then

$$\{f, z\} = \{f, w\} \frac{(ad - bc)^2}{(cz + d)^4}.$$

- (4) For two meromorphic functions f and g on D ,

$$\{f, z\} = \{g, z\} \text{ if and only if } f(z) = \frac{ag(z) + b}{cg(z) + d} \text{ for some } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{C}).$$

- (5) If $w(z)$ is a function of z with $w'(z_0) \neq 0$ for some $z_0 \in D$, then locally near z_0 , we have:

$$\{z, w\} = \{w, z\} (dz/dw)^2.$$

Higher order analogues of the Schwarz derivative that enjoy similar properties than the ordinary Schwarzian have been studied in the literature, notably those defined by Aharonov [1], Tamanoi [16] and Kim and Sugawa [8] among others. In this paper, we will follow Aharonov's construction which we now introduce.

Let f be a meromorphic function on a domain D and $z \in D$ be a point where f is holomorphic and $f'(z) \neq 0$. Define

$$G(w, z) = \frac{f'(z)}{f(w) - f(z)}.$$

As a function of w , the function $G(w, z)$ has a simple pole at $w = z$, and near $w = z$, it admits the expansion:

$$(2.4) \quad G(w, z) = \frac{1}{w - z} - \sum_{n=1}^{\infty} S_n[f](z)(w - z)^{n-1}.$$

The coefficients $S_n[f](z)$, referred to as the Aharonov invariants of f , exhibit several interesting properties shown in [1]. Differentiating $G(w, z)$, with respect to z , we see that G satisfies a Riccati differential equation:

$$(2.5) \quad \frac{\partial G}{\partial z} = G(z)^2 + \frac{f''(z)}{f'(z)} G(z).$$

Using the expansion (2.4) and equating the coefficients of $(w - z)^{-1}$, we get

$$S_1[f](z) = \frac{f''(z)}{2f'(z)},$$

known as the pre-Schwarzian, that is used to construct the Schwarz-Christoffel mapping between the upper half-plane and a polygonal domain. Now, equating the constant term in $w - z$, we obtain:

$$6S_2[f](z) = \{f, z\},$$

the usual Schwarzian.

Finally, equating the coefficients of positive powers of $w - z$ shows that the higher Schwarzians satisfy the recurrence relation:

$$(2.6) \quad (n + 1)S_n[f] = S'_{n-1}[f] + \sum_{k=2}^{n-2} S_k[f]S_{n-k}[f], \quad n \geq 3.$$

Differentiating (2.4) with respect to w , we get

$$(2.7) \quad \frac{f'(w)f'(z)}{(f(w) - f(z))^2} = \frac{1}{(w - z)^2} + \sum_{n=1}^{\infty} (n - 1)S_n[f](z)(w - z)^{n-2}.$$

Since the quantity

$$\frac{f'(w)f'(z)}{(f(w) - f(z))^2}$$

is invariant under a linear fractional transformation of f , and by the uniqueness of the Laurent series expansion around $w = z$, we conclude that for $n \geq 2$ and $ad - bc \neq 0$, we have:

$$S_n \left[\frac{af(z) + b}{cf(z) + d} \right] = S_n[f].$$

This projective invariance of S_n (for $n \geq 2$) justifies calling them higher Schwarzians.

3. QUASIMODULAR FORMS

From now on, we consider the domain D to be the upper half of the complex plane, $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$. Let Γ be a discrete subgroup of $\text{SL}_2(\mathbb{R})$, that is a Fuchsian group of the first kind, acting on \mathbb{H} via linear fractional transformations.

The notion of quasimodular forms was introduced by Kaneko and Zagier in [7] as a generalization of classical modular forms that includes all derivatives of modular forms. Following [17] and [13], we begin with the definition of nearly holomorphic modular forms.

A nonzero complex function F defined on \mathbb{H} is said to be a nearly holomorphic modular form of weight k and depth n on Γ , for nonnegative integers k and n , if it satisfies the following conditions:

- (1) There exist holomorphic functions f_0, f_1, \dots, f_n on \mathbb{H} , with $f_n \neq 0$, such that for all $z \in \mathbb{H}$,

$$(3.1) \quad F(z) = \sum_{r=0}^n \frac{f_r(z)}{(-4\pi y)^r}, \quad y = \text{Im } z.$$

- (2) For all $z \in \mathbb{H}$ and $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$,

$$(3.2) \quad F\left(\frac{az + b}{cz + d}\right) = (cz + d)^k F(z).$$

- (3) F has a finite limit at the cusps of Γ .

A function f on \mathbb{H} is said to be a quasimodular form of weight k and depth r if it is the holomorphic part of a nearly holomorphic form F of weight k and depth r , i.e., $f = f_0$ in (3.1). It turns out that f is a quasimodular form of weight k and depth n if

and only if there exist holomorphic functions f_0, f_1, \dots, f_n on \mathbb{H} , with $f_n \not\equiv 0$ such that for all $z \in \mathbb{H}$, we have:

$$(3.3) \quad (cz + d)^{-k} f\left(\frac{az + b}{cz + d}\right) = \sum_{r=0}^n f_r(z) \left(\frac{c}{cz + d}\right)^r.$$

Here, the coefficients f_r , $r = 0, \dots, n$, are the same as those of (3.1). In particular, a quasimodular form of weight k and depth 0 is an ordinary modular form of weight k , while the derivative of a weight k modular form is a quasimodular form of weight $k + 2$ and depth 1. The polynomial

$$P_{f(z)}(X) = \sum_{r=0}^n f_r(z) X^r$$

is referred as the quasimodular polynomial of f and it is unique.

Furthermore, if f satisfy (3.3), then $f = f_0$, and for each r ($0 \leq r \leq n$), f_r is a quasimodular form of weight $k - 2r$ and depth $n - r$. More precisely,

$$(cz + d)^{-k+2r} f_r\left(\frac{az + b}{cz + d}\right) = \sum_{j=0}^{n-r} \binom{r+j}{j} f_{r+j}(z) \left(\frac{c}{cz + d}\right)^j.$$

In particular, f_n is a modular form of weight $k - 2n$.

The space of quasimodular forms on Γ is of weight k and depth n denoted by $M_k^n(\Gamma)$ or simply M_k^n if there is no confusion. The total space of quasimodular forms M^* is closed under differentiation as

$$D(M_k^n) \subseteq M_{k+2}^{n+1},$$

where D is the differentiation.

As an example, when Γ is the modular group $\mathrm{SL}_2(\mathbb{Z})$, the space M^* has a special structure. Let us recall the Eisenstein series E_2, E_4 and E_6 , defined on \mathbb{H} by:

$$E_2(z) = 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^n,$$

$$E_4(z) = 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n) q^n,$$

and

$$E_6(z) = 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) q^n.$$

Here, $q = \exp(2i\pi z)$, and $\sigma_k(n)$ denotes the sum of the k -th powers of the divisors of n . It turns out that E_2 is a quasimodular form of weight 2 and depth 1, while E_4 and E_6 are modular forms of respective weight 4 and 6.

Furthermore, M^* has a structure of a graded differential ring given by

$$M^* = \mathbb{C}[E_2, E_4, E_6]$$

where the grading correspond to the weights of the Eisenstein series as quasimodular forms. The space M^* can also be viewed as a graded module of the graded ring of modular forms $\mathbb{C}[E_4, E_6]$.

More interestingly, for a general Γ , we define the operator δ on M^* by $\delta(f) = f_1$ where f is a quasimodular form satisfying (3.3). Then δ lowers the weight by 2. Define also the operator E on M^* which multiplies by k a quasimodular form of weight k . These operators δ , E together with differentiation operator D , satisfy the following commutation relations:

$$[E, D] = 2D, \quad [E, \delta] = -2\delta, \quad [D, \delta] = E.$$

These relations endow M^* the structure of an $\mathfrak{sl}_2(\mathbb{C})$ -module.

Remark 3.1.

- (1) In the definition of quasimodular forms given above, one can replace the term holomorphic with meromorphic throughout to define the notion of meromorphic quasimodular forms. All previously stated properties remain valid, except for the ring structure of quasimodular forms for $\mathrm{SL}_2(\mathbb{Z})$. Additionally, unless explicitly stated otherwise, a modular form is assumed to be meromorphic.
- (2) For convenience, even when the discrete group Γ is not a subgroup of the modular group, we will use the terms modular forms and quasimodular forms rather than the more general terms automorphic forms and quasi-automorphic forms.

4. EQUIVARIANT FUNCTIONS

We now introduce the notion of equivariant functions, which are closely related to the Schwarz derivative. Let h be a meromorphic function on \mathbb{H} . By leveraging the properties of the Schwarz derivative, one can show that $f(z) = \{h, z\}$ is a weight 4 modular form for a discrete subgroup Γ of $\mathrm{SL}_2(\mathbb{R})$ if and only if there exists a representation $\varrho : \Gamma \rightarrow \mathrm{GL}_2(\mathbb{C})$ such for all $z \in \mathbb{H}$ and all $\gamma \in \Gamma$,

$$h(\gamma z) = \varrho(\gamma)h(z).$$

Here, the action in both sides is by linear fractional transformation. A functions satisfying this property is called a ϱ -equivariant function for Γ . Throughout this paper, whenever we consider a representation ϱ of a discrete subgroup of $\mathrm{SL}_2(\mathbb{R})$, we assume that $\varrho(-I) \in \mathbb{C}^\times I$ if $-I \in \Gamma$. As for the existence of these type of functions, we have

Theorem 4.1. [15] *For every discrete group Γ and every 2-dimensional representation ϱ of Γ , ϱ -equivariant functions for Γ always exist.*

This result is proven by associating to each pair (Γ, ϱ) a vector bundle whose sections yield the desired ϱ -equivariant function. The construction of this vector bundle relies on sophisticated techniques, including solving various Riemann-Hilbert problems locally and applying the Kodaira Vanishing Theorem. The theorem is still valid if Γ is a Fuchsian group of the first or the second kind.

When ϱ is the constant representation, i.e., $\varrho(\gamma) = I$ for all $\gamma \in \Gamma$, a ϱ -equivariant function is simply an modular function for Γ . Meanwhile, if ϱ is the defining representation of Γ (i.e. the natural embedding of Γ in $\mathrm{GL}_2(\mathbb{C})$), then h commutes with the action of Γ and is simply called an equivariant function. Examples of such functions include the trivial function $h(z) = z$. Additionally, if f be a modular form of weight k for Γ , then

$$h_f(z) = z + k \frac{f(z)}{f'(z)}$$

is an equivariant function for Γ . Functions of this type are referred to as the rational equivariant functions [4].

We now prove the main result of this section. Recall that if f is ϱ -equivariant for Γ , then $\{f, z\}$ is a weight 4 modular form. It turns out that the higher Schwarzians satisfy similar properties.

Theorem 4.2. *Let Γ be a discrete subgroup of $SL_2(\mathbb{R})$. A meromorphic function f is ϱ -equivariant, for some 2-dimensional representation of Γ , if and only if, for each $n \geq 2$, the higher Schwarzian $S_n[f]$ is a quasimodular form of weight $2n$ and depth $n - 2$.*

Proof. We need to determine $S_n[f \circ \gamma](z)$ for $\gamma(z) = \frac{az+b}{cz+d}$, with $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. We have, on one hand

$$\frac{(f \circ \gamma)'}{f \circ \gamma(w) - f \circ \gamma(z)} = \frac{1}{w - z} - \sum_{n=1}^{\infty} S_n[f \circ \gamma](z)(w - z)^{n-1}.$$

On the other hand, direct computation gives

$$\begin{aligned} \frac{(f \circ \gamma)'}{f \circ \gamma(w) - f \circ \gamma(z)} &= \frac{\gamma'(z)f'(\gamma(z))}{f(\gamma(w) - f(\gamma(z)))} \\ &= \frac{\gamma'(z)}{\gamma(w) - \gamma(z)} - \sum_{k=1}^{\infty} \gamma'(z)S_k[f](\gamma(z))(\gamma(w) - \gamma(z))^{k-1} \end{aligned}$$

Meanwhile,

$$\frac{\gamma'(z)}{\gamma(w) - \gamma(z)} = \frac{1}{w - z} + \frac{c}{cz + d},$$

which implies that $S_1[\gamma](z) = \frac{-c}{cz+d}$ and $S_n[\gamma] = 0$ for $n \geq 2$. This is expected since S_n is projectively invariant for $n \geq 2$ and thus $S_n[\gamma] = S_n[z] = 0$ for $n \geq 2$.

Furthermore,

$$\begin{aligned}
& \sum_{k=1}^{\infty} \gamma'(z) S_k[f](\gamma(z)) (\gamma(w) - \gamma(z))^{k-1} \\
&= \gamma'(z) S_1[f](\gamma(z)) + \sum_{k=2}^{\infty} S_k[f](\gamma(z)) \frac{(w-z)^{k-1}}{(cz+d)^{k-1}} \frac{1}{(cw+d)^{k-1}} \\
&= \gamma'(z) S_1[f](\gamma(z)) + \sum_{k=2}^{\infty} S_k[f](\gamma(z)) \frac{(w-z)^{k-1}}{(cz+d)^{k-1}} \sum_{n=0}^{\infty} \frac{1}{n!} \frac{d^n}{dz^n} \left[\frac{1}{(cz+d)^{k-1}} \right] (w-z)^n \\
&= \gamma'(z) S_1[f](\gamma(z)) + \sum_{k=2}^{\infty} S_k[f](\gamma(z)) \sum_{n=0}^{\infty} \binom{n+k-2}{k-2} \frac{(-c)^n}{(cz+d)^{n+2k}} (w-z)^{n+k-1} \\
&= \gamma'(z) S_1[f](\gamma(z)) + \sum_{k=2}^{\infty} S_k[f](\gamma(z)) \sum_{n=k}^{\infty} \binom{n-2}{k-2} \frac{(-c)^{n-k}}{(cz+d)^{n+k}} (w-z)^{n-1} \\
&= \gamma'(z) S_1[f](\gamma(z)) + \sum_{n=2}^{\infty} \sum_{k=2}^n S_k[f](\gamma(z)) \sum_{n=k}^{\infty} \binom{n-2}{k-2} \frac{(-c)^{n-k}}{(cz+d)^{n+k}} (w-z)^{n-1}
\end{aligned}$$

Consequently,

$$S_1[f \circ \gamma](z) = \frac{-c}{cz+d} + \gamma'(z) S_1[f](\gamma(z)),$$

and for $n \geq 2$:

$$\begin{aligned}
S_n[f \circ \gamma](z) &= \sum_{k=2}^n S_k[f](\gamma(z)) \binom{n-2}{k-2} \frac{(-c)^{n-k}}{(cz+d)^{n+k}} \\
&= \sum_{j=0}^{n-2} S_{n-j}[f](\gamma(z)) \binom{n-2}{j} \frac{(-c)^j}{(cz+d)^{2n-j}}.
\end{aligned}$$

Replacing γ with γ^{-1} and substituting z by $\gamma(z)$, we deduce

$$(cz+d)^{-2n} S_n[f \circ \gamma^{-1}](\gamma(z)) = \sum_{j=0}^{n-2} \binom{n-2}{j} S_{n-j}[f](z) \left(\frac{c}{cz+d} \right)^j.$$

If f is ϱ -equivariant then, for $n \geq 2$, we have

$$S_n[f \circ \gamma^{-1}] = S_n[\varrho(\gamma^{-1})f] = S_n[f]$$

since S_n is projectively invariant. Therefore,

$$(4.1) \quad (cz + d)^{-2n} S_n[f](\gamma(z)) = \sum_{j=0}^{n-2} \binom{n-2}{j} S_{n-j}[f](z) \left(\frac{c}{cz + d} \right)^j.$$

Thus, for $n \geq 2$, $S_n[f]$ is a quasimodular form of weight $2n$ and depth $n - 2$.

Conversely if each for each $n \geq 2$, $S_n[f]$, is a quasimodular form of weight $2n$ and depth $n - 2$, then, in particular, the Schwarzian derivative $\{f, z\} = 6S_2[f]$ is a weight 4 modular form for Γ . Consequently, f is ρ -equivariant for some 2-dimensional representation of Γ . \square

We have the following type of converse.

Proposition 4.3. *Suppose that for a meromorphic function f , there exists $n \geq 2$ such that the relation (4.1) holds for all $\gamma \in \Gamma$, then it holds for every $n \geq 2$, and f is ρ -equivariant for some representation ρ of Γ .*

Proof. Suppose that for some $n \geq 2$, the relation (4.1) holds, then $S_n[f]$ is quasimodular of weight $2n$ and depth $n - 2$. In particular, $S_2[f](z) = \{f, z\}$, the term of highest depth corresponding to $j = n - 2$, is a weight 4 modular form. Therefore, f is ρ -equivariant for some ρ and the relation (4.1) holds for all $n \geq 2$ by the above theorem. \square

REFERENCES

- [1] D. Aharonov. A necessary and sufficient condition for univalence of a meromorphic function. *Duke Math. J.* 36 (1969), 599–604.
- [2] L. Ahlfors, *Lectures on quasi-conformal mappings*, Van Nostrand, (1966).
- [3] P. L. Duren, Peter, *Univalent functions*, Grundlehren der Mathematischen Wissenschaften, 259, Springer-Verlag (1983).
- [4] A. Elbasraoui; A. Sebbar. Rational equivariant forms. *Int. J. Number Th.* 08 No. 4(2012), 963–981.
- [5] L. R. Ford. *Automorphic functions*. McGraw-Hill 1929.
- [6] M. Kaneko; M. Koike, On modular forms arising from a differential equation of hypergeometric type. *Ramanujan J.* 7(2003), no. 1–3, 145–164.
- [7] M. Kaneko; D. Zagier. Supersingular j -invariants, hypergeometric series, and Atkin’s orthogonal polynomials. *Computational perspectives on number theory* (Chicago, IL, 1995), 97–126, AMS/IP Stud. Adv. Math., 7, Amer. Math. Soc., Providence, RI, 1998.
- [8] S-A. Kim; T. Sugawa. Invariant Schwarzian derivatives of higher order. *Complex Anal. Oper. Theory* 5 (2011), no. 3, 659–670.
- [9] S. Mathur, S. Mukhi, and A. Sen, On the classification of rational conformal field theories. *Phys. Lett. B* 213(1988), no. 3, 303–308.
- [10] J. McKay; A. Sebbar. Fuchsian groups, automorphic functions and Schwarzians. *Math. Ann.* 318 (2), (2000) 255–275. *Ramanujan J* (2008) 17: 405–427.
- [11] Z. Nehari, (1949), The Schwarzian derivative and schlicht functions, *Bulletin of the American Mathematical Society*, 55 (1949) 545–551.

- [12] V. Ovsienko; S. Tabachnikov. What is the Schwarzian derivative? *AMS Notices*, 56 (01): 34–36
- [13] E. Royer. Quasimodular forms: an introduction. *Ann. Math. Blaise Pascal* 19 (2012), no. 2, 297–306.
- [14] G. Sansone, J. Gerretsen. Lectures on the theory of functions of a complex variable. II, Geometric theory. Wolters–Noordhoff Publishing, Groningen 1969.
- [15] A. Sebbar; H. Saber. On the existence of vector-valued automorphic forms. *Kyushu J. Math.* 71 (2017), no. 2, 271–285.
- [16] H. Tamanoi. Higher Schwarzian operators and combinatorics of the Schwarzian derivative. *Math. Ann.* 305 (1996), no. 1, 127–151.
- [17] D. Zagier. Elliptic modular forms and their applications, in *The 1-2-3 of modular forms*, Universitext, Springer, Berlin, 2008, p. 1–103.

DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, UNIVERSITY OF HA'IL, HA'IL 55473,
KINGDOM OF SAUDI ARABIA

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, OTTAWA ONTARIO
K1N 6N5 CANADA

Email address: `hi.saber@uoh.edu.sa`

Email address: `asebbar@uottawa.ca`