# A Survey on Constructing Parseval Fusion Frames via Scaling Weights

Ehsan Ameli<sup>1</sup>, Ali Akbar Arefijamaal<sup>1\*†</sup>, Fahimeh Arabyani Neyshaburi<sup>2,1†</sup>

<sup>1</sup>Department of Mathematics and Computer Sciences, Hakim Sabzevari University, Sabzevar, Iran. <sup>2</sup>Department of Mathematical Sciences, Ferdowsi University of Mashhad, Mashhad, Iran.

\*Corresponding author(s). E-mail(s): arefijamaal@hsu.ac.ir; Contributing authors: eh.ameli@hsu.ac.ir; fahimeh.arabyani@gmail.com; <sup>†</sup>These authors contributed equally to this work.

#### Abstract

The construction of Parseval fusion frames is highly desirable in a wide range of signal processing applications. In this paper, we study the problem of modifying the weights of a fusion frame in order to generate a Parseval fusion frame. To this end, we extend the notion of the scalability to the fusion frame setting. We then proceed to characterize scalable fusion Riesz bases and 1-excess fusion frames. Furthermore, we provide the necessary and sufficient conditions for the scalability of certain  $\mathbf{k}$ -excess fusion frames,  $\mathbf{k} \geq 2$ . Finally, we present several pertinent examples to confirm the obtained results.

Keywords: Fusion frames, scalable fusion frames, Parseval fusion frames, excess

MSC Classification: Primary 42C15; Secondary 15A12

## 1 Introduction

Over the past few years, the construction of tight frames has emerged as a key concept in various areas of applied mathematics, computer science, and engineering. There have been many works to construct tight frames [7, 9, 12–14]. However, it is preferable to generate tight frames by simply scaling each frame vector, as this modification has no effect on the frame properties such as structure and erasure resilience. Thus, scaling appears to be a natural method for generating a Parseval frame. By the nature of scaling, we are looking for some scalars such that the scaled frame becomes a Parseval frame [13]. These types of frames possess a reconstruction formula analogous to orthonormal bases, and therefore they are easily used in a wide variety of applications such as sampling, signal processing, filtering, image processing, and in the other areas [5, 6, 9, 13, 14].

Fusion frames are a generalization of ordinary frames in separable Hilbert spaces, first proposed by Casazza and Kutyniok in [10]. They are a novel development that provide a mathematical framework in many applications, including sensor networks, coding theory, filter bank theory, signal and image processing and wireless communications and many other fields, where cannot be modeled by ordinary frames [1-5, 8]. One of the crucial properties that Parseval fusion frames have in common with fusion orthonormal bases is their useful role in signal processing. Indeed, they ensure a straightforward reconstruction of a signal and are resilient to noise [8]. In this respect, the construction of Parseval fusion frames has considerable significance. Inspired by this perspective, our objective is to extend the concept of scalability to the fusion frame setting and to generate Parseval fusion frames. A Parseval fusion frame  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  for  $\mathcal{H}$  has a property that every vector  $f \in \mathcal{H}$  can be recovered via the reconstruction formula:  $f = \sum_{i \in I} \omega_i^2 \pi_{W_i} f$ . If a fusion frame is not Parseval, the reconstruction formula depends on the inverse of the fusion frame operator. This may be difficult or even impossible to compute. Now, a central question emerges in the context of fusion frame theory is, how can a Parseval fusion frame be constructed from a given fusion frame without altering the structure of its subspaces?

In this survey, we propose an approach based on scaling weights to address this pivotal question. Moreover, if there exist weights that can turn a given fusion frame into a Parseval fusion frame, we refer to the fusion frame as weight-scalable. It is imperative to note that the weight-scalability process preserves the structure of subspaces, which aligns with the concept of scalability in frame theory. This provides further insight into the analysis and construction of Parseval fusion frames. Furthermore, we observe how the weights can be determined from those of the original fusion frame prior to construction. A natural question to ask is whether the weight-scalability in fusion frame theory enjoys similar properties as scalability in ordinary frames, and whether or not it provides a possibility for convenient reconstruction of signals. In addition, under what conditions is a fusion frame weight-scalable? This article responds to all of these questions.

Our recent work [1] shows that in order to determine the excess of a fusion frame, it is sufficient to compute the excess of its local frame obtained from Riesz bases. Building upon this result, we are interested in investigating the weight-scalability of overcomplete fusion frames. Indeed, it is worthwhile to identify the conditions under which a k-excess fusion frame is weight-scalable for any  $k \in \mathbb{N}$ .

The remainder of the paper is organized as follows. After introducing fusion frames and reviewing known results on them in Section 2, our main result in Section 3 presents a way to construct Parseval fusion frames; so called the weight-scalability. We examine the weight-scalability of fusion frames and present results that are comparable

to those valid for scalable frames. Furthermore, we completely characterize weightscalable fusion Riesz bases and 1-excess fusion frames. In Section 4, we investigate the conditions under which some 2-excess fusion frames are not weight-scalable and give several examples in order to confirm the obtained results. Finally, in Section 5, we turn our attention to the weight-scalability of certain k-excess fusion frames for  $k \geq 2$ and derive an equivalent condition for the weight-scalability of 2-excess fusion frames by utilizing the acquired results.

### 2 Preliminaries and Notations

We briefly review the concept of fusion frames and recall some of their main properties. We also present some results that will be used later in the paper. Throughout this paper, we suppose that  $\mathcal{H}$  is a separable Hilbert space and  $\mathcal{H}_n$  is an *n*-dimensional Hilbert space. Furthermore, I and J denote countable index sets and  $I_{\mathcal{H}}$  denotes the identity operator on  $\mathcal{H}$ . We denote the set of all bounded operators on  $\mathcal{H}$  by  $B(\mathcal{H})$ and the orthogonal projection from Hilbert space  $\mathcal{H}$  onto a closed subspace  $W \subseteq \mathcal{H}$ by  $\pi_W$ . Moreover, we denote the null space of  $T \in B(\mathcal{H})$  by N(T).

**Definition 2.1.** [10] Let  $\{W_i\}_{i \in I}$  be a family of closed subspaces of  $\mathcal{H}$  and  $\{\omega_i\}_{i \in I}$  a family of weights, i.e.  $\omega_i > 0$ ,  $i \in I$ . Then  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  is called a fusion frame for  $\mathcal{H}$  if there exists the constants  $0 < A \leq B < \infty$  such that

$$A\|f\|^{2} \leq \sum_{i \in I} \omega_{i}^{2} \|\pi_{W_{i}} f\|^{2} \leq B\|f\|^{2}, \quad (f \in \mathcal{H}).$$

$$(2.1)$$

The constants A and B are called the *fusion frame bounds*. If we only have the upper bound in (2.1), then  $\mathcal{W}$  is said to be a *fusion Bessel sequence*. A fusion frame is called A-tight if A = B, and Parseval if A = B = 1. If  $\omega_i = \omega$  for all  $i \in I$ , then  $\mathcal{W}$  is called  $\omega$ -uniform and if dim $W_i = n$  for all  $i \in I$ , then  $\mathcal{W}$  is called *n*-equi-dimensional fusion frame. We abbreviate 1-uniform fusion frames as  $\{W_i\}_{i\in I}$ . A family of closed subspaces  $\{W_i\}_{i\in I}$  is said to be a *fusion orthonormal basis* when  $\mathcal{H}$  is the orthogonal sum of the subspaces  $W_i$  and it is a *Riesz decomposition* of  $\mathcal{H}$ , if for every  $f \in \mathcal{H}$  there is a unique choice of  $f_i \in W_i$  such that  $f = \sum_{i\in I} f_i$ . It is obvious that every fusion orthonormal basis is a Riesz decomposition for  $\mathcal{H}$ , and also every Riesz decomposition is a 1-uniform fusion frame for  $\mathcal{H}$  [10]. Moreover, a family  $\{W_i\}_{i\in I}$  of closed subspaces of  $\mathcal{H}$  is a fusion orthonormal basis if and only if it is a 1-uniform Parseval fusion frame [10]. A fusion frame is said to be *exact*, if it ceases to be a fusion frame whenever any of its elements is deleted. A family of closed subspaces  $\{W_i\}_{i\in I}$  is called a *fusion Riesz basis* whenever it is complete for  $\mathcal{H}$  and there exist positive constants C and D such that for every finite subset  $J \subset I$  and arbitrary vector  $f_j \in W_j$   $(j \in J)$ , we have

$$C\sum_{j\in J} \|f_j\|^2 \le \left\|\sum_{j\in J} \omega_j f_j\right\|^2 \le D\sum_{j\in J} \|f_j\|^2.$$

Recall that for each sequence  $\{W_i\}_{i \in I}$  of closed subspaces in  $\mathcal{H}$ , the space

$$\sum_{i \in I} \bigoplus W_i = \left\{ \{f_i\}_{i \in I} : f_i \in W_i \ , \ \sum_{i \in I} \|f_i\|^2 < \infty \right\},\$$

with the inner product

$$\left\langle \{f_i\}_{i\in I}, \{g_i\}_{i\in I}\right\rangle = \sum_{i\in I} \langle f_i, g_i \rangle,$$

constitutes a Hilbert space. For a Bessel sequence  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$ , the synthesis operator  $T_{\mathcal{W}} : \sum_{i \in I} \bigoplus W_i \to \mathcal{H}$  is defined by

$$T_{\mathcal{W}}(\{f_i\}_{i\in I}) = \sum_{i\in I} \omega_i f_i, \quad \{f_i\}_{i\in I} \in \sum_{i\in I} \bigoplus W_i.$$

Moreover, the fusion frame operator  $S_{\mathcal{W}}: \mathcal{H} \to \mathcal{H}$  defined by

$$S_{\mathcal{W}}f = T_{\mathcal{W}}T_{\mathcal{W}}^*f = \sum_{i \in I} \omega_i^2 \pi_{W_i}f,$$

is positive and self-adjoint. It is shown that for a fusion frame  $\mathcal{W}$ , the fusion frame operator  $S_{\mathcal{W}}$  is invertible, and thus we have the following reconstruction formula [10]:

$$f = \sum_{i \in I} \omega_i^2 S_{\mathcal{W}}^{-1} \pi_{W_i} f, \quad (f \in \mathcal{H}).$$

In [10], it has been proved that  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  is a Parseval fusion frame if and only if  $S_{\mathcal{W}} = I_{\mathcal{H}}$ . The family  $\{(S_{\mathcal{W}}^{-1}W_i, \omega_i)\}_{i \in I}$ , which is also a fusion frame, is called the *canonical dual* of  $\mathcal{W}$ . Generally, a Bessel sequence  $\{(V_i, v_i)\}_{i \in I}$  is said to be an *alternate (Gãvruţa) dual* of  $\mathcal{W}$ , whenever

$$f = \sum_{i \in I} \omega_i \upsilon_i \pi_{V_i} S_{\mathcal{W}}^{-1} \pi_{W_i} f, \quad (f \in \mathcal{H}).$$

Let  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  be a fusion frame for  $\mathcal{H}$ . A fusion Bessel sequence  $\mathcal{V} = \{(V_i, v_i)\}_{i \in I}$  is a dual of  $\mathcal{W}$  if and only if [17]

$$T_{\mathcal{V}}\varphi_{\mathcal{V}\mathcal{W}}T_{\mathcal{W}}^*=I_{\mathcal{H}},$$

where the bounded operator  $\varphi_{\mathcal{VW}}: \sum_{i \in I} \bigoplus W_i \to \sum_{i \in I} \bigoplus V_i$  is given by

$$\varphi_{\mathcal{VW}}(\{f_i\}_{i\in I}) = \left\{\pi_{V_i}S_{\mathcal{W}}^{-1}f_i\right\}_{i\in I}.$$

For the basic facts about fusion frames we refer the reader to [4, 10, 11, 15, 16, 19]. The following result establishes the relationship between local and global properties.

**Theorem 2.2.** [10] Let  $\{W_i\}_{i \in I}$  be a family of closed subspaces of  $\mathcal{H}$ ,  $\omega_i > 0$  and  $\{f_{i,j}\}_{j \in J_i}$  be a frame (Riesz basis) for  $W_i$  with frame bounds  $A_i$  and  $B_i$  such that

$$0 < A = \inf_{i \in I} A_i \le \sup_{i \in I} B_i = B < \infty.$$

Then the following conditions are equivalent:

- (i)  $\{(W_i, \omega_i)\}_{i \in I}$  is a fusion frame (fusion Riesz basis) for  $\mathcal{H}$ ,
- (ii)  $\{\omega_i f_{i,j}\}_{i \in I, j \in J}$  is a frame (Riesz basis) for  $\mathcal{H}$ .

Finally, we list two known results that will be utilized in the subsequent sections. **Proposition 2.3.** [11] Let  $U \in B(\mathcal{H})$  and  $W \subset \mathcal{H}$  be a closed subspace. Then the following are equivalent:

(i)  $\pi_{UW}U = U\pi_W$ .

(ii)  $U^*UW \subseteq W$ .

**Proposition 2.4.** [10, 20] Let  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  be a fusion frame for  $\mathcal{H}$ . Then the following are equivalent:

(i)  $\mathcal{W}$  is a fusion Riesz basis. (ii)  $S_{\mathcal{W}}^{-1}W_i \perp W_j$  for all  $i, j \in I$ ,  $i \neq j$ . (iii)  $\omega_i^2 \pi_{W_i} S_{\mathcal{W}}^{-1} \pi_{W_j} = \delta_{i,j} \pi_{W_j}$  for all  $i, j \in I$ .

## 3 Weight-Scalable Fusion Riesz Bases and 1-excess Fusion Frames

In this section, we present a method for generating Parseval fusion frames that relies on preserving subspaces and merely scaling weights. First, we examine the results that are comparable to those observed for scalable frames. Then we provide a characterization for the scalability of fusion Riesz bases and discuss on the scalability of 1-excess fusion frames. In the following, we state the general definition of the weight-scalability for fusion frames.

**Definition 3.1.** A fusion frame  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  of  $\mathcal{H}$  is called *weight-scalable* if there exists a family of weights  $\gamma := \{\gamma_i\}_{i \in I}$  such that  $\mathcal{W}_{\gamma} := \{(W_i, \omega_i \gamma_i)\}_{i \in I}$  is a Parseval fusion frame for  $\mathcal{H}$ .

We use  $\gamma$ -scalable whenever we intend to refer to the family of weights  $\gamma = \{\gamma_i\}_{i \in I}$ that make a fusion frame weight-scalable. Assume that  $\{W_i\}_{i \in I}$  is a family of closed subspaces of  $\mathcal{H}$ . Obviously, any finite fusion frame  $\{W_i\}_{i=1}^n$  together with any sequence of weights  $\{\omega_i\}_{i=1}^n$  is again a fusion frame. This may not be the case in the infinitedimensional situation. However, if  $\{\omega_i\}_{i \in I}$  is a semi-normalized sequence [6], then  $\mathcal{W} =$  $\{(W_i, \omega_i)\}_{i \in I}$  is a fusion frame. Corresponding to a sequence of weights  $\gamma := \{\gamma_i\}_{i \in I}$ , we define the diagonal operator  $D_{\gamma}$  in  $\sum_{i \in I} \bigoplus W_i$  as follows.

$$D_{\gamma}\{f_i\}_{i\in I} = \{\gamma_i f_i\}_{i\in I}, \ \{f_i\}_{i\in I} \in \sum_{i\in I} \bigoplus W_i,$$

where

dom 
$$D_{\gamma} := \left\{ \{f_i\}_{i \in I} \in \sum_{i \in I} \bigoplus W_i : \{\gamma_i f_i\}_{i \in I} \in \sum_{i \in I} \bigoplus W_i \right\}$$

In [18], it was shown that  $D_{\gamma}$  is a self-adjoint operator in  $\sum_{i \in I} \bigoplus W_i$ . Moreover, an equivalent condition for the scalability of fusion frames was provided, see [18, Proposition 2.9] for more details.

**Proposition 3.2.** [18] Let  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  be a fusion frame for  $\mathcal{H}$ . Then the following are equivalent:

(i)  $\mathcal{W}$  is scalable,

(ii) There exists a positive diagonal operator  $D_{\gamma}$  in  $\sum_{i \in I} \bigoplus W_i$  such that

$$\overline{T_{\mathcal{W}}D_{\gamma}}D_{\gamma}T_{\mathcal{W}}^{*} = I_{\mathcal{H}}.$$
(3.1)

It is evident that 1-equi-dimensional fusion frames correspond to ordinary frames, and thereby Proposition 3.2 will align with Proposition 2.4 of [13]. Furthermore, if  $\liminf_{i \in I} \omega_i > 0$ , then the operator  $D_{\gamma}$  as in Proposition 3.2, is bounded and (3.1) can be simplified as  $T_{\mathcal{W}} D_{\gamma}^2 T_{\mathcal{W}}^* = I_{\mathcal{H}}$ . A straightforward consequence of this characterization is that the weight-scalability, like ordinary frames, is invariant under unitary transformations. What is more, if  $U \in B(\mathcal{H})$  satisfies  $U^*UW_i \subseteq W_i$   $(i \in I)$ , then it follows from Proposition 2.3 that

$$S_{UW_{\gamma}} = \sum_{i \in I} (\omega_i \gamma_i)^2 \pi_{UW_i} = U S_{W_{\gamma}} U^{-1}.$$

Therefore,  $\mathcal{W}$  is weight-scalable if and only if  $U\mathcal{W}$  is weight-scalable.

#### 3.1 Fusion Riesz bases

In the following, we provide a characterization for weight-scalable fusion Riesz bases. **Proposition 3.3.** Let  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  be a fusion Riesz basis for  $\mathcal{H}$  and  $\omega = \{\omega_i\}_{i \in I}$ . Then the following are equivalent:

(i)  $\mathcal{W}$  is  $\omega^{-1}$ -scalable, (ii)  $S_{\mathcal{W}}^{-1}\pi_{W_i} = \omega_i^{-2}\pi_{W_i}$ , for all  $i \in I$ , (iii)  $\mathcal{W}$  is orthogonal.

*Proof.*  $(i) \Rightarrow (ii)$  If  $\mathcal{W}$  is  $\omega^{-1}$ -scalable, then  $\mathcal{W}_{\omega^{-1}}$  is a Parseval fusion frame. Hence,

$$S_{\mathcal{W}}^{-1}\pi_{W_i} = S_{\mathcal{W}_{\omega^{-1}}}S_{\mathcal{W}}^{-1}\pi_{W_i}$$
$$= \sum_{j\in I}\pi_{W_j}S_{\mathcal{W}}^{-1}\pi_{W_i}$$
$$= \pi_{W_i}S_{\mathcal{W}}^{-1}\pi_{W_i} = \omega_i^{-2}\pi_{W_i},$$

for all  $i \in I$ , where the last line is obtained by Proposition 2.4. The implications  $(ii) \Rightarrow (iii)$  and  $(iii) \Rightarrow (i)$  are straightforward.

The following example demonstrates a weight-scalable fusion Riesz basis.

**Example 3.4.** Consider  $W_1 = \operatorname{span}\{u\}$  and  $W_2 = \{0\} \times \mathcal{H}_2$ , where  $u = (u_1, u_2, u_3) \in \mathcal{H}_3$  is a unit vector such that  $u_1 \neq 0$ . Then  $\mathcal{W} = \{(W_i, \omega_i)\}_{i=1}^2$  is a fusion Riesz basis for  $\mathcal{H}_3$  with the fusion frame operator

$$S_{\mathcal{W}} = \begin{pmatrix} \omega_1^2 u_1^2 & \omega_1^2 u_1 u_2 & \omega_1^2 u_1 u_3 \\ \omega_1^2 u_1 u_2 & \omega_1^2 u_2^2 + \omega_2^2 & \omega_1^2 u_2 u_3 \\ \omega_1^2 u_1 u_3 & \omega_1^2 u_2 u_3 & \omega_1^2 u_3^2 + \omega_2^2 \end{pmatrix}.$$

Obviously, W is  $\omega^{-1}$ -scalable if and only if  $u_2 = u_3 = 0$ , which confirms Proposition 3.3.

A direct computation analogous to the above example confirms that, without losing the generality, we may consider 1-uniform fusion frames. However, it should be noted that our results are valid for general fusion frames.

#### 3.2 1-excess fusion frames

In what follows, we study the weight-scalability of 1-excess fusion frames. First, we state the concept of excess for fusion frames [15]. Let  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  be a fusion frame for  $\mathcal{H}$  with the synthesis operator  $T_{\mathcal{W}}$ . The *excess* of  $\mathcal{W}$  is defined as

$$e(\mathcal{W}) = \dim N(T_{\mathcal{W}}).$$

In [1], the authors present a novel approach for the excess of fusion frames, which is related to the excess of their local frames.

**Proposition 3.5.** [1] Let  $\mathcal{W} = \{(W_i, \omega_i)\}_{i \in I}$  be a fusion frame for  $\mathcal{H}$  with a local frame  $\mathcal{F} = \{\omega_i f_{i,j}\}_{i \in I, j \in J_i}$ , where  $\{f_{i,j}\}_{j \in J_i}$  is a Riesz basis for  $W_i$ , for all  $i \in I$ . Then

$$e(\mathcal{W}) = e(\mathcal{F}).$$

Suppose that  $\mathcal{W} = \{W_i\}_{i \in I}$  is a fusion frame for  $\mathcal{H}$ . As stated before, if all  $W_i$ 's are one dimensional, then  $\mathcal{W}$  corresponds to an ordinary frame. Accordingly, the notion of scalability for such frames and fusion frames becomes equivalent. For instance, 1-equi-dimensional fusion frames associated with 1-excess tight frames, such as the Mercedes-Benz frame, can easily be made weight-scalable. In this respect, it is crucial to investigate the weight-scalability of fusion frames in which the subspaces are not necessarily one dimensional. Taking this into account, we discuss the weight-scalability of 1-excess fusion frames and provide some equivalent conditions for the weight-scalability of such fusion frames.

**Theorem 3.6.** Let  $\mathcal{V}$  be a 1-excess fusion frame for  $\mathcal{H}$  with the Riesz part  $\mathcal{W}$ . Then the following conditions are equivalent:

- (i)  $\mathcal{V}$  is  $\gamma$ -scalable,  $\gamma = \{\gamma_i\}_{i=0}^{\infty}$ .
- (ii) Every excess element x belongs to a one dimensional subspace and

$$\gamma_0^2 \pi_{W_i} S_{W_\gamma}^{-1} \pi_{V_0} \pi_{W_j} = \left(\gamma_i^{-2} \delta_{i,j} - \pi_{W_i}\right) \pi_{W_j}, \quad (i, j \ge 1),$$
(3.2)

where  $V_0$  is the subspace generated by x.

(iii) Every excess element x belongs to a one dimensional subspace and

$$\gamma_0^2 \left\langle \pi_{W_j} f, x \right\rangle x_i = \left( \delta_{i,j} - \gamma_i^2 \pi_{W_i} \right) \pi_{W_j} f, \quad (i, j \ge 1, \ f \in \mathcal{H}).$$

$$(3.3)$$

*Proof.*  $(i) \Rightarrow (ii)$  Towards a contradiction, assume that the excess element x does not belong to a one dimensional subspace. Thus, without losing the generality,  $\mathcal{V}$  can be expressed as

$$\mathcal{V} = V_1 \cup \{W_i\}_{i=2}^{\infty}$$

where  $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$  is a fusion Riesz basis,  $V_1 = W_1 \oplus \operatorname{span}\{x\}$  and the redundant element  $x = \sum_{i=2}^{\infty} x_i$  is a unit vector of  $\mathcal{H}$  such that  $x \perp W_1$  and  $x_i \in W_i$  for each  $i \geq 2$ . Since  $\mathcal{V}$  is  $\gamma$ -scalable, then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=1}^{\infty}$  such that

$$f = S_{\mathcal{V}_{\gamma}}f = \gamma_1^2 \pi_{W_1}f + \gamma_1^2 \langle f, x \rangle x + \sum_{i=2}^{\infty} \gamma_i^2 \pi_{W_i}f, \quad (f \in \mathcal{H}).$$
(3.4)

Put a non zero vector  $g \in W_1$  in (3.4). It gives  $g = \gamma_1^2 g + \sum_{i=2}^{\infty} \gamma_i^2 \pi_{W_i} g$ . It follows from the Riesz decomposition property of  $\mathcal{W}$  that  $\gamma_1 = 1$  and  $W_1 \perp W_i$  for all  $i \geq 2$ . Moreover, by reformulating (3.4), we get

$$S_{\mathcal{W}_{\gamma}}f = f - \gamma_1^2 \langle f, x \rangle \, x = f - \langle f, x \rangle \, x,$$

for each  $f \in \mathcal{H}$ . Substituting x for f gives  $S_{W_{\gamma}}x = 0$ . Thus, x = 0. That is a contradiction and so x must belong to a one dimensional subspace. As such,  $\mathcal{V}$  must be rewritten as

$$\mathcal{V} = V_0 \cup \mathcal{W}_2$$

where  $V_0 = \operatorname{span}\{x\}$  and  $x = \sum_{i=1}^{\infty} x_i$  is a unit vector of  $\mathcal{H}$ . In light of  $\mathcal{V}$  is  $\gamma$ -scalable, we get  $\gamma_0^2 \pi_{V_0} + S_{\mathcal{W}_{\gamma}} = I_{\mathcal{H}}$ , and hence

$$S_{\mathcal{W}_{\gamma}}^{-1} = \gamma_0^2 S_{\mathcal{W}_{\gamma}}^{-1} \pi_{V_0} + I_{\mathcal{H}}.$$

Thus, for each  $i, j \ge 1$ , we obtain

$$\gamma_{0}^{2}\pi_{W_{i}}S_{\mathcal{W}_{\gamma}}^{-1}\pi_{V_{0}}\pi_{W_{j}} = \gamma_{0}^{2}\pi_{W_{i}}S_{\mathcal{W}_{\gamma}}^{-1}\pi_{V_{0}}\pi_{W_{j}} + \pi_{W_{i}}\pi_{W_{j}} - \pi_{W_{i}}\pi_{W_{j}}$$
$$= \pi_{W_{i}}\left(\gamma_{0}^{2}S_{\mathcal{W}_{\gamma}}^{-1}\pi_{V_{0}} + I_{\mathcal{H}}\right)\pi_{W_{j}} - \pi_{W_{i}}\pi_{W_{j}}$$
$$= \pi_{W_{i}}S_{\mathcal{W}_{\gamma}}^{-1}\pi_{W_{j}} - \pi_{W_{i}}\pi_{W_{j}}$$
$$= \left(\gamma_{i}^{-2}\delta_{i,j} - \pi_{W_{i}}\right)\pi_{W_{j}},$$

where the last equality follows from Proposition 2.4. So, (ii) holds.

 $(ii) \Rightarrow (i)$  Assume that  $\mathcal{V} = V_0 \cup \mathcal{W}$ , where  $V_0 = \operatorname{span}\{x\}$ . In addition, suppose that there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=0}^{\infty}$  such that (3.2) is satisfied. Then, we induce

$$S_{\mathcal{V}_{\gamma}}S_{\mathcal{W}} = \left(\gamma_0^2 \pi_{V_0} + S_{\mathcal{W}_{\gamma}}\right)S_{\mathcal{W}}$$

$$= \sum_{j=1}^{\infty} \left( \gamma_0^2 S_{W_{\gamma}} S_{W_{\gamma}}^{-1} \pi_{V_0} \pi_{W_j} + S_{W_{\gamma}} \pi_{W_j} \right)$$
  
$$= \sum_{i,j=1}^{\infty} \left( (\gamma_0 \gamma_i)^2 \pi_{W_i} S_{W_{\gamma}}^{-1} \pi_{V_0} \pi_{W_j} + \gamma_i^2 \pi_{W_i} \pi_{W_j} \right)$$
  
$$= \sum_{i,j=1}^{\infty} \delta_{i,j} \pi_{W_j} = \sum_{i=1}^{\infty} \pi_{W_i} = S_W,$$

where the last line is derived by applying (3.2). Therefore,  $S_{\mathcal{V}_{\gamma}} = I_{\mathcal{H}}$ , i.e.  $\mathcal{V}$  is  $\gamma$ -scalable.

 $(ii) \Leftrightarrow (iii)$  The equation (3.2) may be simplified as follows:

$$\left( \delta_{i,j} - \gamma_i^2 \pi_{W_i} \right) \pi_{W_j} f = (\gamma_i \gamma_0)^2 \pi_{W_i} S_{\mathcal{W}_{\gamma}}^{-1} \pi_{V_0} \pi_{W_j}$$
  
=  $(\gamma_i \gamma_0)^2 \langle \pi_{W_j} f, x \rangle \pi_{W_i} S_{\mathcal{W}_{\gamma}}^{-1} x$   
=  $\gamma_0^2 \langle \pi_{W_j} f, x \rangle x_i,$ 

where the last equality follows from Proposition 2.4. This completes the proof.  $\Box$ 

This result can be generalized to a certain type of k-excess fusion frames by a similar approach for any  $k \ge 2$ , see Proposition 5.1. In the following, we examine the validity of Theorem 3.6 through an example.

**Example 3.7.** Let  $\{e_i\}_{i\in\mathbb{Z}}$  be an orthonormal basis for  $\mathcal{H}$ ,  $W_1 = \overline{\operatorname{span}}_{i\geq 0}\{e_i\}$  and  $W_2 = \overline{\operatorname{span}}_{i\leq 0}\{e_i\}$ . It is easily observed that  $\mathcal{V} = \{W_i\}_{i=1}^2$  is an exact fusion frame which is not Riesz basis, see [10]. In addition, it is proved in [1] that  $\mathcal{V}$  is a 1-excess fusion frame for  $\mathcal{H}$ . According to Theorem 3.6,  $\mathcal{V}$  is not weight-scalable, as the excess element does not belong to a one dimensional subspace. However, we investigate this matter directly. Suppose  $\mathcal{V}$  is  $\gamma$ -scalable, then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i\in\mathbb{Z}}$  such that

$$\begin{split} f &= S_{\mathcal{V}_{\gamma}} f = \sum_{i \geq 0} \gamma_1^2 \left\langle f, e_i \right\rangle e_i + \sum_{i \leq 0} \gamma_2^2 \left\langle f, e_i \right\rangle e_i \\ &= \sum_{i > 0} \gamma_1^2 \left\langle f, e_i \right\rangle e_i + \sum_{i < 0} \gamma_2^2 \left\langle f, e_i \right\rangle e_i + \left(\gamma_1^2 + \gamma_2^2\right) \left\langle f, e_0 \right\rangle e_0, \end{split}$$

for all  $f \in \mathcal{H}$ . Substituting  $e_{-1}, e_0, e_1$  for f results in  $\gamma_i = 1$  (i = 1, 2) and  $\gamma_1^2 + \gamma_2^2 = 1$ , a contradiction.

In the sequel, in view of Theorem 3.6, we proceed to investigate the weightscalability of 1-excess fusion frames of the form

$$\mathcal{V} = V_0 \cup \mathcal{W},\tag{3.5}$$

where  $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$  is a fusion Riesz basis,  $V_0 = \operatorname{span}\{x\}$  and  $x = \sum_{i \in \sigma} x_i$  is a unit vector of  $\mathcal{H}$  such that  $x_i \in W_i$  and  $\sigma = \{i \ge 1 \mid x_i \ne 0\}$ . The next result provides the essential criteria for the weight-scalability of  $\mathcal{V}$ .

**Corollary 3.8.** Let  $\mathcal{V}$  be the 1-excess fusion frame given by (3.5). If  $\mathcal{V}$  is weightscalable, then the following statements hold:

- (i)  $\gamma_0 < 1$  and x is an eigenvector of  $S_{W_{\gamma}}$  associated with eigenvalue  $1 \gamma_0^2$ .
- (*ii*)  $\langle x_j, x \rangle = \frac{1 \gamma_j^2}{\gamma_0^2}, \ \pi_{W_i} x_j = \frac{\gamma_j^2 1}{\gamma_i^2} x_i \text{ for } j \in \sigma, \ i \neq j.$
- (iii) If  $j \in \sigma^c$ , then  $\gamma_j = 1$ ,  $x \perp W_j$  and  $W_j \perp W_i$  for  $i \neq j$ . (iv) If  $j \in \sigma$ , then  $\gamma_j \neq 1$ ,  $x \not\perp W_j$  and  $\dim W_j = 1$ .

such that

*Proof.* Due to  $\mathcal{V}$  is weight-scalable, there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=0}^{\infty}$ 

 $f = \gamma_0^2 \langle f, x \rangle x + S_{\mathcal{W}_{\alpha}} f, \quad (f \in \mathcal{H}).$ (3.6)

(i) Putting f = x in (3.6), we get  $S_{W_{\gamma}}x = (1 - \gamma_0^2)x$ . If  $\gamma_0 = 1$ , it yields x = 0, a contradiction. Moreover, the positivity of  $S_{W_{\gamma}}$  assures that  $\gamma_0 < 1$ . Thus (i) holds.

(ii) Replacing  $f = x_j$   $(j \in \sigma)$  in (3.3) of Theorem 3.6, we get  $(\delta_{i,j} - \gamma_i^2 \pi_{W_i}) x_j =$  $\gamma_0^2 \langle x_j, x \rangle x_i$ , which leads to

$$\begin{cases} \left(1 - \gamma_0^2 \langle x_j, x \rangle - \gamma_j^2\right) x_j = 0, & (j = i), \\ \gamma_0^2 \langle x_j, x \rangle x_i + \gamma_i^2 \pi_{W_i} x_j = 0, & (j \neq i). \end{cases}$$

The first equation yields  $\langle x_j, x \rangle = \frac{1 - \gamma_j^2}{\gamma_0^2}$ . Moreover, the second equation implies that  $\pi_{W_i} x_j = \frac{\gamma_j^2 - 1}{\gamma_i^2} x_i \text{ for all } i \neq j.$ 

(iii) Setting  $0 \neq g \in W_j$  in (3.3), we have

$$\left(\delta_{i,j} - \gamma_i^2 \pi_{W_i}\right)g = \gamma_0^2 \langle g, x \rangle x_i, \quad (i \ge 1).$$

$$(3.7)$$

Since  $x_j = 0$ , then by assigning i = j in (3.7) yields  $\gamma_j = 1$ . Moreover, after renaming the indices  $i \leftrightarrow j$  and taking  $0 \neq f \in W_i$  in (3.3), we obtain

$$\left(\delta_{j,i} - \pi_{W_j}\right)f = \gamma_0^2 \langle f, x \rangle \, x_j = 0, \quad (i \neq j).$$

It derives that  $\pi_{W_j} f = 0$ , i.e.,  $W_j \perp W_i$  for  $i \neq j$ . Hence,  $x \perp W_j$  because  $\langle x, g \rangle =$  $\sum_{i \in \sigma} \langle x_i, g \rangle = 0$  for each  $0 \neq g \in W_j$ .

(iv) Let  $x_j \neq 0$ . By contrary, assume that  $x \perp W_j$ , then

$$||x_j||^2 + \sum_{i \in \sigma, i \neq j} \langle x_i, x_j \rangle = \langle x, x_j \rangle = 0.$$
(3.8)

If  $i \neq j$  in (3.7), we get  $W_j \perp W_i$ . Thus, it yields from (3.8) that  $||x_j|| = 0$  and thereby  $x_j = 0$ , a contradiction. So,  $x \not\perp W_j$ . That means that there exists a non zero vector  $g \in W_j$  such that  $\langle g, x \rangle \neq 0$ . Putting f = g in (3.7) and considering i = j, we get

$$(1 - \gamma_j^2)g = \gamma_0^2 \langle g, x \rangle x_j.$$

Hence, it follows that  $\gamma_j \neq 1$  and  $g \in \text{span}\{x_j\}$ . We claim that  $W_j = \text{span}\{x_j\}$ . In fact, if there exists  $0 \neq h \in W_j$  such that  $\langle h, x \rangle = 0$ , then from (iii) we obtain  $\gamma_j = 1$ , that is a contradiction. Therefore,  $\dim W_j = 1$ , the desired result.

This result prompts us to characterize all weight-scalable 1-excess fusion frames. **Theorem 3.9.** Every weight-scalable 1-excess fusion frame  $\mathcal{V}$  for  $\mathcal{H}$  is precisely of the form

$$\mathcal{V} = \left\{ \operatorname{span} \left\{ f_i \right\} \right\}_{i \in I_1} \cup \left\{ W_i \right\}_{i \in I_2},$$

where  $\mathcal{F} := \{f_i\}_{i \in I_1}$  is a strictly scalable 1-excess frame sequence of unit vectors and  $\{W_i\}_{i \in I_2}$  is an orthogonal fusion Riesz basis for  $(\operatorname{span} \{f_i\}_{i \in I_1})^{\perp}$ .

*Proof.* Assume that  $\mathcal{F}$  is a strictly scalable frame for some  $\mathcal{H}_1 \subset \mathcal{H}$ , then there exists a sequence of weights  $\{\gamma_i\}_{i \in I_1}$  such that

$$\sum_{i \in I_1} \gamma_i^2 \langle f, f_i \rangle f_i = f, \quad (f \in \mathcal{H}_1).$$
(3.9)

Moreover,  $\{W_i\}_{i \in I_2}$  is a uniform scalable fusion Riesz basis for  $\mathcal{H}_1^{\perp}$ , i.e.,  $\sum_{i \in I_2} \pi_{W_i} = I_{\mathcal{H}_1^{\perp}}$ . We put  $\gamma_i = 1$  for all  $i \in I_2$ . Thus, by considering  $\gamma = \{\gamma_i\}_{i \in I_1 \cup I_2}$  and in view of the fact that for every  $g \in \mathcal{H}$ , there exist unique vectors  $g_1 \in \mathcal{H}_1$  and  $g_2 \in \mathcal{H}_1^{\perp}$  such that  $g = g_1 + g_2$  we obtain

$$S_{\mathcal{V}_{\gamma}}g = \sum_{i \in I_1} \gamma_i^2 \pi_{\text{span}\{f_i\}}(g_1 + g_2) + \sum_{i \in I_2} \pi_{W_i}(g_1 + g_2)$$
$$= \sum_{i \in I_1} \gamma_i^2 \langle g_1, f_i \rangle f_i + \sum_{i \in I_2} \pi_{W_i}g_2$$
$$= g_1 + g_2 = g,$$

where the last line is obtained by using (3.9). Conversely, if the 1-excess fusion frame  $\mathcal{V}$  given by (3.5) is weight-scalable, then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i \in I}$  such that  $S_{\mathcal{V}_{\gamma}} = I_{\mathcal{H}}$ . The conditions (*iii*) and (*iv*) of Corollary 3.8 ensure that  $\mathcal{F} := \{x, x_i\}_{i \in \sigma}$  constitutes a 1-excess ordinary frame for  $\mathcal{H}_1 := \overline{\text{span}} \{x, x_i\}_{i \in \sigma}$  and  $\{W_i\}_{i \in \sigma^c}$  is an orthogonal fusion Riesz basis for  $\mathcal{H}_1^{\perp} = \bigoplus_{i \in \sigma^c} W_i$ . Hence, it follows from the weight-scalability of  $\mathcal{V}$  and Corollary 3.8 that  $\gamma_i = 1$  for  $i \in \sigma^c$ . So, for every  $f \in \mathcal{H}_1$  we get

$$f = S_{\mathcal{V}_{\gamma}} f$$
  
=  $\gamma_0^2 \pi_{\text{span}\{x\}} f + \sum_{i \in \sigma} \gamma_i^2 \pi_{\text{span}\{x_i\}} f + \sum_{i \in \sigma^c} \pi_{W_i} f$   
=  $\gamma_0^2 \langle f, x \rangle x + \sum_{i \in \sigma} \gamma_i^2 \frac{\langle f, x_i \rangle}{\|x_i\|^2} x_i = S_{\gamma \mathcal{F}} f.$ 

Therefore,  $\mathcal{F}$  is strictly scalable.

Now, we present an equivalent condition for the weight-scalability of 1-excess fusion frames containing orthogonal fusion Riesz bases.

**Corollary 3.10.** Let  $\mathcal{V}$  be the 1-excess fusion frame as defined in (3.5) and  $\mathcal{W}$  an orthogonal fusion Riesz basis for  $\mathcal{H}$ . Then the following are equivalent:

- (i)  $\mathcal{V}$  is weight-scalable.
- (ii) There exists a subspace of W generated by the excess element.

*Proof.*  $(i) \Rightarrow (ii)$  One can easily see that there exists a  $j \ge 1$  such that the excess element  $x \not\perp W_j$ . Since  $\mathcal{V}$  is weight-scalable, it gives  $W_j = \operatorname{span}\{x_j\}$  by Corollary 3.8(iv). Putting  $f = x_j$  in (3.3) yields

$$\gamma_0^2 \langle x_j, x \rangle x_i = \left( \delta_{i,j} - \gamma_i^2 \pi_{W_i} \right) x_j = 0, \quad (i \neq j).$$

It can be seen that  $\langle x_j, x \rangle$  is a non zero scalar because of the orthogonality of  $\mathcal{W}$ . Therefore,  $x_i = 0$  for all  $i \neq j$ , which implies that  $W_j = V_0$ .

 $(ii) \Rightarrow (i)$  Assume that there exists a  $j \ge 1$  such that  $W_j = V_0$ . Taking  $\gamma_0, \gamma_j$  satisfying  $\gamma_0^2 + \gamma_j^2 = 1$  and  $\gamma_i = 1$  for  $i \ne j$ , we conclude that

$$S_{\mathcal{V}_{\gamma}} = \left(\gamma_0^2 + \gamma_j^2\right) \pi_{W_j} + \sum_{i=1, i \neq j}^{\infty} \gamma_i^2 \pi_{W_i}$$
$$= \pi_{W_j} + \sum_{i=1, i \neq j}^{\infty} \pi_{W_i} = \pi_{\bigoplus_{i=1}^{\infty} W_i} = I_{\mathcal{H}}.$$

Motivated by the obtained results, we seek a more precise understanding of weightscalable 1-excess fusion frames. To this end, consider the 1-excess fusion frame  $\mathcal{V}$ introduced in (3.5). It is apparent that if dim $W_i > 1$  for all  $i \ge 1$ , then  $\mathcal{V}$  is not weightscalable. Indeed, the redundant element belongs to one of  $W_i$ 's, which is in conflict with the assertion made in Theorem 3.6. Therefore, there is at least a one dimensional subspace in weight-scalable 1-excess fusion frames.

Remark 3.11. (i) The orthogonality condition of  $\mathcal{W}$  can be removed from the assumptions of Corollary 3.10, provided that dim $\mathcal{H} = 3$  and  $\mathcal{W}$  is not a 1-equi-dimensional fusion Riesz basis. In this case, the weight-scalability of  $\mathcal{V}$  implies the orthogonality of  $\mathcal{W}$ . Indeed, without losing the generality, such 1-excess fusion frames for  $\mathcal{H}_3$  are of the form  $\mathcal{V} = V_0 \cup \mathcal{W}$ , where

$$V_0 = \operatorname{span}\{v\}, \ W_1 = \operatorname{span}\{u\}, \ W_2 = \{0\} \times \mathcal{H}_2,$$

such that  $v = (v_1, v_2, v_3)$ ,  $u = (u_1, u_2, u_3)$  and  $u_1 \neq 0$ . Let  $\mathcal{V}$  be weight-scalable and by contrary assume that  $W_1 \not\perp W_2$ . If  $V_0 = W_1$ , then  $V_0 \not\perp W_2$ . Hence, dim $W_2 = 1$ , by Corollary 3.8(iv), a contradiction. Thus, suppose that v and u are linearly independent. Now, if  $\{v, u, e_2\}$  or  $\{v, u, e_3\}$  constitutes a basis for  $\mathcal{H}_3$ , then  $e_3$  or  $e_2$  may be regarded as the excess element. Thus, Theorem 3.6 guarantees that it must belong to a one dimensional subspace, which is a contradiction. Otherwise, if  $\{v, u, e_2\}$  and

 $\{v, u, e_3\}$  are not bases, it easily follows that  $e_2, e_3 \in \text{span}\{v, u\}$ , as v and u are linearly independent. Hence,  $\mathcal{W}$  is not a Riesz basis. That is again a contradiction. Therefore,  $\mathcal{W}$  is orthogonal and  $V_0 = W_1$ , by Corollary 3.10.

(ii) The assertion presented in (i) is not universally applicable to the cases where  $\dim \mathcal{H} \neq 3$  or  $\mathcal{W}$  is 1-equi-dimensional. To illustrate, one can refer to 1-equi-dimensional fusion frames associated with the Mercedes-Benz frames. As another example, let  $\{e_i\}_{i=1}^n$  be the canonical orthonormal basis for  $\mathcal{H}_n$   $(n \geq 4)$ . Consider

$$V_0 = \operatorname{span} \left\{ \alpha e_1 - \sqrt{1 - \alpha^2} e_2 \right\},$$
  

$$W_1 = \operatorname{span} \{ e_1 \}, \ W_2 = \operatorname{span} \{ e_1 + e_2 \}, \ W_3 = \operatorname{span} \{ e_i \}_{i=3}^n,$$

where  $0 < \alpha < \frac{\sqrt{2}}{2}$  and  $\{W_i\}_{i=1}^3$  is a non-orthogonal fusion Riesz basis. Then  $\mathcal{V} = V_0 \cup \{W_i\}_{i=1}^3$  is a 1-excess fusion frame for  $\mathcal{H}_n$ . A straightforward computation shows that  $\mathcal{V}$  is  $\gamma$ -scalable by

$$\begin{cases} \gamma_0 = \sqrt{\frac{1}{1 - \alpha^2 + \alpha\sqrt{1 - \alpha^2}}}, \ \gamma_1 = \sqrt{\frac{1 - 2\alpha^2}{1 - \alpha^2 + \alpha\sqrt{1 - \alpha^2}}},\\ \gamma_2 = \sqrt{\frac{2\alpha\sqrt{1 - \alpha^2}}{1 - \alpha^2 + \alpha\sqrt{1 - \alpha^2}}}, \ \gamma_3 = 1. \end{cases}$$

**Example 3.12.** Let  $\{e_i\}_{i=1}^4$  be the canonical orthonormal basis for  $\mathcal{H}_4$ . Take

$$V_0 = \operatorname{span}\{\alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3 + \alpha_4 e_4\},\$$
  
$$W_1 = \operatorname{span}\{e_1, e_2\}, W_2 = \operatorname{span}\{e_3 + \beta e_1\}, W_3 = \operatorname{span}\{e_4\},\$$

where  $\alpha_i, \beta \in \mathbb{C}$  for  $1 \leq i \leq 4$  and  $\mathcal{W} = \{W_i\}_{i=1}^3$  is a fusion Riesz basis. Then  $\mathcal{V} = V_0 \cup \{W_i\}_{i=1}^3$  is a 1-excess fusion frame for  $\mathcal{H}_4$ . If  $\beta = 0$ , then  $\mathcal{W}$  is orthogonal, and it can be easily seen that  $\mathcal{V}$  is weight-scalable if and only if  $V_0 = W_2$  or  $V_0 = W_3$ , according to Corollary 3.10. Otherwise, let  $\beta \neq 0$ . If either  $\alpha_1$  or  $\alpha_2$  is non zero, then  $\mathcal{V}$  is not weight-scalable, by Corollary 3.8(iv). Also, in the case  $\alpha_1 = \alpha_2 = 0$ , Corollary 3.8(ii) confirms that  $\mathcal{V}$  is not weight-scalable.

## 4 Weight-Scalable 2-excess Fusion Frames

This section is devoted to studying the weight-scalability of 2-excess fusion frames. In general, four distinct situations there exist for the excess elements of a 2-excess fusion frame. However, this investigation focuses on the situation where the excess elements are located in more than one subspace of the Riesz part, as this case allows for the derivation of outcomes that are comparable to those of the other three cases. It should be noted that a 2-excess fusion frame is not weight-scalable if the redundant elements lie in a subspace of its Riesz part, see Corollary 5.2 for the general case. Hence, without losing the generality, every 2-excess fusion frame  $\mathcal{V}$  in which the redundant vectors are

located in different subspaces of its Riesz part can be expressed as

$$\mathcal{V} = \{V_1, V_2, W_i\}_{i=3}^{\infty}, \qquad (4.1)$$

where  $V_{\ell} = W_{\ell} \oplus \text{span}\{x^{(\ell)}\}$  for  $\ell = 1, 2$ , and  $x^{(1)} = \sum_{i=2}^{\infty} x_i^{(1)}$  and  $x^{(2)} = \sum_{i=1, i\neq 2}^{\infty} x_i^{(2)}$  are unit vectors of  $\mathcal{H}$  such that  $x_i^{(\ell)}$ 's are vectors of  $W_i$  and  $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$  is a fusion Riesz basis for  $\mathcal{H}$ . At first, we present some equivalent conditions for the weight-scalability of  $\mathcal{V}$ .

**Proposition 4.1.** Let  $\mathcal{V}$  be the 2-excess fusion frame defined by (4.1) and  $\mathcal{W}$  a fusion Riesz basis for  $\mathcal{H}$ . Then the following conditions are equivalent:

(i)  $\mathcal{V}$  is  $\gamma$ -scalable,  $\gamma = \{\gamma_i\}_{i=1}^{\infty}$ .

(*ii*) 
$$\sum_{\ell=1}^{2} \gamma_{\ell}^{2} \pi_{W_{i}} S_{W_{\gamma}}^{-1} \pi_{\operatorname{span}\{x^{(\ell)}\}} \pi_{W_{j}} = (\gamma_{i}^{-2} \delta_{i,j} - \pi_{W_{i}}) \pi_{W_{j}}, \text{ for all } i, j.$$

(iii)  $\sum_{\ell=1}^{2} \gamma_{\ell}^{2} \langle \pi_{W_{j}} f, x^{(\ell)} \rangle x_{i}^{(\ell)} = (\delta_{i,j} - \gamma_{i}^{2} \pi_{W_{i}}) \pi_{W_{j}} f, \text{ for all } f \in \mathcal{H}.$ 

*Proof.* The proof is derived by using an analogous approach to Theorem 3.6.

In order to simplify the notation, the symbols x and y are used instead of  $x^{(1)}$  and  $x^{(2)}$ , as the excess elements introduced in (4.1). One of our aims is to characterize weight-scalable 2-excess fusion frames. In this regard, we obtain some necessary conditions for the weight-scalability of such fusion frames.

**Theorem 4.2.** Let the 2-excess fusion frame  $\mathcal{V}$  given by (4.1) be weight-scalable. Then the following statements hold:

(1) 
$$\langle x, y \rangle = 0, \ \pi_{W_i} x = \frac{1 - \gamma_1^2}{\gamma_i^2} x_i \text{ for all } i \ge 2.$$

(2) 
$$\pi_{W_i} y = \frac{1}{\gamma_i^2} y_i \text{ for all } i \ge 2.$$

(3)  $\gamma_1, \gamma_2 \neq 1, W_1 \perp W_2 \text{ and } \dim W_i = 1 \text{ for } i = 1, 2.$ 

(4) 
$$\langle x_2, x \rangle = \frac{1 - \gamma_2^2}{\gamma_1^2}, \ \pi_{W_i} x_2 = \frac{\gamma_2^2 - 1}{\gamma_i^2} x_i \text{ for all } i \ge 3.$$

(5) 
$$\langle y_1, y \rangle = \frac{1 - \gamma_1^2}{\gamma_2^2}, \ \pi_{W_i} y_1 = \frac{\gamma_1^2 - 1}{\gamma_i^2} y_i \text{ for all } i \ge 3.$$

(6) 
$$||x_2||^2 = ||y_1||^{-2} = \frac{\gamma_2(1-\gamma_2)}{\gamma_1^2(1-\gamma_1^2)}$$
  
(7)  $1 \le \gamma_1^2 + \gamma_2^2 < 2.$ 

*Proof.* Since  $\mathcal{V}$  is  $\gamma$ -scalable, then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=1}^{\infty}$  such that

$$f = \gamma_1^2 \pi_{V_1} f + \gamma_2^2 \pi_{V_2} f + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} f, \quad (f \in \mathcal{H}).$$
(4.2)

First of all, it turns out that  $y_1, x_2 \neq 0$ . To see this, we only prove that  $x_2 \neq 0$  and the proof of  $y_1 \neq 0$  is analogous. To the contrary, assume that  $x_2 = 0$ . Putting a non zero vector  $f \in W_2$  in (4.2) and using the Riesz decomposition property of  $\mathcal{W}$ , we get

 $(1 - \gamma_2^2)f = 0$ . So,  $\gamma_2 = 1$ . We now investigate two cases: If  $y_1 = 0$ , it similarly follows  $\gamma_1 = 1$ . Moreover, by taking f = x in (4.2), we get

$$\pi_{V_2} x + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} x = x - \pi_{V_1} x = 0.$$

Hence,  $\|\pi_{V_2}x\|^2 + \sum_{i=3}^{\infty} \gamma_i^2 \|\pi_{W_i}x\|^2 = \langle \pi_{V_2}x + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i}x, x \rangle = 0$ , which assures that  $\pi_{V_2}x = 0$ , and  $\pi_{W_i}x = 0$  for all  $i \ge 3$ . Therefore,

$$||x||^{2} = \left\langle x, \sum_{i=3}^{\infty} x_{i} \right\rangle = \sum_{i=3}^{\infty} \left\langle x, x_{i} \right\rangle = 0.$$

This means that x = 0, which is a contradiction. Otherwise, suppose that  $y_1 \neq 0$ . Substituting f = x in (4.2) gives  $x = \gamma_1^2 x + \pi_{W_2} x + \langle x, y \rangle y + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} x$ . By employing the Riesz decomposition property of  $\mathcal{W}$ , we get  $\langle x, y \rangle y_1 = 0$ . Since  $y_1 \neq 0$ , then  $\langle x, y \rangle = 0$ . Thus, by setting f = y in (4.2), it yields

$$y = y + \gamma_1^2 \pi_{W_1} y + \gamma_1^2 \langle y, x \rangle x + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} y$$
  
=  $y + \gamma_1^2 \pi_{W_1} y + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} y.$ 

So, we obtain  $\pi_{W_i} y = 0$  for all  $i \neq 2$ . Since  $y \perp W_2$ , we induce that y = 0, which is again a contradiction. That is why  $x_2 \neq 0$ . Now, we proceed to prove the theorem.

(1) Taking f = x in (4.2) leads to

$$x = \gamma_1^2 x + \gamma_2^2 \pi_{W_2} x + \gamma_2^2 \langle x, y \rangle y + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} x$$

By an analogous method to the first part of the proof, we get  $\langle x, y \rangle = 0$ . Furthermore,  $x_i - \gamma_1^2 x_i - \gamma_i^2 \pi_{W_i} x = 0$  for  $i \ge 2$ , which ensures that  $\pi_{W_i} x = \frac{1 - \gamma_1^2}{\gamma_i^2} x_i$  for all  $i \ge 2$ . Condition (2) is proved similar to (1) by setting f = y in (4.2).

(3) We first note that,  $\gamma_i \neq 1$  for i = 1, 2. Indeed, if  $\gamma_1 = 1$ , then it follows from (1) that  $x \perp W_i$  for all  $i \geq 2$ . Since  $x \perp W_1$ , we get x = 0, which is a contradiction, similarly  $\gamma_2 \neq 1$ . Putting j = 1, i = 1, 2 and  $g \in W_1$  in Proposition 4.1(iii), we obtain  $g - \gamma_1^2 g - \gamma_2^2 \langle g, y \rangle y_1 = 0$  and  $\gamma_2^2 \pi_{W_2} g = 0$ . Hence,  $g = \frac{\gamma_2^2 \langle g, y \rangle}{1 - \gamma_1^2} y_1$  and  $W_1 \perp W_2$ . In particular,  $W_1 = \operatorname{span}\{y_1\}$ . Similarly, taking j = i = 2 and  $f \in W_2$  in Proposition 4.1(iii), gives  $f = \frac{\gamma_1^2 \langle f, x \rangle}{1 - \gamma_2^2} x_2$ , which yields that  $W_2 = \operatorname{span}\{x_2\}$ .

(4) Substituting  $f = x_2$  in (4.2) and applying (3), we get

$$x_{2} = \gamma_{1}^{2} \langle x_{2}, y_{1} \rangle \frac{y_{1}}{\|y_{1}\|^{2}} + \gamma_{1}^{2} \langle x_{2}, x \rangle x + \gamma_{2}^{2} x_{2} + \sum_{i=3}^{\infty} \gamma_{i}^{2} \pi_{W_{i}} x_{2}$$
$$= \gamma_{1}^{2} \langle x_{2}, x \rangle x + \gamma_{2}^{2} x_{2} + \sum_{i=3}^{\infty} \gamma_{i}^{2} \pi_{W_{i}} x_{2}.$$

It follows from the Riesz decomposition property of  $\mathcal W$  that

$$\begin{cases} \left(1 - \gamma_1^2 \langle x_2, x \rangle - \gamma_2^2\right) x_2 = 0, \\ \gamma_1^2 \langle x_2, x \rangle x_i + \gamma_i^2 \pi_{W_i} x_2 = 0, \quad (i \ge 3). \end{cases} \end{cases}$$

The first equation gives  $\langle x_2, x \rangle = \frac{1 - \gamma_2^2}{\gamma_1^2}$ . In addition, the second equation yields that

 $\pi_{W_i} x_2 = \frac{\gamma_2^2 - 1}{\gamma_i^2} x_i \text{ for all } i \ge 3.$ 

(5) Putting  $f = y_1$  in (4.2), the result is obtained by an analogous approach to (4). Condition (6) is also easily proved by using (1), (2), (4) and (5).

(7) Equation (4.2) can be reformulated as follows:

$$S_{\mathcal{W}_{\gamma}}f = f - \gamma_1^2 \langle f, x \rangle \, x - \gamma_2^2 \langle f, y \rangle \, y, \quad (f \in \mathcal{H}).$$

In (3), we observed that  $\gamma_i \neq 1$  for i = 1, 2. Now, according to the positivity of  $S_{W_{\gamma}}$ , we get

$$\gamma_1^2 |\langle f, x \rangle|^2 + \gamma_2^2 |\langle f, y \rangle|^2 \le ||f||^2, \quad (f \in \mathcal{H}).$$

Substituting x and y for f, it derives that  $\gamma_i < 1$  for i = 1, 2. Applying (2) and (5) we obtain

$$\begin{split} \sum_{i=3}^{\infty} \gamma_i^{-2} \|y_i\|^2 &= \sum_{i=3}^{\infty} \gamma_i^{-2} \langle y_i, y_i \rangle \\ &= \sum_{i=3}^{\infty} \gamma_i^{-2} \left\langle \frac{\gamma_i^2}{1 - \gamma_2^2} \pi_{W_i} y, \frac{\gamma_i^2}{\gamma_1^2 - 1} \pi_{W_i} y_1 \right\rangle \\ &= \frac{1}{(1 - \gamma_2^2) (\gamma_1^2 - 1)} \sum_{i=3}^{\infty} \gamma_i^2 \langle \pi_{W_i} y, \pi_{W_i} y_1 \rangle \\ &= \frac{1}{(1 - \gamma_2^2) (\gamma_1^2 - 1)} \left\langle \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} y, y_1 \right\rangle. \end{split}$$

Replacing f by y in (4.2) yields

$$\sum_{i=3}^{\infty} \gamma_i^{-2} \|y_i\|^2 = \frac{1}{(1-\gamma_2^2)(\gamma_1^2-1)} \left\langle \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} y, y_1 \right\rangle$$
  
$$= \frac{1-(\gamma_1^2+\gamma_2^2)}{(1-\gamma_2^2)(\gamma_1^2-1)} \left\langle y, y_1 \right\rangle$$
  
$$= \frac{\gamma_1^2+\gamma_2^2-1}{(1-\gamma_2^2)(1-\gamma_1^2)} \times \frac{1-\gamma_1^2}{\gamma_2^2} = \frac{\gamma_1^2+\gamma_2^2-1}{\gamma_2^2(1-\gamma_2^2)} \ge 0.$$
 (4.3)

Since  $\gamma_i < 1$  (i = 1, 2), it easily follows from the last inequality that  $\gamma_1^2 + \gamma_2^2 \ge 1$ . Therefore, we induce that  $1 \le \gamma_1^2 + \gamma_2^2 < 2$ , which completes the proof.

In light of the proof of Theorem 4.2(7) and analogous to (4.3), it can be concluded that

$$\sum_{i=3}^{\infty} \gamma_i^{-2} \|x_i\|^2 = \frac{\gamma_1^2 + \gamma_2^2 - 1}{\gamma_1^2 (1 - \gamma_1^2)} \ge 0.$$
(4.4)

Thus, if the 2-excess fusion frame  $\mathcal{V}$  is  $\gamma$ -scalable for some  $\gamma$  such that  $\gamma_1^2 + \gamma_2^2 = 1$ , then it follows from (4.3) and (4.4) that  $y_i = x_i = 0$  for all  $i \ge 3$ , and thereby  $x \in W_2$  and  $y \in W_1$ . In this case, we will obtain an equivalent condition for the weight-scalability of  $\mathcal{V}$ , see Corollary 5.6. Now, we provide two examples of 2-excess fusion frames that are weight-scalable.

**Example 4.3.** (1) Let  $\{e_i\}_{i=1}^3$  be the canonical orthonormal basis for  $\mathcal{H}_3$ . Consider  $\mathcal{V} = \{W_1 \oplus W_2, W_2 \oplus W_1, W_3\}$ , where  $W_i = \operatorname{span}\{e_i\}$  for  $1 \le i \le 3$ . Then  $\mathcal{V}$  is a 2-excess fusion frame containing the fusion orthonormal basis  $\{W_i\}_{i=1}^3$ . It is easy to see that  $\mathcal{V}$  is  $\gamma$ -scalable by  $\gamma_i = \sqrt{1/2}$  for i = 1, 2 and  $\gamma_3 = 1$ .

(2) Let  $\{e_i\}_{i=1}^4$  be the canonical orthonormal basis for  $\mathcal{H}_4$ . Take

$$V_{1} = W_{1} \oplus \operatorname{span}\left\{\frac{1}{2}e_{2} + \frac{\sqrt{3}}{2}e_{4}\right\},\$$

$$V_{2} = W_{2} \oplus \operatorname{span}\left\{\frac{1}{2}e_{1} + \frac{\sqrt{3}}{2}e_{3}\right\},\$$

$$W_{3} = \operatorname{span}\left\{\frac{1}{2}e_{2} - \frac{\sqrt{3}}{2}e_{4}\right\},\$$

$$W_{4} = \operatorname{span}\left\{\frac{1}{2}e_{1} - \frac{\sqrt{3}}{2}e_{3}\right\},\$$

where  $W_i = \text{span}\{e_i\}$  for i = 1, 2 and  $\mathcal{W} = \{W_i\}_{i=1}^4$  is a fusion Riesz basis for  $\mathcal{H}_4$ . Then, the 2-excess fusion frame  $\mathcal{V} = \{V_1, V_2, W_3, W_4\}$  is  $\gamma$ -scalable by  $\gamma_i = \sqrt{2/3}$  for  $1 \leq i \leq 4$ . More precisely,

$$S_{\mathcal{V}_{\gamma}} = \sum_{i=1}^{2} \gamma_{i}^{2} \pi_{W_{i}} + \gamma_{1}^{2} \pi_{\operatorname{span}\left\{\frac{1}{2}e_{2} + \frac{\sqrt{3}}{2}e_{4}\right\}} + \gamma_{2}^{2} \pi_{\operatorname{span}\left\{\frac{1}{2}e_{1} + \frac{\sqrt{3}}{2}e_{3}\right\}} + \gamma_{3}^{2} \pi_{\operatorname{span}\left\{\frac{1}{2}e_{2} - \frac{\sqrt{3}}{2}e_{4}\right\}} + \gamma_{4}^{2} \pi_{\operatorname{span}\left\{\frac{1}{2}e_{1} - \frac{\sqrt{3}}{2}e_{3}\right\}}.$$

Thus, for every  $(a, b, c, d) \in \mathcal{H}_4$ , we get

$$S_{\mathcal{V}_{\gamma}}(a, b, c, d) = 2/3 \left[ (a, b, 0, 0) + \left( 0, (b + \sqrt{3}d)/4, 0, (\sqrt{3}b + 3d)/4 \right) + \left( (a + \sqrt{3}c)/4, 0, (\sqrt{3}a + 3c)/4, 0 \right) + \left( 0, (b - \sqrt{3}d)/4, 0, (3d - \sqrt{3}b)/4 \right) + \left( (a - \sqrt{3}c)/4, 0, (3c - \sqrt{3}a)/4, 0 \right) \right] = (a, b, c, d),$$

as required.

In the preceding theorem, it was shown that the 2-excess fusion frame  $\mathcal{V}$  is not weight-scalable whenever  $x_2 = 0$  or  $y_1 = 0$ . The following theorem examines the weight-scalability of  $\mathcal{V}$  in other cases for x and y.

**Theorem 4.4.** Let  $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$  be a fusion Riesz basis for  $\mathcal{H}$ . If  $x_i = 0$   $(y_i = 0)$  for any  $i \geq 3$  and  $y_j \neq 0$   $(x_j \neq 0)$  for some  $j \geq 3$ , then the 2-excess fusion frame  $\mathcal{V}$  introduced in (4.1) is not weight-scalable.

Proof. For the sake of contradiction, assume that  $\mathcal{V}$  is  $\gamma$ -scalable. Then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=1}^{\infty}$  such that (4.2) holds. Suppose that  $x_i = 0$   $(i \ge 3)$ , so  $x_2 \ne 0$ . If  $y_1 = 0$ , then  $\gamma_1 = 1$ , by the proof of Theorem 4.2. Putting f = x in (4.2) leads to  $x = x + \gamma_2^2 x + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} x$ . It follows from the Riesz decomposition property of  $\mathcal{W}$  that  $\gamma_2 = 0$ , a contradiction. Otherwise, assume that  $y_1 \ne 0$ . Since x is a unit vector of  $W_2$ , Theorem 4.2(4) assures that  $\gamma_1^2 + \gamma_2^2 = 1$ . In view of (4.3), it implies that  $y_i = 0$  for all  $i \ge 3$ , which is in conflict with the assumption. Therefore,  $\mathcal{V}$  is not weight-scalable. A similar result is obtained whenever  $y_i = 0$   $(i \ge 3)$ , as desired.  $\Box$ 

**Example 4.5.** Let  $\{e_i\}_{i=1}^{\infty}$  be an orthonormal basis for  $\mathcal{H}$ . Consider

$$\begin{split} V_1 &= \mathrm{span}\{e_1, e_2\} \oplus \mathrm{span}\{e_3\}, \\ V_2 &= \mathrm{span}\{e_3\} \oplus \mathrm{span}\{\alpha_1 e_1 + \alpha_2 e_2 + \beta(\alpha_3 e_1 + \alpha_4 e_4) + \alpha_7 e_6\}, \\ V_3 &= \mathrm{span}\{\alpha_3 e_1 + \alpha_4 e_4, \alpha_5 e_2 + \alpha_6 e_5, e_6\}, \\ V_4 &= \mathrm{span}\{e_\ell\}_{\ell \geq 7}, \end{split}$$

where  $\alpha_i, \beta \in \mathbb{C}$  for all  $1 \leq i \leq 7$  and  $\alpha_i \neq 0$  for i = 4, 6. Then  $\mathcal{V} = \{V_i\}_{i=1}^4$  is a 2-excess fusion frame for  $\mathcal{H}$ . Assume that  $\mathcal{V}$  is  $\gamma$ -scalable, then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=1}^4$  such that

$$f = \sum_{i=1}^{4} \gamma_i^2 \pi_{V_i} f, \quad (f \in \mathcal{H}).$$

$$(4.5)$$

Putting  $f = e_3$  in (4.5) gives  $\gamma_1^2 + \gamma_2^2 = 1$ . Moreover, replacing f by  $e_5$  in (4.5) yields

 $e_5 = \gamma_3^2 \pi_{V_3} e_5 = \gamma_3^2 \pi_{\text{span}\{\alpha_5 e_2 + \alpha_6 e_5\}} e_5$ 

$$=\frac{\gamma_3^2}{\alpha_5^2+\alpha_6^2}\left\langle e_5, \alpha_5 e_2+\alpha_6 e_5\right\rangle \left(\alpha_5 e_2+\alpha_6 e_5\right) =\frac{\gamma_3^2}{\alpha_5^2+\alpha_6^2}\left(\alpha_5 \alpha_6 e_2+\alpha_6^2 e_5\right).$$

Since  $\alpha_6 \neq 0$ , then we get  $\alpha_5 = 0$ . Now, by setting  $f = e_2$  in (4.5), we have

$$\begin{aligned} e_2 &= \gamma_1^2 \pi_{V_1} e_2 + \gamma_2^2 \pi_{V_2} e_2 \\ &= \gamma_1^2 e_2 + \gamma_2^2 \pi_{\text{span}\{(\alpha_1 + \beta \alpha_3)e_1 + \alpha_2 e_2 + \beta \alpha_4 e_4 + \alpha_7 e_6\}} e_2 \\ &= \left(\gamma_1^2 + \frac{\gamma_2^2 \alpha_2^2}{c}\right) e_2 + \frac{\gamma_2^2 \alpha_2}{c} \left((\alpha_1 + \beta \alpha_3)e_1 + \beta \alpha_4 e_4 + \alpha_7 e_6\right), \end{aligned}$$

where  $c = (\alpha_1 + \beta \alpha_3)^2 + \alpha_2^2 + (\beta \alpha_4)^2 + \alpha_7^2$ . If  $\alpha_2 = 0$ , then it follows that  $\gamma_1 = 1$  and thereby  $\gamma_2 = 0$ , which is a contradiction. Otherwise, assume that  $\alpha_2 \neq 0$ . Since  $\alpha_4 \neq 0$ , then  $\alpha_7 = \alpha_1 = \beta = 0$ . Hence, by substituting  $f = e_1$  in (4.5) gives

$$e_{1} = \gamma_{1}^{2} \pi_{V_{1}} e_{1} + \gamma_{3}^{2} \pi_{V_{3}} e_{1}$$

$$= \gamma_{1}^{2} e_{1} + \gamma_{3}^{2} \pi_{\text{span}} \{\alpha_{3} e_{1} + \alpha_{4} e_{4}\} e_{1}$$

$$= \gamma_{1}^{2} e_{1} + \frac{\gamma_{3}^{2}}{\alpha_{3}^{2} + \alpha_{4}^{2}} \langle e_{1}, \alpha_{3} e_{1} + \alpha_{4} e_{4} \rangle (\alpha_{3} e_{1} + \alpha_{4} e_{4})$$

$$= \left(\gamma_{1}^{2} + \frac{\gamma_{3}^{2} \alpha_{3}^{2}}{\alpha_{3}^{2} + \alpha_{4}^{2}}\right) e_{1} + \frac{\gamma_{3}^{2} \alpha_{3} \alpha_{4}}{\alpha_{3}^{2} + \alpha_{4}^{2}} e_{4}.$$

It derives that  $\alpha_3 = 0$ , as  $\alpha_4 \neq 0$ . Thus,  $\gamma_1 = 1$  and thereby  $\gamma_2 = 0$ , which is again a contradiction. Therefore,  $\mathcal{V}$  is not weight-scalable. It is worthwhile to mention that the result can be obtained directly from Theorem 4.4.

In Example 4.3(1), a weight-scalable 2-excess fusion frame is provided for  $\mathcal{H}_3$  that contains an orthogonal fusion Riesz basis. The subsequent result shows that the orthogonality of the Riesz part is a necessary condition for the weight-scalability of certain 2-excess fusion frames in  $\mathcal{H}_3$ .

**Corollary 4.6.** Let the 2-excess fusion frame  $\mathcal{V}$  given by (4.1) be weight-scalable for  $\mathcal{H}_3$ . Then every fusion Riesz basis contained in  $\mathcal{V}$  is orthogonal.

*Proof.* Suppose that  $\mathcal{V} = \{V_i\}_{i=1}^n \ (n \leq 3)$  is a  $\gamma$ -scalable 2-excess fusion frame containing a fusion Riesz basis  $\mathcal{W}$ . Then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=1}^n$  such that

$$f = \sum_{i=1}^{n} \gamma_i^2 \pi_{V_i} f, \quad (f \in \mathcal{H}).$$

$$(4.6)$$

As a result of the proof of Theorem 3.6, the excess elements cannot be contained in only one subspace of  $\mathcal{W}$ , see also Corollary 5.2. By keeping this fact in mind, let  $\{e_i\}_{i=1}^3$  be the canonical orthonormal basis for  $\mathcal{H}_3$ . Notice that, Theorem 4.2(3) ensures that  $\mathcal{W}$  must have more than two subspaces, as otherwise it is incomplete. Hence,  $V_i$ 's are represented as

$$V_1 = W_1 \oplus \operatorname{span}\{x\},$$
  
$$V_2 = W_2 \oplus \operatorname{span}\{y\},$$

1	0
т	J

 $V_3 = W_3,$ 

where x, y are the redundant elements of  $\mathcal{V}$ . Then, according to Theorem 4.2(3) and without losing the generality,  $V_i$ 's can be rewritten as follows:

$$V_{1} = \operatorname{span}\{e_{1}\} \oplus \operatorname{span}\{\alpha e_{2} + \alpha' u\},$$

$$V_{2} = \operatorname{span}\{e_{2}\} \oplus \operatorname{span}\{\beta e_{1} + \beta' u\},$$

$$V_{3} = \operatorname{span}\{u\},$$

$$(4.7)$$

where  $\alpha, \alpha', \beta, \beta' \in \mathbb{C}$  and  $\alpha\beta \neq 0$ . Moreover, the redundant vectors and  $u = (u_1, u_2, u_3)$  are unit vectors of  $\mathcal{H}_3$  such that  $u_3 \neq 0$ . In light of (4.7), we infer that

$$\alpha' u_1 = \beta' u_2 = 0. \tag{4.8}$$

Furthermore, by Theorem 4.2(1) we have  $\langle \alpha e_2 + \alpha' u, \beta e_1 + \beta' u \rangle = 0$ , which yields  $\alpha'\beta' = 0$ . First, assume that  $\alpha' = \beta' = 0$ , then (4.7) is reformulated as follows:

$$V_1 = V_2 = \operatorname{span}\{e_1, e_2\}, V_3 = \operatorname{span}\{u\}.$$

Putting  $f = e_3$  in (4.6) leads

$$e_3 = \gamma_3^2 \pi_{V_3} e_3 = \gamma_3^2 \langle e_3, u \rangle u = \gamma_3^2 \left( u_1 u_3 e_1 + u_2 u_3 e_2 + u_3^2 e_3 \right).$$

Since  $u_3 \neq 0$ , we get  $u_1 = u_2 = 0$ , that means  $\mathcal{W}$  is orthogonal. Now, suppose that  $\alpha' \neq 0$  and  $\beta' = 0$ , then  $u_1 = 0$ , by (4.8). Thus, (4.7), without losing the generality, is represented in the following manner:

$$V_1 = \operatorname{span}\{e_1\} \oplus \operatorname{span}\{ae_2 + be_3\},$$
  
$$V_2 = \operatorname{span}\{e_1, e_2\}, \ V_3 = \operatorname{span}\{u_2e_2 + u_3e_3\},$$

where  $a^2 + b^2 = u_2^2 + u_3^2 = 1$  and  $b \neq 0$ . Substituting  $f = e_i$   $(1 \le i \le 3)$  in (4.6) gives

$$\begin{cases} (1) \ \gamma_1^2 + \gamma_2^2 = 1, \\ (2) \ a^2 \gamma_1^2 + \gamma_2^2 + u_2^2 \gamma_3^2 = 1, \\ (3) \ b^2 \gamma_1^2 + u_3^2 \gamma_3^2 = 1, \\ (4) \ ab \gamma_1^2 + u_2 u_3 \gamma_3^2 = 0. \end{cases}$$

Summing equations (2) and (3) and using (1), we get  $\gamma_3 = 1$ . If  $u_2 \neq 0$ , then by (2), (3) and (4) we obtain

$$\gamma_2^2 = 1 - u_2^2 - a^2 \gamma_1^2 = u_3^2 - \frac{a^2 b^2 \gamma_1^4}{b^2 \gamma_1^2}$$

$$= u_3^2 - \frac{u_2^2 u_3^2}{1 - u_3^2} = u_3^2 - u_3^2 = 0,$$

a contradiction. Thus,  $u_2 = 0$ , which implies that  $\mathcal{W}$  is orthogonal. Also, in the case where  $\alpha' = 0$  and  $\beta' \neq 0$ , the desired result is obtained by a similar argument.

The next example shows that the converse of Corollary 4.6 is not valid, in general. Obviously, the converse holds whenever  $V_1 = V_2$  in (4.7).

**Example 4.7.** Let  $\{e_i\}_{i=1}^3$  be the canonical orthonormal basis for  $\mathcal{H}_3$  and  $W_i = \text{span}\{e_i\}$  for  $1 \leq i \leq 3$ . Then  $\mathcal{V} = \{W_1 \oplus W_2, W_2 \oplus W_3, W_3\}$  is a 2-excess fusion frame containing the fusion orthonormal basis  $\{W_i\}_{i=1}^3$ . However,  $\mathcal{V}$  is not weight-scalable by Theorem 4.4.

## 5 Weight-Scalability: General Case

Finally, we would like to discuss the weight-scalability of fusion frames with higher excess. First of all, we note that tight fusion frames are clearly weight-scalable. Now, we consider fusion frames in which a specific subspace of a fusion Riesz basis is repeated. In the other words, let

$$\mathcal{V} = \{W_\ell, \dots, W_\ell\} \cup \{W_i\}_{i \in I}$$

be a fusion frame with the Riesz part  $\mathcal{W} = \{W_i\}_{i \in I}$  and the redundant subspace  $W_{\ell}$  ( $\ell \in I$ ) occurs *n* times. Then, a direct computation shows that the following conditions are equivalent:

(i)  $\mathcal{V}$  is  $\gamma$ -scalable,  $\gamma = {\gamma_i}_{i \in I}$ .

(ii)  $\mathcal{W}$  is  $\omega$ -scalable,  $\omega_i = 1, i \in I$ .

(iii)  $\mathcal{W}$  is orthogonal,

where

$$\gamma_i = \begin{cases} 1, & i \neq \ell, \\ \frac{1}{\sqrt{n+1}}, & i = \ell. \end{cases}$$

This indicates that weights play an important role in the scalability of such fusion frames. The same result can be extended to the fusion frames with more redundant subspaces. Inspired by Theorem 3.6, we now obtain an analogous result for the weight-scalability of k-excess fusion frames of the form

$$\mathcal{V} = \{V_\ell\}_{\ell=-k}^{-1} \cup \mathcal{W}, \quad (k \ge 2), \tag{5.1}$$

where  $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$  is a fusion Riesz basis,  $V_{\ell} = \operatorname{span}\{x^{(\ell)}\}\)$  and the redundant element  $x^{(\ell)} = \sum_{i=1}^{\infty} x_i^{(\ell)}\)$  is a unit vector of  $\mathcal{H}\)$  such that  $x_i^{(\ell)} \in W_i\)$  for each  $i \ge 1$ . **Proposition 5.1.** Let  $\mathcal{V}\)$  be the k-excess fusion frame defined by (5.1) and  $\mathcal{W}\)$  a fusion

**Proposition 5.1.** Let V be the k-excess fusion frame defined by (5.1) and W a fusion Riesz basis for  $\mathcal{H}$ . Then the following conditions are equivalent:

(i) 
$$\mathcal{V}$$
 is  $\gamma$ -scalable,  $\gamma = \{\gamma_i\}_{i=-k}^{\infty}$ .  
(ii)  $\sum_{\ell=-k}^{-1} \gamma_{\ell}^2 \pi_{W_i} S_{W_{\gamma}}^{-1} \pi_{V_{\ell}} \pi_{W_j} = (\gamma_i^{-2} \delta_{i,j} - \pi_{W_i}) \pi_{W_j}$ , for all  $i, j \ge 1$ .  
(iii)  $\sum_{\ell=-k}^{-1} \gamma_{\ell}^2 \langle \pi_{W_j} f, x^{(\ell)} \rangle x_i^{(\ell)} = (\delta_{i,j} - \gamma_i^2 \pi_{W_i}) \pi_{W_j} f$ , for all  $f \in \mathcal{H}$ .

It is worthy of note that by rearranging the excess elements among the subspaces of  $\mathcal{W}$ , one can achieve similar results to the above proposition. For instance, if the k-excess fusion frame  $\mathcal{V}$  in Proposition 5.1 is represented as

$$\mathcal{V} = V_0 \cup \mathcal{W},$$

where  $V_0 = \bigoplus_{\ell=-k}^{-1} V_{\ell}$ , then (ii) and (iii) are derived as follows:

(ii) 
$$\gamma_0^2 \pi_{W_i} S_{W_\gamma}^{-1} \pi_{V_0} \pi_{W_j} = \left(\gamma_i^{-2} \delta_{i,j} - \pi_{W_i}\right) \pi_{W_j}$$
, for all  $i, j \ge 1$ .  
(iii)  $\gamma_0^2 \sum_{\ell=-k}^{-1} \langle \pi_{W_j} f, x^{(\ell)} \rangle x_i^{(\ell)} = \left(\gamma_i^2 \delta_{i,j} - \gamma_i^2 \pi_{W_i}\right) \pi_{W_j} f$ , for all  $f \in \mathcal{H}$ .

The aim here is not to provide an exhaustive list of analogous conditions for the weight-scalability, as there are numerous situations in which the excess elements may be presented. The next result can be noted as a consequence of the proof of Theorem 3.6.

**Corollary 5.2.** Every k-excess fusion frame, whose all the redundant elements are in only one subspace of its Riesz part is not weight-scalable for any  $k \ge 2$ .

In the sequel, we focus on the weight-scalability of k-excess fusion frames whose the redundant vectors are located in more than one subspace of their Riesz part. **Theorem 5.3.** Let  $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$  be a fusion Riesz basis for  $\mathcal{H}$ . Take

$$\begin{cases} V_1 = W_1 + Z_2, \\ V_2 = W_2 + Z_1, \end{cases}$$

where  $Z_i$  is a non zero closed subspace of  $W_i$  for i = 1, 2 and  $\mathcal{V} = \{V_1, V_2, W_i\}_{i=3}^{\infty}$  constitutes a fusion frame for  $\mathcal{H}$ . Then the following statements hold.

- (i) If  $\mathcal{V}$  is weight-scalable, then  $Z_i = W_i$  for i = 1, 2.
- (ii) If  $Z_i = W_i$  for i = 1, 2,  $\{W_j\}_{j=3}^{\infty}$  is orthogonal and  $(W_1 + W_2) \perp W_j$  for all  $j \ge 3$ , then  $\mathcal{V}$  is weight-scalable.

*Proof.* (i) Since  $\mathcal{V}$  is  $\gamma$ -scalable, then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=1}^{\infty}$  such that (4.2) holds. It implies that  $\gamma_1^2 + \gamma_2^2 = 1$ , as  $V_1 \cap V_2 \neq \{0\}$ . By contrary, assume that  $Z_1 \subsetneq W_1$ . It is not difficult to see that  $W_1 \cap Z_1^{\perp} \neq \{0\}$ . Hence, there exists a non zero vector  $g \in W_1 \cap Z_1^{\perp}$ , so that

$$g = \gamma_1^2 g + \gamma_2^2 \pi_{V_2} g + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} g$$
$$= \gamma_1^2 g + \gamma_2^2 \pi_{V_2} g,$$

where the last equality is due to the Riesz decomposition property of  $\mathcal{W}$ . Hence,  $\pi_{V_2}g = g$  and thereby  $g \in V_2 = W_2 + Z_1$ . Thus, there exist  $w \in W_2$  and  $z \in Z_1$  such that g = w + z. Since  $g - z = w \in W_1 \cap W_2$ , we conclude that g - z = 0. So  $g = z \in Z_1$ , which is a contradiction. Therefore,  $Z_1 = W_1$  and by a similar argument  $Z_2 = W_2$  as well.

(ii) Suppose that  $V_i = W_1 + W_2$  for i = 1, 2. Taking  $\gamma_1, \gamma_2$  satisfying  $\gamma_1^2 + \gamma_2^2 = 1$ and  $\gamma_j = 1$  for  $j \ge 3$ , we get

$$S_{\mathcal{V}_{\gamma}} = (\gamma_1^2 + \gamma_2^2) \pi_{(W_1 + W_2)} + \sum_{j=3}^{\infty} \gamma_j^2 \pi_{W_j}$$
  
=  $\pi_{(W_1 + W_2)} + \sum_{j=3}^{\infty} \pi_{W_j} = \pi_{(W_1 + W_2)} + \pi_{\bigoplus_{j=3}^{\infty} W_j} = I_{\mathcal{H}}.$ 

**Corollary 5.4.** Let  $\mathcal{W}$  be an orthogonal fusion Riesz basis for  $\mathcal{H}$ . Then  $\mathcal{V}$ , introduced in Theorem 5.3 is weight-scalable if and only if  $Z_i = W_i$  for i = 1, 2. **Example 5.5.** Consider the 1-excess fusion frame  $\mathcal{V}$  introduced in Example 3.7. Take

**Example 5.5.** Consider the 1-excess rusion frame  $\nu$  introduced in Example 5.7. Take  $Z_1 = \overline{\text{span}}\{e_i\}_{i \ge -n}$  and  $Z_2 = \overline{\text{span}}\{e_i\}_{i \le m}$ , for every  $n, m \in \mathbb{N}$ . Then  $\mathcal{Z} = \{Z_i\}_{i=1}^2$  is a dual fusion frame of  $\mathcal{V}$  with  $e(\mathcal{Z}) = n + m + 1$ , see [1] for more details. Clearly,  $\mathcal{Z}$  can be expressed as

$$\begin{cases} Z_1 = \overline{\operatorname{span}}\{e_i\}_{i\geq 1} \oplus \operatorname{span}\{e_i\}_{i=-n}^0, \\ Z_2 = \overline{\operatorname{span}}\{e_i\}_{i<1} \oplus \operatorname{span}\{e_i\}_{i=1}^m. \end{cases}$$

Thus, Corollary 5.4 assures that  $\mathcal{Z}$  is weight-scalable if and only if  $n = m = \infty$ .

The following result states that under which conditions the 2-excess fusion frame  $\mathcal{V}$  given by (4.1) is weight-scalable. The proof follows immediately from Theorem 4.2, Theorem 5.3 and Corollary 5.4.

**Corollary 5.6.** Let  $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$  be a fusion Riesz basis for  $\mathcal{H}$ ,  $x \in W_2$  and  $y \in W_1$ . If the 2-excess fusion frame  $\mathcal{V}$  defined by (4.1) is weight-scalable, then  $V_1 = V_2 = \operatorname{span}\{x, y\}$ . In particular, the converse holds if  $\{W_j\}_{j=3}^{\infty}$  is orthogonal and  $V_1 \perp W_j$  for all  $j \geq 3$ .

Obviously, if W is orthogonal in the above corollary, then  $\mathcal{V}$  is weight-scalable if and only if  $V_1 = V_2 = \operatorname{span}\{x, y\}$ . Notice that Theorem 4.2(3) can be generalized to every k-excess fusion frame for k > 2 whose the redundant vectors are in only two subspaces of its Riesz part.

**Theorem 5.7.** Let  $\mathcal{V} = \{V_i\}_{i=1}^2 \cup \{W_i\}_{i=1}^{\infty}$  be a weight-scalable fusion frame for  $\mathcal{H}$  such that  $W_i \subsetneq V_i$  for i = 1, 2 and  $\{W_i\}_{i=1}^{\infty}$  is a fusion Riesz basis. Then  $W_1 \perp W_2$ . Moreover, the following statements hold:

(i) If  $\gamma_1 = 1$ , then  $\gamma_2 = 1$ ,  $V_1 \perp W_2$  and  $W_2 \perp W_i$  for all  $i \geq 3$ .

(ii) If  $\gamma_2 = 1$ , then  $\gamma_1 = 1$ ,  $V_2 \perp W_1$  and  $W_1 \perp W_i$  for all  $i \geq 3$ .

(iii) If  $\gamma_1 \neq 1$  ( $\gamma_2 \neq 1$ ), then dim $W_1 \leq \dim(V_2 \setminus W_2)$  (dim $W_2 \leq \dim(V_1 \setminus W_1)$ ).

*Proof.* Without losing the generality, we may choose subspaces  $Z_1$  and  $Z_2$  such that  $\dim Z_i = \dim(V_i \setminus W_i)$  for i = 1, 2 and

$$\begin{cases} V_1 = W_1 \oplus Z_1, \\ V_2 = W_2 \oplus Z_2. \end{cases}$$

റ	•
$\Delta$	э

Furthermore, the first and second components of the elements of  $Z_1$  and  $Z_2$ , respectively, are equal to zero in the Riesz decomposition. Since  $\mathcal{V}$  is  $\gamma$ -scalable, then there exists a sequence of weights  $\gamma = \{\gamma_i\}_{i=1}^{\infty}$  such that (4.2) holds. For every  $g \in W_1$  in (4.2), we have

$$g = \gamma_1^2 g + \gamma_2^2 \pi_{W_2} g + \gamma_2^2 \pi_{Z_2} g + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} g.$$
 (5.2)

Applying the Riesz decomposition property of  $\mathcal{W}$ , we infer that  $\gamma_2^2 \pi_{W_2} g = 0$  and so  $W_1 \perp W_2$ . Now, for the "moreover" part:

(i) We note that (5.2) also holds for all  $g \in V_1$ . Thus, if  $\gamma_1 = 1$ , then the Riesz decomposition property of  $\mathcal{W}$  implies that  $V_1 \perp W_2$ . Hence, putting a non zero vector  $f \in W_2$  in (4.2), we obtain  $f = \gamma_2^2 f + \sum_{i=3}^{\infty} \gamma_i^2 \pi_{W_i} f$ . It follows that  $\gamma_2 = 1$  and  $W_2 \perp W_i$  for all  $i \geq 3$ . The proof of (ii) is analogous to that of (i).

(iii) If  $\gamma_1 \neq 1$ , then (5.2) gives  $g = \gamma_1^2 g + \gamma_2^2 \pi_{Z_2} g$ . Hence, we get  $g = \frac{\gamma_2^2}{1 - \gamma_1^2} \pi_{Z_2} g$ , which ensures that

$$\dim W_1 \le \dim Z_2 = \dim (V_2 \smallsetminus W_2).$$

In the case where  $\gamma_2 \neq 1$ , by means of an analogous argument, it is proved that  $\dim W_2 \leq \dim Z_1 = \dim (V_1 \setminus W_1)$ . This completes the proof.

Suppose that  $\{e_i\}_{i=1}^{\infty}$  is an orthonormal basis for  $\mathcal{H}$ ,  $W_1 = \operatorname{span}\{e_1\}$  and  $W_2 = \operatorname{span}\{e_2, e_3\}$ . Consider

$$V_1 = W_1 \oplus \overline{\text{span}} \{e_2, e_3, e_{2i} + e_{2i+1}\}_{i=2}^{\infty}$$
  

$$V_2 = W_2 \oplus \overline{\text{span}} \{e_1, e_{2i} - e_{2i+1}\}_{i=2}^{\infty},$$
  

$$W_3 = \text{span} \{e_i\}_{i=4}^{\infty},$$

where  $\{W_i\}_{i=1}^3$  is a fusion orthonormal basis. Then  $\mathcal{V} = \{V_1, V_2, W_3\}$  constitutes an infinite-excess 2-tight fusion frame for  $\mathcal{H}$ . Hence, it follows that  $\mathcal{V}$  is  $\gamma$ -scalable by  $\gamma_i = \sqrt{1/2}$  for  $1 \leq i \leq 3$ . Obviously,  $\dim W_i < \dim(V_j \setminus W_j)$  for distinct indexes i, j = 1, 2, which confirms Theorem 5.7.

Conflict-of-interest. This work does not have any conflicts of interest.

## References

- E. Ameli, A. Arefijamaal and F. Arabyani Neyshaburi, Excess of fusion frames: A comprehensive approach, arXiv:2408.03179 (2024). https://doi.org/10.48550/arXiv.2408.03179
- [2] F. Arabyani Neyshaburi and A. Arefijamaal, Weaving Hilbert space fusion frames, Rocky Mountain J. Math. 51(1) (2021), 55-66.
- [3] F. Arabyani Neyshaburi, A. Arefijamaal, R. Farshchian and R. A. Kamyabi-Gol, Norm retrieval algorithms: A new frame theory approach, *Math. Meth. Appl. Sci.* 47(9) (2024), 7111-7132.

- [4] A. Arefijamaal and F. Arabyani Neyshaburi, Some properties of alternate duals and approximate alternate duals of fusion frames, *Turkish J. Math.* 41(5) (2017), 1191-1203.
- [5] A. Arefijamaal, F. Arabyani Neyshaburi and M. Shamsabadi, On the duality of frames and fusion frames, *Hacet. J. Math. Stat.* 47(1) (2018), 1-10.
- [6] P. Balazs, J. P. Antoine and A. Grybos, Weighted and controlled frames: Mutual relationship and first numerical properties, *Int. J. Wavelets Multiresolut. Inf. Process.* 8(1) (2010), 109-132.
- [7] J. Cahill, M. Fickus, D. G. Mixon, M. J. Poteet and N. K. Strawn, Constructing finite frames of a given sprectrum and set of lengths, *Appl. Comput. Harmon. Anal.* 35(1) (2013), 52-73.
- [8] P.G. Casazza, M. Fickus, D. Mixon, Y. Wang and Z. Zhou, Constructing tight fusion frames, Appl. Comput. Harmon. Anal. 30 (2011), 175-187.
- [9] P. Casazza and X. Chen, Frame scalings: A condition number approach, *Linear Algebra Appl.* 523 (2017), 152-168.
- [10] P. Casazza and G. Kutyniok, Frame of subspaces. Contemp. Math. 345 (2004), 87-114.
- [11] P. Găvruţa, On the duality of fusion frames, J. Math. Anal. Appl. 333 (2007), 871-879.
- [12] B. Heydarpour, A. Arefijamaal and F. Ghaani, Characterization of dual scalable frames, *Complex Anal. Oper. Theory.* 18(3) (2024).
- [13] G. Kutyniok, K. A. Okoudjou, F. Philipp and E. K. Tuley, Scalable frames, Linear Algebra Appl. 438 (2013), 2225-2238.
- [14] G. Kutyniok, K. A. Okoudjou and F. Philipp, Scalable frames and convex geometry, *Contemp. Math.* 626 (2014), 19-32.
- [15] A. R. Mariano and D. Stojanoff, Some properties of frames of subspaces obtained by operator theory methods, J. Math. Anal. Appl. 343 (2008), 366-378.
- [16] N. Q. Nga, Some results on fusion frames and g-frames, *Results Math*, 73(2) (2018), 1-9.
- [17] E. Osgooei and A. Arefijammal, Compare and contrast between duals of fusion and discrete frames, Sahand Commun. Math. Anal. 08(1) (2017), 83-96.
- [18] A. Rahimi, S. Moayyadzadeh, Scalable Fusion Frames, Iran. J. Sci. Technol. Trans. A Sci. (2025), 1-7. https://doi.org/10.1007/s40995-024-01765-y

- [19] A. Rahimi, G. Zandi and B. Daraby, Redundancy of fusion frames in Hilbert spaces, Complex Anal. Oper. Theory, 10(3) (2016), 545-565.
- [20] M. Shamsabadi, A. Arefijamaal and P. Balazs, The invertibility of U-fusion cross Gram matrices of operators. *Mediterr. J. Math.* 17(2) (2020).