LIE ALGEBRAS OF QUOTIENT GROUPS

DAVID MIYAMOTO

ABSTRACT. We give conditions on a diffeological group G and a normal subgroup Hunder which the quotient group G/H differentiates to a Lie algebra, and $\text{Lie}(G/H) \cong$ Lie(G)/Lie(H). Our Lie functor is derived from the tangent structure on elastic diffeological spaces introduced by Blohmann. The requisite conditions on G and H hold when G is a convenient infinite-dimensional Lie group and all iterated tangent bundles T^kH are initial subgroups of T^kG ; in particular, G may be finite-dimensional, or H may be countable. As an application, we integrate some classically non-integrable Banach-Lie algebras to diffeological groups.

Contents

| 1. Introduction | | 2 |
|--|---|----|
| 1.1. | Summary of results | 2 |
| 1.2. | Remarks on the literature | 3 |
| 1.3. | Structure of the paper | 4 |
| 1.4. | Acknowledgements | 5 |
| 2. A tangent structure | | 5 |
| 2.1. | Elastic diffeological spaces | 5 |
| 2.2. | Elastic diffeological groups | 9 |
| 3. Di | ffeological groupoids | 11 |
| 3.1. | Tangent bundles of orbit spaces | 12 |
| 3.2. | Elasticity of quotient groups | 15 |
| 4. The convenient setting and diffeology | | 17 |
| 4.1. | Convenient diffeological vector spaces | 17 |
| 4.2. | From convenient manifolds to diffeology | 22 |
| 5. Int | 25 | |
| 5.1. | Diffeological central extensions | 26 |
| 5.2. | Integrating convenient Lie algebras | 32 |
| Appendix A. Tangent categories | | 35 |
| A.1. | First definitions | 35 |
| A.2. | The Cartesian structure | 38 |
| A.3. | Groups in tangent categories | 39 |
| References | | 41 |

Date: February 17, 2025.

2020 Mathematics Subject Classification. Primary 22E65, 58A40; Secondary 18F40. Key words and phrases. Tangent structure, diffeology, convenient analysis.

1. INTRODUCTION

One of the fundamental results in the study of Lie groups is Lie's third theorem, proved in its present form by Cartan.

Theorem (Lie-Cartan). For every finite-dimensional Lie algebra \mathfrak{g} , there is some finitedimensional Lie group G whose Lie algebra is \mathfrak{g} .

Of course, not every Lie algebra is finite-dimensional, and Lie's third theorem famously fails in the infinite-dimensional case: van Est and Korthagen [EK64] gave the first example of a Banach-Lie algebra which is not the Lie algebra of any Banach-Lie group. We call such Lie algebras \mathfrak{g} non-enlargeable, and otherwise say G integrates or enlarges \mathfrak{g} . Van Est and Korthagen justify their example by identifying a topological obstruction to enlargeability within the centre \mathfrak{z} of \mathfrak{g} . Precisely, they showed that a Banach-Lie algebra \mathfrak{g} is enlargeable if and only if a relevant subgroup $\Pi \leq (\mathfrak{z}, +)$ is topologically discrete. In that case, their construction of an enlarging G requires the Banach-Lie group structure on \mathfrak{z}/Π , and the fact that its Lie algebra is isomorphic to \mathfrak{z} .

While it is only a manifold if Π is topologically discrete, the quotient \mathfrak{z}/Π is always a well-behaved diffeological group. Souriau [Sou80] introduced diffeological groups, and later diffeological spaces,¹ to axiomatize smoothness for non-Lie groups. The structure of a diffeological space is lightweight: a diffeology on a set X is a collection of maps $U \xrightarrow{p} X$ from open subsets of Cartesian spaces, called plots, satisfying three axioms.² Diffeological spaces form a category \mathcal{D} flg which contains the category of Banach manifolds. But unlike a category of manifolds, \mathcal{D} flg is a quasi-topos, meaning it is complete, co-complete, and locally Cartesian closed. In particular, if G is any diffeological group, and H is any normal subgroup, then both H and G/H inherit natural diffeologies, whence \mathfrak{z}/Π is a diffeological group. This motivates us to integrate Banach-Lie algebras to diffeological groups. In doing so, we accomplish three objectives, which we now summarize.

1.1. Summary of results. First, we restrict our attention to the subcategory of elastic diffeological groups, introduced by Blohmann [Blo24a]. Whereas the existence of a Lie functor defined for all diffeological groups is unknown, we have one on the full subcategory of elastic groups:

Lie : {elastic Lie groups} \rightarrow {elastic Lie algebras}.

This Lie functor invokes a tangent functor $T : \mathcal{D}flg \to \mathcal{D}flg$, which we define as the left Kan extension of the usual tangent functor on manifolds. The main result announced in [Blo24a], with details to appear [Blo25], is that when restricted to the subcategory of elastic spaces, T participates in a tangent structure in the sense of Rosický [Ros84]. It is the tangent structure which imparts $\text{Lie}(G) := T_e G$ with its Lie bracket, a fact which has been established in broad generality by Blohmann and Aintablian [AB24]. We show that, under some conditions, the Lie functor is compatible with quotients.

¹K. T. Chen [Che77] gave a similar notion.

²(i) locally constant maps are plots; (ii) if $U' \xrightarrow{F} U$ is smooth, and $U \xrightarrow{p} X$ is a plot, so is pF; (iii) if p is locally a plot, it is a plot. One may recognize that the last two conditions describe a sheaf, and the first states that the sheaf is concrete.

Theorem I. Let G be an elastic group, with a normal subgroup H carrying its induced diffeology. Denote the inclusion by $\iota : H \hookrightarrow G$. If the iterated tangents $T^k \iota : T^k H \to T^k G$ are inductions³ for all $k \ge 0$, then both H and G/H are elastic, and $\text{Lie}(G/H) \cong \text{Lie}(G)/\text{Lie}(H)$.

The theorem applies, for instance, when H is diffeologically discrete (e.g. countable), or if G is any finite-dimensional Lie group.

Second, we embed convenient Lie groups, in the sense of Kriegl and Michor [KM97], into the category of elastic diffeological groups. These infinite-dimensional Lie groups are modelled by convenient locally convex topological vector spaces, and the category of convenient manifolds includes the categories of Banach and Fréchet manifolds. We have found that convenience and elasticity are intimately related. More precisely, we call a diffeological vector space V tangent stable if the map

$$V \times V \to TV$$
, $(u, v) \mapsto \frac{d}{dt}\Big|_{t=0} u + tv$

is an isomorphism. Tangent stability implies elasticity, and we show:

Theorem II. A Hausdorff bornological locally convex topological vector space is convenient if and only if, when viewed as a diffeological space, it is tangent-stable.

We conclude that our Lie functor, and the usual one on convenient Lie groups, coincide. Finally, we treat enlargeability of convenient Lie algebras. We follow the strategy used by Cartan, and extended by Neeb [Nee02], of integrating central extensions of Lie algebras. Let \mathfrak{g} be a convenient Lie algebra, which we assume integrates to a connected and simply-connected convenient Lie group G. Let \mathfrak{a} be an abelian convenient Lie algebra, and take some topologically split central extension $\mathfrak{a} \hookrightarrow \hat{\mathfrak{g}} \twoheadrightarrow \mathfrak{g}$. There is a group of periods $\Pi \leq (\mathfrak{a}, +)$ associated to the central extension. We prove:

Theorem III. A topologically split central extension $\mathfrak{a} \hookrightarrow \hat{\mathfrak{g}} \twoheadrightarrow \mathfrak{g}$, in the presence of a connected and simply-connected convenient Lie group G with $\text{Lie}(G) = \mathfrak{g}$, integrates to a central extension of elastic groups if the period group $\Pi \leq (\mathfrak{a}, +)$ is diffeologically discrete, in which case we have:



As an application, we integrate some non-enlargeable Lie algebras, such as the classical examples given by van Est and Korthagen [EK64] and Douady and Lazard [DL66].

1.2. **Remarks on the literature.** There is a litany of diffeological tangent functors. They broadly fall into three approaches: the internal, representing tangent vectors by curves with equality probed by representations in higher dimensional plots; the external,

³Initial maps in the category of diffeological spaces.

identifying tangent vectors with derivations of rings of functions; and the mixed, representing tangent vectors by curves, but testing equality using smooth functions. We have traced some first-appearances of the various functors below.

| Internal | External | Mixed | | |
|--|--------------------------|-----------------------|--|--|
| Losik [Los92] | Iglesias-Zemmour [Igl13] | Souriau [Sou80] | | |
| Hector $[Hec95]$ | Christensen–Wu [CW16] | Torre [Tor98] | | |
| Christensen–Wu [CW16] | Taho [Tah24] | Leslie [Les03] | | |
| Blohmann [Blo24a] | | Dugmore–Ntumba [DN07] | | |
| | | Vincent [Vin08] | | |
| | | Stacey [Sta13] | | |
| $\mathbf{T}_{\mathbf{T}} = \mathbf{T}_{\mathbf{T}} $ | | | | |

TABLE 1. Some approaches to tangent functors

We also mention Magnot [Mag18], who follows Dugmore and Ntumba, and Goldammer and Welker [GW21], who follow Vincent. We use Blohmann's tangent functor, which coincides with Losik's. Its closest, yet distinct, relative is Christensen and Wu's. Like us, Leslie and Magnot assign Lie algebras to some diffeological groups. However, it was unclear whether our desired examples satisfy their hypotheses.

Tangent structures were introduced by Rosický [Ros84], and Cockett and Cruttwell (e.g. [CC14; CC15]) have substantially expanded the theory. Aintablian and Blohmann [AB24] recently established Lie functors in tangent categories, and more generally differentiated groupoid objects in a tangent category to generalized Lie algebroids.

The standard reference for convenient analysis is Kriegl and Michor's book [KM97], though the categorical treatment of Frölicher and Kriegl [FK88] more directly reveals the connection to diffeology, via Frölicher spaces. Hain [Hai79] showed that the category of Banach manifolds embeds into \mathcal{D} flg. Losik [Los92] showed the same for Fréchet manifolds.

Integrations of central extensions of Lie algebras originate with Cartan [Car52], with advances by van Est and Korthagen [EK64] and Neeb [Nee02]. Our integration into diffeological spaces is new, but the principal of using more flexible categories to avoid obstructions is not. Wockel, individually [Woc11] and later with Zhu [WZ16], integrated central extensions of locally exponential locally convex Lie algebras into central extensions of certain Lie 2-groups. This works without caveats, at the expense of higher categorical structures. Beltita and Pelletier [BP24] integrate Banach-Lie algebras into what they call H-manifolds. The structure of an H-manifold is very close to the diffeological quotient of a Banach manifold, but Beltita and Pelletier do not use diffeology. The precise relationship between H-manifold structures and diffeology remains open.

1.3. Structure of the paper. In Section 2, we introduce the tangent category of elastic spaces, with a particular focus on elastic groups. In Section 3, we address the elasticity of orbit spaces of diffeological groupoids. We then apply our general result to subgroups of elastic groups, and achieve Theorem I. In Section 4, we introduce the convenient setting of analysis from a diffeological perspective, and prove Theorem II. Finally, in Section 5, we integrate some central extensions of convenient Lie algebras, and in particular integrate

specific Banach-Lie algebras to diffeological groups via Theorem III. In Appendix A we review tangent categories.

1.4. Acknowledgements. I am grateful to Christian Blohmann for introducing me to tangent structures and elastic spaces. I would also like to thank Lory Aintablian and David Aretz for helpful discussions on tangent categories and convenient analysis, respectively. Finally, I would like to acknowledge the Max Planck Institute for Mathematics in Bonn, which provided a friendly and supportive environment while I pursued this work.

2. A TANGENT STRUCTURE

This section has two parts. In Subsection 2.1, we introduce our tangent functor T: \mathcal{D} flg $\rightarrow \mathcal{D}$ flg, and the tangent category of elastic spaces \mathcal{E} lst. Many of the basic properties of T are proved in [Blo24b]. In Subsection 2.2, we restrict our focus to elastic diffeological groups and their Lie algebras. We emphasize that it is the entire tangent structure on \mathcal{E} lst that lets us define the Lie functor.

We assume basic familiarity with diffeological spaces: the definition, distinguished maps (inductions and subductions),⁴ and the D-topology. For the necessary background, the reader may refer to the first two chapters of Iglesias-Zemmour's book [Igl13]. For a sheaf-based introduction to diffeology, we recommend Baez and Hoffnung's article [BH11].

2.1. Elastic diffeological spaces. Let \mathcal{C} art denote the category of open subsets of Cartesian (Euclidean) spaces, with smooth maps between them. We have the embedding $y: \mathcal{C}$ art $\rightarrow \mathcal{D}$ flg, which assigns to an open subset of Cartesian space its canonical diffeology. Given a functor $\hat{F}: \mathcal{C}$ art $\rightarrow \mathcal{C}$ art, we denote the left Kan extension of $y\hat{F}$ along y by $\mathbb{L}\hat{F} := \operatorname{Lan}_y y\hat{F}$.

The category \mathcal{C} art is small, and \mathcal{D} flg is co-complete. Therefore, for any functor \hat{F} , the left Kan extension $\mathbb{L}\hat{F}$ exists, and moreover it is given by the colimit

$$(\mathbb{L}\hat{F})X = \operatorname{colim}_{y \downarrow X} y \hat{F} \Pi^X.$$

Here we are taking the colimit over the category of plots of X, which is the comma category $y \downarrow X$. Explicitly,

$$y \downarrow X$$
 has
 $\begin{cases} \text{objects: pairs } (U_p, p) \text{ where } yU_p \xrightarrow{p} X \text{ is a plot} \\ \text{morphisms: smooth maps } f : U_p \to U'_{p'} \text{ such that } p' \circ yf = p \end{cases}$

The functor $\Pi^X : y \downarrow X \to C$ art is the projection sending (U_p, p) to U_p .

In order to compute $(\mathbb{L}\hat{F})X$, we must compute colimits in \mathcal{D} flg. Given any functor $D: \mathcal{J} \to \mathcal{D}$ flg from a small category \mathcal{J} , we may take its colimit to be induced by the co-equalizer

(2.1)
$$\coprod_{f \in \operatorname{Mor}(\mathcal{J})} D(\operatorname{dom} f) \xrightarrow[]{\operatorname{id}} \prod_{U \in \operatorname{Ob}(\mathcal{J})} DU \xrightarrow{\operatorname{pr}} \operatorname{colim}_{\mathcal{J}} D.$$

In detail, the two coproducts have their standard diffeology, and $\operatorname{colim}_{\mathcal{J}} D$ is the diffeological quotient of $\coprod DU$ by the relation generated by $x \sim Df(x)$. Since Df is

⁴These are the strong monomorphisms and strong epimorphisms.

not necessarily invertible, $x \sim y$ if and only if there is a *zig-zag* between them: a sequence $x = x_0, x_1, \ldots, x_{k-1}, x_k = y$, and morphisms f_1, \ldots, f_k , where for each *i*, either $Df_i(x_i) = x_{i+1}$ or $Df_i(x_{i+1}) = x_i$ for $1 \leq i \leq k$. We illustrate this in a diagram later.

We can also take the left Kan extension of a natural transformation. Given a natural transformation $\hat{\alpha} : \hat{F} \to \hat{G}$, we denote the left Kan extension of $y\hat{\alpha} : y\hat{F} \to y\hat{G}$ along y by $\mathbb{L}\hat{\alpha} := \operatorname{Lan}_y y\hat{\alpha}$. This is a natural transformation from $\mathbb{L}\hat{F}$ to $\mathbb{L}\hat{G}$. Under our choice of colimits 2.1, the map $(\mathbb{L}\hat{\alpha})_X$ is uniquely determined by commutativity of:

$$(2.2) \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}(\operatorname{dom} f) \xrightarrow{\operatorname{id}} \coprod_{(U_p, p) \in \operatorname{Ob}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{F})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \downarrow X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \coprod X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \coprod X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \coprod X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \coprod X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G})(X)} \qquad \underbrace{\coprod_{f \in \operatorname{Mor}(y \coprod X)} y \hat{F}U_p}_{\operatorname{pr}}_{(\mathbb{L}\hat{G}$$

Notation 2.3. We write $p_*v \in (\mathbb{L}\hat{F})(X)$ for the equivalence class of $v \in y\hat{F}U_p$, which is also $p_*v = \mathbb{L}\hat{F}(p)(v)$. We can write explicitly

(2.4)
$$(\mathbb{L}\hat{\alpha})_X(p_*v) = \operatorname{pr}\left((y\hat{\alpha}\Pi^X)_{U_p}(v)\right) = \mathbb{L}\hat{F}(p)((y\hat{\alpha}\Pi^X)_{U_p}(v)).$$

We now turn specifically to the tangent functor on \mathcal{D} flg, and its role in a tangent category. We summarize the theory of tangent categories in Appendix A. The tangent structure on \mathcal{C} art is given by the following data: the tangent functor $\hat{T} : \mathcal{C}$ art $\rightarrow \mathcal{C}$ art,

 \triangleleft

$$T(U \subseteq \mathbb{R}^n) := U \times \mathbb{R}^n, \quad Tf(u,v) := (f(u), D_u f(v)),$$

where $D_u f$ is the Jacobian of f at u, together with the natural transformations

| the projection | $\hat{\pi}:\hat{T}\to 1$ | $(u,v)\mapsto u$ |
|-------------------------|--|---|
| the zero-section | $\hat{0}: 1 \to \hat{T}$ | $u \mapsto (u, 0)$ |
| the addition | $\hat{+}:\hat{T}\times_{\hat{\pi}}\hat{T}\to\hat{T}$ | $(u, v, u, w) \mapsto (u, v + w)$ |
| the symmetric structure | $\hat{\tau}:\hat{T}^2\to\hat{T}^2$ | $(u, v_0, v_1, v_{01}) \mapsto (u, v_1, v_0, v_{01})$ |
| the vertical lift | $\hat{\lambda}:\hat{T}\rightarrow \hat{T}^2$ | $(u,v)\mapsto (u,0,0,v).$ |

Formally, we say the tuple $(\hat{T}, \hat{\pi}, \hat{0}, \hat{+}, \hat{\tau}, \hat{\lambda})$ is a tangent structure on \mathcal{C} art. To get a tangent structure on \mathcal{D} flg, we would like to apply \mathbb{L} to this tuple.

Definition 2.5. The tangent functor $T : \mathcal{D}flg \to \mathcal{D}flg$ is defined by $T := \mathbb{L}\hat{T}$.

We use this functor extensively, so we establish some notation.

Notation 2.6. Examining diagram (2.1), we realize TX as a quotient of

$$\prod_{(U_p,p)\in \mathrm{Ob}(y\downarrow X)} y\hat{T}U_p = \prod_p yU_p \times y\mathbb{R}^{n_p}$$

A generic element of this coproduct has the form (u, v, p), where $(u, v) \in yTU_p$ and $yU_p \xrightarrow{p} X$ is a plot. We will often suppress the base-point and write (v, p). A zig-zag witnessing $(v_b, p_b) \sim (v_e, p_e)$ (using b for "beginning" and e for "end") has the generic form.

$$(yU_b, v_b) \xrightarrow{f_b} (yU_1, v_1) \xleftarrow{f_1} (yU_2, v_2) \xrightarrow{f_2} \cdots \xleftarrow{f_e} (yU_e, v_e)$$

where all sub-triangles commute and

$$\hat{T}f_i(v_i) = v_{i+1} \text{ or } \hat{T}f_i(v_{i+1}) = v_i \text{ for } 0 \le i \le k.^5$$

Any arrow may be reversed. In our notation, the diagram is not one of pointed spaces; for instance $f_b(v_b)$ makes no sense. However, the base points u_i are all preserved, and map to the same point in X. We will use p_*v or $p_*(u, v)$ to denote the equivalence class of (v, p), and we will say " p_*v is the tangent vector represented by v in the plot p with base-point u."

For a smooth map $f: X \to Y$, its tangent $Tf: TX \to TY$ is simply given by

$$Tf(p_*v) = (f \circ p)_*(v).$$

If we identify $TyU = y\hat{T}U$, which we henceforth do, then for a plot $yU \xrightarrow{p} X$, its tangent map is given by $Tp(u, v) = p_*v$.

Lemma 2.7. The tangent functor has the following properties:

- ([Blo24b, Proposition 2.2.10]) The tangent functor preserves products, meaning that $T(X \times Y)$ is naturally isomorphic to $TX \times TY$. This is because \hat{T} preserves products.
- ([Blo24b, Proposition 2.2.16]) If $p: X \to Y$ is a subduction, so is $Tp: TX \to TY$. This can be verified directly, using only the fact that \hat{T} is a cosheaf.
- ([Blo24a, Proposition 3.35]) If $U \subseteq X$ is D-open, with inclusion $i: U \hookrightarrow X$, then $Ti: TU \to TX$ is an induction with D-open image. In other words, we identify TU with an open subset of TX.

Another important property, which has a surprisingly subtle proof, is that TX is a quotient of the space of smooth curves into X.

Proposition 2.8 ([Blo24b, Proposition 2.2.20]). The map $\partial : C^{\infty}(y\mathbb{R}, X) \to TX$ defined by sending a smooth path $\gamma : y\mathbb{R} \to X$ to the tangent vector $\gamma_*(0, 1)$ is a subduction.

Let us move on to the natural transformations in the tangent structure.

Definition 2.9. We define

$$\pi := \mathbb{L}\hat{\pi} : T \to 1, \text{ and } 0 := \mathbb{L}0 : 1 \to T.$$

 \diamond

⁵For the purposes of this equation, b = 0 and e = k.

Examining Diagram (2.2) and Equation (2.4), we see that $\pi(p_*(u, v)) = p(u)$. Similarly, $0(x) = p_*(u, 0)$, for any plot $yU \xrightarrow{p} X$ with p(u) = x. Using π , we have the k-fold fiber products $T_k := T \times_{\pi} \cdots \times_{\pi} T$. For a tangent structure, T_2 must be the source of addition, and T^2 must be the source and target of the symmetric structure. This is where we encounter a problem, because in general

 $\mathbb{L}\hat{T}_2$ is not isomorphic to $(\mathbb{L}\hat{T})_2 = T_2$, and $\mathbb{L}(\hat{T}^2)$ is not isomorphic to $(\mathbb{L}\hat{T})^2 = T^2$.

We encounter the former difficulty in Example 3.9. Furthermore, in general

 $T(T_k)$ is not isomorphic to $T^2 \times_T \cdots \times_T T^2$.

Nevertheless, we have the natural transformations, which we write by adapting Notation 2.6 (replacing \hat{T} with \hat{T}_k or \hat{T}^k as needed):

$$\theta_k : (\mathbb{L}T_k) \to T_k \quad \theta_{2,X}(p_*(u,v,w)) = (p_*v, p_*w)$$

$$\theta^k : (\mathbb{L}\hat{T}^k) \to T^k \quad \theta_X^2(p_*(u,v_0,v_1,v_{01})) = T^2p(u,v_0,v_1,v_{01})$$

$$\nu_k : TT_k \to T^2 \times_T \cdots \times_T T^2 \quad \nu_{2,X}((p_1,p_2)_*v) = ((p_1)_*v, (p_2)_*v).$$

We illustrate only k = 2 to save space. To define a tangent structure, Blohmann [Blo24a] has isolated a subcategory of \mathcal{D} flg on which these natural transformations behave properly.

Definition 2.10 ([Blo24a, Definition 4.1]). A diffeological space X is *elastic* if it satisfies the following five axioms:

- (E1) The maps $\theta_{k,X}$ are isomorphisms for all $k \ge 1$.
- (E2) There is map $\tau_X: T^2X \to T^2X$ (the canonical flip) such that

$$\begin{array}{ccc} (\mathbb{L}\hat{T}^2)X \xrightarrow{(\mathbb{L}\hat{\tau})_X} (\mathbb{L}\hat{T}^2)X \\ & & \downarrow^{\theta^2_X} & \downarrow^{\theta^2_X} \\ T^2X \xrightarrow{\tau} T^2X \end{array}$$

commutes.

(E3) The vertical lift

$$\lambda_X : TX \xrightarrow{(\mathbb{L}\hat{\lambda})_X} (\mathbb{L}\hat{T}^2)X \xrightarrow{\theta_X^2} T^2X$$

is an induction.

- (E4) The maps $\nu_{k,X}$ are injective for all $k \ge 1$.
- (E5) For every finite set of positive integers k_1, \ldots, k_n , the spaces $T_{k_1} \cdots T_{k_n} X$ satisfy (E1) through (E4).

 \diamond

A diffeological space X is either elastic, or it is not. The collection of elastic diffeological spaces, with smooth maps between them, is a full sub-category of \mathcal{D} flg, which we denote by \mathcal{E} lst. The raison d'être for these axioms is the following theorem.

Theorem 2.11 ([Blo24a, Theorem 4.2]). On \mathcal{E} lst, the tangent functor T, together with the natural transformations

$$\pi, \quad 0, \quad + := \mathbb{L} \hat{+} \circ \theta_2^{-1}, \quad \tau \text{ from (E2)}, \quad \lambda := \theta^2 \circ \mathbb{L} \hat{\lambda}.$$

forms a tangent structure. Moreover, \mathcal{E} lst is a Cartesian tangent category with scalar \mathbb{R} multiplication (Definition A.9).

Remark 2.12. Example 3.9 shows that (E1) may fail. Similarly, (E4) can fail. We also suspect that (E3) can fail, with a potential example coming from a non-convenient locally convex topological vector space (cf. Section 4). However, it may be that (E2) holds for any diffeological space.

To be a Cartesian tangent category means that the tangent functor preserves products: $T(X \times Y) \cong TX \times TY$. In this setting, partial derivatives make sense. Given a map $f: X \times Y \to Z$ of elastic spaces, set

$$T_{(1)}f := Tf \circ 0 \times 1 : TX \times Y \to TZ$$
$$T_{(2)}f := Tf \circ 1 \times 0 : X \times TY \to TZ.$$

Then both $T_{(1)}T_{(2)}f$ and $T_{(2)}T_{(1)}f$ are maps $TX \times TY \to T^2Z$, and they are related by

$$T_{(2)}T_{(1)}f = \tau_Z T_{(1)}T_{(2)}f.$$

This holds true in a general Cartesian tangent category, not just on \mathcal{E} lst. See Appendix A.2 for details.

2.2. Elastic diffeological groups. It might be laborious to check all of (E1) through (E5). Thankfully, for groups, the conditions simplify.

Theorem 2.13 ([Blo24a, Theorem 5.2]). Diffeological groups satisfy axioms (E1), (E2), and (E4). If they also satisfy (E3), then they are elastic.

We will elucidate the tangent structure on elastic groups. Let G be a diffeological group. When there is no ambiguity, we drop the subscript G from natural transformations, for instance we write $\pi = \pi_G$. Denote by $\mathfrak{g} := T_e G$ the fiber of its tangent space over the unit, equipped with its subspace diffeology. Denote the group multiplication by m and inversion by i. Then we have the left trivialization

$$\varphi_G: G \times \mathfrak{g} \hookrightarrow G \times TG \xrightarrow{T_{(2)}m} TG.$$

Its inverse, denoted ψ_G , is determined by

Remark 2.14. If we write left multiplication by $q \in G$ as

$$L_q: G \times G, \ L_q(h) := gh,$$

then the trivializations are $\varphi_G(g,\xi) = TL_g(\xi)$ and $\psi_G(\xi_g) = (g, TL_{g^{-1}}\xi_g)$. However, writing it this way obscures their smoothness.

In the left trivialization, the projection and zero-section are

$$\pi(g,\xi) = g, \quad 0(g) = (g,0).$$

We can also write down the canonical flip. Because T preserves products, \mathfrak{g} is also a diffeological group, so we have its left-trivialization $\psi_{\mathfrak{g}}: T\mathfrak{g} \to \mathfrak{g} \times T_0\mathfrak{g}$. We then have the isomorphism β :

$$T^{2}G \xrightarrow{T\psi_{G}} TG \times T\mathfrak{g} \xrightarrow{\psi_{G} \times \psi_{\mathfrak{g}}} G \times \mathfrak{g} \times \mathfrak{g} \times T_{0}\mathfrak{g}$$

We define τ by

$$\beta \circ \tau \circ \beta^{-1}(g, \xi_0, \xi_1, \xi_{01}) = (g, \xi_1, \xi_0, \xi_{01}).$$

Finally, the vertical lift can be computed using the compatibility of the natural transformations that will constitute the tangent structure, as described in Appendix A.1.

$$\pi_{TG} \circ \lambda = 0 \circ \pi \implies \beta \circ \lambda \circ \varphi_G(g, \xi) = (g, ?, 0, ?)$$
$$\lambda \circ \tau = \tau \implies \beta \circ \lambda \circ \varphi_G(g, \xi) = (g, 0, 0, \lambda^{\perp}(g, \xi)).$$

Furthermore,

(2.15)

Lemma 2.16 ([Blo25, Proposition 5.7]). The map λ^{\perp} depends only on \mathfrak{g} , and is given by

$$\lambda^{\perp}: \mathfrak{g} \to T_0 \mathfrak{g}, \quad \xi \mapsto \partial[t\xi],$$

where $[t\xi]$ denotes the map $\mathbb{R} \to \mathfrak{g}$ given by $t \mapsto t\xi$.

The vertical lift λ^{\perp} controls elasticity of diffeological groups.

Proposition 2.17 ([Blo24a, Theorem 5.2 continued]). A diffeological group G is elastic (i.e. (E3) is satisfied) if and only if the map λ^{\perp} is an isomorphism.

The condition that λ^{\perp} be an isomorphism, and not merely an induction, may be surprising, so we summarize the proof.

Proof. If λ^{\perp} is an isomorphism, then it is clear from expression (2.15) that λ is an induction. Thus (E3) is satisfied, and G is elastic. Conversely, if G is elastic, then λ is an induction, and by expression (2.15) so is λ^{\perp} . A surjective induction is an isomorphism, so we show λ^{\perp} is surjective. We appeal to the axioms of a tangent category, stated in Appendix A. Specifically, the last axiom implies that the diagram

$$\begin{array}{ccc} T_2G & \xrightarrow{\lambda_2} & T^2G \\ \pi \circ \mathrm{pr}_1 & & & \downarrow_{T\pi} \\ G & \xrightarrow{0} & TG \end{array}$$

is a pullback, where λ_2 is, in the trivializations of TG and T^2G , the map

$$\lambda_2(g,\xi_0,g,\xi_1) := (g,0,\xi_0,\lambda^{\perp}(\xi_1)).$$

Then, since the maps

$$\begin{split} T_0 \mathfrak{g} &\to G, \quad \Xi \mapsto e \\ T_0 \mathfrak{g} &\to T^2 G \quad \Xi \mapsto (e,0,0,\Xi) \end{split}$$

coincide when composed with 0 and $T\pi$, respectively, there is a map $s: T_0\mathfrak{g} \to T_2G$ such that $\lambda_2 \circ s(\Xi) = (e, 0, 0, \Xi)$. In other words, λ^{\perp} has a section, and thus is surjective. \Box

We abstract the condition in Proposition 2.17 to general diffeological vector spaces.

Definition 2.18. Let V be a diffeological vector space. Then V is *tangent-stable* if the map

$$\lambda_V: V \to T_0 V, \quad v \mapsto \partial[tv]$$

is an isomorphism.

With this terminology, the tangent fiber $\mathfrak{g} = T_e G$ of an elastic group is tangent-stable.

Proposition 2.19 ([Blo24a, Proposition 5.5]). Tangent-stable vector spaces are elastic. Moreover, it is equivalent to check that the map

$$\nu: V \times V \to TV, \quad (u, v) \mapsto \partial[u + tv]$$

is an isomorphism.

The axioms of elasticity are chosen so that \mathcal{E} lst is a tangent category. We require a tangent category to invoke the main result of Aintablian and Blohmann [AB24], which for groups states: a group object in a tangent category (Cartesian with scalar \mathbb{R} multiplication) has a Lie algebra. We defer the technical details to Appendix A.3, but the construction is familiar. When G is an elastic group, we can identify \mathfrak{g} with $\mathfrak{X}(G)^L$, the space of left-invariant vector fields on G. The tangent structure equips $\mathfrak{X}(G)$, and also $\mathfrak{X}(G)^L$, with a Lie bracket, which we use to define the Lie algebra structure on \mathfrak{g} . Moreover, this assignment is functorial; given a smooth homomorphism $f: G \to H$, its tangent map $Tf: \mathfrak{g} \to \mathfrak{h}$ is a homomorphism of Lie algebras.

Proposition 2.20. We have a Lie functor

Lie: $\{elastic groups\} \rightarrow \{tangent-stable Lie algebras\},\$

given by $\operatorname{Lie}(G) := \mathfrak{g} \cong \mathfrak{X}(G)^L$.

3. Diffeological groupoids

Having introduced the objects we want to differentiate, we now justify our key examples: certain quotients of elastic groups. This is a two-step process. In Subsection 3.1, we compute the tangent bundle of the orbit space of some diffeological groupoids. In Subsection 3.2, we apply this computation to the action groupoid corresponding to a subgroup acting on its ambient group, in order to conclude elasticity in favourable cases.

 \diamond

3.1. Tangent bundles of orbit spaces.

Definition 3.1. A *diffeological groupoid* is a groupoid object in \mathcal{D} flg. Thus, a diffeological groupoid consists of a groupoid $\mathcal{G} \rightrightarrows X$, and diffeological structures on \mathcal{G} and X for which all structure maps are smooth.

We will denote the source map by α , the target map by β , the multiplication by m, the unit by u, and the inversion by i. As a consequence of the algebraic structure of $\mathcal{G} \rightrightarrows X$, the source and target are subductions, the inversion is a diffeomorphism, and the unit is an induction. We denote the orbit map by

$$\operatorname{pr} = \operatorname{pr}^0 : X \to X/\mathcal{G}.$$

This is a subduction, and specifically co-equalizes the pair (α, β) .

Remark 3.2. Lie groupoids, by contrast, are more than simply groupoid objects in the category of manifolds. For a Lie groupoid, we additionally require that: the base space is Hausdorff and second-countable; the source, hence target, maps are submersions; and the source, hence target, fibers are Hausdorff. Submersions between manifolds are subductions, but the converse does not hold. \triangleleft

Let \mathcal{G}_{\bullet} denote the nerve of a groupoid \mathcal{G} . Thus, at level k, we have

$$\mathcal{G}_k := \mathcal{G}_{\alpha} \times_{\beta} \cdots_{\alpha} \times_{\beta} \mathcal{G}_{\alpha}$$

We also denote $\mathcal{G}_1 := \mathcal{G}$ and $\mathcal{G}_0 := X$.

Definition 3.3 ([AB24, Definition 3.8]). A diffeological groupoid \mathcal{G} is differentiable if the natural morphism

$$T^{n}(\mathcal{G}_{k}) \to T^{n}\mathcal{G}_{T^{n}\alpha} \times_{T^{n}\beta} \cdots {}_{T^{n}\alpha} \times_{T^{n}\beta} T^{n}\mathcal{G}$$

is an isomorphism for each $k \ge 0$ and $n \ge 0$.

When \mathcal{G} is differentiable, we identify $T\mathcal{G}$ as a diffeological groupoid $T\mathcal{G} \rightrightarrows TX$, with source $T\alpha$, target $T\beta$, and multiplication Tm.

Remark 3.4. Strictly speaking, Aintablian and Blohmann define differentiable groupoids only in tangent categories. In that context, the differentiable groupoids are precisely those which differentiate to, in their sense, Lie algebroids. \triangleleft

Example 3.5. Aintablian and Blohmann showed [AB24, Proposition 8.1] that a groupoid object in the category of manifolds is differentiable if and only if it is a Lie groupoid. This result can be used as a justification for the definition of a Lie groupoid being stronger than that of a diffeological groupoid. \triangleright

We are primarily concerned with action groupoids.

Lemma 3.6. Let $X \rtimes H \rightrightarrows X$ be an action groupoid for a diffeological group H acting on a diffeological space X. Then $X \rtimes H$ is differentiable.

 \diamond

Proof. Recall that $X \rtimes H$ has for its source and target

the action map $\mu: X \times H \to X$, $(x, h) \mapsto x \cdot h$

the projection map $pr_1: X \times H \to X$, $(x, h) \mapsto x$.

Thus, for instance,

$$(X \rtimes H)_2 \cong X \times H \times H, \quad (x, h, x' = x \cdot h, h') \mapsto (x, h, h').$$

Therefore differentiability of $X \rtimes H$ follows from the fact that T preserves finite products.

When \mathcal{G} is a differentiable groupoid, the orbit space $TX/T\mathcal{G}$ of $T\mathcal{G}$ is a co-equalizer of the pair $(T\alpha, T\beta)$. On the other hand, the map $T \text{ pr} : TX \to T(X/\mathcal{G})$ also satisfies $T \text{ pr} \circ T\alpha = T \text{ pr} \circ T\beta$. Thus the universal property of co-equalizers gives us a unique smooth map $\eta_{\mathcal{G}}$ fitting in



The map $\eta_{\mathcal{G}}$ descends to the identity on X/\mathcal{G} . We do not anticipate that this map is always an isomorphism. However, it will be an isomorphism in the cases relevant to us.

Proposition 3.7. Suppose that $\mathcal{G} \rightrightarrows X$ is a differentiable groupoid satisfying the following condition:

(3.8) if $(q, q'): U \to X \times_{\mathrm{pr}} X$ is a plot, then $(Tq, Tq'): TU \to TX \times_{\mathrm{pr}^1} TX$ is a plot. Then the natural map $\eta_{\mathcal{G}}: T(X/\mathcal{G}) \to TX/T\mathcal{G}$ is an isomorphism.

Proof. Since the quotient pr is a subduction, so is T pr. The quotient map pr^1 is also a subduction. Therefore $\eta_{\mathcal{G}}$ is a subduction, and we only need to show it is injective. To do this, we show that if $T \operatorname{pr}(w_b) = T \operatorname{pr}(w_e)$ (b for "beginning" and e for "end"), then $\operatorname{pr}^1(w_b) = \operatorname{pr}^1(w_e)$.

Represent w_{\bullet} in a plot $(v_{\bullet}, q_{\bullet})$ for $\bullet \in \{b, e\}$. Then $T\pi(w_b) = T\pi(w_e)$ means there is a zig-zag between $(v_b, \pi q_b)$ and $(v_e, \pi q_e)$. An arbitrary link in such a zig-zag has the form

$$(U, u, v) \xrightarrow{f} (U', u', v')$$

$$p \xrightarrow{p'} Tf(v) = v'.$$

$$X/\mathcal{G},$$

Since pr : $X \to X/G$ is a subduction, we may lift p, p' along π at u, u' to maps q, q' into X. This may involve shrinking U, U' about u, u'. Then we have a plot

 $(q,q'f): U \to X \times_{\mathrm{pr}} X$, implying we have a plot $(Tq,T(q'f)): TU \to TX \times_{\mathrm{pr}^1} TX$. In particular, $\mathrm{pr}^1(q_*v) = \mathrm{pr}^1(q'_*v')$.

At the leftmost arm of a zig-zag witnessing $(v_b, \pi q_b) \sim (v_e, \pi q_e)$, choose q_b as the lift of πq_b . At the rightmost arm, choose q_e as the lift of πq_e . By choosing lifts in

each intermediate link of the zig-zag as we did above, we conclude that $\operatorname{pr}^1((q_b)_*v_b)$ is preserved across the zig-zag, and must therefore coincide with $\operatorname{pr}^1((q_b)_*v_b)$. In other words, $\operatorname{pr}^1(w_b) = \operatorname{pr}^1(w_e)$, and so $\eta_{\mathcal{G}}$ is injective.

Example 3.9. Let \mathbb{Z}_2 act on \mathbb{R} by the involution $x \mapsto -x$. Suppose that $q, q' : U \to \mathbb{R}$ are two plots, such that for each $u \in U$, we have $q(u) = \pm q'(u)$. Set

$$\Delta_{+} := \{ u \in U \mid q(u) = q'(u) \} \text{ and } \Delta_{-} := \{ u \in U \mid q(u) = -q'(u) \}.$$

Then $U = \Delta_+ \cup \Delta_-$ is a finite union of closed sets, and as a consequence of the Baire category theorem, the union $\Delta_+^\circ \cup \Delta_-^\circ$ is dense in U. On each Δ_{\pm}° , we have $Tq = \pm Tp$. Thus, by continuity, for each $(u, v) \in TU|_{\overline{\Delta_{\pm}^\circ}}$ we have $Tq(u, v) = \pm Tp(u, v)$. Since $U = \overline{\Delta_+^\circ} \cup \overline{\Delta_-^\circ}$ (a property of finite unions of sets), we have showed that condition (3.8) holds. Therefore $T(\mathbb{R}/\mathbb{Z}_2) \cong T\mathbb{R}/\mathbb{Z}_2$, where \mathbb{Z}_2 acts on $T\mathbb{R}$ by the involution $(x, v) \mapsto (-x, -v)$.

Using a similar argument, we find that $\mathbb{L}\hat{T}_2(\mathbb{R}/\mathbb{Z}_2) \cong (T_2\mathbb{R})/\mathbb{Z}_2$. The spaces $(T\mathbb{R})/\mathbb{Z}_2$ and $(T_2\mathbb{R})/\mathbb{Z}_2$ are not isomorphic, so \mathbb{R}/\mathbb{Z}_2 is not elastic.

More generally, we can show that if $\mathcal{G} \rightrightarrows M$ is an orbifold groupoid, meaning that it is locally isomorphic to action groupoids of the form $\Gamma \ltimes \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ for finite groups $\Gamma \leq \operatorname{GL}(n)$,⁶ then $T(M/\mathcal{G}) \cong TM/T\mathcal{G}$. Upon embedding the category of orbifolds into \mathcal{D} flg (as done in [IKZ10]), we observe that our tangent functor coincides with the usual orbifold tangent functor. If any of the isotropy groups \mathcal{G}_x are non-trivial, then M/\mathcal{G} is not elastic.

 \triangleright

We are primarily interested in the case that H is a subgroup of a diffeological group G, equipped with its subspace diffeology. Let $\iota : H \hookrightarrow G$ denote the inclusion. Let $\mu : G \times G \to G$ denote the multiplication in G. Right multiplication by H gives the action groupoid $G \rtimes H \rightrightarrows G$. This is differentiable, so we have its tangent groupoid $T(G \rtimes H) \rightrightarrows G$. The source map is $T(\mu \circ (1 \times \iota)) = T\mu \circ (1 \times T\iota)$, and the target map is $T \operatorname{pr}_1$.

The tangent bundle TG of G is also a diffeological group, with multiplication given by $T\mu$. Therefore, as groupoids,

$$T(G \rtimes H) \cong TG \rtimes T\iota(TH),$$

where $T\iota(TH)$ acts on TG from the right.

Corollary 3.10. Let $\iota : H \hookrightarrow G$ be a subgroup of a diffeological group. For the associated action groupoid $G \rtimes H \rightrightarrows G$, the natural map

$$\eta_{G \rtimes H} : T(G/H) \to TG/T(G \rtimes H) = TG/T\iota(TH)$$

is an isomorphism.

⁶The reader may be more comfortable defining an orbifold groupoid as an étale and proper Lie groupoid. Our definition is strictly more general, and allows for non-Hausdorff orbifolds.

Proof. Write the action of H on G by $gh := \mu(g, h)$. Being an action groupoid, $G \rtimes H$ is differentiable. We verify it satisfies condition 3.8 in Proposition 3.7. Let

 $(1,d): G \times_{\mathrm{pr}} G \to G \times H$ denote the inverse of $(1,\mu): G \times H \to G \times_{\mathrm{pr}} G$.

Here d stands for "division." Suppose that $(q, q') : U \to G \times_{\mathrm{pr}} G$ is a plot. Recall that pr^1 is the orbit map for the groupoid $T(G \rtimes H)$, which has source $T\mu$ and target $T \mathrm{pr}_1$. Observe that

$$T\mu \circ (Tq, Td \circ T(q, q')) = T(\mu \circ (q, d \circ (q, q'))) = Tq'$$

$$T\operatorname{pr}_1 \circ (Tq, Td \circ T(q, q')) = T(\operatorname{pr}^1 \circ (q, d \circ (q, q'))) = Tq.$$

Thus $(Tq, Tq'): TU \to TG \times_{\mathrm{pr}^1} TG$ is a plot, as required.

Remark 3.11. The conclusion of Corollary 3.10 generalizes to the case that $\mathcal{G} = X \rtimes H \Rightarrow X$, where we assume that the anchor map $(1, \mu)$ is an isomorphism. In this case, the orbit map pr : $X \to X/H$ is a principal fibration with structure group H in the sense of [Igl13, Article 7.3], and the conclusion writes: the natural map $\eta_{X \rtimes H} : TX/TH \to T(X/H)$ is an isomorphism. The proof is identical.

3.2. Elasticity of quotient groups. Let H be a normal subgroup of a diffeological group G with inclusion $\iota : H \hookrightarrow G$. Then G/H is a diffeological group, and $T\iota(TH) \hookrightarrow TG$ is a normal subgroup, and by Corollary 3.10 we have an isomorphism of groups

$$T(G/H) \cong TG/T\iota(TH).$$

Whenever $T^k \iota : T^k H \to T^k G$ is an induction, we identify $T^k H$ with $T^k \iota(T^k H)$, and write $T^k G/T^k H := T^k G/T^k \iota(T^k H)$.

Theorem I. Let G be an elastic group, with a normal subgroup H carrying its induced diffeology. Denote the inclusion by $\iota: H \hookrightarrow G$. If the iterated tangents $T^k \iota: T^k H \to T^k G$ are inductions for all $k \ge 0$, then H and G/H are elastic, and $\eta_{G \rtimes H} : \text{Lie}(G/H) \to$ Lie(G)/Lie(H) is an isomorphism of Lie algebras.

In fact, we may only assume that $T\iota$ and $T^2\iota$ are inductions.

Proof. We first show that H is elastic. Recall that G being elastic is, by Theorem 2.13, equivalent to the vertical lift λ_G being an induction. Naturality of the vertical lift gives the commutative diagram

$$TG \xrightarrow{\lambda_G} T^2G$$

$$T\iota \int \qquad \int T^2\iota$$

$$TH \xrightarrow{\lambda_H} T^2H.$$

Then λ_H being an induction follows from the universal property of inductions. Therefore H is elastic.

Now we show that G/H is elastic. We identify \mathfrak{h} as a subspace of \mathfrak{g} , and $T_0\mathfrak{h}$ as a subspace of $T_0\mathfrak{g}$, using the inclusions $T\iota$ and $T^2\iota$. Then we have the morphism of

short-exact sequences

$$\begin{split} \mathfrak{h} & \stackrel{T\iota}{\longrightarrow} \mathfrak{g} \stackrel{\mathrm{pr}^{1}}{\longrightarrow} \mathfrak{g}/\mathfrak{h} \\ \downarrow^{\lambda_{H}^{\perp}} & \downarrow^{\lambda_{G}^{\perp}} & \downarrow^{:=\lambda_{0}} \\ T_{0}\mathfrak{h} \stackrel{T^{2}\iota}{\longrightarrow} T_{0}\mathfrak{g} \stackrel{\mathrm{pr}^{2}}{\longrightarrow} T_{0}\mathfrak{g}/T_{0}\mathfrak{h}. \end{split}$$

Since λ_H^{\perp} and λ_G^{\perp} are isomorphisms by Proposition 2.17, so is λ_0 . We now apply Proposition 3.7 twice, and use naturality of λ , to obtain the commutative diagram

$$TG \xrightarrow{\lambda_G} T^2G \xrightarrow{pr^2} T^2G$$

$$\downarrow^{T \text{ pr}} \xrightarrow{T^2 \text{ pr}} T^2 \text{ pr}$$

$$TG/TH \xrightarrow{\cong} T(G/H) \xrightarrow{\lambda_{G/H}} T^2(G/H) \xrightarrow{\cong} T(TG/TH) \xrightarrow{\cong} T^2G/T^2H.$$

Then, identifying $\mathfrak{g}/\mathfrak{h}$ as a subspace of T(G/H), and identifying $T_0\mathfrak{g}/T_0\mathfrak{h}$ as a subspace of $T^2(G/H)$, we see that λ_0 identifies with $\lambda_{G/H}^{\perp}$. Since λ_0 is an isomorphism, so too is $\lambda_{G/H}^{\perp}$, as required. Therefore G/H is elastic.

The next step is to show that \mathfrak{h} is an ideal in \mathfrak{g} . By Lemma A.16, the Lie bracket on \mathfrak{g} is determined by $T_{(1)}T_{(2)}c$, where $c: G \times G \to G$ is conjugation $(g,h) \mapsto ghg^{-1}$. By definition of a normal subgroup, whenever $h \in H$, we have $c(g,h) \in H$. Therefore $T_{(1)}T_{(2)}c$ restricts to a map

$$T_{(1)}T_{(2)}c:TG\times TH\to T^2H$$

This means that whenever $\xi \in \mathfrak{g}$ and $\eta \in \mathfrak{h}$, we have $\delta(\xi, \eta) \in T^2 H$. By definition of the Lie bracket in terms of δ , it follows that $[\xi, \eta] \in \mathfrak{h}$, and so \mathfrak{h} is an ideal in \mathfrak{g} .

Finally, we show that $\eta_{G \rtimes H} : \mathfrak{g}/\mathfrak{h} \to \operatorname{Lie}(G/H)$ is an isomorphism of Lie algebras. Proposition 3.7 shows that $\eta_{G \rtimes H}$ is an isomorphism of vector spaces (and we already used this fact). It preserves the brackets because it fits into the diagram



and the downward arrows are epimorphisms of Lie algebras.

Our first corollary is immediate.

Corollary 3.12. If H is a diffeologically discrete normal subgroup of an elastic group G, then G/H is elastic and $\mathfrak{g} \cong \operatorname{Lie}(G/H)$.

The next corollary is of independent interest, but we do not use it in this paper.

Corollary 3.13. If H is a normal subgroup of a finite-dimensional Lie group G, then G/H is elastic and $\mathfrak{g}/\mathfrak{h} \cong \operatorname{Lie}(G/H)$.

Proof. By Yamabe's theorem [Yam50], the subspace diffeology on H is a manifold diffeology (i.e. H is a Lie group), the inclusion $\iota : H \to G$ is an induction, and $T\iota : TH \to TG$ is injective. In the trivializations, $T\iota$ has the form

$$H \times \mathfrak{h} \xrightarrow{(\iota, T_0 \iota)} G \times \mathfrak{g}.$$

The map ι is an induction, and $T_0\iota : \mathfrak{h} \to \mathfrak{g}$ is a linear injection of finite-dimensional vector spaces equipped with their canonical diffeology, hence is also an induction. Thus $T\iota : TH \to TG$ is an induction. Inductively repeating this argument, we find that we can apply Theorem I to $\iota : H \hookrightarrow G$. The claim follows immediately. \Box

4. The convenient setting and diffeology

Up to this point, our considerations and results have been fairly abstract, the only familiar example being quotients of finite-dimensional Lie groups. We now show that our setting includes the convenient analysis of Kriegl and Michor [KM97]: the category of convenient manifolds embeds into \mathcal{E} lst, and their tangent structures coincide. This allow us freely use all the constructions from the convenient calculus, and their examples become ours.

In Subsection 4.1 we re-cast tangent stability as convenience in the sense of Frölicher and Kriegl [FK88]. In Subsection 4.2, we embed Kriegl and Michor's category of convenient manifolds into \mathcal{E} lst. Doing so, we *a fortiori* embed the categories of Banach and Fréchet manifolds into \mathcal{E} lst. Crucially, we also show that their tangent structures coincide.

4.1. Convenient diffeological vector spaces. The diffeologies we encounter in this section will arise from collections of functions. Let V be a set, and \mathscr{F}_0 be some collection of functions $V \to \mathbb{R}$. The set

$$\Pi \mathscr{F}_0 := \{ p : U \to V \mid U \in \mathcal{C} \text{art}, \ f \circ p \text{ is smooth for all } f \in \mathscr{F}_0 \}$$

is the diffeology on V determined by \mathscr{F}_0 .

Definition 4.1. A diffeology \mathscr{D} on V is *reflexive* if $\mathscr{D} = \prod C^{\infty}(V, \mathbb{R})$.

 \diamond

The terminology "reflexive," and the notation Π , is from [BKW23].

Example 4.2. We have several examples.

- The diffeology $\Pi \mathscr{F}_0$ is always reflexive, and $\mathscr{F}_0 \subseteq C^{\infty}((V, \Pi \mathscr{F}_0), \mathbb{R})$, but equality need not hold. For instance, the inclusion is strict if \mathscr{F}_0 contains only a single constant function.
- If V = M is a finite-dimensional manifold, its canonical diffeology is reflexive.
- Consider the spaghetti diffeology on \mathbb{R}^2 , which is generated by the smooth curves $C^{\infty}(\mathbb{R}, \mathbb{R}^2)$, i.e. it is finest diffeology admitting all smooth curves as plots. Observe that $C^{\infty}((\mathbb{R}^2, \mathscr{D}_{standard}), \mathbb{R})$ and $C^{\infty}((\mathbb{R}^2, \mathscr{D}_{spaghetti}), \mathbb{R})$ coincide. This is by Boman's theorem [Bom67], which grants that a function $\mathbb{R}^2 \to \mathbb{R}$ is smooth if and only if its pullback by all smooth curves is smooth. Thus both collections of functions determine the standard diffeology on \mathbb{R}^2 , and the spaghetti diffeology must not be reflexive.

Reflexive diffeological spaces form a full subcategory of \mathcal{D} flg.

Remark 4.3. Given a reflexive diffeological space (V, \mathcal{D}) , we can form the triple

 $(V, C^{\infty}(\mathbb{R}, V), C^{\infty}(V, \mathbb{R})).$

The data of this triple is an example of a *Frölicher space*. In fact, by [BKW23, Theorem 2.13] this assignment of a Frölicher space to each reflexive diffeological space is an isomorphism between the categories of reflexive diffeological spaces, and Frölicher spaces. Frölicher and Kriegl [FK88] work with the category of Frölicher spaces, which they call $\mathcal{L}ip^{\infty}$ -spaces; the "Lip" stands for "Lipschitz".

The next definition is a reminder, and sets notation.

Definition 4.4. A *diffeological vector space*, abbreviated "dvs," is a vector space V equipped with a diffeology for which addition and scalar multiplication are smooth. A *morphism* of diffeological vector spaces is a smooth linear map.

Diffeological vector spaces, and morphisms between them, form the category \mathcal{D} flgVS. We write $L^{\infty}(V, W) := \mathcal{D}$ flgVS(V, W), and if $W = \mathbb{R}$, then we simply write $L^{\infty}(V)$.

Given a diffeological vector space V, and a smooth curve $c : \mathbb{R} \to V$, we would like to define its velocity $\dot{c} : \mathbb{R} \to V$. For $V = \mathbb{R}^n$, recall that the tangent functor is defined by

$$\hat{T}\mathbb{R}^n = \mathbb{R}^n \times \mathbb{R}^n, \quad \hat{T}c(t_0, t) = (c(t_0), t\dot{c}(t_0)),$$

where $\dot{c}(t_0)$ is the usual velocity vector. Thus $\dot{c}(t_0)$ is the second component of $\hat{T}c(t_0, 1)$. For a general dvs V, we no longer have $TV \cong V \times V$. Instead, we have the natural map

$$\nu: V \times V \to TV, \quad (u, v) \mapsto \partial[u + tv].$$

Recall that under Definition 2.18, we call a dvs tangent-stable if ν is an isomorphism, and by Proposition 2.19, tangent-stable dvs are elastic.

Definition 4.5. Given a smooth curve $c : \mathbb{R} \to V$ into a tangent-stable dvs, its *velocity* curve, denoted $\dot{c} : \mathbb{R} \to V$, is the smooth curve defined by the commutative diagram



Proposition 4.7. Fix a smooth curve $c : \mathbb{R} \to V$ into a tangent-stable dvs. For each $l \in L^{\infty}(V)$, the composition $l \circ c$ is smooth, and $\hat{l \circ c} = l \circ \dot{c}$.

18

 \diamond

Proof. Smoothness of $l \circ c$ is clear. Unravelling definitions, and using the chain rule, we write

$$\widehat{l \circ c}(t_0) = \operatorname{pr}_2 \circ \nu^{-1} \circ T(l \circ c) \circ \nu(t_0, 1)$$

= $\operatorname{pr}_2 \circ (\nu^{-1} \circ Tl \circ \nu) \circ (\nu^{-1} \circ Tc \circ \nu)(t_0, 1).$

But because l is linear, $Tl \circ \nu = \nu \circ (l \times l)$, where $(l \times l)(v, w) := (l(v), l(w))$. For clarity, we confirm this:

$$Tl(\nu(v,w)) = Tl(\partial[v+tw]) = \partial[l(v) + tl(w)] = \nu(l(v), l(w)).$$

Thus

$$\operatorname{pr}_2 \circ (\nu^{-1} \circ Tl \circ \nu) = \operatorname{pr}_2 \circ (l \times l) = l \circ \operatorname{pr}_2.$$

We conclude

$$\dot{l \circ c}(t_0) = l \circ (\operatorname{pr}_2 \circ \nu^{-1} \circ Tc \circ \nu)(t_0, 1) = l \circ \dot{c}(t_0).$$

Proposition 4.7 says that the velocity curve of a smooth curve is also its "weak derivative," which is a notion we can define for a general dvs.

Definition 4.8. A curve $c : \mathbb{R} \to V$ into a dvs is *weakly differentiable* (with respect to $L^{\infty}(V)$), with a *weak derivative* $\dot{c} : \mathbb{R} \to V$, if for all $l \in L^{\infty}(V)$, the composition $l \circ c$ is differentiable, and its derivative satisfies $\hat{l \circ c} = l \circ \dot{c}$.

Weak derivatives need not be unique, but they are for the following spaces.

Definition 4.9. A diffeological vector space V is *linearly separated* if $L^{\infty}(V)$ separates points.

A dvs may, or may not, satisfy the following properties:

- infinitely weakly differentiable curves are smooth;
- weak derivatives, when they exist, are unique;
- smooth curves are infinitely weakly differentiable.

Each property is a consequence of a natural condition imposed on V. We describe these conditions sequentially, and we will call those dvs that satisfy all three "convenient." We have proved that tangent-stable dvs satisfy the third property.

Definition 4.10. A diffeological vector space (V, \mathscr{D}_V) is pre-convenient if $\mathscr{D}_V = \Pi L^{\infty}(V)$.

Pre-convenient dvs are in particular reflexive, but we are not sure if the converse holds. The terminology "pre-convenient" comes from [FK88].

Lemma 4.11. Fix a pre-convenient dvs V. Infinitely weakly differentiable curves into V are smooth.

Proof. This is clear from the definition: since $\mathscr{D}_V = \Pi L^{\infty}(V)$, smoothness of $c : \mathbb{R} \to V$ is equivalent to smoothness of $l \circ c$ for all $l \in L^{\infty}(V)$, which holds for infinitely weakly differentiable curves.

Lemma 4.12. Fix a linearly separated dvs V. Then if a curve is weakly differentiable, its weak derivative is unique.

Proof. This is also clear from the definition: if \dot{c} and d are two weak derivatives of c, then for all $l \in L^{\infty}(V)$,

$$l \circ \dot{c}(t_0) = \widehat{l \circ c}(t_0) = l \circ d(t_0),$$

$$a_0) = d(t_0).$$

thus by linear separation, $\dot{c}(t_0) = d(t_0)$.

Definition 4.13. A pre-convenient and linearly separated dvs V is called *convenient* if every smooth curve is infinitely weakly differentiable. \diamond

This definition of a convenient dvs V is precisely the one used in [FK88], where they view V as a Frölicher space. Specifically, in their notation, it is condition (5) in [FK88, Theorem 2.6.2], with $k = \infty$. We proved in Proposition 4.7 that pre-convenient and linearly separated dvs that are tangent-stable are convenient. Now we prove the converse. We will need the notion of a weak integral.

Definition 4.14. A curve $c : \mathbb{R} \to V$ into a dvs is *weakly integrable*, with a *weak integral* $\int c : \mathbb{R} \to V$, if for all $l \in L^{\infty}(V)$, the composition $l \circ c$ is integrable, and its integral satisfies $\int_0^t l \circ c = l \circ \int c(t)$.

Proposition 4.15 ([FK88, Theorem 2.6.2]). If V is a convenient dvs, then every smooth curve is weakly integrable, and its weak integral is unique.

This is a fundamental result in the theory of convenient calculus. In fact, on a preconvenient and linearly separated dvs, weak differentiability of smooth curves is equivalent to weak integrability.

Lemma 4.16 (Fundamental theorem of calculus). Fix a convenient dvs V. The maps

$$C^{\infty}(\mathbb{R}, V) \to C^{\infty}(\mathbb{R}, V), \quad c \mapsto \dot{c} \text{ and } c \mapsto \int dc$$

are smooth and linear. Furthermore, $t \mapsto c(t) - c(0)$ is the weak integral of \dot{c} .

We will denote $\int_s^t c := \int c(t) - \int c(s)$. Thus $\int c(t) = \int_0^t c$.

Proof. Take any plot $p: U \to C^{\infty}(\mathbb{R}, V)$. We handle differentiation first, so we must show that $(r, t) \mapsto \dot{p_r}(t)$ is smooth. By pre-convenience of V, we may check this in each $l \in L^{\infty}(V)$: we must show that $(r, t) \mapsto l \circ \dot{p_r}(t) = \hat{l \circ p_r}(t)$ is smooth. But this map is simply the partial derivative of the smooth map $(r, t) \mapsto l \circ p_r(t)$ in the t-direction, hence is itself smooth.

For integration, we must show that $(r,t) \mapsto \int p_r(t)$ is smooth. By pre-convenience of V, we may check this in each $l \in L^{\infty}(V)$: we must show that $(r,t) \mapsto l \circ \int p_r(t) = \int_0^t l \circ p_r$ is smooth. But this map is simply the integration of the smooth map $(r,t) \mapsto l \circ p_r(t)$, hence is itself smooth.

For the remaining assertions, simply check all required equalities scalar-wise. \Box

Lemma 4.17 (Hadamard's lemma). Fix a convenient dvs V. If $c : \mathbb{R} \to V$ is a smooth curve, there exists some smooth curve $h : \mathbb{R} \to V$ with c(t) = c(0) + th(t).

Proof. Consider the auxiliary smooth curve H(s) := c(st). Then $\dot{H}(s) = \dot{c}(st)t$ (check scalar-wise, and use the chain rule), and so by the fundamental theorem of calculus and linearity of the integral,

$$c(t) - c(0) = H(1) - H(0) = \int_0^1 \dot{c}(st)tds = t \int_0^1 \dot{c}(st)ds$$

Thus setting $h(t) := \int_0^1 \dot{c}(st) ds$ will work, if h is smooth. This is true because h is the composition of the smooth maps:

$$\mathbb{R} \to C^{\infty}(\mathbb{R}, V)$$
, given by $t \mapsto [s \mapsto \dot{c}(st)]$, and
 $\int : C^{\infty}(\mathbb{R}, V) \to C^{\infty}(\mathbb{R}, V)$, and
 $\operatorname{ev}_1 : C^{\infty}(\mathbb{R}, V) \to V.$

Proposition 4.18. Let V be a convenient vector space. Write $D_0 : C^{\infty}(\mathbb{R}, V) \to V$ for the map $c \mapsto \dot{c}(0)$. In the diagram below,



where we set $\gamma(u, v) := [t \mapsto u + tv]$, we have:

- (a) the maps γ and (ev_0, D_0) are smooth,
- (b) the map γ is a section of (ev_0, D_0) , hence (ev_0, D_0) is a subduction,
- (c) the triangle with downward arrows commutes: $\nu \circ (ev_0, D_0) = \partial$
- (d) the map ν is injective, hence an isomorphism.

Proof. We prove each part in order.

(a) The map γ is smooth if and only if the map

$$(V \times V) \times \mathbb{R} \to V, \quad (v_0, v, t) \mapsto \gamma(u, v)(t) = u + tv$$

is smooth, and this true because V carries a vector space diffeology. The map (ev_0, D_0) is smooth because ev_0 is smooth, and D_0 is the composition of the smooth map ev_0 with the smooth map $c \mapsto \dot{c}$ (Lemma 4.16).

(b) Clearly $ev_0(\gamma(u, v)) = v_0$. As for D_0 , we observe that for each $l \in L^{\infty}(V)$,

$$l \circ D_0(\gamma(u, v)) = \partial_{t=0}(l(u) + tl(v)) = l(v),$$

hence by linear separation $D_0(\gamma(u, v)) = v$. Having a section implies (ev_0, D_0) is a subduction.

(c) Fix a curve c, and denote $(u, v) := (c(0), \dot{c}(0))$. We need to show that $\partial c = \partial [u+tv]$. By applying Hadamard's lemma 4.17 to c, we write $c(t) = u + th_1(t)$ for smooth h_1 with h(0) = v (confirm this scalar-wise). Applying Hadamard's lemma to h_1 , we get $h_1(t) = u + th(t)$ for smooth h. Thus $c(t) = u + tv + t^2h(t)$.

We form the commuting diagram



where H(t, t') := u + tv + t'h(t). This diagram shows that $\partial[u + tv] = \partial c$, as required.

(d) To show ν is injective, consider an arbitrary link in a zig-zag witnessing $\partial [u_b + tv_b] = \partial [u_e + tv_e]$ (the subscripts denote "beginning" and "end"):



Let c, c' denote any curves in U, U' whose derivatives at 0 are v, v'. Then, for any $l \in L^{\infty}(V)$, observe:

$$l(\overrightarrow{pc}(0)) = \overrightarrow{lpc}(0) = \overrightarrow{lp'fc}(0) = \widehat{T}(lp') \circ \overrightarrow{fc}(0) = \widehat{T}(lp') \circ \overrightarrow{c'}(0),$$

because $\widehat{fc}(0) = \dot{c'}(0) = v'$. Thus $\widehat{pc}(0) = \widehat{p'c'}(0)$.

At the beginning of the zig-zag, clearly $p(u_b + tv_b)$ has weak derivative v_b at t = 0, and similarly $p(u_e + tv_e)$ has weak derivative v_e . Since these weak derivatives are preserved across the zig-zag, we find that $v_b = v_e$. As $u_b = u_e$ a necessary condition for $\partial [u_b + tv_b] = \partial [u_e + tv_e]$, we have proved that ν is injective.

The map ν is an isomorphism because it is an injective subduction.

We have now arrived at a main result of this section.

Proposition 4.19. A pre-convenient and linearly separated dvs is convenient if and only if it is tangent stable.

Proof. Fix a pre-convenient and linearly separated dvs V. If it is convenient, Proposition 4.7 says it is tangent stable. If it is tangent-stable, Proposition 4.18 says it is tangent stable.

Remark 4.20. When V is tangent-stable, and $U \subseteq V$ is D-open, we have $TU \cong U \times V$, and the rest of the tangent structure is identical to that on Cart. This also gives the tangent structure on any space that is locally diffeomorphic to V.

4.2. From convenient manifolds to diffeology. Now we import topological vector spaces into diffeology. We recall the notion of locally convex topological vector spaces.

Definition 4.21. A *locally convex topological vector space*, abbreviated "lctvs," is a vector space *E* equipped with a vector space topology that is generated by a family of seminorms. A *morphism* of lctvs is a continuous linear map.

Letvs, and morphisms between them, form the category $\mathcal{L}CTVS$. We will write $L^*(E, E_1) := \underline{\mathcal{L}CTVS}(E, E_1)$, and when $E_1 = \mathbb{R}$, we write simply $L^*(E)$.

Definition 4.22. A lctvs is *separated* if the seminorms defining its topology separate points.

Example 4.23. A *Banach* space is a lctvs whose topology is determined by a complete norm. A *Fréchet* space is a separated lctvs whose topology is generated by a countable collection of seminorms, and which is complete with respect to these seminorms. \triangleright

Given a separated lctvs E, we consider two reasonable diffeologies.

• The diffeology $\mathscr{D}_{c}(E)$. Given a map $c: U \to E$, with $U \subseteq \mathbb{R}$ open, we call c differentiable if for each $t \in U$, the difference quotient

$$c'(t) = c^{(1)}(t) := \lim_{h \to 0} \frac{c(t+h) - c(t)}{h}$$

exists. We inductively define $c^{(k)}$, and declare c to be $\mathscr{D}_{c}(E)$ --smooth⁷ if $c^{(k)}$ exists for each $k = 0, 1, \ldots$. The plots of $\mathscr{D}_{c}(E)$ are those parametrizations $p: U \to E$ for which $p \circ c$ is $\mathscr{D}_{c}(E)$ -smooth for all smooth (in the usual sense) curves $c: V \to U$. • The diffeology $\Pi L^{*}(E)$.

In the diffeology $\mathscr{D}_{c}(E)$, it is apparent and natural that the velocity of the curve c is c'. However, the assignment $E \mapsto (E, \mathscr{D}_{c}(E))$ is not a functor from $\mathcal{L}CTVS$ to \mathcal{D} flg, because continuous linear maps between lctvs need not be \mathscr{D}_{c} -smooth. On the other hand, the assignment $E \mapsto \Pi L^{*}(E)$ is functorial. We always have $\mathscr{D}_{c}(E) \subseteq \Pi L^{*}(E)$. Kriegl and Michor single out those separated lctvs E for which $\mathscr{D}_{c}(E) = \Pi L^{*}(E)$, and call them "convenient."

The condition $\mathscr{D}_{c}(E) = \Pi L^{*}(E)$ is not sufficient to ensure that the diffeological vector space $(E, \Pi L^{*}(E))$ is convenient in the sense of Definition 4.13. This is because, in general,

$$L^*(E) \subseteq L^{\infty}(E, \Pi L^*(E)),$$

and equality need not hold. A remedy is to consider only bornological lctvs.

Definition 4.24. A lctvs E is *bornological* if every bounded seminorm on E is continuous.

Example 4.25. Banach and Fréchet spaces are bornological.

Theorem II. Let E be a Hausdorff bornological locally convex topological vector space. The following are equivalent

- $(E, \Pi L^*(E))$ is convenient.
- $(E, \Pi L^*(E))$ is tangent-stable.
- $\mathscr{D}_c(E) = \Pi L^*(E)$

Proof. Suppose that $(E, \Pi L^*(E))$ is convenient. Then every smooth curve c has a weak-derivative \dot{c} . One proves that $\dot{c} = c'$. This shows that the 1-plots of $\Pi L^*(E)$ agree with the

23

 \triangleright

⁷the "c" stands for "convenient"

1-plots of $\mathscr{D}_{c}(E)$. But both diffeologies are determined by their 1-plots, so the diffeologies agree.

Conversely, if $\mathscr{D}_{c}(E) = \Pi L^{*}(E)$, then given any smooth curve c, we may always form c'. One checks that for every $l \in L^{*}(E)$, we have $\hat{l \circ c} = l \circ c'$. Then, because E is bornological, $L^{\infty}(E, \Pi L^{*}(E)) = L^{*}(E)$, so c' is the required weak-derivative of c.

Since E is Hausdorff and bornological, convenience of $(E, \Pi L^*(E))$ is equivalent to tangent-stability of $(E, \Pi L^*(E))$ by Proposition 4.19.

In light of Theorem II, we will call a separated bornological lctvs E convenient if $(E, \Pi L^*(E))$ is convenient.

Corollary 4.26. Banach and Fréchet spaces are convenient.

We now work towards convenient manifolds. We need the notion of a non-linear smooth map between convenient vector spaces. For two separated lctvs E, F, Kriegl and Michor take a smooth map to be any $f : E \to F$ which is \mathscr{D}_{c} -smooth. Thus if E and F are convenient lctvs, the smooth maps between them are precisely the diffeologically smooth ones, when they are viewed as convenient diffeological vector spaces.

Remark 4.27. There is another notion of smooth map between lctvs. Let E, F be separated lctvs. We call a map $f : E \to F$ Michel-Bastiani smooth if all its iterated directional derivatives

 $Df: E \times E \to F, \quad D^2f: (E \times E) \times (E \times E) \to F, \dots,$

exist. Here $Df(u, v) := [t \mapsto f(u + tv)]'(0)$. Every MB-smooth map is \mathscr{D}_{c} -smooth, but the converse need not hold, even if E and F are convenient lctvs; Glöckner [Glö06] gives a counter-example. However, when E and F are Fréchet spaces, Frölicher [Frö81] proved that every \mathscr{D}_{c} -smooth map is Michel-Bastiani smooth.

Now we need a notion of locality. Fix a convenient lctvs E. In defining manifolds modelled on E, Kriegl and Michor equip E not with its original topology, but with its "c[∞]-topology." This is the final topology with respect to the c-smooth curves $C^{\infty}(\mathbb{R}, (E, \mathscr{D}_{c}(E)))$. This is the same as the *D*-topology of $(E, \mathscr{D}_{c}(E))$.

Lemma 4.28 ([CSW14, Theorem 3.7]). Let X be a diffeological space. Its D-topology is the final topology with respect to its smooth curves $C^{\infty}(\mathbb{R}, X)$.

Kriegl and Michor proceed to define a convenient manifold M modelled on E precisely as one does for finite-dimensional manifolds modelled on \mathbb{R}^n : charts take values in c^{∞} open subsets of E, and transition functions are required to be \mathscr{D}_c -smooth. Smooth maps between convenient manifolds (perhaps modelled on different convenient spaces) are those maps which are \mathscr{D}_c -smooth in charts. They equip M with the final topology induced by the charts, and require this to be Hausdorff. This yields a category of manifolds \mathcal{C} onvMan.

Example 4.29. Banach and Fréchet manifolds are convenient manifolds. This is because the original topology on a Banach or Fréchet space, and their c^{∞} -topology, agree, and because the Michel-Bastiani smooth maps coincide with the \mathscr{D}_{c} -smooth ones (Remark 4.27).

On the other hand, we may also consider those diffeological spaces which are locally diffeomorphic to convenient dvs, and whose D-topology is Hausdorff. Exactly as in the case of finite-dimensional manifolds, there is no difference in the two approaches.

Proposition 4.30. Let ConvMan denote the category of convenient manifolds as described above. The canonical functor

$$ConvMan \to Dflg,$$

which assigns to a convenient manifold M the diffeology consisting of \mathscr{D}_c -smooth maps $U \to M$, is full and faithful. Its essential image consists of those diffeological spaces which are locally diffeomorphic to convenient spaces, and which have Hausdorff D-topology.

Proof. Proceed as in [Igl13, Article 4.3], which treats the finite-dimensional case. \Box

Remark 4.31. We have seen that convenient lctvs are all pre-convenient, hence reflexive, diffeological vector spaces. However, reflexivity is a property that any diffeological space may possess. It is not clear if convenient manifolds embed into the subcategory of reflexive diffeological spaces. \triangleleft

This embedding is also compatible with the relevant tangent functors. Kriegl and Michor define the tangent functor on c^{∞} -open subsets of convenient lctvs by

$$\hat{S}(U \subseteq E) := U \times E, \quad \hat{S}f(e,v) := (f(e), [t \mapsto f(e+tv)]'(0)),$$

and then define $S : ConvMan \to ConvMan$ as one usually does for finite-dimensional manifolds.

Proposition 4.32. When we view C onvMan as a subcategory of \mathcal{E} lst, its tangent structure coincides with ours on \mathcal{E} lst.

Proof. Fix a convenient manifold M, and a some D-open subset U which is diffeomorphic to a D-open subset of a convenient space V. Remark 4.20 details the tangent structure on U. But all of these maps coincide with those from tangent structure used by [KM97], because on open subsets of V, we have S = T. Thus we have the same tangent structure.

From this discussion, it follows that any results or computations for convenient, and in particular Banach or Fréchet, manifolds from the literature are also valid in \mathcal{D} flg. For instance, if G is a Banach-Lie group, meaning a group object in the category of convenient manifolds, modelled on a Banach space, with Banach-Lie algebra \mathfrak{g} , then when we view G as a diffeological group, it is elastic with Lie algebra \mathfrak{g} .

5. Integrating central extensions

In this final section, we integrate some non-enlargeable Banach-Lie algebras to elastic diffeological groups. In Subsection 5.1 we review central extensions of diffeological groups and diffeological Lie algebras, culminating in a differentiation procedure for central extensions of elastic groups. In Subsection 5.2, we restrict our attention to central extensions of convenient Lie algebras, and perform the desired integration into elastic groups.

5.1. Diffeological central extensions. We recall the basic theory of central extensions of groups and Lie algebras. Fix a group G and an abelian group A. We use concatenation to denote multiplication in G, and we use + for the group addition in A. Implicitly we assume that G acts trivially on A.

Definition 5.1. A *central extension* of G by A consists of a group \hat{G} and a short-exact sequence $A \hookrightarrow \hat{G} \twoheadrightarrow G$.

Central extensions of G are classified by the second degree group cohomology. A 2cochain on G valued in A is simply a map $G^2 \to A$. The collection of 2-cochains forms a group with pointwise addition. The 2-cocycles and 2-coboundaries satisfy, respectively

$$f(g_0g_1, g_2) + f(g_0, g_1) = f(g_0, g_1g_2) + f(g_1, g_2)$$
, and
 $f(g_0, g_1) = -h(g_0g_1) + h(g_1) + h(g_0)$ for some $h: G \to A$.

Denote the resulting cohomology group by $H^2(G, A)$. We may insist that our cochains are normalized. Let

$$C^{2}(G, A) := \{ f : G^{2} \to A \mid f(g_{0}, g_{1}) = 0 \text{ if } g_{i} = e \text{ for some } i \},\$$

with subgroups of *n*-cocycles and *n*-coboundaries denoted by $Z^2(G, A)$ and $B^2(G, A)$, respectively. Then we still have $H^2(G, A) = Z^2(G, A)/B^2(G, A)$.

To each cocycle $f \in Z^2(G, A)$, we associate a central extension of G by A, denoted \hat{G}_f or $G \times_f A$. This has underlying set $G \times A$, and multiplication

$$(g, a)(h, b) := (gh, a + b + f(g, h))$$

The inclusion of A into the second factor of \hat{G} , followed by the projection of the first factor to G, forms the required short-exact sequence. It is straightforward to check that \hat{G}_f is a central extension. Moreover, $H^2(G, A)$ classifies central extensions of G by A up to equivalence.

The restriction to diffeological central extensions is fairly straightforward. Let G be a diffeological group, and let A be an abelian diffeological group.

Definition 5.2. A diffeological central extension of G by A consists of a diffeological group \hat{G} and a short-exact sequence $A \stackrel{\iota}{\hookrightarrow} \hat{G} \stackrel{\pi}{\twoheadrightarrow} G$, such that ι is an induction, and π is a subduction.

The conditions on ι and π ensure that A identifies with its image $\iota(A) \subseteq G$, viewed as a diffeological subgroup, and that G identifies with $\hat{G}/\iota(A)$, where the latter carries the quotient diffeology. A fortiori $\pi : \hat{G} \to G$ is a principal A-bundle. We caution that in \mathcal{D} flg, principal bundles are not necessarily locally trivial; π does not necessarily admit a smooth local section.

Example 5.3. Fix some dense and countable subgroup K of \mathbb{R} . Then $K \hookrightarrow \mathbb{R} \twoheadrightarrow \mathbb{R}/K$ expresses \mathbb{R} as a central extension of \mathbb{R}/K by K, where K carries the subspace diffeology, and \mathbb{R}/K carries the quotient diffeology. But the principal bundle $\mathbb{R} \to \mathbb{R}/K$ is not locally trivial.

This complicates the task of classifying diffeological group extensions by cohomology. On the other hand, we do not work in this generality.

Definition 5.4. Let $C^2_{s,\text{loc}}(G, A)$ denote those normalized group cochains which are smooth on a neighbourhood of $(e, \ldots, e) \in G^n$ of the form $U \times \cdots \times U$, for some $U \subseteq G$ open. This gives, suppressing the (G, A) notation, cocycles $Z^2_{s,\text{loc}}$, coboundaries $B^2_{s,\text{loc}}$, and a cohomology $H^2_{s,\text{loc}}$.

The "s" is for "smooth," and "loc" is for "local."

Remark 5.5. The D-topology functor $D: \mathcal{D}flg \to \mathcal{T}$ op does not generally preserve products; the natural map $D(X \times Y) \to D(X) \times D(Y)$ is only guaranteed to be continuous. This means that a neighbourhood of (e, \ldots, e) in $D(G^n)$ need not contain any set of the form $U \times \cdots \times U$. Therefore the elements of $C^2_{s,\text{loc}}(G, A)$ are not merely maps $G^2 \to A$ that are smooth on some neighbourhood of (e, e). Also observe that D(G) need not be a topological group.

For a connected diffeological group G, the cohomology $H^2_{s,\text{loc}}(G, A)$ classifies, up to equivalence, central extensions of G by A that are locally split, i.e. for which the projection π admits a section $\sigma: G \to \hat{G}$ with $\sigma(e) = \hat{e}$ that is smooth in a neighbourhood of the identity. This is a consequence of two lemmas.

Lemma 5.6. Let G be a group, and U be a subset of G equipped with a diffeology \mathscr{D}_U . Suppose we have $V \subseteq U$ a D-open subset containing the identity, such that

(i) $V = V^{-1}$ and $V \cdot V \subseteq U$;

(ii) the multiplication and inversion

$$m: V \times V \to U, \quad i: V \to V$$

are smooth (in the subspace diffeology of V, denoted \mathscr{D}_V);

(iii) V generates G as a group.

Then the finest group diffeology on G containing \mathscr{D}_V is

$$\mathscr{D} := \{ p \mid p \text{ locally has the form } p = g \cdot q, \text{ for } g \in G, q \in \mathscr{D}_V \}.$$

Moreover

- (a) The subset V is a D-open subset of (G, \mathscr{D}) .
- (b) If $V' \subseteq U$ is another open neighbourhood of the identity satisfying (i) through (iii), then \mathscr{D} is also the finest group diffeology containing $\mathscr{D}_{V'}$.

The proof may seem long, but it employs standard techniques.

Proof. The family $\{g \cdot q\}$ covers G. The plots of the diffeology generated by this family are those parametrizations p which locally have the form $p = (g \cdot q) \circ F$, for some smooth map F between open subsets of Cartesian spaces. But $q \circ F$ remains a plot of \mathscr{D}_V , so we see that the diffeology generated by $\{g \cdot q\}$ is precisely \mathscr{D} . This shows that \mathscr{D} is a diffeology. Now we must show it is a group diffeology.

We begin with the following observation. Take a plot q of V, with $r_0 \in \text{dom } q$. Then $q(r_0)^{-1} \cdot q$ is a plot of U (because $m : V \times V \to U$ is smooth), sending r_0 to e. Since V is

a D-open neighbourhood of e, the restriction q_1 of $q(r_0)^{-1} \cdot q$ to some neighbourhood of r_0 is a plot of V. Thus near r_0 , we locally write $q = q(r_0) \cdot q_1$, for q_1 a plot of V pointed at e.

Now consider $g \in G$, and the map $q \cdot g$. Write $g = g_1 \cdots g_l$, with $g_i \in V$, which is possible since V generates G. Repeatedly applying the observation above, first to qyielding q_1 , then to $q_1 \cdot g_1$, yielding q_2 , etc., we find that near $r_0 \in \text{dom } q$,

$$q \cdot g = q(r_0) \cdot (q_1 \cdot g_1) \cdots q_l$$

= $q(r_0) \cdot g_1 \cdot (q_2 \cdot g_2) \cdots q_l$
...
= $q(r_0) \cdot g \cdot q_{l+1}$
=: $\tilde{g} \cdot \tilde{q}$.

The plot \tilde{q} is a plot of V mapping r_0 to e.

We now check that the multiplication $m: G \times G \to G$ is smooth. It suffices to check that $g'q' \cdot gq$ is a plot of \mathscr{D} for plots q', q of V on a shared domain, and $g', g \in G$. Fix $r_0 \in \operatorname{dom} q' = \operatorname{dom} q$. We may locally write $q' \cdot g$ as $\tilde{g} \cdot \tilde{q'}$, where $\tilde{q'}$ is a plot of V mapping r_0 to e. Then, locally

$$g'q' \cdot gq = g'\tilde{g} \cdot q'\tilde{q},$$

and $\tilde{q}'q$ is a plot of U since $m: V \times V \to U$ is smooth. Since $\tilde{q}'q$ maps r_0 to $q(r_0) \in V$, this restricts to a plot of V, and we conclude that $g'q' \cdot gq$ is locally $g'\tilde{g} \in G$ times a plot of V. Therefore the multiplication m is smooth.

We similarly show inversion is smooth. It suffices to show $(g \cdot q)^{-1}$ is a plot. Fix $r_0 \in \text{dom } q$. We have $(g \cdot q)^{-1} = q^{-1} \cdot g^{-1}$, and q^{-1} is a plot of V because $i : V \to V$ is smooth. Near r_0 , the product $q^{-1} \cdot g^{-1}$ therefore has the form $\tilde{g} \cdot q^{-1}$, for a plot q^{-1} of V. This shows $(g \cdot q)^{-1}$ is locally a plot of \mathscr{D} , hence is a plot.

Thus we have verified that \mathscr{D} is a group diffeology containing \mathscr{D}_V . Let \mathscr{D}' denote the finest such diffeology. Then $\mathscr{D}' \subseteq \mathscr{D}$. Conversely, given $g \cdot q$ in the generating family for \mathscr{D} , we have $q \in \mathscr{D}_V \subseteq \mathscr{D}'$. Left multiplication $L_g : (G, \mathscr{D}') \to (G, \mathscr{D}')$ is smooth, so $g \cdot q = L_g \circ q$ is a plot of \mathscr{D}' . It follows that $\mathscr{D} \subseteq \mathscr{D}'$, as required.

Let us now treat (a). To see V is D-open, we may show that $(g \cdot q)^{-1}(V)$ is open for all $g \in G$ and $q \in \mathscr{D}_V$. If $g \in V$, denote by L_g the map $(V, \mathscr{D}_V) \to (U, \mathscr{D}_U)$ given by multiplication by g. This is smooth. Then

$$(g \cdot q)^{-1}(V) = \{r \mid q(r) \in (g^{-1} \cdot V) \cap V\} = \{r \mid q(r) \in L_g^{-1}(V)\},\$$

and $L_g^{-1}(V)$ is D-open in (V, \mathscr{D}_V) by smoothness of L_g . Therefore $(g \cdot q)^{-1}(V)$ is open. For general $g \in G$, at r_0 such that $g \cdot q(r_0) \in V$, write locally $q = q(r_0) \cdot q_1$, for a plot q_1 of V. Then $g \cdot q$ is, locally, $(g \cdot q(r_0)) \cdot q_1$, and we may apply the result of the previous case because $g \cdot q(r_0) \in V$. This provides an open neighbourhood of $r_0 \in \text{dom } q$ which is sent by $g \cdot q$ into V, and thus proves that $(g \cdot q)^{-1}(V)$ is open in dom q. Therefore V is D-open.

Finally, we treat (b). For $V' \subseteq U$ a D-open neighbourhood of the identity satisfying (i) through (iii), let \mathscr{D}' denote the finest group diffeology containing $\mathscr{D}_{V'}$. Suppose first

that $V' \subseteq V$. Then $\mathscr{D}' \subseteq \mathscr{D}$ is clear. Conversely, note that given $q \in \mathscr{D}_V$, and $r_0 \in \text{dom } q$, we may locally write $q = q(r_0)^{-1} \cdot q_1$, where q_1 is a plot of V'. We may do so because V'is D-open in V. This shows that $\mathscr{D} \subseteq \mathscr{D}'$. For general V', consider $V \cap V'$. The finest diffeologies containing \mathscr{D}_V and $\mathscr{D}_{V'}$, respectively, must coincide with the finest diffeology containing $V \cap V'$, hence with each other. \Box

Lemma 5.7. Let $f \in Z^2_{s,loc}(G, A)$ be a locally smooth 2-cocycle. Then $\hat{G}_f = G \times_f A$ admits a group diffeology which makes it a locally split central extension of G by A.

Proof. Say $f: U \times U \to A$ is smooth, for an open neighbourhood U of the identity. We may assume U is connected. Choose an open neighbourhood V of the identity in U such that $V^{-1} = V$, and $V \cdot V \subseteq U$. Being an open neighbourhood of the identity in a topological group, V generates G. Smoothness of f on $V \times V$ implies that in \hat{G} ,

multiplication : $(V \times A) \times (V \times A) \rightarrow U \times A$, and inversion : $(V \times A) \rightarrow (V \times A)$

are smooth. Also, $V \times A$ generates \hat{G} . Then Lemma 5.6, applied to the group \hat{G} , the diffeological space $U \times A$ (with the product diffeology), and the D-open subset $V \times A$, furnishes \hat{G} with a group diffeology for which $V \times A$ is D-open.

The inclusion $A \hookrightarrow \hat{G}$, given by $a \mapsto (e, a)$ is an induction into $V \times A$, hence into \hat{G} . Now we check that the projection pr : $\hat{G} \to G$, given by $(g, a) \mapsto g$, is a subduction. Fix a plot p of G, and take $r_0 \in \text{dom } p$. Near r_0 , we may write p as $p = p(r_0) \cdot q$, for a plot qof V. Then $(p(r_0), 0) \cdot (q, 0)$ is the required plot of \hat{G} which projects to $p(r_0)q = p$.

Finally, the map $V \to V \times A$ given by $g \mapsto (g, 0)$ is smooth, thus it is smooth as a map $V \to \hat{G}$. This is a section of the projection, and its existence corresponds to a local trivialization of $\hat{G} \to G$. Note that this section is not a group homomorphism, so it may not extend to a smooth map beyond V.

Proposition 5.8. Let G be a connected diffeological group, and A an abelian diffeological group. The cohomology $H^2_{s,loc}(G, A)$ classifies, up to equivalence, the locally split central extensions of G by \hat{G} .

Proof. Given $f \in Z^2_{s,\text{loc}}(G, A)$, Lemma 5.7 gives the corresponding locally split central extension of G by A. Conversely, given such an extension, let $\sigma : G \to A$ be a section of the projection that is smooth near e. Then

$$f(g,h) := \sigma(g)\sigma(h)\sigma(gh)^{-1}$$

is the element of $Z^2_{s,\text{loc}}(G,A)$ that corresponds to the original \hat{G} .

Now we turn to Lie algebras. Let \mathfrak{g} be a Lie algebra, and \mathfrak{a} be a trivial Lie algebra, thus merely a vector space. We implicitly assume that \mathfrak{g} acts trivially on \mathfrak{a} .

Definition 5.9. A *central extension* of \mathfrak{g} by \mathfrak{a} consists of a Lie algebra $\hat{\mathfrak{g}}$ and a short-exact sequence of Lie algebras $\mathfrak{a} \hookrightarrow \hat{\mathfrak{g}} \twoheadrightarrow \mathfrak{g}$.

Like with groups, central extensions of Lie algebras are classified by second-degree cohomology. A 2-cocycle for \mathfrak{g} valued in \mathfrak{a} is an anti-symmetric bilinear map $\omega : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{a}$

satisfying $\sum_{\bigcirc} \omega([x, y], z) = 0.^8$ A 2-coboundary is a 2-cocycle which, additionally, is of the form $\omega(x, y) = b([x, y])$, for some linear map $b : \mathfrak{g} \to \mathfrak{a}$. We use $Z^2(\mathfrak{g}, \mathfrak{a}), B^2(\mathfrak{g}, \mathfrak{a})$, and $H^2(\mathfrak{g}, \mathfrak{a})$, to denote the 2-cocycles, 2-coboundaries, and the resulting cohomology group, respectively.

To each $\omega \in Z^2(\mathfrak{g}, \mathfrak{a})$, we associate a central extension of \mathfrak{g} by \mathfrak{a} , denoted $\hat{\mathfrak{g}}_{\omega}$ or $\mathfrak{g} \oplus_{\omega} \mathfrak{a}$. This has underlying vector space $\mathfrak{g} \oplus \mathfrak{a}$, and Lie bracket

$$[(x, a), (y, b)] := ([x, y], \omega(a, b)).$$

The inclusion of \mathfrak{a} into the second factor of $\hat{\mathfrak{g}}_{\omega}$, followed by the projection of the first factor to \mathfrak{g} , forms the required short exact sequence. Like with groups, the second degree Lie algebra cohomology $H^2(\mathfrak{g},\mathfrak{a})$ classifies central extensions of \mathfrak{g} by \mathfrak{a} up to equivalence.

We now let \mathfrak{g} be a diffeological Lie algebra, and \mathfrak{a} be a trivial diffeological Lie algebra.

Definition 5.10. A *diffeological central extension* of \mathfrak{g} by \mathfrak{a} is a Lie algebra central extension of \mathfrak{g} by \mathfrak{a} , which is simultaneously a diffeological central extension of the underlying abelian diffeological groups.

Since central extensions of diffeological Lie algebras are also central extensions of diffeological groups, the same considerations apply. However, we will be content to work with globally split central extensions, and thus do not need versions of Lemmas 5.6 or 5.7. We have, suppressing $(\mathfrak{g}, \mathfrak{a})$ from the notation, the groups Z_s^2 and B_s^2 of smooth 2-cocycles and 2-coboundaries, and this gives the smooth Lie algebra cohomology H_s^2 . Given $\omega \in Z_s^2$, the associated central extension $\hat{\mathfrak{g}}_{\omega}$, equipped with the product diffeology, is a split diffeological central extension of \mathfrak{g} with values in \mathfrak{a} . The cohomology H_s^2 classifies such central extensions up to equivalence.

Proposition 5.11. Let \mathfrak{g} be a diffeological Lie algebra, and \mathfrak{a} be a trivial diffeological Lie algebra. The second degree smooth Lie algebra cohomology $H_s^2(\mathfrak{g},\mathfrak{a})$ classifies split diffeological central extensions of \mathfrak{g} by \mathfrak{a} , up to equivalence.

The following table summarizes our discussion.

| Extension | Cohomology | Correspondence |
|---|------------------------------------|--|
| Central extensions of groups | $H^2(G, A)$ | $f \mapsto \hat{G}_f := G \times_f A$ |
| Locally split central extensions of diffeological groups | $H^2_{s,\mathrm{loc}}(G,A)$ | $f \mapsto \hat{G}_f$, diffeology in Lemma 5.7 |
| Central extensions of Lie algebras | $H^2(\mathfrak{g},\mathfrak{a})$ | $\omega\mapsto \hat{\mathfrak{g}}_\omega:=\mathfrak{g}\oplus_\omega\mathfrak{a}$ |
| Split central extensions of diffeological Lie algebras | $H^2_s(\mathfrak{g},\mathfrak{a})$ | $\omega \mapsto \hat{\mathfrak{g}}_{\omega}$, product diffeology. |

To end this section, we differentiate locally smooth cocycles. Let G be an elastic group, with Lie algebra \mathfrak{g} . Our differentiation of a 2-cocycle $f \in Z^2_{s,\text{loc}}(G, A)$ to a 2-cocycle $L(f) \in Z^2_s(\mathfrak{g}, \mathfrak{a})$ amounts to taking the Hessian of f at (e, e). Suppose that M, N, and P are elastic spaces, and $f: M \times N \to P$ is smooth. Then we have the map

$$T_{(1)}T_{(2)}f:TM \times TN \to T^2P$$

⁸We use \sum_{\bigcirc} to denote a sum over cyclic permutations of the arguments.

If, moreover, we have $m_0 \in M$, and $n_0 \in N$, and $p_0 \in P$ such that $f(m_0, \cdot) = f(\cdot, n_0) = p_0$, then Lemma A.11 implies that, whenever $v \in T_{m_0}M$ and $w \in T_{n_0}N$,

$$T_{(1)}T_{(2)}f(v,w) = T_{(2)}T_{(1)}f(v,w) \in \lambda(T_{p_0}P) \cong T_{p_0}P.$$

The map $T_{(1)}T_{(2)}f$ is also bilinear as a map $T_{m_0}M \times T_{n_0}N \to T_{p_0}P$.

Definition 5.12. Let $f \in Z^2_{s, \text{loc}}(G, A)$ be a 2-cocycle. We denote

 $d^2 f := T_{(1)}T_{(2)}f$ restricted to a map $\mathfrak{g} \times \mathfrak{g} \to \mathfrak{a}$,

and we define $L(f) : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{a}$ by

$$L(f)(\xi,\eta) := T_{(1)}T_{(2)}f(\xi,\eta) - T_{(2)}T_{(1)}f(\eta,\xi).$$

| | ۰. | |
|----|----|----|
| | | ۰. |
| | | |
| ς. | 4 | |
| | ٠ | |
| | | |
| | | |
| | | |

Now we can differentiate cocycles.

Proposition 5.13. Let G be an elastic diffeological group, and A be an abelian elastic diffeological group. Let $f \in Z^2_{s,loc}(G,A)$. Then L(f) is a smooth Lie algebra 2-cocycle for \mathfrak{g} with values in \mathfrak{a} , and the Lie algebra of \hat{G}_f is $\hat{\mathfrak{g}}_{L(f)}$.

Proof. Suppose that f is smooth on $U \times U$, for U an open neighbourhood of the identity in G. Since d^2f is a smooth bilinear map, so is L(f). Anti-symmetry is clear. Now we evaluate the Lie bracket on \hat{G}_f . Recall that the conjugation in \hat{G}_f is explicitly given by

$$(g,z)(h,w)(g,z)^{-1} = (ghg^{-1}, w + f(g,h) - f(ghg^{-1},g)).$$

Applying $T_{(1)}T_{(2)}$ to this expression, and evaluating at $((\xi, V), (\eta, W))$ yields, on the left side, $[(\xi, V), (\eta, W)]$ by Lemma A.16. On the right side, we have $[\xi, \eta]$ in the first slot, and the f(g, h) term gives $T_{(1)}T_{(2)}f(\xi, \eta)$. For the $-f(ghg^{-1}, g)$ term, we consider

$$f \circ (c, \mathrm{pr}_1) : G \times G \to G$$

Here c denotes conjugation. Applying $T_{(1)}T_{(2)}$ to this gives

$$\begin{split} T_{(1)}T_{(2)}(f\circ(c,\mathrm{pr}_1)) &= T_{(1)}(Tf\circ(Tc,\mathrm{pr}_1)\circ(0\times 1)) \\ &= T_{(1)}(Tf\circ(T_{(2)}c,0\circ\mathrm{pr}_1)) \\ &= T_{(1)}(T_{(1)}f\circ(T_{(2)}c,\mathrm{pr}_1)) \\ &= T(T_{(1)}f\circ(T_{(2)}c,\mathrm{pr}_1))\circ(1\times 0) \\ &= TT_{(1)}f\circ(T_{(1)}T_{(2)}c,\mathrm{pr}_1). \end{split}$$

Apply Lemma A.10 to break this into a sum:

$$TT_{(1)}f \circ (T_{(1)}T_{(2)}c, \mathrm{pr}_1) = T_{(1)}T_{(1)}f \circ (T_{(1)}T_{(2)}c, \pi \circ \mathrm{pr}_1) + T_{(2)}T_{(1)}f \circ (\pi \circ T_{(1)}T_{(2)}c, \mathrm{pr}_1)$$

The first summand will vanish, because $f(e, \cdot) = f(\cdot, e) = e$, and the second is $T_{(2)}T_{(1)}f \circ (\operatorname{pr}_2, \operatorname{pr}_1)$. Thus upon differentiating, the $-f(ghg^{-1}, g)$ term gives $T_{(2)}T_{(1)}f(\eta, \xi) =$

 $T_{(1)}T_{(2)}f(\eta,\xi)$. Therefore

$$[(\xi, V), (\eta, W)] = ([\xi, \eta], T_{(1)}T_{(2)}f(\xi, \eta) - T_{(1)}T_{(2)}f(\eta, \xi))$$

= ([\xi, \eta], L(f)(\xi, \eta)).

The fact that the Lie bracket can be expressed in this way implies that L(f) is a Lie algebra 2-cocycle in $Z_s^2(\mathfrak{g}, \mathfrak{a})$, and that $\operatorname{Lie}(\hat{G}_f) = \hat{\mathfrak{g}}_{L(f)}$.

5.2. Integrating convenient Lie algebras. Having established how to differentiate locally smooth 2-cocycles, we turn to the problem of integration. We draw from results in the literature [Nee02; Woc11]. These authors work in the category of infinite-dimensional manifolds modelled on lctvs, and Michel-Bastiani smooth maps between them (cf. Remark 4.27), but we will apply their results to convenient manifolds.

For our preliminary discussion, fix a connected and simply connected convenient Lie group G, with Lie algebra \mathfrak{g} , and a convenient abelian Lie algebra \mathfrak{a} . Also take a 2-cocycle $\omega \in Z_s^2(\mathfrak{g}, \mathfrak{a})$. Let $\omega^l \in \Omega^2(G, \mathfrak{a})$ be the \mathfrak{a} -valued, G-invariant differential 2-form with $\omega^l(e) = \omega$.⁹

Definition 5.14. The period homomorphism is the map

$$\operatorname{per}_{\omega}: H_2(G) \to \mathfrak{a}, \quad [\sigma] \mapsto \int_{\sigma} \omega^l.$$

We denote its image by $\Pi_{\omega} := \operatorname{per}_{\omega}(H_2(G))$, and we have the quotient $\pi : \mathfrak{a} \to \mathfrak{a}/\Pi_{\omega}$. We set $A := \mathfrak{a}/\Pi_{\omega}$.

Neeb, inspired by van Est [Est58], gave the following construction of a locally smooth 2-cocycle integrating ω . While Neeb assumes that Π_{ω} is a topologically discrete subgroup of \mathfrak{a} , we make the less-restrictive assumption that

Assumption 5.15. Π_{ω} is a diffeologically discrete subgroup of \mathfrak{a} .

Lemma 5.16 ([Nee02, Lemmas 6.2, 6.3]). Let $u \subseteq \mathfrak{g}$ be an open convex neighbourhood of the origin, with a chart $\varphi : u \to G$ sending 0 to e and such that $T_0\varphi = \mathrm{id}_{\mathfrak{g}}$. Define, for $x \in u$,

$$\alpha_{\varphi(x)} : [0,1] \to G, \quad \alpha_{\varphi(x)}(t) := \varphi(tx).$$

Let $v \subseteq u$ be an open convex neighbourhood of the origin, such that $\varphi(v)\varphi(v) \subseteq \varphi(u)$, and define, for $x, y \in V$,

$$x * y := \varphi^{-1}(\varphi(x)\varphi(y)).$$

We use this multiplication to define the map $\gamma_{x,y}$ from the standard 2-simplex Δ^2

$$\gamma_{x,y}: \Delta^2 \to u, \quad \gamma_{x,y}(t,s) \mapsto t(x*sy) + s(x*(1-t)y),$$

and we define $\sigma_{x,y} := \varphi \circ \gamma_{x,y} : \Delta^2 \to G$.

⁹For a diffeological vector space V, and a diffeological space X, we have not defined the meaning of a V-valued differential form on X. Instead, to understand ω^l we simply remain in the setting of convenient manifolds.

(a) The function

$$f_0: v \times v \to \mathfrak{a}, \quad f_0(x, y) := \int_{\sigma_{x,y}} \omega^l$$

is smooth, and $d^2 f_0 = \frac{1}{2}\omega$. (b) The function

$$f: \varphi(v) \times \varphi(v) \to A, \quad f(\varphi(x), \varphi(y)) := \pi(f_0(x, y))$$

is a locally smooth 2-cocycle in $Z^2_{s,loc}(G,A)$, and $L(f) = \omega$.

Proof. Part (a) is simply a restatement of [Nee02, Lemma 6.2], under identifying $T_0 v \cong \mathfrak{g}$ and $T_0 \mathfrak{a} \cong \mathfrak{a}$. For (b), we begin with the fact that, since $T_0 \varphi = \mathrm{id}_{\mathfrak{g}}$,

$$L(f_0 \circ (\varphi^{-1}, \varphi^{-1})) = \omega.$$

Now, by Corollary 3.12 the map $T_0\pi$: Lie(\mathfrak{a}) $\cong \mathfrak{a} \to$ Lie(A) is an isomorphism. Therefore $L(f) = \omega$ also holds.

We now arrive at our main result for this section.

Theorem III. Suppose that $\mathfrak{a} \hookrightarrow \hat{\mathfrak{g}} \twoheadrightarrow \mathfrak{g}$ is a topologically split central extension of convenient Lie algebras. Fix $\omega \in Z^2_s(\mathfrak{g}, \mathfrak{a})$ corresponding to this extension. Assume that there is some connected and simply-connected convenient Lie group G with $\operatorname{Lie}(G) = \mathfrak{g}$. If the image Π_{ω} of the associated period homomorphism $\operatorname{per}_{\omega} : H_2(G) \to \mathfrak{a}$ is diffeologically discrete, then there is some elastic diffeological group \hat{G} centrally extending G by $\mathfrak{a}/\Pi_{\omega}$, for which



Proof. Let $A := \mathfrak{a}/\Pi_{\omega}$. Appealing to Lemma 5.16, we get $f \in Z^2_{s,\text{loc}}(G,A)$ with $L(f) = \omega$. Then, by Proposition 5.13, the central extension of G by A corresponding to f differentiates to the central extension of \mathfrak{g} by \mathfrak{a} corresponding to ω . The former gives $\hat{G} := \hat{G}_f$; the latter is $\hat{g} \cong \hat{g}_{\omega}$.

With this machinery, we integrate a class of convenient Lie algebras, as promised. Let \mathfrak{z} denote the centre of \mathfrak{g} , and set $\mathfrak{g}_{\mathrm{ad}} := \mathfrak{g}/\mathfrak{z}$. If $\mathfrak{z} \hookrightarrow \mathfrak{g} \twoheadrightarrow \mathfrak{g}_{\mathrm{ad}}$ is a split central extension of Lie algebras, it corresponds to the cocycle

$$\omega:\mathfrak{g}_{\mathrm{ad}}\times\mathfrak{g}_{\mathrm{ad}}\to\mathfrak{z},\quad (\xi,\eta)\mapsto[\xi,\eta]_{\mathfrak{z}},$$

where $[\cdot, \cdot]_{\mathfrak{z}}$ denotes the \mathfrak{z} -component of the Lie bracket. If there is, furthermore, a connected and simply-connected convenient Lie group G_{ad} with $\text{Lie}(G_{ad}) = \mathfrak{g}_{ad}$, then we have the associated period homomorphism $\text{per}_{\omega} : H_2(G_{ad}) \to \mathfrak{z}$. If this has diffeologically discrete image, then Theorem III gives G with $\text{Lie}(G) = \mathfrak{g}$.

Remark 5.17. We require the assumption that the central extension $\mathfrak{z} \hookrightarrow \mathfrak{g} \twoheadrightarrow \mathfrak{g}_{\mathrm{ad}}$ is split in order to identify the extension with a 2-cocycle in $Z_s^2(\mathfrak{g}_{\mathrm{ad}},\mathfrak{z})$. If the extension is not split, Neeb [Nee06, Section VI.1] has given a two-step process that salvages the strategy. One first integrates the central term in a split-exact sequence

$$\mathfrak{g} \hookrightarrow \hat{\mathfrak{g}} := \mathfrak{g} \oplus_b \mathfrak{g}_{\mathrm{ad}} \twoheadrightarrow |\mathfrak{g}_{\mathrm{ad}}|,$$

where $|\cdot|$ indicates the underlying vector space and $b \in Z_s^2(\mathfrak{g}, \mathfrak{g}_{ad})$ is the 2-cocycle induced by the bracket. Say this integration is \hat{G} . Then, using the fact that \mathfrak{g} is a closed ideal of $\hat{\mathfrak{g}}$, argue that the inclusion $\mathfrak{g} \hookrightarrow \hat{\mathfrak{g}}$ lifts to an inclusion $G \hookrightarrow \hat{G}$ for some G.

Thus, removing our assumption that the central extension is split amounts to understanding when, given an elastic group G and a closed ideal \mathfrak{h} of Lie(G), there is some subgroup H of G with $\text{Lie}(H) = \mathfrak{h}$.

The various pre-conditions of Theorem III, applied to $\mathfrak{z} \hookrightarrow \mathfrak{g} \twoheadrightarrow \mathfrak{g}_{ad}$, hold in the following contexts:

- If \mathfrak{g} is a Banach Lie algebra, and \mathfrak{z} is finite-dimensional, then $\mathfrak{z} \hookrightarrow \mathfrak{g} \twoheadrightarrow \mathfrak{g}_{ad}$ is split. This follows from the facts that, by the Hahn-Banach theorem, \mathfrak{z} has a closed complement, and by the open mapping theorem, the existence of this closed complement is equivalent to the existence of a continuous (hence smooth) section $\mathfrak{g}_{ad} \to \mathfrak{g}$.
- If Π_{ω} is countable, then it is diffeologically discrete.
- If \mathfrak{g}_{ad} is a "locally exponential" convenient Lie algebra, then Neeb [Nee06, Theorem IV.3.8]¹⁰ has constructed a convenient Lie group integrating it, and we may then pass to the simply-connected cover. In particular, since Banach Lie algebras are "locally exponential," when \mathfrak{g} is a Banach Lie algebra we may always find a connected and simply-connected Banach Lie group G_{ad} with $\text{Lie}(G_{ad}) = \mathfrak{g}_{ad}$.

We finish with the classic examples of non-enlargeable Banach Lie algebras, given by van Est and Korthagen [EK64] and Douady and Lazard [DL66], as retold by Neeb [Nee06].

Example 5.18 ([EK64, Section 6]). Consider the Banach-Lie algebra $C^1(S^1, \mathfrak{su}_2(\mathbb{C}))$, and the \mathbb{R} -valued cocycle

$$\omega(f,g) := \int_0^1 \operatorname{trace}(f(t)g'(t))dt$$

For the integral, we identify the continuous maps from S^1 with maps from \mathbb{R} with period 1. Let $\mathfrak{g}_{\mathrm{EK}}$ denote the central extension of $C^1(S^1, \mathfrak{su}_2(\mathbb{C}))$ by ω . One verifies that

$$(\mathfrak{g}_{\mathrm{EK}})_{\mathrm{ad}} \cong C^1(S^1, \mathfrak{su}_2(\mathbb{C})) \text{ and } G_{\mathrm{ad}} \cong C^1(S^1, SU_2(\mathbb{C})),$$

so then $\pi_2(G_{ad}) = \pi_3(SU_2(\mathbb{C})) = \pi_3(S^3) = \mathbb{Z}$. This means that Π_{ω} is at most countable. Therefore Theorem III applies, integrating $\mathfrak{g}_{\mathrm{EK}}$ to an elastic group G_{EK} .

Example 5.19 ([DL66, Section IV]). Let H be a separable infinite-dimensional complex Hilbert space. Consider the Banach-Lie algebra $\mathfrak{u}(H)$ of skew-adjoint bounded linear

¹⁰note that this citation contains no proof

operators on H. This has centre $i\mathbb{R}$. For an irrational number θ , define

$$\mathfrak{n}_{ heta} := (1_H, heta \cdot 1_H) \cdot i\mathbb{R}_{+}$$

This is a closed ideal of the centre $i\mathbb{R} \times i\mathbb{R}$ of $\mathfrak{u}(H) \times \mathfrak{u}(H)$, thus $\mathfrak{g}_{DL} := (\mathfrak{u}(H) \times \mathfrak{u}(H))/\mathfrak{n}_{\theta}$ is a Banach-Lie algebra. The central extension

$$i\mathbb{R} \cong \mathfrak{z}_{\mathrm{DL}} \hookrightarrow \mathfrak{g}_{\mathrm{DL}} \twoheadrightarrow (\mathfrak{g}_{\mathrm{DL}})_{\mathrm{ad}} \cong (\mathfrak{u}(H) \times \mathfrak{u}(H))_{\mathrm{ad}}$$

is split because \mathfrak{z} is finite-dimensional. Furthermore, by the discussion in [Nee06, Pages 415–416], the corresponding period group Π_{ω} is isomorphic to \mathbb{Z}^2 , hence is countable. Therefore, again, we may use Theorem III to integrate \mathfrak{g}_{DL} to an elastic group G_{DL} . \triangleright

Remark 5.20. In [BP24], Beltita and Pelletier recall Banach Q-manifolds¹¹ and generalize these to H-manifolds. Both are sets equipped with atlases of some kind. Beltita and Pelletier integrate every Banach-Lie algebra to an H-group, and remark that it is unclear whether Douady and Lazard's example \mathfrak{g}_{DL} integrates to a Q-group. However, careful examination of our integration, specifically Lemma 5.7, reveals that G_{DL} is locally diffeomorphic to $\mathfrak{z}_{DL}/\Pi_{\omega} \times V$, where V is an open subset of a Banach-Lie group. Since Π_{ω} is a countable subgroup of \mathfrak{z}_{DL} , the set $\mathfrak{z}_{DL}/\Pi_{\omega} \times V$, hence the set G_{DL} , also inherit the structure of a Q-group. We anticipate that this Q-group differentiates to \mathfrak{g}_{DL} under Beltita and Pelletier's Lie functor.

There are natural faithful functors from the categories of Q- and H-manifolds into \mathcal{D} flg. Knowing whether these functors are full, and having a characterization of their essential images, would be helpful in comparing Beltita and Pelletier's approach to the diffeological one.

APPENDIX A. TANGENT CATEGORIES

In this Appendix, we collect some facts about tangent categories. Specifically, Lemmas A.11 and A.16 hold in an arbitrary tangent category, so we prove them in that generality. We include the definitions and axioms of a tangent category for completeness rather than pedagogy. We direct the reader to [CC14; AB24] for a more comprehensive introduction to tangent categories.

A.1. First definitions. Let \mathcal{C} be a category, with a terminal object *. We assume that the over-category $\mathcal{C} \downarrow X$ (hence also $\mathcal{C} \cong \mathcal{C} \downarrow *$) has all finite products. We need three initial notions to define a tangent category.

Definition A.1. A bundle of abelian groups $p : A \to X$ is an abelian group object in $\mathcal{C} \downarrow X$.

The data of such a bundle consists of the projection $p : A \to X$, the addition $+ : A \times_X A \to A$, zero-section $0 : A \to A$, and inversion $i : A \to A$.

Definition A.2. Let $T: \mathcal{C} \to \mathcal{C}$ be an endofunctor, and $\tau: T^2 \to T^2$ a natural transformation. Let

$$\tau_{12} := \tau T$$
 and $\tau_{23} := T \tau$

¹¹Following Barre's [Bar73] definition of finite-dimensional Q-manifolds.

be the trivial extensions of τ to natural transformations $T^3 \to T^3$. We call τ a symmetric structure if

- $\tau_{12} \circ \tau_{23} \circ \tau_{12} = \tau_{23} \circ \tau_{12} \circ \tau_{23}$ (i.e. τ is a braiding on T), and
- $\tau \circ \tau = T^2$.

 \diamond

 \diamond

Definition A.3. We say an endofunctor $T : C \to C$ preserves fiber products of a bundle of abelian groups $p : A \to X$ if for all $k \ge 1$, the natural morphisms

$$\nu_{k,X}: T(A \times_X \cdots \times_X A) \to TA \times_{TX} \cdots \times_{TX} TA$$

are all isomorphisms.

Here is the definition of a tangent structure.

Definition A.4. The data of a *tangent structure* on C consists of an endofunctor $T : C \to C$, and natural transformations

- the projection $\pi: T \to 1$,
- the zero-section $0: 1 \to T$,
- the addition $+: T_2 \to T$,
- the vertical lift $\lambda : T \to T^2$, and
- the symmetric structure $\tau: T^2 \to T^2$.

The axioms are:

- The projection $\pi: T \to 1$ is a bundle of abelian groups over 1, with zero-section 0 and addition +.
- All pullbacks $T_k := T \times_1 \cdots \times_1 T$ exists, and are pointwise preserved by T.
- The transformation τ is a symmetric structure, and also a morphism of bundles of abelian groups.
- The diagrams

$$\begin{array}{cccc} T & \xrightarrow{\lambda} & T^2 & T & \xrightarrow{\lambda} & T^2 \\ \pi & & & \downarrow \pi T & \lambda \\ 1 & \xrightarrow{0} & T & T^2 & \xrightarrow{T\lambda} & T^3 \end{array}$$

commute, and the first is a morphism of abelian groups.

• The diagrams



commute.

• The diagram



is a pointwise pullback. We denote the top transformation by λ_2 .

Remark A.5. The last axiom, in the presence of all others, is by [CC14, Lemma 3.10], equivalent to the easier-to-write condition that

$$\begin{array}{ccc} T & \xrightarrow{\lambda} & T^2 \\ \pi & & \downarrow (\pi T, T\pi) \\ 1 & \xrightarrow{(0,0)} & T_2 \end{array}$$

is a pointwise pullback. However, it is λ_2 that will appear in the definition of the Lie bracket.

The tangent structure lets us define a Lie bracket. Fix v, w vector fields on X, i.e. sections of $\pi_X : TX \to X$. The footprint (in TX) of $Tw \circ v$ is w:

$$\pi_{TX} \circ Tw \circ v = w \circ \pi_X \circ v = w \circ \mathrm{id}_X = w$$

Using the fact $\pi_{TX} \circ \tau_X = T\pi_X$, we see the footprint of $\tau_X \circ Tv \circ w$ is w:

 $\pi_{TX} \circ \tau_X \circ Tv \circ w = T\pi_X \circ Tv \circ w = T \operatorname{id}_X \circ w = \operatorname{id}_{TX} \circ w = w.$

Thus we may subtract them in T^2X . In other words, we have a map

$$\delta(v,w): X \to T^2 X, \quad \delta(v,w) = T w \circ v - \tau_X \circ T v \circ w.$$

On the other hand, again using the chain rule, we can check that

$$T\pi_X \circ \delta(v, w) = v - v = 0_X.$$

Thus $\delta(v, w)$ takes its values in the kernel of $T\pi_X$, which, by the axioms of a tangent category, coincides with the image of $\lambda_{2,X}$.

Definition A.6. We use the notation established above. The Lie bracket of vector fields $v, w \in \mathfrak{X}(X)$ is defined as the unique vector field [v, w] such that

$$\delta(v, w) = \lambda_{2,X} \circ (w, [v, w]).$$

 \diamond

We now add more structure to our tangent category. Recall that \mathcal{C} has finite products.

Definition A.7. The tangent category (\mathcal{C}, T) is *Cartesian* if the natural morphisms

$$T(X \times Y) \to TX \times TY$$

are isomorphisms for all X, Y, and also $T * \cong *$.

 \diamond

 \diamond

Fix a ring object R in a Cartesian tangent category C. The natural transformation $1 \times R \to 1$ inherits from R the structure of a ring object in the category of endofunctors $\operatorname{End}(C)$ over 1.

Definition A.8. An $(1 \times R \to 1)$ -module structure on $(T \to 1) \in \text{End}(\mathcal{C}) \downarrow 1$ is an *R*-module structure on *T*.

Such a structure is given by a natural morphism $\kappa_X : R \times TX \to TX$, which acts as the *R*-multiplication on the fiber. We have a morphism

$$TR \to R \times R$$

If this is an isomorphism of *R*-modules, we call *R* tangent-stable.

Definition A.9. A Cartesian tangent category with scalar R multiplication consists of a Cartesian tangent category C, a commutative ring object R, and an R-module structure on T that is tangent-stable and for which some diagrams commute.

The precise definition appears in [AB24, Definition 2.28]. In a Cartesian tangent category with scalar \mathbb{R} -multiplication, the collection of vector fields $\mathfrak{X}(X)$, equipped with the Lie bracket, is a real Lie algebra.

A.2. The Cartesian structure. Fix a Cartesian tangent category C. The natural morphisms $T(X \times Y) \to TX \times TY$ are all isomorphisms, so we suppress them from notation and simply identify $T(X \times Y) \cong TX \times TY$. Given $f: X \times Y \to Z$, we have the partial tangent morphisms

$$T_{(1)}f: TX \times Y \xrightarrow{1 \times 0} TX \times TY \xrightarrow{Tf} TZ$$
$$T_{(2)}f: X \times TY \xrightarrow{0 \times 1} TX \times TY \xrightarrow{Tf} TZ.$$

Thus $T_{(1)}T_{(2)}f$ and $T_{(2)}T_{(1)}f$ give two morphisms $TX \times TY \to TZ$. They are related by the canonical flip:

$$T_{(1)}T_{(2)}f = \tau_Z \circ T_{(2)}T_{(1)}f.$$

Indeed, by naturality of τ , and its compatibility with the Cartesian structure,

$$\tau_Z \circ T_{(2)} T_{(1)} f = \tau_Z \circ T^2 f \circ (0, T0) = T^2 f \circ (\tau_Z \circ 0, \tau_Z \circ T0)$$

= $T^2 f \circ (T0, 0) = T_{(1)} T_{(2)} f.$

We have the usual expression of the total derivative as the sum of partial derivatives.

Lemma A.10 ([CC14, Proposition 2.10]). For $f: X \times Y \to Z$, we have

$$Tf = T_{(1)}f \circ (1 \times \pi) + T_{(2)}f \circ (\pi \times 1).$$

We also need the following sufficient criterion for $T_{(1)}T_{(2)}f \circ (v,w) = T_{(2)}T_{(1)}f \circ (v,w)$.

Lemma A.11. Suppose that $v : X \to TX$ and $w : Y \to TY$ are vector fields, and $a : A \to X \times Y$ is a morphism. If

$$Tf \circ (0, w) \circ a = Tf \circ (v, 0) \circ a = 0 \circ f \circ a,$$

then

$$T_{(1)}T_{(2)}f \circ (v,w) \circ a = T_{(2)}T_{(1)}f \circ (v,w) \circ a.$$

Proof. Since $\lambda = \lambda \tau$, it suffices to show that $T_{(1)}T_{(2)}f \circ (v,w) \circ a = \lambda \circ \varepsilon$ for map $\varepsilon : A \to TZ$. To do this, we use the fact that λ is a kernel. Compute

$$\pi_{TZ} \circ T_{(1)}T_{(2)}f = \pi_{TZ} \circ T(T_{(2)}f) \circ (1 \times 0)$$

= $T_{(2)}f \circ (\pi_X \times \pi_{TY}) \circ (1 \times 0)$
= $T_{(2)}f \circ (\pi_X \times 1_{TY})$
= $Tf \circ (0 \times 1) \circ (\pi_X \times 1_{TY})$
= $Tf \circ (0 \circ \pi_X, 1_{TY})$
 $T\pi_Z \circ T_{(1)}T_{(2)}f = T(\pi_Z \circ T_{(2)}f) \circ (1 \times 0)$
= $Tf \circ (1 \times T\pi_Y) \circ (1 \times 0)$
= $Tf \circ (1_{TX} \times 0 \circ \pi_Y)$

These evaluate to $Tf \circ (0, w) \circ a$ and $Tf \circ (v, 0) \circ a$ when pulled back along $(v, w) \circ a$, respectively. By assumption, these coincide with $0 \circ f \circ a$. Since λ is a kernel, there must exist $\varepsilon : a \to TZ$ with $\lambda \circ \varepsilon = T_{(1)}T_{(2)}f \circ (v, w) \circ a$. Then

$$T_{(2)}T_{(1)}f \circ (v,w) \circ a = \tau \circ T_{(1)}T_{(2)}f \circ (v,w) \circ a = \tau \circ \lambda \circ \varepsilon = \lambda \circ \varepsilon = T_{(1)}T_{(2)}f \circ (v,w) \circ a.$$

A.3. Groups in tangent categories. Let G be a group object in \mathcal{C} , with multiplication m, unit u, and inversion i. Then TG is also a group object in \mathcal{C} , with multiplication Tm, unit $Tu = 0 \circ u$, and inversion Ti.

Definition A.12. The *left-invariant* vector fields on G are those $v \in \mathfrak{X}(G)$ such that the following commutes:

$$\begin{array}{c} G \times TG \xrightarrow{T_{(2)}m} TG \\ \downarrow \times v \uparrow & v \uparrow \\ G \times G \xrightarrow{m} G. \end{array}$$

We write $\mathfrak{X}(G)^L$ for the collection left-invariant vector fields on G.

Verifying that $\mathfrak{X}(G)^L$ is a Lie subalgebra of $\mathfrak{X}(G)$ takes significant work. It is a consequence of [AB24, Theorem 6.6], which applies more generally to groupoid objects.

Lemma A.13. Let G be a group object in a Cartesian tangent category C with scalar R-multiplication. The set $\mathfrak{X}(G)^L$ is a Lie subalgebra of $\mathfrak{X}(G)$.

 \diamond

Let \mathfrak{g} denote the pullback of TG along u. We have the left trivialization

$$\varphi_G: G \times \mathfrak{g} \hookrightarrow G \times TG \xrightarrow{T_{(2)}m} TG,$$

and its inverse, denoted ψ_G , determined by

Lemma A.14. The $\underline{C}(*, R)$ -module of sections $\Gamma(*, \mathfrak{g})$ is isomorphic to the $\underline{C}(*, R)$ module $\mathfrak{X}(G)^L$. The isomorphism and its inverse are given by

$$\xi \mapsto \varphi_G \circ (1 \times \xi), \quad v \mapsto v \circ u.$$

Remark A.15. If G is an elastic diffeological group, then $\mathfrak{g} = T_e G$ carries the subspace diffeology. Furthermore, since \mathcal{D} flg is locally Cartesian closed, the space of left-invariant vector fields $\mathfrak{X}(G)^L$ inherits the subspace diffeology of the functional diffeology on $\mathfrak{X}(G)$. The isomorphism of \mathbb{R} -modules in Lemma A.14 is also an isomorphism in \mathcal{D} flg.

We let c denote the conjugation

$$c: G \times G \to G, \quad c:= m \circ (m, i \circ \operatorname{pr}_1).$$

Lemma A.16. Let $v, w \in \mathfrak{X}(G)^L$. Then

$$T_{(1)}T_{(2)}c \circ (v,w) \circ u = \delta(v,w) \circ u.$$

This amounts to the classic formulation of the Lie bracket as the second derivative of the conjugation.

Proof. We break the computation into parts. First, by expanding definitions and using associativity of arrow composition,

$$T_{(2)}c = Tm \circ (T_{(2)}m, \ T_{(2)}(i \circ \mathrm{pr}_1))$$

= $Tm \circ (T_{(2)}m, \ 0 \circ i \circ \mathrm{pr}_1)$ using $T_{(2)}(i \circ \mathrm{pr}_1) = Ti \circ 0 \circ \mathrm{pr}_1$
and $Ti \circ 0 = 0 \circ i$
= $T_{(1)}m \circ (T_{(2)}m, i \circ \mathrm{pr}_1).$

Similarly,

$$T_{(1)}T_{(2)}c = TT_{(1)}m \circ \left(T_{(1)}T_{(2)}m, \ T_{(1)}(i \circ \mathrm{pr}_1)\right)$$
$$= TT_{(1)}m \circ \left(T_{(1)}T_{(2)}m, \ T(i \circ 0) \circ \mathrm{pr}_1\right).$$

Using A.10, we decompose the right side into a sum

$$T_{(1)}T_{(2)}c = T_{(1)}T_{(1)}m \circ \left(T_{(1)}T_{(2)}m, \ i \circ 0 \circ \pi \circ \mathrm{pr}_1\right) + T_{(2)}T_{(1)}m \circ \left(\pi \circ T_{(1)}T_{(2)}m, \ T(i \circ 0) \circ \mathrm{pr}_1\right)$$

Now we pre-compose with $(v, w) \circ u$. The most substantive term is

(A.17)

$$T_{(1)}T_{(2)}m \circ (v,w) \circ u = TT_{(2)}m \circ (v,0 \circ w) \circ u$$

$$= TT_{(2)}m \circ (1 \times Tw) \circ (v,0) \circ u$$

$$= Tw \circ Tm \circ (v \circ u, 0 \circ u)$$

(A.18) $= Tw \circ v \circ u.$

For (A.17), apply T to the diagram expressing left-invariance of w; for (A.18), use the fact that $0 \circ u$ is the unit of TG, and Tm is the multiplication. Therefore

$$T_{(1)}T_{(2)}c \circ (v,w) \circ u = T_{(1)}T_{(1)}m \circ (Tw \circ v \circ u,u) + T_{(2)}T_{(1)}m \circ (w \circ u, Ti \circ v \circ u).$$

The first summand expands to

$$T_{(1)}T_{(1)}m \circ (Tw \circ v \circ u, u) = T^2m(Tw \circ v \circ u, 0 \circ 0 \circ u),$$

and $0 \circ 0 \circ u$ is the group unit in T^2G , we get $Tw \circ v \circ u$. For the second summand, recall that $\tau \circ T_2T_{(1)} = T_{(1)}T_{(2)}$. We also have

$$Ti \circ v \circ u = -v \circ u_{i}$$

because Ti is the group inversion on TG, and $Tm \circ (a \circ u, b \circ u) = a \circ u + b \circ u$. Putting these together, the second summand is

$$T_{(2)}T_{(1)}m \circ (w \circ u, Ti \circ v \circ u) = -\tau \circ Tv \circ w \circ u.$$

We conclude that

$$T_{(1)}T_{(2)}c\circ(v,w)\circ u=\delta(v,w)\circ u,$$

as desired.

References

- [AB24] Lory Aintablian and Christian Blohmann. Differentiable groupoid objects and their abstract Lie algebroids. 2024. arXiv: 2412.19697 [math.CT] (cit. on pp. 2, 4, 11, 12, 35, 38, 39).
- [BH11] John C. Baez and Alexander E. Hoffnung. Convenient categories of smooth spaces. Trans. Amer. Math. Soc. 363 (2011), no. 11, pp. 5789–5825. DOI: 10.1090/S0002-9947-2011-05107-X (cit. on p. 5).
- [Bar73] Raymond Barre. De quelques aspects de la théorie des q-variétés différentielles et analytiques. Ann. Inst. Fourier (Grenoble) 23 (1973), no. 3, pp. 227–312 (cit. on p. 35).
- [BKW23] Augustin Batubenge, Yael Karshon, and Jordan Watts. Diffeological, Frölicher, and Differential Spaces. 2023. arXiv: 1712.04576 [math.DG] (cit. on pp. 17, 18).
- [BP24] Daniel Beltiță and Fernand Pelletier. Smooth Banach structure on orbit spaces and leaf spaces. To appear in Dissertationes Mathematicae. 2024. arXiv: 230 5.16167 [math.DG] (cit. on pp. 4, 35).

- [Blo24a] Christian Blohmann. Elastic diffeological spaces. In: Recent advances in diffeologies and their applications. Vol. 794. Contemp. Math. Amer. Math. Soc.,
 [Providence], RI, 2024, pp. 49–86. DOI: 10.1090/conm/794/15925 (cit. on pp. 2, 4, 7–11).
- [Blo24b] Christian Blohmann. Lagrangian field theory. Online lecture notes. 2024 (cit. on pp. 5, 7).
- [Blo25] Christian Blohmann. Tangent structure and Cartan calculus on elastic diffeological spaces. In preparation. 2025 (cit. on pp. 2, 10).
- [Bom67] Jan Boman. Differentiability of a function and of its compositions with functions of one variable. Math. Scand. 20 (1967), pp. 249–268. DOI: 10.7146/ma th.scand.a-10835 (cit. on p. 17).
- [Car52] Elie Cartan. Le troisième théorème fondamental de Lie. In: Oeuvres complètes. Part I. Groupes de Lie. Gauthier-Villars, Paris, 1952, pp. 1143–1148 (cit. on p. 4).
- [Che77] Kuo Tsai Chen. *Iterated path integrals*. Bull. Amer. Math. Soc. **83** (1977), no. 5, pp. 831–879. DOI: 10.1090/S0002-9904-1977-14320-6 (cit. on p. 2).
- [CSW14] J. Daniel Christensen, Gordon Sinnamon, and Enxin Wu. The D-topology for diffeological spaces. Pacific J. Math. 272 (2014), no. 1, pp. 87–110. DOI: 10.2140/pjm.2014.272.87 (cit. on p. 24).
- [CW16] J. Daniel Christensen and Enxin Wu. Tangent spaces and tangent bundles for diffeological spaces. Cah. Topol. Géom. Différ. Catég. 57 (2016), no. 1, pp. 3– 50 (cit. on p. 4).
- [CC14] J. Robin B. Cockett and Geoffrey S. H. Cruttwell. *Differential structure, tan*gent structure, and SDG. Appl. Categ. Structures 22 (2014), no. 2, pp. 331– 417. DOI: 10.1007/s10485-013-9312-0 (cit. on pp. 4, 35, 37, 38).
- [CC15] J. Robin B. Cockett and Geoffrey S. H. Cruttwell. The Jacobi identity for tangent categories. Cah. Topol. Géom. Différ. Catég. 56 (2015), no. 4, pp. 301– 316 (cit. on p. 4).
- [DL66] Adrien Douady and Michel Lazard. Espaces fibrés en algèbres de Lie et en groupes. Invent. Math. 1 (1966), pp. 133–151. DOI: 10.1007/BF01389725 (cit. on pp. 3, 34).
- [DN07] Brett. Dugmore and Patrice P. Ntumba. On tangent cones of Frölicher spaces. Quaest. Math. **30** (2007), no. 1, pp. 67–83. DOI: 10.2989/160736007780205747 (cit. on p. 4).
- [Est58] Willem T. van Est. A group theoretic interpretation of area in the elementary geometries. Simon Stevin **32** (1958), pp. 29–38 (cit. on p. **32**).
- [EK64] Willem T. van Est and Theodorus J. Korthagen. Non-enlargible Lie algebras. Indag. Math. 26 (1964). Nederl. Akad. Wetensch. Proc. Ser. A 67, pp. 15–31 (cit. on pp. 2–4, 34).
- [Frö81] Alfred Frölicher. Applications lisses entre espaces et variétés de Fréchet. C.
 R. Acad. Sci. Paris Sér. I Math. 293 (1981), no. 2, pp. 125–127 (cit. on p. 24).

- [FK88] Alfred Frölicher and Andreas Kriegl. Linear spaces and differentiation theory.
 Pure and Applied Mathematics (New York). A Wiley-Interscience Publication.
 John Wiley & Sons, Ltd., Chichester, 1988 (cit. on pp. 4, 17–20).
- [Glö06] Helge Glöckner. Discontinuous non-linear mappings on locally convex direct limits. Publ. Math. Debrecen **68** (2006), no. 1-2, pp. 1–13. DOI: 10.5486/pmd .2006.2714 (cit. on p. 24).
- [GW21] Nico Goldammer and Kathrin Welker. Towards optimization techniques on diffeological spaces by generalizing Riemannian concepts. 2021. arXiv: 2009.0 4262 [math.OC] (cit. on p. 4).
- [Hai79] Richard M. Hain. A characterization of smooth functions defined on a Banach space. Proc. Amer. Math. Soc. **77** (1979), no. 1, pp. 63–67. DOI: 10.2307/204 2717 (cit. on p. 4).
- [Hec95] Gilbert Hector. Géométrie et topologie des espaces difféologiques. In: Analysis and geometry in foliated manifolds (Santiago de Compostela, 1994). World Sci. Publ., River Edge, NJ, 1995, pp. 55–80 (cit. on p. 4).
- [IKZ10] Patrick Iglesias, Yael Karshon, and Moshe Zadka. Orbifolds as diffeologies. Trans. Amer. Math. Soc. 362 (2010), no. 6, pp. 2811–2831. DOI: 10.1090/S00 02-9947-10-05006-3 (cit. on p. 14).
- [Igl13] Patrick Iglesias-Zemmour. *Diffeology*. Vol. 185. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2013. DOI: 10 .1090/surv/185 (cit. on pp. 4, 5, 15, 25).
- [KM97] Andreas Kriegl and Peter W. Michor. The convenient setting of global analysis. Vol. 53. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1997, pp. x+618. DOI: 10.1090/surv/053 (cit. on pp. 3, 4, 17, 25).
- [Les03] Joshua Leslie. On a diffeological group realization of certain generalized symmetrizable Kac-Moody Lie algebras. J. Lie Theory 13 (2003), no. 2, pp. 427– 442 (cit. on p. 4).
- [Los92] Mark V. Losik. Fréchet manifolds as diffeological spaces. Izv. Vyssh. Uchebn. Zaved. Mat. (1992), no. 5, pp. 36–42 (cit. on p. 4).
- [Mag18] Jean-Pierre Magnot. The group of diffeomorphisms of a non-compact manifold is not regular. Demonstr. Math. **51** (2018), no. 1, pp. 8–16. DOI: 10.1515/de ma-2018-0001 (cit. on p. 4).
- [Nee02] Karl-Hermann Neeb. Central extensions of infinite-dimensional Lie groups. Ann. Inst. Fourier (Grenoble) 52 (2002), no. 5, pp. 1365–1442. DOI: 10.5802 /aif.1921 (cit. on pp. 3, 4, 32, 33).
- [Nee06] Karl-Hermann Neeb. Towards a Lie theory of locally convex groups. Jpn. J. Math. 1 (2006), no. 2, pp. 291–468. DOI: 10.1007/s11537-006-0606-y (cit. on pp. 34, 35).
- [Ros84] Jiří Rosický. Abstract tangent functors. Diagrammes 12 (1984), JR1–JR11 (cit. on pp. 2, 4).
- [Sou80] Jean-Marie Souriau. Groupes différentiels. In: Differential geometrical methods in mathematical physics (Proc. Conf., Aix-en-Provence/Salamanca, 1979).

Vol. 836. Lecture Notes in Math. Springer, Berlin-New York, 1980, pp 91–128 (cit. on pp. 2, 4).

- [Sta13] Andrew Stacey. Yet More Smooth Mapping Spaces and Their Smoothly Local Properties. 2013. arXiv: 1301.5493 [math.DG] (cit. on p. 4).
- [Tah24] Masaki Taho. Tangent spaces of diffeological spaces and their variants. 2024. arXiv: 2406.04703 [math.AT] (cit. on p. 4).
- [Tor98] Carlos A. Torre. A tangent bundle on diffeological spaces. 1998. arXiv: math /9801046 [math.DG] (cit. on p. 4).
- [Vin08] Martin Vincent. Diffeological differential geometry. MSc. Copenhagen, DK: University of Copenhagen, 2008 (cit. on p. 4).
- [Woc11] Christoph Wockel. Categorified central extensions, étale Lie 2-groups and Lie's third theorem for locally exponential Lie algebras. Adv. Math. **228** (2011), no. 4, pp. 2218–2257. DOI: 10.1016/j.aim.2011.07.003 (cit. on pp. 4, 32).
- [WZ16] Christoph Wockel and Chenchang Zhu. Integrating central extensions of Lie algebras via Lie 2-groups. J. Eur. Math. Soc. (JEMS) 18 (2016), no. 6, pp. 1273– 1320. DOI: 10.4171/JEMS/613 (cit. on p. 4).
- [Yam50] Hidehiko Yamabe. On an arcwise connected subgroup of a Lie group. Osaka Math. J. 2 (1950), pp. 13–14 (cit. on p. 17).

D. Miyamoto, Max Planck Institute for Mathematics, Vivatsgasse 7, 53111 Bonn, Germany

Email address: miyamoto@mpim-bonn.mpg.de