

EXPLICIT FORMULAS FOR THE ALEXANDER POLYNOMIAL OF PRETZEL KNOTS

Y. BELOUSOV

ABSTRACT. We provide explicit formulas for the Alexander polynomial of Pretzel knots and establish several immediate corollaries, including the characterization of Pretzel knots with a trivial Alexander polynomial.

INTRODUCTION

A *Pretzel link* $P(q_1, q_2, \dots, q_n)$ is a link obtained by connecting n twisted bands, where each band has q_i half-twists. The sign of q_i determines the direction of the i th twist region (see Fig. 1 and Fig. 3 for examples). A Pretzel link is a knot if and only if either precisely one q_i is even, or all q_i are odd and n is odd.

Pretzel links were introduced by Reidemeister in [Rei32] and have been thoroughly studied since then. The Alexander polynomial of these knots has also been studied before — see, for example, [Nak87], [Lec15], [MMS15], [BL20] — and explicit formulas for some special cases were found. Kim and Lee [KL07] also found explicit formulas for the Conway polynomial for several classes of Pretzel links with at least one even parameter.¹

Nevertheless, general formulas for the Alexander polynomial of Pretzel knots have not appeared in the literature. In this note, we fill this gap (see Theorem 1).




FIGURE 1. Examples of Pretzel knots

In Section 1, we provide an elementary proof of the explicit formulas, relying on the skein relation. In Section 2, we derive some straightforward corollaries, including the characterization of Pretzel knots with a trivial Alexander polynomial and a formula for the knot determinant.

¹The case where all parameters are odd is also discussed in the paper, although the authors do not provide an explicit formula.

Conventions and notation.

- If $K = P(q_1, q_2, \dots, q_n)$ is a Pretzel knot with exactly one even q_i , we may assume, without loss of generality, that q_1 is even.
- We denote the Alexander polynomial of a link L by $\Delta_L(t)$.
- We write $f(t) \doteq g(t)$ to indicate that the Laurent polynomials $f(t)$ and $g(t)$ are equal up to multiplication by $\pm t^k$ for some $k \in \mathbb{Z}$.
- We denote by $\sigma_k(q_1, q_2, \dots, q_n)$ the elementary symmetric polynomial of degree k , that is, the sum of all products of k distinct variables among q_1, q_2, \dots, q_n . For $k < 0$, we set $\sigma_k(q_1, q_2, \dots, q_n) = 0$.

Theorem 1. *Let $K = P(q_1, q_2, \dots, q_n)$ be a Pretzel knot. Then the Alexander polynomial $\Delta_K(t)$ is given by the following cases:*

- (1) *If q_1 is even and n and all other q_i are odd,*

$$\Delta_K(t) \doteq \left(\prod_{i=2}^n \frac{1+t^{q_i}}{1+t} \right) \left(1 + \frac{q_1}{2}(t^{-1} - t) \left(\sum_{i=2}^n \frac{t^{q_i}}{1+t^{q_i}} - \frac{n-1}{2} \right) \right).$$

- (2) *If q_1 and n are even, while all other q_i are odd,*

$$\Delta_K(t) \doteq \left(\prod_{i=2}^n \frac{1+t^{q_i}}{1+t} \right) \left(t^{q_1/2} + (t^{q_1/2} - t^{-q_1/2}) \left(\sum_{i=2}^n \frac{t^{q_i}}{1+t^{q_i}} - \frac{n}{2} \right) \right).$$

- (3) *If both n and all q_i are odd,*

$$\Delta_K(t) \doteq \frac{1}{2^{n-1}} \sum_{k=0}^{(n-1)/2} (t-1)^{2k} (t+1)^{n-1-2k} \sigma_{2k}(q_1, q_2, \dots, q_n).$$

1. PROOF OF THEOREM 1

Recall that the *symmetrized Alexander polynomial* of a link L is a well-defined normalization of the Alexander polynomial, given as a Laurent polynomial in $t^{1/2}$ and characterized by the following two properties: (1) $\Delta_L(t) = \Delta_L(t^{-1})$, and (2) if L is a knot, then $\Delta_L(1) = 1$. The symmetrized Alexander polynomial satisfies the skein relation:

$$\Delta_{L_+}(t) - \Delta_{L_-}(t) = (t^{1/2} - t^{-1/2})\Delta_{L_0}(t),$$

where links L_+ , L_- , and L_0 differ by the local operation shown in Fig. 2.




FIGURE 2. Illustration of the skein relation

To apply the skein relation, we also need explicit formulas for a special case of two-component Pretzel links. The following lemma will be proved simultaneously with Theorem 1.




FIGURE 3. Example of two-component Pretzel link $P(3,3,3,3)$ with different orientations

Lemma 1. *Let $L = P(q_1, q_2, \dots, q_n)$ be an oriented two-component Pretzel link, where n is even and all q_i are odd. Then the Alexander polynomial $\Delta_L(t)$ is given by the following cases:*

- (1) *If the strands in each twist of L are co-directed (as in Fig. 3(A)), then*

$$\Delta_L(t) \doteq \frac{1}{(1+t)^{n-1}} \prod_{i=1}^n (1+t^{q_i}) \left(\sum_{i=1}^n \frac{t^{q_i}}{1+t^{q_i}} - \frac{n}{2} \right).$$

- (2) *If the strands in each twist of L are oppositely directed (as in Fig. 3(B)), then*

$$\Delta_L(t) \doteq \frac{1}{2^{n-1}} \sum_{k=1}^{n/2} (t-1)^{2k-1} (t+1)^{n-2k} \sigma_{2k-1}(q_1, q_2, \dots, q_n).$$

1.1. Case (3) of Theorem 1 and Case (2) of Lemma 1. Let $L = P(q_1, q_2, \dots, q_n)$ be a Pretzel link (or a knot) where all q_i are odd. We will use the following notation:

$$(1) \quad f_o(t; q_1, \dots, q_n) = \frac{t^{-\frac{n-1}{2}}}{2^{n-1}} \sum_{k=0}^{(n-1)/2} (t-1)^{2k} (t+1)^{n-1-2k} \sigma_{2k}(q_1, \dots, q_n),$$

$$(2) \quad f_e(t; q_1, \dots, q_n) = \frac{t^{-\frac{n-1}{2}}}{2^{n-1}} \sum_{k=1}^{n/2} (t-1)^{2k-1} (t+1)^{n-2k} \sigma_{2k-1}(q_1, \dots, q_n).$$

First, observe that

$$\begin{aligned} f_o(t; 1, -1, q_1, \dots, q_n) &= \frac{t^{-\frac{n+1}{2}}}{2^{n+1}} \sum_{k=0}^{(n+1)/2} (t-1)^{2k} (t+1)^{n+1-2k} \underbrace{\sigma_{2k}(1, -1, q_1, \dots, q_n)}_{=\sigma_{2k}(q_1, \dots, q_n) - \sigma_{2k-2}(q_1, \dots, q_n)} = \\ &= \frac{t^{-\frac{n+1}{2}}}{2^{n+1}} \sum_{k=0}^{(n-1)/2} \underbrace{\left((t-1)^{2k} (t+1)^{n+1-2k} - (t-1)^{2k+2} (t+1)^{n-1-2k} \right)}_{4t(t-1)^{2k} (t+1)^{n-1-2k}} \sigma_{2k}(q_1, \dots, q_n) = \\ &= f_o(t; q_1, q_2, \dots, q_n). \end{aligned}$$

A similar argument holds for $f_e(t; q_1, \dots, q_n)$.

If all $|q_i| = 1$ and $m = |\sum_{i=1}^n q_i|$, then L is $(2, m)$ -torus link (or knot), and

$$\begin{aligned} f_o(t; q_1, \dots, q_n) &= \frac{t^{-\frac{m-1}{2}}}{2^{m-1}} \sum_{k=0}^{(m-1)/2} (t-1)^{2k} (t+1)^{m-1-2k} \underbrace{\sigma_{2k}(1, \dots, 1)}_{=\binom{m}{2k}} = \\ &= t^{-\frac{m-1}{2}} \sum_{k=0}^{m-1} (-t)^k = t^{-\frac{m-1}{2}} \frac{t^m + 1}{t + 1}. \end{aligned}$$

By analogous calculation

$$f_e(t; q_1, \dots, q_n) = t^{-\frac{m-1}{2}} \frac{t^m - 1}{t + 1}.$$

These are well-known formulas, confirming our induction base.

Now, assume that the induction hypothesis holds for $K_1 = P(q_1, q_2, \dots, q_n)$ and for $K_0 = P(q_2, \dots, q_n)$. We can apply the skein relation to compute $\Delta_K(t)$ for $K = P(q_1 \pm 2, q_2, \dots, q_n)$. We consider the case $q_1 + 2$, the argument for $q_1 - 2$ is analogous. Applying the induction hypothesis, we obtain the following for odd n :

$$\Delta_{P(q_1+2, q_2, \dots, q_n)}(t) = f_o(t; q_1, q_2, \dots, q_n) + (t^{1/2} - t^{-1/2}) f_e(t; q_2, \dots, q_n).$$

Similarly, for even n we have:

$$\Delta_{P(q_1+2, q_2, \dots, q_n)}(t) = f_e(t; q_1, q_2, \dots, q_n) + (t^{1/2} - t^{-1/2}) f_o(t; q_2, \dots, q_n),$$

Since

$$\sigma_k(q_1 + 2, q_2, \dots, q_n) = \sigma_k(q_1, q_2, \dots, q_n) + 2\sigma_{k-1}(q_2, \dots, q_n)$$

this confirms that f_o and f_e satisfy the desired recursion, completing the proof.

1.2. Cases (1) and (2) of Theorem 1 and Case (1) of Lemma 1. The proofs of these cases share a similar approach, so we present only the key steps, omitting direct computations.

The base case of the induction for Case (1) of Lemma 1 corresponds to the link $P(\pm 1, \pm 1, \dots, \pm 1)$, which is, once again, a two-component $(2, m)$ -torus link, though with a different orientation. The base case of the induction for Cases (1) and (2) of Theorem 1 is $P(0, q_2, \dots, q_n)$, which is a connected sum of $(2, q_i)$ -torus knots for each q_i , regardless of the parity of n . Recall that the Alexander polynomial of a connected sum is the product of the Alexander polynomials of its summands.

Finally, we use the skein relation to confirm the validity of the given formulas.

2. COROLLARIES

2.1. Determinant of a Pretzel Knot. The determinant of a Pretzel knot follows directly from Theorem 1. This result reproduces the formula obtained by Kolay in [Kol23].

Corollary 1. *The determinant of the Pretzel knot $K = P(q_1, q_2, \dots, q_n)$ is given by:*

$$\det(K) = |\sigma_{n-1}(q_1, q_2, \dots, q_n)| = \left| q_1 q_2 \dots q_n \left(\frac{1}{q_1} + \frac{1}{q_2} + \dots + \frac{1}{q_n} \right) \right|.$$

Proof. Recall that $\det(K) = |\Delta_K(-1)|$. Thus, to calculate the determinant of a Pretzel knot, we need to consider the three cases in Theorem 1.

In Case (3), the result follows by direct substitution. Cases (1) and (2) involve routine calculus computations. We perform the computation for Case (1); the argument for Case (2) is completely analogous.

The formula for Case (1) in Theorem 1 consists of two factors. For the first one, we have

$$\lim_{t \rightarrow -1} \prod_{i=2}^n \frac{1+t^{q_i}}{1+t} = \prod_{i=2}^n q_i.$$

For the second factor, using the expansion at $t = -1$, we obtain the following limits:

$$\begin{aligned} & \lim_{t \rightarrow -1} \left(1 + \frac{q_1}{2} (t^{-1} - t) \left(\sum_{i=2}^n \frac{t^{q_i}}{1+t^{q_i}} - \frac{n-1}{2} \right) \right) = \\ & = 1 + \frac{q_1}{2} \lim_{\varepsilon \rightarrow 0} \left(\underbrace{\left(\frac{1}{-1+\varepsilon} + 1 - \varepsilon \right)}_{=-2\varepsilon + O(\varepsilon^2)} \left(\sum_{i=2}^n \underbrace{\left(\frac{(-1+\varepsilon)^{q_i}}{1+(-1+\varepsilon)^{q_i}} - \frac{1}{2} \right)}_{=\frac{1}{2} - \frac{1}{q_i\varepsilon} + O(\varepsilon)} \right) \right) = \\ & = 1 + q_1 \left(\sum_{i=2}^n \frac{1}{q_i} \right) = q_1 \left(\sum_{i=1}^n \frac{1}{q_i} \right). \end{aligned}$$

Combining these results, we obtain $\sigma_{n-1}(q_1, q_2, \dots, q_n)$, as stated. \square

2.2. Degree of the Alexander Polynomial of Pretzel Knots. For Pretzel knots with at least one even parameter, the genus was completely determined in [KL07] using the formulas for the Conway polynomials. The case where all parameters are odd is more subtle (except for alternating knots; see Corollary 2). In this case, the degree of the Alexander polynomial can take any even value from 0 to $n-1$, and its dependence on the parameters is not straightforward.

In this section, we examine two cases: when the Alexander polynomial of such knots has a maximal degree ($n-1$) and when it is trivial. Before analyzing the general case, we first examine the special case of alternating Pretzel knots, where the degree of the Alexander polynomial directly determines the genus.

Corollary 2. *Let $K = P(q_1, q_2, \dots, q_n)$ be a Pretzel knot with all q_i odd and of the same sign. Then its genus satisfies $g(K) = \frac{n-1}{2}$.*

Proof. If all q_i have the same sign:

- (1) The knot is alternating, so $g(K) = \frac{\deg \Delta_K(t)}{2}$. This follows from a classical result of Murasugi [Mur58] and Crowell [Cro59].
- (2) Since $\sigma_{2k}(q_1, q_2, \dots, q_n)$ is positive for all $0 \leq k \leq \frac{n-1}{2}$, it follows that $\deg \Delta_K(t) = n-1$.

\square

Corollary 3. *Let $K = P(q_1, q_2, \dots, q_n)$ be a Pretzel knot where all q_i are odd. Then the following hold:*

(1) The Alexander polynomial satisfies $\deg \Delta_K(t) = n - 1$ if and only if

$$\sum_{k=0}^{(n-1)/2} \sigma_{2k}(q_1, q_2, \dots, q_n) \neq 0.$$

(2) The Alexander polynomial is trivial, i.e., $\Delta_K(t) \doteq 1$, if and only if for each $0 \leq k \leq \frac{n-1}{2}$,

$$\sigma_{2k}(q_1, q_2, \dots, q_n) = (-1)^k \binom{\frac{n-1}{2}}{k}.$$

Proof. To simplify the notation, let us denote $a_k = \sigma_k(q_1, q_2, \dots, q_n)$.

Consider the Alexander polynomial in the form

$$\Delta_K(t) = \frac{1}{2^{n-1}} \sum_{k=0}^{(n-1)/2} (t-1)^{2k} (t+1)^{n-1-2k} a_{2k}.$$

Observe that the coefficient of $a_{2i}t^j$ in $\Delta_K(t)$ equals the coefficient of $a_{2i}t^{n-1-j}$, where $0 \leq i \leq \frac{n-1}{2}$ and $0 \leq j \leq n-1$. Thus, $\deg \Delta_K(t) = n-1$ if the coefficient of t^0 is nonzero. Since $\Delta_K(1) = \sum_{k=0}^{(n-1)/2} a_{2k}$, this completes the proof of the first case.

The condition $\Delta_K(t) \doteq 1$ holds if and only if its only nonzero coefficient appears at $t^{(n-1)/2}$ and equals 1. This requirement can be expressed as a system of linear equations:

$$(3) \quad A \left(a_0, a_2, \dots, a_{\frac{n-1}{2}} \right)^\top = (0, 0, \dots, 0, 1)^\top,$$

where $A \in \mathbb{R}^{\frac{n+1}{2} \times \frac{n+1}{2}}$ is a matrix whose element $A_{i,j}$ is equal to the coefficient of $a_{2i-2}t^{j-1}$ in $\Delta_K(t)$.

The matrix A is non-degenerate. Indeed, if we consider $u = t - 1$, then

$$(t-1)^{2k} (t+1)^{n-1-2k} = u^{2k} (u+2)^{n-1-2k},$$

and such polynomials are clearly linearly independent. Thus, there exists a unique solution to the system (3).

Setting

$$a_{2k} = (-1)^k \binom{\frac{n-1}{2}}{k},$$

we compute:

$$\begin{aligned} \frac{1}{2^{n-1}} \sum_{k=0}^{(n-1)/2} (-1)^k \binom{\frac{n-1}{2}}{k} (t-1)^{2k} (t+1)^{n-1-2k} &= \frac{1}{2^{n-1}} \left((t+1)^2 - (t-1)^2 \right)^{\frac{n-1}{2}} = \\ &= t^{\frac{n-1}{2}}. \end{aligned}$$

□

Remark 1. The Alexander polynomial of a Pretzel knot with an even parameter is never trivial. Hence, Case (2) of Corollary 3 provides a complete characterization of all Pretzel knots with a trivial Alexander polynomial.

Remark 2. For $n = 3$, the condition in Case (2) of Corollary 3 reduces to the well-known equation

$$q_1q_2 + q_1q_3 + q_2q_3 = -1,$$

which has many known solutions. For example, for each odd number p , the triple $(-p, 2p - 1, 2p + 1)$ satisfies the equation.

If odd numbers (q_1, q_2, \dots, q_n) satisfy the equations in Case (2) of Corollary 3, then $(\pm 1, \mp 1, q_1, q_2, \dots, q_n)$ yields a solution in $n + 2$ variables. This is obvious from a geometric point of view — it is nothing but an application of the second Reidemeister move. Thus, it would be interesting to search for non-trivial solutions, i.e., those where no pairs of variables are equal to -1 and $+1$. At present, we know only one non-trivial solution:

$$(-13, 15, 17, -29, -71),$$

which, according to Bar-Natan's Θ -invariant [BNvdV], leads to a non-trivial knot.

REFERENCES

- [BL20] Y. Bae and I. S. Lee. On Alexander polynomials of Pretzel links. *Kyungpook mathematical journal*, 60(2):239–253, June 2020.
- [BNvdV] D. Bar-Natan and R. van der Veen. A very fast, very strong, topologically meaningful and fun knot invariant. Unpublished preprint. Available at <https://drorbn.net/AcademicPensieve/Projects/Theta/>.
- [Cro59] R. Crowell. Genus of alternating link types. *Annals of Mathematics*, 69(2):258–275, 1959.
- [KL07] D. Kim and J. Lee. Some invariants of Pretzel links. *Bulletin of the Australian Mathematical Society*, 75(2):253–271, April 2007.
- [Kol23] S. Kolay. Knot colorings: Coloring and Goeritz matrices. *The American Mathematical Monthly*, 130(4):335–354, April 2023.
- [Lec15] A. Lecuona. On the slice-ribbon conjecture for Pretzel knots. *Algebraic & Geometric Topology*, 15(4):2133–2173, 2015.
- [MMS15] A. Mironov, A. Morozov, and A. Sleptsov. Colored HOMFLY polynomials for the Pretzel knots and links. *Journal of High Energy Physics*, 2015(7):1–35, 2015.
- [Mur58] K. Murasugi. On the genus of the alternating knot, i. *Journal of the Mathematical Society of Japan*, 10(1):94–105, 1958.
- [Nak87] Y. Nakagawa. On the Alexander polynomials of Pretzel links $L(p_1, \dots, p_n)$. *Kobe journal of mathematics*, 3:167–177, 1987.
- [Rei32] K. Reidemeister. *Knotentheorie*. Springer-Verlag, 1932.