

A TATE ALGEBRA VERSION OF THE JACOBIAN CONJECTURE

LUCAS HAMADA, KAZUKI KATO, AND RYO KOMIYA

ABSTRACT. This paper investigates a Tate algebra version of the Jacobian conjecture, referred to as *the Tate-Jacobian conjecture*, for commutative rings R equipped with an I -adic topology. We show that if the I -adic topology on R is Hausdorff and R/I is a subring of a \mathbb{Q} -algebra, then the Tate-Jacobian conjecture is equivalent to the Jacobian conjecture. Conversely, if R/I has positive characteristic, the Tate-Jacobian conjecture fails. Furthermore, we establish that the Jacobian conjecture for \mathbb{C} is equivalent to the following statement: for all but finitely many primes p , the inverse of a polynomial map over \mathbb{C}_p whose Jacobian determinant is an element of \mathbb{C}_p^\times lies in the Tate algebra over \mathbb{C}_p .

INTRODUCTION

The **Jacobian conjecture** (JC), first proposed by O.-H. Keller in 1939 [6] and later included among Smale's problems, concerns the invertibility of polynomial maps over a commutative ring R under the assumption that their Jacobian determinant is a unit in the polynomial ring. Given a polynomial map $F = (f_1, \dots, f_n) \in R[X_1, \dots, X_n]^n$, its Jacobian matrix is defined by

$$JF = \left(\frac{\partial f_i}{\partial X_j} \right)_{1 \leq i, j \leq n}.$$

The conjecture, referred to as $\text{JC}(R, n)$, asserts the following:

$\text{JC}(R, n)$ If $F \in R[X_1, \dots, X_n]^n$ satisfies $\det JF \in R[X_1, \dots, X_n]^\times$, then there exists $G \in R[X_1, \dots, X_n]^n$ such that $F \circ G = G \circ F = (X_1, \dots, X_n)$.

In this paper, $\text{JC}(R)$ refers to the conjunction of $\text{JC}(R, n)$ for all $n \geq 1$. The existing results on the Jacobian conjecture depend on the characteristic of the ring R . Here, R is said to have characteristic c , where c is a nonnegative integer, if the kernel of the unique homomorphism $\mathbb{Z} \rightarrow R$ is the ideal $c\mathbb{Z}$.

- *Characteristic zero case:* E. Connell and L. van den Dries [3], together with H. Bass, E. Mutchnik, and D. Wright [1], proved that for any integral domains R and S of characteristic zero, $\text{JC}(R)$ is equivalent to $\text{JC}(S)$. Later, A. van den Essen

2020 *Mathematics Subject Classification.* Primary 14R15; Secondary 13F20.

Key words and phrases. Jacobian conjecture, Tate algebra.

[4, Theorem 1.1.12] generalized this result to any rings R and S that are subrings of \mathbb{Q} -algebras.

- *Positive characteristic case:* For a ring R of characteristic $c > 0$, $\text{JC}(R, n)$, and hence $\text{JC}(R)$, is known to be false. For instance, the polynomial map $F = (X_1 - X_1^c, \dots, X_n - X_n^c)$ provides a counterexample.

This paper defines and explores the **Tate-Jacobian Conjecture** (TJC), a variation of the Jacobian conjecture in the context of Tate algebras over rings equipped with a topology induced by an ideal. Let I be an ideal of a ring R . The Tate algebra $(R, I)\langle X_1, \dots, X_n \rangle$ consists of formal power series over R whose coefficients converge to zero in the I -adic topology on R . We study the following conjecture, referred to as $\text{TJC}(R, I, n)$, for rings R with an ideal I and integers $n \geq 1$:

If $F \in (R, I)\langle X_1, \dots, X_n \rangle^n$ satisfies $\det JF \in (R, I)\langle X_1, \dots, X_n \rangle^\times$
 $\text{TJC}(R, I, n)$ and $F(0) = 0$, then there exists $G \in (R, I)\langle X_1, \dots, X_n \rangle^n$ such that
 $F \circ G = G \circ F = (X_1, \dots, X_n)$.

Similarly to the Jacobian conjecture, $\text{TJC}(R, I)$ refers to the conjunction of $\text{TJC}(R, I, n)$ for all $n \geq 1$. The following cases illustrate its scope:

- If I is nilpotent, that is, $I^e = (0)$ for some $e \geq 1$, then $\text{TJC}(R, I, n)$ is equivalent to $\text{JC}(R, n)$.
- If $I = R$, then $\text{TJC}(R, I)$ corresponds to the analog of the Jacobian conjecture for formal power series, which is known to be true (see [4, Theorem 1.1.2]).

From the cases above, we see that the Tate-Jacobian conjecture generalizes the Jacobian conjecture. When the I -adic topology on R is Hausdorff, we obtain an equivalence between the two conjectures.

THEOREM 0.1. *Let R be a ring with an ideal I . If $\text{TJC}(R, I, n)$ holds, then $\text{JC}(R/I, n)$ holds. Conversely, if the I -adic topology on R is Hausdorff, then $\text{JC}(R/I, n)$ implies $\text{TJC}(R, I, n)$.*

By the previous theorem, together with the results from [1, 3, 4], we obtain the following corollary:

COROLLARY 0.2. *The following conditions are equivalent:*

- (i) $\text{TJC}(R, I)$ holds for some ring R with an ideal I such that the I -adic topology on R is Hausdorff and R/I is a subring of a \mathbb{Q} -algebra.
- (ii) $\text{JC}(R)$ holds for some ring R which is a subring of a \mathbb{Q} -algebra.
- (iii) $\text{JC}(R)$ holds for all rings R which are subrings of a \mathbb{Q} -algebra.

Additionally, when R/I has positive characteristic, we obtain the following corollary:

COROLLARY 0.3. *Let R be a ring and I an ideal such that R/I has characteristic $c > 0$. Then, $\text{TJC}(R, I, n)$ is false.*

Hence, the Tate-Jacobian conjecture fails for important rings such as the p -adic rings. However, by refining the assumptions of the Tate-Jacobian conjecture and applying a result from A. van den Essen and R. J. Lipton [5], we can establish an equivalence for the Jacobian conjecture within the framework of Tate algebras over p -adic rings:

THEOREM 0.4. *The Jacobian conjecture for \mathbb{C} is equivalent to the following statement: for every $n \geq 1$ and for all but finitely many primes p , if $F \in \mathbb{C}_p[X_1, \dots, X_n]^n$ satisfies $\det JF \in \mathbb{C}_p^\times$, then there exists $G \in \mathbb{C}_p \langle X_1, \dots, X_n \rangle^n$ such that $F \circ G = G \circ F = (X_1, \dots, X_n)$.*

Notation and Conventions:

- All rings are commutative and have a nonzero identity element.
- The notation X usually refers to the n variables X_1, \dots, X_n . For example, $R[X]$ denotes the ring of polynomials in n variables $R[X_1, \dots, X_n]$ (see Section 2).
- If the ideal I defining the I -adic topology on R is clear from context, we write the Tate algebra $(R, I) \langle X_1, \dots, X_n \rangle$ simply as $R \langle X_1, \dots, X_n \rangle$ (see Section 1).

Outline of the Paper. Section 1 provides a brief overview of the definition and notation of Tate algebras. In Section 2, we study the Tate-Jacobian conjecture and prove Theorem 0.1. Finally, Section 3 presents the proof of Theorem 0.4.

Acknowledgments. The authors would like to express their gratitude to Yuichiro Taguchi and Naganori Yamaguchi for their valuable feedback and insightful advice on the first draft of this paper.

1. GENERALITIES ON TATE ALGEBRAS

In this section, we recall the definition and the characterization of the unit group of Tate algebras over topological rings. For a more detailed account, we refer the reader to [2].

DEFINITION 1.1. Let R be a topological ring. The *Tate algebra* over R in n variables, denoted by $R \langle X_1, \dots, X_n \rangle$, is the subalgebra of the R -algebra of formal power series $R[[X_1, \dots, X_n]]$ consisting of power series

$$f(X_1, \dots, X_n) = \sum_{\nu_1, \dots, \nu_n \geq 0} a_{\nu_1, \dots, \nu_n} X_1^{\nu_1} \cdots X_n^{\nu_n} \in R[[X_1, \dots, X_n]]$$

such that the coefficients a_{ν_1, \dots, ν_n} tends to zero in the topology of R . Specifically, let $\{\mathcal{U}_i\}_{i \in I}$ be a system of neighborhood of 0 in the topology of R . Then, for every $i \in I$, we have $a_{\nu_1, \dots, \nu_n} \in \mathcal{U}_i$ for all but finitely many indices (ν_1, \dots, ν_n) .

REMARK 1.2. Let R be a ring with an ideal I . We denote by $(R, I) \langle X_1, \dots, X_n \rangle$ the Tate algebra over the ring R with the I -adic topology, and simply write $R \langle X_1, \dots, X_n \rangle$ when I is clear from context.

When R is equipped with the topology induced by an ideal I , we can determine the group of units $(R, I) \langle X_1, \dots, X_n \rangle^\times$. This is shown by the following criterion:

LEMMA 1.3 (Adapted from Proposition 1, Chapter 5, [2]). *A series $f \in (R, I) \langle X_1, \dots, X_n \rangle$ is a unit if and only if $f(0) \in R^\times$ and $f - f(0)$ has all its coefficients in the radical \sqrt{I} .*

PROOF. Suppose f is a unit in $(R, I) \langle X_1, \dots, X_n \rangle$. Then, f must also be a unit in the ring of formal power series $R[[X_1, \dots, X_n]]$. This implies $f(0) \in R^\times$. Furthermore, the reduction $\bar{f} \in (R/I)[X_1, \dots, X_n]$ is a unit, which forces all the coefficients of $f - f(0)$ to belong in \sqrt{I} . Conversely, assume $f(0) \in R^\times$ and that $f - f(0)$ has all its coefficients in \sqrt{I} . Let $b = f(0)^{-1} \in R$. Then, we can write $bf = 1 + u$, where $u(0) = 0$, $u \in (R, I) \langle X_1, \dots, X_n \rangle$, and the coefficients of u are in \sqrt{I} . Now, we claim that the sum $\sum_{i \geq 0} (-u)^i$, which is well-defined since $u(0) = 0$, belongs to $(R, I) \langle X_1, \dots, X_n \rangle$. Indeed, since u has only finitely many coefficients in $\sqrt{I} \setminus I$, some power of u must have all its coefficients in I . So for every positive integer N , there exists $a_N \in \mathbb{Z}$ such that u^{a_N} has all its coefficients in I^N . Since $u^i \in (R, I) \langle X_1, \dots, X_n \rangle$, there exists a positive integer $A_i^{(N)}$ such that if $\nu_1 + \dots + \nu_n > A_i^{(N)}$, then the coefficient a_{ν_1, \dots, ν_n} belongs to I^N . Now, defining

$$A^{(N)} := \max_{1 \leq i \leq a_N} A_i^{(N)},$$

it follows that for every u^i , the coefficient of any monomial $X^{\nu_1} \cdots X^{\nu_n}$ satisfying $\nu_1 + \dots + \nu_n > A^{(N)}$ is necessarily in I^N . Consequently, the same holds for the sum $\sum_{i \geq 0} (-u)^i$. Therefore, the product $b \sum_{i \geq 0} (-u)^i$ is the inverse of f in $(R, I) \langle X_1, \dots, X_n \rangle$. \square

2. ON THE TATE-JACOBIAN CONJECTURE

In this section, we study the Tate-Jacobian conjecture for rings equipped with a topology induced by an ideal. As stated in the introduction, we denote the n variables X_1, \dots, X_n simply by X .

THEOREM 0.1. *Let R be a ring with an ideal I . If $\text{TJC}(R, I, n)$ holds, then $\text{JC}(R/I, n)$ holds. Conversely, if the I -adic topology on R is Hausdorff, then $\text{JC}(R/I, n)$ implies $\text{TJC}(R, I, n)$.*

The key part of the proof is the following lemma, which relates the existence of the inverse of $F \in R \langle X \rangle^n$ to the existence of the inverse of its projection $\bar{F} \in (R/I)[X]^n$.

LEMMA 2.1. *Let R be a ring and I an ideal such that the I -adic topology on R is Hausdorff. For any $F \in R \langle X \rangle^n$ such that $F(0) = 0$, the following equivalence holds: F is invertible in $R \langle X \rangle^n$ if and only if its projection \bar{F} is invertible in $(R/I)[X]^n$.*

PROOF. To prove the equivalence, we first assume that F is invertible in $R\langle X \rangle^n$, and prove that \overline{F} is invertible in $(R/I)[X]^n$. By assumption, there exists $G \in R\langle X \rangle^n$ such that $F \circ G = G \circ F = X$. Consider the projection \overline{G} of G in $(R/I)[X]^n$. Then, we see that

$$X = \overline{G \circ F} = \overline{G} \circ \overline{F},$$

and conclude that \overline{G} is the inverse of \overline{F} in $(R/I)[X]^n$.

Next, we show the reverse implication. Suppose that \overline{F} is invertible in $(R/I)[X]^n$, and let $\overline{G} = (\overline{G}_1, \dots, \overline{G}_n)$ be its inverse. Let G_i be a lift of \overline{G}_i to $R[X]^n$ such that $G_i(0) = 0$ for every i . Then,

$$G_i(F(X)) - X_i \in IR\langle X \rangle,$$

where $IR\langle X \rangle$ denotes the subset of $R\langle X \rangle$ of power series whose coefficients belong to the ideal I . Hence, there exists an $f \in IR\langle X \rangle$ such that $f(0) = 0$ and

$$X_i = G_i(F(X)) + f \in R\langle F \rangle + IR\langle X \rangle,$$

where $R\langle F \rangle$ is the abbreviation of $R\langle F_1, \dots, F_n \rangle$. Since the choice of i was arbitrary, and the constant term of $G_i(F(X)) + f$ is zero, we obtain $R\langle X \rangle \subseteq R\langle F \rangle + IR\langle X \rangle$. Therefore,

$$R\langle X \rangle \subseteq R\langle F \rangle + IR\langle X \rangle \subseteq R\langle F \rangle + I(R\langle F \rangle + IR\langle X \rangle) \subseteq (R\langle F \rangle + IR\langle F \rangle) + I^2R\langle X \rangle.$$

By repeating this process for any natural number e we obtain:

$$R\langle X \rangle \subseteq (R\langle F \rangle + IR\langle F \rangle + \dots + I^{e-1}R\langle F \rangle) + I^eR\langle X \rangle \subseteq R\langle F \rangle + I^eR\langle X \rangle.$$

Since the I -adic topology is Hausdorff, it satisfies $\bigcap_{i \geq 1} I^i = (0)$. We conclude that $R\langle X \rangle \subseteq R\langle F \rangle$, and therefore $R\langle F \rangle = R\langle X \rangle$. This shows that F is invertible in $R\langle X \rangle^n$. \square

We now proceed to prove Theorem 0.1.

Proof of Theorem 0.1. First, suppose TJC(R, I, n) holds, and let $F \in (R/I)[X]^n$ be such that $\det JF \in (R/I)[X]^\times$. We claim that we may assume, without loss of generality, that $F = X + (\text{degree} \geq 2 \text{ terms})$. Indeed, let F_n denote the homogeneous component of F of degree n , so that F can be decomposed as $F = F_0 + F_1 + F_2 + \dots$. The Jacobian matrix JF can similarly be expressed as $JF = JF_1 + JF_2 + \dots$ where each JF_n is a matrix of homogeneous polynomials of degree $n - 1$. Since $\det JF \in (R/I)[X]^\times$, it follows that $\det JF_1 = \det JF(0) \in (R/I)^\times$. As F_1 is linear, there exists a linear map $G' \in (R/I)[X]^n$ such that $G' \circ F_1 = X$. Define

$$F' := G' \circ (X - F(0)) \circ F = X + (\text{degree} \geq 2 \text{ terms}).$$

If F' has an inverse map G , then the composition $G \circ G' \circ (X - F(0))$ is clearly the inverse map of F .

Next, choose a lift $\tilde{F} = X + (\text{degree} \geq 2 \text{ terms}) \in R[X]^n$ of F . Then, we have

$$\det J\tilde{F} = 1 + (\text{degree} \geq 1 \text{ terms}),$$

and

$$\overline{\det J\tilde{F}} = \det J\overline{\tilde{F}} \in (R/I)[X]^\times.$$

Hence, $\det J\tilde{F}(0) = 1 \in R^\times$ and $\det J\tilde{F} - \det J\tilde{F}(0)$ has coefficients in \sqrt{I} . By Lemma 1.3, we deduce that $\det J\tilde{F} \in (R, I)\langle X \rangle^\times$ and hence there exists $\tilde{G} \in R\langle X \rangle^n$ such that $\tilde{F} \circ \tilde{G} = \tilde{G} \circ \tilde{F} = X$. Therefore, the image G of \tilde{G} in $(R/I)[X]^n$ is the inverse of F .

Conversely, suppose the I -adic topology on the ring R is Hausdorff and $\text{JC}(R/I, n)$ holds. Let $F \in R\langle X \rangle^n$ be such that $F(0) = 0$ and $\det JF \in R\langle X \rangle^\times$, and denote by \overline{F} the image of F in $(R/I)[X]^n$. Since $\det J\overline{F} \in (R/I)[X]^\times$, by assumption, \overline{F} is invertible in $(R/I)[X]^n$. Thus, by Lemma 2.1, F is invertible in $R\langle X \rangle^n$. \square

3. PROOF OF THEOREM 0.4

We conclude this paper with a proof of Theorem 0.4. The main part of the proof was already established by A. van den Essen and R. J. Lipton [5], who showed that the Jacobian conjecture for \mathbb{C} holds if and only if the Unimodular conjecture for \mathbb{Z}_p holds for all but finitely many primes p :

UNIMODULAR CONJECTURE FOR \mathbb{Z}_p . *For every $n \geq 1$, and for every $F \in \mathbb{Z}_p[X]^n$ satisfying $\det JF \in \mathbb{Z}_p^\times$, there exists $b \in \mathbb{Z}_p^n$ such that $F(b) \in \mathbb{Z}_p^n$ is unimodular, that is, the ideal generated by the entries of the vector $F(b)$ is \mathbb{Z}_p .*

The idea of our proof is to relate Theorem 0.4 to the Unimodular conjecture. To proceed, we first establish a key result that enables us to reduce the problem to \mathbb{Z}_p :

LEMMA 3.1. *Let R and S be topological rings such that $R \subseteq S$ and R has the subspace topology induced from S . Let $F \in R\langle X \rangle^n$ be such that $F(0) = 0$ and $\det JF(0) \in R^\times$. If there exists $G \in S\langle X \rangle^n$ such that $F \circ G = G \circ F = X$, then $G \in R\langle X \rangle^n$.*

PROOF. By [4, Theorem 1.1.2], F has a unique inverse H in $R[[X]]^n$, which is also the unique inverse of F in $S[[X]]^n$. Since G is also an inverse of F in $S\langle X \rangle^n \subseteq S[[X]]^n$, the uniqueness of the inverse implies that $G = H$. Therefore, the inverse of F must belong to the intersection

$$R[[X]]^n \cap S\langle X \rangle^n.$$

By the definition of the Tate algebra, and since R has the subspace topology induced from S , this intersection is precisely $R\langle X \rangle^n$. \square

Proof of Theorem 0.4. For simplicity, we denote the following proposition by (\star) :

(\star) *For every $n \geq 0$ and for all but finitely many primes p , if $F \in \mathbb{C}_p[X]^n$ satisfies $\det JF \in \mathbb{C}_p^\times$, then there exists $G \in \mathbb{C}_p\langle X \rangle^n$ such that $F \circ G = G \circ F = X$.*

First, suppose that $\text{JC}(\mathbb{C})$ holds. By the results of [1, 3], $\text{JC}(\mathbb{C}_p)$ holds for all primes p , which clearly implies (\star) .

Conversely, suppose that (\star) holds, and let $F \in \mathbb{Z}_p[X]^n$ be such that $\det JF \in \mathbb{Z}_p^\times$. By a suitable translation, we assume without loss of generality that $F(0) = 0$. Under this assumption, we can apply Lemma 3.1 and conclude the existence of $G \in \mathbb{Z}_p\langle X \rangle^n$ such that $F \circ G = G \circ F = X$. Observe that G converges at every point $x \in \mathbb{Z}_p^n$, in particular, it converges at $\mathbf{1} = (1, \dots, 1)$. Hence, if we set $b = G(\mathbf{1})$, then $F(b) = (1, \dots, 1)$ is unimodular. Since the choice of $n \geq 1$ and the prime p were arbitrary, we conclude that the Unimodular conjecture for \mathbb{Z}_p holds for all but finitely many primes p . Therefore, by [5, Theorem 6], $J\mathcal{C}(\mathbb{C})$ holds. \square

REFERENCES

- [1] H. BASS, E. MUTCHNIK, AND D. WRIGHT, The Jacobian Conjecture: Reduction of Degree and Formal Expansion of the Inverse, *Bull. Amer. Math. Soc.* 7 (1982), 287–330.
- [2] S. BOSCH, U. GÜNTZER, AND R. REMMERT, *Non-Archimedean Analysis: A Systematic Approach to Rigid Analytic Geometry*, Springer Berlin, Heidelberg 1984.
- [3] E. CONNELL, L. VAN DEN DRIES, Injective polynomial maps and the Jacobian conjecture, *J. Pure Appl. Algebra* 28 (1983), 235–239.
- [4] A. VAN DEN ESSEN, *Polynomial Automorphisms and the Jacobian Conjecture*, *Progr. Math.* 190, Birkhäuser 2000.
- [5] A. VAN DEN ESSEN AND R. J. LIPTON, A p -adic approach to the Jacobian Conjecture, *J. Pure Appl. Algebra* 219 (2015), 2624–2628.
- [6] O.-H. KELLER, Ganze Cremona-Transformationen, *Monatsh. f. Mathematik und Physik* 47 (1939), 299–306.

DEPARTMENT OF MATHEMATICS
 INSTITUTE OF SCIENCE TOKYO
 2-12-1, O-OKAYAMA, MEGURO-KU, TOKYO 152-8551
 JAPAN
Email address: lucas.h.r.hamada@gmail.com

DEPARTMENT OF MATHEMATICS
 INSTITUTE OF SCIENCE TOKYO
 2-12-1, O-OKAYAMA, MEGURO-KU, TOKYO 152-8551
 JAPAN
Email address: kato.k.ba@m.titech.ac.jp

DEPARTMENT OF MATHEMATICS
 INSTITUTE OF SCIENCE TOKYO
 2-12-1, O-OKAYAMA, MEGURO-KU, TOKYO 152-8551
 JAPAN
Email address: komiya.r.ab@m.titech.ac.jp