

## CONTENTS

1. Introduction	1
2. Zip shift spaces	3
2.1. Definition of zip shift space and zip shift map	4
2.2. Higher block zip shifts, higher power zip shifts and zip shift sliding block codes	6
3. Pre-images of points under zip shift map	10
3.1. Constructing the labeled and backward labeled graphs for sofic zip shift spaces	11
3.2. Walking on graphs associated to a sofic space $\Sigma_{\mathcal{A}, \mathcal{A}'}$	12
4. Construction of an $N$ -to-1 endomorphism horseshoe	16
4.1. A conjugate zip shift map to the $N$ -to-1 horseshoe	17
5. Stable and unstable sets; periodic, pre-periodic, homoclinic, heteroclinic points and their orbits	20
5.1. Stable and unstable sets and periodic, homoclinic and heteroclinic orbits of the $N$ -to-1 horseshoe	23
References	26

## ZIP SHIFT SPACE

SANAZ LAMEI<sup>\*1</sup>, POUYA MEHDIPOUR<sup>2</sup>

ABSTRACT. We introduce a new extension in symbolic dynamics on two sets of alphabets, called the zip shift space. In finite case, it represents a finite-to-1 local homeomorphism called zip shift map. Such extension, offers a conjugacy between some endomorphisms and some zip shift map over two-sided space with finite sets of alphabets. As an application, the topological conjugacy of an  $N$ -to-1 uniformly hyperbolic horseshoe map with a zip shift map and its orbit structure is investigated. Moreover, the pre-image studies over zip shift space and the concepts of stable and unstable sets and homoclinic orbits, with a precise description for  $N$ -to-1 horseshoe are illustrated.

## 1. INTRODUCTION

Symbolic dynamics is a powerful implement to verify a worth amount of dynamical properties in smooth and topological dynamics. It provides a discrete modeling of the system. Any state of such modeling is represented by finite (or infinite) set of alphabets. The evolution of dynamics is displayed by the shift transformation over the space of one-sided or two-sided sequences of configurations.

---

*Date:* Received: xxxxxx; Revised: yyyyyy; Accepted: zzzzzz.

\* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 37B10; Secondary 37D05, 37C29.

*Key words and phrases.* zip shift, endomorphism horseshoe, endomorphism homoclinic orbit.

The idea of symbolic dynamics goes back to the studies of Hadamard (1899) on geodesic flows of surfaces with negative curvature [9]. After that the theory was further developed by the attempts of mathematicians such as Morse, Artin, Hedlund, Birkhoff, Levinson, Cartwright, Littlewood etc. [1], [3], [11]. Around 1970, the two-sided symbolic dynamics and the Markov partition was developed for hyperbolic toral automorphisms and Axiom A diffeomorphisms (including Anosov diffeomorphisms), by Adler, Weiss, Anosov, Sinai, and Bowen [3], [4], [6], [19]. In particular, R. Bowen used symbolic dynamics and Markov partition to study the Axiom A diffeomorphisms and it allowed the techniques of statistical mechanics to be applied to dynamical systems. Also S. Smale developed a method to determine the chaotic behavior of dynamical systems by giving a symbolic representation to the horseshoe map. Currently, this powerful method is improved in many areas of mathematics, such as ergodic theory, complex dynamics, partially hyperbolic diffeomorphisms, billiards, etc., and has significant applications in other research areas such as information theory, cellular automaton, statistical mechanics, quantum chaos, thermodynamics, etc [2], [18], [24].

In the context of invertible dynamics, the existence of a Markov partition [6], [11] provides a topological (or measure theoretical) conjugacy with some two-sided shift homeomorphism and offers convenient dynamical information. In non-invertible case, when the system admits a Markov partition, one can associate a semi-conjugacy to a shift map over one-sided shift space. In addition, another semi-conjugacy can be obtained via the natural extension (inverse limit) of the map over the (two-sided) orbit space. It is noteworthy that recently, E. Ermerson, Y. Lima and P. Mauricio [8], used the methods introduced in [16], [17], to construct countable Markov partitions for the natural extension of some non-invertible non-uniformly hyperbolic systems. This construction allowed to verify some ergodic and topological properties compatible with the inverse limit of the original map. In fact, there is lots of important data, which is accessible only for the natural extension of the map, and not necessarily for the original dynamics. To the best of our knowledge, outside the natural extension of the map which is a symbolic space over uncountably many symbols [7], [10], [22], no attempt has been done to develop a two-sided symbolic dynamics on finite alphabets for non-invertible dynamics.

In this work, we aim to introduce the zip shift maps over zip shift spaces with finite alphabets, which are extensions of two-sided shift maps over shift spaces. The zip shift map is defined as a local homeomorphism on the zip shift space which is the set of bi-infinite sequences over two sets of alphabets. This is because for non-invertible dynamics, although the forward dynamics is induced by the backward one, yet it can be a completely different dynamics, including important data. For instance, in Section 4 we compare the 2-to-1 horseshoe map and the map  $g(x) = 4x$  modulo 1. As it is known, both of the maps are semi-conjugate to the full one-sided shift over the full-shift in 4 symbols and the shift map can not distinguish these two dynamics. We show that these systems can be identified, by conjugation with different zip shift maps. From a measure theoretical point of view, in [13],[14], we introduce the notion of  $(m, l)$ -Bernoulli property as an extension to the previously known two-sided Bernoulli property.

It illustrates the fundamental role of investigating two-sided ergodic properties in identifying one-sided Bernoulli maps. From a dynamical system point of view we wish that, the zip shift map provides a new approach to the classification of local homeomorphisms and to a better comprehension of the statistical properties of local diffeomorphisms.

In addition, in this work, we present a concise construction of an  $N$ -to-1 horseshoe set, which is an example of admitting a good Markov partition for endomorphisms. In such cases the zip shift map acts like a zipper by taking special bi-infinite sequences and merge or collapse the entries of negative indices and create entries with new alphabet to describe the forward dynamics (Definition 2.1). When a system admits a ‘good’ Markov partition or even a ‘good’ labeling for its different states, (the labeled states in forward and backward should map together by dynamics) it is enough to represent the map by a conjugate zip shift map.

In what follows, Section 2 of this paper contains the introduction to the zip shift maps over zip shift spaces and also some basic definitions, properties and generalizations.

Section 3 proceeds with the study of pre-images of points under zip shift maps. In Section 3.2, for the case when the zip shift space is sofic, we defined a labeled graph where the bi-infinite walks on the graph represent the elements of the zip shift space. Example 3.1 shows that in general case, even for a simple zip shift space, it is not easy to find the exact pre-images of the points. In Theorem 3.5, we provide a way to find all the pre-images of a given point, up to sofic zip shift spaces.

A generalization of the Smale horseshoe to an  $N$ -to-1 case, is given in Section 4. Two natural Markov partitions, one for forward iterates and another for the backward iterates are stated. These partitions are mapped together by horseshoe map and give rise to a zip shift space. In Theorem 4.2 the corresponding conjugacy between a zip shift map and the horseshoe map is stated. In Example 4.3 the 2-to-1 horseshoe map and the map  $g(x) = 4x$  modulo 1 are compared.

In Section 5, the stable-unstable sets; periodic, pre-periodic and homoclinic points and their orbits for the zip shift map, are defined. Furthermore, as an application in Subsection 5.1 the stable-unstable sets for the  $N$ -to-1 horseshoe, also, their periodic and homoclinic points as well as their orbits, are investigated. These studies asserts many new data on the behavior and the importance of the role of the pre-periodic points (see Definition 5.3) homoclinic points and their orbits.

## 2. ZIP SHIFT SPACES

Consider two finite sets of alphabets  $\mathcal{A} = \{a, \dots, b\}$  and  $\mathcal{A}' = \{a', \dots, b'\}$ . Let  $\Sigma_{\mathcal{A}'} \subseteq \{(x_i)_{i \in \mathbb{N} \cup \{0\}}, x_i \in \mathcal{A}'\}$  be a one-sided shift space. A finite sequence  $w = w_{[k, \ell]} = x_k x_{k+1} \dots x_\ell$  which occurs in some point  $(x_i)_{i \in \mathbb{N} \cup \{0\}} \in \Sigma_{\mathcal{A}'}$  is called a *word* or *block* of length  $|w| = \ell - k + 1$ . Let  $\beta_n^{\mathcal{A}'} = \beta_n^{\mathcal{A}'}(\Sigma_{\mathcal{A}'})$  be the set of all admissible words of length  $n$  and  $\beta^{\mathcal{A}'} = \beta^{\mathcal{A}'}(\Sigma_{\mathcal{A}'}) = \bigcup_{n \geq 1} \beta_n^{\mathcal{A}'}$ . Obviously,  $\beta_1^{\mathcal{A}'} = \mathcal{A}'$ .

**2.1. Definition of zip shift space and zip shift map.** Let  $\varphi_n : \beta_n^{\mathcal{A}' } \rightarrow \mathcal{A}$  be a transition factor map which is onto (not necessarily invertible) with domain  $\beta_n^{\mathcal{A}'}$ . Associate to  $\varphi_n$  a 0-1 transition matrix  $T$  such that for the word  $a'_1 \dots a'_n \in \beta_n^{\mathcal{A}'}$  and  $a \in \mathcal{A}$ ,  $T(a'_1 \dots a'_n, a) = 1$  if and only if  $\varphi_n(a'_1 \dots a'_n) = a$ .

Let  $Y = \{y = (y_i)_{i \in \mathbb{Z}}, y_i \in \mathcal{A}'\}$  be a two-sided shift space with its shift map  $\sigma_Y$ . To any point  $y \in Y$ , correspond a point  $x = (x_i)_{i \in \mathbb{Z}} = (\dots, x_{-2}, x_{-1}; x_0, x_1, x_2, \dots)$  such that

$$x_i = \begin{cases} y_i \in \mathcal{A}' & \forall i \geq 0 \\ \varphi_n(y_i \dots y_{i+n-1}) \in \mathcal{A} & \forall i < 0. \end{cases} \quad (2.1)$$

According to set  $Y$ , define the *zip shift space* as the set

$$\Sigma_{\mathcal{A}, \mathcal{A}'} := \{x = (x_i)_{i \in \mathbb{Z}} : x_i \text{ satisfies (2.1)}\}.$$

If  $Y$  is a full shift space, then  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  will be called a *full zip shift space*. Let  $\Sigma_{\mathcal{A}'} = \{(x_0, x_1, \dots) : (x_i)_{i \in \mathbb{Z}} \in \Sigma_{\mathcal{A}, \mathcal{A}'}\}$  and  $\Sigma_{\mathcal{A}}^- = \{(\dots, x_{-2}, x_{-1}) : (x_i)_{i \in \mathbb{Z}} \in \Sigma_{\mathcal{A}, \mathcal{A}'}\}$ . From here on, we denote  $\Sigma_{\mathcal{A}}^-$  by  $\Sigma_{\mathcal{A}}$ . The set  $\Sigma_{\mathcal{A}}$  is a factor of  $\Sigma_{\mathcal{A}'}$  due to the map  $\varphi_n$ . Define the *zip shift map* on  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  with same notation for shift maps as follows

$$(\sigma(x))_i = \begin{cases} \varphi_n(x_0 \dots x_{n-1}) & \text{if } i = -1 \\ x_{i+1} & \text{otherwise.} \end{cases}$$

The zip shift map is a local homeomorphism generalization of a two-sided shift homeomorphism. If one let  $\mathcal{A} = \mathcal{A}'$  and  $\varphi_1 = Id$  then the zip shift map is a two-sided shift map.

Any subspace  $\Sigma$  of  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  which is closed under  $\sigma$  is called a *sub zip shift space* of  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ . Now  $\beta_n^{\mathcal{A}, \mathcal{A}'} = \beta_n^{\mathcal{A}, \mathcal{A}'}(\Sigma)$  is the set of all words of length  $n$  in  $\Sigma$ . The set  $\beta^{\mathcal{A}, \mathcal{A}'} = \beta^{\mathcal{A}, \mathcal{A}'}(\Sigma) = \bigcup_{n \geq 1} \beta_n^{\mathcal{A}, \mathcal{A}'}(\Sigma)$  is the set of all *admissible* words or the *language* of  $\Sigma$ . Let  $\beta_n^{\mathcal{A}}$  be the subset of  $\beta^{\mathcal{A}, \mathcal{A}'}(\Sigma)$  with alphabets in  $\mathcal{A}$  and  $\beta^{\mathcal{A}} = \bigcup_{n \geq 1} \beta_n^{\mathcal{A}}$ . For  $w' = a'_i a'_{i+1} \dots a'_{i+m} \in \beta^{\mathcal{A}'}$  and  $m > n$ , we simply extend the definition of  $\varphi_n$  as

$$\varphi_n(w') = \varphi_n(a'_i \dots a'_{i+n-1}) \varphi_n(a'_{i+1} \dots a'_{i+n}) \dots \varphi_n(a'_{i+m-n+1} \dots a'_{i+m}) \in \beta^{\mathcal{A}}. \quad (2.2)$$

Let  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  be a full zip shift space. For  $\Sigma \subset \Sigma_{\mathcal{A}, \mathcal{A}'}$  with admissible words  $\beta^{\mathcal{A}, \mathcal{A}'}$ , the words in  $\beta^{\mathcal{A}, \mathcal{A}'}(\Sigma_{\mathcal{A}, \mathcal{A}'}) \setminus \beta^{\mathcal{A}, \mathcal{A}'}(\Sigma)$  will not appear in any sequences in  $\Sigma$ . Let  $\mathcal{F}$  be a minimal set of words (a set with fewest and shortest words) such that no word in  $\beta^{\mathcal{A}, \mathcal{A}'}(\Sigma)$  contains a word of  $\mathcal{F}$ . This set is called the set of *forbidden blocks* of  $\Sigma$ . Let  $\mathcal{F}^{\mathcal{A}}$  and  $\mathcal{F}^{\mathcal{A}'}$  be the subsets of  $\mathcal{F}$  determining the forbidden blocks of the sets  $\Sigma_{\mathcal{A}}$  and  $\Sigma_{\mathcal{A}'}$  respectively. Note that if  $w' = a'_1 a'_2 \dots a'_m \in \mathcal{F}$ , then  $\varphi_n$  will not be defined for any word containing  $w'$ .

Recall that a shift space is a subshift of finite type (SFT), if there are only finite forbidden blocks with finite length. For any SFT, there exists a  $K$  such that the SFT is a  $K$ -step Markov space and the length of forbidden blocks is less than  $K + 1$ . The image of an SFT under a factor map is called a sofic space [11]. For a zip shift space  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ , if  $|\mathcal{F}| < \infty$  and the length of longest word in  $\mathcal{F}$  is  $M + 1$ , then call  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  an  *$M$ -step Markov space*. More precisely, suppose that  $\Sigma_{\mathcal{A}'}$  is  $K$ -step and  $\Sigma_{\mathcal{A}}$  is  $N$ -step. Then for  $M = \max\{N, K\}$ ,  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  is an  $M$ -step Markov space. In this case one can associate 3 matrices such that they all together represent the




FIGURE 1. The right graph is the labeled graph  $\mathcal{G}$  of the left graph.

$M$ -step space  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ . As for shift spaces,  $\Sigma_{\mathcal{A}'}$  can be represented by an adjacency matrix  $A'$  such that for words  $v' = a'_1 \dots a'_K, w' = b'_1 \dots b'_K \in \beta_K^{\mathcal{A}'}$ ,  $A'(v', w') = 1$  if and only if  $a'_1 \dots a'_{K-1} = b'_2 \dots b'_K$  and  $a'_1 \dots a'_K b'_K \in \beta_{K+1}^{\mathcal{A}'}$ . The matrices  $A$  and  $T$  are defined as follows. For words  $v = a_1 \dots a_N, w = b_1 \dots b_N \in \beta_N^{\mathcal{A}}$ ,  $A(v, w) = 1$  if and only if  $a_1 \dots a_{N-1} = b_2 \dots b_N$  and  $a_1 \dots a_N b_N \in \beta_{N+1}^{\mathcal{A}}$ . Also,  $T(v, v') = 1$  if and only if  $vv' \in \beta_{N+K}^{\mathcal{A}, \mathcal{A}'}$ . We call  $A$  and  $A'$  the *adjacency matrices* due to alphabets  $\mathcal{A}$  and  $\mathcal{A}'$ . The matrix  $T$  can be defined by using the words of maximum length  $M + 1$  but it would be a larger matrix.

The space  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  is an SFT only if both  $\Sigma_{\mathcal{A}}$  and  $\Sigma_{\mathcal{A}'}$  are SFTs. For an SFT or sofic space  $\Sigma_{\mathcal{A}'}$ , the factor space  $\Sigma_{\mathcal{A}}$  can be either SFT or sofic. If both  $\Sigma_{\mathcal{A}}$  and  $\Sigma_{\mathcal{A}'}$  are sofic spaces (any SFT is sofic), then call  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  a *sofic space*. Such space is called *strictly sofic* if it is not an SFT.

If  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  does not depend on time, then in some cases it can be represented by graphs. Let  $\Sigma_{\mathcal{A}'}$  be a sofic (not necessarily strictly sofic) space with labeled graph  $\mathcal{G} = (G, \mathcal{L})$  (see [11] for definitions). Let  $y = (y_i)_{i \in \mathbb{Z}}$  be a bi-infinite walk on  $\mathcal{G}$ . Then  $x = (x_i)_{i \in \mathbb{Z}}$  is a point in  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  if  $x_i = y_i$  for  $i \geq 0$  and  $x_i = \varphi_n(y_i, \dots, y_{i+n})$  for  $i \leq -1$ .

**Example 2.1.** Let  $G$  be the left graph in Figure 1. Let  $\mathcal{A}' = \{1, 2, 3\}$  be the alphabet set and  $\mathcal{F} = \{11, 13, 21, 33\}$  be the forbidden set. The corresponding labeled graph is the right graph in Figure 1. Suppose  $\mathcal{A} = \{a, b\}$  with  $\varphi_1(1) = a$  and  $\varphi_1(2) = \varphi_1(3) = b$ . A typical sequence in  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  will be

$$\begin{aligned} x &= (\dots, x_{-3}, x_{-2}, x_{-1}; x_0, x_1, x_2, x_3, x_4, \dots) \\ &= (\dots, b, a, b; 3, 1, 2, 2, 3, \dots), \end{aligned}$$

which is obtained by a walk on labeled graph  $\mathcal{G}$ .

**Definition 2.2.** Let  $\Sigma$  be a sub zip shift space with alphabets  $\mathcal{A}$  and  $\mathcal{A}'$ .

- The space  $\Sigma$  is called *irreducible* if for any  $a', b' \in \mathcal{A}'$ , there exists a word  $w' \in \beta^{\mathcal{A}'}$  such that  $a'w'b' \in \beta^{\mathcal{A}'}$ .
- A point  $(x_i)_{i \in \mathbb{Z}} \in \Sigma$  is *periodic* with period  $m$  if there exist  $p'_0, \dots, p'_{m-1} \in \mathcal{A}'$  such that for  $i \geq 0$ ,  $x_{mi} = p'_0, x_{mi+1} = p'_1, \dots, x_{mi+m-1} = p'_{m-1}$  and also,  $x_{mi} = \varphi_n(x_0, \dots, x_{n-1}), x_{mi+1} = \varphi_n(x_1, \dots, x_n), \dots$  and  $x_{mi+m-1} = \varphi_n(x_{m-1}, \dots, x_{m+n-2})$  for  $i \leq -1$ . Denote such a periodic point by  $p = \overline{(p'_0, p'_1, \dots, p'_{m-1})}$ . Let  $\text{Per}_m(\Sigma)$  be the set of all periodic points with period  $m$  and  $\text{Per}(\Sigma) = \cup_{m \geq 0} \text{Per}_m(\Sigma)$ .

*Remark 2.3.* • Since  $\Sigma$  is closed under  $\sigma$ , there is no need to verify the irreducibility for the words in  $\beta^{\mathcal{A}}$ .

- If  $\Sigma$  is representable by graph  $G$  or  $\mathcal{G} = (G, \mathcal{L})$ , then  $\Sigma$  is irreducible if and only if  $G$  is a strongly connected graph.
- If matrices  $A$ ,  $A'$  and  $T$  associate to an SFT zip shift space  $\Sigma$ , then  $\Sigma$  is irreducible if and only if for any  $a', b' \in \mathcal{A}'$ , there exists  $n \geq 0$  with  $A'^n(a', b') \geq 1$ .
- Suppose the factor map  $\varphi_n$  is not one-to-one. Then for some distinct words  $w' = a'_1 \dots a'_n$  and  $w'' = a''_1 \dots a''_n$  both in  $\beta_{\mathcal{A}'}(\Sigma)$ , we may have  $\varphi_n(w') = \varphi_n(w'')$ . For this reason, the repeating part in negative indices of the periodic point may have different length from the repeating part in non-negative indices. For example the point  $y = (\dots, b, b, b; 2, 3, 2, 3, 2, 3, \dots) = (\overline{2, 3})$  is a periodic point in Example 2.1. The repeating part in negative and non-negative indices are different. Also, the distinct points  $y$  and  $z = (\dots, b, b, b; 3, 2, 2, 3, 2, 2, \dots) = (\overline{3, 2, 2})$  have the same negative part.
- If  $A'$  is the adjacency matrix for an SFT space  $\Sigma$ , then  $\text{trace}(A'^n)$  determines the number of points of period  $n$ .

**2.2. Higher block zip shifts, higher power zip shifts and zip shift sliding block codes.** The definitions and theorems in this subsection are generalizations of the similar ones for shift spaces defined in [11].

**Definition 2.4.** Let  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  be a zip shift space. For some natural number  $N$ ,

- i) define the  $N$ th higher block code  $h_N : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow (\beta_N^{\mathcal{A}, \mathcal{A}'}(\Sigma_{\mathcal{A}, \mathcal{A}'}))^{\mathbb{Z}}$  as

$$(h_N(x))_{[i]} := x_{[i, i+N-1]}$$

and the  $N$ th higher block zip shift  $\Sigma_{\mathcal{A}, \mathcal{A}'}^{[N]}$  as  $\Sigma_{\mathcal{A}, \mathcal{A}'}^{[N]} = h_N(\Sigma_{\mathcal{A}, \mathcal{A}'})$ .

- ii) let  $\gamma_N : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow (\beta_N^{\mathcal{A}, \mathcal{A}'}(\Sigma_{\mathcal{A}, \mathcal{A}'}))^{\mathbb{Z}}$  be the  $N$ th higher power code

$$(\gamma_N(x))_{[i]} := x_{[iN, iN+N-1]}$$

and define the  $N$ th higher power zip shift  $\Sigma_{\mathcal{A}, \mathcal{A}'}^N$  as  $\Sigma_{\mathcal{A}, \mathcal{A}'}^N = \gamma_N(\Sigma_{\mathcal{A}, \mathcal{A}'})$ .

Note that  $h_N(x)$  and  $\gamma_N(x)$  are defined with respect to the order of successive blocks of length  $N$  occurring in  $x$ . The next theorem follows from the definition of zip shift space and Theorems 1.4.3 and 1.4.6 in [11].

**Theorem 2.5.** *The higher block zip shift code and higher power zip shift code of a zip shift space are zip shift spaces.*

To define the sliding block code, suppose  $X \subset \mathcal{C}^{\mathbb{Z}}$  and  $Y \subset \mathcal{C}'^{\mathbb{Z}}$  are two shift spaces with alphabets in  $\mathcal{C}$  and  $\mathcal{C}'$ . For these spaces, the block map  $\Phi : \beta_{M+N+1}^{\mathcal{C}}(X) \rightarrow \mathcal{C}'$  maps the admissible blocks of  $X$  with length  $M + N + 1$  to letters in  $\mathcal{C}'$  [11]. This block map factors the points  $x = (\dots x_{-1}, x_0, x_1, \dots) \in X$  to a point  $y = (\dots y_{-1}, y_0, y_1, \dots) \in Y$  and naturally implies a map  $\phi : X \rightarrow Y$  such that  $\phi(x) = y$ . This definition extends to sliding block codes for zip shift spaces. In new definition, we just consider the anticipation  $N$ .

**Definition 2.6.** Let  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  be a zip shift space with transition maps  $\varphi_n^{\mathcal{A}, \mathcal{A}'} : \beta_n^{\mathcal{A}'} \rightarrow \mathcal{A}$ . For two alphabet sets  $\mathcal{C}$  and  $\mathcal{C}'$ , define a map

$$\Psi : \beta_n^{\mathcal{A}, \mathcal{A}'} \cup \beta_{n+1}^{\mathcal{A}, \mathcal{A}'} \rightarrow \mathcal{C} \cup \mathcal{C}'$$

such that

$$\Psi(x_{[-1, n-1]}) = y_{-1} \in \mathcal{C}, \text{ and } \Psi(x_{[0, n-1]}) = y_0 \in \mathcal{C}'. \quad (2.3)$$

For  $m > 0$ , define  $\beta_m^{\mathcal{C}'} := [y_0, \dots, y_{m-1}]$  induced by iterations of  $\Psi$  over subblocks of  $x_{[0, m+n-2]}$ . If there exists a transition map  $\varphi_m^{\mathcal{C}, \mathcal{C}'} : \beta_m^{\mathcal{C}'} \rightarrow \mathcal{C}$  such that

$$y'_{-1} := \varphi_m^{\mathcal{C}, \mathcal{C}'}([y_0, \dots, y_{m-1}]) = \Psi([\varphi_n^{\mathcal{A}, \mathcal{A}'}[x_0, \dots, x_{n-1}], x_1, \dots, x_{n-1}]), \quad (2.4)$$

then call  $\Psi$  a *block map*. If there exists a map  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  such that for any  $x \in \Sigma_{\mathcal{A}, \mathcal{A}'}$ , there exists  $y \in \Sigma_{\mathcal{C}, \mathcal{C}'}$  such that  $\psi(x) = y$  and  $y_i$  is given by  $\Psi$  satisfying in (2.3) and (2.4), then call  $\psi$  the *zip shift sliding block code* induced by block map  $\Psi$ .

*Remark 2.7.* • In Theorem 2.12, we prove that the image of a zip shift sliding block code is in fact a zip shift space. Therefore, we used the notation of the zip shift space  $\Sigma_{\mathcal{C}, \mathcal{C}'}$  for  $\psi(\Sigma_{\mathcal{A}, \mathcal{A}'})$ .

- The equations (2.3) and (2.4) are the result of the following diagram, where  $\Psi$  should induce a “local commutation” rule between the maps  $\varphi_n^{\mathcal{A}, \mathcal{A}'}$ ,  $\varphi_m^{\mathcal{C}, \mathcal{C}'}$ ,  $\sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}}$  and  $\sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}}$  as

$$\begin{array}{ccc} [x_{-1}; x_0, x_1, \dots, x_k] & \longrightarrow & [\varphi_n^{\mathcal{A}, \mathcal{A}'}[x_0, \dots, x_{n-1}]; x_1, \dots, x_{k+1}] \\ \downarrow & & \downarrow \\ [y_{-1}; y_0, \dots, y_{k-n}] & \longrightarrow & [y'_{-1}; y_1, \dots, y_{k-n+1}]. \end{array} \quad (2.5)$$

Here,  $k = n + m - 2$ .

- The zip shift sliding block code can be defined through a block map  $\Theta : \beta_{k+2}^{\mathcal{A}, \mathcal{A}'} \rightarrow \beta_{k-n+2}^{\mathcal{C}, \mathcal{C}'}$  over blocks of length  $k \geq n + m - 2$ , but still any entry of  $\theta(x_{[-1, k]})$  is determined by  $\Psi$  defined in equations (2.3) and (2.4).
- By equation (2.3), the definition of  $\Psi$  naturally extends to blockes with all positive indeces  $i \geq 0$ . For negative indeces, let  $\ell = \max\{n, m\}$ . Then  $\ell$  iteration of diagram (2.5) defines a rule to convert a block in  $\beta_\ell^{\mathcal{A}, \mathcal{A}'}$  to a block in  $\beta_\ell^{\mathcal{C}, \mathcal{C}'}$ .

**Theorem 2.8.** *Let  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  and  $\Sigma_{\mathcal{C}, \mathcal{C}'}$  be two zip shift spaces with transition maps  $\varphi_n^{\mathcal{A}, \mathcal{A}'}$  and  $\varphi_m^{\mathcal{C}, \mathcal{C}'}$  as in Definition 2.6 and  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  be a zip shift sliding block code. Let  $\sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}}$  and  $\sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}}$  be the zip shift maps on spaces  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  and  $\Sigma_{\mathcal{C}, \mathcal{C}'}$  respectively. Then  $\psi \circ \sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}} = \sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}} \circ \psi$ .*

*Proof.* Let  $x = (\dots x_{-1}; x_0, x_1, \dots) \in \Sigma_{\mathcal{A}, \mathcal{A}'}$  and  $y = (\dots y_{-1}; y_0, y_1, \dots) \in \Sigma_{\mathcal{C}, \mathcal{C}'}$ . Suppose  $\psi(x) = y$  according to definition of  $\Psi$  in (2.3) and (2.4). For a word  $x_{[-1, n-1]} \in \beta_{n+1}^{\mathcal{A}, \mathcal{A}'}$ , a word  $x_{[i, i+n-1]} \in \beta_n^{\mathcal{A}, \mathcal{A}'}$  with  $i \geq 0$  or a word  $x_{[i, i+\ell-1]} \in \beta_\ell^{\mathcal{A}, \mathcal{A}'}$  with  $i \leq -\ell$ ,

$$\begin{aligned} \left( (\sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}} \circ \psi)(x) \right)_{[i]} &= \Psi \left( \sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}}(x)_{[i, i+n-1]} \right) \\ &= \Psi(x_{[i+1, i+n]}) \\ &= (\psi \circ \sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}}(x))_{[i]}. \\ &= y_{[i+1]} \\ &= (\sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}} \circ \psi)_{[i]}. \end{aligned}$$

For a block  $x_{[i, i+\ell-1]} \in \beta_\ell^{\mathcal{A}, \mathcal{A}'}$ , where  $-\ell < i < -1$ , at most  $\ell$  iteration of  $x$  and  $y$  will determine the entries with negative indices. Then the above discussion finishes the proof.  $\square$

If  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  is onto, then call it a *zip shift factor code* from  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  onto  $\Sigma_{\mathcal{C}, \mathcal{C}'}$ . If  $\psi$  is invertible, then it is called a *conjugacy* from  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  to  $\Sigma_{\mathcal{C}, \mathcal{C}'}$ . Both maps  $h_N(x)$  and  $\gamma_N(x)$  are conjugacies between the space  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  and spaces  $h_N(\Sigma_{\mathcal{A}, \mathcal{A}'})$  and  $\gamma_N(\Sigma_{\mathcal{A}, \mathcal{A}'})$  respectively.

**Corollary 2.9.** *Let  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  be a sliding block code. Then if  $p \in \Sigma_{\mathcal{A}, \mathcal{A}'}$  is a periodic point with period  $k$ , then the least period of  $\psi(p)$  divides  $k$ . If  $\psi$  is a conjugacy, then  $p$  and  $\psi(p)$  have the same period.*

The following theorem is a generalization of Curtis-Hedlund-Lyndon Theorem to zip shift maps.

**Theorem 2.10.** *Let  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  and  $\Sigma_{\mathcal{C}, \mathcal{C}'}$  be two zip shift spaces with transition maps  $\varphi_n^{\mathcal{A}, \mathcal{A}'} : \beta_n^{\mathcal{A}'} \rightarrow \mathcal{A}$  and  $\varphi_m^{\mathcal{C}, \mathcal{C}'} : \beta_m^{\mathcal{C}'} \rightarrow \mathcal{C}$  respectively. A map  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  is a zip shift sliding block code if and only if  $\psi \circ \sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}} = \sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}} \circ \psi$  and  $\psi(x)_{-1}$  is a function of  $x_{[-1, n-1]}$  and  $\psi(x)_0$  is a function of  $x_{[0, n-1]}$  such that the block map  $\Psi$  induced by  $\psi$  satisfies (2.4).*

*Proof.* We only need to prove the reverse direction. Let  $\psi$  induce a block map  $\Psi$  such that it is determined on blocks  $x_{[-1, n-1]}$  and  $x_{[0, n-1]}$ . Since it is defined on blocks  $x_{[0, n-1]}$ , the definition of  $\Psi$  naturally extends to blocks  $y_i = \Psi(x_{[i, i+n-1]})$ ,  $i \geq 0$ . Now according to (2.4) and Remark 2.7, the definition extends to blocks of length  $n + m + 1$  containing negative indices.  $\square$

**Theorem 2.11.** *Let  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  be a zip shift sliding block code with the block map  $\Psi : \beta_\ell^{\mathcal{A}, \mathcal{A}'} \rightarrow \mathcal{C} \cup \mathcal{C}'$ . Then there exists a higher block zip shift space  $\widetilde{\Sigma_{\mathcal{A}, \mathcal{A}'}}$  of  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ , a conjugacy  $h : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \widetilde{\Sigma_{\mathcal{A}, \mathcal{A}'}}$  and a 1-block code  $\tilde{\psi} : \widetilde{\Sigma_{\mathcal{A}, \mathcal{A}'}} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  such that  $\tilde{\psi} \circ h = \psi$ .*

*Proof.* Take  $\widetilde{\Sigma_{\mathcal{A}, \mathcal{A}'}}$  as  $\Sigma_{\mathcal{A}, \mathcal{A}'}^{[\ell]}$  and define  $h$  to be the  $\ell$ th higher block code  $(h(x))_{[i]} = x_{[i, i+\ell-1]}$ . The map  $h$  is a conjugacy and so  $\tilde{\psi} = \psi \circ h^{-1}$  is a 1-block map.  $\square$

**Theorem 2.12.** i) *The image of a zip shift space under the zip shift sliding block code is a zip shift space.*  
 ii) *A one-to-one and onto zip shift sliding block code has a zip shift sliding block code inverse.*

*Proof.* The proofs of parts i and ii are similar to the proofs of Theorems 1.5.13 and 1.5.14 [11] respectively.

(i). Let  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  and  $\Sigma_{\mathcal{C}, \mathcal{C}'}$  be two zip shift spaces with block map  $\Psi : \beta_\ell^{\mathcal{A}, \mathcal{A}'} \rightarrow \mathcal{C} \cup \mathcal{C}'$ , and  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  be a zip shift sliding block code. According to Definition 2.6 and Remark 2.7, for  $x \in \Sigma_{\mathcal{A}, \mathcal{A}'}$ , any block in  $\psi(x)$  belongs to  $\beta^{\mathcal{C}, \mathcal{C}'}$ . Therefore,  $\psi(\Sigma_{\mathcal{A}, \mathcal{A}'}) \subseteq \Sigma_{\mathcal{C}, \mathcal{C}'}$ .

To prove the inverse, Observe that according to Remark 2.7 and Theorem 2.10, the definition of block map  $\Psi$  can extend to map central blocks  $x_{[-k, k]}$  where



$k \geq n + m - 2$ . Now for any  $k$  the central  $(2k - 2n + 1)$ -block of  $y$  is the  $\Psi$ -image of the central  $(2k + 1)$ -block of some point  $x^{(k)} \in \Sigma_{\mathcal{A}, \mathcal{A}'}$ . Since  $\mathcal{C} \cup \mathcal{C}'$  contains finite symbols, therefore, there exists an infinite set  $S_k$  of integers such that  $x_{[-k, k]}^{(n)}$  is the same for all  $n \in S_k$ . Continuing this procedure and increasing  $k$ , we find nested subsets of infinite sets as  $\cdots \subseteq S_{k+1} \subseteq S_k$ . Define  $x_{[-k, k]} = x_{[-k, k]}^{(n)}$  for all  $n \in S_k$ . Therefore, for any  $k \geq n + m - 2$  we have

$$\Psi(x_{[-k, k]}) = \Psi(x_{[-k, k]}^{(n)}) = \psi(x^{(n)})_{[-k, k]} = y_{[-k, k]}.$$

This means  $\psi(x) = y$  and  $\Sigma_{\mathcal{C}, \mathcal{C}'} \subseteq \psi(\Sigma_{\mathcal{A}, \mathcal{A}'})$ .

(ii) Let  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  be a one-to-one and onto zip shift sliding block code and  $\xi : \Sigma_{\mathcal{C}, \mathcal{C}'} \rightarrow \Sigma_{\mathcal{A}, \mathcal{A}'}$  be the inverse of  $\psi$ . From  $\psi \circ \sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}} = \sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}} \circ \psi$  and  $y = \psi(x)$ , we have

$$\sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}}(\xi(y)) = \sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}}(x) = \xi(\psi(\sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}}(x))) = \xi(\sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}}(\psi(x))) = \xi(\sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}}(y)).$$

Therefore,  $\xi \circ \sigma_{\Sigma_{\mathcal{C}, \mathcal{C}'}} = \sigma_{\Sigma_{\mathcal{A}, \mathcal{A}'}} \circ \psi$ . Now according to Theorem 2.10, it suffices to determine  $\xi(y)_{[-1]}$  and  $\xi(y)_{[0]}$  over blocks  $y_{[-1, m-1]}$  and  $y_{[0, m-1]}$  respectively.

Observe that if there exists a block map  $\Xi$  defined over blocks  $\xi(y)_{[0]}$  and  $\xi(y)_{[-1]}$ , then its definition would extend over any block  $\xi(y)_{[-t, t]}$ ,  $t \geq m - 1$ . We prove the theorem for index 0 and the proof for index -1 is the same. By contradiction suppose that for any  $t \geq 0$ , there are two points  $y^{(t)}$  and  $\tilde{y}^{(t)}$  in  $\Sigma_{\mathcal{C}, \mathcal{C}'}$  such that  $y_{[-t, t]}^{(t)} = \tilde{y}_{[-t, t]}^{(t)}$ , but  $\xi(y^{(t)})_{[0]} \neq \xi(\tilde{y}^{(t)})_{[0]}$ . Let  $\xi(y^{(t)}) = x^{(t)}$  and  $\xi(\tilde{y}^{(t)}) = \tilde{x}^{(t)}$ . Since  $\mathcal{A} \cup \mathcal{A}'$  is finite, there would be distinct symbols  $a \neq b$  and an infinite set  $S_0$  of integers so that  $x_{[0]}^{(t)} = \xi(y^{(t)})_0 = a$  and  $\tilde{x}_{[0]}^{(t)} = \xi(\tilde{y}^{(t)})_0 = b$  for all  $t \in S_0$ . Then choose an infinite subset  $S_1 \subset S_0$  so that  $x_{[-1, 1]}^{(t)}$  are equal for  $t \in S_1$  and  $\tilde{x}_{[-1, 1]}^{(t)}$  are equal for  $t \in S_1$ . Continuing this way, for each  $k \geq 1$ , we would find infinite subset  $S_k \subset S_{k+1}$ .

As in the proof of part (i), we construct points  $x$  and  $\tilde{x}$  in  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  such that  $x_{[-t, t]} = x_{[-t, t]}^{(k)}$  and  $\tilde{x}_{[-t, t]} = \tilde{x}_{[-t, t]}^{(k)}$  for all  $n \in S_k$ . Note that  $x_{[0]} = a \neq b = \tilde{x}_{[0]}$ . Therefore,  $x \neq \tilde{x}$ . Now for  $t \in S_k$  and  $t \geq k$ . we have

$$\begin{aligned} \Psi(x_{[-k, k]}) &= \Psi(x_{[-k, k]}^{(t)}) = \psi(x^{(t)})_{[-k, k]} = y_{[-k, k]}^{(t)} \\ &= \tilde{y}_{[-k, k]}^{(t)} = \psi(\tilde{x}^{(t)})_{[-k, k]} = \Psi(\tilde{x}_{[-k, k]}^{(t)}) = \Psi(\tilde{x}_{[-k, k]}). \end{aligned}$$

This would imply that  $\psi(x) = \psi(\tilde{x})$  which is a contradiction.

If condition (2.4) does not hold for  $\Xi$ , it contradicts the map  $\xi$  to be one-to-one.  $\square$

*Remark 2.13.* • Let  $\psi : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{C}, \mathcal{C}'}$  be a one-to-one sliding block code induced by  $\Psi$ . Note that the map  $\Psi$  does not need to be one-to-one.  
• For  $\ell \geq n$  and  $\ell' \geq m$ , the block map  $\Psi$  can be defined as

$$\Psi : \beta_{\ell}^{\mathcal{A}, \mathcal{A}'} \rightarrow \beta_{\ell'}^{\mathcal{C}, \mathcal{C}'}$$

with  $\Psi(x_{[i, i+\ell-1]}) = y_{[i, i+\ell'-1]}$ . According to the indices, the letters in  $y_{[i, i+\ell'-1]}$  are chosen from  $\mathcal{C} \cup \mathcal{C}'$ .




FIGURE 2. This graph represents a zip shift space  $\Sigma_{\mathcal{A},\mathcal{A}'}$  with  $\mathcal{A}' = \{1, 2, 3, 4, 5, 6\}$  and  $\mathcal{A} = \{a, b, c, d, e\}$ . Walking in direction of edges and considering the numbers on edges, we get entries with non-negative indices and walking in opposite direction of edges and picking the letters written on edges, we get the entries with negative indices of a point  $x \in \Sigma_{\mathcal{A},\mathcal{A}'}$ .

### 3. PRE-IMAGES OF POINTS UNDER ZIP SHIFT MAP

Let  $\Sigma \subseteq \Sigma_{\mathcal{A},\mathcal{A}'}$  be an SFT or strictly sofic space and  $\sigma : \Sigma \rightarrow \Sigma$  be a zip shift map over  $\Sigma$ . For a point  $x \in \Sigma$ , we aim to find the set of all pre-images of it under  $\sigma$ . To do this, we construct a labeled graph called “backward labeled graph” which shows the backward dynamics of the points in  $\Sigma_{\mathcal{A},\mathcal{A}'}$ .

Let  $\varphi_n : \beta_n^{\mathcal{A}'} \rightarrow \mathcal{A}$  be the transition map. If  $a_i \in \mathcal{A}$ ,  $a'_i \in \mathcal{A}'$  and

$$x = (\dots, a_{-2}, a_{-1}; a'_0, a'_1, a'_2, \dots) \in \Sigma, \quad (3.1)$$

then

$$\sigma(x) = (\dots, a_{-2}, a_{-1}, a; a'_1, a'_2, \dots),$$

where  $a = \varphi_n(a'_0 a'_1 \dots a'_{n-1}) \in \mathcal{A}$ . We let

$$\sigma^{-1}(x) = (\dots, a_{-2}; a', a'_0, a'_1, a'_2, \dots), \quad (3.2)$$

where  $a'$  is an “appropriate” pre-image of  $a_{-1} a'_0 a'_1 \dots a'_{n-2}$  under  $\varphi_n$ . Sometimes, deciding which pre-image is appropriate is not easy at all.

**Example 3.1.** Consider the simple graph in Figure 2 with  $\mathcal{A}' = \{1, 2, 3, 4, 5, 6\}$  and  $\mathcal{A} = \{a, b, c, d, e\}$ . Set  $\varphi_1(1) = a$ ,  $\varphi_1(2) = \varphi_1(3) = b$ ,  $\varphi_1(4) = c$ ,  $\varphi_1(5) = d$  and  $\varphi_1(6) = e$ . The space  $\Sigma_{\mathcal{A},\mathcal{A}'}$  is sofic. Let

$$x = (\dots, b, b, b; 1, 6, \dots) \in \Sigma_{\mathcal{A},\mathcal{A}'}$$

Since  $\varphi_1^{-1}(b) = \{2, 3\}$ , we cannot decide which one to choose in order to determine  $\sigma^{-1}(x)$ . In fact, we should trace back the sequece of  $b$ 's in negative indices up to an entry  $x_{-k}$  which is  $c$  or  $e$ . If  $x_{-k} = c$ , then  $y = \sigma^{-1}(x) = (\dots, b, b, 2, 1, 6, \dots)$ , and if  $x_{-k} = e$ , then  $z = \sigma^{-1}(x) = (\dots, b, b, 3, 1, 6, \dots)$ . If there does not exist  $k$  with  $x_{-k} = c$  or  $e$ , then both points  $y$  and  $z$  are pre-images of  $x$ .

In sequel we want to find the pre-images of a given point  $x \in \Sigma_{\mathcal{A},\mathcal{A}'}$  through the backward labeled graph associated to  $\Sigma_{\mathcal{A},\mathcal{A}'}$ . We consider two cases when  $\Sigma_{\mathcal{A}'}$  is an  $M$ -step Markov space or when  $\Sigma_{\mathcal{A}'}$  is a strictly sofic space.

**3.1. Constructing the labeled and backward labeled graphs for sofic zip shift spaces.** In [11],  $M$ -step Markov spaces and sofic spaces are represented by graphs. A vertex graph or an edge graph associates to the first space and a labeled graph represents the second one. Here, we correspond a graph to the sofic zip shift space which carries labels both on its vertices and edges. Call it *labeled graph* for zip shift space. The *backward labeled graph* can be obtained from the labeled graph by reversing the direction of edges. It is easy to construct the labeled graph when  $\Sigma_{\mathcal{A}'}$  is an  $M$ -step Markov space. If  $\Sigma_{\mathcal{A}'}$  is strictly sofic, some modifications will be made.

**Case 1.** Let  $\Sigma_{\mathcal{A}'}$  be an  $M$ -step Markov space. If  $n = M = 1$  ( $n$  is due to map  $\varphi_n$ ), then take the edge graph representing  $\Sigma_{\mathcal{A}'}$ . Here, the labels on the edges are taken from  $\mathcal{A}'$ . If the label of an edge is  $a'$ , then replace it with  $a'/\varphi_1(a')$ .

If  $n = 1$ , set  $\mathcal{V}(H) = \beta_M^{\mathcal{A}'}$  as the vertex set of the labeled graph. There is a directed edge  $e$  from  $v_1 = a'_1 a'_2 \dots a'_M$  to  $v_2 = a'_2 \dots a'_M a'_{M+1}$  if  $a'_1 a'_2 \dots a'_M a'_{M+1} \in \beta_{M+1}^{\mathcal{A}'}$ . Label this edge by

$$a'_{M+1}/\varphi_1(a'_1).$$

If there are  $m$  letters  $b'_1, \dots, b'_m \in \mathcal{A}'$  such that  $b'_j a'_2 a'_3 \dots a'_{M+1} \in \beta_{M+1}^{\mathcal{A}'}$ ,  $1 \leq j \leq m$ , then  $m$  edges enters to the vertex  $a'_2 a'_3 \dots a'_{M+1}$  with labels  $a'_{M+1}/\varphi_1(b'_j)$ . Note that the letters  $b'_j$  are different, however the sofic parts  $\varphi_1(b'_j)$  may be the same for some edges.

If  $n \geq 2$ , let  $\ell = \max\{n - 1, M\}$ . Set  $\mathcal{V}(\mathcal{H}) = \beta_\ell^{\mathcal{A}'}$ . There exists an edge  $e$  from  $v_1 = a'_1 \dots a'_\ell$  to  $v_2 = a'_2 a'_3 \dots a'_{\ell+1}$  if  $a'_1 a'_2 \dots a'_\ell a'_{\ell+1} \in \beta_{\ell+1}^{\mathcal{A}'}$ . Label this edge by  $a'_{\ell+1}/\varphi_n(a'_1 \dots a'_n)$ .

**Case 2.** Let  $\Sigma_{\mathcal{A}'}$  be a strictly sofic shift space. Call a graph a *vertex-like graph* if it is a vertex graph such that some of the vertices carry the same label. See Figure 3.

**Lemma 3.2.** *Let  $X$  be a sofic shift space. There exists a directed vertex-like graph  $H$  which represents  $X$ .*

*Proof.* Let  $G$  be a labeled graph representing  $X$ . Let  $e$  be an edge from vertex  $v'$  to  $w'$ . Denote by  $i(e)$  and  $t(e)$  the initial and terminal vertices of  $e$ , i.e.,  $i(e) = v'$  and  $t(e) = w'$ . The graph  $G$  might have at least two edges  $e_1$  and  $e_2$  with  $i(e_1) = i(e_2) = I$  and  $t(e_1) = t(e_2) = J$ . As described in Section 2.4, [11], we can split  $I$  into two vertices  $I_1$  and  $I_2$  such that  $i(e_1) = I_1$  and  $i(e_2) = I_2$ . For all such vertices we split initial vertices into two or more vertices until all of the edges in the new graph  $G'$  have different initial vertices or different terminal vertices. Then  $X_G = X_{G'}$  [11]. The graph  $G'$  contains some edges with the same label. If  $G'$  has  $k$  such edges, rename all edges with  $e_1$  to  $e_k$ . There exists a vertex graph conjugate to  $G'$  with vertices  $e_1$  to  $e_k$ . Again rename each  $e_i$  by the original label as in  $G'$ . This vertex-like graph  $H$  has some vertices with the same label and represents  $X$ .  $\square$

For  $n \geq 1$ , let  $\mathcal{H}'$  be the vertex-like graph with vertex set  $\mathcal{V}(\mathcal{H}') = \beta_\ell^{\mathcal{A}'}$ ,  $\ell \geq n - 1$  representing  $\Sigma_{\mathcal{A}'}$  (for  $n = 1$ , take  $\ell \geq 1$  and). For  $n = 1$ , one can also consider the labeled graph of  $\Sigma_{\mathcal{A}'}$ . Suppose  $v_1$  and  $v_2$  are two vertices in  $\mathcal{G}'$  and  $e$  is an




FIGURE 3. Graphs (a), (b), (c), and (e) represent the labeled or vertex-like graphs for the even shift  $\Sigma_{\mathcal{A}'}$ . Graphs (d) and (f) represent the labeled graphs for  $\Sigma_{\mathcal{A},\mathcal{A}'}$  with the transition map described in Example 3.3.

edge from  $v_1$  to  $v_2$ . Let  $v_2$  end with letter  $a' \in \mathcal{A}'$ . Then label  $e$  by  $a' / \varphi_n(w_{[1,n]})$  where  $w = v_1 a'$ .

**Example 3.3.** Let  $\mathcal{A}' = \{0, 1\}$ . The space  $\Sigma_{\mathcal{A}'}$  with forbidden block set  $\mathcal{F} = \{10^{2n-1}1 : n \in \mathbb{N}\}$  is called even shift. Its labeled graph is shown in Figure 3 parts a and b. There is a condition on the number of symbols zero between two successive 1's. To show this space with a vertex-like graph, it suffices to duplicate the vertex with label zero. The new graph illustrated in Figure 3c, is a vertex-like graph for the even shift.

If the vertex set is  $\beta_2^{\mathcal{A}'}$ , then another representation of the even shift by a vertex-like graph is as in Figure 3e. Let  $\mathcal{A} = \{a, b, c\}$  and  $\varphi_2(11) = a$ ,  $\varphi_2(10) = \varphi_2(01) = b$ , and  $\varphi_2(00) = c$ . Its labeled graphs are shown in Figure 3, parts d and f.

**3.2. Walking on graphs associated to a sofic space  $\Sigma_{\mathcal{A},\mathcal{A}'}$ .** For a sofic space  $\Sigma_{\mathcal{A},\mathcal{A}'}$  with transition map  $\varphi_n : \beta_n^{\mathcal{A}'} \rightarrow \mathcal{A}$ , let  $\mathcal{H}$  and  $\mathcal{G}$  be a labeled and a backward labeled graph with vertex set  $\beta_\ell^{\mathcal{A}'}$  for  $\ell \geq n - 1$  respectively. Let  $e_j$  connect the vertex  $v_j$  to  $v_{j+1}$ , carrying the label  $a'_j/a_j$ . For a label  $a'_j/a_j$  with  $a'_j \in \mathcal{A}'$  and  $a_j \in \mathcal{A}$ , we call  $a'_j$  and  $a_j$  the *first* and the *sofic part* of the label. Walks or paths on these graphs can have finite or infinite length.

First consider the graph  $\mathcal{H}$ . Any finite segment  $\pi = e_i \cdots e_{i+m}$  of a path in the direction of edges is a concatenation of edges  $e_j$  with  $t(e_j) = i(e_{j+1})$  for

$0 \leq i \leq j \leq i + m - 1$ . This path corresponds to a block  $a'_i \dots a'_{i+m} \in \beta_{m+1}^{\mathcal{A}'}$  where  $a'_i$  is the first label of  $e_i$ .

The graph  $\mathcal{G}$  is constructed from  $\mathcal{H}$  by reversing the direction of edges without changing the labels of edges. Walking in opposite direction of edges in  $\mathcal{H}$  or in the direction of edges in  $\mathcal{G}$ , any finite segment  $\pi = e_{i+m} \dots e_i$  of a path is a concatenation of edges  $e_j$  with  $t(e_j) = i(e_{j+1})$  for  $i, m > 0$  and  $i \leq j \leq i + m - 1$ . This path starts from  $e_i$  and ends at  $e_{i+m}$ . The indices are sorted this way to show their dependency to the entries  $(\dots, a_{-(i+m)}, \dots, a_{-(i+1)}, a_{-i}, \dots; \dots)$ . This path corresponds to a word  $a_{-(i+m)} \dots a_{-i} \in \beta_{m+1}^{\mathcal{A}}$  where  $a_{-i}$  is the sofic label of  $e_i$ . The block  $L := a_{-(i+m)} \dots a_{-i}$  is called the *sofic label* of path  $\pi$  and is denoted by  $\mathcal{L}(\pi)$ . Denote by  $i(\pi)$  and  $t(\pi)$  the initial and terminal vertices of  $\pi$ . For both above paths  $\pi$ ,  $i(\pi) = e_i$  and  $t(\pi) = e_{i+m}$ .

**Definition 3.4.** Let  $\mathcal{G}$  be a labeled backward graph for the sofic space  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ .

- Let  $v'$  be a vertex of  $\mathcal{G}$  and  $L$  be an admissible sofic label. A path  $\pi = e_d e_{d-1} \dots e_1$  emanating from  $v'$  has *delay*  $d$  if  $d$  is the smallest natural number and  $\pi$  is the unique path such that  $\mathcal{L}(\pi) = L$ .
- Let  $v'$  be a vertex of  $\mathcal{G}$  and let  $L$  be an admissible sofic label of paths  $\pi_1, \dots, \pi_m$ ,  $m \geq 2$  all of length  $\ell$  with  $i(\pi_1) = \dots = i(\pi_m) = v'$  and  $t(\pi_1) = \dots = t(\pi_m)$ . Call these paths *left-closing* paths of length  $\ell$  if  $\ell$  is the smallest natural number and  $\pi_j$ 's are the only paths emanating from  $v'$  having sofic label  $L$  (see Figures 4 and 5).
- A sofic label  $L$  is called  $d_L$ -*distinguishable* if the number  $d$  and the label  $L$  determines a unique vertex  $v'$  and a unique path  $\pi$  such that  $i(\pi) = v'$  and  $\pi$  has delay  $d$ .
- A sofic label  $L$  is called  $L$ -*distinguishable* if there exist a unique vertex  $v'$  and some paths  $\xi_j = \dots e_3^j e_2^j e_1^j$ ,  $1 \leq j \leq t$  where  $i(\xi_j) = v'$  and  $\mathcal{L}(\xi_j) = L$ .
- Suppose there are vertices  $v'_1$  to  $v'_s$ ,  $s \geq 2$  with label  $a'_0 \dots a'_{n-1}$  and for sofic label  $L$ , each of these vertices is the initial vertex of at least one of the paths  $\pi_j$ ,  $1 \leq j \leq m$ ,  $m \geq 2$  carrying the label  $L$ . If all of these paths have the same endpoint  $w'$ , then they are called *sofic-left-closing*.

Let  $\mathcal{G}$  be a backward labeled graph for  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ . To obtain the pre-images of a point  $x = (\dots, a_{-2}, a_{-1}; a'_0, a'_1, \dots)$ , first find the vertex  $v'$  or vertices  $v'_1$  to  $v'_s$  which determine the starting state according to the sequence  $(a'_0, a'_1, a'_2, \dots)$ . Suppose there are more than one edge with label  $a_{-1}$  emanating from such vertices. It might happen that only some of the paths starting with these edges have label  $(\dots, a_{-2}, a_{-1})$  to form the pre-images of point  $x$ . In some cases it is very hard to verify if a finite part of  $(\dots, a_{-2}, a_{-1})$  suffices to decide whether the paths carrying such label produce the pre-images of  $x$  or not.

**Theorem 3.5.** Let  $\varphi_n : \beta_n^{\mathcal{A}'} \rightarrow \mathcal{A}$  be an onto factor map. If  $\Sigma_{\mathcal{A}'}$  is a sofic space, then the pre-images of a point  $x \in \Sigma_{\mathcal{A}, \mathcal{A}'}$  under  $\sigma$  can be determined by its backward labeled graph according to the following rules.

- i) If  $\Sigma_{\mathcal{A}'}$  is an  $M$ -step Markov space, and a sofic label  $L$  occurs with delay  $d$  or is the label of left-closing paths, then all of these paths form the

pre-images of  $x$ . Otherwise, all paths with sofic label  $\mathcal{L} = \dots a_{-3} a_{-2} a_{-1}$  produce the pre-images of  $x$ .

- ii) If  $\Sigma_{\mathcal{A}'}$  is a strictly sofic space, and a sofic label  $L$  is  $d_L$ -distinguishable,  $L$ -distinguishable or is the label of sofic-left-closing paths, then all of these paths form the pre-images of  $x$ . Otherwise, all paths with sofic label  $\mathcal{L} = \dots a_{-3} a_{-2} a_{-1}$  produce the pre-images of  $x$ .

*Remark 3.6.* The proof of Theorem 3.5 can be used to determine the pre-images of points in a one-sided sofic shift space with some adjustments.

*Proof.* Let

$$K_n = \{w' \in \beta_n^{\mathcal{A}'} : \exists v' \in \beta_n^{\mathcal{A}'} : \varphi_n(w') = \varphi_n(v')\},$$

where  $n$  is due to the transition map. For  $w' = w'_1 \dots w'_n \in K_n$ , let

$$R_n(w') = \{v' = v'_1 \dots v'_n \in K_n \setminus \{w'\} : w'_2 \dots w'_n = v'_2 \dots v'_n\}$$

and  $R(n) = \cup_{w' \in K_n} R_n(w')$ . If  $\varphi_n$  is one-to-one, then  $K_n$  and  $R(n)$  are empty sets and  $\sigma$  is one-to-one on  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ . Suppose  $\varphi_n$  is not one-to-one. So,  $K_n \neq \emptyset$ . Suppose  $R(n) = \emptyset$ . In this case  $\sigma$  is again invertible. To see this let  $x$  be as in (3.1) and for a letter  $a' \in \mathcal{A}'$ ,  $w' = a' a'_0 a'_1 \dots a'_{n-2} \in K_n$ . Since  $R(n) = \emptyset$ , only for the word  $w'$ , we have  $\varphi_n(w') = a_{-1}$  and so,  $\sigma^{-1}(x)$  is as in (3.2). If  $R(n) \neq \emptyset$ , then  $\sigma$  might be invertible or not as we discuss in sequel.

First let  $\Sigma_{\mathcal{A}'}$  be an  $M$ -step Markov space and  $\mathcal{G}$  be its backward labeled graph. Suppose  $n = 1$ . Without loss of generality, suppose for the vertex  $v' = a'_0 a'_1 \dots a'_{M-1} \in \mathcal{V}(\mathcal{G}) = \beta_M^{\mathcal{A}'}$ , there are at least two edges  $e_1$  and  $e_2$  emanating from  $v'$  with  $\mathcal{L}(e_1) = \mathcal{L}(e_2) = a_{-1}$ . Then one of the following 3 cases happen.

- (1) There exists  $d \geq 1$  and a path  $\pi = e_d \dots e_1$  with sofic label  $\mathcal{L}(\pi) = a_{-d} \dots a_{-1}$ , such that  $i(\pi) = v'$  and  $\pi$  has delay  $d$ .
- (2) There are  $m \geq 2$  left-closing paths  $\pi_j = e_\ell^j e_{\ell-1}^j \dots e_2^j e_1^j$ ,  $1 \leq j \leq m$  of length  $\ell$  with  $i(\pi_j) = v'$ ,  $t(\pi_j) = w'$  and  $\mathcal{L}(\pi_j) = a_{-d} \dots a_{-1}$ .
- (3) There are infinite length paths  $\xi_j = \dots e_3^j e_2^j e_1^j$ ,  $1 \leq j \leq t$  carrying the label  $\mathcal{L}(\dots e_3^j e_2^j e_1^j) = \dots a_{-3} a_{-2} a_{-1}$ , where neither of them happen with delay  $d$  nor is a left-closing path with some length  $\ell$ .

Any vertex  $v'$  admits only one of these cases.

- If Case (1) happens for the vertex  $v'$  and a path  $\pi = e_{-d} \dots e_{-1}$  which starts from  $v'$  with delay  $d$ , then  $\sigma$  is invertible on  $x$  and for  $1 \leq k \leq d$ ,

$$\sigma^{-k}(x) = (\dots, a_{-k-1}; a'_{-k}, a'_{-k+1}, \dots), \quad (3.3)$$

where  $a'_{-k-1}$  is the first part of the label of edge  $e_k$ .

- If  $v'$  satisfies the second case, then  $x$  has  $m$  pre-images along the paths  $\pi_1$  to  $\pi_m$ , i.e., the pre-images of  $x$  are the set of points

$$(\dots, a_{-2}; (a^j)'_{-1}, a'_0, \dots), \quad (3.4)$$

where  $(a^j)'_{-1}$  is the first part of the label of edge  $e_1^j$ . Also, for  $1 \leq k \leq \ell$  and  $1 \leq j \leq m$ ,  $\sigma^{-k}(x)$  is the set of points

$$(\dots, a_{-k-1}; (a^j)'_{-k}, (a^j)'_{-k+1}, \dots, (a^j)'_{-1}, a_0, a_1, \dots), \quad (3.5)$$




FIGURE 4. By definition, the paths ending at  $w'$ , form left-closing paths and the smaller groups of paths ending at  $v'_1$ ,  $v'_2$  or  $v'_3$  are not left-closing.




FIGURE 5. We take all the paths associated to the numbers (1), (2) and (3) to have sofic label  $\dots a_{-3} a_{-2} a_{-1}$ . For number (4), the sofic label of paths are  $a_{-k} \dots a_{-3} a_{-2}, a_{-1}$  up to vertex  $v'_1$ , but the next sofic label is “a” instead of being  $a_{-k-1}$ . This means that none of the paths in number (4) produces the pre-images of  $x$ . In this figure there are not any left-closing paths.

where  $(a^j)'_{-k}$  is the first part of the label of edge  $e_k^j$ . In Figure 4, there are some kinds of left-closing paths.

- If the third case happens, then all paths  $\pi_j = \dots e_3^j e_2^j e_1^j$  with  $i(\pi_j) = v'$  and  $\mathcal{L}(\dots e_3^j e_2^j e_1^j) = \dots a_{-3} a_{-2} a_{-1}$  determine the pre-images of  $x$ . Some cases are shown in Figure 5.

For  $n > 1$ , the argument is similar and one needs to consider the backward labeled graph for case  $n > 1$ . Note that the length of the vertices in  $\mathcal{G}$  should be greater than  $n - 1$  and if so, their lengths effect neither on the sets  $K_n$  and  $R_n$  nor on the Definition 3.4.

Now suppose  $\Sigma_{\mathcal{A}'}$  is a strictly sofic space and  $\mathcal{G}$  is its backward labeled graph as described in Subsection 3.1. Let  $\mathcal{V}(\mathcal{G}') = \beta_n^{\mathcal{A}'}$ . Choose the vertex with label  $v' = a'_0 \dots a'_{n-1}$ . If there is only one vertex with this label, then the argument is similar to the Markov case.

Suppose there are vertices  $v'_1$  to  $v'_s$  with label  $a'_0 \dots a'_{n-1}$  such that starting from these vertices and walking on  $\mathcal{G}$ , the first label of the edges is  $a'_0 a'_1 a'_2 \dots$

- If there exists  $d$  such that the sofic label  $L = a_{-d} \dots a_{-1}$  is  $d_L$ -distinguishable or if the label  $L = \dots a_{-2} a_{-1}$  is  $L$ -distinguishable, then there exists a unique path  $\pi$  with label  $L$  and a unique  $t, 1 \leq t \leq s$ , such that  $i(\pi) = v'_t$



FIGURE 6. The rectangles  $H_a$  and  $H_b$  are formed from the first iterate of  $f$ . The rectangles  $H_{aa}$ ,  $H_{ab}$ ,  $H_{ba}$  and  $H_{bb}$  are shown with labels  $aa$ ,  $ab$ ,  $ba$  and  $bb$ .

and the pre-images of  $x$  satisfy (3.3) (In  $L$ -distinguishable case,  $d \rightarrow \infty$ ). Let for  $1 \leq j \leq k$ ,  $t(\pi_j) = v'_j$ . Then take  $v'_j$  as  $v'$  and the argument is as the Markov case.

- Suppose for  $m \geq 2$  and  $1 \leq j \leq m$ , the paths  $\pi_j = e_\ell^j e_{\ell-1}^j \dots e_2^j e_1^j$  are sofic-left-closing,  $t(\pi_j) = w'$  and  $i(\pi_j) \in \{v'_1, \dots, v'_s\}$ . Then all such paths produce pre-images of  $x$  and they satisfy (3.4) and (3.5). Again take  $w'$  as  $v'$  and the argument is as the Markov case.
- If there are paths  $\pi_j = \dots e_3^j e_2^j e_1^j$  with  $i(\pi_j) \in \{v'_1, \dots, v'_s\}$  and  $\mathcal{L}(\dots e_3^j e_2^j e_1^j) = \dots a_{-3} a_{-2} a_{-1}$ , then all of them determine the pre-images of  $x$ .

□

#### 4. CONSTRUCTION OF AN $N$ -TO-1 ENDOMORPHISM HORSESHOE

Let  $\mathbf{S}$  be the  $[0, 1] \times [0, 1]$  square. We aim to construct an  $N$ -to-1 horseshoe map  $f$  on a subset of  $\mathbf{S}$  according to the following description. Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  linearly expand  $\mathbf{S}$  in the horizontal direction by a factor  $\delta$  greater than  $2N$ ,  $N \in \mathbb{N}$  and linearly contract it in the vertical direction by a factor  $\delta'$  less than  $\frac{1}{N}$  such that  $\delta\delta' = N$ . If  $\delta = 2N + \epsilon$ , then taking modulo by  $2 + \frac{\epsilon}{N}$  or folding the rectangle successively,  $N$  copies of the rectangles  $[0, \frac{\delta}{N}] \times [0, \delta']$  overlay. Now bend these rectangles back inside  $\mathbf{S}$ . Obviously,  $f$  is  $N$ -to-1 on  $\mathbf{S}$ . Note that for any  $k \geq 2$ ,  $f^k(\mathbf{S})$  is **one** horizontal “snake-like” strip lying inside  $f^{k-1}(\mathbf{S})$ . Since  $f$  shrinks  $\mathbf{S}$  in vertical direction by  $\delta'$ ,  $\lim_{k \rightarrow +\infty} f^k(\mathbf{S})$  tends to an infinite-length horizontal curve  $c^u$ .

Inside the square,  $f(\mathbf{S}) \cap \mathbf{S}$  is consisted of two horizontal rectangles named  $H_a$  and  $H_b$ . The map  $f$  has the Markov property, i.e., the iterates of  $f$  on  $\mathbf{S}$  generate thin rectangles such that  $f^k(\mathbf{S}) \cap \mathbf{S} \subset f^{k-1}(\mathbf{S}) \cap \mathbf{S}$ ,  $k \geq 1$ . See Figure 6 for  $N = 2$  and  $k = 1, 2$ . In each iterate,  $f$  shrinks the vertical side by factor  $\delta'$ . Therefore, a sequence of nested horizontal rectangles  $f^k(\mathbf{S})$  tends to a horizontal line segment as  $k \rightarrow \infty$ . This implies that

$$\Lambda_H = \{(x, y) \in \mathbf{S} : (x, y) \in \lim_{k \rightarrow \infty} f^k(\mathbf{S}) \cap \mathbf{S}\} = [0, 1] \times C_2,$$

where  $C_2$  is isomorphic to the Cantor middle third set. Any point in  $\Lambda_H$  remains in  $\Lambda_H$  under the iterations of  $f$ .

The pre-image of  $\mathbf{S}$  under  $f$  is consisted of  $N$  thin vertical rectangles. The place of such rectangles depends on map  $f$ . In Figure 7, since  $f$  takes modulo by  $2 + \frac{\epsilon}{N}$ , the vertical rectangles in backward iterate are placed at equal distances. Each






FIGURE 7. The first backward iterate of the horseshoe map  $f$  for  $N = 2$ .

of these rectangles are the pre-images of  $\mathbf{S}$  which are linearly stretched by factor  $1/\delta'$  and linearly shrank by factor  $1/\delta$  in the vertical and horizontal directions respectively. Bending the rectangles back inside the square,  $f^{-1}(\mathbf{S})$  is consisted of  $N$  bended strips. Then  $f^{-2}(\mathbf{S})$  produces  $N$  vertical snake-like strips inside each of existing strips. This means that  $f^{-k}(\mathbf{S})$  tends to infinitely many vertical snake-like curves  $c^s$  as  $k \rightarrow \infty$  all of them lying inside  $f^{-1}(\mathbf{S})$ .

Also,  $f^{-1}(\mathbf{S}) \cap \mathbf{S}$  produces  $2N$  vertical rectangles  $V_{0^1}, V_{1^1}, V_{0^2}, V_{1^2}, \dots, V_{0^N}, V_{1^N}$ . See Figure 7 for  $N = 2$ . For any  $i$ ,  $1 \leq i \leq N$ ,

$$f(V_{0^i}) = H_a, \quad f(V_{1^i}) = H_b.$$

See Figure 8. For any  $k \geq 2$ ,  $f^{-k}(\mathbf{S}) \cap \mathbf{S} \subset f^{-k+1}(\mathbf{S}) \cap \mathbf{S}$  and  $\lim_{k \rightarrow \infty} f^{-k}(\mathbf{S})$  tends to a vertical line segment in  $\mathbf{S}$ . More precisely,

$$\Lambda_V = \{(x, y) \in \mathbf{S} : (x, y) \in \lim_{k \rightarrow \infty} f^{-k}(\mathbf{S}) \cap \mathbf{S}\} = C_1 \times [0, 1],$$

where  $C_1$  is isomorphic to a Cantor set. Now let

$$\Lambda = \Lambda_H \cap \Lambda_V = C_1 \times C_2, \tag{4.1}$$

and define the restriction of  $f$  to  $\Lambda$ , the horseshoe map. The set  $\Lambda$  is the trapping region for the horseshoe map  $f : \Lambda \rightarrow \Lambda$  and remains invariant under iterations of  $f$  and  $f^{-1}$ .

**4.1. A conjugate zip shift map to the  $N$ -to-1 horseshoe.** To represent this system in symbolic dynamics, we correspond a bi-infinite sequence of determined symbols to any point in  $\Lambda$ . For this purpose, the following observation would be useful. Let  $Z = \{0^1, 0^2, \dots, 0^N\}$ ,  $O = \{1^1, 1^2, \dots, 1^N\}$  and  $\mathcal{A}' = Z \cup O$ . Fix a vertical rectangle  $V_{n_0}$ ,  $n_0 \in \mathcal{A}'$ . Let  $V_{n_0, n_1} = \{(x, y) \in V_{n_0} : f(x, y) \in V_{n_1}, n_1 \in \mathcal{A}'\} = V_{n_0} \cap f^{-1}(V_{n_1})$ . Inductively, define

$$\begin{aligned} V_{n_0, n_1, \dots, n_k} &= \{(x, y) \in V_{n_0, \dots, n_{k-1}} : f^{n_k}(x, y) \in V_{n_k}, n_i \in \mathcal{A}'\} \\ &= V_{n_0, \dots, n_{k-1}} \cap f^{-n_k}(V_{n_k}). \end{aligned} \tag{4.2}$$




FIGURE 8. Let  $\mathcal{A}' = \{0, 1, 0', 1'\}$ ,  $\mathcal{A} = \{a, b\}$ ,  $\varphi_1(0) = \varphi_1(0') = a$  and  $\varphi_1(1) = \varphi_1(1') = b$ . First the upper left unit square is mapped to the upper right rectangle with horizontal and vertical sides  $\delta$  and  $2/\delta$  where  $\delta = 4 + \epsilon$ . Cutting the rectangle from the dashed line and taking modulo by  $2 + \frac{\epsilon}{2}$ , the rectangles with labels 0 and  $0'$  overlay as the ones with labels 1 and  $1'$ . This gives the lower left rectangle. Bending the obtained rectangle back to itself the vertical rectangles are mapped to horizontal rectangle.

The vertical region  $V_{n_0, n_1, \dots, n_k}$  is the intersection of nested rectangles and tend to the vertical line segment  $v_{x_0} = \lim_{k \rightarrow \infty} V_{n_0, n_1, \dots, n_k}$  passing through the point  $x_0$  on the  $x$ -axis. Obviously,  $v_{x_0} \subset c^s$ .

Now consider a horizontal rectangle  $H_{n_{-1}}$ ,  $n_{-1} \in \mathcal{A} = \{a, b\}$ . Set  $H_{n_{-2}, n_{-1}} = \{(x, y) \in H_{n_{-1}} : f^{-1}(x, y) \in H_{n_{-2}}\} = H_{n_{-1}} \cap f(H_{n_{-2}})$ . Inductively define

$$\begin{aligned} H_{n_{-k}, \dots, n_{-1}} &= \{(x, y) \in H_{n_{-(k-1)}, \dots, n_{-1}}, f^{-n_k}(x, y) \in H_{n_{-k}}\} \\ &= H_{n_{-(k-1)}, \dots, n_{-1}} \cap f^{n_k}(H_{n_{-k}}). \end{aligned} \quad (4.3)$$

Obviously,  $\lim_{k \rightarrow \infty} H_{n_{-k}, \dots, n_{-1}}$  tends to the horizontal line segment  $h_{y_0}$  passing through the point  $y_0$  on the  $y$ -axis. Clearly,  $h_{y_0} \subset c^u$ .

Therefore,

$$\left( \lim_{k \rightarrow \infty} V_{n_0, n_1, \dots, n_k} \right) \cap \left( \lim_{k \rightarrow \infty} H_{n_{-k}, \dots, n_{-1}} \right) \quad (4.4)$$

uniquely determines the point  $(x_0, y_0) = v_{x_0} \cap h_{y_0}$ , where  $n_i \in \mathcal{A}'$  for  $i \geq 0$  and  $n_i \in \mathcal{A}$  for  $i \leq -1$ .

The set

$$\Sigma_{\mathcal{A}, \mathcal{A}'} = \{(n_k)_{k \in \mathbb{Z}} : \left( \lim_{k \rightarrow \infty} V_{n_0, n_1, \dots, n_k} \right) \cap \left( \lim_{k \rightarrow \infty} H_{n_{-k}, \dots, n_{-1}} \right) \neq \emptyset\} \quad (4.5)$$

is a zip shift space over alphabets  $\mathcal{A}$  and  $\mathcal{A}'$ . We correspond the bi-infinite sequence  $(n_k)_{k \in \mathbb{Z}}$  to a point  $(x_0, y_0) \in \Lambda$  with respect to the indices obtained in equation (4.4).

Let  $\varsigma : \Lambda \rightarrow \Sigma_{\mathcal{A}, \mathcal{A}'}$  be the *code* map which maps each point in  $\Lambda$  to its unique sequence of symbols in  $\mathcal{A} \cup \mathcal{A}'$ . For  $z \in Z$  and  $o \in O$ , define  $\varphi_1(z) = a$  and  $\varphi_1(o) = b$ . Since  $\varphi_1$  is not one-to-one,  $\varphi_1^{-1}(a) \in Z$  and  $\varphi_1^{-1}b \in O$ . According to equations (4.2) and (5.5),  $\varsigma((x_0, y_0)) = (\dots, n_{-2}, n_{-1}; n_0, n_1, \dots)$  means that for  $i \in \mathbb{N}$ ,

$$f^{-i}(x_0, y_0) \cap H_{n_{-i}} \neq \emptyset, (x_0, y_0) \in V_{n_0}, f^i(x_0, y_0) \in V_{n_i} \quad (4.6)$$

where  $n_i \in \mathcal{A}'$  for  $i \geq 0$  and  $n_i \in \mathcal{A}$  for  $i \leq -1$ . Equation (4.4) indicates that for a point  $(x_0, y_0) \in \Lambda$  all the points on  $v_{x_0} \cap \Lambda$  have the same forward code. But the points on  $h_{y_0} \cap \Lambda$  have different backward codes due to the possibility of different choices of pre-images of  $x_0$ .

The zip shift map on  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  represents the dynamics of  $f$  on  $\Lambda$  and also illustrates the  $n$ -to-1 nature of  $f$ . Fix the point  $(x_0, y_0) \in \Lambda$ . Denote by  $\{(x_{-1}^i, y_{-1})\}$  the set of  $n$  pre-images of  $(x_0, y_0)$  under  $f$ . Each of the points  $(x_{-1}^i, y_{-1})$  lies on the intersection of  $h_{y_{-1}}$  with  $v_{x_{-1}^i}$ . Any segment  $v_{x_{-1}^i}$  is a subset of one of the vertical rectangles  $\{V_0^1, \dots, V_0^n\}$  or  $\{V_1^1, \dots, V_1^n\}$ . Since the point  $(x_0, y_0)$  has  $n$  pre-images, there are  $n$  possible choices for the place  $n_{-1}$  in the corresponding sequence. Selecting a special branch  $(x_{-1}^i, y_{-1})$  of pre-images of  $(x_0, y_0)$ ,  $n_{-1}$  will be  $0_i$  or  $1_i$  with respect to the index of the vertical rectangle  $V_0^i$  or  $V_1^i$  containing  $(x_{-1}^i, f^{-1}(y_0))$ .

**Theorem 4.1.** *The factor code map  $\varsigma : \Lambda \rightarrow \Sigma_{\mathcal{A}, \mathcal{A}'}$  is a homeomorphism.*

*Proof.* By using statements 4.4 and 4.5 any sequence  $(\dots, s_{-1}; s_0, s_1, s_2, \dots) \in \Sigma_{\mathcal{A}, \mathcal{A}'}$  represents the unique point on the intersection of  $v_{x_0} \cap h_{y_0}$  and any such point has a unique coding via  $\varsigma$  which shows that  $\varsigma$  is one-to-one and onto.

To see the continuity of  $\varsigma$ , let  $\{x_n\}_{i=1}^{+\infty}$  be any sequence of points in  $\Lambda$  convergent to some  $x \in \Lambda$ . By definition of  $\varsigma$ , it is obvious that  $\{\varsigma(x_n)\}_{i=1}^{+\infty}$  is convergent to  $\varsigma(x)$ . A similar argument shows that  $\varsigma^{-1}$  is continuous.  $\square$

**Theorem 4.2.** *The maps  $f : \Lambda \rightarrow \Lambda$  and  $\sigma : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{A}, \mathcal{A}'}$  are topologically conjugate via the code map  $\varsigma$ .*

*Proof.* According to the diagram

$$\begin{array}{ccc} \Lambda & \xrightarrow{f} & \Lambda \\ \varsigma \downarrow & & \downarrow \varsigma \\ \Sigma_{\mathcal{A}, \mathcal{A}'} & \xrightarrow{\sigma} & \Sigma_{\mathcal{A}, \mathcal{A}'} \end{array}$$

we need to show that  $f = \varsigma^{-1} \circ \sigma \circ \varsigma$ . Let  $(x_0, y_0) \in \Lambda$  be a point with

$$\varsigma((x_0, y_0)) = (\dots, s_{-2}, s_{-1}; s_0, s_1, \dots).$$

This means that

$$\dots, f^{-1}((x_0, y_0)) \cap H_{s_{-1}} \neq \emptyset, (x_0, y_0) \in V_{s_0}, f((x_0, y_0)) \in V_{s_1}, \dots$$

Equivalently,

$$\dots, f^{-1}((x_0, y_0)) \cap H_{\dots, s_{-2}, s_{-1}} \neq \emptyset, (x_0, y_0) \in V_{s_0, s_1, \dots}, f((x_0, y_0)) \in V_{s_1, s_2, \dots}, \dots$$




FIGURE 9. The way  $f$  maps vertical strips to horizontal ones.

According to the definition of  $V_{s_0, s_1, \dots}$  in (4.2),  $V_{s_0, s_1, \dots} = V_{s_0} \cap f^{-1}(V_{s_1, s_2, \dots})$ . To obtain the code of the point  $f((x_0, y_0))$ , apply  $f$  to all of the horizontal and vertical rectangles containing  $(x_0, y_0)$ . See Figure 9. Then

$$(x_1, y_1) = f((x_0, y_0)) \in f(V_{s_0, s_1, \dots}) = f(V_{s_0}) \cap V_{s_1, s_2, \dots} = H_{\varphi_1(s_0)} \cap V_{s_1, s_2, \dots}.$$

Thus, for  $k \geq 1$ ,  $f^{n_k}((x_1, y_1)) = f^{n_k+1}((x_0, y_0)) \in V_{s_k+1, s_k+2, \dots}$ . Therefore,  $\varsigma(f(x_0, y_0)) = (\dots s_{-2}, s_{-1}, \varphi_1(s_0); s_1, s_2 \dots)$  and  $f(x_0, y_0) = \varsigma^{-1} \circ \sigma \circ \varsigma(x_0, y_0)$  by Theorem 4.1.  $\square$

**Example 4.3.** Let  $f : \Lambda \rightarrow \Lambda$  be a 2 to 1 horseshoe map defined at the beginning of Section 4, where  $\Lambda = C_1 \times C_2$  as in (4.1) for  $n = 1$ . Let  $\mathcal{A}' = \{1, 2, 3, 4\}$ . The usual shift map relevant to this system is the one-sided shift map  $\sigma$  over  $\Sigma_{\mathcal{A}'}$ .

Suppose  $g(x) = 4x$  modulo 1. Again its corresponding shift map is  $\sigma$  over  $\Sigma_{\mathcal{A}'}$ . For both systems, the maps  $f$  and  $g$  are semi-conjugate to  $\sigma : \Sigma_{\mathcal{A}'} \rightarrow \Sigma_{\mathcal{A}'}$ . Therefore, the shift map cannot distinguish these systems. But if we code them into a zip shift space, then they will have different code spaces as follows.

Let  $\mathcal{A} = \{a, b\}$ ,  $\mathcal{B} = \{a\}$  and  $\mathcal{A}' = \{1, 2, 3, 4\}$  be three sets of alphabets. Then  $f$  is conjugate to  $\sigma_f : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{A}, \mathcal{A}'}$  with  $\varphi_1(1) = \varphi_1(3) = a$  and  $\varphi_1(2) = \varphi_1(4) = b$ . The map  $\sigma_f$  illustrates the 2 to 1 feature of  $f$ . The map  $g$  is conjugate to  $\sigma_g : \Sigma_{\mathcal{B}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{B}, \mathcal{A}'}$  with  $\varphi_1(1) = \varphi_1(2) = \varphi_1(3) = \varphi_1(4) = a$  which shows that  $g$  is a 4 to 1 map.

## 5. STABLE AND UNSTABLE SETS; PERIODIC, PRE-PERIODIC, HOMOCLINIC, HETEROCLINIC POINTS AND THEIR ORBITS

In this subsection we take  $n$  to be 1 in the definition of the transition map  $\varphi_n$  for simplicity. All of the definitions and theorems can be adjusted for  $n > 1$ . For points  $s = (\dots, s_{-2}, s_{-1}; s_0, s_1, \dots)$ ,  $t = (\dots, t_{-2}, t_{-1}; t_0, t_1, \dots) \in \Sigma_{\mathcal{A}, \mathcal{A}'}$ , let  $N(s, t) = \min\{|i|, s_i \neq t_i, i \in \mathbb{Z}\}$ ,  $N^+(s, t) = \min\{i, s_i \neq t_i, i \geq 0\}$  and  $N^-(s, t) = \min\{i+1, s_i \neq t_i, i < 0\}$ . Let  $d(s, t) = \frac{1}{2^{N(s, t)}}$ ,  $d^+(s, t) = \frac{1}{2^{N^+(s, t)}}$ ,  $d^-(s, t) = \frac{1}{2^{N^-(s, t)}}$ . Clearly,  $d$  is a metric and  $d^+$ ,  $d^-$  are pseudo-metrics on  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ . For a point  $s \in (\Sigma_{\mathcal{A}, \mathcal{A}'}, d)$ ,  $\mathcal{N}_r(s) = \{t \in \Sigma_{\mathcal{A}, \mathcal{A}'} : d(t, s) < r\}$  is a neighborhood of  $s$  with diameter  $2r$ . For  $i \in \mathbb{Z}$  and  $\ell \in \mathbb{N} \cup \{0\}$ , the set

$$s_i^\ell = [s_i, s_{i+1}, \dots, s_{i+\ell}] = \{(k_n)_{n \in \mathbb{Z}} \in \Sigma_{\mathcal{A}, \mathcal{A}'} : k_i = s_i, \dots, k_{i+\ell} = s_{i+\ell}\}$$

is a *cylinder*. These sets are open and close and form a basis for the topology on  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  induced by the metric  $d$ . It is easy to see that the space  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  is a Cantor set according to  $d$ .

In order to define the notions of stable and unstable sets we equip the zip shift space  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  with an equivalent metric  $d^\pm(s, t) = \frac{1}{2}(d^-(s, t) + d^+(s, t))$ . If  $\Omega_1$  and  $\Omega_2$  are two subsets of  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ , then define  $d^\pm(\Omega_1, \Omega_2) = \inf\{d^\pm(\omega_1, \omega_2) : \omega_1 \in \Omega_1, \omega_2 \in \Omega_2\}$ .

**Definition 5.1.** Let  $p = (\dots, p_{-2}, p_{-1}; p_0, p_1, \dots) \in \Sigma_{\mathcal{A}, \mathcal{A}'}$  be a periodic point. Then the *stable* and *unstable sets* of  $p$  are defined as

$$\begin{aligned} W_{\text{global}}^s(p) &= \{t \in \Sigma_{\mathcal{A}, \mathcal{A}'} : d^\pm((\sigma^n(p)), (\sigma^n(t))) = 0, n \rightarrow +\infty\} \\ &= \{t \in \Sigma_{\mathcal{A}, \mathcal{A}'} : \exists N'(t) \geq 0, \forall n \geq N'(t), t_n = p_n\}, \\ W_{\text{special}}^s(p) &= \{t \in W_{\text{global}}^s(p) : t_n = p_n, \forall n \geq 0\}, \end{aligned} \quad (5.1)$$

and

$$\begin{aligned} W_{\text{global}}^u(p) &= \{r \in \Sigma_{\mathcal{A}, \mathcal{A}'} : d^\pm((\sigma^{-n}(p)), (\sigma^{-n}(r))) = 0, n \rightarrow +\infty\}, \\ &= \{r \in \Sigma_{\mathcal{A}, \mathcal{A}'} : \exists N(r) > 0, \forall n \geq N(r), r_{-n} = p_{-n}\}, \\ W_{\text{special}}^u(p) &= \{r \in W_{\text{global}}^u(p) : r_{-n} = p_{-n}, \forall n > 0\}. \end{aligned} \quad (5.2)$$

*Remark 5.2.* In the definition of  $W_{\text{global}}^u(p)$ , the metric  $d^\pm$  is the distance between two sets of points since  $\sigma$  is a local homeomorphism. Therefore, for a point  $r \in \Sigma_{\mathcal{A}, \mathcal{A}'}$ , the condition

$$\{r \in \Sigma_{\mathcal{A}, \mathcal{A}'} : \exists N(r) > 0, \forall n \geq N(r), r_{-n} = p_{-n}\},$$

is a sufficient condition for  $r$  being in  $W_{\text{global}}^u(p)$ .

For any point  $t$ , define  $D^u(t) = (t_0, t_1, \dots)$  and  $D^s(t) = (\dots, t_{-2}, t_{-1})$ , the stable (forward) and unstable (backward) “*directions*” of  $t$  respectively. Let  $p = (\overline{p_0, p_1, \dots, p_{m-1}})$  be a periodic point with period  $m$  according to Definition 2.2. Note that all points  $p^i = \sigma^i(p)$  for  $0 \leq i \leq m-1$  are periodic points with period  $m$ . They can produce infinitely many stable directions. Howbeit, all of them share the same unstable direction. In fact, by Definition 5.1, the unstable direction of all points  $\sigma^{km}(p^i)$ ,  $k \geq 0$ , is the same and by equation (2.2) is  $D^u(p^i) = \{(\varphi_1(\overline{p_i, \dots, p_{m-i-1}})) : 0 \leq i \leq m-1\}$ . Denote by  $D^u(O(p))$  the union of the unstable directions of  $p^i$  for  $0 \leq i \leq m-1$  i.e.,

$$D^u(O(p)) := \bigcup_{i=0}^{m-1} D^u(p^i).$$

If  $\sigma$  fails to be one to one on  $p$ , then  $p$  has pre-images as

$$(\varphi_1(\overline{p_{m-1}, p_0, \dots, p_{m-2}}); \varphi_1^{-1}(\varphi_1(p_{m-1})), \overline{p_0, p_1, \dots, p_{m-1}}), \quad (5.3)$$

where obviously,  $p_{m-1} \in \varphi_1^{-1}(\varphi_1(p_{m-1}))$ .

**Definition 5.3.** Let  $p = (\overline{p_0, \dots, p_{m-1}})$  be a periodic point of period  $m$  and let  $O_n(p) = f^{-n}(O(p))$ . Then we call  $p' \in O_n(p^i)$  the pre-periodic point of  $p^i = \sigma^i(p)$  if there exists some  $k \geq 1$  such that  $\sigma^{km}(p') = p^i$  (See Figure 10).



Obviously, the homoclinic point  $x$  might have more than one homoclinic orbit to  $p$ . See Example 5.5. The similar procedure works to find heteroclinic orbits.

**5.1. Stable and unstable sets and periodic , homoclinic and heteroclinic orbits of the  $N$ -to-1 horseshoe.** In this subsection, some properties of  $f$  such as stable and unstable sets, periodic, pre-periodic, homoclinic, heteroclinic points and their orbits are verified. Here we take  $n$  to be 1 ( $n$  is due to  $\varphi_n$ ) for simplicity.

**5.1.1. Periodic points and their orbits.** Consider the periodic point  $P = (x_0, y_0)$  and its orbit  $(x_i, y_i) \in \Lambda$ ,  $0 \leq i \leq m - 1$  with least period  $m$ . Since  $f$  is an  $N$ -to-1 endomorphism, any point  $(x_i, y_i)$  has  $N$  pre-images. To have a periodic orbit, we take  $f^{-1}((x_i, y_i))$  to be  $(x_{i-1}, y_{i-1})$  for  $1 \leq i \leq m - 1$  and  $f^{-1}((x_0, y_0)) = (x_{m-1}, y_{m-1})$ . Other pre-images of  $f$  do not form a periodic orbit. Let  $\varsigma((x_0, y_0)) = p = (\overline{p_0, p_1, \dots, p_{m-1}})$ ,  $p_i \in \mathcal{A}'$ . Then for  $0 \leq i \leq m - 1$ ,  $\varsigma((x_i, y_i)) = p^i = (\overline{p_i, p_{i+1}, \dots, p_{i+m-1}})$  where the indices are taken modulo by  $m$ . Similarly,  $p^i$  has  $N$  pre-images.

Any point  $\varsigma(x_i, y_i) = (\dots, \varphi_1(p_{i-1}); p_i, \dots, p_{m-1}, p_0, p_1, \dots)$  has  $N$  pre-images as  $(\dots, \varphi_1(p_{i-2}); t, p_i, \dots, p_{m-1}, p_0, p_1, \dots)$  with  $t \in \varphi_1^{-1}(p_{i-1})$  for  $1 \leq i \leq m - 1$ . To have the periodic orbit select  $t$  to be  $p_{i-1}$ . Then

$$(\dots, \varphi_1(p_{i-2}); p_{i-1}, p_i, \dots, p_{m-1}, p_0, p_1, \dots)$$

coincides with the code of that pre-image of  $f^{-1}(x_i, y_i)$  which belongs to the set  $\{(x_0, y_0), \dots, (x_{m-1}, y_{m-1})\}$ . None of the other pre-images of  $p^i$  are periodic. Let  $\text{Per}_m(f) \subset \Lambda$  and  $\text{Per}_m(\sigma) \subset \Sigma_{\mathcal{A}, \mathcal{A}'}$  be the set of all periodic points with least period  $m$  under  $f$  and  $\sigma$  respectively. Obviously,  $|\text{Per}_m(\sigma)| = (2N)^m$  and the set  $\text{Per}(\sigma) = \cup_{m \geq 0} \text{Per}_m(\sigma)$  is dense in  $\Sigma_{\mathcal{A}, \mathcal{A}'}$ . According to the Theorem 4.1, the same result holds for  $f$  over  $\Lambda$ .

**5.1.2. Stable and unstable sets.** Consider a point  $S_0 = (x_0, y_0) \in \Lambda$ . As mentioned in Subsection 4.1, it lies on the intersection of line segments  $h_{y_0}$  and  $v_{x_0}$  passing through  $x_0$  and  $y_0$  respectively.

In the sense of symbolic dynamics, let  $\varsigma(S_0) = s = (\dots, s_{-2}, s_{-1}; s_0, s_1, \dots)$ . Therefore, by Definition 5.1,  $t = (\dots, t_{-2}, t_{-1}; t_0, t_1, \dots) \in W^s(s)$  if there exists an integer  $N'(t) \geq 0$  with  $t_n = s_n$  for  $n \geq N'(T)$ . Also,  $r = (\dots, r_{-2}, r_{-1}; r_0, \dots) \in W^u(s)$  if there exists an integer  $N(r) > 0$  with  $r_{-n} = s_{-n}$  for  $n \geq N(r)$ . In fact since the zip shift space representing the  $N$ -to-1 horseshoe is a full zip shift, for any letter in  $\mathcal{A} = \{a, b\}$ ,  $\varphi^{-1}\{a\}$  or  $\varphi^{-1}\{b\}$  can choose any symbol in  $Z$  or  $O$  without restrictions respectively. For non full zip shift cases, adjacent entries impose restrictions on the choice of symbols from  $Z$  or  $O$  to avoid the occurrence of forbidden words. Therefore in full zip shift spaces, the statement  $d^\pm((\sigma^{-n}(r)), (\sigma^{-n}(s))) = 0, n \rightarrow +\infty$  is equivalent to  $r_{-n} = s_{-n}$  for  $n \geq N(r)$ .

Consider the vertical line segment  $v_{x_0}$ . All points  $(x_0, y)$  on  $v_{x_0} \cap \Lambda$  have the code  $\varsigma((x_0, y)) \in \{t = (\dots, t_{-2}, t_{-1}; t_0, t_1, \dots), t_n = s_n, n \geq 0\} \subset W^s(s)$ . These points have the same forward code. If  $T \in v_{x_0}$ , then  $d(f^n(S_0), f^n(T)) \rightarrow 0$  when

$n \rightarrow +\infty$ . Therefore, for  $S_0 \in v_{x_0} \cap \Lambda$ , we define

$$\begin{aligned} W_{\text{special}}^s(S_0) &:= v_{x_0} \\ &\supset \{T \in \mathbf{S} : \varsigma(T) = (\dots, t_{-2}, t_{-1}; t_0, t_1, \dots), t_n = s_n, \mathbf{n} \geq \mathbf{0}\}, \\ W_{\text{global}}^s(S_0) &:= \left\{ T \in \bigcup_{k \geq 0} f^{-k}(v_{x_0}) : k \in \mathbb{N} \cup \{0\} \right\} \\ &\supset \{T \in \mathbb{R}^2 : \varsigma(T) \in W_{\text{global}}^s(s)\}. \end{aligned} \quad (5.4)$$

For  $T \in W_{\text{global}}^s(S_0)$ , there exists  $N' = N'(T) \geq 0$  such that for  $n \geq N'$ ,  $d(f^n(S_0), f^n(T)) \rightarrow 0$  when  $n \rightarrow +\infty$ .

For the horizontal line segment  $h_{y_0}$ , all points  $(x, y_0)$  on  $h_{y_0} \cap \Lambda$  have the code  $\varsigma((x, y_0)) \in \{r = (\dots, r_{-2}, r_{-1}; r_0, r_1, \dots), r_{-n} = s_{-n}, n > 0\} \subset W_{\text{special}}^u(s)$ . Such points have the same backward code. Using the notations of the Subsection 4.1, the point  $S_0 = (x_0, y_0)$  on  $h_{y_0}$  has  $N$  pre-images denoted by  $\{(x_{-1}^i, y_{-1}) : 1 \leq i \leq N\}$ . All these points lie on the unique horizontal line segment  $h_{y_{-1}}$  but on different stable line segments passing through  $x_{-1}^i$ . In fact,  $f^{-1}(h_{y_0}) \subset h_{y_{-1}}$ . Therefore, the unstable direction of all points on  $f^{-1}(y_0)$  overlay. We define

$$\begin{aligned} W_{\text{special}}^u(S_0) &= h_{y_0} \\ &\supset \{R \in \mathbf{S} : \varsigma(R) = (\dots, r_{-2}, r_{-1}; r_0, r_1, \dots), r_{-n} = s_{-n}, \mathbf{i} > \mathbf{0}\}, \\ W_{\text{global}}^u(S_0) &= \{R \in f^{\mathbf{N}}(h_{y_0}) : N \in \mathbb{N}\} \\ &\supset \{R \in \mathbb{R}^2 : \varsigma(R) \in W^u(s)\}. \end{aligned} \quad (5.5)$$

If  $R \in W_{\text{global}}^u(S_0)$ , then there exists  $N = N(R) > 0$  such that the  $d^\pm$  distance between the set of points  $f^{-N}(S_0)$  and the set of points  $f^{-N}(R)$  tends to zero as  $n \rightarrow +\infty$ .

The unstable curve and stable curves can also be determined by symbolic labels. As mentioned before,  $f^{-1}(\mathbf{S})$  is consisted of  $N$  vertical strips and in all of them, the rectangle  $V_{1^i}$  is followed by  $V_{0^i}$  for  $1 \leq i \leq N$ . This gives a rule denoted by

$$0_1 \rightarrow 1_1 \quad 0_2 \rightarrow 1_2, \dots, 0_N \rightarrow 1_N. \quad (5.6)$$

For simplicity, take  $N = 2$ . As in Figures 7 and 8, in the first step, the rectangle  $V_1$  is followed by  $V_0$  and also  $V_{1'}$  is followed by  $V_{0'}$ . Therefore,

$$0 \rightarrow 1 \quad 0' \rightarrow 1'. \quad (5.7)$$

In the second backward iterate,  $f^{-2}(\mathbf{S})$  is consisted of 4 vertical strips. By following the labels of the rectangles in each strip, the rules in (5.7) extend to

$$\begin{aligned} 00 \rightarrow 10 \rightarrow 11 \rightarrow 01 & \quad 0'0 \rightarrow 1'0 \rightarrow 1'1 \rightarrow 0'1 \\ 00' \rightarrow 10' \rightarrow 11' \rightarrow 01' & \quad 0'0' \rightarrow 1'0' \rightarrow 1'1' \rightarrow 0'1'. \end{aligned} \quad (5.8)$$

Note that the first item in (5.8) means that the rectangles  $V_{00}$ ,  $V_{10}$ ,  $V_{11}$  and  $V_{01}$  are followed one after another in  $f^{-2}(\mathbf{S})$ . The index  $ij$  in  $V_{ij}$  means that the rectangle  $V_j$  has stretched vertically into the rectangle  $V_i$  under  $f^{-1}$ . In fact, the strings in (5.7) produce the stable strings in (5.8). For example, the strings



$00 \rightarrow 10 \rightarrow 11 \rightarrow 01$  and  $0'0 \rightarrow 1'0 \rightarrow 1'1 \rightarrow 0'1$  are constructed from  $0 \rightarrow 1$  in (5.7) in the following sense.

The first line in (5.7) means  $V_0$  has stretched into  $V_0$  and  $V_1$  to create  $V_{00}$  followed by  $V_{10}$  and then  $V_1$  has stretched in  $V_1$  and  $V_0$  to create  $V_{11}$  followed by  $V_{01}$  (Note that the order of rectangles  $V_0$  and  $V_1$  is changed when  $V_1$  is stretching into them). Similarly,  $V_0$  has stretched in  $V_{0'}$  and  $V_{1'}$ . Afterwards,  $V_1$  has stretched in  $V_{1'}$  and  $V_{0'}$  to form the connected strip of rectangles  $V_{0'0}$ ,  $V_{1'0}$ ,  $V_{1'1}$  and  $V_{0'1}$ . Therefore,  $0 \rightarrow 1$  turns to two sequences. Each sequence has 4 words  $t = t'_1 t'_2$  of length two. The letters  $t'_2$  in the first (resp. second) two words in both sequences equal to 0 (resp. 1) which means  $V_0$  (resp.  $V_1$ ) in  $0 \rightarrow 1$  has been stretched. Because of the way the horseshoe is constructed, the first two letters  $t'_1$  would be 0 then 1 (resp.  $0'$  then  $1'$ ) and the second two letters  $t'_1$  would be 1 then 0 (resp.  $1'$  then  $0'$ ). Likewise, the two other strings in (5.8) are formed.

In the  $k$ th step, there are  $2^k$  disjoint stable strings of length  $2^k$  and its entries belong to  $\beta'_k$ . If a word  $w = s'_1 s'_2 \dots s'_k \in \beta'_k$  is given, we can simply find all of the vertical rectangles which are connected to  $V_{s'_1 \dots s'_k}$  in the stable strip containing  $V_w$  in  $f^{-k}(\mathbf{S})$ . To do this consider a sequence containing  $2^k$  entries of words such as  $t = t'_1 \dots t'_k$ . The words in this sequence resemble the labels of a sequence of connected vertical rectangles in the stable strip containing the rectangle with label  $s'_1 s'_2 \dots s'_k$ . If  $s'_k \in \{0, 1\}$ , then the first  $2^{k-1}$  words get 0 for  $t'_k$  and the other words get 1 for  $t'_k$ . If  $s'_k \in \{0', 1'\}$ , simply replace 0 and 1 by  $0'$  and  $1'$ . Continuing this procedure, for any  $i \in \{1 \dots k\}$ , if  $s'_i \in \{0, 1\}$ , then the first  $2^{i-1}$  words in the string get 0 for  $t'_i$ , the next  $2 \times 2^{i-1}$  words receive 1, the next  $2 \times 2^{i-1}$  words receive 0 and so on. The last  $2^{i-1}$  words receive 0 for  $t'_i$ . If  $s'_i \in \{0', 1'\}$ , replace 0 and 1 by  $0'$  and  $1'$ .

For example let  $w = s'_1 s'_2 s'_3 = 10'1'$ . So,  $s'_1 = 1$ ,  $s'_2 = 0'$  and  $s'_3 = 1'$ . The connected stable string to  $w$  has length  $2^3$  and its entries are in  $\beta'_3$ . Since  $s'_3 = 1'$ , the first  $2^2$  words in the string receive  $0'$  and the other words get  $1'$  in their 3rd positions. To determine the letters in the second position of the words, since  $s'_2 = 0'$ , the first two words receive  $0'$ , the next four words get  $1'$  and the last two words get  $0'$ . Finally,  $s'_1 = 1$ . The first position of the words in the string should be 0, 1, 1, 0, 0, 1, 1 and 0 respectively. Therefore, the connected stable string to  $w$  is as follows:

$$00'0' \rightarrow 10'0' \rightarrow 11'0' \rightarrow 01'0' \rightarrow 01'1' \rightarrow 11'1' \rightarrow w = 10'1' \rightarrow 00'1'.$$

Note that any word  $w \in \beta'$  belongs to a unique stable strip and can be used as the label of that strip.

Due to these strips, we define

$$\begin{aligned} W^s(P) &= \{R \in c_{x_0}^s : f^{km}(R) \subset v_{x_0}, k \geq 0\}, \\ W^u(P) &= \{R \in c_{y_0}^u : f^{-km}(R) \subset h_{y_0}, k \geq 0\}. \end{aligned} \tag{5.9}$$

**5.1.3. Homoclinic and heteroclinic points and their orbits.** Without loss of generality, suppose that  $f$  is a 2-to-1 horseshoe map. Let  $P = (x_0, y_0) \in \Lambda$  be a periodic point with zip shift code  $\zeta(P) = p = (\overline{p_0, p_1, \dots, p_{m-1}})$ . By (5.5),  $P$  lies on the

intersection of stable line segment  $v_{x_0} \cap \Lambda$  and unstable line segment  $h_{y_0} \cap \Lambda$  consisting of points with coding  $(\dots; \overline{p_0}, \overline{p_1}, \dots, \overline{p_{m-1}})$  and  $(\varphi_1(\overline{p_0}, \overline{p_1}, \dots, \overline{p_{m-1}}); \dots)$ , respectively. Therefore, any point  $P'$  in  $W^s(P) \cap W^u(P)$  is a homoclinic point to  $P$ . If  $\varsigma(P') = r \in \Sigma_{\mathcal{A}, \mathcal{A}'}$ , then there are natural numbers  $N(r) > 0$  and  $N'(r) \geq 0$  such that  $p_n = r_n$  for  $n \geq N(r)$  and  $p_{-n} = r_{-n}$  for  $n \geq N'(r)$ . Let  $P = (x_0, y_0)$  be a fixed point. For any  $k \geq 0$ ,  $f^k(P) = P$  and all the forward orbits of  $P$  lie on  $h_{y_0}$ . The point  $P$  has two pre-images under  $f$ , say  $P$  and  $P'$  where  $P'$  is a pre-fixed point. One of them passes through  $P$  and the other passes through  $P'$ . After  $k$  backward iterates,  $2^k$  stable curves passes through the pre-images of  $P$ .

Let  $H = (x, y) \in \Lambda$  be a homoclinic point to  $P$ . Denote by  $h_i(P, H)$ , the point in  $i$ th position in homoclinic orbit of  $(x, y)$  to  $P$  for  $i \in \mathbb{Z}$ . Therefore, the homoclinic orbit of  $(x, y)$  to  $P$  is the set of points

$$P \leftarrow \dots, h_{-2}(P, H), h_{-1}(P, H), H, h_1(P, H), \dots \rightarrow P.$$

Let  $u = \varsigma(H) = (\dots, u_{-2}, u_{-1}; u'_0, u'_1, \dots)$ . Then  $u$  is a homoclinic point to  $p$ . Let  $h_i(p, u)$  be the homoclinic orbit of  $u$  to  $p$ . This orbit can be determined as the argument in Section 5.  $\mathcal{A} = \{a, b\}$ ,  $\mathcal{B} = \{a\}$  and  $\mathcal{A}' = \{1, 2, 3, 4\}$  be three sets of alphabets. Then  $f$  is conjugate to  $\sigma_f : \Sigma_{\mathcal{A}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{A}, \mathcal{A}'}$  with  $\varphi_1(1) = \varphi_1(3) = a$  and  $\varphi_1(2) = \varphi_1(4) = b$ . The map  $\sigma_f$  illustrates the 2 to 1 feature of  $f$ . The map  $g$  is conjugate to  $\sigma_g : \Sigma_{\mathcal{B}, \mathcal{A}'} \rightarrow \Sigma_{\mathcal{B}, \mathcal{A}'}$  with  $\varphi_1(1) = \varphi_1(2) = \varphi_1(3) = \varphi_1(4) = a$  which shows that  $g$  is a 4 to 1 map.

**Example 5.5.** Let  $\mathcal{A}' = \{0, 1, 0', 1'\}$ ,  $\mathcal{A} = \{a, b\}$ ,  $\varphi_1(0) = \varphi_1(0') = a$  and  $\varphi_1(1) = \varphi_1(1') = b$  as in Figure 8. Let  $H$  be the homoclinic point to a periodic point  $P$  with coding  $\varsigma(P) = (\overline{01'})$  and  $\varsigma(H) = h = (\overline{a, b}, b, a; 1, \overline{0}, 1')$ . Obviously, the orbit of  $P$  has two stable directions. To obtain the backward orbit of  $H$  to  $P$  we use the second approach since  $\Sigma_{\mathcal{A}, \mathcal{A}'}$  is a full zip shift. Let  $\sigma^{-1}(h) \in \{h_1, h_2\}$  where  $h_1 = (\overline{a, b}, b; 0, 1, \overline{1}, 0)$  and  $h_2 = (\overline{a, b}, b; 0', 1, \overline{1}, 0)$ . Also,  $\sigma^{-1}(h_1) \in \{h_{11}, h_{12}\}$  and  $\sigma^{-1}(h_2) \in \{h_{21}, h_{22}\}$  where  $h_{11} = (\overline{a, b}; 1, 0, 1, \overline{1}, 0)$ ,  $h_{12} = (\overline{a, b}; 1', 0, 1, \overline{1}, 0)$ ,  $h_{21} = (\overline{a, b}; 1, 0', 1, \overline{1}, 0)$  and  $h_{22} = (\overline{a, b}; 1', 0', 1, \overline{1}, 0)$ . Now, since we have reached to the repeating part of the periodic point, the pre-images are determined according to the point  $P$ . This means that we choose  $\varphi_1^{-1}(a)$  and  $\varphi_1^{-1}(b)$  to be 0 and 1 respectively. Therefore,  $\sigma^{-1}(h_{ij})$  would be  $(\overline{b, a}; 1, \dots)$  where the dotted part is the entries of non-negative indices of  $h_{ij}$ ,  $i, j \in \{1, 2\}$ . The homoclinic point  $H$  produces 4 homoclinic orbits. All of these orbits converges to  $P$  in the past and future.

## REFERENCES

1. Roy L. Adler, Symbolic dynamics and Markov partitions, *Bull. Amer. Math. Soc.*, (N.S.) 35, no. 1, 1-56, 1998.
2. V. I. Arnold, A. Avez, Ergodic problems of classical mechanics, (The Mathematical Physics Monograph Series.) Pp. ix+286, Benjamin: New York and Amsterdam, 1968.
3. R. Adler, L. Flatto, Cross section map for geodesic flow on the modular surface, *Ergodic Theory, Symbolic Dynamics and Hyperbolic Spaces*, Oxford University Press, Oxford, 1991.

4. Roy L. Adler and Benjamin Weiss, *Similarity of automorphisms of the torus*, Memoirs of the American Mathematical Society, No. 98, American Mathematical Society, Providence, R.I., 1970.
5. NL. Balazs, and A. Voros, *The Quantized Baker's transformation*, Annals of Physics 190: 1-31, (1989).
6. R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms.*, Lecture Notes in Mathematics, Vol. 470. Springer-Verlag, Berlin-New York, (1975).
7. S. Bezuglyi, P. E. T. Jorgensen, *Transfer Operators, Endomorphisms, and Measurable Partitions*, Lecture Notes in Mathematics 2217, Springer, (2018)
8. E. Ermerson, Y. Lima and P. Mauricio, Symbolic dynamics for nonuniformly hyperbolic maps with singularities in high dimension, arXiv: Dynamical Systems, 2020.
9. J. Hadamard, *Les surfaces à courbures opposées et leurs lignes géodésiques* J. Math. Pures Appl. 5 (4): 27-73, (1898).
10. F. Przytycki, Anosov endomorphisms. *Studia Math.*, 58, 249-285, (1976).
11. D. Lind, M. Marcus, *An Introduction to Symbolic Dynamics and Coding*, Cambridge University Press, Cambridge (1995).
12. S. Lamei, P. Mehdipour, *An  $N$ -to-1 Smale Horseshoe*, pre-print (2021).
13. S. Lamei, P. Mehdipour,  *$(m, l)$ -Bernoulli Transformations*, pre-print (2022).
14. N. Martins, P. Mehdipour, Encoding  $n$ -to-1 baker's transformations (To appear in Archiv der Mathematik).
15. D. Ornstein, Bernoulli shifts with the same entropy are isomorphic, *Advances in Math.*, 4 (1970), 337-352.
16. S. B. Ovadia, Symbolic dynamics for non-uniformly hyperbolic diffeomorphisms of compact smooth manifolds, *Journal of Modern Dynamics*, 13 (2018), 43-113.
17. O. M. Sarig, Symbolic dynamics for surface diffeomorphisms with positive entropy, *J. Amer. Math. Soc.*, 26 (2013), no. 2, 341-426.
18. Sh. Claude Elwood, *A Mathematical Theory of Communication*, Bell System Technical Journal, 27 (3): 379-423, (July 1948).
19. Y. Sinai, Construction of Markov Partitions, *Func. Anal. and its Appl.*, 2, 245-253, (1968).
20. Smale, S. Differentiable dynamical systems. Bull. A mer. Math. Soc., 73, 747-817, (1967).
21. Shub, M., *All, Most, Some Differentiable Dynamical Systems*, Proceedings of the International Congress of Mathematicians, European Math. Society, 99-120, 2006.
22. Qian, M., Xie, J.sh., Zhu Sh., *Smooth ergodic theory for endomorphisms*, Lect.39 Notes Math, Vol. 1978, Springer-Verlag, Berlin Heidelberg, 2009.
23. D. Ruelle, *Thermodynamic formalism.*, Vol. 5. Encyclopedia of Mathematics and its Applications. Addison-Wesley Publishing Co., Reading, Mass., pp. xix+183, (1978).
24. D. Ruelle, *Thermodynamic formalism*, Vol. 5. Encyclopedia of Mathematics and its Applications. Addison-Wesley Publishing Co., Reading, Mass., pp. xix+183, (1978).
25. D. Ruelle, *Statistical mechanics: Rigorous results*, World Scientific. 1995.
26. P. Walters, *An Introduction to Ergodic Theory*, Springer-Verlag, New York, 1982.

<sup>1</sup> FACULTY OF MATHEMATICAL SCIENCES, UNIVERSITY OF GUILAN, P. O. BOX 1914, RASHT, IRAN.

*Email address:* lamei@guilan.ac.ir

<sup>2</sup> DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE VIÃÇOSA, DMA- BRAZIL.  
*Email address:* Pouya@ufv.br