ABELIAN VARIETIES ANALOGS OF TWO RESULTS ABOUT ALGEBRAIC CURVES

NELSON ALVARADO AND GIUSEPPE PARESCHI

ABSTRACT. We characterize decomposable principally polarized abelian varieties of the form $E \times B$, with E an elliptic curve, in two different ways. The first one is by the non-surjectivity of a certain multiplication map of global sections, i.e. by the non-vanishing of a certain 0-th Koszul cohomology group. The second one is by the non-surjectivity of a certain second order gaussian map. Both results are analogous to well known characterizations of hyperelliptic curves among all smooth curves of given genus. We also show that, according to previous work of the first author, the second characterization can be seen as an effective version of a theorem of Nakamaye characterizing the above decomposable abelian varieties as those of minimal Seshadri constant.

1. INTRODUCTION

In curve theory it is possible to characterize curves of genus ≥ 2 with special Brill-Noether properties (e.g. hyperelliptic curves) by means of algebraic properties of the section ring of the canonical bundle, or of line bundles of suitable high degree. We allude to properties as normal generation, normal presentation, vanishing of Koszul cohomology groups, i.e. "syzygies". Another less known way of characterizing hyperelliptic curves is by the non-surjectivity of certain higher gaussian maps. An interesting feature of this sort of results is that they are in terms of the nonsurjectivity of certainly uniformly defined linear maps.

The aim of this note is to show two results suggesting that there is a chance that similar things can be done, at least in some cases, in the realm of abelian varieties. In this paper we will mainly (but not exclusively) focus on the crucial case of principal polarizations.

To put the main results into perspective, recall that, among smooth curves of fixed genus $g \ge 2$, hyperelliptic curves are the most special ones in many respects (e.g. Brill-Noether theory). As such, they admit many different characterizations. We will show that, in some respects, decomposable g-dimensional p.p.a.v.'s $E \times B$, with dim E = 1 seem to play, among p.p.a.v's,, the same role of hyperelliptic curves among curves, as they can be characterized, among all g-dimensional p.p.a.v.'s , in two different ways, which are analogous to two specific characterizations of hyperelliptic curves.

1.1. First characterization. Our first result is

Theorem A. Let $(A, \underline{\theta})$ be a principally polarized abelian variety and Θ a divisor representing the polarization. The following are equivalent:

(1) A is isomorphic, as polarized variety, to $E \times B$, where dim E = 1.

Both authors were partially supported by the MIUR Excellence Department Project MatMod@TOV awarded to the Department of Mathematics of the University of Rome Tor Vergata. GP was also partially supported by the PRIN 2022 "Moduli spaces and Birational Geometry" and is a member of GNSAGA - INDAM.

(2) For every $x \in A$ there exists $\alpha \in \text{Pic}^0 A$ such that the multiplication map of global sections

 $H^{0}(A, \mathcal{O}_{A}(2\Theta)) \otimes H^{0}(A, \mathcal{I}_{x}(3\Theta) \otimes P_{\alpha}) \to H^{0}(A, \mathcal{I}_{x}(5\Theta) \otimes P_{\alpha})$ (1.1)

is not surjective.

(3) There exists $x \in A$ and $\alpha \in \text{Pic}^0 A$ such that the map (1.1) is not surjective.

A more manageable version Theorem A is given by

Corollary B. Let Θ by a symmetric divisor representing the principal polarization, and let $e \in A$ be the neutral element. Then $A \cong E \times B$, as in Theorem A, if and only if the multiplication map of global sections

$$H^0(A, \mathcal{O}_A(2\Theta)) \otimes H^0(A, \mathcal{I}_e(3\Theta)) \to H^0(A, \mathcal{I}_e(5\Theta))$$

is not surjective.

To put this into perspective, let us recall a classical theorem about smooth projective curves, due to the combined efforts of Mumford, Eisenbud-Koh-Stillman, Green-Lazarsfeld (after a previous result/conjecture of Lange-Martens).

Let C be a curve of genus $g \ge 1$ and L, N two line bundles on C. We consider the multiplication map of global sections

$$m_{L,N}: H^0(C,L) \otimes H^0(C,N) \to H^0(C,L \otimes N).$$

$$(1.2)$$

(a) The map $m_{L,N}$ is always surjective as soon as deg L, deg $N \geq 2g$ and deg L + deg N > 4g([M2, §2, Theorem 6], [G, Corollary 4.2.4]). According to Mukai, this result and the next one can be seen as a Fujita-type statement: we write $L = \omega_C \otimes E$, $N = K_C \otimes F$ so that the condition for the surjectivity of $M_{L,N}$ is: deg E, deg $F \geq 2$ and deg $E + \deg F > 4$.

(b) If deg $L = \deg N = 2g$ then the map $m_{L,N}$ is always surjective if $L \neq N$ ([EKS, Theorem 2]). If L = M the map $m_{L,L}$ is surjective unless L is not very ample (i.e., in the notation above, E is effective) or C is hyperelliptic ([GL1, Corollary 1.4], [LM]).

Part (a) of the previous result was extended to abelian varieties by Koizumi, [Ko, Theorem 4.6], improving earlier results of Mumford, [M1, Corollary p.340]. Koizumi's theorem says that for all $\alpha \in \text{Pic}^0 X$ and n, m such that $n \geq 2, m \geq n$ and n + m > 4 the map

$$H^{0}(A, \mathcal{O}_{A}(n\Theta) \otimes H^{0}(A, \mathcal{O}_{A}(m\Theta) \otimes P_{\alpha}) \to H^{0}(A, \mathcal{O}_{A}((n+m)\Theta \otimes P_{\alpha})$$
(1.3)

is surjective for every $\alpha \in \text{Pic}^0 A$ (see [BL, Proposition 7.3.4], [K2, Theorem 6.8(c)]).

Theorem A can be seen as an analog of part (b), which can be stated as follows: the curve C is elliptic or hyperelliptic if and only if for every line bundle $L = \omega_C \otimes E$ with deg E = 2 and for every point $x \in C$ there exists a line bundle $M = \omega_C \otimes F$ with deg F = 3 such that the map $m_{L,M(-x)}$ is not surjective.

We recall that Theorem A, as well as the above result on curves, can be seen as a result concerning graded modules. In the case of Theorem A the modules are

$$R_{A,x,\alpha}(2,3) := \bigoplus_{n \ge 0} H^0(A, \mathcal{I}_x((2n+3)\Theta) \otimes P_\alpha).$$

Moreover it is worth to recall that, remarkably, the above characterization of hyperelliptic curves is a particular case of a general result of Green-Lazarsfeld on the projective normality of very ample line bundles in function of the Brill-Noether theory of the curve ([GL1]). In turn such result has a (still open) conjectural extension to syzygies (Green-Lazarsfeld's [GL1, Conjecture 3.4]), which is known to hold in certain cases (see e.g. [GL2], [F]). A natural question (to which at this point we have no answer, not even conjectural, see however §5 below) is whether there is something like that in the realm of abelian varieties (having also in mind that Koizumi's result has been extended to defining equations and higher syzygies, see [K2, Theorem 6.13], [P2], [PP2]).

1.2. Second characterization. This makes use of higher gaussian maps (see $\S2(8)$ below). They were introduced, in the context of deformation theory of linear series on curves, in [ACGH, Chapter 9]. Subsequently, they were re-introduced, in a general setting, in [W].

Theorem C. Let $(A, \underline{\theta})$ be a polarized abelian variety (not necessarily principally polarized) and let $\mathcal{O}_A(\Theta)$ any line bundle representing the polarization. Let $p \ge 2$. The following are equivalent

- (1) A is isomorphic, as polarized variety, to $E \times B$, where dim E = 1.
- (2) There exists $\alpha \in \operatorname{Pic}^{0} A$ such that the p-th order gaussian map

$$\operatorname{Rel}_{A}^{p}(\mathcal{O}_{A}(2(p+1)\Theta), \mathcal{O}_{A}(2(p+1)\Theta) \otimes P_{\alpha}) \to H^{0}(A, S^{p}\Omega_{A}^{1} \otimes \mathcal{O}_{A}(4(p+1)\Theta) \otimes P_{\alpha})$$
(1.4)

is not surjective.

(3) The gaussian map

$$\operatorname{Rel}_{A}^{p}(\mathcal{O}_{A}(2(p+1)\Theta), \mathcal{O}_{A}(2(p+1)\Theta)) \to H^{0}(A, S^{p}\Omega_{A}^{1} \otimes \mathcal{O}_{A}(4(p+1)\Theta))$$
(1.5)

is not surjective.

We remark that, for principal polarizations, the theorem is false for p = 0, 1 (see the next Subsection). The crucial case is p = 2, asserting that $A \cong E \times B$ if and only if there exists some $\alpha \in \text{Pic}^0 A$ such that

$$\operatorname{Rel}_{A}^{2}(\mathcal{O}_{A}(6\Theta), \mathcal{O}_{A}(6\Theta) \otimes P_{\alpha}) \to H^{0}(A, S^{2}\Omega_{A}^{1} \otimes \mathcal{O}_{A}(12\Theta) \otimes P_{\alpha})$$

$$(1.6)$$

is not surjective, or the analogous condition (3) holds.

Again, to put thing into perspective, we first recall the following theorem of Bertram-Ein-Lazarsfeld ([BEL, Theorem 1.7]) on gaussian maps on curves:

(a) given a smooth curve C of genus $g \ge 1$, for all pairs of line bundles L and M on C such that $\deg L, \deg M \ge (g+1)(p+1)$ and $\deg L + \deg M > (p+2)(2g-2) + 4(p+1)$, the p-th order gaussian map

$$\operatorname{Rel}^p_C(L,M) \to H^0(C, \omega_C^{\otimes p} \otimes L \otimes M)$$
(1.7)

is surjective,

(b) there exist pairs of line bundles L and M on C with deg $L = \deg M = (p+2)(g-1) + 2(p+1)$ such that the map (1.7) is not surjective if and only if g(C) = 1 or C is hyperelliptic.

Part (a) of the above theorem was (partially) extended to abelian varieties in [P1, Theorem C] (re-proved and improved in [A, Theorem 6.1]), stating that, if $n, m \ge 2(p+1)$ and n+m > 4(p+1) then the k-th order gaussian map

$$\operatorname{Rel}_{A}^{p}(\mathcal{O}_{A}(n\Theta), \mathcal{O}_{A}(m\Theta) \otimes P_{\alpha}) \to H^{0}(A, S^{p}\Omega_{A}^{1} \otimes \mathcal{O}_{A}((n+m)\Theta) \otimes P_{\alpha})$$

is surjective for all $\alpha \in \operatorname{Pic}^0 A$. So Theorem C is analogous to part (b) for $p \geq 2$.

1.3. **p-jets separation.** Contrary to Theorem A, the proof of Theorem C is not direct (in fact it would be interesting to have a direct proof). In fact what we prove is an equivalent characterization in terms of jets separation:

Theorem D. Let $(A, \mathcal{O}_A(\Theta))$ be a polarized abelian variety. The following are equivalent

- (1) A is not isomorphic, as polarized variety, to $E \times B$, where dim E = 1.
- (2) Let p be an integer ≥ 2 . For all $\alpha \in \operatorname{Pic}^0 A$ the line bundle $\mathcal{O}_A((p+1)\Theta) \otimes P_\alpha$ is p-immersive, i.e. separates p-jets at every point of A, i.e. the natural map

$$H^0(A, \mathcal{O}_A((p+1)\Theta) \otimes P_\alpha) \to H^0(A, \mathcal{O}_A(p+1)\Theta) \otimes P_\alpha \otimes (\mathcal{O}_A/\mathcal{I}_x^{p+1}))$$

is surjective for all $x \in A$.

(3) For all $\alpha \in \operatorname{Pic}^0 A$ the line bundle $\mathcal{O}_A(3\Theta) \otimes P_\alpha$ is 2-immersive.

Note that, for principal polarizations, this is not true for p = 0, 1. For example, $\mathcal{O}_A(2\Theta)$ never separates tangent vectors at every point. The equivalence of Theorem D with Theorem C follows the work of the first author. This is recalled in Subsection 4.4 below. Briefly, via the Fourier-Mukai transform, [A, Theorem 1.2] establishes an equivalence between the *p*-jets separation thresholds and surjectivity thresholds of gaussian maps of order *p*.

Possible extensions of Theorems C and D seem (conjecturally) less obscure than Theorem A, as they point to the Sehadri constant. In fact an interesting aspect of Theorem D (and consequently, Theorem C) is that it provides an (optimal) effective version (and a different proof) of Nakamaye's theorem characterizing polarized abelian varieties of the form $E \times B$ as the only ones achieving the minimal Seshadri constant, namely 1 ([N, Theorem 1.1]). This is explained in Section 5, where we have included other remarks, questions and conjectures.

2. NOTATION, TERMINOLOGY AND BASIC FACTS USED IN THE PAPER

We will work over an algebraically closed ground field k of characteristic zero.

(1) Kernel bundle. For a base point free line bundle L on the variety A we denote M_L the locally free sheaf defined by the exact sequence

$$0 \to M_L \to H^0(A, L) \otimes \mathcal{O}_A \to L \to 0.$$

(2) Polarizations and isogenies. Given an ample line bundle L on an abelian variety A, we denote l its class in the Néron-Severi group and

$$\varphi_l : A \to \operatorname{Pic}^0 A$$

the corresponding isogeny. Given a non zero integer k,

 $k_A: A \to A$

denotes the multiplication-by-k isogeny $x \mapsto kx$.

(3) Fourier-Mukai functors. A Fourier-Mukai functor between bounded derived categories, associated to a kernel \mathcal{F} on $X \times Y$ is denoted

$$\Phi_F: D(X) \to D(Y).$$

(4) Cohomological support loci. Given an an abelian variety A, we denote \mathcal{P} the Poincaré line bundle on $A \times \operatorname{Pic}^0 A$. Given $\alpha \in \operatorname{Pic}^0 A$ (respectively $x \in A$) the line bundle on A (resp. on $\operatorname{Pic}^0 A$) parametrized by a closed point α (resp. by x) is denoted P_{α} (resp. P_x). Given a coherent sheaf \mathcal{F} on A, its cohomological support loci are the subvarieties (of $\operatorname{Pic}^0 A$)

$$V^{i}(A,\mathcal{F}) = \{ \alpha \in \operatorname{Pic}^{0}A \mid h^{i}(A,\mathcal{F} \otimes P_{\alpha}) > 0 \}.$$

(5) Vanishing conditions on coherent sheaves. A coherent sheaf \mathcal{F} on an abelian variety A is said to satisfy the index theorem with index zero, or, for short, to be IT(0), if $V^i(A, \mathcal{F}) = \emptyset$ for all i > 0. It is said to satisfy generic vanishing, or, for short, to be a GV sheaf, if $\operatorname{codim}_{\operatorname{Pic}^0 A} V^i(A, \mathcal{F}) \geq i$ for all i > 0. It is known by Hacon and Pareschi-Popa ([H], [PP3]) that \mathcal{F} is a GV sheaf if and only if its derived dual $\mathcal{F}^{\vee} := R\mathcal{H}om(\mathcal{F}, \mathcal{O}_A)$ satisfies the weak index theorem with index g (where $g = \dim A$), i.e. its Fourier-Mukai transform (with kernel \mathcal{P}^{\vee}) is a sheaf in cohomological degree g:

$$\Phi_{\mathcal{P}^{\vee}}(\mathcal{F}^{\vee}) = R^g \Phi_{\mathcal{P}^{\vee}}(\mathcal{F}^{\vee})[-g].$$
(2.1)

Finally, \mathcal{F} is said to be *M*-regular if $\operatorname{codim}_{\operatorname{Pic}^0 A} V^i(A, \mathcal{F}) > i$ for all i > 0. This is equivalent to the fact that (2.1) holds and the sheaf $R^g \Phi_{\mathcal{P}^{\vee}}(\mathcal{F}^{\vee})$ is torsion free. Needless to say, a coherent sheaf \mathcal{F} satisfies conditions IT(0), or GV or M-regularity if and only if $\mathcal{F} \otimes P_{\alpha}$ does, for every $\alpha \in \operatorname{Pic}^0 A$.

- (6) \mathbb{Q} -twisted coherent sheaves. According to [L2, Definitions 6.2.1 and 6.2.2] \mathbb{Q} -twisted coherent sheaves are equivalence classes of pairs $(\mathcal{F}, \underline{\delta})$, where \mathcal{F} is a coherent sheaf on an abelian variety A and $\underline{\delta}$ is a class in $NS(A)_{\mathbb{Q}}$, under the equivalence relation generated by $(\mathcal{F} \otimes L, \delta) \sim (\mathcal{F}, l + \underline{\delta})$ (where l is a polarization and L is a line bundle representing it). The equivalence class of $(\mathcal{F}, \underline{\delta})$, is denoted $\mathcal{F}\langle \underline{\delta} \rangle$.
- (7) Vanishing conditions on \mathbb{Q} -twisted coherent sheaves. According to [JP, §5] the above vanishing conditions extend as follows: given a polarization l, represented by a line bundle L, a coherent sheaf \mathcal{F} , and a rational number $\lambda = \frac{a}{b}$, the \mathbb{Q} -twisted coherent sheaf $\mathcal{F}\langle\lambda l\rangle$ is said to be IT(0) (respectively GV, M-regular) if the coherent sheaf $b_A^* \mathcal{F} \otimes L^{\otimes ab}$ is (see (2) above for the notation). It should be noted that to be IT(0) is an open condition, in the sense that $\mathcal{F}\langle\lambda l\rangle$ is IT(0) if and only if for all sufficiently small $\varepsilon \in \mathbb{Q}^+$, $\mathcal{F}\langle\lambda - \varepsilon l\rangle$ is IT(0)([JP, Theorem 5.2(c)]). On the other hand $\mathcal{F}\langle\lambda l\rangle$ is GV if and only of $\mathcal{F}\langle\lambda + \varepsilon l\rangle$ is IT(0)for all $\epsilon \in \mathbb{Q}^+$ ([JP, Theorem 5.2(a)]).
- (8) Gaussian maps (see e.g. [ACGH, §9], [W], [BEL, §1], [P1, §1]). On a projective variety A, let L and M two line bundles. The p-th order gaussian map is the linear map

$$\gamma^p_{L,M}: H^0(A \times A, \mathcal{I}^p_\Delta \otimes (L \boxtimes M)) \to H^0(A \times A, (\mathcal{I}^p_\Delta \otimes (L \boxtimes M))_{|\Delta}) = H^0(A, S^p\Omega^1_A \otimes L \otimes M)$$

It is customary to refer to the source as the space of p-th order relations between L and M and denote it

$$\operatorname{Rel}_{A}^{p}(L,M) := H^{0}(A \times A, \mathcal{I}_{\Lambda}^{p} \otimes (L \boxtimes M)).$$

3. Proof of Theorem A

3.1. Reduction. What we will prove is the following

Theorem 3.1.1. If dim $A \ge 2$ and Θ is irreducible then the vector bundles $M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(3\Theta) \otimes P_\alpha$ are globally generated for all $\alpha \in \operatorname{Pic}^0 A$.

N.ALVARADO, G.PARESCHI

Assuming this, Theorem A follows by the following standard argument. In the first place we recall that Θ is reducible if and only if the p.p.a.v. $(A, \underline{\theta})$ is *decomposable*, namely the product of lower dimensional p.p.a.v.'s $(A_i, \underline{\theta}_i)$ (e.g. [BL, Theorems 4.3.1-6], [K2, Corollary 10.4]). If this is the case then conditions (2) and (3) of Theorem A hold for the p.p.a.v. A if and only if they they hold on at least one of the factors A_i . Therefore (1) implies (2) (and (3)), because (2) is clearly true on elliptic curves. It remains to prove that (3) implies (1). By the same remark it is sufficient to prove that, under the assumption that dim $A \ge 2$ and Θ irreducible, the multiplication map (1.1) is surjective for all $x \in A$ and $\alpha \in \text{Pic}^0 A$. We have that $\mathcal{I}_x(3\Theta)$ and $\mathcal{I}_x(5\Theta)$ are IT(0) (because $\mathcal{O}_A(3\Theta) \otimes P_{\alpha}$ and $\mathcal{O}_A(5\Theta) \otimes P_{\alpha}$ are base point free for all $\alpha \in \text{Pic}^0 A$). Therefore, taking cohomology in the exact sequence

$$0 \to M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{I}_x(3\Theta) \otimes P_\alpha \to H^0(A, \mathcal{O}_A(2\Theta)) \otimes \mathcal{I}_x(3\Theta) \otimes P_\alpha \to \mathcal{I}_x(5\Theta) \otimes P_\alpha \to 0,$$

we get that the simultaneous surjectivity of all multiplication maps of global sections (1.1) is equivalent to the fact that the coherent sheaves $M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{I}_x(3\Theta)$ are IT(0) for all $x \in A$. The standard exact sequence

$$0 \to M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{I}_x(3\Theta) \to M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(3\Theta) \to (M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(3\Theta))_{|x} \to 0$$

shows that the sheaves $M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{I}_x(3\Theta)$ are IT(0) as soon as $M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(3\Theta)$ is IT(0) and globally generated. But it is well known that $M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(3\Theta)$ is IT(0) (this is an equivalent statement of Koizumi's surjectivity result recalled in the introduction after Corollary B). This concludes the proof that Theorem 3.1.1 implies Theorem A.

3.2. Main Lemma. In the proof of Theorem 3.1.1 the generation properties (in the sense of the paper [P4]) of the vector bundle $M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(2\Theta)$ play an essential role. In fact the main point is

Lemma 3.2.1. Let $(A, \underline{\theta})$ be an indecomposable p.p.a.v. of dimension $g \geq 2$.

- (1) The subvariety $V^0(A, M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(2\Theta))$ (with its reduced structure) is a prime divisor of $\operatorname{Pic}^0 A$ algebraically equivalent to $4^{g-1}\widehat{\underline{\theta}}$ (where $\underline{\hat{\theta}}$ denotes the principal polarization on the dual abelian variety $\operatorname{Pic}^0 A$).
- (2) the map

$$\operatorname{ev}_{U}: \bigoplus_{\beta \in U} H^{0}(A, M_{\mathcal{O}_{A}(2\theta)} \otimes \mathcal{O}_{A}(2\Theta) \otimes P_{\beta}) \otimes P_{\beta}^{-1} \to M_{\mathcal{O}_{A}(2\theta)} \otimes \mathcal{O}_{A}(2\Theta)$$

is surjective for all nonempty open subsets $U \subset \operatorname{Pic}^0 A$ meeting $V^0(A, M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(2\Theta))$.

(we recall that, in the terminology introduced in ([P4, Definition 1.1]) statement (2) means that the vector bundle $M_{\mathcal{O}_A(2\theta)} \otimes \mathcal{O}_A(2\Theta)$ is generated by $\{V^0(A, M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(2\Theta))\}$).

Lemma 3.2.1 implies Theorem 3.1.1. This follows at once from [P4, Proposition 2.3.1]. Let us review explicitly the argument in the case at hand. For $x \in A$ let U_x^{α} be the open set of Pic⁰A defined as $U_x^{\alpha} = \{\beta \in \text{Pic}^0A \mid x \notin \Theta_{\alpha-\beta}\}$, where, for $\gamma \in \text{Pic}^0A$, Θ_{γ} denotes the divisor corresponding to the line bundle $\mathcal{O}_A(\Theta) \otimes P_{\gamma}$ (which is the translate of Θ by γ , where γ is seen as a point of A via the identification induced by the principal polarization). We have that $x \in \Theta_{\alpha-\beta}$ if and only if $\beta \in \Theta_{\alpha-x}$. Therefore U_x^{α} is the complement of a theta divisor of Pic⁰A and therefore, by (1) of the Lemma, U_x^{α} meets the irreducible subvariety $V^0(A, M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(2\Theta))$ as soon as $g \geq 2$. Hence, by (2) of the Lemma, the map

$$\bigoplus_{\beta \in U_x^{\alpha}} H^0(A, M_{\mathcal{O}_A(2\theta)} \otimes \mathcal{O}_A(2\Theta) \otimes P_{\beta}) \otimes H^0(A, \mathcal{O}_A(\Theta) \otimes P_{\alpha-\beta}) \to (M_{\mathcal{O}_A(2\theta)} \otimes \mathcal{O}_A(3\Theta) \otimes P_{\alpha})_{|x|}$$

is surjective. Therefore the vector bundle $\mathcal{M} := M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(3\Theta) \otimes P_\alpha$ is globally generated at x, because the above map factors through the evaluation of global sections of \mathcal{M} at x. Since x is arbitrary, \mathcal{M} is globally generated. This proves that Lemma 3.2.1 imples Theorem 3.1.1.

3.3. Proof of Lemma 3.2.1. (1) Applying the Fourier-Mukai transform $\Phi_{\mathcal{P}}$ to the twisted evaluation map

 $e: H^0(A, \mathcal{O}_A(2\Theta)) \otimes \mathcal{O}_A(2\Theta) \to \mathcal{O}_A(4\Theta)$

we get a map of vector bundles on $\operatorname{Pic}^0 A$ (both of rank 4^g)

$$\Phi_{\mathcal{P}}(e): H^0(\mathcal{O}_A(2\Theta)) \otimes \Phi_{\mathcal{P}}(\mathcal{O}_A(2\Theta)) \to \Phi_{\mathcal{P}}(\mathcal{O}_A(4\Theta))$$

which, by base-change, is fiberwise the multiplication-of-global-sections map

$$m_{\alpha}: H^0(A, \mathcal{O}_A(2\Theta) \otimes H^0(A, \mathcal{O}_A(2\Theta) \otimes P_{\alpha}) \to H^0(\mathcal{O}_A(4\Theta) \otimes P_{\alpha}).$$

Therefore

$$V^0(A, M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(2\Theta))$$

(with the reduced scheme structure) coincides with the support of the divisor of zeroes

$$D := (\det \Phi_{\mathcal{P}}(e))_0.$$

The Néron-Severi class of D is $4^{g-1}\hat{\underline{\theta}}$. This follows from the computation of the first Chern class of the source and target of the map $\Phi_{\mathcal{P}}(e)$, for example using that

$$\varphi_{2\underline{\theta}}^*(\Phi_{\mathcal{P}}(\mathcal{O}_A(2\Theta))) \cong \mathcal{O}_A(-2\Theta)^{\oplus 2^g} \quad \text{and} \quad \varphi_{4\underline{\theta}}^*(\Phi_{\mathcal{P}}(\mathcal{O}_A(4\Theta))) \cong \mathcal{O}_A(-4\Theta)^{\oplus 4^g}$$

([Mu2, Proposition 3.11(1)]), and that the degree of the isogenies $\varphi_{k\underline{\theta}}$ is k^{2g} , (as they are identified, via the principal polarization, to the multiplication-by-k isogenies).

Now we consider the pulled back map

$$\varphi_{2\underline{\theta}}^* \Phi_{\mathcal{P}}(e) : H^0(\mathcal{O}_A(2\Theta)) \otimes \varphi_{2\underline{\theta}}^* \Phi_{\mathcal{P}}(\mathcal{O}_A(2\Theta)) \to \varphi_{2\underline{\theta}}^* \Phi_{\mathcal{P}}(\mathcal{O}_A(4\Theta))$$

As above the class of the divisor of zeroes of the determinant of this map, say E, is $4^{g} \theta$.

If $\underline{\theta}$ is irreducible, assuming, as we can, that the divisor Θ is symmetric, we have that

$$E = \sum_{a \in A[2]} \Theta_a \tag{3.1}$$

(note that, since Θ is assumed to be irreducible, this is a sum of distinct prime divisors). This is shown in [PS, Subsection 3.4]. Briefly, the proof is as follows. Fiberwise the map $\varphi_{2\underline{\theta}}^* \Phi_{\mathcal{P}}(e)$ is the multiplication map of global sections

$$m_x: H^0(A, \mathcal{O}_A(2\Theta)) \otimes H^0(A, \mathcal{O}_A((2\Theta)_x)) \to H^0(A, \mathcal{O}_A(2\Theta + (2\Theta)_x)).$$
(3.2)

A theorem of Kempf ([K1, Theorem 3], see [PS, Theorem 2.1] for the correct statement) says that the dimension of the kernel of the map m_x is the number of points in $A[2] \cap \Theta_x$, where A[2] denotes the group of 2-division points of A. In particular the linear map m_x is singular if and only if $x \in \Theta_a$ for some $a \in A[2]$ (recall that Θ is assumed to be symmetric). This proves that the support of Eis the right hand side of (3.1). Since the class of the right hand side is equal to class of E, namely $4^g \underline{\theta}$, (3.1) is proved. Finally, since $\varphi_{2\underline{\theta}}^*(\varphi_{2\underline{\theta}*}(\Theta)) = \sum_{a \in A[2]} \Theta_a$, it follows that $D = \varphi_{2\underline{\theta}*}(\Theta)$. We recall that it is well known (see e.g. [P3, p.1591]) that $\varphi_{2\underline{\theta}*}(\Theta)$ is a prime divisor (of class 4^{g-1}). Hence it coincides with $V^0(A, M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(2\Theta))$.

(2) Let us denote, for typographical brevity,

$$\mathcal{E} := M_{\mathcal{O}_A(2\Theta)} \otimes \mathcal{O}_A(2\Theta).$$

Let

$$\mathcal{T} := \Phi_{\mathcal{P}^{\vee}}(\mathcal{E}^{\vee})[g] = R^g \Phi_{\mathcal{P}^{\vee}}(\mathcal{E}^{\vee})$$

(the equality holds because \mathcal{E} is a GV sheaf, see §2(5)). Note that, by duality, \mathcal{T} is the dual of the vector bundle $\Phi_{\mathcal{P}}(\mathcal{E})$. By [P4, Corollary 6.2.1] (see below), in order to prove (2) it is enough to prove that the scheme-theoretic support of \mathcal{T} is irreducible and reduced, and that \mathcal{T} is of pure dimension g - 1. But \mathcal{T} sits in the locally free resolution

$$0 \to R^{g} \Phi_{\mathcal{P}^{\vee}}(\mathcal{O}_{A}(-4\Theta)) \to H^{g}(A, \mathcal{O}_{A}(-2\Theta)) \otimes \Phi_{\mathcal{P}^{\vee}}(\mathcal{O}_{A}(-2\Theta)) \to \mathcal{T} \to 0$$

where the first map is the dual of $\Phi_{\mathcal{P}}(e)$. Therefore \mathcal{T} is of pure dimension g-1 supported on the divisor D. This proves (2).

3.4. Appendix: proof of the generation property. For the reader's convenience, we outline the proof of [P4, Corollary 6.2.1] in the case at hand, which is much simpler. We need to show is that the map

$$\bigoplus_{\beta \in U} H^0(A, \mathcal{E} \otimes P_\beta) \otimes H^0(A, P_\beta^{\vee} \otimes k(x)) \to H^0(A, \mathcal{E} \otimes k(x))$$
(3.3)

(where $k(x) \cong k$ denotes the residue field at x) is surjective for all $x \in A$ and for all open sets $U \subset \operatorname{Pic}^0 A$ meeting D. Dualizing the individual maps of (3.3) we get

$$\operatorname{Ext}^{g}(k(x), \mathcal{E}^{\vee}) \to H^{g}(A, \mathcal{E}^{\vee} \otimes P_{\beta}^{\vee}) \otimes H^{0}(A, P_{\beta}^{\vee} \otimes k(x))^{\vee}$$

The Fourier-Mukai equivalence $\Phi_{\mathcal{P}^{\vee}}: D(A) \to D(\operatorname{Pic}^{0} A)$ identifies the source of the above map to $\operatorname{Hom}(P_{x}^{\vee}, \Phi_{\mathcal{P}^{\vee}}(\mathcal{E}^{\vee})) = \operatorname{Hom}(P_{x}^{\vee}, R^{g}\Phi_{\mathcal{P}^{\vee}}(\mathcal{E}^{\vee}))$ (recalling that \mathcal{E} is a GV sheaf, see above). Moreover, by base-change, $H^{g}(A, \mathcal{E}^{\vee} \otimes P_{\beta}^{\vee})$ is identified to fibre at β of the coherent sheaf $R^{g}\Phi_{\mathcal{P}^{\vee}}(\mathcal{E}^{\vee})$ and $H^{0}(A, P_{\beta}^{\vee} \otimes k(x))^{\vee}$ is identified to the fibre at β of the line bundle P_{x} . In conclusion, the Fourier-Mukai equivalence $\Phi_{\mathcal{P}^{\vee}}$ identifies the dual of the map (3.3) to the evaluation map

$$H^{0}(\operatorname{Pic}^{0}A, R^{g}\Phi_{\mathcal{P}^{\vee}}(\mathcal{E}^{\vee}) \otimes P_{x}) \to \coprod_{\beta \in U} R^{g}\Phi_{\mathcal{P}^{\vee}}(\mathcal{E}^{\vee}) \otimes P_{x} \otimes k(\beta)$$
(3.4)

An element of the kernel of this map would be a global section vanishing identically on a non empty open set of the support of the sheaf (pure of dimension g-1) $R^g \Phi_{\mathcal{P}^{\vee}}(\mathcal{E}^{\vee})$, which is a one-codimensional reduced irreducible subvariety of Pic⁰A. Therefore such a global section must be zero.

4. 2-jets separation and proof of Theorems C and D

4.1. **Proof of Theorem D.** We recall that a line bundle L is said to separate p-jets at a point $x \in A$ if the map $H^0(A, L) \to H^0(A, L \otimes \mathcal{O}_A/\mathcal{I}_x^{p+1})$ is surjective, and p-immersive if it separates p-jets at every point.

It is clear, as it happens for elliptic curves, that if $A \cong E \times B$ no line bundle of the form $\mathcal{O}_A((p+1)\Theta) \otimes P_\alpha)$ is p-immersive. To establish the converse we begin with the following

Lemma 4.1.1. The sheaf $\mathcal{I}_x^2(2\Theta) \otimes P_\alpha$ is M-regular for all $x \in A$ and $\alpha \in \operatorname{Pic}^0 A$ if and only if A is not isomorphic, as polarized variety, to $E \times B$, where E is an elliptic curve.

Proof. It is enough to prove the statement for $P_{\alpha} = \mathcal{O}_A$, and we can assume that Θ is symmetric. Since $V^i(A, \mathcal{I}_x(2\Theta))$ is empty for $i \geq 2$, $\mathcal{I}_x(2\Theta)$ is M-regular if and only if $\operatorname{codim}_{\operatorname{Pic}^0 A} V^1(A, \mathcal{I}_x^2(2\Theta)) > 1$. Since the line bundles $\mathcal{O}_A(2\Theta) \otimes P_{\alpha}$ are translates of $\mathcal{O}_A(2\Theta)$ (and conversely) and $\mathcal{I}_x^2(t_y^*\mathcal{O}_A(2\Theta)) \cong t_y^*(\mathcal{I}_{x-y}^2(2\Theta))$ we have that $V^1(A, \mathcal{I}_x^2(2\Theta))$ coincides with the locus $R = \{z \in A, h^1(A, \mathcal{I}_z^2(2\Theta)) > 0\}$, i.e. the ramification locus of the Kummer morphism $f : A :\to \mathbb{P}(H^0(A, \mathcal{O}_A(2\Theta)^{\vee}))$. If the p.p.a.v A is indecomposable then it is well known that R is either 0-dimensional or empty. If $(A, \underline{\theta})$ is decomposed as the product of $(A_1, \underline{\theta}_1) \times (A_2, \underline{\theta}_2)$ then the components of the ramification locus (if any) have dimension equal to dim A_1 or dim A_2 . In conclusion, the ramification locus has codimension 1 if and only if one of the factors is an elliptic curve. \Box

As it is well known, if $\mathcal{I}_x^2(2\Theta)$ is M-regular, then $\mathcal{I}_x^2(3\Theta)$ is globally generated (this is the basic M-regularity criterion of [PP1, Theorem 2.4]). In particular, it is globally generated at the point x itself, i.e. the map $H^0(A, \mathcal{I}_x^2(3\Theta)) \to H^0(A, (\mathcal{I}_x^2/\mathcal{I}_x^3)(3\Theta))$ is surjective, and it is easily seen that this means that $H^0(A, \mathcal{O}_A(3\Theta))$ surjects onto $H^0(A, (\mathcal{O}_A/\mathcal{I}_x^3)(3\Theta))$, i.e. $\mathcal{O}_A(3\Theta)$ separates 2-jets at x. Since x is arbitrary this proves the implication $(1) \Rightarrow (3)$ Theorem D.

The same argument (actually an easier version of it, because it follows that for $p \geq 3$ the sheaf $\mathcal{I}_x^p(p\Theta)$ is IT(0), and not only M-regular) proves that the implication $(3) \Rightarrow (2)$.

4.2. **p-jets separation thresholds.** For the reader's benefit, we recall the notion of *p-jets sepa*ration thresholds and *p*-th gaussian maps surjectivity thresholds from the paper [A]. Let L be an ample line bundle on A. Since the set of all translates t_y^*L coincides with the set of all line bundles of the form $L \otimes P_{\alpha}$, we have that L is *p*-immersive if and only if, fixing a point $e \in A$ (for example the neutral element) the line bundles $L \otimes P_{\alpha}$ separate *p*-jets at *e* for all $\alpha \in \text{Pic}^0 A$. In turn, this is equivalent to the fact that the coherent sheaf $\mathcal{I}_e^{p+1}L$ is an IT(0) sheaf. But (see §2(7)) the IT(0) property can be defined also for \mathbb{Q} -twisted sheaves, as $\mathcal{I}_e^{p+1}\langle\lambda L\rangle$ for $\lambda \in \mathbb{Q}$. Therefore it is natural to define the *p-jets separation thresholds* as

$$\epsilon_p(l) = \inf\{\lambda \in \mathbb{Q} \mid \mathcal{I}_e^{p+1}(\lambda L) \text{ is } \mathrm{IT}(0)\}$$

(note that such invariants depend only on the Néron-Severi class l of L). This generalizes to higher jets the base point freeness threshold introduced in [JP, §8] and [C].

By definition the *p*-jets thresholds are multiplicative, in the sense that, given a polarization $\underline{\theta}$ on A, $\epsilon_p(\underline{k}\underline{\theta}) = \frac{1}{k}\epsilon_p(\underline{\theta})$. Moreover it known ([A, Corollary 3.9]) that

$$\epsilon_p((p+1)\underline{\theta}) = \frac{\epsilon_p(\underline{\theta})}{p+1} \le 1.$$
(4.1)

Moreover equality is achieved, i.e. $\epsilon_p((p+1)\underline{\theta}) = 1$, if and only if $(p+1)\underline{\theta}$ is not *p*-immersive. Thus Theorem D tells that, for $p \ge 2$, this happens if and only if $A \cong E \times B$.

4.3. Gaussian maps thresholds. Following a similar principle, in [A, §5] are defined higher gaussian maps surjectivity thresholds. To recall this, we use the notation of §2(8) on gaussian maps. In the first place, applying p_{2*} to the exact sequence

$$0 \to \mathcal{I}^{p+1}_{\Delta} \otimes (L \boxtimes M) \to \mathcal{I}^{p}_{\Delta} \otimes (L \boxtimes M) \to (\mathcal{I}^{p}_{\Delta}/\mathcal{I}^{p+1}_{\Delta})(L \boxtimes M) \to 0$$

one gets exact sequences

$$0 \to R^p_{A,L} \otimes M \to R^{p-1}_{A,L} \otimes M \to S^p \Omega^1_A \otimes L \otimes M$$

where $R_{A,L}^i := p_{2*}(\mathcal{I}_{\Delta}^{i+1} \otimes L)$, and the gaussian map $\gamma_{L,M}^p$ is the H^0 of the last map. It is easily seen that this is exact also on the right as soon as the line bundle L is p-immersive (see e.g. [A, (5.3)]), i.e., in the above notation, $\epsilon_p(L) < 1$ (this also implies also that the coherent sheaves $R_{A,L}^p$ are locally free for $k \leq p$). If this is the case the condition that the locally free sheaf $R_{A,L}^p$ is IT(0) implies that the gaussian map $\gamma_{L,M\otimes P_{\alpha}}^p$ is surjective for all $\alpha \in \operatorname{Pic}^0 A$. Since the IT(0) condition makes sense for \mathbb{Q} -twisted sheaves, it is natural to define

$$\mu_p = \inf\{t \in \mathbb{Q} \mid R^p_{A,L} \text{ is } \operatorname{IT}(0)\}^1$$

Then [A, Theorem 5.5]² states that, as soon as L is p-immersive then $\mu_p(L)$ depends only on the Néron-Severi class of L, so one can call it $\mu_p(l)$ and

$$\mu_p(l) = \frac{\epsilon_p(l)}{1 - \epsilon_p(l)}.\tag{4.2}$$

4.4. **Proof of Theorem C.** Theorem C follows quickly from (4.2). Indeed, as we saw, Theorem D can be stated as follows: given an integer $p \ge 2$, A is not isomorphic, as polarized variety, to $E \times B$ if and only if $\epsilon_p((p+1)\underline{\theta}) < 1$ i.e. $\epsilon_p(2(p+1)\underline{\theta}) < \frac{1}{2}$. By (4.2) this yields $\mu_p(2(p+1)\underline{\theta}) < 1$ i.e. the p-th gaussian map (1.4) is surjective for all $\alpha \in \operatorname{Pic}^0 A$. This proves the (2) implies (1) in Theorem D. The implication $(1) \Rightarrow (3)$ can be proved in a similar way, or more directly one sees simply that the p-th gaussian map (1.5) is never surjective on elliptic curves (in this case the dimensions of the source and target are equal and the map can't be injective because, as it is well known, for p even (respectively odd) it is zero on skew-symmetric (resp. symmetric) tensors). Hence the map is not surjective on decomposable polarized abelian varieties of the form $E \times B$.

5. VARIANTS, REMARKS AND CONJECTURES

5.1. Variants of the first characterization. Theorem A can be extended to other multiplication maps of global sections, involving semihomogeneous vector bundles. This statements don't seem to have an analog for curves of genus $g \ge 2$. In what follows we will use the notation of the paper [AP] for simple semihomogeneous bundles. Namely, given a polarized abelian variety (A, l), and a rational number $\lambda = \frac{a}{b}$, with b > 0, we will denote $E_{A,\lambda l}$ a simple semihomogeneous vector bundle on A such that $\frac{[\det E]}{\mathrm{rk}E} = \lambda l$ in $NS(A)_{\mathbb{Q}}$ ([Mu1, Theorem 7.11]). Then we have

Theorem 5.1.1. Let $(A, \underline{\theta})$ be a p.p.a.v., and let $n \ge 2$. The following are equivalent:

- (1) A is isomorphic, as polarized variety, to $E \times B$, where dim E = 1.
- (2) For every $x \in A$ there exists $\alpha \in \text{Pic}^0 A$ such that the multiplication map of global sections

$$H^{0}(A, \mathcal{O}_{A}(n\Theta)) \otimes H^{0}(A, \mathcal{I}_{x} \otimes E_{A, \frac{2n-1}{n-1}\underline{\theta}} \otimes P_{\alpha}) \to H^{0}(A, \mathcal{I}_{x}(n\Theta) \otimes E_{A, \frac{2n-1}{n-1}\underline{\theta}} \otimes P_{\alpha})$$
(5.1)

is not surjective.

(3) There exists $x \in A$ and $\alpha \in \text{Pic}^0 A$ such that the map (5.1) is not surjective.

¹The definition given in [A, Definition 5.4] is slightly different, but equivalent (see [A, Lemma 5.3]).

²This theorem generalizes to arbitrary p the relation between surjectivity of multiplication maps of global sections and base point freeness shown in [JP, Theorem D], see also [AP, Proposition A, Corollary 4.3.1] for a simpler formulation.

This is proved in the very same way, replacing the multiplication maps of global sections (3.2) (and related maps) used in the proof of Theorem A with the multiplication maps

$$H^{0}(A, \mathcal{O}_{A}(n\Theta)) \otimes H^{0}(A, E_{A, \frac{n}{n-1}\underline{\theta}} \otimes P_{\alpha}) \to H^{0}(A, \mathcal{O}_{A}(n\Theta) \otimes E_{A, \frac{n}{n-1}\underline{\theta}} \otimes P_{\alpha}),$$

and replacing Kempf's theorem about the dimension of the kernel of such maps ([PS, Theorem 2.1]) with [P3, Theorem A].³

5.2. Remarks about the first characterization. (a) As we said in the introduction, we don't know how to generalize the first characterization, in the spirit of the beautiful and rich theory concerning curves. However the following seems to be a reasonable conjecture, extending Theorem A to higher syzygies (similar to the characterization of hyperelliptic curves given in [GL2]). To this purpose we consider the Koszul cohomology groups $K_p(n, m, x, \alpha)$ defined as the cohomology of the complexes

$$\wedge^{p+1} H^0(\mathcal{O}_A(n\Theta)) \otimes H^0(\mathcal{I}_x(m\Theta) \otimes P_\alpha) \to \wedge^p H^0(\mathcal{O}_A(n\Theta) \otimes H^0(\mathcal{I}_x(n+m)\Theta) \otimes P_\alpha) \to \\ \to \wedge^{p-1} H^0(\mathcal{O}_A(n\Theta)) \otimes H^0(\mathcal{I}_x(2n+m)\Theta) \otimes P_\alpha).$$

It can be shown, using [P2] or [PP2], that, for $n, m \ge p+3$, $K_p(n, m, x, \alpha) = 0$ for all x and α (i.e. the above complex is always exact in the middle), and we expect that, for n = p+2 and m = p+3, there are pairs (x, α) such that $K_p(n, m, x, \alpha) \ne 0$ if and only if A is of the form $E \times B$.

(b) A way to extend Theorem A would be to replace, in the maps of Theorem A(b), the ideal sheaf of one point \mathcal{I}_x with the ideal sheaf of *n* points, or of a subscheme of length *n*. We remark that in a jacobian J(C), where *C* is a curve of genus *g*, if *Z* is a scheme of length *g* contained in (a translate of) *C* then, for a suitable choices of a theta divisor Θ and of $\alpha \in \operatorname{Pic}^0 J(C)$ the map

$$H^{0}(J(C), \mathcal{O}_{J(C)}(2\Theta)) \otimes H^{0}(J(C), \mathcal{I}_{Z}(3\Theta) \otimes P_{\alpha}) \to H^{0}(J(C), \mathcal{I}_{Z}(5\Theta) \otimes P_{\alpha})$$
(5.2)

is not surjective⁴ (therefore the same happens on decomposable varieties of the form $J(C) \times B$).

5.3. Variants of the second characterization. There are many other different second order gaussian maps whose non-surjectivity is equivalent to the non-surjectivity of (1.6), hence characterizing decomposable abelian varieties of the form $E \times B$. This follows from (4.2), coupled with Theorem D. For example, starting with the polarization $l = 4\underline{\theta}$, from Theorem D we have that if A is not of the form $E \times B$ then $\epsilon_2(4\Theta) = \frac{3}{4}\epsilon_2(3\Theta) < \frac{3}{4}$. By (4.2) this implies that $\mu_2(4\Theta) < 3$ which means that the gaussian maps

$$\operatorname{Rel}_{A}^{2}(\mathcal{O}_{A}(4\Theta), \mathcal{O}_{A}(12\Theta) \otimes P_{\alpha}) \to H^{0}(A, S^{2}\Omega_{A}^{1} \otimes \mathcal{O}_{A}(16\Theta) \otimes P_{\alpha})$$

are surjective for all $\alpha \in \text{Pic}^0 A$. Conversely, it can be shown that for elliptic curves this does not happen, hence it does not happen also for decomposable varieties $A \cong E \times B$.

One has similar characterizations even when $\mu_2(k\underline{\theta})$ is not integer. Rather than giving the general formula (left to the reader) let us give an example. Let $l = 5\underline{\theta}$. Then, as above, if A is not of the form $E \times A$ then $\epsilon_2(5\underline{\theta}) < \frac{3}{5}$. Thus, by (4.2), $\mu_2(5\theta) < \frac{3}{2}$, which means that $R^2_{A,\mathcal{O}_A(5\Theta)} \langle \frac{15}{2}\underline{\theta} \rangle$ is an

 $H^{0}(C, \mathcal{O}_{C}(2\Theta)) \otimes H^{0}(C, O_{C}(3\Theta - Z)) \otimes P_{\alpha}) \to H^{0}(C, \mathcal{O}_{C}(5\Theta - Z)) \otimes P_{\alpha})$

³Beware that in [P3] the notation is different, namely the (symmetric) simple semihomogeneous vector bundles $E_{A,\frac{a}{b}\underline{\theta}}$ are denoted, according to [O], $W_{b,a}$).

 $^{^{4}}$ One can reduce the non-surjectivity of (5.2) to the non surjectivity of the map

Choosing Θ such that $\mathcal{O}_C(2\Theta)$ is not very ample and $\alpha \in \operatorname{Pic}^0 J(C)$ such that $\mathcal{O}_C(3\Theta - Z) \otimes P_\alpha \cong \mathcal{O}_C(2\Theta)$ one has the non-surjectivity. Note that if the curve C is hyperelliptic then for any choice of Θ , for $\alpha \in \operatorname{Pic}^0 A$ such that $\mathcal{O}_C(3\Theta - Z) \otimes P_\alpha \cong \mathcal{O}_C(2\Theta)$ the above map is not surjective.

N.ALVARADO, G.PARESCHI

IT(0) Q-twisted coherent sheaf. This can be expressed as the surjectivity of certain (second order) gaussian maps by means of semihomogeneous vector bundles, as in [AP]. Indeed, considering a simple semihomogeneous vector bundle $E := E_{A,\frac{15}{2}\underline{\theta}}$ (notation as in Subsection 5.1), the condition that the Q-twisted coherent sheaf $R^2_{A,\mathcal{O}_A(5\Theta)}\langle \frac{15}{2}\underline{\theta}\rangle$ is IT(0) equivalent to the condition that the locally free sheaf $R^2_{A,\mathcal{O}_A(5\Theta)} \otimes E_{A,\frac{15}{2}\underline{\theta}}$ is IT(0) ([AP, Proposition 2.1.2]). In turns this means the surjectivity of the gaussian maps

$$\operatorname{Rel}_{A}^{2}(\mathcal{O}_{A}(5\Theta), E_{A, \frac{15}{2}\underline{\theta}} \otimes P_{\alpha}) \to H^{0}(A, S^{2}\Omega_{A}^{1} \otimes \mathcal{O}_{A}(5\Theta) \otimes E_{A, \frac{15}{2}\underline{\theta}} \otimes P_{\alpha}).$$

5.4. Relation with the Seshadri constant. In view of (4.1) the normalized *p*-jets separation thresholds of a polarization $\underline{\theta}$ are defined as the quantities $\frac{\epsilon_p(\underline{\theta})}{p+1}$. Let us recall a result of the first author relating them to the Seshadri constant. Let $\epsilon(A,\underline{\theta})$ denote the Seshadri constant of the polarization $\underline{\theta}$ ([L1, p.293]). Then [A, Theorem 3.7] states that $\epsilon_{p+r}(\underline{\theta}) \leq \epsilon_p(\underline{\theta}) + \epsilon_r(\underline{\theta})$ (hence in particular $\epsilon_{kp}(\underline{\theta}) \leq k\epsilon_p(\underline{\theta})$) and

$$\epsilon(A,\underline{\theta}) = \sup_{p \ge 0} \frac{p+1}{\epsilon_p(\underline{\theta})} = \lim_{p \to \infty} \frac{p+1}{\epsilon_p(\underline{\theta})}.$$
(5.3)

This result is in fact a sharpened version, valid for abelian varieties, of Demailly's theorem relating the Seshadri constant to asymptotic p-jets separation ([L1, Theorem 5.1.17], [BS]).

It follows that Theorem D is an effective version of Nakamaye's theorem characterizing polarized abelian varieties of the form $E \times B$ as the only ones achieving the minimal Seshadri constant, namely 1 ([N, Theorem 1.1]). In fact, thanks to (5.3), Nakamaye's theorem characterizes such decomposable abelian varieties as the only ones such that $\sup_{p\geq 0} \frac{p+1}{\epsilon_p(\underline{\theta})} = 1$, but Theorem D shows that they are in fact the only ones such that $\sup_{0\leq p\leq 2} \frac{p+1}{\epsilon_p(\underline{\theta})} = 1$. In other words, in this case the asymptotic description (5.3) is equivalent to an explicit finite condition.

Thus a natural conjecture, already raised in [A, Question 7.2] as a problem, is that something of this sort always happens. Namely for all real numbers $\lambda \geq 1$ there is a positive integer $p_0(\lambda)$ such that $\epsilon(A, \underline{\theta}) \leq \lambda$ if and only if $\sup_{p \leq p_0(\lambda)} \frac{p+1}{\epsilon_p(\underline{\theta})} \leq \lambda$. If true the locus of polarized abelian varieties $(A, \mathcal{O}_A(\underline{\theta}))$ such that $\epsilon(A, \underline{\theta}) \leq \lambda$ would be characterized, via the above mentioned [A, Theorem 1.2], by the non-surjectivity of suitable gaussian map of order $p_0(\lambda)$.

References

- [ACGH] Arbarello, E., Cornalba, M., Griffiths, Ph., Harris, J., Special divisors on algebraic curves, Informal lecture notes distributed in conjunction with the principal lectures at the Regional Algebraic Geometry Conference, Athens, Georgia, May 7-May 12, 1979. Reprinted in "Phillip A. Griffiths, Selected Works of Phillip A. Griffiths with Commentary, volume II, American Mathematical Society, Providence, RI, 2003, pp. 649-778".
- [A] Alvarado, N., Jets-separation thresholds, Seshadri constants and higher Gauss-Wahl maps on abelian varieties, preprint arXiv:2407.20769 [math.AG]
- [AP] Alvarado, N., Pareschi, G., Semihomogeneous vector bundles, Q-twisted sheaves, duality, and linear systems on abelian varieties, preprint 2024, arXiv:2407.20646 [math AG]
- [BL] Birkenhake, Ch., Lange, H., Complex abelian varieties, 2nd edition, Springer, 2004.
- [BS] Bauer, Th., Szemberg, T., Seshadri constants and the generation of jets, Journal of pure and applied algebra 213 (2009) 2134–2140
- [BEL] Bertram, A., Ein, L., Lazarsfeld, R., Surjectivity of the Gaussian maps for line bundles of large degree on curves, in Algebraic Geometry, Proceedings, 1990, Lect. Notes in Math. 1479, 15–25
- [EKS] Eisenbud, D., Koh, J., Stillman, M., Determinantal equations for curves of high degree, Amer. J. Math., 110 (1988) 513–539

- [C] Caucci, F., The basepoint-freeness threshold and syzygies of abelian varieties, Algebra Number Theory 14 (2020) 947–970
- [F] Farkas, G., Difference varieties and the Green-Lazarsfeld Secant Conjecture epiga:11658 Épijournal de Géométrie Algébrique, (2024)
- [H] Hacon, C., A derived category approach to generic vanishing, J. Reine Angew. Math. 575 (2004), 173–187
- [K1] Kempf, G., Multiplication over abelian varieties, Amer. J. Math., 110 (1988), 765–773
- [K2] Kempf, G., Complex Abelian Varieties and Theta Functions, Springer-Verlag (1991)
- [Ko] Koizumi, S., Theta relations and projective normality of abelian variety, Amer. J. Math., 98 (1976) 865-889
- [G] Green, M., Koszul cohomology and the geometry of projective varieties, J. Diff. Geometry 19 (1984) 125–171
- [GL1] Green, M., Lazarsfeld, R., Projective normality of complete linear series on an algebraic curve, Invent. math. 83, 73-90 (1986)
- [GL2] Green, M., Lazarsfeld, R., Some results on the syzygies of finite sets and algebraic curves, Comp. Math 67 (1988) 301–314
- [JP] Jiang, Z., Pareschi, G., Cohomological rank functions on abelian varieties, Ann. Sci. Éc. Norm. Supér. (4) 53 (2020) 815–846
- [LM] Lange, H., Martens, G.: Normal generation and presentation of line bundles of low degree. J. Reine Angew. Math. 356 (1985) 1-18
- [L1] Lazarsfeld, R. Positivity in Algebraic Geometry I, Springer, 2004
- [L2] Lazarsfeld, R. Positivity in Algebraic Geometry II, Springer, 2004
- [Mu1] S. Mukai, Semi-homogeneous vector bundles on abelian varieties J. Math. Kyoto Univ. 18 (1978), 239–272
- [Mu2] S. Mukai, Duality between D(X) and D(X) with its application to Picard sheaves, Nagoya Math. J. 81 (1981), 153–175.
- [M1] Mumford, D., Equations defining abelian varieties I, Inv. Math 1 (1966) 287–354
- [M2] Mumford, D., Varieties defined by quadratic equations, in Questions on Algebraic Varieties, Varenna 1969 C.I.M.E. Summer Schools (1969) Springer, 29–100
- [N] Nakamaye, M., Seshadri constants on abelian varieties, Amer. J. Math. 118 (1996), no. 3, 621–635.
- [O] Oprea, D., The Verlinde bundles and the semihomogeneous Wirtinger duality, J. Reine Angew. Math. 654 (2011), 181–217
- [P1] Pareschi, G., Gaussian maps and multiplication maps on certain projective varieties, Comp. Math. 98 (1995) 219–268
- [P2] Pareschi, G., Syzygies of abelian varieties, Journal of the American Mathematical Society 13 (2000), 651-664
- [P3] Pareschi, G., Torsion points on theta divisors and semihomogeneous vector bundles, Algebra and Number Theory Vol.15 n.6 (2021) 1581 - 1592
- [P4] Pareschi, G., Generation and ampleness of coherent sheaves on abelian varieties, with application to Brill-Noether theory, Pure and Applied Mathematics Quarterly, 20 (2024) 2379-2413
- [PP1] Pareschi, G., Popa M., Regularity of abelian varieties I, Journal of the AMS 16 (2002) 285–302
- [PP2] Pareschi, G., Popa M., Regularity of abelian varieties II: basic results on linear series and defining equations, J. Alg. Geom. (2004) 167–193
- [PP3] Pareschi, G., Popa, M., GV-sheaves, Fourier–Mukai transform, and generic vanishing, Amer. J. Math. 133 (2011), 235–271
- [PS] Pareschi, G., Salvati Manni, R., 2-torsion points on theta divisors, International Mathematics Research Notices, Volume 2021 Issue 19 (2021) p. 14616 – 14628
- [W] Wahl, J., Introduction to Gaussian maps on an algebraic curve, Complex projective geometry, London Math. Soc. Lecture Note Ser., 179, Cambridge Univ. Press, Cambridge, (1992), 304–323.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROME, TOR VERGATA, ITALY

Email address: nelson.alvarado@ug.uchile.it

Email address: pareschi@mat.uniroma2.it