

HEAT KERNELS AND GREEN FUNCTIONS FOR FRACTIONAL SCHRÖDINGER OPERATORS WITH CONFINING POTENTIALS

XIN CHEN, KAMIL KALETA, JIAN WANG

ABSTRACT. We give two-sided, global (in all variables) estimates of the heat kernel and the Green function of the fractional Schrödinger operator with a non-negative and locally bounded potential V such that $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$. We assume that V is comparable to a radial profile with the doubling property. Our bounds are sharp with respect to spatial variables and qualitatively sharp with respect to time. The methods we use combine probabilistic and analytic arguments. They are based on the strong Markov property and the Feynman–Kac formula.

Keywords: fractional Schrödinger operator; confining potential; heat kernel; Green function; Feynman–Kac formula; ground state

MSC 2020: 35J10; 47D08; 35K08; 60G52; 81Q10.

1. INTRODUCTION AND MAIN RESULTS

Let $d \in \mathbb{N} := \{1, 2, \dots\}$ and $\alpha \in (0, 2)$. Let $\mathcal{L} := -(-\Delta)^{\alpha/2}$ be the *fractional Laplacian* of order α – the non-local, pseudo-differential operator which is defined by

$$\mathcal{L}f(x) = C_{d,\alpha} \int_{\mathbb{R}^d} (f(x+z) - f(x) - \langle \nabla f(x), z \rangle) \mathbf{1}_{\{|z| \leq 1\}} \frac{1}{|z|^{d+\alpha}} dz, \quad f \in C_c^\infty(\mathbb{R}^d)$$

with

$$C_{d,\alpha} = \frac{\alpha 2^{\alpha-1} \Gamma((d+\alpha)/2)}{\pi^{d/2} \Gamma(1-\alpha/2)}.$$

It is well known that \mathcal{L} is an infinitesimal generator of the rotationally invariant α -stable Lévy process $X := \{X_t\}_{t \geq 0}$ in \mathbb{R}^d , see e.g. [23] for various definitions of \mathcal{L} and the probabilistic background.

The main goal of the paper is to find global, two-sided estimates of the heat kernel and the Green function associated with the Schrödinger operator

$$(1.1) \quad \mathcal{L}^V = \mathcal{L} - V.$$

Here and in what follows, the potential $V : \mathbb{R}^d \rightarrow [0, \infty)$ is a sufficiently regular, non-negative and locally bounded function such that $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$. Therefore, \mathcal{L}^V extends to a (negative) self-adjoint operator on $L^2(\mathbb{R}^d) := L^2(\mathbb{R}^d, dx)$, see e.g. [25, Proposition 10.22].

The operator $-\mathcal{L}^V$ can be interpreted as a Hamiltonian in the fractional version of the classical model of *oscillator*, where \mathcal{L}^V takes the form $\Delta - V$, with Δ denoting the standard Laplacian and V being a potential as defined above. The most classical case, $V(x) = c|x|^2$ (where $c > 0$), corresponds to the harmonic oscillator. However, potentials exhibiting different growth rates at infinity are also of interest, such as the anharmonic (quartic) [1] or logarithmic [4] oscillators. In these models, particles are confined by forces corresponding to potentials V that grow to infinity at infinity. Such potentials are, therefore, commonly referred to as *confining* potentials in the literature. Fractional operators, especially in the *relativistic* case with $\alpha = 1$, play a significant role in contemporary mathematical physics (see, e.g., [16, 24]).

X. Chen: Department of Mathematics, Shanghai Jiao Tong University, 200240 Shanghai, P.R. China. chenxin217@sjtu.edu.cn.

K. Kaleta: Wrocław University of Science and Technology, Wybrzeże St. Wyspiańskiego 27, 50-370 Wrocław, Poland. kamil.kaleta@pwr.edu.pl.

J. Wang: School of Mathematics and Statistics & Key Laboratory of Analytical Mathematics and Applications (Ministry of Education) & Fujian Provincial Key Laboratory of Statistics and Artificial Intelligence, Fujian Normal University, 350007 Fuzhou, P.R. China. jianwang@fjnu.edu.cn.

The Schrödinger semigroup corresponding to the operator \mathcal{L}^V is denoted by $\{T_t^V\}_{t \geq 0}$; it fulfills the Feynman–Kac formula

$$(1.2) \quad T_t^V f(x) = \mathbb{E}_x \left[\exp \left(- \int_0^t V(X_s) ds \right) f(X_t) \right], \quad f \in L^2(\mathbb{R}^d) \text{ or } f \in C_b(\mathbb{R}^d)$$

(\mathbb{P}_x and \mathbb{E}_x denote the probability measure and the expectation with respect to starting position $x \in \mathbb{R}^d$, respectively). Moreover, there exists a jointly continuous and symmetric integral kernel $p : (0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow (0, \infty)$ such that

$$(1.3) \quad T_t^V f(x) = \int_{\mathbb{R}^d} p(t, x, y) f(y) dy, \quad t > 0, f \in L^2(\mathbb{R}^d) \text{ or } f \in C_b(\mathbb{R}^d),$$

see e.g. the monograph by Demuth and van Casteren [15, Theorem 2.5] for a reference. Since $V \geq 0$,

$$p(t, x, y) \leq q(t, x, y), \quad t > 0, x, y \in \mathbb{R}^d,$$

where $q(t, x, y)$ is the heat kernel associated with the fractional Laplacian \mathcal{L} (or, equivalently, the transition density of the α -stable process X). We remark that there are constants $c_1, c_2 > 0$ such that

$$(1.4) \quad c_1 \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) \leq q(t, x, y) \leq c_2 \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right), \quad t > 0, x, y \in \mathbb{R}^d,$$

see [2, Theorem 2.1] and [3, Proof of Lemma 5].

Throughout the paper we use the following assumption on the regularity and growth of V .

Assumption (H)

There exist a non-decreasing function $g : [0, \infty) \rightarrow [1, \infty)$, satisfying $\lim_{r \rightarrow \infty} g(r) = \infty$, and constants $C_1, C_2, C_3 > 0$ such that

$$(1.5) \quad g(2r) \leq C_1 g(r), \quad r \geq 0,$$

and

$$(1.6) \quad C_2 g(|x|) \leq V(x) \leq C_3 g(|x|), \quad |x| \geq 1.$$

Though the theory of non-local Schrödinger operators with confining potentials and their semigroups is an active area of research, existing results are mainly focused on intrinsic contractivity properties and ground state estimates. For basic definitions, we refer the reader to Davies and Simon [14]. A sharp necessary and sufficient condition for the intrinsic ultracontractivity of the non-local Schrödinger operator \mathcal{L}^V , driven by relativistic and usual rotationally invariant α -stable processes, was provided by Kulczycki and Siudeja [20], and by Kaleta and Kulczycki [18], respectively. Later, Kaleta and Lórinzi [19] extended these results to non-local Schrödinger operators \mathcal{L}^V associated with the class of symmetric Lévy processes possessing the direct jump property (DJP). The authors developed a new iteration technique to establish eigenfunction estimates for \mathcal{L}^V , including two-sided sharp bounds for the ground state, and characterized the asymptotic version (valid for large times) of intrinsic ultracontractivity. Chen and Wang [6] applied the intrinsic super-Poincaré inequality to establish a sharp criterion for intrinsic ultracontractivity and derived two-sided estimates for the ground state of non-local Schrödinger operators associated with general symmetric jump processes. Their analysis accounted for both finite-range and long-range jump behaviors. Furthermore, by employing a similar approach, the authors investigated various hyperboundedness properties of the associated Schrödinger semigroups [5]. More recently, Kaleta and Schilling [20] further analyzed the DJP class, providing large-time heat kernel estimates for the corresponding non-local Schrödinger operators with confining potentials. These estimates were established for $t \geq T_0$, with $T_0 > 0$ properly chosen. However, to the best of our knowledge, *global estimates of the heat kernel $p(t, x, y)$ for the full time regime, as well as estimates for the Green function $G(x, y)$ associated with \mathcal{L}^V , remain unknown.* In this paper, we address this problem comprehensively. We focus on the fractional Schrödinger operator \mathcal{L}^V and prove two-sided estimates for $p(t, x, y)$ over the full time horizon, as well as two-sided estimates for the Green function $G(x, y)$.

To state our first main result, we need to introduce some more notation. Define

$$(1.7) \quad t_0(s) := \inf \left\{ t > 0 : \exp(-tg(s)) = \frac{t}{(1+s)^\alpha} \right\}, \quad s \geq 0.$$

Note that, for any fixed $s \geq 0$, the function $t \mapsto e^{-tg(s)}$ is strictly decreasing and onto the interval $(0, 1)$, while $t \mapsto \frac{t}{(1+s)^\alpha}$ is strictly increasing, onto $(0, \infty)$. Then, $t_0(s)$ is unique in the sense that for any $t < t_0(s)$, $\exp(-tg(s)) > \frac{t}{(1+s)^\alpha}$, and for $t > t_0(s)$, $\exp(-tg(s)) < \frac{t}{(1+s)^\alpha}$. See Lemma 2.3 below for further estimates of $t_0(s)$.

Also, we say that the function $t_0 : [0, \infty) \rightarrow (0, \infty)$ is almost increasing (resp. decreasing), if there exists a continuous, strictly increasing (resp. strictly decreasing) function $h : [0, \infty) \rightarrow (0, \infty)$ and constants $C_*, C_{**} \geq 1$ such that

$$(1.8) \quad C_*^{-1}h(s) \leq t_0(s) \leq C_*h(s), \quad C_{**}^{-1}h(s) \leq h(2s) \leq C_{**}h(s), \quad s \geq 0.$$

For any $a, b \geq 0$, $a \wedge b := \min\{a, b\}$ and $a \vee b := \max\{a, b\}$.

Under Assumption **(H)** it holds that $\lim_{|x| \rightarrow \infty} V(x) = \infty$. Then, according to [27, Corollary 1.3] (see also [6, Proposition 1.1] or [18, Lemma 1]), the operator $T_t^V : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ is compact for every $t > 0$. Hence, by general theory of semigroups of compact operators, there exists an orthonormal basis in $L^2(\mathbb{R}^d)$, consisting of eigenfunctions $\{\phi_n\}_{n \geq 1}$ of the operator $-\mathcal{L}^V$, with associated eigenvalues $\{\lambda_n\}_{n \geq 1}$ satisfying $0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots$ and $\lim_{n \rightarrow \infty} \lambda_n = \infty$. We usually call ϕ_1 the ground state of $-\mathcal{L}^V$. In particular, for all $t > 0$ and $x \in \mathbb{R}^d$,

$$(1.9) \quad T_t^V \phi_1(x) = e^{-\lambda_1 t} \phi_1(x).$$

Furthermore, it is known that $\phi_1 : \mathbb{R}^d \rightarrow \mathbb{R}_+$ has a version which is continuous and strictly positive (see e.g. [6, Proposition 1.2]). Then, by [18, Corollary 3], for all $x \in \mathbb{R}^d$,

$$c_1 H(x) \leq \phi_1(x) \leq c_2 H(x),$$

where

$$H(x) := \frac{1}{g(|x|)(1+|x|)^{d+\alpha}}.$$

This, along with (1.9), yields that

$$(1.10) \quad c_3 e^{-\lambda_1 t} H(x) \leq T_t^V H(x) \leq c_4 e^{-\lambda_1 t} H(x), \quad x \in \mathbb{R}^d, \quad t > 0.$$

The following theorem is the first main result of the paper.

Theorem 1.1. *Let Assumption **(H)** hold. Then there are constants $C_0, \tilde{C}_0, C_1, \dots, C_{12} > 0$ such that for every $x, y \in \mathbb{R}^d$ and $t > 0$ the following statements hold.*

(1) *If $0 < t \leq C_0 t_0(|x| \wedge |y|)$, then*

$$(1.11) \quad \begin{aligned} C_1 \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) \left(\frac{1}{tg(|x| \vee |y|)} \wedge 1 \right) e^{-C_2 tg(|x| \wedge |y|)} &\leq p(t, x, y) \\ &\leq C_3 \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) \left(\frac{1}{tg(|x| \vee |y|)} \wedge 1 \right) e^{-C_4 tg(|x| \wedge |y|)}. \end{aligned}$$

(2-i) *If $t_0(\cdot)$ is almost increasing, then, for $t > C_0 t_0(|x| \wedge |y|)$,*

$$(1.12) \quad C_5 e^{-\lambda_1 t} H(x) H(y) \leq p(t, x, y) \leq C_6 e^{-\lambda_1 t} H(x) H(y).$$

(2-ii) *If $t_0(\cdot)$ is almost decreasing, then, for $C_0 t_0(|x| \wedge |y|) < t \leq \tilde{C}_0$,*

$$(1.13) \quad \begin{aligned} C_7 H(x) H(y) \int_{\{|z| \leq s_0(t)\}} e^{-C_6 tg(|z|)} dz &\leq p(t, x, y) \\ &\leq C_8 H(x) H(y) \int_{\{|z| \leq s_0(C_9 t)\}} e^{-C_{10} tg(|z|)} dz. \end{aligned}$$

where $s_0(t) = h^{-1}(t) \vee 2$, $h^{-1}(t) := \inf\{s \geq 0 : h(s) \leq t\}$ with h given in (1.8), and we use the convention that $\inf \emptyset = -\infty$; for every $t > \tilde{C}_0$,

$$(1.14) \quad C_{11}e^{-\lambda_1 t}H(x)H(y) \leq p(t, x, y) \leq C_{12}e^{-\lambda_1 t}H(x)H(y).$$

We now illustrate our result with the following example.

Example 1.2. *Let*

$$V(x) = \log^\beta(1 + |x|), \quad \beta > 0.$$

Then there are constants $C_0, \dots, C_{10} > 0$ such that for every $x, y \in \mathbb{R}^d$ with $|x| \leq |y|$ and $t > 0$ we have the following estimates.

(1) *If $0 < t \leq C_0 \log^{1-\beta}(2 + |x|)$, then*

$$\begin{aligned} C_1 \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) \left(\frac{1}{t \log^\beta(2 + |y|)} \wedge 1 \right) e^{-C_2 t \log^\beta(2 + |x|)} &\leq p(t, x, y) \\ &\leq C_3 \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) \left(\frac{1}{t \log^\beta(2 + |y|)} \wedge 1 \right) e^{-C_4 t \log^\beta(2 + |x|)}. \end{aligned}$$

(2-i) *If $t \geq C_0 \log^{1-\beta}(2 + |x|)$ and $\beta \leq 1$, then*

$$C_5 e^{-\lambda_1 t} H(x)H(y) \leq p(t, x, y) \leq C_6 e^{-\lambda_1 t} H(x)H(y).$$

Here and in what follows,

$$H(x) = \frac{1}{\log^\beta(2 + |x|)(1 + |x|)^{d+\alpha}}, \quad x \in \mathbb{R}^d.$$

(2-ii) *If $t \geq C_0 \log^{1-\beta}(2 + |x|)$ and $\beta > 1$, then*

$$\begin{aligned} C_7 \left(e^{C_8 t^{-1/(\beta-1)}} \mathbf{1}_{\{t \leq 1\}} + e^{-\lambda_1 t} \mathbf{1}_{\{t > 1\}} \right) H(x)H(y) &\leq p(t, x, y) \\ &\leq C_9 \left(e^{C_{10} t^{-1/(\beta-1)}} \mathbf{1}_{\{t \leq 1\}} + e^{-\lambda_1 t} \mathbf{1}_{\{t > 1\}} \right) H(x)H(y). \end{aligned}$$

Our second main result gives estimates for the Green function $G^V(x, y) := \int_0^\infty p(t, x, y) dt$. In what follows the notation $f(x) \asymp g(x)$, $x \in A$, means that there is a constant $c \geq 1$ such that $(1/c)g(x) \leq f(x) \leq cg(x)$, $x \in A$ (we also write “ $\stackrel{c}{\asymp}$ ” to indicate the comparison constant c).

Theorem 1.3. *Let Assumption (H) hold. If $t_0(\cdot)$ is an almost monotone function, then*

$$G^V(x, y) \asymp \frac{1}{|x-y|^{d-\alpha}} \left(1 \wedge \frac{1}{g(|x||x-y|^\alpha)} \right) \left(1 \wedge \frac{1}{g(|y||x-y|^\alpha)} \right) Q(x, y), \quad x, y \in \mathbb{R}^d, \quad x \neq y.$$

where

$$Q(x, y) := \begin{cases} 1, & \alpha < d, \\ \max \left\{ \log \left(\frac{1}{|x-y|^\alpha g(|x| \wedge |y|)} \right), 1 \right\}, & \alpha = d = 1, \\ \max \left\{ \left(\frac{1}{|x-y|^\alpha g(|x| \wedge |y|)} \right)^{\frac{\alpha-d}{\alpha}}, 1 \right\}, & \alpha > d = 1. \end{cases}$$

Next, we will make some remarks on our main results above.

(1) In order to make the presentation and proofs in this paper as clear and readable as possible, we decided to focus on the case when $X := \{X_t\}_{t \geq 0}$ is the rotationally invariant α -stable process. This choice also makes our contribution accessible to a wider audience. However, we believe that the arguments we propose apply to a large class of symmetric jump processes. In fact, it is straightforward to observe that the expression (2.27) for the heat kernel $p(t, x, y)$ relies solely on the strong Markov property of X and the Feynman-Kac formula (1.2). Using this, along with the Lévy system (2.15) and Dirichlet heat kernel estimates for more general symmetric jump processes established in [17, 21], it is possible to apply suitable iteration arguments (by choosing an appropriate domain

U when applying (2.27)) to derive two-sided estimates for the Schrödinger heat kernel $p(t, x, y)$ associated with such processes.

- (2) As mentioned above, the expression (2.27) serves as the starting point for our proofs. It allows us to derive appropriate estimates for the kernel $p(t, x, y)$ by investigating the probabilities that the process X visits various positions where the potential V takes specific values. In contrast to [5], where the classical Schrödinger semigroup was studied, here we apply the Lévy system formula (2.15) to estimate the joint distribution (τ_U, X_{τ_U}) in (2.27). Our approach here is different because our process X is a jump process. We also note that our argument in this paper differs significantly from that in [20], where the DJP and the estimates for harmonic functions played a crucial role.
- (3) According to Theorem 1.1, we know that there exists a threshold time $C_0 t_0(|x| \wedge |y|)$ for the two-sided estimates of the heat kernel $p(t, x, y)$. Intuitively, if $t \leq C_0 t_0(|x| \wedge |y|)$, the typical path from x to y that maximizes the probability for the process X evolving in the presence of the potential V follows the path of the free process X (i.e., when $V = 0$), and the effect of the potential along this path is expressed only through the local values of V near x and y . When $t > C_0 t_0(|x| \wedge |y|)$, the corresponding path that maximizes the probability will first travel from x to a domain near the origin, where the effect of the potential V is small, and then proceed from this domain to y , where the effect of the potential V is more significant. For example, see the integral terms in (1.13).

The rest of the paper is organized as follows. In the next section, we present some necessary preliminaries for the proof of Theorem 1.1. In particular, we establish an upper bound for $T_t^V 1(x)$ and some rough off-diagonal estimate of $p(t, x, y)$. Section 3 is devoted to two-sided estimates of $p(t, x, y)$ when $0 < t \leq C'_0 t_0(|x| \wedge |y|)$ for any $C'_0 > 0$. In Section 4, we verify two-sided estimates of $p(t, x, y)$ when $t \geq C_0 t_0(|x| \wedge |y|)$ for some $C_0 > 0$. In Section 5, we provide the proof of Example 1.2. Finally, in Section 6, we present the proofs of Theorems 1.1 and 1.3, based on our findings from Sections 3 and 4.

2. PRELIMINARIES

The following identity, called the Lévy system formula, describes the structure of jumps of the (rotationally invariant) α -stable process $X := \{X_t\}_{t \geq 0}$, see e.g. [10, Lemma 4.7].

Lemma 2.1. *Suppose that $f : \mathbb{R}_+ \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$ is a nonnegative measurable function so that $f(s, x, x) = 0$ for all $x \in \mathbb{R}^d$ and $s \in \mathbb{R}_+$. Then, for every $x \in \mathbb{R}^d$ and stopping time τ , it holds that*

$$(2.15) \quad \mathbb{E}_x \left[\sum_{s \leq \tau} f(s, X_{s-}, X_s) \right] = \mathbb{E}_x \left[\int_0^\tau \int_{\mathbb{R}^d} f(s, X_{s-}, z) \frac{C_{d,\alpha}}{|X_{s-} - z|^{d+\alpha}} dz ds \right],$$

where $X_{s-} := \lim_{\substack{t \rightarrow s \\ t < s}} X_t$ denotes the left limit of X . at s .

For any open subset $D \subset \mathbb{R}^d$ and $x \in D$, we define by

$$\delta_D(x) := \inf\{|x - z| : z \in D^c\}$$

the distance from $x \in D$ to the boundary ∂D . Let $\tau_D := \inf\{t \geq 0 : X_t \notin D\}$ be the first exit time of the process X from the set D . Denote by $q_D(t, x, y)$ the transition density of X killed upon exiting D . The following two-sided estimates for $q_{B(x,R)}(t, x, y)$ essentially have been proved in [9].

Lemma 2.2. *For any constant $C'_0 > 0$ there exists a positive constant C_1 such that for every $x \in \mathbb{R}^d$, $R > 0$, $y \in B(x, R)$ and $0 < t \leq C'_0 R^\alpha$,*

$$(2.16) \quad q_{B(x,R)}(t, x, y) \underset{C_1}{\asymp} \left(t^{-d/\alpha} \wedge \frac{t}{|x - y|^{d+\alpha}} \right) \left(\frac{\delta_{B(x,R)}(y)^{\alpha/2}}{\sqrt{t}} \wedge 1 \right).$$

Proof. Due to the spatial homogeneity of the process X , it suffices to prove (2.16) for $x = 0$. According to [9, Theorem 1.1], for every $C'_0 > 0$ there exists a positive constant c_1 such that for all $0 < t \leq C'_0$ and $y \in B(0, 1)$,

$$(2.17) \quad q_{B(0,1)}(t, 0, y) \stackrel{c_1}{\geq} \left(t^{-d/\alpha} \wedge \frac{t}{|y|^{d+\alpha}} \right) \left(\frac{\delta_{B(0,1)}(y)^{\alpha/2}}{\sqrt{t}} \wedge 1 \right).$$

Furthermore, we note that $\delta_{B(0,1)}(y) = 1 - |y|$ for all $y \in B(0, 1)$. Combining this and (2.17) with the scaling property

$$q_{B(0,1)}(t, 0, y) = R^d q_{B(0,R)}(R^\alpha t, 0, Ry), \quad t > 0, y \in B(0, 1),$$

we obtain (2.16) immediately. \square

We also give a lemma which describes the properties of $t_0(\cdot)$ and $g(\cdot)$.

Lemma 2.3. *Let $t_0(\cdot)$ be defined by (1.7) and assume that $g(\cdot)$ satisfies (1.5) and (1.6). Then there are constants $C_2, q_1 > 0$ such that for all $s \geq 0$,*

$$(2.18) \quad g(s) \leq C_2(1+s)^{q_1},$$

and

$$(2.19) \quad C_2^{-1}(1+s)^{-q_1} \leq t_0(s) \leq C_2 \left(1 + \frac{\log(2+s)}{g(s)} \right).$$

Proof. If $s > 1$, then $s \in (2^n, 2^{n+1}]$ for some $n \in \{0, 1, 2, \dots\}$. By the monotonicity of g and (1.5) (we may assume that $c_0 > 1$, where c_0 denotes the comparability constant in (1.5)),

$$g(s) \leq g(2^{n+1}) \leq c_0^{n+1} g(1) = c_0 g(1) 2^{n \log_2 c_0} \leq c_1 s^{q_1} \leq c_1 (1+s)^{q_1}$$

with $c_1 := c_0 g(1)$ and $q_1 := \log_2 c_0$. If $s \in [0, 1]$, then $g(s) \leq g(1) \leq c_1 \leq c_1 (1+s)^{q_1}$. By all of the above estimates we obtain (2.18).

By (1.7) it holds that $\exp(-t_0(s)g(s)) = t_0(s)(1+s)^{-\alpha}$, so $t_0(s)g(s) + \log t_0(s) = \alpha \log(1+s)$. If $t_0(s) \leq c_2 \log 2$ holds for some $c_2 > 0$ that satisfies $c_2 \log 2 \geq 1$, then the second inequality in (2.19) follows. If $t_0(s) \geq c_2 \log 2 (\geq 1)$, then $t_0(s) = \frac{\alpha \log(1+s) - \log t_0(s)}{g(s)} \leq \frac{\alpha \log(1+s)}{g(s)}$ for $s > 0$. Putting both estimates above together, we prove the second inequality in (2.19).

By the fact that $g(r) \geq 1$ for all $r \geq 0$, there is $a_0 > 0$ large enough such that for all $s > 0$,

$$(2.20) \quad (a_0 g(s))^{-1} (1+s)^{-\alpha} \leq e^{-a_0^{-1}} = \exp(-(a_0 g(s))^{-1} g(s)).$$

By (1.7) again, $t_0(s) \geq (a_0 g(s))^{-1} \geq c_3 (1+s)^{-q_1}$ for every $s > 0$. Then, we prove the first inequality in (2.19). \square

Next, we present upper bounds for $T_t^V \mathbf{1}(x)$, which will be frequently used in the paper.

Lemma 2.4. *There exist positive constants C_3, C_4 such that for any $t > 0$ and $x \in \mathbb{R}^d$,*

$$(2.21) \quad T_t^V \mathbf{1}(x) \leq C_3 \left(\exp(-C_4 t g(|x|)) + \frac{t}{(1+|x|)^\alpha} \right).$$

Proof. Since

$$\inf_{x \in B(0,2), t > 0} \left(\exp(-C_4 t g(|x|)) + \frac{t}{(1+|x|)^\alpha} \right) \geq c_0,$$

(2.21) holds trivially for every $x \in \mathbb{R}^d$ with $|x| \leq 2$ by taking C_3 large enough. Then, it suffices to prove (2.21) for every $x \in \mathbb{R}^d$ with $|x| \geq 2$. Set $U := B(x, |x|/2)$. According to (1.2),

$$\begin{aligned} T_t^V \mathbf{1}(x) &= \mathbb{E}_x \left[\exp \left(- \int_0^t V(X_s) ds \right) \right] \\ &= \mathbb{E}_x \left[\exp \left(- \int_0^t V(X_s) ds \right) \mathbf{1}_{\{\tau_U \geq t\}} \right] + \mathbb{E}_x \left[\exp \left(- \int_0^t V(X_s) ds \right) \mathbf{1}_{\{\tau_U < t\}} \right] \\ &=: I_1 + I_2. \end{aligned}$$

Obviously, by (1.5) and (1.6),

$$I_1 \leq \exp \left(-t \inf_{z \in U} V(z) \right) \leq \exp(-c_1 t g(|x|)).$$

On the other hand,

$$\begin{aligned} I_2 &= \mathbb{E}_x \left[\exp \left(- \int_0^t V(X_s) ds \right) \mathbb{1}_{\{\tau_U < t, X_t \in B(x, |x|/3)\}} \right] \\ &\quad + \mathbb{E}_x \left[\exp \left(- \int_0^t V(X_s) ds \right) \mathbb{1}_{\{X_t \notin B(x, |x|/3)\}} \right] \\ &=: I_{21} + I_{22}. \end{aligned}$$

By the strong Markov property,

$$\begin{aligned} I_{21} &\leq \mathbb{E}_x \left[\mathbb{P}_{X_{\tau_U}}(X_{t-\tau_U} \in B(x, |x|/3)) \mathbb{1}_{\{\tau_U < t\}} \right] \leq \sup_{\substack{0 \leq s \leq t \\ z \in B(x, |x|/2)^c}} \int_{B(x, |x|/3)} q(s, z, y) dy \\ &\leq c_2 \sup_{z \in B(x, |x|/2)^c} \int_{B(x, |x|/3)} t |z - y|^{-d-\alpha} dy \leq \frac{c_3 t}{(1 + |x|)^\alpha}, \end{aligned}$$

where in the third inequality we used (1.4). Similarly, by using (1.4) once again,

$$I_{22} \leq \int_{B(x, |x|/3)^c} q(t, x, y) dy \leq \frac{c_4 t}{(1 + |x|)^\alpha}.$$

Combining all the estimates above, we obtain the desired assertion. \square

Corollary 2.5. *There exist positive constants C_5 , C_6 and C_7 such that for all $x \in \mathbb{R}^d$ and $t \geq C_5 t_0(|x|)$,*

$$(2.22) \quad T_t^V 1(x) \leq C_6 \log(2 + |x|)(1 + |x|)^{-\alpha} \leq C_7(1 + |x|)^{-\alpha/2}.$$

Proof. According to (2.21), for every $\tilde{C}_0 > 0$ and $x \in \mathbb{R}^d$,

$$(2.23) \quad T_{\tilde{C}_0 t_0(|x|)}^V 1(x) \leq c_1 \left(\exp \left(-c_2 \tilde{C}_0 t_0(|x|) g(|x|) \right) + \frac{\tilde{C}_0 t_0(|x|)}{(1 + |x|)^\alpha} \right).$$

By (1.7) and (2.19), it holds that for all $x \in \mathbb{R}^d$,

$$\begin{aligned} (2.24) \quad \exp \left(-c_2 \tilde{C}_0 t_0(|x|) g(|x|) \right) &= \left(e^{-t_0(|x|) g(|x|)} \right)^{c_2 \tilde{C}_0} \\ &= \left(\frac{t_0(|x|)}{(1 + |x|)^\alpha} \right)^{c_2 \tilde{C}_0} \leq c_3 [\log(2 + |x|)]^{c_2 \tilde{C}_0} (1 + |x|)^{-c_2 \tilde{C}_0 \alpha}. \end{aligned}$$

On the other hand, by (2.19) again,

$$\frac{t_0(|x|)}{(1 + |x|)^\alpha} \leq c_4 \log(2 + |x|)(1 + |x|)^{-\alpha}.$$

Choosing $\tilde{C}_0 > c_2^{-1}$ and putting both estimates above into (2.23) yield that

$$T_{\tilde{C}_0 t_0(|x|)}^V 1(x) \leq c_5 \log(2 + |x|)(1 + |x|)^{-\alpha}.$$

Therefore, for every $t \geq \tilde{C}_0 t_0(|x|)$,

$$\begin{aligned} T_t^V 1(x) &= T_{\tilde{C}_0 t_0(|x|)}^V \left(T_{t-\tilde{C}_0 t_0(|x|)}^V 1 \right) (x) \\ &\leq T_{\tilde{C}_0 t_0(|x|)}^V 1(x) \leq c_5 \log(2 + |x|)(1 + |x|)^{-\alpha}. \end{aligned}$$

The proof of the first inequality in the desired assertion is complete. The second assertion is a direct consequence of the first one. \square

Remark 2.6. According to (1.7), for every $c_0 > 0$, $x \in \mathbb{R}^d$ and $0 < t \leq c_0 t_0(|x|)$,

$$(2.25) \quad \frac{t}{(1+|x|)^\alpha} = c_0 \frac{c_0^{-1}t}{(1+|x|)^\alpha} \leq c_0 \exp(-c_0^{-1}tg(|x|)).$$

Thus, by (2.21), for any $c_0 > 0$, there exist positive constants c_1 and c_2 (which may depend on c_0) such that for any $x \in \mathbb{R}^d$ and $0 < t \leq c_0 t_0(|x|)$,

$$(2.26) \quad T_t^V 1(x) \leq c_1 \exp(-c_2 tg(|x|)).$$

At the end of this section, we prove an upper bound for $p(t, x, y)$ when $|x - y| \geq Ct^{1/\alpha}$. For this we need the following representation of $p(t, x, y)$. Since the proof is almost the same as that [5, Lemma 2.1] (it only uses the strong Markov property of the process X and the Feynman–Kac formula (1.2)), we omit it here.

Lemma 2.7. *Suppose U is a domain in \mathbb{R}^d . Then, for every $x \in U$ and $y \notin \bar{U}$,*

$$(2.27) \quad p(t, x, y) = \mathbb{E}_x \left[\exp \left(- \int_0^{\tau_U} V(X_s) ds \right) \mathbf{1}_{\{\tau_U \leq t\}} p(t - \tau_U, X_{\tau_U}, y) \right].$$

Lemma 2.8. *For any $C'_0 > 0$ there is a constant $C_8 > 0$ such that for all $x, y \in \mathbb{R}^d$ and $t > 0$ with $|x - y| \geq 2C'_0 t^{1/\alpha}$,*

$$(2.28) \quad p(t, x, y) \leq \frac{C_8 t}{|x - y|^{d+\alpha}} \left(1 \wedge \frac{1}{t \max\{g(|x|), g(|y|)\}} \right).$$

Proof. Without loss of generality throughout the proof we assume that $|x| \leq |y|$. Fix any $C'_0 > 0$, and let $x, y \in \mathbb{R}^d$ and $t > 0$ be such that $|x - y| \geq 2C'_0 t^{1/\alpha}$. Define $U = B(y, c_0 t^{1/\alpha})$ with $c_0 := C'_0/6$ (in particular, $c_0 t^{1/\alpha} = (2C'_0 t^{1/\alpha})/12 \leq |x - y|/12$).

We first consider the case that $|y| \geq 3$. Set $W = B(y, |x - y|/3)$. Since $|x - y| \geq 12c_0 t^{1/\alpha}$, it holds that $U \subset W$. Hence, by (2.27),

$$\begin{aligned} p(t, x, y) &= \mathbb{E}_y \left[\exp \left(- \int_0^{\tau_U} V(X_s) ds \right) p(t - \tau_U, X_{\tau_U}, x) \mathbf{1}_{\{\tau_U \leq t\}} \right] \\ &= \mathbb{E}_y \left[\exp \left(- \int_0^{\tau_U} V(X_s) ds \right) p(t - \tau_U, X_{\tau_U}, x) \mathbf{1}_{\{\tau_U \leq t, X_{\tau_U} \in W \setminus U\}} \right] \\ &\quad + \mathbb{E}_y \left[\exp \left(- \int_0^{\tau_U} V(X_s) ds \right) p(t - \tau_U, X_{\tau_U}, x) \mathbf{1}_{\{\tau_U \leq t, X_{\tau_U} \notin W\}} \right] \\ &=: I_1 + I_2. \end{aligned}$$

By the inclusions $U \subset W \subset B(y, 2|y|/3)$ (the second one follows from $|x - y|/3 \leq (|x| + |y|)/3 \leq 2|y|/3$), (1.5) and (1.6), we have

$$(2.29) \quad \inf_{v \in U} V(v) \geq \inf_{v \in W} V(v) \geq \inf_{v \in B(y, 2|y|/3)} V(v) \geq c_1 g(|y|), \quad |y| \geq 3.$$

Therefore, by the Lévy system formula (2.15), (2.29) and the estimates

$$q_U(s, y, z) \leq c_2 \left(s^{-d/\alpha} \wedge \frac{s}{|y - z|^{d+\alpha}} \right) \left(\frac{\delta_U(z)^{\alpha/2}}{\sqrt{s}} \wedge 1 \right), \quad 0 < s \leq t, \quad z \in U,$$

$$p(t - s, u, x) \leq q(t - s, u, x) \leq \frac{c_3 t}{|u - x|^{d+\alpha}} \leq \frac{c_4 t}{|x - y|^{d+\alpha}}, \quad 0 < s \leq t, \quad u \in W,$$

(cf. (2.16) and (1.4)), we get

$$\begin{aligned} I_1 &\leq c_5 \int_0^t \int_{W \setminus U} \left(\int_U q_U(s, y, z) p(t - s, u, x) \frac{1}{|z - u|^{d+\alpha}} e^{-s \inf_{v \in U} V(v)} dz \right) du ds \\ &\leq \frac{c_6 t}{|x - y|^{d+\alpha}} \int_0^t \int_{W \setminus U} \int_U \left(s^{-d/\alpha} \wedge \frac{s}{|y - z|^{d+\alpha}} \right) \left(\frac{\delta_U(z)^{\alpha/2}}{\sqrt{s}} \wedge 1 \right) \frac{e^{-c_1 s g(|y|)}}{|z - u|^{d+\alpha}} dz du ds \end{aligned}$$

$$\begin{aligned}
 &\leq \frac{c_6 t}{|x-y|^{d+\alpha}} \int_0^t \int_U \left(s^{-d/\alpha} \wedge \frac{s}{|y-z|^{d+\alpha}} \right) \left(\frac{\delta_U(z)^{\alpha/2}}{\sqrt{s}} \wedge 1 \right) e^{-c_1 s g(|y|)} \\
 &\quad \times \int_{B(z, \delta_U(z))^c} \frac{1}{|z-u|^{d+\alpha}} du dz ds \\
 &\leq \frac{c_7 t}{|x-y|^{d+\alpha}} \int_0^t \int_U \left(s^{-d/\alpha} \wedge \frac{s}{|y-z|^{d+\alpha}} \right) \delta_U^{-\alpha}(z) \left(\frac{\delta_U(z)^{\alpha/2}}{\sqrt{s}} \wedge 1 \right) e^{-c_1 s g(|y|)} dz ds.
 \end{aligned}$$

Take now $c_8 := c_0/4$. In particular, $\delta_U(z) \geq c_9 t^{1/\alpha}$ for all $z \in B(y, c_8 s^{1/\alpha})$, and the expression on the right hand side above is less or equal to the sum

$$\begin{aligned}
 &\frac{c_{10} t}{|x-y|^{d+\alpha}} \int_0^t \int_{B(y, c_8 s^{1/\alpha})} s^{-d/\alpha} t^{-1} e^{-c_1 s g(|y|)} dz ds \\
 &+ \frac{c_{10} t}{|x-y|^{d+\alpha}} \int_0^t \int_{B(y, c_0 t^{1/\alpha} - c_8 s^{1/\alpha}) \setminus B(y, c_8 s^{1/\alpha})} \frac{s}{|y-z|^{d+\alpha}} \delta_U^{-\alpha}(z) e^{-c_1 s g(|y|)} dz ds \\
 &+ \frac{c_{10} t}{|x-y|^{d+\alpha}} \int_0^t \int_{B(y, c_0 t^{1/\alpha}) \setminus B(y, c_0 t^{1/\alpha} - c_8 s^{1/\alpha})} \frac{s}{|y-z|^{d+\alpha}} \frac{\delta_U^{-\alpha/2}(z)}{\sqrt{s}} e^{-c_1 s g(|y|)} dz ds \\
 &=: I_{11} + I_{12} + I_{13}.
 \end{aligned}$$

It is clear that

$$I_{11} \leq \frac{c_{11}}{|x-y|^{d+\alpha}} \int_0^t e^{-c_1 s g(|y|)} ds \leq \frac{c_{12} t}{|x-y|^{d+\alpha}} \left(\frac{1}{t g(|y|)} \wedge 1 \right).$$

On the other hand, for every $0 \leq s \leq t$,

$$\begin{aligned}
 &\int_{B(y, c_0 t^{1/\alpha} - c_8 s^{1/\alpha}) \setminus B(y, c_8 s^{1/\alpha})} \frac{\delta_U^{-\alpha}(z)}{|y-z|^{d+\alpha}} dz \\
 &\leq c_{13} \int_{c_8 s^{1/\alpha}}^{c_0 t^{1/\alpha} - c_8 s^{1/\alpha}} r^{-1-\alpha} (c_0 t^{1/\alpha} - r)^{-\alpha} dr \\
 &\leq c_{14} t^{-1} \int_{c_8 s^{1/\alpha}}^{c_0 t^{1/\alpha}/3} r^{-1-\alpha} dr + c_{15} t^{-(1+\alpha)/\alpha} \int_{c_0 t^{1/\alpha}/3}^{c_0 t^{1/\alpha} - c_8 s^{1/\alpha}} (c_0 t^{1/\alpha} - r)^{-\alpha} dr \\
 &\leq \frac{c_{15}}{t s} + \frac{c_{16}}{t^{(1+\alpha)/\alpha}} \left[t^{(1-\alpha)/\alpha} \mathbb{1}_{\{\alpha \in (0,1)\}} + \log \left(1 + \frac{t}{s} \right) \mathbb{1}_{\{\alpha=1\}} + s^{(1-\alpha)/\alpha} \mathbb{1}_{\{\alpha \in (1,2)\}} \right] \\
 &\leq \frac{c_{17}}{t} \left[\frac{1}{s} + \frac{1}{t} \mathbb{1}_{\{\alpha \in (0,1)\}} + \frac{1}{t} \log \left(1 + \frac{t}{s} \right) \mathbb{1}_{\{\alpha=1\}} + \left(\frac{s}{t} \right)^{1/\alpha} \frac{1}{s} \mathbb{1}_{\{\alpha \in (1,2)\}} \right] \leq \frac{c_{18}}{t s},
 \end{aligned}$$

where in the first inequality above we have used the spherical coordinates in $B(y, c_0 t^{1/\alpha})$. This yields

$$I_{12} \leq \frac{c_{19}}{|x-y|^{d+\alpha}} \int_0^t e^{-c_1 s g(|y|)} ds \leq \frac{c_{20} t}{|x-y|^{d+\alpha}} \left(1 \wedge \frac{1}{t g(|y|)} \right).$$

Furthermore, we have for every $0 \leq s \leq t$,

$$\begin{aligned}
 &\int_{B(y, c_0 t^{1/\alpha}) \setminus B(y, c_0 t^{1/\alpha} - c_8 s^{1/\alpha})} \frac{\delta_U^{-\alpha/2}(z)}{|y-z|^{d+\alpha}} dz \\
 &\leq c_{21} t^{-1-d/\alpha} \int_{B(y, c_0 t^{1/\alpha}) \setminus B(y, c_0 t^{1/\alpha} - c_8 s^{1/\alpha})} \delta_U^{-\alpha/2}(z) dz \\
 &\leq c_{22} t^{-1-d/\alpha} \int_{c_0 t^{1/\alpha} - c_8 s^{1/\alpha}}^{c_0 t^{1/\alpha}} (c_0 t^{1/\alpha} - r)^{-\alpha/2} r^{d-1} dr \\
 &\leq c_{23} t^{-1-1/\alpha} s^{1/\alpha-1/2} \leq c_{23} t^{-1} s^{-1/2}.
 \end{aligned}$$

This gives us that

$$I_{13} \leq \frac{c_{24}}{|x-y|^{d+\alpha}} \int_0^t e^{-c_1 s g(|y|)} ds \leq \frac{c_{25} t}{|x-y|^{d+\alpha}} \left(1 \wedge \frac{1}{tg(|y|)}\right).$$

Combining with all estimates above, we arrive at

$$I_1 \leq \frac{c_{26} t}{|x-y|^{d+\alpha}} \left(1 \wedge \frac{1}{tg(|y|)}\right).$$

According to the Lévy system formula (2.15) and (2.29), we get

$$\begin{aligned} I_2 &\leq \int_0^t \int_{W^c} \int_U q_U(s, y, z) p(t-s, u, x) \frac{1}{|z-u|^{d+\alpha}} e^{-s \inf_U V} dz du ds \\ &\leq \frac{c_{27}}{|x-y|^{d+\alpha}} \int_0^t \left(\int_U q_U(s, y, z) dz \right) \left(\int_{\mathbb{R}^d} p(t-s, u, x) du \right) e^{-c_{28} s g(|y|)} ds \\ &\leq \frac{c_{29} t}{|x-y|^{d+\alpha}} \left(1 \wedge \frac{1}{tg(|y|)}\right), \end{aligned}$$

where in the second inequality we used the fact that for $z \in U$ and $u \in W^c$,

$$|z-u| \geq |u-y| - |y-z| \geq |x-y|/3 - c_0 t^{1/\alpha} \geq |x-y|/3 - |x-y|/12 = |x-y|/4.$$

Therefore, we have proved (2.28) for the case $|y| \geq 3$.

When $|y| \leq 3$, the condition $|x-y| \geq 2C'_0 t^{1/\alpha}$ implies that

$$t \leq \left(\frac{|x-y|}{2C'_0} \right)^\alpha \leq c_{30} (|x| + |y|)^\alpha \leq 2^\alpha c_{30} |y|^\alpha \leq c_{31},$$

and so

$$p(t, x, y) \leq q(t, x, y) \leq \frac{c_{32} t}{|x-y|^{d+\alpha}} \leq \frac{c_{33} t}{|x-y|^{d+\alpha}} \left(1 \wedge \frac{1}{tg(|y|)}\right).$$

The proof is finished. \square

3. TWO-SIDED ESTIMATES WHEN $0 < t \leq C'_0 t_0 (|x| \wedge |y|)$ FOR ANY $C'_0 > 0$

Throughout this section, without loss of generality, we assume that $x, y \in \mathbb{R}^d$ are such that $|x| \leq |y|$. We begin with the following statement for $x, y \in \mathbb{R}^d$ with $|x-y| \leq C'_1 t^{1/\alpha}$.

Lemma 3.1. *For any $C'_0, C'_1 > 0$ there exist positive constants C_1, \dots, C_4 such that for all $x, y \in \mathbb{R}^d$ and $0 < t \leq C'_0 t_0 (|x|)$ with $|x-y| \leq C'_1 t^{1/\alpha}$,*

$$(3.30) \quad C_1 t^{-d/\alpha} e^{-C_2 t g(|x|)} \leq p(t, x, y) \leq C_3 t^{-d/\alpha} e^{-C_4 t g(|x|)}.$$

Proof. Suppose that $x, y \in \mathbb{R}^d$ and $0 < t \leq C'_0 t_0 (|x|)$ are such that $|x-y| \leq C'_1 t^{1/\alpha}$. Then,

$$(3.31) \quad \begin{aligned} p(t, x, y) &= \int_{\mathbb{R}^d} p(t/2, x, z) p(t/2, z, y) dz \leq c_1 t^{-d/\alpha} \int_{\mathbb{R}^d} p(t/2, x, z) dz \\ &= c_1 t^{-d/\alpha} T_{t/2}^V 1(x) \leq c_2 t^{-d/\alpha} e^{-c_3 t g(|x|)}, \end{aligned}$$

where in the last inequality we used (2.26).

On the other hand, since $|x-y| \leq C'_1 t^{1/\alpha}$, $y \in B(x, C'_1 t^{1/\alpha})$. Take $U := B(x, 3C'_1 t^{1/\alpha})$ and $f \in C_b(\mathbb{R}^d)$ with $\text{supp}[f] \subset B(x, 2C'_1 t^{1/\alpha})$. We have

$$\begin{aligned} T_t^V f(x) &= \mathbb{E}_x \left[f(X_t) \exp \left(- \int_0^t V(X_s) ds \right) \right] \geq \mathbb{E}_x \left[f(X_t) \exp \left(- \int_0^t V(X_s) ds \right) \mathbf{1}_{\{\tau_U > t\}} \right] \\ &\geq e^{-t \sup_{z \in U} V(z)} \mathbb{E}_x [f(X_t) \mathbf{1}_{\{\tau_U > t\}}] \geq e^{-c_4 t g(|x|)} \int_{B(x, 2C'_1 t^{1/\alpha})} q_U(t, x, z) f(z) dz \\ &\geq c_5 t^{-d/\alpha} e^{-c_4 t g(|x|)} \int_{B(x, 2C'_1 t^{1/\alpha})} f(z) dz = c_5 t^{-d/\alpha} e^{-c_4 t g(|x|)} \|f\|_1. \end{aligned}$$

Here in the third inequality we used the fact that

$$(3.32) \quad \sup_{z \in B(x, 3C'_0 t^{1/\alpha})} V(z) \leq \sup_{z \in B(x, c_6(1+|x|))} V(z) \leq c_7 g(|x|),$$

which is due to that $0 < t \leq t_0(|x|) \leq c_8(1+|x|)^\alpha$ (that can be verified by (2.19)), and the fourth inequality follows from (2.16). Since f is arbitrary, the inequality above yields that

$$p(t, x, y) \geq c_5 t^{-d/\alpha} e^{-c_4 t g(|x|)}.$$

Therefore, the proof of (3.30) is completed. \square

Lemma 3.2. *For any $C'_0, C'_1 > 0$ there exist positive constants C_5 and C_6 such that for $x, y \in \mathbb{R}^d$ and $0 < t \leq C'_0 t_0(|x|)$ with $|x - y| > C'_1 t^{1/\alpha}$,*

$$(3.33) \quad p(t, x, y) \geq C_5 \left(\frac{1}{t g(|y|)} \wedge 1 \right) e^{-C_6 t g(|x|)} \frac{t}{|x - y|^{d+\alpha}}.$$

Proof. Recall that $|x| \leq |y|$. Let $U = B(y, c_0 t^{1/\alpha})$ and $W = B(x, c_0 t^{1/\alpha})$ with $c_0 := C'_1/4$. Since $C'_1 t^{1/\alpha} < |x - y| \leq 2|y|$, we have $c_0 t^{1/\alpha} < |y|/2$. Consequently,

$$(3.34) \quad \sup_{z \in U} V(z) \leq \sup_{z \in B(y, |y|/2)} V(z) \leq c_1 g(|y|).$$

According to (2.27), the Lévy system formula (2.15), (3.30), (3.34) and the fact that

$$|z - u| \leq |x - y| + |z - x| + |u - y| \leq |x - y| + 2c_0 t^{1/\alpha} \leq c_2 |x - y|, \quad z \in W, u \in U,$$

we have

$$\begin{aligned} p(t, y, x) &= \mathbb{E}_y \left[p(t - \tau_U, X_{\tau_U}, x) \mathbf{1}_{\{\tau_U \leq t\}} \exp \left(- \int_0^{\tau_U} V(X_s) ds \right) \right] \\ &\geq \mathbb{E}_y \left[p(t - \tau_U, X_{\tau_U}, x) e^{-\int_0^{\tau_U} V(X_s) ds} \mathbf{1}_{\{\tau_U \leq t/2, X_{\tau_U} \in W\}} \right] \\ &\geq c_3 \left(\inf_{\substack{t/2 \leq s \leq t \\ z \in W}} p(s, z, x) \right) \int_0^{t/2} \int_U \int_W q_U(s, y, z) \frac{1}{|z - u|^{d+\alpha}} e^{-c_1 s g(|y|)} dz du ds \\ &\geq c_4 e^{-c_5 t g(|x|)} t^{-d/\alpha} \frac{|W|}{|x - y|^{d+\alpha}} \int_0^{t/2} \int_U q_U(s, y, z) dz e^{-c_1 s g(|y|)} ds \\ &\geq c_6 e^{-c_5 t g(|x|)} \frac{1}{|x - y|^{d+\alpha}} \int_0^{t/2} \mathbb{P}_y(\tau_{B(y, c_0 t^{1/\alpha})} > t/2) e^{-c_1 s g(|y|)} ds \\ &\geq c_7 e^{-c_5 t g(|x|)} \frac{1}{|x - y|^{d+\alpha}} \left(\frac{1}{g(|y|)} \wedge t \right) \\ &= c_7 \left(\frac{1}{t g(|y|)} \wedge 1 \right) e^{-c_5 t g(|x|)} \frac{t}{|x - y|^{d+\alpha}}. \end{aligned}$$

Here we also used the property

$$\mathbb{P}_y(\tau_{B(y, c_0 t^{1/\alpha})} > t) = \mathbb{P}_0(\tau_{B(0, c_0 t^{1/\alpha})} > t) \geq c_8 > 0,$$

which can be verified directly by (2.16). Hence, the conclusion (3.33) follows. \square

Lemma 3.3. *For any $C'_0, C'_1 > 0$ there exist positive constants C_7, C_8 such that for $x, y \in \mathbb{R}^d$ and $0 < t \leq C'_0 t_0(|x|)$ with $|x - y| > C'_1 t^{1/\alpha}$,*

$$(3.35) \quad p(t, x, y) \leq C_7 \left(\frac{1}{t g(|y|)} \wedge 1 \right) e^{-C_8 t g(|x|)} \frac{t}{|x - y|^{d+\alpha}}.$$

Proof. Recall we assume that $|x| \leq |y|$. The proof is split into two cases.

Case 1: $|y - x| \leq |y|/4$. In particular, $|x| \geq |y| - |y - x| \geq |y|/2$. We write

$$\begin{aligned} p(t, x, y) &= \int_{\{|z-y| \leq |x-y|/2\}} p(t/2, x, z) p(t/2, z, y) dz + \int_{\{|z-y| \geq |x-y|/2\}} p(t/2, x, z) p(t/2, z, y) dz \\ &=: I_1 + I_2. \end{aligned}$$

When $|y - z| \leq |x - y|/2$, we have $|z - x| \geq |x - y| - |y - z| \geq |x - y|/2 \geq C'_1 t^{1/\alpha}/2$ and $|z| \geq |y| - |y - z| \geq |y| - |x - y|/2 \geq |y| - |y|/8 \geq 7|y|/8$. Then, according to (2.28) and (1.5),

$$p(t/2, x, z) \leq c_1 \left(\frac{1}{t \max\{g(|x|), g(|z|)\}} \wedge 1 \right) \frac{t}{|x - z|^{d+\alpha}} \leq c_2 \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}}.$$

So for any $0 < t \leq C'_0 t_0(|x|)$ it holds that

$$\begin{aligned} I_1 &\leq c_2 \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} \int_{\mathbb{R}^d} p(t/2, z, y) dz \\ &\leq c_3 \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} \left(e^{-c_4 tg(|y|)} + \frac{t}{(1 + |y|)^\alpha} \right) \\ &\leq c_5 \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} \left(e^{-c_6 tg(|x|)} + \frac{t}{(1 + |x|)^\alpha} \right) \\ &\leq c_7 \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} e^{-c_8 tg(|x|)}, \end{aligned}$$

where the second inequality follows from (2.21), in the third inequality we have used the fact $|x| \leq |y|$, and the last inequality is due to the fact $0 < t \leq C'_0 t_0(|x|)$ and (2.25).

Similarly, by (2.28), for every $z, y \in \mathbb{R}^d$ with $|z - y| \geq |x - y|/2 \geq C'_1 t^{1/\alpha}/2$,

$$p(t/2, z, y) \leq c_9 \left(\frac{1}{t \max\{g(|y|), g(|z|)\}} \wedge 1 \right) \frac{t}{|z - y|^{d+\alpha}} \leq c_{10} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}}.$$

Hence, due to (2.26), it holds that for every $0 < t \leq C_0 t_0(|x|)$,

$$I_2 \leq c_{10} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} \int_{\mathbb{R}^d} p(t/2, x, z) dz \leq c_{11} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} e^{-c_{12} tg(|x|)}.$$

Therefore, putting all the estimates above for I_1 and I_2 together yields

$$p(t, x, y) \leq c_{13} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} e^{-c_{14} tg(|x|)}.$$

Case 2: $|y - x| > |y|/4$. Since we assume that $|x| \leq |y|$, it holds that $|y|/4 \leq |y - x| \leq |y| + |x| \leq 2|y|$. Then, we set

$$\begin{aligned} p(t, x, y) &= \int_{\{|z-y| \leq |y|/16\}} p(t/2, x, z) p(t/2, z, y) dz + \int_{\{|z-y| \geq |y|/16\}} p(t/2, x, z) p(t/2, z, y) dz \\ &=: J_1 + J_2. \end{aligned}$$

When $|z - y| \leq |y|/16$, we have $|z - x| \geq |x - y| - |y - z| \geq |x - y| - |y|/16 \geq |x - y| - |x - y|/4 \geq 3|x - y|/4 \geq 3C'_1 t^{1/\alpha}/4$ and $|z| \geq |y| - |z - y| \geq |y|/2$. Then, according to (2.28) and (1.5),

$$p(t/2, z, x) \leq c_{15} \left(\frac{1}{t \max\{g(|x|), g(|z|)\}} \wedge 1 \right) \frac{t}{|x - z|^{d+\alpha}} \leq c_{16} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}}.$$

Thus, following the argument for I_1 and using (2.26), we find that

$$J_1 \leq c_{16} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} \int_{\mathbb{R}^d} p(t/2, z, y) dz \leq c_{17} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} e^{-c_{18} tg(|x|)}.$$

Note that, when $|z - y| \geq |y|/16$, $|z - y| \geq |x - y|/32 \geq C'_1 t^{1/\alpha}/32$. Hence, according to (2.28), in this case we have

$$p(t/2, z, y) \leq c_{19} \left(\frac{1}{t \max\{g(|z|), g(|y|)\}} \wedge 1 \right) \frac{t}{|z - y|^{d+\alpha}} \leq c_{20} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}}.$$

Then, applying (2.26) again, we deduce that

$$J_2 \leq c_{20} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} \int_{\mathbb{R}^d} p(t/2, x, z) dz \leq c_{21} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x - y|^{d+\alpha}} e^{-c_{22} tg(|x|)}.$$

Putting the estimates for J_1 and J_2 together, we arrive at

$$p(t, x, y) \leq c_{23} \left(\frac{1}{tg(|y|)} \wedge 1 \right) \frac{t}{|x-y|^{d+\alpha}} e^{-c_{24}tg(|x|)}.$$

Therefore, the assertion follows. \square

Now we can summarise the results in this section as follows.

Proposition 3.4. *For any $C'_0 > 0$ there exist positive constants C_9, \dots, C_{12} such that for all $x, y \in \mathbb{R}^d$ with $0 < t \leq C'_0 t_0(|x|)$,*

$$(3.36) \quad \begin{aligned} C_9 \left(\frac{1}{tg(|y|)} \wedge 1 \right) e^{-C_{10}tg(|x|)} \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) &\leq p(t, x, y) \\ &\leq C_{11} \left(\frac{1}{tg(|y|)} \wedge 1 \right) e^{-C_{12}tg(|x|)} \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right). \end{aligned}$$

Proof. Let $C'_1 > 0$. When $x, y \in \mathbb{R}^d$ and $0 < t \leq C'_0 t_0(|x|)$ are such that $|x-y| \geq C'_1 t^{1/\alpha}$, the assertion follows directly from Lemmas 3.2 and 3.3. On the other hand, by Lemma 3.1, for all $x, y \in \mathbb{R}^d$ with $|x-y| \leq C'_1 t^{1/\alpha}$ and $0 < t \leq C'_0 t_0(|x|)$,

$$p(t, x, y) \geq c_1 e^{-c_2 tg(|x|)} t^{-d/\alpha} \geq c_1 \left(\frac{1}{tg(|y|)} \wedge 1 \right) e^{-c_2 tg(|x|)} t^{-d/\alpha}.$$

Furthermore, for all $x, y \in \mathbb{R}^d$ with $|x-y| \leq C'_1 t^{1/\alpha}$,

$$(3.37) \quad \begin{aligned} g(|x|) &\leq g(|y|) \leq g(|x| + |x-y|) \leq g(|x| + C'_1 t^{1/\alpha}) \\ &\leq g(|x| + C'_1 C_0'^{1/\alpha} t_0(|x|)^{1/\alpha}) \leq g(c_3(1 + |x|)) \leq c_4 g(|x|), \end{aligned}$$

and so

$$(3.38) \quad \frac{1}{tg(|y|)} \wedge 1 \geq c_5 \left(\frac{1}{tg(|x|)} \wedge 1 \right) \geq c_6 e^{-c_7 tg(|x|)},$$

where we used the monotonicity of $g(\cdot)$ and the fact that $t_0(|x|)^{1/\alpha} \leq c_8(1 + |x|)$. Thus, using (3.38) and Lemma 3.1 again, we obtain that for all $x, y \in \mathbb{R}^d$ with $|x-y| \leq C'_1 t^{1/\alpha}$ and $0 < t \leq C'_0 t_0(|x|)$,

$$p(t, x, y) \leq c_9 t^{-d/\alpha} e^{-c_{10}tg(|x|)} \leq c_{11} t^{-d/\alpha} \left(\frac{1}{tg(|y|)} \wedge 1 \right) e^{-c_{12}tg(|x|)}.$$

Putting all estimates above together yields the desired assertion (3.36). \square

4. TWO-SIDED ESTIMATES WHEN $t \geq C_0 t_0(|x| \wedge |y|)$ FOR SOME $C_0 > 0$

In this section, we will present two-sided estimates of $p(t, x, y)$ when $t > C_0 t_0(|x| \wedge |y|)$ for some properly chosen constant $C_0 > 0$. The statements are split into two parts, according to two different almost monotonicity properties of $t_0(\cdot)$. Without loss of generality, we still assume that $|x| \leq |y|$. Recall that

$$H(x) := \frac{1}{g(|x|)(1 + |x|)^{d+\alpha}}, \quad x \in \mathbb{R}^d.$$

We first give a lemma for a rough upper bound of the heat kernel $p(t, x, y)$ when $t > C_0 t_0(|x|)$ for some $C_0 > 0$.

Lemma 4.1. *There exist $N_1, C_1 > 0$ such that for all $z, u \in \mathbb{R}^d$ and $t > 0$ with $t > N_1 C_5 t_0(|z|)$,*

$$(4.39) \quad p(t, z, u) \leq C_1 H(z),$$

where $C_5 > 0$ is the constant given in Corollary 2.5.

Proof. Let $R_0 > 0$. For every $|z| \leq R_0$ and $t > C_5 t_0(|z|)$, we have $t \geq c_1$ by (2.19), and so

$$p(t, z, u) \leq c_2 t^{-d/\alpha} \leq \frac{c_3}{g(|z|)(1+|z|)^{d+\alpha}}, \quad z, u \in \mathbb{R}^d \text{ with } |z| \leq R_0, t > C_5 t_0(|z|).$$

Here $C_5 > 0$ is the constant from Corollary 2.5. Next, we prove (4.39) for every $|z| > R_0$ with R_0 large enough so that

$$\inf_{|z| > R_0} (|z|^\alpha - C_5 t_0(|z|)) > 0;$$

we note that such R_0 can be found due to (2.19).

For $C_5 t_0(|z|) \leq t \leq |z|^\alpha$,

$$(4.40) \quad \begin{aligned} p(2t, z, u) &\leq \begin{cases} \int_{\mathbb{R}^d} p(t, z, w) p(t, w, u) dw, & |u| \geq |z|/4 \\ c_4 \left(\frac{1}{tg(|z|)} \wedge 1 \right) \frac{t}{|z-u|^{d+\alpha}}, & |u| < |z|/4 \end{cases} \\ &\leq \begin{cases} c_5 t^{-d/\alpha} \int_{\mathbb{R}^d} p(t, z, w) dw, & |u| \geq |z|/4 \\ \frac{c_4}{g(|z|)(1+|z|)^{d+\alpha}}, & |u| < |z|/4 \end{cases} \\ &\leq \begin{cases} c_6 (1+|z|)^{dq_1/\alpha} (1+|z|)^{-\alpha/2}, & |u| \geq |z|/4 \\ \frac{c_4}{g(|z|)(1+|z|)^{d+\alpha}}, & |u| < |z|/4 \end{cases} \\ &\leq \frac{c_7}{g(|z|)(1+|z|)^{d+\alpha}} \left(1 + (1+|z|)^{dq_1/\alpha + d + \alpha + q_1 - \alpha/2} \right). \end{aligned}$$

Here in the first inequality we have used Lemma 2.8 and the fact that $|z-u| \geq |z|/2 \geq t^{1/\alpha}/2$ for every $|u| < |z|/4$ and $t \leq |z|^\alpha$, the third inequality follows from (2.22), and the fact $t > C_5 t_0(|z|) \geq c_8 (1+|z|)^{-q_1}$ (which has been established by (2.19)), and the last inequality is due to (2.18).

Furthermore, according to (4.40), we get

$$\begin{aligned} p(3t, z, u) &\leq \begin{cases} \int_{\mathbb{R}^d} p(2t, z, w) p(t, w, u) dw, & |u| \geq |z|/4 \\ c_9 \left(\frac{1}{tg(|z|)} \wedge 1 \right) \frac{t}{|z-u|^{d+\alpha}}, & |u| < |z|/4 \end{cases} \\ &\leq c_{10} \begin{cases} \frac{(1+(1+|z|)^{dq_1/\alpha + d + \alpha + q_1 - \alpha/2})}{g(|z|)(1+|z|)^{d+\alpha}} \int_{\mathbb{R}^d} p(t, w, u) dw, & |u| \geq |z|/4 \\ \frac{1}{g(|z|)(1+|z|)^{d+\alpha}}, & |u| < |z|/4 \end{cases} \\ &\leq c_{11} \begin{cases} \frac{1}{g(|z|)(1+|z|)^{d+\alpha}} (1 + (1+|z|)^{dq_1/\alpha + d + \alpha + q_1 - \alpha}), & |u| \geq |z|/4 \\ \frac{1}{g(|z|)(1+|z|)^{d+\alpha}}, & |u| < |z|/4 \end{cases} \\ &\leq \frac{c_{12}}{g(|z|)(1+|z|)^{d+\alpha}} \left(1 + (1+|z|)^{dq_1/\alpha + d + \alpha + q_1 - \alpha} \right), \end{aligned}$$

where in the first inequality we used Lemma 2.8 (similarly as in the proof of the first inequality in (4.40)), the second inequality follows from (4.40), and the third inequality is due to the fact that for all $|u| \geq |z|/4$ and $C_5 t_0(|z|) \leq t \leq |z|^\alpha$,

$$\begin{aligned} \int_{\mathbb{R}^d} p(t, w, u) dw &= T_t^V 1(u) = T_{C_5 t_0(|z|)} (T_{t-C_5 t_0(|z|)} 1)(u) \leq T_{C_5 t_0(|z|)} 1(u) \\ &\leq c_{13} \left[\exp(-c_{14} C_5 t_0(|z|) g(|u|)) + \frac{C_5 t_0(|z|)}{(1+|u|)^\alpha} \right] \\ &\leq c_{15} \left[\exp(-c_{16} C_5 t_0(|z|) g(|z|)) + \frac{C_5 t_0(|z|)}{(1+|z|)^\alpha} \right] \leq c_{17} (1+|z|)^{-\alpha/2} \end{aligned}$$

thanks to (2.21) and the arguments used for (2.24) (note that one can take the constant C_5 large enough to ensure the procedure in (2.24)).

Now, by repeating the procedure above for N_1 times, where $dq_1/\alpha + d + \alpha + q_1 - (N_1 - 1)\alpha/2 < 0$, we obtain that

$$(4.41) \quad p(N_1 t, z, u) \leq \frac{c_{18}}{g(|z|)(1 + |z|)^{d+\alpha}}, \quad C_5 t_0(|z|) \leq t \leq |z|^\alpha.$$

Hence, for any $N_1 |z|^\alpha \geq t \geq N_1 C_5 t_0(|z|)$, we can find $t_1 \in [C_5 t_0(|z|), |z|^\alpha]$ such that $t = N_1 t_1$. Then, by (4.41),

$$p(t, z, u) = p(N_1 t_1, z, u) \leq \frac{c_{18}}{g(|z|)(1 + |z|)^{d+\alpha}}.$$

On the other hand, for any $t > N_1 |z|^\alpha$,

$$\begin{aligned} p(t, z, u) &= \int_{\mathbb{R}^d} p(N_1 |z|^\alpha, z, w) p(t - N_1 |z|^\alpha, w, u) dw \\ &\leq \frac{c_{19}}{g(|z|)(1 + |z|)^{d+\alpha}} \cdot \int_{\mathbb{R}^d} p(t - N_1 |z|^\alpha, w, u) dw \\ &\leq \frac{c_{20}}{g(|z|)(1 + |z|)^{d+\alpha}} = c_{20} H(z). \end{aligned}$$

The proof is complete. \square

4.1. $t_0(\cdot)$ is almost decreasing. In this subsection, we always assume that $t_0(\cdot)$ is almost decreasing, i.e., there is a continuous and strictly decreasing function $h : [0, \infty) \rightarrow (0, \infty)$ such that (1.8) is satisfied. Define

$$s_0(t) = h^{-1}(t) \vee 2, \quad t \geq 0,$$

where $h^{-1}(t) := \inf\{s \geq 0 : h(s) \leq t\}$ and here we use the convention that $\inf \emptyset = -\infty$. Below, set $T_0 := h(2) > 0$ and $T_\infty := h(\infty) = \lim_{s \rightarrow \infty} h(s) \geq 0$. As $h(\cdot)$ is continuous and strictly decreasing, it is easy to verify that

$$(4.42) \quad s_0(t) = h^{-1}(t) \geq 2 \quad \text{for } t \in (T_\infty, T_0];$$

moreover, for every $t \in (T_0, \infty)$, and for every $t \in (0, T_\infty]$ when $T_\infty > 0$, it holds that $s_0(t) = 2$.

Lemma 4.2. *For every sufficiently large $C_0 > 0$ there are constants $C_2, C_3 > 0$ such that for all $x, y \in \mathbb{R}^d$ and $C_0 h(|x|) < t \leq C_0 T_0$, it holds that*

$$(4.43) \quad p(t, x, y) \geq C_2 H(x) H(y) \int_{\{|z| \leq s_0(t)\}} e^{-C_3 t g(|z|)} dz.$$

Proof. Let $C_0 \geq C_{**}$, where C_{**} is the constant in (1.8), and assume first that $C_0 h(|x|) < t \leq T_0$. We have

$$T_0 = h(2) \geq t > C_0 h(|x|) \geq C_0 C_{**}^{-1} h(|x|/2) \geq h(|x|/2) > h(\infty) = T_\infty.$$

This, along with (4.42) and the fact that $h(\cdot)$ is continuous and strictly decreasing, yields that $|x| \geq 2h^{-1}(t) = 2s_0(t) \geq 4$ for every $C_0 h(|x|) < t \leq T_0$. On the other hand, for every $T_\infty < t \leq T_0$ and $z \in \mathbb{R}^d$ with $|z| \leq s_0(t)$, by (1.8) and (4.42), it holds that

$$(4.44) \quad t_0(|z|) \geq c_1 h(|z|) \geq c_1 h(s_0(t)) = c_1 t.$$

Hence, according to (3.36) and (4.44), for every $z \in \mathbb{R}^d$ with $|z| \leq s_0(t)$ (also by noting here that $|z| \leq |x|/2 \leq |y|/2$ since $|x| \geq 2s_0(t)$)

$$p(t/2, z, x) \geq c_2 \left(\frac{1}{tg(|x|)} \wedge 1 \right) e^{-c_3 t g(|z|)} \frac{t}{|z - x|^{d+\alpha}} \geq \frac{c_4}{g(|x|)} e^{-c_3 t g(|z|)} \frac{1}{(1 + |x|)^{d+\alpha}},$$

where we used the facts that for all $C_0 h(|x|) < t \leq T_0$,

$$c_5 t^{1/\alpha} \leq 2 \leq s_0(t) \leq |x|/2 \leq |z - x| \leq c_6(1 + |x|), \quad |x| \geq 2s_0(t) \text{ and } |z| \leq s_0(t)$$

and

$$(4.45) \quad tg(|x|) \geq c_7 t_0(|x|) g(|x|) \geq c_8,$$

thanks to (1.7) (see the second to last line in the proof of Lemma 2.3).

Since $|y| \geq |x| \geq 2s_0(t)$, similarly we can obtain that for every $z \in \mathbb{R}^d$ with $|z| \leq s_0(t)$,

$$p(t/2, z, y) \geq c_9 \left(\frac{1}{tg(|y|)} \wedge 1 \right) e^{-c_{10}tg(|z|)} \frac{t}{|y-z|^{d+\alpha}} \geq \frac{c_{11}}{g(|y|)} e^{-c_{12}tg(|z|)} \frac{1}{(1+|y|)^{d+\alpha}}.$$

Therefore, for all $C_0h(|x|) \leq t \leq T_0$,

$$\begin{aligned} p(t, x, y) &\geq \int_{\{|z| \leq s_0(t)\}} p(t/2, x, z) p(t/2, z, y) dz \\ &\geq \frac{c_{13}}{g(|x|)g(|y|)(1+|x|)^{d+\alpha}(1+|y|)^{d+\alpha}} \int_{\{|z| \leq s_0(t)\}} e^{-c_{14}tg(|z|)} dz, \end{aligned}$$

which is the claimed estimate.

Suppose now that $t > C_0h(|x|)$ and $T_0 < t \leq C_0T_0$. Since $s_0(t) = 2$ for this range of t , we can follow the argument above to show the same estimate when $|y| \geq |x| \geq 4 = 2s_0(t)$. When $|x| < 4$ and $|y| \geq |x|$, by the fact $t \leq C_0T_0 \leq c_{15}t_0(|x|)$ (since $|x| < 4$), we can directly apply (3.36) to obtain

$$\begin{aligned} p(t, x, y) &\geq c_{16} \left(\frac{1}{tg(|y|)} \wedge 1 \right) e^{-c_{17}tg(|x|)} \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) \\ &\geq \frac{c_{18}}{g(|x|)g(|y|)(1+|x|)^{d+\alpha}(1+|y|)^{d+\alpha}} \\ &\geq \frac{c_{18}}{g(|x|)g(|y|)(1+|x|)^{d+\alpha}(1+|y|)^{d+\alpha}} \int_{\{|z| \leq s_0(t)\}} e^{-c_{17}tg(|z|)} dz. \end{aligned}$$

Here in the second inequality we have used the facts

$$g(|x|) \asymp (1+|x|) \asymp 1, \quad |x| \leq 4$$

and

$$\left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) \geq c_{20}(1+|y|)^{-d-\alpha}, \quad T_0 < t \leq C_0T_0, \quad |y| \geq |x|, \quad |x| \leq 4,$$

and the last inequality is due to the property

$$\int_{\{|z| \leq s_0(t)\}} e^{-c_{17}tg(|z|)} dz = \int_{\{|z| \leq 2\}} e^{-c_{17}tg(|z|)} dz \leq c_{21}, \quad T_0 < t \leq C_0T_0.$$

By now the proof of (4.43) is complete. \square

Lemma 4.3. *For sufficiently large $C_0 > 0$ there are constants C_4, C_5, C_6 such that for every $C_0h(|x|) < t \leq C_0T_0$,*

$$(4.46) \quad p(t, x, y) \leq C_4H(x)H(y) \int_{\{|z| \leq s_0(C_5t)\}} e^{-C_6tg(|z|)} dz.$$

Proof. According to (1.8), (2.22) and (4.39), there exists a constant $\tilde{C}_0 > 1$ such that

$$(4.47) \quad T_t^V 1(z) \leq c_1(1+|z|)^{-\alpha/2}, \quad z \in \mathbb{R}^d, \quad t > 0 \text{ with } t \geq \tilde{C}_0h(|z|)$$

and

$$(4.48) \quad p(t, z, u) \leq c_1H(z), \quad u, z \in \mathbb{R}^d, \quad t > 0 \text{ with } t \geq \tilde{C}_0h(|z|).$$

Furthermore, by (1.8) and the fact that $h(\cdot)$ is continuous and strictly decreasing, as well as that $s_0(t) = h^{-1}(t)$ when $T_\infty \leq h(|x|) < t \leq T_0$, we obtain that for all $\tilde{C}_0h(|x|) < t \leq \tilde{C}_0T_0$ the following properties hold:

$$(4.49) \quad \begin{aligned} t &\leq \tilde{C}_0h(|z|), \quad z \in \mathbb{R}^d \text{ with } |z| \leq s_0\left(\frac{t}{\tilde{C}_0}\right) =: \tilde{s}_0(t), \\ t &> \tilde{C}_0h(|z|), \quad z \in \mathbb{R}^d \text{ with } |z| > s_0\left(\frac{t}{\tilde{C}_0}\right) =: \tilde{s}_0(t). \end{aligned}$$

Now, for every $\tilde{C}_0 h(|x|) < t \leq \tilde{C}_0 T_0$ and $|z| > \tilde{s}_0(t)$,

$$(4.50) \quad \begin{aligned} p(2t, z, u) &= \int_{\mathbb{R}^d} p(t, z, w) p(t, w, u) dw \\ &\leq c_2 H(z) T_t^V 1(u) \\ &\leq c_3 H(z) \begin{cases} e^{-c_4 t g(|u|)}, & |u| \leq \tilde{s}_0(t) \\ (1 + |u|)^{-\alpha/2}, & |u| > \tilde{s}_0(t), \end{cases} \end{aligned}$$

where we have used (2.26), (4.47), (4.48) and (4.49). In the same way, we can obtain that for every $c_5 \geq 2$, $\tilde{C}_0 h(|x|) < t \leq \tilde{C}_0 T_0$, $|z| > \tilde{s}_0(t)$ and $|u| \leq \tilde{s}_0(t)$,

$$(4.51) \quad p(c_5 t, z, u) = \int_{\mathbb{R}^d} p((c_5 - 1)t, z, w) p(t, w, u) dw \leq c_6 H(z) e^{-c_4 t g(|u|)}.$$

For any $\tilde{C}_0 h(|x|) < t \leq \tilde{C}_0 T_0$ and $z, u \in \mathbb{R}^d$ with $|z| \geq \tilde{s}_0(t)$ and $|u| \geq \tilde{s}_0(t)$, we write

$$\begin{aligned} p(4t, z, u) &= \int_{\mathbb{R}^d} p(2t, z, w) p(2t, w, u) dw \\ &= \int_{\{|w| \leq \tilde{s}_0(t)\}} + \int_{\{\tilde{s}_0(t) < |w| \leq |u|\}} + \int_{\{|w| > |u|\}} p(2t, z, w) p(2t, w, u) dw \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

According to (4.50),

$$p(2t, z, w) \leq c_3 H(z) e^{-c_4 t g(|w|)}, \quad p(2t, u, w) \leq c_3 H(u) e^{-c_4 t g(|w|)}$$

for all $|w| \leq \tilde{s}_0(t)$, and so

$$I_1 \leq c_7 H(z) H(u) \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-2c_4 t g(|w|)} dw.$$

Similarly, by (4.50), we have

$$\begin{aligned} I_2 &\leq c_8 H(z) H(u) \int_{\{\tilde{s}_0(t) < |w| \leq |u|\}} (1 + |w|)^{-\alpha} dw \\ &\leq c_9 H(z) H(u) \max\{(1 + |u|)^{d-\alpha}, 1\} (1 + \log(1 + |u|)) \mathbb{1}_{\{d=\alpha=1\}}. \end{aligned}$$

Furthermore, by (4.50) again, for all $|w| \geq |u| \geq \tilde{s}_0(t)$,

$$p(2t, z, w) \leq c_3 H(z) (1 + |w|)^{-\alpha/2}, \quad p(2t, w, u) \leq c_3 H(w) (1 + |u|)^{-\alpha/2}.$$

This yields (by noting that $H(w) = \frac{1}{g(|w|)(1+|w|)^{d+\alpha}}$)

$$\begin{aligned} I_3 &\leq c_{10} H(z) (1 + |u|)^{-\alpha/2} \int_{\{|w| > |u|\}} (1 + |w|)^{-\alpha/2} H(w) dw \\ &\leq c_{11} H(z) (1 + |u|)^{-2\alpha} g(|u|)^{-1} \leq c_{12} H(z) H(u) (1 + |u|)^{d-\alpha}. \end{aligned}$$

Combining with all the estimates above for I_1 , I_2 and I_3 , we get

$$(4.52) \quad \begin{aligned} p(4t, z, u) &\leq c_{13} H(z) H(u) \max\{(1 + |u|)^{d-\alpha}, 1\} (1 + \log(1 + |u|)) \mathbb{1}_{\{d=\alpha=1\}} \\ &\quad \times \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_4 t g(|w|)} dw \end{aligned}$$

for all $z, u \in \mathbb{R}^d$, $t > 0$ with $|z| \geq \tilde{s}_0(t)$, $|u| \geq \tilde{s}_0(t)$ and $\tilde{C}_0 h(|x|) < t \leq \tilde{C}_0 T_0$. When $d \neq \alpha$, this along with (4.51) yields that for all $\tilde{C}_0 h(|x|) < t \leq \tilde{C}_0 T_0$ and $|z| \geq \tilde{s}_0(t)$,

$$(4.53) \quad p(4t, z, u) \leq c_{15} H(z) \begin{cases} e^{-c_{16} t g(|u|)}, & |u| \leq \tilde{s}_0(t), \\ (1 + |u|)^{-\min\{d+\alpha, 2\alpha\}} \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{17} t g(|w|)} dw, & |u| > \tilde{s}_0(t). \end{cases}$$

Here we have used the fact that

$$\int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{17}tg(|w|)} dw \geq \int_{\{|w| \leq c_{18}\}} e^{-c_{17}tg(|w|)} dw \geq c_{19}.$$

When $d = \alpha = 1$, we can use the following estimate instead of (4.47):

$$T_t^V 1(z) \leq c_{20}(1 + |z|)^{-\alpha} \log(1 + |z|), \quad z \in \mathbb{R}^d, \quad t > 0 \text{ with } t \geq \tilde{C}_0 h(|z|),$$

thanks to (2.22). Then, by the same arguments as above, we can prove that (4.53) still holds.

In the next step, we obtain the estimate of $p(6t, z, u)$ for every $\tilde{C}_0 h(|x|) < t \leq \tilde{C}_0 T_0$ and $z, u \in \mathbb{R}^d$ with $|z| > \tilde{s}_0(t)$ and $|u| > \tilde{s}_0(t)$ via the iterated estimate (4.53). We have

$$\begin{aligned} p(6t, z, u) &= \int_{\mathbb{R}^d} p(4t, z, w)p(2t, w, u) dw \\ &= \int_{\{|w| \leq \tilde{s}_0(t)\}} + \int_{\{\tilde{s}_0(t) < |w| \leq |u|\}} + \int_{\{|w| > |u|\}} p(4t, z, w)p(2t, w, u) dw \\ &=: J_1 + J_2 + J_3. \end{aligned}$$

Applying (4.50) and (4.53) directly, we derive

$$J_1 \leq c_{21}H(z)H(u) \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{22}tg(|w|)} dw.$$

Using (4.50) and the second inequality of (4.53) instead of the first one for the estimate of $p(4t, z, w)$, we get

$$\begin{aligned} J_2 &\leq c_{23}H(z)H(u) \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{24}tg(|w|)} dw \cdot \int_{\{\tilde{s}_0(t) < |w| \leq |u|\}} (1 + |w|)^{-\min\{d+\alpha, 2\alpha\}-\alpha/2} dw \\ &\leq c_{25}H(z)H(u) \max\{(1 + |u|)^{d-2\alpha}, 1\} \cdot \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{24}tg(|w|)} dw \end{aligned}$$

and

$$\begin{aligned} J_3 &\leq c_{26}H(z)(1 + |u|)^{-\min\{d+\alpha, 2\alpha\}} \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{27}tg(|w|)} dw \cdot \int_{\{|w| > |u|\}} (1 + |w|)^{-\alpha/2} H(w) dw \\ &\leq c_{28}H(z)H(u) \max\{(1 + |u|)^{d-2\alpha}, 1\} \cdot \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{27}tg(|w|)} dw. \end{aligned}$$

Combining with all the estimates above for J_1 – J_3 yields that for all $\tilde{C}_0 T_\infty < t \leq \tilde{C}_0 T_0$ and $z, u \in \mathbb{R}^d$ with $|z| > \tilde{s}_0(t)$ and $|u| > \tilde{s}_0(t)$,

$$(4.54) \quad p(6t, z, u) \leq c_{29}H(z)H(u) \max\{(1 + |u|)^{d-2\alpha}, 1\} \cdot \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{30}tg(|w|)} dw.$$

Furthermore, according to the decomposition

$$p(8t, z, u) = \int_{\{|w| \leq \tilde{s}_0(t)\}} + \int_{\{\tilde{s}_0(t) < |w| \leq |u|\}} + \int_{\{|w| > |u|\}} p(6t, z, w)p(2t, w, u) dw,$$

one can apply (4.54) for the estimate of $p(6t, z, w)$ and follow the arguments for (4.54) to prove that for every $\tilde{C}_0 h_0(|x|) < t \leq \tilde{C}_0 T_0$, $|z| > \tilde{s}_0(t)$ and $|u| > \tilde{s}_0(t)$,

$$p(8t, z, u) \leq c_{31}H(z)H(u) \max\{(1 + |u|)^{d-3\alpha}, 1\} \cdot \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{32}tg(|w|)} dw.$$

Then, following this procedure iteratively for N_2 times, where N_2 is such that $d - N_2\alpha < 0$, we can show that for all $\tilde{C}_0 h_0(|x|) < t \leq \tilde{C}_0 T_0$, $|z| > \tilde{s}_0(t)$ and $|u| > \tilde{s}_0(t)$,

$$\begin{aligned} p(2(N_2 + 1)t, z, u) &\leq c_{33}H(z)H(u) \int_{\{|w| \leq \tilde{s}_0(t)\}} e^{-c_{34}tg(|w|)} dw \\ &= c_{33}H(z)H(u) \int_{\{|w| \leq s_0(\tilde{C}_0^{-1}t)\}} e^{-c_{34}tg(|w|)} dw. \end{aligned}$$

Combining this estimate with the fact that $\tilde{C}_0 h(|x|) < t \leq \tilde{C}_0 T_0$ implies $|y| \geq |x| > \tilde{s}_0(t)$, we prove the desired bound for $2(N_2 + 1)\tilde{C}_0 h(|x|) < t \leq 2(N_2 + 1)\tilde{C}_0 T_0$, i.e., (4.46) holds with $C_0 = 2(N_2 + 1)\tilde{C}_0$ and $C_5 = \tilde{C}_0^{-1}$. \square

Lemma 4.4. *For sufficiently large $C_0 > 0$ there are constants $C_7, C_8 > 0$ such that for all $t > C_0 T_0$,*

$$(4.55) \quad C_7 e^{-\lambda_1 t} H(x)H(y) \leq p(t, x, y) \leq C_8 e^{-\lambda_1 t} H(x)H(y).$$

Proof. For every positive constants c_1, c_2 ,

$$c_3 \leq \int_{\{|z| \leq 2\}} e^{-c_1 g(|z|)} dz \leq \int_{\{|z| \leq s_0(c_2)\}} e^{-c_1 g(|z|)} dz \leq c_4.$$

Then, according to (4.43), (4.46) and Proposition 3.4, it holds that

$$(4.56) \quad c_5 H(z)H(u) \leq p(C_0 T_0, z, u) \leq c_6 H(z)H(u), \quad z, u \in \mathbb{R}^d.$$

Indeed, when $h(\min\{|z|, |u|\}) < T_0$ and C_0 is large enough, (4.56) follows from (4.43) and (4.46) by taking $t = C_0 T_0$. On the other hand, if $h(\min\{|z|, |u|\}) \geq T_0$, then $\min\{|z|, |u|\} \leq 2$, and so $C_0 T_0 \leq c_7 t_0(\min\{|z|, |u|\})$. Thus, by Proposition 3.4, we have

$$\begin{aligned} p(C_0 T_0, z, u) &\asymp \left(\frac{1}{C_0 T_0 g(\max\{|u|, |z|\})} \wedge 1 \right) e^{-c_7 C_0 T_0 g(\min\{|u|, |z|\})} \left((C_0 T_0)^{-d/\alpha} \wedge \frac{C_0 T_0}{|u - z|^{d+\alpha}} \right) \\ &\asymp \frac{1}{g(|u|)} \frac{1}{g(|z|)} \frac{1}{(1 + |u|)^{d+\alpha}} \frac{1}{(1 + |z|)^{d+\alpha}} = H(u)H(z). \end{aligned}$$

Here the second step is due to the facts that for every $z, u \in \mathbb{R}^d$ with $\min\{|z|, |u|\} \leq 2$,

$$\begin{aligned} g(\max\{|z|, |u|\}) &\asymp g(|z|)g(|u|), \quad g(\min\{|z|, |u|\}) \asymp 1, \\ \left((C_0 T_0)^{-d/\alpha} \wedge \frac{C_0 T_0}{|u - z|^{d+\alpha}} \right) &\asymp (1 + |u|)^{-d-\alpha} (1 + |z|)^{-d-\alpha}. \end{aligned}$$

This implies that (4.56) is still true when $h(\min\{|z|, |u|\}) \geq T_0$.

Therefore, by (1.10) and (4.56), for every $t > C_0 T_0$,

$$\begin{aligned} p(t, x, y) &= \int_{\mathbb{R}^d} p(t - C_0 T_0, x, w) p(C_0 T_0, w, y) dw \\ &\leq c_6 H(y) \int_{\mathbb{R}^d} p(t - C_0 T_0, x, w) H(w) dw \\ &= c_6 H(y) T_{t-C_0 T_0}^V H(x) \leq c_8 e^{-\lambda_1 t} H(x)H(y), \end{aligned}$$

where in the last inequality we have used (1.10).

Applying (1.10) and (4.56) again, we derive that for every $t > C_0 T_0$,

$$\begin{aligned} p(t, x, y) &= \int_{\mathbb{R}^d} p(t - C_0 T_0, x, w) p(C_0 T_0, w, y) dw \\ &\geq c_5 H(y) \int_{\mathbb{R}^d} p(t - C_0 T_0, x, w) H(w) dw \\ &= c_5 H(y) T_{t-C_0 T_0}^V H(x) \geq c_9 e^{-\lambda_1 t} H(x)H(y). \end{aligned}$$

Then, we prove (4.55). \square

4.2. $t_0(\cdot)$ is almost increasing. In this subsection, we will assume that $t_0(\cdot)$ is almost increasing. In particular, $c_{t_0} := \inf_{z \in \mathbb{R}^d} t_0(|z|) > 0$. Set

$$s_1(t) := \inf\{s > 0 : h(s) \geq t\} \vee 2, \quad t > 0,$$

where $h(\cdot)$ is a strictly increasing and continuous function given in (1.8) and we use the convention that $\inf \emptyset = -\infty$. Recall that $T_0 = h(2) > 0$ and $T_\infty = h(\infty) = \lim_{s \rightarrow \infty} h(s) \in (0, \infty]$. In this part, we are looking for estimates of $p(t, x, y)$ when $t > C_0 t_0(|x|)$. Thus, by taking C_0 large enough, we only need to consider the case that $t > T_0$. In particular, in this case we have $s_1(t) = \inf\{s > 0 : h(s) \geq t\}$.

Lemma 4.5. *There exist constants $C_0, C_9 > 0$ such that for every $z, u \in \mathbb{R}^d$, $t > 0$ with $t \geq C_0 t_0(|z|)$,*

$$(4.57) \quad p(t, z, u) \leq C_9 H(z) H(u).$$

Proof. We first note that when $t_0(\cdot)$ is a bounded function, i.e., $T_\infty = h(\infty) < \infty$, $t_0(\cdot)$ is almost increasing and almost decreasing at the same time. Indeed, we can simply modify the profile $h(\cdot)$ to be a continuous and decreasing function and keep the value $h(2)$ at the cost of comparability constants in (1.8). In this case, the estimate (4.57) follows directly from the upper bound estimate in Lemma 4.4. Therefore, for the rest of the proof we assume that $t_0(\cdot)$ is an almost increasing function and the corresponding profile h satisfies that $T_\infty = \infty$.

According to (1.8), (2.22) and (4.39), there is a constant $\tilde{C}_0 > 0$ so that

$$(4.58) \quad T_t^V 1(z) \leq c_1 (1 + |z|)^{-\alpha/2}, \quad z \in \mathbb{R}^d, \quad t > 0 \text{ with } t \geq \tilde{C}_0 h(|z|),$$

and

$$(4.59) \quad p(t, z, u) \leq c_2 H(z), \quad u, z \in \mathbb{R}^d, \quad t > 0 \text{ with } t \geq \tilde{C}_0 h(|z|).$$

Without loss of generality in the following we can assume that $\tilde{C}_0 > T_0/c_{t_0} = h(2)/c_{t_0}$. On the other hand, by (1.8), the definition of $s_1(\cdot)$ and the fact that $h(\cdot)$ is continuous and strictly increasing, we know that there exists $\tilde{C}_1 > 0$ large enough such that for all $t \geq \tilde{C}_1 T_0$,

$$(4.60) \quad \begin{aligned} t \geq \tilde{C}_1 h(|z|/6) &\geq \tilde{C}_0 h(|z|), & z \in \mathbb{R}^d \text{ with } |z| \leq 6s_1\left(\frac{t}{\tilde{C}_1}\right) &=: 6\tilde{s}_1(t), \\ t \leq \tilde{C}_1 h(|z|/6) &\leq \tilde{C}_1 h(|z|), & z \in \mathbb{R}^d \text{ with } |z| > 6s_1\left(\frac{t}{\tilde{C}_1}\right) &=: 6\tilde{s}_1(t). \end{aligned}$$

The proof below is split into three steps.

Step 1: According to (4.58), (4.59) and (4.60), we obtain that for all $z, u \in \mathbb{R}^d$ and $t > \tilde{C}_1 T_0$ with $|z| \leq \tilde{s}_1(t)$ and $|u| \leq 6\tilde{s}_1(t)$,

$$(4.61) \quad p(2t, z, u) = \int_{\mathbb{R}^d} p(t, z, w) p(t, w, u) dw \leq c_3 H(z) T_t^V 1(u) \leq c_4 H(z) (1 + |u|)^{-\alpha/2}.$$

When $t > \tilde{C}_1 T_0$, $|z| \leq \tilde{s}_1(t)$ and $|u| > \tilde{s}_1(t)$ with $|u| > 6|z|$, set

$$p(2t, z, u) = \left(\int_{\{|z-w| \leq |u|/3\}} + \int_{\{|z-w| > |u|/3\}} \right) p(t, z, w) p(t, w, u) dw =: I_1 + I_2.$$

If $|z - w| \leq |u|/3$, then $|u - w| \geq |u - z| - |z - w| \geq |u| - |z| - |u|/3 \geq |u|/2 \geq c_5 t^{1/\alpha}$, where in the last inequality we used the fact that for $t \geq \tilde{C}_1 T_0$,

$$|u| \geq \tilde{s}_1(t) \geq e^{c_6(1+t)} \geq c_7 t^{1/\alpha},$$

thanks to the fact that $h(s) \leq c_8 \log(2 + s)$ (which can be verified by (2.19) directly). By (2.28) and (4.58), we get

$$\begin{aligned} I_1 &\leq \int_{\{|z-w| \leq |u|/3\}} \left(\frac{c_9}{t \max\{g(|u|), g(|w|)\}} \wedge 1 \right) \frac{t}{|u - w|^{d+\alpha}} p(t, z, w) dw \\ &\leq \frac{c_{10}}{g(|u|)(1 + |u|)^{d+\alpha}} T_t^V 1(z) \leq c_{11} H(u) (1 + |z|)^{-\alpha/2}. \end{aligned}$$

Similarly, using (2.28) again we have

$$\begin{aligned} I_2 &\leq \int_{\{|z-w| > |u|/3\}} \left(\frac{c_{12}}{t \max\{g(|z|), g(|w|)\}} \wedge 1 \right) \frac{t}{|z - w|^{d+\alpha}} p(t, w, u) dw \\ &\leq \frac{c_{13}}{g(|u|)(1 + |u|)^{d+\alpha}} T_t^V 1(u) \\ &\leq \frac{c_{14}}{g(|u|)(1 + |u|)^{d+\alpha}} \left(e^{-c_{15}tg(|u|)} + \frac{t}{(1 + |u|)^\alpha} \right) \end{aligned}$$

$$\leq c_{14}H(u) \left(e^{-c_{15}tg(|z|)} + \frac{t}{(1+|z|)^\alpha} \right) \leq c_{16}H(u)(1+|z|)^{-\alpha/2},$$

where the second inequality follows from the fact that $g(|w|) \geq g(|z-w|-|z|) \geq g(|u|/6) \geq c_{17}g(|u|)$ for all $z, w \in \mathbb{R}^d$ with $|z-w| > |u|/3$ and $|z| \leq |u|/6$, the third inequality is due to (2.21), the fourth inequality follows from the fact $|u| \geq 6|z|$, and in the last inequality we have used the arguments from the proof of (2.24) (by taking \tilde{C}_1 larger if necessary). Hence, by all above estimates, for every $|z| \leq \tilde{s}_1(t)/6$,

$$(4.62) \quad \begin{aligned} p(2t, z, u) &\leq \begin{cases} c_{18}H(z)(1+|u|)^{-\alpha/2}, & |u| \leq \tilde{s}_1(t), \\ c_{18}H(u)(1+|z|)^{-\alpha/2}, & |u| > \tilde{s}_1(t), \end{cases} \\ &= \begin{cases} c_{18}H(z)H(u)(1+|u|)^{d+\alpha/2}g(|u|), & |u| \leq \tilde{s}_1(t), \\ c_{18}H(z)H(u)(1+|z|)^{d+\alpha/2}g(|z|), & |u| > \tilde{s}_1(t); \end{cases} \end{aligned}$$

and for every $\tilde{s}_1(t)/6 \leq |z| \leq \tilde{s}_1(t)$,

$$(4.63) \quad \begin{aligned} p(2t, z, u) &\leq \begin{cases} c_{19}H(z)H(u)(1+|u|)^{d+\alpha/2}g(|u|), & |u| \leq 6\tilde{s}_1(t), \\ c_{19}H(z)H(u)(1+|z|)^{d+\alpha/2}g(|z|), & |u| > 6\tilde{s}_1(t), \end{cases} \\ &\leq \begin{cases} c_{20}H(z)H(u)(1+|u|)^{d+\alpha/2}g(|u|), & |u| \leq \tilde{s}_1(t), \\ c_{20}H(z)H(u)(1+|z|)^{d+\alpha/2}g(|z|), & |u| > \tilde{s}_1(t), \end{cases} \end{aligned}$$

where the last inequality we have used the fact that $|u| \asymp |z|$ and $H(u) \asymp H(z)$ for every $\tilde{s}_1(t)/6 \leq |z| \leq \tilde{s}_1(t)$ and $\tilde{s}_1(t) \leq |u| \leq 6\tilde{s}_1(t)$. Then, we show that the inequality (4.62) holds for every $|z| \leq \tilde{s}_1(t)$.

Below we assume that $t > \tilde{C}_1 T_0$ and consider the case that $|z| > \tilde{s}_1(t)$. According to (4.62) (by noting that (4.62) has been verified for every $|z| \leq \tilde{s}_1(t)$ and exchanging the position of z and u),

$$(4.64) \quad p(2t, z, u) \leq c_{21}H(z)(1+|u|)^{-\alpha/2}, \quad |z| > \tilde{s}_1(t), \quad |u| \leq \tilde{s}_1(t).$$

When $|z| > \tilde{s}_1(t)$ and $|u| > \tilde{s}_1(t)$, by (4.60) and its proof, there is a constant $c_{22} > 0$ such that $t \leq c_{22} \min\{t_0(|z|), t_0(|u|)\}$. Thus, it follows from (3.36) that

$$p(2t, z, u) \leq c_{23} \left(\frac{1}{t \max\{g(|z|), g(|u|)\}} \wedge 1 \right) \left(t^{-d/\alpha} \wedge \frac{t}{|z-u|^{d+\alpha}} \right) e^{-c_{21}t \min\{g(|z|), g(|u|)\}}.$$

Hence, for every $t > \tilde{C}_1 T_0$ and $|z| > \tilde{s}_1(t)$,

$$(4.65) \quad \begin{aligned} &p(2t, z, u) \\ &\leq \begin{cases} c_{21}H(z)(1+|u|)^{-\alpha/2}, & |u| \leq \tilde{s}_1(t), \\ c_{23}e^{-c_{21}t \min\{g(|z|), g(|u|)\}} \left(\frac{1}{t \max\{g(|z|), g(|u|)\}} \wedge 1 \right) \left(t^{-d/\alpha} \wedge \frac{t}{|z-u|^{d+\alpha}} \right), & |u| > \tilde{s}_1(t). \end{cases} \end{aligned}$$

Step 2: In what follows, we will make use of (4.62) and (4.65) to obtain some improved estimates for $p(t, z, u)$, which will be applied to the iteration procedure later on. We always assume that $t > \tilde{C}_1 T_0$.

We first assume that $d \neq \alpha$. For every $|z| \leq \tilde{s}_1(t)$ and $|u| \leq \tilde{s}_1(t)$, let

$$\begin{aligned} p(4t, z, u) &= \left(\int_{\{|w| \leq |u|\}} + \int_{\{|u| \leq |w| \leq 6\tilde{s}_1(t)\}} + \int_{\{|w| > 6\tilde{s}_1(t)\}} \right) p(2t, z, w)p(2t, w, u) dw \\ &=: J_1 + J_2 + J_3. \end{aligned}$$

By (4.62) (which has been proven for every $|z| \leq \tilde{s}_1(t)$),

$$\begin{aligned} J_1 &\leq c_{24}H(z)H(u) \int_{\{|w| \leq |u|\}} H^2(w)(1+|w|)^{2(d+\alpha/2)}g^2(|w|) dw \\ &\leq c_{25}H(z)H(u) \max\{(1+|u|)^{d-\alpha}, 1\}. \end{aligned}$$

According to (4.61) and the proof of (4.62),

$$\begin{aligned} J_2 &\leq c_{26}H(z)(1+|u|)^{-\alpha/2} \int_{\{|u|\leq|w|\leq 6\tilde{s}_1(t)\}} (1+|w|)^{-\alpha/2}H(w) dw \\ &\leq c_{27}H(z)H(u)(1+|u|)^{d-\alpha}. \end{aligned}$$

On the other hand, $|w| \geq 6 \max\{|z|, |u|\}$ if $|w| > 6\tilde{s}_1(t)$. Applying (4.62) again, we derive

$$\begin{aligned} J_3 &\leq c_{28}(1+|z|)^{-\alpha/2}(1+|u|)^{-\alpha/2} \int_{\{|w|>6\max\{|z|,|u|\}\}} H^2(w) dw \\ &\leq c_{28}(1+|z|)^{-\alpha/2}(1+|u|)^{-\alpha/2}(g(|z|)g(|u|))^{-1} \int_{\{|w|>6\max\{|z|,|u|\}\}} (1+|w|)^{-2(d+\alpha)} dw \\ &\leq c_{29}H(z)H(u)(1+|u|)^{d-\alpha}. \end{aligned}$$

Hence, combining all above estimates together yields that

$$(4.66) \quad p(4t, z, u) \leq c_{30}H(z)H(u) \max\{(1+|u|)^{d-\alpha}, 1\}, \quad t > \tilde{C}_1 T_0, |z| \leq \tilde{s}_1(t), |u| \leq \tilde{s}_1(t).$$

When $d = \alpha$, we can use the following estimate instead of (4.58):

$$T_t^V 1(z) \leq c_{31}(1+|z|)^{-\alpha} \log(1+|z|), \quad z \in \mathbb{R}^d, t > 0 \text{ with } t \geq \tilde{C}_0 h(|z|),$$

thanks to (2.22). Then, by the same arguments as above, we can prove that (4.66) still holds true.

On the other hand, for every $|z| \leq \tilde{s}_1(t)$ and $|u| > \tilde{s}_1(t)$, set

$$\begin{aligned} p(6t, z, u) &= \left(\int_{\{|w|\leq\tilde{s}_1(t)\}} + \int_{\{|w|>\tilde{s}_1(t)\}} \right) p(4t, z, w)p(2t, w, u) dw \\ &=: K_1 + K_2. \end{aligned}$$

Applying (4.66) and (4.64) to $p(4t, z, w)$ and $p(2t, w, u)$ respectively, we have

$$\begin{aligned} K_1 &\leq c_{32}H(z)H(u) \max\{(1+|z|)^{d-\alpha}, 1\} \int_{\{|w|\leq\tilde{s}_1(t)\}} (1+|w|)^{-\alpha/2}H(w) dw \\ &\leq c_{33}H(z)H(u) \max\{(1+|z|)^{d-\alpha}, 1\}. \end{aligned}$$

Meanwhile, by (4.60), (4.65) and both of the second inequalities in (4.62) and (4.63), we have

$$\begin{aligned} K_2 &\leq \frac{c_{34}}{(1+|z|)^{\alpha/2}} \int_{\{|w|>\tilde{s}_1(t)\}} \frac{e^{-c_{35}t \min\{g(|w|), g(|u|)\}} H(w)}{t \max\{g(|w|), g(|u|)\}} \left(t^{-d/\alpha} \wedge \frac{t}{|w-u|^{d+\alpha}} \right) dw \\ &\leq \frac{c_{34}e^{-c_{35}tg(|z|)}}{g(|u|)(1+|z|)^{\alpha/2}} \int_{\{|w|>\tilde{s}_1(t)\}} H(w)t^{-1} \left(t^{-d/\alpha} \wedge \frac{t}{|w-u|^{d+\alpha}} \right) dw \\ &= \frac{c_{34}e^{-c_{35}tg(|z|)}}{g(|u|)(1+|z|)^{\alpha/2}} \cdot \left[\sum_{i=1}^3 \int_{\{|w|>\tilde{s}_1(t)\} \cap A_i} H(w)t^{-1} \left(t^{-d/\alpha} \wedge \frac{t}{|w-u|^{d+\alpha}} \right) dw \right] \\ &=: \frac{c_{34}e^{-c_{35}tg(|z|)}}{g(|u|)(1+|z|)^{\alpha/2}} (K_{21} + K_{22} + K_{23}), \end{aligned}$$

where $A_1 := \{w \in \mathbb{R}^d : |w-u| \leq c_{36}t^{1/\alpha}\}$, $A_2 := \{w \in \mathbb{R}^d : c_{36}t^{1/\alpha} < |w-u| \leq |u|/2\}$ and $A_3 := \{w \in \mathbb{R}^d : |w-u| > |u|/2\}$. Here we used again the fact that

$$|u| \geq \tilde{s}_1(t) \geq e^{c_{37}(1+t)} \geq 8c_{36}t^{1/\alpha},$$

which can be verified by the property that $h(s) \leq c_{38} \log(2+s)$.

By (4.60), we know that $t \geq \tilde{C}_0 h(|z|)$ when $|z| \leq \tilde{s}_1(t)$. Increasing the constant \tilde{C}_0 if necessary and following the argument for (2.24), we obtain

$$(4.67) \quad e^{-c_{35}tg(|z|)} \leq e^{-c_{35}\tilde{C}_0 h_0(|z|)g(|z|)} \leq c_{39}g(|z|)^{-1}(1+|z|)^{-d-\alpha/2}, \quad |z| \leq \tilde{s}_1(t),$$

where in the last step we have used the lower bound of $g(\cdot)$ in (2.19). Furthermore, by direct computations we can find that

$$K_{21} \leq \sup_{w \in A_1} H(w) \cdot \left(\int_{\{|w-u| \leq c_{36}t^{1/\alpha}\}} t^{-d/\alpha-1} dw \right) \leq c_{40}H(u),$$

$$K_{22} \leq \sup_{w \in A_2} H(w) \cdot \left(\int_{\{c_{36}t^{1/\alpha} < |w-u| \leq |u|/2\}} \frac{1}{|w-u|^{d+\alpha}} dw \right) \leq c_{40}H(u),$$

and

$$K_{23} \leq \frac{c_{40}}{(1+|u|)^{d+\alpha}} \int_{\{|w-u| > |u|/2\}} H(w) dw \leq c_{41}H(u).$$

Putting all the estimates above together yields

$$K_2 \leq c_{42}H(z)H(u).$$

By combining this bound with the estimate of K_1 above, we obtain that

$$p(6t, z, u) \leq c_{43}H(z)H(u) \max\{(1+|z|)^{d-\alpha}, 1\}, \quad |z| \leq \tilde{s}_1(t), |u| > \tilde{s}_1(t).$$

This, along with (4.66) (for the case that $|z| \leq \tilde{s}_1(t)$ and $|u| \leq \tilde{s}_1(t)$), yields that for every $t > \tilde{C}_1 T_0$ and $|z| \leq \tilde{s}_1(t)$,

$$(4.68) \quad p(6t, z, u) \leq \begin{cases} c_{44}H(z)H(u) \max\{(1+|u|)^{d-\alpha}, 1\}, & |u| \leq \tilde{s}_1(t), \\ c_{44}H(z)H(u) \max\{(1+|z|)^{d-\alpha}, 1\}, & |u| > \tilde{s}_1(t). \end{cases}$$

Indeed, by (4.66),

$$p(6t, z, u) = p\left(4 \times \frac{3t}{2}, z, u\right) \leq c_{44}H(z)H(u) \max\{(1+|u|)^{d-\alpha}, 1\}$$

when $|u| \leq \tilde{s}_1(3t/2)$ and $|z| \leq \tilde{s}_1(3t/2)$. According to this and the fact that $\tilde{s}_1(3t/2) \geq \tilde{s}_1(t)$, we obtain the desired assertion in (4.68).

Step 3: Now we will apply the iteration procedure and use the improved estimate (4.68). In fact, for every $t > \tilde{C}_1 T_0$, $|z| \leq \tilde{s}_1(t)$ and $|u| \leq \tilde{s}_1(t)$, we set

$$p(8t, z, u) = \left(\int_{\{|w| \leq |u|\}} + \int_{\{|u| \leq |w| \leq 6\tilde{s}_1(t)\}} + \int_{\{|w| > 6\tilde{s}_1(t)\}} \right) p(6t, z, w)p(2t, w, u) dw$$

$$=: L_1 + L_2 + L_3.$$

Using (4.62), (4.63) and (4.68) (for the estimate of $p(6t, z, w)$), we obtain

$$L_1 \leq c_{45}H(z)H(u) \int_{\{|w| \leq |u|\}} H^2(w)g(|w|) \max\{(1+|w|)^{2d-\alpha/2}, (1+|w|)^{d+\alpha/2}\} dw$$

$$\leq c_{46}H(z)H(u) \max\{(1+|u|)^{d-2\alpha}, 1\},$$

$$L_2 \leq c_{47}H(z)(1+|u|)^{-\alpha/2} \int_{\{|u| \leq |w| \leq 6\tilde{s}_1(t)\}} H^2(w) \max\{(1+|w|)^{d-\alpha}, 1\} dw$$

$$\leq c_{48}H(z)H(u)(1+|u|)^{d-2\alpha}$$

and

$$L_3 \leq c_{49}H(z) \max\{(1+|z|)^{d-\alpha}, 1\}(1+|u|)^{-\alpha/2} \int_{\{|w| > \max\{|z|, |u|\}\}} H^2(w) dw$$

$$\leq c_{50}H(z)H(u)(1+|u|)^{d-2\alpha}.$$

Combining with all above estimates yields that

$$(4.69) \quad p(8t, z, u) \leq c_{51}H(z)H(u) \max\{(1+|u|)^{d-2\alpha}, 1\}, \quad |z| \leq \tilde{s}_1(t), |u| \leq \tilde{s}_1(t).$$

On the other hand, when $|z| \leq \tilde{s}_1(t)$ and $|u| > \tilde{s}_1(t)$,

$$\begin{aligned} p(10t, z, u) &= \left(\int_{\{|w| \leq \tilde{s}_1(t)\}} + \int_{\{|w| > \tilde{s}_1(t)\}} \right) p(8t, z, w) p(2t, w, u) dw \\ &=: M_1 + M_2. \end{aligned}$$

According to (4.69) for $p(8t, z, w)$ and (4.62) for $p(2t, w, u)$ respectively, we can verify directly that

$$\begin{aligned} M_1 &\leq c_{52} H(z) H(u) \max\{(1 + |z|)^{d-2\alpha}, 1\} \int_{\{|w| \leq \tilde{s}_1(t)\}} (1 + |w|)^{-\alpha/2} H(w) dw \\ &\leq c_{53} H(z) H(u) \max\{(1 + |z|)^{d-2\alpha}, 1\}. \end{aligned}$$

Applying the arguments for the estimate of K_2 in **Step 2** (by choosing \tilde{C}_0 large enough if necessary), we can derive

$$M_2 \leq c_{54} H(z) H(u).$$

Thus, for all $|z| \leq \tilde{s}_1(t)$ and $|u| > \tilde{s}_1(t)$,

$$p(10t, z, u) \leq c_{55} H(z) H(u) \max\{(1 + |z|)^{d-2\alpha}, 1\}.$$

This, along with (the proof of) (4.69), yields that for every $t > \tilde{C}_1 T_0$ and $|z| \leq \tilde{s}_1(t)$,

$$(4.70) \quad p(10t, z, u) \leq \begin{cases} c_{56} H(z) H(u) \max\{(1 + |u|)^{d-2\alpha}, 1\}, & |u| \leq \tilde{s}_1(t), \\ c_{56} H(z) H(u) \max\{(1 + |z|)^{d-2\alpha}, 1\}, & |u| > \tilde{s}_1(t). \end{cases}$$

Furthermore, by using the improved estimate (4.70) and iterating the same arguments as above for N_0 times with $N_0\alpha > d$, we can find that

$$p((6 + 4N_0)t, z, u) \leq c_{57} H(z) H(u), \quad z, u \in \mathbb{R}^d, \quad t > \tilde{C}_1 T_0 \text{ with } |z| \leq \tilde{s}_1(t).$$

We can now combine this estimate with the fact that the inequalities $t \geq N_1 C_* \tilde{C}_1 t_0(|x|) \geq N_1 \tilde{C}_1 h(|x|)$ (with N_1 large enough to ensure $N_1 h(0) \geq T_0$) imply that $|x| \leq \tilde{s}_1(t)$. This shows that the desired bound (4.57) holds with $C_0 = N_1 C_* \tilde{C}_1 (6 + 4N_0)$ since N_1 can be chosen large enough so that $C_0 t_0(|x|) > \tilde{C}_1 T_0$. \square

Next, we can further improve the upper estimate for $p(t, x, y)$ when $t > C_0 t_0(|x|)$.

Lemma 4.6. *There exist constants $C_0, C_{10} > 0$ such that for every $t > C_0 t_0(|x|)$,*

$$(4.71) \quad p(t, x, y) \leq C_{10} e^{-\lambda_1 t} H(x) H(y).$$

Proof. For every $z, u \in \mathbb{R}^d$ and $t > 0$ with $t > C_0 t_0(|z|)$,

$$\begin{aligned} (4.72) \quad p(t, z, u) &= \int_{\mathbb{R}^d} p(t - C_0 t_0(|z|), u, w) p(C_0 t_0(|z|), w, z) dw \\ &\leq c_1 H(z) \int_{\mathbb{R}^d} p(t - C_0 t_0(|z|), u, w) H(w) dw \\ &= c_1 H(z) T_{t - C_0 t_0(|z|)}^V H(u) \\ &\leq c_2 e^{-\lambda_1 t} e^{\lambda_1 C_0 t_0(|z|)} H(z) H(u). \end{aligned}$$

Here the constant C_0 is the same as that in Lemma 4.5, the first inequality is due to (4.57), and the last inequality follows from (1.10). By (4.72), we know that for every $R_0 > 0$, (4.71) holds true for all $t > C_0 t_0(|x|)$ and $|x| \leq R_0$. Hence, it suffices to show (4.71) for any $t > C_0 t_0(|x|)$ and $|x| > R_0$ with R_0 to be determined later.

For every $t > 2C_0 \max\{t_0(|x|), t_0(|y|)\}$,

$$\begin{aligned}
 (4.73) \quad p(t, x, y) &= \int_{\mathbb{R}^d} p(t/2, x, w)p(t/2, w, y) dw \\
 &\leq c_3 e^{-\lambda_1 t} H(x)H(y) \int_{\mathbb{R}^d} H^2(w)e^{c_4 t_0(|w|)} dw \\
 &\leq c_5 e^{-\lambda_1 t} H(x)H(y).
 \end{aligned}$$

Here the first inequality follows from (4.72), and in the last inequality we have used the fact

$$(4.74) \quad \int_{\mathbb{R}^d} H^2(w)e^{c_4 t_0(|w|)} dw < +\infty,$$

which can be verified directly by (2.19), thanks to the fact that $g(r) \rightarrow \infty$ as $r \rightarrow \infty$. On the other hand, similarly as in (4.60), we can find $\tilde{C}_1 > 0$ large enough such that for $t \geq \tilde{C}_1 T_0$,

$$\begin{aligned}
 (4.75) \quad t \geq \tilde{C}_1 h(|z|) &\geq 2C_0 t_0(|z|), \quad z \in \mathbb{R}^d \text{ with } |z| \leq s_1 \left(\frac{t}{\tilde{C}_1} \right) =: \tilde{s}_1(t), \\
 t \leq \tilde{C}_1 h(|z|) &\leq c_6 t_0(|z|), \quad z \in \mathbb{R}^d \text{ with } |z| > s_1 \left(\frac{t}{\tilde{C}_1} \right) =: \tilde{s}_1(t).
 \end{aligned}$$

Below we choose the constant C_0 large enough so that $C_0 \inf_{x \in \mathbb{R}^d} t_0(|x|) \geq \tilde{C}_1 T_0$. Then, according to (4.75) and (4.72), we have

$$(4.76) \quad p(t, z, u) \leq c_7 e^{-\lambda_1 t} e^{c_8 t_0(|z|)} H(z)H(u), \quad t \geq C_0 t_0(|x|), \quad |z| \leq \tilde{s}_1(t);$$

by (4.75) and (4.73), we know that (4.71) is true for every $|x| \leq \tilde{s}_1(t)$ and $|y| \leq \tilde{s}_1(t)$, whenever $t > C_0 t_0(|x|)$.

Next, it remains to consider the case that $|x| \leq \tilde{s}_1(t)$, $|y| > \tilde{s}_1(t)$ and $t \geq C_0 t_0(|x|)$. Set

$$\begin{aligned}
 p(2t, x, y) &= \left(\int_{\{|w| \leq \tilde{s}_1(t)\}} + \int_{\{|w| > \tilde{s}_1(t)\}} \right) p(t, x, w)p(t, w, y) dw \\
 &=: I_1 + I_2.
 \end{aligned}$$

By (4.76), it holds that

$$\begin{aligned}
 I_1 &\leq c_9 e^{-2\lambda_1 t} H(x)H(y) \int_{\{|w| \leq \tilde{s}_1(t)\}} H^2(w)e^{c_{10} t_0(|w|)} dw \\
 &\leq c_{11} e^{-2\lambda_1 t} H(x)H(y),
 \end{aligned}$$

where the last inequality is due to (4.74).

When $|w| > \tilde{s}_1(t)$ and $|y| > \tilde{s}_1(t)$, it follows from (4.75) that there is a constant $c_{12} > 0$ such that $t \leq c_{12} \min\{t_0(|w|), t_0(|y|)\}$. Hence, applying (3.36), we obtain

$$p(t, w, y) \leq c_{13} \left(\frac{1}{t \max\{g(|w|), g(|y|)\}} \wedge 1 \right) \left(t^{-d/\alpha} \wedge \frac{t}{|w-y|^{d+\alpha}} \right) e^{-c_{14} t \min\{g(|w|), g(|y|)\}}.$$

By (4.76) and the fact that $|x| \leq \tilde{s}_1(t) \leq \min\{|w|, |y|\}$,

$$\begin{aligned}
 I_2 &\leq c_{15} H(x) e^{c_{16} t_0(|x|)} \int_{\{|w| > \tilde{s}_1(t)\}} \frac{e^{-2c_{17} t \min\{g(|w|), g(|y|)\}} H(w)}{t \max\{g(|w|), g(|y|)\}} \left(t^{-d/\alpha} \wedge \frac{t}{|w-y|^{d+\alpha}} \right) dw \\
 &\leq c_{15} e^{c_{16} t_0(|x|) - 2c_{17} t g(|x|)} H(x) g(|y|)^{-1} \cdot \int_{\{|w| > \tilde{s}_1(t)\}} H(w) t^{-1} \left(t^{-d/\alpha} \wedge \frac{t}{|w-y|^{d+\alpha}} \right) dw.
 \end{aligned}$$

Note that

$$e^{c_{16} t_0(|x|) - 2c_{17} t g(|x|)} = e^{-c_{17} t g(|x|)} e^{c_{16} t_0(|x|) - c_{17} t g(|x|)}.$$

Since $\lim_{r \uparrow \infty} g(r) = \infty$, we can find $R_0 > 0$ large enough so that for all $|x| \geq R_0$,

$$e^{-c_{17} t g(|x|)} \leq e^{-2\lambda_1 t}, \quad t > 0$$

and

$$\begin{aligned} e^{c_{16}t_0(|x|)-c_{17}tg(|x|)} &\leq \exp\left(-t_0(|x|)(C_0c_{17}g(|x|)-c_{16})\right) \leq \exp\left(-\frac{C_0c_{17}t_0(|x|)g(|x|)}{2}\right) \\ &\leq c_{18}, \quad t > C_0t_0(|x|). \end{aligned}$$

Putting both estimates above together, we obtain

$$e^{c_{16}t_0(|x|)-2c_{17}tg(|x|)} \leq c_{19}e^{-2\lambda_1t}, \quad t > C_0t_0(|x|), \quad |z| > R_0.$$

On the other hand, according to the procedures for the estimate of the term K_2 in the proof of Lemma 4.5, we find that

$$\int_{\{|w|>\tilde{s}_1(t)\}} H(w)t^{-1} \left(t^{-d/\alpha} \wedge \frac{t}{|w-y|^{d+\alpha}} \right) dw \leq c_{20}g(|y|)H(y).$$

Hence,

$$I_2 \leq c_{21}e^{-2\lambda_1t}H(x)H(y), \quad t > 2C_0t_0(|x|), \quad |x| \leq \tilde{s}_1(t), \quad |y| > \tilde{s}_1(t), \quad |x| > R_0.$$

Finally, by the estimates of I_1 and I_2 , we obtain

$$p(2t, x, y) \leq c_{22}e^{-2\lambda_1t}H(x)H(y), \quad t > 2C_0t_0(|x|), \quad |x| \leq \tilde{s}_1(t), \quad |y| > \tilde{s}_1(t), \quad |x| > R_0,$$

whenever $t > C_0t_0(|x|)$. This implies that (4.71) holds for every $|x| \leq \tilde{s}_1(t)$ and $|y| > \tilde{s}_1(t)$, and the proof of the assertion (4.71) is finished. \square

Lemma 4.7. *There exist constants $C_0, C_{11} > 0$ such that for every $t > C_0t_0(|x|)$,*

$$(4.77) \quad p(t, x, y) \geq C_{11}e^{-\lambda_1t}H(x)H(y).$$

Proof. Since $t_0(\cdot)$ is almost increasing, we know that $\inf_{x \in \mathbb{R}^d} t_0(|x|) \geq c_1 > 0$. According to (3.36), for all $w \in B(0, 2)$ and $z \in \mathbb{R}^d$,

$$\begin{aligned} p(1, z, w) &\geq c_2 \left(1 \wedge \frac{1}{|w-z|^{d+\alpha}} \right) \left(\frac{1}{\max\{g(|w|), g(|z|)\}} \wedge 1 \right) e^{-c_3 \min\{g(|w|), g(|z|)\}} \\ &\geq \frac{c_4}{g(|z|)(1+|z|)^{d+\alpha}}. \end{aligned}$$

Hence, for every $z, u \in \mathbb{R}^d$,

$$p(2, z, u) = \int_{\mathbb{R}^d} p(1, z, w)p(1, w, u) dw \geq \int_{B(0,2)} p(1, z, w)p(1, w, u) dw \geq c_5H(z)H(u).$$

Therefore, for every $t > C_0t_0(|x|) > 2$ (by taking C_0 large enough),

$$\begin{aligned} p(t, x, y) &= \int_{\mathbb{R}^d} p(t-2, x, z)p(2, z, y) dz \geq c_5H(y) \int_{\mathbb{R}^d} p(t-2, x, z)H(z) dz \\ &= c_5H(y)T_{t-2}^V H(x) \geq c_6e^{-\lambda_1t}H(x)H(y), \end{aligned}$$

where the last inequality follows from (1.10). So, the proof of (4.77) is finished. \square

We can now summarise the results in this section as follows.

Proposition 4.8. *There are constants $C_0, \tilde{C}_0, C_{12}, \dots, C_{20} > 0$ such that the following statements hold.*

(1) *If $t_0(\cdot)$ is almost decreasing, then, for every $C_0t_0(|x|) < t \leq \tilde{C}_0$,*

$$\begin{aligned} C_{12}H(x)H(y) \int_{\{|z| \leq s_0(t)\}} e^{-C_{13}tg(|z|)} dz &\leq p(t, x, y) \\ &\leq C_{14}H(x)H(y) \int_{\{|z| \leq s_0(C_{15}t)\}} e^{-C_{16}tg(|z|)} dz; \end{aligned}$$

for every $t > \tilde{C}_0$,

$$C_{17}e^{-\lambda_1t}H(x)H(y) \leq p(t, x, y) \leq C_{18}e^{-\lambda_1t}H(x)H(y).$$

(2) If $t_0(\cdot)$ is almost increasing, then for every $t > C_0 t_0(|x|)$

$$C_{19}e^{-\lambda_1 t}H(x)H(y) \leq p(t, x, y) \leq C_{20}e^{-\lambda_1 t}H(x)H(y).$$

Remark 4.9. Note that in Proposition 3.4 the time interval can be $0 < t \leq C'_0 t_0(|x|)$ with any $C'_0 > 0$. Therefore, according to (1.8), the estimates in Proposition 4.8 still hold true for $C_0 t_0(|x|) < t \leq \tilde{C}_0$ by taking a different constant C_0 if necessary. On the other hand, by the statement (1) in Proposition 4.8, the estimates in (1) hold for any \tilde{C}_0 with $\tilde{C}_0 > C_0 t_0(|x|)$.

5. JUSTIFICATION OF EXAMPLE 1.2

By assumption we know that V satisfies (1.6) with $g(s) = \log^\beta(e + s)$. So, by (1.7), it is easy to check that $t_0(s)$ satisfies (1.8) with $h(s) = \log^{1-\beta}(2 + s)$. Then the estimates (1) and (2-i) follow directly from Theorem 1.1.

It remains to verify (2-ii). Let $\beta > 1$. Since $h(s) = \log^{1-\beta}(2 + s)$,

$$s_0(t) := h^{-1}(t) \vee 2 = \begin{cases} 2, & t \geq \log^{1-\beta} 4, \\ \exp(t^{-1/(\beta-1)}) - 2, & t \in (0, \log^{1-\beta} 4]. \end{cases}$$

For every $0 < t \leq 2$, first observe that for every positive constants c_0, c_1 and c_2 ,

$$\int_{\{|z| \leq c_1 s_0(c_2 t)\}} e^{-c_0 t \log^\beta(2+|z|)} dz \leq c_3 (c_1 s_0(c_2 t))^d \leq c_4 \exp(c_5 t^{-1/(\beta-1)}).$$

On the other hand, for every $0 < t \leq 2$,

$$\int_{\{|z| \leq s_0(t)\}} e^{-c_0 t \log^\beta(2+|z|)} dz \geq \int_{\{|z| \leq 2\}} e^{-c_0 t \log^\beta(2+|z|)} dz \geq c_5.$$

Hence, in order to get the desired lower bound it suffices to consider the time interval $0 < t \leq t_0$ for some fixed $t_0 \in (0, \log^{1-\beta} 4]$. Using spherical coordinates and the substitution $s = \log^\beta(2+r)$, we get

$$\begin{aligned} \int_{\{|z| \leq s_0(t)\}} e^{-c_0 t \log^\beta(2+|z|)} dz &= c_6 \int_0^{s_0(t)} e^{-c_0 t \log^\beta(2+r)} r^{d-1} dr \\ &= c_6 \beta^{-1} \int_{\log^\beta 2}^{t^{-\beta/(\beta-1)}} e^{-c_0 t s} \left(e^{s^{1/\beta}} - 2 \right)^{d-1} e^{s^{1/\beta}} s^{-(\beta-1)/\beta} ds. \end{aligned}$$

Choose $\theta \in (0, 1/2)$ and $t_0 \in (0, \log^{1-\beta} 4]$ such that

$$(5.78) \quad \theta t_0^{-\beta/(\beta-1)}/2 \geq \log^\beta 2, \quad d \left(\frac{\theta}{2} \right)^{1/\beta} - c_0 \theta \geq c_0 \theta / 2;$$

this is possible because $\beta > 1$. Therefore, for every $0 < t \leq t_0$,

$$\begin{aligned} &\int_{\{|z| \leq s_0(t)\}} e^{-c_0 t \log^\beta(2+|z|)} dz \\ &\geq c_6 \beta^{-1} \int_{\theta t^{-\beta/(\beta-1)}/2}^{\theta t^{-\beta/(\beta-1)}} e^{-c_0 t s} \left(e^{s^{1/\beta}} - 2 \right)^{d-1} e^{s^{1/\beta}} s^{-(\beta-1)/\beta} ds \\ &\geq c_7 t \int_{\theta t^{-\beta/(\beta-1)}/2}^{\theta t^{-\beta/(\beta-1)}} \exp \left(ds^{1/\beta} - c_0 t s \right) ds \\ &\geq c_8 t^{1-\beta/(\beta-1)} \cdot \exp \left(t^{-1/(\beta-1)} \left(d \left(\frac{\theta}{2} \right)^{1/\beta} - c_0 \theta \right) \right) \\ &\geq c_9 \exp \left(c_0 \theta t^{-1/(\beta-1)}/2 \right), \end{aligned}$$

where the last inequality follows from (5.78). Thus, the proof is complete.

6. PROOFS OF MAIN RESULTS

Proof of Theorem 1.1. The conclusion is a direct consequence of Proposition 3.4 and Proposition 4.8. See Remark 4.9 for the additional comments. \square

Proof of Theorem 1.3. Let $x, y \in \mathbb{R}^d$ with $|x| \leq |y|$ and $x \neq y$. We write

$$G^V(x, y) = \int_0^{C_0 t_0(|x|)} p(t, x, y) dt + \int_{C_0 t_0(|x|)}^\infty p(t, x, y) dt =: I + II,$$

where the constant $C_0 > 0$ comes from Theorem 1.1.

We first apply the estimates from Theorem 1.1(1) to estimate the integral I . An analysis is divided into three cases.

Case 1: $2|x - y|^\alpha \leq C_0 t_0(|x|)$ and $2/g(|x|) < |x - y|^\alpha$.

We write

$$I = \left(\int_0^{1/g(|y|)} + \int_{1/g(|y|)}^{|x-y|^\alpha} + \int_{|x-y|^\alpha}^{C_0 t_0(|x|)} \right) p(t, x, y) dt =: I_1 + I_2 + I_3.$$

According to (1.11), we have

$$I_1 \asymp |x - y|^{-d-\alpha} \int_0^{1/g(|y|)} t e^{-ctg(|x|)} dt \asymp |x - y|^{-d-\alpha} \int_0^{1/g(|y|)} t dt \asymp |x - y|^{-d-\alpha} g(|y|)^{-2}.$$

We point out that all of the comparability formulas in this proof contain the exponential terms with a generic constant c , but this is only a shorthand notation. The exponential term $e^{-ctg(|x|)}$ comes from the two-sided bound in Theorem 1.1(1) and the reader should note that we always have $c = C_2$ in lower estimates and $c = C_4$ in upper estimates, for different C_2, C_4 .

Similarly, by (1.11),

$$\begin{aligned} I_2 &\asymp |x - y|^{-d-\alpha} g(|y|)^{-1} \int_{1/g(|y|)}^{|x-y|^\alpha} e^{-ctg(|x|)} dt \\ &\asymp |x - y|^{-d-\alpha} g(|x|)^{-1} g(|y|)^{-1} \int_{g(|x|)/g(|y|)}^{|x-y|^\alpha g(|x|)} e^{-cs} ds \\ &\asymp |x - y|^{-d-\alpha} g(|x|)^{-1} g(|y|)^{-1}, \end{aligned}$$

where in the third step we have used the fact that (since $g(|x|)/g(|y|) \leq 1$ and $|x - y|^\alpha g(|x|) \geq 2$)

$$c_1 \leq \int_1^2 e^{-cs} ds \leq \int_{g(|x|)/g(|y|)}^{|x-y|^\alpha g(|x|)} e^{-cs} ds \leq \int_0^\infty e^{-cs} ds \leq c_2.$$

Moreover, by applying (1.11) and using substitution $t = |x - y|^\alpha s$, we get

$$\begin{aligned} I_3 &\asymp g(|y|)^{-1} \int_{|x-y|^\alpha}^{C_0 t_0(|x|)} t^{-d/\alpha-1} e^{-ctg(|x|)} dt \\ &\asymp |x - y|^{-d} g(|y|)^{-1} \int_1^{C_0 t_0(|x|)/|x-y|^\alpha} s^{-d/\alpha-1} e^{-c|x-y|^\alpha g(|x|)s} ds \\ &\asymp |x - y|^{-d} g(|y|)^{-1} e^{-c|x-y|^\alpha g(|x|)}, \end{aligned}$$

where the third step follows from the estimates

$$\begin{aligned} \int_1^{C_0 t_0(|x|)/|x-y|^\alpha} s^{-d/\alpha-1} e^{-c|x-y|^\alpha g(|x|)s} ds &\leq e^{-c|x-y|^\alpha g(|x|)} \int_1^\infty s^{-d/\alpha-1} ds \\ &\leq c_3 e^{-c|x-y|^\alpha g(|x|)} \end{aligned}$$

and

$$\int_1^{C_0 t_0(|x|)/|x-y|^\alpha} s^{-d/\alpha-1} e^{-c|x-y|^\alpha g(|x|)s} ds \geq e^{-2c|x-y|^\alpha g(|x|)} \int_1^2 s^{-d/\alpha-1} ds$$

$$\geq c_4 e^{-2c|x-y|^\alpha g(|x|)}.$$

In particular,

$$I_3 \leq c_5 |x-y|^{-d-\alpha} g(|x|)^{-1} g(|y|)^{-1}.$$

Consequently, combining all the estimates above with the facts $g(|y|) \geq g(|x|)$ and $Q(x, y) \asymp 1$ (since $2/g(|x|) < |x-y|^\alpha$), we have

$$I \asymp |x-y|^{-d-\alpha} g(|x|)^{-1} g(|y|)^{-1} \asymp |x-y|^{-d+\alpha} \cdot \left(\frac{1}{g(|y|)|x-y|^\alpha} \wedge 1 \right) \left(\frac{1}{g(|x|)|x-y|^\alpha} \wedge 1 \right) Q(x, y).$$

Case 2: $2|x-y|^\alpha \leq C_0 t_0(|x|)$ and $2/g(|x|) \geq |x-y|^\alpha$. In this case, we will use the simpler decomposition

$$I = \left(\int_0^{|x-y|^\alpha/2} + \int_{|x-y|^\alpha/2}^{C_0 t_0(|x|)} \right) p(t, x, y) dt =: J_1 + J_2.$$

Indeed, $|x-y| \leq (2/g(|x|))^{1/\alpha} \leq c_6$, so it holds that $g(|x|) \asymp g(|y|)$ for this range of x and y , which implies that

$$(6.79) \quad (tg(|y|))^{-1} \wedge 1 e^{-ctg(|x|)} \asymp e^{-tg(|x|)}, \quad 0 \leq t \leq C_0 t_0(|x|).$$

With this and (1.11),

$$J_1 \asymp |x-y|^{-d-\alpha} \int_0^{|x-y|^\alpha/2} t e^{-ctg(|x|)} dt \asymp |x-y|^{-d-\alpha} \int_0^{|x-y|^\alpha/2} t dt \asymp |x-y|^{-d+\alpha},$$

where we have used the fact that $|x-y|^\alpha g(|x|) \leq 2$.

Meanwhile, by (1.11), (6.79) and substitution $t = |x-y|^\alpha s/2$, we derive

$$J_2 \asymp \int_{|x-y|^\alpha/2}^{C_0 t_0(|x|)} t^{-d/\alpha} e^{-ctg(|x|)} dt \asymp |x-y|^{-d+\alpha} \int_1^{\frac{2C_0 t_0(|x|)}{|x-y|^\alpha}} s^{-d/\alpha} e^{-csg(|x|)|x-y|^\alpha} ds.$$

If $\alpha < d$, then, by the fact $2|x-y|^\alpha \leq C_0 t_0(|x|)$ and $|x-y|^\alpha g(|x|) \leq 2$, it holds that

$$c_7 \leq c_8 \int_1^2 s^{-d/\alpha} ds \leq c_9 \int_1^{\frac{2C_0 t_0(|x|)}{|x-y|^\alpha}} s^{-d/\alpha} e^{-csg(|x|)|x-y|^\alpha} ds \leq c_{10} \int_1^\infty s^{-d/\alpha} ds = \frac{c_{10}\alpha}{d-\alpha}.$$

If $\alpha = d = 1$, then, by noting from (1.7) and (2.20) that $2C_0 t_0(|x|) \geq c_{11}g(|x|)^{-1}$,

$$\begin{aligned} \int_1^{\frac{2C_0 t_0(|x|)}{|x-y|^\alpha}} s^{-d/\alpha} e^{-csg(|x|)|x-y|^\alpha} ds &\geq c_{12} \int_1^{\max\left\{\frac{c_{11}}{g(|x|)|x-y|^\alpha}, 4\right\}} s^{-1} ds \\ &\geq c_{13} \max\left\{\log\left(\frac{1}{|x-y|^\alpha g(|x|)}\right), 1\right\} \end{aligned}$$

and

$$\begin{aligned} &\int_1^{\frac{2C_0 t_0(|x|)}{|x-y|^\alpha}} s^{-d/\alpha} e^{-csg(|x|)|x-y|^\alpha} ds \\ &\leq c_{14} \int_1^{\max\left\{\frac{c_{11}}{g(|x|)|x-y|^\alpha}, 1\right\}} s^{-1} ds + c_{14} g(|x|) |x-y|^\alpha \cdot \int_{\max\left\{\frac{c_{11}}{g(|x|)|x-y|^\alpha}, 1\right\}}^\infty e^{-csg(|x|)|x-y|^\alpha} ds \\ &\leq c_{15} \max\left\{\log\left(\frac{c_{11}}{g(|x|)|x-y|^\alpha}\right), 1\right\}. \end{aligned}$$

If $\alpha > d = 1$, then

$$\begin{aligned} \int_1^{\frac{2C_0 t_0(|x|)}{|x-y|^\alpha}} s^{-d/\alpha} e^{-csg(|x|)|x-y|^\alpha} ds &\geq c_{16} \int_1^{\max\left\{\frac{c_{11}}{g(|x|)|x-y|^\alpha}, 4\right\}} s^{-d/\alpha} ds \\ &\geq c_{17} \left(\frac{1}{|x-y|^\alpha g(|x|)}\right)^{(\alpha-d)/\alpha} \end{aligned}$$

and

$$\begin{aligned}
& \int_1^{\frac{2C_0t_0(|x|)}{|x-y|^\alpha}} s^{-d/\alpha} e^{-csg(|x|)|x-y|^\alpha} ds \\
& \leq c_{18} \int_1^{\max\left\{\frac{c_{11}}{g(|x|)|x-y|^\alpha}, 1\right\}} s^{-d/\alpha} ds \\
& \quad + c_{18} (g(|x|)|x-y|^\alpha)^{-d/\alpha} \cdot \int_{\max\left\{\frac{c_{11}}{g(|x|)|x-y|^\alpha}, 1\right\}}^\infty e^{-csg(|x|)|x-y|^\alpha} ds \\
& \leq c_{19} \left(\frac{1}{|x-y|^\alpha g(|x|)} \right)^{(\alpha-d)/\alpha}.
\end{aligned}$$

Hence, putting all above estimates together, we obtain

$$J_2 \asymp |x-y|^{-d+\alpha} Q(x, y),$$

which implies immediately that

$$I \asymp |x-y|^{-d+\alpha} Q(x, y) \asymp |x-y|^{-d+\alpha} \cdot \left(\frac{1}{g(|y|)|x-y|^\alpha} \wedge 1 \right) \left(\frac{1}{g(|x|)|x-y|^\alpha} \wedge 1 \right) Q(x, y),$$

where we have used the fact that $|x-y|^\alpha \leq 2/g(|x|)$ and $g(|x|) \asymp g(|y|)$.

Case 3: $2|x-y|^\alpha > C_0t_0(|x|)$.

Similarly, it follows from (1.7) and (2.20) that $C_0t_0(|x|) > 2c_{20}g(|x|)^{-1} \geq 2c_{20}g(|y|)^{-1}$. Let

$$I = \left(\int_0^{c_{20}/g(|y|)} + \int_{c_{20}/g(|y|)}^{C_0t_0(|x|)} \right) p(t, x, y) dt =: K_1 + K_2.$$

So by (1.11), we get

$$K_1 \asymp |x-y|^{-d-\alpha} \int_0^{c_{20}/g(|y|)} t dt \asymp |x-y|^{-d-\alpha} g(|y|)^{-2}$$

and

$$\begin{aligned}
K_2 & \asymp |x-y|^{-d-\alpha} g(|y|)^{-1} \int_{c_{20}/g(|y|)}^{C_0t_0(|x|)} e^{-ctg(|x|)} dt \\
& \asymp |x-y|^{-d-\alpha} g(|x|)^{-1} g(|y|)^{-1},
\end{aligned}$$

where in the last step we have used the fact that

$$\begin{aligned}
c_{21}g(|x|)^{-1} & \leq \int_{c_{20}/g(|y|)}^{2c_{20}/g(|x|)} e^{-ctg(|x|)} dt \leq \int_{c_{20}/g(|y|)}^{C_0t_0(|x|)} e^{-ctg(|x|)} dt \\
& \leq \int_{c_{20}/g(|y|)}^\infty e^{-ctg(|x|)} dt \leq c_{22}g(|x|)^{-1}.
\end{aligned}$$

Putting all the estimates above together and combining them with the facts that $g(|y|) \geq g(|x|)$ and $Q(x, y) \asymp 1$ (since $c_{20}/g(|x|) < |x-y|^\alpha$), we have

$$I \asymp |x-y|^{-d-\alpha} g(|x|)^{-1} g(|y|)^{-1} \asymp |x-y|^{-d+\alpha} \cdot \left(\frac{1}{g(|y|)|x-y|^\alpha} \wedge 1 \right) \left(\frac{1}{g(|x|)|x-y|^\alpha} \wedge 1 \right) Q(x, y).$$

By now we have obtained that for all of these cases,

$$I \asymp \frac{1}{|x-y|^{d-\alpha}} \left(1 \wedge \frac{1}{g(|x|)|x-y|^\alpha} \right) \left(1 \wedge \frac{1}{g(|y|)|x-y|^\alpha} \right) Q(x, y).$$

Next, we are going to estimate the integral II . We will show that there is a constant $c_{23} > 0$ such that $II \leq c_{23}I$, which will lead us immediately to the desired conclusion. Recall we assume that $|x| \leq |y|$.

When $t_0(\cdot)$ is almost increasing, it follows directly from (1.12) that

$$(6.80) \quad II \leq \frac{c_{24}}{g(|x|)g(|y|)(1+|x|)^{d+\alpha}(1+|y|)^{d+\alpha}}.$$

Suppose that $t_0(\cdot)$ is almost decreasing. Then we set

$$II = \left(\int_{C_0 t_0(|x|)}^{\tilde{C}_0} + \int_{\tilde{C}_0}^{\infty} \right) p(t, x, y) dt =: L_1 + L_2,$$

where \tilde{C}_0 is the same constant in Theorem 1.1. By (1.14) it is direct to see that

$$L_2 \leq \frac{c_{25}}{g(|x|)g(|y|)(1+|x|)^{d+\alpha}(1+|y|)^{d+\alpha}}.$$

By tracking the proof of Theorem 1.1(2-ii) we can take $C_0 > C_* C_{10}^{-1}$, where C_* , C_{10} are positive constants in (1.8) and (1.13), respectively. Hence, according to (1.8), for every $C_0 t_0(|x|) \leq t \leq \tilde{C}_0$,

$$s_0(C_{10}t) \leq s_0(C_0 C_{10} t_0(|x|)) \leq s_0(C_0 C_{10} C_*^{-1} h(|x|)) \leq s_0(h(|x|)) \leq (|x| \vee 2) \leq 2(1+|x|).$$

Then, by (1.13),

$$\begin{aligned} L_1 &\leq c_{26} H(x) H(y) \int_{C_0 t_0(|x|)}^{\tilde{C}_0} \int_{\{|z| \leq c_{27} s_0(C_{10}t)\}} e^{-c_{28} t g(|z|)} dz dt \\ &\leq c_{29} H(x) H(y) (1+|x|)^d. \end{aligned}$$

Consequently, combining the estimates of L_1 and L_2 with (6.80), we can conclude that if $t_0(\cdot)$ is an almost monotone function and $C_0 t_0(|x|) \leq h(2)$, then

$$II \leq \frac{c_{30}}{g(|x|)g(|y|)(1+|y|)^{d+\alpha}}.$$

On the other hand, when $C_0 t_0(|x|) > h(2)$, $II \leq L_2$ and so it is clear that the inequality above holds true.

Observe now that for the **Case 1** and **Case 3** above, it holds that $|x-y|^\alpha > c_{31} g(|x|)^{-1} \geq c_{31} g(|y|)^{-1}$. Then, we can write

$$\begin{aligned} II &\leq \frac{c_{30}}{g(|x|)g(|y|)(1+|y|)^{d+\alpha}} \\ &= \frac{c_{30} |x-y|^{d+\alpha}}{|x-y|^{d-\alpha} g(|x|) |x-y|^\alpha g(|y|) |x-y|^\alpha (1+|y|)^{d+\alpha}} \\ &\leq \frac{c_{32}}{|x-y|^{d-\alpha}} \left(1 \wedge \frac{1}{g(|x|) |x-y|^\alpha} \right) \left(1 \wedge \frac{1}{g(|y|) |x-y|^\alpha} \right) \frac{|x-y|^{d+\alpha}}{(1+|y|)^{d+\alpha}} \\ &\leq \frac{c_{32}}{|x-y|^{d-\alpha}} \left(1 \wedge \frac{1}{g(|x|) |x-y|^\alpha} \right) \left(1 \wedge \frac{1}{g(|y|) |x-y|^\alpha} \right) \frac{(2|y|)^{d+\alpha}}{(1+|y|)^{d+\alpha}} \\ &\leq c_{33} I. \end{aligned}$$

For the remaining **Case 2** above, it holds that $|x-y|^\alpha \leq c_{34} g(|x|)^{-1} \leq c_{35}$ and $g(|x|) \asymp g(|y|)$. By this we obtain immediately that

$$II \leq c_{36} \leq c_{37} I.$$

So now we have finished the proof. \square

Acknowledgements. The research of Xin Chen is supported by the National Natural Science Foundation of China (No. 12122111). The research of Kamil Kaleta is supported by the National Science Centre, Poland, project OPUS-18 2019/35/B/ST1/02421. The research of Jian Wang is supported by the National Key R&D Program of China (2022YFA1006003) and the National Natural Science Foundation of China (No. 12225104).

Conflict of Interest The authors have no competing interests to declare that are relevant to the content of this article.

REFERENCES

- [1] Bender, C.M. and Wu, T.T.: Anharmonic Oscillator, *Phys. Rev.*, **184** (1969), 1231.
- [2] Blumenthal, R.M. and Gettoor, R.K.: Some theorems on stable processes, *Trans. Amer. Math. Soc.*, **9** (1960), 263–273.
- [3] Bogdan, K. and Jakubowski, T.: Estimates of heat kernel of fractional Laplacian perturbed by gradient operators, *Comm. Math. Phys.*, **271** (2007), 179–198.
- [4] Campisi, M., Zhan, F., Talkner, P. and Hänggi, P.: Logarithmic oscillators: ideal Hamiltonian thermostats, *Phys. Rev. Lett.*, **108** (2012), 250601.
- [5] Chen, X. and Wang, J.: Intrinsic contractivity properties of Feynman–Kac semigroups for symmetric jump processes with infinite range jumps, *Front. Math. China*, **10** (2015), 753–776.
- [6] Chen, X. and Wang, J.: Intrinsic ultracontractivity of Feynman–Kac semigroups for symmetric jump processes, *J. Funct. Anal.*, **270** (2016), 4152–4195.
- [7] Chen, X. and Wang, J.: Two-sided heat kernel estimates for Schrödinger semigroups with unbounded potential, *Ann. Probab.*, **52** (2024), 1016–1047.
- [8] Chen, Z.-Q., Kim, P. and Song, R.M.: Stability of Dirichlet heat kernel estimates for non-local operators under Feynman–Kac perturbation, *Trans. Amer. Math. Soc.*, **367** (2015), 5237–5270.
- [9] Chen, Z.-Q., Kim, P. and Song, R.M.: Heat kernel estimates for the Dirichlet fractional Laplacian, *J. Eur. Math. Soc.*, **12** (2010), 1307–1329.
- [10] Chen, Z.-Q. and Kumagai, T.: Heat kernel estimates for stable-like processes on d -sets, *Stochastic Process. Appl.*, **108** (2003), 27–62.
- [11] Chen, Z.-Q. and Kumagai, T.: Heat kernel estimates for jump processes of mixed types on metric measure spaces, *Probab. Theory Related Fields*, **140** (2008), 277–317.
- [12] Cho, S., Kim, P., Song, R.M. and Vondracek, Z.: Factorization and estimates of Dirichlet heat kernels for non-local operators with critical killings, *J. Math. Pures Appl.*, **143** (2020), 208–256.
- [13] Chung, K.L. and Zhao, Z.: *From Brownian Motion to Schrödinger’s Equation*, Grundlehren der mathematischen Wissenschaften, **312**, Springer-Verlag, Berlin, 1995.
- [14] Davies, E.B. and Simon, B.: Ultracontractivity and heat kernels for Schrödinger operators and Dirichlet Laplacians, *J. Funct. Anal.*, **59** (1984), 335–395.
- [15] Demuth, M. and van Casteren, J.A.: *Stochastic Spectral Theory for Self-Adjoint Feller Operators. A Functional Analysis Approach*, Birkhäuser, Basel, 2000.
- [16] Frank, R., Merz, K. and Siedentop, H.: Relativistic strong Scott conjecture: a short proof, in: *Density Functionals for Many-Particle Systems. Mathematical Theory and Physical Applications of Effective Equations*, Englert, B.-G., et al. (eds.), Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap., vol. **41**, 69–79, World Scientific, Singapore, 2023.
- [17] Grzywny, T., Kim, K. and Kim, P.: Estimates of Dirichlet heat kernel for symmetric Markov processes, *Stochastic Process. Appl.*, **130** (2020), 431–470.
- [18] Kaleta, K. and Kulczycki, T.: Intrinsic ultracontractivity for Schrödinger operators based on fractional Laplacians, *Potential Anal.*, **33** (2010), 313–339.
- [19] Kaleta, K. and Lőrinczi, J.: Pointwise eigenfunction estimates and intrinsic ultracontractivity-type properties of Feynman–Kac semigroups for a class of Lévy processes, *Ann. Probab.*, **43** (2015), 1350–1398.
- [20] Kaleta, K. and Schilling, R.L.: Progressive intrinsic ultracontractivity and heat kernel estimates for non-local Schrödinger operators, *J. Funct. Anal.*, **279** (2020), 108606, 69 pp.
- [21] Kim, K. and Kim, P.: Two-sided estimates for the transition densities of symmetric Markov processes dominated by stable-like processes in $C^{1,\eta}$ open sets, *Stochastic Process. Appl.*, **124** (2014), 3055–3083.
- [22] Kulczycki, T. and Siudeja, B.: Intrinsic ultracontractivity of the Feynman–Kac semigroup for relativistic stable processes, *Trans. Amer. Math. Soc.*, **358** (2006), 5025–5057.
- [23] Kwaśnicki, M.: Ten equivalent definitions of the fractional Laplace operator, *Fract. Calc. Appl. Anal.*, **20** (2017), 7–51.
- [24] Lieb, E. and Seiringer, R.: *The Stability of Matter in Quantum Mechanics*, Cambridge University Press, Cambridge, 2009.
- [25] Schmüdgen, K.: *Unbounded Self-adjoint Operators on Hilbert Space*, Graduate Texts in Mathematics, vol. **265**, Springer, Dordrecht, 2012.
- [26] Song, R.M.: Two-sided estimates on the density of the Feynman–Kac semigroups of stable-like processes, *Electron. J. Probab.*, **11** (2006), 146–161.
- [27] Wang, F.-Y. and Wu, J.-L.: Compactness of Schrödinger semigroups with unbounded below potentials, *Bull. Sci. Math.*, **132** (2008), 679–689.