NONLINEAR YOSIDA APPROXIMATION AND MULTI-VALUED STOCHASTIC EVOLUTION INCLUSIONS[†]

WUJING FAN^a, WEI HONG^b, WEI LIU^{b1}

a. School of Statistics and Data Science, Nankai University, Tianjin 300071, China

b. School of Mathematics and Statistics, Jiangsu Normal University, Xuzhou 221116, China

ABSTRACT. In this paper, we develop a nonlinear version of the Yosida approximation to establish the existence and uniqueness of both (probabilistically) weak and strong solutions and demonstrate the continuous dependence on initial values for a class of multivalued stochastic evolution inclusions within the variational framework. Furthermore, leveraging this generalized Yosida approximation, we derive the finite-time extinction of solutions with probability one for multi-valued stochastic evolution inclusions perturbed by linear multiplicative noise. The main results are applicable to various examples, including multi-valued stochastic porous media equations, stochastic Φ -Laplace equations and stochastic evolution inclusions involving subdifferentials.

Keywords: Generalized Yosida approximation; Stochastic evolution inclusions; Variational framework; Maximal-monotone operators. Mathematics Subject Classification (2020): 60H15, 35R60

CONTENTS

1. Introduction	2
1.1. Variational framework	2
1.2. Finite-time extinction	4
2. Preliminaries	6
2.1. Maximal-monotone and pseudo-monotone operators	6
2.2. Generalized Yosida approximation	8
3. Well-posedness	10
3.1. Main results	10
3.2. Approximating sequences	13
3.3. Pre-compactness	16
3.4. Proof of Theorem 3.1	20
3.5. Proof of Theorem 3.3	29
3.6. Proof of Theorem 3.4	30
4. Finite time extinction	32
4.1. Main result	32

[†] This work is supported by National Key R&D program of China (No. 2023YFA1010101). The research of W. Hong is also supported by NSFC (No. 12401177) and NSF of Jiangsu Province (No. BK20241048). The research of W. Liu is also supported by NSFC (No. 12171208, 12090011, 12090010).

¹ Corresponding author: weiliu@jsnu.edu.cn

4.2. Approximating sequences	33
4.3. Proof of main result	35
5. Examples/Applications	38
5.1. Multi-valued stochastic porous media equations	38
5.2. Multi-valued stochastic Φ -Laplacian equations	41
5.3. SEIs with subdifferentials	45
6. Appendix	47
References	49

1. INTRODUCTION

The multi-valued stochastic evolution inclusion (SEI), which is given by the following form

$$dX(t) \in \mathcal{A}(t, X(t))dt + \sigma(t, X(t))dW(t),$$
(1.1)

is closely linked to a wide range of physical phenomena and models, such as self-organized criticality, fluid flow in porous media, the dynamics of non-Newtonian fluids and phase transition processes (cf. e.g. [6,8–10,20,21] and references therein). Moreover, it plays a pivotal role in the analysis of the Skorohod problem, which serves as a model for a class of reflected stochastic differential equations (SDEs) and constrained stochastic processes (cf. [60]).

As far as we know, the first result addressing multi-valued SEIs was conducted by Krée [27]. Subsequently, Cépa [19] investigated a class of SEIs associated with the Skorohod problem, while Petterson [41] employed the Yosida approximation technique to establish the existence of solutions to SEIs involving a maximal-monotone operator. Zhang [60] further generalized Cépa's results to infinite-dimensional settings and relaxed the Lipschitz continuity assumption for the single-valued operator to a global monotonicity condition. Additionally, Ren et al. [45, 46] investigated the long-time behaviour and the regularity of invariant measures to multi-valued SDEs.

On the other hand, the theory of maximal-monotone operators plays a crucial role in the analysis of nonlinear partial differential equations. In this paper, we focus on SEIs in infinite-dimensional spaces, where the multi-valued mapping \mathcal{A} in (1.1) is decomposed into a combination of a multi-valued maximal-monotone operator and a single-valued pseudomonotone operator. To address this problem, we will adopt a variational framework to investigate this class of SEIs.

1.1. Variational framework. Let H be a separable Hilbert space, with its inner product and norm denoted by $\langle \cdot, \cdot \rangle_H$ and $\|\cdot\|_H$, respectively. The dual space of H is denoted by H^* , which is identified with H through the Riesz isomorphism. Let V and its dual space V^* be uniformly convex Banach spaces such that the embedding $V \subset H$ is compact, continuous, and dense. Under these conditions, V and V^* are also strictly convex and reflexive (cf. [5]). The norms of V and V^* are donated by $\|\cdot\|_V$ and $\|\cdot\|_{V^*}$, respectively. Furthermore, the dual pairing between $v \in V^*$ and $u \in V$ is represented by $\langle v, u \rangle$. Through the Riesz

 $\mathbf{2}$

isomorphism, we construct the following Gelfand triple

$$V \subset H \simeq H^* \subset V^*. \tag{1.2}$$

Consequently, we have

$$\langle z, v \rangle = \langle z, v \rangle_H$$
, for all $z \in H, v \in V$.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space equipped with a normal filtration $\{\mathcal{F}_t\}_{t\geq 0}$. Let $(U, \langle \cdot, \cdot \rangle_U)$ be a separable Hilbert space, and let $(L_2(U, H), \|\cdot\|_{\mathcal{L}_2})$ denote the space of all Hilbert-Schmidt operators from U to H.

Let T > 0. In this paper we focus on the following type multi-valued SEIs

$$dX(t) + B(t, X(t))dt \in -\mathcal{A}(t, X(t))dt + \sigma(t, X(t))dW(t), \ X(0) = x \in H,$$
(1.3)

where the multi-valued (maximal-monotone) mapping \mathcal{A} and the single-valued mappings B, σ are defined as follows

$$\mathcal{A} : [0,T] \times V \to 2^{V^*},$$

$$B : [0,T] \times V \to V^*,$$

$$\sigma : [0,T] \times V \to L_2(U,H)$$

Here W(t) denotes a cylindrical Wiener process on U w.r.t. the filtration $\{\mathcal{F}_t\}_{t\geq 0}$. Furthermore, the mappings \mathcal{A}, B, σ are assumed to be measurable.

The classical variational framework and the theory of monotone operators were first introduced by Minty [35]. This foundational work was further developed by Browder [14,15], Leray and Lions [29] as well as Hartman and Stampacchia [24]. Afterwards, the variational approach for stochastic partial differential equations (SPDEs) was originally developed by Pardoux [39,40]. This approach was subsequently explored by Krylov and Rozovskii [28] and Gyongy [23]. Based on these developments, Liu and Röckner [31] and Liu [32] extended the variational framework to encompass a class of locally monotone operators, broadening the range of nonlinear SPDEs, including e.g. stochastic 2D Navier-Stokes equations, that can be analyzed. Very recently, Röckner et al. [47] further generalized the results of [31, 32] to SPDEs with multiplicative noises and fully local monotonicity conditions. For additional applications and advancements of the variational approach to SPDEs, we refer the reader to [34, 42, 44, 48, 50, 61].

Notice that all the aforementioned works focus on the single-valued case, there are much less results in the literature that can be applied to multi-valued SEIs. As previously mentioned, Zhang [60] investigated the multi-valued SEIs (1.3) within the Gelfand triple (1.2). However, the multi-valued operator \mathcal{A} in their study is restricted to the space H (i.e., $\mathcal{A} : H \to 2^H$), which excludes some model of nonlinear SEIs, such as multi-valued stochastic porous media equations. Liu and Stephan [33] made a progress by establishing the existence and uniqueness of solutions to a class of multi-valued SEIs (1.3) driven by Lévy noises. In their work, the multi-valued operator \mathcal{A} is assumed to be coercive with an exponent $\alpha \in (1, 2]$, and the single-valued operator B is required to be Lipschitz continuous. Gess and Tölle [21] studied multi-valued, monotone, weakly dissipative SEIs, which extend to cases where the coercivity exponent $\alpha \in [1, 2]$, thereby covering certain stochastic singular diffusion equations.

However, the well-posedness result of multi-valued SEIs within the variational framework for coercivity exponent $\alpha > 2$ remains an open problem. We would like to highlight the main challenges in studying multi-valued SEIs with general coercivity exponent $\alpha > 1$. Specifically, the classical Yosida approximation (cf. [5]), commonly employed to handle maximal-monotone operators, only works for the case of $\alpha \in (1, 2]$ due to certain technical limitations, as pointed out in [33]. In particular, the coercivity of the Yosida approximation operators fails to hold when $\alpha > 2$ (cf. Lemma 3.10 in [33]). Consequently, to address the general case $\alpha > 2$, several new techniques involving the generalized Yosida approximation are required.

More precisely, we introduce a nonlinear version of the Yosida approximation by employing a novel duality mapping (cf. Definition 2.2 and (2.5)), which is also of independent interest in analysis. To achieve this, we first establish the existence and uniqueness of solutions to an associated resolvent equation (cf. (2.2) below). Subsequently, we derive several key properties of the generalized Yosida approximation, which are crucial for the convergence of the approximating sequence.

To develop a more general theory for the existence of (probabilistically) weak solutions to multi-valued SEIs (1.3), we assume the single-valued operator B to be pseudo-monotone. While the monotonicity technique is effective for SPDEs with single-valued monotone operators (cf. [34]), they fail in addressing pseudo-monotone operators. To solve this problem, we combine stochastic compactness arguments with pseudo-monotonicity techniques. However, due to the joint perturbation caused by multi-valued maximal-monotone operators, several non-trivial difficulties arise in proving the convergence of the coefficient $B(t, X_{\lambda}(t))$ and the Yosida approximation operator $\mathcal{A}_{\lambda}(t, X_{\lambda}(t))$, as detailed in Lemmas 3.8 and 3.9. Furthermore, we establish the existence of a unique (probabilistically) strong solution and demonstrate the continuous dependence on initial values for (1.3) under fully local monotonicity conditions on single-valued operators.

1.2. Finite-time extinction. Self-organized criticality (SOC) is a widely studied framework in physics for modeling complex systems, including phenomena such as earthquakes and neuronal activity. The SOC behavior observed in the Bak-Tang-Wiesenfeld (BTW) sandpile model corresponds to the finite-time extinction of solutions to multi-valued stochastic diffusion equations (cf. [6]). Similarly, the Stefan problem, which describes phase transitions in porous media or heat conduction, can be formulated as a class of multi-valued stochastic porous media equations, see also [6]. The time evolution of these systems exhibits SOC, underscoring the fundamental role of finite-time extinction in the analysis of multi-valued SEIs.

Several previous works have explored the finite-time extinction of solutions to multivalued SEIs perturbed by linear multiplicative noises. For example, Barbu et al. [6] proved that, for multi-valued stochastic porous media equations in one spatial dimension, finitetime extinction occurs with positive probability for small initial values. In their subsequent work [7], the restriction to one dimension was relaxed, extending the results to all dimensions. In [9], the asymptotic extinction of solutions to multi-valued stochastic porous media equations was established with probability one in dimensions $d = \{1, 2, 3\}$. Röckner and Wang [49] generalized the results of [6] to encompass a broader class of operators, including fractional Laplacians. They proved the exponential integrability of the extinction time for the Zhang model, provided the noise is small enough. Gess [20] showed that the finite-time extinction of the BTW model holds with probability one in all dimensions.

Concerning finite time extinction, we distinguish the following situations:

- (i) Extinction with positive probability for small initial conditions;
- (ii) Extinction with positive probability for all initial values;
- (iii) Finite time extinction with probability one for all initial values.

While both Situations (i) and (ii) are mathematically intriguing, the robustness of the relaxation into subcritical states in SOC is of fundamental importance in physics. Consequently, Situation (iii) aligns most closely with the perspective of SOC.

In this work, we establish finite-time extinction with probability one for all initial values $x \in H$ (i.e., Situation (iii)) for a class of multi-valued SEIs perturbed by linear multiplicative noise. As a consequence in the proof, we demonstrate the L^2 -convergence of the Yosida approximation sequence (see Lemma 4.1). Additionally, we provide a quantitative characterization of the probability that the extinction time τ_e is less than any given time T, i.e.

$$\mathbb{P}(\tau_e \leqslant T) \ge 1 - \frac{c^* \|x\|_H^{2-\alpha}}{T},$$

for multi-valued SEIs in the variational framework with any coercivity exponent $\alpha \in (1, 2)$. In comparison to existing works [6–8,10], our results extend the finite-time extinction from cases with positive probability and small initial conditions to cases with probability one for all initial values. Moreover, we also remove the restriction on spatial dimension.

To illustrate the generality of the present framework, we apply the main results to several concrete examples. The first application concerns the multi-valued stochastic porous media equations

$$\begin{cases} dX(t) \in \Delta \Psi(X(t))dt + \sigma(t, X(t))dW(t), \\ X(0) = x \in H^{-1}(\Lambda). \end{cases}$$

A typical example of the multi-valued mapping Ψ is given by

$$\Psi(s) = \begin{cases} \rho + \delta s^{p-1}, & \text{if } s > 0, \\ [-\rho, \rho], & \text{if } s = 0, \\ -\rho + \delta (-s)^{p-1}, & \text{if } s < 0, \end{cases}$$

where ρ, δ are some positive constants. This example was involved in [33] for $p \in (1, 2]$. In this work, we extend the well-posedness result to all p > 1 and further establish finite-time extinction of solutions with probability one for all initial values when $p \in (1, 2)$.

Next, we investigate the multi-valued stochastic Φ -Laplace equations

$$\begin{cases} dX(t) \in \operatorname{div}\Phi(\nabla X(t))dt + \sigma(t, X(t))dW(t), \\ X(0) = x \in L^2(\Lambda), \end{cases}$$
(1.4)

which have been studied in e.g. [21, 52] for specific potential functions Φ (see Subsection 5.2 for details). We develop a general well-posedness result and also for the first time establish finite-time extinction results for (1.4), which is of independent interest.

Finally, we apply our results to SEIs with subdifferentials

$$\begin{cases} \partial_t u(t,x) \in [\Delta u(t,x) - g(t,x,u(t,x),\nabla u(t,x)) - \partial \varphi(u(t,x))] \, dt + \sigma(t,u(t,x)) dW(t), \\ u(0,x) = u_0(x) \in L^2(\Lambda). \end{cases}$$

This type of SEI with single-valued monotone operators was previously studied in [60]. Within our framework, we extend the results of [60] to more general operators, broadening their applicability to a wider class of nonlinear parabolic equations.

The rest of the paper is structured as follows: Section 2 gives some preliminaries, including the definition and properties of the maximal-monotone operator and the generalized Yosida approximation. Section 3 presents the weak and strong well-posedness results. Section 4 discusses the finite-time extinction of solutions to multi-valued SEIs. Section 5 illustrates several concrete applications of our general framework. Finally, we provide several useful results related to multi-valued maximal-monotone operators in the Appendix.

2. Preliminaries

In this section, we recall some definitions and basic results regarding maximal monotone and pseudo monotone operators on Banach spaces. Then, we construct a nonlinear version of Yosida approximation for maximal monotone operators on Banach spaces by solving a newly formulated resolvent equation.

We denote the power set of V^* by 2^{V^*} . A multi-valued mapping $\mathcal{A} : V \to 2^{V^*}$ maps $x \in V$ to a subset $\mathcal{A}(x) \subset V^*$. The domain of \mathcal{A} is defined as the set $\mathcal{D}(\mathcal{A}) := \{x \in V \mid \mathcal{A}(x) \neq \emptyset\}$. For a multi-valued map \mathcal{A} , its graph is given by

$$\mathcal{G}(\mathcal{A}) := \{ [x, y] \in V \times V^* \mid y \in \mathcal{A}(x) \} \,.$$

2.1. Maximal-monotone and pseudo-monotone operators. First, we recall the definition of maximal-monotone operators.

Definition 2.1. (i) A multi-valued operator $\mathcal{A}: V \to 2^{V^*}$ is called monotone if

$$\langle v_1 - v_2, u_1 - u_2 \rangle \ge 0$$
, for all $[v_i, u_i] \in \mathcal{G}(\mathcal{A}), i = 1, 2$.

(ii) A monotone operator $\mathcal{A}: V \to 2^{V^*}$ is called maximal-monotone if it is not properly contained in any other monotone extension $\tilde{\mathcal{A}}$ such that $\mathcal{G}(\mathcal{A}) \subsetneq \mathcal{G}(\tilde{\mathcal{A}})$.

(iii) The minimal section $\mathcal{A}^0 : \mathcal{D}(\mathcal{A}) \subset V \to 2^{V^*}$ of a maximal-monotone operator \mathcal{A} is defined for $x \in \mathcal{D}(\mathcal{A})$ as

$$\mathcal{A}^{0}(x) := \Big\{ y \in \mathcal{A}(x) \mid \|y\|_{V^{*}} = \min_{z \in \mathcal{A}(x)} \|z\|_{V^{*}} \Big\}.$$

Remark 2.1. We note that since V^* is strictly convex, then \mathcal{A}^0 is single-valued (cf. [5]).

The following lemma concerns the convergence of maximal-monotone operators.

Lemma 2.1. (cf. [5, Lemma 2.3]) Let $\mathcal{A}: V \to 2^{V^*}$ be maximal-monotone. Let $[u_n, v_n] \in \mathcal{G}(\mathcal{A})$ be such that $u_n \rightharpoonup u$, $v_n \rightharpoonup v$, and either

$$\limsup_{n,m\to\infty} \left\langle u_n - u_m, v_n - v_m \right\rangle \leqslant 0$$

or

$$\limsup_{n \to \infty} \langle u_n - u, v_n - v \rangle \leq 0.$$

Then, $[u, v] \in \mathcal{G}(\mathcal{A})$ and $\langle u_n, v_n \rangle \to \langle u, v \rangle$, as $n \to \infty$.

In the sequel, we introduce the definition of duality mapping, which differs from the classical one presented in the existing works (e.g. [5]) and plays an important role in

classical one presented in the existing works (e.g. [5]) and plays an important role in applications to examples of multi-valued nonlinear stochastic evolution inclusions.

Definition 2.2. Let $\alpha > 1$. The duality mapping $J : V \to 2^{V^*}$ on the space V is defined by

$$J(u) := \left\{ v \in V^* \mid \langle v, u \rangle = \|u\|_V^\alpha = \|v\|_{V^*}^{\frac{\alpha}{\alpha-1}} \right\}, \quad \text{for each } u \in V.$$

Proposition 2.1. (i) The duality mapping J is monotone, locally bounded and coercive.

(ii) J is single-valued, demicontinuous and odd, and the set J(x) is closed and convex for any $x \in V$.

Proof. **Proof of (i)**: Let $u_1, u_2 \in V$ and $v_1 \in J(u_1), v_2 \in J(u_2)$. In view of Definition 2.2 and $\alpha > 1$, we have

$$\langle v_1 - v_2, u_1 - u_2 \rangle \geq \langle v_1, u_1 \rangle + \langle v_2, u_2 \rangle - \|v_1\|_{V^*} \|u_2\|_V - \|v_2\|_{V^*} \|u_1\|_V$$

$$= \|u_1\|_V^{\alpha} + \|u_2\|_V^{\alpha} - \|u_1\|_V^{\alpha-1} \|u_2\|_V - \|u_2\|_V^{\alpha-1} \|u_1\|_V$$

$$= (\|u_1\|_V^{\alpha-1} - \|u_2\|_V^{\alpha-1})(\|u_1\|_V - \|u_2\|_V)$$

$$\geq 0.$$

$$(2.1)$$

Thus, J is monotone.

Let $u \in V$ and $v \in J(u)$. Since $\alpha > 1$, $||v||_{V^*} = ||u||_V^{\alpha-1}$ and $\langle v, u \rangle = ||u||_V^{\alpha}$, we deduce that J is locally bounded and coercive.

Proof of (ii): The proof is postponed in Proposition 6.1 in the Appendix.

Remark 2.2. Note that, applying the Hahn-Banach theorem, J(u) is non-empty for every $u \in V$, which implies that $\mathcal{D}(J) = V$. Recall that J is a single-valued mapping. Since J is monotone, demicontinuous and coercive, it follows from Theorem 26.A in [59] that J is surjective. Consequently, the mapping $J: V \to V^*$ is a bijection.

Furthermore, since J is monotone and demicontinuous, we can conclude that J is maximal-monotone by Proposition 32.7 in [59]. Additionally, since V^* is uniformly convex, it implies from Proposition 21.23 (d) in [59] that J is continuous.

Definition 2.3. The inverse mapping $J^{-1}: V^* \to 2^V$ defined by

$$J^{-1}(v) := \{ u \in V \mid v \in J(u) \}$$

satisfies

$$J^{-1}(v) = \left\{ u \in V \mid \langle v, u \rangle = \|u\|_{V}^{\alpha} = \|v\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} \right\}, \text{ for each } v \in V^{*}.$$

Remark 2.3. Since V is reflexive, J^{-1} is the duality mapping on V^* and $\mathcal{D}(J^{-1}) = V^*$. Moreover, due to the strict convexity of V, we know that J^{-1} is also single-valued (from V^* to V) and demicontinuous.

In what follows, we also recall the definition of pseudo-monotone operators, which was first proposed by Brézis in [12].

Definition 2.4. The operator $B: V \to V^*$ is called pseudo-monotone if $u_n \rightharpoonup u$ in V and

$$\limsup_{n \to \infty} \langle B(u_n), u_n - u \rangle \leqslant 0,$$

then for any $v \in V$,

$$\langle B(u), u-v \rangle \leq \liminf_{n \to \infty} \langle B(u_n), u_n-v \rangle.$$

Remark 2.4. There is an alternative definition of pseudo-monotonicity introduced by Browder in [16]. Specifically, an operator $B: V \to V^*$ is pseudo-monotone iff $u_n \rightharpoonup u$ in V and

$$\limsup_{n \to \infty} \langle B(u_n), u_n - u \rangle \leqslant 0$$

then $B(u_n) \rightharpoonup B(u)$ in V^* and

 $\lim_{n \to \infty} \langle B(u_n), u_n \rangle = \langle B(u), u \rangle.$

It has been proved that these two definitions are equivalent, which can be referred to Remark 5.2.12 in [34].

2.2. Generalized Yosida approximation. To construct the generalized Yosida approximation of maximal-monotone operators on Banach spaces, we first consider the following resolvent equation

$$0 \in J(x_{\lambda} - x) + \lambda \mathcal{A}(x_{\lambda}), \tag{2.2}$$

where the operator \mathcal{A} is maximal-monotone, $\lambda > 0$ and $x \in V$.

Remark 2.5. It should be pointed out that since the definition of the duality mapping J differs from the classical one (see e.g. [5]), the well-posedness of the resolvent equation (2.2) and the properties of the generalized Yosida approximation require to be established.

The existence and uniqueness of solutions to Eq. (2.2) are presented as follows.

Proposition 2.2. For all $x \in V$, there exists a unique solution x_{λ} to (2.2).

Proof. Recall that V is a reflexive Banach space, $\mathcal{A}: V \to 2^{V^*}$ is maximal-monotone and $J: V \to V^*$ is demicontinuous and monotone. It follows from Corollary 2.6 in [5] that for any $\lambda > 0$ and $x \in V$, the operator

$$J(\cdot - x) + \lambda \mathcal{A}$$

is maximal-monotone. Furthermore, since J is coercive and \mathcal{A} is monotone, we deduce that $J(\cdot - x) + \lambda \mathcal{A}$ is also coercive for any $\lambda > 0$ and $x \in V$. Thus, by Corollary 2.2 in [5], there exists $x_{\lambda} \in V$ such that

$$0 \in J(x_{\lambda} - x) + \lambda \mathcal{A}(x_{\lambda}).$$

Now, we show the uniqueness of solutions to Eq. (2.2) by contradiction. We assume that the solutions of Eq. (2.2) are not unique, i.e., there exist $x_{\lambda}^1, x_{\lambda}^2$ that satisfy Eq. (2.2). Then, there exist $y_{\lambda}^1 \in \mathcal{A}(x_{\lambda}^1)$ and $y_{\lambda}^2 \in \mathcal{A}(x_{\lambda}^2)$ such that

$$J(x_{\lambda}^{1} - x) + \lambda y_{\lambda}^{1} = J(x_{\lambda}^{2} - x) + \lambda y_{\lambda}^{2}.$$

It follows that

$$\langle J(x_{\lambda}^{1}-x) - J(x_{\lambda}^{2}-x), x_{\lambda}^{1}-x - (x_{\lambda}^{2}-x) \rangle = -\lambda \langle y_{\lambda}^{1}-y_{\lambda}^{2}, x_{\lambda}^{1}-x_{\lambda}^{2} \rangle \leqslant 0.$$
(2.3)

On the other hand, by the definition of J we have

$$\langle J(x_{\lambda}^{1} - x) - J(x_{\lambda}^{2} - x), x_{\lambda}^{1} - x - (x_{\lambda}^{2} - x) \rangle \geq (\|x_{\lambda}^{1} - x\|_{V}^{\alpha - 1} - \|x_{\lambda}^{2} - x\|_{V}^{\alpha - 1})(\|x_{\lambda}^{1} - x\|_{V} - \|x_{\lambda}^{2} - x\|_{V}) \geq 0.$$

$$(2.4)$$

Combining (2.3) and (2.4), we can deduce that

$$\langle J(x_{\lambda}^1 - x), x_{\lambda}^2 - x \rangle = \langle J(x_{\lambda}^2 - x), x_{\lambda}^1 - x \rangle = \|x_{\lambda}^1 - x\|_V^{\alpha} = \|x_{\lambda}^2 - x\|_V^{\alpha}.$$

In view of the definition of J and the fact that J is single-valued, it follows that

$$J(x_{\lambda}^{1} - x) = J(x_{\lambda}^{2} - x).$$

Finally, since J^{-1} is single-valued, we conclude that

$$x_{\lambda}^1 = x_{\lambda}^2.$$

The proof is complete.

Now, let $\mathcal{R}_{\lambda}: V \to V$ be the resolvent operator of \mathcal{A} , which is defined by

$$\mathcal{R}_{\lambda}(x) := x_{\lambda},$$

where x_{λ} is the unique solution of (2.2). Then, the generalized Yosida approximation $\mathcal{A}_{\lambda}: V \to V^*$ of \mathcal{A} is defined by

$$\mathcal{A}_{\lambda}(x) := \frac{1}{\lambda} J(x - \mathcal{R}_{\lambda}(x)), \qquad (2.5)$$

where $\lambda > 0$ and $x \in V$. By Definition 2.2 and Proposition 2.2, it is known that for each $x \in V$ and $\lambda > 0$, the mapping $\mathcal{A}_{\lambda}(\cdot)$ is single-valued.

Remark 2.6. Since Definition 2.2 extends the classical duality mapping, which reduces to the special case of $\alpha = 2$, the generalized Yosida approximation (2.5) offers a more natural extension compared to the standard Yosida approximation. Specifically, for any $\alpha > 1$, if the maximal monotone operator \mathcal{A} is coercive, its generalized Yosida approximation operator \mathcal{A}_{λ} retains coercivity. In contrast, the standard Yosida approximation is limited to $\alpha \in (1, 2]$, as demonstrated in Lemma 3.10 in [33]. Furthermore, the current definition is pivotal in establishing the convergence of \mathcal{A}_{λ} for the general case $\alpha > 1$, as illustrated in the proof of Lemma 3.9.

The properties of the Yosida approximation (2.5) are collected in the following.

Proposition 2.3. The generalized Yosida approximation \mathcal{A}_{λ} has the following properties: (i) \mathcal{A}_{λ} is monotone, bounded on bounded sets, and demicontinuous from V to V^{*}.

- (ii) $\|\mathcal{A}_{\lambda}(x)\|_{V^*} \leq \|\mathcal{A}^0(x)\|_{V^*}$, for each $x \in \mathcal{D}(\mathcal{A})$ and $\lambda > 0$.
- (iii) $\mathcal{A}_{\lambda}(x) \to \mathcal{A}^{0}(x)$ in V^{*} , as $\lambda \to 0$, for each $x \in \mathcal{D}(\mathcal{A})$.
- (iv) For any $x \in V$, $\mathcal{A}_{\lambda}(x) \in \mathcal{A}(\mathcal{R}_{\lambda}(x))$.

Proof. Since J is odd, it follows from (2.2), (2.5), and the definition of \mathcal{R}_{λ} that $\mathcal{A}_{\lambda}(x) \in \mathcal{A}(\mathcal{R}_{\lambda}(x))$ for all $x \in V$, which yields claim (iv). We now proceed to prove claims (i), (ii), and (iii) in order.

Proof of (i): First, it clear that $\mathcal{A}_{\lambda} : V \to V^*$ is monotone and demicontinuous. Let $[u, v] \in \mathcal{G}(\mathcal{A})$. By the definition of $\mathcal{A}_{\lambda}(x)$ and the monotonicity of \mathcal{A} , we can obtain

$$\langle J(\mathcal{R}_{\lambda}(x)-x), \mathcal{R}_{\lambda}(x)-u \rangle \leq \lambda \langle v, u - \mathcal{R}_{\lambda}(x) \rangle,$$

which implies

$$\begin{aligned} &\|\mathcal{R}_{\lambda}(x) - x\|_{V}^{\alpha} \\ &\leqslant \|x - u\|_{V} \|\mathcal{R}_{\lambda}(x) - x\|_{V}^{\alpha-1} + \lambda \|x - u\|_{V} \|v\|_{V^{*}} + \lambda \|v\|_{V^{*}} \|\mathcal{R}_{\lambda}(x) - x\|_{V} \\ &\leqslant \frac{1}{2} \|\mathcal{R}_{\lambda}(x) - x\|_{V}^{\alpha} + C \Big(\|x - u\|_{V}^{\alpha} + \|x - u\|_{V} \|v\|_{V^{*}} + \|v\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} \Big). \end{aligned}$$

Then, it follows that the operators \mathcal{R}_{λ} and \mathcal{A}_{λ} are bounded on bounded sets.

Proof of (ii): Let $[x, y] \in \mathcal{G}(\mathcal{A})$. By the monotonicity of \mathcal{A} , we have

$$0 \leqslant \langle y - \mathcal{A}_{\lambda}(x), x - \mathcal{R}_{\lambda}(x) \rangle \leqslant \|y\|_{V^*} \|x - \mathcal{R}_{\lambda}(x)\|_V - \frac{1}{\lambda} \|x - \mathcal{R}_{\lambda}(x)\|_V^{\alpha}.$$

It follows that

$$\lambda \|\mathcal{A}_{\lambda}(x)\|_{V^*} = \|x - \mathcal{R}_{\lambda}(x)\|_{V}^{\alpha - 1} \leq \lambda \|y\|_{V^*}, \ \forall y \in \mathcal{A}(x).$$

Thus, we deduce that for any $x \in \mathcal{D}(\mathcal{A})$ and $\lambda > 0$,

$$\|\mathcal{A}_{\lambda}(x)\|_{V^*} \leqslant \|\mathcal{A}^0(x)\|_{V^*}.$$

Proof of (iii): The proof of claim (iii) is similar to that of Proposition 2.2 (v) in [5], we outline it here for reader's convenience. Let $x \in \mathcal{D}(\mathcal{A})$ and consider $\lambda \to 0$ such that

$$\mathcal{A}_{\lambda}(x) \rightharpoonup y \text{ in } V^*.$$

As shown in the proof of Proposition 2.2 (iv) in [5], we have $y \in \mathcal{A}(x)$. Furthermore, since $\|\mathcal{A}_{\lambda}(x)\|_{V^*} \leq \|\mathcal{A}^0(x)\|_{V^*}$, it implies that $y = \mathcal{A}^0(x)$. Since V^* is uniformly convex, by Lemma 1.1 in [5] it follows that $\mathcal{A}_{\lambda}(x) \to \mathcal{A}^0(x)$ in V^* . This completes the proof. \Box

3. Well-posedness

3.1. Main results. In this subsection, we show the existence and uniqueness of weak and strong solutions and the continuous dependence on initial values to Eq. (1.3). We first recall the definition of (probabilistically) weak solutions.

Definition 3.1. A couple (X, η, W) is called a (probabilistically) weak solution to Eq. (1.3), if there exists a stochastic basis $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,T]}, \mathbb{P})$ such that $X \in L^{\alpha}([0,T] \times \Omega, V) \cap$ $L^2([0,T] \times \Omega, H), \eta \in L^1([0,T] \times \Omega, V^*)$ and W is an U-valued cylindrical Wiener process on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,T]}, \mathbb{P})$, which satisfy

- (*i*) $X \in C([0,T],H), \mathbb{P}$ -a.s.;
- (ii) X and $\int_0^{\cdot} \eta(s) ds$ are (\mathcal{F}_t) -adapted;
- (*iii*) $\eta \in \mathcal{A}(\cdot, X(\cdot)), dt \otimes \mathbb{P}$ -a.e.;
- (iv) for all $t \in [0,T]$, the following equality holds in V^*

$$X(t) = x - \int_0^t (\eta(s) + B(s, \bar{X}(s))) ds + \int_0^t \sigma(s, \bar{X}(s)) dW(s), \ \mathbb{P}\text{-}a.s.,$$

where \overline{X} is any V-valued progressively measurable $dt \otimes \mathbb{P}$ -version of X.

We assume that there exist $f \in L^1([0,T], [0,\infty))$, $C, \delta > 0, \beta \ge 0$ and $\alpha > 1$ such that the following conditions hold for a.e. $t \in [0,T]$.

More precisely, for the multi-valued operator \mathcal{A} , we suppose that

 $(\mathbf{H}^1_{\mathbf{A}})$ (Maximal monotonicity): For any $x, y \in V$,

 $\langle v - w, x - y \rangle \ge 0$, for any $v \in \mathcal{A}(t, x), w \in \mathcal{A}(t, y)$,

and $\mathcal{A}(t, \cdot)$ is maximal-monotone.

 $(\mathbf{H}^2_{\mathcal{A}})$ (Coercivity): For any $x \in V$ and $v \in \mathcal{A}(t, x)$,

$$\langle v, x \rangle \ge \delta \|x\|_V^{\alpha} - f(t).$$

 $(\mathbf{H}^3_{\mathcal{A}})$ (Growth): For any $x \in V$,

$$\|\mathcal{A}^{0}(t,x)\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} \leq (f(t)+C\|x\|_{V}^{\alpha})(1+\|x\|_{H}^{\beta}).$$

For mappings B and σ , we suppose that

 (\mathbf{H}_B^1) (Weak coercivity): For any $x \in V$,

$$2\langle B(t,x),x\rangle \ge -f(t)(1+\|x\|_H^2)$$

 (\mathbf{H}_B^2) (Growth): For any $x \in V$,

$$||B(t,x)||_{V^*}^{\frac{\alpha}{\alpha-1}} \leq (f(t) + C||x||_V^{\alpha})(1 + ||x||_H^{\beta}).$$

 (\mathbf{H}_{σ}) For any sequence $\{x_n\}_{n=1}^{\infty}$ and x in V satisfying $||x_n - x||_H \to 0$,

$$\|\sigma(t, x_n) - \sigma(t, x)\|_{\mathcal{L}_2} \to 0.$$

Moreover, for any $x \in V$,

$$\|\sigma(t,x)\|_{\mathcal{L}_2}^2 \leqslant f(t)(1+\|x\|_H^2).$$

In the sequel, we state the first main result of this work concerning the existence of weak solutions to Eq. (1.3).

Theorem 3.1. Assume that the operator $B(t, \cdot)$ is pseudo-monotone for a.e. $t \in [0, T]$, and that $(\mathbf{H}^{1}_{\mathcal{A}}) \cdot (\mathbf{H}^{3}_{\mathcal{A}})$, (\mathbf{H}^{1}_{B}) , (\mathbf{H}^{2}_{B}) and (\mathbf{H}_{σ}) hold. Then for any $x \in H$, there exists a (probabilistically) weak solution to Eq. (1.3). In addition, for any $p \ge 2$, the following moment estimate holds

$$\mathbb{E}\left[\sup_{t\in[0,T]}\|X(t)\|_{H}^{p}\right] + \mathbb{E}\left\{\left(\int_{0}^{T}\|X(s)\|_{V}^{\alpha}ds\right)^{\frac{p}{2}}\right\} < \infty.$$
(3.1)

Remark 3.2. The classical Skorohod problem was investigated in foundational works such as [30, 51, 53, 55]. Cépa [19] demonstrated that the Skorohod problem is equivalent to a multi-valued maximal monotone operator equation and established an existence result for its generalized form. Further developments in this topic can be found in [45, 57, 60] and reference therein. In this work, we present a more general existence result of weak solutions to SEIs, extending the existing works by incorporating multi-valued maximal-monotone operators and single-valued pseudo-monotone operators within a variational framework. This generalization encompasses a broader class of nonlinear SEIs, as illustrated in Section 5.

Next, we consider the existence and uniqueness of (probabilistically) strong solutions to Eq. (1.3). The definitions of the (probabilistically) strong solutions and the pathwise uniqueness are presented as follows.

Definition 3.2. A couple (X, η) is called a (probabilistically) strong solution to Eq. (1.3), if for every probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,T]}, \mathbb{P})$ with an U-valued cylindrical Wiener process W, we have $X \in L^{\alpha}([0,T] \times \Omega, V) \cap L^2([0,T] \times \Omega, H)$ and $\eta \in L^1([0,T] \times \Omega, V^*)$ satisfy (i)-(iv) in Definition 3.1.

Definition 3.3. We say that pathwise uniqueness holds for Eq. (1.3), if whenever the probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,T]}, \mathbb{P})$ and the U-valued cylindrical Wiener process W are fixed, two solutions X and X' such that $X(0) = X'(0) \mathbb{P}$ -a.s., then \mathbb{P} -a.s.

$$X(t) = X'(t), t \in [0, T].$$

In order to guarantee the existence and uniqueness of (probabilistically) strong solutions to Eq. (1.3), we assume the hemicontinuity and local monotonicity conditions instead of the pseudo-monotonicity as follows.

- (\mathbf{H}_B^3) (Hemicontinuity): The map $s \mapsto \langle B(t, x + sy), v \rangle$ is continuous on \mathbb{R} for any $x, y, v \in V$.
- (\mathbf{H}_{B}^{4}) (Local monotonicity): For any $x, y \in V$,

$$-2\langle B(t,x) - B(t,y), x - y \rangle + \|\sigma(t,x) - \sigma(t,y)\|_{\mathcal{L}_2}^2 \leq (f(t) + \rho(x) + \zeta(y))\|x - y\|_H^2,$$

where ρ and ζ are measurable functions from V to \mathbb{R}_+ satisfying

$$\rho(x) + \zeta(x) \leqslant C(1 + \|x\|_V^{\alpha})(1 + \|x\|_H^{\beta}).$$

The following discusses the pseudo-monotonicity of B under assumptions (\mathbf{H}_B^3) and (\mathbf{H}_B^4) , whose proof can refer to [32].

Lemma 3.1. Assume that (\mathbf{H}_B^3) and (\mathbf{H}_B^4) hold. Then $B(t, \cdot)$ is pseudo-monotone from V to V^* for a.e. $t \in [0, T]$.

Now, we state the existence and uniqueness of (probabilistically) strong solutions to Eq. (1.3).

Theorem 3.3. Assume that $(\mathbf{H}^{1}_{\mathcal{A}})$ - $(\mathbf{H}^{3}_{\mathcal{A}})$, (\mathbf{H}^{1}_{B}) - (\mathbf{H}^{4}_{B}) and (\mathbf{H}_{σ}) hold. Then for any $x \in H$, there exists a unique (probabilistically) strong solution to Eq. (1.3) and the estimate (3.1) holds.

The continuous dependence on initial data is also given as follows.

Theorem 3.4. Suppose that the assumptions in Theorem 3.3 hold. Let $\{x_n\}_{n=1}^{\infty}$ and x be a sequence with $||x_n - x||_H \to 0$ and X(t, x) be the unique solution of Eq. (1.3) with initial value x. Then, for any p > 0,

$$\lim_{n \to \infty} \mathbb{E} \Big[\sup_{t \in [0,T]} \left\| X(t,x_n) - X(t,x) \right\|_H^p \Big] = 0.$$

Remark 3.5. (i) The existence and uniqueness of solutions to multi-valued SEIs driven by Lévy noise were established in [33]. Theorems 3.3 and 3.4 extend this result by allowing arbitrary coercivity exponents $\alpha > 1$, whereas [33] is limited to $\alpha \in (1, 2]$. Moreover, we establish the well-posedness of multi-valued SEIs with single-valued fully local monotone operators, relaxing the Lipschitz continuity assumption imposed in [33]. This result is applicable to a broader class of multi-valued stochastic porous media equations, thereby extending the previous works [6, 7] (see Subsection 5.1).

(ii) Theorem 3.3 also generalizes the results in [21] by permitting a general polynomial growth condition for the maximal-monotone operator (i.e., condition $(\mathbf{H}_{\mathcal{A}}^3)$), whereas [21] is restricted to the linear growth case. This generalization encompasses a wider class of multi-valued stochastic Φ -Laplace equations, such as p-Laplace equations (with p > 1) and Non-Newtonian fluid models, as detailed in Subsection 5.2.

Remark 3.6. Note that Theorem 3.4 implies the Feller property of the transition semigroup $P_t : C_b(H) \to C_b(H)$ associated with Eq. (1.3). Building on this result, in forthcoming work, we aim to investigate the existence and uniqueness of invariant probability measures for Eq. (1.3) by employing the generalized Yosida approximation developed in this work.

3.2. Approximating sequences. In order to prove the existence of (probabilistically) weak solutions to Eq. (1.3), we consider the following approximating equations

$$\begin{cases} dX_{\lambda}(t) + [\mathcal{A}_{\lambda}(t, X_{\lambda}(t)) + B(t, X_{\lambda}(t))] dt = \sigma(t, X_{\lambda}(t)) dW(t), \\ X_{\lambda}(0) = x \in H, \end{cases}$$
(3.2)

where \mathcal{A}_{λ} is the generalized Yosida approximation of \mathcal{A} given by (2.5).

The following lemma gives the existence of (probabilistically) weak solutions to Eq. (3.2).

Lemma 3.2. Suppose that the assumptions in Theorem 3.1 hold. For any $\lambda \in (0, \delta^{-1})$ and initial value $x \in H$, there exists a weak solution X_{λ} to Eq. (3.2). *Proof.* It suffices to justify the assumptions in Corollary 2.7 in [47]. Let $\lambda \in (0, \delta^{-1})$. Denote

$$A := -\mathcal{A}_{\lambda} - B.$$

Since $\mathcal{A}_{\lambda}(t, \cdot)$ is monotone and demicontinuous and $B(t, \cdot)$ is pseudo-monotone for a.e. $t \in [0, T]$, it follows from Proposition 27.7 (c) in [59] that $B(t, \cdot) + \mathcal{A}_{\lambda}(t, \cdot)$ is pseudo-monotone for a.e. $t \in [0, T]$. According to Lemma 6.1, the Yosida approximation \mathcal{A}_{λ} is coercive. From this, using (\mathbf{H}_{B}^{1}) and (\mathbf{H}_{σ}) , we deduce that for any $v \in V$ and a.e. $t \in [0, T]$,

$$2\langle A(t,v),v\rangle + \|\sigma(t,v)\|_{\mathcal{L}_{2}}^{2} \leqslant Cf(t)(1+\|v\|_{H}^{2}) - \delta 2^{-\alpha+1}\|v\|_{V}^{\alpha}.$$
(3.3)

Thus, Hypothesis (H3) in [47] is satisfied. Since $\mathcal{D}(\mathcal{A}) = V$ and $\|\mathcal{A}_{\lambda}(\cdot, x)\|_{V^*} \leq \|\mathcal{A}^0(\cdot, x)\|_{V^*}$ for all $x \in \mathcal{D}(\mathcal{A})$, the conditions $(\mathbf{H}^3_{\mathcal{A}})$ and (\mathbf{H}^2_B) ensure that for any $v \in V$ and $\alpha > 1$, there exist constants $\beta \geq 0$ and C > 0 such that

$$\|A(t,v)\|_{V^*}^{\frac{\alpha}{\alpha-1}} \leqslant \left(f(t) + C\|v\|_V^{\alpha}\right) \left(1 + \|v\|_H^{\beta}\right).$$
(3.4)

Therefore, Hypothesis (H4) in [47] is also satisfied. This completes the proof.

Remark 3.7. Since $B(t, \cdot) + \mathcal{A}_{\lambda}(t, \cdot)$ is pseudo-monotone for a.e. $t \in [0, T]$ and locally bounded, we find that $B(t, \cdot) + \mathcal{A}_{\lambda}(t, \cdot)$ is demicontinuous (hence, hemicontinuous) for a.e. $t \in [0, T]$ by Proposition 27.7 (b) in [59], which implies Hypothesis (H1) in [47] directly.

Lemma 3.3. There exists a constant $C_T > 0$ such that for any $0 < \lambda < \delta^{-1}$ we have

$$\mathbb{E}\Big[\sup_{t\in[0,T]} \|X_{\lambda}(t)\|_{H}^{p}\Big] + \mathbb{E}\int_{0}^{T} \|X_{\lambda}(s)\|_{V}^{\alpha}\|X_{\lambda}(s)\|_{H}^{p-2}ds \leq C_{T}(1+\|x\|_{H}^{p}).$$

Proof. By Itô's formula we have

$$\begin{split} \|X_{\lambda}(t)\|_{H}^{p} &= \|x\|_{H}^{p} - \frac{p}{2} \int_{0}^{t} \|X_{\lambda}(s)\|_{H}^{p-2} \left[2 \left\langle \mathcal{A}_{\lambda}(s, X_{\lambda}(s)) + B(s, X_{\lambda}(s)), X_{\lambda}(s) \right\rangle \right. \\ &+ \|\sigma(s, X_{\lambda}(s))\|_{\mathcal{L}_{2}}^{2} \right] ds + \frac{p(p-2)}{2} \int_{0}^{t} \|X_{\lambda}(s)\|_{H}^{p-4} \|\sigma(s, X_{\lambda}(s))^{*} X_{\lambda}(s)\|_{U}^{2} ds \\ &+ p \int_{0}^{t} \|X_{\lambda}(s)\|_{H}^{p-2} \left\langle X_{\lambda}(s), \sigma(s, X_{\lambda}(s)) dW(s) \right\rangle_{H}. \end{split}$$

Then by (3.3) and let $c = \delta 2^{-\alpha+1} > 0$, it follows that

$$\|X_{\lambda}(t)\|_{H}^{p} + \frac{pc}{2} \int_{0}^{t} \|X_{\lambda}(s)\|_{V}^{\alpha} \|X_{\lambda}(s)\|_{H}^{p-2} ds$$

$$\leq \|x\|_{H}^{p} + C \int_{0}^{t} f(s)(1 + \|X_{\lambda}(s)\|_{H}^{p}) ds$$

$$+ p \int_{0}^{t} \|X_{\lambda}(s)\|_{H}^{p-2} \langle X_{\lambda}(s), \sigma(s, X_{\lambda}(s)) dW(s) \rangle_{H}.$$
(3.5)

 Set

$$\tau_{\lambda}^{M} := \inf \left\{ t \ge 0 : \|X_{\lambda}(t)\|_{H} > M \right\} \wedge T.$$

We have $\tau_{\lambda}^{M} \to T, \mathbb{P}$ -a.s., as $M \to \infty$, for each $\lambda \in (0, \delta^{-1})$. Taking the supremum over $t \in [0, T \land \tau_{\lambda}^{M}]$ and then taking expectation on both sides of (3.5), we get

$$\mathbb{E}\Big[\sup_{t\in[0,T\wedge\tau_{\lambda}^{M}]}\|X_{\lambda}(t)\|_{H}^{p}\Big] + \frac{pc}{2}\mathbb{E}\int_{0}^{T\wedge\tau_{\lambda}^{M}}\|X_{\lambda}(s)\|_{V}^{\alpha}\|X_{\lambda}(s)\|_{H}^{p-2}ds$$

$$\leqslant \|x\|_{H}^{p} + C\int_{0}^{T}f(s)ds + C\mathbb{E}\int_{0}^{T\wedge\tau_{\lambda}^{M}}f(s)\|X_{\lambda}(s)\|_{H}^{p}ds$$

$$+ p\mathbb{E}\left[\sup_{t\in[0,T\wedge\tau_{\lambda}^{M}]}\left|\int_{0}^{t}\|X_{\lambda}(s)\|_{H}^{p-2}\langle X_{\lambda}(s),\sigma(s,X_{\lambda}(s))dW(s)\rangle_{H}\right|\Big].$$
(3.6)

By B-D-G's inequality, Young's inequality and (\mathbf{H}_{σ}) , we have

$$p\mathbb{E}\left[\sup_{t\in[0,T\wedge\tau_{\lambda}^{M}]}\left|\int_{0}^{t}\|X_{\lambda}(s)\|_{H}^{p-2}\langle X_{\lambda}(s),\sigma(s,X_{\lambda}(s))dW(s)\rangle_{H}\right|\right]$$

$$\leq C\mathbb{E}\left[\int_{0}^{T\wedge\tau_{\lambda}^{M}}\|X_{\lambda}(s)\|_{H}^{2p-2}\|\sigma(s,X_{\lambda}(s))\|_{\mathcal{L}_{2}}^{2}ds\right]^{\frac{1}{2}}$$

$$\leq C\mathbb{E}\left[\sup_{s\in[0,T\wedge\tau_{\lambda}^{M}]}\|X_{\lambda}(s)\|_{H}^{p}\cdot\int_{0}^{T\wedge\tau_{\lambda}^{M}}\|X_{\lambda}(s)\|_{H}^{p-2}\|\sigma(s,X_{\lambda}(s))\|_{\mathcal{L}_{2}}^{2}ds\right]^{\frac{1}{2}}$$

$$\leq \frac{1}{2}\mathbb{E}\left[\sup_{s\in[0,T\wedge\tau_{\lambda}^{M}]}\|X_{\lambda}(s)\|_{H}^{p}\right]+C\mathbb{E}\int_{0}^{T\wedge\tau_{\lambda}^{M}}\|X_{\lambda}(s)\|_{H}^{p-2}\|\sigma(s,X_{\lambda}(s))\|_{\mathcal{L}_{2}}^{2}ds$$

$$\leq \frac{1}{2}\mathbb{E}\left[\sup_{s\in[0,T\wedge\tau_{\lambda}^{M}]}\|X_{\lambda}(s)\|_{H}^{p}\right]+C\int_{0}^{T}f(s)ds+C\mathbb{E}\int_{0}^{T\wedge\tau_{\lambda}^{M}}f(s)\|X_{\lambda}(s)\|_{H}^{p}ds.$$
(3.7)

Combining inequalities (3.6)-(3.7) and applying Gronwall's inequality, we obtain

$$\mathbb{E}\Big[\sup_{t\in[0,T\wedge\tau_{\lambda}^{M}]}\|X_{\lambda}(t)\|_{H}^{p}\Big] + C\mathbb{E}\int_{0}^{T\wedge\tau_{\lambda}^{M}}\|X_{\lambda}(s)\|_{V}^{\alpha}\|X_{\lambda}(s)\|_{H}^{p-2}ds$$
$$\leqslant C\left(\|x\|_{H}^{p} + \int_{0}^{T}f(s)ds\right)\exp\Big\{C\int_{0}^{T}f(s)ds\Big\}.$$

Letting $M \to \infty$ and applying Fatou's lemma, due to $f \in L^1([0,T], \mathbb{R}_+)$, it follows that Lemma 3.3 holds.

Lemma 3.4. For any $\lambda \in (0, \delta^{-1})$, the following inequality holds

$$\mathbb{E}\int_0^T \left(\|\mathcal{A}_{\lambda}(s, X_{\lambda}(s))\|_{V^*}^{\frac{\alpha}{\alpha-1}} + \|B(s, X_{\lambda}(s))\|_{V^*}^{\frac{\alpha}{\alpha-1}} + \|\sigma(s, X_{\lambda}(s))\|_{\mathcal{L}_2}^2 \right) ds$$

$$\leq C_T (1 + \|x\|_H^{\beta+2}).$$

Proof. Recall the fact that the operator \mathcal{A} has the full domain (i.e. $\mathcal{D}(\mathcal{A}) = V$). By $(\mathbf{H}_{\mathcal{A}}^3)$, $(\mathbf{H}_{\mathcal{B}}^2)$, (\mathbf{H}_{σ}) and Proposition 2.3 (ii), we have

$$\mathbb{E}\int_{0}^{T} \left(\left\| \mathcal{A}_{\lambda}(s, X_{\lambda}(s)) \right\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} + \left\| B(s, X_{\lambda}(s)) \right\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} + \left\| \sigma(s, X_{\lambda}(s)) \right\|_{\mathcal{L}_{2}}^{2} \right) ds$$

$$\leq \mathbb{E}\int_{0}^{T} \left((f(s) + C \| X_{\lambda}(s) \|_{V}^{\alpha}) (1 + \| X_{\lambda}(s) \|_{H}^{\beta}) + f(s) (1 + \| X_{\lambda}(s) \|_{H}^{2}) \right) ds.$$
(3.8)

Combining Lemma 3.3 with (3.8), we complete the proof.

3.3. **Pre-compactness.** In this part, we take a subsequence $\{\lambda_k\}_{k=1}^{\infty}$ of $\{\lambda\}_{\lambda \in (0,\delta^{-1})}$, where $\lambda_k \to 0$ as $k \to \infty$. Along this subsequence, we denote X_{λ_k} by X_k . Define the following stopping times

$$\tau_k^M := \inf\left\{t \ge 0 : \|X_k(t)\|_H^2 > M\right\} \wedge \inf\left\{t \ge 0 : \int_0^t \|X_k(s)\|_V^\alpha ds > M\right\} \wedge T,$$

with the convention $\inf \emptyset = \infty$. By Chebyshev's inequality, Lemma 3.3 implies that

$$\lim_{M \to \infty} \sup_{k \in \mathbb{N}} \mathbb{P}(\tau_k^M < T) = 0.$$
(3.9)

We present the tightness of the laws of $\{X_k\}_{k=1}^{\infty}$.

Lemma 3.5. The sequence $\{X_k\}_{k=1}^{\infty}$ is tight in the space $C([0,T], V^*) \cap L^{\alpha}([0,T], H)$.

Proof. In order to get the lemma, we will prove that $\{X_k\}_{k=1}^{\infty}$ is tight in $C([0,T], V^*)$ and in $L^{\alpha}([0,T], H)$ separately. Then for any sequence $\{\mathcal{L}_{X_k}\}_{k=1}^{\infty}$ we can find a subsequence, still denoted by $\{\mathcal{L}_{X_k}\}_{k=1}^{\infty}$, such that $\{\mathcal{L}_{X_k}\}_{k=1}^{\infty}$ converges weakly in $C([0,T], V^*) \cap L^{\alpha}([0,T], H)$.

Step 1. In this step, we prove that $\{X_k\}_{k=1}^{\infty}$ is tight in $C([0,T], V^*)$. Notice that H is compactly embedded into V^* , it is sufficient to show that for every $e \in P_m H$, $m \in \mathbb{N}$, $\{\langle X_k, e \rangle\}_{k=1}^{\infty}$ is tight in the space $C([0,T], \mathbb{R})$ by Theorem 3.1 in [25]. In terms of the Aldou's tightness criterion, the result will be proved if we can show that for any stopping time $0 \leq \zeta_k \leq T$ and for any $\varepsilon > 0$,

$$\lim_{\delta \to 0} \sup_{k \in \mathbb{N}} \mathbb{P}\left(\left| \left\langle X_k(\zeta_k + \delta) - X_k(\zeta_k), e \right\rangle \right| > \varepsilon \right) = 0,$$
(3.10)

where $\zeta_k + \delta := T \wedge (\zeta_k + \delta) \vee 0$. Set $X_k^M(t) := X_k(t \wedge \tau_k^M)$. By Chebyshev's inequality, it implies that

$$\mathbb{P}\left(\left|\left\langle X_{k}(\zeta_{k}+\delta)-X_{k}(\zeta_{k}),e\right\rangle\right|>\varepsilon\right)$$

$$\leq \mathbb{P}\left(\left|\left\langle X_{k}(\zeta_{k}+\delta)-X_{k}(\zeta_{k}),e\right\rangle\right|>\varepsilon,\tau_{k}^{M}\geqslant T\right)+\mathbb{P}(\tau_{k}^{M}
$$\leq \frac{1}{\varepsilon^{\alpha}}\mathbb{E}\left|\left\langle X_{k}^{M}(\zeta_{k}+\delta)-X_{k}^{M}(\zeta_{k}),e\right\rangle\right|^{\alpha}+\mathbb{P}(\tau_{k}^{M}
(3.11)$$$$

Using (3.2) and B-D-G's inequality, we have

$$\mathbb{E} \left| \left\langle X_k^M(\zeta_k + \delta) - X_k^M(\zeta_k), e \right\rangle \right|^{\alpha} \\ \leqslant C \mathbb{E} \left(\int_{\zeta_k \wedge \tau_k^M}^{(\zeta_k + \delta) \wedge \tau_k^M} \left| \left\langle \mathcal{A}_{\lambda_k}(s, X_k(s)) + B(s, X_k(s)), e \right\rangle \right| ds \right)^{\alpha}$$

YOSIDA APPROXIMATION AND MULTI-VALUED SEI

$$+ C\mathbb{E}\left(\int_{\zeta_k \wedge \tau_k^M}^{(\zeta_k + \delta) \wedge \tau_k^M} \|e\|_H^2 \|\sigma(s, X_k(s))\|_{\mathcal{L}_2}^2 ds\right)^{\frac{\alpha}{2}}$$
$$=: I_k + II_k.$$
(3.12)

According to Hölder's inequality, $(\mathbf{H}_{\mathcal{A}}^3)$, (\mathbf{H}_{B}^2) and Proposition 2.3 (ii), we deduce that

$$I_{k} \leq C|\delta|\mathbb{E}\left(\int_{\zeta_{k}\wedge\tau_{k}^{M}}^{(\zeta_{k}+\delta)\wedge\tau_{k}^{M}}|\langle\mathcal{A}_{\lambda_{k}}(s,X_{k}(s))+B(s,X_{k}(s)),e\rangle|^{\frac{\alpha}{\alpha-1}}ds\right)^{\alpha-1}$$
$$\leq C|\delta|\mathbb{E}\left(\int_{0}^{T\wedge\tau_{k}^{M}}\|e\|_{V}^{\frac{\alpha}{\alpha-1}}(f(s)+C\|X_{k}(s)\|_{V}^{\alpha})(1+\|X_{k}(s)\|_{H}^{\beta})ds\right)^{\alpha-1}$$
$$\leq C_{M}|\delta|.$$

Thus, we have

$$\lim_{\delta \to 0} \sup_{k \in \mathbb{N}} I_k = 0. \tag{3.13}$$

Similarly, by (\mathbf{H}_{σ}) it follows that

$$II_{k} \leqslant C\mathbb{E} \left(\int_{\zeta_{k} \wedge \tau_{k}^{M}}^{(\zeta_{k}+\delta) \wedge \tau_{k}^{M}} \|e\|_{H}^{2} f(s)(1+\|X_{k}(s)\|_{H}^{2}) ds \right)^{\frac{\alpha}{2}}$$
$$\leqslant C_{M} \left(\int_{\zeta_{k} \wedge \tau_{k}^{M}}^{(\zeta_{k}+\delta) \wedge \tau_{k}^{M}} f(s) ds \right)^{\frac{\alpha}{2}}.$$

Since $f \in L^1([0,T], \mathbb{R}_+)$, by the absolute continuity of the Lebesgue integral, we have

$$\lim_{\delta \to 0} \sup_{k \in \mathbb{N}} II_k = 0. \tag{3.14}$$

Combining (3.12)-(3.14) yields

$$\lim_{\delta \to 0} \sup_{k \in \mathbb{N}} \mathbb{E} \left| \left\langle X_k^M(\zeta_k + \delta) - X_k^M(\zeta_k), e \right\rangle \right|^{\alpha} = 0.$$
(3.15)

According to (3.9) and (3.15), letting $\delta \to 0$ and then letting $M \to \infty$ in (3.11) yield (3.10).

Step 2. In this step, we show that $\{X_k\}_{k=1}^{\infty}$ is tight in $L^{\alpha}([0,T],H)$. By Lemma 5.2 in [47] and

$$\sup_{k\in\mathbb{N}}\mathbb{E}\int_0^T \|X_k(t)\|_V^\alpha dt < \infty,$$

it suffices to prove that for any $\varepsilon > 0$

$$\lim_{\delta \to 0^+} \sup_{k \in \mathbb{N}} \mathbb{P}\left(\int_0^{T-\delta} \|X_k(t+\delta) - X_k(t)\|_H^\alpha dt > \varepsilon\right) = 0.$$
(3.16)

Let $X_k^M(t) := X_k(t \wedge \tau_k^M)$. Then we have

$$\mathbb{P}\left(\int_0^{T-\delta} \|X_k(t+\delta) - X_k(t)\|_H^\alpha dt > \varepsilon\right)$$

$$\leq \mathbb{P}\left(\int_{0}^{T-\delta} \|X_{k}(t+\delta) - X_{k}(t)\|_{H}^{\alpha} dt > \varepsilon, \tau_{k}^{M} \geq T\right) + \mathbb{P}\left(\tau_{k}^{M} < T\right)$$

$$\leq \frac{1}{\varepsilon} \mathbb{E}\int_{0}^{T-\delta} \|X_{k}^{M}(t+\delta) - X_{k}^{M}(t)\|_{H}^{\alpha} dt + \mathbb{P}(\tau_{k}^{M} < T).$$

$$(3.17)$$

Once we can prove that for any fixed M > 0,

$$\lim_{\delta \to 0^+} \sup_{k \in \mathbb{N}} \mathbb{E} \int_0^{T-\delta} \|X_k^M(t+\delta) - X_k^M(t)\|_H^\alpha dt = 0,$$
(3.18)

letting $\delta \to 0$ and $M \to \infty$ in (3.17), then we deduce that (3.16) holds by (3.9). Next, we prove (3.18).

Firstly, we consider the case $\alpha \in (1, 2]$. By Itô's formula, we have

$$\mathbb{E} \|X_k^M(t+\delta) - X_k^M(t)\|_H^2$$

= $\mathbb{E} \int_{t\wedge\tau_k^M}^{(t+\delta)\wedge\tau_k^M} \left[-2\langle \mathcal{A}_{\lambda_k}(r, X_k(r)) + B(r, X_k(r)), X_k(r) - X_k(t\wedge\tau_k^M)\rangle\right] dr$
+ $\mathbb{E} \int_{t\wedge\tau_k^M}^{(t+\delta)\wedge\tau_k^M} \|\sigma(r, X_k(r))\|_{\mathcal{L}_2}^2 dr.$

Thus, it implies that

$$\mathbb{E} \int_{0}^{T-\delta} \|X_{k}^{M}(t+\delta) - X_{k}^{M}(t)\|_{H}^{2} dt$$

$$= \mathbb{E} \int_{0}^{T-\delta} dt \int_{t\wedge\tau_{k}^{M}}^{(t+\delta)\wedge\tau_{k}^{M}} \left[-2 \left\langle \mathcal{A}_{\lambda_{k}}(r, X_{k}(r)) + B(r, X_{k}(r)), X_{k}(r) \right\rangle + \|\sigma(r, X_{k}(r))\|_{\mathcal{L}_{2}}^{2} \right] dr$$

$$+ 2\mathbb{E} \int_{0}^{T-\delta} dt \int_{t\wedge\tau_{k}^{M}}^{(t+\delta)\wedge\tau_{k}^{M}} \left\langle \mathcal{A}_{\lambda_{k}}(r, X_{k}(r)) + B(r, X_{k}(r)), X_{k}(t\wedge\tau_{k}^{M}) \right\rangle dr$$

$$=: III_{k} + IV_{k}. \qquad (3.19)$$

By Fubini's theorem and (3.3), it follows that

$$III_{k} = \mathbb{E} \int_{0}^{T \wedge \tau_{k}^{M}} \left[-2 \left\langle \mathcal{A}_{\lambda_{k}}(r, X_{k}(r)) + B(r, X_{k}(r)), X_{k}(r) \right\rangle + \|\sigma(r, X_{k}(r))\|_{\mathcal{L}_{2}}^{2} \right] dr$$

$$\cdot \int_{0 \vee (r-\delta)}^{r} \mathbf{1}_{\{\tau_{k}^{M} > t\}} dt$$

$$\leq C \delta \mathbb{E} \int_{0}^{T \wedge \tau_{k}^{M}} f(s) \left(1 + \|X_{k}(s)\|_{H}^{2}\right) ds$$

$$\leq C_{M} \delta.$$
(3.20)

Similarly, by (3.4) we can deduce that

$$|IV_k| = 2 \left| \mathbb{E} \int_0^{T \wedge \tau_k^M} dr \int_{0 \lor (r-\delta)}^r \mathbf{1}_{\{\tau_k^M > t\}} \left\langle \mathcal{A}_{\lambda_k}(r, X_k(r)) + B(r, X_k(r)), X_k(t \land \tau_k^M) \right\rangle dt \right|$$

$$\leq C \mathbb{E} \int_{0}^{T \wedge \tau_{k}^{M}} \|\mathcal{A}_{\lambda_{k}}(r, X_{k}(r)) + B(r, X_{k}(r))\|_{V^{*}} dr \int_{0 \vee (r-\delta)}^{r} \|X_{k}(t \wedge \tau_{k}^{M})\|_{V} dt$$

$$\leq C \delta^{\frac{\alpha-1}{\alpha}} \mathbb{E} \int_{0}^{T \wedge \tau_{k}^{M}} \|\mathcal{A}_{\lambda_{k}}(r, X_{k}(r)) + B(r, X_{k}(r))\|_{V^{*}} dr \left(\int_{0}^{T \wedge \tau_{k}^{M}} \|X_{k}(t)\|_{V}^{\alpha} dt\right)^{\frac{1}{\alpha}}$$

$$\leq C_{M} \delta^{\frac{\alpha-1}{\alpha}}.$$

$$(3.21)$$

Combining (3.19)-(3.21) yields

$$\sup_{k\in\mathbb{N}}\mathbb{E}\int_0^{T-\delta} \|X_k^M(t+\delta) - X_k^M(t)\|_H^2 dt \leqslant C_M(\delta+\delta^{\frac{\alpha-1}{\alpha}}).$$

Since $\alpha \in (1, 2]$, we can apply Hölder's inequality to obtain

$$\lim_{\delta \to 0^+} \sup_{k \in \mathbb{N}} \mathbb{E} \int_0^{T-\delta} \|X_k^M(t+\delta) - X_k^M(t)\|_H^\alpha ds$$
$$\leqslant C \left(\lim_{\delta \to 0^+} \sup_{k \in \mathbb{N}} \mathbb{E} \int_0^{T-\delta} \|X_k^M(t+\delta) - X_k^M(t)\|_H^2 ds\right)^{\frac{\alpha}{2}} = 0.$$

We have completed the proof of (3.18) for $\alpha \in (1, 2]$.

Now we consider the case $\alpha > 2$. Note that

$$\begin{split} & \mathbb{E} \|X_k^M(t+\delta) - X_k^M(t)\|_H^{\alpha} \\ &= \frac{\alpha}{2} \mathbb{E} \int_{t\wedge\tau_k^M}^{(t+\delta)\wedge\tau_k^M} \|X_k(r) - X_k(t\wedge\tau_k^M)\|_H^{\alpha-2} \\ & \cdot \left[-2\left\langle \mathcal{A}_{\lambda_k}(r, X_k(r)) + B(r, X_k(r)), X_k(r) - X_k(t\wedge\tau_k^M) \right\rangle + \|\sigma(r, X_k(r))\|_{\mathcal{L}_2}^2 \right] dr \\ & + \frac{\alpha(\alpha-2)}{2} \mathbb{E} \int_{t\wedge\tau_k^M}^{(t+\delta)\wedge\tau_k^M} \|X_k(r) - X_k(t\wedge\tau_k^M)\|_H^{\alpha-4} \\ & \cdot \|(\sigma(r, X_k(r)))^*(X_k(r) - X_k(t\wedge\tau_k^M))\|_U^2 dr. \end{split}$$

Using Fubini's theorem, we have

$$\begin{split} & \mathbb{E} \int_{0}^{T-\delta} \|X_{k}^{M}(t+\delta) - X_{k}^{M}(t)\|_{H}^{\alpha} dt \\ &= \frac{\alpha}{2} \mathbb{E} \int_{0}^{T-\delta} dt \int_{t\wedge\tau_{k}^{M}}^{(t+\delta)\wedge\tau_{k}^{M}} \|X_{k}(r) - X_{k}(t\wedge\tau_{k}^{M})\|_{H}^{\alpha-2} \\ & \cdot \left[-2 \left\langle \mathcal{A}_{\lambda_{k}}(r, X_{k}(r)) + B(r, X_{k}(r)), X_{k}(r) \right\rangle + \|\sigma(r, X_{k}(r))\|_{\mathcal{L}_{2}}^{2}\right] dr \\ & + \alpha \mathbb{E} \int_{0}^{T-\delta} dt \int_{t\wedge\tau_{k}^{M}}^{(t+\delta)\wedge\tau_{k}^{M}} \|X_{k}(r) - X_{k}(t\wedge\tau_{k}^{M})\|_{H}^{\alpha-2} \\ & \cdot \left\langle \mathcal{A}_{\lambda_{k}}(r, X_{k}(r)) + B(r, X_{k}(r)), X_{k}(t\wedge\tau_{k}^{M}) \right\rangle dr \\ & + \frac{\alpha(\alpha-2)}{2} \mathbb{E} \int_{0}^{T-\delta} dt \int_{t\wedge\tau_{k}^{M}}^{(t+\delta)\wedge\tau_{k}^{M}} \|X_{k}(r) - X_{k}(t\wedge\tau_{k}^{M})\|_{H}^{\alpha-4} \end{split}$$

YOSIDA APPROXIMATION AND MULTI-VALUED SEI

$$\cdot \|(\sigma(r, X_k(r)))^* (X_k(r) - X_k(t \wedge \tau_k^M))\|_U^2 dr$$

=: $I_k + II_k + III_k.$ (3.22)

Applying the similar arguments as in (3.20) and (3.21), it implies that

$$I_k \leqslant C_M \delta, \tag{3.23}$$

$$|II_k| \leqslant C_M \delta^{\frac{\alpha-1}{\alpha}}.$$
(3.24)

On the other hand, by (\mathbf{H}_{σ}) and Fubini's theorem it follows that

$$III_{k} \leq C_{M} \mathbb{E} \int_{0}^{T \wedge \tau_{k}^{M}} f(r) \|X_{k}(r) - X_{k}(t \wedge \tau_{k}^{M})\|_{H}^{\alpha - 2} \left(1 + \|X_{k}(r)\|_{H}^{2}\right) dr$$

$$\cdot \int_{0 \vee (r - \delta)}^{r} \mathbf{1}_{\{\tau_{k}^{M} > t\}} dt$$

$$\leq C_{M} \delta.$$
(3.25)

Combining (3.22)-(3.25), we obtain

$$\sup_{k\in\mathbb{N}}\mathbb{E}\int_0^{T-\delta} \|X_k^M(t+\delta) - X_k^M(t)\|_H^\alpha dt \leqslant C_M(\delta+\delta^{\frac{\alpha-1}{\alpha}}).$$

Therefore, (3.18) is proved. We complete the proof of the lemma.

3.4. Proof of Theorem 3.1. Let

$$\Upsilon := [L^{\alpha}([0,T],H) \cap C([0,T],V^*)] \times C([0,T],U_1),$$

where U_1 is a Hilbert space and the embedding $U \subset U_1$ is Hilbert-Schmidt. According to Lemma 3.5, we note that the family of the laws $\mathcal{L}(X_k, W)$ is tight in Υ . By the Prohorov theorem and the generalized version of Skorohod representation theorem (cf. Theorem C.1 in [17]), there exists a new probability space $(\widetilde{\Omega}, \widetilde{\mathcal{F}}, \widetilde{\mathbb{P}})$, along with a sequence of Υ -valued random vectors $\{(\widetilde{X}_k, \widetilde{W}_k)\}$ and $(\widetilde{X}, \widetilde{W})$, such that

(i)
$$\widetilde{W}_{k} = \widetilde{W}$$
 for any $k \in \mathbb{N}$, $\widetilde{\mathbb{P}}$ -a.s.;
(ii) $\mathcal{L}(\widetilde{X}_{k}, \widetilde{W}_{k}) = \mathcal{L}(X_{k}, W);$ (3.26)

(iii) $\widetilde{\mathbb{P}}$ -a.s.,

$$\|\widetilde{X}_{k} - \widetilde{X}\|_{L^{\alpha}([0,T],H)} + \|\widetilde{X}_{k} - \widetilde{X}\|_{C([0,T],V^{*})} \to 0.$$
(3.27)

Let $\widetilde{\mathcal{F}}_t$ be the usual filtration generated by $\{\widetilde{X}_k(s), \widetilde{X}(s), \widetilde{W}(s) : s \leq t\}$. It is clear that \widetilde{W} is an $\{\widetilde{\mathcal{F}}_t\}$ -cylindrical Wiener process on U and

$$\widetilde{X}_{k}(t) = x - \int_{0}^{t} \mathcal{A}_{\lambda_{k}}(s, \widetilde{X}_{k}(s)) ds - \int_{0}^{t} B(s, \widetilde{X}_{k}(s)) ds + \int_{0}^{t} \sigma(s, \widetilde{X}_{k}(s)) d\widetilde{W}(s), \quad t \in [0, T].$$
(3.28)

20

In view of the Claim (ii) above and Lemma 3.2, we know that $\{\tilde{X}_k\}$ satisfies the same moment estimates, i.e., for any $p \ge 2$,

$$\sup_{k\in\mathbb{N}}\left\{\widetilde{\mathbb{E}}\left[\sup_{t\leqslant T}\|\widetilde{X}_{k}(t)\|_{H}^{p}\right]+\widetilde{\mathbb{E}}\left(\int_{0}^{T}\|\widetilde{X}_{k}(t)\|_{V}^{\alpha}dt\right)^{\frac{p}{2}}\right\}<\infty.$$
(3.29)

Due to the fact that $\|\cdot\|_H$ and $\|\cdot\|_V$ are lower semicontinuous in V^* , by (3.27) and Fatou's lemma, it follows that

$$\widetilde{\mathbb{E}}\left[\sup_{t\in[0,T]}\|\widetilde{X}(t)\|_{H}^{p}\right] \leqslant \widetilde{\mathbb{E}}\left[\sup_{t\in[0,T]}\liminf_{k\to\infty}\|\widetilde{X}_{k}(t)\|_{H}^{p}\right] \\
\leqslant \widetilde{\mathbb{E}}\left[\liminf_{k\to\infty}\sup_{t\in[0,T]}\|\widetilde{X}_{k}(t)\|_{H}^{p}\right] \\
\leqslant \liminf_{k\to\infty}\widetilde{\mathbb{E}}\left[\sup_{t\in[0,T]}\|\widetilde{X}_{k}(t)\|_{H}^{p}\right] < \infty.$$
(3.30)

Similarly, we have

$$\mathbb{E}\left(\int_{0}^{T} \|\widetilde{X}(s)\|_{V}^{\alpha} ds\right)^{\frac{p}{2}} < \infty.$$
(3.31)

Then, by Lemma 3.4 we can establish the following estimates.

Lemma 3.6. The following estimates hold

$$\sup_{k\in\mathbb{N}} \mathbb{E} \int_{0}^{T} \|\mathcal{A}_{\lambda_{k}}(t,\widetilde{X}_{k}(t))\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} dt < \infty;$$

$$\sup_{k\in\mathbb{N}} \mathbb{E} \int_{0}^{T} \|B(t,\widetilde{X}_{k}(t))\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} dt < \infty;$$

$$\sup_{k\in\mathbb{N}} \mathbb{E} \int_{0}^{T} \|\sigma(t,\widetilde{X}_{k}(t))\|_{\mathcal{L}_{2}}^{2} dt < \infty.$$
(3.32)

In the sequel, we aim to prove that $(\widetilde{X}, \widetilde{W})$ is a solution to equation (1.3). Since the spaces $L^{\alpha}(\widetilde{\Omega} \times [0,T], V), L^{\frac{\alpha}{\alpha-1}}(\widetilde{\Omega} \times [0,T], V^*)$ and $L^2(\widetilde{\Omega} \times [0,T], L_2(U,H))$ are reflexive, by the estimate (3.29), Lemma 3.6 and the Banach-Alaoglu's theorem, it implies that there exist $\widehat{X} \in L^{\alpha}(\widetilde{\Omega} \times [0,T], V), \ \widetilde{\eta} \in L^{\frac{\alpha}{\alpha-1}}(\widetilde{\Omega} \times [0,T], V^*), \ \widetilde{\mathcal{B}} \in L^{\frac{\alpha}{\alpha-1}}(\widetilde{\Omega} \times [0,T], V^*)$ and $\widetilde{\mathcal{D}} \in L^2(\widetilde{\Omega} \times [0,T], L_2(U,H))$ such that

$$\widetilde{X}_k \to \widehat{X} \quad \text{in } L^{\alpha}(\widetilde{\Omega} \times [0,T],V);$$

$$(3.33)$$

$$\mathcal{A}_{\lambda_k}(\cdot, \widetilde{X}_k(\cdot)) \rightharpoonup \widetilde{\eta} \quad \text{in } L^{\frac{\alpha}{\alpha-1}}(\widetilde{\Omega} \times [0, T], V^*); \tag{3.34}$$

$$B(\cdot, \widetilde{X}_k(\cdot)) \rightharpoonup \widetilde{\mathcal{B}} \quad \text{in } L^{\frac{\alpha}{\alpha-1}}(\widetilde{\Omega} \times [0, T], V^*);$$

$$(3.35)$$

$$\sigma(\cdot, \widetilde{X}_k(\cdot)) \rightharpoonup \widetilde{\mathcal{D}} \quad \text{in } L^2(\widetilde{\Omega} \times [0, T], L_2(U, H));$$
(3.36)

$$\int_{0}^{\cdot} \sigma(s, \widetilde{X}_{k}(s)) d\widetilde{W}(s) \rightharpoonup \int_{0}^{\cdot} \widetilde{\mathcal{D}}(s) d\widetilde{W}(s) \text{ in } L^{\infty}([0, T], L^{2}(\widetilde{\Omega}, H)).$$
(3.37)

$$\bar{X}(t) := x - \int_0^t \tilde{\eta}(s)ds - \int_0^t \tilde{\mathcal{B}}(s)ds + \int_0^t \tilde{\mathcal{D}}(s)d\widetilde{W}(s).$$
(3.38)

Then, it is clear that

$$\widetilde{X} = \widehat{X} = \overline{X}, \quad dt \otimes \widetilde{\mathbb{P}}$$
-a.e.. (3.39)

Indeed, the left equality in (3.39) follows from the uniqueness of the limits and the equality on the right follows from e.g. page 58 in [54]. Moreover, applying Theorem 4.2.5 in [42], it follows that \overline{X} is an *H*-valued continuous process. According to (3.27) and (3.30), \widetilde{X} is *H*-valued and weakly continuous in *H*. Hence, \widetilde{X} and \overline{X} are indistinguishable.

Throughout the remaining part of this subsection, we work on the new filtered probability space $(\widetilde{\Omega}, \widetilde{\mathcal{F}}, \{\widetilde{\mathcal{F}}_t\}_{t \ge 0}, \widetilde{\mathbb{P}})$, however, we drop all the superscripts for sake of simplicity. The following lemma characterizes the limit of $\sigma(\cdot, X_k(\cdot))$.

Lemma 3.7. $\mathcal{D}(\cdot) = \sigma(\cdot, X(\cdot)), dt \otimes \mathbb{P}$ -a.e..

Proof. Since $||X_k - X||_{L^{\alpha}([0,T],H)} \to 0$, \mathbb{P} -a.s., by (3.29), (3.31) and Hölder's inequality, we can use the dominated convergence theorem to obtain

$$\lim_{k \to \infty} \mathbb{E} \int_0^T \|X_k(t) - X(t)\|_H^\kappa = 0 \text{ for all } \kappa \in [1, \alpha).$$

Then there exists a subsequence still denoted by $\{X_k\}$ such that for a.e. (t, ω)

$$\lim_{k \to \infty} \|X_k(t) - X(t)\|_H = 0.$$
(3.40)

Thus by (\mathbf{H}_{σ}) , (3.29) and (3.30), we can obtain

$$\lim_{k \to \infty} \mathbb{E} \int_0^T \|\sigma(t, X_k(t)) - \sigma(t, X(t))\|_{\mathcal{L}_2}^2 dt = 0.$$
(3.41)

Combining (3.36) and the uniqueness of limits implies that $\mathcal{D}(\cdot) = \sigma(\cdot, X(\cdot))$.

The following lemma characterizes the limit of $B(\cdot, X_k(\cdot))$. To this end, we recall that X_k and X satisfy the equation (3.28), (3.38) respectively.

Lemma 3.8. If we have

$$X_k \rightharpoonup X \text{ in } L^{\alpha}([0,T] \times \Omega, V);$$

$$\mathcal{A}_{\lambda_k}(\cdot, X_k(\cdot)) \rightharpoonup \eta \text{ in } L^{\frac{\alpha}{\alpha-1}}([0,T] \times \Omega, V^*);$$
 (3.42)

$$B(\cdot, X_k(\cdot)) \rightharpoonup \mathcal{B} \text{ in } L^{\frac{\alpha}{\alpha-1}}([0,T] \times \Omega, V^*);$$

$$(3.43)$$

$$\limsup_{k \to \infty} \mathbb{E} \int_0^T \langle \mathcal{A}_{\lambda_k}(t, X_k(t)) + B(t, X_k(t)), X_k(t) \rangle dt \leqslant \mathbb{E} \int_0^T \langle \eta(t) + \mathcal{B}(t), X(t) \rangle dt, \quad (3.44)$$

then $\mathcal{B}(\cdot) = B(\cdot, X(\cdot)), dt \otimes \mathbb{P}$ -a.e..

Proof. According to the proof of Lemma 3.2 and the Young's inequality, it follows that

$$\langle \mathcal{A}_{\lambda_{k}}(t, X_{k}(t)) + B(t, X_{k}(t)), X_{k}(t) - X(t) \rangle$$

$$\geq C \| X_{k}(t) \|_{V}^{\alpha} + Cf(t)(1 + \| X_{k}(t) \|_{H}^{2}) - \| \mathcal{A}_{\lambda_{k}}(t, X_{k}(t)) + B(t, X_{k}(t)) \|_{V^{*}} \| X(t) \|_{V}$$

$$\geq C \| X_{k}(t) \|_{V}^{\alpha} + Cf(t)(1 + \| X_{k}(t) \|_{H}^{2})$$

$$- \left[f(t) + C \| X_{k}(t) \|_{V}^{\alpha} \right]^{\frac{\alpha - 1}{\alpha}} \left[1 + \| X_{k}(t) \|_{H}^{\beta} \right]^{\frac{\alpha - 1}{\alpha}} \| X(t) \|_{V}$$

$$\geq \frac{C}{2} \| X_{k}(t) \|_{V}^{\alpha} - Cf(t)(1 + \| X_{k}(t) \|_{H}^{2}) - C \| X(t) \|_{V}^{\alpha} - C \| X_{k}(t) \|_{H}^{\beta(\alpha - 1)} \| X(t) \|_{V}^{\alpha}.$$

$$(3.45)$$

We denote

$$g_k(t,\omega) := \langle \mathcal{A}_{\lambda_k}(t, X_k(t,\omega)) + B(t, X_k(t,\omega)), X_k(t,\omega) - X(t,\omega) \rangle,$$

$$F_k(t,\omega) := C \Big(f(t)(1 + \|X_k(t,\omega)\|_H^2) + \|X(t,\omega)\|_V^\alpha + \|X_k(t,\omega)\|_H^{\beta(\alpha-1)} \|X(t,\omega)\|_V^\alpha \Big).$$

Then, (3.45) reduces to

$$g_k(t,\omega) \ge \frac{C}{2} \|X_k(t,\omega)\|_V^\alpha - F_k(t,\omega).$$
(3.46)

We divide the rest of our proof in three steps.

Step 1. In this part, we intend to prove that for a.e. (t, ω) ,

$$\liminf_{k \to \infty} g_k(t, \omega) \ge 0. \tag{3.47}$$

Due to (3.40) and Lemma 3.1, there exists a measurable subset Γ of $[0,T] \times \Omega$ such that $([0,T] \times \Omega) \setminus \Gamma$ is an $dt \otimes \mathbb{P}$ -null set,

$$\lim_{k \to \infty} \|X_k(t,\omega) - X(t,\omega)\|_H = 0, \quad \forall (t,\omega) \in \Gamma,$$
(3.48)

and $B(t, \cdot)$ is pseudo-monotone for any $(t, \omega) \in \Gamma$. Taking any fixed $(t, \omega) \in \Gamma$ and assuming that

 $\liminf_{k \to \infty} g_k(t, \omega) < 0.$

Therefore, there exists a subsequence $\{k_i\}_{i\in\mathbb{N}}$ such that

$$\lim_{i \to \infty} g_{k_i}(t, \omega) < 0. \tag{3.49}$$

Combining (3.46) and (3.48) yields that $\{\|X_{k_i}(t,\omega)\|_V^{\alpha}\}_{i\in\mathbb{N}}$ is bounded. Thus, we have an element $z \in V$ such that $X_{k_i}(t,\omega)$ converges weakly to z in V. According to (3.48), it is clear that $z = X(t,\omega)$ and

$$X_{k_i}(t,\omega) \rightharpoonup X(t,\omega)$$
 in V as $i \to \infty$.

By the monotonicity of $A_{\lambda_{k_i}}$ and (3.49), it follows that

$$\langle B(t, X_{k_i}(t, \omega), X_{k_i}(t, \omega) - X(t, \omega)) \rangle$$

$$< - \langle A_{\lambda_{k_i}}(t, X_{k_i}(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \rangle$$

$$\leq - \langle A_{\lambda_{k_i}}(t, X(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \rangle.$$
(3.50)

By Proposition 2.3 (iii), we know that $A_{\lambda_{k_i}}(x) \to A^0(x)$ for $x \in \mathcal{D}(A)$. Since $\mathcal{D}(A) = V$ and $X_{k_i}(t,\omega) \to X(t,\omega)$ in V for a.e. (t,ω) , in view of Proposition 21.23 (j) in [58] we have

$$\langle A_{\lambda_{k_i}}(t, X(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \rangle \to 0 \text{ as } i \to \infty$$

Thus, we can get

$$\liminf_{i \to \infty} \langle A_{\lambda_{k_i}}(t, X_{k_i}(t, \omega), X_{k_i}(t, \omega) - X(t, \omega) \rangle \ge 0.$$
(3.51)

Then, by (3.50), we have

$$\limsup_{i \to \infty} \langle B(t, X_{k_i}(t, \omega), X_{k_i}(t, \omega) - X(t, \omega) \rangle \leq 0.$$

Applying the pseudo-monotonicity of B, it follows that

$$\liminf_{i \to \infty} \langle B(t, X_{k_i}(t, \omega), X_{k_i}(t, \omega) - X(t, \omega) \rangle \ge 0.$$
(3.52)

Combining (3.51) and (3.52) yields that

$$\liminf_{i \to \infty} \langle A_{\lambda_{k_i}}(t, X_{k_i}(t, \omega)) + B(t, X_{k_i}(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \ge 0,$$

that is

$$\liminf_{i \to \infty} g_{k_i}(t, \omega) \ge 0$$

which contradicts to (3.49). Hence, we complete the proof of (3.47).

Step 2. In this step, we prove that there exists a subsequence $\{k_i\}$ such that

$$\lim_{i \to \infty} g_{k_i}(t,\omega) = 0, \quad \text{for a.e. } (t,\omega).$$
(3.53)

In view of (3.48), we note that $F_k(t, \omega)$ is convergent for a.e. (t, ω) . By (3.29), we also know that F_k is uniformly integrable. Therefore, in view of the generalized Fatou's lemma (see e.g. [18, p10]), (3.46) and Step 1, it follows that

$$\liminf_{k \to \infty} \mathbb{E} \int_0^T g_k(t) dt \ge \mathbb{E} \int_0^T \liminf_{k \to \infty} g_k(t) dt \ge 0.$$
(3.54)

Using conditions (3.42)-(3.44) yields

$$\limsup_{k \to \infty} \mathbb{E} \int_0^T g_k(t) dt \leqslant 0.$$
(3.55)

Hence, combining (3.54) and (3.55) reduces

$$\lim_{k \to \infty} \mathbb{E} \int_0^T g_k(t) dt = 0.$$
(3.56)

Set $g_k^-(t,\omega) := \min\{g_k(t,\omega), 0\}$. From Step 1, we know that for a.e. (t,ω) ,

$$\lim_{k \to \infty} g_k^-(t,\omega) = 0.$$

By (3.46) and the uniform integrability of F_k , it follows that

$$\lim_{k \to \infty} \mathbb{E} \int_0^T g_k^-(t) dt = 0.$$

Then, applying $|g_k| = g_k - 2g_k^-$ and (3.56), we obtain

$$\lim_{k \to \infty} \mathbb{E} \int_0^T |g_k(t)| dt = 0.$$

Thus, (3.53) follows.

Step 3. In this step, we prove that $\mathcal{B}(\cdot) = B(\cdot, X(\cdot)), dt \otimes \mathbb{P}$ -a.e..

From (3.46) and Step 2, it follows that

$$\sup_{i \in \mathbb{N}} \|X_{k_i}(t,\omega)\|_V < \infty \quad \text{for a.e. } (t,\omega).$$
(3.57)

Combining (3.48) and (3.57) yields for a.e. (t, ω) , as $i \to \infty$,

$$X_{k_i}(t,\omega) \rightharpoonup X(t,\omega)$$
 in V.

By the monotonicity of $A_{\lambda_{k_i}}$ and Step 2, we have a.e. (t, ω) ,

$$\begin{split} &\limsup_{i \to \infty} \langle B(t, X_{k_i}(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \rangle \\ &= -\liminf_{i \to \infty} \langle A_{\lambda_{k_i}}(t, X_{k_i}(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \rangle \\ &\leqslant -\liminf_{i \to \infty} \langle A_{\lambda_{k_i}}(t, X(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \rangle \end{split}$$

Using the same argument as in the proof of Step 1, we can get that for a.e. (t, ω) , as $i \to \infty$,

$$\langle A_{\lambda_{k_i}}(t, X(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \rangle \to 0.$$

Hence, it follows that

$$\limsup_{i \to \infty} \langle B(t, X_{k_i}(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \rangle \leq 0.$$
(3.58)

The inequality (3.58) together with the pseudo-monotonicity of B yields that for a.e. (t, ω) , as $i \to \infty$,

$$B(t, X_{k_i}(t, \omega)) \rightharpoonup B(t, X(t, \omega))$$
 in V^* ,

and

$$\lim_{i \to \infty} \left\langle B(t, X_{k_i}(t, \omega)), X_{k_i}(t, \omega) - X(t, \omega) \right\rangle = 0.$$
(3.59)

According to (3.35) and the uniqueness of limits, we deduce that $\mathcal{B}(\cdot) = B(\cdot, X(\cdot))$, which completes the proof of Lemma 3.8.

By combining the above lemmas, we present the proof of Theorem 3.1 as follows.

Proof of Theorem 3.1. We intend to prove that X is a weak solution to (1.3) in the sense of Definition 3.1. By Lemma 3.7, we have already proved $\mathcal{D}(\cdot) = \sigma(\cdot, X(\cdot)), dt \otimes \mathbb{P}$ -a.e.. Now, we show that

$$\mathcal{B}(\cdot) = B(\cdot, X(\cdot)), \ dt \otimes \mathbb{P}\text{-a.e.}.$$
(3.60)

Note that if we can prove (3.44), by Lemma 3.8 the result follows immediately. According to Lemma 3.7, we know that for any $t \in [0, T]$,

$$\mathbb{E} \int_{0}^{t} \|\mathcal{D}(s)\|_{\mathcal{L}_{2}}^{2} ds = \mathbb{E} \int_{0}^{t} \|\sigma(s, X(s))\|_{\mathcal{L}_{2}}^{2} ds.$$
(3.61)

Applying Itô's formula to the equation (3.28) and (3.38), which satisfied by X_k and X respectively, and taking expectations, by (3.61) we obtain that for any $t \in [0, T]$,

$$\mathbb{E} \|X_{k}(t)\|_{H}^{2} = \|x\|_{H}^{2} - 2\mathbb{E} \int_{0}^{t} \langle \mathcal{A}_{\lambda_{k}}(s, X_{k}(s)), X_{k}(s) \rangle \, ds \\ - 2\mathbb{E} \int_{0}^{t} \langle B(s, X_{k}(s)), X_{k}(s) \rangle \, ds + \mathbb{E} \int_{0}^{t} \|\sigma(s, X_{k}(s))\|_{\mathcal{L}_{2}}^{2} ds, \qquad (3.62)$$

$$\mathbb{E}\|X(t)\|_{H}^{2} = \|x\|_{H}^{2} - 2\mathbb{E}\int_{0}^{t} \langle \eta(s), X(s) \rangle \, ds$$
$$- 2\mathbb{E}\int_{0}^{t} \langle \mathcal{B}(s), X(s) \rangle \, ds + \mathbb{E}\int_{0}^{t} \|\sigma(s, X(s))\|_{\mathcal{L}_{2}}^{2} ds. \tag{3.63}$$

Since $||X_k - X||_{C([0,T],V^*)} \to 0$, by the lower semicontinuity of $|| \cdot ||_H$ in V^* and Fatou's lemma, it follows that for any $t \in [0,T]$,

$$\mathbb{E}\|X(t)\|_{H}^{2} \leq \mathbb{E}\left[\liminf_{k \to \infty} \|X_{k}(t)\|_{H}^{2}\right] \leq \liminf_{k \to \infty} \mathbb{E}\|X_{k}(t)\|_{H}^{2}.$$
(3.64)

Collecting (3.62)-(3.64), in view of (3.41) it implies that for any $t \in [0, T]$,

$$2 \limsup_{k \to \infty} \mathbb{E} \int_0^t \langle \mathcal{A}_{\lambda_k}(s, X_k(s)) + B(s, X_k(s)), X_k(s) \rangle ds$$

$$\leq 2 \mathbb{E} \int_0^t \langle \eta(s) + \mathcal{B}(s), X(s) \rangle ds + \liminf_{k \to \infty} \mathbb{E} \int_0^t \left(\|\sigma(s, X_k(s))\|_{\mathcal{L}_2}^2 - \|\sigma(s, X(s))\|_{\mathcal{L}_2}^2 \right) ds$$

$$\leq 2 \mathbb{E} \int_0^t \langle \eta(s) + \mathcal{B}(s), X(s) \rangle ds + \liminf_{k \to \infty} \mathbb{E} \int_0^t \|\sigma(s, X_k(s)) - \sigma(s, X(s))\|_{\mathcal{L}_2}^2 ds$$

$$= 2 \mathbb{E} \int_0^t \langle \eta(s) + \mathcal{B}(s), X(s) \rangle ds.$$

Thus, we can conclude that

$$\limsup_{k \to \infty} \mathbb{E} \int_0^T \langle \mathcal{A}_{\lambda_k}(t, X_k(t)) + B(t, X_k(t)), X_k(t) \rangle dt \leq \mathbb{E} \int_0^T \langle \eta(t) + \mathcal{B}(t), X(t) \rangle dt,$$

which implies that (3.60) holds.

Finally, once we can show that $\eta \in \mathcal{A}(\cdot, X)$, $dt \otimes \mathbb{P}$ -a.e. whose proof is provided in the following lemma, we can deduce that X is a weak solution to Eq. (1.3). The moment estimates (3.1) for X are derived from the estimates (3.30) and (3.31). Thus, we complete the proof.

The following lemma characterizes the limit of $\mathcal{A}(\cdot, X_k(\cdot))$.

Lemma 3.9. $\eta \in \mathcal{A}(\cdot, X), dt \otimes \mathbb{P}$ -a.e..

Proof. Claim 1. the multi-valued operator

$$L^{\alpha}([0,T] \times \Omega, V) \ni x \mapsto \Psi(x) := \mathcal{A}(\cdot, x(\cdot)) \in 2^{L^{\frac{\alpha}{\alpha-1}}([0,T] \times \Omega, V^*)}$$

is maximal-monotone.

Let $x_1, x_2 \in L^{\alpha}([0,T] \times \Omega, V)$ and $v_i \in \Psi(x_i), i = 1, 2$. In view of the monotonicity of \mathcal{A} , it follows that

$${}_{L^{\frac{\alpha}{\alpha-1}}([0,T]\times\Omega,V^*)}\langle v_1 - v_2, x_1 - x_2 \rangle_{L^{\alpha}([0,T]\times\Omega,V)} = \mathbb{E}\int_0^T \langle v_1(t) - v_2(t), x_1(t) - x_2(t) \rangle dt \ge 0.$$

Hence, Ψ is monotone.

Since for a.e. $t \in [0,T]$ the operator $\mathcal{A}(t,\cdot)$ is maximal-monotone, by Proposition 3.14 in [33], we have that for any $Y \in L^{\frac{\alpha}{\alpha-1}}([0,T] \times \Omega, V^*)$ there exists a progressively measurable process X(t) such that for a.e. $t \in [0,T]$ and any $\lambda > 0$,

$$Y(t,\omega) \in \mathcal{A}(t,X(t,\omega)) + \lambda J(X(t,\omega)), \ \omega \in \Omega.$$

Let $Z(t,\omega) \in \mathcal{A}(t,X(t,\omega))$ such that

$$Y(t,\omega) = Z(t,\omega) + \lambda J(X(t,\omega))$$
 for any $\omega \in \Omega$ and a.e. $t \in [0,T]$.

Taking the dualization product with $X(t,\omega)$, due to $(\mathbf{H}^2_{\mathcal{A}})$ we can get for a.e. $(t,\omega) \in [0,T] \times \Omega$

$$\langle Y(t,\omega), X(t,\omega) \rangle = \langle Z(t,\omega), X(t,\omega) \rangle + \lambda \langle J(X(t,\omega)), X(t,\omega) \rangle \geq (\delta + \lambda) \| X(t,\omega) \|_{V}^{\alpha} - f(t).$$

Since $Y \in L^{\frac{\alpha}{\alpha-1}}([0,T] \times \Omega, V^*)$, according to Young's inequality, we have $X \in L^{\alpha}([0,T] \times \Omega, V)$. By Theorem 6.1 in Appendix below, we deduce that the mapping Ψ is maximal-monotone. The claim follows.

Claim 2. $\mathcal{R}_{\lambda_k}(X_k)$ converges weakly to X in $L^{\alpha}([0,T] \times \Omega, V)$, as $k \to \infty$.

By the definition of the generalized Yosida approximation (2.5), it follows that for any $(t, \omega) \in [0, T] \times \Omega$

$$\lambda_k \mathcal{A}_{\lambda_k}(t, X_k(t, \omega)) = J(X_k(t, \omega) - \mathcal{R}_{\lambda_k}(X_k(t, \omega))),$$

which implies

$$\|\mathcal{R}_{\lambda_k}(X_k(t,\omega)) - X_k(t,\omega)\|_V^{\alpha} = \lambda_k^{\frac{\alpha}{\alpha-1}} \|\mathcal{A}_{\lambda_k}(t,X_k(t,\omega))\|_{V^*}^{\frac{\alpha}{\alpha-1}}.$$

Hence, in terms of Lemma 3.4 we can deduce that

$$\mathbb{E}\int_0^T \|\mathcal{R}_{\lambda_k}(X_k(s)) - X_k(s)\|_V^\alpha ds = \lambda_k^{\frac{\alpha}{\alpha-1}} \mathbb{E}\int_0^T \|\mathcal{A}_{\lambda_k}(s, X_k(s))\|_{V^*}^{\frac{\alpha}{\alpha-1}} ds \xrightarrow{k \to \infty} 0,$$

that is,

$$\lim_{k \to \infty} \|\mathcal{R}_{\lambda_k}(X_k) - X_k\|_{L^{\alpha}([0,T] \times \Omega, V)} = 0.$$
(3.65)

Thus, we have for any $\varphi \in L^{\frac{\alpha}{\alpha-1}}([0,T] \times \Omega, V^*)$,

$$\mathbb{E}\int_0^T \langle \varphi(s), \mathcal{R}_{\lambda_k}(X_k(s)) - X_k(s) \rangle ds \xrightarrow{k \to \infty} 0.$$

Since $X_k \to X$ in $L^{\alpha}([0,T] \times \Omega, V)$, we can get that for any $\varphi \in L^{\frac{\alpha}{\alpha-1}}([0,T] \times \Omega, V^*)$,

$$\mathbb{E} \int_0^T \langle \varphi(s), \mathcal{R}_{\lambda_k}(X_k(s)) - X(s) \rangle ds$$

= $\mathbb{E} \int_0^T \langle \varphi(s), \mathcal{R}_{\lambda_k}(X_k(s)) - X_k(s) \rangle ds + \mathbb{E} \int_0^T \langle \varphi(s), X_k(s) - X(s) \rangle ds \xrightarrow{k \to \infty} 0.$

The claim follows.

Claim 3. $\limsup_{k \to \infty} \mathbb{E} \int_0^T \langle \mathcal{A}_{\lambda_k}(s, X_k(s)), \mathcal{R}_{\lambda_k}(X_k(s)) \rangle \, ds \leq \mathbb{E} \int_0^T \langle \eta(s), X(s) \rangle \, ds.$

By (3.62) and using the fact $a^2 = (a-b)^2 - b^2 + 2ab$, we deduce that for any $t \in [0,T]$,

$$\mathbb{E}\|X_k(t)\|_H^2 + 2\mathbb{E}\int_0^t \left\langle \mathcal{A}_{\lambda_k}(s, X_k(s)), X_k(s) \right\rangle ds + I_1 + I_2 = \|x\|_H^2, \tag{3.66}$$

where

$$I_{1} := \mathbb{E} \int_{0}^{t} \left(2 \langle B(s, X_{k}(s)) - B(s, X(s)), X_{k}(s) - X(s) \rangle - \|\sigma(s, X_{k}(s)) - \sigma(s, X(s))\|_{\mathcal{L}_{2}}^{2} \right) ds,$$

$$I_{2} := \mathbb{E} \int_{0}^{t} \left(2 \langle B(s, X(s)), X_{k}(s) \rangle + 2 \langle B(s, X_{k}(s)) - B(s, X(s)), X(s) \rangle + \|\sigma(s, X(s))\|_{\mathcal{L}_{2}}^{2} - 2 \langle \sigma(s, X_{k}(s)), \sigma(s, X(s)) \rangle_{\mathcal{L}_{2}} \right) ds.$$

By (3.32), (3.33) and (3.59), there exists a subsequence, still denoted by k, such that for all $t \in [0, T]$, we have

$$\lim_{k \to \infty} \mathbb{E} \int_0^t \left\langle B(s, X_k(s)), X_k(s) - X(s) \right\rangle ds = 0$$
(3.67)

Using the convergence properties (3.33)-(3.37), we conclude

$$\lim_{k \to \infty} I_2 = \mathbb{E} \int_0^t \left(2 \left\langle B(s, X(s)), X(s) \right\rangle + 2 \left\langle \mathcal{B}(s) - B(s, X(s)), X(s) \right\rangle \right. \\ \left. + \left\| \sigma(s, X(s)) \right\|_{\mathcal{L}_2}^2 - 2 \left\langle \mathcal{D}(s), \sigma(s, X(s)) \right\rangle_{\mathcal{L}_2} \right) ds.$$
(3.68)

Substituting (3.64) and (3.68) into (3.66) and recalling (3.63), we obtain

$$\limsup_{k \to \infty} 2\mathbb{E} \int_0^T \left(\langle \mathcal{A}_{\lambda_k}(s, X_k(s)), X_k(s) \rangle - \langle \eta(s), X(s) \rangle \right) ds$$

$$\leqslant \mathbb{E} \int_0^T \left(2 \langle \mathcal{B}(s) - B(s, X(s)), X(s) - X(s) \rangle - \|\sigma(s, X(s)) - \mathcal{D}(s)\|_{\mathcal{L}_2}^2 \right) ds$$

$$-\limsup_{k \to \infty} I_1.$$
(3.69)

In view of (3.33), (3.41) and (3.67), it follows that

$$\lim_{k \to \infty} I_1 = 0.$$

Recall that $\mathcal{B} = B(\cdot, X)$ and $\mathcal{D} = \sigma(\cdot, X)$. From (3.69), we conclude that

$$\limsup_{k \to \infty} \mathbb{E} \int_0^T \left\langle \mathcal{A}_{\lambda_k}(s, X_k(s)), X_k(s) \right\rangle ds \leqslant \mathbb{E} \int_0^T \left\langle \eta(s), X(s) \right\rangle ds.$$
(3.70)

Applying Hölder's inequality, Lemma 3.4, (3.65) and (3.70), we deduce that

$$\begin{split} &\limsup_{k \to \infty} \mathbb{E} \int_{0}^{T} \left\langle \mathcal{A}_{\lambda_{k}}(s, X_{k}(s)), \mathcal{R}_{\lambda_{k}}(X_{k}(s)) \right\rangle ds \\ &\leqslant \limsup_{k \to \infty} \left(\left(\mathbb{E} \int_{0}^{T} \|\mathcal{A}_{\lambda_{k}}(s, X_{k}(s))\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} ds \right)^{\frac{\alpha-1}{\alpha}} \cdot \|\mathcal{R}_{\lambda_{k}}(X_{k}) - X_{k}\|_{L^{\alpha}([0,T] \times \Omega, V)} \right) \\ &+ \limsup_{k \to \infty} \mathbb{E} \int_{0}^{T} \left\langle \mathcal{A}_{\lambda_{k}}(s, X_{k}(s)), X_{k}(s) \right\rangle ds \\ &\leqslant \mathbb{E} \int_{0}^{T} \left\langle \eta(s), X(s) \right\rangle ds. \end{split}$$

The claim follows.

Now, we proceed to prove the lemma. Since $\mathcal{A}_{\lambda_k}(\cdot, X_k(\cdot))$ converges weakly to η in $L^{\frac{\alpha}{\alpha-1}}([0,T]\times\Omega, V^*)$, by **Claims 1-3** along with the fact that $\mathcal{A}_{\lambda_k}(\cdot, X_k(\cdot)) \in \mathcal{A}(\cdot, \mathcal{R}_{\lambda_k}(X_k(\cdot)))$ (cf. Proposition 2.3 (iv)), Lemma 2.1 implies that

$$\eta \in \Psi(X), \ dt \otimes \mathbb{P}\text{-a.e.},$$

that is, $\eta(t,\omega) \in \mathcal{A}(t,X(t,\omega))$ for a.e. $(t,\omega) \in [0,T] \times \Omega$.

3.5. **Proof of Theorem 3.3.** It suffices to prove the pathwise uniqueness of solutions to (1.3). Then, Theorem 3.3 follows directly from the Yamada-Watanabe theorem.

Let $(X_1, \eta_1), (X_2, \eta_2)$ be two solutions of Eq. (1.3) with initial values $X_1(0) = X_2(0) = x$. Set

$$\varphi(t) := \exp\left\{-\int_0^t \left(f(s) + \rho(X_1(s)) + \zeta(X_2(s))\right) ds\right\}.$$

Applying Itô's formula and by $(\mathbf{H}^1_{\mathcal{A}})$ and (\mathbf{H}^4_B) , it follows that

$$\begin{aligned} \varphi(t) \|X_{1}(t) - X_{2}(t)\|_{H}^{2} \\ &= \int_{0}^{t} \varphi(s) \bigg\{ -2 \langle \eta_{1}(s) - \eta_{2}(s), X_{1}(s) - X_{2}(s) \rangle \\ &- 2 \langle B(s, X_{1}(s)) - B(s, X_{2}(s)), X_{1}(s) - X_{2}(s) \rangle + \|\sigma(s, X_{1}(s)) - \sigma(s, X_{2}(s))\|_{\mathcal{L}_{2}}^{2} \\ &- [f(s) + \rho(X_{1}(s)) + \zeta(X_{2}(s))] \|X_{1}(s) - X_{2}(s)\|_{H}^{2} \bigg\} ds \\ &+ 2 \int_{0}^{t} \varphi(s) \langle X_{1}(s) - X_{2}(s), (\sigma(s, X_{1}(s)) - \sigma(s, X_{2}(s))) dW(s) \rangle_{H} \\ \leqslant 2 \int_{0}^{t} \varphi(s) \langle X_{1}(s) - X_{2}(s), (\sigma(s, X_{1}(s)) - \sigma(s, X_{2}(s))) dW(s) \rangle_{H} . \end{aligned}$$
(3.71)

Let $\{\tau_n\}$ be a sequence of stopping times such that the local martingale in (3.71) becomes a martingale up to τ_n . Then, by taking the expectation on both sides of (3.71), it implies that

$$\mathbb{E}\left[\varphi(t\wedge\tau_n)\|X_1(t\wedge\tau_n) - X_2(t\wedge\tau_n)\|_H^2\right] \leqslant 0.$$
(3.72)

Letting $n \to \infty$ and applying Fatou's lemma, we deduce that

$$\mathbb{E}\left[\varphi(t)\|X_{1}(t) - X_{2}(t)\|_{H}^{2}\right] \leqslant 0.$$
(3.73)

Since

$$\int_0^T \left(f(r) + \rho(X_1(r)) + \zeta(X_2(r)) \right) dr < \infty, \quad \mathbb{P}\text{-a.s.},$$

then (3.73) yields the pathwise uniqueness of solutions to Eq. (1.3) by the pathwise continuity of solutions in H.

3.6. Proof of Theorem 3.4. Let the stopping time τ_n^M be defined as follows

$$\begin{aligned} \tau_n^M &:= T \wedge \inf \left\{ t \ge 0 : \|X(t, x_n)\|_H > M \right\} \\ & \wedge \inf \left\{ t \ge 0 : \int_0^t \|X(s, x_n)\|_V^\alpha ds > M \right\} \\ & \wedge \inf \left\{ t \ge 0 : \|X(t, x)\|_H > M \right\} \\ & \wedge \inf \left\{ t \ge 0 : \int_0^t \|X(s, x)\|_V^\alpha ds > M \right\}, \ M > 0 \end{aligned}$$

First, we note that by the moment estimates (3.1) for the solutions, it follows that

$$\lim_{M \to \infty} \sup_{n \in \mathbb{N}} \mathbb{P}(\tau_n^M < T) = 0.$$
(3.74)

Let $\varphi_n(t) := \exp\left\{-\int_0^t \left(f(r) + \rho(X(r, x_n)) + \zeta(X(r, x))\right) dr\right\}$. Combining (3.72) and (3.73), we obtain

$$\mathbb{E}\Big[\varphi_n(t\wedge\tau_n^M)\|X(t\wedge\tau_n^M,x_n)-X(t\wedge\tau_n^M,x)\|_H^2\Big] \leqslant \|x_n-x\|_H^2.$$
(3.75)

In light of Chebyshev's inequality and (3.75), there exists a constant $C_M > 0$ such that for any $\varepsilon > 0$,

$$\mathbb{P}(\|X(t,x_n) - X(t,x)\|_H \ge \varepsilon)
\leqslant \mathbb{P}(\|X(t,x_n) - X(t,x)\|_H \ge \varepsilon, \tau_n^M \ge T) + \mathbb{P}(\tau_n^M < T)
\leqslant \frac{1}{\varepsilon^2 C_M} \mathbb{E}\Big[\varphi_n(t \wedge \tau_n^M) \|X(t \wedge \tau_n^M, x_n) - X(t \wedge \tau_n^M, x)\|_H^2\Big] + \mathbb{P}(\tau_n^M < T)
\leqslant \frac{1}{\varepsilon^2 C_M} \|x_n - x\|_H^2 + \sup_{n \in \mathbb{N}} \mathbb{P}(\tau_n^M < T).$$
(3.76)

Taking the limit as $n \to \infty$ and $M \to \infty$ in (3.76) and using (3.74), we conclude that for any $t \in [0, T]$,

$$||X(t,x_n) - X(t,x)||_H \xrightarrow{n \to \infty} 0$$
 in probability.

Furthermore, by (3.1) we have for any $p \ge 2$,

$$\sup_{n \in \mathbb{N}} \mathbb{E} \Big[\sup_{t \in [0,T]} \| X(t,x_n) \|_H^p \Big] < \infty.$$
(3.77)

Then, the Vitali's theorem implies that

$$\lim_{n \to \infty} \mathbb{E} \int_0^T \|X(t, x_n) - X(t, x)\|_H^2 dt = 0,$$

which yields that

$$|X(t,x_n) - X(t,x)||_H \xrightarrow{n \to \infty} 0, \quad dt \otimes \mathbb{P}\text{-a.e.}.$$

Hence, in view of (\mathbf{H}_{σ}) and (3.77), we have

$$\lim_{n \to \infty} \mathbb{E} \int_0^T \|\sigma(t, X(t, x_n)) - \sigma(t, X(t, x))\|_{\mathcal{L}_2}^2 dt = 0.$$
(3.78)

Then, using (3.71), B-D-G's inequality and Young's inequality, we obtain

$$\mathbb{E} \left[\sup_{t \in [0, T \wedge \tau_n^M]} \left(\varphi_n(t) \| X(t, x_n) - X(t, x) \|_H^2 \right) \right] \\
\leq \|x_n - x\|_H^2 + 2\mathbb{E} \left[\sup_{t \in [0, T \wedge \tau_n^M]} \left| \int_0^t \varphi_n(s) \langle X(s, x_n) - X(s, x), \left(\sigma(s, X(s, x_n)) - \sigma(s, X(s, x_n)) \right) \right| \right] \\
= \sigma(s, X(s, x)) dW(s) \| \\
= \|x_n - x\|_H^2 \\
+ C\mathbb{E} \left(\int_0^{T \wedge \tau_n^M} \varphi_n(t)^2 \| X(t, x_n) - X(t, x) \|_H^2 \| \sigma(t, X(t, x_n)) - \sigma(t, X(t, x)) \|_{\mathcal{L}_2}^2 dt \right)^{\frac{1}{2}} \\
\leq \|x_n - x\|_H^2 + \frac{1}{2} \mathbb{E} \left[\sup_{t \in [0, T \wedge \tau_n^M]} \left(\varphi_n(t) \| X(t, x_n) - X(t, x) \|_H^2 \right) \right] \\
+ C\mathbb{E} \int_0^{T \wedge \tau_n^M} \varphi_n(t) \| \sigma(t, X(t, x_n)) - \sigma(t, X(t, x)) \|_{\mathcal{L}_2}^2 dt.$$
(3.79)

Combining (3.78) and (3.79) implies

$$\lim_{n \to \infty} \mathbb{E} \Big[\sup_{t \in [0, T \land \tau_n^M]} \big(\varphi_n(t) \| X(t, x_n) - X(t, x) \|_H^2 \big) \Big] = 0.$$

A similar argument as in the proof of (3.76) leads to

$$\sup_{t \in [0,T]} \|X(t,x_n) - X(t,x)\|_H \xrightarrow{n \to \infty} 0 \quad \text{in probability.}$$

Furthermore, from (3.77) we have

$$\lim_{n \to \infty} \mathbb{E} \Big[\sup_{t \in [0,T]} \| X(t,x_n) - X(t,x) \|_H^p \Big] = 0,$$

which completes the proof.

YOSIDA APPROXIMATION AND MULTI-VALUED SEI

4. FINITE TIME EXTINCTION

Self-organized criticality is extensively studied in physics from various perspectives (cf. [56]). For example, the authors in [7] have suggested that the continuum limit of the Bak-Tang-Wiesenfeld sandpile model, as introduced in [2], with a stochastic force can be interpreted as a type of multi-valued stochastic porous media equations. They have established that the finite time extinction results for this type of multi-valued SPDEs are particularly relevant to the self-organized critical behavior of the BTW model. Subsequently, further investigations have been conducted on the finite time extinction of stochastic (sign) fast-diffusion equations perturbed by linear multiplicative noises (cf. [7, 9, 10, 20, 49]).

In this section, we demonstrate that the finite time extinction of solutions holds with probability one for all initial value $x \in H$, for a class of multi-valued stochastic evolution inclusions perturbed by linear multiplicative noise.

4.1. Main result. Building on the existence and uniqueness of solutions to Eq. (1.3) (i.e. Theorem 3.3), in this part, we mainly focus on the case of the linear multiplicative noise, i.e.,

$$\sigma(t,x)v := \sum_{k=1}^{\infty} h_k(t)x \langle v, g_k \rangle_U, \qquad (4.1)$$

where $\{g_k\}_{k\in\mathbb{N}}$ is an orthonormal basis on U, $\{h_k(t)\}_{k\in\mathbb{N}}$ is a sequence of real-valued functions. More precisely, we consider the noise of the following form

$$\int_0^t \sigma(s, X(s)) dW(s) = \sum_{k=1}^\infty \int_0^t h_k(s) X(s) d\beta_k(s),$$

where $\{\beta_k\}_{k\in\mathbb{N}}$ are independent standard Brownian motions defined on a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t\geq 0}, \mathbb{P})$.

We assume that there are constants $\alpha \in (1,2)$ and $\delta > 0$ such that the following conditions are satisfied for a.e. $t \in [0,\infty)$.

 $(\mathbf{H}_{\mathcal{A}}^{2*})$ For any $x \in V$ and $v \in \mathcal{A}(t, x)$,

$$\langle v, x \rangle \ge \delta \|x\|_V^{\alpha}$$

 (\mathbf{H}_B^{1*}) There exists $f \in L^1([0,\infty), [0,\infty))$ such that for any $x \in V$,

$$2\langle B(t,x),x\rangle \ge -f(t)\|x\|_{H}^{2}.$$

 $(\mathbf{H}_{\sigma}^{*})$ The function

$$h(t) := \sum_{k=1}^{\infty} |h_k(t)|^2 \in L^1([0,\infty), [0,\infty)),$$

which satisfies

$$(\alpha - 1)h(t) \ge f(t).$$

Remark 4.1. If the diffusion coefficient in (4.1) satisfies (\mathbf{H}_{σ}^*) , then it is easy to verify that (\mathbf{H}_{σ}) also holds.

Let τ_e be the extinction time as follows

$$\tau_e := \inf \{ t \ge 0 : \| X(t) \|_H = 0 \},\$$

where $X(t), t \ge 0$, is the solution of Eq. (1.3) with the initial value $x \in H$.

The following result shows the finite time extinction of solutions to Eq. (1.3).

Theorem 4.2. Suppose that $(\mathbf{H}^1_{\mathcal{A}})$, $(\mathbf{H}^{2*}_{\mathcal{A}})$, $(\mathbf{H}^3_{\mathcal{A}})$, (\mathbf{H}^{1*}_{B}) , (\mathbf{H}^2_{B}) - (\mathbf{H}^4_{B}) , and (\mathbf{H}^*_{σ}) hold. For any $x \in H$, we have for any $t \ge \tau_e$,

$$||X(t)||_{H} = 0, \ \mathbb{P}\text{-}a.s., \tag{4.2}$$

and there is a constant $c^* > 0$ such that for any T > 0,

$$\mathbb{P}(\tau_e \leqslant T) \ge 1 - \frac{c^* \|x\|_H^{2-\alpha}}{T}.$$
(4.3)

In particular, we have

$$\mathbb{P}(\tau_e < \infty) = 1.$$

Remark 4.3. In Theorem 4.2, we provide a quantitative characterization of the probability for the extinction time being less than any given time, which particularly implies the finite time extinction with probability one. This result is more general compared to previous works (cf. e.g. [7-10]). Additionally, we drop the requirement of small initial values, which is typically assumed in [7-10], and instead consider any initial value $x \in H$ (see Section 5 for concrete applications).

4.2. Approximating sequences. In order to prove Theorem 4.2, we first recall the approximating equations (3.2) constructed in the proof of Theorem 3.1, i.e.,

$$\begin{cases} dX_{\lambda}(t) + \left[\mathcal{A}_{\lambda}(t, X_{\lambda}(t)) + B(t, X_{\lambda}(t))\right] dt = \sum_{k=1}^{\infty} h_k(t) X_{\lambda}(t) d\beta_k(t), \\ X_{\lambda}(0) = x \in H, \end{cases}$$

$$(4.4)$$

where \mathcal{A}_{λ} is the generalized Yosida approximation of \mathcal{A} . Moreover, in view of the proof of Theorem 3.1 we take a subsequence $\{\lambda_n\}_{n=1}^{\infty}$, for which $\lambda_n \to 0$ as $n \to \infty$. Along this subsequence, we denote X_{λ_n} by X_n .

We establish the following convergence of the sequence $\{X_n(t)\}_{n\in\mathbb{N}}$, possibly along with a subsequence, to X(t).

Lemma 4.1. For any $t \in [0, \infty)$, we have

$$\mathbb{E}||X_n(t) - X(t)||_H^2 \to 0, \ as \ n \to \infty.$$

Proof. Recall the proof of Theorem 3.1, it is known that there exist a probability space $(\widetilde{\Omega}, \widetilde{\mathcal{F}}, \widetilde{\mathbb{P}})$ and sequences $\{(\widetilde{X}_n, \widetilde{W}_n)\}$ and $(\widetilde{X}, \widetilde{W})$ such that along a subsequence, still denoted by $\{\widetilde{X}_n\}$, we have

$$\lim_{n \to \infty} \|\widetilde{X}_n(t) - \widetilde{X}(t)\|_H = 0, \ dt \otimes \widetilde{\mathbb{P}}\text{-a.e.},$$

which combining with (3.29) and (3.30) implies

$$\lim_{n \to \infty} \widetilde{\mathbb{E}} \int_0^t \left(h(s) \| \widetilde{X}_n(s) - \widetilde{X}(s) \|_H^2 \right) ds = 0.$$
(4.5)

Applying Itô's formula for $\|\widetilde{X}_n(t) - \widetilde{X}(t)\|_H^2$ and in view of (3.28), (3.38) and (\mathbf{H}_{σ}^*) , it follows that

$$\begin{split} \|\widetilde{X}_{n}(t) - \widetilde{X}(t)\|_{H}^{2} \\ &= \int_{0}^{t} \left\{ -2\langle \mathcal{A}_{\lambda_{n}}(s, \widetilde{X}_{n}(s)) - \widetilde{\eta}(s), \widetilde{X}_{n}(s) - \widetilde{X}(s) \rangle \\ &- 2\langle B(s, \widetilde{X}_{n}(s)) - B(s, \widetilde{X}(s)), \widetilde{X}_{n}(s) - \widetilde{X}(s) \rangle + h(s) \|\widetilde{X}_{n}(s) - \widetilde{X}(s)\|_{H}^{2} \right\} ds \\ &+ 2\sum_{k=1}^{\infty} \int_{0}^{t} \left(h_{k}(s) \|\widetilde{X}_{n}(s) - \widetilde{X}(s)\|_{H}^{2} \right) d\beta_{s}^{k}. \end{split}$$

Then, taking the expectation on both sides, we have

$$\widetilde{\mathbb{E}} \|\widetilde{X}_{n}(t) - \widetilde{X}(t)\|_{H}^{2} = \widetilde{\mathbb{E}} \int_{0}^{t} \left\{ -2\langle \mathcal{A}_{\lambda_{n}}(s, \widetilde{X}_{n}(s)) - \widetilde{\eta}(s), \widetilde{X}_{n}(s) - \widetilde{X}(s) \rangle - 2\langle B(s, \widetilde{X}_{n}(s)) - B(s, \widetilde{X}(s)), \widetilde{X}_{n}(s) - \widetilde{X}(s) \rangle + h(s) \|\widetilde{X}_{n}(s) - \widetilde{X}(s)\|_{H}^{2} \right\} ds. \quad (4.6)$$

By the monotonicity of \mathcal{A} , Lemma 3.4 and (3.65), we deduce that

$$\lim_{n \to \infty} \widetilde{\mathbb{E}} \int_{0}^{t} \langle \widetilde{\eta}(s) - \mathcal{A}_{\lambda_{n}}(s, \widetilde{X}_{n}(s)), \widetilde{X}_{n}(s) - \widetilde{X}(s) \rangle ds$$

$$\leq \sup_{n \in \mathbb{N}} C \left\{ \widetilde{\mathbb{E}} \int_{0}^{t} \left(\|\mathcal{A}_{\lambda_{n}}(s, \widetilde{X}_{n}(s))\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} + \|\widetilde{\eta}(s)\|_{V^{*}}^{\frac{\alpha}{\alpha-1}} \right) ds \right\}^{\frac{\alpha-1}{\alpha}}$$

$$\cdot \lim_{n \to \infty} \left(\widetilde{\mathbb{E}} \int_{0}^{t} \|\mathcal{R}_{\lambda_{n}}(\widetilde{X}_{n}(s)) - \widetilde{X}_{n}(s)\|_{V}^{\alpha} ds \right)^{\frac{1}{\alpha}}$$

$$- \lim_{n \to \infty} \widetilde{\mathbb{E}} \int_{0}^{t} \langle \mathcal{A}_{\lambda_{n}}(s, \widetilde{X}_{n}(s)) - \eta(s), \mathcal{R}_{\lambda_{n}}(\widetilde{X}_{n}(s)) - \widetilde{X}(s) \rangle ds$$

$$\leq 0.$$

$$(4.7)$$

According to the proof of Lemma 3.8, it follows that there exists a subsequence, still denoted by $\{\tilde{X}_n\}$, such that for a.e. (t, ω) ,

$$\lim_{n \to \infty} \langle B(t, \widetilde{X}_n(t, \omega)), \widetilde{X}_n(t, \omega) - \widetilde{X}(t, \omega) \rangle = 0.$$

By applying the dominated convergence theorem and using (3.29) and (3.32), we conclude that for any $t \in [0, \infty)$,

$$\lim_{n \to \infty} \widetilde{\mathbb{E}} \int_0^t \langle B(s, \widetilde{X}_n(s)) - B(s, \widetilde{X}(s)), \widetilde{X}_n(s) - \widetilde{X}(s) \rangle ds = 0.$$
(4.8)

Combining (4.5)-(4.8) and using Gronwall's lemma implies that for any $t \in [0, \infty)$,

$$\lim_{n \to \infty} \widetilde{\mathbb{E}} \| \widetilde{X}_n(t) - \widetilde{X}(t) \|_H^2 = 0$$

In light of (3.26) and the uniqueness of solutions, we obtain that for any $t \in [0, \infty)$,

$$\lim_{n \to \infty} \mathbb{E} \|X_n(t) - X(t)\|_H^2 = 0$$

Hence, we complete the proof.

4.3. Proof of main result. The following lemma plays an important role in the derivation of (4.2) and (4.3).

Lemma 4.2. There exists a constant $\rho > 0$ such that for any $0 \leq r < t$,

$$||X(t)||_{H}^{2-\alpha} + \delta\rho^{\alpha}(1-\frac{\alpha}{2})\int_{r}^{t} \mathbf{1}_{\{||X(s)||_{H}>0\}}ds$$

$$\leq ||X(r)||_{H}^{2-\alpha} + 2(1-\frac{\alpha}{2})\sum_{k=1}^{\infty}\int_{r}^{t} \left(h_{k}(s)||X(s)||_{H}^{2-\alpha}\mathbf{1}_{\{||X(s)||_{H}>0\}}\right)d\beta_{s}^{k}.$$
 (4.9)

Proof. By Itô's formula for $||X_n(t)||_H^2$, which is defined in (4.4), and due to (\mathbf{H}_{σ}^*) , we have that for any $0 \leq r < t$,

$$||X_n(t)||_H^2 = ||X_n(r)||_H^2 + \int_r^t \left(-2\langle \mathcal{A}_{\lambda_n}(s, X_n(s)) + B(s, X_n(s)), X_n(s) \rangle + h(s) ||X_n(s)||_H^2 \right) ds + 2\sum_{k=1}^\infty \int_r^t \left(h_k(s) ||X_n(s)||_H^2 \right) d\beta_s^k.$$

For any $\varepsilon > 0$, we apply Itô's formula based on the auxiliary function $V^{\varepsilon}(x) := (\varepsilon + x)^{1-\frac{\alpha}{2}}$ to get

$$\begin{aligned} &(\varepsilon + \|X_n(t)\|_H^2)^{1-\frac{\alpha}{2}} + \delta 2^{-\alpha} (1 - \frac{\alpha}{2}) \int_r^t \frac{\|X_n(s)\|_V^{\alpha}}{(\varepsilon + \|X_n(s)\|_H^2)^{\frac{\alpha}{2}}} ds \\ &\leqslant (\varepsilon + \|X_n(r)\|_H^2)^{1-\frac{\alpha}{2}} + (1 - \frac{\alpha}{2}) \int_r^t \frac{(f(s) + h(s))\|X_n(s)\|_H^2}{(\varepsilon + \|X_n(s)\|_H^2)^{\frac{\alpha}{2}}} ds \\ &- \frac{\alpha}{2} (1 - \frac{\alpha}{2}) \int_r^t \frac{2h(s)\|X_n(s)\|_H^4}{(\varepsilon + \|X_n(s)\|_H^2)^{\frac{\alpha}{2} + 1}} ds + \mathcal{M}_{r,t}^{\varepsilon,n} \\ &= (\varepsilon + \|X_n(r)\|_H^2)^{1-\frac{\alpha}{2}} + (1 - \frac{\alpha}{2}) \int_r^t \frac{(f(s) + h(s))(\varepsilon + \|X_n(s)\|_H^2)\|X_n(s)\|_H^2}{(\varepsilon + \|X_n(s)\|_H^2)^{\frac{\alpha}{2} + 1}} ds \\ &- (1 - \frac{\alpha}{2}) \int_r^t \frac{\alpha h(s)\|X_n(s)\|_H^4}{(\varepsilon + \|X_n(s)\|_H^2)^{\frac{\alpha}{2} + 1}} ds + \mathcal{M}_{r,t}^{\varepsilon,n}, \end{aligned}$$

where

$$\mathcal{M}_{r,t}^{\varepsilon,n} := 2(1 - \frac{\alpha}{2}) \sum_{k=1}^{\infty} \int_{r}^{t} \frac{h_{k}(s) \|X_{n}(s)\|_{H}^{2}}{(\varepsilon + \|X_{n}(s)\|_{H}^{2})^{\frac{\alpha}{2}}} d\beta_{s}^{k},$$

and we used assumptions $(\mathbf{H}_{\mathcal{A}}^{2*})$, (\mathbf{H}_{B}^{1*}) and Lemma 6.1 in the first inequality. Since there exists a constant $\rho > 0$ such that $||x||_{V} \ge 2\rho ||x||_{H}$, it implies that

$$(\varepsilon + \|X_n(t)\|_H^2)^{1-\frac{\alpha}{2}} + \delta\rho^{\alpha} (1-\frac{\alpha}{2}) \int_r^t \frac{\|X_n(s)\|_H^{\alpha}}{(\varepsilon + \|X_n(s)\|_H^2)^{\frac{\alpha}{2}}} ds \leq (\varepsilon + \|X_n(r)\|_H^2)^{1-\frac{\alpha}{2}} + \mathcal{M}_{r,t}^{\varepsilon,n} + (1-\frac{\alpha}{2}) \int_r^t \frac{(f(s) + h(s))(\varepsilon + \|X_n(s)\|_H^2)\|X_n(s)\|_H^2}{(\varepsilon + \|X_n(s)\|_H^2)^{\frac{\alpha}{2}+1}} ds - (1-\frac{\alpha}{2}) \int_r^t \frac{\alpha h(s)\|X_n(s)\|_H^4}{(\varepsilon + \|X_n(s)\|_H^2)^{\frac{\alpha}{2}+1}} ds.$$

$$(4.10)$$

By Lemma 4.1 and (3.29), we can take $n \to \infty$ (possibly along with a subsequence) to (4.10) and apply the dominated convergence theorem. As a result, it follows that

$$\begin{split} &(\varepsilon + \|X(t)\|_{H}^{2})^{1-\frac{\alpha}{2}} + \delta\rho^{\alpha}(1-\frac{\alpha}{2})\int_{r}^{t}\frac{\|X(s)\|_{H}^{\alpha}}{(\varepsilon + \|X(s)\|_{H}^{2})^{\frac{\alpha}{2}}}ds\\ \leqslant &(\varepsilon + \|X(r)\|_{H}^{2})^{1-\frac{\alpha}{2}} + 2(1-\frac{\alpha}{2})\sum_{k=1}^{\infty}\int_{r}^{t}\frac{h_{k}(s)\|X(s)\|_{H}^{2}}{(\varepsilon + \|X(s)\|_{H}^{2})^{\frac{\alpha}{2}}}d\beta_{s}^{k}.\\ &+ (1-\frac{\alpha}{2})\int_{r}^{t}\frac{(f(s)+h(s))(\varepsilon + \|X(s)\|_{H}^{2})\|X(s)\|_{H}^{2}}{(\varepsilon + \|X(s)\|_{H}^{2})^{\frac{\alpha}{2}+1}}ds\\ &- (1-\frac{\alpha}{2})\int_{r}^{t}\frac{\alpha h(s)\|X(s)\|_{H}^{4}}{(\varepsilon + \|X(s)\|_{H}^{2})^{\frac{\alpha}{2}+1}}ds. \end{split}$$

Thus, we deduce that

$$\begin{split} & (\varepsilon + \|X(t)\|_{H}^{2})^{1-\frac{\alpha}{2}} + \delta\rho^{\alpha}(1-\frac{\alpha}{2})\int_{r}^{t} \left(\frac{\|X(s)\|_{H}^{\alpha}}{(\varepsilon + \|X(s)\|_{H}^{2})^{\frac{\alpha}{2}}} \mathbf{1}_{\{\|X(s)\|_{H}>0\}}\right) ds \\ \leqslant & (\varepsilon + \|X(r)\|_{H}^{2})^{1-\frac{\alpha}{2}} + 2(1-\frac{\alpha}{2})\sum_{k=1}^{\infty}\int_{r}^{t} \left(\frac{h_{k}(s)\|X(s)\|_{H}^{2}}{(\varepsilon + \|X(s)\|_{H}^{2})^{\frac{\alpha}{2}}} \mathbf{1}_{\{\|X(s)\|_{H}>0\}}\right) d\beta_{s}^{k} \\ & + (1-\frac{\alpha}{2})\int_{r}^{t} \left(\frac{(f(s)+h(s))(\varepsilon + \|X(s)\|_{H}^{2})\|X(s)\|_{H}^{2}}{(\varepsilon + \|X(s)\|_{H}^{2})^{\frac{\alpha}{2}+1}} \mathbf{1}_{\{\|X(s)\|_{H}>0\}}\right) ds \\ & - (1-\frac{\alpha}{2})\int_{r}^{t} \left(\frac{\alpha h(s)\|X(s)\|_{H}^{4}}{(\varepsilon + \|X(s)\|_{H}^{2})^{\frac{\alpha}{2}+1}} \mathbf{1}_{\{\|X(s)\|_{H}>0\}}\right) ds. \end{split}$$

Letting $\varepsilon \to 0$, it implies that

$$||X(t)||_{H}^{2-\alpha} + \delta \rho^{\alpha} (1 - \frac{\alpha}{2}) \int_{r}^{t} \mathbf{1}_{\{||X(s)||_{H} > 0\}} ds$$

$$\leq \|X(r)\|_{H}^{2-\alpha} + \mathcal{M}_{r,t} \\ + (1 - \frac{\alpha}{2}) \int_{r}^{t} \left(\frac{(f(s) + h(s) - \alpha h(s)) \|X(s)\|_{H}^{4}}{\|X(s)\|_{H}^{\alpha+2}} \mathbf{1}_{\{\|X(s)\|_{H} > 0\}} \right) ds,$$

where

$$\mathcal{M}_{r,t} := 2(1 - \frac{\alpha}{2}) \sum_{k=1}^{\infty} \int_{r}^{t} \left(h_{k}(s) \| X(s) \|_{H}^{2-\alpha} \mathbf{1}_{\{\|X(s)\|_{H} > 0\}} \right) d\beta_{s}^{k}.$$

Finally, according to assumption (\mathbf{H}_{σ}^*) , we conclude that

$$\|X(t)\|_{H}^{2-\alpha} + \delta\rho^{\alpha}(1-\frac{\alpha}{2})\int_{r}^{t} \mathbf{1}_{\{\|X(s)\|_{H}>0\}} ds \leq \|X(r)\|_{H}^{2-\alpha} + \mathcal{M}_{r,t},$$

which completes the proof.

Proof of Theorem 4.2. In view of Lemma 4.2, we have that for any $0 \leq r < t$,

$$\mathbb{E}[\|X(t)\|_{H}^{2-\alpha}|\mathcal{F}_{r}] \leqslant \|X(r)\|_{H}^{2-\alpha} + \mathbb{E}[\mathcal{M}_{r,t}|\mathcal{F}_{r}] = \|X(r)\|_{H}^{2-\alpha}$$

Therefore, $t \to ||X(t)||_{H}^{2-\alpha}$ is a nonnegative (\mathcal{F}_t) -supermartingale. This combining with (3.1) implies that for every pair of stopping times $\tau_1 < \tau_2$,

$$\mathbb{E} \|X(\tau_2)\|_H^{2-\alpha} \leqslant \mathbb{E} \|X(\tau_1)\|_H^{2-\alpha}.$$

In particular, for any $t > \tau_e$ it clear that

$$\mathbb{E} \|X(t)\|_{H}^{2-\alpha} \leq \mathbb{E} \|X(\tau_{e})\|_{H}^{2-\alpha} = 0$$

It implies that for any $t \ge \tau_e$,

$$||X(t)||_H = 0$$
, \mathbb{P} -a.s..

Thus, (4.2) follows.

On the other hand, letting r = 0 and taking the expectation on both sides of (4.9), we obtain that for any T > 0,

$$\mathbb{E}||X(T)||_{H}^{2-\alpha} + \delta\rho^{\alpha}(1-\frac{\alpha}{2})\int_{0}^{T} \mathbb{P}(\tau_{e} > s)ds \leqslant ||x||_{H}^{2-\alpha}.$$

This implies that

$$\left(\delta\rho^{\alpha}(1-\frac{\alpha}{2})\right)\mathbb{P}(\tau_{e}>T)T\leqslant\delta\rho^{\alpha}(1-\frac{\alpha}{2})\int_{0}^{T}\mathbb{P}(\tau_{e}>s)ds\leqslant\|x\|_{H}^{2-\alpha}.$$

Then, we deduce that

$$\mathbb{P}(\tau_e > T) \leqslant \left\|x\right\|_H^{2-\alpha} / \left(\delta\rho^{\alpha}(1-\frac{\alpha}{2})\right)T.$$

Thus, (4.3) follows.

Finally, letting $T \to \infty$, it follows that

$$\mathbb{P}(\tau_e < \infty) = 1$$

Thus, we complete the proof.

37

YOSIDA APPROXIMATION AND MULTI-VALUED SEI

5. Examples/Applications

In this section, we illustrate our general results to some examples of multi-valued stochastic evolution inclusions, e.g., multi-valued stochastic porous media equations, multivalued stochastic Φ -Laplacian equations and stochastic differential inclusions involving subdifferentials.

5.1. Multi-valued stochastic porous media equations. The porous media equation models the flow and diffusion of fluids, such as liquids or gases, through porous materials. It has applications in various fields, including gas flow, heat transfer, and groundwater flow (cf. [11]). In this context, we investigate the well-posedness and the finite time extinction of multi-valued stochastic porous media equations.

Let $\Lambda \subset \mathbb{R}^d$ be an open and bounded domain with $d \ge 3$ and $p > \frac{2d}{d+2}$. We consider the multi-valued stochastic porous media equations as follows

$$\begin{cases} dX(t) \in \Delta \Psi(X(t))dt + \sigma(t, X(t))dW(t), \\ X|_{\partial \Lambda} = 0, X(0) = x, \end{cases}$$
(5.1)

where W(t) is a cylindrical Wiener process defined on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \ge 0}, \mathbb{P})$ taking values in U.

Let $H_0^1(\Lambda) := H_0^{1,2}(\Lambda)$ and $H^{-1}(\Lambda)$ denote the dual space of $H_0^1(\Lambda)$. If $d \ge 3$, by the classical Sobolev embedding theorem, we have $H_0^1(\Lambda) \subset L^{\frac{2d}{d-2}}(\Lambda)$ continuously and densely. Since $\frac{p}{p-1} < \frac{2d}{d-2}$, it follows that $L^p(\Lambda) = (L^{\frac{p}{p-1}}(\Lambda))^* \subset (H_0^1(\Lambda))^* = H^{-1}(\Lambda)$ continuously and densely.

By Lemma 4.1.12 in [34], the map $(-\Delta)^{-1} : H^{-1}(\Lambda) \to H^1_0(\Lambda)$ is the Riesz isomorphism on $H^{-1}(\Lambda)$. Thus, we can identify $H^{-1}(\Lambda)$ with its dual $(H^{-1}(\Lambda))^* = H^1_0(\Lambda)$ by the Riesz map $(-\Delta)^{-1}$.

Define $V := L^p(\Lambda), H := H^{-1}(\Lambda)$ and $V^* := (L^p(\Lambda))^*$, we have the Gelfand triple $V \subset H \subset V^*$,

and the embedding $V \subset H$ is compact.

Lemma 5.1. (cf. [34, Lemma 4.1.13]) The map

$$\Delta: H^1_0(\Lambda) \to (L^p(\Lambda))^*$$

extends to a linear isometry

$$\Delta: L^{\frac{p}{p-1}}(\Lambda) \to (L^p(\Lambda))^* = V^*$$

and for all $u \in L^{\frac{p}{p-1}}(\Lambda), v \in L^{p}(\Lambda)$,

$$_{V^*}\langle -\Delta u, v \rangle_V =_{L^{\frac{p}{p-1}}} \langle u, v \rangle_{L^p} = \int_{\Lambda} u(\xi)v(\xi)d\xi.$$
(5.2)

Let $\Psi : \mathbb{R} \to 2^{\mathbb{R}}$ be a function satisfying the following assumptions

- $(\Psi 1) \Psi$ is maximal-monotone.
- (Ψ 2) For some $p \in (1, \infty)$, there exist $c_1 > 0, c_2 \ge 0$ such that for every $x \in \Psi(s)$,

$$s \cdot x \geqslant c_1 |s|^p - c_2$$

(Ψ 3) There exists c > 0 such that for every $x \in \Psi(s)$ and every $s \in \mathbb{R}$,

$$|x| \leqslant c|s|^{p-1} + c$$

Here are some examples from physics models that satisfy $(\Psi 1)$ - $(\Psi 3)$.

Example 5.1. The stochastic BTW model and the Zhang model play significant roles in the study of self-organized criticality. Both can be described as multi-valued stochastic porous media equations involving the following operator

$$\Psi(s) = \begin{cases} 1 + \delta s, & \text{if } s > 0\\ [0, 1], & \text{if } s = 0,\\ 0, & \text{if } s < 0, \end{cases}$$

with $\delta = 0$ in the BTW model (cf. [2]) and $\delta > 0$ in the Zhang model (cf. [62]).

In this work, for p > 1 and $\nu > 0$, we can consider the function

$$\Psi(s) = sign(s)(\nu + |s|^{p-1}), \tag{5.3}$$

which satisfies $(\Psi 1)$ - $(\Psi 3)$. Here, the sign function is defined as follows

$$sign(s) = \begin{cases} \frac{s}{|s|}, & \text{if } s \neq 0, \\ [-1,1], & \text{if } s = 0. \end{cases}$$
(5.4)

For p = 2, (5.3) can be seen as a modification of the Zhang model. Moreover, we also extend the well-posedness result established in [33] for $p \in (1, 2]$ to the case p > 1.

Example 5.2. For p > 1 and $\rho > 0$, we consider the function

$$\Psi(s) = \rho \cdot \operatorname{sign}(s)|s|^{p-1} + \tilde{\Psi}(s),$$

where $\tilde{\Psi}$ is a continuous monotonically nondecreasing function with growth of order p-1. Our results extend those in [7] from $p \in (1,2)$ to p > 1, which can be applied to the slow diffusion models for p > 2 and the fast diffusion models for $1 (in particular, the plasma fast diffusion model if <math>p = \frac{3}{2}$).

Example 5.3. For $\rho > 0$, we consider the function

$$\Psi(s) = \rho \cdot \operatorname{sign}(s) + \Psi(s),$$

where $\tilde{\Psi} \in C^1(\mathbb{R}\setminus\{0\})$ and there exist constants $\delta, c > 0$ and p > 1 such that $\tilde{\Psi}'(s) \ge \delta |s|^{p-2}$ and $|\tilde{\Psi}(s)| \le c|s|^{p-1} + c$ for all $s \in \mathbb{R}\setminus\{0\}$. For every $x \in \tilde{\Psi}(0)$, assume that there exists a constant c > 0 such that $|x| \le c$ and that condition (Ψ 1) holds. Then, conditions (Ψ 1)-(Ψ 3) are satisfied. The authors [6] investigated the case where p = 2 in this example and also pointed out that the finite time extinction of solutions to this example exhibits self-organized criticality behavior.

Theorem 5.1. Assume that Ψ satisfies $(\Psi 1)$ - $(\Psi 3)$ and that (\mathbf{H}_{σ}) holds. Then for any initial value $x \in H$, the multi-valued stochastic porous media equation (5.1) has a unique

(probabilistically) strong solution. In addition, for any $q \ge 2$, we have the following estimate

$$\mathbb{E}\left[\sup_{t\in[0,T]}\|X(t)\|_{H}^{q}\right] + \mathbb{E}\left\{\left(\int_{0}^{T}\|X(s)\|_{V}^{p}ds\right)^{\frac{q}{2}}\right\} < \infty$$

Furthermore, let X(t, x) be the unique solution of Eq. (5.1) with the initial value x, $\{x_n\}_{n \in \mathbb{N}}$ and x be a sequence with $||x_n - x||_H \to 0$. Then for any q > 0,

$$\lim_{n \to \infty} \mathbb{E} \Big[\sup_{t \in [0,T]} \| X(t,x_n) - X(t,x) \|_H^q \Big] = 0.$$

Proof. We define the multi-valued porous media operator $\mathcal{A}: V \to 2^{V^*}$ by

$$\mathcal{A}(u) := -\Delta \Psi(u), \quad u \in L^p(\Lambda).$$

According to condition (Ψ 3), we know that $\Psi(u) \in L^{\frac{p}{p-1}}(\Lambda)$ for any $u \in L^{p}(\Lambda)$. Therefore, the operator \mathcal{A} is well-defined by Lemma 5.1.

In view of (5.2) and condition (Ψ 2), we have that for any $u \in L^p(\Lambda)$ and $v \in \Psi(u)$,

$$\langle -\Delta v, u \rangle = \int_{\Lambda} v(\xi) u(\xi) d\xi \ge c_1 \int_{\Lambda} |u(\xi)|^p d\xi - c_2 |\Lambda| = c_1 ||u||_V^p - C.$$

Therefore, the multi-valued porous media operator \mathcal{A} satisfies $(\mathbf{H}_{\mathcal{A}}^2)$ with $\delta := c_1$ and $\alpha = p$. In addition, let $u \in L^p(\Lambda)$ and $v \in \Psi(u)$ be such that $-\Delta v = \mathcal{A}^0(u)$. According to Lemma 5.1 and condition (Ψ 3), we deduce that

$$\|\mathcal{A}^{0}(u)\|_{V^{*}} \leq C \|u\|_{V}^{p-1} + C.$$

Hence, $(\mathbf{H}_{\mathcal{A}}^3)$ holds for the multi-valued porous media operator \mathcal{A} . By the proof in Section 6 of [33], we know that $-\Delta\Psi(\cdot)$ is maximal-monotone.

Since \mathcal{A} satisfies $(\mathbf{H}^1_{\mathcal{A}})$ - $(\mathbf{H}^3_{\mathcal{A}})$, B = 0 and σ satisfies (\mathbf{H}_{σ}) , we can directly obtain the desired results by applying Theorem 3.3 and Theorem 3.4.

Next, let us focus on the finite time extinction of the multi-valued stochastic porous media equations perturbed by linear multiplicative noise, i.e.,

$$\begin{cases} dX(t) \in \Delta \Psi(X(t))dt + \sum_{k=1}^{\infty} h_k(t)X(t)d\beta_k(t), \\ X|_{\partial \Lambda} = 0, X(0) = x, \end{cases}$$
(5.5)

where $\{\beta_k\}$ is a sequence of standard Brownian motions on a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \ge 0}, \mathbb{P})$ and $h(t) := \sum_{k=1}^{\infty} |h_k(t)|^2 \in L^1([0, \infty), [0, \infty)).$

Theorem 5.2. Assume that $(\Psi 1)$ - $(\Psi 3)$ hold for Eq. (5.5) with $p \in (1,2)$ and $c_2 = 0$ in $(\Psi 2)$. For any $x \in H$ and $t \ge \tau_e$, where τ_e is the extinction time, we have

$$||X(t)||_{H} = 0, \mathbb{P}$$
-a.s.,

and there is a constant $c^* > 0$ such that for any T > 0,

$$\mathbb{P}(\tau_e \leqslant T) \ge 1 - \frac{c^* \|x\|_H^{2-\alpha}}{T}$$

In particular, we have

$$\mathbb{P}(\tau_e < \infty) = 1.$$

Remark 5.3. We point out that if d = 1, 2, one could apply stronger embedding from Theorems 7.10 and 7.15 in [22] to obtain the well-posedness and the finite time extinction for all p > 1 and $p \in (1, 2)$, respectively.

5.2. Multi-valued stochastic Φ -Laplacian equations. The Φ -Laplacian equations, which generalize the *p*-Laplacian equations, were studied by Gess and Tölle [21] under the weak coercivity condition for $p \in [1, 2)$. In this subsection, we focus on the well-posedness and finite time extinction of multi-valued stochastic Φ -Laplacian equations for p > 1. These include the (singular/degenerate) *p*-Laplacian equations and the models of non-Newtonian fluids governed by the Φ -Laplacian operator.

Let $\Lambda \subset \mathbb{R}^d$ be an open and bounded domain, and either p > 1 if d = 1, 2 or $p > \frac{2d}{d+2}$ if $d \ge 3$. Considering the following multi-valued stochastic Φ -Laplacian equations

$$\begin{cases} dX(t) \in \operatorname{div}\Phi(\nabla X(t))dt + \sigma(t, X(t))dW(t), \\ X|_{\partial\Lambda} = 0, X(0) = x. \end{cases}$$
(5.6)

Define $V := W_0^{1,p}(\Lambda), H := L^2(\Lambda)$ and $V^* = (W_0^{1,p}(\Lambda))^*$. Then, we have the Gelfand triple

$$V \subset H \subset V^*$$

compactly and densely.

Let $\Phi : \mathbb{R} \to 2^{\mathbb{R}}$ be a function satisfying the following conditions.

(Φ 1) Φ is maximal-monotone.

($\Phi 2$) For $p \in (1, \infty)$, there exist $c_1 > 0, c_2 \ge 0$ such that for any $x \in \Phi(s)$,

$$s \cdot x \geqslant c_1 |s|^p - c_2$$

(Φ 3) There exist c > 0 such that for any $x \in \Phi(s)$ and $s \in \mathbb{R}$,

$$|x| \leqslant c|s|^{p-1} + c$$

Example 5.4. (i) p-Laplacian equations: For p > 1, the operator

$$\Phi(s) = |s|^{p-1} \operatorname{sign}(s)$$

satisfies $(\Phi 1)$ - $(\Phi 3)$ with $c_1 = 1$ and $c_2 = 0$, where the sign function is defined in (5.4).

(ii) Non-Newtonian fluid models: For p > 1, the operator

$$\Phi(s) = (1+s^2)^{(p-2)/2}s$$

satisfies $(\Phi 1)$ - $(\Phi 3)$, which has been considered in [52].

Theorem 5.4. Assume that (Φ_1) - (Φ_3) and (\mathbf{H}_{σ}) holds. Then for any initial value $x \in H$, the multi-valued stochastic Φ -Laplacian equations (5.6) have a unique (probabilistically) strong solution. In addition, for any $q \ge 2$, we have the following estimate

$$\mathbb{E}\bigg[\sup_{t\in[0,T]}\|X(t)\|_{H}^{q}\bigg] + \mathbb{E}\left\{\left(\int_{0}^{T}\|X(s)\|_{V}^{p}ds\right)^{\frac{q}{2}}\right\} < \infty.$$

Furthermore, let X(t, x) be the unique solution of Eq. (5.6) with the initial value x, $\{x_n\}_{n \in \mathbb{N}}$ and x be a sequence with $||x_n - x||_H \to 0$. Then for any q > 0,

$$\lim_{n \to \infty} \mathbb{E} \Big[\sup_{t \in [0,T]} \| X(t,x_n) - X(t,x) \|_H^q \Big] = 0.$$

Proof. Define the Φ -Laplace operator $\mathcal{A}: V \to 2^{V^*}$ by

$$\mathcal{A}(u) := -\operatorname{div}\Phi(\nabla u), \quad u \in W_0^{1,p}(\Lambda),$$

more precisely, given $u \in W_0^{1,p}(\Lambda)$, for any $\tilde{v} \in \Phi(\nabla u)$, we define $v := -\operatorname{div}(\tilde{v}) \in \mathcal{A}(u)$ such that

$$\langle v, w \rangle := \int_{\Lambda} \tilde{v}(\xi) \nabla w(\xi) d\xi \text{ for all } w \in W_0^{1,p}(\Lambda).$$
 (5.7)

In view of condition (Φ 3), for any $u \in H_0^{1,p}(\Lambda)$ we note that each $v \in \mathcal{A}(u)$ is a well-defined element on $(H_0^{1,p}(\Lambda))^*$, and

$$||v||_{V^*} \leqslant C ||u||_V^{p-1} + C$$

Thus, $(\mathbf{H}^3_{\mathcal{A}})$ holds for the multi-valued Φ -Laplace operator \mathcal{A} with $\alpha = p$.

According to condition ($\Phi 2$), we deduce that for any $u \in H_0^{1,p}(\Lambda)$,

$$\langle -\operatorname{div}\Phi(\nabla u), u \rangle = \int_{\Lambda} \Phi(\nabla u(\xi)) \nabla u(\xi) d\xi \ge c_1 \int_{\Lambda} |\nabla u(\xi)|^p d\xi - c_2 |\Lambda| \ge c_1 ||u||_V^p - C,$$

then the operator \mathcal{A} satisfies $(\mathbf{H}_{\mathcal{A}}^2)$ with $\delta := c_1$ and $\alpha = p$.

Next, we prove that \mathcal{A} is maximal-monotone. Let $x, y \in V, \tilde{v} \in \Phi(\nabla x), \tilde{u} \in \Phi(\nabla y)$ and $v := -\operatorname{div}(\tilde{v}), u := -\operatorname{div}(\tilde{u})$. By condition (Φ 1), we have

$$\langle v - u, x - y \rangle = \int_{\Lambda} (\tilde{v}(\xi) - \tilde{u}(\xi)) (\nabla x(\xi) - \nabla y(\xi)) d\xi \ge 0.$$

This shows that \mathcal{A} is monotone.

In order to get the maximal monotonicity \mathcal{A} , according to Theorem 6.1, it is sufficient to show that if for any $y \in V^*$, there exists $x \in V$ such that

$$J(x) - \operatorname{div}(\tilde{v}) = y, \tag{5.8}$$

where $\tilde{v} \in \Phi(\nabla x)$ and J is the duality mapping from V to V^* .

We begin by considering the approximating equation

$$J(x) - \operatorname{div}\Phi_{\lambda}(\nabla x) = y, \qquad (5.9)$$

where $\Phi_{\lambda} : \mathbb{R} \to \mathbb{R}$ is the generalized Yosida approximation of Φ defined by (2.5).

Let $u_n \to u$ in V, then we have $\nabla u_n \to \nabla u$ in $L^p(\Lambda)$. By the demicontinuity of Φ_{λ} , we deduce that $\Phi_{\lambda}(\nabla u_n) \rightharpoonup \Phi_{\lambda}(\nabla u)$ in $L^{\frac{p}{p-1}}(\Lambda)$. Therefore, for any $w \in V$, we have

$$\lim_{n \to \infty} \langle -\operatorname{div} \Phi_{\lambda}(\nabla u_n) + \operatorname{div} \Phi_{\lambda}(\nabla u), w \rangle = \lim_{n \to \infty} \lim_{L^{\frac{p}{p-1}}} \langle \Phi_{\lambda}(\nabla u_n) - \Phi_{\lambda}(\nabla u), \nabla w \rangle_{L^p} = 0.$$

This shows that $-\operatorname{div}\Phi_{\lambda}(\nabla \cdot)$ is also demicontinuous.

Next, since $-\operatorname{div}\Phi_{\lambda}(\nabla \cdot)$ is monotone and demicontinuous, and J is maximal-monotone, we conclude that the operator $x \mapsto J(x) - \operatorname{div}\Phi_{\lambda}(\nabla x)$ is maximal-monotone. According

to condition ($\Phi 2$) and Lemma 6.1, there exist constants $c_1 > 0, c_2 \ge 0$ such that for any $0 < \lambda < c_1^{-1}, x \in \Phi_{\lambda}(s)$ and $s \in \mathbb{R}$, we have

$$s \cdot x \geqslant c_1 2^{-p} |s|^p - c_2.$$

Hence, for any $u \in V$, we obtain

$$\langle -\operatorname{div}\Phi_{\lambda}(\nabla u), u \rangle = \int_{\Lambda} \Phi_{\lambda}(\nabla u(\xi)) \cdot \nabla u(\xi) d\xi \ge c_1 2^{-p} \|u\|_V^p - C.$$
(5.10)

This shows that $-\operatorname{div}\Phi_{\lambda}(\nabla \cdot)$ is coercive. Then, it is clear that $J - \operatorname{div}\Phi_{\lambda}(\nabla \cdot)$ is also coercive. Finally, by Corollary 2.2 in [5] there exists a solution x_{λ} to (5.9).

By (5.10), we can deduce that

$$c_1 2^{-p} \|x_\lambda\|_V^p \leqslant \langle -\operatorname{div} \Phi_\lambda(\nabla x_\lambda), x_\lambda \rangle + C$$

= $\langle y - J(x_\lambda), x_\lambda \rangle + C$
 $\leqslant c_1 2^{-(p+1)} \|x_\lambda\|_V^p + C(\|y\|_{V^*}^{\frac{p}{p-1}} + 1).$

Therefore, for any $0 < \lambda < c_1^{-1}$, we have

$$\sup_{\lambda} \|x_{\lambda}\|_{V} < \infty.$$
(5.11)

Denote the operator $\mathcal{A}_{\lambda}(\cdot) := -\operatorname{div}\Phi_{\lambda}(\nabla \cdot)$, it follows from Proposition 2.3 (ii) and condition (Φ 3) that

$$\left\|\Phi_{\lambda}(\nabla x_{\lambda})\right\|_{L^{\frac{p}{p-1}}(\Lambda)} \leqslant C(\left\|x_{\lambda}\right\|_{V}^{p}+1)$$
(5.12)

and

$$\begin{aligned} \|\mathcal{A}_{\lambda}(x_{\lambda})\|_{V^{*}} &= \sup_{\|x\|_{V}=1} |\langle -\operatorname{div}\Phi_{\lambda}(\nabla x_{\lambda}), x\rangle| \\ &\leqslant \sup_{\|x\|_{V}=1} \int_{\Lambda} |\Phi_{\lambda}(\nabla x_{\lambda}(\xi))| \cdot |\nabla x(\xi)| d\xi \\ &\leqslant C(\|x_{\lambda}\|_{V}^{p}+1). \end{aligned}$$

By the definition of J, we can conclude that for any $0 < \lambda < c_1^{-1}$, the following estimate holds

$$\sup_{\lambda} \left(\|\Phi_{\lambda}(\nabla x_{\lambda})\|_{L^{\frac{p}{p-1}}(\Lambda)} + \|\mathcal{A}_{\lambda}(x_{\lambda})\|_{V^*} + \|J(x_{\lambda})\|_{V^*} \right) < \infty.$$
(5.13)

Hence, in terms of (5.11)-(5.13), there exists a subsequence λ_k of λ ($0 < \lambda < c_1^{-1}$), we denote x_{λ_k} by x_k , such that as $k \to \infty$,

$$x_k \rightharpoonup x \text{ in } V, \ \Phi_{\lambda_k}(\nabla x_k) \rightharpoonup x_{\Phi} \text{ in } L^{\frac{p}{p-1}}(\Lambda), \ \mathcal{A}_{\lambda_k}(x_k) \rightharpoonup x_{\mathcal{A}}, J(x_k) \rightharpoonup x_J \text{ in } V^*.$$

By (5.9), it implies that for any $m, n \in \mathbb{N}$,

$$\langle J(x_n) - J(x_m), x_n - x_m \rangle + \langle \mathcal{A}_{\lambda_n}(x_n) - \mathcal{A}_{\lambda_m}(x_m), x_n - x_m \rangle = \langle y - y, x_n - x_m \rangle = 0.$$

Since J is monotone, we have

$$\limsup_{n,m\to\infty} \langle \mathcal{A}_{\lambda_n}(x_n) - \mathcal{A}_{\lambda_m}(x_m), x_n - x_m \rangle \leqslant 0.$$

This implies that

$$\lim_{n,m\to\infty} \sup_{\Lambda} \int_{\Lambda} (\Phi_{\lambda_n}(\nabla x_n(\xi)) - \Phi_{\lambda_m}(\nabla x_m(\xi))) (\nabla x_n(\xi) - \nabla x_m(\xi)) d\xi \leq 0.$$
(5.14)

In view of $x_k \to x$ in V, we have $\nabla x_k \to \nabla x$ in $L^p(\Lambda)$. Since Φ is maximal-monotone, we know that $\Phi_{\lambda_k}(\nabla x_k) \in \Phi(\mathcal{R}_{\lambda_k}(\nabla x_k))$ and $\|\mathcal{R}_{\lambda_k}(\nabla x_k) - \nabla x_k\|_{L^p(\Lambda)} \to 0$, as $k \to \infty$. By (5.14) and Lemma 2.1, we deduce that $x_{\Phi} \in \Phi(\nabla x)$. Then, according to (5.7), we have that for all $w \in V$,

$$\lim_{k \to \infty} \langle \mathcal{A}_{\lambda_k}(x_k) + \operatorname{div}(x_{\Phi}), w \rangle = \lim_{k \to \infty} \sum_{L^{\frac{p}{p-1}}} \langle \Phi_{\lambda_k}(\nabla x_k) - x_{\Phi}, \nabla w \rangle_{L^p} = 0.$$

This implies that $\mathcal{A}_{\lambda_n}(x_k) \rightarrow -\operatorname{div}(x_{\Phi}) \in \mathcal{A}(x)$. Consequently,

$$\limsup_{n,m\to\infty} \langle J(x_n) - J(x_m), x_n - x_m \rangle \leqslant 0,$$

which implies that $x_J = J(x)$. Thus, we obtain that $x \in V$ is the solution of (5.8). We complete the proof of (5.8).

Since \mathcal{A} satisfies $(\mathbf{H}^1_{\mathcal{A}})$ - $(\mathbf{H}^3_{\mathcal{A}})$, B = 0 and σ satisfies (\mathbf{H}_{σ}) , the conclusion follows directly from Theorem 3.3 and Theorem 3.4.

We consider the finite time extinction for the following multi-valued stochastic Φ -Laplacian equations perturbed by linear multiplicative noise

$$\begin{cases} dX(t) \in \operatorname{div}\Phi(\nabla X(t))dt + \sum_{k=1}^{\infty} h_k(t)X(t)d\beta_k(t), \\ X|_{\partial\Lambda} = 0, X(0) = x, \end{cases}$$
(5.15)

where $h(t) := \sum_{k=1}^{\infty} |h_k(t)|^2 \in L^1([0,\infty), [0,\infty)).$

Theorem 5.5. Assume that $(\Phi 1)$ - $(\Phi 3)$ hold for Eq. (5.15) with $p \in (1, 2)$ and $c_2 = 0$ in $(\Phi 2)$. For any $x \in H$ and $t \ge \tau_e$, we have

$$||X(t)||_{H} = 0, \mathbb{P}$$
-a.s.,

and there is a constant $c^* > 0$ such that for any T > 0,

$$\mathbb{P}(\tau_e \leqslant T) \ge 1 - \frac{c^* ||x||_H^{2-\alpha}}{T}.$$

In particular, we have

$$\mathbb{P}(\tau_e < \infty) = 1.$$

Remark 5.6. To the best of our knowledge, this is the first result in the literature concerning the finite time extinction of solutions to the multi-valued stochastic Φ -Laplacian equations, which is of independent interest. 5.3. SEIs with subdifferentials. Evolution inclusions with subdifferentials were first introduced by Brézis in [13], originated from simplified models employed in the description of porous medium combustion, chemical reactor theory and game theory (cf. [38]). These evolution inclusions have contributed to the theory of nonlinear evolution equations (cf. [43]). Further research on evolution inclusions with subdifferentials has been conducted in [1, 26, 36, 37]. In this part, we intend to present a more general result for the existence of weak solutions to a class of SEIs involving subdifferentials with single-valued pseudo-monotone operators.

Let K be a uniformly convex Banach space and $\varphi : K \to \overline{\mathbb{R}} := (-\infty, \infty]$ be a lower semicontinuous, proper and convex function on K. Let $V \subset K$ be a domain of $\partial \varphi$, where the mapping $\partial \varphi : K \to 2^{K^*}$ defined by

$$\partial \varphi(v) := \left\{ v^* \in K^* \mid \varphi(v) \leqslant \varphi(w) + \langle v^*, v - w \rangle, \forall w \in K \right\}$$

is called the subdifferential of φ . We denote $\mathcal{D}(\partial \varphi)$ as the set of all $x \in K$ for which $\partial \varphi(x) \neq \emptyset$.

Proposition 5.1. (cf. Theorem 2.8 in [5]) For a lower semicontinuous, proper and convex function φ , the subdifferential $\partial \varphi : K \to 2^{K^*}$ is maximal-monotone.

Let $\Lambda \subset \mathbb{R}^d$ be a bounded domain with smooth boundary $\partial \Lambda$. We consider the following multi-valued stochastic partial differential equation:

$$\begin{cases} \partial_t u(t,x) \in [\Delta u(t,x) - g(t,x,u(t,x),\nabla u(t,x)) - \partial \varphi(u(t,x))] \, dt \\ + \sigma(t,u(t,x)) dW(t), \\ u(t,\cdot)|_{\partial \Lambda} = 0, \ u(0,x) = u_0(x). \end{cases}$$
(5.16)

We define $H := L^2(\Lambda)$ and $V := W_0^{1,2}(\Lambda)$. This forms the Gelfand triple

$$V \subset H \subset V^*.$$

The space V is a uniformly convex Banach space and the embedding $V \subset H$ is compact.

We assume that φ is a lower semicontinuous, proper and convex function with $\varphi(0) = 0$, and that $\mathcal{D}(\partial \varphi) = V$. Additionally, there exist constants $c_1 > 0, c_2 \ge 0$ such that

$$\varphi(u) \ge c_1 \|u\|_V^2 - c_2 \text{ for all } u \in V,$$

and a constant c > 0 such that

$$|\varphi(u) - \varphi(u-z)| \leq c_3 ||u||_V + c_4 \text{ for all } u, z \in V \text{ with } ||z||_V = 1.$$

We further assume that g satisfies the following conditions:

(i) g satisfies the Carathéodory conditions: for a.e. $(t,x) \in [0,T] \times \Lambda$, g(t,x,u,z) is continuous in $(u,z) \in \mathbb{R} \times \mathbb{R}^d$, and for each $(u,z) \in \mathbb{R} \times \mathbb{R}^d$, g(t,x,u,z) is measurable with respect to $(t,x) \in [0,T] \times \Lambda$.

(ii) There exist a constant c > 0 and a function $f_1 \in L^1([0,T] \times \Lambda, [0,\infty))$ such that for a.e. $(t,x) \in [0,T] \times \Lambda$ and all $(u,z) \in \mathbb{R} \times \mathbb{R}^d$,

$$g(t, x, u, z)u \ge -c|u|^2 - f_1(t, x).$$

(iii) There exist a nonnegative constant c and a function $f_2 \in L^2([0,T] \times \Lambda, [0,\infty))$ such that for a.e. $(t, x) \in [0, T] \times \Lambda$ and all $(u, z) \in \mathbb{R} \times \mathbb{R}^d$,

$$|g(t, x, u, z)| \leq c|z| + c|u|^{\frac{d+2}{d}} + f_2(t, x).$$

Theorem 5.7. Suppose that φ and g satisfy the above assumptions and that (\mathbf{H}_{σ}) holds. Then for any initial value $u_0 \in H$, the multi-valued stochastic evolution inclusions (5.16) have a (probabilistically) weak solution. Moreover, for any $p \ge 2$, we have the following estimate

$$\mathbb{E}\Big[\sup_{t\in[0,T]}\|X(t)\|_{H}^{p}\Big] + \mathbb{E}\left\{\left(\int_{0}^{T}\|X(s)\|_{V}^{2}ds\right)^{\frac{p}{2}}\right\} < \infty.$$

Proof. We define the operator $\mathcal{A}: V \to 2^{V^*}$ by

$$\mathcal{A}(u) := \partial \varphi(u), \quad u \in W_0^{1,2}(\Lambda),$$

it follows from Proposition 5.3 that \mathcal{A} satisfies the conditions $(\mathbf{H}^1_{\mathcal{A}})$.

By the definition and assumptions of φ , we have that for any $u \in V$ and $u^* \in \mathcal{A}(u)$,

$$\langle u^*, u \rangle \geqslant \varphi(u) - \varphi(0) \geqslant c_1 \|u\|_V^2 - c_2,$$

and

$$||u^*||_{V^*} \leq \sup_{||z||_V=1} |\langle u^*, z \rangle| \leq \sup_{||z||_V=1} |\varphi(u) - \varphi(u-z)| \leq c_3 ||u||_V + c_4.$$

Thus, \mathcal{A} satisfies the conditions $(\mathbf{H}^2_{\mathcal{A}})$ and $(\mathbf{H}^3_{\mathcal{A}})$. Define the operator $B: V \to V^*$ by

$$B(u) := -\Delta u + g(u, \nabla u), \quad u \in W_0^{1,2}(\Lambda).$$

From Example 4.1 in [47], we know that B is pseudo-monotone and satisfies the growth condition (\mathbf{H}_B^2) . We have

$$\langle -\Delta u + g(u, \nabla u), u \rangle \ge -c \|u\|_{H}^{2} + \|u\|_{V}^{2} - \|f(t, \cdot)\|_{L^{1}}$$

Hence, (\mathbf{H}_B^1) holds for *B*. Then, by Theorem 3.1 the result follows.

Remark 5.8. Let $h : \mathbb{R} \to (-\infty, +\infty]$ be a lower semicontinuous, proper and convex function, and $\varphi: L^2(\Lambda) \to (-\infty, +\infty)$ be defined by

$$\varphi(u) := \begin{cases} \int_{\Lambda} h(u(x))dx, & \text{if } h(u) \in L^{1}(\Lambda), \\ +\infty, & \text{otherwise.} \end{cases}$$

By Proposition 2.8 in [3], φ is a lower semicontinuous, proper and convex function and $\mathcal{D}(\partial \varphi) = L^2(\Lambda).$

6. Appendix

In this section, we present some results of the multi-valued maximal-monotone operator.

Proposition 6.1. Let V be a Banach space and the dual space V^* is strictly convex, then the duality mapping $J: V \to V^*$ is single-valued, demicontinuous and odd, the set J(x) is closed and convex for any $x \in V$.

Proof. We first prove that J is single-valued. Let $v_1, v_2 \in J(u)$, we have

$$\langle v_1, u \rangle = \langle v_2, u \rangle = \|v_1\|_{V^*}^{\frac{\alpha}{\alpha-1}} = \|v_2\|_{V^*}^{\frac{\alpha}{\alpha-1}} = \|u\|_V^{\alpha}.$$

Hence, according to (2.1), it follows that

$$2\|v_1\|_{V^*}\|u\|_V \leq \langle v_1 + v_2, u \rangle \leq \|v_1 + v_2\|_{V^*}\|u\|_V \text{ for all } u \in V,$$

which implies that $||v_1||_{V^*} \leq \frac{1}{2} ||v_1 + v_2||_{V^*}$. Since $||v_1||_{V^*} = ||v_2||_{V^*}$ and V^* is strictly convex, we deduce that $v_1 = v_2$.

Next, we show that J is demicontinuous. Let $\{u_n\}_{n\in\mathbb{N}}\in V$ be strongly convergent to u and let u^* be any weak-* limit point of $\{J(u_n)\}_{n\in\mathbb{N}}$. According to the definition of $J(u_n)$, we have

$$||u_n||_V^{\alpha} = ||J(u_n)||_{V^*}^{\frac{\alpha}{\alpha-1}}$$

Thus, by the above conditions and the weak lower semicontinuity of the norm in V^* , we conclude that

$$\begin{aligned} \|u^*\|_{V^*}^{\frac{\alpha}{\alpha-1}} &\leqslant \liminf_{n \to \infty} \|J(u_n)\|_{V^*}^{\frac{\alpha}{\alpha-1}} \leqslant \lim_{n \to \infty} \|u_n\|_V^{\alpha} (= \|u\|_V^{\alpha}) \\ &= \lim_{n \to \infty} \langle J(u_n), u_n \rangle = \langle u^*, u \rangle \leqslant \|u^*\|_{V^*} \|u\|_V. \end{aligned}$$

Then, it follows that

$$\langle u^*, u \rangle = \|u\|_V^{\alpha} = \|u^*\|_{V^*}^{\frac{\alpha}{\alpha-1}}.$$

By the definition of J(u), we deduce that $u^* = J(u)$. Consequently, the sequence $\{J(u_n)\}_{n\in\mathbb{N}}$ converges weak-* to J(u). Therefore, $J: V \to V^*$ is demicontinuous.

According to the definition of J, it implies that

$$\langle J(-u), -u \rangle = \|J(-u)\|_V^{\frac{\alpha}{\alpha-1}} = \|u\|_V^{\alpha} = \langle -J(u), -u \rangle \text{ for all } u \in V.$$

Since J is single-valued, we conclude

$$J(-u) = -J(u),$$

which shows that J is odd.

Moreover, since J is maximal-monotone, we note that J(x) is closed and convex for any $x \in V$ by Proposition 1.1 in [4]. We complete the proof.

Theorem 6.1. Let V and V^{*} be reflexive and strictly convex Banach spaces. Let $\mathcal{A} : V \to 2^{V^*}$ be a monotone operator and $J : V \to V^*$ denote the duality mapping of V. Then \mathcal{A} is maximal-monotone if and only if for any (or some) $\lambda > 0$, the range condition

$$\mathcal{R}(\mathcal{A} + \lambda J) = V$$

is satisfied.

Proof. Step 1: Suppose that $\mathcal{R}(\mathcal{A} + \lambda J) = V^*$ for some $\lambda > 0$. If \mathcal{A} is not maximalmonotone, then there exists $[x_1, y_1] \in V \times V^*$ such that $[x_1, y_1] \notin \mathcal{G}(\mathcal{A})$ and

$$\langle y - y_1, x - x_1 \rangle \ge 0, \quad \forall [x, y] \in \mathcal{G}(\mathcal{A}).$$
 (6.1)

By assumption, we can choose $[x_2, y_2] \in \mathcal{G}(\mathcal{A})$ such that

$$\lambda J(x_2) + y_2 = \lambda J(x_1) + y_1.$$

Then in view of (6.1), we can deduce that

$$\langle J(x_2) - J(x_1), x_2 - x_1 \rangle \leqslant 0.$$

According to the definition of J, we obtain

$$||x_1||_V^{\alpha} + ||x_2||_V^{\alpha} \leq \langle J(x_2), x_1 \rangle + \langle J(x_1), x_2 \rangle \leq ||x_2||^{\alpha - 1} ||x_1||_V + ||x_1||^{\alpha - 1} ||x_2||_V,$$

which implies

$$(\|x_1\|_V^{\alpha-1} - \|x_2\|_V^{\alpha-1})(\|x_1\|_V - \|x_2\|_V) \le 0.$$

In light of $\alpha > 1$, we deduce that

$$\langle J(x_2), x_1 \rangle = \langle J(x_1), x_2 \rangle = ||x_1||_V^{\alpha} = ||x_2||_V^{\alpha}$$

Thus, it follows that $J(x_1) = J(x_2)$. Since the duality mapping J^{-1} of V^* is single-valued, it implies that $x_1 = x_2$. Then it is clear that $[x_1, y_1] = [x_2, y_2] \in \mathcal{G}(\mathcal{A})$, which contradicts to the hypothesis. Thus, \mathcal{A} is maximal-monotone,

Step 2: Suppose that \mathcal{A} is maximal-monotone. Let y be an arbitrary element of V^* and let $\lambda > 0$. Define

$$B(u) := \lambda J(u) - y, \quad \forall u \in V.$$

Since

$$\lim_{\|u\|_V \to \infty} \frac{\langle B(u), u \rangle}{\|u\|_V} \ge \lim_{\|u\|_V \to \infty} (\lambda \|u\|_V^{\alpha-1} - \|y\|_{V^*}) = \infty,$$

we note that B is coercive. From the definition of J, it follows that B is monotone, singlevalued and hemicontinuous. Then, according to Theorem 2.1 in [5], there exists $x \in V$ such that

$$\langle \lambda J(x) - y + v, u - x \rangle \ge 0, \quad \forall [u, v] \in \mathcal{G}(\mathcal{A}).$$

In view of the maximal monotonicity of \mathcal{A} , we deduce that $[x, -\lambda J(x) + y] \in \mathcal{G}(\mathcal{A})$, which implies that $y \in \lambda J(x) + \mathcal{A}(x)$. We complete the proof.

Lemma 6.1. Let $\alpha > 1$ and $\mathcal{A} : V \to 2^{V^*}$ be a maximal-monotone operator. Let \mathcal{A}_{λ} denote its generalized Yosida approximation (i.e. (2.5)). Suppose that there exist $\delta > 0$ and $C \in \mathbb{R}$ such that

$$\langle v, x \rangle \ge \delta \|x\|_V^{\alpha} + C \quad \text{for all } x \in \mathcal{D}(\mathcal{A}) \text{ and } v \in \mathcal{A}(x).$$
 (6.2)

Then, for any $0 < \lambda < \delta^{-1}$, we have

$$\langle \mathcal{A}_{\lambda} x, x \rangle \ge \delta 2^{-\alpha} \|x\|_V^{\alpha} + C \quad for \ all \ x \in V.$$

Proof. For any $x \in V$, by the definition of \mathcal{A}_{λ} and J, we have

$$\langle \mathcal{A}_{\lambda}x, x - \mathcal{R}_{\lambda}x \rangle = \frac{1}{\lambda} \langle J(x - \mathcal{R}_{\lambda}x), x - \mathcal{R}_{\lambda}x \rangle = \frac{1}{\lambda} \|x - \mathcal{R}_{\lambda}x\|_{V}^{\alpha}.$$

Since $\mathcal{A}_{\lambda}x \in \mathcal{A}(\mathcal{R}_{\lambda}x)$, by (6.2) it follows that

$$\begin{aligned} \langle \mathcal{A}_{\lambda}x, x \rangle &= \langle \mathcal{A}_{\lambda}x, \mathcal{R}_{\lambda}x \rangle + \frac{1}{\lambda} \|x - \mathcal{R}_{\lambda}x\|_{V}^{\alpha} \\ &\geq \delta \|\mathcal{R}_{\lambda}x\|_{V}^{\alpha} + \frac{1}{\lambda} \|x - \mathcal{R}_{\lambda}x\|_{V}^{\alpha} + C \\ &= \delta \left(\|\mathcal{R}_{\lambda}x\|_{V}^{\alpha} + \|x - \mathcal{R}_{\lambda}x\|_{V}^{\alpha}\right) + \left(\frac{1}{\lambda} - \delta\right) \|x - \mathcal{R}_{\lambda}x\|_{V}^{\alpha} + C. \end{aligned}$$

Noting that $0 < \lambda < \delta^{-1}$, we conclude

$$\langle \mathcal{A}_{\lambda} x, x \rangle \ge \delta 2^{-\alpha} \|x\|_V^{\alpha} + C.$$

We complete the proof.

Data availability Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Statements and Declarations On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- H. Attouch, and A. Damlamian, On multivalued evolution equations in Hilbert spaces, Israel J. Math. 12 (1972), 373-390.
- [2] P. Bak, C. Tang, and K. Wiesenfeld, Self-organized criticality, Phys. Rev. A (3) 38 (1988), 364-374.
- [3] V. Barbu, Nonlinear Semigroups and Differential Equations in Banach Spaces, Editura Academiei Republicii Socialiste România, Bucharest; Noordhoff International Publishing, Leiden, 1976.
- [4] V. Barbu, Analysis and Control of Nonlinear Infinite-Dimensional Systems, Math. Sci. Eng., vol. 190, Academic Press, 1993.
- [5] V. Barbu, Nonlinear Differential Equations of Monotone Types in Banach Spaces, Springer Monogr. Math., Springer, New York, 2010.
- [6] V. Barbu, G. Da Prato, and M. Röckner, Stochastic porous media equations and self-organized criticality, Comm. Math. Phys. 285 (2009), 901-923.
- [7] V. Barbu, G. Da Prato, and M. Röckner, Finite time extinction for solutions to fast diffusion stochastic porous media equations, C. R. Math. Acad. Sci. Paris 347 (2009), 81-84.
- [8] V. Barbu, G. Da Prato, and M. Röckner, Finite time extinction of solutions to fast diffusion equations driven by linear multiplicative noise, J. Math. Anal. Appl. 389 (2012), 147-164.
- [9] V. Barbu, and M. Röckner, Stochastic porous media equations and self-organized criticality: convergence to the critical state in all dimensions, Comm. Math. Phys. 311 (2012), 539-555.
- [10] V. Barbu, M. Röckner, and F. Russo, Stochastic porous media equations in R^d, J. Math. Pures Appl. 103 (2015), 1024-1052.
- [11] J. Boussinesq, Recherches théoriques sur l'écoulement des nappes d'eau infiltrées dans le sol et sur le débit des sources, J. Math. Pures Appl. 10 (1904), 5-78.
- [12] H. Brézis, Équations et inéquations non linéaires dans les espaces vectoriels en dualité, Ann. Inst. Fourier (Grenoble) 18 (1968), 115-175.
- [13] H. Brézis, Opérateurs Maximaux Monotones et Semi-groupes de Contractions dans les Espaces de Hilbert, North-Holland Math. Stud. 5, North-Holland, Amsterdam, London; Elsevier, New York, 1973.
- [14] F.E. Browder, Nonlinear elliptic boundary value problems, Bull. Amer. Math. Soc. 69 (1963), 862-874.

- [15] F.E. Browder, Non-linear equations of evolution, Ann. Math. 80 (1964), 485-523.
- [16] F.E. Browder, Pseudo-monotone operators and nonlinear elliptic boundary value problems on unbounded domains, Proc. Nat. Acad. Sci. 74 (1977), 2659-2661.
- [17] Z. Brzeźniak, E. Hausenblas, and P.A. Razafimandimby, Stochastic reaction-diffusion equations driven by jump processes, Potential Anal. 49 (2018), 131-201.
- [18] R. Cairoli, and R. C. Dalang, Sequential Stochastic Optimization, Wiley, New York, 1996.
- [19] E. Cépa, Problème de Skorohod multivoque, Ann. Probab. 26 (1998), 500-532.
- [20] B. Gess, Finite time extinction for stochastic sign fast diffusion and self-organized criticality, Comm. Math. Phys. 335 (2015), 309-344.
- [21] B. Gess, and J.M. Tölle, Multi-valued, singular stochastic evolution inclusions, J. Math. Pures Appl. 101 (2014), 789-827.
- [22] D. Gilbarg, and N.S. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer, Berlin, 1983.
- [23] I. Gyöngy, On stochastic equations with respect to semimartingales III, Stochastics 7 (1982), 231-254.
- [24] P. Hartman, and G. Stampacchia, On some nonlinear elliptic differential equations, Acta. Math. 115 (1966), 271-310.
- [25] A. Jakubowski, On the Skorokhod topology, Ann. Inst. H. Poincaré Probab. Statist. 22 (1986), 263-285.
- [26] D. Kravvaritis, and N.S. Papageorgiou, Multivalued perturbations of subdifferential type evolution equations in Hilbert spaces, J. Differential Equations 76 (1988), 238-255.
- [27] P. Krée, Diffusion for multivalued stochastic differential equations, J. Funct. Anal. 49 (1982), 73-90.
- [28] N.V. Krylov, and B.L. Rozovskii, Stochastic evolution equations, J. Soviet Math. (1979) 71-147 (in Russian); transl. 16 (1981) 1233-1277.
- [29] J. Leray, and J.L. Lions, Quelques resultats de Visik sur les problèmes elliptiques non linéaires par les méthodes de Minty-Browder, Bull. Soc. Math. France 93 (1965), 97-107.
- [30] P.L. Lions and A.S. Sznitman, Stochastic differential equations with reflecting boundary conditions, Comm. Pure Appl. Math. 37 (1984), 511-537
- [31] W. Liu, and M. Röckner, SPDE in Hilbert space with locally monotone coefficients, J. Funct. Anal. 259 (2010), 2902-2922.
- [32] W. Liu, Existence and uniqueness of solutions to nonlinear evolution equations with locally monotone operators, Nonlinear Anal. 74 (2011), 7543-7561.
- [33] W. Liu, and M. Stephan, Yosida approximations for multivalued stochastic partial differential equations driven by Lévy noise on a Gelfand triple, J. Math. Anal. Appl. 410 (2014), 158-178.
- [34] W. Liu, and M. Röckner, Stochastic Partial Differential Equations: An Introduction, Universitext, Springer, Berlin (2015).
- [35] G.J. Minty, Monotone (non-linear) operators in Hilbert space, Duke. Math. J. 29 (1962), 341-346.
- [36] M. Ôtani, Nonmonotone perturbations for nonlinear parabolic equations associated with subdifferential operators, Cauchy problems, J. Differential Equations 46 (1982), 268-299.
- [37] M. Ötani, Nonmonotone perturbations for nonlinear parabolic equations associated with subdifferential operators, Periodic problems, J. Differential Equations 54 (1984), 248-273.
- [38] M. Otani, and V. Staicu. On some quasilinear parabolic equations with non-monotone multivalued terms, Commun. Pure Appl. Anal. 22 (2023), 1831-1865.
- [39] E. Pardoux, Sur des équations aux dérivées partielles stochastiques monotones, C. R. Acad. Sci. Paris Sér. A-B 275 (1972), A101-A103.
- [40] E. Pardoux, Équations aux dérivées partielles stochastiques non linéaires monotones, Ph.D. thesis, Université Paris XI, 1975.
- [41] R. Pettersson, Yosida approximations for multivalued stochastic differential equations, Stoch. Stoch. Rep. 52 (1995), 107-120.
- [42] C. Prévôt, and M. Röckner, A Concise Course on Stochastic Partial Differential Equations, Lecture Notes in Math. vol. 1905, Springer, Berlin, 2007.
- [43] S. Qin, and X. Xue, Periodic solutions for nonlinear differential inclusions with multivalued perturbations, J. Math. Anal. Appl. 424 (2015), 988-1005.

- [44] J. Ren, M. Röckner, and F.-Y. Wang, Stochastic generalized porous media and fast diffusion equations, J. Differential Equations 238 (2007), 118-152.
- [45] J. Ren, and J. Wu, On regularity of invariant measures of multivalued stochastic differential equations, Stochastic Process. Appl. 122 (2012), 93-105.
- [46] J. Ren, J. Wu, and X. Zhang Exponential ergodicity of non-Lipschitz multivalued stochastic differential equations, Bull. Sci. Math. 134 (2010), 391-404.
- [47] M. Röckner, S. Shang, and T. Zhang, Well-posedness of stochastic partial differential equations with fully local monotone coefficients, Math. Ann. 390 (2024), 3419-3469.
- [48] M. Röckner, and F.-Y. Wang, Non-monotone stochastic generalized porous media equations, J. Differential Equations 245 (2008), 3898-3935.
- [49] M. Röckner, and F.-Y. Wang, General extinction results for stochastic partial differential equations and applications, J. Lond. Math. Soc. 87 (2013), 545-560.
- [50] M. Röckner, and X. Zhang, Stochastic tamed 3D Navier-Stokes equations: existence, uniqueness and ergodicity, Probab. Theory Related Fields 145 (2009), 211-267.
- [51] Y. Saisho, Stochastic differential equations for multidimensional domains with reflecting boundary, Prob. Theory Related Fields 74 (1987), 455-477.
- [52] F. Seib, W. Stannat, and J.M. Tölle, Stability and moment estimates for the stochastic singular Φ-Laplace equation, J. Differential Equations 377 (2023), 663-693.
- [53] A.V. Skorohod, Stochastic equations for diffusions in a bounded region, Theory Prob. Appl. 6 (1961), 264-274.
- [54] M. Stephan, Yosida approximations for multivalued stochastic differential equations on Banach spaces via a Gelfand triple, Ph.D. thesis, Universität Bielefeld, 2012.
- [55] H. Tanaka, Stochastic differential equations with reflecting boundary conditions in convex regions, Hiroshima Math. J. 9 (1979), 163-177.
- [56] D. L. Turcotte, Self-organized criticality, Rep. Progr. Phys. 62 (1999), 1377-1429.
- [57] J. Wu and M. Zhang, On approximations of the Euler-Peano scheme for multivalued stochastic differential equations, Stochastics 91 (2019), 215-234.
- [58] E. Zeidler, Nonlinear Functional Analysis and Its Applications, II/A, Springer-Verlag, New York, 1990.
- [59] E. Zeidler, Nonlinear Functional Analysis and Its Applications, II/B, Springer-Verlag, New York, 1990.
- [60] X. Zhang, Skorohod problem and multivalued stochastic evolution equations in Banach spaces, Bull. Sci. Math. 131 (2007), 175-217.
- [61] X. Zhang, On stochastic evolution equations with non-Lipschitz coefficients, Stoch. Dyn. 9 (2009), 549-595.
- [62] Y. Zhang, Scaling theory of self-organized criticality, Phys. Rev. Lett. 63 (1989), 470-473.