

# Fundamental Origin of Viscosity in 2D Simple Liquids

Dong Huang<sup>1</sup>, Shaoyu Lu<sup>1</sup>, Chen Liang<sup>1</sup>, Matteo Baggioli<sup>2‡</sup>, and Yan Feng<sup>1‡</sup>

<sup>1</sup>Institute of Plasma Physics and Technology, Jiangsu Key Laboratory of Frontier Material Physics and Devices, School of Physical Science and Technology, Soochow University, Suzhou 215006, China

<sup>2</sup>Wilczek Quantum Center, School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China & Shanghai Research Center for Quantum Sciences, Shanghai 201315, China

‡b.matteo@sjtu.edu.cn; fengyan@suda.edu.cn

## ABSTRACT


Shear viscosity plays a fundamental role in liquid dynamics from heavy-ion collisions to biological processes. Still, its physical origin at the individual particle kinetic level remains strongly debated. In this work, we systematically investigate the shear viscosity ( $\eta$ ) of two-dimensional (2D) simple liquids using computer simulations of Lennard-Jones, Yukawa, and one-component plasma systems. By combining Frenkel's liquid description, consisting of solid-like quasi-harmonic vibrations interrupted by thermally activated hops, with the concept of lifetime of local atomic connectivity  $\tau_{LC}$ , we find a surprisingly simple formula for the kinematic viscosity that is solely determined by  $\tau_{LC}$  and the average kinetic particle speed  $\bar{v}_p$ . The derived analytical expression provides a direct link between macroscopic and microscopic dynamics, which shows excellent agreement with the simulation data in all the 2D liquids considered. Moreover, it is discovered that,  $\tau_{LC}$  in 2D liquids is universally determined by the effective potential difference between the first peak and valley of the pair correlation function, implying a direct connection between macroscopic shear transport and microscopic structure. Finally, we demonstrate that the characteristic length scale  $l_p = \bar{v}_p \tau_{LC}$ , which governs the macroscopic shear viscosity, aligns with the elastic length-scale that defines the propagation limit of collective shear waves in liquids. These findings establish that shear viscosity in 2D liquids arises from the diffusive transport of average particle momentum across the elastic length scale. Moreover, they highlight that shear dynamics are fundamentally governed by localized configurational excitations within the atomic connectivity network.

As first described by Newton in his 1687 *Principia*, the shear viscosity ( $\eta$ ) characterizes the momentum flux transport in fluids, defined as the ratio of shear stress to shear rate [1,2] (see Fig. 1(a)). Newton's law of viscosity provides a macroscopic framework that can be translated into an operative definition using the Green-Kubo formalism [3], nevertheless, it does not offer any physical insights about the microscopic and kinetic origin of viscosity.

In gases, a microscopic description of viscosity has been already achieved more than 200 years ago with the formulation of kinetic theory, where momentum transport is accomplished via molecular collisions [4]. This led to the known relation of  $\eta \propto \rho \bar{v}_p l_{\text{mfp}}$  and the typical temperature dependence of  $\eta \propto \sqrt{T}$  in the dilute gas regime, where  $\rho$  is the mass density,  $\bar{v}_p$  is the mean speed of particles, and  $l_{\text{mfp}}$  is the mean free path. Here,  $l_{\text{mfp}}$  is the average distance between two consecutive collisions, which is independent of temperature. However, in liquids, molecular dynamics are inherently collective and cooperative, rendering traditional kinetic theory inapplicable as potential energy contributions play a significant role. This leads to a completely different and much stronger temperature dependence  $\eta \propto \exp(U/T)$ , indicating that viscosity in liquids is governed by the activation energy barrier for molecular rearrangements  $U$ , while not by thermal collisions.

In line with the original picture of liquid dynamics proposed by Frenkel [5], consisting of solid-like quasi-harmonic vibrations in the local basins interrupted by thermally activated hops over potential barriers, numerous studies have attempted a description of liquid viscosity based on activated-rate theory of chemical reactions. Eyring theory [6] is the most famous example of this sort, predicting that  $\eta = A \exp(\Delta G/k_B T)$ . Here,  $A$  is an undetermined pre-factor and  $\Delta G$  is the atomic hopping potential barrier (see Fig. 1(c)). The Eyring equation for viscosity correctly captures the experimentally observed exponential decrease in liquid viscosity with temperature [7]. However, it remains a semi-empirical theory, since both  $A$  and  $\Delta G$  are left undetermined.

On the other hand, in Frenkel's description of liquids dynamics [5], the potential barrier  $\Delta G$  appearing in Eyring's formula is interpreted as the energy necessary to re-arrange the nearest-neighbor cage. The typical timescale of these configurational re-arrangements was formally defined by Egami [8] using the concept of lifetime of local atomic connectivity  $\tau_{LC}$ . By definition,  $\tau_{LC}$  (see Fig. 1(b)) refers to the average time in which one particle loses or gains one neighbor within the surrounding cage. It was realized that above the Arrhenius temperature this timescale coincides with the collective Maxwell relaxation timescale



**Figure 1. Viscosity, particle motion and collective shear dynamics in liquids:** (a) The shear viscosity  $\eta$  determines the macroscopic resistance to shear flow in fluids. (b) A local configurational excitation consisting in losing or gaining one neighbor.  $\tau_{LC}$ , the lifetime of local connectivity, is the average timescale associated to this microscopic process. (c) Structural rearrangements in liquids are governed by localized events in which one or few particles hop a potential barrier ( $\Delta G$ ), as assumed in Eyring and Frenkel theories of liquid viscosity. This activated process happens with an averaged rate  $\tau_F^{-1}$ , where  $\tau_F$  is the microscopic Frenkel time. (d) Collective shear waves in liquids propagate only up to a length-scale  $l \sim 1/k_g$ , with  $k_g$  the wave-vector gap in their dispersion. According to Maxwell viscoelasticity theory,  $k_g \sim 1/(v_T \tau_M)$  where  $v_T$  is the instantaneous speed of propagation for shear waves and  $\tau_M$  is the collective Maxwell relaxation time.

$\tau_M = \eta/G_\infty$  (with  $G_\infty$  the instantaneous shear modulus), providing a microscopic interpretation of structural relaxation and viscosity in high-temperature liquids. This observation suggests a connection between viscosity and local configurational excitations in liquids that was further discussed in [9, 10], revealing surprising similarities with the Drude model for electric transport in metals [11]. By looking at the atomic scale stress correlation function, the relevant length-scale for viscosity was also related to the range of propagation of shear waves [10] (see Fig. 1(d)), providing another link to a modern version of Frenkel's ideas known as  $k$ -gap theory [12, 13], and also to the dual model of liquid viscosity [14].

It is interesting to notice that a microscopic expression for the viscosity in terms of the liquid structure was already proposed by Born and Green using the radial distribution function [15], but never related to the local liquid configurations nor to dynamical properties. Moreover, the numerical prefactor in Born-Green formula is left undetermined, reducing significantly the utility of this formula. Another microscopic formula for liquid viscosity was recently suggested based on viscoelastic non-affine motion [16] but the identification of the relevant degrees of freedom was not properly clarified. A follow-up analysis [17] showed that, within that framework, only unstable localized normal modes contribute to viscosity with a possible crossover to a stable mode dominated regime at low temperatures. The results of [17] align with the idea of Egami that viscosity in liquids is governed by localized events. Interestingly, for water the timescale relevant for shear viscosity was proven to correlate with the connectivity of the fluctuating hydrogen bond network and the evolution of the first and second nearest neighbors [18, 19].

More recently, the microscopic origin of viscosity for 2D Yukawa liquids was revisited under the view of Egami's idea and discovered to coincide with the momentum transfer process of losing/gaining nearest neighbors for individual particles [20], leading to the phenomenological expression  $\eta = nm\bar{v}_p^2\tau_{LC}$  (with  $\bar{v}_p = \sqrt{2k_B T/m}$  the averaged particle speed and  $\rho = mn$ , where  $n$  is the number density). Nevertheless, the validity of this expression was verified only in one specific system and was not derived using any physical argument.

In summary, despite viscosity is a fundamental property of liquids whose microscopic origin has been disputed since centuries, a successful and universal theory from particle level motion, *akin* to kinetic theory for dilute gases, is still missing. Building on the '*shoulders of the giants*', in this work we combine previous theoretical ideas and propose a simple, and yet universal, theoretical framework to understand the fundamental origin of viscosity and collective shear dynamics in 2D simple liquids. We present two analytical and closed-form formulae for the liquid viscosity based on (I) single particle motion and

local configurational excitations and (II) pure structural information, encoded in the short range behavior of the pair correlation function. We demonstrate the validity of these expressions using extensive simulations in 2D Lennard-Jones (L-J), Yukawa, and one-component plasma (OCP) liquids, proving the universality of our findings. Finally, we demonstrate that the length-scale relevant for liquid viscosity aligns with the propagation length of collective shear waves, bridging particle level motion to collective dynamics in liquids, and unifying the previous theoretical frameworks for liquid viscosity.

## Liquid viscosity from microscopic particle motion

Following Frenkel's liquid description [5], self diffusion can be regarded as a hopping process between potential minima for individual atoms/molecules, where the average distance between two potential minima is assumed to be  $\xi$ , while the average hopping period is given by the Frenkel time  $\tau_F$ . By assuming simple random walk motion for the liquid constituents, the diffusion constant  $D$  can be written in terms of these two parameters as  $D = \xi^2/(4\tau_F)$ . At the same time, the mobility  $\alpha$  in a 2D liquid is expressed using Stokes law as  $\alpha = 1/(4\pi\eta)$ , and the Einstein's relation implies  $D = \alpha k_B T$ . By combining these expressions, we can obtain a simple formula for the shear viscosity of 2D liquids

$$\eta = \frac{k_B T \tau_F}{\pi \xi^2}. \quad (1)$$

This equation is the direct generalization of Frenkel's result (see Ref. [5]) to 2D liquids. Despite the elegance of this formula, its usefulness is questionable since nor  $\tau_F$  or  $\xi$  are explicitly defined and hence they cannot be estimated from simulation or experimental data.

In order to overcome this problem, we make two further assumptions. First, we assume that the Frenkel time  $\tau_F$  is given by the average lifetime of local connectivity  $\tau_F \equiv \tau_{LC}$ . This is reasonable since particle hopping in the potential landscape corresponds to local structural rearrangements described in real space by changes in the short-range topology. Second, we assume that the length-scale  $\xi$  is roughly independent of temperature and, at constant density, is approximately the same in the liquid and gas states. This assumption is corroborated *a posteriori* by direct numerical and analytical computations (see below). Following this hypothesis,  $\xi$  can be computed in the high-temperature gas state. There, the diffusion constant is well approximated by the kinetic theory formula [21]  $D = 1/2 v_{th} \xi$ , where  $v_{th} = \sqrt{k_B T/m}$  is the thermal average speed. Moreover, in the high-temperature regime, the diffusion process can be regarded as a traditional random walk [2], *i.e.*,  $D = \xi^2/(4\tau_0)$ , with  $\tau_0$  the corresponding collision time-scale. Combining these expressions, we obtain  $\xi = 2v_{th}\tau_0$ . From the definition of the average particle velocity  $\bar{v}_p = \sqrt{2k_B T/m}$ , *i.e.*,  $\bar{v}_p/\sqrt{2} = v_{th}$ , we then derive  $\xi = \sqrt{2}\bar{v}_p\tau_0$ .


Following the arguments just described, and defining a dimensionless parameter  $\lambda \equiv 1/(4\pi\bar{v}_p^2\tau_0^2)$ , Eq. (1) can be rewritten in a more concise form,

$$\eta = \lambda \rho \bar{v}_p^2 \tau_{LC}. \quad (2)$$

In summary, we have shown analytically that Eq. (2) is equivalent to Frenkel's expression for the liquid viscosity Eq. (1) upon identifying the Frenkel time with  $\tau_{LC}$ . We notice that, despite the simplifying assumptions, we are not able to derive the value of  $\lambda$ , which we expect to be strongly dependent on the microscopic details of the liquid considered and therefore not universal. Interestingly, as will be presented below, we find  $\lambda \approx 1$  in all the systems considered, suggesting an even simpler formula  $\eta = \rho \bar{v}_p^2 \tau_{LC}$ .

To validate Eq. (2), we perform numerical simulations of 2D L-J, Yukawa, and OCP liquids. In 2D L-J systems [22], the interparticle interaction  $\phi(r) = 4\epsilon [(\sigma/r)^{12} - (\sigma/r)^6]$  consists of both repulsive and attractive terms, where  $\epsilon$  and  $\sigma$  are the energy and distance parameters. In 2D Yukawa systems, the interaction between particles is a screened Coulomb repulsion  $\phi(r) = Q^2 \exp(-r/\lambda_D)/4\pi\epsilon_0 r$ , where  $\lambda_D$  is the Debye length and  $Q$  is the particle charge. In 2D OCP systems, the interaction is the classical Coulomb repulsion  $\phi(r) = Q^2/4\pi\epsilon_0 r$ . For our simulated 2D systems under various conditions, the reduced temperature  $T/T_m$  is specified from  $\geq 1.1$  to 20 at most, where the corresponding melting points  $T_m$  of these systems are obtained from the previous investigations [23, 24]. Besides the temperature  $T$ , we also vary the number density  $n$  of 2D L-J systems and the screening parameter  $\kappa = a/\lambda_D$  of 2D Yukawa systems, where  $a$  is the Wigner-Seitz radius [25–29]. All simulation details are provided in the Methods section. Our simulations provide a rather large sample of 2D liquids with remarkably different particle interactions under various conditions, allowing us to test in detail the universality of our findings.

Our numerical results obtained using the Green-Kubo formalism are presented in Fig. 2(a). Following Ref. [30], we present the obtained viscosity in a dimensionless form  $\eta/(nmk_B T)^{1/2}$ . Interestingly, both the L-J data and the Yukawa and OCP ones collapse into two universal curves as a function of the reduced temperature  $T/T_m$ . For all systems, the shear viscosity in the liquid phase decreases with temperature, as expected.



**Figure 2. Microscopic origin of viscosity as diffusive transport of average particle momentum:** (a) Dimensionless viscosity as a function of reduced temperature. (b) Dimensionless local connectivity time as a function of reduced temperature. (c) Test of the universal formula for viscosity proposed in Eq. (2) of the main text.

To reveal the fundamental origin of viscosity at the particle level motion, we calculate the lifetime of the local atomic connectivity  $\tau_{LC}$  [8, 31] for the simulated 2D liquids by tracking the neighbors of each particle (see Methods), as presented in Fig. 2(b). By employing the same dimensionless form,  $\tau_{LC}/(nk_B T/m)^{-1/2}$ , all the data collapse into two universal curves as well. We notice the discrepancy between these two curves, as in the case of viscosity in Fig. 2(a), probably due to the attractive contribution to the potential that is present in the L-J systems but absent in the Yukawa and OCP liquids.

The similarity in the shape of the normalized viscosity and the normalized local connectivity time is striking, suggesting a direct relation between these two quantities. In order to investigate this point, we rewrite Eq. (2) in dimensionless units [30],

$$\frac{\eta}{(nmk_B T)^{1/2}} = 2\lambda \frac{\tau_{LC}}{(nk_B T/m)^{-1/2}}. \quad (3)$$


This identity can now be directly tested using the simulation data presented in Figs. 2(a) and 2(b). In Fig. 2(c), we test directly our proposed expression Eq. (3) for 2D L-J, Yukawa and OCP liquids. Our numerical results confirm the validity of Eq. (2) with  $\lambda \approx 1.03$  for 2D Yukawa and OCP liquids, and  $\lambda \approx 1.18$  for 2D L-J liquids. As already anticipated, these values are surprisingly close to  $\lambda = 1$ .

In summary, our analysis supports the physical idea that, in 2D simple liquids, the momentum transport process responsible for the macroscopic shear viscosity does originate from losing/gaining neighbors at the individual particle level. Since  $\eta/\rho$  in liquids controls the diffusive transport of transverse collective momentum, our results also indicate that this macroscopic dynamical process is associated to a characteristic microscopic time-scale  $\tau_{LC}$  and a microscopic length-scale  $l_p = \bar{v}_p \tau_{LC}$ . We also emphasize that Eq. (2) establishes a direct link between macroscopic dynamics and microscopic particle motion, providing a fundamental understanding of shear viscosity in liquids at all scales.

## Connecting microscopic particle motion with collective shear dynamics in liquids

We now take a step back and reconsider the collective shear dynamics in liquid under Maxwell's perspective [32]. Combining Maxwell approach with Navier Stokes equations, it has been shown [12, 13] that the dynamics of collective shear waves in liquids are described by the following telegrapher equation:

$$\omega^2 + i\omega/\tau_M = v_T^2 k^2. \quad (4)$$



**Figure 3. Bridging collective shear dynamics to particle-level motion:** (a) Spectra of transverse modes in a 2D Yukawa liquid with  $\kappa = 1$ , and the corresponding dispersion relation marked as dots. Frequencies are normalized by the nominal dusty plasma frequency  $\omega_{pd} \equiv (Q^2/2\pi m \epsilon_0 m a^3)^{1/2}$  [37], while wave-vectors are normalized using the Wigner-Seitz radius  $a$ . The obtained dispersion relations under different reduced temperatures are presented in the inset. (b) Same analysis for 2D L-J liquids with particle density  $n = 1$  at different reduced temperatures  $T/T_m$ . (c) Universal linear relation between the dimensionless cutoff wave-vector  $k_g$  and the inverse dimensionless microscopic length-scale  $l_p = \bar{v}_p \tau_{LC}$ . The slopes of the two fitting lines are  $\approx 0.98$  and  $0.88$ , respectively.

This equation implies that the real part of the shear wave dispersion relation presents a gap in momentum space,

$$\text{Re}(\omega) = v_T \sqrt{k^2 - k_g^2}, \quad k_g = \frac{1}{2v_T \tau_M}, \quad (5)$$

where the expression of  $k_g$  relies on Maxwell viscoelasticity theory (see [33] for one alternative interpretation). This  $k$ -gap feature has been confirmed in many simulation works (*e.g.*, [34]) and also in a few experimental setups [35, 36]. Most importantly, this result implies that the propagation of collective shear waves in liquids is confined up to an elastic length-scale given by  $l \equiv 1/k_g$ . In other words, one could construct an idealized model of a liquid as composed of elastic patches of size  $l$  in which the dynamics are solid-like and mainly composed of quasi-harmonic oscillations localized at the bottom of the potential [13] (see Fig. 1(d)). Beyond this elastic length-scale, elastic forces get screened, shear stresses are not supported anymore and the dynamics become liquid-like, *i.e.* dominated by shear diffusive transport rather than coherent wave-like excitations as in solids. Moreover, withing Maxwell's theory, the average size of these solid-like regions shrinks with temperature, as a direct consequence of  $\tau_M$  decreasing rapidly with  $T$ .

Following this idea, one could ask whether this collective elastic length-scale bears any relation to the particle-level length-scale governing viscosity,  $l_p = \bar{v}_p \tau_{LC}$ , connecting somehow the macroscopic Maxwell view with the particle level Frenkel's description. In this direction, we notice that in Frenkel's picture of liquid dynamics (see Fig. 1(a)), the collective elastic length-scale corresponds to the average length made by one particle hopping across potential barriers that is directly related to the viscosity of the system (see Eq. (1)).

In Fig. 3(a) we present the numerical results for the dispersion relation of collective shear waves of the 2D Yukawa liquids with  $\kappa = 1$ . The presence of a cutoff wave-vector  $k_g$  is evident and its size grows with temperature, as expected. By tracking the position at which  $\text{Re}(\omega) \rightarrow 0$ , we are able to derive the temperature dependence of  $k_g$ . A similar analysis has been performed for the 2D L-J liquids and the corresponding results are presented in Fig. 3(b).

Finally, in Fig. 3(c), we plot the dimensionless cutoff wave-vector  $k_g a$  as a function of the dimensionless inverse length-scale  $a/l_p$ , with  $l_p = \bar{v}_p \tau_{LC}$ . For all the systems considered, we find a universal linear relation:

$$k_g = \frac{\beta}{l_p}, \quad (6)$$

where  $\beta$  is a constant of order one:  $\beta \approx 0.88$  for 2D L-J systems and  $\beta \approx 0.98$  for 2D Yukawa and OCP systems.

This result implies a direct proportionality between the elastic length-scale  $l \sim k_g^{-1}$ , relevant for collective shear dynamics, and the microscopic length-scale  $l_p = \bar{v}_p \tau_{LC}$  that governs the macroscopic shear viscosity through Eq. (2). It also suggests that the propagation of collective shear waves in liquids is hindered by local configurational excitations that drive structural

rearrangements within the nearest neighbor cage. Furthermore, it is important to stress that, despite a lot of discussions in the past [12, 13], a formal connection between Frenkel's ideas and  $k$ -gap theory has never been achieved before. In fact, despite  $k$ -gap theory was strongly motivated by Frenkel's intuition, the relevant timescale has been always identified with the collective Maxwell relaxation time that, in first approximation, bears no connection with the microscopic Frenkel's time. Eq. (6) provides the missing link between collective shear dynamics, as envisaged by  $k$ -gap theory, and microscopic particle hops à la Frenkel, formalized using the concept of local connectivity time proposed by Egami. Motivated by this observation, we proposed that the length-scale associated to the  $k$ -gap should not be associated to the Maxwell length-scale  $l_m = \nu_T \tau_M$  but rather to the particle length-scale  $l_p$ , in better alignment with Frenkel's initial proposal.

## Structural origin of liquid viscosity

So far, we have succeeded in connecting the local connectivity time with the macroscopic shear viscosity and showed that the length-scale associated to it aligns (up to an order one constant) with the propagation length of collective shear waves in liquids. In Frenkel's liquid description [5], fluidity and viscosity arise from particles' hopping over potential barriers, as depicted in Fig. 1(c). The average hopping time  $\tau_F$  can be expressed as  $\tau_F = \tau_0 \exp(U/k_B T)$ , where  $U$  is the potential energy barrier, while  $\tau_0$  is the corresponding time in the limit of very high temperatures, or in the gas-like state.

From a structural point of view, a particle hopping over a single potential barrier corresponds to a re-arrangement of its cage or its neighbors. We have therefore advanced the idea that the single particle Frenkel time should be taken to coincide with  $\tau_{LC}$ . This idea is also supported by the validity of Eq. (6) that has been directly verified in Fig. 3. Following this hypothesis, the energy barrier  $U$  should correspond to the energy for one particle hopping outside the cage formed by its neighboring particles. As a result, the pair correlation function  $g(r)$  should encode the information about  $U$ .

In Fig. 4(a), we plot the calculated  $g(r)$  and the corresponding effective potential  $w(r)/k_B T = -\ln(g(r))$  [2] for a typical 2D L-J liquid with  $n = 1$  and  $T/T_m = 5$ . We propose that the energy barrier  $U$  governing the hopping of individual particles in the Frenkel description of liquids is given by

$$U \equiv \Delta w = k_B T \ln [g(r)_{\max}/g(r)_{\min}], \quad (7)$$

where  $\max$  and  $\min$  correspond respectively to the position of the first maximum and first minimum in  $g(r)$ . The magnitude of  $U$  in Eq. (7) is represented with the vertical black arrows in Fig. 4(a).

We then calculate  $\exp(\Delta w/k_B T) = g(r)_{\max}/g(r)_{\min}$  for different 2D liquids under various conditions and present these results as a function of the reduced temperature in Fig. 4(b). These obtained data points for  $\exp(\Delta w/k_B T)$  collapse into two universal curves and present similar variation trends as those for  $\eta$  and  $\tau_{LC}$  in Fig. 2, clearly indicating the strong correlations between these three physical quantities.

To further elucidate this connection, in Fig. 4(c) we plot the dimensionless local connectivity time as a function of  $\exp(\Delta w/k_B T)$ . We find that these quantities present a robust linear relation independently of the thermodynamic conditions, *i.e.* the value of  $T/T_m$ . This suggests a simple and striking relation between the local connectivity time and the short-range structural properties of liquids that can be formalized as


$$\tau_{LC} = \tau_0 \exp\left(\frac{\Delta w}{k_B T}\right) = \tau_0 \frac{g(r)_{\max}}{g(r)_{\min}}, \quad (8)$$

where  $\tau_0 = \gamma(nk_B T/m)^{-1/2}$  with  $\gamma \approx 0.181$  for 2D L-J liquids and  $\gamma \approx 0.201$  for 2D Yukawa and OCP liquids. Here,  $\tau_0$  represents the relevant time-scale in the decorrelated gas-like regime in which particles undergo independent dynamics. The correction coming from  $g(r)_{\max}/g(r)_{\min}$  takes into account the short-range and mid-range correlations that become important in the liquid state upon decreasing temperature. This term indeed vanishes if the pair correlation function loses its first peak and first minimum, as expected in the ideal gas state. It is immediate to verify that the parameter  $\gamma$  is related to  $\lambda$  in Eq. (2) via  $\lambda = 1/(8\pi\gamma^2)$ , as verified by directly numerical comparison.

Eq. (8) allows us to re-write our formula for the viscosity in the following form

$$\eta = \frac{m}{4\pi\tau_0} \frac{g(r)_{\max}}{g(r)_{\min}}, \quad (9)$$

in terms of the mass of each particle  $m$ , the high-temperature relaxation time  $\tau_0$ , and the pure short-range structural information based on the pair correlation function  $g(r)$ .



**Figure 4. Structural definition of the local connectivity time:** (a) Calculated pair-correlation function  $g(r)$  of a 2D L-J liquid (red line) and corresponding effective potential  $w(r)/k_B T$  (blue dashed line). The vertical black arrows indicate the potential difference between the first maximum and first minimum that is identified with the potential barrier  $U$  in Frenkel’s description, Eq. (7). (b) The temperature dependence of the potential factor  $\exp(\Delta w/k_B T)$  as a function of the reduced temperature  $T/T_m$  for the various systems studied. (c) The universal linear relation between the dimensionless local connectivity time  $\tau_{LC}/(nk_B T/m)^{-1/2}$  and  $\exp(\Delta w/k_B T)$  for all systems considered.

Eq. (9) completes in a sense Eyring’s expression,  $\eta = A \exp(\Delta G/k_B T)$  [6], by providing a clear definition of the pre-factor  $A$  and the energy barrier  $\Delta G$ . Indeed, in 2D liquids, in view of our results

$$A \equiv \frac{m}{4\pi\tau_0}, \quad \Delta G = \Delta w = k_B T \ln [g(r)_{\max}/g(r)_{\min}]. \quad (10)$$

In fact,  $A = m/(4\pi\tau_0)$  is just the viscosity at extremely high temperatures, *i.e.* in the gas-like state, and the potential barrier  $\Delta G$  is directly defined from the short-range order properties of the liquid.


After proving that the energy barrier governing liquid viscosity is the one between the first maximum and first minimum in  $g(r)$ , it comes naturally to identify the distance between these two as the length-scale associated to the potential hops in Frenkel’s description of liquid dynamics. To confirm this hypothesis, we have computed the pair correlation functions  $g(r)$  for 2D Yukawa and L-J fluids in a wide range of conditions, as presented in Figs. 5(a) and 5(b).

We have then computed the distance  $\Delta$  between the first peak and first valley in  $g(r)$  as a function of the reduced temperature  $T/T_m$ , as presented in Fig. 5(c). Interestingly, when normalizing  $\Delta$  using the Wigner-Seitz radius  $a$ , we find that this length-scale is approximately constant in temperature, showing mild deviations only at low temperatures. For both systems, we find that  $\Delta \approx 0.87a$ , perfectly aligning with the idea that the length-scale  $\xi$  in Frenkel’s approach does not depend on temperature but only on the density  $n$  (as kept constant in Fig. 5), as a result,  $\xi$  has to be identified with  $\Delta$  as defined above. To prove this further, by combining  $\xi = \sqrt{2}\bar{v}_p\tau_0$  with the definitions  $\tau_0 = \gamma(nk_B T/m)^{-1/2}$  and  $\bar{v}_p = \sqrt{2k_B T/m}$ , we immediately obtain  $\xi = 2\gamma/\sqrt{n}$ , confirming that  $\xi$  is a quantity that does not depend on temperature but only on the particle’s number density  $n$ . By expressing  $n$  in terms of the Wigner-Seitz radius as  $n = 1/(\pi a^2)$ , we can rewrite the above expression as  $\xi = 2\gamma\sqrt{\pi}a$ . Finally, by using the values of  $\gamma$  extrapolated from the previous data analysis, our theoretical estimate gives  $\xi \approx 0.71a$  for 2D Yukawa and OCP liquids, and  $\xi \approx 0.64a$  for 2D L-J liquids.

## Discussion

In this work, we have considered the long-standing problem of deriving a microscopic and predictive formula for the shear viscosity of 2D simple fluids, challenging the famous Landau argument that is “*impossible to derive any general formulae giving a quantitative description of the properties of a liquid*” [38]. Our findings defy this paradigm and prove that, at least for 2D simple liquids, a microscopic formula for the viscosity can be found, in excellent agreement with the simulation data in several systems characterized by profoundly different particle interactions.

In fact, our microscopic formula for the viscosity has been achieved not only at the particle level motion, but also from direct information of the short-range structural correlation of the liquid, which is encoded in its pair correlation function. Our



**Figure 5.** Calculated pair correlation functions  $g(r)$  of 2D Yukawa (a) and L-J (b) fluids under various conditions, as well as the distance  $\Delta$  between the first peak and first valley of  $g(r)$  (c).

derived viscosity equation connects the microscopic motion of particles, the macroscopic dynamics of collective shear waves, and the liquid structure of  $g(r)$ .

Importantly, our results complete three of the most successful frameworks to describe liquid dynamics and viscosity: Frenkel’s theory, Maxwell’s theory, and Eyring’s theory. First, we propose and confirm the idea of identifying the microscopic Frenkel’s time with the lifetime of local connectivity. Second, we find that the elastic length-scale below which collective shear waves propagate in liquids according to Maxwell model and  $k$ -gap theory can be directly connected to a single particle length-scale governing the diffusive transport of particle’s momentum. Finally, we provide a precise definition of the undetermined parameters in Eyring’s formula for viscosity and in particular we propose a simple method to obtain the hopping potential barrier from the short-range properties of the pair correlation function.

Do similar simple arguments apply to 3D liquids? Do our formulae provide an accurate estimate of the shear viscosity also for complex liquids and glass forming systems? This remains to be seen.

## Methods

### Simulation Method for 2D simple liquids

We perform equilibrium molecular dynamics (MD) simulations of 2D Lennard-Jones (L-J), Yukawa, and Coulomb one-component plasma (OCP) liquids. For all these three simple liquids, the equation of motion for each particle is

$$m\ddot{\mathbf{r}}_i = -\nabla\Sigma\phi_{ij}, \quad (11)$$

where  $-\nabla\Sigma\phi_{ij}$  is the particle-particle interaction, while  $\mathbf{r}_i$  is the position vector for the  $i$ -th particle. In our current investigation, we always simulate  $N = 4096$  particles constrained in a 2D simulation box with the length ratio of  $L_x : L_y = 2 : \sqrt{3}$  with periodic boundary conditions.

For each simulation run, first we integrate the equation of motion for all particles with a thermostat for  $N_1$  steps, so that the simulation system reaches the specified conditions. Then, we turn off the thermostat to integrate the equation of motion for the next  $N_2$  steps, and the obtained data are used for the data analysis presented in the main text. In our simulations, we specify the reduced temperature value  $T/T_m$ , where  $T$  is the temperature of the simulated 2D system, while  $T_m$  is the corresponding melting point. Also, we truncate the interparticle potential at  $r_c$  to ensure that the potential energy of the simulation system does not change significantly with the increase of  $r_c$  any more. We also verify that, for each simulation run, our time step is always chosen to be small enough, so that energy conservation is adequately obeyed.



## 2D L-J liquids

For 2D L-J liquids, the interparticle interaction is

$$\phi(r) = 4\epsilon \left[ (\sigma/r)^{12} - (\sigma/r)^6 \right], \quad (12)$$

where  $\epsilon$  and  $\sigma$  are the energy and distance parameters. Here, we normalize the length and time using  $\sigma$  and  $\sqrt{m\sigma^2/\epsilon}$ , respectively. In our simulations of 2D L-J liquids, we specify the values of both the number density  $n = N/A'$  and the reduced temperature  $T/T_m$ , where  $A'$  is the area of the simulated box.

Here are other simulation details. In our simulations, the number density are specified as  $n = 0.8, 0.85, 0.9, 0.95$ , and  $1.00$ . For each specified value of  $n$ , we vary the reduced temperature  $T/T_m$  from 2 to 20. Note, we choose the melting points of 2D L-J systems  $T_m$  for different  $n$  values from [23]. In our 2D L-J simulations, the cutoff radius is chosen as  $r_c = 2.5\sigma$ , while the corresponding integration steps are specified as  $N_1 = 2 \times 10^6$  and  $N_2 = 10^9$ , respectively.

## 2D Yukawa liquids

For 2D Yukawa liquids, the interparticle interaction is the Yukawa repulsion

$$\phi(r) = Q^2 \exp(-r/\lambda_D) / 4\pi\epsilon_0 r, \quad (13)$$

where  $\lambda_D$  is the Debye length and  $Q$  is the charge on each particle. Besides the reduced temperature  $T/T_m$ , we also use the screening parameter  $\kappa = a/\lambda_D$  to characterize the simulated 2D Yukawa liquids.

In order to mimic the conditions of most 2D dusty plasma experiments, we vary the  $\kappa$  value from 0.75 to 2. For each  $\kappa$  value, we vary the value of the reduced temperature  $T/T_m$  from 1.11 to 10, where the values for the melting point  $T_m$  are taken from [24]. In our 2D Yukawa simulations, the cutoff radius is chosen as  $r_c = 22a$ , less than one half of each side of the simulation box. The integration steps are specified as  $N_1 = 1 \times 10^7$  and  $N_2 = 10^8$ , respectively.

## 2D Coulomb OCP liquids

For 2D Coulomb OCP liquids, the interparticle interaction is

$$\phi(r) = Q^2 / 4\pi\epsilon_0 r. \quad (14)$$

Unlike L-J and Yukawa liquids above, the interaction between particles in 2D Coulomb OCP liquids is long-range. To avoid the Ewald summation [39], we use the approximate potential [40]

$$\phi(r) = \frac{Q^2}{4\pi\epsilon_0} \left[ \frac{\text{erfc}(\alpha_1 r)}{r} - \frac{\text{erfc}(\alpha_1 r_c)}{r_c} + \left( \frac{\text{erfc}(\alpha_1 r_c)}{r_c^2} + \frac{2\alpha_1 \exp(-\alpha_1^2 r_c^2)}{\sqrt{\pi} r_c} \right) (r - r_c) \right], \quad (15)$$

where  $\alpha_1$  is the ‘‘damping’’ parameter,  $r_c$  is the cutoff radius, and  $\text{erfc}()$  is the complementary error function. From previous studies, the choice of  $\alpha_1 = 0.2$  enables the energy and forces of the simulated system to quickly converge to the long-range Coulomb system. In our 2D Coulomb OCP simulations, we choose  $\alpha_1 = 0.2$  and  $r_c = 10 a$ , respectively. Other simulation details are the same as those for 2D Yukawa simulations described above.

## Lifetime of local connectivity from simulations

In our current investigation, to calculate the lifetime of local connectivity  $\tau_{LC}$  of our simulated 2D liquids, we need to track the neighbor list of all particles at each moment. For each studied particle  $i$ , its neighbors are defined as its pairing particle  $j$  with their distance  $r_{ij}$  less than the separation of the first minimum of the radial distribution function  $g(r)$  [8, 31]. For example, in the initial configuration, there are  $N(t_0)$  neighbors for the studied particle  $i$ . As the time goes from the initial time  $t_0$  to  $t_0 + t$ , the neighbors of the studied particle  $i$  change, i.e., some of the initial neighbors are not its neighbors any more. We may use  $N(t_0 + t)$  to label the number of the initial neighbors which are still its neighbors at the time of  $t_0 + t$ . Thus, the lifetime of local connectivity  $\tau_{LC}$  is defined as the time duration, relative to  $t_0$ , for the number of initial neighbors falls by 1 in the ensemble average, i.e.,  $\langle N(t_0) \rangle - \langle N(t_0 + t) \rangle = 1$  [8, 31], for all studied particles and varying the different initial times of  $t_0$ . In fact, if one neighbor leaves the studied particle  $i$  for a while, then comes back as a neighbor again, it is still regarded as a new neighbor for the studied particle  $i$ . In summary,  $\tau_{LC}$  can be regarded as the averaged time for the first of the initial neighbors of one particle  $i$  goes beyond the distance of the first minimum of  $g(r)$ , i.e., the coordination number falls by 1 [31], or equivalently a new particle enter the range of one particle  $i$  within the distance of the first minimum of  $g(r)$  while none of the initial neighbors goes beyond.

## Data availability

The datasets generated and analyzed during the current study are available upon reasonable request by contacting the corresponding authors.

## Code availability

The codes that support the findings of this study is available upon reasonable request by contacting the corresponding authors.

## Acknowledgements

This work was supported by the National Natural Science Foundation of China under Grants No. 12175159, No. 12305220, No. 12347110, the Excellent Postdoctoral Program of Jiangsu Province, the 1000 Youth Talents Plan, and the Priority Academic Program Development (PAPD) of Jiangsu Higher Education Institutions. M.B. acknowledges the support of the Shanghai Municipal Science and Technology Major Project (Grant No.2019SHZDZX01) and the sponsorship from the Yangyang Development Fund.

## Author contributions

Y. F. and D. H. conceived the idea of this project. Y. F. and M. B. supervised the project. D. H., S. L. and C. L. performed the simulations and the data analysis. All authors contributed to the writing of the manuscript and the theoretical interpretation of the results.

## Competing interests

The authors declare no competing interests.

## References

1. Landau, L. D. & Lifshitz, E. M. *Fluid Mechanics (Second Edition)* (Pergamon, 1987).
2. Hansen, J.-P. & McDonald, I. R. *Theory of simple liquids: with applications to soft matter* (Academic press, 2013).
3. Evans, D. J. & Morriss, G. *Statistical Mechanics of Nonequilibrium Liquids* (Cambridge University Press, 2008), 2 edn.
4. Loeb, L. B. *The kinetic theory of gases* (Courier Corporation, 2004).
5. Frenkel, J. *Kinetic theory of liquids*. International series of monographs on physics (Clarendon Press Oxford, Oxford, 1946).
6. Eyring, H. The activated complex in chemical reactions. *The J. Chem. Phys.* **3**, 107–115 (1935).
7. Touloukian, Y. S., Saxena, S. C. & Hestermans, P. *Thermophysical properties of matter - the TPRC data series. Volume 11. Viscosity. (Reannouncement). Data book* (1975).
8. Iwashita, T., Nicholson, D. M. & Egami, T. Elementary excitations and crossover phenomenon in liquids. *Phys. Rev. Lett.* **110**, 205504, DOI: [10.1103/PhysRevLett.110.205504](https://doi.org/10.1103/PhysRevLett.110.205504) (2013).
9. Shinohara, Y. *et al.* Viscosity and real-space molecular motion of water: Observation with inelastic x-ray scattering. *Phys. Rev. E* **98**, 022604, DOI: [10.1103/PhysRevE.98.022604](https://doi.org/10.1103/PhysRevE.98.022604) (2018).
10. Levashov, V. A., Morris, J. R. & Egami, T. The origin of viscosity as seen through atomic level stress correlation function. *The J. Chem. Phys.* **138**, 044507, DOI: [10.1063/1.4789306](https://doi.org/10.1063/1.4789306) (2013).
11. Bellissard, J. & Egami, T. Simple theory of viscosity in liquids. *Phys. Rev. E* **98**, 063005, DOI: [10.1103/PhysRevE.98.063005](https://doi.org/10.1103/PhysRevE.98.063005) (2018).
12. Trachenko, K. & Brazhkin, V. V. Collective modes and thermodynamics of the liquid state. *Reports on Prog. Phys.* **79**, 016502, DOI: [10.1088/0034-4885/79/1/016502](https://doi.org/10.1088/0034-4885/79/1/016502) (2015).
13. Baggioli, M., Vasin, M., Brazhkin, V. & Trachenko, K. Gapped momentum states. *Phys. Reports* **865**, 1–44, DOI: <https://doi.org/10.1016/j.physrep.2020.04.002> (2020). Gapped momentum states.
14. Peluso, F. The viscosity of liquids in the dual model (2024). [2409.15502](https://arxiv.org/abs/2409.15502).
15. Born, M. & Green, H. S. A general kinetic theory of liquids iii. dynamical properties. *Proc. Royal Soc. London. Ser. A. Math. Phys. Sci.* **190**, 455–474, DOI: [10.1098/rspa.1947.0088](https://doi.org/10.1098/rspa.1947.0088) (1947).

16. Zaccone, A. General theory of the viscosity of liquids and solids from nonaffine particle motions. *Phys. Rev. E* **108**, 044101, DOI: [10.1103/PhysRevE.108.044101](https://doi.org/10.1103/PhysRevE.108.044101) (2023).
17. Huang, L.-Z., Cui, B., Vaibhav, V., Baggioli, M. & Wang, Y.-J. Microscopic origin of liquid viscosity from unstable localized modes (2024). [2408.07937](https://arxiv.org/abs/2408.07937).
18. Iwashita, T. *et al.* Seeing real-space dynamics of liquid water through inelastic x-ray scattering. *Sci. Adv.* **3**, e1603079, DOI: [10.1126/sciadv.1603079](https://doi.org/10.1126/sciadv.1603079) (2017).
19. Yahya, A. *et al.* Molecular origins of bulk viscosity in liquid water. *Phys. Chem. Chem. Phys.* **22**, 9494–9502, DOI: [10.1039/D0CP01560A](https://doi.org/10.1039/D0CP01560A) (2020).
20. Huang, D., Lu, S., Murillo, M. S. & Feng, Y. Origin of viscosity at individual particle level in yukawa liquids. *Phys. Rev. Res.* **4**, 033064, DOI: [10.1103/PhysRevResearch.4.033064](https://doi.org/10.1103/PhysRevResearch.4.033064) (2022).
21. Pitaevskii, L. P. & Lifshitz, E. *Physical Kinetics: Volume 10*, vol. 10 (Butterworth-Heinemann, 2012).
22. Barker, J. A., Henderson, D. & Abraham, F. F. Phase diagram of the two-dimensional Lennard-Jones system; Evidence for first-order transitions. *Phys. A Stat. Mech. its Appl.* **106**, 226–238, DOI: [10.1016/0378-4371\(81\)90222-3](https://doi.org/10.1016/0378-4371(81)90222-3) (1981).
23. Li, Y.-W. & Ciamarra, M. P. Phase behavior of lennard-jones particles in two dimensions. *Phys. Rev. E* **102**, 062101, DOI: [10.1103/PhysRevE.102.062101](https://doi.org/10.1103/PhysRevE.102.062101) (2020).
24. Hartmann, P., Kalman, G. J., Donkó, Z. & Kutasi, K. Equilibrium properties and phase diagram of two-dimensional yukawa systems. *Phys. Rev. E* **72**, 026409, DOI: [10.1103/PhysRevE.72.026409](https://doi.org/10.1103/PhysRevE.72.026409) (2005).
25. Melzer, A., Homann, A. & Piel, A. Experimental investigation of the melting transition of the plasma crystal. *Phys. Rev. E* **53**, 2757–2766, DOI: [10.1103/PhysRevE.53.2757](https://doi.org/10.1103/PhysRevE.53.2757) (1996).
26. Morfill, G. E. & Ivlev, A. V. Complex plasmas: An interdisciplinary research field. *Rev. Mod. Phys.* **81**, 1353, DOI: [10.1103/RevModPhys.81.1353](https://doi.org/10.1103/RevModPhys.81.1353) (2009).
27. Donkó, Z., Goree, J., Hartmann, P. & Kutasi, K. Shear viscosity and shear thinning in two-dimensional yukawa liquids. *Phys. Rev. Lett.* **96**, 145003, DOI: [10.1103/PhysRevLett.96.145003](https://doi.org/10.1103/PhysRevLett.96.145003) (2006).
28. Daligault, J. Liquid-state properties of a one-component plasma. *Phys. Rev. Lett.* **96**, 065003, DOI: [10.1103/PhysRevLett.96.145003](https://doi.org/10.1103/PhysRevLett.96.145003) (2006).
29. Baalrud, S. D. & Daligault, J. Effective potential theory for transport coefficients across coupling regimes. *Phys. Rev. Lett.* **110**, 235001, DOI: [10.1103/PhysRevLett.110.235001](https://doi.org/10.1103/PhysRevLett.110.235001) (2013).
30. Rosenfeld, Y. Quasi-universal melting-temperature scaling of transport coefficients in yukawa systems. *J. Physics: Condens. Matter* **13**, L39, DOI: [10.1088/0953-8984/13/2/101](https://doi.org/10.1088/0953-8984/13/2/101) (2001).
31. Ashwin, J. & Sen, A. Microscopic origin of shear relaxation in a model viscoelastic liquid. *Phys. Rev. Lett.* **114**, 055002, DOI: [10.1103/PhysRevLett.114.055002](https://doi.org/10.1103/PhysRevLett.114.055002) (2015).
32. Maxwell, J. C. IV. On the dynamical theory of gases. *Philos. transactions Royal Soc. Lond.* 49–88 (1867).
33. Baggioli, M., Landry, M. & Zaccone, A. Deformations, relaxation, and broken symmetries in liquids, solids, and glasses: A unified topological field theory. *Phys. Rev. E* **105**, 024602, DOI: [10.1103/PhysRevE.105.024602](https://doi.org/10.1103/PhysRevE.105.024602) (2022).
34. Yang, C., Dove, M. T., Brazhkin, V. V. & Trachenko, K. Emergence and evolution of the  $k$  gap in spectra of liquid and supercritical states. *Phys. Rev. Lett.* **118**, 215502, DOI: [10.1103/PhysRevLett.118.215502](https://doi.org/10.1103/PhysRevLett.118.215502) (2017).
35. Nosenko, V., Goree, J. & Piel, A. Cutoff wave number for shear waves in a two-dimensional yukawa system (dusty plasma). *Phys. Rev. Lett.* **97**, 115001, DOI: [10.1103/PhysRevLett.97.115001](https://doi.org/10.1103/PhysRevLett.97.115001) (2006).
36. Jiang, C., Zheng, Z., Chen, Y., Baggioli, M. & Zhang, J. Experimental observation of gapped shear waves and liquid-like to gas-like dynamical crossover in active granular matter. *arXiv preprint arXiv:2403.08285* (2024).
37. Kalman, G. J., Hartmann, P., Donkó, Z. & Rosenberg, M. Two-dimensional yukawa liquids: Correlation and dynamics. *Phys. Rev. Lett.* **92**, 065001, DOI: [10.1103/PhysRevLett.92.065001](https://doi.org/10.1103/PhysRevLett.92.065001) (2004).
38. Landau, L. D. & Lifshitz, E. M. *Statistical Physics: Volume 5*, vol. 5 (Pergamon, 1980).
39. LeBard, D. N. *et al.* Self-assembly of coarse-grained ionic surfactants accelerated by graphics processing units. *Soft Matter* **8**, 2385–2397 (2012).
40. Fennell, C. J. & Gezelter, J. D. Is the ewald summation still necessary? pairwise alternatives to the accepted standard for long-range electrostatics. *The J. Chem. Phys.* **124**, 234104, DOI: [10.1063/1.2206581](https://doi.org/10.1063/1.2206581) (2006).