UNIQUE EXPANSIONS IN NUMBER SYSTEMS VIA SOLUTIONS OF REFINEMENT EQUATION

SERGEI V. KONYAGIN, VLADIMIR YU. PROTASOV, AND ALEXEY L. TALAMBUTSA

ABSTRACT. Using the subdivision schemes theory, we develop a criterion to check if any natural number has at most one representation in the *n*-ary number system with a set of non-negative integer digits $A = \{a_1, a_2, \ldots, a_n\}$ that contains zero. This uniqueness property is shown to be equivalent to a certain restriction on the roots of the trigonometric polynomial $\sum_{k=1}^{n} e^{-2\pi i a_k t}$. From this criterion, under a natural condition of irreducibility for A, we deduce that in case of prime n the uniqueness holds if and only if the digits of A are distinct modulo n, whereas for any composite n we show that the latter condition is not necessary. We also establish the connection of this uniqueness to the semigroup freeness problem for affine integer functions of equal integer slope; this together with the two criteria allows to fill the gap in the work of D.Klarner on Erdös question about densities of affine integer orbits and establish a simple algorithm to check the freeness and the positivity of density when the slope is a prime number.

1. Number systems and semigroups freeness

Given an integer $n \ge 2$, we consider a set of nonnegative integer numbers $A = \{a_1, \ldots, a_m\}$, which are called digits of the *(non-standard) n-ary number system* A.

For an integer $k \ge 0$, we call its A-expansion any tuple of digits $(\alpha_0, \ldots, \alpha_J)$ such that

(1)
$$k = \sum_{j=0}^{J} \alpha_j n^j$$
, where $\alpha_J \neq 0$.

In particular, for k = 0 its A-expansion is an empty tuple. We say that the *n*-ary number system A has *uniqueness property* if any integer $k \ge 0$ has at most one A-expansion.

The uniqueness property has been studied in the literature mainly because of two applications. The first is the theory of self-similar tiles, where uniqueness gives the existence criterion for the tile. See [14, 20, 21] and references therein for the general multivariate case. The second one is related to the coding theory, but as we shall show, it also turns out to be connected to the freeness of particular semigroups and densities of integer orbit sets arising from the action of integer affine functions.

The active analysis of number systems with positive digits was initiated in early 1980's by H.A. Maurer, A. Salomaa and D. Wood in [22], where they introduced L-codes and discovered that if the code alphabet is unary, then an L-code corresponds to a non-standard number system¹. They have proved that such unary code can be uniquely decoded if and only if the corresponding number system has uniqueness property. It was also proved that this property necessarily fails if m > n, while if all α_i are distinct modulo n, then the property holds. The authors also suggested a general decision problem of checking a number system,

Date: February 19, 2025.

²⁰²⁰ Mathematics Subject Classification. 11B75, 20M05, 39A06.

Key words and phrases. number system, semigroup freeness, refinement equation, subdivision scheme, Borel measure.

¹The condition (1) for the most significant digit is set to generalize the definition of uniqueness from [22] to the number systems containing zero, for example the standard digit system $\{0, 1, 2, ..., n-1\}$.

whether it has the uniqueness property. As it was shown by J. Honkala in [11], this problem is decidable. Some more results concerning number systems were obtained within the next decade; for their description we refer the reader to [12].

Even a little earlier, in [15] D. Klarner came up to a study of another decision problem, which is closely related to the uniqueness of number system. The motivation was coming from Erdös question on when the density of an orbit set $\langle F : S \rangle$ is positive (see [9, p.23] and [19]). Here the set $\langle F : S \rangle$ contains all images of a finite set of non-negative integers Sunder an application of any number of integer-valued affine functions taken from the set

(2)
$$F = \{f_i = n_i x + a_i, i = 1, \dots, m\}, \text{ with } n_i \ge 2 \text{ and } a_i \in \mathbb{N} \cup \{0\}$$

Klarner studied a particular case when all slopes n_i are equal to some integer $n \ge 2$ and noted that for m = n the (upper) density

$$\delta(\langle F:S\rangle) = \lim \sup_{T \to \infty} \frac{\langle F:S \rangle \cup \{0, \dots T\}}{T+1}$$

is equal to 0 if and only if the set F has a non-trivial semigroup relation, i.e.

(3)
$$f_{u_1} \circ \ldots \circ f_{u_p} = f_{v_1} \circ \ldots \circ f_{v_q},$$

for some $u_1, ..., u_p, v_1, ..., v_q \in \{1, ..., m\}$ such that $(u_1, ..., u_p) \neq (v_1, ..., v_s)$.

The decision problem in which, for an input set F, one needs to answer whether a relation of the form (3) exists, is known as a *freeness problem for a semigroup*.²

Klarner noted, that if all slopes in (2) are equal to n, the inequality m > n implies that there exists a relation of type (3); for non-equal slopes this was generalized in [16] and [18]. The important subcase m = n was considered separately, but for its resolution the reader was referred to a preprint, which has never appeared. However, it was noted that the freeness can be obtained once the set

$$\{0, (a_2 - a_1)/d, \dots, (a_m - a_1)/d\}$$

is a complete residue system modulo n, where $d = \text{gcd}(a_2 - a_1, \ldots, a_m - a_1)$. As we will see in Section 4, in the case of prime n = m, the freeness and this condition for residues are actually equivalent.

The uniqueness problem for a number system and the freeness problem for a set of integervalued affine functions of the same slope can be related by the following observation.

Proposition 1. Let $F = \{f_i(x) = nx + a_i \mid i = 1, ..., m\}$ be the set of affine functions such that $n, a_2, ..., a_m \in \mathbb{N}$ with $n \ge 2$ and $a_1 = 0$. Then the n-ary system $A = \{a_1, ..., a_m\}$ has uniqueness property if and only if the set F is a free semigroup basis.

Proof. Suppose first that there is a semigroup relation (3). A straightforward computation shows that $f_{u_1} \circ \ldots \circ f_{u_p} = n^p x + c = f_{v_1} \circ \ldots \circ f_{v_q} = n^q x + d$, where

(4)
$$c = a_{u_1} + a_{u_2}n + a_{u_3}n^2 + \ldots + a_{u_p}n^{p-1}$$
$$d = a_{v_1} + a_{v_2}n + a_{v_2}n^2 + \ldots + a_{v_n}n^{q-1}.$$

Since c = d, from (4) we get two non-equal A-expansions of the same number.

Suppose that there exist two tuples $(a_{u_1}, \ldots, a_{u_p})$ and $(a_{v_1}, \ldots, a_{v_q})$ giving A-expansions of the same number c = d in (4). Without loss of generality, we suppose $p \ge q$ and consider $S = f_1^{(p-q)} = n^{p-q}x$. Then we immediately obtain, that $L = f_{u_1} \circ \ldots \circ f_{u_p} = n^p + c$ and $R = f_{v_1} \circ \ldots \circ f_{v_q} \circ S = n^p + d$, and L = R is a non-trivial semigroup relation since $u_p \ne 0$.

²This problem is known to be undecidable for integer upper-triangular 3×3 matrices (see [17] and [1]), and it is still open for integer affine functions, which can be presented by 2×2 upper-triangular matrices.

The restriction $a_1 = 0$ in the statement of Proposition 1 is actually somewhat artificial, and it can be bypassed by the following tool:

Proposition 2. Let $F = \{f_i(x) = nx + a_i \mid i = 1, ..., m\}$ be a set of affine functions such that $n, a_1, a_2, ..., a_m \in \mathbb{Z}$ and $n \ge 2$. Then for any $s \in \mathbb{Z}$ the set F is a free semigroup basis if and only if the set $F = \{\tilde{f}_i(x) = nx + (a_i - s) \mid i = 1, ..., m\}$ is a free semigroup basis.

Proof. For the function g(x) = x - c, we have $g^{-1} \circ f_i \circ g = nx + (a_i - cn + c)$. Let us take c = s/(n-1), then $g^{-1} \circ f_i \circ g = \tilde{f}_i$. Consequently, the existence of a group relation

$$f_{u_1} \circ \ldots \circ f_{u_p} = f_{v_1} \circ \ldots \circ f_{v_q}.$$

is equivalent to the existence of the conjugated relation

$$g^{-1} \circ f_{u_1} \circ \ldots \circ f_{u_p} \circ g = g^{-1} \circ f_{v_1} \circ \ldots \circ f_{v_q} \circ g$$

which can be also written as

$$\tilde{f}_{u_1} \circ \ldots \circ \tilde{f}_{u_p} = \tilde{f}_{v_1} \circ \ldots \circ \tilde{f}_{v_q}.$$

Propositions 1 and 2 together with Theorem 2 from Section 2 will provide an algorithm to check freeness for the set F in the case m = n, which was missing in Klarner's considerations. If n is prime, the freeness check for the set F (and for the density $\delta(\langle F : S \rangle)$) to be positive) can be done much easier by the use of Theorem 4 from Section 4.

2. Subdivision scheme and transition operator

Subdivision schemes are iterative algorithms for linear approximation of functions from their values on a mesh. They originated in late 1980s with N.Dyn, D.Levin, J.Gregory, C.De Boor, S.Dubuc, etc., see [2, 7, 8] for many references. They are widely used now for interpolation and approximation of smooth functions and in modelling of curves and surfaces. Some special cases of subdivisions appeared earlier in works of G. De Rham (cutting angle scheme) [4], G.Chaikin [3], etc. We consider only stationary univariate schemes defined by an integer contraction factor $n \ge 2$ and by a finitely supported sequence of real numbers $\{c_i\}_{i\in\mathbb{Z}}$. After a suitable shift of numeration it can be assumed that $c_i = 0$ for all $i \notin \{0, \ldots, N\}$ and $c_0 c_N \ne 0$, when $N \ge 1$. We denote by l_{∞} the space of bounded sequences $\{x_k\}_{k\in\mathbb{Z}}$ and by $\delta \in l_{\infty}$ the sequence $(\delta)_k = \delta_{0k}$ (the Kronecker symbol).

The subdivision operator $S: l_{\infty} \to l_{\infty}$ acts on l_{∞} as follows:

(5)
$$(Sg)_k = \sum_{i \in \mathbb{Z}} c_{k-ni} g_i$$

where $g = (g_i)_{i \in \mathbb{Z}} \in l_{\infty}$. Furthermore, we consider the *transition operator* T on the space of compactly supported tempered distributions \mathcal{S}'_0 :

(6)
$$[Tf](t) = \sum_{k \in \mathbb{Z}} c_k f(nt-k), \qquad f \in \mathcal{S}'_0.$$

There is a simple relation between the subdivision scheme and the transition operator. For every $j \in \mathbb{N}$, we have:

(7)
$$[T^{j}f](t) = \sum_{k \in \mathbb{Z}} \left(S^{j}\delta \right)_{k} f(n^{j}t - k), \qquad f \in \mathcal{S}'.$$

The proof can be found, for example, in [2] or verified by a direct computation. The following theorem is well known [24]:

Theorem A. Assume $\sum_k c_k = n$. Then for every $f \in \mathcal{S}'_0$ such that $(f, \mathbf{1}) = 1$, the sequence $T^j f$ converges in \mathcal{S}' to a unique solution $\varphi \in \mathcal{S}'_0$ of the functional equation

(8)
$$\varphi(t) = \sum_{k \in \mathbb{Z}} c_k \varphi(nt - k)$$

such that $(\varphi, \mathbf{1}) = 1$ This solution is supported on the segment [0, N] and satisfies the equation $T\varphi = \varphi$.

Of course, we could have omitted the normalization condition (f, 1) = 1 and obtain by homogeneity: for every $f \in S'_0$ the sequence $T^j f$ converges to $(f, 1)\varphi$. The function φ is called in the literature *refinable function* and equation (8) is called a *refinement equation*.

To every refinement equation we associate the trigonometric polynomial

$$\boldsymbol{c}(\xi) \quad = \quad \frac{1}{n} \sum_{k \in \mathbb{Z}} c_k e^{-2\pi i k \xi}$$

called *mask*. This is the characteristic function of the sequence $\{c_k\}_{k\in\mathbb{Z}}$. Computing the Fourier transform of both parts of the refinement equation gives the following equation in the frequency domain:

(9)
$$\widehat{\varphi}(\xi) = c(\xi/n)\,\widehat{\varphi}(\xi/n)$$

Iterating j times we get

(10)
$$\widehat{\varphi}(\xi) = \boldsymbol{c} \left(n^{-1} \xi \right) \cdots \boldsymbol{c} \left(n^{-j} \xi \right) \widehat{\varphi}(n^{-j} \xi).$$

Now we focus on the case of nonnegative coefficients c_k . Some of the results below are known [26], we give their proofs for the convenience of the reader. The following simple observation has been made independently in a number of papers [6, 7, 26, 28].

Proposition 3. If all c_k are nonnegative, then φ is a Borel measure of pure type, namely, it is either absolutely continuous (i.e., $\varphi \in L_1$) or purely singular.

Proof. Choosing $f = \chi_{[0,1]}$, we see that the functions $T^j f$ are all non-negative and, hence, so is their limit. Thus, φ is a nonnegative distribution, that is a Borel measure.

According to the Lebesgue theorem, there exists a unique representation of φ in the sum of absolutely continuous and singular measures $\varphi = \varphi_{cont} + \varphi_{sing}$. The uniqueness implies that both φ_{cont} and φ_{sing} satisfy the refinement equation (8). However, this equation possesses a unique solution up to normalization. Therefore, one of those functions is zero.

Proposition 3 rises the question of separating the cases of absolute continuity and singularity of the measure φ . The following criterion is proved by applying (10) and the Poisson summation formula:

Proposition 4. Suppose all c_k are nonnegative; then $\varphi \in L_1$ if and only if $\widehat{\varphi}(m) = 0$ for all integer $m \neq 0$. Moreover, in this case $\varphi(t) \leq 1$ almost everywhere.

Proof. (Necessity). Assume that there exists $m \in \mathbb{Z} \setminus \{0\}$ such that $\widehat{\varphi}(m) \neq 0$. Taking an arbitrary number $j \in \mathbb{N}$ and substituting $\xi = n^j m$ to equation (10), we get

$$\widehat{\varphi}(n^{j}m) = c(n^{j-1}m)\cdots c(m)\widehat{\varphi}(m) = \widehat{\varphi}(m)$$

The latter equality holds because all the numbers $n^{j-s}m$, $s = 1, \ldots, j$, are integer and therefore, $\mathbf{c}(n^{j-s}m) = \mathbf{c}(0) = 1$. Thus, $\widehat{\varphi}(n^{j}m) = \widehat{\varphi}(m)$ for all $j \in \mathbb{N}$. On the other hand, if $\varphi \in L_1(\mathbb{R})$, then $\widehat{f}(\xi) \to 0$ as $\xi \to \infty$. This is not true for $\xi = n^j m, j \to \infty$, hence $\varphi \notin L_1$. (Sufficiency). If $\widehat{\varphi}(m) = \delta_{m0}$, then applying the Poisson summation formula to the function $\varphi(t - \cdot)$ we get

(11)
$$\sum_{k \in \mathbb{Z}} \varphi(t-k) = \sum_{m \in \mathbb{Z}} \widehat{\varphi}(m) e^{2\pi i m t} = \widehat{\varphi}(0) = 1$$

Thus, $\sum_{k\in\mathbb{Z}}\varphi(t-k) = \mathbf{1}$. Since $\varphi \geq 0$, we see that $\varphi \leq \mathbf{1}$, i.e., for every nonnegative test function $f \in \mathcal{S}$, we have $(\varphi, f) \leq \int_{\mathbb{R}} f dt$. Hence, φ is majorized by the Lebesgue measure and, therefore $\varphi \in L_1$.

The criterion of Proposition 4 has a disadvantage that it involves the function $\widehat{\varphi}$, which is a priori unknown. A criterion in terms of the coefficients $\{c_k\}_{k\in z}$ exploits the *n*-ary tree. We define the tree \mathcal{T} as follows. The root is associated to zero and has n-1 children with the numbers $\frac{k}{n}$, $k = 1, \ldots, n-1$. The further construction is by induction: every vertex α , apart from the root, has *n* children $\frac{\alpha+k}{n}$, $k = 0, \ldots, n-1$. The root has level zero, its children form the first level, etc. Thus, the *j*th root consists of $(n-1)n^{j-1}$ numbers $n^{-j}k$, k = $1, \ldots, n^j - 1$, $k \not\equiv 0 \pmod{n}$. A subset \mathcal{A} of vertices of \mathcal{T} is called a *minimal cut set* if every infinite path along the tree from the root (all paths are without backtracking, the root is not in the path) has exactly one common vertex with \mathcal{A} . All minimal cut sets are finite. The simplest one is the set of vertices of the first level: $\mathcal{A} = \{\frac{1}{n}, \ldots, \frac{n-1}{n}\}$.

Proposition 5. Suppose all c_k are nonnegative; then $\varphi \in L_1$ if and only if there exists a minimal cut set of the n-ary tree \mathcal{T} that consists of roots of the mask c.

Proof. (Necessity). Since φ is compactly supported, it follows from the Paley-Wiener theorem that \widehat{f} is an entire function. Therefore, it has finitely many zeros, if any, on the unit disc. This implies that there exists $j \geq 1$ such that $\widehat{\varphi}$ does not have zeros on the *j*th level of \mathcal{T} , t.e., $\mathbf{c}(n^{-j}m) \neq 0$ for every natural $m < n^j$, $m \not\equiv 0 \pmod{n}$. Let $m = \sum_{k=0}^j d_k n^k = d_j \dots d_0$ be the standard *n*-adic expansion of *m*, possibly, starting with zeros, $d_0 \neq 0$. Add the digit $d_{j+1} = 0$ and substitute $\xi = m$ to equation (10). We obtain

(12)
$$\widehat{\varphi}(m) = \widehat{\varphi}(n^{-j}m) \prod_{q=0}^{j} c \left(d_{j+1} \dots d_{j-q+1} \dots d_{j-q} \dots d_{0} \right) = \widehat{\varphi}(n^{-j}m) \prod_{q=0}^{j} c \left(0 \dots d_{j-q} \dots d_{0} \right).$$

Since $\widehat{\varphi}(m) = 0$ and $\widehat{\varphi}(n^{-j}m) \neq 0$, we see that one of the numbers $n^{q-j}m = 0.d_{j-q}\ldots d_0$ must be a root of \boldsymbol{c} . Those numbers form a finite path of length j. Thus, every path of length j contains a root of the polynomial \boldsymbol{c} .

(Sufficiency). Assume there exists a minimal cut set $\mathcal{A} \subset \mathcal{T}$ such that $\mathbf{c}(\mathcal{A}) = 0$. Then taking an arbitrary natural number m and applying formula (12) we obtain $\widehat{\varphi}(m) = 0$, which completes the proof for positive m. The proof for negative m is the same.

3. CRITERION OF UNIQUENESS FOR A NUMBER SYSTEM

Now we consider a set of digits $A = \{a_1, \ldots, a_n\}$ of the (non-standard) *n*-ary number system. We will assume that $0 = a_1 < \ldots < a_n$. For any integer $k \ge 0$ we denote by b(k) the total number of its A-expansions having form (1) and set formally b(k) = 0 for any integer k < 0.

The uniqueness expansion property means that $b(k) \leq 1$ for all $k \in \mathbb{Z}$. We are going to see that the sequence $\{b(k)\}_{k\in\mathbb{Z}}$ is generated by a subdivision operator S with the following

coefficients:

(13)
$$c_i = \begin{cases} 1 & , i \in A \\ 0 & , \text{ otherwise} \end{cases}$$

This sequence will be referred to as an *indicator sequence of* A. We start with the following simple observation:

Lemma 1. The sequence b(k) satisfies the following recurrent relations:

(14)
$$b(nq+d) = \sum_{s \in \mathbb{Z}} c_{ns+d} b(q-s), \quad d = 0, \dots, n-1, \qquad q \ge 0$$

Proof. In expansion (1) for k = nq+d, the digit $\alpha_0 \in A$ must be such that $\alpha_0 \equiv d \pmod{n}$, hence $\alpha_0 = ns + d$ for some s. Subtracting d from both sides of (1) and dividing by n, we get $q - s = \sum_j \alpha_j n^{j-1}$, provided that $ns + d \in A$, i.e., $\alpha_{ns+d} = 1$. Thus, the total number of expansions of the number nq + d is equal to the sum of numbers of expansions of q - s over all s such that $\alpha_{ns+d} = 1$. This completes the proof.

If we denote k = nq + d and i = q - s, then the equation (14) becomes

(15)
$$b(k) = \sum_{i \in \mathbb{Z}} c_{k-ni} b(i).$$

The right hand side is precisely the subdivision operator (5). Thus, we obtain

Corollary 1. Let $\mathbf{b}^{(j)} \in l_{\infty}$ be the sequence $\{b(k)\}_{k=0}^{n^{j}-1}$, complemented by zeros for k < 1 and for $k \geq n^{j}$. Then $\mathbf{b}^{(j)} = S\mathbf{b}^{(j-1)}$.

Theorem 1. For every $j \ge 1$, we have

(16)
$$b(k) = (S^j \delta)_k, \quad k = 0, \dots, n^j - 1$$

where S is the subdivision operator defined by the indicator sequence of A.

Thus, the sequence $\boldsymbol{b}^{(j)}$ defined in Corollary 1 coincides with the sequence $S^{j}\delta$.

Proof. We argue by induction in j. For j = 0, we have b(0) = 1 and $(S^0\delta)_0 = (\delta)_0 = 1$, so, this is true. The transfer $j - 1 \mapsto j$ is provided by Corollary 1.

Now we formulate the fundamental result. Let $\{c_k\}_{k\in\mathbb{Z}}$ be the indicator sequence of A, $c(\cdot)$ be the mask of this sequence, \mathcal{T} be the *n*-ary tree.

Theorem 2. The following assertions are equivalent

a) Every natural number possesses at most one n-ary expansion with the digit set A;

b) The compactly supported solution of the refinement equation $\varphi(t) = \sum_{i=1}^{n} \varphi(nt-a_i)$ belongs to $L_1(\mathbb{R})$;

c) The n-ary tree \mathcal{T} possesses a minimal cut set that consists of roots of the polynomial $\mathbf{c}(t) = \frac{1}{n} \sum_{k=1}^{n} e^{-2\pi i a_k t}$.

Proof. The equivalence of b) and c) follows from Proposition 5. Let us prove the equivalence of a) and b).

a) \Rightarrow **b**). Assume the contrary: the property a) holds, i.e., $b(k) \leq 1, k \in \mathbb{Z}$, but $\varphi \notin L_1$. The latter means that φ is a purely singular Borel measure (Proposition 3). Consider the transition operator $[Tf](t) = \sum_{i=1}^{n} \varphi(nt - a_i)$ and take $f = \chi_{[0,1]}$. By equation (7), we have

$$[T^{j}f](t) = \sum_{k \in \mathbb{Z}} \left(S^{j}\delta\right)_{k} f(n^{j}t - k) = \sum_{k \in \mathbb{Z}} \left(S^{j}\delta\right)_{k} \chi_{[n^{-j}k, n^{-j}(k+1)]}.$$

On the other hand, $T^j f \to \varphi$ due to Theorem A. The convergence is in \mathcal{S}' , which, for nonnegative distributions, means the convergence in measure. Applying now Theorem 1 we obtain $(S^j \delta)_k = b(k)$ for all $k \leq n^j - 1$. Since the segments $[n^{-j}k, n^{-j}(k+1)]$ cover the segment [0, 1] when k runs from zero to $n^j - 1$, we conclude that the restriction of the function

$$\sum_{k \in \mathbb{Z}} \left(S^j \delta \right)_k \chi_{[n^{-j}k, n^{-j}(k+1)]}$$

to the segment [0, 1] is equal to

$$f_j = \sum_{k=0}^{n^j - 1} b(k) \, \chi_{[n^{-j}k, n^{-j}(k+1)]} \, .$$

Therefore,

(17)
$$f_j = T^j f \Big|_{[0,1]} \to \varphi \Big|_{[0,1]} \quad \text{as} \quad j \to \infty.$$

By Proposition 3, the function φ is of pure type, hence, the assumption $\varphi \notin L_1$ implies that φ is purely singular. So, its restriction to the segment [0, 1] is purely singular as well. On the other hand, if $b(x) \leq 1$ for all k, then the function on the right hand side of (17) does not exceed one and hence, so its limit $\varphi|_{[0,1]}$. Thus, the function $\varphi|_{[0,1]}$ is majorized by the Lebesgue measure and is not purely singular.

b) \Rightarrow **a**). Assume that there is a number that has at least two *A*-expansions, the longest of which contains *r* digits. Let $A_r = \left\{ \sum_{j=0}^{r-1} \alpha_j n^j, \alpha_j \in A, j \leq r \right\}$. Since $a_1 = 0$, the set A_r contains all *A*-presentable numbers of length at most *r*. Hence, at least two elements of A_r coincide, so we have $|A_r| < n^r$. Consider the *r*th power of the transition operator:

(18)
$$[T^r f](t) = \sum_{k \in A_r}^n b_r(k)\varphi(n^r t - k),$$

where $b_r(k)$ is the total number of A-expansions of the number k with at most r digits. Clearly, $T^r \varphi = \varphi$, hence, the refinement equation with the transition operator T_r possesses the same solution φ . By the Hutchinson theorem [13], there is a unique compact set $Q \subset \mathbb{R}$ such that $Q = \bigcup_{k \in A_r} n^{-r}(Q+k)$. This is a fractal generated by affine contractions $n^{-r}(\cdot + k)$. The Lebesgue measure $\mu(Q)$ does not exceed $\sum_{k \in A_r} n^{-r} \mu(Q+k) = |A_r| n^{-r} \mu(Q)$. Since $n^{-r}|A_r| < 1$, it follows that $\mu(Q) = 0$. On the other hand, T_r respects the set of Borel measures supported on Q. Taking an arbitrary such measure ψ normalized by the condition $(\psi, \mathbf{1}) = 1$ and applying Theorem A, we obtain $T^{rj}\psi \to \varphi$ as $j \to \infty$. Therefore, φ is also supported on Q and hence, is purely singular.

Remark 1. Now we see that Theorem 2 together with Proposition 1 and Proposition 2 provide an algorithm to check freeness for a set $F = \{f_i = nx + a_i, i = 1, ..., n\}$, with $n, a_1, \ldots, a_n \in \mathbb{Z}$ and $n \geq 2$. Indeed, the degree of the trigonometric polynomial $\sum_{k=1}^{n} e^{-2\pi i a_k t}$ is bounded from above, so is the cardinality of its roots on the segment [0, 1]. Thus, the possible minimal cut sets of the *n*-ary tree in Theorem 2 also have bounded size, which in turn bounds the height of possible tree vertices in the minimal cut. It follows that the check of all such vertices can be done through their finite enumeration and polynomial evaluation in the corresponding points.

4. UNIQUENESS OF THE EXPANSION AND DIGITS RESIDUES

In this Section we will use the criterion from Theorem 2 to study the uniqueness property for the *n*-ary number system consisting of exactly *n* digits. Let again $A = \{a_1, \ldots, a_n\}$ be the set of nonnegative digits such that $0 \le a_1 < \cdots < a_n$. We will say that *n*-ary number system *A* is *irreducible* if

(19)
$$gcd(a_2 - a_1, \dots, a_n - a_1, n) = 1.$$

Since zero is allowed to be present in the set of digits, we remind that A-expansion of a number k has the form

(20)
$$k = \sum_{j=0}^{J} \alpha_j n^j, \text{ where } a_J \neq 0.$$

If k has at least one such representation, we will say that k is a representable number.

We will say that n-ary number system A has weak uniqueness property if two A-expansions of the same length represent different numbers. The following statement is useful for working with sets of digits that contain only positive numbers.

Proposition 6. Let $A = \{a_1, \ldots, a_m\}$ be a set of digits such that $0 < a_1 < a_2 < \ldots < a_m$. Then the n-ary system A has weak uniqueness property if and only if the n-ary system $B = \{0, a_2 - a_1, \ldots, a_m - a_1\}$ has uniqueness property.

Proof. We prove both directions by contraposition. First, if A does not have weak uniqueness property, then there exist two tuples $U = (u_0, u_1, \ldots, u_s)$ and $V = (v_0, v_1, \ldots, v_s)$ from A^s such that

$$\sum_{j=0}^{s} u_j n^j = \sum_{j=0}^{s} v_j n^j,$$

and by subtraction of $a_0(1 + n + \ldots + a^s)$ from both sides we obtain

$$\sum_{j=0}^{s} (u_j - a_0) n^j = \sum_{j=0}^{s} (v_j - a_0) n^j,$$

which gives two distinct tuples U' and V' from B^p . By removing most significant digits until they are non-zero, we obtain two tuples refuting the uniqueness property for the system B.

Now, suppose that the *n*-ary system B does not have the uniqueness property and there exists $U = (u_0, u_1, \ldots, u_s) \in B^s$ and $V = (v_0, v_1, \ldots, v_t) \in B^t$ such that

(21)
$$\sum_{j=0}^{s} u_j n^j = \sum_{j=0}^{t} v_j n^j.$$

Without loss of generality, we may suppose that $s \ge t$. Consider two tuples $U', V' \in A^{s+1}$ obtained as $U' = (u_0+a_0, u_1+a_0, \ldots, u_s+a_0)$ and $(v_0+a_0, v_1+a_0, \ldots, v_t+a_0, a_0, \ldots, a_0)$. They represent the same number in the system A, obtained from (21) by adding $a_0(1+n+\ldots+a^s)$.

Now we are ready to prove two results on how the uniqueness property of the n-ary system with n digits is related to their residues modulo n.

Theorem 3. Let n be a composite number. Then there exists an irreducible set A consisting of n non-negative digits such that the n-ary number system A possesses the uniqueness property, but A contains two numbers congruent modulo n.

Proof. Since n is a composite number, there exist integers n_1, n_2 , both larger than 1 such that $n = n_1 n_2$. We construct the following set of digits

(22)
$$A = \{un_1n + v : 0 \le u < n_2, 0 \le v < n_1\}.$$

As $n_1 > 1$, then by taking u = 0 we get digits $a_1 = 0$, $a_2 = 1$, so the condition (19) holds. Since n_1 divides un_1n and $0 \le v < n_1$ in (22), no digit can be congruent to n_1 modulo n. Hence, the digits do not form the complete system of residues modulo n.

From (22) it follows that no digits a, a' can satisfy the congruence $a - a' \equiv un_1 \pmod{n}$ for $1 \leq u < n_2$. Then, considering the difference of two expansions of type (20) modulo nwe obtain that if y, y' are representable numbers, we also have

(23)
$$y - y' \not\equiv un_1 \pmod{n}$$
, when $1 \le u < n_2$.

Now we will prove that any positive integer has at most one expansion of type (20). Assume the contrary, and let x be the least positive integer having at least two expansions

$$x = \sum_{j=0}^{J} \alpha_j n^j = \sum_{j=0}^{J'} \alpha'_j n^j \text{ with } \alpha_J \neq 0 \text{ and } \alpha'_{J'} \neq 0.$$

Then the last digits α_0 and α'_0 must be distinct: otherwise the number $(x - \alpha_0)/n$ also has two distinct expansions, but this contradicts the supposition of minimality for chosen x. Hence, $\alpha_0 \neq \alpha'_0$. Without loss of generality we will presume that $\alpha_0 < \alpha'_0$. Then we have

$$x = ny + \alpha_0 = ny' + \alpha'_0$$

and y > y' are some representable numbers. Thus, $\alpha'_0 - \alpha_0 = n(y - y')$, hence

(24)
$$\alpha'_0 - \alpha_0 = 0 \pmod{n}.$$

Then, if the digits α'_0 and α_0 were formed in A as $\alpha'_0 = u_1n_1n + v_1$ and $\alpha' = u_2n_1n + v_2$, from (24) we obtain $v_1 = v_2$, hence $\alpha'_0 - \alpha_0 = (u_2 - u_1)n_1n$. Therefore, $y - y' = (u_2 - u_1)n_1$ with $1 \le u_2 - u_1 < n_2$, which contradicts (23). This proves the uniqueness of the expansion.

Theorem 4. For an arbitrary prime number p the following holds: an irreducible p-ary number system A has a uniqueness property, if and only if the set A does not contain two numbers which are congruent modulo p.

Proof. The sufficiency is straightforward (e.g., [22, Theorem 4]), so we prove the necessity.

Let $a_1 < \cdots < a_p$ be an irreducible set of non-negative digits such that every number has at most one representation in base p with digits a_1, \ldots, a_p . We will show that in this case a_1, \ldots, a_p form a complete system of residues modulo p. To do so, we will use Theorem 2. For any tuple $\mathbf{c} = (c_1, \ldots, c_p) \in \mathbb{Z}_{>0}^p$ we define the polynomial

$$P_{\boldsymbol{c}}(z) = \sum_{j=1}^{p} z^{c_j}$$

For $m \in \mathbb{N}$, a complex number z is a primitive p^m th root of unity if $z^{p^m} = 1$, but $z^{p^{m-1}} \neq 1$. Denote $\mathbf{a} = (a_1, \ldots, a_p)$ and $\mathbf{b} = (0, a_2 - a_1, \ldots, a_p - a_1)$. Using Proposition 6 we can obtain that the uniqueness property for the *n*-ary system A implies the uniqueness for the system $B = \{0, a_2 - a_1, \ldots, a_p - a_1\}$, to which we can apply Theorem 2. The direction $(a) \Rightarrow (c)$ implies that some root of the polynomial $P_{\mathbf{b}}$ is a primitive p^m th root of unity for some $m \in \mathbb{N}$, from which it follows that for some m all primitive p^m th roots of unity are roots of $P_{\mathbf{b}}$. Since $P_{\mathbf{a}} = e^{-2\pi i a_0} P_{\mathbf{b}}$, the latter is also true for the polynomial $P_{\mathbf{a}}$. This will suffice for our purpose. Recall that z is a p^m th primitive root of unity if and only if it is a root of the cyclotomic polynomial

$$Q(z) = Q_{p^m}(z) = \sum_{j=0}^{p-1} z^{jp^{m-1}}.$$

Next, we observe that for any $a, \tilde{a} \in \mathbb{Z}_+$ such that $a \geq \tilde{a}$ and $a \equiv \tilde{a} \pmod{p^m}$, the polynomial $z^a - z^{\tilde{a}}$ is divisible by $z^{p^m} - 1$ which in turn is divisible by Q. Hence, all primitive p^m th roots of unity are roots of $z^a - z^{\tilde{a}}$. For any $j = 1, \ldots, p$ we take $0 \leq \tilde{a}_j < p^m$ so that $\tilde{a}_j \equiv a_j \pmod{p^m}$. Denote

$$\tilde{P}(z) = \sum_{j=1}^{p} z^{\tilde{a}_j}$$

Then the primitive p^m th roots of unity are roots of \tilde{P} . This implies that $\tilde{P} = QR$, where R is an integral polynomial of degree $< p^{m-1}$. Moreover, all coefficients of R are nonnegative. We have $\tilde{P}(1) = Q(1) = p$. Hence, R(1) = 1, and $R(z) = z^u$ for some u with $0 \le u < p^{m-1}$.

We have $\tilde{P}(1) = Q(1) = p$. Hence, R(1) = 1, and $R(z) = z^u$ for some u with $0 \le u < p^{m-1}$. If m = 1, then R(z) = 1, $\tilde{P}(z) = \sum_{j=0}^{p-1} z^j$. We see that the multiset $\{\tilde{a}_1, \ldots, \tilde{a}_p\}$ is actually the set $\{0, \ldots, p-1\}$. This means that a_1, \ldots, a_p form the complete system of residues modulo p as desired.

Let m > 1. Then all numbers \tilde{a}_j are congruent to u modulo p^{m-1} . Consequently, all numbers a_j are also congruent to u modulo p^{m-1} . This does not agree with the irreducibility of the set $\{a_1, \ldots, a_p\}$.

Remark 2. If *n* is prime, Theorem 4 provides an easy freeness test for a set of functions $F = \{f_i = nx + a_i, i = 1, ..., m\}$, with $n \ge 2$ and $a_i \in \mathbb{N} \cup \{0\}$. First, we increasingly order the set $A = \{a_1, \ldots, a_n\}$ and compute $d = \gcd(a_2 - a_1, \ldots, a_p - a_1, p)$. Then, the set $\{0, (a_2 - a_1)/d, \ldots, (a_p - a_1)/d\}$ is an irreducible *p*-ary number system, and we are left to check whether its elements are all distinct modulo *p*. Here, the correctness of scaling the coefficients by *d* can be proved similarly to the correctness of the shift in Proposition 2 by the use of conjugation function g(x) = dx.

5. Acknowledgements

The work of the second author and the third author was prepared with the support of the Theoretical Physics and Mathematics Advancement Foundation "BASIS", grants No. 22-7-1-20-1 and No. 22-7-2-32-1 respectively.

References

 J. Cassaigne, T. Harju, J. Karhumäki, On the Undecidability of Freeness of Matrix Semigroups, Int. J. Algebra Comput. 9 (3–4), 295–305 (1999)

[2] A. S. Cavaretta, W. Dahmen, and C.A. Micchelli, *Stationary subdivision*, Mem. Amer. Math. Soc. 93 (1991), no. 453.

- [3] G.M. Chaikin, An algorithm for high speed curve generation, Comp. Graphics and Image. Proc., 3 (1974), 346–349.
- [4] G. de Rham, Sur une courbe plane, J. Math. Pures Appl. 35 (1956), 25-42.
- [5] G. de Rham, Sur les courbes limites de polygones obtenus par trisection, Enseignement Math. 5 (1959), 29-43.
- [6] G.A. Derfel, Probabilistic method for a class of functional-differential equations Ukrain. Math. J., 41 (1989), 1137--1141.

 [7] N. Dyn, J.A. Gregory, and D. Levin, Analysis of uniform binary subdivision schemes for curve design, Constr. Approx. 7 (1991), 127–147.

[8] N. Dyn and D. Levin, Subdivision schemes in geometric modelling, Acta Numer., 11 (2002), 73–144.

- [9] P. Erdos, R. Graham, Old and new problems and results in combinatorial number theory, L'Enseignement Math., Monogr., Vol. 28, 1980.
- [10] D.-J. Feng and N. Sidorov, Growth rate for beta-expansions, Monatsh. Math. 162 (2011), 41-60
- [11] J. Honkala, Unique representation in number systems and L codes, Discrete Applied Mathematics 4 (1982), 229–232.
- [12] J. Honkala, On Number Systems with Finite Degree of Ambiguity, Information and Computation 145 (1998), 51–63.
- [13] J. E. Hutchinson, Fractals and self-similarity, Indiana Univ. Math. J., 30 (1981), no 5, 713–747.
- [14] J. Jankauskas and J.M. Thuswaldner, *Rational matrix digit systems*, Linear and Multilinear Alg., 71 (2023), no 10, 1606—1639.
- [15] D. A. Klarner, An Algorithm to Determine When Certain Sets Have O-Density, Journal of Algorithms 2 (1981), 31–43.
- [16] D.A. Klarner, A Sufficient Condition for Certain Semigroups to Be Free" J. Algebra 74, 140–148 (1982).
- [17] D.A. Klarner, J.C. Birget, W. Satterfield, On the undecidability of the freeness of integer matrix semigroups, International J. of Algebra and Computation 1 (1991) 223-226.
- [18] A. Kolpakov, A. Talambutsa, On free semigroups of affine maps on the real line, Proc. Amer. Math. Soc., 150:6 (2022), 2301–2307.
- [19] J. C. Lagarias, "Erdös, Klarner, and the 3x+1 problem", Amer. Math. Monthly 123 (8), 753–776 (2016)
- [20] J. C. Lagarias and Y.Wang, Integral self-affine tiles in \mathbb{R}^n . I. Standard and nonstandard digit sets J. London Math. Soc., 54 (1996), no 1, 161–179.
- [21] J.-L. Li, *Digit sets of integral self-affine tiles with prime determinant*, Studia Mathematica, 177 (2006), no 2, 183–194.
- [22] H.A. Maurer, A. Salomaa, D. Wood, *L codes and number systems*, Theoretical Computer Science 22 (1983), 331–346.
- [23] C.A. Micchelli and H. Prautzsch, Uniform refinement of curves, Lin. Alg. Appl. 114/115 (1989), 841–870.
- [24] I.Ya. Novikov, V.Yu. Protasov, and M.A. Skopina, Wavelets theory, AMS, Translations Mathematical Monographs, 239 (2011).
- [25] V. Yu. Protasov, Asymptotic behaviour of the partition function, Sb. Math., 191 (2000), No 3-4, 381-414.
- [26] V. Yu. Protasov, *Refinement equations with nonnegative coefficients*, J. Fourier Anal. Appl., 6 (2000), no 1, 55–78.
- [27] V.Yu. Protasov, On the asymptotics of the binary partition function, Math. Notes, 76 (2004), no 1, 151–156.
- [28] O.K. Zakusilo, Some properties of classes Lc of limit distribution, Teoria Veroyatnosti i Mat. Statistika, 15 (1976), 68–73.

STEKLOV MATHEMATICAL INSTITUTE OF RAS, 8 GUBKINA ST., 119991 MOSCOW, RUSSIA *Email address:* konyagin@mi-ras.ru

DISIM, UNIVERSITY OF L'AQUILA, 67100 L'AQUILA, ITALY Moscow State University, Moscow, 119991, Russia *Email address*: vladimir.protasov@univaq.it

STEKLOV MATHEMATICAL INSTITUTE OF RAS, 8 GUBKINA ST., 119991 MOSCOW, RUSSIA HSE UNIVERSITY, LABORATORY OF THEORETICAL COMPUTER SCIENCE, 11 POKROVSKY BLVD., 109028 MOSCOW, RUSSIA *Email address*: altal@mi-ras.ru