Near-Optimal Private Learning in Linear Contextual Bandits

Fan Chen fanchen@mit.edu Jiachun Li jiach334@mit.edu Alexander Rakhlin rakhlin@mit.edu David Simchi-Levi dslevi@mit.edu

February 19, 2025

Abstract

We analyze the problem of private learning in generalized linear contextual bandits. Our approach is based on a novel method of *re-weighted* regression, yielding an efficient algorithm with regret of $\tilde{O}\left(d^2\sqrt{T} + \frac{d^{5/2}}{\alpha}\right)$ and $\tilde{O}\left(\sqrt{d^5T}/\alpha\right)$ in the joint and local model of privacy, respectively. Further, we provide near-optimal private procedures that achieve dimension-independent rates in private linear models and linear contextual bandits. In particular, our results imply that joint privacy is almost "for free" in all the settings we consider, partially addressing the open problem posed by Azize and Basu [2024].

1 Introduction

Contextual bandits provide a natural framework for interactive decision making, applicable to numerous real-world domains. In this setting, the decision maker (or, the algorithm) sequentially observes a context, selects an action, and receives a reward [Abbasi-Yadkori et al., 2011, Auer et al., 2002, Simchi-Levi and Xu, 2020, Foster and Rakhlin, 2020]. The central challenge is to balance exploration (learning the reward structure) and exploitation (maximizing the cumulative rewards). Further, in many applications, there are additional privacy concerns, as the contexts often involve sensitive personal information—such as past purchase histories, credit scores, or physcial data—information not meant for public disclosure [Dwork et al., 2014, Lei et al., 2024, Chen et al., 2022]. Despite extensive research on interactive decision making, it is not yet well-understood how to achieve the optimal privacy-utility trade-offs even in the fundamental setting of contextual bandits—arguably one of the simplest and most commonly considered models of online learning [Shariff and Sheffet, 2018, Azize and Basu, 2024].

Formally, in the setting of contextual bandits, the learner observes a context $x_t \in \mathcal{X}$ at each step $t \in [T]$, drawn stochastically as $x_t \sim P$. Based on the context x_t (and the history up to step t), the learner selects an action $a_t \in \mathcal{A}$ and observes a reward $r_t \in [-1, 1]$ with expected value $\mathbb{E}[r_t|x_t, a_t] = f^*(x_t, a_t)$. Here, $f^* : \mathcal{X} \times \mathcal{A} \to [-1, 1]$ is the underlying mean reward function. The performance of the learner is typically measured by its *regret*, defined as

$$\mathbf{Reg} := \mathbb{E}\left[\sum_{t=1}^{T} \max_{a_t^{\star} \in \mathcal{A}} f^{\star}(x_t, a_t^{\star}) - f^{\star}(x_t, a_t)\right],$$

which measures the gap between the learner's cumulative rewards and that of an optimal policy with the full knowledge of f^* . In generalized linear contextual bandits, a widely studied model, the ground-truth f^* is assumed to take the form

$$f^{\star}(x,a) = \nu(\langle \phi(x,a), \theta^{\star} \rangle), \qquad \forall (x,a) \in \mathcal{X} \times \mathcal{A},$$

where $\nu : [-1,1] \to [-1,1]$ is a known link function, $\phi : \mathcal{X} \times \mathcal{A} \to \mathbb{B}^d(1)$ is a known feature map, and $\theta^* \in \mathbb{B}^d(1)$ is the unknown underlying parameter.

Even in linear contextual bandits (where $\nu(t) = t$ is the identity function and f^* is linear), there is limited understanding of how to design privacy-preserving procedures that attain optimal regret. Under the *joint*

Setting	Results	Regret bound	Gen	Adv
Joint DP	Shariff and Sheffet [2018]	$d\sqrt{T} + \frac{d^{3/4}\sqrt{T}}{\sqrt{lpha}}$	~	~
	Theorem 5.1	$\sqrt{d^5T} + \frac{d^{5/2}}{\alpha}$	1	×
	Theorem 5.2 $(\mathcal{A} = O(1))$	$\sqrt{dT} + \frac{d^{3/2}}{\alpha}$	×	×
	Lower bound [He et al., 2022]	$\sqrt{dT \log \mathcal{A} } + \frac{d}{\alpha}$	/	/
Local DP	Zheng et al. [2020]	$(dT)^{3/4}/lpha$	~	>
	Han et al. $[2021]^{\dagger}$	$\sqrt{d\log \mathcal{A} \cdot T}/(\lambda_{\min}^{\star}\alpha)$	~	~
	Li et al. [2024]	$ \mathcal{A} ^2 \log^d(T) \cdot \sqrt{T}/\alpha$	×	×
	Chen and Rakhlin [2025]	$\sqrt{d^3T}/lpha$	~	~
	Theorem 5.3	$\sqrt{d^5T}/lpha$	1	X
	Lower bound [Chen and Rakhlin, 2025]	$\sqrt{d^2T}/lpha$	/	/

Table 1: Summary of the existing results for private learning in (generalized) linear contextual bandits. The "Gen" column indicates whether similar regret bounds can be obtained under generalized linear contextual bandits. The "Adv" column indicates whether the contexts are allowed to be adversarial (as opposed to stochastic contexts considered in the present paper). †The assumptions of Han et al. [2021] are slightly stronger than $\lambda_{\min}^* > 0$, and their regret bound is always lower bounded by $\sqrt{d \log |\mathcal{A}| \cdot T}/(\lambda_{\min}^* \alpha)$ stated in the table. Note that $\lambda_{\min}^* \leq \frac{1}{d}$ always holds.

differential privacy (JDP) model [Dwork et al., 2006] with privacy parameter α , the only known regret upper bound—due to Shariff and Sheffet [2018]—scales as $\sqrt{T/\alpha}$. This regret scaling is particularly undesirable in the high-privacy regime (i.e., $\alpha \ll 1$), and Azize and Basu [2024] have posed the question of identifying the rate-optimal regret in this setting as an open problem.

In the *local differential privacy* (LDP) model [Duchi et al., 2013], the best-known regret bound until recently was $\sqrt{T}/(\alpha \lambda_{\min}^*)$ [Han et al., 2021], where

$$\lambda_{\min}^{\star} := \min_{\pi} \lambda_{\min}(\mathbb{E}^{\pi} \phi(x, a) \phi(x, a)^{\top})$$

is the minimum eigenvalue of the covariance matrix over all linear policies.¹ Assuming λ_{\min}^{\star} being lower bounded implies that the learner can effectively estimate the ground-truth parameter θ^{\star} while executing greedy policies, and hence it essentially removes the difficulty of *exploration*. Such a condition is typically called "explorability" ("diversity") because it assumes the actions taken by any greedy policy are diverse enough. The quantity $1/\lambda_{\min}^{\star}$ can be prohibitively large for most scenarios of interest, e.g., when there exists a direction $v \in \mathbb{R}^d$ that cannot be sufficiently explored, i.e., $\phi(x, a) \perp v$ for each $a \in \mathcal{A}$ with high probability over $x \sim P$. However, it is known that for algorithms based on squared loss regression—such as variants of the LinUCB algorithm [Abbasi-Yadkori et al., 2011]—to achieve the optiaml rates, a dependence on $1/\lambda_{\min}^{\star}$ can be unavoidable (cf. Section 3). Therefore, algorithmic innovations are necessary to attain the optimal regret without assuming such an explorability condition.

Towards the optimal regret under LDP, the recent work of Li et al. [2024] develops an alternative approach based on regression with confidence interval (L_1 -regression). However, as their algorithm involves iteratively performing PCA, the final regret bound scales with $\log^d(T)\sqrt{T}$, i.e., the dependence on d is exponential. On the other hand, Chen and Rakhlin [2025] prove a regret bound of $\sqrt{d^3T}/\alpha$ by controlling the Decision-Estimation Coefficient (DEC) [Foster et al., 2021, 2023], achieved by the Exploration-by-Optimization algorithm [Lattimore and Szepesvári, 2020, Foster et al., 2022] that operates in at least exponential-time.

Building upon these recent advances, we propose a new method called private *re-weighted regression*, which first privately learns a *normalization matrix* U from data, and then performs regression on the loss function,

¹A policy π is *linear* if there exists $\theta \in \mathbb{R}^d$ so that $\pi(x) \in \arg \max_{x \in \mathcal{X}} \langle \theta, \phi(x, a) \rangle$. For generalized linear contextual bandits, it is clear that the optimal policy must be linear.

re-weighted according to the matrix U. In the generalized linear model, the re-weighted regression provides a near-optimal convergence rate in the L_1 -error. Further, using this method as a subroutine, we propose a computationally efficient algorithm that achieves regret bounds of $\tilde{O}\left(d^2\sqrt{T} + \frac{d^{5/2}}{\alpha}\right)$ and $\tilde{O}\left(\sqrt{d^5T}/\alpha\right)$ in the joint and local privacy models, respectively. In particular, our results imply that the joint privacy is almost "for free" in generalized linear contextual bandits, partially resolving the open problem of Azize and Basu [2024].

Furthermore, under the setting where the dimension d is prohibitively large or even unbounded, we develop private estimation procedures that achieve *dimension-independent* rates in linear models. As application, we provide nearly minimax-optimal dimension-free regret bounds in private linear contextual bandits.

Existing results. In Table 1, we summarize the known regret bounds for learning in (generalized) linear contextual bandits under the joint or local model of privacy. For simplicity of presentation, all results are specialized to linear contextual bandits with stochastic contexts, and we omit poly-logarithmic factors.

Organization. In Section 3, we discuss the main difficulty of achieving optimal rates while preserving privacy in linear contextual bandits, motivating the approach of L_1 -regression. We then present our reweighted regression method (Section 4) for private generalized linear models. In Section 5, we apply our method to provide private regret guarantees in generalized linear contextual bandits. Further, in Section 6, we investigate *dimension-free* private learning in linear models and linear contextual bandits. For succinctness, we mostly focus on the JDP setting in the main body of the paper and present the algorithms for LDP setting in the appendices.

2 Preliminaries

Differential privacy (DP) is a widely adopted framework for ensuring privacy in data analysis. In this section, we will introduce the definition of joint DP and local DP algorithms for both offline and interactive settings. We first review the notion of differentially private (DP) channels.

Definition 1 (DP channel) For the latent observation space Z and the noisy observation space O, a channel Q is a measurable map $Z \to \Delta(O)$. A channel Q is (α, β) -DP if for $z, z' \in Z$, any measurable set $E \subseteq O$,

$$\mathsf{Q}(E|z) \le e^{\alpha} \mathsf{Q}(E|z') + \beta.$$

In this paper, we focus on the regime $\alpha, \beta \in (0, 1)$.

The Gaussian channel is a standard example of the (α, β) -DP channels (see e.g., [Balle and Wang, 2018]).

Definition 2 (Gaussian channel) Suppose that $\alpha, \beta \in (0,1)$, and denote $\sigma_{\alpha,\beta} = \frac{2\sqrt{\log(1.25/\beta)}}{\alpha}$. For any given function $F : \mathcal{Z} \to \mathbb{R}^d$, we let $\Delta(F) := \sup_{z,z'} \|F(z) - F(z')\|_2$. Then, for any $\Delta \ge \Delta(F)$, the channel $\mathbb{Q}(\cdot|z) = \mathbb{N}(F(z), \sigma^2_{\alpha,\beta}\Delta^2)$ is a (α, β) -DP channels.

In the following, we denote by $\operatorname{Priv}_{\Delta}(v) = \mathsf{N}(v, 4\sigma_{\alpha,\beta}^2 \Delta^2)$ the Gaussian channel with sensitivity Δ . It is guaranteed that for any function F with $||F(z)|| \leq \Delta$, the channel $z \mapsto \operatorname{Priv}_{\Delta}(F(z))$ is (α, β) -DP. Further, for any symmetric matrix $V \in \mathbb{R}^{d \times d}$, we denote $\operatorname{SymPriv}_{\Delta}(V)$ to be the distribution of V + Z, where Z is a symmetric Gaussian random matrix, i.e., $Z_{ij} = Z_{ji} \sim \mathsf{N}(0, 4\sigma_{\alpha,\beta}^2 \Delta^2)$ independently.

Joint DP. We first recall the definition of (joint) DP algorithms for *non-interactive* problems. In this setting, an *algorithm* maps the dataset $\mathcal{D} = \{z_1, \dots, z_T\} \in \mathcal{Z}^T$ to a distribution over the output decision Π . The two dataset $\mathcal{D} = (z_1, \dots, z_T), \mathcal{D}' = (z'_1, \dots, z'_T) \in \mathcal{Z}^T$ are neighbored if there is at most one index i such that $z_i \neq z'_i$.

Definition 3 (JDP for non-interactive algorithms) An algorithm Alg preserves (α, β) -JDP if for any neighbored dataset $\mathcal{D}, \mathcal{D}'$ and any measurable set $E \subseteq \Delta(\Pi)$,

$$\mathsf{Alg}(E|\mathcal{D}) \le e^{\alpha} \mathsf{Alg}(E|\mathcal{D}') + \beta.$$

On the other hand, a *T*-round (interactive) algorithm Alg for contextual bandit problem is specified by a sequence of mapping $\{\pi_t(\cdot \mid \cdot)\}$, where the *t*-th mapping $\pi_t(\cdot \mid \mathcal{H}_{t-1}, x_t)$ specifies the distribution of $a_t \in \mathcal{A}$ based on the history $\mathcal{H}_{t-1} = \{z_s = (x_s, a_s, r_s)\}_{s \leq t-1}$ and the current context x_t . As the algorithm has to ensure the whole sequence (a_1, \dots, a_T) protects privacy, it's a more challenging task compared to the non-interactive setting. In this work, we consider the following notion of joint DP with interaction.

Definition 4 (Anticipating JDP) An algorithm Alg is said to preserve (α, β) -JDP (or simply be (α, β) -JDP) if for every round $t \in [T]$, any two neighbored history $\mathcal{H}_t, \mathcal{H}'_t$ differing only at round t, and any sequence of possible future observations $\mathcal{D}_{t+1:T} = \{(x_{t'}, r_{t'})\}_{t' \in [t+1,T]}$, we have

$$\mathbb{P}^{\operatorname{Alg}}((a_{t+1},\cdots,a_T)\in E|\mathcal{H}_t,\mathcal{D}_{t+1:T})\leq e^{\alpha}\mathbb{P}^{\operatorname{Alg}}((a_{t+1},\cdots,a_T)\in E|\mathcal{H}'_t,\mathcal{D}_{t+1:T})+\beta,\qquad\forall E\subseteq\mathcal{A}^{T-t},$$

where the probability \mathbb{P}^{Alg} only accounts the randomness of Alg, i.e., for $\mathcal{H}_t = \{(x_t, a_t, r_t)\}_{t \in [T]}$,

$$\mathbb{P}^{\text{Alg}}(a_{t+1}, \cdots, a_T | \mathcal{H}_t, \mathcal{D}_{t+1:T}) = \prod_{t'=t+1}^T \pi_{t'}(a_{t'} | (x_s, a_s, r_s)_{s < t'}, x_{t'}).$$

Definition 4 is the most widely-adopted definition of privacy-preserving procedures in the literature of contextual bandits [Shariff and Sheffet, 2018, Azize and Basu, 2024] and episodic RL [Vietri et al., 2020], and it can be interpreted as following. Assume that a malicious adversary is trying to identify the private information (x_t, a_t, r_t) of the unit which is treated at round $t \in [T]$, and it can adversarially design the context x_s and reward r_s after round t. An algorithm is private if the adversary cannot infer the private information (x_t, a_t, r_t) from the output actions $a_{>t}$ of the algorithm no matter what the history \mathcal{H}_{t-1} is and how the agent designs the input $\{x_{>t}, r_{>t}\}$. Since the algorithm is *non-anticipating*, the history output $a_{<t}$ will also not be impacted by (x_t, a_t, r_t) . In the meanwhile, the output a_t will unavoidably contain information about x_t , otherwise no non-trivial regret guarantee could be attained, as proved by Shariff and Sheffet [2018].

Local DP. Parallel to the above model of DP, a line of work [Kasiviswanathan et al., 2011, Duchi et al., 2013, 2016] studies the *local differential privacy* (LDP) model that provides stronger protection of privacy. It is adopted in scenarios where each individual wants to protect their personal privacy and does not fully trust the data collector, so users locally perturb or add noise to their data before sending it, ensuring that even the collecting party cannot learn the exact original information, but only privatized observation $o \in \mathcal{O}$.

For a *T*-round algorithm Alg operating on observation space \mathcal{Z} and decision space Π , Alg is said to preserve (α, β) -LDP if it adopts the following protocol for each round t = 1, ..., T:

- Alg selects a decision $\pi_t \in \Pi$ and a (α, β) -DP channel Q_t based on the history $\mathcal{H}_{t-1} = \{\pi_1, o_1, \cdots, \pi_{t-1}, o_{t-1}\}$.
- The environment generates an observation $z_t \in \mathcal{Z}$ sampled via $z_t \sim M^*(\pi_t)$, where M^* is the underlying *model* of the environment.
- Alg receives a noisy observation $o_t \sim Q_t(\cdot|z_t)$.

In other words, an algorithm that preserves local DP never has direct access to the observation $z_t \in \mathbb{Z}$, and only the "privatized" observation $o_t \sim Q_t(\cdot|z)$ is revealed. Therefore, an LDP algorithm has to adaptively select both the decision π_t and also the private channel Q_t to obtain information from the environment.

Generalized linear models. In this paper, we also study the generalized linear models, an important sub-problem of the generalized linear contextual bandits.

Definition 5 In generalized linear models (GLM), a covariate space $C \subseteq \mathbb{B}^d(1)$ and a link function $\nu : [-1,1] \rightarrow [-1,1]$ is given, and the latent observation space is $\mathcal{Z} = \mathcal{C} \times [-1,1]$. The ground truth model $M^* \in \Delta(\mathcal{Z})$ is specified as

$$(\phi, y) \sim M^{\star}: \quad \phi \sim p, \ \mathbb{E}[y|\phi] = \nu(\langle \phi, \theta^{\star} \rangle),$$

where p is a covariate distribution over \mathcal{C} , and $\theta^* \in \mathbb{B}^d(1)$ is the ground truth parameter.

Canonical examples of GLM include the *linear models*, where $\nu(t) = t$, and the *logistic models*, where $\nu(t) = \frac{1}{1+e^{-t}}$. In this paper, we assume the link function ν is well-conditioned.

Assumption 1 There exists constant $0 < \mu_{\nu} \leq L_{\nu}$ such that $\mu_{\nu} \leq \nu'(t) \leq L_{\nu}$ for all $t \in [-1,1]$, and we denote $\kappa_{\nu} := \frac{L_{\nu}}{\mu_{\nu}}$ to be the condition number of the link function ν .

3 Motivation

We start by reviewing why most existing private algorithms for linear contextual bandits fail to achieve an optimal regret rate without the strong explorability condition $\lambda_{\min}^{\star} > 0$. Such insufficiency motivates the confidence intervals-based approach of Li et al. [2024] discussed in Section 3.2.

3.1 Insufficiency of standard regression

In general, the existing algorithmic principles for learning contextual bandits mostly rely (either explicitly or implicitly) on the regression subroutines that, given a sequence of observation $\{(x_t, a_t, r_t)\}_{t \in [N]}$, produce an reward estimation \hat{f} with bounded mean-square error:

$$\mathbb{E}_{(x,a)\sim\mathcal{D}}(\hat{f}(x,a) - f^{\star}(x,a))^2 \le \mathcal{E}(N)^2.$$

For non-private linear contextual bandits, it is well-known that regression-based estimators achieve the optimal rate of $\mathcal{E}(N)^2 \simeq \frac{d}{N}$, and such a convergence guarantee of N^{-1} -rate is essential in the regret analysis of the classical LinUCB algorithm and its variants [Abbasi-Yadkori et al., 2011, Li et al., 2019, Bastani and Bayati, 2020]. Further, for contextual bandits with a general reward function class, the recent regression-oracle based algorithms [Simchi-Levi and Xu, 2020, Foster and Rakhlin, 2020] achieve regret bounds scaling with $\tilde{O}(T \cdot \mathcal{E}(T))$, and hence a T^{-1} -rate of convergence under L_2 -error metric is also crucial to obtain a regret of order $\tilde{O}(\sqrt{T})$.

Therefore, for regression-based algorithms, achieving rate-optimal regret essentially relies on the L_2 -error guarantee of the regression subroutine. However, it is known that in linear models, privacy leads to slower convergence under the L_2 -error if the covariate distribution is ill-conditioned, as the following folklore lemma indicates.

Proposition 3.1 (Lower bounds for ill-conditioned linear regression) Suppose that $T \ge 1$, $\alpha \in (0,1]$, $\lambda \in [0,1]$, d = 1, and the link function $\nu(t) = t$ is identity. Let the covariate distribution $p \in \Delta([-1,1])$ be known and given by $p(1) = \lambda$, $p(0) = 1 - \lambda$. Then

(1) For any T-round (α, β) -JDP algorithm Alg with output $\hat{\theta}$, it holds that

$$\sup_{\theta^{\star} \in [-1,1]} \mathbb{E}^{\theta^{\star},\operatorname{Aig}} \Big[\mathbb{E}_{\phi \sim p} \langle \phi, \hat{\theta} - \theta^{\star} \rangle^2 \Big] \gtrsim \min \left\{ \frac{1}{\lambda(\alpha + \beta)^2 T^2}, \lambda \right\}.$$

(2) For any T-round (α, β) -LDP algorithm Alg with output $\hat{\theta}$, as long as $\beta \leq \frac{\lambda}{T^2}$, it holds that

$$\sup_{\theta^{\star} \in [-1,1]} \mathbb{E}^{\theta^{\star},\operatorname{Aig}} \Big[\mathbb{E}_{\phi \sim p} \langle \phi, \hat{\theta} - \theta^{\star} \rangle^2 \Big] \gtrsim \min \left\{ \frac{1}{\lambda \alpha^2 T}, \lambda \right\}$$

Note that in the above construction, $\mathbb{E}_{\phi\sim p}[\phi\phi^{\top}] = \lambda$. Hence, in linear contextual bandits, the oracle-based regret bounds described above will scale with $\tilde{O}\left(\sqrt{T} + \frac{1}{\lambda_{\min}^{\star}\alpha}\right)$ under joint DP model, and $\tilde{O}\left(\sqrt{T}/(\lambda_{\min}^{\star}\alpha)\right)$ under the local DP model, where λ_{\min}^{\star} is the minimum eigenvalue over any policy that the algorithm may play. Further, if there is not a lower bound on $\lambda > 0$, then Proposition 3.1 provides the worst-case lower bounds of $\Omega\left(\frac{1}{(\alpha+\beta)T}\right)$ and $\Omega\left(\frac{1}{\alpha\sqrt{T}}\right)$ for L_2 -regression under the JDP model and LDP model, respectively, implying significant degradation under privacy.

To sum it up, without a new analysis framework, the existing algorithms that only rely on the regression oracles with L_2 -error guarantee might not avoid the strong explorability condition $\lambda_{\min}^{\star} > 0$. Therefore, to achieve the optimal regret rates for (generalized) linear contextual bandits, we cannot use the standard linear regression primitives to estimate ground-truth parameter θ^{\star} .

3.2 Alternative approach: Regression with confidence intervals

As an alternative to the standard regression based approach [Foster et al., 2018, Foster and Rakhlin, 2020], Li et al. [2024] propose an action elimination framework based on regression with L_1 -error guarantee and the additional confidence interval structures. The key observation is that, while the negative results (Proposition 3.1) do rule out the regression oracles with O(1/T) convergence rate under L_2 -error, such oracles are not necessary for designing algorithm. Specifically, Li et al. [2024] consider the regression subroutine with confidence intervals (L_1 -regression for short), which is defined as following:

Definition 6 (L_1 -regression oracle) Let $N \ge 1$, $\delta \in (0, 1)$. In contextual bandits, a L_1 -regression oracle is a N-round algorithm Alg that outputs an estimate of reward function $\hat{f} : \mathcal{X} \times \mathcal{A} \to [-1, 1]$ and an confidence bound $\hat{b} : \mathcal{X} \times \mathcal{A} \to \mathbb{R}_{\ge 0}$, such that for any fixed policy $\pi : \mathcal{X} \to \Delta(\mathcal{A})$, given data (x_t, a_t, r_t) generated independently as

$$x_t \sim P, \quad a_t \sim \pi(\cdot | x_t), \quad \mathbb{E}[r_t | x_t, a_t] = f^*(x_t, a_t), \quad t \in [N]$$

the following holds with probability at least $1 - \delta$:

- (1) (Valid confidence interval) $\left| \hat{f}(x,a) f^{\star}(x,a) \right| \leq \hat{b}(x,a)$ for all $(x,a) \in \mathcal{X} \times \mathcal{A}$.
- (2) (L₁-performance bound) $\mathbb{E}_{x \sim P, a \sim \pi(x)} \Big[\hat{b}(x, a) \Big] \leq \mathcal{E}_{\delta}(N).$

The above conditions on L_1 -regression oracle only imply that the L_1 -error is bounded as $\mathbb{E}|\hat{f}(x,a) - f^*(x,a)| \leq \mathcal{E}(N)$. This is arguably weaker than the mean-square (L_2) convergence $\mathbb{E}(\hat{f}(x,a) - f^*(x,a))^2 \leq \mathcal{E}(N)^2$ by the common L_2 -regression.

With a private L_1 -regression oracle, Li et al. [2024] adopt an algorithm based on action elimination that achieves a regret of $\tilde{O}(T \cdot \mathcal{E}_{1/T}(T))$ (for details, see also Appendix D.2). Therefore, the framework opens the door for a \sqrt{T} -regret by developing L_1 -regression oracle with $\mathcal{E}_{\delta}(T) = \tilde{O}(1/\sqrt{T})$. However, the L_1 -regression oracle of Li et al. [2024] is based on iterative private PCA and layered private linear regression, achieving the rate $\mathcal{E}_{\delta}(T) \leq \tilde{O}\left(\frac{\log^d(T)}{\alpha\sqrt{T}}\right)$ and hence leading to a $\log^d(T)\sqrt{T}$ -regret that is exponential of the dimension d. This regret bound is meaningful only when the dimension d = O(1) is of constant order.

On the other hand, Chen and Rakhlin [2025] provides a significantly improved regret of $\sqrt{d^3T}/\alpha$. While the Decision-Estimation Coefficient (DEC) approach is much different from the aforementioned ones, the way they upper bound the DEC implicitly utilizes the confidence interval bounds with L_1 guarantee. Indeed, for linear regression, Chen and Rakhlin [2025] also provide a near-optimal $T^{-1/2}$ -rate under L_1 -error. For achieving such guarantees, they propose a novel normalization method based on a re-weighting matrix U (detailed discussion in Section 4). However, this method is introduced purely for upper bounding the DEC, and it is unknown whether it provides a more efficient algorithm.

Inspired by the insights from Li et al. [2024] and Chen and Rakhlin [2025], in Section 4, we develop an efficient and near-optimal L_1 -regression procedure that applies to both joint DP and local DP settings. Then, in Section 5, we adopt the proposed L_1 -regression oracle in the action elimination framework of Li et al. [2024] to provide rate-optimal private regret bounds.

4 Private Reweighted Regression

In this section, we build upon the techniques of Chen and Rakhlin [2025] to provide an optimal estimation guarantee under L_1 -error. For linear regression, Chen and Rakhlin [2025] provides a near-optimal convergence

rate under L_1 -error, based on the DEC framework and the Exploration-by-Optimization algorithm. For each round, the Exploration-by-Optimization algorithm in some sense estimates a "worst-case" context distribution p, and Chen and Rakhlin [2025] utilize such a context distribution to compute a normalization matrix U:

$$\mathbb{E}_{\phi \sim p} \left[\frac{U \phi \phi^{\top} U}{\|U \phi\|} \right] + \lambda U = \mathbf{I}, \tag{1}$$

where $\lambda > 0$ is a regularization parameter. Chen and Rakhlin [2025] then bound the DEC based on the normalization (U, λ) , hence providing the desired convergence rate through the DEC framework.

However, it is unclear whether the normalization matrix U can be useful algorithmically. As the main motivation of our approach, we first discuss in Section 4.1 how such a normalization matrix U, satisfying the following relaxed equation,

$$\frac{1}{2}\mathbf{I} \preceq \mathbb{E}_{\phi \sim p} \left[\frac{U\phi\phi^{\top}U}{\|U\phi\|} \right] + \lambda U \preceq 2\mathbf{I},\tag{2}$$

can be helpful in L_1 -regression. We then provide a private procedure (Section 4.2) that learns a normalization (U, λ) satisfying Eq. (2), and develop our private reweighted regression method with near-optimal convergence guarantee in Section 4.3.

4.1 Key idea: Reweighting based on normalization matrix

For generalized linear models with link function ν , we can consider the following loss objective,

$$\mathcal{L}_{\mathsf{GLM}}(\theta) := \mathbb{E}_{(\phi, y) \sim M^{\star}}[\ell_{\nu}(\langle \phi, \theta \rangle, y)],$$

where the *integral loss* ℓ_{ν} associated with ν is defined as $\ell_{\nu}(t, y) := -yt + \int_{0}^{t} \nu(s) ds$. The basic property of $\mathcal{L}_{\mathsf{GLM}}$ is that $\nabla \mathcal{L}_{\mathsf{GLM}}(\theta) = \mathbb{E}_{(\phi, y) \sim M^{\star}}[(\nu(\langle \phi, \theta \rangle) - y) \cdot \phi]$, and hence $\nabla \mathcal{L}_{\mathsf{GLM}}(\theta^{\star}) = 0$, i.e., θ^{\star} is a global minimizer of $\mathcal{L}_{\mathsf{GLM}}$.

Reweighted objective. Given a normalization matrix U and a parameter together satisfying Eq. (2), we may reweigh and regularize the objective function $\mathcal{L}_{\mathsf{GLM}}$ according to

$$\mathcal{L}_{\mathsf{weighted}}(\theta) \mathrel{\mathop:}= \mathbb{E}_{(\phi,y) \sim M^{\star}} \bigg[\frac{\ell_{\nu}(\langle \phi, \theta \rangle, y)}{\|U\phi\|} \bigg] + \frac{L_{\nu}\lambda}{2} \left\|\theta\right\|_{U^{-1}}^2.$$

By Assumption 1 and Eq. (2), we know that $\frac{1}{2}\mu_{\nu} \cdot U^{-2} \preceq \nabla^2 \mathcal{L}_{weighted}(\theta) \preceq 2L_{\nu} \cdot U^{-2}$ for any $\|\theta\| \leq 1$. Therefore, the objective $\mathcal{L}_{weighted}$ is well-conditioned after suitable linear transformation $\theta = Uw$. Specifically, we define

$$\mathcal{L}_{U}(w) := \mathbb{E}_{(\phi,y)\sim M^{\star}}\left[\frac{\ell_{\nu}(\langle U\phi, w\rangle, y)}{\|U\phi\|}\right] + \frac{L_{\nu}\lambda}{2} \|w\|_{U}^{2},$$
(3)

and then \mathcal{L}_U is $(2L_\nu)$ -smooth and $(\mu_\nu/2)$ -strongly convex over the domain $\mathcal{W}_U := \{w \in \mathbb{R}^d : ||Uw|| \le 1\}$. Further, the gradient of \mathcal{L}_U can be derived as

$$\nabla \mathcal{L}_U(w) = \mathbb{E}_{(\phi,y)\sim M^\star} \left[\frac{U\phi}{\|U\phi\|} (\nu(\langle U\phi, w\rangle) - y) \right] + \lambda L_\nu \cdot Uw.$$

The following lemma indicates that, any approximate minimizer of \mathcal{L}_U is provides a good approximation of the ground truth parameter (under the L_1 -error).

Lemma 4.1 Suppose that (U, λ) satisfies Eq. (2). We let $\widehat{w}_U^* := \arg \min_{w: \|Uw\| \le 1} \mathcal{L}_U(w)$ and $w^* := U^{-1}\theta^*$. Then it holds that $\|\widehat{w}_U^* - w^*\| \le 4\lambda$. Further, the following holds:

(1) Estimation error: for any $w \in W$, we have

$$\mathbb{E}_{\phi \sim p} |\nu(\langle \phi, Uw \rangle) - \nu(\langle \phi, \theta^* \rangle)| \le L_{\nu} \mathbb{E}_{\phi \sim p} |\langle \phi, Uw - \theta^* \rangle| \le 2\sqrt{d}L_{\nu} \|w - w^*\|$$
(4)

(2) Confidence interval: $\forall \phi \in \mathbb{R}^d$, $|\langle \phi, Uw \rangle - \langle \phi, \theta^* \rangle| \le ||U\phi|| \cdot ||w - w^*||$, and $\mathbb{E}_{\phi \sim p}[||U\phi||] \le 2d$.

Algorithm 1 Subroutine JDP_Learning_Normalization

Input: Dataset $\mathcal{D} = \{(\phi_t, y_t)\}_{t \in [T]}$, parameter $\delta \in (0, 1)$. **Input:** Epoch $K \ge 1$, batch size $N = \lfloor \frac{T}{K} \rfloor$, parameters $\lambda_{(0)}, \dots, \lambda_{(K)}$. Initialize $U_{(0)} = \mathbf{I}$. **for** $k = 0, \dots, K - 1$ **do** Compute the estimate on the split dataset $\mathcal{D}_{(k)} = \{(\phi_t, y_t)\}_{t \in [kN+1, (k+1)N]}$:

 $1 \sum_{i=1}^{(k+1)N} U_{i,i}^{1/2} \phi_t \phi_t^{\top} U_{i,i}^{1/2}$

$$H_{(k)} = \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \frac{U_{(k)}^{1/2} \phi_t \phi_t^\top U_{(k)}^{1/2}}{\|U_{(k)} \phi_t\|}.$$
(6)

Privatize $\widetilde{H}_{(k)} \sim \mathsf{SymPriv}_{1/N}(H_{(k)})$. Update

$$\mathbf{\Sigma}_{(k)} = U_{(k)}^{1/2} \widetilde{H}_{(k)} U_{(k)}^{1/2} + \lambda_{(k)} U_{(k)}, \qquad U_{(k+1)} = \operatorname{sym}(\mathbf{\Sigma}_{(k)}^{-1/2} U_{(k)}).$$

Output: Normalization matrix $U = U_{(K)}$ and normalization parameter $\lambda = \lambda_{(K)}$.

Therefore, with N samples, the privatized gradient descent produces a solution \hat{w} with $\|\hat{w} - \hat{w}_U^{\star}\| \leq \tilde{O}(N^{-1/2})$ hence providing the desired convergence rate under L_1 -error through Eq. (4). Note that this bypasses the lower bounds of Proposition 3.1, because the objective \mathcal{L}_U is different from the standard squared loss objective.

4.2 Learning normalization matrix privately

We start by describing how we can learn the normalization matrix U satisfying Eq. (2) from data privately. Indeed, even when the covariate distribution p is known, it is not clear how to compute a normalization matrix U, and Chen and Rakhlin [2025] have to invoke a fixed-point argument (Brouwer's theorem) to prove its existence. Our key observation is that the following spectral iterates converge to solution to Eq. (1):

$$\boldsymbol{\Sigma}_{(k)} = \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k)} \phi \phi^{\top} U_{(k)}}{\|U_{(k)} \phi\|} \right] + \lambda U_{(k)}, \qquad U_{(k+1)} = \operatorname{sym}(\boldsymbol{\Sigma}_{(k)}^{-1/2} U_{(k)}), \tag{5}$$

with the initial point $U_{(0)} = \mathbf{I}$. Specifically, it holds that

$$\lambda_{\min}(\mathbf{\Sigma}_{(k)})^{1/2} \leq \lambda_{\min}(\mathbf{\Sigma}_{(k+1)}) \leq \lambda_{\max}(\mathbf{\Sigma}_{(k+1)}) \leq \lambda_{\max}(\mathbf{\Sigma}_{(k)})^{1/2}.$$

Therefore, if the iteration (5) is exact, the matrix $\Sigma_{(k)}$ converges to the identity matrix at a quadratic rate, implying that $O(\log \log(1/\lambda))$ iterations are enough to achieve sufficient accuracy.

Approximate spectral iteration. In general, the covariate distribution p is also not known, and we have to approximately implement the update rule (5). To this end, we propose Algorithm 1, which privately approximates Eq. (5) by Eq. (6) with batched samples.

Algorithm 1 preserves (α, β) -JDP by the composition property of joint DP mechanisms, as proved in Appendix C.2.

Lemma 4.2 Subroutine JDP_Learning_Normalization (Algorithm 1) preserves (α, β) -JDP.

We demonstrate that the iterates of Algorithm 1 converge to a solution of Eq. (2), as follows. The details are deferred to Appendix C.3.

Proposition 4.3 Let $T \ge 1, K \ge 1, \delta \in (0,1)$, and $\varepsilon_N := C_0 \left(\sqrt{\frac{\log(K/\delta)}{N}} + \frac{\sigma_{\alpha,\beta}\sqrt{d + \log(K/\delta)}}{N} \right)$, where C_0 is an absolute constant chosen according to Lemma C.2. Suppose that Algorithm 1 is instantiated with

Algorithm 2 JDP_L1_Regression

Input: Dataset $\mathcal{D} = \{(\phi_t, y_t)\}_{t \in [T]}$, parameter $\delta \in (0, 1)$. 1: Split the dataset $\mathcal{D} = \mathcal{D}_0 \cup \mathcal{D}_1$ equally.

- 2: Set $(U, \lambda) \leftarrow \mathsf{JDP_Learning_Normalization}(\mathcal{D}_0, \delta)$.
- 3: Set $\widehat{w} \leftarrow \mathsf{JDP}_\mathsf{Batch}_\mathsf{SGD}(\mathcal{L}_U, \mathcal{D}_1)$.

Output: Normalization (U, λ) , estimator $\hat{\theta} = U\hat{w}$, and estimation error $\overline{\epsilon}$.

parameters $\lambda_{(k)} = (2k+5)\varepsilon_N$, and then with probability at least $1-\delta$,

$$\exp\left(-\frac{\log(1/\lambda_0)}{2^{k-1}}\right)\mathbf{I} \preceq \mathbb{E}_{\phi \sim p}\left[\frac{U_{(k)}\phi\phi^{\top}U_{(k)}}{\|U_{(k)}\phi\|}\right] + \lambda_{(k)}U_{(k)} \preceq \exp\left(\frac{12}{k}\right)\mathbf{I}.$$

In particular, as long as $K \ge \max \{ \log \log(N), 20 \}$, Algorithm 1 outputs (U, λ) that satisfies Eq. (2) with probability at least $1 - \delta$, where $\lambda := \lambda_{(K)} = (2K + 5)\varepsilon_N$.

4.3Private regression with reweighting

In the following, we present Algorithm 2 for private L_1 -regression, which is based on (1) first learning the normalization (U, λ) by the subroutine JDP_Learning_Normalization (Algorithm 1), and then (2) running the private batched SGD subroutine (Algorithm 7) on the reweighted objective \mathcal{L}_U defined in Eq. (3). The batched SGD subroutine is standard and hence deferred to Appendix C.4.

Theorem 4.4 (Generalized linear regression with JDP) Let $T \geq 1, \delta \in (0, 1)$, and the subroutines of Algorithm 2 are suitably instantiated according to Appendix C.4. Then, under generalized linear model, Algorithm 2 preserves (α, β) -JDP, and the following holds:

(1) With probability at least $1-\delta$, the normalization (U,λ) satisfies Eq. (2), and the returned estimator \hat{w} satisfies $L_{\nu} \| \widehat{w} - \widehat{w}_U^{\star} \| \leq \underline{\epsilon}$, where

$$\lambda := \lambda(T, \delta) = \tilde{O}\left(\sqrt{\frac{\log(1/\delta)}{T}} + \sigma_{\alpha,\beta} \frac{\sqrt{d\log(1/\delta)}}{T}\right),$$

$$\underline{\boldsymbol{\epsilon}} := \underline{\boldsymbol{\epsilon}}(T, \delta) = \tilde{O}\left(\kappa_{\nu}^{3/2} \sqrt{\frac{\log(1/\delta)}{T}} + \sigma_{\alpha,\beta}(\kappa_{\nu}^{3/2} + \kappa_{\nu}\sqrt{d}) \frac{\sqrt{\log(1/\delta)}}{T}\right),$$

are defined in Proposition 4.3 and Proposition C.6, respectively, and $\hat{O}(\cdot)$ hides polynomial factors of $\log(T)$. The overall estimation error $\overline{\boldsymbol{\epsilon}}$ is defined as $\overline{\boldsymbol{\epsilon}}(T, \delta) := 4L_{\nu}\lambda(T, \delta) + \boldsymbol{\epsilon}(T, \delta)$.

(2) In particular, under the success event of (1), we have (by Lemma 4.1)

$$\mathbb{E}_{\phi \sim p} |\nu(\langle \phi, \hat{\theta} \rangle) - \nu(\langle \phi, \theta^{\star} \rangle)| \le L_{\nu} \cdot \mathbb{E}_{\phi \sim p} |\langle \phi, \hat{\theta} - \theta^{\star} \rangle| \le 2\sqrt{d} \cdot \overline{\epsilon},$$

and for all $\phi \in \mathbb{R}^d$, the confidence bound holds: $|\nu(\langle \phi, \hat{\theta} \rangle) - \nu(\langle \phi, \theta^* \rangle)| \leq ||U\phi|| \overline{\epsilon}$.

Therefore, Algorithm 2 is an L_1 -regression oracle in the sense of Definition 6. In particular, for linear models, we have $\kappa_{\nu} = 1$, and hence the convergence rate can further be simplified to $\tilde{O}\left(\sqrt{\frac{d}{T}} + \frac{d}{\alpha T}\right)$. The first term matches the minimax-optimal convergence rate of non-private L_1 -regression (given by the least squares), and the second term ("price-of-privacy") is of lower order compared to the first term.

Extension: Local DP. As an LDP extension of Algorithm 2, we propose Algorithm 8 that preserves local DP while achieving near-optimal L_1 -regression guarantee.

Theorem 4.5 (Generalized linear regression with LDP) Let $T \ge 1, \delta \in (0, 1)$. Algorithm 8 (described in Appendix C.5) preserves (α, β) -LDP, and it returns estimator $\hat{\theta}$ such that with probability at

Algorithm 3 Action Elimination Algorithm with JDP

Input: Round $T \ge 1$, parameter $\delta \in (0, 1)$, epoch schedule $1 = T^{(0)} < T^{(1)} < \cdots < T^{(J)} = T$. Initialize $\hat{f}^{(0)} \equiv 0, \hat{b}^{(0)} \equiv 1$. **for** $j = 0, 1, \cdots, J - 1$ **do** Set $\pi^{(j)} \leftarrow \text{Confidence_based_Planning}(\{(\hat{f}^{(\tau)}, \hat{b}^{(\tau)})\}_{\tau < j})$ and initialize the dataset $\mathcal{D}^{(j)} = \{\}$. **for** $t = T^{(j)}, \cdots, T^{(j+1)} - 1$ **do** Observe context $x_t \sim P$, select action $a_t \sim \pi^{(j)}(x_t)$, and receive reward r_t . Update the dataset $\mathcal{D}^{(j)} \leftarrow \mathcal{D}^{(j)} \cup \{(\phi(x_t, a_t), r_t)\}$. Set $(U^{(j)}, \lambda^{(j)}), \hat{\theta}^{(j)}, \overline{\boldsymbol{\epsilon}}^{(j)} \leftarrow \text{JDP_L1_Regression}(\mathcal{D}^{(j)}, \delta)$. Update the estimated reward function and confidence bound as

$$\hat{f}^{(j)}(x,a) = \nu(\langle \phi(x,a), \hat{\theta}^{(j)} \rangle), \qquad \hat{b}^{(j)}(x,a) = \overline{\epsilon}^{(j)} \cdot \left\| U^{(j)}\phi(x,a) \right\|, \qquad \forall (x,a) \in \mathcal{X} \times \mathcal{A}.$$

least $1 - \delta$,

$$\mathbb{E}_{\phi \sim p} |\nu(\langle \phi, \hat{\theta} \rangle) - \nu(\langle \phi, \theta^{\star} \rangle)| \leq \tilde{O}\left(\sigma_{\alpha, \beta}(\kappa_{\nu}d + \sqrt{\kappa_{\nu}^{3}d})\sqrt{\frac{\log(1/\delta)}{T}}\right),$$

where $O(\cdot)$ hides polynomial factors of $\log(T)$. Further, the total number of switches (or, changes) of the deployed private channels is bounded by $O(\kappa_{\nu} \log T)$.

For linear models, the upper bound of Theorem 4.5 simplifies to $\tilde{O}\left(\frac{d}{\alpha\sqrt{T}}\right)$. Such a nearly minimax-optimal convergence rate is first derived by Chen and Rakhlin [2025], using an algorithm that solves an *exponential-time* optimization problem for each round.

5 Private Learning in Generalized Linear Contextual Bandits

In this section, we use our L_1 -regression method from Section 4 as a subroutine for the action elimination framework [Li et al., 2024], providing rate-optimal private regret bounds for generalized linear contextual bandits.

Our algorithm (Algorithm 3) is epoch-based (with a given epoch schedule $1 = T_0 < T_1 < T_2 < \cdots < T_J = T$), and it iteratively builds estimations of the ground truth reward function f^* and plans according to the estimations. The algorithm, which consists of an estimation procedure and a planning procedure described as follows, is similar to that of Li et al. [2024].

Estimation procedure. For *j*th epoch, the estimate reward function $\hat{f}^{(j)}$ and the confidence radius $\hat{b}^{(j)}$ are produced by the subroutine JDP_L1_Regression. Then, by Theorem 4.4, it holds that with high probability

$$f^{\star}(x,a) \in \left[\hat{f}^{(j)}(x,a) - \hat{b}^{(j)}(x,a), \hat{f}^{(j)}(x,a) + \hat{b}^{(j)}(x,a) \right], \qquad \forall x \in \mathcal{X}, a \in \mathcal{A}.$$
(7)

Planning procedure. The policy $\pi^{(j)}$ of the *j*th epoch is built upon the confidence intervals (7) given by the estimations $(\hat{f}^{(0)}, \hat{b}^{(0)}), \dots, (\hat{f}^{(j-1)}, \hat{b}^{(j-1)})$ from previous epochs. Given the estimations, subroutine **Confidence_based_Planning** (Algorithm 4) eliminates the sub-optimal arms for each context $x \in \mathcal{X}$ according to Eq. (8), and output the $\pi^{(j)}(x)$ based on a *spanner* of the remaining actions.² This procedure implicitly encourages exploration, as it uses optimistic estimation (UCB) of the value of each arm.

²Defined in Definition 7. Particularly, when $|\mathcal{A}| = O(1)$, we can directly takes $\mathcal{A}_{sp}^{(j)}(x) = \mathcal{A}^{(j-1)}(x)$ and set $\pi^{(j)}(x) = \text{Unif}(\mathcal{A}^{(j-1)}(x))$.

Algorithm 4 Subroutine Confidence_based_Planning

Input: Confidence interval descriptions $(\hat{f}^{(0)}, \hat{b}^{(0)}), \cdots, (\hat{f}^{(j-1)}, \hat{b}^{(j-1)})$ up to the *j*th epoch. for context $x \in \mathcal{X}$ do Set $\mathcal{A}^{(0)}(x) = \mathcal{A}$. for $\tau = 1, \cdots, j - 1$ do Set $\mathcal{A}^{(\tau)}(x) = \left\{ a \in \mathcal{A}^{(\tau-1)}(x) : \hat{f}^{(\tau)}(x, a) + \hat{b}^{(\tau)}(x, a) \ge \max_{a' \in \mathcal{A}} \hat{f}^{(\tau)}(x, a') - \hat{b}^{(\tau)}(x, a') \right\}.$ (8) Compute a spanner $\mathcal{A}_{sp}^{(j)}(x)$ of $\mathcal{A}^{(j-1)}(x)$ (Definition 7). Set $\pi^{(j)}(x) = \text{Unif}(\mathcal{A}_{sp}^{(j)}(x))$.

Output: Policy $\pi^{(j)}$.

Regret guarantee. We state the upper bound of Algorithm 3 in terms of the dimensionality of the per-context action space:

$$d_{\mathcal{A}} := \max_{x \in \mathcal{X}} \dim \left(\{ \phi(x, a) : a \in \mathcal{A} \} \right).$$

Note that $d_{\mathcal{A}} \leq \min\{d, |\mathcal{A}|\}.$

Theorem 5.1 (Regret upper bound under JDP) Algorithm 3 preserves (α, β) -JDP. With the epoch schedule be $T^{(j)} = 2^j$ for $j = 0, 1, \cdots$ and $\delta = \frac{1}{T}$, it holds that

$$\mathbf{Reg} \leq \tilde{O} \Big(d_{\mathcal{A}} d\kappa_{\nu}^{3/2} \sqrt{T} + \sigma_{\alpha,\beta} d_{\mathcal{A}} d(\kappa_{\nu}^{3/2} + \kappa_{\nu} \sqrt{d}) \Big).$$

This provides a regret of order $\sqrt{T} + \frac{1}{\alpha}$ (omitting poly(d)-factors and logarithmic terms), which partially resolves the open problem stated by Azize and Basu [2024].

As a remark, we note that we state Algorithm 4 for the sake of clarity: In the implementation of Algorithm 3, we do *not* need to range over every context $x \in \mathcal{X}$ to form the policy $\pi^{(j)}$. It is sufficient to compute $\pi^{(j)}(x_t)$ for each round t in jth epoch (according to Eq. (8)). Note that a spanner can be computed in time poly $(d, |\mathcal{A}|)$, and hence Algorithm 3 can be implemented in time poly $(d, |\mathcal{A}|) \cdot T$. Further discussions are deferred to Appendix D.1.

Regret for linear contextual bandits. In linear contextual bandits with a bounded action space \mathcal{A} , we can use a slightly different subroutine for producing confidence intervals (detailed in Appendix D.3), which provides the following refined upper bound.

Theorem 5.2 (Regret upper bound in linear contextual bandits) Algorithm 3 (with the estimation subroutine replaced by Algorithm 12) preserves (α, β) -JDP and guarantees that

$$\mathbf{Reg} \leq \tilde{O}\left(d_{\mathcal{A}}\sqrt{dT\log|\mathcal{A}|} + \sigma_{\alpha,\beta}d_{\mathcal{A}}d^{3/2}\right)$$

In particular, when $|\mathcal{A}| = O(1)$, we obtain a upper bound of $\tilde{O}\left(\sqrt{dT} + \frac{d^{3/2}}{\alpha}\right)$. Note that the \sqrt{dT} -term matches the optimal non-private regret bound for $|\mathcal{A}| = O(1)$ (up to logarithmic factors). Therefore, in this case, privacy is almost *for free*, as the second term (the "price of privacy") grow as $\tilde{O}(1/\alpha)$ and is of lower order as $T \to \infty$.

Extension: Local DP. By using LDP_L1_Regression (Algorithm 8) as the estimation subroutine, we can easily adapt Algorithm 3 so that it preserves local DP. We state the corresponding regret bound as follows and defer the details to Appendix D.4.

Theorem 5.3 (Regret upper bound under LDP) A variant of Algorithm 3 with estimation subroutine Algorithm 8 (detailed in Appendix D.4) preserves (α, β) -LDP and

$$\mathbf{Reg} \leq \tilde{O}\left(\sigma_{\alpha,\beta}d_{\mathcal{A}}d(\kappa_{\nu}\sqrt{d} + \kappa_{\nu}^{3/2})\sqrt{T}\right).$$

Particularly, in linear contextual bandits, Theorem 5.3 provides a regret bound of $\sqrt{d^5T}/\alpha$. While this is a *d*-factor worse than the result of Chen and Rakhlin [2025], our algorithm is computationally efficient, and the number of switches (or, changes) of the deployed decision-channel pair (π_t, \mathbf{Q}_t) is bounded by $O(\kappa_{\nu} \log^2(T))$.

6 Dimension-free Linear Regression

In Section 4 and Section 5, we provided a somewhat satisfactory picture of the optimal rates of private learning in contextual bandits with generalized linear models, when the dimension d is *bounded*. In this section, we turn our focus to the setting where the dimension d is prohibitively large or *unbounded*, e.g., when the linear function parametrization is in fact given by a Reproducing Kernel Hilbert Space (RKHS).³

This setting is fundamentally more challenging, as the following lower bounds indicates. The proof can be found in [Chen and Rakhlin, 2025, Appendix C].

Proposition 6.1 Let $d \ge 1$, covariate space $C = \mathbb{S}^{d-1}$ be the d-dimensional unit sphere. For each $\theta \in \mathbb{S}^{d-1}$, we consider the linear model M_{θ} :

$$(\phi, y) \sim M_{\theta}: \qquad \phi = \theta, y = 1$$

For any parameter $R \in [1, c\sqrt{d}]$ (c > 0 is a small absolute constant), the following holds:

(a) Suppose that Alg is a T-round $(\alpha, 0)$ -JDP algorithm with output $\|\hat{\theta}\| \leq R$. Then it holds that

$$\sup_{\theta^{\star} \in \mathbb{S}^{d-1}} \mathbb{E}^{M_{\theta^{\star}}, \mathsf{Alg}} |\langle \theta^{\star}, \hat{\theta} \rangle - 1| \gtrsim 1, \qquad unless \ T \gtrsim \frac{d}{R^2 \alpha}.$$

(b) Suppose that Alg is a T-round (α, β) -LDP algorithm with output $\|\hat{\theta}\| \leq R$. Then it holds that

$$\sup_{\theta^{\star} \in \mathbb{S}^{d-1}} \mathbb{E}^{M_{\theta^{\star}}, \mathsf{Alg}} |\langle \theta^{\star}, \hat{\theta} \rangle - 1| \gtrsim 1, \qquad unless \ T \gtrsim \min\left\{\frac{d}{R^{2}\alpha^{2}}, \frac{1}{\beta}\right\}.$$

Note that for each linear model \mathbb{P}_{θ} , the covariance matrix $\mathbb{E}_{\mathbb{P}_{\theta}}\phi\phi^{\top} = \theta\theta^{\top}$ is of rank 1. Therefore, Proposition 6.1 has two implications for *pure JDP* (and also LDP) regression with unbounded dimension *d* and unknown covariate distribution:

- (1) Estimating the covariance matrix requires $\Omega(d)$ samples, even when the covariance matrix is known to have rank 1.
- (2) Proper estimator of the parameter θ^* also requires $\Omega(d)$ samples to achieve a non-trivial error. Conversely, any non-trivial estimator $\hat{\theta}$ must have norm $\|\hat{\theta}\| \ge \Omega(\sqrt{d/T})$ (with non-trivial probability).

Therefore, with unbounded dimension d, to achieved estimation guarantees, the estimator in consideration has to be highly *improper*, and it also cannot rely on estimating the covariance matrix.

Based on the observations above, in this section, we develop improper private procedures with *dimension-free* bounds in private linear regression. We then apply the proposed methods to provide dimension-free regret bounds in private linear contextual bandits.

 $^{^{3}}$ Our approach naturally applies to learning in RKHS. However, to avoid measure-theoretic issues, we only present our algorithms for finite dimensional space.

Algorithm 5 JDP_Improper_BatchSGD

Input: Dataset $\mathcal{D} = \{(\phi_t, y_t)\}_{t \in [T]}$. **Input:** Epoch $K \ge 1$, batch size $N = \lfloor \frac{T}{K} \rfloor$, stepsize $\eta = 1$, parameter R = 2. 1: Initialize $\theta_{(0)} = \mathbf{0}$. 2: **for** $k = 0, 1, \dots, K - 1$ **do**

3: Compute gradient estimate

$$g_{(k)} = \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \phi_t(\langle \phi_t, \theta_{(k)} \rangle - y_t)$$

4: Update

$$\theta_{(k+1)} = \operatorname{Proj}_{\mathbb{B}^d(R)} (\theta_{(k)} - \eta g_{(k)}).$$

5: Privatize $\hat{\theta} \sim \mathsf{Priv}_{\eta(R+1)/N}(\theta_{(K)})$ **Output:** Estimator $\hat{\theta}$.

6.1 Private improper batched SGD

We begin with the joint DP setting. For any non-private estimator $\hat{\theta}$ with *sensitivity* s, it is well-known that the estimator $\hat{\theta}' = \hat{\theta} + \zeta$ ensures (α, β) -JDP with noise $\zeta \sim \mathsf{N}(0, \varepsilon^2 \mathbf{I})$ and parameter $\varepsilon := s \cdot \sigma_{\alpha,\beta}$. The key idea is that, while $\|\zeta\| \simeq \varepsilon \sqrt{d}$, we have $\|\zeta\|_{\Sigma} \lesssim \varepsilon$ with high probability, where $\Sigma := \mathbb{E}_{\phi \sim p} \phi \phi^{\top}$ is the covariance matrix. Therefore, to ensure JDP, it is sufficient to privatize a non-private estimator with low sensitivity.

Based on these observations, we consider the projected batched SGD on the standard square-loss:

$$\mathcal{L}_{\mathsf{Sq}}(\theta) = \frac{1}{2} \mathbb{E}_{(\phi, y) \sim M^{\star}} \Big[\left(\langle \phi, \theta \rangle - y \right)^2 \Big].$$

By directly privatizing its last iterate (Algorithm 5), we can achieve a near-optimal convergence rate (detailed in Appendix E.1).

Theorem 6.2 (Dimension-free JDP regression) Let $T \ge 1, \delta \in (0, 1)$. Algorithm 5 preserves (α, β) -JDP, and with a suitably chosen parameter K, it ensures that

$$\mathbb{E}_{\phi \sim p} \langle \phi, \hat{\theta} - \theta^* \rangle^2 = \left\| \hat{\theta} - \theta^* \right\|_{\Sigma}^2 \lesssim \left(\frac{\log T \log(1/\delta)}{T} \right)^{1/2} + \left(\frac{\sigma_{\alpha,\beta} \sqrt{\log(1/\delta)}}{T} \right)^{2/3}$$

Therefore, the estimator $\hat{\theta}$ achieves the *dimension-independent* convergence rate of $\frac{1}{\sqrt{T}} + \frac{1}{(\alpha T)^{2/3}}$. In non-private linear models, the $T^{-1/2}$ -rate of convergence is known to be minimax-optimal and can be achieved by vanilla gradient descent.

Extension: Local DP. The situation under the local DP model is much more subtle. Indeed, one may expect that Algorithm 5 naturally extends to this setting. However, to ensure local privacy in (batched) SGD, for every step, the gradient estimator has to be privatized. To this end, it is typically necessary to add a noise vector ζ that has norm scaling with $\Omega(\sqrt{d})$ (e.g., when ζ is the Gaussian noise). In other words, the privatized gradient estimator has norm scaling with \sqrt{d} . Hence, after a single step of gradient descent, the iterate falls outside the unit ball $\mathbb{B}^d(1)$, and projection back to $\mathbb{B}^d(1)$ can lead to large bias. Therefore, the method of projected gradient descent may not be applied here.

Instead, we consider performing privatized batch SGD directly, replacing the projection operation with a careful clipping on the gradient estimator (Algorithm 6). This is based on extending the aforementioned

Algorithm 6 LDP_Improper_BatchSGD

Input: Round $T \ge 1$. **Input:** Epoch $K \ge 1$, batch size $N = \lfloor \frac{T}{K} \rfloor$, stepsize $\eta = 1$, parameter R = 2.

1: Initialize $\theta_{(0)} = \mathbf{0}$.

- 2: for $k = 0, \cdots, K 1$ do
- 3: **for** $t = kN + 1, \cdots, (k+1)N$ **do**
- 4: Observe $(\phi_t, y_t) \sim p$ and form the gradient estimator

$$g_t = \phi_t(\mathsf{clip}_R(\langle \theta_{(k)}, \phi_t \rangle) - y_t) \tag{9}$$

5: Privatize $\tilde{g}_t \sim \mathsf{Priv}_{R+1}(g_t)$.

6: Compute batched gradient estimator $\tilde{g}_{(k)} = \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \tilde{g}_t$ and update

$$\theta_{(k+1)} = \theta_{(k)} - \eta \widetilde{g}_{(k)}$$

Output: $\hat{\theta} = \theta_{(K)}$.

observation under JDP: When $\zeta \sim \mathsf{N}(0, \varepsilon^2 \mathbf{I})$, while $\|\zeta\| \sim \varepsilon \sqrt{d}$, for the covariate $\phi \sim p$ that is independent of ζ , the random variable $\langle \phi, \zeta \rangle \sim \mathsf{N}(0, \varepsilon^2 \|\phi\|^2)$ is a zero-mean Gaussian random variable conditional on ϕ . Particularly, it holds that $|\langle \phi, \zeta \rangle| \lesssim \varepsilon \|\phi\|$ with high probability with respect to the randomness of the noise ζ and $\phi \sim p$. Using this idea, we can show that Eq. (9) provides an estimator of $\nabla \mathcal{L}_{\mathsf{Sq}}(\theta_{(k)})$ with small bias for all epochs. Here, the clipping operation is defined as

$$\mathsf{clip}_R(v) := \max\left\{\min\left\{v, R\right\}, -R\right\} \in [-R, R], \qquad \forall v \in \mathbb{R}.$$

With a careful analysis that bounds the bias introduced by clipping (9), we provide the following guarantee for Algorithm 6.

Theorem 6.3 (Dimension-free LDP regression) Algorithm 6 preserves (α, β) -LDP. For $T \ge 1, \delta \in (0, 1)$, with a suitable number of epochs $K \ge 1$, Algorithm 6 returns $\hat{\theta}$ that with probability at least $1 - \delta$,

$$\mathbb{E}_{\phi \sim p} \langle \phi, \hat{\theta} - \theta^* \rangle^2 = \left\| \hat{\theta} - \theta^* \right\|_{\Sigma}^2 \lesssim \left(\frac{\sigma_{\alpha, \beta} \log(T/\delta)}{T} \right)^{1/3}$$

This establishes a convergence rate of $\tilde{O}((\alpha^2 T)^{-1/3})$ under the square loss. As shown by Chen and Rakhlin [2025, Corollary I.8], any LDP algorithm must incur an L_1 -error of

$$\Omega\left(\min\left\{\frac{d}{\sqrt{\alpha^2 T}}, \left(\frac{1}{\alpha^2 T}\right)^{1/6}\right\}\right).$$

Therefore, Algorithm 6 achieves the optimal dimension-free convergence rate under L_1 -error, and hence it is optimal under the L_2 -error. Further, Theorem 4.5 and Theorem 6.3 together provide the near-optimal L_1 -estimation error for the full range of $T \in [1, \infty)$ under LDP.

6.2 Application: Linear contextual bandits with dimension-free regret

As an application, we use the dimension-free procedures developed in Section 6.1 as subroutines for learning linear contextual bandits. We invoke the SquareCB algorithm [Abe and Long, 1999, Foster and Rakhlin, 2020, Simchi-Levi and Xu, 2020], which has a regret guarantee given any offline regression oracle with L_2 -error bound. In particular, by instantiating the regression oracle as JDP_Improper_BatchSGD (Algorithm 5) or LDP_Improper_BatchSGD (Algorithm 6), we obtain the following private regret bounds. The details are presented in the Appendix E.3.

Theorem 6.4 (Dimension-free regret bounds) Let $T \ge 1$ and suppose \mathcal{A} is finite.

(1) Suppose that SquareCB (Algorithm 13) is instantiated by the regression subroutine JDP_Improper_BatchSGD (Algorithm 5). Then SquareCB preserves (α, β) -JDP and it ensures

$$\operatorname{\mathbf{Reg}} \leq \sqrt{|\mathcal{A}|} \cdot \tilde{O}\Big(T^{3/4} + \sigma_{\alpha,\beta}^{1/3}T^{2/3}\Big).$$

(2) Suppose that SquareCB (Algorithm 13) is instantiated by the regression subroutine LDP_Improper_BatchSGD (Algorithm 6). Then SquareCB preserves (α, β) -LDP and it ensures

$$\operatorname{\mathbf{Reg}} \le \sqrt{|\mathcal{A}|} \cdot \tilde{O}\left(\sigma_{\alpha,\beta}^{1/3} T^{5/6}\right)$$

To the best of our knowledge, such dimension-free regret bounds are new in private contextual bandits. Under JDP, the regret rate is $\tilde{O}(T^{3/4} + \alpha^{-1/3}T^{2/3})$, and the first term matches the optimal dimension-free $T^{3/4}$ -regret in non-private linear contextual bandits [Abe and Long, 1999, Foster and Rakhlin, 2020], implying that privacy is almost "for free" in this setting. Furthermore, the LDP regret bound scales as $\alpha^{-1/3}T^{5/6}$, which nearly matches the minimax lower bound [Chen and Rakhlin, 2025, Corollary I.15].

Conclusion

In this work, we propose a novel method of private re-weighted regression for private (generalized) linear regression, achieving near-optimal convergence rates under L_1 -error. Based on this method, we provide efficient algorithms for (generalized) linear contextual bandits with near-optimal regret bounds in both the joint and local model of differential privacy. Furthermore, we also develop the improper private procedures with near-optimal, dimension-independent rates in linear models and linear contextual bandits.

Acknowledgements

FC and AR acknowledge support from ARO through award W911NF-21-1-0328, as well as Simons Foundation and the NSF through awards DMS-2031883 and PHY-2019786.

References

- Y. Abbasi-Yadkori, D. Pál, and C. Szepesvári. Improved algorithms for linear stochastic bandits. In Advances in Neural Information Processing Systems, pages 2312–2320, 2011.
- N. Abe and P. M. Long. Associative reinforcement learning using linear probabilistic concepts. In *Proceedings* of the Sixteenth International Conference on Machine Learning, pages 3–11. Morgan Kaufmann Publishers Inc., 1999.
- P. Auer, N. Cesa-Bianchi, and P. Fischer. Finite-time analysis of the multiarmed bandit problem. Machine Learning, 47(2):235–256, 2002.
- B. Awerbuch and R. Kleinberg. Online linear optimization and adaptive routing. Journal of Computer and System Sciences, 74(1):97–114, 2008.
- A. Azize and D. Basu. Open problem: What is the complexity of joint differential privacy in linear contextual bandits? In *The Thirty Seventh Annual Conference on Learning Theory*, pages 5306–5311. PMLR, 2024.
- B. Balle and Y.-X. Wang. Improving the gaussian mechanism for differential privacy: Analytical calibration and optimal denoising. In *International Conference on Machine Learning*, pages 394–403. PMLR, 2018.
- H. Bastani and M. Bayati. Online decision making with high-dimensional covariates. Operations Research, 68(1):276–294, 2020.
- F. Chen and A. Rakhlin. Decision making in changing environments: Robustness, query-based learning, and differential privacy. arXiv preprint arXiv:2501.14928, 2025.

- X. Chen, D. Simchi-Levi, and Y. Wang. Privacy-preserving dynamic personalized pricing with demand learning. *Management Science*, 68(7):4878–4898, 2022.
- J. C. Duchi, M. I. Jordan, and M. J. Wainwright. Local privacy and statistical minimax rates. In 2013 IEEE 54th annual symposium on foundations of computer science, pages 429–438. IEEE, 2013.
- J. C. Duchi, J. Lafferty, Y. Zhu, et al. Local minimax complexity of stochastic convex optimization. Advances in Neural Information Processing Systems, 29, 2016.
- C. Dwork, F. McSherry, K. Nissim, and A. Smith. Calibrating noise to sensitivity in private data analysis. In Theory of Cryptography: Third Theory of Cryptography Conference, TCC 2006, New York, NY, USA, March 4-7, 2006. Proceedings 3, pages 265–284. Springer, 2006.
- C. Dwork, A. Roth, et al. The algorithmic foundations of differential privacy. Foundations and Trends® in Theoretical Computer Science, 9(3–4):211–407, 2014.
- D. J. Foster and A. Rakhlin. Beyond UCB: Optimal and efficient contextual bandits with regression oracles. arXiv preprint arXiv:2002.04926, pages 3199–3210, 2020.
- D. J. Foster, A. Agarwal, M. Dudík, H. Luo, and R. E. Schapire. Practical contextual bandits with regression oracles. arXiv preprint arXiv:1803.01088, 2018.
- D. J. Foster, S. M. Kakade, J. Qian, and A. Rakhlin. The statistical complexity of interactive decision making. arXiv preprint arXiv:2112.13487, 2021.
- D. J. Foster, A. Rakhlin, A. Sekhari, and K. Sridharan. On the complexity of adversarial decision making. Advances in Neural Information Processing Systems, 35:35404–35417, 2022.
- D. J. Foster, N. Golowich, and Y. Han. Tight guarantees for interactive decision making with the decisionestimation coefficient. In *The Thirty Sixth Annual Conference on Learning Theory*, pages 3969–4043. PMLR, 2023.
- Y. Han, Z. Liang, Y. Wang, and J. Zhang. Generalized linear bandits with local differential privacy. Advances in Neural Information Processing Systems, 34:26511–26522, 2021.
- E. Hazan and Z. Karnin. Volumetric spanners: an efficient exploration basis for learning. The Journal of Machine Learning Research, 17(1):4062–4095, 2016.
- J. He, J. Zhang, and R. Q. Zhang. A reduction from linear contextual bandits lower bounds to estimations lower bounds. In *International Conference on Machine Learning*, pages 8660–8677. PMLR, 2022.
- V. Karwa and S. Vadhan. Finite sample differentially private confidence intervals. arXiv preprint arXiv:1711.03908, 2017.
- S. P. Kasiviswanathan, H. K. Lee, K. Nissim, S. Raskhodnikova, and A. Smith. What can we learn privately? SIAM Journal on Computing, 40(3):793–826, 2011.
- T. Lattimore and C. Szepesvári. Exploration by optimisation in partial monitoring. In Conference on Learning Theory, pages 2488–2515. PMLR, 2020.
- Y. Lei, S. Miao, and R. Momot. Privacy-preserving personalized revenue management. *Management Science*, 70(7):4875–4892, 2024.
- J. Li, D. Simchi-Levi, and Y. Wang. On the optimal regret of locally private linear contextual bandit. arXiv preprint arXiv:2404.09413, 2024.
- Y. Li, Y. Wang, and Y. Zhou. Nearly minimax-optimal regret for linearly parameterized bandits. In Conference on Learning Theory, pages 2173–2174. PMLR, 2019.
- V. Perchet, P. Rigollet, S. Chassang, and E. Snowberg. Batched bandit problems. 2016.
- R. Shariff and O. Sheffet. Differentially private contextual linear bandits. Advances in Neural Information Processing Systems, 31, 2018.

- D. Simchi-Levi and Y. Xu. Bypassing the monster: A faster and simpler optimal algorithm for contextual bandits under realizability. arXiv preprint arXiv:2003.12699, 2020.
- R. Vershynin. High-dimensional probability, 2019.
- G. Vietri, B. Balle, A. Krishnamurthy, and S. Wu. Private reinforcement learning with pac and regret guarantees. In *International Conference on Machine Learning*, pages 9754–9764. PMLR, 2020.
- K. Zheng, T. Cai, W. Huang, Z. Li, and L. Wang. Locally differentially private (contextual) bandits learning. Advances in Neural Information Processing Systems, 33:12300–12310, 2020.

A Technical tools

Lemma A.1 (Gaussian concentration) For the random vector $z \in \mathbb{R}^d$ with *i.i.d.* $\mathsf{N}(0, \sigma^2)$ entries, with probability at least $1 - \delta$,

$$\frac{\|z\|^2}{\sigma^2} \le d + 2\sqrt{d\log(1/\delta)} + 2\log(1/\delta) \le 2d + 3\log(1/\delta).$$

For the random symmetric matrix $Z \in \mathbb{R}^{d \times d}$ with i.i.d. Gaussian entries $Z_{ij} = Z_{ji} \sim \mathsf{N}(0, \sigma^2)$ $(1 \le i \le j \le d)$, with probability at least $1 - \delta$,

$$\|Z\|_{\rm op} \le C\sigma\sqrt{d + \log(1/\delta)},$$

where C is an absolute constant.

Lemma A.2 (Matrix Bernstein's inequality (Vershynin [2019, Theorem 5.4.1])) Let X_1, X_2, \ldots, X_n be i.i.d. zero-mean symmetric matrices in $\mathbb{R}^{d \times d}$. Suppose that $\|X_1\|_{op} \leq R$ almost surely, and let $\sigma^2 = \|\mathbb{E}[X_1^2]\|_{op}$. Then, for all $t \geq 0$,

$$\mathbb{P}\left(\left\|\sum_{i=1}^{n} X_{i}\right\| \geq t\right) \leq 2d \exp\left(-\frac{t^{2}/2}{n\sigma^{2} + Rt/3}\right).$$

Lemma A.3 Suppose that V_1, \dots, V_n are *i.i.d.* positive semi-definite random matrices such that $\mathbb{E}[V_i] = \Sigma$ and $\|V_i\|_{op} \leq 1$ almost surely. Then for any fixed parameter $c \in [0, 1]$, with probability at least $1 - \delta$:

$$(1-c)\Sigma - \frac{4\log(2d/\delta)}{nc}\mathbf{I} \preceq \frac{1}{n}\sum_{i=1}^{n}V_i \preceq (1+c)\Sigma + \frac{4\log(2d/\delta)}{nc}$$

Proof. Fix a $\lambda > 0$ to be specified later. Let $\widetilde{V}_i = (\Sigma + \lambda \mathbf{I})^{-1/2} V_i (\Sigma + \lambda \mathbf{I})^{-1/2}$. Note that $\|\widetilde{V}_i\|_{\text{op}} \leq \frac{1}{\lambda}$ and $\mathbb{E}\widetilde{V} \leq \mathbf{I}$, and hence

$$\mathbb{E}\widetilde{V}_i^2 \preceq \mathbb{E}\frac{1}{\lambda}\widetilde{V}_i \preceq \frac{1}{\lambda}\mathbf{I}$$

Then by Lemma A.2, with probability at least $1 - \delta$,

$$\left\|\sum_{i=1}^{n} \widetilde{V}_{i} - \mathbb{E}\widetilde{V}_{i}\right\|_{\text{op}} \leq \sqrt{\frac{2n}{\lambda}\log(2d/\delta)} + \frac{2\log(2d/\delta)}{3\lambda} =: c_{\lambda}.$$

Under this event, we have $v^{\top} \left(\sum_{i=1}^{n} \widetilde{V}_{i} - \mathbb{E}\widetilde{V}_{i} \right) v \leq c_{\lambda} \|v\|^{2}$ for all $v \in \mathbb{R}^{d}$, and hence

$$n\Sigma - c_{\lambda}(\Sigma + \lambda \mathbf{I}) \preceq \sum_{i=1}^{n} V_i \preceq n\Sigma + c_{\lambda}(\Sigma + \lambda \mathbf{I}).$$

Therefore, we choose $\lambda = \frac{4 \log(2d/\delta)}{nc^2}$, which ensures $c_{\lambda} \leq nc$ and

$$(1-c)\Sigma - c\lambda \mathbf{I} \preceq \frac{1}{n} \sum_{i=1}^{n} V_i \preceq (1+c)\Sigma + c\lambda \mathbf{I},$$

with probability at least $1 - \delta$.

Lemma A.4 Suppose that $(X_t)_t$ is a vector-valued martingale-difference sequence adapted to the filtration $(\mathcal{F}_t)_t$. Assume that $||X_t||_2 \leq c_t$ almost surely. Then it holds that $\forall x \geq 0$

$$\mathbb{P}\left(\left\|\sum_{t=1}^{n} X_i\right\|_2 \ge \sqrt{\sum_{t=1}^{n} c_t^2 + x}\right) \le \exp\left(-\frac{x^2}{2\sum_{t=1}^{n} c_t^2}\right)$$

Proof. This is a direct corollary of Azuma's inequality by considering the martingale (Z_t) given by $Z_t = \mathbb{E}[||S_n||_2|\mathcal{F}_t]$, where $S_n = \sum_{t=1}^n X_i$. We have $Z_0 = \mathbb{E}[||S_n||_2]$, $Z_n = ||S_n||_2$, and $|Z_{t+1} - Z_t| \leq c_{t+1}$ for $0 \leq t \leq n-1$. Therefore, Azuma's inequality implies that

$$\mathbb{P}(Z_n - Z_0 \ge x) \le \exp\left(-\frac{x^2}{2\sum_{t=1}^n c_t^2}\right).$$

The desired inequality follows from the fact that $Z_0 = \mathbb{E}[||S_n||_2] \le \sqrt{\mathbb{E}||S_n||_2^2} \le \sqrt{\sum_{t=1}^n c_t^2}$.

Lemma A.5 For any vector $\phi \in \mathbb{R}^d$ and positive-definite matrix U, the matrix $\Phi = \frac{U^{1/2}\phi\phi^{\top}U^{1/2}}{\|U\phi\|}$ is bounded under the Frobenius norm: $\|\Phi\|_{\mathbf{F}} \leq \|\phi\|$.

Proof. Denote $U^{1/2}\phi = (v_1, \cdots, v_d)$, then $\Phi = ||U\phi||^{-1}(v_iv_j)_{i,j} \in \mathbb{R}^{d \times d}$, and hence

$$\|\Phi\|_F = \|U\phi\|^{-1} \left(\sum_{i=1}^d v_i^2\right) = \|U\phi\|^{-1} \left\|U^{1/2}\phi\right\|^2 = \|U\phi\|^{-1}\phi^T U\phi \le \|U\phi\|^{-1} \|U\phi\| \|\phi\| = \|\phi\|.$$

Lemma A.6 (Monotone matrix operation) Suppose that $A, B \in \mathbb{R}^{d \times d}$ are PSD matrices, B is invertible, and $A \leq B$. Then it holds that $A^{-1} \succeq B^{-1}$ and $A^{1/2} \leq B^{1/2}$.

Proof. Because $A \leq B$, we have $(A^{1/2}B^{-1/2})^{\top}(A^{1/2}B^{-1/2}) = B^{-1/2}AB^{-1/2} \leq \mathbf{I}$, and hence $A^{1/2}B^{-1}A^{1/2} = (A^{1/2}B^{-1/2})(A^{1/2}B^{-1/2})^{\top} \leq \mathbf{I}$, which implies $B^{-1} \leq A^{-1}$.

Similarly, we have $||A^{1/2}v|| \leq ||B^{1/2}v||$ for all $v \in \mathbb{R}^d$, and hence $||A^{1/2}B^{-1/2}||_{\text{op}} \leq 1$. Let $D = B^{-1/4}A^{1/2}B^{-1/4}$. Because D is symmetric, all eigenvalues of D are real numbers. For any eigenvalue λ of D, we take an eigenvector v of λ , i.e., $Dv = \lambda v$. Then, $A^{1/2}B^{-1/2}(B^{1/4}v) = \lambda(B^{1/4}v)$, and hence $|\lambda| \leq 1$. Therefore, we can conclude that all eigenvalues of D are bounded by 1, and hence $||D||_{\text{op}} \leq 1$, which implies $A^{1/2} \leq B^{1/2}$. \Box

Lemma A.7 Let Σ be a positive semi-definite matrix with $\Sigma \leq \mathbf{I}$, and $\eta \in [0,1]$. For any integer $k \geq 0$, we have $\|\Sigma^{1/2}(\mathbf{I} - \eta\Sigma)^k\|_{\text{op}} \leq \sqrt{\frac{2e}{\eta k}}$.

Proof. Let the $1 \ge \lambda_1 \ge \cdots \ge \lambda_d \ge 0$ be the spectrum of Σ . Then

$$\left\|\boldsymbol{\Sigma}^{1/2}(\mathbf{I}-\eta\boldsymbol{\Sigma})^{k}\right\|_{\mathrm{op}} = \max_{i\in[d]}\sqrt{\lambda_{i}}(1-\eta\lambda_{i})^{k} \leq \sup_{\lambda\in[0,1]}\sqrt{\lambda}e^{-\eta k\lambda} \leq \sqrt{\frac{2e}{\eta k}}.$$

B Proofs from Section 3

B.1 Proof of Proposition 3.1

Proposition 3.1 (2) is a well-known result (see e.g., Lemma 25 and Appendix B.2 of [Chen and Rakhlin, 2025]). In the following, we prove Proposition 3.1 (1) using the same idea. We invoke the following lemma of Karwa and Vadhan [2017].

Lemma B.1 Fix a non-interactive (α, β) -JDP algorithm $\operatorname{Alg} : \mathcal{Z}^T \to \Delta(\Pi)$. For any model $M \in \Delta(\mathcal{Z})$, we let $\mathbb{P}^{M,\operatorname{Alg}} \in \Delta(\Pi)$ be the distribution of $\pi \sim \operatorname{Alg}(z_1, \cdots, z_T)$ under i.i.d $z_1, \cdots, z_T \sim M$. Then for any models $M, \overline{M} \in \Delta(\mathcal{Z})$, it holds that

$$\mathbb{P}^{M,\mathsf{Alg}}(E) \le e^{\alpha_T} \mathbb{P}^{\overline{M},\mathsf{Alg}}(E) + \beta_T, \qquad \forall E \subseteq \Pi,$$

where $\alpha_T = 6\alpha T D_{\text{TV}}(M, \overline{M})$ and $\beta_T = 4e^{\alpha_T} T \beta D_{\text{TV}}(M, \overline{M})$.

Now, for any $\theta \in [-1, 1]$, we let M_{θ} be the distribution of (x, y) under $\phi \sim p, y | \phi \sim \text{Rad}(\phi \theta)$. Then, by definition,

$$D_{\mathrm{TV}}(M_{\theta}, M_0) = \mathbb{E}_{\phi \sim p} \frac{1}{2} |\phi \theta| = \frac{\lambda |\theta|}{2}.$$

Let $\theta = \min\left\{1, \frac{1}{20\lambda T(\alpha+\beta)}\right\}$. Then, by Lemma B.1, we have

$$\mathbb{P}^{M_{\theta}, \operatorname{Alg}}(E) \le 1.4 \ \mathbb{P}^{M_{0}, \operatorname{Alg}}(E) + \frac{1}{4}, \qquad \forall E \subseteq \Pi.$$

In this problem, the decision space is $\Pi = [-1, 1]$, and the loss function is $L(\theta, \pi) = \mathbb{E}_{\phi \sim p} \left[|\phi(\pi - \theta)|^2 \right] = \lambda^2 |\pi - \theta|^2$. We consider the event $E = \{\pi : \pi > \frac{\theta}{2}\}$. By definition,

$$\mathbb{E}^{_{M_{\theta}},\mathrm{Alg}}L(\theta,\pi) \geq \frac{\lambda^{2}\theta^{2}}{4} \mathbb{P}^{_{M_{\theta}},\mathrm{Alg}}(E^{c}), \qquad \mathbb{E}^{_{M_{0}},\mathrm{Alg}}L(0,\pi) \geq \frac{\lambda^{2}\theta^{2}}{4} \mathbb{P}^{_{M_{0}},\mathrm{Alg}}(E).$$

Thus,

$$\max \left\{ \mathbb{E}^{M_{\theta}, \operatorname{Alg}} L(\theta, \pi), \mathbb{E}^{M_{0}, \operatorname{Alg}} L(0, \pi) \right\} \geq \frac{\lambda^{2} \theta^{2}}{8} (\mathbb{P}^{M_{\theta}, \operatorname{Alg}}(E^{c}) + \mathbb{P}^{M_{0}, \operatorname{Alg}}(E))$$
$$\geq \frac{\lambda^{2} \theta^{2}}{8} \left(\mathbb{P}^{M_{\theta}, \operatorname{Alg}}(E^{c}) + 0.7 \ \mathbb{P}^{M_{\theta}, \operatorname{Alg}}(E) - \frac{1}{5} \right)$$
$$\geq \frac{\lambda^{2} \theta^{2}}{16} \geq c_{0} \min \left\{ \lambda, \frac{1}{\lambda T^{2}(\alpha + \beta)^{2}} \right\}.$$

This is the desired lower bound.

C Proofs from Section 4

C.1 Proof of Lemma 4.1

By definition, $\nabla \mathcal{L}_U(w^*) = \lambda \mu_{\nu} \cdot Uw^*$, and hence $\|\nabla \mathcal{L}_U(w^*)\| \leq \mu_{\nu} \lambda$. Then, because \mathcal{L}_U is $(\mu_{\nu}/2)$ -strongly convex over $w \in \mathcal{W}$, we have

$$\mathcal{L}_U(w) \ge \mathcal{L}_U(w^*) + \langle \nabla \mathcal{L}_U(w^*), w - w^* \rangle + \frac{\mu_{\nu}}{4} \|w - w^*\|^2, \qquad w \in \mathcal{W}.$$

In particular, using $\mathcal{L}_U(\widehat{w}_U^{\star}) \leq \mathcal{L}_U(w^{\star})$, we have

$$\frac{\mu_{\nu}}{4} \left\| \widehat{w}_{U}^{\star} - w^{\star} \right\|^{2} \leq -\langle \nabla \mathcal{L}_{U}(w^{\star}), \widehat{w}_{U}^{\star} - w^{\star} \rangle \leq \mu_{\nu} \lambda \left\| \widehat{w}_{U}^{\star} - w^{\star} \right\|,$$

and hence $\|\widehat{w}_U^{\star} - w^{\star}\| \leq 4\lambda$.

Next, for any vector $v \in \mathbb{R}^d$, we have

$$\mathbb{E}_{\phi \sim p}[|\langle \phi, Uv \rangle|]^2 \le \mathbb{E}_{\phi \sim p}[||U\phi||] \cdot \mathbb{E}_{\phi \sim p}\left[\frac{\langle \phi, Uv \rangle^2}{||U\phi||}\right].$$

Note that $\mathbb{E}_{\phi \sim p} \left[\frac{U \phi \phi^{\top} U}{\|U \phi\|} \right] \preceq 2\mathbf{I}$, and hence we have $\mathbb{E}_{\phi \sim p} [\|U \phi\|] \leq 2d$ and $\mathbb{E}_{\phi \sim p} \left[\frac{\langle \phi, U v \rangle^2}{\|U \phi\|} \right] \leq 2 \|v\|^2$. Therefore, it holds that

$$\mathbb{E}_{\phi \sim p}[|\langle \phi, Uv \rangle|] \le 2\sqrt{d} \, \|v\| \, .$$

Substituting $v = w - w^*$ completes the proof.

C.2 Proof of Lemma 4.2

We first note that in Algorithm 1, the dataset \mathcal{D} is split equally as $\mathcal{D} = \mathcal{D}_{(0)} \sqcup \cdots \sqcup \mathcal{D}_{(K-1)}$, and iteration at the *k*th epoch can be regarded as a random function $(U_{(k)}; \mathcal{D}_{(k)}) \mapsto U_{(k+1)}$. Therefore, using the composition property of joint DP (Lemma C.1), we only need to verify that $(U_{(k)}; \mathcal{D}_{(k)}) \mapsto U_{(k+1)}$ preserves (α, β) -JDP (with respect to $\mathcal{D}_{(k)}$).

For the data (ϕ_t, y_t) in the *k*th epoch, the quantity $\Phi_t = \frac{U_{(k)}^{1/2} \phi_t \phi_t^\top U_{(k)}^{1/2}}{\|U_{(k)} \phi_t\|}$ can be bounded as $\|\Phi_t\|_{\rm F} \leq 1$ (Lemma A.5). Thus, $H_{(k)}$ defined in Eq. (6) has sensitivity $\Delta = 1/N$ under Frobenius norm. Therefore, by the privacy guarantee of Gaussian channels (Definition 2), the mechanism $(U_{(k)}; \mathcal{D}_{(k)}) \mapsto \tilde{H}_{(k)}$ preserves (α, β) -JDP with respect to $\mathcal{D}_{(k)}$. Consequently, by the post-processing property, $(U_{(k)}; \mathcal{D}_{(k)}) \mapsto U_{(k)}$ also preserves (α, β) -JDP. Therefore, by applying the composition property (Lemma C.1) inductively, we have shown that Algorithm 2 also preserves (α, β) -JDP.

Lemma C.1 (Iterative composition of JDP) Suppose the algorithm $Alg : \mathbb{Z}^{N_1+N_2} \to \Delta(\Pi)$ outputs $\pi \sim Alg(z_1, \dots, z_{N_1+N_2})$ generated as

$$\pi_1 \sim \mathsf{Alg}_1(z_1, \cdots, z_{N_1}), \quad \pi \sim \mathsf{Alg}_2(\pi_1; z_{N_1+1}, \cdots, z_{N_1+N_2}),$$

where the algorithm $\operatorname{Alg}_1 : \mathbb{Z}^{N_1} \to \Delta(\Pi_1)$ preserves (α, β) -JDP, and $\operatorname{Alg}_2(\pi_1; \cdot) : \mathbb{Z}^{N_2} \to \Delta(\Pi)$ preserves (α, β) -JDP for any $\pi_1 \in \Pi_1$. Then Alg preserves (α, β) -JDP.

C.2.1 Proof of Lemma C.1

For ease of presentation, we only consider the case both Π and Π_1 are discrete. For two neighbored dataset $\mathcal{D} = \{z_i\}_{i=1}^{N_1+N_2}$ and $\mathcal{D}' = \{z'_i\}_{i=1}^{N_1+N_2}$, denote

$$\mathcal{D}_1 = \{z_1, \cdots, z_{N_1}\}, \qquad \mathcal{D}_2 = \{z_{N_1+1}, \cdots, z_{N_1+N_2}\}, \\ \mathcal{D}'_1 = \{z'_1, \cdots, z'_{N_1}\}, \qquad \mathcal{D}'_2 = \{z'_{N_1+1}, \cdots, z'_{N_1+N_2}\}.$$

Assume that \mathcal{D} and \mathcal{D}' differ at index $t \in [N_1 + N_2]$, i.e., $z_j = z'_j$ for $j \neq t$. We consider two cases.

Case 1: $t \leq N_1$. Because Alg₁ preserves (α, β) -JDP, we have

τ

$$\mathsf{Alg}_1(E_1|\mathcal{D}_1) \le e^{\alpha}\mathsf{Alg}_1(E_1|\mathcal{D}_1) + \beta, \qquad \forall E_1 \subseteq \Pi_1,$$

and hence, equivalently, it holds that

$$\sum_{\mathbf{r}_1\in\Pi_1} \left[\mathsf{Alg}_1(\pi_1|\mathcal{D}_1) - e^{\alpha}\mathsf{Alg}_1(\pi_1|\mathcal{D}_1')\right]_+ \leq \beta.$$

Note that for any $E \subseteq \Pi$,

$$\mathsf{Alg}(E|\mathcal{D}) = \sum_{\pi_1 \in \Pi_1} \mathsf{Alg}_1(\pi_1|\mathcal{D}_1) \cdot \mathsf{Alg}_2(E|\pi_1;\mathcal{D}_2),$$

and therefore,

$$\begin{split} \mathsf{Alg}(E|\mathcal{D}) - e^{\alpha}\mathsf{Alg}(E|\mathcal{D}') &= \sum_{\pi_1 \in \Pi_1} \left[\mathsf{Alg}_1(\pi_1|\mathcal{D}_1) - e^{\alpha}\mathsf{Alg}_1(\pi_1|\mathcal{D}'_1)\right] \cdot \mathsf{Alg}_2(E|\pi_1;\mathcal{D}_2) \\ &\leq \sum_{\pi_1 \in \Pi_1} \left[\mathsf{Alg}_1(\pi_1|\mathcal{D}_1) - e^{\alpha}\mathsf{Alg}_1(\pi_1|\mathcal{D}'_1)\right]_+ \leq \beta, \end{split}$$

where we use the fact that $Alg_2(E|\pi_1; \mathcal{D}_2) \in [0, 1]$ for any $\pi_1 \in \Pi_1, E \subseteq \Pi$.

Case 2: $t > N_1$. In this case, because $Alg_2(\pi_1; \cdot)$ preserves (α, β) -JDP for any $\pi_1 \in \Pi_1$, for any $E \subseteq \Pi$, we have

$$\begin{aligned} \mathsf{Alg}(E|\mathcal{D}) &= \sum_{\pi_1 \in \Pi_1} \mathsf{Alg}_1(\pi_1|\mathcal{D}_1) \cdot \mathsf{Alg}_2(E|\pi_1;\mathcal{D}_2) \\ &\leq \sum_{\pi_1 \in \Pi_1} \mathsf{Alg}_1(\pi_1|\mathcal{D}_1) \cdot [e^{\alpha}\mathsf{Alg}_2(E|\pi_1;\mathcal{D}_2') + \beta] \\ &= e^{\alpha}\mathsf{Alg}(E|\mathcal{D}') + \beta. \end{aligned}$$

Combining the inequalities above from both cases completes the proof.

C.3 Proof of Proposition 4.3

The following lemma is a standard concentration result (following from taking union bounds with Lemma A.1 and Lemma A.4).

Lemma C.2 In Algorithm 1, with probability at least $1 - \delta$, the following inequalities hold simultaneously for all $k = 0, \dots, K - 1$:

$$\left\| H_{(k)} - \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k)}^{1/2} \phi \phi^{\top} U_{(k)}^{1/2}}{\| U_{(k)} \phi \|} \right] \right\|_{\mathcal{F}} \le C_0 \sqrt{\frac{\log(K/\delta)}{N}},\tag{10}$$

$$\left\| \widetilde{H}_{(k)} - H_{(k)} \right\|_{\text{op}} \le C_0 \frac{\sigma_{\alpha,\beta} \sqrt{d + \log(K/\delta)}}{N},\tag{11}$$

where C_0 is a large absolute constant. In the following, we denote this event as \mathcal{E} .

Therefore, we can simplify the iterations in Algorithm 1 as follows: $U_{(0)} = \mathbf{I}$, and for $k = 0, 1, \dots, K$:

$$\widetilde{H}_{(k)} = \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k)}^{1/2} \phi \phi^{\top} U_{(k)}^{1/2}}{\|U_{(k)} \phi\|} \right] + E_{(k)},$$
(12)

$$\Sigma_{(k)} = U_{(k)}^{1/2} \widetilde{H}_{(k)} U_{(k)}^{1/2} + \lambda_{(k)} U_{(k)}, \qquad (13)$$

$$U_{(k+1)} = \operatorname{sym}(\Sigma_{(k)}^{-1/2}U_{(k)}), \tag{14}$$

where sym $(A) = (A^{\top}A)^{1/2}$, $E_{(k)}$ is a symmetric matrix. We note that Algorithm 1 does not actually compute $(H_{(K)}, \Sigma_{(K)})$, and they only appear in our analysis (where we can regard $E_{(K)} = 0$).

Proposition C.3 Suppose that the sequence of matrices $\{(U_{(k)}, H_{(k)}, \Sigma_{(k)})\}$ is defined recursively by Eq. (12) - (14), with $||E_{(k)}||_{op} \leq \varepsilon$. Suppose that $\lambda_{(k)} = (2k+5)\varepsilon$, and $\varepsilon \leq 0.1$. Then, for any $k \geq 1$, it holds that

$$\lambda_{\min}(\boldsymbol{\Sigma}_{\scriptscriptstyle (k)}) \geq \exp{\left(-\frac{\log(1/\varepsilon)}{2^k}\right)}, \qquad \lambda_{\max}(\boldsymbol{\Sigma}_{\scriptscriptstyle (k)}) \leq \exp{\left(\frac{12}{k}\right)}.$$

In particular, $K \geq \log \log(1/\varepsilon) \vee 20$, we have

$$\frac{1}{2}\mathbf{I} \preceq \mathbb{E}_{\phi \sim p} \left[\frac{U_{\scriptscriptstyle (K)} \phi \phi^{\top} U_{\scriptscriptstyle (K)}}{\|U_{\scriptscriptstyle (K)} \phi\|} \right] + \lambda_{\scriptscriptstyle (K)} U_{\scriptscriptstyle (K)} \preceq 2\mathbf{I}.$$

Proof of Proposition 4.3. By Lemma C.2, under the event \mathcal{E} , the matrix $E_{(k)} = \widetilde{H}_{(k)} - \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k)}^{1/2} \phi \phi^\top U_{(k)}^{1/2}}{\|U_{(k)} \phi\|} \right]$ is bounded as $\|E_{(k)}\|_{\text{op}} \leq \varepsilon_N$. Therefore, under \mathcal{E} , we can apply Proposition C.3, which gives the desired results.

C.3.1 Proof of Lemma C.2

In Algorithm 1, the matrix $\widetilde{H}_{(k)}$ is given by $\widetilde{H}_{(k)} = H_{(k)} + Z_{(k)}$,

$$H_{(k)} = \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \Phi_t, \qquad \Phi_t = \frac{U_{(k)}^{1/2} \phi_t \phi_t^\top U_{(k)}^{1/2}}{\|U_{(k)} \phi_t\|},$$

where $Z_{(k)}$ has i.i.d entries $Z_{ij} = Z_{ji} \sim \mathsf{N}(0, \frac{4\sigma_{\alpha,\beta}^2}{N})$, and $\Phi_{kN+1}, \cdots, \Phi_{(k+1)N}$ are independent (conditional on $U_{(k)}$). Further, by Lemma A.5, we have $\|\Phi_t\|_{\mathrm{F}} \leq 1$, and $\mathbb{E}[\Phi_t|U_{(k)}] = \mathbb{E}_{\phi \sim p}\left[\frac{U_{(k)}^{1/2}\phi\phi^\top U_{(k)}^{1/2}}{\|U_{(k)}\phi\|}\right]$. Therefore, using Lemma A.4, we have with probability at least $1 - \delta$,

$$\left\| H_{(k)} - \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k)}^{1/2} \phi \phi^{\top} U_{(k)}^{1/2}}{\| U_{(k)} \phi \|} \right] \right\|_{\mathcal{F}} \le \frac{1 + \sqrt{2 \log(1/\delta)}}{\sqrt{N}}$$

By Lemma A.1, we also have $\|Z_{(k)}\|_{\text{op}} \leq C \frac{\sigma_{\alpha,\beta} \sqrt{d + \log(1/\delta)}}{N}$ with probability at least $1 - \delta$. Taking the union bound over $k = 0, 1, \dots, K - 1$ and rescaling $\delta \leftarrow \frac{\delta}{2K}$ completes the proof.

C.3.2 Proof of Proposition C.3

By definition (14), for each $k \ge 1$, there exists an orthogonal matrix $V_{(k+1)}$ such that $U_{(k+1)} = V_{(k+1)} \Sigma_{(k)}^{-1/2} U_{(k)}$. Therefore,

$$\begin{split} \mathbf{I} &= V_{(k+1)} V_{(k+1)}^{\top} = V_{(k+1)} \boldsymbol{\Sigma}_{(k)}^{-1/2} \left(U_{(k)}^{1/2} \widetilde{H}_{(k)} U_{(k)}^{1/2} + \lambda_{(k)} U_{(k)} \right) \boldsymbol{\Sigma}_{(k)}^{-1/2} V_{(k+1)}^{\top} \\ &= \mathbb{E}_{\phi \sim p} \left[\frac{V_{(k+1)} \boldsymbol{\Sigma}_{(k)}^{-1/2} U_{(k)} \phi \phi^{\top} U_{(k)} \boldsymbol{\Sigma}_{(k)}^{-1/2} V_{(k+1)}^{\top}}{\|U_{(k)} \phi\|} \right] + V_{(k+1)} \boldsymbol{\Sigma}_{(k)}^{-1/2} U_{(k)}^{1/2} (E_{(k)} + \lambda_{(k)} \mathbf{I}) U_{(k)}^{1/2} \boldsymbol{\Sigma}_{(k)}^{-1/2} V_{(k+1)}^{\top} \\ &= \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right] + U_{(k+1)} U_{(k)}^{-1/2} (E_{(k)} + \lambda_{(k)} \mathbf{I}) U_{(k)}^{-1/2} U_{(k+1)}. \end{split}$$

Therefore, using $-\varepsilon \mathbf{I} \preceq E_{(k)} \preceq \varepsilon \mathbf{I}$ (because $\|E_{(k)}\|_{\mathrm{op}} \leq \varepsilon$),

$$\mathbf{I} - (\lambda_{(k)} + \varepsilon) U_{(k+1)} U_{(k)}^{-1} U_{(k+1)} \preceq \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right] \preceq \mathbf{I} - (\lambda_{(k)} - \varepsilon) U_{(k+1)} U_{(k)}^{-1} U_{(k+1)}.$$

Notice that for any $\phi \in \mathbb{R}^d$, it holds that $\|U_{(k+1)}\phi\| = \|\mathbf{\Sigma}_{(k)}^{-1/2}U_{(k)}\phi\|$, and hence we have

$$\sqrt{\lambda_{\min}(\boldsymbol{\Sigma}_{(k)})} \cdot \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right] \preceq \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k+1)} \phi\|} \right] \preceq \sqrt{\lambda_{\max}(\boldsymbol{\Sigma}_{(k)})} \cdot \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right].$$

Now, combining the inequalities above and using the definition of $\Sigma_{(k+1)}$, we can lower bound

$$\begin{split} \mathbf{\Sigma}_{(k+1)} &= \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k+1)} \phi\|} \right] + U_{(k+1)}^{1/2} E_{(k+1)} U_{(k+1)}^{1/2} + \lambda_{(k+1)} U_{(k+1)} \\ &\succeq \sqrt{\lambda_{\min}(\mathbf{\Sigma}_{(k)})} \cdot \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right] + (\lambda_{(k+1)} - \varepsilon) U_{(k+1)} \\ &\succeq \sqrt{\lambda_{\min}(\mathbf{\Sigma}_{(k)})} \left(\mathbf{I} - (\lambda_{(k)} + \varepsilon) U_{(k+1)} U_{(k)}^{-1} U_{(k+1)} \right) + (\lambda_{(k+1)} - \varepsilon) U_{(k+1)} \end{split}$$

Notice that $U_{(k+1)}^2 = U_{(k)} \Sigma_{(k)}^{-1} U_{(k)} \preceq \frac{1}{\lambda_{\min}(\Sigma_{(k)})} U_{(k)}^2$, and the matrix function $U \mapsto -U^{-1/2}$ is matrix monotone (cf. Lemma A.6), and hence we have

$$\frac{1}{\sqrt{\lambda_{\max}(\mathbf{\Sigma}_{(k)})}} U_{(k+1)}^{-1} \preceq U_{(k)}^{-1} \preceq \frac{1}{\sqrt{\lambda_{\min}(\mathbf{\Sigma}_{(k)})}} U_{(k+1)}^{-1}.$$
(15)

In particular, we have proven $\Sigma_{(k+1)} \succeq \sqrt{\lambda_{\min}(\Sigma_{(k)})} \mathbf{I}$, which implies

$$\lambda_{\min}(\mathbf{\Sigma}_{(k)}) \ge \lambda_{\min}(\mathbf{\Sigma}_{(0)})^{\frac{1}{2^k}} \ge \exp\left(-\frac{\log(1/\varepsilon)}{2^k}\right),$$

where we use $\lambda_{\min}(\mathbf{\Sigma}_{(0)}) \ge \lambda_{(0)} - \varepsilon \ge \varepsilon$.

Similarly,

$$\begin{split} \mathbf{\Sigma}_{(k+1)} &\preceq \sqrt{\lambda_{\max}(\mathbf{\Sigma}_{(k)})} \cdot \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right] + (\lambda_{(k+1)} + \varepsilon) U_{(k+1)} \\ &\preceq \sqrt{\lambda_{\max}(\mathbf{\Sigma}_{(k)})} (\mathbf{I} - (\lambda_{(k)} - \varepsilon) U_{(k+1)} U_{(k)}^{-1} U_{(k+1)}) + (\lambda_{(k+1)} + \varepsilon) U_{(k+1)} \\ &\preceq \sqrt{\lambda_{\max}(\mathbf{\Sigma}_{(k)})} \mathbf{I} + 4\varepsilon U_{(k+1)}. \end{split}$$

Note that we have also shown that $\Sigma_{(k+1)} \succeq (\lambda_{(k+1)} - \varepsilon)U_{(k+1)}$, and hence

$$U_{(k+1)} \preceq \frac{1}{\lambda_{(k+1)} - 5\varepsilon} \sqrt{\lambda_{\max}(\boldsymbol{\Sigma}_{(k)})} \mathbf{I}, \qquad \boldsymbol{\Sigma}_{(k+1)} \preceq \left(1 + \frac{4\varepsilon}{\lambda_{(k+1)} - 5\varepsilon}\right) \sqrt{\lambda_{\max}(\boldsymbol{\Sigma}_{(k)})} \mathbf{I}.$$

Therefore, it holds that

$$\begin{split} \log \lambda_{\max}(\boldsymbol{\Sigma}_{(k+1)}) &\leq \frac{4\varepsilon}{\lambda_{(k+1)} - 5\varepsilon} + \frac{1}{2} \log \lambda_{\max}(\boldsymbol{\Sigma}_{(k)}) \\ &\leq \sum_{j=0}^{k} \frac{1}{2^{j}} \cdot \frac{4\varepsilon}{\lambda_{k+1-j} - 5\varepsilon} + \frac{\log \lambda_{\max}(\boldsymbol{\Sigma}_{(0)})}{2^{k+1}} \end{split}$$

Note that $\lambda_{\max}(\Sigma_{(0)}) \leq 1 + \varepsilon + \lambda_{(0)}$, and we also have

$$\frac{1}{2^{k+1}} + \sum_{j=0}^{k} \frac{1}{2^j} \cdot \frac{1}{k+1-j} \le \frac{1}{2^{k+1}} + \frac{2}{k+1} + \sum_{j=0}^{k} \frac{1}{2^j} \cdot \frac{j}{(k+1)(k+1-j)} \le \frac{6}{k+1}$$

Therefore, we have proven that as long as $\varepsilon \in [0, 0.1]$,

$$\log \lambda_{\max}(\mathbf{\Sigma}_{(k+1)}) \leq \frac{12}{k+1}.$$

This is the desired result.

C.3.3 Tighter rate with a different parameter schedule

In the following, we show that we can in fact choose $\lambda_{(k)} \approx \frac{\sqrt{d\sigma_{\alpha,\beta}}}{N}$ in Algorithm 1. This result is useful for getting the refined regret bound in Theorem 5.2 (cf. Appendix D.3).

Specifically, our analysis is based on the following concentration result. Its proof is essentially the same as the proof of Lemma C.2, except that we apply Lemma A.3.

Lemma C.4 Suppose that the sequence $\{(U_{(k)}, \tilde{H}_{(k)})\}$ is generated by Algorithm 1. For each $k = 0, 1, \dots, K-1$, we define

$$H_{(k)}^{\star} := \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k)}^{1/2} \phi \phi^{\top} U_{(k)}^{1/2}}{\|U_{(k)} \phi\|} \right]$$

Then, for any fixed parameter c > 1, with probability at least $1 - \delta$, the following inequality holds for all $k = 0, \dots, K - 1$:

$$c^{-1}H_{(k)}^{\star} - \varepsilon_{N}\mathbf{I} \preceq \widetilde{H}_{(k)} \preceq cH_{(k)}^{\star} + \varepsilon_{N}\mathbf{I},$$
(16)

where $\varepsilon_N = C_0 \left(\frac{\log(dK/\delta)}{(c-1)N} + \frac{\sigma_{\alpha,\beta}\sqrt{d+\log(K/\delta)}}{N} \right)$, and C_0 is a large absolute constant. In the following, we denote this event as \mathcal{E} and condition on \mathcal{E} .

Therefore, following the analysis from Proposition C.3, we prove the following result.

Proposition C.5 Let c > 1 be a constant. Suppose that Algorithm 1 is instantiated with $\lambda_{(k)} = \lambda = \frac{c^2+1}{c^2-1}\varepsilon_N$, where ε_N is defined in Lemma C.4. Then, under the event \mathcal{E} of Lemma C.4, for any $k \ge 1$, it holds that

$$\lambda_{\min}(\mathbf{\Sigma}_{(k)}) \ge c^{-4} \exp\left(-rac{\log(1/arepsilon_N)}{2^k}
ight), \qquad \lambda_{\max}(\mathbf{\Sigma}_{(k)}) \le c^4 \exp\left(rac{\lambda_{(0)} + arepsilon_N}{2^k}
ight).$$

In particular, for c = 1.1, $\varepsilon_N \leq 0.1$, $K \geq \max \{ \log \log(1/\varepsilon_N), 10 \}$, we have

$$\frac{1}{2}\mathbf{I} \preceq \mathbb{E}_{\phi \sim p} \left[\frac{U_{(K)} \phi \phi^{\top} U_{(K)}}{\|U_{(K)} \phi\|} \right] + \lambda U_{(K)} \preceq 2\mathbf{I}.$$

Proof of Proposition C.5. In the following proof, we abbreviate $\varepsilon := \varepsilon_N$. Recall that we can simplify the iterations in Algorithm 1 as follows: $U_{(0)} = \mathbf{I}$, and for $k = 0, 1, \dots, K - 1$:

$$\boldsymbol{\Sigma}_{(k)} = U_{(k)}^{1/2} \widetilde{H}_{(k)} U_{(k)}^{1/2} + \lambda_{(k)} U_{(k)}, \qquad U_{(k+1)} = \operatorname{sym}(\boldsymbol{\Sigma}_{(k)}^{-1/2} U_{(k)}),$$

and we regard $\widetilde{H}_{\scriptscriptstyle (K)}=H^\star_{\scriptscriptstyle (K)}.$

Then, for each $k \ge 1$, there exists an orthogonal matrix $V_{(k+1)}$ such that $U_{(k+1)} = V_{(k+1)} \Sigma_{(k)}^{-1/2} U_{(k)}$, and hence

$$\begin{split} \mathbf{I} &= V_{(k+1)} V_{(k+1)}^{\top} = V_{(k+1)} \boldsymbol{\Sigma}_{(k)}^{-1/2} \Big(U_{(k)}^{1/2} \widetilde{H}_{(k)} U_{(k)}^{1/2} + \lambda_{(k)} U_{(k)} \Big) \boldsymbol{\Sigma}_{(k)}^{-1/2} V_{(k+1)}^{\top} \\ &= U_{(k+1)} U_{(k)}^{-1/2} \big(\widetilde{H}_{(k)} + \lambda_{(k)} \mathbf{I} \big) U_{(k)}^{-1/2} U_{(k+1)}. \end{split}$$

Using Eq. (16), we have

$$\mathbf{I} \preceq U_{(k+1)} U_{(k)}^{-1/2} (cH_{(k)}^{\star} + (\lambda_{(k)} + \varepsilon_N) \mathbf{I}) U_{(k)}^{-1/2} U_{(k+1)}$$

= $c \cdot \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right] + (\lambda_{(k)} + \varepsilon) U_{(k+1)} U_{(k)}^{-1} U_{(k+1)}$

Using Eq. (16) again, we can bound

$$\begin{split} \boldsymbol{\Sigma}_{(k+1)} &= U_{(k+1)}^{1/2} \widetilde{H}_{(k+1)} U_{(k+1)}^{1/2} + \lambda_{(k+1)} U_{(k+1)} \\ &\succeq c^{-1} U_{(k+1)}^{1/2} H_{(k+1)}^{*} U_{(k+1)}^{1/2} + (\lambda_{(k+1)} - \varepsilon) U_{(k+1)} \\ &= c^{-1} \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k+1)} \phi\|} \right] + (\lambda_{(k+1)} - \varepsilon) U_{(k+1)} \\ &\succeq c^{-1} \sqrt{\lambda_{\min}(\boldsymbol{\Sigma}_{(k)})} \cdot \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right] + (\lambda_{(k+1)} - \varepsilon) U_{(k+1)} \\ &\succeq c^{-2} \sqrt{\lambda_{\min}(\boldsymbol{\Sigma}_{(k)})} \left(\mathbf{I} - (\lambda_{(k)} + \varepsilon) U_{(k+1)} U_{(k)}^{-1} U_{(k+1)} \right) + (\lambda_{(k+1)} - \varepsilon) U_{(k+1)} \\ &\succeq c^{-2} \sqrt{\lambda_{\min}(\boldsymbol{\Sigma}_{(k)})}, \end{split}$$

where the second inequality follows from $\|U_{(k+1)}\phi\| = \|\mathbf{\Sigma}_{(k)}^{-1/2}U_{(k)}\phi\| \le \sqrt{\lambda_{\max}(\mathbf{\Sigma}_{(k)}^{-1/2})} \|U_{(k)}\phi\| = \frac{1}{\sqrt{\lambda_{\min}(\mathbf{\Sigma}_{(k)})}} \|U_{(k)}\phi\|$, and the last inequality uses Eq. (15) and the fact that $\lambda_{(k+1)} - \varepsilon \ge c^{-2}(\lambda_{(k)} + \varepsilon)$. Similarly, we have

miniarly, we have

$$\begin{split} \boldsymbol{\Sigma}_{(k+1)} &\preceq c U_{(k+1)}^{1/2} H_{(k+1)}^{\star} U_{(k+1)}^{1/2} + (\lambda_{(k+1)} + \varepsilon) U_{(k+1)} \\ &= c \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k+1)} \phi\|} \right] + (\lambda_{(k+1)} + \varepsilon) U_{(k+1)} \\ &\preceq c \sqrt{\lambda_{\max}(\boldsymbol{\Sigma}_{(k)})} \cdot \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k+1)} \phi \phi^{\top} U_{(k+1)}}{\|U_{(k)} \phi\|} \right] + (\lambda_{(k+1)} + \varepsilon) U_{(k+1)} \end{split}$$

$$\leq c^2 \sqrt{\lambda_{\max}(\boldsymbol{\Sigma}_{(k)})} \left(\mathbf{I} - (\lambda_{(k)} - \varepsilon) U_{(k+1)} U_{(k)}^{-1} U_{(k+1)} \right) + (\lambda_{(k+1)} + \varepsilon) U_{(k+1)} }$$

$$\leq c^2 \sqrt{\lambda_{\max}(\boldsymbol{\Sigma}_{(k)})} \mathbf{I}.$$

where the last inequality uses Eq. (15) and the fact that $\lambda_{(k+1)} + \varepsilon \leq c^2 (\lambda_{(k)} - \varepsilon)$. Therefore, we have shown

$$c^{-2}\sqrt{\lambda_{\min}(\mathbf{\Sigma}_{(k)})} \leq \lambda_{\min}(\mathbf{\Sigma}_{(k+1)}) \leq \lambda_{\max}(\mathbf{\Sigma}_{(k+1)}) \leq c^2\sqrt{\lambda_{\max}(\mathbf{\Sigma}_{(k)})}.$$

Using this inequality recursively, we then have

$$\lambda_{\min}(\boldsymbol{\Sigma}_{(k)}) \ge c^{-4} \lambda_{\min}(\boldsymbol{\Sigma}_{(0)})^{\frac{1}{2^{k}}}, \qquad \lambda_{\max}(\boldsymbol{\Sigma}_{(k)}) \le c^{4} \lambda_{\max}(\boldsymbol{\Sigma}_{(0)})^{\frac{1}{2^{k}}}$$

The desired conclusion follows by recalling that we regard $H_{(K)} = H_{(K)}^{\star}$ and hence

$$\boldsymbol{\Sigma}_{(K)} = \mathbb{E}_{\phi \sim p} \left[\frac{U_{(K)} \phi \phi^{\top} U_{(K)}}{\|U_{(K)} \phi\|} \right] + \lambda U_{(K)}.$$

C.4 Details of Algorithm 2

In the following, we present the detailed description of the subroutine JDP_Batch_SGD (Algorithm 7).

Algorithm 7 Subroutine JDP_Batch_SGD: Batched SGD under JDP

Input: Loss \mathcal{L}_U under normalization (U, λ) , dataset $\mathcal{D} = \{(\phi_t, y_t)\}_{t \in [T]}$. **Input:** Epoch $K \ge 1$, batch size $N = \lfloor \frac{T}{K} \rfloor$, stepsize $\eta = \frac{1}{4L_{\nu}}$. Initialize $w_{(0)} = \mathbf{0}$.

for $k = 0, 1, \dots, K - 1$ do

Compute gradient estimate

$$g_{(k)} = \lambda \mu_{\nu} \cdot U w_{(k)} + \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \frac{U\phi_t}{\|U\phi_t\|} (\nu(\langle U\phi_t, w_{(k)}\rangle) - y_t).$$

Privatize $\widetilde{g}_{(k)} \sim \mathsf{Priv}_{2/N}(g_{(k)})$ and update

$$w_{(k+1)} = \operatorname{Proj}_{\mathcal{W}} (w_{(k)} - \eta \widetilde{g}_{(k)}).$$

Output: Estimator $\hat{\theta} = Uw_{(K)}$.

By the standard analysis of stochastic approximation, we provide the following guarantee of Algorithm 7.

Proposition C.6 (Convergence of batched SGD; JDP) Suppose that the input normalization (U, λ) satisfies Eq. (2), and $\lambda \geq \frac{1}{T}$. Then Algorithm JDP_Batch_SGD preserves (α, β) -JDP achieves with probability at least $1 - \delta$

$$\|w_{(k)} - \widehat{w}_{U}^{\star}\|^{2} \leq \left(1 - \frac{1}{8\kappa_{\nu}}\right)^{k} \|\widehat{w}_{U}^{\star}\|^{2} + O\left(\frac{\log(K/\delta)}{\mu_{\nu}^{2}N}\right) + O\left(\frac{1}{\mu_{\nu}^{2}} + \frac{d}{\mu_{\nu}L_{\nu}}\right) \cdot \frac{\sigma_{\alpha,\beta}^{2}\log(K/\delta)}{N^{2}},$$

where we recall that $\widehat{w}_U^{\star} = \operatorname{arg\,min}_{w \in \mathcal{W}} \mathcal{L}_U(w)$. In particular, when we take $K = 16\kappa_{\nu}\log(T)$, we have

$$L_{\nu} \cdot \|w_{(\kappa)} - \widehat{w}_{U}^{\star}\| \leq C_{1} \left[\kappa_{\nu}^{3/2} \sqrt{\frac{\log(T) \log(\log(T)/\delta)}{T}} + \sigma_{\alpha,\beta} (\kappa_{\nu}^{3/2} + \kappa_{\nu} \sqrt{d}) \frac{\log(T) \sqrt{\log(\log(T)/\delta)}}{T} \right] =: \underline{\epsilon}(T, \delta).$$
⁽¹⁷⁾

Algorithm 9 Subroutine LDP Learning Normalization

$$\begin{split} & \text{Input: Round } T \geq 1, \text{ parameter } \delta \in (0, 1). \\ & \text{Input: Epoch } K \geq 1, \text{ batch size } N = \lfloor \frac{T}{K} \rfloor. \\ & \text{Initialize } U_{(0)} = \mathbf{I}. \\ & \text{for } k = 0, \cdots, K-1 \text{ do} \\ & \text{for } t = kN+1, \cdots, (k+1)N \text{ do} \\ & \text{Observe } \phi_t \text{ and compute } V_t = \frac{U_{(k)}^{1/2}\phi_t\phi_t^\top U_{(k)}^{1/2}}{\|U_{(k)}\phi_t\|}. \\ & \text{Privatize } \widetilde{V}_t \sim \text{SymPriv}_1(V_t). \\ & \text{Compute } \widetilde{H}_{(k)} = \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \widetilde{V}_t. \\ & \text{Update} \\ & \boldsymbol{\Sigma}_{(k)} = U_{(k)}^{1/2} \widetilde{H}_{(k)} U_{(k)}^{1/2} + \lambda_{(k)} U_{(k)}, \qquad U_{(k+1)} = \text{sym}(\boldsymbol{\Sigma}_{(k)}^{-1/2} U_{(k)}). \end{split}$$

Output: Normalization matrix $U = U_{(K)}$.

Proof of Theorem 4.4. Let the subroutine JDP_Learning_Normalization be instantiated as in Proposition 4.3. By combining Proposition 4.3 and Proposition C.6, we know that with probability at least $1 - 2\delta$, normalization (U, λ) satisfies Eq. (2), and the estimator $\hat{w} = \hat{w}_{(K)}$ satisfies $L_{\nu} \| w_{(K)} - \hat{w}_{U}^{\star} \| \leq \underline{\epsilon}$. Applying Lemma 4.1 gives Theorem 4.4 (2).

C.5 Details of the LDP regression algorithm

In this section, we describe the details of LDP_L1_Regression (Algorithm 8), a LDP analogue of Algorithm 2.

Algorithm 8 LDP_L1_Regression Input: Round $T \ge 1$, parameter $\delta \in (0, 1)$. 1: Run the subroutine LDP_Learning_Normalization $(T/2, \delta)$ for the first T/2 rounds and receive (U, λ) .

2: Run the subroutine LDP_Batch_SGD($\mathcal{L}_U, T/2$) for the remaining T/2 rounds and receive \hat{w} and the error bound $\underline{\epsilon}$ (defined in Eq. (18)).

Output: Normalization (U, λ) , estimator $\hat{\theta} = U\hat{w}$, and overall error $\overline{\epsilon} = \lambda + \underline{\epsilon}$.

In the following, we specifying the details of the subroutine LDP Learning Normalization and LDP Batch SGD.

LDP learning normalization. By adapting Algorithm 1, we can derive a similar algorithm for learning normalization matrix under JDP.

Then, similar to Proposition 4.3, we have the following guarantee of Algorithm 9.

Proposition C.7 Let $T \ge 1, K \ge 1, \delta \in (0, 1)$, and $\varepsilon_N := C_0 \sigma_{\alpha, \beta} \sqrt{\frac{d + \log(K/\delta)}{N}}$, where C_0 is an absolute constant chosen according to Lemma C.10. Suppose that Algorithm 9 is instantiated with parameters $\lambda_{(k)} = (2k+5)\varepsilon_N$, and then with probability at least $1-\delta$,

$$\exp\left(-\frac{\log(1/\lambda_0)}{2^{k-1}}\right)\mathbf{I} \preceq \mathbb{E}_{\phi \sim p}\left[\frac{U_{(k)}\phi\phi^{\top}U_{(k)}}{\|U_{(k)}\phi\|}\right] + \lambda_{(k)}U_{(k)} \preceq \exp\left(\frac{12}{k}\right)\mathbf{I}.$$

In particular, as long as $K \ge \max \{ \log \log(N), 20 \}$, Algorithm 9 output (U, λ) satisfying Eq. (2) with probability at least $1 - \delta$, with

$$\lambda = (2K+5)\varepsilon_N = \tilde{O}(K) \cdot \sigma_{\alpha,\beta} \sqrt{\frac{d + \log(1/\delta)}{N}}$$

LDP batched SGD. Similarly, we use the following LDP batched SGD subroutine (Algorithm 10).

Algorithm 10 Subroutine LDP_Batch_SGD Input: Loss \mathcal{L}_U under normalization (U, λ) , round $T \ge 1$. Input: Epoch $K \ge 1$, batch size $N = \lfloor \frac{T}{K} \rfloor$, stepsize $\eta = \frac{1}{4L_{\nu}}$. for $k = 0, 1, \dots, K - 1$ do for $t = kN + 1, \dots, (k + 1)N$ do Observe (ϕ_t, y_t) and form gradient estimator $g_t = \frac{U\phi_t}{||U\phi_t||} (\nu(\langle U\phi_t, w_{(k)} \rangle) - y_t) + \lambda \mu_{\nu} \cdot Uw_{(k)}.$

Privatize $\tilde{g}_t \sim \mathsf{Priv}_2(g_t)$. Compute $\tilde{g}_{(k)} = \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \tilde{g}_t$ and update

$$w_{(k+1)} = \operatorname{Proj}_{\mathcal{W}} (w_{(k)} - \eta \widetilde{g}_{(k)}).$$

Output: $\widehat{w} = w_{(K)}$.

Proposition C.8 Suppose that the input normalization (U, λ) satisfies Eq. (2). Then subroutine LDP_Batch_SGD (Algorithm 10) preserves (α, β) -LDP and achieves with probability at least $1 - \delta$

$$\|w_{(k)} - \widehat{w}_{U}^{\star}\|^{2} \leq \left(1 - \frac{1}{8\kappa_{\nu}}\right)^{k} \|\widehat{w}_{U}^{\star}\|^{2} + O\left(\frac{1}{\mu_{\nu}^{2}} + \frac{d}{\mu_{\nu}L_{\nu}}\right) \cdot \frac{\sigma_{\alpha,\beta}^{2}\log(K/\delta)}{N}$$

In particular, when $\lambda \geq \frac{1}{T}$ and we take $K = 16\kappa_{\nu}\log T$, the output of Algorithm 10 $\hat{w} = w_{(K)}$ satisfies

$$L_{\nu} \cdot \|\widehat{w} - \widehat{w}_{U}^{\star}\| \le C_{1} \sigma_{\alpha,\beta} (\kappa_{\nu}^{3/2} + \kappa_{\nu} \sqrt{d}) \sqrt{\frac{\log(T) \log(\log(T)/\delta)}{T}} =: \underline{\epsilon}(T, \delta).$$
(18)

Guarantees of LDP_L1_Regression. By combining Proposition C.7 and Proposition C.8, we have the following result, as claimed in Theorem 4.5.

Theorem C.9 Let $T \ge 1, \delta \in (0, 1)$, and the subroutines of Algorithm 8 instantiated as in Proposition C.7 and Proposition C.8. Then, Algorithm 8 preserves (α, β) -LDP, and the following holds with probability at least $1 - 2\delta$:

(1) The normalization (U, λ) satisfies Eq. (2), and the estimator \hat{w} satisfies $L_{\nu} \| \hat{w} - \hat{w}_{U}^{\star} \| \leq \underline{\epsilon}$, where

$$\lambda = \lambda(T, \delta) = \tilde{O}\left(\sigma_{\alpha,\beta}\sqrt{\frac{d\log(1/\delta)}{T}}\right),$$
$$\underline{\boldsymbol{\epsilon}} = \underline{\boldsymbol{\epsilon}}(T, \delta) = \tilde{O}\left(\sigma_{\alpha,\beta}(\kappa_{\nu}^{3/2} + \kappa_{\nu}\sqrt{d})\sqrt{\frac{\log(1/\delta)}{T}}\right).$$

are defined in Proposition C.7 and Eq. (18), and $\tilde{O}(\cdot)$ hides polynomial factors of $\log(T)$. The overall error is defined as $\overline{\epsilon}(T, \delta) := 4L_{\nu}\lambda(T, \delta) + \underline{\epsilon}(T, \delta)$.

(2) By Lemma 4.1, the estimator $\hat{\theta} = U\hat{w}$ satisfies

$$\mathbb{E}_{\phi \sim p} |\nu(\langle \phi, \hat{\theta} \rangle) - \nu(\langle \phi, \theta^* \rangle)| \le L_{\nu} \cdot \mathbb{E}_{\phi \sim p} |\langle \phi, \hat{\theta} - \theta^* \rangle| \le 2\sqrt{d}\overline{\epsilon},$$

and for all $\phi \in \mathbb{R}^d$, we have the confidence bound $|\nu(\langle \phi, \hat{\theta} \rangle) - \nu(\langle \phi, \theta^* \rangle)| \le ||U\phi|| \overline{\epsilon}$.

C.5.1 Proof of Proposition C.7

The following lemma can be proven by Lemma A.1 and Lemma A.4, similar to Lemma C.2.

Lemma C.10 In Algorithm 9, with probability at least $1 - \delta$, the following inequality holds for all $k = 0, \dots, K - 1$:

$$\left\| \widetilde{H}_{(k)} - \mathbb{E}_{\phi \sim p} \left[\frac{U_{(k)}^{1/2} \phi \phi^{\top} U_{(k)}^{1/2}}{\| U_{(k)} \phi \|} \right] \right\|_{\text{op}} \le C_0 \sigma_{\alpha,\beta} \sqrt{\frac{d + \log(K/\delta)}{N}} =: \varepsilon_N,$$
(19)

where C is a large absolute constant. In the following, we denote this event as \mathcal{E} .

The proof of Proposition C.7 is then completed by combining Lemma C.10 and Proposition C.3. \Box

C.6 Proof of Proposition C.6 and Proposition C.8

In the following, we present the analysis of Algorithm 7 and Algorithm 10. We first state the following standard convergence result of batched SGD on a strongly convex function.

Proposition C.11 Suppose that $W \subseteq \mathbb{R}^d$ is a closed, convex domain, and F is a smooth convex function over W such that $\mu \mathbf{I} \preceq \nabla^2 F(w) \preceq L\mathbf{I}$ for all $w \in W$. Consider the following iteration of approximate gradient descent:

$$w_{(k+1)} = \operatorname{Proj}_{\mathcal{W}} (w_{(k)} - \eta (\nabla F(w_{(k)}) + Z_{(k)})), \tag{20}$$

where $Z_{(k)}$ is the noise vector at step k that is σ -sub-Gaussian (conditional on $w_{(1:k)}$), and the stepsize $\eta \in (0, \frac{1}{2L}]$. Then with probability at least $1 - \delta$, it holds that for all $k \in [K]$,

$$\|w_{(k)} - w^{\star}\|^{2} \leq \left(1 - \frac{\eta\mu}{2}\right)^{k} \|w_{(0)} - w^{\star}\|^{2} + \frac{4\eta}{\mu} \max_{j < k} \|Z_{(j)}\|^{2} + O\left(\frac{\sigma^{2}\log(K/\delta)}{\mu^{2}}\right),$$

where $w^* := \arg\min_{w \in \mathcal{W}} F(w)$.

Proof of Proposition C.6. For the *k*th epoch of Algorithm 7, we denote

$$g_t = \frac{U\phi_t}{\|U\phi_t\|} (\nu(\langle U\phi_t, w_{(k)}\rangle) - y_t).$$

Then, it is clear that $\mathbb{E}[g_t|w_{(k)}] = \nabla \mathcal{L}_U(w_{(k)}) - \lambda \mu_{\nu} \cdot Uw_{(k)}$. Therefore, Algorithm JDP_Batch_SGD (Algorithm 7) is an instantiation of Eq. (20) with $F = \mathcal{L}_U$, $w_{(0)} = 0$, $\mu = \frac{\mu_{\nu}}{2}$, $L = 2L_{\nu}$, $\eta = \frac{1}{2L}$.

As we have $||g_t|| \leq 2$, it is clear that Algorithm 7 preserves (α, β) -JDP. Furthermore, we can decompose

$$Z_{(k)} = g_{(k)} - \nabla \mathcal{L}_U(w_{(k)}) = G_{(k)} + \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} (g_t - \mathbb{E}[g_t|w_{(k)}]).$$

where $G_{(k)} \in \mathbb{R}^{d \times d}$ has i.i.d entries drawn from $\mathsf{N}(0, \frac{16\sigma_{\alpha,\beta}^2}{N^2})$. Therefore, using Hoeffding's bound, $Z_{(k)}$ is σ -sub-Gaussian with $\sigma^2 \leq O\left(\frac{1}{N} + \frac{\sigma_{\alpha,\beta}^2}{N^2}\right)$. Using Lemma A.1 and Lemma A.4, we also have $\max_k ||Z_{(k)}||^2 \leq O\left(\frac{1}{N} + \frac{d\sigma_{\alpha,\beta}^2}{N^2}\right) \log(K/\delta)$ with probability at least $1 - \delta$. Applying Proposition C.11 gives the desired upper bounds.

Proof of Proposition C.8. Similarly, for the *k*th epoch of Algorithm 7, it is clear that $||g_t - \lambda \mu_{\nu} \cdot Uw_{(k)}|| \le 2$ and $\mathbb{E}[g_t|w_{(k)}] = \nabla \mathcal{L}_U(w_{(k)})$. Therefore, Algorithm 7 preserves (α, β) -LDP, and it is also an instantiation of Eq. (20) with $F = \mathcal{L}_U$, $w_{(0)} = 0$, $\mu = \frac{\mu_{\nu}}{2}$, $L = 2L_{\nu}$, $\eta = \frac{1}{2L}$.

To proceed, we decompose

$$Z_{(k)} = g_{(k)} - \nabla \mathcal{L}_U(w_{(k)}) = \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} \left[(g_t - \mathbb{E}[g_t|w_{(k)}]) + G_t \right],$$

where $(G_t \in \mathbb{R}^{d \times d})_{t=kN+1,\dots,(k+1)N}$ are i.i.d Gaussian random vectors, and each G_t has i.i.d entries drawn from N(0, $16\sigma_{\alpha,\beta}^2$). Therefore, we denote $G_{(k)} = \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} G_t$, and $G_{(k)}$ has i.i.d entries drawn from N(0, $\frac{16\sigma_{\alpha,\beta}^2}{N}$). Therefore, using Hoeffding's bound, $Z_{(k)}$ is σ -sub-Gaussian with $\sigma^2 \leq O\left(\frac{\sigma_{\alpha,\beta}^2}{N}\right)$. Using Lemma A.1 and Lemma A.4, we also have $\max_k ||Z_{(k)}||^2 \leq O\left(\frac{d\sigma_{\alpha,\beta}^2 \log(K/\delta)}{N}\right)$ with probability at least $1 - \delta$. Applying Proposition C.11 gives the desired upper bounds.

C.6.1 Proof of Proposition C.11

By definition, we have $w^{\star} = \operatorname{Proj}_{\mathcal{W}}(w^{\star} - \eta \nabla F(w^{\star}))$, and hence

$$\begin{aligned} &\|w_{(k+1)} - w^{\star}\|^{2} \\ &= \|\operatorname{Proj}_{\mathcal{W}}(w_{(k)} - \eta(\nabla F(w_{(k)}) + Z_{(k)})) - \operatorname{Proj}_{\mathcal{W}}(w^{\star} - \eta\nabla F(w^{\star}))\|^{2} \\ &\leq \|(w_{(k)} - \eta(\nabla F(w_{(k)}) + Z_{(k)}) - w^{\star}) - (w^{\star} - \nabla F(w^{\star}))\|^{2} \\ &= \|w_{(k)} - w^{\star}\|^{2} - 2\eta \langle \nabla F(w_{(k)}) - \nabla F(w^{\star}) + Z_{(k)}, w_{(k)} - w^{\star} \rangle + \eta^{2} \|\nabla F(w_{(k)}) - \nabla F(w^{\star}) + Z_{(k)}\|^{2}. \end{aligned}$$

Thus, using the fact that F is μ -strongly-convex and L-smooth, we have

$$\mu \|w - w^{\star}\|^{2} \le \langle \nabla F(w) - \nabla F(w^{\star}), w - w^{\star} \rangle \le \frac{1}{L} \|\nabla F(w) - \nabla F(w^{\star})\|^{2}.$$

Hence, it holds that

$$\begin{aligned} \|w_{(k+1)} - w^{\star}\|^{2} \\ &\leq \|w_{(k)} - w^{\star}\|^{2} - 2\eta \left\langle \nabla F(w_{(k)}) - \nabla F(w^{\star}) + Z_{(k)}, w_{(k)} - w^{\star} \right\rangle + 2\eta^{2} \left\| \nabla F(w_{(k)}) - \nabla F(w^{\star}) \right\|^{2} + 2\eta^{2} \left\| Z_{(k)} \right\|^{2} \\ &\leq \|w_{(k)} - w^{\star}\|^{2} - 2\eta (1 - \eta L) \left\langle \nabla F(w_{(k)}) - \nabla F(w^{\star}), w_{(k)} - w^{\star} \right\rangle - 2\eta \left\langle Z_{(k)}, w_{(k)} - w^{\star} \right\rangle + 2\eta^{2} \left\| Z_{(k)} \right\|^{2} \\ &\leq (1 - \eta \mu) \left\| w_{(k)} - w^{\star} \right\|^{2} - 2\eta \left\langle Z_{(k)}, w_{(k)} - w^{\star} \right\rangle + 2\eta^{2} \left\| Z_{(k)} \right\|^{2}, \end{aligned}$$

where we use $\eta \leq \frac{1}{2L}$.

In the following, we condition on the following event:

$$\mathcal{E} := \{ \forall k \in [K] : \langle Z_{(k)}, w_{(k)} - w^* \rangle \le Z \| w_{(k)} - w^* \| \},\$$

where we denote $Z := C\sigma \sqrt{\log(K/\delta)}$, C is a large universal constant. By sub-Gaussian concentration, we know $\mathbb{P}(\mathcal{E}) \ge 1 - \delta$.

Under \mathcal{E} , we can bound

$$\begin{aligned} \|w_{(k+1)} - w^{\star}\|^{2} &\leq (1 - \eta\mu) \|w_{(k)} - w^{\star}\|^{2} + 2\eta Z \|w_{(k)} - w^{\star}\| + 2\eta^{2} \|Z_{(k)}\|^{2} \\ &\leq \left(1 - \frac{\eta\mu}{2}\right) \|w_{(k)} - w^{\star}\|^{2} + 2\frac{\eta}{\mu} Z^{2} + 2\eta^{2} \|Z_{(k)}\|^{2}, \qquad \forall k \in [K], \end{aligned}$$

where the last step is by Cauchy-Schwarz. Then, we know for any $k \in [K]$,

$$\|w_{(k)} - w^{\star}\|^{2} \leq \left(1 - \frac{\eta\mu}{2}\right)^{k} \|w_{(0)} - w^{\star}\|^{2} + \frac{4}{\mu^{2}}Z^{2} + \frac{4\eta}{\mu} \max_{j < k} \|Z_{(j)}\|^{2}$$

This is the desired result.

Algorithm 11 Meta Batch Elimination Algorithm

Input: Round $T \ge 1$, epoch schedule $1 = T_0 < T_1 < T_2 < \cdots < T_J = T$. Input: Subroutine Regression_with_Confidence. Initialize $\hat{f}^{(0)} \equiv 0, \hat{b}^{(0)} \equiv 1$. for $j = 0, 1, \cdots, J - 1$ do Set $\pi^{(j)} \leftarrow$ Confidence_based_Planning $(\{(\hat{f}^{(\tau)}, \hat{b}^{(\tau)})\}_{\tau < j})$. Initialize the subroutine Regression_with_Confidence^(j) with round $N^{(j)} = T^{(j+1)} - T^{(j)}$. for $t = T_j, \cdots, T_{j+1} - 1$ do Receive context x_t , take action $a_t \sim \pi^{(j)}(x_t)$, and receive reward r_t . Feed $(\phi(x_t, a_t), r_t)$ into Regression_with_Confidence^(j). Receive estimation $(\hat{f}^{(j)}, \hat{b}^{(j)})$ from Regression_with_Confidence^(j).

D Proofs from Section 5

D.1 Spanner

In Confidence_based_Planning (Algorithm 4), we utilize the following notion of the *spanner* of a set of actions.

Definition 7 Given a context $x \in \mathcal{X}$ and a set \mathcal{A}_1 of actions, a subset $\mathcal{A}' \subseteq \mathcal{A}_1$ is a spanner of \mathcal{A}_1 if for any $a \in \mathcal{A}$, there exists weights $(\gamma_{a'} \in [-1,1])_{a' \in \mathcal{A}'}$ such that

$$\phi(x,a) = \sum_{a' \in \mathcal{A}'} \gamma_{a'} \phi(x,a').$$

It is well-known that a spanner with size bounded by the dimension exists, known as the *barycentric spanner* [Awerbuch and Kleinberg, 2008].

Lemma D.1 (Barycentric spanner) For any context $x \in \mathcal{X}$ and set \mathcal{A}_1 , there exists a spanner of size

$$\dim(\mathcal{A}_1, x) := \dim(\{a \in \mathcal{A}_1 : \phi(x, a)\}).$$

An approximate barycentric spanner can be computed in time $poly(d, |\mathcal{A}_1|)$. Further, given a linear optimization oracle over the set $\{a \in \mathcal{A}_1 : \phi(x, a)\}$, the time complexity can further be reduced to poly(d) [Hazan and Karnin, 2016, Perchet et al., 2016].

D.2 Proof of Theorem 5.1

To provide a unified analysis framework for Algorithm 3 with different private regression subroutines, we first present a general action elimiation algorithm (Algorithm 11) that additionally takes an L_1 -regression subroutine Regression_with_Confidence as input.

Similar to our argument in Appendix C.2, we can show that Algorithm 11 preserves (α, β) -JDP $((\alpha, \beta)$ -LDP) if the subroutine Regression_with_Confidence preserves (α, β) -JDP $((\alpha, \beta)$ -LDP). Furthermore, we state the regret guarantee of Algorithm 11 under the following assumption on the subroutine Regression with Confidence.

Assumption 2 For each *j*, the subroutine Regression_with_Confidence returns $(\hat{f}^{(j)}, \hat{b}^{(j)})$ such that the following holds with probability at least $1 - \delta$.

(1) The function $\hat{b}^{(j)}(x, a)$ provides a valid confidence bound:

$$\mathbb{P}_{x \sim \mu}\Big(\forall a \in \mathcal{A}, \left|\hat{f}^{(\tau)}(x,a) - f^{\star}(x,a)\right| \leq \hat{b}^{(\tau)}(x,a)\Big) \geq 1 - \delta_0.$$

(2) The function $\hat{b}^{(j)}(x,a) = b^{(j)}(\phi(x,a))$ is given by a norm function $b^{(j)}$ over \mathbb{R}^d .

Theorem D.2 (Meta regret guarantee) Under Assumption 2, Algorithm 11 ensures that

$$\mathbf{Reg} \le \mathbb{E}\left[\sum_{j=0}^{J-2} 4d_{\mathcal{A}} \cdot N^{(j+1)} \mathbb{E}^{\pi^{(j)}} \left[\hat{b}^{(j)}(x,a) \right] \right] + 2N^{(0)} + 2TJ\delta + 2T^2\delta_0,$$

where $N^{(j)} = T^{(j+1)} - T^{(j)}$ is the batch size of the *j*th epoch. Further, if the subroutine Regression_with_Confidence preserves (α, β) -JDP (or correspondingly (α, β) -LDP), then Algorithm 11 preserves (α, β) -JDP (or correspondingly (α, β) -LDP).

With Theorem D.2, we can prove Theorem 5.1 easily from the guarantee of subroutine JDP_L1_Regression (Theorem 4.4).

Proof of Theorem 5.1. To see how Algorithm 11 generalizes Algorithm 3, we consider instantiate it with the subroutine Regression_with_Confidence be specified by JDP_L1_Regression (Algorithm 2), and for the output $(U^{(j)}, \lambda^{(j)}), \hat{\theta}^{(j)}, \overline{\epsilon}^{(j)}$ of the instance JDP_L1_Regression^(j), we set

$$\hat{f}^{(j)}(x,a) = \nu(\langle \phi(x,a), \hat{\theta}^{(j)} \rangle), \qquad \hat{b}^{(j)}(x,a) = \overline{\epsilon}^{(j)} \left\| U^{(j)}\phi(x,a) \right\|, \qquad \forall (x,a) \in \mathcal{X} \times \mathcal{A},$$

where $\overline{\epsilon}^{(j)} = \overline{\epsilon}(N^{(j)}, \delta)$ is defined in Theorem 4.4. Then, it is clear that under these specifications, Algorithm 11 agrees with Algorithm 3.

Further, we have $\mathbb{E}^{\pi^{(j)}}\left[\hat{b}^{(j)}(x,a)\right] \leq 2d\overline{\epsilon}^{(j)}$, and Assumption 2 holds with $\delta_0 = 0$, because under the success event of Theorem 4.4, we have

$$\left|\hat{f}^{(j)}(x,a) - f^{\star}(x,a)\right| \le \hat{b}^{(j)}(x,a), \qquad \forall x \in \mathcal{X}, a \in \mathcal{A}.$$

Then, Theorem D.2 yields (with $\delta = \frac{1}{T}$)

$$\begin{aligned} \mathbf{Reg} &\lesssim d_{\mathcal{A}} d \sum_{j=0}^{J-2} N^{(j+1)} \overline{\epsilon}^{(j)} + N^{(0)} \\ &\leq \operatorname{poly}(\log T) \cdot d_{\mathcal{A}} d \sum_{j=0}^{J-2} \left(\kappa_{\nu}^{3/2} \frac{N^{(j+1)}}{\sqrt{N^{(j)}}} + \sigma_{\alpha,\beta} (\kappa_{\nu}^{3/2} + \kappa_{\nu} \sqrt{d}) \frac{N^{(j+1)}}{N^{(j)}} \right) + N^{(0)}. \end{aligned}$$

In particular, under the choice $T^{(j)} = 2^j$, we have

$$\operatorname{\mathbf{Reg}} \leq \tilde{O}\left(d_{\mathcal{A}}d\kappa_{\nu}^{3/2}\sqrt{T} + \sigma_{\alpha,\beta}d_{\mathcal{A}}d(\kappa_{\nu}^{3/2} + \kappa_{\nu}\sqrt{d})\right).$$

D.2.1 Proof of Theorem D.2

For any policy $\pi : \mathcal{X} \to \Delta(\mathcal{A})$, we define its sub-optimality as

$$\mathsf{Reg}(\pi) = \mathbb{E}_{x \sim P, a \sim \pi(x)} [f^{\star}(x, \pi^{\star}(x)) - f^{\star}(x, a)],$$

where we recall that $\pi^{\star}(x) := \arg \max_{a \in \mathcal{A}} f^{\star}(x, a)$ is the optimal policy under f^{\star} . Then, by definition, for Algorithm 11,

$$\mathbf{Reg} = \mathbb{E}\left[\sum_{j=0}^{J-1} \sum_{t=T^{(j)}}^{T^{(j+1)}-1} \operatorname{Reg}(\pi^{(j)})\right] = \mathbb{E}\left[\sum_{j=0}^{J-1} N^{(j)} \cdot \operatorname{Reg}(\pi^{(j)})\right].$$

In the following, we work with the following quantity:

$$\operatorname{\mathbf{Reg}}^{+} := \sum_{j=0}^{J-1} N^{(j)} \cdot \operatorname{\mathsf{Reg}}(\pi^{(j)}),$$

which is a random variable measuring the cumulative sub-optimality of the algorithm.

We assume the success event of Assumption 2 and define

$$\mathcal{X}^{(j)} := \left\{ x \in \mathcal{X} : \forall \tau \le j, a \in \mathcal{A}, \left| \hat{f}^{(\tau)}(x, a) - f^{\star}(x, a) \right| \le \hat{b}^{(\tau)}(x, a) \right\}.$$

Then, for any $x \in \mathcal{X}^{(j)}$ and $\tau \leq j$, we have

$$\hat{f}^{(\tau)}(x,\pi^{\star}(x)) + \hat{b}^{(\tau)}(x,\pi^{\star}(x)) \ge f^{\star}(x,\pi^{\star}(x)) = \max_{a\in\mathcal{A}} f^{\star}(x,a) \ge \max_{a\in\mathcal{A}} \hat{f}^{(\tau)}(x,a) - \hat{b}^{(\tau)}(x,a),$$

and hence $\pi^{\star}(x) \in \mathcal{A}^{(j)}(x)$. Further, for any $a \in \mathcal{A}^{(j)}(x)$,

$$\hat{f}^{(j)}(x,a) + \hat{b}^{(j)}(x,a) \ge \hat{f}^{(j)}(x,\pi^{\star}(x)) - \hat{b}^{(j)}(x,\pi^{\star}(x)),$$

and hence

$$\begin{aligned} f^{\star}(x,\pi^{\star}(x)) - f^{\star}(x,a) &\leq \hat{f}^{(j)}(x,\pi^{\star}(x)) + \hat{b}^{(j)}(x,\pi^{\star}(x)) - \hat{f}^{(j)}(x,a) + \hat{b}^{(j)}(x,a) \\ &\leq 2\hat{b}^{(j)}(x,\pi^{\star}(x)) + 2\hat{b}^{(j)}(x,a) \\ &\leq 4 \max_{a' \in \mathcal{A}^{(j)}(x)} \hat{b}^{(j)}(x,a'), \qquad \forall x \in \mathcal{X}^{(j)}, a \in \mathcal{A}^{(j)}(x). \end{aligned}$$

Note that $\pi^{(j+1)}(x)$ is supported on $\mathcal{A}^{(j)}(x)$, and hence

$$\operatorname{Reg}(\pi^{(j+1)}) = \mathbb{E}_{x \sim P, a \sim \pi^{(j+1)}(x)}[f^{\star}(x, \pi^{\star}(x)) - f^{\star}(x, a)] \le 4\mathbb{E}_{x \sim P} \max_{a \in \mathcal{A}^{(j)}(x)} \hat{b}^{(j)}(x, a)$$

Further, $\mathcal{A}_{sp}^{(j)}(x)$ is a barycentric spanner of $\mathcal{A}^{(j-1)}(x)$, and hence for any $a \in \mathcal{A}^{(j)}(x) \subseteq \mathcal{A}^{(j-1)}(x)$, there exists parameters $(\gamma_{x,a,a'} \in [-1,1])_{a' \in \mathcal{A}_{sp}^{(j)}(x)}$, such that

$$\phi(x,a) = \sum_{a' \in \mathcal{A}_{\rm sp}^{(j)}(x)} \gamma_{x,a,a'} \phi(x,a')$$

Hence, by Assumption 2 (2), for any $x \in \mathcal{X}^{(j)}$, $a \in \mathcal{A}^{(j)}(x)$,

$$\hat{b}^{(j)}(x,a) = b^{(j)} \left(\sum_{\substack{a' \in \mathcal{A}_{sp}^{(j)}(x) \\ a' \in \mathcal{A}_{sp}^{(j)}(x)}} \gamma_{x,a,a'} \phi(x,a') \right) \le \sum_{\substack{a' \in \mathcal{A}_{sp}^{(j)}(x) \\ a' \in \mathcal{A}_{sp}^{(j)}(x)}} |\gamma_{x,a,a'}| b^{(j)}(\phi(x,a')) = |\mathcal{A}_{sp}^{(j)}(x)| \cdot \mathbb{E}_{a' \sim \pi^{(j)}(x)} \hat{b}^{(j)}(x,a').$$

Therefore, for any $x \in \mathcal{X}^{(j)}$,

$$\max_{a \in \mathcal{A}^{(j)}(x)} \hat{b}^{(j)}(x,a) \le d_{\mathcal{A}} \cdot \mathbb{E}_{a' \sim \pi^{(j)}(x)} \hat{b}^{(j)}(x,a'),$$

and thus,

$$\operatorname{\mathsf{Reg}}(\pi^{(j+1)}) \le 4d_{\mathcal{A}} \cdot \mathbb{E}_{x \sim P, a \sim \pi^{(j)}(x)} \hat{b}^{(j)}(x, a) + 2P(x \notin \mathcal{X}^{(j)}).$$

Algorithm 12 Subroutine JDP_Reweighted_Linear_Regression

Input: Dataset $\mathcal{D} = \overline{\{(\phi_t, y_t)\}_{t \in [T]}}$ with size T = 2N, parameter $\delta \in (0, 1), \delta_0 \in (0, 1)$.

- 1: Split $\mathcal{D} = \mathcal{D}_0 \cup \mathcal{D}_1$ equally.
- 2: Set $(U, \lambda) \leftarrow \mathsf{JDP_Learning_Normalization}(\mathcal{D}_0, \delta/2)$
- 3: Compute the following estimates on \mathcal{D}_1 :

$$\xi = \frac{1}{N} \sum_{t=N+1}^{T} \frac{U\phi_t}{\|U\phi_t\|} y_t, \qquad \Xi = \frac{1}{N} \sum_{t=N+1}^{T} \frac{U\phi_t}{\|U\phi_t\|} \phi_t^{\scriptscriptstyle \top} + \lambda \mathbf{I}, \qquad W = \frac{1}{N} \sum_{t=N+1}^{T} \frac{U\phi_t\phi_t^{\scriptscriptstyle \top}U}{\|U\phi_t\|^2} + \lambda \mathbf{I}.$$

4: Privatize $[\tilde{\xi}; \tilde{\Xi}] \sim \operatorname{Priv}_{3/N}([\xi; \Xi])$ and $\widetilde{W} \sim \operatorname{SymPriv}_{3/N}(W)$. **Output:** Estimators $\hat{\theta} = \widetilde{\Xi}^{-1}\widetilde{\xi}$ and confidence bound

$$\hat{b}(\phi) = 16\lambda \|U\phi\| + \underline{\epsilon} \|\widetilde{\Xi}^{-\top}\phi\|_{\widetilde{W}}.$$

By Assumption 2, $P(x \notin \mathcal{X}^{(j)}) \leq T\delta_0$, and hence taking summation over $j = 0, 1, \dots, J-2$ gives

$$\sum_{j=1}^{J-1} N^{(j)} \cdot \mathsf{Reg}(\pi^{(j)}) \le 4d_{\mathcal{A}} \sum_{j=0}^{J-2} N^{(j+1)} \cdot \mathbb{E}^{\pi^{(j)}} \Big[\hat{b}^{(j)}(x,a) \Big] + 2T^2 \delta_0.$$

Note that the above inequality holds under the success event of Assumption 2, which holds with probability at least $1 - J\delta$. Then taking expectation gives the desired upper bound on **Reg**.

We also remark that a high-probability upper bound on the regret follows similarly (with an extra step of applying martingale concentration).

D.3 Proof of Theorem 5.2

In this section, we present an adaption of Algorithm 3 based on the subroutine JDP_Reweighted_Linear_Regression (Algorithm 12), which provides a tighter rate of convergence.

The subroutine JDP_Reweighted_Linear_Regression utilizes the linear structure of the linear models. Instead of running a batched SGD procedure, it directly computes an estimator of θ^* by solving a privatized linear equation $\Xi \hat{\theta} = \tilde{\xi}$. It is clear that Algorithm 12 preserves (α, β) -JDP, and we also show that the confidence bound $|\langle \phi, \hat{\theta} - \theta^* \rangle| \leq \hat{b}(\phi)$ holds true in a *distributional* sense.

Proposition D.3 Let $T \ge 1$, $\delta \in (0, 1)$, P' be a fixed distribution over $\mathbb{B}^d(1)$, and the subroutine JDP_Learning_Normalization of Algorithm 12 be instantiated as in Proposition C.5. Then Algorithm 12 preserves (α, β) -JDP and ensures the following holds with probability at least $1 - \delta$:

(1) The normalization (U, λ) satisfies Eq. (2), with λ given by

$$\lambda := \lambda(T, \delta) = C \frac{\sigma_{\alpha, \beta} \sqrt{d + \log(K/\delta)}}{T},$$
(21)

where C is a large absolute constant.

(2) With $\underline{\epsilon} := \underline{\epsilon}(T, \delta) = C_0 \sqrt{\frac{\log(1/(\delta \delta_0)}{T}}$ for a large absolute constant C_0 , it holds that

$$\mathbb{P}_{\phi' \sim P'} \left(|\langle \phi', \hat{\theta} - \theta^* \rangle| \ge \hat{b}(\phi') \right) \le \delta_0$$

(3) It holds that $\mathbb{E}_{\phi \sim p} \left[\hat{b}(\phi) \right] \leq 16\sqrt{d} \cdot \underline{\epsilon} + 32d(\lambda + \underline{\epsilon}^2).$

Theorem 5.2 then follows from the general guarantee of Theorem D.2.

Proof of Theorem 5.2. We instantiate the subroutine Regression_with_Confidence to be JDP_Reweighted_Linear_Regression in Algorithm 11, with parameter $\delta_0 = \frac{1}{\delta T^2|\mathcal{A}|}$. We also let P' be the distribution of $\phi(x, a)$ under $x \sim P$, $a \sim \text{Unif}(\mathcal{A})$.

Then, for the *j*th epoch, we let $\lambda^{(j)} := \lambda(N^{(j)}, \delta), \lambda_0^{(j)} := \lambda_0(N^{(j)}, \delta)$. Proposition D.3 guarantees that with probability at least $1 - \delta$,

$$\begin{aligned} \mathbb{P}_{x\sim P}\Big(\exists a\in\mathcal{A}, \left|\langle\phi(x,a),\hat{\theta}^{(j)}-\theta^{\star}\rangle\right|\geq \hat{b}^{(j)}(\phi(x,a))\Big)\\ \leq |\mathcal{A}|\cdot\mathbb{P}_{\phi'\sim P'}\Big(\left|\langle\phi',\hat{\theta}-\theta^{\star}\rangle\right|\geq \hat{b}^{(j)}(\phi')\Big)\leq |\mathcal{A}|\delta_{0}\leq\frac{\delta}{T},\end{aligned}$$

and we also have

$$\mathbb{E}^{\pi^{(j)}} \left[\hat{b}^{(j)}(x,a) \right] \lesssim \sqrt{d} \underline{\epsilon}^{(j)} + d(\lambda^{(j)} + (\underline{\epsilon}^{(j)})^2) \\ \lesssim \sqrt{\frac{d \log(1/\delta_0)}{N^{(j)}}} + \frac{\sigma_{\alpha,\beta} d^{3/2} \sqrt{\log(N^{(j)}/\delta)} + d \log(1/\delta_0)}{N^{(j)}}.$$

Therefore, Assumption 2 holds, and Theorem D.2 yields

$$\begin{split} \mathbf{Reg} &\lesssim d_{\mathcal{A}} \sqrt{d \log(1/\delta_0)} \sum_{j=0}^{J-2} \frac{N^{(j+1)}}{\sqrt{N^{(j)}}} + d_{\mathcal{A}} \Big(\sigma_{\alpha,\beta} d^{3/2} \sqrt{\log(T/\delta)} + d \log(1/\delta_0) \Big) \sum_{j=0}^{J-2} \frac{N^{(j+1)}}{N^{(j)}} \\ &+ N^{(0)} + T J \delta. \end{split}$$

In particular, with the choice $T^{(j)} = 2^{j+1}$ and $\delta = \frac{1}{T}$, we have

$$\operatorname{\mathbf{Reg}} \leq \tilde{O}\Big(d_{\mathcal{A}}\sqrt{dT\log|\mathcal{A}|} + d_{\mathcal{A}}(\sigma_{\alpha,\beta}d^{3/2} + d\log|\mathcal{A}|)\Big),$$

where $\tilde{O}(\cdot)$ hides poly(log T) factors.

D.3.1 Proof of Proposition D.3

By Proposition C.5, the subroutine JDP_Learning_Normalization can be suitably instantiated so that the output (U, λ) satisfies Eq. (2) with probability at least $1 - \frac{\delta}{2}$, with epoch size $K = \max \{ \log \log(T), 10 \}$, and λ given by Eq. (21).

In the following, we proceed to prove (2). By definition, we have

$$\widetilde{\xi} = \xi + Z_{\xi}, \qquad \widetilde{\Xi} = \Xi + Z_{\Xi}, \qquad \widetilde{W} = W + Z_{W_{\xi}}$$

where entries of $Z = [Z_{\xi}; Z_{\Xi}; Z_W]$ are independent zero-mean Gaussian random variable with variance $\frac{36\sigma_{\alpha,\beta}^2}{N^2}$. Further, we also have

$$y_t = \langle \phi_t, \theta^* \rangle + \zeta_t, \qquad \mathbb{E}[\zeta_t | \phi_t] = 0,$$

and conditional on the random variables $U, (\phi_t)_{t \in [N+1,T]}$ and Z, the random variables $(\zeta_t)_{t \in [N+1,T]}$ are independent.

In particular, the following event \mathcal{E}_1 holds with probability at least $1 - \frac{\delta}{6}$:

$$\mathcal{E}_1: \quad \max\left\{ \left\| Z_{\xi} \right\|, \left\| Z_{\Xi} \right\|_{\mathrm{op}}, \left\| Z_W \right\|_{\mathrm{op}} \right\} \leq \frac{\lambda}{16}.$$

Further, by Lemma A.3, the following event \mathcal{E}_2 holds with probability at least $1 - \frac{\delta}{6}$:

$$\mathcal{E}_{2}: \quad \frac{1}{N} \sum_{t=N+1}^{T} \frac{U^{1/2} \phi_{t} \phi_{t}^{\top} U^{1/2}}{\|U \phi_{t}\|} + \frac{1}{2} \lambda \mathbf{I} \succeq \frac{1}{2} \mathbb{E}_{\phi \sim p} \frac{U^{1/2} \phi \phi^{\top} U^{1/2}}{\|U \phi\|}.$$

c	
L	_

In the following, we condition on $\mathcal{E}_1 \cap \mathcal{E}_2$.

Under \mathcal{E}_2 , we have

$$D := \frac{1}{N} \sum_{t=N+1}^{T} \frac{U \phi_t \phi_t^{\mathsf{T}} U}{\|U \phi_t\|} + \lambda U \succeq \frac{1}{4} \mathbf{I}.$$

Note that $\Xi = DU^{-1}$. For any vector $v \in \mathbb{R}^d$, we can bound

$$\left\| U \widetilde{\Xi}^{\top} v \right\| = \left\| (D + U Z_{\Xi}) v \right\| \ge \left\| D v \right\| - \left\| U Z_{\Xi} v \right\| \ge \frac{1}{4} \left\| v \right\| - \left\| U \right\|_{\text{op}} \left\| Z_{\Xi} \right\|_{\text{op}} \left\| v \right\| \ge \frac{1}{8} \left\| v \right\|,$$

where we use $\|U\|_{\text{op}} \leq \frac{2}{\lambda}$ and $\|Z_{\Xi}\|_{\text{op}} \leq \frac{\lambda}{16}$. Therefore, we have $\|\widetilde{\Xi}^{-\top}U^{-1}\|_{\text{op}} \leq 8$.

Now, we decompose

$$\widetilde{\xi} = \frac{1}{N} \sum_{t=N+1}^{T} \frac{U\phi_t}{\|U\phi_t\|} \phi_t^{\mathsf{T}} \theta^\star + \frac{1}{N} \sum_{t=N+1}^{T} \frac{U\phi_t}{\|U\phi_t\|} \zeta_t + Z_{\xi}, \qquad \widetilde{\Xi} = \frac{1}{N} \sum_{t=N+1}^{T} \frac{U\phi_t}{\|U\phi_t\|} \phi_t^{\mathsf{T}} + \lambda \mathbf{I} + Z_{\Xi}.$$

Hence, we can re-write

$$\widetilde{\xi} = \left(\widetilde{\Xi} - Z_{\Xi} - \lambda \mathbf{I}\right) \theta^* + \frac{1}{N} \sum_{t=N+1}^T \frac{U\phi_t}{\|U\phi_t\|} \zeta_t + Z_{\xi},$$

and we denote $e = \frac{1}{N} \sum_{t=N+1}^{T} \frac{U\phi_t}{\|U\phi_t\|} \zeta_t$. Note that under \mathcal{E}_1 , we have $\|\widetilde{\xi} - \widetilde{\Xi}\theta^* - e\| \leq 2\lambda$, and hence for any vector $\phi \in \mathbb{R}^d$,

$$\left|\langle\phi,\hat{\theta}-\theta^{\star}\rangle\right| \leq \left|\langle\widetilde{\Xi}^{-\top}\phi,\widetilde{\xi}-\widetilde{\Xi}\theta^{\star}-e\rangle\right| + \left|\langle\phi,\Xi^{-1}e\rangle\right| \leq 16\lambda \left\|U\phi\right\| + \left|\langle\phi,\Xi^{-1}e\rangle\right|$$

where we also use $\left\|\widetilde{\Xi}^{-\top}U^{-1}\right\|_{\text{op}} \leq 8.$

It remains to upper bound $|\langle \phi', \Xi^{-1}e \rangle|$ under the fixed distribution $\phi' \sim P'$. The following lemma asserts e is a sub-Gaussian random vector, conditional on $\phi_{N+1}, \cdots, \phi_T$. The proof is a direct corollary of Hoeffding's bound and is deferred to the end of this section.

Lemma D.4 There is an absolute constant $c_0 > 0$ such that for any $v \in \mathbb{B}^d(1)$,

$$\mathbb{E}_{\zeta}\left[\exp\left(c_0 N \frac{\langle v, e \rangle^2}{\|v\|_{W_0}^2}\right) \middle| U, \phi_{N+1}, \cdots, \phi_T, Z\right] \le 2,$$

where $W_0 = \frac{1}{N} \sum_{t=N+1}^{T} \frac{U\phi_t \phi_t^\top U}{\|U\phi_t\|^2}$.

In particular, taking expectation over $v = \widetilde{\Xi}^{-\top} \phi'$, with $\phi' \sim P$, we have

$$\mathbb{E}_{\phi'\sim P}\mathbb{E}_{\zeta}\left[\exp\left(c_0N\frac{\langle v,\widetilde{\Xi}^{-1}e\rangle^2}{\left\|\widetilde{\Xi}^{-\tau}\phi'\right\|_{W_0}^2}\right)\right] \leq 2.$$

By Markov's inequality, the following event \mathcal{E}_3 holds with probability at least $1 - \delta$:

$$\mathcal{E}_3: \quad \mathbb{E}_{\phi' \sim P}\left[\exp\left(c_0 N \frac{\langle \phi', \widetilde{\Xi}^{-1} e \rangle^2}{\left\|\widetilde{\Xi}^{-\top} \phi'\right\|_{W_0}^2}\right)\right] \leq \frac{6}{\delta}.$$

Note that under \mathcal{E}_3 , using Markov's inequality again, we have

$$\mathbb{P}_{\phi'\sim P'}\left(\frac{|\langle \phi', \widetilde{\Xi}^{-1}e\rangle|}{\left\|\widetilde{\Xi}^{-\top}\phi'\right\|_{W_0}} \ge \sqrt{\frac{\log(6/(\delta\delta_0))}{c_0 N}}\right) \le \delta_0.$$

Note that under \mathcal{E}_1 , we have $\widetilde{W} = W + Z_W = W_0 + \lambda \mathbf{I} + Z_W \succeq W_0$. Therefore, we can conclude that

$$\mathbb{P}_{\phi'\sim P'}\left(|\langle \phi', \hat{\theta} - \theta^{\star}\rangle| \ge 16\lambda \, \|U\phi'\| + \sqrt{\frac{\log(6/(\delta\delta_0))}{c_0 N}} \|\widetilde{\Xi}^{-\tau}\phi'\|_{\widetilde{W}}\right) \le \delta_0.$$

It remains to prove (3). by Cauchy inequality,

$$\begin{split} \left(\mathbb{E}_{\phi \sim p} \left\| \widetilde{\Xi}^{-\top} \phi \right\|_{\widetilde{W}} \right)^2 &\leq \mathbb{E}_{\phi \sim p} \left\| U \phi \right\| \cdot \mathbb{E}_{\phi \sim p} \left[\frac{\left\| \widetilde{\Xi}^{-\top} \phi \right\|_{\widetilde{W}}^2}{\left\| U \phi \right\|} \right] \\ &\leq 2d \cdot \left\langle \widetilde{W}, \mathbb{E}_{\phi \sim p} \frac{\widetilde{\Xi}^{-\top} \phi \phi^{\top} \widetilde{\Xi}^{-1}}{\left\| U \phi \right\|} \right\rangle \\ &\leq 4d \left\| \widetilde{\Xi}^{-\top} U^{-1} \right\|_{\text{op}}^2 \operatorname{tr}(\widetilde{W}), \end{split}$$

where we use the fact that $\mathbb{E}_{\phi \sim p} \left[\frac{U \phi \phi^{\top} U}{\|U \phi\|} \right] \preceq 2\mathbf{I}$. Using $\operatorname{tr}(\widetilde{W}) = \operatorname{tr}(W_0) + \lambda d + \operatorname{tr}(Z_W) \leq 1 + 2\lambda d$, we have $\mathbb{E}_{\phi \sim p} \left\| \widetilde{\Xi}^{-\top} \phi \right\|_{\widetilde{W}} \leq 16\sqrt{d(1+2\lambda d)}$. Combining the inequalities above completes the proof. \Box

Proof of Lemma D.4. In the following, we condition on $(U, \phi_{N+1}, \dots, \phi_T, Z)$. By Hoeffding's bound, for any $\lambda \in \mathbb{R}$,

$$\mathbb{E}_{\zeta}[\exp\left(\lambda\langle v, Ne\rangle\right)] = \mathbb{E}_{\zeta}\left[\exp\left(\sum_{t=N+1}^{T}\lambda\langle v, \frac{U\phi_t}{\|U\phi_t\|}\rangle\zeta_t\right)\right]$$
$$\leq \exp\left(\frac{\lambda^2}{2}\sum_{t=N+1}^{T}\langle v, \frac{U\phi_t}{\|U\phi_t\|}\rangle^2\right)$$
$$= \exp\left(\frac{N\lambda^2}{2}\|v\|_{W_0}^2\right).$$

Therefore, $\langle v, Ne \rangle$ is a σ_v -sub-Gaussian random variable with $\sigma_v^2 \leq N \|v\|_{W_0}^2$. Hence, there is an absolute constant $c_0 > 0$ such that for any $v \in \mathbb{R}^d$,

$$\mathbb{E}_{\zeta}\left[\exp\left(c_0 N \frac{\langle v, e \rangle^2}{\|v\|_{W_0}^2}\right)\right] \le 2.$$

D.4 Proof of Theorem 5.3

In Algorithm 11, we instantiate the subroutine Regression_with_Confidence with LDP_L1_Regression (Algorithm 8). For the *j*th epoch, we let $(U^{(j)}, \lambda^{(j)}), \hat{\theta}^{(j)}, \overline{\epsilon}^{(j)}$ be the output of the subroutine LDP_L1_Regression, and we let

$$\hat{f}^{(j)}(x,a) = \nu(\langle \phi(x,a), \hat{\theta}^{(j)} \rangle), \qquad \hat{b}^{(j)}(x,a) = \overline{\epsilon}^{(j)} \cdot \left\| U^{(j)}\phi(x,a) \right\|, \qquad \forall (x,a) \in \mathcal{X} \times \mathcal{A}$$

following Algorithm 3. Recall that for the subroutine LDP_L1_Regression, we have

$$\overline{\boldsymbol{\epsilon}}^{(j)} := \overline{\boldsymbol{\epsilon}}(N^{(j)}, \delta) = \tilde{O}\left(\sigma_{\alpha,\beta}(\kappa_{\nu}^{3/2} + \kappa_{\nu}\sqrt{d})\sqrt{\frac{\log(1/\delta)}{T}}\right).$$

which is defined in Theorem C.9.

Note that Assumption 2 holds with $\delta_0 = 0$ under the above specifications, and we also have $\mathbb{E}^{\pi^{(j)}} \left[\hat{b}^{(j)}(x,a) \right] \leq 2d\overline{\epsilon}^{(j)}$. Therefore, applying Theorem D.2 yields

$$\mathbf{Reg} \lesssim d_{\mathcal{A}} d \sum_{j=0}^{J-2} N^{(j+1)} \cdot \overline{\boldsymbol{\epsilon}}^{(j)} + N^{(0)} + T J \delta.$$

In particular, with the epoch schedule $T^{(j)} = 2^j$ and $\delta = \frac{1}{T}$, we have

$$\mathbf{Reg} \le \tilde{O}\left(d_{\mathcal{A}}d(\kappa_{\nu}^{3/2} + \kappa_{\nu}\sqrt{d})\sqrt{T}\right).$$

This is the desired upper bound.

E Proofs from Section 6

E.1 Proof of Theorem 6.2

Privacy guarantee. To make the presentation clear, we re-write the iteration of Algorithm 5 as follows. For any sub-dataset $\mathcal{D}_{sub} = \{(\phi_t, y_t)\}$, we denote

$$g(\theta; \mathcal{D}_{\mathsf{sub}}) := \frac{1}{|\mathcal{D}_{\mathsf{sub}}|} \sum_{(\phi, y) \in \mathcal{D}_{\mathsf{sub}}} \phi(\langle \phi, \theta \rangle - y),$$

$$F(\theta; \mathcal{D}_{\mathsf{sub}}) := \operatorname{Proj}_{\mathbb{B}^{d}(R)} (\theta - \eta g(\theta; \mathcal{D}_{\mathsf{sub}})).$$

Then, the iteration of Algorithm 5 can be re-written as follows: $\theta_{(0)} = 0$, and for $k = 0, 1, \dots, K-1$,

$$\theta_{(k+1)}(\mathcal{D}_{(0:k)}) := F(\theta_{(k)}(\mathcal{D}_{(0:k-1)}); \mathcal{D}_{(k)})).$$

Note that for any \mathcal{D}_{sub} ,

$$g(\theta; \mathcal{D}_{\mathsf{sub}}) = \nabla \left(\frac{1}{2|\mathcal{D}_{\mathsf{sub}}|} \sum_{(\phi, y) \in \mathcal{D}_{\mathsf{sub}}} (\langle \phi, \theta \rangle - y)^2 \right)$$

is the gradient of a 1-Lipschitz convex function, and hence $\theta \mapsto F(\theta; \mathcal{D})$ is a contraction under $\|\cdot\| = \|\cdot\|_2$:

$$\|\theta - \theta'\| \le \|F(\theta; \mathcal{D}) - F(\theta'; \mathcal{D})\|, \quad \forall \theta, \theta'.$$

Further note that, for neighbored sub-dataset $\mathcal{D}_{\mathsf{sub}}$ and $\mathcal{D}'_{\mathsf{sub}}$ of size N, we have

$$\|g(\theta; \mathcal{D}_{\mathsf{sub}}) - g(\theta; \mathcal{D}'_{\mathsf{sub}})\| \le \frac{2(R+1)}{N}, \quad \forall \|\theta\| \le R,$$

and hence

$$\|F(\theta; \mathcal{D}_{\mathsf{sub}}) - F(\theta; \mathcal{D}'_{\mathsf{sub}})\| \le \frac{2(R+1)\eta}{N}.$$

Therefore, for any neighbored dataset \mathcal{D} and \mathcal{D}' , using the inequalities above, we have

$$\|\theta_{(\kappa)}(\mathcal{D}) - \theta_{(\kappa)}(\mathcal{D}')\| \le \frac{2(R+1)\eta}{N}$$

This immediately show that Algorithm 5 preserves (α, β) -JDP by the privacy of Gaussian channels (Definition 2).

Convergence guarantee. We state the following convergence guarantee.

Proposition E.1 Let $K, N \ge 2, \delta \in (0, 1)$. We denote $B_{\delta} := 10(R+1)\sqrt{\frac{K \log(K/\delta)}{N}}$. Suppose that the parameters (η, R) are chosen so that

$$R \ge 1 + B_\delta \eta. \tag{22}$$

Then it holds that with probability at least $1 - \delta$

$$\left\|\hat{\theta} - \theta^{\star}\right\|_{\Sigma} \lesssim \frac{1}{\sqrt{\eta K}} + \sqrt{\frac{\eta \log K \log(1/\delta)}{N}} + \frac{\eta (R+1)\sigma_{\alpha,\beta}\sqrt{\log(1/\delta)}}{N}$$

In particular, we may choose $R = 2, \eta = 1$, and

$$K = \min\left\{c\sqrt{\frac{T}{\log T \log(1/\delta)}}, \left(\frac{T}{\sigma_{\alpha,\beta}\sqrt{\log(1/\delta)}}\right)^{2/3}\right\} \lor 1, \qquad N = \frac{T}{K},$$

where c > 0 is a small absolute constant so that Eq. (22) holds. Then, it holds that

$$\left\|\hat{\theta} - \theta^{\star}\right\|_{\Sigma} \lesssim \left(\frac{\log T \log(1/\delta)}{T}\right)^{1/4} + \left(\frac{\sigma_{\alpha,\beta}\sqrt{\log(1/\delta)}}{T}\right)^{1/3}.$$

This completes the proof of Theorem 6.2.

E.1.1 Proof of Proposition E.1

We denote

$$\bar{g}(\theta) := \mathbb{E}_{(\phi, y) \sim M^{\star}}[\phi(\langle \phi, \theta \rangle - y)] = \mathbf{\Sigma}(\theta - \theta^{\star}).$$

We also write

$$\operatorname{err}_{(k)} := g(\theta_{(k)}; \mathcal{D}_{(k)}) - \bar{g}(\theta_{(k)}),$$
$$E_{(k)} := \sum_{i=0}^{k} (\mathbf{I} - \eta \mathbf{\Sigma})^{k-i} \operatorname{err}_{(i)}.$$

We first invoke the following concentration result.

Lemma E.2 The following holds with probability at least $1 - \delta$: For all $k = 0, 1, \dots, K - 1$:

$$||E_{(k)}|| \le 5(R+1)\sqrt{\frac{K\log(K/\delta)}{N}} := B_{\delta},$$

and

$$||E_{(K)}||_{\Sigma} \le 10(R+1)\sqrt{\frac{(\eta^{-1}\log K+2)\log(2/\delta)}{N}} =:\varepsilon_N.$$

Proof of Lemma E.2. By definition,

$$\begin{split} N \cdot E_{\scriptscriptstyle (k)} &= N \sum_{i=0}^{k} (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k-i} \mathsf{err}_{\scriptscriptstyle (i)} \\ &= \sum_{i=0}^{k} \sum_{t=iN+1}^{(i+1)N} (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k-i} (g_t - \bar{g}(\boldsymbol{\theta}_{\scriptscriptstyle (i)})), \end{split}$$

where $g_t := \phi_t(\langle \phi_t, \theta_{(i)} \rangle - y_t)$ is the gradient at $t \in [iN+1, (i+1)N]$. Note that $\mathbb{E}[g_t|\theta_{(i)}] = \bar{g}(\theta_{(i)})$ for round t in ith epoch. Thus, we denote $Z_t := (\mathbf{I} - \eta \mathbf{\Sigma})^{k-i}(g_t - \bar{g}(\theta_{(i)}))$ for $t \in [iN+1, (i+1)N]$. Then, the sequence Z_t is a martingale difference sequence with respect to the filtration $\mathcal{F}_t = \sigma(\{(\phi_s, y_s)\}_{s \leq t})$. Further, note that $||Z_t|| \leq 2(R+1)$, and hence using Lemma A.4 with a union bound, we have with probability at least $1 - \delta$ for all $k \in [K]$,

$$N \|E_{(k)}\| = \left\| \sum_{t=1}^{(k+1)N} Z_t \right\| \le 2(R+1)\sqrt{NK} \cdot \left(1 + \sqrt{2\log(K/\delta)}\right).$$

Further, by Lemma A.7, we also have

$$\|\mathbf{\Sigma}^{1/2} Z_t\| \le 2(R+1) \|\mathbf{\Sigma}^{1/2} (\mathbf{I} - \eta \mathbf{\Sigma})^{k-i}\|_{\text{op}} \le 5(R+1) \min\left\{\frac{1}{\sqrt{\eta(k-i)}}, 1\right\}$$

Therefore, using Lemma A.4, we have with probability at least $1 - \delta$

$$N \|E_{(K-1)}\|_{\Sigma} = \left\| \sum_{t=1}^{KN} (\Sigma^{1/2} Z_t) \right\| \le 5(R+1) \sqrt{N \cdot \left(\frac{\log K}{\eta} + 2\right)} \cdot \left(1 + \sqrt{2\log(1/\delta)}\right).$$

Taking the union bound again and rescaling $\delta \leftarrow \frac{\delta}{2}$ completes the proof.

We denote \mathcal{E}_1 to be the success event of Lemma E.2. In the following, we work under \mathcal{E}_1 .

We inductively show that for all i < K:

$$\theta_{(i+1)} = \theta_{(i)} - \eta g_{(i)}.$$
(23)

The base case is trivial: Eq. (23) holds for all i < 0.

Now, we assume that for some k < K, Eq. (23) holds for all i < k. Then, we only need to prove Eq. (23) for the case i = k. We denote $\theta^+_{(k+1)} = \theta_{(k)} - \eta g_{(k)}$. Using $\bar{g}(\theta) = \Sigma(\theta - \theta^*)$ and Eq. (23) recursively for i < k, we know

$$\theta^+_{\scriptscriptstyle (k+1)} - \theta^{\star} = (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k+1} (\theta_{\scriptscriptstyle (0)} - \theta^{\star}) - \eta \sum_{i=0}^k (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k-i} \mathsf{err}_{\scriptscriptstyle (i)}$$

Therefore,

$$\left\|\theta_{(k+1)}^{+}\right\| \leq \left\|\left(\mathbf{I} - (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k+1}\right)\theta^{\star}\right\| + \eta \left\|E_{(k)}\right\| \leq 1 + \eta B_{\delta} \leq R.$$

Therefore, $\theta^+_{(k+1)} \in \mathbb{B}^d(R)$, and hence

$$\theta_{(k+1)} = \operatorname{Proj}_{\mathbb{B}^d(R)}(\theta_{(k+1)}^+) = \theta_{(k+1)}^+$$

This completes the proof of case i = k.

Therefore, by induction, Eq. (23) indeed holds for all j < K. In particular, we have

$$\begin{aligned} \boldsymbol{\theta}_{(K)} - \boldsymbol{\theta}^{\star} &= (\mathbf{I} - \eta \boldsymbol{\Sigma})^{K} (\boldsymbol{\theta}_{(0)} - \boldsymbol{\theta}^{\star}) - \eta \sum_{i=0}^{K-1} (\mathbf{I} - \eta \boldsymbol{\Sigma})^{K-1-i} \mathsf{err}_{(i)} \\ &= (\mathbf{I} - \eta \boldsymbol{\Sigma})^{K} (\boldsymbol{\theta}_{(0)} - \boldsymbol{\theta}^{\star}) - \eta E_{(K-1)}. \end{aligned}$$

Hence, it holds that (by Lemma A.7 and under \mathcal{E}_1):

$$\|\theta_{(K)} - \theta^{\star}\|_{\Sigma} \leq \left\| (\mathbf{I} - \eta \boldsymbol{\Sigma})^{K} \theta^{\star} \right\|_{\Sigma} + \eta \|E_{(K-1)}\|_{\Sigma} \lesssim \frac{1}{\sqrt{\eta K}} + \sqrt{\frac{\eta \log K \log(1/\delta)}{N}}$$

Finally, we know $\hat{\theta} = \theta_{(K)} + Z$, where $Z \sim \mathsf{N}(0, \frac{4(R+1)^2 \sigma_{\alpha,\beta}^2}{N^2})$ is a Gaussian random vector. Therefore, with probability at least $1 - \delta$, $\|Z\|_{\Sigma} \lesssim \frac{(R+1)\sigma_{\alpha,\beta}\sqrt{\log(1/\delta)}}{N}$. Combining the inequalities above and rescaling $\delta \leftarrow \frac{\delta}{3}$ complete the proof.

E.2 Proof of Theorem 6.3

We prove the following convergence rate of Algorithm 6, under general choice of (η, R) .

Proposition E.3 Let $K, N \ge 2, \delta \in (0, 1)$. We denote $B_{\delta} := 6(R+1)\sqrt{\frac{K \log(K/\delta)}{N}}$ and $\varepsilon_N = (R+1)\sqrt{\frac{K \log(K/\delta)}{N}}$. Suppose that the parameters (η, R) are chosen so that

$$R \ge 1 + \eta \cdot (B_{\delta} + 4\varepsilon_N). \tag{24}$$

Then it holds that with probability at least $1 - \delta$

$$\|\theta_{(K)} - \theta^{\star}\|_{\Sigma} \lesssim \frac{1}{\sqrt{\eta K}} + R\eta \sigma_{\alpha,\beta} \sqrt{\frac{K \log(K/\delta)}{N}}.$$

Proof of Theorem 6.3. For Algorithm 6, we choose $\eta = 1$, R = 2, and

$$K = c \left(\frac{T}{\sigma_{\alpha,\beta}^2 \log(T/\delta)} \right)^{1/3} \vee 1, \qquad N = \frac{T}{K},$$

where c > 0 is an absolute constant so that Eq. (24) holds. Then, by Proposition E.3, Algorithm 5 achieves

$$\|\theta_{\scriptscriptstyle (K)} - \theta^{\star}\|_{\Sigma} \lesssim \left(\frac{\sigma_{\alpha,\beta}^2 \log(T/\delta)}{T}\right)^{1/6}$$

This is the desired upper bound.

E.2.1 Proof of Proposition E.3

In Algorithm 6, for epoch $k = 0, 1, \dots, K - 1$, we have

$$\widetilde{g}_{\scriptscriptstyle (k)} = \zeta_{\scriptscriptstyle (k)} + \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} g_t, \qquad \theta_{\scriptscriptstyle (k+1)} = \theta_{\scriptscriptstyle (k)} - \eta \widetilde{g}_{\scriptscriptstyle (k)},$$

where $\{\zeta_{(0)}, \dots, \zeta_{(K-1)}\}$ are i.i.d samples from $\mathsf{N}(0, \sigma_N^2)$ with $\sigma_N = \frac{(R+1)\sigma_{\alpha,\beta}}{\sqrt{N}}$ and independent of the dataset $\{(\phi_t, y_t)\}_{t \in [T]}$.

To begin with, we denote $\mathbf{\Sigma} := \mathbb{E}_{\phi \sim p}[\phi \phi]$ (the covariance matrix),

$$\bar{g}_{(k)} := \mathbb{E}_{(x,y)\sim p}[x(\mathsf{clip}_R(\langle \theta_{(k)}, x \rangle) - y)],$$

and define the error vectors

$$\mathrm{err}_{0}^{(k)} \coloneqq \nabla \mathcal{L}_{\mathsf{Sq}}(\theta_{\scriptscriptstyle (k)}) - \bar{g}_{\scriptscriptstyle (k)} \qquad \mathrm{err}_{1}^{(k)} \coloneqq \bar{g}_{\scriptscriptstyle (k)} - \frac{1}{N} \sum_{t=kN+1}^{(k+1)N} g^{t}, \qquad \mathrm{err}_{2}^{(k)} = -\zeta_{\scriptscriptstyle (k)}$$

Then, we can decompose the error of the estimator $g_{\scriptscriptstyle (k)}$ as

$$\operatorname{err}^{(k)} := \nabla \mathcal{L}_{\mathsf{Sq}}(\theta_{(k)}) - g_{(k)} = \operatorname{err}_0^{(k)} + \operatorname{err}_1^{(k)} + \operatorname{err}_2^{(k)}.$$

Notice that by definition, we have

$$\mathcal{L}_{\mathsf{Sq}}(\theta_{(k)}) = \frac{1}{2} \left\| \theta_{(k)} - \theta^{\star} \right\|_{\Sigma}^{2}, \qquad \nabla \mathcal{L}_{\mathsf{Sq}}(\theta_{(k)}) = \Sigma(\theta_{(k)} - \theta^{\star}).$$

Therefore, recursively using $g_{(k)} = \nabla \mathcal{L}_{\mathsf{Sq}}(\theta_{(k)}) - \mathsf{err}^{(k)}$ and $\theta_{(k+1)} = \theta_{(k)} - \eta g_{(k)}$, we have

$$\theta_{(k)} - \theta^{\star} = (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k} (\theta_{(0)} - \theta^{\star}) + \eta \sum_{i=0}^{k-1} (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k-i-1} \operatorname{err}^{(i)}.$$
(25)

We bound the three types of error separately: For each $j \in \{0, 1, 2\}$, denote

$$E_j^{(k)} := \sum_{i=0}^{k-1} (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k-i-1} \operatorname{err}_j^{(i)}$$

Lemma E.4 With probability at least $1 - \delta$, the following holds: For all $k \in [K]$, it holds that

$$\left\|E_1^{(k)}\right\| \le 6(R+1)\sqrt{\frac{K\log(K/\delta)}{N}} =: B_{\delta}.$$

The proof of Lemma E.4 is essentially similar to that of Lemma E.2, and hence we omit it for succintness. Lemma E.5 Denote $\varepsilon_N := \sigma_N \sqrt{K \log(2K/\delta)}$. With probability at least $1 - \delta$, for all $k = 0, 1, \dots, K - 1$, it holds that

$$\mathbb{P}_{\phi \sim p}\Big(|\langle E_2^{(k)}, \phi \rangle| > 3\varepsilon_N\Big) \leq \frac{1}{K^6}, \qquad \mathbb{E}_{\phi \sim p}\langle E_2^{(k)}, \phi \rangle^2 \leq 4\varepsilon_N^2.$$

where C_2 is an absolute constant. We denote this success event at \mathcal{E}_2 .

Proof of Lemma E.5. Fix a $k \in [K]$. Then by definition

$$E_2^{(k)} = \sum_{i=0}^k (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k-i} \zeta_{(i)}$$

where $\zeta_{(i)} \sim \mathsf{N}(0, \sigma_N^2 \mathbf{I})$. Therefore, because $\zeta = (\zeta_{(0)}, \cdots, \zeta_{(K)})$ is a sequence of independent Gaussian random variables, we have

$$E_2^{(k)} \sim \mathsf{N}(0, C_k), \qquad C_k = \sigma_N^2 \sum_{i=0}^k (\mathbf{I} - \eta \boldsymbol{\Sigma})^{2(k-i)} \preceq K \sigma_N^2 \mathbf{I}.$$

Therefore, for any fixed $\phi \in \mathbb{B}^d(1)$, $\langle E_2^{(k)}, \phi \rangle \sim \mathsf{N}(0, \|\phi\|_{C_k}^2)$ is a zero-mean Gaussian random variable with variance $\|\phi\|_{C_k}^2 \leq K\sigma_N^2$. This immediately implies that

$$\forall \phi \in \mathbb{B}^d(1), \qquad \mathbb{E}_{\zeta}\left[\exp\left(\frac{|\langle E_2^{(k)}, \phi \rangle|^2}{4K\sigma_N^2}\right)\right] \le 2,$$

where the expectation is taken over the sequence $\zeta = (\zeta_{(0)}, \dots, \zeta_{(K)})$ of independent Gaussian random vectors. Therefore, we have

$$\mathbb{E}_{\zeta}\left[\mathbb{E}_{\phi\sim p}\exp\left(\frac{|\langle E_{2}^{(k)},\phi\rangle|^{2}}{4K\sigma_{N}^{2}}\right)\right] \leq 2, \quad \forall k.$$

Therefore, by Markov's inequality and taking the union bound, we have

$$\mathbb{P}_{\zeta}\left(\forall k \in [K] : \mathbb{E}_{\phi \sim p} \exp\left(\frac{|\langle E_2^{(k)}, \phi \rangle|^2}{4K\sigma_N^2}\right) \le \frac{2K}{\delta}\right) \ge 1 - \delta.$$

Let this event be \mathcal{E}_2 . Then, under \mathcal{E}_2 , using Markov inequality's again, we have

$$\mathbb{P}_{\phi \sim p}\Big(|\langle E_2^{(k)}, \phi \rangle| \ge 3\sigma_N \sqrt{K \log(K/\delta)}\Big) \le \frac{1}{K^6}, \qquad \forall k.$$

Similarly, under \mathcal{E}_2 , using Jensen's inequality, we also have

$$\mathbb{E}_{\phi \sim p} |\langle E_2^{(k)}, \phi \rangle|^2 \le 4\sigma_N^2 K \log(2K/\delta)$$

This is the desired result.

Lemma E.6 Under the success event of Lemma E.5 and assuming that $K \ge 2$ and

$$R \ge 1 + \eta \cdot (B_{\delta} + 4\varepsilon_N).$$

Then, we have for all $k = 0, 1, \dots, K - 1$:

$$\left\|\operatorname{err}_{0}^{(k)}\right\| \leq \frac{2\eta\varepsilon_{N}}{K^{3}}.$$
(26)

In particular, it holds that $\left\|E_0^{(k)}\right\| \leq \frac{2\eta\varepsilon_N}{K^2}$.

Proof of Lemma E.5. We prove by induction. The base case k = 0 is trivial because $err_0^{(0)} = 0$. Now, suppose that Eq. (26) holds for all $k' \le k$. In the following, we proceed to prove Eq. (26) for k + 1. By Eq. (25), we have

$$\theta_{(k+1)} = \left(\mathbf{I} - (\mathbf{I} - \eta \mathbf{\Sigma})^k \right) \theta^* + \eta \left(E_0^{(k)} + E_1^{(k)} + E_2^{(k)} \right),$$

and under $\mathcal{E}_1 \cap \mathcal{E}_2$, we have $\|\mathcal{E}_1\| \leq C_1 \varepsilon_N$,

$$\mathbb{P}_{\phi \sim p}\Big(|\langle E_2^{(k)}, \phi \rangle| > 3\varepsilon_N\Big) \le \frac{1}{K^6}.$$

By induction hypothesis,

$$\left\|E_0^{(k)}\right\| = \left\|\sum_{i=0}^k (\mathbf{I} - \eta \boldsymbol{\Sigma})^{k-i} \mathrm{err}_0^{(i)}\right\| \le \sum_{i=0}^k \left\|\mathrm{err}_0^{(i)}\right\| \le \frac{2\varepsilon_N}{K^2}.$$

Combining all the equations above, we have

$$\left|\langle\phi,\theta_{(k+1)}\rangle\right| \le 1 + \eta \left(\frac{2\eta\varepsilon_N}{K^2} + B_{\delta} + \left|\langle E_2^{(k)},\phi\rangle\right|\right).$$

By our assumption on R, we know that $|\langle \phi, \theta_{(k+1)} \rangle| - R \leq \eta \Big(|\langle E_2^{(k)}, \phi \rangle| - 3\varepsilon_N \Big)$, and hence

$$\mathbb{P}_{\phi \sim p}(|\langle \phi, \theta_{(k+1)} \rangle| \ge R) \le \mathbb{P}_{\phi \sim p}\Big(|\langle E_2^{(k)}, \phi \rangle| > 3\varepsilon_N\Big) \le \frac{1}{K^6}$$

Finally, by the definition of $err_0^{(k+1)}$, we have

$$\operatorname{err}_{0}^{(k+1)} = \nabla \mathcal{L}_{\mathsf{Sq}}(\theta_{(k+1)}) - \bar{g}_{(k+1)} = \mathbb{E}_{\phi \sim p}[\phi(\langle \theta_{(k+1)}, \phi \rangle - \operatorname{clip}_{R}(\langle \theta_{(k+1)}, \phi \rangle))].$$

Hence,

$$\begin{aligned} \left\| \mathsf{err}_{0}^{(k+1)} \right\| &\leq \mathbb{E}_{\phi \sim p} [\mathbf{1} \left\{ \left| \left\langle \theta_{(k+1)}, \phi \right\rangle \right| > R \right\} \cdot \left(\left| \left\langle \theta_{(k+1)}, \phi \right\rangle \right| - R \right) \right] \\ &\leq \sqrt{\mathbb{P}_{\phi \sim p} (\left| \left\langle \phi, \theta_{(k+1)} \right\rangle \right| \ge R) \cdot \mathbb{E}_{\phi \sim p} (\left| \left\langle \theta_{(k+1)}, \phi \right\rangle \right| - R) \right\}_{+}^{2}} \end{aligned}$$

Algorithm 13 SquareCB [Foster and Rakhlin, 2020, Simchi-Levi and Xu, 2020]

Input: Round $T \ge 1$, epoch schedule $1 = T_0 < T_1 < T_2 < \cdots < T_J = T$. **Input:** Oracle Regression, parameter $\delta \in (0, 1)$. Initialize $\hat{f}^{(0)} \equiv 0$. **for** $j = 0, 1, \cdots, J - 1$ **do** Initialize the subroutine Regression^(j) with round $N^{(j)} = T^{(j+1)} - T^{(j)}$ and confidence $\delta' = \frac{\delta}{2J^2}$. **for** $t = T_j, \cdots, T_{j+1} - 1$ **do** Receive context x_t . Let $\hat{a}_t := \arg \max_{a \in \mathcal{A}} \hat{f}^{(j)}(x_t, a)$, and set

$$p_t(a) := \begin{cases} \frac{1}{|\mathcal{A}| + \gamma^{(j)} \left(\hat{f}^{(j)}(x_t, a_t) - \hat{f}^{(j)}(x_t, a)\right)}, & a \neq \hat{a}_t, \\ 1 - \sum_{a \neq \hat{a}_t} p_t(a), & a = \hat{a}_t. \end{cases}$$

Take action $a_t \sim p_t$, and receive reward r_t . Feed $(\phi(x_t, a_t), r_t)$ into Regression^(j). Receive estimation $\hat{f}^{(j+1)}$ from Regression^(j).

$$\leq \sqrt{\frac{\eta^2}{K^6} \mathbb{E}_{\phi \sim p} \langle E_2^{(k)}, \phi \rangle^2} \leq \frac{2\eta \varepsilon_N}{K^3}.$$

This completes the proof of the step k + 1.

Now, we prove Proposition E.1 by combining the lemmas above. By definition,

$$\left\|\theta_{(K)} - \theta^{\star}\right\|_{\Sigma} \leq \left\|\left(\mathbf{I} - \eta \boldsymbol{\Sigma}\right)^{K} \theta^{\star}\right\| + \eta \left\|E_{0}^{(K)}\right\|_{\Sigma} + \eta \left\|E_{1}^{(K)}\right\|_{\Sigma} + \eta \left\|E_{2}^{(K)}\right\|_{\Sigma}.$$

Under the success event of Lemma E.4 and Lemma E.5, we have

$$\left\|E_1^{(K)}\right\|_{\Sigma} \le \left\|E_1^{(K)}\right\| \le B_{\delta} \lesssim \varepsilon_N, \qquad \left\|E_2^{(K)}\right\|_{\Sigma} \le 2\varepsilon_N,$$

and $\left\|E_0^{(K)}\right\| \leq \frac{2\eta\varepsilon_N}{K^2}$. Further, by Lemma A.7, we also have

$$\left\| (\mathbf{I} - \eta \boldsymbol{\Sigma})^{K} \boldsymbol{\theta}^{\star} \right\| \leq \sqrt{\frac{2e}{\eta K}}.$$

Combining the inequalities above gives

$$\|\theta_{(K)} - \theta^{\star}\|_{\Sigma} \lesssim \frac{1}{\sqrt{\eta K}} + \eta \varepsilon_N.$$

This is the desired upper bound.

E.3 Algorithm SquareCB

Assumption 3 For any policy $\pi : \mathcal{X} \to \Delta(\mathcal{A})$ and any linear reward function f^* , given N independent samples $\{(x_t, a_t, r_t)\}$ generated as

$$x_t \sim P, \quad a_t \sim \pi(x_t), \quad \mathbb{E}[r_t | x_t, a_t] = f^{\star}(x_t, a_t),$$

the regression oracle Regression (initialized with round N and confidence δ) returns an estimated mean function $\hat{f} : \mathcal{X} \times \mathcal{A} \to \mathbb{R}$ such that with probability at least $1 - \delta$,

$$\mathbb{E}_{x \sim P, a \sim \pi(x)} \left(\hat{f}(x, a) - f^{\star}(x, a) \right)^2 \leq \mathcal{E}_{\delta}(N)^2,$$

where \mathcal{E}_{δ} is a non-increasing function of N.

Theorem E.7 (Guarantee of SquareCB) Suppose that Assumption 3 holds. Then, with parameters

$$\gamma^{(0)} = 1, \qquad \gamma^{(j)} := \frac{\sqrt{|\mathcal{A}|}}{\mathcal{E}_{\delta'}(N^{(j-1)})}, \quad j = 1, \cdots, J-1,$$

SquareCB (Algorithm 13) achieves

$$\operatorname{\mathbf{Reg}} \le C\sqrt{|\mathcal{A}|} \sum_{j=0}^{J-1} \mathcal{E}_{\delta'}(N^{(j)}) \cdot N^{(j+1)} + N^{(0)} + T\delta.$$

Furthermore, the privacy guarantee of Algorithm 13 can be implied by the privacy guarantee of the regression oracle (similar to Algorithm 11).

Lemma E.8 If the oracle Regression preserves (α, β) -JDP (or correspondingly (α, β) -LDP), then Algorithm 13 preserves (α, β) -JDP (or correspondingly (α, β) -LDP)

Proof of Theorem 6.4 (1). For JDP learning, we consider instantiating the regression oracle Regression with the algorithm JDP_Improper_BatchSGD (Algorithm 5). For the output $\hat{\theta}$ of JDP_Improper_BatchSGD given a dataset of size N, we consider the estimated mean function $\hat{f}(x, a) := \langle \hat{\theta}, \phi(x, a) \rangle$ for all $(x, a) \in \mathcal{X} \times \mathcal{A}$. Then, by Theorem 6.2, Assumption 3 holds with

$$\mathcal{E}_{\delta}(N) \lesssim \left(\frac{\log N \log(1/\delta)}{N}\right)^{1/4} + \left(\frac{\sigma_{\alpha,\beta} \log(1/\delta)}{N}\right)^{1/3}$$

Therefore, we choose $T^{(j)} = 2^j$ and $\delta = \frac{1}{T}$, and Theorem E.7 provides the following regret bound:

$$\operatorname{\mathbf{Reg}} \le \sqrt{|\mathcal{A}|} \cdot \tilde{O} \Big(T^{3/4} + \sigma_{\alpha,\beta}^{1/3} T^{2/3} \Big)$$

This is the desired result.

Proof of Theorem 6.4 (2). Similarly, for LDP learning, we consider instantiating the regression oracle Regression with the algorithm LDP_Improper_BatchSGD (Algorithm 6). Then, by Theorem 6.3, Assumption 3 holds with

$$\mathcal{E}_{\delta}(N) \lesssim \left(\frac{\sigma_{\alpha,\beta}^2 \log(N/\delta)}{N}\right)^{1/6}$$

Therefore, we choose $T^{(j)} = 2^j$ and $\delta = \frac{1}{T}$, and Theorem E.7 provides the following regret bound:

$$\mathbf{Reg} \le \sqrt{|\mathcal{A}|} \cdot \tilde{O}\left(\sigma_{\alpha,\beta}^{1/3} T^{5/6}\right).$$

This is the desired regret bound.