

UNIRULEDNESS OF SOME MODULI SPACES OF POINTED SPIN CURVES

BOGDAN-PETRU CARASCA

ABSTRACT. The moduli space $\mathcal{S}_{g,2n}$ parametrizes pointed curves with spin structure. We prove that $\mathcal{S}_{g,2}$, $\mathcal{S}_{g,4}$ and $\mathcal{S}_{g,6}$ are uniruled for particular values of g .

1. INTRODUCTION

The moduli space $\mathcal{S}_{g,2n}$ of $2n$ -pointed spin curves of genus g parametrizes tuples $(X, x_1, \dots, x_{2n}, \eta)$ where (X, x_1, \dots, x_{2n}) is a $2n$ -pointed smooth curve of genus g and η is a line bundle on X such that $\eta^{\otimes 2} = \omega_X(-x_1 - \dots - x_{2n})$. The natural forgetful map $\pi_{g,2n} : \mathcal{S}_{g,2n} \rightarrow \mathcal{M}_{g,2n}$ is finite of degree 2^{2g} .

These moduli spaces are in close connection with the classical moduli spaces of spin curves. A spin curve of genus g is a pair $[X, \eta]$ where η is a line bundle on X such that $\eta^{\otimes 2} = \omega_X$, and the moduli space of spin curves of genus g is denoted by \mathcal{S}_g . We say that a spin curve $[X, \eta]$ is even (resp. odd) if $h^0(X, \eta)$ is even (resp. odd). In the simultaneous papers [Mum71] and [Ati71] it was shown that even and odd spin curves do not mix. This means that given a family of spin curves over a connected base, the parity is constant. As a consequence of this, \mathcal{S}_g splits into two connected components \mathcal{S}_g^+ and \mathcal{S}_g^- , the moduli spaces of even and odd spin curves respectively. We denote by $\overline{\mathcal{S}}_g^+$ and $\overline{\mathcal{S}}_g^-$ the Cornalba compactifications of these moduli spaces, cf. [Cor89].

Given an irreducible n -nodal curve $X = \tilde{X}/\{x_i \sim y_i\}_{1 \leq i \leq n}$, a spin structure on X is equivalent to a spin structure on $[\tilde{X}, x_1, y_1, \dots, x_n, y_n]$ plus some gluing data. Therefore $\mathcal{S}_{g,2n}$ occurs naturally in the boundary of $\overline{\mathcal{S}}_{g+n}^+$ and $\overline{\mathcal{S}}_{g+n}^-$, up to a finite factor given by the gluing data, and the problem of describing their birational geometry is quite natural.

The problem of the Kodaira classification of \mathcal{M}_g has a rich history. It was known from [Sev15] that \mathcal{M}_g is unirational for $g \leq 10$, and great progress has been made in recent years. Currently, we know from [Ser81], [CR84] and [Ver05] that \mathcal{M}_g is unirational for $g \leq 14$ and that it is rationally connected for $g = 15$. In the other direction, by [HM82] and [EH87], $\overline{\mathcal{M}}_g$ is of general type for $g \geq 24$. In the recent paper [FJP20], the long-lasting open cases of $g = 22$ and $g = 23$ have been shown to be of general type. Naturally, the same problems have been posed for $\overline{\mathcal{M}}_{g,n}$, and Logan in [Log03] shows that all but finitely many $\overline{\mathcal{M}}_{g,n}$ are of general type for $g \geq 4$. There remain, nonetheless, some open cases, and are especially worth mentioning the cases of $16 \leq g \leq 21$.

Date: February 20, 2025.

On the other hand, for moduli spaces of spin curves $\overline{\mathcal{S}}_g^\pm$, the Kodaira classification has been completed by Farkas and Verra in the papers [Far10], [FV12] and [FV14]. For $\overline{\mathcal{S}}_g^+$, it has been shown that it is uniruled for $g \leq 7$, of Kodaira dimension 0 for $g = 8$ and of general type for $g \geq 9$, whereas $\overline{\mathcal{S}}_g^-$ is uniruled for $g \leq 11$ and of general type for $g \geq 12$.

This paper is concerned with the uniruledness of some of the moduli spaces of pointed spin curves $\mathcal{S}_{g,2n}$. More precisely, we show the following:

Theorem 1. *The moduli spaces $\mathcal{S}_{2,4}$, $\mathcal{S}_{2,6}$, $\mathcal{S}_{3,2}$, $\mathcal{S}_{3,4}$, $\mathcal{S}_{3,6}$, $\mathcal{S}_{4,2}$, $\mathcal{S}_{4,4}$, $\mathcal{S}_{5,2}$ and $\mathcal{S}_{5,4}$ are uniruled.*

The method we employ to show the uniruledness of $\mathcal{S}_{g,2n}$ for small g and n relies on the results of Lelli-Chiesa, Knutsen and Verra in [LCKV23], which revolve around Nikulin surfaces of non-standard type. In that paper, the moduli spaces $\mathcal{R}_{g,2n}$ of double covers of smooth curves of genus g ramified at $2n$ points are shown to be (uni)rational for $2 \leq g \leq 5$ and n sufficiently small.

2. PRELIMINARIES

Let us now recall some definitions.

Definition 2. A *Nikulin surface of genus h* is the datum of (S, M, H) where (S, H) is a $K3$ surface of genus h and $M \in \text{Pic}(S)$ is a line bundle on S such that $2M \sim N_1 + \dots + N_8$ for eight disjoint (-2) -curves on S . Write

$$\mathbf{N} := \bigoplus \mathbb{Z}[N_i] \subset \text{Pic}(S),$$

for the *Nikulin lattice* and set $\Lambda_h := \mathbf{N} \oplus \mathbb{Z}[H]$. A Nikulin surface is of *standard type* if the embedding $\Lambda_h \subset \text{Pic}(S)$ is primitive, i.e. $\text{Pic}(S)/\Lambda_h$ is torsion-free. Otherwise it is called of *non-standard type*.

It is shown in [vGS07] that if (S, M, H) is a Nikulin surface of non-standard type, then the genus h is odd and $\Lambda_h \subset \text{Pic}(S)$ has index two.

Based on this observation, we introduce the following definition. Let $R, R' \in \text{Pic}(S)$ be such that, up to reordering,

$$R = \frac{H - N_1 - N_2}{2}, \quad R' = \frac{H - N_3 - \dots - N_8}{2} \quad \text{if } h \equiv 3 \pmod{4}$$

and

$$R = \frac{H - N_1 - N_2 - N_3 - N_4}{2}, \quad R' = \frac{H - N_5 - N_6 - N_7 - N_8}{2} \quad \text{if } h \equiv 1 \pmod{4}.$$

Set $g := g(R)$ and $g' := g(R')$ so that by adjunction $g = \frac{h+1}{4}$ and $g' = \frac{h-3}{4}$ if $h \equiv 3 \pmod{4}$, and $g = g' = \frac{h-1}{4}$ if $h \equiv 1 \pmod{4}$.

We will retain the notation in [LCKV23]. To recall, let $\mathcal{F}_h^{\mathbf{N}, \text{ns}}$ be the moduli space of Nikulin surfaces of non-standard type of genus h . A slight modification of this moduli space, introduced to address the fact that R and R' are numerically identical in the case $h \equiv 1 \pmod{4}$, is the moduli space

$$\widehat{\mathcal{F}}_h^{\mathbf{N}, \text{ns}} := \left\{ (S, M, H, R) : (S, M, H) \in \mathcal{F}_h^{\mathbf{N}, \text{ns}} \text{ and } R \in \text{Pic}(S) \right. \\ \left. \text{such that } H - 2R \text{ is a sum of } (-2)\text{-curves} \right\}$$

It is clear that $\widehat{\mathcal{F}}_h^{\mathbf{N},\text{ns}}$ is an étale double cover of $\mathcal{F}_h^{\mathbf{N},\text{ns}}$.

Define the following projective bundles over these varieties. Over $\mathcal{F}_h^{\mathbf{N},\text{ns}}$ we let

$$\mathcal{P}_g := \{(S, M, H, X) : (S, M, H) \in \mathcal{F}_h^{\mathbf{N},\text{ns}} \text{ and } X \in |R|\}$$

and

$$\mathcal{P}'_{g'} := \{(S, M, H, X') : (S, M, H) \in \mathcal{F}_h^{\mathbf{N},\text{ns}} \text{ and } X' \in |R'|\},$$

and over $\widehat{\mathcal{F}}_h^{\mathbf{N},\text{ns}}$ we let

$$\widehat{\mathcal{P}}_g := \{(S, M, H, R, X) : (S, M, H, R) \in \widehat{\mathcal{F}}_h^{\mathbf{N},\text{ns}} \text{ and } X \in |R|\}.$$

Finally, let $r_{g,2} : \mathcal{P}_g \rightarrow \mathcal{R}_{g,2}$, $r'_{g',6} : \mathcal{P}'_{g'} \rightarrow \mathcal{R}_{g',6}$ and $r_{g,4} : \widehat{\mathcal{P}}_g \rightarrow \mathcal{R}_{g,4}$ be the maps defined as follows. Let $(S, M, H, X) \in \mathcal{P}_g$, and set $\{x_i\} = X \cap N_i$. The associated point in $\mathcal{R}_{g,2}$ is

$$r_{g,2}(S, M, H, X) = [X, x_1 + x_2, \mathcal{O}_X(M^\vee)].$$

Similarly, for $(S, M, H, X') \in \mathcal{P}'_{g'}$, we set

$$r'_{g',6}(S, M, H, X') = [X, x_1 + x_2 + x_3 + x_4 + x_5, +x_6, \mathcal{O}_X(M^\vee)]$$

and for $(S, M, H, R, X) \in \widehat{\mathcal{P}}_g$ set

$$r_{g,4}(S, M, H, R, X) = [X, x_1 + x_2 + x_3 + x_4, \mathcal{O}_X(M^\vee)].$$

Theorem 3 (cf. [LCKV23]). *The maps $r_{2,4}$, $r'_{2,6}$, $r_{3,2}$, $r_{3,4}$, $r'_{3,6}$, $r_{4,2}$, $r_{4,4}$, $r_{5,2}$ and $r_{5,4}$ are dominant.*

2.1. Irreducibility of $\mathcal{S}_{g,2n}$. This subsection is devoted to proving the following result.

Proposition 4. *The moduli spaces of pointed spin curves $\mathcal{S}_{g,2n}$ are irreducible for $n > 0$.*

Before starting the proof, let us recall some definitions and notation regarding $\overline{\mathcal{S}}_g^\pm$. A *quasi-stable $2n$ -pointed curve* is a pointed curve $[X, x_1, \dots, x_{2n}]$ obtained by blowing up some of the nodes of a nodal pointed curve at most once. The rational components arising in this way are called *exceptional components*. A *stable spin curve* of genus g is a triple $[X, \eta, \beta]$ where X is a quasi-stable curve of genus g , η is a line bundle on X of degree $g-1$ such that $\eta|_E = \mathcal{O}_E(1)$ for every exceptional component E of X , and $\beta : \eta^{\otimes 2} \rightarrow \omega_X$ is homomorphism of line bundles on X which is generically non-zero on the non-exceptional components.

Finally, recall the divisors $A_0^\pm \subset \overline{\mathcal{S}}_g^\pm$, whose generic point is an irreducible even (resp. odd) spin curve $[\widetilde{X}/x_1 \sim x_2, \eta]$. They are irreducible by [Cor89].

Proof. We proceed by induction on n . If $n = 1$, consider the map from $A_0^+ \cup A_0^-$ to $\mathcal{S}_{g,2/\mathbb{Z}_2}$ given by

$$[X, \eta] \mapsto [\widetilde{X}, x_1 + x_2, \nu^* \eta(-x_1 - x_2)],$$

where $\nu : \widetilde{X} \rightarrow X$ is the normalization map, $p \in X$ is the node and $\nu^{-1}(p) = \{x_1, x_2\}$. This is clearly surjective, and by [Cor89] (6.1), both restrictions to A_0^+ and A_0^- are surjective. As these boundary divisors are irreducible, it follows that $\mathcal{S}_{g,2/\mathbb{Z}_2}$ is irreducible as well. From this we can easily deduce the irreducibility of $\mathcal{S}_{g,2}$. Let $\gamma : [0, 1] \rightarrow \mathcal{S}_{g,2/\mathbb{Z}_2}$ be a non-contractible loop based at $[X, x_1 + x_2, \eta]$, $\gamma(t) = [X(t), x_1(t) + x_2(t), \eta_t]$. This can be lifted to a path $\gamma'(t)$ from $[X, x_1, x_2, \eta]$ to

$[X, x_2, x_1, \eta]$ by continuously lifting $x_1(t) + x_2(t) \in X(t)_2$ to $X(t) \times X(t)$. Therefore, the monodromy of $\mathcal{S}_{g,2} \rightarrow \mathcal{S}_{g,2/\mathbb{Z}_2}$ is transitive, implying that $\mathcal{S}_{g,2}$ is irreducible.

As in [Cor89], we can construct a compactification $\overline{\mathcal{S}}_{g,2n}$ of $\mathcal{S}_{g,2n}$ by means of stable pointed spin curves. Consider the inclusion

$$\begin{aligned} \mathcal{C}^1\mathcal{S}_{g,2n-2} &:= \overline{\mathcal{M}}_{g,1} \times_{\overline{\mathcal{M}}_g} \overline{\mathcal{S}}_{g,2n-2} \subset \overline{\mathcal{S}}_{g,2n}, \\ [X, x_1, \dots, x_{2n-2}, t_1, \eta_X] &\mapsto [X \cup_{t_1 \sim q_1} E \cup_{t_2 \sim q_2} \mathbb{P}^1, x_1, \dots, x_{2n}, \eta] \end{aligned}$$

where $\{[\mathbb{P}^1, x_{2n-1}, x_{2n}, t_2]\} = \mathcal{M}_{0,3}$, E is the exceptional component, and η is given by $\eta|_X = \eta_X$, $\eta|_E = \mathcal{O}_E(1)$ and $\eta|_{\mathbb{P}^1} = \mathcal{O}(-2)$. Suppose that $\overline{\mathcal{S}}_{g,2n}$ is not irreducible. On one hand, for a generic point $[X, x_1, \dots, x_{2n-2}, t_1, \eta_X] \in \mathcal{C}^1\overline{\mathcal{S}}_{g,2n}$ the corresponding point in $\overline{\mathcal{S}}_{g,2n}$ is smooth, on the other hand, $\mathcal{C}^1\overline{\mathcal{S}}_{g,2n-2}$ is irreducible by induction, so it is contained in the intersection of all the irreducible components of $\overline{\mathcal{S}}_{g,2n}$. This implies that every point in the image is singular, a contradiction. \square

3. PROOF OF THE MAIN THEOREM

This section contains the proof of the main theorem.

Proof of Theorem 1. We proceed case by case. Throughout $[X, x_1, \dots, x_{2n}, \eta] \in \mathcal{S}_{g,2n}$ is a generic spin curve and $\mathcal{O}_X(Z)$ is an odd theta characteristic on X with corresponding effective divisor Z .

$g = 2$ and $2n = 4$: Since the map $r_{2,4} : \widehat{\mathcal{P}}_2 \rightarrow \mathcal{R}_{2,4}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H, R) \in \widehat{\mathcal{F}}_9^{\mathbf{N},\text{ns}}$ such that $X \in |R|$, $\{x_i\} = X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. Consider the linear system $|R \otimes \mathcal{I}_Z|$, and as $h^0(S, R) = g+1$ and $\deg Z = g-1$, it follows that there exists a pencil $P \subset |R \otimes \mathcal{I}_Z|$ containing X . The map

$$P \dashrightarrow \mathcal{S}_{2,4}, \quad X' \mapsto [X', x'_1, x'_2, x'_3, x'_4, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ exhibits a rational curve through $[X, x_1, x_2, \eta]$.

$g = 2$ and $2n = 6$: Since the map $r'_{2,6} : \mathcal{P}'_2 \rightarrow \mathcal{R}_{2,6}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H) \in \mathcal{F}_{11}^{\mathbf{N},\text{ns}}$ such that $X \in |R'|$, $\{x_i\} = X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. Consider the linear system $|R' \otimes \mathcal{I}_Z|$, and as before there exists a pencil $P \subset |R' \otimes \mathcal{I}_Z|$ containing X . The map

$$P \dashrightarrow \mathcal{S}_{2,6}, \quad X' \mapsto [X', x'_1, x'_2, x'_3, x'_4, x'_5, x'_6, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ gives a rational curve through $[X, x_1, x_2, x_3, x_4, x_5, x_6, \eta]$.

$g = 3$ and $2n = 2$: Since the map $r_{3,2} : \mathcal{P}_3 \rightarrow \mathcal{R}_{3,2}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H) \in \mathcal{F}_{11}^{\mathbf{N},\text{ns}}$ such that $X \in |R|$, $\{x_i\} = X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. Considering the linear system $|R \otimes \mathcal{I}_Z|$, there exists a pencil $P \subset |R \otimes \mathcal{I}_Z|$ which contains X . The map

$$P \dashrightarrow \mathcal{S}_{2,4}, \quad X' \mapsto [X', x'_1, x'_2, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ exhibits a rational curve through $[X, x_1, x_2, \eta]$.

$g = 3$ and $2n = 4$: Since the map $r_{3,4} : \widehat{\mathcal{P}}_3 \rightarrow \mathcal{R}_{3,4}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H, R) \in \widehat{\mathcal{F}}_{13}^{\mathbf{N},\text{ns}}$ such that $X \in |R|$, $\{x_i\} =$

$X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. As before, there exists a pencil $P \subset |R \otimes \mathcal{I}_Z|$ containing X . The map

$$P \dashrightarrow \mathcal{S}_{3,4}, \quad X' \mapsto [X', x'_1, x'_2, x'_3, x'_4, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ exhibits a rational curve through $[X, x_1, x_2, x_3, x_4, \eta]$.

$g = 3$ and $2n = 6$: Since the map $r'_{3,6} : \mathcal{P}'_3 \rightarrow \mathcal{R}_{3,6}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H) \in \mathcal{F}_{15}^{\mathbf{N}, \text{ns}}$ such that $X \in |R'|$, $\{x_i\} = X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. Considering the linear system $|R' \otimes \mathcal{I}_Z|$, there exists a pencil $P \subset |R' \otimes \mathcal{I}_Z|$ containing X . The map

$$P \dashrightarrow \mathcal{S}_{3,6}, \quad X' \mapsto [X', x'_1, x'_2, x'_3, x'_4, x'_5, x'_6, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ exhibits a rational curve through $[X, x_1, x_2, x_3, x_4, x_5, x_6, \eta]$.

$g = 4$ and $2n = 2$: Since the map $r_{4,2} : \mathcal{P}_4 \rightarrow \mathcal{R}_{4,2}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H) \in \mathcal{F}_{15}^{\mathbf{N}, \text{ns}}$ such that $X \in |R|$, $\{x_i\} = X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. Considering again the linear system $|R \otimes \mathcal{I}_Z|$, there exists a pencil $P \subset |R \otimes \mathcal{I}_Z|$ containing X . The map

$$P \dashrightarrow \mathcal{S}_{4,2}, \quad X' \mapsto [X', x'_1, x'_2, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ exhibits a rational curve through $[X, x_1, x_2, \eta]$.

$g = 4$ and $2n = 4$: Since the map $r_{4,4} : \widehat{\mathcal{P}}_4 \rightarrow \mathcal{R}_{4,4}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H, R) \in \widehat{\mathcal{F}}_{17}^{\mathbf{N}, \text{ns}}$ such that $X \in |R|$, $\{x_i\} = X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. Looking at the linear system $|R \otimes \mathcal{I}_Z|$, there exists a pencil $P \subset |R \otimes \mathcal{I}_Z|$ containing X . The map

$$P \dashrightarrow \mathcal{S}_{4,4}, \quad X' \mapsto [X', x'_1, x'_2, x'_3, x'_4, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ exhibits a rational curve through $[X, x_1, x_2, x_3, x_4, \eta]$.

$g = 5$ and $2n = 2$: Since the map $r_{5,2} : \mathcal{P}_5 \rightarrow \mathcal{R}_{5,2}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H) \in \mathcal{F}_{19}^{\mathbf{N}, \text{ns}}$ such that $X \in |R|$, $\{x_i\} = X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. Considering the linear system $|R \otimes \mathcal{I}_Z|$, there exists a pencil $P \subset |R \otimes \mathcal{I}_Z|$ containing X . The map

$$P \dashrightarrow \mathcal{S}_{2,4}, \quad X' \mapsto [X', x'_1, x'_2, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ exhibits a rational curve through $[X, x_1, x_2, \eta]$.

$g = 5$ and $2n = 4$: Since the map $r_{5,4} : \widehat{\mathcal{P}}_5 \rightarrow \mathcal{R}_{5,4}$ is dominant, there exists a non-standard Nikulin surface $(S, M, H, R) \in \widehat{\mathcal{F}}_{21}^{\mathbf{N}, \text{ns}}$ such that $X \in |R|$, $\{x_i\} = X \cap N_i$ and $\eta \otimes \mathcal{O}_X(-Z) = \mathcal{O}_X(M^\vee)$. Considering the linear system $|R \otimes \mathcal{I}_Z|$, there exists a pencil $P \subset |R \otimes \mathcal{I}_Z|$ containing X . The map

$$P \dashrightarrow \mathcal{S}_{5,4}, \quad X' \mapsto [X', x'_1, x'_2, x'_3, x'_4, M^\vee \otimes \mathcal{O}_{X'}(Z)]$$

where $\{x'_i\} = X' \cap N_i$ exhibits a rational curve through $[X, x_1, x_2, x_3, x_4, \eta]$. \square

Acknowledgements. I would like to thank Prof. Gavril Farkas for suggesting this problem to me and all the valuable discussions and advice that he gave me along the way.

REFERENCES

- [Ati71] Michael F. Atiyah, *Riemann surfaces and spin structures*, Ann. Sci. Éc. Norm. Supér. (4) **4** (1971), 47–62.
- [Cor89] Maurizio Cornalba, *Moduli of curves and theta-characteristics*, Proceedings of the first college on Riemann surfaces held in Trieste, Italy, November 9 - December 18, 1987, Teaneck, NJ: World Scientific Publishing Co., 1989, pp. 560–589.
- [CR84] Mei-Chu Chang and Ziv Ran, *Unirationality of the moduli spaces of curves of genus 11, 13 (and 12)*, Invent. Math. **76** (1984), 41–54.
- [EH87] David Eisenbud and Joe Harris, *The Kodaira dimension of the moduli space of curves of genus ≥ 23* , Invent. Math. **90** (1987), 359–387.
- [Far10] Gavril Farkas, *The birational type of the moduli space of even spin curves*, Advances in Mathematics **223** (2010), no. 2, 433–443.
- [FJP20] Gavril Farkas, David Jensen, and Sam Payne, *The kodaira dimensions of $\overline{\mathcal{M}}_{22}$ and $\overline{\mathcal{M}}_{23}$* , arXiv preprint arXiv:2005.00622 (2020).
- [FV12] Gavril Farkas and Alessandro Verra, *Moduli of theta-characteristics via nikulin surfaces*, Mathematische Annalen **354** (2012), no. 2, 465–496.
- [FV14] ———, *The geometry of the moduli space of odd spin curves*, Ann. Math. (2) **180** (2014), no. 3, 927–970.
- [HM82] Joe Harris and David Mumford, *On the Kodaira dimension of the moduli space of curves*, Invent. Math. **67** (1982), 23–86.
- [LCKV23] Margherita Lelli-Chiesa, Andreas Leopold Knutsen, and Alessandro Verra, *(uni)rational parametrizations of $\overline{\mathcal{R}}_{g,2}$, $\overline{\mathcal{R}}_{g,4}$ and $\overline{\mathcal{R}}_{g,6}$ in low genera*, arXiv preprint arXiv:2310.16635 (2023).
- [Log03] Adam Logan, *The Kodaira dimension of moduli spaces of curves with marked points.*, Am. J. Math. **125** (2003), no. 1, 105–138.
- [Mum71] David Mumford, *Theta characteristics of an algebraic curve*, Ann. Sci. Éc. Norm. Supér. (4) **4** (1971), 181–192.
- [Ser81] Edoardo Sernesi, *L'unirazionalità della varietà dei moduli delle curve di genere dodici*, Ann. Sc. Norm. Super. Pisa, Cl. Sci., IV. Ser. **8** (1981), 405–439.
- [Sev15] Francesco Severi, *Sulla classificazione delle curve algebriche e sul teorema d'esistenza di riemann*, Rendiconti della R. Accad. Naz. Lincei (1915), no. 24, 877–888.
- [Ver05] Alessandro Verra, *The unirationality of the moduli spaces of curves of genus 14 or lower*, Compos. Math. **141** (2005), no. 6, 1425–1444.
- [vGS07] Bert van Geemen and Alessandra Sarti, *Nikulin involutions on $k3$ surfaces*, Mathematische Zeitschrift **255** (2007), no. 4, 731–753.

INSTITUT FÜR MATHEMATIK, HUMBOLDT UNIVERSITÄT ZU BERLIN UNTER DEN LINDEN 6, 10099 BERLIN, GERMANY

Email address: bogdan-petru.carasca@hu-berlin.de